The relationship between oxytocin pathway genes and personality traits and psychosis characteristics

Marit Haram MD

Division of Mental Health and Addiction, Oslo University Hospital

and

Institute of Clinical Medicine, University of Oslo

Oslo, 2016

© Marit Haram, 2017

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-002-5

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

Table of contents

1. ACKNOWLEDGEMENTS.....	5
2. LIST OF STUDIES.....	7
3. ABBREVIATIONS.....	8
4. ABSTRACT.....	10
5. INTRODUCTION.....	13
5.1 Psychotic Disorders.....	13
5.1.1 History.....	13
5.1.2 Definition.....	14
5.1.3 Human genetics.....	16
5.1.4 Etiology.....	17
5.2 Oxytocin.....	18
5.2.1 Structure and function.....	18
5.2.2 Structure of oxytocin pathway genes.....	21
5.2.3 Study methods and theoretical framework.....	22
5.3 Personality traits.....	27
5.3.1 Definition and correlates with psychotic disorders.....	27
5.3.2 Genetic modulation.....	28
5.3.3 Oxytocin.....	28
5.4 Positive and negative symptoms.....	30
5.4.1 Symptom course and treatment.....	30
5.4.2 Oxytocin.....	31
5.5 Affective face perception in psychotic disorders.....	33
5.5.1 Definition.....	33
5.5.2 Amygdala activation.....	33
5.5.3 Oxytocin.....	35
5.6 Aims of the thesis.....	36
6. MATERIAL AND METHODS.....	38
6.1 Participants.....	38
6.1.1 Patient sample.....	38
6.1.2 Healthy controls.....	40

6.2 Clinical Assessment.....	41
6.2.1 Diagnostic assessment.....	41
6.2.2 Assessment of positive and negative symptoms.....	41
6.2.3 Assessment of personality traits.....	42
6.3 Selection of genetic variants.....	43
6.4 Genotyping and Imputation.....	44
6.5 Polygenic risk score.....	45
6.6 fMRI analyses.....	45
6.6.1 Experimental paradigm.....	45
6.6.2 BOLD fMRI data acquisition	46
6.6.3 fMRI Data Analysis.....	46
6.7 Statistics.....	47
7. SUMMARY OF RESULTS.....	50
8. DISCUSSION.....	53
8.1. Main results.....	53
8.1.1 The relationship between personality traits and oxytocin pathway genes.....	53
8.1.2 The effect of oxytocin pathway genes on positive and negative symptoms and diagnosis in psychotic disorders.....	55
8.1.3 The effect of oxytocin pathway genes on affective face perception in psychotic disorders.....	59
8.1.4 Oxytocin pathway genes – biological modulators of social dysfunction in psychotic disorders?.....	60
8.2. Strength and limitations.....	66
8.2.1 Sample and assessments.....	66
8.2.2 Methods.....	67
8.3. Future directions.....	70
9. CONCLUSION.....	72
10. REFERENCES.....	74

Appendix

Errata

1. ACKNOWLEDGMENTS

The work in this thesis was carried out at NORMENT, the KG Jebsen Center for psychosis research, Division of Mental Health and Addiction, Oslo University Hospital, and the Institute of Clinical Medicine, University of Oslo. The present studies were part of the larger TOP (Thematically Organized Psychosis) study, and were supported by the Kristian Gerhard Jebsen Foundation, the Research Council of Norway, and the Regional Health Authority for South-Eastern Norway.

First and foremost, I would like to give my deepest gratitude to the participants involved in this project. Despite of often challenging life circumstances, they have with patience and altruism completed demanding interviews and tests. Without their endurance, this work could never have taken place. I am sincerely grateful to my main supervisor Martin Tesli, who has guided me safely through the field of psychiatric genetics. His enthusiasm and impressive ability to find overview in a complex research field has been invaluable. With steady academic supervision and everlasting support, he is an example to follow. I am forever thankful to my co supervisor Ingrid Melle, who has introduced me to all levels of psychiatric research. With continuous warm encouragement, always available for guidance, she has shown me a way through the realm of clinical, biological, environmental and statistical sides of this research field. Her care for the patients, massive experience and knowledge has been an enormous inspiration in both the research- and clinical work. I am sincerely thankful to my co supervisor Ole A. Andreassen who has supported me through all aspects of this research project. With an impressive ability to be thorough, efficient, available, creative and enthusiastic, he has always provided me with to-the-point feedback and firm academic guidance.

I want to thank Srdjan Djurovic for always being available for questions and his kind responses. Francesco Bettella deserves a great appreciation. His kindness and programming skills are beyond imagination, and I am forever thankful for his contributions. Furthermore, I want to thank all my colleagues in the TOP research group who has created an inspiring, creative and fun working place that I am so grateful for. A special thanks to Thomas for skillful technical support, interesting discussions and many laughs, and to Ragnhild and Eivind for keeping this place in order and creating a warm atmosphere. And of course- to my good friend Mari- whom I want to thank for all the good conversations about everything and nothing, laughs, warmth and support.

I am forever grateful to my parents for support, scientific interest, patience and encouragement. And finally, the biggest hug goes to Eivind, Mikkel and Jonas- who's support I could not have done without.

2. LIST OF STUDIES

Study I

An Attempt to Identify Single Nucleotide Polymorphisms Contributing to Possible Relationships between Personality Traits and Oxytocin-Related Genes

Marit Haram, Martin Tesli, Ingrid Dieset, Nils Eiel Steen, Jan Ivar Røssberg, Srdjan Djurovic, Ole A. Andreassen, Ingrid Melle

Neuropsychobiology (2014). 69, pp.25–30

Study II

Association between genetic variation in the oxytocin receptor gene and emotional withdrawal, but not between oxytocin pathway genes and diagnosis in psychotic disorders

Marit Haram, Martin Tesli, Francesco Bettella, Srdjan Djurovic, Ole A. Andreassen, Ingrid Melle

Frontiers in Human Neuroscience (2015). 9, pp.9

Study III

Contribution of oxytocin receptor polymorphisms to amygdala activation in schizophrenia spectrum disorders

Marit Haram, Francesco Bettella, Christine Lycke Brandt, Daniel S. Quintana, Mari Nerhus, Thomas Bjella, Srdjan Djurovic, Lars T. Westlye, Ole A. Andreassen, Ingrid Melle, Martin Tesli

Accepted in British Journal of Psychiatry Open

3. ABBREVIATIONS

AD	Affective Spectrum Disorder
ASD	Autism Spectrum Disorders
AVP	gene coding for Vasopressin
BBB	the Blood Brain Barrier
BOLD	Blood-Oxygen-Level Dependent
CD38	gene coding for CD38
CSF	Cerebrospinal Fluid
CNS	Central Nervous System
COPE	Contrast Parameter Estimates
DNA	Deoxyribonucleic Acid
DRD2	gene coding for the Dopamine receptor D2
DSM	Diagnostic and Statistical Manual
eQTL	expression Quantitative Trait Loci
FEAT	FMRI Expert Analysis Tool
fMRI	functional Magnetic Resonance Imaging
GWAS	Genome Wide Association studies
ICD	International Classification of Diseases
Kb	Kilobases
LD	Linkage Disequilibrium
mRNA	messenger Ribonucleic Acid
NEO-FFI	Neuroticism-Extraversion-Openness Five-Factor Inventory
NEO-PI-R	Neuroticism Extroversion Openness Personality Inventory Revised Version
NMDA	N-Methyl-D-Aspartate
NOS	Not Otherwise Specified
OR	Odds Ratio

OXT	gene coding for oxytocin
OXTR	gene coding for the oxytocin receptor
PANSS	Positive and Negative Syndrome Scale
PGC	Psychiatric Genomics Consortium
PGRS	Polygenic Risk Score
PVN	Paraventricular Nuclei
PPI	Prepulse Inhibition
RDoC	Research Domain Criteria project
ROI	Regions of Interest
SCID-1	Structured Clinical Interview for DSM-IV Axis 1 Disorders
SCZ	Schizophrenia spectrum disorder
SNP	Single Nucleotide Polymorphisms
SON	Supraoptic Nuclei
SPECT	Single-Photon Emission Computed Tomography
TOP	Thematically Organized Psychosis Research
UCSC	University of California Santa Cruz

4. ABSTRACT

Oxytocin is a neuropeptide hormone well known for its importance in parturition and milk let down in women. An increasing amount of research has demonstrated that this hormone is involved in several aspects of social behavior, and oxytocin's role in social memory, pair bonding and parental behavior is well established in animal research. This knowledge has led to an augmented interest in oxytocin as a potential therapeutic target for psychiatric illnesses, including psychotic disorders. Several trials with intranasal oxytocin have been conducted in both healthy and patient samples, but the results from oxytocin trials are inconsistent in general. This inconsistency has been attributed to individual differences, including genetically based differences in the oxytocin system, and a poor understanding of drug targets and mechanisms.

The overall aim of the current PhD work was to identify minor genetic alterations in the oxytocin pathway genes regulating oxytocin, which could lead to deficits in features that are important for social functioning in patients with psychotic disorders, with possible implications for treatment and the understanding of pathophysiological mechanisms. We focused on common variations in genes related to oxytocin pathway, especially the genes coding for oxytocin (OXT), vasopressin (AVP), the oxytocin receptor (OXTR) and the transmembrane receptor CD38 (CD38), and investigated:

1. The association between oxytocin pathway polymorphisms and personality traits correlated with trust and social behavior in healthy subjects; Neuroticism, Extraversion and Agreeableness.
2. The association between oxytocin pathway genes and psychotic disorders per se in a case-control design as well as between oxytocin pathway genes and specific psychopathological features associated with trust and social behavior, in particular

suspiciousness/persecutory delusions, hostility, emotional withdrawal and passive/apathetic social withdrawal.

3. The association between oxytocin receptor polymorphisms and amygdala activation during affective face perception of faces expressing anger or fear, and possible disorder specific associations in patients with psychotic disorders.

We were not able to detect associations between oxytocin pathway genes and personality traits in healthy subjects. There was a significant association between symptoms of emotional withdrawal and the A allele in rs53576 located on OXTR. No significant associations between oxytocin pathway gene variants and suspiciousness/persecutory delusions, hostility or passive/apathetic social withdrawal, or a diagnosis of psychotic disorder were found. In participants with schizophrenia spectrum disorders, the rs237902 G allele was associated with low amygdala activation and interaction analyses showed that this association was disorder specific. There were no associations between oxytocin receptor polymorphisms and amygdala activation in the total sample, among patients with affective spectrum disorders or healthy controls.

Taken together, our findings indicate that i) it is less likely that oxytocinergic signaling influences the expression of the personality traits Neuroticism, Extraversion and Agreeableness ii) oxytocinergic signaling influences the expression of emotional withdrawal in patients with psychotic disorders. It is less likely that oxytocinergic signaling influences suspiciousness/persecutory delusions, hostility or passive/apathetic social withdrawal or that oxytocinergic signaling is important in the pathogenesis of psychotic disorder per se iii) oxytocinergic signaling influences amygdala activation during affective face perception in patients with schizophrenia spectrum disorders, and in a different way than in patients with

affective spectrum disorders and healthy controls. This supports the idea that alterations in the endogenous oxytocin system are present in patients with schizophrenia spectrum disorders.

5. INTRODUCTION

5.1 Psychotic disorders

5.1.1 History

Delirious behavior has been recorded since ancient times, but was long considered to be a sign of mania, a somatic illness not localized in the brain, or a psychological result of wrongdoing (Beer, 1996b). The symptoms of mania and depression were first described as separate entities, and while Hippocrates gave depressive symptoms the clinical term melancholia, Arateus of Cappadocia (1st century AD) uniformed the concepts to manic-depressive illness with a common etiology (Berrios, 1996). This understanding was not established in its current time, but rediscovered in the middle of the 19th century. In its renewal, several contributors helped define the clinical presentations, and the German psychiatrist Emil Kraepelin (1856-1926) first categorized and studied the natural course of manic-depression (Kraepelin, 1910).

The concept of psychosis as a brain disorder with mechanisms not yet discovered was first described in the late 19th century, and Kraepelin dichotomized for the first time endogenous psychosis into manic-depression and dementia praecox (Beer, 1996a). Dementia praecox was initially a term for deteriorating psychotic disorders, while manic-depression mainly was thought of as an intermittent illness, with symptom-free intervals (Adityanjee et al., 1999). During his career, Kraepelin's concept of dementia praecox developed, and he eventually acknowledged the fact that some patients recovered (Adityanjee et al., 1999).

The term schizophrenia was first introduced by Eugene Bleuler in 1908, derived from the greek words *schizein* and *phren*, denoting *splitting* and *mind* respectively (Bleuler, 1911). Bleuler criticized the term “dementia praecox”, arguing that psychotic illness onset and course varied, and did not necessarily end in disintegration (Bleuler, 1911). Inspired by Sigmund

Freud (1856-1939) and C.G. Jung (1875-1965), Bleuler focused on schizophrenia as a thought-disorder, and described four core symptoms; looseness of associations, affective flattening, autism, and ambivalence (today known as Bleuler's 4 A's) (Bleuler, 1911). In the mid 20th century, the importance of positive symptoms (i.e. delusions and hallucinations, also defined as psychotic symptoms) was emphasized, symptoms that lay the foundation for diagnostic criteria today (Nordgaard et al., 2008). In the same era, manic-depression was described in detail and a distinction between depressions with hypomania and depressions with mania was made (Kleist, 1953, Dunner et al., 1976). This division is still given in the current diagnostic manual.

Today, the WHO has classified schizophrenia as one of the world's most costly diseases, only 8% of the patients are in employment, and the average life-expectancy is 15-20 years shorter than in healthy individuals (Schizophrenia Commission, 2012). Most of the patients and their families suffer from a life-long burden of the disorder, and there are several unresolved issues regarding both treatment-efficacy and -side-effects, as well as social care.

5.1.2 Definition

The diagnostic criteria for psychotic disorders today are categorical and defined by clusters of symptoms described in the *Diagnostic and Statistical Manual of Mental Disorders (DSM)* or the *WHO International Classification of Diseases (ICD)*. Schizophrenia is primarily described by the presence of positive symptoms, and delusions with bizarre content (i.e. beliefs that are not possible physically and implausible in context of culture), weighs heavily (Cermolacce et al., 2010). Other domains like disorganized thinking, disorganized motor behavior and negative symptoms (i.e. diminished emotional expression, impaired motivation and reduction in spontaneous speech), may also be present.

Patients with positive symptoms that do not fully fulfill the criteria for schizophrenia, either because of short duration of illness, the dominance of depressive/manic episodes, not markedly loss of function or less severe symptoms, may be diagnosed with psychotic disorder not otherwise specified (NOS), schizophreniform, schizoaffective, delusional or brief psychotic disorder (American Psychiatric Association, 2000). Bipolar disorder is defined by the presence of at least one manic episode (bipolar I) or one hypomanic episode and one major depressive episode (bipolar II). Patients with affective episodes that do not fully fulfill these criteria (e.g. recurrent hypomanic episodes without depression) may be diagnosed with bipolar NOS. Patients with bipolar and also Major Depressive Disorders may experience psychotic symptoms, but in these cases always during an affective episode (American Psychiatric Association, 2000).

This strict categorization of disorders and the dichotomization of bipolar disorder and schizophrenia introduced by Kraepelin, have in the recent years been challenged (Owen et al., 2016). A rich and growing literature provide insight into the genetic overlap between these disorders (Craddock and Owen, 2010, Andreassen et al., 2013), shared clinical symptoms, neurocognitive impairments (Simonsen et al., 2010), social cognitive impairments (Gur and Gur, 2016) as well as brain abnormalities (Brandt et al., 2014, Rimol et al., 2010). For these reasons, it has been suggested that dimensional approaches to diagnoses, treatment, and research questions may replace the strict categorical approaches applied today (Owen, 2014). Recently, the National Institute of Mental Health proposed a precision medicine model for psychiatry, i.e. the Research Domain Criteria project (RDoC) (Insel, 2014). This is a research working frame encouraging investigation of the biological and psychosocial basis of core features in mental health, regardless of diagnostic categories (Insel, 2014). Subsequently, it might be possible to improve the classification of psychotic disorders in the future based on

the newest research findings, and thus provide more personalized medicine and develop novel medical treatments.

In this thesis, we wanted to investigate features with implications for social dysfunction often occurring in patients with psychotic disorders, and often with most severe impairments in schizophrenia. Based on the theory that these disorders could be better classified with a dimensional approach, we investigated these features in both healthy individuals as well as in a wide range of diagnoses (i.e. schizophrenia, psychotic disorder NOS, schizophreniform, schizoaffective, delusional and brief psychotic disorder, bipolar disorder I, II and NOS, Major Depressive disorders with psychotic symptoms), hereby named “psychotic disorders”.

5.1.3 Human genetics

In 2001, the human genome was mapped for the first time (Lander et al., 2001), and since then about 21 000 protein coding genes have been discovered (Pennisi, 2012). Not more than approximately 0,2 % of the genes mapped differ between human individuals, and a large proportion of this variance is by common alterations in the single base-pairs, with population frequencies above 1%, also known as single nucleotide polymorphisms (SNPs) (Nature.com, 2016, Sebat, 2007). Complex traits (like diabetes, cardiovascular diseases, psychiatric disorders and personality traits) are polygenic, meaning that several genetic variants contribute to the trait, each with small effects (Dudbridge, 2016). The measurement of genetically mediated differences will then provide us with important information about underlying biological mechanisms to the trait of investigation.

SNPs tend to be inherited together in clusters, and linkage disequilibrium (LD) is defined as the phenomenon of two or more alleles at different loci having a stronger association than what would be expected to occur if they were randomly combined (Slatkin, 2008). LD could

be expressed by D' or r^2 (both ranging from 0 to 1), and a value of 0 indicates linkage equilibrium, i.e. statistical independence, while a value of 1 indicates complete/perfect LD, i.e. statistical dependence. A tag SNPs is then defined as a polymorphism that captures other polymorphisms by LD and can be investigated to reduce the number of independent test. The Affymetrix Chip is a DNA microarray that has predefined SNPs to test, and serve as an efficient way of investigating multiple polymorphisms. Due to Affymetrix's selection protocol, only half of the variants on their array are tag SNPs. However, the remaining polymorphisms have been shown to provide high genomic coverage (Perkel, 2008).

5.1.4 Etiology

The etiology of psychotic disorders is complex, mostly unknown, and subject to continuous research (Owen et al., 2016). The extensive contribution of genetic factors has been acknowledged for over 50 years (Gottesman and Shields, 1967), and a biopsychosocial model which includes environmental risk factors has been a theoretical research frame for several decades (Engel, 1977). Even though this “stress-vulnerability” theory has been difficult to prove (van Os et al., 2010), there is convincing evidence that socioeconomic risk factors such as childhood adversity and migration are strongly associated with the risk of developing a psychotic disorder (Varese et al., 2012, Cantor-Graae and Pedersen, 2013).

Based on family studies, the heritability of psychotic disorders is found to be approximately 60-80% (Bienvenu et al., 2011, Lichtenstein et al., 2009). With new genotyping techniques and advanced statistical methods emerging rapidly, the identification of susceptibility genes are continuously aimed for, hoping to shed light on the biological pathways involved in the disorders. Genome wide association studies (GWAS) are theoretically capable of identifying several relevant SNPs with a *hypothesis-free* approach, as the whole genome is screened for

association with a particular phenotype (Zondervan and Cardon, 2007). This method has been applied to large European samples (36,989 cases with schizophrenia and 113,075 controls involved in the Psychiatric Genomics Consortium (PGC2)) and 108 susceptibility loci for schizophrenia are detected. Immune-related genes seem to contribute to a great extent (Schizophrenia Working Group of the Psychiatric Genomics, 2014). However, the currently identified susceptibility loci account for only a few percent of the estimated heritability, leaving a large “missing heritability”. Dopamine has been hypothesized to be involved in the pathogenesis for over 60 years, and the first dopamine related gene (the dopamine receptor D2 gene, DRD2) was just recently detected as a susceptibility gene (Schizophrenia Working Group of the Psychiatric Genomics, 2014). Previous studies with smaller samples involved were however not able to detect this gene, illustrating the need for enormous samples to detect differences between patients and controls (Genome-wide association study identifies five new schizophrenia loci, 2011a). Several other explanations for this missing heritability have been suggested, such as the need for new statistical tools (Andreassen et al., 2014), gene-gene and gene-environment interactions as well as epigenetic factors (Harrison, 2015).

5.2 Oxytocin

5.2.1 Structure and function

Oxytocin is a polypeptide released from the posterior pituitary gland, and is interestingly active both as a peripheral hormone and a neuropeptide in the central nervous system (CNS). Fairly all vertebrates have a related peptide represented, and oxytocin’s important role in parturition and milk let-down has been known for over 60 years (du Vigneaud, 1960, Gimpl and Fahrenholz, 2001). Oxytocin consists of nine amino acids, differing from vasopressin, a related neuropeptide also released from the posterior pituitary gland with only two amino acid positions (National Center for Biotechnology Information, 2010). This structural difference

enables each peptide to bind to its corresponding receptor, but at the same time display overlapping pharmacological properties (Gimpl and Fahrenholz, 2001).

The oxytocin receptor belongs to the G-coupled protein receptor family and is a polypeptide consisting of 388 amino acids (Kimura et al., 1992). With seven transmembrane domains, an external signal of oxytocin activates a G protein inside the cell, which in turn affects further signal pathways and transduces cell functioning (Qin et al., 2011). The oxytocin receptor is distributed widely throughout the brain, including the olfactory system, cortical areas, basal ganglia, limbic system, thalamus, hypothalamus and brain stem as well as the body, including the uterus, ovary, testis, prostate gland, thymus, pancreas, the cardiovascular system and mammary tissues (Gimpl and Fahrenholz, 2001, Loup et al., 1991, Loup et al., 1989).

Oxytocin for peripheral release is mainly synthesized in magnocellular neurons in the hypothalamic paraventricular (PVN) and supraoptic nuclei (SON). After transportation to the pituitary gland it is released to the blood stream and act as a hormone throughout the body.

Oxytocin for central release is mainly synthesized in parvocellular neurons in the PVN, and terminate in different areas of the brain through axonal transport (McEwen, 2004b) (Figure 1), but there is to some extent also dendritic release of oxytocin from magnocellular neurons into central regions, which is possibly self-sustaining and long-lasting (Ludwig, 1998, Ludwig and Leng, 2006). Functionally, oxytocin has been proposed to act as a modulator in synaptic transmission in brain areas involved in social behavior (Mitre et al., 2016).

Up- and down-regulation of the oxytocin receptor is highly dynamic, complex and tissue specific processes, and factors in the promotor region of the oxytocin receptor gene as well as epigenetic factors matter (Kimura et al., 2003). We know that psychological stressors induce secretion of oxytocin, both peripherally and centrally (Ludwig, 1998, Nishioka et al., 1998).

Pregnancy, parturition, suckling and positive social interaction are probably important central

regulators (Landgraf et al., 1992, Uvnas-Moberg, 1998). Furthermore, estrogen can modulate the transcription of oxytocin receptors in the amygdala, suggesting that steroids are important for central regulation (Choleris et al., 2003). At a molecular level, the trans-membrane receptor CD38, that is recognized as important for several complex system, e.g. the reproductive-, immune- and cardiovascular system (Malavasi et al., 2008), has recently been found to be important for central release of oxytocin and hence behavioral effects (Jin et al., 2007, Munesue et al., 2010).

It is not clear whether the peripheral and central secretion of oxytocin are synchronous (Gimpl and Fahrenholz, 2001, Ludwig and Leng, 2006, Leng et al., 2008, Neumann et al., 1993), and most likely the hormone does not cross the blood brain barrier (BBB) (McEwen, 2004a).

Additionally, oxytocin receptor ligands are just emerging and have not been implemented in human research yet (Karpenko et al., 2015, Smith et al., 2016), making the investigation of human central pathways of oxytocin challenging (Leng et al., 2015, Leng and Ludwig, 2016).

The rich literature from animal studies however applies convincing evidence for oxytocin's important role in social behavior, and several study methods in humans have been introduced (Meyer-Lindenberg et al., 2011, Leng et al., 2015).

Figure 1

The site for production, release and central pathways of oxytocin. Oxytocin travels in the brain via axonal projections and dendritic release into several brain areas. Here they act as neuromodulators and affect neurotransmissions. The oxytocin receptor sites are mainly mapped through animal research and are not fully elucidated (Meyer-Lindenberg et al., 2011, Grinevich et al., 2016). ACC = anterior cingulate cortex; BNST = bed nucleus of the stria terminalis; NAc = nucleus accumbens; SCN = suprachiasmatic nucleus. Reprinted by permission from Macmillan Publishers Ltd: Nature Reviews Neuroscience (Meyer-Lindenberg et al., 2011), copyright 2011.

5.2.2 Structure of oxytocin pathway genes

The oxytocin pathway genes are often defined as the genes coding for oxytocin (OXT), vasopressin (AVP), the oxytocin receptor (OXTR), and CD38 (CD38) (Feldman et al., 2016). AVP is of interest since its gene location is separated from OXT with only 12 kilobases (kb), and produces hormones with similar polypeptide structure (Manning et al., 2008, Wallis, 2012). OXT and AVP are located on chromosome 20, and positioned in opposite

transcriptional directions (Summar et al., 1990). Both genes have three coding exons each with transcriptions lengths of only 898 and 2170 base pairs, respectively (Ensembl.org., 2016, Flicek et al., 2014). The OXTR spans over 19.22 kb and consist of 4 exons and 3 introns. Coding regions are located in exon 3 and 4 alone, and the largest gene region is the third intron which spans over approximately 12 kb (Inoue et al., 1994). CD38 is located at chromosome 4. With a span of 74.96 kb and 8 coding exons (Ensembl.org., 2016, Flicek et al., 2014), this is the largest gene studied in the present thesis. For central effects, the gene is expressed both intracellularly and on the plasma membrane, and both in neurons and glial cells (Salmina et al., 2010).

RNA, and ultimately protein, is expressed after the reading of coding regions, but a rich literature has shown that polymorphisms in non-coding regions are of equal importance regarding protein expression, subsequently with potential clinical consequences (Pagani and Baralle, 2004). In particular deep intronic variants and synonymous SNPs in untranslated regions of the exon can affect splicing and thus be an important contributing factor in disease development (Pagani and Baralle, 2004, Sauna and Kimchi-Sarfaty, 2011).

5.2.3 Study methods and theoretical framework

Animal studies

Important information about oxytocin's behavioral effects has been revealed through animal research. Maternal behavior in rats can be induced by central injection of exogenous oxytocin, and decreased by oxytocin antagonists (Insel and Young, 2001). The difference in central oxytocin receptor distribution between prairie voles and montane voles is interesting as prairie voles demonstrate a "partner preference" after mating, and montane voles do not. After giving birth however, the distribution of oxytocin receptors change temporarily in the montane vole

brain (Insel and Young, 2001). The formation of close bonds with relatives in rodents could serve as a correlate with behavior related to social cognition (Rosenfeld et al., 2011), and interestingly, mice that are genetically engineered to lack the gene for oxytocin (oxytocin knockout mice) have a deficit in recognizing familiar mice, which can be recovered by infusion of oxytocin into the medial amygdala (Ferguson et al., 2001). CD38 knockout mice also show defects in maternal nurturing and social behavior (Jin et al., 2007).

Based on the knowledge from animal literature, a sizeable number of studies investigating the role of oxytocin in human social behavior has been conducted over the past decade. The main research tools used in human studies are measures of oxytocin in plasma/cerebrospinal fluid (CSF), treatment trials with intranasal delivery of oxytocin and measurement of genetically mediated differences.

Plasma/CSF levels

Oxytocin plasma levels have been measured in several psychiatric populations, hypothesizing that low endogenous levels are correlated with impaired social cognition and disease characteristics related to social behavior (Rubin et al., 2010, Goldman et al., 2008, Yuen et al., 2014). A recent meta-analysis involving 52 studies of plasma level (as well as urine, saliva and CSF) were however not able to detect any association with psychiatric disorders (Rutigliano et al., 2016). Together with findings of un-correlation between oxytocin levels in plasma and CSF (Leng and Ludwig, 2016), this lends support to the understanding that the plasma oxytocin levels do not reflect central effects in any certain way.

Intranasal oxytocin

Due to the challenge of identifying oxytocin biomarkers, and the limitations of the BBB, oxytocin administered intranasally has become a central research method. This administration route is hypothesized to have a direct pathway into the brain, and has over the past decade

been proposed as a possible treatment for a wide range of symptoms and deficits related to social behavior in several psychiatric disorders, including schizophrenia (Quintana et al., 2016). In 2005, Kosfeld and colleagues found that intranasal oxytocin increased trust in a money-transferring game performed by healthy subjects (Kosfeld et al., 2005) and showed for the first time that this administration route could elicit behavioral effects. Since then, numerous trials with intranasal oxytocin have been conducted, and some of the effects seen are lower anxiety levels, increased generosity, increased performance on tests of theory of mind and better ability to classify positive facial emotions correctly (Macdonald and Macdonald, 2010, Bartz et al., 2011b, Churchland and Winkielman, 2012). Brain activation measured by functional Magnetic Resonance Imaging (fMRI) during social cognitive tasks has been included as a research endpoint, and reduced activation in amygdala during presentation of fearful visual images has repeatedly been seen after intranasal oxytocin administration (Bethlehem et al., 2013). The effects on amygdala activation seem to be task-dependent, and possibly with diverse direction of effect between males and females (Shamay-Tsoory and Abu-Akel, 2016, Rilling et al., 2014, Domes et al., 2010).

The trial results have however been inconsistent (Herpertz and Bertsch, 2016), and both nonsignificant results and even adverse effects are reported (for review see (Bartz et al., 2011b)). While oxytocin initially was thought of as an unambiguously pro-social neuropeptide, effects like envy, gloating and aggression towards competitors appeared (De Dreu et al., 2010, Shamay-Tsoory et al., 2009). The suggested travel pathways from the nasal cavities to the brain (either through internalization of the peptide into olfactory neurons and thus with axonal transport to the brain, or by intercellular clefts into the subarachnoidal space) have been questioned (Leng and Ludwig, 2016), and to date it is uncertain whether intranasal oxytocin actually penetrate the brain (Bakermans-Kranenburg and van, 2013, Leng and

Ludwig, 2016). Recent critical reviews have also pointed to the small sample sizes involved in oxytocin trials, with the following low power and risk of both type I and type II errors (Walum et al., 2016, Button et al., 2013).

Oxytocin pathway genes

The measurement of genetically mediated differences has become an approach aiming to provide new knowledge of the endogenous oxytocin system in relation to social behavior and brain structure/function, as the genetic make-up is identical both in the individual's peripheral and central oxytocin system. Numerous studies have linked OXTR polymorphisms to parental attachment, empathy, theory of mind, stress response, social cognition and amygdala volume in healthy subjects (Feldman et al., 2016, Bakermans-Kranenburg and van Ijzendoorn, 2008, Rodrigues et al., 2009, Inoue et al., 2010), and to the risk for developing autism spectrum disorders (ASD) (LoParo and Waldman, 2015). Through candidate gene approaches, rs53576 and rs2254298 have received particular attention (Feldman et al., 2016). Rs53576 has additionally been shown to modulate the effect of intranasal oxytocin (Chen et al., 2015). However, recent conflicting meta-analyses disagree about their importance (Li et al., 2015, Bakermans-Kranenburg and van Ijzendoorn, 2014). While no essential associations between AVP and social behavior are shown (Meyer-Lindenberg et al., 2011), OXT and CD38 have both been associated with ASD, parental attachment and stress-coping (Ebstein et al., 2009, Ebstein et al., 2012, Feldman et al., 2016).

Theoretical framework

Initially, oxytocin was seen as uniformly related to increased trust, reduced anxiety and activation of affiliative behavior. But as ambiguous effects of exogenous oxytocin were found, new hypotheses about the neuropeptide arose. Bartz and colleagues proposed an interactionist model in 2011, where the importance of individual and contextual factors was

included (Bartz et al., 2011b). They accentuated the idea that oxytocin alters the processing of social cues, also known as the social salience hypothesis. Thus, the response of exogenous oxytocin would then be dependent on predisposed cues (Bartz et al., 2011b). In light of this theory, they saw that intranasal oxytocin showed strongest effects on emotion recognition only when the tasks were demanding for the individual (Bartz et al., 2010) and that negative emotions like mistrust, envy and attachment insecurity could be induced in relation to out-groups (Declerck et al., 2010, Bartz et al., 2011a, De Dreu, 2012, Bartz et al., 2011b, Shamay-Tsoory et al., 2009). Recent results have strengthened this interactionist model, stressing that oxytocin given in addition to social cues considered as “safe” by the participants enhances pro-social behavior, whereas anti-social emotion and behavior could be induced when the social cues are considered to be “unsafe” , or in individuals with aggressive tendencies (Olff et al., 2013, DeWall et al., 2014). Recently, Shamay and Tsoory integrated cumulative evidence from animal and human research and boosted the social salience hypothesis (Shamay-Tsoory and Abu-Akel, 2016). Building on previous research, claiming that dopamine plays an important role in the information processing of social cues (Berridge, 2007), they suggest a modulating role of oxytocin on dopamine signaling in the mesocorticolimbic system when exposed to an external cue (Shamay-Tsoory and Abu-Akel, 2016). In rodents, dopamine release has shown influenced by central oxytocin injections during pup grooming (Shahrokh et al., 2010), and stimulation of both dopamine and oxytocin receptors in the nucleus accumbens was in one study necessary for inducing pair bonding formation (Liu and Wang, 2003). Furthermore, oxytocin receptor expression in the amygdala seems dependent on dopamine (Bale et al., 2001). Although stating that more research is needed to clarify the role of oxytocin in social salience, these authors provide a theoretical framework within which new hypotheses might be generated.

5.3 Personality traits

5.3.1 Definition and correlates with psychotic disorders

The term “personality” was introduced in the late 18th century, accompanied with an attempt to understand its neuroanatomical basis (Crocq, 2013). Kraepelin emphasized the existence of a gradual limit between normal and pathological personality, thus the investigation of personality became an integrated part of research on psychiatric disorders (Kendler and Jablensky, 2011). Today, personality traits are often defined as individual characteristics in cognition, emotion and behavior that are stable over time. Several personality scales have been validated, and the Neuroticism Extroversion Openness Personality Inventory Revised Version (NEO-PI-R) is widely used (Costa and McCrae, 1992). A rich body of literature support a five-factor model of personality, and in addition to the traits; Neuroticism, Extraversion and Openness, the NEO-PI-R captures Agreeableness and Conscientiousness (Briggs, 1992).

Personality traits are shown to predict a wide range of behaviors (Hopwood et al., 2009) and have repeatedly been related to psychiatric disorders (Barnett et al., 2011). Both high scores on Neuroticism and low scores on Extraversion are shown associated with depression, anxiety, phobias, as well as a diagnosis of schizophrenia and bipolar disorder (Brandes and Bienvenu, 2006, Klein et al., 2011, Solyom et al., 1986, Van Os and Jones, 2001, Barnett et al., 2011). While low levels on Conscientiousness and Openness have been correlated with schizophrenia as well in a recent meta-analysis (Ohi et al., 2016), these traits may not predict social dysfunction as much as Neuroticism, Extraversion and Agreeableness (Hopwood et al., 2009). In patients with psychotic disorders, personality traits have been associated with symptom course and functioning levels (Gleeson et al., 2005, Compton et al., 2015) and premorbid high scores on Neuroticism and low scores on Extraversion might serve as risk

factors for the development of psychotic disorders (Van Os and Jones, 2001, Lonnqvist et al., 2009). The biological basis of personality traits is largely unknown, but the strong associations found between extreme personality scores and psychotic disorders implies that the NEO-PI-R measures features that have shared underlying biological mechanisms.

5.3.2 Genetic modulation

Based on twin, family and adoption studies, the heritability of personality traits is estimated to be approximately 40% (Vukasovic and Bratko, 2015), and several GWAS for personality traits have been conducted (Balestri et al., 2014). The results are in general mixed (Montag and Reuter, 2014), and most associations found have failed to be replicated (Kim et al., 2013, Terracciano et al., 2010, Kim et al., 2015). As with psychotic disorders, this “missing heritability” has encouraged researchers to perform other complementary genetic approaches, such as using new statistical tools on the GWAS data sets (Kim et al., 2015, Power and Pluess, 2015, Montag and Reuter, 2014), and candidate gene studies based on hypotheses about neurobiological underpinnings (Montag and Reuter, 2014).

5.3.3 Oxytocin

In Bartz and colleague’s summary of trials with intranasal oxytocin in 2011, they reported that situational or individual characteristics moderated the outcome in approximately 60% of the cases (Bartz et al., 2011b). The moderating effect of personality traits was shown in one study investigating performance on the Reading the Mind in the Eyes test after intranasal oxytocin administration, where only participants with high degree of alexithymia showed prosocial benefit of the neuropeptide (Luminet et al., 2011). Alexithymia is previously shown to correlate positively with Neuroticism and negatively with Extraversion and Openness (Luminet et al., 1999). Furthermore, Cardoso and colleagues found that intranasal oxytocin

was particularly beneficial on affective response to stress in women showing high levels of emotion-oriented coping (Cardoso et al., 2012b), and in a separate study they found effects of intranasal oxytocin to be associated with higher scores on Openness and Extraversion measured by the NEO-PI-R compared to participants that received placebo (Cardoso et al., 2012a). This corresponds with the association found between high plasma oxytocin levels and high scores on Extraversion (Andari et al., 2014).

The candidate gene approach to personality research has traditionally focused on serotonin and dopamine, but the results are however mainly inconclusive (Montag and Reuter, 2014). Oxytocin pathway genes have received augmented interest in the personality research field, due to their potential role in social behavior. Rs53576 has been associated with an anxiety-related temperamental trait measured by the Tridimensional Personality Questionnaire (Wang et al., 2014) as well as an interaction effect between neuroticism (measured by the Maudsley Personality Inventory) and dopamine transporter availability (measured by photon emission computed tomography) (Chang et al., 2014). Also, an interaction effect between a CD38 polymorphism and chronic interpersonal stress has been related to social anxiety (Tabak et al., 2016).

While a rich literature has shown associations between oxytocin pathway genes and social behavior and psychiatric symptoms, only a few studies have investigated trait effects of inter-individual signaling (Aspé Sánchez et al., 2016, Kumsta and Heinrichs, 2013). In addition, no investigation of the association between oxytocin pathway genes and personality traits measured by the NEO-PI-R has yet been performed (Balestri et al., 2014).

5.4 Positive and negative symptoms

5.4.1 Symptom course and treatment

Symptoms of psychotic disorders usually emerge in adolescence or early adulthood (Leboyer et al., 2005, Owen et al., 2016), and a diagnosis of schizophrenia or bipolar disorder is first set after a clear psychotic and/or affective episode, respectively. While patients with bipolar disorder seem to have preserved cognitive and functional levels in the pre-illness period (Lewandowski et al., 2011), many patients with schizophrenia experience a period with unspecific symptoms before the onset of the first psychotic episode. This period, also called the prodrome, often involves social withdrawal (Lieberman et al., 2001) and subsequently negative symptoms emerge. Social- and neurocognitive deficits are common as well in this period (Lee et al., 2015, Fusar-Poli et al., 2012). Eventually, attenuated psychotic symptoms such as suspiciousness, perceptual abnormalities and ideas of reference appear.

The only psychopharmacological treatments for schizophrenia offered today are dopamine receptor antagonists (antipsychotics), which mainly have an effect on positive symptoms. Antipsychotic drugs have little effect on negative symptoms and neurocognitive deficits, in fact they may even in some cases have a direct adverse effect (Harvey et al., 2004). Cognitive functioning deficits, poor pre-morbid functioning, earlier age of onset, longer duration of untreated psychosis and male gender are baseline factors that are found to be associated with poor treatment response (Case et al., 2010), but why these factors contribute to poor outcome is not well elucidated (Zhang and Malhotra, 2011). In lack of further knowledge about factors predicting treatment response, intolerable side effects and inefficacy are common reasons for discontinuation or switch of assigned treatment (Lieberman et al., 2005). For these reasons, a growing body of literature focus on genetic variants that may affect antipsychotic drug response (McClay et al., 2011), and development of novel drugs (Gopalakrishna et al., 2016, Feifel et al., 2016).

5.4.2 Oxytocin

The interest for associations between oxytocin and psychotic disorders started over thirty years ago (Bujanow, 1972). In animal models for schizophrenia, rats dosed with the N-methyl-D-aspartate (NMDA) antagonist phencyclidine, displayed altered social behavior, reduced oxytocin mRNA in the anterior hypothalamus, and increased oxytocin receptor (OXTR) binding in the amygdala (Lee et al., 2005). In the same study, direct injection of oxytocin in the amygdala restored social interaction behavior. Furthermore, oxytocin gene knockout mice have shown larger pre-pulse inhibitions (PPI) deficits after NMDA administration than mice with the oxytocin gene intact (Caldwell et al., 2009), and reduced dopaminergic hyperactivity in the striatum and nucleus accumbens after oxytocin administration has been shown as well (Qi et al., 2007).

A few studies have focused on plasma and CSF oxytocin levels in patients with psychotic disorders. As mentioned above, these methods are criticized for not reflecting oxytocin levels in the CNS. Goldman and colleagues divided 22 patients with schizophrenia into subgroups of polydipsic and non-polydipsic forms of the disorder, and found diminished plasma oxytocin levels in the polydipsic hyponatremic group (Goldman et al., 2008). Another study measured plasma levels after neutral and trust-related interpersonal interactions. In a sample of 50 patients with schizophrenia and 50 healthy controls, the plasma levels did not increase after trust-related interactions in patients in the same way as they did in controls (Keri et al., 2009). In addition, high plasma oxytocin levels have been associated with less severe positive symptoms in schizophrenia (Rubin et al., 2010). CSF oxytocin levels, however, have been found to be not different or even higher in patients with schizophrenia and bipolar disorders compared to healthy controls (Linkowski et al., 1984, Glover et al., 1994).

To date, 15 add-on clinical trials using intranasal oxytocin involving patients with schizophrenia have been performed, and treatment effects on positive and negative symptoms have been suggested (Feifel et al., 2016). However, a recent meta-analysis including six of the studies were not able to detect any effect on these symptoms (Heringa et al., 2015). No clinical trials with oxytocin administration have been performed in patients with bipolar disorder yet (Mercedes Perez-Rodriguez et al., 2015).

Association studies of oxytocin pathway polymorphisms in patients with schizophrenia have been conducted in seven samples (Bartholomeusz et al., 2015), with patient samples ranging from 74 to 406. Both tag SNPs and hypothesis based SNPs previously associated with social phenotypes have been selected, but solely OXT and OXTR have been investigated. While rs2268493 (OXTR) has been associated with mentalizing abilities and social perception (Davis et al., 2014), rs2740204 (OXT) with clozapine treatment response (Souza et al., 2010a), rs4813626 (OXT) and rs9840864 (OXTR) with the risk of developing schizophrenia (Watanabe et al., 2012, Teltsh et al., 2012), only the OXTR polymorphisms rs53576, rs2254298 and rs237902 have been significantly associated with symptoms in Caucasian schizophrenia samples. Montag and colleagues found the rs2254298A and rs53576G allele to be associated with higher general psychopathology scores and rs237902G to be associated with higher negative symptom scores measured by the Positive and Negative Symptom Scale (PANSS) (Montag et al., 2012, Montag et al., 2013). In addition, rs53576A and rs237885T (OXTR) have been associated with a diagnosis of schizophrenia in a Caucasian sample (Montag et al., 2013).

5.5 Affective face perception in psychotic disorders

5.5.1 Definition

Social cognition comprise the information processing and regulation of social stimuli, and is often divided into four separate processes; perception of social cues (including affective face perception), experience of sharing, mentalizing, and experiencing and regulating emotions (Green et al., 2015). Deficits in these domains have been shown prominent, chronic and with long-term effects on daily-life functioning in patients with psychotic disorders, and may mediate the functional loss from neurocognitive deficits (Fett et al., 2011). Social cognitive deficits are also shown to be the strongest predictor for transition to psychosis from a prodromal state (Corcoran et al., 2015). Despite having an important role in functional impairment, social cognitive deficits are not part of the diagnostic criteria of psychotic disorders.

Affective face perception is widely investigated, and both impairments in the ability to identify emotional face expressions and deficits in neural activity during face perception tasks have been found consistently in psychotic disorders (Green et al., 2015). Interestingly, patients with schizophrenia seem to display more severe deficits in affective face perception compared to patients with schizoaffective disorders, that in turn show more severe deficits than patients with bipolar disorders, even though impairments are present in all these patient groups (Ruocco et al., 2014, Bora and Pantelis, 2016).

5.5.2 Amygdala activation

Several studies have shown that social cognition rely on distinct neural circuits from those involved in other cognitive processes (Meyer-Lindenberg et al., 2011). A large meta-analysis including 100 fMRI reports on affective face perception in healthy subjects showed that the

parahippocampal gyrus, inferior frontal gyrus, medial prefrontal gyrus and fusiform gyrus as well as the amygdala are important brain regions involved in this social cognitive domain (Sabatinelli et al., 2011). In social salience and threat-related signal processing, amygdala in particular seems to have a critical role (Phelps and LeDoux, 2005).

A recent large-scale structural MRI study established that the amygdala volume was significantly smaller in patients with schizophrenia compared to healthy controls (van Erp et al., 2016), while reduced amygdala volume is possibly most prominent during youth and normalized in adulthood due to medical treatment in bipolar disorder (Wijeratne et al., 2013, Foland et al., 2008, Phillips and Swartz, 2014). Additionally, amygdala dysfunction has been associated with altered affective face perception in schizophrenia in several studies (Takahashi et al., 2004, Brunet-Gouet and Decety, 2006), and threat-related facial affect perception is particularly altered in this patient group (Kohler et al., 2010, Edwards et al., 2002, Kohler et al., 2003). Three meta-analyses suggest altered amygdala activation in patients with schizophrenia in response to negative emotional stimuli, either in itself (Li et al., 2010), or in contrast to neutral stimuli (i.e. neutral faces or other neutral stimuli) (Taylor et al., 2012, Anticevic et al., 2012), and the authors propose that patients with schizophrenia show an over-activation in response to neutral emotional stimuli more than an under-activation in response to negative emotional stimuli. Interestingly, a meta-analysis of 1040 patients with bipolar disorder and 1074 controls, including six fMRI studies, found an over-activation in the amygdala during both negative and positive faces, regardless of affective state (Chen et al., 2011). The explanation for this difference in amygdala response to emotional stimuli between patients with schizophrenia and bipolar disorder remains unclear, but could ultimately reflect the differences in social cognitive deficits.

Despite of an increasing knowledge of the neural circuits involved in affective face perception, there are no pharmacological treatments available to date targeting these deficits, and there are no compelling evidence that antipsychotics ameliorate these deficits either (Kucharska-Pietura and Mortimer, 2013).

5.5.3 Oxytocin

Due to the rich literature stressing the central role of oxytocin in social cognition, deficits in these domains have been targeted in trials with intranasal oxytocin in patients with schizophrenia. While some improvements in detecting empathic accuracy (Davis et al., 2013), fear recognition (Gibson et al., 2014) and Theory of Mind (Pedersen et al., 2011) have been seen, the results have been mixed (Cacciotti-Saija et al., 2015) and found too heterogeneous to be included in a recent meta-analysis (Heringa et al., 2015). Shin and colleagues measured in 2015 for the first time brain activation during oxytocin administration in patients with psychotic disorders, and interestingly found decreased amygdala activation when the participants underwent an affective face perception paradigm (Shin et al., 2015). Recently, Rubin and colleagues investigated DNA methylation of OXTR in 167 patients with both schizophrenia and bipolar disorder, and found higher levels to be associated with both smaller volumes in temporal-limbic and prefrontal regions as well as poorer affective face perception (only in females), and higher levels were seen in patients with schizophrenia compared to patients with bipolar disorder (Rubin et al., 2016). To date, no other association studies of oxytocin pathway genes in patients with bipolar disorder have been performed.

Combined, there is support from animal and genetic studies that the endogenous oxytocin system is altered in patients with schizophrenia. However, the evidence is still mixed, and the research field mainly struggles with small sample sizes and most studies investigating

oxytocin pathway genes include mixed ethnic populations. Further, the associations with clinical symptomatology have been inconsistent (Davis et al., 2014, Watanabe et al., 2012), and no study investigating the association between oxytocin pathway genes and brain activation has yet been performed in patients with psychotic disorders.

5.6 Aims of the thesis

The overall aim of this PhD study was to gain more knowledge of the relationship between the endogenous oxytocin system and features that are important for social dysfunction in patients with psychotic disorders; personality traits related to trust and social behavior, positive symptoms related to trust, negative symptoms related to social withdrawal and amygdala activation related to affective face perception.

We sought to identify minor genetic alterations in oxytocin pathway genes regulating oxytocin, which could lead to deficits in core social features, with possible implications for treatment and the understanding of pathophysiological mechanisms of these features. The studies were conducted in both a patient and a healthy control sample with the following subaims:

- i) To identify oxytocin pathway polymorphisms contributing to the relationship between personality traits in healthy subjects correlated with trust and social behavior that have previously been associated with inter-individual effects of oxytocin, and oxytocin pathway genes.
- ii) To identify associations between oxytocin pathway genes and psychotic disorders per se in a case-control design as well as between oxytocin pathway genes and specific psychopathological features associated with trust and social behavior, in

particular suspiciousness/persecutory delusions, hostility, emotional withdrawal and passive/apathetic social withdrawal.

- iii) To identify associations between three genetic polymorphisms of the OXTR gene (rs53576, rs2254298 and rs237902) and amygdala activation during affective face perception of negative faces, and determine possible disorder specific associations in patients with psychotic disorders.

6. MATERIAL AND METHODS

6.1 Participants

Participants were included in the Norwegian multi-center Thematically Organized Psychosis Research (TOP) study recruiting patients from in- and out-patient clinics in the greater Oslo area in South-Eastern Norway. In this part of the study, a total of 734 with a psychotic disorder and 417 healthy individuals were included. All participants were Caucasian to avoid confounding by population stratification, and met an adequate level of spoken Scandinavian language. None had a history of head injury, neurological disorders, autoimmune or infectious disorders or malignancies, or an IQ below 70. All participants gave written informed consent and the study is approved by the Norwegian Scientific-Ethical Committees and the Norwegian Data Protection Agency.

Table 1 Sample overview. All participants in Study I and III are represented in Study II.

	Study I	Study II	Study III
Patient sample, N	0	734	204
Healthy controls, N	196	417	142
Overlap, N (%):			
Study I	196 (100)	196 (17)	136 (39)
Study II	196 (100)	1151 (100)	346 (100)
Study III	136 (69)	346 (30)	346 (100)

6.1.1 Patient sample

Participants in the patient sample were referred from a clinician and received treatment during inclusion in the project. They all underwent clinical and physical examination by a physician, extensive neuropsychological testing by a psychologist, collection of blood samples for somatic screening and DNA analyses. The patients included were interviewed, investigating sociodemographic history, medical history, substance abuse, psychiatric symptoms, medication and potentially side effects. They all underwent diagnostic interviews based on the

Structured Clinical Interview for DSM-IV Axis 1 Disorders (SCID-1) and were diagnosed with a psychotic disorder. All underwent symptoms assessments including the PANSS.

Study II: Of the total sample of 734, all had a schizophrenia polygenic risk score (PGRS) and genotyped or imputation data for OXTR SNPs of interest. 725 individuals (364 women and 361 men) had available symptom scores of p6 measured by the PANSS, while 724 patients had available symptom scores of p7, n2 and n4.

The sample of 734 patients covered the following diagnoses: schizophrenia ($n=265$, out of which 263 had symptom scores), schizophreniform disorder ($n=23$), schizoaffective disorder ($n=62$), bipolar disorder I ($n=180$), bipolar disorder II ($n=72$), bipolar disorder NOS ($n=18$), depressive psychosis ($n=21$) and other psychosis ($n=93$).

Study III: The group of 104 patients (41 women and 63 men) with a schizophrenia spectrum disorder (SCZ) included the following diagnoses: schizophrenia ($n = 62$), schizophreniform disorder ($n = 6$), schizoaffective disorder ($n = 12$), and psychotic disorder NOS ($n = 24$). The affective spectrum (AD) group ($n= 100$, 57 women and 43 men) consisted of patients with bipolar disorder I ($n = 50$), bipolar disorder II ($n = 38$), bipolar disorder NOS ($n = 6$), and depressive psychosis ($n = 6$). In addition to a standard fMRI protocol, these participants underwent an emotional faces matching paradigm, and task-induced blood-oxygen-level dependent (BOLD) amygdala response to emotional faces was measured. Complete behavioral data (response time and accuracy rate) were available for 89 participants with SCZ and 94 participants with AD. For the remaining individuals an accuracy rate (i.e. combined rate for negative faces and shapes) was calculated.

6.1.2 Healthy controls

Healthy participants were selected randomly from statistical records of persons from the same catchments areas in South-Eastern Norway as the hospitals giving services to the TOP study's case sample. All subjects were screened for illness using the Primary Care Evaluation of Mental Disorders and interviewed about severe psychiatric disorders, drug abuse, and somatic disease. Healthy participants were excluded in case of any history of severe psychiatric disorders (major depression, bipolar disorders, and schizophrenia) in the participants or in any of their first-degree relatives, substance or alcohol abuse or dependency. Blood samples for genotyping were collected for all healthy participants.

Study I: A total of 196 healthy subjects were included (86 women and 110 men). Personality traits were assessed by the Neuroticism-Extraversion-Openness Five-Factor Inventory (NEO-FFI).

Study II: The control sample consisted of 417 healthy individuals, 209 women and 208 men. All had genotyped or imputation data for SNPs of interest as well as PGRSs.

Study III: 142 healthy individuals were included, 58 women and 84 men. As within the patient sample in this study, these participants underwent an emotional faces matching paradigm during fMRI, and task-induced BOLD amygdala response to emotional faces was measured. Complete behavioral data (response time and accuracy rate) were available for 140 healthy individuals. For the two remaining individuals an accuracy rate was calculated.

6.2 Clinical Assessments

6.2.1 Diagnostic assessment

The SCID-1 was used for diagnostic purposes in the patient sample (First, 2002). Trained psychologists and physicians performed all the diagnostic interviews, based on modules A-E. The ratings are based on both the patients' thought content as conveyed during the interview, information from informants, and clinical observation during the interview. Interviewers underwent regularly supervision and consensus meetings with experienced clinicians and investigators. Directed by the South-Eastern Norway Regional Health Authority, all interviewers completed a mandatory SCID-1 Training and Quality Assurance Program for researchers (Ventura et al., 1998). The mean overall inter-rater variability for study patients was 82%, $\kappa = 0.77$ (95% CI 0.60-0.94).

6.2.2 Assessment of positive and negative symptoms

We used the PANSS; that is a 30-item, 7-point rating instrument (Kay et al., 1987) for symptoms assessments in the patient sample. This interview-based scale measures psychopathology that is common in psychotic disorders: positive, negative, depressive, disorganized and excitative symptoms for the last week (Langeveld et al., 2013). The ratings are based both on the patients' thought content as conveyed during the interview, information from informants and clinical observation during the interview. For the current study we selected four items that are closely related to trust and social function for investigation: suspiciousness/persecution (p6); e.g. unrealistic or exaggerated ideas of persecution, as reflected in guardedness, a distrustful attitude, suspicious hypervigilance or frank delusions that others mean harm, hostility (p7); e.g. verbal and nonverbal expressions of anger and resentment, including sarcasm, passive-aggressive behavior, verbal abuse and assaultiveness, emotional withdrawal (n2); e.g. lack of interest in, involvement with, and affective

commitment to life's events and passive/apathetic social withdrawal (n4); e.g. diminished interest and initiative in social interactions due to passivity, apathy, anergy or avolition that leads to reduced interpersonal involvements and neglect of activities of daily living.

6.2.3 Assessment of personality traits

In the healthy control sample in study I, we used the Neuroticism-Extraversion-Openness-Five Factor Inventory (NEO-FFI); that is a 60 item short-version of the NEO-PI-R. Both versions measure five personality traits: Neuroticism (anxiousness, impulsiveness, depression, hostility, vulnerability to stress, self-consciousness), Extraversion (excitement-seeking, gregariousness, assertiveness, warmth, activity, positive emotion), Openness to experience (fantasy, aesthetics, feelings, actions, openness to ideas and values), Agreeableness (trust, modesty, altruism, compliance, straightforwardness, tender-mindedness) and Conscientiousness (self-discipline, order, achievement striving, competence, dutifulness, deliberation) (McCrae et al., 2005). The NEO-FFI is a self-report inventory with 12 items for each of the five personality domains. The items comprise descriptions of behavior with answers on a five-point scale, ranging from “strongly agree” to “strongly disagree”. For the current study we pre-selected three personality domains with relation to trust and social behavior for investigation: Agreeableness, Neuroticism and Extraversion. We thus obtained a total of three domain scores based on the mean of the 12 items representing each personality domain.

6.3 Selection of genetic variants

We used three different approaches for the selection of genetic variants to answer our research questions:

- i. The first was to investigate SNPs covering all the four oxytocin pathway genes (i.e. OXT, AVP, OXTR, CD38) from the Affymetrix chip. This was performed in Study I since no previous studies regarding oxytocin pathway genes and personality traits measured by the NEO-PI-R had been performed previously, and we wanted to identify possible contributing oxytocin pathway polymorphisms.
- ii. The second was to investigate an overall effect of oxytocin pathway genes by calculating a PGRS score for schizophrenia for OXT, AVP, CD38 and OXTR polymorphisms using the results from the PGC schizophrenia study (Ripke et al., 2011) as a discovery sample for generating PGRS. This study was performed before the release of PGC2 and the results from the PGC1 were used, involving 9146 schizophrenia cases and 12111 controls. This method was applied in study II and represented a cumulative risk contribution of oxytocin pathway genes to schizophrenia.
- iii. The third was to investigate SNPs with previous association to either the diagnosis of schizophrenia (OXTR SNP rs53576), to social behavior, or to related symptoms in schizophrenia based on previous studies i.e. the four OXTR SNPs (rs1042778, rs53576, rs2254298 and rs237902). Rs237902 was genotyped on the Affymetrix chip, while rs53576, rs2254298 and rs1042778 were not available on this chip and hence imputed. This was performed in all three studies, however study II and III did not include rs1042778 since this polymorphism had not been related to symptoms in psychotic disorders previously.

6.4 Genotyping and Imputation

The TOP sample was genotyped using the Affymetrix Genome-Wide Human SNP Array 6.0 (Affymetrix Inc., Santa Clara, CA, USA). Samples were excluded through quality control if they had low-yield (call rate below 95%), if they were duplicates of other samples included in the study, if they had a sex determined by X chromosome marker homozygosity different from their reported sex or if they were calculated to have other ancestry than European. Markers with severe deviation from Hardy–Weinberg equilibrium, low-yield (<95%) or with minor allele frequencies below 0.01 were excluded. Following the above mentioned quality control, the candidate SNPs were imputed with MaCH (Sph.umich.edu., 2016a) using the European samples in the Phase I release of the 1000 Genomes Project (SNPs not present in the 1000 Genomes reference, and SNPs with ambiguous strand alignments (A/T and G/C SNPs), were removed from the sample data sets). Imputation was a three stage process, involving (I) ChunkChromosome where the data set was broken into 2500 SNP pieces with 500 SNP overlap (Genome.sph.umich.edu., 2016a), (II) MaCH where each piece was phased (40 rounds and 400 states) (Sph.umich.edu., 2016b), and (III) Minimac where each phased piece was imputed to the 1000 Genomes European reference panel (20 rounds and 400 states) (Genome.sph.umich.edu., 2016b). In the third stage, all imputed SNPs were provided with an estimated r^2 score as quality metric. Exclusions were made of SNPs with an r^2 score <0.5 leaving 9,584,802 SNPs. Imputed OXT, OXTR, AVP and CD38 SNPs based on University of California Santa Cruz (UCSC) coordinates ± 20 kb (hg19) were extracted for inclusion in the current analyses and exclusions were made of SNPs with a minor allele frequency <0.01.

6.5 Polygenic risk score

A PGRS for the schizophrenia phenotype was computed based on imputed SNPs following the method developed by Purcell (Purcell et al., 2009b). With the aid of PLINK version 1.07 (Purcell, 2009a, Purcell et al., 2007), we performed a meta-analysis including all PGC sub studies (2011) except ours (TOP3) (n= 9146 schizophrenia cases and 12111 controls), to obtain risk allele effect sizes (the natural logarithm of odds ratios, $\ln(\text{OR})$) for SNPs in the oxytocin pathway genes: OXT [OMIM 167050], OXTR [OMIM 167055], CD38 [OMIM 107270] and AVP [OMIM 192340]. The SNPs within each gene were subsequently pruned by grouping them in LD ($r^2 < 0.25$) sets and selecting from each set the representative with the highest absolute effect size. This procedure left us with 32 SNPs. A cumulative risk score was then computed for each individual in our sample summing up the effect sizes of the selected SNPs multiplied by the number of risk alleles carried by that individual. Such a cumulative risk score is assumed to represent a sort of oxytocin pathway cumulative risk for schizophrenia.

6.6 fMRI analyses

6.6.1 Experimental paradigm

A widely used and validated paradigm was employed to elicit amygdala activation in study III (Carre et al., 2012, Hariri et al., 2002, Ousdal et al., 2012, Tesli et al., 2015). In this task, participants selected which of two stimuli (displayed at the bottom of the screen) matched a target stimulus (displayed at the top). In the faces-matching task, the images displayed were either human faces expressing anger or fear ("negative faces"), or happiness ("positive faces"). In the sensorimotor control task, geometrical shapes were displayed. The negative and positive faces were presented in two separate runs, each with 4 blocks of faces interleaved

by 5 blocks of geometrical shapes. The order of the runs was counterbalanced between participants. Within blocks of negative faces, fearful and angry faces were intermixed. Each block of faces consisted of 6 emotion-specific face trios derived from a standard set of facial affect pictures (Tottenham et al., 2009). Each trial (face/shape stimulus) was presented for 5.4 seconds with no inter-stimulus interval, for a total block length of 32.6 seconds. Each run lasted 294 seconds. The E-prime software (version 1.0 Psychology Software Tools, Inc, Pittsburgh, PA, USA) controlled the presentations of the stimuli using VisualSystem (NordicNeuroLab, Bergen, Norway). Measures of reaction times and task accuracy were recorded through MRI-compatible ResponseGrips (NordicNeuroLab, Bergen, Norway).

6.6.2 BOLD fMRI data acquisition

MRI was performed on a 1.5 T Siemens Magnetom Sonata scanner (Siemens Medical Solutions, Erlangen, Germany) supplied with a standard head coil. We collected 152 functional volumes per run using a T2*-weighted echo-planar imaging sequence. Each volume consisted of 24 axial slices with a pixel size of 3 mm in the axial plane and a slice thickness of 4 mm with 1 mm gap between slices (TR=2040 ms, TE=50ms, FA=90°, matrix 64 x 64, FOV 224 mm). The first seven volumes and the last one were discarded, leaving 144 volumes for analyses. Prior to fMRI, a sagittal T1-weighted 3D Magnetization Prepared Rapid Gradient Echo scan was collected (160 slices, TR= 2730 ms, TE=3.93 ms, FA =7°, matrix 192 x 256, FOV 240 mm, voxel size 1.33 x 0.94 x 1 mm³), here used for co-registration purposes.

6.6.3 fMRI Data Analysis

The functional MRI data went through an initial quality check procedure, to detect images with poor quality due to excessive head motion, slice dropout, radiofrequency artifacts or other noise. It was subsequently processed and analyzed using FEAT (FMRI Expert Analysis

Tool) version 6.0, part of FSL (Fmrib.ox.ac.uk., 2016, Smith et al., 2004). Conventional preprocessing included motion correction (Jenkinson et al., 2002), non-brain removal (Smith, 2002), spatial smoothing with a 6 mm full-width half-maximum kernel (Smith and Brady, 1997) and a high pass filter with a 90 s window. A first-level general linear model analysis was performed for each run, where the onset and duration of the on-blocks (positive faces and negative faces, respectively) was modeled with that of the off-blocks (geometrical shapes) as implicit baseline. The design matrix was filtered and convolved with a hemodynamic response function before the model fit. This analysis resulted in individual contrast parameter estimates (COPEs) reflecting the faces vs. shapes contrast within runs. In study III, we focused on participants' amygdala responses to negative stimuli (the negative faces > shapes contrast), due to the implication of altered threat-related facial affect perception in psychotic disorders (Kohler et al., 2003, Hall et al., 2008) and since this contrast has been reported to reliably engage the amygdala (Tesli et al., 2015, Costafreda et al., 2008, Zald, 2003). Amygdala regions of interest (ROIs) were defined in accordance with the probabilistic Harvard-Oxford subcortical atlas provided with FSL, and were thresholded at 25% probability. Mean BOLD signal changes (parameter estimates) across all ROI voxels were extracted from FSL and used in statistical analyses.

6.7 Statistics

We performed genetic association analyses between oxytocin pathway polymorphisms and phenotypes using linear regression analyses in all three studies. The phenotypes were set as dependent variables, and genotypes in allelic dosage models were set at independent variables (i.e. entering the number of minor alleles with the possibility of 0, 1 or 2). For imputed SNPs, the expected allele count was entered as the independent

variable. Possible confounding variables were included as covariates. In study II, an allelic logistic regression model was used to test for case/control association. For these purposes, the PLINK software version 1.07 (Purcell, 2009a, Purcell et al., 2007) was used.

In study II, a multiple regression model was performed with all three SNPs in question included in the model to test for potential specificity of each SNP. In study III, interaction effects between SNP and diagnosis as well as between SNP and sex were tested for. Only one interaction term was included in the model at the time. For these purposes, the software R Version 3.2.2 (Cran.r-project.org., 2015) was used.

The software SPSS Version 22 (IBM, Armonk, N.Y) was used for inspection of distribution of the variables, PGRS analyses, group comparisons and correlation analyses. In study II, potential differences in mean schizophrenia oxytocin pathway PGRS between cases and healthy controls were first investigated with t-tests for independent samples. Additionally, we performed multiple linear regression analyses for each PANSS scores with the PGRS as an independent variable in the total patient sample, correcting for potential confounders. In study III, potential differences in amygdala activation, response time and accuracy rate between subgroups (SCZ, AD, and healthy controls) were examined using Kruskal-Wallis tests. Sex effects for amygdala activation were tested for in the total group with Mann-Whitney tests. The correlation between right and left amygdala activation was examined using bivariate correlation analyses (Spearman's rho).

Genetic statistical power was estimated with the Genetic Power Calculator (Purcell et al., 2003, Pngu.mgh.harvard.edu., 2016) assuming a disease prevalence of 0.01 for schizophrenia and 0.02 for psychotic disorders, an additive allelic model and a type I error rate of 0.05. To

correct for multiple testing in the genetic association analyses, Bonferroni corrections for the number of independent tests were performed in all studies.

7. SUMMARY OF RESULTS

Study I

The purpose of this study was to identify oxytocin pathway polymorphisms contributing to the relationship between three personality traits related to trust and social behavior; Agreeableness, Neuroticism and Extraversion, and oxytocin pathway genes.

69 SNPs in the oxytocin pathway genes OXT, AVP, OXTR and CD38 were genotyped from the Affymetrix chip and three pre-selected OXTR SNPs (rs53576, rs2254298 and rs1042778) found associated with social behavior in previous literature were imputed in 196 healthy subjects.

We found seven nominally significant associations for personality traits; Agreeableness (rs857240 (AVP, $p=0.0075$), rs2270463 (OXTR, $p=0.047$)), Neuroticism (rs3756242 (CD38, $p=0.024$), rs13104011 (CD38, $p=0.024$), rs6816486 (CD38, $p=0.024$), rs7655635 (CD38, $p=0.034$) and Extraversion (rs237878 (OXTR, $p=0.019$)). None of these associations remained significant after Bonferroni correction ($P\text{-threshold} = 2.31 \times 10^{-4}$), and none of these seven polymorphisms had been found associated with social phenotypes in the literature previously.

Our results do not support the notion of significant contributions of oxytocin pathway genes to personality traits and we find it less likely that they are central biological modulators of the individual levels of Agreeableness, Neuroticism and Extraversion.

Study II

We performed association analyses between four oxytocin pathway genes (OXT, OXTR, AVP, CD38) and four areas of social behavior-related psychopathology as measured by the

PANSS; suspiciousness/persecutory delusions, hostility, emotional withdrawal and passive/apathetic social withdrawal. Additionally, we tested for genetic case/control differences. For these purposes, we investigated both a PGRS and single OXTR candidate SNPs previously reported in the literature (rs53576, rs237902, rs2254298). A total of 734 subjects with DSM-IV psychotic spectrum disorders and 417 healthy controls were included.

There was a significant association between higher symptom levels of emotional withdrawal and the OXTR risk allele A in rs53576 ($p = 0.007$, $p = 0.084$ after Bonferroni correction). This finding survived Bonferroni corrections when all three SNPs were added in the same multiple regression model ($p = 0.002$, $p = 0.009$ after Bonferroni correction). No significant associations between rs53576 and suspiciousness/persecutory delusions, hostility, or passive/apathetic social withdrawal were found and rs53576 failed to differ between cases and controls. No significant associations were found between rs237902 or rs2254298 and the PANSS measures under investigation, nor between PGRS and symptom levels or case/control status.

Our findings indicate that oxytocinergic signaling is involved in the pathophysiology of the level of emotional withdrawal in patients with psychotic disorders, a feature related to social dysfunction in this patient group. Oxytocin pathway genes do however appear to not be important in the pathogenesis of psychotic disorders per se.

Study III

346 participants (104 cases with SCZ, 100 cases with AD, and 142 healthy controls) underwent genotyping and fMRI during an emotional faces matching paradigm. Genetic association analyses were performed to test the possible effects of the OXTR SNPs rs53576, rs237902 and rs2254298 on task-induced BOLD amygdala response to fearful/angry faces.

Potential differences in amygdala activation, response time and accuracy rate between subgroups (SCZ, AD, and healthy controls) as well as interaction effects between diagnoses and amygdala activation were examined as well.

In participants with SCZ, the rs237902 G allele was associated with low amygdala activation (left hemisphere, $b = -4.99$, Bonferroni corrected $p = 0.04$) and interaction analyses showed that this association was disorder specific (left hemisphere; Bonferroni corrected $p = 0.003$, right hemisphere; Bonferroni corrected $p = 0.03$). There were no associations between oxytocin receptor polymorphisms and amygdala activation in the total sample, among AD patients or healthy controls. We did not detect differences in amygdala activation (left hemisphere: $\chi^2 = 0.73$, $p = 0.69$, right hemisphere: $\chi^2 = 0.19$, $p = 0.91$) or accuracy rate ($\chi^2 = 0.29$, $p = 0.87$) between diagnostic subgroups, but in response time ($\chi^2 = 13.6$, $p = 0.001$).

Our findings indicate that minor genetic alterations in the OXTR could serve as associated factors in biological underpinnings of affective face perception, a social cognitive trait important for social functioning in patients with psychotic disorders. The disorder-specific association found suggests that oxytocinergic signaling influences affective face perception in a different way in patients with SCZ than in patients with AD or healthy controls.

8. DISCUSSION

8.1 Main results

The three studies in the present thesis sought to determine if genetic alterations in oxytocin pathway genes could be associated with features important for social dysfunction in patients with psychotic disorders. The main findings are significant associations between an intronic single oxytocin receptor polymorphism and emotional withdrawal in patients with psychotic disorders and between an exonic single oxytocin receptor polymorphism and amygdala activation during affective face perception of negative faces in patients with schizophrenia spectrum disorders. Oxytocin pathway genes however, failed to explain any association to a diagnosis of psychotic disorder per se, or to variation in personality traits in healthy subjects. In this discussion, these findings will firstly be presented in light of previous literature. Secondly, these findings will be discussed in light of a new insight into the social-salience hypothesis with potential biological and clinical implications. At last, methodological strength and limitations, as well as future directions will be discussed.

8.1.1 The relationship between personality traits and oxytocin pathway genes

In Study I, we found seven nominally significant associations between variants in oxytocin pathway genes (OXT, OXTR, CD38 and AVP) and personality traits related to trust and social behavior (Agreeableness, Neuroticism and Extraversion), but none of these associations remained significant after correction for multiple testing.

Ten of the 72 investigated SNPs included for analysis in the present study have been found in previous literature to be associated with trust and/or social behavior (eight located at OXTR: rs11131149, rs237900, rs237902, rs2268494, rs7632287, rs53576, rs2254298, rs1042778 and

two located at CD38: rs6449197, rs3796863). Rs1042778 is an exonic variant in the 3 prime untranslated region of the OXTR, with uncertain predictive value. However, Feldman and colleagues found this polymorphism, in addition to rs2254298 located at the OXTR, to be associated with oxytocin plasma levels in a study of 352 healthy individuals, making further investigation of this SNP interesting (Feldman et al., 2012). In the same study, the rs1042778 TT genotype was associated with less parental touch. In other previous studies, this polymorphism has been associated with autism, antisocial behavior and early signs of psychopathy (Dadds et al., 2014, Malik et al., 2012). Rs53576 has been associated with a range of social features (Li et al., 2015), and recently an interaction effect between this polymorphism and insecure childhood attachment was shown to modulate the level of alexithymia in adulthood, as well as grey matter volume in brain areas related to social salience processing (Schneider-Hassloff et al., 2016). Based on this literature, we imputed rs53576, rs2254298 and rs1042778 that are not available on the Affymetrix chip, and included these polymorphisms in the analyses. We did however not find any association between these three polymorphisms and personality traits, hence we were not able to replicate earlier findings.

Concerning other OXTR SNPs, we examined rs237900 that has been associated with Harm Avoidance, measured by the Temperament and Character Inventory (Stankova et al., 2012), a personality trait positively correlated with Neuroticism and negatively correlated with Extraversion in the five-factor model (De Fruyt et al., 2000). Further, rs11131149 has been associated with depressive temperaments (Kawamura et al., 2010), rs2268494 with the ability to cooperate in a Prisoner's Dilemma Game (Tabak et al., 2014), and rs7632287 with pair-bonding behavior (Walum et al., 2012). As with the imputed polymorphisms in our study,

none of these genotyped polymorphisms showed associations with personality traits in our sample.

Variants in OXT, AVP and CD38 are less extensively examined in general regarding social behavior, and to our knowledge, the relationship between these three genes and personality traits has not been examined before. Feldman and colleagues found rs3796863 located at CD38 to be associated with oxytocin plasma levels in the same study detecting association between oxytocin plasma levels and rs2254298 and rs1042778, highlighting this particular polymorphism for further investigation (Feldman et al., 2012). No associations with personality traits were found with this particular polymorphism in the present study, indicating that the potential expression of oxytocin associated with this SNP is less likely to influence the level of Agreeableness, Neuroticism and Extraversion.

In summary, we were not able to detect associations between variants in oxytocin related genes and personality traits expected to be strongly correlated with trust and social behavior, and our results do not support the notion of significant contributions of oxytocin pathway genes to personality traits. We were not able to replicate earlier findings, and all the nominally significant associations found were for SNPs with no previous reports in the literature.

8.1.2 The effect of oxytocin pathway genes on positive and negative symptoms and diagnosis in psychotic disorders

We found in study II that rs53576 A-allele carriers had higher scores on emotional withdrawal measured by the PANSS compared to G-allele carriers, while we were not able to detect associations between symptom scores and rs237902 or rs2254298 in patients with psychotic disorders. Neither did we detect any predictive value of rs53576 to a diagnosis of any

psychotic disorder or associations between oxytocin pathway gene PGRS and symptoms/diagnosis of any psychotic disorder.

Rs53576 and rs2254298, two intronic polymorphisms, have been extensively studied across social features, psychiatric symptoms and ethnic groups, and have in separate studies been associated with having an autism spectrum disorder, levels of empathy, parental sensitivity and amygdala volumes (Wu et al., 2005, Bakermans-Kranenburg and van Ijzendoorn, 2008, Rodrigues et al., 2009, Meyer-Lindenberg et al., 2011, Inoue et al., 2010, Wang et al., 2014, Kumsta and Heinrichs, 2013). However, results are inconsistent to the degree that different meta-analyses seem to disagree about their importance (LoParo and Waldman, 2015, Li et al., 2015, Bakermans-Kranenburg and van Ijzendoorn, 2014). In psychotic disorders these variants have been associated with both general psychopathology (rs53576 and rs2254298), and negative symptoms as measured by the PANSS (rs237902) and empathic concern (rs2254298) (Montag et al., 2012, Montag et al., 2013).

Emotional withdrawal is characterized by the lack of interest in, involvement with, and affective commitment to life's events, and is seen in a wide range of psychiatric disorders. Thus, the association we found with rs53576 in the present thesis supports previous studies of a negative association with social behavior in the A-allele for areas covering maternal sensitivity (Bakermans-Kranenburg and van Ijzendoorn, 2008), empathy (Rodrigues et al., 2009), sociability (Tost et al., 2010), trust behavior in healthy subjects (Krueger et al., 2013) and seeking of emotional support in times of distress (Kim et al., 2010). Contrasting studies showing that patients with schizophrenia carrying the G-allele had higher general psychopathology scores measured by the PANSS (Montag et al., 2013) could be in line with a “flip-flop phenomenon” (Lin et al., 2007). We were not able to detect associations between rs2254298 and positive or negative symptoms in patients with psychotic disorders, a finding

that corresponds to a previous result in a smaller sample (Montag et al., 2012). Neither did we find any associations between rs237902 and symptom measures, in contrast to Montag and colleagues who found a higher risk of negative symptoms in rs237902G carriers with schizophrenia (Montag et al., 2013). Montag and colleagues however used a combined score of all PANSS negative sub-items and our conflicting finding could be explained by rs237902 being associated more with other negative sub-items than emotional withdrawal or passive/apathetic social withdrawal.

Only a few case-control studies have investigated the relationship between the diagnosis of schizophrenia and oxytocin related genes. Previous studies have used different strategies for SNP selection in the oxytocin pathway genes and seem to be rather inconclusive (Souza et al., 2010b, Watanabe et al., 2012, Teltsh et al., 2012). Montag and colleagues found an association between the diagnosis of schizophrenia and rs53576 in a case-control study with 406 patients and 406 healthy controls, with the A-allele occurring more frequent in the patient group compared to the controls (Montag et al., 2013). We were not able to replicate this finding indicating that this association is not robust. With a combined psychosis sample twice the size as in Montag and colleagues' study, we find it unlikely that this particular polymorphism contribute significantly to the risk of developing schizophrenia or any other psychotic disorder.

In addition, we used a polygenic risk approach, a measure derived from associations with risk in a large case-control sample, to capture more of the small effects to risk for a psychotic disorder across the oxytocin pathway genes (OXT, OXTR, CD38 and AVP). Based on recent reports about overlapping symptomatology and susceptibility genes across different severe mental disorders (Craddock and Sklar, 2013, Kidd, 2013, Levy and Manove, 2012) we here sought to include a wide range of diagnoses in the psychosis spectrum. To our knowledge, no

previous studies have investigated their relationship to a diagnosis of any psychotic disorder. We were not able to find any significant associations between PGRS over the four oxytocin-related genes and the diagnosis of any psychotic disorder in our sample. We also did not find any associations between PGRS and relevant symptom measures. The calculated polygenic risk scores that include a cumulative risk profile across the whole genome, i.e. thousands of common genetic variants each with small effect, are shown significantly higher in patients with schizophrenia and bipolar disorder than in cases (Purcell et al., 2009b, Agerbo et al., 2015, Tesli et al., 2014). This indicates both that psychotic disorders are highly polygenic and that there is genetic overlap across psychotic disorders, but also that the measure of PGRSs is useful in detecting case-control status. The calculated PGRS based on oxytocin pathway genes is then likely to not involve a sufficient amount of risk variants to predict case-control status, reflecting that common genetic variants in these genes do not contribute substantially to the genetic risk of a psychotic disorder. This finding is in line with results in the large PGC study (Schizophrenia Working Group of the Psychiatric Genomics, 2014). The correlation between PGRS based on common genetic variants across the whole genome and quantitative phenotypes has however been more difficult to detect. Derks and colleagues measured positive, negative, disorganized, depressive and manic symptoms in 715 schizophrenia cases and 643 controls, but were not able to detect any association with the calculated PGRS (Derks et al., 2012). Corresponding results were found when PGRSs based on risk loci for bipolar disorder were calculated, and not found associated with positive, negative or depressive symptoms, but, however with the level of mania in patients with schizophrenia (Ruderfer et al., 2014). Derck and colleagues explain their non-significant findings with low statistical power, that other alleles, more rare variants or epigenetic variation may be more involved in dimensional symptom scales than risk alleles (Derks et al., 2012). This could certainly explain our non-significant association between oxytocin PGRS and symptom measures as well, and

most likely we also included too few genetic variants in the calculated PGRSs to detect significant associations. To our knowledge, this is the first study using an oxytocin candidate gene polygenic approach to investigate phenotypes in psychotic disorders, and replication in other samples could be useful.

8.1.3 The effect of oxytocin pathway genes on affective face perception in psychotic disorders

In Study III we showed attenuated amygdala activation in response to emotionally negative faces in rs237902G allele carriers among patients with schizophrenia spectrum disorders, but not in patients with affective spectrum disorders or healthy subjects. In this study, we were not able to detect associations between amygdala activation during affective face perception of negative faces and rs53576 or rs2254298 in any subsample.

While the present study is the first fMRI study investigating oxytocin pathway polymorphisms in patients with psychotic disorders, several investigations have been performed in healthy subjects. The OXTR gene has been of particular interest, and rs2254298 as well as rs53576 have previously been associated with amygdala activation in response to affective face perception in healthy individuals with samples sizes ranging from 55 to 228 (Tost et al., 2010, Tost et al., 2011, Marusak et al., 2015). We were not able to replicate these findings, with a healthy sample size of 142, and this could be due to lack of statistical power. However, our results are in line with findings from a large study investigating 23 tag SNPs in the OXTR in 1445 healthy adolescents (Loth et al., 2014). Loth and colleagues detected an association between rs237915 (that could serve as a proxy for rs237902) and activation in the ventral striatum, but not in amygdala, in response to threat perception (Loth et al., 2014), indicating that oxytocinergic signaling does play an important role in affective face

perception. The ventral striatum most likely interact with the amygdala in the social salience process (Shamay-Tsoory and Abu-Akel, 2016) and combined with our disorder specific association between rs237902 and amygdala activation in response to negative faces, it is possible that oxytocinergic signaling is altered in patients with schizophrenia compared to healthy individuals.

8.1.4 Oxytocin pathway genes – biological modulators of social dysfunction in psychotic disorders?

Despite the rich literature proposing oxytocin as a potential therapeutic target for schizophrenia, less is known of the underlying biological mechanisms of the endogenous oxytocin system (Leng and Ludwig, 2016). The rapidly growing amount of trials conducted with intranasal oxytocin is understandable, as few side-effects of the drug, negative subjective reactions or adverse effects are associated with this study method (MacDonald et al., 2011). In addition, oxytocin is already in clinical use for women in fertile age and is easy to administer. However, since no oxytocin receptors ligands are implemented in human clinical trials yet, the central pathways for this neuropeptide are difficult to investigate. This knowledge is of importance, since oxytocin administration has been associated with inconsistent treatment effects in this patient group and adverse effects are observed in subsets of samples. Also the biological mechanisms underlying the social disabilities that this hormone is supposed to alleviate are mostly unknown. Understanding the role of individual variation in the endogenous oxytocin system in these social disabilities would then be a step towards a more personalized medicine. We have used a translational approach measuring underlying genetics, brain activation and clinical symptomatology to understand this role better. Summarized, we found a subset of individuals that are more prone to emotional withdrawal (patients with

psychotic disorders carrying the OXTR rs53576A allele compared to rs53576G carriers) and a subset of individuals that are more prone to low amygdala activation during affective face perception of negative faces (patients with a schizophrenia spectrum disorder carrying the OXTR rs237902G allele compared to rs237902A carriers). This suggests that the endogenous oxytocin system has a role in biological underpinnings of negative symptoms and affective face perception. In study III, we found that this role differed between patients with schizophrenia spectrum disorders, affective spectrum disorders and healthy controls. In study II we did not test for disorder specificity of symptoms, but the more severe presence of negative symptoms in patients with schizophrenia compared to patients with affective disorders lies within the diagnostic criteria.

Our findings lend support from studies of schizophrenia animal models (Bartholomeusz et al., 2015). Rats dosed with the NMDA antagonist phencyclidine, displayed altered social behavior, reduced oxytocin mRNA in the anterior hypothalamus, and increased oxytocin receptor binding in the amygdala (Lee et al., 2005). In the same study, direct injection of oxytocin in the amygdala restored social interaction behavior. Other preclinical studies have shown that oxytocin may modulate PPI (Caldwell et al., 2009) and restore social deficits induced by dopamine agonists (Young et al., 2014).

As part of the social salience hypothesis, Rosenfeld and colleagues theorized that altered oxytocinergic signaling combined with dysfunctional dopamine signaling in the mesocorticolimbic system in patients with schizophrenia, leads to disrupted emotional salience to external cues in the environment (Rosenfeld et al., 2011). According to this theory, this aberrant emotion processing would lead to misinterpretations of social cues, which in turn would lead to negative symptoms (emotional withdrawal and isolation) and positive symptoms (suspicion and paranoia). Based on findings from animal studies and trials with

intranasal oxytocin as well as fMRI studies in patients with schizophrenia, they hypothesized that the oxytocinergic modulation of the amygdala is of greatest importance for this aberrant social salience (Rosenfeld et al., 2011).

Our findings support this theory, since OXTR polymorphisms showed association with amygdala activation during affective face perception and negative symptoms. Why different OXTR polymorphisms showed association with the investigated phenotypes in study II and III is challenging to understand, but could hypothetically be explained by that these polymorphisms both are important for oxytocin receptor expression in the mesocorticolimbic system, but at different brain sites. In addition, as mentioned above, other PANSS negative sub-items could possibly be more associated with rs237902 than the selected sub-items in this thesis. Negative symptoms in schizophrenia are clinical characteristics found to be correlated with emotion processing and face recognition in a meta-analysis (Ventura et al., 2013), strengthening the idea that they share biological underpinnings. But how the individual variations in oxytocin receptor polymorphisms in patients with psychotic disorders affect oxytocin receptor expression and oxytocin receptor distribution in amygdala, interaction with dopamine signaling and response to oxytocin administration remains unclear. To date, only a single trial with intranasal oxytocin measuring brain activation during an affective face perception task in patients with schizophrenia has been conducted (Shin et al., 2015). Shin and colleagues found that patients responded on oxytocin treatment with attenuated amygdala activation when presented to affective faces (combined result of negative, positive and neutral faces) compared to placebo, whereas healthy participants showed increased amygdala activation in the same paradigms. When patients and controls were investigated together (n=31), decreased amygdala activation for fearful emotion and increased activity for happy faces after oxytocin administration was seen (Shin et al., 2015). This finding needs to be replicated,

but if it holds true, even more questions about the modulating role of oxytocin in affective face perception emerge. If patients with schizophrenia with affective face perception deficits show lower amygdala activation when presented to fearful/angry emotional faces, as presented in three meta-analysis (Li et al., 2010, Taylor et al., 2012, Anticevic et al., 2012), a beneficial effect of even more decreased amygdala activation after oxytocin administration seems unlikely. Positive treatment effects have been difficult to detect in this patient group (Heringa et al., 2015), but importantly, no adverse effects have been reported (Feifel et al., 2016). In comparison, an attenuated effect on amygdala activation of oxytocin administration when presented to negative facial expressions has been seen in patients with generalized social anxiety disorder that in general show over-activation in the amygdala during affective face perception at baseline (Gorka et al., 2015), and an increased effect on amygdala activation in patients with ASD, that is abnormally under-activated at baseline (Domes et al., 2013).

A possible mechanism proposed by Shin and colleagues is that oxytocin reduces sensitivity to affective faces and social anxiety in patients with schizophrenia (Shin et al., 2015), and this explanation may be more in line with a hypothesis of anxiety reduction effects of oxytocin more than the social salience hypothesis. Our results neither support nor contradict this hypothesis, since we have not investigated how the participants in the present thesis respond to oxytocin administration. But one possible explanation for diagnostic differences in response to oxytocin administration on amygdala activation could be that oxytocinergic signaling influences this response. Based on our results combined with Shin and colleagues finding of a disorder specific response to oxytocin administration, and with the social salience hypothesis stating that individual characteristics influence treatment response, oxytocinergic signaling may influence this response in a different way in patients with schizophrenia

spectrum than in patients with affective spectrum disorders and healthy individuals. In fact, a recent study found that a haplotype block in the promotor region and intron 3 of OXTR, that comprise rs237897 (that is in moderate LD with rs237902) and rs53576 modulated oxytocin sensitivity to behavioral response (Chen et al., 2015). Thus, it is also possible that oxytocin administration would display different effects on affective face perception in patients with schizophrenia spectrum disorders carrying the rs237902G allele (that in our sample displayed a larger risk for low amygdala activation during affective face perception) compared to rs237902A carriers in the same patient group, and on the level of emotional withdrawal in patients with psychotic disorders carrying the rs53576A allele compared to rs53576G carriers.

Theoretically, genetic- or environmental risk factors can be associated with the risk for a particular disease phenotype in addition to – or independent of - the risk for the disorder per se. The first seen for the role of early adverse events in schizophrenia (Varese et al., 2012, Aas et al., 2012) and the second in coronary heart disease, another complex illness with strong genetic risk factors (Pranavchand and Reddy, 2013). Psychotic disorders are highly polygenic (Biological insights from 108 schizophrenia-associated genetic loci, 2014), hence the probability of detecting risk variants is minimal in a hypothesis based candidate gene approach. This is reflected by multiple negative results from dopamine candidate genes studies (Berry et al., 2003). Combined, this could explain why we were not able to detect associations with oxytocin pathway genes and a diagnosis of psychotic disorder per se.

The understanding that complex traits like psychiatric disorders are polygenic, implies that there are overlapping genetic variants between healthy individuals and patients. The genetic risk of disease is based on several hundred common genetic variants to some extent also carried by healthy individuals, and what combination of variants that ultimately is associated to a phenotype, e.g., extreme scores on personality traits, positive or negative symptoms,

affective face perception or a psychotic disorder per se, is still unknown. This accumulating genetic evidence is much of the basis of the dimensional theory to psychiatric disorders initiated by RDoC. Based on the correlation found between social dysfunction in both healthy individuals and patients with psychotic disorders and personality traits, the correlation seen between social dysfunction and oxytocin in the previous literature, and the inter-individual response seen in oxytocin administration, we hypothesized that common genetic variants in the oxytocin pathway genes could serve as such overlapping genetic variants. Our findings however failed to support this hypothesis. This could be explained by that oxytocin pathway genes are not central biological modulators of the investigated personality traits, an explanation that questions the social salience hypothesis. If the endogenous oxytocin system does not vary among individuals with different levels of Neuroticism, Extraversion or Agreeableness, it becomes less likely that oxytocinergic signaling influences anxiousness, hostility, vulnerability to stress, positive emotions, or trust in an inter-individual way. But, possibly could more rare variants in the oxytocin pathway genes or epigenetic variation may be more involved than the investigated alleles.

To conclude, we have found oxytocin pathway genes to be biological modulators of features related to social dysfunction in psychotic disorders to some extent- however with mixed results. Oxytocinergic signaling may contribute to affective face perception deficits and negative symptoms that are aspects of social salience, but are unlikely to explain most of the biological variance of social dysfunction in patients with psychotic disorders.

8.2 Strength and limitations

8.2.1 Sample and assessments

One of the main strengths of this thesis is the well characterized sample. This enabled us to use a translational approach measuring underlying genetics, brain activation and clinical symptomatology to better understand the endogenous oxytocin system across several psychotic disorders. These are phenotypes that are not available in the large PGC, where only information about diagnosis is present. This has provided us with the ability to explore pathophysiological mechanisms of these features which is highly requested in this research field (Leng and Ludwig, 2016). A second main strength of this thesis is the large patient sample. The field of oxytocin research is troubled by small sample sizes and our patient sample is to our knowledge the largest investigating symptom levels and oxytocin pathway genes. To date, Study III holds the first investigation of brain activation in relation to oxytocin pathway genes in patients with psychotic disorders.

To be included as a patient in the TOP-project, participants must be in a stable state of the disorder, for ethical and practical considerations. This has given us the opportunity to provide reliable measures of diagnosis, symptoms and brain activation, as these are demanding examinations for the patient. However, this is also reflected in the phenotypes that we targeted for investigation. In Study II, the mean symptom values for the PANSS were 2.33 for suspiciousness/persecution (p6), 1.22 for hostility (p7), 2.04 for emotional withdrawal (n2) and 2.19 for passive/apathetic social withdrawal (n4). A value score of 7 is rated as having the largest symptom burden in this 7-point rating instrument, which our patients are well below. In our study, we sought to investigate the relation to symptom severity, not the presence of symptoms, since only patients with a valid symptom measure were included. With overall low scores, our sample may not be representable for patients in less stable states. On the other

hand, the PANNS is considered a reliable tool for symptom assessment that is little prone to type II errors (Kay et al., 1987).

All patients described are recruited from in- and out-patient clinics in the greater Oslo area in South-Eastern Norway, and all participants were Caucasian. That is of importance, since the minor allele frequency of at least rs2254298 and rs53576 has shown correlations with ethnicity (Bakermans-Kranenburg and van Ijzendoorn, 2014). Population stratification, defined as the presence of a systematic difference in allele frequencies between subpopulations, is generally a possible confounder. If not corrected for, this might lead to both type I and type II errors. This issue seems however to trouble the oxytocin research field, since several studies regarding oxytocin pathway genes show population admixture (Bakermans-Kranenburg and van Ijzendoorn, 2014). For this reason we selected hypothesized SNPs for investigation that had been associated with schizophrenia phenotypes in Caucasian samples. Our sample had also fairly equally distributed gender in all studies which is an advantage considering the possible confounding effect between oxytocin pathway polymorphisms and phenotypes.

8.2.2 Methods

A major strength of this thesis is the investigation of symptoms and features across diagnostic categories. This has been stressed for future research, since the genetic overlap shown between psychotic disorders today indicate that the division of these heterogenic disorders does not reflect the etiological mechanisms (Lawrie et al., 2016). Then, the investigation of phenotypes in one categorical psychotic disorder could potentially reduce the statistical power to detect genetic risk. We approached this issue by exploring possible associations between oxytocin pathway genes and the phenotype itself, regardless of diagnosis. Further in study III,

to test if this association differed between diagnoses, we performed interaction analyses involving diagnosis and the associated polymorphism.

We chose a *hypothesis-based* approach and investigated oxytocin pathway genes that may be biological relevant to psychotic disorders, although they have not shown significant as susceptibility genes for development of psychotic disorders in the latest GWAS (Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near ODZ4, 2011b, Schizophrenia Working Group of the Psychiatric Genomics, 2014). With the missing heritability in mind, this investigation is still useful. In a case-control design, as we performed in study II, this however raises several issues. Firstly, and as mentioned above, psychotic disorders are highly polygenic and in a hypothesis based candidate gene approach, the probability of detecting risk variants is thus minimal. The need for a very large sample size is also prominent to reach sufficient statistical power, and we used the Genetic Power Calculator in study II. This calculator takes into account the risk allele frequency, disease prevalence, relative risk for homozygous/heterozygous carriers of the risk allele, LD between the investigated marker and the true risk allele, sample size, control/case ratio and if controls are screened for disease or not. Our calculated probabilities that the case/control analysis for rs53576 would reject a false null hypothesis were 0.51 for psychotic disorders and 0.34 for schizophrenia. There is obviously a need for a larger sample size to state that this polymorphism does not serve risk for a diagnosis per se, but with the largest patient sample to date investigating this issue, combined with the PGRS results, and the GWAS results from the large PGC, we find it plausible to state that there is little evidence that oxytocin pathway genes serve as risk variants to a diagnosis of any psychotic disorder per se.

With the large amount of risk alleles contributing to risk of psychotic disorders, and most likely to the social phenotypes investigated in this thesis, the effect-size estimated per SNP is

as a consequence low (Biological insights from 108 schizophrenia-associated genetic loci, 2014). This is reflected in the association between rs53576 and emotional withdrawal ($\beta = -0.203$) which should be taken into consideration for clinical implications.

Multiple tests were performed and corrected for in study I, leaving a stringent level of evidence required to detect a true positive finding ($p\text{-threshold} = 2.31 \times 10^{-4}$). This in turn reduced the statistical power of the study, with a high risk of making type II errors. Thus, the finding of no associations between personality traits and oxytocin pathway genes should be interpreted with caution. In study II, the association between rs53576 and emotional withdrawal did not survive Bonferroni corrections for 12 independent tests (4 PANSS items $\times$ 3 SNPs). We found this correction too strict, since all variables were pre-selected based on findings in previous literature, and thus all association tests performed were not independent tests that we screened for genetic association, but were based in pre-defined hypothesis. When all three SNPs were included in the same multiple regression model, the association between rs53576 and emotional withdrawal was still significant, and here also survived Bonferroni corrections. This strengthened the results and we find a significant association between rs53576 and emotional withdrawal plausible. In study III, the significance threshold was set to $p = 0.017$ due to Bonferroni corrections for three SNPs. The number of subgroups or ROI was not corrected for, due to a hypothesis based approach to diagnostic differences, and a strong correlation between left and right amygdala activation.

As no oxytocin related polymorphisms have been detected as risk variants for schizophrenia in large genome-wide association studies, we chose a hypothesis based approach for SNP selection. Rs53576, rs2254298 and rs237902 are the only SNPs previously found associated with disease characteristics in Caucasian schizophrenia samples (Bartholomeusz et al., 2015, Montag et al., 2012, Montag et al., 2013), hence we were interested in these polymorphisms

in particular. However, only a few studies have focused on the oxytocin pathway genes in patients with psychotic disorders, and the possibility of publication bias must be taken into account. The knowledge of biological consequences and functions of these variants are also limited, and their influence on brain function should be interpreted with caution.

8.3 Future directions

There are several unanswered questions to the main aim of this thesis; how the endogenous oxytocin system relate to features that are important for social dysfunction in patients with psychotic disorders. Firstly, replication of our findings is necessary in large samples. Secondly, further imaging studies involving other brain circuits and functional connectivity to amygdala as well as meta-analyses of oxytocin pathway genes in both healthy subjects and psychotic disorders should be performed. Additionally, several issues have not been addressed in this thesis, and would be of future interest.

Individuals constantly interact with the environment, and this will without doubt interact with the endogenous oxytocin system and in turn most likely influence cognitive, emotional and behavioral reactions. Epigenetic factors, defined as the genetic control by factors other than an individual's DNA sequence (Simmons, 2008), are to some extent under investigation, by gene-environment interactions and the measure of methylation states of the oxytocin pathway genes (Feldman et al., 2016). Methylation of the gene modulates gene expression, and individual variance in gene modulation could to some extent be addressed. To our knowledge, methylation of oxytocin pathway genes has been investigated in only one study in patients with psychotic disorders (Rubin et al., 2016), and not in relation to personality traits, which would be of interest for future research.

In terms of gene expression, mRNA analyses are generally also lacking in the field. Since plasma levels of oxytocin are considered unreliable measures of oxytocin levels in the brain, mRNA measured in the peripheral blood, could provide new information about individual and diagnostic differences in large samples. Further, regions of the genome containing DNA sequence variants that influence the expression level of the genes (expression quantitative trait loci, eQTL) could be identified.

Dopaminergic nuclei, including nucleus accumbens and ventral tegmental area, and prefrontal cortical areas most likely interact with the amygdala in affective face perception (Phillips et al., 2003, Gibbs et al., 2007, Green and Phillips, 2004), and dopamine antagonism from antipsychotic medication may disturb and attenuate this important function. If oxytocin should serve as a treatment option to these deficits among patients that already are medicated with dopamine-antagonists, it should be addressed how these neurotransmitter interact in humans. In a single-photon emission computed tomography (SPECT) study of healthy subjects, rs53576 interestingly predicted dopamine-transporter availability (Chang et al., 2014), but this has to our knowledge not been investigated in patients with psychotic disorders. Several other neurotransmitters influence oxytocin expression (Meyer-Lindenberg et al., 2011), and estrogen could be of particular interest in this regard. Sex differences in response to oxytocin administration are seen (Shamay-Tsoory and Abu-Akel, 2016), but little is known if estrogen levels could predict this response.

Ultimately, if individual differences in the genetic oxytocin system are better understood, they could possibly predict treatment response. This knowledge would however require a pharmacogenetic approach, and has been tested to our knowledge in one single study on healthy men (Chen et al., 2015). In order to be used in clinical settings, this must be replicated in several large studies, in patient samples and in both men and women.

9. CONCLUSION

Psychotic disorders and personality traits are complex and heterogeneous in their nature, and several contributing biological and environmental factors are involved. Based on a rich literature from animal studies as well as clinical trials, several hypotheses of oxytocin's role in these traits and disorders are generated, which we have scoped in this thesis.

We have used a dimensional approach to diagnosis, investigating features that are important for social dysfunction in patients with psychotic disorders both in a healthy sample and a broad set of psychotic disorders. We have provided new insight into the understanding of pathophysiological mechanisms of these features with possibly clinical implications. In particular we have shown:

1. That oxytocin pathway genes are not associated with the personality traits Neuroticism, Extraversion or Agreeableness and we find it less likely that oxytocinergic signaling influences the expression of these traits in an inter-individual way.
2. An association between the OXTR polymorphism rs53576 and emotional withdrawal in patients with psychotic disorders, a core feature related to social dysfunction in this patient group, which suggests that oxytocinergic signaling influences expression of this feature. We were however not able to detect associations between OXTR and suspiciousness/persecutory delusions, hostility or passive/apathetic social withdrawal or a predictive value of oxytocin pathway genes on case/control status. Our findings indicate that oxytocinergic signaling is not important in the pathogenesis of psychotic disorder per se.

3. An increased risk of low amygdala activation during affective face perception of negative faces in a subset of patients with schizophrenia spectrum disorders carrying the rs237902G allele compared to rs237902A carriers. This finding was disorder specific and indicates that oxytocinergic signaling influences affective face perception in this patient group in a different way than in patients with affective spectrum disorders or healthy controls.

10. REFERENCES

- AAS, M., STEEN, N. E., AGARTZ, I., AMINOFF, S. R., LORENTZEN, S., SUNDET, K., ANDREASSEN, O. A. & MELLE, I. (2012). Is cognitive impairment following early life stress in severe mental disorders based on specific or general cognitive functioning? *Psychiatry Res*, 198, pp.495-500.
- ADITYANJEE, ADERIBIGBE, Y. A., THEODORIDIS, D. & VIEWEG, V. R. (1999). Dementia praecox to schizophrenia: the first 100 years. *Psychiatry Clin Neurosci*, 53, pp.437-48.
- AGERBO, E., SULLIVAN, P. F., VILHJALMSSON, B. J., PEDERSEN, C. B., MORS, O., BORGLUM, A. D., HOUGAARD, D. M., HOLLEGAARD, M. V., MEIER, S., MATTHEISEN, M., RIPKE, S., WRAY, N. R. & MORTENSEN, P. B. (2015). Polygenic Risk Score, Parental Socioeconomic Status, Family History of Psychiatric Disorders, and the Risk for Schizophrenia: A Danish Population-Based Study and Meta-analysis. *JAMA Psychiatry*, 72, pp.635-41.
- AMERICAN PSYCHIATRIC ASSOCIATION (2000). *Diagnostic and statistical manual of mental disorders: DSM-IV-TR*. 4th ed., text rev. Washington, DC: American Psychiatric Association.
- ANDARI, E., SCHNEIDER, F. C., MOTTOLESE, R., VINDRAS, P. & SIRIGU, A. (2014). Oxytocin's fingerprint in personality traits and regional brain volume. *Cereb Cortex*, 24, pp.479-86.
- ANDREASSEN, O. A., THOMPSON, W. K. & DALE, A. M. (2014). Boosting the Power of Schizophrenia Genetics by Leveraging New Statistical Tools. *Schizophr Bull*, 40, pp.13-17.
- ANDREASSEN, O. A., THOMPSON, W. K., SCHORK, A. J., RIPKE, S., MATTINGSDAL, M., KELSOE, J. R., KENDLER, K. S., O'DONOVAN, M. C., RUJESCU, D., WERGE, T., SKLAR, P., RODDEY, J. C., CHEN, C. H., MCEVOY, L., DESIKAN, R. S., DJUROVIC, S. & DALE, A. M. (2013). Improved detection of common variants associated with schizophrenia and bipolar disorder using pleiotropy-informed conditional false discovery rate. *PLoS Genet* [Online], 9, pp.e1003455. DOI: 10.1371/journal.pgen.1003455.
- ANTICEVIC, A., VAN SNELLENBERG, J. X., COHEN, R. E., REPOVS, G., DOWD, E. C. & BARCH, D. M. (2012). Amygdala recruitment in schizophrenia in response to aversive emotional material: a meta-analysis of neuroimaging studies. *Schizophr Bull*, 38, pp.608-21.
- ASPÉ SÁNCHEZ, M., MORENO, M., RIVERA, M.-I., ROSSI, A. & EWER, J. (2016). Oxytocin and vasopressin receptor gene polymorphisms: role in social and psychiatric traits. *Front Neurosci* [Online], 9, pp.510. DOI: 10.3389/fnins.2015.00510.
- BAKERMANS-KRANENBURG, M. J. & VAN, I. J. M. H. (2013). Sniffing around oxytocin: review and meta-analyses of trials in healthy and clinical groups with implications for pharmacotherapy. *Transl Psychiatry* [Online], 3, pp.e258. DOI: 10.1038/tp.2013.34.
- BAKERMANS-KRANENBURG, M. J. & VAN IJZENDOORN, M. H. (2008). Oxytocin receptor (OXTR) and serotonin transporter (5-HTT) genes associated with observed parenting. *Soc Cogn Affect Neurosci*, 3, pp.128-34.
- BAKERMANS-KRANENBURG, M. J. & VAN IJZENDOORN, M. H. (2014). A sociability gene? Meta-analysis of oxytocin receptor genotype effects in humans. *Psychiatr Genet*, 24, pp.45-51.
- BALE, T. L., DAVIS, A. M., AUGER, A. P., DORSA, D. M. & MCCARTHY, M. M. (2001). CNS region-specific oxytocin receptor expression: importance in regulation of anxiety and sex behavior. *J Neurosci*, 21, pp.2546-52.
- BALESTRI, M., CALATI, R., SERRETTI, A. & DE RONCHI, D. (2014). Genetic modulation of personality traits: a systematic review of the literature. *Int Clin Psychopharmacol*, 29, pp.1-15.
- BARNETT, J. H., HUANG, J., PERLIS, R. H., YOUNG, M. M., ROSENBAUM, J. F., NIERENBERG, A. A., SACHS, G., NIMGAONKAR, V. L., MIKLOWITZ, D. J. & SMOLLER, J. W. (2011). Personality and bipolar disorder: dissecting state and trait associations between mood and personality. *Psychol Med*, 41, pp.1593-604.
- BARTHOLOMEUSZ, C. F., GANELLA, E. P., LABUSCHAGNE, I., BOUSMAN, C. & PANTELIS, C. (2015). Effects of oxytocin and genetic variants on brain and behaviour: Implications for treatment in schizophrenia. *Schizophr Res*, 168, pp.614-27.

- BARTZ, J., SIMEON, D., HAMILTON, H., KIM, S., CRYSTAL, S., BRAUN, A., VICENS, V. & HOLLANDER, E. (2011a). Oxytocin can hinder trust and cooperation in borderline personality disorder. *Soc Cogn Affect Neurosci*, 6, pp.556-63.
- BARTZ, J. A., ZAKI, J., BOLGER, N., HOLLANDER, E., LUDWIG, N. N., KOLEVZON, A. & OCHSNER, K. N. (2010). Oxytocin selectively improves empathic accuracy. *Psychol Sci*, 21, pp.1426-8.
- BARTZ, J. A., ZAKI, J., BOLGER, N. & OCHSNER, K. N. (2011b). Social effects of oxytocin in humans: context and person matter. *Trends Cogn Sci*, 15, pp.301-9.
- BEER, M. D. (1996a). The dichotomies: psychosis/neurosis and functional/organic: a historical perspective. *Hist Psychiatry*, 7, pp.231-55.
- BEER, M. D. (1996b). Psychosis: a history of the concept. *Compr Psychiatry*, 37, pp.273-91.
- BERRIDGE, K. C. (2007). The debate over dopamine's role in reward: the case for incentive salience. *Psychopharmacology (Berl)*, 191, pp.391-431.
- BERRIOS, G. E. (1996). *The History of Mental Symptoms* [Online]. Cambridge: Cambridge University Press. Available from: <http://dx.doi.org/10.1017/CBO9780511526725> [Accessed 24 Aug. 2016].
- BERRY, N., JOBANPUTRA, V. & PAL, H. (2003). Molecular genetics of schizophrenia: a critical review. *J Psychiatry Neurosci*, 28, pp.415-429.
- BETHLEHEM, R. A., VAN HONK, J., AUYEUNG, B. & BARON-COHEN, S. (2013). Oxytocin, brain physiology, and functional connectivity: a review of intranasal oxytocin fMRI studies. *Psychoneuroendocrinology*, 38, pp.962-74.
- BIENVENU, O. J., DAVYDOW, D. S. & KENDLER, K. S. (2011). Psychiatric 'diseases' versus behavioral disorders and degree of genetic influence. *Psychol Med*, 41, pp.33-40.
- BIOLOGICAL INSIGHTS FROM 108 SCHIZOPHRENIA-ASSOCIATED GENETIC LOCI. (2014). *Nature*, 511, pp.421-7.
- BLEULER, E. (1911). *Dementia praecox, oder Gruppe der Schizophrenien*. Leipzig: Deuticke.
- BORA, E. & PANTELIS, C. (2016). Social cognition in schizophrenia in comparison to bipolar disorder: A meta-analysis. *Schizophr Res*, 175, pp.72-8.
- BRANDES, M. & BIENVENU, O. J. (2006). Personality and anxiety disorders. *Curr Psychiatry Rep*, 8, pp.263-9.
- BRANDT, C. L., EICHELE, T., MELLE, I., SUNDET, K., SERVER, A., AGARTZ, I., HUGDAHL, K., JENSEN, J. & ANDREASSEN, O. A. (2014). Working memory networks and activation patterns in schizophrenia and bipolar disorder: comparison with healthy controls. *Br J Psychiatry*, 204, pp.290-8.
- BRIGGS, S. R. (1992). Assessing the five-factor model of personality description. *J Pers*, 60, pp.253-93.
- BRUNET-GOUET, E. & DECETY, J. (2006). Social brain dysfunctions in schizophrenia: a review of neuroimaging studies. *Psychiatry Res*, 148, pp.75-92.
- BUJANOW, W. (1972). Hormones in the treatment of psychoses. *Br Med J*, 4, pp.298.
- BUTTON, K. S., IOANNIDIS, J. P., MOKRYSZ, C., NOSEK, B. A., FLINT, J., ROBINSON, E. S. & MUNAFO, M. R. (2013). Power failure: why small sample size undermines the reliability of neuroscience. *Nat Rev Neurosci*, 14, pp.365-76.
- CACCIOTTI-SAIJA, C., LANGDON, R., WARD, P. B., HICKIE, I. B., SCOTT, E. M., NAISMITH, S. L., MOORE, L., ALVARES, G. A., REDOBLADO HODGE, M. A. & GUASTELLA, A. J. (2015). A double-blind randomized controlled trial of oxytocin nasal spray and social cognition training for young people with early psychosis. *Schizophr Bull*, 41, pp.483-93.
- CALDWELL, H. K., STEPHENS, S. L. & YOUNG, W. S., 3RD (2009). Oxytocin as a natural antipsychotic: a study using oxytocin knockout mice. *Mol Psychiatry*, 14, pp.190-6.
- CANTOR-GRAAE, E. & PEDERSEN, C. B. (2013). Full spectrum of psychiatric disorders related to foreign migration: a Danish population-based cohort study. *JAMA Psychiatry*, 70, pp.427-35.
- CARDOSO, C., ELLENBOGEN, M. A. & LINNEN, A. M. (2012a). Acute intranasal oxytocin improves positive self-perceptions of personality. *Psychopharmacology (Berl)*, 220, pp.741-9.

- CARDOSO, C., LINNEN, A. M., JOOBER, R. & ELLENBOGEN, M. A. (2012b). Coping style moderates the effect of intranasal oxytocin on the mood response to interpersonal stress. *Exp Clin Psychopharmacol*, 20, pp.84-91.
- CARRE, J. M., FISHER, P. M., MANUCK, S. B. & HARIRI, A. R. (2012). Interaction between trait anxiety and trait anger predict amygdala reactivity to angry facial expressions in men but not women. *Soc Cogn Affect Neurosci*, 7, pp.213-21.
- CASE, M., STAUFFER, V. L., ASCHER-SVANUM, H., CONLEY, R., KAPUR, S., KANE, J. M., KOLLACK-WALKER, S., JACOB, J. & KINON, B. J. (2010). The heterogeneity of antipsychotic response in the treatment of schizophrenia. *Psychol Med*, pp.1-10.
- CERMOLACCE, M., SASS, L. & PARNAS, J. (2010). What is bizarre in bizarre delusions? A critical review. *Schizophr Bull*, 36, pp.667-79.
- CHANG, W. H., LEE, I. H., CHEN, K. C., CHI, M. H., CHIU, N. T., YAO, W. J., LU, R. B., YANG, Y. K. & CHEN, P. S. (2014). Oxytocin receptor gene rs53576 polymorphism modulates oxytocin-dopamine interaction and neuroticism traits--a SPECT study. *Psychoneuroendocrinology*, 47, pp.212-20.
- CHEN, C. H., SUCKLING, J., LENNOX, B. R., OOI, C. & BULLMORE, E. T. (2011). A quantitative meta-analysis of fMRI studies in bipolar disorder. *Bipolar Disord*, 13, pp.1-15.
- CHEN, F. S., KUMSTA, R., DVORAK, F., DOMES, G., YIM, O. S., EBSTEIN, R. P. & HEINRICHS, M. (2015). Genetic modulation of oxytocin sensitivity: a pharmacogenetic approach. *Transl Psychiatry* [Online], 5, pp.e664. DOI: 10.1038/tp.2015.163.
- CHOLERIS, E., GUSTAFSSON, J. A., KORACH, K. S., MUGLIA, L. J., PFAFF, D. W. & OGAWA, S. (2003). An estrogen-dependent four-gene micronet regulating social recognition: a study with oxytocin and estrogen receptor-alpha and -beta knockout mice. *Proc Natl Acad Sci U S A*, 100, pp.6192-7.
- CHURCHLAND, P. S. & WINKIELMAN, P. (2012). Modulating social behavior with oxytocin: how does it work? What does it mean? *Horm Behav*, 61, pp.392-9.
- COMPTON, M. T., BAKEMAN, R., ALOLAYAN, Y., BALDUCCI, P. M., BERNARDINI, F., BROUSSARD, B., CRISAFIO, A., CRISTOFARO, S., AMAR, P., JOHNSON, S. & WAN, C. R. (2015). Personality domains, duration of untreated psychosis, functioning, and symptom severity in first-episode psychosis. *Schizophr Res*, 168, pp.113-9.
- CORCORAN, C. M., KEILP, J. G., KAYSER, J., KLIM, C., BUTLER, P. D., BRUDER, G. E., GUR, R. C. & JAVITT, D. C. (2015). Emotion recognition deficits as predictors of transition in individuals at clinical high risk for schizophrenia: a neurodevelopmental perspective. *Psychol Med*, 45, pp.2959-73.
- COSTA, P. T. & MCCRAE, R. R. (1992). Normal personality assessment in clinical practice: The NEO Personality Inventory. *Psychol Assess*, 4, pp.5-13.
- COSTAFREDA, S. G., BRAMMER, M. J., DAVID, A. S. & FU, C. H. Y. (2008). Predictors of amygdala activation during the processing of emotional stimuli: A meta-analysis of 385 PET and fMRI studies. *Brain Res Rev*, 58, pp.57-70.
- CRADDOCK, N. & OWEN, M. J. (2010). The Kraepelinian dichotomy – going, going... but still not gone. *Br J Psychiatry*, 196, pp.92-95.
- CRADDOCK, N. & SKLAR, P. (2013). Genetics of bipolar disorder. *Lancet*, 381, pp.1654-62.
- CRAN.R-PROJECT.ORG. (2015). *The Comprehensive R Archive Network*. [Online]. Vienna. Available from: <https://cran.r-project.org> [Accessed 24 Aug. 2016].
- CROCQ, M.-A. (2013). Milestones in the history of personality disorders. *Dialogues Clin Neurosci*, 15, pp.147-153.
- DADDS, M. R., MOUL, C., CAUCHI, A., DOBSON-STONE, C., HAWES, D. J., BRENNAN, J., URWIN, R. & EBSTEIN, R. E. (2014). Polymorphisms in the oxytocin receptor gene are associated with the development of psychopathy. *Dev Psychopathol*, 26, pp.21-31.

- DAVIS, M. C., HORAN, W. P., NURMI, E. L., RIZZO, S., LI, W., SUGAR, C. A. & GREEN, M. F. (2014). Associations between oxytocin receptor genotypes and social cognitive performance in individuals with schizophrenia. *Schizophr Res*, 159, pp.353-7.
- DAVIS, M. C., LEE, J., HORAN, W. P., CLARKE, A. D., MCGEE, M. R., GREEN, M. F. & MARDER, S. R. (2013). Effects of single dose intranasal oxytocin on social cognition in schizophrenia. *Schizophr Res*, 147, pp.393-7.
- DE DREU, C. K. (2012). Oxytocin modulates cooperation within and competition between groups: an integrative review and research agenda. *Horm Behav*, 61, pp.419-28.
- DE DREU, C. K., GREER, L. L., HANDGRAAF, M. J., SHALVI, S., VAN KLEEF, G. A., BAAS, M., TEN VELDEN, F. S., VAN DIJK, E. & FEITH, S. W. (2010). The neuropeptide oxytocin regulates parochial altruism in intergroup conflict among humans. *Science*, 328, pp.1408-11.
- DE FRUYT, F., VAN DE WIELE, L. & VAN HEERINGEN, C. (2000). Cloninger's Psychobiological Model of Temperament and Character and the Five-Factor Model of Personality. *Pers Individ Dif*, 29, pp.441-452.
- DECLERCK, C. H., BOONE, C. & KIYONARI, T. (2010). Oxytocin and cooperation under conditions of uncertainty: the modulating role of incentives and social information. *Horm Behav*, 57, pp.368-74.
- DERKS, E. M., VORSTMAN, J. A., RIPKE, S., KAHN, R. S. & OPHOFF, R. A. (2012). Investigation of the genetic association between quantitative measures of psychosis and schizophrenia: a polygenic risk score analysis. *PLoS One* [Online], 7, pp.e37852. DOI: 10.1371/journal.pone.0037852.
- DEWALL, C. N., GILLATH, O., PRESSMAN, S. D., BLACK, L. L., BARTZ, J. A., MOSKOVITZ, J. & STETLER, D. A. (2014). When the Love Hormone Leads to Violence: Oxytocin Increases Intimate Partner Violence Inclinations Among High Trait Aggressive People. *Soc Psychol Personal Sci*, 5, pp.691-697.
- DOMES, G., HEINRICHS, M., KUMBIER, E., GROSSMANN, A., HAUENSTEIN, K. & HERPERTZ, S. C. (2013). Effects of intranasal oxytocin on the neural basis of face processing in autism spectrum disorder. *Biol Psychiatry*, 74, pp.164-171.
- DOMES, G., LISCHKE, A., BERGER, C., GROSSMANN, A., HAUENSTEIN, K., HEINRICHS, M. & HERPERTZ, S. C. (2010). Effects of intranasal oxytocin on emotional face processing in women. *Psychoneuroendocrinology*, 35, pp.83-93.
- DU VIGNEAUD, V. (1960). EXPERIENCES IN THE POLYPEPTIDE FIELD: INSULIN TO OXYTOCIN. *Ann N Y Acad Sci*, 88, pp.537-548.
- DUDBRIDGE, F. (2016). Polygenic Epidemiology. *Genet Epidemiol*, 40, pp.268-72.
- DUNNER, D. L., FLEISS, J. L. & FIEVE, R. R. (1976). The course of development of mania in patients with recurrent depression. *Am J Psychiatry*, 133, pp.905-8.
- EBSTEIN, R. P., ISRAEL, S., LERER, E., UZEFOVSKY, F., SHALEV, I., GRITSENKO, I., RIEBOLD, M., SALOMON, S. & YIRMIYA, N. (2009). Arginine vasopressin and oxytocin modulate human social behavior. *Ann N Y Acad Sci*, 1167, pp.87-102.
- EBSTEIN, R. P., KNAFO, A., MANKUTA, D., CHEW, S. H. & LAI, P. S. (2012). The contributions of oxytocin and vasopressin pathway genes to human behavior. *Horm Behav*, 61, pp.359-79.
- EDWARDS, J., JACKSON, H. J. & PATTISON, P. E. (2002). Emotion recognition via facial expression and affective prosody in schizophrenia: a methodological review. *Clin Psychol Rev*, 22, pp.789-832.
- ENGEL, G. L. (1977). The need for a new medical model: a challenge for biomedicine. *Science*, 196, pp.129-36.
- ENSEMBL.ORG. (2016). *Ensembl genome browser 85*. [Online] Available from: <http://www.ensembl.org/index.html> [Accessed 24 Aug. 2016].
- FEIFEL, D., SHILLING, P. D. & MACDONALD, K. (2016). A Review of Oxytocin's Effects on the Positive, Negative, and Cognitive Domains of Schizophrenia. *Biol Psychiatry*, 79, pp.222-33.

- FELDMAN, R., MONAKHOV, M., PRATT, M. & EBSTEIN, R. P. (2016). Oxytocin Pathway Genes: Evolutionary Ancient System Impacting on Human Affiliation, Sociality, and Psychopathology. *Biol Psychiatry*, 79, pp.174-84.
- FELDMAN, R., ZAGOORY-SHARON, O., WEISMAN, O., SCHNEIDERMAN, I., GORDON, I., MAOZ, R., SHALEV, I. & EBSTEIN, R. P. (2012). Sensitive Parenting Is Associated with Plasma Oxytocin and Polymorphisms in the OXTR and CD38 Genes. *Biol Psychiatry*, 72, pp.175-81.
- FERGUSON, J. N., ALDAG, J. M., INSEL, T. R. & YOUNG, L. J. (2001). Oxytocin in the medial amygdala is essential for social recognition in the mouse. *J Neurosci*, 21, pp.8278-85.
- FETT, A. K., VIECHTBAUER, W., DOMINGUEZ, M. D., PENN, D. L., VAN OS, J. & KRABBENDAM, L. (2011). The relationship between neurocognition and social cognition with functional outcomes in schizophrenia: a meta-analysis. *Neurosci Biobehav Rev*, 35, pp.573-88.
- FIRST, M. B. (2002). The DSM series and experience with DSM-IV. *Psychopathology*, 35, pp.67-71.
- FLICEK, P., AMODE, M. R., BARRELL, D., BEAL, K., BILLIS, K., BRENT, S., CARVALHO-SILVA, D., CLAPHAM, P., COATES, G., FITZGERALD, S., GIL, L., GIRÓN, C. G., GORDON, L., HOURLIER, T., HUNT, S., JOHNSON, N., JUETTEMANN, T., KÄHÄRI, A. K., KEENAN, S., KULESHA, E., MARTIN, F. J., MAUREL, T., MCLAREN, W. M., MURPHY, D. N., NAG, R., OVERDUIN, B., PIGNATELLI, M., PRITCHARD, B., PRITCHARD, E., RIAT, H. S., RUFFIER, M., SHEPPARD, D., TAYLOR, K., THORMANN, A., TREVANION, S. J., VULLO, A., WILDER, S. P., WILSON, M., ZADISSA, A., AKEN, B. L., BIRNEY, E., CUNNINGHAM, F., HARROW, J., HERRERO, J., HUBBARD, T. J. P., KINSELLA, R., MUFFATO, M., PARKER, A., SPUDICH, G., YATES, A., ZERBINO, D. R. & SEARLE, S. M. J. (2014). Ensembl 2014. *Nucleic Acids Res*, 42, pp.D749-D755.
- FMRIB.OX.AC.UK. (2016). *FSL - FslWiki*. [Online] Available from: <http://www.fmrib.ox.ac.uk/fsl> [Accessed 24 Aug. 2016].
- FOLAND, L. C., ALTSHULER, L. L., SUGAR, C. A., LEE, A. D., LEOW, A. D., TOWNSEND, J., NARR, K. L., ASUNCION, D. M., TOGA, A. W. & THOMPSON, P. M. (2008). Increased volume of the amygdala and hippocampus in bipolar patients treated with lithium. *Neuroreport*, 19, pp.221-4.
- FUSAR-POLI, P., DESTE, G., SMIESKOVA, R. & ET AL. (2012). Cognitive functioning in prodromal psychosis: A meta-analysis. *Arch Gen Psychiatry*, 69, pp.562-571.
- GENOME.SPH.UMICH.EDU. (2016a). *ChunkChromosome - Genome Analysis Wiki*. [Online] Available from: <http://genome.sph.umich.edu/wiki/ChunkChromosome> [Accessed 24 Aug. 2016].
- GENOME.SPH.UMICH.EDU. (2016b). *Minimac - Genome Analysis Wiki*. [Online] Available from: <http://genome.sph.umich.edu/wiki/Minimac> [Accessed 24 Aug. 2016].
- GENOME-WIDE ASSOCIATION STUDY IDENTIFIES FIVE NEW SCHIZOPHRENIA LOCI. (2011a). *Nat Genet*, 43, pp.969-76.
- GIBBS, A. A., NAUDTS, K. H., SPENCER, E. P. & DAVID, A. S. (2007). The role of dopamine in attentional and memory biases for emotional information. *Am J Psychiatry*, 164, pp.1603-9; quiz 1624.
- GIBSON, C. M., PENN, D. L., SMEDLEY, K. L., LESERMAN, J., ELLIOTT, T. & PEDERSEN, C. A. (2014). A pilot six-week randomized controlled trial of oxytocin on social cognition and social skills in schizophrenia. *Schizophr Res*, 156, pp.261-5.
- GIMPL, G. & FAHRENHOLZ, F. (2001). The oxytocin receptor system: structure, function, and regulation. *Physiol Rev*, 81, pp.629-83.
- GLEESON, J. F., RAWLINGS, D., JACKSON, H. J. & MCGORRY, P. D. (2005). Agreeableness and neuroticism as predictors of relapse after first-episode psychosis: a prospective follow-up study. *J Nerv Ment Dis*, 193, pp.160-9.
- GLOVINSKY, D., KALOGERAS, K. T., KIRCH, D. G., SUDDATH, R. & WYATT, R. J. (1994). Cerebrospinal fluid oxytocin concentration in schizophrenic patients does not differ from control subjects and is not changed by neuroleptic medication. *Schizophr Res*, 11, pp.273-6.

- GOLDMAN, M., MARLOW-O'CONNOR, M., TORRES, I. & CARTER, C. S. (2008). Diminished plasma oxytocin in schizophrenic patients with neuroendocrine dysfunction and emotional deficits. *Schizophr Res*, 98, pp.247-55.
- GOPALAKRISHNA, G., ITHMAN, M. H. & LAURIELLO, J. (2016). Update on New and Emerging Treatments for Schizophrenia. *Psychiatr Clin North Am*, 39, pp.217-38.
- GORKA, S. M., FITZGERALD, D. A., LABUSCHAGNE, I., HOSANAGAR, A., WOOD, A. G., NATHAN, P. J. & PHAN, K. L. (2015). Oxytocin modulation of amygdala functional connectivity to fearful faces in generalized social anxiety disorder. *Neuropsychopharmacology*, 40, pp.278-86.
- GOTTESMAN, I. I. & SHIELDS, J. (1967). A polygenic theory of schizophrenia. *Proc Natl Acad Sci U S A*, 58, pp.199-205.
- GREEN, M. F., HORAN, W. P. & LEE, J. (2015). Social cognition in schizophrenia. *Nat Rev Neurosci*, 16, pp.620-31.
- GREEN, M. J. & PHILLIPS, M. L. (2004). Social threat perception and the evolution of paranoia. *Neurosci Biobehav Rev*, 28, pp.333-42.
- GRINEVICH, V., KNOBLOCH-BOLLMANN, H. S., ELIAVA, M., BUSNELLI, M. & CHINI, B. (2016). Assembling the Puzzle: Pathways of Oxytocin Signaling in the Brain. *Biol Psychiatry*, 79, pp.155-64.
- GUR, R. C. & GUR, R. E. (2016). Social cognition as an RDoC domain. *Am J Med Genet B Neuropsychiatr Genet*, 171b, pp.132-41.
- HALL, J., WHALLEY, H. C., MCKIRDY, J. W., ROMANIUK, L., MCGONIGLE, D., MCINTOSH, A. M., BAIG, B. J., GOUNTOUNA, V.-E., JOB, D. E., DONALDSON, D. I., SPRENGELMEYER, R., YOUNG, A. W., JOHNSTONE, E. C. & LAWRIE, S. M. (2008). Overactivation of Fear Systems to Neutral Faces in Schizophrenia. *Biol Psychiatry*, 64, pp.70-73.
- HARIRI, A. R., MATTAY, V. S., TESSITORE, A., KOLACHANA, B., FERA, F., GOLDMAN, D., EGAN, M. F. & WEINBERGER, D. R. (2002). Serotonin transporter genetic variation and the response of the human amygdala. *Science*, 297, pp.400-3.
- HARRISON, P. J. (2015). Recent genetic findings in schizophrenia and their therapeutic relevance. *J Psychopharmacol*, 29, pp.85-96.
- HARVEY, P. D., GREEN, M. F., KEEFE, R. S. & VELLIGAN, D. I. (2004). Cognitive functioning in schizophrenia: a consensus statement on its role in the definition and evaluation of effective treatments for the illness. *J Clin Psychiatry*, 65, pp.361-72.
- HERINGA, S. M., BEGEMANN, M. J., GOVERDE, A. J. & SOMMER, I. E. (2015). Sex hormones and oxytocin augmentation strategies in schizophrenia: A quantitative review. *Schizophr Res*, 168, pp.603-13.
- HERPERTZ, S. C. & BERTSCH, K. (2016). Oxytocin Effects on Brain Functioning in Humans. *Biol Psychiatry*, 79, pp.631-632.
- HOPWOOD, C. J., MOREY, L. C., ANSELL, E. B., GRILO, C. M., SANISLOW, C. A., MCGLASHAN, T. H., MARKOWITZ, J. C., GUNDERSON, J. G., YEN, S., SHEA, M. T. & SKODOL, A. E. (2009). THE CONVERGENT AND DISCRIMINANT VALIDITY OF FIVE-FACTOR TRAITS: CURRENT AND PROSPECTIVE SOCIAL, WORK, AND RECREATIONAL DYSFUNCTION. *J Pers Disord*, 23, pp.466-476.
- INOUE, H., YAMASUE, H., TOCHIGI, M., ABE, O., LIU, X., KAWAMURA, Y., TAKEI, K., SUGA, M., YAMADA, H., ROGERS, M. A., AOKI, S., SASAKI, T. & KASAI, K. (2010). Association between the oxytocin receptor gene and amygdalar volume in healthy adults. *Biol Psychiatry*, 68, pp.1066-72.
- INOUE, T., KIMURA, T., AZUMA, C., INAZAWA, J., TAKEMURA, M., KIKUCHI, T., KUBOTA, Y., OGITA, K. & SAJI, F. (1994). Structural organization of the human oxytocin receptor gene. *J Biol Chem*, 269, pp.32451-6.
- INSEL, T. R. (2014). The NIMH Research Domain Criteria (RDoC) Project: Precision Medicine for Psychiatry. *Am J Psychiatry*, 171, pp.395-397.

- INSEL, T. R. & YOUNG, L. J. (2001). The neurobiology of attachment. *Nat Rev Neurosci*, 2, pp.129-36.
- JENKINSON, M., BANNISTER, P., BRADY, M. & SMITH, S. (2002). Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. *Neuroimage*, 17, pp.825-841.
- JIN, D., LIU, H. X., HIRAI, H., TORASHIMA, T., NAGAI, T., LOPATINA, O., SHNAYDER, N. A., YAMADA, K., NODA, M., SEIKE, T., FUJITA, K., TAKASAWA, S., YOKOYAMA, S., KOIZUMI, K., SHIRAISHI, Y., TANAKA, S., HASHII, M., YOSHIHARA, T., HIGASHIDA, K., ISLAM, M. S., YAMADA, N., HAYASHI, K., NOGUCHI, N., KATO, I., OKAMOTO, H., MATSUSHIMA, A., SALMINA, A., MUNESUE, T., SHIMIZU, N., MOCHIDA, S., ASANO, M. & HIGASHIDA, H. (2007). CD38 is critical for social behaviour by regulating oxytocin secretion. *Nature*, 446, pp.41-5.
- KARPENKO, I. A., MARGATHE, J. F., RODRIGUEZ, T., PFLIMLIN, E., DUPUIS, E., HIBERT, M., DURROUX, T. & BONNET, D. (2015). Selective nonpeptidic fluorescent ligands for oxytocin receptor: design, synthesis, and application to time-resolved FRET binding assay. *J Med Chem*, 58, pp.2547-52.
- KAWAMURA, Y., LIU, X., AKIYAMA, T., SHIMADA, T., OTOWA, T., SAKAI, Y., KAKIUCHI, C., UMEKAGE, T., SASAKI, T. & AKISKAL, H. S. (2010). The association between oxytocin receptor gene (OXTR) polymorphisms and affective temperaments, as measured by TEMPS-A. *J Affect Disord*, 127, pp.31-7.
- KAY, S. R., FISZBEIN, A. & OPLER, L. A. (1987). The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr Bull*, 13, pp.261-76.
- KENDLER, K. S. & JABLENSKY, A. (2011). Kraepelin's concept of psychiatric illness. *Psychol Med*, 41, pp.1119-1126.
- KERI, S., KISS, I. & KELEMEN, O. (2009). Sharing secrets: oxytocin and trust in schizophrenia. *Soc Neurosci*, 4, pp.287-93.
- KIDD, S. A. (2013). From social experience to illness experience: reviewing the psychological mechanisms linking psychosis with social context. *Can J Psychiatry*, 58, pp.52-8.
- KIM, H. N., KIM, B. H., CHO, J., RYU, S., SHIN, H., SUNG, J., SHIN, C., CHO, N. H., SUNG, Y. A., CHOI, B. O. & KIM, H. L. (2015). Pathway analysis of genome-wide association datasets of personality traits. *Genes Brain Behav*, 14, pp.345-56.
- KIM, H. N., ROH, S. J., SUNG, Y. A., CHUNG, H. W., LEE, J. Y., CHO, J., SHIN, H. & KIM, H. L. (2013). Genome-wide association study of the five-factor model of personality in young Korean women. *J Hum Genet*, 58, pp.667-74.
- KIM, H. S., SHERMAN, D. K., SASAKI, J. Y., XU, J., CHU, T. Q., RYU, C., SUH, E. M., GRAHAM, K. & TAYLOR, S. E. (2010). Culture, distress, and oxytocin receptor polymorphism (OXTR) interact to influence emotional support seeking. *Proc Natl Acad Sci U S A*, 107, pp.15717-21.
- KIMURA, T., SAJI, F., NISHIMORI, K., OGITA, K., NAKAMURA, H., KOYAMA, M. & MURATA, Y. (2003). Molecular regulation of the oxytocin receptor in peripheral organs. *J Mol Endocrinol*, 30, pp.109-15.
- KIMURA, T., TANIZAWA, O., MORI, K., BROWNSTEIN, M. J. & OKAYAMA, H. (1992). Structure and expression of a human oxytocin receptor. *Nature*, 356, pp.526-9.
- KLEIN, D. N., KOTOV, R. & BUFFERD, S. J. (2011). Personality and depression: explanatory models and review of the evidence. *Annu Rev Clin Psychol*, 7, pp.269-95.
- KLEIST, K. (1953). [Classification of nervous and mental diseases]. *Monatsschr Psychiatr Neurol*, 125, pp.526-54.
- KOHLER, C. G., TURNER, T. H., BILKER, W. B., BRENSINGER, C. M., SIEGEL, S. J., KANES, S. J., GUR, R. E. & GUR, R. C. (2003). Facial emotion recognition in schizophrenia: intensity effects and error pattern. *Am J Psychiatry*, 160, pp.1768-74.
- KOHLER, C. G., WALKER, J. B., MARTIN, E. A., HEALEY, K. M. & MOBERG, P. J. (2010). Facial emotion perception in schizophrenia: a meta-analytic review. *Schizophr Bull*, 36, pp.1009-19.

- KOSFELD, M., HEINRICHS, M., ZAK, P. J., FISCHBACHER, U. & FEHR, E. (2005). Oxytocin increases trust in humans. *Nature*, 435, pp.673-6.
- KRAEPELIN, E. (1910). *Psychiatrie: ein Lehrbuch für Studierende und Ärzte* [Online]. Leipzig: Barth. Available from:
<https://ia802602.us.archive.org/26/items/psychiatrieeinle04krae/psychiatrieeinle04krae.pdf>
 [Accessed 24 Aug. 2016].
- KRUEGER, F., PARASURAMAN, R., MOODY, L., TWIEG, P., DE VISSER, E., MCCABE, K., O'HARA, M. & LEE, M. R. (2013). Oxytocin selectively increases perceptions of harm for victims but not the desire to punish offenders of criminal offenses. *Soc Cogn Affect Neurosci*, 8, pp.494-8.
- KUCHARSKA-PIETURA, K. & MORTIMER, A. (2013). Can antipsychotics improve social cognition in patients with schizophrenia? *CNS Drugs*, 27, pp.335-43.
- KUMSTA, R. & HEINRICHS, M. (2013). Oxytocin, stress and social behavior: neurogenetics of the human oxytocin system. *Curr Opin Neurobiol*, 23, pp.11-6.
- LANDER, E. S., LINTON, L. M., BIRREN, B., NUSBAUM, C., ZODY, M. C., BALDWIN, J., DEVON, K., DEWAR, K., DOYLE, M., FITZHUGH, W., FUNKE, R., GAGE, D., HARRIS, K., HEAFORD, A., HOWLAND, J., KANN, L., LEHOCZKY, J., LEVINE, R., MCEWAN, P., MCKERNAN, K., MELDRIM, J., MESIROV, J. P., MIRANDA, C., MORRIS, W., NAYLOR, J., RAYMOND, C., ROSETTI, M., SANTOS, R., SHERIDAN, A., SOUGNEZ, C., STANGE-THOMANN, Y., STOJANOVIC, N., SUBRAMANIAN, A., WYMAN, D., ROGERS, J., SULSTON, J., AINSCOUGH, R., BECK, S., BENTLEY, D., BURTON, J., CLEE, C., CARTER, N., COULSON, A., DEADMAN, R., DELOUKAS, P., DUNHAM, A., DUNHAM, I., DURBIN, R., FRENCH, L., GRAFHAM, D., GREGORY, S., HUBBARD, T., HUMPHRAY, S., HUNT, A., JONES, M., LLOYD, C., MCMURRAY, A., MATTHEWS, L., MERCER, S., MILNE, S., MULLIKIN, J. C., MUNGALL, A., PLUMB, R., ROSS, M., SHOWNKEEN, R., SIMS, S., WATERSTON, R. H., WILSON, R. K., HILLIER, L. W., MCPHERSON, J. D., MARRA, M. A., MARDIS, E. R., FULTON, L. A., CHINWALLA, A. T., PEPIN, K. H., GISH, W. R., CHISSOE, S. L., WENDL, M. C., DELEHAUNTY, K. D., MINER, T. L., DELEHAUNTY, A., KRAMER, J. B., COOK, L. L., FULTON, R. S., JOHNSON, D. L., MINX, P. J., CLIFTON, S. W., HAWKINS, T., BRANSCOMB, E., PREDKI, P., RICHARDSON, P., WENNING, S., SLEZAK, T., DOGGETT, N., CHENG, J. F., OLSEN, A., LUCAS, S., ELKIN, C., UBERBACHER, E., FRAZIER, M., et al. (2001). Initial sequencing and analysis of the human genome. *Nature*, 409, pp.860-921.
- LANDGRAF, R., NEUMANN, I., RUSSELL, J. A. & PITTMAN, Q. J. (1992). Push-pull perfusion and microdialysis studies of central oxytocin and vasopressin release in freely moving rats during pregnancy, parturition, and lactation. *Ann N Y Acad Sci*, 652, pp.326-39.
- LANGEVELD, J., ANDREASSEN, O. A., AUESTAD, B., FAERDEN, A., HAUGE, L. J., JOA, I., JOHANNESSEN, J. O., MELLE, I., RUND, B. R., ROSSBERG, J. I., SIMONSEN, E., VAGLUM, P. & LARSEN, T. K. (2013). Is there an optimal factor structure of the Positive and Negative Syndrome Scale in patients with first-episode psychosis? *Scand J Psychol*, 54, pp.160-5.
- LARGE-SCALE GENOME-WIDE ASSOCIATION ANALYSIS OF BIPOLAR DISORDER IDENTIFIES A NEW SESCEPTIBILITY LOCUS NEAR ODZ4. (2011b). *Nat Genet*, 43, pp.977-983.
- LAWRIE, S. M., O'DONOVAN, M. C., SAKS, E., BURNS, T. & LIEBERMAN, J. A. (2016). Towards diagnostic markers for the psychoses. *The Lancet Psychiatry* [Online], 3, pp.375-385. DOI: 10.1016/S2215-0366(16)00021-3.
- LEBOYER, M., HENRY, C., PAILLIERE-MARTINOT, M. L. & BELLIVIER, F. (2005). Age at onset in bipolar affective disorders: a review. *Bipolar Disord*, 7, pp.111-8.
- LEE, P. R., BRADY, D. L., SHAPIRO, R. A., DORSA, D. M. & KOENIG, J. I. (2005). Social interaction deficits caused by chronic phencyclidine administration are reversed by oxytocin. *Neuropsychopharmacology*, 30, pp.1883-94.
- LEE, T. Y., HONG, S. B., SHIN, N. Y. & KWON, J. S. (2015). Social cognitive functioning in prodromal psychosis: A meta-analysis. *Schizophr Res*, 164, pp.28-34.

- LENG, G. & LUDWIG, M. (2016). Intranasal Oxytocin: Myths and Delusions. *Biol Psychiatry*, 79, pp.243-50.
- LENG, G., MEDDLE, S. L. & DOUGLAS, A. J. (2008). Oxytocin and the maternal brain. *Curr Opin Pharmacol*, 8, pp.731-4.
- LENG, G., PINEDA, R., SABATIER, N. & LUDWIG, M. (2015). 60 YEARS OF NEUROENDOCRINOLOGY: The posterior pituitary, from Geoffrey Harris to our present understanding. *J Endocrinol*, 226, pp.T173-85.
- LEVY, B. & MANOVE, E. (2012). Functional outcome in bipolar disorder: the big picture. *Depress Res Treat*, 2012, pp.949248.
- LEWANDOWSKI, K. E., COHEN, B. M. & ÖNGUR, D. (2011). Evolution of neuropsychological dysfunction during the course of schizophrenia and bipolar disorder. *Psychol Med*, 41, pp.225-241.
- LI, H., CHAN, R. C., MCALONAN, G. M. & GONG, Q. Y. (2010). Facial emotion processing in schizophrenia: a meta-analysis of functional neuroimaging data. *Schizophr Bull*, 36, pp.1029-39.
- LI, J., ZHAO, Y., LI, R., BROSTER, L. S., ZHOU, C. & YANG, S. (2015). Association of Oxytocin Receptor Gene (OXTR) rs53576 Polymorphism with Sociality: A Meta-Analysis. *PLoS One* [Online], 10, pp.e0131820. DOI: 10.1371/journal.pone.0131820.
- LICHTENSTEIN, P., YIP, B. H., BJORK, C., PAWITAN, Y., CANNON, T. D., SULLIVAN, P. F. & HULTMAN, C. M. (2009). Common genetic determinants of schizophrenia and bipolar disorder in Swedish families: a population-based study. *Lancet*, 373, pp.234-9.
- LIEBERMAN, J. A., PERKINS, D., BELGER, A., CHAKOS, M., JARSKOG, F., BOTEVA, K. & GILMORE, J. (2001). The early stages of schizophrenia: Speculations on pathogenesis, pathophysiology, and therapeutic approaches. *Biol Psychiatry*, 50, pp.884-897.
- LIEBERMAN, J. A., STROUP, T. S., MCEVOY, J. P., SWARTZ, M. S., ROSENHECK, R. A., PERKINS, D. O., KEEFE, R. S., DAVIS, S. M., DAVIS, C. E., LEBOWITZ, B. D., SEVERE, J. & HSIAO, J. K. (2005). Effectiveness of antipsychotic drugs in patients with chronic schizophrenia. *N Engl J Med*, 353, pp.1209-23.
- LIN, P. I., VANCE, J. M., PERICAK-VANCE, M. A. & MARTIN, E. R. (2007). No gene is an island: the flip-flop phenomenon. *Am J Hum Genet*, 80, pp.531-8.
- LINKOWSKI, P., GEENEN, V., KERKHOFS, M., MENDLEWICZ, J. & LEGROS, J. J. (1984). Cerebrospinal fluid neurophysins in affective illness and in schizophrenia. *Eur Arch Psychiatry Neurol Sci*, 234, pp.162-5.
- LIU, Y. & WANG, Z. X. (2003). Nucleus accumbens oxytocin and dopamine interact to regulate pair bond formation in female prairie voles. *Neuroscience*, 121, pp.537-44.
- LONNQVIST, J. E., VERKASALO, M., HAUKKA, J., NYMAN, K., TIIHONEN, J., LAAKSONEN, I., LESKINEN, J., LONNQVIST, J. & HENRIKSSON, M. (2009). Premorbid personality factors in schizophrenia and bipolar disorder: results from a large cohort study of male conscripts. *J Abnorm Psychol*, 118, pp.418-23.
- LOPARO, D. & WALDMAN, I. D. (2015). The oxytocin receptor gene (OXTR) is associated with autism spectrum disorder: a meta-analysis. *Mol Psychiatry*, 20, pp.640-6.
- LOTH, E., POLINE, J. B., THYREAU, B., JIA, T., TAO, C., LOURDUSAMY, A., STACEY, D., CATTRELL, A., DESRIVIERES, S., RUGGERI, B., FRITSCH, V., BANASCHEWSKI, T., BARKER, G. J., BOKDE, A. L., BUCHEL, C., CARVALHO, F. M., CONROD, P. J., FAUTH-BUEHLER, M., FLOR, H., GALLINAT, J., GARAVAN, H., HEINZ, A., BRUEHL, R., LAWRENCE, C., MANN, K., MARTINOT, J. L., NEES, F., PAUS, T., PAUSOVA, Z., POUSTKA, L., RIETSCHER, M., SMOLKA, M., STRUVE, M., FENG, J. & SCHUMANN, G. (2014). Oxytocin receptor genotype modulates ventral striatal activity to social cues and response to stressful life events. *Biol Psychiatry*, 76, pp.367-76.

- LOUP, F., TRIBOLLET, E., DUBOIS-DAUPHIN, M. & DREIFUSS, J. J. (1991). Localization of high-affinity binding sites for oxytocin and vasopressin in the human brain. An autoradiographic study. *Brain Res*, 555, pp.220-32.
- LOUP, F., TRIBOLLET, E., DUBOIS-DAUPHIN, M., PIZZOLATO, G. & DREIFUSS, J. J. (1989). Localization of oxytocin binding sites in the human brainstem and upper spinal cord: an autoradiographic study. *Brain Res*, 500, pp.223-30.
- LUDWIG, M. (1998). Dendritic release of vasopressin and oxytocin. *J Neuroendocrinol*, 10, pp.881-95.
- LUDWIG, M. & LENG, G. (2006). Dendritic peptide release and peptide-dependent behaviours. *Nat Rev Neurosci*, 7, pp.126-36.
- LUMINET, O., BAGBY, R. M., WAGNER, H., TAYLOR, G. J. & PARKER, J. D. (1999). Relation between alexithymia and the five-factor model of personality: a facet-level analysis. *J Pers Assess*, 73, pp.345-58.
- LUMINET, O., GRYNBERG, D., RUZETTE, N. & MIKOLAJCZAK, M. (2011). Personality-dependent effects of oxytocin: greater social benefits for high alexithymia scorers. *Biol Psychol*, 87, pp.401-6.
- MACDONALD, E., DADDS, M. R., BRENNAN, J. L., WILLIAMS, K., LEVY, F. & CAUCHI, A. J. (2011). A review of safety, side-effects and subjective reactions to intranasal oxytocin in human research. *Psychoneuroendocrinology*, 36, pp.1114-26.
- MACDONALD, K. & MACDONALD, T. M. (2010). The peptide that binds: a systematic review of oxytocin and its prosocial effects in humans. *Harv Rev Psychiatry*, 18, pp.1-21.
- MALAVASI, F., DEAGLIO, S., FUNARO, A., FERRERO, E., HORENSTEIN, A. L., ORTOLAN, E., VAISITTI, T. & AYDIN, S. (2008). Evolution and Function of the ADP Ribosyl Cyclase/CD38 Gene Family in Physiology and Pathology. *Physiol Rev*, 88, pp.841-886.
- MALIK, A. I., ZAI, C. C., ABU, Z., NOWROUZI, B. & BEITCHMAN, J. H. (2012). The role of oxytocin and oxytocin receptor gene variants in childhood-onset aggression. *Genes Brain Behav*, 11, pp.545-51.
- MANNING, M., STOEV, S., CHINI, B., DURROUX, T., MOUILLAC, B. & GUILLON, G. (2008). Peptide and non-peptide agonists and antagonists for the vasopressin and oxytocin V1a, V1b, V2 and OT receptors: research tools and potential therapeutic agents. *Prog Brain Res*, 170, pp.473-512.
- MARUSAK, H. A., FURMAN, D. J., KURUVADI, N., SHATTUCK, D. W., JOSHI, S. H., JOSHI, A. A., ETKIN, A. & THOMASON, M. E. (2015). Amygdala responses to salient social cues vary with oxytocin receptor genotype in youth. *Neuropsychologia*, 79, pp.1-9.
- MCCLAY, J. L., ADKINS, D. E., ABERG, K., BUKSZAR, J., KHACHANE, A. N., KEEFE, R. S., PERKINS, D. O., MCEVOY, J. P., STROUP, T. S., VANN, R. E., BEARDSLEY, P. M., LIEBERMAN, J. A., SULLIVAN, P. F. & VAN DEN OORD, E. J. (2011). Genome-wide pharmacogenomic study of neurocognition as an indicator of antipsychotic treatment response in schizophrenia. *Neuropsychopharmacology*, 36, pp.616-26.
- MCCRAE, R. R., COSTA JR, P. T. & MARTIN, T. A. (2005). The NEO-PI-3: a more readable revised NEO Personality Inventory. *J Pers Assess*, 84, pp.261-70.
- MCEWEN, B. B. (2004a). Brain-fluid barriers: relevance for theoretical controversies regarding vasopressin and oxytocin memory research. *Adv Pharmacol*, 50, pp.531-92, 655-708.
- MCEWEN, B. B. (2004b). General introduction to vasopressin and oxytocin: structure/metabolism, evolutionary aspects, neural pathway/receptor distribution, and functional aspects relevant to memory processing. *Adv Pharmacol*, 50, pp.1-50, 655-708.
- MERCEDES PEREZ-RODRIGUEZ, M., MAHON, K., RUSSO, M., UNGAR, A. K. & BURDICK, K. E. (2015). Oxytocin and social cognition in affective and psychotic disorders. *Eur Neuropsychopharmacol*, 25, pp.265-82.
- MEYER-LINDENBERG, A., DOMES, G., KIRSCH, P. & HEINRICHS, M. (2011). Oxytocin and vasopressin in the human brain: social neuropeptides for translational medicine. *Nat Rev Neurosci*, 12, pp.524-38.

- MITRE, M., MARLIN, B. J., SCHIAVO, J. K., MORINA, E., NORDEN, S. E., HACKETT, T. A., AOKI, C. J., CHAO, M. V. & FROEMKE, R. C. (2016). A Distributed Network for Social Cognition Enriched for Oxytocin Receptors. *J Neurosci*, 36, pp.2517-35.
- MONTAG, C., BROCKMANN, E. M., BAYERL, M., RUJESCU, D., MULLER, D. J. & GALLINAT, J. (2013). Oxytocin and oxytocin receptor gene polymorphisms and risk for schizophrenia: a case-control study. *World J Biol Psychiatry*, 14, pp.500-8.
- MONTAG, C., BROCKMANN, E. M., LEHMANN, A., MULLER, D. J., RUJESCU, D. & GALLINAT, J. (2012). Association between oxytocin receptor gene polymorphisms and self-rated 'empathic concern' in schizophrenia. *PLoS One* [Online], 7, pp.e51882. DOI: 10.1371/journal.pone.0051882.
- MONTAG, C. & REUTER, M. (2014). Disentangling the molecular genetic basis of personality: From monoamines to neuropeptides. *Neurosci Biobehav Rev*, 43, pp.228-239.
- MUNESUE, T., YOKOYAMA, S., NAKAMURA, K., ANITHA, A., YAMADA, K., HAYASHI, K., ASAKA, T., LIU, H. X., JIN, D., KOIZUMI, K., ISLAM, M. S., HUANG, J. J., MA, W. J., KIM, U. H., KIM, S. J., PARK, K., KIM, D., KIKUCHI, M., ONO, Y., NAKATANI, H., SUDA, S., MIYACHI, T., HIRAI, H., SALMINA, A., PICHUGINA, Y. A., SOUMAROKOV, A. A., TAKEI, N., MORI, N., TSUJII, M., SUGIYAMA, T., YAGI, K., YAMAGISHI, M., SASAKI, T., YAMASUE, H., KATO, N., HASHIMOTO, R., TANIKE, M., HAYASHI, Y., HAMADA, J., SUZUKI, S., OOI, A., NODA, M., KAMIYAMA, Y., KIDO, M. A., LOPATINA, O., HASHII, M., AMINA, S., MALAVASI, F., HUANG, E. J., ZHANG, J., SHIMIZU, N., YOSHIKAWA, T., MATSUSHIMA, A., MINABE, Y. & HIGASHIDA, H. (2010). Two genetic variants of CD38 in subjects with autism spectrum disorder and controls. *Neurosci Res*, 67, pp.181-91.
- NATIONAL CENTER FOR BIOTECHNOLOGY INFORMATION (2010). *PubChem Compound Database*; CID=439302 [Online]. Bethesda, MD. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/439302> [Accessed 24 August 2016].
- NATURE.COM. (2016). *single nucleotide polymorphism / SNP | Learn Science at Scitable*. [Online] Available from: <http://www.nature.com/scitable/definition/single-nucleotide-polymorphism-snp-295> [Accessed 24 Aug. 2016].
- NEUMANN, I., LUDWIG, M., ENGELMANN, M., PITTMAN, Q. J. & LANDGRAF, R. (1993). Simultaneous microdialysis in blood and brain: oxytocin and vasopressin release in response to central and peripheral osmotic stimulation and suckling in the rat. *Neuroendocrinology*, 58, pp.637-45.
- NISHIOKA, T., ANSELMO-FRANCI, J. A., LI, P., CALLAHAN, M. F. & MORRIS, M. (1998). Stress increases oxytocin release within the hypothalamic paraventricular nucleus. *Brain Res*, 781, pp.56-60.
- NORDGAARD, J., ARNFRED, S. M., HANDEST, P. & PARNAS, J. (2008). The Diagnostic Status of First-Rank Symptoms. *Schizophr Bull*, 34, pp.137-154.
- OHI, K., SHIMADA, T., NITTA, Y., KIHARA, H., OKUBO, H., UEHARA, T. & KAWASAKI, Y. (2016). The Five-Factor Model personality traits in schizophrenia: A meta-analysis. *Psychiatry Res*, 240, pp.34-41.
- OLFF, M., FRIJLING, J. L., KUBZANSKY, L. D., BRADLEY, B., ELLENBOGEN, M. A., CARDOSO, C., BARTZ, J. A., YEE, J. R. & VAN ZUIDEN, M. (2013). The role of oxytocin in social bonding, stress regulation and mental health: an update on the moderating effects of context and interindividual differences. *Psychoneuroendocrinology*, 38, pp.1883-94.
- OUSDAL, O. T., ANAND BROWN, A., JENSEN, J., NAKSTAD, P. H., MELLE, I., AGARTZ, I., DJUROVIC, S., BOGDAN, R., HARIRI, A. R. & ANDREASSEN, O. A. (2012). Associations between variants near a monoaminergic pathways gene (PHOX2B) and amygdala reactivity: a genome-wide functional imaging study. *Twin Res Hum Genet*, 15, pp.273-85.
- OWEN, M. J. (2014). New approaches to psychiatric diagnostic classification. *Neuron*, 84, pp.564-571.
- OWEN, M. J., SAWA, A. & MORTENSEN, P. B. (2016). Schizophrenia. *Lancet*, 388, pp.86-97.
- PAGANI, F. & BARALLE, F. E. (2004). Genomic variants in exons and introns: identifying the splicing spoilers. *Nat Rev Genet*, 5, pp.389-396.

- PEDERSEN, C. A., GIBSON, C. M., RAU, S. W., SALIMI, K., SMEDLEY, K. L., CASEY, R. L., LESERMAN, J., JARSKOG, L. F. & PENN, D. L. (2011). Intranasal oxytocin reduces psychotic symptoms and improves Theory of Mind and social perception in schizophrenia. *Schizophr Res*, 132, pp.50-3.
- PENNISI, E. (2012). ENCODE Project Writes Eulogy for Junk DNA. *Science*, 337, pp.1159-1161.
- PERKEL, J. (2008). SNP genotyping: six technologies that keyed a revolution. *Nat Meth*, 5, pp.447-453.
- PHELPS, E. A. & LEDOUX, J. E. (2005). Contributions of the amygdala to emotion processing: from animal models to human behavior. *Neuron*, 48, pp.175-87.
- PHILLIPS, A. G., AHN, S. & HOWLAND, J. G. (2003). Amygdalar control of the mesocorticolimbic dopamine system: parallel pathways to motivated behavior. *Neurosci Biobehav Rev*, 27, pp.543-54.
- PHILLIPS, M. L. & SWARTZ, H. A. (2014). A critical appraisal of neuroimaging studies of bipolar disorder: toward a new conceptualization of underlying neural circuitry and a road map for future research. *Am J Psychiatry*, 171, pp.829-43.
- PNGU.MGH.HARVARD.EDU. (2016). *Genetic Power Calculator*. [Online] Available from: <http://pngu.mgh.harvard.edu/~purcell/gpc/> [Accessed 24 Aug. 2016].
- POWER, R. A. & PLUESS, M. (2015). Heritability estimates of the Big Five personality traits based on common genetic variants. *Transl Psychiatry* [Online], 5, pp.e604. DOI: 10.1038/tp.2015.96.
- PRANAVCHAND, R. & REDDY, B. M. (2013). Current status of understanding of the genetic etiology of coronary heart disease. *J Postgrad Med*, 59, pp.30-41.
- PURCELL, S., CHERNY, S. S. & SHAM, P. C. (2003). Genetic Power Calculator: design of linkage and association genetic mapping studies of complex traits. *Bioinformatics*, 19, pp.149-50.
- PURCELL, S., NEALE, B., TODD-BROWN, K., THOMAS, L., FERREIRA, M. A., BENDER, D., MALLER, J., SKLAR, P., DE BAKKER, P. I., DALY, M. J. & SHAM, P. C. (2007). PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*, 81, pp.559-75.
- PURCELL, S. (2009a). *Whole genome association analysis toolset. PLINK: Whole genome data analysis toolset* [Online]. Available from: <http://pngu.mgh.harvard.edu/purcell/plink> [Accessed 23 August 2016].
- PURCELL, S. M., WRAY, N. R., STONE, J. L., VISSCHER, P. M., O'DONOVAN, M. C., SULLIVAN, P. F. & SKLAR, P. (2009b). Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature*, 460, pp.748-52.
- QI, J., YANG, J.-Y., SONG, M., LI, Y., WANG, F. & WU, C.-F. (2007). Inhibition by oxytocin of methamphetamine-induced hyperactivity related to dopamine turnover in the mesolimbic region in mice. *Naunyn Schmiedeberg's Arch Pharmacol*, 376, pp.441-448.
- QIN, K., DONG, C., WU, G. & LAMBERT, N. A. (2011). Inactive-state preassembly of G(q)-coupled receptors and G(q) heterotrimers. *Nat Chem Biol*, 7, pp.740-7.
- QUINTANA, D. S., GUASTELLA, A. J., WESTLYE, L. T. & ANDREASSEN, O. A. (2016). The promise and pitfalls of intranasally administering psychopharmacological agents for the treatment of psychiatric disorders. *Mol Psychiatry*, 21, pp.29-38.
- RILLING, J. K., DEMARCO, A. C., HACKETT, P. D., CHEN, X., GAUTAM, P., STAIR, S., HAROON, E., THOMPSON, R., DITZEN, B., PATEL, R. & PAGNONI, G. (2014). Sex differences in the neural and behavioral response to intranasal oxytocin and vasopressin during human social interaction. *Psychoneuroendocrinology*, 39, pp.237-248.
- RIMOL, L. M., HARTBERG, C. B., NESVAG, R., FENNEMA-NOTESTINE, C., HAGLER, D. J., JR., PUNG, C. J., JENNINGS, R. G., HAUKVIK, U. K., LANGE, E., NAKSTAD, P. H., MELLE, I., ANDREASSEN, O. A., DALE, A. M. & AGARTZ, I. (2010). Cortical thickness and subcortical volumes in schizophrenia and bipolar disorder. *Biol Psychiatry*, 68, pp.41-50.
- RIPKE, S., SANDERS, A. R., KENDLER, K. S., LEVINSON, D. F., SKLAR, P., HOLMANS, P. A., LIN, D. Y., DUAN, J., OPHOFF, R. A., ANDREASSEN, O. A., SCOLNICK, E., CICHON, S., ST CLAIR, D., CORVIN, A., GURLING, H., WERGE, T., RUJESCU, D., BLACKWOOD, D. H., PATO, C. N., MALHOTRA, A. K., PURCELL, S., DUDBRIDGE, F., NEALE, B. M., ROSSIN, L., VISSCHER, P. M.,

- POSTHUMA, D., RUDERFER, D. M., FANOUS, A., STEFANSSON, H., STEINBERG, S., MOWRY, B. J., GOLIMBET, V., DE HERT, M., JONSSON, E. G., BITTER, I., PIETILAINEN, O. P., COLLIER, D. A., TOSATO, S., AGARTZ, I., ALBUS, M., ALEXANDER, M., AMDUR, R. L., AMIN, F., BASS, N., BERGEN, S. E., BLACK, D. W., BORGLUM, A. D., BROWN, M. A., BRUGGEMAN, R., BUCCOLA, N. G., BYERLEY, W. F., CAHN, W., CANTOR, R. M., CARR, V. J., CATTS, S. V., CHOUDHURY, K., CLONINGER, C. R., CORMICAN, P., CRADDOCK, N., DANOY, P. A., DATTA, S., DE HAAN, L., DEMONTIS, D., DIKEOS, D., DJUROVIC, S., DONNELLY, P., DONOHOE, G., DUONG, L., DWYER, S., FINK-JENSEN, A., FREEDMAN, R., FREIMER, N. B., FRIEDL, M., GEORGIEVA, L., GIEGLING, I., GILL, M., GLENTHOJ, B., GODARD, S., HAMSHERE, M., HANSEN, M., HANSEN, T., HARTMANN, A. M., HENSKENS, F. A., HOUGAARD, D. M., HULTMAN, C. M., INGASON, A., JABLENSKY, A. V., JAKOBSEN, K. D., JAY, M., JURGENS, G., KAHN, R. S., KELLER, M. C., KENIS, G., KENNY, E., KIM, Y., KIROV, G. K., KONNERTH, H., KONTE, B., KRABBENDAM, L., KRASUCKI, R., et al. (2011). Genome-wide association study identifies five new schizophrenia loci. *Nat Genet*, 43, pp.969-976.
- RODRIGUES, S. M., SASLOW, L. R., GARCIA, N., JOHN, O. P. & KELTNER, D. (2009). Oxytocin receptor genetic variation relates to empathy and stress reactivity in humans. *Proc Natl Acad Sci U S A*, 106, pp.21437-41.
- ROSENFELD, A. J., LIEBERMAN, J. A. & JARSKOG, L. F. (2011). Oxytocin, dopamine, and the amygdala: a neurofunctional model of social cognitive deficits in schizophrenia. *Schizophr Bull*, 37, pp.1077-87.
- RUBIN, L. H., CARTER, C. S., DROGOS, L., POURNAJAFI-NAZARLOO, H., SWEENEY, J. A. & MAKI, P. M. (2010). Peripheral oxytocin is associated with reduced symptom severity in schizophrenia. *Schizophr Res*, 124, pp.13-21.
- RUBIN, L. H., CONNELLY, J. J., REILLY, J. L., CARTER, C. S., DROGOS, L. L., POURNAJAFI-NAZARLOO, H., RUOCCO, A. C., KEEDY, S. K., MATTHEW, I., TANDON, N., PEARLSON, G. D., CLEMENTZ, B. A., TAMMINGA, C. A., GERSHON, E. S., KESHAVAN, M. S., BISHOP, J. R. & SWEENEY, J. A. (2016). Sex and diagnosis specific associations between DNA methylation of the oxytocin receptor gene with emotion processing and temporal-limbic and prefrontal brain volumes in psychotic disorders. *Biol Psychiatry Cogn Neurosci Neuroimaging*, 1, pp.141-151.
- RUDERFER, D. M., FANOUS, A. H., RIPKE, S., MCQUILLIN, A., AMDUR, R. L., GEJMAN, P. V., O'DONOVAN, M. C., ANDREASSEN, O. A., DJUROVIC, S., HULTMAN, C. M., KELSOE, J. R., JAMAIN, S., LANDEN, M., LEBOYER, M., NIMGAONKAR, V., NURNBERGER, J., SMOLLER, J. W., CRADDOCK, N., CORVIN, A., SULLIVAN, P. F., HOLMANS, P., SKLAR, P. & KENDLER, K. S. (2014). Polygenic dissection of diagnosis and clinical dimensions of bipolar disorder and schizophrenia. *Mol Psychiatry*, 19, pp.1017-1024.
- RUOCCO, A. C., REILLY, J. L., RUBIN, L. H., DAROS, A. R., GERSHON, E. S., TAMMINGA, C. A., PEARLSON, G. D., HILL, S. K., KESHAVAN, M. S., GUR, R. C. & SWEENEY, J. A. (2014). Emotion recognition deficits in schizophrenia-spectrum disorders and psychotic bipolar disorder: Findings from the Bipolar-Schizophrenia Network on Intermediate Phenotypes (B-SNIP) study. *Schizophr Res*, 158, pp.105-12.
- RUTIGLIANO, G., ROCCHETTI, M., PALOYELIS, Y., GILLEEN, J., SARDELLA, A., CAPPUCCIATI, M., PALOMBINI, E., DELL'OSSO, L., CAVERZASI, E., POLITI, P., MCGUIRE, P. & FUSAR-POLI, P. (2016). Peripheral oxytocin and vasopressin: Biomarkers of psychiatric disorders? A comprehensive systematic review and preliminary meta-analysis. *Psychiatry Res*, 241, pp.207-220.
- SABATINELLI, D., FORTUNE, E. E., LI, Q., SIDDIQUI, A., KRAFFT, C., OLIVER, W. T., BECK, S. & JEFFRIES, J. (2011). Emotional perception: Meta-analyses of face and natural scene processing. *Neuroimage*, 54, pp.2524-2533.

- SALMINA, A. B., LOPATINA, O., EKIMOVA, M. V., MIKHUTKINA, S. V. & HIGASHIDA, H. (2010). CD38/cyclic ADP-ribose system: a new player for oxytocin secretion and regulation of social behaviour. *J Neuroendocrinol*, 22, pp.380-92.
- SAUNA, Z. E. & KIMCHI-SARFATY, C. (2011). Understanding the contribution of synonymous mutations to human disease. *Nat Rev Genet*, 12, pp.683-691.
- SCHIZOPHRENIA COMMISSION (2012). *The abandoned illness: a report from the Schizophrenia Commission* [Online]. London: Rethink Mental Illness. Available from: <https://www.rethink.org/about-us/the-schizophrenia-commission> [Accessed 24th August 2016].
- SCHIZOPHRENIA WORKING GROUP OF THE PSYCHIATRIC GENOMICS, C. (2014). Biological insights from 108 schizophrenia-associated genetic loci. *Nature*, 511, pp.421-427.
- SCHNEIDER-HASSLOFF, H., STRAUBE, B., JANSEN, A., NUSCHELER, B., WEMKEN, G., WITT, S. H., RIETSCHER, M. & KIRCHER, T. (2016). Oxytocin receptor polymorphism and childhood social experiences shape adult personality, brain structure and neural correlates of mentalizing. *Neuroimage*, 134, pp.671-684.
- SEBAT, J. (2007). Major changes in our DNA lead to major changes in our thinking. *Nat Genet*, 39, pp.S3-5.
- SHAHROKH, D. K., ZHANG, T. Y., DIORIO, J., GRATTON, A. & MEANEY, M. J. (2010). Oxytocin-dopamine interactions mediate variations in maternal behavior in the rat. *Endocrinology*, 151, pp.2276-86.
- SHAMAY-TSOORY, S. G. & ABU-AKEL, A. (2016). The Social Salience Hypothesis of Oxytocin. *Biol Psychiatry*, 79, pp.194-202.
- SHAMAY-TSOORY, S. G., FISCHER, M., DVASH, J., HARARI, H., PERACH-BLOOM, N. & LEVKOVITZ, Y. (2009). Intranasal administration of oxytocin increases envy and schadenfreude (gloating). *Biol Psychiatry*, 66, pp.864-70.
- SHIN, N. Y., PARK, H. Y., JUNG, W. H., PARK, J. W., YUN, J. Y., JANG, J. H., KIM, S. N., HAN, H. J., KIM, S. Y., KANG, D. H. & KWON, J. S. (2015). Effects of Oxytocin on Neural Response to Facial Expressions in Patients with Schizophrenia. *Neuropsychopharmacology*, 40, pp.1919-27.
- SIMMONS, D. (2008). Epigenetic influence and disease. *Nature Education* [Online], 1, pp.6. Available from: <http://www.nature.com/scitable/topicpage/epigenetic-influences-and-disease-895> [Accessed 24 August 2016].
- SIMONSEN, C., SUNDET, K., VASKINN, A., UELAND, T., ROMM, K. L., HELLVIN, T., MELLE, I., FRIIS, S. & ANDREASSEN, O. A. (2010). Psychosocial function in schizophrenia and bipolar disorder: Relationship to neurocognition and clinical symptoms. *J Int Neuropsychol Soc*, 16, pp.771-83.
- SLATKIN, M. (2008). Linkage disequilibrium [mdash] understanding the evolutionary past and mapping the medical future. *Nat Rev Genet*, 9, pp.477-485.
- SMITH, A. L., FREEMAN, S. M., BARNHART, T. E., ABBOTT, D. H., AHLERS, E. O., KUKIS, D. L., BALES, K. L., GOODMAN, M. M. & YOUNG, L. J. (2016). Initial investigation of three selective and potent small molecule oxytocin receptor PET ligands in New World monkeys. *Bioorg Med Chem Lett*, 26, pp.3370-5.
- SMITH, S. M. (2002). Fast robust automated brain extraction. *Hum Brain Mapp*, 17, pp.143-155.
- SMITH, S. M. & BRADY, J. M. (1997). SUSAN—A New Approach to Low Level Image Processing. *Int J Comput Vis*, 23, pp.45-78.
- SMITH, S. M., JENKINSON, M., WOOLRICH, M. W., BECKMANN, C. F., BEHRENS, T. E., JOHANSEN-BERG, H., BANNISTER, P. R., DE LUCA, M., DROBNJAK, I., FLITNEY, D. E., NIAZY, R. K., SAUNDERS, J., VICKERS, J., ZHANG, Y., DE STEFANO, N., BRADY, J. M. & MATTHEWS, P. M. (2004). Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage*, 23 Suppl 1, pp.S208-19.
- SOLYOM, L., LEDWIDGE, B. & SOLYOM, C. (1986). Delineating social phobia. *Br J Psychiatry*, 149, pp.464-70.

- SOUZA, R. P., DE LUCA, V., MELTZER, H. Y., LIEBERMAN, J. A. & KENNEDY, J. L. (2010a). Schizophrenia severity and clozapine treatment outcome association with oxytocinergic genes. *Int J Neuropsychopharmacol*, 13, pp.793-8.
- SOUZA, R. P., ISMAIL, P., MELTZER, H. Y. & KENNEDY, J. L. (2010b). Variants in the oxytocin gene and risk for schizophrenia. *Schizophr Res*, 121, pp.279-80.
- SPH.UMICH.EDU. (2016a). *1000G Phase1 Alpha Download*. [Online] Available from: <http://www.sph.umich.edu/csg/abecasis/MACH/download/1000G-Phase1-Interim.html> [Accessed 24 Aug. 2016].
- SPH.UMICH.EDU. (2016b). *MACH Download*. [Online] Available from: <http://www.sph.umich.edu/csg/abecasis/MaCH/download/> [Accessed 24 Aug. 2016].
- STANKOVA, T., EICHHAMMER, P., LANGGUTH, B. & SAND, P. G. (2012). Sexually dimorphic effects of oxytocin receptor gene (OXTR) variants on Harm Avoidance. *Biol Sex Differ*, 3, pp.17.
- SUMMAR, M. L., JOHN A. PHILLIPS, I., BATTEY, J., CASTIGLIONE, C. M., KIDD, K. K., MANESS, K. J., WEIFFENBACH, B. & GRAVIUS, T. C. (1990). Linkage Relationships of Human Arginine Vasopressin-Neurophysin-II and Oxytocin-Neurophysin-I to Prodynorphin and Other Loci on Chromosome 20. *Mol Endocrinol*, 4, pp.947-950.
- TABAK, B. A., MCCULLOUGH, M. E., CARVER, C. S., PEDERSEN, E. J. & CUCCARO, M. L. (2014). Variation in oxytocin receptor gene (OXTR) polymorphisms is associated with emotional and behavioral reactions to betrayal. *Soc Cogn Affect Neurosci*, 9, pp.810-6.
- TABAK, B. A., VRSHEK-SCHALLHORN, S., ZINBARG, R. E., PRENOVEAU, J. M., MINEKA, S., REDEI, E. E., ADAM, E. K. & CRASKE, M. G. (2016). Interaction of CD38 Variant and Chronic Interpersonal Stress Prospectively Predicts Social Anxiety and Depression Symptoms Over Six Years. *Clin Psychol Sci*, 4, pp.17-27.
- TAKAHASHI, H., KOEDA, M., ODA, K., MATSUDA, T., MATSUSHIMA, E., MATSUURA, M., ASAI, K. & OKUBO, Y. (2004). An fMRI study of differential neural response to affective pictures in schizophrenia. *Neuroimage*, 22, pp.1247-54.
- TAYLOR, S. F., KANG, J., BREGE, I. S., TSO, I. F., HOSANAGAR, A. & JOHNSON, T. D. (2012). Meta-analysis of functional neuroimaging studies of emotion perception and experience in schizophrenia. *Biol Psychiatry*, 71, pp.136-45.
- TELTSH, O., KANYAS-SARNER, K., RIGBI, A., GREENBAUM, L., LERER, B. & KOHN, Y. (2012). Oxytocin and vasopressin genes are significantly associated with schizophrenia in a large Arab-Israeli pedigree. *Int J Neuropsychopharmacol*, 15, pp.309-19.
- TERRACCIANO, A., SANNA, S., UDA, M., DEIANA, B., USALA, G., BUSONERO, F., MASCHIO, A., SCALLY, M., PATRICIU, N., CHEN, W. M., DISTEL, M. A., SLAGBOOM, E. P., BOOMSMA, D. I., VILLAFUERTE, S., SLIWERSKA, E., BURMEISTER, M., AMIN, N., JANSSENS, A. C., VAN DUIJN, C. M., SCHLESSINGER, D., ABECASIS, G. R. & COSTA, P. T., JR. (2010). Genome-wide association scan for five major dimensions of personality. *Mol Psychiatry*, 15, pp.647-56.
- TESLI, M., ESPESETH, T., BETTELLA, F., MATTINGSDAL, M., AAS, M., MELLE, I., DJUROVIC, S. & ANDREASSEN, O. A. (2014). Polygenic risk score and the psychosis continuum model. *Acta Psychiatr Scand*, 130, pp.311-7.
- TESLI, M., KAUPPI, K., BETTELLA, F., BRANDT, C. L., KAUFMANN, T., ESPESETH, T., MATTINGSDAL, M., AGARTZ, I., MELLE, I., DJUROVIC, S., WESTLYE, L. T. & ANDREASSEN, O. A. (2015). Altered Brain Activation during Emotional Face Processing in Relation to Both Diagnosis and Polygenic Risk of Bipolar Disorder. *PLoS One* [Online], 10, pp.e0134202. DOI: 10.1371/journal.pone.0134202.
- TOST, H., KOLACHANA, B., HAKIMI, S., LEMAITRE, H., VERCHINSKI, B. A., MATTAY, V. S., WEINBERGER, D. R. & MEYER-LINDENBERG, A. (2010). A common allele in the oxytocin receptor gene (OXTR) impacts prosocial temperament and human hypothalamic-limbic structure and function. *Proc Natl Acad Sci U S A*, 107, pp.13936-41.

- TOST, H., KOLACHANA, B., VERCHINSKI, B. A., BILEK, E., GOLDMAN, A. L., MATTAY, V. S., WEINBERGER, D. R. & MEYER-LINDENBERG, A. (2011). Neurogenetic effects of OXTR rs2254298 in the extended limbic system of healthy Caucasian adults. *Biol Psychiatry* [Online], 70, pp.e37-9; author reply e41-2. DOI: 10.1016/j.biopsych.2011.06.034.
- TOTTENHAM, N., TANAKA, J. W., LEON, A. C., MCCARRY, T., NURSE, M., HARE, T. A., MARCUS, D. J., WESTERLUND, A., CASEY, B. J. & NELSON, C. (2009). The NimStim set of facial expressions: judgments from untrained research participants. *Psychiatry Res*, 168, pp.242-9.
- UVNAS-MOBERG, K. (1998). Oxytocin may mediate the benefits of positive social interaction and emotions. *Psychoneuroendocrinology*, 23, pp.819-35.
- VAN ERP, T. G., HIBAR, D. P., RASMUSSEN, J. M., GLAHN, D. C., PEARLSON, G. D., ANDREASSEN, O. A., AGARTZ, I., WESTLYE, L. T., HAUKVIK, U. K., DALE, A. M., MELLE, I., HARTBERG, C. B., GRUBER, O., KRAEMER, B., ZILLES, D., DONOHUE, G., KELLY, S., MCDONALD, C., MORRIS, D. W., CANNON, D. M., CORVIN, A., MACHIELSEN, M. W., KOENDERS, L., DE HAAN, L., VELTMAN, D. J., SATTERTHWAITE, T. D., WOLF, D. H., GUR, R. C., GUR, R. E., POTKIN, S. G., MATHALON, D. H., MUELLER, B. A., PREDA, A., MACCIARDI, F., EHRLICH, S., WALTON, E., HASS, J., CALHOUN, V. D., BOCKHOLT, H. J., SPONHEIM, S. R., SHOEMAKER, J. M., VAN HAREN, N. E., POL, H. E., OPHOFF, R. A., KAHN, R. S., ROIZ-SANTIANEZ, R., CRESPO-FACORRO, B., WANG, L., ALPERT, K. I., JONSSON, E. G., DIMITROVA, R., BOIS, C., WHALLEY, H. C., MCINTOSH, A. M., LAWRIE, S. M., HASHIMOTO, R., THOMPSON, P. M. & TURNER, J. A. (2016). Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Mol Psychiatry*, 21, pp.585.
- VAN OS, J. & JONES, P. B. (2001). Neuroticism as a risk factor for schizophrenia. *Psychol Med*, 31, pp.1129-34.
- VAN OS, J., KENIS, G. & RUTTEN, B. P. F. (2010). The environment and schizophrenia. *Nature*, 468, pp.203-212.
- VARESE, F., SMEETS, F., DRUKKER, M., LIEVERSE, R., LATASTER, T., VIECHTBAUER, W., READ, J., VAN OS, J. & BENTALL, R. P. (2012). Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective- and cross-sectional cohort studies. *Schizophr Bull*, 38, pp.661-71.
- VENTURA, J., LIBERMAN, R. P., GREEN, M. F., SHANER, A. & MINTZ, J. (1998). Training and quality assurance with the Structured Clinical Interview for DSM-IV (SCID-I/P). *Psychiatry Res*, 79, pp.163-73.
- VENTURA, J., WOOD, R. C., JIMENEZ, A. M. & HELLEMANN, G. S. (2013). Neurocognition and symptoms identify links between facial recognition and emotion processing in schizophrenia: meta-analytic findings. *Schizophr Res*, 151, pp.78-84.
- VUKASOVIC, T. & BRATKO, D. (2015). Heritability of personality: A meta-analysis of behavior genetic studies. *Psychol Bull*, 141, pp.769-85.
- WALLIS, M. (2012). Molecular evolution of the neurohypophysial hormone precursors in mammals: Comparative genomics reveals novel mammalian oxytocin and vasopressin analogues. *Gen Comp Endocrinol*, 179, pp.313-8.
- WALUM, H., LICHTENSTEIN, P., NEIDERHISER, J. M., REISS, D., GANIBAN, J. M., SPOTTS, E. L., PEDERSEN, N. L., ANCKARSATER, H., LARSSON, H. & WESTBERG, L. (2012). Variation in the oxytocin receptor gene is associated with pair-bonding and social behavior. *Biol Psychiatry*, 71, pp.419-26.
- WALUM, H., WALDMAN, I. D. & YOUNG, L. J. (2016). Statistical and Methodological Considerations for the Interpretation of Intranasal Oxytocin Studies. *Biol Psychiatry*, 79, pp.251-7.
- WANG, J., QIN, W., LIU, B., ZHOU, Y., WANG, D., ZHANG, Y., JIANG, T. & YU, C. (2014). Neural mechanisms of oxytocin receptor gene mediating anxiety-related temperament. *Brain Struct Funct*, 219, pp.1543-54.

- WATANABE, Y., KANEKO, N., NUNOKAWA, A., SHIBUYA, M., EGAWA, J. & SOMEYA, T. (2012). Oxytocin receptor (OXTR) gene and risk of schizophrenia: case-control and family-based analyses and meta-analysis in a Japanese population. *Psychiatry Clin Neurosci*, 66, pp.622.
- WIJERATNE, C., SACHDEV, S., WEN, W., PIGUET, O., LIPNICKI, D. M., MALHI, G. S., MITCHELL, P. B. & SACHDEV, P. S. (2013). Hippocampal and amygdala volumes in an older bipolar disorder sample. *Int Psychogeriatr*, 25, pp.54-60.
- WU, S., JIA, M., RUAN, Y., LIU, J., GUO, Y., SHUANG, M., GONG, X., ZHANG, Y., YANG, X. & ZHANG, D. (2005). Positive association of the oxytocin receptor gene (OXTR) with autism in the Chinese Han population. *Biol Psychiatry*, 58, pp.74-7.
- YOUNG, K. A., LIU, Y., GOBROGGE, K. L., WANG, H. & WANG, Z. (2014). Oxytocin reverses amphetamine-induced deficits in social bonding: evidence for an interaction with nucleus accumbens dopamine. *J Neurosci*, 34, pp.8499-506.
- YUEN, K. W., GARNER, J. P., CARSON, D. S., KELLER, J., LEMBKE, A., HYDE, S. A., KENNA, H. A., TENNAKON, L., SCHATZBERG, A. F. & PARKER, K. J. (2014). Plasma oxytocin concentrations are lower in depressed vs. healthy control women and are independent of cortisol. *J Psychiatr Res*, 51, pp.30-36.
- ZALD, D. H. (2003). The human amygdala and the emotional evaluation of sensory stimuli. *Brain Res Brain Res Rev*, 41, pp.88-123.
- ZHANG, J. P. & MALHOTRA, A. K. (2011). Pharmacogenetics and antipsychotics: therapeutic efficacy and side effects prediction. *Expert Opin Drug Metab Toxicol*, 7, pp.9-37.
- ZONDERVAN, K. T. & CARDON, L. R. (2007). Designing candidate gene and genome-wide case-control association studies. *Nat. Protocols*, 2, pp.2492-2501.

Appendix

Appendix 1. Regional linkage disequilibrium plots of rs53576 and rs237902.

Abbreviations: cM = centiMorgan, Mb = Megabases, kb = kilobases

Errata

Study I, II and III:

Section Methods, Genotyping: “Samples were excluded through quality control if they had low-yield (call rate below 97%)” should be “Samples were excluded through quality control if they had low-yield (call rate below 95%)”. “Markers with.....minor allele frequencies below 0.05 were excluded” should be “Markers with.....minor allele frequencies below 0.01 were excluded”. The correct numbers are reported in the thesis.

Study II:

Page 2, section Sample Characteristics and Table 1: 734 patients with a psychotic disorder and 417 healthy individuals were included (not 420). All had both genotyped/imputation data as well as schizophrenia PGRS. 725 individuals in the patient group had available symptom scores of p6 measured by the PANSS, while 724 patients had available symptom scores of p7, n2 and n4. The correct numbers are reported in the thesis.