Drug Discovery, Design and Synthesis of
Polycyclic Regulators of Cancer Cells and Nuclear Receptors:
Phenazine 5,10-dioxides and Oxysterols

Dissertation for the degree of *Philosophiae Doctor*

By

Elvar Örn Viktorsson

Faculty of Mathematics and Natural Sciences

University of Oslo

2017

© Elvar Örn Viktorsson, 2017

*Series of dissertations submitted to the
Faculty of Mathematics and Natural Sciences, University of Oslo
No. 1893*

ISSN 1501-7710

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

To my father, Viktor Daði Bóasson (24.08.1952-19.12.2012†)

Acknowledgements

The work presented in this thesis was carried out between autumn 2013 and spring 2017 at the Department of Pharmaceutical Chemistry, School of Pharmacy, University of Oslo. Liver X receptors were the primary focus of attention from autumn 2013 until spring 2014 and thereafter, the scope was turned to phenazines until spring 2017. A six-week research period was undertaken in summer 2015 at University Claude Bernard Lyon 1, France.

As the end of this four-year journey at the UiO's School of Pharmacy seems to be drawing near, it is only appropriate to thank those who have contributed to this work in many ways.

First of all, I would like to express my deepest gratitude to Prof. Pål Rongved, my main supervisor. First and foremost, for giving me the opportunity to take on this challenge and for placing your trust in me. All the exciting talks about chemistry, biology, guitar gear and life in general have been a great pleasure so far, and I hope we can continue to collaborate in the unseen future.

A special gratitude to my good friend and colleague, Dr. Alexander H. Åstrand for all his input and all the good times. You are the one who helped me the most in the beginning, and for that I am grateful.

I would like acknowledge the talented biologists at the University of Bergen. Prof. Lars Herfindal, Dr. Reidun Æsøy and Prof. Stein Ove Døskeland, for all their contribution to the phenazine part.

Dr. Eili Tranheim Kase as well for the supervision and design of all biological evaluations performed in the Liver X receptor project, along with helpful discussions on LXR-related biology.

Assoc. Prof. Tore Bonge-Hansen from the Department of Chemistry and Bendik Grøthe also deserve large credit for their important contributions to the phenazine project.

Professor II Ørjan Samuelsen and Christopher Fröhlich are acknowledged for antimicrobial evaluations in Tromsø.

I also wish to thank some other current and former colleagues and staff at the University of Oslo's Department of Pharmaceutical chemistry, all of which have contributed to such a

motivating and fun place to work at: Christian, Geir, Renate, Karoline, Åsmund, Marius, Jørn, Anthony, Eirik, Vegard, Lisa, Anders, Jens, Trond, Bora, Martin H. and Peter. “Legemiddelkjemi Blastoff!”

The department engineers, Iuliana Johansen and Anne Bjerke are also thanked for general lab support and Halvor Aandal as well for his important help with formalities and shipment of samples.

The master degree candidates I’ve had the pleasure of supervising and work with; Sindre Støa and Rasha Haseeb.

This work could probably not have been finished without funding from the following sources: A NOVO exploratory Pre Seed Grant no. 4462, a BIOTEK2021 program grant from the Norwegian research council, a grant from University of Oslo’s innovation fund in 2015, Norwegian PhD School of Pharmacy grant 2015 and last but not least, the School of pharmacy, University of Oslo for employing me.

It also seems impossible not to mention Prof. Frode Rise and Dirk Petersen for maintaining an outstanding and an impressive NMR-facility. Prof. Carl Henrik Görbitz and Dr. Sigurd Øien-Ødegaard are also acknowledged for the X-ray crystallographic work related to this thesis.

The people who helped me proof read this thesis deserve a special mention: Pål Rongved, Alexander Åstrand, Åsmund Kaupang, Eili Kase, Christian Schnaars, Geir Kildahl-Andersen, and Þorsteinn Viktorsson. Thank you all.

Outermost gratitude to Guðrún Þorsteinsdóttir and Eva Viktorsdóttir for all the support and encouragement throughout my studies.

And finally, my better half, Sigríður Ólafsdóttir. Thank you for always supporting me, and for moving with me to Oslo. Your care, love and patience have made this dream of mine come true.

Oslo, June 25th, 2017

Elvar Örn Viktorsson

Abstract

The work presented in this thesis employed conventional structure-activity relationship studies to guide the rational design of compounds that target acute myeloid leukemia, infections caused by resistant bacteria and metabolic diseases such as atherosclerosis, diabetes mellitus type 2 and obesity.

Recent literature concludes that the phenazine 5,10-dioxide natural product iodinin (**5**) targets human leukemic cells potently and selectively. In this study, we synthesized iodinin from commercially available reagents and used structure-activity relationship studies as a tool to rationally guide the design and discovery of new antileukemic agents. In this process, the known antibiotic myxin (**8**) was identified to be equally potent to- and as selective towards leukemic cells as iodinin. This finding enabled us to conclude that one of iodinin's two phenol groups can be *O*-functionalized without any loss of cytotoxic potency at the same time the selectivity towards AML cells over healthy cells is preserved (Paper I). Furthermore, synthesis and biological evaluation of iodinin- and myxin prodrugs revealed carbamate side chains to be an advantageous functionality to attach to the phenolic function. These prodrugs displayed enhanced potency, solubility and more efficient cell membrane penetration compared to iodinin (Paper II). By altering the substitution patterns of the phenazine scaffold in positions 6, 7 and 8, we could identify 7,8-dihalogenated- and 7,8-dimethylated carbamate-functionalized phenazine 5,10-dioxides with considerably enhanced potency, compared to that of iodinin and myxin, while preserving the selectivity towards leukemic cells.

Phenazine 5,10-dioxides were also tested against strains of both Gram-positive and -negative bacteria with different profiles of resistance to antibiotics used in the clinic. Results retrieved from these experiments revealed clear trends that can guide further development of phenazine 5,10-dioxides for use in the treatment of severe infections.

We were also interested in targeting the transcriptional machinery of cells through the liver X nuclear receptors (LXRs). The LXRs play a crucial role in the metabolism and homeostasis of cholesterol, lipids, bile acids and carbohydrates. Therefore, their potential to serve as drug targets in the context of metabolic diseases is immense.

Individual reports state that the natural oxysterol 22(*R*)-hydroxycholesterol (**22RHC**) and its unnatural synthetic epimer 22(*S*)-hydroxycholesterol (**22SHC**) can promote opposite

biological effects in terms of LXR-regulated gene expression. Through virtual screening, synthesis of analogues and biological evaluations that assessed the expression of key LXR target genes, we managed to identify two new leads that mimicked the activities of **22SHC**. One compound (**LXR11**) was found to be very promising *in vitro* and was investigated further in rats (Paper **III**). This compound did however fail to provide the desired outcome *in vivo*, most likely due to pharmacokinetic problems, such as rapid metabolism in the rat's gastrointestinal tract. SAR studies were also conducted *in vitro*, and involved the synthetic preparation of analogues with different functionalities in position 22 of the cholesterol scaffold. The results from this study gave an enhanced understanding of how LXR-regulated gene expression can be manipulated by synthetic transformations in position 22 of the cholesterol scaffold – the compound 22-ketocholesterol (**22KCH**) was found to upregulate the expression of *ABCA1* selectively, rendering it a promising new lead compound for further anti-atherosclerotic evaluations (Paper **IV**).

List of publications and patents

Paper I

Total synthesis and antileukemic evaluations of the phenazine 5,10-dioxide natural products iodinin, myxin and their derivatives

Viktorsson, E. Ö.; Grøthe, B. M.; Aesoy, R.; Sabir, M.; Snellingen, S.; Prandina, A.; Åstrand, O. A. H.; Bonge-Hansen, T.; Døskeland, S. O.; Herfindal, L. and Rongved, P.

Bioorg. Med. Chem. **2017**, 25 (7): 2285-2293.

Paper II

New prodrugs and analogues of the phenazine 5,10-dioxide natural products iodinin and myxin promote selective cytotoxicity towards human acute myeloid leukemia cells.

Viktorsson, E. Ö.; Aesoy, R.; Støa, S.; Døskeland, S. O.; Herfindal, L. and Rongved, P.

Prepared manuscript

Paper III

Synthesis, *in vitro* and *in vivo* biological evaluation of new oxysterols as modulators of the liver X receptors

Åstrand, O. A. H.; Viktorsson, E. Ö.; Kristensen, A. L.; Ekeberg, D.; Røberg-Larsen, H.; Wilson, S. R.; Gabrielsen, M.; Sylte, I.; Rustan, A.; Thoresen, H.; Rongved, P.; Kase, E. T.

J. Steroid. Biochem. **2017**, 165, 323-330.

Paper IV

Regulation of liver X receptor target genes by 22-functionalized oxysterols. Synthesis, *in silico* and *in vitro* evaluations

Viktorsson, E. Ö.; Gabrielsen, M.; Kumarachandran, N.; Sylte, I.; Rongved, P.; Åstrand, O. A. H.; Kase, E.

Steroids **2017**, 118, 119-127.

Paper V

Crystal structure of (S)-2-[(3S,8S,9S,10R,13S,14S,17R)-3-hydroxy-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-17-yl]-N-methoxy-N-methylpropanamide (Fernholz Weinreb amide)

Viktorsson, E. Ö.; Åstrand, O. A. H.; Haseeb, R. S.; Görbitz, C. H. and Rongved, P.

Acta. Crystallogr. E. **2015**, 71 (3): 275-277

Patent application 1

Preparation of phenazine derivatives and cyclodextrin nanoparticle delivery thereof for treatment of cancer and infection.

Døskeland, S. O.; Rongved, P.; Herfindal, L.; Le Borgne, M.; Viktorsson, E. Ö. and Åstrand, O. A. H.

International publication number: WO 2015/063516 A2 2015.05.07.

Graphical abstracts

Paper I

Total synthesis and antileukemic evaluations of the phenazine 5,10-dioxide natural products iodinin, myxin and their derivatives (a front cover of Bioorganic and Medicinal Chemistry, volume 25, issue 7)

Paper II

New prodrugs and analogues of the phenazine 5,10-dioxide natural products iodinin and myxin promote selective cytotoxicity towards human acute myeloid leukemia cells

Paper III

Synthesis, *in vitro* and *in vivo* biological evaluation of new oxysterols as modulators of the liver X receptors

Paper IV

Regulation of liver X receptor target genes by 22-functionalized oxysterols. Synthesis, *in silico* and *in vitro* evaluations

List of abbreviations

1-HP	1-Hydroxyphenazine
22KC	22-Ketocholesterol
22RHC	22(<i>R</i>)-Hydroxycholesterol
22SHC	22(<i>S</i>)-Hydroxycholesterol
ABC	ATP Binding cassettes (ATP : Adenosine triphosphate)
ADIC	2-amino-2-desoxyisochorismic acid
ADME	Absorption, distribution, metabolism and elimination
AML	Acute myeloid leukemia
AMR	Antimicrobial resistance
AOCHC	6-amino-5-oxocyclohex-2-ene-1-carboxylic acid
APL	Acute promyelocytic leukemia
CR	Complete remission
DHHA	<i>Trans</i> -2,3-dihydro-3-hydroxyanthranilic acid
DHMCa	<i>N,N</i> -dimethyl-3 β -hydroxy-choleamide
DHPDC	Dehydrophenazine 1,6-dicarboxylic acid
DHPH _z	Dehydrophenazine
DIPEA	Diisopropyl ethyl amine
DMAP	4-(Dimethylamino)pyridine
DMP	Dess-Martin periodinane
Eq	Molar equivalent
FASN	Fatty acid synthase
FDA	U.S. Food and Drug Administration
GSH	Glutathione
HATU	1-[Bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5- <i>b</i>]pyridinium 3-oxide PF ₆ ⁻
HBTU	<i>O</i> -(Benzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate
HHPDC	Hexahydrophenazine-1,6-dicarboxylic acid
HOBt	1-Hydroxybenzotriazole
HTS	High-throughput screening
IPR	Intellectual property right
LDA	Lithiumdiisopropylamide
LXR	Liver X receptor
<i>m</i> CBA	<i>meta</i> -chlorobenzoic acid
<i>m</i> CPBA	<i>meta</i> -chloroperbenzoic acid
NADPH	Nicotineamide adenine dinucleotide phosphate
NAS	Nucleophilic aromatic substitution
NBS	<i>N</i> -Bromosuccinimide
NCS	<i>N</i> -chlorosuccinimide
NIS	<i>N</i> -Iodosuccinimide
NR	Nuclear receptor
PCA	Phenazine 1-carboxylic acid
PDC	Phenazine 1,6-dicarboxylic acid
PH _z	Phenazine

PYO	Pyocyanin
RCT	Reverse cholesterol transport
ROS	Reactive oxygen species
SCD	Stearoyl CoA reductase
SREBP-1	Sterol regulatory element-binding protein 1
T2DM	Diabetes mellitus type 2
TAG	Triacylglycerol
TBAF	Tetrabutylammonium fluoride
TBS	<i>tert</i> -butyldimethylsilyl
THPCA	Tetrahydrophenazine-1,6-dicarboxylic acid
TPZ	Tirapazamine

Contents

Acknowledgements	III
Abstract.....	V
List of publications and patents.....	VII
Graphical abstracts.....	IX
List of abbreviations	XI
Contents	XIII
Chapter 1 – General introduction	1
1.1 Drug discovery.....	1
1.2 Natural products	4
1.3 Medicinal Chemistry.....	6
1.4 Cancer	7
1.4.1 Tumor hypoxia	9
1.4.2 Leukemia.....	9
1.5 References	12
Chapter 2 – Overall aims of study	15
2.1 References	16
Chapter 3 – Introduction to phenazines	17
3.1 Biosynthesis	20
3.2 Biological functions and modes of action.....	22
3.2.1 Redox cycling of natural phenazines	22
3.2.2 Lomofungin, polynucleotide interaction and metal chelation.....	24
3.2.3 Iodinin	25
3.2.4 Myxin	27
3.2.5 Phenazine 5,10-dioxides, tirapazamine and cancer.....	28
3.3 Synthesis of phenazines.....	32
3.3.1 The Wohl-Aue condensation.....	32
3.3.2 The Bamberger-Ham reaction.....	33
3.3.3 The Beirut-reaction	34
3.3.4 Condensations of 1,2-benzoquinones and O-phenylenediamines	35
3.3.5 Cyclizations of O-nitrodiphenylamines	36
3.3.6 Buchwald-Hartwig C-N cross-coupling.....	37
3.3.7 Miscellaneous modern methods	39
3.4 References	40
Chapter 4 – Phenazines and aim of study	45
Chapter 5 – Phenazines results and discussion	47
5.1 Paper I – Synthesis and biological evaluations of iodinin, myxin and alkylated derivatives...47	
5.1.1 Synthesis	47
5.1.2 Biological evaluations of iodinin, myxin and alkylated analogs.....	52
5.2 Paper II – Prodrug development and further SAR studies on iodinin and myxin.....	55
5.2.1 Functionalization of iodinin and myxin phenols.....	55
5.2.2 Biological evaluations of iodinin and myxin prodrugs	58
5.2.3 Synthesis of 1-hydroxyphenazine 5,10-dioxide and analogues	59
5.2.4 Biological evaluations of 1-hydroxyphenazine 5,10-dioxide analogues.....	61
5.2.5 Synthesis of 7,8-disubstituted analogues of 1-hydroxyphenazine 5,10-dioxide and quinoxaline 1,4-dioxides	63

5.2.6	Biological evaluations of 7,8-disubstituted phenazines and quinoxaline 1,4-dioxides...	66
5.3	Synthesis of miscellaneous compounds and antimicrobial evaluations	69
5.3.1	Synthesis of miscellaneous phenazine 5,10-dioxides.....	69
5.3.2	Antimicrobial evaluations	71
5.4	Conclusions and summary of structure-activity relationships	73
5.5	Future perspectives.....	76
5.5.1	Synthesis	76
5.5.2	Biology	78
5.6	References	79
Chapter 6	- Introduction to liver X receptors	81
6.1	Nuclear receptors.....	81
6.2	Liver X receptors and biological functions.....	83
6.2.1	Target genes and therapeutic potentials.	84
6.2.2	Ligands.....	87
Chapter 7	- Oxysterols and aim of study	91
Chapter 8	- Oxysterols results and discussion.....	93
8.1	Papers III-V	93
8.1.1	Synthesis and molecular modelling.....	93
8.1.2	Biological evaluations – paper III	98
8.1.3	Biological evaluations – paper IV	100
8.2	Future perspectives.....	103
8.3	Conclusions	104
8.4	References	105
Chapter 9	- Concluding remarks	111
Appendixes		115
Appendix A - Experimental procedures and characterization data from section 5.3.1		
Appendix B - Paper I and supporting information		
Appendix C - Paper II and supporting information		
Appendix D - Paper III and supporting information		
Appendix E - Paper IV and supporting information		
Appendix F - Paper V and supporting information		

Chapter 1 – General introduction

1.1 Drug discovery

“The universe is hardwired to be an organic chemist. It’s not a very clean or tidy one, but it has really big beakers and plenty of time.” – Scott Sandford, NASA

Drug discovery is devoted to the identification of new substances, so-called lead compounds (or hits) that exhibit attractive biological features in terms of treating an illness by affecting a specific biological target of interest. Modern drug discovery is a complex, time-consuming and multidisciplinary process. Moreover, it is enormously expensive and its success rates are extremely low. New drug molecules are seldom discovered directly. Therefore, lead compounds are substances that have the potential to be further developed via medicinal chemistry to achieve the end goal; a marketed drug that provides a positive impact on human health. Typically, the biological targets of interest are proteins that according to our current understanding and knowledge play a central role in a disease of interest.¹

There are several strategies to approach the discovery of a new lead compound. In the 1940s, following Alexander Fleming’s discovery of benzylpenicillin (penicillin G),² Selman Waksman introduced a screening platform in the 1940s that suddenly drew a line between an active discovery platform and discoveries that were simply based on luck.³ The concept was successful for a time, based on the screening of soil-derived *Streptomyces* for antimicrobial activity. This approach was incorporated into the pharmaceutical industry and led to the discoveries of streptomycin and the first aminoglycoside. However, this particular platform was abandoned after a while as the same compounds were continuously rediscovered again and again.⁴

Nowadays, the pharmaceutical industry is fond of high-throughput methods, e.g. high-throughput synthesis and high-throughput screening (HTS). The first step of the drug discovery process applying the HTS method is target identification and validation. Technological advances in molecular biology have enabled cloning and expression of protein targets to become a routine practice in the pharmaceutical industry. Consequently, compounds can be evaluated and even optimized *in vitro* before they are exposed to animals.⁵ After target validation is established, the screening process is initiated, and once a HTS-hit is identified,

the chance of further developing this specific molecule to a marketed drug-brand is somewhere around one in a million.⁶ To conduct these types of screenings, such facilities are in possession of chemical libraries counting typically well over one million compounds, and the screening capacity is truly enormous; several million compounds can be screened in a matter of a few weeks against a given biological target. Despite these and many other technological advances, the industry has simply failed to increase its output of marketed drugs per year.⁷⁻⁸

The increased efficiency of a drug discovery process based on HTS has thus been a topic for debate and still, no obvious solutions have been provided. However, a large focus has been turned towards the design of the chemical libraries screened. Hypothetically, there are some disadvantages to using synthetic libraries, as their chemical diversity is simply not large enough. This can be exploited though, in some cases, if structural knowledge regarding the drug target is already in hand.⁹ Scientists at Merck pursued a rational approach, choosing the 2-aryl-indole chemical scaffold as a starting point for constructing a G protein-coupled receptor (GPCR)-targeted library through combinatorial synthesis. In this project alone, a total of 128.000 analogues were prepared and further screened for various GPCR targets. Multiple compounds were identified to display potent activities across a variety of different receptors.¹⁰ This is rational drug design and targeted screening. Some of these compounds were optimized via medicinal chemistry and this eventually led to the identification of a potent clinical candidate – an antagonist of the human neurokinin-1 receptor (hNK₁).¹¹⁻¹²

Murcko et al. investigated the chemical diversity of 5120 drugs from the Comprehensive Medicinal Chemistry database.¹³ Simplifying these chemical structures down to skeletons of carbon frameworks, 32 of the most common scaffolds accounted for approximately half of all the drugs that were investigated.¹³ Although these findings are built on simplified chemical structures, the chemical diversity seems surprisingly low among marketed drugs.⁹ Evans et al. introduced the concept of “privileged structures” in 1986,¹⁴ as it was recognized that certain chemical scaffolds have the potential to interact with diverse biological targets upon modification of their functional groups.⁹ The 2-arylidole scaffold earlier mentioned is a good example of such a structure.

High-throughput screening of several hundred thousand compounds can afford multiple hits. This calls for the necessity to prioritize the most promising ones for further investigations and to ensure the hit was not a false-positive. This selection is done by medicinal chemists evaluating “druglikeness” in terms of privileged scaffolds. Lipinski and colleagues at Pfizer

proposed a simple rule of thumb to evaluate druglikeness, referred to as the Lipinski's Rule of Five. This set of four (not five) guidelines was found to account for 90% of all orally active drugs that have reached phase II clinical status.¹⁵

- $M_w \leq 500$ Daltons
- Hydrogen bond donors ≤ 5
- Hydrogen bond acceptors ≤ 10
- $\log P \leq 5$

Due to an ever-increasing pressure on the industry to deliver new types of medicine in particular due to enormous research costs and lowered product output, the focus has also been turned towards the overall quality of discovered lead compounds. As a result, the term "leadlike" has also been coined as the "Rule of Three":¹⁵

- $M_w \leq 300$ Daltons
- H-bond donors and acceptors ≤ 3
- $\log P \leq 3$
- Number of rotatable bonds ≤ 3

This set of guidelines is stricter than those proposed by the Lipinski's Rule of Five, but this stems from the fact that the development process of a lead compound via medicinal chemistry is most certainly followed by increase in M_w and $\log P$.^{6, 15}

Use of computational power in drug design has gained popularity as technological advances of computers have emerged over the course of the past three decades. Unlike the brute-force HTS-method which requires minimum knowledge about the interactions between the target and the ligand, virtual screenings can also be performed *in silico* if detailed knowledge of the target is already available. This is computer-aided drug discovery (CADD).¹⁶ Rule-based filters that sort out compounds with undesirable properties from the chemical library can also be employed to reduce the number of compounds that are relevant for conventional HTS (e.g. an inappropriate molecular size or $\log P$).¹⁷ A huge advantage of CADD is that neither the actual proteins, nor the ligands need actually to be in hand for conducting virtual screenings of ligand interactions with biological targets.

Other approaches have proven successful in drug discovery and particularly the evaluation of drug metabolites. A dye called Prontosil rubrum was investigated by Gerhard Domagk at Bayer in the 1930s. This compound was shown to be active against streptococcal infections in mice. Interestingly, this red dye was only active *in vivo*, not *in vitro*. Prontosil was later shown to be metabolized by liver enzymes to the active ingredient sulfanilamide which proved useful in treating syphilis.¹⁸ Other classic examples include morphine, a metabolite of codeine,¹⁹ and oxazepam (Sobril®), an oxidized metabolite of diazepam (Valium®).²⁰

Clinical observations can also result in new drugs for unforeseen indications. Examples include sildenafil which was first evaluated for its antihypertensive effects, but later showed an interesting side effect and was developed into a drug against male erectile dysfunction.²¹ Another example is minoxidil, also first clinically evaluated to treat high blood pressure, but revealed properties as a drug to prevent hair loss.²² Obviously, the disadvantage of this approach is that the molecules must have gone through expensive and time consuming pre-clinical studies before entering clinical trials in humans. On a positive note, these compounds have in addition passed expensive pre-clinical toxicity studies in animals.

1.2 Natural products

Nature has been the most prominent source of medicinal products since the beginning of recorded history and still continues to have a great impact on drug discovery and -development.²³ Records from Mesopotamia (~2600 BC) explain the medicinal use of over one thousand different plant-derived crude materials, and the “Ebers papyrus” from ancient Egypt (~1500 BC) contains quite detailed descriptions of more than 700 drugs and their use.²⁴

Looking at marketed anticancer agents alone, from the 1940s to 2014, in a detailed analysis by Newman and Cragg, 49% were found to be natural products or their close derivatives.²⁵ Figure 1 (next page) provides a selected overview over unaltered natural products that have served humans as drugs. Terrestrial microorganisms have particularly been a fruitful source of compounds, giving us drug prototypes for important classes of antibiotics (penicillins, cephalosporins and tetracyclines), cholesterol-lowering drugs (statins) and anti-cancer agents like daunorubicin (structure shown later in page 11).¹⁸

Figure 1: Natural products that have been used as drugs.

Natural products are in many ways privileged structures since they are produced by living organisms and have been selected through time and evolution to occupy three-dimensional spaces provided by nature. These cavities arrive in the forms of receptors, enzymes and in the metabolic machineries of living cells.^{12, 26} A report from Novartis showed that screening of library subsets counting only natural products resulted in significantly higher hit-rates compared to conventional synthetic- or combinatorial libraries.²⁷ Thus, screening libraries should be assessed rather by quality than quantity when it comes to the number of structures in the discovery process.²⁶ Hypothetically speaking, the chances of a unique and chemically complex structure like paclitaxel (Taxol®, Figure 1) being discovered from a library only consisting exclusively of synthetic compounds seems literally impossible. Nature is a vast source that can provide the type of structural diversity needed to drive revolutionary drug discoveries, as well as challenge organic chemists with new difficult synthetic targets.

Most of the time, unaltered natural products are simply not fit to serve as drugs in humans or pass through today's strict regulatory protocols. They require specific chemical tailoring to tune in properties such as potency, target affinity, pharmacokinetics, and to eliminate or reduce others, such as toxicity and unwanted side effects. This is where the practice of medicinal chemistry comes in.

1.3 Medicinal Chemistry

Medicinal chemistry is engaged in the early stages of drug discovery and is devoted to the selection and optimization of new lead compounds. As a part of the lead optimization process, synthetic alterations to the molecular structure of a lead compound are usually carried out systematically to map structure-activity relationships (SAR). SAR-studies aim to provide information about which part of the molecule is essential to retain bioactivity (i. e. its pharmacophore), and what parts of the structure can be substituted chemically or removed. Quantitative Structure-activity relationships (QSARs) are the mathematical relationships determined between the structural properties of a molecule and another feature, such as specific activity in a biological assay or a physicochemical property. Relationships determined from QSAR experiments can be of aid to either optimize a lead compound or to guide the rational design of new analogues.²⁸

Medicinal chemistry has the purpose of making the drug safer and more effective for therapeutic use in humans, in other words more druglike. Chemical transformations of the lead candidate can influence important factors such as potency, toxicity, stability, water solubility and target selectivity in addition to absorption, distribution, metabolism and elimination (ADME). The chemical structures of many natural products are also simply too complex to be synthesized efficiently on an industrial scale. In this context, simplifications of the molecular structure or semi-synthetic strategies often prove to be necessary. Figure 2 shows the core structure and pharmacophore of the quinolone class of antibiotics, discovered in 1980s – a late-coming end to the golden era of antibiotic drug discovery.

Figure 2: The core structure of quinolones exemplified by ciprofloxacin, a synthetic broad-spectrum antimicrobial agent derived from structure-activity relationship studies. Quinolones inhibit bacterial DNA gyrase and topoisomerase IV.¹⁸

Medicinal chemistry is uniquely placed at the interface between several scientific disciplines, heavily relying on organic chemistry and synthesis, pharmacology and the biology of eukaryotic- and prokaryotic cells. Therefore, the modern medicinal chemist is a member of an interdisciplinary research team where he or she is required, not only to thoroughly understand the field of synthetic chemistry, but to gain knowledge of several other related research disciplines. Otherwise, his/her decision making regarding the selection and further development of lead compounds is at high risk of being flawed or inefficient.⁷ Immense technological advances in computational equipment, purification techniques and analytical instrumentation have enabled humans to move forward from consumption of impure crude materials towards the use of well-documented active ingredients of very high purity. The role of the medicinal chemist in modern drug discovery has thus changed rapidly over the course of last decades, much owing to technological advances and changed perspectives in drug discovery. Despite the altered landscapes, the goal of the journey remains intact – to find and develop new and safer drugs to treat threatening diseases.

1.4 Cancer

Cancer is a term for more than 100 heterogeneous diseases of different etiology. All of them appear to have at least three factors in common: Unhindered cell-growth, invasion of adjacent tissues by tumor and finally, tumor dislocation and migration to sites distant from the point of origin. The most common cancers in humans are those associated with the lungs, the prostate, the breasts, and the colorectum.²⁹ The World health organization (WHO) classifies cancer as a leading cause of death worldwide, with over 14 million confirmed new cases and 8.2 million deaths in 2012 alone.³⁰

In the majority of cases, options for cancer treatment include tumor resection, chemotherapy and radiation therapy. These options are often used in combination, depending on the diagnosis and stage of tumor development.³¹ Cancer chemotherapy is in many cases heavily reliant on the use of antibiotics. This results from the immunosuppressive effects promoted by many chemotherapeutic agents. The emergence of antibiotic resistance therefore poses a drastic threat in all cancer therapies.³²

Despite massive efforts in cancer research in the past decades, with build-up of knowledge and complex literature, the mechanisms behind carcinogenesis are still not fully understood. Lines of evidence suggest that the stepwise evolution required for a normal cell to become a

cancerous one is a multistep path driven by dynamic changes in the genome.³³ This process can be divided into phases of *initiation*, *promotion*, *transformation* and *progression*. During the *initiation* phase, normal, healthy cells are exposed to carcinogenic substances. This results in DNA-damages that, if not repaired, lead to an irreversibly mutated cell. The *promotion* phase is reversible and the cell's microenvironment will be exposed to carcinogens or other unknown factors that may favor the growth of the mutated cells over normal ones. At some point, these cells mutate further to become malignant (*transformation*). Further mutations result in a *progression* phase characterized by increased replicative potentials. This phase involves invasion into local tissues and progression of metastasis.²⁹

To provide a joint framework over the complex term of cancer, six hallmarks have been coined which seem to account for the clear majority of cancerous diseases (Figure 3). These hallmarks represent functional properties acquired by tumors and are as follows: Self-sufficiency in growth signals, insensitivity to growth-suppressing signals, resistance to cell death by apoptosis, unlimited replicative potential, angiogenetic and metastatic activity.³³⁻³⁴

Figure 3: The six hallmarks of cancer. A modified figure from Hanahan et al.³⁴

1.4.1 Tumor hypoxia

It is a well-established fact that solid tumors in humans are less oxygenated than the normal tissues surrounding them.³⁵ This characteristic feature is the result of a rapid, “out-of-control” growth and a disorganized vascular network surrounding tumors.³⁶ Most tumors of a volume larger than 1 mm³ contain hypoxic regions located relatively distant from blood vessels. Insufficient flow of oxygen, nutrition and even conventional chemotherapeutic agents contribute to the hypoxic nature of solid tumors.³⁷ These factors may result in the tumors resistance against chemotherapy, as drugs are not able to enter the tumor in sufficient concentrations. Moreover, a prolonged hypoxic state in a tumor tissue may also result in necrosis, another characteristic feature of solid tumors.³⁷

The difference in oxygen saturation can possibly be exploited for selectivity between cancerous cells and normal healthy cells. Several chemical scaffolds are prone to act more potently in hypoxic environments. These scaffolds depend on molecular oxygen for the detoxification (back-oxidation) of radicals formed after an enzymatic reduction. Examples include *N*-oxidized derivatives of phenazines,³⁸ quinoxalines³⁹ and 1,2,4-benzotriazines.⁴⁰ These scaffolds will be more thoroughly addressed in section 3.2.5, page 28.

1.4.2 Leukemia

Millions of cells are formed every minute in the bone marrow and all types of specific blood components evolve from stem cells located therein. Normally, these cells mature and develop in a stepwise manner towards more sophisticated cell types that carry out specific tasks in the normal blood economy.

Leukemia is a group of cancerous diseases that involve cells that normally should develop and differentiate further into more specific blood components (platelets, red- and white blood cells etc.). Four major classes of leukemia are generally referred to, although multiple sub-types have been assigned depending on the classification system used. These are acute myeloid leukemia (AML), acute lymphoblastic leukemia (ALL), chronic myeloid leukemia (CML) and chronic lymphocytic leukemia (CLL).²⁹ In acute leukemia, the cancerous cells are immature blasts, whereas chronic leukemia is characterized by more mature cells of cancerous origin. The terms myeloid and lymphocytic refer to which cell line is affected (see Figure 4, next page).

Figure 4: Development process from stem cells towards blood cells with more specific purposes. Picture from the National Cancer Institute.⁴¹

1.4.2.1. Acute myeloid leukemia

Acute myeloid leukemia (AML) is an aggressive hematopoietic stem cell disorder characterized by infiltration of the bone marrow, blood and other tissues by clonal, proliferative and abnormally differentiated myeloid blast cells. These cancerous cells have acquired specific mutations that result in hampered apoptotic and differentional capabilities. Moreover, these mutations give rise to abilities for uncontrolled proliferation and other survival advantages.⁴² Physiologically, this rapid accumulation of immature myeloid blasts leads to a “crowd-out” effect in the bone marrow which interferes with all normal blood cell production.

The exact cause of acute leukemia remains unknown. However, several major risk factors have been identified including benzene exposure, chemotherapeutic alkylating agents, ionizing radiation, maternal ethanol consumption and genetic factors, e.g. Down’s syndrome. The Epstein-Barr- and Human T-lymphocyte viruses have also been acknowledged as risk factors.²⁹

Diagnosis of AML is established when $\geq 20\%$ of a total blood count consists of myeloid blasts.⁴³ However, AML is a heterogeneous disease and subtype diagnosis is complex.⁴⁴ Two main classification systems are used for the diagnosis of myeloid cancers. One is the so-called French-American-British (FAB) system, which defines eight subtypes of AML based on morphological and cytochemical characteristics. Another classification system has more recently been defined by WHO and considers combinations of morphological, genetic and immunologic characteristics.^{29, 45}

AML is the most common leukemia in adults and continues to be accompanied by the lowest survival rates.⁴⁶ Statistics from 2015, in the USA alone, estimated some 20.830 new cases diagnosed and 10.000 deaths. Advances in AML treatments have led to significantly improved survival rates in younger patients, while the prognoses for the elderly remain poor.⁴⁵ Incurable 50 years ago, AML is now considered cured in 35 - 40% of cases in patients younger than 60 years. These numbers drop drastically in patients over 60 years of age, where the disease is cured in only 5 - 15% cases.⁴⁷ Even with current treatments, 70% of all patients 65 years and older will not survive one year from their point of diagnosis.⁴⁵ This is due to the very intensive chemotherapy that must be initiated immediately after diagnosis, often in an aged patient with other comorbid illnesses. The chemotherapy is followed by serious side effects, even lethal ones such as cardiotoxicity after anthracycline exposure.⁴⁸ If left untreated, death will typically occur within few weeks or months.⁴² Therefore, new alternative chemotherapies that target this disease in a milder and more selective way are utterly called for.

Through intensive induction chemotherapy, the treatment goals are to achieve rapid onset of complete remission (CR). The AML induction therapy consists of so-called “7+3” cycles of a high-dose combination of the cell cycle inhibitor cytarabine (first 7 days) and the anthracycline daunorubicin (first 3 days) (see Figure 5). A complete remission is achieved in the majority of cases (65%) using this combination.⁴⁹ However, CR will typically not last for longer than 4 - 8 months if an intensive post-remission therapy is not initiated immediately after the induction phase. The goal of the post-remission therapy is thus to eradicate any remaining cancerous cells that might have survived the induction phase. The post-remission therapy has been shown to clearly improve survival rates, but the exact duration and number of therapy cycles remain a topic for debate.²⁹

Figure 5: The chemotherapeutic agents cytarabine (Ara-C) and daunorubicin.

1.5 References

1. Kapetanovic, I. M., Overview of Current Drug Discovery and Development with an Eye towards the Future. In *Drug Discovery and Development - Present and Future*, Kapetanovic, I. M., Ed. InTech: InTech, 2011.
2. Fleming, Sir Alexander. A. Noble lecture - Penicillin. https://www.nobelprize.org/nobel_prizes/medicine/laureates/1945/fleming-lecture.html. (accessed 14.06.2017).
3. Kresge, N.; Simoni, R. D.; Hill, R. L., Selman Waksman: the Father of Antibiotics. *J. Biol. Chem.* **2004**, 279 (48), e7.
4. Lewis, K., Platforms for antibiotic discovery. *Nat. Rev. Drug. Discov.* **2013**, 12 (5), 371-387.
5. Dolle, R. E., Historical overview of chemical library design. *Methods Mol Biol* **2011**, 685, 3-25.
6. Oprea, T. I., Current trends in lead discovery: Are we looking for the appropriate properties? *Mol. Divers.* **2000**, 5 (4), 199-208.
7. Lombardino, J. G.; Lowe, J. A., The role of the medicinal chemist in drug discovery – then and now. *Nat. Rev. Drug. Discov.* **2004**, 3 (10), 853-862.
8. Smietana, K.; Siatkowski, M.; Moller, M., Trends in clinical success rates. *Nat Rev Drug Discov* **2016**, 15 (6), 379-380.
9. DeSimone, R. W.; Currie, K. S.; Mitchell, S. A.; Darrow, J. W.; Pippin, D. A., Privileged Structures: Applications in Drug Discovery. *Comb. Chem. High. T. Scr.* **2004**, 7 (5), 473-493.
10. Willoughby, C. A.; Hutchins, S. M.; Rosauer, K. G.; Dhar, M. J.; Chapman, K. T.; Chicchi, G. G.; Sadowski, S.; Weinberg, D. H.; Patel, S.; Malkowitz, L.; Di Salvo, J.; Pacholok, S. G.; Cheng, K., Combinatorial synthesis of 3-(Amidoalkyl) and 3-(Aminoalkyl)-2-arylindole derivatives: discovery of potent ligands for a variety of G-protein coupled receptors. *Bioorg. Med. Chem. Lett.* **2002**, 12 (1), 93-96.
11. Dinnell, K.; Chicchi, G. G.; Dhar, M. J.; Elliott, J. M.; Hollingworth, G. J.; Kurtz, M. M.; Ridgill, M. P.; Rycroft, W.; Tsao, K.-L.; Williams, A. R.; Swain, C. J., 2-Aryl indole NK1 receptor antagonists: optimisation of the 2-Aryl ring and the indole nitrogen substituent. *Bioorg. Med. Chem. Lett.* **2001**, 11 (9), 1237-1240.
12. Welsch, M. E.; Snyder, S. A.; Stockwell, B. R., Privileged scaffolds for library design and drug discovery. *Curr. Opin. Chem. Biol.* **2010**, 14 (3), 347-361.
13. Bemis, G. W.; Murcko, M. A., The Properties of Known Drugs. 1. Molecular Frameworks. *J. Med. Chem.* **1996**, 39 (15), 2887-2893.
14. Evans, B. E.; Rittle, K. E.; Bock, M. G.; DiPardo, R. M.; Freidinger, R. M.; Whitter, W. L.; Lundell, G. F.; Veber, D. F.; Anderson, P. S.; Chang, R. S. L.; Lotti, V. J.; Cerino, D. J.; Chen, T. B.; Kling, P. J.; Kunkel, K. A.; Springer, J. P.; Hirshfield, J., Methods for drug discovery: development of potent, selective, orally effective cholecystokinin antagonists. *J. Med. Chem.* **1988**, 31 (12), 2235-2246.
15. Lipinski, C. A., Lead- and drug-like compounds: the rule-of-five revolution. *Drug. Discov. Today Technol.* **2004**, 1 (4), 337-341.
16. Sliwoski, G.; Kothiwale, S.; Meiler, J.; Lowe, E. W., Computational Methods in Drug Discovery. *Pharmacol. Rev.* **2014**, 66 (1), 334-395.
17. Kapetanovic, I. M., Computer-aided drug discovery and development (CADD): *In silico*-chemico-biological approach. *Chem.-Biol. Interact.* **2008**, 171 (2), 165-176.
18. Lemke, T. L.; Foye, W. O., *Foye's principles of medicinal chemistry*. 6th ed. / Thomas L. Lemke. ed.; Lippincott Williams & Wilkins: Philadelphia, Pa. ; London, 2008.
19. Smith, H. S., Opioid Metabolism. *Mayo Clin. Proc.* **2009**, 84 (7), 613-624.
20. Andersson, T.; Miners, J. O.; Veronese, M. E.; Birkett, D. J., Diazepam metabolism by human liver microsomes is mediated by both S-mephenytoin hydroxylase and CYP3A isoforms. *Br. J. Clin. Pharmacol.* **1994**, 38 (2), 131-137.
21. Osterloh, I. H., The discovery and development of Viagra® (sildenafil citrate). In *Sildenafil*, Dunzendorfer, U., Ed. Birkhäuser Basel: Basel, **2004**; pp 1-13.
22. Zappacosta, A. R. Reversal of Baldness in Patient Receiving Minoxidil for Hypertension. *N. Engl. J. Med.* **1980**, 303 (25), 1480-1481.

23. Cragg, G. M.; Newman, D. J., Natural products: A continuing source of novel drug leads. *Biochim. Biophys. Acta (BBA) - General Subjects* **2013**, *1830* (6), 3670-3695.
24. Cragg, G. M.; Grothaus, P. G.; Newman, D. J., Impact of Natural Products on Developing New Anti-Cancer Agents. *Chem. Rev.* **2009**, *109* (7), 3012-3043.
25. Newman, D. J.; Cragg, G. M., Natural Products as Sources of New Drugs from 1981 to 2014. *J. Nat. Prod.* **2016**, *79* (3), 629-661.
26. Harvey, A. L.; Edrada-Ebel, R.; Quinn, R. J., The re-emergence of natural products for drug discovery in the genomics era. *Nat. Rev. Drug. Discov.* **2015**, *14* (2), 111-129.
27. Sukuru, S. C. K.; Jenkins, J. L.; Beckwith, R. E. J.; Scheiber, J.; Bender, A.; Mikhailov, D.; Davies, J. W.; Glick, M., Plate-Based Diversity Selection Based on Empirical HTS Data to Enhance the Number of Hits and Their Chemical Diversity. *J. Biomol. Screen.* **2009**, *14* (6), 690-699.
28. Jitender, V.; Vijay, M. K.; Evans, C. C., 3D-QSAR in Drug Design - A Review. *Curr. Top. Med. Chem.* **2010**, *10* (1), 95-115.
29. DiPiro, J. T., *Pharmacotherapy : a pathophysiologic approach*. 7th ed. ed.; McGraw-Hill Medical ; London : McGraw-Hill: New York, 2008.
30. Cancer fact sheets: All cancer excluding non-melanoma skin cancer. <https://gco.iarc.fr/today/fact-sheets-cancers?cancer=29&type=0&sex=0> (accessed 13.05.2017).
31. Rang, H. P., *Pharmacology*. 5th ed. ed.; Churchill Livingstone: Edinburgh, 2003.
32. Antimicrobial resistance. <http://www.who.int/mediacentre/factsheets/fs194/en/> (accessed 13.05.2014).
33. Hanahan, D.; Weinberg, R. A., The hallmarks of cancer. *Cell* **2000**, *100* (1), 57-70.
34. Hanahan, D.; Weinberg, R. A., Hallmarks of cancer: the next generation. *Cell* **2011**, *144* (5), 646-74.
35. Wilson, W. R.; Hay, M. P., Targeting hypoxia in cancer therapy. *Nat. Rev. Cancer* **2011**, *11* (6), 393-410.
36. Brown, J. M.; Wilson, W. R., Exploiting tumour hypoxia in cancer treatment. *Nat. Rev. Cancer* **2004**, *4* (6), 437-447.
37. Brown, J. M., Tumor Hypoxia in Cancer Therapy. In *Methods Enzymol.*, Academic Press: 2007; Vol. Volume 435, pp 295-321.
38. Cerecetto, H.; González, M.; Lavaggi, M. L.; Azqueta, A.; López de Cerain, A.; Monge, A., Phenazine 5,10-Dioxide Derivatives as Hypoxic Selective Cytotoxins. *J. Med. Chem.* **2005**, *48* (1), 21-23.
39. Junnotula, V.; Rajapakse, A.; Arbillaga, L.; López de Cerain, A.; Solano, B.; Villar, R.; Monge, A.; Gates, K. S., DNA strand cleaving properties and hypoxia-selective cytotoxicity of 7-chloro-2-thienylcarbonyl-3-trifluoromethylquinoxaline 1,4-dioxide. *Bioorg. Med. Chem.* **2010**, *18* (9), 3125-3132.
40. Hay, M. P.; Gamage, S. A.; Kovacs, M. S.; Pruijn, F. B.; Anderson, R. F.; Patterson, A. V.; Wilson, W. R.; Brown, J. M.; Denny, W. A., Structure-Activity Relationships of 1,2,4-Benzotriazine 1,4-Dioxides as Hypoxia-Selective Analogues of Tirapazamine. *J. Med. Chem.* **2003**, *46* (1), 169-182.
41. Cancer Stat Facts: Leukemia. <https://seer.cancer.gov/statfacts/html/leuks.html> (accessed 02.05.2017).
42. Myhren, L.; Nygaard, G.; Gausdal, G.; Sletta, H.; Teigen, K.; Degnes, K.; Zahlsen, K.; Brunsvik, A.; Bruserud, Ø.; Døskeland, S.; Selheim, F.; Herfindal, L., Iodinin (1,6-Dihydroxyphenazine 5,10-Dioxide) from *Streptosporangium* sp. Induces Apoptosis Selectively in Myeloid Leukemia Cell Lines and Patient Cells. *Mar. Drugs* **2013**, *11* (2), 332.
43. Gilliland, D. G.; Tallman, M. S., Focus on acute leukemias. *Cancer Cell* **2002**, *1* (5), 417-420.
44. Arber, D. A.; Orazi, A.; Hasserjian, R.; Thiele, J.; Borowitz, M. J.; Le Beau, M. M.; Bloomfield, C. D.; Cazzola, M.; Vardiman, J. W., The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood* **2016**, *127* (20), 2391-2405.
45. De Kouchkovsky, I.; Abdul-Hay, M., 'Acute myeloid leukemia: a comprehensive review and 2016 update'. *Blood Cancer J.* **2016**, *6*, e441.
46. Deschler, B.; Lübbert, M., Acute myeloid leukemia: Epidemiology and etiology. *Cancer* **2006**, *107* (9), 2099-2107.

47. Döhner, H.; Weisdorf, D. J.; Bloomfield, C. D., Acute Myeloid Leukemia. *N. Engl. J. Med.* **2015**, *373* (12), 1136-1152.
48. Simunek, T.; Sterba, M.; Popelova, O.; Adamcova, M.; Hrdina, R.; Gersl, V., Anthracycline-induced cardiotoxicity: overview of studies examining the roles of oxidative stress and free cellular iron. *Pharmacol. Rep.* **2009**, *61* (1), 154-71.
49. Mayer , R. J.; Davis , R. B.; Schiffer , C. A.; Berg , D. T.; Powell , B. L.; Schulman , P.; Omura , G. A.; Moore , J. O.; McIntyre , O. R.; Frei , E., Intensive Postremission Chemotherapy in Adults with Acute Myeloid Leukemia. *N. Engl. J. Med.* **1994**, *331* (14), 896-903.

Chapter 2 – Overall aims of study

The purpose of this work was to discover new drug leads for improved therapies of acute myeloid leukemia, infections caused by resistant bacteria and metabolic diseases such as atherosclerosis, diabetes mellitus type 2 (T2DM) and obesity. We set off with the aim of investigating two related, yet fundamentally different drug targets, namely the DNA (of leukemic cells and bacteria) and the liver X receptors (LXRs). We already had ideas related to i) heterocyclic compounds with potential effects on acute myeloid leukemia (AML) and infectious diseases caused by resistant bacteria, and ii) oxygenated steroid derivatives showing activity towards nuclear receptors controlling gene products related to T2DM, obesity and atherosclerosis.

Our interest in phenazine 5,10-dioxides was based on recent reports in which the natural product iodinin (**5**) was shown to be a potent and selective inducer of apoptotic cell death in human AML cell lines.¹⁻² Our initial plan in this context was to establish a new total synthesis of the lead compound, which would give access to sufficient amounts of the product for further lead development and biological evaluations. The synthesis of analogues was considered essential to map structure-activity relationships, in order to gain a more detailed insight into their DNA-damaging mechanism of action and to guide a rational design of new antileukemic drug candidates.

In the light of counteracting atherosclerosis, agonists of the LXRs have been pursued in drug discovery due to their desirable upregulation of the expression of ATP-binding cassettes.³ However, it is also well documented that long-term exposure to full agonists of the LXRs results in undesirable side-effects, such as hypertriglyceridemia, upregulated lipogenesis and hepatic steatosis.⁴ Our enthusiasm in this field was based on reports demonstrating that 22(*S*)-hydroxycholesterol is an LXR-antagonist, with a potential to selectively downregulate the expression of genes involved in lipid formation, while simultaneously increasing glucose uptake.⁵⁻⁷ Interestingly, its natural epimer, the endogenous compound 22(*R*)-hydroxycholesterol is an LXR full agonist.⁸⁻⁹ We envisioned that virtual screening, synthesis and biological evaluations of different oxysterols functionalized in position 22 could provide new knowledge about the ligand-dependent regulation of LXR target gene expression and possibly guide us towards discoveries of new drug candidates for the treatment of LXR-related diseases like atherosclerosis, T2DM and obesity.

The two projects described above differ in terms of ligand structures and targets. However, they are connected by aspects like drug discovery, medicinal chemistry and structure-activity relationship studies, in which small-molecular compounds, derived from natural products, are in focus.

More specific descriptions of other aims of the work described herein are given in Chapters 4 and 7.

2.1 References

1. Sletta, H.; Degnes, K. F.; Herfindal, L.; Klinkenberg, G.; Fjaervik, E.; Zahlén, K.; Brunsvik, A.; Nygaard, G.; Aachmann, F. L.; Ellingsen, T. E.; Doskeland, S. O.; Zotchev, S. B., Anti-microbial and cytotoxic 1,6-dihydroxyphenazine-5,10-dioxide (iodinin) produced by *Streptosporangium* sp. DSM 45942 isolated from the fjord sediment. *Appl. Microbiol. Biotechnol.* **2014**, *98* (2), 603-10.
2. Myhren, L. E.; Nygaard, G.; Gausdal, G.; Sletta, H.; Teigen, K.; Degnes, K. F.; Zahlén, K.; Brunsvik, A.; Bruserud, O.; Doskeland, S. O.; Selheim, F.; Herfindal, L., Iodinin (1,6-dihydroxyphenazine 5,10-dioxide) from *Streptosporangium* sp. induces apoptosis selectively in myeloid leukemia cell lines and patient cells. *Mar. Drugs* **2013**, *11* (2), 332-49.
3. Hong, C.; Tontonoz, P., Liver X receptors in lipid metabolism: opportunities for drug discovery. *Nat. Rev. Drug. Discov.* **2014**, *13* (6), 433-444.
4. Schultz, J. R.; Tu, H.; Luk, A.; Repa, J. J.; Medina, J. C.; Li, L.; Schwendner, S.; Wang, S.; Thoolen, M.; Mangelsdorf, D. J.; Lustig, K. D.; Shan, B., Role of LXRs in control of lipogenesis. *Genes Dev.* **2000**, *14* (22), 2831-2838.
5. Kase, E. T.; Thoresen, G. H.; Westerlund, S.; Højlund, K.; Rustan, A. C.; Gaster, M., Liver X receptor antagonist reduces lipid formation and increases glucose metabolism in myotubes from lean, obese and type 2 diabetic individuals. *Diabetologia* **2007**, *50* (10), 2171-2180.
6. Kase, E. T.; Andersen, B.; Nebb, H. I.; Rustan, A. C.; Thoresen, G. H., 22-Hydroxycholesterols regulate lipid metabolism differently than T0901317 in human myotubes. *Biochim. Biophys. Acta, Mol. Cell Biol. Lipids* **2006**, *1761* (12), 1515-1522.
7. Kase, E. T.; Nikolić, N.; Hessvik, N. P.; Fjeldheim, Å.-K.; Jensen, J.; Thoresen, G. H.; Rustan, A. C., Dietary Supplementation with 22-S-Hydroxycholesterol to Rats Reduces Body Weight Gain and the Accumulation of Liver Triacylglycerol. *Lipids* **2012**, *47* (5), 483-493.
8. Janowski, B. A.; Willy, P. J.; Devi, T. R.; Falck, J. R.; Mangelsdorf, D. J., An oxysterol signalling pathway mediated by the nuclear receptor LXR[alpha]. *Nature* **1996**, *383* (6602), 728-731.
9. Viennois, E.; Mouzat, K.; Dufour, J.; Morel, L.; Lobaccaro, J.-M.; Baron, S., Selective liver X receptor modulators (SLiMs): What use in human health? *Mol. Cell. Endocrinol.* **2012**, *351*, 129-141.

Chapter 3 – Introduction to phenazines

The phenazine (PHz) literature is vast, spanning more than 150 years. A detailed overview covering all aspects of current knowledge regarding the physiological roles of phenazines, their biosynthesis, modes of action and synthetic methods is far beyond the scope of the presented introduction. However, some key aspects regarding the current knowledge on their biosynthesis, modes of action and synthetic preparations will be addressed. Specific PHz molecules of interest to the presented work will also be discussed with the aim to provide the reader with a better understanding of which biological effects can be expected by looking at different PHzs.

Phenazines (PHzs) are a large class of colorful aromatic *N*-heterocyclic compounds. They are found widely in nature, both in terrestrial and water-habited environments, as well as in surroundings that are both rich in and deprived of oxygen. The biosynthesis of PHzs has been confirmed to take place in a wide array of Gram-positive (G⁺)¹ and Gram-negative (G⁻)² bacteria, and even in the archaeal species *Methanosarcona*.³ Natural production is thus limited to bacteria and archaea of which the sub-types of *Streptomyces* (G⁺) and *Pseudomonas* (G⁻) are generally referred to as the richest natural sources. Several excellent comprehensive reviews have been published that highlight their occurrence, physiological roles, modes of action, synthesis and biosynthesis.³⁻⁶

The phenazine core structure consists of a central pyrazine ring annulated by two benzene rings (**1**, Figure 6) and the colors of these bacterial metabolites range over the whole visible spectrum depending on which functional groups decorate the scaffold. Each nitrogen atom can also be found covalently bound to oxygen, forming *N*-oxides, as exemplified by phenazine 5,10-dioxide (**2**) (Figure 6). The colorful nature of these compounds therefore facilitated an early discovery of the first PHz natural products within bacterial cultures.⁵

Figure 6. Phenazine (**1**) and phenazine 5,10-dioxide (**2**).

Pyocyanin (**3**, PYO, Figure 7), isolated by Fordos in 1859 is the oldest known phenazine and the one most extensively studied to date.⁶ This substance (**3**) was originally isolated by chloroform extraction of blue colored wound dressings probably containing the opportunistic pathogen *Pseudomonas aeruginosa*, later known for its extensive production of the blue dye. Pyocyanin (**3**) is essential for survival of *Pseudomonas aeruginosa* at late stages of growth and an important virulence factor that worsens condition of the lung tissue in patients suffering from cystic fibrosis.⁷ Its mode of action will be discussed in more detail in section 3.2.1, page 22.

Figure 7: The natural products pyocyanin (**3**), chlororaphine (**4**) and iodinin (**5**)

The discoveries of other PHzs followed and among these were the green pigment chlororaphine (**2**) reported in 1894 and the purple, glittering iodinin (**5**) reported in 1938 (Figure 7).¹ The current literature includes accounts of over 180 different PHzs of natural origin and if all synthetic structures are contained within, this number rises to well over 6000 reported compounds.⁵ Hundreds of these compounds display a range of biological effects, attesting to the immense structural variety of the PHzs. In many cases, anti-infective and antitumor effects have been recognized in addition to several other biological functions. Examples of natural products include the antibiotics griseolutein A (**6**), lomofungin (**7**) and myxin (**8**) as well as mycomethoxin (**9**) which is active against *Mycobacterium tuberculosis*, hence its name (Figure 8, top row, next page). Selected examples of compounds that promote the death of cancerous cells include saphenamycin (**10**) and esmeraldin B (**11**). A variety of other functions have been reported, such as the potent radical scavenging effect of benthocyanin A (**12**) (Figure 8, bottom row, next page).⁸

Figure 8: A selection of phenazine natural products

To the authors' best knowledge, no PHz has to this date gained market approval for therapeutic use in humans. This statement might come as a surprise if seen in the context of the long-term interest in these compounds from both academia and the pharmaceutical industry, especially regarding growth inhibition of microbes and cancerous cells. To approach a logical explanation to this paradox, several aspects should be considered and the first of these is toxicity. Phenazines are small-molecular compounds and able to interfere with an array of biological functions. The planar aromatic nature of this compound class has especially been feared to result in unselective binding or interference with eukaryotic DNA. Secondly, due to the multiple modes of action involved and the relatively poor understanding concerning physiological roles, other antibiotics with more appealing safety profiles were certainly ranked higher during the golden era of antibiotics, when antimicrobial research was as a top priority in the pharmaceutical industry. A lack of efficient synthetic methods in the past has also provided a barrier for commercialized production, and in some cases purification, due to the formation of regioisomers. Due to the immense variety in terms of chemical structures and biological actions, it is extremely difficult to conclude on any general structure-activity relationships of this compound class.

Despite the considerations mentioned above, the PHz scaffold comes from nature and should thus be considered as a privileged scaffold with a potential to interact with an array of extra- and intracellular targets. Accordingly, if a PHz lead compound is identified, it is an excellent starting point for drug discovery and lead optimization via medicinal chemistry.

3.1 Biosynthesis

All PHz-producing species studied have been found to possess a specific conserved set of genes clustered in a single operon.⁹ Earlier studies on PHz biosynthesis were mostly performed on *Pseudomonas* subtypes, which produce relatively simple structures such as pyocyanin (**3**), PHz 1-carboxylic acid (PCA, **13**) and PHz 1,6-dicarboxylic acid (PDC, **14**).⁴ The discovery of this operon at the end of last century has paved the way for a more detailed and stepwise understanding of the PHz biosynthetic process.¹⁰ Moreover, the gene cluster can most likely be transferred via horizontal gene transfer⁹ and includes a set of at least 5 enzymes that are encoded therein (PhzB, PhzD, PhzE, PhzF and PhzG, the structures of all of which are available in the RCSB Protein Databank). These enzymes are responsible for the biocatalyzed production

Scheme 1: Key steps representing the current knowledge of the phenazine biosynthetic pathway.⁵

of the two core phenazines from which all other functionalized PHzs are believed to be derived, namely PDC (**13**) and PCA (**14**).⁵ The PHz operon also contains extensions of the gene cluster so that the two key building blocks can be further derivatized into more strain-specific and complex PHzs (Scheme 1).¹⁰

The biosynthesis of PHzs has recently been reviewed.^{5, 10} In a brief summary, chorismic acid (**15**) is transformed into the highly unstable 6-amino-5-oxocyclohex-2-ene-1-carboxylic acid (AOCHC, **18**) in 4 steps (see Scheme 1). The first intermediate of this path is 2-amino-2-desoxyisochorismic acid (ADIC, **16**) produced by PhzE-mediated, stereoselective amination and dehydration of **15**. ADIC (**16**) is then converted to *trans*-2,3-dihydro-3-hydroxyanthranilic acid (DHHA, **17**) by PhzD. This substrate is then further transformed into AOCHC (**18**). As mentioned earlier, the intermediate **18** is highly unstable and is therefore subject to rapid self-condensation, in which two identical molecules condense to form hexahydrophenazine-1,6-dicarboxylic acid (HHPDC, **19**). Compound **19** is also unstable and can aromatize in PhzG to give tetrahydrophenazine-1-carboxylic acid (**20**) or spontaneously undergo non-enzymatic, oxidative decarboxylation to form tetrahydrophenazine-1,6-dicarboxylic acid (THPCA, **21**). The two compounds **20** and **21** are oxidized further to afford dehydrophenazine 1-carboxylic acid (DHPCA, **22**) and dehydrophenazine 1,6-dicarboxylic acid (DHPDC, **23**). These are the true precursors for all other phenazines with more complex structures. Compounds **22** and **23** can also undergo further oxidation to simply give PCA (**13**) and PDC (**14**), respectively. The tautomeric form of **21**, THPCAa (**21a**), can also undergo oxidative carboxylation once more to afford dehydrophenazine (DHPH_z, **24**) which is oxidized further to **1**.

Strain-specific phenazines of great chemical diversity have been identified from bacterial sources that thrive in different environments and it is fair to say that there are large gaps in our general understanding of these biosynthetic processes.

3.2 Biological functions and modes of action

Phenazines are redox-active molecules and can therefore accept- and donate electrons. The direction of this process depends on the redox potential of the phenazine compared with other molecules that can transport electrons, and that are within range of the phenazine.¹¹ Hundreds of PHzs reported throughout the literature display various interesting biological features.⁴ Many promote biological events that result in anti-infective-¹²⁻¹³ and antitumor effects.^{6, 14-16} In addition, biofilm eradicating-¹⁷⁻¹⁸ anti-inflammatory-¹⁹ radical scavenging-⁸ and immunosuppressive effects²⁰ been acknowledged as part of their effects just to name few examples. As a result of that, various modes of action have been assigned for their individual biological effects such as DNA-intercalation,¹⁵ inhibition of topoisomerases,²¹ production of reactive oxygen species (ROS),²²⁻²³ metal chelation²⁴ and radical scavenging.⁸ Several modes of action should thus be considered when a PHz displays biological effects.

Like many other natural products, phenazines have undergone the term “secondary metabolites” although that word can be controversial in its nature.²⁵ The classical definition of a secondary metabolite accounts for organic molecules that serve no specific role in the growth, reproduction or replication of a living organism. In reflection of this, the exact physiological roles of phenazines have been a mystery for a long time.⁴ However, some important questions have to be addressed such as: Why should nutrient-deprived organisms, in the late stages of their growth in dense colonies, spend their energy, time and effort producing large amounts of PHzs² and why would they conserve the genes for their PHz-producing protein machinery through evolution, spanning millions of years or longer?²⁵ It must be that these genes are of critical importance to their hosts. Reflecting on this notion, it has been shown that PHz-producing bacteria survive longer compared with mutant strains that cannot produce them.⁴

3.2.1 Redox cycling of natural phenazines

The most studied phenazine to date is the blue pigment PYO (**3**) (see Figure 9, next page), produced by the Gram-negative bacteria *Pseudomonas aeruginosa*, which gives their colonies a characteristic blue-green appearance. As for the other phenazines, PYO (**3**) is secreted during late stages of growth in colonies of high cell densities and can be isolated from the sputum of patients with cystic fibrosis in relatively large quantities.²⁶ This compound, which is readily able to penetrate biological membranes due to its zwitterionic nature,² promotes oxidative stress to the pathogen's host cells and disrupts catalase and electron transport in its mitochondria.²⁷ PYO has been shown to induce apoptotic cell death in neutrophils thus

providing suppressive effects to the human immune system which eventually results in enhanced bacterial virulence.²⁸ *Pseudomonas aeruginosa* produces many other virulence factors, including the phenazines PCA (**13**) and 1-hydroxyphenazine (**25**) (Figure 9) as well as multiple other non-PHz factors which contribute to the virulence. This makes the pathophysiology of these infections highly complex.²⁹ There is evidence to indicate that PYO (**3**) production significantly benefit the growth and survival of *Pseudomonas* strains in mice infections.³⁰

Figure 9: Pyocyanin (**3**) and 1-hydroxyphenazine (**25**), virulence factors of *Pseudomonas aeruginosa*.

A major impact factor in PYO's (**3**) biological actions in eukaryotic cells is the production of reactive oxygen species (ROS) and this feature promotes oxidative stress.⁷ The production of ROS takes place through a process called redox cycling. This process is initiated by a reduction of PYO (**3**) by various reducing agents such as NADH, NADPH or glutathione.²² The reduction of PYO affords a PYO-radical (activation), which can oxidize molecular oxygen to yield the superoxide radical ($O_2^{\cdot-}$). This superoxide radical can further undergo a cascade of reactions to form other ROS, such as hydrogen peroxide (H_2O_2), and even the hydroxyl radical ($\cdot OH$).³¹ PYO has also been shown to deplete cellular storage of glutathione (GSH), an important cell anti-oxidant that detoxifies harmful reactive substances, including ROS.^{5, 32} All these actions combined impact the host's redox balance in a negative way.²⁵ The ROS-production can also inhibit the growth of other competing microorganisms, especially those that display low activity of superoxide dismutase.³³ Same mode of action (redox cycling) was also later confirmed for 1-hydroxyphenazine (**25**)²² (see Scheme 2).

Scheme 2: Production of reactive oxygen species by 1-hydroxyphenazine (**25**).²²

3.2.2 Lomofungin, polynucleotide interaction and metal chelation

The PHz antibiotic lomofungin (methyl 6-formyl-4,7,9-trihydroxyphenazine-1-carboxylate; **7**, Figure 10) was isolated from *Streptomyces lomodensis*.³⁴⁻³⁵ This compound received attention in the 1960s - 70s for its effects on polynucleotide synthesis in microorganisms. Lomofungin (**7**) was found to rapidly inhibit DNA and RNA synthesis in yeast.³⁶⁻³⁷ These effects were believed to stem from inhibition of DNA-dependent RNA polymerase. However, Fraser et al. found lomofungin (**7**) to inhibit RNA synthesis in a manner analogous to that of 8-hydroxyquinoline (**26**, Figure 10), a well-known metal cation chelating agent.³⁸ Thus, lomofungin (**7**) may be able to chelate divalent cations, such as Mn^{2+} , Mg^{2+} and Zn^{2+} , and thereby interfere with RNA-polymerase activity (in *E. choli*).³⁹ Indeed, Pavletich et al. proposed that the mode of lomofungin's action involves chelation of the tightly bound Zn^{2+} of the DNA- and RNA-polymerase.⁴⁰ Although the mode of action has not been unraveled in clear detail, these publications suggest that lomofungin (**7**) strongly interferes with both RNA- and DNA metabolism. In line with that hypothesis, lomofungin (**7**) was more recently found to be the most potent hit out of ~280.000 compounds in an HTS conducted to identify inhibitors of the binding of MBNL1-CUG to RNA.⁴¹ MBNL1-CUG is associated with the genetic disease myotonic dystrophy type 1, characterized by extensive repetitions of cytosine-thymine-guanine

Figure 10: The structures of Lomofungin (**7**), 8-Hydroxyquinoline (**26**) and Dilomofungin (**27**).

(CTG) in the genetic material. Interestingly, lomofungin (**7**) was found to dimerize spontaneously in a solution of DMSO to transform into the dimer dilomofungin (**27**, Figure 10), which in turn proved to be 17-times more potent than the parent compound against this target.⁴¹ No clarification nor any mechanistic insights were given to account for how **7** dimerizes and forms a new carbon-carbon bond.⁴¹ Pavletic et al. also mention that DMSO solutions of **7** rapidly lose their potency when polymerase activity was studied.⁴⁰ To the best of the authors' knowledge, no total synthesis of lomofungin (**7**) has ever been reported.

3.2.3 Iodinin

Iodinin (**5**, Figure 11) or 1,6-dihydroxyphenazine 5,10-dioxide is among the first described phenazines⁶ and was identified by Davies in 1939, as purple glittering crystals floating on the surface of old *Brevibacterium iodinum* cultures.⁴² The chemical structure of iodinin (**5**) is quite special as two intramolecular ion-dipole bonds drastically reduce solubility of the pigment in both common organic solvents, as well as in aqueous media below pH ~12. As heterocyclic, aromatic *N*-oxides have later proved to be of very rare natural occurrence,⁴³ iodinin (**5**) is thus most likely the first isolated natural product in possession of such functional groups.

Figure 11: Iodinin and its reduced form 1,6-dihydroxyphenazine

The antimicrobial effects of this compound were first noted by McIlwain, although the molecular structure was not known in detail at the time.⁴⁴ A variety of bacteria have later been shown to produce iodinin (**5**).^{4, 6} Tanabe and co-workers conducted a detailed investigation into the spectrum of iodinin's (**5**) antimicrobial effects, revealing growth inhibition of various bacteria, fungi and microalgae at concentrations varying from 1 - 30 µg/mL. The same study also showed that the reduced form of iodinin (**5**), namely 1,6-dihydroxyphenazine (**28**) which is also its biosynthetic precursor,⁴³ possessed no antimicrobial effects.⁴⁵

The first indications of anticancer activity were probably revealed by Hollstein, who showed that both iodinin (**5**) and myxin (**8**) intercalate in double-stranded DNA. The studies were based on shifts in UV-absorption and no interference with single-stranded polynucleotides was observed.¹⁵ Moreover, Hano et al. showed that iodinin (**5**) extends the lifetime of mice with Ehrlich sarcoma compared with placebo control, where other non-oxidized phenazines were studied as well and were not found to possess comparable activities.⁴⁶

More recently, iodinin was isolated from marine *actinomycetes* collected from sediment samples in the Trondheim fjord of Norway.⁴⁷⁻⁴⁸ Screening in various cell lines revealed that

iodinin (**5**) possesses cytotoxic properties and high selectivity towards human cell lines of acute myeloid leukemia (AML) and acute promyelocytic leukemia (APL) – the EC₅₀ values for apoptotic cell death were found to be up to 40 times lower than for various normal cell lines. Moreover, iodinin (**5**) was found to induce apoptosis in cell lines characterized by genetic markers of poor patient prognosis.^{14, 48} Figure 12 describes some features of the nuclear changes in morphology and the signaling events induced by iodinin (**5**). (Figure 12A obtained from the Herfindal/Døskeland laboratory at the University of Bergen, Figure 12B from Myhren et al., 2013).¹⁴

Figure 12: a) Nuclear morphology of AML cell lines treated with 0.3 or 0.6 μ M iodinin (**5**) for 24 hours. The cell lines are the human AML cells MV4-11, MOLM-13, as well as MOLM-13 with silencing of the anti-oncogen p53 (FigureA was obtained from the Herfindal/Døskeland laboratory at UiB, unpublished data) b) IPC-81 rat AML cells were treated with iodinin (**5**) or daunorubicin and the cell lysates were immunoblotted to detect procaspase 3, cleaved caspase 3, and phosphorylation of γ H2AX. Caspase 3 is an important element in the apoptotic machinery, and γ H2AX is phosphorylated if DNA strand-breaks occur. (Figure 12B was obtained from Myhren *et.al.*, 2013)¹⁴

3.2.4 Myxin

Peterson et al. reported the isolation of an unusually potent antibiotic from a *Sorangium* species in 1966. Therein, the newly discovered cherry-red compound was shown to inhibit the growth of a broad range of microorganisms including Gram-positive and -negative bacteria, in addition to fungi, actinomycetes and yeasts.⁴⁹ The structure was later elucidated and shown to be 1-hydroxy-6-methoxyphenazine 5,10-dioxide and was given the trivial name myxin (**8**, Figure 13).⁵⁰⁻⁵¹ Lesley and Behki further displayed myxin's (**8**) potency on *Escherichia coli* revealing that synthesis of new bacterial DNA was completely inhibited within few minutes at concentration of 5 µg/mL.⁵²⁻⁵³ Due to these potent antibiotic activities, myxin (**8**) is likely to be the phenazine that has been most thoroughly investigated after PYO (**3**).^{12-13, 53-55}

Myxin (**8**) was investigated by the pharmaceutical company Hoffmann-La Roche in the 1970s where *in vitro* and *in vivo* studies were conducted.⁵⁶ In addition, **8** was developed as a cupric complex (cuprimyxin, **29**) to a product that gained FDA-approval (Unitrop®) for the treatment of veterinary skin infections.^{6, 57} The manufacturing process was based on semi synthesis from iodinin (**5**) cultured from *Brevibacterium iodinum*. Weigele and Leimgruber also prepared synthetic derivatives and performed a SAR-study.¹² As a general trend observed in this study, the antimicrobial effects were shown to drop when the structure was enlarged with substituents. On the contrary, two compounds that can be considered to have less complex structures than myxin (**8**) were found to be very active; 1-methoxyphenazine 5,10-dioxide (**30**) and 1-methylphenazine 5,10-dioxide (**31**, Figure 13).¹²

Concerning its biological effects on eukaryotic cells, several lines of evidence suggest that Myxin's (**8**) mode of action stems from several features. First, Hollstein et al. demonstrated

Figure 13: The antibiotics myxin (**8**), cuprimyxin (**29**, Unitrop®), 1-methoxyphenazine 5,10-dioxide (**30**) and 1-methylphenazine 5,10-dioxide (**31**).

that myxin (**8**) interferes with polynucleotides via DNA intercalation, especially in guanine-cytosine-rich regions of the DNA double-strand. A similar affinity was observed for myxin (**8**) as for other known intercalators, such as ethidium bromide and actinomycin.^{15, 55} Shortly after, the same authors conducted experiments showing that myxin (**8**) affected the rate terms for the incorporation of the nucleoside triphosphates CTP and GTP into RNA. This effect was not observed for ATP and UTP.⁵⁵

More recently, the group of Gates showed that myxin (**8**) causes radical-mediated DNA strand cleavage in eukaryotic cells.⁵⁴ The process is initiated by an intracellular, single-electron enzymatic reduction (see Scheme 3). The reduction affords an unstable intermediate radical of myxin (**8b**) which can further collapse and release the highly reactive and cytotoxic hydroxyl radical ($\cdot\text{OH}$), as well as the reduced myxin analog **32**. The hydroxyl radical is well known for causing DNA damage and being able to abstract protons from the DNA backbone.⁵⁸⁻⁵⁹ Gates et al. also demonstrated the cytotoxic activities of myxin (**8**) against the cancerous human colorectal cell line HCT-116 under both aerobic and anaerobic conditions ($\text{IC}_{50}=1.75\ \mu\text{M}$, at normoxia).⁶⁰ Although this work did not indicate any hypoxia-selectivity for myxin (**8**) in this particular cell line, the reported mode of action of myxin (**8**) is strikingly similar to that of a well-known hypoxia-selective cytotoxic agent, namely tirapazamine (**35**), discussed further in section 3.2.5 below.

Scheme 3: The discharge of reactive oxygen species by myxin (**8**).⁵⁴

3.2.5 Phenazine 5,10-dioxides, tirapazamine and cancer

As emphasized earlier, in section 1.4.1 in page 9, the oxygen saturation level varies considerably between normal healthy cells and cells of solid tumors. The microenvironment of solid tumors is characterized by abnormal vascularization as a consequence of chaotic growth.⁶¹ This phenomenon affords regions of tumors that are severely hypoxic. The hypoxic

cells are also often resistant to radiation and chemotherapy, a result that can enhance disease progression and metastasis.⁶²

Cellular hypoxia may be exploited and offers a potential window to differentiate selectively between healthy and cancerous cells, using hypoxia-selective drugs. Such drugs are dependent on molecular oxygen for a detoxifying back-oxidation, after they are reduced by reducing agents to form radicals. Three scaffolds possessing a potential to act more potently under hypoxic conditions have been identified: Phenazine 5,10-dioxides (**2**),⁶³ di-*N*-oxidized derivatives of quinoxalines (**33**)⁶⁴ and 1,2,4-benzotriazines (**34**) (see Figure 14).⁶⁵ In addition, Cimmino et al. reviewed phenazines and anticancer activity, concluding that phenazine 5,10-dioxides are among the most attractive phenazines to be exploited for anticancer purposes.¹⁶

Figure 14. The core structures of known hypoxia selective scaffolds and tirapazamine (**35**).

Tirapazamine (3-amino-1,2,4-benzotriazine 1,4-dioxide; TPZ **35**, Figure 14) is the most well-documented hypoxia-selective cytotoxic agent to date and has been the focal point of several phase I, II and III clinical trials attempting to treat various forms of cancer. In some cases, clinical efficacy has been confirmed and especially in combination with *cis*-platin.⁶⁶ Tirapazamin's (**35**) hypoxia-selective cell-killing ability stems from its mode of action which is influenced by the level of molecular oxygen in its surroundings. The mechanism is initiated with an enzymatic single-electron reduction of the substrate, which in turn affords an unstable radical intermediate (**35b**) of the TPZ (**4**) (see Scheme 4, next page). If the cell is well oxygenated (normoxic), the drug-radical **35b** is rapidly converted back to the mother compound **35** at the cost of one superoxide radical ($O_2^{\cdot-}$) from molecular O_2 .⁶⁷⁻⁶⁸ In a hypoxic environment however, the lifetime of the drug-radical **35b** is prolonged which allows it to form highly cytotoxic species, such as the hydroxyl radical ($\cdot OH$) and an oxidizing benzotriazinyl radical (**35c**). It should be pointed out, however, that the detailed mechanistic nature of this

process has not been fully elucidated and it has been proposed that the radicals formed in this process inhibit topoisomerase II.⁶⁹

Scheme 4: Proposed mode of action for tirapazamine (35), a hypoxia selective lead compound.⁷⁰

Gates et al. investigated the degradation products formed upon TPZ-promoted DNA damage. Therein, the data suggested that the damage occurred via proton abstraction from three different positions on the deoxyriboses of the DNA double-strand helix. In addition, the degradation products were found to be of the same identity as those produced by the hydroxyl radical ($\cdot\text{OH}$).⁷¹

Some phenazine 5,10-dioxides have been shown to selectively become bioreduced analogously to TPZ (35) and can therefore produce cytotoxic ROS selectively in hypoxic cells.⁷²⁻⁷³ This feature makes the phenazine 5,10-dioxide subclass the most promising in terms of anticancer applications.¹⁶

Few publications have pursued the phenazine 5,10-dioxides as hypoxia-selective cytotoxins for anticancer purposes. Most of the work has been reported by the group of Cerecetto et al.^{63, 72, 74-75} The same group synthesized and biologically evaluated several synthetic phenazine 5,10-dioxides, with some of the resulting compounds displaying high hypoxic selectivity in studies conducted in V79-cells (Chinese hamster, lung fibroblasts). Interestingly, the compounds tested were regioisomeric mixtures of 7- and 8-substituted phenazine 5,10-dioxides, to use of which is inevitable due to the chromatographic inseparability of these isomers, formed in the synthetic process. These studies demonstrated a transformation of the phenazine 5,10-dioxides to their corresponding mono- and double-reduced compounds after *N*-oxide disposal, indicating discharge of the hydroxyl radical ($\cdot\text{OH}$). For example, compound 36 (Figure 15) caused 80% survival of cells under normoxic conditions, while 0% survived in hypoxia (at 20 μM concentration).^{72, 75} The hypoxia-selectivity effect was considerably reduced

only by replacing the -OH group by -NH₂ in position 2 of the scaffold (Compound **37**, Figure 15).⁷²

Figure 15. Hypoxia selective phenazine 5,10-dioxides.⁷²

3.3 Synthesis of phenazines

The following discussion will highlight the most used synthetic methods to construct the phenazine scaffold, their advantages and limitations. These methods have been reviewed in several places⁴⁻⁵ and will only be addressed shortly in the following subsections. When constructing a phenazine, the most important aspect to consider is the resulting location on the phenazine scaffold of any desired substituents. In addition, some of the potential synthetic routes will involve reactions that afford chromatographically inseparable regioisomers, if the starting materials are not chosen wisely. Scheme 5 provides an overview over different approaches to synthesize phenazines.

Scheme 5: Common synthetic routes to the phenazine scaffold.

3.3.1 The Wohl-Aue condensation

The Wohl-Aue reaction was reported in 1901 by Wohl and Aue⁷⁶ and describes the condensation of nitrobenzenes and anilines. The method has severe disadvantages which are represented by very low yields (<15%),^{18, 77-78} limited substrate scope and harsh reaction conditions (strongly alkaline). Despite the disappointing yields and tedious workups, the method has proven to be of particular value in the construction of 1,6-*O*-substituted phenazines – a substitution pattern found in many phenazine natural products. Pachter and Kloetzel, and more recently Chowdhury et al., used the Wohl-Aue reaction to synthesize 1,6-dimethoxyphenazine (**39**) from 2-nitroanisole (**38**, Scheme 6, left).^{54, 77} Another recent example

includes the synthesis of 8-bromo-1-methoxyphenazine (**39**) by Garrison et al. from the same starting material, 2-nitroanisole **38** (Scheme 6, right).¹⁸

Scheme 6: Wohl-Aue synthesis of 1,6-dimethoxyphenazine by Pachter and Kloetzel in 1951⁷⁷ and of 8-bromo-1-methoxyphenazine by Garrison et al. in 2016.¹⁸

3.3.2 The Bamberger-Ham reaction

The Bamberger-Ham reaction is another old method (1911) that is based on the dimerization of two nitrosobenzenes under strongly acidic conditions.⁷⁹ Like the Wohl-Aue procedure, the method is plagued by low yields, harsh conditions and of an even more limited substrate scope, as a result of the mandatory electron-donating groups in *para*-positions to the nitroso groups.⁸⁰ Thus, this method appears to be of very limited value unless 2,7-substituted phenazines are to be synthesized (Scheme 7). Furthermore, an additional reductive deoxygenation step is required if the phenazine *N*-oxides are not wanted. The general procedure is depicted in Scheme 7, as no literature reports including detailed experimental data were found.

Scheme 7: The Bamberger-Ham phenazine synthesis

3.3.3 The Beirut-reaction

Haddadin and co-workers at the American University of Beirut introduced a new method for synthesizing the phenazine scaffold in 1971. The procedure involves a base-catalyzed condensation between phenols and benzofuroxan (**41**) (Scheme 8, left).⁸¹ This method gave access to phenazine 5,10-dioxides with substitution patterns that were inaccessible by other methods reported in prior. Six years later, the same group had developed a modification of the reaction, and showed how 1,2-diketones could be reacted with benzofuroxan (**41**) in the presence of a base, thereby providing access to 1-hydroxysubstituted phenazines applying this method (Scheme 8, right).⁸²

Scheme 8: The Beirut synthesis of phenazines. Examples of when benzofuroxan (**41**) is reacted with catechol (left) or cyclohexane-1,2-diones (right).

These methods are of high value since they provide synthetic routes to phenazines that are not accessible efficiently by other methods. The majority of the reactions reported in these two publications afforded several grams of recrystallized product.⁸¹⁻⁸² The reaction mechanism for the diketone condensation starts with nucleophilic attack on the electrophilic nitrogen of benzofuroxan by the enolate of cyclohexane-1,2-dione, formed in neat diethylamine (Scheme 9, next page). If this method is to be applied for the synthesis of phenazines without *N*-oxides, reductions may be easily conducted, using e.g. sodium borohydride (NaBH_4) or sodium dithionite ($\text{Na}_2\text{S}_2\text{O}_4$). If the intention is to synthesize phenazine 5,10-dioxides however, oxidations with peracids often prove necessary to increase the yields of the phenazine 5,10-dioxides, since the crude reaction mixtures consist of both mono- and dioxidized phenazines.⁸²

Scheme 9: A plausible reaction mechanism for the Beirut diketone application.

3.3.4 Condensations of 1,2-benzoquinones and *O*-phenylenediamines

1,2-benzoquinones can be condensed with *O*-phenylenediamines to give phenazines. The earliest example of this concept dates back to 1886, when *ortho*-phenylenediamine was condensed with catechol in the presence of lead dioxide (PbO_2).⁴ Modern applications of the reaction involve other catechol-oxidizing agents to afford their corresponding diketones, such as silver(I) oxide (Ag_2O),⁸³ and *ortho*-chloranil.^{17, 83} Typically, the produced unstable diketones are reacted further as crudes with *ortho*-phenylenediamines giving the phenazine skeleton upon condensation and aromatization. This method can be hampered by the formation of regioisomers if the *ortho*-phenylenediamine contains an unsymmetrically substituted aromatic ring.

Recent SAR reports on phenazines from the groups of Cushman (Scheme 10)⁸³ and Huigens III¹⁷ have exploited this synthetic strategy extensively to construct 7,8-disubstituted PHz analogues.

Scheme 10: Synthesis of 7,8-dimethyl-1-methoxyphenazine (45) by Conda-Sheridan *et al.* in 2010.⁸³

3.3.5 Cyclizations of *O*-nitrodiphenylamines

Reductive cyclization of *ortho*-nitrodiphenylamines (Holliman cyclization) provides another route towards phenazines and has probably served as the most common method of constructing the scaffold to date. With an *ortho*-nitrodiphenylamine in hand, the transformation can be undertaken in a one-pot procedure with the aid of an alkali base and a reductive agent such as NaBH₄. The main limitation of this approach is that the cyclization can occur from both positions *ortho* to the aniline-nitrogen. This can be avoided however by appropriate substitutions, exemplified in Holliman's synthesis of the natural product Griseoluteic acid (**49**) where the methoxy group of **48** eliminates the possibility of an electrophilic attack on the nitro group from two *ortho*-positions simultaneously (Scheme 11).⁸⁴

Scheme 11: The synthesis of Griseoluteic acid (**49**) by Holliman and co-workers in 1970.⁸⁴

The reaction mechanism for the reductive base-catalyzed cyclization is depicted in Scheme 12. The substrates for the cyclization reaction (2-nitrodiphenylamines) can typically be synthesized via nucleophilic aromatic substitution (NAS) on *ortho*-halonitrobenzenes by anilines⁸⁴ or via Jourdan-Ullman Cu(I)-catalyzed couplings. Yet another alternative is the more modern Buchwald-Hartwig *C-N* cross-coupling reaction (Pd-catalyzed) reviewed in more detail in Section 3.3.6.

Scheme 12: General reaction mechanism for reductive cyclization of *ortho*-nitrodiphenylamines.

3.3.6 Buchwald-Hartwig C-N cross-coupling

The Buchwald-Hartwig cross-coupling reaction is a palladium-catalyzed reaction that affords aryl amines from aryl halides and amines (primary or secondary). The breakthrough prototype of this reaction was reported in 1983 by Migita *et al.*, as depicted in Scheme 13.⁸⁵ The name of the reaction, however, stems from the groups of Buchwald and Hartwig that individually explored the utility and scope of this reaction more extensively, starting in the 1990s.⁸⁶⁻⁸⁷

Scheme 13: A prototype of the reaction that later became known as the Buchwald-Hartwig C-N cross-coupling, reported by Migita *et al.* in 1983.⁸⁵

The reaction has been heavily investigated over the past two decades in which several generations of phosphine ligands and pre-catalysts have been developed to broaden the scope of the reaction. Moreover, user guides have been published and systems have been developed in an attempt to generalize the procedure and make it more “user-friendly”, for example by developing air-stable catalysts.⁸⁸ Nevertheless, labor-intensive screenings of ligands, bases, pre-catalysts and solvents may still be needed to achieve a satisfactory outcome. The Buchwald-Hartwig C-N cross coupling thus offers a new, exciting path towards substituted phenazines and the method is characterized by milder conditions and a higher selectivity than described for the above mentioned classic routes. A simplified mechanism for the catalytic cycle of the reaction is given in Scheme 14. A Pd(0)-ligand complex undergoes oxidative addition with an aryl halide in the first step. Thereafter, a nucleophilic amine coordinates to the Palladium and the metal amide is deprotonated by a base (preferably a negatively charged base, e.g. NaOtBu).⁸⁹ Finally, a reductive elimination yields an arylated amine product, along with the regenerated palladium catalyst.

Scheme 14: A simplified catalytic cycle for the Buchwald-Hartwig *C-N* cross-coupling.

Emoto and colleagues were the first to publish the synthesis of a phenazine scaffold using this approach. Their synthesis involved three steps, first an intermolecular Buchwald-Hartwig reaction, then a reduction of the nitro group and finally the second Pd-catalyzed ring-closing step (Scheme 15).⁹⁰

Scheme 15: A New route to phenazines by Emoto et al. in 2000.⁹⁰

Winkler *et al.* further demonstrated the potential of this reaction by condensing functionalized 2-bromoanilines into phenazines in a one-pot fashion (Scheme 16, top).⁹¹ In a similar manner, Laha and colleagues reported a double Buchwald-Hartwig *C-N* cross-coupling that involved the condensation of *o*-phenylenediamines and *o*-phenylenedibromides (Scheme 16, bottom).⁹²

Scheme 16: Synthesis of 2,7-dimethoxyphenazine and 2-methoxyphenazine via Pd-catalyzed Buchwald-Hartwig C-N cross-coupling.⁹¹⁻⁹²

3.3.7 Miscellaneous modern methods

Over the course of the last decade, several new “exotic” synthetic routes to phenazines have been introduced, clearly demonstrating that this research field is alive and that phenazines are molecules of interest to chemists. A review has recently been published by Guttenberger *et al.*⁵ and the detailed coverage provided therein goes beyond the scope of this introduction.⁵ Selected examples are however depicted in Scheme 17. Lian *et al.* reported a Rh(III)-catalyzed method reacting diphenyl diazenes and aryl azides,⁹³ Yu and colleagues showed how phenazines can be synthesized in water using a Cu-catalyzed homocoupling of 2-halo-anilines,⁹⁴ and Laha *et al.* explored a novel procedure converting 10,11-dihydro-5H-dibenzo[b,e][1,4]diazepines to phenazines through a radical cascade using potassium persulfate.⁹⁵

Scheme 17: Selected overview of modern phenazine-yielding synthetic transformations.

3.4 References

1. Clemo, G. R.; Daglish, A. F., 307. The phenazine series. Part VIII. The constitution of the pigment of *Chromobacterium iodinum*. *J. Chem. Soc.* **1950**, (0), 1481-1485.
2. Wilson, R.; Sykes, D. A.; Watson, D.; Rutman, A.; Taylor, G. W.; Cole, P. J., Measurement of *Pseudomonas aeruginosa* phenazine pigments in sputum and assessment of their contribution to sputum sol toxicity for respiratory epithelium. *Infect. Immun.* **1988**, 56 (9), 2515-2517.
3. Abken, H.-J.; Tietze, M.; Brodersen, J.; Bäumer, S.; Beifuss, U.; Deppenmeier, U., Isolation and Characterization of Methanophenazine and Function of Phenazines in Membrane-Bound Electron Transport of *Methanosarcina mazei* Gö1. *J. Bacteriol.* **1998**, 180 (8), 2027-2032.
4. Laursen, J. B.; Nielsen, J., Phenazine Natural Products: Biosynthesis, Synthetic Analogues, and Biological Activity. *Chem. Rev.* **2004**, 104 (3), 1663-1686.
5. Guttenberger, N.; Blankenfeldt, W.; Breinbauer, R., Recent developments in the isolation, biological function, biosynthesis, and synthesis of phenazine natural products. *Bioorg. Med. Chem.* **2017**, (in press).
6. Turner, J. M.; Messenger, A. J., Occurrence, biochemistry and physiology of phenazine pigment production. *Adv. Microb. Physiol.* **1986**, 27, 211-75.
7. Mavrodi, D. V.; Parejko, J. A.; Mavrodi, O. V.; Kwak, Y.-S.; Weller, D. M.; Blankenfeldt, W.; Thomashow, L. S., Recent insights into the diversity, frequency and ecological roles of phenazines in fluorescent *Pseudomonas* spp. *Environ. Microbiol.* **2013**, 15 (3), 675-686.
8. Shin-ya, K.; Furihata, K.; Hayakawa, Y.; Seto, H.; Kato, Y.; Clardy, J., The structure of benthocyanin A. A new free radical scavenger of microbial origin. *Tetrahedron Lett.* **1991**, 32 (7), 943-946.
9. Mavrodi, D. V.; Peever, T. L.; Mavrodi, O. V.; Parejko, J. A.; Raaijmakers, J. M.; Lemanceau, P.; Mazurier, S.; Heide, L.; Blankenfeldt, W.; Weller, D. M.; Thomashow, L. S., Diversity and evolution of the phenazine biosynthesis pathway. *Appl. Environ. Microbiol.* **2010**, 76 (3), 866-79.
10. Blankenfeldt, W.; Parsons, J. F., The structural biology of phenazine biosynthesis. *Curr. Opin. Struct. Biol.* **2014**, 29, 26-33.
11. Pierson, L. S.; Pierson, E. A., Metabolism and function of phenazines in bacteria: impacts on the behavior of bacteria in the environment and biotechnological processes. *Appl. Microbiol. Biotechnol.* **2010**, 86 (6), 1659-1670.
12. Weigele, M.; Maestroni, G.; Mitrovic, M.; Leimgruber, W., Antimicrobial agents structurally related to myxin. *Antimicrob. Agents. Chemother.* **1970**, 10, 46-9.
13. Behki, R. M.; Lesley, S. M., Deoxyribonucleic Acid Degradation and the Lethal Effect by Myxin in *Escherichia coli*. *J. Bacteriol.* **1972**, 109 (1), 250-261.
14. Myhren, L. E.; Nygaard, G.; Gausdal, G.; Sletta, H.; Teigen, K.; Degnes, K. F.; Zahlsen, K.; Brunsvik, A.; Bruserud, O.; Doskeland, S. O.; Selheim, F.; Herfindal, L., Iodinin (1,6-dihydroxyphenazine 5,10-dioxide) from *Streptosporangium* sp. induces apoptosis selectively in myeloid leukemia cell lines and patient cells. *Mar. Drugs* **2013**, 11 (2), 332-49.
15. Hollstein, U.; Van Gemert, R. J., Interaction of phenazines with polydeoxyribonucleotides. *Biochemistry* **1971**, 10 (3), 497-504.
16. Cimmino, A.; Evidente, A.; Mathieu, V.; Andolfi, A.; Lefranc, F.; Kornienko, A.; Kiss, R., Phenazines and cancer. *Nat. Prod. Rep.* **2012**, 29 (4), 487-501.
17. Garrison, A. T.; Abouelhassan, Y.; Kallifidas, D.; Bai, F.; Ukhanova, M.; Mai, V.; Jin, S.; Luesch, H.; Huigens, R. W., Halogenated Phenazines that Potently Eradicate Biofilms, MRSA Persister Cells in Non-Biofilm Cultures, and *Mycobacterium tuberculosis*. *Angew. Chem. Int. Ed.* **2015**, 54 (49), 14819-14823.
18. Garrison, A. T.; Abouelhassan, Y.; Norwood, V. M.; Kallifidas, D.; Bai, F.; Nguyen, M. T.; Rolfe, M.; Burch, G. M.; Jin, S.; Luesch, H.; Huigens, R. W., Structure-Activity Relationships of a Diverse Class of Halogenated Phenazines That Targets Persistent, Antibiotic-Tolerant Bacterial Biofilms and *Mycobacterium tuberculosis*. *J. Med. Chem.* **2016**, 59 (8), 3808-3825.

19. Kondratyuk, T. P.; Park, E.-J.; Yu, R.; van Breemen, R. B.; Asolkar, R. N.; Murphy, B. T.; Fenical, W.; Pezzuto, J. M., Novel Marine Phenazines as Potential Cancer Chemopreventive and Anti-Inflammatory Agents. *Mar. Drugs* **2012**, *10* (2), 451.
20. Zhao, J.; Wu, Y.; Alfred, A. T.; Wei, P.; Yang, S., Anticancer effects of pyocyanin on HepG2 human hepatoma cells. *Lett. Appl Microbiol.* **2014**, *58* (6), 541-548.
21. Zhuo, S.-T.; Li, C.-Y.; Hu, M.-H.; Chen, S.-B.; Yao, P.-F.; Huang, S.-L.; Ou, T.-M.; Tan, J.-H.; An, L.-K.; Li, D.; Gu, L.-Q.; Huang, Z.-S., Synthesis and biological evaluation of benzo[a]phenazine derivatives as a dual inhibitor of topoisomerase I and II. *Org. Biomol. Chem.* **2013**, *11* (24), 3989-4005.
22. Sinha, S.; Shen, X.; Gallazzi, F.; Li, Q.; Zmijewski, J. W.; Lancaster, J. R.; Gates, K. S., Generation of Reactive Oxygen Species Mediated by 1-Hydroxyphenazine, a Virulence Factor of *Pseudomonas aeruginosa*. *Chem. Res. Toxicol.* **2015**, *28* (2), 175-181.
23. Hassan, H. M.; Fridovich, I., Mechanism of the antibiotic action pyocyanine. *J. Bacteriol.* **1980**, *141* (1), 156-163.
24. Fraser, R. S.; Creanor, J., The mechanism of inhibition of ribonucleic acid synthesis by 8-hydroxyquinoline and the antibiotic lomofungin. *Biochemical Journal* **1975**, *147* (3), 401-410.
25. Price-Whelan, A.; Dietrich, L. E. P.; Newman, D. K., Rethinking 'secondary' metabolism: physiological roles for phenazine antibiotics. *Nat. Chem. Biol.* **2006**, *2* (2), 71-78.
26. Lau, G. W.; Hassett, D. J.; Ran, H.; Kong, F., The role of pyocyanin in *Pseudomonas aeruginosa* infection. *Trends Mol. Med.* **2004**, *10* (12), 599-606.
27. Lau, G. W.; Hassett, D. J.; Britigan, B. E., Modulation of lung epithelial functions by *Pseudomonas aeruginosa*. *Trends Microbiol.* **2005**, *13* (8), 389-397.
28. Usher, L. R.; Lawson, R. A.; Geary, I.; Taylor, C. J.; Bingle, C. D.; Taylor, G. W.; Whyte, M. K. B., Induction of Neutrophil Apoptosis by the *Pseudomonas aeruginosa* Exotoxin Pyocyanin: A Potential Mechanism of Persistent Infection. *J. Immunol.* **2002**, *168* (4), 1861-1868.
29. Lyczak, J. B.; Cannon, C. L.; Pier, G. B., Establishment of *Pseudomonas aeruginosa* infection: lessons from a versatile opportunist. *Microbes Infect.* **2000**, *2* (9), 1051-1060.
30. Lau, G. W.; Ran, H.; Kong, F.; Hassett, D. J.; Mavrodi, D., *Pseudomonas aeruginosa* Pyocyanin Is Critical for Lung Infection in Mice. *Infect. Immun.* **2004**, *72* (7), 4275-4278.
31. Britigan, B. E.; Roeder, T. L.; Rasmussen, G. T.; Shasby, D. M.; McCormick, M. L.; Cox, C. D., Interaction of the *Pseudomonas aeruginosa* secretory products pyocyanin and pyochelin generates hydroxyl radical and causes synergistic damage to endothelial cells. Implications for *Pseudomonas*-associated tissue injury. *J. Clin. Invest.* **1992**, *90* (6), 2187-2196.
32. Ballatori, N.; Krance Suzanne, M.; Notenboom, S.; Shi, S.; Tieu, K.; Hammond Christine, L., Glutathione dysregulation and the etiology and progression of human diseases. In *Biol. Chem.*, **2009**; Vol. 390, p 191.
33. Baron, S. S.; Rowe, J. J., Antibiotic action of pyocyanin. *Antimicrob. Agents Chemother.* **1981**, *20* (6), 814-820.
34. Johnson, L. E.; Dietz, A., Lomofungin, a New Antibiotic Produced by *Streptomyces lomondensis* sp. n. *Appl. Microbiol.* **1969**, *17* (5), 755-759.
35. Bergy, M. E.; Johnson, L. R. E., Antibiotic lomondomycin and method of production. Google Patents: **1967**.
36. Kuo, S. C.; Cano, F. R.; Lampen, J. O., Lomofungin, an Inhibitor of Ribonucleic Acid Synthesis in Yeast Protoplasts: Its Effect on Enzyme Formation. *Antimicrob. Agents Chemother.* **1973**, *3* (6), 716-722.
37. Cannon, M.; Davies, J. E.; Jimenez, A., Inhibition by lomofungin of nucleic acid and protein synthesis in *Saccharomyces cerevisiae*. *FEBS Lett.* **1973**, *32* (2), 277-80.
38. Prachayasittikul, V.; Prachayasittikul, S.; Ruchirawat, S.; Prachayasittikul, V., 8-Hydroxyquinolines: a review of their metal chelating properties and medicinal applications. *Drug Des. Dev. Ther.* **2013**, *7*, 1157-1178.
39. Fraser, R. S.; Creanor, J., The mechanism of inhibition of ribonucleic acid synthesis by 8-hydroxyquinoline and the antibiotic lomofungin. *Biochem. J.* **1975**, *147* (3), 401.
40. Pavletich, K.; Kuo, S. C.; Lampen, J. O., Chelation of divalent cations by lomofungin: Role in inhibition of nucleic acid synthesis. *Biochem. Biophys. Res. Commun.* **1974**, *60* (3), 942-950.

41. Hoskins, J. W.; Ofori, L. O.; Chen, C. Z.; Kumar, A.; Sobczak, K.; Nakamori, M.; Southall, N.; Patnaik, S.; Marugan, J. J.; Zheng, W.; Austin, C. P.; Disney, M. D.; Miller, B. L.; Thornton, C. A., Lomofungin and dilomofungin: inhibitors of MBNL1-CUG RNA binding with distinct cellular effects. *Nucleic Acids Res.* **2014**, *42* (10), 6591-6602.
42. Davis, J. G., *Chromobacterium iodinum* (n. sp.). *Zentralbl. Bakteriolog., Parasitenkd., Infektionskrankh. Hyg., Abt. 2, Naturwiss. Allg., Landwirtsch. Tech. Mikrobiol.* **1939**, *100* (II Abt.), 273-6.
43. Zhao, Y.; Qian, G.; Ye, Y.; Wright, S.; Chen, H.; Shen, Y.; Liu, F.; Du, L., Heterocyclic Aromatic N-Oxidation in the Biosynthesis of Phenazine Antibiotics from *Lysobacter antibioticus*. *Org. Lett.* **2016**, *18* (10), 2495-2498.
44. McIlwain, H., The anti-streptococcal action of iodinin. Naphthaquinones and anthraquinones as its main natural antagonists. *Biochem. J.* **1943**, *37* (2), 265-71.
45. Tanabe, I.; Iio, N.; Imanura, R.; Matsumoto, T.; Obayashi, A., Studies on the antibiotic action of Iodinin, and Its reduced from 1,6-phenazinediol. *Memories of the Faculty of Agriculture, Kagoshima University* **1976**, *10*, 127-125.
46. Hano, K.; Iwata, H.; Nakajima, K., Studies on the Anticancer Activity of Phenazine Derivatives. *Chem. Pharm. Bull.* **1965**, *13* (2), 107-113.
47. Oftedal, L.; Selheim, F.; Wahlsten, M.; Sivonen, K.; Doskeland, S. O.; Herfindal, L., Marine benthic cyanobacteria contain apoptosis-inducing activity synergizing with daunorubicin to kill leukemia cells, but not cardiomyocytes. *Mar. Drugs* **2010**, *8* (10), 2659-72.
48. Sletta, H.; Degnes, K. F.; Herfindal, L.; Klinkenberg, G.; Fjaervik, E.; Zahlsen, K.; Brunsvik, A.; Nygaard, G.; Aachmann, F. L.; Ellingsen, T. E.; Doskeland, S. O.; Zotchev, S. B., Anti-microbial and cytotoxic 1,6-dihydroxyphenazine-5,10-dioxide (iodinin) produced by *Streptosporangium* sp. DSM 45942 isolated from the fjord sediment. *Appl. Microbiol. Biotechnol.* **2014**, *98* (2), 603-10.
49. Peterson, E. A.; Gillespie, D. C.; Cook, F. D., A wide-spectrum antibiotic produced by a species of sorangium. *Can. J. Microbiol.* **1966**, *12* (2), 221-230.
50. Hanson, A. W.; Huml, K., Crystal structure of iodinin. *Acta Crystallogr., Sect. B* **1969**, *25* (9), 1766-74.
51. Weigle, M.; Leimgruber, W., The structure of myxin. *Tetrahedron Lett.* **1967**, *8*, 715-8.
52. Behki, R. M.; Lesley, S. M., Deoxyribonucleic acid degradation and the lethal effect by myxin in *Escherichia coli*. *J. Bacteriol.* **1972**, *109* (1), 250-61.
53. Lesley, S. M.; Behki, R. M., Mode of Action of Myxin on *Escherichia coli*. *J. Bacteriol.* **1967**, *94* (6), 1837-1845.
54. Chowdhury, G.; Sarkar, U.; Pullen, S.; Wilson, W. R.; Rajapakse, A.; Fuchs-Knotts, T.; Gates, K. S., DNA Strand Cleavage by the Phenazine Di-N-oxide Natural Product Myxin under Both Aerobic and Anaerobic Conditions. *Chem. Res. Toxicol.* **2012**, *25* (1), 197-206.
55. Hollstein, U.; Butler, P. L., Inhibition of ribonucleic acid synthesis by myxin. *Biochemistry* **1972**, *11* (8), 1345-1350.
56. Grunberg, E.; Berger, J.; Beskid, G.; Cleeland, R.; Prince, H. N.; Titsworth, E., Studies on the in vitro and in vivo Chemotherapeutic Properties of the Antibiotic Myxin. *Chemotherapy* **1967**, *12* (5), 272-281.
57. Manius, G.; Tscherne, R.; Venteicher, R.; Secker, A., High-performance liquid chromatographic determination of free myxin and its reduction product as impurities in cuprimyxin-containing creams. *J. Pharm. Sci.* **1981**, *70* (9), 1024-1026.
58. Junnotula, V.; Sarkar, U.; Sinha, S.; Gates, K. S., Initiation of DNA Strand Cleavage by 1,2,4-Benzotriazine 1,4-Dioxide Antitumor Agents: Mechanistic Insight from Studies of 3-Methyl-1,2,4-benzotriazine 1,4-Dioxide. *J. Am. Chem. Soc.* **2009**, *131* (3), 1015-1024.
59. Balasubramanian, B.; Pogozelski, W. K.; Tullius, T. D., DNA strand breaking by the hydroxyl radical is governed by the accessible surface areas of the hydrogen atoms of the DNA backbone. *PNAS* **1998**, *95* (17), 9738-9743.
60. Chowdhury, G.; Sarkar, U.; Pullen, S.; Wilson, W. R.; Rajapakse, A.; Fuchs-Knotts, T.; Gates, K. S., DNA Strand Cleavage by the Phenazine Di-N-oxide Natural Product Myxin under Both Aerobic and Anaerobic Conditions. *Chem. Res. Toxicol.* **2011**, *25* (1), 197-206.

61. Brown, J. M., Tumor Hypoxia in Cancer Therapy. In *Methods Enzymol.*, Academic Press: **2007**; Vol. Volume 435, pp 295-321.
62. Padhani, A. R.; Krohn, K. A.; Lewis, J. S.; Alber, M., Imaging oxygenation of human tumours. *Eur. Radiol.* **2007**, *17* (4), 861-872.
63. Lavaggi, M. L.; Nieves, M.; Cabrera, M.; Olea-Azar, C.; López de Cerain, A.; Monge, A.; Cerecetto, H.; González, M., Structural modifications on the phenazine *N,N'*-dioxide-scaffold looking for new selective hypoxic cytotoxins. *Eur. J. Med. Chem.* **2010**, *45* (11), 5362-5369.
64. Junnotula, V.; Rajapakse, A.; Arbillaga, L.; López de Cerain, A.; Solano, B.; Villar, R.; Monge, A.; Gates, K. S., DNA strand cleaving properties and hypoxia-selective cytotoxicity of 7-chloro-2-thienylcarbonyl-3-trifluoromethylquinoxaline 1,4-dioxide. *Bioorg. Med. Chem.* **2010**, *18* (9), 3125-3132.
65. Hay, M. P.; Gamage, S. A.; Kovacs, M. S.; Pruijn, F. B.; Anderson, R. F.; Patterson, A. V.; Wilson, W. R.; Brown, J. M.; Denny, W. A., Structure–Activity Relationships of 1,2,4-Benzotriazine 1,4-Dioxides as Hypoxia-Selective Analogues of Tirapazamine. *J. Med. Chem.* **2003**, *46* (1), 169-182.
66. von Pawel, J.; von Roemeling, R.; Gatzemeier, U.; Boyer, M.; Elisson, L. O.; Clark, P.; Talbot, D.; Rey, A.; Butler, T. W.; Hirsh, V.; Olver, I.; Bergman, B.; Ayoub, J.; Richardson, G.; Dunlop, D.; Arcenas, A.; Vescio, R.; Viallet, J.; Treat, J., Tirapazamine Plus Cisplatin Versus Cisplatin in Advanced Non–Small-Cell Lung Cancer: A Report of the International CATAPULT I Study Group. *J. Clin. Oncol.* **2000**, *18* (6), 1351-1359.
67. Laderoute, K.; Wardman, P.; Rauth, A. M., Molecular mechanisms for the hypoxiaindependent activation of 3-amino-1,2,4-benzotriazine-1, 4-dioxide (SR 4233). *Biochem. Pharmacol.* **1988**, *37* (8), 1487-1495.
68. Silva, J. M.; O'Brien, P. J., Molecular mechanisms of SR 4233-induced hepatocyte toxicity under aerobic versus hypoxic conditions. *Br. J. Cancer* **1993**, *68* (3), 484-491.
69. Peters, K. B.; Brown, J. M., Tirapazamine. *A Hypoxia-activated Topoisomerase II Poison* **2002**, *62* (18), 5248-5253.
70. Brown, J. M.; Wilson, W. R., Exploiting tumour hypoxia in cancer treatment. *Nat. Rev. Cancer* **2004**, *4* (6), 437-447.
71. Chowdhury, G.; Junnotula, V.; Daniels, J. S.; Greenberg, M. M.; Gates, K. S., DNA Strand Damage Product Analysis Provides Evidence That the Tumor Cell-Specific Cytotoxin Tirapazamine Produces Hydroxyl Radical and Acts as a Surrogate for O₂. *J. Am. Chem. Soc.* **2007**, *129* (42), 12870-12877.
72. Cerecetto, H.; González, M.; Lavaggi, M. L.; Azqueta, A.; López de Cerain, A.; Monge, A., Phenazine 5,10-Dioxide Derivatives as Hypoxic Selective Cytotoxins. *J. Med. Chem.* **2005**, *48* (1), 21-23.
73. Lavaggi, M. L.; Cabrera, M.; González, M.; Cerecetto, H., Differential Enzymatic Reductions Governing the Differential Hypoxia-Selective Cytotoxicities of Phenazine 5,10-Dioxides. *Chem. Res. Toxicol.* **2008**, *21* (9), 1900-1906.
74. Cerecetto, H.; Gonzalez, M.; Lavaggi, M. L.; Aravena, M. A.; Rigol, C.; Olea-Azar, C.; Azqueta, A.; Cerain, A. L. d.; Monge, A.; Bruno, A. M., Phenazine 5,10-Dioxide Derivatives as Hypoxic Selective Cytotoxins: Part II. Structure-Activity Relationship Studies. *Med. Chem.* **2006**, *2* (5), 511-521.
75. Lavaggi, M.; Cabrera, M.; Pintos, C.; Arredondo, C.; Pachon, G.; Rodriguez, J.; Raymondo, S.; Pacheco, J.; Cascante, M.; Olea-Azar, C.; Cerain, A.; Monge, A.; Cerecetto, H.; Gonzalez, M., Novel Phenazine 5,10-Dioxides Release •OH in Simulated Hypoxia and Induce Reduction of Tumour Volume In Vivo. *ISRN Pharmacol.* **2011**, *2011*, 11.
76. Wohl, A.; Aue, W., Ueber die Einwirkung von Nitrobenzol auf Anilin bei Gegenwart von Alkali. *Ber. Dtsch. Chem. Ges.* **1901**, *34* (2), 2442-2450.
77. Pachter, I. J.; Kloetzel, M. C., The Wohl-Aue Reaction. I. Structure of Benzo [a] phenazine Oxides and Syntheses of 1,6-Dimethoxyphenazine and 1,6-Dichlorophenazine. *J. Am. Chem. Soc.* **1951**, *73* (10), 4958-4961.
78. Yang, H.; Abouelhassan, Y.; Burch, G. M.; Kallifidas, D.; Huang, G.; Yousaf, H.; Jin, S.; Luesch, H.; Huigens, R. W., A Highly Potent Class of Halogenated Phenazine Antibacterial and Biofilm-

- Eradicating Agents Accessed Through a Modular Wohl-Aue Synthesis. *Sci. Rep.* **2017**, 7 (1), 2003.
79. Bamberger, E.; Ham, W., Behavior of p-Substituted Nitrosobenzenes towards Concentrated Sulfuric Acid. *Liebigs Ann. Chem.* **1911**, 382, 82-128.
 80. Sawhney, S. N.; Boykin, D. W., A reinvestigation of the bamberger-ham phenazine. Synthesis: A limitation. *J. Heterocycl. Chem.* **1979**, 16 (2), 397-400.
 81. El-Haj, M. J. A.; Dominy, B. W.; Johnston, J. D.; Haddadin, M. J.; Issidorides, C. H., New route to phenazine 5,10-dioxides and related compounds. *J. Org. Chem.* **1972**, 37 (4), 589-93.
 82. Issidorides, C. H.; Atfah, M. A.; Sabounji, J. J.; Sidani, A. R.; Haddadin, M. J., Application of 1,2-diketones in the synthesis of phenazine oxides. *Tetrahedron* **1978**, 34 (2), 217-221.
 83. Conda-Sheridan, M.; Marler, L.; Park, E.-J.; Kondratyuk, T. P.; Jermihov, K.; Mesecar, A. D.; Pezzuto, J. M.; Asolkar, R. N.; Fenical, W.; Cushman, M., Potential Chemopreventive Agents Based on the Structure of the Lead Compound 2-Bromo-1-hydroxyphenazine, Isolated from *Streptomyces* Species, Strain CNS284. *J. Med. Chem.* **2010**, 53 (24), 8688-8699.
 84. Challand, S. R.; Herbert, R. B.; Holliman, F. G., A new phenazine synthesis. The synthesis of griseoluteic acid, griseolutein A, and methyl diacetylgriseolutein B. *J. Chem. Soc. D* **1970**, (21), 1423-1425.
 85. Kosugi, M.; Kameyama, M.; Migita, T., Palladium-Catalyzed Aromatic Amination of Aryl Bromides with N,N-Diethylamino-Tributyltin. *Chem. Lett.* **1983**, (6), 927-928.
 86. Guram, A. S.; Rennels, R. A.; Buchwald, S. L., A Simple Catalytic Method for the Conversion of Aryl Bromides to Arylamines. *Angew. Chem. Int. Ed.* **1995**, 34 (12), 1348-1350.
 87. Paul, F.; Patt, J.; Hartwig, J. F., Palladium-catalyzed formation of carbon-nitrogen bonds. Reaction intermediates and catalyst improvements in the hetero cross-coupling of aryl halides and tin amides. *J. Am. Chem. Soc.* **1994**, 116 (13), 5969-5970.
 88. Surry, D. S.; Buchwald, S. L., Dialkylbiaryl Phosphines in Pd-Catalyzed Amination: A User's Guide. *Chem. sci. (Royal Society of Chemistry : 2010)* **2011**, 2 (1), 27-50.
 89. Sunesson, Y.; Limé, E.; Nilsson Lill, S. O.; Meadows, R. E.; Norrby, P.-O., Role of the Base in Buchwald-Hartwig Amination. *J. Org. Chem.* **2014**, 79 (24), 11961-11969.
 90. Emoto, T.; Kubosaki, N.; Yamagiwa, Y.; Kamikawa, T., A new route to phenazines†. *Tetrahedron Lett.* **2000**, 41 (3), 355-358.
 91. Winkler, J. D.; Twenter, B. M.; Gendrineau, T., A new approach to the synthesis of substituted phenazines via palladium-catalyzed aryl ligation. *Heterocycles* **2012**, 84 (2), 10.3987/COM-11-S(P)76.
 92. Laha, J. K.; Tummalapalli, K. S. S.; Gupta, A., Palladium-Catalyzed Domino Double *N*-Arylations (Inter- and Intramolecular) of 1,2-Diamino(hetero)arenes with o,o'-Dihalo(hetero)arenes for the Synthesis of Phenazines and Pyridoquinoxalines. *Eur. J. Org. Chem.* **2013**, 2013 (36), 8330-8335.
 93. Lian, Y.; Hummel, J. R.; Bergman, R. G.; Ellman, J. A., Facile Synthesis of Unsymmetrical Acridines and Phenazines by a Rh(III)-Catalyzed Amination/Cyclization/Aromatization Cascade. *J. Am. Chem. Soc.* **2013**, 135 (34), 12548-12551.
 94. Yu, L.; Zhou, X.; Wu, D.; Xiang, H., Synthesis of phenazines by Cu-catalyzed homocoupling of 2-halogen anilines in water. *J. Organomet. Chem.* **2012**, 705, 75-78.
 95. Laha, J. K.; Tummalapalli, K. S. S.; Gupta, A., Transition-Metal-Free Tandem Oxidative Removal of Benzylic Methylene Group by C-C and C-N Bond Cleavage Followed by Intramolecular New Aryl C-N Bond Formation under Radical Conditions. *Org. Lett.* **2014**, 16 (17), 4392-4395.

Chapter 4 – Phenazines and aim of study

Prior to this work, iodinin (**5**) isolated from marine *actinomycetes*, has shown promise by inhibiting the growth of cancerous human- and rat AML cell-lines *in vitro*.¹⁻² Iodinin was thus considered a lead candidate and an attractive SAR-scaffold from the start of the presented work. However, the natural product iodinin (**5**) itself, like many other compounds discovered in nature, is not fit to serve as a drug. Part of the reason for this is the very poor solubility of **5** in general. In addition, neither commercial sources of iodinin (**5**), nor efficient synthetic routes were available. Thus, the aims of this project can be summarized as follows;

- To establish a robust synthetic methodology for the preparation of iodinin (**5**), ensuring that gram-scale quantities could be made available to further lead optimization towards novel antileukemic drugs.
- Synthesize and test myxin (**8**) *in vitro*, due to its close structural similarity to iodinin (**5**). The antileukemic activity of myxin (**8**) was entirely unknown from the start.
- Explore the structure-activity relationships of the hydroxylated phenazine 5,10-dioxide scaffold, involving various substitution patterns in positions 6, 7 and 8, to see how changes in the scaffold would affect potency against- and selectivity among human leukemic cell lines.
- Evaluate the antibacterial potential of the synthesized compounds in Gram-positive and Gram-negative bacteria resistant to various types of antibiotics.

Figure 16. Graphical representation of the aims of study.

Chapter 5 – Phenazines results and discussion

Paper I and paper II (a prepared manuscript) are discussed and summarized in the following sections with some additional results from antimicrobial evaluations of selected compounds.

5.1 Paper I – Synthesis and biological evaluations of iodinin, myxin and alkylated derivatives

5.1.1 Synthesis

The first crucial aim was to establish a reliable synthetic route to iodinin (**5**) as this was considered mandatory for further biological evaluation and lead development employing SAR-studies.

To construct the 1,6-*O*-disubstituted phenazine scaffold, there are strikingly few methods available if (i) reasonable yields, and (ii) if formation of inseparable regioisomers are to be avoided (see section 3.3, page 32). 1,6-Dimethoxyphenazine (**39**) was chosen as target-precursor for iodinin (**5**) since the *O*-methyl functionalities can serve as phenol protecting groups under alkaline conditions applied to construct the three-ring scaffold.

First, a double Buchwald-Hartwig *C-N* cross coupling was explored with the aim to self-condense 2-bromo-3-methoxyaniline (**58**, Scheme 18, left). This was achieved using the Pd(II)-Brettphos precatalyst (3 mol%) in presence of KHMDS (3 mol%) and Cs₂CO₃ in refluxing toluene. Similar work had already been performed by Winkler et al. in 2012,³ for example on 2,7-dimethoxyphenazine (**54**, see Scheme 16, page 39), but no attempts to synthesize 1,6-dioxy-substituted phenazines using palladium chemistry were known in prior art.

Scheme 18: Two methods were explored to synthesize 1,6-dimethoxyphenazine (**39**)

The best entry gave 79% yield on a single gram scale. Attempts to scale up the reaction resulted in lowered yields. Because of the reaction mechanism, two Br-atoms are expelled as byproducts (see Scheme 14, page 38) and this results in a considerable weight loss of material. Therefore, the 79% yield only stands for 480 mg of **39** from a single gram of **58**. This was not considered enough material as we wanted to provide gram quantities of iodinin (**5**) and in order to ensure that, we turned our attention to the old Wohl-Aue approach as a second option.⁴ This procedure involves the condensation of *o*-anisidine (**59**) and 2-nitroanisole (**60**) under harsh alkaline conditions. (Scheme 18, right). After tedious extractions and recrystallizations, 4.5g (6%) of yellow needles were pulled out of the crude black viscous tar.

In the following step, complete demethylation of **39** was achieved by exposure of an excess of BBr₃ (neat, reflux) (Scheme 19) in accordance with earlier reports from Alonso et al.⁵

The final step towards iodinin (**5**) was anticipated to be the most challenging one, a double *N*-oxidation of 1,6-dihydroxyphenazine (**28**). Early literature reports 21% yield utilizing H₂O₂ and acetic anhydride, thus forming peracetic acid *in situ*.⁶ This method did not prove to be reproducible in our hands. A considerable time and effort was spent on tuning in reaction conditions and entries are summarized in Table S-2 for paper 1 (supplementary information). Key bullet points summarizing the progress of this work are listed below:

- No entries based on H₂O₂, peracetic acid, *t*BuOOH or KMnO₄ proved successful
- Adding *m*CPBA (5 eq) in one portion in DCM at 40 °C gave 7-10% yield
- Adding *m*CPBA (5 eq) portion wise in DCE at 80 °C gave 21% yield
- Adding *m*CPBA (5-7 eq) portion wise in toluene at 80 °C over 6 hours gave 76% yield

Scheme 19: Synthesis of iodinin (**5**) from 1,6-dimethoxyphenazine (**39**)

Scheme 20: A plausible reaction mechanism for *m*CPBA oxidation of 1,6-dihydroxyphenazine (**28**).

A phenol functionality was observed to be more favorable than a methoxy group in terms of *N*-oxidation which about two main reasons were expected to cause. First, a methoxy group does not support the mechanism proposed in Scheme 21. Secondly, iodinin's (**5**) *N*-oxides are both stabilized by intramolecular ion-dipole/hydrogen forces, a feature that makes the compound more tolerant to elevated temperatures, which again proved necessary to force the oxidation forward in good yields. These suggestions are in line with Weigele and Leimgrubers findings that reported myxin (**8**), which only has the *N*-oxide in position 10 stabilized in this way, to fully decompose after 4h reflux in CHCl_3 .⁷ In further support of these suggestions, the oxidation of 1-hydroxy-6-methoxyphenazine (**60**) was also attempted in order to prepare myxin (**8**), yet only afforded the orange mono-oxidized analogue **61** and not myxin (**8**) (Scheme 20).

Scheme 21: An oxidation attempted on 1-hydroxyphenazine-6-methoxyphenazine (**60**) did not afford myxin (**8**).

The intramolecular ion-dipole/hydrogen bonds result in high lipophilicity, followed by tailing on silica that caused problems for chromatographic purifications, extractions and recrystallizations. Accordingly, a purification strategy was developed based on extensive washing of the crude product with solvents that do not solute iodinin (**5**), but dissolve *m*CPBA, *m*CBA and other unknown impurities. Eventually, this strategy gave purified iodinin in 76% yield and this result could be reproduced multiple times on a gram scale. The low solubility of iodinin (**5**) was confirmed by ^1H -NMR of **5** in saturated solution of $\text{DMSO}-d_6$ (see Figure 17, next page).

As expected, inevitable over-alkylated and deoxygenated products were formed during the reactions as well (see compounds **61-63**, Scheme 22). However, chromatographic isolation of these compounds was simple and all of them were of great interest for upcoming SAR-studies in cancerous cell lines, especially to generate data bringing insight into the mechanism of action related to the level of *N*-oxidation and *O*-alkylation.

Next, several alkylated derivatives were prepared as iodinin (**5**) was reacted with ethyl-bromoacetate affording the 1-hydroxy-6-(2-ethoxy-2-oxoethoxy)phenazine 5,10-dioxide (**64**) and the dialkylated analogue **65** (Scheme 23). This step was also repeated with *tert*-butyl bromoacetate which gave the more lipophilic 1-hydroxy-6-(2-(*tert*-butoxy)-2-oxoethoxy)phenazine 5,10-dioxide (**66**) and the corresponding di-functionalized analog (**67**). The *tert*-butyl esters **66** and **67** were thereafter individually treated with H₃PO₄ (85% aq.), a mild approach¹⁰ that gave carboxymethoxy analogues **68** and **69** (Scheme 23) in decent yields.

Scheme 23: Alkylated analogues of iodinin (**5**). The X-ray structure of compound **64** was recorded by Dr. Sigurd Øien-Ødegaard.

5.1.2 Biological evaluations of iodinin, myxin and alkylated analogs

This section describes the biological results presented in paper I which solely were conducted on cancerous AML cells (MOLM-13, human). After the publication was accepted, biological studies of most central compounds were extended to cell lines derived from healthy tissues as well to address the cancer cell selectivity. Accordingly, these new results are included in the discussion below.

Prior to the biological evaluations described within this section, iodinin (**5**) was the only *N*-oxidized phenazine derivative, isolated from fermentation broths, that had been evaluated by the collaborating group of Herfindal/Døskeland in Bergen, Norway.¹ We were therefore eager to see what effects myxin (**5**) and the other prepared derivatives would display *in vitro*.

Synthesized iodinin (**5**) was evaluated where apoptotic cell death in human AML cells (MOLM-13) was assessed. The results from these experiments showed that iodinin (**5**) had an EC₅₀ value of 2.0 μM under aerobic conditions (19% O₂, see Table 1). The data were acquired employing WST-1 proliferation assay and confirmed by microscopic evaluation of cell death. In hypoxic environment (2% O₂) however, iodinin (**5**) was found to kill AML-cells more potently with an EC₅₀ value of 0.79 μM. Iodinin was also screened in cells from healthy tissues in order to address cancer cell selectivity, the NRK cell-line (*Rattus norvegicus*, rat, kidney) and H9c2 cells (*Rattus norvegicus*, rat heart myoblast). Exposure of **5** to these two cell lines did not show any toxic events below a concentration of 50 μM, further demonstrating a promising therapeutic window for this lead candidate.

As the focus was turned to myxin (**8**), potent cytotoxicity was observed and it was found to have low toxicity in healthy cell lines as well (Table 1). Thus, it showed a very comparable potency and selectivity profile to that of iodinin (**5**) (Table 1). In terms of cytotoxicity at hypoxic levels, myxin (**8**) was found to be more selective towards AML cells by a factor of ~100 when compared with the NRK cell line (Table 1). This finding was considered particularly important, since compared with iodinin (**5**), myxin (**8**) is lacking one intramolecular ion-dipole bond and this results in drastically improved solubility. Accordingly, the results presented within Table 1 strongly suggest that the physicochemical properties of the lead candidate iodinin (**5**) can be improved whereas cytotoxicity and selectivity remained unaltered. These findings, showing iodinin and myxin to be equally active according to the results, also strongly suggest that iodinin's (**5**) discharge of reactive oxygen species (ROS) is

Table 1: EC₅₀ values (±SEM) of phenazine-analogues on MOLM-13, NRK and H9c2 cells.

Compound	EC ₅₀ (μM) MOLM-13		EC ₅₀ (μM) NRK		EC ₅₀ (μM) H9c2
	19% O ₂	2% O ₂	19% O ₂	2% O ₂	19% O ₂
5 (iodinin)	2.0 ± 0.07	0.79 ± 0.10	>50	>50	>50
8 (myxin)	1.4 ± 0.30	0.77 ± 0.13	77 ± 11	77 ± 14	43 ± 2
28	100-120*	110-140*	n.d.	n.d.	n.d.
35 (TPZ)	95 ± 8*	22 ± 2*	>200	88 ± 2.4	107 ± 1
39	72 ± 14*	84 ± 6*	n.d.	n.d.	n.d.
60	>200	39 ± 9	n.d.	n.d.	n.d.
61	>200	22 ± 5	n.d.	n.d.	n.d.
62	4.0 ± 0.48	3.1 ± 0.83	n.d.	n.d.	n.d.
63	5.4 ± 0.68	4.1 ± 0.38	n.d.	n.d.	n.d.
64	0.57 ± 0.06	0.49 ± 0.12	11 ± 1.7	18 ± 1.3	15 ± 1
65	2.0 ± 0.47	0.79 ± 0.06	>20	>20	>20
66	2.9 ± 0.34	2.4 ± 0.16	n.d.	n.d.	n.d.
67	4.2 ± 0.20	3.3 ± 0.22	n.d.	n.d.	n.d.
68	54 ± 7	61 ± 4	n.d.	n.d.	n.d.
69	18 ± 3	39 ± 9	n.d.	n.d.	n.d.

Cell viability was assessed by the WST-1 proliferation assay and Hoechst staining 24 hrs after treatment of cells with various doses of iodinin (**3**) or its analogues. Regression analyses were performed using SigmaPlot in order to calculate the given EC₅₀ values. The EC₅₀ values of tirapazamine (**10**; TPZ) are based on microscopic evaluation (marked with *), whereas the EC₅₀ values of iodinin (**5**) and its analogues are based on the WST-1 results, and all are adjusted relative to untreated control cells in each experiment, *n* = 4-6 for MOLM-13 and *n* = 2 for NRK and H9c2 cells. n.d. = not determined

initiated by involvement of reducing agents or enzymes, as it is less likely that this is a spontaneous process, since the intrinsic stability of **5** would prohibit it (see stability discussion of iodinin versus myxin in 5.1.1).

Removal of one *N*-oxide from myxin's position 5 (**8**) leads to the structure of compound **61**. This simple change results in a drop in cytotoxic potency by a factor of more than one hundred (19% O₂, Table 1). Despite lacking comparable potency to myxin (**5**), **61** was found to be over nine-times more active under hypoxic environment and harmless even at 200 μM concentration (19% O₂). The myxin scaffold devoid of any *N*-oxides (compound **60**, Scheme 21) behaved similarly. Interestingly, this compound **60** shows hypoxia selectivity although the compound lacks both *N*-oxides, and is therefore physically unable to discharge any cytotoxic ·OH radicals. Factors like redox cycling, iron chelation, Fenton-type reactions and depletion of antioxidants

like intracellular glutathione might contribute to this hypoxia selectivity, but this requires further investigation. Comparing the three molecules, myxin (**8**), **60** and **61**, demonstrates that both *N*-oxides are clearly indispensable for optimal cytotoxic potency in MOLM-13 cells.

The effect of phenol alkylation was investigated further. If both phenols of iodinin (**5**) were methylated (resulting in compound **62**), the cytotoxic potency in MOLM-13 (19% O₂) dropped by a factor of ~2 (Table 1). The same trend was also observed comparing mono- and dialkylated compounds **64** vs. **65** and **66** vs. **67** (Table 1, page 53 and structures in Scheme 23 page 51). These results led us to conclude that only a single hydroxy group of iodinin should be functionalized in further lead development steps.

The highest potency from these initial studies was observed for 1-hydroxy-6-(2-ethoxy-2-oxoethoxy)phenazine 5,10 dioxide (**64**), an EC₅₀ of 0.49 ± 0.12 μM at 2% O₂. Compound **64** was also found to be the most active in cell lines derived from healthy tissues, inhibiting growth of NRK and H9c2 cells in the range of 11-18 μM EC₅₀. Compound **64** did thus not outperform myxin (**8**) in terms of selectivity towards the MOLM-13 cell line. The dialkylated analogue of **64**, compound **65** was not as active, yet displaying approximately the same potency as iodinin (**5**) under both normal and low O₂ saturation. The potency of **64** and **65** was observed to be reduced significantly when the ester functionalities were replaced for free carboxylic acids (forming **68** and **69**). An obvious explanation is that the carboxylic groups carry a negative charge at physiological pH and are thus far too polar to penetrate biological membranes efficiently.

Tirapazamine (**35**) was also included in these experiments out of curiosity, as this molecule has served as a hypoxia-selective lead compound in clinical trials earlier.¹¹ TPZ (**35**) did however reveal low potency when exposed to MOLM-13 cells, not nearly comparable to the lead iodinin (**5**). Despite the low potency, these measurements confirmed TPZ (**35**) to be more than four-times as potent at 2% O₂, compared with 19% O₂ respectively. Correspondingly, this ratio was ~2.5 for iodinin (**5**).

Taken together, these findings confirmed that iodinin was a worthy lead candidate to pursue further in due to several factors. Biological evaluations of alkylated iodinin derivatives like myxin and compound **64** demonstrated that the lead can be made more soluble without drop in potency and selectivity between cancerous and healthy cell lines. This work did also gather knowledge of what transformations of the lead should not be undertaken, that is, reducing the level of *N*-oxidation and to alkylate both phenols. These findings combined led us to conclude

that prodrugs of iodinin and myxin would define the next phase of this study as a rational approach to the identification of a new and improved drug candidate.

5.2 Paper II – Prodrug development and further SAR studies on iodinin and myxin

5.2.1 Functionalization of iodinin and myxin phenols

After alkylated analogs of iodinin (**5**) had been synthesized and evaluated *in vitro* (sections 5.1.1 and 5.1.2), the focus was turned to the synthesis of iodinin (**5**) and myxin (**8**) analogues with prodrug-functionalities. The rational design aimed at increased cell penetration with retained potency and preserved cancerous cell selectivity. Synthesis of a small prodrug catalogue was initiated starting with esters and carbonates. Ester analogues **70** and **71** were prepared separately by reacting pivaloyl chloride and valeric anhydride with iodinin (**5**) under basic conditions catalyzed by DMAP (Scheme 24, next page). In a similar way, mono- and di-substituted ethyl iodinin carbonates (**72** and **73**) were isolated after iodinin was reacted with ethyl chloroformate at -40 °C and 0 °C respectively. No attempts were made to optimize these reactions as the focus at this point was to secure as many analogues as possible for the upcoming SAR-studies.

The synthesis of iodinin carbamates required screening of reaction conditions including several combinations of bases, acids, catalysts and electrophiles. Some of these entries are listed in Table 2, page 57. Iodinin (**5**) was not found to show any nucleophilicity towards isocyanates in order to obtain primary carbamates of iodinin (**5**). DMAP wasn't either found to be appropriate catalyst to react iodinin with carbamoyl chlorides to give secondary carbamates. After many unsuccessful reactions, an impatient attempt where iodinin (**5**) and DABCO (1,4-diazabicyclo-[2.2.2] octane) were exposed to neat dimethyl carbamoyl chloride resulted in a mixture that began to smoke – a clear indication of high reactivity. When THF was added as a solvent in the following entries, these reactions began to run smoothly. Iodinin (**5**) was in consequence of that functionalized with three different carbamoyl chlorides resulting in structures **74-76** (Scheme 24, next page). Overall, yields from these reactions aimed to mono-functionalize iodinin (**5**) proved disappointing (9-33% yields), but when identical reactions were undertaken with myxin (**8**), the yields improved considerably (Scheme 24). This can be explained by greater solubility of myxin, compared with that of iodinin, and only a single phenol to functionalize.

Scheme 24: Synthesis of iodinin (5) and myxin (8) prodrugs. Reagents and conditions: a) Pivaloyl chloride, Et₃N, DMAP, PhMe -40° C, 30 min, 39%. b) Valeric anhydride, Et₃N, DMAP, PhMe, 0° C, 16 h, 39%. c) ethyl chloroformate, Et₃N, DMAP, PhMe -40° C, 30 min, 25%. d) Ethyl chloroformate, Et₃N, DMAP, PhMe, 0° C, 90 min, 67%. e) A corresponding carbamoyl chloride (or HCl salt), DABCO, THF, rt,

Scheme 25: A proposed reaction mechanism for DABCO enforced carbamoylation of iodinin (**5**).

Interestingly, DABCO does not deprotonate the phenols of neither iodinin (**5**) nor myxin (**8**) as such deprotonation would clearly be observed by color changes in the reaction mixture (like for the K_2CO_3 /18-crown-6 system used for alkylations). This indicates that DABCO activates the carbamoyl chloride before the neutral phenol can act as a nucleophile. A proposed reaction mechanism is depicted in Scheme 25.

Table 2: Screening of reaction conditions to transform iodinin (**5**) into mono-functionalized carbamate analogues.

<div style="display: flex; align-items: center;"> <div style="margin-left: 20px;"> <p>Electrophiles:</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;"> <p>E1</p> </div> <div style="text-align: center;"> <p>E2</p> </div> <div style="text-align: center;"> <p>E3</p> </div> <div style="text-align: center;"> <p>E4</p> </div> </div> </div> </div>						
Entry	E+	Solvent	Additives	Temp (°C)	Time (h)	Yield
1	E1 (1.5 eq)	PhMe	Et_3N (1.5eq), DMAP (0.2 eq)	0-40	3h	Trace ^a
2	E1 (5 eq)*	DCM	Et_3N (1.5eq), DMAP (0.2 eq)	45	3h	Trace ^a
3	E2 (1.2 eq)	AcOH	-	100	2h	-
4	E2 (5 eq)	HMPA/DMF	KOtBu	rt	2h	Trace ^a
5	E2 (2 eq)	HMPA/DMF	NaH	rt	overnight	-
6	E1 (3 eq)	ACN	K_2CO_3	85	1.5	-
7	E1 (2 eq)	TFA	-	80	2.5	Trace [‡]
8	E3 (1.5 eq)	DCM	Et_3N , DMAP (0.3 eq)	45	overnight	-
9	E3 (2.5 eq)	C_6H_6	KOtBu/18-Crown-6 (0.2 eq)	50	overnight	-
10	E4 (2 eq)	PhMe	Et_3N (cat)	70	overnight	Trace ^a
11	E4 (7 eq)	PhMe	K_2CO_3 (5 eq)	rt	overnight	-
12	E4 (4 eq)	THF	Et_3N (2 eq), DMAP (0.2 eq)	60	overnight	Trace ^a
13	E2	neat	DABCO (3 eq)	rt	30 min	5% ^{b, c}
14	E2	THF	DABCO (3 eq)	rt	overnight	9% ^b

^aA cherry red product observed by TLC in trace amounts among many other colorful spots, ^bmajor product was the di-*O*-substituted product, ^cSmoke was observed after electrophile addition.

5.2.2 Biological evaluations of iodinin and myxin prodrugs

Iodinin ester and carbonate analogs **70-73** (Scheme 24) were evaluated in MOLM-13 cells and displayed low activities compared with that of iodinin (**5**) and myxin (**8**) (see Table 3). Low potency was also found to be the case with the mono-functionalized ethyl iodinin carbonate **72** ($EC_{50} \sim 50 \mu M$ normoxia) and no hypoxia selectivity. This result came as a surprise due to structural similarities between **72** and the most potent analogue observed so far; compound **64**. However, the di-*O*-substituted carbonate **73** (Scheme 24) proved to be more potent with an EC_{50} of $6 \mu M$. This was not in line with earlier trends from Table 1 (page 53) where di-*O*-alkylated derivatives were found to be less active in every case compared with their corresponding mono-alkylated analogues. To confirm this result, compound **72** was resynthesized, recrystallized and tested again, and gave the same result. These findings thus expelled further interest, at least temporarily, in ester- and carbonate prodrugs of iodinin (**5**).

When it came to the biological evaluation of mono-substituted iodinin carbamates, these compounds **74-76** were all assessed more potent than the lead iodinin in aerobic environment (see Table 3). Similar potency was also observed for the myxin carbamates **77-79**, ranging from ~ 1 - $2 \mu M$ EC_{50} in MOLM-13 (19% O_2). Nevertheless, the activity of these compounds (**74-79**) did not appear to augment in hypoxic cell cultures (see Table 3, MOLM-13 at 2% O_2).

Table 3. EC_{50} values ($\pm$ SEM) of phenazine-analogues on MOLM-13-, NRK- and H9c2 cells. The measured values of the lead iodinin (**5**) and myxin (**8**) from Table 1 are included as reference.

Cpd #	EC_{50} (μM) MOLM-13		EC_{50} (μM) NRK		EC_{50} H9c2 (n=2)
	19% O_2	2% O_2	19% O_2	2% O_2	19% O_2 (n=2)
5 (iodinin)	2.0 ± 0.07	0.79 ± 0.10	>50	>50	>50
8 (myxin)	1.4 ± 0.30	0.77 ± 0.13	77 ± 11	77 ± 14	43 ± 2
70	11.0 ± 0.7	8.9 ± 1.7	n.d.	n.d.	n.d.
71	35.4 ± 0.02	17.1 ± 1.2	n.d.	n.d.	n.d.
72	49.5 ± 0.4	66.1 ± 2.2	n.d.	n.d.	n.d.
73	6.1 ± 0.1	4.0 ± 0.3	n.d.	n.d.	n.d.
74	1.50 ± 0.16	1.41 ± 1.41	53 ± 2	59 ± 1	88 ± 10
75	0.86 ± 0.05	0.79 ± 0.09	9.6 ± 0.4	15 ± 1	12 ± 1
76	0.89 ± 0.03	0.75 ± 0.04	49 ± 2	62 ± 1	64 ± 9
77	1.01 ± 0.06	0.98 ± 0.08	27.9 ± 1.6	42.7 ± 3.4	33.2 ± 4.0
78	1.95 ± 0.11	1.84 ± 0.17	31.6 ± 2.5	34.9 ± 1.5	38.0 ± 0.1
79	1.88 ± 0.07	1.73 ± 0.09	22.3 ± 1.7	29.1 ± 0.7	39.2 ± 0.6

Experimental description is given below table 1 page 53

A possible explanation for the non-apparent hypoxia selectivity of these compounds (**74-79**) might originate from the electron withdrawing effects of the carbamate group, which consist of three electro-negative atoms and one very electron deficient carbon atom centered in between. This electronegative feature might result in lessened electron density in the aromatic ring system. However, these are only speculations, nevertheless, SAR-studies conducted on tirapazamine (TPZ, **35**) showed that incorporation of weakly electron donating groups to the aromatic ring system of **35** resulted in a higher hypoxia selectivity in most cases (with the exception of -CF₃).¹²

Due to the high potency of **74-79** observed in MOLM-13 cells, we also conducted experiments on these compounds in NRK- and H9c2 cells. As presented in Table 3, the healthy cell lines were not affected by **74-79** unless the concentration was increased considerably, in some cases by a factor of 70 like observed for compound **76** in MOLM-13 vs H9c2 (19% O₂). The pyrrolidine iodinin analogue **75** was exceptionally toxic among those – inhibiting growth of NRK and H9c2 cells at concentrations in the range of ~10-15 µM (EC₅₀). However, the EC₅₀ in MOLM-13 for this compound was found to be below 1 µM (Table 3). The piperazine-analogues **74** and **77** were considered particularly attractive due to good solubility observed in slightly acidified aqueous phase and in DMSO.

The preceding findings led us to conclude that carbamates were the most attractive functional group to be attached to the phenols of iodinin and myxin, as they appear to provide higher solubility to the lead scaffold and increase potency in some cases. With several promising alkyl- and carbamate side chains identified at this stage, our thoughts were driven towards further SAR studies of the PHz 5,10-dioxide scaffold, starting with the removal of one oxygen substituent from the 6-position.

5.2.3 Synthesis of 1-hydroxyphenazine 5,10-dioxide and analogues

In continuation of our previous findings described so far, analogues of 1-hydroxyphenazine 5,10-dioxide (**42**) were scheduled for synthesis. This strategy was anticipated to ease synthetic efforts as the two phenol functionalities of iodinin (**5**) were now known to result in selectivity difficulties when only one phenol out of two was to be functionalized. Accordingly, the intention here was to explore what influence removal of one -OH from the scaffold would have on MOLM-13 cell death and selectivity towards leukemic cell lines.

The main building block, 1-hydroxyphenazine 5,10-dioxide (**42**) was prepared by two different synthetic strategies (see Scheme 26), a pulse-wise *m*CPBA oxidation of the natural product 1-hydroxyphenazine (**25**, 1-HP) and a Beirut reaction (diketone modification). The *m*CPBA oxidation of **25** was performed on a 0.7g scale and gave the dioxide in 49% yield after chromatographic purification. The Beirut method¹³ gave larger quantities of **42** after cyclohexane-1,2-dione was condensed with benzofuroxan (**41**) in neat diethylamine, followed by a subsequent *m*CPBA oxidation step. This protocol gave 40% yield (2.0g isolated) of **42** over two steps.

The 1-hydroxyphenazine 5,10-dioxide building block was further functionalized to prepare analogues of the most active compounds identified in the project so far. A reaction of **42** with ethyl bromoacetate in the presence of K₂CO₃ and 18-Crown-6 gave compound **80** in 81% yield after chromatographic purification (see Scheme 26, right). The same step was also repeated with *tert*-butyl bromoacetate which gave **81**, and again with 2-chloro-*N,N*-diethylacetamide affording **82** after chromatography and subsequent recrystallization.

Scheme 26: The synthesis and functionalization of 1-hydroxyphenazine 5,10-dioxide (**42**). Reagents and conditions: a) *m*CPBA, PhMe, 80 °C, 5h, 49% b) 1,2-cyclohexanedione, diethylamine, 0 °C → rt, 90 min, then filtration and *m*CPBA oxidation (as in step a), 40% over 2 steps c) ethyl-/ or *tert*-butyl bromoacetate (or 2-chloro-*N,N*-diethylacetamide and KI), K₂CO₃, 18-Crown-6, DMF, rt, 2-5 hours d) A corresponding carbamoyl chloride (or the hydrochloride salt), DABCO, THF, rt, 2-5 hours.

1-hydroxyphenazine 5,10-dioxide (**42**) was also exposed to carbamoyl chlorides and DABCO forming **83-85** in good yields (see Scheme 26, left).

Apparently, all these analogues presented in Scheme 26 were observed to be more polar than their corresponding analogues with a hydroxyl group in the 6-position. This conclusion was based on lowered R_f values on silica gel, resulting in relatively polar mixtures of solvents needed to elute them (typically 2-5% MeOH in DCM), compared with their 6-OH analogues. We were therefore curious to see how the increased polarity would affect factors like potency and cell selectivity of the compounds.

5.2.4 Biological evaluations of 1-hydroxyphenazine 5,10-dioxide analogues

1-Hydroxyphenazine 5,10-dioxide (**42**) was assessed to be less potent in MOLM-13 cells by a factor of more than five, compared with that of iodinin (**5**). Based on the polarity of solvents required to elute compound **42** through silica, compound **42** is more polar than iodinin. However, **42** is also much better soluble in common organic solvents as a result of only one intramolecular ion-dipole/hydrogen bond instead of two. Furthermore, the 1-hydroxyphenazine 5,10-dioxide (**42**) proved to be less potent at low O_2 than at normal O_2 , oppositely to both iodinin (**5**) and myxin (**8**). However, the potency of **42** was restored by introduction of both alkyl- and carbamate side-chains to the phenol oxygen (see EC_{50} values of **42** versus compounds **80-85** in table 4 (MOLM-13, 20% and 2% O_2)).

Table 4. EC_{50} values ($\pm$ SEM) of 1-hydroxyphenazine 5,10-dioxide (**80**) analogues on MOLM-13-, NRK- and H9c2 cells. Determined values of the lead compound iodinin (**5**) and myxin (**8**) are included for reference.

Compound	EC_{50} (μ M) MOLM-13		EC_{50} (μ M) NRK		EC_{50} (μ M) H9c2
	19% O_2	2% O_2	19% O_2	2% O_2	19% O_2
5 (iodinin)	2.0 $\pm$ 0.07	0.79 $\pm$ 0.10	>50	>50	>50
8 (myxin)	1.4 $\pm$ 0.30	0.77 $\pm$ 0.13	77 $\pm$ 11	77 $\pm$ 14	43 $\pm$ 2
42	10.8 $\pm$ 1.8	11.5 $\pm$ 0.04	77 $\pm$ 6	>200	n.d.
80	1.51 $\pm$ 0.14	1.74 $\pm$ 0.29	24.7 $\pm$ 0	25.1 $\pm$ 0.3	n.d.
81	2.18 $\pm$ 0.09	1.88 $\pm$ 0.11	26.8 $\pm$ 0.1	26.4 $\pm$ 0.1	n.d.
82	1.17 $\pm$ 0.08	1.38 $\pm$ 0.15	12.9 $\pm$ 0.01	16.1 $\pm$ 1.0	n.d.
83	1.47 $\pm$ 0.18	1.94 $\pm$ 0.05	24.8 $\pm$ 0.3	34.8 $\pm$ 0.9	n.d.
84	1.33 $\pm$ 0.15	2.07 $\pm$ 0.16	19.5 $\pm$ 1.7	25.3 $\pm$ 0.4	n.d.
85	1.51 $\pm$ 0.18	1.21 $\pm$ 0.15	17.2 $\pm$ 0.7	23.5 $\pm$ 0.8	n.d.

Experimental description is given below Table 1, page 53.

Compounds **80-85** all showed similar EC₅₀ values in MOLM-13, in the range of 1.1-2.2 μ M (Table 4). On average basis, these analogues were 14 times more potent in leukemic cells compared with the healthy NRK cell line. Comparing the activity of 1-hydroxyphenazine 5,10-dioxide (**42**) with its alkylated- and carbamated analogues **80-85** in MOLM-13 cells shows how **42** is most certainly not capable of penetrating cell membranes as efficiently as its analogues **80-85**, as the result of two high polarity. The data presented in Table 4 also suggest that the carbamate analogues **83-85** do not hydrolyze before they enter the cells. The flexible side chains introduced are likely to lower intercalative effects and therefore, it seems plausible that the mode of action for compounds **80-85** predominantly stems from ROS generation rather than DNA-intercalation.

Carbamates have been used extensively as pesticide agents in agriculture and their breakdown has been a subject to various degradation studies to assess environmental pollution caused by them. Wolfe et al. investigated the hydrolysis of both primary and secondary carbamates using SAR-studies,¹⁴ concluding that secondary carbamates do neither hydrolyze readily in natural environment nor in non-basic solutions.

Carbamates have been used extensively as prodrugs and peptide surrogates in medicinal chemistry.¹⁵ Examples of marketed drugs with carbamate functional groups include zopiclone, ritonavir, irinotecan (**86**) and bambuterol (**88**). Irinotecan (**87**) and bambuterol (**89**) do both have carbamated phenol functional groups (see Figure 18). Irinotecan (**87**) serves as a prodrug for SN-38 (**88**), a phenol-camptothecin analogue and a topoisomerase I inhibitor for the

Figure 18: Selected examples of marketed carbamate-prodrugs, irinotecan (**87**) and bambuterol (**89**)

treatment of cancer (see Figure 18). Human liver microsomal carboxylesterases (CES1A1, CES2, and CES3) are known to convert irinotecan (**87**) to its active metabolite *in vivo*.¹⁶ Bambuterol (**89**) is a double carbamate prodrug of terbutaline (**90**), a β 2-agonist and a bronchodilator. The carbamates in bambuterol (**89**, see Figure 18) serve as protecting groups as terbutaline is rapidly metabolized via first-pass metabolism in the liver. This allows once-daily dosage of bambuterol (**89**) instead of 3 times daily for terbutaline (**90**).¹⁵ Enzymes like pChE (plasma cholinesterase)¹⁷ and liver microsomal carboxylesterases¹⁸ have been shown to metabolize bambuterol to terbutaline *in vivo*.¹⁷ Although the data presented in Table 4 suggest that the phenazine 5,10-dioxide carbamates **83-85** penetrate cell membranes before being hydrolyzed, the metabolic process these compounds must be evaluated at later stages through membrane passage investigations and pharmacokinetic studies *in vivo*.

Another concern for enhanced activity of analogues **80-85** compared with **42** is stabilization of the *N*-oxide (PHz scaffold, 10th position) via intramolecular ion-dipole/hydrogen forces. The lack of such stabilization in analogues **80-85** might enhance intrinsic reactivity of the *N*-oxide in position 10. This does however not explain the difference in cytotoxic activity between **42** and iodinin (**5**).

To briefly summarize this section, the findings herein confirm that a single oxygen-based substituent in position 6 on the scaffold can be removed without significant drop in cytotoxic potency. However, absence of the 6-positioned phenol seemed to result in less consistent trend concerning hypoxia selectivity, and to somewhat reduced selectivity towards MOLM-13, especially compared with iodinin (**5**) and myxin (**8**). The analogues prepared in Scheme 26 did therefore not outperform neither the iodinin nor myxin carbamates **74-79** in terms of potency or selectivity towards MOLM-13 cells. These findings led us to explore if potency and selectivity of the scaffold could be further improved by adding substituents to the 7 and 8 position of the 1-hydroxyphenazine 5,10-dioxide scaffold.

5.2.5 Synthesis of 7,8-disubstituted analogues of 1-hydroxyphenazine 5,10-dioxide and quinoxaline 1,4-dioxides

This section describes the synthesis of 7,8-disubstituted analogues of 1-hydroxyphenazine 5,10-dioxide. This strategy was intended to reveal if cytotoxic potency could be influenced by introduction of functional groups to the 7 and 8 positions of the PHz scaffold. In addition, similar 2-ring systems were scheduled for synthesis to reveal if one aromatic ring can be removed from the PHz scaffold without reducing the cytotoxic potency towards MOLM-13.

The synthesis of 7,8-disubstituted methoxyphenazines **45** and **93-95** was achieved applying the classic condensation procedure of 1,2-benzoquinones and *O*-phenylenediamines (see section 3.3.4 page 35). Oxidation of 3-methoxycatechol (**43**) using *ortho*-chloranil in cold ether gave a filtered crude material of the corresponding 1,2-benzoquinone **43b** (see Scheme 27). The collected crude was immediately transferred to an acidic toluene solution containing the corresponding *o*-phenylenediamine (**44** and **90-92**). These efforts gave the 1-methoxyphenazines **45** and **93-95** isolated in various yields. Similar work had been performed earlier by Cusman et al.¹⁹ and later Huigens and coworkers.²⁰ The 1-methoxyphenazines (**45**,

Scheme 27: The synthesis of 7,8-disubstituted phenazine 5,10-dioxides **100-103** and their following functionalizations either by alkyl- or carbamate side chains. *Rapid decompose of the compound observed in NMR analysis (CDCl₃ solution).

93-95) were all treated with BBr₃ under refluxing conditions to give their corresponding 1-hydroxyphenazines **96-99** in excellent yields (Scheme 27).

The 1-hydroxyphenazines **96-99** were exposed to a pulse-wise addition of *m*CPBA in toluene at 80 °C. This gave 7,8-disubstituted-1-hydroxyphenazine 5,10-dioxides **100-103** in acceptable yields. Neither were the reaction conditions nor the purification methods optimized in any way for the reactions depicted in Scheme 27 as the project focus was still valid, to provide as many analogues as possible for *in vitro* studies. Especially, 7,8-dibromo and 7,8-dichloro substituted analogs **102** and **103** turned out to be difficult to purify due to very poor solubility in organic media, high lipophilicity and tailing on silica gel. The crudes could however be used easily for further synthesis.

7,8-disubstituted-1-hydroxyphenazine 5,10-dioxides **100-103** were further functionalized with one alkyl side chain and the most promising carbamate side chain to prepare 1-(2-ethoxy-2-oxoethoxy)phenazine 5,10-dioxide analogs **104-107** and *N*-methylpiperazine carbamates **108-111** (Scheme 28). The four-ring analogue **105** turned out to be exceptionally unstable. Although TLC-analysis indicated pure fractions obtained after flash column chromatography on silica, the compound decomposed rapidly in CDCl₃ during NMR analysis. This compound was therefore not subjected to biological testing. This problem was not observed for any of the other compounds. The 7,8-dimethyl-1-hydroxyphenazine 5,10-dioxide (**100**) was also used to construct diethylamide-analogue **112** and pyrrolidine carbamate analogue **113** (Scheme 27, bottom).

Now, turning to synthesis of 2-ring quinoxalines, 2,3-dimethylquinoxalin-5-ol (**115**) was prepared by reducing 3-amino-2-nitrophenol (**114**) with sodium dithionite refluxing in a basic dispersion of MeOH and H₂O. The crude mixture was then concentrated to a slur, diluted with

Scheme 28: Synthesis of functionalized two-cyclic aromatic *N*-oxides. Reaction conditions: i) Na₂S₂O₄, Na₂CO₃, H₂O/MeOH, reflux, 2h. ii) diacetyl, AcOH/PhMe, 20h, 79% over two steps. iii) *m*CPBA, PhMe, 80 °C, 4.5h, 80%. iv) corresponding carbamoyl chloride, DABCO, THF.

AcOH-PhMe (3:2) and subsequently exposed to the diketone compound diacetyl (2,3-diketo-butane). This method gave **115** in 79% yield over 2 steps and the subsequent *m*CPBA oxidation gave the corresponding dioxide **116** conveniently in 80% yield after flash column chromatography. Further functionalizations with carbamoyl chlorides were then undertaken to obtain carbamate analogues **117** and **118** were performed as well (Scheme 28).

5.2.6 Biological evaluations of 7,8-disubstituted phenazines and quinoxaline 1,4-dioxides

Biological evaluations of compounds **100-113** (except **102**, **103** and **105**) and **116-118** are presented within Table 5, next page.

Comparing 1-hydroxyphenazine 5,10-dioxide (**42**) and its 7,8-dimethyl 1-hydroxyphenazine (**100**), an increased potency could be observed (from 10.8 to 2.8 μ M EC₅₀ in MOLM-13, normoxia). The 7,8-dimethylated analogue **100** proved also to be significantly more potent at low O₂, which **42** was not. Based on that comparison, increased cell membrane penetration due to higher lipophilicity of **100** compared with **42** is likely to be the case. Incorporation of the two methyl groups in the 7 and 8 positions therefore appears to increase the hypoxia selectivity and that would be in line with the trends observed from SAR-studies on TPZ (**35**) as earlier mentioned in page 55.¹²

Further on, extending the aromatic scaffold with one annulated benzene ring to positions 7 and 8 (compound **101**) resulted in a highly insoluble substance, that failed to give any response in any biological assays, most likely because of very poor solubility in the cell media or insufficient penetration of cell membranes. However, when this compound was functionalized with a piperazine side chain, resulting in compound **109**, the compound became moderately active (EC₅₀ = 2.75 μ M). This result exemplifies dramatic increase in solubility and potency provided by the piperazine side chain introduced. In addition, the 4-ring candidate **109** turned out to be the most toxic compound tested in healthy NRK cells to this point, closing this chapter to further interest in these ring-systems.

Four new analogues turned out to be particularly potent in MOLM-13 cells, the 7,8-dimethylated- and dihalogenated piperazine analogues **108**, **110** and **111** and the 7,8-dichloro analogue **106** bearing the 2-ethoxy-2-oxoethoxy side chain (see Table 5). These results clearly demonstrate that synthetic alterations in the 7 and 8 position of the PHz 5,10-dioxied scaffold can influence potency dramatically. Due to very high potency of these compounds, it was considered very important to address the toxicity of these compounds in other healthy cell lines as well, and more studies currently ongoing in H9c2 cells. 7,8-dichloropiperazine analogue **110**

Table 5. EC₅₀ values (±SEM) of analogues on MOLM-13-, NRK- and H9c2 cells. The measured values of iodinin (5), myxin (8) and 1-hydroxyphenazine 5,10-dioxide (42) are included for reference.

Cpd #	EC ₅₀ (μM) MOLM-13		EC ₅₀ (μM) NRK		EC ₅₀ (μM) H9c2
	19% O ₂	2% O ₂	19% O ₂	2% O ₂	19% O ₂
5 (iodinin)	2.0 ± 0.07	0.79 ± 0.10	>50 [†]	>50	>50
8 (myxin)	1.4 ± 0.30	0.77 ± 0.13	77 ± 11[†]	77 ± 14	43 ± 2
42	10.8 ± 1.8	11.5 ± 0.04	77 ± 6	>200	n.d.
100	2.8 ± 0.3	2.3 ± 0.1	>50 [†]	>50 [†]	>50
101	-**	>10 [†]	n.d.	n.d.	n.d.
102	n.d.	n.d.	n.d.	n.d.	n.d.
103	n.d.	n.d.	n.d.	n.d.	n.d.
104	8.3 ± 0.8	6.3 ± 0.3	140 ± 2 [†]	>200	194
105	n.d.	n.d.	n.d.	n.d.	n.d.
106	0.41 ± 0.05	0.38 ± 0.04	3.14 ± 0.02	4.67 ± 0.09	n.d.
107	2.44 ± 0.03	1.57 ± 0.09	10.4 ± 0.7	9.7 ± 2.4	n.d.
108	0.63 ± 0.05	0.54 ± 0.06	12.5 ± 0.6	13.9 ± 0.5	10
109	2.75 ± 0.18	1.76 ± 0.10	2.02 ± 0.06	7.52 ± 0.38	n.d.
110	0.34 ± 0.04	0.32 ± 0.06	10.4 ± 1.5	12.9 ± 0.4	n.d.
111	0.41 ± 0.04	0.23 ± 0.03	6.2 ± 0.02	7.6 ± 0.6	n.d.
112	1.88 ± 0.37	3.39 ± 0.78	20.6 ± 0.3	51.5 ± 1.2	n.d.
113	2.03 ± 0.30	2.01 ± 0.61	6.63 ± 0.04	8.89 ± 0.06	n.d.
116	>200	>200	n.d.	n.d.	n.d.
117	>200	>200	n.d.	n.d.	n.d.
118	>200	>200	n.d.	n.d.	n.d.

Experimental description is given below Table 1 page 53.

was assessed to be more cytotoxic in leukemic cell lines compared with the NRK-line by a factor of 30 at normoxia (19% O₂). This ratio even increased to a factor of 40 at hypoxia (2% O₂). The corresponding 7,8-dibromo-analogue **111** proved to be the most potent compound measured so far at 2% O₂ in MOLM-13 but high activity of this analogue in the NRK cell line excluded further interest in it (see Table 5).

Daunorubicin is a standard in current treatment of AML.²¹⁻²² As discussed and demonstrated by Herfindal et al. *in vitro*,¹ daunorubicin has severe cardiotoxic side effects, and was found to introduce cell-death to 50% of rat cardio myoblasts (H9c2) at 1 μM concentration.¹ The most toxic compound from this project measured in this cell line so far is compound **109** which displays an EC₅₀ value of 10 μM concentration. However, it has to be pointed out clearly that these results from both NRK and H9c2 are preliminary and only represent the average EC₅₀ values based on two parallel experiments. Other phenazine 5,10-dioxide compounds presented

herein range from an EC₅₀ of 13-97 μ M in H9c2 (iodinin, myxin and **42** only known to have >50 μ M EC₅₀).

Overall, the results point towards a very promising technology for improved AML-treatment regimes. Clear trends have been identified, increasing the probability of identifying promising drug candidates with improved performance compared to the standard treatment regimens with anthracyclines. In particular, the far less cardiotoxic effects of the identified phenazine candidates in H9c2, compared with the anthracycline daunorubicin encourages further development.

It cannot be put to doubt that the anthracyclines are a highly successful class of antineoplastic antibiotics derived from nature. Both doxo- and daunorubicin have been utilized in chemotherapies for multiple types of cancer for a long time.²¹ Anthracyclines are supposed to induce their notorious cardiotoxicity (at least partly) through iron mediated production of ROS, thereby introducing oxidative stress to cardiac cells.²² This mechanism is not known in clear detail and remains a topic for debate after decades of use in the clinic.²³ In many ways, anthracyclines have similarities to phenazines and phenazine 5,10-dioxides. First, the trigonal planar structure of both scaffolds is ensured by aromaticity and this feature is necessary for DNA-intercalation.²⁴ Secondly, their overlapping structural features, indicates that the anthracycline quinone moiety and the *N*-oxide functionalities of the phenazine 5,10-dioxides may induce attractive forces at the same points in target receptors. Last but not least, both can produce ROS and can undergo redox-cycling.²⁵⁻²⁶ This mode of action was explained for naturally occurring phenazines in page 22. The reduction of doxorubicin by NADH dehydrogenase in mitochondria forms a hydroquinone radical able to react with molecular O₂. This process can further produce the superoxide radical (O₂^{•-}) also generating hydrogen peroxide (H₂O₂). Consequently, Fe²⁺ can transform H₂O₂ to the hydroxyl radical ([•]OH) in a Fenton reaction like shown here below in equation 1.

Equation 1: Fenton reaction.

5.3 Synthesis of miscellaneous compounds and antimicrobial evaluations

5.3.1 Synthesis of miscellaneous phenazine 5,10-dioxides

Experimental procedures for this section are in Appendix 1. Some of the compounds presented herein have been tested in antimicrobial evaluations.

This section describes synthetic efforts that were conducted in continuation of results that showed that halogens in the 7- and 8 position of the phenazine scaffold can enhance cytotoxic effects on MOLM-13 and still be selective. Therefore, our interest in halogenations of other positions on the PHz scaffold was enhanced. Some of the new compounds described within this section have not yet been evaluated in MOLM-13 cells but investigations are ongoing.

1-Hydroxyphenazine 5,10-dioxide (**42**) was halogenated by exposure to either *N*-iodosuccinimide (NIS) or *N*-bromosuccinimide (NBS). The iodination proved highly efficient affording 98% yield after chromatography and the corresponding bromination attempt gave 73% respectively (Scheme 29). This reaction was also performed with *N*-chlorosuccinimide (NCS) but low reactivity compared with NBS and NIS was observed and resulted in a mixture of unreacted starting material, mono-chlorinated, and di-chlorinated compounds that proved difficult to separate by chromatography. As NBS was found to be less reactive than NIS, a di-brominated compound was also observed on TLC. This compound was found to be highly insoluble in organic solvents and difficult to purify.

Scheme 29: Halogenations and subsequent carbamoylations of 1-hydroxyphenazine 5,10-dioxide.

To ensure what position was halogenated first, a crystal was grown of **119** (DCM as inner solvent, hexane as outer; vapor diffusion) and the structure determined by X-ray crystallography. This effort indeed revealed the structure of 1-hydroxy-2-iodophenazine 5,10-dioxide (**119**). The crystal structure of a di-brominated compound (structure not shown) was

also determined by X-ray crystallography and revealed the structure of 2,4-dibromo-1-hydroxyphenazine 5,10-dioxide (experimental data not shown). These results confirm that halogens react first *ortho* to the 1-hydroxy group and thereafter into the position *para* to the 1-hydroxy group.

Figure 19: Crystal structure of 1-hydroxy-2-iodophenazine 5,10-dioxide (**119**). Determined by Prof. Carl Henrik Görbitz.

As mentioned earlier in section 3.2.4, page 27, Weigle et al. had shown 1-methoxyphenazine 5,10-dioxide (**30**) to be potent in terms of antimicrobial effects.⁹ It was therefore decided to prepare compound **30** in addition to three other methoxy analogues (compounds **125-127**) for upcoming antimicrobial evaluations (see Scheme 30). Dimethyl sulfate (Me_2SO_4) was used for methylation of the *ortho*-halogenated compounds as attempts involving MeI on compounds **119** and **120** were characterized by low reactivity.

Scheme 30: Methylations of 1-hydroxyphenazine 5,10-dioxides.

The piperazine carbamate compound **130** was also prepared (Scheme 31). ethyl piperazine-1-carboxylate was reacted with chloroacetyl chloride to give the α -chloro piperazine carbamate **129** which was only purified via extraction and reacted further with 1-hydroxyphenazine 5,10-dioxide in the presence of K_2CO_3 , 18-crown-6 and KI in DMF. Purification of **130** by recrystallization from hot EtOH gave **130** in 30% yield from **42** (Scheme 31).

Scheme 31: Preparation of piperazine carbamate compound **130** from **42**.

5.3.2 Antimicrobial evaluations

Antimicrobial resistance (AMR) is considered an ongoing and increasing threat to human healthcare worldwide. Defined by the WHO; “*AMR is resistance of a microorganism to an antimicrobial drug that was originally effective for treatment of infections caused by it*”. WHO’s report on global surveillance of antimicrobial resistance from 2014 reveals that AMR is happening right now and spreading fast across the world. Treatment options for common types of infections that have been treatable for decades are therefore at risk as the arsenal of antimicrobial drugs is getting thinner and thinner. In addition, other treatments such as cancer chemotherapy, organ transplantation and surgical operations rely heavily on use of antibiotic treatments.²⁷ As a consequence of AMR, new classes of antimicrobial drugs with new modes of action are utterly called for. Many phenazines have been shown to possess antimicrobial activities and especially myxin (**8**).^{9, 28-29} Phenazines and phenazine 5,10-dioxides might therefore have a significant potential as starting points for drug development of new antimicrobial agents, especially as no molecules of this class have earlier been used to treat infectious diseases in humans.

As a part of this project, synthesized compounds were submitted to Tromsø, Norway (Prof. Ørjan Samuelsen lab) for antimicrobial evaluations in highly resistant strains of both Gram-positive (*E. faecium* and *S. aureus*) and -negative (*P. aeruginosa* and *K. pneumoniae*) bacteria.

To the best of authors' knowledge, the only SAR-study that has been reported on phenazine 5,10-dioxides in microbes, including iodinin (**5**) and myxin (**8**) was conducted by researchers from Hoffman-La Roche in the 1970s.⁹ This was briefly discussed in section 3.2.4, page 27 and as emphasized earlier, the general trend observed therein was that antimicrobial activities and spectra of the myxin (**8**) analogues tested generally dropped when the structure was enlarged or made more complicated than myxin (**8**). Two compounds that can be considered simplified structures of myxin (**8**), were however found to possess potent and broad spectrum activity: 1-methoxyphenazine 5,10-dioxide (**30**) and 1-methylphenazine 5,10-dioxide (**31**).

Results from our antimicrobial screening of heterocyclic aromatic *N*-oxides are presented in Table 6 (next page). These data must be considered preliminary as further investigations are ongoing.

The most obvious trend observed after collection of the data was that all the analogues which were most potent in inhibiting growth of both Gram-positive and -negative bacteria had either methyl group or an alkyl side-chain attached to the phenol (see Figure 20). Carbamate-analogues **75**, **83** and **85** and showed activity as well, but only towards Gram-positive strains.

Figure 20. Alkylated analogues observed to be potent in antimicrobial evaluations.

Table 6. Antimicrobial evaluations of selected phenazine 5,10-dioxides. Values are presented as minimum inhibitory concentration (MIC) values in μM concentrations.

Ref. no <i>Species</i>	Gram+		Gram-	
	A1-22 <i>E. faecium</i> VanA	A4-37 <i>S. aureus</i> MRSA	K34-7 <i>P. aeruginosa</i> VIM-2	K66-45 <i>K. pneumoniae</i> NDM-1
5 (Iodinin)	62.5	>250	>250	>250
8 (myxin)	4 to 8	<0,5	16	16
30	31.25	1 to 2	16	8
42	12	6	>100	>100
66	8	1 - < 0,5	>250	>250
68	63	125	1000	1000
74	>250	125	>250	>250
75	8	4	>250	>250
76	Not soluble in 10 mM DMSO			
77	125 to 250	>250	31.25	>250
78	62.5	2 to 4	>250	>250
79	62.5	2 to 4	>250	>250
80	4	< 0,5	125	62.5
82	31	≤ 2	250	250
83	8	2 to 4	>250	250
84	Not soluble in 10 mM DMSO			
85	4	≤ 2	1000	>100
104	25	3	>100	>100
106	≤ 0.5	≤ 0.5	125	62.5
107	8	16	>250	>250
108	Not soluble in 10 mM DMSO			
109	50	-	>100	>100
110	Not soluble in 10 mM DMSO			
112	Not soluble in 10 mM DMSO			
113	Not soluble in 10 mM DMSO			
116	>1000	>1000	>1000	>1000
117	1000	>1000	1000	1000
118	>1000	>1000	>1000	>1000
126	0.4	0.8	25	12.5
129	Not soluble in 10 mM DMSO			
130	31	4	1000	500

5.4 Conclusions and summary of structure-activity relationships

A new efficient total synthesis of the phenazine 5,10-dioxide natural products iodinin (**5**) and myxin (**8**) and many new compounds derived from them has been established in few and reliable steps. The key synthetic step; oxidation of phenazines to phenazine 5,10-dioxides has enabled the synthesis of a library of structurally related phenazine 5,10-dioxides that to authors' best knowledge has no precedence in published literature.

Figure 21, depicts and summarizes the potency ranking of heterocyclic aromatic *N*-oxides attached to the most promising side-chain identified. It can be observed that iodinin (**5**) with an EC₅₀ of 2.0 μM in MOLM-13 has been functionalized with a piperazine carbamate side chain to result in iodinin-carbamate **74**. This functionalization increased the potency of the lead candidate by ~25%. Removal of the phenol -OH (position 6) results in compound **84** that is equally potent to **74**. This 1-hydroxyphenazine 5,10-dioxide piperazine analogue **84** can be synthesized with much more ease than **74** in terms of yields, purifications and overall labor intensity. However, the selectivity ratio between MOLM-13 and the healthy NRK cell has been reduced (from 35 to 17) which was not the goal. Installation of a methoxy group in the 6th position (myxin analogue **77**) increased potency to 1 μM EC₅₀ in MOLM-13 and this transformation also resulted in higher selectivity between MOLM-13 and NRK cells compared with **84**. Furthermore, installation of two methyl groups into positions 7 and 8 of compound **84** gives the dimethyl analogue **109** that is 230% more potent than **84**, and has superior selectivity between cancerous and healthy cells compared with that of **84**. Substitution of the methyl groups for two -Br atoms (compound **112**) gave higher potency but at the cost of reduced selectivity between MOLM-13 and NRK. Finally, the 7,8-dichloro-analogue **111** gave the most potent compound in MOLM-13 (at 19% O₂) with an EC₅₀ of 0.34 μM. This analogue (**111**) serves as a good example in the summary of the presented work, as potency to introduce cell

Figure 21: Structure activity relationships – potency ranking of synthetic analogues. *ratio: Determined EC₅₀ in MOLM-13 divided by EC₅₀ in NRK (both at 19% O₂).

death to cancerous leukemic cells (MOLM-13) was enhanced by a factor of ~6 compared with iodinin (**5**) (normoxia). Additionally, this analogue (**111**) was found to be 30 times more selective towards MOLM-13 cells than healthy NRK cells at normoxic levels, and 40 times at hypoxic ones. This analogue is currently being investigated in the H9c2 cell line as well.

The lead optimization process provided much knowledge of phenazine 5,10-dioxides and structure-activity relationships. For example, 1-hydroxyphenazine 5,10-dioxide (**80**) is over 5-times less potent than iodinin (**5**). When this compound was functionalized with various side-chains on its 1-hydroxy moiety, the potency was restored. This strongly suggests that an appropriate side chain enhances cell-membrane permeability which results in increased potency.

The lead optimization process was undertaken using classic medicinal chemistry. An interdisciplinary cross-talk and decision-making between chemists and biologists has afforded new compounds like **74-77**, **79**, **108** and **110** which all outperform iodinin (**5**) in terms of potency and solubility.

Furthermore, this work has unravelled some transformations of the lead that should not be pursued further for anti-AML therapy, such as addition of, or removal of one condensed aromatic ring from the scaffold (compounds **110** and **118**) (Figure 21).

Preliminary screenings in resistant Gram-positive and -negative strains reveal a trend that phenazine 5,10-dioxides characterized by either methyl- or alkyl group attached to the phenol -OH are more potent and broad spectrum compared with the carbamate analogues tested so far. Some carbamates were however shown to be active against the Gram-positive vancomycin resistant *E. faecium* and the methicillin resistant *S. aureus* (MRSA) (Table 6). The myxin carbamates **77-79** did not perform as well as myxin (**8**) in these studies, clearly suggesting that these compounds do not hydrolyse to unleash their parent compound myxin (**8**). Therefore, alkylated analogues hold most promise in terms of microbe growth inhibition at this stage, whether they are halogenated or not in the 2-position.

5.5 Future perspectives

5.5.1 Synthesis

The most important aspect for future development of phenazine 5,10-dioxides as antileukemic agents would be to identify novel compounds that have a high selectivity ratio between MOLM-13 and healthy cells. As summarized in section 5.4. This work has already provided knowledge regarding the SAR of phenazine 5,10-dioxides in various human and non-human cells. It is of high interest for this project to synthesize and perform biological evaluations of more halogenated compounds *in vitro* where modification of other positions on the phenazine scaffold are explored. Fluorinations would be the next important step since fluorine is isosteric to hydrogen, and such compounds have not been tested before in this project. Hypoxia selectivity would be preferred but not considered obligatory if compounds show good selectivity towards leukemic cells. The results presented in this thesis strongly suggests that this could be achievable.

7,8-difluoro-1-hydroxyphenazine 5,10-dioxide can be synthesized either via the Beirut diketone application as depicted in Scheme 32 or employing the classic condensation between 1,2-benzoquinones and *o*-diphenylamines. All reagents to achieve this transformation are easily accessible from commercial suppliers.

Scheme 32: The synthesis of 7,8-difluoro-1-hydroxyphenazine 5,10-dioxide and analogues.

Trifluoromethylations have not been explored yet, but are of high interest for both anti-cancer and antimicrobial evaluations. As an example, iodinin (**5**) could be trifluoromethylated to afford the halogenated myxin analogue 1-hydroxy-6-(trifluoromethoxy)phenazine 5,10-dioxide as depicted in Scheme 36. It would be surprising if this molecule and analogues derived from it would not perform well *in vitro* based on results presented in Table 1, page 53.

Scheme 33: Trifluoromethylation of iodinin (**5**)

Section 5.3.1, page 69 describes initial efforts on late stage halogenations of the phenazine 5,10-dioxide scaffold by NCS, NBS or NIS. What we learned from these reactions is that the *ortho* position to the phenol -OH is most reactive towards radical reactions. Other types of free radical reactions could therefore be investigated, such as reactions of phenazine 5,10-dioxides and Zinc sulphinates (Baran reagents).³⁰ These types of reagents could give access to various functional groups on the phenazine scaffold and influence important factors such as log P and cell membrane penetration (Scheme 37).³¹

Scheme 34: Reactions between phenazine 5,10-dioxides and zinc sulphinates could be explored

The halogenated compounds synthesized in Section 5.3.1, page 69 could also be used as substrates for Buchwald-Hartwig reactions to afford novel aryl-amines as depicted in scheme 35.

Scheme 35: Buchwald-Hartwig C-N cross coupling for novel phenazine aryl-amines.

5.5.2 Biology

This work has so far led to identification of several promising candidates ready for *in vivo* biological evaluation, more specifically starting with leukemic mice or -rat models. To start with, this could be performed by comparing survival of leukemic mice that either will have administered a chosen phenazine derivative compared to negative (placebo) and positive (anthracycline therapy) controls. However, more *in vitro* screens should first be conducted in a broader spectrum of cell types, both cancerous and healthy ones. In this context, selectivity is the key. Accordingly, it is extremely important to select a candidate wisely, that is potent and has a large difference in EC₅₀ between cancerous and healthy cells. The results discussed within this chapter show that design and synthesis of such candidate is most likely well achievable, if not already done.

5.6 References

1. Myhren, L. E.; Nygaard, G.; Gausdal, G.; Sletta, H.; Teigen, K.; Degnes, K. F.; Zahlén, K.; Brunsvik, A.; Bruserud, O.; Doskeland, S. O.; Selheim, F.; Herfindal, L., Iodinin (1,6-dihydroxyphenazine 5,10-dioxide) from *Streptosporangium* sp. induces apoptosis selectively in myeloid leukemia cell lines and patient cells. *Mar. Drugs* **2013**, *11* (2), 332-49.
2. Sletta, H.; Degnes, K. F.; Herfindal, L.; Klinkenberg, G.; Fjaervik, E.; Zahlén, K.; Brunsvik, A.; Nygaard, G.; Aachmann, F. L.; Ellingsen, T. E.; Doskeland, S. O.; Zotchev, S. B., Anti-microbial and cytotoxic 1,6-dihydroxyphenazine-5,10-dioxide (iodinin) produced by *Streptosporangium* sp. DSM 45942 isolated from the fjord sediment. *Appl. Microbiol. Biotechnol.* **2014**, *98* (2), 603-10.
3. Winkler, J. D.; Twenter, B. M.; Gendrineau, T., A new approach to the synthesis of substituted phenazines via palladium-catalyzed aryl ligation. *Heterocycles* **2012**, *84* (2), 10.3987/COM-11-S(P)76.
4. Wohl, A.; Aue, W., Ueber die Einwirkung von Nitrobenzol auf Anilin bei Gegenwart von Alkali. *Ber. Dtsch. Chem. Ges.* **1901**, *34* (2), 2442-2450.
5. Mateo Alonso, A.; Horcajada, R.; Groombridge, H. J.; Chudasama, R.; Motevalli, M.; Utley, J. H. P.; Wyatt, P. B., Synthesis of phenazine derivatives for use as precursors to electrochemically generated bases. *Org. Biomol. Chem.* **2005**, *3* (15), 2832-2841.
6. Yosioka, I.; Kidani, Y., Studies on Phenazines. II Synthesis of Iodinin. *Yakugaku Zasshi* **1952**, *72* (10), 1301-1303.
7. Leimgruber, W. M., (NJ), Weigele, Manfred (North Caldwell, NJ) Alkylation of iodinin. 1975.
8. Lavaggi, M. L.; Nieves, M.; Cabrera, M.; Olea-Azar, C.; López de Ceráin, A.; Monge, A.; Cerecetto, H.; González, M., Structural modifications on the phenazine N,N'-dioxide-scaffold looking for new selective hypoxic cytotoxins. *Eur. J. Med. Chem.* **2010**, *45* (11), 5362-5369.
9. Weigele, M.; Maestroni, G.; Mitrovic, M.; Leimgruber, W., Antimicrobial agents structurally related to myxin. *Antimicrob. Agents Chemother.* **1970**, *10*, 46-9.
10. Li, B.; Berliner, M.; Buzon, R.; Chiu, C. K. F.; Colgan, S. T.; Kaneko, T.; Keene, N.; Kissel, W.; Le, T.; Leeman, K. R.; Marquez, B.; Morris, R.; Newell, L.; Wunderwald, S.; Witt, M.; Weaver, J.; Zhang, Z.; Zhang, Z., Aqueous Phosphoric Acid as a Mild Reagent for Deprotection of tert-Butyl Carbamates, Esters, and Ethers. *J. Org. Chem.* **2006**, *71* (24), 9045-9050.
11. Brown, J. M.; Wilson, W. R., Exploiting tumour hypoxia in cancer treatment. *Nat. Rev. Cancer* **2004**, *4* (6), 437-447.
12. Hay, M. P.; Gamage, S. A.; Kovacs, M. S.; Pruijn, F. B.; Anderson, R. F.; Patterson, A. V.; Wilson, W. R.; Brown, J. M.; Denny, W. A., Structure-Activity Relationships of 1,2,4-Benzotriazine 1,4-Dioxides as Hypoxia-Selective Analogues of Tirapazamine. *J. Med. Chem.* **2003**, *46* (1), 169-182.
13. Issidorides, C. H.; Atfah, M. A.; Sabounji, J. J.; Sidani, A. R.; Haddadin, M. J., Application of 1,2-diketones in the synthesis of phenazine oxides. *Tetrahedron* **1978**, *34* (2), 217-221.
14. Lee Wolfe, N.; Zepp, R. G.; Paris, D. F., Use of structure-reactivity relationships to estimate hydrolytic persistence of carbamate pesticides. *Water Res.* **1978**, *12* (8), 561-563.
15. Ghosh, A. K.; Brindisi, M., Organic Carbamates in Drug Design and Medicinal Chemistry. *J. Med. Chem.* **2015**, *58* (7), 2895-2940.
16. Sanghani, S. P.; Quinney, S. K.; Fredenburg, T. B.; Davis, W. I.; Murry, D. J.; Bosron, W. F., Hydrolysis of irinotecan and its oxidative metabolites, 7-ethyl-10-(4-N-(5-aminopentanoic acid)-1-piperidino) carbonyloxycamptothecin and 7-ethyl-10-(4-(1-piperidino)-1-amino)-carbonyloxycamptothecin, by human carbocylesterases CES1A1, CES2, and newly expressed carboxylesterase isoenzyme, CES3. *Drug Metab. Dispos.* **2004**, *32* (5), 505-511.
17. Tunek, A.; Levin, E.; Svensson, L.-Å., Hydrolysis of 3H-bambuterol, a carbamate prodrug of terbutaline, in blood from humans and laboratory animals in vitro. *Biochem. Pharmacol.* **1988**, *37* (20), 3867-3876.
18. Svensson, L.-Å.; Tunek, A., The design and bioactivation of presystemically stable prodrugs. *Drug Metab. Rev.* **1988**, *19* (2), 165-194.

19. Conda-Sheridan, M.; Marler, L.; Park, E.-J.; Kondratyuk, T. P.; Jermihov, K.; Mesecar, A. D.; Pezzuto, J. M.; Asolkar, R. N.; Fenical, W.; Cushman, M., Potential Chemopreventive Agents Based on the Structure of the Lead Compound 2-Bromo-1-hydroxyphenazine, Isolated from *Streptomyces* Species, Strain CNS284. *J. Med. Chem.* **2010**, *53* (24), 8688-8699.
20. Garrison, A. T.; Abouelhassan, Y.; Kallifidas, D.; Bai, F.; Ukhanova, M.; Mai, V.; Jin, S.; Luesch, H.; Huigens, R. W., Halogenated Phenazines that Potently Eradicate Biofilms, MRSA Persister Cells in Non-Biofilm Cultures, and *Mycobacterium tuberculosis*. *Angew. Chem. Int. Ed.* **2015**, *54* (49), 14819-14823.
21. Gewirtz, D., A critical evaluation of the mechanisms of action proposed for the antitumor effects of the anthracycline antibiotics adriamycin and daunorubicin. *Biochem. Pharmacol.* **1999**, *57* (7), 727-741.
22. Minotti, G.; Cairo, G.; Monti, E., Role of iron in anthracycline cardiotoxicity: new tunes for an old song? *The FASEB Journal* **1999**, *13* (2), 199-212.
23. Simunek, T.; Sterba, M.; Popelova, O.; Adamcova, M.; Hrdina, R.; Gersl, V., Anthracycline-induced cardiotoxicity: overview of studies examining the roles of oxidative stress and free cellular iron. *Pharmacol. Rep.* **2009**, *61* (1), 154-71.
24. Lemke, T. L.; Foye, W. O., *Foye's principles of medicinal chemistry*. 6th ed. / Thomas L. Lemke. ed.; Lippincott Williams & Wilkins: Philadelphia, Pa. ; London, **2008**.
25. Goodman, J.; Hochstein, P., Generation of free radicals and lipid peroxidation by redox cycling of adriamycin and daunomycin. *Biochem. Biophys. Res. Commun.* **1977**, *77* (2), 797-803.
26. Sinha, S.; Shen, X.; Gallazzi, F.; Li, Q.; Zmijewski, J. W.; Lancaster, J. R.; Gates, K. S., Generation of Reactive Oxygen Species Mediated by 1-Hydroxyphenazine, a Virulence Factor of *Pseudomonas aeruginosa*. *Chem. Res. Toxicol.* **2015**, *28* (2), 175-181.
27. Antimicrobial resistance. <http://www.who.int/mediacentre/factsheets/fs194/en/> (accessed 13.05.2014).
28. Behki, R. M.; Lesley, S. M., Deoxyribonucleic Acid Degradation and the Lethal Effect by Myxin in *Escherichia coli*. *J. Bacteriol.* **1972**, *109* (1), 250-261.
29. Lesley, S. M.; Behki, R. M., Mode of Action of Myxin on *Escherichia coli*. *J. Bacteriol.* **1967**, *94* (6), 1837-1845.
30. Fujiwara, Y.; Dixon, J. A.; O'Hara, F.; Funder, E. D.; Dixon, D. D.; Rodriguez, R. A.; Baxter, R. D.; Herle, B.; Sach, N.; Collins, M. R.; Ishihara, Y.; Baran, P. S., Practical and innate carbon-hydrogen functionalization of heterocycles. *Nature* **2012**, *492* (7427), 95-99.
31. Gerebtzoff, G.; Li-Blatter, X.; Fischer, H.; Frentzel, A.; Seelig, A., Halogenation of Drugs Enhances Membrane Binding and Permeation. *Chem. Bio. Chem.* **2004**, *5* (5), 676-684.

Chapter 6 - Introduction to liver X receptors

6.1 Nuclear receptors

The nuclear receptors (NRs) are a large superfamily of protein receptors that recognize lipophilic ligands and interact with DNA directly to influence gene transcription. They are therefore also known as transcription factors.¹ Nuclear receptors play an essential role in all mammalian growth, development, metabolism, aging, function and dysfunction.¹⁻² Dysfunction of nuclear receptors can cause a wide array of malfunctions such as infertility, but also diseases such as diabetes, obesity, atherosclerosis and cancer.³ There are 48 NRs in humans⁴ and all of them have been identified through genome sequencing efforts. Accordingly, some NRs were discovered even before their endogenous ligands were known, terming them so-called orphan receptors.⁵ Known endogenous ligands include steroids, thyroid hormones, retinoic acid (vitamin A) and calcitriol (vitamin D) among others (see Figure 22). Since NRs are activated by small molecular lipophilic ligands, they are often termed as “ideal” drug targets.⁴ Endogenous ligands such as those depicted in Figure 22 can in most, if not all cases be replaced by modified drug molecules. Many of the nuclear receptors have specific isoform sub-types that regulate different genes, are activated by different ligands and are expressed more selectively in specific tissues. Thus, the fact that ligands can be designed specifically not as pure agonists or antagonists of these receptors, but even to be partial agonists- or antagonists regulating expression for a specific set of target genes in a distinct tissue makes the NRs exceptionally attractive as drug targets.⁴ For example, ligands like tamoxifen, raloxifene and

Figure 22: Selected overview of nuclear receptor ligands

clomiphene act as selective estrogen receptor modulators (SERMs) with tissue specific agonistic/antagonistic function.⁶

The NRs are complex structures composed of several functional domains. A typical NR structure (Figure 23) contains a variable -NH₂-terminal region (A/B), a DNA-binding domain (C) which is highly evolutionary conserved and contains two zinc fingers that target and bind to specific DNA-elements.⁵ Thereafter is a linker region (D) followed by the ligand binding domain (LBD). The LBD (E) is conserved as well and consists of 12 helices that form a lipophilic pocket where the ligand binding site is located. Some receptors also contain a -COOH terminal region (F) where the function is less clear. Upon agonistic ligand binding, conformational changes are promoted to alter interactions with other regulatory proteins: Co-activators are assembled and co-repressors released in order to initiate gene transcription.^{2, 7}

Regulation of gene transcription by NRs is controlled by several distinct mechanisms where endogenous ligands, co-repressors and co-activators are involved. These factors combined make the biochemical pathways of NR-controlled gene expression complex beyond measure. Despite the complexity, NR drug discovery has proven to be very successful so far. In fact, 13% of all FDA-approved drugs on the market in 2005 targeted NRs⁸ and estimations from 2017 state 16%.⁹

Figure 23: Structural overview of Nuclear receptors represented by the Estrogen (ER) receptor

6.2 Liver X receptors and biological functions

The liver X receptors (LXRs) are ligand-activated transcription factors and a members of the NR superfamily.¹⁰ Two isoforms of the receptors are known, LXR α (encoded by gene *NR1H3*) and $-\beta$ (*NR1H2*) and the biological pathways they regulate depend on activation by endogenous oxysterols (discussed below, see Figure 26, page 88).¹¹⁻¹² Expression of LXR α is mainly restricted to tissues known to play an important role in lipid metabolism, such as liver, adipose tissue, macrophages, kidney, skeletal muscle and small intestine.¹³ LXR α was the first isotype to be cloned and shown to be highly expressed in hepatocytes, hence its name.¹⁴ The β -isoform is however expressed more widely and unselectively and has been identified in most tissues throughout the human body.¹⁵⁻¹⁷ The two isoforms α and $-\beta$ are highly similar and share 74% of the same amino acid sequence in the ligand binding domain, only differing by two amino acid residues at the ligand binding site.¹⁸

LXRs play a crucial role in the metabolism and homeostasis of cholesterol, lipids, bile acids and carbohydrates.¹⁹⁻²² Therefore, their potential to serve as drug targets is immense. LXR drug discovery has particularly been pursued in diseases like atherosclerosis,²³ diabetes mellitus type 2 (T2DM) and obesity,²⁴⁻²⁵ cancer,²⁶⁻²⁷ inflammation²⁸, skin disorders²⁹ and Alzheimer's disease.³⁰⁻³¹ These potential indications stem from the genes that LXRs regulate the expression of.

The LXRs form an obligatory heterodimer with Retinoid X Receptor (RXR) that binds to a specific LXR responsive element (LXRE) characterized direct by repetitions separated by four nucleotides spaced in between the repetitions. Conformational changes of this heterodimer-complex are promoted by ligands that bind to the binding pockets of LXR (oxysterols), RXR (9-*cis*-retinoic acid) or both (see Figure 24, next page). This structural change leads to release of co-repressor proteins that inhibit gene transcription and promote assembly of co-activators that induce gene transcription.¹⁴ RXR is special among NRs since it serves as a heterodimeric partner with many other NR types including farnesoid X receptors (FXRs) and peroxisome proliferator-activated receptors (PPARs).³² The heterodimers can be further classified as permissive or non-permissive, all depending on the role RXR plays in the dimer complex – either as an active transcription partner or a silent one (Figure 24).^{2, 32-33}

Figure 24: LXR regulated gene transcription in heterodimeric partnership with RXR.

6.2.1 Target genes and therapeutic potentials.

The LXRs regulate expression for cohorts of genes involved in lipid metabolism (a selected overview is presented in Table 7, next page). Moreover, novel genes expressed upon LXR activation are still being discovered.^{10, 34} Agonists of LXRs promote effects in the liver that upregulate proteins of the ATP-binding cassette (ABC) superfamily of transporter proteins and others of well-documented importance in lipid metabolism, like sterol regulatory element-binding protein-1c gene (SREBP1c), fatty acid synthase (FASN) and stearyl-CoA desaturase (SCD).³⁵⁻³⁶ Additionally, lipidation of apolipoprotein E (ApoE) which is highly expressed in the human brain and is linked to pathogenesis of Alzheimer's disease is LXR-regulated.³⁰

Reverse cholesterol transport (RCT) is a process where excess cholesterol from peripheral tissues is transported back to the liver for secretion and elimination through bile.³⁴ Several genes regulated by LXRs are highly involved in this process, key players being the ABC-transport proteins. Members of this family are expressed in an array of tissues, such as ABCG5/G8 in liver and intestine³⁷ and ABCA1/G1 in macrophages.³⁸⁻³⁹ Induced expression of ABCA1 and ABCG1 upon oxysterol LXR-activation suggest important roles in macrophage lipid loading, a process important for avoiding atherosclerosis.³⁸

Table 7. Important target genes of LXR in metabolism of lipids and cholesterol (recreated from Edwards et al.)⁴⁰

Target genes	Human/ Mouse	LXR responsive element	Function	Ref
↑ SREBP-1C	H, M	GGGTTA nx4 CGGTCA	Fatty acid synthesis	Repa et al. ⁴¹
↑ FASN	H, M	GGGTTA nx4 CGGTCA		Joseph et al. ⁴²
↑ CYP7A1	M	TGGTCA nx4 AGTTCA	Cholesterol clearance	Lehmann et al. ⁴³
↑ ApoE	H, M	GGGTCA nx4 CGGTCA		Laffitte et al. ⁴⁴
↑ LXRα	H	AGGTTA nx4 TGGTCA	Lipid homeostasis (transcriptional control)	Laffitte et al. ⁴⁵ Whitney et al. ⁴⁶
↑ CETP	H, M	GGGTCA nx4 CGGGCA	Triglycerides metabolism	Luo et al. ⁴⁷
↑ LPL	H, M	TGGTCA nx4 CGGTCA		Zhang et al. ⁴⁸
↑ ABCA1	H, M	AGGTTA nx4 CGGTCA	Cholesterol and phospholipid efflux	Costet et al. ⁴⁹
↑ ABCG1	H, M	TGGTCA nx4 AGTTCA AGTTTA nx4 AGTTCA		
↑ ABCG5	H	-	Cholesterol and phytosterol efflux	Berge et al. ⁵⁰
↑ ABCG8	H	-		
↑ ABCG4	H	-		Engel et al. ⁵¹

The process of LXR-induced gene expression (by agonists) of ABCA1, ABCG5 and ABCG8 also prevents absorption of cholesterol from the gastrointestinal tract, resulting in a net loss of cholesterol from the body. These LXR-involved features point towards a promising therapeutic application in terms of hypercholestermia.³⁷ Moreover, mice that lack ABCG5 or ABCG8 have been demonstrated to contain increased cholesterol concentrations in the liver and reduced levels of bile acids, further suggesting a crucial role in secretion of sterols through bile.⁵²

Steroyl-CoA desaturases (SCDs) are rate-limiting enzymes in cellular biosynthesis of monounsaturated fatty acids from saturated fatty acids - a critical step in production of triglycerides. Chronic upregulation of SCD activity (provided by LXR-agonists) is followed by increased synthesis of triglycerides and esters of cholesterol that are transported out of the liver. This process leads to increased plasma concentration of lipids followed by elevated risks of cardiovascular diseases.⁵³

Statins are the current choice of medicine used to treat hypocholesteremia. Statins, including simvastatin and atorvastatin as examples, inhibit HMG-CoA-reductase, the rate limiting step in cholesterol biosynthesis. Although statins decrease cholesterol concentrations in the blood circulation, they do not provide reversible effects on established atherosclerotic plaques. The liver X receptors therefore provide an exciting alternative to target cholesterol metabolism as they upregulate genes involved in reverse cholesterol transport and inhibit absorption of cholesterol from the intestines.¹⁸

The development of mice lacking LXR (LXR-knockout mice) has greatly enhanced our understanding concerning LXRs. For example, critical involvement of LXR α in cholesterol homeostasis was demonstrated by the group of Mangelsdorf et al. utilizing LXR α knockout mice.²² Therein, LXR α lacking mice appeared healthy and normal as long as they were exposed to diet containing <0.02% cholesterol. This was determined by both histological and morphological studies in combination with measurement of factors such as hepatic- and serum cholesterol and lipoproteins.²² When these mice were fed cholesterol-rich diet (2% cholesterol), normal wild-type mice were shown to compensate for this cholesterol increment by upregulation of cholesterol 7 α -hydroxylase (CYP7A1), another important LXR target gene. This enzyme converts cholesterol to 7 α -hydroxycholesterol,⁵⁴ the first step in bile acid synthesis and a rate-limiting one (Figure 25). Upregulation of this enzyme led to increased production of cholesterol-derived bile acids that were eliminated by the fecal route. In mice lacking LXR α however, they were not found able to increase their bile acid synthesis and elimination, leading to accumulation of hepatic cholesteryl esters, that eventually led to liver failure.²²

Figure 25: The rate limiting step of bile acid synthesis

A similar knockout study on mice that do not express LXR β was conducted by Alberti and co-workers. Results from therein showed these mice to be able to resist cholesterol rich diet, strongly suggesting that LXR α and LXR β do not serve comparable roles in hepatic cholesterol metabolism.⁵⁵

LXRs have also been shown to play important roles of lipid- and carbohydrate metabolism in muscle cells.⁵³ Their role in carbohydrate metabolism specifically is complex, as they interfere with- and influence multiple factors, like target genes that promote expression of key-enzymes, both directly and indirectly. This literature is highly complex, but the role of LXRs in carbohydrate metabolism has been elegantly reviewed by Gustavsson et al.⁵³ Stulnig et al. showed that down-regulation of essential enzymes of gluconeogenesis in liver possibly point towards beneficial effects of LXR-agonists for the treatment of T2DM.⁵⁶ Feeding mice with a LXR-agonist has also demonstrated significant inhibition of enzyme expression of key players in gluconeogenesis including the enzymes phosphoenolpyruvate carboxykinase (PECK), fructose biphosphatase-1 and glucose-6-phosphatase.⁵⁶⁻⁵⁷

Several LXR target genes like SCD and SREBP1c are not only LXR-regulated but are also affected by insulin as well.⁵³ Further, Tobin et al. demonstrated that LXR α expression is upregulated by insulin in liver cells.⁵⁸

Gustavsson et al. concluded that when LXR-mediated mechanisms are defected or impaired, they promote higher risk for diabetes in most cases. The LXRs are therefore a very interesting drug target for drug development and research concerning T2DM and its “twin disease” obesity.⁵³

6.2.2 Ligands

A total of seven LXR α and eleven LXR β crystal structures are available currently in the protein data-bank (PDB). All these structures include an agonist co-crystallized within the ligand binding domain. These structures vary considerably as they differ in co-crystallized ligands (steroidal or non-steroidal) and heterodimer partnership with RXR. As elegantly reviewed by Tice et al.,¹⁸ all of the reported structures include a ligand occupying the ligand binding site between helix 3, and this allows helix 12 to adapt agonistic conformation and press against the LBD to allow co-activator binding.¹⁸

6.2.2.1 Natural ligands – Oxysterols

A wide array of ligands, both natural and synthetic, have been explored in terms of LXR-modulation and such compounds have been reviewed elsewhere.^{18, 59} The structural diversity of compounds that bind and affect LXRs also reflects how large and flexible the ligand binding pocket of LXRs are compared with other NRs.^{18, 59} Drug discovery involving LXRs has also suffered from lack of ligands that differentiate in selectivity between the two isoforms α and β that are highly similar at the ligand binding site.¹⁸

The discovery of oxysterols as endogenous ligands marked a breakthrough in LXR-research demonstrating compounds like 22(*R*)-hydroxycholesterol, 24(*S*)-hydroxycholesterol, 24(*S*)-25-epoxycholesterol and 27-hydroxycholesterol to bind LXR receptors and activate them (see Figure 26).^{12, 60-61}

Figure 26: Examples of endogenous oxysterols that agonize LXRs

6.2.2.2 Synthetic agonists

Potent full-agonists of LXR like **T0901317** and **GW3965** (Figure 27) upregulate genes for ABC transporter proteins and therefore display promising effects in terms of anti-atherosclerotic activity. However, they also upregulate factors like FAS and SCD potently. This results in unwanted side effects like hypertriglyceridemia, enhanced lipogenesis and hepatic steatosis. In other words; fatty liver by triglyceride-overload. These side effects have proved particularly difficult to avoid in LXR drug discovery, leaving full LXR agonists like **T0901317** and **GW3965** behind only as valuable research tools that have facilitated discoveries of many LXR regulated genes.^{24, 62}

Interesting LXR modulating effects of *N,N*-dimethyl-3 β -hydroxy-choleamide (**DHMCA**) were reported in 2009 (see structure, Figure 27).⁶³ In contrast to potent LXR-agonists like **T0901317** and **GW3965**, **DHMCA** was shown to circumvent the unwanted side effects of other potent LXR-agonists *in vitro*. Further, **DHMCA** did not increase hepatic triglycerides when administered to apoE-deficient mice. An upregulation of ABCA1, ABCG1 and CYP7A1 was found to occur while mRNA expression of SREBP1c was left intact. Therefore, **DHMCA** was among the first agonistic LXR modulators that promoted selective gene expression of target genes and a promising potential for the LXR-directed treatment of atherosclerosis.⁶⁴

Figure 27: Synthetic LXR-agonists

6.2.2.3 22(*S*)-Hydroxycholesterol and selective LXR-antagonism

As emphasized in sections 6.2.1 and 6.2.2.2, LXR agonists are attractive therapeutic targets in terms of anti-atherosclerotic effects. The pharmaceutical industry has therefore mostly focused its efforts towards the discovery of new LXR-agonists. Very few natural and synthetic compounds have been identified to antagonize LXRs. These compounds are anticipated first and foremost to enhance our understanding of the physiological roles of the LXRs.¹⁰

Research on 22(*S*)-Hydroxycholesterol (**22SHC**) revealed somewhat of mixed results to begin with. Earlier findings concluded **22SHC** to be inactive,⁶⁵⁻⁶⁶ although binding the receptor with

Figure 28: 22(*S*)-Hydroxycholesterol, a synthetic LXR-antagonist

high affinity. As stated earlier, the natural epimer **22RHC** (Figure 26, page 88) is a well-known endogenous LXR agonist. More recent studies report that **22SHC** may possess antagonistic effects on LXRs and have the ability to selectively down-regulate the expression of genes involved in lipid formation, that is: Reduce lipid metabolism and increase glucose uptake.¹⁹ These biological findings are of interest as they suggest that a single change in stereochemistry of the steroidal side chain in the 22-position manipulates gene expression of key LXR target genes “oppositely” in many ways. Spencer et al. conducted a SAR study on various oxysterols and showed **22SHC** to antagonize effects of the agonist **24S25EHC** (Figure 26).⁶⁷ Based on *in silico* studies, the unnatural stereo center at the 22-position is on the “wrong” side, thus not being able to form a hydrogen bond with Tryptophan residue 433 (Trp433) in the ligand binding pocket. Instead, **22SHC** was anticipated to rather create a hydrogen bond with Histidine 421 (His421), resulting in ligand binding without recruitment of co-activators necessary to initiate gene transcription like a normal LXR-agonist.⁶⁷

Kase et al. reported **22SHC** to counteract the potent agonistic effects of **T0901317** strongly suggesting an antagonistic effect for **22SHC**.¹⁹ In addition, the authors showed **22SHC** to reduce the formation of diacylglycerol, cholesteryl esters and free cholesterol from acetate below baseline in human myotubes, at the same time as glucose uptake and oxidation was enhanced.¹⁹ The same authors further conducted an *in vivo* study where **22SHC** was administered to rats receiving either standard chow diet or high-fat diet in combination with **22SHC**. Results showed that rats given **22SHC** gained less weight than their control parallels in both cases whether rats were given high fat diet or a regular one. Control rats on high fat diet showed increased levels of triacylglycerols whereas rats on high fat diet in combination with **22SHC** did not.⁶⁸ These findings are of clear interest in terms of new alternatives to target diabetes mellitus type II and obesity.

Recently, Lee et al. showed **22SHC** to hinder development of ethanol-induced fatty liver injury in mice.⁶⁹ The mode of action was concluded to include LXR α as a regulator for expression of monocyte chemoattractant protein-1 (MCP-1), a hypoxia sensing response factor that recruits monocytes to the site of injury.⁶⁹

Chapter 7 - Oxysterols and aim of study

Our interest in this field is largely based on earlier reports stating the synthetic oxysterol **22SHC** to have attractive effects that might be exploited to counteract diabetes and obesity via LXR-modulation.⁶⁸ The **22SHC** has shown LXR-antagonistic features, selectively being capable of down-regulating expression of LXR target genes involved in lipid formation which leads to reduction of *de novo* lipogenesis simultaneously as glucose uptake is increased in skeletal muscle.^{19, 70} The natural endogen epimer **22RHC** is considered a full agonist, and chronic exposure of such compounds leads to unacceptable side-effects, due to constant upregulation of key genes important in lipid synthesis.¹⁰

Since both epimers **22SHC** and **22RHC** possess documented biological effects on LXRs, compounds specifically functionalized in the 22-position of the steroidal scaffold were prioritized in this project. An important parameter for synthesized compounds were *in vitro* evaluations measuring LXR influenced gene expression of key LXR target genes. The avoidance of the synthesized compounds to downregulate expression of ABC-transporters was considered of special importance, as this would increase the risk for development of atherosclerosis. Moreover, they should not upregulate key target genes of LXR related to lipogenesis, such as SCD-1 and FASN. The synthesis of ligands was to be aided by molecular modelling to gain more insight into the binding properties of ligands within the LXR ligand binding domain.

As of overall structural complexity concerning stereochemistry and synthesis of steroids, semisynthetic strategies were to be applied from commercially available reagents.

Figure 29: Numbered positions and ring nomenclature of the cholestane steroid scaffold and the structure of cholesterol

Chapter 8 - Oxysterols results and discussion

8.1 Papers III-V

8.1.1 Synthesis and molecular modelling

In order to synthesize **22SHC** and potential mimics, the commercially available Fernholtz acid (**LXR1**, Scheme 36) was identified as an excellent precursor for synthesis of an array of 22-functionalized steroids. The TBS-aldehyde **LXR5** (Scheme 36), regardless of hydroxyl protection in the 3-position has been utilized for synthesis of multiple steroid compounds for various purposes.⁷¹⁻⁷⁴ This compound was thus considered an important target to serve as a main precursor molecule for other 22-functionalized compounds.

A synthetic route previously published by Guerlet *et. al.*⁷⁵ was repeated starting from Fernholtz acid (**LXR1**) and a Fischer esterification in MeOH gave the Fernholtz-methylester (**LXR2**). Subsequent TBS-protection step gave TBS-Fernholtz methyl ester (**LXR3**), which was further reduced using DIBALH. The last step towards the key aldehyde **LXR5** was achieved by IBX oxidation in EtOAc. This route was undertaken multiple times where the overall yield from **LXR1** to **LXR5** stretched up to 76% over 4 steps. The last step of this path, the IBX oxidation did however prove to be difficult to reproduce. Attempts with Dess-Martin periodinane (DMP)

Scheme 36: Synthesis of TBS-aldehyde **LXR5** by two separate pathways. Reagents and conditions: a) H₂SO₄, MeOH, reflux, 17h, 78%, b) TBS-OTf, 2,6-lutidine, DCM, -40 °C→rt, 91%, c) DIBALH, THF, -78 °C, 98% d) IBX, EtOAc, 96% e) MeNHOMe·HCl, HATU, HOBT, DIPEA, DMF, rt, overnight, 77% f) TBS-OTf, 2,6-lutidine, DCM, -40 °C, 78%, g) DIBALH, THF, -78 °C - rt, 83 %.

at room temperature in DCM resulted in “greasy” TLC plates with multiple inseparable products detected. Swern-oxidation gave full racemization of the methyl group in 21-position judged by $^1\text{H-NMR}$.

Exploration of another route towards the TBS-aldehyde **LXR5** was therefore initiated to circumvent the problematic oxidation step from **LXR4** to **LXR5** (Scheme 36). Fernholtz acid (**LXR1**) was transformed into its corresponding Weinreb amide (**LXR6**) using HATU (1-[Bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxide PF_6^-) as a peptide coupling agent. This step proved highly efficient affording 77% yield on a 1 gram scale. The instant activation of the carboxyl acid could clearly be observed when the poorly soluble Fernholtz acid (**LXR1**) formed an activated ester that resulted in clear transparent solution (see Scheme 37).

To ensure that the 21-position had retained its stereospecific configuration after exposure to basic peptide coupling conditions, the compound was recrystallized from hot EtOAc and the crystal structure determined (Paper 5). The result showed that the 21-position was still intact (see Figure 30, next page for structure of **LXR6** determined with X-ray crystallography).

Scheme 37: A reaction mechanism for the transformation of **LXR1** to **LXR6**

Figure 30: The structure of Weinreb Amide **LXR6** was determined by X-ray crystallography.

The 3-positioned alcohol of the Weinreb amide **LXR6** was TBS-protected in the 2nd step to afford TBS-Weinreb amide **LXR7**, followed by DIBALH reduction of the Weinreb amide which gave the aldehyde **LXR5** upon hydrolysis of the chelate formed in the reaction.

Yields of this pathway proved to be high, robust and reliable. All the steps gave relatively clean reaction mixtures that were easy to purify. In fact, the aldehyde **LXR5** could be obtained in sufficient purity for further synthesis after simple extraction of the aqueous phase.

Scheme 38: Synthesis of 22SHC, 22KC and analogues. Synthesis outline and reaction conditions: a) isopentyl-MgBr, THF, -78 °C, 52% b) TBAF, THF, rt, 67% c) DMP, DCM, rt, 78% d) TBAF, THF, rt, 70% e) *N,N*-Dimethylacetamide, LDA, THF, -78 °C, 73%, then recrystallization from hot EtOAc f) TBAF, THF, rt, 86%. g) DMP, DCM, rt., 69%. h) TBAF, THF, rt., 23%.

Next, TBS-protected 22(*S*)-hydroxycholesterol (**LXR8**) was synthesized by exposure of aldehyde **LXR5** to isopentylmagnesium bromide under standard Grignard conditions. That step gave **LXR8** in 52% yield, and the subsequent TBAF-mediated deprotection step gave **22SHC** in moderate yields (Scheme 41). The asymmetric induction in the Grignard reaction is consistent with typical Felkin-Ahn stereoselectivity. Biological evidence also confirmed at a later stage that the compound was an antagonist (not an agonist like 22RHC). Moreover, the ^1H -NMR of the compound matched perfectly with a purchased standard of 22SHC.

Of special interest was 22-ketocholesterol (**22KC**) since both **22RHC** and **22SHC** activate LXRs in different ways. We anticipated that this molecule would show LXR-modulation of some kind, although not known in advance. Oxidation of **LXR8** with DMP gave **LXR9** in good yield (78%), followed by TBAF deprotection to afford **22KC** (Scheme 41).

Reacting the lithium enolate of *N,N*-dimethylacetamide with aldehyde **LXR5**, gave the β -hydroxy amide **LXR10**, a close structurally related analogue of the known selective agonist **DHMCA** (see Figure 27). The ^{13}C -NMR of **LXR10** showed a diastereomeric ratio of approximately 2:1. The compound could be recrystallized from hot EtOAc and X-ray crystallography revealing the correct *S*-configuration in the 22nd position of **LXR10**. An additional TBAF deprotection step then gave **LXR11**. DMP oxidation of **LXR10** gave the β -keto amide **LXR12**. Part of the synthesis depicted in scheme 38 was also undertaken by another member of the group. An unsaturated analogue of **LXR11** (and **DHMCA**) was also prepared through standard Wittig chemistry. This molecule, **LXR14** (Scheme 39) was supplied by Synthetica AS.

Scheme 39: Synthesis of **LXR14**. Reaction conditions: a) (Ethoxycarbonylmethyl)-triphenylphosphonium bromide, b) 1) LiOH, THF, MeOH, H₂O, 0 °C. 2) DIPEA, dimethyl hydrochloride, HATU, HOBT, DMF, 0 °C, 35%

Molecular modelling was undertaken as a part of this project where a catalogue of steroid based ligands constituting of different side chains was plotted and docked into the ligand binding pocket of LXR β (PDB code: 1P8D).⁷⁶ The choice of this crystal structure for molecular modelling is based on structural similarity of the ligand complexed in this structure, that is the oxysterol compound **24,25EHC** (see Figure 26). In addition, the ligand binding pocket of LXR α and β are highly similar as earlier explained, where difference in amino acid sequence at the ligand binding site is only 2 amino acid residues.¹⁸ A selected overview of compounds and docking scores is given within table 8 after docking of structures into the LXR β ligand binding pocket after removal **24,25EHC**.

As observed in Table 8, the two last amide compounds **LXR15** and **LXR16** afforded good docking scores, better or comparable with known modulators such as **22SHC** and **22RHC**. The Fernholtz isobutyl amide **LXR15** and Fernholtz benzylamide **LXR16** were synthesized among other amides (**LXR17-LXR20**) as depicted in Scheme 40 (next page). These compounds were considered rather attractive due to a simple one-step synthetic protocol developed for the Weinreb amide **LXR6** (see Scheme 36 and Scheme 37).

Table 8. Docking scores for selected compounds in LXR β (1P8D)

Compound	Chemical structure	Docking score (LXR β)	Compound	Chemical structure	Docking score (LXR β)
T0901317		-22.60	LXR11		-42.00
22RHC		-36.76	LXR14		-42.61
22SHC		-35.49	LXR15		-36.65
22KC		-38.91	LXR16		-40.68

Scheme 40: Synthesis of LXR amides in one step. Reagents and conditions. a) corresponding R-NH₂, HBTU, HOBt, DIPEA, DMF, rt.

8.1.2 Biological evaluations – paper III

This section summarizes the biological findings that were achieved in the first phase of this project, including *in vitro* studies of selected compounds and *in vivo* study of **LXR11**.

Initial biological efforts involved screening of the newly synthesized compounds for lipogenesis in human myotubes and HepG2 liver cells. Compounds **22SHC**, **LXR11** and **LXR14** were shown to down-regulate de novo lipogenesis in myotubes relative to control. Moreover, all three compounds were found to neutralize the potent agonistic effects provided by **T0901317** in these cells. This is a strong indication that the oxysterol compounds **22SHC**, **LXR11** and **LXR14** compete with **T0901317** for the same point in the ligand binding pocket of the LXR receptor. In addition, compound **LXR11** and **LXR14** were shown to reduce lipogenesis in HepG2 liver cells in a dose-responsive manner (see Figure 31). After lipogenesis screening, the most promising compounds were further evaluated in terms of three well-known LXR-target genes; FASN, SCD1 and ABCA1. As expected, the results showed **T0901317** to act as a potent LXR-agonist for upregulation of all three target genes. Compounds **LXR11** and **LXR14** were shown to have similar gene regulating profile to that of **22SHC**. In particular, the

LXR11 showed attractive effects, by slightly increasing expression of ABCA1 while significantly reducing de novo lipogenesis. As emphasized earlier, ABCA1 is important for reverse cholesterol transport so it was considered essential that expression of this gene would not be down-regulated, hence avoiding development of atherosclerotic events in humans and rodents.

If **22SHC** and the mimics **LXR11** and **LXR14** were true agonists, they would strongly enhance the expression of key LXR genes like ABCA1, SCD1 and FASN. The mechanisms behind why they do bind LXR-receptors but do not enhance expression of ABCA1, SCD1 and FASN like true agonists are not understood. Nevertheless, these consequences are likely to result from different recruitment/release profiles of co-activators and co-repressors attached to the LXR-RXR heterodimer complex in comparison to full-agonists such as **22RHC** and **T0901317**.

Due to similar *in vitro* profile of **22SHC**, **LXR11** was chosen for an *in vivo* rat study. However, **LXR14** was not chosen as of very close structural resemblance to **DHMCA**, a known literature compound. Earlier findings by Kase *et.al.* showed that **22SHC** reduced both gain of body weight and triacylglycerol (TAG) levels in rats given high-fat diet.²⁵ Unfortunately, **LXR11** failed to produce similar effects like **22SHC** *in vivo* (Wistar rats). Earlier, 22SHC had been shown to possess oral bioavailability of ~50%,⁶⁸ but plasma analysis of **LXR11** from samples collected on the last day of the experiment revealed that concentrations of **LXR11** were hardly

Figure 31: De novo lipogenesis in human myotubes and HepG2 cells was reduced by **22SHC**, **LXR11** and **LXR14**. Figures obtained from paper 3.

detectable. The cLogP for **LXR11** (4.6) is considerably lower than of **22SHC** (7.3) (calculated in ChemDraw Ultra 13.0, CamebridgeSoft) which might be an important factor concerning bioavailability. The amide function of **LXR** might also be susceptible to amide hydrolyzing enzymes in the gastrointestinal tract of the rats.

8.1.3 Biological evaluations – paper IV

This section summarizes the biological findings that were made during the second phase of this project, including *in silico* and *in vivo* studies in addition to a SAR-study of compounds with different functionalities in the 22-position of the cholesterol scaffold. To the best of the authors knowledge, 22-Ketocholesterol (**22KC**) has not been evaluated before in terms of LXR-modulation, although 22SHC has shown antagonistic features while 22RHC is a well-known endogenous LXR-agonist.

First screening for lipogenesis in HepG2 cells, **22KC** did not show any effects while **22SHC** behaved as an antagonist as expected. However, when **22KC** was evaluated in terms of expression of key LXR-target genes, it was found to be a potent inducer of ABCA1 expression (~20-fold increase relative to control) (Figure 32A). Further, expression of FASN and SCD1 was mostly left intact by **22KC**, but upregulation of FASN and SCD1 would normally be expected for an LXR-full agonist (see Figure 32B and -C).

Figure 32: Effects of **22SHC**, **22RHC**, **22KC** and **LXR15** on basal and T0901317-induced gene expression in skeletal muscle cells. Human skeletal muscle cells were treated for 4 days, respectively, with DMSO (0.1%) control, 1 μ M **T0901317** and 10 μ M **22SHC**, **22RHC**, **22KC**, **LXR15** $\pm$ 1 μ M **T0901317**. Total RNA was then isolated from human skeletal muscle cells and analyzed by qPCR as described in experimental procedures for paper 4. Gene expressions were normalized to 36B4. Values represent fold change relative to control given as means $\pm$ SEM (n = 3–6). Analyzed LXR target genes were (A) ATP-binding cassette transporter A1 (ABCA1), (B) fatty acid synthase (FASN) and (C) stearoyl-CoA desaturase 1 (SCD1).

Moreover, **22KC**, abolished the effects of **T0901317**-induced expression of FAS and SCD1. These *in vitro* data from skeletal muscle cells strongly suggest that **22KC** should be investigated further in terms of anti-atherosclerotic activity and the compound should be tested in more cell lines and screened for expression of more LXR target genes, starting with expression of ABC family members in liver cells.

The new amide structures depicted in Scheme 40 were also screened for lowered lipogenesis in HepG2 liver cells (Figure 33A). All except one, the Fernholtz-cyclohexylamide **LXR21** (see Figure 33A), proved to be inactive. This molecule was further shown to reduce lipogenesis *in vitro* in a dose-responsive way (see Figure 33B).

Figure 33: *De novo* lipogenesis and effects of **22SHC** and amide analogs on basal and T0901317-induced gene expression. Human skeletal muscle and HepG2 cells were treated for 4 days and 24 h, respectively, with DMSO (0.1%) control, 1 μ M **T0901317** and 10 μ M **22SHC**, **LXR6**, **LXR16**, **LXR22** $\pm$ 1 μ M **T0901317**. Total RNA was then isolated from human skeletal muscle cells and analyzed by qPCR as described in experimental procedures. For lipogenesis, HepG2 cells were incubated with [$1\text{-}^{14}\text{C}$]acetate (1 μ Ci/ml, 100 μ M) for 4 h before lipids were isolated by filtration through hydrophobic MultiScreen® HTS plate. The levels of lipids were determined by scintillation counting. Values represent fold change relative to control for total lipid synthesis given as means $\pm$ SEM n = 4 separate experiments from (A) lipogenesis (B) dose-response for **LXR21**. Values represent fold change relative to control for gene expression is given as means $\pm$ SEM (n = 3–6). Gene expressions were normalized to 36B4. *P < 0.05 vs. control and #P < 0.05 for T0901317 vs. treatment + T0901317.

Figure 35: Compound **LXR16** docked to the ligand binding pocket of LXR β including relevant amino acid side chains. The Ligand Surface (mesh) will be displayed colored by binding property, white = neutral surface, green = hydrophobic surface, blue = hydrogen bonding acceptor potential, and red = hydrogen bond donor potential.

As mentioned before (Table 8, page 97), some of these amides gave good docking scores, especially the benzylic amide **LXR16** (Figure 34). However, they were characterized by higher polarity than for example **22SHC** and **22KC**. cLogP calculations for Fernholtz isobutyl amide **LXR15**, sharing same carbon framework as **22SHC** and **22KC** proved to be lower by a factor of ~1.5-2 (see Figure 34). This might suggest that most of the amides tested are of too high polarity for the very hydrophobic binding pocket of LXR. Additionally, some of these structures were characterized by low solubility in DMSO as in the case of **LXR16**.

Figure 34: Calculated LogP values of compounds **LXR15**, **22KC** and **22SHC**.

8.2 Future perspectives

Based on the findings presented within section 8.1, a keto-functional group in the 22 position (exemplified by **22KC**) can result in increased expression of ABCA1 while expression of other main LXR-target genes was mostly unaffected. This selective upregulation could prove useful to guide rational design of new LXR-modulators that target atherosclerosis.

Robust and efficient synthetic strategies have been established for preparation the key TBS-aldehyde **LXR5** and TBS-Weinreb amide **LXR6**, two valuable steroid precursors allowing many more analogues to be synthesized from these two precursors to form both 22S and 22-keto functionalities, for example using Grignard reagents. Choice of Grignard reagents could further be aided by molecular modelling investigations on more LXR-crystal structures from the protein databank.

Despite promising results showed by **LXR11** *in vitro*, it failed to provide any detectable effects *in vivo*. In fact, measurement of blood plasma showed hardly detectable traces of the compound in samples collected on the last day of the experiment and this result strongly points towards pharmacokinetic problems (ADME). Accordingly, the alcohols of **LXR11** could be protected, for example by converting them into cleavable esters, carbonates or carbamates. Carbamates have been shown to prolong biological half-live of compounds that are rapidly metabolized, such as the carbamate prodrug functionalities of irinotecan and bambuterol.⁷⁷

Compound **22KC** is an endogenous substance, but the biological half-live and other key parameters are unknown. Therefore, the first step would be to conduct more *in vitro* studies, followed by an *in vivo* study in an atherosclerotic rodent model. Further, ketones could be transformed into their corresponding ketoximes to serve as prodrugs. This has been efficiently demonstrated on the drugs ketoprofen and nabumethone to form prodrugs of the ketone-

Scheme 41: Proposed structures. A library of 22-ketostructures and ketoxime prodrugs.

functionalities. Furthermore, the prodrugs were shown to be selectively oxidized to their corresponding ketones by CYP450.⁷⁸

The biological literature of LXR has grown fast over the last 3 decades. Two aspects that would enhance overall and better understanding of these complex functioning structures are: 1) an accessible crystal structure of LXR α or LXR β co-crystallized with an LXR-antagonist since current available structures only represent motifs characterized by agonistic binding and 2) More understanding concerning LXR co-activators and co-suppressors would be of great interest, since these co-factors also have potential to be manipulated by drug molecules and influence the LXRs.

8.3 Conclusions

This chapter briefly summarizes the results obtained from our development and testing of new LXR-modulators that mimic the activity of **22SHC**. Overall, a diverse set of synthetic sterols functionalized at the 22-position have been synthesized and biologically evaluated, aided by molecular modelling. Synthetic methods have been developed to efficiently reach biologically active target molecules such as **22SCH** and **22KC** and other mimics. As a part of this project, more than total of 100 molecules have been synthesized and evaluated biologically. Especially, the synthesis of Fernholtz Weinreb amide **LXR6** proved to be a new important route towards the key intermediate aldehyde **LXR5** and 22-ketone compounds as Grignard reagents can be reacted directly with Weinreb amides to form ketones.

We have shown that the transcription of key LXR-target genes involved in lipid metabolism is influenced greatly by stereochemistry and functionalities in the 22nd position on the cholesterol scaffold. These results might therefore enhance further interest in the use of oxysterols for LXR-related diseases.

8.4 References

1. Germain, P.; Staels, B.; Dacquet, C.; Spedding, M.; Laudet, V., Overview of Nomenclature of Nuclear Receptors. *Pharmacol. Rev.* **2006**, *58* (4), 685-704.
2. Aranda, A.; Pascual, A., Nuclear Hormone Receptors and Gene Expression. *Physiol. Rev.* **2001**, *81* (3), 1269-1304.
3. Gronemeyer, H.; Gustafsson, J.-A.; Laudet, V., Principles for modulation of the nuclear receptor superfamily. *Nat. Rev. Drug. Discov.* **2004**, *3* (11), 950-964.
4. Sladek, F. M., Nuclear receptors as drug targets: New developments in coregulators, orphan receptors and major therapeutic areas. *Expert Opin. Ther. Targets* **2003**, *7* (5), 679-684.
5. Mangelsdorf, D. J.; Thummel, C.; Beato, M.; Herrlich, P.; Schütz, G.; Umesono, K.; Blumberg, B.; Kastner, P.; Mark, M.; Chambon, P.; Evans, R. M., The nuclear receptor superfamily: The second decade. *Cell* **1995**, *83* (6), 835-839.
6. Mirkin, S.; Pickar, J. H., Selective estrogen receptor modulators (SERMs): A review of clinical data. *Maturitas* **2015**, *80* (1), 52-57.
7. Steinmetz, A. C. U.; Renaud, J. P.; Moras, D., Binding of ligands and activation of transcription by nuclear receptors. *Annu. Rev. Biophys. Biomol. Struct.* **2001**, *30*, 329-359.
8. Overington, J. P.; Al-Lazikani, B.; Hopkins, A. L., How many drug targets are there? *Nat. Rev. Drug. Discov.* **2006**, *5* (12), 993-996.
9. Santos, R.; Ursu, O.; Gaulton, A.; Bento, A. P.; Donadi, R. S.; Bologa, C. G.; Karlsson, A.; Al-Lazikani, B.; Hersey, A.; Oprea, T. I.; Overington, J. P., A comprehensive map of molecular drug targets. *Nat. Rev. Drug. Discov.* **2017**, *16* (1), 19-34.
10. Viennois, E.; Mouzat, K.; Dufour, J.; Morel, L.; Lobaccaro, J.-M.; Baron, S., Selective liver X receptor modulators (SLiMs): What use in human health? *Mol. Cell. Endocrinol.* **2012**, *351*, 129-141.
11. Jakobsson, T.; Treuter, E.; Gustafsson, J.-Å.; Steffensen, K. R., Liver X receptor biology and pharmacology: new pathways, challenges and opportunities. *Trends Pharmacol. Sci.* **2012**, *33* (7), 394-404.
12. Janowski, B. A.; Willy, P. J.; Devi, T. R.; Falck, J. R.; Mangelsdorf, D. J., An oxysterol signalling pathway mediated by the nuclear receptor LXR α . *Nature* **1996**, *383* (6602), 728-731.
13. Auboeuf, D.; Rieusset, J.; Fajas, L.; Vallier, P.; Frering, V.; Riou, J. P.; Staels, B.; Auwerx, J.; Laville, M.; Vidal, H., Tissue distribution and quantification of the expression of mRNAs of peroxisome proliferator-activated receptors and liver X receptor- α in humans: no alteration in adipose tissue of obese and NIDDM patients. *Diabetes* **1997**, *46* (8), 1319-27.
14. Willy, P. J.; Umesono, K.; Ong, E. S.; Evans, R. M.; Heyman, R. A.; Mangelsdorf, D. J., LXR, a Nuclear Receptor That Defines a Distinct Retinoid Response Pathway. *Genes Dev.* **1995**, *9* (9), 1033-1045.
15. Viennois, E.; Pommier, A. J. C.; Mouzat, K.; Oumeddour, A.; El Hajjaji, F.-Z.; Dufour, J.; Caira, F.; Volle, D. H.; Baron, S.; Lobaccaro, J.-M. A., Targeting liver X receptors in human health: Deadlock or promising trail? *Expert Opin. Ther. Targets* **2011**, *15* (2), 219-232.
16. de Boussac, H.; Pommier, A. J. C.; Dufour, J.; Trousson, A.; Caira, F.; Volle, D. H.; Baron, S.; Lobaccaro, J.-M. A., LXR, prostate cancer and cholesterol: the Good, the Bad and the Ugly. *Am. J. Cancer Res.* **2013**, *3* (1), 58-69.
17. Jakobsson, T.; Treuter, E.; Gustafsson, J.-Å.; Steffensen Knut, R., Liver X receptor biology and pharmacology: new pathways, challenges and opportunities. *Trends Pharmacol. Sci.* **2012**.
18. Tice, C. M.; Noto, P. B.; Fan, K. Y.; Zhuang, L.; Lala, D. S.; Singh, S. B., The Medicinal Chemistry of Liver X Receptor (LXR) Modulators. *J. Med. Chem.* **2014**, *57* (17), 7182-7205.
19. Kase, E. T.; Thoresen, G. H.; Westerlund, S.; Höjlund, K.; Rustan, A. C.; Gaster, M., Liver X receptor antagonist reduces lipid formation and increases glucose metabolism in myotubes from lean, obese and type 2 diabetic individuals. *Diabetologia* **2007**, *50* (10), 2171-2180.
20. Calkin, A. C.; Tontonoz, P., Transcriptional integration of metabolism by the nuclear sterol-activated receptors LXR and FXR. *Nat. Rev. Mol. Cell Biol.* **2012**, *13* (4), 213-224.
21. Mitro, N.; Mak, P. A.; Vargas, L.; Godio, C.; Hampton, E.; Molteni, V.; Kreusch, A.; Saez, E., The nuclear receptor LXR is a glucose sensor. *Nature* **2007**, *445* (7124), 219-223.

22. Peet, D. J.; Turley, S. D.; Ma, W.; Janowski, B. A.; Lobaccaro, J.-M. A.; Hammer, R. E.; Mangelsdorf, D. J., Cholesterol and Bile Acid Metabolism Are Impaired in Mice Lacking the Nuclear Oxysterol Receptor LXR α . *Cell* **1998**, *93* (5), 693-704.
23. Calkin, A. C.; Tontonoz, P., Liver X Receptor Signaling Pathways and Atherosclerosis. *Arterioscler., Thromb., Vasc. Biol.* **2010**, *30* (8), 1513-1518.
24. Hessvik, N. P.; Bakke, S. S.; Smith, R.; Ravna, A. W.; Sylte, I.; Rustan, A. C.; Thoresen, G. H.; Kase, E. T., The liver X receptor modulator 22(S)-hydroxycholesterol exerts cell-type specific effects on lipid and glucose metabolism. *J. Steroid Biochem. Mol. Biol.* **2012**, *128*, 154-164.
25. Kase, E. T.; Nikolic, N.; Hessvik, N. P.; Fjeldheim, A. K.; Jensen, J.; Thoresen, G. H.; Rustan, A. C., Dietary Supplementation with 22-S-Hydroxycholesterol to Rats Reduces Body Weight Gain and the Accumulation of Liver Triacylglycerol. *Lipids* **2012**, *47* (5), 483-493.
26. Pommier, A. J. C.; Alves, G.; Viennois, E.; Bernard, S.; Communal, Y.; Sion, B.; Marceau, G.; Damon, C.; Mouzat, K.; Caira, F.; Baron, S.; Lobaccaro, J. M. A., Liver X Receptor activation downregulates AKT survival signaling in lipid rafts and induces apoptosis of prostate cancer cells. *Oncogene* **2010**, *29* (18), 2712-2723.
27. Lin, C.-Y.; Gustafsson, J.-A., Targeting liver X receptors in cancer therapeutics. *Nat. Rev. Cancer* **2015**, *15* (4), 216-224.
28. Castrillo, A.; Tontonoz, P., Nuclear receptors in macrophage biology: At the Crossroads of Lipid Metabolism and Inflammation. *Annu. Rev. Cell Dev. Biol.* **2004**, *20* (1), 455-480.
29. Fowler, A. J.; Sheu, M. Y.; Schmuth, M.; Kao, J.; Fluhr, J. W.; Rhein, L.; Collins, J. L.; Willson, T. M.; Mangelsdorf, D. J.; Elias, P. M.; Feingold, K. R., Liver X receptor activators display anti-inflammatory activity in irritant and allergic contact dermatitis models: Liver-X-receptor-specific inhibition of inflammation and primary cytokine production. *J. Invest. Dermatol.* **2003**, *120* (2), 246-255.
30. Jiang, Q.; Lee, C. Y. D.; Mandrekar, S.; Wilkinson, B.; Cramer, P.; Zelcer, N.; Mann, K.; Lamb, B.; Willson, T. M.; Collins, J. L.; Richardson, J. C.; Smith, J. D.; Comery, T. A.; Riddell, D.; Holtzman, D. M.; Tontonoz, P.; Landreth, G. E., ApoE Promotes the Proteolytic Degradation of A β . *Neuron* **2008**, *58* (5), 681-693.
31. Riddell, D. R.; O'Neill, D. J., Chapter 24 Targeting ApoE in Alzheimer's Disease: Liver X Receptor Agonists as Potential Therapeutics. In *Emerging Drugs and Targets for Alzheimer's Disease: Volume 2: Neuronal Plasticity*, The Royal Society of Chemistry: **2010**; Vol. 2, pp 191-212.
32. Evans, Ronald M.; Mangelsdorf, David J., Nuclear Receptors, RXR, and the Big Bang. *Cell.* **2014**, *157* (1), 255-266.
33. de Lera, Á. R.; Krezel, W.; Rühl, R., An Endogenous Mammalian Retinoid X Receptor Ligand, At Last! *Chem. Med. Chem.* **2016**, *11* (10), 1027-1037.
34. Hong, C.; Tontonoz, P., Liver X receptors in lipid metabolism: opportunities for drug discovery. *Nat. Rev. Drug. Discov.* **2014**, *13* (6), 433-444.
35. Joseph, S. B.; Laffitte, B. A.; Patel, P. H.; Watson, M. A.; Matsukuma, K. E.; Walczak, R.; Collins, J. L.; Osborne, T. F.; Tontonoz, P., Direct and indirect mechanisms for regulation of fatty acid synthase gene expression by liver X receptors. *J. Biol. Chem.* **2002**, *277* (13), 11019-25.
36. Repa, J. J.; Liang, G.; Ou, J.; Bashmakov, Y.; Lobaccaro, J. M.; Shimomura, I.; Shan, B.; Brown, M. S.; Goldstein, J. L.; Mangelsdorf, D. J., Regulation of mouse sterol regulatory element-binding protein-1c gene (SREBP-1c) by oxysterol receptors, LXR α and LXR β . *Genes Dev.* **2000**, *14* (22), 2819-30.
37. Repa, J. J.; Berge, K. E.; Pomajzl, C.; Richardson, J. A.; Hobbs, H.; Mangelsdorf, D. J., Regulation of ATP-binding Cassette Sterol Transporters ABCG5 and ABCG8 by the Liver X Receptors α and β . *J. Biol. Chem.* **2002**, *277* (21), 18793-18800.
38. Venkateswaran, A.; Laffitte, B. A.; Joseph, S. B.; Mak, P. A.; Wilpitz, D. C.; Edwards, P. A.; Tontonoz, P., Control of cellular cholesterol efflux by the nuclear oxysterol receptor LXR α . *Proc. Natl. Acad. Sci. U. S. A.* **2000**, *97* (22), 12097-12102.
39. Kennedy, M. A.; Venkateswaran, A.; Tarr, P. T.; Xenarios, I.; Kudoh, J.; Shimizu, N.; Edwards, P. A., Characterization of the Human *ABCG1* Gene: Liver X receptor activates and internal

- promoter that produces a novel transcript encoding an alternative form of the protein. *J. Biol. Chem.* **2001**, *276* (42), 39438-39447.
40. Edwards, P. A.; Kennedy, M. A.; Mak, P. A., LXRs; Oxysterol-activated nuclear receptors that regulate genes controlling lipid homeostasis. *Vasc. Pharmacol.* **2002**, *38* (4), 249-256.
 41. Repa, J. J.; Liang, G.; Ou, J.; Bashmakov, Y.; Lobaccaro, J.-M. A.; Shimomura, I.; Shan, B.; Brown, M. S.; Goldstein, J. L.; Mangelsdorf, D. J., Regulation of mouse sterol regulatory element-binding protein-1c gene (SREBP-1c) by oxysterol receptors, LXR α and LXR β . *Genes Dev.* **2000**, *14* (22), 2819-2830.
 42. Joseph, S. B.; Laffitte, B. A.; Patel, P. H.; Watson, M. A.; Matsukuma, K. E.; Walczak, R.; Collins, J. L.; Osborne, T. F.; Tontonoz, P., Direct and Indirect Mechanisms for Regulation of Fatty Acid Synthase Gene Expression by Liver X Receptors. *J. Biol. Chem.* **2002**, *277* (13), 11019-11025.
 43. Lehmann, J. M.; Kliewer, S. A.; Moore, L. B.; Smith-Oliver, T. A.; Oliver, B. B.; Su, J.-L.; Sundseth, S. S.; Winegar, D. A.; Blanchard, D. E.; Spencer, T. A.; Willson, T. M., Activation of the Nuclear Receptor LXR by Oxysterols Defines a New Hormone Response Pathway. *J. Biol. Chem.* **1997**, *272* (6), 3137-3140.
 44. Laffitte, B. A.; Repa, J. J.; Joseph, S. B.; Wilpitz, D. C.; Kast, H. R.; Mangelsdorf, D. J.; Tontonoz, P., LXRs control lipid-inducible expression of the apolipoprotein E gene in macrophages and adipocytes. *PNAS* **2001**, *98* (2), 507-512.
 45. Laffitte, B. A.; Joseph, S. B.; Walczak, R.; Pei, L.; Wilpitz, D. C.; Collins, J. L.; Tontonoz, P., Autoregulation of the Human Liver X Receptor α Promoter. *Mol. Cell. Biol.* **2001**, *21* (22), 7558-7568.
 46. Whitney, K. D.; Watson, M. A.; Goodwin, B.; Galardi, C. M.; Maglich, J. M.; Wilson, J. G.; Willson, T. M.; Collins, J. L.; Kliewer, S. A., Liver X Receptor (LXR) Regulation of the LXR α Gene in Human Macrophages. *J. Biol. Chem.* **2001**, *276* (47), 43509-43515.
 47. Luo, Y.; Tall, A. R., Sterol upregulation of human CETP expression in vitro and in transgenic mice by an LXR element. *J. Clin. Invest.* **2000**, *105* (4), 513-520.
 48. Zhang, Y.; Repa, J. J.; Gauthier, K.; Mangelsdorf, D. J., Regulation of Lipoprotein Lipase by the Oxysterol Receptors, LXR α and LXR β . *J. Biol. Chem.* **2001**, *276* (46), 43018-43024.
 49. Costet, P.; Luo, Y.; Wang, N.; Tall, A. R., Sterol-dependent Transactivation of the ABC1 Promoter by the Liver X Receptor/Retinoid X Receptor. *J. Biol. Chem.* **2000**, *275* (36), 28240-28245.
 50. Berge, K. E.; Tian, H.; Graf, G. A.; Yu, L.; Grishin, N. V.; Schultz, J.; Kwiterovich, P.; Shan, B.; Barnes, R.; Hobbs, H. H., Accumulation of Dietary Cholesterol in Sitosterolemia Caused by Mutations in Adjacent ABC Transporters. *Science* **2000**, *290* (5497), 1771-1775.
 51. Engel, T.; Lorkowski, S.; Lueken, A.; Rust, S.; Schlüter, B.; Berger, G.; Cullen, P.; Assmann, G., The Human ABCG4 Gene Is Regulated by Oxysterols and Retinoids in Monocyte-Derived Macrophages. *Biochem. Biophys. Res. Commun.* **2001**, *288* (2), 483-488.
 52. Yu, L.; York, J.; von Bergmann, K.; Lutjohann, D.; Cohen, J. C.; Hobbs, H. H., Stimulation of Cholesterol Excretion by the Liver X Receptor Agonist Requires ATP-binding Cassette Transporters G5 and G8. *J. Biol. Chem.* **2003**, *278* (18), 15565-15570.
 53. Steffensen, K. R.; Gustafsson, J.-Å., Putative Metabolic Effects of the Liver X Receptor (LXR). *Diabetes* **2004**, *53* (suppl 1), S36-S42.
 54. Pullinger, C. R.; Eng, C.; Salen, G.; Shefer, S.; Batta, A. K.; Erickson, S. K.; Verhagen, A.; Rivera, C. R.; Mulvihill, S. J.; Malloy, M. J.; Kane, J. P., Human cholesterol 7 α -hydroxylase (CYP7A1) deficiency has a hypercholesterolemic phenotype. *J. Clin. Invest.* **2002**, *110* (1), 109-117.
 55. Alberti, S.; Schuster, G.; Parini, P.; Feltkamp, D.; Diczfalussy, U.; Rudling, M.; Angelin, B.; Björkhem, I.; Pettersson, S.; Gustafsson, J. Å., Hepatic cholesterol metabolism and resistance to dietary cholesterol in LXR β -deficient mice. *J. Clin. Invest.* **2001**, *107* (5), 565-573.
 56. Stulnig, T. M.; Steffensen, K. R.; Gao, H.; Reimers, M.; Dahlman-Wright, K.; Schuster, G. U.; Gustafsson, J.-Å., Novel Roles of Liver X Receptors Exposed by Gene Expression Profiling in Liver and Adipose Tissue. *Mol. Pharmacol.* **2002**, *62* (6), 1299-1305.

57. Stulnig, T. M.; Oppermann, U.; Steffensen, K. R.; Schuster, G. U.; Gustafsson, J.-Å., Liver X Receptors Downregulate 11 β -Hydroxysteroid Dehydrogenase Type 1 Expression and Activity. *Diabetes* **2002**, *51* (8), 2426-2433.
58. Tobin, K. A. R.; Ulven, S. M.; Schuster, G. U.; Steineger, H. H.; Andresen, S. M.; Gustafsson, J.-Å.; Nebb, H. I., Liver X Receptors as Insulin-mediating Factors in Fatty Acid and Cholesterol Biosynthesis. *J. Biol. Chem.* **2002**, *277* (12), 10691-10697.
59. Loren, J.; Huang, Z.; Laffitte, B. A.; Molteni, V., Liver X receptor modulators: A review of recently patented compounds (2009 – 2012). *Expert Opin. Ther. Pat.* **2013**, *23* (10), 1317-1335.
60. Forman, B. M.; Ruan, B.; Chen, J.; Schroepfer, G. J., Jr.; Evans, R. M., The orphan nuclear receptor LXRA is positively and negatively regulated by distinct products of mevalonate metabolism. *Proc. Natl. Acad. Sci. U. S. A.* **1997**, *94* (20), 10588-93.
61. Janowski, B. A.; Willy, P. J.; Devi, T. R.; Falck, J. R.; Mangelsdorf, D. J., An oxysterol signalling pathway mediated by the nuclear receptor LXR alpha. *Nature* **1996**, *383* (6602), 728-31.
62. Hessvik, N. P.; Bakke, S. S.; Fredriksson, K.; Boekschoten, M. V.; Fjoerkenstad, A.; Koster, G.; Hesselink, M. K.; Kersten, S.; Kase, E. T.; Rustan, A. C.; Thoresen, G. H., Metabolic switching of human myotubes is improved by n-3 fatty acids. *J. Lipid Res.* **2010**, *51*, 2090-2104.
63. Kratzer, A.; Buchebner, M.; Pfeifer, T.; Becker, T. M.; Uray, G.; Miyazaki, M.; Miyazaki-Anzai, S.; Ebner, B.; Chandak, P. G.; Kadam, R. S.; Calayir, E.; Rathke, N.; Ahammer, H.; Radovic, B.; Trauner, M.; Hoefler, G.; Kompella, U. B.; Fauler, G.; Levi, M.; Levak-Frank, S.; Kostner, G. M.; Kratky, D., Synthetic LXR agonist attenuates plaque formation in apoE^{-/-} mice without inducing liver steatosis and hypertriglyceridemia. *J. Lipid Res.* **2009**, *50* (2), 312-326.
64. Thomas, P.; Marlene, B.; Prakash, G. C.; Jay, P.; Adelheid, K.; Sascha, O.; Gerald, N. R.; Rajendra, S. K.; Uday, B. K.; Gerhard, M. K.; Dagmar, K.; Sanja, L.-F., Synthetic LXR Agonist Suppresses Endogenous Cholesterol Biosynthesis and Efficiently Lowers Plasma Cholesterol. *Curr. Pharm. Biotechnol.* **2011**, *12* (2), 285-292.
65. Janowski, B. A.; Grogan, M. J.; Jones, S. A.; Wisely, G. B.; Kliewer, S. A.; Corey, E. J.; Mangelsdorf, D. J., Structural requirements of ligands for the oxysterol liver X receptors LXRA and LXR β . *Proc. Natl. Acad. Sci. U. S. A.* **1999**, *96* (1), 266-271.
66. Lehmann, J. M.; Kliewer, S. A.; Moore, L. B.; Smith-oliver, T. A.; Oliver, B. B.; Su, J.-l.; Scott, S.; Winegar, D. A.; Daniel, E.; Spencer, T. A.; Willson, M.; Lehmann, M.; Sundseth, S. S.; Blanchard, D. E.; Willson, T. M., Activation of the Nuclear Receptor LXR by Oxysterols Defines a New Hormone Response Pathway. *J. Biol. Chem.* **1997**, *272*, 3137-3140.
67. Spencer, T. A.; Li, D.; Russel, J. S.; Collins, J. L.; Bledsoe, R. K.; Consler, T. G.; Moore, L. B.; Galardi, C. M.; McKee, D. D.; Moore, J. T.; Watson, M. A.; Parks, D. J.; Lambert, M. H.; Willson, T. M., Pharmacophore Analysis of the Nuclear Oxysterol Receptor LXRA. *J. Med. Chem.* **2001**, *44* (6), 886-897.
68. Kase, E. T.; Nikolić, N.; Hessvik, N. P.; Fjeldheim, Å.-K.; Jensen, J.; Thoresen, G. H.; Rustan, A. C., Dietary Supplementation with 22-S-Hydroxycholesterol to Rats Reduces Body Weight Gain and the Accumulation of Liver Triacylglycerol. *Lipids* **2012**, *47* (5), 483-493.
69. Na, T.-Y.; Han, Y.-H.; Ka, N.-L.; Park, H.-S.; Kang, Y. P.; Kwon, S. W.; Lee, B.-H.; Lee, M.-O., 22-S-Hydroxycholesterol protects against ethanol-induced liver injury by blocking the auto/paracrine activation of MCP-1 mediated by LXRA. *J. Pathol.* **2015**, *235* (5), 710-720.
70. Åstrand, O. A. H.; Sandberg, M.; Sylte, I.; Görbitz, C. H.; Thoresen, G. H.; Kase, E. T.; Rongved, P., Synthesis and initial biological evaluation of new mimics of the LXR-modulator 22(S)-hydroxycholesterol. *Bioorg. Med. Chem.* **2014**, *22* (1), 643-650.
71. Mikami, K.; Kawamoto, K.; Nakai, T., Stereocontrolled synthesis of either (22S)- or (22R)-hydroxy-23-acetylenic steroid side chains via [2,3]-Wittig sigmatropic rearrangement. *Tetrahedron Lett.* **1985**, *26* (47), 5799-5802.
72. Nagano, H.; Matsuda, M.; Yajima, T., Stereoselective synthesis of 24-alkyl-22-hydroxysterols based on chelation-controlled radical reactions. *J. Chem. Soc., Perkin Trans. 1* **2001**, (2), 174-182.
73. Cui, J. G.; Lin, C. W.; Zeng, L. M.; Su, J. Y., Synthesis of polyhydroxysterols (III): synthesis and structural elucidation of 24-methylenecholest-4-en-3 β ,6 α -diol. *Steroids* **2002**, *67* (13-14), 1015-1019.

74. Gros, L.; Lorente, S. O.; Jimenez; Yardley, V.; Rattray, L.; Wharton, H.; Little, S.; Croft, S. L.; Ruiz-Perez, L. M.; Gonzalez-Pacanowska, D.; Gilbert, I. H., Evaluation of Azasterols as Anti-Parasitics. *J. Med. Chem.* **2006**, *49* (20), 6094-6103.
75. Guerlet, G.; Spangenberg, T.; Mann, A.; Faure, H.; Ruat, M., Synthesis and biological evaluation of desmethylveramiline, a micromolar Hedgehog inhibitor. *Bioorg. Med. Chem. Lett.* **2011**, *21* (12), 3608-3612.
76. Williams, S.; Bledsoe, R. K.; Collins, J. L.; Boggs, S.; Lambert, M. H.; Miller, A. B.; Moore, J.; McKee, D. D.; Moore, L.; Nichols, J.; Parks, D.; Watson, M.; Wisely, B.; Willson, T. M., X-ray Crystal Structure of the Liver X Receptor β Ligand Binding Domain: Regulation by a Histidine-Tryptophan Switch. *J. Biol. Chem.* **2003**, *278* (29), 27138-27143.
77. Ghosh, A. K.; Brindisi, M., Organic Carbamates in Drug Design and Medicinal Chemistry. *J. Med. Chem.* **2015**, *58* (7), 2895-2940.
78. Kumpulainen, H.; Mähönen, N.; Laitinen, M.-L.; Jaurakkajärvi, M.; Raunio, H.; Juvonen, R. O.; Vepsäläinen, J.; Järvinen, T.; Rautio, J., Evaluation of Hydroxyimine as Cytochrome P450-Selective Prodrug Structure. *J. Med. Chem.* **2006**, *49* (3), 1207-1211.

Chapter 9 – Concluding remarks

The intention of the present study has been to identify new drug candidates, either related to DNA-damaging effects in leukemic cells and bacteria, or to candidates that bind to the liver X receptors and consequently influence the transcription of genes relevant to diseases such as atherosclerosis, T2DM and obesity.

We started with a rather broad list of aims which, at least partly, can be related to how complex, unpredictable and multidisciplinary the process of drug discovery can be. New discoveries have been made during this process, mostly as the result of analogue synthesis, biological evaluations and lead optimizations of natural compounds. We have also exploited these discoveries in order to guide a rational development of new drug leads with increased therapeutic potential.

Several new compounds with therapeutic potentials have been identified. For instance, carbamate prodrugs of the natural products iodinin (**5**) and myxin (**8**) have been designed and synthesized, and hold promise as new potential drug candidates, as they were found to possess more attractive physicochemical properties than the natural products themselves, higher activity towards leukemic cells and similar cancer cell selectivity. We also explored structure-activity relationship of the phenazine scaffold in positions 6, 7 and 8. Biological evaluation of these compounds led us to discover new carbamate analogues of 7,8-dimethyl- and 7,8-dichloro-1-hydroxyphenazine 5,10-dioxide (**109** and **111**) that were substantially more potent than iodinin and myxin, and still selective towards leukemic cells. The above mentioned compounds have a potential to enter more advanced pre-clinical trials in the form of animal studies, and funding has already been secured to do so. Intellectual property rights have also been secured for the phenazine compounds presented herein, essential for the transfer of data out of academia and into the industry at a later stage. What is particularly attractive with this class of compounds is that they appear to target leukemic cells selectively and preliminary studies demonstrate that they are considerably less cardiotoxic than the anthracycline daunorubicin, which is a standard part of current AML-induction therapy and which is notorious for its severe cardiotoxicity.

New classes of antibiotics are utterly called for in society due to the ever-emerging threat of antimicrobial resistance to the current last-resort antibiotics. In this context, we felt obligated to test the new phenazine compounds in various strains of resistant bacteria. The antimicrobial

evaluations of the compounds described herein showed clear trends for what kind of phenazine structures should be pursued further in lead development for such a purpose. Alkylated analogues with or without halogens in the 2-position hold most promise at this stage.

Leukemia often leads to fatal infections. Accordingly, we believe that a phenazine 5,10-dioxide compound could be developed, that targets both the leukemia and the infecting bacteria simultaneously. However, more SAR studies are necessary to achieve such a goal.

Our efforts towards targeting the liver X receptors has acquired new knowledge on how key LXR target genes are expressed differently as the result of synthetic modifications in the 22-position of oxysterol ligands. In this context, we identified the compound 22-ketocholesterol (**22KC**) as a selective upregulator of the ABCA1 transport protein. This finding is of particular interest, since LXR agonists usually not only promote an upregulation of ABC-transporter expression, but also of key LXR-target genes such as SCD-1 and FASN. Therefore, LXR full agonists cannot be used in the clinic due to the undesirable side effects they promote through the upregulation of these genes. We believe that **22KC** may be able to bypass these side-effects through a selective upregulation of ABC-transporters and thus guide the development of new therapeutic agents to treat atherosclerosis in a reversible manner, unlike the drug class of the statins. Furthermore, other new drug leads for the treatment of T2DM and obesity have been identified among the compounds that mimic the activity of the LXR-antagonist 22(*S*)-hydroxycholesterol. Although one compound was investigated more thoroughly and failed to provide the desired outcome in rats, this new lead could be chemically developed via prodrug strategies, as discussed in the future perspectives section for LXR modulators.

From a broader perspective, we believe the synthetic methods and routes presented in this thesis could prove to be of great value for other researchers within the field of ligand-dependent LXR modulation and in that of phenazines. For instance, very few SAR-studies have been reported on phenazine 5,10-dioxides to date and we believe this may be related to the lack of efficient synthetic protocols to oxidize the phenazine scaffold. To reflect on this note, we have developed a synthetic method that can afford iodinin in 76% yield on scales exceeding one gram. We have also developed a new synthetic route to the TBS-Fernholtz Weinreb amide, which can serve as a common steroid precursor for multiple 22(*S*)- and 22-keto-functionalized oxysterols. In an overall context, the above mentioned synthetic methods might pave the way for more diverse classes of phenazine 5,10-dioxides and LXR modulators in the future.

Our work is highly dependent on biological assays to evaluate the compounds we synthesize in our projects. These results could therefore only have been achieved through a close collaboration with biochemists, pharmacologist, chemists and structural biologists. Some of these results have generated intellectual property rights (IPR), resulting in patent applications. This is essential to ensure a successful transfer of these discoveries out of academia for further research and development in the industry

Given the diversity of the work presented in this thesis, there are no other words that could summarize this work better than *drug discovery*.

Appendix A

Experimental procedures from section 5.3.1

Appendix A

Experimental procedures from section 5.3.1

General procedure A1, methylation of 1-hydroxyphenazine 5,10-dioxides (synthesis of compounds 30, 125, 126 and 127)

Either iodomethane or dimethyl sulfate (0.6 mmol, 3 eq.) was added to a stirring solution of a corresponding 1-hydroxyphenazine 5,10-dioxide (0.2 mmol, 1 eq.), K₂CO₃ (0.6 mmol, 3 eq.) and 18-Crown-6 (0.6 mmol, 3 eq.) in DMF (5 mL) under argon atm. The resulting mixture allowed to stir overnight (unless stated otherwise) at rt before it was quenched with H₂O (50 mL) and ~5 drops of HCl (1M aqueous sol.). The aqueous phase was extracted with DCM (3 x 15 mL). Combined organic phases were dried over MgSO₄, filtered and concentrated *in vacuo*. Further purification was undertaken as stated for each individual compound synthesized in accordance to this procedure.

1-methoxyphenazine 5,10-dioxide (30)

Synthesized in accordance to general procedure A-1 starting from compound **42** (15 mg, 0.07 mmol) using MeI as a methylating reagent. Flash column chromatography on silica (1% MeOH/DCM) afforded 10 mg (63%) of an orange solid. R_f: 0.13 (1% MeOH/DCM). ¹H NMR (400 MHz, CDCl₃) δ 8.69 (td, *J* = 8.3, 1.1 Hz, 2H), 8.34 (dd, *J* = 9.0, 1.1 Hz, 1H), 7.86 – 7.74 (m, 2H), 7.67 (dd, *J* = 9.1, 7.9 Hz, 1H), 7.13 – 7.05 (m, 1H), 4.10 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 154.2, 138.5, 137.5, 135.7, 131.8, 131.4, 131.0, 130.1, 120.7, 120.2, 112.0, 110.0, 57.4. HRMS (ESI⁺): Exact mass calculated for C₁₃H₁₀N₂O₃Na [M+Na]⁺:265.0584, found 265.0584 (0.1 ppm).

1-hydroxy-2-iodophenazine 5,10-dioxide (119)

A round bottomed flask was charged with 1-hydroxyphenazine 5,10-dioxide (200 mg, 0.87 mmol) dissolved in DCM (9 mL). NIS (217 mg, 0.96 mmol) was added and the mixture stirred at rt under open air for 2 hours. The reaction mixture was diluted with 20 mL DCM and absorbed onto silica *in vacuo*. Flash column chromatography on silica (100% DCM) afforded 305 mg (98%) of a deep-purple powder. R_f: 0.61 (100% DCM). ¹H NMR (400 MHz, CDCl₃)

δ 15.75 (s, 1H), 8.72 – 8.55 (m, 2H), 8.03 (d, J = 9.4 Hz, 1H), 7.84 (d, J = 9.4 Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 153.54, 141.49, 137.19, 136.13, 133.65, 132.43, 131.92, 125.47, 120.11, 119.59, 109.92, 83.15. HRMS (ESI^+): Exact mass calculated for $\text{C}_{12}\text{H}_7\text{N}_2\text{O}_3\text{INa}$ $[\text{M}+\text{Na}]^+$: 376.9394, found 376.9394 (-0.1 ppm).

1-hydroxy-2-bromophenazine 5,10-dioxide (120)

A round bottomed flask was charged with 1-hydroxyphenazine 5,10-dioxide (200 mg, 0.87 mmol) dissolved in DCM (9 mL). NBS (171 mg, 0.96 mmol) and the mixture stirred at rt under open air for 2 hours. The reaction mixture was diluted with 20 mL DCM and absorbed onto silica *in vacuo*. Flash column chromatography on silica (100% DCM) afforded 196 mg (73%) of a deep-red powder. R_f : 0.34 (100% DCM). ^1H NMR (400 MHz, CDCl_3) δ 15.54 (s, 1H), 8.73 – 8.60 (m, 2H), 7.97 (d, J = 9.6 Hz, 1H), 7.91 – 7.82 (m, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ 151.0, 136.5, 136.4, 136.09, 134.0, 132.5, 131.9, 126.4, 120.2, 119.5, 109.3, 109.0. HRMS (ESI^+): Exact mass calculated for $\text{C}_{12}\text{H}_7\text{N}_2\text{O}_3\text{BrNa}$ $[\text{M}+\text{Na}]^+$: 328.9532, found 328.9533 (-0.1 ppm).

General procedure A2, carbamoylation of 1-hydroxyphenazine 5,10-dioxides (synthesis of compounds 121-124)

A dry round bottomed flask was charged with a corresponding 2-halo-1-hydroxyphenazine 5,10-dioxide (0.17 mmol, 1.0 eq.), DABCO (1.04 mmol, 3-6 eq.)* and a magnetic stir bar under argon atm. The mixed solids were dispersed in anhydrous THF (10 mL) and the resulting mixture cooled on ice water-bath. A corresponding carbamoyl chloride was added (0.52 mmol, 3 eq) before the cooling source was removed and the mixture allowed to stir for a period of 90-180 min. If starting material was still observed after 3 hours from start (judged by TLC analysis typically using 2-5% MeOH/DCM as eluent), DABCO (0.34 mmol, 2.0 eq.) and a carbamoyl chloride (0.17 mmol, 1.0 eq.) were added and the mixture allowed to rotate at rt for 1-2 hours. The resulting mixture was diluted with 50 mL NaHCO_3 (saturated aqueous sol.) and the aqueous layer extracted with DCM (4 x 20 mL). Combined organic phases were washed with brine, dried over MgSO_4 and filtered before solvents were removed *in vacuo*. Obtained crude material was further purified as stated for each individual compound.

*6 eq. of DABCO were used if the corresponding carbamoyl chloride was a HCl-salt, otherwise 3 eq.

2-bromo-1-((4-methylpiperazine-1-carbonyl)oxy)phenazine 5,10-dioxide (121)

Synthesized according to general procedure A2 starting from 53 mg (0.17 mmol) of compound **120**. Flash column chromatography on silica (5% MeOH/DCM) afforded 74 mg (>99%) of an orange solid. R_f : 0.20 (5% MeOH/DCM). ^1H NMR (400 MHz, CDCl_3) δ 8.67 – 8.57 (m, 2H), 8.51 (d, J = 9.7 Hz, 1H), 7.89 (d, J = 9.7 Hz, 1H), 7.86 – 7.73 (m, 2H), 4.09 – 3.59 (m, 4H), 2.93 – 2.49 (m, 4H), 2.43 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ 152.0, 141.4, 137.7, 136.9, 135.8, 134.4, 132.3, 131.9, 131.6, 121.1, 120.6, 120.2, 118.7, 54.8, 54.7, 46.3, 45.1, 44.5. HRMS (ESI^+): Exact mass calculated for $\text{C}_{18}\text{H}_{18}\text{N}_4\text{O}_4\text{BrNa}$ $[\text{M}+\text{Na}]^+$: 433.0506, found 433.0506 (0.1 ppm).

2-bromo-1-((pyrrolidine-1-carbonyl)oxy)phenazine 5,10-dioxide (122)

Synthesized according to general procedure A2 starting from 54 mg (0.18 mmol) of compound **120**. Flash column chromatography on silica (1% MeOH/DCM) and subsequent recrystallization from hot EtOH afforded 37 mg (52%) of red-orange glittering flakes. R_f : 0.21 (5% MeOH/DCM). ^1H NMR (300 MHz, CDCl_3) δ 8.74 – 8.60 (m, 2H), 8.52 (d, J = 9.7 Hz, 1H), 7.90 (d, J = 9.7 Hz, 1H), 7.87 – 7.72 (m, 2H), 4.06 – 3.70 (m, 2H), 3.70 – 3.43 (m, 2H), 2.24 – 1.90 (m, 4H). ^{13}C NMR (101 MHz, CDCl_3) δ 151.5, 141.6, 137.9, 137.0, 135.8, 134.5, 132.5, 131.8, 131.5, 121.1, 120.7, 120.3, 118.5, 47.0, 46.9, 26.0, 25.3. HRMS (ESI^+): Exact mass calculated for $\text{C}_{17}\text{H}_{14}\text{N}_3\text{O}_4\text{BrNa}$ $[\text{M}+\text{Na}]^+$: 426.0060, found 426.0060 (0.0 ppm).

2-iodo-1-((4-methylpiperazine-1-carbonyl)oxy)phenazine 5,10-dioxide (123)

Synthesized according to general procedure A2 starting from 27 mg (0.08 mmol) of compound **119**. Flash column chromatography on silica afforded 20 mg (54%) of an orange solid. R_f : 0.16 (5% MeOH/DCM). ^1H NMR (300 MHz, CDCl_3) δ 8.71 – 8.58 (m, 2H), 8.39 (d, J = 9.4 Hz, 1H), 8.08 (d, J = 9.6 Hz, 1H), 7.89 – 7.74 (m, 2H), 4.06 – 3.59 (m, 4H), 2.94 – 2.50 (m, 4H), 2.45 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ 151.9, 144.3, 139.6, 137.7, 137.5, 135.9, 131.9,

131.8, 131.6, 120.8, 120.3, 119.0, 96.9, 54.9, 54.7, 46.3, 45.2, 44.5. (ESI⁺): Exact mass calculated for C₁₈H₁₈N₄O₄I [M+H]⁺: 481.0367, found 481.0366 (0.3 ppm).

2-iodo-1-((pyrrolidine-1-carbonyl)oxy)phenazine 5,10-dioxide (124)

Synthesized according to general procedure A2 starting from 53 mg (0.15 mmol) of compound **119**. Flash column chromatography on silica (1% MeOH/DCM) afforded an orange solid which was dissolved in minimum amount of a mixture of CHCl₃ and heptane. The resulting mixture was then concentrated *in vacuo* removing CHCl₃. The orange precipitate was then filtered and dried affording 35 mg (52%) of orange powder. R_f: 0.38 (5% MeOH/DCM). ¹H NMR (400 MHz, CDCl₃) δ 8.69 – 8.56 (m, 2H), 8.38 (d, *J* = 9.5 Hz, 1H), 8.08 (d, *J* = 9.5 Hz, 1H), 7.90 – 7.72 (m, 2H), 3.97 – 3.74 (m, 2H), 3.71 – 3.49 (m, 2H), 2.22 – 1.92 (m, 4H). ¹³C NMR (101 MHz, CDCl₃) δ 151.4, 144.4, 139.6, 137.7, 137.6, 135.9, 132.0, 131.9, 131.5, 120.8, 120.2, 118.8, 97.0, 47.0, 46.9, 26.0, 25.3. HRMS (ESI⁺): Exact mass calculated for C₁₇H₁₄N₃O₄INa [M+Na]⁺: 473.9921, found 473.9920 (0.2 ppm).

2-iodo-1-methoxyphenazine 5,10-dioxide (125)

Synthesized in accordance to general procedure A-1 starting from compound **119** (29 mg, 0.08 mmol) using dimethyl sulfate as a methylating reagent. Flash column chromatography on silica (100 % DCM) afforded 13 mg (43%) of an orange solid. R_f: 0.13. ¹H NMR (600 MHz, CDCl₃) δ 8.72 – 8.69 (m, 2H), 8.69 – 8.64 (m, 2H), 8.30 (d, *J* = 9.4 Hz, 1H), 8.08 (d, *J* = 9.5 Hz, 1H), 7.86 – 7.79 (m, 2H), 4.08 (s, 3H). ¹³C NMR (151 MHz, CDCl₃) δ 152.7, 140.4, 138.2, 137.7, 135.8, 133.3, 131.9, 131.7, 120.8, 120.3, 117.6, 96.7, 63.2.

2-bromo-1-methoxyphenazine 5,10-dioxide (126)

Synthesized in accordance to general procedure A-1 starting from compound **120** (55 mg, 0.18 mmol) using dimethyl sulfate as a methylating reagent. Flash column chromatography on silica

(0-10% EtOAc/DCM) afforded 26 mg (45%) of an orange solid. R_f : 0.21. ^1H NMR (600 MHz) δ 8.71 – 8.63 (m, 2H), 8.43 (d, J = 9.6 Hz, 1H), 7.87 (d, J = 9.6 Hz, 1H), 7.85 – 7.79 (m, 2H), 4.11 (s, 3H). ^{13}C NMR (151 MHz, CDCl_3) δ 150.1, 137.9, 137.4, 135.7, 135.2, 134.0, 131.8, 131.7, 121.2, 120.6, 120.3, 117.1, 63.1.

1-methoxy-7,8-dimethylphenazine 5,10-dioxide (127)

Synthesized in accordance to general procedure A-1 starting from compound **100** (36 mg, 0.14 mmol) using MeI as a methylating reagent. The reaction time was 3h. Flash column chromatography on silica (1% MeOH/DCM) afforded 26 mg (69%) of an orange solid. R_f : 0.07 (1% MeOH/DCM). ^1H NMR (600 MHz, CDCl_3) δ 8.43 (s, 1H), 8.41 (s, 1H), 8.32 (dd, J = 9.0, 1.1 Hz, 1H), 7.62 (dd, J = 9.0, 7.9 Hz, 1H), 7.05 (d, J = 7.8 Hz, 1H), 4.08 (s, 3H), 2.63 – 2.46 (m, 6H). ^{13}C NMR (151 MHz, CDCl_3) δ 154.1, 143.6, 142.7, 138.0, 136.1, 134.4, 130.8, 129.6, 119.4, 118.9, 112.0, 109.7, 57.4, 20.7, 20.6. HRMS (ESI⁺): Exact mass calculated for $\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$: 293.0897, found 293.0896 (0.1 ppm).

1-(2-(4-(ethoxycarbonyl)piperazin-1-yl)-2-oxoethoxy)phenazine 5,10-dioxide (130)

ethyl piperazine-1-carboxylate 2.9 mL (20 mmol) was dissolved in DCM (30 mL) under argon atm and the resulting solution cooled to 0 °C. To this solution, chloroacetyl chloride (1.4 mL, 18 mmol), dissolved in DCM (40 mL) was added drop-wise via dropping funnel. The resulting mixture was stirred overnight before transferred to a separatory funnel and diluted by 250 mL HCl (0.1 M aqueous sol). The organic phase was collected followed by extraction of the aqueous phase using DCM (3 x 30 mL). The pooled organic phases were washed with brine (200 mL), dried over MgSO_4 and filtered before solvent were removed *in vacuo*. This gave 2.78 g the crude product **129** as a clear transparent viscous oil (approx. 70% purity judged by ^1H NMR). Next, 0.75 g of the crude compound **129** was added to a stirring solution of 1-hydroxyphenazine 5,10-dioxide (**42**) (150 mg, 0.66 mmol), KI (33 mg, 0.20 mmol), K_2CO_3 (274 mg, 1.98 mmol) and 18-Crown-6 (523 mg, 1.98 mmol) in DMF (15 mL). The resulting solution was left stirring overnight at room temperature. The day after, the reaction mixture was diluted with H_2O (200 mL) and 2 mL of HCl (1M aqueous sol.) and the aqueous phase was extracted using DCM (4 x 30 mL). The pooled organic phases were washed with brine (200 mL), dried over MgSO_4 and filtered before solvents were removed *in vacuo*. Flash column chromatography on silica (5% MeOH/DCM) and subsequent recrystallization from hot EtOH gave 84 mg (30% from **42**)

of 1-(2-(4-(ethoxycarbonyl)piperazin-1-yl)-2-oxoethoxy)phenazine 5,10-dioxide (**130**) as a red powder. R_f: 0.16 (5% MeOH/DCM). ¹H NMR (400 MHz, CDCl₃) δ 8.76 – 8.56 (m, 2H), 8.41 (dd, *J* = 9.0, 1.1 Hz, 1H), 7.89 – 7.73 (m, 2H), 7.64 (dd, *J* = 9.0, 7.9 Hz, 1H), 7.29 (dd, *J* = 7.9, 1.2 Hz, 1H), 4.97 (s, 2H), 4.13 (q, *J* = 7.1 Hz, 2H), 3.89 – 3.75 (m, 2H), 3.68 – 3.43 (m, 6H), 1.24 (t, *J* = 7.1 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 166.0, 155.4, 151.7, 138.4, 137.5, 135.8, 131.8, 131.2, 131.1, 130.3, 120.6, 120.3, 114.0, 113.9, 70.6, 61.8, 45.7, 44.2, 43.6, 42.2, 14.8.

^1H - and ^{13}C -NMR spectra of synthesized compounds

(Appendix A)

Figure A-1: ¹H-NMR spectra of compound 30

Figure A-2: ¹³C-NMR spectra of compound 30

Figure A-3: ¹H-NMR spectra of compound **119**

Figure A-4: ¹³C-NMR spectra of compound **119**

Figure A-5: ¹H-NMR spectra of compound 120

Figure A-6: ¹³C-NMR spectra of compound 120

Figure A-7: ¹H-NMR spectra of compound **121**

Figure A-8: ¹³C-NMR spectra of compound **121**

Figure A-9: ¹H-NMR spectra of compound **122**

Figure A-10: ¹³C-NMR spectra of compound **122**

Figure A-11: ^1H -NMR spectra of compound **123**

Figure A-12: ^{13}C -NMR spectra of compound **123**

Figure A-13: ¹H-NMR spectra of compound **124**

Figure A-14: ¹³C-NMR spectra of compound **124**

Figure A-15: ¹H-NMR spectra of compound 125

Figure A-16: ¹³C-NMR spectra of compound 125

Figure A-17: ¹H-NMR spectra of compound 126

Figure A-18: ¹³C-NMR spectra of compound 126

Figure A-19: ¹H-NMR spectra of compound **127**

Figure A-20: ¹³C-NMR spectra of compound **127**

Figure A-21: ¹H-NMR spectra of compound **130**

Figure A-20: ¹³C-NMR spectra of compound **130**

Appendix F

Paper V and supporting information

Crystal structure of (S)-2-[(3S,8S,9S,10R,13S,14S,17R)-3-hydroxy-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1*H*-cyclopenta[*a*]phenanthren-17-yl]-*N*-methoxy-*N*-methylpropanamide (Fernholz Weinreb amide)

Viktorsson, E. Ö.; Åstrand, O. A. H.; Haseeb, R. S.; Görbitz, C. H. and Rongved, P.

Acta. Crystallogr. E. **2015**, 71 (3): 275-277

Crystal structure of (S)-2-[(3S,8S,9S,10R,13S,14S,-17R)-3-hydroxy-10,13-dimethyl-2,3,4,7,8,9,10,11,-12,13,14,15,16,17-tetradecahydro-1H-cyclopenta-[a]phenanthren-17-yl]-N-methoxy-N-methylpropanamide (Fernholz Weinreb amide)

Elvar Ørn Viktorsson,^a Ove Alexander Høgmoen Åstrand,^a Rasha Sabah Haseeb,^a Carl Henrik Görbitz^b and Pål Rongved^{a*}

^aSchool of Pharmacy, University of Oslo, PO Box 1068 Blindern, N-0316 Oslo, Norway, and ^bDepartment of Chemistry, University of Oslo, PO Box 1033 Blindern, N-0315 Oslo, Norway. *Correspondence e-mail: pal.rongved@farmasi.uio.no

Received 15 December 2014
Accepted 27 January 2015

Edited by W. T. A. Harrison, University of Aberdeen, Scotland

Keywords: crystal structure; Liver X receptor; obesity; steroid

CCDC reference: 1045692

Supporting information: this article has supporting information at journals.iucr.org/e

The literature compound 3 β -hydroxy-bisnor-5-cholenic aldehyde is an important intermediate for the synthesis of new modulators of the nuclear oxysterol receptor Liver X. As part of our ongoing search for new LXR antagonists, the title compound, C₂₄H₃₉NO₃, has proven to be an important intermediate in our new synthetic pathway, giving the corresponding aldehyde in high yield and in only three steps from the commercially available 3 β -hydroxy-bisnor-5-cholenic acid. The title amide crystallized with two molecules in the asymmetric unit, linked into helices by O—H...O hydrogen bonds involving the hydroxy and carbonyl groups.

1. Chemical context

In the nuclear receptor (NR) family, the two isoforms of the nuclear oxysterol receptor Liver X (LXR α and LXR β) are emerging new drug targets. They are key players for a number of important processes related to disease, such as metabolic and cardiovascular diseases, lipid metabolism, inflammation and cancer (Steffensen & Gustafsson, 2006; Laffitte *et al.*, 2003). LXR modulators have been investigated as potential drugs in the therapy of cardiovascular diseases, metabolic syndrome, regulation of inflammatory response and immunity, skin diseases and are effective in the treatment of murine models of atherosclerosis, diabetes and Alzheimer's disease (Viennois *et al.*, 2011, 2012; Jakobsson *et al.*, 2012). Further, such agents have been shown to affect anti-inflammatory activity (Zhu & Bakovic, 2008; Zhu *et al.*, 2012; Solan *et al.*, 2011) and cell proliferation in a number of major cancer forms such as LNCaP human prostate cancer cells. (Viennois *et al.*, 2012; Jakobsson *et al.*, 2012). The ligand-binding pocket (LBP) of LXR allows binding of side-chain-oxygenated sterols (OHCs).

Recently, OHCs with a specific stereochemistry at the 23-hydroxylated side-chain carbon have also been shown to regulate the Hedgehog signalling pathway (Hh), a key developmental pathway playing multiple roles in embryonic development, including stem-cell differentiation (Corman *et al.*, 2012). In our drug-design programme, our retrosynthetic analysis for the establishment of synthetic routes to the pharmacophores in different OHCs revealed that the aldehyde analogue of the title compound [Fernholz aldehyde, (II)] is a key compound leading to a number of new library

OPEN ACCESS

Figure 1
(a) The asymmetric unit of (I), showing the two molecules *A* (light grey C atoms and atomic labels included) and *B* (dark C atoms). (b) An overlay between molecules *A* (blue) and *B* (red), with an r.m.s. value of 0.300 Å. H atoms have been omitted in (b).

candidates for biological testing (Åstrand *et al.*, 2014*a,b*). We have now identified the title compound, Fernholz Weinreb amide (I), as a new key intermediate to the Fernholz aldehyde, reducing the number of steps in the stereoselective synthesis. The O-TBDMS-protected Weinreb amide (I) may be used to prepare (II) using DIBALH, transferred to ketones with Grignard reagents or used for other synthetic transformations (Sivaraman *et al.*, 2009; Davies *et al.*, 2013).

2. Structural commentary

The asymmetric unit of (I), with two independent molecules *A* and *B*, is depicted in Fig. 1*a*. The macrocyclic part of (I) is also found in the naturally occurring hormone cholesterol and in close to 250 other steroids in the Cambridge Structural Database (CSD; Version 5.35 of November 2013; Groom & Allen, 2014). The molecular conformation of this part of the molecule is rigid, as shown from the overlay between *A* and *B* in Fig. 1*b*. If the substituent at C17 is not included, the fit

Figure 2
(a) Unit-cell and crystal packing viewed along the *a* axis. The colour coding is as in Fig. 1. The orange circles highlight a series of methyl groups; the blue area shades a hydrogen-bonded chain in shape of a flat helix. The chain, as a pink shape, is shown in more detail in (b) (the view is along the *b* axis).

improves from 0.300 to 0.173 Å. Compound (I) also shares the hydroxy group at C3 with cholesterol, but the *N*-methoxy-*N*-methylpropanamide functionality has not previously been introduced into steroids; only the structure of the parent carboxylic acid has been reported previously (CSD refcode HAHSAL; Kurek-Tyrlik *et al.*, 2004).

3. Supramolecular features

The unit-cell and the molecular packing of (I) are shown in Fig. 2. As a class, steroids display a pronounced tendency to form crystal structures with more than one molecule in the asymmetric unit; *e.g.* for about 35% of the 250 compounds mentioned above. The maximum *Z'* value of 16 is reached for the high-temperature polymorph of cholesterol itself (CHOEST21; Hsu *et al.*, 2002). Compound (I) has a *Z'* value of 2, the two molecules differing in the way the hydroxy groups make intermolecular hydrogen bonds (Table 1). Only the

Table 1
Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O3A—H3A...O3B ⁱ	0.86 (4)	1.93 (4)	2.782 (4)	180 (5)
O3B—H3B...O1A ⁱⁱ	0.83 (4)	1.95 (4)	2.768 (3)	169 (4)

Symmetry codes: (i) $-x + \frac{3}{2}, -y + 1, z - \frac{1}{2}$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$.

carbonyl group of molecule *A* is an acceptor, while the hydroxy groups of the *B* molecules are both donors and acceptors and thus serve to link adjacent *A* molecules along the *a* axis. In this process, stacks of either *A* or *B* molecules along the *a* axis expose all the methyl groups on the outside, giving distinct regions with methyl–methyl interactions (Fig. 2*a*). This is not a common molecular aggregation pattern for steroids, but some related $Z' = 2$ structures were found in the CSD, all hydrates without additional hydrogen-bond donors or acceptors in their C17 substituents (KESNAX: Sheng-Zhi *et al.*, 1990; ZZZNVG01: Jiang *et al.*, 2001; XOSLOH: Subash-Babu *et al.*, 2009).

4. Synthesis and crystallization

Compound (I) (348 mg) was dissolved in a minimum amount of boiling EtOAc (40 ml). The flask containing the solution was wrapped in aluminium foil and left overnight at room temperature to afford colourless crystalline needles.

5. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. Coordinates were refined for hydroxyic H atoms; other H atoms were positioned with idealized geometry with fixed C–H = 0.95 (aromatic), 0.98 (methyl), 0.99 (methylene) or 1.00 Å (methine) Å. U_{iso} values were set to $1.2U_{\text{eq}}$ of the carrier atom, or $1.5U_{\text{eq}}$ for methyl and hydroxy groups.

References

- Åstrand, O. A. H., Gikling, I., Sylte, I., Rustan, A. C., Thoresen, G. H., Rongved, P. & Kase, E. T. (2014*a*). *Eur. J. Med. Chem.* **74**, 258–263.
 Åstrand, O. A. H., Sandberg, M., Sylte, I., Görbitz, C. H., Thoresen, G. H., Kase, E. T. & Rongved, P. (2014*b*). *Bioorg. Med. Chem.* **22**, 643–650.
 Bruker (2013). *APEX2*, *SAINT-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Corman, A., DeBerardinis, A. M. & Hadden, M. K. (2012). *ACS Med. Chem. Lett.* **3**, 828–833.
 Davies, S. G., Fletcher, A. M. & Thomson, J. E. (2013). *Chem. Commun.* **49**, 8586–8598.
 Groom, C. R. & Allen, F. H. (2014). *Angew. Chem. Int. Ed.* **53**, 662–671.
 Hsu, L.-Y., Kampf, J. W. & Nordman, C. E. (2002). *Acta Cryst.* **B58**, 260–264.
 Jakobsson, T., Treuter, E., Gustafsson, J.-Å. & Steffensen, K. R. (2012). *Trends Pharmacol. Sci.* **33**, 394–404.
 Jiang, R.-W., Ma, S.-C., But, P. P.-H. & Mak, T. C. W. (2001). *J. Nat. Prod.* **64**, 1266–1272.
 Kurek-Tyrlik, A., Michalak, K., Urbanczyk-Lipkowska, Z. & Wicha, J. (2004). *Tetrahedron Lett.* **45**, 7479–7482.
 Laffitte, B. A., Chao, L. C., Li, J., Walczak, R., Hummasti, S., Joseph, S. B., Castrillo, A., Wilpitz, D. C., Mangelsdorf, D. J., Collins, J. L., Saez, E. & Tontonoz, P. (2003). *PNAS*, **100**, 5419–5424.

Table 2

Experimental details.

Crystal data	
Chemical formula	C ₂₄ H ₃₉ NO ₃
M_r	389.56
Crystal system, space group	Orthorhombic, $P2_12_12_1$
Temperature (K)	105
a, b, c (Å)	7.7256 (4), 19.0030 (9), 29.8162 (15)
V (Å ³)	4377.3 (4)
Z	8
Radiation type	Mo $K\alpha$
μ (mm ^{−1})	0.08
Crystal size (mm)	0.65 × 0.21 × 0.10
Data collection	
Diffractometer	Bruker D8 Vantage single-crystal CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2013)
$T_{\text{min}}, T_{\text{max}}$	0.852, 1.000
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	44800, 7739, 5760
R_{int}	0.089
$(\sin \theta/\lambda)_{\text{max}}$ (Å ^{−1})	0.596
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.048, 0.098, 1.04
No. of reflections	7739
No. of parameters	511
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ^{−3})	0.21, −0.19

Computer programs: *APEX2* and *SAINT-Plus* (Bruker, 2013), *SHELXS2013* (Sheldrick, 2008), *SHELXL2013* (Sheldrick, 2015) and *Mercury* (Macrae *et al.*, 2008).

- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
 Sheng-Zhi, H., You-Qing, H., Kai-Liang, S., Rui-Sheng, L. & Li, D. (1990). *Jiegou Huaxue (Chin. J. Struct. Chem.)*, **9**, 90.
 Sivaraman, B., Manjunath, B. N., Senthilmurugan, A., Harikrishna, K. & Singh, A. I. (2009). *Indian J. Chem. Sect. B*, **48**, 1749–1756.
 Solan, P. D., Piraino, G., Hake, P. W., Denenberg, A., O'Connor, M., Lentsch, A. & Zingarelli, B. (2011). *Shock*, **35**, 367–374.
 Steffensen, K. R. & Gustafsson, J. Å. (2006). *Future Lipidol.* **1**, 181–189.
 Subash-Babu, P., Ignacimuthu, S., Agastian, P. & Varghese, B. (2009). *Bioorg. Med. Chem.* **17**, 2864–2870.
 Viennois, E., Mouzat, K., Dufour, J., Morel, L., Lobaccaro, J.-M. & Baron, S. (2012). *Mol. Cell. Endocrinol.* **351**, 129–141.
 Viennois, E., Pommier, A. J. C., Mouzat, K., Oumeddour, A., El Hajjaji, F.-Z., Dufour, J., Caira, F., Volle, D. H., Baron, S. & Lobaccaro, J.-M. A. (2011). *Expert Opin. Ther. Targets*, **15**, 219–232.
 Zhu, L. & Bakovic, M. (2008). *Res. Lett. Biochem.* Article ID 801849.
 Zhu, R., Ou, Z., Ruan, X. & Gong, J. (2012). *Mol. Med. Rep.* **5**, 895–900.

supporting information

Acta Cryst. (2015). E71, 275-277 [doi:10.1107/S2056989015001747]

Crystal structure of (S)-2-[(3S,8S,9S,10R,13S,14S,17R)-3-hydroxy-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-17-yl]-N-methoxy-N-methylpropanamide (Fernholz Weinreb amide)

Elvar Örn Viktorsson, Ove Alexander Høgmoen Åstrand, Rasha Sabah Haseeb, Carl Henrik Görbitz and Pål Rongved

Computing details

Data collection: *APEX2* (Bruker, 2013); cell refinement: *SAINT-Plus* (Bruker, 2013); data reduction: *SAINT-Plus* (Bruker, 2013); program(s) used to solve structure: *SHELXS2013* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2015); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL2013* (Sheldrick, 2015).

(S)-2-[(3S,8S,9S,10R,13S,14S,17R)-3-Hydroxy-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-17-yl]-N-methoxy-N-methylpropanamide

Crystal data

C₂₄H₃₉NO₃

M_r = 389.56

Orthorhombic, *P*2₁2₁2₁

a = 7.7256 (4) Å

b = 19.0030 (9) Å

c = 29.8162 (15) Å

V = 4377.3 (4) Å³

Z = 8

F(000) = 1712

D_x = 1.182 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 9981 reflections

θ = 2.3–24.9°

μ = 0.08 mm⁻¹

T = 105 K

Flat needle, colourless

0.65 × 0.21 × 0.10 mm

Data collection

Bruker D8 Vantage single-crystal CCD diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 8.3 pixels mm⁻¹

Sets of exposures each taken over 0.5° ω rotation scans

Absorption correction: multi-scan (*SADABS*; Bruker, 2013)

T_{min} = 0.852, *T_{max}* = 1.000

44800 measured reflections

7739 independent reflections

5760 reflections with *I* > 2σ(*I*)

R_{int} = 0.089

θ_{max} = 25.1°, θ_{min} = 2.3°

h = −9→9

k = −22→22

l = −35→35

*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.048$ $wR(F^2) = 0.098$ $S = 1.04$

7739 reflections

511 parameters

0 restraints

Hydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0448P)^2 + 0.1501P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.21 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. No constraints or restraints applied

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1A	0.8266 (3)	0.67664 (11)	0.71479 (7)	0.0314 (6)
O2A	0.7377 (3)	0.50014 (11)	0.73658 (6)	0.0268 (6)
O3A	0.7896 (3)	0.69062 (13)	0.28850 (7)	0.0321 (6)
H3A	0.893 (6)	0.697 (2)	0.2790 (13)	0.048*
N1A	0.8127 (4)	0.56722 (14)	0.74205 (8)	0.0262 (7)
C1A	0.6249 (5)	0.67785 (17)	0.40586 (10)	0.0234 (8)
H1A	0.5099	0.6633	0.4171	0.028*
H2A	0.6438	0.7272	0.4153	0.028*
C2A	0.6223 (5)	0.67502 (18)	0.35440 (10)	0.0254 (8)
H4A	0.5928	0.6268	0.3444	0.030*
H5A	0.5327	0.7075	0.3428	0.030*
C3A	0.7964 (5)	0.69551 (17)	0.33605 (10)	0.0250 (8)
H31A	0.8210	0.7454	0.3446	0.030*
C4A	0.9375 (5)	0.64851 (17)	0.35544 (10)	0.0256 (8)
H41A	0.9215	0.6001	0.3440	0.031*
H42A	1.0515	0.6655	0.3449	0.031*
C5A	0.9374 (5)	0.64668 (16)	0.40622 (10)	0.0202 (8)
C6A	1.0815 (5)	0.65764 (16)	0.42912 (11)	0.0243 (8)
H61A	1.1832	0.6685	0.4126	0.029*
C7A	1.0964 (4)	0.65414 (17)	0.47918 (10)	0.0233 (8)
H71A	1.1122	0.7023	0.4911	0.028*
H72A	1.2005	0.6264	0.4872	0.028*
C8A	0.9385 (4)	0.62097 (16)	0.50128 (10)	0.0172 (7)
H81A	0.9438	0.5689	0.4968	0.021*
C9A	0.7721 (4)	0.64902 (16)	0.47901 (10)	0.0195 (7)
H91A	0.7785	0.7015	0.4811	0.023*
C10A	0.7637 (4)	0.63144 (15)	0.42821 (10)	0.0190 (7)
C11A	0.6082 (4)	0.62733 (18)	0.50481 (10)	0.0242 (8)

H11A	0.5089	0.6543	0.4928	0.029*
H11C	0.5855	0.5769	0.4990	0.029*
C12A	0.6168 (4)	0.63881 (17)	0.55576 (10)	0.0230 (8)
H12A	0.6176	0.6899	0.5621	0.028*
H12C	0.5122	0.6184	0.5699	0.028*
C13A	0.7778 (4)	0.60512 (15)	0.57639 (10)	0.0183 (7)
C14A	0.9350 (4)	0.63605 (16)	0.55146 (10)	0.0181 (8)
H14A	0.9255	0.6883	0.5545	0.022*
C15A	1.0903 (4)	0.61434 (16)	0.57969 (10)	0.0224 (8)
H15A	1.1881	0.6473	0.5754	0.027*
H15C	1.1289	0.5662	0.5719	0.027*
C16A	1.0214 (4)	0.61751 (17)	0.62834 (11)	0.0229 (8)
H16A	1.0735	0.6578	0.6445	0.027*
H16C	1.0508	0.5737	0.6446	0.027*
C17A	0.8214 (4)	0.62640 (16)	0.62533 (10)	0.0202 (8)
H17A	0.7948	0.6776	0.6287	0.024*
C18A	0.7693 (5)	0.52436 (15)	0.57226 (10)	0.0219 (8)
H18A	0.7630	0.5112	0.5405	0.033*
H18C	0.6664	0.5069	0.5879	0.033*
H18D	0.8732	0.5036	0.5857	0.033*
C19A	0.7198 (5)	0.55331 (16)	0.42009 (10)	0.0277 (9)
H19A	0.7365	0.5420	0.3883	0.042*
H19C	0.5991	0.5445	0.4284	0.042*
H19D	0.7961	0.5237	0.4384	0.042*
C20A	0.7291 (4)	0.58734 (16)	0.66317 (10)	0.0223 (8)
H20A	0.7572	0.5361	0.6608	0.027*
C21A	0.5318 (4)	0.59589 (18)	0.66345 (11)	0.0285 (9)
H21A	0.4833	0.5750	0.6362	0.043*
H21C	0.5025	0.6460	0.6645	0.043*
H21D	0.4836	0.5721	0.6898	0.043*
C22A	0.7925 (4)	0.61417 (17)	0.70820 (10)	0.0234 (8)
C23A	0.8338 (5)	0.58740 (19)	0.78883 (11)	0.0340 (9)
H23A	0.9119	0.5541	0.8037	0.051*
H23C	0.7209	0.5868	0.8038	0.051*
H23D	0.8829	0.6349	0.7905	0.051*
C24A	0.8716 (5)	0.44840 (17)	0.73243 (12)	0.0329 (9)
H24A	0.8205	0.4012	0.7337	0.049*
H24C	0.9545	0.4540	0.7570	0.049*
H24D	0.9314	0.4546	0.7037	0.049*
O1B	0.3248 (3)	0.29218 (11)	0.34061 (7)	0.0241 (6)
O2B	0.1000 (3)	0.43877 (11)	0.30237 (7)	0.0244 (6)
O3B	0.3740 (3)	0.28938 (12)	0.75771 (7)	0.0275 (6)
H3B	0.308 (5)	0.2590 (19)	0.7683 (11)	0.041*
N1B	0.2268 (4)	0.38484 (12)	0.30187 (8)	0.0212 (6)
C1B	0.1604 (4)	0.30886 (16)	0.64465 (9)	0.0203 (8)
H1B	0.0411	0.3224	0.6362	0.024*
H2B	0.1808	0.2606	0.6333	0.024*
C2B	0.1737 (4)	0.30804 (17)	0.69563 (10)	0.0222 (8)

H4B	0.1448	0.3552	0.7076	0.027*
H5B	0.0897	0.2739	0.7080	0.027*
C3B	0.3537 (4)	0.28808 (16)	0.70983 (10)	0.0204 (8)
H31B	0.3798	0.2396	0.6987	0.024*
C4B	0.4852 (4)	0.33881 (16)	0.69055 (10)	0.0216 (8)
H41B	0.6031	0.3220	0.6980	0.026*
H42B	0.4695	0.3855	0.7047	0.026*
C5B	0.4690 (4)	0.34639 (16)	0.63992 (10)	0.0167 (8)
C6B	0.6090 (4)	0.34157 (15)	0.61433 (10)	0.0190 (8)
H61B	0.7165	0.3331	0.6288	0.023*
C7B	0.6103 (4)	0.34850 (16)	0.56420 (10)	0.0181 (7)
H71B	0.6346	0.3018	0.5508	0.022*
H72B	0.7053	0.3806	0.5554	0.022*
C8B	0.4407 (4)	0.37637 (15)	0.54505 (10)	0.0148 (7)
H81B	0.4377	0.4287	0.5487	0.018*
C9B	0.2861 (4)	0.34433 (15)	0.56994 (9)	0.0148 (7)
H91B	0.2973	0.2922	0.5666	0.018*
C10B	0.2883 (4)	0.35925 (15)	0.62135 (9)	0.0164 (7)
C11B	0.1142 (4)	0.36437 (16)	0.54751 (9)	0.0186 (8)
H11B	0.0926	0.4151	0.5525	0.022*
H11E	0.0194	0.3379	0.5622	0.022*
C12B	0.1090 (4)	0.34958 (17)	0.49675 (10)	0.0195 (8)
H12B	0.1125	0.2981	0.4917	0.023*
H12E	−0.0013	0.3675	0.4843	0.023*
C13B	0.2603 (4)	0.38405 (15)	0.47190 (10)	0.0159 (7)
C14B	0.4271 (4)	0.35862 (16)	0.49529 (10)	0.0159 (7)
H14B	0.4238	0.3061	0.4937	0.019*
C15B	0.5736 (4)	0.38110 (16)	0.46425 (10)	0.0185 (8)
H15B	0.6739	0.3489	0.4670	0.022*
H15E	0.6119	0.4296	0.4711	0.022*
C16B	0.4928 (4)	0.37679 (17)	0.41675 (10)	0.0190 (8)
H16B	0.5038	0.4225	0.4011	0.023*
H16E	0.5516	0.3402	0.3987	0.023*
C17B	0.2998 (4)	0.35775 (15)	0.42354 (9)	0.0163 (7)
H17B	0.2924	0.3052	0.4242	0.020*
C18B	0.2422 (5)	0.46448 (14)	0.47231 (10)	0.0205 (8)
H18B	0.2399	0.4813	0.5034	0.031*
H18E	0.1344	0.4779	0.4572	0.031*
H18F	0.3407	0.4856	0.4566	0.031*
C19B	0.2362 (5)	0.43602 (15)	0.63154 (10)	0.0238 (8)
H19B	0.1118	0.4420	0.6264	0.036*
H19E	0.3007	0.4679	0.6118	0.036*
H19F	0.2632	0.4470	0.6629	0.036*
C20B	0.1861 (4)	0.38264 (15)	0.38456 (10)	0.0171 (7)
H20B	0.1963	0.4349	0.3819	0.021*
C21B	−0.0056 (4)	0.36319 (18)	0.38993 (10)	0.0241 (8)
H21B	−0.0159	0.3127	0.3961	0.036*
H21E	−0.0680	0.3745	0.3622	0.036*

H21F	−0.0554	0.3899	0.4149	0.036*
C22B	0.2502 (4)	0.34904 (16)	0.34128 (10)	0.0180 (7)
C23B	0.2346 (5)	0.34870 (17)	0.25866 (10)	0.0297 (9)
H23B	0.3161	0.3093	0.2606	0.045*
H23E	0.2735	0.3817	0.2355	0.045*
H23F	0.1194	0.3309	0.2509	0.045*
C24B	0.1771 (5)	0.50416 (17)	0.28946 (11)	0.0323 (9)
H24B	0.2626	0.5182	0.3120	0.048*
H24E	0.0872	0.5404	0.2872	0.048*
H24F	0.2342	0.4987	0.2603	0.048*

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1A	0.0366 (16)	0.0248 (14)	0.0329 (14)	0.0014 (12)	−0.0019 (12)	−0.0128 (10)
O2A	0.0243 (14)	0.0266 (13)	0.0295 (13)	−0.0025 (12)	0.0005 (11)	−0.0031 (10)
O3A	0.0298 (16)	0.0442 (15)	0.0224 (14)	−0.0058 (13)	0.0051 (12)	0.0016 (11)
N1A	0.0296 (19)	0.0282 (16)	0.0207 (16)	−0.0023 (14)	0.0012 (14)	−0.0059 (13)
C1A	0.018 (2)	0.027 (2)	0.025 (2)	0.0001 (16)	0.0036 (16)	0.0005 (14)
C2A	0.021 (2)	0.028 (2)	0.027 (2)	−0.0012 (17)	0.0015 (16)	0.0008 (15)
C3A	0.030 (2)	0.0219 (18)	0.0228 (19)	−0.0021 (16)	0.0047 (17)	−0.0006 (14)
C4A	0.024 (2)	0.0242 (19)	0.028 (2)	−0.0035 (17)	0.0075 (16)	−0.0017 (15)
C5A	0.020 (2)	0.0133 (17)	0.028 (2)	0.0011 (16)	0.0059 (17)	−0.0014 (14)
C6A	0.016 (2)	0.0227 (19)	0.034 (2)	0.0007 (16)	0.0084 (18)	0.0027 (15)
C7A	0.0145 (19)	0.0252 (18)	0.030 (2)	−0.0010 (16)	0.0016 (16)	0.0002 (15)
C8A	0.0149 (18)	0.0132 (16)	0.0235 (19)	0.0002 (15)	0.0008 (15)	−0.0028 (13)
C9A	0.0172 (19)	0.0161 (16)	0.0253 (18)	0.0005 (15)	0.0024 (16)	−0.0020 (14)
C10A	0.0140 (18)	0.0156 (16)	0.0274 (18)	−0.0007 (15)	0.0015 (16)	−0.0008 (13)
C11A	0.016 (2)	0.033 (2)	0.024 (2)	−0.0011 (17)	−0.0005 (16)	0.0009 (15)
C12A	0.015 (2)	0.0266 (19)	0.027 (2)	0.0000 (16)	0.0030 (16)	−0.0003 (15)
C13A	0.0190 (19)	0.0149 (16)	0.0210 (18)	0.0003 (15)	0.0005 (16)	−0.0040 (13)
C14A	0.0151 (19)	0.0112 (17)	0.0279 (19)	−0.0008 (15)	0.0012 (16)	−0.0009 (14)
C15A	0.019 (2)	0.0187 (18)	0.030 (2)	0.0024 (16)	0.0004 (16)	0.0000 (14)
C16A	0.021 (2)	0.0176 (18)	0.030 (2)	0.0004 (15)	−0.0036 (16)	−0.0021 (14)
C17A	0.022 (2)	0.0126 (16)	0.0259 (19)	0.0026 (15)	−0.0009 (16)	−0.0042 (13)
C18A	0.024 (2)	0.0205 (17)	0.0214 (18)	−0.0060 (16)	0.0004 (16)	−0.0035 (13)
C19A	0.033 (2)	0.0234 (18)	0.0266 (19)	−0.0068 (17)	0.0025 (18)	−0.0024 (14)
C20A	0.023 (2)	0.0185 (17)	0.0254 (19)	0.0032 (16)	0.0014 (17)	−0.0062 (14)
C21A	0.026 (2)	0.031 (2)	0.028 (2)	−0.0026 (17)	0.0015 (17)	−0.0005 (16)
C22A	0.018 (2)	0.026 (2)	0.026 (2)	0.0031 (16)	0.0052 (16)	−0.0074 (15)
C23A	0.028 (2)	0.050 (2)	0.024 (2)	0.0018 (19)	0.0016 (18)	−0.0092 (17)
C24A	0.029 (2)	0.028 (2)	0.042 (2)	0.0008 (18)	0.0009 (18)	0.0000 (16)
O1B	0.0315 (15)	0.0202 (12)	0.0206 (12)	0.0058 (11)	−0.0002 (11)	−0.0011 (9)
O2B	0.0215 (14)	0.0249 (12)	0.0269 (13)	0.0064 (11)	0.0017 (11)	0.0063 (10)
O3B	0.0290 (16)	0.0349 (15)	0.0184 (14)	−0.0094 (12)	−0.0034 (11)	0.0058 (10)
N1B	0.0221 (17)	0.0238 (14)	0.0177 (15)	0.0091 (14)	0.0002 (13)	0.0021 (11)
C1B	0.0157 (19)	0.0251 (18)	0.0199 (18)	0.0001 (16)	−0.0014 (15)	0.0018 (14)
C2B	0.021 (2)	0.0228 (18)	0.0226 (19)	−0.0033 (15)	0.0003 (16)	0.0018 (14)

C3B	0.026 (2)	0.0195 (17)	0.0153 (18)	−0.0002 (16)	−0.0022 (15)	0.0001 (14)
C4B	0.021 (2)	0.0216 (19)	0.0224 (19)	−0.0043 (15)	−0.0039 (16)	−0.0014 (14)
C5B	0.019 (2)	0.0121 (17)	0.0191 (18)	−0.0028 (15)	−0.0009 (15)	0.0001 (13)
C6B	0.016 (2)	0.0168 (17)	0.025 (2)	−0.0021 (15)	−0.0064 (16)	0.0027 (14)
C7B	0.0160 (19)	0.0149 (16)	0.0233 (19)	0.0013 (15)	−0.0013 (15)	−0.0015 (13)
C8B	0.0138 (18)	0.0111 (16)	0.0196 (18)	0.0002 (15)	−0.0015 (14)	−0.0002 (13)
C9B	0.0146 (19)	0.0116 (15)	0.0183 (17)	0.0006 (14)	−0.0006 (14)	−0.0002 (12)
C10B	0.0142 (19)	0.0160 (16)	0.0189 (17)	−0.0010 (14)	−0.0013 (15)	−0.0006 (13)
C11B	0.0130 (19)	0.0255 (19)	0.0173 (18)	−0.0004 (15)	0.0012 (15)	−0.0004 (14)
C12B	0.0125 (19)	0.0246 (18)	0.0213 (19)	−0.0004 (16)	−0.0011 (15)	−0.0006 (14)
C13B	0.0136 (18)	0.0156 (16)	0.0184 (17)	0.0006 (15)	−0.0038 (14)	−0.0019 (13)
C14B	0.0148 (18)	0.0118 (16)	0.0210 (18)	−0.0003 (15)	−0.0011 (15)	−0.0009 (13)
C15B	0.0156 (19)	0.0181 (17)	0.0217 (19)	−0.0012 (16)	0.0017 (15)	0.0015 (13)
C16B	0.0165 (19)	0.0193 (18)	0.0214 (18)	−0.0002 (15)	0.0005 (15)	−0.0013 (14)
C17B	0.0161 (19)	0.0133 (16)	0.0195 (17)	0.0013 (14)	−0.0010 (15)	0.0005 (13)
C18B	0.022 (2)	0.0203 (17)	0.0193 (17)	0.0061 (17)	−0.0020 (16)	−0.0027 (13)
C19B	0.028 (2)	0.0229 (17)	0.0202 (18)	0.0055 (18)	0.0004 (16)	−0.0013 (13)
C20B	0.0175 (19)	0.0151 (16)	0.0188 (18)	−0.0015 (14)	0.0012 (15)	0.0011 (13)
C21B	0.019 (2)	0.033 (2)	0.0197 (18)	0.0022 (16)	−0.0010 (15)	0.0005 (15)
C22B	0.0123 (18)	0.0189 (17)	0.0229 (18)	−0.0049 (17)	−0.0040 (15)	−0.0017 (14)
C23B	0.034 (2)	0.036 (2)	0.0190 (18)	0.0055 (19)	−0.0005 (17)	−0.0012 (15)
C24B	0.038 (2)	0.0243 (19)	0.035 (2)	0.0023 (17)	0.0045 (18)	0.0064 (16)

Geometric parameters (Å, °)

O1A—C22A	1.232 (4)	O1B—C22B	1.225 (4)
O2A—N1A	1.410 (3)	O2B—N1B	1.418 (3)
O2A—C24A	1.433 (4)	O2B—C24B	1.431 (4)
O3A—C3A	1.422 (4)	O3B—C3B	1.436 (4)
O3A—H3A	0.86 (4)	O3B—H3B	0.83 (4)
N1A—C22A	1.356 (4)	N1B—C22B	1.370 (4)
N1A—C23A	1.456 (4)	N1B—C23B	1.461 (4)
C1A—C2A	1.535 (4)	C1B—C2B	1.523 (4)
C1A—C10A	1.540 (4)	C1B—C10B	1.541 (4)
C1A—H1A	0.9900	C1B—H1B	0.9900
C1A—H2A	0.9900	C1B—H2B	0.9900
C2A—C3A	1.503 (5)	C2B—C3B	1.503 (4)
C2A—H4A	0.9900	C2B—H4B	0.9900
C2A—H5A	0.9900	C2B—H5B	0.9900
C3A—C4A	1.523 (5)	C3B—C4B	1.513 (4)
C3A—H31A	1.0000	C3B—H31B	1.0000
C4A—C5A	1.514 (4)	C4B—C5B	1.522 (4)
C4A—H41A	0.9900	C4B—H41B	0.9900
C4A—H42A	0.9900	C4B—H42B	0.9900
C5A—C6A	1.322 (5)	C5B—C6B	1.327 (4)
C5A—C10A	1.521 (5)	C5B—C10B	1.522 (4)
C6A—C7A	1.499 (4)	C6B—C7B	1.501 (4)
C6A—H61A	0.9500	C6B—H61B	0.9500

C7A—C8A	1.523 (5)	C7B—C8B	1.524 (4)
C7A—H71A	0.9900	C7B—H71B	0.9900
C7A—H72A	0.9900	C7B—H72B	0.9900
C8A—C14A	1.524 (4)	C8B—C14B	1.525 (4)
C8A—C9A	1.542 (4)	C8B—C9B	1.532 (4)
C8A—H81A	1.0000	C8B—H81B	1.0000
C9A—C11A	1.538 (4)	C9B—C11B	1.535 (4)
C9A—C10A	1.552 (4)	C9B—C10B	1.559 (4)
C9A—H91A	1.0000	C9B—H91B	1.0000
C10A—C19A	1.542 (4)	C10B—C19B	1.544 (4)
C11A—C12A	1.536 (4)	C11B—C12B	1.540 (4)
C11A—H11A	0.9900	C11B—H11B	0.9900
C11A—H11C	0.9900	C11B—H11E	0.9900
C12A—C13A	1.528 (4)	C12B—C13B	1.531 (4)
C12A—H12A	0.9900	C12B—H12B	0.9900
C12A—H12C	0.9900	C12B—H12E	0.9900
C13A—C14A	1.541 (4)	C13B—C18B	1.535 (4)
C13A—C18A	1.541 (4)	C13B—C14B	1.543 (4)
C13A—C17A	1.551 (4)	C13B—C17B	1.556 (4)
C14A—C15A	1.522 (4)	C14B—C15B	1.523 (4)
C14A—H14A	1.0000	C14B—H14B	1.0000
C15A—C16A	1.546 (4)	C15B—C16B	1.550 (4)
C15A—H15A	0.9900	C15B—H15B	0.9900
C15A—H15C	0.9900	C15B—H15E	0.9900
C16A—C17A	1.557 (5)	C16B—C17B	1.548 (4)
C16A—H16A	0.9900	C16B—H16B	0.9900
C16A—H16C	0.9900	C16B—H16E	0.9900
C17A—C20A	1.527 (4)	C17B—C20B	1.531 (4)
C17A—H17A	1.0000	C17B—H17B	1.0000
C18A—H18A	0.9800	C18B—H18B	0.9800
C18A—H18C	0.9800	C18B—H18E	0.9800
C18A—H18D	0.9800	C18B—H18F	0.9800
C19A—H19A	0.9800	C19B—H19B	0.9800
C19A—H19C	0.9800	C19B—H19E	0.9800
C19A—H19D	0.9800	C19B—H19F	0.9800
C20A—C22A	1.517 (4)	C20B—C22B	1.523 (4)
C20A—C21A	1.533 (5)	C20B—C21B	1.535 (4)
C20A—H20A	1.0000	C20B—H20B	1.0000
C21A—H21A	0.9800	C21B—H21B	0.9800
C21A—H21C	0.9800	C21B—H21E	0.9800
C21A—H21D	0.9800	C21B—H21F	0.9800
C23A—H23A	0.9800	C23B—H23B	0.9800
C23A—H23C	0.9800	C23B—H23E	0.9800
C23A—H23D	0.9800	C23B—H23F	0.9800
C24A—H24A	0.9800	C24B—H24B	0.9800
C24A—H24C	0.9800	C24B—H24E	0.9800
C24A—H24D	0.9800	C24B—H24F	0.9800

N1A—O2A—C24A	109.5 (2)	N1B—O2B—C24B	109.7 (2)
C3A—O3A—H3A	107 (3)	C3B—O3B—H3B	107 (2)
C22A—N1A—O2A	117.5 (2)	C22B—N1B—O2B	116.2 (2)
C22A—N1A—C23A	123.6 (3)	C22B—N1B—C23B	121.2 (2)
O2A—N1A—C23A	113.3 (2)	O2B—N1B—C23B	112.2 (2)
C2A—C1A—C10A	114.9 (3)	C2B—C1B—C10B	114.4 (3)
C2A—C1A—H1A	108.5	C2B—C1B—H1B	108.7
C10A—C1A—H1A	108.5	C10B—C1B—H1B	108.7
C2A—C1A—H2A	108.5	C2B—C1B—H2B	108.7
C10A—C1A—H2A	108.5	C10B—C1B—H2B	108.7
H1A—C1A—H2A	107.5	H1B—C1B—H2B	107.6
C3A—C2A—C1A	110.0 (3)	C3B—C2B—C1B	110.3 (3)
C3A—C2A—H4A	109.7	C3B—C2B—H4B	109.6
C1A—C2A—H4A	109.7	C1B—C2B—H4B	109.6
C3A—C2A—H5A	109.7	C3B—C2B—H5B	109.6
C1A—C2A—H5A	109.7	C1B—C2B—H5B	109.6
H4A—C2A—H5A	108.2	H4B—C2B—H5B	108.1
O3A—C3A—C2A	108.2 (3)	O3B—C3B—C2B	112.1 (3)
O3A—C3A—C4A	111.5 (3)	O3B—C3B—C4B	107.1 (2)
C2A—C3A—C4A	110.5 (3)	C2B—C3B—C4B	110.7 (2)
O3A—C3A—H31A	108.9	O3B—C3B—H31B	109.0
C2A—C3A—H31A	108.9	C2B—C3B—H31B	109.0
C4A—C3A—H31A	108.9	C4B—C3B—H31B	109.0
C5A—C4A—C3A	113.1 (3)	C3B—C4B—C5B	112.5 (3)
C5A—C4A—H41A	109.0	C3B—C4B—H41B	109.1
C3A—C4A—H41A	109.0	C5B—C4B—H41B	109.1
C5A—C4A—H42A	109.0	C3B—C4B—H42B	109.1
C3A—C4A—H42A	109.0	C5B—C4B—H42B	109.1
H41A—C4A—H42A	107.8	H41B—C4B—H42B	107.8
C6A—C5A—C4A	120.8 (3)	C6B—C5B—C4B	119.8 (3)
C6A—C5A—C10A	123.3 (3)	C6B—C5B—C10B	123.4 (3)
C4A—C5A—C10A	115.8 (3)	C4B—C5B—C10B	116.8 (3)
C5A—C6A—C7A	124.9 (3)	C5B—C6B—C7B	124.9 (3)
C5A—C6A—H61A	117.5	C5B—C6B—H61B	117.6
C7A—C6A—H61A	117.5	C7B—C6B—H61B	117.6
C6A—C7A—C8A	112.8 (3)	C6B—C7B—C8B	113.5 (3)
C6A—C7A—H71A	109.0	C6B—C7B—H71B	108.9
C8A—C7A—H71A	109.0	C8B—C7B—H71B	108.9
C6A—C7A—H72A	109.0	C6B—C7B—H72B	108.9
C8A—C7A—H72A	109.0	C8B—C7B—H72B	108.9
H71A—C7A—H72A	107.8	H71B—C7B—H72B	107.7
C7A—C8A—C14A	111.2 (3)	C7B—C8B—C14B	110.3 (3)
C7A—C8A—C9A	109.8 (2)	C7B—C8B—C9B	110.5 (2)
C14A—C8A—C9A	110.1 (3)	C14B—C8B—C9B	109.3 (2)
C7A—C8A—H81A	108.6	C7B—C8B—H81B	108.9
C14A—C8A—H81A	108.6	C14B—C8B—H81B	108.9
C9A—C8A—H81A	108.6	C9B—C8B—H81B	108.9
C11A—C9A—C8A	112.2 (2)	C8B—C9B—C11B	111.4 (2)

C11A—C9A—C10A	113.3 (3)	C8B—C9B—C10B	113.3 (2)
C8A—C9A—C10A	112.4 (2)	C11B—C9B—C10B	113.1 (2)
C11A—C9A—H91A	106.1	C8B—C9B—H91B	106.1
C8A—C9A—H91A	106.1	C11B—C9B—H91B	106.1
C10A—C9A—H91A	106.1	C10B—C9B—H91B	106.1
C5A—C10A—C1A	108.6 (2)	C5B—C10B—C1B	108.9 (2)
C5A—C10A—C19A	108.0 (3)	C5B—C10B—C19B	108.6 (3)
C1A—C10A—C19A	109.3 (3)	C1B—C10B—C19B	109.4 (3)
C5A—C10A—C9A	110.1 (3)	C5B—C10B—C9B	109.8 (3)
C1A—C10A—C9A	109.1 (2)	C1B—C10B—C9B	108.9 (2)
C19A—C10A—C9A	111.7 (2)	C19B—C10B—C9B	111.3 (2)
C12A—C11A—C9A	114.9 (3)	C9B—C11B—C12B	113.9 (3)
C12A—C11A—H11A	108.5	C9B—C11B—H11B	108.8
C9A—C11A—H11A	108.5	C12B—C11B—H11B	108.8
C12A—C11A—H11C	108.5	C9B—C11B—H11E	108.8
C9A—C11A—H11C	108.5	C12B—C11B—H11E	108.8
H11A—C11A—H11C	107.5	H11B—C11B—H11E	107.7
C13A—C12A—C11A	112.0 (3)	C13B—C12B—C11B	112.2 (3)
C13A—C12A—H12A	109.2	C13B—C12B—H12B	109.2
C11A—C12A—H12A	109.2	C11B—C12B—H12B	109.2
C13A—C12A—H12C	109.2	C13B—C12B—H12E	109.2
C11A—C12A—H12C	109.2	C11B—C12B—H12E	109.2
H12A—C12A—H12C	107.9	H12B—C12B—H12E	107.9
C12A—C13A—C14A	106.7 (2)	C12B—C13B—C18B	110.6 (3)
C12A—C13A—C18A	110.5 (3)	C12B—C13B—C14B	106.6 (2)
C14A—C13A—C18A	112.0 (3)	C18B—C13B—C14B	112.6 (3)
C12A—C13A—C17A	116.5 (3)	C12B—C13B—C17B	117.4 (2)
C14A—C13A—C17A	100.5 (3)	C18B—C13B—C17B	110.2 (2)
C18A—C13A—C17A	110.1 (2)	C14B—C13B—C17B	98.9 (2)
C15A—C14A—C8A	118.6 (3)	C15B—C14B—C8B	118.5 (3)
C15A—C14A—C13A	104.6 (2)	C15B—C14B—C13B	105.0 (2)
C8A—C14A—C13A	114.6 (3)	C8B—C14B—C13B	115.3 (3)
C15A—C14A—H14A	106.1	C15B—C14B—H14B	105.6
C8A—C14A—H14A	106.1	C8B—C14B—H14B	105.6
C13A—C14A—H14A	106.1	C13B—C14B—H14B	105.6
C14A—C15A—C16A	103.7 (3)	C14B—C15B—C16B	103.9 (3)
C14A—C15A—H15A	111.0	C14B—C15B—H15B	111.0
C16A—C15A—H15A	111.0	C16B—C15B—H15B	111.0
C14A—C15A—H15C	111.0	C14B—C15B—H15E	111.0
C16A—C15A—H15C	111.0	C16B—C15B—H15E	111.0
H15A—C15A—H15C	109.0	H15B—C15B—H15E	109.0
C15A—C16A—C17A	107.0 (3)	C17B—C16B—C15B	106.3 (2)
C15A—C16A—H16A	110.3	C17B—C16B—H16B	110.5
C17A—C16A—H16A	110.3	C15B—C16B—H16B	110.5
C15A—C16A—H16C	110.3	C17B—C16B—H16E	110.5
C17A—C16A—H16C	110.3	C15B—C16B—H16E	110.5
H16A—C16A—H16C	108.6	H16B—C16B—H16E	108.7
C20A—C17A—C13A	117.8 (3)	C20B—C17B—C16B	112.4 (2)

C20A—C17A—C16A	111.6 (3)	C20B—C17B—C13B	119.5 (2)
C13A—C17A—C16A	104.0 (3)	C16B—C17B—C13B	103.6 (2)
C20A—C17A—H17A	107.7	C20B—C17B—H17B	106.9
C13A—C17A—H17A	107.7	C16B—C17B—H17B	106.9
C16A—C17A—H17A	107.7	C13B—C17B—H17B	106.9
C13A—C18A—H18A	109.5	C13B—C18B—H18B	109.5
C13A—C18A—H18C	109.5	C13B—C18B—H18E	109.5
H18A—C18A—H18C	109.5	H18B—C18B—H18E	109.5
C13A—C18A—H18D	109.5	C13B—C18B—H18F	109.5
H18A—C18A—H18D	109.5	H18B—C18B—H18F	109.5
H18C—C18A—H18D	109.5	H18E—C18B—H18F	109.5
C10A—C19A—H19A	109.5	C10B—C19B—H19B	109.5
C10A—C19A—H19C	109.5	C10B—C19B—H19E	109.5
H19A—C19A—H19C	109.5	H19B—C19B—H19E	109.5
C10A—C19A—H19D	109.5	C10B—C19B—H19F	109.5
H19A—C19A—H19D	109.5	H19B—C19B—H19F	109.5
H19C—C19A—H19D	109.5	H19E—C19B—H19F	109.5
C22A—C20A—C17A	109.9 (3)	C22B—C20B—C17B	109.1 (3)
C22A—C20A—C21A	106.3 (3)	C22B—C20B—C21B	107.5 (3)
C17A—C20A—C21A	114.6 (3)	C17B—C20B—C21B	113.6 (2)
C22A—C20A—H20A	108.6	C22B—C20B—H20B	108.8
C17A—C20A—H20A	108.6	C17B—C20B—H20B	108.8
C21A—C20A—H20A	108.6	C21B—C20B—H20B	108.8
C20A—C21A—H21A	109.5	C20B—C21B—H21B	109.5
C20A—C21A—H21C	109.5	C20B—C21B—H21E	109.5
H21A—C21A—H21C	109.5	H21B—C21B—H21E	109.5
C20A—C21A—H21D	109.5	C20B—C21B—H21F	109.5
H21A—C21A—H21D	109.5	H21B—C21B—H21F	109.5
H21C—C21A—H21D	109.5	H21E—C21B—H21F	109.5
O1A—C22A—N1A	119.4 (3)	O1B—C22B—N1B	119.1 (3)
O1A—C22A—C20A	122.3 (3)	O1B—C22B—C20B	122.5 (3)
N1A—C22A—C20A	118.3 (3)	N1B—C22B—C20B	118.4 (3)
N1A—C23A—H23A	109.5	N1B—C23B—H23B	109.5
N1A—C23A—H23C	109.5	N1B—C23B—H23E	109.5
H23A—C23A—H23C	109.5	H23B—C23B—H23E	109.5
N1A—C23A—H23D	109.5	N1B—C23B—H23F	109.5
H23A—C23A—H23D	109.5	H23B—C23B—H23F	109.5
H23C—C23A—H23D	109.5	H23E—C23B—H23F	109.5
O2A—C24A—H24A	109.5	O2B—C24B—H24B	109.5
O2A—C24A—H24C	109.5	O2B—C24B—H24E	109.5
H24A—C24A—H24C	109.5	H24B—C24B—H24E	109.5
O2A—C24A—H24D	109.5	O2B—C24B—H24F	109.5
H24A—C24A—H24D	109.5	H24B—C24B—H24F	109.5
H24C—C24A—H24D	109.5	H24E—C24B—H24F	109.5
C24A—O2A—N1A—C22A	112.3 (3)	C24B—O2B—N1B—C22B	122.9 (3)
C24A—O2A—N1A—C23A	−93.4 (3)	C24B—O2B—N1B—C23B	−91.7 (3)
C10A—C1A—C2A—C3A	−57.2 (4)	C10B—C1B—C2B—C3B	−58.0 (3)

C1A—C2A—C3A—O3A	178.6 (3)	C1B—C2B—C3B—O3B	177.7 (2)
C1A—C2A—C3A—C4A	56.3 (4)	C1B—C2B—C3B—C4B	58.2 (3)
O3A—C3A—C4A—C5A	−174.3 (3)	O3B—C3B—C4B—C5B	−176.0 (3)
C2A—C3A—C4A—C5A	−53.9 (4)	C2B—C3B—C4B—C5B	−53.5 (4)
C3A—C4A—C5A—C6A	−129.7 (3)	C3B—C4B—C5B—C6B	−131.5 (3)
C3A—C4A—C5A—C10A	50.5 (4)	C3B—C4B—C5B—C10B	48.4 (4)
C4A—C5A—C6A—C7A	−177.5 (3)	C4B—C5B—C6B—C7B	−179.6 (3)
C10A—C5A—C6A—C7A	2.3 (5)	C10B—C5B—C6B—C7B	0.5 (5)
C5A—C6A—C7A—C8A	12.6 (5)	C5B—C6B—C7B—C8B	11.6 (4)
C6A—C7A—C8A—C14A	−164.5 (3)	C6B—C7B—C8B—C14B	−160.5 (2)
C6A—C7A—C8A—C9A	−42.5 (3)	C6B—C7B—C8B—C9B	−39.6 (3)
C7A—C8A—C9A—C11A	−170.4 (3)	C7B—C8B—C9B—C11B	−172.9 (3)
C14A—C8A—C9A—C11A	−47.7 (3)	C14B—C8B—C9B—C11B	−51.4 (3)
C7A—C8A—C9A—C10A	60.5 (3)	C7B—C8B—C9B—C10B	58.2 (3)
C14A—C8A—C9A—C10A	−176.8 (2)	C14B—C8B—C9B—C10B	179.7 (2)
C6A—C5A—C10A—C1A	133.4 (3)	C6B—C5B—C10B—C1B	135.2 (3)
C4A—C5A—C10A—C1A	−46.8 (4)	C4B—C5B—C10B—C1B	−44.7 (3)
C6A—C5A—C10A—C19A	−108.2 (3)	C6B—C5B—C10B—C19B	−105.8 (3)
C4A—C5A—C10A—C19A	71.6 (3)	C4B—C5B—C10B—C19B	74.3 (3)
C6A—C5A—C10A—C9A	14.0 (4)	C6B—C5B—C10B—C9B	16.1 (4)
C4A—C5A—C10A—C9A	−166.2 (3)	C4B—C5B—C10B—C9B	−163.8 (2)
C2A—C1A—C10A—C5A	50.5 (4)	C2B—C1B—C10B—C5B	49.4 (3)
C2A—C1A—C10A—C19A	−67.1 (4)	C2B—C1B—C10B—C19B	−69.1 (3)
C2A—C1A—C10A—C9A	170.5 (3)	C2B—C1B—C10B—C9B	169.1 (3)
C11A—C9A—C10A—C5A	−173.5 (2)	C8B—C9B—C10B—C5B	−45.2 (3)
C8A—C9A—C10A—C5A	−45.0 (3)	C11B—C9B—C10B—C5B	−173.2 (2)
C11A—C9A—C10A—C1A	67.4 (3)	C8B—C9B—C10B—C1B	−164.3 (2)
C8A—C9A—C10A—C1A	−164.1 (3)	C11B—C9B—C10B—C1B	67.7 (3)
C11A—C9A—C10A—C19A	−53.5 (4)	C8B—C9B—C10B—C19B	75.1 (3)
C8A—C9A—C10A—C19A	75.0 (3)	C11B—C9B—C10B—C19B	−52.9 (3)
C8A—C9A—C11A—C12A	47.0 (4)	C8B—C9B—C11B—C12B	51.4 (3)
C10A—C9A—C11A—C12A	175.6 (3)	C10B—C9B—C11B—C12B	−179.6 (2)
C9A—C11A—C12A—C13A	−52.4 (4)	C9B—C11B—C12B—C13B	−54.1 (3)
C11A—C12A—C13A—C14A	56.0 (3)	C11B—C12B—C13B—C18B	−68.3 (3)
C11A—C12A—C13A—C18A	−66.0 (3)	C11B—C12B—C13B—C14B	54.4 (3)
C11A—C12A—C13A—C17A	167.3 (3)	C11B—C12B—C13B—C17B	164.0 (3)
C7A—C8A—C14A—C15A	−56.5 (4)	C7B—C8B—C14B—C15B	−54.4 (4)
C9A—C8A—C14A—C15A	−178.3 (3)	C9B—C8B—C14B—C15B	−176.1 (3)
C7A—C8A—C14A—C13A	179.2 (2)	C7B—C8B—C14B—C13B	179.9 (2)
C9A—C8A—C14A—C13A	57.4 (3)	C9B—C8B—C14B—C13B	58.3 (3)
C12A—C13A—C14A—C15A	167.5 (2)	C12B—C13B—C14B—C15B	168.6 (2)
C18A—C13A—C14A—C15A	−71.4 (3)	C18B—C13B—C14B—C15B	−69.9 (3)
C17A—C13A—C14A—C15A	45.5 (3)	C17B—C13B—C14B—C15B	46.4 (3)
C12A—C13A—C14A—C8A	−61.0 (3)	C12B—C13B—C14B—C8B	−59.0 (3)
C18A—C13A—C14A—C8A	60.1 (3)	C18B—C13B—C14B—C8B	62.4 (3)
C17A—C13A—C14A—C8A	177.0 (2)	C17B—C13B—C14B—C8B	178.8 (2)
C8A—C14A—C15A—C16A	−164.3 (3)	C8B—C14B—C15B—C16B	−162.5 (3)
C13A—C14A—C15A—C16A	−35.2 (3)	C13B—C14B—C15B—C16B	−32.0 (3)

C14A—C15A—C16A—C17A	11.0 (3)	C14B—C15B—C16B—C17B	4.4 (3)
C12A—C13A—C17A—C20A	83.7 (3)	C15B—C16B—C17B—C20B	154.5 (2)
C14A—C13A—C17A—C20A	-161.5 (3)	C15B—C16B—C17B—C13B	24.2 (3)
C18A—C13A—C17A—C20A	-43.2 (4)	C12B—C13B—C17B—C20B	77.7 (3)
C12A—C13A—C17A—C16A	-152.3 (3)	C18B—C13B—C17B—C20B	-50.2 (4)
C14A—C13A—C17A—C16A	-37.4 (3)	C14B—C13B—C17B—C20B	-168.3 (3)
C18A—C13A—C17A—C16A	80.9 (3)	C12B—C13B—C17B—C16B	-156.4 (3)
C15A—C16A—C17A—C20A	144.9 (2)	C18B—C13B—C17B—C16B	75.7 (3)
C15A—C16A—C17A—C13A	16.9 (3)	C14B—C13B—C17B—C16B	-42.5 (3)
C13A—C17A—C20A—C22A	178.5 (3)	C16B—C17B—C20B—C22B	58.9 (3)
C16A—C17A—C20A—C22A	58.3 (3)	C13B—C17B—C20B—C22B	-179.5 (3)
C13A—C17A—C20A—C21A	-62.0 (4)	C16B—C17B—C20B—C21B	178.9 (3)
C16A—C17A—C20A—C21A	177.9 (3)	C13B—C17B—C20B—C21B	-59.5 (4)
O2A—N1A—C22A—O1A	166.4 (3)	O2B—N1B—C22B—O1B	162.6 (3)
C23A—N1A—C22A—O1A	14.9 (5)	C23B—N1B—C22B—O1B	20.5 (5)
O2A—N1A—C22A—C20A	-14.0 (4)	O2B—N1B—C22B—C20B	-18.9 (4)
C23A—N1A—C22A—C20A	-165.5 (3)	C23B—N1B—C22B—C20B	-161.0 (3)
C17A—C20A—C22A—O1A	37.4 (4)	C17B—C20B—C22B—O1B	27.8 (4)
C21A—C20A—C22A—O1A	-87.2 (4)	C21B—C20B—C22B—O1B	-95.8 (4)
C17A—C20A—C22A—N1A	-142.2 (3)	C17B—C20B—C22B—N1B	-150.7 (3)
C21A—C20A—C22A—N1A	93.2 (3)	C21B—C20B—C22B—N1B	85.8 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O3 <i>A</i> —H3 <i>A</i> $\cdots$ O3 <i>B</i> ⁱ	0.86 (4)	1.93 (4)	2.782 (4)	180 (5)
O3 <i>B</i> —H3 <i>B</i> $\cdots$ O1 <i>A</i> ⁱⁱ	0.83 (4)	1.95 (4)	2.768 (3)	169 (4)

Symmetry codes: (i) $-x+3/2, -y+1, z-1/2$; (ii) $-x+1, y-1/2, -z+3/2$.