Faults affecting the Triassic Barents Shelf: Syn-kinematic deposition, deformation mechanisms and driving forces

Mark Joseph Mulrooney

Faculty of Mathematics and Natural Sciences
Department of Geosciences
University of Oslo
Norway

A thesis submitted for the degree of
Philosophiae Doctor (PhD)
November 2017

© Mark Joseph Mulrooney, 2018

*Series of dissertations submitted to the
Faculty of Mathematics and Natural Sciences, University of Oslo
No. 1958*

ISSN 1501-7710

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

PREFACE

This thesis has been submitted to the Department of Geosciences at the University of Oslo (UiO) in accordance with the requirements for the degree of Philosophiae Doctor (PhD). The work presented herein was conducted between August 2013 and July 2017 during which time I was employed as a PhD research fellow at the University Centre in Svalbard (UNIS) and enrolled in a PhD program at UiO.

The fieldwork aspect of the study was supported by the Trias North project which is funded under grant 234152 from the Research Council of Norway (RCN) and an Arctic Field Grant from the Svalbard Science Forum (SSF). The principal supervisor of this work is Prof. Alvar Braathen (UiO/UNIS). Co-supervision was provided by Prof. Snorre Olaussen (UNIS) and Prof. Jan Inge Faleide (UiO). The work was also conducted in collaboration with Johan Leutscher, Geophysical Operations Manager at ENI Norge AS. The work primarily took place at UNIS (3.5 years) and at UiO (0.5 years) as well as shorter stays at the University in Parma, Italy and at ENI Norge AS, Stavanger. Svalbard based fieldwork was conducted in central and northern Spitsbergen, Nordaustlandet, Edgeøya and Wilhelmøya. Time dedicated to the PhD consisted of 75% research and 25% teaching and outreach.

This thesis serves to synthesise the results of this four year study and comprises three first author articles (one published, one in press, and one in review at time of thesis submission) and a single second author paper (in press). Further, four articles which I co-authored (one published, two in press, and one in review at time of thesis submission) are contained in the appendix of this body of work.

The thesis is structured in two parts, the first part provides a scientific background to the concepts dealt with in the study, and outlines the motivations, methods, objectives behind the study, as well as a synthesising the main results. The second part of the thesis consists of the collection of articles that form the main body of this work. Finally, the appendices provide supplementary material including articles, conference proceedings and details of other works undertaken during the PhD program.

*Mark Mulrooney,
Longyearbyen, July 31st 2017*

ABSTRACT

This thesis addresses the Triassic development of the Norwegian Barents shelf. As a frontier petroleum province with only the area south of 74° 30' N open for petroleum activities before 2017; to date, only two fields have come on stream, the Snøhvit (gas) and Goliat (oil) fields. Both are located in the Hammerfest Basin in the southwestern Barents Sea. The exploration associated with these developments has led to a relatively good understanding of the tectonostratigraphy of the southwestern Norwegian Barents shelf, the north however, remains relatively enigmatic.

The Triassic successions on the Barents shelf represent large deltaic systems that prograded out from the Uralide orogeny and the Baltic shield towards the west and northwest. These successions are partly sandstone prone and envisaged to host significant undiscovered resources (NPD, 2017). Moreover, the distal parts of these successions are being appraised for potential CO₂ storage onshore Svalbard, the emergent north-western limit of the Barents shelf. Until recently, the Triassic Barents shelf has been described as a period of relative tectonic quiescence with ongoing regional subsidence in a shallow epicontinental basin. Newer studies, however, suggest this to be an over simplification, a position which is strengthened by this contribution.

This study attempts to bridge some of the knowledge gaps that exists in respect to the Triassic evolution of the Barents shelf, in particular this body of work investigates evidence for Late Triassic extensional tectonism affecting the Barents shelf, the driving mechanisms and styles of this faulting, and how the faulting affects the distribution and petrophysical properties of sediments. Furthermore, we investigate later (Cenozoic) phases of brittle deformation affecting the Triassic succession onshore Svalbard and the impact of associated structural discontinuities on potential CO₂ storage.

This body of work consists of four scientific articles, which were conducted using an expansive range of datasets and methodologies. This includes two articles, which investigate the structural complexity of the Goliat field using a multi-azimuth (MAZ) seismic survey, core, and wireline data. The combined study addresses the spatio-temporal evolution of the multiple trending fault populations in the area and the evolution of the large anticlinal structure that forms the Goliat trapping structure. Furthermore, palaeogeographic reconstructions for the Fruholmen and Tubåen formations that comprise the Realgrunnen Subgroup are presented. Evidence from core, amplitude vs angle seismic attributes, and fault analysis are synthesised and together advocate a phase of Norian to Rhaetian extension (normal faulting) took place in the Hammerfest Basin and on the basin-platform delineating Troms-Finnmark Fault complex (TFFC).

The third article comprises detailed structural descriptions of a syn-kinematic growth fault system that affects the Late Triassic successions on Edgeøya, Eastern Svalbard. The study focuses on several scales combining outcrop based observations with micro-structural analyses. In addition, virtual outcrop models were used to establish gross geometries of large structures and sandstone bodies, and to provide measurements from otherwise inaccessible localities. The importance of early cementation and differential compaction is highlighted in the formation of listric growth faults and the active deformation mechanisms (i.e., hydro-plastic to brittle), while fault orientation and dip direction are related to slope direction and reactivation of basement structures.

The fourth article consists of a structural analysis of the Late Triassic to Early Jurassic succession onshore Svalbard with the purpose of assessing the target formations as a potential CO₂ storage unit for the arctic settlement, Longyearbyen. The study utilises core, wireline, and outcrop data from extensive field campaigns. The results of well injection tests are presented and indicate that barriers to fluid flow are present in the reservoir. The study details the heterogeneities affecting the reservoir and credits sub-seismic extensional faults that developed oblique to the transport direction of the Palaeogene Western Spitsbergen Fold-and-Thrust-Belt (WSFTB) to the observed lack of reservoir communication.

Together, these collections of articles indicate that fault activity was more prominent during deposition of the Triassic successions of the Barents shelf than previously reported. Furthermore, far-field stress was capable of instigating extensional failure, jointing and fracturing of rocks throughout the region during shelf-wide tectonic events.

ACKNOWLEDGEMENTS

First and foremost, I would like to acknowledge my principal supervisor, Alvar Braathen (UiO) for creating this fantastic PhD opportunity. Alvar has been crucial to the smooth progression of this study, I have learned a great deal from his organised, matter-of fact-approach to research, and am very thankful for the experience I have gained from accompanying him on numerous field campaigns both on Svalbard and in Utah. Next, I want to thank my co-supervisor, Snorre Olaussen (UNIS) for his good humour, Sci-Fi related chats and his library-like mind on all things Barents shelf related. I also extend gratitude to my co-supervisor Jan Inge Faleide (UiO) for his guidance during the initial stages of this research.

I would like to thank my UNIS Arctic Geology colleagues past and present for providing a friendly working environment for four years, Aleksandra for showing me the ropes in my first few months in Svalbard and for providing an outlet for my need to talk about faults, Kim for his enthusiasm and work ethic which helped bulk-up this thesis, Bjarte for the scientific collaboration and company in the field, Hanna Rósa for the hallway chats and ensuring I was sufficiently caffeinated at all times, and Tom and Martin for adding an element of comedy to my work days. Very special thanks to Oscar and Anne for being my partners in crime during the last number of years. I would also like to extend my gratitude to the UNIS logistics for facilitating four years of fieldwork and teaching excursions.

For my travels outside of Svalbard, I want to thank Åsgeir Seland for facilitating my trips to UiO, Kei Ogata and his colleagues at the University of Parma for my time spent at the Elisabetta Costa Analogue Modelling Laboratory, Johan Leutscher and the Goliat team for the ENI Norge research collaboration and the time I spent working with them in-house in Stavanger and finally, Ivar Midtkandal for leading the fantastic GEO4011 Utah field course.

The highlights of this experience have been the three summer field seasons to Edgeøya and other eastern parts of the Svalbard archipelago. I want to extend gratitude to all my colleagues and co-authors that I spent time in the field with as part of the Trias North project.

Finally, I would like to thank my parents, Pat and Angela, who have supported me throughout my academic endeavours.

CONTENTS

PREFACE	I
ABSTRACT	III
ACKNOWLEDGEMENTS	V
1. INTRODUCTION	1
1.1 MOTIVATION	1
1.2 OUTLINE.....	2
1.3 AUTHORSHIP AND CONTRIBUTION	4
1.4 AIMS AND OBJECTIVES	6
2. SCIENTIFIC BACKGROUND: FAULTS AND FAULT GROWTH	9
2.1 INTRODUCTION	9
2.2 DRIVING MECHANISMS	10
2.3 EARTH QUAKE SLIP AND THE IDEAL FAULT MODEL	10
2.4 SHORTCOMINGS OF THE CONVENTIONAL GROWTH MODEL.....	12
2.5 STUDYING FAULT KINEMATICS.....	12
2.6 DISPLACEMENT TERMINOLOGY	13
2.7 ISOLATED AND COHERENT FAULT MODELS	14
2.8 LINKAGE AND SEGMENTATION	14
2.9 INFLUENCE OF BASEMENT STRUCTURES	16
2.10 INFLUENCE OF MECHANICAL STRATIGRAPHY.....	16
2.11 STAGES OF FAULT GROWTH.....	17
2.12 IDENTIFYING LINKAGE HISTORY	19
2.13 DEFORMATION OF THE SURROUNDING WALL ROCK	19
2.14 FAULT GEOMETRIES	20
2.15 INVERSION OF NORMAL FAULTS.....	22
2.16 FAULT CONTROL ON SEDIMENTATION	23
2.17 IMPLICATIONS ON RESERVOIR PROPERTIES.....	24
3. GEOLOGICAL CONTEXT – THE BARENTS SHELF	27
3.1 STRUCTURAL ELEMENTS	27
3. PALAEOZOIC TO CENOZOIC SHELF EVOLUTION	28
4 DATA AND METHODS	34
4. OFFSHORE INVESTIGATIONS	34
4.1.1 Seismic data.....	34
4.1.2 Well logs	34
4.2 ONSHORE INVESTIGATIONS	35
4.2.1 Discontinuity mapping.....	35
4.2.2 Borehole and outcrop plug data	36

4.2.3 Virtual outcrops	36
4.2.4 Additional Methods.....	37
5. ARTICLE SUMMARIES	38
5.1 ARTICLE I.....	38
5.2 ARTICLE II.....	39
5.3 ARTICLE III.....	41
5.4 ARTICLE IV.....	42
6. DISCUSSION	44
6.1 RESOLUTION: BRIDGING THE GAP.....	45
6.2 FAULT GROWTH	46
6.3 INFLUENCE OF FAULTS IN RESERVOIRS	48
6.4 LATE TRIASSIC TECTONISM IN THE BARENTS SEA	50
7. CONCLUSIONS AND FUTURE WORK.....	53
7.1 MAIN FINDINGS OF THE THESIS	53
7.2 FURTHER STUDIES	54
8. REFERENCES	56
9. ARTICLES	79
9.1 ARTICLE I: A 3D structural analysis of the Goliat field, Barents Sea, Norway	79
9.2 ARTICLE II: Impacts of small-scale faults on continental to coastal plain deposition: Evidence from the Realgrunnen Subgroup in the Goliat field, Southwest Barents Sea, Norway	103
9.3 ARTICLE III: Architecture, deformation style and petrophysical properties of a Late Triassic growth fault system in southern Edgeøya (East Svalbard).....	161
9.4 ARTICLE IV: Fluid flow properties of a potential unconventional CO ₂ storage unit in central Spitsbergen: the Upper Triassic to Middle Jurassic Wilhelmøya Subgroup.....	203
10. APPENDICES.....	261
10.1 APPENDIX 1: Supplementary Article – Braathen et al. (2017)	261
10.2 APPENDIX 2: Supplementary Article – Rismyhr et al. (In press).....	283
10.3 APPENDIX 3: Supplementary Article – Maher et al. (In review).....	337
10.4 APPENDIX 4: Conference Proceedings	387
10.5 APPENDIX 5: Conference Proceedings	395
10.6 APPENDIX 6: Conference Proceedings	399
10.7 APPENDIX 7: Conference Proceedings	403
10.8 APPENDIX 8: Conference Proceedings	407
10.9 APPENDIX 9: Conference Proceedings	415
10.10 APPENDIX 10: Conference Proceedings	415
10.11 APPENDIX 11: Internal ENI Norge Report (Unpublished).....	419
10.12 APPENDIX 12: Photogrammetric model.....	423
10.13 APPENDIX 13: Teaching and Outreach	424

1. INTRODUCTION

1.1 MOTIVATION

The Norwegian part of the Barents shelf is a relatively under-explored, expansive continental shelf. It is estimated that approximately half of the undiscovered resources on the Norwegian continental shelf are located within. Although exploration activities here have been ongoing for more than 30 years, the area is still considered a frontier petroleum province with only the area south of 74° 30' N open for petroleum activities prior to 2017. The only fields in production in the Barents shelf are Snøhvit (gas) and Goliat (oil), which came on stream in 2007 and 2016, respectively, both of which are located in the Hammerfest Basin (NPD, 2017). This exploration has increased knowledge regarding the southwestern Barents Shelf whereas the northern parts of the shelf remain relatively enigmatic.

This thesis forms part of the Trias North project “Reconstructing the Triassic northern Barents shelf”, a multi-disciplined (geoscience) knowledge-building project aimed at gaining insight into the sedimentary and structural history of the Barents shelf. The Triassic deposits of the Barents shelf are of particular interest given that during this time a larger epicontinental depositional regime persisted and was subject to shelf-wide expansive deltaic sequences which prograded across the entire Barents shelf (Glørstad-Clark et al., 2010, 2011; Anell et al., 2014). These were sourced from the uplifted Uralides and the Baltic Shield (Riis et al., 2008; Mørk, 2009). Svalbard also experienced some sedimentary input from the west and the north during the Early to Middle Triassic. The Triassic deposits investigated in this thesis host the reservoir units in the Goliat field (Mulrooney et al., 2017) and comprise potential CO₂ storage units (along with the Early Jurassic) onshore central Spitsbergen, Svalbard (Braathen et al., 2012).

Given the underexplored nature of this frontier petroleum province, the Goliat field and the Svalbard outcrops (central Spitsbergen and Edgeøya) provide two study areas at opposing ends of the shelf where the tectonostratigraphy is well documented. Given the epicontinental nature of the Triassic Barents shelf and the scale of the deltaic systems active during this time, Late Triassic outcrops onshore Svalbard provide high resolution analogues for the offshore Goliat field. Further, virtual outcrop model studies onshore Svalbard provide an intermediate resolution that can link large scale features identified in seismic (i.e., clinoforms and faults) to that observed in outcrop.

Key to this study is the availability of the EN0901 MAZ survey that covers the Goliat area. This excellent high-resolution seismic survey gives detailed insight into the structurally complex Goliat field which is compartmentalised by several fault populations and located in a zone of complexity within a jog on the TFFC. The survey can be used to determine the evolution of the Goliat trapping

structure and investigate the tectonostratigraphy of the Triassic successions. Knowledge from the offshore seismic imagery of this compartmentalised field can in turn be used to inform subsurface structures in the CO₂ target reservoir on Svalbard where seismic data is poor and where baffles to fluid flow are evident from fluid injection tests (Larsen, 2010, 2012). Further, these subsurface studies of fault-affected reservoirs are supported by conducting detailed mapping of fault architecture in equivalent outcropping units in order to determine the effects of these faults on the fluid flow properties of the host rock.

The research presented herein concerning Triassic tectonism in the Barents shelf is motivated by provisional investigations by Anell et al. (2013) who postulated a far field driving force for listric growth faults spectacularly exposed on southern Edgeøya, Svalbard, which contradicted the interpretation of the Triassic Barents shelf as a period of general tectonic quiescence (Riis et al., 2008; Worsley, 2008; Glørstad-Clark et al., 2011; Høy & Lundschieen, 2011). Further, these listric faults, previously described as gravitational collapse features (Edwards, 1976), but later reinterpreted as tectonically controlled by Anell et al. (2013) merit further study in order to better understand how the style of listric faults can reflect factors such as the causal driving mechanisms, the slope attitude, depositional environment, diagenetic history etc.

Additionally, channel stacking patterns and potential footwall derived alluvial fans described in the Late Triassic of the Goliat field by Rismyhr (2012), indicate tectonic activity, and merited further study. Since beginning this research, additional zones of local fault activity in the Late Triassic have been described, including the Fingerdjupet Sub-basin (Serck et al., 2017), the southern Loppa High Fault System (Indrevær et al., 2016) and the Hoop Graben (Mahajan et al., 2014). The recognition of this phase of activity affecting Triassic deltaic sediments in the Barents shelf is important in future exploration as fault related subsidence will create accommodation space and influence the distribution of potential reservoir facies.

Finally, the outcrop and borehole related research on the Late Triassic successions in central Spitsbergen is motivated by a proposed carbon capture and storage project, which envisages sequestering CO₂ emitted by a local coal burning power-plant and injecting it within a fracture-dominated subsurface reservoir. Faults and additional heterogeneities observed in outcrop and core are investigated to establish their impacts on the fluid flow properties of the host rock.

1.2 OUTLINE

This thesis is divided into ten chapters; chapter 1 provides an overview of the thesis content including the motivation for undertaking the work and the main aims and objectives that the study aspires towards. The authorship and contribution of authors to each study is outlined. The scientific background to the research topics undertaken herein is detailed in Chapter 2, with particular focus

on the spatio-temporal evolution of normal faults, their influence on sedimentation and their influence on petrophysical properties of reservoirs. Chapter 3 establishes the geographical and geological context of the Barents shelf localities subject to the respective articles. The various data sets and methods used are summarised in Chapter 4. Chapter 5 summarises the main findings from the four articles that form the thesis. A discussion on the contribution of this work to our broader understanding of several key topics is presented in chapter 6, this includes i) how various investigative techniques of contrasting scales can provide a more complete understanding of fault processes, ii) how observations of fault geometries from a MAZ survey herein reflect fault growth models, iii) how the various faults observed in the studies affect reservoir geometries and internal properties, and finally, iv) the significance of late Triassic tectonism on the Barents shelf. Chapter 7 summarises the main findings from the thesis and discusses some future work that can build on the information herein. A list of references for chapters 1 to 7 can be found in chapter 8. Chapter 9 comprises four scientific articles that form the main body of this study; the list of the articles is as follows:

1. **Mulrooney, M.J.**, Leutscher, J., Braathen, A., 2017. A 3D structural analysis of the Goliat field, Barents Sea, Norway. *Mar. Pet. Geol.* doi:10.1016/j.marpetgeo.2017.05.038
2. **Mulrooney, M.J.**, Rismyhr, R., Yenwongfai, H. D., Leutscher, J., Olaussen, S., Braathen, A., (In Press). Impacts of small-scale faults on continental to coastal plain deposition: Evidence from the Realgrunnen Subgroup in the Goliat field, Southwest Barents Sea, Norway.
3. Ogata, K., **Mulrooney, M.J.**, Braathen, A., Maher, H., Osmundsen, P.T.O., Anell, I., Balsamo, F. (In press). Architecture, deformation style and petrophysical properties of a Late Triassic growth fault system in southern Edgeøya (East Svalbard).
4. **Mulrooney, M.J.**, Van Stappen, J., Cnudde, V., Senger, K., Rismyhr, B., Braathen, A., Olaussen, S., Mørk, M.B.E., Ogata, K., (In review). Fluid flow properties of a potential unconventional CO₂ storage unit in central Spitsbergen: the Upper Triassic to Middle Jurassic Wilhelmøya Subgroup.

The first two articles detail the tectonostratigraphy of the Goliat field using well cores and a MAZ survey. Article I provides a detailed structural description of the Goliat area with emphasis on the evolution of fault systems and large fold structures whereas article II focuses on the development of the Late Triassic Realgrunnen Subgroup. Article III details structural descriptions of a Late Triassic growth fault system on eastern Svalbard. Article IV details heterogeneities affecting a potential Late Triassic CO₂ storage site on central Spitsbergen, Svalbard. Chapter 10 comprises an appendix to the thesis which include three additional peer-reviewed articles that were contributed to during the PhD

study and supplement the findings of the main body of the text. The appendix also contains conference proceedings abstracts/short papers, in addition to details of teaching, outreach and miscellaneous activities.

1.3 AUTHORSHIP AND CONTRIBUTION

Mark Mulrooney is the sole author of Part I of this PhD thesis, and the principal author of three of four manuscripts presented in Part II. The approximate contribution of Mark Mulrooney and the co-authors to each manuscript is presented in table format below.

Article I: A 3D structural analysis of the Goliat field, Barents Sea, Norway.

Principal author	Mark Mulrooney
Co-authors	Johan Leutscher, Alvar Braathen
Text	Mulrooney
Figures	Mulrooney
Concept	Mulrooney, Leutscher, Braathen
Editing	Leutscher, Braathen
Data processing and interpretation	Mulrooney, Leutscher (seismic interpretation) Mulrooney (fault analysis)
Approximate Contribution	Mulrooney: 85% Leutscher: 10% Braathen: 5%
Status of manuscript	Published in Marine and Petroleum Geology Volume 86, September 2017, Pages 192-212

Article II: Impacts of small-scale faults on continental to coastal plain deposition.

Principal author	Mark Mulrooney
Co-authors	Bjarte Rismyhr, Honore Yenwongfai, Johan Leutscher, Snorre Olaussen, Alvar Braathen
Text	Mulrooney Rismyhr (Facies analysis + sequence stratigraphy) Yenwongfai (Amplitude vs Angle)
Figures	Mulrooney (Figs 1, 2, 3, 4, 5, 12, 13, parts of 14, 15, 16) Rismyhr (Figs 6, 7, 8, 9, 10, 11, parts of 16) Yenwongfai (parts of Fig 14)
Tables	Rismyhr (all tables)
Concept	Mulrooney, Rismyhr, Olaussen
Editing	all authors
Data processing and interpretation	Mulrooney, Leutscher (seismic interpretation) Mulrooney (Fault analysis) Yenwongfai, Mulrooney (AVA analysis and interpretation) Rismyhr (core logging, petrophysical log interpretation, facies analysis, sequence stratigraphy)
Approximate Contribution	Mulrooney: 45% Rismyhr: 40%

Status of manuscript	Yenwongfai: 5% Other co-authors: 10% in total Marine and Petroleum Geology (In press)
Article III: Late Triassic growth fault system in southern Edgeøya	
Principal author	Kei Ogata
Co-authors	Mark Mulrooney, Alvar Braathen, Harmon Maher, Fabrizio Balsamo
Text	Ogata, Mulrooney
Figures	Ogata
Concept	Ogata, Braathen, Mulrooney
Editing	all authors
Fieldwork and sampling	Ogata, Mulrooney, Braathen, Maher (structural measurements, stratigraphic logging and sampling from outcrop) Mulrooney (Photogrammetry)
Data processing and interpretation	Ogata (meso and macro structural analysis) Mulrooney (Virtual outcrop model generation and analysis)
Approximate Contribution	Ogata: 60% Mulrooney 20% Other co-authors: 20% in total
Status of manuscript	Basin Research in October (In press)
Article IV: Fluid flow properties of the Wilhelmøya Subgroup	
Principal author	Mark Mulrooney
Co-authors	Leif Larsen, Jeroen Van Stappen, Veerle Cnudde, Kim Senger, Bjarte Rismyhr, Alvar Braathen, Snorre Olaussen, Mai Britt E. Mørk, Kei Ogata
Text	Mulrooney Olaussen Senger (natural fractures) Van Stappen (X-ray micro-CT) Larsen (Fluid injection tests)
Figures	Mulrooney (parts of Fig 1,2, Fig 6, 7, 8, parts of Fig 12, Fig 13, parts of Fig 14) Ogata (parts of Fig 1,2) Senger (Figs 9, 10, 14) Van Stappen (Figs 3, 11) Larsen (Figs 4, 5) Rismyhr (parts of Fig 12)
Tables	Mulrooney
Concept	Braathen, Olaussen, Mulrooney
Editing	Mulrooney, Olaussen, Ogata, Braathen
Fieldwork and sampling	Mulrooney (photogrammetry, structural measurements and fault gouge sampling in outcrop) Ogata, Senger (structural and measurements and sampling from core and outcrop) Rismyhr (stratigraphic logging) Van Stappen, Cnudd (core sampling)
Data processing and interpretation	Mulrooney (virtual outcrop model generation and analysis, structural interpretation, study synthesis)

	Van Stappen, Cnudd (Pore-scale characterization using X-ray micro-CT)
	Senger (Structural interpretation)
	Rismyhr (sequence stratigraphic and facies interpretation)
	Larsen (interpretation of fluid injection data)
	Mørk (Diagenesis interpretation)
	Olaussen (aid with synthesis of study data)
Approximate Contribution	Mulrooney: 25%
	Leif Larsen: 10%
	Van Stappen/Cnudde: 10%
	Senger 10%
	Olaussen: 10%
	Ogata: 10%
	Mørk: 10%
	Rismyhr: 10%
	Braathen: 5%
Status of manuscript	Submitted to the Norwegian Journal of Geology (CO ₂ special issue) on 10 October, 2017

1.4 AIMS AND OBJECTIVES

This structural geology study aims to enhance our regional understanding of the tectono-stratigraphy of the Triassic Barents shelf. The work addresses fault-affected hydrocarbon and CO₂ subsurface reservoirs with emphasis on fault evolution, sediment dispersion during fault activity, and perturbations to the petrophysical properties of the host rock. More specifically, the thesis addresses these topics by attempting to answer the following questions:

1. Did regional tectonic and sediment loading forces trigger faulting in the Late Triassic Barents Shelf?
2. Can we better constrain the evolution of the Goliat field trapping structures and fault network using multi-azimuth 3D data?
3. How did faulting in the Late Triassic impact the distribution of sedimentary facies in the Goliat field and on Edgeøya?
4. Can the geometry and deformation mechanisms of growth faults inform the causal driving mechanisms? i.e., tectonic 'vs' gravity driven.
5. How will outcrop scale normal faults and other discontinuities affect fluid flow in the Late Triassic to Early Jurassic Longyearbyen CO₂ lab storage unit?

To address these five questions, offshore studies of the Goliat field area were conducted with the aims of discerning whether fault movement (postulated by Rismyhr, 2012) occurred in the

Hammerfest Basin and on the basin-platform delineating TFFC during the Late Triassic, contemporaneous to other parts of the shelf (e.g., Anell et al., 2013; Mahajan et al., 2014; Indrevær et al., 2016; Serck et al., 2017). The study aimed to define the sequence stratigraphy of the Late Triassic Realgrunnen Subgroup and determine potential circum-arctic tectonic events responsible for their generation. An additional objective in this offshore investigation was to describe the gross structural geology of the Goliat field and determine the evolutionary history and mechanisms responsible for development of the Goliat anticline and the population of faults affecting it. This aspect of the investigation serves to gain knowledge of a structurally complex hydrocarbon field that is situated within a bend on a basin–platform delineating fault complex (TFFC), which to date has not been adequately described owing to single azimuth survey related resolution issues created, in part, by fault shadow. Furthermore, this investigation of the structurally compartmentalised Goliat field can be used as an analogue for onshore Svalbard where the approximate time equivalent Wilhelmøya Subgroup is also compartmentalised but poorly imaged by terrestrial seismic (Bælum et al., 2012).

Secondly, the thesis attempts to determine how Triassic faulting affected the distribution of sediments, especially sand prone reservoir quality facies, using both the offshore Goliat case and the approximate time equivalent Triassic successions onshore Kvalpynten, Edgeøya. The onshore, listric faulting at Kvalpynten is investigated due to observations that they rerouted and captured sand prone facies in an otherwise fine grained pro-delta environment. Here, the excellent cliff exposures allow both micro- and meso-scale investigations to be conducted in order to characterise fault architecture and to discern the main deformation mechanisms. Moreover, this work had the ambition to discern the spatio-temporal evolution of the faults (including the history of diagenesis and compaction) in order to understand how the control of fault damage zones on fluid flow properties can change throughout the evolution of a growth fault system. Furthermore, given the non-conventional characteristics of the Kvalpynten growth fault system (counter regional dips, non curvilinear fault traces) the study had the objective investigating an apparent tectonic triggering mechanism. Offshore, the study aimed to construct palaeogeographic reconstructions of the depositional environment during deposition of the late Norian to Rhaetian Realgrunnen Subgroup, with emphasis on fault related rerouting of sand prone (proven reservoir in the Goliat field) channelised and alluvial fan facies deposited in terrestrial and coastal plain environments. Here the goal is to further inform the architecture of the uppermost Goliat reservoir with a vision towards enhancing production.

Thirdly, the thesis aimed to characterise the Late Triassic to Early Jurassic Longyearbyen CO₂ lab storage unit (Wilhelmøya Subgroup, the counterpart of the offshore Realgrunnen Subgroup), onshore Svalbard, with a goal of subdividing the target reservoir unit based on varying fluid flow

properties by conducting petrophysical and petrological analysis of drill cores and outcrops. Structural discontinuities were investigated in drill core and outcrop in order to characterise flow-baffling features in the subsurface, which are evident from fluid injection tests. This study will inform potential future CO₂ injection in the subsurface of Svalbard as well as forming an analogue for offshore investigations of the Realgrunnen Subgroup which has proven reservoir potential in the Barents Shelf.

2. SCIENTIFIC BACKGROUND: FAULTS AND FAULT GROWTH

2.1 INTRODUCTION

Faults, fault geometries, and displacement characteristics are essential to this study. Therefore, the following section is used to establish a fundamental scientific basis for the context of the thesis. A fault can be defined as a sharp structural discontinuity defined by slip planes (surfaces of discontinuous shear displacement) and/or thin tabular shear zones and other related structures commonly ascribed to fault core and damage zones that formed during the evolution of the structure (Schultz & Fossen, 2008). The Earth's crust is fragmented into an intricate array of discontinuous faults which grow incrementally over time amounting displacement, in most cases during earth quake rupture events (Peacock, 1991; Peacock and Sanderson, 1991; Walsh and Watterson, 1991; Cartwright et al., 1995; Cartwright et al., 1996; Childs et al., 1995, 1996; Willemse et al., 1996; Meyer et al., 2002; Wilkins and Gross, 2002; Childs et al., 2002; Soliva & Benedicto, 2004; Walsh et al., 2003; Kim & Sanderson, 2005). Faults can be driven by tectonic stresses or by gravity. Fault expressions on profile maps generally show intricate arrays of discontinuous overlapping fault segments which vary in size by two or greater orders of magnitude on average and have a consistent strike direction associated with regional stress tensors (or slope profile in the case of gravity driven faults). Systematic variations in displacement to length ratios along individual faults and variations in fault displacement where adjacent faults overlap, suggest all faults within a population form a geometrically coherent system which accommodates strain variation (Mansfield and Cartwright, 2001). Fault growth has also been shown to consist of a complex process of lateral and vertical linkage of individual segments as well as bifurcation into additional segments over the history of activity (Childs et al., 1995, 1996, 2002; Cartwright et al., 1995, 1996; Marchal et al., 2003; Walsh et al., 2003).

Faulting can be broadly classified into 3 different types (normal, reverse and strike-slip, i.e., Andersonian fault types, after Anderson, 1951) based on kinematics which in turn relate to the orientation of the principal stresses responsible for their deformation. In this contribution, we focus on normal (extensional) faulting which is associated with rift basins. The distribution, geometries, and segmentation history of extensional faults has significant implications for hydrocarbon prospecting and CO₂ storage. During rifting, the surface expression of faulting in a basin has a strong local effect on geomorphology and thereby the routing and site of deposition of reservoir quality sand bodies. Post deposition, faults can act as conduits, baffles, and trapping structures within hydrocarbon reservoirs (Fossen & Hammerfest, 1998). Their importance has facilitated investigative studies resulting in major advances in understanding fault growth in recent years. The current

understanding of fault evolution is summarised in this section and underpins the scientific articles herein.

2.2 DRIVING MECHANISMS

Tectonically induced normal faults are generally associated with tensile stresses during lithospheric stretching and thinning, the processes responsible for eventual continental breakup and seafloor spreading (Wilson, 1966; Burke & Dewey, 1973; Mckenzie, 1978; Barr, 1987 Duncan & Turcotte, 1994). The continental crust accommodates the up-rise and decompression of the underlying asthenosphere (the process known as knecking) mainly by brittle strain. Commonly, early rift sediments are down-faulted into the developing rift graben while erosion takes place contemporaneously on uplifting margins.

Faulting can also be gravity driven, i.e., the result of gravitational collapse in depositional environments such as deltas where sediments build out over a slope and becomes critically unstable. These faults are commonly observed in modern and ancient delta tops and fronts and are generally linked to a toe thrust zone in the corresponding prodelta (Hooper et al., 2002; Imber et al., 2003; Fort et al., 2004; Morely et al., 2011). This thin skinned deformation is usually facilitated by the presence of a weak detachment layer such as evaporates or over-pressured muds. The resulting faults are often scoop-shaped and show transport directions in the downslope direction. A deltaic setting makes fault-created accommodation prone to routing and capturing sandy facies (Osmundsen et al., 2014).

2.3 EARTH QUAKE SLIP AND THE IDEAL FAULT MODEL

Displacement accrues systematically on faults during individual earth quake ruptures (Watterson, 1986; Walsh & Watterson, 1988; Marrett & Allmendinger, 1991; Cowie & Scholz, 1992; Dawers et al., 1993; Cartwright et al., 1995; Dawers & Anders, 1995; Bull et al., 2006; Nicol et al., 2010). Quantitative constraints from ancient faults indicate that the maximum displacement (D) of a fault scales with fault trace length (L), which forms the basis of the conventional growth model.

$$D = cL^n$$

where c is a constant and n is a variable that typically ranges from 1.0 – 1.5.

Systematic maximum displacement to Length ratios ($D:L$ in Fig. 1) have resulted in a simple earthquake-slip model which suggests fault geometries are directly related to the summation of earthquake slip during rupture events. The resultant ideal fault model (Fig. 2) is a discontinuous

Figure 1 Plots of maximum displacement (d_{max}) against fault length (L) from a collection of publications (compiled by Kim & Sanderson, 2005).

Figure 2 A) A rock volume with an idealised elliptical fault surface exhibits maximum displacement towards the centre and deformation in the surrounding rock volume (modified from Walsh & Watterson, 1991). B) Graphical representation of maximum displacement (d_{max}) with fault length (L).

elliptical structure with maximum displacement at the centre that dissipates laterally. The discontinuous nature of faults requires deformation in the surrounding rock volume which is often referred to as 'drag' (see section 2.13 below). Tip-line propagation occurs contemporaneously to

displacement accumulation with a systematic relationship being maintained throughout evolution. The ideal fault model has proven very useful in understanding fault growth but does not realistically occur in nature as will be demonstrated below.

2.4 SHORTCOMINGS OF THE CONVENTIONAL GROWTH MODEL

Despite the usefulness of the conventional fault growth model, it fails to reconcile inconsistencies between observed fault and earth quake D:L ratios, with the former often exhibiting higher ratios (Walsh et al., 2002). In addition, displacement studies (e.g., Mouslopoulou et al., 2009) reveal short term fault displacement rates (≤ 20 kyr) to depart from million year average rates by up to three orders of magnitude. Larger time scale sampling results in smoothing of displacement rates. Variation in displacement rates is not consistent with the conventional earthquake slip model. Displacement rates and fluctuations in earthquake recurrence intervals are suggested to result from fault interactions (e.g., linkage and segmentation; see section 2.8). Problems in modelling fault displacement rates are primarily caused by inconsistent sampling rates, i.e., outcrop and seismic data provide million year sampling intervals while modern seismicity provides near instantaneous sampling. Synthesis of both these timescales is required to adequately describe fault displacement rates, with precisely age controlled syn-kinematic sediments proving the most useful in such ongoing studies.

2.5 STUDYING FAULT KINEMATICS

Variation in hanging wall and footwall thickness can be used to record throw accumulation of each interval (bound by traceable horizons) and throw subsequent to deposition. Back-stripping is the process where the total displacement accumulation on a fault is restored incrementally by successive marker horizons allowing the history of fault activity (i.e., initiation, hiatus, reactivation and arrest) to be determined. Displacement rate analysis is mainly derived from displacement measurements from seismic interpretation as well as from outcrop geology, trenching and gravity modelling. Ages of displaced horizons are defined by radiocarbon dating, tephrochronology, fission track dating and biozonation. Average displacement rates are calculated by the vertical displacement accrued on a reference formation over the time interval of deposition. Seismic displacement rate analysis requires preservation of syn-kinematic sediments and precise age control of horizons over a significant span of the fault's kinematic life and requires a delicate balance to be maintained between sedimentation rates, erosion, regional tectonic uplift and displacement through long periods of time. Sedimentation rates must be greater than displacement rates and footwall erosion must be minimal. Error in such analyses occurs due to footwall erosion,

differential compaction and variations in interval velocities. Moreover, displacement measurements are limited by the accuracies of horizon and fault interpretation.

2.6 DISPLACEMENT TERMINOLOGY

General confusion can occur when dealing with the subject of fault geometry and kinematics mainly due to the inconsistent use of terminology; Fig. 3 attempts to synthesise some common fault

Figure 3 A synthesis of terminology used to describe fault displacement (modified from Lohr et al., 2008).

nomenclature. The main confusion arises from the description of displacement. Real displacement is measured between a reference horizon along the fault plane while heave and throw are linear horizontal and vertical measurements of horizon separation, the latter often being used as a proxy for real displacement (Lohr et al., 2008). This is in parts due to the simpler calculation algorithm of throw in fault analysis software. Terms referring to the size of a fault (i.e., a small or a large fault) are usually a proxy for length (along axis of fault tip propagation), or throw. The orientation of displacement measurement is important when interpreting segmentation and linkage history in 2D and 3D seismic data. Large variations occur in real, vertical and horizontal displacement especially where former isolated segments have linked to form a larger through-going fault. In these areas, real displacement (measured along the fault plane) can vary dramatically, represented as undulations in D vs L plots (Alan diagrams). Throw in these cases is a poor approximation of real displacement and does not exhibit the same dramatic variations which results in smoothing of D:L curves and possible loss of information. Furthermore, fault geometry affects the relationship between heave and throw,

e.g., listric faults will have much larger heaves than planar faults with the same real displacement (Wheeler, 1987; Dula Jr, 1991).

2.7 ISOLATED AND COHERENT FAULT MODELS

Fault arrays generally consist of multiple fault segments which can vary in size and even orientation. The classic 'isolated fault model' (e.g., Cartwright et al., 1996) involves individual faults growing in separation similar to the ideal fault discussed above (section 2.3). In this model, kinematically independent faults nucleate radially followed by incidental overlap with neighbouring segments which consequently form relay zones. The isolated fault model is mainly derived from 2D data and assumes that the fault propagation direction is in the plane of inspection. In reality, fault propagation is a complex 3D phenomenon and cannot be properly described or understood by 2D observations. Detailed studies of fault arrays have shown individual faults may have up to 5 orders of population size (Walsh and Watterson, 1991). Displacement contours plotted on strike projections of single and multiple faults show simple systematic patterns and spacing. Displacement profiles of an array of faults along fault trace lengths exhibit the same systematic displacement distribution as one large fault suggesting kinematic coherency which requires synchronous movement on all faults in contrast to a sequential order suggested by the isolated fault model (Mansfield and Cartwright, 2001; Walsh et al., 2003). The 'coherent fault model' suggests individual fault segments initiate and grow as kinematically related components of a larger array.

2.8 LINKAGE AND SEGMENTATION

Relay zones form when neighbouring faults overlap and transfer displacement (Peacock & Sanderson, 1991; Trudgill & Cartwright, 1994; Huggins et al., 1995; Childs et al., 1995, 1996, 2016; Kristensen et al., 2008). This process can occur due to interaction between initially isolated faults or by bifurcation of a single fault. Bifurcation (also known as segmentation) is the process where faults splay into two or more segments and is usually credited to changes in host rock mechanical properties or changes in local stress due to interactions with neighbouring faults which causes retardation of fault tip propagation (Kristensen et al., 2008). Bifurcation can result in apparent segmentation in certain planes of investigation but in reality the bifurcated faults likely link in three dimensions (Fig. 4). Progressive overlap of fault segments usually results in breaching and linkage of the segments. Soft linkage (Fig. 5) is the term used to describe ductile strain (continuous deformation) within the relay zone of two interacting faults where displacement is transferred without physical connection of the two faults. This form of linkage requires the host rock to be able

to accommodate shear strain across the relay without failure of the structure and will not occur in rocks with high shear strength. Hard linkage or breaching (Fig. 5) occurs when displacement transfer is accommodated by the propagation of the footwall and/or the hanging wall or a new breaching fault. When both faults propagate, an isolated fault bound lens or ‘eye structure’ is formed. Bifurcation and relay zone breach is a 3D phenomenon and the modes of overlap are reflected by the 3D geometry (Fig. 4). Relay zones can be unconnected in 3D (soft linkage) or can be linked in 3D at branch lines or branch point (hard linkage in both cases). Walsh and Watterson (1991) suggest the identification of soft linkage may be limited by data resolution and that higher resolution investigations may often reveal brittle faults breaching relay zones. Relay zones are continually formed and destroyed throughout fault growth and so record all stages of evolution when arrest occurs (Childs et al., 1995).

Figure 4 Schematic diagram of the three possible structures underlying a segmented fault zone. Individual segments may be (A) unconnected in 3D, B) linked at depth along branch lines or (C) linked at depth at branch points (modified from Childs et al., 1995).

Figure 5 Three possible ways in which a relay zone can be breached. A) Propagation of the hanging wall or the footwall fault (shown). B) Propagation of both the footwall and hanging wall faults. C) Development of a secondary linking fault (modified from Childs et al., 1995).

2.9 INFLUENCE OF BASEMENT STRUCTURES

Meyer et al. (2002), Walsh et al. (2002) and Giba et al. (2012) support an alternative model for fault growth which contradicts the conventional growth model where maximum displacement and length are believed to accumulate systematically. The alternative model suggests reactivated basement faults propagate upwards and result in quick (instantaneous on geological timescales) establishment of fault length with subsequent fault growth dominated by increase in displacement. The model predicts a progressive increase in D:L ratios over time which reconciles problems with the scaling properties of faults and earth quakes where faults (especially larger faults) often exhibit higher D:L ratios than earth quakes. Retardation of fault-tip propagation is credited to interaction between fault tips which alter local stresses. The time taken to reach interaction stages is dependent on a number of factors including strain rates and the rheology. The model also predicts a larger variation in fault sizes during initial stages of extension. Application of the alternative fault growth model to basins which have not experienced re-activation is still unknown and remains a popular topic of debate.

2.10 INFLUENCE OF MECHANICAL STRATIGRAPHY

Further complicating D:L scaling laws is the effects of strong mechanical stratigraphic contrasts on fault propagation (e.g., Rippon, 1985; Gross & Engelder, 1995; Gross et al., 1997; Ackermann et al., 2001; Schultz & Fossen, 2002; Wilkins and Gross, 2002). As faults encounter and breach multiple mechanical layers, populations may exhibit different scaling as mechanical properties and contacts between adjacent lithologic units can create effective barriers that inhibit fault propagation. This is highlighted by Soliva & Benedicto (2005) where small normal faults were shown to be vertically restricted to brittle carbonates by overlying and underlying plastic claystone layers. Displacement profiles for these faults were shown to evolve from a linear (during propagation through the carbonates) to a flat-topped D:L distribution once the faults become vertically restricted by the plastic clay layers, i.e., fault growth becomes dominated by propagation of length and relative stasis of displacement accumulation. Further observations show large vertically restricted faults accommodate less relative strain than unrestricted faults and so for a given strain, vertically restricted normal fault systems require more faults than unrestricted systems. Ackermann et al. (2001) suggested systems may oscillate between different D:L distributions as they progressively encounter and eventually breach confining stratigraphy (Fig. 6).

Figure 6 Faults propagating radially through stiff sandstone until encountering shale layers (B) where faults are temporarily arrested from propagating in the vertical dimension and layer parallel growth becomes temporarily dominant. C) Faults eventually cross-cut the shale and (D) continue to grow radially (modified from Fossen, 2016).

2.11 STAGES OF FAULT GROWTH

Analysis of fault throw populations in syn-kinematic horizons (e.g., Meyer et al., 2002) show general constant D:L ratio slopes for each horizon, however an up-sequence shallowing of slope and geometric moment populations demonstrate a progressive concentration of strain onto fewer and larger faults over time. Continued extension will ultimately results in a single active through-going fault. Kinematic analyses show that higher strain rates result in faster moving faults and that strain rate has no effect on fault population numbers. As a consequence of strain localisation, propagation strain accommodated by smaller faults becomes obsolete and is consistent with the observation that smaller faults have higher termination (mortality) rates (Fig. 7). Meyer et al. (2002) have used the temporal changes in strain localisation to construct a generalised 3 stage model for fault growth where, 1) Initial growth is dominated by lateral tip propagation resulting in long faults with relatively small amounts of displacement (low D:L ratios). Displacement rates are established quickly and remain constant. As larger faults move faster due to strain localisation, the hierarchy of faults is established quickly. 2) Tip line propagation results in fault interactions. Changes in local stress fields

within fault relay zones results in retardation of lateral growth (stasis). This stage in growth development is dominated by fault displacement and fault populations exhibit higher D:L ratios. 3) Late stages in fault growth are often characterised by an up section reduction in the trace of active

Figure 7 Schematic diagram showing progressive fault growth by linkage.

faults and has been linked to slowing in regional extension rates. Strain is accommodated by larger faults with small faults having higher mortality rates. Late stage increase in fault length is mainly due to amalgamation of existing coeval faults by fault capture and relay zone failure. Extensional fault systems are described by the soft-domino model (Walsh and Watterson, 1991) where each block has a systematic nonlinear variation in displacement and ductile strain plays an important role in displacement transfer. Fault blocks rotate over time resulting in shallower fault dips than when they were initiated (Fig. 8).

Figure 8 Conceptual block diagram showing the principal features of the soft-domino model (modified from Walsh and Watterson, 1991).

2.12 IDENTIFYING LINKAGE HISTORY

As discussed above, fault growth by means of segmentation and linkage is determined by syn-kinematic packages which can be used to study the spatio-temporal history of individual fault segments. 3D seismic imaging can reveal subtle changes in syn-kinematic packages associated with the segmented nature of a fault array (Walsh et al., 2003). In addition, scaled Analogue Sandbox Modelling and Discrete Element Modelling (DEM) have been used to observe fault growth in laboratory conditions (McClay & Ellis, 1987; McClay, 1990; Cartwright et al., 1995; Finch et al., 2004; Imber et al., 2004; Schopfer et al., 2006). Determining the former history of segmentation and linkage in mature fault systems may be desirable in order to understand the structural evolution of a rifted basin and to assess fault sealing capacities where they act as potential structural traps. Fault displacement analysis often reveals the existence of sub horizontal anomalies in distribution of throw along a fault trace which deviate from the expected elliptical displacement on an ideal fault. Sub horizontal anomalies are interpreted by Mansfield and Cartwright (1996) as sites of former linkage. Displacement minima along a fault trace are interpreted as points where fault breach and linkage has occurred. Displacement maxima are said to represent the centre of former segments where displacement rates are highest. In addition, D:L profile plots are shown to change in character where linkage has occurred. Lohr et al. (2008) suggest isolated fault segments exhibit a characteristic triangular shaped displacement to length profile where maximum displacement is represented by a sharp peak, while linkage of several segments results in a more gradual elliptical D:L profile. In addition to displacement analysis, Lohr et al. (2008) demonstrate the use of fault attribute analysis which can reveal undulations or corrugations in a fault plane which can also be used to identify sites of former segmentation and linkage. Attributes including strike, dip, throw accumulation and curvature can highlight anomalous undulations along a fault plane and are shown to complement interpretations made from displacement analysis.

2.13 DEFORMATION OF THE SURROUNDING WALL ROCK

The rock volume surrounding faults is geometrically and coherently linked to fault movement. Footwall uplift and hanging wall subsidence results in net rotation of fault blocks (Barr, 1987; Wernicke & Axen, 1988; Yielding, 1990; Long & Imber, 2010). The resultant geometries of this phenomenon can be described as longitudinal (sub-parallel to fault) and transverse (orientated at high angles to faults) folding in the hanging wall and footwall that form at the same time as or just prior to faulting (Schlische, 1995; summarised in Fig. 9). Drag folds form longitudinal forced folds or fault propagation folds that form hanging wall synclines and footwall anticlines (Withjack et al., 1990; Schlische, 1995; Corfield and Sharp, 2000; Khalil & McClay, 2002; Sharp et al., 2000; Willsey et

al., 2002; Finch et al., 2004; Jackson et al., 2006; Braathen et al., 2011; Maher & Braathen, 2011). More locally, drag folds can form from frictional drag and differential compaction. Minor monoclinal folding occurs at the trace tips of faults. Reverse drag folds result from a decrease in displacement away from the fault and form hanging wall anticlines and footwall synclines. Hanging wall roll-over folds form monoclines as a response to movement over listric faults (Gibbs, 1984; Williams & Vann, 1987; Ellis & McClay 1988; McClay & Scott, 1991, Imber et al., 2003). Transverse folds result from along-strike variation in fault displacement. In the hanging wall, broad synclines form with maximum displacement towards the centre of segments, whereas narrow anticlines occur at segment boundaries, i.e., relay zones, which remain even after breach (Childs et al., 2016). Conversely, broad anticlines, and narrow synclines are observed at corresponding locations in the footwall. Interference folds result from the superposition of longitudinal and transverse folds, and include domes, basins, culminations, and saddles.

Figure 9 Geometric relationships among a segmented normal fault system, relay ramps, transverse folds, and rider blocks (modified from Schlische, 1995).

2.14 FAULT GEOMETRIES

Further to deformation in the immediate surrounding rock volume, on a crustal scale, the geometry of large fault systems and the underlying detachment system exerts fundamental control on structures produced in the hanging wall. The progressive development of extensional fault systems in the upper most 10 km of lithosphere has been studied compressively by analogue modelling (e.g., McClay & Ellis, 1987; Ellis & McClay, 1988; McClay, 1990; Fossen & Gabrielsen,

1996; Dooley et al., 2003). In these experiments (summarised in Fig. 10), extension above a horizontal basal detachment which undergoes stretching over a limited area produces an asymmetric rift graben bound by planar faults and in which the internal deformation is accommodated by rotated fault blocks, consistent with structures common in intra-continental rifts. Extension above a basal detachment which undergoes extension under the whole model, and is not laterally constrained, produces domino-style fault arrays comparable to highly extended rift basins. Extension above listric faults produce hanging wall roll-over and crestal collapse graben systems, whereas more complex ramp-flat listric extensional fault systems produce an upper roll-over crestal collapse system, a ramp syncline and reverse fault/fold zone, and a lower roll-over crestal collapse system (Ellis and McClay, 1988; McClay, 1990).

Figure 10 Geometric relationships between fault geometries and hanging wall deformation. A) Planar faults. B) Simple listric faults. C) Ramp-flat listric faults (simplified from analogue model experiments by McClay, 1990).

2.15 INVERSION OF NORMAL FAULTS

Positive inversion, referred to herein as basin inversion (e.g., Bally et al., 1981; Cooper et al., 1989; Williams et al., 1989; Bonini et al., 2012) refers to a switch in tectonic mode, i.e., from extension to contraction and can result in basins becoming positive structural features (Cooper et al., 1989). Essential to the concept of inversion tectonics is a component of fault reactivation such that extensional faults become reactivated as contractional faults (Bally et al., 1981; Cooper et al., 1989). Driving mechanisms of basin inversion are diverse and can result from changes in plate motions, ridge/plume push, isostatic uplift, halokinesis and the onset of subduction. The concept of a regional elevation, extrapolated from outside the area of tectonic deformation, is vital for recognition of inversion tectonics, i.e., inversion is apparent where syn-rift successions are observed to be elevated above this datum. This process is also recognisable by characteristic “harpoon” or “arrow-head” structures where growth packages in half grabens are inverted (Fig. 11). As normal

Figure 11 Common basin inversion features of a pre-rift and syn-rift succession. A) Initial basin configuration before inversion. B) Harpoon or arrow-head structure. C) Footwall short-cut fault. D) Hanging wall bypass fault.

faults are not ideally orientated to facilitate reverse motions (Anderson, 1951), inversion of high angle dip-slip faults is unfavourable. Deeper parts of listric faults (with low dip angles) are often reactivated whereas upsection, new faults are commonly generated either in the hanging wall (hanging wall bypass faults), in the footwall (footwall short-cut faults) or splaying from the tip points of the original extensional faults (Fig. 11). The generation of these structures has been demonstrated in analogue models (McClay, 1989; Buchanan & McClay, 1991). In addition, inversion of domino block faulting showed dramatic back-rotation and steepening of faults, harpoon structures and a dominance of footwall short cut faults. Inversion of simple listric faults showed reactivation of the main detachment, tightening of the crestal collapse graben, formation of new thrusts from tips of crestal collapse faults and generation of harpoon structures. Finally, inversion of ramp-flat listric

faults showed reactivation of parts of the more preferentially orientated main detachment, an abundance of hanging wall by-pass faults, and a complete bypass of the upper roll-over structure. Inversion buttressing (e.g., Dart et al., 1995; McClay, 1995; Baum et al., 2008) is the formation of folds in the hanging walls of non-planar normal faults during basin inversion due to steep upper parts of the fault plane inhibiting slip. These folds commonly occur as single hanging wall anticlines adjacent to the fault zone or as a couplet of a hanging-wall syncline and anticline.

2.16 FAULT CONTROL ON SEDIMENTATION

The sedimentary response to the aforementioned tectonic topography varies with regard to the stage of rift development, depositional setting, i.e., continental vs marine and climatic conditions (Prosser, 1993; Gawthorpe & Leeder, 2000). Of importance for this body of work, is that initial stages of rifting in a continental setting are dominated by antecedent drainage. The aforementioned transverse hanging wall synclines form individual isolated basins associated with hanging walls of individual fault segments which over time coalesce as faults progressively link. Contemporary to increase in hanging wall accommodation space, the corresponding upwards movement in the footwall leads to erosion, and establishment of semi-circular drainage catchments most developed at the centres of segments and characterised by erosional linear channels fed by larger dendritic channel systems (Leeder, 1991; Leeder & Jackson, 1993; Eliet & Gawthorpe, 1995; Densmore et al., 2003, 2004; Foster et al., 2010; Elliott et al. 2011).

In fluvial and deltaic environments, axial through-flowing river channels and delta lobes tend to migrate or avulse towards the axis of maximum subsidence, i.e., parallel to the structural strike (Bridge and Leeder, 1979; Miall, 1981). Preferential flooding of the down-tilted side of the floodplain also occurs. Channels become more confined due to tilting, and stacked channel geometries become dominant (Gupta et al., 1999; Young et al., 2002; Garcia-Garcia et al., 2006). Individually, tilting can cause sandstone bodies to become abnormally wide and asymmetric (Alexander & Leeder, 1987).

Relay zones and erosional fault scarps act as loci for drainage entering rift basins. Typical early stage rift deposits include continental alluvial fans (Hooke, 1967; Nemec & Postma, 1993; Blair & McPherson, 1994; Harvey et al., 2005) or fan deltas (Holmes, 1965; Rust, 1979; Friedman & Sanders, 1978; Ethridge & Wescott, 1984; Nemec & Steel 1988; Postma, 1990). Both deposits consist of coarse siliciclastics that are usually immature to sub-mature both texturally and mineralogically. A schematic of tectonic influence on continental facies during the initial phase of rift basin development is summarised in Fig. 12.

Figure 12 Sediment routing in a continental early rift system. Antecedent drainage is diverted by footwall uplift and enters hanging wall basins via unbreached relay zones. Isolated sub-basins develop at the centres of fault segments and amalgamate over time as faults link. Hanging walls are characterised by axial drainage where rivers avulse or migrate down dip towards the fault plane.

2.17 IMPLICATIONS ON RESERVOIR PROPERTIES

Faults can dramatically impact reservoir properties where they can act as transmitters of, and/or barriers to, fluid flow and pressure communication (Knipe, 1992; Knipe et al., 1998). Fault seal can arise from two primary scenarios: 1) Juxtaposition seals where reservoir units are juxtaposed against a low-permeability unit (e.g., shale) with a high entry pressure, and 2) membrane seals, i.e., where fault rock exhibits high capillary entry pressure. The dominant control on failure of membrane seals is the capillary entry pressure of the seal rock; that is, the pressure required for hydrocarbons to enter the largest interconnected pore throat of the seal. When the entry pressure has to exceed the strength of the rock in order to breach the seal, the seal is considered a hydraulic seal (Knipe, 1992). Several variable types of fault rocks can lead to sealing faults and are listed in order of importance, i) clay smear, i.e., the entrainment of intact clay or shale layers into the fault plane, ii) cataclasis, the crushing of sand grains to produce a fault gouge of finer grained material, and iii) diagenesis, when preferential cementation along an originally permeable fault plane may partially or completely remove porosity, ultimately creating a hydraulic seal. These parameters vary over the fault surface, implying that faults cannot simply be designated sealing or non-sealing. The 3D geometry, linkage history and distribution of displacement on surfaces influence the sites of

reservoir juxtaposition between hanging walls and footwalls which has implications for sealing and permeability capacity across faults. Zones of linkage can form gaps in otherwise sealing faults, while breached relay ramps can often form small structural traps. In addition, understanding the intrinsic nature of fault evolution can help predict the spatial and temporal occurrence of seismicity, and the criteria for reactivation of faults.

The common division of faults (irrespective of throw) into core and damage zone reflects the distribution of strain, from a high-strain fault core in which most primary features of the rock is erased, to the damage zone offering discrete structures in otherwise mostly intact host-rock. Apart from juxtaposition, fault elements can affect fluid flow within a reservoir. Faults incorporate a volume of deformed rocks that hosts structures that approximately parallel the fault. The fault core commonly includes structural elements (or facies) such as membranes (fault-parallel layers) of gouges (of variable composition), breccias (of progressive intensity), and smears (of different continuity), as well as lenses (of host rock and fault rock), and discrete structures, e.g., deformation bands, fractures, and minor slip surfaces. The latter discrete structures make up strain in the rock volume around the core, known as a damage zone (e.g., Caine et al., 1996; Kim et al., 2004; Schueller et al., 2013; Choi et al., 2016); the thickness (or width) of which correlates with fault displacement (e.g., Knott et al., 1996; Shipton and Cowie, 2001). Damage zones (Fig. 13) also form ahead of fault

Figure 13 Conceptual block-diagram showing fault zone architecture (from Choi et al., 2016).

tips, i.e., within the process zone (Shipton and Cowie, 2001; Braathen et al., 2004; Braathen et al., 2013) and as such will affect fluid flow over a larger area than the fault core. Deformation bands (Fossen et al., 2007 and references therein), i.e., tabular structural discontinuities having a continuous change in strength or stiffness across a relatively narrow zone, occur in porous rocks. They can be ascribed to grain reorganisation, e.g., grain sliding, rotation and/or fracturing during

overall dilation, shearing, and/or compaction. Classified by either their kinematics or based on deformation mechanisms, deformation bands can increase or diminish porosity and permeability in otherwise good reservoir units. The dominant deformation mechanisms are (1) granular flow (disaggregation bands; grain boundary sliding and grain rotation), (2) cataclasis (grain fracturing and grinding or abrasion), (3) phyllosilicate smearing, and (4) dissolution and cementation. In general, cataclastic, phyllosilicate, and cemented deformation with shear and/or compression are detrimental to reservoir permeability whereas disaggregation and dissolution bands with or without components of dilation can increase permeability. Shear-compaction cataclastic deformation bands that lower porosity and permeability are, however, most common (e.g., Fossen et al., 2007).

An increase in fracture intensity in relation to background intensity is also a common feature of faults and damage zones. Ogata et al., (2014) identify various types of fracture corridors, including (i) damage zones of faults and (ii) fault-tip process zones, both in low-porosity rocks not prone to formation of deformation bands. Further, (iii) fold-crest fracture corridors are identified in footwall rebound anticlines caused by fault-growth. Open fracturing parallel/subparallel to faults may enhance along-fault fluid flow, whereas healed or mineralised fractures may reduce the permeability of the host rock.

3. GEOLOGICAL CONTEXT – THE BARENTS SHELF

The scientific articles herein focus on aspects of the structural evolution of the Barents shelf and as such section 3.1 summarises the Palaeozoic to Recent evolution of this large shelfal area and the geological context of the studies.

3.1 STRUCTURAL ELEMENTS

The Barents shelf is a large area of submerged Eurasian continental crust north of the Norwegian and Russian coastlines. The shelf stretches from Svalbard and Franz Josef Land in the north, Novaya Zemlya to the east, and the Norwegian-Greenland Sea to the west (Smelror et al., 2009). Primary structural elements including basins and plateaus are summarised in Fig. 14.

Figure 14 Bathymetric map of the Arctic Ocean showing the location of the western Barents Shelf, sourced from the IBCAO bathymetric model (2012). B) Main structural elements of the western Barents shelf and adjacent areas modified from Faleide et al. (202009, 2010). Abbreviations: BB = Bjørnøya Basin, FSB = Fingerdjupet Sub-basin, GH = Gardarbanken High, HB = Harstad Basin, HfB = Hammerfest Basin, HFZ = Hornsund Fault Zone, KFC = Knølegga Fault Complex, KR = Knipovich Ridge, LH = Loppa High, MB = Maud Basin, MH = Mercurius High, MR = Mohns Ridge, NB = Nordkapp Basin, NH = Nordsel High, OB = Ottar Basin, PSP = Polheim Sub-platform, SB = Sørvestsnaget Basin, SFZ = Senja Fracture Zone, SH = Stappen High, SR = Senja Ridge, TB = Tromsø Basin, TFP = Troms-Finnmark Platform, VH = Veslemøy High, VVP = Vestbakken Volcanic Province.

Onshore investigations (articles III and IV) herein, were conducted on the Svalbard Archipelago which represents part of the emergent, uplifted northwest Barents shelf where the uplifted strata provide outcrop analogues for the petroleum systems of the southwestern Barents

shelf currently targeted for hydrocarbon exploration (Elverhøi et al., 1988; Riis & Fjeldskaar, 1992; Faleide et al., 1993, 2008; Henriksen et al., 2011). Offshore investigations (articles I and II) were conducted on the Goliat field on the southern margin of the Hammerfest Basin.

The Hammerfest Basin (Rønnevik & Jacobsen, 1984; Berglund et al., 1986; Gabrielsen et al., 1990; Knutsen & Vorren, 1991; Ostanin et al., 2012) is a relatively shallow Mesozoic basin, 50–75 km wide, elongate in the ENE–WSW direction and situated on the south-western flank of the Barents Shelf. The basin is separated from the Finnmark Platform to the southeast by the Troms-Finnmark Fault Complex (TFFC). Separation from the Tromsø Basin to the west is defined by the approximately N–S striking Ringvassøy-Loppa Fault Complex (RLFC). The basin can be divided into sub regions, east and west, possibly defined by the offshore extension of the Trollfjord-Komagelv Fault Zone (TKFZ). The western region exhibits a gentle, westerly dip with an open central dome. The central part of the basin is characterised by an elongate gentle central dome which strikes parallel to the basin axis and is underpinned by the E–W trending Hammerfest Basin Fault System (Gabrielsen, 1984), which trends slightly oblique to the TFFC (ENE–WSW to NE–SW). Local fault populations adjacent to the TFFC and the RLFC exhibit ENE–WSW to NE–SW and N–S to NNE–SSW orientations, respectively, i.e., parallel to subparallel to the fault complexes.

The TFFC (Øvebro & Talleraas, 1977; Rønnevik et al., 1982; Gabrielsen et al., 1984, 2010; Gabrielsen & Færseth, 1988, 1989; Indrevær et al., 2013) is a major basement involved, basin delineating fault system that exploits a Caledonian trend (Faleide et al., 1984; Gudlaugsson et al., 1998; Worsley, 2008). The fault complex parallels the shoreline of the Troms and Finnmark counties and divides a landward plateau (Finnmark Platform) in the south from basins in the north, e.g., the Harstad, Tromsø and Hammerfest basins (Figure 14). The complex, or fault system, consists of several hard linked segments which exhibit orientations of NNE–SSW to NE–SW in the south, ENE–WSW in the centre, and ENE–WSW towards the north-eastern extent of the structure. The complex terminates to the northeast against the offshore extension of the TKFZ.

3.2 PALAEOZOIC TO CENOZOIC SHELF EVOLUTION

Geophysical studies (Ritzman & Faleide, 2007; Marelllo et al., 2010) suggest the Barents Shelf is underlain by a collage of Baltican, Laurentian and Siberian plates along with smaller continental blocks. The framework for development of the Barents Shelf initiated in the Palaeoproterozoic with the onset of the Svecofennian orogeny (Nironen, 1997; Korja et al., 2006). Subsequent orogenic deformation of the Timanian (Baikalian) occurred in the Middle to Late Vendian (Ediacaran), in which Neoproterozoic magmato-sedimentary assemblages were telescoped and accreted against the northeastern margin of Baltica (Roberts & Siedlecka, 2002). Timanian terrains are known to exist under parts of the Barents Shelf comprising NW–SE fabrics, however, their full extent and

significance in the Arctic is still poorly constrained (Olovyanishnikov et al., 2000; Roberts & Siedlecka, 2002; Pease and Scott, 2009). The structural grain of the Hammerfest basin, especially the basin–platform delineating TFFC is suggested to reflect a Caledonian grain. Separation of the Hammerfest Basin from the Finnmark Platform is envisaged to have taken place in the Late Carboniferous. (Gabrielsen, 1984; Gabrielsen et al., 2010; Gabrielsen & Færseth, 1988; 1989).

The Caledonian orogeny resulted in closure of the Iapetus Ocean during collision of Laurentia, Baltica and Avalonia and created a wide suture zone trending NE–SW in northern Norway which swings towards a N–S trend further north in Svalbard (Ritzmann & Faleide, 2007; Faleide et al., 2008; Gernigon & Brønner, 2012). In the Svalbard Archipelago, the metamorphic basement is dominated by the Devonian aged Scandian phase of the Caledonian orogeny (McKerrow et al., 2000), as well as earlier tectonic events, e.g., the Svecofennian and Timanian orogenies (Nironen, 1997; Ritzman & Faleide, 2007; Faleide et al., 2008; Marelli et al., 2010; Andresen et al., 2014). There are also older units preserved on Svalbard which are comprised of the Precambrian to Early Palaeozoic lower and partly middle Hecla Hoek series (Harland et al., 1966; Ohta, 1982).

Following the Caledonian orogeny, uplifted areas were subject to sub-aerial erosion, with deposition of Old Red Sandstone taking place within a fan shaped array of post orogenic extensional collapse basins (Faleide et al., 1993, 2008; Braathen et al., 2000; Braathen et al., 2004; Ritzmann & Faleide, 2007; Souche et al., 2012; Braathen et al., in press), including the large basin in northern Spitsbergen (Braathen et al., submitted). In Svalbard, this was followed by (Piepjohn, 2000; McCann, 2000) or interlinked with (Braathen et al., submitted) a phase of east–west crustal shortening during the latest Devonian to earliest Carboniferous Svalbardian-Ellesmerian deformation event.

In the Middle–Late Carboniferous, narrow rift grabens formed, again reactivating Caledonian lineaments, e.g., the Billefjorden Fault Zone (Johannessen & Steel, 1992; Braathen et al., 2011; Maher & Braathen, 2011). These half-grabens were filled by a mixture of siliciclastic and evaporite deposits of the Gipsdalen Group. Extension also took place in the SW Barents Shelf in the Carboniferous (Berglund et al., 1986; Faleide et al., 2008) where rift basins, including the Hammerfest Basin, accommodated continental siliciclastic sediments.

Extensional activity continued through the Late Carboniferous and into the Early Permian, (Høy & Lundschie, 2011; Faleide et al., 2017), and is marked by the deposition of the upper part of the Gipsdalen Group representing a transition from continental red bed to carbonate basin facies to more open water shallow marine carbonates and evaporates consistent with semi-arid to arid climates (Cutbill and Challinor, 1965; Steel & Worsley, 1984; Dallmann, 1999; Worsley et al., 2001). The Permian Bjarmeland Group records a shift to temperate cool water environments with deposition dominated by carbonates (Stemmerik & Worsley, 1995; Worsley et al., 2001). A renewed

pulse of extension occurred in the western Barents shelf during the Middle Permian (Faleide et al., 2017).

The Late Carboniferous to Permian also marked the onset of the Uralian orogeny where the Siberian Plate moved southwards accreting Kazakhstan and colliding with the nearly assembled Pangaeian Supercontinent (Rickard & Belbin, 1980; Ziegler, 1988; Gee et al., 2006; Brown et al., 2006; Puchkov, 2009; Pease, 2011). The entire Barents Shelf experienced a regional deep seated response to the Uralian orogeny with increased subsidence rates especially in the east (Worsley, 2008; Gac et al., 2014; Drachev, 2016). With time, the orogeny provided a prominent sediment source and large deltaic systems prograded from the southeast, and across the Barents Shelf from the Late Permian and onwards (Riis et al., 2008; Glørstad-Clark et al., 2010; Høy & Lundschieen, 2011; Klausen et al., 2016). Structurally-confined basins that were initially fault-controlled experienced rapid infill and subsidence without much movement along associated faults (Gudlaugsson et al., 1998).

The western parts of the Barents Sea experienced another phase of extension in the Late Permian to Early Triassic (Gudlaugsson et al., 1998; Serck et al., 2017) as evident by growth packages in the Bjørnøya Basin, Fingerdjupet Sub-basin and southern Stappen High (Glørstad-Clark et al., 2011; Faleide et al., 2015; Kamp, 2016). Furthermore, a growth sequence has been recognised in the Bjørnøya Basin along the Leirdjupet Fault Complex and correlates to a similar package in the Fingerdjupet Sub-basin (Serck et al., 2017).

The Triassic in the Barents Shelf is described generally as a period of tectonic quiescence (Riis et al., 2008; Worsley, 2008; Glørstad-Clark et al., 2011; Høy & Lundschieen, 2011) with ongoing regional subsidence, however, local zones of fault activity persisted, e.g., the southern Loppa High Fault System (Gabrielsen, 1984; Indræver et al., 2016), the Hoop Graben (Mahajan et al., 2014), eastern Svalbard (Anell et al., 2013; Osmundsen et al., 2014), and segments of the TFFC (this thesis). During this time, the Hammerfest Basin likely formed part of a larger epicontinental depositional regime, although Berglund et al., (1986) identify it as a separate depocentre from the Early Triassic onwards. The latest Permian to Middle Triassic is dominated by deltaic sequences which prograded across the entire Barents shelf (Glørstad-Clark et al., 2010, 2011; Anell et al., 2014). The Ingøydjupet Subgroup was primarily sourced from the recently uplifted Uralides, but the Baltic Shield also contributed and is likely a more prominent source for the southern Hammerfest Basin including the Goliat area (Riis et al., 2008; Mørk, 2009). The Middle to Late Triassic (comprising of the Ingøydjupet and Storfjorden subgroups) saw the shelf edge prograde north-westwards as far as Svalbard (Anell et al., 2014) where analogue outcrops of the Barents shelf strata can be observed. On Svalbard, siliciclastic wedges also prograded from the west, and possibly the north. The overlying Realgrunnen Subgroup was deposited in the Late Triassic to Middle Jurassic. It is most completely developed in the south-western Barents Sea where it consists of mature sandstones deposited in coastal plain and

deltaic to shallow marine environments (Olaussen et al., 1984; Berglund et al., 1986; Worsley, 2008; Henriksen et al., 2011; Rismyhr, 2012; Klausen et al., 2017). Dominant lithologies comprise sandstones and shales with subordinate conglomerates, carbonates and coals. The Realgrunnen and Ingøydjupet (underlying) subgroups host the target reservoirs in the Goliat field. Only the lower part of the Realgrunnen Subgroup is preserved in the Goliat field; the Norian Fruholmen Formation and the lower part of the Norian? to Rhaetian Tubåen Formation. Together these formations define a 67–118 m thick unit and form the upper Goliat reservoir. Here, the subgroup is interpreted as a fluvial dominated delta complex where reservoir units comprise of fluvial channels, tidally influenced channels, overbank lobes, crevasse splays and alluvial fan deposits all inter-bedded with silty mudstones (Leutscher & Nerby, 2011; Rismyhr, 2012; Leutscher, 2013). Palaeodrainage during the Rhaetian to Hettangian, i.e., during deposition of the Tubåen Formation is envisaged to have followed a general course from east to west within the central axis of the Hammerfest Basin, while in the Goliat field, a more northwest to west-northwest paleo-direction is evident.

On Svalbard, the targeted reservoir section envisaged for the Longyearbyen CO₂ Lab (article IV) belongs to the Upper Triassic to Middle Jurassic Kapp Toscana Group and consist of shales and sandstones of the De Geerdalen Formation (N/G approx. 0.3) and the overlying more sandy Wilhelmøya Subgroup (i.e., the Knorringfjellet Formation; Worsley, 1973, 2008; Knarud, 1980; Mørk et al., 1982; Harland & Geddes, 1997; Mørk & Worsley, 2006; Mørk, 2013). These units are interpreted to have been deposited in near-shore, paralic environments.

A regional phase of Mesozoic rifting took place in the Mid/Late Jurassic to Cretaceous (Gabrielsen et al., 1990; Faleide et al., 1993) and extended from the Rockall Trough west of Ireland to the Barents shelf. This event was closely affiliated with rifting in the North Atlantic and Amerasia basins. Primary fault complexes such as the TFFC developed large accommodation zones adjacent to active segments. A large domal structure formed in the hanging wall of the TFFC as it moved which Gabrielsen et al. (1990) describe as a roll-over structure (See article I). The central dome in the Hammerfest Basin is also believed to have developed at this time (Gabrielsen et al., 1990). This main phase of fault activity is associated with the deposition of shales, siltstones and sandstones throughout the Barents shelf which comprise the Adventdalen Group. Dark marine mudstones that dominate this group form important hydrocarbon source rocks including the Hekkingen Formation. In the Goliat field the Realgrunnen Subgroup reservoir is sealed by middle to upper Jurassic shales of the Hekkingen Formation and Cretaceous shales of the Nordvestbanken Group (Olaussen et al., 2010; Leutscher & Nerby, 2011). On Svalbard, the potential CO₂ reservoir is effectively sealed by the time equivalent 450 m-thick, shale-dominated succession belonging to the Middle Jurassic – Lower Cretaceous Agardhfjellet Formation (Koevoets et al., 2016) and the Lower Cretaceous Rurikfjellet Formation (Dypvik et al., 1991).

In the SW Barents shelf, structural elements including the TFFC, the Asterias Fault Complex and the Ringvassøya–Loppa Fault Complex were subject to inversion during the Barremian (Gabrielsen & Færseth, 1989; Indrevær et al., 2016) and may have comprised of a sinistral oblique component (Berglund et al., 1986; Sund, 1984; Gabrielsen and Færseth, 1989; Gabrielsen et al., 1990). A common driving force, possibly the uplift of the Loppa High during the Hauterivian–Barremian, has been suggested where compressive stresses are localised along the margins of the highs due to space accommodation problems (Indrevær et al., 2016). Following the inversion phase, subsidence resumed in the SW Barents shelf, but now thermal subsidence prevailed with only minor reactivations of fault systems continuing into the Early Cenozoic.

On Svalbard, the transition between the offshore to lower shoreface Hauterivian Rurikfjellet Formation and the fluvial to deltaic deposits of the Barremian to Aptian Helvetiafjellet Formation (Gjelberg and Steel, 1995; Midtkandal et al., 2007, Grundvåg & Olaussen, 2017) is marked by an erosional unconformity related to crustal up-doming driven by the HALIP event (Maher, 2001; Nejbirt et al., 2011; Minakov et al., 2012; Corfu et al., 2013; Senger et al., 2014; Polteau et al., 2016). Mafic igneous intrusions (approx. 122.2–124.5 Ma) associated with the HALIP played an important role in terms of thermal, localised deposition and perhaps compartmentalisation of the Mesozoic sedimentary succession (Corfu et al., 2013; Senger et al., 2013; 2014; Brekke et al., 2014).

During the Late Cretaceous, sea level transgression resulted in shelf starvation. This coupled with localised uplift are the conditions in which the Nygrunnen Group (Nøttvedt et al., 1993) was deposited. In the Hammerfest Basin, these deposits consist of condensed calcareous units and claystones that reflect marine, bathyal to deep/shallow shelf environments (Nøttvedt et al., 1993; Mørk, 1999). The top of the group is marked by a regional seismic reflector representing an unconformity, possibly related to phases of Late Cretaceous uplift and delineates the Cretaceous–Cenozoic boundary (Gabrielsen et al., 1990; Worsley, 2008; Ostanin et al., 2013). The entire Barents Shelf was transgressed in the Mid-Paleocene and a uniform basin of outer sublittoral to deep shelf was established. Accordingly, the Palaeocene and Eocene Hammerfest Basin deposition (Knutsen & Vorren, 1991; Vorren et al., 1991) is dominated by grey claystones of the Torsk Formation. The Eocene was punctuated by the opening of the Atlantic and major sedimentation in the western Barents Shelf whereas central and eastern regions experienced uplift.

In the latest Cretaceous, a dextral transform fault zone known as the De Geer Zone (i.e., the palaeo-Hornsund Fault Zone) developed between Greenland and the western Barents Sea (Talwani & Eldholm, 1977; Gaina et al., 2009). Initial stages of break-up and seafloor spreading were accompanied by a phase of Palaeogene transpression, which led to the development of the Western Spitsbergen Fold-and-Thrust Belt (WSFTB; Andresen et al., 1992; Braathen & Bergh, 1995; Bergh et al., 1997; Braathen et al., 1999; Leever et al., 2011). The WSFTB is characterised by a western thick-

skinned province where structures are basement-involved, and an eastern thin-skinned fold-thrust belt with several detachment levels developed along weak evaporite and shale intervals. Crustal flexure induced by orogenic loading of the WSFTB created the Palaeogene Central Spitsbergen Basin (CSB), a foreland basin that accommodated sediments from the uplifted western hinterland which comprises of Cenozoic marine to continental facies (Steel & Worsley, 1984; Braathen et al., 1999; Helland-Hansen, 2010; Anell et al., 2014).

Sea floor spreading in the Norwegian to Greenland Sea in the Early Eocene migrated northwards in the Mid-Eocene (Faleide et al., 2008). Subsidence in the western Barents Shelf, including the Hammerfest Basin continued until approximately the Mid-Eocene (Knutsen & Vorren, 1991), while the east and north experienced uplift. This is reflected in the fact that much of the Quaternary is missing on the Finnmark Platform. On the Barents Shelf the Nordland Group (Pavlidis & Polyakova, 1997; Sarnthein et al., 2003) consists of sandstones and claystones with the sand content increasing upwards. Cobbles and boulders of quartzite, granite and different metamorphic rocks occur with clay in the upper parts of the group. Depositional age of the group is envisaged as Late Pliocene to Pleistocene/Holocene in the Hammerfest Basin.

Neogene uplift and erosion (Våagnes et al., 1998; Laberg et al., 2012) in the western Barents Shelf was largely controlled by northern hemisphere glaciations and resulted in significant erosion and deposition of major submarine fans along the western margin of the shelf (Eidvin et al., 1998). In other parts of the shelf Neogene deposits are rare due to uplift. Three major glacial events are noted by Knies et al. (2009), at 3.6-2.1 ma, 2.4-1.0 ma and 1.0 Ma onwards, after which a shift to 100,000 year cycles was established.

4. DATA AND METHODS

The articles that form this body of work present results from a variety of geological methods including offshore seismic and well log investigations, onshore outcrop studies and a range of laboratory testing carried out on outcrop and core samples. A concise description of the data sets and methods used are given in this section and summarised in Figs. 15 and 16. More in-depth descriptions can be found in the respective articles.

4.1 OFFSHORE INVESTIGATIONS

4.1.1 Seismic data

Seismic investigations of the Goliat field (articles I and II) were conducted using the EN0901 multi-azimuth (MAZ) survey made available by Eni Norge. The survey consists of a pre-stack, depth migrated dataset covering an area of approximately 207 km² and providing imagery down to depths of approximately 4000 m. The survey was shot at 3 azimuths, 7°, 67° and 127° with a shot point interval of 12.5 m, and a streamer length of 4000 m. Two-dimensional regional transects sourced from the Norwegian Petroleum Directorate were examined in conjunction with the 3D survey and provide subsurface imagery from 8000 m depth.

Seismic workflows were carried out primarily utilising the Petrel E&P Software Platform and focused on identifying prominent seismic reflectors and high resolution mapping of the fault populations affecting the Goliat region (article I). Article II concentrated on depositional geometries within the Realgrunnen Subgroup and high resolution mapping of faults affecting the interval. Displacement analysis was conducted on fault polygons and seismic horizons utilising Badley's T7 software. AVA (Amplitude versus Angle) attribute analysis was conducted over a 10 ms window below the top Realgrunnen Subgroup using HampsonRussel AVO analysis & Modelling software in order to highlight the gross sand prone areas and depositional geometries of the Realgrunnen Subgroup.

4.1.2 Well logs

Both the Goliat studies in articles I and II utilised six exploration wells, 7122/7-1, 7122/7-2, 7122/7-3, 7122/7-4 S, 7122/7-5 (5A deviated within the Snadd Formation) and 7122/7-6. The dataset for the sedimentological study comprised wireline logs and 235 m of slabbed core from the six exploration wells. Sedimentological studies focused on grouping the deposits into facies associations reflecting discrete depositional elements or environments. Composite wireline log suites, which included gamma ray, neutron-porosity and bulk density logs, were used for a more

simplistic recognition and interpretation of lithologies, boundaries and relating non-cored intervals to the sedimentary facies framework. A well-to-well correlation was performed using facies architecture and stacking patterns, and guided by in-house (Eni Norge) palynological data.

Figure 15 Summary of data and methods used for describing the offshore studies (Goliat field) described herein.

4.2 ONSHORE INVESTIGATIONS

4.2.1 Discontinuity mapping

Articles III and IV feature outcrop based investigations which mapped out structural discontinuities including faults, fractures, joints, veins and deformation bands. Discontinuity terminology is based on Schultz and Fossen (2008), unless otherwise stated. These investigations took place in central Spitsbergen (Deltaneset locality) and eastern Svalbard (SW Edgeøya). Fault architecture was mapped in terms of breccia series and gouge thickness/composition after Braathen et al. (2004). Physical characteristics and frequency distribution of natural discontinuities were established by conducting scan lines along beds, i.e., the 1-D line intersection method after Singhal and Gupta (2010). Measurements recorded include azimuth, dip and spatial distribution. Additional

fracture features, continuity (whether the fracture is bed-confined or continues through multiple beds or the entire section, i.e., through-going) and mineral infill, if present, where recorded. Fractures and joints were also mapped using oriented and scaled plan view photographs, both taken in the field, and available online e.g., the Norsk Polarinstitut online maps (<http://toposvalbard.npolar.no/>).

4.2.2 Borehole and outcrop plug data

The Longyearbyen CO₂ reservoir study (article IV) also made use of core and wireline log data from three closely spaced wells in Adventdalen (DH4, DH5R, DH7A) and an additional well 7 km to the northwest (DH2), which fully penetrated the Wilhelmøya Subgroup. A suite of wireline logs from DH4, including natural gamma, caliper, velocity, and density, were used to further constrain the interpretations. A televIEWER was used for downhole fracture mapping (frequency only, as not orientated). Core logging was tied to the aforementioned outcrop logging (section 4.2.2).

4.2.3 Virtual outcrops

Outcrop investigations were complimented by the creation of virtual outcrop models which allowed remote measurements and large scale depositional and structural geometries to be discerned. Virtual outcrops were utilised in articles III and IV. Models were generated using structure-from-motion (Photogrammetry), i.e., the generation of 3D models from a series of highly overlapping geo-referenced photographs. Images were captured with a full-frame DSLR camera (Canon 6D) which contained an integrated GPS module, and a range of prime lenses (35, 50, 80 and 100 mm) depending on distance to outcrops. In addition, averaged GPS positions of recognisable outcrop features were noted in order to increase the spatial accuracy of the model. Photographs were taken at 21 megapixel resolution and in RAW format. The focal length of the lens was fixed to insure similar focus in all pictures. An approximate 80% overlap was maintained between each image in order to ensure the same features are covered by multiple images. Model generation was conducted using “Agisoft Photoscan” software which builds point clouds from similar matched features in multiple images. A 3D framework depicting outcrop geometry is produced upon which the original photograph textures can be draped. Models were imported into “Lime” software (developed by the Centre for Integrated Petroleum Research, Bergen) which facilitates remote outcrop measurements including distance and orientated surface azimuth/dip calculated from 3 point selections. The 3D models allow accurate measurements to be recorded whereas panoramic imagery is subject to barrel distortion (i.e., measurements from image extremities will be underestimated).

4.2.4 Additional Methods

Article IV built on an extensive reservoir appraisal campaign focused on discerning the reservoir properties of the Late Triassic Succession in Central Spitsbergen. As such, article IV made use of injection and monitoring test data to obtain permeability information and to test lateral communication between wells in addition to a multitude of laboratory investigations, including high resolution X-ray computed tomography (micro-CT), Mercury Intrusion Porosimetry (MIP) and He-Porosimetry for pore network characterisation. Clay gouge from 5 extensional faults affecting the reservoir unit was sampled from outcrops and subject to X-Ray diffraction (XRD) mineralogical composition analyses. Borehole plug analyses were also conducted in order to measure porosities and permeabilities of the reservoir interval. Additional permeability measurements were collected using a Miniperm tool.

Figure 16 Summary of data and methods used for describing the onshore studies (Deltanaset area, outside Longyearbyen and SW Edgeøya, East Svalbard)

5. ARTICLE SUMMARIES

5.1 ARTICLE I

Mulrooney, M.J., Leutscher, J., Braathen, A., 2017. A 3D structural analysis of the Goliat field, Barents Sea, Norway. *Mar. Pet. Geol.* doi:10.1016/j.marpetgeo.2017.05.038

The first article in this collection comprises a detailed structural description of the Goliat field and provides the first published account of the area utilising a 3D seismic survey. Moreover, the seismic data set used, the EN0901, is a multi-azimuth (MAZ) survey which reconciled several resolution problems encountered with vintage single azimuth surveys of the area. Poor resolution for earlier data related to a dense configuration of multiple trending fault populations, the hard nature of the substrate, and the presence of complex glacial morphology.

The article identifies three unique fault populations, two of which exhibit orientations parallel/sub-parallel to segments of the Troms-Finnmark Fault Complex (TFFC), the Goliat–Tornerose (GT; 030–210°) and the Alke–Goliat (AG; 085–265°) populations. These populations are shown to consist of basement involved master faults and a large number of subsidiary faults. The GT1 master fault delineates the Hammerfest Basin from the Finnmark Platform. A third population, the Hammerfest Regional (HR; 105–285°) is identified and possibly represents the offshore extension of the Trollfjord-Komagelv Fault Zone (as suggested by Berglund et al., 1986; Gabrielsen & Færseth, 1989; Gernigon & Bronner, 2012).

The article describes the geometry of the Goliat field which exploits an anticlinal trapping structure that abuts the northern segment of the TFFC. The highest point of the Goliat anticline, informally named the Goliat crest, coincides with a displacement minimum on the GT1 segment of the TFFC. An additional NW–SE striking minor fault population which is referred to as, the Goliat Central (GC) fault population affects the crest of the Goliat anticline and partitions the occurrence of AG and GT subsidiary faults which exhibit curvilinear traces to the west of the Goliat area.

Interpretation of regional 2D seismic transects reveals the TFFC exists as two discrete, generally planar segments at depth, the GT1P and the GT1M, contrary to previous works which describe listric/ramp-flat geometries (e.g., Ahmed, 2012; Indrevær et al., 2015). The article outlines that differential subsidence on these two vertical segments is primarily responsible for generation of the Goliat anticline. Significant post-kinematic subsidence on the GT1P segment formed the prominent western limb of the Goliat anticline, whereas the eastern limb is suggested to form by hanging wall roll-over established by moderate ramp-flat-ramp and listric geometries on the GT1M fault segment as deduced by fault attribute analysis. The area of the GT1M that exhibits a ramp-flat-ramp geometry is likely the result of dip linkage of two vertical fault segments.

Sedimentary package geometries suggest movement on the GT1P ceased in the Late Palaeozoic, however, possible minor reactivations persisted as suggested by downward propagation of subsidiary faults that coalesce towards the upper tip-line of the structure. The GT1M became active in the late-most Palaeozoic and controlled deposition of the Mesozoic succession.

Depth structure maps along with fault attribute and displacement analysis reveal large scale fault jogs on the GT1 master fault, as well as displacement minima which identify the sites of 3 former segments that interacted and linked during the Late Permian to Early Triassic. The AG1 master fault is shown to bifurcate up section from a through-going structure at depth into a series of en-echelon segments at the level of the upper Triassic. The orientation of the through-going structure is likely inherited from a reactivated basement structure, whereas the overlying en-echelon segments and the captured adjacent subsidiary faults may reflect the stress regime contemporary to their formation suggesting the tensile stress field during the main phase of Mesozoic rifting is oblique to the AG1.

Seismic interpretation of the Late Jurassic and Early Cretaceous syn-kinematic units in the hanging wall of the GT1 are consistent with geometries indicative of a breached fault propagation monocline or a "forced fold"; these observations include very steep dipping syn-kinematic units, lenticular syn-rift basins, along-strike geometric variation, over-turned faults and relatively undeformed footwall strata.

Folds affecting the Late Jurassic to Early Cretaceous succession display positive relief, i.e., the crests are raised above the regional elevation and are accordingly interpreted as inversion structures. Folding generally overlies reactivated extensional faults or footwall short-cut thrust faults suggesting they are the result of fault propagation.

Polygonal faults affecting the Early Cretaceous to Early Cenozoic successions are identified in variance depth slices. These faults show a low level of organisation with preference for orientations similar to that of the GC and the AG fault populations.

The article concludes with a summary of fault activity for the largest faults representing each fault population in the Goliat area including the subsidiary faults. Timings are related to the main phases of tectonism that affected the Barents shelf during the Phanerozoic.

5.2 ARTICLE II

Mulrooney, M.J., Rismyhr, R., Yenwongfai, H. D., Leutscher, J., Olaussen, S., Braathen, A., (In press). Impacts of small-scale faults on continental to coastal plain deposition: Evidence from the Realgrunnen Subgroup in the Goliat field, Southwest Barents Sea, Norway.

Building on the tectonostratigraphic framework of the Goliat area described in article I

(Mulrooney et al., 2017), the second article in this collection focuses on the depositional environment of the Realgrunnen Subgroup (Fruholmen and Tubåen formations) which forms the uppermost reservoir interval in the Goliat field. Key to this article is the identification of a previously undocumented Norian to Rhaetian tectonic event affecting the southern Hammerfest Basin. This finding is evident from a synthesis of information from sedimentary cores and petrophysical data, seismic interpretation, fault displacement analysis, stratigraphic thickness variations adjacent to faults, and Amplitude versus Angle (AVA) attribute analysis of the EN0901 MAZ survey. The article presents detailed palaeogeographic reconstructions for the Fruholmen and Tubåen formations with emphasis on the role of tectonism on the distribution of syn-kinematic facies in continental and coastal plain environments.

Core based facies analysis suggests the Fruholmen Formation was deposited in an upper delta plain environment which was characterised by tidal and river-dominated distributary channels. Towards the west, this environment transitioned towards a lower delta plain and a prodelta. Evidence for fault activity during deposition of the Fruholmen Formation includes individual channel features and ribbons of gross sand prone areas identified by AVA analysis (interpreted as rivers/distributary channels) that are aligned with fault traces (i.e., axial drainage). Furthermore, a lateral variation in channel deposit densities in the upper part of the Fruholmen Formation may reflect some differential movements already at this time.

The overlying Tubåen Formation is coarser grained and represents a widespread braid plain with localised alluvial fans. The transition from the Fruholmen Formation into the Tubåen Formation is signified by a sequence boundary representing a regressive event caused by a drop in relative sea level, and/or an increase in sediment supply from the source. A general correlation of fault displacement maxima, footwall thickness minima and hanging wall thickness maxima are suggested to represent syn-kinematic footwall erosion, and preferential deposition in isolated sub-basins along a segmented, immature fault system. Minor evidence of growth wedging is evident from seismic profiles. Alluvial fan deposits in wells coincide with large sand prone bodies identified in the AVA attribute map, some of which fan out from relay zones on large active faults. These features are interpreted to have been derived from semi-circular drainage catchments that develop in the footwalls of active fault segments. Conceptual depositional models for key sites of sediment and fault interaction are displayed in a series of block diagrams.

Four sequence boundaries were identified in the Goliat well cores and referred to as SB1, SB2, SB3 and SB4 in article II. The article concludes with a discussion on the regional significance of the sequence boundaries, some of which can be correlated throughout the Circum-arctic region, including the Sverdrup Basin, onshore Svalbard, the Barents shelf area and eastern Siberia. The article discusses two synchronous tectonic episodes possibly responsible for the Late Triassic faulting

event, 1) Rejuvenation of Fennoscandia to the south and/or vertical movements related to the early Cimmerian tectonic phase in northern Europe and 2) a tectonic episode which could correlate to movement on the Novaya Zemlya Segment of the Pai-Khoi–Novaya Zemlya–South Taimyr fold belt to the east. The article also discusses whether sediment loading from the aforementioned rejuvenation of Fennoscandia could have driven, or contributed to the fault event.

5.3 ARTICLE III

Ogata, K., Mulrooney, M.J., Braathen, A., Maher, H., and Balsamo, F., (In press.). Architecture, deformation style and petrophysical properties of a Late Triassic growth fault system in southern Edgeøya (East Svalbard).

Article III presents a multi-scaled investigation of a well exposed growth fault system which outcrops on southern Edgeøya, eastern Svalbard. The fault system records Late Triassic syn-kinematic sedimentation that took place in the pro-delta of an overall north-westward prograding deltaic system. Faults have been studied on the micro, meso and macro scale utilising thin sections, measurements from outcrop, and measurements from a virtual outcrop model spanning an 8 km long cliff section, respectively.

Three distinct hierarchical orders of growth faults were recognised based on fault plane geometry, deformation mechanisms and stratigraphic locations of their respective detachment interval. In general, faults show SW to SE dip directions which implies the system had an overall landward (counter-regional) dip with respect to the north to north-westward progradation direction inferred for the associated delta system.

The three orders of faulting formed as part of a cohesive system and show complex spatial and temporal interference and progressive reworking of inherited structures. The 2nd and 3rd order faults exhibit listric geometries and controlled the lower (silty-shaly) and upper (sandy) growth packages, respectively. Planar, 1st order faults influenced the accommodation in both the growth stages. Fault slip was apparently accommodated by a polyphase deformation style where the different fault architectural elements record a transition from fluidal-hydro-plastic to ductile-brittle deformation, from the smaller (3rd order) to the larger-scale (1st order) end-member faults, respectively. This transition records a shift from strain diffusion to strain localisation towards fault cores which can be observed at the meso- to micro-scale (<1mm). In addition, a meso-scale variation in petrophysical parameters of the litho-structural facies across and along the fault envelope is observed where bulk porosity, density, pore size and micro-crack intensity varies in accordance to the intensity of deformation and reworking of inherited structural fabrics.

In fault hanging walls where sandstones have accumulated, deformation mechanisms vary from liquidisation processes and independent particulate flow at low confining pressures causing shear-dilation to diffused disaggregation and compaction banding with localised incipient cataclasis, eventually leading to localised shear and extensional fracturing. In the footwall where silt and shalestone facies are more common, deformation mechanisms vary from hydro-plastic deformation and disaggregation-phyllonitic tabular shear zones, to diffuse brittle-ductile shearing and later localised shear failure and extensional fracturing. This evolving rheological-mechanical behavior is credited to the transition of deformation mechanisms along with burial and shallow diagenesis. The micro-scale, fabric-related structural anisotropy generated during the early deformation phases appears to control the bulk pore size distribution promoting the development of a bimodal porosity system characterised by a background primary (sedimentary) matrix pore network that has been overprinted by a secondary (tectonic) pore network produced by micro-cracks which have localised along pre-existing micro-fabric discontinuities. The cementation patterns also confirm the systematic occurrence of pervasive Fe-sulfide impregnation halos in the sheared hanging wall sandstones along fault cores that likely formed due to the enhanced dilation-related fluid circulation in the early to intermediate phases of growth faulting.

The 1st order planar faults are through-going and cut down into the underburden. The driving mechanism responsible for these faults is envisaged to be tectonic reactivation of Palaeozoic structures in a far field stress regime of the paleo-Uralide orogeny as well as differential compaction associated with increased sand-prone sedimentary input in a fine-grained, water-saturated, low-accommodation depositional environment. These planar faults are inferred to be responsible for the creation of topographic relief atop the Botnheia Formation (the underburden) during the deposition of the Tchemakfjellet Formation, which established localised depozones and consequent differential compaction, thus preconditioning the formation of the 3rd and then 2nd order faults. Furthermore, listric faults appear to be localised above fault tip monoclines that deform the top of the underburden.

Superimposed on this large-scale tectonic driving force, small-scale causative factors partially responsible for 2nd and 3rd order faulting seem to be related to mechanical-rheological instabilities related to the localised shallow diagenesis and liquidisation fronts.

5.4 ARTICLE IV

Mulrooney, M.J., Van Stappen, J., Cnudde, V., Senger, K., Rismyhr, B., Braathen, A., Olaussen, S., Mørk, M.B.E., Ogata, K., (In review). Fluid flow properties of a potential unconventional CO₂ storage unit in central Spitsbergen: the Upper Triassic to Middle Jurassic Wilhelmøya Subgroup.

The fourth article in this collection investigates the impacts of sub-seismic heterogeneities on the Late Triassic to Early Jurassic Wilhelmøya Subgroup (northern equivalent of the Realgrunnen Subgroup), which has been identified as one of the most prospective reservoir units in a carbon capture and storage pilot study on the Norwegian Arctic archipelago of Svalbard. The unit is severely under-pressured (approx. 45 bar), and displays characteristics of an unconventional, tight and compartmentalised reservoir.

A main premise of the article is that water injection and monitoring tests in adjacent wells attests to the presence of barriers or severe flow restrictions. Despite this, water injection tests also show hydraulic fractures have been induced during injection (Larsen, 2010, 2012; Boholi et al., 2014). These fractures will enhance an extensive existing natural fracture network observed in core and outcrops that will contribute both to fluid injectivity and storage potential (Ogata et al., 2012, 2014). Two main natural fracture sets have been identified, an ENE–WSW trending principal fracture set and a NNW–SSE trending subordinate fracture set. Preferred orientations and frequency of fracturing varies with contrasting mechanical properties. Reservoir sandstones will act as preferential vertical fluid migration pathways, while shale-dominated successions will preferentially form lateral fluid migration pathways. Micro-CT analysis combined with Mercury Intrusion Porosimetry (Van Stappen et al., 2014) provides reservoir information at the pore scale and allows for a quantification of the fracture apertures at the micro-scale and their contribution to porosity.

The discussion in article IV considers three sub-seismic reservoir heterogeneities that can contribute towards the flow restrictions observed in the aforementioned well tests. The article proposes that an extensional fault system comprised of relatively small (less than 10 m displacement) but numerous segments is located between the investigated wells, and oriented ENE–WSW in accordance to fault systems identified in outcrop. Dolerite dykes and stratigraphic closures may also contribute to lateral pressure compartmentalisation.

Finally, the article divides the potential reservoir unit into three zones. Injection and storage potential of the reservoir zones are evaluated based on net to gross (NTG) ratios and published reservoir property data (i.e., porosity and permeability) from (Farokhpour et al., 2010; Magnabosco et al., 2014). Variation in properties are caused by a combination of multistage depositional and diagenetic processes including quartz and calcite cementation, authigenic clay growth and secondary porosity dissolution (Mørk, 2013). Porosities of 8.7–19.6% and permeabilities of 0–1.8mD are measured from core plugs (Farokhpour et al., 2010).

6. DISCUSSION

The four articles contained herein investigate Late Triassic faulting in the Barents Shelf and the controls of syn- and post-kinematic faulting on reservoir properties by 1) controlling the distribution of sand-prone facies and by 2) changing the fluid flow properties of the pristine country rock. The articles comprise case studies from the Barents Shelf combining onshore and offshore investigations from Svalbard and the Hammerfest Basin, respectively. The block diagram in Fig. 17 summarises the geological processes explored in the articles.

Figure 17 Conceptual block diagram showing primary topics dealt with in this body of work. 1) Basement involved fault systems, 2) en-echelon fault segments showing progressive linkage and existing as a through-going fault at depth (3), 4) dip-linkage of faults establishing vertical fault kinks that set up roll-over structures, 5) forced folds in the overburden of upward propagating faults, 6) damage zones surrounding faults affecting reservoir properties, 7) igneous bodies and corresponding aureoles record information on stress conditions during emplacement, 8) gravity driven deltaic collapse faults with diapirism of over-pressured shales, 9) corresponding down-slope toe thrust system, 10) relay zones in immature fault systems acting as point sources for sediments to enter the hanging wall, 11) axial drainage where rivers migrate towards fault induced tilt and 12) footwall uplift driving erosion and creating drainage areas.

In this section, the main study results are discussed in context of existing work. The topics explored include, i) how investigations of contrasting scales provide a more complete understanding of fault processes, ii) how observations of fault geometries from a MAZ survey herein reflect fault growth models, iii) the different influences of faults on reservoirs with respect to variables such as timing and mechanisms of deformation, and finally iv) the regional significance of fault events during the Late Triassic evolution of the Barents Shelf.

6.1 RESOLUTION: BRIDGING THE GAP

The articles herein, use various geological datasets including seismic surveys, virtual outcrop models, core and wireline data, aerial photography, traditional outcrop measurements and micro-structural data sets to investigate outcropping and subsurface fault systems. These data sets span a wide range of scales, the vertical and horizontal resolutions of which are plotted in Fig. 18.

Figure 18 Vertical and horizontal resolution spread of the datasets used in this study (Inspired by Matt Hall, Agile).

Article I and II utilise a high resolution multi-azimuth survey which is controlled by well core and petrophysical data. The apparent gap in data of intermediate resolutions between these two datasets requires inferences, which add a significant source of error. The resolution plot in Fig. 18 shows the importance that outcrop geology (articles III and IV) plays in structural and sedimentological investigations owing to the wide spread of spatial resolution it can encapsulate, in sharp contrast to core and wireline data, which provide a good range of vertical resolution, but

laterally are very limited. Aerial photography also accommodates some of these intermediate scales and has been used to identify regional lineaments in article III. Aerial photography is, however, a mostly 2D data set and has very restricted vertical resolution. On the finer end of the spectrum, microscopic analyses, e.g., articles III and IV, are required for discerning reservoir properties, e.g., porosity, permeability and fracture aperture. Some fault elements studied in article III in thin section are fractal (scale invariant) as they show similarities to larger structures identified in outcrop.

The advance in virtual outcrop geology in recent years (e.g., Buckley et al., 2016) has played an important role in integrating outcrop scale features with that of seismic. Virtual outcrop geology has been used in articles III and IV (also see appendix, section 9.1 and 9.3) to identify large-scale structural discontinuities and sedimentological bodies, and to measure orientations in otherwise inaccessible localities. Automated fracture mapping using virtual outcrops is detailed in the appendix (section 9.6). The virtual outcrop model of southwest Edgeøya (article III), Svalbard has been used by Anell et al. (2016) as a basis for generating synthetic seismic which affectively shows that the faults, lenticular shaped sandstone bodies and horizontally orientated igneous bodies (sills) at that locality would indeed be identifiable in high resolution seismic studies.

6.2 FAULT GROWTH

Fault growth as outlined in section 2, is a complex three-dimensional phenomenon best described with high resolution 3D seismic data. The availability of the EN0901 MAZ survey herein (articles I and II) provides fantastic insight into a complicated zone of brittle crustal deformation located within a major bend on the basin-platform delineating TFFC. Structures imaged by the survey include two master segments of the TFFC, a master segment of the Hammerfest Regional Fault System, a large domal structure seemingly genetically linked to fault movement and many subsidiary faults which parallel master faults, and in cases laterally splay or vertically branch from the larger structures. Some key observations which may help inform current growth models (section 2) are outlined below, in particular regarding fault displacement to length ratios (D:L), inheritance of characteristics from reactivated basement structures and preservation of anomalies (i.e., displacement maxima/minima and undulations in attitude) after early fault segments link to form through-going structures.

A notable discrepancy is observed in the Goliat area between D:L ratios of the master faults (e.g., GT1, AG1, AG2, AG3 and HR1; article I) and the large population of subsidiary faults, with the latter exhibiting much smaller ratios. In other words, subsidiary faults have long trace lengths and low to moderate displacement values. This observation is in line with Meyer et al. (2002)'s description of fault growth where the hierarchy of faults is established quickly and early fault

propagation is dominated by rapid accrual of length. The master fault segments, some of which are basement involved (Gabrielsen, 1984; Indrevær et al., 2013; Indrevær and Bergh, 2014), show similar trace lengths, but have accrued far greater displacement which suggests strain has been localised. For the non basement involved faults, the rapid (instantaneous on geological timescales) establishment of fault length is credited by Meyer et al. (2002), Walsh et al. (2002) and Giba et al. (2012) to reflect the upward propagation of reactivated basement faults. This affiliation with basement faults is reflected in Fig. 4 of article I, where subsidiary faults affecting the Goliat anticline show spatial affinity to an underlying vertical segment of the TFFC, i.e., they appear to branch from the tip point of the deeper structure (possibly soft-linked) and have comparable strike directions. These observations support a fault growth model dominated by reactivation of older structures.

Another principal of fault growth models outlined in section 2 is that fault interactions, i.e., within relay zones when faults propagate and overlap (Peacock & Sanderson, 1991; Trudgill & Cartwright, 1994; Huggins et al., 1995; Childs et al., 1995, 1996, 2016; Kristensen et al., 2008) cause changes in local stress fields which can result in retardation of lateral growth, perturbation of fault strike, or instigate faults to splay/bifurcate (Kristensen et al., 2008). Several such local stress perturbations are observed within the Goliat field where subsidiary faults of the AG fault population can be seen (article I; Fig. 6) striking parallel to the AG1 master fault, before suddenly turning sharply towards the master fault, where they tip-out or intersect with it. The phenomenon is termed, “fault capture” herein, after Browning & Gudmundsson (2015). Moreover, the master fault is an amalgamation of a series of en-echelon fault segments in various stages of breach. At depth this fault exists as a single through-going structure. In this case, the en-echelon faulting and fault capture are credited to an extensional stress direction (σ_3) that is oblique to the AG1 master fault, which reactivates, none-the-less, and propagates up-section where en-echelon fault segments form at more preferential orientations to the ambient stress field. In this scenario, the GT1 master fault which is close to parallel to the en-echelon segment of AG1 is more preferentially orientated to accommodate extensional strain, and is the largest fault as a result. This observation also shows that normal faults can be reactivated by extensional stresses up to at least 30–35° oblique to the existing structure.

The model proposed for the evolution of the hanging wall geometries related to the GT1 segment of the TFFC (article I), where strata shows severe “normal drag”, support the interpretation that “drag” (e.g., Grasemann, 2005) is a fault propagation feature, and in this case the result of a fault propagation monocline which was subsequently breached by continued upward propagation of a blind fault.

The final discussion point on fault growth concerns the preservation of geometric signatures of former segments in through-going mature fault systems, i.e., displacement, strike and dip

anomalies (Mansfield and Cartwright, 1995; Lohr et al., 2008; Mulrooney, 2011). The preservation of displacement minima along a fault trace where fault breach and linkage has occurred has been well documented (e.g., Lohr et al., 2008). Moreover, the presence of lateral and vertical kinks in fault planes is also consistent with sites of linkage, and is easily identified in the master faults affecting the Goliat area, especially the GT1 segment of the TFFC. Some authors (Lohr et al., 2008; Mulrooney, 2011) have identified linkage sites by collating evidence from fault displacement profiles, and fault plane kinks and curvature. In article II, this approach was complimented by overlaying this type of data with strata thickness profiles immediately adjacent and parallel to fault planes. This approach uses thickness anomalies as a proxy for areas of increased deposition in hanging walls, and areas where erosion has taken place in footwalls. While these interpretations are not unique, where correlation occurs, i.e., where displacement maxima coincide with increased and decreased hanging wall and footwall thickness, respectively, the fault portion likely represents a former segment.

6.3 INFLUENCE OF FAULTS IN RESERVOIRS

A reoccurring topic in this body of work is that faults impact reservoir properties. Articles II and III demonstrate how syn-kinematic sediment distribution can be controlled by fault related morphology and can therefore impact the geometry and connectivity of reservoir quality sandstones.

Article II treats this concept on a regional scale, where fault related geomorphology is shown to source immature sediments which feed into an evolving continental to coastal rift basin via relay zones and across-fault scarps. Antecedent drainage is shown to be rerouted by the fault controlled geomorphology. Channelised features exhibit axial drainage in the hanging walls of faults which result in stacked and amalgamated channel facies. Article III describes sediments that have been rerouted and accommodated by individual growth basins controlled by thin-skinned listric faulting. The article displays how faulting can affectively capture sandy facies in a pro-delta environment otherwise typically dominated by finer grained facies.

Within reservoirs, petrophysical properties (including porosity and permeability) can locally be significantly perturbed by the presence of fault zones. The nature of fault zones, i.e., the individual fault elements encountered in an envelope impacted by the fault, depend heavily on the existing properties of the reservoir rock including country rock lithology, primary matrix porosity, water saturation, pore fluid pressure (which may reflect depth) and diagenesis.

Article III provides new information regarding the characteristics of damage zones within poorly lithified syn-kinematic sediments, in particular on the topic of the mixed (or mixing) zones, i.e., variably deformed, largely homogenised zones with lithological mixing down to the grain scale that develop mostly within hanging walls. The concept of mixed zones forming in the inner damage

zone of faults in poorly lithified sand-shale successions has been discussed before by Heynekamp et al. (1999) and Rawling and Goodwin (2006). Braathen et al. (2014) further investigated this phenomenon and reasoned these features form as a product of fluid mobilisation along the fault core, which along with differential compaction, induces weakening along the fault plane which facilitates creep and smear-shear band deformation. New insight into the formation of these features is discussed herein (article III), and relates to the temporal and causative nature of this deformation style. The observed transition from hydro-plastic strain diffusion to brittle strain localisation along faults is credited, in part, to early calcite cementation. In this scenario, sand grains cemented by calcite facilitate grain fracturing in parallel with calcite cement breakage rather than grain-contact sliding (in the earlier non-cemented hydro-plastic regime). In more shale dominated facies, early dewatering reflecting differential load by sand is a more likely causal factor of the transition. Moreover, article III suggests that high fluid pressure and liquefaction potentially associated with fluid escape or ground shaking (earthquakes) played a key role in the generation of this fault rock.

Article IV demonstrates more brittle fault zones which affect older lithified successions onshore Svalbard. These faults formed during evolution of the Palaeogene Western Spitsbergen Fold-and-Thrust-Belt (WSFTB). In this scenario, fault damage zones in the primarily heterolithic successions (see appendix, section 9.2) are more prone to fracturing and brecciating than deformation banding, which is more common in highly porous rocks (Fossen et al., 2007). Knowledge of the nature of the fault zones in the Wilhelmøya Subgroup onshore Svalbard will be used to enhance reservoir models for potential CO₂ injection, and builds on studies that characterised the natural fracture networks within the stratigraphic successions (Ogata et al., 2012, 2014) and igneous bodies emplaced in the target reservoir (Senger et al., 2014).

Furthermore, the Goliat case study from the Hammerfest Basin, (articles I and II) and the onshore Svalbard CO₂ reservoir study (article IV) show how faults can compartmentalise reservoirs by forming impermeable barriers to fluid flow. In the Goliat field, differential pressures and oil-water contacts are present across faults which affect the Goliat anticline (domal trapping structure; article I). In this case, the compartments are complex owing to the presence of 4 unique fault trends making horizontal drilling necessary for production (Johan Leutscher, pers. comm., 2 March 2017). Articles I and II provide the first published descriptions of the Goliat area (i.e., parts of the TFFC, the southern margin of the Hammerfest Basin, and parts of the Finnmark Platform) based on three-dimensional data. Before this, descriptions of the area were limited to 2D regional seismic data (e.g., Gabrielsen, 1984; Gabrielsen & Færseth, 1989; Faleide et al., 1993; Ahmed, 2012). The studies herein provide better insight into the temporal development of the Goliat anticline, and therefore can be used to refine the petroleum system, i.e., defining the relative timing of trap structure generation with

regard to the timings of hydrocarbon generation and migration. Furthermore, the recognition of an earlier Mesozoic rift event which influenced the type and dispersion of sediments (mapped out in article II) will inform future development of the field.

Compartmentalisation in the Late Triassic to Middle Jurassic succession onshore Svalbard (article IV) is a simpler case where uni-directional faults likely baffle communication between two appraisal wells. The baffles have been related to a minor stage of extensional faulting that strike ENE–WSW and formed during the late stages of the Western Spitsbergen Fold-and-Thrust Belt (WSFTB). The description of the Goliat field herein, specifically that of the Realgrunnen Subgroup, can form an analogue for the Late Triassic Wilhelmøya Subgroup on Svalbard given their approximate time and facies equivalence, and given the fact that seismic imagery onshore Svalbard is of poor quality (Bælum et al., 2012). Moreover, the fault architecture observed in outcrops onshore Svalbard can inform subsurface faults in the Goliat field, as they likely share similar characteristics due to similarities in the host rock, albeit the Subgroup was likely buried to greater depths onshore Svalbard than the equivalent offshore in the Hammerfest Basin.

6.4 LATE TRIASSIC TECTONISM IN THE BARENTS SEA

A commonality that links article III (investigations onshore Svalbard) and article II (offshore southwestern Barents Shelf), is the apparent importance of far field (intraplate) stresses acting on the Barents shelf in the Late Triassic, related in part, to the Uralian orogeny (Also see appendix, section 9.3). The Triassic in the Barents Shelf has generally been described as a period of tectonic quiescence (Riis et al., 2008; Worsley et al., 2008; Glørstad-Clark et al., 2011; Høy & Lundschieen, 2011) with ongoing regional subsidence. The observations, however, of extensional tectonics herein and in several recent publications suggests this description is an over-simplification.

Onshore Svalbard, in addition to the Carnian aged Triassic growth faults in SW Edgeøya described herein (article III; Anell et al., 2013; Osmundsen et al., 2014). Osmundsen et al., (2014) also described similar shallow-rooted normal faults at Skrukkefjellet and Klinkhamaren in NW Edgeøya and on Hopen, approx. 100 km to the southeast. These faults are again dated to the Carnian. A normal fault that strikes in a similar direction to those on Edgeøya and Hopen (i.e., ESE–WNW) is mapped in Triassic strata on Barentsøya (Winsnes & Worsley, 1980), further increasing the likely footprint of this deformation. More tentatively, Harland et al. (1974) reported possible evidence for an eastward thickening Mid-Triassic Package (Botneheia and Tschermakfjellet formations) towards the Flowerdalen Fault (Parker, 1966; Major & Nagy, 1972) which is inferred as a southern continuation of the Billefjorden Fault Zone in central Spitsbergen.

Offshore, apart from the Triassic activity in the Hammerfest basin described herein (article II),

additional zones of local fault activity have been described. They include the Fingerdjupet Sub-basin (Serck et al., 2017), the southern Loppa High Fault System (Gabrielsen, 1984; Indræver et al., 2016) and the Hoop Graben (Mahajan et al., 2014), several of which have been credited to far field stresses related to the Uralian orogeny (Svalbard, the southern Loppa High Fault System and perhaps the Hammerfest Basin). This new information gives insight into the Triassic evolution of the Barents shelf. The aforementioned Structural elements of the Barents Shelf where Triassic activity has been identified are highlighted in Fig. 19.

Figure 19 Positions and directions of Late Triassic extension across the Barents Shelf. The data is derived from articles II, III and appendix - section 9.3, in addition to previously published literature as listed. The base structural elements map is from Worsley, (2008). Bottom right: Strain ellipse showing ideal formation of tensile and compressional brittle structures given the orientation of maximum horizontal stress (σ_H) for the Uralides during the Middle to Late Triassic (derived from appendix - section 9.3).

The entire Barents shelf experienced a regional deep seated response to the Uralian orogeny which instigated increased subsidence rates, especially in the east (Worsley, 2008; Gac et al., 2014). Furthermore, the orogeny provided a prominent sediment source for large deltaic systems that

prograded from the southeast, and across the Barents shelf from the Late Permian and onwards (Riis et al., 2008; Glorstad-Clark et al., 2010; Høy & Lundschieen, 2011). For Svalbard, Hopen and the northern Barents shelfal area, article III (also see appendix, section 9.3) relates the driving mechanism of Late Triassic growth faults to the model first proposed by Anell et al. (2013) where indentation stresses driven by the Uralian orogeny reactivated deep seated faults of Carboniferous-Permian age (or older) that define the Edgeøya platform (northeast–southwest to east–west fabrics). The age of the event is constrained between the Early Carnian and the Early Norian by syn-kinematic geometries adjacent to faults, and onlap of the Tschermakfjellet Formation upon a fault monocline within the Botneheia Fm (appendix - section 9.3).

The subsequent progradation of a deltaic system (sourced from the Uralides) towards the northwest and against the southern flank of the fault bounded Edgeøya platform resulted in differential clinoform advancement rates and delta lobe indentation. Differential compaction-driven subsidence between the pro-delta and the thicker delta top and plain may be credited for the development of a counter regional dipping slope and a subsequent listric growth fault system at the delta toe (article III). A normal fault showing a similar orientation was mapped in Triassic strata on Barentsøya by Winsnes & Worsley (1980), further increasing the likely footprint of this deformation. Anell et al. (2013) also reported the presence of these faults in offshore seismic imagery where they propagate up-section and interact with turbulent zones consistent with listric fault bound growth basins, although on the limits of seismic resolution (see section 6.1). The Cenozoic Alpine foreland has been put forward as an analogue where reactivation of structures as far away as 1300 km has been reported (Ziegler, 1987; Zielger et al., 1995).

In the Hammerfest Basin (articles I and II), a similar aged phase of extension (Norian–Rhaetian) can be tentatively linked to the Uralian orogeny based on correlation of a sequence boundary that coincides with the Taimyr deformation of the Uralides (article II), the timing of which has been constrained to post-Mid Triassic to Late Triassic times (Torsvik and Anderson, 2002; Klausen et al., 2016). It is important, however, to note that rejuvenation of Fennoscandia to the south of the Hammerfest Basin and/or vertical movements related to the early Cimmerian tectonic phase in northern Europe (Ziegler, 1982; Mayall, 1983; Zonenshain et al., 1990; Golonka, 2004) also occurred contemporaneously to this phase of Norian to Rhaetian tectonism, and may have contributed as driving forces.

7. CONCLUSIONS AND FUTURE WORK

7.1 MAIN FINDINGS OF THE THESIS

This thesis investigated Late Triassic faulting in the Barents shelf and the controls of syn- and post-kinematic faulting on reservoir properties by 1) controlling the distribution of sand-prone facies and by 2) changing the fluid flow properties of the pristine country rock. For specific conclusions of each study, the reader is referred to the respective articles. The thesis can, however, be distilled into three main findings:

1) The Triassic Barents shelf was more tectonically active than previously described; far field (intraplate) stresses (the maximum horizontal stress was orientated WNW–ESE) related to the Uralian orogeny drove an Early Carnian to Early Norian episode of extension that instigated growth basins in a pro-delta environment (eastern Svalbard). A slightly later phase of ?Norian to Rhaetian extension took place in the Hammerfest Basin and on the basin-platform delineating Troms-Finnmark Fault Complex and may have been triggered by one or a combination of three events, i) rejuvenation of Fennoscandia to the south, ii) vertical movements related to the early Cimmerian tectonic phase in northern Europe and iii) to movement on the Novaya Zemlya Segment of the Pai-Khoi–Novaya Zemlya–South Taimyr fold-and-thrust belt to the east. The findings herein compliment recent evidence for Triassic tectonism in the Fingerdjupe Subbasin, the southern Loppa High Fault System, the Hoop Graben and onshore Svalbard.

2) Growth faults in deltaic settings are driven/influenced by a multitude of processes which include the generation of excess pore pressures by rapid deposition in highly under consolidated marine sediments, differential compaction, reactivation of basement structures, localised slope over steepening, and earthquake induced stresses. Deltaic growth faults typically exhibit three-dimensional concave (cusate) fault geometry. Less curvilinear fault traces such as those described in Kvalpynten, Edgeøya, may reflect influence of reactivation of underlying (soft linked) basement faults. Toe-thrust faults that occur in a pro-delta environment in response to extension higher up on the delta front will not necessarily form where basement reactivation is a driving force, given they do not have to coincide with gravitational driven movements down-slope, and may form in less sand-prone environments.

Early calcite cementation and compaction driven dewatering within syn-kinematic sediments are shown to play a significant role regarding the deformation mechanisms active within growth fault systems, the onset of which control a transition from hydro-plastic strain diffusion (grain-contact sliding) to brittle strain localisation (grain fracturing) along faults. Differential compaction-driven

subsidence and subsequent sediment loading by prograding delta systems can induce localised tilting, the direction of which has a controlling factor on fault dip directions (preferentially down-slope).

3) To date, many authors (e.g., Gawthrope & Leeder, 2000) have demonstrated that syn-kinematic sediment distribution can be controlled by fault related geomorphology and can therefore impact the geometry and connectivity of reservoir quality sandstones. Fault related uplift and erosion during a previously undocumented Norian to Rhaetian phase of extension in the southern Hammerfest Basin (Goliat area) sourced immature sediments that were deposited in an evolving rift basin (continental to coastal plain environments) and were routed towards depocentres via relay zones and across-fault scarps. Gross sand prone bodies have been delineated using amplitude versus angle seismic attribute analysis and reveal channelised features that were rerouted by fault controlled geomorphology and exhibit axial drainage in the hanging walls of faults which resulted in stacked and amalgamated channel facies. The recognition of this event and associated sand body distribution will have consequences for the continued development of the Goliat field for which the Realgrunnen Subgroup forms a reservoir unit.

7.2 FURTHER STUDIES

The superb EN0901 MAZ seismic survey is a fantastic resource for investigating fault growth given it images a complex zone of deformation within a bend on the TFFC and interactions between four fault populations which result in intricate compartmentalisation of the Goliat field, especially within the Realgrunnen Subgroup reservoir. The work herein outlined the structural framework of the Goliat field and the facies associations of the Realgrunnen Subgroup. This study can form the basis of a more detailed investigation of fault compartmentalisation, i.e., the detailed knowledge of the facies can be combined with the mapped fault surfaces in order to conduct detailed juxtaposition and membrane seal analysis.

In the Goliat study, the evolution of the Goliat anticline was described primarily as the consequence of differential subsidence controlled by two vertical segments of the TFFC. The recognition of an inversion event during the Barremian, in part manifested as buttress folding in the hanging walls of large faults, suggests the Goliat anticline may in part have been accentuated during the inversion. In order to more accurately differentiate the causal mechanisms, a structural restoration (balancing) of the Goliat area based on the seismic interpretation is merited in order to quantify the extensional and compressional strain (stretching and shortening) that has occurred since the early Mesozoic. Due to assumptions made by 1D and 2D balancing techniques (e.g., Gibbs, 1983; Rowan, 1989) that restoration of sections must be conducted in the plane of the deformation

transport direction in order to preserve both line length and area, the multiple trending fault populations affecting the Goliat field will require a thorough 3D balancing workflow.

To date, 2D seismic surveys acquired in Adventdalen (central Spitsbergen) by snow-streamer technology (Bælum et al., 2012) have failed to resolve the baffling subsurface features identified in outcrop (faults and possibly igneous dykes and stratigraphic pinch-outs). The virtual outcrop models of the target reservoir and cap rock sections can form the basis for generating velocity models in order to synthetically model the seismic signature of the discontinuities and heterogeneities affecting the target units (e.g., Lecomte et al., 2015; Anell et al., 2016). The synthetic seismic parameters can then be modified in order to determine (i) the adequate future survey methods required to image these features (if possible), and (ii) predict the potential contribution of sub-seismic features in similar investigations based on the quality of seismic data available.

The identification of polyphase deformation mechanisms within growth fault systems controlled by progressive cementation and compaction driven dewatering is important to understand as petrophysical properties in damage zones will vary largely between the two identified end-members (diffused hydro-plastic strain to brittle strain localisation). Understanding the precise controls on this progressive change, and how they can be up-scaled into reservoir models is critical for plays in such sand-prone facies. Further outcrop studies of growth fault systems are merited and potential high quality outcrops include, the Western Irish Namurian Basin, Co. Clare (e.g., Wignall & Best, 2000) and the Makran foreland, SW Pakistan (e.g., Back & Morley, 2006).

8. REFERENCES

- Ackermann, R. V., Schlische, R.W., Withjack, M.O., 2001. The geometric and statistical evolution of normal fault systems: An experimental study of the effects of mechanical layer thickness on scaling laws. *J. Struct. Geol.* 23, 1803–1819. doi:10.1016/S0191-8141(01)00028-1
- Ahmed, W., 2012. Structural Analysis of the Troms-Finnmark Fault Complex, SW Barents Sea, Masters Thesis, University of Oslo.
- Ahmed, S., Bhattacharya, J.P., Garza, D.E., Li, Y., 2014. Facies Architecture and Stratigraphic Evolution of A River-Dominated Delta Front, Turonian Ferron Sandstone, Utah, U.S.A. *J. Sediment. Res.* 84, 97–121. doi:10.2110/jsr.201 4.6
- Alexander, J., Leeder, M.R., 1987. Active Tectonic Control on Alluvial Architecture. *Recent Dev. Fluv. Sedimentol.* 243–252. doi:10.2110/pec.87.39.0243
- Anderson, E.M., 1951. The dynamics of faulting and dyke formation with applications to Britain. Hafner Pub. Co.
- Anderson, C.R., Ponce, A.M., Price, R.J., 2004. Immunohistochemical identification of an extracellular matrix scaffold that microguides capillary sprouting in vivo. *J. Histochem. Cytochem.* 52, 1063–1072.
- Andresen, A., Bergh, S.G., Haremo, P., 1992. Basin inversion and thin-skinned deformation associated with the Tertiary transpressional west Spitsbergen orogen, in: *International Conference on Arctic Margins*. pp. 1–6.
- Andresen, A., Agyei-Dwarko, N.Y., Kristoffersen, M., Hanken, N.-M., 2014. A Timanian foreland basin setting for the late Neoproterozoic–Early Palaeozoic cover sequences (Dividal Group) of northeastern Baltica. *Geol. Soc. London, Spec. Publ.* 390. doi:10.1144/SP390.29
- Anell, I., Braathen, A., Olaussen, S., Osmundsen, P.T., 2013. Evidence of faulting contradicts a quiescent northern Barents Shelf during the Triassic. *First Break* 31, 67–76. doi:10.3997/1365-2397.2013017
- Anell, I., Braathen, A., Olaussen, S., 2014. The Triassic – Early Jurassic of the northern Barents Shelf: a regional understanding of the Longyearbyen CO₂ reservoir. *Nor. J. Geol.* 94, 83–98.
- Anell, I., Lecomte, I., Braathen, A., Buckley, S.J., 2016. Synthetic seismic illumination of small-scale growth faults, paralic deposits and low-angle clinoforms: A case study of the Triassic successions on Edgeøya, NW Barents Shelf. *Mar. Pet. Geol.* 77, 625–639. doi:10.1016/j.marpetgeo.2016.07.005
- Armstrong, F.C., Oriel, S.S., 1965. Tectonic development of Idaho-Wyoming thrust belt. *Am. Assoc. Pet. Geol. Bull.* 49, 1847–1861. doi:10.1306/A663386E-16C0-11D7-8645000102C1865D
- Back, S., Morley, C.K., 2016. Growth faults above shale–Seismic-scale outcrop analogues from the Makran foreland, SW Pakistan. *Marine and Petroleum Geology* 70, 144–162.

- Bælum, K., Johansen, T.A., Johnsen, H., Rød, K., Ruud, B.O., Braathen, A., 2012. Subsurface structures of the longyearbyen CO₂ lab study area in central spitsbergen (arctic Norway), as mapped by reflection seismic data. *Norsk Geologisk Tidsskrift* 92, 377–389.
- Bally, A., Bernoulli, D., Davis, A., Montadert, L., 1981. Listric normal faults. *Oceanol. Acta*, Spec. issue 87–101.
- Barr, D., 1987. Lithospheric stretching, detached normal faulting and footwall uplift. *Cont. Extensional Tectonics* 28, 75–94. doi:10.1144/GSL.SP.1987.028.01.07
- Baum, M.S., Withjack, M.O., Schlische, R.W., 2008. The Ins and Outs of Buttress Folds: Examples from the Inverted Fundy Rift Basin, Nova Scotia and New Brunswick, Canada. *Cent. Atl. Conjug. Margins* 53–61.
- Bergh, S.G., Braathen, A., Andresen, A., 1997. Interaction of basement-involved and thin-skinned tectonism in the Tertiary fold-thrust belt of central Spitsbergen, Svalbard. *Am. Assoc. Pet. Geol. Bull.* 81, 637–661. doi:10.1306/522B43F7-1727-11D7-8645000102C1865D
- Berglund, L.T., Augustson, G., Færseth, R., Ramberg-Moe, H., 1986. The evolution of the Hammerfest Basin. *Habitat Hydrocarb. Nor. Cont. Margin* 319–338.
- Bhattacharya, J.P., Davies, R.K., 2001. Growth faults at the prodelta to delta-front transition, Cretaceous Ferron sandstone, Utah. *Mar. Pet. Geol.* 18, 525–534. doi:10.1016/S0264-8172(01)00015-0
- Bhattacharya, J.P., MacEachern, J., 2000. Hyperpycnal Rivers and Prodeltaic Shelves in the Cretaceous Seaway of North America. *J. Sediment. Res.* 79, 184–209. doi:10.1111/joa.12085
- Bhattacharya, J.P., Tye, R.S., 2004. Searching for modern Ferron analogs and application to subsurface interpretation. *Reg. to wellbore Analog fluvial-deltaic Reserv. Model. Ferron Sandstone Utah AAPG Stud. Geol.* 50, 39–57.
- Blair, T.C., McPherson, J.G., 1994. Alluvial fans and their natural distinction from rivers based on morphology, hydraulic processes, sedimentary processes, and facies assemblages. *J. Sediment. Res.* 64.
- Bohloli, B., Skurtveit, E., Grande, L., Titlestad, G.O., Børresen, M.H., Johnsen, Ø., Braathen, A., Braathen, A., 2014. Evaluation of reservoir and cap-rock integrity for the longyearbyen CO₂ storage pilot based on laboratory experiments and injection tests. *Norsk Geologisk Tidsskrift* 94, 171–187.
- Bonini, M., Sani, F., Antonielli, B., 2012. Tectonophysics Basin inversion and contractional reactivation of inherited normal faults : A review based on previous and new experimental models. *Tectonophysics* 522–523, 55–88. doi:10.1016/j.tecto.2011.11.014
- Braathen, A., Bergh, S.G., 1995. Kinematics of Tertiary deformation in the basement-involved fold-thrust complex, western Nordenskiöld Land, Svalbard: tectonic implications based on fault-slip data analysis. *Tectonophysics* 249, 1–29. doi:10.1016/0040-1951(95)00036-M

- Braathen, A., Bergh, S.G., Maher, H.D., 1999. Application of a critical wedge taper model to the tertiary transpressional fold-thrust belt on Spitsbergen, Svalbard. *Bull. Geol. Soc. Am.* 111, 1468–1485. doi:10.1130/0016-7606(1999)111<1468:AOACWT>2.3.CO;2
- Braathen, A., Nordgulen, O., Osmundsen, P.T., Andersen, T.B., Solli, A., Roberts, D., 2000. Devonian, orogen-parallel, opposed extension in the Central Norwegian Caledonides. *Geology* 28, 615–618. doi:10.1130/0091-7613(2000)28<615:DOOEIT>2.0.CO;2
- Braathen, A., Osmundsen, P.T., Gabrielsen, R.H., 2004. Dynamic development of fault rocks in a crustal-scale detachment: An example from western Norway. *Tectonics* 23. doi:10.1029/2003TC001558
- Braathen, A., Bælum, K., Maher, H., Buckley, S.J., 2011. Growth of extensional faults and folds during deposition of an evaporite-dominated half-graben basin; the Carboniferous Billefjorden Trough, Svalbard. *Nor. Geol. Tidsskr.* 91, 137–161.
- Braathen, A., Bælum, K., Christiansen, H.H., Dahl, T., Eiken, O., Elvebakk, H., Hansen, F., Hanssen, T.H., Jochmann, M., Johansen, T. a., Johnsen, H., Larsen, L., Lie, T., Mertes, J., Mørk, A., Mørk, M.B., Nemec, W., Olaussen, S., Oye, V., Rød, K., Titlestad, G.O., Tveranger, J., Vagle, K., 2012. The Longyearbyen CO₂ Lab of Svalbard, Norway — initial assessment of the geological conditions for CO₂ sequestration. *Nor. J. Geol.* 92, 353–376.
- Braathen, A., Osmundsen, P.T., Hauso, H., Semshaug, S., Fredman, N., Buckley, S.J., 2013. Fault-induced deformation in a poorly consolidated, siliciclastic growth basin: A study from the Devonian in Norway. *Tectonophysics* 586, 112–129.
- Braathen, A., Erambert, M., 2014. Structural and metamorphic history of the Engebøfjellet Eclogite and the exhumation of the Western Gneiss Region, Norway. *Norsk Geologisk Tidsskrift* 94, 53–76.
- Brandes, C., Polom, U., Winsemann, J., 2011. Reactivation of basement faults: Interplay of ice-sheet advance, glacial lake formation and sediment loading. *Basin Res.* 23, 53–64. doi:10.1111/j.1365-2117.2010.00468.x
- Brekke, T., Krajewski, K.P., Hubred, J.H., 2014. Organic geochemistry and petrography of thermally altered sections of the Middle Triassic Botneheia Formation on south-western Edgeøya, Svalbard. *Norwegian Petroleum Directorate Bulletin* 11, 111–128.
- Bridge, J.S., Leeder, M.R., 1979. A simulation model of alluvial stratigraphy. *Sedimentology* 26, 617–644.
- Brown, D., Puchkov, V., Alvarez-Marron, J., Bea, F., Perez-Estaún, A., 2006. Tectonic processes in the Southern and Middle Urals: an overview. *Geol. Soc. London, Mem.* 32, 407–419. doi:10.1144/gsl.mem.2006.032.01.24
- Browning, J., Gudmundsson, A., 2015. Caldera faults capture and deflect inclined sheets: an alternative mechanism of ring dike formation. *Bulletin of Volcanology* 77, 4. doi:10.1007/s00445-014-0889-4
- Buchanan, P.G., McClay, K.R., 1991. Sandbox experiments of inverted listric and planar fault systems. *Tectonophysics* 188, 97–115. doi:10.1016/0040-1951(91)90317-L

- Buckley, S. J., Naumann, N., Kurz, T. H., Eide, C.H., 2016. 2nd Virtual Geoscience Conference, Proceedings Volume, Uni Research CIPR, Bergen, Norway: 212 pages.
- Bull, J.M., Barnes, P.M., Lamarche, G., Sanderson, D.J., Cowie, P.A., Taylor, S.K., Dix, J.K., 2006. High-resolution record of displacement accumulation on an active normal fault: implications for models of slip accumulation during repeated earthquakes. *J. Struct. Geol.* 28, 1146–1166. doi:10.1016/j.jsg.2006.03.006
- Burke, K., Dewey, J.F., 1973. Plume-Generated Triple Junctions: Key Indicators in Applying Plate Tectonics to Old Rocks. *J. Geol.* 81, 406–433. doi:10.1086/627882
- Caine, J.S., Evans, J.P., Forster, C.B., 1996. Fault zone architecture and permeability structure. *Geology* 24, 1025–1028. doi:10.1130/0091-7613(1996)024<1025
- Cartwright, J.A., Trudgill, B.D., Mansfield, C.S., 1995. Fault growth by segment linkage: an explanation for scatter in maximum displacement and trace length data from the Canyonlands Grabens of SE Utah. *J. Struct. Geol.* 17, 1319–1326. doi:10.1016/0191-8141(95)00033-A
- Cartwright, J. A., Mansfield, C., Trudgill, B., 1996. The growth of normal faults by segment linkage. *Geol. Soc. London, Spec. Publ.* 99, 163–177. doi:10.1144/GSL.SP.1996.099.01.13
- Childs, C., Manzocchi, T., Nicol, A., Walsh, J.J., Soden, A.M., Conneally, J.C., Delogkos, E., 2016. The relationship between normal drag, relay ramp aspect ratio and fault zone structure. *Geol. Soc. London, Spec. Publ.* 439. doi:10.1144/SP439.16
- Childs, C., Watterson, J., Walsh, J.J., 1995. Fault overlap zones within developing normal fault systems. *J. Geol. Soc. London.* 152, 535–549. doi:10.1144/gsjgs.152.3.0535
- Childs, C., Nicol, A., Walsh, J.J., Watterson, J., 1996. Growth of vertically segmented normal faults. *J. Struct. Geol.* 18, 1389–1397. doi:10.1016/S0191-8141(96)00060-0
- Childs, C., Nicol, A., Walsh, J.J., Watterson, J., 2002. The growth and propagation of synsedimentary faults. *J. Struct. Geol.* 25, 633–648. doi:10.1016/S0191-8141(02)00054-8
- Choi, J.H., Edwards, P., Ko, K., Kim, Y.S., 2016. Definition and classification of fault damage zones: A review and a new methodological approach. *Earth-Science Rev.* 152, 70–87. doi:10.1016/j.earscirev.2015.11.006
- Coleman, J.M., Prior, D.B., 1982. Deltaic environments of deposition. Sandstone depositional environments *Mem.* 31, 139–178.
- Cooper, M. a., Williams, G.D., de Graciansky, P.C., Murphy, R.W., Needham, T., de Paor, D., Stoneley, R., Todd, S.P., Turner, J.P., Ziegler, P. a., 1989. Inversion tectonics -- a discussion. *Geol. Soc. London, Spec. Publ.* 44, 335–347. doi:10.1144/GSL.SP.1989.044.01.18
- Copeland, P., Currie, C.A., Lawton, T.F., Murphy, M.A., 2017. Location, location, location: The variable lifespan of the Laramide orogeny. *Geology* 45, 223–226. doi:10.1130/G38810.1
- Corfield, S., Sharp, I.R., 2000. Structural style and stratigraphic architecture of fault propagation folding in extensional settings: a seismic example from the Smørbukk area, Halten Terrace, Mid-Norway. *Basin Res.* 12, 329–341. doi:10.1111/j.1365-2117.2000.00133.x

- Corfu, F., Polteau, S., Planke, S., Faleide, J.I., Svensen, H., Zayoncheck, A., Stolbov, N., 2013. U–Pb geochronology of Cretaceous magmatism on Svalbard and Franz Josef Land, Barents Sea Large Igneous Province. *Geol. Mag.* 150, 1127–1135. doi:10.1017/S0016756813000162
- Cotter, E., 1975. Late Cretaceous sedimentation in a low-energy coastal zone: the Ferron Sandstone of Utah. *J. Sediment. Res.* 45, 669–685.
- Cowie, P.A., Scholz, C.H., 1992. Displacement-length scaling relationship for faults: data synthesis and discussion. *J. Struct. Geol.* 14, 1149–1156. doi:10.1016/0191-8141(92)90066-6
- Cutbill, J.L., Challinor, A., 1965. Revision of the Stratigraphical Scheme for the Carboniferous and Permian Rocks of Spitsbergen and Bjørnøya. *Geol. Mag.* 102, 418–439. doi:10.1017/S0016756800053693
- Dallmann, W.K., 1999. Lithostratigraphic Lexicon of Svalbard: review and recommendations for nomenclature use: Upper Paleozoic to Quaternary Bedrock. Committee on the Stratigraphy of Svalbard. Norsk Polarinstitut.
- Dart, C.J., McClay, K., Hollings, P.N., 1995. 3D analysis of inverted extensional fault systems, southern Bristol Channel basin, UK. *Geol. Soc. London Spec. Publ.* 88, 393–413. doi:10.1144/GSL.SP.1995.088.01.21
- Dawers, N.H., Anders, M.H., 1995. Displacement-length scaling and fault linkage. *J. Struct. Geol.* 17, 607–614. doi:10.1016/0191-8141(94)00091-D
- Dean, W.E., Arthur, M.A., 1989. Iron-sulfur-carbon relationships in organic-carbon-rich sequences I: Cretaceous Western Interior seaway. *American Journal of Science* 289, 708–743. doi:10.2475/ajs.289.6.708
- DeCelles, P.G., Coogan, J.C., 2006. Regional structure and kinematic history of the Sevier fold-and-thrust belt, central Utah. *Geol. Soc. Am. Bull.* 118, 841–864. doi:10.1130/B25759.1
- Dengo, C.A., Røssland, K.G., 1992. Extensional tectonic history of the western Barents Sea. *Structural Tecton. Model. its Appl. to Pet. Geol.* 1, 91–107.
- Densmore, A.L., Dawers, N.H., Gupta, S., Allen, P.A., Gilpin, R., 2003. Landscape evolution at extensional relay zones. *J. Geophys. Res.* 108, 1–15. doi:10.1029/2001JB001741
- Densmore, A.L., Dawers, N.H., Gupta, S., Guidon, R., Goldin, T., 2004. Footwall topographic development during continental extension. *J. Geophys. Res.* 109, 1–16. doi:10.1029/2003JF000115
- Deveugle, P.E.K., Jackson, M.D., Hampson, G.J., Farrell, M.E., Sprague, A.R., Stewart, J., Calvert, C.S., 2011. Characterization of stratigraphic architecture and its impact on fluid flow in a fluvial-dominated deltaic reservoir analog: Upper Cretaceous Ferron Sandstone Member, Utah. *Am. Assoc. Pet. Geol. Bull.* 95, 693–727.
- Dooley, T., McClay, K.R., Pascoe, R., 2003. 3D analogue models of variable displacement extensional faults: applications to the Revfallet Fault system, offshore mid-Norway. *New Insights into Struct. Interpret. Model.* 212, 151–167. doi:10.1144/Gsl.Sp.2003.212.01.10

- Drachev, S.S., 2016. Fold belts and sedimentary basins of the Eurasian Arctic. *Arktos* 2, 21. doi:10.1007/s41063-015-0014-8
- Dula, W.F., 1991. Geometric models of listric faults and rollover folds. *Am. Assoc. Pet. Geol. Bull.* 75, 1609–1625. doi:10.1306/0C9B29B1-1710-11D7-8645000102C1865D
- Duncan, C.C., Turcotte, D.L., 1994. On the breakup and coalescence of continents. *Geology* 22, 103–106. doi:10.1130/0091-7613(1994)022<0103:OTBACO>2.3.CO
- Dypvik, H., Nagy, J., Eikeland, T.A., Backer-Owe, K., Johansen, H., 1991. Depositional conditions of the Bathonian to Hauterivian Janusfjellet Subgroup, Spitsbergen. *Sediment. Geol.* 72, 55–78. doi:10.1016/0037-0738(91)90123-U
- Edwards, M.B., 1976. Growth faults in Upper Triassic deltaic sediments, Svalbard. *AAPG Bulletin* 60, 341–355.
- Eidvin, T., Goll, R.M., Grogan, P., Smelror, M., Ulleberg, K., 1998. The Pleistocene to Middle Eocene stratigraphy and geological evolution of the western Barents Sea continental margin at well site 7316/5-1 (Bjørnøya West area). *Nor. Geol. Tidsskr.* 78, 99–124.
- Eliet, P.P., Gawthorpe, R.L., 1995. Drainage development and sediment supply within rifts, examples from the Sperchios basin, central Greece. *J. Geol. Soc. London.* 152, 883–893. doi:10.1144/gsjgs.152.5.0883
- Elliott, G.M., Wilson, P., Jackson, C.A.L., Gawthorpe, R.L., Michelsen, L., Sharp, I.R., 2011. The linkage between fault throw and footwall scarp erosion patterns: An example from the Bremstein Fault Complex, offshore Mid-Norway. *Basin Res.* 24, 180–197. doi:10.1111/j.1365-2117.2011.00524.x
- Ellis, P.G., McClay, K.R., 1988. Listric extension fault systems; results of analogue model experiments. *Basin Res.* 1, 55.
- Elverhøi, A., Antonsen, P., Flood, S.B., Solheim, A., Vullstad, A.A., 1988. The physical environment, western Barents Sea 1: 1,500 000, Shallow bedrock geology. *Nor. Polarinstitutt Skr.* 179, 40.
- Enge, H.D., Howell, J.A., 2010. Impact of deltaic clinoforms on reservoir performance: Dynamic studies of reservoir analogs from the Ferron Sandstone Member and Panther Tongue, Utah. *Am. Assoc. Pet. Geol. Bull.* 94, 139–161. doi:10.1306/07060908112
- Ethridge, F.G., Wescott, W.A., 1984. Tectonic Setting, Recognition and Hydrocarbon Reservoir Potential of Fan-Delta Deposits In; Koster, E.M. and Steel, R.J. (Eds.) *Sedimentology of Gravels and Conglomerates*. Canadian Society of Petroleum Geologists, Memoir 10 (1984), p. 217–235. *Can. Soc. Pet. Geol.* 0, 217–235.
- Faleide, J.I., Gudlaugson, S.T., Jacquart, G., 1984. Evolution of the western Barents Sea. *Mar. Pet. Geol.* 1, 123–150.
- Faleide, J.I., Vågnes, E., Gudlaugsson, S.T., 1993. Late Mesozoic-Cenozoic evolution of the southwestern Barents Sea in a regional rift-shear tectonic setting. *Mar. Pet. Geol.* 10, 186–214. doi:http://dx.doi.org/10.1016/0264-8172(93)90104-Z

- Faleide, J.I., Tsikalas, F., Breivik, a J., Mjelde, R., Ritzmann, O., Engen, O., Wilson, J., Eldholm, O., 2008. Structure and evolution of the continental margin off Norway and the Barents Sea. *Episodes* 31, 82–91. doi:10.1016/j.strusafe.2006.11.005
- Faleide, J.I., Bjørlykke, K., Gabrielsen, R.H., 2015. Geology of the Norwegian continental shelf, in: *Petroleum Geoscience*. Springer, pp. 603–637.
- Faleide, J.I., Pease, V., Curtis, M., Klitzke, P., Minakov, A., Scheck-Wenderoth, M., Kostyuchenko, S., Zayonchek, A., 2017. Tectonic implications of the lithospheric structure across the Barents and Kara shelves. Geological Society, London, Special Publications SP460.18. doi:10.1144/SP460.18
- Farokhpoor, R., Torsaeter, O., Baghbanbashi, T., Mørk, A., Lindeberg, E., 2010. Experimental and Numerical Simulation of CO₂ Injection Into Upper-Triassic Sandstones in Svalbard, Norway. *Society of Petroleum Engineers SPE* 139524 1-11.
- Fielding, C.R., 2010. Planform and Facies Variability in Asymmetric Deltas: Facies Analysis and Depositional Architecture of the Turonian Ferron Sandstone in the Western Henry Mountains, South-Central Utah, U.S.A. *J. Sediment. Res.* 80, 455–479. doi:10.2110/jsr.2010.047
- Fielding, C.R., 2015. Anatomy of falling-stage deltas in the Turonian Ferron Sandstone of the western Henry Mountains Syncline, Utah: Growth faults, slope failures and mass transport complexes. *Sedimentology* 62, 1–26. doi:10.1111/sed.12136
- Finch, E., Hardy, S., Gawthorpe, R., 2004. Discrete-element modelling of extensional fault-propagation folding above rigid basement fault blocks. *Basin Res.* 16, 489–506. doi:10.1111/j.1365-2117.2004.00241.x
- Fort, X., Brun, J.P., Chauvel, F., 2004. Salt tectonics on the Angolan margin, synsedimentary deformation processes. *Am. Assoc. Pet. Geol. Bull.* 88, 1523–1544. doi:10.1306/06010403012
- Fossen, H., Gabrielsen, R.H., 1996. Experimental modeling of extensional fault systems by use of plaster. *J. Struct. Geol.* 18, 673–687. doi:10.1016/S0191-8141(96)80032-0
- Fossen, H., Schultz, R. A., Shipton, Z.K., Mair, K., 2007. Deformation bands in sandstone: a review. *J. Geol. Soc. London.* 164, 755–769. doi:10.1144/0016-76492006-036
- Fossen, H., 2016. *Structural Geology*. Cambridge University Press.
- Foster, D., Brocklehurst, S.H., Gawthorpe, R.L., 2010. Glacial-topographic interactions in the Teton Range, Wyoming. *J. Geophys. Res. Earth Surf.* 115, 1–20. doi:10.1029/2008JF001135
- Friedman, G.M., Sanders, J.E., 1978. *Principles of sedimentology*. Wiley, New York. doi:10.1002/esp.3290040317
- Gabrielsen, R.H., 1984. Long-lived fault zones and their influence on the tectonic development of the southwestern Barents Sea. *J. Geol. Soc. London.* 141, 651–662. doi:10.1144/gsjgs.141.4.0651
- Gabrielsen, R.H., Foersth, R., Hamar, G., Rønnevik, H., 1984. Nomenclature of the main structural features on the Norwegian Continental Shelf north of the 62nd parallel BT - Petroleum Geology of the North European Margin: Proceedings of the North European Margin Symposium

- (NEMS '83), organized by the Norwegian Petr, in: Spencer, A.M. (Ed.), . Springer Netherlands, Dordrecht, pp. 41–60. doi:10.1007/978-94-009-5626-1_5
- Gabrielsen, R.H., Færseth, R.B., 1988. Cretaceous and Tertiary reactivation of master fault zones of the Barents Sea. *Nor. Polarinstitut Rapp* 46, 93–97.
- Gabrielsen, R.H., Færseth, R.B., 1989. The inner shelf of North Cape, Norway and its implications for the Barents Shelf-Finnmark Caledonide boundary. A comment. *Nor. Geol. Tidsskr.* 69, 57–62.
- Gabrielsen, R.H., Færseth, R.B., Jensen, L.N., Kalheim, J.E., Riis, F., 1990. Structural Elements of the Norwegian continental shelf. Part 1: The Barents Sea Region. *Nor. Pet. Dir. Bull.* No.6, 33 pp.
- Gabrielsen, R.H., Faleide, J.I., Pascal, C., Braathen, A., Nystuen, J.P., Etzelmuller, B., O'Donnell, S., 2010. Latest Caledonian to Present tectonomorphological development of southern Norway. *Mar. Pet. Geol.* 27, 709–723. doi:http://dx.doi.org/10.1016/j.marpetgeo.2009.06.004
- Gac, S., Huismans, R.S., Simon, N.S.C., Faleide, J.I., Podladchikov, Y.Y., 2014. Effects of lithosphere buckling on subsidence and hydrocarbon maturation: A case-study from the ultra-deep East Barents Sea basin. *Earth Planet. Sci. Lett.* 407, 123–133. doi:10.1016/j.epsl.2014.09.029
- Gaina, C., Gernigon, L., Ball, P., 2009. Palaeocene-Recent plate boundaries in the NE Atlantic and the formation of the Jan Mayen microcontinent. *J. Geol. Soc. London.* 166, 601–616. doi:10.1144/0016-76492008-112
- García-García, F., Fernández, J., Viseras, C., Soria, J.M., 2006. Architecture and sedimentary facies evolution in a delta stack controlled by fault growth (Betic Cordillera, southern Spain, late Tortonian). *Sediment. Geol.* 185, 79–92. doi:10.1016/j.sedgeo.2005.10.010
- Gardner, M.H., 1995. Tectonic and eustatic controls on the stratal architecture of mid-Cretaceous stratigraphic sequences, central western interior foreland basin of North America. *Stratigr. Evol. Forel. Basins Stratigrap*, 243–281. doi:10.2110/pec.95.52.0243
- Gardner, M.H., Barton, M.D., Tyler, N., Fisher, R.S., 1992. Architecture and Permeability Structure of Fluvial-Deltaic Sandstone, Ferron Utah, East-central, in: *Rocky Mountain Section SEPM. Rocky Mountain Section (SEPM)*.
- Garrison Jr, J.R., van den Bergh, T.C. V, 1997. Coal zone and high-resolution depositional sequence stratigraphy of the Upper Ferron Sandstone. *Mesozoic to Recent Geol. Utah Brigham Young Univ. Geol. Stud.* 42, 160–178.
- Garrison, J.R.J., van der Bergh, T.C. V, Barker, C.E., Tabet, D.E., 1997. Depositional sequence stratigraphy and architecture of the Cretaceous Ferron Sandstone: implications for coal and coalbed methane resources - A field excursion. 2. *Mesozoic to Recent Geol. Utah* 42, 155–202.
- Gawthorpe, R.L., Leeder, M.R., 2000. Tectono-sedimentary evolution of active extensional basins. *Basin Res.* 12, 195–218. doi:10.1111/j.1365-2117.2000.00121.x
- Garrison, J.R., Bergh, T.C.V. Van Den, 2006. Effects of sedimentation rate, rate of relative rise in sea level, and duration of sea-level cycle on the filling of incised valleys: examples of filled and “overfilled” incised valleys from the upper Ferron Sandstone, Last Chance Delta, east-central Utah. *Incised Val. Time Sp.* 239–279.

- Gee, D.G., Bogolepova, O.K., Lorenz, H., 2006. The Timanide, Caledonide and Uralide orogens in the Eurasian high Arctic, and relationships to the palaeo-continent Laurentia, Baltica and Siberia. *Geol. Soc. London, Mem.* 32, 507–520. doi:10.1144/GSL.MEM.2006.032.01.31
- Gernigon, L., Bronner, M., 2012. Late Palaeozoic architecture and evolution of the southwestern Barents Sea: insights from a new generation of aeromagnetic data. *J. Geol. Soc. London.* 169, 449–459. doi:10.1144/0016-76492011-131
- Giba, M., Walsh, J.J., Nicol, A., 2012. Segmentation and growth of an obliquely reactivated normal fault. *J. Struct. Geol.* 39, 253–267. doi:10.1016/j.jsg.2012.01.004
- Gibbs, A.D., 1983. Balanced cross-section construction from seismic sections in areas of extensional tectonics. *Journal of structural geology* 5, 153–160.
- Gibbs, A.D., 1984. Structural evolution of extensional basin margins. *J. Geol. Soc. London.* 141, 609–620. doi:10.1144/gsjgs.141.4.0609
- Glørstad-clark, E., Inge, J., Anders, B., Petter, J., 2010. Triassic seismic sequence stratigraphy and paleogeography of the western Barents Sea area. *Mar. Pet. Geol.* 27, 1448–1475. doi:10.1016/j.marpetgeo.2010.02.008
- Glørstad-Clark, E., Birkeland, E.P., Nystuen, J.P., Faleide, J.I., Midtkandal, I., 2011. Triassic platform-margin deltas in the western Barents Sea. *Mar. Pet. Geol.* 28, 1294–1314. doi:10.1016/j.marpetgeo.2011.03.006
- Golonka, J., 2004. Plate tectonic evolution of the southern margin of Eurasia in the Mesozoic and Cenozoic. *Tectonophysics* 381, 235–273. doi:10.1016/j.tecto.2002.06.004
- Grasemann, B., Martel, S., Passchier, C., 2005. Reverse and normal drag along a fault. *Journal of Structural Geology* 27, 999–1010. doi:10.1016/j.jsg.2005.04.006
- Gross, M.R., Engelder, T., 1995. Strain accommodated by brittle failure in adjacent units of the Monterey Formation, U.S.A.: scale effects and evidence for uniform displacement boundary conditions. *J. Struct. Geol.* 17. doi:10.1016/0191-8141(95)00011-2
- Gross, M.R., Gutierrez-Alonso, G., Bai, T., Wacker, M.A., Collinsworth, K.B., Behl, R.J., 1997. Influence of mechanical stratigraphy and kinematics on fault scaling relations. *J. Struct. Geol.* 19, 171–183. doi:10.1016/S0191-8141(96)00085-5
- Grundvåg, S.-A., Olausson, S., 2017. Sedimentology of the Lower Cretaceous at Kikutodden and Keilhaufjellet, southern Spitsbergen: implications for an onshore–offshore link. *Polar Research* 36, 1302124.
- Gudlaugsson, S.T., Faleide, J.I., Johansen, S.E., Breivik, A.J., 1998. Late Palaeozoic structural development of the South-western Barents Sea. *Mar. Pet. Geol.* 15, 73–102. doi:10.1016/S0264-8172(97)00048-2
- Gupta, S., Underbill, J.R., Sharp, I.R., Gawthorpe, R.L., 1999. Role of fault interactions in controlling synrift sediment dispersal patterns: Miocene, Abu Alaq Group, Suez Rift, Sinai, Egypt. *Basin Res.* 11, 167–189. doi:10.1046/j.1365-2117.1999.00300.x

- Fossen, H., Schultz, R.A., Shipton, Z.K., Mair, K., 2007. Deformation bands in sandstone: a review. *Journal of the Geological Society* 164, 755–769. doi:10.1144/0016-76492006-036
- Hale, L.A., 1972. Depositional history of the Ferron Formation, central Utah, Plateau - Basin and Range Transition Zone, Central Utah, 1972: 29–40.
- Harland, W.B., Wallis, R.H., Gayer, R.A., 1966. A revision of the lower Hecla Hoek succession in central north Spitsbergen and correlation elsewhere. *Geol. Mag.* 103, 70–97. doi:10.1017/S0016756800050433
- Harland, W.B., Cutbill, J.L., Friend, P.F., Gobbett, D.J., Holliday, D.W., Maton, P.I., Parker, J.R., Wallis, R.H., 1974. The Billefjorden Fault Zone, Spitsbergen: the long history of a major tectonic lineament.
- Harland, W.B., Geddes, I., 1997. Triassic history. *Geol. Svalbard* 2, 340–362. doi:10.1144/GSL.MEM.1997.017.01.18
- Harvey, A.M., Mather, A.E., Stokes, M., 2005. Alluvial fans: geomorphology, sedimentology, dynamics — introduction. A review of alluvial-fan research. *Geol. Soc. London, Spec. Publ.* 251, 1–7. doi:10.1144/GSL.SP.2005.251.01.01
- Helland-Hansen, W., 2010. Facies and stacking patterns of shelf-deltas within the Palaeogene Battfjellet Formation, Nordenskiöld Land, Svalbard: Implications for subsurface reservoir prediction. *Sedimentology* 57, 190–208. doi:10.1111/j.1365-3091.2009.01102.x
- Hendriks, B.W.H., Andriessen, P.A.M., 2002. Pattern and timing of the post-Caledonian denudation of northern Scandinavia constrained by apatite fission-track thermochronology. *Geol. Soc. London, Spec. Publ.* 196, 117–137. doi:10.1144/gsl.sp.2002.196.01.08
- Hendriks, B., 2003. Cooling and denudation of the Norwegian and Barents Sea Margin, North Scandinavia, constrained by apatite fission track and (U-Th)/He thermochronology. doi:10.13140/RG.2.1.4347.9127
- Henriksen, E., Bjornseth, H.M., Hals, T.K., Heide, T., Kiryukhina, T., Klovjan, O.S., Larssen, G.B., Ryseth, A.E., Ronning, K., Sollid, K., Stoupakova, A., 2011. Uplift and erosion of the greater Barents Sea: impact on prospectivity and petroleum systems. *Geol. Soc. London, Mem.* 35, 271–281. doi:10.1144/M35.17
- Heynekamp, M.R., Goodwin, L.B., Mozley, P.S., Haneberg, W.C., 1999. Controls on Fault-Zone Architecture in Poorly Lithified Sediments, Rio Grande Rift, New Mexico: Implications for Fault-Zone Permeability and Fluid Flow. *Faults and subsurface fluid flow in the shallow crust* 27–49.
- Hintze, L.F., 1974. Preliminary geologic map of the Notch Peak Quadrangle, Millard County, Utah.
- Holmes, a., 1965. Principles of physical geology, Geologiska Föreningen i Stockholm. Föreläsningar. New York : Ronald Press Co. doi:10.1080/11035894509446436
- Hooke, R.L., 1967. Processes on Arid-Region Alluvial Fans. *The Journal of Geology* 75, 438–460. doi:10.1086/627271

- Hooper, R.J., Fitzsimmons, R.J., Grant, N., Vendeville, B.C., 2002. The role of deformation in controlling depositional patterns in the south-central Niger Delta, West Africa. *J. Struct. Geol.* 24, 847–859.
- Houmark-Nielsen, M., Henrik Kjaer, K., 2003. Southwest Scandinavia, 40–15 kyr BP: Palaeogeography and environmental change. *J. Quat. Sci.* 18, 769–786.
- Høy, T., Lundschie, B.A., 2011. Triassic deltaic sequences in the northern Barents Sea. *Geol. Soc. London, Mem.* 35, 249–260. doi:10.1144/M35.15
- Huggins, P., Watterson, J., Walsh, J.J., Childs, C., 1995. Relay zone geometry and displacement transfer between normal faults recorded in coal-mine plans. *J. Struct. Geol.* 17, 1741–1755. doi:10.1016/0191-8141(95)00071-K
- Imber, J., Childs, C., Nell, P.A.R., Walsh, J.J., Hodgetts, D., Flint, S., 2003. Hanging wall fault kinematics and footwall collapse in listric growth fault systems. *J. Struct. Geol.* 25, 197–208. doi:10.1016/S0191-8141(02)00034-2
- Imber, J., Tuckwell, G.W., Childs, C., Walsh, J.J., Manzocchi, T., Heath, a. E., Bonson, C.G., Strand, J., 2004. Three-dimensional distinct element modelling of relay growth and breaching along normal faults. *J. Struct. Geol.* 26, 1897–1911. doi:10.1016/j.jsg.2004.02.010
- Indrevær, K., Bergh, S.G., Koehl, J.B., Hansen, J.A., Schermer, E.R., Ingebrigtsen, A., 2013. Post-Caledonian brittle fault zones on the hyperextended SW Barents Sea margin: New insights into onshore and offshore margin architecture. *Nor. Geol. Tidsskr.* 93, 167–188.
- Indrevær, K., Stunitz, H., Bergh, S.G., 2014. On Palaeozoic–Mesozoic brittle normal faults along the SW Barents Sea margin: fault processes and implications for basement permeability and margin evolution. *Journal of the Geological Society* 171, 831–846.
- Indrevær, K., Faleide, J.I., Gabrielsen, R.H., Mulrooney, M., Braathen, A., 2015. Inversion along the Troms-Finnmark Fault Complex, SW Barents Sea - Structural character and timing, in: Oral Presentation, AAPG 3P Arctic Meeting, October, 2015, Stavanger.
- Indrevær, K., Gabrielsen, R.H., Faleide, J.I., 2016. Early Cretaceous synrift uplift and tectonic inversion in the Loppa High area, southwestern Barents Sea, Norwegian shelf. *J. Geol. Soc. London.* jgs2016-066. doi:10.1144/jgs2016-066
- Jackson, C.A.L., Gawthorpe, R.L., Sharp, I.R., 2006. Style and sequence of deformation during extensional fault-propagation folding: Examples from the Hammam Faraun and El-Qaa fault blocks, Suez Rift, Egypt. *J. Struct. Geol.* 28, 519–535. doi:10.1016/j.jsg.2005.11.009
- Johannessen, E.P., Steel, R.J., 1992. Mid-Carboniferous extension and rift-infill sequences in the Billefjorden Trough, Svalbard. *Norsk Geologisk Tidsskrift* 72, 35–48.
- Kamp, W.K., 2016. Permian to Late Triassic Structural and Stratigraphic Evolution of the Fingerdjupet Subbasin. MSc Thesis. University of Oslo.
- Kauffman, E.G., 1977. Geological and biological overview: Western Interior Cretaceous basin. *Mt. Geol.* 14, 75–99.

- Khalil, S.M., McClay, K.R., 2002. Extensional fault-related folding, northwestern Red Sea, Egypt. *J. Struct. Geol.* 24, 743–762. doi:10.1016/S0191-8141(01)00118-3
- Kim, Y.-S., Peacock, D.C.P., Sanderson, D.J., 2004. Fault damage zones. *J. Struct. Geol.* 26, 503–517. doi:http://dx.doi.org/10.1016/j.jsg.2003.08.002
- Kim, Y.S., Sanderson, D.J., 2005. The relationship between displacement and length of faults: A review. *Earth-Science Rev.* 68, 317–334. doi:10.1016/j.earscirev.2004.06.003
- Klausen, T.G., Müller, R., Slama, J., Helland-Hansen, W., 2016. Evidence for Late Triassic provenance areas and Early Jurassic sediment supply turnover in the Barents Sea Basin of northern Pangea. *Lithosphere* L556.1. doi:10.1130/L556.1
- Klausen, T.G., Müller, R., Sláma, J., Olaussen, S., Rismyhr, B., Helland-Hansen, W., 2017. Depositional history of a condensed shallow marine reservoir succession: stratigraphy and detrital zircon geochronology of the Jurassic Stø Formation, Barents Sea. *Journal of the Geological Society.* doi:10.1144/jgs2017-024
- Knarud, R., 1980. En sedimentologisk og diagenetisk undersøkelse av Kapp Toscana Formasjonens sedimenter på Svalbard. Univ. Oslo 208.
- Knies, J., Matthiessen, J., Vogt, C., Sverre, J., Hjelstuen, B.O., Smelror, M., Larsen, E., Andreassen, K., Eidvin, T., Vorren, T.O., 2009. The Plio-Pleistocene glaciation of the Barents Sea – Svalbard region : a new model based on revised chronostratigraphy. *Quat. Sci. Rev.* 28, 812–829. doi:10.1016/j.quascirev.2008.12.002
- Knipe, R.J., 1992. Faulting processes and fault seal, IN: Structural and Tectonic Modelling and its Application to Petroleum Geology: Proceedings of Norwegian Petroleum Society Workshop, 18-20 October 1989, Stavanger, Norway, 325–342.
- Knipe, R.J., Jones, G., Fisher, Q.J., 1998. Faulting, fault sealing and fluid flow in hydrocarbon reservoirs: an introduction. *Geol. Soc. London, Spec. Publ.* 147, vii–xxi. doi:10.1144/GSL.SP.1998.147.01.01
- Knott, S.D., Beach, A., Brockbank, P.J., Brown, J.L., McCallum, J.E., Welbon, A.I., 1996. Spatial and mechanical controls on normal fault populations. *J. Struct. Geol.* 18, 359–372. doi:http://dx.doi.org/10.1016/S0191-8141(96)80056-3
- Knutsen, S.M., Vorren, T.O., 1991. Early Cenozoic sedimentation in the Hammerfest Basin. *Mar. Geol.* 101, 31–48. doi:10.1016/0025-3227(91)90061-8
- Koevoets, M.J., Abay, T.B., Hammer, Olaussen, S., 2016. High-resolution organic carbon-isotope stratigraphy of the Middle Jurassic-Lower Cretaceous Agardhfjellet Formation of central Spitsbergen, Svalbard. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 449, 266–274. doi:10.1016/j.palaeo.2016.02.029
- Korja, A., Lahtinen, R., Nironen, M., 2006. The Svecofennian orogen: a collage of microcontinents and island arcs. *Geol. Soc. London, Mem.* 32, 561–578. doi:10.1144/GSL.MEM.2006.032.01.34

- Kristensen, M.B., Childs, C.J., Korstgård, J.A., 2008. The 3D geometry of small-scale relay zones between normal faults in soft sediments. *J. Struct. Geol.* 30, 257–272. doi:<http://dx.doi.org/10.1016/j.jsg.2007.11.003>
- Laberg, J.S., Andreassen, K., Vorren, T.O., 2012. Late Cenozoic erosion of the high-latitude southwestern barents sea shelf revisited. *Bull. Geol. Soc. Am.* 124, 77–88. doi:[10.1130/B30340.1](https://doi.org/10.1130/B30340.1)
- Lahr, K.M., Lahr, J.C., Lindh, A.G., Bufe, C.G., Lester, F.W., 1976. The August 1975 Oroville Earthquakes. *Bull. Seismol. Soc. Am.* 66, 1085–1099.
- Larsen, L., 2010. Analyses of Injection and Falloff Data from DH4, Aug. 12 - Sept. 4, 2010. Longyearbyen CO2 Lab UNIS Report.
- Larsen, L., 2012. Analyses of Injection and Falloff Data from DH4, Aug. 12, 2010 – Dec. 28, 2011. Longyearbyen CO2 Lab UNIS Report.
- Lecomte, I., Lubrano Lavadera, P., Anell, I., Buckley, S.J., Schmid, D.W., Heeremans, M., 2015. Ray-based seismic modeling of geologic models: Understanding and analyzing seismic images efficiently. *Interpretation* 3, SAC71-SAC89. doi:[10.1190/INT-2015-0061.1](https://doi.org/10.1190/INT-2015-0061.1)
- Leeder, M.R., 1991. Denudation, vertical crustal movements and sedimentary basin infill. *Geol. Rundschau* 80, 441–458. doi:[10.1007/BF01829376](https://doi.org/10.1007/BF01829376)
- Leeder, M.R., Jackson, J.A., 1993. The interaction between normal faulting and drainage in active extensional basins, with examples from the western United States and central Greece. *Basin Res.* 5, 79–102. doi:[10.1111/j.1365-2117.1993.tb00059.x](https://doi.org/10.1111/j.1365-2117.1993.tb00059.x)
- Leever, K.A., Gabrielsen, R.H., Sokoutis, D., Willingshofer, E., 2011. The effect of convergence angle on the kinematic evolution of strain partitioning in transpressional brittle wedges: Insight from analog modeling and high-resolution digital image analysis. *Tectonics* 30, 1–25. doi:[10.1029/2010TC002823](https://doi.org/10.1029/2010TC002823)
- Leutscher, J., Nerby, A.M., 2011. Seismic Interpretation Report, EN0901 MAZ Survey, Unpublished Eni Norge internal report.
- Leutscher, J., 2013. Goliat Field, a step-by-step improvement of the Seismic Image, in: *Production Geoscience Society of Norway, Stavanger*, 06/11/2013.
- Li, Z., Schieber, J., Bhattacharya, J.P., 2016. Facies Characteristics and Sequence Stratigraphy of the Upper Cretaceous Tununk Shale Member, Henry Mountains Region, Utah, in: *AAPG Annual Convention and Exhibition*.
- Lohr, T., Krawczyk, C.M., Oncken, O., Tanner, D.C., 2008. Evolution of a fault surface from 3D attribute analysis and displacement measurements. *J. Struct. Geol.* 30, 690–700. doi:[10.1016/j.jsg.2008.02.009](https://doi.org/10.1016/j.jsg.2008.02.009)
- Long, J.J., Imber, J., 2010. Geometrically coherent continuous deformation in the volume surrounding a seismically imaged normal fault-array. *J. Struct. Geol.* 32, 222–234. doi:[10.1016/j.jsg.2009.11.009](https://doi.org/10.1016/j.jsg.2009.11.009)

- Magnabosco, C., Braathen, A., Ogata, K., 2014. Permeability model of tight reservoir sandstones combining core-plug and Miniperm analysis of drillcore; Longyearbyen CO. *Norwegian Journal of Geology* 94, 189–200.
- Mahajan, A., Gabrielsen, R. H., Faleide, J.I., 2014. Structural Evolution using 3D Seismic of Hoop Fault Complex, SW Barents Sea Norway. *Conf. Arct. Energy Conf. Tromso*, Vol. NGF Abstr. Proc. 2014, Vol 2.
- Maher, Jr., H.D., 2001. Manifestations of the Cretaceous High Arctic Large Igneous Province in Svalbard. *J. Geol.* 109, 91–104. doi:10.1086/317960
- Maher, H.D.J., Braathen, A., 2011. Løvehovden fault and Billefjorden rift basin segmentation and development, Spitsbergen, Norway. *Geol. Mag.* 148, 154–170. doi:10.1017/S0016756810000567
- Maher, H.D., Ogata, K.E.I., Braathen, A., 2017. Cone-in-cone and beef mineralization associated with Triassic growth basin faulting and shallow shale diagenesis, Edgeøya, Svalbard. *Geological Magazine* 154, 201–216. doi:DOI: 10.1017/S0016756815000886
- Major, H., Nagy, J., 1972. Geology of the Adventdalen map area: with a geological map, Svalbard C9G 1: 100 000.
- Mansfield, C.S., Cartwright, J.A., 1996. High resolution fault displacement mapping from three-dimensional seismic data: evidence for dip linkage during fault growth. *Journal of Structural Geology* 18, 249–263.
- Mansfield, C., Cartwright, J., 2001. Fault growth by linkage: Observations and implications from analogue models. *J. Struct. Geol.* 23, 745–763. doi:10.1016/S0191-8141(00)00134-6
- Marchal, D., Guiraud, M., Rives, T., 2003. Geometric and morphologic evolution of normal fault planes and traces from 2D to 4D data. *J. Struct. Geol.* 25, 135–158. doi:10.1016/S0191-8141(02)00011-1
- Marello, L., Ebbing, J., Gernigon, L., 2010. Magnetic basement study in the Barents Sea from inversion and forward modelling. *Tectonophysics* 493, 153–171. doi:10.1016/j.tecto.2010.07.014
- Marrett, R., Allmendinger, R.W., 1991. Estimates of strain due to brittle faulting: sampling of fault populations. *J. Struct. Geol.* 13, 735–738. doi:10.1016/0191-8141(91)90034-G
- Mayall, M.J., 1983. An earthquake origin for synsedimentary deformation in a late Triassic (Rhaetian) lagoonal sequence, southwest Britain. *Geol. Mag.* 120, 613–622. doi:10.1017/S001675680002776X
- McCann, A.J., 2000. Deformation of the Old Red Sandstone of NW Spitsbergen; links to the Ellesmerian and Caledonian orogenies. *Geol. Soc. London, Spec. Publ.* 180, 567–584. doi:10.1144/GSL.SP.2000.180.01.30
- McClay, K.R., Ellis, P.G., 1987. Geometries of extensional fault systems developed in model experiments. *Geology* 15, 341–344. doi:10.1130/0091-7613(1987)15<341

- McClay, K.R., 1989. Analogue models of inversion tectonics. *Geol. Soc. London, Spec. Publ.* 44, 41–59. doi:10.1144/GSL.SP.1989.044.01.04
- McClay, K.R., 1990. Extensional fault systems in sedimentary basins: a review of analogue model studies. *Mar. Pet. Geol.* 7, 206–233. doi:10.1016/0264-8172(90)90001-W
- McClay, K.R., Scott, A.D., 1991. Experimental models of hangingwall deformation in ramp-flat listric extensional fault systems. *Tectonophysics* 188, 85–96. doi:10.1016/0040-1951(91)90316-K
- McClay, K.R., 1995. The geometries and kinematics of inverted fault systems: a review of analogue model studies. *Geol. Soc. London, Spec. Publ.* 88, 97–118. doi:10.1144/GSL.SP.1995.088.01.07
- McKenzie, D., 1978. Some remarks on the development of sedimentary basins. *Earth Planet. Sci. Lett.* 40, 25–32.
- McKerrow, W.S., Mac Niocaill, C., Dewey, J.F., 2000. The Caledonian Orogeny redefined. *J. Geol. Soc. London.* 157, 1149–1154. doi:10.1144/jgs.157.6.1149
- Meyer, V., Nicol, A., Childs, C., Walsh, J.J., Watterson, J., 2002. Progressive localisation of strain during the evolution of a normal fault population. *J. Struct. Geol.* 24, 1215–1231. doi:10.1016/S0191-8141(01)00104-3
- Miall, A.D., 1981. Sedimentation and tectonics in alluvial basins., *Geol.Assoc.Can.Spec.Pap.* Geological Association of Canada, Department of Earth Sciences, University of Waterloo. doi:10.1016/0012-8252(83)90024-7
- Midtkandal, I., Nystuen, J.P., Nagy, J., 2007. Paralic sedimentation on an epicontinental ramp shelf during a full cycle of relative sea-level fluctuation; the Helvetiafjellet Formation in Nordenskiöld Land, Spitsbergen. *Nor. J. Geol.* 87, 343–359.
- Minakov, A., Faleide, J.I., Glebovsky, V.Y., Mjelde, R., 2012. continental margin from integrated analysis of potential fields , 79–102. doi:10.1111/j.1365-246X.2011.05258.x
- Mørk, A., Knarud, R., Worsley, D., 1982. Depositional and diagenetic environments of the Triassic and lower Jurassic succession of Svalbard. *Arct. Geol. Geophys.* 8, 371–398.
- Mørk, A., Worsley, D., 2006. Triassic of Svalbard and the Barents shelf, NGF Abstracts and Proceedings, no. 3,.
- Mørk, M.B.E., 1999. Compositional variations and provenance of Triassic sandstones from the Barents Shelf. *J. Sediment. Res.* 69, 690–710. doi:10.2110/jsr.69.690
- Mørk, M.B.E., 2013. Diagenesis and quartz cement distribution of low-permeability Upper Triassic–Middle Jurassic reservoir sandstones, Longyearbyen CO2 lab well site in Svalbard, Norway. *Am. Assoc. Pet. Geol. Bull.* 97, 577–596. doi:10.1306/10031211193
- Morley, C.K., King, R., Hillis, R., Tingay, M., Backe, G., 2011. Deepwater fold and thrust belt classification, tectonics, structure and hydrocarbon prospectivity: A review. *Earth-Science Rev.* 104, 41–91.

- Mouslopoulou, V., Walsh, J.J., Nicol, A., 2009. Fault displacement rates on a range of timescales. *Earth Planet. Sci. Lett.* 278, 186–197. doi:10.1016/j.epsl.2008.11.031
- Mulrooney, M.J., 2011. Fault Segmentation & Linkage Analysis of the Exmouth Sub-basin, NW Shelf, Australia. MSc Thesis, Department of Earth Sciences, Royal Holloway, University of London.
- Mulrooney, M.J., Leutscher, J., Braathen, A., 2017. A 3D structural analysis of the Goliat field, Barents Sea, Norway. *Mar. Pet. Geol.* 86, 192–212. doi:10.1016/j.marpetgeo.2017.05.038
- Nejbert, K., Krajewski, K.P., Dubińska, E., Pécskay, Z., 2011. Dolerites of Svalbard, north-west Barents Sea Shelf: age, tectonic setting and significance for geotectonic interpretation of the High-Arctic Large Igneous Province. *Polar Res. Vol 30 incl Suppl. 1*, 1–24. doi:10.3402/polar.v30i0.7306
- Nemec, W., Steel, R.J., 1988. What is a fan delta and how do we recognize it? Fan deltas; *Sedimentol. Tecton. settings.* 3–13. doi:10.1111/j.1365-2117.1988.tb00021.x
- Nemec, W., Postma, G., 1993. Quaternary alluvial fans in southwestern Crete: sedimentation processes and geomorphic evolution, in: *Alluvial Sedimentation*. IAS Special Publication, pp. 235–276.
- Nicol, A., Walsh, J.J., Villamor, P., Seebeck, H., Berryman, K.R., 2010. Normal fault interactions, paleoearthquakes and growth in an active rift. *J. Struct. Geol.* 32, 1101–1113. doi:10.1016/j.jsg.2010.06.018
- Nironen, M., 1997. The Svecofennian Orogen: a tectonic model. *Precambrian Res.* 86, 21–44. doi:10.1016/S0301-9268(97)00039-9
- Norwegian Petroleum Directorate Resource Report for Fields and Discoveries, 2017.
- Nøttvedt, A., Cecchi, M., Gjelberg, J.G., Kristensen, S.E., Lønøy, A., Rasmussen, A., Rasmussen, E., Skott, P.H., Van Veen, P.M., 1993. Svalbard-Barents Sea correlation: a short review. *Arct. Geol. Pet. Potential, Nor. Pet. Soc. (NPF), Spec. Publ. 2*, 363–375.
- Ogata, K., Senger, K., Braathen, A., Tveranger, J., Olaussen, S., 2012. Geological Society , London , Special Publications Online First The importance of natural fractures in a tight reservoir for potential CO₂ storage : a case study of the upper Triassic – middle Jurassic Kapp Toscana Group (Spitsbergen , Arctic Norway) T. doi:10.1144/SP374.9
- Ogata, K., Senger, K., Braathen, A., Tveranger, J., Olaussen, S., 2014. Fracture systems and mesoscale structural patterns in the siliciclastic Mesozoic reservoir-caprock succession of the Longyearbyen CO₂ Lab project : Implications for geological CO₂ sequestration in Central Spitsbergen , Svalbard. *Nor. J. Geol.* 4, 121–154.
- Ohta, Y., 1982. Hecla Hoek Rocks in Central and Western Nordaustlandet, SKRIFTER N. ed, Norskpolarinstitut skrifter. Oslo.
- Olaussen, S., Dalland, A., Gloppen, T.G., Johannessen, E., 1984. Depositional environment and diagenesis of Jurassic reservoir sandstones in the eastern part of Troms I area, in: Spencer, A.M. (Ed.), *Petroleum Geology of the North European Margin: Proceedings of the North European Margin Symposium (NEMS '83)*, Organized by the Norwegian Petroleum Society and

- Held at the Norwegian Institute of Technology (NTH) in Trondheim 9--11 May, 1983. Springer Netherlands, Dordrecht, pp. 61–79. doi:10.1007/978-94-009-5626-1_6
- Olaussen, S. Galimberti, R.F., Grindhaug, J.K. Heskestad, B. , Leutscher, J. Johnsen, E. Johnsen, S., Opsahl, E., Seldal, J. and Stensland, D., 2010. Hydrocarbon pools banked to the basin margin faults: harvesting a successful play model in the south-western Barents Sea, in: 29th Nordic Geological Winter Meeting. Abstracts and Proceedings of the Geological Society of Norway. Pp 139 -140.
- Olovyanishnikov, V.G., Roberts, D., Siedlecka, A., 2000. Tectonics and sedimentation of the meso- to neoproterozoic Timan-Varanger Belt along the northeastern margin of Baltica. *Polarforschung* 68, 267–274.
- Osmundsen, P.T., Braathen, A., Rød, R.S., Hynne, I.B., 2014. Styles of normal faulting and fault-controlled sedimentation in the Triassic deposits of Eastern Svalbard. *Nor. Pet. Dir. Bull.* 11, 61–79.
- Ostanin, I., Anka, Z., di Primio, R., Bernal, A., 2012. Identification of a large Upper Cretaceous polygonal fault network in the Hammerfest basin: Implications on the reactivation of regional faulting and gas leakage dynamics, SW Barents Sea. *Mar. Geol.* 332–334, 109–125. doi:10.1016/j.margeo.2012.03.005
- Ostanin, I., Anka, Z., di Primio, R., Bernal, A., 2013. Hydrocarbon plumbing systems above the Snøhvit gas field: Structural control and implications for thermogenic methane leakage in the Hammerfest Basin, SW Barents Sea. *Mar. Pet. Geol.* 43, 127–146. doi:10.1016/j.marpetgeo.2013.02.012
- Øvrebø, O., Talleraas, E., 1977. The Structural geology of the Troms Area (Barents-Sea). *GeoJournal* 1, 47–54. doi:10.1007/BF00189603
- Parker, J.R., 1966. Folding, faulting and dolerite intrusions in the Mesozoic rocks of the fault zone of central Spitsbergen. *Norsk Polarinstitutt Årbok* 1964, 47–55.
- Paul, J., Wemmer, K., Wetzell, F., 2009. Keuper (Late Triassic) sediments in Germany - indicators of rapid uplift of Caledonian rocks in southern Norway. *Nor. Geol. Tidsskr.* 89, 193–202.
- Pavlidis, Y.A., Polyakova, E.I., 1997. Late Pleistocene and Holocene depositional environments and paleoceanography of the Barents Sea: evidence from seismic and biostratigraphic data. *Mar. Geol.* 143, 189–205.
- Peacock, D.C.P., 1991. Displacements and segment linkage in strike-slip fault zones. *J. Struct. Geol.* 13, 1025–1035. doi:10.1016/0191-8141(91)90054-M
- Peacock, D.C.P., Sanderson, D.J., 1991. Displacements, segment linkage and relay ramps in normal fault zones. *J. Struct. Geol.* 13, 721–733. doi:10.1016/0191-8141(91)90033-F
- Pease, V., Scott, R.A., 2009. Crustal affinities in the Arctic Uralides, northern Russia: significance of detrital zircon ages from Neoproterozoic and Palaeozoic sediments in Novaya Zemlya and Taimyr. *J. Geol. Soc. London.* 166, 517–527. doi:10.1144/0016-76492008-093

- Pease, V., 2011. Eurasian orogens and Arctic tectonics: an overview. *Geol. Soc. London, Mem.* 35, 311–324. doi:10.1144/M35.20
- Piepjohn, K., 2000. The Svalbardian-Ellesmerian deformation of the Old Red Sandstone and the pre-Devonian basement in NW Spitsbergen (Svalbard). *Geol. Soc. London, Spec. Publ.* 180, 585–601. doi:10.1144/GSL.SP.2000.180.01.31
- Polteau, S., Hendriks, B.W.H., Planke, S., Ganerød, M., Corfu, F., Faleide, J.I., Midtkandal, I., Svensen, H.S., Myklebust, R., 2016. The Early Cretaceous Barents Sea Sill Complex: Distribution, $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology, and implications for carbon gas formation. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 441, 83–95. doi:10.1016/j.palaeo.2015.07.007
- Postma, G., 1990. Depositional Architecture and Facies of River and Fan Deltas: A Synthesis In: Colella, A., Prior, D.B., Eds., *Coarse-Grained Deltas*, Vol. 10, Blackwell Publishing Ltd., Oxford, 13–28. doi:10.1002/9781444303858.ch2
- Pratt, L.M., Arthur, M.A., Dean, W.E., Scholle, P.A., 1993. Paleo-oceanographic cycles and events during the Late Cretaceous in the Western Interior Seaway of North America, in: *Evolution of the Western Interior Basin*. *Geol. Ass. Canada, Spec. Pap.*, pp. 333–354.
- Prosser, S., 1993. Rift-related linked depositional systems and their seismic expression. *Geol. Soc. London, Spec. Publ.* 71, 35–66. doi:10.1144/GSL.SP.1993.071.01.03
- Puchkov, V.N., 2009. The diachronous (step-wise) arc-continent collision in the Urals. *Tectonophysics* 479, 175–184. doi:10.1016/j.tecto.2009.01.014
- Rawling, G.C., Goodwin, L.B., 2006. Structural record of the mechanical evolution of mixed zones in faulted poorly lithified sediments, Rio Grande rift, New Mexico, USA. *Journal of Structural Geology* 28, 1623–1639.
- Rickard, M.J., Belbin, L., 1980. A new continental assembly for Pangaea. *Tectonophysics* 63, 1–12. doi:10.1016/0040-1951(80)90104-3
- Riis, F., Fjeldskaar, W., 1992. On the magnitude of the Late Tertiary and Quaternary erosion and its significance for the uplift of Scandinavia and the Barents Sea. *Norwegian Petroleum Society Special Publications* 1, 163–185. doi:10.1016/B978-0-444-88607-1.50016-4
- Riis, F., Lundschie, B. A., Høy, T., Mørk, A., Mørk, M.B.E., 2008. Evolution of the Triassic shelf in the northern Barents Sea region. *Polar Res.* 27, 318–338. doi:10.1111/j.1751-8369.2008.00086.x
- Rippon, J.H., 1985. Contoured patterns of the throw and hade of normal faults in the Coal Measures (Westphalian) of north-east Derbyshire, in: *Proceedings of the Yorkshire Geological Society. Geological Society of London*, pp. 147–161. doi:10.1144/pygs.45.3.147
- Rismyhr, B., 2012. Sedimentological Facies Models of the Lower Part of the Realgrunnen Subgroup in the Goliath Field Compared with Outcrops, MSc Thesis. Department of Earth Science, University of Bergen.
- Ritzmann, O., Faleide, J.I., 2007. Caledonian basement of the western Barents Sea. *Tectonics* 26, 1–20. doi:10.1029/2006TC002059

- Roberts, D., Siedlecka, A., 2002. Timanian orogenic deformation along the northeastern margin of Baltica, Northwest Russia and Northeast Norway, and Avalonian-Cadomian connections. *Tectonophysics* 352, 169–184. doi:10.1016/S0040-1951(02)00195-6
- Rønnevik, H., Beskow, B., Jacobsen, H.P., 1982. Evolution of the Barents Sea. In Embry, A.F. & Balkwill, H.R. (eds.): *Arct. Geol. Geophys.* 432–440. *Can. Soc. Pet. Geologists, Memoir* 8.
- Rønnevik, H.C., Jacobsen, H.P., 1984. Structural highs and basins in the western Barents Sea. In: Spencer, A.M. (ed.) *Petroleum geology of the north European margin*, in: Spencer, A.M. (Ed.), *Norsk Petroleum Forening, London, Graham and Trotman. Springer Netherlands, Dordrecht*, pp. 19–32. doi:10.1007/978-94-009-5626-1_3
- Rowan, M.G., Kligfield, R., 1989. Cross section restoration and balancing as aid to seismic interpretation in extensional terranes. *AAPG bulletin* 73, 955–966.
- Royse, F., 1993. Case of the phantom foredeep: early Cretaceous in west-central Utah. *Geology* 21, 133–136. doi:10.1130/0091-7613(1993)021<0133:COTPFE>2.3.CO;2
- Rust, N.R., 1979. Facies models 2: Coarse alluvial deposits. In: Walker, R.G. (Ed.), *Facies Models. Geoscience Canada Reprint Series* 1, p. 9-23.
- Ryer, T.A., 1981. Deltaic coals of Ferron Sandstone Member of Mancos Shale: predictive model for Cretaceous coal-bearing strata of Western Interior. *Am. Assoc. Pet. Geol. Bull.* 65, 2323–2340.
- Ryer, T.A., Anderson, P.B., 2004. *AAPG Studies in Geology* No. 50, *The Ferron Sandstone: Overview and Reservoir Analog*, Chapter 3: Facies of the Ferron Sandstone, East-Central Utah.
- Ryseth, A., 2014. Sedimentation at the Jurassic-Triassic boundary, south-west Barents Sea: Indication of climate change. *From Depos. Syst. to Sediment. Successions Nor. Cont. Margin* 46, 187–214. doi:10.1002/9781118920435.ch9
- Sarnthein, M., Van Kreveld, S., Erlenkeuser, H., Grootes, P.M., Kucera, M., Pflauman, U., Schulz, M., 2003. Centennial-to-millennial-scale periodicities of Holocene climate and sediment injections off the western Barents shelf, 75 N. *Boreas* 32, 447–461. doi:10.1111/j.1502-3885.2003.tb01227.x
- Schlische, R.W., 1995. Geometry and origin of fault-related folds in extensional settings. *Am. Assoc. Pet. Geol. Bull.* 79, 1661–1678. doi:10.1306/7834DE4A-1721-11D7-8645000102C1865D
- Schöpfer, M.P.J., Childs, C., Walsh, J.J., 2006. Localisation of normal faults in multilayer sequences. *J. Struct. Geol.* 28, 816–833. doi:10.1016/j.jsg.2006.02.003
- Schueller, S., Braathen, A., Fossen, H., Tveranger, J., 2013. Spatial distribution of deformation bands in damage zones of extensional faults in porous sandstones: Statistical analysis of field data. *Journal of Structural Geology* 52, 148–162.
- Schultz, R.A., Fossen, H., 2002. Displacement-length scaling in three dimensions: The importance of aspect ratio and application to deformation bands. *J. Struct. Geol.* 24, 1389–1411. doi:10.1016/S0191-8141(01)00146-8

- Schultz, R.A., Fossen, H., 2008. Terminology for structural discontinuities. *Am. Assoc. Pet. Geol. Bull.* 92, 853–867. doi:10.1306/02200807065
- Senger, K., Roy, S., Braathen, A., Buckley, S.J., Bælum, K., Gernigon, L., Mjelde, R., Noormets, R., Ogata, K., Olaussen, S., Planke, S., Ruud, B.O., Tveranger, J., 2013. Geometries of doleritic intrusions in central Spitsbergen, Svalbard: an integrated study of an onshore-offshore magmatic province with implications for CO₂ sequestration 143–166.
- Senger, K., Planke, S., Polteau, S., Ogata, K., Svensen, H., Senger, K., Senger, K., Planke, S., Svensen, H., 2014. Sill emplacement and contact metamorphism in a siliciclastic reservoir on Svalbard, Arctic Norway. *Nor. Geol. Tidsskr.* 94, 155–169.
- Serck, C.S., Faleide, J.I., Braathen, A., Kjølhamar, B., Escalona, A., 2017. Jurassic to Early Cretaceous basin configuration(s) in the Fingerdjupet Subbasin, SW Barents Sea. *Mar. Pet. Geol.* doi:10.1016/j.marpetgeo.2017.06.044
- Sharp, I.R., Gawthorpe, R.L., Underhill, J.R., Gupta, S., 2000. Fault-propagation folding in extensional settings: Examples of structural style and synrift sedimentary response from the Suez rift, Sinai, Egypt. *Bull. Geol. Soc. Am.* 112, 1877–1899. doi:10.1130/0016-7606(2000)112<1877:FPFIES>2.0.CO;2
- Shipton, Z.K., Cowie, P.A., 2001. Damage zone and slip-surface evolution over μm to km scales in high-porosity Navajo sandstone, Utah. *J. Struct. Geol.* 23, 1825–1844. doi:10.1016/S0191-8141(01)00035-9
- Singhal, B.B.S., Gupta, R.P., 2010. Applied hydrogeology of fractured rocks: Second edition. *Appl. Hydrogeol. Fract. Rocks Second Ed.* doi:10.1007/978-90-481-8799-7
- Smelror, M., Petrov, O. V., Larssen, G.B., Werner, S.C., Asov, V.A., Ebbing, J., Gernigon, L., Korchinskaya, M. V., Koren, T., Kosteva, N. V., Kotljar, G. V., Litvinova, T., Negrov, O.B., Olesen, O., Pascal, C., Pchelina, T.M., Petrov, Y.O., Sjulstad, H.-I., Sobolev, N.N., Vasiliev, V., 2009. Geological history of the Barents Sea, *Norges Geol. undersøkelse*. Geological Survey of Norway.
- Soliva, R., Benedicto, A., 2004. A linkage criterion for segmented normal faults. *J. Struct. Geol.* 26, 2251–2267. doi:10.1016/j.jsg.2004.06.008
- Soliva, R., Benedicto, A., 2005. Geometry, scaling relations and spacing of vertically restricted normal faults. *J. Struct. Geol.* 27, 317–325. doi:10.1016/j.jsg.2004.08.010
- Souche, A., Beyssac, O., Andersen, T.B., 2012. Thermal structure of supra-detachment basins: a case study of the Devonian basins of western Norway. *J. Geol. Soc. London.* 169, 427–434. doi:10.1144/0016-76492011-155
- Steel, R.J., Worsley, D., 1984. Svalbard's post-Caledonian strata - an atlas of sedimentational patterns and palaeogeographic evolution, in: Spencer, A.M. (Ed.), *Petroleum Geology of North European Margin*. Springer Netherlands, pp. 109–135. doi:10.1007/978-94-009-5626-1_9
- Stemmerik L., W.D., 1995. Permian History of the Barents Shelf Area. In: Scholle P.A., Peryt T.M., Ulmer-Scholle D.S. (eds) *The Permian of Northern Pangea*. Springer, Berlin, Heidelberg. doi:https://doi.org/10.1007/978-3-642-78590-0_5

- Sund, 1984. Tectonic Development and Hydrocarbon Potential Offshore Troms, Northern Norway: ABSTRACT. *Am. Assoc. Pet. Geol. Bull.* 68, 7–8. doi:10.1306/AD46178C-16F7-11D7-8645000102C1865D
- Talwani, M., Eldholm, O., 1977. Evolution of the Norwegian-Greenland Sea. *Geol. Soc. Am. Bull.* 88, 969–999.
- Taylor, A.M., Goldring, R., 1993. Description and analysis of bioturbation and ichnofabric. *J. Geol. Soc. London.* 150, 141–148. doi:10.1144/gsjgs.150.1.0141
- Taylor, K.G., Gawthorpe, R.L., Curtis, C.D., Marshall, J.D., Awwiller, D.N., 2000. Carbonate cementation in a sequence-stratigraphic framework: Upper Cretaceous sandstones, Book Cliffs, Utah-Colorado. *J. Sediment. Res.* 70, 360–372.
- Torrence, C., Compo, G.P., 1998. A Practical Guide to Wavelet Analysis. *Bull. Am. Meteorol. Soc.* 79, 61–78. doi:10.1175/1520-0477(1998)079<0061:APGTWA>2.0.CO;2
- Torsvik, T.H., Andersen, T.B., 2002. The Taimyr fold belt, Arctic Siberia: Timing of prefold remagnetisation and regional tectonics. *Tectonophysics* 352, 335–348. doi:10.1016/S0040-1951(02)00274-3
- Trudgill, B., Cartwright, J., 1994. Relay-ramp forms and normal-fault linkages, Canyonlands National Park, Utah. *Geol. Soc. Am. Bull.* 106, 1143–1157. doi:10.1130/0016-7606(1994)106<1143:RRFANF>2.3.CO;2
- Våagnes, E., Gabrielsen, R.H., Haremo, P., 1998. Late Cretaceous-Cenozoic intraplate contractional deformation at the Norwegian continental shelf: Timing, magnitude and regional implications. *Tectonophysics* 300, 29–46. doi:10.1016/S0040-1951(98)00232-7
- Van Stappen, J., Kock, T. De, Boone, M.A., Olaussen, S., Cnudde, V., 2014. Pore-scale characterisation and modelling of CO₂ flow in tight sandstones using X-ray micro-CT ; Knorringfjellet Formation of the Longyearbyen CO₂ Lab , Svalbard.
- Vorren, T.O., Richardsen, G., Knutsen, S.M., Henriksen, E., 1991. Cenozoic erosion and sedimentation in the western Barents Sea. *Marine and Petroleum Geology* 8, 317–340. doi:10.1016/0264-8172(91)90086-G
- Walsh, J.J., Watterson, J., 1988. Analysis of relationship between displacement and dimensions of faults. *J. Struct. Geol.* 10, 239–247.
- Walsh, J.J., Watterson, J., 1991. Geometric and kinematic coherence and scale effects in normal fault systems. *Geol. Soc. London, Spec. Publ.* 56, 193–203. doi:10.1144/GSL.SP.1991.056.01.13
- Walsh, J.J., Nicol, A., Childs, C., 2002. An alternative model for the growth of faults. *J. Struct. Geol.* 24, 1669–1675. doi:10.1016/S0191-8141(01)00165-1
- Walsh, J.J., Bailey, W.R., Childs, C., Nicol, A., Bonson, C.G., 2003. Formation of segmented normal faults: A 3-D perspective. *J. Struct. Geol.* 25, 1251–1262. doi:10.1016/S0191-8141(02)00161-X
- Watterson, J., 1986. Fault Dimension, Displacements and Growth. *Pure Appl. Geophys.* 124, 365–373.

- Wernicke, B., Axen, G.J., 1988. On the role of isostasy in the evolution of normal-fault systems. *Geology* 16, 848–851. doi:10.1130/0091-7613(1988)016<0848
- Wheeler, J., 1987. Variable-heave models of deformation above listric normal faults: the importance of area conservation. *J. Struct. Geol.* 9, 1047–1049. doi:10.1016/0191-8141(87)90013-7
- Wignall, P.B., Best, J.L., 2000. The western Irish Namurian basin reassessed. *Basin Research* 12, 59–78.
- Wilkins, S.J., Gross, M.R., 2002. Normal fault growth in layered rocks at Split Mountain, Utah: Influence of mechanical stratigraphy on dip linkage, fault restriction and fault scaling. *J. Struct. Geol.* 24, 1413–1429. doi:10.1016/S0191-8141(01)00154-7
- Willemse, E.J.M., Pollard, D.D., Aydin, A., 1996. Three-dimensional analyses of slip distributions on normal fault arrays with consequences for fault scaling. *Journal of Structural Geology* 18, 295–309. doi:10.1016/S0191-8141(96)80051-4
- Williams, G., Vann, I., 1987. The geometry of listric normal faults and deformation in their hangingwalls. *J. Struct. Geol.* 9, 789–795. doi:10.1016/0191-8141(87)90080-0
- Williams, G.D., Powell, C.M., Cooper, M.A., 1989. Geometry and kinematics of inversion tectonics. *Geol. Soc. London, Spec. Publ.* 44, 3–15. doi:10.1144/GSL.SP.1989.044.01.02
- Willsey, S.P., Umhoefer, P.J., Hilley, G.E., 2002. Early evolution of an extensional monocline by a propagating normal fault: 3D analysis from combined field study and numerical modeling. *J. Struct. Geol.* 24, 651–669. doi:10.1016/S0191-8141(01)00120-1
- Wilson, J.T., 1966. Some rules for continental drift. *R. Soc. Canada Spec. Publ.* 9, 1–17.
- Winsnes, T.S., Worsley, D., 1981. Geological Map of Svalbard 1: 500,000: Sheet 2G, Edgeøya. Norsk Polarinstitut.
- Withjack, M.O., Olson, J., Peterson, E., 1990. Experimental models of extensional forced folds. *Am. Assoc. Pet. Geol. Bull.* 74, 1038–1054. doi:10.1306/OC9B23FD-1710-11D7-8645000102C1865D
- Wolstencroft, M., Shen, Z., Törnqvist, T.E., Milne, G.A., Kulp, M., 2014. Understanding subsidence in the Mississippi Delta region due to sediment, ice, and ocean loading: Insights from geophysical modeling. *Journal of Geophysical Research: Solid Earth* 119, 3838–3856.
- Worsley, D., 1973. The Wilhelmøya Formation—a new lithostratigraphical unit from the Mesozoic of eastern Svalbard. *Arb. Nor. Polarinst* 1971, 7–16.
- Worsley, D., Agdestein, T., Gjelberg, J.G., Kirkemo, K., Mørk, A., Nilsson, I., Olaussen, S., Steel, R.J., Stemmerik, L., 2001. The geological evolution of Bjørnøya, Arctic Norway: Implications for the Barents Shelf. *Nor. Geol. Tidsskr.* 81, 195–234.
- Worsley, D., 2008. The post-Caledonian development of Svalbard and the western Barents Sea. *Polar Res.* 27, 298–317. doi:10.1111/j.1751-8369.2008.00085.x

- Yielding, G., 1990. Footwall uplift associated with Late Jurassic normal faulting in the northern North Sea. *J. Geol. Soc. London*. 147, 219–222. doi:10.1144/gsjgs.147.2.0219
- Young, M.J., Gawthorpe, R.L., Sharp, I.R., 2002. Architecture and evolution of syn-rift clastic depositional systems towards the tip of a major fault segment, Suez rift, Egypt. *Basin Res.* 14, 1–23. doi:10.1046/j.1365-2117.2002.00162.x
- Ziegler, P.A., 1982. Triassic rifts and facies pattern in Western and Central Europe. *Int. J. Earth Sci.* 71, 747–772.
- Ziegler, P.A., 1987. Late Cretaceous and Cenozoic intra-plate compressional deformations in the Alpine foreland—a geodynamic model. *Tectonophysics* 137, 389–420.
- Ziegler, P.A., 1988. Post-Hercynian plate reorganization in the Tethys and Arctic-North Atlantic domains. *Dev. Geotecton.* 22, 711–755. doi:10.1016/B978-0-444-42903-2.50035-X
- Ziegler, P.A., Cloetingh, S., van Wees, J.D., 1995. Dynamics of intra-plate compressional deformation: the Alpine foreland and other examples. *Tectonophysics* 252, 7–59. doi:10.1016/0040-1951(95)00102-6
- Zonenshain, L.P., 1990. Geology of the USSR: a plate-tectonic synthesis. *Geodyn. Ser.* 21, 120p.

9. ARTICLES

9.1 ARTICLE I

Mulrooney, M.J., Leutscher, J., Braathen, A.

A 3D structural analysis of the Goliat field, Barents Sea, Norway

Published in Marine and Petroleum Geology Volume 86, September 2017, Pages 192-212

doi:10.1016/j.marpetgeo.2017.05.038

9.2 ARTICLE II

Mulrooney, M.J., Rismyhr, R., Yenwongfai, H. D., Leutscher, J.,
Olaussen, S., Braathen, A.

***Impacts of small-scale faults on continental to coastal plain
deposition: Evidence from the Realgrunnen Subgroup in the Goliat
field, Southwest Barents Sea, Norway***

Marine and Petroleum Geology (In press)

9.3 ARTICLE III

Ogata, K., **Mulrooney, M.J.**, Braathen, A., Maher, H., Osmundsen, P.T.O., Anell, I., Balsamo, F.

Architecture, deformation style and petrophysical properties of a Late Triassic growth fault system in southern Edgeøya (East Svalbard)

Basin Research (In press)

9.4 ARTICLE IV

Mulrooney, M.J., Van Stappen, J., Cnudde, V., Senger, K., Rismyhr, B., Braathen, A., Olaussen, S., Mørk, M.B.E., Ogata, K.

Fluid flow properties of a potential unconventional CO₂ storage unit in central Spitsbergen: the Upper Triassic to Middle Jurassic Wilhelmøya Subgroup

Submitted to Norwegian Journal of Geology on 10 October, 2017

10. APPENDICES

10.1 APPENDIX 1

Supplementary Article 1:

Braathen, A., Midtkandal, I., Mulrooney, M.J., Appleyard, T.R., Haile, B., Van Yperen, A.E.

Growth-faults from delta collapse – structural and sedimentological investigation of the Last Chance delta, Ferron Sandstone, Utah.

Published in Basin Research 1–20, <https://doi.org/10.1111/bre.12271>

10.2 APPENDIX 2**Supplementary Article 2:**

Rismyhr, B., Bjærke, T., Olausen, S., Mulrooney, M.J., Senger, K.

Facies, palynostratigraphy and sequence stratigraphy of the Wilhelmøya Subgroup (Upper Triassic–Middle Jurassic) in western central Spitsbergen, Svalbard

Norwegian Journal of Geology (In press)

10.3 APPENDIX 3

Supplementary Article 3:

Maher, H., Senger, K., Ogata, K., Braathen, A., Mulrooney, M. J., Smyrak-Sikora, A., Osmundsen, P.T.

Mesozoic-Cenozoic regional stress field evolution in Svalbard

Submitted to Tectonics on 15 August 2017

10.4 APPENDIX 4

Conference Proceedings 1:

Senger, K., Mulrooney, M., Schaaf, N., Braathen, A., Ogata, K., Olaussen, S.

Characterization and modelling of a naturally fractured reservoir-caprock unit targeted for CO₂ storage in Arctic Norway

Conference proceedings.

Fourth EAGE Workshop on Rock Physics 11 - 13 November 2017, Abu Dhabi

Accepted for oral presentation

10.5 APPENDIX 5

Conference abstract 2:

Mulrooney, M.J., Leutscher, J. and Braathen, A.

Structural Architecture of the Goliat Field: Application of a high resolution multi-azimuth 3D seismic survey in the southern Hammerfest Basin of the Barents Shelf, Norway

Conference proceedings.

Vinterkonferansen 2017, Oslo, January 9th-11th, 2017.

ISBN: 978-82-8347-018-5

Accepted for poster presentation

10.6 APPENDIX 6

Conference abstract 3:

Mulrooney, M.J., & Senger, S.

Automated mapping of discontinuities within Cretaceous dolerite sills, central Spitsbergen, Arctic Norway

Conference proceedings.

2nd Virtual Geoscience Conference, Bergen, Norway, 22-23 September 2016

ISBN: 978-82-8361-005-5

Accepted for poster presentation

10.7 APPENDIX 7**Conference abstract 4:**

Harmon Jr., H., Senger, K., Ogata, K., Braathen, A., Mulrooney, M., Smyrak-Sikora, A., Osmundson, P. T.O.

Mesozoic Regional Stress Field Evolution in Svalbard

Conference proceedings.

GSA Annual Meeting in Denver, Colorado, USA – 27 September 2016

doi: 10.1130/abs/2016AM-281384

Accepted for oral presentation

10.8 APPENDIX 8

Conference abstract 5:

Senger, K, Mulrooney, M., Braathen, A., Ogata, K., and Olausson, S.

Integrated Characterization of an Organic-rich Caprock Shale, Svalbard, Arctic Norway

Conference proceedings.

Fifth EAGE Shale Workshop 2-4 May 2016, Catania, Italy

DOI: 10.3997/2214-4609.201600430

Accepted for oral presentation

10.9 APPENDIX 9**Conference abstract 6:**

Mulrooney, M. and Braathen, A.

Outcrop Scale Normal Faults affecting the Longyearbyen CO₂ Reservoir, Svalbard

Conference proceedings.

31st Geological Winter Meeting, Stavanger, January 12-14 2015

ISBN 978-82-92-39497-7

Accepted for poster presentation

10.10 APPENDIX 10**Conference abstract 7:**

Mulrooney, M., Braathen, A. and Leutscher, J.

Evidence for late Triassic Tectonism affecting the Goliat Field, South Hammerfest Basin, SW Barents Shelf

Conference proceedings.

31st Geological Winter Meeting, Stavanger, January 12-14 2015

ISBN 978-82-92-39497-7

Accepted for oral presentation

10.11 APPENDIX 11**Internal ENI Norge Report (Unpublished)**

Mulrooney, M.

Structural Architecture and Kinematics of the Goliat Field, Southern Hammerfest Basin.

The following abstract is derived from a 115 page report on the structural geology of the Goliat field for ENI Norge which described a detailed seismic investigation of the EN0901 multi-azimuth seismic survey. The work is unpublished and subject to a confidentiality agreement but formed the basis of articles 1 and 2 herein.

Unpublished internal ENI Norge Report