Deciphering molecular interactions of AQP4 at the gliovascular interface in brain

© Eystein Hellstrøm Hoddevik, 2019

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-430-6

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

Table of contents

Acknowledgements	3
Abstract	4
List of abbreviations	5
List of articles included in the thesis	6
Other articles published during the Ph.D. period	7
The candidate's contributions	8
1. Introduction	9
2. Background and definitions	10
3. Hypotheses and aims of the thesis	22
4. Methodological considerations	23
5. Summary of individual articles	28
6. Discussion	30
7. Conclusions and future perspectives	41
8. References	41

Acknowledgements

The work presented in this thesis was carried out at the Department of Anatomy, Institute of Basic Medical Science, University of Oslo. Financial support was received jointly from the Centre for Molecular Biology and Neuroscience (CMBN) and the Institute. I am grateful to the University of Oslo for funding my Ph.D. project. I also thank the Department of Pathology and colleagues at the Oslo University Hospital for generously accommodating my need to continue my scientific work during my training as a registrar.

First and foremost, I wish to express my sincere gratitude to Professor Mahmood Reza Amiry-Moghaddam for his dedication and persistence throughout my Ph.D. period and for excellent supervision. I look forward to our continued and future scientific collaborations. I also want to convey special thanks to Professor Ole Petter Ottersen whose help and support has been invaluable, in the very meaning of that word. I thank my co-supervisor Professor Vidar Gundersen, Professor emeritus Per Brodal and Professor Erlend Nagelhus for their helpful and frank advice in all matters. I have had the pleasure of befriending and doing science with wonderful colleagues and fellow scientists. You have all been a source of inspiration and support, and for this I am very grateful. I therefore thank Shreyas Rao, Henning Bünsow Boldt, Laura Maria Azzurra Camassa, Lisa Lunde, Faraz Hameed Khan, Soulmaz Rahmani, Shirin Katoozi, Agnete Prytz, Gry Helen Enger Syverstad and Katja Stahl. I extend particular thanks to Tom Tallak Solbu for sharing his extensive laboratory knowledge and thus helping me refine all the methods I used in my Ph.D. I thank Torgeir Holen for countless motivating discussions and for sharing his insight into experimental work. I would not have been able to complete my work without the help of laboratory engineers Karen Marie Gujord, Jorunn Knutsen and in particular Bjørg Riber. The same applies to Bashir Hakim, whose abilities and dedication always ensured that the electron microscopes worked flawlessly, and to Gunnar Lothe and Carina Knudsen who helped with the visual presentation of my findings.

My greatest discovery from my first year at the institute was the daily, anatomical coffee break at 9:30 AM where no problems were ever too big or too small for plenary discussions. I want to thank all regular attendants for their impeccable sense of humour, pragmatism and ability to shed light upon my scientific path when all seemed lost. In addition to previously mentioned colleagues I therefore also thank Reidun Torp, Finn-Mogens Smedja Haug, Sigbjørn Fossum, Wendi Jensen, Annabjörg Bore, Grazyna Babinska and posthumously Professors Eric Rinvik and Bent Rolstad.

I thank my parents, Gunnar Martin Hoddevik and Grethe Hellstrøm Hoddevik, for always being a source of inspiration and for all the help they have given me since I embarked upon this scientific endeavour.

I am forever indebted to my wonderful wife, Hilde Steinfeld, for her unwavering support. Together, we have surpassed my feeble, laboratory based efforts a total of four times. Thus, our joint and greatest successes in molecular biology are called Alma, Signe, Thorvald and Haakon. The inherent fearlessness with which our children explore the unknown is an attribute I envy them, that I wish they will never lose, and that will enable them to become excellent scientists should they ever wish to pursue such a path. On a continuous basis do they also remind me that the true cornerstone of all science is *curiosity*.

Oslo, 6th of November 2018

Eystein Hellstrøm Hoddevik

Abstract

The gliovascular interface is crucial for homeostasis of the central nervous system (CNS) and is perturbed in several neuropathological states with vascular involvement. At this site, astrocyte endfeet ensheath the vasculature completely and contain an abundance of the water channel aquaporin-4 (AQP4) in highly specialised membrane microdomains. Loss of this protein from endfoot membranes is found in several neurological conditions and the first study in the present work demonstrates how AQP4 loss precedes the development of chronic seizures in an animal model of temporal lobe epilepsy. For these reasons, deciphering the molecular mechanisms that regulate expression of AQP4 in astrocyte endfeet at the gliovascular interface stands out as a project with ramifications and direct consequences for several pathophysiological states. Both localisation and distribution of AQP4 has been shown to depend on intracellular as well as extracellular proteins. My thesis thus explores the hypothesis of whether extracellular matrix (ECM) molecules have an instructive and/or regulatory role on the composition of astrocyte membrane domains and on AQP4 levels in particular. Evidence is provided in favour of such a regulatory role for ECM proteins agrin and laminin as well as for the intracellular anchoring protein α -syntrophin (α -Syn). A case in point is the immediate surroundings of perivascular pericytes, where astrocyte proteins AQP4 and α -Syn as well as ECM proteins laminin and agrin co-exist and co-distribute. In the systematic approach towards examining the aforementioned mechanisms, we examined several different anatomical regions of rodent CNS. Evidence is therefore also provided that the gliovascular interface displays significant heterogeneity across the brain with regards to ECM proteins and astrocyte endfeet. The potential implications for pathophysiology and physiology is discussed and may in particular be of relevance for epilepsy, neuromyelitis optica (NMO) and waste clearance from CNS.

List of abbreviations

ALS	Acute lateral sclerosis
AD	Alzheimer's disease
α -Syn	α -syntrophin protein
α -Syn	α -syntrophin gene (<i>SNTA1</i>)
AQPs	aquaporin proteins
AQP4	aquaporin-4 protein
<i>Aqp4</i>	aquaporin-4 gene
<i>Aqp4</i> ^{-/-}	aquaporin-4 genetic knockout
BBB	blood-brain barrier
β -DG	beta-dystroglycan
BOLD fMRI	blood oxygen level-dependent functional magnetic resonance imaging
BL	basal lamina
BM	basement membrane
CNS	central nervous system
CSF	cerebrospinal fluid
DAG-1	dystroglycan associated gene- 1
DAPC	dystrophin associated protein complex
DGC	dystrophin glycoprotein complex
Dp-71	dystrophin (short isoform)
EBJC	endfoot-basal lamina junctional complex
ECM	extracellular matrix
ECS	extracellular space
EEA-1	early endosome antigen-1
GBS	Guillan-Barré syndrome
GFAP	glial fibrillary acidic protein
Kir 4.1	K ⁺ inwards rectifying channel 4.1
MCAO	middle cerebral artery occlusion
MCTs	monocarboxylate transporters
MIPs	membrane intrinsic proteins
MS	multiple sclerosis
NMO	neuromyelitis optica
NMOSDs	neuromyelitis optica spectrum disorders
OAPs	orthogonal arrays of particles
PC	pericyte coverage
PFA	paraformaldehyde
RT-qPCR	quantitative real-time polymerase chain-reaction
rCBF	regional cerebral blood flow
TJs	tight junctions, from latin <i>zonulae occludentes</i>
TLE	temporal lobe epilepsy

List of articles included in the thesis

This thesis is based on the following articles, which will be referred to by their roman numerals or as ordinary citations where appropriate:

- I** Alvestad S, Hammer J, **Hoddevik EH**, Skare Ø, Sonnewald U, Amiry-Moghaddam M, Ottersen OP. Mislocalization of AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy. *Epilepsy Research* 2013 Jul;105(1-2):30-41
- II** **Hoddevik EH**, Kahn FH, Rahmani S, Ottersen OP, Boldt HB, Amiry-Moghaddam M. Factors determining the density of AQP4 water channel molecules at the brain-blood interface. *Brain Struct Funct.* 2017 May;222(4):1753-1766. doi: 10.1007/s00429-016-1305-y. Epub 2016 Sep 15
- III** Camassa LM, Lunde LK, **Hoddevik EH**, Stensland M, Boldt HB, De Souza GA, Ottersen OP, Amiry-Moghaddam M. Mechanisms underlying AQP4 accumulation in astrocyte endfeet. *Glia.* 2015 Nov;63(11):2073-2091. doi: 10.1002/glia.22878. Epub 2015 Jun 29.
- IV** Lunde LK, Camassa LM, **Hoddevik EH**, Khan FH, Ottersen OP, Boldt HB, Amiry-Moghaddam M. Postnatal development of the molecular complex underlying astrocyte polarization. *Brain Struct Funct.* 2015 Jul;220(4):2087-101. doi: 10.1007/s00429-014-0775-z. Epub 2014 Apr 29.
- V** **Hoddevik EH**, Rahmani S, Rao S, Boldt HB, Ottersen OP, Amiry-Moghaddam M. Organisation of extracellular matrix proteins laminin and agrin in pericapillary basal laminae: insensitivity to targeted deletion of α -syntrophin. *Manuscript.*

Other articles published during the Ph.D. period

Heuser K, **Hoddevik EH**, Taubøll E, Gjerstad L, Indahl U, Kaczmarek L, Berg PR, Lien S, Nagelhus EA, Ottersen OP. Temporal lobe epilepsy and matrix metalloproteinase 9: a tempting relation but negative genetic association. *Seizure*. 2010 Jul;19(6):335-8. Epub 2010 Jun 3.

Amiry-Moghaddam M, **Hoddevik EH**, Ottersen OP. Aquaporins: multifarious rôles in brain. *Neuroscience*. 2010 Jul 28;168(4):859-61. Epub 2010 May 5.

Eilert-Olsen M, Haj-Yasein NN, Vindedal GF, Enger R, Gundersen GA, **Hoddevik EH**, Petersen PH, Haug FM, Skare Ø, Adams ME, Froehner SC, Burkhardt JM, Thoren AE, Nagelhus EA. Deletion of aquaporin-4 changes the perivascular glial protein scaffold without disrupting the brain endothelial barrier. *Glia*. 2012 Mar;60(3):432-40. doi: 10.1002/glia.22277. Epub 2011 Nov 30.

The candidate's contributions

Article I: Mislocalisation of AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy:

- Electron microscopy consisting of immunogold labelling of α -syntrophin and subsequent image acquisition, analysis and quantitation
- Edited the manuscript

Article II: Factors determining the density of AQP4 water channel molecules at the brain-blood interface.

- Designed study and experiments (except RT-qPCR).
- Performed all aspects of electron microscopy analysis (except AQP4 imaging and analysis), including all transcardial perfusions, CNS dissections, initial sectioning, immunogold labelling on ultrathin sections and subsequent image acquisition, analysis and quantitation
- Breeding and maintenance of *Aqp4*^{-/-} colony (control sections)
- Immunofluorescence and immunostains on thick sections
- All aspects of western blotting including homogenate preparation and casting of gels
- Prepared the manuscript

Article III: Mechanisms underlying AQP4 accumulation in astrocyte endfeet:

- Performed all aspects of electron microscopy, including embedding in durcupan resins, ultrathin serial sectioning and immunogold labelling of experimental and validating nature
- Breeding and maintenance of *Aqp4*^{-/-} colony
- Wrote parts of the manuscript, edited the manuscript

Article IV: Postnatal development of the molecular complex underlying astrocyte polarisation:

- Transcardial perfusions of mice from all milestone ages (P0 to adults) for thick section experiments
- CNS dissections
- Immunostains on thick sections
- Breeding and maintenance of *Aqp4*^{-/-} and β -DG ^{-/-} (*DAG1-flox*GFAP-cre*) colonies (control tissue)
- Antibody testing and optimisation (agrin and laminin) for light and electron microscopy
- Wrote parts of the manuscript, edited the manuscript

Article V: Organisation of extracellular matrix proteins laminin and agrin in pericapillary basal laminae: insensitivity to targeted deletion of alpha-syntrophin:

- Designed study and experiments (except RT-qPCR).
- Performed all aspects of electron microscopy analysis, including transcardial perfusions, CNS dissections, sectioning, immunogold labelling on ultrathin sections and subsequent image acquisition, analysis and quantitation
- Immunostains on thick sections
- Prepared the manuscript

1. Introduction

Neuroscience is the study of all facets of the human brain, animal brains and nervous tissue. The spectrum included therein is vast. Examples span from the unresolved issues of consciousness, how information is stored and processed to research into the pathogenesis of both rare and common brain diseases. My thesis is best viewed in the context of pathophysiology but also carries potential implications for neurophysiology. The included experiments are focused on the gliovascular interface in brain at the capillary level. Here, astrocytes ensheath the entire vasculature with their specialised projections, termed endfeet (Mathiisen et al. 2010). These star-shaped cells are thus uniquely positioned to link brain parenchyma to the circulation. Transport and signalling mechanisms at the gliovascular interface are critically involved in a number of brain functions (Iliff et al. 2012; Nagelhus and Ottersen 2013; Verkhratsky and Nedergaard 2018). Sustained interest has been devoted to the water channel AQP4, which is abundantly expressed in astrocyte endfoot membranes. The present study is a systematic approach towards deciphering how this abundant and highly polarised expression is induced and regulated. Targeted deletion of *Aqp4* protects against brain oedema formation (Amiry-Moghaddam et al. 2003a; Manley et al. 2000) and delays clearance of waste products from brain neuropil (Iliff et al. 2012). Loss of AQP4 from endfoot membranes is found in several neurological conditions (Eid et al. 2005; Frydenlund et al. 2006; Yang et al. 2011), but the mechanisms underlying this downregulation remain unclear, despite evidence of regulatory mechanisms and tempting hypotheses (Heuser et al. 2010a; Madrid et al. 2001). In the first work presented in this thesis, perivascular levels of AQP4 were examined in an animal model of temporal lobe epilepsy (TLE). This demonstrated that endfoot reduction of AQP4 levels precedes the development of chronic seizures (Alvestad et al. 2013) and correlates with reductions in the AQP4 anchoring protein α -syntrophin (α -Syn). The latter study addressed one of the most challenging questions in current research: whether changes observed are causes or consequences of neurological disease.

This initial study prompted further examination of the interdependence between AQP4 and α -Syn. The next article therefore makes use of a transgenic mouse model with targeted deletion of α -Syn and examines the *in situ* occurrence of both proteins. During this investigation, evidence of significant regional differences within the rodent brain became evident. Across all examined CNS regions, AQP4 and α -Syn are constitutively expressed around all blood vessels. Also, levels were significantly higher in the proximity of perivascular pericytes. For AQP4, this proved resistant to targeted deletion of α -Syn. These phenomena were quantified, and also constitute examples ideally suited to examine AQP4 regulation.

Interlaced between the aforementioned, cellular elements of the gliovascular interface is a distinct, perivascular basal lamina (BL). This is nothing short of an extraordinary assembly of extracellular matrix (ECM) constituents. It has long been known that acquired or inherited defects of ECM cause diseases in many organs. An illustrative example is Alport syndrome (Alport 1927), where a defect in the widely expressed collagen IV protein results in symptoms from kidney, ears and eyes. It is quite surprising, therefore, that few researchers have pursued the idea that similar 'matricopathies' might also cause brain disease. But evidence exists that ECM proteins influence astrocytes and AQP4 (Guadagno and Moukhles 2004; Noel et al. 2009; Noell et al. 2009; Tham et al. 2016). The third article is therefore a tailor made cell culture system that permits the examination of astrocyte endfeet in a systematic fashion. Here, evidence is provided that several factors impact on AQP4 levels, including co-culture of astrocytes with brain derived endothelial cells, and incubation with ECM proteins agrin and laminin. In the fourth article, therefore, the ontogenic profiles of all the aforementioned and implicated proteins were assessed. This included several proteins in the intracellular dystrophin-dystroglycan protein complex (Blake et al. 2002) which is a key organiser of endfoot proteins and architecture (Noell et al. 2011). Dystrophin anchors AQP4 through α -Syn and is coupled to the endothelial basal lamina through α - and β -dystroglycan (β -DG; (Blake et al. 2002; Culligan et al. 2001). These molecular interactions likely explain the highly specific

accumulation of AQP4 in perivascular endfoot membranes. The conclusions from this fourth study were that ECM proteins agrin and laminin have an inductive role in establishing the polarised expression of AQP4 in astrocyte endfeet. Finally, the fifth article examines agrin and laminin *in situ* in mouse CNS in both WT and α -Syn^{-/-} genotypes. These findings constitute the first quantitative assessment of agrin and pan-laminin at ultrastructural level. In accordance with the AQP4 and α -Syn data from article II (Hoddevik et al. 2017), agrin and laminin also display significant regional differences. Surprisingly, this is also the case at local, pericapillary level where the distribution of these proteins overlaps surprisingly well with variations of AQP4 and α -Syn. This is consistent with the idea that agrin and laminin both regulate AQP4 levels and provides a potential, mechanistic explanation of how AQP4 is enriched in astrocyte membrane domains abutting pericytes. Further, targeted deletion of α -Syn leaves laminin and agrin distribution and measured protein levels unaltered. This supports the hypothesis that α -Syn affects AQP4 redistribution via a direct effect rather than indirectly via ECM proteins agrin and laminin.

In the first part, the discussion details the provided evidence on induction and regulation of AQP4 in a context with other known mechanisms. Findings also indicate that the gliovascular interface displays significant regional heterogeneity. The second part of the discussion therefore addresses physiological implications thereof, including waste clearance from CNS via paravascular flow. Finally, the findings are discussed in the context of epilepsy and neuromyelitis optica (NMO). Mention is also made of select neuropathologies that may benefit from an understanding of AQP4 regulation.

Collectively, the experiments and articles touch on a broad range of topics. This limits the depth to which one can dive into them all, but makes the thesis relevant for an equally broad audience. An initial background chapter is therefore included to render the work more accessible to readers, irrespective of prior knowledge in the field.

2. Background and definitions

This chapter introduces the gliovascular interface and defines notions around which the ensuing parts of the thesis are structured. The cellular compartments that have been studied are introduced first: astrocytes, pericytes and endothelium. Relevant aspects of astrocytes are defined. These include astrocyte endfeet, astrocyte membrane proteins, dystrophin associated glycoprotein (DAP) complex and AQP4. ECM proteins and structure in CNS is then reviewed. Finally, mention is made of a demyelinating disease that is particularly relevant for AQP4 in the CNS. Other pathologies of interest are mentioned in the discussion.

Astrocytes

Ramon Y Cajal was among the first to employ Golgi staining on sections from the CNS. To date do his drawings remain impressive and provide more than historical data. Importantly, his observations were the first to highlight the true diversity of astrocytes. A mere 22 pages of his two-volume compendium “The histology of the central nervous system” are used to describe glial cells (Ramon y Cajal 1911). According to him, *la cellule neuroglie* is one of the two specialised elements constituting the nervous tissue – the other being “*la cellule nerveuse avec ses prolongements*”. Cajal highlights that neuroglial cells are also called “spider cells” due to their resemblance to this animal, or “corpuscles of Deiter” in honour of the scholar who - according to Cajal - first discovered them. The aforementioned 22 pages are truly exemplary in their scientific approach. Not only do they contain drawings of such quality that they remain useful and are still cited but the accompanying text also elaborates on the morphological findings and what they imply for cell function. In that regard, an exciting

hypothesis proposed by Andriezen (Andriezen 1893) is that astrocytes provide true cuffs for blood vessels with the ability and function to dampen or absorb physical pressure propagated along blood vessels. Of particular note to this intriguing hypothesis is a small, but interesting, finding in one of the articles included in the current thesis potentially regarding pressure exerted on endfeet (Camassa et al. 2015).

Astrocytes still retain the name that was proposed for them in 1893 by Michael von Lenhossek (Lenhossek 1893) for their stellate appearance in histological stains. Arguably, astrocytes represent the most diverse class of cells in the CNS. They perform homeostatic functions of the CNS at molecular and whole organ level. For further background on astrocyte physiology, the reader is referred to the recent and thorough review by Verkhratsky and Nedergaard (Verkhratsky and Nedergaard 2018). In another review on functions of mature, mammalian astrocytes, Harold Kimelberg recapitulates those eight criteria that hitherto have been used to recognise a mature astrocyte (Kimelberg 2010), namely:

1. Nonexcitability; the absence of electrical excitability
2. A very negative membrane potential (-80 to -90 mV) determined by the transmembrane K^+ gradient (because of a prevalence of K^+ permeability of the plasmalemma)
3. Functional expression of GABA and glutamate transporters, thus GABA and glutamate uptake
4. A large number of intermediate filament bundles (sites of the astrocyte specific protein glial fibrillary acidic protein - GFAP)
5. Glycogen granules
6. Processes from each cell contacting and surrounding blood vessels
7. Elaborated perisynaptic processes
8. Linkage to other astrocytes by gap junctions consisting of connexin 43 and/or 30

Many astroglial cells do conform to these criteria, but there are exceptions. Kimelberg's proposed division of astroglia into either star-shaped- (protoplasmic and fibrous astrocytes) or elongated astroglia (retinal Müller glia and cerebellar Bergmann glia) excludes cells that should have been included. For this reason, Verkhratsky and Butt propose that the family of astrocytes can be *functionally* defined as "*the true homeostatic cells of the CNS that provide for molecular, cellular and organ homeostasis*" (Verkhratsky 2013). Neither definition may be perfect, the former – despite its rigorous approach – due to the number of astroglial cells that it unintentionally excludes and the latter – despite its elegant brevity – for not being applicable as inclusion/exclusion criteria in studies relying on morphology. The title of the referenced review (Kimelberg 2010) emphasises that there are differences between immature and mature astrocytes. Indeed, article IV (Lunde et al. 2015) in this thesis provides evidence on postnatal development of the dystrophin associated protein complex (DAPC) in neonatal, perivascular astrocytes that might differ from their mature counterparts. Further, this thesis is limited to the study of astrocytes in mouse and rat brain. Kimelberg correctly highlights that astrocytes in other species are quite different from their mammalian equivalents. As an intriguing example of particular relevance to the gliovascular interface is that astrocytes/neuroglia in shark brains express tight junction proteins and thus constitute endocellular vessels (Long et al. 1968). Recently, it has also been shown that reactive astrocytes from mice create tight junctions (TJs) in inflammatory lesions and constitute a proposed barrier to migration of immune cells and humoral immune response into the brain parenchyma (Horng et al. 2017). Of more relevance are the acknowledged and significant morphological differences between rodent and human astrocytes, whereby human astrocytes are 15-20 times larger than their rodent counterparts (Oberheim et al. 2012). Excitingly, a recent study showed that implantation of human glial cell progenitors into mouse brain

resulted in humanised, chimeric mice that outperform their littermates (Han et al. 2013). Such differences pose challenges for how animal studies, such as the ones proposed herein, can be extrapolated to the human case.

Astrocyte morphology and relationship to neurons, capillaries and parenchyma-to-fluid interfaces define these cells more so than their molecular signature (Oberheim et al. 2012). Nonetheless, the intermediate filament protein GFAP has been used for more than 30 years as a marker for identifying astrocytes. It has become clear, however, that not all GFAP positive cells are astrocytes, and conversely, not all astrocytes are GFAP positive on immunohistochemical preparations (Kimelberg 2004; Mishima and Hirase 2010).

In a tissue such as the CNS, whose function crucially depends on cellular organisation rather than just sheer numbers, the importance of cells involved in homeostasis and structure is not to be underestimated. With evidence that astrocytes ensheath and provide an almost complete covering of vasculature in the CNS (Mathiisen et al. 2010), they are an excellent example thereof - uniquely situated to link brain parenchyma to the vasculature. And even though pathologies such as Guillan-Barré syndrome (GBS), acute lateral sclerosis (ALS) and multiple sclerosis (MS) manifest due to damaged neurons or oligodendrocytes, it is quite possible - even probable - that the respective pathogeneses also involve compensatory mechanisms of adjoining astrocytes. A bold statement regarding neuropathologies would therefore be that no disease manifests symptomatically unless astrocytes fail in maintaining or restoring perturbed homeostasis of affected regions or microenvironment (Verkhratsky 2013). The dedicated study of all aspects of the astrocyte is therefore crucial to a more in depth understanding of neuropathology. Included in this approach would be a systematic investigation of the proteins expressed by astrocytes. This includes the water channel AQP4, which is expressed in abundance in astrocyte endfeet.

Aquaporins

Aquaporins (AQPs) constitute an ancient family of proteins that is present in all living organisms and thus all three kingdoms of life; Eukarya, Bacteria and Archaea (Finn et al. 2014; Laloux et al. 2018). This suggests a common ancient ancestor. AQPs are membrane intrinsic proteins (MIPs) that facilitate the passive movement of water and small solutes across biological membranes. There are 13 mammalian aquaporin genes. They can both functionally and phylogenetically be divided into four categories based on their amino acid sequence. First, the channels uniquely permeable to water (*Aqp0*, 1, 2, 4, 5 and 6) are the water selective, classical or orthodox aquaporins. Second, *Aqp8* can be termed an ammoniaporin due to the ability to transport water, ammonium and urea (Jahn et al. 2004), although AQP8 mediated ammonium transport may not be physiologically significant (Yang et al. 2006a). Third, the aquaglyceroporins (*Aqp3*, 7, 9 and 10) facilitate the transport of H₂O, arsenic, urea and polyols such as glycerol. Finally, a relatively recently described group comprising *Aqp11* and *Aqp12* is called superaquaporins by some and *unorthodox aquaporins* by others (Ishibashi 2009; Ishibashi et al. 2009). Their permeation properties have yet to be identified. As their names imply, the functions of orthodox aquaporins and aquaglyceroporins are related to water transport, but evidence of other physiological properties certainly exists. Evidence towards ammonia channelling via a fifth, central pore resulting from the assembly of AQP1 into tetramers is one example (Ip et al. 2013; Nakhoul et al. 2001).

To date, expression of seven aquaporins has been described in rodent brain. These are *Aqp1*, -3, -4, -5, -8, -9 and -11 (Agre et al. 2002; Badaut et al. 2001; Elkjaer et al. 2000; Gorelick et al. 2006; Nielsen et al. 1997; Venero et al. 1999; Yamamoto et al. 2001). Two of these have known physiological functions in the human brain (AQP1 and AQP4). There is evidence both for and against the localisation and potential physiological role of AQP9

(Amiry-Moghaddam et al. 2005; Badaut et al. 2004; Mylonakou et al. 2009; Yang et al. 2006b). Studies on a transgenic knock out mouse strain have also implicated the protein in the pathogenesis of Parkinson's disease (Stahl et al. 2018). For the remaining three, the evidence for their presence in CNS has been reviewed previously (Badaut et al. 2001, Elkjaer et al. 2000; Nielsen et al 1997, Venero et al 1999; Yamamoto et al 2001b). Although mRNA for *Aqp3*, *Aqp5* and *Aqp8* is detected in rodent CNS, the localisation studies at protein level are limited by the relative lack of functioning antibodies. To what extent these latter aquaporins are expressed at protein level and provide functional properties to the CNS can therefore be debated.

AQP4 in the CNS

The scientific exploration of AQP4 begins long before the gene was cloned. Of particular note is the freeze-fracture technique and how this was employed as early as the 1970s to study membrane domains not just in astrocyte endfeet but also in other organs. Immunocytochemistry enabled Nielsen and co-workers (Nielsen et al. 1997) to characterise the distribution of AQP4 in the CNS and other tissues. But the original work by Dermietzel was the first to visualise OAPs, which we now know to be composed uniquely of AQP4 molecules (Dermietzel 1973). Landis and Reese (Landis and Reese 1974) then showed these particles to be organised into “assemblies” and Rash et.al showed that they were not gap junctions (Rash et al. 1974). They have since been referred to as either square arrays or orthogonal arrays of particles (OAPs). The topic was thoroughly reviewed by Wolburg in 1995 (Wolburg 1995). The molecular composition of OAPs has been studied and evidence suggests AQP4 to be the sole constituent of OAPs. But the functional significance of such supramolecular assemblies remains enigmatic (Sorbo et al. 2008). Frydenlund and co-workers demonstrated how AQP4 is progressively lost following brain ischemia (Frydenlund et al. 2006). But this very phenomenon was also observed by Landis & Reese as early as in 1981 (Landis and Reese 1981a). The authors described how “*assemblies are progressively replaced by amorphous clumps and then disappear as the interval between decapitation and rapid freezing increases. Nearly normal numbers of assemblies may be maintained in cerebellar slices in vitro, but there too they disappear at low PO₂ [...]. No other neuronal or glial membrane specialization exhibits a comparable lability*”. A review written by the same authors in 1981 uses a wording which is surprisingly similar to what authors of this millennium employ when summarising the role of aquaporin-4 in the CNS: “*The function of assemblies is unknown, but their positioning suggests that they may have a role in the transport of some material into or out of the blood and cerebrospinal fluid compartments*” (Landis and Reese 1981b). The relevance of different AQP4 isoforms to form square arrays has been studied (Rossi et al. 2012; Verbavatz et al. 1997; Yang et al. 1996). There are significant species differences to be taken into account. In the rat, a total of 6 isoforms have been identified (Moe et al. 2008) and termed AQP4 a-f, where a and c correspond to the M1 and M23 isoforms in mice. As shown in the western blots from mouse brain in article II (figure 5B, C) three distinct bands appear upon incubation with anti-AQP4 antibodies directed against the C-terminal, these are M1, M23 and Mz. Each of these bands disappear following *Aqp4* knockout.

Article III in this thesis makes use of a cell culture system, tailor made so as to study the particular expression pattern of AQP4, whereby most of this protein is localised to astrocyte endfeet abutting either vasculature or forming the glia limitans proper. This phenomenon of protein polarisation was also noted by scientists employing the freeze fracture technique (Landis and Reese 1982). As a hallmark of perivascular astrocytes, functional polarisation should perhaps have been included in the defining criteria proposed by Kimelberg. Arguably, though, neurons and oligodendrocytes are also functionally polarised and the criterion would have been a necessary rather than sufficient factor in the definition.

The discovery of aquaporins in brain resulted in further insights into pathophysiology compared to physiology (Nagelhus and Ottersen 2013). The contrast to the situation in kidney is obvious. The physiological role of AQP4 proved elusive, but evidence now exists of contribution to both physiology and pathophysiology. Inevitably, due to the localisation of AQP4 and the functional polarisation of astrocytes, this literature relates in large part to the gliovascular interface (Nagelhus and Ottersen 2013). One recent and exciting report implicates AQP4 in clearance of waste products from brain (Ilf et al. 2012). Parts of this hypothesis have been contested by others showing no difference in paracellular diffusion when comparing WT to *Aqp4* knockout (*Aqp4*^{-/-}) genotypes (Smith et al. 2017).

AQP4 localisation in brain is not limited to perivascular spaces, however, and is also found in astrocytes of olfactory bulb and in basolateral membrane of ependymal cells lining the entire ventricular system. The first accounts of immunogold labelling (Nielsen et al. 1997) also show distinct labelling in astrocyte processes in contact with synapses which spawned further investigations related to neurophysiology and the so-called tripartite synapse (Araque et al. 1999; Newman 2003). Further, an exciting report on lamellar stack astrocytes (Holen 2011) expanded on initial observations from Nielsen et.al in the supraoptic nucleus and demonstrated distinct structures termed “circular lamellar stacks” that envelop neuronal structures and are enriched in AQP4. Such structures are found scattered throughout the CNS (unpublished data, Hoddevik). They define an astrocyte–neurone interaction that merits further investigation.

Studying AQP4 expression in other organs provides equally interesting clues on regulation as do studies from the CNS. Mechanisms regulating AQP4 expression in skeletal muscle have been poorly understood, but include both neuronal innervation (Ishido and Nakamura 2018) and exercise induced changes in protein level. AQP4 supports muscle contractile activity and metabolic changes that occur in fast-twitch skeletal muscle during prolonged exercise (Basco et al. 2013). Controversy exists, though, since not all fast-twitch fibres seem to express AQP4 (Kaakinen et al. 2007).

The literature on AQP4 is vast. A literature search on the US National Library of Medicine website (PubMed.gov) on the keyword “aquaporin-4 OR AQP4” currently yields 3737 scientific articles of interest. Not all are relevant for AQP4 regulation and localisation, but the search result is provided to illustrate the extent to which this protein has been studied.

Astrocyte endfeet

The perivascular processes of astrocytes are perhaps best described as lamellopodia. In the literature, however, they are referred to as end-feet or endfeet. In 1955, Glees used the word “end plate” to describe astrocyte processes terminating on a cerebral vessel wall. He referred to the envelopment of the vascular wall by endfeet as the glial barrier and termed the ensemble of end plates as “membrana limitans gliae perivascularis (Glees 1955). But endfeet are not coupled by tight junctions and the clefts between them (Korogod et al. 2015; Long et al. 1968) are not likely to restrict movement of solutes to the extent that brain endothelium does. They can be viewed as specialised projections from the cell soma and perikaryon. They establish the concept of astrocyte polarity, a reason for which astrocytes bear similarities to neurons (Barres 2008; Nedergaard et al. 2003; Seifert et al. 2006). Despite the importance of endfeet, a review article listing a complete “inventory list” of endfoot constituents has yet to be written. Consequently, this paragraph is included to briefly review some select and important examples of endfoot composition.

Endfoot cytoskeletons do contain GFAP (El-Khoury et al. 2006) and intermediate filaments, albeit in modest amounts compared with projections of fibrous astrocytes. Filaments were described morphologically long before the protein was identified in 1972 (Eng et al. 1971). At

the ultrastructural level, the classical intermediary filament bundles are generally not seen, but GFAP immunogold labelling in endfeet is sufficiently strong for immunogold quantitation (Hoddevik, unpublished data on GFAP in WT vs α -Syn^{-/-} mice). In cell culture assays, GFAP is successfully employed as an astrocyte marker labelling endfeet, like we show in article III (Camassa et al. 2015). In reactive astrocytes, AQP4 localisation to endfeet has been suggested to have an active involvement in migration – a process that must involve changes in the underlying cytoskeleton (Saadoun et al. 2005).

Among the transmembrane proteins known to link endfeet to the adjoining ECM are β -integrin and β -DG (Campbell and Stull 2003; Mayer 2003). In turn, the latter is coupled to the proteins of the DAP complex, which includes dystrobrevin, α -, β 1 and β 2 syntrophins. The handling of K⁺ is of essence in an excitable tissue such as the CNS and endfeet contain the inwards rectifying K⁺ channel Kir4.1 (Nagelhus et al. 1999). As defined by Kimelberg, one of the characteristic features of astrocytes is their ability to form a glial syncytium through expression of gap junction proteins such as connexin 43 and connexin 30. Their presence is likely to serve several functions, including a better resilience and increased capacity for buffering K⁺ levels (Katoozi et al. 2017; Nagy and Rash 2000). Considering the fundamental role of neurotransmitters in the CNS it is perhaps not surprising that glutamate transporters are also found in endfeet (Lehre et al. 1995). The transient receptor potential vanilloid-related channel 4 (TRPV-4) is particularly abundant in astrocyte membranes at the interface between brain and extracerebral liquid spaces, and thus also in endfeet. TRPV-4 has been suggested to play a role in astrocyte swelling which in turn promotes intracellular Ca²⁺ elevation through Ca²⁺ influx (Benfenati et al. 2011).

Involvement of endfeet in metabolism is inevitable, considering how astrocytes link with the vasculature. Neurons and the brain proper depend on glycolysis. In humans, the brain constitutes 2 % of the body weight, but is responsible for 20 % of the body's energy consumption at rest (Rolfe and Brown 1997). Endothelial cells and astrocyte endfeet alike therefore express GLUT-1 (Allen and Messier 2013; Simpson et al. 2007). Further, monocarboxylate transporters (MCT1) transporting lactate are also found in endfeet (Simpson et al. 2007). Astrocytes regulate metabolic homeostasis through glycogen synthesis and the supply of energy substrates to neurones. Endfeet also release vasoactive substances and contribute to functional hyperaemia (Nippert et al. 2018).

The above list does not cover nearly all the proteins present in endfeet but should serve its purpose of illustrating why and how astrocyte endfeet are crucial for CNS function and homeostasis.

Astrocyte polarity, AQP4 regulation and dystrophin associated glycoprotein complex

Immunolabelling techniques have revealed a multitude of different membrane proteins in endfoot membranes, some of which display a highly polarised expression pattern. This includes AQP4 (Nielsen et al. 1997). In articles III and IV we use AQP4 as a marker of functional astrocyte polarity. This is a hallmark of astrocytes whose underlying regulation, induction and maintenance have been incompletely understood. There is ample evidence that AQP4 in perivascular endfeet co-localises with the DAP complex, consisting of DG, α -Syn, dystrophin and dystrobrevin (Amiry-Moghaddam et al. 2003a; Inoue et al. 2002; Neely et al. 2001; Noell et al. 2011). The reader is referred to the thorough review by Blake for further reference (Blake et al. 2002). Articles II, III, IV and V all address anchoring mechanisms of transmembrane proteins via the dystrophin-dystroglycan protein complex. Evidence suggests that α -Syn has a direct interaction with AQP4 in brain to fulfil the role as an anchoring protein to the DAP complex. β -DG is the transmembrane part of the DG protein and the extracellular and heavily glycosylated α -dystroglycan binds to laminin (Culligan et al. 2001). Together with integrins, β -DG therefore contributes as an ECM receptor for astrocytes. This interaction is

what we hypothesised to contribute as both an inductive and regulative mechanisms for AQP4 levels and astrocyte polarity.

Syntrophins are cytoplasmic peripheral adapter proteins of the DAP complex (Neely et al. 2001). The five syntrophin isoforms ($\alpha 1$, $\beta 1$, $\beta 2$, $\gamma 1$, $\gamma 2$) are encoded for by distinct genes. Each contains two pleckstrin homology (PH) domains, a syntrophin-unique (SU) domain, and a PDZ domain. The latter acronym stems from the first letter of those three proteins found to contain repeats of this domain, namely PSD-95, Drosophila discs large protein, and the zonula occludens protein 1. PDZ domains in other proteins are known to bind to the C termini of ion channels and neurotransmitter receptors that contain the consensus sequence (S/T)XV-COOH (where S = Serine, T = Threonine, X is any amino acid and V = Valine) and mediate the clustering or synaptic localisation of these proteins (Gee et al. 1998; Peters et al. 1998). AQP4 has the C-terminal sequence Ser-Ser-Val (-SSV). This sequence potentially binds to PDZ domains (Fanning and Anderson 1999; Gee et al. 1998; Kornau et al. 1995), present in syntrophins and therefore also in α -Syn.

Since the present work was undertaken, other mechanisms of AQP4 regulation have been unravelled, but their role *in vivo* is not fully established. Quite recently, a pivotal study provided evidence of nascent protein synthesis in these distal astrocyte processes (Boulay et al. 2017). In addition, the authors found smooth- and rough endoplasmic reticulum as well as Golgi apparatus to be present in endfeet. They suggest that the resulting protein synthesis may sustain the functional polarisation phenomenon. Similarly, Sakers and co-workers have also shown that astrocytes locally translate transcripts in their peripheral processes (Sakers et al. 2017). Together, these findings open for exciting, new hypotheses regarding control of astrocyte polarity. The thought is not new, however, and polarisation phenomena of AQP4 were early on studied in cell cultures (Madrid et al. 2001).

Another recent publication based on primary astrocyte cultures (rat) sheds light on astrocyte-ECM interactions and potential mechanisms of regulation. Thus, Tham et.al. examined AQP4 cell surface expression and turnover (Tham et al. 2016) and found that cell-surface expression of AQP4 is dependent on DG and potentiated by laminin. Excitingly, the authors found evidence of an intracellular, early endosome antigen-1 (EEA-1) associated pool of AQP4 from which the protein was recruited to the cell surface by presence of laminin. Further, they demonstrated an association between DG and dynamin where β -DG preferentially bound to inactive dynamin and was proposed to inhibit endocytosis. In turn, dynamin was shown to function in conjunction with clathrin to regulate surface AQP4 levels. It remains to be seen if such findings are replicated *in vivo*.

In Figure 1 I have attempted to provide an illustration of the putative mechanisms involved in AQP4 regulation. These mechanisms include anchoring to the DAP complex via α -Syn (Amiry-Moghaddam et al. 2004; Neely et al. 2001), local protein translation in astrocyte endfeet (Boulay et al. 2017), vesicle mobility (Potokar et al. 2013), local recruitment from an intracellular pool of AQP4 and local vesicular internalisation (Tham et al. 2016).

Figure 1: Putative mechanisms for AQP4 regulation in astrocyte endfeet are shown schematically at the gliovascular interface, where astrocyte endfeet (ast) abut the basal lamina (BL), pericytes (Per) and endothelium (End). Inset in 1) shows anchoring of AQP4 via α -Syn to DAP complex, a mechanism underlying astrocyte polarity and AQP4 localisation to astrocyte endfeet. The DAP complex in turn is coupled to ECM via α - and β -DG. Shown in 2) is the laminin-induced, local recruitment of AQP4 to the membrane from an intracellular EEA-1 associated pool (arrow). β -DG preferentially binds to the inactive form of dynamin and may inhibit endocytosis of AQP4 (bottom of 2). Vesicular transport may also be present in cell soma, underlying the potential membrane diffusion of AQP4 shown in 3) (curved arrow). AQP4 would diffuse along the cellular membrane and adhere to the adluminal endfoot membrane upon interaction with α -Syn, enabling functional polarisation for AQP4. Finally in 4) is a schematic of the local protein translation that has been shown to exist in astrocyte endfeet following mRNA transport from cell soma. Common for the proposed mechanisms 1-3 is the reliance upon ECM proteins laminin and agrin in the specialised, perivascular BL. Visual presentation is adapted from diagrams in Nagelhus and Ottersen 2013, Tham et.al. 2016.

Pericytes

The term “pericytes” was introduced by Zimmermann in 1923 (Zimmermann 1923) and the first descriptions of such cells can be found in the works of Eberth (Eberth 1871) and Rouget (Rouget 1873). Pericytes have since been given much attention and in 1979, active pericytic contraction was shown to narrow the capillary lumen (Tilton et al. 1979). It is often stated that microvessels are ubiquitously associated with pericytes, but there are quantitative differences in pericyte coverage (PC) of vessels in the CNS. An example is the albino rat, where PC is greater in retinal than in cerebral capillaries (Frank et al. 1987). Pericytes are located on pre-capillary arterioles, capillaries and post-capillary venules of many organs (Dalkara et al. 2011) but they occur very frequently on microvessels of retina and brain. For instance, the pericyte-to-endothelial cell ratio is 1:100 in striated muscle, but as high as 1:3 in brain and 1:1 in retina (Dalkara et al. 2011; Shepro and Morel 1993). The impressive variation in stoichiometry between these tissues implies physiological differences. Indeed, there is evidence that pericytes contribute to the integrity and formation of the blood to brain barrier (BBB) (Armulik et al. 2010; Bell et al. 2010; Daneman et al. 2010). There is also evidence of pericyte contractility and neurotransmitter control, with subsequent implications for flow regulation in the capillary bed of the CNS (Hall et al. 2014).

There is evidence that pericytes play a role in regulating regional cerebral blood flow (rCBF) to the very focal demand of nearby, active cell populations. The very basis of blood oxygen level-dependent (BOLD) and functional magnetic resonance imaging (fMRI) in CNS is that

neural activity increases local blood flow. How active neurons signal to blood vessels is termed neurovascular coupling, and the phenomenon by which active brain regions induce a local increase in blood flow is named functional hyperaemia. BOLD fMRI is currently the leading method for non-invasive imaging of human brain activity, but it does not measure brain activity directly – rather, it relies on neurovascular coupling mechanisms (Rungta and Charpak 2016). Pericytes are thus implicated and Peppiatt and co-workers provided evidence that these cells contribute to this phenomenon by altering blood flow in the capillary bed rather than in arterioles (Peppiatt et al. 2006). An excellent review on this topic was written by Hamilton et al. in 2010 (Hamilton et al. 2010) and recent publications expand our understanding of neurovascular coupling (Mishra et al. 2016). It has also been proposed that capillary flow dynamics constitute a passive phenomenon which is significantly affected by the architectural properties of capillaries *per se* (Nakada et al. 2017). The latter review also puts forward the intriguing hypothesis that functional hyperaemia has the primary function of regulating temperature, rather than simply supplying glucose to an active cerebral tissue.

Pericytes might also contribute to or influence the composition of the ECM organised into the surrounding basal lamina. An example is provided by Robel et al (Robel et al. 2009). The composition of BL between astrocytes and pericytes is thus likely to be influenced by both cell populations.

Brain endothelium and the blood brain barrier

If astrocytes constitute the most diverse cell type of the CNS, endothelial cells represent perhaps the most diverse cell of the entire organism. In the brain, they are an integral part of the gliovascular unit. One of the well-known characteristics of brain endothelium is its impressive impermeability to many substances, a phenomenon still known as the blood-to-brain barrier (BBB), succinctly reviewed by Ingo Bechmann in 2007 (Bechmann et al. 2007). That astrocytes surround endothelium was described by Andriezen as early as in 1893 and astrocytes have long been known to influence the formation of the BBB via endothelial tight junction (TJ) production (Arthur et al. 1987).

As a concept, the BBB knows its origin from Berlin in 1885. The unwanted observation made by Paul Ehrlich, which ruined his experiment at the time, has since spawned an entire field that continues to grow. Three other works are key papers in defining the notion of the BBB. Lewandowski's work 'on the cerebrospinal fluid' from 1900 (Lewandowski 1900). Then Goldmann from 1913 with 'Intravital labelling of the central nervous system' (Goldmann 1913) and finally Reese and Karnowski's work; 'Fine structural localization of a blood-brain barrier to exogenous peroxidase' from 1967 (Reese and Karnovsky 1967). In a logical manner do these four studies zoom in on the structural correlates of the BBB. From a special 'surface' as defined by Ehrlich in 1885, to tight junctions, membrane channels and endothelial transport systems. When Reese and Karnovsky defined and introduced the notion of tight junctions in 1967, the premises for modern BBB research had been laid (Reese and Karnovsky 1967). Today, we know that the blood-brain barrier consists of several separate phenomena: 1) the presence of cellular interactions named tight junctions that restrict (or possibly regulate) paracellular transport, and 2) specialised membrane domains in endothelial cells that permit selective transport in both directions across the endothelium. The literature is packed with unspecific and vague terminology, including terms such as "leaky BBB". There is no doubt, however, that transport mechanisms at the BBB are closely regulated phenomena.

Lymphatic drainage of the CNS

The experimental evidence reported in this thesis is more descriptive than what it involves registrations of physiology. But the implications of AQP4 regional heterogeneity are potentially interesting when discussed in the context of vascular physiology in brain, which

includes perivascular spaces and what has recently been characterised as “glymphatic” flow. In a seminal work, Iliff and co-workers injected tracers into the cisterna magna and were able to map out a flow via perivascular spaces back into the neuropil (Iliff et al. 2012). In another study from the same group, the authors were able to assess extracellular space in mice that were sleeping and thus provided the first historical visualisation of sleeping brains (Xie et al. 2013). A finding of increased tracer flow through the neuropil of sleeping mice led to the deduction that the extracellular space (ECS) expands and cells shrink during sleep. The cellular mechanisms underlying this observation are intriguing but must include osmotic forces and osmolytes that do not interfere with brain functions. Studies have been reported on glymphatic flow in the ageing brain (Kress et al. 2014), brain trauma (Plog et al. 2015), and potentially Alzheimer’s disease (Iliff et al. 2012).

Excitingly, cerebrospinal fluid (CSF) characteristics and glymphatic function in humans was recently studied using MRI detection of intrathecally administered gadobutrol (Ringstad et al. 2017).

Solute transport through the ECS is a phenomenon which is critical for many aspects of brain homeostasis. It ensures delivery of nutrients to brain cells, clearance of metabolites, neurotransmitters and toxic macromolecules. ECS in brain includes CSF containing spaces (ventricular system and subarachnoid space) on the one hand, and parenchymal extracellular space as well as para- and perivascular spaces on the other.

With reference to the above historical account of the BBB, the tracers that Goldmann injected seemingly did not cross brain vessel walls. However, further examination revealed that the neuropil remained unstained, but that the dyes accumulated in cells of the choroid plexus, the leptomeninges and along the perivascular spaces. These perivascular spaces were rightly regarded by Goldmann and his contemporaries as lymphatic clefts. The labelling was stored in granules of cells which Goldmann recognised as susceptible for chemotactic signals and capable of migration and phagocytosis. With humour has this been referred to by some as “the dirty little secret of the anatomists” (Galea et al. 2007). It is mentioned here to illustrate that the phenomenon of perivascular and paravascular drainage and transport is not new, but has been reported and studied previously.

The experimental data published by Iliff et.al has been investigated by others. A mathematical modelling based on 3D reconstructed models of CNS thus finds diffusion to be more prominent than bulk flow for interstitial transport of solutes in CNS (Holter et al. 2017). This is evidence against parts of the discussion offered by Iliff and co-workers to explain their findings. Similarly, Smith et.al. provided evidence against the proposed glymphatic mechanism by convective solute transport in brain parenchyma (Smith et al. 2017). The exact mechanisms underlying the experimental data provided by Iliff and co-workers are perhaps not fully mapped out, but the article remains seminal. Smith and co-workers also provided other experimental data on the diffusion of dextrans of different molecular weights in *Aqp4*^{-/-} vs WT genotypes. They found no difference between genotypes and thus argue against the role of AQP4 in these physiological processes. There are discrepancies in the experimental setup by Smith compared with that of Iliff, however, but controversy now exists as to the involvement of AQP4 in paravascular- and glymphatic flow. The first account of glymphatic flow studied by MRI in a human patient cohort (Ringstad et al. 2017) also included patients with normal pressure hydrocephalus, known to exhibit reduced intracranial compliance and restricted intracranial artery pulsations (Eide and Sorteberg 2010). Gadolinium conjugates have previously been used to study rat, where such pulsations were interpreted as driving paravascular CSF-interstitial fluid exchange (Iliff et al. 2013). Further, the study of such disease states in humans will undoubtedly improve our understanding of glymphatic flow and intracranial fluid dynamics in humans. Such methodology could shed further light on the role

of AQP4 in glymphatic flow if used to study patients with polymorphisms in – or known, disease causing mutations of – the human *Aqp4* gene (Nicchia et al. 2011; Sorani et al. 2008a; Sorani et al. 2008b; Yadav et al. 2014). Of particular interest would be those patients with phenotypes arising from the CNS (Berland et al. 2018).

The brain lacks conventional lymph vessels and the removal of solutes through the brain interstitium thus has to rely on other anatomical structures and mechanisms. It is perhaps surprising that our understanding of such fundamental notions are only now beginning to unravel. Only time will tell, however, to what extent this may be of clinical use. AQP4 is certainly not the only protein involved, but the substantial amount of scientific articles studying this particular protein may now find unforeseen, practical applications.

Extracellular matrix, basal lamina, laminins and agrin

In view of how fundamental the ECM is in biology and the multitude of constituents it is assembled from, it is challenging to provide but a brief introduction. I have attempted to limit this paragraph to those concepts necessary to understand the 5 articles on which my thesis rests. I will therefore first focus on nomenclature. Thereafter, I will review the composition of basal laminae and emphasise the functions of those proteins I have examined.

The terms basement membrane (BM) and basal lamina (BL) are often used synonymously, depending on whether researchers employ light- or electron microscopy, respectively. For consistency, I have adhered to employing BL in this thesis. Bowman defined the concept in his first description from muscle tissue, he rightly called it a “membranous sheath of the most exquisite delicacy” (Bowman 1840). The ECM is not limited to basement membranes, however, and the term does mean different things to different people. For the purpose of the proposed work, I wish to limit the discussion to ECM in brain. The best understood components of the ECM comprise collagens, proteoglycans and major glycoproteins. Based on prior evidence suggesting that agrin and laminin (Fallier-Becker et al. 2011; Guadagno and Moukhles 2004; Tham et al. 2016) impact on AQP4 membrane levels, we hypothesised that protein levels of AQP4 and α -Syn would correlate with those of both agrin and laminin. But the proposed study is limited to these two, latter proteins. They are but two on a long list of ECM constituents. This highlights the limitations of article V when interpreting heterogeneity of the capillary BL. Focus of this particular work was to examine potential correlations between interaction molecules of α -dystroglycan that might in turn explain the observed variations in endfeet levels of AQP4 and α -Syn. This dictated the choice of proteins to study, namely those two ECM proteins respectively known to influence membrane AQP4 and OAP formation (agrin), and known to link DAP complex in place (laminin) with secondary influence on AQP4.

The heterotrimeric laminins are a defining component of all BLs and self-assemble into a cell-associated network (Hohenester and Yurchenco 2013). The laminin molecule is cross-shaped and the three short arms constitute the network nodes, with a strict requirement for one α , one β and one γ arm (Aumailley et al. 2005; Parsons et al. 2002). The α -subunits (two of which are examined in article V) are largely responsible for cell surface adhesion and receptor interactions, but also contribute to self-assembly (Yurchenco 2011).

The globular domain at the end of laminins long arm binds to cellular receptors, including integrins, α -dystroglycan, heparan sulfates and sulfated glycolipids. Both integrins and dystroglycans are transmembrane proteins that interact with the cytoskeleton (Campbell and Stull 2003; Mayer 2003). This series of direct protein interactions therefore constitutes a bridge which links the basal lamina to the very cytoskeleton of neighbouring cells, reviewed

by Baressi and Campbell (Barresi and Campbell 2006). Collateral anchorage of the laminin network is then provided by the proteoglycans perlecan and agrin. A second network is thereafter formed by type IV collagen, which interacts with the laminin network through the heparan sulfate chains of perlecan and agrin and additional linkage by nidogen.

BLs have been deduced to assemble through a multistep process substantially driven by mass-action. This is termed self-assembly and is initiated by the binding of laminins to cell surfaces (Li et al. 2002; Li et al. 2005; McKee et al. 2009; McKee et al. 2007; Smyth et al. 1998). Networks of both collagens IV and laminin are capable of such self-assembly (Sanes 2003). The reader is encouraged to cast a glance at Fig 4 in article V at this point, where collagen fibres in pia mater are displayed in an impressive and systematic pattern. I find this image fully in line with Bowman's initial description of a tissue of exquisite delicacy.

Devic's disease:

For the practicing neurologist, any study of AQP4 in the CNS now yields an immediate association to neuromyelitis optica (NMO). The name was coined by Eugène Devic and his student Fernand Gault who described the disease as a different version of "*la sclerose en plaque*" as early as in 1894 (Devic 1894) and described the condition as "*La neuromyélite aiguë du nerf optique*". Even earlier, a review written by the German oculist Friedrich Albin Schanz (1863–1923) points to most of the clinical and pathological features nowadays considered characteristic for NMO. Some would claim this to be the first account of NMO in the literature (Jarius and Wildemann 2017). Historical roots set aside, its existence was refuted by some during the 20th century but not forgotten by others. Excitingly, a potential molecular basis for the disease was shown by Vanda Lennon and co-workers in 2005 (Lennon et al. 2005) upon isolating antibodies from sera of multiple sclerosis (MS) patients displaying the characteristic symptoms described by Devic. These autoantibodies were shown to be directed against AQP4, and thus not against epitopes present in oligodendrocytes. The key control to Lennons experiment was the use of transgenic *Aqp4*^{-/-} mice generated by Ma and co-workers (Ma et al. 1997). The perivascular labelling pattern resulting from incubation of NMO-Ig on tissue sections disappeared entirely in *Aqp4*^{-/-} mice. The AQP4 protein was already cloned and well characterised at the time when Lennon and co-workers published their 2005 report. This facilitated the further exploration of the epitope(s) recognised by the NMO auto-antibodies. Evidence shows that NMO-IgG bind to extracellular loops of AQP4, monomers, tetramers and higher order structures (Iorio et al. 2013). Another report, using pull-down experiments, show assemblies of AQP4 into OAPs as the NMO-IgG target, and demonstrate that antigen-antibody binding does not impede water transport (Nicchia et al. 2009).

Whether the development of anti-AQP4 auto-antibodies in NMO patients truly holds the key to treating the disorder remains unclear, but the coupling of NMO to AQP4 has been seminal. The discovery of anti-AQP4 monoclonal antibodies now provides the medical community with the means to test this hypothesis. The current thesis provides quantitative data from several regions in the CNS on AQP4 expression and localisation. In that regard, it may be relevant to several aspects of NMO development, which is elaborated upon in the third part of the discussion. Current view is that AQP4 is not expressed in oligodendrocytes or neurons, albeit for one report from the vomeronasal organ (Ablimit et al. 2008). The discovery of auto-antibodies from human sera able to bind to this protein has therefore not only been seminal for publications on NMO, but indeed also for CNS autoimmunity. It prompts the question of whether the pathogenesis in other demyelinating disorders also involves astrocytes.

Anti-aquaporin-4 auto-antibodies are specific for neuromyelitis optica spectrum disorders (NMOSD), but there is evidence both supporting and refuting causality. Evidence does suggest complement involvement (Duan et al. 2018). But as another example, anti-aquaporin-4 auto-antibodies have been detected in sera of patients suffering from systemic

lupus erythematosus (SLE) but who did not have CNS symptoms for more than a decade following initial detection (Alexopoulos et al. 2015). Such findings might suggest that the presence of anti-aquaporin-4 auto-antibodies is a necessary but not sufficient factor to develop NMO.

Also, the titer of the anti-AQP4 antibody ELISA assay has been shown not to be predictive in the disease course for patients with NMOSD. Low titer patients experience the same disease course as medium-titer and high-titer anti-AQP4 antibody patients with NMOSD (Kessler et al. 2017). Seemingly, therefore, the diagnostic value of NMO-IgG is currently limited to a positive/borderline/negative result. AQP4 is also found in other tissues that are spared in NMO, which sheds light on the pathogenesis. Kidney and muscle are appropriate examples to be studied. The rare case of myositis does exist in NMO (Cosgrove et al. 2014), where IgG binding and muscle injury has been reported (Malik et al. 2014). As a rule of thumb, however, NMO only involves CNS. Why skeletal muscle is spared in NMO can be considered an evolving mystery (Verkman et al. 2018), also since intravenously administered AQP-IgG has been shown to bind rapidly to mouse skeletal muscle (Ratelade et al. 2011). It has been proposed that NMO-IgG binds differentially according to differing size of the supramolecular aggregates (Rosito et al. 2018). Freeze fracture data on healthy, human skeletal muscle show OAPs of similar size to what has been demonstrated in brain, however (Furman et al. 2003; Wakayama et al. 1986). And conceptually, an NMO/AQP4-IgG molecule bound anywhere within an AQP4 cluster away from the cluster edge is unable to sense cluster size. The explanation of inflammatory sparing in peripheral tissues may reside in factors other than epitope availability and OAP structure. Differential expression of complement regulator molecules may be one such possibility (Yao and Verkman 2017a; Yao and Verkman 2017b).

The deciphering of causal mechanisms for any disease provides the possibility of understanding general disease mechanisms. Such is also the case for NMO: the field of neuroimmunology has not been left unchanged by the increasing number of publications on patients suffering from this disorder.

3. Hypotheses and aims of the thesis

- Hypothesis 1: The specialised extracellular matrix found in basal laminae governs the composition of the adjoining astrocyte membranes and regulates the level of proteins expressed therein.
- Hypothesis 2: Membrane levels of AQP4 are regulated both by intra- and extracellular factors.
- Aim 1: To explore the molecular bridging between endfeet and basement membranes
- Aim 2: To unravel those mechanisms responsible for accumulation and localisation of AQP4 in astrocyte endfeet abutting perivascular basement membranes
- Aim 3: To unravel examples of heterogeneous AQP4 expression in order to assess potential regulatory mechanisms under different circumstances.

4. Methodological considerations

Quantitative immunogold labelling

The following is a schematic of the eight-step approach to the conducted, quantitative immunogold experiments. In article II and V, we used at least 4 mice in each group. For article II, we dissected out 6 brain regions with a maximal diameter of 1 mm. Images were sampled from all identifiable subregions (in hippocampus, for instance) and 20-30 images were sampled for each.

Figure 2: The eight-step approach in performing quantitative immunogold labelling is shown.

Immunogold labelling is a potent technique. It allows for quantitative assessments of cellular microdomains at ultrastructural levels. The quantitation for AQP4 and α -Syn was done as linear alignment whereby gold particles in proximity of the cell membrane were counted. In the case of article V, immunogold labelling of agrin and laminin was assessed in capillary BLs. An arbitrary line was drawn in the middle of the cross-sectioned basal lamina. Histograms were made for both proteins and did not show any convincing sign of increased expression towards the endfeet or endothelium (histograms, article V).

Immunogold labelling is also associated with significant challenges. In principle, only two datapoints showing different values would be sufficient to provide evidence towards regional difference for a given protein. Images in articles II, IV and V were uniquely sampled from capillaries and the data does not, therefore, allow discussions concerning endfeet surrounding veins, venules, arterioles or arteries. To avoid bias, images were systematically taken from the upper right quadrant of all identified vessels. A prerequisite for the statistical analysis is that the sampling is representative of the examined phenomenon, i.e. that the membrane distribution of AQP4 in each capillary endfoot is uniform. The scope of the experiments in article III was to create an experimental approach that allowed quantitation. This proved too difficult. Unlike the sampling of capillaries in brain parenchyma, astrocyte processes in cell cultures proved difficult to identify in sufficient numbers. Many reasons accounted for this, including the practical issue of embedding this tailor made cell culture system. 3D orientation had to be preserved and ultrathin sections from the resin blocks had to be taken perpendicularly to the plastic membrane which was technically challenging. Transmission electron microscopy therefore proved suboptimal. Scanning electron microscopy or freeze-fracture replica with immunogold labelling might have been a better choice in order to examine morphology and immunohistochemistry. Below is an image of some of the rare processes detected inside of the porous filter. The image also illustrates the sparse, anti-AQP4 immunogold labelling.

Figure 3: Shown in A) are processes found inside the porous filter used in article III (Unpublished data), sectioned on formvar single hole grids and incubated with anti-AQP4 immunogold particles. Shown in B) is a magnification of inset in A of one process. The immunogold particle associated with the membrane is illustrative of the sparse labelling observed. Note unspecific labelling over the filter proper in A).

Choice of regions

We considered several hypotheses to test when choosing regions. Our choice reflected evolutionary older vs newer parts, highly metabolically active regions (inferior colliculus), areas prone to epilepsy development (hippocampus), sensory regions (piriform cortex and olfactory tract), those two regions primarily affected in NMO (spinal cord and optic nerve) and cerebellum in order to allow comparison with the previous study to quantify regional differences in AQP4 (Amiry-Moghaddam et al. 2004). Both white matter and grey matter regions were also included. Several studies from the 1980s and 1970s evaluated glucose concentrations in various regions of rat and mouse brains (Hawkins 1979, Borowsky 1989). We hypothesised that glucose metabolism, and hence metabolic activity, would display correlations with the heterogeneity of AQP4.

Correlations between signal intensity and protein content – the case of immunology and Aqp4^{+/-} mice

The astute reader will have noticed that “signal strength” from the immunohistochemically based experiments is often used synonymously as “protein levels” in articles I-V. This issue prompts clarification. Linear alignment of the AQP gene family clearly demonstrates the particularities of AQP4 compared to the other 12 genes. No other aquaporin has a C-terminal of similar length as AQP4, nor is any other aquaporin able to form supramolecular assemblies such as the AQP4 square arrays. The AQP4 antibodies used in all of the provided quantitative immunogold data are polyclonal and directed at C-terminal epitopes. If the very reason for which AQP4 is regulated as a separate gene is its capacity to form square arrays, then question arises whether the size and composition of these assemblies may also be a confounding factor in interpreting the regional differences as measured by immunogold labelling. Antibody based assays rely upon availability of epitopes, and the methods employed in articles I-V do not permit an in-depth examination of whether assembly into square arrays might impact on the measured signal. M1 and M23 isoforms have been shown to play different roles in forming square arrays (Crane et al. 2009; Furman et al. 2003) but the AQP4 antibodies used cannot differentiate between them. The western blot in article II with urea cast gels clearly shows both corresponding bands, and it would be possible to compare M1:M23 ratio between CX and IC samples by doing densitometry.

Alas, despite the beauty and accuracy of immunogold cytochemistry, it is probably not possible to provide an exact, mathematical correlation between signal strength and protein levels for either of the quantified immunogold experiments. Attempts at resolving this question have been made (Amiry-Moghaddam and Ottersen 2013), but need to be individualised for each antibody and epitope in question. A theoretical possibility for establishing a correlation between anti-AQP4 immunogold signal and AQP4 levels does exist, and I want to make mention of it here. In the article detailing the creation of an *Aqp4*^{-/-} strain (Ma et al. 1997), Ma and co-workers provide evidence of a consistent reduction of AQP4 levels in brain by ~50% in *Aqp4* heterozygous mice (*Aqp4*^{+/-}). We confirmed these findings in our *Aqp4*^{-/-} strain (Thrane

et al. 2011) through quantitative RT-PCR (unpublished data, courtesy of H.Boldt). To my awareness, no other studies have thus far examined these heterozygous animals. Whether regional heterogeneity within their CNS still exists and how local regulatory mechanisms apply to endfeet remains to be seen. But if the observed 50% reduction in AQP4 protein/*Aqp4* mRNA levels also apply to endfeet, it is possible to make a correlation curve for endfeet labelling based on those three datapoints from *Aqp4* WT, *Aqp4*^{+/-} (50% of WT labelling) and *Aqp4*^{-/-} mice (no residual labelling). Ideally, one final datapoint from mice overexpressing AQP4 should be included – in such a way that AQP4 levels are a set multiple of levels in WT mice. Albeit an interesting model, the overexpressing AQP4 mice linked to the GFAP promotor do not satisfy this criterion (Yang et al. 2008).

That *Aqp4* heterozygous mice express half the level of *Aqp4* mRNA - and correspondingly only half of the AQP4 protein - is indeed also an intriguing phenomenon. Implicitly, there does not seem to be any upregulation in the intact allele despite that half of the AQP4 pool is missing. To my knowledge, up- and downregulation phenomena have not been studied in the *Aqp4*^{+/-} mice and the qPCR data reported here are only examined under physiological conditions. But it may support the hypothesis that regulatory mechanisms resulting in increased *Aqp4* gene transcription are independent of AQP4 levels. This may also complicate the proposed explanation that local translation of *Aqp4* mRNA in endfeet is what sets the AQP4 heterogeneity (Boulay et al. 2017).

One important limitation of the immunogold technique is availability of antibodies that yield a quantifiable signal. The tissue processing inevitably interferes with epitopes thus reducing sensitivity of most antibodies. Some few examples of antibodies I tested during my PhD period with negative results included several anti-CD31, anti-fibulin, anti-collagen XVIII (endostatin), commercially available anti-agrin antibodies, NMO-IgG autoantibodies and antibodies directed at tight junction proteins. A desire for studying a given protein or epitope cannot therefore always be realised through immunogold labelling.

Transgenic mouse strains used

The pitfalls of using transgenic mouse strains are numerous when trying to deduce physiological and functional roles of the studied genes. The flanking gene problem, genetic redundancy and influence of genetic background upon disease phenotypes are but some illustrative examples (Crusio 2004; Nowak et al. 1997; Schafer and Crabbe 1996; Utriainen et al. 2004). We had anticipated the use of one such transgenic mouse strain (DAG-1 x GFAP-cre for conditional deletion of β -dystroglycan) but the data are of preliminary character and a detailed discussion of aforementioned pitfalls is therefore not included. In articles II, III, IV and V the transgenic mouse strains are primarily employed as specificity controls for immunoreactivity. It is difficult to envisage any better controls. Particularly is this true for AQP4 labelling since all immunoreactivity is abolished upon incubation on control tissue/sections. Any potential secondary effect of up- or downregulation of other genes thus becomes irrelevant. The *Aqp4*^{-/-} mouse strain used was first reported in Thrane et.al. 2011 (Thrane et al. 2011) and is one of three available *Aqp4*^{-/-} strains (Ma et al. 1997). Along the lines of the initial postulate regarding pitfalls, findings have been made in but one of these strains that AQP4 gene deletion resulted in altered BBB integrity (Zhou et al. 2008). Similar experiments in the other two strains provided evidence against such effects (Eilert-Olsen et al. 2012; Saadoun et al. 2009). But regardless of the cause, the findings reported by Zhou and co-workers on their Nanjing *Aqp4*^{-/-} strain are intriguing.

Immunoreactivity, antibody specificity and sensitivity

The specificity of the antisera used was confirmed by using transgenic knockout animals for *Aqp4*, α -Syn and β -dystroglycan. In the case of anti-agrin antibody, prior testing on tissue from *agrin* knockout animals is reported in Eusebio et al. 2003 (Eusebio et al. 2003). No knock out tissue was available for laminin. For all immunogold and IF experiments, labelling with primary antibody was done so as to reach equilibrium in the antigen-antibody reaction, according to the following, reversible equation whereby antigen and antibody bind to form a complex:

Equilibrium is reached when the forward and reverse speeds are equal. Factors that affect this reaction can be classified into two groups; those that act on the equilibrium constant (temperature, pH, ionic strength) or not ([antigen], [antibody], duration of incubation) (Reverberi and Reverberi 2007). Normally, equilibrium is reached quickly. To ensure that equilibrium was reached, however, sections were systematically left in primary antibody solution overnight. Due to the importance of immunocytochemistry for the proposed thesis, several aspects are considered further in the discussion. In the following, I include a comment on selected antibodies of importance used in the thesis – particularly those used for immunogold experiments. Complementary information can also be found in the method sections of the respective articles (I-V).

Anti-AQP4 antibody (Sigma A5971)

In the case of the commercially available anti-AQP4 antibody used in the immunogold experiments, it is directed at the C-terminal of AQP4 and does not, therefore, differentiate between M1 and M23 isoforms, nor would it selectively label any of the other proposed AQP4 isoforms (Moe et al. 2008). No labelling is detected upon incubation of sections from *Aqp4* $-/-$ mice, suggesting a near complete absence of unspecific fractions in the dilutions used. In addition, freeze fracture replica immunogold studies have shown these antibodies to label square arrays (Rash et al. 1998).

All antibodies employed were polyclonal, with the accompanying uncertainty to which epitopes they bind to and convey information about. A comparison between regions examined in article II is therefore based on the assumption that epitope availability is identical across the examined regions. With the methods at hand, no control for this possibility is available. When reference is made to *AQP4 levels* in the included articles and the ensuing discussion, more often than not, this really means *signal strength of anti-AQP4 immunohistochemistry*. This is perhaps the most significant uncertainty in all of the reported quantitative immunogold experiments; how does signal strength correspond to protein level. This is a difficult issue to tackle for all four immunogold experiments (AQP4, α -Syn, laminin, agrin), and no absolute correlation can be made with the controls at hand. Theoretically, it might be possible to provide such a correlation using radiolabelled proteins. This would require a systematic study in its own right, the success of which would among other variables depend on the sensitivity of radiodetection. In the case of AQP4, an alternative approach is proposed in the first part of the discussion. At light microscopy level, findings from the above reported antibody were also reproduced (not shown) using the commercially available anti-AQP4 (C-19) antibody from Santa Cruz Biotechnology (Cat# sc-9888).

Anti-pan-laminin

Knock out lines were not available at the time when the current work was carried out. The anti-pan-laminin signal observed on immunoperoxidase labelling of thick sections and immunogold labelling of ultrathin sections is similar, whereby perivascular labelling is confined to the perivascular basal lamina (article IV; Lunde et al. 2015). Similar labelling is also seen in

glia limitans (article V, fig 4). The precise localisation of anti-pan-laminin signal to the basal lamina indicates absence of unspecific labelling.

Anti-agrin

Three other commercially available antibodies directed against agrin yielded no labelling in the described assays, despite protein recovery. This included: anti-Agrin Millipore cat# MAB5204, anti-Agrin (H-300) from Santa Cruz Cat# sc-25528 and anti-Agrin mAb (Agr 13) from Enzo life sciences cat#ADI-AGR-530. The anti-agrin antibody used in article IV and V was a kind gift from Professor Markus A Rüegg at the University of Basel.

Anti- α -Syn (SYN 259)

The antibodies were a kind gift from Marvin E. Adams and were developed in that laboratory. Transgenic α -Syn^{-/-} mice were used as negative controls for all assays. At the light microscopy level, this antibody does label sections from α -Syn^{-/-} and the antibody does, therefore, contain an unspecific fraction. The perivascular labelling is completely abolished in α -Syn^{-/-} stained sections, however, and this is the only localisation where the anti- α -Syn signal was quantified. Because of the unspecific fractions, however, I quantified the signal detected on α -Syn^{-/-} sections, and was also blinded for genotype upon conducting the experiments using this antibody (Hoddevik et al. 2017). As shown in the α -Syn immunogold experiments reported under discussion (Figure 6, graph with heading “synKO control”), there is no detectable anti- α -Syn signal in α -Syn^{-/-} tissue, irrespective of the signal strength in WT tissue. In article IV we also validated the antibodies by employing commercially available anti- α -Syn antibodies on western blot (Lunde et al. 2015).

Antigen retrieval techniques

The ultrastructural examinations of laminin and agrin were conducted without any antigen retrieval, and ample labelling was identified in the anticipated locations (BLs). A striking feature of both anti-agrin and anti-pan-laminin antibodies when used on thick sections, however, was the reliance upon pepsin digestion to amplify signal strength in adult mouse brains. Without any antigen recovery, only sections from P0-P7 developmental stages will stain, while hardly any labelling is visible in adult mouse brains. This phenomenon has been described previously for other vasculature markers, including collagen IV, laminin and isolectin B4 (Franciosi et al. 2007). In a systematic approach I tested other conditions (pH, temperature, citric acid, NaCl concentrations) where no other parameter had any significant impact. Aldehyde-based fixatives, the likes of those used in all the included articles, react with basic amino acids, and principally lysine and arginine. This generates intra- and intermolecular interactions that crosslink and stabilise proteins in situ. Such interactions may alter existing, antigenic epitopes and inhibit antibody access (MacIntyre 2001; Shi et al. 1997). One might speculate that the pepsin-induced proteolysis exposes epitopes that would not otherwise have bound to the antibodies. Further, it is not possible to rule out that the same proteolysis might generate new, unspecific epitopes.

Controls for immunohistochemistry

The studies of AQP4 and α -Syn were both conducted with negative controls provided for by the respective transgenic knock out strains. In contrast, we did not have available knock out tissue for agrin and laminin. Thus, the immunohistochemistry directed against agrin and pan-laminin lack optimal negative controls, albeit that the anti-agrin antibodies were tested by Eusebio and co-workers on knockout material previously (Eusebio et al. 2003). Previous reports on agrin-deficient mice have shown lacking postsynaptic specialisations and animals die at birth of respiratory failure (Gautam et al. 1996; Lin et al. 2001), and knockout material from adult mouse brain is therefore not available. It should be taken into account that cross-reactivity for antibodies is common and that identity of the cross reacting molecular species

are rarely determined (Danbolt et al. 2016; MacIntyre 2001). The implication is that pitfalls in immunohistochemistry occur commonly. In the absence of robust, negative controls, one cheap and simple strategy is to use at least two different antibodies directed against the same protein/epitopes. Similar results would indeed strengthen the conclusion(s). To this end, I tested several commercially available anti-agrin antibodies, without any resulting labelling strong enough to enable further use. The antibodies were also assessed through SDS-PAGE and western blotting, reported in the respective articles (anti-pan-laminin and anti-agrin in article IV).

SDS-PAGE and western blotting

Homogenates are reported as either “whole” brain or regional. Dissections were made under a dissection microscope but carry a certain risk of variation. Depending on the antigen in question, I used either pre-cast gels (Biorad) or gels cast with the mini-protean system where both stacking- and resolving gel contained 9M urea as described in Sørnbø et al from 2007 (Sorbo et al. 2007). This latter strategy permits separation of 3 distinct bands detected by probing with anti-AQP4 antibodies, corresponding to M1, M23 and Mz isoforms, respectively. These bands merge into one single band upon using non-urea containing, pre-cast gels.

PCR analysis

We performed quantitative real-time PCR on regional dissections to provide a quantitative assessment of mRNA levels. qPCR is not a substitute for western blotting, but in article II, findings of qPCR and western blot directed at AQP4/*Aqp4* were consistent. There have been reports of isoform specific *Aqp4* probes. We chose probes that do not differentiate between M1 and M23 isoforms.

With regards to laminin probes, we limited our assays to α -1 and α -2 isoforms. This choice was made due to the α -subunits being largely responsible for cell surface adhesion and receptor interactions (Yurchenco 2011). One particular challenge for the use of RT-qPCR in article IV resided in how to choose reference genes, and relate findings to total protein content in this study of the developing mouse brain at different milestones.

5. Summary of individual articles

Article I: Mislocalisation of AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy. This study was undertaken to assess the temporal aspect in AQP4 and α -Syn changes shown in other epilepsy materials. The study thus investigated whether expression and distribution of AQP4 and α -Syn were altered prior to or after the onset of chronic epileptic seizures in the kainate (KA) model of mesial temporal lobe epilepsy. Quantitative immunogold labelling revealed that AQP4 density in adluminal endfoot membranes was reduced in KA treated rats already in the latent phase, while the AQP4 density in the abluminal endfoot membrane was stable or slightly increased. The decrease in adluminal AQP4 immunogold labelling was accompanied by a reduction in the density of AQP4's anchoring protein α -Syn. The latent and chronic phases were associated with an upregulation of the M1 isoform of AQP4, as judged by semi-quantitative Western blot analysis. Taken together, the findings suggest that a mislocalisation of AQP4 is an integral part of the epileptogenic process but it remains to explain why and how AQP4 redistributes in the latent phase of this experimental model.

Article II: Factors determining the density of AQP4 water channel molecules at the brain-blood interface. This study was conducted in order to examine the relationship between AQP4 and α -Syn levels in adult mouse CNS. This article provides evidence that the

size of perivascular AQP4 differs considerably between brain regions, roughly mirroring regional differences in *Aqp4* mRNA copy numbers. Quantitative immunogold analyses of AQP4 and α -Syn in pericapillary astrocyte membranes indicate that there is a fixed stoichiometry between these two molecules. Findings show that a targeted deletion of α -Syn removes a substantial and fairly constant proportion (79–94 %) of the perivascular AQP4 pool across the CNS. Both molecules occur at higher densities in endfoot membrane domains facing pericytes than in endfoot membrane domains facing endothelial cells. Thus, several factors seem to influence the content of AQP4 in astrocyte endfeet.

Article III: Mechanisms underlying AQP4 accumulation in astrocyte endfeet. This study was commenced in order to study astrocyte polarisation and unravel mechanisms underlying AQP4 accumulation in astrocyte endfeet. To this end, the study makes use of a tailor made co-culture system. Evidence suggests that endothelial cells promote AQP4 accumulation by exerting an inductive effect through ECM components such as agrin, as well as through a direct mechanical interaction with the endfoot processes. Through the compounds they secrete, the endothelial cells also increase AQP4 expression. The present data suggest that the highly specialised gliovascular interface is established through inductive processes that include both chemical and mechanical factors.

Article IV: Postnatal development of the molecular complex underlying astrocyte polarisation. This study was undertaken in order to test whether observations from cell culture experiments are consistent with in vitro experiments. The article therefore examines the developmental profile of those proteins thought to contribute to the endfoot-basal lamina junctional complex (EBJC). Different members of this complex exhibit distinct ontogenic profiles; ECM proteins laminin and agrin appear earlier than the other members of the complex. Specifically, while laminin and agrin expression peak at P7, quantitative immunoblot analyses indicate that AQP4, α -Syn, and the inwardly rectifying K⁺ channel Kir4.1 expression increases towards adulthood. Findings are consistent with ECM having an instructive role in establishing astrocyte polarisation.

Article V: Organisation of extracellular matrix proteins laminin and agrin in pericapillary basal laminae: insensitivity to targeted deletion of α -syntrophin. This study was done on brains from adult mice to assess regions and subregions with known heterogeneous expression of AQP4 and α -Syn for potential correlations with ECM proteins. The study thus provides quantitative immunogold labelling of agrin and pan-laminin in cortex and cerebellum in mouse brain and compares different BL domains at the pericapillary level. Data from WT mice show that agrin and laminin are constitutively expressed in pericapillary BLs and in meninges immediately overlying glia limitans. Evidence is provided of significant regional differences in both agrin and laminin levels. Quantitative real-time PCR demonstrates significant regional differences for *laminin α -2* and *agrin* mRNA when comparing cortex to cerebellum. Similar differences for laminin and agrin are seen at the local, pericapillary level and correlate well with previously measured levels of AQP4 and α -Syn. Notably, the increased laminin levels in the proximity of pericytes constitute a possible mechanistic explanation of how AQP4 levels are increased in the adjoining astrocyte endfoot membrane domain. Overall, findings are consistent with the idea that ECM proteins agrin and laminin enable membrane compartmentalisation in astrocyte endfeet. Furthermore, targeted deletion of α -Syn leaves laminin and agrin distribution unaltered. This supports the hypothesis that targeted deletion of α -Syn affects AQP4 redistribution via a direct effect rather than indirectly via changes in ECM proteins agrin and laminin.

6. Discussion

The findings are presented and discussed in three parts. First, I review factors that regulate perivascular AQP4 membrane density and influence localisation. The second part concerns the potential heterogeneity of the gliovascular interface, what evidence the current work provides in that regard and a brief account of potential implications for neurophysiology. Finally, the third part is a synthesis aimed at discussing the relevance of findings for selected neuropathologies, particularly NMO and epilepsy.

I: Factors regulating perivascular AQP4 density and that influence localisation to astrocyte endfeet

The challenges, limitations and possible ameliorations of the reported immunogold experiments are listed and discussed under methodological considerations. They apply to all elements of the discussion. When these factors are taken into account, the data should still be interpreted and possible explanatory models offered. The role of endothelial cells, pericytes and the basal lamina is discussed first, thereafter ECM proteins laminin and agrin along with the intracellular α -Syn. Mention is also made of mechanisms that may alter AQP4 levels but which have not been examined directly in articles I-V.

1. Do endothelial cells, pericytes and basal lamina regulate the perivascular AQP4 density in astrocyte endfeet?

1.1: Brain endothelium

Perivascular astrocyte endfeet surround endothelium. Further, AQP4 is constitutively expressed in the endfoot membrane abutting the BL and endothelium. Upon reflecting away from this BL, the AQP4 membrane density immediately decreases. For these reasons did we assess what effect brain derived endothelial cells had on astrocytes in culture. We used the cancer cell line bEND3 as a proxy of endothelium and co-cultured the two cell types in our tailor made cell culture system. As shown in article III, the plasma membrane compartmentalisation of AQP4 increased more than threefold in the presence of bEND3. No such difference was observed when employing astrocytes with a targeted deletion of α -syn, suggesting that the effect endothelial cells exert on astrocytes depend on this latter protein, and by implication the dystrophin-dystroglycan complex. One might speculate that the effect is mediated through the ECM proteins that endothelia secrete. In the same article, we replaced the bEND3 cells with HeLa cells and cells from the umbilical cord (ECV). Interestingly, each cell line induces polarised AQP4 expression in astrocytes, but only co-culture with the bEND3 cell line induced a statistically significant change compared with control experiments. The data cannot determine whether the observed effect of HeLa and ECV cells is mediated in a similar manner to that of bEND3 cells. In our cell culture based assay, the effect of bEND3 cells was limited to endfeet. No change was observed in AQP4 levels in soma, thus suggesting a local effect on endfeet alone.

Kimelberg debates the extent to which findings from astrocyte cell cultures can be extrapolated to the in vivo situation (Kimelberg 2010). Cultured astrocytes are not identical to astrocytes in intact brain, not even primary cultures. One example to illustrate differences between parenchymal- and cultured astrocytes is the pronounced cytoplasmic occurrence of AQP4 in the latter, which is not present in vivo, but which does carry advantages when studying membrane localisation and vesicular transport (Potokar et al. 2013). Further, our cell culture based assay could perhaps have been ameliorated by employing primary cultures of endothelium rather than bEND3 cells as a proxy. This latter, immortalised cell line has been described thoroughly, however, and it maintains BBB characteristics (Omidi et al. 2003). With regards to the role of endothelium towards increasing perivascular AQP4 membrane density,

evidence in article II is in line with what is shown in article III. We did not examine characteristics of endothelia in the brain regions reported on in article II, however, and the evidence is therefore of limited value when evaluating any effect of endothelium.

1.2 Pericytes

The first article to report increased levels of AQP4 in the vicinity of pericytes examined astrocyte endfeet of cortical astrocytes (Gundersen et al. 2013). In article II did we provide evidence that this enrichment is statistically significant and present in all examined regions. Both references made use of the same anti-AQP4 antibodies, and the same experimental protocol in performing the immunogold labelling. Showing the same phenomenon with several antibodies would have strengthened the findings, but the lack of residual labelling on *Aqp4*^{-/-} tissue suggests absence of unspecific fractions. The reader is referred to figure 6 displayed in this discussion (part III.2 under considerations on NMO) which demonstrates the consistent increase of α -Syn in vicinity of pericytes across all examined brain regions.

As measured by quantitative immunogold labelling in WT mice, the overall AQP4 linear density is 15% higher for endfoot membrane domains abutting pericytes than for endfoot membrane domains abutting endothelium (Article II, Fig. 6b, $p < 0.001$). Gundersen and co-workers demonstrated that the increase of α -Syn in vicinity of pericytes persists upon genetic deletion of *Aqp4*. We show that this is also the case for AQP4 increase following α -Syn knockout. Overall AQP4 linear density is 16 % higher in endfeet abutting pericytes compared with adjacent membranes abutting endothelium ($p < 0.05$) in α -Syn^{-/-} animals. The nominal difference of linear density between these two membrane domains is decreased in α -Syn^{-/-} mice, however, and a statistically significant difference ($p < 0.05$) was only observed in CX.

In article V do we provide a potential mechanistic explanation of how the enrichment in AQP4 and α -Syn can be achieved. Anti-pan-laminin immunogold labelling is significantly more intense in the BL interlaced between astrocytes and pericytes compared with the other perivascular BL. A similar trend is also seen for agrin (Fig 2, Article V). In the same article, however, we have interpreted the data as evidence that the source of these two proteins may be astrocytes themselves. A case in point is the labelling at the glia limitans where no pericytes are present. The study design does not allow a firm conclusion in this regard and the data does not exclude the possibility that pericytes also secrete laminin and agrin. Finally, none of the provided evidence can shed light on why the levels of AQP4, α -Syn, agrin and laminin should be increased in the vicinity of pericytes. AQP4 is present in abundance also in endfeet abutting endothelium. The presence of pericytes is therefore not a necessary factor for AQP4 localisation to endfeet. But the systematic, relative increase of about 15 % compared with neighbouring endfeet is puzzling. If the physiological needs of pericytes can be met with a mere 15 % increase, it suggests a finely tuned molecular regulation. What those physiological needs may be cannot be determined based on articles II and V. It would have been interesting, however, to study pericytes and the aforementioned effects in the cell culture system employed in article III. At the time when scientific work for article III was undertaken, no knowledge of the pericyte effect had been reported. Further, other endfoot proteins should be studied as internal controls to AQP4 in order to establish the validity of the findings. Possibility exists that the presence of pericytes constitutes an additional diffusion barrier for water and that this underlies the changes observed in article II and V.

1.3. Pericapillary basal lamina

We focused on the link between DAP and ECM. We therefore acquired and crossed two commercially available transgenic mouse strains to create GFAP-cre x floxed DAG1 (Cohn et al. 2002; Moore et al. 2002) animals. Data on the DAG1 knockout animals is not presented in this thesis but mentioned as an example of the scientific approach towards studying astrocyte-ECM interactions. The aim was to study the impact of β -DG gene deletion in astrocytes. Alas, only two of the acquired breeding animals yielded offspring, and the litters

were much smaller than previously reported. Breeding challenges may have been related to the crossing of several genetic backgrounds (129 x C57Bl6 x FVB), effectively yielding a substrain. At the time, GFAP-cre mice were not commercially available on a C57Bl6 background. Backcrossing onto C57Bl6 background for 2 generations markedly increased litter size and eradicated unexpected death like that seen initially in a subpopulation of animals (white fur). Further backcrossing was not possible due to closure of the local animal facility and the colony had to be terminated. The data acquired from these experiments remains unpublished but was in line with prior freeze-fracture and IF analyses that perivascular AQP4 levels were very low or even absent following selective β -DG (*DAG1*) knockout (Moore et al. 2002; Noell et al. 2011). Quite like the previous examples of endothelium and pericytes, the pericapillary BL is present in all of the imaged capillaries. And as alluded to in the background chapter, the BL is a specialised matrix, composed of a multitude of constituents. It was beyond the scope of the current project to study them all, but articles III, IV and V provide evidence of select ECM components.

2. Proteins that may subserve a regulatory role for perivascular AQP4 density in astrocyte endfeet

Neither articles II nor III attempt to answer why AQP4 is enriched in the proximity of endothelial cells, pericytes and the pericapillary basal lamina. They do, however, attempt to provide an explanation of how it is possible. A summary of findings for candidate proteins α -Syn, agrin and laminin is thus provided in the following paragraph.

2.1 α -Syn

Prior reports have shown that α -Syn serves as a molecular anchor of endfoot AQP4 (Neely et al. 2001) but article II is the first study to show that the densities of α -Syn and AQP4 appear to be stoichiometrically coupled. The nominal ratio between these two molecules as measured by their respective, immunogold labelling intensities varies little among regions. A limitation to this data resides in challenges of translating labelling intensity into exact protein levels. This would necessitate titration curves and linearity between the latter, two factors.

The quantitative RT-PCR analysis in article II revealed pronounced regional differences in the amount of *Aqp4* mRNA. The *Aqp4* transcript levels were high in those regions that displayed high levels of AQP4 in perivascular endfeet and were not sensitive to α -Syn deletion.

That AQP4 and α -Syn covary seems not to be a coincidental correlation. Rather, we chose to interpret this as causality: removal of α -Syn via genetic deletion removes a substantial fraction of endfoot AQP4 that varies little across regions. The possibility that α -Syn regulates *Aqp4* expression through regulation at the transcriptional level is ruled out by our finding that the number of *Aqp4* mRNA copies remains constant in α -Syn^{-/-} animals. We interpret these findings as evidence that the number of α -Syn molecules targeted to the endfoot membrane is the main determinant for AQP4 density around the CNS capillary bed.

Experiments in article II show that the level of AQP4 in endfeet is correlated to the level of *Aqp4* mRNA in the corresponding region. This observation does not necessarily imply that the level of endfoot AQP4 is controlled at the transcriptional level. But recent evidence of local protein translation poses interesting questions regarding the role of transcriptional regulation compared with those other mechanisms that exist. The transcriptional regulation of *Aqp4* has been shown to be complex (Pisani et al. 2011; Rossi et al. 2010) and requires further studies. Despite the evidence provided that α -Syn is a factor involved in regulating AQP4 levels, interactions between these two proteins remain incompletely understood. As an example, it is not known how many α -Syn proteins bind to each AQP4 molecule, be that M1, M23 or Mz. In line with previous methodology to study OAPs, it would seem natural to employ freeze fracture techniques on α -Syn^{-/-} tissue, while no such report is hitherto published. The role of α -Syn in OAP assembly is therefore unresolved.

2.2 laminin

In article III we applied laminin to primary astrocyte cultures. This did not induce any change in AQP4 compartmentalisation, nor did the co-application of laminin with bEND3. The relative increase in proximity of pericytes is mentioned above.

Laminin and agrin are the only two ECM proteins I have studied, and the interpretation of the data rests on the strength of the immunosignal detected. To strengthen this evidence, article V would have benefited from internal controls. Ideally suited candidates would have been ECM proteins shown not to alter AQP4 membrane density. Collagen IV could perhaps have been one such control. The same reasoning applies to article III, although the cell culture experiments provide stronger experimental evidence to establish agrin and laminin as candidates for AQP4 membrane density regulation than do the descriptive data of article V.

2.3 agrin

As shown in article III (Fig 7), agrin by itself did not induce any significant change in AQP4 compartmentalisation. But when co-applied with bEND3, cells gave rise to a more pronounced polarisation than what was seen with bEND3 cells alone. The presence of endothelial cells (bEND3) therefore is a necessary factor for agrin to exert its effect on AQP4 membrane density in astrocytes. Interestingly, agrin on its own was able to induce the same magnitude of AQP4 membrane enrichment that did the application of Matrigel. In contrast, Matrigel contains a host of ECM proteins, including proteoglycans. This response to agrin has been shown by others (Noell et al. 2007).

Article IV demonstrates that agrin is indeed present in the capillary BL from even early stages of development in mouse brain. And in article V do we provide a quantitative assessment of agrin in the capillary bed of three distinct brain regions. The difference in agrin levels when comparing the three perivascular BL microdomains only shows a trend towards higher levels in the vicinity of pericytes. Although not statistically significant, this is still consistent with the idea that it exerts an effect on neighbouring endfeet membranes. Negative results from appropriate, internal controls in articles III, IV and VI would have strengthened these findings.

3. Other potential factors and mechanisms

3.1 Mechanical stimulation

Pericytes are contractile cells. The slightly increased protein levels (α -Syn, AQP4, laminin and agrin) in their vicinity may occur so as to compensate for the physical strain this imposes on neighbouring cells and basal lamina. Mechanical stress exerted by pericytes on astrocyte endfeet may also be the mechanism by which the enrichment in AQP4, α -Syn, agrin and laminin are induced. This might involve the existence of mechanoreceptors in endfeet that in turn would have to be coupled to one/more of the gene transcription, protein translation and secretion in question.

Alterations in agrin and particularly in laminin levels would potentially also alter the functional properties of the BL proper. Basal laminae that are self-assembled through interactions between the constituent proteins and BLs can be considered as stabilising extensions of the plasma membrane, which protects tissues from disruptive physical stresses. There is also evidence to suggest that the organisation of the BL, including its density and surface patterning, can be dynamically modulated by the cytoskeleton through its receptor connections (Colognato et al. 1999).

3.2 Protein turnover

The degradation of AQP4 has been tested previously. Evidence suggests the half-life of wild-type AQP4 expressed in cell cultures is approximately 24 hours, a degradation which increases with removal of SSV (PDZ-binding motif) near the C-terminus (Neely et al. 2001). With the proposed binding of α -Syn to this PDZ domain, this has been taken as evidence to show how DAPs influence AQP4 levels also through protein turnover. Thus, turnover and

production of the other dystrophin-associated proteins can also be a way in which to down- or perhaps also upregulate AQP4.

Under ischemic conditions, the temporary loss of perivascular AQP4 occurred without a similar change in dystrophin and α -Syntrophin (Frydenlund et al. 2006). This suggests AQP4 loss is due to altered protein-protein interactions at the local level. Findings from article I are a contrast to this. The immunogold analysis in article I revealed that AQP4 density in adluminal endfoot membranes was reduced in KA treated rats already in the latent phase, while AQP4 density in the abluminal endfoot membrane was stable or slightly increased. The decrease in adluminal AQP4 immunogold labelling was accompanied by a reduction in the density of α -Syn. Thus, the KA induced changes in AQP4 membrane density may reside in changes related to protein production. And what determines the reduction in perivascular AQP4 levels is – or involves/correlates with – α -Syntrophin.

Frydenlund et.al reported regional differences in recovery of perivascular AQP4 that differ between the neocortical and the striatal part of the ischemic core. What underlies this regional difference may reside in differential regulatory mechanisms and concomitant variations in response to ischemia.

3. Limitations of the conducted experiments

The reported studies examine a selection of components at the gliovascular interface, but only at the capillary level. A further limitation is that most of the data is descriptive and thus void of functional assessments. In studying interactions between the ECM and astrocyte endfeet, the provided experiments only examine dystrophin-dystroglycan related proteins. A rigorous approach in studying astrocyte- ECM interaction would have included the other known transmembrane receptor family known to bind laminin, namely integrins. The extrapolation of data from rodents to the human case was described by Oberheim et.al, whereby human astrocytes are significantly larger than their rodent counterparts. A recent study also found that human and mouse cortical astrocytes differ in their polarisation towards microvessels (Eidsvaag et al. 2017). But since the adluminal membrane of human endfeet is enriched in OAPs and AQP4 like that in mouse and rat CNS, this is rather evidence to support that extrapolation to the human case is possible.

While considerable data is included from the gliovascular interface, AQP4 is also expressed in abundance in periventricular ependymal cells and at the glia limitans. Only 50% of the AQP4 at the glia limitans is lost following α -Syn deletion (Amiry-Moghaddam et al. 2004). The findings from article II on gliovascular AQP4 are thus not applicable to the entire mouse CNS.

AQP4 expression is not limited to the CNS. This defines possibilities for studying mechanisms of regulation and physiological roles. As an example from muscle, both neuronal innervation (Ishido and Nakamura 2018) and exercise induces AQP4 modulation. Here, AQP4 supports muscle contractile activity and metabolic changes that occur in fast-twitch skeletal muscle during prolonged exercise (Basco et al. 2013). Controversy still exists, though, since not all fast-twitch fibres seem to express AQP4 (Kaakinen et al. 2007). We did not include any non-CNS tissue expressing AQP4 in the proposed articles I-V, but this could have represented ameliorations to our studies.

II: Evidence for and against regional heterogeneity of the gliovascular interface; implications for neurophysiology

In clinical practice, a neurologist will first attempt to answer the question of *where* the lesion is located, before considering *what* the diagnosis is. This is in contrast to all other diagnostic approaches in clinical medicine. In neurological practice, the site of the disease is what dictates the list of differential diagnoses. For this reason, this second part of the discussion is included. Emphasis is placed on the evidence suggesting regional heterogeneity of AQP4. A brief comment is included also on ECM proteins agrin and laminin.

1. From representative examples to generalisations

The interpretation of whether the gliovascular interface displays regional heterogeneity is limited to the study of endfeet and the perivascular BL. Further, the study of these structures is based on and limited to the data on AQP4, α -Syn, laminin and agrin. Endfeet serve a myriad of functions, however. These are dependent on proteins other than AQP4 and α -Syn that may or may not correlate with the capacity to transport water and thus AQP4 levels. With regards to K^+ siphoning, the Kir4.1 channel has been shown to co-localise with AQP4 (Nagelhus et al. 1999) and might be affected by AQP4 levels. This has also been shown for TRPV-4 (Benfenati et al. 2011). Apart from these two examples, each physiological function of endfeet would have to be examined separately in order to conclude whether there exists a similar, regional heterogeneity thereof.

2. Evidence of regional heterogeneity for AQP4 and differences in rapid water transport

In article II (Hoddevik et al. 2017) we provide evidence of regional heterogeneity by the use of western blot, RT-qPCR and quantitative immunogold labelling. Collectively, the experiments establish that pericapillary AQP4 levels vary across the mouse CNS and are already established at birth. The sagittal sections of adult and newborn mouse brain (Fig 1, article II) illustrate this well. Such regional differences in endfoot AQP4 imply regional differences in water transport capacity. But immunogold experiments in article II also show how AQP4 is constitutively expressed around the entire capillary bed.

That AQP4 is abundantly expressed in astrocyte endfeet has been shown by freeze fracture replicas (Landis and Reese 1974; Wolburg 1995). Reference articles contain images where impressive portions of endfeet are covered by square arrays. But any quantitative assessment must question what protein level is necessary to fulfil the physiological function of the studied candidate protein, i.e. provide a functional assessment. This prompts the discussion of what the function(s) of AQP4 is (are), and how effective the protein is at fulfilling it or them (Nagelhus and Ottersen 2013). The crystal structure of AQP4 has been resolved (Ho et al. 2009), and the water permeability studied. AQP4 has a very high water permeability when compared with AQP1. Thus, a further specification of the physiological role of AQP4 is implied: AQP4 is not merely a water channel that enables the specific transport of water - rather, it is a channel with very high water permeability and ability to convey rapid fluxes of water (Yang et al. 1997). This molecular property of the channel begs the question of why such pronounced water transport is needed in the CNS, and in the studied pericapillary regions. Three different mechanisms of how water crosses cell membranes have been proposed: 1) by diffusion, 2) via aquaporins, but also 3) via co-transporters and uniports where H_2O transport is coupled to the flux of substrates (MacAulay et al. 2004; Zeuthen 2010). The provided experiments do not measure actual water transport and we have not assessed any co-transporters. We can therefore not ascertain whether there is a strict correlation between pericapillary AQP4 expression and water permeability/transport capacity. Nor can we not rule out the possibility that AQP4 is differentially regulated, depending on region, subregion or membrane microdomain. Several factors have been identified that affect water flux through AQP4 (Gunnarson et al. 2005; Yukutake et al. 2008; Yukutake and Yasui 2009; Zelenina 2010) but direct evidence for gating is still missing (Nagelhus and Ottersen 2013).

In summary, the evidence provided by article II on regional differences in AQP4 levels is hard to dismiss. But a functional correlate to explain why such differences should be necessary is not obvious. The suggested involvement in paravascular flow is a tempting explanation (Iliff et al. 2012). But the immunogold experiments in article II are limited to the capillary bed and disregard venules and arterioles where such paravascular flow has been shown to occur. Should AQP4 prove to be important for waste clearance and sleep as was suggested (Iliff et

al. 2012; Xie et al. 2013), then the regional differences in AQP4 levels shown in article II might still find their functional correlate.

3. How do the examined regions compare to one another?

Studying different brain regions carries the inherent obligation of comparing them to each other. One obvious parameter to discuss is neuronal density. The cerebellar granular layer contains an impressive density of neurons and its capillary bed comparably contains significantly more perivascular AQP4 than does cerebral cortex. This also holds true for hippocampal regions (Article I, Fig 2B). The hypothesis was alluded to by Badaut et.al in a review article. What underlying mechanisms of regulation this implies remains unexplored. Cell density could correlate inversely with extracellular space size, and thus correlate positively with a need to buffer K^+ (Badaut et al. 2002).

Overall, the possibility exists that the reason for heterogeneous AQP4 expression resides in internal factors, i.e., that the perivascular astrocytes themselves differ from one another depending on the examined region. Differential expression of connexins 43/30 with subsequent variability in glial syncytiae and gap junction expression may constitute one such situation. It is a limitation of the present study that the astroglial syncytium was not evaluated simultaneously as pericapillary AQP4 membrane density. Reciprocity in Cx43/30 and AQP4 expression has been shown (Katoozi et al. 2017) and *Aqp4* gene deletion results in upregulation of gap junctions. This constitutes a potential confounding factor and networks of astrocytes might differ from one region to the other. Larger networks may increase the buffering capacity for perivascular astrocytes and thus reduce the need for fast water transport at the gliovascular interface. Conversely, smaller networks may increase reliance upon vascular coupling and the need for fast water transport. Figure 4 below illustrates this hypothesis.

Figure 4: Putative confounding factor for regional heterogeneity is shown through illustration of astrocytes coupled via an endfoot to a brain capillary. A) shows a theoretical glial syncytium composed of one perivascular and 3 parenchymal (protoplasmic) astrocytes. Buffering capacity of this network would have a relative, 4-fold increase compared to the lone perivascular astrocyte in B). Adluminal AQP4 membrane density (red dots) might thus be reduced in the larger glial network, inversely proportional to the networks buffering capacity. L, capillary lumen.

The other possibility is that the observed variation in AQP4 is a result of external factors, from the environment surrounding the cells. Differences in metabolic activity may be one such example. There is evidence of interdependency between local capillary density, blood flow and metabolism in rat brains (Klein et al. 1986; Kuschinsky 1990). Klein et.al. report the inferior colliculus with 811 ± 20.7 capillary sections per mm^2 , the frontal cortex as 456 ± 25.3 , hippocampus as 361 ± 11.0 and cerebellar white matter as 178 ± 6.7 . Cerebellar granular- and molecular layers are alas not included in these reported studies and we did not assess capillary density in article II or V. Also, article II included parietal and not frontal cortex. But with these limitations, the reported capillary density may still correlate with anti-AQP4 immunogold signal and *Aqp4* RT-qPCR data. Even if article II demonstrates statistically

significant differences in AQP4 and α -Syn levels across the CNS, question remains to what extent this is of functional relevance.

4. Does the pericapillary basal lamina display regional heterogeneity?

Extracellular matrix and the basal laminae have traditionally been viewed as rather static entities. The immunogold labelling of ECM proteins agrin and laminin reported in article V illustrates how linear density of these molecules varies substantially depending on which cell type adjoins the BL in question. This was an unanticipated finding. The case has already been made that increased levels of both agrin and laminin might alter the properties of the BL proper, and that this might compensate for increased mechanical stress exerted by pericytes. The changes within the basal lamina proper are greater than the regional differences. Although the immunogold experiments do demonstrate statistically significant differences between cerebral cortex and cerebellum (Fig 3, article V), the difference in labelling intensity is not impressive. Whether they have biological importance or not cannot be ascertained based on these experiments. The RT-qPCR experiments against two laminin isoforms and agrin also demonstrate regional differences. It should be noted, then, that neither of these proteins are specific to perivascular regions and that this approach yields evidence on the general tissue expression, including both parenchymal and perivascular protein pools.

5. Protein correlations at regional level

Previous publications have shown that redistribution of AQP4 correlates with loss of agrin in a human glioblastoma cell line (Warth et al. 2004). A positive correlation of agrin and AQP4 protein levels would have suggested that agrin might not just be a necessary factor, but also serve a regulatory role. Correlations between proteins are not sufficient to deduce functional relationships, however, but may give important indications thereof. Of particular note is the data provided in article IV from developing cortex, consistent with the idea that laminin and agrin influence AQP4 levels *in vivo*. Both agrin and laminin levels peak prior to levels of AQP4, Dp-71 and DAG1. The postnatal development of AQP4 has also been studied previously (Fallier-Becker et al. 2014).

In Figure 5 is a graph displaying the 4 immunogold experiments in articles II and V, organised according to region and arbitrarily normalised to cortical levels. Here, there is no apparent correlation between ECM proteins (agrin and pan-laminin) and astrocyte proteins (AQP4 and α -Syn). The data constitutes evidence in favour of the null hypothesis - that ECM proteins are not instructive to endfeet - but seems insufficient to confirm it.

Normalised data of 4 immunogold experiments

Figure 5: Four immunogold experiments from articles II and V are shown, directed respectively against AQP4, α -Syn, pan-laminin and agrin. The mean linear density (immunogold particle per unit length) for each subregion is normalised (arbitrary value = 1) according to levels measured in cerebral cortex (CX), and shown for molecular- (CB-mol) and granular (CB-gran) cell layers of the cerebellum. Mean linear density for AQP4 and α -Syn covary, while no such correlation is seen when comparing laminin and agrin with AQP4.

Error bars represent standard error of the mean (SEM).

This is substantiated by the fact that such regional assessment cannot differentiate between which molecular mechanisms for AQP4 regulation is involved (agrin, laminin, local protein translation, vesicular transport and DAPs). That the immunogold signal of both AQP4 and α -

Syn correlates with the respective mRNA suggests that endfoot levels of both proteins correlate with protein transcription. At first glance, this sounds obvious, but the hypothesis that ECM in the adjoining basal lamina served a regulatory role would potentially be independent of transcription. Question remains therefore, whether the observed correlation is indeed also evidence against the instructive role of ECM and rather supports other mechanisms involving transcription factors.

III: Potential ramifications of presented findings for select neuropathologies

In this third part I review two clinical conditions to which the findings are of particular relevance. The list of other brain disorders and pathophysiological states involving astrocyte endfeet and the basal lamina is extensive and some important examples are provided.

1. Epilepsy

Article I examines AQP4 and the kainate model of epilepsy on rat samples. Of note to the current thesis is that the reductions of AQP4 and α -Syn levels correlate positively in the latent phase. This is consistent with the fixed stoichiometry between AQP4 and α -Syn as demonstrated in article II. Interestingly, reduction of endfoot AQP4 occurring after cerebral ischemia, in a model of MCAO, occurs with no concomitant change in α -Syn levels (Frydenlund et al. 2006). The results from Frydenlund et.al suggest that the bridging between α -Syn and the cytoplasmic tail of AQP4 is sensitive to ischemia. Question remains, therefore, what (or which) regulatory effect(s) are responsible for the AQP4 and concomitant α -Syn reductions observed in article I, but this may be independent of reduced oxygen levels. The regulatory mechanism(s) at play in the kainate model might be same as the one(s) responsible for the observed regional heterogeneity (article II), providing a potential proof of concept that this might be of relevance in epilepsy.

An overall aim for the presented, five articles, was to make extrapolations from *mus musculus* and *rattus norvegicus* to *homo sapiens sapiens*. Article I on the kainate model of epilepsy is a case in point. Two major challenges exist in this regard. Firstly, the kainate model is a pharmacological way of inducing epilepsy and one may question to what extent this is comparable to the idiopathic and spontaneously occurring temporal lobe epilepsy. Secondly, epilepsy is found in neither mice nor rats, only in certain inbred strains. The human case and that of the dog¹ set aside, the Mongolian gerbil constitutes perhaps the only documented example of naturally occurring epileptic seizures. Such seizures manifest following several stimuli, including stress. It has been hypothesised that the seizures of Mongolian gerbils are perceived by predators as signs of disease thus rendering it an uninteresting prey. It was introduced as an animal model of epilepsy in the 1960s and has since been extensively studied (Fujisawa et al. 2003; Norris 1987). The use of animal models in the study of epilepsy has been reviewed and is not without pitfalls (Avanzini 1995; Fariello 1995; Kupferberg 2001). While post-traumatic epilepsy induced by lateral-percussion fluid induced injury is arguably more representative as a disease model, it also has disadvantages in that the number of study animals needed is much greater since only 43-50 % of injured animals develop the disease, and the latency period varies between 7 weeks to a year following the mechanical insult (Bolkvadze and Pitkanen 2012; Kharatishvili et al. 2006). The kainate model is but one of many ways in which to induce seizures in animals, but it carries the advantage of having a well-defined latent phase and was hence ideal in order to assess the temporal aspect of AQP4 changes that we set out to explore.

¹ The breeding of dogs has been characterised as the biggest biological experiment undertaken by humanity; exemplified by how wolves were bred into Chihuahuas. Of note is that wolves do not suffer epileptic seizures, and the systematic in-breeding set in motion by humans is not a natural occurring evolutionary pressure.

Studies conducted on human hippocampal resections have shown changes in astrocytes, and in AQP4 localisation and levels (Eid et al. 2005; Lee et al. 2004; Salman et al. 2017). The findings from article I cannot ascertain whether the observed changes in AQP4 and α -Syn levels are compensatory or causal in disease development. The only study on epileptic seizure propensity in α -Syn^{-/-} mice (Amiry-Moghaddam et al. 2003b) employed a hyperthermic model and is not comparable to the pharmacological kainate model. Our conclusion that mislocalisation of AQP4 – reflecting a loss of astrocyte polarisation – is an integral part of the epileptogenic process must be tempered by the fact that polymorphisms in *Aqp4* and *α -Syn* genes have not been reported to induce epilepsy per se. But one report exists with evidence that variants of the genes encoding AQP4 and Kir4.1 (as assessed using SNP analysis) are associated with subgroups of mesial temporal lobe epilepsy (Heuser et al. 2010b). Also, *Aqp4* knockout reduced seizure thresholds in another pharmacological model (Binder et al. 2004). This evidence implicates AQP4 in the pathogenesis of epilepsy. But the reported findings in article I are likely to represent but a few of many that together result in development of chronic seizures.

2. NMO

In neurological practice, all neurologists are trained to first answer the question of *where* a lesion is located, before considering *what* the diagnosis is. It is the anatomical site that dictates the list of differential diagnoses. A case in point is neuromyelitis optica (NMO), a demyelinating disorder affecting optic nerve and spinal cord in particular. It has been speculated that the relative abundance and localisation of AQP4 high order arrays in distinct CNS regions may explain this regional preference and variability in clinical phenotype of NMO spectrum disorders (Iorio et al. 2013). This is based on an assumption of how AQP4 levels vary across the CNS. The proposed thesis, as an account of regional heterogeneity, therefore allows discussion in this regard. Figure 6 below is an expanded overview of data from the α -Syn immunogold labelling experiment reported in article II. Note in particular the sparse labelling in optic nerve (far left), compared with other regions. Included is a quantitation from KO tissue where two different regions are included (labelling is not detectable in either, suggesting specificity and lack of unspecific labelling despite increasing epitope levels).

Figure 6: Regions are shown in ascending order according to anti- α -Syn signal strength from immunogold labelling: optic nerve (optic-n), cerebellar molecular layer (cb-mol), olfactory tract (olfract), molecular layer of gyrus dentatus (hcgyrml), cortex piriformis (pirilam1&2), cortex (cx), corpus callosum (corpcall), stratum radiatum (hc-rad), stratum

pyramidale (hc-pyr), gyrus dentatus (hc-gyr), grey matter of spinal cord (spinal-g), stratum oriens (hc-or), colliculus inferior (col-inf), white matter of spinal cord (spinal-w), cerebellar granular layer (cb-gran), cerebellar white matter (cb-white). Blue circles signify mean of linear density (detected immunogold particles per μm cell membrane) in astrocyte membranes facing endothelium, Green circles for astrocyte membranes facing pericytes. *Error bars* represent 95 % confidence intervals. Sagittal section from mouse brain shows regions embedded for immunogold analysis (red circles). Anti- α -Syn immunogold labelling on α -Syn^{-/-} tissue is shown (control, right).

Visualising cross-sectioned white matter on mesh grids is challenging. For this reason, the AQP4 experiment published in article II did not include sufficient images from optic nerve for statistical purposes. The data does open for speculation, however, since both proteins had a fixed stoichiometry in all CNS regions where included images were sufficient for statistical analysis, including white matter of spinal cord. By extrapolation, therefore, pericapillary AQP4 levels in optic nerve (distal to chiasma opticus) are likely to be lower than any other region/subregion mentioned in article II. Notably, pericapillary AQP4 levels in both grey and white matter of spinal cord were among the highest recorded. Thus, a temporary conclusion is that high, perivascular AQP4 levels do not explain the regional affection in NMO spectrum disorders. This scientific problem is perhaps comparable to the enigmatic sparing of skeletal muscle and kidney in NMO patients. On a similar note, the way in which the patient's immune system is sensitised to AQP4, resulting in anti-AQP4 autoantibodies, remains unclear. AQP4 expression is not limited to the CNS. The protein is found in skeletal muscle (Frigeri et al. 1998; Shibuya et al. 2006), cardiomyocytes (Butler et al. 2006; Rutkovskiy et al. 2012) in the collecting duct principal cells of kidney (Terris et al. 1995) and occurs in abundance in nasal/olfactory epithelium (Sorbo et al. 2007). AQP4 protein from any one of these sites could be responsible for sensitisation.

If NMO-IgG binding to AQP4 is indeed causal, the explanation of why NMO patients primarily develop lesions in optic nerve and spinal cord must have an alternate explanation. The experiments conducted in article II do not permit conclusions regarding differences in epitope availability, however. In-depth studies of NMO-IgG labelling is required to tackle this problem. We therefore acquired two batches of such antibodies from collaborators in Lyon and Italy with the hope to examine this at the ultrastructural level, on mouse and rat samples. Unfortunately, hardly any labelling was seen using either antibody, despite implemented optimisation for conserving epitopes and increasing antibody-epitope binding (pH-shift fixation using paraformaldehyde only, NaCl concentrations). Any tissue exposure to paraformaldehyde abolished the labelling, and sufficient PFA is a pre-requisite for conducting immunogold experiments. As this paragraph alludes to, the exact pathogenesis of NMO remains to be fully explored. Particularly so the role of astrocytes and endfeet containing the antigen that is now integral to the diagnosis (Lennon et al. 2005). The explanation may rather reside in the cascade of events that follow antigen-epitope binding, such as complement regulation and activation (Yao and Verkman 2017a). By implication – the neuroimmunological response might differ according to brain region. Either way, the combination of scientific publications on AQP4 together with those on demyelinating diseases may herald ground-breaking knowledge for the field of neuroimmunology and the involvement of astrocytes in demyelinating disorders.

3. Other neuropathological states

AQP4 has been studied in the context of many neurological disorders. The potential for making therapeutic use of AQP4 was recently reviewed (Tradtrantip et al. 2017; Verkman et al. 2017). The list of clinical conditions in which changes in AQP4 levels have been documented grows long and among others include Alzheimer's disease (Yang et al. 2011; Yang et al. 2017), epilepsy (Binder et al. 2004; Eid et al. 2005; Lee et al. 2004), post-ischemic oedema formation (Frydenlund et al. 2006; Manley et al. 2000), obstructive and normal pressure hydrocephalus (Bloch et al. 2006; Eide and Hansson 2017; Feng et al. 2009; Hasan-Olive et al. 2018), glioblastoma (Noell et al. 2012; Warth et al. 2004), bacterial meningitis (Papadopoulos and Verkman 2005) and cerebral malaria (Promeneur et al. 2013).

Data has also emerged on polymorphisms and mutations in the human AQP4 gene that may cause disease directly (Berland et al. 2018), and both predispose or protect against oedema development (Sorani et al. 2008b).

There is thus a broad and compelling evidence base which implicates AQP4 as a potential drug target. The challenges, however, are equally well documented. A good example is oedema, where AQP4 contributes to both resolution and formation (Manley et al. 2004; Papadopoulos et al. 2004). As but one example, the temporal aspect of any AQP4 blockade must be resolved to realise any therapeutic benefit.

7. Conclusions and future perspectives

The proposed thesis provides evidence of hitherto unexplored and significant heterogeneities of the gliovascular interface across the mouse CNS. The case of AQP4 is particularly intriguing and implications for physiology remain to be fully explored. While no conclusive evidence towards a unifying, regulatory mechanism for membrane AQP4 levels is identified, the present work provides evidence that α -Syn, agrin and laminin all contribute to regulation of perivascular AQP4 membrane density. In addition, proximity to pericytes seems to upregulate AQP4 concentrations – which may be mediated through ECM proteins laminin and agrin. The original hypothesis underlying this work was that ECM is instructive to membrane composition of endfeet. Data from the developing and mature mouse brain give indications that this is a viable hypothesis.

The physiological role of AQP4 is still being investigated and the exciting finding that AQP4 is involved in sleep and waste clearance has rejuvenated efforts to solve the problem. The role of AQP4 in NMO pathogenesis may culminate in the long anticipated coming of therapies directed at this protein. The findings presented herein may have ramifications for both of these topics. It will be interesting to see if and when the regional differences in AQP4 levels shown in article II find their functional correlate.

8. References

- Ablimit A, Aoki T, Matsuzaki T, Suzuki T, Hagiwara H, Takami S, Takata K (2008) Immunolocalization of water channel aquaporins in the vomeronasal organ of the rat: expression of AQP4 in neuronal sensory cells *ChemSenses* 33:481-488 doi:bjn015 [pii];10.1093/chemse/bjn015 [doi]
- Agre P et al. (2002) Aquaporin water channels--from atomic structure to clinical medicine *J Physiol* 542:3-16
- Alexopoulos H et al. (2015) Anti-aquaporin-4 autoantibodies in systemic lupus erythematosus persist for years and induce astrocytic cytotoxicity but not CNS disease *J Neuroimmunol* 289:8-11 doi:10.1016/j.jneuroim.2015.10.007
- Allen A, Messier C (2013) Plastic changes in the astrocyte GLUT1 glucose transporter and beta-tubulin microtubule protein following voluntary exercise in mice *Behav Brain Res* 240:95-102 doi:10.1016/j.bbr.2012.11.025
- Alport AC (1927) Hereditary Familial Congenital Haemorrhagic Nephritis *Br Med J* 1:504-506
- Alvestad S, Hammer J, Hoddevik EH, Skare O, Sonnewald U, Amiry-Moghaddam M, Ottersen OP (2013) Mislocalization of AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy *Epilepsy research* 105:30-41 doi:10.1016/j.eplepsyres.2013.01.006
- Amiry-Moghaddam M et al. (2005) Brain mitochondria contain aquaporin water channels: evidence for the expression of a short AQP9 isoform in the inner mitochondrial membrane *FASEB J* 19:1459-1467 doi:19/11/1459 [pii];10.1096/fj.04-3515com [doi]
- Amiry-Moghaddam M et al. (2003a) An alpha-syntrophin-dependent pool of AQP4 in astroglial end-feet confers bidirectional water flow between blood and brain *Proceedings of the National Academy of Sciences of the United States of America* 100:2106-2111 doi:10.1073/pnas.0437946100
- Amiry-Moghaddam M, Ottersen OP (2013) Immunogold cytochemistry in neuroscience *Nature neuroscience* 16:798-804 doi:10.1038/nn.3418
- Amiry-Moghaddam M et al. (2003b) Delayed K⁺ clearance associated with aquaporin-4 mislocalization: phenotypic defects in brains of alpha-syntrophin-null mice *Proceedings of the National*

- Academy of Sciences of the United States of America 100:13615-13620
doi:10.1073/pnas.2336064100
- Amiry-Moghaddam M et al. (2004) Alpha-syntrophin deletion removes the perivascular but not endothelial pool of aquaporin-4 at the blood-brain barrier and delays the development of brain edema in an experimental model of acute hyponatremia *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* 18:542-544 doi:10.1096/fj.03-0869fje
- Andriezen WL (1893) The Neuroglia Elements in the Human Brain *Br Med J* 2:227-230
- Araque A, Parpura V, Sanzgiri RP, Haydon PG (1999) Tripartite synapses: glia, the unacknowledged partner *Trends in neurosciences* 22:208-215
- Armulik A et al. (2010) Pericytes regulate the blood-brain barrier *Nature* 468:557-561 doi:10.1038/nature09522
- Arthur FE, Shivers RR, Bowman PD (1987) Astrocyte-mediated induction of tight junctions in brain capillary endothelium: an efficient in vitro model *Brain Res* 433:155-159
- Aumailley M et al. (2005) A simplified laminin nomenclature *Matrix Biol* 24:326-332 doi:10.1016/j.matbio.2005.05.006
- Avanzini G (1995) Animal models relevant to human epilepsies *Ital J Neurol Sci* 16:5-8
- Badaut J, Hirt L, Granziera C, Bogousslavsky J, Magistretti PJ, Regli L (2001) Astrocyte-specific expression of aquaporin-9 in mouse brain is increased after transient focal cerebral ischemia *J Cereb Blood Flow Metab* 21:477-482 doi:10.1097/00004647-200105000-00001
- Badaut J, Lasbennes F, Magistretti PJ, Regli L (2002) Aquaporins in brain: distribution, physiology, and pathophysiology *J Cereb Blood Flow Metab* 22:367-378 doi:10.1097/00004647-200204000-00001
- Badaut J, Petit JM, Brunet JF, Magistretti PJ, Charriaut-Marlangue C, Regli L (2004) Distribution of Aquaporin 9 in the adult rat brain: preferential expression in catecholaminergic neurons and in glial cells *Neuroscience* 128:27-38 doi:10.1016/j.neuroscience.2004.05.042 [doi];S0306452204003951 [pii]
- Barres BA (2008) The mystery and magic of glia: a perspective on their roles in health and disease *Neuron* 60:430-440 doi:10.1016/j.neuron.2008.10.013
- Barresi R, Campbell KP (2006) Dystroglycan: from biosynthesis to pathogenesis of human disease *Journal of cell science* 119:199-207 doi:10.1242/jcs.02814
- Basco D et al. (2013) AQP4-dependent water transport plays a functional role in exercise-induced skeletal muscle adaptations *PLoS One* 8:e58712 doi:10.1371/journal.pone.0058712
- Bechmann I, Galea I, Perry VH (2007) What is the blood-brain barrier (not)? *Trends Immunol* 28:5-11 doi:S1471-4906(06)00329-2 [pii];10.1016/j.it.2006.11.007 [doi]
- Bell RD, Winkler EA, Sagare AP, Singh I, LaRue B, Deane R, Zlokovic BV (2010) Pericytes control key neurovascular functions and neuronal phenotype in the adult brain and during brain aging *Neuron* 68:409-427 doi:10.1016/j.neuron.2010.09.043
- Benfenati V, Caprini M, Dovizio M, Mylonakou MN, Ferroni S, Ottersen OP, Amiry-Moghaddam M (2011) An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex is essential for cell-volume control in astrocytes *Proceedings of the National Academy of Sciences of the United States of America* 108:2563-2568 doi:10.1073/pnas.1012867108
- Berland S et al. (2018) A de novo Ser111Thr variant in aquaporin-4 in a patient with intellectual disability, transient signs of brain ischemia, transient cardiac hypertrophy, and progressive gait disturbance *Cold Spring Harb Mol Case Stud* 4 doi:10.1101/mcs.a002303
- Binder DK, Oshio K, Ma T, Verkman AS, Manley GT (2004) Increased seizure threshold in mice lacking aquaporin-4 water channels *Neuroreport* 15:259-262 doi:00001756-200402090-00009 [pii]
- Blake DJ, Weir A, Newey SE, Davies KE (2002) Function and genetics of dystrophin and dystrophin-related proteins in muscle *Physiological reviews* 82:291-329 doi:10.1152/physrev.00028.2001
- Bloch O, Auguste KI, Manley GT, Verkman AS (2006) Accelerated progression of kaolin-induced hydrocephalus in aquaporin-4-deficient mice *J Cereb Blood Flow Metab* 26:1527-1537 doi:10.1038/sj.jcbfm.9600306
- Bolkvadze T, Pitkanen A (2012) Development of post-traumatic epilepsy after controlled cortical impact and lateral fluid-percussion-induced brain injury in the mouse *J Neurotrauma* 29:789-812 doi:10.1089/neu.2011.1954
- Boulay AC et al. (2017) Translation in astrocyte distal processes sets molecular heterogeneity at the gliovascular interface *Cell Discov* 3:17005 doi:10.1038/celldisc.2017.5
- Bowman W (1840) On the minute structure and movements of voluntary muscle. *Philos Trans R Soc Lon Biol Sci* 130:457-494
- Butler TL et al. (2006) Cardiac aquaporin expression in humans, rats, and mice *Am J Physiol Heart Circ Physiol* 291:H705-713 doi:10.1152/ajpheart.00090.2006
- Camassa LM et al. (2015) Mechanisms underlying AQP4 accumulation in astrocyte endfeet *Glia* doi:10.1002/glia.22878
- Campbell KP, Stull JT (2003) Skeletal muscle basement membrane-sarcolemma-cytoskeleton interaction minireview series *The Journal of biological chemistry* 278:12599-12600 doi:10.1074/jbc.R300005200

- Cohn RD et al. (2002) Disruption of DAG1 in differentiated skeletal muscle reveals a role for dystroglycan in muscle regeneration *Cell* 110:639-648
- Colognato H, Winkelman DA, Yurchenco PD (1999) Laminin polymerization induces a receptor-cytoskeleton network *The Journal of cell biology* 145:619-631
- Cosgrove J, Alli S, Ramadan H, Ford HL (2014) Myocarditis and diffuse skeletal muscle oedema: new features of neuromyelitis optica spectrum disorder? A case report *Mult Scler* 20:120-122 doi:10.1177/1352458513495939
- Crane JM, Tajima M, Verkman AS (2009) Live-cell imaging of aquaporin-4 diffusion and interactions in orthogonal arrays of particles *Neuroscience* doi:S0306-4522(09)01392-X [pii];10.1016/j.neuroscience.2009.08.034 [doi]
- Crusio WE (2004) Flanking gene and genetic background problems in genetically manipulated mice *Biol Psychiatry* 56:381-385 doi:10.1016/j.biopsych.2003.12.026
- Culligan K, Glover L, Dowling P, Ohlendieck K (2001) Brain dystrophin-glycoprotein complex: persistent expression of beta-dystroglycan, impaired oligomerization of Dp71 and up-regulation of utrophins in animal models of muscular dystrophy *BMC Cell Biol* 2:2
- Dalkara T, Gursay-Ozdemir Y, Yemisci M (2011) Brain microvascular pericytes in health and disease *Acta Neuropathol* 122:1-9 doi:10.1007/s00401-011-0847-6
- Danbolt NC, Zhou Y, Furness DN, Holmseth S (2016) Strategies for immunohistochemical protein localization using antibodies: What did we learn from neurotransmitter transporters in glial cells and neurons *Glia* 64:2045-2064 doi:10.1002/glia.23027
- Daneman R, Zhou L, Kebede AA, Barres BA (2010) Pericytes are required for blood-brain barrier integrity during embryogenesis *Nature* 468:562-566 doi:10.1038/nature09513
- Dermietzel R (1973) Visualization by freeze-fracturing of regular structures in glial cell membranes *Naturwissenschaften* 60:208
- Devic E (1894) Myélite subaigue compliquée de neurite optique. *Bull Med* 1894;8:1033-1034 *Bull Med* 8:1033-1034
- Duan T, Smith AJ, Verkman AS (2018) Complement-dependent bystander injury to neurons in AQP4-IgG seropositive neuromyelitis optica *J Neuroinflammation* 15:294 doi:10.1186/s12974-018-1333-z
- Eberth CJ (1871) *Handbuch der Lehre von den Geweben des Menschen und der Tiere*, ed. S. Stricker, Bd. 1, Leipzig
- Eid T et al. (2005) Loss of perivascular aquaporin 4 may underlie deficient water and K⁺ homeostasis in the human epileptogenic hippocampus *Proceedings of the National Academy of Sciences of the United States of America* 102:1193-1198 doi:10.1073/pnas.0409308102
- Eide PK, Hansson HA (2017) Astrogliosis and impaired aquaporin-4 and dystrophin systems in idiopathic normal pressure hydrocephalus *Neuropathol Appl Neurobiol* doi:10.1111/nan.12420
- Eide PK, Sorteberg W (2010) Diagnostic intracranial pressure monitoring and surgical management in idiopathic normal pressure hydrocephalus: a 6-year review of 214 patients *Neurosurgery* 66:80-91 doi:10.1227/01.NEU.0000363408.69856.B8
- Eidsvaag VA, Enger R, Hansson HA, Eide PK, Nagelhus EA (2017) Human and mouse cortical astrocytes differ in aquaporin-4 polarization toward microvessels *Glia* 65:964-973 doi:10.1002/glia.23138
- Eilert-Olsen M et al. (2012) Deletion of aquaporin-4 changes the perivascular glial protein scaffold without disrupting the brain endothelial barrier *Glia* 60:432-440 doi:10.1002/glia.22277
- El-Khoury N, Braun A, Hu F, Pandey M, Nedergaard M, Lagamma EF, Ballabh P (2006) Astrocyte end-feet in germinal matrix, cerebral cortex, and white matter in developing infants *Pediatr Res* 59:673-679 doi:10.1203/01.pdr.0000214975.85311.9c
- Elkjaer M et al. (2000) Immunolocalization of AQP9 in liver, epididymis, testis, spleen, and brain *BiochemBiophysResCommun* 276:1118-1128 doi:10.1006/bbrc.2000.3505 [doi];S0006291X00935058 [pii]
- Eng LF, Vanderhaeghen JJ, Bignami A, Gerstl B (1971) An acidic protein isolated from fibrous astrocytes *Brain research* 28:351-354
- Eusebio A, Oliveri F, Barzaghi P, Ruegg MA (2003) Expression of mouse agrin in normal, denervated and dystrophic muscle *Neuromuscul Disord* 13:408-415
- Fallier-Becker P, Sperveslage J, Wolburg H, Noell S (2011) The impact of agrin on the formation of orthogonal arrays of particles in cultured astrocytes from wild-type and agrin-null mice *Brain research* 1367:2-12 doi:10.1016/j.brainres.2010.09.092
- Fallier-Becker P, Vollmer JP, Bauer HC, Noell S, Wolburg H, Mack AF (2014) Onset of aquaporin-4 expression in the developing mouse brain *International journal of developmental neuroscience : the official journal of the International Society for Developmental Neuroscience* 36:81-89 doi:10.1016/j.ijdevneu.2014.06.001
- Fanning AS, Anderson JM (1999) PDZ domains: fundamental building blocks in the organization of protein complexes at the plasma membrane *J Clin Invest* 103:767-772 doi:10.1172/JCI16509
- Fariello RG (1995) Critical review of the animal models of generalized epilepsies *Ital J Neurol Sci* 16:69-72
- Feng X et al. (2009) Sporadic obstructive hydrocephalus in Aqp4 null mice *JNeurosciRes* 87:1150-1155 doi:10.1002/jnr.21927 [doi]

- Finn RN, Chauvigne F, Hlidberg JB, Cutler CP, Cerda J (2014) The lineage-specific evolution of aquaporin gene clusters facilitated tetrapod terrestrial adaptation PLoS One 9:e113686 doi:10.1371/journal.pone.0113686
- Franciosi S et al. (2007) Pepsin pretreatment allows collagen IV immunostaining of blood vessels in adult mouse brain Journal of neuroscience methods 163:76-82 doi:10.1016/j.jneumeth.2007.02.020
- Frank RN, Dutta S, Mancini MA (1987) Pericyte coverage is greater in the retinal than in the cerebral capillaries of the rat Invest Ophthalmol Vis Sci 28:1086-1091
- Frigeri A, Nicchia GP, Verbavatz JM, Valenti G, Svelto M (1998) Expression of aquaporin-4 in fast-twitch fibers of mammalian skeletal muscle J Clin Invest 102:695-703 doi:10.1172/JCI2545
- Frydenlund DS et al. (2006) Temporary loss of perivascular aquaporin-4 in neocortex after transient middle cerebral artery occlusion in mice Proceedings of the National Academy of Sciences of the United States of America 103:13532-13536 doi:10.1073/pnas.0605796103
- Fujisawa N, Maeda Y, Yamamoto Y, Sato NL, Niimura S (2003) Newly established low seizure susceptible and seizure-prone inbred strains of Mongolian gerbil Exp Anim 52:169-172
- Furman CS, Gorelick-Feldman DA, Davidson KG, Yasumura T, Neely JD, Agre P, Rash JE (2003) Aquaporin-4 square array assembly: opposing actions of M1 and M23 isoforms ProcNatlAcadSciUSA 100:13609-13614 doi:10.1073/pnas.2235843100 [doi];2235843100 [pii]
- Galea I, Bechmann I, Perry VH (2007) What is immune privilege (not)? Trends Immunol 28:12-18 doi:S1471-4906(06)00326-7 [pii];10.1016/j.it.2006.11.004 [doi]
- Gautam M, Noakes PG, Moscoso L, Rupp F, Scheller RH, Merlie JP, Sanes JR (1996) Defective neuromuscular synaptogenesis in agrin-deficient mutant mice Cell 85:525-535
- Gee SH, Madhavan R, Levinson SR, Caldwell JH, Sealock R, Froehner SC (1998) Interaction of muscle and brain sodium channels with multiple members of the syntrophin family of dystrophin-associated proteins The Journal of neuroscience : the official journal of the Society for Neuroscience 18:128-137
- Glees P (1955) Neuroglia. Morphology and function. Oxford: Blackwell,
- Goldmann E (1913) Vitalfärbungen am Zentralnervensystem. Beitrag zur Physio-Pathologie des Plexus Choroideus und der Hirnhäute. Akademie der Wissenschaften Physikalisch-Mathematische Classe 1:1-64
- Gorelick DA, Praetorius J, Tsunenari T, Nielsen S, Agre P (2006) Aquaporin-11: a channel protein lacking apparent transport function expressed in brain BMCBiochem 7:14 doi:1471-2091-7-14 [pii];10.1186/1471-2091-7-14 [doi]
- Guadagno E, Moukhles H (2004) Laminin-induced aggregation of the inwardly rectifying potassium channel, Kir4.1, and the water-permeable channel, AQP4, via a dystroglycan-containing complex in astrocytes Glia 47:138-149 doi:10.1002/glia.20039
- Gundersen GA, Vindedal GF, Skare O, Nagelhus EA (2013) Evidence that pericytes regulate aquaporin-4 polarization in mouse cortical astrocytes Brain StructFunct doi:10.1007/s00429-013-0629-0 [doi]
- Gunnarson E, Axehult G, Baturina G, Zelenin S, Zelenina M, Aperia A (2005) Lead induces increased water permeability in astrocytes expressing aquaporin 4 Neuroscience 136:105-114 doi:S0306-4522(05)00768-2 [pii];10.1016/j.neuroscience.2005.07.027 [doi]
- Hall CN et al. (2014) Capillary pericytes regulate cerebral blood flow in health and disease Nature 508:55-60 doi:10.1038/nature13165
- Hamilton NB, Attwell D, Hall CN (2010) Pericyte-mediated regulation of capillary diameter: a component of neurovascular coupling in health and disease Front Neuroenergetics 2 doi:10.3389/fnene.2010.00005
- Han X et al. (2013) Forebrain engraftment by human glial progenitor cells enhances synaptic plasticity and learning in adult mice Cell Stem Cell 12:342-353 doi:10.1016/j.stem.2012.12.015
- Hasan-Olive MM, Enger R, Hansson HA, Nagelhus EA, Eide PK (2018) Loss of perivascular aquaporin-4 in idiopathic normal pressure hydrocephalus Glia doi:10.1002/glia.23528
- Heuser K et al. (2010a) Temporal lobe epilepsy and matrix metalloproteinase 9: a tempting relation but negative genetic association Seizure 19:335-338 doi:10.1016/j.seizure.2010.05.003
- Heuser K et al. (2010b) Variants of the genes encoding AQP4 and Kir4.1 are associated with subgroups of patients with temporal lobe epilepsy Epilepsy Res 88:55-64 doi:S0920-1211(09)00286-1 [pii];10.1016/j.epilepsyres.2009.09.023 [doi]
- Ho JD et al. (2009) Crystal structure of human aquaporin 4 at 1.8 Å and its mechanism of conductance ProcNatlAcadSciUSA 106:7437-7442 doi:0902725106 [pii];10.1073/pnas.0902725106 [doi]
- Hoddevik EH, Khan FH, Rahmani S, Ottersen OP, Boldt HB, Amiry-Moghaddam M (2017) Factors determining the density of AQP4 water channel molecules at the brain-blood interface Brain structure & function 222:1753-1766 doi:10.1007/s00429-016-1305-y
- Hohenester E, Yurchenco PD (2013) Laminins in basement membrane assembly Cell Adh Migr 7:56-63 doi:10.4161/cam.21831
- Holen T (2011) The ultrastructure of lamellar stack astrocytes Glia 59:1075-1083 doi:10.1002/glia.21180
- Holter KE et al. (2017) Interstitial solute transport in 3D reconstructed neuropil occurs by diffusion rather than bulk flow Proceedings of the National Academy of Sciences of the United States of America 114:9894-9899 doi:10.1073/pnas.1706942114

- Hornig S et al. (2017) Astrocytic tight junctions control inflammatory CNS lesion pathogenesis *J Clin Invest* 127:3136-3151 doi:10.1172/JCI91301
- Iliff JJ et al. (2012) A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid beta *Science translational medicine* 4:147ra111 doi:10.1126/scitranslmed.3003748
- Iliff JJ et al. (2013) Cerebral arterial pulsation drives paravascular CSF-interstitial fluid exchange in the murine brain *The Journal of neuroscience : the official journal of the Society for Neuroscience* 33:18190-18199 doi:10.1523/JNEUROSCI.1592-13.2013
- Inoue M, Wakayama Y, Liu JW, Murahashi M, Shibuya S, Oniki H (2002) Ultrastructural localization of aquaporin 4 and alpha1-syntrophin in the vascular feet of brain astrocytes *Tohoku J Exp Med* 197:87-93
- Iorio R, Fryer JP, Hinson SR, Fallier-Becker P, Wolburg H, Pittcock SJ, Lennon VA (2013) Astrocytic autoantibody of neuromyelitis optica (NMO-IgG) binds to aquaporin-4 extracellular loops, monomers, tetramers and high order arrays *J Autoimmun* 40:21-27 doi:10.1016/j.jaut.2012.07.008
- Ip YK et al. (2013) Molecular characterization of branchial aquaporin 1aa and effects of seawater acclimation, emersion or ammonia exposure on its mRNA expression in the gills, gut, kidney and skin of the freshwater climbing perch, *Anabas testudineus* *PLoS One* 8:e61163 doi:10.1371/journal.pone.0061163
- Ishibashi K (2009) New members of mammalian aquaporins: AQP10-AQP12 *Handb Exp Pharmacol*:251-262 doi:10.1007/978-3-540-79885-9_13
- Ishibashi K, Hara S, Kondo S (2009) Aquaporin water channels in mammals *ClinExpNephrol* 13:107-117 doi:10.1007/s10157-008-0118-6 [doi]
- Ishido M, Nakamura T (2018) The expression of aquaporin-4 is regulated based on innervation in skeletal muscles *J Muscle Res Cell Motil* 39:17-23 doi:10.1007/s10974-018-9494-z
- Jahn TP et al. (2004) Aquaporin homologues in plants and mammals transport ammonia *FEBS letters* 574:31-36 doi:10.1016/j.febslet.2004.08.004
- Jarius S, Wildemann B (2017) Devic's disease before Devic: On the contribution of Friedrich Albin Schanz (1863-1923) *J Neurol Sci* 379:99-102 doi:10.1016/j.jns.2017.05.044
- Kaakinen M, Salmela P, Zelenin S, Metsikko K (2007) Distribution of aquaporin 4 on sarcolemma of fast-twitch skeletal myofibres *Cell and tissue research* 329:529-539 doi:10.1007/s00441-007-0442-2
- Katoozi S, Skauli N, Rahmani S, Camassa LMA, Boldt HB, Ottersen OP, Amiry-Moghaddam M (2017) Targeted deletion of Aqp4 promotes the formation of astrocytic gap junctions *Brain structure & function* 222:3959-3972 doi:10.1007/s00429-017-1448-5
- Kessler RA et al. (2017) Anti-aquaporin-4 titer is not predictive of disease course in neuromyelitis optica spectrum disorder: A multicenter cohort study *Mult Scler Relat Disord* 17:198-201 doi:10.1016/j.msard.2017.08.005
- Kharatishvili I, Nissinen JP, McIntosh TK, Pitkanen A (2006) A model of posttraumatic epilepsy induced by lateral fluid-percussion brain injury in rats *Neuroscience* 140:685-697 doi:10.1016/j.neuroscience.2006.03.012
- Kimelberg HK (2004) The problem of astrocyte identity *Neurochem Int* 45:191-202 doi:10.1016/j.neuint.2003.08.015
- Kimelberg HK (2010) Functions of mature mammalian astrocytes: a current view *The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry* 16:79-106 doi:10.1177/1073858409342593
- Klein B, Kuschinsky W, Schrock H, Vetterlein F (1986) Interdependency of local capillary density, blood flow, and metabolism in rat brains *Am J Physiol* 251:H1333-1340 doi:10.1152/ajpheart.1986.251.6.H1333
- Kornau HC, Schenker LT, Kennedy MB, Seeburg PH (1995) Domain interaction between NMDA receptor subunits and the postsynaptic density protein PSD-95 *Science* 269:1737-1740
- Korogod N, Petersen CC, Knott GW (2015) Ultrastructural analysis of adult mouse neocortex comparing aldehyde perfusion with cryo fixation *Elife* 4 doi:10.7554/eLife.05793
- Kress BT et al. (2014) Impairment of paravascular clearance pathways in the aging brain *Ann Neurol* 76:845-861 doi:10.1002/ana.24271
- Kupferberg H (2001) Animal models used in the screening of antiepileptic drugs *Epilepsia* 42 Suppl 4:7-12
- Kuschinsky W (1990) Coupling of blood flow and metabolism in the brain *J Basic Clin Physiol Pharmacol* 1:191-201
- Laloux T, Junqueira B, Maistriaux LC, Ahmed J, Jurkiewicz A, Chaumont F (2018) Plant and Mammal Aquaporins: Same but Different *Int J Mol Sci* 19 doi:10.3390/ijms19020521
- Landis DM, Reese TS (1974) Arrays of particles in freeze-fractured astrocytic membranes *Journal of Cell Biology* 60:316-320
- Landis DM, Reese TS (1981a) Astrocyte membrane structure: changes after circulatory arrest *The Journal of cell biology* 88:660-663
- Landis DM, Reese TS (1981b) Membrane structure in mammalian astrocytes: a review of freeze-fracture studies on adult, developing, reactive and cultured astrocytes *J Exp Biol* 95:35-48

- Landis DM, Reese TS (1982) Regional organization of astrocytic membranes in cerebellar cortex *Neuroscience* 7:937-950
- Lee TS, Eid T, Mane S, Kim JH, Spencer DD, Ottersen OP, de Lanerolle NC (2004) Aquaporin-4 is increased in the sclerotic hippocampus in human temporal lobe epilepsy *Acta Neuropathol* 108:493-502 doi:10.1007/s00401-004-0910-7 [doi]
- Lehre KP, Levy LM, Ottersen OP, Storm-Mathisen J, Danbolt NC (1995) Differential expression of two glial glutamate transporters in the rat brain: quantitative and immunocytochemical observations *The Journal of neuroscience : the official journal of the Society for Neuroscience* 15:1835-1853
- Lenhossek M (1893) *Der feinere Bau des Nervensystems im Lichte neuester Forschung*. Fischer's Medicinische Buchhandlung; Berlin
- Lennon VA, Kryzer TJ, Pittock SJ, Verkman AS, Hinson SR (2005) IgG marker of optic-spinal multiple sclerosis binds to the aquaporin-4 water channel *JExpMed* 202:473-477 doi:jem.20050304 [pii];10.1084/jem.20050304 [doi]
- Lewandowski M (1900) Zur Lehre von der Cerebrospinalflüssigkeit. *Z Klin Forsch* 40:480-494
- Li S, Harrison D, Carbonetto S, Fassler R, Smyth N, Edgar D, Yurchenco PD (2002) Matrix assembly, regulation, and survival functions of laminin and its receptors in embryonic stem cell differentiation *The Journal of cell biology* 157:1279-1290 doi:10.1083/jcb.200203073
- Li S, Liquari P, McKee KK, Harrison D, Patel R, Lee S, Yurchenco PD (2005) Laminin-sulfatide binding initiates basement membrane assembly and enables receptor signaling in Schwann cells and fibroblasts *The Journal of cell biology* 169:179-189 doi:10.1083/jcb.200501098
- Lin W, Burgess RW, Dominguez B, Pfaff SL, Sanes JR, Lee KF (2001) Distinct roles of nerve and muscle in postsynaptic differentiation of the neuromuscular synapse *Nature* 410:1057-1064 doi:10.1038/35074025
- Long DM, Bodenheimer TS, Hartmann JF, Klatzo I (1968) Ultrastructural features of the shark brain *Am J Anat* 122:209-236 doi:10.1002/aja.1001220204
- Lunde LK, Camassa LM, Hoddevik EH, Khan FH, Ottersen OP, Boldt HB, Amiry-Moghaddam M (2015) Postnatal development of the molecular complex underlying astrocyte polarization *Brain structure & function* 220:2087-2101 doi:10.1007/s00429-014-0775-z
- Ma T, Yang B, Gillespie A, Carlson EJ, Epstein CJ, Verkman AS (1997) Generation and phenotype of a transgenic knockout mouse lacking the mercurial-insensitive water channel aquaporin-4 *J Clin Invest* 100:957-962 doi:10.1172/JCI231
- MacAulay N, Hamann S, Zeuthen T (2004) Water transport in the brain: role of cotransporters *Neuroscience* 129:1031-1044 doi:10.1016/j.neuroscience.2004.06.045
- MacIntyre N (2001) Unmasking antigens for immunohistochemistry *Br J Biomed Sci* 58:190-196
- Madrid R, Le Maout S, Barrault MB, Janvier K, Benichou S, Merot J (2001) Polarized trafficking and surface expression of the AQP4 water channel are coordinated by serial and regulated interactions with different clathrin-adaptor complexes *EMBO J* 20:7008-7021 doi:10.1093/emboj/20.24.7008
- Malik R et al. (2014) Transient hyperckemia in the setting of neuromyelitis optica (NMO) *Muscle Nerve* 50:859-862 doi:10.1002/mus.24298
- Manley GT, Binder DK, Papadopoulos MC, Verkman AS (2004) New insights into water transport and edema in the central nervous system from phenotype analysis of aquaporin-4 null mice *Neuroscience* 129:983-991 doi:10.1016/j.neuroscience.2004.06.088
- Manley GT et al. (2000) Aquaporin-4 deletion in mice reduces brain edema after acute water intoxication and ischemic stroke *NatMed* 6:159-163 doi:10.1038/72256 [doi]
- Mathiisen TM, Lehre KP, Danbolt NC, Ottersen OP (2010) The perivascular astroglial sheath provides a complete covering of the brain microvessels: an electron microscopic 3D reconstruction *Glia* 58:1094-1103 doi:10.1002/glia.20990
- Mayer U (2003) Integrins: redundant or important players in skeletal muscle? *The Journal of biological chemistry* 278:14587-14590 doi:10.1074/jbc.R200022200
- McKee KK, Capizzi S, Yurchenco PD (2009) Scaffold-forming and Adhesive Contributions of Synthetic Laminin-binding Proteins to Basement Membrane Assembly *The Journal of biological chemistry* 284:8984-8994 doi:10.1074/jbc.M809719200
- McKee KK, Harrison D, Capizzi S, Yurchenco PD (2007) Role of laminin terminal globular domains in basement membrane assembly *The Journal of biological chemistry* 282:21437-21447 doi:10.1074/jbc.M702963200
- Mishima T, Hirase H (2010) In vivo intracellular recording suggests that gray matter astrocytes in mature cerebral cortex and hippocampus are electrophysiologically homogeneous *The Journal of neuroscience : the official journal of the Society for Neuroscience* 30:3093-3100 doi:10.1523/JNEUROSCI.5065-09.2010
- Mishra A, Reynolds JP, Chen Y, Gourine AV, Rusakov DA, Attwell D (2016) Astrocytes mediate neurovascular signaling to capillary pericytes but not to arterioles *Nature neuroscience* 19:1619-1627 doi:10.1038/nn.4428
- Moe SE, Sorbo JG, Sogaard R, Zeuthen T, Petter OO, Holen T (2008) New isoforms of rat Aquaporin-4 *Genomics* 91:367-377 doi:S0888-7543(07)00293-5 [pii];10.1016/j.ygeno.2007.12.003 [doi]
- Moore SA et al. (2002) Deletion of brain dystroglycan recapitulates aspects of congenital muscular dystrophy *Nature* 418:422-425 doi:10.1038/nature00838 [doi];nature00838 [pii]

- Mylonakou MN et al. (2009) Analysis of mice with targeted deletion of AQP9 gene provides conclusive evidence for expression of AQP9 in neurons *JNeurosciRes* 87:1310-1322 doi:10.1002/jnr.21952 [doi]
- Nagelhus EA et al. (1999) Immunogold evidence suggests that coupling of K⁺ siphoning and water transport in rat retinal Muller cells is mediated by a coenrichment of Kir4.1 and AQP4 in specific membrane domains *Glia* 26:47-54
- Nagelhus EA, Ottersen OP (2013) Physiological roles of aquaporin-4 in brain *Physiological reviews* 93:1543-1562 doi:10.1152/physrev.00011.2013
- Nagy JI, Rash JE (2000) Connexins and gap junctions of astrocytes and oligodendrocytes in the CNS *Brain Res Brain Res Rev* 32:29-44
- Nakada T, Kwee IL, Igarashi H, Suzuki Y (2017) Aquaporin-4 Functionality and Virchow-Robin Space Water Dynamics: Physiological Model for Neurovascular Coupling and Glymphatic Flow *Int J Mol Sci* 18 doi:10.3390/ijms18081798
- Nakhoul NL, Hering-Smith KS, Abdulnour-Nakhoul SM, Hamm LL (2001) Transport of NH₃/NH in oocytes expressing aquaporin-1 *Am J Physiol Renal Physiol* 281:F255-263 doi:10.1152/ajprenal.2001.281.2.F255
- Nedergaard M, Ransom B, Goldman SA (2003) New roles for astrocytes: redefining the functional architecture of the brain *Trends in neurosciences* 26:523-530 doi:10.1016/j.tins.2003.08.008
- Neely JD, Amiry-Moghaddam M, Ottersen OP, Froehner SC, Agre P, Adams ME (2001) Syntrophin-dependent expression and localization of Aquaporin-4 water channel protein *Proceedings of the National Academy of Sciences of the United States of America* 98:14108-14113 doi:10.1073/pnas.241508198
- Newman EA (2003) New roles for astrocytes: regulation of synaptic transmission *Trends in neurosciences* 26:536-542 doi:10.1016/S0166-2236(03)00237-6
- Nicchia GP et al. (2011) D184E mutation in aquaporin-4 gene impairs water permeability and links to deafness *Neuroscience* 197:80-88 doi:10.1016/j.neuroscience.2011.09.023
- Nicchia GP et al. (2009) Aquaporin-4 orthogonal arrays of particles are the target for neuromyelitis optica autoantibodies *Glia* 57:1363-1373 doi:10.1002/glia.20855
- Nielsen S, Nagelhus EA, Amiry-Moghaddam M, Bourque C, Agre P, Ottersen OP (1997) Specialized membrane domains for water transport in glial cells: high-resolution immunogold cytochemistry of aquaporin-4 in rat brain *The Journal of neuroscience : the official journal of the Society for Neuroscience* 17:171-180
- Nippert AR, Biesecker KR, Newman EA (2018) Mechanisms Mediating Functional Hyperemia in the Brain *The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry* 24:73-83 doi:10.1177/1073858417703033
- Noel G, Tham DK, Moukhles H (2009) Interdependence of laminin-mediated clustering of lipid rafts and the dystrophin complex in astrocytes *The Journal of biological chemistry* 284:19694-19704 doi:10.1074/jbc.M109.010090
- Noell S, Fallier-Becker P, Beyer C, Kroger S, Mack AF, Wolburg H (2007) Effects of agrin on the expression and distribution of the water channel protein aquaporin-4 and volume regulation in cultured astrocytes *The European journal of neuroscience* 26:2109-2118 doi:10.1111/j.1460-9568.2007.05850.x
- Noell S, Fallier-Becker P, Deutsch U, Mack AF, Wolburg H (2009) Agrin defines polarized distribution of orthogonal arrays of particles in astrocytes *Cell and tissue research* 337:185-195 doi:10.1007/s00441-009-0812-z
- Noell S et al. (2011) Evidence for a role of dystroglycan regulating the membrane architecture of astroglial endfeet *The European journal of neuroscience* 33:2179-2186 doi:10.1111/j.1460-9568.2011.07688.x
- Noell S et al. (2012) Dynamics of expression patterns of AQP4, dystroglycan, agrin and matrix metalloproteinases in human glioblastoma *Cell and tissue research* 347:429-441 doi:10.1007/s00441-011-1321-4
- Norris ML (1987) *THE UFAW handbook on the care and management of laboratory animals.* (Poole T.B. edit.). 6th edn. Longman Scientific & Technical, Essex, England,
- Nowak MA, Boerlijst MC, Cooke J, Smith JM (1997) Evolution of genetic redundancy *Nature* 388:167-171 doi:10.1038/40618
- Oberheim NA, Goldman SA, Nedergaard M (2012) Heterogeneity of astrocytic form and function *Methods Mol Biol* 814:23-45 doi:10.1007/978-1-61779-452-0_3
- Omid Y, Campbell L, Barar J, Connell D, Akhtar S, Gumbleton M (2003) Evaluation of the immortalised mouse brain capillary endothelial cell line, b.End3, as an in vitro blood-brain barrier model for drug uptake and transport studies *Brain research* 990:95-112
- Papadopoulos MC, Manley GT, Krishna S, Verkman AS (2004) Aquaporin-4 facilitates reabsorption of excess fluid in vasogenic brain edema *FASEB J* 18:1291-1293 doi:10.1096/fj.04-1723fje [doi];04-1723fje [pii]
- Papadopoulos MC, Verkman AS (2005) Aquaporin-4 gene disruption in mice reduces brain swelling and mortality in pneumococcal meningitis *JBiolChem* 280:13906-13912 doi:M413627200 [pii];10.1074/jbc.M413627200 [doi]

- Parsons MJ, Campos I, Hirst EM, Stemple DL (2002) Removal of dystroglycan causes severe muscular dystrophy in zebrafish embryos *Development* 129:3505-3512
- Peppiatt CM, Howarth C, Mobbs P, Attwell D (2006) Bidirectional control of CNS capillary diameter by pericytes *Nature* 443:700-704 doi:10.1038/nature05193
- Peters MF et al. (1998) Differential membrane localization and intermolecular associations of alpha-dystrobrevin isoforms in skeletal muscle *The Journal of cell biology* 142:1269-1278
- Pisani F, Rossi A, Nicchia GP, Svelto M, Frigeri A (2011) Translational regulation mechanisms of aquaporin-4 supramolecular organization in astrocytes *Glia* 59:1923-1932 doi:10.1002/glia.21234
- Plog BA et al. (2015) Biomarkers of traumatic injury are transported from brain to blood via the glymphatic system *The Journal of neuroscience : the official journal of the Society for Neuroscience* 35:518-526 doi:10.1523/JNEUROSCI.3742-14.2015
- Potokar M, Stenovec M, Jorgacevski J, Holen T, Kreft M, Ottersen OP, Zorec R (2013) Regulation of AQP4 surface expression via vesicle mobility in astrocytes *Glia* 61:917-928 doi:10.1002/glia.22485
- Promeneur D, Lunde LK, Amiry-Moghaddam M, Agre P (2013) Protective role of brain water channel AQP4 in murine cerebral malaria *Proceedings of the National Academy of Sciences of the United States of America* 110:1035-1040 doi:10.1073/pnas.1220566110
- Ramon y Cajal S (1911) *Histologie du Systeme Nerveux de l'Homme et des Vertebres* vol Volumes I and II.
- Rash JE, Staehelin LA, Ellisman MH (1974) Rectangular arrays of particles on freeze-cleaved plasma membranes are not gap junctions *Experimental cell research* 86:187-190
- Rash JE, Yasumura T, Hudson CS, Agre P, Nielsen S (1998) Direct immunogold labeling of aquaporin-4 in square arrays of astrocyte and ependymocyte plasma membranes in rat brain and spinal cord *Proceedings of the National Academy of Sciences of the United States of America* 95:11981-11986
- Ratelade J, Bennett JL, Verkman AS (2011) Intravenous neuromyelitis optica autoantibody in mice targets aquaporin-4 in peripheral organs and area postrema *PLoS One* 6:e27412 doi:10.1371/journal.pone.0027412
- Reese TS, Karnovsky MJ (1967) Fine structural localization of a blood-brain barrier to exogenous peroxidase *JCell Biol* 34:207-217
- Reverberi R, Reverberi L (2007) Factors affecting the antigen-antibody reaction *Blood Transfus* 5:227-240 doi:10.2450/2007.0047-07
- Ringstad G, Vatnehol SAS, Eide PK (2017) Glymphatic MRI in idiopathic normal pressure hydrocephalus *Brain* 140:2691-2705 doi:10.1093/brain/awx191
- Robel S et al. (2009) Conditional deletion of beta1-integrin in astroglia causes partial reactive gliosis *Glia* 57:1630-1647 doi:10.1002/glia.20876
- Rolfe DF, Brown GC (1997) Cellular energy utilization and molecular origin of standard metabolic rate in mammals *Physiological reviews* 77:731-758 doi:10.1152/physrev.1997.77.3.731
- Rosito S et al. (2018) Supramolecular aggregation of aquaporin-4 is different in muscle and brain: correlation with tissue susceptibility in neuromyelitis optica *J Cell Mol Med* 22:1236-1246 doi:10.1111/jcmm.13401
- Rossi A, Moritz TJ, Ratelade J, Verkman AS (2012) Super-resolution imaging of aquaporin-4 orthogonal arrays of particles in cell membranes *Journal of cell science* 125:4405-4412 doi:10.1242/jcs.109603
- Rossi A, Pisani F, Nicchia GP, Svelto M, Frigeri A (2010) Evidences for a leaky scanning mechanism for the synthesis of the shorter M23 protein isoform of aquaporin-4: implication in orthogonal array formation and neuromyelitis optica antibody interaction *The Journal of biological chemistry* 285:4562-4569 doi:10.1074/jbc.M109.069245
- Rouget C (1873) Memoire sur le développement, la structure et les propriétés physiologiques des capillaires sanguins et lymphatiques *Arch Physiol Norm Path*:603-663
- Rungta RL, Chappak S (2016) Astrocyte endfeet march to the beat of different vessels *Nature neuroscience* 19:1539-1541 doi:10.1038/nn.4446
- Rutkovskiy A et al. (2012) Aquaporin-4 in the heart: expression, regulation and functional role in ischemia *Basic Res Cardiol* 107:280 doi:10.1007/s00395-012-0280-6
- Saadoun S, Papadopoulos MC, Watanabe H, Yan D, Manley GT, Verkman AS (2005) Involvement of aquaporin-4 in astroglial cell migration and glial scar formation *JCell Sci* 118:5691-5698 doi:jcs.02680 [pii];10.1242/jcs.02680 [doi]
- Saadoun S, Tait MJ, Reza A, Davies DC, Bell BA, Verkman AS, Papadopoulos MC (2009) AQP4 gene deletion in mice does not alter blood-brain barrier integrity or brain morphology *Neuroscience* 161:764-772 doi:S0306-4522(09)00462-X [pii];10.1016/j.neuroscience.2009.03.069 [doi]
- Sakers K, Lake AM, Khazanchi R, Ouwenga R, Vasek MJ, Dani A, Dougherty JD (2017) Astrocytes locally translate transcripts in their peripheral processes *Proceedings of the National Academy of Sciences of the United States of America* 114:E3830-E3838 doi:10.1073/pnas.1617782114
- Salman MM et al. (2017) Transcriptome analysis suggests a role for the differential expression of cerebral aquaporins and the MAPK signalling pathway in human temporal lobe epilepsy *The European journal of neuroscience* 46:2121-2132 doi:10.1111/ejn.13652

- Sanes JR (2003) The basement membrane/basal lamina of skeletal muscle *The Journal of biological chemistry* 278:12601-12604 doi:10.1074/jbc.R200027200
- Schafer GL, Crabbe JC (1996) Sensitivity to ethanol-induced ataxia in HOT and COLD selected lines of mice *Alcohol Clin Exp Res* 20:1604-1612
- Seifert G, Schilling K, Steinhäuser C (2006) Astrocyte dysfunction in neurological disorders: a molecular perspective *Nature reviews Neuroscience* 7:194-206 doi:10.1038/nrn1870
- Shepro D, Morel NM (1993) Pericyte physiology *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* 7:1031-1038
- Shi SR, Cote RJ, Taylor CR (1997) Antigen retrieval immunohistochemistry: past, present, and future *J Histochem Cytochem* 45:327-343 doi:10.1177/002215549704500301
- Shibuya S, Wakayama Y, Inoue M, Kojima H, Oniki H (2006) The relationship between intramembranous particles and aquaporin molecules in the plasma membranes of normal rat skeletal muscles: a fracture-label study *J Electron Microscop* (Tokyo) 55:63-68 doi:10.1093/jmicro/dfi017
- Simpson IA, Carruthers A, Vannucci SJ (2007) Supply and demand in cerebral energy metabolism: the role of nutrient transporters *J Cereb Blood Flow Metab* 27:1766-1791 doi:10.1038/sj.jcbfm.9600521
- Smith AJ, Yao X, Dix JA, Jin BJ, Verkman AS (2017) Test of the 'glymphatic' hypothesis demonstrates diffusive and aquaporin-4-independent solute transport in rodent brain parenchyma *Elife* 6 doi:10.7554/eLife.27679
- Smyth N, Vatansever HS, Meyer M, Frie C, Paulsson M, Edgar D (1998) The targeted deletion of the LAMC1 gene *Annals of the New York Academy of Sciences* 857:283-286
- Sorani MD, Manley GT, Giacomini KM (2008a) Genetic variation in human aquaporins and effects on phenotypes of water homeostasis *Hum Mutat* 29:1108-1117 doi:10.1002/humu.20762 [doi]
- Sorani MD, Zador Z, Hurowitz E, Yan D, Giacomini KM, Manley GT (2008b) Novel variants in human Aquaporin-4 reduce cellular water permeability *Hum Mol Genet* 17:2379-2389 doi:ddn138 [pii];10.1093/hmg/ddn138 [doi]
- Sorbo JG, Moe SE, Holen T (2007) Early upregulation in nasal epithelium and strong expression in olfactory bulb glomeruli suggest a role for Aquaporin-4 in olfaction *FEBS Lett* 581:4884-4890 doi:S0014-5793(07)00994-5 [pii];10.1016/j.febslet.2007.09.018 [doi]
- Sorbo JG, Moe SE, Ottersen OP, Holen T (2008) The molecular composition of square arrays *Biochemistry* 47:2631-2637 doi:10.1021/bi702146k [doi]
- Stahl K et al. (2018) Targeted deletion of the aquaglyceroporin AQP9 is protective in a mouse model of Parkinson's disease *PLoS One* 13:e0194896 doi:10.1371/journal.pone.0194896
- Stephan A et al. (2008) Neurotrophin cleaves agrin locally at the synapse *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* 22:1861-1873 doi:10.1096/fj.07-100008
- Terris J, Ecelbarger CA, Marples D, Knepper MA, Nielsen S (1995) Distribution of aquaporin-4 water channel expression within rat kidney *Am J Physiol* 269:F775-785 doi:10.1152/ajprenal.1995.269.6.F775
- Tham DK, Joshi B, Moukhles H (2016) Aquaporin-4 Cell-Surface Expression and Turnover Are Regulated by Dystroglycan, Dynamin, and the Extracellular Matrix in Astrocytes *PLoS One* 11:e0165439 doi:10.1371/journal.pone.0165439
- Thrane AS et al. (2011) Critical role of aquaporin-4 (AQP4) in astrocytic Ca²⁺ signaling events elicited by cerebral edema *Proceedings of the National Academy of Sciences of the United States of America* 108:846-851 doi:10.1073/pnas.1015217108
- Tilton RG, Kilo C, Williamson JR (1979) Pericyte-endothelial relationships in cardiac and skeletal muscle capillaries *Microvasc Res* 18:325-335
- Tradtrantip L, Jin BJ, Yao X, Anderson MO, Verkman AS (2017) Aquaporin-Targeted Therapeutics: State-of-the-Field *Adv Exp Med Biol* 969:239-250 doi:10.1007/978-94-024-1057-0_16
- Utriainen A et al. (2004) Structurally altered basement membranes and hydrocephalus in a type XVIII collagen deficient mouse line *Hum Mol Genet* 13:2089-2099 doi:10.1093/hmg/ddh213
- Venero JL, Vizuete ML, Ilundain AA, Machado A, Echevarria M, Cano J (1999) Detailed localization of aquaporin-4 messenger RNA in the CNS: preferential expression in periventricular organs *Neuroscience* 94:239-250
- Verbavatz JM, Ma T, Gobin R, Verkman AS (1997) Absence of orthogonal arrays in kidney, brain and muscle from transgenic knockout mice lacking water channel aquaporin-4 *Journal of cell science* 110 (Pt 22):2855-2860
- Verkhatsky A, Nedergaard M (2018) Physiology of Astroglia *Physiological reviews* 98:239-389 doi:10.1152/physrev.00042.2016
- Verkhatsky AB, A. (2013) *Glial Physiology and Pathophysiology*. Wiley-Blackwell, A John Wiley & Sons, Ltd., Publication,
- Verkman AS, Smith AJ, Phuan PW, Tradtrantip L, Anderson MO (2017) The aquaporin-4 water channel as a potential drug target in neurological disorders *Expert Opin Ther Targets* 21:1161-1170 doi:10.1080/14728222.2017.1398236
- Verkman AS, Yao X, Smith AJ (2018) The evolving mystery of why skeletal muscle is spared in seropositive neuromyelitis optica *J Cell Mol Med* 22:2039-2040 doi:10.1111/jcmm.13482

- Wakayama Y, Kumagai T, Jimi T (1986) Small size of orthogonal array in muscle plasma membrane of Fukuyama type congenital muscular dystrophy *Acta Neuropathol* 72:130-133
- Warth A, Kroger S, Wolburg H (2004) Redistribution of aquaporin-4 in human glioblastoma correlates with loss of agrin immunoreactivity from brain capillary basal laminae *Acta Neuropathol* 107:311-318 doi:10.1007/s00401-003-0812-0
- Wolburg H (1995) Orthogonal arrays of intramembranous particles: a review with special reference to astrocytes *Journal fur Hirnforschung* 36:239-258
- Xie L et al. (2013) Sleep drives metabolite clearance from the adult brain *Science* 342:373-377 doi:10.1126/science.1241224
- Yadav BK, Oh SY, Kim NK, Shin BS (2014) Association of rs2075575 and rs9951307 polymorphisms of AQP-4 gene with leukoaraiosis *J Stroke Cerebrovasc Dis* 23:1199-1206 doi:10.1016/j.jstrokecerebrovasdis.2013.10.017
- Yamamoto N et al. (2001) Alterations in the expression of the AQP family in cultured rat astrocytes during hypoxia and reoxygenation *Brain Res Mol Brain Res* 90:26-38
- Yang B, Brown D, Verkman AS (1996) The mercurial insensitive water channel (AQP-4) forms orthogonal arrays in stably transfected Chinese hamster ovary cells *JBiolChem* 271:4577-4580
- Yang B, van Hoek AN, Verkman AS (1997) Very high single channel water permeability of aquaporin-4 in baculovirus-infected insect cells and liposomes reconstituted with purified aquaporin-4 *Biochemistry* 36:7625-7632 doi:10.1021/bi970231r
- Yang B, Zador Z, Verkman AS (2008) Glial cell aquaporin-4 overexpression in transgenic mice accelerates cytotoxic brain swelling *The Journal of biological chemistry* 283:15280-15286 doi:10.1074/jbc.M801425200
- Yang B, Zhao D, Solenov E, Verkman AS (2006a) Evidence from knockout mice against physiologically significant aquaporin 8-facilitated ammonia transport *Am J Physiol Cell Physiol* 291:C417-423 doi:10.1152/ajpcell.00057.2006
- Yang B, Zhao D, Verkman AS (2006b) Evidence against functionally significant aquaporin expression in mitochondria *The Journal of biological chemistry* 281:16202-16206 doi:10.1074/jbc.M601864200
- Yang J et al. (2011) Loss of astrocyte polarization in the tg-ArcSwe mouse model of Alzheimer's disease *Journal of Alzheimer's disease : JAD* 27:711-722 doi:10.3233/JAD-2011-110725
- Yang J et al. (2017) AQP4 Association with Amyloid Deposition and Astrocyte Pathology in the Tg-ArcSwe Mouse Model of Alzheimer's Disease *Journal of Alzheimer's disease : JAD* 57:157-169 doi:10.3233/JAD-160957
- Yao X, Verkman AS (2017a) Complement regulator CD59 prevents peripheral organ injury in rats made seropositive for neuromyelitis optica immunoglobulin G *Acta Neuropathol Commun* 5:57 doi:10.1186/s40478-017-0462-4
- Yao X, Verkman AS (2017b) Marked central nervous system pathology in CD59 knockout rats following passive transfer of Neuromyelitis optica immunoglobulin G *Acta Neuropathol Commun* 5:15 doi:10.1186/s40478-017-0417-9
- Yukutake Y et al. (2008) Mercury chloride decreases the water permeability of aquaporin-4-reconstituted proteoliposomes *Biol Cell* 100:355-363 doi:10.1042/BC20070132
- Yukutake Y, Yasui M (2009) Regulation of water permeability through aquaporin-4 *Neuroscience* doi:S0306-4522(09)01713-8 [pii];10.1016/j.neuroscience.2009.10.029 [doi]
- Yurchenco PD (2011) Basement membranes: cell scaffoldings and signaling platforms *Cold Spring Harb Perspect Biol* 3 doi:10.1101/cshperspect.a004911
- Zelenina M (2010) Regulation of brain aquaporins *Neurochem Int* 57:468-488 doi:10.1016/j.neuint.2010.03.022
- Zeuthen T (2010) Water-transporting proteins *J Membr Biol* 234:57-73 doi:10.1007/s00232-009-9216-y
- Zhou J, Kong H, Hua X, Xiao M, Ding J, Hu G (2008) Altered blood-brain barrier integrity in adult aquaporin-4 knockout mice *Neuroreport* 19:1-5 doi:10.1097/WNR.0b013e3282f2b4eb [doi];00001756-200801080-00001 [pii]
- Zimmermann K (1923) Der feinere Bau der Blutkapillaren *Z Anat Entwicklungsgesch* 68:29-109

ELSEVIER

journal homepage: www.elsevier.com/locate/epilepsyres

Mislocalization of AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy

Silje Alvestad^{a,1}, Janniche Hammer^{b,1}, Eystein Hellstrøm Hoddevik^b, Øivind Skare^c, Ursula Sonnewald^a, Mahmood Amiry-Moghaddam^{b,*}, Ole Petter Ottersen^b

^a Department of Neuroscience, Norwegian University of Science and Technology (NTNU), N-7489 Trondheim, Norway

^b Laboratory of Molecular Neuroscience, Centre for Molecular Biology and Neuroscience, and Department of Anatomy, Institute of Basic Medical Sciences, University of Oslo, N-0317 Oslo, Norway

^c Department of Nutrition, Institute of Basic Medical Sciences, University of Oslo, N-0317 Oslo, Norway

Received 19 January 2012; received in revised form 4 January 2013; accepted 6 January 2013

Available online 26 January 2013

KEYWORDS

Aquaporin-4;
Epilepsy;
Syntrophin;
Kainic acid;
Astrocyte polarity

Summary It has been suggested that loss of the astrocytic water channel aquaporin-4 (AQP4) from perivascular endfeet in sclerotic hippocampi contributes to increased seizure propensity in human mesial temporal lobe epilepsy (MTLE). Whether this loss occurs prior to or as a consequence of epilepsy development remains to be resolved. In the present study, we investigated whether the expression and distribution of AQP4 was altered prior to (i.e., in the latent phase) or after the onset of chronic epileptic seizures (i.e., in the chronic phase) in the kainate (KA) model of MTLE. Immunogold electron microscopic analysis revealed that AQP4 density in adluminal endfoot membranes was reduced in KA treated rats already in the latent phase, while the AQP4 density in the abluminal endfoot membrane was stable or slightly increased. The decrease in adluminal AQP4 immunogold labeling was accompanied by a reduction in the density of AQP4's anchoring protein alpha-syntrophin. The latent and chronic phases were associated with an upregulation of the M1 isoform of AQP4, as judged by semi-quantitative Western blot analysis. Taken together, the findings in this model suggest that a mislocalization of AQP4 – reflecting a loss of astrocyte polarization – is an integral part of the epileptogenic process. © 2013 Elsevier B.V. All rights reserved.

* Corresponding author at: University of Oslo, Laboratory of Molecular Neuroscience, Centre for Molecular Biology and Neuroscience, Institute of Basic Medical Science, Department of Anatomy, Sognsvannsveien 9, P.O. Box 1105, Blindern, 0317 Oslo, Norway. Tel.: +47 91742177; fax: +47 22851278.

E-mail address: mahmo@medisin.uio.no (M. Amiry-Moghaddam).

¹ These authors contributed equally to this work.

Introduction

Mesial temporal lobe epilepsy (MTLE) is one of the most common and also one of the most therapy resistant types of epilepsy. To date, there is no efficient way to prevent the development of epilepsy after a potential epileptogenic insult. The underlying causes of MTLE are incompletely understood, but hyperexcitability in neuronal networks is considered a key feature.

Over the past decades, astrocytes have emerged as central regulators of neuronal excitability by releasing neuroactive agents that modulate synaptic transmission (Papura et al., 1994; Tian et al., 2005; Thrane et al., 2011), as well as by controlling ion and water homeostasis (Amiry-Moghaddam and Ottersen, 2003). Neuronal firing causes a rise in extracellular potassium and shrinkage of the perisynaptic extracellular space. It has been proposed that the subsequent uptake of K⁺ into surrounding astrocytes is coupled with a concomitant flux of water through the astrocytic water channel aquaporin-4 (AQP4) (Nagelhus et al., 1999; Østby et al., 2009; Haj-Yasein et al., 2012). The water flux via perisynaptic AQP4 is likely linked to water redistribution via astrocytic endfeet whose adluminal membranes are enriched in AQP4. Notably, loss of perivascular AQP4 leads to swelling of the perivascular endfeet, delayed clearance of extracellular potassium, and enhanced seizure intensity in an animal model where alpha-syntrophin – the AQP4 anchoring protein – is deleted (Amiry-Moghaddam et al., 2003a,b).

However, recent research has shown that the interaction between AQP4 and K⁺ handling is more complex than originally anticipated (Lee et al., 2012; Jin et al., 2013). The idea that epileptogenicity might be coupled to glial dysfunction and perturbed water and K⁺ homeostasis is supported by genetic analyses. Thus, variants of the genes encoding AQP4 and Kir4.1 are associated with subgroups of patients with MTLE (Heuser et al., 2010). In sclerotic hippocampi of MTLE patients, T2 signal hyperintensity and increased apparent diffusion coefficient on magnetic resonance imaging signify increased water content (Bronen et al., 1991; Hugg et al., 1999). Previous studies have shown an overall increase in the level of AQP4 in sclerotic MTLE hippocampi compared to non-MTLE hippocampi (Lee et al., 2004; Eid et al., 2005). However, in MTLE hippocampi, the density of AQP4 along the perivascular endfoot membrane was decreased, while no difference was found in the endfoot membrane facing the neuropil. It was proposed that the loss of perivascular AQP4 contributes to increased seizure propensity via impaired K⁺ and water homeostasis (Eid et al., 2005). Whether this change in AQP4 distribution precedes development of epilepsy, or is a consequence thereof, remains to be resolved.

In the present study, we examined AQP4 expression in rat hippocampal formation following status epilepticus (SE) induced by systemic kainate injections. This widely used model of MTLE allows the latent phase after SE to be distinguished from the chronic phase – the latter characterized by spontaneous recurrent seizures. We aimed to resolve whether a change in AQP4 distribution occurs in this model, and if so, whether such a change precedes chronic seizures, as would be expected if AQP4 redistribution contributes to epileptogenicity. This was addressed by use of quantitative immunogold cytochemistry. Differences between subregions

of the hippocampal formation were assessed by studying CA1 and CA3 separately. We also investigated the contents of AQP4 isoforms by semi-quantitative Western blotting.

Materials and methods

Animal experiments

Male Sprague Dawley rats (~250 g) were obtained from Taconic M&B, Copenhagen, Denmark. All animals were treated in accordance with the European Convention (ETS 123 of 1986), and all protocols were approved by the Norwegian National Animal Research Authority. The animals were maintained under standard laboratory conditions at 22 °C, humidity of 66%, and a 12/12 h light/dark cycle, with free access to food and water. They were adapted to the above conditions for one week prior to the experiments. Body weight and general condition were monitored throughout the study period and in case of weight loss or lack of weight gain they were fed with water soaked chow.

Status epilepticus was induced by intraperitoneal injection of 10 mg/kg kainate (KA; from Sigma–Aldrich, St. Louis, MO, USA) in buffered saline. A total of 58 rats received KA, whereas 26 rats received the same volume of saline. The rats were placed in separate plastic cages immediately after the injections and monitored for at least 4 h for behavioral seizure manifestation. The seizure activity was described using the Racine score (Racine, 1972): Stage 1, facial clonus; Stage 2, nodding; Stage 3, forelimb clonus; Stage 4, forelimb clonus with rearing; Stage 5, rearing and falling. Status epilepticus was defined as continuous limbic seizures scored as class 4 or 5 and lasting for minimum 90 min. Out of the 58 KA injected rats, 23 fulfilled the criteria for status epilepticus, 15 died, and 20 did not reach status epilepticus. Rats that displayed status epilepticus were selected for further analysis. Two weeks after the injections, i.e., in the latent phase, 23 rats were sacrificed (12 KA injected rats and 11 controls). The rats were not monitored by EEG, and therefore we cannot rule out the possibility that some of the rats had electrographic epileptiform discharges, or even seizures, during the two weeks following SE. Spontaneous seizures were not observed by the animal caretakers or by the researchers at any point. The remaining 26 rats (11 KA injected rats and 15 controls) were video monitored 8–12 h per day, 7 days per week. They were sacrificed eleven weeks after the injections, i.e., in the chronic phase, when the KA injected rats had exhibited a minimum of two spontaneous generalized seizures. The rats were either prepared for Western blotting or for immunocytochemistry.

Immunohistochemistry

17 rats were used for immunocytochemistry (latent phase: 6 KA injected and 4 controls; chronic phase: 3 KA injected and 4 controls). The rats were deeply anesthetized with Equithesin (0.4 ml/100 g, i.p.) and perfused transcardially with a flush of approximately 30 ml of 2% dextran in 0.1 M phosphate buffer followed by 1 l of a fixative containing 4% formaldehyde and 0.5% glutaraldehyde in the same buffer. They were kept overnight at 4 °C before the brains were removed and stored in the above fixative

diluted 1:10 in 0.1 M phosphate buffer at 4°C. The brain was divided at the midline. The right hemisphere was used for light microscopy. It was cut in 50 µm serial sagittal sections on a vibratome (Technical Products International, St. Louis, MO, USA) and stored in 0.1 M phosphate buffer with 0.1% azide. The left hemisphere was used for electron microscopy. It was cut in 300 µm horizontal vibratome slices. Slices from the septotemporal middle of the hippocampal formation were used for microdissection of subregions and subjected to freeze substitution as described previously (Amiry-Moghaddam et al., 2003a). Briefly, the tissue blocks were cryoprotected in glycerol and rapidly frozen in liquid propane at −170°C. The frozen tissue was immersed into anhydrous methanol containing 0.5% uranyl acetate at −90°C in an automatic freeze substitution unit (EMAFS, Leica, Vienna, Austria), infiltrated with Lowicryl HM20 Resin (Lowi, Waldkraiburg, Germany) at −30°C, polymerized by ultraviolet light, sectioned ultrathin, transferred to one hole grids, and used for immunogold labeling.

Immunogold electron microscopy

On-grid immunogold labeling for AQP4 was carried out in CA1 and CA3 of KA injected rats (latent phase $n = 4$, chronic phase $n = 3$, both subregions) and age-matched controls ($n = 4$, both phases and subregions, except $n = 3$ for CA1 controls latent phase). The immunogold procedure included the following steps: (1) Etching with 1% H_2O_2 for 30 min; (2) 48 mM glycine in Tris buffer containing 46 mM NaCl and 0.1% Triton X-100 (TBST); (3) 2% human serum albumin (HSA) in TBST; (4) incubation overnight (room temperature) with rabbit polyclonal anti-AQP4 from Chemicon Millipore (Temecula, CA, USA; 1:100) for CA3 and from Sigma–Aldrich (1:500) for CA1; (5) 2% human serum albumin (HSA) in TBST; (6) incubation for 2 h with a 10 nm colloidal gold-conjugated secondary antibody from Abcam (Cambridge, UK) diluted 1:20 in TBST with 2% HSA and 5 mg/ml polyethyleneglycol. The sections were counterstained with uranyl acetate followed by lead citrate before being examined in a transmission electron microscope (Tecnai 12, FEI Company, Eindhoven, the Netherlands). For quantification, 15 capillaries were randomly selected (at low magnification) within the stratum radiatum of each hippocampal subregion. Images were acquired and gold particle linear densities (particles/µm) were quantified in clearly delineated ad- and abluminal membranes of perivascular astrocytic endfeet. The analysis was performed using analySIS Pro version 3.2 (Soft Imaging system GmbH, Münster, Germany). For the analysis of alpha-syntrophin, we used a rabbit affinity-purified polyclonal antibody to alpha-syntrophin (Syn259) diluted 1:150 (Peters et al., 1997). The procedure was similar to that used for the analysis of AQP4, with amendments in the following steps: (1) No H_2O_2 etching; (2) all buffers contained 154 mM NaCl; (6) the secondary antibody was a 15 nm colloidal gold-conjugated anti-rabbit antibody from Abcam (AB27236). For quantification, 15–20 capillaries from each section were selected at random. Images were obtained at 20,500× magnification, and gold particle linear densities were quantified in clearly delineated adluminal membranes of perivascular astrocytic endfeet in CA3 stratum radiatum.

Immunofluorescence microscopy

For double labeling immunocytochemistry (immunofluorescence), vibratome sections of hippocampal formation of KA injected rats (latent phase $n = 6$, chronic phase $n = 3$) and controls ($n = 4$, both phases) were simultaneously processed free-floating. The sections were treated with 1% Na-borohydride in Tris-saline buffer (50 mM Tris, 120 mM NaCl, pH 7.4) for 10 min to quench autofluorescence from glutaraldehyde. The sections were rinsed in Tris-saline buffer before incubation over night (4°C) in a mixture of two primary antibodies (goat polyclonal anti-AQP4, Santa Cruz, 1:300 and rabbit polyclonal anti-alpha-syntrophin, 1:1000, Abcam) in Tris-saline buffer. After rinsing in Tris-saline buffer, the sections were incubated for 1 h in a mixture of two secondary antibodies (Alexa 488 donkey anti-goat together with Cy3 donkey anti-rabbit, diluted 1:1000, Invitrogen – Molecular Probes, Eugene, OR, USA) in Tris-saline buffer followed by rinsing and mounting with ProLong-Gold Antifade Reagent (Invitrogen – Molecular Probes, Eugene, OR, USA). The sections were analyzed using an Axioplan2 microscope (Carl Zeiss, Jena, Germany) equipped with a Pascal LSM5 confocal unit (Carl Zeiss), and the micrographs were obtained with the same parameters to allow reliable comparison between sections.

Western blotting

The 32 rats used for Western blotting (latent phase: 6 KA injected and 7 age-matched controls; chronic phase: 8 KA injected and 11 age-matched controls) were decapitated, the brain removed from the skull immediately, and the hippocampal formation isolated on both sides. The tissue was stored at −80°C.

Frozen whole hippocampal formations and microdissected subregions were sonicated on ice in homogenization buffer (1% SDS, 10 mM sodium phosphate buffer, pH 7.4, 150 mM NaCl, 1 mM phenylmethylsulphonyl fluoride (PMSF), 10 mM ethylenediamine tetraacetic acid (EDTA)). Protein concentrations were determined with a BCA Protein Assay Kit and measured on a microplate reader at 570 nm (Tecan, Grödig, Austria).

Protein homogenates were mixed with sample loading buffer (62.5 mM Tris–HCl, 10% glycerol, 2% SDS, 5% 2-mercaptoethanol, 0.025% bromophenol blue), and 5 µg protein per lane were applied. Homogenized rat cerebellum was used to make a standard curve with increasing protein concentrations for each blot. The proteins were separated by electrophoresis on 1.5 mm 10% acrylamide gels (urea) and transferred onto PVDF membranes.

The membranes were blocked with 5% blocking agent (Enhanced Chemi-fluorescence kit) before incubation overnight with primary antibody against AQP4 (1:20,000, polyclonal, Santa Cruz) diluted in Tris-buffered saline (20 mM Tris, 137 mM NaCl, pH 7.6) with 0.05% Tween 20 (TBS-T). After extensive washing in TBS-T, the membranes were incubated with alkaline-phosphatase-conjugated secondary antibody to goat immunoglobulin (1:10,000, Sigma) for 1 h and then washed again before application of substrate. The chemifluorescent signals were detected on a Typhoon scanner 9410, and quantified using ImageQuant TL (GE

Figure 1 Electron micrographs showing AQP4 immunogold labeling (10nm gold particles). Sections from CA1 and CA3 stratum radiatum of control (A and D) and kainate-treated rats 14 days (latent phase; B and E) and 11 weeks (chronic phase; C and F) after kainate-induced status epilepticus. In age-matched controls, AQP4 was clearly enriched in the astrocytic endfoot membrane facing capillaries (*adluminal* membrane, single arrows), and to a lesser extent in the *abluminal* membranes facing neuropil (double arrows). (*) Astrocytic endfoot; (e) endothelial cell; (l) lumen of capillary. Scale bar 500 nm.

Figure 2 Quantitative analysis of the polarity of AQP4 immunogold labeling in CA1 stratum radiatum and CA3 stratum radiatum for the latent (A) and chronic phase (B). Polarity is defined as the ratio between adluminal/abluminal gold particle density. Median data points with confidence interval error bars for the adluminal/abluminal ratios are indicated. Asterisks, significantly different from control, *** $p < 0.001$ ($n = 3-4$ per group, ratio of estimated medians within groups, taking into account the number of repetitions; see Materials and Methods). KA, kainate injected rat.

Healthcare, Buckinghamshire, UK). The pixel intensity (volume) was transformed to relative level by using the standard curve based on cerebellar homogenate.

After detection of chemifluorescent signals, the membranes were stripped of bound antibodies and reprobed for GDH (1:100,000; Rothe et al., 1995). Briefly, the membranes were submerged in stripping buffer (100 mM mercaptoethanol, 2% SDS, 62.5 mM Tris-HCl, pH 6.7) and incubated at 50°C for 60 min with agitation. Then, the membranes were washed in TBS-T before incubation with antibodies and immunodetection and quantification as described above.

Statistical analysis

Count data from semi-quantitative immunogold analysis were analyzed by a Poisson mixed model, with a log link, using the glmer function in R. Fixed effects were included for every combination of phase, region, group and membrane. We assumed a dependency structure with random components on animals, images and replications, where images were nested inside animals, and replications inside images. Parameter estimates were obtained by maximum likelihood using a Laplace approximation of the likelihood function. The differences in the adluminal:abluminal ratio were analyzed by a Bayesian Poisson mixed model, using the MCMCglmm package in R. Data from Western blotting were analyzed by a linear mixed model, using the lme function in R (Pinheiro and Bates, 2002). Separate analyses were done for the M1 and M23 data. Fixed effects were included for every combination of phase and group. We assumed a dependency structure with random components on animals. Log transformations were applied as quantile-quantile plots showed these transformations to give the best fits to the Gaussian distribution. Parameter estimates were obtained by maximum likelihood for the fixed effects and by restricted maximum likelihood for the variances of the random effects.

Results

Immunogold analysis of AQP4 and alpha-syntrophin

An immunogold approach was used to analyze the distribution of AQP4 in astrocytic endfoot membranes before and after development of spontaneous seizures, i.e. in the latent phase (two weeks after KA-induced status epilepticus) and in the chronic phase (11 weeks after KA). All animals that experienced status epilepticus went on to display recurrent seizures in the chronic phase. The linear density of gold particles was recorded in the endfoot membrane facing capillaries (adluminal membrane) and in the endfoot membrane facing neuropil (abluminal membrane; see Fig. 1), followed by an assessment of the ratio between the two membrane domains. This ratio (adluminal:abluminal) ranged from 4:1 to 6:1 in CA1, and from 8:1 to 10:1 in CA3 control groups (Fig. 2). The ratio was decreased in KA treated animals in both latent and chronic phases and in both regions, compared with controls; $p < 0.001$ (Fig. 2A). Actual adluminal:abluminal signal ratios were 3:1 and 2:1 (CA1), and 4:1 and 2:1 (CA3), in latent and chronic phase, respectively. The decreased ratio reflected a reduced labeling density in adluminal membranes (CA1: latent $p < 0.0001$, chronic $p = 0.0336$; CA3: latent $p = 0.0079$), combined with a stable or slightly increased labeling in the abluminal membranes (CA1: latent $p = 0.0386$, chronic $p = 0.0313$; CA3: chronic 0.0018; Fig. 4A–D). Endfeet appeared swollen in KA-treated animals compared with controls, and more prominently so in the latent phase (Fig. 1B and E) than in the chronic phase (Fig. 1C and F).

The immunogold signal for alpha-syntrophin (a dystrophin associated protein responsible for anchoring AQP4 in adluminal endfoot membranes) was reduced in KA treated animals (Figs. 3 and 4E). Statistical significance was reached in the latent phase ($p = 0.0135$), but not in the chronic phase ($p = 0.0555$).

The control animals were age-matched to the KA treated animals. To check for age-dependent changes, we compared

Figure 3 Electron micrographs showing alpha-syntrophin immunogold labeling (15 nm gold particles). Sections from CA3 stratum radiatum of the age-matched controls (A, B), kainate-treated rats 14 days (latent phase; C) and 11 weeks (chronic phase; D) after kainate-induced status epilepticus. Alpha-syntrophin is enriched in the astrocytic endfoot membrane facing capillaries (*adluminal* membrane, single arrows). (e) Endothelial cell; (l) lumen of capillary. Scale bar 1 μ m.

the two control groups (latent vs. chronic) with respect to AQP4 and alpha-syntrophin immunogold labeling. This analysis showed that the density of AQP4 increases slightly with age – but only in the abluminal endfoot membranes of CA1 (Fig. 4A and B; $p=0.042$). The signal for alpha-syntrophin increased in the *adluminal* membrane (Fig. 4E; $p=0.026$). The two control groups did not differ in regard to other parameters, i.e., *adluminal*/*abluminal* ratios for AQP4 (Fig. 2), the *adluminal* linear density of AQP4 immunogold labeling in CA1 (Fig. 4A and B), or linear density of AQP4 immunogold labeling in either membrane of CA3 (Fig. 4C and D).

Immunofluorescence analysis of AQP4 and alpha-syntrophin

In control animals, immunofluorescence signals for AQP4 and alpha-syntrophin were found in association with brain microvessels (Figs. 5A–C and 6A–C). This is consistent with the immunogold data showing a concentration of either molecule in astrocytic endfoot membranes. AQP4 immunofluorescence also occurs in the neuropil, albeit at lower intensities (Figs. 5A and 6A). Previous immunogold analyses (Nielsen et al., 1997) revealed that the neuropil labeling reflects AQP4 pools in astrocytic membranes,

Figure 4 Top: Quantitative immunogold analysis of AQP4 gold particle density in the stratum radiatum of CA1 and CA3. Box plot representation of the gold particle linear densities in adluminal and abluminal membranes of astrocytic endfeet in the latent (A and C) and chronic phase (B and D) after kainate-induced status epilepticus, compared with age-matched controls. Box, inter quartile range (IQR); line across box, median; whiskers, the lowest datum still within 1.5 IQR of the lower quartile, and the highest datum still within 1.5 IQR of the upper quartile; asterisks, significantly different from control, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ ($n = 3-4$ per group, taking into account the number of repetitions; see Materials and Methods). The data points indicating mean (dot) and standard error (lines ending with arrows) are shown beside each box. KA, kainate injected rat. Bottom: Quantitative immunogold analysis of alpha-syntrophin in the stratum radiatum of CA3. Box plot representation of the gold particle linear density, along with the data points indicating the mean and standard error, in the adluminal membranes of astrocytic endfeet in the latent and chronic phase (E).

Figure 5 Immunofluorescence micrographs of septal CA1 (stratum radiatum) in sagittal sections in control (A–C), latent (D–F) and chronic phase (G–I) double labeled for AQP4 (green) and alpha-syntrophin (red). KA treated animals were compared with age-matched controls. Arrowheads indicate the perivascular labeling. Scale bar 20 μ m. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of the article.)

including those facing synapses and other neuronal elements. Judged by the immunogold data, kainate treated animals display a redistribution of AQP4 from adluminal end-foot membranes. In agreement, the immunofluorescence analysis revealed an increased neuropil labeling of AQP4, compared with controls (Figs. 5 and 6). Alpha-syntrophin immunoreactivity was used as a marker for adluminal membranes.

Quantification of AQP4 by Western blotting

The level of AQP4 protein in the hippocampal formation was analyzed by semi-quantitative Western blotting, using GDH as a reference. The latter enzyme is expressed mainly in glia and was shown previously to be unchanged in the same set of animals (Hammer et al., 2008). AQP4 occurs in two major isoforms: M1 and M23. In addition, the rat brain also contains a third isoform termed Mz. The high molecular weight band (Fig. 7A and B) probably represents an AQP4 dimer. The

expression of M1 was increased in the latent phase compared with age-matched controls, whereas M23 was not significantly different (Fig. 7C). Both the M1 and M23 isoforms showed significant increases in the chronic phase, with a more pronounced increase for M1 than for M23 (Fig. 7D). The high molecular weight band ($\sim$ 55 kDa) similarly showed an increase in the chronic phase (Fig. 7E and F).

Discussion

Loss of perivascular AQP4 has been reported in human MTLE hippocampi (Eid et al., 2005) but it has not been resolved whether this loss occurs prior to or as a consequence of epilepsy development. In the present study, we took advantage of an epilepsy model with a well defined latent phase and showed that loss of perivascular AQP4 precedes the development of chronic seizures. Taken together, the immunogold, immunofluorescence and Western data suggest that a mislocalization of AQP4 – i.e., a

Figure 6 Immunofluorescence micrographs of septal CA3 (stratum radiatum) in sagittal sections in control (A–C), latent (D–F) and chronic phase (G–I) double labeled for AQP4 (green) and alpha-syntrophin (red). KA treated animals were compared with age-matched controls. Arrowheads indicate the perivascular labeling. Scale bar 20 μ m. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of the article.)

redistribution of AQP4 from perivascular membranes to other astrocytic plasma membrane domains – is an integral part of the epileptogenic process. The possibility must be considered that the redistribution of AQP4 along the astrocyte membrane is indicative of a more general loss of functional polarization, reflecting a pervasive dysfunction of hippocampal astrocytes.

The fact that AQP4 mislocalization precedes the onset of chronic seizures does not necessarily imply any causal relation. However, it has been shown previously that a mislocalization of AQP4, similar to that found here, increases the severity of seizures in a hyperthermia model (Amiry-Moghaddam et al., 2003b). In the latter study, AQP4 mislocalization was induced by genetic deletion of alpha-syntrophin which normally serves to anchor AQP4 to adluminal endfoot membranes (for a review of anchoring mechanisms, see Amiry-Moghaddam and Ottersen, 2003). The same genetic model showed a delayed clearance of K^+ from activated hippocampal neuropil, thus providing a

possible mechanistic link between AQP4 mislocalization and seizure propensity (Amiry-Moghaddam et al., 2003b; review: Binder et al., 2012).

It remains to explain why AQP4 redistributes in astrocytic membranes in the latent phase of the present experimental model. Our immunogold analysis indicates that the latent phase is associated with a significant downregulation of alpha-syntrophin. Any decrease in alpha-syntrophin expression would be expected to promote a loss of AQP4 from endfoot membranes (Amiry-Moghaddam et al., 2003a). However, recent data indicate that the two molecules are mutually dependent, as deletion of AQP4 causes a partial loss of alpha-syntrophin (Eilert-Olsen et al., 2012). This implies that we cannot resolve what is the primary event – loss of alpha-syntrophin or loss of AQP4. Efforts should now be made to characterize in more detail the factors that promote and regulate the binding between AQP4 and its anchoring protein alpha-syntrophin. The nature of this binding could hold important clues as to the mechanisms

Figure 7 Semi-quantitative analysis of Western blots from the latent and chronic phase. Top: Representative Western blots of whole hippocampal formation in the latent (A) and chronic phase (B) showing AQP4 isoforms at 32 kDa (M23), 34 kDa (M1) and ~37 kDa (Mz), the high molecular weight band of AQP4 (possibly a dimer) and GDH at 55 kDa. Bottom: Quantitative analysis of the immunoblots for the different AQP4 isoforms and the high molecular weight band of AQP4 in the latent (C, E) and chronic phase (D, F). Each sample represented by the mean of two to three separate blots. The data points indicating mean (dot) and standard error (lines ending with arrows) are shown beside each box. The pixel intensity (volume) is transformed to relative level by using a standard curve based on cerebellar homogenate (see Materials and Methods). The statistical analysis is based on raw data taking into account the number of repetitions (see Materials and Methods). Box, inter quartile range; line across box, median; whiskers, maximum and minimum; open circle, outlier; asterisk, significantly different from control, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ ($n = 6-11$ per group; see Materials and Methods). Co, control; KA, kainate injected rat.

underlying epileptogenesis in humans as well as in experimental models.

A comment should be made when it comes to the observation of a relatively higher increased level of the M1 isoform of AQP4 compared to the M23 isoform in KA treated animals. As pointed out, AQP4 is largely confined to astrocytic membranes (Nielsen et al., 1997). Thus, an increased immunoblot signal for AQP4, combined with a loss of AQP4 from perivascular endfeet, is likely to reflect a redistribution of AQP4 from endfoot to non-endfoot astrocytic membranes. A gliosis, coupled with a tangential displacement of AQP4 along the astrocytic membranes, would account for such a redistribution. We showed previously that gliosis is a feature of the latent as well as the chronic phase of the kainate model (Hammer et al., 2008). If the redistribution of AQP4 were fully explained by tangential movement of AQP4 along the astrocyte plasma membrane, one would not expect any shift in the relative abundance of the two main AQP4 isoforms, M1 and M23. In contrast, such a shift would occur if an increased synthesis of AQP4 was initiated in order to match the increased expanse of astrocytic membranes in the gliotic areas. Thus, several lines of evidence indicate that the M1 isoform is the predominant isoform in non-endfoot membranes while M23 is the major constituent of the square arrays that populate the adluminal endfoot membranes (Sorbo et al., 2008; Nicchia et al., 2010).

Taking advantage of the fact that the two control groups differed in age by nine weeks, we looked for putative age dependent changes in AQP4 and alpha-syntrophin immunolabelling. We found that the concentration of AQP4 increases slightly with age in the abluminal endfoot membrane and that the signal for alpha-syntrophin increases in the adluminal membrane. The biological significance of these findings will have to be explored in future studies.

The present study is the first to employ electron microscopy to analyze AQP4 expression in the kainate model of epilepsy. Electron microscopy is required to distinguish between different astrocytic plasma membrane domains and to provide quantitative estimates of AQP4 mislocalization. Previous studies of AQP4 in this model were based solely on light microscopic analyses (Kim et al., 2009, 2010), precluding direct comparisons.

Conclusion

Mislocalization of astrocytic AQP4, as reflected by a loss of perivascular AQP4 combined with a stable or increased AQP4 pool in non-endfoot membranes, is found in patients with MTE (Eid et al., 2005) and in animals with targeted depletion of the gene encoding alpha-syntrophin (Amiry-Moghaddam et al., 2003b). In both settings, the AQP4 mislocalization is associated with increased seizure propensity. Here, we show that a similar mislocalization occurs in the kainate model of epilepsy. The use of this model allowed us to conclude that AQP4 mislocalization precedes the chronic phase of epilepsy, thus presenting the possibility that the perturbed AQP4 expression is of pathophysiological relevance. A deranged water and K⁺ homeostasis may constitute the link between the present observations and the epilepsy phenotype. Our findings warrant further

studies to characterize in more detail how loss of astrocyte polarization affects neuronal excitability and seizure propensity.

Acknowledgements

The authors thank Karen Marie Gujord, Bjørge Riber, Jorunn Knutsen, and Bente Urfjell for technical assistance and Carina V. Knudsen for the artwork. This project was supported by the Letten Foundation and the Polish-Norwegian Research Fund grant PNR-96.

References

- Amiry-Moghaddam, M., Ottersen, O.P., 2003. The molecular basis of water transport in the brain. *Nat. Rev. Neurosci.* 4 (12), 991–1001.
- Amiry-Moghaddam, M., et al., 2003a. An alpha-syntrophin-dependent pool of AQP4 in astroglial end-feet confers bidirectional water flow between blood and brain. *Proc. Natl. Acad. Sci. U. S. A.* 100, 2106–2111.
- Amiry-Moghaddam, M., et al., 2003b. Delayed K⁺ clearance associated with aquaporin-4 mislocalization: phenotypic defects in brains of alpha-syntrophin-null mice. *Proc. Natl. Acad. Sci. U. S. A.* 100, 13615–13620.
- Binder, et al., 2012. Aquaporin-4 and epilepsy. *Glia* 60 (8), 1203–1214.
- Brønen, R.A., et al., 1991. Imaging findings in hippocampal sclerosis: correlation with pathology. *Am. J. Neuroradiol.* 12, 933–940.
- Eid, T., et al., 2005. Loss of perivascular aquaporin 4 may underlie deficient water and K⁺ homeostasis in the human epileptogenic hippocampus. *Proc. Natl. Acad. Sci. U. S. A.* 102, 1193–1198.
- Eilert-Olsen, M., et al., 2012. Deletion of aquaporin-4 changes the perivascular glial protein scaffold without disrupting the brain endothelial barrier. *Glia* 60 (3), 432–440.
- Haj-Yasein, N.N., et al., 2012. Aquaporin-4 regulates extracellular space volume dynamics during high-frequency synaptic stimulation: a gene deletion study in mouse hippocampus. *Glia* 60 (6), 867–874.
- Hammer, J., et al., 2008. Expression of glutamine synthetase and glutamate dehydrogenase in the latent phase and chronic phase in the kainate model of temporal lobe epilepsy. *Glia* 56, 856–868.
- Heuser, K., et al., 2010. Variants of the genes encoding AQP4 and Kir4.1 are associated with subgroups of patients with temporal lobe epilepsy. *Epilepsy Res.* 88 (1), 55–64.
- Hugg, J.W., Butterworth, E.J., Kuzniecky, R.I., 1999. Diffusion mapping applied to mesial temporal lobe epilepsy: preliminary observations. *Neurology* 53, 173–176.
- Jin, B.J., et al., 2013. Aquaporin-4-dependent K⁺ and water transport modeled in brain extracellular space following neuroexcitation. *J. Gen. Physiol.* 141 (1), 119–132.
- Kim, J.E., et al., 2009. Differential expressions of aquaporin subtypes in astroglia in the hippocampus of chronic epileptic rats. *Neuroscience* 163, 781–789.
- Kim, J.E., et al., 2010. Astroglial loss and edema formation in the rat piriform cortex and hippocampus following pilocarpine-induced status epilepticus. *J. Comp. Neurol.* 518, 4612–4628.
- Lee, D.J., et al., 2012. Decreased expression of the glial water channel aquaporin-4 in the intrahippocampal kainic acid model of epileptogenesis. *Exp. Neurol.* 235 (1), 246–255.
- Lee, T.S., et al., 2004. Aquaporin-4 is increased in the sclerotic hippocampus in human temporal lobe epilepsy. *Acta Neuropathol.* 108, 493–502.

- Nagelhus, E.A., et al., 1999. Immunogold evidence suggests that coupling of K⁺ siphoning and water transport in rat retinal Muller cells is mediated by a coenrichment of Kir4.1 and AQP4 in specific membrane domains. *Glia* 26, 47–54.
- Nielsen, S., et al., 1997. Specialized membrane domains for water transport in glial cells: high-resolution immunogold cytochemistry of aquaporin-4 in rat brain. *J. Neurosci.* 17 (1), 171–180.
- Nicchia, G.P., et al., 2010. Higher order structure of aquaporin-4. *Neuroscience* 168, 903–914.
- Parpura, V., et al., 1994. Glutamate-mediated astrocyte-neuron signalling. *Nature* 369, 744–747.
- Peters, M.F., et al., 1997. Differential association of syntrophin pairs with the dystrophin complex. *J. Cell Biol.* 138, 81–83.
- Pinheiro, J.C., Bates, D.M., 2002. *Mixed-Effects Models in S and S-Plus*, Business & Economics, 1st ed. Springer.
- Racine, R.J., 1972. Modification of seizure activity by electrical stimulation. II. Motor seizure. *Electroencephalogr. Clin. Neurophysiol.* 32 (3), 281–294.
- Rothe, F., Brosz, M., Storm-Mathisen, J., 1995. Quantitative ultrastructural localization of glutamate dehydrogenase in the rat cerebellar cortex. *Neuroscience* 64 (4), iii–xvi.
- Sorbo, J.G., et al., 2008. The molecular composition of square arrays. *Biochemistry* 47, 2631–2637.
- Tian, G.F., et al., 2005. An astrocytic basis of epilepsy. *Nat. Med.* 11 (9), 973–981.
- Thrane, A.S., et al., 2011. Critical role of aquaporin-4 (AQP4) in astrocytic Ca²⁺ signaling events elicited by cerebral edema. *Proc. Natl. Acad. Sci. U. S. A.* 108 (2), 846–851.
- Østby, I., et al., 2009. Astrocytic mechanisms explaining neural-activity-induced shrinkage of extraneuronal space. *PLoS Comput. Biol.*, e1000272.

Factors determining the density of AQP4 water channel molecules at the brain–blood interface

Eystein Hellstrøm Hoddevik^{1,2} · Faraz Hameed Khan¹ · Soulmaz Rahmani¹ · Ole Petter Ottersen¹ · Henning Bünsow Boldt¹ · Mahmood Amiry-Moghaddam¹

Received: 14 April 2016 / Accepted: 4 September 2016

© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract Perivascular endfeet of astrocytes are enriched with aquaporin-4 (AQP4)—a water channel that is critically involved in water transport at the brain–blood interface and that recently was identified as a key molecule in a system for waste clearance. The factors that determine the size of the perivascular AQP4 pool remain to be identified. Here we show that the size of this pool differs considerably between brain regions, roughly mirroring regional differences in *Aqp4* mRNA copy numbers. We demonstrate that a targeted deletion of α -syntrophin—a member of the dystrophin complex responsible for AQP4 anchoring—removes a substantial and fairly constant proportion (79–94 %) of the perivascular AQP4 pool across the central nervous system (CNS). Quantitative immunogold analyses of AQP4 and α -syntrophin in perivascular membranes indicate that there is a fixed stoichiometry between these two molecules. Both molecules occur at higher densities in endfoot membrane domains facing pericytes than in endfoot membrane domains facing endothelial cells. Our data suggest that irrespective of region, endfoot targeting of α -syntrophin is the single most important factor determining the size of the perivascular AQP4 pool and hence the capacity for water transport at the brain–blood interface.

Keywords Astrocyte heterogeneity · Pericyte · Aquaporin-4 · α -Syntrophin · Immunogold histochemistry

Introduction

The water channel aquaporin-4 (AQP4) occurs in abundance at the brain–blood interface and may well be the most plentiful membrane molecule at this site, forming distinctive, orthogonal arrays that make up a substantial areal fraction of the perivascular endfoot membrane (Amiry-Moghaddam and Ottersen 2003; Nielsen et al. 1997; Rash et al. 1998). These arrays were identified in freeze fracture preparations long before AQP4 was cloned and characterised (Dermietzel 1973; Landis and Reese 1974; Rash et al. 1974). Orthogonal arrays are absent from *Aqp4*^{−/−} mice (Verbavatz et al. 1997) and are formed in oocytes transfected with AQP4 cDNA (Yang et al. 1996), bolstering the notion that these arrays are made up of AQP4.

The perivascular pool of AQP4 serves a number of important functions. Mimicking the effects of global AQP4 knockout (Manley et al. 2000; Papadopoulos et al. 2004), a selective depletion of perivascular AQP4 is rate limiting for water flux in the build-up phase and resolution phase of brain oedema (Amiry-Moghaddam et al. 2003a, 2004; Nakayama et al. 2016; Vajda et al. 2002). The perivascular AQP4 pool is critically involved in osmosensing and volume control (Benfenati et al. 2011; Jo et al. 2015; Mola et al. 2016). Recent studies suggest that perivascular AQP4 also plays a pivotal role in the clearance of waste products from brain, possibly by propelling a paravascular drainage route (Iloff et al. 2012). In addition, AQP4 has been implicated in K⁺ homeostasis and regulation of excitability (Amiry-Moghaddam et al. 2003b; Binder et al. 2004; Haj-Yasein et al. 2012), and loss of perivascular

✉ Mahmood Amiry-Moghaddam
mahmo@medisin.uio.no

¹ Division of Anatomy, Department of Molecular Medicine, Institute of Basic Medical Sciences, University of Oslo, Post box 1105, Blindern, 0317 Oslo, Norway

² Department of Pathology, Oslo University Hospital, Oslo, Norway

AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy (Alvestad et al. 2013). Mislocalisation of AQP4 or changes in the expression level of this water channel have been reported in hippocampi resected from patients with mesial temporal lobe epilepsy (Eid et al. 2005), in mouse models of Alzheimer's disease (Yang et al. 2011), and in several other pathological and experimental conditions where homeostatic functions are lost or compromised (Nagelhus and Ottersen 2013; Promeneur et al. 2013).

Given the pivotal role of perivascular AQP4 in brain physiology and pathophysiology the question arises as to how the size of this AQP4 pool is determined. There is no unequivocal evidence of AQP4 channel gating (Nagelhus and Ottersen 2013). Thus, current data favour the idea that the water transport capacity at the brain–blood interface is tightly coupled to the number of AQP4 molecules at this site. Here we show that the density of perivascular AQP4 channels varies considerably between regions, roughly matching regional differences in the abundance of *Aqp4* mRNA. All regions showed a substantial loss of perivascular AQP4 following targeted deletion of α -syntrophin. This loss was fairly constant across the central nervous system (CNS), ranging from 79 % in the spinal cord to 94 % in the neocortex. Consistent with these data, quantitative immunogold analyses pointed to a fixed stoichiometry between AQP4 and α -syntrophin in perivascular membranes. Our findings provide new insight in the factors that regulate the size of the perivascular AQP4 pool and hence the capacity for water transport and waste clearance in brain.

Methods

Animals

We used adult male and newborn C57BL/6 mice (Jackson Laboratories, Boulder, CO) and two previously described

transgenic mouse strains. *Aqp4*^{−/−} animals were generated as described (Thrane et al. 2011) and backcrossed 10 times on a C57BL/6 background; α -*syntrophin*^{−/−} (α -*Syn*^{−/−}) animals were generated as described (Adams et al. 2000). The mice were allowed ad libitum access to food and drinking water. Animal experiments were performed according to the European Council law on protection of laboratory animals, with the approval of the University of Oslo's Animal Care and Use Committee. Every effort was made to minimise the number of animals.

Antibodies

A list of all antibodies used in this study is shown in Table 1.

Transcardial perfusion fixation

Adult animals were anaesthetised with a single intraperitoneal injection of Equithesin (150 μ L) and transcardially perfused using 4 % formaldehyde in 0.1 M phosphate buffer (PB) for 15 min with an initial 20 s perfusion with ice cold 2 % dextran in 0.1 M PB. Prompt neck stiffness was used as an indication of satisfactory perfusion. Brains were removed, left in the fixation solution overnight and stored in a 1:10 dilution of the same solution in 0.1 M PB. For newborn (P0) animals, both perfusion velocity and anaesthesia dosage were titrated according to weight and animals were initially sedated using chloroform gas. We employed a modified version of the pH-shift perfusion technique (Eilert-Olsen et al. 2012), initially defined by Berod et al. (1981), for those animals processed further for high resolution electron microscopy to obtain better ultrastructure and minimise epitope loss. This allowed for one half of the brain being used for light and immunofluorescence microscopy while the other was used for embedding in Lowicryl resin.

Table 1 Antibodies

Method	Primary antibody	Secondary antibody
Immunogold	AQP4, host: rabbit, Sigma A5971, 1,0 mg/ml diluted 1:400	Goat anti-rabbit 15 nm, Abcam 1:20
	α -syntrophin (SYN259), host: rabbit, 6 mg/ml diluted 1:200, Gift from Dr. Marvin E. Adams	Goat anti-rabbit 15 nm, Abcam 1:20
Immunofluorescence	AQP4, host: rabbit, Sigma A5971, 1,0 mg/ml diluted 1:400	Alexa fluor 488, donkey anti-rabbit, Invitrogen, A21206
Immunoperoxidase	AQP4, host: rabbit, Sigma A5971; 1,0 mg/ml diluted 1:400	Biotin Donkey anti-rabbit IgG 31821, Pierce, Thermo Scientific, 1:100
		Streptavidin-biotinylated HRP, RPN1051 V, GE Healthcare, 1:100
Western blot	AQP4, anti-rabbit, Sigma A5971; 1,0 mg/ml diluted 1:5000	Mouse monoclonal anti-Rabbit IgG-alkaline phosphatase, A2556, 1:5000
	Actin, anti-rabbit, Sigma, A2066; 0,67 mg/ml diluted 1:200	

Table 2 Analysed CNS regions according to experimental assay

Method	Analysed regions and subregions
Immunogold histochemistry	Cortex; parietofrontal cortex, piriform cortex Hippocampus; stratum pyramidale, stratum radiatum, dentate gyrus Inferior colliculus Cerebellum; molecular layer, granular layer, white matter Spinal cord (thoracic); white matter, grey matter
Quantitative RT-PCR	Parietofrontal cortex, hippocampus, inferior colliculus, cerebellum, whole brain samples
Western blot	Parietofrontal cortex, inferior colliculus

Post-embedding immunogold electron microscopy

Brains ($n = 4$ per group, $n = 2$ for each specificity control) were cut into 0.5–1.0 mm slices, regions were dissected, cryoprotected, quick-frozen in liquid propane ($-170\text{ }^{\circ}\text{C}$), and subjected to freeze substitution. Specimens were embedded in methacrylate resin (Lowicryl HM20) and polymerised by UV light below $0\text{ }^{\circ}\text{C}$. Ultrathin sections (70–100 nm) were cut using an Ultratome (Reichert Ultracut S, Leica) and placed on 300 mesh grids. Immunogold labelling was carried out as previously described (Promeneur et al. 2013). Briefly, sections were rinsed in Tris-buffered saline with Triton X-100 (TBS-T; 5 mM Tris-HCl, 0.3 % NaCl, 0.1 % Triton X-100), incubated in 2 % human serum albumin (HSA), followed by primary antibody (anti-AQP4 or anti- α -syntrophin) overnight, secondary antibody (15 nm gold) for 90 min, and contrasted with 2 % uranyl acetate for 90 s and 0.3 % lead citrate for 90 s. The sections were examined using a Tecnai 12 electron microscope at 80 kV. The examiner was blinded for animal genotype. Appropriate knockout tissue was used as negative control (*Aqp4*^{−/−} and *α -Syn*^{−/−} mice) for the two immunogold experiments.

A list of the regions dissected and analysed in this study is shown in Table 2.

Immunogold quantitation

Quantitative analysis was performed as previously described (Amiry-Moghaddam et al. 2004; Lunde et al. 2015). Briefly, images of 20–30 capillaries were acquired from each subregion present on each section. Care was taken to acquire images so that a similar distance of astrocyte membrane abutting on endothelium and pericytes was shown on each picture. We defined inclusion and exclusion criteria for capillaries, astrocyte endfeet and pericytes prior to image collection. In particular, a pericyte was defined as a perivascular cell surrounded by a clearly defined ad- and

abluminal basal lamina. Linear densities of gold particles over astrocyte membranes were determined by an extension of analysis [Soft Imaging Systems (SIS), Münster, Germany]. Linear densities were determined semi-automatically and transferred to SPSS Version 22 (SPSS, Chicago, IL) for statistical analysis.

Immunofluorescence

Mouse brains ($n = 4$ per study group, other half of brains processed for electron microscopy) were cryoprotected in increasing levels of saccharose dissolved in 0.1 M PB (10–20–30 %), and snap frozen in 30 % saccharose on a freeze microtome (Leica). Parasagittal brain sections (30 μm) were cut and collected in wells of sterile cell culture plates containing 0.4 % PFA diluted in 0.1 PB and stored in cold room ($4\text{ }^{\circ}\text{C}$). For immunofluorescence, sections were washed with 0.01 M phosphate buffered saline (PBS) and permeabilised for 10 min with 0.1 % Triton X-100 in PBS. After blocking with 2 % bovine serum albumin (BSA) in PBS, sections were incubated overnight with primary antibodies diluted in blocking solution. Following three washes with PBS, sections were incubated with secondary antibodies diluted in blocking solution for 90 min. After a further wash with PBS, sections were incubated for 5 min with a Hoechst preparation, thereafter washed with PBS and mounted using Prolong Gold anti-fade reagent (Thermo Fisher). Confocal images were acquired using a Leica fluorescence microscope.

Immunoperoxidase

Mouse brains ($n = 3$ per group) were cryoprotected and sectioned as described above. Sections were washed with PBS, exposed to 2 % H_2O_2 for 10 min to deplete endogenous peroxidase activity, and washed three times with PBS. Blocking was done using 2 % BSA in PBS-T (PBS, 0.1 % Triton X-100). Incubation with primary antibody diluted in blocking buffer was done overnight in cold room ($4\text{ }^{\circ}\text{C}$), followed by three washes with PBS-T. A 1 h incubation with secondary antibody in blocking buffer ensued, followed by three washes with PBS-T, incubation for 1 h with biotinylated-streptavidin horseradish peroxidase complex (GE healthcare UK limited, Rpn 1051 V) diluted 1:100 in PBS-T, three washes with PBS-T, then three washes with PBS. Sections were then incubated for 5 min with 0.5 mg/mL diaminobenzidine (3,3-diaminobenzidine tetrahydrochloride 10 mg/tab, Sigma D5905, DAB) dissolved in 0.1 M PB and exposed to DAB solution containing 0.01 % H_2O_2 (90 s). The DAB/HRP reaction was stopped by three washes with 0.1 M PB. Sections were mounted using glycerine gelatine, scanned

using a Mirax scanner (Zeiss) and examined using a light microscope.

Lysates and tissue processing for Western blotting and quantitative RT-PCR

WT and α -Syn^{-/-} mice were anaesthetised with isoflurane and decapitated. Brains were dissected out and split in two with a midline, sagittal cut. Regional dissections ($n = 4$) and whole brain samples ($n = 4$) were processed for quantitative RT-PCR analysis by overnight incubation in RNeasy Lysis Buffer (Qiagen) and storage at -80°C until further processing. Regional dissections ($n = 4$) from parietal cortex and colliculus inferior were obtained and processed for western blotting. Samples were put into 2 mL Lysing matrix D (MP Biomedicals) Eppendorf tubes containing RIPA (radio-immunoprecipitation assay) homogenisation buffer with protease inhibitor cocktail (Roche) added immediately before use, samples were homogenised in a FastPrep FP120 Cell Disrupter (MP Biomedicals), shaken at intervals and intermittently cooled on ice. Supernatant of a $9300 \times g$ spin for 30 min at 4°C was used for immunoblotting; pellets were frozen at -80°C . Protein concentrations were determined with a DC protein assay (Bio-Rad) according to the manufacturer's instructions. Equal numbers of right and left hemispheres were included in each analysis.

Western blot

Gels were cast using the Mini-Protean III cell from Bio-Rad as previously described (Sorbo et al. 2007). Both stacking and resolving gels were cast with added 3 M urea. 12 % SDS-PAGE gels were run at 150 V for 60 min, blotted with the Criterion cell (Bio-Rad), equilibrated with Towbin buffer (25 mM Tris, 192 mM glycine) supplemented with 20 % methanol and blotted onto 0.2 μm pore size PVDF membranes (Bio-Rad) at 100 V for 40 min. Blots were rinsed briefly in TBS-T (20 mM Tris; 137 mM NaCl; 0.05 % Tween-20 (Sigma), 0.05 % NaN_3 ; pH = 7.6) prior to incubation for 30 min with 5 % non-fat dried milk powder (Applichem) dissolved in TBS-T (blocking buffer). Blots were incubated with rabbit anti-AQP4 (Sigma) diluted in blocking buffer overnight at 4°C , then washed in TBS-T, incubated in a 1:5000 dilution in TBS-T of alkaline phosphatase conjugated to mouse monoclonal anti-Rabbit IgG (A2556 Sigma) for 1 h at room temperature and subsequently washed in TBS-T for 3 h. Finally, blots were incubated for 15 min with the ECF substrate (Amersham, ECF Kit), rinsed briefly in TBS-T, dried and scanned with a Kodak desktop scanner (Kodak Image Station 4000 MM PRO). Certain blots were re-probed with an anti-actin antibody (Sigma). Briefly, blots

were immersed in methanol, incubated with blocking buffer for 30 min, then incubated with anti-actin diluted in blocking buffer at 4°C overnight, rinsed and scanned as described above for the initial primary antibody. All samples were heat treated for 10 min at 37°C , spun at 13,000 rpm for 10 min, resuspended and loaded onto the gel. All samples were assayed for total protein content using the DC-kit (Bio-Rad) with BSA diluted in H_2O as standard. 5 μg of protein was loaded in each well, except for control tissue (*Aqp4*^{-/-}) where 10 μg was added. Precision plus protein standards Dual colour (Bio-rad) was used as Mw ladder.

Quantitative RT-PCR

Quantitative RT-PCR was carried out as described previously (Lunde et al. 2015). In brief, total RNA was extracted using RNeasy Lipid Tissue kit (Qiagen). 2 μg of RNA from whole brain and 100–1000 ng of RNA from regions were reverse-transcribed using oligo d(T)18 primer and RevertAid H Minus First Strand cDNA transcription reagents (Fermentas). The real-time PCR was set up using 2 μL template, corresponding to 5 ng cDNA, in a reaction volume of 20 μL containing AB Power SYBR Green PCR Master Mix (Applied Biosystems) and primers (200 nM) for detection of mouse *Aqp4*: TTTGGACCCGCAGT-TATCAT (forward) and GTTGTCTCCACCTCCATGT (reverse). Expression was assessed by absolute quantification using the StepOnePlus real-time PCR system (Applied Biosystems). Tbp (ACGGACAACCTGCGTTGATTT (forward); CAAGGCCTTCCAGCCTTATAG (reverse)) was used as endogenous control for normalisation of gene expression. SPSS was used for statistical analysis. *T* test was used for comparing *Aqp4* expression in WT and α -Syn^{-/-} mice. For comparing expression among regions, one-way ANOVA Bonferroni post hoc was used.

Results

AQP4 expression displays regional heterogeneity in adult and newborn murine brain

Light microscopic immunohistochemistry shows distinct caudo-rostral and ventro-dorsal gradients in AQP4 immunolabelling. Thus, labelling is notably stronger in cerebellum, olfactory bulb, spinal cord and brainstem than in cerebral cortex and striatum (Fig. 1). In the adult brain, AQP4 labelling is concentrated around blood vessels and subjacent to pia. A regional heterogeneity in AQP4 distribution is evident in the adult brain (Fig. 1a) as well as postnatally (Fig. 1b; also see Lunde et al. 2015). However, at this early stage the AQP4 immunolabelling is not

concentrated around vessels or subpially, but rather evenly distributed in the neuropil. Absence of labelling in *Aqp4*^{-/-} mice attests to the selectivity of the immunolabelling (Fig. 1b inset).

Evidence of non-uniform expression of AQP4 in pericapillary endfeet

Quantitative, high-resolution AQP4 immunogold histochemistry on ultrathin sections confirms and expands observations from light microscopy. AQP4 is found to be constitutively expressed in pericapillary endfeet and enriched in the endfoot membranes where levels display significant regional variation (Fig. 2a). Cortex (CX) and molecular layer of cerebellum (CB-m) display sparse labelling in contrast to the intense labelling of regions such as inferior colliculus (IC) and cerebellar granular layer (CB-gran). Linear density of gold particles along perivascular endfoot membranes was determined for capillaries of cortex, hippocampus, cerebellum, inferior colliculus and spinal cord and provides evidence of significant region-

specific variations. Heterogeneity is also present within subregions of the same anatomical region, shown here for hippocampus and cerebellum (Fig. 2b). No statistically significant difference could be found between grey and white matter regions of the spinal cord. In cerebellum, AQP4 density in the molecular layer is considerably lower than that in the granular layer and white matter. Within the hippocampal region, perivascular immunogold density is significantly lower in dentate gyrus than in stratum pyramidale and stratum radiatum. When the immunogold data is grouped according to region, the spinal cord and neocortex show the highest and lowest mean linear densities, respectively (14.80 vs. 9.82 particles per μm ; Fig. 4a, left, b, top).

Evidence of an α -syntrophin independent pericapillary pool of AQP4

Due to prior evidence implicating α -syntrophin as an anchoring molecule of endfoot AQP4 (Neely et al. 2001), we assessed AQP4 localisation in *α -Syn*^{-/-} mice.

Fig. 1 AQP4 expression displays regional heterogeneity in adult and newborn mouse brain. Sagittal sections from an adult (**a**) and a newborn (**b**) mouse brain demonstrate distinct regional differences in AQP4 immunoperoxidase labelling intensity. In both sections, neocortex and striatum demonstrate the least pronounced staining. Heterogeneity in labelling intensity is seen in a caudo-rostral and ventro-dorsal developmental pattern which is already established at birth (**b**). *Insets (right panel)* show higher magnification of cortex (CX) and brainstem (BS) for both ages. Cortex in adult mice exhibit

distinct labelling of glia limitans (*arrow*) and of vasculature (*arrowheads*). The overall labelling intensity is less pronounced in the newborn cortex, where the labelling is located mainly at the glia limitans (*arrow*) and vessels (*arrowheads*) close to the brain surface. The strong labelling observed in newborn brainstem and cerebellar white matter is located primarily in brain parenchyma, and to a much lesser extent around blood vessels (*arrowheads*). Specificity control (*left inset, Aqp4*^{-/-}) is shown on a sagittal section from an adult *Aqp4*^{-/-} mouse where no residual labelling is seen

Fig. 2 Regional and subregional perivascular heterogeneity of AQP4 immunogold labelling. Shown in **a** are portions of perivascular structures and cells from four CNS subregions. Each electron microscopic picture shows endothelium (end), the capillary lumen (L) and part of a pericyte (coloured green and annotated with asterisk) surrounded by a basal lamina on both sides. Immunogold particles can be seen along the length of the perivascular astrocyte endfoot membrane, but fewer gold particles per membrane length are seen in the membrane abutting endothelium (arrow) compared to the membrane domain abutting pericytes. Perivascular labelling is notably less intense in the endfoot from cortex (CX, upper left) compared to that of the inferior colliculus (IC, upper right). This heterogenic pattern is also present in layers within the same region, shown here for cerebellum (CB-m, CB-gran, lower panel), whereby immunogold linear density of granular cell layer is remarkably more pronounced than in cerebellar molecular layer. **b** Adjacent subregions within the hippocampal (HC-PYR; stratum pyramidale, HC-SR; stratum radiatum, HC-DGM; molecular layer of the dentate gyrus) and cerebellar formation (CB-G; cerebellar granular layer, CB-W; cerebellar white matter, CB-M; cerebellar molecular layer) demonstrate statistically significant differences in mean linear AQP4 density when compared within each region. In cerebellum, AQP4 density of the molecular layer (mean 9.45) is considerably lower than that of both granular layer (mean 15.84) and white matter (14.23). Within the hippocampal formation, perivascular immunogold density was significantly less pronounced in dentate gyrus (mean 9.24) than in stratum pyramidale (11.59) and stratum radiatum (13.86). Parieto-frontal cortex (CX-PF) also exhibits weaker labelling than piriform cortex (CX-PIR). No statistically significant difference was found between white (SC-W, mean 15.11) and grey (SC-G, mean 14.68) matter of spinal cord. Bar 100 nm. * and ** show $p < 0.05$ and $p < 0.001$, respectively. Error bars represent 95 % confidence interval

Genetic deletion of α -syntrophin leads to a pronounced reduction in perivascular AQP4 labelling in all regions and subregions analysed, as exemplified by images from cerebellum (Fig. 3). The representative image from an α -Syn $^{-/-}$ mouse (Fig. 3b) suggests a redistribution of AQP4 whereby perivascular and subpial AQP4 immunostaining in α -Syn $^{-/-}$ animals is not detectable, but staining of astrocyte processes surrounding Purkinje cells appears more intense than in wild-type mice (Fig. 3a). Immunogold detection of AQP4 is more sensitive than immunofluorescence, and—albeit for the notable reduction (79–94 %)—demonstrates residual

pericapillary AQP4 immunogold labelling in α -Syn $^{-/-}$ animals (Fig. 4a). Immunogold labelling was abolished in control sections from *Aqp4* $^{-/-}$ mice (specificity control, not shown).

The present analysis is restricted to the perivascular labelling. It should be noted, however, that the granular layer of the cerebellum contains a sizeable non-endfoot AQP4 pool that is largely independent of α -syntrophin and dystrophin (Fig. 3, see Amiry-Moghaddam et al. 2004; Nicchia et al. 2008b).

The size of residual AQP4 around capillaries is subject to regional variations

Residual AQP4 labelling around capillaries was quantified by immunogold histochemistry (Fig. 4a–c). Strikingly, in α -Syn $^{-/-}$ mice, perivascular AQP4 immunogold linear density in spinal cord is more than fourfold higher than in cortex (Fig. 4c). Following α -syntrophin gene deletion, cortex remains the region of lowest AQP4 linear density (mean 0.58 particles/μm) while spinal cord still displays the highest perivascular labelling (3.16 particles/μm). There is thus compelling evidence for retained, heterogeneous expression of AQP4 in α -Syn $^{-/-}$ mice.

Fig. 3 Impact of α -syntrophin gene deletion on AQP4 distribution. Immunofluorescence micrographs of cerebellar folia ($\times 20$ magnification) from a WT (a) and α -Syn $^{-/-}$ mouse (b) are shown, depicting the molecular (M), Purkinje (P), granular (G) and white matter (W) layers of the cerebellum. Both images show AQP4

immunostaining (green) and nuclear DAPI staining (blue). Genetic deletion of α -syntrophin leads to loss of perivascular (arrow head) and subpial (arrow) AQP4 labelling (arrow head). Staining of astrocyte processes surrounding Purkinje cells appears more intense in sections from α -Syn $^{-/-}$ mice (b) compared to WT controls (a)

Genetic deletion of α -syntrophin results in elimination of a similar proportion of AQP4 in all endfoot membranes, irrespective of the size of the AQP4 pool at this site

The relative AQP4 loss following α -syntrophin gene deletion is most pronounced in CX (94 %) followed by IC (89 %), CB (87 %), HC (83 %) and SC (79 %). The nominal loss in AQP4 linear density following α -syntrophin deletion, calculated as difference between means of linear density of AQP4 in knockout vs. wildtype (Fig. 4b), is greatest for IC and SC (11.71 and 11.64 particles/ μ m, respectively), followed by CB (10.96), HC (9.83) and CX (9.25).

Linear density of perivascular α -syntrophin correlates positively with that of AQP4

Perivascular α -syntrophin immunogold labelling (Fig. 4d) in WT mice shows how α -syntrophin levels correlate positively with that of AQP4 density (Fig. 4b) in endfoot membranes. Capillaries of cortical regions display the lowest density of α -syntrophin followed by hippocampus, cerebellum, spinal cord and inferior colliculus. As illustrated (Fig. 4b, d), the ratio of AQP4 to α -syntrophin varies little between regions. The calculated nominal ratio between AQP4 and α -syntrophin (in terms of immunogold

particles) is 5.82 for cortex, 5.89 for hippocampus, 4.84 for cerebellum, 4.54 for inferior colliculus and 5.87 for spinal cord. Labelling was abolished in control sections from α -Syn $^{-/-}$ mice (specificity control, not shown).

Quantitative RT-PCR demonstrates regional differences in gene expression

Shown in Fig. 5a is quantitative evidence of regional differences in *Aqp4* mRNA for both WT and α -Syn $^{-/-}$ genotypes. For WT animals, the mean copy number of *Aqp4* mRNA is highest in cerebellum, followed by inferior colliculus, whole brain, then hippocampus and cortex. Similarly, for α -Syn $^{-/-}$ animals, the mean copy number of *Aqp4* mRNA is highest in cerebellum followed by inferior colliculus, whole brain, hippocampus and cortex. For WT animals, the number of mRNA copies detected in cerebellum is fourfold that detected in cortex ($p < 0.001$), and the amount of mRNA copies detected in inferior colliculus is twice as high as in cortex ($p < 0.001$). We find no statistical difference between hippocampus and cortex in regard to total copy numbers of *Aqp4* mRNA. This is true for WT as well as α -Syn $^{-/-}$ mice. Notably, there is no difference between *Aqp4* expression in whole brain samples ($p = 0.188$) when comparing WT and α -Syn $^{-/-}$. The same holds true for all regions (from CB, IC, HC and CX).

Fig. 4 Regional perivascular heterogeneity is retained in $\alpha\text{-Syn}^{-/-}$ mice. **a** Perivascular AQP4 immunogold labelling in WT and $\alpha\text{-Syn}^{-/-}$ mice is illustrated by representative images from hippocampus (HC) and spinal cord (SC). Each picture shows endothelium (end), the capillary lumen (L) and part of a pericyte (coloured green and annotated with asterisk). For both regions, AQP4 labelling intensity is dramatically reduced in $\alpha\text{-Syn}^{-/-}$ mice compared to controls (WT) but some residual gold particles are observed. **b** Diagram (top) showing quantitation of AQP4 mean linear density in perivascular astrocyte domains for five brain regions. Significant differences in perivascular AQP4 labelling between the examined regions can be seen: cortex displays the lowest density (mean 9.82), followed by hippocampus (11.82), cerebellum (12.53), inferior colliculus (13.22) and spinal cord (14.80). Cortex has a statistically significant lower density when compared to the other regions ($p < 0.001$) and the density in spinal cord is significantly higher compared to the other four regions ($p < 0.001$). Following $\alpha\text{-Syn}^{-/-}$

syntrophin gene deletion, a substantial portion of membrane bound AQP4 in endfeet is lost. The calculated mean loss for each region is shown in the bottom diagram and appears proportional to the AQP4 density in the WT brain. The relative loss of AQP4 is also less pronounced in spinal cord compared to other regions. **c** Diagram showing mean AQP4 linear density from the five quantified regions in $\alpha\text{-Syn}^{-/-}$ mice. All regions in $\alpha\text{-Syn}^{-/-}$ mice lose most of the perivascular AQP4 labelling, but the regional heterogeneity pattern is retained: cortex remains the region of lowest AQP4 linear density (mean 0.58) and spinal cord still displays the highest perivascular labelling (3.16). No statistical difference is found between CB (1.59), HC (2.00) and IC (1.51). **d** Diagram showing quantitative analysis of perivascular $\alpha\text{-syntrophin}$ immunogold labelling in five regions. Regional heterogeneity of perivascular $\alpha\text{-syntrophin}$ labelling follows that of AQP4. Cortex displays the lowest linear density of $\alpha\text{-syntrophin}$ (mean 1.69) followed by HC (2.01), SC (2.52), CB (2.59) and IC (2.91). Error bars represent 95 % confidence intervals

Western blot analysis of AQP4 shows no difference between genotypes and demonstrates qualitative differences in AQP4 levels of CX and IC

Immunoblots from inferior colliculus (IC) and cortex demonstrate two major bands at the expected molecular weight, corresponding to M1 and M23 isoforms of AQP4, respectively (Fig. 5b, c). Both bands appear stronger in the immunoblot containing homogenates from inferior colliculus (Fig. 5b) compared to those from cortex (Fig. 5c) and provide qualitative data in accordance with the quantitative real-time PCR analysis. Anti-actin immunoblot on cortical samples is included to demonstrate presence of protein in *Aqp4*^{-/-} mice. Importantly, no difference in AQP4 labelling patterns or intensity can be seen between WT and α -Syn^{-/-} mice, as assessed for cortex and inferior colliculus (Fig. 5b, c). This is in line with earlier studies, indicating that deletion of α -syntrophin causes a mislocalisation rather than a loss of AQP4 from brain tissue (Neely et al. 2001).

Pericapillary astrocyte endfeet abutting pericytes are enriched in both AQP4 and α -syntrophin

As measured by quantitative immunogold labelling in WT mice, the overall AQP4 linear density is 15 % higher for endfoot membrane domains abutting pericytes than for endfoot membrane domains abutting endothelium (Fig. 6b, $p < 0.001$). This enrichment is statistically significant and present in all examined regions ($p < 0.05$ for CX, HC, SC and $p < 0.001$ for CB and IC), ranging from a factor of 1.26 higher in cortex to 1.08 in spinal cord. Nominally, the difference in linear density ranges from 2.60 (IC) to 1.01 (HC) gold particles per μm . We next investigated to what extent the regional, non-uniform expression of AQP4 in pericapillary endfeet was dependent on α -syntrophin. The difference between adjacent membrane domains remains following α -syntrophin gene deletion. Overall AQP4 linear density is 16 % higher in endfeet abutting pericytes compared to adjacent membranes abutting endothelium (Fig. 6c, $p < 0.05$) in α -Syn^{-/-} animals. The nominal difference of linear density between these two membrane domains is decreased in α -Syn^{-/-} mice, however, and a statistically significant difference ($p < 0.05$) was only observed in CX (Fig. 6c). As is the case for AQP4, α -syntrophin is also enriched in endfoot domains abutting pericytes compared to those abutting endothelium, a difference which reaches statistical significance for all examined regions (Fig. 6d, $p < 0.001$). The relative difference in α -syntrophin linear density between pericyte and endothelium abutting endfoot membrane domains is constant across examined regions, by a factor ranging from 1.27 (CB) to 1.38 (HC).

Fig. 5 Differential *Aqp4* mRNA expression and AQP4 protein in brain regions from WT and α -Syn^{-/-} mice. **a** Sagittal section of a mouse brain indicating the regions analysed. qRT-PCR analysis shown in **a** demonstrates that for WT animals, the mean copy number of *Aqp4* mRNA per nanogram of total RNA is the highest in CB (7928) followed by IC (4521), HC (1873) and CX (1920) while mean copy number for whole brain was 2787. For α -Syn^{-/-} animals, the mean copy number of *Aqp4* mRNA is the highest in CB (7115), followed by IC (3876), HC (1783) and CX (1368), mean copy number for whole brain was 2495. No statistically significant difference between the two genotypes is seen for any of the regional dissections, or for the whole brain samples. Cerebellum (CB) and inferior colliculus (IC) show higher expression of *Aqp4* mRNA than cortex (CX) and hippocampus (HC). Shown in **b**, **c** (top) are western blots showing anti-AQP4 labelling for two brain regions from WT and α -Syn^{-/-} (KO) mice. No difference in AQP4 labelling between genotypes is seen for either cortex or inferior colliculus (**b**, **c** top). Samples from wild-type and α -Syn^{-/-} mice ($n = 4$ for each genotype) are included. Two major bands, corresponding to M1 and M23 isoforms of AQP4, are detected in samples from both regions. Samples from inferior colliculus (**b**) consistently display more intense AQP4 immunostaining than those from cortex (**c**, top). No labelling is seen in the samples from *Aqp4*^{-/-} mice incubated with anti-AQP4 (specificity control). Error bars represent 95 % confidence intervals

Discussion

Given the emerging evidence pointing to the involvement of pericapillary AQP4 in a range of brain functions and pathologies there is a need to identify the factors that

Fig. 6 Pericapillary astrocyte endfeet abutting pericytes are enriched in both AQP4 and α -syntrophin. **a** Sagittal section of mouse brain indicating the dissected regions subjected to post-embedding immunogold histochemistry against AQP4 and α -syntrophin. Piriform and parietal cortex were grouped and reported as cortex. **b** Quantitative immunogold analysis demonstrates statistically significant differences in the mean linear density of AQP4 between perivascular astrocyte membrane domains abutting pericytes (per) and endothelium (end) in all the regions examined. To the far right is a graphic representation of all regions grouped together, annotated 'overall'. AQP4 linear density is 15 % higher in endfoot membrane domains abutting pericytes than for endfoot membrane domains abutting endothelium, **c** The overall difference between the AQP4 density in

the endfeet abutting pericytes compared to adjacent membranes abutting endothelium is retained in α -Syn^{-/-} mice. AQP4 linear density is 16 % higher in *per* compared to *end* membrane domains. Within subregions, the difference in AQP4 linear density of astrocyte membranes abutting endothelium compared to those abutting pericytes only reaches statistical significance in cortex. **d** Quantitative analysis of sections from WT animals shows that the linear density of α -syntrophin immunogold particles in astrocyte endfeet membrane domains abutting pericytes is consistently higher than in the endfeet abutting endothelium for all regions examined. Error bars represent 95 % confidence intervals. * and ** show $p < 0.05$ and $p < 0.001$, respectively

determine the size of this pool of AQP4. We decided to use the following, two step approach: first, we assessed the range by which the perivascular density of AQP4 varies across regions and subregions. Second, we aimed to identify the molecular basis of this heterogeneity.

The present study is restricted to the pericapillary AQP4 pool since this is the one that is most directly involved in regulating water flux across the brain–blood interface. Other AQP4 pools in brain (perisynaptic and subpial) have been investigated in previous studies (reviews: Amiry-Moghaddam and Ottersen 2003; Manley et al. 2004; Nagelhus and Ottersen 2013). It should also be emphasised that no attempt was made to discriminate between the two main isoforms of AQP4—M1 and M23. Thus, even if they differentially engage in supramolecular assemblies (square

arrays, Furman et al. 2003; Hiroaki et al. 2006; Nicchia et al. 2008a), these two isoforms flux water at the same rate (Jung et al. 1994). The AQP4 antibody presently used is directed to the C-terminus and recognises isoform M1 as well as M23.

Little is known about the extent to which the perivascular AQP4 pool differs in size between regions and subregions. In the only study that has addressed this issue thus far (Amiry-Moghaddam et al., 2004) we showed that the perivascular pool of AQP4 increased in size on moving from the molecular layer to the granular cell layer of the cerebellum. This study left us with several important questions that now emerge as highly important and relevant, given the recent breakthroughs in our understanding of water transport in brain. In particular, the notion that

perivascular AQP4 might be a key element in the brain's waste clearance system has drawn much attention and calls for new analyses (Igarashi et al. 2014; Iliff et al. 2012; Nagelhus and Ottersen 2013; Papadopoulos and Verkman 2013).

First, we ask whether the differences that were uncovered within the cerebellum (Amiry-Moghaddam et al. 2004) are a reflection of large scale differences between regions and subregions in the brain. Second, we ask whether such differences are rooted in differential transcription rates of the *Aqp4* gene and/or differential targeting of α -syntrophin to perivascular endfeet. It has long been known that α -syntrophin contributes to the anchoring of the perivascular AQP4 pool (Neely et al. 2001). However, it remains to be explored whether any heterogeneity in perivascular AQP4 expression can be attributed to differential expression of α -syntrophin.

Regional differences in the abundance of endfoot AQP4 imply regional differences in water transport capacity

Based on our quantitative immunogold data, we conclude that perivascular endfoot membranes differ considerably with respect to the amount of AQP4 they contain. Differences occur between regions and subregions, as well as on the level of individual capillaries (with endfoot membranes facing pericytes containing more AQP4 than endfoot membranes facing endothelia). This suggests that there is a corresponding non-uniformity in the capacity of water transport. AQP4 is the only aquaporin that is known to be concentrated in endfoot membranes. However, there is not necessarily a strict correlation between pericapillary AQP4 expression and water transport capacity. Thus, we cannot rule out the possibility that AQP4 is differentially regulated, depending on region, subregion or membrane microdomain. Several factors have been identified that affect water flux through AQP4 (Yukutake et al. 2008; Yukutake and Yasui 2010; Zelenina 2010) although direct evidence for bona fide gating is still missing (Nagelhus and Ottersen 2013). Available evidence suggests that the capacity for water transport at the brain–blood interface determines the propensity for oedema formation as well as the capability for oedema resolution (Amiry-Moghaddam et al. 2003a; Papadopoulos et al. 2004; Zeynalov et al. 2008).

Evidence for a stoichiometric coupling of AQP4 and α -syntrophin

By use of quantitative immunogold analyses in combination with genetic manipulation we showed that a major fraction of the AQP4 pool in endfoot membranes is

dependent on the presence of α -syntrophin. While it has long been known that α -syntrophin serves as a molecular anchor of endfoot AQP4 (Neely et al. 2001), this study is the first to show that the densities of α -syntrophin and AQP4 appear to be stoichiometrically coupled. Thus, the nominal ratio between the two molecules in terms of their immunogold labelling intensities varies little among regions (4.54 for inferior colliculus, 4.84 for cerebellum, 5.82 for cortex, 5.87 for spinal cord and 5.89 for hippocampus). Obviously, this ratio does not necessarily reflect the actual molecular ratios, as AQP4 and α -syntrophin may differ in regard to their labelling efficiency (defined as number of gold particle per target molecule). A more direct assessment of the actual molecular ratio (through biochemical approaches) must await the development of techniques for selective isolation of endfoot membranes. However, the uniformity in terms of AQP4/ α -syntrophin gold particle ratios is consistent with the idea that the complement of AQP4 in the endfoot membrane is governed by the number of α -syntrophin molecules that are targeted to this membrane.

Targeted disruption of the α -syntrophin gene deletes a similar fraction of the astrocyte endfoot pool of AQP4, irrespective of brain region

Our finding of a fairly uniform AQP4/ α -syntrophin ratio does not necessarily imply that α -syntrophin is the only anchoring molecule for AQP4. Indeed, confirming previous observations (Amiry-Moghaddam et al. 2004), we show that there is a residual pool of the astrocyte endfoot AQP4 following targeted deletion of α -syntrophin. Here we extend these observations by demonstrating that the size of the residual pool varies in proportion to the amount of AQP4 in WT endfeet. In other words, the percentage of perivascular AQP4 removed by deletion of α -syntrophin does not differ significantly between regions. The most probable explanation is that the residual AQP4 pool is anchored by a molecule whose expression is proportional to that of α -syntrophin. This unknown anchoring protein may well be another syntrophin, such as β -syntrophin (Puwarawuttipanit et al. 2006). This hypothesis will be explored once β -syntrophin knockout animals are made available.

α -Syntrophin and β -syntrophin are parts of a multi-member molecular assembly—the dystrophin associated protein complex (DAPC; Frigeri et al. 2001; Neely et al. 2001). Mdx mice that lack the brain dystrophin DP71 also lack a sizeable endfoot pool of AQP4 (Frigeri et al. 2001; Vajda et al. 2004). The DAP complex is held in place through its attachment to laminin (Culligan et al. 2001; Neely et al. 2001). This explains why there is such a sudden drop in AQP4 as soon as the endfoot plasma membrane is reflected away from the capillary.

In line with the important role of DAPC in AQP4 anchoring, deficient *O*-glycosylation of α -dystroglycan (a member of DAPC) causes a reduction in perivascular AQP4 (Michele et al. 2002; Rurak et al. 2007). Further, agrin (which interacts with α -dystroglycan) is required for normal expression of AQP4 in brain endfeet. It is important to note, however, that the brain also contains DAPC-independent pools of AQP4. One such pool is found in cerebellar astrocytes Nicchia et al. 2008b). This complexity notwithstanding, the present data show that α -syntrophin is a major player in controlling the expression of AQP4 at the brain–blood interface.

The level of AQP4 mRNA covaries with the level of perivascular AQP4

Our quantitative RT-PCR analysis revealed pronounced regional differences in the amount of *Aqp4* mRNA. The *Aqp4* transcript levels were high in those regions that displayed high levels of AQP4 in perivascular endfeet and were not sensitive to α -syntrophin deletion.

Heterogeneous distribution of AQP4 within individual endfoot membranes

Superimposed on the regional differences in the size of the endfoot AQP4 pool there is a heterogeneity in AQP4 distribution within individual endfoot membrane domains. Specifically, the densities of AQP4 and α -syntrophin immunogold particles are higher where endfoot membranes abut pericytes than where they abut endothelium. This is true not only for cortical astrocytes (Gundersen et al. 2013) but for all regions included in the present analysis. The local increase in the amount of AQP4 and α -syntrophin could be a direct effect of pericytes—through secretion of soluble factors or extracellular matrix molecules. As pericytes are contractile, they could also affect contiguous membranes mechanically. This is of relevance, as a recent cell-culture study from our laboratory showed increased AQP4 levels in astrocyte processes subjected to mechanical stress (Camassa et al. 2015).

The mechanistic basis for the non-uniform expression of AQP4 in astrocyte endfeet

Taken together, our data suggest that α -syntrophin is a key factor in regulating the AQP4 pool of perivascular endfoot membranes. Drawing on quantitative, high resolution immunogold analyses, we provide evidence that the number of AQP4 molecules is stoichiometrically coupled to that of α -syntrophin. A definition of the exact molecular ratio between these molecules must await the development

of appropriate techniques for isolating endfoot membranes. That AQP4 and α -syntrophin covary is not a coincidental correlation but reflects a causality: removal of α -syntrophin via genetic deletion removes a substantial fraction of endfoot AQP4 that varies little across regions. What determines the size of the AQP4 pool and hence the endfoot's water fluxing capacity, therefore, is the number of α -syntrophin molecules that are targeted to the endfoot membrane. The alternative explanation—that α -syntrophin regulates *Aqp4* expression through regulation at the transcriptional level—is ruled out by our finding that the number of *Aqp4* mRNA copies remains constant in α -syntrophin knockout animals.

Our experiments show that the level of AQP4 in endfeet is correlated to the level of *Aqp4* mRNA in the corresponding region. This observation does not necessarily imply that the level of endfoot AQP4 is controlled at the transcriptional level. Thus, the transcriptional regulation of *Aqp4* is complex (Pisani et al. 2011; Rossi et al. 2010), requiring further studies.

Conclusion

Three seminal findings have provided new insight in the mechanisms that control the size of the perivascular AQP4 pool, and thus the capacity for water transport and waste clearance in brain. First, targeting of α -syntrophin to endfoot membranes is the most important factor determining the size of the perivascular AQP4 pool. Second, those regions that show high densities of α -syntrophin in endfoot membranes also show high levels of *Aqp4* transcripts. Third, regional heterogeneities in *Aqp4* transcript levels persist after α -syntrophin deletion. While important in their own right, these observations point to the existence of a regulatory factor, upstream of α -syntrophin expression, that controls the size of the perivascular AQP4 pool. The identification of this factor stands as an important goal for future studies.

Acknowledgments The authors would like to thank Stan Froehner and Marv Adams for providing α -Syn^{-/-} animals; Gry Fluge Vindegal for help with animal breeding; Bjørge Riber, Karen Marie Gujord, Jorunn Knutsen, Yiqing Cai and Bashir Hakim for technical assistance; Carina Knutsen for graphics and Trygve Leergard for use of the Mirax Scanner.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Adams ME, Kramarcy N, Krall SP, Rossi SG, Rotundo RL, Sealock R, Froehner SC (2000) Absence of alpha-syntrophin leads to structurally aberrant neuromuscular synapses deficient in utrophin. *J Cell Biol* 150:1385–1398
- Alvestad S, Hammer J, Hoddevik EH, Skare O, Sonnewald U, Amiry-Moghaddam M, Ottersen OP (2013) Mislocalization of AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy. *Epilepsy Res* 105:30–41. doi:[10.1016/j.eplepsyres.2013.01.006](https://doi.org/10.1016/j.eplepsyres.2013.01.006)
- Amiry-Moghaddam M, Ottersen OP (2003) The molecular basis of water transport in the brain. *Nat Rev Neurosci* 4:991–1001. doi:[10.1038/nrn1252](https://doi.org/10.1038/nrn1252)
- Amiry-Moghaddam M et al (2003a) An alpha-syntrophin-dependent pool of AQP4 in astroglial end-feet confers bidirectional water flow between blood and brain. *Proc Natl Acad Sci USA* 100:2106–2111. doi:[10.1073/pnas.0437946100](https://doi.org/10.1073/pnas.0437946100)
- Amiry-Moghaddam M et al (2003b) Delayed K⁺ clearance associated with aquaporin-4 mislocalization: phenotypic defects in brains of alpha-syntrophin-null mice. *Proc Natl Acad Sci USA* 100:13615–13620. doi:[10.1073/pnas.2336064100](https://doi.org/10.1073/pnas.2336064100)
- Amiry-Moghaddam M et al (2004) Alpha-syntrophin deletion removes the perivascular but not endothelial pool of aquaporin-4 at the blood–brain barrier and delays the development of brain edema in an experimental model of acute hyponatremia. *FASEB J: Off Publ Fed Am Soc Exp Biol* 18:542–544. doi:[10.1096/fj.03-0869tje](https://doi.org/10.1096/fj.03-0869tje)
- Benfenati V, Caprini M, Dovizio M, Mylonakou MN, Ferroni S, Ottersen OP, Amiry-Moghaddam M (2011) An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex is essential for cell-volume control in astrocytes. *Proc Natl Acad Sci USA* 108:2563–2568. doi:[10.1073/pnas.1012867108](https://doi.org/10.1073/pnas.1012867108)
- Berod A, Hartman BK, Pujol JF (1981) Importance of fixation in immunohistochemistry: use of formaldehyde solutions at variable pH for the localization of tyrosine hydroxylase. *J Histochem Cytochem* 29:844–850
- Binder DK, Oshio K, Ma T, Verkman AS, Manley GT (2004) Increased seizure threshold in mice lacking aquaporin-4 water channels. *Neuroreport* 15(2):259–262
- Camassa LM et al (2015) Mechanisms underlying AQP4 accumulation in astrocyte endfeet. *Glia*. doi:[10.1002/glia.22878](https://doi.org/10.1002/glia.22878)
- Culligan K, Glover L, Dowling P, Ohlendieck K (2001) Brain dystrophin-glycoprotein complex: persistent expression of beta-dystroglycan, impaired oligomerization of Dp71 and up-regulation of utrophins in animal models of muscular dystrophy. *BMC Cell Biol* 2:2
- Dermietzel R (1973) Visualization by freeze-fracturing of regular structures in glial cell membranes. *Naturwissenschaften* 60:208
- Eid T et al (2005) Loss of perivascular aquaporin 4 may underlie deficient water and K⁺ homeostasis in the human epileptogenic hippocampus. *Proc Natl Acad Sci USA* 102:1193–1198. doi:[10.1073/pnas.0409308102](https://doi.org/10.1073/pnas.0409308102)
- Eilert-Olsen M et al (2012) Deletion of aquaporin-4 changes the perivascular glial protein scaffold without disrupting the brain endothelial barrier. *Glia* 60:432–440. doi:[10.1002/glia.22277](https://doi.org/10.1002/glia.22277)
- Frigeri A, Nicchia GP, Nico B, Quondamatteo F, Herken R, Roncali L, Svelto M (2001) Aquaporin-4 deficiency in skeletal muscle and brain of dystrophic mdx mice. *FASEB J: Off Publ Fed Am Soc Exp Biol* 15(1):90–98. doi:[10.1096/fj.00-0260com](https://doi.org/10.1096/fj.00-0260com)
- Furman CS, Gorelick-Feldman DA, Davidson KG, Yasumura T, Neely JD, Agre P, Rash JE (2003) Aquaporin-4 square array assembly: opposing actions of M1 and M23 isoforms. *Proc Natl Acad Sci USA* 100:13609–13614. doi:[10.1073/pnas.2235843100](https://doi.org/10.1073/pnas.2235843100)
- Gundersen GA, Vindedal GF, Skare O, Nagelhus EA (2013) Evidence that pericytes regulate aquaporin-4 polarization in mouse cortical astrocytes. *Brain Struct Funct*. doi:[10.1007/s00429-013-0629-0](https://doi.org/10.1007/s00429-013-0629-0)
- Haj-Yasein NN et al (2012) Aquaporin-4 regulates extracellular space volume dynamics during high-frequency synaptic stimulation: a gene deletion study in mouse hippocampus. *Glia* 60:867–874. doi:[10.1002/glia.22319](https://doi.org/10.1002/glia.22319)
- Hiroaki Y et al (2006) Implications of the aquaporin-4 structure on array formation and cell adhesion. *J Mol Biol* 355:628–639. doi:[10.1016/j.jmb.2005.10.081](https://doi.org/10.1016/j.jmb.2005.10.081)
- Igarashi H, Suzuki Y, Kwee IL, Nakada T (2014) Water influx into cerebrospinal fluid is significantly reduced in senile plaque bearing transgenic mice, supporting beta-amyloid clearance hypothesis of Alzheimer's disease. *Neurol Res* 36(12):1094–1098. doi:[10.1179/1743132814Y.0000000434](https://doi.org/10.1179/1743132814Y.0000000434)
- Iiiff JJ et al (2012) A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid beta. *Sci Transl Med* 4:147ra111. doi:[10.1126/scitranslmed.3003748](https://doi.org/10.1126/scitranslmed.3003748)
- Jo AO, Ryskamp DA, Phuong TT, Verkman AS, Yarishkin O, MacAulay N, Krizaj D (2015) TRPV4 and AQP4 channels synergistically regulate cell volume and calcium homeostasis in retinal muller glia. *J Neurosci* 35:13525–13537. doi:[10.1523/JNEUROSCI.1987-15.2015](https://doi.org/10.1523/JNEUROSCI.1987-15.2015)
- Jung JS, Preston GM, Smith BL, Guggino WB, Agre P (1994) Molecular structure of the water channel through aquaporin CHIP. The hourglass model. *J Biol Chem* 269:14648–14654
- Landis DM, Reese TS (1974) Arrays of particles in freeze-fractured astrocytic membranes. *J Cell Biol* 60:316–320
- Lunde LK, Camassa LM, Hoddevik EH, Khan FH, Ottersen OP, Boldt HB, Amiry-Moghaddam M (2015) Postnatal development of the molecular complex underlying astrocyte polarization. *Brain Struct Funct* 220:2087–2101. doi:[10.1007/s00429-014-0775-z](https://doi.org/10.1007/s00429-014-0775-z)
- Manley GT et al (2000) Aquaporin-4 deletion in mice reduces brain edema after acute water intoxication and ischemic stroke. *Nat Med* 6:159–163. doi:[10.1038/72256](https://doi.org/10.1038/72256)
- Manley GT, Binder DK, Papadopoulos MC, Verkman AS (2004) New insights into water transport and edema in the central nervous system from phenotype analysis of aquaporin-4 null mice. *Neuroscience* 129:983–991. doi:[10.1016/j.neuroscience.2004.06.088](https://doi.org/10.1016/j.neuroscience.2004.06.088)
- Michele DE et al (2002) Post-translational disruption of dystroglycan-ligand interactions in congenital muscular dystrophies. *Nature* 418:417–422. doi:[10.1038/nature00837](https://doi.org/10.1038/nature00837)
- Mola MG et al (2016) The speed of swelling kinetics modulates cell volume regulation and calcium signaling in astrocytes: a different point of view on the role of aquaporins. *Glia* 64:139–154. doi:[10.1002/glia.22921](https://doi.org/10.1002/glia.22921)
- Nagelhus EA, Ottersen OP (2013) Physiological roles of aquaporin-4 in brain. *Physiol Rev* 93:1543–1562. doi:[10.1152/physrev.00011.2013](https://doi.org/10.1152/physrev.00011.2013)
- Nakayama S, Migliati E, Amiry-Moghaddam M, Ottersen OP, Bhardwaj A (2016) Osmotherapy with hypertonic saline attenuates global cerebral edema following experimental cardiac arrest via perivascular pool of aquaporin-4. *Crit Care Med* 44:e702–e710. doi:[10.1097/CCM.0000000000001671](https://doi.org/10.1097/CCM.0000000000001671)
- Neely JD, Amiry-Moghaddam M, Ottersen OP, Froehner SC, Agre P, Adams ME (2001) Syntrophin-dependent expression and localization of aquaporin-4 water channel protein. *Proc Natl Acad Sci USA* 98:14108–14113. doi:[10.1073/pnas.241508198](https://doi.org/10.1073/pnas.241508198)
- Nicchia GP, Cogotzi L, Rossi A, Basco D, Brancaccio A, Svelto M, Frigeri A (2008a) Expression of multiple AQP4 pools in the plasma membrane and their association with the dystrophin complex. *J Neurochem* 105:2156–2165. doi:[10.1111/j.1471-4159.2008.05302.x](https://doi.org/10.1111/j.1471-4159.2008.05302.x)

- Nicchia GP, Rossi A, Nudel U, Svelto M, Frigeri A (2008b) Dystrophin-dependent and -independent AQP4 pools are expressed in the mouse brain. *Glia* 56:869–876. doi:[10.1002/glia.20661](https://doi.org/10.1002/glia.20661)
- Nielsen S, Nagelhus EA, Amiry-Moghaddam M, Bourque C, Agre P, Ottersen OP (1997) Specialized membrane domains for water transport in glial cells: high-resolution immunogold cytochemistry of aquaporin-4 in rat brain. *J Neurosci: Off J Soc Neurosci* 17:171–180
- Papadopoulos MC, Verkman AS (2013) Aquaporin water channels in the nervous system. *Nat Rev Neurosci* 14:265–277. doi:[10.1038/nrn3468](https://doi.org/10.1038/nrn3468)
- Papadopoulos MC, Manley GT, Krishna S, Verkman AS (2004) Aquaporin-4 facilitates reabsorption of excess fluid in vasogenic brain edema. *FASEB J* 18:1291–1293. doi:[10.1096/fj.04-1723fje](https://doi.org/10.1096/fj.04-1723fje)
- Pisani F, Rossi A, Nicchia GP, Svelto M, Frigeri A (2011) Translational regulation mechanisms of aquaporin-4 supramolecular organization in astrocytes. *Glia* 59:1923–1932. doi:[10.1002/glia.21234](https://doi.org/10.1002/glia.21234)
- Promeneur D, Lunde LK, Amiry-Moghaddam M, Agre P (2013) Protective role of brain water channel AQP4 in murine cerebral malaria. *Proc Natl Acad Sci USA* 110:1035–1040. doi:[10.1073/pnas.1220566110](https://doi.org/10.1073/pnas.1220566110)
- Puwarawuttipanit W et al (2006) Differential effect of alpha-syntrophin knockout on aquaporin-4 and Kir4.1 expression in retinal macroglial cells in mice. *Neuroscience* 137:165–175. doi:[10.1016/j.neuroscience.2005.08.051](https://doi.org/10.1016/j.neuroscience.2005.08.051)
- Rash JE, Staehelin LA, Ellisman MH (1974) Rectangular arrays of particles on freeze-cleaved plasma membranes are not gap junctions. *Exp Cell Res* 86:187–190
- Rash JE, Yasumura T, Hudson CS, Agre P, Nielsen S (1998) Direct immunogold labeling of aquaporin-4 in square arrays of astrocyte and ependymocyte plasma membranes in rat brain and spinal cord. *Proc Natl Acad Sci USA* 95:11981–11986
- Rossi A, Pisani F, Nicchia GP, Svelto M, Frigeri A (2010) Evidences for a leaky scanning mechanism for the synthesis of the shorter M23 protein isoform of aquaporin-4: implication in orthogonal array formation and neuromyelitis optica antibody interaction. *J Biol Chem* 285:4562–4569. doi:[10.1074/jbc.M109.069245](https://doi.org/10.1074/jbc.M109.069245)
- Rurak J, Noel G, Lui L, Joshi B, Moukhles H (2007) Distribution of potassium ion and water permeable channels at perivascular glia in brain and retina of the Large(myd) mouse. *J Neurochem* 103:1940–1953. doi:[10.1111/j.1471-4159.2007.04886.x](https://doi.org/10.1111/j.1471-4159.2007.04886.x)
- Sorbo JG, Moe SE, Holen T (2007) Early upregulation in nasal epithelium and strong expression in olfactory bulb glomeruli suggest a role for aquaporin-4 in olfaction. *FEBS Lett* 581:4884–4890. doi:[10.1016/j.febslet.2007.09.018](https://doi.org/10.1016/j.febslet.2007.09.018)
- Thrane AS et al (2011) Critical role of aquaporin-4 (AQP4) in astrocytic Ca^{2+} signaling events elicited by cerebral edema. *Proc Natl Acad Sci USA* 108:846–851. doi:[10.1073/pnas.1015217108](https://doi.org/10.1073/pnas.1015217108)
- Vajda Z et al (2002) Delayed onset of brain edema and mislocalization of aquaporin-4 in dystrophin-null transgenic mice. *Proc Natl Acad Sci USA* 99:13131–13136. doi:[10.1073/pnas.192457099](https://doi.org/10.1073/pnas.192457099)
- Vajda Z, Pedersen M, Doczi T, Sulyok E, Nielsen S (2004) Studies of mdx mice. *Neuroscience* 129:993–998. doi:[10.1016/j.neuroscience.2004.08.055](https://doi.org/10.1016/j.neuroscience.2004.08.055)
- Verbavatz JM, Ma T, Gobin R, Verkman AS (1997) Absence of orthogonal arrays in kidney, brain and muscle from transgenic knockout mice lacking water channel aquaporin-4. *J Cell Sci* 110(Pt 22):2855–2860
- Yang B, Brown D, Verkman AS (1996) The mercurial insensitive water channel (AQP-4) forms orthogonal arrays in stably transfected Chinese hamster ovary cells. *J Biol Chem* 271:4577–4580
- Yang J et al (2011) Loss of astrocyte polarization in the tg-ArcSwe mouse model of Alzheimer's disease. *J Alzheimer's Dis: JAD* 27:711–722. doi:[10.3233/JAD-2011-110725](https://doi.org/10.3233/JAD-2011-110725)
- Yukutake Y, Yasui M (2010) Regulation of water permeability through aquaporin-4. *Neuroscience* 168:885–891. doi:[10.1016/j.neuroscience.2009.10.029](https://doi.org/10.1016/j.neuroscience.2009.10.029)
- Yukutake Y et al (2008) Mercury chloride decreases the water permeability of aquaporin-4-reconstituted proteoliposomes. *Biol Cell* 100:355–363. doi:[10.1042/BC20070132](https://doi.org/10.1042/BC20070132)
- Zelenina M (2010) Regulation of brain aquaporins. *Neurochem Int* 57:468–488. doi:[10.1016/j.neuint.2010.03.022](https://doi.org/10.1016/j.neuint.2010.03.022)
- Zeynalov E, Chen CH, Froehner SC, Adams ME, Ottersen OP, Amiry-Moghaddam M, Bhardwaj A (2008) The perivascular pool of aquaporin-4 mediates the effect of osmotherapy in postischemic cerebral edema. *Crit Care Med* 36:2634–2640. doi:[10.1097/CCM.0b013e3181847853](https://doi.org/10.1097/CCM.0b013e3181847853)

Mechanisms Underlying AQP4 Accumulation in Astrocyte Endfeet

Laura Maria Azzurra Camassa,¹ Lisa K. Lunde,¹ Eystein H. Hoddevik,¹ Maria Stensland,² Henning B. Boldt,¹ Gustavo A. De Souza,² Ole P. Ottersen,¹ and Mahmood Amiry-Moghaddam¹

The brain–blood interface holds the key to our understanding of how cerebral blood flow is regulated and how water and solutes are exchanged between blood and brain. The highly specialized astrocytic membranes that enwrap brain microvessels are salient constituents of the brain–blood interface. These endfoot membranes contain a distinct set of molecules that is anchored to the subendothelial basal lamina forming an endfoot–basal lamina junctional complex. Here we explore the mechanisms underpinning the formation of this complex. By use of a tailor made model system we show that endothelial cells promote AQP4 accumulation by exerting an inductive effect through extracellular matrix components such as agrin, as well as through a direct mechanical interaction with the endfoot processes. Through the compounds they secrete, the endothelial cells also increase AQP4 expression. The present data suggest that the highly specialized gliovascular interface is established through inductive processes that include both chemical and mechanical factors.

GLIA 2015;00:000–000

Key words: astroglia, agrin, aquaporin, extracellular matrix, polarization, compartmentalization

Introduction

There is now compelling evidence that astrocytes are polarized with respect to protein expression and function and that they in this regard bear similarities to central neurons (Barres, 2008; Nedergaard et al., 2003; Seifert et al., 2006). Astrocytes that populate grey matter typically show a rich complexity of processes that engage in contact with thousands of synapses in their respective domains (Bushong et al., 2004; Iadecola and Nedergaard, 2007; Pekny et al., 2007; Wilhelmsson et al., 2006). However, most astrocytes also extend processes that abut on cerebral microvessels and form perivascular endfeet. The latter processes show a distinct repertoire of membrane proteins and serve specific roles. Evidence is accruing that the polarity of astrocytes is critical for astrocyte function and for their multiple homeostatic roles (Etienne-Manneville, 2008; Nagelhus and Ottersen, 2013; Wolburg et al., 2009). In agreement, a growing list of pathological conditions have been coupled to loss of astrocyte polarity,

including mesial temporal lobe epilepsy, ischemic brain damage, and Alzheimer's disease (Alvestad et al., 2013; Eid et al., 2005; Frydenlund et al., 2006; Yang et al., 2011).

Transport of water, K⁺, and glutamate are examples of processes that are strongly polarized in brain astrocytes. The water channel AQP4 is concentrated in perivascular endfeet where this protein serves as influx routes for water in pathological conditions associated with brain edema (Amiry-Moghaddam et al., 2003, 2004; Manley et al., 2004). Perivascular endfeet are also enriched in the K⁺ channel Kir4.1 which is involved in K⁺ spatial buffering and in K⁺ siphoning in particular (Nagelhus et al., 1999; Nagelhus and Ottersen, 2013; Newman et al., 1984; Strohschein et al., 2011). Glutamate transporters, on the other hand, are preferentially expressed in those delicate astrocytic processes that partially ensheath central excitatory synapses (Chaudhry et al., 1995).

In the absence of tight junctions, functionally distinct membrane domains must be defined by specific anchoring

View this article online at wileyonlinelibrary.com. DOI: 10.1002/glia.22878

Published online Month 00, 2015 in Wiley Online Library (wileyonlinelibrary.com). Received Dec 17, 2014, Accepted for publication June 5, 2015.

Address correspondence to Mahmood Amiry-Moghaddam, University of Oslo, Institute of Basic Medical Sciences, Laboratory of Molecular Neuroscience, Oslo, Norway. E-mail: mahmo@medisin.uio.no

From the ¹Laboratory of Molecular Neuroscience, Department of Molecular Medicine, Institute of Basic Medical Sciences, University of Oslo, Norway; ²Laboratory of Proteomic Research, Department of Immunology, University of Oslo, Norway

Additional Supporting Information may be found in the online version of this article.

mechanisms. Available data indicate that the dystrophin associated protein (DAP) complex is of particular importance when it comes to the definition of astrocyte polarity. This complex has been implicated in the anchoring of AQP4 and Kir4.1 to perivascular endfeet membranes. Notably, AQP4 is lost from endfeet membranes in animals that lack the dystrophin complex (mdx) and in animals with targeted deletion of the gene encoding the dystrophin associated protein α -syntrophin (Frigeri et al., 2001; Neely et al., 2001).

Given the fact that astrocytic polarity and loss thereof figure prominently in current research on astrocytic physiology and pathophysiology, it is essential to identify those factors that target the dystrophin complex to the endfoot membranes. Specifically, there is a need to characterize the conditions that are required to reconstitute polarity in cultured astrocytes. Normally, astrocytes in culture show no polarization and typically internalize many of the molecules that normally are restricted to the plasma membrane (Nicchia et al., 2004).

Here we have designed a model system that allows experimental manipulation of identified astrocytic processes. This model enables differentiation between astrocytic polarization (heterogeneity between different membrane domains) and membrane compartmentalization (localization to plasma membrane vs. intracellular membrane compartments). Our findings indicate that astrocytic polarization depends on a number of factors and that endothelial cells and extracellular matrix molecules play an instructive role in targeting AQP4 and the dystrophin complex to endfoot membranes.

Materials and Methods

Primary Culture of Astrocytes

Primary cultures of cortical astrocytes from newborn (1–2 days) Wistar rats and from wild type (WT) and α -syntrophin-null (α -SYN-KO) mice (Adams et al., 2000) were prepared according to the modified method of McCarthy and deVellis (McCarthy and deVellis, 1980; Schwartz and Wilson, 1992).

After decapitation and removal of meninges, cerebral cortices were collected in Falcon tubes containing 2 ml of fresh medium (FM) containing Dulbecco's modified Eagle's medium high glucose (DMEM-glutamax) with 10% fetal bovine serum (FBS) and penicillin/streptomycin (100 U mL^{-1}) (Gibco-Invitrogen). After mechanical dissociation and passage through cell strainers (70 μm Nylon from BD Falcon), undifferentiated rat brain cells were plated on poly-L-lysine (PLL) (Sigma-Aldrich) coated 75 cm^2 flasks (Corning® Flasks) with 15 mL of FM.

The flasks were maintained in an incubator with humidified atmosphere containing 5% CO_2 , at 37°C. The medium was changed every 2 days. To detach the microglial and precursor cells, the flasks were shaken for 2 min each time changing the medium during the first week. Immunostaining for glial fibrillary acidic protein (GFAP)

indicated that more than 95% of the cells were astrocytes. Immunostaining for β III-tubulin didn't show any neuronal contamination.

Co-culture on Filter/Cell Culture Inserts

One week after the preparation (when the cultured astrocytes reached confluence; see above), the cells were split and seeded at a concentration of $12 \times 10^4 \text{ cells cm}^{-2}$ on the bottom of cell culture inserts (BD Falcon), 1- μm pore size, and maintained in the incubator for 6 h to allow the cells to attach to the filter. Successively, inserts were turned in a six-multiwell plate filled with FM.

When the astrocytes reached 80% of confluence (around 7 days of culture), bEND3 cells, a mouse brain immortalized endothelial cell line (Omidi et al., 2003), were seeded on the top aspect of the filters, at the concentration of $2 \times 10^4 \text{ cells cm}^{-2}$ (Fig. 1A).

The co-cultures were kept in the incubator for 7–10 days, until the endothelial cells had formed a monolayer, and then immersion fixed with 4% formaldehyde (FA).

For assessing the inductive effects of extracellular matrix, the top aspects of the filters were coated with the extracellular matrix components, laminin- α 1, Matrigel (Sigma-Aldrich) or neuronal agrin (R&D System). In some experiments the bEND3 cells were replaced with HeLa cells (Scherer et al., 1953) or ECV304 cells, which is an immature human umbilical endothelia derived cell line (Barar et al., 2010).

Co-culture Conditioned Medium (CM)

Conditioned medium (CM) was prepared by cultivating the primary astrocytes and bEND3 cells in FM without FBS for 2–3 days.

The co-culture was prepared as described above. One day after seeding the bEND3 cells the co-culture was washed twice in PBS and FM was added. New FM was added to the co-cultures and the CM was collected five times with 2- to 3-days interval (CM1-5).

The CM was filtered through 0.22- μm filter (Millipore) and centrifuged at 2100 rpm to remove debris.

To study the effect of CM on primary cultures of astrocytes, CM was mixed 1:1 with FM and applied on the cells. Astrocytes were harvested for western blotting and qPCR (see below). The last fraction of CM (CM5) was used for mass spectrometry (see below).

CM and Soaked Beads

Polystyrene beads (PS03N, Bangs Laboratories) were centrifuged three times at 2,200g in Washing Buffer (WB, Bangs Laboratories) for 15 min at room temperature and the supernatant discarded. The washed beads were divided in two portions. One portion was kept as control and the other portion soaked in CM as described below:

Polystyrene beads and CM were kept at 4°C in a Falcon tube on a rotating table for 2 days. The CM-soaked beads were centrifuged for 15 min at 2,200g and the supernatant discarded. The sequence was repeated five times.

The beads soaked in CM were applied on the top of six filters (instead of bEND3 cells) and co-cultured with astrocytes seeded on the bottom. An aliquot of washed beads was used as negative control in another six filters. The two co-culture combinations were analyzed and compared with six filters on which bEND3 cells had been co-cultured with astrocytes.

FIGURE 1: Schematic representation of the co-culture system. **A:** bEND3 endothelial cells (End, yellow) are seeded on the top and primary culture of astrocytes (Ast, green) on the bottom of the filter. Extracellular matrix components, such as Matrigel (red), are applied on the top of the filter with or without bEND3. **B:** confocal microscopy images show astrocytic cell bodies on the bottom (Ast), and astrocytic endfeet (e) immunostained with GFAP (green), on the top, in proximity of an endothelial cell (End). Nuclei are visible in DAPI. Scale bar: 50 μm . **C:** left panel: electron microscopic image of an astrocytic foot process (e) extending from the cell body (Ast) through a pore in the filter toward the endothelial cells side (End), scale bar: 2 μm . Top, and bottom, right panel: higher magnification electron microscopic images of bEND3 cells (End), and astrocyte processes (Ast), respectively. Scale bars: 2 μm (top), and 1 μm (bottom). [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

Electron Microscopy on Co-culture

Cultures were immersion fixed in a mixture of 4% FA and 0.1% glutaraldehyde (GA) for 30 min. Filters were cut into squares of $\sim 1 \text{ mm}^3$, then cryoprotected in increasing concentrations of glycerol

(10, 20, and 30%, respectively) dissolved in 0.1M phosphate buffered saline (PBS), quick-frozen in liquid propane (-170°C), and subjected to freeze substitution (Neely et al., 2001; Takumi et al., 1999). Specimens were then embedded in methacrylate resin

TABLE 1: Antibodies

Method	Primary antibody	Secondary antibody
Immunofluorescence	AQP4 C19 (sc-9888), anti-goat, 1:300, Santa Cruz Biotech. Inc.	Alexa 594 donkey-anti goat, 1:1,000
		Alexa 488 donkey-anti goat; 1:1,000
	AQP4, anti-rabbit, 1:300, Sigma-Aldrich; Syntrophin-alpha 1 anti-rabbit, 1:1,000, Abcam	Cy3 donkey-anti rabbit, 1:1,000, Jackson Immuno Research Laboratories, Inc; Cy3 donkey-anti rabbit, 1:1,000, Jackson Immuno Research Laboratories, Inc.
	β -Dystroglycan (H-242), anti-rabbit, 1:200, Santa Cruz Biotech. Inc.	Cy3 donkey-anti rabbit, 1:1,000, Jackson Immuno Research Laboratories, Inc.
	Dystrophin, anti-rabbit, 1:40, Abcam	Cy3 donkey-anti rabbit, 1:1000, Jackson Immuno Research Laboratories, Inc.
	GFAP, anti-rabbit, 1:1,000, Dako Cytomation	Alexa 488 donkey-anti; 1:1000
	GFAP, anti-chicken 1:800, Nordic Biosite	Cy2 donkey-anti chicken, 1:1,000, Jackson Immuno Research Laboratories, Inc.
Western blot	AQP4, anti-rabbit, 1:1,500, Chemicon, Millipore	Anti-rabbit IgG Alkaline Phosphatase, 1:10,000; Sigma-Aldrich
	β -Actin (A 2066), anti-Rabbit, 1:500, Sigma-Aldrich	Anti-rabbit IgG Alkaline Phosphatase, 1:10,000; Sigma-Aldrich

(Lowicryl HM20; Polysciences, Warrington, PA) as previously described (Van Lookeren Campagne et al., 1991) and polymerized by exposure to UV light below 0°C.

Ultrathin sections were cut at 70–80 nm using a Leica Reichert Ultracut S Ultratome and a Diatome knife. Blocks were re-orientated then cut perpendicularly to the filter surface, so that both endothelial cells and astrocytes were present on the same section. Sections were collected on single hole grids (Bae et al., 2000) to ensure the integrity of the filter pores. Sections were then contrasted with 2% uranyl acetate for 90 s, rinsed in ddH₂O, exposed to 0.3% lead citrate for 90 s, rinsed in ddH₂O and dried. Sections were examined with a Tecnai electron microscope at 60 kV.

Immunofluorescence

The filters were cut out from the plastic support, fixed for 30 min in 4% FA in 0.01 M PBS and then rinsed in PBS. Co-cultures were then permeabilized by immersion in 0.1% Triton X-100 (Sigma-Aldrich) in PBS for 10 minutes.

After blocking with 2% Bovine Serum Albumin (BSA, Sigma-Aldrich) dissolved in PBS, the co-cultures were incubated overnight with primary antibodies (Table 1). After rinse in PBS, the co-cultures were incubated for 60 min with secondary antibodies (Table 1).

The secondary antibodies were all diluted 1:1,000 in 2% BSA dissolved in PBS. After washing in PBS, the co-cultures were mounted on glass with mounting solution Prolong® Gold antifade reagent with DAPI (Molecular Probes-Invitrogen).

Quantitative Analysis

The images were acquired using a Zeiss LSM 510 confocal microscope, at magnification 40×. Six experiments were conducted for each different combination of co-culture (for example: bEND3 with primary astrocytes or HeLa with primary astrocytes, etc.). Ten to 15 images were acquired from each side of the filter (top aspect for the foot processes and bottom aspect for the astrocyte cell bodies). Images were taken in a blinded manner and GFAP immunofluorescence labelling was used to select the processes and cell bodies. The images were then analysed using LSM image browser with intensity profile mode.

The fluorescence intensity of AQP4 was measured. Separate measurements were made for astrocyte endfeet at the endothelial side (top of the filter) and astrocyte cell bodies (bottom of the filter). For both types of profile, we further quantified, with LSM image browser, the intensity of the AQP4 staining in the cytosol and in the cellular membrane.

For each image six different intensity (I) values were chosen: two in the membrane (membI) and four in the cytosol (cytI).

AQP4 compartmentalization to the plasma membrane was assessed as the ratio between the mean AQP4 immunofluorescence intensity in the plasma membrane and the mean AQP4 fluorescence intensity in the cytosol:

$$R (AQP4 \text{ compartmentalization}) = \frac{[mean \text{ of } AQP4 \text{ membI}]}{[mean \text{ of } AQP4 \text{ cytI}]}$$

The degree of polarization (polarized expression of AQP4) was assessed as the ratio between the mean AQP4 immunofluorescence

intensity in the plasma membrane of the cell bodies (bottom aspect of the filter) and the plasma membrane of the foot processes (top aspect of the filter):

$$R \text{ (AQP4 polarization)} = \frac{[\text{mean of AQP4 endfoot membI}]}{[\text{mean of AQP4 bodies membI}]}$$

In either of the cases, ratios above 1 were considered as a sign of increased membrane compartmentalization or polarization.

We used SPSS to analyse the data. The co-cultures combinations were compared with One Way ANOVA (Multiple Comparison), post hoc test Scheffe and Bonferroni were used with $\pm 2SE$ and $P \leq 0.05$ was considered as significant.

SDS-page and Western Blot on Primary Astrocytes Treated with CM

Every 2 days for 6 days, co-culture CM was collected, mixed 1:1 with FM and applied on astrocytes seeded in a six cell culture multi-well (TRP); as negative control, we used astrocytes grown in FM. At day 7, the astrocytes were harvested by a cell scraper in PBS and spun down in a bench centrifuge.

The astrocyte pellet was lysed in RIPA (Radio-Immunoprecipitation Assay) Buffer (50 mM Tris-HCl pH 7.4; 150 mM NaCl; 5mM EDTA; 1% Triton X-100; 0.5% sodium deoxycholate; 0.1% SDS) and protease inhibitors cocktail tablets purchased from Roche (Complete, Roche), with a mechanical dissociation using a pipette ($3 \times$ every 10 min) and centrifuged at 9,300g for 30 min.

The supernatant was collected and the protein content was measured with a DC protein assay from Bio-Rad. The samples were analyzed by immunoblotting. The samples were diluted in loading buffer (0.875% SDS, 5% Glycerol, 50 mM Tris-HCl pH 6.8, 0.83% β -mercaptoethanol and trace amount of bromophenol blue), denaturated at 37°C for 10 min following centrifugation at 13,000 rpm for 10 min and 2 μ g of protein was added per lane, separated by electrophoresis on 12.5% Tris-HCl 1.0 mm Criterion Precast Gel (Bio-Rad) and transferred onto a PVDF membrane (Criterion gel blotting sandwich, Bio-Rad).

The membrane was blocked for 2 h in Tris-buffered saline (20 mM Tris, 137 mM NaCl) plus 0.1% Tween (TBS-T) plus 5% skim milk powder (Sigma-Aldrich) followed by overnight incubation in primary antibody for AQP4 (Table 1). The membrane was subsequently incubated for 60 min in secondary anti-rabbit antibody (Table 1) diluted 1:10000 in TBS-T and detected with the ECF kit according to the manufacturer's instructions (GE Healthcare). After developing, the membrane was washed and incubated with primary antibody for β -actin (Table 1), followed by the secondary antibody and detected as described above.

Quantitation

The immunofluorescence intensity of the two proteins β -actin and AQP4 was recorded by a Kodak scanner (Kodak Image Station 4000MM PRO) and quantified with a commercially available image analysis program (Kodak MI SE, Carestream Health, Rochester, NY). The ratio between the mean of intensity of AQP4 on β -actin is an indication of AQP4 expression.

Quantitative Real-time PCR

Primary astrocytes were seeded at the concentration of 1×10^7 cell cm^{-2} in 10 cm petri dishes for cell culture. For each group, control and experimental conditioned media (CM), were considered five Petri dishes ($n = 5$). The astrocytes were treated after 24 h either with FM, control, or with CM 1:1 FM not to deprive cells of nutrients. After another 24 h, the astrocytes were collected.

Total RNA was extracted using RNeasy Mini kit according to the manufacturer's instructions (Qiagen). One μ g of RNA was reverse-transcribed using oligo d(T)18 primers and GoScript Reverse transcription System (Promega) in a reaction volume of 20 μ L. The cDNA reaction mixture was diluted five-fold in TE buffer (10 mM Tris-HCl, 1mM EDTA, pH 8.0). Real-time PCR was set up using 2 μ L template, corresponding to 20 ng cDNA, in a reaction mixture of 20 μ L containing AB Power SYBR Green PCR Master Mix (Applied Biosystems) and specific primers (200 nM) for amplification of target genes. Expression levels of target genes were assessed by absolute quantification using StepOnePlus real-time PCR system (Applied Biosystems). Amplicons were generated in a two-step PCR (95°C for 15 s, 60°C for 60 s, 40 cycles). *HPRT1* was used as an endogenous control for normalization of gene expression. The primers used to detect *Aqp4* gene were as follows: 5'-GAATCCAGC TCGATCCTTTG-3' and 5'-CTCCATGTAGCTCCCTTTTCG-3'. The primers used to detect *HPRT1* gene were as follow: 5'-CCC AGCGTCGTGATTAGTGA-3' and 5'-TGGCCTCCCATCTCCTT CAT-3'.

Statistical Analysis Western Blot and qPCR

The obtained values were transferred to SPSS Version 16.0 (SPSS, Chicago, IL) and mean values of the groups were compared using independent *T* test. Data are presented as mean $\pm 2SE$. $P \leq 0.05$ was considered as significant.

Proteomics of the CM

Deglycosylation, Protein Separation, and in Gel Trypsin Digestion. Approximately 25 mL of CM5 was centrifuged at 15,000g for 30 min to remove cell debris. Proteins were precipitated by adding 4 $\times$ cold acetone volume and stored at -20°C overnight. Precipitate was recovered after centrifugation at 15,000g for 20 min, and resuspended in 4 mL of deglycosylation buffer (150 mM NaCl, 50 mM sodium acetate, 10 mM EDTA, pH 6.8) (Didangelos et al., 2010). Protein concentration measured by nanodrop was 2.5 mg mL^{-1} .

For deglycosylation, 250 μ g of sample protein was submitted to reaction against 0.5 U Chondroitinase ABC from *Proteus vulgaris*, 0.5 U Heparitinase II from *Flavobacterium heparinum* and 5U Endo- β -galactosidase from *Bacteroides fragilis* (Sigma-Aldrich) for 16 h at 37°C. Deglycosylated samples were then vacuum concentrated to reduce sample volume, and was mixed 4:1 to NuPAGE SDS Sample buffer (Invitrogen), and submitted in duplicate to electrophoretic separation for 40 min using a 4–12% Bis-Tris NuPAGE minigel in MES buffer. Gel were then fixed and stained with colloidal blue Coomassie kit from Invitrogen, as suggested by manufacturer.

FIGURE 2: Dystrophin associated protein complex in the co-culture model. Confocal microscopic immunofluorescence analysis shows that β -dystroglycan (A), dystrophin (B) and α -syntrophin (C) (red) are similarly distributed and colocalized with AQP4 (green) in endfoot cell membranes (top of filter). GFAP is stained in blue. Membrane compartmentalization of β -dystroglycan and AQP4 is not evident in the astrocyte cell bodies in filter bottom (A, upper panel). Scale bar: 50 μ m. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

Gel lanes from samples were cut in 10 fractions each, and gel pieces were prepared for in gel digestion using trypsin. Shortly, each fraction was washed three times in 50% acetonitrile in water for removal of Coomassie and SDS. This was followed by 1-h reduction using 10 mM DTT at 58°C, and 45 min alkylation using 55 mM iodoacetamide at room temperature and in the dark. Fractions were then washed three times with 50 mM ammonium bicarbonate pH 8.0, dehydrated with 100% acetonitrile and vacuum concentrated to complete dryness. Each gel piece was then rehydrated with 20 μ L of 50 mM ammonium bicarbonate containing 0.1 μ g trypsin (Promega) for 20 min, and then 200 μ L of 50 mM

ammonium bicarbonate was added to cover the pieces and stored at 37°C overnight. Enzyme reaction was stopped with by acidification using 10% v/v trifluoroacetic acid in water. Tryptic peptides were finally recovered from the solution by using C18 resin disks from Empore/3M in a STAGE-TIP setup (Rappsilber et al., 2003).

Mass Spectrometry Analysis and Protein Identification. All experiments were performed on a Dionex Ultimate 3000 nano-LC system (Sunnyvale CA, USA) connected to a quadrupole–Orbitrap (QExactive) mass spectrometer (ThermoElectron, Bremen, Germany)

equipped with a nanoelectrospray ion source (Proxeon/Thermo). For liquid chromatography separation we used an Acclaim PepMap 100 column (C18, 3 μm , 100 $\text{\AA}$) (Dionex, Sunnyvale CA) capillary of

15 cm bed length. The flow rate was $0.3 \mu\text{L min}^{-1}$ for the nano column, and the solvent gradient was 7% B to 35% B in 112 minutes, then 40–80% B in 8 min. Solvent A was aqueous 2% acetonitrile in

0.1% formic acid, whereas solvent B was aqueous 90% acetonitrile in 0.1% formic acid.

The mass spectrometer was operated in the data-dependent mode to automatically switch between MS and MS/MS acquisition. Survey full scan MS spectra (from m/z 300 to 1,750) were acquired in the Orbitrap with resolution $R = 60,000$ at m/z 400, after accumulation to a target of 1,000,000 charges in the quadrupole. The method used allowed sequential isolation of the most intense ions, up to ten, depending on signal intensity, for fragmentation on the HCD cell using high-energy collision dissociation at a target value of 1,000,000 charges or maximum acquisition time of 128 ms (Kelstrup et al., 2012). MS/MS scans were collected at 35,000 resolution at the Orbitrap. Target ions already selected for MS/MS were dynamically excluded for 60 s. General mass spectrometry conditions were: electrospray voltage, 2.4 kV; no sheath and auxiliary gas flow. Ion selection threshold was 500 counts for MS/MS.

All acquired data was processed and analyzed using MaxQuant (version 1.4.0.5) (Cox and Mann, 2008). Protein identification was performed by searching the data against an in house compiled database containing all rodent entries from NCBI. MaxQuant submitted multiple searches on its Andromeda search engine. The search parameters were: Enzyme specificity: trypsin with no proline restriction; Maximum missed cleavages: 2; Carbamidomethyl (C) as fixed modification; *N*-acetyl (Protein), Oxidation (M), pyro-glu (Gln to pyro-Glu) and pyro-glu (Glu to pyro-Glu) as variable modifications; Peptide mass tolerance of ± 20 parts per million for first search and 6 ppm for main search after recalibration; MS/MS mass tolerance of 6 ppm. A search versus reversed sequences was allowed to infer false discovery rates in the results. Peptides were further filtered accordingly to their score by MaxQuant in order to allow a maximum of 1% FDR in the data ($P < 0.01$).

Results

Astrocytes and endothelial cells were segregated to the bottom and top aspects, respectively, of the porous filter (Fig. 1A). GFAP immunocytochemistry revealed that the filter pores served as passageways for astrocytic processes that extended to the top aspect of the filter where they branched out in close apposition to the GFAP negative endothelial cells (Fig. 1B). At the electron microscopic level, astrocytic processes were found

in the filter pore and on either surface of the filter (Fig. 1C). Astrocytic cell bodies were not observed at the top aspect, indicating that the pores were too narrow for these to pass through (Demeuse et al., 2002). Thus the model system reproduces the structural polarization typical of astrocytes *in vivo*.

AQP4 and the DAP Complex Aggregate in Astrocyte Processes on the Endothelial Side of the Filter

The astrocytic processes that extended to the top (endothelial) aspect of the filter expressed the DAP complex, as shown by use of antibodies to three different members of this complex (Fig. 2).

Thus, the two dystrophin associated proteins β -dystroglycan and α -syntrophin, as well as dystrophin itself, were concentrated in the plasma membranes of the astrocytic processes at the filter top (Fig. 2A,C). In contrast, the astrocytic cell bodies at the opposite aspect of the filter showed a rather homogeneous distribution of β -dystroglycan (Fig. 2A) as well as α -syntrophin and dystrophin (not illustrated).

The expression of AQP4 mimicked that of the DAP complex (Fig. 2A–C), consistent with a number of studies that have pointed to a role of the DAP complex in AQP4 anchoring.

The plasma membrane compartmentalization of AQP4 depended on the presence of endothelial cells (Fig. 3). Thus, only when endothelial cells were present did astrocytic processes show a distinct peak of AQP4 immunoreactivity corresponding to the plasma membrane (compare Fig. 3A with Fig. 3B). Membrane compartmentalization was evident only in the top processes. No peak of immunoreactivity was found corresponding to astrocytic membranes at the lower surface of the filter (Fig. 3C), regardless of whether astrocytes were grown alone or in co-culture with endothelial cells.

Quantitative Analysis of AQP4 Compartmentalization and Polarization

In quantitative terms, plasma membrane compartmentalization of AQP4 was defined as the ratio between mean plasma

FIGURE 3: Intensity profiles of AQP4 immunofluorescence labeling in astrocyte endfeet co-cultured with endothelial cells (A), in astrocyte endfeet grown in the absence of endothelial cells (B) and in an astrocyte cell body at the bottom of a filter supporting an endothelial cell-astrocyte co-culture (C). Representative immunofluorescent images with AQP4 (red) and GFAP (green) labeling are shown in the left panels and the intensity profiles (red and green respectively) are in the right panels. An arbitrary red arrow drawn on the profile of interest indicates where intensity has been measured. The two peaks in the intensity profile of AQP4 correspond to the points where the red line crosses the plasma membranes of the astrocyte endfeet in A. There are no peaks corresponding to the plasma membranes in B and C indicating equal distribution of AQP4 immunofluorescent intensity in the plasma membrane and the cytosol. Scale bar: 50 μ m. D and E show AQP4 membrane compartmentalization and polarization. D: quantitative analysis based on the ratio of mean intensity of AQP4 in plasma membrane and in cytosol of astrocytes endfeet and cell bodies in presence of endothelial cells (+bEnd3) or in the absence of endothelial cells (-bEnd3). A ratio greater than 1.0 is indicative of AQP4 compartmentalization in the plasma membrane. E: quantitative analysis based on the ratio between the mean intensity of AQP4 in plasma membrane of the astrocytes endfeet and the mean intensity of AQP4 plasma membrane labeling in the astrocyte cell bodies in the presence (+bEnd3) or in the absence of endothelial cells (-bEnd3). A ratio >1 is indicative of a polarized expression of AQP4. Statistics: ANOVA and independent t test between groups; **significantly different from the control (astrocytes grown on the filter in absence of bEND3); "x" significantly different from previous value. Error bars indicate ± 2 SE, $P \leq 0.05$, $n = 6$. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

FIGURE 4: Orthogonal projection of z-stacks through astrocytic foot processes co-immunostained with antibodies to GFAP (green) and AQP4 (red) in the presence (A) or absence (B) of bEND3 cells. Corresponding 3D reconstruction of the z-stacks is shown in the lower panel of each figure. AQP4 immunostaining is stronger and more compartmentalized to the astrocyte plasma membrane in the presence of the bEND3 cells. The foot process in the absence of the bEND3 is flatter (B; $z = 12 \mu\text{m}$) than in the presence of the bEND3 cells (A, $z = 16 \mu\text{m}$). Scale bar: $10 \mu\text{m}$. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

membrane labelling and mean cytosol labelling (Fig. 3D). This ratio increased more than threefold in the presence of brain endothelial cells. The quantitative analysis confirmed that the effect of endothelial cells was restricted to the astrocytic processes at the filter top (Fig. 3D).

We chose to use the AQP4 immunosignal as a marker for astrocyte polarization. Thus, polarization was defined as the ratio between mean AQP4 signal in membranes of top processes and mean AQP4 signal in membranes of astrocytic cell bodies on the bottom aspect of the filter (Fig. 3E). Defined in this way, astrocyte polarity was contingent on the presence of brain endothelial cells (Fig. 3E).

3D reconstruction of the z-stack through the astrocytic foot processes on the endothelial aspect of the filter showed a

stronger and more compartmentalized AQP4 immunosignal in the plasma membranes, when bEND3 cells were present (Fig. 4A), compared with the condition without bEND3 (Fig. 4B).

Effects of Brain Endothelial Cells Depend on the Presence of α -syntrophin

A priori, endothelial cells could cause AQP4 to accumulate at the membrane through a hypothetical interaction with AQP4 that does not involve the DAP complex. To rule out this possibility, we knocked out α -syntrophin, a dystrophin associated protein that serves as the immediate anchor of AQP4 (Neely et al., 2001). In cells from α -SYN-KO mice the top processes of astrocytes showed no membrane accumulation of AQP4 in

FIGURE 5: **A:** confocal microscopy images showing AQP4 (red) and GFAP (green) immunofluorescence labelling of astrocyte endfeet from wild type (WT) and α -syntrophin null-mice (α -SYN-KO) co-cultured with bEND3. A distinct membrane localization of AQP4 is found in endfeet of WT astrocytes, but is missing in endfeet of α -SYN-KO astrocytes. **B:** quantitative analysis shows that knockout of α -SYN abolishes the effect of bEND3 cells on astrocyte compartmentalization (left panel) and polarization (right panel). Scale bar: 20 μ m. Statistics: Anova; **significantly different from the ctrl (astrocytes grown on the filter in absence of bEND3); "x" significantly different from previous value. Error bars indicate ± 2 SE, $P \leq 0.05$, $n = 6$. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

the presence of endothelial cells (Fig. 5A,B). This is consistent with the idea that endothelial cells dictate AQP4 expression through recruitment of the DAP complex to which AQP4 is normally tethered.

We then asked whether other cell types could substitute for brain endothelial cells. AQP4 membrane compartmentalization was stronger with brain endothelial cells than with HeLa cells or with endothelial cells from the umbilical cord (Fig. 6A,B). The three cell types did not differ significantly in

terms of their effect on astrocyte polarization, as judged by distribution of AQP4 along the astrocyte membrane.

Effects of Extracellular Matrix Molecules

Laminin or Matrigel was applied in the presence or absence of brain endothelia (Fig. 7). Laminin had no effect of its own on AQP4 compartmentalization or polarization, nor did it potentiate the effect of brain endothelial cells (Fig. 7B). In contrast, co-application of Matrigel with endothelial cells gave

FIGURE 6: Quantitative analysis of membrane compartmentalization (A) and polarized expression (B) of AQP4 in primary astrocytes co-cultured with HeLa, ECV and bEND3 cell line. A: the bEND3 cell line induces a significantly higher degree of AQP4 membrane compartmentalization in the astrocyte endfeet than the HeLa or ECV cell lines. B: each cell line induces polarized AQP4 expression in the astrocytes. Statistics: ANOVA; **significantly different from the ctrl (astrocytes co-cultured with ECV); "x" significantly different from previous value. Error bars indicate ± 2 SE, $P \leq 0.05$, $n = 6$.

rise to a more pronounced polarization and membrane compartmentalization than that seen with endothelial cells alone (Fig. 7A,B).

The effect of Matrigel could be reproduced by agrin, a proteoglycan of the extracellular matrix (Fig. 7C).

CM Increases AQP4 Expression and Compartmentalization

In a next series of experiments we tested whether the effects of brain endothelial cells could be mimicked by application of beads soaked in medium harvested from the brain endothelia/astrocyte co-cultures (Fig. 8A). These beads induced a more pronounced membrane compartmentalization of AQP4 than did control beads. However, even control beads showed a significant effect on astrocyte polarization when added to the top of the filter (Fig. 8B). Thus, mechanical interaction per se appears to have an inductive effect.

Medium harvested from co-cultures and added to the standard medium not only caused AQP4 to accumulate in the plasma membrane but also increased the total level of AQP4 and AQP4 mRNA in the cultured cells (Fig. 8C–E).

Mass Spectrometry

The above experiments with beads and conditioned medium indicated that brain endothelial cells secrete components that increase AQP4 expression and accentuate its membrane compartmentalization. Direct applications of Matrigel or agrin onto the co-cultures suggested that the latter molecule could be one of components responsible for the observed effects.

Thus it was deemed of importance to resolve whether agrin is among the compounds present in the CM.

Mass spectrometry showed that the CM contains more than 13,000 unique peptides, corresponding to $\sim 1,900$ protein groups. Agrin figured prominently among the identified proteins, which also included laminin (Fig. 9 and Table 2, and Supp. Info. Table 1).

Discussion

That astrocytes are structurally polarized was recognized already in the classical histological studies of the central nervous system (Ramon and Cajal, 1894). Specifically, it was found that astrocytes extend processes towards microvessels or pial surface and that these processes differed morphologically from the delicate processes that abutted on synapses and neuronal membranes. With the advent of freeze fracture techniques it was demonstrated that endfoot membranes were unique by showing dense aggregates called orthogonal arrays of particles (Rash and Ellisman, 1974; Rash et al., 1974; Wolburg, 1995; Wolburg et al., 2011). This was the first hint that the endfeet serve specific functions and that astrocytes are polarized also in functional terms. Careful analyses of a specialized class of retinal astrocytes (Müller cells) confirmed that astrocytic processes were functionally heterogeneous with subvitreous processes showing much higher K^+ conductance than processes abutting on the retinal neuropil (Newman, 1986a,b; Newman et al., 1984; Orchard and Finlayson, 1977). Immunogold studies unravelled the molecular substrate for this functional heterogeneity and showed that subvitreous endfeet of Müller cells are enriched

FIGURE 7: Effect of different extracellular matrix components on membrane compartmentalization and polarized expression of AQP4. **A:** confocal immunofluorescence image showing AQP4 (red) and GFAP (green) immunostaining at astrocyte endfeet in co-culture in the presence of (+bEnd3; left) or absence of bEND3 cells (-bEnd3) on a filter coated with Matrigel. Scale bar: 50 μ m. **B:** quantitative analysis shows that laminin have no effect on its own on AQP4 compartmentalization or polarization, while Matrigel seems to potentiate the effect of the bEND3 cells on both polarization and membrane compartmentalization. Matrigel also show a modest inductive effect on polarization when astrocytes are grown in absence of bEND3. **C:** agrin has the same potentiating effect as Matrigel on both polarization and membrane compartmentalization. Statistics: ANOVA and independent t test between groups; "x" significantly different from previous value. Error bars indicate ± 2 SE, $P \leq 0.05$, $n = 6$. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

FIGURE 8: Effect of the conditioned medium (CM) on membrane compartmentalization (A) and polarization (B) of AQP4. Endfeet were exposed to beads soaked with CM (BeadsCM) or control beads. **A:** beads soaked with CM induce significantly higher membrane compartmentalization of AQP4 than control beads. **B:** polarization is induced also by control beads, suggesting an inductive effect of mechanical contact. Statistics: ANOVA; **significantly different from control (astrocytes exposed to control beads); "x" significantly different from previous value. Error bars indicate ± 2 SE, $P \leq 0.05$, $n = 6$. **C:** immunoblot showing AQP4 protein expression in primary astrocytes treated with the CM or the control medium (CTRL). **D:** quantitative analysis of the immunoblot shows significantly increased level of AQP4 protein in the astrocytes treated with the CM compared with the control medium (CTRL). Statistics: Independent t-test; **significantly different from control (astrocytes CTRL); Error bars indicate ± 2 SE, $P \leq 0.05$, $n = 6$. **E:** quantitative analysis of AQP4 mRNA copy number on ng of total mRNA of astrocytes treated with the CM or the control medium (CTRL). Treatment with CM leads to a significant increase in the AQP4 mRNA. Statistics: Independent t test; **significantly different from the ctrl (astrocytes CTRL); Error bars indicate ± 2 SE, $P \leq 0.05$, $n = 5$.

in the inwardly rectifying K^+ channel Kir4.1, as well as the water channel AQP4 (Nagelhus et al., 1999). Now we know that the orthogonal arrays identified in early freeze fracture

studies correspond to the tightly packed endfoot pool of AQP4 (Rash et al., 1998; Verbavatz et al., 1997) and that endfeet are characterized by a high transport capacity for water as

FIGURE 9: Typical MS/MS spectrum identified in the dataset. It represents the peptide FDTGSGPAVLTSVLPVEPGR (m/z 1000.026, mass error of 2.37 ppm) of agrin, which was scored by MaxQuant with a value of 59.

well as certain ions and organic solutes (Benfenati et al., 2007, 2011). Endfeet are also critically involved in gliovascular signaling (Nedergaard et al., 2002; Takano et al., 2014).

Recent studies demonstrate that loss of astrocyte polarization is a common denominator of a number of neurological conditions such as mesial temporal lobe epilepsy, Alzheimer's disease and stroke (Alvestad et al., 2013; Frydenlund et al., 2006; Yang et al., 2011). This emphasizes the need to identify those factors that are responsible for the specialization of perivascular endfoot membranes.

Astrocytes in primary culture are not representative of astrocytes in the intact brain. Most importantly, when cultured under standard conditions, astrocytes contain a significant amount of cytoplasmic AQP4, as evidenced by our own experiments and previous studies in other laboratories. Most notably, in a description of immunostained astrocyte primary cultures, Nicchia et al. (2004) found that a large proportion of the AQP4 staining occurred in the cell cytoplasm, indicating that the polarized expression of AQP4 observed *in vivo* is not maintained in culture (Nicchia et al., 2004). This is an appropriate reference to the starting point of our own study: namely that astrocytes in culture show an AQP4 compartmentation very different from that observed *in vivo*.

Another paper by Nicchia et al. (2008) addresses mechanisms by which AQP4 is directed to the membrane and underlines the difference between the *in vivo* and *in vitro* situation when it comes to subcellular AQP4 localization (Nicchia et al., 2008). This is in line with the study of Noell et al. (2007) who described an evenly and diffusely distributed pool of AQP4 in cultured astrocytes (Noell et al. 2007).

Our *in vitro* experimental model was designed so as to reproduce aspects of AQP4 distribution typical of the *in vivo*

situation. Most significantly, the polarized expression of AQP4 is re-established in this model system. However, some AQP4 remains in the cytoplasm. Whether this pool is larger or smaller than that occurring *in vivo* is difficult to judge. Immunogold studies reveal very little cytoplasmic AQP4 *in vivo* (Amiry-Moghaddam and Ottersen, 2013; Nielsen et al., 1997), but the immunofluorescence technique used here is likely to pick up even very small concentrations of AQP4.

We show that processes with structural and biochemical features akin to native endfeet could be generated *in vitro*, by allowing cultured astrocytes to establish contact with endothelial cells at a distance from the astrocytic cell bodies. Technically this was achieved by growing astrocytes and endothelial cells on opposite sides of a Millipore filter with pores small enough to prevent cells from entering, but large enough to permit passage of thin processes. When exposed to endothelial cells, the latter processes formed specialized membrane domains with the hallmarks of endfoot membranes: a dense complement of dystrophin and dystrophin associated proteins as well as a high concentration of the water channel AQP4.

With this model system in hand we could experimentally manipulate the endfeet so as to identify those factors that govern the build-up of their molecular assembly. Guiding our experimental approach was our assumption that the factors in question act locally, as elements of an inductive process that is set in motion when astrocyte processes establish contact with microvessels.

This assumption is a plausible one, given the remarkable precision by which the endfoot specialization coincides with the endothelial basal lamina contact (Amiry-Moghaddam and Ottersen, 2003; Nielsen et al., 1997). Notably, AQP4 is abruptly reduced from the astrocyte endfoot membrane once the latter membrane is reflected away from the endothelial

TABLE 2: Selected Components Detected in the Co-culture Conditioned Medium

Protein ID	Protein name	Gene ID	Peptides	Unique peptides
A2ASQ1	Agrin	Agrn	23	3
P39876	Metalloproteinase inhibitor 3	Timp3	2	3
P04202	Transforming growth factor beta-1	Tgfb1	3	3
O08665	Semaphorin-3A	Sema3a	7	7
O88668	Protein CREG1	Creg1	3	3
P98063	Bone morphogenetic protein 1	Bmp1	6	6
E9Q718	Procollagen-lysine,2-oxoglutarate 5-dioxygenase 2	Plod2	11	11
P10493	Nidogen-1	Nid1	28	28
P42703	Leukemia inhibitory factor receptor	Lifr	4	4
F1MA59	Collagen alpha-1(IV) chain	Col4a1	4	3
P14844	C-C motif chemokine 2	Ccl2	3	2
P35585	AP-1 complex subunit mu-1	Ap1m1	10	10
P21674	Follistatin	Fst	6	6
Q3UVZ1	Interleukin-1 receptor accessory protein	Il1rap	3	3
Q02788	Collagen alpha-2(VI) chain	Col6a2	16	16
P15800	Laminin subunit beta-2	Lamb2	8	8
B2CQD6	Fibulin-1	Fbln1	13	5
B2RQQ8	Collagen alpha-2(IV) chain	Col4a2	8	8
Q80YX1	Tenascin	Tnc	48	48
B1AWB9	Collagen alpha-1(V) chain	Col5a1	15	15
F8WIP8	Osteopontin	Spp1	2	2
B1AUP1	Metalloproteinase inhibitor 1	Timp1	10	10
Q544G5	Dystroglycan	Dag1	8	8
E9QN70	Laminin subunit beta-1	Lamb1	36	36
O35206	Collagen alpha-1(XV) chain	Col15a1	2	2
D3ZYI8	SPARC	Sparc	11	11
Q05895	Thrombospondin-3	Thbs3	5	4
E9PZ16	Basement membrane-specific heparan sulfate proteoglycan core protein	Hspg2	34	34
E9QPX1	Collagen alpha-1(XVIII) chain	Col18a1	6	6
E9QLW5	Insulin-like growth factor II	Igf2	2	2
E0CZ58	Proteoglycan 4	Prg4	4	4
P21744	Insulin-like growth factor-binding protein 4	Igfbp4	8	8
P35441	Thrombospondin-1	Thbs1	39	22
Q3UIP2	Procollagen C-endopeptidase enhancer 1	Pcolce	16	16
P18572	Basigin	Bsg	3	3
F8WH23	Insulin-like growth factor-binding protein 7	Igfbp7	14	6

TABLE 2: Continued

Protein ID	Protein name	Gene ID	Peptides	Unique peptides
O35598	Disintegrin and metalloproteinase domain-containing protein 10	Adam10	1	1
Q62894	Extracellular matrix protein 1	Ecm1	1	8
O08859	Tumor necrosis factor-inducible gene 6 protein	Tnfaip6	4	4
Q2KHP2	Matrix metalloproteinase-19	Mmp19	10	10
Q61001	Laminin subunit alpha-5	Lama5	13	13
P97927	Laminin subunit alpha-4	Lama4	14	14
D4ACU9	Integrin alpha-5	Itga5	4	4
B1AYG5	Integrin beta-6	Itgb6	1	1
G3V991	Integrin alpha-6	Itga6	4	4
G5E8M2	Fibronectin	Fn1	2	101
Q8CJ53	Cdc42-interacting protein 4	Trip10	2	2
Q80X19	Collagen alpha-1(XIV) chain	Col14a1	1	2
F8WHK2	Thrombospondin-2	Thbs2	35	33
P02454	Collagen alpha-1(I) chain	Col1a1	6	6
Q04857	Collagen alpha-1(VI) chain	Col6a1	27	27
A2RTD1	Interleukin-6	Il6	1	1
P49134	Integrin beta-1	Itgb1	5	5
P41245	Matrix metalloproteinase-9	Mmp9	12	12
Q8BPP5	EGF-containing fibulin-like ECM protein 1	Efemp1	14	8
G3V6B1	Transforming growth factor beta-2	Tgfb2	4	4
P28653	Biglycan	Bgn	19	19
Q5DTG2	Collagen alpha-1(III) chain	Col3a1	11	8
P12843	Insulin-like growth factor-binding protein 2	Igfbp2	6	6
Q61554	Fibrillin-1	Fbn1	51	1
Q61508	Extracellular matrix protein 1	Ecm1	24	1
P97298	Pigment epithelium-derived factor	Serpinf1	14	3
P43406	Integrin alpha-V	Itgav	3	3
Q62059	Versican core protein	Vcan	10	10
Q922W6	Stromelysin-1	Mmp3	29	22

basal lamina (Nielsen et al., 1997). Quantitative immunogold analysis of the retinal Müller cells shows that there is a 1:10 ratio of AQP4 immunogold labelling between perisynaptic and perivascular processes (Nagelhus et al., 1999).

Polarization, as here defined, is the extent to which the endfoot and soma membranes differ in terms of their complement of proteins. For quantitative analyses, AQP4 was used as a proxy for astrocyte polarization. Two additional param-

eters, inextricably coupled to astrocyte polarization, are membrane compartmentalization and overall protein expression. Again using AQP4 as a proxy, it is clear that the extent to which this protein is concentrated in the endfoot membrane depends on its recruitment to the plasma membrane from intracellular stores (membrane compartmentalization) as well as on the rate of transcription and translation (determining its overall expression).

The first and salient observation is that endfeet indeed behave as separate compartments. Notably, endothelial cells cause AQP4 to accumulate in endfoot membranes relative to endfoot cytosol while causing no changes to AQP4 compartmentalization in cell bodies.

We hypothesized that the effect of endothelial cells could be mediated by an extracellular matrix component. An obvious candidate is laminin. Laminin is a main constituent of the endothelial basal lamina and is among the first matrix molecules to appear at the endothelial-endfoot interface (Lunde et al., 2014). However, in the present experiments, laminin did not induce any polarization on its own, nor did it enhance the polarizing effect of brain endothelial cells. This was in contrast to Matrigel, which augmented the effect of endothelial cells with respect to polarization as well as compartmentalization.

Matrigel contains a host of extracellular matrix proteins, including proteoglycans.

Our experiments showed that agrin, a proteoglycan, mimicked Matrigel in terms of its effect on AQP4 compartmentalization and polarization. In the case of polarization, Matrigel was more powerful than agrin in potentiating the effect of endothelial cells, and also exerted a modest inductive effect on its own. The observed effect of agrin in our experiments is supported by Noell et al. (2007) who observed that astrocytes cultured in the presence of agrin-conditioned medium, show an increase in the intensity of AQP4-specific membrane-associated staining (Noell et al., 2007).

The effect demonstrated for agrin would be relevant only to the extent that agrin is actually present at the endfoot-endothelium interface. Recent immunogold studies do show that agrin occurs in the perivascular basal lamina from very early stages in development (Lunde et al., 2014). Mass spectrometry was carried out to resolve whether agrin also occurs in the medium to which the endfeet are exposed. The conditioned medium—harvested from the co-cultures—contained several agrin fragments. As expected, this medium also contained thousands of other peptides, including several growth factors (Moore et al. 2009).

Agirin is endowed with several properties that support its candidacy as an inductive factor. It has a binding site for α -dystroglycan and may thus serve to anchor the entire dystrophin associated protein complex in endfoot membranes (Gee et al., 1994; Peng et al., 1998). In addition, agrin is known to act as a cofactor for growth factors (Parthy and Nikolics, 1993)—several of which are found in the co-culture medium (Parthy and Nikolics, 1993). Growth factors are also constituents of Matrigel, possibly explaining why Matrigel potentiates the inductive effect of endothelial cells more effectively than does agrin alone.

The above experiments should not be taken to suggest that agrin is the only factor underpinning astrocytic polarization. Indeed, agrin knockout animals display a residual pool of endfoot AQP4, indicating that other factors also contribute (Fallier-Becker et al., 2011). It is relevant in this regard that astrocyte processes reaching the top surface of the filter are exposed not only to the soluble factors that the endothelial cells secrete, but also to the mechanical stress they impose. Thus we asked whether mechanical interaction per se could favor membrane compartmentalization and polarization of AQP4. In other systems, stretch and membrane displacement are known to effect changes in protein expression (Dehouck et al., 1990; Meyer et al., 1991; Nicchia et al., 2004; Wolburg et al., 1994).

If mechanical factors play a role, one would expect the effects of brain endothelial cells to be reproduced, at least in part, by other cell types. While other cell types failed to mimic brain endothelial cells in terms of their effect on compartmentalization, they did cause a significant polarization. Application of beads rather than cells to the top of the filter similarly induced polarization. In sum, these experiments point to the involvement of mechanical factors. Beads alone were as efficient as beads soaked in conditioned medium when it comes to their ability to induce polarization. Thus, the effect of the mechanical factors is strong enough to obscure any polarizing effect of the conditioned medium. In contrast, the effect of conditioned medium is revealed when recordings are made of the impact of beads on membrane compartmentalization.

In the intact brain, small molecules released from endothelial cells are likely to reach large expanses of the astrocyte plasma membrane. Mimicking this situation, we exposed cultured astrocytes to the conditioned medium and found that this increased significantly the expression of the AQP4 gene, at the protein as well as the transcript levels.

Conclusion

The design of an *in vitro* model that imposes on astrocytes a structural polarity mirroring that seen *in vivo* has made it possible to identify factors that govern the buildup of the endfoot-basal lamina junctional complex responsible for the functional specialization of perivascular endfoot membranes. Using AQP4 as a proxy for this complex, we conclude that endothelial cells exert their effects through agrin and other soluble factors, as well as through a direct mechanical interaction with the endfoot processes. Through the compounds they secrete, the endothelial cells also increase AQP4 expression. The present data suggest that the highly specialized gliovascular interface is established through a series of inductive processes akin to those that underlie the polarization of central neurons.

Acknowledgment

Grant sponsor: EMBO, Heidelberg, Germany.

The authors are grateful to Ms. Bjørg Riber, Karen Marie Gujord, and Jorunn Knutsen for the excellent technical assistance and Carina Knudsen and Gunnar F. Lothe for the artwork. The ECV 304 cells, used in this study, were a kind gift from Professor Stefano Ferroni.

References

- Adams ME, Kramarcy N, Krall SP, Rossi SG, Rotundo RL, Sealock R, Froehner SC. 2000. Absence of alpha-syntrophin leads to structurally aberrant neuromuscular synapses deficient in utrophin. *J Cell Biol* 150:1385–1398.
- Alvestad S, Hammer J, Hoddevik EH, Skare O, Sonnewald U, Amiry-Moghaddam M, Ottersen OP. 2013. Mislocalization of AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy. *Epilepsy Res* 105:30–41.
- Amiry-Moghaddam M, Otsuka T, Hurn PD, Traystman RJ, Haug FM, Froehner SC, Adams ME, Neely JD, Agre P, Ottersen OP. 2003. An alpha-syntrophin-dependent pool of AQP4 in astroglial end-feet confers bidirectional water flow between blood and brain. *Proc Natl Acad Sci USA* 100:2106–2111.
- Amiry-Moghaddam M, Ottersen OP. 2003. The molecular basis of water transport in the brain. *Nat Rev Neurosci* 4:991–1001.
- Amiry-Moghaddam M, Ottersen OP. 2013. Immunogold cytochemistry in neuroscience. *Nat Neurosci* 16:798–804.
- Amiry-Moghaddam M, Xue R, Haug FM, Neely JD, Bhardwaj A, Agre P, Adams ME, Froehner SC, Mori S, Ottersen OP. 2004. Alpha-syntrophin deletion removes the perivascular but not endothelial pool of aquaporin-4 at the blood-brain barrier and delays the development of brain edema in an experimental model of acute hyponatremia. *FASEB J* 18:542–544.
- Bae YC, Ihn HJ, Park MJ, Ottersen OP, Moritani M, Yoshida A, Shigenaga Y. 2000. Identification of signal substances in synapses made between primary afferents and their associated axon terminals in the rat trigeminal sensory nuclei. *J Comp Neurol* 418:299–309.
- Barar J, Gumbleton M, Asadi M, Omid Y. 2010. Barrier functionality and transport machineries of human ECV304 cells. *Med Sci Monit* 16:BR52–BR60.
- Barres BA. 2008. The mystery and magic of glia: A perspective on their roles in health and disease. *Neuron* 60:430–440.
- Benfenati V, Amiry-Moghaddam M, Capriani M, Mylonakou MN, Rapisarda C, Ottersen OP, Ferroni S. 2007. Expression and functional characterization of transient receptor potential vanilloid-related channel 4 (TRPV4) in rat cortical astrocytes. *Neuroscience* 148:876–892.
- Benfenati V, Capriani M, Dovizio M, Mylonakou MN, Ferroni S, Ottersen OP, Amiry-Moghaddam M. 2011. An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex is essential for cell-volume control in astrocytes. *Proc Natl Acad Sci USA* 108:2563–2568.
- Bushong EA, Martone ME, Ellisman MH. 2004. Maturation of astrocyte morphology and the establishment of astrocyte domains during postnatal hippocampal development. *Int J Dev Neurosci* 22:73–86.
- Chaudhry FA, Lehre KP, van Lookeren Campagne M, Ottersen OP, Danbolt NC, Storm-Mathisen J. 1995. Glutamate transporters in glial plasma membranes: Highly differentiated localizations revealed by quantitative ultrastructural immunocytochemistry. *Neuron* 15:711–720.
- Cox J, Mann M. 2008. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol* 26:1367–1372.
- Dehouck MP, Meresse S, Delorme P, Fruchart JC, Cecchelli R. 1990. An easier, reproducible, and mass-production method to study the blood-brain barrier in vitro. *J Neurochem* 54:1798–1801.
- Demeuse P, Kerkhofs A, Struys-Ponsar C, Knoops B, Remacle C, van den Bosch de Aguilar P. 2002. Compartmentalized coculture of rat brain endothelial cells and astrocytes: A syngenic model to study the blood-brain barrier. *J Neurosci Methods* 121:21–31.
- Didangelos A, Yin X, Mandal K, Baumert M, Jahangiri M, Mayr M. 2010. Proteomics characterization of extracellular space components in the human aorta. *Mol Cell Proteom* 9:2048–2062.
- Eid T, Lee TS, Thomas MJ, Amiry-Moghaddam M, Bjornsen LP, Spencer DD, Agre P, Ottersen OP, de Lanerolle NC. 2005. Loss of perivascular aquaporin 4 may underlie deficient water and K⁺ homeostasis in the human epileptogenic hippocampus. *Proc Natl Acad Sci USA* 102:1193–1198.
- Etienne-Manneville S. 2008. Polarity proteins in glial cell functions. *Curr Opin Neurobiol* 18:488–494.
- Fallier-Becker P, Sperveslage J, Wolburg H, Noell S. 2011. The impact of agrin on the formation of orthogonal arrays of particles in cultured astrocytes from wild-type and agrin-null mice. *Brain Res* 1367:2–12.
- Frigeri A, Nicchia GP, Nico B, Quondamatteo F, Herken R, Roncali L, Svelto M. 2001. Aquaporin-4 deficiency in skeletal muscle and brain of dystrophic mdx mice. *FASEB J* 15:90–98.
- Frydenlund DS, Bhardwaj A, Otsuka T, Mylonakou MN, Yasumura T, Davidson KG, Zeynalov E, Skare O, Laake P, Haug FM, et al. 2006. Temporary loss of perivascular aquaporin-4 in neocortex after transient middle cerebral artery occlusion in mice. *Proc Natl Acad Sci USA* 103:13532–13536.
- Gee SH, Montanaro F, Lindenbaum MH, Carbonetto S. 1994. Dystroglycan-alpha, a dystrophin-associated glycoprotein, is a functional agrin receptor. *Cell* 77:675–686.
- Iadecola C, Nedergaard M. 2007. Glial regulation of the cerebral microvasculature. *Nat Neurosci* 10:1369–1376.
- Kelstrup CD, Young C, Lavalley R, Nielsen ML, Olsen JV. 2012. Optimized fast and sensitive acquisition methods for shotgun proteomics on a quadrupole orbitrap mass spectrometer. *J Proteome Res* 11:3487–3497.
- Lunde LK, Camassa LM, Hoddevik EH, Khan FH, Ottersen OP, Boldt HB, Amiry-Moghaddam M. 2014. Postnatal development of the molecular complex underlying astrocyte polarization. *Brain Struct Funct* [Epub ahead of print].
- Manley GT, Binder DK, Papadopoulos MC, Verkman AS. 2004. New insights into water transport and edema in the central nervous system from phenotype analysis of aquaporin-4 null mice. *Neuroscience* 129:983–991.
- McCarthy KD, de Vellis J. 1980. Preparation of separate astroglial and oligodendroglial cell cultures from rat cerebral tissue. *J Cell Biol* 85:890–902.
- Meyer J, Rauh J, Galla HJ. 1991. The susceptibility of cerebral endothelial cells to astroglial induction of blood-brain barrier enzymes depends on their proliferative state. *J Neurochem* 57:1971–1977.
- Moore NH, Costa LG, Shaffer SA, Goodlett DR, Guizzetti M. 2009. Shotgun proteomics implicates extracellular matrix proteins and protease systems in neuronal development induced by astrocyte cholinergic stimulation. *J Neurochem* 108:891–908.
- Nagelhus EA, Horio Y, Inanobe A, Fujita A, Haug FM, Nielsen S, Kurachi Y, Ottersen OP. 1999. Immunogold evidence suggests that coupling of K⁺ siphoning and water transport in rat retinal Muller cells is mediated by a coenrichment of Kir4.1 and AQP4 in specific membrane domains. *Glia* 26:47–54.
- Nagelhus EA, Ottersen OP. 2013. Physiological roles of aquaporin-4 in brain. *Physiol Rev* 93:1543–1562.
- Nedergaard M, Ransom B, Goldman SA. 2003. New roles for astrocytes: Redefining the functional architecture of the brain. *Trends Neurosci* 26:523–530.
- Nedergaard M, Takano T, Hansen AJ. 2002. Beyond the role of glutamate as a neurotransmitter. *Nat Rev Neurosci* 3:748–755.
- Neely JD, Amiry-Moghaddam M, Ottersen OP, Froehner SC, Agre P, Adams ME. 2001. Syntrophin-dependent expression and localization of Aquaporin-4 water channel protein. *Proc Natl Acad Sci USA* 98:14108–14113.

- Newman EA. 1986a. High potassium conductance in astrocyte endfeet. *Science* 233:453–454.
- Newman EA. 1986b. Regional specialization of the membrane of retinal glial cells and its importance to K⁺ spatial buffering. *Ann N Y Acad Sci* 481:273–286.
- Newman EA, Frambach DA, Odette LL. 1984. Control of extracellular potassium levels by retinal glial cell K⁺ siphoning. *Science* 225:1174–1175.
- Nicchia GP, Nico B, Camassa LM, Mola MG, Loh N, Dermietzel R, Spray DC, Svelto M, Frigeri A. 2004. The role of aquaporin-4 in the blood-brain barrier development and integrity: Studies in animal and cell culture models. *Neuroscience* 129:935–945.
- Nicchia GP, Rossi A, Mola MG, Procino G, Frigeri A, Svelto M. 2008. Actin cytoskeleton remodeling governs aquaporin-4 localization in astrocytes. *Glia* 56:1755–1766.
- Nielsen S, Nagelhus EA, Amiry-Moghaddam M, Bourque C, Agre P, Ottersen OP. 1997. Specialized membrane domains for water transport in glial cells: High-resolution immunogold cytochemistry of aquaporin-4 in rat brain. *J Neurosci* 17:171–180.
- Noell S, Fallier-Becker P, Beyer C, Kroger S, Mack AF, Wolburg H. 2007. Effects of agrin on the expression and distribution of the water channel protein aquaporin-4 and volume regulation in cultured astrocytes. *Eur J Neurosci* 26:2109–2118.
- Omid Y, Campbell L, Barar J, Connell D, Akhtar S, Gumbleton M. 2003. Evaluation of the immortalised mouse brain capillary endothelial cell line, b.End3, as an in vitro blood-brain barrier model for drug uptake and transport studies. *Brain Res* 990:95–112.
- Orchard I, Finlayson LH. 1977. Electrically excitable neurosecretory cell bodies in the periphery of the stick insect, *Carausius morosus*. *Experientia* 33:226–228.
- Patthy L, Nikolics K. 1993. Functions of agrin and agrin-related proteins. *Trends Neurosci* 16:76–81.
- Pekny M, Wilhelmsson U, Bogestal YR, Pekna M. 2007. The role of astrocytes and complement system in neural plasticity. *Int Rev Neurobiol* 82:95–111.
- Peng HB, Ali AA, Daggett DF, Rauvala H, Hassell JR, Smalheiser NR. 1998. The relationship between perlecan and dystroglycan and its implication in the formation of the neuromuscular junction. *Cell Adhes Commun* 5:475–489.
- Ramon Y, Cajal S. 1894. Les Novells Ideas sur la structure du systeme nerveux chez l'homme et les vertebres. Paris: Reinwald.
- Rappsilber J, Ishihama Y, Mann M. 2003. Stop and go extraction tips for matrix-assisted laser desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in proteomics. *Anal Chem* 75:663–670.
- Rash JE, Ellisman MH. 1974. Studies of excitable membranes. I. Macromolecular specializations of the neuromuscular junction and the nonjunctional sarcolemma. *J Cell Biol* 63:567–586.
- Rash JE, Staehelin LA, Ellisman MH. 1974. Rectangular arrays of particles on freeze-cleaved plasma membranes are not gap junctions. *Exp Cell Res* 86:187–190.
- Rash JE, Yasumura T, Hudson CS, Agre P, Nielsen S. 1998. Direct immunogold labeling of aquaporin-4 in square arrays of astrocyte and ependymocyte plasma membranes in rat brain and spinal cord. *Proc Natl Acad Sci USA* 95:11981–11986.
- Scherer WF, Syverton JT, Gey GO. 1953. Studies on the propagation in vitro of poliomyelitis viruses. IV. Viral multiplication in a stable strain of human malignant epithelial cells (strain HeLa) derived from an epidermoid carcinoma of the cervix. *J Exp Med* 97:695–710.
- Schwartz JP, Wilson DJ. 1992. Preparation and characterization of type 1 astrocytes cultured from adult rat cortex, cerebellum, and striatum. *Glia* 5:75–80.
- Seifert G, Schilling K, Steinhauser C. 2006. Astrocyte dysfunction in neurological disorders: A molecular perspective. *Nat Rev Neurosci* 7:194–206.
- Strohschein S, Huttman K, Gabriel S, Binder DK, Heinemann U, Steinhauser C. 2011. Impact of aquaporin-4 channels on K⁺ buffering and gap junction coupling in the hippocampus. *Glia* 59:973–980.
- Takano T, He W, Han X, Wang F, Xu Q, Wang X, Oberheim Bush NA, Cruz N, Dienel GA, Nedergaard M. 2014. Rapid manifestation of reactive astrogliosis in acute hippocampal brain slices. *Glia* 62:78–95.
- Takumi Y, Ramirez-Leon V, Laake P, Rinivik E, Ottersen OP. 1999. Different modes of expression of AMPA and NMDA receptors in hippocampal synapses. *Nat Neurosci* 2:618–624.
- Van Lookeren Campagne M, Oestreicher AB, Buma P, Verkleij AJ, Gispén WH. 1991. Ultrastructural localization of adrenocorticotrophic hormone and the phosphoprotein B-50/growth-associated protein 43 in freeze-substituted, Lowicryl HM20-embedded mesencephalic central gray substance of the rat. *Neuroscience* 42:517–529.
- Verbavatz JM, Ma T, Gobin R, Verkman AS. 1997. Absence of orthogonal arrays in kidney, brain and muscle from transgenic knockout mice lacking water channel aquaporin-4. *J Cell Sci* 110 (Part 22):2855–2860.
- Wilhelmsson U, Bushong EA, Price DL, Smarr BL, Phung V, Terada M, Ellisman MH, Pekny M. 2006. Redefining the concept of reactive astrocytes as cells that remain within their unique domains upon reaction to injury. *Proc Natl Acad Sci USA* 103:17513–17518.
- Wolburg H. 1995. Orthogonal arrays of intramembranous particles: A review with special reference to astrocytes. *J Hirnforsch* 36:239–258.
- Wolburg H, Neuhaus J, Kniessel U, Krauss B, Schmid EM, Ocalan M, Farrell C, Risau W. 1994. Modulation of tight junction structure in blood-brain barrier endothelial cells. Effects of tissue culture, second messengers and cocultured astrocytes. *J Cell Sci* 107 (Part 5):1347–1357.
- Wolburg H, Noell S, Wolburg-Buchholz K, Mack A, Fallier-Becker P. 2009. Agrin, aquaporin-4, and astrocyte polarity as an important feature of the blood-brain barrier. *Neuroscientist* 15:180–193.
- Wolburg H, Wolburg-Buchholz K, Fallier-Becker P, Noell S, Mack AF. 2011. Structure and functions of aquaporin-4-based orthogonal arrays of particles. *Int Rev Cell Mol Biol* 287:1–41.
- Yang J, Lunde LK, Nuntagij P, Oguchi T, Camassa LM, Nilsson LN, Lannfelt L, Xu Y, Amiry-Moghaddam M, Ottersen OP, et al. 2011. Loss of astrocyte polarization in the tg-ArcSwe mouse model of Alzheimer's disease. *J Alzheimers Dis* 27:711–722.

Postnatal development of the molecular complex underlying astrocyte polarization

Lisa K. Lunde · Laura M. A. Camassa · Eystein H. Hoddevik ·
Faraz H. Khan · Ole Petter Ottersen · Henning B. Boldt ·
Mahmood Amiry-Moghaddam

Received: 5 November 2013 / Accepted: 4 April 2014 / Published online: 29 April 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract Astrocytes are highly polarised cells with processes that ensheath microvessels, cover the brain surface, and abut synapses. The endfoot membrane domains facing microvessels and pia are enriched with aquaporin-4 water channels (AQP4) and other members of the dystrophin associated protein complex (DAPC). Several lines of evidence show that loss of astrocyte polarization, defined by the loss of proteins that are normally enriched in astrocyte endfeet, is a common denominator of several neurological diseases such as mesial temporal lobe epilepsy, Alzheimer's disease, and stroke. Little is known about the mechanisms responsible for inducing astrocyte polarization *in vivo*. Here we introduce the term endfoot-basal lamina junctional complex (EBJC) to denote the proteins that consolidate and characterize the gliovascular interface. The present study was initiated in order to resolve the developmental profile of the EBJC in mouse brain. We show that the EBJC is established after the first week postnatally. Through a combination of methodological approaches, including light microscopic and high resolution immunogold cytochemistry, quantitative RT-PCR, and Western blotting, we demonstrate that the different members of this complex exhibit distinct ontogenic profiles—with the extracellular matrix (ECM) proteins laminin and agrin appearing earlier than the other members of the complex. Specifically, while laminin

and agrin expression peak at P7, quantitative immunoblot analyses indicate that AQP4, α -syntrophin, and the inwardly rectifying K⁺ channel Kir4.1 expression increases towards adulthood. Our findings are consistent with ECM having an instructive role in establishing astrocyte polarization in postnatal development and emphasize the need to explore the involvement of ECM in neurological disease.

Keywords Astrocyte polarization · Aquaporin-4 · Extracellular matrix · Dystrophin associated protein complex · Brain development · Immunogold cytochemistry

Introduction

A hallmark of astrocytes is their structural polarization. Most if not all astrocytes are equipped with endfeet that contact pia or brain microvessels, and with fine velate processes that abut on central synapses. These morphological characteristics were identified over 100 years ago, soon after astrocytes were first discovered (Somjen 1988). Only over the past few decades it was realized that the structural polarization of astrocytes reflects a distinct functional and biochemical compartmentalization. Notably, endfeet were found to exhibit much higher K⁺ conductance than other parts of the membrane, underpinning the concept of K⁺ siphoning (Newman et al. 1984). The fine perisynaptic astrocyte processes, on the other hand, were shown to harbor a distinct molecular profile mirroring their specific roles in transmitter uptake (Chaudhry et al. 1995).

The astroglial endfeet provide a near complete covering of cortical capillaries (Mathiisen et al. 2010) and contain a unique arrangement of proteins that sets them apart from other astroglial compartments. High resolution immunogold analyses (Amiry-Moghaddam and Ottersen 2013)

Electronic supplementary material The online version of this article (doi:10.1007/s00429-014-0775-z) contains supplementary material, which is available to authorized users.

L. K. Lunde · L. M. A. Camassa · E. H. Hoddevik ·
F. H. Khan · O. P. Ottersen · H. B. Boldt ·
M. Amiry-Moghaddam (✉)
Laboratory of Molecular Neuroscience, Institute of Basic
Medical Sciences, University of Oslo, Oslo, Norway
e-mail: mahmo@medisin.uio.no

indicate that endfeet membranes are enriched with a number of transporters and channels, including the inwardly rectifying K⁺ channel, Kir4.1 (Nagelhus et al. 1999); AQP4 (Nielsen et al. 1997), and the transient receptor potential vanilloid 4, TRPV4 (Benfenati et al. 2007). Kir4.1 is the main molecular correlate of the high K⁺ conductance in endfeet, while AQP4 confers a high water permeability on endfeet membranes. The endfoot pool of AQP4 represents an influx route of water in brain edema (Amiry-Moghaddam et al. 2003), but also has been implicated in physiological functions including water resorption postnatally (Haj-Yasein et al. 2011). In addition, recent data suggest that perivascular AQP4 might propel a paracapillary drainage route that helps clear the brain of β -amyloid and other molecules (Iliff et al. 2012). Two of the endfoot proteins (AQP4 and TRPV4) appear to form a molecular complex involved in osmosensing and regulatory volume control (Benfenati et al. 2011).

Loss of astrocyte polarization—defined as a loss of the proteins that normally are enriched in endfoot membranes—occurs in a number of pathophysiological and clinical conditions. Both AQP4 and Kir4.1 are reduced in CA1 endfoot membranes of patients with mesial temporal lobe epilepsy (MTLE) (Eid et al. 2005; Heuser et al. 2012). In an animal model of MTLE, loss of AQP4 was found to precede the development of chronic seizures (Alvestad et al. 2013). Further, the endfoot pool of AQP4 is strongly reduced after experimental stroke (Frydenlund et al. 2006) and in a mouse model of Alzheimer's disease (Yang et al. 2011). Loss of astrocyte polarization will interfere with important homeostatic processes and clearance mechanisms and could thus contribute to the progress and symptomatology of diverse neurological conditions.

Given the importance of astroglial polarization in health and disease, a key question is how polarization is induced and maintained. Several lines of evidence indicate that dystrophin isoform DP71 and its associated proteins play a special role in this regard. The first data pointing to the involvement of a dystrophin associated protein complex (DAPC) came with studies of mice deficient in dystrophin (Frigeri et al. 2001; Vajda et al. 2002) or α -syntrophin (Neely et al. 2001).

α -Syntrophin is known to bind to DP71 as part of the DAPC. Mice deficient in dystrophin or α -syntrophin concurred in showing reduced expression of AQP4 in astrocytic endfeet (Frigeri et al. 2001; Neely et al. 2001; Amiry-Moghaddam et al. 2003). The dramatic loss of AQP4 observed after knockout of α -syntrophin suggested that the latter molecule could be the immediate anchor of AQP4. Biochemical analyses and co-immunoprecipitation experiments bolstered the idea that DAPC is expressed in endfoot membranes and essential for maintaining astrocyte polarization (Neely et al. 2001). It

was proposed that the DAPC is kept in place primarily through its attachment to laminin and agrin, which are key constituents of the pericapillary basal lamina (Neely et al. 2001; Amiry-Moghaddam and Ottersen 2003; Guadagno and Moukhles 2004; Wolburg et al. 2011). This would explain why the level of AQP4 drops abruptly once the endfoot plasma membrane is reflected away from the basal lamina of brain microvessels (Nagelhus et al. 1999).

The present study was initiated in order to resolve the developmental profile of the DAPC and its binding partners—hereafter collectively termed the endfoot-basal lamina junctional complex (EBJC). Through a combination of methodological approaches, including high resolution immunogold cytochemistry (Amiry-Moghaddam and Ottersen 2013), we could show that the different members of this complex exhibit distinct ontogenetic profiles. The earliest proteins to appear were laminin and agrin, and these proteins reached peak levels as early as P7. In contrast, aquaporin-4 and the inwardly rectifying K⁺ channel Kir4.1 were weakly expressed at P7 and showed a sharp increase in expression towards adulthood. The present data provide new insight in the sequence of events that underlies the assembly of the gliovascular interface.

Materials and methods

Animals

Adult male and pregnant female C57BL/6 mice (Jackson Laboratories, Boulder, CO) were used in this study. The mice were allowed ad libitum access to food and drinking water. Animal experiments were performed according to the European Council law on protection of laboratory animals, with the approval of the University of Oslo's Animal Care and Use Committee. Every effort was made to minimize the number of animals.

Perfusion fixation

Animals from the postnatal groups (P0, P4, P7, P10, P13, P21) as well as adult (8 weeks) animals were anaesthetized with Equitisin (150 μ l for adult mice) and transcardially perfused using 4 % formaldehyde (FA) and 0.1 % glutaraldehyde in 0.1 M phosphate buffer (PB) for at least 10 min, with an initial 20 s of ice cold 2 % dextran in 0.1 M PB. For immunofluorescence and peroxidase based immunocytochemistry, glutaraldehyde was omitted from the fixation solution. Brains were removed and postfixed in the fixation solution overnight and stored in a 1:10 dilution of the same solution in 0.1 M PB.

Post-embedding immunogold electron microscopy

Brains were cut into 0.5–1.0 mm slices, different regions were dissected, cryoprotected, quick-frozen in liquid propane (-170°C), and subjected to freeze substitution. Specimens were embedded in methacrylate resin (Lowicryl HM20) and polymerized by UV light below 0°C . Ultrathin sections (70–100 nm) were cut using an ultratome (Reichert Ultracut S, Leica).

Immunogold labeling was carried as previously described (Yang et al. 2011; Promeneur et al. 2013). Briefly, The sections were rinsed in Tris-buffered saline with Triton X-100 (TBST, 5 mM Tris-HCl, 0.3 % NaCl, 0.1 % Triton X-100), incubated in 2 % human serum albumin, followed by primary antibody (AQP4) overnight, secondary antibody (15 nm gold) for 90 min, and contrasted with 2 % uranyl acetate for 90 s and 0.3 % lead citrate for 90 s. The sections were examined using a Tecnai 12 electron microscope at 60 kV.

Immunogold quantitation

Quantitative analysis was performed as described elsewhere (Amiry-Moghaddam et al. 2004; Mathiisen et al. 2006). Briefly 19–20 digital images of capillaries or pial surface were acquired from each section. Linear densities of gold particles over astrocyte membranes were determined by an extension of analysis [Soft Imaging Systems (SIS), Münster, Germany]. Linear densities were determined semi-automatically and transferred to SPSS Version 16 (SPSS, Chicago, IL) for statistical analysis.

Light microscopic immunocytochemistry

Immunofluorescence

The brains were cryoprotected using increasing levels of sucrose dissolved in 0.1 M PB (10–20–30 %), and snap frozen in optimal cutting temperature (OCT) compound on dry ice. Sagittal brain sections (14 μm) were collected on object glasses and stored at -80°C .

For immunofluorescence the sections were rinsed with 0.01 M phosphate buffered saline (PBS) and permeabilized for 10 min with 0.1 % Triton X-100 in PBS. After blocking with 2 % bovine serum albumin (BSA) in PBS, the brain sections were incubated overnight with primary antibodies diluted in blocking solution. After washing in PBS, the brain sections were incubated with secondary antibodies diluted in blocking solution for 60 min. After a new wash in PBS, the brain sections were mounted using Prolong[®] Gold antifade reagent with DAPI (Invitrogen, Molecular Probes). Confocal images were acquired using a Leica fluorescent microscope.

Immunoperoxidase

Cryoprotection was performed as described above. Cryoprotected brains were snap frozen in 30 % sucrose and free-floating sections (30 μm) were rinsed in PBS, exposed to 2 % H_2O_2 for 10 min to deplete endogenous peroxidase activity, followed by three rinses in PBS. All sections were left in double distilled H_2O at 37°C for 10 min prior to pepsin (Dako Ref S3002, 1 mg/mL 0.2 M HCl) exposure at 37°C for 10 min, followed by three rinses in PBS, first of which for 15 min at 27°C , thereafter at room temperature. Blocking was done using 2 % BSA in PBS-T (PBS, 0.1 % Triton X-100). The sections were then incubated in primary antibody diluted in blocking buffer overnight in cold room (4°C), followed by three times rinse in PBS-T and 1 h incubation in secondary antibody in blocking buffer was followed by three times rinses in PBS-T. The sections were then incubated for 1 h with biotinylated-streptavidin horseradish peroxidase complex (GE healthcare UK limited, Rpn 1051 V) diluted 1:100 in PBS-T and rinsed in PBS-T $\times 3$, followed by PBS $\times 3$ and 0.5 mg/mL diaminobenzidine (3,3'-Diaminobenzidine Tetrahydrochloride 10 mg/tab, Sigma D5905, DAB) dissolved in 0.1 M PB for 5 min. Finally, sections were incubated for a titrated time to DAB solution with added 0.01 % H_2O_2 (1 min for Laminin and 1.5 min for agrin). The DAB/HRP reaction was stopped by 3 thorough rinses in 0.1 M PB. Thereafter, sections were mounted using glycerin gelatine and examined using a light microscope.

SDS-PAGE and Western blotting

The mice ($n = 3$ for each age group) were anesthetized (Chloroform), decapitated and the brains were immediately frozen in liquid nitrogen and stored at -80°C . Frozen mouse brains were homogenized in homogenization buffer (50 mM HEPES, 0.32 M Sucrose, 5 mM EDTA) using a mini-pestle (2 $\times$ 1 min). Immediately before homogenization, one tablet of [Protease Inhibitor Cocktail (PIC), Roche] was dissolved in 50 mL of the homogenization buffer. The supernatant of a $1,000 \times g$ spin was used for immunoblotting. Protein concentrations were determined with a DC protein assay (Bio-Rad) according to the manufacturer's instructions. SDS-PAGE and Western blots were carried out as described previously (Yang et al. 2011) with the following exceptions: 5–30 μg of protein (agrin: 2 μg , α -syntrophin: 15 μg , AQP4 (P0–P21): 15 μg , AQP4 (adult): 1 μg , β -dystroglycan: 5 μg , dystrophin: 10 μg , Kir4.1: 5 μg and laminin: 30 μg) was added per lane and separated by electrophoresis on 7.5, 10, 12.5 or 4–20 % polyacrylamide gels (Criterion Precast Gel, Bio-Rad) depending on the proteins examined.

Table 1 Antibodies

Method	Primary antibody	Secondary antibody
Immunogold	AQP4, anti-rabbit, 1:500, Sigma-Aldrich	Goat anti rabbit 15 nm, 1:40, Amersham Bioscience
	Agrin, anti-rabbit, 1:500, Gift from Professor Markus A. Ruegg, University of Basel	Goat anti rabbit 15 nm, 1:40, Abcam
	Laminin (L9393), anti-rabbit, 1:100, Sigma-Aldrich	Goat anti rabbit 15 nm, 1:40, Abcam
Immunofluorescence	Agrin, anti-mouse, 1:100, Chemicon, Millipore, Med Probe	Alexa 488 donkey anti-mouse, 1:1,000, Molecular Probes
	α -Syntrophin (SYN259), anti-rabbit, 1:200, Gift from Dr. Marvin E. Adams	Cy3 donkey-anti rabbit, 1:1,000, Jackson Immuno Research Laboratories, Inc.
	AQP4, anti-rabbit, 1:200, Chemicon, Millipore	Cy3 donkey-anti rabbit, 1:1,000, Jackson Immuno Research Laboratories, Inc.
	AQP4, anti-rabbit, 1:400, Sigma-Aldrich	Cy3 donkey-anti rabbit, 1:1,000, Jackson Immuno Research Laboratories, Inc.
	β -Dystroglycan (H-242), anti-rabbit, 1:200, Santa Cruz Biotech. Inc.	Cy3 donkey-anti rabbit, 1:1,000, Jackson Immuno Research Laboratories, Inc.
	CD 31 (PECAM-1), anti-rat, 1:800, BD Biosciences	Cy5 donkey-anti-rat, 1:1,000, Jackson Immuno Research Laboratories, Inc.
	Dystrophin (DP-71), anti-rabbit, 1:100, Abcam	Cy3 donkey-anti rabbit, 1:1,000, Jackson Immuno Research Laboratories, Inc.
Immunoperoxidase	Agrin, anti-rabbit, 1:1,000, Gift from Professor Markus A. Ruegg, University of Basel	Biotinylated donkey-anti-rabbit, 1:100, Pierce
	Laminin (L9393), anti-rabbit, 1:100, Sigma-Aldrich	Biotinylated donkey-anti-rabbit, 1:100, Pierce
Western blot	Agrin, anti-rabbit, 1:1,000, Gift from Professor P. Sonderegger	Anti-rabbit AP antibody, 1:10,000, Sigma-Aldrich
	α -Syntrophin, anti-rabbit, 1:1,000, Abcam	Anti-rabbit AP antibody, 1:10,000, Sigma-Aldrich
	AQP4, anti-rabbit, 1:1,500, Chemicon, Millipore	Anti-rabbit AP antibody, 1:10,000, Sigma-Aldrich
	β -Actin (A 2,066), anti-Rabbit, 1:500, Sigma-Aldrich	Anti-rabbit AP antibody, 1:10,000, Sigma-Aldrich
	β -Dystroglycan (H-242), anti-rabbit, 1:750, Santa Cruz Biotech. Inc.	Anti-rabbit AP antibody, 1:10,000, Sigma-Aldrich
	Dystrophin, anti-rabbit, 1:500, Abcam	Anti-rabbit AP antibody, 1:10,000, Sigma-Aldrich
	Kir4.1, anti-rabbit, 1:400, Alomone	Anti-rabbit AP antibody, 1:10,000, Sigma-Aldrich
	Laminin (L9393), anti-rabbit, 1:2,000, Sigma-Aldrich	Anti-rabbit AP antibody, 1:10,000, Sigma-Aldrich

For normalization, the membranes were washed several times in TBS-T [17 mM Trizma base (Sigma), 140 mM NaCl, 0.05 % Tween] following incubation with primary and secondary antibody as above.

Actin was used as a loading control. Membranes for which actin could not be used, Ponceau S staining was used instead (Aldridge et al. 2008; Romero-Calvo et al. 2010). This was the case for Kir4.1 and β -dystroglycan, both of which have MWs very close to that of actin, and in the case of laminin and agrin, which have MWs that are too high to be included in the same blots as actin (Aldridge et al. 2008; Romero-Calvo et al. 2010). Briefly, the membranes were incubated in Ponceau S stain for 3 min, washed until the colored protein bands could be visualized clearly and then scanned in a table top scanner at 600 dpi. The picture was analyzed in Photoshop using the Histogram function. The lane was marked with a thin stripe and the value was read off in the histogram. Linearity was obtained for Ponceau S staining as well as for actin controls.

Quantitation

For the quantification of the bands in SDS-PAGE and Western blotting two methods were used:

1. Samples from each age were loaded and compared to a standard curve on the same membrane. The fluorescent signal was detected with a Typhoon 9410 Workstation (Typhoon 9410, Amersham Biosciences, Piscataway, NJ, USA) and quantified with a commercially available image analysis program (ImageQuant™ TL, Amersham Biosciences, Piscataway, NJ, USA).
2. All samples were loaded on the same membrane. The fluorescent signal was detected with a Kodak scanner (Kodak Image Station 4000 MM PRO) and quantified with a commercially available image analysis program (Kodak MI SE, Carestream Health, Inc., Rochester, NY, USA).

Table 2 Primer sequences for real time PCR

Gene	Forward	Reverse
<i>Agrn</i>	CAGTGGGGGACCTAGAAACA	ATGGCCAGAGCCATGTAGTC
<i>Aqp4</i>	TTTGGACCCGCAGTTATCAT	GTTGTCTCCACCTCCATGT
<i>Dag1</i>	CCGAGAAGAGCAGTGAGGAC	AGCTCATCCGAAAGATGAT
<i>Dp71</i>	CAAGCTTACTCCTCCGCTCT	GAGCCTTCTGAGCTTCATGG
<i>Kcnj10</i>	TTATCAGAGCAGCCACTTCACC	TCTCTGTCTGAGTCGTCTGAC
<i>Lama1</i>	TGGATAAAGACAGGCCCTTG	ACTTTGGCACTGCTGATTCC
<i>Lama2</i>	ACCAGCCTACCTCCAGCTTT	CCCATTCCATCCATCTTCTG
<i>Snta1</i>	GCTGGCTGACAGAACAGTTG	TTCTGCATCATAGGGCACTG
<i>Tbp</i>	ACGGACAACCTGCGTTGATT	CAAGGCCTTCCAGCCTTATAG

Fig. 1 Members of the EBJC complex accumulate at brain surface and around vessels. Confocal images showing immunofluorescence labeling (red) of AQP4, α -syntrophin and β -dystroglycan in neocortex of postnatal mouse brain. These three members of the EBJC appear first at the brain surface and around penetrating vessels (lining the Virchow-Robins spaces), then around brain microvessels. Vessels (arrowheads) are identified by use of an endothelial marker (CD 31; blue). At P0, there was weak or no perivascular labeling for AQP4. At P7, AQP4 starts to

appear around vessels (arrowheads), and at P13 and P21 there is AQP4 labeling around vessels of all calibers. Subpial AQP4 labeling (arrows) is present already at P0 and increases in strength towards P21. α -Syntrophin follows the same labeling pattern as AQP4. Weak labeling is observed for β -dystroglycan at P0. At P7, β -dystroglycan labeling is present at the brain surface (arrows) and around vessels of all calibers (arrow heads), thus being more extensive at this stage than the labeling for AQP4 and α -syntrophin. Scale bar 50 μ m

Antibodies

For the complete list of primary and secondary antibodies used in this study see Table 1.

Quantitative real-time PCR

The mice ($n = 4$ for each age group) were anesthetized, decapitated and the brain was immediately frozen in liquid

Fig. 2 Immunogold analysis shows that subpial endfeet are the first to accumulate AQP4. **a–h** Postnatal immunogold labeling of AQP4 at the perivascular (**a–d**) and subpial (**e–h**) astrocyte membranes in mouse neocortex. Perivascular and subpial membrane domains are indicated by arrows. **a** At P0, the pericapillary basal lamina (asterisk) is immature, and there is no perivascular AQP4 immunogold labeling. **b** At P7, the basal lamina is distinct, and AQP4 appears in the perivascular membranes. **c, d** The AQP4 immunogold density increases further at P13 and P21. **e–h** The subpial basal lamina is well developed already at P0, and is associated with distinct AQP4

immunogold labeling in subpial astrocyte membranes. At P13 and P21, the AQP4 labeling extends into the glial lamellae beneath the pial surface. *E* endothelial cells, *L* vessel lumen, *A* astrocyte (Scale bar 200 nm). **i, j** Quantitative analysis of AQP4 immunogold labeling in perivascular (**i**) and subpial (**j**) membranes. At P4 and P7, the linear density of gold particles (no. of particles per μm membrane) is higher in subpial membranes than in perivascular ones. **Significantly different from P0; 'x' significantly different from previous value. Error bars indicate ± 2 SE, $p = 0.05$

nitrogen and stored at -80°C . Whole mouse brains were isolated, treated with RNA later (Ambion) and stored at -80°C until further processing. Quantitative real-time PCR was carried out the same way as by (Boldt and Conover 2011) with the following exceptions. Total RNA was extracted using RNeasy Lipid Tissue kit (Qiagen). 5 mg of RNA was reverse-transcribed using oligo d(T)18 primers and RevertAid H

Minus First Strand cDNA transcription reagents (Fermentas) in a reaction volume of 20 μl . The cDNA reaction mixture was diluted 100-fold in 10 mM Tris-HCl (pH 8.0), and real-time PCR was set up using 2 μl template in a reaction mixture of 20 μl containing AB Power SYBR Green PCR Master Mix (Applied Biosystems) and specific primers (200 nM) for amplification of target genes. Primers were designed using

Fig. 3 Agrin and laminin are present already at P0. Light microscopic pictures showing immunoperoxidase labeling of agrin (*left column*) and laminin (*right column*) in postnatal mouse neocortex. Agrin and laminin are strongly expressed at the pial surface (*arrows*) and around vessels (*arrow heads*) at P0. The labeling for both molecules becomes weaker towards P28. Labeling is removed by omission of primary antibodies (lower 2 rows; sections from P28 and P0, respectively). Scale bar 100 μ m

Primer3 software. Expression levels of target genes were assessed by absolute quantification. Real-time PCR was performed with plates using StepOnePlus real-time PCR system (Applied Biosystems). Amplicons (~ 200 bps) were generated in a two-step PCR (95 $^{\circ}$ C for 15 s, 60 $^{\circ}$ C for 60 s, 40 cycles). TATA box binding protein (TBP) was used as an endogenous control for normalization of gene expression. TBP was selected among several candidate housekeeping genes evaluated for this study because it was found to have the most constant level of expression across the developmental sample groups. The lower limit of detection was identified through amplification of targets known not to be expressed in mouse brain resulting in Ct values >32 corresponding to copy numbers <10 per ng total RNA.

For the complete list of primers see Table 2.

Statistical analysis

Obtained values were transferred to SPSS Version 16.0 (SPSS, Chicago, IL) and mean value of the groups were compared using ANOVA test. Data are presented as mean $\pm$ SE.

Results

Immunohistochemistry

The immunofluorescence analysis showed similar developmental profiles for AQP4, α -syntrophin, and β -dystroglycan (Fig. 1). These molecules are all members of the EBJC. None of these molecules were associated with the endothelial marker CD31 at P0. At P7, immunosignals for AQP4, α -syntrophin and β -dystroglycan were found subpially and around brain microvessels, with somewhat more prominent signals in the former site. β -Dystroglycan leads the other two molecules when it comes to accumulation around the endothelial lining (Fig. 1, second row, P7). At later developmental stages (P13 and P21) the diffuse neuropil labeling (pronounced for α -syntrophin and β -dystroglycan, less so for AQP4) subsides, leaving behind a rather distinct subpial and perivascular labeling. At P21, the subpial zone of immunofluorescence is broader for AQP4 than for the other members of the EBJC.

Fig. 4 Agrin and laminin are confined to the perivascular basal lamina. Immunogold labeling confirms localization of agrin (**a–d**) and laminin (**e–h**) to the perivascular basal lamina (arrows). Both proteins are present throughout the postnatal period. Electron micrographs of postnatal mouse neocortex. *E* endothelial cells, *L* vessel lumen, *asterisk* tight junction. Scale bar 0.5 μ m

Immunogold cytochemistry confirmed and extended the immunofluorescence analysis. AQP4 could not be detected at P0, when the basal lamina was indistinct, but appeared at P4 with stronger signals subpially than perivascularly (Fig. 2). The immunogold signal for AQP4 increased from P4 to P21 (Fig. 2). AQP4 labeling of non-endfoot membranes occurred primarily in the subpial zone (particularly pronounced in Fig. 2g, h; Supplementary Fig. S1), as predicted from the immunofluorescence data (above).

In contrast to AQP4, α -syntrophin, and β -dystroglycan, the two basal lamina molecules agrin and laminin were strongly expressed already at P0 (Fig. 3). Both molecules occurred at the pial surface as well as perivascularly, and showed stronger DAB immunosignals at P7 than at later postnatal stages. The perivascular labeling, in particular, appeared to abate with age.

To verify that the DAB signal represented laminin and agrin in the appropriate location (i.e., in the basal lamina, consistent with their being members of the EBJC), an

immunogold analysis was performed. Immunogold particles for agrin and laminin were superimposed on the perivascular and subpial basal laminae as early as P0 (Figs. 4 and 5, respectively).

The specificity of the antisera was confirmed by using knockout animals in case of AQP4, α -syntrophin and β -dystroglycan (not shown). The antibody to agrin has been tested previously on agrin knockout mice (Stephan et al. 2008). Knockout lines are not available for laminin. The precise localization of laminin and agrin immunosignals to the basal lamina (Figs. 4 and 5) indicates absence of unspecific labeling. Labeling was abolished after omission of primary antibodies, ruling out nonselective binding of the secondary antibody.

Quantitative Real-Time PCR

The molecules under investigation formed two distinct groups in regard to the developmental profile of their

Fig. 5 Agrin and laminin also occur in subpial basal lamina. Electron micrographs of immunogold labeling of agrin (**a–d**) and laminin (**e–h**). Both proteins are present in the basal lamina (*arrows*) opposed to subpial astrocyte endfeet. Labeling is distinct already at P0 and persists throughout the postnatal period. Scale bar 0.5 μ m

respective messengers (Fig. 6). mRNAs encoding AQP4 and other members of the dystrophin complex (α -syntrophin, β -dystroglycan, and the dystrophin isoform DP71) were scarce at P0 and increased in abundance towards a distinct peak at P13 (AQP4 and α -syntrophin) or a broader peak at P7–P13/21 (β -dystroglycan and DP71). DP71 and Kir4.1 (both being members of the DAPC) stood out as the only molecules whose messengers continue to increase until adulthood.

The second group of mRNAs (i.e., those encoding laminin and agrin) showed entirely different developmental profiles. Agrin-mRNA and Lama1 (encoding the α 1 isoform of laminin) were more abundant at early postnatal stages than at later stages in development (Fig. 6f, g). Both mRNAs peaked near P4. Lama2 (encoding the α 2 isoform of laminin) was relatively weakly expressed at P0 and peaked at P13. Lama2 differed from Lama1 by showing a rather stable expression level towards adulthood (Fig. 6g, h).

Immunoblots

The molecules under study segregate in three groups in regard to their expression at the protein level (Figs. 7 and 8).

AQP4 mirrored α -syntrophin and Kir4.1 in showing a continuous increase from being close to undetectable at P0 to being strongly expressed at adult stages. DP71 and β -dystroglycan, on the other hand, are rather stable throughout postnatal development. Laminin and agrin formed a third group that peaked at P7 and thereafter displayed a sharp decline towards adulthood.

Discussion

Evidence is accruing that the EBJC plays a central role in brain function and neurological disease. Hence, the integrity of the EBJC is a *conditio sine qua non* for maintaining the functional and biochemical polarization of astroglial cells. By orchestrating the molecular organization of the end-foot membranes of astrocytes, the EBJC serves a role akin to that of the protein complex that dictates the organization of pre- and postsynaptic membranes of neurons.

While the properties and developmental profiles of proteins responsible for neuronal polarization have been subjected to intensive studies (Arimura and Kaibuchi 2007)

much less is known about the proteins of the EBJC. Notably, the sequence by which the different EBJC proteins appear and reach peak expression has not been

investigated in detail. Insight in the developmental profile of the EBJC is likely to inform our understanding of the mechanisms that underpin the organization of the

Fig. 6 Different members of the EBJC complex have different mRNA signatures during development. **a–h** Quantitative real time PCR analysis of mouse brains at different stages of development. *Graphs* illustrate the copy number of different mRNA species, compared to the total amount of RNA (ng). The different EBJC members segregate in several groups, in regard to the developmental profile of their respective mRNAs. mRNAs encoding AQP4 and α -syntrophin are weakly expressed at birth and peak before adulthood, while mRNAs encoding agrin and Lama1 are abundant at birth with decreasing levels towards the adult stage. The remaining mRNA species show a rather stable expression throughout postnatal development (dystrophin, Dag1, Lama2) or a sharp increase in expression towards adulthood (Kir4.1). Dag1 encodes both α - and β -dystroglycan. **Significantly different from P0 and 'x' significantly different from previous value. *Error bars* indicate ± 2 SE, $p = 0.05$. **i** Representative DNA agarose gel electrophoresis showing the EBJC expression profile during postnatal development of mouse brain. PCR products were generated using representative cDNA samples from developmental stage P0, P4, P7, P13, P21 and adult (A) as indicated above each lane on the gel. A DNA marker (M) was included in the first lane. The EBJC sample set includes from top to bottom: AQP4, Kir4.1, α -syntrophin, Dp71, Dag1, agrin, Lama1, Lama2 and TBP. The latter was included to verify equivalent amounts of cDNA template across samples in the 30-cycle endpoint PCRs using GoTaq Green polymerase (Promega) and specific primers (Table 2). PCR product sizes in base pairs are shown to the right of each gel insert. Two bands are visible in the gel insert for agrin, which is explained by co-expression of two mRNA species resulting from alternative splicing

gliovascular interface and of the glia limitans at the brain surface. The gliovascular interface attracts particular interest in the context of cerebral blood flow regulation (Iadecola and Nedergaard 2007) while new data indicate that glia limitans is specifically implicated in volume control (Papadopoulos et al. 2004; Benfenati et al. 2011).

Unravelling the developmental profile of the EBJC requires a high anatomical resolution that matches the spatially confined expression of this junctional complex. At the same time, there must be possibilities for quantitation in order to accurately determine the peak of expression. Post-embedding immunogold procedures are uniquely suited to the task at hand. The lateral resolution of this procedure is in the range of ~ 30 nm, which compares favorably with the lateral dimension of the endfeet (in the order of 1000 nm) and the thickness of the basal lamina (in the order of 100 nm). Gold particles are easily identified and their number bears a well-defined relation—often close to linear (Mathiisen et al. 2006; Amiry-Moghaddam and Ottersen 2013)—to the number of target proteins. To the best of our knowledge this study represents the first successful immunogold application in brain of antibodies to pan-laminin and agrin. The basal lamina proteins laminin and agrin were the first proteins to appear at the gliovascular interface, according to the present analysis. Both proteins were detected at P0, predating the appearance of AQP4. However, a possible caveat needs to be considered:

the threshold for detection might differ between different proteins. Thus, proteins may be differentially affected by steric hindrance or by the relatively harsh conditions that prevail during tissue preparation for immunogold cytochemistry. It is of note that the freeze substitution approach currently applied is more “antigen friendly” than traditional procedures for tissue preparation (Amiry-Moghaddam and Ottersen 2013). Also, the three complementary techniques that were used indicate that the specific sequence of labeling intensity that we observed at the immunogold level reflects a *bona fide* developmental profile. Thus, immunofluorescence, immunoblots, and qRT-PCR concurred in showing early expression of laminin and agrin. The data also pointed to an abatement of laminin and agrin expression after the first postnatal week. This expression profile of agrin and laminin is consistent with these proteins having an instructive role in establishing the EBJC.

In the case of laminin and agrin, we did not perform a quantitative analysis of the immunogold labelling. For these proteins, the quantitative data obtained with Western blots and qRT-PCR complement the qualitative data obtained with the immunogold approach.

As the expression level of several of the proteins normally present at the perivascular astrocyte endfeet is very low at P0, establishment of astrocyte polarization must depend on two factors: (1) increased postnatal synthesis of perivascular endfoot proteins, and (2) postnatal insertion and stabilization of the proteins in perivascular endfoot membranes.

In vitro evidence suggests that agrin plays a role in the establishment of astrocyte polarization. A recent study has shown that exposure to agrin isoforms leads to increased AQP4 expression, membrane localization and formation of orthogobal arrays of particles (OAP) in wild type and agrin-null mouse astrocytes (Fallier-Becker et al. 2011). The same effect was achieved by replacing exogenous agrin with ECM purified from Engelbreth-Holm-Swarm (EHS) mouse sarcoma, where laminin, collagen IV, heparan sulfate proteoglycan (a class of molecules that includes agrin), and nidogen/entactin are the major components. Our study shows that expression of agrin is at its highest level in the first postnatal week, prior to the peak of AQP4 expression, supporting the idea that agrin stimulates expression of AQP4. We did not distinguish between the neuronal and endothelial isoforms of agrin and cannot say whether these isoforms play distinct roles. Of note, agrin-null astrocytes did express AQP4 and OAPs, though at lower levels than did wild type controls, indicating that AQP4 expression and OAP formation do not depend solely on the presence of agrin. This conclusion finds support in a recent analysis of Rauch et al. (2011). These authors demonstrated a residual pool of AQP4 following targeted

Fig. 7 AQP4, Kir4.1, and α -syntrophin increase towards adulthood. Western blots of whole brain homogenates from the postnatal day 0 to 21 (P0–P21) and adult (A) mice. Representative immunoblots for AQP4, Kir4.1, and α -syntrophin (left panels) and corresponding quantitation (densitometric values; right panels). **a** The AQP4 antibody labelled two bands at about 30 kDa corresponding to the M1 and M23 isoforms of AQP4. A third band around 35 kDa was not included in the quantitative analysis. β -Actin was used as loading control. The densitometric analysis revealed an increasing immunosignal for AQP4 protein in the postnatal period. **b** Immunoblot of Kir4.1 revealed a major band at ≈ 200 kDa which corresponds to the

tetrameric form of Kir4.1 (Connors and Kofuji 2006; Olsen et al. 2006). Ponceau red staining was used as loading control (not shown). The developmental pattern mimics that of AQP4 (**a**). **c** Immunoblot of α -syntrophin revealed a major band at 59 kDa and a weaker band at slightly higher molecular weight. The major band—absent from α -syntrophin knockout brains—was used for quantitative analysis. β -Actin was used as loading control. The expression pattern for α -syntrophin was similar to those of AQP4 and Kir4.1. **Significantly different from P0 and 'x' significantly different from previous value. Error bars indicate ± 2 SE, $p = 0.05$

deletion of endothelial agrin. This residual pool might be kept in place by laminin, as discussed below.

Laminin has been suggested to play an important role in membrane localization of AQP4 in astrocyte endfeet. In vitro studies have shown that laminin clusters AQP4 and Kir4.1 in astrocyte membranes, and stabilizes these proteins at the basolateral domain of epithelial Madine-Darby canine kidney (MDCK) cells (Guadagno and Moukhles 2004; Tham and Moukhles 2011). The laminin α chain comes in two major isoforms in brain: α -1 and α -2. The antibodies used in this study are raised against full-length laminin and hence cannot distinguish between the different isoforms. However, our qRT-PCR data indicate that the expression of α -1 leads the expression of α -2. α -1 is the laminin isoform that predominates at the pial surface while α -2 predominates perivascularly. This developmental profile is entirely consistent with an instructive role of laminin, as our data suggest that AQP4 appears at an earlier stage subpially than perivascularly.

The developmental profile of α -syntrophin closely mimics that of AQP4. This finding bolsters the increasing body of evidence pointing to α -syntrophin as the immediate anchor of the perivascular AQP4 pool (Amiry-Moghaddam and Ottersen 2003; Nagelhus and Ottersen 2013). Dystrophin and dystroglycan display yet another developmental profile. These proteins can be detected early in the postnatal phase, but are then rather diffusely distributed. Only at day 7 do they begin to show a distinct perivascular accumulation. This coincides with the initial expression of AQP4.

The most parsimonious explanation of our findings is that laminin and agrin cause AQP4 and α -syntrophin to aggregate at endfoot membranes, through recruiting dystrophin and β -dystroglycan to this membrane domain. Previous studies have demonstrated that both dystrophin and β -dystroglycan are required for perivascular expression of AQP4 (Frigeri et al. 2001; Noell et al. 2011).

We also included an analysis of the inwardly rectifying K^+ channel Kir4.1. At the protein level, Kir4.1 showed a

(a) Dystrophin (Dp-71) **(b) β -Dystroglycan** **(c) Agrin** **(d) Laminin**

Fig. 8 Agrin and laminin decrease with postnatal age. Western blots of whole mouse brain homogenate. Representative immunoblots of DP71, β -dystroglycan, agrin and laminin at different postnatal ages and adult (A) mice (*left panels*). The immunoblots were subjected to densitometric analysis (*right panels*). **a** Immunoblot of DP71 revealed a band at ≈ 71 kDa. There were also bands at ≈ 65 kDa and ≈ 55 kDa which could represent other dystrophin isoforms or degradation products. β -Actin was used as loading control. The immunosignal for DP71 was rather stable in the postnatal period. **b** Immunoblot of β -dystroglycan revealed a major band at ≈ 42 kDa. Ponceau red staining was used as loading control (not shown). The protein level is stable in the postnatal period. **c** Immunoblot of agrin revealed a major band at ≈ 300 kDa as shown previously (Stephan

et al. 2008). Ponceau red staining was used as loading control (not shown). The immunosignal for agrin reached a peak at P7 (significantly higher than P0) with a sharp decline towards the adult level. **d** Immunoblot of laminin with whole brain homogenate revealed a major band at ≈ 200 kDa and two weaker bands at ≈ 400 kDa and ≈ 600 – 700 kDa. The 200 kDa band corresponds to the β - and γ -chain of laminin while the 400 kDa band corresponds to the $\alpha 1$ -chain (Zhang et al. 2007). Ponceau red staining was used as loading control (not shown). Quantitative analysis of the 200 kDa band shows a pattern similar to that of agrin. Quantitation of the other two bands revealed no significant differences between the different postnatal stages. **Significantly different from P0 and 'x' significantly different from previous value. Error bars indicate ± 2 SE, $p = 0.05$

developmental profile similar to that of AQP4. Previously it has been shown that these two proteins are strictly colocalized in macroglial membranes (Nagelhus et al. 1999). However, Kir4.1 mRNA differs from AQP4 mRNA in showing a sharp increase with age. Pending data on Kir4.1 regulation and turnover we cannot explain this intriguing difference between AQP4 and Kir4.1.

To what extent are the present findings relevant for the human brain? A study by El-Khoury et al. (2006) shows that AQP4 positive perivascular endfeet appear between gestational weeks 19–22 in humans (El-Khoury et al. 2006). Given that term in mice corresponds to gestational day ~120 in humans with respect to brain development, the finding of El-Khoury et al. (2006) is in good agreement with the situation in mice in which AQP4 positive endfeet emerge during the first postnatal days. Further studies are needed to establish whether the junctional complex in humans mimics that in mice in regard to the order of appearance of its molecular constituents.

There is now a growing awareness that loss of astroglial polarization occurs in a number of pathophysiological conditions, including temporal lobe epilepsy, ischemia and Alzheimer's disease (Eid et al. 2005; Frydenlund et al. 2006; Yang et al. 2011; Heuser et al. 2012; Alvestad et al. 2013). What these diverse conditions have in common is a deficiency in homeostatic mechanisms that can easily be attributed to defunct handling of K^+ and/or water. Notably, each of these conditions is characterized by neuronal hyperexcitability and increased proneness to seizures. Given the results obtained in the present study, there is a need to consider the possibility that the extracellular matrix (ECM) is more directly involved in cerebral pathophysiology than previously believed. Specifically, a loss of the instructive role of laminin or agrin could figure prominently in neurological disease. Proteolytic activities in the extracellular space, such as metalloproteinase activation (Michaluk et al. 2007), could be an upstream factor. These issues merit further investigation.

Conclusion

Taken together with previous in vitro data, the present in vivo analysis suggests that the molecular complex of the gliovascular interface is assembled through a specific sequence of events. The basal lamina proteins laminin and agrin appear first, and stabilize β -dystroglycan and dystrophin at endfoot membranes. AQP4 and Kir4.1 are then recruited to the dystrophin complex in amounts that continue to increase towards adulthood. The developmental profile of α -syntrophin mimics that of AQP4, consistent with the idea that α -syntrophin serves as the immediate anchor of this water channel. We conclude that the ECM is likely to play a pivotal

role in the development of the molecular assembly at the gliovascular interface. The pathophysiological roles of ECM warrant closer examination.

Acknowledgments This work was supported by the Nordic Council and Polish-Norwegian Research Fund Grant PNR-96. Thanks to Ms. Bjørg Riber, Karen Marie Gujord and Jorunn Knutsen for technical assistance and Carina Knudsen and Gunnar F. Lothe for the artwork. The two different agrin antibodies used in this study were kind gifts from Professor P. Sonderegger and Professor Markus A. Ruegg.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Aldridge GM, Podrebarac DM, Greenough WT, Weiler IJ (2008) The use of total protein stains as loading controls: an alternative to high-abundance single-protein controls in semi-quantitative immunoblotting. *J Neurosci Methods* 172(2):250–254
- Alvestad S, Hammer J, Hoddevik EH, Skare O, Sonnewald U, Amiry-Moghaddam M, Ottersen OP (2013) Mislocalization of AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy. *Epilepsy Res* 105(1–2):30–41
- Amiry-Moghaddam M, Ottersen OP (2003) The molecular basis of water transport in the brain. *Nat Rev Neurosci* 4(12):991–1001
- Amiry-Moghaddam M, Ottersen OP (2013) Immunogold cytochemistry in neuroscience. *Nat Neurosci* 16(7):798–804
- Amiry-Moghaddam M, Otsuka T, Hurn PD, Traystman RJ, Haug FM, Froehner SC, Adams ME, Neely JD, Agre P, Ottersen OP, Bhardwaj A (2003) An alpha-syntrophin-dependent pool of AQP4 in astroglial end-feet confers bidirectional water flow between blood and brain. *Proc Natl Acad Sci USA* 100(4):2106–2111
- Amiry-Moghaddam M, Xue R, Haug FM, Neely JD, Bhardwaj A, Agre P, Adams ME, Froehner SC, Mori S, Ottersen OP (2004) Alpha-syntrophin deletion removes the perivascular but not endothelial pool of aquaporin-4 at the blood-brain barrier and delays the development of brain edema in an experimental model of acute hyponatremia. *FASEB J* 18(3):542–544
- Arimura N, Kaibuchi K (2007) Neuronal polarity: from extracellular signals to intracellular mechanisms. *Nat Rev Neurosci* 8(3):194–205
- Benfenati V, Amiry-Moghaddam M, Caprini M, Mylonakou MN, Rapisarda C, Ottersen OP, Ferroni S (2007) Expression and functional characterization of transient receptor potential vanilloid-related channel 4 (TRPV4) in rat cortical astrocytes. *Neuroscience* 148(4):876–892
- Benfenati V, Caprini M, Dovizio M, Mylonakou MN, Ferroni S, Ottersen OP, Amiry-Moghaddam M (2011) An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex is essential for cell-volume control in astrocytes. *Proc Natl Acad Sci USA* 108(6):2563–2568
- Boldt HB, Conover CA (2011) Overexpression of pregnancy-associated plasma protein-A in ovarian cancer cells promotes tumor growth in vivo. *Endocrinology* 152(4):1470–1478
- Chaudhry FA, Lehre KP, van Lookeren Campagne M, Ottersen OP, Danbolt NC, Storm-Mathisen J (1995) Glutamate transporters in glial plasma membranes: highly differentiated localizations revealed by quantitative ultrastructural immunocytochemistry. *Neuron* 15(3):711–720

- Connors NC, Kofuji P (2006) Potassium channel Kir4.1 macromolecular complex in retinal glial cells. *Glia* 53(2):124–131
- Eid T, Lee TS, Thomas MJ, Amiry-Moghaddam M, Bjornsen LP, Spencer DD, Agre P, Ottersen OP, de Lanerolle NC (2005) Loss of perivascular aquaporin 4 may underlie deficient water and K⁺ homeostasis in the human epileptogenic hippocampus. *Proc Natl Acad Sci USA* 102(4):1193–1198
- El-Khoury N, Braun A, Hu F, Pandey M, Nedergaard M, Lagamma EF, Ballabh P (2006) Astrocyte end-feet in germinal matrix, cerebral cortex, and white matter in developing infants. *Pediatr Res* 59(5):673–679
- Fallier-Becker P, Sperveslage J, Wolburg H, Noell S (2011) The impact of agrin on the formation of orthogonal arrays of particles in cultured astrocytes from wild-type and agrin-null mice. *Brain Res* 1367:2–12
- Frigeri A, Nicchia GP, Nico B, Quondamatteo F, Herken R, Roncali L, Svelto M (2001) Aquaporin-4 deficiency in skeletal muscle and brain of dystrophic mdx mice. *Faseb J* 15(1):90–98
- Frydenlund DS, Bhardwaj A, Otsuka T, Mylonakou MN, Yasumura T, Davidson KG, Zeynalov E, Skare O, Laake P, Haug FM, Rash JE, Agre P, Ottersen OP, Amiry-Moghaddam M (2006) Temporary loss of perivascular aquaporin-4 in neocortex after transient middle cerebral artery occlusion in mice. *Proc Natl Acad Sci USA* 103(36):13532–13536
- Guadagno E, Moukles H (2004) Laminin-induced aggregation of the inwardly rectifying potassium channel, Kir4.1, and the water-permeable channel, AQP4, via a dystroglycan-containing complex in astrocytes. *Glia* 47(2):138–149
- Haj-Yasein NN, Vindedal GF, Eilert-Olsen M, Gundersen GA, Skare Ø, Laake P, Klungland A, Thorén AE, Burkhardt JM, Ottersen OP, Nagelhus EA (2011) Glial-conditional deletion of aquaporin-4 (Aqp4) reduces blood-brain water uptake and confers barrier function on perivascular astrocyte endfeet. *Proc Natl Acad Sci U S A* 108(43):17815–17820
- Heuser K, Eid T, Lauritzen F, Thoren AE, Vindedal GF, Tauboll E, Gjerstad L, Spencer DD, Ottersen OP, Nagelhus EA, de Lanerolle NC (2012) Loss of perivascular Kir4.1 potassium channels in the sclerotic hippocampus of patients with mesial temporal lobe epilepsy. *J Neuropathol Exp Neurol* 71(9):814–825
- Iadecola C, Nedergaard M (2007) Glial regulation of the cerebral microvasculature. *Nat Neurosci* 10(11):1369–1376
- Iliff JJ, Wang M, Liao Y, Plogg BA, Peng W, Gundersen GA, Benveniste H, Vates GE, Deane R, Goldman SA, Nagelhus EA, Nedergaard M (2012) A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid beta. *Sci Transl Med* 4(147):147ra111
- Mathiisen TM, Nagelhus EA, Jouleh B, Torp R, Frydenlund DS, Mylonakou M-N, Amiry-Moghaddam M, Covolan L, Utvik JK, Riber B, Gujard KM, Knutsen J, Skare Ø, Laake P, Davanger S, Haug F-M, Rinvik E, Ottersen OP (2006) Postembedding immunogold cytochemistry of membrane molecules and amino acid transmitters in the central nervous system. *Neuroanat Tract-Tracing* 3:37
- Mathiisen TM, Lehre KP, Danbolt NC, Ottersen OP (2010) The perivascular astroglial sheath provides a complete covering of the brain microvessels: an electron microscopic 3D reconstruction. *Glia* 58(9):1094–1103
- Michalak P, Kolodziej L, Mioduszevska B, Wilczynski GM, Dzwonek J, Jaworski J, Gorecki DC, Ottersen OP, Kaczmarek L (2007) Beta-dystroglycan as a target for MMP-9, in response to enhanced neuronal activity. *J Biol Chem* 282(22):16036–16041
- Nagelhus EA, Ottersen OP (2013) Physiological roles of aquaporin-4 in brain. *Physiol Rev* 93(4):1543–1562
- Nagelhus EA, Horio Y, Inanobe A, Fujita A, Haug FM, Nielsen S, Kurachi Y, Ottersen OP (1999) Immunogold evidence suggests that coupling of K⁺ siphoning and water transport in rat retinal Muller cells is mediated by a coenrichment of Kir4.1 and AQP4 in specific membrane domains. *Glia* 26(1):47–54
- Neely JD, Amiry-Moghaddam M, Ottersen OP, Froehner SC, Agre P, Adams ME (2001) Syntrophin-dependent expression and localization of aquaporin-4 water channel protein. *Proc Natl Acad Sci USA* 98(24):14108–14113
- Newman EA, Frambach DA, Odette LL (1984) Control of extracellular potassium levels by retinal glial cell K⁺ siphoning. *Science* 225(4667):1174–1175
- Nielsen S, Nagelhus EA, Amiry-Moghaddam M, Bourque C, Agre P, Ottersen OP (1997) Specialized membrane domains for water transport in glial cells: high-resolution immunogold cytochemistry of aquaporin-4 in rat brain. *J Neurosci* 17(1):171–180
- Noell S, Wolburg-Buchholz K, Mack AF, Beedle AM, Satz JS, Campbell KP, Wolburg H, Fallier-Becker P (2011) Evidence for a role of dystroglycan regulating the membrane architecture of astroglial endfeet. *Eur J Neurosci* 33(12):2179–2186
- Olsen ML, Higashimori H, Campbell SL, Hablitz JJ, Sontheimer H (2006) Functional expression of Kir4.1 channels in spinal cord astrocytes. *Glia* 53(5):516–528
- Papadopoulos MC, Manley GT, Krishna S, Verkman AS (2004) Aquaporin-4 facilitates reabsorption of excess fluid in vasogenic brain edema. *Faseb J* 18(11):1291–1293
- Promeneur D, Lunde LK, Amiry-Moghaddam M, Agre P (2013) Protective role of brain water channel AQP4 in murine cerebral malaria. *Proc Natl Acad Sci USA* 110(3):1035–1040
- Rauch SM, Huen K, Miller MC, Chaudry H, Lau M, Sanes JR, Johanson CE, Stopa EG, Burgess RW (2011) Changes in brain beta-amyloid deposition and aquaporin 4 levels in response to altered agrin expression in mice. *J Neuropathol Exp Neurol* 70(12):1124–1137
- Romero-Calvo I, Ocon B, Martinez-Moya P, Suarez MD, Zarzuelo A, Martinez-Augustin O, de Medina FS (2010) Reversible Ponceau staining as a loading control alternative to actin in Western blots. *Anal Biochem* 401(2):318–320
- Somjen GG (1988) Nervenkitz: notes on the history of the concept of neuroglia. *Glia* 1(1):2–9
- Stephan A, Mateos JM, Kozlov SV, Cinelli P, Kistler AD, Hettwer S, Rulicke T, Streit P, Kunz B, Sonderegger P (2008) Neurotrypsin cleaves agrin locally at the synapse. *Faseb J* 22(6):1861–1873
- Tham DK, Moukles H (2011) Regulation of Kir4.1 and AQP4 expression and stability at the basolateral domain of epithelial MDCK cells by the extracellular matrix. *Am J Physiol Renal Physiol* 301(2):F396–F409
- Vajda Z, Pedersen M, Fuchtbauer EM, Wertz K, Stodkilde-Jorgensen H, Sulyok E, Doczi T, Neely JD, Agre P, Frokiaer J, Nielsen S (2002) Delayed onset of brain edema and mislocalization of aquaporin-4 in dystrophin-null transgenic mice. *Proc Natl Acad Sci USA* 99(20):13131–13136
- Wolburg H, Wolburg-Buchholz K, Fallier-Becker P, Noell S, Mack AF (2011) Structure and functions of aquaporin-4-based orthogonal arrays of particles. *Int Rev Cell Mol Biol* 287:1–41
- Yang J, Lunde LK, Nuntagij P, Oguchi T, Camassa LM, Nilsson LN, Lannfelt L, Xu Y, Amiry-Moghaddam M, Ottersen OP, Torp R (2011) Loss of astrocyte polarization in the tg-ArcSwe mouse model of Alzheimer's disease. *J Alzheimers Dis* 27(4):711–722
- Zhang X, Winkles JA, Gongora MC, Polavarapu R, Michaelson JS, Hahn K, Burkly L, Friedman M, Li XJ, Yepes M (2007) TWEAK-Fn14 pathway inhibition protects the integrity of the neurovascular unit during cerebral ischemia. *J Cereb Blood Flow Metab* 27(3):534–544

Errata:

Page	Paragraph	Line	Original text	Corrected text
4	1	14	...EMC proteins...	...ECM proteins...
5	1	38	...tight Junctions...	...tight junctions...
10	4	2	...thesis are...	...the thesis are...
12	4	3	...Laloux et.al 2018)	...Laloux et.al 2018).
13	2	13	...remain enigmatic...	...remains enigmatic...
14	3	1	...other organ...	...other organs...
14	3	4	...exercise induces...	...exercise induced...
15	1	31	...et al. 2018). "	...et al. 2018).
17	Fig 1	legend, 19	...may be also...	...may also be...
18	3	6	...as early in in...	...as early as in...
25	1	4	...in AQP4/Aqp4...	...in AQP4 protein/Aqp4 mRNA...
26	1	3	...is reported in Stephan et al. 2008 (Stephan et al. 2008)...	...is reported in Eusebio et al. 2003 (Eusebio et. al 2003)...
26	4	16	...antibody From Santa Cruz...	...antibody from Santa Cruz...
26	5	4	...(article IV; (Lunde et al.).	...(article IV; Lunde et.al.).
27	5	5	...tested by Stephan...	...tested by Eusebio...
27	5	5	...(Stephan et al. 2008)...	...(Eusebio et. al 2003)...
30	3	overskrift	...do endothelial cell...	...do endothelial cells...
31	2	7	...tisseu...	...tissue...
31	2	38	...presence of pericytes constitute...	...presence of pericytes constitutes...
32	3	23	...in regulation AQP4...	...in regulating AQP4...
33	3	1	...increased proteins...	...increased protein...
36	3	6	...confounding and...	...confounding factor and...
36	Fig 4	legend, 7	...would hvae...	...would have...
37	1	13	...this approach	... this approach yields evidence on the general tissue expression, including both parenchymal and perivascular protein pools.
37	2	14	...null hypothesis that ECM proteins are instructive to endfeet,...	null hypothesis - that ECM proteins are not instructive to endfeet -...
38	4	12	...extensively studies...	...extensively studied...
43		46	...	...Eusebio A, Oliveri F, Barzaghi P, Ruegg MA (2003) Expression of mouse agrin in normal, denervated and dystrophic muscle Neuromuscul Disord 13:408-415...