Measuring psychodynamic process and change in psychotherapy

Elisabeth Ness

Institute of Clinical Medicine
Faculty of Medicine

UNIVERSITY OF OSLO

2019

© Elisabeth Ness, 2019

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-459-7

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

Table of contents

Acknowledgements.....	3
Abbreviations.....	4
List of papers.....	5
Summary.....	6
1. Introduction.....	10
1.1 Research on psychotherapy.....	10
1.1.1 Psychotherapy outcome research.....	11
1.1.1.1 Instruments to measure outcome.....	12
1.1.2 Psychotherapy process research.....	13
1.1.2.1 Instruments to measure process.....	15
1.1.3 Operationalization.....	16
1.2 Measuring change in psychotherapy.....	17
1.2.1 Validity.....	19
1.2.2 Reliability.....	21
1.2.3 Interrater reliability.....	22
1.2.4 Clinical significance.....	23
1.3 Dynamic psychotherapy.....	24
1.3.1 Transference and transference work.....	27
1.4 Study designs.....	29
1.4.1 Single case studies.....	30
1.4.2 Randomized controlled trials.....	31
1.4.2.1 Dismantling design.....	32
2. Objectives.....	33
3. Material and methods.....	34
3.1 The First Experimental Study of Transference-interpretations (FEST).....	34
3.1.1 Patients.....	34
3.1.2 Treatment.....	34
3.1.3 Therapists and evaluators.....	35
3.1.4 Ethics.....	35

3.1.5 Assessments and outcome measures.....	36
3.1.6 Main results from FEST.....	37
3.2 The First Experimental Study of Transference work – In Teenagers (FEST-IT).....	37
3.2.1 Patients.....	37
3.2.2 Treatment.....	38
3.2.3 Therapists and evaluators.....	39
3.2.4 Ethics.....	40
3.2.5 Assessments and outcome measures.....	40
3.3 Measures used in FEST and FEST-IT.....	41
3.4 Some statistical considerations.....	47
4. Summary of results.....	50
4.1 Paper I.....	50
4.2 Paper II.....	52
4.3 Paper III.....	55
5. Discussion.....	57
5.1 Discussion of the main results.....	57
5.1.1 Methodological discussion.....	62
5.2 Strengths and limitations of the present study.....	63
6. Conclusion.....	64
7. Clinical implications and future research.....	65
7.1 Implications for clinical practice.....	65
7.2 Implications for future research.....	65
8. References.....	67
9. Appendix.....	76
9.1 Adolescent Relationship Scale (ARS).....	76
9.2 Quality of Object Relations (QOR).....	76
9.3 The factors from Feeling Word Checklist (FWC-24).....	77
10. Papers.....	81
Paper I	
Paper II	
Paper III	

Acknowledgements

This thesis is dedicated to my sister-in-law Eva who passed away. I miss you.

I am grateful to my parents Gerd and Arne, and my siblings Gro and Hans Jørgen. You were in different ways all supporting although it is likely you never grasped what I was actually doing in the field of research.

A special thanks to my husband Jon who has supported me along the way.

I am grateful to University of Oslo (UiO) and Vestfold Hospital Trust (Sykehuset i Vestfold) for funding and support. A special thanks to Research Unit, Division of Mental Health (KPR) and Department of Child and Adolescent Psychiatry (BUPA) in Vestfold.

Thanks to everyone I met at Vinderen (Adult Psychiatry Unit) and especially those who attended the interrater reliability-sessions. You were a group of open-minded, humorous, professional and welcoming researchers and collaborators eager to discuss clinical material.

I am also most grateful to my supervisors; Randi Ulberg, Hanne-Sofie Johnsen Dahl and Per Høgland. Randi, I still cannot understand how you can literally be available 24/7 for support, assistance and guidance – always with a smile.

Thanks again to everyone involved in FEST and FEST-IT, including the almost 200 patients.

Last, but by no means least, to my children Fredrik, Erik and Mari. You keep teaching me the importance of relationships and illustrate the everyday ups and downs that make everyone's life unique in experience, process – and change.

Abbreviations

APQ	Adolescent Psychotherapy Q-set
ARS	Adolescent Relationship Scale
BDI	Beck Depression Inventory
CBT	Cognitive Behavioural Therapy
DSM-IV	Diagnostic and Statistical Manual of Mental Disorders
FEST	First Experimental Study of Transference Interpretations
FEST-IT	First Experimental Study of Transference Work – In Teenagers
FWC	Feeling Word Checklist
GAF	Global Assessment of Functioning
GSI	Global Severity Index
ICC	Intraclass Correlation Coefficient
IIP-C	Inventory of Interpersonal Problems-Circumplex
IRR	Interrater reliability
MADRS	Montgomery-Asberg Depression Rating Scale
M.I.N.I.	Mini International Neuropsychiatric Interview
PFS	Psychodynamic Functioning Scales
QOR	Quality of Object Relations
RCI	Reliable Change Index
RCT	Randomized Controlled Trial
RWS	Relational Work Scale
SASB	Structural Analysis of Social Behaviour
SCL-90	Symptom Checklist-90
SIDP-IV	Structured Interview for DSM-IV Personality
STPP	Short-term psychoanalytical psychotherapy
TI	Transference Intervention
TWS	Transference Work Scale
WAI	Working Alliance Inventory

List of papers

- I.** Ness, E., Dahl, H.S.J., Critchfield, K., Ulberg, R. (2018). Exploring in-session process with qualitative and quantitative methods in psychotherapy with an adolescent. *Journal of Infant, Child, and Adolescent Psychotherapy* 17:4,310-327 <https://doi.org/10.1080/15289168.2018.1526021>
- II.** Ness, E., Dahl, H.S.J., Tallberg, P., Amlo, S., Høglend, P., Thorén, A., Egeland, J., Ulberg, R. Assessment of Dynamic Change in Psychotherapy with Adolescents. *BMC Child and Adolescent Psychiatry and Mental Health* 2018**12**:39 <https://doi.org/10.1186/s13034-018-0246-z>
- III.** Ulberg, R., Ness, E., Dahl, H.S.J., Høglend, P., Critchfield, K., Blayvas, P., Amlo, S. (2016). Relational interventions in psychotherapy: development of a therapy process rating scale. *BMC Psychiatry* (2016) 16:310. DOI 10.1186/s12888-016-1021-4 <http://www.biomedcentral.com/1471-244X/16/310>

Summary

Measuring psychodynamic process and change in psychotherapy is a way of adding knowledge to the evidence base of what leads to change. Studying mediators and processes leading to change is complex. Process-outcome studies have so far yielded little definite evidence for the mechanism of specific psychotherapies.

There is emerging evidence for the effect of psychotherapy in adolescents. Research indicates that therapists often use techniques from different theoretical models when working with adults. However, research on the therapy process in adolescents is scarce. Several factors make psychotherapy different with adolescents than with adults. Diagnostic interviews and questionnaires are commonly used in the assessment of adolescents referred to child and adolescent mental health services. Many of these rating scales are constructed for adults and focus on symptoms related to diagnosis. Although specific measures for assessment of dynamic change in psychodynamic therapies exist, there is limited research on adolescent therapies. Similar study designs permit some comparison to be explored in two studies of psychodynamic psychotherapy, the former with adults and the latter with adolescents.

Case studies have contributed to psychotherapy theory and practice since the beginning of the 20th century. A case study makes it possible to explore in-depth the therapist's interventions, the process and the patient's development and dynamic change to occur in an adolescent therapy.

There is a need to develop more methods to reveal what kind of treatment and treatment techniques work best for different patients, and how the various psychotherapy modes work to promote the best possible treatment effect. Interventions of interpreting the patient's relationships with other people are considered key components to enhance change in psychotherapy. Relational interventions, when the therapist focuses on the patient's relationships *outside* therapy, are by nature pan-theoretical. A rating tool for in-session process with transference interventions, when the therapist focuses on the therapist-patient relationship, already exist.

Objectives:

The main objective of this thesis was to investigate the use of different methods for measuring process and change in psychodynamic therapy with adults and adolescents. Data from two randomized studies adjusted for adults and adolescents were used to capture

different aspects of psychodynamic therapy through several measurements being specific and non-specific to psychodynamic therapy. Specific aims were:

- To explore the in-session process of an adolescent therapy with qualitative and quantitative methods to capture the process in terms of therapist interventions, changes in symptomatology, interpersonal problems, and alliance.
- To evaluate a psychometric instrument for assessment of change in dynamic psychotherapy (Psychodynamic Functioning Scales) for adolescents. The aim was to test the interrater reliability of five subscales as well as the reliability of change ratings and the discriminability from general symptoms.
- To develop a measure designed to identify defined categories of therapists' interventions focusing on the patients' relationships outside therapy (Relational Interventions).

Material and methods:

The three papers included in this thesis are based on one adolescent and one adult study, both designed to measure specific long-term effects of transference interpretations in dynamic psychotherapy, using an experimental dismantling design (FEST-IT and FEST). The adolescent study included depressed adolescents. In the adult study, one hundred psychiatric out-patients suffering from mood, anxiety, and personality disorders or interpersonal problems not due to a mental disorder, were randomized to one year (maximum 40 sessions) weekly dynamic psychotherapy, with or without transference interpretations. The randomization was made similar in both studies, although with a treatment manual adjusted for adolescents and a maximum of 28 weekly sessions in FEST-IT.

Paper I is a case study from an adolescent therapy. To focus on the process leading to change, i.e. including the therapist's interventions, different measures were used including clinician-rated and self-reported general symptom-scales, diagnose-specific scales (depression) and in-session rating tools used on audio-recorded sessions.

Paper II evaluates the use of an existing instrument primarily developed for adults, for assessment of dynamic change in adolescents. The interrater reliability was calculated from two or more raters with appropriate statistics. The reliability of change ratings and discriminability from general symptoms were also calculated statistically based on audio-recorded interviews and self-reports.

Paper III describes the development of a new therapy process rating scale. Two independent raters scored transcripts to determine interrater agreement on different items. A total of 24 items were included to primarily identify the defined “relational interventions” and explore the interaction of timing, category, and valence of the interventions.

Results:

The detailed exploration of audio-recorded therapy sessions of an adolescent girl in psychodynamic psychotherapy revealed that the therapist flexibly adjusted different psychotherapeutic techniques in the therapeutic work. The different tools of analysing seemed to pick up different aspects of the adolescent’s development. Only patient-rated questionnaires revealed the mid-treatment crises.

When evaluating the instrument for assessment of dynamic change in psychodynamic therapy with adolescents, the Psychodynamic Functioning Scales (PFS), the interrater reliability was on average good on all five subscales. The subscale *insight* was the most difficult subscale to rate reliably in both the adolescent and the adult study. The reliability of change ratings was adequate and equal for the mean value of the instrument and Global Assessment of Functioning (GAF).

The instrument for measuring relational interventions, Relational Work Scale (RWS), showed good interrater reliability for almost every item. On most items, the ratings were normally distributed for both raters.

Conclusion:

The results from the three studies in the present thesis indicate a promising way of adding knowledge to the research field of understanding psychodynamic process and measurement of change in psychotherapy. Although some of the instruments are designed to be technique-specific, the exploration of a single case revealed that the therapist adjusted and included different techniques through psychodynamic therapy with an adolescent even when adherence to the treatment manual was kept.

A single case design may contribute to provide a bridge between research and clinical practise. The lack of generalisability of a single case is nevertheless a major limitation. As for assessment of dynamic change in psychodynamic psychotherapy, general and specific potential differences between adults and adolescents regarding dynamic capacities should be discussed.

With specific tools measuring therapist's intervention with regard to timing, content and valence, the relational interventions are possible to measure with a new scale (RWS). This scale might be used in all kinds of psychotherapies.

Combining different tools in analysis seems to be fruitful in further investigations of process and change leading to more evidence-based clinical recommendations and individualized psychotherapies.

1. Introduction

1.1 Research on psychotherapy

Psychotherapy might be understood as use of psychological methods aiming at improving an individual's well-being and mental health, resolving troublesome thoughts, emotions and/or behaviours, and improve relationships and social skills (Kennair & Hagen, 2014; M. Lambert, 2013). From Freud and Breuer presenting case studies at the end of 19th century (Breuer & Freud, 1957) and Freud introducing the term “psychoanalysis” in the beginning of the 20th century (Freud, 1905), the field of psychotherapy has developed and a variety of psychotherapeutic methods exist (Norcross, VandenBos, Freedheim, & Association, 2011). Although psychotherapy research was published already in the 1920th (Bergin & Garfield, 1971), the research field has been growing and advances made in the last decades. However, although evidence for the effectiveness of psychotherapy is present (M. J. Lambert, 2013; N. Midgley, Hayes, & Cooper, 2017), conflicting views on treatment methods are indeed present.

The "Dodo Bird Verdict", first suggested in the 1930s by the American psychologist Saul Rosenzweig (Rosenzweig, 1936), proposes that the many and various forms of psychological therapy are all equally effective. This brought a discussion about “common factors” in psychotherapy (e.g. the therapeutic relationship and alliance). The dodo may still not prove right for all mental disorders, e.g. trauma-focused cognitive-behavioural therapy seem to have a better efficacy compared to other treatments for PTSD (Bisson et al., 2007; Ehlers et al., 2010). The most effective therapies should be ensured and available.

Developing new methods of treatments or improving “old” treatments might be a dilemma. An overview on meta-analyses of four decades of outcome research on psychotherapies for adult depression suggested *not* to develop new psychotherapies for depression because all new therapies appeared to be effective and the evidence indicates that all formats with human involvement are effective in all specific target groups (Pim Cuijpers, 2017). Research on outcome and process of existing psychotherapeutic methods is necessary for making recommendations for future practice to improve individual's mental health.

As for children and adolescents, the psychotherapy research field has been suffering from lagging behind compared to adults, and although the knowledge base of “what works for whom?” is gradually building up (Fonagy P, 2014), there is a funding gap in research, e.g. in the UK only 30% of the total mental health budget is put towards child and adolescent mental health (N. Midgley et al., 2018).

1.1.1 Psychotherapy outcome research

Outcome in psychotherapy research refers to short- or long-term changes that occur as a result of therapy. Eysenck's criticism of the effect of psychotherapy (Eysenck, 1952) probably inspired psychotherapists to refine their research methods. Today there are literally thousands of studies, reviews and meta-analyses showing that psychotherapy is generally effective. The general or average effects of psychotherapy are widely accepted to be significant and moderate (M. J. Lambert, 2013; Smith & Glass, 1977; Wampold, 2001). Although the effect is quite constant across most diagnostic conditions, variations in outcome seem to be influenced by patient characteristics (e.g., comorbidity, chronicity, social support) and also by therapist variables and context factors (Larry E Beutler, 2009; Larry E. Beutler, Someah, Kimpara, & Miller, 2016; Wampold, 2001).

When it comes to children and adolescents, the number of psychotherapy studies and trials have increased rapidly over the last five decades (Weisz et al., 2017). This might reflect the recognition of the unique nature of the young person relating to developmental state and context, and arising relationships in therapy.

Psychotherapy outcome research might be diagnose specific. Depression is the inclusion criteria in the adolescent study that will be presented in this thesis. Therefore, some of the relevant studies related to treatment of depression in adolescents will be mentioned. Depression is a common mental health disorder in adolescents as well as adults. The optimal treatment for adolescent depression in children and adolescents is not clear. According to the national report from Norway (Folkehelseinstituttet, 2016), it is not possible to say anything definitely about the effect of psychological therapy compared with antidepressants alone or antidepressant medication in combination with psychological therapy for children and adolescents with depression or depressive symptoms. However, substantial data implies that psychological treatments derived from different theoretical perspectives (Cognitive Behavioural Therapy (CBT), Interpersonal Psychotherapy and Short-term Psychoanalytic Psychotherapy) are efficacious and clinically effective in alleviating depressive symptoms and improving social function in the short term in at least 50% of depressed adolescents (Ian M. Goodyer et al., 2016; I. M. Goodyer et al., 2011). Not many studies are conducted on depressed adolescents to determine the effectiveness in medium terms (i.e. 12-18 months after treatment). Concerns exist regarding the efficacy and safety of the newer-generation antidepressants.

In order to improve the evidence base for effective therapies, outcome research is essential. *Efficacy studies* focus on specific therapies to study causal relationships between

treatment and outcome under optimal conditions, regardless of practicality or cost-effectiveness. Randomized controlled trials are seen as the gold standard due to focus on causality, comparing findings from a treatment group with results from a comparison or control group.

Effectiveness studies measure potential effect/change in ordinary clinical settings. A comparison group may or may not be present. Effectiveness studies have some methodological challenges, making it difficult to interpret their results (e.g. confounders and unblinded treatment conditions) - however, the inclusion criteria are less strict and the external validity higher (higher generalizability) (Möller, 2011).

1.1.1.1 Instruments to measure outcome

The psychotherapy research field altogether presents a large number of measures, rating scales and instruments to measure change and outcome (Froyd Michael J. Lambert Jeff D. Froyd, 1996), including child and adolescent psychotherapy (Johnston & Gowers, 2005). Some guidelines were given to select outcome measures (Fitzpatrick, Davey, Buxton, & Jones, 1998) with regard to e.g. feasibility, psychometric properties and comparability. Choosing the suitable outcome measure(s) may still be challenging, balancing between the instrument's validity and reliability, e.g. if the assessment tool measures what it is intended to measure. Some psychological attributes (such as relationships and emotions) are by nature more ambiguous than measures compared in other research areas. The number of existing measures disclose that a lot of researcher decide to design a new instrument or rating-scale suitable for their study.

Two aspects may organize the available outcome measures in psychotherapy:

- Who assess the outcome?
- What is being measured?

Therapist-rated outcome instruments may measure general symptoms (e.g. Global Assessment of Functioning Scale (GAF) (Aas, 2011), specific symptoms related to diagnosis (e.g. Montgomery-Åsberg Depression Rating Scale (MADRS) (Montgomery & Asberg, 1979), or personality traits (e.g. Structured Interview for DSM-IV Personality, SIDP-IV (Phohl, Blum, & Zimmermann, 1997)) .

Rating-scales assessed by the therapist or an independent expert/rater are often used in research and clinical practice. The advantage is that the rater (therapist or independent rater)

usually know the instrument. Ratings from therapists and blinded raters may differ. At least the therapist's ratings may not be coherent with the patient's ratings (Elkjaer, Mortensen, Poulsen, Kristensen, & Lau, 2012). There are other possible sources of errors when using rating-scales. The rater has to receive enough relevant information from the patient and the rater's style and personal values may influence the info and hence the data available for rating. If the overall impression of a patient influences the rating we may talk about a "halo effect". Especially in domains difficult to operationalize the training of the rater is essential, e.g. inadequate training might be a source of error in the ratings. Although clinician-rated measures are time-consuming (and hence expensive), expert judgement may have advantages over self-reports in providing relevant measures (Cousineau & Shedler, 2006).

Patient-rated outcome instruments may measure general symptoms (e.g. Symptom Checklist-90 (SCL-90) (Derogatis, 1983), interpersonal functioning (e.g. Inventory of Interpersonal Problems (IIP-64) (Alden, Wiggins, & Pincus, 1990), specific symptoms (e.g. Beck Depression Inventory (BDI) (Beck, Steer, Ball, & Ranieri, 1996) or personality traits (e.g. Structural Analyses of Social Behavior, SASB (Benjamin, Rothweiler, & Critchfield, 2006)) .

Self-reporting instruments are commonly used in research projects and clinical practice. The feasibility is high and the patient may report upon own thoughts, feelings and symptoms in a unique way. Possible sources of error with self-reports is the context, meaning the patient deliberately deliver a higher symptom score to stress their symptom load or hide symptoms for opposite reasons. The personal style of the patient influences the ratings, for instance consequently use of the middle of the scales for all questions. Complicated or ambiguous questions could make the interpretation difficult and sometimes the ratings are misplaced due to lay-out with too many items squeezed into each other. In a study of short-term psychodynamic group psychotherapy, the patient's reported effect differed from the therapist's reported effect (Elkjaer et al., 2012).

The majority of rating-scales were made for adults, and although scales for children and adolescents exist, the questions in adult scales may include jargon or expressions young people are not familiar with.

1.1.2 Psychotherapy process research

Process in psychotherapy research refers to the means by which psychotherapy produces its effects (Hardy & Llewelyn, 2015). Research on process is linked to outcome, i.e. process-outcome research. In the process of a variety of psychotherapy methods, general

factors common in all psychotherapy methods are important. Especially *alliance* is an important aspect of process-outcome research (Crits-Christoph, Gibbons, Hamilton, Ring-Kurtz, & Gallop, 2011; Falkenström, Granström, & Holmqvist, 2013; D. N. Klein et al., 2003; Webb et al., 2011). An increasing number of studies focus also on other factors that might explain change in therapy. Psychotherapy methods often involve different techniques representing different ways of structuring therapy sessions. Exploring how therapy is performed with focus on specific therapist's techniques may contribute to the evidence base of what leads to therapeutic change. The overall question is *why* the psychotherapeutic intervention works. Lambert proposed a pie-chart model of main factors explaining the variance in treatment effect (M. Lambert, 2013; Lambert, 1992) where specific techniques accounted for only 15% and common factors and client (and the world external to treatment) made the largest contribution (30% + 40%). Although other researcher would support common factors explaining more than specific factors related to techniques (e.g. homework in Cognitive Behavior Therapy) (P. Cuijpers et al., 2012; Wampold, 2001), it might not be easy to separate the therapist's technique from the relationship between the therapist and the patient, i.e. what exactly are the common factors and what are the specific factors. A suggestion of solving this is to separate the patient's experience of the therapy (common factors) and the therapist's behavior and interventions (specific factors) (Ryum, 2012).

The search for evidence of *how, for whom and under what conditions* treatment works will give a broader and more solid empirical foundation to base our treatment efforts on – in the interest of the individual's need.

In contrast to the outcome measures, few of the process measures have been used repeatedly. Process measures may be designed to fit specific research projects. The focus is on instruments primarily used in treatment of patients with diagnosable disorders. As with outcome measures, the process measures have different perspectives of observation (patient, therapist, independent expert/observer/researcher). Other dimensions for categorizing process measures suggested (Elliott, 1991) involves who is observed (client, therapist, or dyad), aspect (e.g. content, style, or quality), the unit level (e.g. segments of dialogue or whole sessions) and the sequential phase (to understand context, process, or impact).

Psychotherapy process research are exploratory (pan-theoretical and aware of many perspectives) or theory-based (testing hypotheses derived from clinical theory about how psychotherapy operates) (Orlinsky, Heinonen, & Hartmann, 2015).

To focus on specific techniques in psychotherapy research might be more reliable when we can assure the therapist(s) adhere(s) to a treatment/treatment manual (Town et al.,

2012; Trijsburg et al., 2002). The list of instruments for different techniques are long and will not be described in detail. Some instruments are complex and detailed rating system, but time-consuming (and hence expensive). Data available for process ratings are usually video- or audio-recorded sessions and/or transcription from sessions. In the following, a few examples of process measures are given:

1.1.2.1 Instruments to measure process

Clinician-rated process instruments, e.g. SASB or the Psychotherapy Process Q-set (PPQ) (Jones & Windholz, 1990), designed to record a single treatment session as the unit of observation. The Q-set comprises 100 items that capture a wide range of phenomena (pan-theoretical) and with Q-sort procedure the items are sorted into 9 piles from “least characteristic” to “most characteristic”. The later development of The Child Psychotherapy Q-Set (CPQ) (Schneider, 2004) and Adolescent Psychotherapy Q-set (APQ) (Calderon, Schneider, Target, & Midgley, 2017) were adapted from the PPQ. The instrument enables description of the psychotherapy process in a form suitable for quantitative comparison and analysis. To my knowledge there are not numerous clinician/observer-rated process rating instruments developed for or adjusted to children and adolescents.

“Common factors” instruments focusing on e.g. the therapeutic alliance (Working Alliance Inventory; WAI (Horvath & Greenberg, 1989)) with both a patient and therapist form covering the aspects on subscales Bond, Task and Goal. WAI may represent a process measure when used through therapies. When aiming at measuring the therapist empathy, the Barrett-Lennard Relationship Inventory; BLRI (Barrett-Lennard, 1962) may be used.

Micro-analytic process instruments are sequential process research, coding client and therapist responses on pre-defined categories or rating scales. Examples are the Structural Analysis of Social Behavior; (SASB) (Benjamin et al., 2006), and the Transference Work Scale; TWS (Randi Ulberg, Amlo, & Høglend, 2014). The TWS explores direct, immediate influence of therapeutic interventions on within-session processes or the effect of patient actions on the processing and planning activities of the therapist. The major limitation of these instruments are that they are more difficult to use compared to other instruments (specifically trained raters necessary) and time consuming (selecting and transcribing relevant segments from therapy sessions and then rating). However, combining sequential process research with process-outcome research can partially help filling in the process-to-outcome gap.

Having both outcome and process in mind is perhaps the most fruitful way to provide evidence to support a causal mediation and minimize the likelihood that other potential background or mediating variables do not better account for outcome (Kazdin, 2009). With the lack of proven large outcome differences among different methods of psychotherapy, using many perspectives to examine data makes sense. As the knowledge of what therapist techniques work best for the individual patients is building, more work needs to be done to relate patient (personality) characteristics to what therapist techniques are offered, how techniques are responded to by patients, and how patients experience the therapy process (Hill, 1990).

1.1.3 Operationalization

Operationalization is translating a theoretical concept or construct into a functioning and operating reality. In psychotherapy research, ordinal measurements are most common (measures of non-numeric concepts). Operationalization determines how the researcher is going to measure an emotion or concept, e.g. level of distress or problem-solving capacity. The problem is that these measurements are arbitrary. They are still needed for making the concepts under study available for statistical analysis. Operationalization defines the exact measuring method used, and allows other scientists to follow exactly the same methodology. There is a need to be concerned about how well the translation is performed. Designing new instruments involves the researcher to select definable and measurable variables. Some guidelines are available (Anna Clark & Watson, 1995; Barker & Pistrang, 2005; DeVellis, 2016) and involve numerous subtasks, e.g. review literature, assemble a pool of items potentially relevant and then reduce the pool after statistical analyses, evaluate reliability and validity and finally generate norms for a variety of respondents. How this might be done is partially a topic of this thesis.

In general, some core concepts of psychotherapy, like insight and transference, may be hard to grasp or agree on even for clinicians in the psychotherapy field. In psychotherapy with children and adolescent we often have multiple perspectives, e.g. child, parent and teacher. As brain development is phenomenal in children and adolescents (De Luca et al., 2003), most outcome measures questionnaires are designed for a narrow age range. Traditionally, transitioning age at 18 divide child and adolescent mental health services versus adult services. There is a shift in increasingly focus on early intervention in adolescence and a “research gap” with only hypothesis existing about why adolescence seem to be a risky period

for onset of mood disorders like depression and anxiety. To assess and search for helpful treatment processes leading to good outcomes, there is a need for instruments capturing the concepts we want to measure and that contribute to evidence base for measuring reliable change.

1.2 Measuring change in psychotherapy

To evaluate the effectiveness of psychological interventions, measurement instruments that can assess change is necessary. The purpose of psychotherapy is to facilitate positive change. It is of importance to investigate the impact of therapy upon lives of adults and/or children and adolescents. Therapy outcomes may provide information about the effectiveness of the therapy and supplement or add new knowledge to the evidence-based practice of therapeutic work. Studies devoted to the advancement of outcome assessment and the interest in developing the newest, improved treatment for a given disorder have so far outnumbered studies devoted to testing of mediators, moderators, or process of change (*Handbook of psychotherapy and behavior change*, 6th ed, 2013). Randomized controlled studies (RCTs) are considered the “gold standard” of research evidence and important studies in evaluating efficacy of therapy. From a philosophical point of view the assignment of numerical scores to people and quantifying patients and their attributes may be criticized. However, most psychotherapy researchers have accepted this way of measurement tradition. Statistical analysis are of little use without any form of quantification of the clinical material.

To know if an individual improves in therapy we must examine if the change is a result of attending therapy or whether the change is just due to chance. By use of statistics we may find the change to be statistical significant, and calculating the effect size would indicate the magnitude of differences between two treatment groups. Research findings cannot fully capture the complexity of what happens in the therapeutic relationship. We also should bear in mind that applying statistical methods and interpreting data might be at the expense of the individual, e.g. that the statistical change measured is valid only for the sample in the study (non-generalisability) or that the results on group-level does not necessarily pay attention to important individual differences.

The field of psychotherapy research is vital and especially growing in the field of child and adolescent psychiatry. Research findings from the past 50 years of youth psychological therapy suggest both benefits of psychological therapy, but also important directions for the future in youth therapy research and clinical practice (Pim Cuijpers, 2017; Weisz et al., 2017). Research focus is slowly shifting from adding to the list of evidence-supported therapies to

what *processes* lead to change within psychotherapy (Barber, Muran, McCarthy, & Keefe, 2013; Levitt, Pomerville, & Surace, 2016).

Psychological health is not only the absence of symptoms, but the positive presence of inner (dynamic) capacities and resources that allow people to live their lives with freedom and possibility. Symptom-oriented outcome measures commonly used in outcome studies do not attempt to assess such inner capacities.

However, we should understand what types of changes we are attempting to capture, e.g. change in symptoms or assessment for specific disorders. The primary sources to obtain information about treatment results are self-reports and clinician-rated instruments. A review of research literature identified more than 100 different measures for considering outcome (Johnston & Gowers, 2005), and as for all outcome- and process-research, it might be difficult to select which measure, instrument or rating scale to use in different circumstances. No core battery is widely accepted. A review of child self-report measures in child and adolescent mental health services (CAMHS) identified 11 measures having potential for use as outcome measures in routine practice. However, none of these measures had sufficient psychometric evidence available to demonstrate that they could reliably measure both severity and change over time (Deighton et al., 2014). There are limitations in the existing global impairment measures. Most measures are unidimensional and incorporate symptomatology into the measurement, i.e. do not differentiate what is specific for psychodynamic psychotherapy, for instance the quality of relations to close others, and the ability to think about and handle problems, as well as toleration of affects. More “fine-graded” scales are needed to measure change in psychotherapy and track the improvement during and following the therapy.

To my knowledge, there are none existing instruments designed to specifically capture dynamic capacities and assess dynamic change in psychotherapy with adolescents. Assessing change might be evaluated differently according to different perspectives. For children and adolescents there might be more perspectives on whether therapeutic change has occurred, e.g. the perspective of the young person, his/her parents or teacher. Ethical considerations are naturally equally important to keep in mind also when dealing with non-adult patients.

The symptoms themselves are not the focus when assessing change and outcome in dynamic psychotherapy. The overall question in psychotherapy research is *what works for whom and how?* Reviews of treatments for adults and adolescents exist (Fonagy, 2015; Fonagy P, 2014; Roth & Fonagy, 1996).

A *mediator* is an intervening variable that may account (statistically) for the relationship between the independent and dependent variable whereas the *mechanism* is the

reason why change occurred, i.e. the basis for the effect (Kazdin, 2007). Mediator research is difficult because a series of statistical criteria must be fulfilled (Baron & Kenny, 1986; Kraemer, Wilson, Fairburn, & Agras, 2002). A *moderator* might be easier to establish if a characteristic is to be found that influences the direction or the strength of a relationship between dependent variable (outcome) and independent variable (treatment). Moderators could help clinicians clarify which patients might be most responsive to which treatments. E.g. gender would be considered a moderator, though still be related to mechanisms and mediators, if suggesting different processes have been involved for males and females. To evaluate all potential mediators and mechanisms of change, studies should be designed to include assessments during the course of treatment. Evaluating mediators and mechanisms of therapeutic change is important to understand the change produced of different treatments, clarify diverse outcome effects from therapy, what strategies to use or what to focus on in therapy, what are the optimal conditions and components to achieve change, identify moderators, and also to understand human functioning beyond the context of therapy (Kazdin, 2007).

1.2.1 Validity

Validity refers to the credibility or believability of the research. Researches would like to prove their findings genuine. The answer depends on the amount of research support for such a relationship. A test cannot be valid unless it is reliable. A weight scale that is 2 kg off is reliable but not valid. Validity is also dependent on the measurement measuring what it was designed to measure and not something else. Different types of validity have been described (Shadish, Cook, & Campbell, 2002). There are two main aspects of validity; internal and external.

Internal validity refers to the causal relationship between the independent and dependent variable, e.g. whether the observed effect (outcome) is due to the manipulation of an independent variable (specific treatment). If there is no control group, like in naturalistic studies, it might be difficult to address the various factors that *may* have led to change. Drop-out could be different in the groups involved in an RCT and affect the internal validity.

External validity refers to the extent to which the results of a study can be generalized to other patients and settings. Naturalistic studies may therefore have higher external than internal validity; whereas the opposite is true for RCTs. Random sampling to select participants would improve the external validity of RCTs.

In relation to quantitative research methods, the types of validity of tests relates to content or criterion:

Face validity is a basic measure that would mean only that the *responders* agree on that the test measures what the *researcher* intends to measure. It reflects the subjective perception of what the assessment is trying to measure. With high face validity an individual may however manipulate their response to present a more positive/negative image of themselves (on a depression rating scale the patient may deliberately score higher on the items they “know” would alarm the therapist/researcher).

Content validity (Lawshe, 1975; Wilson, Wei, & Donald, 2012) refers to if the items in an instrument reflect the domain of meaning that is intended to be measured, e.g. items relating to all aspects of depression are present. In practise, an independent group of subject matter experts is often asked to review an assessment and compare the questions included on the assessment against a “blueprint”.

Construct validity (Salkind, 2010) refers to the extent to which a test captures a specific theoretical construct, the distinct dimension, it is intended to measure. Scores on the test would correlate with scores on other tests in accordance with theoretical predictions. There is no single method of determining the construct validity of a test, but factor analysis and correlational methods can be used.

Criterion validity (Salkind, 2010) refers to concurrent validity (does the test relate to an existing similar measure?) and predictive validity (does the test predict a criterion that will occur in the future?).

Dilemmas arising from qualitative research design is beyond the scope of the present thesis. Adding to and complementing the knowledge base from measurement-oriented research, qualitative research yields contextualized knowledge (Frost, 2011). In qualitative research, some procedures may enhance the aspects of validity (J. McLeod, 2011):

- Transparency and clarity around the way in which data were analysed
- Providing examples of themes and categories
- Disclosure of relevant aspects of the identifying and experience of the researcher(s)

- External auditing of data analysis

One major source of confounding arises from non-random patterns in the participants in a study, or within groups in the study. This can affect internal and external validity in a variety of ways, none of which are necessarily predictable.

In FEST and FEST-IT the drop-outs were/will be included in the intention-to-treat outcome analysis. There was no drop-out at 3-year follow-up evaluation in FEST. The FEST-IT study is on-going. There might be other factors than treatment, e.g. important life-events, or maturation or spontaneous remission, leading to observed change in adolescents. Raters may have more experience after repeatedly ratings through the study-period and gradually rate in a new way. This would affect the internal validity.

Regularly meetings among therapists and regularly meetings among raters were held. This should diminish the possible threat of expectations of the therapists and/or raters, e.g. favouring one treatment. In FEST and FEST-IT there is no untreated control group. A “researcher allegiance effect” is still a critical problem for all comparative studies. Another way to counteract a possible threat to external validity in FEST and FEST-IT was using both self-rating scales (GSI (SCL-90), IIP) and clinician-rated scales (GAF, PFS). Other objective ratings concerning school/work or use of health care service could give relevant information, and some of these ratings are recorded in the FEST and FEST-IT studies.

1.2.2 Reliability

Reliability refers to the degree to which scores from an instrument (a test, questionnaire or rating scale) are stable and results are consistent. A measure is said to have a high reliability if it produces similar results under consistent conditions. When constructs are not reliably measured the obtained scores will not approximate a true value in relation to the psychological variable being measured. Observed test scores are considered to be composed of true and error elements. A standard error of measurement is often presented to describe, within a level of confidence (e.g., 95 %), that a given range of test scores contains a person's true score. Reliability is generally assessed in four ways (Medicine, 2015):

- Test-retest: Consistency of test scores over time (stability, temporal consistency)
- Parallel or alternate forms: Consistency of scores across different forms of the test (stability and equivalence)

- Internal consistency: Consistency of different items intended to measure the same thing within the test (homogeneity)
- Inter-rater: Consistency of test scores among independent judges (will be described)

A measure being reliable is not necessarily valid. A test cannot be valid unless it is reliable.

1.2.3 Interrater reliability

In two of the present studies, we were interested in the degree of agreement among raters on specific measuring scales. If various raters do not agree, either the scale is defective or the raters need to be re-trained. We therefore concentrated on the Interrater reliability (IRR). IRR measures the level of agreement between two or more observers (often referred to as raters). If everyone agrees, IRR is 1 (or 100%) and if everyone disagrees, IRR is 0 (0%). Which method used to estimate IRR depends on what type of data, and the number of raters.

- Kappa statistics is commonly used for categorical (nominal) variables. A categorical variable has two or more categories, but there is no intrinsic ordering to the categories (e.g. gender, psychiatric diagnosis). Cohen's kappa (for two raters) and Fleiss' kappa (for any fixed number of raters), consider the amount of agreement that could be expected to occur through chance. A modified version of the kappa statistics, the weighted kappa, is calculated allowing one to assign different weights at the different levels (>2). Weighted kappa is the same as simple kappa when there are only two ordered categories.
- Intraclass correlations (ICC) are used for data measured on a continuous scale (e.g. GAF).

For ICC estimations, the model and type of as many as 10 forms of ICC are defined (McGraw & Wong, 1996). To choose the right form, you must know if you have consistent raters for all rates (i.e. the same raters making ratings on every subject) and if you have a sample or a population of raters. The model selection could be *one-way random-effect* (rare unless multicentre study, each subject is rated by a different set of raters randomly chosen from a population of possible raters), *two-way random-effect* (raters are randomly selected from a population of raters with similar characteristics), or *two-way mixed-effect* (selected raters are

the only raters of interest). The type of ICC is selected for single rater (“Single Measure”) or mean of k raters (Average Measures). The definitions “absolute agreement” or “consistency” refers to how important the absolute agreement of one rater’s score to another is, or to what degree one rater’s score can be equated to another rater’s score plus a systematic error. For continuous scales used in FEST and FEST-IT (e.g. PFS, GAF, GSI, BDI) we are interested in the degree of consistency between raters. Formulas for calculating ICC include mean square for residual sources of variance and mean square for error. Calculations are done using statistical software. In SPSS, ICC calculation is based on this terminology (McGraw & Wong, 1996).

1.2.4 Clinical significance

When calculating statistical significance between two groups or pre-/post-treatment, the researcher sets an acceptable level of risk that the observed change was due to chance (typically 0.05 or 0.01). The “magical 5%-number” have some limitations (Cohen, 1969; Gigerenzer, 2004). Some of the philosophical concerns about validity of outcome measurement are related to reducing the included patients in psychotherapy studies as numbers (Foster & Mash, 1999; Ponterotto, 2005).

The individual’s change during psychotherapy is important. The question might be how many points on a rating scale the individual must change for the result to make a difference in the everyday life of the individual. A way of relating to this issue is assessing statistical reliable change. The most commonly used measure is known as the Reliable Change Index (RCI) (Jacobson & Truax, 1991). This is based on the social validity idea that patients make meaningful change when that change is sufficiently large to be noticed by others (Wolf, 1978). The calculation of RCI is based on the difference between the post-treatment and pre-treatment score divided by the standard error of difference between the two test scores. The calculations are made for each subject and when greater than 1.96 it is usually regarded as unlikely ($p < .05$) to occur without any actual change, and therefore indication of the individual’s reliable change.

Formula for Reliable Change Index:

$$RCI = \frac{X_{\text{post}} - X_{\text{pre}}}{S_{\text{diff}}}$$

X_{pre} = the pretest score, X_{post} = the posttest score, S_{diff} = the standard error of the difference between the two test scores. $S_{diff} = \sqrt{2(SE_m^2)}$ and the Standard Error of Measurement $SE_m = s\sqrt{1-r_{xx}}$ where r_{xx} = reliability of instrument.

In psychotherapy research an important distinction is what is clinically significant and what is only statistically significant.

1.3 Dynamic psychotherapy

The terms “psychoanalytic” and “psychodynamic” are both used to describe psychotherapy based on psychoanalytic principles. Seven features reliably distinguished psychodynamic therapy from other therapies (Glen O. Gabbard, 2004; Shedler, 2010):

1. Focus on affect and expression of emotion.
2. Exploration of attempts to avoid distressing thoughts and feelings.
3. Identification of recurring themes and patterns.
4. Discussion of past experience (developmental focus).
5. Focus on interpersonal relations.
6. Focus on the therapy relationship.
7. Exploration of fantasy life.

Dynamic psychotherapy may be of long or short duration. A central concept of leading schools of brief therapy, often referred to as short-term psychodynamic psychotherapy (STPP), is that there should be one major focus for the therapy rather than the more traditional psychoanalytic practice of allowing the client to associate freely and discuss unconnected issues. STPP is typically defined as less than 40 treatment sessions. In brief therapy, the central focus is developed during the initial evaluation process during the first sessions, and the therapist is often more active and encourages problem-solving strategies. Psychodynamic psychotherapy offered in out-patients clinics tends to be individual, using the verbal technique or play technique. Other types of psychodynamic therapies exist, for instance group therapy, family approaches or expressive therapy forms (art/music/drama).

The treatments in the two RCTs in this thesis are psychodynamic psychotherapy. In psychodynamic psychotherapy, the focus of the work on dynamics of the patient's internal world is external realities and relationships. The focus is however also, as in traditionally analytic work, on the patient/therapist relationship. Attentiveness to unconscious phenomena

is specific to psychodynamic psychotherapy. Dynamic oriented therapists work with the assumption that individuals have motives and feelings that are formed and expressed through relations to others, and that these emotions and feelings may, consciously or unconsciously, be conflicted (e.g. love/hatred towards the same person) (Glen O Gabbard, 2014). The developmental history of relationships from childhood is important for how the individual interpret and understand his/her feelings. The quality and patterns of early relationships influence *if*, and to what degree, the individual is aware of inner conflicts, or if the feelings experienced are predominantly threatening and overwhelming. The idea of object relations was first described by Melanie Klein in the fifties (M. Klein, 1988), although there are later contributors to Object relations theory (Zachrisson, 2013).

The Quality of Object Relations scale (QOR) was later developed as an instrument of a dimension supposed to predetermine suitability for dynamic psychotherapy (Azim, Piper, Segal, Nixon, & Duncan, 1991). Other theories, like developmental psychology and attachment theory, are part of the fundament of dynamic psychotherapy.

Psychodynamic therapy might be described as promoting dynamic changes achieved by a technique of active interpretation by the therapist. Studies of change mechanisms relating to process and outcome in dynamic therapy are not numerous. They have focused on psychodynamic constructs as self-understanding (Connolly Gibbons et al., 2009; Grande, Rudolf, Oberbracht, & Pauli-Magnus, 2003; P. Hoglend, Engelstad, Sorbye, Heyerdahl, & Amlo, 1994; Johansson et al., 2010; Kivlighan Jr, Multon, & Patton, 2000), reflective functioning (K. N. Levy et al., 2006; Vermote et al., 2010) and defences (Bond & Perry, 2004; Johansen, Krebs, Svartberg, Stiles, & Holen, 2011; Kramer, Despland, Michel, Drapeau, & de Roten, 2010; Winston, Samstag, Winston, & Muran, 1994). Summarized, the literature suggest that changes in self-understanding, in accordance with psychodynamic theory, is an important part of the therapeutic process of dynamic psychotherapy - and that the change in reflective functioning and defence might be specific to dynamic psychotherapy, however, there is a need for more investigation (Crits-Christoph, Connolly Gibbons, & Mukherjee, 2013).

Although measuring mechanism variables are time-consuming (and costly), more research is needed to understand the process of change and the specific techniques leading to change in dynamic psychotherapy (Crits-Christoph et al., 2013). The dynamic capacities, e.g. relationship skills, insight and planning, may not change independently from psychopathology through therapy, and a psychometric measurement will be described in detail later.

Some factors make psychotherapy different with youths than with adults. Adolescents are dependent on their environments: their family, neighbourhood and school contexts. Most treatment of adolescents is prompted by parents or other adults who also serve as information sources to therapists. Adolescents need to undertake several developmental tasks to make a successful transition to adulthood including grow in size, sexual maturity, emotional development and thinking capacity.

The trial Improving mood with psychoanalytic and cognitive therapies (IMPACT) is to my knowledge the only high-quality, fully powered, superiority and cost-effectiveness study assessing the medium-term effects and costs of psychological treatments on maintenance of reduced depression symptoms in adolescents. IMPACT concluded with Short Term Psychoanalytic Psychotherapy (SSTP) was as effective as CBT and brief psychosocial intervention, i.e. adding psychodynamic therapy to the evidence base for treatment of depression in adolescence (Ian M. Goodyer et al., 2016). An updated review of the evidence base for psychodynamic psychotherapy for children and adolescence concludes that psychodynamic therapy in general is effective also for this age group(s) (Midgley, O’Keeffe, French, & Kennedy, 2017), but although more research is rapidly added to the field, there is still a lack of research evaluating the effectiveness of psychodynamic therapies with children and young people. The review concludes with the need of increased focus on the mechanisms underlying treatment effectiveness and the particular characteristics and circumstances of those likely to respond to treatment. This may potentially lead to new approaches in psychodynamic psychotherapy for young people and the training of psychodynamic child/adolescent psychotherapists in the future.

To investigate the mechanisms of treatment effect we need reliable instruments to measure the concepts we are interested in. When instruments already exist, we have to consider whether adjustment need to be made for the population under investigation. Although adult and youth psychotherapy share historical background and theoretical orientation, there are key differences in psychotherapy with youths versus adults. The knowledge of the effect of psychodynamic psychotherapy for young people is growing. However, to my knowledge there are however none specific instruments designed to measure psychodynamic change like insight, problem-solving capacity or tolerance for affects, in adolescents.

When instruments to measure the concepts we want to measure do not exist, or at least not in the format we need, we may review the literature and design a new instrument.

1.3.1 Transference and transference work

Transference work, or analysis of the patient-therapist relationship, is considered a core active technique in dynamic psychotherapy (Diamond, Yeomans, Clarkin, & Levy, 2008). Freud introduced the concept of transference, hypothesizing that the patient transferred onto the analyst feelings and thoughts resulting from a reactivation of previous psychological experiences. Freud first described transference in treatment in the case study “Dora” (Freud, 1905). Although Freud's treatment of Dora was disastrous according to criticist, mainly pointing out the ignorance of likely sexual abuse, the concept of transference today in psychodynamic therapy is considered to be adaptive solutions to earlier life circumstances (Gullestad & Killingmo, 2013; K. Levy & Scala, 2012). If the solutions are made fixed assumptions, the assumptions remain although life circumstances change, and the assumptions do not adapt to the real life experiences. Typically, transference could involve a situation when someone transfers their feelings that they have towards someone else onto another person, instead of the original person. In psychodynamic therapy the therapist is prepared to play the role as “another person” so that feeling, conflicts and themes arising in the therapeutic relationship may bring unconscious and fixed assumptions into direct communication and help the patient to recognize what is real in the therapeutic relationship and what are enactments influenced by transference. Transference intervention is believed to set in motion a chain of events assumed to bring about insight and dynamic change (J Ogrodniczuk & Piper, 2004).

The research on transference phenomenon in psychotherapy has shown ambiguity about which type of patients may benefit from analysis of the patient-therapist relationship (transference work; TW) in therapy and under what circumstances. Several studies have suggested that only patients with a history of good quality relationships (high Quality of Object Relations; QOR) would benefit from transference interpretations (Connolly et al., 1999; JS Ogrodniczuk & Piper, 1999; Piper et al., 1991). In a study aimed to measure the effects of TW in dynamic psychotherapy, no overall main effect of transference work was found (P. Hoglend et al., 2008). However, especially women with *difficult interpersonal relationships* (R Ulberg, Høglend, Marble, & Johansson, 2012) and patients with *personality disorders* (P. Hoglend, Dahl, Hersoug, Lorentzen, & Perry, 2011) were reported to benefit significantly more from therapy with transference work than without transference work. In the review of the empirical work on transference work, the conclusion tended to suggest that the effects of transference interventions depend on several features of the interpretations themselves, the context and not only the patient characteristics (P Hoglend & Gabbard, 2012).

An instrument for exploring the transference work in therapy sessions has been developed; the Transference Work Scale (TWS) (Randi Ulberg et al., 2014). To identify and analyse in-session effects of therapist techniques in psychodynamic psychotherapy, and empirically establish their links to outcome, adequate process scales are needed. TWS has sub-scales that rate timing, content, and valence of the transference interventions, as well as response from the patient. In TWS the transference interventions are organized in five categories. Category 1, 2, and 3 are interventions pointing at the transaction between the patient and the therapist and exploring the patient's thoughts and feelings about the therapist and the therapy. Category 4 and 5 are interventions including connections between repetitive elements in the patient's relationships with other persons out of therapy and the patient's relationship with the therapies:

1. The therapist addresses transactions in the patient-therapist relationship
2. The therapist encourages patients' exploration of thoughts and feelings about the therapy and the therapist's style and behaviour
3. The therapist encourages patients to discuss how they believe the therapist might feel or think about them
4. The therapist includes him-/herself explicitly in interpretive linking of dynamic elements (conflicts), direct manifestations of transference, and allusions to the transference
5. The therapist interprets repetitive interpersonal patterns (including relationships with parents/genetic interpretations) and links these patterns to transactions between the patient and the therapist

Research on transference work is more profound studied in adults compared to adolescents and may reflect the use of such interventions. An assumption is that interpretations based on the "here and now" aspect of transference is more immediate and vivid to adolescents because it relates current attitudes and relations outside therapy to the ongoing treatment experience itself (Swift & Wonderlich, 1990). The patient's feelings in the therapist-patient relationship might be of discomfort, but still have particular significance in work with teenagers which are in a developmental state and highly relating to issues as autonomy, dependency, and identity. Disagreements exist as to how direct "transference work" with children and adolescents should be explored. Psychotherapists sometimes find it challenging to talk about the transference to adolescents, and likewise adolescents may find it

awkward to become aware or discuss their feelings toward their therapists (Coren, 1996; Golombek & Korenblum, 1995; Sarles, 1994; Swift & Wonderlich, 1990). In a study of children suffering from anxiety, transference and extra-transference interpretations seemed to be an affective technique (Luzzi, Bardi, Ramos, & Slapak, 2015). The cognitive maturation of adolescents may allow mid-adolescents (15-18) to be better at abstract thinking and more easily adjust to a therapeutic relationship than those in early adolescents (12-14) (Ginsburg & Oppen, 1988).

Transference work is more common in the middle and late phase of dynamic therapy, when the therapeutic frame and alliance is set (Samberg & Marcus, 2005). Because transference work is perhaps not as common in brief or short-term psychodynamic psychotherapy in ordinary out-patient clinics, a similar rating tool for in-session process to be used when therapists' interventions focus on the patients' relationships *outside* therapy should be considered. In psychotherapy, interpreting the patient's relationships with other people are key components to enhance change. Those interventions might be categorized as *Transference interventions* (TI) representing transference work in the therapist-patient relationship and *Relational Interventions* (RI) when the therapist focuses on the patient's relationships outside therapy. The patient's relationships outside the therapy room, i.e. with friends, relatives or colleagues, are therefore also of interest.

1.4 Study designs

In psychotherapy research the clinical researchers should consider what design would benefit the study. Although the casual impact of the therapeutic intervention might be difficult to prove, the non-intervention factors responsible for observed change should be controlled (such as passage of time and therapist attention). The objective is to distinguish intervention effects from other factors (*Handbook of psychotherapy and behavior change, 6th ed, 2013*). Although randomized controlled trials (RCT) are commonly considered "gold standard" of research evidence, other study designs exist. Practice-based or naturalistic studies often use brief self-reports to monitor subject outcome through therapy. The limitation is the lack of level on control and how to interpret results. Designs include *descriptive designs* (e.g. case study, naturalistic observation, or survey), *correlational studies* (e.g. cohort study, case-control study), *semi-experimental designs* (e.g. field experiment, twin studies), and *experimental designs* aiming at determining causes (e.g. RCT). In (semi-)experimental studies an intervention is applied. The methodological approaches may be qualitative, quantitative, or mixed methods research design.

Quantitative research focuses on gathering numerical data and generalizing it across groups of people or to explain a particular phenomenon. Quantitative methods emphasize objective measurements and the statistical analysis of data collected using computational techniques (Babbie, 2010).

Qualitative Research is primarily exploratory and used to gain an understanding of underlying reasons, opinions, and motivations (Frost, 2011). It provides insights into the problem or helps to develop ideas or hypotheses for potential quantitative research. Common methods include individual interviews or observations. Respondents are selected, and the sample size is typically small.

Quantitative and qualitative methods were used in the work with the adult study and the adolescent study described in this thesis.

1.4.1 Single case studies

Case studies have contributed to psychotherapy theory and practice since the beginning of the 20th century. Well-known examples are Sigmund Freud's cases of "Little Hans" and "Dora". These cases in particular have been re-read and re-interpreted later with quite different interpretations (N. Midgley, 2006). This might have contributed to the negative view of case reports, with the thought of this design being unscientifically and subjectively biased. All psychotherapy researches would most likely agree on that $n=1$ is a small number. However, guidelines and protocols have been developed for therapy case study researchers (Fishman, 2001; John McLeod, 2010). Some of the criticisms of the clinical case studies are related to the data (the observations or data used may be unreliable), the analysis of the data (the validity of the analysis are hard to check), and the generalizability (even if the case studies are reliable, and analysed so that interpretation is credible, it is still not possible to generalize to other human being) (Nick Midgley, 2006).

The aim of the case study research may be related to effectiveness of therapy, theoretical understanding of the process, what the therapist do that contributed to the outcome, or to tell the story from the patient's view. Case studies could then have a hermeneutic efficacy design, or be theory-building, pragmatic or narrative. High standards of ethical good practice are especially important in case study research.

Case-based evidence represents a form of practice-based evidence and is considered to generate a unique type of knowledge with scientific and/or educational value. The field of single-cases to enhance the field of psychotherapy practice is growing, utilizing the advantages from both psychology's traditional/quantitative and alternative/qualitative

approaches (Fishman, 2001). With a combination of quantitative and qualitative measures in analysing a case, we might have a way of studying the in-session process and subjective change process without the risk of focusing only on the narratives.

1.4.2 Randomized controlled trials

Randomized controlled trials (RCTs) are experimental studies in which participants are randomly assigned to two or more groups. When an intervention is applied to one group, the efficacy of the intervention can be identified. More than one intervention or variable may be of interest. The only expected difference between the control group and the experimental group is the outcome variable being studied. Clinical evaluators should be blind to which group the patients belong. The quality of an RCT depends on an appropriate study question and study design, the prevention of systematic errors, and the use of proper analytical techniques. All of these aspects must be attended to in the planning, conductance, analysis, and reporting of RCTs (Kabisch, Ruckes, Seibert-Grafe, & Blettner, 2011). RCTs must also meet ethical and legal requirements. Although written consent from participants are common in most scientific research, a concern is that a lot of involved researchers seemed to have been aware that patients may not have fully understood what they were doing, and that the informed consent seemed little more than a ritual (Edwards, Lilford, & Hewison, 1998). In a review the proportion of participants in clinical trials who understood different components of informed consent varied from about 50-75% (Tam et al., 2015).

Both studies described in this thesis are RCTs. In psychotherapy research with long-term follow-up, a no-treatment (placebo) control group would be difficult and considered unethical to establish. The therapist cannot be blind or ignorant to the therapy they are offering (e.g. transference versus non-transference) and a double-blind experiment is therefore not possible. Although the patients in the described studies were not explained the nature of randomization, the therapists' expectations about the effectiveness might influence the therapies and the results. The evaluators in the adult and adolescent studies were blind to treatment group. However, sometimes the evaluators would have a qualified guess when interviewing the patients.

Advantages of RCTs are e.g. that good randomization will "wash out" population bias and that it is easier to blind than in other study designs. The patients or individuals are clearly identified, and the results can be analysed with well-known statistical tools.

Disadvantages could be the time (and money) required to fulfil the study included follow-ups and a volunteer bias, i.e. the patients may not represent the population the

researcher intended to study. Confounding variables may be present between the groups and the RCT may not reveal causation.

1.4.2.1 Dismantling design

The design of the FEST and FEST-IT studies is a so-called dismantling design in which a single component is added and/or varied to an existent treatment method (psychodynamic psychotherapy). The efficacy of a specific technique (transference interventions) can then be identified. A dismantling design is one type of therapy outcome study. A dismantling design is optimal to control for nonspecific factors kept constant at treatments to investigate the causal effect of a putatively active technique or intervention (Borkovec, 1993). However, child psychotherapy outcome research presents special challenges later raised by Borkovec and Miranda (Borkovec & Miranda, 1999), e.g. concerning the multitude of influences on young persons that affect their functioning and therapy outcomes. Also defining the referral problem accurately, identifying appropriate outcome measures and reliable informants of change, and considering developmental processes that influence therapeutic effectiveness, might be more challenging with children and adolescents (Donenberg, 1999).

In the research field of psychotherapy there is a need for identifying the key elements of therapeutic interventions that drive an intervention's effect. Dismantling studies are also called component studies. When one component is left out or a component is added to an existing multicomponent therapy, it is possible to examine how therapies work and identify the active ingredients of psychotherapies. For adult depression a systematic review of component studies examining the effects of psychological treatments has been published, but the currently available component studies do not have the statistical power nor the quality to draw conclusion about key ingredients of psychotherapies for adult depression (Pim Cuijpers, Cristea, Karyotaki, Reijnders, & Hollon, 2017).

2. Objectives

The main objective of this thesis was to investigate the use of different methods for measuring process and change in psychodynamic therapy with adults and adolescents. Data from two studies on psychodynamic therapy, one for adults and one for adolescents, was used to capture different aspects of psychodynamic therapy through several measurements being specific and non-specific to psychodynamic therapy. The idea of combining pan-theoretical measurements and rating-scales designed for psychodynamic psychotherapy was held high.

- With a case-study (paper I), we aimed at exploring the in-session process of a depressed adolescent in psychodynamic therapy (FEST-IT). Both quantitative and qualitative measures were used to capture the process both in terms of changes in symptomatology, interpersonal problems, and alliance.
- We aimed at evaluating a psychometric instrument for assessment of change in dynamic psychotherapy (Psychodynamic Functioning Scales, PFS) for adolescents. Data from a FEST-IT was used. The interrater reliability of five subscales was tested (paper II) as well as the reliability of change ratings and the discriminability from global functioning and subjective distress.
- To be able to explore the in-session process in non-transference and non-dynamic therapies, we aimed at developing a simple measure specially designed to identify defined categories of therapists' interventions focusing on the patients' relationships outside therapy (Relational Interventions). We explored the process in sessions in FEST including the timing, content, and valence of each intervention, as well as the response from the patient (paper III). The interrater reliability of different items from the rating-scale was tested.

3. Material and methods

3.1 The First Experimental Study of Transference-interpretations (FEST)

In this thesis data from the First Experimental Study of Transference (FEST) were used in the third paper (P. Hoglend et al., 2008). FEST is a randomized controlled trial with a dismantling design. The study is the first experimental investigation designed to measure long-term effects of transference interpretations in brief dynamic psychotherapy. The main hypothesis was that patients treated with transference interpretations would improve more than patients treated without transference interpretations (comparison group) on the two primary outcome measures, the Psychodynamic Functioning Scale and Inventory of Interpersonal Problems-Circumplex version. The second hypothesis was that patients with mature object relations and/or absence of personality disorders would benefit more from therapy with transference interpretations than from therapy without. Patients were included from 1993-2001. All follow-up evaluations were completed by 2005.

3.1.1 Patients

Hundred patients were included in the study and randomized to one of the two treatment groups. Patients were referred to the study by primary care physicians, private specialist practices, and public outpatient departments (N=122). Included patients were seeking psychotherapy for depression, anxiety, personality disorders, and interpersonal problems. There were 56 women and 44 men with age ranging from 21-57. Inclusion criteria were liberal. Patients with psychosis, bipolar illness, organic mental disorder, substance abuse, and long-term inability to work were excluded. Written informed consent was obtained from all patients. The patients were informed that the aim of the study was long-term effects of dynamic psychotherapy. There were no drop-outs in the 3-year follow-up evaluation. However, 4 patients in the comparison group dropped out of therapy before session 15.

3.1.2 Treatment

The patients were offered weekly sessions of 45 minutes for up to one year. The sessions were audio-recorded. Fifty-two patients were assigned to dynamic psychotherapy with low to moderate use of transference work (transference group). Forty-eight patients were assigned to dynamic psychotherapy without transference work (comparison group). The patients were not informed about which technique was studied or the study hypotheses. A treatment manual in Norwegian was used. Both treatment groups were exploratory in nature.

The therapists should abstain from giving advice or reassurance. For the transference group the specific techniques were prescribed to the therapists: 1) address transaction in the patient-therapist relationship; 2) encourage the patient to explore thoughts and feelings about the therapy and the therapist; 3) encourage the patient to discuss how he/she believes the therapist might feel or think about him/her; 4) include the therapist in interpretive linking to dynamic elements (conflicts) and direct manifestations of transference; 5) interpret and link repetitive interpersonal patterns to transaction between patient and therapist.

In the non-transference group, specific techniques were prescribed to the therapists for relationships excluding the therapist him-/herself: 1) address interpersonal transactions with other(s); 2) actively encourage the patient to explore thoughts and feelings about his/her relationships to other(s) including their style and behavior; 3) encourage the patient to discuss how other(s) might feel or think about him/her; 4) make interpretive linking of dynamic elements (conflicts) in the patient's relationships with other(s); 5) attempt to explore interpersonal repetitive patterns with important other(s) and with parental figures. The techniques from the non-transference group was also used in the transference-group.

Patients were assigned to one of seven therapists based on availability.

3.1.3 Therapists and evaluators

The clinical research team consisted of six psychiatrists and one clinical psychologist, all experienced in practicing psychodynamic psychotherapy. Of the seven therapists, five were men and two were women, all of whom had 10-25 years of experience in practicing psychodynamic psychotherapy. The therapists were trained for up to 4 years to enable them to provide treatment with moderate frequency of transference interpretations and treatment without such interpretations.

Each therapist treated 10-17 patients and all therapists treated patients from both groups. Therapists served as clinical evaluators of the other patients, therefore no ratings of own patients were included in statistical analyses.

3.1.4 Ethics

The Regional Ethics Committee for health region 1 in Norway approved the study protocol (FEST307/95). FEST was registered at ClinicalTrials.gov Identifier: NCT00423462. The random assignment code was kept on a separate computer belonging to the research assistant. The comparison-group was by no means a placebo-group. Therapists in both treatment groups based the therapy on general psychodynamic principles. The patients were

asked to give written consent to participate and information was given by the therapists. The patients were able to withdraw their consent at any time.

3.1.5 Assessments and outcome measures

The patients were assessed with an audio-recorded psychodynamic interview prior to randomization. In addition, the patients completed a number of self-reports. Diagnosis were determined before randomization. The axis I diagnoses, according to DSM-III-R criteria, were discussed and based on consensus of the clinicians. Axis II diagnoses were determined by the patient's therapist using Structural Clinical Interview for DSM-III-R (SCID-II).

The primary outcome measures were the Psychodynamic Functioning Scales (PFS) and the total mean score on the Inventory of Interpersonal Problems-Circumplex Version (IIP-C). The two secondary outcome measures were the Global Assessment of Functioning (GAF) and Symptom Checklist-90-R (GSI). For the PFS, three clinical raters, blind to treatment group, evaluated at baseline and follow-ups. PFS and IIP-C are designed to capture dynamic and interpersonal change. Other life events and additional treatment were recorded at follow-ups. Relevant measures used in this thesis will be described more detailed.

Figure 1. Simplified flowchart in FEST

3.1.6 Main results from FEST

No significant differences were observed between the transference group and the comparison group on the baseline variables. No differences were detected in effectiveness between therapists. No significant treatment differences for any outcome variable at any time point were detected. Patients in both treatment groups improved significantly on outcome measures. However, patients with a lifelong pattern of poor object relations showed sustained positive treatment effect of transference interpretations (P. Hoglend et al., 2008), women responded better to therapy with transference interpretations (R. Ulberg, Johansson, Marble, & Hoglend, 2009), and this was especially true for women with poor relational functioning (R. Ulberg, Marble, & Høglend, 2009). Gender and the Quality of Object Relations Scale (QOR) thus seemed to be important moderators of the effect of transference interventions.

3.2 The First Experimental Study of Transference work – In Teenagers (FEST-IT)

The First Experimental Study on Transference Work-In Teenagers (FEST-IT) is a RCT designed to explore effects of transference work in dynamic psychotherapy in adolescents with major depressive disorder (MDD). The design of FEST-IT is adjusted to depressed adolescents. With a dismantling design the efficacy of a specific technique (transference interventions) can be identified.

3.2.1 Patients

The patients were recruited among adolescents with symptoms of depression referred either to private practice or child and adolescent outpatient mental health clinics in the South-Eastern Health Region in Norway, representing mainly urban and some rural areas. The originally aim was to include one hundred adolescents aged 16-18. The study is ongoing. Among the 66 patients included in this doctoral thesis (paper II), there were 12 boys and 54 girls. All patients were attending classes in lower or upper secondary school.

Adolescents with current unipolar major depressive disorder according to Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) were included. Adolescents with generalized learning difficulties, pervasive developmental disorder, psychosis, or substance addiction were excluded. Comorbidity was expected to be frequent.

Axis I and II diagnosis were based on the Mini International Neuropsychiatric Interview (M.I.N.I.) and Structured Interview for DSM-IV Personality (SIDP-IV). Axis I diagnoses beside depression were mostly affective disorders and anxiety disorders. Almost half of the adolescents had one or more Axis II disorders – primarily depressive or avoidant

personality disorders. The distribution of mean pre-treatment scores indicated that the sample of 66 patients was a group of moderately depressed adolescents, representative of typical outpatients offered dynamic psychotherapy.

Table 1

Pre-treatment characteristic of the 66 included adolescents in paper II from FEST-IT

Total (n=66)	
	Mean (SD)
Age	17.3 (0.7)
PFS	59.7 (6.1) ^a
GAF	58.0 (6.1) ^a
IIP-C	1.36 (0.4) *
GSI	1.33 (0.5) *
BDI	28.7 (9.0)
	N (%)
Female	54 (82)
Axis I diagnosis:	
Depressive disorder	100 (100)
Social phobia	19 (29)
Panic disorder	13 (20)
General anxiety	17 (26)
More than two axis I diagnoses	17 (26)
Axis II diagnosis:	
Depressive	30 (45)
Avoidant	24 (36)
More than one axes II diagnoses	19 (29)
	17 (26)

^abased on evaluator's rating from baseline assessment

*preliminary, not available data for analysis from all 66 patients

3.2.2 Treatment

Short-term psychodynamic/psychoanalytic psychotherapy (STPP) with the STPP manual from the IMPACT study (Midgley, Rhode, Rustin, Hughes, & Cregeen, 2010) is used as the manual for the treatment. The manual combines aspects of STPP that focus principally on techniques aimed at helping young people overcome developmental problems, as well as emphasizing the role of the interpretation of unconscious conflicts, attachment theory and the concepts of internal working models. With the agreement of the adolescent, parallel work

with parents is delivered, however, not included in the data collection. Antidepressant medication may be added in severe cases according to the national guidelines in Norway.

The patients are randomized to two treatment groups. The treatment in both groups will use general psychodynamic techniques. The patients will be given 45-minute sessions. The treatment period is 28 weeks. Therapies lasting less than 12 sessions will be defined as dropouts. In the transference group, specific techniques are prescribed, including the therapist 1) addressing transaction in the patient-therapist relationship; 2) encouraging the patient to explore thoughts and feelings about the therapy and the therapist; 3) encouraging the patient to discuss how he/she believes the therapist might feel or think about him/her; 4) including the therapist in interpretive linking to dynamic elements (conflicts) and direct manifestations of transference; 5) interpret and link repetitive interpersonal patterns to transaction between patient and therapist.

In the comparison group, the non-transference group, these techniques are proscribed. In this group, the therapist will consistently use material about interpersonal relationships outside therapy as the basis for similar interventions: 1) address interpersonal transactions with other(s); 2) actively encourage the patient to explore thoughts and feelings about his/her relationships to other(s) including their style and behaviour; 3) encourage the patient to discuss how other(s) might feel or think about him/her; 4) make interpretive linking of dynamic elements (conflicts) in the patient's relationships with other(s); 5) attempt to explore interpersonal repetitive patterns with important other(s) and with parental figures. The techniques from the non-transference group was also used in the transference-group.

3.2.3 Therapists and evaluators

A total of twelve therapists are involved. All therapists have at least two years of formal education in psychodynamic psychotherapy and a 1-year program with regard to offer therapy with a moderate frequency of transference interventions (1-3 per sessions) and dynamic psychotherapy without transference work. Eight therapists are psychiatrists and four clinical psychologists. There are six men and six women. Each therapist should treat at least four patients with psychodynamic psychotherapy: two patients with and two patients without transference interventions. The therapists meet regularly for peer supervision in groups.

Four individual evaluators conduct the patient interviews at baseline (pre-treatment), at the end of therapy (post-treatment) and at 1-year follow-up.

Two independent raters without direct contact with the patients, rate the audio-recorded interview for all patients. All evaluators and raters have long clinical experience

ranging from 12-30 years. The evaluators and raters meet on regular basis for training before and during the course of the study. Meetings also involve plenary discussions after individual scorings of audio-recorded interviews.

3.2.4 Ethics

The Central Norway Regional Ethics Health Committee approved the study protocol (REK: 2011/1424 FEST-IT). The study is an adjusted replication of the adult study FEST in which patients in the two treatment groups responded, on average, equally well.

3.2.5 Assessments and outcome measures

Before randomization, each patient is diagnosed according to DSM-IV criteria (M.I.N.I.6.0.0, SIDP-IV). Based on the information derived from an interview about present symptoms and situation, background and development history, as well as from information obtained from the psychodynamic interview, the pre-treatment (baseline) assessment battery consist of self-reported questionnaires and therapist questionnaires (R. Ulberg, Hersoug, & Hoglend, 2012). The primary outcome variables are the Psychodynamic Functioning Scales (PFS). The secondary outcome measure is the Global Assessment of Functioning (GAF) Assessments would also be undertaken at four time-points using the Inventory of Interpersonal Problems–Circumplex Version (IIP–C) and the total mean score of Symptom Checklist–90 (Global Severity Index; GSI).

The randomization-codes have not yet been broken.

Figure 2. Preliminary simplified flowchart in FEST-IT

3.3 Measures used in FEST and FEST-IT

The adolescent study FEST-IT is an adjusted replication of the adult study FEST, however, the inclusion criteria and treatment manuals partly differ. General psychodynamic techniques was used in both studies for both treatment-groups (transference-group and comparison-group). In both studies the patients were assigned to the available local therapist, and the therapists treated patients from both treatment-groups. The length of treatment was weekly sessions for 1 year (FEST) versus 28 weekly sessions (FEST-IT). The outcome measures are intentionally similar. The assessment batteries have minor differences. The quality of adolescents' relationships will be a central focus of FEST-IT, and the Adolescent Relationship Scale (ARS) is part of self-reports at pre-/post-treatment as well as follow-up.

A short description of the included measurements from the papers in this thesis follows:

Outcome measures

Global Assessment of Functioning (GAF)

The GAF (DSM.3rd ed. 1987) (Aas, 2011) is a clinician-rated measure which captures symptom relief. GAF represents a numeric scale 1-100 with ten descriptive levels assigning a clinical judgment to the individual's overall functioning level over the last 2 weeks. GAF recorded values used in FEST-IT are separate scores for symptoms (GAF-S) and functioning

(GAF-F). Since GAF was an outcome measure in the adult study FEST, it was chosen also in FEST-IT instead of Children's Global Assessment Scale (CGAS).

Psychodynamic Functioning Scales (PFS)

PFS (P. Hoglend et al., 2000) were developed to capture evaluator-rated change in dynamic and interpersonal functioning. Current functioning was rated on the basis of a semi-structured dynamic interview. Six scales were included in FEST: Quality of Family Relations, Quality of Friendships, Quality of Romantic/Sexual Relationships, and Tolerance for Affects, Insight and Problem-Solving Capacity. The scale *romantic/sexual relationships* was omitted for adolescents (FEST-IT) because the included adolescents (aged 16-18) were not presumed to have developed intimate relationship patterns. The scales have the same format as the Global Assessment of Functioning, with ten descriptive levels and scale points ranging from 1 to 100. The interrater reliability for average scores of three raters in FEST were good to excellent (P. Hoglend et al., 2000). The interrater reliability in FEST-IT is described in paper II.

Symptom Checklist-90 (SCL-90)

The SCL-90 (Derogatis, 1983) is a self-report psychometric instrument (questionnaire) designed to measure severity of psychiatric psychopathology and symptom distress. It is also used in measuring the progress and outcome of psychiatric and psychological treatments or for research purposes. The SCL-90-R is normed on individuals 13 years and older. It consists of 90 items and takes 12–15 minutes to administer. The Global Severity Index (GSI) is the mean of SCL-90.

Beck Depression Inventory (BDI-II)

The BDI-II (Beck et al., 1996) is a widely used 21-item self-report inventory composed of items relating to symptoms of depression. The BDI-II is designed for individuals aged 13 and over, thus measuring the severity of depression in adolescents and adults. Since adolescent depression was the inclusion criterion in the adolescent study, BDI-II was used as a secondary outcome measure in FEST-IT.

Inventory of Interpersonal Problems-Circumplex version (IIP-C)

The IIP-C (Alden et al., 1990) is a 64-item self-report measure designed to assess interpersonal problems. The total mean score measures psychodynamic change and change in

interpersonal functioning. The circumplex model with mean and the 8 sub-sales were used in FEST. In FEST-IT the total mean score is of interest, but subscale scores are recorded.

Predictors and moderators

Quality of Object Relations Scale (QOR)

QOR (Piper et al., 1991; Piper & Duncan, 1999) was the preselected primary moderator in the study protocol in both studies. The QOR measures the patient's life long tendency to establish relationships with others from mature to primitive on an 8-point scale. The predetermined cut-off for low versus high QOR was 5. Low QOR score (1-4) indicates a history of less gratifying relationships, relational difficulties and absence of mutuality in relationships. The *quality of interpersonal relationships* and *friendships* were included in FEST-IT. The *quality of sexual intimate relationships* was included in FEST and omitted in FEST-IT.

In-session rating scales

Transference Work Scale (TWS)

The TWS is a micro-analytical tool for therapy process analyses (R. Ulberg, Amlo, Critchfield, Marble, & Hoglend, 2014; Randi Ulberg et al., 2014). TWS has 26 items and 4 sub-scales that rate timing, content, and valence of the transference interventions in the patient-therapist relationship. Five categories of transference interventions were defined. Category 1: The therapist addresses transaction in the patient-therapist relationship. Category 2: The therapist encourages patient's exploration of thoughts and feelings about the therapy and the therapist's style and behaviour. Category 3: The therapist encourages the patients to discuss how the patient believes the therapist might feel or think about them. Category 4: The therapist includes him-/herself explicitly in interpretive linking of dynamic elements (conflicts), direct manifestations of transference, and allusions to the transference. Category 5: The therapist interprets repetitive interpersonal patterns (including relationships with parents/genetic interpretations) and links these patterns to transaction between the patient and the therapist.

Examples:

Category 1: (Therapist): "You smile a little now when you're saying it to me"

Category 2: (Therapist): "What can I tell you that would help you?"

Category 3: (Therapist): “Perhaps you might have some thoughts about my possible expectations of you?”

Category 4: (Therapist): “You become annoyed when those closest to you say things like that, but you aren’t capable of mobilizing that amount of annoyance towards me because you think that this is what psychiatrists should say”

Category 5: (Patient, P): “He says he’s very fond of me and could very well have a relationship with me”. (Therapist, T): “However, you’re afraid of being let down, that he loses interest in you or something like that? You did think that I too didn’t have any interest in you” (P): Yeah, I know that logically speaking, you do care, but I felt something else. I didn’t know what. It’s just sort of a feeling I get. (T): This fear, that people that mean something to you won’t care—have you always felt that way or has it appeared lately? (P): I think I have had it for a long time, but I have actually never thought about it before.

Relational Work Scale (RWS)

The development and interrater reliability analysis of RWS are described in paper III. The RWS is a micro-analytical tool for therapy process analyses, to identify and categorize Relational Interventions (RI’s) and analyze the process between therapist and patient after an RI. RWS is modelled as in the TWS. The RWS has 24 items and sub-scales that rate timing, content, and valence of the relational interventions in psychotherapy. Five categories of relational interventions were defined, all relating to “other(s)” as everyone except the therapist (i.e. friends, relatives, colleagues). Category 1: The therapist addresses interpersonal transactions with other(s). Category 2: The therapist actively encourages the patient to explore thoughts and feelings about his/her relationships to other(s) including their style and behavior. Category 3: The therapist encourages the patient to discuss how other(s) might feel or think about the patient. Category 4: The therapist makes interpretive linking of dynamic elements (conflicts) in the patient’s relationships with other(s). Category 5: The therapist attempts to explore interpersonal repetitive patterns with important other(s) and with parental figures.

Examples

Category 1: (Therapist): “She tells you. How was that?”

Category 2: (Therapist): “How did you feel when she started to cry?”

Category 3: (Therapist): “What does he mean when he tells you it was nice that you got in contact?”

Category 4: (Therapist): “Perhaps you are afraid you are going to be like your mother?”

Category 5: (Patient): “I don’t want to be a father”. (Therapist): “If you become a father, could you end up like your father?”

Structural Analysis of Social Behaviour (SASB)

The SASB (L. S. Benjamin, 1996; Benjamin et al., 2006) is a method of fine-grained analysis of the patient-therapist interaction on a “unit-coded” micro-analytic level, incorporating all forms of interpersonal behaviour into a theoretical model. The SASB framework characterizes interpersonal behaviour according to three underlying distinctions: focus (other, self, or introjective self-treatment), degree of affiliation (hostile vs. friendly), and degree of interdependence (autonomous vs. enmeshed interpersonal behaviours). These three distinctions combine to locate a specific behaviour within the circumplex space defined by the model. The process between patient and therapist, as well as narrative content involving other figures, and the self, are included. The SASB cluster model is used in paper I.

Figure 3. Simplified Cluster version of the Structural Analysis of Social Behavior (SASB-model).

From: Benjamin, L.S. (1996). *Interpersonal Diagnosis and Treatment of Personality disorders*, second edition. N.Y.: Guilford Press (L.S. Benjamin, 1996). Reprinted with permission of Guilford Press.

Adolescent Psychotherapy Q-set (APQ)

APQ (Calderon et al., 2017) is a Q-set composed of 100 items that describe the psychotherapy treatment process for quantitative comparison and analysis. The APQ is composed of three types of items: (1) items describing young person’s emotional states, attitudes, behaviour, or

experience; (2) items reflecting the therapist's actions and attitudes; and (3) items attempting to capture the nature of the interaction of the dyad, or the climate or atmosphere of the encounter. The hundred items are revised after studying a psychotherapeutic session. Items are sorted into nine categories from extremely characteristic to extremely uncharacteristic. APQ is intended to be neutral with respect to any particular theory of psychotherapy and should permit the portrayal of a wide range of events, interventions, and processes in the psychotherapy process. The validation of APQ was conducted with ratings of audio-recorded youth psychotherapy sessions from different therapeutic approaches (Short-Term Psychoanalytic Psychotherapy and Cognitive Behavioural Therapy) (Calderon et al., 2017).

Other relevant measures used in FEST and/or FEST-IT

Feeling Word Checklist (FWC-58)

The FWC-58 (Rossberg, Hoffart, & Friis, 2003) is a self-report questionnaire comprising 58 feeling words measuring therapists' emotional reactions towards patients during a therapeutic interaction. Factor analysis in FEST revealed 4 main factors: Confident, Inadequate, Paternal and Disengaged (H. S. Dahl, Rossberg, Bogwald, Gabbard, & Hoglend, 2012). FWC-58 was used in FEST.

Feeling Word Checklist-28 (FWC)

The FWC-24 (Holmqvist & Armelius, 1994) is a self-report questionnaire comprising 24 feeling words measuring therapists' emotional reactions towards patients during a therapeutic interaction. Previous factor analysis revealed four meaningful clinical factors: Confident, Inadequate, Disengaged, and Neutral (Ulberg Randi et al., 2013). FWC-28 including four more items (HVILKE VAR DISSE?) was used in FEST-IT.

Working Alliance Inventory (WAI)

The WAI (Horvath & Greenberg, 1989) is a measure for the assessment of the therapeutic alliance. It is pan theoretical and captures three key alliance aspects on a 7-point Likert scale: (a) agreement on the tasks of therapy, (b) agreement on the goals of therapy and (c) development of an affective bond. Working Alliance Inventory-Short Revised (WAI-SR) (Hatcher & Gillaspay, 2006) is a 12-item measure for the therapist and patient respectively.

Adolescent Relationship Scale (ARS)

ARS (Anne Grete Hersoug & Ulberg, 2012) is a visual analogue scale (0-10), measures the importance of the relationships to friends, siblings and parents, as well as the experienced quality of life. ARS is a self-report measure meant to capture the current quality and reciprocity of the different relationships and the development over time with repeated measures.

Table 2. Relevant assessments administered in FEST-IT described in this thesis (Ulberg and colleagues, 2013)

Assessment points	Interviews	Measures	Self-report questionnaires	Therapist questionnaires
Baseline (0 weeks)	M.I.N.I., SIDP-IV, PI	QOR-2, PFS, GAF	BDI, IIP-C, SCL-90, ARS	MADRS
After session 3			WAI	WAI, FWC-24
After session 12			WAI, BDI, IIP-C, SCL-90, ARS	WAI, FWC-24, MADRS
After session 20			WAI, BDI	WAI, FWC-24, MADRS
Post-treatment	M.I.N.I., SIDP-IV, PI	PFS, GAF	WAI, BDI, IIP-C, SCL-90, ARS	WAI, FWC-24, MADRS
1-year follow-up	M.I.N.I., SIDP-IV, PI, MADRS	PFS, GAF	BDI, IIP-C, SCL-90, ARS	

ARS, adolescent relationship scale; BDI, Beck Depression Inventory; FWC-24, Feeling Word Checklist-24; GAF, Global Assessment of Functioning; IIP-C; Inventory of Interpersonal Problems-Circumplex Version; MADRS, Montgomery-Aasberg Depression Rating Scale; M.I.N.I., Mini-International Neuropsychiatric Interview; PI, psychodynamic interview; PFS, Psychodynamic Functioning Scale; QOR-2, Quality of Object Relation Score; SCL-90, Symptom Checklist-90; SIDP-IV, Structured Interview for DSM-IV Personality; WAI, Working Alliance Inventory.

3.4 Some statistical considerations

Statistics are needed when a researcher has formed a hypothesis about the outcome of his/her study. The null hypothesis (H_0), which is a statistical hypothesis that states there will be no difference between observed and expected data, is proposed before completing a study. The H_0 can be supported by data or disproved in favour of an alternative hypothesis. In the adult study FEST, the primary hypothesis was that the transference group would have a more favourable course over the whole study period compared to the comparison group. The hypothesis was rejected when the statistical analysis showed no significant treatment differences between the two groups (P. Hoglend et al., 2006; P. Hoglend et al., 2008). However, in the moderator analyses, women with poor relational functioning and men with good relational functioning showed sustained different treatment-response to transference interpretations (R Ulberg et al., 2009).

Testing hypothesis involves identify null and alternative hypotheses. Then determine the appropriate test statistic and its distribution under the assumption that the null hypothesis is true. The significance level must be specified, and test statistics calculated from the data.

Then the “magical” P-value should be investigated. The P value is defined as the probability under the assumption of no effect or no difference (null hypothesis), of obtaining a result equal to or more extreme than what was observed. The threshold value, often $P < 0.05$, is arbitrary. Factors influencing the P-value are e.g. effect size, size of sample and spread of data. Larger samples are more likely to find a statistical difference than a small sample size, and in small samples important and genuine differences might be missed. Statistical significant tests are vulnerable to type I error (incorrect rejection of a true null hypothesis, "false positive"). Effect sizes indicate the magnitude of differences between groups and were calculated at planning stage of the adult and adolescent studies described in this thesis to find the sample size required for sufficient power of the study.

Confidence intervals (CI) often make it easier to determine whether a finding has any clinical importance. If 95% CIs does not include the null value (usually zero for difference in means,) null hypothesis is rejected at 0.05 levels.

The focus in this thesis is the interrater reliability, analysed with Kappa and ICC. For the development of a therapy process rating scale, transcribed segments from the audio-recorded therapies of different patients were used (FEST). Audio-recorded evaluations (interviews) were the source of statistical material yet ready for analysis in the adolescent study. There are some obvious limitations when new rating scales are developed and investigated by the very same research group that defined the interventions and some of the researchers participating in both studies as therapists, raters and/or researcher. Some of the in-session process-rating scales are based on short segments (10-min) of therapies and cannot be generalized. The raters used were independent. In the adult study, a large group of potential raters were present at regularly plenary sessions for ratings of interviews and discussion of the results after individual ratings. Two fixed raters were used for the studies of interrater reliability of an in-session rating scales from the adult study (Paper III) and the assessment of dynamic change in the adolescent study (Paper II). In the adolescent study, FEST-IT, a total of four individual evaluators conducted the patient interviews at baseline (pre-treatment) and at the end of therapy (post-treatment). These ratings were also included in some of the analysis (in addition to the two fixed raters) although the evaluators assessed a variable number of subjects. The evaluators rated directly from interviews and the fixed raters rated from the audio-recorded interviews.

The paired sample t-test is a statistical procedure used to determine whether the mean difference between two sets of observations is zero. When evaluating the rating scale in the adolescent study, each subject was measured twice, resulting in pairs of observations.

When evaluating the rating scale with adolescents, we added also the *Reliable Change Index* (RCI) to estimate if the individual change was considered reliable, or unlikely to be a product of measurement error:

$$RCI = \frac{X_{\text{post}} - X_{\text{pre}}}{S_{\text{diff}}}$$

Where X_{pre} = the pretest score, X_{post} = the posttest score, S_{diff} = the standard error of the difference between the two test scores. $S_{\text{diff}} = \sqrt{S(\text{SE}_m^2)}$ and the Standard Error of the Measurement $\text{SE}_m = s\sqrt{1 - r_{xx}}$ where r_{xx} = reliability coefficient of the instrument.

The use of RCI is frequently used for examining improvement in psychotherapy (Moleiro & Beutler, 2009; Ogles, Lunnen, & Bonesteel, 2001).

4. Summary of results

4.1 Paper I

Exploring in-session process with qualitative and quantitative methods in psychotherapy with an adolescent.

The purpose of this paper was to explore the therapist's action, the patient's response and the interaction between the therapist and patient through a psychodynamic therapy with an adolescent. Research on therapy process in adolescents is scarce. With a good-outcome case from FEST-IT, i.e. time-limited psychodynamic psychotherapy for adolescents with depression, we explored the therapy of a depressed girl with a mid-treatment crisis. The assessment battery from FEST-IT was supplemented with 3 different rating-scales for analysing in-session relational process; the Structural Analysis of Social Behaviour (SASB), the Adolescent Psychotherapy Q-set (APQ) and the Transference Work Scale (TWS). The patient quite surprisingly became worse with self-harming behaviour at mid-treatment. However, only self-reported questionnaires revealed an increase in interpersonal problems and general symptoms at mid-treatment. The depression ratings, both clinician-rated (MADRS) and self-reported (BDI-II), were at mid-treatment decreasing from baseline values. Comorbidity was not present, i.e. no other Axis I diagnosis besides depression and no personality disorder (Axis II) was present at baseline. Possible differences between adults and adolescents in therapy were discussed. Depression and self-harm in adolescence may represent a developmental crisis. Adolescents need to undertake developmental tasks, e.g. physical growth, sexual and intellectual maturity and renegotiation of relationships to close others. Adolescents in therapy might experience the dilemma of trying to keep their parents out of the therapy room and still be occupied with renegotiating the relationship to their parents. Adolescents are dependent on their environment with e.g. family and school. Important relations explored in this case included relationships with the mother, the father, her boyfriend and with friends.

The detailed exploration of audio-recorded therapy sessions rated with SASB made the therapy process possible to be visually illustrated (figure 4). In the beginning of the therapy the adolescent girl was blaming herself and on the left-side of the circumplex. A shift towards a possible healthier position of self-control was visible before a relapse to neglect and self-blame at the time of a mid-treatment crisis. At the end of therapy a steady shift towards trust/self-protect and disclose/self-affirm was made.

A variety of rating-scales and self-reports revealed that the therapist flexibly adjusted different psychotherapeutic techniques in the therapeutic work. In clinical practise the therapists may integrate different techniques without compromising the treatment manual. A single case design may contribute to providing a bridge between research and clinical practise. The generalisability of a single case is nevertheless a major limitation.

Figure 4. In-session process change during therapy (Paper I) rated with SASB

Visualising the therapy process with SASB-clusters. At mid-treatment there seemed to be a relapse to the clusters Self-neglect and Self-blame.

4.2 Paper II

Assessment of Dynamic Change in Psychotherapy with Adolescents.

In this paper a specific outcome-measure related to dynamic functioning in adolescents is evaluated and the interrater-reliability tested. Few measure of dynamic change exist. Especially for children and adolescents, outcome measures seldom have sufficient psychometric evidence to prove themselves as reliable measures in severity of mental health problems and psychodynamic change over time. Psychodynamic Functioning Scales (PFS) were developed to assess change in adults (in FEST), consistent with the concepts of dynamic psychotherapy. The “dynamic capacity” of patients differs from general symptom-measures and global assessment of functioning. PFS is used for the first time as measure of change in adolescent psychotherapy in FEST-IT. PFS has not previously been reliability-tested for adolescents. PFS consist of 6 subscales:

- Quality of family relations
- Quality of friendships
- Romantic/sexual relationships
- Tolerance for affects
- Insight
- Problem solving and adaptive capacity

The relational scales, *quality of family relations* and *quality of friendships* are based on evaluating the degree of mutuality and adequacy of the commitment in relationships, the ability to take other’s perspective, to describe close others across an external and internal dimension, feeling of being needed and a sense of belonging and the capacity to reconcile parent’s or friends’ shortcomings and make the best of the relationship. The *tolerance for affects* covers the ability to experience, differentiate and express various affects verbally and nonverbally, and to what degree disappointments lead to symptoms like avoidance, anxiety, depression or restrictions of goals. *Insight* covers mainly cognitive understanding of the main dynamics of inner conflicts, related inter-personal patterns and connection to the past. Insight also includes the ability to describe and understand own vulnerability and reactions to stress. The *problem-solving capacity* covers the ability to handle any difficult situation without developing symptoms, avoidance or inadequate actions. Self-observation, planning, ability to explore new areas and enjoy recreation and pursue meaningful goals are parts of this scale.

The *romantic/sexual relationships* involve also the capacity to establish long-term relationships characterized by love, trust, reciprocal mature dependency and active, flexible sexual pleasure. This scale was omitted in the adolescent study. Adolescents 16-18 years are not expected to yet have developed intimate relationship patterns.

Based on audio-recorded semi-structured psychodynamic interviews, raters with different clinical background rated all the available interviews at pre-treatment (n = 66) and post-treatment (n = 30) using PFS. Interrater reliability, the reliability of change ratings and the discriminability from general symptoms were calculated in SPSS.

The interrater reliability of two fixed raters were on average good on all five subscales of PFS. The subscale *insight* was the most difficult to rate reliably. When we included the evaluator for each case in the reliability-testing, the reliability increased. However, the evaluators represent 4 different clinicians/researchers and the interrater-reliability-testing of evaluators is not reported. General and specific potential differences between adults and adolescents are discussed, including comments on the subscale *romantic/sexual relationships* that was omitted in the adolescent study. The dynamic subscales (*tolerance for affects*, *insight* and *problem-solving capacity*) were rated with lower reliability than the relational subscales (*family* and *friends*) for adolescents. This might reflect on the developing mental and dynamic capacity relating to the adolescent brain. The reliability of change ratings was adequate and equal for the mean value of PFS and Global Assessment of Functioning (GAF). However, the statistics included only the adolescents available for analyses and may only give an indication of what the main results from FEST-IT will look like.

Table 3. Pre-treatment interrater reliability estimates (intraclass correlations; ICC), with 95% confidence intervals (CI), for 2 raters

PFS pre-treatment (n=66)		
Scale	ICC	95% CI
Family	0.82	0.71-0.90
Friendships	0.75	0.59-0.84
Tolerance for affect	0.69	0.49-0.81
Insight	0.61	0.36-0.76
Problem-solving capacity	0.70	0.50-0.81
PFS mean	0.81	0.69-0.89
GAF (n=47)	0.80	0.65-0.89

Note: In the paper the interrater estimates for single raters randomly drawn from a group of six (4 evaluator and 2 raters) are reported. The interrater reliability of the two raters are higher, i.e. excellent reliability on the relational scales, *family* and *friendship*, and good reliability on the dynamic subscales, tolerance for *affects*, *insight*, and *problem-solving* scale. The reliability was excellent also on the GAF.

4.3 Paper III

Relational interventions in psychotherapy: development of a therapy process rating scale.

The aim of the third paper was to develop a rating-scale for in-session process to be used when therapists' interventions focus on the patients' relationships outside therapy (e.g. the non-transference group in FEST and FEST-IT). There is a need to develop more methods to reveal what kind of treatment and what treatment techniques work best for different patients and how the various psychotherapy modes work to promote the best possible treatment effect (outcome) for each patient. It has proved difficult to find ways to examine the in-session process and interventions contributing to the patient-therapist interaction. Understanding the process of therapy and the impact on outcome will increase our ability to make treatments more effective, and to individualize them. Relational interventions are used in different psychotherapy models. The therapy process rating scale Relational Work Scale (RWS) helps identify relational interventions and explore the interaction of timing, category, and valence of relational work in psychotherapies without being restricted to any specific theory model.

The RWS is based on a preceding rating-scale on transference interventions by the same research group in FEST. The RWS in this paper rely on transcribed segments from the middle of a mid-therapy session of 20 patients from FEST. The segments were scored by two independent raters trained in different psychoanalytic/psychodynamic institutes and with dissimilar work experience. Each turn of talk in the dialogue between the patient and the therapist was given consecutive index numbers in the transcript. Items in RWS were scored using Likert scales; 0, 1 (not at all, low degree) to 4 (high degree). Items identifying presence or absence of the five categories of RI (Item 8–12) were rated with “Yes” or “No”. A total of 24 items on the RWS are presented.

The IRR was calculated with Kappa-statistics for category scores and with the Intra Class Correlation (ICC) for ordinal scores. The interrater agreement for nearly all the RWS items in the present study was good to high. RWS is a focused and short process measure and is seemingly not too time consuming. Together with Transference Work Scale (TWS), RWS may be used for creating datasets from a larger number of sessions or segments of sessions.

So far, the process-rating-scale has not been used in other studies. As friends, relatives and other persons outside the therapy-room are especially important for adolescents, by nature in separation and rebuilding relational phase, the focus on relational interventions might

deliver more knowledge to the research on *process of change* in adolescent therapies. The preceding Transference Work Scale (TWS) is used in Paper I.

5. Discussion

5.1 Discussion of the main results

The depressed teenage girl (Paper I) recovered from depression and her capacity to deal with the challenges of everyday-situations and relationships increased when measured with the Psychodynamic Functioning Scales. What mediated positive change and how did change come about? A *mediator* is an intervening variable that may account for the relationship between the independent (treatment) and dependent (outcome) variable, but something that mediates change may not explain the process of how change came about. The *mechanism* is the reason why change occurred or how change came about (Kazdin, 2007). A *moderator* is a characteristic that influences the direction or magnitude of the relationship between the independent and the dependent variable. If suggesting different processes have been involved for males and females, gender would be considered a moderator. One hypothesis in FEST-IT is that female adolescents have better treatment effect of transference work than male adolescents do. FEST-IT can be regarded as translation research where results from FEST (adults) are implemented in FEST-IT (adolescents). Gender differences were reported from FEST-IT, i.e. women with poor relational functioning responded better to therapy with transference interpretations (R. Ulberg et al., 2009). The good-outcome of the single case from FEST-IT, an adolescent girl with poor relational functioning (low QOR) given transference interventions during therapy, would fit the hypothesis of female gender and poor relational functioning predicting outcome. Since the randomization code was not to be broken, analysis on gender differences related to transference work in therapy has not yet been performed statistically in FEST-IT. We cannot generalize the result.

Although not all participants from FEST-IT are included in the evaluation of Psychodynamic Functioning Scales (PFS, Paper II), the gender imbalance seem far larger in the adolescent group compared to adults in FEST-IT. This observation complies with the general data from outpatient clinics for children and adolescents. Adolescent girls outnumber the boys in seeking help for internalizing disorders. In a review of 18 trend studies (year 1970-2013) in 10 countries, European countries and Canada, more than half of the studies reported increased reported symptoms of mental health problems for adolescents (Sletten & Bakken, 2016). In several of the studies the reported increase is valid for girls only. Interestingly, a Danish study found that parents and teachers rated children (11-16) significantly lower in problem score in 2010 than in 1996, whereas the young rated themselves higher than previously (Henriksen, Nielsen, & Bilenberg, 2012). Discussions and

ideas about subjective symptom load and psychopathology might be fruitful. Adolescents are dependent on their environments; family, peers, neighbourhood and school. A survey from Norway found remarkably higher depressive scores by adolescents in the capital Oslo (Ung i Oslo) compared to the rest of the country (Ung i Norge) (Andersen & Bakken, 2015). Still, there were some instrumental differences in design and differences in age-group, timing and response rate that may explain some of the regional differences. Adolescents in FEST-IT were recruited from Oslo and rural areas.

Equal for all adolescents worldwide is the need to undertake several developmental tasks to make a successful transition to adulthood. These tasks may include general growth, sexual maturity, emotional development and thinking capacity. Process research into psychotherapies with adult patients is relatively prolific (Kennedy & Midgley, 2007). There is, however, a gap in current knowledge about the processes that take place in psychotherapy with adolescents. This may be due in part to the paucity of tools designed to quantify adolescent psychotherapy process. Designing age customized process- and treatment measures may help evaluating psychotherapy for children and adolescents. The outcome domains relevant for adults may not serve as the best domains for youth psychotherapy, and mediators and moderator variables might differ.

The process scale developed to identify and analyse relational interventions in psychotherapy (Paper III) has comparable items as the previously published Transference Work Scale. The scales are in nature pan-theoretic, but the terms on timing of therapist interventions, content, and the interaction between patient and therapist are based on the interventions used and described in the manual for the adult study (FEST). Transference interventions might however also be non-verbal and not caught in transcribed dialogues. In the case study (Paper I), transference interventions of category 1 and 2 (according to the Transference Work Scale) are in majority, although far from present as often as 1-3 per session suggested by the manual. Interventions of categories 3 and 4 are present mainly before mid-treatment. After the self-harming episode at mid-treatment, the transference interventions are less frequent. The effectiveness of transference interpretations has not yet been assessed in meta-analytic studies. In another study of depressed adolescents, the results show that transference tended to be discussed between the therapist and patient through the use of metaphors (E. Della Rosa, 2016). When discussing with clinicians in outpatient (or inpatient) departments, my personal impression is that transference interpretations, especially those referred to as category 4 and 5 in Transference Work Scale (TWS), are not commonly used in therapy with adolescents. The technique is in general often more typically found to be used in

the later phase of therapy also with adults, as the more profound interpretations or interventions rely on a thorough knowledge of the patient's history, i.e. drawing the lines from the past (relationships) to the present (relationships). In FEST the use of transference intervention was explored in connection to the patient's previous history of relationships (poor/low QOR versus good/high QOR). In another study of adolescents exploring whether it is helpful to link discussions of endings to the interpretation of transference, the results indicate other patient characteristics that make the young emphasize or diminish the importance of their relationship to their therapist (Elena Della Rosa & Midgley, 2017). It would be interesting to explore whether these differences are connected, i.e. that adolescents with poor relational history are "dramatizing" or "down-playing" the therapeutic relationship.

Relational interventions might be more easily depicted by audio-recordings than transference in the therapist-patient relationship. Talking about relations outside the therapy room may serve as more benign or non-confronting to the patients. When exploring the in-session process with a teenage girl (Paper I), the treatment group was revealed and might easily have led to ideas about the impact of transference work. Comparing to the other in-session process-rating scales, APQ and SASB, the TWS complied with the other measures and still seemed to pick up on other aspects. Although the designed measure of the transference-group (TWS) and non-transference group (RWS) in FEST differ in nature of relationship foci, the idea of combining them might not be as unscientifically as it might seem. If transference work, working in the patient-therapist relationship, is less present in adolescent psychotherapy, relational interventions (outside-therapy relationships) might be a better way of exploring the process and gather more scientific ideas about what the therapists are actually doing during sessions in good-outcome therapies. Another interesting question is what the therapists do in non-successful therapies. Meta-analyses of experimental design research suggest that 75% of children and young people who attend therapy benefit from it (Weisz, Sandler, Durlak, & Anton, 2005). Still, the drop-out rates are higher with adolescents than with adults. In a study of predicting dropout in adolescents receiving therapy for depression, 37% of the adolescents dropped out of therapy and therapy did not even start in about 10% (O'Keeffe et al., 2017). We cannot know if the adolescents dropping out of therapy are getting worse in therapy or they drop out because they were feeling better or for some other reason. Clinicians rarely accurately predict who will not benefit from psychotherapy, but identifying clients at risk for treatment failure has been demonstrated (Hannan et al., 2005). As for children and adolescents suffering from depression, youth age is considered to be a moderator, however not evident in all studies. Older children (adolescents)

may be better able cognitively to grasp concepts in therapy than younger children. Mediators have been challenging to agree on in studies of children and adolescent suffering from depression, and future research will be needed to identify mediators of change (*Handbook of psychotherapy and behavior change*, 6th ed, 2013).

Paper II revealed that the subscale *insight* of Psychodynamic Functioning Scales was the most difficult scale to rate reliably. Dynamic insight is a measure of psychological functioning related to subjective interpretation of symptoms. The subjective symptoms may differ from symptoms reported from other informants (parents, teacher, others). Also, a discussion about the concept of insight might be necessary. In relation to psychoanalytic theory, insight is a dimension covering the ability to account for most inner conflicts, related problems and repetitive behaviours and connections to earlier experience. The subscales *insight* and *problem-solving capacity* were rated with only fair to good reliability. The latter subscale covers the ability to flexibly handle any difficult situation without developing symptoms, avoidance or inadequate actions. This might not be the stereotype of an adolescent even without mental health problems. However, the mechanisms and the differentiation on a scale may be relevant also for young people in therapy. Research suggest that changes in insight, or self-understanding, are an important part of, and predictive of change in symptoms, in the therapeutic process of psychodynamic psychotherapy (Gibbons et al., 2009; P. Hoglend et al., 1994; Johansson et al., 2010; Kivlighan Jr et al., 2000). Although these two subscales seemed harder to score than the relational scales (family and friendships), the changes from pre- to post-treatment were larger. Research on mechanism variables are costly and time-consuming. However, more research is needed to understand the processes of change for specific patient samples. The measurement tools should be designed to capture statistically and clinically significant changes in psychotherapy. In paper II the subscale *romantic/sexual relationships* is omitted in the adolescent study. Intimate relationships are of course important to adolescents, but few adolescents have already established a pattern of intimate relationships. Discussions about the ratings of the dynamic subscales and the “missing” intimate scale were present during the plenary meetings.

Therapists' emotional reactions to their patients (countertransference) seem to matter when relating to the treatment process and outcome of psychotherapy. For the Feeling Word Checklist (FWC), previous factor analyses did not find the exact same categories for the observed items. For FEST-IT the four clinical factors from FWC-24 were named: Confident, Inadequate, Disengaged, and Neutral. In another study of FWC-24, heterogeneous in psychotherapy method, diagnoses and age-group (Lindqvist et al., 2017), the items were

found to be best represented by the four factors: Engaged, Inadequate, Relaxed, and Moved. It is to believe that some factors are more therapist dependent and some more patient dependent. From FEST was reported an adverse impact on the effect of transference work for all patients when the therapist had a heightened level of disengaged feelings over a treatment period, especially with patients with a history of poor relationships (H. J. Dahl et al., 2017). In the case from FEST-IT (Paper I), the therapist's reported feelings were categorized as *inadequate* in the beginning of therapy and *confident* at every assessment point later during therapy. The therapist and the adolescent patient both scored the alliance higher in the beginning of therapy and lower at mid-treatment before gradually returning to the initial ratings at the end of therapy. The therapist seemed to be confident although the working alliance rated with Working Alliance Inventory (WAI) was dropping, however not rupturing. Patient's ratings of alliance and therapist's self-evaluations are independent measures. In a multisite project on process and outcome of psychotherapy, therapists' experience, training, skill, and progress as therapists did not have any significant impact on alliance as rated by patients (A. G. Hersoug, Hoglend, Monsen, & Havik, 2001). A confident therapist at mid-treatment, especially when non-confident at the beginning of therapy, would indicate a therapist experienced in fluctuations through therapies with ups and downs that might be typical for adolescent therapies and also in general for adult psychotherapy with sudden gains and sudden losses during the course of psychotherapy (Lutz et al., 2013).

Many parental characteristics influence aspects of the adolescents' regulation of emotions, including parents' own attachment styles and family experiences, levels of stress and social support, and parental mental health (Morris, Silk, Steinberg, Myers, & Robinson, 2007). Influences from parents and peers are important and contribute to what is described as the individual's "dynamic capacity". In absence of existing instruments designed to assess dynamic change in adolescent psychodynamic therapy, adjusting a measure designed for adults seems to work adequately with the PFS. During reliability-testing it was however revealed how different clinicians may interpret the data (audio-recording) and the question about what was *not* on the recording frequently came up. The interrater reliability of the evaluators are not reported in the article. Preliminary analysis indicate the reliability to be lower when including evaluator rating less than 10 interviews.

The statistical approach when working with psychometric measures may contrast the day-to-day work of the clinicians as therapist. The integration of research findings in clinical practice is however a way of bridging the perspectives or narrowing the gap. The assessment with Psychodynamic Functioning Scale described in this thesis should be possible to use for

therapists treating adults or adolescents with psychodynamic therapy. The micro-analytic tools require more training and is more time-consuming. When efficacious therapeutic interventions are identified, the integration of research findings with clinical practice may be possible. After all, the psychodynamic psychotherapy with an adolescent (Paper I), revealed that the focus assessed with APQ form whole therapy sessions were not extremely typical for dynamic techniques although the therapist adhered to the treatment manual.

5.1.1 Methodological discussion

Most measurements in psychotherapy research are subject to some sort of error (John & Benet-Martinez, 2000; J. McLeod, 2015). In FEST-IT and FEST the *treatment outcome*, as dependent variable, is operationalized with a vast number of self-reports and interviews. The dependent variable *transference work* is operationalized as the therapist's focus on the relationship between the therapist and the patient. Some concepts, e.g. insight, alliance or countertransference, are challenging to translate into countable units with reliable measures.

In Paper I, two rating scales for analysing in-session process were used. Because the micro-analytical measures were time-consuming and applied only to limited segments of the therapy, another measure for rating whole therapy sessions was added (APQ). Combining several rating scales seemed to expose different aspects of the therapy process. However, the search for even more suitable measures might have continued. A characteristics of recent research in therapy has been a search for ways of combining information that has been collected using different methods (J. McLeod, 2015). Although a growing consensus on how to collect, analyse and report case material, the selection of a specific case is by nature biased. The questionnaires, i.e. self-reports, were administered at pre-selected time-points for research purposes. The therapists were not aware of the patient's ratings although the ratings in the reported case might have given the therapist important cues to the therapy process.

In Paper II, the evaluators and raters should be blind to treatment group. However, the audio-recorded interviews might reveal the treatment group. The selection of only two fixed raters was made based on the highly variable number of ratings from the rest of the group of raters (including the evaluators). When plenary meetings concerning interrater reliability are more numerous (individual ratings followed by plenary discussion), the interrater reliability will be calculated with different statistical methods, e.g. ICC Two-way random effect, consistency - when each subject is assessed by each rater and the raters have been randomly selected.

In Paper III the Kappa statistics was used for category scores and the ICC for ordinal scores. Since there were only two raters the model of ICC was in fact Two-way mixed effects. The interrater reliability should be tested on transcripts from more therapies and/or research groups.

5.2 Strengths and limitations of the present study

The results are relating to the adult study FEST and the adolescent study FEST-IT. Both studies are randomized controlled trials of dynamic psychotherapy designed to determine the long-term impact of moderate level of transference interventions (1-3 per session). FEST was completed in 2005 with follow-up evaluations and have resulted in several dissertations and numerous published articles in a variety of international journals. FEST-IT is an ongoing study and the main results will be published when the study is completed and the data-set is ready for the main analysis.

Exploring a single case is a case-based knowledge rather opposite to the knowledge based on group experiments like randomised controlled trials. It is however closer to the therapy knowledge of practitioners and clinicians in real-life therapy with adults and adolescents. The exploration of the in-session process in the therapy with a depressed adolescent gave rich narrative material that complied with the quantitative material arriving from the controlled design and also the assessment battery of the therapy setting. Results from a case can however never be generalized.

The sample of adolescents suffering from depression in the data used from FEST-IT in this thesis reflects gender differences known from out-patient clinics (82 % girls). This does however not reflect the gender balance from the adult study and might bias the results from evaluating the Psychodynamic Functioning Scales for adolescents.

The data used in assessment and in developing a new rating-scale (RWS) are all from the related studies FEST and FEST-IT. The items in RWS were defined by interventions prescribed in the treatment manual developed within the same research group.

The on-going adolescent study (FEST-IT) did not allow the randomization-codes to be broken. The code was however broken for the patient described in the case-study to enable the in-depth in-session exploration.

The ratings used for micro-analytic analysis were performed on short segments from therapies (usually 10-min segments); relational interventions (RWS), transference interventions (TWS) and the SASB.

6. Conclusion

This thesis aims at contributing to the measurement of psychodynamic process and change in psychotherapy with adults and adolescents.

“The scientific study of mechanisms of change is complex. A given treatment might work for multiple reasons. Two patients in the same treatment conceivably could respond for different reasons. The best patient care will come from ensuring that the optimal variation of treatment is provided. Understanding mechanisms of treatment is the path toward improved treatment (Kazdin, 2007)”.

When applying quantitative and qualitative methods, these methods may be complementary in order to enhance our understanding of what works in psychodynamic therapy. A contribution to the shortcoming of instruments measuring specific therapy techniques were made; the Relational Work Scale (RWS) to be used in pan-theoretical psychotherapies, and the Psychodynamic Functioning Scales (PFS) for assessment of dynamic change in psychodynamic therapies with adolescents were evaluated.

The way of exploring a case with a combination of different and pan-theoretical in-session rating-scales on here and now interventions (TWS and SASB), and whole sessions (APQ), as well as clinician-rated measures and a variety of self-reports, may add to the knowledge of integration of different psychotherapeutic techniques without compromising the treatment modality.

The instruments designed and evaluated in this thesis would hopefully add to the list of measures to be used in clinical practice and/or research to build on the knowledgebase of what works – and for whom (Fonagy P, 2014; Roth & Fonagy, 2005).

7. Clinical implications and future research

7.1 Implications for clinical practice

According to the Norwegian Institute of Public Health, at any given time, one in 20 Norwegian children have so severe depressive symptoms that they could be diagnosed with depression (Folkehelseinstituttet, 2016). In our sample of adolescents, the girls outnumbered the boys. The latest report from Norway is coherent with findings in other countries; a worrisome increase in adolescent girls reporting mental health problems and predominantly depressive symptoms (FHI, 2018). In a cross-sectional study of outpatient clinic adolescent patients, the girls were overall more severely ill and personality disorders were more prevalent in girls compared to boys (Korsgaard, Torgersen, Wentzel-Larsen, & Ulberg, 2016). How to monitor and pay attention to comorbidity is not always clear in outpatient clinics where “New Public Management” and diagnose-related groups are related to income. Investment in adolescent mental health is beneficial for the present and the future (Patton et al., 2016).Bad

The integration of different technique-specific and pan-theoretical tools enriched the understanding of a single adolescent therapy and may be used more in therapy and training of therapists to let research and research instruments make better sense and be helpful in clinical work.

The Psychodynamic Functioning Scales are made more assessable to clinicians when the instrument is now being translated to Norwegian. Since it is not too time-consuming, the PFS might be an awaited supplement to general symptom-scales that do not capture dynamic capacities.

7.2 Implications for future research

FEST-IT aims at exploring the effect of transference interventions in psychodynamic psychotherapy. When the results are ready, the study will hopefully add more knowledge to the evidence base for treatment of adolescents with major depression (R. Ulberg et al., 2012).

The head investigator in FEST-IT is also chair of the COST action “European Network on Individualized Psychotherapy: Treatment of Young People with Mental Disorders” (TREATME) (http://www.cost.eu/COST_Actions/ca/CA16102?management). TREATME was initiated to advance individualized psychotherapy for young people with mental health problems. As a management committee substitute from Norway, I will be able to participate in Working Group 2 (WG2): Mechanisms of change (mediators). This working

group focuses on identifying putative mechanisms of change in treatment of mental disorders in young people (aged 14-30). WG1 focuses on the empirical research on and theoretical and clinical descriptions of moderators (specific markers) interacting with psychotherapy for adolescents and young adults. WG3 focuses on assessing the need for appropriate and validated process and outcome measures for research on specific markers and mechanisms of change in psychotherapy. The overall aim of WG4 is to review research methods for examining youth (process and outcome) psychotherapy research from systematic reviews and expert consultation, with the particular focus to be identified from WG1 and WG2. Dissemination of Results and Communication with Stakeholders is the goal of WG5. A network of researchers on individualized youth psychotherapy is a way to effectively gather research findings and eventually share the knowledge with clinicians and stakeholders to improve psychotherapeutic interventions in the interest of the patients. As the age-group also includes young adults, the division between adolescents and adults are naturally “washed out”.

8. References

- Aas, I. H. (2011). Guidelines for rating Global Assessment of Functioning (GAF). *Ann Gen Psychiatry*, 10, 2. doi: 10.1186/1744-859x-10-2
- Alden, L., Wiggins, J., & Pincus, A. (1990). Construction of circumplex scales for the inventory of interpersonal problems. *J Pers Assess*, 55, 521 - 536.
- Andersen, P. L., & Bakken, A. (2015). Ung i Oslo 2015 *NOVA Rapport 8/15*.
- Anna Clark, L., & Watson, D. (1995). *Constructing validity: Basic issues in objective scale development* (Vol. 7).
- Azim, H. F., Piper, W. E., Segal, P. M., Nixon, G. W., & Duncan, S. C. (1991). The Quality of Object Relations Scale. *Bull Menninger Clin*, 55(3), 323-343.
- Babbie, E. R. (2010). *The practice of social research*. Belmont, CA; London: Wadsworth ; Cengage Learning [distributor].
- Barber, J., Muran, C., McCarthy, K., & Keefe, J. (2013). Research on dynamic therapies *Bergin and Garfield's handbook of psychotherapy and behavior change* (6th ed., pp. 443-494). New York, NY: John Wiley & Sons.
- Barker, C., & Pistrang, N. (2005). Quality criteria under methodological pluralism: Implications for conducting and evaluating research. *American Journal of Community Psychology*, 35(3-4), 201-212.
- Baron, R. M., & Kenny, D. A. (1986). The Moderator- Mediator Variable Distinction in Social Psychological Research: Conceptual, Strategic, and Statistical Considerations. *Journal of Personality and Social Psychology*, 51(6), 1173-1182. doi: 10.1037/0022-3514.51.6.1173
- Barrett-Lennard, G. T. (1962). Dimensions of therapist response as causal factors in therapeutic change. *Psychological Monographs: General and Applied*, 76(43), 1-36. doi: 10.1037/h0093918
- Beck, A. T., Steer, R. A., Ball, R., & Ranieri, W. (1996). Comparison of Beck Depression Inventories -IA and -II in psychiatric outpatients. *J Pers Assess*, 67(3), 588-597. doi: 10.1207/s15327752jpa6703_13
- Benjamin, L. S. (1996). *Interpersonal Diagnosis and Treatment of Personality disorders, second edition*. N.Y.: Guilford Press
- Benjamin, L. S. (1996). Introduction to the special section on structural analysis of social behavior. *J Consult Clin Psychol*, 64(6), 1203-1212.
- Benjamin, L. S., Rothweiler, J. C., & Critchfield, K. L. (2006). The use of structural analysis of social behavior (SASB) as an assessment tool. *Annu Rev Clin Psychol*, 2, 83-109. doi: 10.1146/annurev.clinpsy.2.022305.095337
- Bergin, A. E., & Garfield, S. L. (1971). *Handbook of psychotherapy and behavior change: An empirical analysis*. Oxford, England: John Wiley.
- Beutler, L. E. (2009). Making science matter in clinical practice: Redefining psychotherapy. *Clinical Psychology: Science and Practice*, 16(3), 301-317.
- Beutler, L. E., Someah, K., Kimpara, S., & Miller, K. (2016). Selecting the most appropriate treatment for each patient. *International Journal of Clinical and Health Psychology*, 16(1), 99-108. doi: <https://doi.org/10.1016/j.ijchp.2015.08.001>
- Bisson, J. I., Ehlers, A., Matthews, R., Pilling, S., Richards, D., & Turner, S. (2007). Psychological treatments for chronic post-traumatic stress disorder. Systematic review and meta-analysis. *Br J Psychiatry*, 190, 97-104. doi: 10.1192/bjp.bp.106.021402
- Bond, M., & Perry, J. C. (2004). Long-term changes in defense styles with psychodynamic psychotherapy for depressive, anxiety, and personality disorders. *Am J Psychiatry*, 161(9), 1665-1671. doi: 10.1176/appi.ajp.161.9.1665
- Borkovec, T. D. (1993). Between-group therapy outcome research: design and methodology. *NIDA Res Monogr*, 137, 249-289.
- Borkovec, T. D., & Miranda, J. (1999). Between-group psychotherapy outcome research and basic science. *J Clin Psychol*, 55(2), 147-158.

- Breuer, J., & Freud, S. (1957). *Studies on hysteria*. New York: Basic Books.
- Calderon, A., Schneider, C., Target, M., & Midgley, N. (2017). The Adolescent Psychotherapy Q-Set (APQ): A Validation Study. *Journal of Infant, Child, and Adolescent Psychotherapy*, 16(1), 106-120. doi: 10.1080/15289168.2016.1255499
- Cohen, J. (1969). *Statistical power analysis for the behavioral sciences*. New York: Academic Press.
- Connolly Gibbons, M. B., Crits-Christoph, P., Barber, J. P., Stirman, S. W., Gallop, R., Goldstein, L. A., . . . Ring-Kurtz, S. (2009). Unique and Common Mechanisms of Change across Cognitive and Dynamic Psychotherapies. *J Consult Clin Psychol*, 77(5), 801-813. doi: 10.1037/a0016596
- Connolly, M., Crits-Christoph, P., Shappell, S., Barber, J., Luborsky, L., & Shaffer, C. (1999). Relation of Transference Interpretations to Outcome in the Early Sessions of Brief Supportive-Expressive Psychotherapy. *Psychotherapy Research*, 9(4), 485-495. doi: 10.1080/10503309912331332881
- Coren, A. (1996). Brief therapy - base metal or pure gold? *Psychodynamic Counselling*, 2(1), 22-38. doi: 10.1080/14753639608411261
- Cousineau, T. M., & Shedler, J. (2006). Predicting physical health: implicit mental health measures versus self-report scales. *J Nerv Ment Dis*, 194(6), 427-432. doi: 10.1097/01.nmd.0000221373.74045.51
- Crits-Christoph, P., Connolly Gibbons, M. B., & Mukherjee, D. (2013). Psychotherapy process-outcome research. In M. J. Lambert (Ed.), *Bergin and Garfield's handbook of psychotherapy and behavior change*. NJ: Wiley: Hoboken.
- Crits-Christoph, P., Gibbons, M. B., Hamilton, J., Ring-Kurtz, S., & Gallop, R. (2011). The dependability of alliance assessments: the alliance-outcome correlation is larger than you might think. *J Consult Clin Psychol*, 79(3), 267-278. doi: 10.1037/a0023668
- Cuijpers, P. (2017). Four decades of outcome research on psychotherapies for adult depression: An overview of a series of meta-analyses. *Canadian Psychology/psychologie canadienne*, 58(1), 7.
- Cuijpers, P., Cristea, I. A., Karyotaki, E., Reijnders, M., & Hollon, S. D. (2017). Component studies of psychological treatments of adult depression: A systematic review and meta-analysis. *Psychotherapy Research*, 1-15. doi: 10.1080/10503307.2017.1395922
- Cuijpers, P., Driessen, E., Hollon, S. D., van Oppen, P., Barth, J., & Andersson, G. (2012). The efficacy of non-directive supportive therapy for adult depression: a meta-analysis. *Clin Psychol Rev*, 32(4), 280-291. doi: 10.1016/j.cpr.2012.01.003
- Dahl, H. J., Hoglend, P., Ulberg, R., Amlo, S., Gabbard, G. O., Perry, J. C., & Christoph, P. C. (2017). Does Therapists' Disengaged Feelings Influence the Effect of Transference Work? A Study on Countertransference. *Clin Psychol Psychother*, 24(2), 462-474. doi: 10.1002/cpp.2015
- Dahl, H. S., Rossberg, J. I., Bogwald, K. P., Gabbard, G. O., & Hoglend, P. A. (2012). Countertransference feelings in one year of individual therapy: an evaluation of the factor structure in the Feeling Word Checklist-58. *Psychother Res*, 22(1), 12-25. doi: 10.1080/10503307.2011.622312
- De Luca, C. R., Wood, S. J., Anderson, V., Buchanan, J. A., Proffitt, T. M., Mahony, K., & Pantelis, C. (2003). Normative data from the CANTAB. I: development of executive function over the lifespan. *J Clin Exp Neuropsychol*, 25(2), 242-254. doi: 10.1076/jcen.25.2.242.13639
- Deighton, J., Croudace, T., Fonagy, P., Brown, J., Patalay, P., & Wolpert, M. (2014). Measuring mental health and wellbeing outcomes for children and adolescents to inform practice and policy: a review of child self-report measures. *Child and Adolescent Psychiatry and Mental Health*, 8, 14-14. doi: 10.1186/1753-2000-8-14
- Della Rosa, E. (2016). *An exploration of transference interpretations in short term psychoanalytic psychotherapy with adolescents suffering from depression*. Unpublished doctoral dissertation. Travistock NHS Trust and University of East London.
- Della Rosa, E., & Midgley, N. (2017). Adolescent Patients' Responses to Interpretations Focused on Endings in Short-Term Psychoanalytic Psychotherapy. *Journal of Infant, Child, and Adolescent Psychotherapy*, 16(4), 279-290. doi: 10.1080/15289168.2017.1378531

- Derogatis, L. (1983). *SCL-90-R: Administration, scoring and procedure manual*: Clinical Psychometric Research, Towson, Md.
- DeVellis, R. F. (2016). *Scale development: Theory and applications* (Vol. 26): Sage publications.
- Diamond, D., Yeomans, F., Clarkin, J., & Levy, K. (2008). Mentalization and Attachment in Borderline Patients in Transference Focused Psychotherapy. *Mind to Mind: Infant Research, Neuroscience and Psychoanalysis*, 167 - 201.
- Donenberg, G. R. (1999). Reconsidering "between-group psychotherapy outcome research and basic science:" applications to child and adolescent psychotherapy outcome research. *J Clin Psychol*, 55(2), 181-190; discussion 191-200.
- Edwards, S. J. L., Lilford, R. J., & Hewison, J. (1998). The ethics of randomised controlled trials from the perspectives of patients, the public, and healthcare professionals. *BMJ : British Medical Journal*, 317(7167), 1209-1212.
- Ehlers, A., Bisson, J., Clark, D. M., Creamer, M., Pilling, S., Richards, D., . . . Yule, W. (2010). Do all psychological treatments really work the same in posttraumatic stress disorder? *Clin Psychol Rev*, 30(2), 269-276. doi: 10.1016/j.cpr.2009.12.001
- Elkjaer, H., Mortensen, E., Poulsen, S., Kristensen, E., & Lau, M. (2012). *(U)overensstemmelse mellem behandler- og patient-vurdering af effekt af gruppeterapi* (Vol. 29).
- Elliott, R. (1991). Five Dimensions of Therapy Process. *Psychotherapy Research*, 1(2), 92-103. doi: 10.1080/10503309112331335521
- Eysenck, H. J. (1952). The effects of psychotherapy: an evaluation. *journal of Consulting Psychology*, 16(5), 319.
- Falkenström, F., Granström, F., & Holmqvist, R. (2013). *Working alliance predicts psychotherapy outcome even while controlling for prior symptom improvement* (Vol. 24).
- FHI. (2018). Psykisk helse i Norge. In A. L. Reneflot A, Aase H, Reichborn-Kjennerud T, Tambs K, Øverland S. Psykisk helse i Norge, Folkehelseinstituttet. Rapport Januar 2018. (Ed.).
- Fishman, D. B. (2001). From single case to database: A new method for enhancing psychotherapy, forensic, and other psychological practice. *Applied and Preventive Psychology*, 10(4), 275-304. doi: 10.1016/S0962-1849(01)80004-4
- Fitzpatrick, R., Davey, C., Buxton, M. J., & Jones, D. R. (1998). Evaluating patient-based outcome measures for use in clinical trials. *Health Technol Assess*, 2(14), i-iv, 1-74.
- Folkehelseinstituttet. (2016). Effects of psychological therapies compared to antidepressant medication alone, or in combination with psychological therapies for depression in children and adolescents: a systematic review.
- Fonagy, P. (2015). The effectiveness of psychodynamic psychotherapies: An update. *World Psychiatry*, 14(2), 137-150. doi: 10.1002/wps.20235
- Fonagy P, C. D., Phillips J. (2014). *What works for whom? A critical review of treatments for children and adolescents* (2nd ed.). New York: Guilford Press.
- Foster, S. L., & Mash, E. J. (1999). Assessing social validity in clinical treatment research: issues and procedures. *J Consult Clin Psychol*, 67(3), 308-319.
- Freud, S. (1905). Fragment of an analysis of a case of hysteria (Vol. Standard Edition, 7: 1-122). London: Hogarth Press.
- Frost, N. (2011). *Qualitative research methods in psychology: Combining core approaches*: McGraw-Hill Education (UK).
- Froyd Michael J. Lambert Jeff D. Froyd, J. E. (1996). A review of practices of psychotherapy outcome measurement. *Journal of Mental Health*, 5(1), 11-16. doi: 10.1080/09638239650037144
- Gabbard, G. O. (2004). *Long-term psychodynamic psychotherapy: A basic text*. Arlington, VA, US: American Psychiatric Publishing, Inc.
- Gabbard, G. O. (2014). *Psychodynamic psychiatry in clinical practice*: American Psychiatric Pub.
- Gibbons, M. B., Crits-Christoph, P., Barber, J. P., Wiltsey Stirman, S., Gallop, R., Goldstein, L. A., . . . Ring-Kurtz, S. (2009). Unique and common mechanisms of change across cognitive and dynamic psychotherapies. *J Consult Clin Psychol*, 77(5), 801-813. doi: 10.1037/a0016596

- Gigerenzer, G. (2004). Mindless statistics. *The Journal of Socio-Economics*, 33(5), 587-606. doi: <https://doi.org/10.1016/j.socec.2004.09.033>
- Ginsburg, H. P., & Oppen, S. (1988). *Piaget's theory of intellectual development*, 3rd ed. Englewood Cliffs, NJ, US: Prentice-Hall, Inc.
- Golombek, H., & Korenblum, M. (1995). Brief psychoanalytic psychotherapy with adolescents. *Adolesc Psychiatry*, 20, 307-324.
- Goodyer, I. M., Reynolds, S., Barrett, B., Byford, S., Dubicka, B., Hill, J., . . . Fonagy, P. (2016). Cognitive behavioural therapy and short-term psychoanalytical psychotherapy versus a brief psychosocial intervention in adolescents with unipolar major depressive disorder (IMPACT): a multicentre, pragmatic, observer-blind, randomised controlled superiority trial. *The Lancet Psychiatry*. doi: 10.1016/S2215-0366(16)30378-9
- Goodyer, I. M., Tsancheva, S., Byford, S., Dubicka, B., Hill, J., Kelvin, R., . . . Fonagy, P. (2011). Improving mood with psychoanalytic and cognitive therapies (IMPACT): a pragmatic effectiveness superiority trial to investigate whether specialised psychological treatment reduces the risk for relapse in adolescents with moderate to severe unipolar depression: study protocol for a randomised controlled trial. *Trials*, 12, 175. doi: 10.1186/1745-6215-12-175
- Grande, T., Rudolf, G., Oberbracht, C., & Pauli-Magnus, C. (2003). PROGRESSIVE CHANGES IN PATIENTS' LIVES AFTER PSYCHOTHERAPY: WHICH TREATMENT EFFECTS SUPPORT THEM? *Psychotherapy Research*, 13(1), 43-58. doi: 10.1093/ptr/kpg006
- Gullestad, S. E., & Killingmo, B. (2013). *Underteksten. Psykoanalytisk terapi i praksis*. Oslo: Universitetsforlaget.
- Handbook of psychotherapy and behavior change*, 6th ed. (2013). (M. Lambert Ed. 6th ed.). Oxford, England: John Wiley & Sons.
- Hannan, C., Lambert, M. J., Harmon, C., Nielsen, S. L., Smart, D. W., Shimokawa, K., & Sutton, S. W. (2005). A lab test and algorithms for identifying clients at risk for treatment failure. *J Clin Psychol*, 61(2), 155-163. doi: 10.1002/jclp.20108
- Hardy, G. E., & Llewelyn, S. (2015). Introduction to psychotherapy process research *Psychotherapy research: Foundations, process, and outcome*. (pp. 183-194). New York, NY, US: Springer-Verlag Publishing.
- Hatcher, R. L., & Gillasp, J. A. (2006). Development and validation of a revised short version of the working alliance inventory. *Psychotherapy Research*, 16(1), 12-25. doi: 10.1080/10503300500352500
- Henriksen, J., Nielsen, P. F., & Bilenberg, N. (2012). New Danish standardization of the Child Behaviour Checklist. *Dan Med J*, 59(7), A4462.
- Hersoug, A. G., Hoglend, P., Monsen, J. T., & Havik, O. E. (2001). Quality of working alliance in psychotherapy: therapist variables and patient/therapist similarity as predictors. *J Psychother Pract Res*, 10(4), 205-216.
- Hersoug, A. G., & Ulberg, R. (2012). Siblings, friends, and parents: Who are the most important persons for adolescents? A pilot study of Adolescent Relationship Scale. *Nordic Psychology*, 64(2), 77-86. doi: 10.1080/19012276.2012.726813
- Hill, C. E. (1990). Exploratory in-session process research in individual psychotherapy: A review. *J Consult Clin Psychol*, 58(3), 288-294. doi: 10.1037/0022-006X.58.3.288
- Hoglend, P., Amlo, S., Marble, A., Bogwald, K. P., Sorbye, O., Sjaastad, M. C., & Heyerdahl, O. (2006). Analysis of the patient-therapist relationship in dynamic psychotherapy: an experimental study of transference interpretations. *Am J Psychiatry*, 163(10), 1739-1746. doi: 10.1176/ajp.2006.163.10.1739
- Hoglend, P., Bogwald, K. P., Amlo, S., Heyerdahl, O., Sorbye, O., Marble, A., . . . Bentsen, H. (2000). Assessment of change in dynamic psychotherapy. *J Psychother Pract Res*, 9(4), 190-199.
- Hoglend, P., Bogwald, K. P., Amlo, S., Marble, A., Ulberg, R., Sjaastad, M. C., . . . Johansson, P. (2008). Transference interpretations in dynamic psychotherapy: do they really yield sustained effects? *Am J Psychiatry*, 165(6), 763-771. doi: 10.1176/appi.ajp.2008.07061028

- Hoglend, P., Dahl, H. S., Hersoug, A. G., Lorentzen, S., & Perry, J. C. (2011). Long-term effects of transference interpretation in dynamic psychotherapy of personality disorders. *Eur Psychiatry*, 26(7), 419-424. doi: 10.1016/j.eurpsy.2010.05.006
- Hoglend, P., Engelstad, V., Sorbye, O., Heyerdahl, O., & Amlo, S. (1994). The role of insight in exploratory psychodynamic psychotherapy. *Br J Med Psychol*, 67 (Pt 4), 305-317.
- Hoglend, P., & Gabbard, G. (2012). When is Transference Work Useful in Dynamic Psychotherapy. *Psychodynamic Psychotherapy Research: Evidence Based Practice and Practice Based Evidence*, 449 - 467.
- Hoglend, P., Sorlie, T., Heyerdahl, O., Sorbye, O., & Amlo, S. (1993). Brief dynamic psychotherapy : patient suitability, treatment length, and outcome. *J Psychother Pract Res*, 2(3), 230-241.
- Holmqvist, R., & Armelius, B. A. (1994). Emotional reactions to psychiatric patients. Analysis of a feeling checklist. *Acta Psychiatr Scand*, 90(3), 204-209.
- Horvath, A. O., & Greenberg, L. S. (1989). Development and validation of the Working Alliance Inventory. *J Couns Psychol*, 36(2), 223-233. doi: 10.1037/0022-0167.36.2.223
- Jacobson, N. S., & Truax, P. (1991). Clinical significance: a statistical approach to defining meaningful change in psychotherapy research. *J Consult Clin Psychol*, 59(1), 12-19.
- Johansen, P. O., Krebs, T. S., Svartberg, M., Stiles, T. C., & Hølen, A. (2011). Change in defense mechanisms during short-term dynamic and cognitive therapy in patients with cluster C personality disorders. *J Nerv Ment Dis*, 199(9), 712-715. doi: 10.1097/NMD.0b013e318229d6a7
- Johansson, P., Hoglend, P., Ulberg, R., Amlo, S., Marble, A., Bogwald, K. P., . . . Heyerdahl, O. (2010). The mediating role of insight for long-term improvements in psychodynamic therapy. *J Consult Clin Psychol*, 78(3), 438-448. doi: 10.1037/a0019245
- John, O. P., & Benet-Martinez, V. (2000). *Handbook of research methods in social and personality psychology. Measurement: Reliability, construct validation, and scale construction*. New York, NY, US: Cambridge University Press.
- Johnston, C., & Gowers, S. (2005). Routine Outcome Measurement: A Survey of UK Child and Adolescent Mental Health Services. *Child and Adolescent Mental Health*, 10(3), 133-139. doi: 10.1111/j.1475-3588.2005.00357.x
- Jones, E. E., & Windholz, M. (1990). The psychoanalytic case study: toward a method for systematic inquiry. *J Am Psychoanal Assoc*, 38(4), 985-1015. doi: 10.1177/000306519003800405
- Kabisch, M., Ruckes, C., Seibert-Grafe, M., & Blettner, M. (2011). Randomized Controlled Trials: Part 17 of a Series on Evaluation of Scientific Publications. *Dtsch Arztebl Int*, 108(39), 663-668. doi: 10.3238/arztebl.2011.0663
- Kazdin, A. E. (2007). Mediators and mechanisms of change in psychotherapy research. *Annu Rev Clin Psychol*, 3, 1-27. doi: 10.1146/annurev.clinpsy.3.022806.091432
- Kazdin, A. E. (2009). Understanding how and why psychotherapy leads to change. *Psychother Res*, 19(4-5), 418-428. doi: 10.1080/10503300802448899
- Kennair, L. E. O., & Hagen, R. (2014). *Psykoterapi : tilnærminger og metoder*. Oslo: Gyldendal Akademisk.
- Kennedy, E., & Midgley, N. (2007). *Process and outcome research in child, adolescent and parent-infant psychotherapy: A thematic review*. London: NHS London.
- Kivlighan Jr, D. M., Multon, K. D., & Patton, M. J. (2000). Insight and symptom reduction in time-limited psychoanalytic counseling. *J Couns Psychol*, 47(1), 50-58. doi: 10.1037/0022-0167.47.1.50
- Klein, D. N., Schwartz, J. E., Santiago, N. J., Vivian, D., Vocisano, C., Castonguay, L. G., . . . Keller, M. B. (2003). Therapeutic alliance in depression treatment: controlling for prior change and patient characteristics. *J Consult Clin Psychol*, 71(6), 997-1006. doi: 10.1037/0022-006x.71.6.997
- Klein, M. (1988). *Envy and gratitude : and other works 1946-1963*: London : Virago, 1988.
- Korsgaard, H. O., Torgersen, S., Wentzel-Larsen, T., & Ulberg, R. (2016). Personality disorders and Axis I comorbidity in adolescent outpatients with ADHD. *BMC Psychiatry*, 16, 175. doi: 10.1186/s12888-016-0871-0

- Kraemer, H. C., Wilson, G. T., Fairburn, C. G., & Agras, W. S. (2002). Mediators and Moderators of Treatment Effects in Randomized Clinical Trials. *Arch Gen Psychiatry*, 59(10), 877-883. doi: 10.1001/archpsyc.59.10.877
- Kramer, U., Despland, J. N., Michel, L., Drapeau, M., & de Roten, Y. (2010). Change in defense mechanisms and coping over the course of short-term dynamic psychotherapy for adjustment disorder. *J Clin Psychol*, 66(12), 1232-1241. doi: 10.1002/jclp.20719
- Lambert, M. (2013). The Efficacy and Effectiveness of Psychotherapy *Bergin and Garfiels's Handbook of Psychotherapy and Behavior Change* (6th ed.). New Jersey: John Wiley & Sons, Inc.
- Lambert, M. J. (1992). Psychotherapy outcome research: Implications for integrative and eclectic therapists *Handbook of psychotherapy integration*. (pp. 94-129). New York, NY, US: Basic Books.
- Lambert, M. J. (2013). Outcome in psychotherapy: the past and important advances.
- Lawshe, C. H. (1975). A quantitative approach to content validity. *Personnel Psychology*, 28(4), 563-575. doi: 10.1111/j.1744-6570.1975.tb01393.x
- Levitt, H. M., Pomerville, A., & Surace, F. I. (2016). A qualitative meta-analysis examining clients' experiences of psychotherapy: A new agenda. *Psychol Bull*, 142(8), 801-830. doi: 10.1037/bul0000057
- Levy, K., & Scala, J. (2012). Transference, transference interpretations, and transference-focused psychotherapies. *Psychother Theory Res Pract*, 49, 391 - 403.
- Levy, K. N., Meehan, K. B., Kelly, K. M., Reynoso, J. S., Weber, M., Clarkin, J. F., & Kernberg, O. F. (2006). Change in attachment patterns and reflective function in a randomized control trial of transference-focused psychotherapy for borderline personality disorder. *J Consult Clin Psychol*, 74(6), 1027-1040. doi: 10.1037/0022-006x.74.6.1027
- Lindqvist, K., Falkenström, F., Sandell, R., Holmqvist, R., Ekeblad, A., & Thoren, A. (2017). Multilevel Exploratory Factor Analysis of the Feeling Word Checklist-24. *Assessment*, 24(7), 907-918. doi: 10.1177/1073191116632336
- Lutz, W., Ehrlich, T., Rubel, J., Hallwachs, N., Rottger, M. A., Joras, C., . . . Tschitsaz-Stucki, A. (2013). The ups and downs of psychotherapy: sudden gains and sudden losses identified with session reports. *Psychother Res*, 23(1), 14-24. doi: 10.1080/10503307.2012.693837
- Luzzi, A., Bardi, D., Ramos, L., & Slapak, S. (2015). *A study of process in psychoanalytic psychotherapy with children: The development of a method* (Vol. 18).
- McGraw, K., & Wong, S. P. (1996). *Forming Inferences About Some Intraclass Correlation Coefficients* (Vol. 1).
- McLeod, J. (2010). Case study research in counselling and psychotherapy. London: SAGE Publications Ltd.
- McLeod, J. (2011). *Qualitative Research in Counselling and Psychotherapy* (2nd ed.). London: Sage.
- McLeod, J. (2015). *Doing Research in Counselling and Psychotherapy*. Los Angeles, London, New Delhi, Singapore, Washington DC: SAGE.
- Medicine, I. o. (2015). *Psychological Testing in the Service of Disability Determination*. Washington, DC: The National Academies Press.
- Midgley, O'Keeffe, S., French, L., & Kennedy, E. (2017). Psychodynamic psychotherapy for children and adolescents: an updated narrative review of the evidence base. *Journal of Child Psychotherapy*, 1-23. doi: 10.1080/0075417X.2017.1323945
- Midgley, Rhode, M., Rustin, M., Hughes, C., & Cregeen, S. (2010). *IMPACT Group: Short-Term Psychoanalytic Psychotherapy (STPP) for Adolescents with Moderate or Severe Depression: A Treatment Manual*. Cambridge: IMPACT trial Office.
- Midgley, N. (2006). The 'inseparable bond between cure and research': clinical case study as a method of psychoanalytic inquiry. *Journal of Child Psychotherapy*, 32(2), 122-147. doi: 10.1080/00754170600780273
- Midgley, N. (2006). Re-reading "Little Hans": Freud's case study and the question of competing paradigms in psychoanalysis. *J Am Psychoanal Assoc*, 54(2), 537-559. doi: 10.1177/00030651060540021601

- Midgley, N., Capella, C., Goodman, G., Lis, A., Noom, M., Tishby, O., & Weitkamp, K. (2018). *Introduction to the special section on child and adolescent psychotherapy research* (Vol. 28).
- Midgley, N., Hayes, J., & Cooper, M. (2017). *Essential research findings in child and adolescent counselling and psychotherapy*. London: SAGE.
- Moleiro, C., & Beutler, L. E. (2009). Clinically significant change in psychotherapy for depressive disorders. *J Affect Disord*, 115(1-2), 220-224. doi: 10.1016/j.jad.2008.09.009
- Montgomery, S. A., & Asberg, M. (1979). A new depression scale designed to be sensitive to change. *Br J Psychiatry*, 134, 382-389.
- Morris, A. S., Silk, J. S., Steinberg, L., Myers, S. S., & Robinson, L. R. (2007). The Role of the Family Context in the Development of Emotion Regulation. *Soc Dev*, 16(2), 361-388. doi: 10.1111/j.1467-9507.2007.00389.x
- Möller, H.-J. (2011). Effectiveness studies: advantages and disadvantages. *Dialogues in Clinical Neuroscience*, 13(2), 199-207.
- Norcross, J. C., VandenBos, G. R., Freedheim, D. K., & Association, A. P. (2011). *History of Psychotherapy: Continuity and Change*: American Psychological Association.
- O'Keeffe, S., Martin, P., Goodyer, I. M., Wilkinson, P., Consortium, I., & Midgley, N. (2017). Predicting dropout in adolescents receiving therapy for depression. *Psychotherapy Research*, 1-14. doi: 10.1080/10503307.2017.1393576
- Ogles, B. M., Lunnen, K. M., & Bonesteel, K. (2001). Clinical significance: history, application, and current practice. *Clin Psychol Rev*, 21(3), 421-446.
- Ogrodniczuk, J., & Piper, W. (1999). Measuring therapist technique in psychodynamic psychotherapies: development and use of a new scale. *J Psychother Pract Res*, 8, 142 - 154.
- Ogrodniczuk, J., & Piper, W. (2004). The Evidence: Transference Interpretations and Patient Outcomes-a Comparison of Types of Patients. *Core Process in Brief Psychodynamic Psychotherapy. Advancing Effective Practice*, 165 - 184.
- Orlinsky, D., Heinonen, E., & Hartmann, A. (2015). Psychotherapy Process Research. In J. D. Wright (Ed.), *International Encyclopedia of the Social & Behavioral Sciences (Second Edition)* (pp. 515-520). Oxford: Elsevier.
- Patton, G. C., Sawyer, S. M., Santelli, J. S., Ross, D. A., Afifi, R., Allen, N. B., . . . Viner, R. M. (2016). Our future: a Lancet commission on adolescent health and wellbeing. *Lancet*, 387(10036), 2423-2478. doi: 10.1016/s0140-6736(16)00579-1
- Phohl, B., Blum, N., & Zimmermann, M. (1997). Structured Interview for DSM-IV Personality: SIDP-IV. *Washington, DC: American Psychiatric Press (Norwegian edition: Helgeland MW: Strukturert intervju for DSM-IV (SIDP-IV))*.
- Piper, W. E., Azim, H. F., Joyce, A. S., McCallum, M., Nixon, G. W., & Segal, P. S. (1991). Quality of object relations versus interpersonal functioning as predictors of therapeutic alliance and psychotherapy outcome. *J Nerv Ment Dis*, 179(7), 432-438.
- Piper, W. E., & Duncan, S. C. (1999). Object relations theory and short-term dynamic psychotherapy: findings from the Quality of Object Relations Scale. *Clin Psychol Rev*, 19(6), 669-685.
- Ponterotto, J. G. (2005). Qualitative research in counseling psychology: A primer on research paradigms and philosophy of science. *J Couns Psychol*, 52(2), 126-136. doi: 10.1037/0022-0167.52.2.126
- Rosenzweig, S. (1936). Some implicit common factors in diverse methods of psychotherapy. *American Journal of Orthopsychiatry*, 6(3), 412-415. doi: doi:10.1111/j.1939-0025.1936.tb05248.x
- Rossberg, J. I., Hoffart, A., & Friis, S. (2003). Psychiatric staff members' emotional reactions toward patients. A psychometric evaluation of an extended version of the Feeling Word Checklist (FWC-58). *Nord J Psychiatry*, 57(1), 45-53. doi: 10.1080/08039480310000257
- Roth, A., & Fonagy, P. (1996). *What Works For Whom? A Critical Review of Psychotherapy Research*.
- Roth, A., & Fonagy, P. (2005). *What works for whom: A critical review of psychotherapy research, 2nd ed*. New York, NY, US: Guilford Publications.

- Ryum, T. (2012). Contribution of the therapeutic bond, therapist interventions and their interaction to outcome in psychotherapy.
- Salkind, N. (2010). Encyclopedia of Research Design. Thousand Oaks, California.
- Samberg, E., & Marcus, E. R. (2005). Process, Resistance, and Interpretation *The American psychiatric publishing textbook of psychoanalysis*. (pp. 229-240). Arlington, VA, US: American Psychiatric Publishing, Inc.
- Sarles, R. M. (1994). Transference-countertransference issues with adolescents: personal reflections. *Am J Psychother*, 48(1), 64-74.
- Schneider, C. (2004). *The development of the child psychotherapy Q-set*. (65), ProQuest Information & Learning, US.
- Shadish, W. R., Cook, T. D., & Campbell, D. T. (2002). *Experimental and quasi-experimental designs for generalized causal inference*. Boston, MA, US: Houghton, Mifflin and Company.
- Shedler, J. (2010). The efficacy of psychodynamic psychotherapy. *American Psychologist*, 65(2), 98-109. doi: 10.1037/a0018378
- Sletten, M. A., & Bakken, A. (2016). Psykiske helseplager blant ungdom – tidstrenger og samfunnsmessige forklaringer *NOVA Notat 4/16*.
- Smith, M. L., & Glass, G. V. (1977). Meta-analysis of psychotherapy outcome studies. *American Psychologist*, 32(9), 752-760. doi: 10.1037/0003-066X.32.9.752
- Swift, W. J., & Wonderlich, S. A. (1990). Interpretation of transference in the psychotherapy of adolescents and young adults. *J Am Acad Child Adolesc Psychiatry*, 29(6), 929-935. doi: 10.1097/00004583-199011000-00016
- Tam, N. T., Huy, N. T., Thoa le, T. B., Long, N. P., Trang, N. T., Hirayama, K., & Karbwang, J. (2015). Participants' understanding of informed consent in clinical trials over three decades: systematic review and meta-analysis. *Bull World Health Organ*, 93(3), 186-198h. doi: 10.2471/blt.14.141390
- Town, J. M., Diener, M. J., Abbass, A., Leichsenring, F., Driessen, E., & Rabung, S. (2012). A meta-analysis of psychodynamic psychotherapy outcomes: evaluating the effects of research-specific procedures. *Psychotherapy (Chic)*, 49(3), 276-290. doi: 10.1037/a0029564
- Trijsburg, R. W., Frederiks, G. C. F. J., Gorlee, M., Klouwer, E., den Hollander, A. M., & Duivenvoorden, H. J. (2002). Development of the Comprehensive Psychotherapeutic Interventions Rating Scale (CPIRS). *Psychotherapy Research*, 12(3), 287-317. doi: 10.1093/ptr/12.3.287
- Ulberg Randi, Falkenberg, A. A., Naerdal, T. B., Johannessen, H., Olsen, J. E., Eide, T. K., . . . Johnsen, s. (2013). Countertransference Feelings when Treating Teenagers. A Psychometric Evaluation of the Feeling Word Checklist–24. *Am J Psychother*, 67(4), 347-358.
- Ulberg, R., Amlo, S., Critchfield, K. L., Marble, A., & Hoglend, P. (2014). Transference interventions and the process between therapist and patient. *Psychotherapy (Chic)*, 51(2), 258-269. doi: 10.1037/a0034708
- Ulberg, R., Amlo, S., & Høglend, P. (2014). Manual for transference work scale; a micro-analytical tool for therapy process analyses. *BMC Psychiatry*, 14(1), 291.
- Ulberg, R., Hersoug, A. G., & Hoglend, P. (2012). Treatment of adolescents with depression: the effect of transference interventions in a randomized controlled study of dynamic psychotherapy. *Trials*, 13, 159. doi: 10.1186/1745-6215-13-159
- Ulberg, R., Høglend, P., Marble, A., & Johansson, P. (2012). Women respond more favorably to transference intervention than men: a randomized study of long-term effects. *J Nerv Ment*, 200, 223 - 229.
- Ulberg, R., Johansson, P., Marble, A., & Hoglend, P. (2009). Patient sex as moderator of effects of transference interpretation in a randomized controlled study of dynamic psychotherapy. *Can J Psychiatry*, 54(2), 78-86.
- Ulberg, R., Marble, A., & Høglend, P. (2009). Do gender and level of relational functioning influence the long-term treatment response in dynamic psychotherapy? *Nord J Psychiatry*, 63, 412 - 419.

- Vermote, R., Lowyck, B., Luyten, P., Vertommen, H., Corveleyn, J., Verhaest, Y., . . . Peuskens, J. (2010). Process and outcome in psychodynamic hospitalization-based treatment for patients with a personality disorder. *J Nerv Ment Dis*, 198(2), 110-115. doi: 10.1097/NMD.0b013e3181cc0d59
- Wampold, B. E. (2001). *The Great Psychotherapy Debate: Models, Methods, and Findings*: L. Erlbaum Associates.
- Webb, C. A., DeRubeis, R. J., Amsterdam, J. D., Shelton, R. C., Hollon, S. D., & Dimidjian, S. (2011). Two aspects of the therapeutic alliance: differential relations with depressive symptom change. *J Consult Clin Psychol*, 79(3), 279-283. doi: 10.1037/a0023252
- Weisz, J. R., Kuppens, S., Ng, M. Y., Eckshtain, D., Ugueto, A. M., Vaughn-Coaxum, R., . . . Fordwood, S. R. (2017). What five decades of research tells us about the effects of youth psychological therapy: A multilevel meta-analysis and implications for science and practice. *Am Psychol*, 72(2), 79-117. doi: 10.1037/a0040360
- Weisz, J. R., Sandler, I. N., Durlak, J. A., & Anton, B. S. (2005). Promoting and protecting youth mental health through evidence-based prevention and treatment. *Am Psychol*, 60(6), 628-648. doi: 10.1037/0003-066x.60.6.628
- Wilson, F. R., Wei, P., & Donald, A. S. (2012). Recalculation of the Critical Values for Lawshe's Content Validity Ratio. *Measurement and Evaluation in Counseling and Development*, 45(3), 197-210. doi: 10.1177/0748175612440286
- Winston, B., Samstag, L. W., Winston, A., & Muran, J. C. (1994). Patient defense/therapist interventions. *Psychotherapy: Theory, Research, Practice, Training*, 31(3), 478-491. doi: 10.1037/0033-3204.31.3.478
- Wolf, M. M. (1978). Social validity: the case for subjective measurement or how applied behavior analysis is finding its heart. *Journal of Applied Behavior Analysis*, 11(2), 203-214. doi: 10.1901/jaba.1978.11-203
- Zachrisson, A. (2013). The Internal/External Issue: What is an Outer Object? Another Person as Object and as Separate Other in Object Relations Models. *The Psychoanalytic Study of the Child*, 67(1), 249-274. doi: 10.1080/00797308.2014.11785497

9. Appendix

9.1 Adolescent Relationship Scale (ARS)

The questions in ARS are separately scored on a visual analogue scale (0-10):

How much:

Do your **friends** mean to you?

Do you mean to your **friends**?

Does your **mother** mean to you?

Does your **father** mean to you?

Do you mean to your **mother**?

Do you mean to your **father**?

Do your **parents** mean to you?

Do you mean to your **parents**?

Do your **siblings** mean to you?

Do you mean to your **siblings**?

What is:

Your **quality of life** like now?

9.2 Quality of Object Relations (QOR)

Lifelong pattern on relationships from primitive to mature assessed with QOR (Azim et al., 1991) is a dimension supposed to predetermine suitability for dynamic psychotherapy (P. Hoglend, Sorlie, Heyerdahl, Sorbye, & Amlo, 1993). The reliability and construct validity is adequate (Piper & Duncan, 1999). In FEST the average of the three QOR 8-point scales were used. In FEST-IT the average of two scales were used, omitting the scale *Quality of sexual intimate relationships*.

Quality of interpersonal relationships:

- 8-5 A history of relationships characterized by stability, gratification and mutuality in most important relations. Others are seen as whole, autonomous persons. Recent interpersonal functioning may be poorer, but the patient can give detailed examples from at least one earlier important high quality relationship. Conflictual feelings may be seen towards same sex persons and fears of loss of opposite sex persons.
- 4-3 Mostly stable but less gratification and mutuality in most important relations. Passivity, dependency or need to control others are predominant due to separation anxiety.

- 2-1 Mostly unstable relations of little valued persons. Others are seen as need-gratifying objects. Stable overly dependent relations only with parental objects.

Quality of sexual intimate relationships:

- 8-5 Ability to establish a deep mutual and stable relationship which includes sexual gratification and mutual support. Willingness to allow emotional vulnerability.
- 4-3 Submissiveness, controlling traits or low self-esteem interfere with sexual gratification and deep mutual contact. Lack of full commitment or preoccupation with possible loss.
- 2-1 Superficial unstable sexual relationships or unable to initiate sexual relationships.

Quality of friendships:

- 8-5 Give-and-take relations with at least two friends.
- 4-3 Superficial friendships or one mutual friendship.
- 2-1 No more than one confident outside near family.

9.3 The factors from Feeling Word Checklist (FWC-24)

Factor	Items
Confident	: Playful, Vigorous, Enthusiastic, Happy, Open, Warm, Content
Inadequate	: Overwhelmed, Tense, Nervous, Thunderstruck, Helpless
Disengaged	: Indifferent, Tired of, Cold, Irritated
Neutral	: Prudent, Neutral

(Ulberg Randi et al., 2013)

Papers

RESEARCH ARTICLE

Open Access

Assessment of dynamic change in psychotherapy with adolescents

Elisabeth Ness^{1,2*} , Hanne-Sofie Johnsen Dahl^{1,2}, Peter Tallberg^{1,3}, Svein Amlo⁴, Per Høglend¹, Agneta Thorén⁵, Jens Egeland^{2,6} and Randi Ulberg^{1,2}

Abstract

Background: Diagnostic interviews and questionnaires are commonly used in the assessment of adolescents referred to child and adolescent mental health services. Many of these rating scales are constructed for adults and focus on symptoms related to diagnosis. Psychodynamic Functioning Scales (PFS) focus on relational aspects and how the patients handle affects and solve problems, rather than manifest symptoms. As these aspects are considered important for mental health, the PFS were developed to assess change in adults, consistent with the relational and intrapsychic concepts of dynamic psychotherapy. The scales describe internal predispositions and psychological resources that can be mobilized to achieve adaptive functioning and life satisfaction. PFS consist of six subscales; the relational subscales *Family*, *Friends* and *Romantic/Sexual relationships* and the dynamic subscales *Tolerance for Affects*, *Insight* and *Problem-solving Capacity*. PFS has been used for the first time as a measure of change in adolescent psychotherapy. This study examines the reliability of PFS when used to assess adolescents' level of relational functioning, affective tolerance, insight, and problem-solving capacities.

Methods: Outpatient adolescents 16–18 years old with a major depressive disorder were included in the First Experimental Study of Transference work in Teenagers (FEST-IT). They were evaluated before and after time-limited psychodynamic psychotherapy with an audio-recorded semi-structured psychodynamic interview. Based on the audio-tapes, raters with different clinical background rated all the available interviews at pre-treatment ($n = 66$) and post-treatment ($n = 30$) using PFS. Interrater reliability, the reliability of change ratings and the discriminability from general symptoms were calculated in SPSS.

Results: The interrater reliability was on average good on the relational subscales and fair to good on the dynamic subscales. All pre-post changes were significant, and the analyses indicated discriminability from general symptoms. The interrater reliability on PFS (mean) and Global Assessment of Functioning were good to excellent.

Conclusion: Based on the interrater reliability in our study, PFS could be recommended in psychotherapy with adolescents by experienced clinicians without extensive training. From the post-treatment evaluations available, the scales seem to capture statistically and clinically significant changes. However, the interrater reliability on dynamic subscales indicates that subscales of PFS might be considered revised or adjusted for adolescents.

Trial registration First Experimental Study of Transference-Work-In Teenagers (2011/1424 FEST-IT). ClinicalTrials.gov Identifier: NCT01531101

Keywords: Rating scales, Outcome, Adolescent, Short-term psychodynamic psychotherapy (STTP)

*Correspondence: elisabeth.ness@siv.no

² Research Unit, Division of Mental Health, Vestfold Hospital Trust, PO Box 2168, 3103 Tønsberg, Norway
Full list of author information is available at the end of the article

© The Author(s) 2018. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Background

Assessing psychological growth in adolescents is important to identify whether a specific treatment is effective. The average treatment effectiveness is important, although to individualize treatment and help those who don't improve, research should also focus on the mechanisms underlying treatment effectiveness [1].

Therapy with adolescents presents the therapist with specific challenges that might be different from psychotherapy with adults. Establishing a therapeutic alliance with adolescents could be comprised by the patients seeing the therapist as just another authority figure in their lives [2]. Adolescents are at a stage in their development in which they are struggling with autonomy and individuation, and they need to undertake several developmental tasks to make a successful transition to adulthood. The ability to recognize and process emotions is under development [3, 4]. In addition, dropout from treatment is significant, especially for adolescents [5, 6].

There is emerging evidence of the efficacy of psychodynamic psychotherapy for children and adolescents [7, 8]. A recent randomized controlled superiority trial in England (IMPACT-study) for adolescents with unipolar major depressive disorder compared Cognitive Behavioural Therapy (CBT) and short-term psychoanalytical psychotherapy (STPP) versus a brief psychosocial intervention [7]. They concluded that none were superior to the others.

Psychodynamic psychotherapy focuses on relational and internal psychological growth. The dynamic processes one seeks to enhance during therapy includes emotional growth, development, and maturation. The normal development in young people, like growth in size, sexual maturity, emotional development, and cognitive capacity, may be potential triggers or amplifiers of psychiatric disorder, or a potential for the adolescents' subjective quality of life.

Psychodynamic therapy aims at helping patients understand more of the origin of their symptoms, and the function these symptoms may play in their life. In addition, self-understanding of interpersonal patterns is seen as a central change mechanism in dynamic psychotherapy [9, 10]. The achievement of a more nuanced understanding of self and others might enhance psychological flexibility without developing symptoms.

Clinicians and researchers are interested in the therapeutic effect on recurrence risk and the long-term effectiveness of existing treatments. Patients who receive psychodynamic therapy seem on average to maintain therapeutic gains and appear to continue to improve after treatment ends [11, 12]. Since psychodynamic therapy aims at endowing patients with healthier relationships, greater insight and increased awareness of their affects,

psychodynamic therapy may contribute to the prevention of recurrent symptoms also in therapy with adolescents.

Diagnostic interviews and questionnaires are commonly used in the assessment of young people referred to child and adolescent mental health services. They are mainly concerned with measuring symptoms to establish diagnoses. Many psychiatric rating scales were originally constructed for adult patients and have not been tested for reliability or validity in adolescents. Over 100 different measures for evaluation of outcomes exist (reflecting upon progress in therapy, overall outcome or specific symptoms) [13]. As for self-reports, a review of child self-report measures in child and adolescent mental health services (CAMHS) identified 11 measures having potential for use as outcome measures in routine practice. However, none of these measures had sufficient psychometric evidence available to demonstrate that they could reliably measure both severity and change over time [14].

In a review of the evidence base of psychodynamic psychotherapy for children and adolescents [8], several outcome measures were used in multiple studies. There are however limitations in the existing global impairment measures. Most are unidimensional and many incorporate symptomatology into the measurement, mixing severity of psychopathology with functional impairment. Some are lengthy and thus impractical for clinical or research use. Overall global functioning measures may not differentiate what is specific for psychodynamic psychotherapy, for instance the quality of relations to close others, and the ability to think about and handle problems, as well as toleration of affects.

Fine-graded scales are needed to measure change in psychotherapy. The scales need to capture the status prior to treatment, ideally also track the improvement during therapy, and after the psychotherapy. Psychodynamic psychotherapy aims at gaining insight into the patients' life histories and their present-day problems and to recognize non-healthy recurring patterns. The symptoms themselves are not the main focus when assessing change and outcome in dynamic psychotherapy. Although outcome measures related to dynamic capacities already exist, they tend to include a defined capacity (e.g. Reflective Functioning Scale [15]), or capacities as one aspect of comprehensive diagnostic systems (e.g. Mental Functioning Scale of the Psychodynamic Diagnostic Manual (PDM [16]), the Operationalized Psychodynamic Diagnoses (OPD [17] and the Shedler-Westen Assessment Procedure with 200 items (SWAP-200) [18]). The Wallerstein's Scales of Psychological Capacities (SPC) is an instrument developed to meet clinical and research needs in assessing change in patients who have undergone long-term psychodynamic or psychoanalytic therapy [19]. The SPC, though rather comprehensive

with 17 defined capacities, have been adapted to adolescents (Ad-SPC) [20].

To our knowledge there is a lack of brief clinician-rated instruments to assess dynamic capacities with adolescents. The Youth Outcome Questionnaire (YOQ) [21] is a 64 item report for children and adolescents (ages 4–17) completed by the parent/guardian. A self-report version also exists. A comprehensive clinician-rated instrument to assess intrapsychic processes in children and adolescents is the Operationalized Psychodynamic Diagnoses in Children and Adolescents (OPD-CA-2) [17], a multi-axial diagnostic and classification system based on psychodynamic principles based on four axes (interpersonal, conflict, structure, and prerequisites for treatment). The diagnostic way of thinking does not require training, but the rating should ideally be done by certified raters.

Psychodynamic Functioning Scales

In the present study we seek to test the reliability of an instrument which is developed to capture change after psychodynamic therapy. Høglend and colleagues developed a set of scales measuring psychological functioning, the Psychodynamic Functioning Scales (PFS) [22]. PFS are meant to discriminate from general symptoms or global functioning and capture the complexity of changes that potentially can occur during and after psychodynamic therapy. Ratings are based on a semi-structured dynamic interview. Current functioning within the last 3 months are rated. The clinician rated scales describe internal predispositions, psychological resources, capacities, or aptitudes that can be mobilized by the individual in order to achieve adaptive functioning and life satisfaction. The six scales are: *quality of family relationships*; *quality of friendships*; *romantic/sexual relationships*; *tolerance for affects*; *insight*; and *problem-solving capacity*. The scale format has been modelled after the Global Assessment of Functioning (GAF), with ten descriptive levels and scale points ranging from 1 to 100. Each of the six scales therefore covers the entire range of functioning, from superior (100) to extremely poor (1). The use of a well-known scale format should make the scales easier to learn. The intention was to make the scales “fine-grained” enough to capture reliable changes during psychotherapy. The content validity and Guttman scale structure have been tested with Q-sort methodology [23–25] performed by a large number of psychotherapists from Norway, Finland, and Germany [26]. PFS has been deemed as a reliable instrument to assess mental health and change after therapy in adults [22]. Using the Psychodynamic Functioning Scales as an outcome measure in a study of adults revealed that insight was the most difficult scale to rate reliably, especially at pre-treatment

[22]. The Psychodynamic Functioning Scale has not until now been reliability-tested for adolescents.

Aims

The present study tests the interrater reliability of five scales from PFS: Quality of Family Relations, Quality of Friendships, Tolerance for Affects, Insight and Problem-Solving Capacity. The reliability of change ratings, and the discriminability from global functioning (GAF; Global Assessment of Functioning [27]) and subjective distress (GSI; Global Severity Index from the Symptom Checklist-90 [28]), during brief dynamic psychotherapy with adolescents is also tested.

Methods

The First Experimental Study of Transference Work-In Teenagers (FEST-IT)

Data from FEST-IT are used. FEST-IT is a randomized, controlled study on psychodynamic psychotherapy for adolescents with depression [29].

Patients

The patients were the first 70 adolescents included in FEST-IT. One patient withdrew the consent and three interviews were lost due to technical problems with the audio-recording. Hence, 66 patients were included in the analyses in the present study. There were 12 boys and 54 girls aged 16–18 years. The patients were recruited among adolescents with symptoms of depression referred either to private practice or child and adolescent outpatient mental health clinics in the South-Eastern Health Region, representing mainly urban and some rural areas. All patients were attending classes in lower or upper secondary school.

Adolescents with current unipolar major depressive disorder according to Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV; American Psychiatric Association, 2000) were included. Adolescents with generalized learning difficulties, pervasive developmental disorder, psychosis, or substance addiction were excluded. Comorbidity was expected to be frequent.

Axis I and II diagnosis were based on the Mini International Neuropsychiatric Interview (M.I.N.I.) and Structured Interview for DSM-IV Personality (SIDP-IV). Table 1 shows some of the pre-treatment characteristics. Axis I diagnoses beside depression were mostly social phobia, panic disorder and general anxiety. A total of 31 patients had one or more Axis II disorders—primarily depressive or avoidant personality disorders. The patient sample had, on the average, mild to moderate symptoms and dysfunctions. The mean GAF score at the initial psychodynamic evaluation (PFS) was 58.0 (SD=6.1 range

Table 1 Pre-treatment characteristic of the 66 patients included

	Total (n = 66) Mean (SD)
Age	17.3 (0.7)
PFS	59.7 (6.1)
GAF	58.0 (6.1)
IIP-C	1.36 (0.4)
GSI	1.33 (0.5)
BDI	28.7 (9.0)
	N (%)
Female	54 (82)
Axis I diagnoses	
Depressive disorder	100 (100)
Social phobia	19 (29)
Panic disorder	13 (20)
General anxiety	17 (26)
Eating disorder	2 (3)
PTSD	2 (3)
More than two axis I diagnoses	17 (26)
Axis II diagnoses	30 (45)
Depressive	24 (36)
Avoidant	19 (29)
Negativistic	4 (6)
Obsessive compulsive	3 (5)
Paranoid	3 (5)
Dependent	2 (3)
Borderline	1 (2)
Histrionic	1 (2)
Schizoid	1 (2)
More than one axis II diagnoses	17 (26)

PFS Psychodynamic Functioning Scale; GAF Global Assessment of Functioning (n = 47), IIP-C inventory of interpersonal problems—circumplex version, GSI Global Severity Index (SCL-90), BDI Beck Depression Inventory

44.2–73.2). The mean GSI score (from SCL-90) was 1.3 (SD = 0.5, range 0.5–2.7). The mean BDI score was 28.7 (SD = 9.0, range 10–58). The distribution of mean pre-treatment scores indicated that the sample of 66 patients was a group of moderately depressed adolescents, representative of typical outpatients offered dynamic psychotherapy. The range of the pre-treatment scores of the five scales of PFS covered the area of functioning from relatively severe and chronic disturbances to moderate and intermittent problems of living (range 45.6–71.0). Only one patient reported taking antidepressant medication at baseline, i.e. at the beginning of therapy. One patient was taking antidepressants at the end of therapy. This was, however, not the same patient. One patient was taking antipsychotics throughout the study period. One patient

was taking sleeping medicine at pre-treatment and 4 patients were taking sleeping medicine at post-treatment.

Therapists

The twelve therapists worked in out-patient clinics and/or in private practice. Eight were psychiatrists and four were clinical psychologists. There were six men and six women. All therapists were trained therapists and had at least 2 years of formal training in psychodynamic psychotherapy.

Treatment

Short-term psychodynamic/psychoanalytic psychotherapy (STPP) based on the STPP manual from the IMPACT study [30] was used as the manual for the treatment. The manual combines aspects of STPP that focus principally on techniques aimed at helping young people overcome developmental problems, as well as emphasizing the role of the interpretation of unconscious conflicts, attachment theory and the concepts of internal working models. With the agreement of the adolescent, parallel work with parents was included. Antidepressant medication could be added in severe cases according to the national guidelines in Norway [31]. The patients were randomized to two treatment groups. In both groups general psychodynamic techniques [30] were used. The patients were offered 28 weekly sessions.

A 1-year training program prepared the therapists for treating patients in the study. Peer supervision in groups with material from the audio-recorded therapies was offered regularly during the study to help maintain the quality of the therapies and adherence check to the manualised therapies.

Evaluators and raters

Four individual evaluators conducted the patient interviews at baseline (pre-treatment) and at the end of therapy (post-treatment). The four evaluators and the two raters were clinical psychologists or psychiatrists and had their clinical training from different psychodynamic institutes. The four evaluators were females, while the two raters in this study were males. All had long clinical experience ranging from 12 to 30 years. One of the two raters had his main clinical background from out-patient adults, while the other rater had been working with adolescents from an in-patient department over the last decade.

Both the evaluators and raters were blind to treatment. They met on regular basis for group supervision both before and during the study. Meetings also involved plenary discussions after individual scorings of audio-recorded interviews.

Measures in the present study

Psychodynamic Functioning Scales (PFS)

PFS [22] were developed to capture evaluator-rated change in dynamic and interpersonal functioning. Current functioning was rated on the basis of a semi-structured dynamic interview. Five of the six scales were used: Quality of Family Relations, Quality of Friendships, Tolerance for Affects, Insight and Problem-Solving Capacity. The five subscales used in the analysis are presented in the Additional file 1. Each of the scales covers the entire range of functioning, with ten descriptive levels and scale points ranging from 1 to 100. The relational scales, *quality of family relations* and *quality of friendships* and *romantic/sexual relationships*, cover the mutuality and emotional responsiveness in relationships. The ratings of the two scales related to family and friends are based on evaluating the degree of mutuality and adequacy of the commitment in relationships, the ability to take other's perspective, to describe close others across an external and internal dimension, feeling of being needed and a sense of belonging and the capacity to reconcile parent's or friends' shortcomings and make the best of the relationship. If parents are not alive the evaluation is based on memory of them or internalized object relations. The *romantic/sexual relationships* involve also the capacity to establish long-term relationships characterized by love, trust, reciprocal mature dependency and active, flexible sexual pleasure. The *tolerance for affects* covers the ability to experience, differentiate and express various affects verbally and nonverbally, and to what degree disappointments lead to symptoms like avoidance, anxiety, depression or restrictions of goals. *Insight* covers mainly cognitive understanding of the main dynamics of inner conflicts, related inter-personal patterns and connection to the past. Also, the ability to describe and understand own vulnerability and reactions to stress. The *problem-solving capacity* covers the ability to handle any difficult situation without developing symptoms, avoidance or inadequate actions. Self-observation, planning, ability to explore new areas and enjoy recreation and pursue meaningful goals are parts of this scale. The PFS is deemed to be reliable [22, 26]. Although most adolescents have some experience of intimate relations, the minority have yet established more definite intimate relationships patterns. Thus, the scale *romantic/sexual relationships* was omitted for adolescents in the present study. The scales (Additional file 1) were developed with descriptive levels in English. In FEST-IT the English version was used although the semi-structured interview with anchor points was in Norwegian.

Global Assessment of Functioning (GAF)

The GAF (DSM.3rd ed. 1987) [27] is a numeric scale (1 through 100) with ten descriptive levels assigning a clinical judgment to the individual's overall functioning level. GAF recorded values used in FEST-IT are separate scores for symptoms (GAF-S) and functioning (GAF-F). For both the GAF-S and GAF-F scales, there are 100 scoring possibilities (1-100). Impairments in psychological, social and occupational/school functioning are considered, but those related to physical or environmental limitations are not. GAF seek to capture symptom relief. GAF was an outcome measure in the adult study FEST and therefore chosen also in the adolescent study instead of Children's Global Assessment Scale (CGAS). The GAF-scale can be scored reliably although the limitations as a single instrument has been discussed. [32, 33].

Symptom Checklist-90 (SCL-90)

The SCL-90 [28] is a self-report psychometric instrument (questionnaire) designed to evaluate a broad range of psychological problems and symptoms of psychopathology. It is also used in measuring the progress and outcome of psychiatric and psychological treatments or for research purposes. The SCL-90-R is normed on individuals 13 years and older. It consists of 90 items and takes 12–15 min to administer. The SCL-90 is used as an outcome measure in many studies. In the present study we use the General Symptom Index, which is the mean of the 90 items. Its psychometric properties have been examined and described [34, 35].

Beck depression inventory (BDI-II)

The BDI-II [36] is a widely used 21-item self-report inventory composed of items relating to symptoms of depression. The BDI-II is designed for individuals aged 13 and over, thus measuring the severity of depression in adolescents and adults. Psychometric properties have been described with high reliability and a capacity to discriminate between depressed and non-depressed subjects and high content and structural validity [37].

Evaluation and rating

Each patient was interviewed by one evaluator at pre- and post-treatment with a semi-structured GAF interview and a psychodynamic interview modified after Malan [38] and Sifneos [39]. The psychodynamic interview lasted approximately 45–60 min and the therapist was present if possible. However, the rater did not discuss or clarify questions with the therapist during the interviews or before rating the scales. No therapist ratings were included in the analysis. Ratings on the five dynamic scales and GAF were done by the evaluator. After the interviews, the patients filled out the SCL-90-R and the

BDI-II. All interviews were audio-recorded and independently assessed by two additional raters. During plenary calibration meetings after the individual ratings were recorded, the ratings and quality of the interview was discussed.

Statistical analysis

The raters and evaluators assessed the patients before and after therapy. From this group of six we estimated the interrater reliability (IRR) for single raters at pre-treatment. Assessments by the two raters were used to determine the IRR at pre- and post-treatment. Ratings of audio-recorded interviews rated by the same two raters for all subjects (66 at pre-treatment and 30 at post-treatment) were used for the Intra Class Correlation-analyses (ICC) [40] (two-way mixed consistency) for ordinal scores. This is represented in SPSS as “Two-Way Mixed” because it models both an effect of rater and of ratee (i.e. two effects) and assumes a random effect of ratee but a fixed effect of rater (i.e. a mixed effect model). The statistical analyses were done using SPSS version 23 SPSS. Inc. 2016. Ratings of GAF were only available for analysis in 47 patients pre-treatment due to missing data. Only 30 patients were rated on both occasions by both raters. The pre-/post-ratings include the same 30 patients for all instruments including GAF.

We also estimated the ICC for average scores of 3 raters, including the evaluator for each subject as the third rater, at pre-treatment. The model was then “Two-Way Random” in SPSS.

Average pre-treatment scores on each scale were compared with average post-treatment scores, by use of paired t-tests, on the 30 patients evaluated before and after therapy from 3 raters.

Guidelines for evaluating assessment instruments in psychology developed by Cicchetti and Sparrow [41], closely resembled by guidelines by Fleiss [42] and by Landis and Koch [43], state that when the reliability coefficient is below 0.40, the level of clinical significance is poor; when it is between 0.40 and 0.59, it is fair; when it is between 0.60 and 0.74, it is good; and when it is above 0.75 the level of clinical significance is excellent.

Jacobson and Truax [44] have developed a commonly used measure of assessing statistically reliable change—the Reliable Change Index (RCI). The RC coefficient is equivalent to the difference between two scores divided by the standard error of the difference between the scores, which is derived from test–retest reliability of a measure and standard deviation of pre-treatment scores on that measure ($RCI = (X_{\text{post}} - X_{\text{pre}}) / S_{\text{diff}}$) where S_{diff} = the standard error of the difference between the two test scores. $S_{\text{diff}} = \sqrt{S(\text{SE}_m^2)}$ and the Standard Error of the Measurement $\text{SE}_m = s\sqrt{1 - r_{xx}}$ where r_{xx} = reliability coefficient of

the instrument (in this study the ICC was used). For the GAF, GSI and BDI the S_{diff} were calculated from applying the denominator from the t test formula with s_1 and s_2 as variance of the pretest scores and posttest scores. An RC coefficient that is larger than 1.96 is usually regarded as unlikely ($p < 0.05$) to occur without any actual change and an indication of the individual's reliable change.

We used a SPSS correlation test (Pearson correlation) to estimate if the average scores of pre-treatment variables were discriminable from general symptoms (GSI) or dysfunction (GAF).

Results

The interrater reliability estimates of all the patients available for analysis at pre-treatment for single raters are shown in Table 2.

To study reliability in subscales we report the ICC for each subscale of PFS, mean values of PFS, and GAF. The lower bounds of the confidence intervals were unsatisfactory (< 0.40) for the subscales *tolerance for affects* and *insight*. With the average scores of the two raters who rated all subjects, the interrater reliability estimates of the 66 patients pre-treatment and of 30 of the same patients from post-treatment are shown in Table 3. At pre-treatment, we achieved excellent average reliability on the scales *family*, *friendships*, *PFS mean* and *GAF*, and good reliability on *insight*, *tolerance for affects* and *problem-solving capacity*. The lower bound of the confidence interval was unsatisfactory for only one of the single scales at pre-treatment: *insight*.

At post-treatment, also Table 3, with the average scores of the two raters, the ICC measures were all above 0.60 except for the subscale *insight* (0.59). However, the lower bounds of the confidence intervals were less than 0.40 for 2 subscales: *family* and *insight*.

The two raters based their ratings on audio-recorded interviews only. The evaluators, on the contrary, met the patients for interviews as part of the assessment. At

Table 2 Interrater reliability estimates (intraclass correlation; ICC) for single raters randomly drawn from a group of six (4 evaluators and 2 raters)

PFS pre-treatment (n = 66)	ICC	95% CI
Family	0.60	0.46–0.71
Friendships	0.56	0.43–0.69
Tolerance for affect	0.44	0.29–0.58
Insight	0.46	0.31–0.60
Problem-solving capacity	0.56	0.42–0.69
PFS mean	0.61	0.47–0.72
GAF pre-treatment (n = 47)	0.66	0.52–0.78

PFS Psychodynamic Functioning Scales, GAF Global Assessment of Functioning

Table 3 Pre-treatment and post-treatment interrater reliability estimates (intraclass correlations; ICC), for average scores of two raters

PFS	Pre-treatment (n = 66) ICC	Post-treatment (n = 30) ICC
Family	0.82**	0.66
Friendships	0.75*	0.73*
Tolerance for affect	0.69*	0.80*
Insight	0.61	0.59
Problem-solving capacity	0.70*	0.71*
PFS mean	0.81**	0.80*
GAF	0.80*** ^a	0.87**

* Lower bounds of 95% confidence interval (CI) > 0.40 ** lower bounds of 95% CI > 0.60

^a Pre-treatment n = 47

pre-treatment, the two raters differed in rating the relational and dynamic subscales respectively. The one rater tended to rate the patients higher than the other rater on the relational scales; *family* and *friendships*. However, the situation was quite opposite for the dynamic scales; *tolerance for affects*, *insight* and *problem-solving capacity*. On all the dynamic subscales, the other rater was rating the patients with higher scores.

With ratings also from the evaluators interviewing the patients at pre-treatment, the interrater reliability estimates increased. For all subscales of PFS, lower bounds of the 95% confidence intervals were above 0.50 and all ICC-values ≥ 0.70 . The same was true for GAF (Table 4).

Table 5 presents the mean scores on all subscales of PFS at pre-treatment and post-treatment for the 30 patients evaluated on both occasions. The post-treatment values of PFS and GAF indicate less severe problems in psychodynamic functioning and global symptoms at the end of therapy. The decreased post-treatment values of BDI and GSI indicate less depressive symptoms and less symptoms of psychopathology respectively. All changes were statistically significant at $p < 0.01$ paired t-test (two-tailed). The largest amount of change on the PFS subscales during the individual psychotherapy, and the highest ratio of patients with reliable changes according to the Reliable Change Index (RCI) [44], tended to be for the *tolerance for affects* (10 patients). The RCI was equal for PFS mean and GAF. The cut off score for reliable change was 6.1 for PFS mean and 6.8 for GAF, meaning a patient would have to improve with more than 6.1 points on the mean rating of PFS for the change to be considered reliable. The individual improvement seemed to be most reliable for the self-reported depression scale BDI (74% of all cases analysed).

Table 4 Pre-treatment interrater reliability estimates (intraclass correlations; ICC), for average scores of 3 raters (evaluator and 2 raters)

PFS pre-treatment (n = 66)	ICC	95% CI
Family	0.82	0.72–0.88
Friendships	0.79	0.69–0.87
Tolerance for affect	0.70	0.55–0.81
Insight	0.72	0.58–0.82
Problem-solving capacity	0.79	0.69–0.87
PFS mean	0.82	0.73–0.89
GAF pre-treatment (n = 47)	0.85	0.76–0.91

PFS Psychodynamic Functioning Scales, GAF Global Assessment of Functioning

A correlation matrix was made to evaluate whether the subscales from PFS could be differentiated from global functioning and subjective distress. Table 6 shows the results. *Tolerance for affects* and *problem-solving capacity* seemed to be the subscales with the highest correlation with GAF. The correlation with GSI was weak or moderate for all subscales. The PFS mean had a strong correlation (0.72) with GAF and a weak correlation (–0.29) with GSI.

Discussion

The interrater reliability of the PFS for assessment of change in psychodynamic therapy with adolescents was on average good on the subscales *family* and *friends* (relational subscales), and fair to good on the subscales *tolerance for affects*, *insight* and *problem-solving capacity* (dynamic subscales).

The two raters at pre-treatment differed in rating the relational subscales and the dynamic subscales. They tended to rate respectively relational and dynamic scales higher or lower than the other rater. The difference was not so clear at post-treatment, although the interrater reliability did not change. The interviews regularly revealed whether the adolescents had been in therapy or not. The raters were therefore not totally blind regarding whether the evaluation was pre-treatment or post-treatment. However, the two raters were scoring pre- and post-treatment interviews randomly and the chronology of ratings is not parallel to the therapies.

With the average scores of three raters (evaluator and 2 raters), the IRR was good to excellent for all subscales. Høglend and colleagues in the adult FEST-study achieved good results for all single scales at pre-treatment as well as post-treatment with average scores of three raters [22]. In the present study, *insight* was, as in the adult study, the most difficult scale to rate reliably. In relation to psychoanalytic theory, dynamic insight is a measure related to subjective interpretation and understanding

Table 5 Changes from pre-treatment to post-treatment with average scores of 3 raters (evaluator and 2 raters), (n = 30)

Scale	Mean $\pm$ SD		Sig. (2-tailed)	Reliable change index (% of all cases)
	Pre-treatment	Post-treatment		
Family	63.3 $\pm$ 7.1	66.3 $\pm$ 7.1	< 0.001	17
Friendships	66.1 $\pm$ 7.6	70.2 $\pm$ 6.5	0.008	17
Tolerance for affect	56.8 $\pm$ 4.7	62.8 $\pm$ 5.7	< 0.001	33
Insight	59.3 $\pm$ 5.9	65.3 $\pm$ 5.8	< 0.001	30
Problem-solving capacity	59.8 $\pm$ 5.1	64.7 $\pm$ 5.9	< 0.001	30
PFS mean	60.8 $\pm$ 5.2	66.0 $\pm$ 5.3	< 0.001	40
GAF	59.2 $\pm$ 5.5	67.1 $\pm$ 7.4	< 0.001	40
BDI	29.2 $\pm$ 7.3	15.1 $\pm$ 12.5	< 0.001	74
GSI	1.31 $\pm$ 0.5	0.81 $\pm$ 0.5	< 0.001	60

PFS Psychodynamic Functioning Scales, GAF Global Assessment of Functioning, BDI Beck Depression Inventory, GSI Global Severity Index

Table 6 Pearson's correlations of PFS subscales (n = 66), GAF and GSI

Variable	1	2	3	4	5	6	7	8
1. Family	–							
2. Friendships	0.52**	–						
3. Tolerance for affects	0.55**	0.65**	–					
4. Insight	0.42**	0.47**	0.77**	–				
5. Problem-solving capacity	0.46**	0.62**	0.81**	0.73**	–			
6. PFS mean	0.75**	0.83**	0.90**	0.80**	0.85**	–		
7. GAF (n = 47)	0.48**	0.48**	0.74**	0.63**	0.73**	0.72**	–	
8. GSI (n = 57)	– 0.33*	– 0.08	– 0.32*	– 0.04	– 0.41**	– 0.29*	0.39**	–

PFS Psychodynamic Functioning Scales, GAF Global Assessment of Functioning, GSI Global Severity Index

* Correlation is significant at the 0.05 level (2-tailed)

** Correlation is significant at the 0.01 level (2-tailed)

of symptoms, vulnerabilities and strengths. Adolescents progress at varying rates in developing their abstract and reflective thinking ability. They may also be more reluctant to talk about the past and link today's problems with past experiences. Only fair to good reliability was achieved also on the scale *problem-solving capacity*. Some adolescents may be able to apply logical operations long before they are able to apply them to personal dilemmas. When emotional issues arise, they often interfere with the young's ability to think in more complex ways. The ability to consider possibilities, as well as facts, may influence decision-making, in either positive or negative ways. This might result in incoherent information and descriptions from the adolescents themselves.

The ratings of subscales were based on the original English version of PFS. Although the dynamic interviews with anchor points relating to each subscale were conducted in Norwegian, i.e. the mother tongue of the evaluators and raters, the use of scales in English might influence the ratings and the interrater reliability.

The dynamic scales seemed harder to score in agreement than the relational scales. However, the changes from pre- to post-treatment were larger for the dynamic scales. An explanation might be that more nuanced post-treatment information contributed to relatively lower ratings of relationships to family and friends. However, the larger change on dynamic scales would be consistent with the concepts of dynamic psychotherapy. It might be explained by the possibility that the patients had been better acquainted with the concepts of therapy and that the therapy mirrored new ways of understanding and talking about perceived problems. The time-period during therapy is though important time for development. From the post-treatment evaluations available, the scales seem to be sufficiently fine-graded to capture statistically and clinically significant pre-post changes during time-limited psychodynamic psychotherapy with depressed adolescents. However, because of small sample size the results may be unstable.

The dynamic scales seem to measure a construct that may prove discriminable from general symptoms and

dysfunctions. The PFS mean value and GAF correlated strongly. The correlation with GSI (global symptoms) was weak.

The reliability change estimates for individual scales in our study is similar to the adult study. This is however true for the PFS. The RCI was larger for the self-reported symptom scores (GSI) in our study. The mean GSI was higher at pre-treatment for adolescents (1.3 vs 1.0). It may indicate that adolescents on self-reports tend to be more extreme in reporting symptoms compared to adults. The evidence base for psychodynamic psychotherapy for children and adolescents is building up [45], but e.g. there are few instruments for assessment of intrapsychic processes specific to the treatment. Outcome evaluation forms may add information to the therapists about their effectiveness and also be of value to researchers examining underlying mechanisms to explain outcome [46].

Although the PFS is used as an outcome measure, it might be useful in further process-outcome studies regarding mechanisms and moderators through which treatment interventions operate [47]. Variations in outcome seem to be influenced by patient characteristics and by the therapist variables and context factors [48–50]. PFS has no parallel forms to compare patient's and therapist's ratings, often used in process research (e.g. the rather newly developed Individual Therapy Process Questionnaire (ITPQ) [51]). Observer-ratings might still be a strength. In other studies of depression, both clinician-rated and self-reported instruments are recommended, although primarily related to symptoms [52, 53].

Reliability is important for outcome assessment. Although an assumption that stable and durable changes in personality characteristics are the most difficult to detect, changes in less stable variables such as mood and affect may have larger effect sizes in response to treatment—but lower reliability estimates. Following measurements administered during and after treatment with statistically significant differences would implicate measures sensitive to change. However, if not compared to other outcome measures there might be a lack of evidence that the real change occurred. Assessment of change makes further investigation of change possible and contributes to the field of research linking to clinical utility in the ideal personalization of psychotherapy treatment for adolescents. The ratings on PFS are based on the last 3–4 months and therefore the instrument might serve best as a follow-up in therapies lasting 6 months or more.

In plenary meetings trainees and students were often present. They were seldom the “outliers” and we were surprised how relatively easy it was for them to grasp the concept of the rating scales. The interrater reliability in

this study on adolescents suggests that the PFS could be used with adolescents in psychotherapy by experienced clinicians without extensive training.

Romantic and/or intimate relationships are important to adolescents and this scale should be considered revised and adjusted for adolescents to capture the ability to establish and stay in intimate mutual relationships.

Limitations

The small sample size and the fact that data from only half of the included patients was available for post-treatment analysis are both limitations of this study. The interrater reliability of additional raters attending plenary meetings are not analysed and reported. However, preliminary results are promising. Analyses on a full dataset would improve the statistical power.

Conclusion

Based on the interrater reliability (IRR) results in our study, the PFS could be recommended for use in psychotherapy with adolescents by experienced clinicians without extensive training. The IRR was good to excellent for all five subscales with the average scores of three raters. The scales seem to capture statistically and clinically significant changes. However, the IRR on the dynamic subscales *tolerance for affects*, *insight* and *problem-solving capacity* indicates that subscales of PFS might be considered revised or adjusted for adolescents.

Additional file

Additional file 1. Scales developed to assess change in dynamic psychotherapy.

Abbreviations

PFS: Psychodynamic Functioning Scales; FEST-IT: First Experimental Study of Transference Work-In Teenagers; GAF: Global Assessment of Functioning; CBT: cognitive behavioural therapy; STPP: short-term psychoanalytical psychotherapy; GSI: Global Severity Index; CAHMS: Child and Adolescent Mental Health Services; SDQ: Strengths and Difficulties Questionnaire; SIS-CA: Severity of Impairment Score for Children and Adolescents; CGAS: Children's Global Assessment Scale; GBOM: Goal Based Outcome Measure; CBCL: Child Behaviour Checklist; BIS: Brief Impairment Scale; M.I.N.I.: Mini International Neuropsychiatric Interview; SIDP-IV: Structured Interview for DSM-IV Personality; SCL-90: Symptom Checklist-90; BDI(-II): Beck Depression Inventory; IRR: interrater reliability; ICC: Intraclass Correlation Coefficient; RCI: Reliable Change Index; CI: confidence interval; FEST: First Experimental Study of Transference Interpretations.

Authors' contributions

EN is the first author of this study and has the main responsibility for analyses of data as well as the writing of the present manuscript. RU is the principal investigator in FEST-IT. PH is the principal investigator in FEST. SA is the clinical director in FEST. SA, PT, RU, H-SJD and EN have participated in providing and analysing treatment data. PH, AT and JE also contributed with statistics, drafting and revising the manuscript. All authors read and approved the final manuscript.

Author details

¹ Division of Mental Health and Addiction, University of Oslo, Oslo, Norway. ² Research Unit, Division of Mental Health, Vestfold Hospital Trust, PO Box 2168, 3103 Tønsberg, Norway. ³ Research Unit, Division of Mental Health, Østfold Hospital Trust, PO box 300, 1714 Grålum, Norway. ⁴ Dragonveien 24, 1396 Billingstad, Norway. ⁵ The Erica Foundation, Stockholm, Sweden. ⁶ Department of Psychology, University of Oslo, Oslo, Norway.

Acknowledgements

A special thanks to everyone involved in the FEST-IT study including patients, therapists and raters including Anne Grete Hersoug.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

Data from the First Experimental Study of Transference-Work-In Teenagers (FEST-IT) was used. Trial registration: ClinicalTrials.gov Identifier: NCT01531101. The data set supporting the results of this article is available on request from the last author, Randi Ulberg.

Consent for publication

The ethical approval and consent to participate included consent to publish. Patient material and data collected were accepted for use in research and publishing as well as teaching.

Ethics approval and consent to participate

The Regional Ethics Committee for health region in Norway approved the study protocol and the information given to the patients (REK: 2011/1424 FEST-IT). Written informed consent was obtained from each participant.

Funding

This study was supported by grants from The University of Oslo, Vestfold Hospital Trust, the MRK Foundation, Josef and Halldis Andresen's Foundation and Solveig and Johan P. Sommer's Foundation.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 31 January 2018 Accepted: 20 July 2018

Published online: 30 July 2018

References

- Green J. Editorial: process to progress? Investigative trials, mechanism and clinical science. *J Child Psychol Psychiatry*. 2015;56(1):1–3. <https://doi.org/10.1111/jcpp.12377>.
- Von Below C. When psychotherapy does not help...and when it does: lessons from young adults' experiences of psychoanalytic psychotherapy. Stockholm: Stockholm University; 2017.
- Arain M, Haque M, Johal L, Mathur P, Nel W, Rais A, et al. Maturation of the adolescent brain. *Neuropsychiatr Dis Treat*. 2013;9:449–61. <https://doi.org/10.2147/NDT.S39776>.
- Reyna VF, Chapman SB, Dougherty MR, Confrey JE. The adolescent brain: Learning, reasoning, and decision making. Washington, DC: American Psychological Association; 2012.
- O'Keeffe S, Martin P, Goodyer IM, Wilkinson P, Consortium I, Midgley N. Predicting dropout in adolescents receiving therapy for depression. *Psychother Res*. 2017. <https://doi.org/10.1080/10503307.2017.1393576>.
- Ormhaug SM, Jensen TK. Investigating treatment characteristics and first-session relationship variables as predictors of dropout in the treatment of traumatized youth. *Psychother Res*. 2018;28(2):235–49. <https://doi.org/10.1080/10503307.2016.1189617>.
- Goodyer IM, Reynolds S, Barrett B, Byford S, Dubicka B, Hill J, et al. Cognitive behavioural therapy and short-term psychoanalytical psychotherapy versus a brief psychosocial intervention in adolescents with unipolar major depressive disorder (IMPACT): a multicentre, pragmatic, observer-blind, randomised controlled superiority trial. *The Lancet Psychiatry*. 2016. [https://doi.org/10.1016/S2215-0366\(16\)30378-9](https://doi.org/10.1016/S2215-0366(16)30378-9).
- Midgley N, O'Keeffe S, French L, Kennedy E. Psychodynamic psychotherapy for children and adolescents: an updated narrative review of the evidence base. *J Child Psychother*. 2017. <https://doi.org/10.1080/0075417x.2017.1323945>.
- Crits-Christoph P, Connolly Gibbons MB, Mukherjee D. Psychotherapy process-outcome research. In: Lambert MJ, editor. *Bergin and Garfield's handbook of psychotherapy and behavior change*. Hoboken: Wiley; 2013.
- Johansson P, Hoglend P, Ulberg R, Amlo S, Marble A, Bogwald KP, et al. The mediating role of insight for long-term improvements in psychodynamic therapy. *J Consult Clin Psychol*. 2010;78(3):438–48. <https://doi.org/10.1037/a0019245>.
- Shedler J. The efficacy of psychodynamic psychotherapy. *Am Psychol*. 2010;65(2):98–109. <https://doi.org/10.1037/a0018378>.
- Abbass AA, Rabung S, Leichsenring F, Refseth JS, Midgley N. Psychodynamic psychotherapy for children and adolescents: a meta-analysis of short-term psychodynamic models. *J Am Acad Child Adolesc Psychiatry*. 2013;52(8):863–75. <https://doi.org/10.1016/j.jaac.2013.05.014>.
- Johnston C, Gowers S. Routine outcome measurement: a survey of UK child and adolescent mental health services. *Child Adolesc Ment Health*. 2005;10(3):133–9. <https://doi.org/10.1111/j.1475-3588.2005.00357.x>.
- Deighton J, Croudace T, Fonagy P, Brown J, Patalay P, Wolpert M. Measuring mental health and wellbeing outcomes for children and adolescents to inform practice and policy: a review of child self-report measures. *Child Adolesc Psychiatry Ment Health*. 2014;8:14. <https://doi.org/10.1186/1753-2000-8-14>.
- Fonagy P, Target M, Steele H, Steele M. *Reflective-functioning manual version 5.0 for application to adult attachment interviews*. London: University College London; 1998. p. 161–2.
- Lingiardi V, McWilliams N, Bornstein RF, Gazzillo F, Gordon RM. The Psychodynamic Diagnostic Manual Version 2 (PDM-2): assessing patients for improved clinical practice and research. *Psychoanal Psychol*. 2015;32(1):94–115. <https://doi.org/10.1037/a0038546>.
- Arbeitskreis OPDKJ. OPD-KJ-2-Operationalisierte Psychodynamische Diagnostik im Kindes-und Jugendalter. Grundlagen und Manual. Bern: Huber; 2013.
- Shedler J, Westen D. The Shedler–Westen assessment procedure (SWAP): making personality diagnosis clinically meaningful. *J Pers Assess*. 2007;79(1):41–55.
- DeWitt KN, Milbrath C, Simon NM. Wallerstein's Scales of Psychological Capacities: a clinically useful measure of character change. *Psychoanal Psychol*. 2018;35(1):115–26. <https://doi.org/10.1037/pap0000139>.
- Greenfield B, Filip C, Schiffrin A, Bond M, Amsel R, Zhang X. The Scales of psychological capacities: adaptation to an adolescent population. *Psychother Res*. 2012;23:232–46.
- Wells MG, Burlingame GM, Lambert MJ, Hoag MJ, Hope CA. Conceptualization and measurement of patient change during psychotherapy: development of the Outcome Questionnaire and Youth Outcome Questionnaire. *Psychotherapy*. 1996;33(2):275–83. <https://doi.org/10.1037/0033-3204.33.2.275>.
- Hoglend P, Bøgdal KP, Amlo S, Heyerdahl O, Sørbye O, Marble A, et al. Assessment of change in dynamic psychotherapy. *J Psychother Pract Res*. 2000;9(4):190–9.
- Jones E. Manual for the psychotherapy process Q-set. Unpublished manuscript. Berkeley: University of California; 1985.
- Jones EE, Windholz M. The psychoanalytic case study: toward a method for systematic inquiry. *J Am Psychoanal Assoc*. 1990;38(4):985–1015. <https://doi.org/10.1177/000306519003800405>.
- Ablon JS, Jones EE. On analytic process. *J Am Psychoanal Assoc*. 2005;53(2):541–68. **discussion 69–78**.
- Bøgdal KP, Dahlbender RW. Procedures for testing some aspects of the content validity of the Psychodynamic Functioning Scales and the Global Assessment of Functioning Scale. *Psychother Res*. 2004;14(4):453–68. <https://doi.org/10.1093/ptr/kph038>.
- Aas IH. Guidelines for rating Global Assessment Of Functioning (GAF). *Ann Gen Psychiatry*. 2011;10:2. <https://doi.org/10.1186/1744-859x-10-2>.
- Derogatis L. SCL-90-R: Administration, scoring and procedure manual. Towson: Clin Psychom Research; 1983.
- Ulberg R, Hersoug AG, Hoglend P. Treatment of adolescents with depression: the effect of transference interventions in a randomized

- controlled study of dynamic psychotherapy. *Trials*. 2012;13:159. <https://doi.org/10.1186/1745-6215-13-159>.
30. Midgley N, Rhode M, Rustin M, Hughes C, Cregeen S. IMPACT group: short-term psychoanalytic psychotherapy (STPP) for adolescents with moderate or severe depression: a treatment manual. Cambridge: IMPACT trial Office; 2010.
 31. Grøholt B. Medikamentell behandling av depresjon hos barn og ungdom. *Tidsskrift Nor Legeforen*. 2011;131(22):2243–5. <https://doi.org/10.4045/tidsskr.10.0282>.
 32. Aas M. Global Assessment of Functioning (GAF): properties and frontier of current knowledge. *Ann Gen Psychiatry*. 2010. <https://doi.org/10.1186/1744-859x-9-20>.
 33. Hilsenroth MJ, Ackerman SJ, Blagys MD, Baumann BD, Baity MR, Smith SR, et al. Reliability and validity of DSM-IV axis V. *Am J Psychiatry*. 2000;157(11):1858–63. <https://doi.org/10.1176/appi.ajp.157.11.1858>.
 34. Ryttilä-Manninen M, Fröjd S, Haravuori H, Lindberg N, Marttunen M, Kettunen K, et al. Psychometric properties of the symptom checklist-90 in adolescent psychiatric inpatients and age-and gender-matched community youth. *Child Adolesc Psychiatry Ment Health*. 2016;10:23. <https://doi.org/10.1186/s13034-016-0111-x>.
 35. Derogatis LR, Unger R. Symptom checklist-90-revised. Hoboken: Wiley; 2010.
 36. Beck AT, Steer RA, Ball R, Ranieri W. Comparison of Beck Depression Inventories—IA and -II in psychiatric outpatients. *J Pers Assess*. 1996;67(3):588–97. https://doi.org/10.1207/s15327752jpa6703_13.
 37. Wang YP, Gorenstein C. Psychometric properties of the Beck Depression Inventory-II: a comprehensive review. *Revista brasileira de psiquiatria*. 2013;35(4):416–31. <https://doi.org/10.1590/1516-4446-2012-1048>.
 38. Malan D. The frontier of brief psychotherapy. New York: Plenum Press; 1976.
 39. Sifneos PE. Short-term anxiety-provoking psychotherapy: a treatment manual. New York: Plenum Press; 1992.
 40. Shrout PE, Fleiss JL. Intraclass correlations: uses in assessing rater reliability. *Psychol Bull*. 1979;86(2):420–8.
 41. Cicchetti D. Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instrument in psychology. *Psychol Assess*. 1994;6:284.
 42. Fleiss JL. Statistical methods for rates and proportions. Hoboken: Wiley; 1981.
 43. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33(1):159–74.
 44. Jacobson NS, Truax P. Clinical significance: a statistical approach to defining meaningful change in psychotherapy research. *J Consult Clin Psychol*. 1991;59(1):12–9.
 45. Midgley N, Kennedy E. Psychodynamic psychotherapy for children and adolescents: a critical review of the evidence base. *J Child Psychother*. 2011;37(3):232–60.
 46. Kazdin AE. Understanding how and why psychotherapy leads to change. *Psychother Res*. 2009;19(4–5):418–28. <https://doi.org/10.1080/10503300802448899>.
 47. Kazdin AE. Mediators and mechanisms of change in psychotherapy research. *Annu Rev Clin Psychol*. 2007;3:1–27. <https://doi.org/10.1146/annurev.clinpsy.3.022806.091432>.
 48. Beutler LE. Making science matter in clinical practice: redefining psychotherapy. *Clin Psychol Sci Pract*. 2009;16(3):301–17.
 49. Beutler LE, Someah K, Kimpara S, Miller K. Selecting the most appropriate treatment for each patient. *Int J Clin Health Psychol*. 2016;16(1):99–108. <https://doi.org/10.1016/j.ijchp.2015.08.001>.
 50. Wampold BE. The Great psychotherapy debate: models, methods, and findings. Mahwah: L. Erlbaum Associates; 2001.
 51. Mander J. The individual therapy process questionnaire: development and validation of a revised measure to evaluate general change mechanisms in psychotherapy. *Clin Psychol Psychother*. 2015;22(4):328–45. <https://doi.org/10.1002/cpp.1892>.
 52. Cuijpers P, Li J, Hofmann SG, Andersson G. Self-reported versus clinician-rated symptoms of depression as outcome measures in psychotherapy research on depression: a meta-analysis. *Clin Psychol Rev*. 2010;30(6):768–78. <https://doi.org/10.1016/j.cpr.2010.06.001>.
 53. Uher R, Perlis RH, Placentino A, Dernovšek MZ, Henigsberg N, Mors O, et al. Self-report and clinician-rated measures of depression severity: can one replace the other? *Depress Anxiety*. 2012;29(12):1043–9. <https://doi.org/10.1002/da.21993>.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

TECHNICAL ADVANCE

Open Access

Relational interventions in psychotherapy: development of a therapy process rating scale

Randi Ulberg^{1,2*} , Elisabeth Ness², Hanne-Sofie Johnsen Dahl², Per Andreas Høglend¹, Kenneth Critchfield³, Phelix Blayvas⁴ and Svein Amlø⁴

Abstract

Background: In psychodynamic psychotherapy, one of the therapists' techniques is to intervene on and encourage exploration of the patients' relationships with other people. The impact of these interventions and the response from the patient are probably dependent on certain characteristics of the context in which the interventions are given and the interventions themselves. To identify and analyze in-session effects of therapists' techniques, process scales are used. The aim of the present study was to develop a simple, not resource consuming rating tool for in-session process to be used when therapists' interventions focus on the patients' relationships outside therapy.

Methods: The present study describes the development and use of a therapy process rating scale, the Relational Work Scale (RWS). The scale was constructed to identify, categorize and explore therapist interventions that focus on the patient's relationships to family, friends, and colleagues. Relational Interventions and explore the impact on the in-session process. RWS was developed with sub scales rating timing, content, and valence of the relational interventions, as well as response from the patient. For the inter-rater reliability analyzes, transcribed segments (10 min) from 20 different patients were scored with RWS by two independent raters. Two clinical vignettes of relational work are included in the paper as examples of how to rate transcripts from therapy sessions with RWS.

Results: The inter-rater agreement on the RWS items was good to excellent.

Conclusion: Relational Work Scale might be a potentially useful tool to identify relational interventions as well as explore the interaction of timing, category, and valence of relational work in psychotherapies. The therapist's interventions on the patient's relationships with people outside therapy and the following patient-therapist interaction might be explored.

Trial registration: First Experimental Study of Transference-interpretations (FEST307/95)
Registration number: ClinicalTrials.gov Identifier: NCT00423462.

Keywords: Relational interventions, Transference, Psychodynamic, In-session process

Abbreviation: FEST, First experimental study of transference-interpretations; IRR, Inter rater reliability; QOR, Quality of object relation score; RI, Relational interventions; RW, Relational work; RWS, Relational work scale; TI, Transference intervention; TW, Transference work; TWS, Transference work scale

* Correspondence: Randi.Ulberg@medisin.uio.no

¹Institute of Clinical Medicine, Division of Mental Health and Addiction, University of Oslo, Box 85 Vinderen, 0319 Oslo, Norway

²Research Unit, Division of Mental Health and Addiction, Vestfold Hospital Trust, PO Box 2169, 3125 Tønsberg, Norway

Full list of author information is available at the end of the article

© 2016 The Author(s). **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Background

There is a growing emphasis on the development of personalized treatment for mental disorders [1]. Several studies have shown effect of different psychotherapies for the most frequent disorders as anxiety and depression. Two of the mostly used and researched therapy modes for these disorders are Cognitive Behavior Therapy (CBT) and Psychodynamic Therapy (PDT). Both CBT and PDT are repetitively shown to be more effective than no treatment. Different psychotherapy modalities work, on average, equally well [2–4]. However, little is known of what kind of patients would profit most from different psychotherapy modes and how different therapist techniques and interventions contribute to the patient-therapist interaction. Also, it has proved difficult to find ways to examine this in-session process. Therefore, there is a need to develop more methods to reveal what kind of treatment and what treatment techniques work best for different patients and how the various psychotherapy modes work to promote the best possible treatment effect (outcome) for each patient. Understanding the process of therapy and the impact on outcome will increase our ability to develop further our treatments, to make them more effective, and to individualize them for the needs of specific individuals [5].

Process-outcome research that examines the action of specific psychotherapies would help developing more personalized treatment. *Psychotherapy process* is primarily the actions, experiences and relatedness of patient and therapist in therapy sessions [6]. Change process might be referred to as “short-term outcome”. *Outcome* is improvements in patient-presenting problems, symptom, and functioning during therapy. Thus, process studies exploring what happens during psychotherapy sessions (e.g. single case studies, qualitative and quantitative process studies) are essential to understand the nature of psychotherapy and would be helpful when outcome is also part of the empirical investigation. On this background, there is a need for more theoretically informed quantitative assessment tools enabling researchers to explore the in-session process in psychotherapy.

In dynamic psychotherapy a theoretical assumption is that people project representational aspects of important others onto relationship with other people at present. Therefore, interpreting the patient’s relationships with other people are key components to enhance change. Those interventions might be categorized in two groups: 1) Transference interventions (TI) are when the therapist focuses on the patient’s relationship to and experience of the therapist. 2) Relational Interventions (RI) are when the therapist focuses on the patient’s relationships outside therapy. Different definitions of RI and TI are given in the literature. Some definitions of RI for example, will include TI as a variant of RI. The different definitions of these constructs are, however, beyond the scope of this paper. For

our purposes, RI and TI are defined in concrete behavioral terms as an explicit interpretive reference to the patient’s ongoing relationship with other people outside therapy or the relationship with the therapist (Table 1).

One example of a relational intervention might be: “You really don’t want to go to that party while your boyfriend does. Since you don’t agree with him, you feel guilt in preventing him from going. During childhood, you had to get out of your way in order not to prevent your mother reaching her goals in life. It might be that your previous feeling in relation to your mother is connected with your present feeling in relation to your boyfriend.” An extension of the RI might include an explicit reference to the patient–therapist interaction (TI) by adding: “Perhaps you accepted my suggestion of a short and time-limited therapy because you felt guilty to occupy my time here?”

In psychodynamic theory especially interpretation of the patient’s relationship to and experience of the therapist (TI), has been considered an important and frequently used specific technique [7, 8]. However, this assumption might be proved wrong as a general guideline for all patients. Transference phenomenon is also present in non-dynamic treatments, but the research in this field is sparse [9]. Patients may improve differently according to various characteristics. Some recent studies indicate that although patients on average might respond equally well in dynamic psychotherapy with or without TI, some sub groups of patients respond better when the focus is on the patient’s relations outside therapy [10, 11]. Working with the

Table 1 Five categories of Relational Interventions and Transference Interventions defined in FEST^a

Relational interventions (RI):

1. The therapist addresses interpersonal transactions with other (s)^b.
2. The therapist actively encourages the patient to explore thoughts and feelings about his/her relationships to other (s)^b including their style and behavior.
3. The therapist encourages the patient to discuss how other (s)^b might feel or think about the patient.
4. The therapist makes interpretive linking of dynamic elements (conflicts) in the patient’s relationships with other (s)^b.
5. The therapist attempts to explore interpersonal repetitive patterns with important other (s)^b and with parental figures.

Transference Interventions (TI):

1. The therapist addresses transactions in the patient–therapist relationship.
2. The therapist encourages exploration of thoughts and feelings about the therapy and therapist’s style and behavior.
3. The therapist encourages patients to discuss how they believed the therapist might feel or think about them.
4. The therapist includes him-/herself explicitly in interpretive linking of dynamic elements (conflicts), direct manifestations of transference, and allusions to the transference.
5. The therapist was to interpret repetitive interpersonal patterns (including genetic interpretations) and link these patterns to transactions between the patient and the therapist.

Note. ^aFirst Experimental Study of Transference-interpretations (FEST). ^bother (s) is/are everyone except the therapist (i.e. friends, relatives, colleagues)

transference (Transference work; TW) more than working with relations outside therapy (Relational work; RW) is expected to trigger defense mechanisms [12, 13]. However, as far as we know, there has been little empirical exploration of the different impact for different patients on the in session process and treatment outcome of therapy with RI and TI combined versus therapy only focusing on RW.

To identify and analyze in-session effects of therapist technique and empirically establish links to outcome, we have to rely on adequate process scales [7, 14]. Different instruments for identification of therapist techniques have been developed and should be acknowledged [15]. Previously the development and use of a therapy process rating scale (Transference Work Scale; TWS) has been described and a manual was published [15, 16].

TWS was constructed to identify, categorize and explore in-session work with the transference. TWS has sub scales that rate timing, content, and valence of the transference interventions, as well as response from the patient. The inter-rater agreement on the TWS items was good to excellent. TWS might be a useful tool to explore the interaction of timing, content, and valence of transference work in predicting in-session patient response as well as treatment outcome.

Aim

To be able to explore the in-session process in non-transference and non-dynamic therapies, the aim of the present study was, to develop a simple measure specially designed to identify the five RI categories and explore the timing, content, and valence of each intervention, as well as the response from the patient. Furthermore, the aim was to test the inter-rater reliability of the different items. Use of the scale is illustrated with two clinical vignettes.

Method

First experimental study of transference-interpretations

Relational Work Scale (RWS) was developed by the First Experimental Study of Transference-interpretations (FEST) research group. Clinical examples in the present paper are from therapies in FEST. The Regional Ethics Committee for health region 1 in Norway approved the study protocol and the information given to the patients. Patient material and data collected including case material were accepted for use in research and publishing as well as teaching. Written informed consent was obtained from each participant. Data are anonymized. It is not possible to reconnect the clinical material to patient identity.

The purpose of the First Experimental Study of Transference-interpretations [10, 17] was to measure the effects of Transference Work (TW) in dynamic psychotherapy. FEST was a randomized controlled trial where one hundred patients seeking psychotherapy for depression, anxiety, and personality disorders, were allocated to

psychodynamic psychotherapy with low to moderate levels of TIs (the transference group; $N = 52$) or psychodynamic psychotherapy with no TIs, only RIs (the comparison group; $N = 48$). Patients with psychosis, bipolar illness, organic mental disorder, or substance abuse were excluded. The random assignment of patients was conducted after the pre-treatment ratings were completed. Only the patients' therapists learned the result of the random assignment procedure. Patients were assigned to one of the seven therapists, depending on availability. The clinical evaluators and therapists consisted of six psychiatrists and one clinical psychologist, all of whom had 10–25 years of experience in practicing psychodynamic psychotherapy. In the pilot phase of this study, the therapists were trained with group supervision for as long as 4 years in order to be able to administer treatment with a moderate frequency of transference interpretations (1–3 per session) as well as treatment without such interpretations with equal ease and mastery. All therapists treated patients from both groups.

The treatment was 45 min once weekly, maximum 40 sessions (1 year) [18]. In both treatment modes the therapists used RIs focusing on the patient's relations outside of therapy. The specific techniques (i.e. categories of TI) were prescribed, for the transference group only (Table 1).

To quantify the use of the five different RI and TI categories in the two treatment groups, a global rating method using a 5-point Likert scale ranging from 0 (not at all) to 4 (very much) was used. This was chosen rather than rating the exact frequency of interventions [19, 20]. The use of the techniques differed significantly between the treatment groups. The average score of TI was 1.7 ($SD = 0.7$) in the transference group, and 0.1 ($SD = 0.2$) in the comparison group ($t = 14.8$, $df = 58.2$, $p < 0.0005$) [10].

The outcome measures were the Psychodynamic Functioning Scales [21], Inventory of Interpersonal Problems-Circumplex version [22], Global Assessment of Functioning Scale (GAF; Diagnostic and statistical manual of mental disorders, 1987), and Symptom Checklist-90-R [23].

Based on a psychodynamic interview modified after Malan [8] and Sifneos [24], at least three clinicians rated PFS and GAF (pre-, posttreatment, and at 1- and 3-year follow-ups) and at pre-treatment also the patient's relational functioning with Quality of Object Relations Scale (QOR) [25]. The QOR score was determined from the following three 8-point scales: evidence of at least one stable and mutual interpersonal relationship in the patient's life, history of adult sexual relationships, and history of nonsexual adult relationships. The QOR measures the patient's lifelong tendency to establish certain kinds of relationships with others, from mature to primitive. Inter-rater reliability for average scores of three raters was 0.84 in this study. The predetermined cutoff score for high versus low QOR was 5.00 [14].

Both treatment groups showed statistically significant change from pre-treatment to 3 year follow-up. Contrary to expectation, no significant between-group differences were revealed in FEST. Moderator analyses showed that patients with a life-long pattern of poor relational functioning measured with the QOR [17] and/or patients with personality disorders profited more in therapy with TI than without TI [10, 17, 26]. This was especially true for women, while men responded better to therapy without TI compared to women [11].

Development of relational work scale (RWS)

Based on psychodynamic theory [13, 27, 28] and years of combined clinical practice and research experience with psychodynamic psychotherapy, the FEST-research group has developed the TWS and subsequently the RWS. Several previous adherence- and process measures have influenced the development process, and resemblance is intentional. However, the main source was the Manual for process rating from FEST [20] and the TWS [15]. While TWS was developed for identifying and categorizing TI and the interaction between therapist and patient in the Transference work, RWS was developed in order to identify and categorize RI's and analyze the process between therapist and patient after an RI. RWS has items on the timing, content, and valence of each RI, as well as the response from the patient modelled as in the TWS. Five researchers and clinicians have participated in the theoretical discussions on constructions of items in RWS.

Two independent raters scored transcripts during the scale developmental period to determine inter-rater agreement on different items. The two raters one female rater, M.D., Ph.D. with 30 years of clinical experience and one male rater, M.D. with more than 50 years of clinical experience. Both were specialists in psychiatry. However, they were trained in different psychoanalytic/psychodynamic institutes and had dissimilar work experience. The male rater had participated in FEST from the beginning of the study while the other joined the research group after the four year study evaluations for all patients were completed.

Items concerning identification and categorization of RI were based on the five categories of RI defined in FEST (Table 1). Predefined non-transference interventions [20] and comparable process exploring items as used in TWS were included in RWS (Additional file 1).

Items assessing the degree to which RIs connect naturally to the preceding clinical material and how precise and striking the RI is [28] and items describing content of therapist interventions, such as dynamic conflict components (anxiety, defense, impulse/motives) and person components (parents, others) were included [8]. Also RWS was developed with items concerning the degree of whether the therapist was challenging or supportive,

and the patient's attempts to avoid themes, level of emotional engagement, and whether the patient showed associations or self-reflections in the response, were included [28].

Items in RWS were scored using Likert scales; 0, 1 (not at all, low degree) to 4 (high degree). Items identifying presence or absence of the five categories of RI (Item 8–12) were rated with “Yes” or “No”. The segments used for the pilot project were from previously transcribed sessions for analyses of gender differences in FEST [29]. Since the two raters had previously rated transcripts for the inter-rater analyses with TWS, and the rating method using RWS was comparable, little additional training was needed. However, first, the raters rated 10 segments from different therapies not included in the IRR analyses. During the training, ratings of one segment were discussed before next segment from another therapy was rated. No further training was conducted before rating of the segments used in the IRR-analyses.

Ratings with RWS for inter-rater reliability analyses

Ten min segments from the middle of the mid-therapy session (session 16) were chosen to explore the RW. Transcripts from 20 of the patients in the non-transference group (10 low QOR women and 10 high QOE men), were used. Each turn of talk in the dialogue between the patient and the therapist was given consecutive index numbers in the transcript [22].

The version of RWS used for the IRR-analysis comprised 26 items (Table 2), and included two items concerning timing of the RI with the highest category score in the transcript.

These two items are included in the table reporting the results from the inter-rater reliability analyses and marked I (Is the timing of the RI with the highest category score in the transcript naturally connected with the preceding clinical material?) and II (How and precise and striking is the therapist's RI?) in Table 2.

Statistical analyses

The IRR was estimated with the coefficient Kappa [30] for category scores and with the Intra Class Correlation (ICC) [31] (two-way random consistency) for ordinal scores. According to Landis and Koch kappa values greater than 0.75 are interpreted as excellent agreement, 0.60–0.75 as good, between 0.40 and 0.60 as fair and below 0.40 indicates poor agreement [30]. The statistical analyses were done using SPSS version 18 SPSS Inc., 2011).

Results

The RWS showed good inter-rater reliability for almost every item. The raw *N* and frequencies for occurrence of each of the categorical variables are listed in Table 2. The raters identified RI in 17 of the 20 segments

Table 2 Rater reliability of 2 raters for the Relational Work Scale (RWS) scored on one segment from the middle of a mid-therapy session of the patients in the non-transference group ($N = 20$)^a

Item			Rater 1 Mean (SD) <i>N</i> (%)	Rater 2 <i>N</i> (%)
1	RI ^b in transcript	Kappa	17 (85 %)	<i>N</i> = 17 (85 %)
Identification		ICC		
2	Beginning IRI ^d	S ^c	Mean (SD) 11.82 (10.67)	Mean (SD) 12.06 (10.84)
		Kappa	<i>N</i> = 17 (85 %)	<i>N</i> = 17 (85 %)
3	IRI first interaction	1	2 (10 %)	2 (10 %)
4	IRI last interaction	1	1 (5 %)	1 (5 %)
		ICC	<i>N</i> = 17 (85 %)	<i>N</i> = 17 (85 %)
5	Category IRI	S	Mean (SD) 2.53 (1.12)	Mean (SD) 2.17 (1.29)
Timing IRI				
6	IRI connect naturally	0.33	3.00 (0.76)	2.80 (0.86)
7	IRI precise/striking	0.82	2.60 (1.12)	2.73 (1.03)
Category RI		Kappa	<i>N</i> (%)	<i>N</i> (%)
8	Category 1	0.47	3 (17.65)	7 (41.18)
9	Category 2	0.77	15 (88.24)	14 (82.35)
10	Category 3	0.85	5 (29.41)	4 (23.53)
11	Category 4	0.76	3 (17.65)	2 (11.76)
12	Category 5	1	3 (17.65)	3 (17.65)
Content				
Timing high category ^e RI		ICC	<i>N</i> = 13 (65 %)	<i>N</i> = 13 (65 %)
		S	Mean (SD)	Mean (SD)
I ^f	Connect naturally	-0.16	3.23 (0.60)	3.08 (0.49)
II ^f	Precise/striking	0.73	2.46 (0.97)	2.54 (0.78)
		ICC	<i>N</i> = 17 (85 %)	<i>N</i> = 17 (85 %)
		S	Mean (SD)	Mean (SD)
13	T ^g Relation other	0.89	2.35 (1.11)	2.35 (0.10)
14	P ^h Relation other	0.96	2.82 (1.86)	2.71 (1.10)
15	T Relation parent	0.95	1.23 (1.35)	1.17 (1.33)
16	P Relation parent	0.96	0.76 (1.25)	0.76 (1.25)
17	Avoid themes	0.61	0.29 (0.69)	0.23 (0.56)
18	Symptoms T	0.63	1.35 (0.10)	1.47 (0.80)
19	Symptoms P	0.85	1.41 (0.80)	1.65 (0.79)
Valence				
20	Supportive	0.52	1.94 (0.83)	2.00 (0.87)
21	Challenging	0.66	2.11 (0.85)	1.82 (0.81)
Response				
22	Associations/self refl	0.89	2.29 (0.91)	2.35 (0.93)
23	Cooperative	0.64	2.65 (0.70)	2.47 (0.80)
24	Emotional involvement	0.62	2.47 (0.71)	2.24 (0.56)

Note ^a20 patients from FEST NT group; 10 high QOR men and 10 low QOR women. 17 segments ($n = 17$, 85 %) were found to contain Relational Interventions. There was complete agreement with regard to presence versus absence of Relational work in these 17 segments (Item 1 on RWS). ^bRelational Interventions (RI). ^cSingle measure (S). ^dInitial Relational Intervention (IRI). ^eTiming of the first Relational Intervention with the highest category score. Deleted from final version of RWS. ^fItem deleted from the final version of RWS. ^gTherapist focusing on (T). ^hPatient focusing on (P)

(kappa = 1). Except for Item 6 (whether the therapist's initial RI was connected naturally to the preceding clinical material, content and time line), and the Item marked "I" in Table 2 (whether therapist's RI with the highest category score in the transcript connect naturally to the preceding clinical material), the scoring agreement was good to excellent. On item 6 the IRR was low. On Item I the IRR was negative, however, the two raters scored nearly identical.

On most items, the ratings were normally distributed for both raters. Ratings on the items concerning whether the therapist (Item 13) or patient (Item 14) refer to the patient's relationship to parents, and whether the therapist points out that the patient attempts to avoid themes (Item 17) the percent of 0-scores was high for both raters; 41.2 and 47.1 % on Item 13 for rater 1 and 2 respectively, 64.7 and 82.4 % on Item 14 and 17 respectively for both raters.

Because of the negative IRR on one of the items concerning timing of the RI with the highest category score in the transcript, both Item I and Item II (Table 2) (Is the timing of the RI with the highest category score in the transcript naturally connected with the preceding material (I) and precise and striking (II) (Table 2) were deleted in the final version of RWS (Additional file 1).

During the rating process, examples of the five different categories of RIs were identified and are provided in Table 3.

Two vignettes illustrating the use of the relational work scale

Case 1 (Low QOR woman treated with TI)

Annie Lowland (not her real name) worked as a headmaster. She was 52 years, lived in a stable marriage with four grown up children. Annie Lowland grew up with three younger sisters. She was from the northern part of Norway. Her father was stable, but distant. Her mother was perceived as unpredictable and mean. In three periods during her first four years, Annie lived with relatives because her mother was "tired". When Annie was a grown up, her mother was treated for periodic alcohol abuse.

She emphasized that she wanted to raise and educate her children in another way than she was brought up herself. Her first child died, from meningo-encephalitis, as a toddler. Through recurrent episodes of depression, she accused herself for the boy's death. Three years ago she was successfully treated for breast cancer with breast conserving surgery. However, after the operation, she felt she had failed at work, as a mother, wife and woman. She felt insecure, with low self-esteem. She was depressed, felt her quality of life was low and hoped therapy would help.

Reflecting her interpersonal problems, Annie Lowland was rated with low QOR.

Table 3 Examples of Relational Interventions (RI)^a

RI Category 1: The therapist addresses interpersonal transactions with others without any linking to analogous transactions in session.	
Therapist	However you will keep up appearances. Pretending that everything is okay with you when you meet him.
RI Category 2: The therapist actively encourages the patient to explore thoughts and feelings about his/her relationships to other (s) including their style and behavior.	
Patient	Our children prefer being with my wife. It often seems fine and lively when they are together, while it often becomes awkward when they are with me.
Therapist	How do you feel about that experience?
Patient	It is sad when I think about these situations and what I lose.
RI Category 3: The therapist encourages the patient to discuss how other (s) might feel or think about the patient.	
Patient	When I unexpectedly come home to visit a Sunday, my mother would say: "it's been a long time not seeing you".
Therapist	Could it be an indication that she is happy? That she misses you more than that she blames you?
Patient	Yes, I think so. She actually is quite supportive. However, I don't really need to meet my parents as often now. It is more like I meet them when I want to. But if my relationship to my girlfriend ends, I think I would visit my parents. Then I will need some kind of comfort, as I always get there.
RI Category 4: The therapist makes interpretive linking of dynamic elements (conflicts) in the patient's relationships with other (s)	
Patient	He says he is very fond of me and would really like to have a relationship with me.
Therapist	So you are afraid of being let down, that he would lose interest in you when he gets to know you better?
Patient	Mmm mmm (confirms)
Therapist	This fear that people who matter to you would not bother about you. Did you always have this fear or did it come gradually?
Patient	I think I have had this feeling for a long time, as long as I can remember, but I have never really thought about it before.
RI Category 5: The therapist attempts to explore interpersonal repetitive patterns with important other (s) and with parental figures.	
Therapist	You say that your boss, in this case, is breaking the regulations. Still you do not confront him, rather you feel like it's your fault; that you have been leading him to it. You defend your boss and suppress your protest. Could it be helpful for you to see this as sort of an extension of the relationship with your father? When facing your father you did not share your opinions or views, that is, if you assumed that your father did not agree. You were afraid of being rejected and losing your father's strong support and positive recognition.
Patient	Yes. It is good to bring up my anger against my boss in this matter, because I put the blame on my anger easily and then I experience that he could not have acted in any other way and that it is almost my fault. It is of course not logical, but that is how I feel. Yes, it is a bit like with my father, I never really expressed my disagreement or criticism towards him. So what I feel now is discomfort, like a knot in my stomach, I want to put the whole case aside.

^aAll clinical examples are identified in the First Experimental Study of Transference-interpretations (FEST). Written informed consent was obtained from each participant

Course of treatment

Annie Lowland was randomized to dynamic psychotherapy without TI. She attended 39 sessions. The focus for the therapy was expected to be her relationship with her mother and her low self-esteem.

For a short example from the dialog late in therapy, please see Table 4. Table 4 illustrates how the IRR-raters would evaluate the presence and timing of RIs in this segment, as well as the themes in the dialogue and the patient's response.

Table 4 shows only a very short excerpt from a session and ratings of the segment are solely meant for illustrating the use of RWS. The two raters identified RIs of category 2 (first RI in transcript) and 5 (second RI). Timing is only rated for the second RI because the first RI is the first patient-therapist interaction in the transcript (Item 3). The first RI was delivered in a less supportive and challenging way rated 0 and 2 (average 1) and 2 and 4 (average 3) respectively by the two raters. The RI in category 5 might have been experienced as quite demanding by Annie Lowland. The therapist asked Annie to explore her interpersonal repetitive patterns towards her colleagues and also compared this pattern with how Annie experienced her mother's negative influence on Annie's self-image. Thus, the therapist focused on other

(average 2.5) and mother (2). The patient responded with associations/self-reflections, however, more freely after the first than the second RI (rated 3 and 1 respectively). She also seemed to be more cooperative and emotional involved after the first (3 vs. 4) than the second (1 vs. 2) RI. After the challenging category 5 intervention the patient's response was to a little degree focused on other (2) and Annie did not follow the therapist's intervention by exploring her relationship with her mother. After the second intervention, the patient seemed to be more distanced. She responded with talking more about others, was less emotional involved, focused more on her own symptoms, and was rated less cooperative.

After treatment, Annie Lowland vaguely described that she experienced improvement and fewer symptoms. She still struggled with her collegial relations at work. However, she described that her self-esteem in relation to her husband had improved a little.

Case 2 (High QOR man treated without TI)

Randolph Highland (not his real name) was a 43 year old geologist working in the petrol industry in Norway.

He married young and had two children 10 and 14 years old. Over the years, he felt more and more rejected by his wife. She was verbally strong and he responded by

Table 4 Segments with Relational Interventions (RI) from the patient-therapist dialogue in a session from late in therapy rated with Relational Work Scale

Individual speaking	Parts of dialogue	(RI): Category 1–5 Rated 0–4: (T): Timing (C): Content (V): Valence (R): Response
Therapist	Your relationships with your colleagues seem to have changed.	RI:2 C: Other (3) V: Supportive (0), Challenging (2)
Patient:	In my job, control and management is really important. However, I have a feeling that I exaggerate the control-part of my work. I don't just control and organize the school; I control the teachers in detail too. I don't like this side of myself.	C: Other (1) R: Associations/self reflections (3), Cooperative (3), Emotional (4)
Therapist	Maybe you feel that you're not any longer a colleague, but only a manager and boss. You might see similarities between yourself and the way your mother acted. She was controlling and interfered in others business. I wonder what impact the way you have experienced your mother has had on your self-image.	RI: 5 T: Connect natural (3) precise/striking (3) C: Other (2) Parents (2) V: Supportive (1), Challenging (4)
Patient:	It gives me a feeling of not being good enough. After the cancer operation I doubt my value as a woman even though I know it's irrational. However, when receiving positive response at work, I get a good feeling of being kind, clever and pretty.	C: Other (2), Symptoms (4) R: Associations/self reflections (1), Cooperative (1), Emotional (2)

withdrawing himself from discussions. He had two good mutual friendships and was highly valued as a colleague because he was always helpful and took on more than his share of the work load.

Randolph Highland grew up as the youngest of four on a farm in the eastern part of Norway. The oldest sibling was a sister proud to be the one inheriting the farm. He also had two older brothers. However, he seldom was welcomed to participate in their activities. His mother was a dominating woman who punished the family members with rejection if they were not submissive to her. His father did not oppose to Randolph's mother. Randolph Highland developed a behavior characterized by low assertiveness and a tendency to avoid conflicts.

He felt that his quality of life was low. However, he expected therapy would be helpful. He was rated with high QOR.

Course of treatment

Randolph Highland was also randomized to dynamic psychotherapy without TI. He came to 36 sessions. Based on the background information it was expected

that the therapist would focus on his inhibited aggression, fear of rejection and guilt if he insisted on his own opinion.

A short segment from mid-treatment is shown in Table 5. Table 5 illustrates how the IRR-raters would evaluate the presence and timing of RIs, the themes in the dialogue and the patient's response.

The raters of the segment from Randolph Highland's session identified RIs in category 1, 2 and 3. The timing of the first RI is not decided since this is the first patient-therapist interaction in the transcript. The timing of the second and third RI is on average 3. The first RI was a preparing intervention where the therapist addressed previous interpersonal transactions with other people. In the first RI the therapist was a little supportive as well as a little challenging (1), in the second RI the therapist was more supportive (3) and still a little challenging (1). In the third RI the therapist was less supportive (1) and more challenging (3). This is the last patient-therapist intervention. Therefore we don't know how the patient responded to this intervention in the transcript. In the second RI the therapist actively

Table 5 Segments with Relational Interventions from the patient-therapist dialogue in a mid-therapy session rated with Relational Work Scale

Individual speaking	Parts of dialogue	(RI): Category 1–5 Rated 0–4: (T): Timing (C): Content (V): Valence (R): Response
Therapist	You have previously focused on whether you wanted to continue to work with your colleagues in that committee.	RI: 1 C: Other (4) V: Supportive (1), Challenging (1)
Patient	I planned to refuse reelection to the committee. However, I was persuaded to continue by the other members. I kind of regret accepting the position because it takes time from other things I want to do like participate in the literature group with my friends.	C: Other (3) R: Associations/self-reflections (3), Cooperative (3), Emotional (3)
Therapist	Then the other members in the committee think you are a great chap. You care for your family, colleagues and friends, but do you care about your own needs?	RI: 2 and 3 T: Connect natural (3) Precise/striking (4) C: Other (4) V: Supportive (3) Challenging (1)
Patient	At home my kids know my wife's opinion, but not mine. They want me to be more explicit. On the other side, I get guilty conscience and fear of them getting angry if I am too opinionated.	C: Other (3) R: Associations/self reflections (2), Cooperative (3), Emotional (2)
Therapist	Yes, they have even told you to stop care so much about them and start prioritize what is necessary for yourself, for example to buy yourself some new clothes. Do you think that is a common concern among children and teenagers nowadays?	RI: 2 T: Connect natural (2) Precise/striking (3) C: Others (3) V: Supportive (1), Challenging (3)

encouraged Randolph Highland to explore thoughts and feelings about his relationships to others (colleagues at work; rated 4) and how he imagined they might feel or think about him (RI category 2 and 3). The timing of the intervention was good (3.5). Randolph Highland responded with associations and self-reflection (rated on average 3), was cooperative (3) and emotionally involved (2.5). Neither the therapist, nor the patient mentioned the patient's symptoms directly. They both focused on Randolph Highland's relationships with other people.

After therapy Mr. Highland described that he had made huge changes in his work load and his relations at work. He still participated as a good team worker, but did not any longer compensate when colleagues did not complete their duties. He was intellectually aware of his still somehow submissive relational pattern with his wife, and articulated that he was satisfied and that he did not want to change this further.

There are obvious limitations in drawing any conclusions from ratings with RWS of these short segments from the two vignettes. However, the available material rated with RWS indicates different processes and therapist–patient interactions in the two therapies. Annie Lowland, who at pre-treatment showed difficult interpersonal relationships, experienced more challenging and less supportive therapist interventions. In the segment from Randolph Highland's therapy both the therapist and the patient talked about Randolph's relations with other people, however, not about his relationships with his parents. When Annie's therapist pointed at the patient's relationship with her mother and compared elements in the mother's and Annie's, relational patterns, Annie refused to explore this, while she began talking about her symptoms. These ratings might be hints that point towards a possible understanding of why Annie, while being challenged by her therapist, preferred talking about symptoms rather than relationships and did not experience improved relations with other people. Randolph might have experienced the therapist interventions more supportive and less challenging, and managed to explore interpersonal relationships to a greater extent. This may have resulted in improved relations with other people. No exploration of his image of his dominating mother was present in this short segment. However, if this interaction was representative for the therapy, the lack of focus on the repetitive interpersonal pattern might have been a hindrance for him in further resolving his relationship with his dominating wife.

Discussion

The Relational Work Scale was specifically developed to identify and explore in more detail Relational Interventions and the in-session process after such interventions and thus, explore the relational work. Five categories of RIs were used in RWS, to delineate and operationalize

the out of therapy relational constructs. Items on timing of therapist interventions, content, and the interaction between patient and therapist were used in the scale which has comparable items as the previously published Transference Work Scale.

A limitation is that RWS was developed within the FEST research group on items defined by interventions prescribed in the treatment manual [20]. The manual was also developed by the same research group. One of the researchers had participated in the study also as a rater and therapist and knew the study and the interventions very well. In the present study, the adherence to the manual was high. Another limitation might be that the ratings with RWS are performed on 10-min segments. To know whether RWS might be a useful tool for other research groups, IRR of RWS needs to be tested on transcripts from other therapies in other studies also with longer transcripts. As with TWS [16], RWS was promising with regard to achievement of inter-rater agreement. Due to limited resources, however, the number of rated sessions in the inter-rater analyses of RWS was limited.

Even though the two raters differed in psychodynamic training, theoretical background, and work experience the rater agreement for nearly all the RWS items in the present study was good to high. RWS is a focused and short process measure and is seemingly not too time consuming. Together with TWS, RWS may be used for creating datasets from a larger number of sessions or segments of sessions. The two vignettes also indicate that through rating of transcribed segments or whole sessions, use of differing therapist interventions and patient responses might be revealed. Different categories of RI and other features/aspects of relational work may relate differently to outcome. Previously reported interaction effects of RI/TI and sub-groups of patients need to be further explored with process-outcome studies. A focus on the unexpected positive effect of RI for some patient groups could be further investigated [11, 26, 32]. RWS might prove especially suitable for intensive case studies combining quantitative and narrative data.

Conclusion

The Relational Work Scale was specifically developed to identify and explore in more detail Relational Interventions and the in-session process after such interventions. The inter-rater agreement was good to high. RWS is a focused and short process measure. This study suggests that RWS might be a potentially useful tool to identify RI as well as explore the interaction of timing, category, and valence of relational work in psychotherapies that focus on the patient's relations with other people than the therapist.

Additional file

Additional file 1: Relational Work Scale (RWS). (DOCX 16 kb)

Acknowledgement

The authors thank; Kjell Petter Bøgwald, MD, PhD, Øystein Sørbye, MD, Oscar Heyerdahl MD, Alice Marble, PsyD, and Mary Cosgrove Sjaastad MD for their contribution in peer supervision and for providing treatment data to the study. They are all psychotherapists in private practice.

Funding

The present study is funded by the Division of Mental Health and Addiction, University of Oslo, Oslo, Norway and the Research Unit, Division of Mental Health, Vestfold Hospital Trust University. The FEST-study has also received funding from The Norwegian Research Council, The Norwegian Council of Mental Health, Health and Rehabilitation, and Diakonhjemmet Hospital Trust. The funding sources had no influence in the design or the decision to publish the study.

Availability of supporting data

The data set supporting the results of this article is available on request from the first author, Randi Ulberg.

Authors' contributions

RU is the first author of this study. She is a researcher in the FEST-research group with responsibility for micro-process analyses. She has the main responsibility for the design, the collection and analyses of data as well as the writing of the present manuscript. SA is the clinical director in FEST. PH is the principal investigator in FEST. SA and PH have participated in designing the present study. SA, EN, PH, H-SJD, KC and PB have participated in providing and analyzing treatment data, and drafting and revising the manuscript critically for important intellectual content. All authors have given final approval.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

The ethical approval and consent to participate included consent to publish. Patient material and data collected including case material were accepted for use in research and publishing as well as teaching.

Ethics approval and consent to participate

The Regional Ethics Committee for health region 1 in Norway approved the study protocol and the information given to the patients: First Experimental Study of Transference-interpretations (FEST307/95). Written informed consent was obtained from each participant.

Author details

¹Institute of Clinical Medicine, Division of Mental Health and Addiction, University of Oslo, Box 85 Vinderen, 0319 Oslo, Norway. ²Research Unit, Division of Mental Health and Addiction, Vestfold Hospital Trust, PO Box 2169, 3125 Tønsberg, Norway. ³James Madison University, Harrisonburg, Virginia, USA. ⁴Department of Psychiatry, Vestre Viken Hospital Trust, Drammen, Norway.

Received: 31 March 2016 Accepted: 29 August 2016

Published online: 06 September 2016

References

- Wykes T, Haro JM, Belli SR, Obradors-Tarrago C, Arango C, Ayuso-Mateos JL, Bitter I, Brunn M, Chevreul K, Demotes-Mainard J, et al. Mental health research priorities for Europe. *Lancet Psychiatry*. 2015;2:1036–42.
- Cuijpers P, Berking M, Andersson G, Quigley L, Kleiboer A, Dobson KS. A meta-analysis of cognitive-behavioural therapy for adult depression, alone and in comparison with other treatments. *Can J Psychiatry*. 2013;58(7):376–85.
- Driessen E, Hollon SD, Bockting CL, Cuijpers P, Turner EH. Does publication bias inflate the apparent efficacy of psychological treatment for major depressive disorder? a systematic review and meta-analysis of US National Institutes of Health-Funded Trials. *PLoS One*. 2015;10(9):e0137864.
- Kazdin AE. Understanding how and why psychotherapy leads to change. *Psychother Res*. 2009;19(4–5):418–28.
- Crits-Christoph P, Connolly Gibbons MB, Mukherjee D. Psychotherapy process-outcome research. In: Lambert MJ, editor. *Bergin and Garfield's handbook of psychotherapy and behavior change*. 6th ed. NJ: Wiley; Hoboken; 2013. p. 298–340.
- Orlinsky DE, Rønnestad MH, Willutzki U. Fifty years of psychotherapy process-outcome research: continuity and change. In: Lambert M, editor. *Bergin and Garfield's handbook of psychotherapy and behavior change*. 5th ed. NJ: Wiley; Hoboken; 2004. p. 307–89.
- Levy K, Scala J. Transference, transference interpretations, and transference-focused psychotherapies. *Psychother Theory Res Pract*. 2012;49:391–403.
- Malan D. *The frontier of brief psychotherapy*. New York: Plenum Press; 1976.
- Høglend P. Exploration of the patient-therapist relationship in psychotherapy. *Am J Psychiatry*. 2014;171(10):1056–66.
- Høglend P, Bøgwald K, Amlo S, Marble A, Ulberg R, Sjaastad M, Sørbye O, Heyerdahl O, Johansson P. Transference interpretations in dynamic psychotherapy: do they really yield sustained effects? *Am J Psychiatry*. 2008;165:763–71.
- Ulberg R, Høglend P, Marble A, Johansson P. Women respond more favorably to transference intervention than men: a randomized study of long-term effects. *J Nerv Ment*. 2012;200:223–9.
- Gabbard G, Westen D. Rethinking therapeutic action. *Int J Psychoanal*. 2003;84:823–41.
- Samberg E, Marcus E. Process, Resistance, and Interpretation. *Textbook of Psychoanalysis*. 2005. p. 229–237.
- Town J, Diener M, Abbass A, Leichsenring F, Driessen E, Rabung S. A meta-analysis of psychodynamic psychotherapy outcomes: evaluating the effects of research-specific procedures. *Psychotherapy*. 2012;49:276–90.
- Ulberg R, Amlo S, Høglend P. Manual for transference work scale; a micro-analytical tool for therapy process analyses. *BMC Psychiatry*. 2014;14(1):291.
- Ulberg R, Amlo S, Høglend P. Transference interventions and the process between therapist and patient. *Psychotherapy*. 2014;51:258–69.
- Høglend P, Amlo S, Marble A, Bøgwald K-P, Sørbye O, Sjaastad M, Heyerdahl O. Analysis of the patient-therapist relationship in dynamic psychotherapy: an experimental study of transference interpretations. *Am J Psychiatry*. 2006;163:1739–46.
- Høglend P. Dynamisk korttidssterapi (brief dynamic psychotherapy). In: Alnes R, Ekern P, Jarval P, editors. *Poliklinikkens psykiatrisk klinikk 25 år*. Oslo: Psykiatrisk klinikk Vinderen; 1990. p. 27–38.
- Bøgwald K-P, Høglend P, Sørbye O. Measurement of transference interpretations. *J Psychother Prac Res*. 1999;8:264–73.
- Høglend P. Manual for process ratings of general skill, supportive interventions and specific techniques. Unpublished manual: Oslo, Norway: Department of Psychiatry, University of Oslo; 1994.
- Høglend P, Bøgwald KP, Amlo S, Heyerdahl O, Sørbye O, Marble A, Sjaastad MC, Bentsen H. Assessment of change in dynamic psychotherapy. *J Psychother Pract Res*. 2000;9(4):190–9.
- Alden L, Wiggins J, Pincus A. Construction of circumplex scales for the inventory of interpersonal problems. *J Pers Assess*. 1990;55:521–36.
- Derogatis L. SCL-90-R: administration, scoring and procedure manual. Towson: Clinical Psychometric Research; 1983.
- Sifneos PE. Short-term anxiety-provoking psychotherapy: a treatment manual. New York: Plenum Press; 1992.
- Piper WE, Duncan SC. Object relations theory and short-term dynamic psychotherapy: findings from the quality of object relations scale. *Clin Psychol Rev*. 1999;19(6):669–85.
- Høglend P, Dahl H, Hersoug A, Lorentzen S, Perry J. Long-term effects of transference interpretation in dynamic psychotherapy of personality disorders. *Eur Psychiatry*. 2011;26:419–24.
- Strachey J. The nature of the therapeutic action of psycho-analysis. *J Psychother Pract Res*. 1934. 1999;8:66–82.
- Greenson R. The technique and practice of psychoanalysis, vol. I. New York: International Universities Press Inc; 1975. p. 372–3.
- Ulberg R, Marble A, Høglend P. Do gender and level of relational functioning influence the long-term treatment response in dynamic psychotherapy? *Nordic J Psychiatry*. 2009;63:412–9.
- Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33(1):159–74.
- Shrout PE, Fleiss JL. Intraclass correlations: uses in assessing rater reliability. *Psychol Bull*. 1979;86(2):420–8.
- Høglend P, Hersoug A, Bøgwald K, Amlo S, Marble A, Sørbye O, Røssberg J, Ulberg R, Gabbard G, Crits-Christoph P. Effects of transference work in the context of therapeutic alliance and quality of object relations. *J Consult Clin Psychol*. 2011;79:697–706.