**Decidual acute atherosclerosis:
immunohistochemical definition, immune cell
involvement, and tissue heterogeneity**

PhD thesis

by

Patji Alnæs-Katjavivi,

MB BS, BSc

2020

**Division of Obstetrics and Gynaecology,
Oslo University Hospital, Ullevål
University of Oslo, Norway**

**UNIVERSITY
OF OSLO**

© **Patji Alnæs-Katjavivi, 2020**

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-550-1

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

PhD thesis 2019

Patji H. Alnæs-Kajavivi, MB BS (MD) & BSc , PhD candidate

Division of Obstetrics and Gynaecology

Oslo University Hospital, Ullevål

and University of Oslo, Oslo, Norway

Dedication

This thesis is dedicated to young Arion Alaj, and to the memory of Matthew Cohen, Grant Norman, and Mark Strode. Rather Use Than Fame. Lastly, to my mother, “*kaende naua... tjinene, tjinene*”.

TABLE OF CONTENTS

1. ACKNOWLEDGEMENTS	5
2. LIST OF PAPERS (Papers I-III)	7
3. ABBREVIATIONS	9
4. ABSTRACT	11
5. INTRODUCTION	15
6. AIMS OF THE THESIS	45
7. MATERIALS AND METHODS	47
8. SUMMARY OF RESULTS	65
9. GENERAL DISCUSSION	75
10. CONCLUSIONS	97
11. FURTHER STUDIES	99
12. ERRATUM	101
13. OTHER PUBLICATIONS CO-AUTHORED BY CANDIDATE DURING PhD STUDIES	103
14. REFERENCE LIST	105
15. APPENDIX 1: Microscopy worksheet for Paper I	115
16. APPENDIX 2: Microscopy worksheet for Paper III	117

PAPER I and Paper I erratum

PAPER II

PAPER III

1. ACKNOWLEDGEMENTS

The studies contributing to the production of this thesis would not have been possible without access and utilisation of the Oslo Pregnancy Biobank. I would like to thank all the women and children participants that have kindly provided their biological samples to the biobank since its conception in 2000, by Professor Annetine Staff. I would also like to thank all those research staff that have secured the recruitment of participants, collected biosamples, and followed the meticulous process required for correct storage and archiving. I would like to thank Lise Levy, Tone Haug and Michael Pilemand Hjørnholm for their tireless dedication to the supervision of the Oslo Pregnancy Biobank, and the administration of the Research Center for Obstetrics and Gynaecology at Ullevål, Oslo University Hospital.

First and foremost, I would like to express my gratitude to professor Annetine Staff, my supervisor, for offering me the opportunity to join her research group in her investigation of the most intriguing conditions of pregnancy: preeclampsia. The current study of acute atherosclerosis is the continuation of the doctoral work undertaken by Dr Nina Harsem. Professor Annetine Staff's ever-striving vision, guidance and instruction are equalled by her generous acknowledgement, shown to both her co-workers and collaborators alike. I would also like to extend this heart-felt gratitude to Professor Borghild Roald, my mentor and navigator through the landscape of decidua, viewed down the lens of the pathologist's microscope.

I would like to thank my two co-supervisors, Professor Christopher Redman (Nuffield Department of Women's and Reproductive Health, Oxford, UK) and Dr Ralf Duchend (Experimental and Clinical Research Center, a cooperation of Charité-Universitätsmedizin Berlin, Germany). Both have contributed invaluable support to the research undertaken, intellectually, personally, and by making available to me the laboratories and collaborators with whom they work.

I am indebted to the Department of Pathology at Ullevål Oslo University Hospital for undertaking the preparation and histological staining of tissue specimens, with particular thanks to the pathology assistant Linn Buer and laboratory technician Giang Huong Nguyen. The workload was never "too much" and the perfectionism with which they executed their work is a credit to their professions.

Post-doctorate scientists Guro Mørk Johnsen and Gro Leite Størvold have been instrumental in their contribution to the evolution, precision, criticism and publication of the research undertaken. This extends to the others in our research group that have been a continuous source of support, advice and ideas, Meryam Sugulle, Kjartan Moe, Ingrid Fossheim, Amalie Bjerke

Rieber-Mohn, Ana Rita Sequeira de Sousa, Hanne Guldsten, Kristina Alme Gardli, Nina Julie Verket, Heidi Fjeldstad, Michaela Golic (Berlin).

I have been most fortunate to receive guidance from experts in the field of placental pathology and preeclampsia. Thank you Fiona Lyall (Glasgow, UK), Robert Pijnenborg (visiting Charité-Universitätsmedizin Berlin, Germany), Graham Burton (Cambridge, UK), and my co-supervisor Christopher Redman.

I would like to thank my heads of department, Linda Helgadottir following on from Anne Flem Jacobsen, both of whom have supported the duality of my clinical work combined with research. My thanks to you both for facilitating this balancing act. My thanks to Bjørn Busund, head of the Department of Obstetrics and Gynaecology at Oslo university hospital, and to the Faculty of Medicine at the University of Oslo, for providing me with unique opportunity to further my academic education, as well as make a contribution to scientific research. My thanks to these institutions, and to the funders for these studies, The Research Council of Norway, and the South-Eastern Norway Regional Health Authority.

My family and friends are my source of meaning and purpose. Without them there is neither. Thank you Maja for being the first “outsider” to read my thesis, and doing so with your even professional critique. Thank you Peter Jourdan for giving me the benefit of your experience working on a similar project. Thorbjørn, trusted wingman. To my patient wife, Heidi, and to my children Egil and Meri, thank you for your Finnish forbearance, and for always believing I could complete this task.

Oslo, 6th December 2019

2. LIST OF PAPERS (Papers I-III)

Paper I: Alnaes-Katjavivi P, Lyall F, Roald B, Redman CWG, Staff AC

Acute atherosclerosis in vacuum suction biopsies of decidua basalis: An evidence based research definition

Placenta 2016; 37:26-33

Paper I erratum: Alnaes-Katjavivi P, Lyall F, Roald B, Redman CWG, Staff AC

Corrigendum to “Acute atherosclerosis in vacuum suction biopsies of decidua basalis: An evidence based research definition” [Placenta 37C (2016) 26-33]

Placenta 2017; 52:114-115

Paper II: Alnaes-Katjavivi P, Roald B, Staff AC

Uteroplacental acute atherosclerosis in preeclamptic pregnancies: rates and clinical outcomes differ by tissue collection methods

Submitted for publication

Paper III: Johnsen GM*, Leite GS*, Alnaes-Katjavivi PH, Roald B, Redman CWG, Staff AC

*Shared first authorship

Acute atherosclerosis and perivascular lymphocytes in decidual spiral arteries: quantification and comparison following preeclamptic and normotensive pregnancies

J Reprod. Immunol. 2019; 132:42-48

3. ABBREVIATIONS

AA	acute atherosclerosis
ABC	avidin-biotin-peroxidase
ANCA	anti-neutrophil cytoplasmic antibody
BMI	body mass index
BPS	Basal Plate (maternal) Surface
DAB	diaminobenzidine
DSM	Decidua vacuum Suction Method
ECM	extracellular matrix
EVT	extravillous trophoblast
FGR	fetal growth restriction
H&E	hematoxylin and eosin
HRP	horseradish peroxidase
IHC	immunohistochemistry
MR	(fetal) Membrane Roll
OPB	Oslo Pregnancy Biobank
PAS	Periodic acid Schiff
PhD	philosophiae doctor
PI	pulsatile index
REK	Regional Committee for Medical and Health Research Ethics
SD	standard deviation
uNK	uterine Natural Killer cell
VSMC	vascular smooth muscle cell(s)
WHO	World Health Organization

4. ABSTRACT

Background

Acute atherosclerosis is an understudied histological lesion of maternal uteroplacental spiral arteries in pregnancy. The rate is reported as particularly high in the condition known as preeclampsia (40%), a potentially dangerous hypertensive complication of pregnancy. Also, the arterial lesions are more common in the decidua than in the myometrial parts of the uterine wall. Acute atherosclerosis was first described in 1945, demonstrating a morphological resemblance to the “fatty streaks” of early stage atherosclerosis, with lipid-laden foam cells within the vessel wall being the prominent feature in both instances. Clear defining lesion criteria are however lacking, as well as the component and localization of a “perivascular infiltrate”. Acute atherosclerosis lesions may be obstructive lesions, and the rate of placental necrosis downstream of these placenta-supplying arteries of the decidua basalis are reported as increased, as are the rates of poorer pregnancy outcomes in some studies.

Women with a history of previous pregnancies affected by preeclampsia have, as a group, an increased future risk of developing atherosclerotic cardiovascular disease, although the biological mechanisms are unclear. Our hypothesis has been whether the presence of acute atherosclerosis could be a histological hallmark useful in stratifying a young woman’s risk for developing remote cardiovascular disease, decades prior to the clinical manifestation of atherosclerosis. To better answer such a question in future longitudinal clinical studies after a pregnancy, where the presence or absence of acute atherosclerosis could be tested for, this PhD thesis aimed to provide an improved morphological characterization of the lesion’s features, across tissue sampling techniques and patient groups, and its associations to clinical short-term outcomes of pregnancy.

Methods

Decidual tissue samples and extensive clinical data from the 297 women with heterogeneous pregnancy outcomes, scheduled for elective cesarean section, were recruited (2001-16) to our ongoing Oslo Pregnancy Biobank research studies and included in this study. Three methods were used to collect decidual (endometrium of pregnancy) samples; 1) our own previously developed decidual vacuum suction method, yielding decidua basalis tissue; 2) biopsies from the basal plate surface of the placenta, also yielding decidua basalis tissue; and 3) fetal membrane rolls, yielding decidua parietalis tissue.

Representative serial tissue slides were used for immunohistochemical staining following standardized preparation by pathologist technicians. All tissue sections were evaluated by light microscopy by the PhD candidate, supported by collaborators and a placenta pathologist specialist. Employing a systematic predefined approach, smooth muscle (Desmin) and/or presence of fibrinoid (Periodic acid Schiff) were used to detect decidual spiral arteries. Scavenger receptor (CD68) confirmed the morphological appearance of intramural foam cells. Standardized immunostains detected subclasses of decidua basalis perivascular lymphocytes, quantified for each artery according to prescribed “zones”. Kappa scores confirmed intra- and inter-observer satisfactory reproducibility of microscopy assessments. Non-parametric tests were used for testing statistical significance of acute atherosclerosis associations with histological findings and clinical parameters.

Results

In keeping with previous research undertaken by our group, the spiral artery detection rate was highest in tissue samples obtained using the decidual vacuum suction method. In the group of preeclamptic women (n=107), the spiral artery detection rate was 88%, compared to 78% in membrane rolls, and only 56% in placental basal plate tissue samples. We concluded that the

simplest (and most reproducible) criteria for defining acute atherosclerosis in decidua basalis spiral arteries was the presence of at least two adjacent lipid-laden intramural cells (in a hematoxylin & eosin stained section sample), confirmed by CD68 positivity on a serial section slide. Presence of arterial wall fibrinoid was common to nearly all spiral arteries of the decidua basalis. Periodic acid Schiff positivity did not clearly discriminate the fibrinoid necrosis of spiral arteries from the almost ubiquitous fibrinoid of physiological remodelling. Perivascular infiltrate was neither consistent nor exclusive to foam cell presence in spiral arteries, though acute atherosclerosis detected in preeclamptics was associated with higher concentrations of CD3+CD8- cells (assumed to represent CD4 T-lymphocytes) in both perivascular and surrounding decidual interstitial infiltrates. Decidual NK-cells (CD56+) did not associate with presence of acute atherosclerosis, and T-regulatory lymphocytes were scarce or absent in samples of both normotensive and preeclamptic pregnancies, irrespective of acute atherosclerosis presence. Presence of luminal thrombosis and absence of endovascular extravillous trophoblasts were associated with artery sections affected by acute atherosclerosis.

Sample sections of vacuum suctioned decidua basalis tissue showed higher rates of acute atherosclerosis compared to samples of placental basal plate samples and membrane rolls collected from the same preeclamptic patients. The rate of acute atherosclerosis was highest among preeclamptic (in 37% of preeclamptics without diabetes, in 29% of preeclamptics with diabetes) but also detectable in normotensive pregnancies (in 11% of normotensive without diabetes, and in 10% of normotensives with diabetes). Normotensive multiparous women with a previous history of preeclampsia were more likely to develop acute atherosclerosis than multiparous women with previously normotensive pregnancies. Acute atherosclerosis lesions in preeclamptics contained more foam cells than in lesions of normotensive controls. In preeclamptics,

acute atherosclerosis associated with fetoplacental syndromes (preterm delivery, fetal growth restriction), but did not associate with maternal parameters (elevated age, BMI, or blood pressure), which are known to be risk factors for both preeclampsia and cardiovascular disease.

Conclusions

Our systematic approach, utilizing endometrium of pregnancy (decidua) tissue samples obtained from heterogeneous pregnancy groups, provided a definition for decidual acute atherosclerosis that is both simple, reproducible and applicable to both vacuum-suctioned tissue samples as well as samples taken in routine clinical pathology (e.g. from the maternal surface of the delivered placenta and fetal membrane tissue rolls). In preeclamptics, vacuum tissue collection of decidua basalis tissue yielded higher amount of spiral arteries, as well as higher rates of acute atherosclerosis when compared to the two other decidual collection methods.

Although smaller lesions are detectable in a small proportion of normotensive as well as diabetic pregnancies, acute atherosclerosis is an arterial lesion most highly represented in preeclamptics, where it associates with adverse fetoplacental parameters. Though our findings can demonstrate differences between the acute atherosclerosis of spiral arteries, and the atherosclerosis of larger arteries, both lesions share an association with the increased presence of the adaptive immune system, specifically CD4⁺ T-lymphocytes. In future studies, the detection of acute atherosclerosis could be tested in early life cardiovascular risk stratification for parous women, although its feasibility is reduced by our documented optimal tissue collection method (vacuum suction of decidua basalis) being less accessible clinically in vaginal deliveries.

5. INTRODUCTION

5.1 Spiral arteries: the blood supply to the placenta intervillous space

Acute atherosclerosis is a lesion specific to the uterine spiral artery in the pregnant state. Spiral arteries are the terminal branches of the arterial supply to the uterus, commonly known as the “womb”, the woman’s organ of reproduction. As such, the spiral arteries of pregnancy represent the conduit for new life.

Figure 1. Illustrating the arterial supply of the uterus from the aorta ending in the spiral arteries of the endometrium (insert), becoming decidua in pregnancy. Reprinted with permission from Panopto.com (image left) and Journal of clinical pathology (for insert (1))

As illustrated in Figure 1, the blood supply of the uterus is delivered by the uterine arteries, arising from the internal iliac arteries, joined by the supply delivered from the ovarian arteries. On penetrating the muscular wall of the uterus, the supply continues through the arcuate arteries, which arch around the body of the uterus, giving rise to radial arteries that penetrate the myometrium, and extend towards the myometrial-endometrial junction (2-4).

Branching from the distal ends of the radial arteries within the cavity-near myometrium arise two types of smaller arteries, or – as the terminal branches of the arterial tree – arterioles. The basal arteries, measuring less than 100-120µm in diameter, supply the basal portion of the endometrium, the mucosal

lining of the uterine cavity. Spiral arteries are non-branching end arteries with a typical corkscrew shape, emerging from the radial arteries embedded in the myometrium (5, 6). They extend beyond the basal layer, supplying the upper functional layer of the endometrium that undergoes hormone-dependent structural changes during a woman's menstrual cycle.

In the non-pregnant endometrium, the spiral arteries are 50-100 μ m in diameter and contribute to the total peripheral resistance in the arterial bed. Uterine blood flow rises from 50ml/min at the outset of pregnancy, to 500-750ml/min by term, increasing from less than 2% of the body's total cardiac output in the non-pregnant state (7) to 10-15% of cardiac output at term (8). This enormous increase in blood flow exists by virtue of an impressive architectural transformation of spiral arteries supplying the placenta. These "physiological changes" (9) became better known as artery "remodelling" (10). The resulting high arterial flow in pregnancy is well suited to the high demands of the growing fetus and its source of nutrition, the placenta. Hemodynamic changes secondary to a major decrease in uterine vascular resistance results from a combination of expansive remodelling and enhanced vasodilation, as well as functional changes in arterial wall reactivity (11).

5.2 Placentation and spiral artery remodelling

A healthy pregnancy outcome is dependent upon a sufficient maternal blood supply delivered to the placenta. Successful placentation is dependent on successful blastocyst implantation, and spiral artery remodelling. The mechanisms involved in how this transformation comes about, and the problems that arise when insufficient spiral artery remodelling occurs, are outlined below.

5.2.1 Normal human placentation

5.2.1.1 Setting the stage: changes prior to implantation

During the female reproductive cycle, ovulation heralds a rise in progesterone levels that coincides with the differentiation of proliferated endometrium into a secretory glandular tissue primed for receiving a blastocyst (12, 13) (implantation). The time of ovulation also coincides with the influx of specialized immune cells to the endometrial stroma that concentrate around spiral arteries and glandular tissue (14-16). The rise in stromal numbers of CD56+ uterine Natural Killer (uNK) cells is proportional to circulating progesterone concentrations, and occurs in preparation for – rather than as a result of – implantation of the blastocyst (13). Without conception resulting in implantation, the cell death of endometrial uNK cells precedes all other signs of impending menstrual breakdown and bleeding (4).

5.2.1.2 Placentation following blastocyst implantation

Decidua is the name given to the specialized endometrium in pregnancy. Following blastocyst apposition and adhesion to the decidualised endometrial epithelium, the decidua becomes invaded by cells of the trophoectoderm (13). Those cells of the trophoectoderm, closest to the inner cell mass, differentiate into two categories of trophoblasts: syncytiotrophoblasts and cytotrophoblasts. These cells are involved in the formation of villi, the development of the placenta, with cytotrophoblasts being the source of extravillous trophoblasts (EVT), that migrate into the surrounding decidua, and junctional myometrium (17). The (fetal) EVTs interact with maternal leukocytes in the decidual tissue, leading to the trophoblast-dependent transformation of decidual and myometrial spiral arteries (18).

The term *decidua basalis* denotes the region of decidua in which EVT-driven spiral artery transformation results in – together with the underlying junctional

zone of the myometrium – the placental bed perfusing the intervillous space of the placenta. The term *decidua parietalis* denotes the decidua where spiral arteries have not undergone EVT-driven transformation, and thus do not contribute to the perfusion of the intervillous space of the placenta (19).

5.2.1.3 Trophoblast-independent spiral artery remodelling

Non-destructive structural changes occur in the vascular smooth muscle cells (VSMC) of the spiral arteries prior to the arrival of invading trophoblasts. This has been demonstrated by examining biopsies of decidualised endometrium sampled in women with ectopic (extrauterine) pregnancies (and thereby an absence of trophoblasts in the uterine wall). Craven describes how the VSMC located in the media of decidual spiral arteries undergoes similar changes even in ectopic pregnancies as VSMC observed early in intrauterine pregnancies (20). Further studies have demonstrated that uNK cells, together with macrophages, temporarily localise around and within the wall of the spiral arteries, prior to trophoblast invasion. This cell combination secretes soluble factors that mediate morphological changes to both the extracellular matrix and VSMC of spiral arteries (18). The absence of uNK cells in murine experiments results in failure of spiral artery remodelling, an outcome that can be reversed if uNK deficient mice are replenished by bone marrow transplantation (18).

Interestingly, not only the spiral arteries are altered in pregnancy, even the uterine arteries (that are never invaded by trophoblasts) are dilated. During human pregnancy, the passive (fully dilated) diameter of the human uterine artery is approximately doubled, illustrating another non-trophoblast mediated cardiovascular adaptation to pregnancy (21). Indeed, the reduction in vascular resistance of the uterine artery in mid-pregnancy has often been ascribed to the remodelling effects of trophoblasts on the VSMC of placental bed spiral arteries. However, as demonstrated by a case of abdominal pregnancy,

remodeling alone could not explain low resistance in the uterine arteries (measured by blood flow Doppler) supplying a uterus devoid of a placental bed (22).

5.2.1.4 Trophoblast-dependent spiral artery remodelling

In his seminal work describing trophoblastic invasion from 8 to 18 weeks of gestation, Professor Pijnenborg describes how extravillous trophoblasts extend in cell columns into the decidua, proliferating from their progenitor cells where the anchoring villi attach to the decidua. Invasion of what is to become the placental bed (decidua basalis and first third of the myometrium) occurs by two separate routes, the interstitial, and the endovascular course (23). With the production of metalloproteinases by EVTs, interstitial invasion (through the decidual tissue) penetrates the extracellular matrix of the decidua, extending to the superficial myometrium (24). The depth of this penetration varies, but it is deepest at end of pregnancy corresponding to the central (sub-umbilical) cotyledon of the normal term placenta, and is least penetrating at the peripheral margins of the placenta (25, 26). The extent of the decidual invasion by “interstitial” extravillous trophoblasts is delineated by their fusion into the non-migratory multi-nucleated syncytia, or “giant cells”, first observed by Klliker in 1879 (4).

The retrograde endovascular invasion of extravillous trophoblasts follows the initial plugging of the distal tips of spiral arteries from cell columns extending into the decidua from the anchoring villi of cytotrophoblasts. The exclusive plugging of spiral arteries (not of veins or nutrient arteries (10)) maintains a low oxygen tension in the site of the placental bed by delaying perfusion of the decidua with maternal oxygen rich blood from high resistance (high velocity) arterioles. EVT plugging of spiral arteries occurs early following blastocyst implantation, providing a state that limits the embryonic (placental cells)

exposure to the cytotoxic oxygen free radicals resulting from aerobic metabolism (27). The extension of extravillous trophoblasts through the lumen and into the walls of spiral arteries is thought to initiate at 8 weeks from the date of the first day of last menses. These endovascular trophoblasts represent the fetal cells responsible for invasion of the spiral artery wall first along its decidual course followed by further extension along the arterial and myometrial course.

In coordination with maternal uNK cells, endovascular EVT's continue the disruption and destruction of the spiral artery VSMC, with gradual replacement of VSMC with a fibrinoid extracellular matrix that is secreted by EVT's (18). These endovascular EVT's are observed to replace the maternal endothelial cells of the spiral artery walls. Some authors believed this replacement to be permanent (28), which was later emphasized as being a temporary phenomenon, given that spiral arteries at delivery are lined by endothelial cells, not trophoblasts (29). It is still debated whether the intramural trophoblast cells – visible in the spiral arteries at delivery – are derived from the luminal advancement and extravasation of cells originating from the trophoblast plugs (endovascular EVT invasion); or, whether they derive from the interstitial invasion of EVT's, passing into the spiral artery wall by intravasation. Some researchers suggest that both intravasation and extravasation of EVT's may play a role (30).

The process of spiral artery transformation by EVT's is a process that evolves gradually (23), removing the resistance capabilities in the transformed spiral artery wall, increasing its calibre by 5-10 fold (31). This process begins within the first trimester involving the spiral arteries of decidua basalis (from 8 weeks gestation) expanding horizontally and vertically into the junctional myometrium throughout the second trimester (23, 26). The result is an increase

in blood flow, with a reduction in perfusion pressure (and the potentially damaging effects) of the blood passing through to the newly developed intervillous space of the placenta.

A necessary balance must be attained between the fetoplacental needs for nutrition and the maternal resources available. This balance is addressed by the interaction between EVTs (bearing paternal/foreign antigens), decidual extracellular matrix, and decidual cells belonging to the maternal immune system - principally tolerogenic uNK cells that are a rich source of angiogenic factors. The interaction of these three players modifies the invasiveness of EVTs, in addition to the essential role of oxygen levels in affecting EVT invasion (27). Lastly, the integrity and thickness of the spiral artery VSMC that is to be disrupted, and replaced, by EVT invasion will also influence the extent of spiral artery remodelling. This is partly influenced by the extent of decidual interactions prior to EVT arrival, as discussed above; and to the extent of non-reversed prior pregnancy remodelling of myometrial spiral artery wall. That spiral arteries are “less work” to remodel after a previous successful placentation, and thereby, easier to transfer in to high calibre low resistance conduits, has been suggested as a contribution to the observation that the birth weights of offspring to multiparous women are often higher than that of their first born (32). This may explain some of the biological cause for less subsequent preeclampsia risk in parous women (provided same partner). The nutrient arteries, or “straight” arteries, of the decidua basalis do not undergo the remodelling changes that manifest on neighbouring spiral arteries (9).

5.2.2 Abnormal placentation and placenta function in preeclampsia

Abnormal placentation is important to the development of the pregnancy dependent syndrome preeclampsia, of which both processes are summarized below.

Preeclampsia (PE) is a leading cause of maternal death, perinatal morbidity and mortality (33). One of its signs, hypertension, complicates up to 10% of all pregnancies. **Hypertension** is defined as systolic blood pressure (BP) ≥ 140 and/or diastolic BP ≥ 90 mm Hg. PE is a syndrome affecting 3-5% of all pregnancies, classically defined as new hypertension and new proteinuria developing in the second half of pregnancy (34). In recognition of the syndromic nature of preeclampsia, proteinuria is no longer a mandatory requirement for a preeclampsia definition in several updated guidelines (35), but can be replaced with other signs of maternal organ dysfunction, such as liver involvement, neurological or hematological complications (e.g. eclampsia, disseminated intravascular coagulation (DIC), and hemolysis). Some guidelines accept uteroplacental dysfunction signs in form of fetal growth restriction or abnormal fetal Doppler findings as such a new-onset feature of organ dysfunction (34). This feature is however not included as a criteria for diagnosing preeclampsia (PE) in most medical birth registries and many national clinical guidelines, such as the American College of Obstetricians and Gynecologists (36).

Preeclampsia is responsible for more than 50 000 maternal deaths annually, worldwide, and is most lethal in low-income settings with poor antenatal and obstetric care (37, 38). PE has also been identified as one of the leading direct causes of maternal death in Norway (39), where the last preeclampsia-related maternal mortality was 7 years ago (personal communication Professor Siri Vangen, Oslo University Hospital). Perinatal morbidity and mortality is also significantly increased in preeclampsia, with fetal demise resulting from either

placental abruption (40), or from chronic placental insufficiency through undersupply of adequate nutrients and gaseous exchange (41, 42). Both placental abruption and chronic insufficiency are clinical manifestations of ischemic placental disease, with which preeclampsia is associated (43). The only present “cure” for preeclampsia is delivery, by removal of the placenta, often necessitating a premature delivery, with added potential morbidity for the surviving newborns (44-46).

An appreciation that vascular abnormalities of the placenta associated with preeclampsia has been described multiple times in the course of the last century. Brosens was the first to demonstrate that preeclampsia was associated with less extensive remodelling in the myometrial segments of spiral arteries compared to remodelling in normotensive pregnancies (47). These seminal findings have been replicated by investigators examining specimens of placental bed biopsies, as well as continuing to utilize whole uteri with the placenta still in situ (23, 48, 49). Several authors also concluded that fetal growth restriction is associated with greater absence of remodelling of myometrial spiral arteries, either combined with or without preeclampsia (50-52). Most of these studies concluded that the morphological features of the placental bed in preeclampsia were secondary to a lack of trophoblast invasion, seemingly visible by the persistence of narrow myometrial spiral artery segments, with intact VSMC and elastic lamina.

Many of the pioneer spiral artery studies were conducted before the availability of immunohistochemical staining (IHC). Ivo Brosens comments in chapter 3 of the 2010 book “Placental Bed Disorders”; *“It can be concluded that in preeclampsia, whether or not it is a complication of essential hypertension, the failure of the spiral arteries to respond adequately to placentation must result in their suboptimal distension. The fetus is then*

subjected to poor intervillous blood flow from early gestation and not only during the period when preeclampsia is clinically manifest” (4). Though he points to hypoxia being the cause of fetal growth restriction, this has subsequently been replaced by another model of malperfusion through the placental bed (spiral artery) of the intervillous space (31), resulting in endoplasmic reticulum (ER) and oxidative stress in the placenta villi from the *high* velocity and the *low* flow of maternal blood entering the intervillous space (Figure 2. below, reprinted from (31)).

G.J. Burton et al. / Placenta 30 (2009) 473–482

Figure 2. Diagrammatic representation of the effects of spiral artery conversion on the inflow of maternal blood into the intervillous space and on lobule architecture predicted by modelling. CC: villous central cavity, ECL: echogenic cystic lesions, SMC: smooth muscle cells. Reprinted from Burton et al. Placenta (2009), with reuse through Open Access.

Though plausible, the concept that the extent of deep EVT invasion into the myometrium is proportional to the success of spiral artery remodelling, and subsequent placental perfusion, has been discussed by Pijnenborg (48, 53) and

Meekins (53). The concept has also been conflicted by contemporary investigation, utilizing more antibody-specific immunohistochemistry (IHC) for identifying invasion of both interstitial and endovascular EVT. Lyall et al. show that presence of both endovascular and interstitial EVTs do not differ when comparing the myometrial spiral artery sections (from placental bed biopsies) of preeclamptic and normotensive pregnancies. What is shown to be demonstrably different in the placental bed biopsies between preeclamptic and normotensive pregnancies is a reduced number of *intramural* EVTs in the myometrial sections of preeclamptic spiral arteries. In addition, the quantity of fibrinoid detected using Periodic acid Schiff (PAS), proportional to the observed disruption of VSMC, was reduced in the myometrial artery segments of preeclampsics, compared to normotensives (29). Whether the difference in fibrinoid/VSMC disruption is related to presence of intramural EVTs, or whether the difference arises from decidual mechanisms (independent of trophoblast invasion) is, as yet, unresolved.

Unlike other somatic cells, EVTs do not express class A or class B of the classical Human Leukocyte Antigen (HLA) system, allowing them to bypass the normal surveillance of the maternal (adaptive) immune system. An important variable that may modify EVT invasion, and the extent of remodelling of spiral arteries, is the interaction of HLA class C, G and E (HLA-C, HLA-G, and HLA-E) expressed by EVTs (54, 55), with uNK cells expressing both inhibitory and stimulatory Killer-cell Immunoglobulin-like Receptors (KIR) (56). The expanding knowledge base in this field is contributing to our understanding of how clinical pregnancy outcomes can be so dissimilar, when the histological appearance of EVT invasion is equivocal (as demonstrated by Lyall et al (29)). The polymorphisms of EVT HLA-C (C1 and C2) have been shown to strongly predict the development of preeclampsia, as well as large-for-gestational-age offspring, in a given pregnancy dependent

upon EVT interaction with the KIR-receptor haplotype A and B profile, expressed by maternal uNK cells (57-60).

5.2.3 An integrating 2-stage model of preeclampsia

The role of the placenta is crucial in preeclampsia (PE). Even a normal pregnancy and a normal placenta will impose increasing inflammatory stress on the maternal circulation. PE differs in that the inflammatory burden is even more excessive (61-64).

The exact etiology for PE remains unknown, but a 2-stage model is suggested (65), and has later been further developed by its creator, Professor Redman of the University of Oxford (66, 67) As summarized, dysregulated tolerance to foreign fetal (paternal) antigens (early pregnancy) underlies abnormal placentation with defect remodelling of maternal uteroplacental spiral arteries. These are early pathophysiological events. Dysfunctional placental perfusion ensues (Stage 1), with placental intracellular stress and thereby a release of placental stress signals to the maternal circulation. The stress signals are inflammatory signals that induce excessive maternal vascular inflammation with endothelial dysfunction, and the maternal features of PE including hypertension and proteinuria (stage 2 the clinical stage). This basic model has been subsequently updated and modified to fit with late-onset preeclampsia features, that mostly exhibit normal spiral artery transformation, but still causing endoplasmic reticulum (ER) and oxidative stress, due to malperfusion intraplacentally in the growing placenta reaching its limitations (67). This concept is presented in Figure 3 (reprinted from (68), with permission from the Journal).

Figure 3. Reprinted with permission from the Journal (68), by Staff and Redman. Revised two-stage preeclampsia (PE) model: all PE is dependent on placenta syncytiotrophoblast (STB) stress, and maternal risk factors may impact on several levels and both stages. Pathway A (associated with early-onset PE) illustrates the ‘extrinsic’ cause, and pathway B (associated with late-onset PE) illustrates the ‘intrinsic’ cause for placental malperfusion and dysfunction (stage 1), leading to the clinically recognized maternal syndrome of pre-eclampsia (stage 2). Pathway B is what is seen in postmature placentas, where the size of the term placenta may restrict intervillous perfusion. These two pathways (A and B) to placental stress are further detailed previously by us (67).

5.3 Increased long-term maternal cardiovascular risk after pregnancies complicated by placental dysfunction

Cardiovascular disease (CVD) is the number one non-communicable cause of death worldwide, affecting women and men alike. Atherosclerotic CVD represents a substantial burden to both individuals and societies (69), and its

prevention remains a global challenge. The early stages of atherosclerosis evolve asymptotically, identifiable in the post-mortem examination of the middle and large arteries of children and young adults, preceding the onset of classical risk factors (70). It takes several years before the accumulation of fatty streaks leads to symptomatic occlusive atherosclerotic cardiovascular disease. The preclinical stages of CVD, with arterial foam cell development and early stages of atherosclerosis, are affected by modifiable risk factors (71) making the condition amenable to potential amelioration through early (primary and secondary) intervention.

In addition to classic, gender-independent risk factors for CVD (tobacco smoking, diabetes mellitus, obesity, hypertension), pregnancy complications present a gender-specific risk. In 2011, The American Heart Association added preeclampsia (PE) and delivery of a growth-restricted child to the list of risk factors for developing CVD (69). Based on large population-based studies, compared to women with normotensive pregnancy, uncomplicated history of preeclampsia at term doubles the risk of future atherosclerotic cardiovascular disease. The risk may be as high as 8-fold if preeclampsia is complicated by a preterm delivery and by fetal growth restriction (72-74).

Placental dysfunction and preeclampsia is associated with long-term morbidity in the offspring, with increased incidence of both cardiovascular disease and diabetes mellitus type 2 in adulthood (75-77).

Placental dysfunction is also associated with the development of gestational diabetes mellitus (78, 79). Population-based data demonstrate that gestational diabetes mellitus (GDM) increases the risk of developing diabetes type 2 within the first ten years following the index pregnancy. This risk is increased

even further if the GDM pregnancy was associated with gestational hypertension or preeclampsia (80).

Two major theories are proposed to link maternal atherosclerotic CVD with placental syndromes such as preeclampsia (73). Firstly, preeclampsia and atherosclerosis share risk factors for systemic inflammation and endothelial dysfunction, such as obesity, dyslipidemia, diabetes, insulin resistance, and hypertension. A second possible mechanism, which does not exclude the contribution of common risk factors, is that the pregnancy, and especially preeclampsia or other placental dysfunction complications, may cause long-lasting effects on the maternal cardiovascular system, including accelerated arterial wall inflammation and myocardial dysfunction, which fail to resolve after delivery, thereby mediating increased long-term CVD risk (73).

Though it is undisputable that the placenta is a mandatory prerequisite for preeclampsia, a major response to pregnancy by the maternal cardiovascular system has been proposed as contributing to the development of preeclampsia (81). This, given pregnancy's substantial cardiovascular load on the maternal heart, and that preeclampsia and adult cardiovascular disease share same risk factors. The revised 2-stage placental model of preeclampsia (see figure 3. page 27) does indeed take into account the maternal cardiovascular system, which is preset at a higher inflammatory and potential dysfunctional stage in many women with epidemiological preeclampsia risk factors.

5.4. Acute atherosclerosis

5.4.1 Acute atherosclerosis: previous definitions with unclear criteria

“Acute atherosclerosis” is the term applied to a particular arteriopathy affecting the spiral arteries of the uteroplacental circulation. Acute atherosclerosis affecting the spiral arteries of the decidua basalis was first documented by Hertig in 1945

(82). In the subsequent publication by Zeek and Assali (1950), acute atherosclerosis was demonstrated affecting spiral arteries in decidua parietalis, as well as in decidua basalis (83). The morphological features of acute atherosclerosis presented in the literature typically include presence of subintimal (intramural) lipid-laden foam cells, fibrinoid necrosis, and perivascular infiltrate of mononuclear leukocytes.

The word *atherosis* alludes to lots of (“osis”) atheroma. “Atheroma” is a Greek word referring to the porridge-like gruel found in large arteries affected by atherosclerotic hardening (84, 85). The word *acute* recognizes that the development of this spiral artery lesion occurs in a limited time-scale, as a pregnancy lasts a limited number of weeks, in contrast to the longer history of atherosclerosis, to which acute atherosclerosis bears some morphological similarity. Identification of spiral artery acute atherosclerosis has been described in nearly all the sampling methods used for the investigation of uteroplacental pathology. Acute atherosclerosis has been detected in postpartum hysterectomy samples (9, 86), as well as in uteroplacental biopsies obtained from the maternal surface of placentas, fetal membranes, and placental bed tissue obtained by sampling techniques developed for research purposes, such as placental bed excision biopsies and decidual vacuum suction collection method (86-89).

A consistent feature of acute atherosclerosis is its lack of uniform distribution across spiral arteries within samples (10), affecting some parts of an artery – but not others (the decidua compared to the myometrium), its detection varying according to the topographic location of the spiral artery affected (myometrium, decidua, parietalis or basalis). Given that investigations of the spiral arteries rarely can be complete, this “patchy” nature of acute atherosclerosis lesion means that its absence on evaluation of relevant tissue samples can never fully exclude its existence in the remaining tissue from which the samples were taken.

Prior to immunohistochemical staining availability, the intramural foam cells of acute atherosclerosis were recognized by morphological appearance (Figure 4), with the lipid content being confirmed through the application of oil-red-O (82, 90). With the advancement of histopathological investigation by applied immunohistochemistry (IHC), acute atherosclerosis foam cells could be identified through their expression of CD68 (see Figure 5), the same scavenger receptor expressed by phagocytosing macrophages (91). These acute atherosclerosis foam cells lesions are heterogeneous in size, and previous literature has not addressed what the minimum foam cell requirements are for defining the presence of a foam cell lesion at light microscopy.

Figure 4. Decidual spiral artery section following H&E staining (collected using the Decidual vacuum -suction method (DSM)) x10 magnification. The artery wall contains both fibrinoid (confirmed using PAS) and lipid-laden foam cells (detailed in insert x40 magnification). The lumen is filled with thrombus. Perivascular infiltrate is not obvious (Photos by PhD candidate PAK).

Fibrinoid located in the walls of in spiral arteries is associated with acute atherosclerosis. However, fibrinoid is a recognized part and parcel of normal spiral artery remodelling. By light microscopy, detection of fibrinoid is confirmed through histochemical staining, most commonly Periodic acid Schiff (PAS), rather than specific antibody detection by IHC. As the term *fibrinoid* suggests, histochemical stains assist in the detection of fibrin-*like* components within the

extracellular matrix of the artery wall. The term “fibrinoid necrosis”, used to describe the fibrinoid of acute atherosclerosis lesions, has no uniform validated criteria. The term dates back to a period when histological descriptions were based upon morphology - assisted with non-specific staining techniques, such as PAS (92). In 1880, Neumann described fibrinoid as “*a chemical change of the intercellular substance of connective tissue, causing swelling, homogenization and conversion into a fibrin-like substance*”(93). This “process” was seen as *degenerative*, as it often resulted in the loss of cellular structures (e.g. smooth muscle), and the process of necrosis was assumed to be one of the many ways in which this cellular loss could come about (as supposed to apoptosis). The terms *fibrinoid degeneration* and *fibrinoid necrosis* have been used interchangeably in describing the same histological feature, where it has been assumed that necrosis is manifest (rightly or wrongly, often without additional qualification).

In descriptions of fibrinoid necrosis in the non-reproductive systemic circulation, the intramural detection of the blood-borne fibrin molecule is suggestive of a loss of integrity in the artery luminal wall, whereby defects within the endothelial layer allow for extravasation, or “leak”, of the relatively large fibrin molecule into the substance of the artery wall. This loss of artery wall integrity, proposed as a result of degenerative processes such as necrosis, has been the explanatory model for the presence of *fibrinoid necrosis* in a number of systemic small vessel arteriopathies, including post-streptococcal glomerula nephritis, polyarteritis nodosa, systemic lupus erythematosus, and anti-neutrophil cytoplasmic antibody (ANCA)-associated systemic vasculitis (94).

The composition of the fibrinoid lesion is not straightforward. Firstly, candidate constituents include pre-existing components of the spiral artery wall, namely smooth muscle, collagens, ECM ground substances such as

fibronectin and elastins, as well as blood borne substance (fibrin), and nucleoproteins (93). Secondly, placental bed tissues sampled in the third trimester contain spiral arteries that have undergone physiological remodelling almost universally at the decidual level (29, 47), and at the outer level of the myometrium. The fibrinoid deposited from remodelling (either trophoblast or non-trophoblast mediated) may share properties with that of fibrinoid necrosis, making the origin of the fibrinoid observed difficult to identify. Though Pijnenborg has claimed there to be a difference in the colour of PAS staining for the fibrinoid related to remodelling (“*brilliant pink*”) and for fibrinoid necrosis (“*greyish-blue*”) (88), there is no validated qualitative, nor quantitative, method for discerning fibrinoid necrosis from the extracellular fibrinoid of physiological remodelling.

According to some investigators, including Hertig (82), the formation of lipid-laden foam cells precede the appearance of fibrinoid deposition. However, Pijnenborg supports other investigators who claim that the first sign of acute atherosclerosis is the presence of fibrinoid (90, 95), or in some cases, the appearance of a perivascular leukocyte infiltrate (96). The problem facing all investigators is that the longitudinal histological development of acute atherosclerosis is not possible to follow antenatally, with the investigation of acute atherosclerosis necessitating delivery. The subsequent histological observation provides only a “snap-shot in time”, from which speculation around the preceding antepartum events cannot undergo vigorous testing). In response to this conundrum, some investigators in recent times have accepted the presence of intramural fibrinoid necrosis as a marker of so-called “decidual vasculopathy”, of which acute atherosclerosis (defined by the coexistence of lipid-laden foam cells) is a specific sub-type (89, 97-99). This pragmatic approach leaves the question open as to what comes first, fibrinoid or foam cells, instead allowing for the possibility that not all arteries demonstrating fibrinoid necrosis undergo foam cell

development of acute atherosclerosis. This suggests that despite similar morphological characteristics, and indeed similar clinical associations, there may be different etiological factors at play in the development of these spiral artery pathologies.

Perivascular infiltrate (PVI) is the characteristic feature least consistently mentioned in descriptions of acute atherosclerosis (Figure 5). PVI is included in many – but not all (100-102) – descriptions of acute atherosclerosis (1, 83, 90, 96, 103, 104). PVI is, for example, not mentioned in Hertig's original description (82). The significance of PVI, and whether it is associated with the development of acute atherosclerosis, or whether it is a consequence of the intramural changes leading to acute atherosclerosis, or even both, is not established. Discrepancies in the literature make it unclear whether the “perivascular” area to which PVI is designated is up to and including the luminal wall of the vessel (90, 105), or whether it involves only the decidua surrounding and adjacent to the outer margin of the artery wall.

Also, descriptions of the “infiltrate” vary from “pleomorphic”, “acute inflammatory”, to “mononuclear” and “lymphocytic”, suggesting both heterogeneity of cells between subjects and even within the same patient (1, 83, 90, 96, 103, 105). Many of these descriptions lack current available IHC markers that, if used, could elucidate the specific immune cell types that make up the leukocyte infiltrate associated with the other two features of acute atherosclerosis (foam cells and fibrinoid necrosis). Though IHC has been used to clarify the heterogeneity of leukocytes in the decidua (106-110), there is a lack of IHC descriptions of decidual immune cells localised in, and around, spiral arteries, and in particular their association with acute atherosclerosis.

Figure 5. Two parallel sections (left: H&E, right: IHC against CD68) showing the same decidual spiral artery (collected using the Decidual vacuum suction method (DSM)). The lumen (L) is obliterated by lipid-laden foam cells (F) which appear brown following IHC. Perivascular mononuclear leukocytic cells are clearly seen (*) x10 magnification (Photos by PAK).

In summary, though there has been some consistency in the three features that have characterized descriptions of acute atherosclerosis over the last 75 years, the criteria by which these features are defined lack both consistency and consensus. Acute atherosclerosis remains defined by its subjective visual appearance by microscopy, with both the etiology and the consequence of its manifestation remaining partly unresolved.

5.4.2 Acute atherosclerosis: association with clinical features of pregnancy

Over the last 75 years, acute atherosclerosis has been associated with the following clinical outcomes; preeclampsia, abruption and infarction of the placenta, occlusive thrombosis of spiral arteries, and fetal growth restriction. Being first described in third trimester uteroplacental samples of pathological pregnancies affected by preeclampsia and placental abruption (82), acute atherosclerosis quickly became synonymous with preeclampsia (47, 90), and assumedly the causative lesion underlying placental abruption. Relatively small and selective studies, with sample numbers often in the tens – to a few hundred - have consistently confirmed the association of acute atherosclerosis and preeclampsia, with a detection

rate of between 5-40% of preeclamptic samples (53, 89, 103, 111-114). The detection rate varies from study to study, likely due to heterogeneous patient selection and heterogeneous tissue sampling methods, and hampered by relatively small sample numbers. Detection rates differ according to the topography of the spiral artery, with higher rates in the decidua basalis arteries as compared to myometrial arteries and decidua parietalis arteries (114, 115). Some authors demonstrate the frequency of the lesion as proportional to the severity of preeclampsia, including preterm delivery and birthweight percentile (97, 116).

Acute atherosclerosis was eventually demonstrated in normotensive pregnancies, associated particularly with those affected by fetal growth restriction (FGR), with authors postulating that the lesion was not caused by hypertension per se (111, 117) but related to dysfunctional spiral artery remodelling and placentation (which in itself was the suggested mechanism underlying preeclampsia). The association of acute atherosclerosis with the finding of thrombotic infarcts in overlying decidua or placental villi was suggested as secondary to the spiral artery luminal narrowing by intimal arterial wall thickening due to the acute atherosclerosis lesions (82, 83, 86, 103). Samples containing spiral arteries from diabetic mothers were “remarkably” free of vascular disease (114), suggesting that development of acute atherosclerosis, as an arteriopathy, is not present in all types of pre-existing maternal co-morbidity.

The development of acute atherosclerosis could be demonstrated prior to the third trimester in case studies of abnormal pregnancies, such as hydatiform moles, or first-trimester pregnancy loss in pregnancies complicated by Systemic Lupus Erythematosus (SLE) (118, 119). Such case reports suggest to us that acute atherosclerosis could arise from states associated with excessive states of inflammation (120).

5.4.3 Acute atherosclerosis: resemblance to other arterial diseases?

With a broader perspective, what has been intriguing to investigators is the morphological similarity that acute atherosclerosis bears with the early stages of atherosclerotic cardiovascular disease, as well as the similarity to vessel disease seen in allograft transplant rejection (90, 96, 114). Though differing in the calibre of artery affected, and the time-course over which the lesion develops, both presence of intramural foam cells and associated perivascular infiltrate are features that acute atherosclerosis shares with graft-vessel disease and evolving atherosclerosis.

Whilst the etiology of acute atherosclerosis remains to be determined, the mechanistic processes by which both atherosclerosis and graft rejection develop are somewhat better understood. Some attention to these mechanistic processes may lead to a better understanding of the development of acute atherosclerosis, its significance in pregnancy, and its possible role as a determinant of future health.

5.4.3.1 AA may share immune mechanisms with atherosclerosis

Atherosclerosis is an inflammatory disease of the arterial wall (121), in which dyslipidemia plays a major part, involving both local and systemic inflammatory processes. Inflammatory activation generates foam cells by down-regulating cholesterol efflux from macrophages and other cells, with inhibited reverse cholesterol transport (122). Features of atherosclerosis not shared by AA include plaque formation and rupture, probably owing to the longer time course of atherosclerosis compared to AA.

Innate and adaptive immunity both contribute to the progression of atherosclerosis (123). Adaptive immunity depends on how T cells differentiate into subsets of T helper (Th1, Th2 and Th17) cells and T regulatory (Treg)

cells that restrain the immune response. Atherosclerosis involves systemic and local Th1 bias (pro-atherogenic interferon gamma producing Th1 cells) and Th17 (IL-17 producing) cells, while Tregs (124) and some B cell subsets are supposed to be protective. Treg numbers are reduced in the circulation and in the lesions of individuals with more severe atherosclerosis. Immune changes of PE are similar, causing excessive systemic inflammatory response (125), with a Th1-bias, unlike the Th2-bias of normal pregnancies. In PE, circulating and decidual Tregs are decreased, whereas circulating Th17 cells are increased (126). Decidual (maternal) Tregs protect fetal cells from immune rejection (127).

5.4.3.2 Similarities to Graft Vessel Disease of transplants

Immunological pathways leading to AA have been postulated by several authors (90, 96, 105, 114), due to its histological resemblance to the rejection observed in graft vessels during acute rejection. Atherosclerosis occurs in transplant vasculopathy, consistent with the hypothesis that inflammation underlies the lesion (121). This may also be relevant in the circulation of the placental semi-allograft with respect to AA. But there are differences. Transplantation identity of the placental allograft is more restricted than that of an organ transplant. Placental villous syncytiotrophoblast is HLA-null, whereas invasive (extravillous) trophoblast, which is in tissue contact with maternal decidual immune cells and spiral arteries, expresses only HLA-C, not HLA-A, B or D. Paternal HLA-C of extravillous trophoblasts stimulates maternal immune responses from both decidual T cells and the specialized uterine NK (uNK) cells, which express HLA-C binding KIR receptors. The higher the number of HLA-C maternal-fetal disparities, the greater is the number of activated decidual T cells (128). Both renal allograft rejection and PE are associated with circulating agonistic angiotensin II receptor AT1 autoantibodies (AT-1 AA) (129, 130).

5.5 Suggested molecular pathways to acute atherosclerosis

Our research group has, together with Professor Redman of University of Oxford, proposed a multistep pathway to acute atherosclerosis lesions (120). We have suggested that AA could represent the end-stage of different pathways (separately or in combination) that are immunological, inflammatory, genetic and hemodynamic (secondary to perturbed laminar blood flow in insufficiently remodeled uteroplacental spiral arteries), which converge to cause excessive inflammation around the distal ends of the spiral arteries in the decidua (the maternal part of the maternal-placental frontier).

Though pregnancy is a state of relative hyper- and dyslipidemia (131), foam cell formation can be stimulated in vitro by factors that promote inflammation, without the need for hyperlipidemia (132). Given the absence of an infectious organism, two drivers of inflammation exist in pregnancy, the maternal immune system – occupied by its continuous surveillance of fetal alloantigen, and the placenta - when it is in a state of dysfunction. We have proposed that excessive inflammation generated at the materno-fetal interface could possibly contribute to a common pathway in the development of acute atherosclerosis (120).

Figure 6 illustrates the development of preeclampsia by a multi-stage theory including the co-evolution and potential interaction with acute atherosclerosis, reprinted from Staff et al. (120).

Figure 6. Reprinted from Staff et al (120). with permission from the publisher. Acute atherosclerosis (Aa) and a five-stage model of preeclampsia. Previous models (Redman, C.W. Baillieres Clin. Obstet. Gynaecol 1992; 6 (3): 601–615 and Redman, C.W., Sacks, G.P., Sargent, I.L., Am.J.Obstet.Gynecol.1999;180:499–506) of preeclampsia have been extended to include Aa as a late stage in some pregnancies and to include the possible impact of pre-existing maternal chronic vascular inflammation. Continuous lines illustrate current consensus. Interrupted lines mark hypothetical pathways, yet to be demonstrated. We suggest four potential routes to Aa (A–D): (A) Poor placenta-tion (“placental preeclampsia”): Stage 1 comprises inadequate pre- and post-conceptual immune tolerisation leading to poor placenta-tion (Stage 2). Ensuing dysfunctional placental perfusion (Stage 3) is associated with placental oxidative, endoplasmic reticulum and inflammatory stress. The clinical signs of preeclampsia (hypertension and proteinuria) represent stage 4. Aa (Stage 5) can exacerbate dysfunctional placental perfusion and worsen the clinical disorder in a positive feed-forward loop (red arrows). (B) Normal placenta-tion (“maternal preeclampsia”). Chronic maternal vascular inflammation (such as in obesity, diabetes mellitus etc.) may generate preeclampsia and Aa, the latter then becoming a primary cause of placental abnormalities. In addition,

maternal vascular inflammation may hypothetically impair placentation without affecting tolerisation (interrupted arrow). (C) Poor tolerisation may be combined with chronic vascular inflammation leading to mixed syndromes, which appear to comprise the most severe preeclampsia presentations (see text). (D) An alternative (hypothetical) route to Aa. Partial tolerisation may not be enough to cause inadequate placentation, but still contributes to acute excessive decidual inflammation and the development of Aa, more likely in combination with B.

5.5.1 Acute atherosclerosis: linked to future maternal cardiovascular disease?

The manifestation of acute atherosclerosis following the convergence of states of excessive inflammatory and hemodynamic dysfunction justifies an interest in shared mechanistic pathways in the development of acute atherosclerosis and the development of atherosclerosis. We have proposed (120) that AA studies may assist in elucidating origins of arterial disease in general, the pathogenesis of preeclampsia in particular, and the interactions between the two conditions.

The association between acute atherosclerosis and preeclampsia is established. The association of preeclampsia and future risk of atherosclerotic cardiovascular disease is established at the epidemiological level (72, 74). The question remains, as discussed above in section 5.4.3; what are the mechanisms by which preeclampsia increases a woman's risk of future CVD?

Another question we pose is; could the histological manifestation of acute atherosclerosis, associating both with a dysfunctional placenta function, and with pregnancies affected by preeclampsia and excessive inflammation, help to identify those women most at CVD risk (73, 120, 133)?

Figure 7, redrawn from Sattar and Greer (134), illustrates that the “failed stress test of pregnancy”, resulting in preeclampsia, identifies women at risk for CVD due to vascular risk factors. Our hypothesis has been that AA may be a

better positive stress test to identify parous women at high future atherosclerotic CVD risk (120, 131).

Redrawn from:

Sattar, N & Greer IA. *BMJ* 2002

Figure 7. Redrawn by us and reprinted from Sattar and Greer (134) with permission from the Journal. The figure illustrates the constitutional “*stress test*” of pregnancy for women, relating incidence of preeclampsia in pregnancy to an increased risk of future maternal CVD. Our hypothesis is that acute atherosclerosis (images added to the original illustration) may provide a histological marker, indicating an increased risk of CVD among those with – or without a previous history of preeclampsia.

5.5.2 Why study acute atherosclerosis?

At the time of writing, our HAPPY PATH (Health after pregnancy complications and acute atherosclerosis) studies explore our novel hypotheses to develop better ways of identifying women at risk for severe CVD, up to decades before clinical disease, enabling follow-up and targeted intervention. Our group has proposed the novel concept that the study of placental arterial atherosclerosis can give vital information about the mechanisms of human

atherosclerosis in general besides identifying young women at CVD risk in particular (135).

In order to determine whether acute atherosclerosis indeed represents an immunohistochemical “stress test” for future maternal CVD, it is of pressing importance to have validated diagnostic criteria for the acute atherosclerosis lesion, as well as to identify optimal uteroplacental tissue from clinical practice to identify the lesions. Uniform criteria and conclusions of these issues are today lacking, and **represent the motivation for this PhD thesis.**

6. AIMS OF THE THESIS

Preeclampsia, a hypertensive complication of pregnancy, is associated with future risk of atherosclerotic cardiovascular disease, though the biological mechanism is unclear. The pregnancy specific arterial lesion of acute atherosclerosis of the uteroplacental circulation is an understudied female-specific arterial lesion. Acute atherosclerosis bears a morphological resemblance to the “fatty streaks” of early stage atherosclerosis, with intramural lipid-laden foam cells being a prominent feature in both lesions. Our hypothesis has been that acute atherosclerosis could be a histological hallmark helpful in stratifying a woman’s risk for developing future cardiovascular disease, decades prior to clinical manifestation of atherosclerosis. In order to test this hypothesis, the current PhD aimed to identify reliable criteria for lesion identification and location, as well as to characterize important morphological and immunological features of decidual spiral artery lesions of the pregnant uterus, across complicated (preeclamptic and diabetic) and uncomplicated pregnancies.

Specifically, the aims of the study were to answer the following questions:

- A. “*What*” are the useful criteria for defining acute atherosclerosis? We aimed to identify simple criteria that would allow reproducible comparability in future studies (**Paper I**).
- B. “*Where*” in the uteroplacental unit is acute atherosclerosis most readily detected? We aimed to identify which of three available non-invasive decidual (basalis and parietalis) sampling techniques was the optimal method for artery visualization and acute atherosclerosis detection, in preeclamptic pregnancies, where the rate of acute atherosclerosis is high (**Paper II**).
- C. “*Why*” are the acute atherosclerosis foam cells present in the arterial wall? We aimed to test our hypothesis of acute atherosclerosis as an inflammatory lesion,

with localized presence of increased numbers of perivascular lymphocytes, akin to atherosclerotic lesions of larger arteries (**Paper III**).

D. “*How*” does the presence of decidual (uteroplacental) acute atherosclerosis associate with preeclampsia, both clinically and immunologically? We aimed to identify if clinical pre- and pregnancy characteristics differed in relation to presence of acute atherosclerosis in either of the decidual locations and tissue sampling techniques (**Papers I, II and III**).

7. MATERIALS AND METHODS

7.1 Patient recruitment

This PhD study is based on information and biological samples from women recruited prior to elective cesarean section at the Department of Obstetrics, Oslo University Hospital, location Ullevål, which is the largest delivery unit in Norway with 7000 deliveries annually.

7.1.1 The Oslo Pregnancy biobank (OPB)

Since 2001 pregnancy sampling at cesarean delivery, and subsequent processing and storage of biological material has been a consent-based, protocol-led research initiative, supervised by Professor of Obstetrics and Gynaecology Annetine Staff. The Oslo Pregnancy Biobank, facilitating translation research within the clinical setting (www.oslo-universitetssykehus.no/opb), is approved by local authorities and Regional Committee for Medical and Health Research Ethics (REK) in South-Eastern Norway. Tissue samples from placenta (biopsies after delivery) and decidua (Decidual vacuum Suction Method), from maternal subcutaneous fat, the pyrimidalis muscle, amniotic fluid, and blood samples from the mother as well as from the fetus (umbilical cord: separate artery and vein samples) are collected at cesarean delivery after informed written consent from the patient.

Patient recruitment to the Oslo Pregnancy Biobank is an ongoing, prospective process. The women with preeclampsia, diabetes mellitus and normotensive (control) pregnancies used for the publications included in the current PhD study were recruited from 2001 to 2016. The decidua basalis tissue sampled with Staff's own developed technique of vacuum suction (62, 87), provides sufficient tissue volume, enabling studies from the maternal-fetal interface. The biological samples are processed and stored according to strict protocols.

All formalities in relation to sampling of biological material have been arranged by the main supervisor Professor Staff. The indication for cesarean delivery is clinician based, occurring independently of recruitment to the OPB research studies.

7.1.2 OPB patients selected for this PhD thesis

The pregnant women (n=297) contributing to this PhD study were recruited to the OPB between 2001 and 2016. Many patients from the 2012-16 period were recruited by the PhD candidate. Figure 8 illustrates the selection of patient samples used in **Papers I-III** of this PhD study.

Figure 8. Overview of the selection of patients recruited to the Oslo Pregnancy Biobank providing decidual tissue samples in **Papers I-III** of this PhD study. PE: preeclampsia, NC: normotensive control, DM: pregnancy affected by diabetes mellitus (all types, normotensive), DM+PE: diabetic, preeclamptic pregnancy. *spiral arteries not identified.

All samples are from pregnant women that underwent cesarean delivery with intact membranes, in the absence of labouring contractions, and without signs of infection. The majority of deliveries were elective, the women having fasted for at least 6-8 hours. Recruited women were categorised clinically according to the final pregnancy obstetric outcome into three main groups; controls (uncomplicated pregnancy), preeclampsia, and diabetics. The subgroups were divided into: normotensive controls (NC, n=114), preeclampsia (PE, n=117), diabetes mellitus (all types, pregestational or gestational, without preeclampsia, DM, n=45) and diabetes mellitus (all types) with superimposed preeclampsia (DM+PE, n=21).

Preeclampsia was defined “classically” as new onset hypertension (blood pressure $\geq 140/90$ mmHg) and new onset proteinuria ($\geq 1+$ on dipstick, or ≥ 30 protein/creatinine ratio) ≥ 20 weeks of pregnancy (34). Pregnancies complicated by diabetes mellitus (pregestational and gestational) were defined according to WHO 1999 guidelines (136).

Gestational week (GW) at delivery was defined by routine ultrasound screening at GW 17-20, offered as part of the free of charge antenatal routine care in Norway and attended by almost 100%. Newborn gender specific baby weight percentiles were calculated according to Norwegian fetal growth curves (137). Normotensive controls had uncomplicated pregnancies without clinical evidence of fetal growth restriction (FGR). FGR (used in **Paper II** and **III**) was defined as a birth weight percentile below 3% (either with or without antenatal pathological fetal Dopplers). Pathological fetal Dopplers were defined as abnormal PI (pulsatile index) in umbilical artery ($>2SD$), and/or cerebroplacental ratio <1.08 (138, 139). Maternal parameters included age at delivery (years), pre- or early conception and delivery BMI (body mass index; kg/m^2), nulliparity, previous history of preeclampsia, and blood pressures (mmHg; both during the first half of pregnancy, and at delivery).

7.2 Decidual tissue sampling methods

Figure 9, reprinted from Harsem et al. (87), demonstrates the three locations of the collection of endometrial (named decidua in pregnancy) tissue for this PhD study. This decidual tissue is the source for spiral artery identification and acute atherosclerosis evaluation of this PhD study. We utilised 3 decidua-providing tissue collecting methods;

1. DSM: Decidua vacuum Suction Method (yields decidua basalis)
2. BPS: Basal (maternal) Placental Surface (yields decidua basalis)
3. MR: (fetal) Membrane Roll (yields decidua parietalis).

Figure 9.Reprinted (with an added perforated red line for Membrane Roll illustration in Fig. 9a) from Harsem et al. (87), with permission from the publisher.

(a) The placenta in relation to the uterine wall, and the location of the decidua parietalis tissue harvested after the delivery of the placenta. MR: placental fetal membrane rolls.

(b) The uteroplacental circulation in the second half of pregnancy. Spiral arteries are the terminal branches of the uterine arteries; decidua basalis is the gestationally transformed endometrium with invading trophoblasts, situated between the placenta and the muscular uterine wall (myometrium).

(c) The location of decidual basalis tissue harvested after the delivery of the placenta: BPS, basal plate section; DSM, decidual vacuum suction method. PBB (placental bed biopsy) was not tested in the present PhD study.

7.2.1 Decidua vacuum Suction Method (DSM)

The decidual vacuum suction method samples tissue from the placental bed following the delivery of the placenta at cesarean section. Careful and extensive vacuum aspiration of the placental bed is performed as described previously by Staff et al. (62), illustrated in Figure 10.

During cesarean delivery, an intravenous bolus of oxytocin is given to the mother once the baby has been delivered, according to the Departmental routines for reducing postpartum hemorrhage. The placenta location is known from medical charts from regular antenatal ultra sound, and is again specifically located by manual palpation after delivery of the baby, then allowed to separate spontaneously from the uterine wall.

Tissue from the identified placental bed (the area underneath the placenta) is harvested, after removal of the placenta, with the vacuum suction force into a tissue-sampling box (using the vacuum curette routinely employed for surgical uterine evacuations). The sampling vial is and the collected tissue is flushed with 500 mL sterile saline solution at room temperature to remove excess blood from the decidual tissue.

The harvested decidual tissue (visually confirmed that no contamination with blood or membranes) is immediately fixed in neutral buffered 4% formaldehyde. Further, the tissue is processed and embedded in a paraffin block by the pathologist assistants at the Department of Pathology, according to routine. Decidua basalis tissue was finally confirmed on tissue sections by light microscopy and immunohistochemistry (IHC), where extravillous trophoblasts (only invading decidua basalis, not decidua parietalis) were identified following immunostaining with cytokeratin 7 (CK7).

Figure 10. A step-wise illustration of the Decidual vacuum Suction Method (DSM). A. Fetus in utero prior to cesarean delivery. The placental bed is located underneath the placenta. B. Abdominal incision in Cesarean section. C. The delivery of the fetus and placenta, exposing the placental bed. D. Vacuum curette inserted through the hysterotomy, and tissue “vacuum cleaned” from the placental bed is suctioned into a collecting pot, followed by saline water irrigation.

7.2.2 Basal Placental (maternal) Surface (BPS)

The delivered placenta, together with fetal membranes attached, is fixed in neutral buffered 4%v formaldehyde on arrival at the department of pathology. During routine processing, Basal Plate placenta Section (BPS) were excised from macroscopically normal looking placental tissue, followed by embedding in a paraffin block by the pathology assistants.

7.2.3 Fetal Membrane Roll (MR)

Following formaldehyde fixation, a segment of the fetal membranes (with the placenta attached) are rolled from the point where they have been surgically

breached (to allow delivery of the fetus) up to, and including, a small segment of the placental margin. The placental margin is continuous with the outer margin of the fetal membrane “roll”, whilst the fetal membrane furthest away from the placenta is at the centre. The roll is then transected like a loaf of bread and the membrane roll segment is embedded in a paraffin block.

7.3 Light microscopy of decidual tissue samples

7.3.1 Decidual tissue samples sectioning

For histopathological evaluation, 3µm thick serial sections were sliced from the paraffin embedded tissue sample blocks (one per decidual tissue collecting method, per woman), using a Micron HM 355 microtome (Thermo Fisher Scientific Inc., Waltham, Massachusetts, USA). Serial sections were mounted on SuperFrost slides (Menzel-Gläser, Braunschweig, Germany). Sections were de-paraffinised, then rehydrated according to Departmental routines, prior to immuno- and histochemical staining.

For DSM derived decidual tissue samples, at least 11 (and up to 21) serial sections were prepared. For BPS and MR derived decidual tissue samples, 5 serial sections were prepared respectively. The first 5 slides were used for acute atherosclerosis identification (**Paper I-II**), whereas the following slides 6,7,8 and 9 were used for immune cell typing (**Paper III**). The remaining slides were used if more slides were needed to perform the planned evaluations.

7.3.2 Immuno- and histochemical staining

Table 1 shows the overview of which immunohistochemical (IHC) stainings that were used for the numbered serial decidual sections, and the section slide numbers used for the separate **Papers I-III**.

Table 1. Overview of the immunohistochemical (IHC) stainings used for serial sections prepared from each decidual tissue sample evaluated, showing target antigen, purpose of stain, per 3 PhD **Papers (I, II, and III)**. Slides 2-4, 6-9 were also stained with PAS following IHC.

Serial section	Stain (n=nuclear, cy=cytoplasmic, tm=transmembranal, m=membranous)	Target antigen	Cell types identified in decidua	Purpose / identifying	Used in Paper I, II, III	Clone/Producer
1	Hemotoxylin&Eosin	Non-specific	Decidual tissue morphology	Decidual architecture, spiral arteries, thrombosis	I, II	Routine departmental staining
2	Monoclonal mouse anti-human Desmin (cy)	Desmin, intermediate between myofibrils	Smooth muscle cells, myometrial cells, myofibroblasts	Identification of spiral arteries, remodelling	I, II	DE R-11 / Roche Ventana Mediscal Systems, Inc, Arizona, USA
3	Monoclonal mouse anti-human Cytokeratin 7 (cy)	Cytokeratin (CK7), cytoskeleton	Trophoblasts, decidual glandular epithelium	Extravillous trophoblasts, perivascular leukocytes	I, II	OV-TL 12/30 / Dako Denmark AS, Glostrup, Denmark
4	Monoclonal mouse anti-human CD68 (cy, m)	CD68 scavenger receptor	Lipid-laden foam cells, macrophages	Foam cells,	I, II	KP1 / Dako Denmark AS, Glostrup, Denmark
5	Hemotoxylin&Eosin	as for section 1	as for section 1	as for section 1	I, II, III	as for section 1
6	Monoclonal mouse anti-human (n)	FOXP3, forkhead transcription factor family	Regulatory T-lymphocytes ("T-regs")	Lymphocytes of perivascular & interstitial decidua basalis	III	236A/E7 / Abcam plc, Cambridge, UK
7	Monoclonal rabbit anti-human CD3 (cy)	CD3, T-cell receptor	All T-lymphocytes		III	2GV6 / Roche Ventana Mediscal Systems, Inc, Arizona, USA
8	Monoclonal mouse anti-human CD8 (cy, tm)	CD8 binds HLA1	Cytotoxic /suppressor T-lymphocytes		III	C8/144B / Dako Denmark AS, Glostrup, Denmark
9	Monoclonal rabbit anti-human (cy, m)	CD56, adhesion molecule (NCAM)	Decidual natural killer cells		III	MRQ-42 / Cell Marque, Merck KGaA Darmstadt, Germany

The first and the fifth slide of every tissue section series underwent hemotoxylin & eosin staining according to standardized departmental routine. Every tissue slide that underwent immunohistochemical staining was

counterstained with Periodic acid Schiff (PAS) to aid in the identification of spiral arteries, and to aid identify presence of fibrinoid (140).

The immunohistochemical stains were performed by pathology assistants Giang Nygen and Linn Buer, either manually or utilizing the BenchMark XT (Ventana Medical Systems Inc., Tuscon Arizona, USA; an automated immunostaining system based on the “ABC” avidin-biotin-peroxidase method).

De-paraffination of tissue sections precedes antigen targeting, by incubation with the antibody (Table 1). Incubation with a secondary polyclonal antibody, conjugated to biotin molecules that can recruit avidin protein bound enzymes, allows for amplification. Colour detection was through enzyme-mediated horseradish peroxidase (HRP) labeled to diaminobenzidine (DAB) reporter molecules.

The immunostains were selected according to their specificity for the target antigen of interest (Table 1), availability and optimization in the department of pathology. Positive immunostaining for antibodies was confirmed using the following human tissue as controls: tonsillar lymphatic tissue (positive control for CD3, CD8 and FoxP3), and ileum small bowel (positive control for CD56).

All decidual tissue sections used in the PhD study were evaluated by the PhD candidate, after training with an experienced placenta Pathologist, using a Motic BA-410 LED light microscope under x4, x10 and x40 objectives. Images were recorded for documentation, or for further quantification of decidual lymphocytes, using a Moticom 2500 digital camera and Motic Images software version 2.0, Pictures were obtained using an Axio Scan slide scanner and Zen Blue software (Zeiss) (**Paper III**)

The systematic approach to the identification of spiral arteries was developed using the decidual tissue collected by DSM. This approach was then applied, with some modifications, to the decidual tissue collected with the two remaining methods, BPS and MR.

The worksheet developed for the systematic light microscopy examination across tissue samples and spiral arteries per patient is presented in Appendix 1: Microscopy worksheet for Paper I.

Figure 11. Illustration representing the sequence of slide-mounted serial sections of decidual samples developed using DSM-collected decidua basalis (Slides 1-5, **Paper I**), adapted for both decidua parietalis collected from fetal membrane rolls (Slides 1-5, **Paper II**), and decidua basalis collected from samples of placental basal plate surface (Slides 5-9, **Paper III**). Spiral arteries are represented by a dark red circle on each slide. The **Slide** is numbered, with both description of the stain used (s:), summarizing the target structure for identification (i:). Utilization of each slide, for each of the samples examined is explained in more detail from section 7.3.3-7.3.5 below.

7.3.3 The utilization of DSM sections 1-5 according to the PhD study aims

(Paper I)

7.3.3.1 Serial sections 1-2

The first step in evaluating presence of decidual acute atherosclerosis in decidual tissue was to detect the presence of decidual tissue spiral arteries. The topography of the decidua basalis tissue obtained from DSM is disorganized, containing decidual fragments from different areas of the placental bed over which the vacuum suction curette has passed. To ensure that spiral arteries were found on more than one serial slide (which was needed in order to diagnose acute atherosclerosis with more than one marker), a spiral artery cross-sections observed on the first slide of the tissue sample sections would be confirmed in the next (second slide, representing a depth of 3µm beneath the first).

Spiral arteries of the placental bed deliver maternal blood to the placental villi, whereas decidual nutrient arteries (“straight arteries”) do not. The nutrient arteries are typically <100µm in diameter and do not undergo the physiological transformation in the first half of pregnancy, in contrast to the spiral arteries of the decidua basalis (2, 88).

In order to avoid overlap with nutrient arteries, we required that a spiral artery section should fulfill the following criteria:

- a minimum outer-outer vessel wall diameter of >140µm
- the artery's media should contain either smooth muscle and/or uptake of amorphous PAS staining (positivity for presence of fibrinoid).

If no representative spiral arteries could be identified in the DSM tissue, then the sample could not be further evaluated for presence of acute atherosclerosis.

The presence of one or more representative spiral artery in a given tissue sample would lead to the search for vacuolated cells within every representative artery identified, from the subendothelial aspect on the luminal

margin, to the outer limit of the media, bordering with decidual cells (shown in figures 4 and 5, respectively).

Presence of fibrinoid and presence of smooth muscle contained within the confines of the artery wall was noted, modified from a categorization developed by Fiona Lyall (29).

PAS staining within the artery wall was graded according uptake by proportion of artery circumference (0: No PAS stain, 1: <25%, 2: 25% -49%, 3: 50%-74%, 4: $\geq$ 75%- 100).

Lastly, the sign of partial or complete thrombotic luminal occlusion for each artery section in each woman sample was noted in decidua basalis slides 1-2. Morphological appearance of thrombosis was defined as identification of red blood cell within a fibrin meshwork occupying >10% of the vessel lumen.

7.3.3.2 Serial section 3

Immunostaining with anti-CK7 antibodies allowed for identification of extravillous trophoblasts, thereby confirming samples of DSM decidua as representative of decidua basalis (88), arising from the placental bed, and thereby the spiral arteries visualized being integral in the uteroplacental circulation.

Presence of intramural CK7+ cells denotes endovascular extravillous trophoblasts, indicative of trophoblast presence and likely trophoblast associated remodelling in those arteries affected. Endothelial cells lining decidual glands would also stain as CK7+ (141), which would prevent glandular structures being misinterpreted as decidual blood vessels. Decidual gland and trophoblasts are easily morphologically distinct in the IHC sections.

Though present in all slides, perivascular mononuclear leucocytes were categorized as being greater to their background concentration in the surrounding interstitial decidua. We categorized this dichotomously per decidual tissue section (and per pregnancy) in **Paper I** as perivascular infiltrate (PVI) present or PVI absent.

7.3.3.3 Serial section 4

We used CD68 positivity corresponding to intramural vacuolated cells identified on serial section slide 1 (H&E) as confirmatory of lipid-laden foam cells (142). The CD68 positivity was quantified by number of CD68+ vacuolated cells per artery section, as well as by the CD68 stained surface area (μm^2) proportional to the entire artery wall area.

7.3.3.4 Serial section 5

Since section slides 1 and 5 were both stained with H&E, this provided in essence an ‘H&E’ sandwich”, allowing confirmation that both the artery and the intramural vacuolated cells it contained as a continuum from serial section slide 1 through to, and including, serial section slide 4.

7.3.4 Modifications to our systematic approach (7.3.3) for the examination of serial section slides utilized for **Paper II**: BPS and MR decidual tissue sections

For the acute atherosclerosis studies on decidual basalis tissue derived from the BPS samples used in **Paper II**, no modification was needed from the DSM procedure described in 7.3.3.

Tissue sections went through both placenta villi and through the decidua basalis represented in the Nitabuch’s layer that merges with the anchoring villi of the placenta. Spiral arteries could be identified within the decidual tissue.

Under the conditions of normal light microscopy, discerning a spiral artery from a nutrient artery in the decidua parietalis may be challenging, given that spiral arteries of the decidua parietalis are not expected to have undergone the EVT-dependent transformation of those spiral arteries located in the placental bed (decidua basalis) (2, 88). Therefore, when evaluating decidual sections from MR samples we included all arteries, both $<100\mu\text{m}$ as well as those $>100\mu\text{m}$ in diameter.

The location of a given artery along the length of the membrane roll was categorized according the “number of turns” (rolls) it was situated from the outer margin of the MR (always marked by the membranes originating from the placental margin). We reported acute atherosclerosis location according to number of full circles of tissue winding from the point furthest away from the placenta (placenta representing the first circle and the outer membrane border representing the inner circle of the roll).

7.3.5 The purpose of DSM sections 6-10 according to the PhD study aims (Paper III): lymphocyte quantification in decidua basalis

Table 1 illustrates the choice of immune cell staining with IHC used in Paper III. Following immunostaining, for each woman sample of decidua basalis (serial sections 6-9) immunopositive cells, denoting specific lymphocyte populations (summarized in Figure 12), were counted manually by the PhD candidate. Since the CD3 stained slide was adjacent in the sequence of slides to the slide stained with CD8 ($3\mu\text{m}$). In the absence of our use of a CD4 immunostain, it allowed for estimation of a CD3+CD8-T cells (presumptive CD4+ cell population) by subtracting the concentration of CD8+ T-cells from CD3+ T-cells, in the same tissue zone investigated.

Reproducible concentrations of lymphocytes (cells/mm²) were obtained by performing cell counts to predefined areas, named Zone 1-3, illustrated in Figure 11. The three predefined spiral artery related areas included:

- **Zone 1**, “*intramural*”, represents the entire area for each of the decidua basalis spiral artery wall, starting from the endothelium lining the lumen, and radiating outward to the outer limit of the smooth muscle/PAS-positive margin of the spiral artery. (Figure 12A and B).

- **Zone 2**, “*perivascular*”, represents the immediate adjoining area for each of the surrounding the spiral artery walls, demarcating a “cuff” of 100µm decidua tissue width, radiating from the outer limit of the artery wall (Figure 12A and C).

- **Zone 3**, “*decidual infiltrate*”, represents the area containing the largest visible concentration of mononuclear leucocytes present in the surrounding decidual tissue continuous with the spiral artery(-ies) evaluated. The 3 largest clusters of CD3+ cells were identified on low power magnification (x4) and demarcated by a counting disc (100µm in diameter, representing a surface area of 7853µm²) (Figure 12A and D). For each immunostain for each woman sample, counting of positively stained cells in Zone 3 was performed at x40 magnification.

Figure 12. Illustration and photos of predefined Zones 1-3 for quantification of lymphocytes in and around decidua basalis spiral arteries (All PAK).

A) Schematic representation of a decidual spiral artery and the 3 different zones used for quantification of decidual lymphocyte concentrations.

The star (*) denotes the artery lumen.

B-D); Examples of the applied zones for a decidual spiral artery section with acute atherosclerosis stained for CD3.

B) Zone 1: “intramural”, spiral artery wall (red with solid black perimeter).
C) Zone 2; “perivascular”, area surrounding the intramural zone. The outer perimeter is set at a depth of 100µm from the outer margin of the artery wall (orange with interrupted black perimeter).

D) Zone 3; “interstitial infiltrate”, circle-shaped area of 100µm in diameter in surrounding decidual tissue not directly in contact with the spiral artery but with a continuous connection to the spiral artery (circle with interrupted black perimeter).

7.3.6 Testing the reproducibility of morphological observations (**Paper I**) and lymphocyte cell counts (**Paper III**)

Morphological observations leading to the new classification of acute atherosclerosis in decidua basalis were tested for inter-observer reproducibility between the PhD candidate (PAK), Professor Fiona Lyall (FL, Professor of Maternal and Fetal Health, specializing in spiral artery research), and Professor Borghild Roald (BR, Professor and placenta pathologist).

Decidual slides with spiral arteries from 16 different women were evaluated on one occasion (PAK versus FL) and from 58 women on two occasions (PAK versus BR). Intra-observer agreement was evaluated for 58 women with spiral artery sections on 2 occasions 8 weeks apart (PAK vs PAK).

The inter-observer reproducibility of lymphocyte cell counts for **Paper III** (using CD3+ cells) was evaluated by the author (PAK) and placenta pathologist professor Borghild Roald (BR). Evaluating a total of 31 spiral arteries (10% of total arteries studied), and a corresponding number of decidual infiltrates, the Kappa scores (allowing for a margin of error of +/-1 between cell counts) showed intra- and inter-observer agreement(143) for Zone 1 as *good* and *very good* (respective Kappa score 0.69 and 0.81); for Zone 2 *fair-moderate* and *moderate* (respective Kappa score 0.40 and 0.55); for Zone 3 *good* (respective Kappa score 0.63 and 0.62).

7.4 Statistical analyses

Data were analysed using SPSS version 22. The Fisher Exact test was used for dichotomous variables, and for non-parametric variables the Mann-Whitney, or Wilcoxon, test was applied. A probability of ≤ 0.05 was considered statistically significant.

Spearman's rank correlation coefficient was used when analysing associations between continuous variables (lymphocyte cell concentrations) and continuous clinical variables.

7.5 Ethical considerations

The pregnancy biobank studies were approved in 2001 by the Regional Committee for Medical Research Ethics, Eastern Norway (REK Sør-Øst), permission from the Norwegian Data Inspectorate (Datatilsynet) has been obtained, and the biobank has been approved by the Directorate for Health and Social Affairs: Sosial- og helsedirektoratet and REK Øst. Subsequent permission for the OPB as a thematic research biobank was given in 2015 (by the Regional Committee for Medical Research Ethics, South-Eastern Norway (REK Sør-Øst)).

All participating women have signed a voluntary informed consent and they could withdraw from the study at any time without giving any reason and without any consequences regarding their relationship to their medical care at Oslo university hospital.

8. SUMMARY OF RESULTS

8.1 Redefinition of acute atherosclerosis in DSM collected decidua basalis tissue (Paper I)

8.1.1 Identification of spiral arteries, fibrinoid, thrombosis and perivascular infiltrate

Using decidua basalis DSM serial sections 1&2, one or more representative artery sections (diameter $>140\mu\text{m}$, as defined in 7.3.3.1) were identified in 85% (237 of 278) pregnancy samples of DSM decidua basalis. The evaluations of these 237 samples in which spiral arteries were identified provides the tissue basis for the redefinition work of acute atherosclerosis.

The median number spiral artery sections, identified per decidua basalis tissue section (based on serial sections 1 and 2), was 3.5 (range 1 - 11). Presence of artery wall acellular fibrinoid (PAS positive) was almost universal for those arteries identified, involving 75-100% of the circumference of the artery section evaluated (and corresponding absence of smooth muscle (Desmin staining) in 3/4 of pregnancy samples. This finding was expected since, in late pregnancy, decidua basalis arteries have undergone remodelling distally, and remodelling is accompanied by fibrinoid deposition (144). Using only PAS staining alone, it was difficult to measure the quality or intensity of the fibrinoid, thereby reliably discerning “physiological fibrinoid” from fibrinoid “necrosis”.

The morphological appearance of partial or complete thrombosis in at least one artery section per decidua basalis tissue sample was present in 37% (88 of 237) of participant samples.

Using the serial section slide 3 derived from DSM decidua basalis, a categorical definition of perivascular infiltrate (PVI) was established in at least one representative artery section in 26% of the participant samples (62 of 237).

8.1.2 Criteria for defining a foam cell lesion

Using the serial section slide 1 and 4 from DSM decidua basalis serial sections, the hallmark of acute atherosclerosis was identified as the (minimum) presence of at least two vacuolated intramural cells adjacent to one another (serial section slide 1), with a confirmatory CD68+ staining in serial section slide 4.

Intracellular vacuolation on H&E staining alone, in the absence of CD68 positivity, is not specific to foam cell formation. This morphological feature was found to be present in at least one spiral artery section in 81% of the decidua basalis tissue samples. Intramural CD68 positivity alone was also a common finding (≥ 1 artery sections in 50% of decidua basalis samples). In the absence of vacuolation (serial section 1), CD68 positivity (in serial section slide 4) was considered as indicative of macrophages, not foam cells, localized in the artery wall.

8.1.3 PAS positivity is not a criterion for defining “fibrinoid necrosis”

In decidua basalis tissue there was a largely universal presence of PAS, and the almost complete replacement of decidual spiral artery smooth muscle with intramural fibrinoid (Desmin combined with PAS staining on serial section slide 2). We found that intramural fibrinoid (intramural PAS positivity) was ubiquitous in the spiral arteries of the decidua basalis, and never absent from spiral arteries containing foam cell lesions.

8.1.4 Perivascular infiltrate (PVI)

Based on a subjective impression of leukocyte infiltrates surrounding decidual spiral arteries (serial section 3, so as exclude fetal EVT's from counts), there was a statistically significant difference in leukocyte numbers (assessed by PAK, counting mononuclear leukocytes according to a prescribed perivascular area) between those arteries identified with PVI as compared to those without PVI (data not included in **Paper I**).

PVI was present in at least one artery section in 26% of decidua basalis samples, but was not found to co-localise with intramural foam cell lesions in **Paper I**.

8.2 Criteria for a revised diagnosis of decidua basalis acute atherosclerosis (Paper I)

As previous definitions have suggested three criteria to define acute atherosclerosis, we reviewed the presence of these criteria in our cohort of 237 decidua basalis DSM tissue sections:

1) Foam cell lesions were identifiable morphologically, and could be confirmed using CD68 positivity. However by requiring a minimum of ≥ 2 adjacent intramural CD68⁺ vacuolated cells, we reduced the risk of including incidental solitary vacuolated cells / macrophages mistaken as foam cells. The intra-observer agreement (PhD candidate PAK vs PAK) for this criterion was almost perfect (Kappa score of 0.86). Inter-observer agreement was “substantial” with Kappa scores between PAK vs FL, and PAK vs BR (0.64 and 0.70, respectively).

2) “Fibrinoid necrosis” was not possible to reliably discern from presence of fibrinoid secondary to the physiological remodelling of the decidual spiral

arteries evaluated. In the setting of the decidua basalis, fibrinoid identified through PAS staining was not specific to foam cell lesions.

3) Perivascular infiltrate was manifest surrounding a proportion of the spiral artery sections, but there was no exclusive co-localisation of PVI with spiral arteries manifesting foam cell lesions.

8.3 Additional morphological features of decidua basalis associated with acute atherosclerosis according to our simplified redefinition (Paper I)

The morphological appearance of spiral artery thrombosis (partial or complete, in at least one artery section) had a higher rate in pregnancy samples where AA was detected. This was true for all pregnancy categories, the incidence being significantly higher in non-diabetics (both PE and normotensive controls).

AA detection was also associated with the absence of endovascular (intramural) extravillous trophoblasts in the affected artery section.

8.4 Clinical features associated with revised acute atherosclerosis definition (Paper I)

The overall rate of acute atherosclerosis in the heterogeneous pregnancy study group was 22% (51 of 237 pregnancy samples). The diagnosis of at least one foam cell lesion (comprising at least two CD68+ vacuolated adjacent intramural cells) contained in one spiral artery section was highest in those pregnancy samples affected by preeclampsia (preeclampsia 37%, diabetes – all forms - with superimposed preeclampsia 29%), but not diabetes (normotensive controls 11%, normotensive diabetics 10%). In non-diabetics the incidence in preeclamptics was significantly higher ($p < 0.001$). The number of affected artery sections per pregnancy sample was greater in those pregnancies affected by PE compared to normotensive controls, as was the median number of nuclei per foam cell lesion.

The preeclamptic group had both a greater rate of PVI, and artery sections with 75-100% intramural smooth muscle replaced by fibrinoid (PAS) as compared to normotensive controls, but these differences did not relate to presence or absence of decidua basalis AA.

Presence of acute atherosclerosis in a pregnancy sample was not associated with maternal parameters (age, BMI, BP) when compared to the same pregnancy category, where AA was absent. However, presence of acute atherosclerosis was associated with parous normotensive controls with a history of previous preeclampsia.

The fetoplacental parameters of gestational age at delivery and birthweight percentiles showed lower median values in pregnancies where AA was detected compared to where AA was absent. Though non-significant, this trend was present in all pregnancy categories of our original study cohort (**Paper I**).

When the cohort was expanded for more preeclamptic pregnancies (**Paper II**), both gestational age and birthweight percentile, was significantly lower in preeclamptic pregnancies where decidua basalis AA was detected (see [8.5.2](#)).

8.5 Comparison of decidua collecting methods for acute atherosclerosis identification in preeclampsia (Paper II)

Preeclamptic women (n=107) from the OPB were selected to the study if as they had decidual tissue collected by three different sampling methods; Decidual vacuum Suction Method (DSM: decidua basalis), Basal Plate Surface (BPS: decidua basalis) and fetal Membrane Roll (MR: decidua parietalis).

8.5.1 Identification of spiral arteries and rate of acute atherosclerosis

Of the three collecting methods, DSM sampled decidual tissue provided the highest yield of spiral artery sections and the highest detection rate of at least one artery section containing acute atherosclerosis (n=33 of 94). MR decidual tissue collection provided a higher yield of spiral artery sections as compared to BPS, but the number of samples where AA was detected was the same for these two clinical routine collection methods (n=8 of 76 vs n=8 of 53). Table (2) (reproduced from Paper II, with kind permission of the publishers) shows the rates of spiral artery identification and AA detection according to decidual collection method.

Table 2. (Reprinted from **Paper II** submitted to Virchows Archives) Rates of decidual spiral artery and acute atherosclerosis presence, according to decidual issue type for the 107 preeclamptic women included in the study. Results are expressed in numbers and rates (%). DSM: decidual vacuum suction method of placental (yielding decidua basalis); BPS: placental basal plate section method (yielding decidua basalis); MR: membrane roll (yielding decidua parietalis).

Tissue	Decidua basalis		Decidua parietalis	Decidua basalis+parietalis	
	DSM	BPS		MR+BPS	DSM+BPS+MR
Spiral artery detection rate	94/107 (87%)	53/107 (50%)	76/107 (71%)	85/107 (79%)	102/107 (95%)
Acute atherosclerosis rate (of samples with spiral arteries present)	33/94 (35%)	8/53 (15%)	8/76 (11%)	15/85 (18%)	37/102 (36%)

The Venn diagram in Figure 13 (reproduced from **Paper II**, submitted to Virchows Archives) represents those women in whose pregnancy samples AA was detected, demonstrating those cases where AA detection was detected by either a single collection method, or where AA was detected by more than one collecting method for the same woman. Of the 37 women where AA was detected with at least one tissue sampling method, 33 of the AA were identified in DSM tissue, of which 22 exclusively. AA detected in BPS was

always represented in one or both of the other two collecting methods. In four cases, AA identification was limited to MR collected decidual tissue.

Figure 13. Venn diagram representation of AA detection in uteroplacental tissue types for the 107 preeclamptic women included in **Paper II**.

DSM (bright red): Decidual vacuum suction method (collecting decidua basalis), BPS (dark red): Basal plate placenta section (collecting decidua basalis), MR (green): Membrane Rolls (collecting decidua parietalis).

Though in the minority, two-thirds of MR spiral arteries with a diameter $>100\mu\text{m}$ were localized proximal to the placental margin (within the first complete “turn” of the membrane roll). None of the MR spiral arteries contained intramural trophoblasts.

8.5.2 Association of acute atherosclerosis with clinical features of preeclamptic pregnancies, according to decidual collection method (DSM, BPS, MR)

When decidual collection methods were evaluated separately, only the presence of AA in DSM samples (as compared to absence of AA in DSM samples) associated significantly with lower gestational age at delivery and

lower birthweight percentile ($p=0.025$ and $p=0.039$ respectively). When combining BPS (decidua basalis) and MR (decidua parietalis) collection methods, presence of AA associated with a significantly lower gestational age at delivery (but not birthweight percentile), when compared to BPS+MR samples where AA was absent ($p=0.025$). When combining samples of all three collection methods (DSM+BPS+MR) from each woman, AA presence was significantly associated with a lower gestational age at delivery ($p=0.007$), and with a lower birthweight percentile ($p=0.034$), compared to those women in whom tissue arteries were detected, but where AA was absent.

Rates of fetal growth restriction (FGR; defined in [7.1.2](#)) were high in preeclamptic pregnancies with AA detected as compared to when AA was absent, irrespective of decidual collection method. Pathological fetal Dopplers, regardless of birth weight percentile, were significantly more frequent in preeclamptic women with AA in BPS+MR as compared to women without AA in these decidual tissues.

We found no significant associations between presence of AA (identified in either single or combinations of decidual tissue collection methods) and other clinical parameters associated with preeclampsia risk: maternal age, maternal BMI, mean blood pressure registered in the first half of pregnancy, infant sex, primigravity, or previous history of preeclampsia.

8.6 Lymphocyte quantification in prespecified DSM collected decidua basalis tissue zones (Paper III)

In general, scattered lymphocytes staining of CD3+, CD8+, and CD56+ were observed throughout the decidual tissue in both preeclamptic and normotensive pregnancies. Lymphocytes were observed in all three predefined zones, across decidual tissue samples with and without AA. The highest concentrations of

immunopositive cells were observed in the decidual infiltrates (Zone 3) continuous with spiral arteries, followed by the perivascular area (Zone 2). In Zone 1, numbers of CD3+, CD8+, and CD56+ cells were very low or absent. FoxP3+ staining was scarce or absent in all zones investigated, both in PE and normotensive pregnancies.

8.6.1 Zone 1 (intramural)

Though lymphocyte numbers were low, both decidua basalis tissue samples and individual arteries affected by acute atherosclerosis were associated with higher concentrations of intramural CD3+ cells, both in preeclamptics and normotensives. Also the CD3+CD8- cell concentrations (inferred CD4+ T-cells) were higher in acute atherosclerosis arteries, as was CD8+ cells, though the latter difference was restricted to preeclamptics only. CD56+ cells significantly lower in concentration compared to the targeted T-lymphocytes. T-reg (FOXP3+) cells were most absent.

8.6.2 Zone 2 (perivascular)

Limited to preeclamptic pregnancy samples only, CD3+ and CD3+CD8- (inferred CD4+) concentrations were significantly higher in spiral artery sections with AA, compared to spiral arteries without AA. In Zone 2 concentrations of CD8+ cells were not associated with presence of acute atherosclerosis in individual artery sections, nor in a AA-affected decidual tissue sample. Concentrations of CD56+ cells were similar to those of CD8+ cells, but bore no association with artery or decidual tissue sample section with or without acute atherosclerosis. The distribution of lymphocytes in the decidua immediately surrounding spiral arteries was uneven, and bore no relation in proximity to the presence of intramural foam cell lesions.

8.6.3 Zone 3 (decidual infiltrate)

Decidua basalis tissue samples with acute atherosclerosis were associated with higher concentrations of CD3+CD8- (inferred CD4+) cells located in the decidual infiltrates surrounding spiral arteries. Like in Zone 2, this was a finding limited to the preeclamptic pregnancies.

9. GENERAL DISCUSSION

9.1 Methodology discussion

9.1.1 Patient inclusion to the Oslo Pregnancy Biobank

Patients were selected to this PhD study from the Oslo Pregnancy Biobank (OPB), ensuring standardization of both tissue sampling and clinical parameters recording. Pregnant women were not recruited to the OPB only for the purpose of this PhD study, rather, they were originally recruited for multiple studies *“to determine the mechanisms underlying why some women develop pregnancy-related complications..”* and to *“study whether there is a connection between placental conditions and the future health of both mother and child”* (text from the English version of the patient informed consent form). Pregnant women were approached for inclusion based on the presence of pregnancy pathology (e.g. diabetes, preeclampsia) or normality (normotensive controls). The biological samples and data from the OPB are the basis for a multiple of studies and publications (87, 111, 130, 145-155).

9.1.2 Patient selection from the Oslo Pregnancy Biobank for **Papers I-III**

Only women included to OPB who were delivered by cesarean section were suitable for the investigations of this PhD. This is because the decidual suction method (DSM) for collecting decidua basalis is only available in the cesarean section setting. Any form of deliveries recruited to OPB could have contributed routine collection of placenta basal plate surface (BPS) (decidua basalis) and decidua parietalis from fetal membrane roll (MR), but only women with cesarean section were included, as we wished to compare the three decidual tissue types based on a uniform delivery setting (**Paper III**).

The rates of pregnancy complications among the selected pregnancy samples for **Paper I** do not reflect the general rates of the obstetric population in

Norway or at Oslo University Hospital, Ullevål. Preeclampsia (PE) is overrepresented in **Paper I** (36% compared to 3.4% preeclampsia in Norway during the patient recruitment period (156)), as are those pregnancies complicated by diabetes (24% compared to 5% (157)). Pregnancies were selected from OPB so as to maximize the number of pathological pregnancies, especially preeclamptics, as acute atherosclerosis is especially prevalent in this pregnancy group (86, 89, 95, 111, 158, 159). This PhD study also included a large number of normotensive pregnancies as controls for comparison since there are only a few reports of acute atherosclerosis in uncomplicated and normotensive pregnancies (91, 111).

Figure 8 (Methods section [7.1.2](#)) illustrates the participants and sampling included in **Papers I-III**. In **Paper II**, we wished to compare which of the three decidual collection methods in order to determine which was best for detecting spiral arteries and acute atherosclerosis. Given that three collecting methods were to be compared, we limited the patient selection to the single pregnancy complication where acute atherosclerosis had the highest detection rate: preeclampsia. Furthermore, only preeclamptic women with tissue samples collected by all three methods were included. All preeclamptics from **Paper I** fitting these criteria were included in **Paper II**. In order to increase the number of pregnancies for analysis, tissues from preeclamptic women subsequently recruited to the OPB (following recruitment of **Paper I** patients to OPB) were added to the pregnancies evaluated in **Paper II**.

The aim in **Paper III** was to determine a potential association between presence of previously defined acute atherosclerosis (**Paper I**) and decidual lymphocyte concentrations, in both preeclamptic and normotensive pregnancies (totaling four groups for comparison). Diabetic pregnancies were excluded in **Paper III** due to low sample numbers, and for the possible

confounding effect the diabetic environment might have on decidual lymphocyte concentrations. Given that we had previously categorized samples according to presence of perivascular infiltrate (PVI, in **Paper I**), an effort was made to ensure that these PVI samples were distributed equally between those identified with acute atherosclerosis, and between those where the lesion was absent. Additional pregnancies and tissue samples were added to each group in order to increase the total number, from those pregnancies and decidual samples where PVI was not previously investigated. Selection of additional patients (pregnancies) was random, but pregnancy samples with higher numbers of previously identified artery sections (**Paper I**) were favored, so as to increase the total number of spiral arteries available for evaluation of acute atherosclerosis lesions in an otherwise relatively small study group (n=94).

9.1.3 Patient clinical parameters and pregnancy category

Patient details regarding clinical parameters were recorded in the OPB at recruitment. There was a thorough extraction of information from the available hospital journal and the woman's national antenatal chart, reflecting the measurements from her antenatal visits (predominantly by the primary health sector). The thorough and comprehensive clinical data collection implied almost no missing data. The PhD study only included one pregnancy where the preconception BMI could not be calculated due to missing bodyweight. There was only one pregnancy included in the PhD study where a mean value for blood pressure prior to 20 weeks of gestation could not be calculated in the normotensive controls.

Fetal Doppler values, used in the assessment of fetal growth restriction, vary in their standard deviations according to the reference group used. Though it is arguable whether this reference group was the most appropriate for the mixed maternity population at Ullevål, the reference group of Arduini and Rizzo

(138) was used for the clinical management of pregnancies during the time of recruitment to OPB (2001-2016).

The pregnancy category to which a patient recruited to the OPB was classified was not only decided by the recruiting investigator, but also by a review panel within the research group (a Diagnosis Revisory Group), including experienced obstetric clinicians, in order to reduce misclassification errors to a minimum.

9.1.4 Methods of tissue sample collection and preparation

This study of acute atherosclerosis is founded upon the investigation of spiral arteries located in decidual tissue (**Paper I**), the interface between the mother's immune system and fetal cells. Our primary choice of Decidua vacuum Suction Method of the placental bed has been previously shown to yield the highest number of spiral arteries compared to placental basal plate surface and placental bed biopsies (87), although this study was not done on all three tissue types from the same women. DSM collection limits investigation of uteroplacental spiral arteries to their decidual segments, the myometrium being largely untouched using this method. DSM limits sample collection to cesarean delivery, as collection following vaginal delivery of the fetus and placenta would require a degree of invasiveness that would probably be considered unacceptable by most mothers (and ethical committees).

Some authors have argued that acute atherosclerosis is a manifestation of the myometrial arteries alone (48), whilst others have demonstrated that the acute atherosclerosis lesion is found in greater abundance in the decidua (53, 104). Two sampling methods allow for examination of myometrial decidual arteries. A total hysterectomy performed during delivery can allow for individual spiral arteries to be followed from their superficial level in the decidua basalis, to

their deep myometrial origins at the radial branches of the uterine artery. This “gold standard” of spiral artery examination has been described by Brosens (160), though the availability of such whole uteri is sporadic and limited. The risks of surgical hysterectomy during delivery are high, and ethically not permissible if only done for research purposes. However, historical tissue samples have been investigated in series where hysterectomy was done during delivery, with the placenta in situ, as a mean for sterilization in a Catholic hospital (9, 10, 26).

Comparable to total uterine specimens, placental bed biopsies following the delivery of the baby and placenta also allow for examination of the spiral artery in the myometrium, in addition to decidual spiral artery assessment. The drawback with this method is however the low yield of spiral arteries, as described previously by us (87), and others (53, 161, 162), along with the small yet additional risk of uterine perforation with excision sampling the placental bed. We did therefore not perform placental bed biopsies in any of the patients recruited to OPB, and such tissue samples as a source of myometrial spiral arteries were not available for this PhD thesis.

We included decidual tissue from the BPS and MR collection methods for this PhD study (**Paper II**), in order to directly compare the detection rate of acute atherosclerosis in decidual spiral arteries using DSM vs BPS/MR. The rationale was that BPS and MR collected decidual tissues are not dependent on cesarean section for their collection. If these methods were shown to be comparable to DSM in detecting acute atherosclerosis, this would mean that any delivery could easily be assessed for presence of decidual acute atherosclerosis.

9.1.5 Decidual tissue sample staining

9.1.5.1 Sectioning and immunohistochemical staining

All tissue samples collected were processed by the same Department of Pathology at Oslo University Hospital, Ullevål. Formalin fixation and paraffin embedding ensured the longevity of sample blocks, but at the same time put restrictions on the staining methods that could be utilized in our studies.

Direct detection of lipid laden foam cells has been described using oil red-O staining (90). Best suited to frozen sections, such staining must be undertaken prior to paraffin embedding, as the subsequent process of deparaffinization results in loss of intracellular lipid, leaving behind empty cell vacuolation(s). Intracellular vacuolation may also arise from other intracellular processes/pathologies unrelated to foam cell development.

It was therefore necessary, without direct detection using oil red-O being amenable, to employ CD68 (a lipid scavenger receptor) as a marker for identifying foam cells, thereby confirming the morphological characteristic of cell vacuolation (due to tissue processing effect of lipid extraction) as indicative of foam cell presence. Evidence of vacuolation without confirmatory CD68 staining suggests a different lesion than foam cell formation. Presence of CD68+ intramural cells in the absence of vacuolation is also not consistent with foam cells and suggests macrophages.

The process of formalin fixation and paraffin embedding of samples reduced the suitability of using immunofluorescence as a staining technique for quantifying targeted cell types. In addition, the need to visualize the decidual architecture whilst quantifying cells within prescribed tissue zones necessitated the need for direct visualization using light microscopy. For this reason we chose immunostaining using colour detection through enzyme-mediated

horseradish peroxidase (HRP) labeled to diaminobenzidine (DAB) reporter molecules.

Single antibody staining on thin (3µm) serial sections of decidua samples was chosen over double/multiple antibody staining on the same sample section, a method also employed by van Dijk (163). On testing, we found that our purchased CD4 immunostain (from Novocastra Leica Biosystems CD4 clone 1F6) showed too much background staining to allow for the accurate counting of individual immunopositive cells. In addition, it would stain more than the cells of interest (CD4+ T-lymphocytes), as it also stains dendritic cells and CD4+ macrophages. Instead, we indirectly calculated CD4+ T-lymphocytes by subtracting CD8 immunopositive cell from the CD3 immunopositive cell counts ($CD4 = CD3 - CD8$), in the same prescribed zone. This indirect method has also been used by Stallmach 1999 (106) and Sasaki 2007 (164). A disadvantage with this method is that it is only an indirect quantitative evaluation of CD4+ T-lymphocytes. Nor does it distinguish the heterogeneous subgroups of CD4+ T-lymphocytes that vary in their biological action. Thus differences in quantity do not necessarily reflect differences in immune response.

9.1.5.2 Choice of histochemical staining for fibrinoid: some background and controversy

Periodic acid Schiff (PAS) has previously been described to confirm presence of fibrinoid (88, 92, 140). However, the identification of fibrinoid in the presence of positive PAS staining alone provides no clear distinction of the causes or the pathophysiological processes involved, nor the speed of its evolution.

Fibrinoid was described by Neumann in 1880, “*as a chemical change of the intercellular substance of connective tissue, causing swelling, homogenization*

and conversion into a fibrin-like substance” (Movat, 1958(93)). The ”process” of fibrinoid was seen as degenerative as it often resulted in the loss of cellular structures (e.g. smooth muscle cells, or extravasation of plasma). Necrosis was *assumed* to be one of the many ways in which this cellular loss could come about (as supposed to apoptosis).

In the over 70 years since the first description of acute atherosclerosis (82, 83), the presence of (PAS positive) fibrinoid has been characterized as arising from “degeneration” (82) or “necrosis” (1, 83, 103). In the last fifty years, there has been a growing understanding regarding the transformation of spiral arteries located in the placental bed (myometrial and decidua basalis), characterized by the physiological replacement of their smooth muscle walls with fibrinoid and endovascular extravascular trophoblasts (165). On the basis of PAS positive staining alone, it is a challenge for investigators to differentiate fibrinoid arising from physiological remodelling of spiral arteries of the decidua basalis, from the “fibrinoid necrosis” associated with acute atherosclerosis lesions; intramural foam cells located in spiral arteries deep in the myometrium or decidua parietalis. Respected placenta expert, Robert Pijnenborg, suggested a subtle colour difference in PAS staining, distinguishing the fibrinoid associated with acute atherosclerosis (“grey-blue”), from the fibrinoid associated with spiral artery remodelling (“brilliant pink”) (88). However, this visual difference has not been standardized nor validated across or within evaluators, nor well documented in colour pictures in previous publications.

Differences in the appearance of PAS positive fibrinoid may not necessarily indicate different etiologies. Following comparison of the presence of fibrinoid in a variety of different vasculopathies, Jellinek commented that the *“development of vascular fibrinoid depends on the intensity and duration of vascular damage, but is unrelated to the etiology”*, concluding that, *“the morphological appearance of the lesions depends on the duration and intensity*

of the etiological factor, but is unrelated to its nature” (166). Jellinek’s conclusions support this thesis evaluation of the difficulty in assuming differing etiologies underlying differing fibrinoid amount, texture, and colour. Differentiation of fibroid etiology has been investigated using antibody staining of the components of placental fibrinoid (e.g. collagen, fibrin, fibronectins). By this immunohistochemical approach, Kaufmann et al. distinguish between “matrix-type” fibrinoid (assumed trophoblast in origin), and “fibrin-type” fibrinoid (assumed maternal in origin) (167). This study did however not mention decidual spiral arteries in tested samples. Antibody detection of decidual basalis spiral artery wall fibrinoid components was not attempted in our study, due to limitations in technical and budgetary resources.

The ubiquity of fibrinoid known to be present normally in remodeled decidual basalis spiral arteries rendered the detection of intramural PAS positivity redundant in this thesis as an aid in the detection of decidual basalis acute atherosclerosis (**Paper I**). However, this thesis showed that foam cell lesions were never detected in the absence of decidual spiral artery wall PAS positivity, in any of the three decidual tissue types investigated (**Paper II**).

Following the work on this thesis, we have added another histochemical stain in order to improve future decidual spiral artery fibrinoid necrosis detection (168). The study investigated endothelial activation of decidual spiral arteries, where decidual basalis is stained additionally with Martius Scarlet Blue (MSB; a histochemical stain). In that study we showed that, similar to positive PAS staining, the MSB stain (red) would appear in the area of the spiral artery wall occupied by foam cell lesions. However, differently to PAS positive staining, MSB would not stain the fibrinoid replacing the smooth muscle in spiral arteries demonstrating remodelling by endovascular extravillous trophoblasts. This subsequent finding led us to the appreciation that PAS is useful for

detecting fibrinoid irrespective of etiology, but that MSB is specifically associated with the deposition of fibrinoid associated with acute atherosclerosis; fibrinoid necrosis.

9.1.6 Developing a systematic method for aiding spiral artery evaluation.

*9.1.6.1 Systematic scoring sheet for AA diagnosis (**Paper I**)*

A systematic approach to sample section evaluation was employed in order to reduce intra- and inter-observer conclusions of AA presence in decidua basalis samples (**Paper I**). We wished to create a systematic evaluation approach to be applied across studies of acute atherosclerosis, and between researchers, given the previous lack of a gold standard by which the defining characteristics of acute atherosclerosis are explained in detail.

Our developed worksheet used to assess sample sections (described in [7.3.3](#), and fully presented in Appendix 1) provided a structured approach for recording the important morphological features of each of the spiral arteries identified, and the surrounding decidual tissue. In the absence of an existing and validated “scoring sheet”, the features of decidual spiral arteries were proposed by our research group, and evaluated in 2013 together with specialist experts in both acute atherosclerosis and uteroplacental pathology (Professor and pathologists Borghild Roald, Oslo, and Professor Fiona Lyall, Glasgow). The scoring sheet aimed to reduce subjectivity of the conclusions of acute atherosclerosis. The utility of the systematic work sheet before concluding on the presence or absence of acute atherosclerosis is reflected in the Kappa value assessing the intra- and inter-observer assessments of AA criteria (“almost perfect” and “substantial”, respectfully) (**Paper I**).

*9.1.6.2 Systematic approach for PVI (perivascular inflammation) quantification (**Paper III**)*

In our study revising the diagnostic criteria for acute atherosclerosis (**Paper I**), the presence of “perivascular” mononuclear leucocytes was defined as categorical, according to the impression of an infiltrate in the decidua in immediate contact with the outer wall of the spiral artery (perivascular). To be defined as a PVI, the apparent density of mononuclear leukocytes around the artery section had to be greater than that in the general distribution of the surrounding decidua. This definition of PVI was based on a subjective impression, and was made for each artery section irrespective of the presence or absence of foam cell lesions. This categorical definition gave no indication the quality of constituent leukocytes, nor any actual quantities. Therefore, in order to characterize type of immune cells involved in these perivascular infiltrates, for **Paper III** we developed a quantitative immunohistochemical approach. The challenge of quantifying lymphocyte populations in relation to decidua basalis spiral arteries is that they do not have a uniform distribution across the decidua basalis. The risk of bias is large if random regions of decidua basalis tissue are assessed when comparing lymphocyte concentrations between samples. Furthermore, selecting random areas of decidua would not allow for comparison of our principle area of interest, namely the relation of lymphocytes populations to spiral artery wall structures and acute atherosclerosis foam cell lesions. The predefined zones for counting decidual lymphocytes, shown in Figure 12, ensured that our counts would enable direct comparisons in leukocyte populations between artery sections (Zone 1 and 2), and between decidual infiltrates (Zone 3). A similar approach was used by Van Dijk et al (2015) in their quantification of lymphocyte populations in and around samples of human abdominal aorta, for their studies of large artery atherosclerosis (169).

A worksheet was also developed and employed for measuring both the surface area for each predefined tissue zone, for each previously evaluated artery section, in each sample, and for the lymphocyte counts they contained (Appendix 2). The inner and outer boundaries of Zone 1 (artery wall) are self-evident. The outer boundary for direct following Zone 2 (decidua basalis in contact with the outer wall of the artery section) was extended to a “perivascular zone” with a width of 100µm, as opposed to the 50µm width in our definition from **Paper I**. We used a wider perivascular zone (100 µm) in **Paper III** in order to yield a greater cell count, but the calculated concentration (cells/mm²) did not alter significantly as compared to using the narrower zone (50 µm, data not shown).

In order to assess the accuracy of our original categorical definition of PVI from **Paper I**, we outlined Zone 2 for the same artery sections, from the same slides, and counted the mononuclear leukocytes. We found that spiral arteries previously categorized as demonstrating PVI were associated with significantly higher concentrations leukocytes in Zone 2 when compared to those arteries where PVI was “absent” (data not shown). This post hoc assessment provides some validity to the categorical definition of PVI employed in **Paper I**.

Our selection of decidual infiltrates (Zone 3, independent of acute atherosclerosis spiral artery lesions) was made on the serial tissue section stained with antibody for CD3, as CD3⁺ cells were found to be the most abundant of the cell types immunostained. The maximal decidual infiltrates chosen for Zone 3 (**Paper III**) were mirrored in the assumed leukocyte populations observed on H&E staining in **Paper I** (serial section slide 3 also stained for CK7 so as to prevent accidental inclusion of EVT in the impression of PVI).

We found agreement of immunostained lymphocyte cell counts in **Paper III**, allowing for a difference of +/-1 cell count between inter and intra-observer observation. We concluded that systematic visual counting of immunostained cells, according to predefined decidua basalis tissue zones, was a reliable method of calculating zone-based decidual lymphocyte concentrations. An alternative would have involved employing digital image analysis to assist in quantification of immunostained cells, an example being the software Image-J, a Java image processing application downloaded free of charge from the website of the National Institutes of Health (NIH, Bethesda, MD) (170).

9.1.7 Kappa scoring and choice of statistical methods

We employed Kappa scoring to evaluate both inter- and intra-observer agreement of morphological findings in **Paper I** and **III** to ensure that our investigations were adequately standardized and reproducible. In Paper I we used Kappa scoring for testing the reproducibility of decidua basalis spiral artery acute atherosclerosis definition. In **Paper III** we employed Kappa scoring for testing the reliability of the quantification of lymphocyte populations within predefined decidua basalis tissue zones of interest. Without this form of evaluation, we could not claim their applicability between investigators and other acute atherosclerosis studies.

Non-parametric testing was employed for all statistical calculations, following convention that considers a p value of <0.05 as statistically significant. Non-parametric testing was chosen given that variables did not always follow a normal distribution. Regression analysis would have been preferable, but of limited merit considering the numbers in each of the groups being compared were low.

9.2 Results discussion

9.2.1 Optimal tissue for uteroplacental acute atherosclerosis diagnosis

Our study demonstrated DSM to be superior to BPS (representing the sample collection method used in routine clinical practice) for successfully identifying decidua basalis spiral arteries (**Paper II**). This result confirmed our previous work of heterogeneous pregnancy groups (87), although that previous paper did not compare DSM and BPS tissue from the same women. We also show in this thesis that, when comparing samples with spiral arteries present, the median number of artery sections was higher in DSM samples compared to BPS (**Paper II**). In addition, we found that, by a small margin, DSM was slightly better than MR in identifying decidual spiral arteries. Though advantageous in being used in clinical practice, the spiral arteries detected in MR samples represent the spiral arteries of the decidua parietalis, not the remodelled arteries of the placental bed supplying the intervillous space (and, indirectly, the fetus).

The disadvantage of the DSM collection method is the lacking topographical orientation of the observed decidual spiral artery. It is challenging to discern whether the decidual spiral artery section observed lies proximal or distal to its myometrial source, or whether it lies central or peripheral in the placental bed. Also, no visual relation to the downstream intervillous space is possible. The presence of trophoblasts do, however, ensure placenta bed location and thereby confirm decidua basalis. The clinically relevant question of *where* (peripheral/central, superficial/deep) is left unanswered for DSM collected decidua basalis tissue samples.

We show for the first time, by direct comparison of BPS and MR decidua tissue collected from the same patients, that DSM has the highest rate of acute atherosclerosis (**Paper II**). This finding might explain the discrepancy in our

identified rate of acute atherosclerosis, compared to, for example, the paper by Kim et al., whose acute atherosclerosis rates are based on BPS and MR collected decidua basalis tissue samples, stained with H&E, and without supplementary IHC (112).

9.2.2 Revised criteria for AA diagnosis and reproducibility

The presence of singular intramural cells demonstrating vacuolation on H&E staining was common in DSM samples containing spiral arteries, for all pregnancy categories evaluated. The intracellular vacuolation in foam cells observed on H&E staining is the empty space previously occupied by phagocytosed lipids that have been removed during the process of deparaffinization, in the preparation of sections cut from paraffin-embedded sample blocks. Intracellular vacuolation may also arise from other intracellular processes/pathologies unrelated to foam cell development.

In the absence of having stained the lipid content of all decidua samples directly using oil red-O (that must be undertaken prior to paraffin embedding), to employ CD68 was chosen as a recognized marker to identify foam cells, thereby confirming the morphological observation of vacuolation as consistent with foam cell formation (91).

One challenge was to distinguish between artefactual and non-specific changes mimicking potential acute atherosclerosis, and genuine acute atherosclerosis. To address this, observed vacuolation alone, without confirmatory CD68 staining, was considered an etiological process other than – and therefore not - foam cell formation.

The presence of CD68+ staining in intramural cells lacking the morphological feature of vacuolation is indicative of an activated phagocyte (for example a

scavenging macrophage), and not specific enough to be considered foam cells. As shown in **Paper I**, Table 3, independent of one another, we observed commonly singular intramural cells demonstrating vacuolation or CD68 positivity. The rate of two adjacent cells demonstrating vacuolation was significantly lower in incidence, and the combination of two adjacent vacuolated cells with confirmatory CD68 positivity even less frequent. The rate of *three* adjacent vacuolated cells was less still, though not significantly less than only two. Because of the dramatic difference in observed rates of vacuolation including solitary cells, compared to observed rates limited to a minimum of two adjacent cells or more, we adopted two adjacent vacuolated cells as the minimum number for suspecting acute atherosclerosis, but only accepted as such when they demonstrated confirmatory CD68 positivity. We used this definition when testing the inter- and intra-observer agreement of acute atherosclerosis definition between all three observers (PAK, BR and FL). The Kappa scores demonstrated a satisfactory reliability (**Paper I**), allowed for this definition of decidua basalis acute atherosclerosis to become our standard in all 3 thesis publications (**Papers I-III**).

9.2.3 AA rate, location, and lesion size and relevance to clinical characteristics

This thesis aimed to determine whether a diagnosis of AA was associated with maternal or pregnancy outcome measures, and, in addition, whether this association varied according to the location of AA: DSM, BPS, MR (alone or in combination).

In **Paper 1**, preeclampsia (as a pregnancy group) biased the association of acute atherosclerosis with both maternal and fetal complications of pregnancy given that these complications are frequent among preeclamptics, irrespective of acute atherosclerosis detection. Interestingly, despite diabetes mellitus being undisputed a risk factor for developing hypertension in pregnancy, as a

pregnancy group, diabetes - unlike preeclampsia - did not associate with AA. When comparing samples with, and without, acute atherosclerosis within their designated pregnancy group (preeclampsia, diabetics with preeclampsia, normotensive controls, normotensive diabetics), the only association with acute atherosclerosis among normotensives was a previous pregnancy affected by PE. Among the preeclamptics, the association of lower gestational age of delivery, birth weight percentile and fetal growth restriction (including pathological fetal Dopplers) was only apparent when the pregnancy group numbers were expanded by the additional participants introduced in **Paper II**. Acute atherosclerosis associates with clinical features that signify dysfunction of the uteroplacental environment. Acute atherosclerosis does not associate with prepregnancy maternal features. Acute atherosclerosis arises during pregnancy, through mechanisms specific to pregnancy, and how prepregnancy maternal features impact the risk of AA, is unknown. The association of acute atherosclerosis to uteroplacental dysfunction relate to its topographical location, namely when it is found in DSM samples, mirrored to some degree in BPS samples (abnormal fetal Dopplers as supposed to FGR), but not fetal MR. We did not find that the detection of acute atherosclerosis in more than one tissue type was associated with clinical pregnancy outcome or complications. This supports the nature of acute atherosclerosis in so much as not being uniform in its expression, in arteries of the same sample, and along the length and circumference of the artery it affects. The exclusion of fetal MR (representing the non-remodelled spiral arteries of the decidua parietalis) from clinical associations suggests that the presence of acute atherosclerosis in decidua basalis is more biologically significant for the pregnancy. This question raises the possibility that acute atherosclerosis may have a different etiology dependent upon its topography. The relevance of the location of acute atherosclerosis detection may be of interest should the artery lesion, in the future, be confirmed to be a marker of increased future atherosclerosis risk.

9.2.4 Acute atherosclerosis, early atherosclerosis and graft vascular disease: common lymphocyte profiles?

Our further study into the quantification of decidual lymphocyte populations associated with spiral arteries (**Paper III**) sought to establish whether presence of acute atherosclerosis was reflected in the quantities of decidual lymphocyte populations integral to (Zone 1), and surrounding spiral artery sections (Zone 2 and Zone 3). An association between lymphocyte populations and presence of acute atherosclerosis (intramural foam cell lesions) might provide an immune cell “signature” that could lead provide clues as to the mechanisms leading to foam cell development, or alternatively the immune response to their presence.

The morphological appearance of acute atherosclerosis has been likened to both early stage atherosclerosis, as well as the vasculopathy seen in transplant graft rejection (96). It is reasonable to speculate on the possibility of them sharing a common immunological activity. Presence of Tregs has been associated with better outcomes for graft recipients (171). Sasaki has proposed that with the lack of Tregs comes a loss of the inhibitory influence on the propagation of proatherogenic immune activity (124). The number of Tregs in human atherosclerosis has been reported as low compared to their relatively high numbers found in relation to inflammatory skin lesions (172). This is a finding in kind with the universal paucity of FOXP3+ cells we found in association with decidual spiral arteries, irrespective of pregnancy category or the presence/absence of acute atherosclerosis (**Paper III**).

Our third trimester findings are at odds with investigations of Treg numbers in the first trimester of viable pregnancies, where decidual Tregs are more abundant (173). This would suggest that, even in normal pregnancies, there is a transition from a state with high numbers of decidual Tregs (1st and 2nd trimester), to a state close to delivery where Tregs are lower in numbers (174). Perhaps the reduction in decidual Treg numbers reduces their inhibitory effect on proatherogenic immune activity, akin to what has been observed in the

organ dysfunction resulting from the graft vessel disease of transplant rejection. An animal model developed by our collaborators, mimicking preeclampsia, has demonstrated the potentially therapeutic effect of Treg expansion in reversing the growth restriction associated with this particular pregnancy phenotype (175). The therapeutic effects of Tregs have also been discussed in the setting of disease progression in atherosclerosis (176).

In our studies comparing concentrations of decidual lymphocytes between preeclamptics and normotensive pregnancies (**Paper III**), it is important to note that although concentrations were comparable between PE samples with AA and normotensives without AA, the impact of the lower gestational age of the PE group at delivery as compared to controls may have impacted these findings. As elective cesareans in normotensive pregnancies are rarely performed before gestational week 39, due to the risk of fetal complications (such as transient tachypnoea of the newborn, known in Norway as “wet lungs”) with delivery at earlier gestations, stringent gestational age matching to PE pregnancies is a challenge for all studies in this field, and not easily solved. Our decidual findings were also in kind with the findings of higher CD3, CD8 and CD4 concentrations are described by van Dijk et al in early stages of disease progression for human aortic atherosclerosis. Van Dijk’s findings (169) of cadaver specimens of abdominal aorta cannot however be directly compared to ours examining much smaller arteries; the spiral arteries.

Interesting as these similarities are, we are a long way from being able to confirm common mechanistic origins for acute atherosclerosis, atherosclerosis and graft rejection. In the recent paper from our group (168) we were unable to demonstrate endothelial activation in spiral arteries containing acute atherosclerosis (intramural foam cells). This is at odds with endothelial activation being accepted as the initiating event, preceding migration of circulating

macrophages, in the development of atherosclerotic disease in large arteries (177). Nonetheless, it is encouraging to be able to demonstrate that the decidual tissue of women, most likely to suffer atherosclerotic morbidity in the future, share immunological proportions similar to those found in specimens of atherosclerotic human aorta. In addition, our studies have found that the natural allograft status of the pregnant condition demonstrates a loss of tolerance-inducing Tregs close to delivery, which is compelling to compare with the finding of loss of Treg inducing tolerance heralding graft rejection (rising rejection activity index (178)).

9.2.5 Hypercoagulable spiral arteries in AA or postdelivery effects?

In **Paper I**, investigation of spiral artery thrombosis was added to our systematic evaluation of decidual spiral artery sections, as it has been pointed out by previous investigators that acute atherosclerosis was associated with downwards placental infarcts (89, 90, 103). Our morphological findings support both previous reports of increased thrombosis in acute atherosclerosis, and indirectly provide a clue as to one mechanism by which abnormal umbilical artery Doppler findings (indicative of dysfunctional placental function secondary to uteroplacental circulatory dysfunction) were associated with preeclampsia with acute atherosclerosis in the decidua basalis tissue.

Our definition of spiral artery thrombosis in **Paper I** was based on the presence of red blood cells incorporated into a meshwork of fibrin strands, suggestive of thrombosis formation in vivo. However, it is difficult to be certain that some of these thromboses were not triggered by the manipulation during cesarean section and under the extraction of the placenta. However, the contribution of this artifact would affect all samples collected, therefore the *difference* in rates is what we wish to highlight in this result. Pregnancy is a hypercoagulable state, especially among those affected by preeclampsia,

irrespective of acute atherosclerosis, which could add to the finding of high thrombosis rates. We plan future studies to further investigate the hypercoagulability of women with PE, with or without AA, in comparison with normotensive pregnancies (179).

9.2.6 Absence of trophoblasts in presence of decidua basalis AA (DSM)

Like all histological evaluations, our evaluations are taken from a snap shot of a myriad of time dependent processes ongoing within a single localization of the placental bed. Like Stallmach et al. (106), we too found an association between presence of acute atherosclerosis and the absence of endovascular trophoblasts. We could also demonstrate that acute atherosclerosis (foam cell lesions) could co-exist in the walls of spiral arteries in which endovascular trophoblasts were also identified (**Paper I**, also illustrated in our later publication (168). However, in these particular cases, there were fewer endovascular trophoblasts present compared to spiral arteries where acute atherosclerosis was absent.

Without decidual specimens taken from the same patient at an earlier juncture during the pregnancy (not ethically compatible with an ongoing pregnancy), it is difficult to conclude whether the absence of endovascular trophoblasts predisposes to acute atherosclerosis development, or whether the development of acute atherosclerosis results in the destruction / disappearance of endovascular trophoblasts. These two speculations would support two opposing theories as to the development of acute atherosclerosis. The first would suggest that dysfunctional remodelling early on in the pregnancy predisposes to the later development of acute atherosclerosis. The other would suggest that acute atherosclerosis can occur in normally remodeled spiral arteries due to later changes in the fetal-maternal interface, subsequent to spiral artery remodelling. There is no reason why both speculations could be true, as suggested by the model by Staff

& Redman (2014) (133). Both pathways are compatible with the concepts of a dysfunctional maternal tolerance to fetal antigen.

10. CONCLUSIONS

This PhD study provided several answers to the acute atherosclerosis questions posed in this PhD study, although the lesion still remains one of the mysteries of Obstetric Medicine. The thesis provides following answers:

A. “*What*” are the useful criteria for defining acute atherosclerosis?

Presence of an intramural foam cell lesion, comprising of two vacuolated cells with confirmatory CD68 positivity, demonstrated itself as a reproducible minimum requirement in the identification of acute atherosclerosis in the spiral arteries of decidua basalis tissue. Fibrinoid (identified using PAS) was ubiquitous in the spiral arteries of the decidua basalis making the classical “fibrinoid necrosis” difficult to discern from the fibrinoid associated with spiral artery remodelling in pregnancy. PAS identified fibrinoid was never absent in arteries containing foam cell lesions. Collections of mononuclear perivascular infiltrates, in the absence of targeted immunohistochemical staining, were not exclusive to spiral arteries containing foam cell lesions. **(Paper I).**

B. “*Where*” in the uteroplacental unit is acute atherosclerosis most readily detected?

Acute atherosclerosis is most frequently detected in decidua basalis tissue, using the Decidual vacuum Suction Method applied to the placental bed of the uterine wall. Spiral arteries and acute atherosclerosis were less frequently detected in the remains of the placental bed attached to the Basal Plate Surface of the delivered placenta. Acute atherosclerosis is also detectable in the spiral arteries of decidua parietalis proximal to the placenta. **(Paper II).**

C. “*Why*” are the acute atherosclerosis foam cells present in the arterial wall? Higher concentrations of decidual T-lymphocytes (specifically assumed CD4+ T-cells) associated with both sample sections and individual spiral arteries where foam cell lesions are identified. Whether a cause or a result, presence of acute atherosclerosis in the decidua basalis coincides with a quantifiably increased presence of T-cells cells associated with the acquired immune response (**Paper III**).

D. “*How*” does the presence of decidual (uteroplacental) acute atherosclerosis associate with preeclampsia, both clinically and immunologically? Our revised definition of acute atherosclerosis shows significant association with preeclamptic pregnancies, in keeping with previous studies, but not with diabetes. The association of disease severity and acute atherosclerosis relates to clinical parameters reflecting failing placental function (lower gestational age of delivery and features of fetal growth restriction) as supposed to preconceptional maternal parameters, irrespective of the topographical location (decidua basalis and parietalis). This suggests that, rather than pregnancy just being a platform upon which prepregnancy maternal risk factors culminate in acute atherosclerosis development, that processes integral to pregnancy itself are driving the development of this enigmatic arterial lesion. (**Papers I, II**).

11. FURTHER STUDIES

As acute atherosclerosis lesions still remain one of the mysteries of Obstetric Medicine, our plan is to pursue our research with the following studies:

1. Downstream (placental) consequence of acute atherosclerosis (AA) presence in the decidua basalis: Morphological features of placental dysfunction according to Redline (180, 181), systematic evaluation by placenta pathologists of placental samples belonging to patients already evaluated for AA.
2. A follow-up of our recently published genetic fetal and maternal risk combination in preeclampsia with AA (182): Maternal KIR genotypes in combination with fetal HLA C and HLA G alleles, and, presence of acute atherosclerosis in decidual spiral arteries.
3. Assessing the hypercoagulability of women with PE, with or without AA; in comparison with normotensive pregnancies (179).

12. ERRATUM

13. OTHER PUBLICATIONS CO-AUTHORED BY CANDIDATE DURING PhD STUDIES

1. Yung HW, **Alnaes-Katjavivi P**, Jones CJ, El-Bacha T, Golic M, Staff AC, Burton GJ. Placental endoplasmic reticulum stress in gestational diabetes: the potential for therapeutic intervention with chemical chaperones and antioxidants. *Diabetologia*. 2016 Oct;59(10):2240-50.
2. Golic M, Haase N, Herse F, Wehner A, Vercruysse L, Pijnenborg R, Balogh A, Saether PC, Dissen E, Luft FC, Przybyl L, Park JK, **Alnaes-Katjavivi P**, Staff AC, Verlohren S, Henrich W, Muller DN, Dechend R. Natural Killer Cell Reduction and Uteroplacental Vasculopathy. *Hypertension*. 2016 Oct;68(4):964-73.
3. Johnsen GM, Størvold GL, Drabbels JJM, Haasnoot GW, Eikmans M, Spruyt-Gerritse MJ, **Alnaes-Katjavivi P**, Scherjon SA, Redman CWG, Claas FHJ, Staff AC. The combination of maternal KIR-B and fetal HLA-C2 is associated with decidua basalis acute atherosclerosis in pregnancies with preeclampsia. *J Reprod Immunol*. 2018 Jul 25;129:23-9.
4. Rieber-Mohn AB, Sugulle M, Wallukat G, **Alnaes-Katjavivi P**, Leite Størvold G, Bolstad N, Redman CW, Dechend R, Staff AC. Auto-antibodies against the angiotensin II type I receptor in women with uteroplacental acute atherosclerosis and preeclampsia at delivery and several years postpartum. *J Reprod Immunol*. 2018 Aug;128:23-9.
5. Moe K, **Alnaes-Katjavivi P**, Størvold GL, Sugulle M, Johnsen GM, Redman CWG, Dechend R, Staff AC. Classical Cardiovascular Risk Markers in Pregnancy and Associations to Uteroplacental Acute Atherosclerosis. *Hypertension*. 2018 Sep;72(3):695-702.
6. Fosheim IK, **Alnaes-Katjavivi P**, Redman C, Roald B, Staff AC, Størvold GL. Acute atherosclerosis of decidua basalis; characterization of spiral arteries, endothelial status and activation. *Placenta*. 2019 Jul;82:10-16.

14. REFERENCE LIST

1. Robertson WB, Brosens I, Dixon G. Maternal uterine vascular lesions in the hypertensive complications of pregnancy. *Perspect Nephrol Hypertens*. 1976;5:115-27.
2. Ramsey EM DM. Placental vasculature and circulation. *Anatomy, physiology, radiology, clinical aspects atlas textbook*. Stuttgart: Georg Thieme Publishers; 1980. p. p.20, p.13.
3. E.C.B. Hall-Craggs ibDA. *Anatomy as a Basis for Clinical Medicine*. Second ed. Baltimore, Mayland, USA: Urban&Schwartzberg, Inc.; 1985 1990. 330-332
4. Placental Bed Disorders - Basic Science and its Translation to Obstetrics. 1st ed. Cambridge: Cambridge University Press; 2010. 43
5. Lyall F. The human placental bed revisited. *Placenta*. 2002;23(8-9):555-62.
6. Brosens I, Benagiano M, Puttemans P, D'Elis MM, Benagiano G. The placental bed vascular pathology revisited: a risk indicator for cardiovascular disease. *J Matern Fetal Neonatal Med*. 2017;1-9.
7. Ueland K, Akamatsu TJ, Eng M, Bonica JJ, Hansen JM. Maternal cardiovascular dynamics. VI. Cesarean section under epidural anesthesia without epinephrine. *Am J Obstet Gynecol*. 1972;114(6):775-80.
8. Williams Obstetrics. 22nd ed: McGraw-Hill (Medical Publishing Division); 2005. p.403
9. Brosens I, Robertson WB, Dixon HG. The physiological response of the vessels of the placental bed to normal pregnancy. *J Pathol Bacteriol*. 1967;93(2):569-79.
10. Pijnenborg R, Vercruysse L, Hanssens M. The uterine spiral arteries in human pregnancy: facts and controversies. *Placenta*. 2006;27(9-10):939-58.
11. Mandala M, Osol G. Physiological remodelling of the maternal uterine circulation during pregnancy. *Basic Clin Pharmacol Toxicol*. 2012;110(1):12-8.
12. Dimitriadis E, Nie G, Hannan NJ, Paiva P, Salamonsen LA. Local regulation of implantation at the human fetal-maternal interface. *Int J Dev Biol*. 2010;54(2-3):313-22.
13. James JL, Carter AM, Chamley LW. Human placentation from nidation to 5 weeks of gestation. Part I: What do we know about formative placental development following implantation? *Placenta*. 2012;33(5):327-34.
14. Bulmer JN, Longfellow M, Ritson A. Leukocytes and resident blood cells in endometrium. *Ann N Y Acad Sci*. 1991;622:57-68.
15. Hamperl H, Hellweg G. Granular endometrial stroma cells. *Obstet Gynecol*. 1958;11(4):379-87.
16. Von Numers C. On the specific granular cells (globular leukocytes) of the human endometrium, with special reference to their occurrence in different pathological conditions and to their staining reactions. *Acta Pathol Microbiol Scand*. 1953;33(3):250-6.
17. Kam EP, Gardner L, Loke YW, King A. The role of trophoblast in the physiological change in decidual spiral arteries. *Hum Reprod*. 1999;14(8):2131-8.
18. Harris LK. IFPA Gabor Than Award lecture: Transformation of the spiral arteries in human pregnancy: key events in the remodelling timeline. *Placenta*. 2011;32 Suppl 2:S154-8.
19. Benirschke K, Kaufmann P. *Pathology of the Human Placenta*. 3rd ed. New York, USA: Springer-Verlag; 1995.
20. Craven CM, Morgan T, Ward K. Decidual spiral artery remodelling begins before cellular interaction with cytotrophoblasts. *Placenta*. 1998;19(4):241-52.
21. Palmer SK, Zamudio S, Coffin C, Parker S, Stamm E, Moore LG. Quantitative estimation of human uterine artery blood flow and pelvic blood flow redistribution in pregnancy. *Obstet Gynecol*. 1992;80(6):1000-6.

22. Acacio GL. Uterine artery Doppler patterns in abdominal pregnancy. *Ultrasound Obstet Gynecol.* 2002;20(2):194-6.
23. Pijnenborg R, Dixon G, Robertson WB, Brosens I. Trophoblastic invasion of human decidua from 8 to 18 weeks of pregnancy. *Placenta.* 1980;1(1):3-19.
24. Kaufmann P, Black S, Huppertz B. Endovascular trophoblast invasion: implications for the pathogenesis of intrauterine growth retardation and preeclampsia. *Biol Reprod.* 2003;69(1):1-7.
25. Burton GJ, Jauniaux E. Pathophysiology of placental-derived fetal growth restriction. *Am J Obstet Gynecol.* 2018;218(2S):S745-S61.
26. Pijnenborg R, Bland JM, Robertson WB, Dixon G, Brosens I. The pattern of interstitial trophoblastic invasion of the myometrium in early human pregnancy. *Placenta.* 1981;2(4):303-16.
27. Jauniaux E, Watson AL, Hempstock J, Bao YP, Skepper JN, Burton GJ. Onset of maternal arterial blood flow and placental oxidative stress. A possible factor in human early pregnancy failure. *Am J Pathol.* 2000;157(6):2111-22.
28. Genbacev O, Zhou Y, Ludlow JW, Fisher SJ. Regulation of human placental development by oxygen tension. *Science.* 1997;277(5332):1669-72.
29. Lyall F, Robson SC, Bulmer JN. Spiral artery remodeling and trophoblast invasion in preeclampsia and fetal growth restriction: relationship to clinical outcome. *Hypertension.* 2013;62(6):1046-54.
30. Kaufmann P, Black S, Huppertz B. Endovascular trophoblast invasion: implications for the pathogenesis of intrauterine growth retardation and preeclampsia. *Biol Reprod.* 2003;69(1):1-7.
31. Burton GJ, Woods AW, Jauniaux E, Kingdom JC. Rheological and physiological consequences of conversion of the maternal spiral arteries for uteroplacental blood flow during human pregnancy. *Placenta.* 2009;30(6):473-82.
32. Khong TY, Adema ED, Erwich JJ. On an anatomical basis for the increase in birth weight in second and subsequent born children. *Placenta.* 2003;24(4):348-53.
33. Duley L. Pre-eclampsia and the hypertensive disorders of pregnancy. *Br Med Bull.* 2003;67:161-76.
34. Roberts JM, Pearson G, Cutler J, Lindheimer M. Summary of the NHLBI Working Group on Research on Hypertension During Pregnancy. *Hypertension.* 2003;41(3):437-45.
35. Brown MA, Magee LA, Kenny LC, Karumanchi SA, McCarthy FP, Saito S, Hall DR, Warren CE, Adoyi G, et al. Hypertensive Disorders of Pregnancy: ISSHP Classification, Diagnosis, and Management Recommendations for International Practice. *Hypertension.* 2018;72(1):24-43.
36. Hypertension in pregnancy. Report of the American College of Obstetricians and Gynecologists' Task Force on Hypertension in Pregnancy. *Obstet Gynecol.* 2013;122(5):1122-31.
37. Duley L. Maternal mortality associated with hypertensive disorders of pregnancy in Africa, Asia, Latin America and the Caribbean. *Br J Obstet Gynaecol.* 1992;99(7):547-53.
38. Duley L. The global impact of pre-eclampsia and eclampsia. *Semin Perinatol.* 2009;33(3):130-7.
39. Vangen S, Ellingsen L, Andersgaard AB, Jacobsen AF, Lorentzen B, Nyflot LT, Rygh AB, Skulstad SM, Tappert C, et al. Maternal deaths in Norway 2005-2009. *Tidsskr Nor Laegeforen.* 2014;134(8):836-9.
40. Ananth CV, Wilcox AJ. Placental abruption and perinatal mortality in the United States. *Am J Epidemiol.* 2001;153(4):332-7.
41. Backes CH, Markham K, Moorehead P, Cordero L, Nankervis CA, Giannone PJ. Maternal preeclampsia and neonatal outcomes. *J Pregnancy.* 2011;2011:214365.

42. Resnik R. Intrauterine growth restriction. *Obstet Gynecol.* 2002;99(3):490-6.
43. Ananth CV, Smulian JC, Vintzileos AM. Ischemic placental disease: maternal versus fetal clinical presentations by gestational age. *J Matern Fetal Neonatal Med.* 2010;23(8):887-93.
44. Ananth CV, Basso O. Impact of pregnancy-induced hypertension on stillbirth and neonatal mortality. *Epidemiology.* 2010;21(1):118-23.
45. Harmon QE, Huang L, Umbach DM, Klungsoyr K, Engel SM, Magnus P, Skjaerven R, Zhang J, Wilcox AJ. Risk of fetal death with preeclampsia. *Obstet Gynecol.* 2015;125(3):628-35.
46. van Esch JJA, van Heijst AF, de Haan AFJ, van der Heijden OWH. Early-onset preeclampsia is associated with perinatal mortality and severe neonatal morbidity. *J Matern Fetal Neonatal Med.* 2017;30(23):2789-94.
47. Brosens IA, Robertson WB, Dixon HG. The role of the spiral arteries in the pathogenesis of preeclampsia. *Obstet Gynecol Annu.* 1972;1:177-91.
48. Pijnenborg R, Anthony J, Davey DA, Rees A, Tiltman A, Vercruysse L, van AA. Placental bed spiral arteries in the hypertensive disorders of pregnancy. *Br J Obstet Gynaecol.* 1991;98(7):648-55.
49. Brosens IA. The Utero-Placental Vessels at Term — the Distribution and Extent of Physiological Changes. In: Kaufmann P, Miller RK, editors. *Placental Vascularization and Blood Flow: Basic Research and Clinical Applications.* Boston, MA: Springer US; 1988. p. 61-7.
50. Khong TY, De WF, Robertson WB, Brosens I. Inadequate maternal vascular response to placentation in pregnancies complicated by pre-eclampsia and by small-for-gestational age infants. *Br J Obstet Gynaecol.* 1986;93(10):1049-59.
51. Gerretsen G, Huisjes HJ, Elema JD. Morphological changes of the spiral arteries in the placental bed in relation to pre-eclampsia and fetal growth retardation. *Br J Obstet Gynaecol.* 1981;88(9):876-81.
52. De Wolf F, De Wolf-Peeters C, Brosens I, Robertson WB. The human placental bed: electron microscopic study of trophoblastic invasion of spiral arteries. *Am J Obstet Gynecol.* 1980;137(1):58-70.
53. Meekins JW, Pijnenborg R, Hanssens M, McFadyen IR, van AA. A study of placental bed spiral arteries and trophoblast invasion in normal and severe pre-eclamptic pregnancies. *Br J Obstet Gynaecol.* 1994;101(8):669-74.
54. King A, Burrows TD, Hiby SE, Bowen JM, Joseph S, Verma S, Lim PB, Gardner L, Le Bouteiller P, et al. Surface expression of HLA-C antigen by human extravillous trophoblast. *Placenta.* 2000;21(4):376-87.
55. Apps R, Murphy SP, Fernando R, Gardner L, Ahad T, Moffett A. Human leucocyte antigen (HLA) expression of primary trophoblast cells and placental cell lines, determined using single antigen beads to characterize allotype specificities of anti-HLA antibodies. *Immunology.* 2009;127(1):26-39.
56. Hilton HG, Parham P. Missing or altered self: human NK cell receptors that recognize HLA-C. *Immunogenetics.* 2017;69(8-9):567-79.
57. Hiby SE, Walker JJ, O'shaughnessy KM, Redman CW, Carrington M, Trowsdale J, Moffett A. Combinations of maternal KIR and fetal HLA-C genes influence the risk of preeclampsia and reproductive success. *J Exp Med.* 2004;200(8):957-65.
58. Moffett A, Hiby SE. How Does the maternal immune system contribute to the development of pre-eclampsia? *Placenta.* 2007;28 Suppl A:S51-S6.
59. Hiby SE, Apps R, Sharkey AM, Farrell LE, Gardner L, Mulder A, Claas FH, Walker JJ, Redman CW, et al. Maternal activating KIRs protect against human reproductive failure mediated by fetal HLA-C2. *J Clin Invest.* 2010;120(11):4102-10.

60. Hiby SE, Apps R, Chazara O, Farrell LE, Magnus P, Trogstad L, Gjessing HK, Carrington M, Moffett A. Maternal KIR in Combination with Paternal HLA-C2 Regulate Human Birth Weight. *J Immunol*. 2014;192(11):5069-73.
61. Staff AC, Halvorsen B, Ranheim T, Henriksen T. Elevated level of free 8-iso-prostaglandin F2alpha in the decidua basalis of women with preeclampsia. *Am J Obstet Gynecol*. 1999;181(5 Pt 1):1211-5.
62. Staff AC, Ranheim T, Khoury J, Henriksen T. Increased contents of phospholipids, cholesterol, and lipid peroxides in decidua basalis in women with preeclampsia. *Am J Obstet Gynecol*. 1999;180(3 Pt 1):587-92.
63. Hung TH, Skepper JN, Burton GJ. In vitro ischemia-reperfusion injury in term human placenta as a model for oxidative stress in pathological pregnancies. *Am J Pathol*. 2001;159(3):1031-43.
64. Jauniaux E, Poston L, Burton GJ. Placental-related diseases of pregnancy: Involvement of oxidative stress and implications in human evolution. *Hum Reprod Update*. 2006;12(6):747-55.
65. Redman CW. Current topic: pre-eclampsia and the placenta. *Placenta*. 1991;12(4):301-8.
66. Redman CW, Sargent IL. Immunology of Pre-Eclampsia. *Am J Reprod Immunol*. 2010;63:534-43.
67. Redman CW, Sargent IL, Staff AC. IFPA Senior Award Lecture: Making sense of pre-eclampsia - Two placental causes of preeclampsia? *Placenta*. 2014;35 Suppl:S20-S5.
68. Staff AC, Redman CWG. The Differences Between Early- and Late-Onset Preeclampsia. In: Saito S, editor. *Preeclampsia Basic, Genomic, and Clinical. Comprehensive Gynecology and Obstetrics*. 1 ed. Singapore: Springer; 2018. p. 157-72.
69. Mosca L, Benjamin EJ, Berra K, Bezanson JL, Dolor RJ, Lloyd-Jones DM, Newby LK, Pina IL, Roger VL, et al. Effectiveness-based guidelines for the prevention of cardiovascular disease in women--2011 update: a guideline from the American Heart Association. *J Am Coll Cardiol*. 2011;57(12):1404-23.
70. Strydom HC, Chandler AB, Glagov S, Guyton JR, Insull W, Jr., Rosenfeld ME, Schaffer SA, Schwartz CJ, Wagner WD, et al. A definition of initial, fatty streak, and intermediate lesions of atherosclerosis. A report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association. *Circulation*. 1994;89(5):2462-78.
71. Weintraub WS, Daniels SR, Burke LE, Franklin BA, Goff DC, Jr., Hayman LL, Lloyd-Jones D, Pandey DK, Sanchez EJ, et al. Value of primordial and primary prevention for cardiovascular disease: a policy statement from the American Heart Association. *Circulation*. 2011;124(8):967-90.
72. Bellamy L, Casas JP, Hingorani AD, Williams DJ. Pre-eclampsia and risk of cardiovascular disease and cancer in later life: systematic review and meta-analysis. *BMJ*. 2007;335(7627):974.
73. Staff AC, Redman CW, Williams D, Leeson P, Moe K, Thilaganathan B, Magnus P, Steegers EA, Tsigas EZ, et al. Pregnancy and Long-Term Maternal Cardiovascular Health: Progress Through Harmonization of Research Cohorts and Biobanks. *Hypertension*. 2016;67(2):251-60.
74. Lykke JA, Langhoff-Roos J, Sibai BM, Funai EF, Triche EW, Paidas MJ. Hypertensive pregnancy disorders and subsequent cardiovascular morbidity and type 2 diabetes mellitus in the mother. *Hypertension*. 2009;53(6):944-51.

75. Kajantie E, Eriksson JG, Osmond C, Thornburg K, Barker DJ. Pre-eclampsia is associated with increased risk of stroke in the adult offspring: the Helsinki birth cohort study. *Stroke*. 2009;40(4):1176-80.
76. Kajantie E, Osmond C, Eriksson JG. Gestational hypertension is associated with increased risk of type 2 diabetes in adult offspring: the Helsinki Birth Cohort Study. *Am J Obstet Gynecol*. 2017;216(3):281 e1- e7.
77. Kajantie E, Osmond C, Barker DJ, Forsen T, Phillips DI, Eriksson JG. Size at birth as a predictor of mortality in adulthood: a follow-up of 350 000 person-years. *Int J Epidemiol*. 2005;34(3):655-63.
78. Cao SS, Kaufman RJ. Targeting endoplasmic reticulum stress in metabolic disease. *Expert Opin Ther Targets*. 2013;17(4):437-48.
79. Yung HW, Alnaes-Katjavivi P, Jones CJ, El-Bacha T, Golic M, Staff AC, Burton GJ. Placental endoplasmic reticulum stress in gestational diabetes: the potential for therapeutic intervention with chemical chaperones and antioxidants. *Diabetologia*. 2016;59(10):2240-50.
80. Feig DS, Zinman B, Wang X, Hux JE. Risk of development of diabetes mellitus after diagnosis of gestational diabetes. *CMAJ*. 2008;179(3):229-34.
81. Melchiorre K, Sutherland GR, Liberati M, Thilaganathan B. Preeclampsia is associated with persistent postpartum cardiovascular impairment. *Hypertension*. 2011;58(4):709-15.
82. Hertig AT. Vascular pathology in hypertensive albuminuric toxemias of pregnancy. *Clinics*. 1945;4:602-14.
83. Zeek PM, Assali NS. Vascular changes in the decidua associated with eclamptogenic toxemia of pregnancy. *Am J Clin Pathol*. 1950;20(12):1099-109.
84. Reyman TA, Barraco RA, Cockburn A. Histopathological examination of an egyptian mummy. *Bull N Y Acad Med*. 1976;52(4):506-16.
85. Sandison AT. Degenerative vascular disease in the Egyptian mummy. *Med Hist*. 1962;6(1):77-81.
86. Sheppard BL, Bonnar J. An ultrastructural study of utero-placental spiral arteries in hypertensive and normotensive pregnancy and fetal growth retardation. *Br J Obstet Gynaecol*. 1981;88(7):695-705.
87. Harsem NK, Staff AC, He L, Roald B. The decidual suction method: a new way of collecting decidual tissue for functional and morphological studies. *Acta Obstet Gynecol Scand*. 2004;83(8):724-30.
88. Pijnenborg R, Ball E, Bulmer JN, Hanssens M, Robson SC, Vercruysse L. In vivo analysis of trophoblast cell invasion in the human. *Methods Mol Med*. 2006;122:11-44.
89. Stevens DU, Al-Nasiry S, Bulten J, Spaanderman ME. Decidual vasculopathy and adverse perinatal outcome in preeclamptic pregnancy. *Placenta*. 2012;33(8):630-3.
90. Robertson WB, Brosens I, Dixon HG. The pathological response of the vessels of the placental bed to hypertensive pregnancy. *J Pathol Bacteriol*. 1967;93:581-92.
91. Katabuchi H, Yih S, Ohba T, Matsui K, Takahashi K, Takeya M, Okamura H. Characterization of macrophages in the decidual atherotic spiral artery with special reference to the cytology of foam cells. *Med Electron Microsc*. 2003;36(4):253-62.
92. Fox H. Fibrinoid necrosis of placental villi. *J Obstet Gynaecol Br Commonw*. 1968;75(4):448-52.
93. Movat HZ. The concept of fibrinoid. *Am J Med Sci*. 1958;236(3):373-82.
94. Bajema IM, Bruijn JA. What stuff is this! A historical perspective on fibrinoid necrosis. *J Pathol*. 2000;191(3):235-8.
95. Khong TY. Acute atherosclerosis in pregnancies complicated by hypertension, small-for-gestational-age infants, and diabetes mellitus. *Arch Pathol Lab Med*. 1991;115(7):722-5.

96. Labarrere CA. Acute atherosclerosis. A histopathological hallmark of immune aggression? *Placenta*. 1988;9(1):95-108.
97. Stevens DU, Al-Nasiry S, Bulten J, Spaanderman ME. Decidual vasculopathy in preeclampsia: lesion characteristics relate to disease severity and perinatal outcome. *Placenta*. 2013;34(9):805-9.
98. Hecht JL, Zsengeller ZK, Spiel M, Karumanchi SA, Rosen S. Revisiting decidual vasculopathy. *Placenta*. 2016;42:37-43.
99. Zhang P. Decidual Vasculopathy in Preeclampsia and Spiral Artery Remodeling Revisited: Shallow Invasion versus Failure of Involution. *AJP reports*. 2018;8(4):e241-e6.
100. Hertig A, Liere P. New markers in preeclampsia. *Clin Chim Acta*. 2010;411(21-22):1591-5.
101. Meekins JW, Pijnenborg R, Hanssens M, van AA, McFadyen IR. Immunohistochemical detection of lipoprotein(a) in the wall of placental bed spiral arteries in normal and severe preeclamptic pregnancies. *Placenta*. 1994;15(5):511-24.
102. De Wolf F, Robertson WB, Brosens I. The ultrastructure of acute atherosclerosis in hypertensive pregnancy. *Am J Obstet Gynecol*. 1975;123(2):164-74.
103. Maqueo M, Chavezazuela J, Dosaldelavega M. Placental pathology in eclampsia and preeclampsia. *Obstet Gynecol*. 1964;24:350-6.
104. Robertson WB, Brosens I, Dixon G. Uteroplacental vascular pathology. *Eur J Obstet Gynecol Reprod Biol*. 1975;5(1-2):47-65.
105. Althabe O, Labarrere C, Telenta M. Maternal vascular lesions in placentae of small-for-gestational-age infants. *Placenta*. 1985;6(3):265-76.
106. Stallmach T, Hebisch G, Orban P, Lu X. Aberrant positioning of trophoblast and lymphocytes in the fetal-maternal interface with pre-eclampsia. *Virchows Arch*. 1999;434(3):207-11.
107. Baltajian K, Hecht JL, Wenger JB, Salahuddin S, Verlohren S, Perschel FH, Zsengeller ZK, Thadhani R, Karumanchi SA, et al. Placental lesions of vascular insufficiency are associated with anti-angiogenic state in women with preeclampsia. *Hypertens Pregnancy*. 2014;33(4):427-39.
108. Bulmer JN, Williams PJ, Lash GE. Immune cells in the placental bed. *Int J Dev Biol*. 2010;54(2-3):281-94.
109. Williams PJ, Searle RF, Robson SC, Innes BA, Bulmer JN. Decidual leucocyte populations in early to late gestation normal human pregnancy. *J Reprod Immunol*. 2009;82(1):24-31.
110. Williams PJ, Bulmer JN, Searle RF, Innes BA, Robson SC. Altered decidual leucocyte populations in the placental bed in pre-eclampsia and foetal growth restriction: a comparison with late normal pregnancy. *Reproduction*. 2009;138(1):177-84.
111. Harsem NK, Roald B, Braekke K, Staff AC. Acute atherosclerosis in decidual tissue: not associated with systemic oxidative stress in preeclampsia. *Placenta*. 2007;28(8-9):958-64.
112. Kim YM, Chaemsathong P, Romero R, Shaman M, Kim CJ, Kim JS, Qureshi F, Jacques SM, Ahmed AI, et al. The frequency of acute atherosclerosis in normal pregnancy and preterm labor, preeclampsia, small-for-gestational age, fetal death and midtrimester spontaneous abortion. *J Matern Fetal Neonatal Med*. 2014:1-9.
113. Zhang P, Schmidt M, Cook L. Maternal vasculopathy and histologic diagnosis of preeclampsia: poor correlation of histologic changes and clinical manifestation. *Am J Obstet Gynecol*. 2006;194(4):1050-6.
114. Hustin J, Foidart JM, Lambotte R. Maternal vascular lesions in pre-eclampsia and intrauterine growth retardation: light microscopy and immunofluorescence. *Placenta*. 1983;4 Spec No:489-98.

115. Sheppard BL, Bonnar J. The ultrastructure of the arterial supply of the human placenta in pregnancy complicated by fetal growth retardation. *Br J Obstet Gynaecol.* 1976;83(12):948-59.
116. Khong TY. Acute atherosclerosis in pregnancies complicated by hypertension, small-for-gestational-age infants, and diabetes mellitus. *Arch Pathol Lab Med.* 1991;115(7):722-5.
117. De WF, Brosens I, Renaer M. Fetal growth retardation and the maternal arterial supply of the human placenta in the absence of sustained hypertension. *Br J Obstet Gynaecol.* 1980;87(8):678-85.
118. Nayar R, Lage JM. Placental changes in a first trimester missed abortion in maternal systemic lupus erythematosus with antiphospholipid syndrome; a case report and review of the literature. *Hum Pathol.* 1996;27(2):201-6.
119. Hayati AR, Azizah A, Wahidah A. Incidence of acute atherosclerosis in complete molar pregnancy. *Malays J Pathol.* 1998;20(2):113-4.
120. Staff AC, Johnsen GM, Dechend R, Redman CW. Preeclampsia and uteroplacental acute atherosclerosis: immune and inflammatory factors. *J Reprod Immunol.* 2014;101-102:120-6.
121. Libby P. Inflammation in atherosclerosis. *Arterioscler Thromb Vasc Biol.* 2012;32(9):2045-51.
122. Esteve E, Ricart W, Fernandez-Real JM. Dyslipidemia and inflammation: an evolutionary conserved mechanism. *Clin Nutr.* 2005;24(1):16-31.
123. Major AS, Harrison DG. What fans the fire: insights into mechanisms of inflammation in atherosclerosis and diabetes mellitus. *Circulation.* 2011;124(25):2809-11.
124. Sasaki N, Yamashita T, Takeda M, Hirata K. Regulatory T cells in atherogenesis. *J Atheroscler Thromb.* 2012;19(6):503-15.
125. Redman CW, Sargent IL. Latest advances in understanding preeclampsia. *Science.* 2005;308(5728):1592-4.
126. Saito S, Nakashima A, Shima T, Ito M. Th1/Th2/Th17 and regulatory T-cell paradigm in pregnancy. *Am J Reprod Immunol.* 2010;63(6):601-10.
127. Aluvihare VR, Kallikourdis M, Betz AG. Regulatory T cells mediate maternal tolerance to the fetus. *Nat Immunol.* 2004;5(3):266-71.
128. Tilburgs T, Scherjon SA, van der Mast BJ, Haasnoot GW, Versteeg V, Roelen DL, van Rood JJ, Claas FH. Fetal-maternal HLA-C mismatch is associated with decidual T cell activation and induction of functional T regulatory cells. *J Reprod Immunol.* 2009;82(2):148-57.
129. Rieber-Mohn AB, Sugulle M, Wallukat G, Alnaes-Katjavivi P, Leite Storvold G, Bolstad N, Redman CW, Dechend R, Staff AC. Auto-antibodies against the angiotensin II type I receptor in women with uteroplacental acute atherosclerosis and preeclampsia at delivery and several years postpartum. *J Reprod Immunol.* 2018;128:23-9.
130. Herse F, Staff AC, Hering L, Muller DN, Luft FC, Dechend R. AT1-receptor autoantibodies and uteroplacental RAS in pregnancy and pre-eclampsia. *J Mol Med.* 2008;86(6):697-703.
131. Staff AC, Dechend R, Pijnenborg R. Learning from the placenta: acute atherosclerosis and vascular remodeling in preeclampsia-novel aspects for atherosclerosis and future cardiovascular health. *Hypertension.* 2010;56(6):1026-34.
132. Majdalawieh A, Ro HS. LPS-induced suppression of macrophage cholesterol efflux is mediated by adipocyte enhancer-binding protein 1. *Int J Biochem Cell Biol.* 2009;41(7):1518-25.
133. Staff AC, Redman CW. IFPA Award in Placentology Lecture: preeclampsia, the decidual battleground and future maternal cardiovascular disease. *Placenta.* 2014;35 Suppl:S26-S31.

134. Sattar N, Greer IA. Pregnancy complications and maternal cardiovascular risk: opportunities for intervention and screening? *BMJ*. 2002;325(7356):157-60.
135. Staff AC, Johnsen GM, Dechend R, Redman CWG. Preeclampsia and uteroplacental acute atherosclerosis: immune and inflammatory factors. *J Reprod Immunol*. 2013;In press.
136. WHO E, IDF. Definition and diagnosis of diabetes mellitus and intermediate hyperglycemia: report of a WHO/IDF consultation. WHO Press, Geneva, Switzerland: **WHO Library Cataloguing-in-Publication Data**; 2006. p. 1-50.
137. Johnsen SL, Rasmussen S, Wilsgaard T, Sollien R, Kiserud T. Longitudinal reference ranges for estimated fetal weight. *Acta Obstet Gynecol Scand*. 2006;85(3):286-97.
138. Arduini D, Rizzo G. Normal values of Pulsatility Index from fetal vessels: a cross-sectional study on 1556 healthy fetuses. *J Perinat Med*. 1990;18(3):165-72.
139. DeVore GR. The importance of the cerebroplacental ratio in the evaluation of fetal well-being in SGA and AGA fetuses. *Am J Obstet Gynecol*. 2015;213(1):5-15.
140. McManus JF. Histological and histochemical uses of periodic acid. *Stain Technol*. 1948;23(3):99-108.
141. Pereira L, Petitt M, Fong A, Tsuge M, Tabata T, Fang-Hoover J, Maidji E, Zydek M, Zhou Y, et al. Intrauterine growth restriction caused by underlying congenital cytomegalovirus infection. *J Infect Dis*. 2014;209(10):1573-84.
142. de Villiers WJ, Smart EJ. Macrophage scavenger receptors and foam cell formation. *J Leukoc Biol*. 1999;66(5):740-6.
143. Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. *Fam Med*. 2005;37(5):360-3.
144. Brosens I, Pijnenborg R, Vercruysse L, Romero R. The "Great Obstetrical Syndromes" are associated with disorders of deep placentation. *Am J Obstet Gynecol*. 2011;204(3):193-201.
145. Ranheim T, Haugen F, Staff AC, Braekke K, Harsem NK, Drevon CA. Adiponectin is reduced in gestational diabetes mellitus in normal weight women. *Acta Obstet Gynecol Scand*. 2004;83(4):341-7.
146. Staff AC, Braekke K, Harsem NK, Lyberg T, Holthe MR. Circulating concentrations of sFlt1 (soluble fms-like tyrosine kinase 1) in fetal and maternal serum during preeclampsia. *Eur J Obstet Gynecol Reprod Biol*. 2005;122(1):33-9.
147. Braekke K, Holthe MR, Harsem NK, Fagerhol MK, Staff AC. Calprotectin, a marker of inflammation, is elevated in the maternal but not in the fetal circulation in preeclampsia. *Am J Obstet Gynecol*. 2005;193(1):227-33.
148. Braekke K, Harsem NK, Staff AC. Oxidative Stress and Antioxidant Status in Fetal Circulation in Preeclampsia. *Pediatr Res*. 2006.
149. Staff AC, Braekke K, Johnsen GM, Karumanchi SA, Harsem NK. Circulating concentrations of soluble endoglin (CD105) in fetal and maternal serum and in amniotic fluid in preeclampsia. *Am J Obstet Gynecol*. 2007;197(2):176-.
150. Hering L, Herse F, Verlohren S, Park JK, Wellner M, Qadri F, Pijnenborg R, Staff AC, Huppertz B, et al. Trophoblasts reduce the vascular smooth muscle cell proatherogenic response. *Hypertension*. 2008;51(2):554-9.
151. Sugulle M, Dechend R, Herse F, Weedon-Fekjaer MS, Johnsen GM, Brosnihan KB, Anton L, Luft FC, Wollert KC, et al. Circulating and placental growth-differentiation factor 15 in preeclampsia and in pregnancy complicated by diabetes mellitus. *Hypertension*. 2009;54(1):106-12.
152. Weedon-Fekjaer MS, Dalen KT, Solaas K, Staff AC, Duttaroy AK, Nebb HI. Activation of LXR increases acyl-CoA synthetase activity through direct regulation of ACSL3 in human placental trophoblast cells. *J Lipid Res*. 2010;51(7):1886-96.

153. Kvehaugen AS, Andersen LF, Staff AC. Anthropometry and cardiovascular risk factors in women and offspring after pregnancies complicated by preeclampsia or diabetes mellitus. *Acta Obstet Gynecol Scand*. 2010;89(11):1478-85.
154. Kvehaugen AS, Dechend R, Ramstad HB, Troisi R, Fugelseth D, Staff AC. Endothelial function and circulating biomarkers are disturbed in women and children after preeclampsia. *Hypertension*. 2011;58(1):63-9.
155. Sugulle M, Herse F, Hering L, Mockel M, Dechend R, Staff AC. Cardiovascular Biomarker Midregional Proatrial Natriuretic Peptide During and After Preeclamptic Pregnancies. *Hypertension*. 59:395-401.
156. Sole KB, Staff AC, Laine K. The association of maternal country of birth and education with hypertensive disorders of pregnancy: A population-based study of 960 516 deliveries in Norway. *Acta Obstet Gynecol Scand*. 2018;97(10):1237-47.
157. Roland MC, Friis CM, Voldner N, Godang K, Bollerslev J, Haugen G, Henriksen T. Fetal growth versus birthweight: the role of placenta versus other determinants. *PLoS One*. 2012;7(6):e39324.
158. Stevens DU, Al-Nasiry S, Bulten J, Spaanderman ME. Decidual vasculopathy and adverse perinatal outcome in preeclamptic pregnancy. *Placenta*. 2012.
159. Zhang P, Schmidt M, Cook L. Maternal vasculopathy and histologic diagnosis of preeclampsia: poor correlation of histologic changes and clinical manifestation. *Am J Obstet Gynecol*. 2006;194(4):1050-6.
160. Brosens I, Renaer M. On the pathogenesis of placental infarcts in pre-eclampsia. *J Obstet Gynaecol Br Commonw*. 1972;79(9):794-9.
161. Gerretsen G, Huisjes HJ, Elema JD. Morphological changes of the spiral arteries in the placental bed in relation to pre-eclampsia and fetal growth retardation. *Br J Obstet Gynaecol*. 1981;88(9):876-81.
162. Robson SC, Simpson H, Ball E, Lyall F, Bulmer JN. Punch biopsy of the human placental bed. *Am J Obstet Gynecol*. 2002;187(5):1349-55.
163. van Dijk RA, Rijs K, Wezel A, Hamming JF, Kolodgie FD, Virmani R, Schaapherder AF, Lindeman JH. Systematic Evaluation of the Cellular Innate Immune Response During the Process of Human Atherosclerosis. *Journal of the American Heart Association*. 2016;5(6).
164. Sasaki Y, rmochwal-Kolarz D, Suzuki D, Sakai M, Ito M, Shima T, Shiozaki A, Rolinski J, Saito S. Proportion of peripheral blood and decidual CD4(+) CD25(bright) regulatory T cells in pre-eclampsia. *Clin Exp Immunol*. 2007;149(1):139-45.
165. Pijnenborg R, Vercruysse L, Brosens I. Deep placentation. *Best Pract Res Clin Obstet Gynaecol*. 2011;25(3):273-85.
166. Jellinek H. Failure of etiological differentiation of human diseases associated with fibrinoid necrosis. *Angiology*. 1970;21(11):691-9.
167. Kaufmann P, Huppertz B, Frank HG. The fibrinoids of the human placenta: origin, composition and functional relevance. *Ann Anat*. 1996;178(6):485-501.
168. Fosheim IK, Alnaes-Katjavivi P, Redman C, Roald B, Staff AC, Størvold GL. Acute atherosclerosis of decidua basalis; characterization of spiral arteries, endothelial status and activation. *Placenta*. 2019;82:10-6.
169. van Dijk RA, Duiniveld AJ, Schaapherder AF, Mulder-Stapel A, Hamming JF, Kuiper J, de Boer OJ, van der Wal AC, Kolodgie FD, et al. A change in inflammatory footprint precedes plaque instability: a systematic evaluation of cellular aspects of the adaptive immune response in human atherosclerosis. *Journal of the American Heart Association*. 2015;4(4).

170. Jourdan PM, Holmen SD, Gundersen SG, Roald B, Kjetland EF. HIV target cells in *Schistosoma haematobium*-infected female genital mucosa. *Am J Trop Med Hyg*. 2011;85(6):1060-4.
171. Taflin C, Nochy D, Hill G, Frouget T, Rioux N, Verine J, Bruneval P, Glotz D. Regulatory T cells in kidney allograft infiltrates correlate with initial inflammation and graft function. *Transplantation*. 2010;89(2):194-9.
172. de Boer OJ, van der Meer JJ, Teeling P, van der Loos CM, van der Wal AC. Low numbers of FOXP3 positive regulatory T cells are present in all developmental stages of human atherosclerotic lesions. *PLoS One*. 2007;2(8):e779.
173. Mjosberg J, Berg G, Jenmalm MC, Ernerudh J. FOXP3+ regulatory T cells and T helper 1, T helper 2, and T helper 17 cells in human early pregnancy decidua. *Biol Reprod*. 2010;82(4):698-705.
174. Quinn KH, Parast MM. Decidual regulatory T cells in placental pathology and pregnancy complications. *Am J Reprod Immunol*. 2013;69(6):533-8.
175. Przybyl L, Ibrahim T, Haase N, Golic M, Rugor J, Luft FC, Bendix I, Serdar M, Wallukat G, et al. Regulatory T cells ameliorate intrauterine growth retardation in a transgenic rat model for preeclampsia. *Hypertension*. 2015;65(6):1298-306.
176. Foks AC, Lichtman AH, Kuiper J. Treating atherosclerosis with regulatory T cells. *Arterioscler Thromb Vasc Biol*. 2015;35(2):280-7.
177. Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med*. 2005;352(16):1685-95.
178. Wang Y, Zhang M, Liu ZW, Ren WG, Shi YC, Sun YL, Wang HB, Jin L, Wang FS, et al. The ratio of circulating regulatory T cells (Tregs)/Th17 cells is associated with acute allograft rejection in liver transplantation. *PLoS One*. 2014;9(11):e112135.
179. Kell DB, Pretorius E. To What Extent Are the Terminal Stages of Sepsis, Septic Shock, Systemic Inflammatory Response Syndrome, and Multiple Organ Dysfunction Syndrome Actually Driven by a Prion/Amyloid Form of Fibrin? *Semin Thromb Hemost*. 2018;44(3):224-38.
180. Redline RW, Boyd T, Campbell V, Hyde S, Kaplan C, Khong TY, Prashner HR, Waters BL. Maternal vascular underperfusion: nosology and reproducibility of placental reaction patterns. *Pediatr Dev Pathol*. 2004;7(3):237-49.
181. Redline RW, Heller D, Keating S, Kingdom J. Placental diagnostic criteria and clinical correlation--a workshop report. *Placenta*. 2005;26 Suppl A:S114-S7.
182. Johnsen GM, Storvold GL, Drabbels JJM, Haasnoot GW, Eikmans M, Spruyt-Gerritse MJ, Alnaes-Katjavivi P, Scherjon SA, Redman CWG, et al. The combination of maternal KIR-B and fetal HLA-C2 is associated with decidua basalis acute atherosclerosis in pregnancies with preeclampsia. *J Reprod Immunol*. 2018;129:23-9.

15. APPENDIX 1:

Microscopy worksheet for Paper I

DSM Decidua basalis		M-ID:	Histology ID:	Date of Study:												
Slide no.	Stain	Findings		Vessel wall section							Local interstitium					
1	H&E	Vessel outer-outer diameter not specified		1	2	3	4	5	6	7	No (0), Yes (1)					
		No? STOP! Yes (1)									xxxxxxxxxxxxxxxxxxxxxxxxxx					
		Solitary mural "foam" cells No (0), Yes (1)									xxxxxxxxxxxxxxxxxxxxxxxxxx					
		"Foam" cells, 2 or more, No (0), Yes (1)														
		Perivasculature decidua present No (0), Yes (1)									xxxxxxxxxxxxxxxxxxxxxxxxxx					
		Evidence of intraluminal thrombus grade 0,1,2,3									xxxxxxxxxxxxxxxxxxxxxxxxxx					
2	Desmin	Positive cells No (0), Yes (1)									xxxxxxxxxxxxxxxxxxxxxxxxxx					
		Smooth muscle layer intact No (0), Yes (1)														
		Separate No (0), Yes (1)									xxxxxxxxxxxxxxxxxxxxxxxxxx					
		Disrupted No (0), Yes (1)									xxxxxxxxxxxxxxxxxxxxxxxxxx					
		Disorganized No (0), Yes (1)									xxxxxxxxxxxxxxxxxxxxxxxxxx					
		Perivascular myometrial cells No (0), Yes (1)									Desmin positive cells					
	Periodic Acid Schiff (PAS)	PAS + No (0), 25% (1), 50% (2), 75% (3), 100% (4)									PAS present					
3	CK7 - PAS	No. of trophoblast cells lining the lumen									InterstitialCK 7+ cells					
		No. of intramural trophoblast cells									xxxxxxxxxxxxxxxxxxxxxxxxxx					
		No. of perivascular trophoblast cells									xxxxxxxxxxxxxxxxxxxxxxxxxx					
		Giant cells recognized? 2+ nuclei No (0), Yes (1)									Giant cells recognized?					
	Impression of Perivasc. mononuclear cell infiltrate?	No (0), Unsure (1), Yes (2)									Mononuclear cell 25u r					
4	CD68 - PAS	No. of cells in largest intramural cluster									CD68 positive cells 0/1					
		Perivasculature CD68 positive cells No (0), Yes (1)									xxxxxxxxxxxxxxxxxxxxxxxxxx					
5	H&E	Not included here									xxxxxxxxxxxxxxxxxxxxxxxxxx					
											xxxxxxxxxxxxxxxxxxxxxxxxxx					
6 - 10	Stored for future IHC															
Vessel type (1,2,3,4,5,6) - please see separate key.																
Comments regarding abnormal features, difficulties with defining specific findings:																
Conclusion: Please tick: ☐ The slides do not demonstrate any features pertaining to the presence of acute atherosclerosis. ☐ The slides demonstrate the presence of acute atherosclerosis according to criteria: 1. 3. 4. 5. (please encircle)																

Microscopy worksheet for Paper III

Sample comments:

Acute atherosclerosis in vacuum suction biopsies of decidua basalis: An evidence based research definition

Patji Alnaes-Katjavivi ^a, Fiona Lyall ^b, Borghild Roald ^c, Christopher W.G. Redman ^d, Anne Cathrine Staff ^{a,*}

^a Department of Obstetrics and Department of Gynecology, Oslo University Hospital, Ullevål and Faculty of Medicine, University of Oslo, Norway

^b University of Glasgow, School of Medicine, Glasgow, UK

^c Department of Pathology, Oslo University Hospital, Ullevål and Faculty of Medicine, University of Oslo, Norway

^d Department of Obstetric Medicine, University of Oxford, Oxford, UK

ARTICLE INFO

Article history:

Received 8 June 2015

Received in revised form

26 October 2015

Accepted 27 October 2015

Keywords:

Placenta

Decidua

Acute atherosclerosis

Morphology

Immunohistochemistry

Criteria

Preeclampsia

ABSTRACT

Background: Acute atherosclerosis (AA) of the uteroplacental spiral arteries has been characterised by sub-endothelial lipid-laden foam cells, perivascular leukocyte infiltrates (PVI) and fibrinoid necrosis. Because precise diagnostic criteria are not available for comparative research studies we developed and tested new simplified criteria based on 237 cases.

Methods: Decidual basalis samples were collected by vacuum suction at elective cesarean deliveries. Spiral arteries were evaluated in serial decidual tissue sections from women with normal pregnancy, preeclampsia, and diabetes. Features of AA were sought in parallel sections stained with H&E and immunostained for CD68, cytokeratin CK7 and desmin, and costained with Periodic Acid Schiff (PAS).

Results: Foam cell lesions were defined as two or more adjacent, intramural, vacuolated CD68 positive cells, PVI as a focal perivascular lymphocyte accumulation, more dense than in the surrounding decidua. Increased fibrinoid (PAS positive) was identified if present in $\geq 75\%$ of the arterial wall circumference. PVI and increased fibrinoid were significantly associated with preeclampsia but not specifically associated with the presence of foam cell lesions. Hence we diagnosed decidua basalis AA lesions solely by the presence of foam cell lesions, occurring in preeclampsia (37%), diabetes (10%) and healthy normotensive women (11%). The simplified criterion was reproducible by different investigators. Decidua basalis AA occurred most commonly and extensively in preeclampsia, but did not distinguish between preterm and term disease.

Discussion: Our evidence based criterion for decidua basalis AA diagnosis in vacuum suction biopsies may not apply to myometrial or decidua parietalis arteries. In decidual basalis samples it should facilitate comparisons between research studies, to improve pathophysiological understanding of AA and preeclampsia.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Acute atherosclerosis (AA) of the uteroplacental spiral arteries is commonly described in, but not confined to, preeclampsia [1]. Maternal hypertension is not mandatory, as it occurs in autoimmune diseases, diabetes mellitus and even normal pregnancy [2]. AA is associated with spiral artery thrombosis and downstream

placental infarction [3,4]. The etiology and pathophysiology of AA is not established, and may have implications for the general understanding of artery physiology and diseases, far beyond pregnancy and preeclampsia. AA may share pathophysiological features [1,5] with allograft arterial lesions during immune rejection [6] and early atherosclerosis [7], which are similar but not identical artery lesions.

The first descriptions [8,9] identified lipid laden foam cells in the arterial wall, together with fibrinoid necrosis and perivascular infiltrates (PVI) as the hallmarks of acute atherosclerosis [10–12]. Foam cells of AA lesions have been assumed to be mostly lipid-laden macrophages [13]. Criteria to identify “fibrinoid necrosis” or PVI

* Corresponding author. Department of Obstetrics and Department of Gynecology, Women and Children's Division, Oslo University Hospital, Ullevål, PO Box 4956 Nydalen, NO-0424 Oslo, Norway.

E-mail address: uxnnaf@ous-hf.no (A.C. Staff).

are lacking. In some reports PVI was not a diagnostic criterion [8,12,14,15]. Moreover, the extent to which the three components may separately contribute to AA lesion quantification and severity has not been considered.

AA occurs more often in the decidual than myometrial segments of the spiral arteries [16]. The lesions in the unremodelled muscular segments may differ from those in the fully remodelled decidual basalis segments or indeed in the decidual parietalis [9,10]. In this study we only consider decidual basalis. Our vacuum suction technique of sampling the decidual basalis, which has a higher yield of decidual spiral arteries than traditional placental bed and basal plate biopsies [17], was used to develop and test diagnostic criteria for AA that could be reliably used for research purposes and for comparison between studies. Our aim was, for the first time, to define evidence based criteria and test them for inter- and intra-observer reproducibility. We then used the revised criteria to quantify AA and its component features in preeclampsia, diabetes and normotensive control pregnancies. We analysed spiral artery thrombi [11] and their relation to pregnancy outcome and AA. Throughout this paper the unqualified term ‘spiral artery’ refers only to their decidual basalis segments unless otherwise specified. Likewise “acute atherosclerosis” (AA) refers to the lesion in decidual basalis spiral arteries. In consequence it is important to recognize that all our conclusions relate to AA in the spiral arteries in the vacuum suction biopsies of decidual basalis. The lesion may differ in other locations (myometrial segments or in the decidual parietalis).

2. Methods

2.1. Patient selection

Pregnant women ($n = 278$) were recruited from January 2000 to December 2013 to the Oslo Pregnancy Biobank (www.oslo-universitetssykehus.no/opb). They were scheduled for elective caesarean section, with intact membranes, and absence of regular uterine contractions and no signs of infection. All women provided informed written consent and the study was approved by the Regional Committee for Medical and Health Research Ethics. Forty-one women were excluded because their decidual samples lacked sufficient spiral arteries. The remaining 237 (85%) were allocated to one of four pregnancy groups (Table 1); normotensive controls ($n = 94$), preeclampsia ($n = 87$), diabetes mellitus (all types, pregestational or gestational, without preeclampsia, $n = 39$) and diabetes mellitus (all types) with superimposed preeclampsia ($n = 17$). Preeclampsia was defined as new onset hypertension (blood pressure $\geq 140/90$ mmHg) and new onset proteinuria ($\geq 1 +$ on

dipstick, or ≥ 30 protein/creatinine ratio) ≥ 20 weeks of pregnancy [18]. Diabetic women included 23 with pregestational diabetes mellitus (17 of type 1 and 6 of type 2) and 16 with gestational diabetes, as defined by the WHO 1999 guidelines [19].

Gestational week (GW) at delivery was defined by routine ultrasound screening at GW 17–20. Newborn gender specific baby weight percentiles were calculated according to Norwegian fetal growth curves [20]. Normotensive controls had uncomplicated pregnancies without clinical evidence of fetal growth restriction (and all birth weights were above the 5th birth weight percentile).

2.2. Collection of decidual basalis tissue by vacuum suction, stainings and spiral artery examinations

Following placental delivery, the uterine wall of the placental bed was aspirated by vacuum suction, as previously described [17,21]. Parts of the harvested tissue were formalin fixed, routinely processed and paraffin embedded. Four $3 \mu\text{m}$ serial sections were stained (Supplementary Table 1), with Haematoxylin and Eosin (H&E) and immunostained for smooth muscle cells (Desmin, Clone DE R-11, Roche Ventana Medical system Inc, Arizona, USA, ready-made), trophoblasts (Cytokeratin 7 (CK7), Clone OV-TL 12/30, Dako AS, Glostrup, Denmark, dilution 1/300) and for CD68 (CD68, Clone KP1, Dako AS, Glostrup, Denmark, dilution 1/1000, identifying monocytes, macrophages and foam cells). The immunostained sections were costained with Periodic Acid Schiff (PAS) [22] for fibrinoid detection. The sections were evaluated with a Motic BA-410 LED light microscope (objectives 4, 10 and 40) and recorded with a Moticam 2500 digital camera.

Decidual spiral arteries were evaluated for acute atherosclerosis (AA) only if their diameter was $\geq 140 \mu\text{m}$ including the outer vessel wall, which could contain desmin positive smooth muscle cells and/or PAS positive fibrinoid. The presence of CK7 interstitial trophoblasts were also noted [23].

In all 237 patients, the decidual sections included at least one cross-section of one spiral artery. Spiral arteries were first identified on the H&E section, and followed in the three immunostained serial sections (Supplementary Table 1). The median number of spiral artery sections identified and evaluated for each patient per slide was two (healthy control women), three (preeclampsia) and four (diabetes with or without superimposed preeclampsia). The examiners were blinded to the pregnancy characteristics during their microscopic tissue evaluations and recordings.

We focused on the three histological features of AA that have been previously emphasised. Detailed, morphological criteria were defined for intramural foam cells, PVI and fibrinoid of the arterial

Table 1

Clinical characteristics of women providing a decidual sample with spiral arteries identified ($n = 237$), according to pregnancy groups. Data are presented as median values, percentages, or rates.

	Normotensive controls $n = 94$	PE $n = 87$	DM $n = 39$	DM + PE $n = 17$	Total $n = 237$
Age at delivery (years)	33	32	33	32	32
BMI, pre-pregnancy (kg/m^2)	22.1	23.6	27.6	25.1	23.2
BMI, at delivery (kg/m^2)	27.2	29.8	32.2	33.2	29.0
First pregnancy (gravida 1)	29%	49%	21%	53%	37%
First delivery (nulliparous)	49%	63%	36%	76%	54%
Previous PE (of women para ≥ 1)	4% (2/48)	39% (12/31)	16% (4/25)	75% (3/4)	19% (21/108)
Gestational age at delivery (weeks ⁺ days)	38 ⁺⁵	33 ⁺⁰	38 ⁺⁰	34 ⁺⁵	37 ⁺⁵
^a Mean Systolic BP < 20 GW (mmHg)	110	116	115	120	115
^a Mean Diastolic BP < 20 GW (mmHg)	65	72	71	70	70
Systolic BP at delivery (mmHg)	120	160	135	160	140
Diastolic BP at delivery (mmHg)	70	100	79	95	85

PE: preeclampsia, DM: diabetes mellitus (all types), DM + PE: diabetes mellitus with superimposed preeclampsia, BMI: body mass index, BP: blood pressure, GW: gestational week.

^a This value is the median value for the mean of all blood pressures registered before the 20th week of gestation, per woman.

Table 2
Study definitions for the investigated decidual spiral artery morphology.

Definition	Description
Artery wall margins	The inner margin identified by the luminal endothelial surface. Outer margin identified by the contact between vessel wall components and surrounding decidual tissue
Vacuolated cells	≥2 adjacent vacuolated cells (in H&E sections), no more than one cell width apart, localized within the vessel wall margins
CD68 positivity	CD68 + in ≥2 adjacent vacuolated cells (as described above) was considered confirmatory for lipid-laden foam cell lesion
Perivascular	The tissue area surrounding the spiral artery cross section within 50 µm perpendicular to its outer margin
Perivascular infiltrate (PVI)	A perivascular (as defined above) focal or uniform presence of mononuclear lymphocytic cells with a density greater than that of the surrounding decidual tissue
Fibrinoid	PAS staining within the vessel wall, graded by proportion of vessel circumference: 0: No PAS stain, 1: <25%, 2: 25%–49%, 3: 50%–74%, 4: ≥75%–100%
Desmin positivity	Identifying smooth muscle cells
Vessel lumen thrombosis	H&E identification of red blood cell within a fibrin meshwork occupying >10% of the vessel lumen
Surrounding decidual tissue	Decidual tissue with exception of the spiral arteries and perivascular area
Trophoblasts, extravillous	CK7+ cells, usually as single cells: <u>Intramural</u> : within the margins of the vessel wall, counted by eye <u>Interstitial</u> : in the surrounding decidual tissue, as defined above

wall to allow their consistent and accurate recognition. A major criterion used by all previous AA studies is the presence of mural foam cells. In formalin fixed, routinely processed tissue material using ethanol (used in most studies), intracellular lipid droplets disappear in the preparation process and thus cannot be demonstrated directly. The lipid laden cells (“foam cells”) will thus appear “vacuolated”. The scavenger receptor CD68, one of several groups of scavenger receptors important in the pathogenesis of atherosclerosis, is present in monocytes, macrophages and foam cells [24]. We identified vacuolated cells that were CD68 positive (CD68+) as “foam cells” (Table 2), providing there were least two such intramural cells adjacent to each other, representing a ‘foam cell lesion’. We decided on this definition in order to exclude possible over-registrations. In H&E sections, cells may have vacuolated cytoplasm due to degeneration (hydropic degeneration) and non-vacuolated monocytes (CD68 + cells) are regularly seen in the decidua and arterial wall. Our foam cell definition thus diminished these pitfalls by combining the morphological and immunohistochemical criteria and the close spatial localisation of at least two such cells.

In the absence of a published definition, PVI was defined as a focal perivascular accumulation of lymphocytes (within 50 µm to the outer margin of the spiral artery), more dense than in the surrounding tissue (Table 2). It is reported that fibrinoid necrosis may appear more bluish rather than red on PAS staining [23]. In the present research study we could not consistently quantify this. Instead, the presence of fibrinoid was noted and graded according to its intramural extent (from absent to ≥75% of the vessel circumference, Table 2).

There is no gold standard of AA diagnosis except that, to justify the name, foam cells must be present. Nor is AA diagnostic of a specific clinical condition. Using our defined criteria for the morphological features, we analysed how they could contribute to a reproducible definition and morphological diagnosis of AA. Our starting point was thus to identify one or more spiral arteries with one or more foam cell lesions per “case”. A “case” in our study represents a woman providing a single sample of decidua basalis, in which one or more artery cross-sections yielded evidence or absence of foam cell lesions. The number of decidual artery sections affected by foam cell lesions was registered. The number of CD68 + vacuolated cells (at least 2 adjacent cells to be identified as “foam cells”) in the largest foam cell lesion was counted, and the maximum cross-sectional area of an arterial wall foam cell lesion was measured as a percentage of the total arterial wall surface area (using Motic images version 2.0). Per case (one decidual sample per

case/woman), a single count included the four serial slides, as these parallel four slides identified the same arterial cross-section(s) and decidual area. PVI and fibrinoid were recorded, as defined, and whether or not PVI and/or fibrinoid co-localised with a foam cell lesion.

2.3. Reproducibility of revised classification

Inter- and intra-observer agreement was tested after each observer independently had evaluated whether any vessel section fulfilled the defined criteria. Decidual slides with spiral arteries from 16 different women were evaluated on one occasion (PAK versus FL) and from 58 women on two occasions (PAK versus BR). Intra-observer agreement was evaluated for 58 women with spiral artery sections on 2 occasions with 8 weeks apart (PAK vs PAK).

2.4. Statistical analyses

Data were analysed using IBM Statistical Package for the Social Sciences software (SPSS) version 22. The Fisher Exact test was used for dichotomous variables, and the non-parametric Mann-Whitney- Wilcoxon test otherwise. A probability of ≤0.05 was considered statistically significant.

3. Results

Only a few of the vacuum suction specimens contained microscopic fragments of myometrium, the focus of this study was thus exclusively on decidual artery morphology.

3.1. Decidual artery morphology

At least one vacuolated cell in the spiral artery wall, identified in H&E section, was common in all pregnancy groups (81% for the total group, Table 3), without significant differences between the groups. Two or more adjacent vacuolated mural cells (regardless of CD68 labelling) was less common (35% for the total group, Table 3), but significantly more common in preeclampsia as compared to normotensive controls (45% vs 26%, $p = 0.005$). Similarly, per case, presence of a single intramural CD68 + staining cell was more common than were two or more adjacent CD68 + staining intramural cells, for the total pregnancy group (50% versus 32%, $p < 0.001$). Two or more adjacent vacuolated cells (35%) usually represented ≥2 adjacent CD68 + cells (32%) in at least one of the same artery serial cross-sections. For the total patient group, 22%

Table 3

Decidual acute atherosclerosis morphological criteria, and morphology rates. Data are given in numbers (and percentages) according to pregnancy groups. One case within a pregnancy group represents the parallel sections from same woman, identifying the same spiral arteries across the 4 parallel sections investigated per sample (1 decidual sample per woman).

	Normotensive controls n = 94	PE n = 87	DM n = 39	DM + PE n = 17	Total n = 237
Solitary vacuolated cell in vessel wall	47 (50%)	36 (41%)	16 (41%)	7 (41%)	106 (45%)
≥1 vacuolated cells	70 (74%)	77 (89%) ^a	31 (79%)	14 (82%)	192 (81%)
≥2 vacuolated cells	23 (24%)	39 (45%) ^a	15 (38%)	7 (41%)	84 (35%)
Solitary CD68 positive vessel wall cell	58 (62%)	39 (45%) ^a	13 (33%) ^a	8 (47%)	118 (50%)
≥1 CD68 positive vessel wall cell	70 (74%)	76 (87%) ^a	21 (54%) ^a	16 (94%)	194 (82%)
≥2 CD68 positive vessel wall cell	22 (23%)	37 (43%) ^a	9 (23%)	8 (47%) ^a	76 (32%)
CD68 positivity of ≥2 vessel wall vacuolated cells ("foam cells")	10 (11%)	32 (37%) ^a	4 (10%)	5 (29%) ^a	51 (22%)
Presence of perivascular infiltrate (PVI) per sample	19 (20%)	30 (34%) ^a	6 (15%)	7 (41%) ^a	62 (26%)
Colocalization of "foam cells" and PVI in same decidual sample	1 (1%)	16 (18%) ^a	1 (2%)	3 (18%) ^a	21 (9%)
Colocalization of "foam cells" and PVI in same artery	1 (1%)	12 (14%) ^a	1 (2%)	3 (18%) ^a	17 (7%)
PAS positivity grade 4 (75%–100% of total vessel wall circumference)	62 (66%)	72 (83%) ^a	26 (67%)	12 (70%)	172 (73%)
PAS positivity grade 1–3 (1%–74% of total vessel wall circumference)	32 (34%)	15 (17%) ^a	13 (33%)	5 (30%)	65 (27%)

PE: preeclampsia, DM: diabetes mellitus (all types), DM + PE: diabetes mellitus with superimposed preeclampsia, PAS: Periodic Acid Schiff.

^a Represents p-values <0.05 (comparing the PE or DM or D + PE clinical groups to the normotensive control group).

had ≥2 adjacent intramural CD68 + cells that were vacuolated. Thus foam cells lesions were identified by at least two adjacent vacuolated CD68 + cells in the arterial wall (Fig. 1a and b). The highest rates of foam cells were found in the preeclampsia group (37%) and diabetes with superimposed preeclampsia (29%) group. The lowest rates were in normotensive controls and diabetes without preeclampsia (11% and 10% respectively, Table 3).

PVI was present in at least one cross-section of a spiral artery in 26% of the total pregnancy group (Fig. 1b), and significantly more often in preeclampsia than in normotensive control pregnancies (Table 3; 34% vs 20%, $p = 0.04$). PVI and foam cell lesions did not colocalise in the same spiral artery section in 93% of the total group, and there was no significant colocalisation of PVI and foam cell lesions for the same artery segment in any of the pregnancy groups (Table 3). There was also a poor colocalisation of PVI and foam cells when considering the same decidual section (9%, Table 3), not requiring presence of PVI and foam cells in the same cross artery segment.

No decidual spiral artery had grade 0 fibrinoid, as defined in Table 2. This is as expected since, in late pregnancy, these arteries are all remodelled distally, and remodelling is accompanied by fibrinoid deposition [25]. As stated earlier it was not possible to judge the quality of the fibrinoid (including fibrinoid "necrosis") by systematic assessment of staining colour or intensity. Table 3 shows that most patients in all pregnancy groups had PAS + fibrinoid of at least 75% of their arterial wall circumference (grade 4 fibrinoid), but its presence was not significantly colocalised with foam cell lesions (data not shown). Women with preeclampsia had significantly more arterial wall fibrinoid than controls (grade 4 fibrinoid: 83% vs 66%, $p = 0.011$, Table 3). Therefore increased fibrinoid was a feature of decidual spiral arteries associated with preeclampsia, but not specifically with acute atherosclerosis.

3.2. Developing a classification of decidual acute atherosclerosis

We defined a foam cell lesion (≥2 adjacent intramural vacuolated, CD68 + cells) as a mandatory feature of AA. Foam cell lesions did not specifically colocalise with either PVI or fibrinoid excess. However both the latter features were significantly associated with preeclampsia and may, instead, be part of the decidual pathology of preeclampsia but not mandatory for AA development. Therefore we conclude that neither PVI nor mural fibrinoid excess are helpful features for identifying AA, but may have uses for sub-classifying the preeclampsia syndrome. Therefore we simplified our definition of AA to be the presence of at least one foam cell lesion (≥2

adjacent intramural vacuolated, CD68 + cells) in the spiral arterial wall (Fig. 1a and b). It should be noted that this definition is for AA of the decidua basalis only and may not apply to that observed in the myometrial segments of spiral arteries or to atherosclerosis and atherosclerosis-like lesions described in the decidua parietalis [4,26].

The classification of a pregnancy (a woman) being affected by decidual AA in this study was not altered even if other spiral arterial sections in the same decidual specimen section did not fulfil the diagnostic AA criteria.

3.3. Reproducibility of revised acute atherosclerosis (AA) classification

There was good intra- and inter-observer reproducibility for the revised criterion of AA between the three first authors. For the few discrepant cases among the three evaluators, consensus was obtained for the final AA categorisation.

Intra-observer agreement (PAK vs PAK) was almost perfect (Kappa score of 0.86) [27]. Inter-observer agreement was "substantial" [27] with Kappa scores between authors of 0.64 (PAK vs FL) and 0.70 (PAK vs BR).

3.4. Prevalence of AA across the study groups and relation to clinical features

AA rates across the pregnancy groups are presented in Table 4, and parallel the foam cell prevalence in Table 3, due to the simplified AA definition. In total, 22% of the 237 patients were diagnosed with decidual AA. The rate of AA according to the new criterion was highest for the preeclampsia group (37%) and lowest for the normotensive control group (11%) and the normotensive diabetes group (10%). The diabetes group with superimposed preeclampsia had a prevalence of AA almost as high as the preeclampsia group (29%).

As shown in Table 4, there were few statistical significant differences in clinical characteristics between women with or without AA for each pregnancy group and even for the total group of 237 women. For the total study group however, there was a higher rate of primigravidae (first pregnancy) amongst women diagnosed with AA compared to those without AA (51% vs 33%, $p = 0.02$). Among parous women, those with previous preeclampsia were more likely to have AA diagnosis in the present study than those without a previous history of PE (43% vs 13%, $p = 0.024$). For the total study group, also the median systolic and diastolic blood pressures were higher for the AA group as compared to the group without AA (158/100 vs 138/80, $p < 0.001$). In addition, for the total study cohort, AA

Fig. 1. Parallel sections from decidual basalis collected by the vacuum suction technique in 3 patients. **a–b** demonstrate decidual spiral arteries with acute atherosclerosis (AA). **c** demonstrates an artery without acute atherosclerosis. For each patient, the sequence of 4 parallel sections demonstrate the same decidual spiral artery stained with (i) hematoxylin and eosin, (ii) desmin and Periodic Acid Schiff (PAS), (iii) Cytokeratin 7 (CK7) and PAS, (iv) CD68 and PAS. **a.** (i) Luminal thrombosis (T) and intramural vacuolated cells (V). (ii) Lack of desmin positive smooth muscle layer (large black arrows) and presence of fibrinoid (PAS-positive areas*). (iii) Interstitial CK7+ extravillous trophoblasts present (black squares), but no intramural trophoblasts. (iv) CD68+ (black squares, inset) in same locations as intramural vacuolated cells (i). Perivascular infiltrate is lacking. Objective $\times 10$, $\times 40$ (inset: objective $\times 40$). **b.** Decidual AA with associated perivascular infiltrate (i: black circles). Spiral artery lumen (L) is nearly obliterated by extensive acute atherosclerosis, shown as vacuolated (i: V) and CD68+ cells (iv: black

was more frequent in the SGA groups than the remaining pregnancies without SGA (Table 4).

In the present cohort, AA was not associated with the clinical severity of preeclampsia, determined by the ACOG criteria [28]. Nor was AA more common in women with preterm preeclampsia (evaluated as delivered either prior to 34 or 37 weeks), or with birth weights ≤ 10 th or ≤ 3 rd percentiles within the preeclampsia group (Table 4).

3.5. Quantification of AA

The pathological pregnancy groups yielded more decidual tissue artery cross-sections to evaluate per decidual sample (Table 5). The median artery numbers identified with AA was 2 for the total study group. Preeclamptic patients diagnosed with AA had more arteries affected than normotensive women diagnosed with AA (median artery number affected being 2 vs 1, $p = 0.02$) (Table 5). When quantifying AA by number of foam cells in the largest lesion for a given sample (Fig. 1b), the preeclamptic groups (with or without diabetes) had the more extensive AA lesions (Table 5), with largest AA lesions containing significantly more foam cells than in the normotensive control group. No significant difference in artery wall surface area affected by AA was however found when comparing pregnancy groups (Table 5).

3.6. Additional morphological features in relation to AA

Absence of intramural trophoblasts was significantly more common in arteries containing AA for the total study group, as compared to arteries without AA (71% versus 32%, $p < 0.001$). Similar findings were also found for the three pregnancy pathology groups (Table 5). The same trend for control patients was not statistically significant (data not shown). In the arteries where intramural trophoblasts and foam cell lesions coexisted, the number of intramural trophoblasts was significantly lower than in arteries without AA (data not shown). Presence or not of extravillous trophoblasts in the surrounding interstitial decidual tissue (as defined in Table 2) did not differ between the various pregnancy groups, nor was it related to the presence or absence of AA (data not shown).

Spiral artery thrombosis (Fig. 1a) was common for all pregnancy groups. However, significantly more samples diagnosed with AA had at least one thrombosed artery as compared to samples without AA, both for the total study group (92% vs 52%, $p = 0.001$) and for the normotensive and preeclampsia groups (Table 5).

4. Discussion

This is a large systematic study of decidual AA. It involves the distal ends of uteroplacental spiral arteries in the decidua basalis, collected by vacuum suction of the placental bed following elective cesarean delivery. In line with other vascular lesions, AA is a patchy arterial lesion. AA may affect parts of some, but not all arteries [29]. Targeted tissue sampling is thus essential. The vacuum suction method is superior to traditional biopsies from the placental bed or the maternal surface of placenta, in terms of numbers of decidual spiral arteries and volume of decidual tissue obtained [17]. The

squares). Foam cells dominate the artery wall, where there is presence of fibrinoid (PAS+, ii-iv: *). Desmin-positive smooth muscle cells are absent (ii). Extravillous trophoblasts (CK7+) are seen in the interstitial decidua (iii: black squares), but are absent within the vessel wall. Objective $\times 10$. **c.** Decidua without AA. (i) Spiral artery lumen (L). (ii) Artery wall with scattered peripheral smooth muscle cells (large black arrows, desmin positive) and PAS-positive fibrinoid (*). (iii) Intramural trophoblasts (black squares). (iv) Solitary CD68+, non-vacuolated cells that may represent monocytes (black squares). Objective $\times 10$.

Table 4

Clinical features according to presence or absence of acute atherosclerosis (AA) for each of the pregnancy groups (and the total study group). AA is defined according to the simplified evidence-based criterion of intramural foam cell lesions (defined as ≥ 2 adjacent intramural vacuolated and CD68 positive cells). Values are expressed as medians, percentages, and rates.

Acute atherosclerosis (AA) (No AA or AA diagnosis)	Normotensive controls n = 94		PE n = 87		DM n = 39		DM + PE n = 17		Total n = 237	
	No AA n = 84 (89%)	AA n = 10 (11%)	No AA n = 55 (63%)	AA n = 32 (37%)	No AA n = 35 (90%)	AA n = 4 (10%)	No AA n = 12 (71%)	AA n = 5 (29%)	No AA n = 186 (78%)	AA n = 51 (22%)
Age at delivery (years)	34	33	32	32	33	31	33	31	33	32
First pregnancy (gravida 1)	26%	50%	51%	47%	17%	50%	42%	43%	33%	51% ^a
BMI: pre-pregnancy (kg/m ²)	22.1	22.0	23.1	24.3	24.3	27.6	24.4	27.7	22.7	24.6
BMI: at delivery (kg/m ²)	27.2	27.5	30.1	29.7	33.3	29.6	31.5	36.7	29.0	29.3
^b Median Systolic BP < 20weeks (mmHg)	108	110	119	115	117	115	130	119	117	113 ^a
^b Median Diastolic BP < 20weeks (mmHg)	67	65	72	71	72	71	75	70	72	70
BP at delivery (mmHg) (Systolic BP/Diastolic BP)	120/71	119/70	162/100	160/104	135/79	132/77	160/95	162/100	138/80	158 ^a /100 ^a
Gestational age at delivery (weeks ⁺ days)	38 ⁺ 5	38 ⁺ 5	33 ⁺ 5	32 ⁺ 4	38 ⁺ 0	37 ⁺ 2	34 ⁺ 4	36 ⁺ 5	38 ⁺ 0	34 ⁺ 0 ^a
Delivery <37 ⁺ 0	2%	10%	87%	91%	20%	25%	92%	60%	37%	67% ^a
Delivery <34 ⁺ 0	1%	0%	55%	69%	3%	25%	71%	40%	26%	51% ^a
Birth percentile	66%	50%	6%	2%	98%	55% ^a	62%	66%	55%	16% ^a
Small for gestational age (birthweight $\leq 10\%$)	2%	0%	58%	69%	3%	25%	25%	0%	20%	45% ^a
Small for gestational age (birthweight $\leq 3\%$)	0%	0%	42%	59%	3%	25%	17%	0%	14%	39% ^a
Previous PE (amongst parous women)	0% (0/44)	67% ^a (2/3)	33% (6/18)	46% (6/13)	16% (4/25)	0% (0/4)	67% (2/3)	100% (1/1)	13% (12/90)	43% ^a (9/21)
Severe PE according to ACOG (28)	0%	0%	82%	78%	0%	0%	67%	75%	29%	54% ^a

PE: preeclampsia, DM: diabetes mellitus (all types), DM + PE: diabetes mellitus with superimposed preeclampsia, BMI: body mass index.

BP: blood pressure.

^a p-values <0.05 when comparing the variable investigated (e.g. rate of primiparity) for the “AA present” and “no AA” groups within the specific pregnancy group (e.g. PE or DM) or within the total group (last column: comparing all women with AA to all women without AA; independent of clinical group).

^b Presents the median blood pressure (systolic or diastolic) value for the specific pregnancy group before gestation week 20 (the mean per woman is first calculated of all blood pressures registered per woman before gestation week 20).

Table 5

Quantification of acute atherosclerosis (AA) and additional morphological features for each of the pregnancy groups (and the total study group). AA is defined according to the simplified evidence-based criterion of intramural foam cell lesions (defined as ≥ 2 adjacent intramural vacuolated and CD68 positive cells). Values are expressed as medians, percentages, and ranges.

Acute atherosclerosis (AA) (No AA or AA diagnosis)	Normotensive Controls n = 94		PE n = 87		DM n = 39		DM + PE n = 17		Total n = 237	
	No AA n = 84 (89%)	AA n = 10 (11%)	No AA n = 55 (63%)	AA n = 32 (37%)	No AA n = 35 (90%)	AA n = 4 (10%)	No AA n = 12 (71%)	AA n = 5 (29%)	No AA n = 186 (78%)	AA n = 51 (22%)
Number of artery cross-sections per decidual tissue sample	2 (1–12)	5 (1–5)	2 (1–8)	4 (1–11)	2 (1–8)	6 (2–7)	2 (1–8)	4 (4–10)	2 (1–12)	3.5 (1–11)
Number of artery cross-sections diagnosed with AA per sample	–	1 (1–2)	–	2 (1–7)	–	2 (2–4)	–	3 (1–6)	–	2 (1–7)
Largest number of foam cells per single lesion of AA	–	3 (2–4)	–	4 (2–25)	–	2.5 (2–5)	–	11 (4–23)	–	4 (2–25)
Largest surface area of artery wall filled by foam cells	–	3.3% (0.4–21.7)	–	8.2% (0.6–62.4)	–	4.6% (1.2–18.8)	–	12.5% (5.9–28.6)	–	7.3% (0.4–62.3)
Decidual sample containing arteries without intramural extravillous trophoblasts	30%	80% ^a	42%	69% ^a	20%	75% ^a	33%	60%	32%	71% ^a
Decidual samples with at least one partially of fully thrombosed artery lumen	56%	100% ^a	62%	94% ^a	57%	75%	50%	80%	52%	92% ^a

PE: preeclampsia, DM: diabetes mellitus (all types), DM + PE: diabetes mellitus with superimposed preeclampsia.

^a Represents p-values <0.05 (comparing, where relevant, AA and no AA groups within the specific pregnancy group or within the total group).

vacuum sample may also be more unbiased, being derived from the entire placental bed surface, rather than the restricted numbers of random tissue biopsies. Our rates of decidual AA are therefore likely to be more accurate than those derived from placental bed [17] or basal plate biopsies [4]. However the samples in this study were derived only from cesarean delivered women who may not be fully representative of all delivered women of our included pregnancy groups. AA lesions in decidua basalis may also differ from those in myometrial segments, where the arteries are known to retain more elements of their original structure (including vascular smooth muscle, elastic laminae and intima). AA lesions in the decidual

basalis may also differ from atherosclerosis [9] and atherosclerosis-like [3,30] lesions previously described in the decidua parietalis.

Presence of foam cells was our minimum requirement for identifying AA. The present study has set a simple criterion for recognising them in relation to arteries of a defined size and has shown that this criterion is reliably reproducible by different observers. Only arteries with a diameter ≥ 140 μ m were examined so as to exclude smaller basal arteries (<100 μ m in diameter). Basal arteries do not perfuse the intervillous space and their remodelling by trophoblasts has not been described [23,31]. Foam cells have been identified in basal arteries-by Sheppard and Bonnar [32],

however the vessel calibre was not documented. In our study, including the smaller arteries of $<140\ \mu\text{m}$, would not have altered the AA rates, as mural foam cells were not identified in such arteries, without foam cells also being identified in the same samples within arteries $>140\ \mu\text{m}$.

The two other classical morphological features of AA, perivascular infiltrates and fibrinoid necrosis, presented problems. PVI, as defined, did not specifically colocalise with foam cells. Identifying fibrinoid necrosis in spiral arteries of the decidua basalis artery wall is also a problem, as smooth muscle vascular wall cells in the decidua portions of the artery are expected to be replaced by fibrinoid following placentation. We found that it was not possible to distinguish fibrinoid deposition from fibrinoid necrosis in decidua basalis spiral arteries by grading the colour and intensity of PAS staining. Even if this distinction can be made by expert perinatal pathologists [33], there are no validated, objective criteria which can be applied to standardise its measurement and allow different studies to be compared. In this study we could only comment on the presence of fibrinoid, but not its etiology or degree of “necrosis”. Instead we quantified fibrinoid in the arterial wall by a new method of grading the proportion of the arterial vessel circumference which demonstrated PAS staining. Although fibrinoid did not colocalise with foam cell lesions, we believe that its presence in the spiral arterial wall warrants further investigation as it is not well described in the literature. This should include better identification of the molecular composition of “fibrinoid necrosis” as described by placental pathologists [33]. We suggest that the presence of fibrinoid and its composition may still be central features of spiral artery physiology and pathology. Its relation to acute atherosclerosis needs further investigation in decidua and myometrial segments of the spiral arteries. As for PVI, it did not colocalise with foam cells. On the other hand, both PVI and fibrinoid did occur significantly more often in preeclampsia (Table 3).

Per case, more arterial sections showed AA lesions in preeclampsia than in normotensive controls, confirming previous reports that the lesion is more widespread in preeclampsia [2,34,35]. However, in keeping with Khong et al. [36], we could not confirm previous reports that a diagnosis of AA is more associated with severe than moderate preeclampsia [3,30,37,38], neither in terms of preterm vs term delivery nor in terms of maximum systolic blood pressures.

AA was not confined to preeclampsia but occurred in normotensive and diabetic women without preeclampsia, as we - and others - have shown previously [2,13]. The large variations in rates of AA lesions reported in preeclampsia (4–50%) [3,4,39] may in part be due to different diagnostic criteria, sampling methods or patient selection. Our AA rate of 37% for preeclampsia sample in this study is consistent with that of 40% in our earlier study, which included different decidua sections from some of the same patients, but from fewer patients and using different diagnostic criteria [2]. Our diabetic group is small and the subgroups are heterogeneous in both etiology and severity. We did not find increased rates of decidua basalis AA lesions in any of the diabetic subgroups, as compared to controls, which suggests that the metabolic disturbances associated with the different types of diabetes do not increase the rates of AA. It is interesting that we only found increased AA lesion rate as compared to normotensive in the diabetic pregnancies that had developed superimposed preeclampsia. These issues warrant further and more detailed investigation. Whether acute atherosclerosis in relation to pregestational and gestational diabetes mellitus is an important risk for future cardiovascular disease [1] certainly deserves attention.

AA was significantly associated with spiral artery thrombosis, as previously reported [11], for both preeclampsia and normotensive control groups. Our rate of spiral artery thrombosis seems high.

Such thrombi are unlikely to lead to massive downstream placental infarcts, as more than one spiral artery supplies one cotyledon. Our finding of a significant difference in thrombosis rates between pregnancy groups, which depends on the presence or absence of acute atherosclerosis, is an important observation, for which the mechanisms need further research.

The significantly higher rate of AA in primigravid women, who by definition never have experienced decidual trophoblast invasion, was interesting, as was that of AA being more prevalent in parous (currently normotensive) women with a history of preeclampsia. The association of primigravida with AA confirms previous observations also made by Zhang et al. [39]. A first pregnancy is also associated with increased risk of preeclampsia that has high rates of AA, also in parous women. A first pregnancy may present problems of maternal immune adaptation to the genetically foreign fetus and we speculate that these may contribute to the development of both AA as well as the development of preeclampsia [5]. A possible mechanism may arise from abnormal remodelling of spiral arteries by endovascular trophoblasts. This may also relate to our findings of significantly reduced numbers, or absence of, mural trophoblasts in arteries diagnosed with AA, a finding previously described by Stallmach [40].

In conclusion: for the first time evidence based criteria for histopathological definition of decidual AA have been developed. Other important features of this study include: unbiased tissue sampling using the vacuum suction technique; measuring the reproducibility of the new criteria and developing the criteria without the assumption that AA is necessarily specific for any clinical diagnosis. It will make spiral artery research easier and allow better comparison between different studies. We will use this stringent definition in a future study to test the hypothesis that AA could be a marker for future atherosclerotic cardiovascular disease [41].

Conflict of interest statement

The authors declare that they do not have any conflict of interest.

Funding

The Research Council of Norway funded part of this work (the PATH study: Pregnancy acute atherosclerosis and future cardiovascular disease, grant number ref. 230652) as did South-Eastern Norway Regional Health Authority (the HAPPY PATH study, grant number ref. 2014026). Oslo University Hospital provided PhD salary (PAK) and further research support.

Acknowledgements

The authors greatly acknowledge the assistance from pathology assistants Linn Buer, Lene Nordenstrøm and Giang Huong Nguyen (Dept of Pathology, Oslo University Hospital). The authors are very grateful for the laboratory assistance and patient recruitment by Lise Øhra Levy (Oslo University Hospital), as well as for patient recruitment to Oslo Pregnancy Biobank (Oslo University Hospital) by previous PhD students, including Kristin Brække, Nina K. Harssem and Meryam Sugulle.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.placenta.2015.10.020>.

References

- [1] A.C. Staff, C.W. Redman, IFPA Award in Placentology Lecture: preeclampsia, the decidual battleground and future maternal cardiovascular disease, *Placenta* 35 (Suppl:S26–S31) (2014 Feb).
- [2] N.K. Harsem, B. Roald, K. Braekke, A.C. Staff, Acute atherosclerosis in decidual tissue: not associated with systemic oxidative stress in preeclampsia, *Placenta* 28 (8–9) (2007 Aug) 958–964.
- [3] D.U. Stevens, S. Al-Nasiry, J. Bulten, M.E. Spaanderman, Decidual vasculopathy and adverse perinatal outcome in preeclamptic pregnancy, *Placenta* 33 (8) (2012 Aug) 630–633.
- [4] Y.M. Kim, P. Chaemsaitong, R. Romero, M. Shaman, C.J. Kim, J.S. Kim, et al., Placental lesions associated with acute atherosclerosis, *J. Matern. Fetal Neonatal Med.* (2014 Sep 3) 1–29.
- [5] A.C. Staff, G.M. Johnsen, R. Dechend, C.W. Redman, Preeclampsia and uteroplacental acute atherosclerosis: immune and inflammatory factors, *J. Reprod. Immunol.* 101–102 (2014 Mar) 120–126.
- [6] C.A. Labarrere, Acute atherosclerosis. A histopathological hallmark of immune aggression? *Placenta* 9 (1) (1988 Jan) 95–108.
- [7] H.C. Stary, A.B. Chandler, S. Glagov, J.R. Guyton, W. Insull Jr., M.E. Rosenfeld, et al., A definition of initial, fatty streak, and intermediate lesions of atherosclerosis. A report from the committee on vascular lesions of the council on arteriosclerosis, American heart association, *Circulation* 89 (5) (1994 May) 2462–2478.
- [8] A.T. Hertig, Vascular pathology in hypertensive albuminuric toxemias of pregnancy, *Clinics* 4 (1945) 602–614.
- [9] P.M. Zeek, N.S. Assali, Vascular changes in the decidua associated with eclamptogenic toxemia of pregnancy, *Am. J. Clin. Pathol.* 20 (12) (1950 Dec) 1099–1109.
- [10] W.B. Robertson, I. Brosens, H.G. Dixon, The pathological response of the vessels of the placental bed to hypertensive pregnancy, *J. Pathol. Bacteriol.* 93 (1967) 581–592.
- [11] M. Maqueo, J. Chavezazuela, M. Dosaldelavega, Placental pathology in eclampsia and preeclampsia, *Obstet. Gynecol.* 24 (1964 Sep) 350–356.
- [12] J.W. Meekins, R. Pijnenborg, M. Hanssens, A.A. van, I.R. McFadyen, Immunohistochemical detection of lipoprotein(a) in the wall of placental bed spiral arteries in normal and severe preeclamptic pregnancies, *Placenta* 15 (5) (1994 Jul) 511–524.
- [13] H. Katabuchi, S. Yih, T. Ohba, K. Matsui, K. Takahashi, M. Takeya, et al., Characterization of macrophages in the decidual atherotic spiral artery with special reference to the cytology of foam cells, *Med. Electron Microsc.* 36 (4) (2003 Dec) 253–262.
- [14] F. De Wolf, W.B. Robertson, I. Brosens, The ultrastructure of acute atherosclerosis in hypertensive pregnancy, *Am. J. Obstet. Gynecol.* 123 (2) (1975 Sep 15) 164–174.
- [15] J.W. Meekins, R. Pijnenborg, M. Hanssens, I.R. McFadyen, van AA, A study of placental bed spiral arteries and trophoblast invasion in normal and severe pre-eclamptic pregnancies, *Br. J. Obstet. Gynaecol.* 101 (8) (1994 Aug) 669–674.
- [16] W.B. Robertson, I. Brosens, G. Dixon, Uteroplacental vascular pathology, *Eur. J. Obstet. Gynecol. Reprod. Biol.* 5 (1–2) (1975) 47–65.
- [17] N.K. Harsem, A.C. Staff, L. He, B. Roald, The decidual suction method: a new way of collecting decidual tissue for functional and morphological studies, *Acta Obstet. Gynecol. Scand.* 83 (8) (2004 Aug) 724–730.
- [18] J.M. Roberts, G. Pearson, J. Cutler, M. Lindheimer, Summary of the NHLBI working group on research on hypertension during pregnancy, *Hypertension* 41 (3) (2003 Mar) 437–445.
- [19] WHO, IDF (Ed.), Definition and Diagnosis of Diabetes Mellitus and Intermediate Hyperglycemia: Report of a WHO/IDF Consultation, WHO Press, Geneva, Switzerland, 2006, pp. 1–50 (WHO Library Cataloguing-in-Publication Data).
- [20] S.L. Johnsen, S. Rasmussen, T. Wilsgaard, R. Sollien, T. Kiserud, Longitudinal reference ranges for estimated fetal weight, *Acta Obstet. Gynecol. Scand.* 85 (3) (2006) 286–297.
- [21] A.C. Staff, T. Ranheim, J. Khoury, T. Henriksen, Increased contents of phospholipids, cholesterol, and lipid peroxides in decidua basalis in women with preeclampsia, *Am. J. Obstet. Gynecol.* 180 (3 Pt 1) (1999 Mar) 587–592.
- [22] J.F. McManus, Histological and histochemical uses of periodic acid, *Stain Technol.* 23 (3) (1948 Jul) 99–108.
- [23] R. Pijnenborg, E. Ball, J.N. Bulmer, M. Hanssens, S.C. Robson, L. Vercruysse, In vivo analysis of trophoblast cell invasion in the human, *Methods Mol. Med.* 122 (2006) 11–44.
- [24] W.J. de Villiers, E.J. Smart, Macrophage scavenger receptors and foam cell formation, *J. Leukoc. Biol.* 66 (5) (1999 Nov) 740–746.
- [25] I. Brosens, R. Pijnenborg, L. Vercruysse, R. Romero, The “Great Obstetrical Syndromes” are associated with disorders of deep placentation, *Am. J. Obstet. Gynecol.* 204 (3) (2011 Mar) 193–201.
- [26] D.U. Stevens, S. Al-Nasiry, J. Bulten, M.E. Spaanderman, Decidual vasculopathy in preeclampsia: lesion characteristics relate to disease severity and perinatal outcome, *Placenta* 34 (9) (2013 Sep) 805–809.
- [27] A.J. Viera, J.M. Garrett, Understanding interobserver agreement: the kappa statistic, *Fam. Med.* 37 (5) (2005 May) 360–363.
- [28] Hypertension in pregnancy, Report of the American college of obstetricians and gynecologists’ task force on hypertension in pregnancy, *Obstet. Gynecol.* 122 (5) (2013 Nov) 1122–1131.
- [29] R. Pijnenborg, L. Vercruysse, M. Hanssens, The uterine spiral arteries in human pregnancy: facts and controversies, *Placenta* 27 (9–10) (2006 Sep) 939–958.
- [30] Y.M. Kim, P. Chaemsaitong, R. Romero, M. Shaman, C.J. Kim, J.S. Kim, et al., The frequency of acute atherosclerosis in normal pregnancy and preterm labor, preeclampsia, small-for-gestational age, fetal death and midtrimester spontaneous abortion, *J. Matern. Fetal Neonatal Med.* (2014 Nov 11) 1–9.
- [31] E.M.D.M. Ramsey, Placental Vasculature and Circulation. Anatomy, Physiology, Radiology, Clinical Aspects Atlas Textbook, Georg Thieme Publishers, Stuttgart, 1980.
- [32] B.L. Sheppard, J. Bonnar, The ultrastructure of the arterial supply of the human placenta in early and late pregnancy, *J. Obstet. Gynaecol. Br. Commonw.* 81 (7) (1974 Jul) 497–511.
- [33] R.W. Redline, T. Boyd, V. Campbell, S. Hyde, C. Kaplan, T.Y. Khong, et al., Maternal vascular underperfusion: nosology and reproducibility of placental reaction patterns, *Pediatr. Dev. Pathol.* 7 (3) (2004 May) 237–249.
- [34] T.Y. Khong, Acute atherosclerosis in pregnancies complicated by hypertension, small-for-gestational-age infants, and diabetes mellitus, *Arch. Pathol. Lab. Med.* 115 (7) (1991 Jul) 722–725.
- [35] B.L. Sheppard, J. Bonnar, An ultrastructural study of utero-placental spiral arteries in hypertensive and normotensive pregnancy and fetal growth retardation, *Br. J. Obstet. Gynaecol.* 88 (7) (1981 Jul) 695–705.
- [36] T.Y. Khong, J.M. Pearce, W.B. Robertson, Acute atherosclerosis in preeclampsia: maternal determinants and fetal outcome in the presence of the lesion, *Am. J. Obstet. Gynecol.* 157 (2) (1987 Aug) 360–363.
- [37] W.T. Parks, Placental hypoxia: the lesions of maternal malperfusion, *Semin. Perinatol.* 39 (1) (2015 Feb) 9–19.
- [38] K. Baltajian, J.L. Hecht, J.B. Wenger, S. Salahuddin, S. Verlohren, F.H. Perschel, et al., Placental lesions of vascular insufficiency are associated with antiangiogenic state in women with preeclampsia, *Hypertens. Pregnancy* 33 (4) (2014 Nov) 427–439.
- [39] P. Zhang, M. Schmidt, L. Cook, Maternal vasculopathy and histologic diagnosis of preeclampsia: poor correlation of histologic changes and clinical manifestation, *Am. J. Obstet. Gynecol.* 194 (4) (2006 Apr) 1050–1056.
- [40] T. Stallmach, G. Hebisch, P. Orban, X. Lu, Aberrant positioning of trophoblast and lymphocytes in the feto-maternal interface with pre-eclampsia, *Virchows Arch.* 434 (3) (1999 Mar) 207–211.
- [41] A.C. Staff, R. Dechend, C.W. Redman, Review: Preeclampsia, acute atherosclerosis of the spiral arteries and future cardiovascular disease: two new hypotheses, *Placenta* 34 (Suppl:S73–S78) (2013 Mar).

Paper I (erratum)

Corrigendum to “Acute atherosclerosis in vacuum suction biopsies of decidua basalis: An evidence based research definition” [Placenta 37C (2016) 26–33]

Patji Alnaes-Katjavivi^a, Fiona Lyall^b, Borghild Roald^c, Christopher W.G. Redman^d, Anne Cathrine Staff^{a,*}

^a Department of Obstetrics and Department of Gynecology, Oslo University Hospital, Ullevål and Faculty of Medicine, University of Oslo, Norway

^b University of Glasgow, School of Medicine, Glasgow, UK

^c Department of Pathology, Oslo University Hospital, Ullevål and Faculty of Medicine, University of Oslo, Norway

^d Department of Obstetric Medicine, University of Oxford, Oxford, UK

The authors regret that some of the column data entries were incorrectly placed according to pregnancy groups. Pasted below is the table as it appears published in Placenta, followed by the same table with corrections highlighted in grey.

P. Alnaes-Katjavivi et al. / Placenta 37 (2016) 26–33

31

Table 4

Clinical features according to presence or absence of acute atherosclerosis (AA) for each of the pregnancy groups (and the total study group). AA is defined according to the simplified evidence-based criterion of intramural foam cell lesions (defined as ≥ 2 adjacent intramural vacuolated and CD68 positive cells). Values are expressed as medians, percentages, and rates.

Acute atherosclerosis (AA) (No AA or AA diagnosis)	Normotensive controls n = 94		PE n = 87		DM n = 39		DM + PE n = 17		Total n = 237	
	No AA n = 84 (89%)	AA n = 10 (11%)	No AA n = 55 (63%)	AA n = 32 (37%)	No AA n = 35 (90%)	AA n = 4 (10%)	No AA n = 12 (71%)	AA n = 5 (29%)	No AA n = 186 (78%)	AA n = 51 (22%)
Age at delivery (years)	34	33	32	32	33	31	33	31	33	32
First pregnancy (gravida 1)	26%	50%	51%	47%	17%	50%	42%	43%	33%	51% ^a
BMI: pre-pregnancy (kg/m ²)	22.1	22.0	23.1	24.3	24.3	27.6	24.4	27.7	22.7	24.6
BMI: at delivery (kg/m ²)	27.2	27.5	30.1	29.7	33.3	29.6	31.5	36.7	29.0	29.3
^b Median Systolic BP < 20weeks (mmHg)	108	110	119	115	117	115	130	119	117	113 ^a
^b Median Diastolic BP < 20weeks (mmHg)	67	65	72	71	72	71	75	70	72	70
BP at delivery (mmHg) (Systolic BP/Diastolic BP)	120/71	119/70	162/100	160/104	135/79	132/77	160/95	162/100	138/80	158 ^a /100 ^a
Gestational age at delivery (weeks ⁺ days)	38 ⁺⁵	38 ⁺⁵	33 ⁺⁵	32 ⁺⁴	38 ⁺⁰	37 ⁺²	34 ⁺⁴	36 ⁺⁵	38 ⁺⁰	34 ^{+0a}
Delivery <37 ⁺⁰	2%	10%	87%	91%	20%	25%	92%	60%	37%	67% ^a
Delivery <34 ⁺⁰	1%	0%	55%	69%	3%	25%	71%	40%	26%	51% ^a
Birth percentile	66%	50%	6%	2%	98%	55% ^a	62%	66%	55%	16% ^a
Small for gestational age (birthweight $\leq 10\%$)	2%	0%	58%	69%	3%	25%	25%	0%	20%	45% ^a
Small for gestational age (birthweight $\leq 3\%$)	0%	0%	42%	59%	3%	25%	17%	0%	14%	39% ^a
Previous PE (amongst parous women)	0% (0/44)	67% ^a (2/3)	33% (6/18)	46% (6/13)	16% (4/25)	0% (0/4)	67% (2/3)	100% (1/1)	13% (12/90)	43% ^a (9/21)
Severe PE according to ACOG (28)	0%	0%	82%	78%	0%	0%	67%	75%	29%	54% ^a

PE: preeclampsia, DM: diabetes mellitus (all types), DM + PE: diabetes mellitus with superimposed preeclampsia, BMI: body mass index.

BP: blood pressure.

^a p-values <0.05 when comparing the variable investigated (e.g. rate of primiparity) for the “AA present” and “no AA” groups within the specific pregnancy group (e.g. PE or DM) or within the total group (last column: comparing all women with AA to all women without AA; independent of clinical group).

^b Presents the median blood pressure (systolic or diastolic) value for the specific pregnancy group before gestation week 20 (the mean per woman is first calculated of all blood pressures registered per woman before gestation week 20).

DOI of original article: <http://dx.doi.org/10.1016/j.placenta.2015.10.020>.

* Corresponding author.

E-mail address: uxnnaf@ous-hf.no (A.C. Staff).

<http://dx.doi.org/10.1016/j.placenta.2017.03.002>

0143-4004/© 2017 Elsevier Ltd. All rights reserved.

Below is a copy of the Table 4 above with the corrected values highlighted in grey. The corrected values do not entail any alterations to the statistical significant values presented. The only relevant change in the data presentation is that, compared to those without, women with AA have slightly higher systolic blood pressure (BP) in the first half of pregnancy (median value for mean blood pressure taken prior to 20 weeks of gestation).

These changes for Table 4 have no bearing on the text contained in the article.

The replacement table.

Table 4. Clinical features according to presence or absence of acute atherosclerosis (AA) for each of the pregnancy groups (and the total study group). AA is defined according to the simplified evidence-based criterion of intramural foam cell lesions (defined as ≥ 2 adjacent intramural vacuolated and CD68 positive cells). Values are expressed as medians, percentages, and rates.

*Represents p-values <0.05 (comparing AA and no AA groups within the specific pregnancy group or within the total group).

Acute atherosclerosis (AA) (No AA or AA diagnosis)	Normotensive controls n = 94		PE n = 87		DM n = 39		DM+PE n = 17		Total n = 237	
	No AA n = 84 (89%)	AA n=10 (11%)	No AA n = 55 (63%)	AA n=32 (37%)	No AA n = 35 (90%)	AA n=4 (10%)	No AA n = 12 (71%)	AA n=5 (29%)	No AA n = 186 (78%)	AA n=51 (22%)
Age at delivery (years)	33	34	32	32	33	31	33	31	33	32
First pregnancy (gravida 1)	26%	50%	51%	47%	17%	50%	42%	80%	33%	51%*
BMI: pre-pregnancy (kg/m ²)	22.1	22.0	23.1	24.3	27.6	25.4	24.4	27.7	22.7	24.6
BMI: at delivery (kg/m ²)	27.2	27.5	30.1	29.7	33.3	29.6	31.5	36.7	29.0	29.3
Mean ^a Systolic BP < 20weeks (mmHg)	110	108	115	119	115	117	119	130	113	117*
Mean ^a Diastolic BP < 20weeks (mmHg)	65	67	72	72	71	72	70	75	70	72
BP at delivery (mmHg) (Systolic BP/Diastolic BP)	120/71	119/70	162/100	160/104	135/79	132/77	163/95	160/100	138/80	158*/100*
Gestational age at delivery (weeks+days)	38 ⁺⁵	38 ⁺⁵	33 ⁺⁵	32 ⁺⁴	38 ⁺¹	37 ⁺²	34 ⁺⁴	36 ⁺⁵	38 ⁺¹	34 ^{+0*}
Delivery <37 ⁺⁰	2%	10%	87%	91%	20%	50%	92%	60%	35%	69%*
Delivery <34 ⁺⁰	1%	0%	59%	69%	3%	25%	42%	40%	15%	49%*
Birth percentile	66%	50%	6%	2%	98%	55%*	62%	66%	55%	16%*
Small for gestational age (BW = 10%)	2%	0%	60%	66%	3%	25%	25%	0%	20%	45%*
Small for gestational age (BW = 3%)	0%	0%	40%	53%	3%	25%	17%	0%	14%	37%*
Previous PE (amongst parous women)	0%	67%*	33%	46%	16%	0%	67%	100%	13%	43%*
	(0/44)	(2/3)	(6/18)	(6/13)	(4/25)	(0/4)	(2/3)	(1/1)	(12/90)	(9/21)
Severe PE according to ACOG (28)	0%	0%	82%	78%	0%	0%	67%	75%	29%	54%*

PE: preeclampsia, DM: diabetes mellitus (all types), DM+PE: diabetes mellitus with superimposed preeclampsia, BMI: body mass index, BP: blood pressure, BW: birth weight.

^aThese values represent the median of the mean of all systolic/diastolic blood pressure (BP) readings registered before gestation week 20 per woman.

The authors would like to apologise for any inconvenience caused.

Uteroplacental acute atherosclerosis in preeclamptic pregnancies: rates and clinical outcomes differ by tissue collection methods

Alnaes-Katjavivi P.H.^{a}, Roald B.^b, Staff A.C.^a*

^aDivision of Obstetrics and Gynaecology, Oslo University Hospital, Ullevål and Faculty of Medicine, University of Oslo, Norway, ^bDepartment of Pathology, Oslo University Hospital, Ullevål and Faculty of Medicine, University of Oslo, Norway,

*Corresponding author: Patji Alnaes-Katjavivi MB BS

Division of Obstetrics and Gynaecology, Oslo University Hospital, Ullevål, PO Box 4956 Nydalen, NO-0424 Oslo, Norway

Business telephone: + 47- 22 11 98 00, Personal telephone number: + 47 98401818

E-mail: alkp@ous-hf.no

Abstract

Acute atherosclerosis (AA) is a uteroplacental spiral artery lesion, identified by intramural lipid-laden foam cells. This study compares AA detection rates in preeclampsia (PE) across three different decidual spiral artery collection methods that were performed in 107 preeclamptic women. Routine sampled basal surface placenta (decidua basalis, DB) and fetal membrane roll (decidua parietalis, DP) biopsies were compared with decidual vacuum suction biopsies (DB) measuring spiral artery rate and AA detection, and whether AA presence associated with PE-related clinical parameters. Spiral arteries and AA were identified using predefined criteria, utilizing IHC on serial sections. Detection of spiral arteries (87%) and AA rate (35%) was highest in DB samples collected by vacuum suction compared to the two other methods. Pregnancies with AA detected in vacuum suctioned DB had lower gestational age at delivery, lower birth weight percentile and more often fetal growth restriction. Basal plate DB samples demonstrating AA associated with pregnancies affected by pathological fetal Dopplers, whereas AA detected in DP membrane rolls did not. We conclude that AA associates with clinical parameters related to placental dysfunction rather than to prepregnancy maternal risk factors for preeclampsia. As collection of DB tissue by placental bed vacuum suction provides the highest yield of arteries and AA, underestimation of AA is possible when limited to routine clinical sections of basal plate and membrane roll. For the first time, rate of decidual artery AA-detection in the same patient is compared using the decidual suction collection method of DB versus routine sampling of DB and DP.

Keywords:

Placenta, decidua, acute atherosclerosis, morphology, immunohistochemistry, preeclampsia

Background

Acute atherosclerosis (AA) is a histological lesion of uteroplacental arteries first characterized by Hertig in 1945 [1]. AA is found in both myometrial [2] and decidual parts [1, 3] of the spiral arteries.

AA is most commonly associated with preeclamptic (PE) pregnancies [4], with rates varying between studies, from 10% to 38-41% [5-7]. This variation may be due to different tissue collection

methods, heterogeneous patient groups, as well as lack of uniform and clear criteria for AA diagnosis. AA remains a lesion of unclear etiology and clinical significance, although its association to downstream placenta lesions is documented [3].

As well as being associated with pregnancy complications (e.g. placental insufficiency, placental abruption, eclampsia), preeclampsia is also associated with health complications later in life, in

particular morbidity from hypertension and atherosclerosis [8-10]. Acute atherosclerosis bears some morphological similarities to early stage atherosclerosis [11], and our group has proposed that AA could represent an accelerated form of vasculopathy, developing during the short time of pregnancy at the fetal-maternal interface, and arising from local excessive inflammation as the final common pathway [12].

What remains to be clarified is whether the presence of decidual AA across various sampling sites differs in the association to pregnancy-related morbidity, as well as to the increased long-term maternal cardiovascular complications associated with preeclampsia.

We have previously defined minimum inclusion criteria for a predefined and simple diagnosis of AA, allowing for reliable comparison of AA rates between different sample types, patient groups, and between different investigators [6]. Decidua is the name of the endometrium of pregnancy, and can be found underneath the placenta (decidua basalis) or on the uterine wall outside the placenta (decidua parietalis). Our criteria for AA diagnosis were developed studying vacuum suction samples of decidua basalis tissue (DB) from the placental bed following the removal of the placenta during pre-labor cesarean section [7]. Previously, comparing these tissue collection methods from different pregnancy groups, we have shown that our vacuum suction method provides a higher yield of decidua basalis spiral arteries (present in 86% of tissue sections evaluated), than in decidua basalis derived from placenta bed biopsies and maternal placental plate sections (61% and 48%, respectively [7]). We have therefore suggested that vacuum suction represents a better opportunity to identify acute atherosclerosis lesions than decidual tissue sampled by other methods routinely used in evaluating the placenta postpartum. However, DB tissue sampling by the

vacuum suction method remains clinically less feasible for vaginal deliveries, thereby biasing the AA investigations to pregnancies with clinical indications for a cesarean section.

Most preeclamptic women will deliver vaginally, making vacuum suction of decidua basalis not easily feasible. Consequently, decidual AA detection in placental and fetal membrane samples collected for clinical routine pathology might be more appropriate and feasible in the everyday clinical setting. No previous studies, including our own [7] have, however, compared AA rates for these tissue methods across the same pregnancies.

The aim of this paper was therefore to compare the rates of spiral artery and AA presence across the same cohort of preeclamptic pregnancies, comparing decidua samples from these three collection sites. We hypothesized that the rate of AA differs in the same group of preeclamptic women between these three decidual tissues evaluated, and that the location(s) for where the AA is detected differs in association with severity of PE and placental function.

Method

Patient recruitment, biopsy tissue sampling and preparation

We included 107 preeclamptic women from the Oslo Pregnancy Biobank (2000-2016, www.oslo-universitetssykehus.no/opb) to the present study. Eighty-eight had also participated in a previous decidua basalis (vacuum suction) acute atherosclerosis study [6]. Preeclampsia (PE) was defined as new onset hypertension (blood pressure ≥ 140 and/or ≥ 90 mmHg) and new onset proteinuria ($\geq 1+$ on dipstick, or ≥ 30 protein/creatinine ratio) ≥ 20 weeks of pregnancy [13]. None of the women participating in the present study had any known chronic disease. Gestational week (GW) at delivery was defined by routine ultrasound screening at GW 17-20.

Newborn gender specific baby weight percentiles were calculated according to Norwegian fetal growth curves [14]. Pathological fetal Dopplers were defined as abnormal PI in umbilical artery and/or middle cerebral artery ($>2SD$), and/or cerebroplacental ratio <1.08 [15, 16]. All women provided informed written consent. The study was approved by the Regional Committee for Medical and Health Research Ethics.

Decidual tissue was sampled following delivery by pre-labor cesarean section [6], with intact membranes. Fig. 1 illustrates the three different sampling sites for decidual tissue in all 107 patients. Decidua basalis (DB) tissue from the placental bed was collected both by the decidual vacuum suction method (DSM) developed by our group [7, 17], and by routine basal plate placenta section (BPS) excised from macroscopically normal looking maternal plate placental tissue. From the same deliveries, decidua parietalis (DP) was collected through routinely prepared placental fetal membrane rolls (MR). Both BPS and MR were prepared by the same pathology technician and according to the standardized routine for our University Hospital's Department of Pathology. All samples were fixated in 4% buffered formaldehyde and preserved in paraffin blocks from which slides were prepared [18].

Decidual spiral artery identification by immunohistochemistry (IHC)

For each pregnancy ($n=107$), slides from each of the three tissue types were firstly studied microscopically in H&E stained sections for identification of spiral arteries, followed by the identification of mural foam cells (acute atherosclerosis). The stains and antibodies for IHC are listed in Supplementary material Table 1. Serial slides ($3\mu m$ thick) were prepared from each biopsy tissue-type, minimizing alteration in tissue topography between the first H&E stained slides (used for initial

identification of arteries and vacuolated arterial wall cells), and the subsequent immunostaining of slides with CD68 (used to confirm presence of lipid scavenger receptor, typically of macrophages and foam cells). Thus, at least two serial decidual slides per pregnancy were evaluated to confirm the presence or absence of spiral arteries and, thereafter, the presence or absence of acute atherosclerosis, for each of the three tissue sampling methods.

Spiral arteries were detected first on H&E-stained slides according to morphological features, assisted – if necessary – by identification of intramural smooth muscle (Desmin, Clone DE R-11, Roche Ventana Medical system Inc, Arizona, USA, readymade), or its replacement by fibrinoid change (intramural PAS-positive staining detected beyond the normal PAS staining of subendothelial basement, see Fig. 2) [19].

For decidua basalis samples (either collected by vacuum suction or from the placental basal plate sections), spiral arteries were considered representative if $\geq 140\mu m$ in diameter, avoiding inclusion of the smaller decidual nutrient (“straight”) arteries that do not undergo trophoblast remodeling (typically $<100\mu m$ in diameter) [20, 21]. Inter- and intra-observer scoring for AA agreement between authors in decidua basalis tissue, according to our predefined criteria, has been shown previously to be substantial [6].

Spiral arteries belonging to the decidua parietalis lie beyond the area of endovascular trophoblast invasion and therefore are not expected to demonstrate features of trophoblast-assisted arterial remodeling [22]. Hence, for the biopsy tissues of placental fetal membrane rolls, smaller arteries of $<140\mu m$ in diameter were also included in our evaluation for AA in the decidua parietalis.

Acute atherosclerosis was identified as previously [6], where ≥ 2 adjacent vacuolated intramural cells visualized by H&E were confirmed as foam cells by

positive immunostaining with CD68 on serial slides (Clone KP1, Dako AS, Glostrup, Denmark, dilution 1/1000, for the detection of the scavenger receptor expressed by foam cells and macrophages). Representative images for normal and AA-affected spiral arteries are demonstrated, by tissue-type category, in Fig. 2.

Rates of AA detection per tissue type sampled were based on samples where at least one artery section with AA was identified.

Statistical analysis

Data were analyzed using SPSS version 22. The Chi squared test was used for dichotomous variables, and the Mann-Whitney-Wilcoxon test for continuous variables. Associations between AA presence per tissue category and clinical outcome were tested using Spearman correlation and bivariate logistic regression. A probability of ≤ 0.05 was considered statistically significant.

Results

Table 1 presents spiral artery and AA detection rates for the three uteroplacental tissue types.

In five women no arteries were detectable in any of their three decidual tissue samples. Of the 107 preeclamptic women, spiral arteries were thus detected in 102 of the 107 preeclamptic women, with acute atherosclerosis detected in 36% (37 of 102), when all three tissue types sampled were combined for evaluation of AA.

Decidua basalis sections from vacuum tissue sampling detected the highest rate of spiral arteries (DSM: at least one spiral artery section detected in 87% of sections), followed by decidua parietalis tissue sections from fetal membranes (MR: 71%). These were significantly higher than decidual spiral artery detection rate in decidua basalis tissue from placental basal plate sections (BPS: 50%), all $p < 0.05$ (Table 1). In addition to identifying the highest rate of spiral arteries, vacuum collection samples

of decidua basalis also identified the significantly highest rate of AA (Table 1. DSM: 33 of 94; 35%), supporting our previous conclusions of this sample type representing the superior method for decidua basalis spiral artery AA investigation [7]. Combining DSM with MR tissue increased both the rate of women where decidual spiral arteries are detected, and the rate of women where AA is detected (Table 1).

There is overlap in AA detection between the three tissue collection methods for the 102 preeclamptic women where spiral arteries were found. Demonstrated in Supplementary material Fig.1, sampling from 37 women showed acute atherosclerosis detected in the spiral artery wall(s) of at least one tissue sample type. Among the 37 PE women with AA, only one woman had AA lesions present in all three decidual samples types. All 8 women with AA presence in basal plate sections also had AA in their decidua basalis vacuum suction sample. However, the basal plate sections missed identifying 25 of the AA lesions present in the decidua vacuum suction collected samples.

Despite a satisfactory rate of decidua parietalis spiral artery detection in the placental fetal membrane roll (MR) samples (one or more arteries detected in 76 of 107 samples = 71%), the AA rate in the MR was only 11% (one or more AA affected artery in 8 of 76 samples, Table 1). In four women, AA was detected in MR tissue alone (Supplementary material Fig.1). Taking membrane samples as a whole, 80% (317 out of 337) of arteries counted had diameters of $< 100\mu\text{m}$, in keeping with the arteries not having undergone trophoblast remodeling (data not shown).

Most of MR arteries containing AA were $> 100\mu\text{m}$ (in 7 of 8 women, and in 15 of 20 arteries counted containing AA), as opposed to only 17% (53 of 317) of MR arteries without AA, counted in the total of decidua parietalis samples (data not shown). Irrespective of AA, 69% (47 of

68) of MR artery sections with a diameter $>100\mu\text{m}$ were located proximal to the placental margin (within the first complete “turn” of the membrane roll, data not shown). None of the MR arterial walls contained endovascular trophoblasts.

Association of AA with clinical pregnancy parameters, according to tissue-type

Table 2 presents the clinical parameters according to the detection of AA in one or more of the three tissue types evaluated. When combining samples from all three tissue types for each woman (DSM+BPS+MR), AA presence was significantly associated with a lower gestational age at delivery ($p=0.007$), and with a lower birthweight percentile ($p=0.034$), compared to those women in whom tissue arteries were detected, but where AA was absent (Fig. 3a). When evaluated separately, only presence of AA in DSM samples of decidua basalis associated significantly with lower gestational age at delivery and lower birthweight percentile ($p=0.025$ and $p=0.039$ respectively, Fig. 3b). When combining decidua basalis and parietalis tissue types collected from the two tissue types utilized in routine clinical histopathology (BPS+MR), presence of AA associated with a significantly lower gestational age at delivery (but not birthweight percentile), compared to women where AA was absent ($p=0.025$, Fig. 3c).

Rates of fetal growth restriction (FGR; defined as a birth weight percentile below 3% and/or antenatal pathological fetal Dopplers) were high in pregnancies with AA presence either in BPS (100%) and MR (75%), but not significantly higher compared to samples where AA is absent in either BPS or MR (or combination thereof). Only in DSM decidua basalis samples was FGR significantly associated with AA presence (AA present vs AA absent: 79% vs 40% FGR, $p=0.0004$, Table 2).

Pathological fetal Dopplers antenatally, regardless of birth weight percentile, was significantly more frequent in preeclamptic women with AA in combined BPS and MR as compared to lack of AA in these tissues (53% vs 23% FGR $p=0.0279$, Table 2), but not in DSM (39% vs 20% FGR, $p=0.088$). In subanalyses, this decidual AA and pathological fetal Dopplers association was only valid for BPS decidua basalis AA ($p=0.0328$), not for MR decidua parietalis AA (data not shown).

We found no significant associations between AA presence (identified in either single or combinations of decidual tissue collection methods) and other clinical parameters associated with preeclampsia risk, such as maternal age, maternal BMI, mean blood pressure in the first half of pregnancy, infant sex, primigravidity, or previous history of PE.

Discussion

Our findings in a relatively large preeclampsia group from one single hospital confirm our previous conclusion, based on comparing different groups of women, namely that the decidua vacuum suction method is the method of choice for the detection of decidual spiral arteries [7], in the absence of available postpartum hysterectomies with placentas still attached to the uterine wall [23]. Compared to our previous study (7), the present study included all three tissue sites from the same large group of women. After comparing tissue from three different collection methods - all obtained from every patient - our conclusion was in line with our previous work demonstrating DSM as superior to BPS and placenta bed biopsies [6], the latter not investigated in the present study of DSM, BPS, and MR. The vacuum suctioned decidua basalis tissue had highest detection rate of AA (37%), as compared to the two other spiral artery sampling methods (basal plate placenta sections and membrane roll sections). The homogeneity and

completeness of the sample tissue types is an advantage of this study, as is the predefined definition for detection of AA in the tissues evaluated, with previously demonstrated satisfactory inter- and intra-observer agreement for AA detection in decidua basalis spiral arteries [6].

Though the detection rate of spiral arteries was lowest in the placental basal plate section method (50%), AA was found in an equal rate of pregnancies from BPS samples (with detected arteries) as with MR samples (8 of 53 compared to 8 of 76). That we have no cases of AA detection exclusive to the basal plate section tissue alone is in keeping with the fact that the basal plate placental section (from the maternal plate of the delivered placenta) reflects the same placental bed (where the remaining decidual tissue after placenta removal is collected by the vacuum suction method). The lower detection rate for AA in samples used for standard clinical pathology alone (BPS and MR) may explain the disparity between AA detection rate reported between research groups examining solely BPS and MR, and the results from our decidua basalis vacuum suction studies.

An 18% AA rate in BPS and MR samples in the current study is closer to the more conservative detection rate of 10% reported by Kim et al in the hitherto largest published study of AA detection using H&E stained BPS and MR tissues. Furthermore, Kim et al suggest a potential additional under-reporting of AA due to not immunostaining decidual sections with CD68, nor the use of oil-red O confirming foam cells [5]. Our MR observations demonstrate characteristic AA lesions detected in predominantly larger ($>100\mu\text{m}$ diameter) spiral arteries, located close to the junction of the fetal membranes and the basal plate, at the margin of the placenta.

Khong et al have demonstrated *en face* basal plate sections as an alternative method of sampling placentas for the assessment of uteroplacental vasculature [24], demonstrating an increased detection

of spiral arteries, albeit AA detection was not quantified. Our decidual sections were taken from only one biopsy, as opposed to possible multiple sampling sites, which would have been desirable for identifying more spiral arteries for AA evaluation. However, as the same method was performed for all three decidual sampling procedures, it is likely that the difference in AA detection rates would remain the same.

Our study showed a significant association between AA presence and three fetoplacental clinical outcomes that are common in early-onset and clinically severe preeclampsia: low gestational age at delivery, low birthweight percentile, and pre-delivery pathological fetal Dopplers. Low birthweight percentile and pathological Dopplers are also features of fetal growth restriction and of placental dysfunction in general. Women with a history of severe preeclampsia, especially when complicated by preterm delivery and/or fetal growth restriction, represent epidemiologically the group with highest risk for premature cardiovascular mortality, up to 8-fold [25]. Another study of decidual AA found a significant association to maternal vascular under perfusion [26], which is in line with our conclusion of decidual AA being associated with placental dysfunction. Our sub analysis showing no association between pathological fetal Dopplers and AA presence in MR decidual tissue alone, is consistent with the findings of Khong et al [27], where AA was not found in MR decidual samples in pregnancies with growth restricted babies [27].

Our group has recently shown that decidual AA is significantly associated with the combination of maternal KIR-B haplotype and fetal HLA-C2 [28], suggestive of an impact of alloimmune interactions at the decidual materno-fetal interface, promoting decidual acute atherosclerosis vascular lesions and affecting placental function. In general, there is an intimate relationship between the immune

and the cardiovascular system, both during [29] and outside pregnancy [30, 31]. We have suggested that AA may serve as a positive screening test for parous women with increased risk for premature CVD [12, 32-34]. This is a new concept being tested by ourselves, which is also gaining interest among other investigators [35]. The American Heart Association has since 2011 included a history of pregnancy hypertension as a major risk factor for cardiovascular disease (CVD) [36]. Our future aim is to investigate whether decidual acute atherosclerosis, either documented in decidua basalis or parietalis, could serve as a personalized targeting of young women at highest future cardiovascular risk, facilitating intensive personalized prevention and follow-up programs. Should such a predictive role of acute atherosclerosis be found, then AA identification by clinical pathologists postpartum could represent an early tool contributing to the improvement of future CVD risk stratification in parous women.

Conclusion

Decidua basalis tissue, obtained using the vacuum suction technique at cesarean section remains the most reliable method for the identification of decidual spiral arteries and subsequent detection of spiral artery acute atherosclerosis, when compared to routine clinical decidual sampling from basal plate placenta sections (decidua basalis) and the membrane roll sections (decidua parietalis). Compared to preeclampsia without identified AA lesions, AA presence in the spiral arteries of decidua basalis was associated with lower gestational age, lower birthweight percentile, and pathological fetal Dopplers. Whether presence and quantity of AA in decidua basalis and/or parietalis identifies preeclamptic women with high risk of

longterm atherosclerotic CVD, remains to be seen.

Acknowledgements:

We would like to thank administrator for the Oslo Pregnancy Biobank, Lise Levy, for assisting with sample procurement and biobank handling. We would also like to thank Linn Buer and Giang Nguyen, Department of Pathology, Oslo University Hospital, for the sectioning and immunostaining of decidual tissue samples. A special thanks to Professor Christopher WG Redman (Oxford, UK) for his guidance in the planning of this study. Lastly, thanks to the support of Lien My Diep, Biostatistician/Statistician, Oslo Centre for Biostatistics and Epidemiology, Oslo University Hospital, Oslo, Norway.

Compliance with Ethical Standards:

This study has the approval of the Regional Committee of Medical Ethics in Norway (Region Eastern Norway) and is conducted in line with the declaration of Helsinki. All participating pregnant women provided informed consent.

Funding:

The Research Council of Norway funded part of this work (the PATH study: Pregnancy acute atherosclerosis and future cardiovascular disease, grant ref. 230652) as did South-Eastern Norway Regional Health Authority (the HAPPY PATH study, grant ref. 2014026). Oslo University Hospital provided PhD salary (PAK) and further research support.

Conflict of interest statement:

The authors have no conflict of interest to declare.

Reference List

1. Hertig AT (1945) Vascular pathology in hypertensive albuminuric toxemias of pregnancy Clinics 4:602-614

2. De Wolf F, Robertson WB, Brosens I (1975) The ultrastructure of acute atherosclerosis in hypertensive pregnancy *Am. J. Obstet. Gynecol* 123:164-174
3. Zeek PM, Assali NS (1950) Vascular changes in the decidua associated with eclamptogenic toxemia of pregnancy *Am. J. Clin. Pathol* 20:1099-1109
4. Stevens DU, Al-Nasiry S, Bulten J, Spaanderman ME (2012) Decidual vasculopathy and adverse perinatal outcome in preeclamptic pregnancy *Placenta* 33:630-633. doi: 10.1016/j.placenta.2012.04.013
5. Kim YM, Chaemsaitong P, Romero R, Shaman M, Kim CJ, Kim JS, Qureshi F, Jacques SM, Ahmed AI, Chaiworapongsa T, Hassan SS, Yeo L, Korzeniewski SJ (2015) The frequency of acute atherosclerosis in normal pregnancy and preterm labor, preeclampsia, small-for-gestational age, fetal death and midtrimester spontaneous abortion *J. Matern. Fetal Neonatal Med.* 28:2001-2009. doi: 10.3109/14767058.2014.976198
6. Alnaes-Katjavivi P, Lyall F, Roald B, Redman CW, Staff AC (2016) Acute atherosclerosis in vacuum suction biopsies of decidua basalis: An evidence based research definition *Placenta* 37:26-33. doi: 10.1016/j.placenta.2015.10.020
7. Harsem NK, Staff AC, He L, Roald B (2004) The decidual suction method: a new way of collecting decidual tissue for functional and morphological studies *Acta Obstet. Gynecol. Scand* 83:724-730
8. Bellamy L, Casas JP, Hingorani AD, Williams DJ (2007) Pre-eclampsia and risk of cardiovascular disease and cancer in later life: systematic review and meta-analysis *BMJ* 335:974. doi: 10.1136/bmj.39335.385301.BE
9. Lykke JA, Langhoff-Roos J, Sibai BM, Funai EF, Triche EW, Paidas MJ (2009) Hypertensive pregnancy disorders and subsequent cardiovascular morbidity and type 2 diabetes mellitus in the mother *Hypertension* 53:944-951
10. Staff AC, Redman CW, Williams D, Leeson P, Moe K, Thilaganathan B, Magnus P, Steegers EA, Tsigas EZ, Ness RB, Myatt L, Poston L, Roberts JM (2016) Pregnancy and Long-Term Maternal Cardiovascular Health: Progress Through Harmonization of Research Cohorts and Biobanks *Hypertension* 67:251-260. doi: 10.1161/hypertensionaha.115.06357
11. Sary HC, Chandler AB, Glagov S, Guyton JR, Insull W, Jr., Rosenfeld ME, Schaffer SA, Schwartz CJ, Wagner WD, Wissler RW (1994) A definition of initial, fatty streak, and intermediate lesions of atherosclerosis. A report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association *Circulation* 89:2462-2478
12. Staff AC, Redman CW (2014) IFPA Award in Placentology Lecture: preeclampsia, the decidual battleground and future maternal cardiovascular disease *Placenta* 35 Suppl:S26-S31
13. Roberts JM, Pearson G, Cutler J, Lindheimer M (2003) Summary of the NHLBI Working Group on Research on Hypertension During Pregnancy *Hypertension* 41:437-445
14. Johnsen SL, Rasmussen S, Wilsgaard T, Sollien R, Kiserud T (2006) Longitudinal reference ranges for estimated fetal weight *Acta Obstet. Gynecol. Scand* 85:286-297
15. Arduini D, Rizzo G (1990) Normal values of Pulsatility Index from fetal vessels: a cross-sectional study on 1556 healthy fetuses *J. Perinat. Med.* 18:165-172
16. DeVore GR (2015) The importance of the cerebroplacental ratio in the evaluation of fetal well-being in SGA and AGA fetuses *Am. J. Obstet. Gynecol.* 213:5-15. doi: 10.1016/j.ajog.2015.05.024
17. Staff AC, Ranheim T, Khoury J, Henriksen T (1999) Increased contents of phospholipids, cholesterol, and lipid peroxides in decidua basalis in women with preeclampsia *Am. J. Obstet. Gynecol* 180:587-592
18. Turowski G, Berge LN, Helgadottir LB, Jacobsen EM, Roald B (2012) A new, clinically oriented, unifying and simple placental classification system *Placenta* 33:1026-1035. doi: 10.1016/j.placenta.2012.10.002
19. McMANUS JF (1948) Histological and histochemical uses of periodic acid Stain Technol. 23:99-108

20. Pijnenborg R, Ball E, Bulmer JN, Hanssens M, Robson SC, Vercruyssen L (2006) In vivo analysis of trophoblast cell invasion in the human *Methods Mol. Med* 122:11-44
21. Ramsey EM DM (1980) Placental vasculature and circulation *Anatomy, physiology, radiology, clinical aspects atlas textbook*. Georg Thieme Publishers, Stuttgart, pp.
22. Pijnenborg R, Vercruyssen L, Hanssens M (2006) The uterine spiral arteries in human pregnancy: facts and controversies *Placenta* 27:939-958
23. Robertson WB, Brosens I, Dixon HG (1967) The pathological response of the vessels of the placental bed to hypertensive pregnancy *J. Pathol. Bacteriol.* 93:581-592
24. Khong TY, Chambers HM (1992) Alternative method of sampling placentas for the assessment of uteroplacental vasculature *J. Clin. Pathol* 45:925-927
25. Staff AC, Redman CW, Williams D, Leeson P, Moe K, Thilaganathan B, Magnus P, Steegers EA, Tsigas EZ, Ness RB, Myatt L, Poston L, Roberts JM, Global Pregnancy C (2016) Pregnancy and Long-Term Maternal Cardiovascular Health: Progress Through Harmonization of Research Cohorts and Biobanks *Hypertension* 67:251-260. doi: 10.1161/HYPERTENSIONAHA.115.06357
26. Kim YM, Chaemsaitong P, Romero R, Shaman M, Kim CJ, Kim JS, Qureshi F, Jacques SM, Ahmed AI, Chaiworapongsa T, Hassan SS, Yeo L, Korzeniewski SJ (2014) Placental Lesions Associated with Acute Atherosclerosis *J. Matern. Fetal Neonatal Med*:1-29
27. Khong TY (1991) Acute atherosclerosis in pregnancies complicated by hypertension, small-for-gestational-age infants, and diabetes mellitus *Arch. Pathol. Lab Med* 115:722-725
28. Johnsen GM, Størvold GL, Drabbe JMM, Haasnoot GW, Eikmans M, Spruyt-Gerritse MJ, Alnaes-Katjavivi P, Scherjon SA, Redman CWG, Claas FHJ, Staff AC (2018) The combination of maternal KIR-B and fetal HLA-C2 is associated with decidua basalis acute atherosclerosis in pregnancies with preeclampsia *J. Reprod. Immunol.* 129:23-29. doi: 10.1016/j.jri.2018.07.005
29. Care AS, Bourque SL, Morton JS, Hjartarson EP, Robertson SA, Davidge ST (2018) Reduction in Regulatory T Cells in Early Pregnancy Causes Uterine Artery Dysfunction in Mice *Hypertension* 72:177-187. doi: 10.1161/hypertensionaha.118.10858
30. Sasaki N, Yamashita T, Takeda M, Hirata K (2012) Regulatory T cells in atherogenesis *J. Atheroscler. Thromb* 19:503-515
31. Chistiakov DA, Orekhov AN, Bobryshev YV (2016) Immune-inflammatory responses in atherosclerosis: Role of an adaptive immunity mainly driven by T and B cells *Immunobiology* 221:1014-1033. doi: 10.1016/j.imbio.2016.05.010
32. Staff AC, Dechend R, Pijnenborg R (2010) Learning from the placenta: acute atherosclerosis and vascular remodeling in preeclampsia—novel aspects for atherosclerosis and future cardiovascular health *Hypertension* 56:1026-1034
33. Staff AC, Dechend R, Redman CW (2013) Review: Preeclampsia, acute atherosclerosis of the spiral arteries and future cardiovascular disease: two new hypotheses *Placenta* 34 Suppl:S73-S78
34. Staff AC, Johnsen GM, Dechend R, Redman CW (2014) Preeclampsia and uteroplacental acute atherosclerosis: immune and inflammatory factors *J Reprod. Immunol* 101-102:120-126
35. Stevens DU, Al-Nasiry S, Fajta MM, Bulten J, van Dijk AP, van der Vlugt MJ, Oyen WJ, van Vugt JM, Spaanderman ME (2014) Cardiovascular and thrombotic risk of decidual vasculopathy in preeclampsia *Am. J. Obstet. Gynecol* 210:545-546
36. Mosca L, Benjamin EJ, Berra K, Bezanson JL, Dolor RJ, Lloyd-Jones DM, Newby LK, Pina IL, Roger VL, Shaw LJ, Zhao D, Beckie TM, Bushnell C, D'Armiento J, Kris-Etherton PM, Fang J, Ganiats TG, Gomes AS, Gracia CR, Haan CK, Jackson EA, Judelson DR, Kelepouris E, Lavie CJ, Moore A, Nussmeier NA, Ofili E, Oparil S, Ouyang P, Pinn VW, Sherif K, Smith SC, Jr., Sopko G, Chandra-Strobos N, Urbina EM, Vaccarino V, Wenger NK (2011) Effectiveness-based guidelines for the prevention of cardiovascular disease in women—2011 update: a guideline from the American Heart Association *J. Am. Coll. Cardiol.* 57:1404-1423. doi: 10.1016/j.jacc.2011.02.005

Fig. 1 (a) The placenta in relation to the uterine wall, and the location of the decidua parietalis tissue harvested after the delivery of the placenta. MR: placental fetal membrane rolls.

(b) The uteroplacental circulation in the second half of pregnancy. Spiral arteries are the terminal branches of the uterine arteries; decidua basalis is the gestationally transformed endometrium with invading trophoblasts, situated between the placenta and the muscular uterine wall (myometrium).

(c) The location of decidua basalis tissue harvested after the delivery of the placenta: BPS, basal plate section; DSM, decidual vacuum suction method. PBB (placental bed biopsy) was not tested in our present study. Revised from our previous Figure [7]; with permission from the Journal).

Fig. 2 Decidual spiral arteries for each of the three *tissue-type* sample collection methods; decidual vacuum suction (collecting *decidua basalis*), basal placental surface (collecting *decidua basalis*), and membrane rolls (collecting *decidua parietalis*). On the top row are arteries where acute atherosclerosis (AA) is *present*, the bottom row where acute atherosclerosis is *absent*. Decidual sections are IHC-stained brown for foam cells/macrophages (CD68). Inserts show focal accumulation of CD68+ intramural lipid-laden foam cells (x40). Intramural fibrinoid is identified using histochemical staining with Periodic acid Schiff (PAS)

	Decidua basalis tissue		Decidua parietalis tissue
	Decidual vacuum suction (DSM)	Basal placental surface (BPS)	Membrane rolls (MR)
AA present	x10	x5	x1.25
AA absent	x5	x5	x10

Table 1 Rates of decidual spiral artery and acute atherosclerosis presence, according to *decidual tissue type* for the 107 preeclamptic women included in the study. Results are expressed in numbers and rates (%).
DSM: decidual vacuum suction method of placental (yielding *decidua basalis*); BPS: placental basal plate section method (yielding *decidua basalis*); MR: membrane roll (yielding *decidua parietalis*).

Tissue	Decidua basalis		Decidua parietalis	Decidua basalis+parietalis	
	DSM	BPS		MR+BPS	DSM+BPS+MR
Spiral artery detection rate	94/107 (87%)	53/107 (50%)	76/107 (71%)	85/107 (79%)	102/107 (95%)
Acute atherosclerosis rate (of samples with spiral arteries present)	33/94 (35%)	8/53 (15%)	8/76 (11%)	15/85 (18%)	37/102 (36%)

Statistical significance of spiral artery detection rate differences; DSM vs BPS p<0.001, DSM vs MR p=0.007, BPS vs MR p=0.001.
Statistical significance of AA detection differences; DSM vs BPS p=0.012, DSM vs MR p<0.001, BPS vs MR p=0.583.

Table 2. PE-related clinical parameters, according to acute atherosclerosis (AA) detection or not, absence of arteries, per collection method of *tissue-type* evaluated. DSM: decidua suction method (*decidua basalis*), BPS: basal plate (placental) surface (*decidua basalis*), MR: membrane roll (*decidua parietalis*). Values are presented by a median, or by a proportion (%).

<u>Clinical parameters</u>	DSM tissue-type			BPS+MR tissue-types			DB+BPS+MR			All samples
	AA+	AA-	No arteries	AA+	AA-	No arteries	AA+	AA-	No arteries	
<u>Age (years)</u>	N=33	N=60	N=14	N=15	N=73	N=19	N=37	N=65	N=5	N=107
<u>BMI pre-pregnancy (kg/m²)</u>	33.54	32.96	32.49	33.54	33.76	31.23	33.53	33.26	30.45	33.38
<u>BMI delivery (kg/m²)</u>	24.84	22.65	24.89	23.67	23.17	29.08	24.79	22.72	24.52	23.67
<u>Gestational age at delivery (days)</u>	30.84	29.14	31.35	29.04	30.67	29.08	30.84	29.76	29.07	29.76
<u>Birth weight percentile</u>	30+5 § (215)	34+0 (238)	32+2 (226)	29+5§ (208)	34+0 (238)	34+1 (239)	29+6 § (209)	34+0 (238)	34+1 (239)	33+0 (231)
<u>BP@<20weeks</u>	0.12	3.07	0.99	0.20	3.07	2.23	0.12 §	2.74	4.53	1.63
<u>Fetal growth restriction*</u>	120/74	115/72.5	117/77.5	116/73	115/74	115/71	120/74	115/71	113/75	117/73
<u>Pat. fetal Dopplers</u>	26/33 § (79%)	24/60 (40%)	9/14 (64%)	13/15 (87%)	45/73 (62%)	11/18 (61%)	29/37 (78%)	38/65 (58%)	2/5 (40%)	69/107 (64%)
<u>Female infant</u>	13/33 (39%)	12/60 (20%)	4/14 (29%)	8/15 § (53%)	17/73 (23%)	4/18 (22%)	15/37 (41%)	14/65 (22%)	0/5	29/107 (27%)
<u>Primigravida</u>	15/33 (45%)	34/60 (57%)	8/14 (57%)	10/15 (66%)	39/73 (53%)	10/18 (56%)	18/37 (49%)	36/65 (55%)	3/5 (60%)	57/107 (53%)
<u>Previous PE**</u>	15/33 (45%)	29/60 (48%)	3/13 (23%)	7/15 (47%)	26/73 (36%)	9/18 (50%)	17/37 (46%)	29/65 (45%)	1/5 (20%)	47/107 (44%)
	8/14 (57%)	5/20 (25%)	3/5 (60%)	1/4 (25%)	12/30 (40%)	3/5 (60%)	8/14 (57%)	6/23 (26%)	2/4 (50%)	16/41 (39%)

*Fetal growth restriction defined as a birthweight percentile >3, and /or pathological fetal Doppler

**previous nulliparous women (P0) excluded

§ indicates statistically significant (p<0.05) difference in values between samples where AA is detected, and where AA is not detected.

Fig. 3. Clinical parameters (gestational age and birthweight percentile) according to tissue samples categorized according to presence of acute atherosclerosis, AA (° no, • yes). 3a: all three tissue samples per woman are evaluated together (DSM+BPS+MR). 3b: DSM decedua basalis are evaluated separately. 3c: BPS decedua basalis and MR decedua parietalis are evaluated together. Median values shown in horizontal bar.

p values shown when $p < 0.05$. DSM: decidual vacuum suction method. BPS: placental basal plate section. MR: membrane roll.

Supplementary material Table 1 Staining protocol for serial sections (3µm) of decidua basalis (derived from vacuum suction samples of placental bed, and placenta basal plate) and decidua parietalis (derived from fetal membrane rolls). IHC= immunohistochemistry

Slide number	Identifying in decidual tissue:	Staining technique *	Antibody details	Producer
<u>1</u>	Tissue morphology	Hematoxylin and Eosin (H&E)	Routine departmental staining	
<u>2</u> (<i>not always included</i>)	Desmin, component of smooth muscle cells	IHC: Primary antibody monoclonal mouse anti-human Desmin	Clone DE R-11 (Ventana)	Roche Ventana Medical Systems, Inc, Arizona, USA
<u>3</u>	Scavenger receptor common to macrophages and macrophage-derived foam cells	IHC: Primary antibody monoclonal mouse anti-human CD68	Clone KP1 (Dako)	Dako Denmark AS, Glostrup, Denmark

* Slides 2&3 were co-stained with Periodic acid Schiff, using protocol according to Mc Manus (19).

Supplementary material Fig.1 Venn diagram representation of AA detection in uteroplacental tissue types for the 37 preeclamptic women with AA present. No AA was detected in any of the 3 decidual tissue types for the remaining 65 preeclamptic pregnancies with spiral arteries detected. DSM: Decidual vacuum suction method (collecting decidua basalis), BPS: Basal plate placenta section (collecting decidua basalis), MR: membrane rolls (collecting decidua parietalis).

Lymphocyte characterization of decidua basalis spiral arteries with acute atherosclerosis in preeclamptic and normotensive pregnancies

G.M. Johnsen^{a,b,c,1}, G.L. Størvold^{a,b,c,1}, P.H. Alnaes-Katjavivi^{a,b}, B. Roald^{b,d}, M. Golc^{e,f}, R. Dechend^g, C.W.G. Redman^h, A.C. Staff^{a,b,*}

^a Division of Obstetrics and Gynaecology, Oslo University Hospital, Norway

^b Faculty of Medicine, University of Oslo, Norway

^c Institute for Experimental Medical Research, Oslo University Hospital, Norway

^d Department of Pathology, Oslo University Hospital, Norway

^e Department of Obstetrics, Charité - Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany

^f Experimental and Clinical Research Center, a cooperation between the Max Delbrück Center for Molecular Medicine in the Helmholtz Association and the Charité, Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany

^g HELIOS Clinic Berlin Buch, Germany

^h Department of Obstetric Medicine, University of Oxford, Oxford, UK

ARTICLE INFO

Keywords:

Acute atherosclerosis

Decidua

Lymphocytes

Perivascular infiltrate

Preeclampsia

ABSTRACT

Uteroplacental acute atherosclerosis (AA) is a common spiral arterial lesion in preeclampsia, characterized by intramural foam cells, fibrinoid necrosis, and a perivascular immune cell infiltrate. A clear definition of this infiltrate is lacking. Therefore, our aim was to characterize lymphocytes in pre-defined zones regarding spiral arteries with or without AA, from preeclamptic and normotensive pregnancies.

Lymphocytes were characterized in decidua basalis samples (n = 91), previously evaluated for AA, around spiral arteries in three pre-defined zones; 1) intramural, 2) perivascular and 3) interstitial. Adjacent serial sections were immunostained to identify different T-cell populations (CD3+, CD8+, FOXP3+), and NK-cells (CD56+). CD3+CD8- T-cells were also identified. These were presumed to be largely CD4+ T-cells.

AA was associated with significantly higher intramural CD3+ cell concentrations in Zone 1, in both normotensives and preeclamptics. In preeclamptics only, this difference extended into Zone 2. Similar results were observed for CD3+CD8- cells. AA was also associated with increased intramural CD8+ concentration; however, the number of cells was low. Regulatory T-cells (FOXP3+) were generally scarce or absent in all pre-defined zones. Although intramural NK-cells (CD56+) were scarce, the intramural concentration was significantly lower in spiral arteries with AA compared to without AA in preeclamptics.

Our main finding was that CD3+CD8-FoxP3- T-cells were associated with AA. We therefore suggest that T-cells, of a non-regulatory CD4+ subtype, could be involved in the formation of spiral artery AA in the decidua basalis. Whether AA gives rise to, or is partly mediated by increased T-cell concentration around the lesions, remains to be determined.

1. Introduction

Acute atherosclerosis (AA) of the uteroplacental spiral arteries is a pregnancy-specific lesion predominantly found in preeclampsia, also occurring in a lower proportion of uncomplicated pregnancies (Alnaes-Katjavivi et al., 2016). The mechanisms behind AA development are not established, but we have suggested that it is an inflammatory lesion

caused by excessive decidual inflammation (Staff et al., 2013).

Spiral artery AA has been characterized by presence of fibrinoid necrosis in the artery media, sub-endothelial lipid-laden “foam cells”, and the presence of a perivascular infiltrate (PVI) (Robertson et al., 1976). PVI is included in many, but not all descriptions of AA, but has not been clearly defined (Hertig, 1945; Zeek and Assali, 1950; Robertson et al., 1967; Meekins et al., 1994). Most of these descriptions

* Corresponding author at: Department of Obstetrics and Department of Gynaecology, Women's Division, Oslo University Hospital, Ullevål, PO Box 4956 Nydalen, NO-0424, Oslo, Norway.

E-mail address: annetne.staff@ous-hf.no (A.C. Staff).

¹ These authors contributed equally.

<https://doi.org/10.1016/j.jri.2019.03.003>

Received 4 January 2019; Received in revised form 13 March 2019; Accepted 21 March 2019

0165-0378/© 2019 Elsevier B.V. All rights reserved.

Fig. 1. Representative examples of immunohistochemistry lymphocyte staining.

Representative examples of lymphocyte staining of decidua basalis tissue sections. First row demonstrate a) H&E, b) CD3, and c) CD8 staining of tissue sections from a late-onset preeclampsia patient (gestational week > 34) without AA (nonAA artery/nonAA sample). Second row demonstrate d) H&E, e) CD3, and f) CD8 staining of tissue sections from an early-onset preeclampsia patient (gestational week < 34).

are based on simple hematoxylin and eosin staining. No explicit definition or consensus exists regarding the phenotype of the constituent PVI leukocytes. Therefore, we sought to describe the presence of lymphocytes around decidual spiral arteries in a more systematic manner by using pre-defined zones for their quantification in, and around the arteries, and specifically in relation to AA. Additionally, we sought to identify specific lymphocyte subsets.

Decidual lymphocytes are of maternal origin and are phenotypically different compared to circulating immune cells. In most previous studies, NK-cells have been emphasized as the predominant immune cells in the early stages of pregnancy, where they have essential roles in regulation of extravillous trophoblast (EVT) invasion and spiral artery remodeling (Pijnenborg et al., 2006; Cartwright et al., 2017). Towards term, the number of decidual NK-cells is significantly reduced and their role is undefined. Reduced frequency of decidual NK-cells has been reported in preeclampsia and fetal growth restriction (Williams et al., 2009a), but has not been reported in AA.

We are studying AA in third trimester decidual samples, where most decidual leukocytes are T-cells (Tilburgs et al., 2009; Williams et al., 2009b). The role of these decidual T-cells is generally unknown (Nancy and Erlebacher, 2014). They can respond to paternal/placental antigens (Sharkey et al., 2008) but could also be important to control infection (Lissauer et al., 2017). However, mice studies indicate that regulatory T-cells (Tregs) are important in implantation and fetomaternal immunotolerance (Erlebacher, 2013). Also, Tregs appear to be reduced in the circulation and at the fetomaternal interface in women with preeclampsia (Nancy and Erlebacher, 2014). The role of T-cells and Tregs in relation to AA is not known.

Here, our aim was to characterize the immune cell infiltrate present around spiral arteries with AA compared to spiral arteries without AA in preeclamptic and normotensive pregnancies, and to describe both its

localization and its composition of lymphocyte subtypes.

2. Methods

2.1. Patient inclusion

Pregnant women were recruited, after written consent, as described in Supplemental Methods. Clinical characteristics of the pregnancies are presented in Supplemental Table 1. From our cohort of 237 pregnancies previously evaluated for AA in the decidua basalis collected by vacuum suctioning of the placental bed (Alnaes-Katjavivi et al., 2016), we included 91 pregnancies without diabetes and with the highest number of spiral artery sections per decidual slide. Of these, 54 had preeclampsia (27 with AA) and 37 were normotensive (10 with AA). Three pregnancies were excluded, owing to lack of spiral arteries in the tissue sections for the present study.

2.2. Acute atherosclerosis evaluation

AA in decidua basalis was evaluated as described in Supplemental Methods and Supplemental Fig. 1. AA was defined as presence of minimum two intramural vacuolated CD68+ cells (foam cell lesion) in the wall of at least one spiral artery (Alnaes-Katjavivi et al., 2016).

2.3. Serial immunostaining of lymphocytes in decidua basalis

For each woman four 3 μm adjacent tissue sections from one decidua basalis sample were analyzed with immunohistochemistry (IHC). The sections were in continuation with those slides previously used to evaluate AA (Alnaes-Katjavivi et al., 2016). The slides were stained as follows: 1) regulatory T-cells (FOXP3: Abcam 236 A/E7 Monoclonal

mouse nuclear stain), 2) CD3+ T-cells (CD3: Roche 2GV6 Monoclonal rabbit anti-human CD3 cytoplasmic stain), 3) CD8+ T-cells (CD8: Dako Clone C8/144B Monoclonal mouse anti-human CD8 cytoplasmic domain stain), and 4) NK-cells (CD56: Cell Marque MRO-42 Monoclonal rabbit anti-human) (Figs. 1 & 4). The sections were immunostained using a Ventana Benchmark XT autostainer with DAB staining, followed by Periodic Acid Schiff counterstain for identification of spiral arteries. Positive immunostaining was confirmed using human tissue as controls: tonsillar lymphatic tissue (CD3, CD8 and FoxP3), and ileum small bowel (CD56).

2.4. Estimation of CD4+ T-cells

The decidua basalis tissue contained many CD4+ macrophages, consequently we estimated the CD4+ T-cell concentration based on the results from the CD3 and CD8 staining, similar to a previous publication (Stallmach et al., 1999). The CD3 pan T-cell marker stains CD3+ T-cells that are mainly either CD8+ or CD8-. The vast majority of T-cells are committed to either a CD4+ or CD8+ lineage, however minor subsets (1–5%) of double negative (CD3+CD4-CD8-) cells also exist (Juvet and Zhang, 2012). As far as we are aware double negative T-cells have not been documented in the human decidua. Since the CD3 stained slide was adjacent in the sequence of slides to the slide stained with CD8 (3 µm), it allowed for estimation of a CD3+CD8- T-cells (presumptive CD4+ cell population) by subtracting the concentration of CD8+ T-cells from CD3+ T-cells, in the same tissue zone investigated.

2.5. Quantification of lymphocytes in decidua basalis

Cells were counted manually using a Motic BA-410 LED light microscope, without knowledge of the pregnancy outcome. Pictures of each spiral artery section on each slide were recorded digitally using a Moticam 2500 digital camera and Motic Images vs 2.0. Lymphocyte concentrations were determined for each spiral artery, by counting positively stained cells in pre-defined counting zones. To ensure reproducibility, two researchers independently evaluated ten percent of the tissue sections.

2.6. Pre-defined lymphocyte counting zones

We defined three specific and reproducible counting areas, named Zone 1–3 (Fig. 2). We measured the area of each zone for each artery, and calculated the lymphocyte concentrations in each zone as the number of immunopositive cells within the zone investigated, divided

by the area of the zone (cells/mm²). This enabled comparison of cell density between arteries of different sizes, from different decidua basalis samples, or different arteries within the same samples. It also enabled comparison of cell density in the three different zones. **Zone 1**, “intramural”, represents the entire area of the spiral artery wall, from the endothelium to the outer limit of the smooth muscle/PAS + margin of the artery (Fig. 2). **Zone 2**, “perivascular”, represents the immediate adjoining area surrounding the spiral artery wall, demarcating a “cuff” of 100 µm tissue width, from the outer limit of the artery wall (Fig. 2). In Zone 2 we examined potential co-localization of perivascular lymphocytes and AA by dividing Zone 2 into quadrants originating from the center of the artery lumen (Fig. 2a). **Zone 3**, “interstitial infiltrate”, represents the area containing the largest visible concentration of T-cells (in the CD3 stained slide, Fig. 1) present in the surrounding decidual tissue continuous with the spiral artery evaluated. The three largest clusters of CD3+ cells were identified at x4 magnification, and demarcated by a counting disc (100 µm in diameter) (Fig. 2), the cells were counted at x40 magnification.

2.7. Definition of spiral artery AA categories

AA is typically “patchy” (Pijnenborg et al., 2006). When AA was identified in a sample “AA sample” it often contained spiral arteries with “AA artery/AA sample”, and without AA “nonAA artery/AA sample”. For Zone 1 and 2, we applied this categorization to determine whether lymphocyte patterns associate with AA presence in the sample, or rather associate more specifically to the lesion of the individual artery. Samples where none of the spiral arteries had AA “nonAA sample” were defined as “nonAA artery/nonAA sample”. For patient samples with more than one spiral artery in the same category, the mean lymphocyte concentration across these arteries was calculated and used in the statistical analyses.

2.8. Statistical analyses

Data were analyzed using SPSS version 22. The Fisher Exact test was used for dichotomous variables and the non-parametric Mann-Whitney test otherwise. A p-value of ≤ 0.05 was considered statistically significant.

Fig. 2. Schematic illustration and examples of the pre-defined Zones for quantification of lymphocytes in and around decidua basalis spiral arteries. a) Schematic representation of a decidual spiral artery and the three different Zones used for quantification of decidual lymphocytes. Zone 1 “intramural”, spiral artery wall (red with solid black perimeter). Zone 2, “perivascular”, area surrounding the intramural zone. The outer perimeter is set at a depth of 100 µm from the outer margin of the artery wall (orange with interrupted black perimeter). Zone 3 “interstitial infiltrate”, circle-shaped area of 100 µm in diameter in surrounding decidual tissue not directly in contact with the spiral artery but with a continuous connection to the spiral artery (circle with interrupted black perimeter). The star (*) denotes the artery lumen. b) Examples of the applied zones for a decidual spiral artery section with AA stained for CD3.

tery (circle with interrupted black perimeter). The star (*) denotes the artery lumen. b) Examples of the applied zones for a decidual spiral artery section with AA stained for CD3.

3. Results

3.1. Lymphocyte distribution across all zones

The decidua basalis samples contained a median of three spiral arteries (range 1–10). In general, scattered staining of CD3+ and CD8+ lymphocytes was observed throughout the decidual tissue. Lymphocytes were observed in all three pre-defined zones, across samples with and without AA. The highest concentration of immunopositive cells (cells/mm²) was observed in the surrounding interstitial infiltrates (Zone 3), followed by the perivascular area (Zone 2). In Zone 1, CD3+, CD8+, and CD56+ cells were scarce or absent (Supplemental Table 2). FoxP3+ staining was scarce or absent in all zones investigated (Supplemental Table 2), both in preeclampsia and normotensive pregnancies.

3.2. Intramural T-cell distribution (Zone 1)

Although the number of cells in Zone 1 (Figs. 1 & 2) was low, there were significantly higher concentration of CD3+ cells in the artery wall in spiral arteries with AA (AA artery/AA sample) compared to those in samples without AA (nonAA artery/nonAA sample) (Fig. 3a). This was the case both in preeclampsia (Fig. 3a) and in normotensives (Supplemental Fig. 2a). In the preeclampsia, the concentration of CD3+ cells was also significantly higher in arteries with AA (AA artery/AA sample) than in arteries without AA (nonAA artery/AA sample) in samples with AA (Fig. 3a). The results were similar for CD8+ cells. The

intramural CD8+ concentration was significantly higher in spiral arteries with AA (AA artery/AA sample) than without AA (nonAA artery/nonAA sample), but only in preeclampsia (Fig. 3b). There were no differences in CD8+ concentration in the normotensives (Supplemental Fig. 2b). The CD3+CD8- cells (inferred CD4+ T-cells) followed the same pattern as CD3+ cells. The CD3+CD8- concentration was significantly higher in spiral arteries with AA (AA artery/AA sample) compared to those in samples without AA (nonAA artery/nonAA sample) in both in preeclampsia (Fig. 3c) and in normotensives (Supplemental Fig. 2c). In preeclampsia, the concentration of CD3+ cells was also significantly higher in arteries with AA (AA artery/AA sample) than in arteries without AA (nonAA artery/AA sample) in samples with AA (Fig. 3c).

3.3. Perivascular T-cell distribution (Zone 2)

In Zone 2 (Figs. 1 & 2) the CD3+ and CD3+CD8- (inferred CD4+) concentration was significantly higher in preeclampsia in spiral arteries with AA (AA artery/AA sample) compared to spiral arteries in samples without AA (nonAA artery/nonAA sample) (Fig. 3d, f). There was no difference in CD8+ cells in Zone 2 in preeclampsia (Fig. 3e). In normotensives, there were no differences in CD3+, CD8+ or CD3+CD8- (inferred CD4+) cells across the spiral artery categories (Supplemental Fig. 2d-f). The T-cells in Zone 2 were distributed unevenly in clusters, between the different artery quadrants. However, we found no significant co-localization of foam cells and artery quadrants demonstrating the highest T-cell concentrations (data not shown).

Fig. 3. T-cell concentration in the pre-defined Zones 1–3 in preeclamptic pregnancies. Dot plots of median T-cell concentrations per patient in the pre-defined counting zones. Each plot shows the T-cell concentration in three categories of spiral arteries: 1) arteries without AA in samples without AA (nonAA artery/nonAA sample, white dots), 2) arteries without AA in a sample with other arteries affected by AA (nonAA artery/AA sample, grey dots), and finally 3) arteries with AA lesions (AA artery/AA sample, black dots). The different plots show: a) CD3+ cells, b) CD8+ cells, and c) CD3+CD8- cells (inferred CD4+ cells) in Zone 1 (intramural area); d) CD3+ cells, e) CD8+ cells, and f) CD3+CD8- cells (inferred CD4+ cells) in Zone 2 (perivascular area); and g) CD3+ cells, h) CD8+ cells, and i) CD3+CD8- cells (inferred CD4+ cells) in Zone 3 (interstitial infiltrate area). The p-values were calculated using Mann Whitney U-test, a p-value < .05 was considered statistically significant.

Fig. 4. NK-cell concentration in the pre-defined counting Zones 1–3 in preeclamptic pregnancies. Dot plots of median NK-cell concentrations per patient in the pre-defined counting zones. Each plot shows the NK-cell in three categories of spiral arteries: 1) arteries without AA in samples without AA (nonAA artery/nonAA sample, white dots), 2) arteries without AA in a sample with other arteries affected by AA (nonAA artery/AA sample, grey dots), and finally 3) arteries with AA lesions (AA artery/AA sample, black dots). The different plots show: a) CD56+ cells in Zone 1 (intramural area), b) CD56+ cells in Zone 2 (perivascular area), and c) CD56+ cells in Zone 3 (interstitial infiltrate area). The p-values were calculated using Mann Whitney U-test, a p-value < .05 was considered statistically significant. d) Representative serial staining of CD56 in decidual tissue sections from a late-onset preeclampsia patient (nonAA artery/nonAA sample). e) Representative staining of CD56 in decidual tissue sections from an early-onset preeclampsia patient (AA artery/AA sample).

3.4. Interstitial T-cell distribution (Zone 3)

For Zone 3 (Figs. 1 & 2), the concentration of CD3+CD8- cells (inferred CD4+) in samples with AA was higher than in samples from women without AA in preeclampsia (Fig. 3i). A similar trend (but not significant) was observed for CD3+ cells in preeclampsia (Fig. 3g). None of the T-cell markers were significantly changed for AA in Zone 3 in normotensives (Supplemental Fig. 2g-i).

3.5. NK-cell distribution across all zones

CD56+ NK-cells (Figs. 1 & 4 d-e) were less abundant than CD3+ cells in all zones (Supplemental Table 2), with the highest concentration of NK-cells in Zone 3. In Zone 1 there was a significantly lower concentration of cells in arteries with AA as compared to arteries without AA, however the NK-cells were generally scarce or absent in this zone (Fig. 4a). In Zone 2 and 3, the NK cell concentration was not significantly different between samples with AA compared to without AA in either preeclamptics (Fig. 4a-c) or normotensives (Supplemental Fig. 3a-c).

3.6. Preeclamptic vs. normotensive pregnancies

We found no significant differences for any of the lymphocyte

markers assessed (CD3, CD8, FOXP3 and CD56) when comparing pre-eclamptic and normotensive pregnancies in Zone 2 and 3 (Supplemental Fig. 4).

4. Discussion

A perivascular infiltrate (PVI) has been considered a classical characteristic of uteroplacental AA lesions. However, its definition and characterization is lacking. Decidua basalis contains many different maternal immune cells and the composition of these cells change during the course of pregnancy. Lymphocytes are always present in the tissue surrounding the spiral arteries, and the presence of perivascular leukocytes is not a unique feature of AA (Robertson et al., 1967; Stevens et al., 2012; Alnaes-Katjavivi et al., 2016). Here, we characterized the lymphocyte concentration in pre-defined zones without considering whether the cells should be termed “PVI” or not, and found a general significant increase in T-cell concentration around spiral arteries with AA. We rarely observed dense clusters of perivascular lymphocytes around spiral arteries with or without AA. Therefore, the term “PVI” may not be useful as a mandatory prerequisite for identifying AA in decidua basalis spiral arteries. However, knowledge about the phenotype of the lymphocytes present could improve understanding of AA lesions, and their relation to pregnancy complications such as pre-eclampsia.

We found that in pregnancies affected by preeclampsia, AA in spiral arteries was associated with higher concentrations of CD3+ cells in both intramural and perivascular zones compared to spiral arteries without AA. However, the cell counts in the intramural area (Zone 1) were low and therefore we propose that T-cells in the perivascular area (Zone 2) may be more important in the context of AA development. More specifically, this increase in CD3+ cells in the perivascular zone was due to an increase in the CD3+CD8- subset, which we use as a surrogate measure of CD4+ cells. These cells were also FOXP3-. In addition, CD4+ T-cells were increased in interstitial infiltrates (Zone 3) in preeclampsia samples with AA. This increase in perivascular and interstitial T-cells could indicate a localized decidual response to the AA affected artery. We found that decidual NK-cells were few in all three zones investigated, with a lower intramural concentration in AA in our 3rd trimester samples. However, we cannot exclude a role for decidual NK-cells at an earlier point in the development of AA.

Stevens et al (2012) observed that a perivascular infiltrate was present more often in arteries with AA, however not distinguishing between decidual basalis and parietalis (Stevens et al., 2012). In a large study of over 14,000 pregnancies, Kim et al (2015) used the presence of PVI in their AA definition, without defining PVI (Kim et al., 2015). This could explain their low frequency of AA as compared to our cohort (Alnaes-Katjavivi et al., 2016). In these studies, only H&E staining was used, and hence they cannot distinguish the types of immune cells present. Hecht et al (2016) studied decidual parietalis tissue from six patients with preeclampsia, identifying with CD45 (pan-lymphocyte marker) small lymphocyte aggregation around arteries with AA, however without comparing to arteries without AA (Hecht et al., 2016). Interestingly, these lymphocytes also stained for nitrotyrosine, a marker of oxidative injury (Baltajian et al., 2014).

Our results could indicate that some AA lesions are associated with an adaptive immune response that involves CD4 helper T-cells. However, as our samples were collected at the time of delivery, when AA is well established, we cannot determine whether the higher concentrations of CD3+CD8- cells (inferred CD4+ T-cells) are a cause or a consequence of AA. CD4+ T-cells consist of several subtypes with different inflammatory and suppressive properties, and exist in several differentiation stages. However, the T-cells we observed probably do not have a regulatory phenotype since FOXP3+ staining was almost absent in our samples. Based on the markers used in our study, we cannot conclude regarding the function or the differentiation stage of the CD4+ cells that we observed. However, recent studies show that a majority of decidual T-cells are highly differentiated effector and memory cells (Feyaerts et al., 2017; Lissauer et al., 2017). This means that they are experienced antigen-specific T-cells, that probably have been in contact with fetal antigens (Powell et al., 2017). Decidual T-cells have a unique transcriptional profile with high expression of genes involved in inflammatory interferon signaling (Powell et al., 2017). However, the number of T-cells we observe are few, and do not imply a massive tissue inflammatory reaction.

We previously reported a lack of intramural (Zone 1) EVT in spiral arteries with AA lesions (Alnaes-Katjavivi et al., 2016). The absence of intramural EVTs could influence the activation and signaling of nearby lymphocytes. Interaction of decidual NK-cells with EVTs regulate their secretion of cytokines and angiogenic factors, besides cytotoxicity, and is an important part of the placentation process (Cartwright et al., 2017). However, AA is also found in basal arteries of the decidual parietalis where there is no trophoblast invasion (Pijnenborg et al., 2006). Trophoblasts appear to be able to regulate the induction of Tregs in the decidual by direct interaction with T-cells and by secreted factors (Svensson-Arvelund et al., 2015). Here, we found that Tregs, defined as FOXP3+ cells, were scarce in all the three zones investigated. Hence, we speculate that simultaneous increase in T-cells and lack of EVTs in the spiral arteries, could potentially lead to local decidual inflammation that is not regulated by EVTs or Tregs. This in turn could contribute to foam cell development and AA formation.

We recently found that a combination of maternal activating KIR haplotype (KIR-B) and fetal HLA-C2 genes was associated with AA in preeclampsia (Johnsen et al., 2018). In the decidual KIR receptors are predominantly expressed by NK-cells, but a subset of T-cells can also express KIRs (Tilburgs et al., 2009). These results indicate a role for decidual NK-cells and possibly T-cells in AA. Based on our findings here we find increased support for the involvement of T-cells in AA etiology.

A limitation of our study is that the results are based on IHC staining of formalin-fixed-paraffin-embedded tissue, which limits the number of markers used. Conversely, IHC has the advantage of showing the specific location of the cells in connection with spiral arteries. The indirect estimation of CD4+ T-cells is likely less accurate than direct counting of stained cells, and might be an overestimation as there could be CD3+ cells present that are neither CD8+ nor CD4+, such as NKT-cells or other 'double-negative' subsets. However, we believe that any under- or overestimation of the CD4+ cells would be the same regardless of AA, and therefore not affect our conclusions.

The cause of the spiral artery PVI is still unknown. However, here we give a more detailed description of its composition and location in and around spiral arteries in the decidual basalis. We conclude that AA is associated with an increased concentration of T-cells (inferred CD4+) in the intramural, perivascular and possibly the interstitial zones in pregnancies with preeclampsia. This increase in T-cells coincides with a lack of intramural EVTs in the spiral arteries (Alnaes-Katjavivi et al., 2016). We speculate that this leads to a lack of interaction between fetal and maternal cells that would be important for regulation of immunological processes in decidual basalis. We suggest that this could be one of several factors contributing to the formation of AA.

Funding

The Research Council of Norway funded part of this work (the PATH study: Pregnancy acute atherosclerosis and future cardiovascular disease, grant number ref. 230,652) as did South-Eastern Norway Regional Health Authority (the HAPPY PATH study, grant number ref. 2,014,026). Oslo University Hospital provided PhD salary (PAK) and further research support.

Conflict of interest statement

The authors declare no conflicts of interest.

Acknowledgements

The authors greatly acknowledge the assistance from pathology assistants Linn Buer, Lene Nordenstrøm and Giang Huong Nguyen, and the laboratory assistance and patient recruitment by Lise Øhra Levy, as well as patient recruitment by previous PhD students, including Kristin Brække, Nina K. Harsem and Meryam Sugulle (all Oslo University Hospital).

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.jri.2019.03.003>.

References

- Alnaes-Katjavivi, P., et al., 2016. Acute atherosclerosis in vacuum suction biopsies of decidual basalis: an evidence based research definition. *Placenta* 37, 26–33.
- Baltajian, K., et al., 2014. Placental lesions of vascular insufficiency are associated with anti-angiogenic state in women with preeclampsia. *Hypertens. Pregnancy* 33, 427–439.
- Cartwright, J.E., et al., 2017. The role of decidual NK cells in pregnancies with impaired vascular remodelling. *J. Reprod. Immunol.* 119, 81–84.
- Erlebacher, A., 2013. Mechanisms of T cell tolerance towards the allogeneic fetus. *Nat.*

- Rev. Immunol. 13, 23–33.
- Feyaerts, D., et al., 2017. Human uterine lymphocytes acquire a more experienced and tolerogenic phenotype during pregnancy. *Sci. Rep.* 7, 2884.
- Hecht, J.L., et al., 2016. Revisiting decidual vasculopathy. *Placenta* 42, 37–43.
- Hertig, A.T., 1945. Vascular pathology in hypertensive albuminuric toxemias of pregnancy. *Clinics* 4, 602–614.
- Johnsen, G.M., et al., 2018. The combination of maternal KIR-B and fetal HLA-C2 is associated with decidua basalis acute atherosclerosis in pregnancies with preeclampsia. *J. Reprod. Immunol.* 129, 23–29.
- Juvet, S.C., Zhang, L., 2012. Double negative regulatory T cells in transplantation and autoimmunity: recent progress and future directions. *J. Mol. Cell Biol.* 4, 48–58.
- Kim, Y.M., et al., 2015. The frequency of acute atherosclerosis in normal pregnancy and preterm labor, preeclampsia, small-for-gestational age, fetal death and midtrimester spontaneous abortion. *J. Matern. Fetal Neonatal Med.* 28, 2001–2009.
- Lissauer, D., et al., 2017. Maternal effector T cells within decidua: The adaptive immune response to pregnancy? *Placenta* 60, 140–144.
- Meekins, J.W., et al., 1994. Immunohistochemical detection of lipoprotein(a) in the wall of placental bed spiral arteries in normal and severe preeclamptic pregnancies. *Placenta* 15, 511–524.
- Nancy, P., Erlebacher, A., 2014. T cell behavior at the maternal-fetal interface. *Int. J. Dev. Biol.* 58, 189–198.
- Pijnenborg, R., et al., 2006. The uterine spiral arteries in human pregnancy: facts and controversies. *Placenta* 27, 939–958.
- Powell, R.M., et al., 2017. Decidual T cells exhibit a highly differentiated phenotype and demonstrate potential fetal specificity and a strong transcriptional response to ifn. *J. Immunol.* 199, 3406–3417.
- Robertson, W.B., et al., 1967. The pathological response of the vessels of the placental bed to hypertensive pregnancy. *J. Pathol. Bacteriol.* 93, 581–592.
- Robertson, W.B., et al., 1976. Maternal uterine vascular lesions in the hypertensive complications of pregnancy. *Perspect. Nephrol. Hypertens.* 5, 115–127.
- Sharkey, A.M., et al., 2008. Killer Ig-like receptor expression in uterine NK cells is biased toward recognition of HLA-C and alters with gestational age. *J. Immunol.* 181, 39–46.
- Staff, A.C., et al., 2013. Review: Preeclampsia, acute atherosclerosis of the spiral arteries and future cardiovascular disease: Two new hypotheses. *Placenta* 34 (Suppl), S73–8.
- Stallmach, T., et al., 1999. Aberrant positioning of trophoblast and lymphocytes in the feto-maternal interface with pre-eclampsia. *Virchows Arch.* 434, 207–211.
- Stevens, D.U., et al., 2012. Decidual vasculopathy and adverse perinatal outcome in preeclamptic pregnancy. *Placenta* 33, 630–633.
- Svensson-Arvelund, J., et al., 2015. The human fetal placenta promotes tolerance against the semiallogeneic fetus by inducing regulatory T cells and homeostatic M2 macrophages. *J. Immunol.* 194, 1534–1544.
- Tilburgs, T., et al., 2009. Expression of NK cell receptors on decidual T cells in human pregnancy. *J. Reprod. Immunol.* 80, 22–32.
- Williams, P.J., et al., 2009a. Altered decidual leucocyte populations in the placental bed in pre-eclampsia and foetal growth restriction: a comparison with late normal pregnancy. *Reproduction (Cambridge, England)* 138, 177–184.
- Williams, P.J., et al., 2009b. Decidual leucocyte populations in early to late gestation normal human pregnancy. *J. Reprod. Immunol.* 82, 24–31.
- Zeek, P.M., Assali, N.S., 1950. Vascular changes in the decidua associated with eclamptogenic toxemia of pregnancy. *Am. J. Clin. Pathol.* 20, 1099–1109.