Medication appropriateness, complexity and adherence in older adults with chronic kidney disease

Maria Krystina Parker, MD

Department of Nephrology

Division of Medicine

Akershus University Hospital

and

Institute of Clinical Medicine

Faculty of Medicine

University of Oslo

September 2019

© Maria Krystina Parker, 2020

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-607-2

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

“Take a little aspirin, add one part low-dose cholesterol medicine and three parts low-dose blood pressure medicine. Put it in a single pill and give to everybody older than age 45.

What do you get?”

John Fauber, Journal Sentinel, March 2009

TABLE OF CONTENTS

1	ACKNOWLEDGEMENTS	I
2	LIST OF THE STUDIES	1
3	ABBREVIATIONS	2
4	INTRODUCTION.....	4
5	BACKGROUND	6
5.1	The aging population	6
5.2	Old age and chronic diseases	8
5.3	Age-related changes in physiology, pharmacokinetics and pharmacodynamics	8
5.4	Chronic kidney disease	11
5.4.1	Changes in pharmacokinetics and pharmacodynamics	13
5.4.2	Prevalence	14
5.4.3	Treatment modality	15
5.5	Prescribing medication	18
5.5.1	Polypharmacy	19
5.5.2	Potentially inappropriate medication	21
5.5.3	Assessment of potentially inappropriate medication	21
5.5.4	Prescribing of PIPs in older adults with CKD.....	22
5.6	Medication regimen complexity	22
5.7	Medication adherence.....	23
5.7.1	Assessment of medication adherence.....	25

5.7.2	Medication adherence in older adults.....	26
5.7.3	Medication adherence in CKD patients	27
6	AIMS OF THE THESIS.....	28
7	MATERIALS AND METHODS	29
7.1	Potentially inappropriate medications with the STOPP criteria (Study I).....	29
7.1.1	Study population	29
7.1.2	Data collection.....	29
7.1.3	Instruments	29
7.1.4	Statistical analysis	31
7.2	Medication complexity regimen and adherence (Study II)	32
7.2.1	Study population	32
7.2.2	Enrolment and data collection at baseline.....	32
7.2.3	Instruments	33
7.2.4	Statistical analysis	35
7.3	Effectiveness of using the STOPP/START criteria (Study III).....	36
7.3.1	Study population	36
7.3.2	Randomization and intervention	36
7.3.3	Follow-up after six months with data collection.....	36
7.3.4	Instruments	38
7.3.5	Statistical analysis	39
7.4	Ethics	40

7.4.1	Potentially inappropriate medications with the STOPP criteria (Study I)	40
7.4.2	Medication regimen complexity and adherence (Study II) and Effectiveness of using the STOPP/START criteria (Study III).....	40
8	RESULTS	41
8.1	Potentially inappropriate medications with the STOPP criteria (Study I).....	41
8.2	Medication regimen complexity and adherence (Study II)	42
8.3	Effectiveness of using the STOPP/START criteria (Study III).....	43
9	DISCUSSION	44
9.1	General findings	44
9.2	Discussion of the methodology	45
9.2.1	Study design	45
9.2.2	Reliability and validity	46
9.2.3	Instruments	47
9.2.4	Confounding and bias.....	49
9.2.5	Missing data	51
9.2.6	Statistical considerations	52
9.3	Discussion of the main findings	53
9.3.1	Potentially inappropriate medications with the STOPP criteria (Study I)	53
9.3.2	Medication regimen complexity and adherence (Study II).....	55
9.3.3	Effectiveness of using the STOPP/START criteria (Study III)	58
10	CONCLUSIONS.....	61

11	CLINICAL IMPLICATIONS.....	62
12	SUGGESTIONS FOR FUTURE RESEARCH	64
13	REFERENCES	66
14	LIST OF ERRATA	77
	PAPERS	78
	APPENDIX	

1 ACKNOWLEDGEMENTS

I want to thank the University of Oslo for employing me as a PhD student. Through this opportunity, I learned much, both as a researcher and a clinician, and met amazing people who inspired me.

It is to Knut Stavem that I first wish to express my utmost gratitude for his patience and acceptance in taking on the task of being my mentor. Truthfully, in having me as a novice in every step of these scientific studies, he did not have the easier part of this arrangement. He guided me through many obstacles and his door was always open. Thank you!

Ingrid Os, I hope that you realise how greatly I appreciate the role you played with your positivity and support. I enjoyed working with you and always appreciated your valuable suggestions.

Beginning with my early attempts (before I was accepted as a PhD student), I received never-ending support from my friend, Willy Aasebø. He always had time to discuss the challenges of the PhD project and the writing process. Thank you!

For her help and unconditional trust in dozens of ways, I thank my friend Toril Enger.

My thanks go to every colleague in the nephrology department, Akershus University Hospital, for their support during inclusion and follow-up of patients, especially in offering convenient logistical solutions so that the elderly could participate in this project.

I thank Aud Stenehjem and her team at OUS Ullevål, nephrology department, for their unlimited support and engagement in my study project. It has been a joy to meet her, Christa, Berit and all the doctors and nurses. A special thank you goes to Heidi for her interest in my work, especially for the many open and encouraging discussions we had.

My deepest thanks go to Morten Reier-Nilsen, Ingrid, Elin, and all the nephrologists from Vestre Viken, Drammen Hospital. They welcomed me so warmly, included me in their professional discussions and provided me with warm and funny memories.

I also want to thank to the members of our “PhD-club”: Anke, Kjersti, Nanna, Nina, and Viera, for all the lively discussions about quality of life, validation, accuracy, and English grammar. I am grateful to each of you for providing laughter along the way. Unfortunately, I must announce the end of the club. I will take all the warm memories with me, Thank you!

I am grateful for the friendship, fun and support provided by Andre, Berit, Ellen, and Vibeke. While struggling with writing, our lunch themes about family, gardening, baking, and diet/exercising were always a welcome respite. Furthermore, I would like to thank Prof. Torbjørn Omland and Anna B. Frengen for their support.

I wish to thank all the patients, who were positive towards my project and gave their unconditional support. I have learned so much from you all.

I am also grateful for the opportunity to teach and meet medical students. I loved every moment of it and I will definitely miss teaching.

My sincere thanks to my family and friends in Germany and England for their interest in my work and their love. My thanks also go to my friends Annelise, Sonja, Therese, and Toril for their help and support. I offer a special thank you to my editor-in-chief and father-in-law: Vernon Parker.

Most helpful of all, the most encouraging, inspiring and indispensable, as always, and most deserving of my wholehearted gratitude, is my guiding star and my husband Glyn. In addition, our marvellous and “never stop to amaze me” children, Emilie Marie and Karl Vernon. Your

laughter and joy of life comforted me and reminded me of what is already my most significant success.

September 2019

Krystina Parker

2 LIST OF THE STUDIES

I. **Potentially inappropriate medications in elderly haemodialysis patients using the STOPP criteria**

Krystina Parker, Willy Aasebø, Knut Stavem

Drugs – Real World Outcome. 2016 Sep; 3(3): 359–363.

II. **Medication regimen complexity and medication adherence in elderly patients with chronic kidney disease**

Krystina Parker, Ingrid Bull-Engelstad, Willy Aasebø, Nanna von der Lippe, Morten Reier-Nilsen, Ingrid Os, Knut Stavem

Hemodialysis International. 2019 Jul;23(3):333-342.

III. **Effectiveness of using STOPP/START criteria to identify potentially inappropriate medication in people aged ≥ 65 years with chronic kidney disease: a randomized clinical trial**

Krystina Parker, Ingrid Bull-Engelstad, Jūratė Šaltytė Benth, Willy Aasebø, Nanna von der Lippe, Morten Reier-Nilsen, Ingrid Os, Knut Stavem

European Journal of Clinical Pharmacology, 2019 Jul. doi: 10.1007/s00228-019-02727-9. [Epub ahead of print]

3 ABBREVIATIONS

ADE	Adverse Drug Event
CCI	Charlson Comorbidity Index
CI	Confidence Interval
CKD	Chronic Kidney Disease
DDD	Defined Daily Dose
ESRD	End Stage Renal Disease
GFR	Glomerular Filtration Rate
HD	Haemodialysis
HRQoL	Health-related Quality of Life
KDOQI	Kidney Disease Outcomes Quality Initiative
KDQOL-SF	Kidney Disease Quality Of Life-Short Form
MCS	Mental Component Summary
MMAS-8	Eight-item Morisky Medication Adherence Scale
MMSE-NR	Mini-Mental State Examination - Norwegian Revision
MRCI	Medication Regimen Complexity Index
NICE	National Institute for Health and Care Excellence
NRC	National Research Council
OR	Odds Ratio
PCS	Physical Component Summary
PD	Peritoneal Dialysis
PIM	Potentially Inappropriate Medication
PIP	Potentially Inappropriate Prescribing
PPO	Potential Prescribing Omission
RCT	Randomized Clinical Trial
RRT	Renal Replacement Treatment

START	Screening Tool to Alert doctors to Right Treatment
STOPP	Screening Tool of Older Persons` Prescriptions
WHO	World Health Organization

4 INTRODUCTION

“To understand Homer, learn Greek. To understand old people, learn their language.” Bernard Isaacs

From breath-taking discoveries in treatments and ground-breaking basic scientific work to modernizing technology, supported by technological development in treatment, diagnostic and prevention, the world of medicine has leaped forward and made it today much more likely to reach old age.

The report “An Aging Word” shows the older population is projected to increase substantially, and actually expected to represent 17% of the total population in the world in 2050. New and less apparent challenges occur in the daily medical routine after many centuries of primarily focusing on what once seemed not treatable. Today we have evidence-based medicine and guidelines to guide treatment in acute and chronic diseases. However, older people rarely have only one treatable disease, many have extensive comorbidity which increases the age-related challenges.

Chronic kidney disease is common in the older adults, and is treated almost exclusively and extensively with pharmacological treatment. Five or more different pharmacological treatments are often simultaneously used, and the older adults are exposed to polypharmacy that can lead to negative health effects and the high complexity of treatment enhances the risk of inappropriateness. Any medication list should mirror the necessity and appropriateness of each treatment at any time. Specific screening tools such as Screening Tool to Older Persons’ Prescription/Screening Tool to Alert doctors to Right Treatment (STOPP/START) criteria might be essential and necessary. These tools, developed by experts and agreed by a consensus, enable doctors to identify potential inappropriateness in pharmacological treatments. The STOPP/START criteria are evidence-based comments to avoid commonly encountered potentially inappropriate prescribing and potential prescribing omissions.

Based on this background, we tested the STOPP criteria in a haemodialysis population. As a result, we extended and planned a prospective, randomized, clinical trial of older adults with advanced chronic kidney disease. We assumed that STOPP criteria would highlight potentially inappropriate medication and tested the hypothesis that a screening tool could improve appropriateness, medication adherence and health-related quality of life in this same patient group. Furthermore, we performed a cross-sectional study to investigate the complexity of medication regimen and medication adherence in older adults with chronic kidney disease.

5 BACKGROUND

“Age is not a diagnosis”. Atul Gawande

5.1 The aging population

Old age in humans describes the final stage of the normal life span. There is a wide range of definitions for “old age”. However, there is no general agreement on which age a person becomes old (1). An age of 65 years has been suggested as a threshold for old age, but that was from a time when the life expectancy was much shorter than it is today. For statistical, administrative and epidemiologic purposes the start of old age is frequently defined as between 60 and 65 years of age (2). Defining a person above 65 years as “old” may also not be an accurate term, because many will still have a very good physical health and an active social life.

Figure 1 Population in Norway in 1986 and 2019, data from Statistic Norway, www.ssb.no, yr. = years

Therefore, specialists in geriatric medicine allocate “old” people into sub-groups, i.e. “young-old” 65 to 74 years, “old” 75 to 84 years and “oldest old” beyond 85 years (3). An aging population is not only a Norwegian phenomenon (Fig. 1) as all estimation reports from Europe and around the world show an increase especially in the age group of 70 years and older. It is estimated that the European population above 65 years will increase to 24% in 2030, from 17% in 2008 (4). In the report, “An Aging World”, estimation suggests about 17% of the world total population in 2050 is represented by older adults (Fig. 2). Global life expectancy at birth is estimated to reach 76.2 years in 2050 compared to 68.6 years in 2015.

World Population by Age Group: 2015 to 2050

Source: U.S. Census Bureau, 2013; International Data Base.

Figure 2 World Population by Age Group (in millions): 2015 to 2050, published by US. Census Bureau

In this thesis, the terms “older” and “aging” will be used in their most common sense as synonyms for persons 65 years of age and older. Older adults are a heterogeneous group vary

from healthy and physical fit to frail with multiple chronic diseases. This heterogeneity describes a vulnerable aging population by acknowledging inter-individual differences in health, diseases and disabilities (5). Prescribing medications in this population has to take into consideration these individual variabilities and ongoing physiological changes related to aging including consequential pharmacodynamic and pharmacokinetic changes. These physiological changes expose the older adults to 1) drug-drug interactions, 2) drug-disease interactions, 3) potentially inappropriate medication (PIM) and 4) adverse drug effects (ADEs) (6-9).

5.2 Old age and chronic diseases

Old age is often associated with chronic diseases. There are several different definitions of chronic disease by World Health Organization (WHO), Centers for Disease Control and Prevention and MedicineNet. Bernell and colleagues advocate a simpler approach by using the definition from Merriam Webster Dictionary: “chronic” as something that is “*continuing or occurring again and again for a long time*” to describe chronic disease (10, 11).

The most common chronic diseases in older adults are hypertension, hyperlipidaemia, ischemic heart disease, diabetes mellitus, anaemia, atrial fibrillation and chronic kidney disease (CKD) (12). Older adults with comorbidities are facing diverse challenges compared to a person with one single chronic disease (13).

5.3 Age-related changes in physiology, pharmacokinetics and pharmacodynamics

Aging people face physiological changes in their organ functions and in body composition. The lean body mass, such as total body water, skeletal muscle, organ mass, and bone mineral tend to decrease, while total body fat increases and is redistributed more towards abdominal

adiposity than to peripheral adipose tissues (14). All organs undergo a slow age-related process of change that has an impact on pharmacokinetics and pharmacodynamics.

In a steady-state, the reduced function in an organ system may barely affect the organ, but the organ's functional reserve is reduced which increases the body's vulnerability to stress (Fig. 3) (15).

Figure 3 Organ system functional reserve. Geriatric individuals who appear to be physiologically “young” or “old” have lesser or greater rates of decline, respectively, than average, but they also have significantly different maximum capacities than young adults. Organ system functional reserve is defined as the difference between maximal (broken lines) and basal (solid line) function. Reprinted from Muravchick, S., *Anesthesia for the Elderly*, 2000

Pharmacokinetics is the process of absorption, distribution, metabolism and excretion of the drug and its metabolites (Fig. 4).

Figure 4 Passage of a medication through the body. Adapted from Comprehensive Clinical Nephrology, J Floege, RJ Johnson, J Feehally (ed), 4th edition.

The volume of distribution is the theoretical volume into which a given dose of a drug would have to be distributed in order to obtain the same plasma concentration (16). Older adults differ from younger individuals in the change in volume of distribution, and renal and hepatic clearance. Changes in renal and hepatic clearance increase the distribution volume of lipid-soluble medication, and decrease the excretion of lipid-soluble and water soluble medication. The consequence is prolonged plasma elimination half time. Furthermore, the medication sensitivity is changed due to pharmacodynamical changes (17).

Pharmacodynamics describes the connection between the concentration of a drug and its response, e.g. on cell receptors, drug-receptor interactions (depending receptor number

variations, receptor affinity, second messenger response and cellular response) and homeostatic regulations such as body temperature. Pharmacodynamics response differs between young and old adults, and between organs. The knowledge about age-related changes regarding pharmacodynamics is based primarily on cross-sectional studies, and is linked to the simultaneous age-related changes in pharmacokinetics. Older adults compared to younger persons may experience ADE with lower concentration of a given drug. This response may be attributed to the age-related decline of an organs baseline (Fig. 5A) or the differences in pharmacodynamics sensitivity (Fig. 5B) (18).

Figure 5 Baseline (A) and sensitivity (B) differences between young and elderly adults. E_0 = baseline effect before drug administration when drug concentration are zero. Reprinted from publication The American Journal of Geriatric Pharmacotherapy, 2007, Bowie and Slattum, PW. Pharmacodynamics in older adults: a review, with permission from Elsevier.

5.4 Chronic kidney disease

CKD is common in older adults. The definition is: established kidney damage for ≥ 3 months, with or without a decreased glomerular filtration rate (GFR), that can lead to a decrease in GFR, manifested by either: a) pathologic abnormalities, or b) markers of kidney damage, including

abnormalities in the composition of the blood or urine, or abnormalities in imaging tests, or GFR <60 mL/min/1.73m² for ≥ 3 months with or without kidney damage (19). The most common causes of CKD are diabetes mellitus type I and II, hypertension, glomerulonephritis, interstitial nephritis, polycystic kidney disease, post renal diseases and recurrent kidney infections.

CKD can be divided into five stages based on GFR according to Kidney Disease Outcomes Quality Initiative (KDOQI) guidelines (Table 1). The most severe, or advanced stage of CKD is also called in the literature “end stage renal disease” (ESRD). Advanced CKD defines all patients with an estimated GFR <15 mL/min/1.73m², but not necessarily with uremic symptoms or need for a renal replacement therapy (RRT). ESRD in literature and in the clinic is often associated with patients in advanced CKD receiving RRT. Until now a good clinical definition of ESRD has been lacking, but it is suggested that two criteria must be fulfilled to define ESRD: establishing of uremia and the need for RRT (20). The term “uremia” means “urine in the blood” and was first proposed by Piorry in 1840 (21), describing the clinical condition in patients with renal failure. No exact diagnostic guideline exists to diagnose symptomatic uremia, but it is easy to recognize bedside by nephrologists, at least in its final stages (20). Therefore, the National Institute for Health and Care Excellence (NICE) guidelines recommend the follow up of CKD stage 4 and 5 by a nephrologist (22).

Table 1 Classification of chronic kidney disease

Stage	Description	GFR (mL/min/1.73m ²)	Related terms
1	Kidney damage with normal or ↑ GFR	≥ 90	
2	Kidney damage with mild ↓ GFR	60-89	
3	Moderate ↓ GFR	30-59	Chronic renal insufficiency, early renal insufficiency
4	Severe ↓ GFR	15-29	Chronic renal insufficiency, late renal insufficiency, pre- end stage renal disease (ESRD)
5	Kidney failure	< 15	Renal failure, uremia, ESRD

Reprinted from publication Kidney International, Levey, Andrew S. et al. Definition and classification of chronic kidney disease: A position statement from Kidney Disease: Improving Global Outcomes (KDIGO), with permission from Elsevier

5.4.1 Changes in pharmacokinetics and pharmacodynamics

It is important to understand the pharmacokinetic changes in advanced CKD. The absorption is impaired, but difficult to quantify; however, this is only relevant for a limited number of medications. Gastrointestinal edema can reduce oral absorption. Phosphate binders can interact with medications minimizing also the oral uptake. The distribution of medicines can be increased for water-soluble substances due to edema. Metabolism or hepatic clearance is affected by renal impairment through the downregulation of cytochrome P450 and decreased activity of drug transporter (23). The kidney is the other main organ for excretion of medications. The impairment of the kidney function influences directly the renal clearance. Haemodialysis (HD) and peritoneal dialysis (PD) have also an impact on the removal of medications. Factors influencing this removal are dialysis frequency, blood flow, pore size and membrane type of the filter, and dialysis duration. HD can be more efficient than PD in removal of medications. Furthermore, CKD may alter the pharmacodynamics response on cell

membrane, which will affect the clinical response, i.e. platelet function is impaired leading to higher risk of bleeding under antiplatelet therapy.

5.4.2 Prevalence

The American National Health and Nutrition Examination Survey reported an increased prevalence of CKD from 10% in the period 1988-1994 to 13% in 1999-2004 (24). The increase is caused partly by an aging population, and partly by a higher prevalence of hypertension and diabetes mellitus (25). On the other hand, data from UK, published 2014, showed a decline of the prevalence of CKD, with uncertainties about the causes of this decline (26). Regarding the prevalence of CKD in the Norwegian population there is a significant rise of high risk CKD among the persons older than 75 years of age over a 10-year period. (27). Furthermore, data from the Norwegian Renal Registry showed an increasing number of older patients receiving RRT treatment (Fig. 6).

Figure 6 Prevalence in renal replacement therapy, sorted by treatment mode, for patients >65 years from Norway 2005-2018, with courtesy of Anders Åsberg.

An estimation of the future burden of CKD in the United States suggests an increase from 13% in 2015 to 17% in 2030 (28) and a further climb on the ranking list of causes of death.

5.4.3 Treatment modality

Any treatment of advanced CKD encompass fluid balance, treatment of anaemia, correction of acidosis and hyperkalaemia, including management of symptoms, preserve residual renal function and blood pressure, as well the phosphorous/calcium balance and giving diet advice.

5.4.3.1 *Conservative (predialytic) treatment*

Conservative management is an alternative treatment option in advanced CKD without initiating RRT. This management focuses mainly on preventing a worsening of the kidney function and on early detection of complications. The focus is a pharmacological support therapy; however, patients can still start dialysis treatment if symptoms and CKD condition worsen, and if patients wish to start dialysis treatment. Conservative treatment can slowly change to palliative management as the kidney function declines. The transition from conservative to palliative management may go unnoticed as both treatments focus on a person-centred management and symptom management to maximize the quality of life (29).

5.4.3.2 *Renal replacement therapy*

RRT is often necessary when patients reach advanced CKD with symptomatic uraemia. Under RRT, we include the following treatment modalities: dialysis and kidney transplantation.

5.4.3.2.1 *Dialysis*

When patients reach the advanced stage of CKD and are showing symptoms of increased uraemia and subjective non-wellbeing, dialysis treatment can be initiated. Today there are two forms of dialysis treatment: HD and PD. Dialysis treatment is based on two physiological

principles: diffusion and osmosis. Blood and dialytic fluid are separated by a semipermeable membrane (Fig. 7).

Figure 7 Physiological principles of dialysis.

HD treatment is mainly performed in a hospital, attending by a specialised nurse and a nephrologist, although self-administered home haemodialysis is increasing in the recent years. HD is extra-corporeal and requires an access to the vessel system, either by a catheter or by an arteriovenous fistula. The fistula is a special vessel created through surgery by connecting an artery with a vein. It requires a dialysis machine and a special filter, called artificial kidney or dialysis filter, to purify the uremic blood. The dialysis machine pumps the blood from the patient through this filter. The filter consists of two parts divided by a semipermeable membrane. One part is for the patient's blood and the other is for the fluid, called dialysate, which counter-current flows in the filter. The semipermeable membrane allows diffusion and osmosis of electrolytes and small to moderate-size molecules. This procedure is performed in a hospital dialysis centre, three times per week for four hours, whereas home-HD is more

frequent, and is done by the patients` themselves, or in future by trained home attending nurses (Fig. 8).

PD is a form of dialysis that is performed outside of hospitals, mainly at patients` homes, either by the patients` themselves or by a trained home attending nurse. PD is para-corporeal and starts with adding dialytic fluid via a peritoneal catheter into the abdomen (Fig. 8). In this treatment, the semipermeable membrane is actually the vessels in the peritoneum. The dialytic fluid has to dwell for a pre-defined time period in the abdomen, allowing diffusion of the small molecules and transport of fluid by osmotic pressure to happen. After the dwell time the dialytic fluid will be removed from the abdomen, and a new cycle starts again. PD can be performed manually or by an automated process. Patients, home attending nurse or nurses from homes will be trained in at least one of these treatment forms. Patients with PD will meet up in the outpatient clinic regularly every four to six weeks.

Thanks to the courtesy of Torun Nilsen

Figure 8 Illustration of peritoneal dialysis (left) and of haemodialysis (right) treatment

5.4.3.3 *Kidney transplantation*

The first kidney transplantation took place in Norway in 1958, initially with short survival times. Through the implementation of new immunosuppressive agents in 1960s the survival after transplantation improved dramatically. Kidney transplantation often removes symptoms of the renal failure, restores the kidney function above 60ml/min/1.73m² and increases survival and quality of life (30-32). In Norway, there is so far no upper age limit for patients, and the published results indicated that age *per se* seems not to be a contraindication (33). Each case is treated individually and needs to be well considered for its benefit or harm on an individual basis. After receiving a kidney transplant all patients are still dependent on long-term pharmacological therapy related to immunosuppression and comorbidities.

5.5 Prescribing medication

Today's definition of "medication", we find with the WHO stating: "*Traditional medicine is used in the maintenance of health as well as in the prevention, diagnosis, improvement or treatment of physical and mental illness*" (34). Prescribing of medication in older adults aims to 1) cure disease, 2) eliminate or reduce symptoms and 3) improve functional capacity.

In a 5-year U.S. survey, prescriptions marginally increased from 84% to 88% among adults aged 62 to 85 years. Further, the prescriptions of at least five medications or more substantially increased from 31% to 36%, and about 15% of the older adults were at risk for a major potential drug-drug interactions (35).

The Norwegian Institute of Public Health published an annual rapport based on data from the national prescription database in Norway. In 2017, 92% of the people of 65 years or older had a least one medication dispensed on prescription. Adults over 90 years used most both regarding

multiple medications and higher quantity. People over 65 years of age were dispensed about 48% of the total number of defined daily doses (DDD) on prescription in 2017 (Table 2) (36). Thus, with the increasing proportions of older people worldwide there is a concern for quality and safety regarding prescribed medicines (5, 36, 37).

Table 2 Number of individuals having a prescription dispensed in 2017 in the major ATC groups and the corresponding sales in total number of DDDs in Norway

	Total number of individuals	Proportion (%) ≥ 65 years	Total million DDDs	Proportion (%) ≥65 years
A- Alimentary tract and metabolism	1 081 420	39	357	47
B- Blood and blood forming organs	699 782	61	252	65
C- Cardiovascular system	1 117 743	55	790	64
G- Genito-urinary system and sex hormones	876 600	24	213	20
H- Systemic hormonal preparations, excl. sex hormones and insulins	463 794	39	78	43
J- Antiinfectives for systemic use	1 177 724	24	32	36
M- Musculo-skeletal system	963 353	26	96	42
N- Nervous system	1 450 941	31	380	35
R- Respiratory system	1 374 981	22	297	33
Total (all ATCs)	3 688 097	23	2572	48

ATC=anatomical therapeutic chemical, DDD=defined daily doses, Proportion (%) in the age group 65 years and older

5.5.1 Polypharmacy

Polypharmacy is common and increasing particularly in the elderly, often due to comorbidities. The use of medication is high according to reports from UK, Denmark and Sweden (38-40). In the literature, however, there is a large heterogeneity regarding the definition of polypharmacy. It has been defined as number of drugs, as use of inappropriate drugs or as the concurrent use of multiple drugs (41). Merriam Webster dictionary medical defines: “*the concurrent use of*

multiple medications by a patient to treat usually coexisting conditions and which may result in adverse drug interactions” (42). Hence, there is no consensus on the definition. Most commonly, polypharmacy is defined numerically, as the use of five or more drugs daily (43).

Many studies have shown an association between polypharmacy and negative outcomes such as falls, mortality, ADE, prolonged hospitalization or readmission to hospital soon after discharge (44). Additionally, polypharmacy is associated with reduced medication adherence and increase the risk for harm and inappropriateness (45-47).

Older adults with CKD reach over time an average use of medication up to 10 to 12 medications daily (48, 49). Each medication mirrors the patient’s variety of medical conditions. The numbers of prescriptions do not necessarily reflect clinical appropriateness. If multiple prescriptions are based on evidence-based medicine or established guidelines, older patients could benefit from appropriate polypharmacy (Table 3). However, many physicians do not recognize the hazards with polypharmacy, and its risk of prescription-related problems and inappropriateness. That’s why it is of importance, when physicians meet older adults with complex and multiple chronic diseases that they keep a good overview while accepting that it is a time consuming and overwhelming challenge.

Table 3 Advantages and disadvantages with polypharmacy

Benefits	Improved disease management Optimised medicines management
Outcome	Reduced risk of disease complications and mortality Evidence based prescribing
Harms	Increased drug interaction Increased risk adverse drug outcomes Inappropriate prescribing and medication errors Lack of monitoring
Consequences	Electrolyte disturbances; potentiation of drug effects Patient morbidity and mortality Unscheduled contacts with healthcare providers Safety risks to patients

Molokhia M, Majeed A, 2017, BMC Family Practice

5.5.2 Potentially inappropriate medication

Potentially inappropriate medication (PIM) encompasses all medication where the risks outweigh the clinical benefit, lacking a clear evidence-based indication, carrying a higher risk for causing ADE or economic costs (8, 50, 51). PIMs include medications that increase the risk of drug-drug or drug-disease interactions, incorrect dosage, or frequency and duration (overprescribing), not using safer alternatives (misprescribing), as well as clinically indicated medicines, but missing (underprescribing) (5, 52). The prevalence of PIMs is high in older patients, and has been associated with ADE, hospitalisation, reduced mobility and increased mortality, and resource wastage (53-55).

5.5.3 Assessment of potentially inappropriate medication

Over the last decades many measurements have been developed to identify inappropriateness, both implicit (judgement-based) and explicit (criteria-based) measurement tools. For implicit criteria, the best known one is “Medication Appropriateness Index” (52). Implicit measurements require solid knowledge of medication pharmacokinetics and

pharmacodynamics, and rely on an expert's judgement to assess the appropriateness of medications. Explicit or criteria measurements have been developed through an expert consensus. The expert groups create lists of medication that should be avoided in older people both for specific comorbidities and in general. The most known explicit criteria are the Beers criteria (51) or the Screening tool of Older Persons' Prescriptions (STOPP) and Screening to Alert doctors to Right Treatment (START) criteria (56). Criteria-based use of medication has been implemented in a number of countries (57-59), including Norway (60). Explicit criteria are easy to use in clinical practice and research, partly because the medication list is limited.

In this thesis we used the STOPP/START criteria. With the STOPP criteria it is possible to identify potentially inappropriate prescriptions (PIPs), whereas the START criteria identify potential prescribing omissions (PPOs). PIPs include overprescribing or misprescribing, while PPOs comprise underprescribing.

5.5.4 Prescribing of PIPs in older adults with CKD

The prevalence of inappropriate prescriptions in patients with CKD ranges from 9% to 81% in hospitals and outpatient clinics (61-63). Most frequent predictors for inappropriate prescription are patients' characteristics such as age and sex, number of medication and comorbidities (62, 64).

5.6 Medication regimen complexity

Polypharmacy can result in a complex medication regimen. Patients use tablets, capsules, injections, creams, drops, sprays and patches with each of them having own instructions, dosages and frequencies. To define the complexity of any medication regimen has been a challenge due to lack of consensus in the definition. In the literature, complexity of a medication regime was often definite by reference to the number of medications, but it can also take into

account the number of times per day or “doses” taking by the patients (multiple doses schedule), ignoring patient-related factors such as dexterity and cognitive function. In 2004, Georg et al published Medication Regimen Complexity Index (MRCI) based on a medication complexity index tool developed in 1988 (65). MRCI is not a disease-specific tool, and aims to assess objectively the complexity of any regimen. Furthermore, several studies reported a positive link between a higher MRCI score and readmission to hospital, ADE and mortality (66-69).

5.7 Medication adherence

The adherence project of WHO (2003) described adherence in the context of long-term therapy as follows: *“the extent to which a person’s behaviour – taking medication, following a diet, and/or executing lifestyle changes, corresponds with agreed recommendations from a health care provider”* (70). The definition of adherence is not synonymous with terms such as compliance or concordance. The definition of compliance is *“the extent to which a person’s behaviour coincides with medical and health advice”* (71, 72). Behind “compliance” is the paternalistic authority of clinicians, while patients have a more passive role. Over the years, the patient role has changed by having access to internet, social media and libraries. An alternative was offered in 1997, so called “concordance”. Concordance” was *“based on the notion that the work of prescriber and patient in the consultation is a negotiation between equals and that therefore the aim is a therapeutic alliance between them. Its strength lies in a new assumption of respect for the patient’s agenda”* (73). This ideal implies that patients take active responsibility in management, but it is not clear if everyone likes to do so.

The term “adherence” is most commonly used. This term recognises patients’ beliefs, barriers and problems, but at the same time includes clinicians’ behaviour. Adherence related to taking medication is not only based on pharmacological components, but also influenced by individual

factors such as psychological, social and economic factors (72) (Fig. 9). There is another factor influencing adherence and self-management so called health literacy. Health literacy “*entails people’s knowledge, motivation and competences to access, understand, appraise and apply health information*” (74). In a recent review, the authors concluded that because there was a limited health literacy in patients with CKD (75), this could affect their medication adherence.

Figure 9 The five dimensions of adherence. WHO, published with permission

Several studies have described different behaviour theories regarding taking medication with its focus on patients’ attitudes, beliefs and intentions (76). The model, “common sense self-regulation model”, has been used in many diseases to describe or predict patients’ adherence to treatment (76). Horne and colleagues extended this theory by showing that the patients’ belief of necessity and their wider general concerns over medication are important (77).

In the literature, adherence is often presented in two possibilities: adherent/non-adherent. Furthermore, non-adherence can be classified as intentional or unintentional. Intentional is based on behaviour, opinions and beliefs from each person. Intentional non-adherence includes avoiding to take medicine due to ADE or not achieving the benefit the patients has hoped for, disappearing of symptoms without realizing that discontinuation will bring symptoms back, not

picking up medicine from the pharmacy and personal opinion regarding medicines or doctors. In addition stockpiling, drug holidays, white-coat adherence belongs to this group of non-adherence (78-81). Unintentional adherence is forgetting to take the drugs due to cognitive impairment, difficulties to open bottles and boxes, visual problems, as well as misunderstanding between clinician and patients (81).

5.7.1 Assessment of medication adherence

There is “no gold standard” in how to assess medication adherence. Many studies have used 1) self-reported questionnaires), 2) patient medication records, 3) direct questioning or 4) electronic devices (82).

Subjective ratings by using standardised self-reported questionnaires or direct questioning have shown that patients overestimate adherence (83). For example, people unequivocally admitting to following recommendations will ordinarily describe their behaviour, whereas patients not following the advice they have been given, may inaccurately state their behaviour.

Objective measurements are electronic medical monitor devices and patients electronic medical records. By counting tablets or updating prescriptions on the pharmacy database there is an indirect feedback on adherence, but still the pitfall of overestimation. Electronic medical monitor devices have the advantage that they register the time and date a container was opened, but cannot confirm if the medicine was really taken. These devices are expensive and so far have mostly been used in study populations (84). Another objective alternative measurement is the biochemical measurement of medicine, both for the drug itself and its metabolites. The disadvantage of using this method is the inter-individual difference of absorption, metabolism and excretion (85).

5.7.2 Medication adherence in older adults

A huge variety of factors and diversity of study designs make a comparison of adherence-studies difficult. Of interest are the conclusions of Sturges and colleagues when comparing three methods to assess adherence. They found that the use of two methods, patients' medical records and direct questioning, was a good way to identify non-adherence (86). As said earlier adherence cannot in reality be divided into adherent/non-adherent. There is a huge "grey-zone" and many factors to be considered in an older population (Table 4).

Table 4 Factors which may affect medication adherence in older adults

Drug-related factors	Administration regimens
	Number of drugs
	Adverse effects
	Packaging
Patient-related factors	Changing physiology
	Multiple morbidities
	Cognitive ability
	Health beliefs
	Psycho-social profile
Other factors	Patient-prescriber relationship
	Access to medication (insurance, restrictive formularies)
	Social support

Hughes CM, 2004, Drugs Aging, published with permission from Springer.

These predisposing factors are risk factors for older adults to be non-adherent, and make the research field a minefield for poor design studies and bias. Another challenge is the cut-off for "old". The review from 1998 concluded that age *per se* was not an important predictor to negative medication adherence (87). Weintraub and colleagues showed that older adults altered dosage of treatment on a rational basis to avoid side effects, but managed to achieve therapeutic effects (88). Still, there is the possibility that non-adherence gives a higher risk for poor disease control and polypharmacy. There is a wide range of studies focused on different diseases, conditions, and possible factors to describe adherence of older adults. Basically, the conclusion

is that all factors, in addition to age, need to be considered when trying to understand adherence in an older population (84).

5.7.3 Medication adherence in CKD patients

Adherence in CKD patients has been a topic in a variety of publication over the past decades. Patients with CKD need to be adherent not only to prescribed medications, but to recommended diet and fluid restriction. Studies of adherence in CKD patients have reported a wide range of results depending on the angle of study question, the definition of adherence, and the methods chosen for evaluation. Thus, it is no surprise that non-adherence was found in 13% to 99% in HD patients, 4% to 85% in PD patients, and 18% in predialytic patients (89-91).

Factors identified as influencing patients with CKD can be grouped in the following categories: patient-related factors, socioeconomic status, psychological factors, therapy related factors and disease related factors (92). Mehta and colleagues' report captured three considerations when it comes to adherence in patients with CKD: logistics, benchmarking the need for medication and the quality of the patient-physician relationship. First, the practical side of having a chronic disease with complex medication regime and many appointments (logistics) challenges patients. Second, the individual patients prioritize medication on how important it is to relief symptoms, the knowledge of side effects or the lack of understanding (benchmarking). Third, the lack of continuity, time and trust, and the acknowledgement of patients' wish of involvement and their concern or belief regarding medication decisions might disturb the patient and physician relationship (93).

6 AIMS OF THE THESIS

The overall aim of this thesis was to present the medication use in older adults with advanced CKD with particular focus on medication appropriateness, medication complexity and medication adherence.

The specific aims were:

- To describe the prevalence of potentially inappropriate medications by using the STOPP criteria and the Beers criteria in a haemodialysis population
- To determine risk factors such as age, sex, number of medications and comorbidity for potentially inappropriate medications according to the STOPP criteria
- To identify factors associated with medication complexity and medication adherence in older adults with CKD
- To determine the association between medication complexity and medication adherence
- To identify potentially inappropriate prescribing and potential prescribing omission by using the STOPP/START criteria in older patients with CKD stage 5, either treated conservatively or by dialysis as a randomized clinical trial study design
- The hypothesis was that a medication review with the STOPP/START criteria would lead to reduction of inappropriate prescriptions and number of medications, and improve the health-related quality of life and medication adherence

7 MATERIALS AND METHODS

“We can follow the data, be objective, and be human at the same time.” Jonathan Foley

7.1 Potentially inappropriate medications with the STOPP criteria (Study I)

7.1.1 Study population

This cross-sectional study included patients over 65 years of age from the haemodialysis centre at Akershus University Hospital from July through December 2012. A total of 102 patients received HD treatment during the inclusion time. And 52 patients over 65 years of age consented to participate.

7.1.2 Data collection

Data on age, sex, number and types of prescribed medications, cause of kidney disease, comorbidity, time on haemodialysis and dialysis treatment quality index (quantified as the urea clearance [Kt/V]) were collected from the electronic medical records. Additionally, all participants answered a pre-specified list of questions regarding experienced side effects such as vertigo and chronic obstipation.

7.1.3 Instruments

7.1.3.1 Beers criteria

The Beers criteria, developed by the Delphi consensus based on nursing home residents and published in 1991, are the first explicit criteria for identifying inappropriate prescribing (51). From the 1990s, the Beers criteria were expanded and updated (50, 94, 95). The criteria focused mainly on inappropriate prescribing and less on underprescribing of clinically relevant medications. In 2015, these criteria were updated again to be clinically in ambulatory, acute, and institutional settings (6). The 2015 update added two areas of recommendations: 1)

medications and their dosage adjustments in cases of kidney impairment, and 2) drug-drug interactions. Moreover, the panel performing the update stated that the Beers criteria cannot account for all individuals and special populations (6). During the last decades, these criteria have been worldwide to study prescribing management and its effects on the associated health outcomes, falls and mortality (96-99). The Beers criteria is presented in the Appendix I.

7.1.3.2 *Charlson Comorbidity Index*

We assessed comorbidities using the Charlson Comorbidity Index (CCI) (100), which consists of 19 weighted comorbidity items. All items were added to produce a total score (Table 5). The original version adds 1 extra point to the total score for each decade >40 years of age. We did not add age into our total CCI score because age was a separate variable in the statistical model. Additionally, the CCI has previously been validated in dialysis patients (101).

Table 5 Charlson Comorbidity Index

Score	Condition
1	Myocardial infarction (history, not ECG changes only)
	Congestive heart failure
	Peripheral vascular disease (includes aortic aneurysm ≥ 6 cm)
	Cerebrovascular disease: CVA with mild or no residua or TIA
	Dementia
	Chronic pulmonary disease
	Connective tissue disease
	Peptic ulcer disease
	Mild liver disease (without portal hypertension, includes chronic hepatitis)
	Diabetes without end-organ damage (excludes diet-controlled alone)
	Hemiplegia
2	Moderate or severe renal disease
	Diabetes with end-organ damage (retinopathy, neuropathy, nephropathy, or brittle diabetes)
	Tumour without metastases (exclude if > 5y from diagnosis)
	Leukaemia (acute or chronic)
3	Lymphoma
	Moderate or severe liver disease
6	Metastatic solid tumour
	AIDS (not just HIV positive)

ECG: electrocardiogram; CVA: cerebrovascular accident; TIA: transient ischemic attack; AIDS: acquired immunodeficiency syndrome; HIV: human immunodeficiency virus

7.1.3.3 STOPP/START criteria version 1

These criteria were developed and validated in 2004 through a Delphi consensus process by 18 experts in geriatric pharmacotherapy. They include indicators to identify important drug-drug and drug-disease interactions (56). The STOPP/START criteria exhibit a high sensitivity for detecting potentially inappropriate prescriptions and good inter-rater reliability with a median κ -coefficient of 0.93 for STOPP criteria and 0.85 for START criteria (102). The criteria consist of 65 STOPP items that highlight medications defined as overprescribed or misprescribed. Each item comes with a brief clarification as to why it is inappropriate or not. In this study, we applied only the STOPP criteria to the medication lists of all participants. The STOPP criteria version 1 is presented in the Appendix II.

7.1.4 Statistical analysis

We used mean and standard deviation for normal distributed data, median and range/min-max for skewed data for the descriptive statistic. Histograms visualised the data distribution. We compared the continuous variables with the Student's t-test and the categorical variables with the χ^2 -test or Fisher's exact test.

We assessed the inter-rater agreement in assessments between the two raters using the STOPP and Beers screening tools with % agreement (proportion of cases in which the raters agree) and kappa coefficient for 2 x 2 tables. Strength of reliability coefficients was defined using the following nomenclature (103): <0.20 poor, 0.21-0.40 fair, 0.41-0.60 moderate, 0.61-0.80 good, 0.81-1.00 very good.

7.2 Medication complexity regimen and adherence (Study II)

7.2.1 Study population

The study population was a prospective cohort of 180 patients included from June 2015 to January 2017. All patients were ≥ 65 years of age and had advanced CKD with estimated GFR $< 15 \text{ ml/min/1.73 m}^2$ in either conservative/predialytic or dialysis treatment. The estimated GFR was calculated by each laboratory using the Chronic Kidney Disease Epidemiology Collaboration equation. We recruited from three different Norwegian nephrology centres (Akershus University Hospital, Oslo University Hospital, Ullevål, and Vestre Viken, Drammen Hospital). All dialysis patients were asked to participate during a scheduled dialysis session. Patients with conservative/predialytic treatment were recruited during scheduled ambulatory visits. We excluded patients with

- Severe hearing loss or visual impairment,
- Diagnosed dementia or known Mini-Mental State Examination - Norwegian Revision (MMSE-NR) score < 23 , or
- Unsatisfactory knowledge of the Norwegian language.

7.2.2 Enrolment and data collection at baseline

The same investigator (KP) visited the hospitals and introduced the study design prior to enrolment. The selection of patients was discussed with attending physicians or research nurses at the respective hospitals. All patients received oral and written information, that included the aims of the study, and they signed written consent. The next step was a semi-structured interview with closed and open questions about prescribed medications, over the counter medications, side effects, and administration. As part of the semi-structured interview, each

patient performed a cognitive evaluation. At the end of the interview, each patient received questionnaires about medication adherence and HRQoL to be answered at home. The answered questionnaires were sent by postage with prepaid envelopes to the interviewer. In cases in which questionnaires were not returned, the investigator contacted the patient directly once by phone.

7.2.3 Instruments

7.2.3.1 *Charlson Comorbidity Index*

As described in Study I.

7.2.3.2 *Medication Regimen Complexity Index*

MRCI was used to assess medication regimen complexity. This tool consists of 65 items and is designed to quantify the complexity of any prescribed medication regimen (104). There are three sections: A (dosage forms), B (dosing frequency), and C (additional direction), with 32, 23, and 10 items, respectively. We coded each medication item according to the weighted scoring system and added the aggregate score to the total score (104). There is no maximum score because the total score increases continuously when adding dosage forms, dosing frequency, or additional directions. As of now there is no definition of the line between a “high” and low” score. Validation of the index was done using a population with chronic obstructive pulmonary disease with inter-rater and test-retest reliabilities on MRCI of ≥ 0.9 (104). A few studies have used MRCI scores in chronic diseases, including CKD (105, 106). The MRCI instrument is presented in Appendix III.

7.2.3.3 *Mini Mental State Examination – Norwegian Revision*

The MMSE-NR is based on the original Mini-Mental State Examination published by Folstein et al in 1975 (107). Folstein and colleagues developed a screening tool to evaluate a person’s cognitive function using short questions. The reliability and validation were tested. In Norway,

many versions of this instrument were available with a high diversity of scoring guidelines which made the re-testing challenging. At the beginning of 2000, Strobel and Engedal developed the MMSE-NR with standardized questions and guidelines for testing and scoring (108). The total score is linear from 0 to 30, with 30 representing the best result. The results are a guide to evaluating the cognitive function of patients without establishing a diagnosis. The test recommended the following interpretation of scores: A total score of 28 or higher shows no indication of cognitive impairment. A total score from 25 to 27 could indicate cognitive impairment, but further tests are recommended. A total score of 24 or less indicates cognitive impairment, although other explanations such as low motivation and, problems with reading or writing are possible influential factors. The MMSE-NR is presented in the Appendix IV.

7.2.3.4 Eight-item Morisky Medication Adherence scale

Morisky and colleagues developed and published a tool that assesses patients' behaviour to study medication-taking in a hypertensive population. It began as a four-item self-reporting measurement and was used widely for numerous chronic diseases (109). Some years later, four additional items were added to cover more factors that influence medication-taking behaviour. The internal consistency was 0.83 in a hypertensive population (110-113). The eight-item Morisky Medication Adherence Scale (MMAS-8) was used in RCTs investigating heart disease, hypertension, diabetes mellitus and cancer. This tool has also been applied in CKD populations (114, 115). The questionnaire has seven items requiring "yes" or "no" answers and one item with an ordinal Likert scale where "0" is a low score, and "4" is the best score. Items 5 and 8 were transformed in accordance with the given scoring algorithm. All items were combined in a total score that graded adherence on a 0 to 8 scale with "0" signifying no adherence and "8" meaning high adherence. Furthermore, the total score can be interpreted as

low (<6), moderate (6 to <7.9), or high ($=8$) adherence (111). The MMAS-8 is presented in the Appendix V.

7.2.4 Statistical analysis

The descriptive statistics were performed as described in Study I.

In this study, we combined all dialysis patients, both HD and PD, in one group due to the small number of PD patients. The MMSE-NR scores were dichotomised using a cut-off point for normal cognitive function (≥ 28) (108). The CCI total scores were divided into three categories: 2-3, 4-5, and >5 . Phosphate binders were divided into five categories: none, and the quartile of phosphate binder use (in g/day).

For the multivariate analysis, the selection of explanatory variables was based on the literature, perceived clinical relevance and the use of a directed acyclic graph. Furthermore, correlations between variables were tested with Spearman rank correlation. There was a strong correlation between pill burden and MRCI, thus, we chose to include only the use of phosphate binders in the regression analyses. To assess the association between the selected variables and medication adherence, we used multivariable ordinal logistic regression analysis with the dependent variable, MMAS-8 scores, in ordinal categories: low (<6), moderate (6 to <8), and high ($=8$). Ordinal logistic regression analysis can be interpreted as the odds of being in groups greater than k versus being in groups less than or equal to k , where k is the level of the response variable. This odds ratio is constant. Thus, the proportional odds ratio represents the odds of moderate/high adherence (MMAS-8 score ≥ 6) versus low (<6) and the odds of high adherence (MMAS-8 $=8$) versus moderate/low (MMAS-8 <8). We tested the proportional odds assumption with the Brant test (116).

7.3 Effectiveness of using the STOPP/START criteria (Study III)

7.3.1 Study population

The same study population was used as in “Medication complexity regimen and adherence (Study II)”. The collection of data, the question of participation, the semi-structured interviews, the giving out of questionnaires, and the follow-up routine were all done by one investigator to ensure consistency and uniformity.

7.3.2 Randomization and intervention

The assignment to either the intervention or control group was 1:1 based on a computer-generated allocation (Fig. 10). The allocation for the assignment was stored in sealed envelopes and prepared by another investigator (KS). The sealed envelopes were opened after the first interview with the participants was carried out. Throughout the entire study, the participants were blinded regarding the allocation, but attending physicians and investigators were not blinded.

A medication review with the STOPP/START criteria was applied to each group by one investigator. In the intervention group, recommended medication changes were added to the electronic medical records. The attending physician could choose to implement the recommendations or not. For the control group, no notifications of recommendations were recorded, but in cases of severe inappropriateness, the investigator would note such in the medical record or inform the attending physician.

7.3.3 Follow-up after six months with data collection

After six months, each participant was invited to a second semi-structured interview regarding medication and changes that occurred during the follow-up time. We asked specifically about

new comorbidity and number of hospitalizations. Each participant received the same questionnaires to be complete at home and returned to the investigator by mail.

Figure 10 Flowchart of participants for Study II and III.

7.3.4 Instruments

7.3.4.1 *Health-related quality of life*

Health related quality of life was evaluated using the Kidney Disease and Quality of Life Short Form version 1.3 (KDQOL-SF). This self-reported questionnaire has been translated into Norwegian and validated for dialysis and kidney transplant patients (117, 118). The internal consistency or reliability, assessed with Cronbach's alpha, ranged from 0.74-0.91 (119). All coding and recording of data was conducted according to the "Manual for Use and Scoring" (120). The overall score after transformation ranged from "0" (worst possible health) to "100" (best possible health). Two summary scores, the physical component summary (PCS) and the mental component summary (MCS), can be obtained from the subscales (120). The subscales consist of the following items: physical functioning, role limitations due to physical health, bodily pain, general health, vitality, social functioning, role limitations due to emotional problems, and mental health.

For the adjustment of PCS and MCS we used the general U.S. population norm with a mean of 50 and standard deviation of 10 (121). In cases of missing items, we followed the recommended "half-rule", meaning that the missing items were replaced by the mean of the answered item on a subscale if at least 50% of items on that subscale had been answered. The KDQOL-SF is presented in the Appendix VI.

7.3.4.2 *STOPP/START criteria version 2*

Due to a widening of knowledge in evidence-based medicine, less accurate or relevant criteria, and lack of clinical importance, the criteria were updated to version 2 in 2014 by a panel of 19 experts from 13 European countries. The updated version consists of 114 criteria: 80 STOPP criteria and 34 START criteria (122). Presence of PIPs and potential prescribing omissions (PPOs) identified by the STOPP/START criteria version 2 were associated with readmission

and mortality (123). The structure of the STOPP/START version 2 is the same as that of version 1. The Norwegian version of the STOPP/START criteria is presented in Appendix VII.

7.3.5 Statistical analysis

The descriptive statistics were performed as described in Study I.

Before conducting the study, we calculated the sample size for the study population using medication adherence as a continuous variable (110). We needed to detect a difference of 0.5 standard deviation in the MMAS-8 score between the intervention and control groups with a statistical power of 80% and at a significance level of 5%. Thus, each group would need 63 participants.

In this population, we observed a high percentage of missing data. To make use of as much information as possible, we decided to use mixed model analyses. No imputation was done. The differences between the allocation groups at follow-up was calculated for the continuous variables (number of medications, PCS and MCS scores) using a linear mixed model. The number of PIPs and PPOs according to the STOPP/START criteria and medication adherence was dichotomous meaning for PIPs and PPOs (none versus ≥ 1) and medication adherence (adherent versus non-adherent). The statistical analysis model was the generalized linear mixed model for the binary variables. These models cover the fixed effects regarding measurement time point (baseline or six months) and the interaction between the time points and allocation groups. The interaction term in such a model quantifies differences between groups at follow-up adjusted for baseline. Random effects for patients were included. The models were estimated for cases with observations available at baseline. As sensitivity analyses, we conducted complete-case analysis using longitudinal analysis of covariance for the continuous outcomes

(fixed effects for group and baseline values) and binary logistic regression analysis for the dichotomous outcomes (fixed effects for group) (124).

All statistical analyses of all three studies were performed using STATA 15.1 (StataCorp, College Station, TX, USA) or SAS version 9.4 (SAS Institute Inc., NC, USA). All tests were two-sided with any result reaching p-value below 0.05 being defined as statistically significant.

7.4 Ethics

All three studies were conducted in line with the Helsinki Declaration. Prior to participation all eligible patients received information, both orally and written, about the study aim, the handling of personal data during the participation time, regarding to the database and after the end of the studies. Each patient gave a written consent prior to participation.

7.4.1 Potentially inappropriate medications with the STOPP criteria (Study I)

The data protection officer of Akershus University Hospital (No: SV12-086) reviewed and approved this study.

7.4.2 Medication regimen complexity and adherence (Study II) and Effectiveness of using the STOPP/START criteria (Study III)

The Regional Committee for Research Ethics (South/East, No: 2014/1255) and the Akershus University Hospital data protection officer (No: 15-068) reviewed and approved these studies. Furthermore, the study information was registered with www.clinicaltrials.gov (NCT 02424786).

8 RESULTS

“The young physician starts life with 20 drugs for each disease, and the old physician ends life with one drug for 20 diseases.”

Sir William Osler

8.1 Potentially inappropriate medications with the STOPP criteria (Study I)

In this cross-sectional study we included 50 patients age ≥ 65 years from a single HD centre. Hypertension, cardiovascular disease, and diabetes mellitus were the most common comorbidities in the patient population. A median of 13 medications per day (range 7–21) were used by all patients. A total of 652 medications were identified, and 32 patients (63%) used PIMs. In the medication lists of 21 patients (42%), one PIM was observed, and 11 patients (22%) had two PIMs. According to the STOPP criteria, the most prevalent PIMs were proton pump inhibitors with full therapeutic dosage use ($n=11$, 22%), calcium-channel blockers ($n = 7$, 14%) in patients with chronic constipation, and benzodiazepines ($n = 6$, 12%) in patients prone to falls. The number of PIMs did not differ significantly with age (≤ 74 years versus >74 years, $p = 0.39$), sex (male versus female, $p = 0.74$), comorbidity severity ($CCI \leq 5$ versus >5 , $p = 1.00$), time in haemodialysis (≤ 8 months versus >8 months, $p = 0.78$), or number of medications (≤ 13 versus >13 , $p = 0.56$). The Beers criteria identified PIMs in 22 patients (43%); the most common medication category was proton pump inhibitor. The following side effects were registered: 19 patients (37%) had falls during the previous three months, nine patients (18%) suffered chronic constipation before start of dialysis, and three patients (6%) had dizziness.

8.2 Medication regimen complexity and adherence (Study II)

In this prospective study, we included 180 patients of whom 157 patients aged 76 ± 7.2 years (mean $\pm$ SD) from all three centres could be included in the analysis. Participating were 64 predialytic patients, 19 PD patients, and 73 patients in HD without nocturnal HD. We used MRCI to assess the complexity of the medication regimen. The overall MRCI score was 22.8 ± 7.7 . Patients in HD reached the highest MRCI score, 24.7 ± 7.9 , followed by PD at 23.3 ± 6.4 and predialytic patients at 20.5 ± 7.3 . The univariate linear regression analysis showed a difference of the mean MRCI score between the dialysis and predialytic populations (coefficient=3.86, 95% CI 1.47–6.25, $p=0.002$). In multivariable linear regression analysis, being female (coefficient=2.44, 95% CI -0.07–4.81, $p=0.044$), having a Charlson Comorbidity Index of 4 or 5 (coefficient=2.56, 95% CI 0.27–4.85, $p=0.029$) and using several categories of phosphate binders ($p<0.001$ to 0.04) were associated with MRCI. The medication adherence measured with MMAS-8 with a skewed distribution had a median score of 8.0 (range 1.5–8.0). In total, 83% of the patients were adherent, defined as moderate/high adherence (MMAS-8 score ≥ 6). A more detailed division of the proportion of patients with moderate/high adherence showed 78%, 100% and 83% among the HD, PD and predialytic patients, respectively. The multivariable logistic regression analysis found no association of medication complexity, age, dialysis vintage, or other variables with medication adherence. However, there was a tendency towards being non-adherent in patients with higher educational levels and increased numbers of comorbidities.

8.3 Effectiveness of using the STOPP/START criteria (Study III)

We included and randomized 180 patients with advanced CKD in this single-blind, multicentre, parallel-group randomized, clinical trial. All patients were allocated either to an intervention or a control group. Among the 180 patients, 11% were lost due to death and 17% due to incomplete or lack of response to questionnaires. These “lost to follow-up” patients differed from the respondents in their baseline characteristics by exhibiting lower MMSE-NR scores and experiencing more hospitalisations during follow-up time. Otherwise, there was no difference in the baseline characteristics between allocation groups.

A total of 265 inappropriate medications were identified by the STOPP/START criteria at baseline. The medications most frequently prescribed inappropriately were proton pump inhibitors, benzodiazepine, and first generation antihistamines, whereas the most frequent omissions were angiotensin-converting enzyme inhibitors, statins and vitamin D. At baseline, the prevalence of PIPs was 54% in the intervention group and 55% in the control group. The PPO's prevalence at baseline was 50% in the intervention group and 56% in the control group.

For the primary outcomes, we found no statistically significant decrease in the number of patients with PIPs in the intervention group. Regarding the PPOs, the odds at follow-up were lower in the intervention group compared to the control group (OR 0.42, 95% CI 0.19–0.92, $p=0.032$). There was no difference between the groups regarding medication adherence. Moreover, for the secondary outcomes, there was no difference in average number of medications and HRQoL scores between the groups.

All sensitivity analyses verified no difference in PIPs, medication adherence, average number of medications or HRQoL scores, including the lower odds for omissions in the intervention group ($n=161$, OR 0.46, 95% CI 0.24–0.87, $p=0.017$).

9 DISCUSSION

“We shall not cease from exploration, and the end of all our exploring will be to arrive where we started and know the place for the first time”. T.S. Eliot

9.1 General findings

The central finding in the current papers was inappropriate prescribing as identified by the STOPP/START criteria, which were more sensitive than the Beers criteria in detecting inappropriate prescriptions. In addition, older patients with advanced CKD were characterized by polypharmacy and complex medication regimens, and in addition they had high medication adherence.

In extensive polypharmacy, it is not easy to assess underprescribing, while at the same time over- and misprescribing are occurring. By using a screening tool, this dilemma of recognition can be envisaged. Studies I and III showed that the medication lists of older patients with advanced CKD contained inappropriate medications. Several studies reported similar observations (125, 126). There are many reasons for missing clinically relevant medications; for instance, because of a lack of evidence-based knowledge due to older adults with comorbidities often being excluded in clinical trials, polypharmacy is mostly the rule, and age discrimination and concerns about ADEs may occur (127). In Study III, we showed that medication review can increase the awareness of inappropriateness. On the other hand, our intervention could not show an improvement in medication adherence, number of medications, or HRQoL.

In the following chapter, methodological issues and the main findings from all three studies will be discussed.

9.2 Discussion of the methodology

9.2.1 Study design

We chose a cross-sectional study design in two of the three papers to answer questions about prevalence and predictors for PIMs and to assess a possible association between medication complexity and adherence. The advantage of a cross-sectional study was that none would be lost to follow-up. However, it does not provide a “cause and effect” relationship. Cross-sectional studies offer an opportunity to establish links or associations. They are easy to conduct and provide data in a relatively short period of time.

For the RCT study to assess the impact of a medication intervention, we used a prospective, longitudinal study design. The advantages of a longitudinal study design include that all individuals in a cohort 1) can be observed for exposure or outcome, 2) can be followed over time with repeated measurements, 3) can possibly show cause and effect relationships, and 4) can exclude recall bias by collecting data prospectively. Some disadvantages of a longitudinal study design are the time factor, the effort needed to include enough data for robust statistical analysis and the higher chances of missing data through drop-outs (128).

We aimed to include a representative population. This was especially important if we were to transfer our findings to the real-world population. Trials often have restrictive inclusion and exclusion criteria, frequently excluding patients with high comorbidity burden, older age, or advanced CKD. This stringent policy leads to a homogenous group with good internal validity. When all representatives of a patient population are included, it is more likely that good external validity will be achieved. External validity relates to the generalization of the study, asking how likely it is that observed effects could occur outside the study population. Dekker and colleagues summarized it like this: *“From a clinician’s point of view the generalizability of study results*

is of paramount importance. According to the CONSORT statement external validity should be addressed in reporting RCT” (129). Potential factors influencing external validity are inclusion and exclusion criteria and any factors, e.g., concerns that may affect participation in a trial (130). The patient population of Studies II and III could count as a representative sample because there were few exclusion criteria. In fact, after the PRECIS-2 tool, this RCT could be understood as “pragmatic” (130). Still, when comparing the non-respondents with the respondents, the results revealed a frailer group with lower MMSE-NR scores and more hospitalisations. This underlines the continually present issue of other unknown factors affecting study participation. Therefore, caution is needed when generalizing results.

9.2.2 Reliability and validity

We understand reliability to be the overall consistency of a measurement, meaning that every time a questionnaire is used, it should deliver similar scores/answers. Reliability can be measured in different ways, e.g., inter-rater reliability (equivalence), test-retest methods (stability) and split halves reliability (homogeneity). Inter-rater reliability measures the agreement of independent raters. Jacob Cohen developed a statistical measurement of inter-rater reliability based on the theoretical assumption that there is some level of agreement between raters when the correct answer is unknown (131). The inter-rater reliability of Study I was moderate (kappa: 0.42 (0.24-0.56, 95% CI)) compared to previous studies (56, 102, 132), but this was in line with other findings (133). The moderate inter-rater reliability should be interpreted with caution because items from the STOPP criteria version 1 were not always clearly inferred by nephrologists, e.g., chronic constipation can be caused by multiple factors in a population with CKD.

Validity is the ability of an instrument to measure what it is intended to measure. The predictive validity of the Beers criteria and the STOPP criteria version 1 were tested in an older,

community-dwelling population (134). The conclusion was that there was low agreement and no significant difference between these criteria for outcome measurements such as ADEs, emergency department visits, and hospitalizations. Although the STOPP criteria slightly outperformed the Beers criteria regarding hospitalizations (134), the predictive validity has not been tested in patients with CKD. Furthermore, there are medications that might be more relevant for patients with CKD which are not covered in the STOPP/START criteria. Taking these points into consideration, there is a need to perform a pilot study to determine the validity for the CKD population, and by doing so perhaps more acceptance by nephrologists will be achieved.

9.2.3 Instruments

9.2.3.1 Eight-item Morisky Medication Adherence scale

Initially developed as a four-item questionnaire for patients with hypertension the Morisky Medication Adherence scale was updated to an eight-item questionnaire (110). This questionnaire has been used worldwide in a wide range of diseases, including CKD (135, 136). The advantages of using a self-reported questionnaire are that they are brief; they apply to many diseases; they are often inexpensive, and they provide immediate feedback. Any self-report questionnaire can result in incorrect estimates or response biases, but these questionnaires still can provide reasonably truthful adherence (78). The predictive and concurrent validity of this questionnaire was tested on a hypertensive population with good results (110). But validating any adherence questionnaire is difficult due to the lack of a “gold standard”. In a recent meta-analysis, the authors highlighted the acceptable internal consistency and reproducibility of MMAS-8 in a few diseases such as diabetes mellitus type 2. Furthermore, the authors concluded that a criterion validity with a cut-off score of 6 was not good enough to determinate medication adherence. This meta-analysis did not include any studies on patients with CKD. The authors

suggest a reassessment of the criterion validity for other scores and diseases (137). Even Morisky and colleagues expressed the need for further refinement of MMAS-8 with repeated demonstrations of validity and reliability in different populations (110). MMAS-8 may be relevant for dialysis patients and patients with CKD because it addresses a few behavioural issues that may be of importance to physicians. Another point supporting the use of self-reporting tools is to be found in the recommendations for clinical practice in NICE guidelines.

9.2.3.2 Mini Mental State Evaluation - Norwegian Revision

This test has been shown to be reliable in different disease populations. Its disadvantages are that it relies too much on memory, which may lead to a poor test-retest performance, and its failure to assess the executive functions, which are often altered in CKD patients (138, 139).

We included all patients with a MMSE-NR score of 23 or more to ensure a representative population. Regarding the cut-off score in this older population, there are some relevant considerations. There appears to be a connection between the various uremic toxins and pathogenesis of cognitive impairment, but the mechanisms are poorly understood (140-142). The timing of the testing is important because it seems to have an impact on performance. One study showed that performance varied over the course of a dialysis session by being worse during the session and best before or on the day after (141, 143). This observation could be attributed to the fluctuation of blood flow during a dialysis session. We performed the MMSE-NR test before the HD session, meaning that all patients had high levels of uremic toxins at testing time. Thus, the results could be compared to the group with conservative treatment and PD that may have equal uremic toxin levels. It is established that there is mild cognitive impairment in HD patients, which is in line with the reported average MMSE score of 23 in an older HD population (144, 145).

9.2.3.3 Medication Regimen Complexity Index

The MRCI describes any medication regimen in terms of complexity independent of disease or age, taking into account not only pill count, but also administration instructions and dosage timing, and this might represent a more realistic picture faced by patients with polypharmacy. Unfortunately, there is no defined cut-off score for “high” and “low” complexity, which makes inter-study comparisons difficult and its transfer to clinical practice unrealistic due to interpretation issues. Only three studies reported a different cut-off score of MRCI for predicting unplanned hospitalizations (67). Thus, Wimmer and colleagues suggested that the cut-offs are specific to the populations in which MRCI is applied (67). In addition, a recent literature review found that MRCI was not a better predictor of non-adherence and unplanned hospitalizations (146).

9.2.4 Confounding and bias

The common definition of “confounder” is considered to be a variable that was associated with exposure and outcome, also possibly conditional to other covariates (147). As confounding is linked to both dependent and independent variables, the outcome can be affected. In dealing with confounding, the epidemiologist can respond in two ways: 1) through study design and data collection, and 2) through statistical analysis. The most common ways to adjust this in study designs are randomization, restriction, and matching. Stratification and multivariate methods are commonly used after collection for confounding adjustments by statistical analysis (148). In modern epidemiology, the use of directed acyclic graphics to create visual and causal diagrams to identify confounders from observational data is increasing (149).

The definition of bias given in the Merriam Webster dictionary is: “*systematic error introduced into sampling or testing by selecting or encouraging one outcome or answer over others*” (150).

Bias occurs all the time, even in well-planned and executed studies. Statistical analysis cannot adjust for bias, and in case of bias all results should be interpreted with caution.

In all three studies, we recognized one or more biases such as selection bias, transfer bias, attrition bias, recall bias, interview bias, and many more. All studies might have selection bias. We included only adults ≥ 65 years of age. In addition, a preselection of patients at inclusion time was done by doctors, nurses and the investigators. This preselection excluded all older adults with impaired cognitive function and all non-Norwegian speaking patients. By using a cut-off in the evaluation of cognitive function, we ignored patients who already had more daily challenges and were more vulnerable. Older adults with unsatisfactory Norwegian language skills were not included due to the lack of validated questionnaires in other languages, which was unavoidable. Yet, one of many questions remains: Would adults of 55 years of age with advanced CKD differ compared to the selected population? Many research questions asked in the thesis are of importance for any person of any age with advanced CKD.

Transfer bias occurs in clinical trials when subjects are lost unequally to follow-up. We expected this bias in the population of Studies II and III. Thus, as a preventive step to reduce this bias, we arranged convenient appointments in the outpatient clinic or dialysis centre, both at baseline and follow up. Furthermore, payment for visits was waived; necessary transportation was organized and facilitated, and telephone calls were made to remind non-respondents (151).

When participants leave an ongoing study, it is called “attrition”. Attrition bias occurs when there is a systemic difference between the participants who stayed and those who left the study. Awareness of this bias needs to be taken into account when interpreting results; some would call it non-response bias. We realized that this type of bias appeared during our study when analysing the data for Study III. We had a high death rate of 11%, and 17% of questionnaires

were not returned or completed. The analysis between the respondents versus non-respondents highlighted that the non-responder group was more vulnerable based on lower MMSE-NR scores and increased hospitalizations.

9.2.5 Missing data

Missing data are defined as values that are not available and that could be meaningful for the analysis if they were observed (152).

In Study III, there was missing data from the included patients, e.g., resulting from them not responding to questionnaires. In the literature, there is no consensus about what proportion of missing data is acceptable without compromising the validity of a trial. The validity of a trial can be jeopardized if missing data reaches 20% or more, but less than 5% is negligible. In general, missing data in the 5% to 20% range can impact the results and raise statistical concerns (153). Regarding prevention and treatment of missing data in clinical trials, the National Research Council (NRC) published a report addressing the ability to draw correct conclusions (152).

In Study III, we had about 11% missing data due to death of patients, which would be expected in this population as the mortality hazard ratio was 3.6 for CKD (95% CI, 3.3 to 4.0), 9.2 for PD (95% CI, 6.6 to 12.7) and 12.6 for HD (95% CI, 10.8 to 14.6) (154). About an additional 17% of data were missing due to questionnaires not being returned even after a reminder. Multiple reasons for this included worsening of the general health state, losing the questionnaires, changing life condition by moving to a long-term facility, and loss of interest in participating in this study due to increasing frailty.

9.2.6 Statistical considerations

The amount of missing data in our longitudinal follow-up study (Study III) represented a challenge for statistical analysis. In longitudinal studies, we are repeatedly measuring a single outcome for each individual. For the continuous variables, several statistical models are available to address this, but for categorical or binary variables, little guidance for longitudinal data analysis is offered in the literature (155). A high proportion of missing data may influence the results and the choice of statistical methods (156). Longitudinal RCTs usually have less than 25% missing data (156), which is less than in the present Study III. The NRC report is helpful regarding these difficulties in handling missing data and choosing the appropriate statistical methods (152). The panel preferred estimating-equation methods and methods that are based on statistical models. Furthermore, they recommended sensitivity analysis to assess robustness (152). A generalized linear mixed model is one of many statistical models being used increasingly in medical literature (155). It takes into account all data at baseline, both continuous and binary variables that involve random effects.

9.3 Discussion of the main findings

9.3.1 Potentially inappropriate medications with the STOPP criteria (Study I)

The study found a high prevalence of PIPs according to the STOPP criteria in this older population with HD. This is in contrast with another older HD population showing a lower prevalence, which might be explained by a lower average number of medications and CCI scores (157). Yet, the prevalence of PIPs in this present study is in agreement with two studies on patients with CKD using the Beers criteria (63, 158). Across all types of populations, the prevalence of inappropriate medications ranges from 23% to 79% (125). This is understandable due to differences in study populations, study designs, number of medications and comorbidities.

We did not find an association between the number of PIPs and age, sex, comorbidity, time in HD or number of medications. This could be explained by the small sample size, heterogeneity of the study population, and the chosen cut-off points for each predictor that are consistent with another study on older adults with metabolic disease (159).

Several studies compared explicit criteria such as the STOPP and the Beers criteria. The authors of two reviews concluded that the STOPP criteria were more sensitive than the Beers criteria (125, 160), which is in line with our conclusion. However, a European study highlighted that half the medications registered in the Beers criteria were not approved in most European countries (161). The percentage of approved medications on the Beers criteria list in Norway was 32%, in the Netherlands 48%, in Iceland 51%, in Denmark 52%, in Finland and the UK 56%, and in Italy 71%. Conversely, medications approved for the European market were not listed in the Beers criteria. Thus, it is problematic to see the suitability of comparing the Beers criteria with the STOPP/START criteria as it “may be a case of comparing apples with

oranges”. Logically, it would be more appropriate to use a physician-pharmacist team or other European explicit criteria such as the Norwegian General Practice Criteria for comparison (162, 163).

9.3.2 Medication regimen complexity and adherence (Study II)

Many studies and reviews reported an association between the complexity of a regimen and negative health outcomes such as declining medication adherence (146). A previous review suggested that simplifying a medication regimen may improve adherence (79). A standardized agreement on how to assess a medication regimen's complexity is still missing. So far, studies have shown an inverse association between number of pills, mainly phosphate binders, and adherence (49, 164). Consistent with our results, a lack of association between medication complexity as measured by MRCI and medication adherence has been observed in a HD population (105). In this population, less than half the patients were defined as non-adherent, which contrasts with to our high medication adherence. The reason for the lack of association in both studies remains unclear, and is in contrast to another study on HD patients (164). An explanation for this difference might lie in the definition and assessment of medication complexity. Furthermore, there has been reported a high variability regarding the association between medication complexity and adherence (165).

Our findings regarding the MRCI score line up with several other studies in the geriatric population and other chronic diseases such as diabetes mellitus and heart failure (68, 166, 167). There was uncertainty as to how to interpret the item "dialysate" in Section A of the MRCI. As a consequence our total MRCI score was lower when compared to previous studies on HD populations (105, 106). Another explanation for a lower score might be the inter-centre prescription traditions and availability of medications. However, it is challenging to define any MRCI score as high or low. This definition is missing, and it might be that a cut-off score needs to be defined for each chronic disease (67).

We preselected predictors that might influence the medication regimens' complexity. The predictor *age* does not seemed to have an influence (168), but the results of previous studies

diverge regarding age as a predictor (169). Female sex seemed to have an impact, not only in this study, but also in other reports regarding different diseases (170, 171). Women might present symptoms and concerns about their own health differently than men, resulting in a different prescribing attitude and management by physicians (172).

It is known that the pill burden in patients with advanced CKD is high, and phosphate binders contribute significantly to this. These binders prescribed to most patients with CKD, are essential in the management of hyperphosphatemia, which may play a role in cardiovascular calcification and cardiovascular mortality (173, 174). Moreover, we found a strong correlation between pill burden and phosphate binders: so, we decided to use phosphate binders because of their major role in treatment of CKD. Administering phosphate binders involves additional instructions and individual administration requirements that are more accurately captured by MRCI than by pill burden.

In other studies of older patients with CKD, the medication adherence rate was high, agreeing with our findings (175). Phosphate binders are negatively associated with medication adherence (49, 105). However, we could not confirm such an association. The probability of having a high adherence seemed to decline with an increasing amount of phosphate binders, expressed through the changing gradient in the odds ratios. This phenomenon could indicate a bidirectional relationship between prescriber and patients. If non-adherence towards a medical treatment is present, the expected effect will not occur, leading the physician to a false assumption of a failed treatment. This will trigger a maximisation of the established treatment. However, an increased number of pills can cause non-adherence because of the medication load. For the other predictors such as age, comorbidity, and educational level, we did not report an association with medication adherence.

In general, age has been the variable most frequently associated with adherence, and younger people have been associated with non-adherence independently of the assessment tool (89). Older people have a different understanding of the statement “*I need to take medicine*“. Their understanding recognises distinctive levels of importance. Important medicines are defined by complex inter-related factors: drug-related, patient-related, and external (176, 177). Older people gain knowledge about dealing with medications over the years. They believe in the importance of their disease management and may have more support related to their medication regimens (178, 179). Still, there are concerns about the continuation of a medication when test results are normal. Improving patient-physician relationships may increase medication adherence and reduce medication regimens’ complexity by altering patients’ perception of medications, enhancing knowledge, and supporting a shared decision-making process.

9.3.3 Effectiveness of using the STOPP/START criteria (Study III)

In this study, we performed a medication review using the STOPP/START criteria to identify over- and underprescribing in an older population with advanced CKD. This medication review reduced the number of omissions, but it had no effect on PIPs, medication adherence, or HRQoL.

Applying the STOPP/START criteria to medication lists of people with advanced CKD highlighted possible PIMs, and this is consistent with findings reported in other studies with older adults in HD (61, 180). The prevalence of PIMs in this study group was high and in accordance with other studies (125), but was lower when compared only to one study on dialysis patients using the same version of the STOPP/START criteria (180). The prevalence of PIMs differs between studies due to differences in included study populations, medications, prescribing practices or traditions, and the use of different versions of the STOPP/START criteria.

Medication groups such as benzodiazepines and proton pump inhibitors were identified frequently as PIPs in several other studies (61, 125, 181). Furthermore, the present findings support previous reports showing that underprescribing is common in older adults and people with CKD. About 50% of all participants lacked recommended preventive or evidence-based medications. This is particularly important for the medication group treating cardiovascular diseases and complications, which occur frequently in this population. Especially, the underprescribing of angiotensin-converting-enzyme inhibitors has been raised in previous studies (182-184). This medication group was one of the most commonly detected omissions in the present study. Moreover, evidence showed an association between underprescribing and mortality and hospitalization rates in a cohort of older adults (185), which makes PPOs just as important as PIPs. There are reasons for underprescribing in older adults, such as trying to avoid

adding medications to polypharmacy lists, uncertainty or ignorance regarding evidence-based medications, and the aversion of older adults towards new treatment suggestions (185).

The intervention was associated with a reduction in numbers of PPOs, but not PIPs, which contrasts with a recent review that reported improvements for both PIPs and PPOs. Some possible causes could be raised. First, the implementation of recommended medication changes was left completely to the judgement of the attending physician. Each physician has barriers and enablers regarding prescribing, especially when it comes to stopping or adapting prescriptions. Physicians' behaviour and perspectives need to be overcome to minimize inappropriate medications (186, 187). Furthermore, it is unclear to what degree single blinding has diminished the effect of the difference between allocation groups. Physicians tend to be biased towards the intervention group. Steps taken to reduce this bias were using the objective questionnaires as outcome variables, and applying standardizing follow-up care for all participants (188). Second, there may be a lack of acceptance of these criteria due to the lack of validity in a nephrology population. After all, of all medications taken by patients with CKD, about 26% are specific to renal disease (189). Third, despite a possible initial acceptance of medication recommendations, there may be a gradual increase of PIPs over the follow-up time of six months as observed by Gallagher and colleagues (190). These authors suggested a medication review every six months with a screening tool, which is in line with the recommendations of the National Service Framework for Older People for those taking four or more medications (191). Another point is the length of follow-up time being six months. The length of follow-up was based on previously published RCTs (126) on the high mortality risk in people with ESRD (154) and as an attempt to decrease missing data for statistical analysis (152).

The medication review with the STOPP/START criteria in the present study had no effect on medication adherence or HRQoL. Medication adherence was high at baseline and at six months follow-up in both groups, supporting previous reports. The lack of improvement in adherence could not be detected based on the ceiling effect of the questionnaire or the chosen definition of cut-off. Pharmacist-led intervention in ambulatory and community care populations have also failed to demonstrate an improvement in medication adherence, but comparison between studies is difficult due to the high diversity of the study population, study design, and chosen interventions (192). One RCT study of community-dwelling older people, including a pharmacist-led review of medications, comes to the same conclusion of no improvement of medication adherence when measured by the four-item MMAS (193). Similar inconsistent findings could be reported about HRQoL after intervention by a pharmacist (192).

The present study found no change in the number of medications, which is in line with some previous reports (194, 195). This observation highlights that an intervention with the STOPP/START criteria is not necessarily associated with an increase in number of prescriptions, but could even lead to a reduction of polypharmacy (196).

10 CONCLUSIONS

“The desire to take medicine is perhaps the greatest feature which distinguishes man from animals”. Sir William Osler

The major conclusions based on the findings of all three studies of older adults with advanced CKD, included in this thesis, are:

- Inappropriate medications in medication lists of older adults in HD could be identified using the STOPP criteria or the Beers criteria.
- Risk factors such as age, sex, number of medications and comorbidity were not associated with inappropriate medications in older adults in HD.
- Being female, multimorbid or use of phosphate binders were all factors associated with an increasing medication regimen complexity as measured by MRCL.
- Older adults with advanced CKD showed a high level of medication adherence, but there was no association between the medication regimen complexity and medication adherence.
- Explicit criteria such as the STOPP/START criteria were able to identify inappropriate medications, both over- and underprescribing, in older adults with advanced CKD.
- After applying the STOPP/START criteria in the intervention group, this group had a lower number of PPOs, but not in the number of PIPs, medication adherence, number of medications and in the HRQoL-scores than the control group.

11 CLINICAL IMPLICATIONS

“It is much more important to know what sort of a patient has a disease than what sort of a disease a patient has.” Sir William Osler

Epidemiologic data point to an increase in the aging population over the coming years. This is in line with an increasing number of people with advanced CKD. Pharmacological therapy is the basic treatment for impaired renal function independent of the stage of CKD, treatment modality, and complications. The challenge is that older adults with or without CKD are often excluded from studies, leaving a knowledge gap on how strictly a physician should follow these guidelines. Moreover, independent of how many medications a person takes, every medication prescribed should be necessary and appropriate, and the administration should be as simple as possible.

Our studies added new knowledge regarding this older population on how to approach a complex polypharmacy list. In clinical practice, few HD centres have a pharmacist available to support the attending physicians with medication reviews. Nephrologists have a challenging task in dealing with the extensive use of medications in older adults with a wide range of comorbidities. In addition, they must understand the interactions between different drugs as well as drug-disease interactions with simultaneously impaired kidney function. Through medication reviews, physicians can engage in important and necessary conversations to understand their patients’ needs, and to keep the medication lists updated while continuously adjusting to physical changes or declining renal function, and this might prevent unplanned hospitalisations. Therefore, the implementation of screening tools, i.e., explicit criteria, may be useful in clinical practice. Future research is needed to develop explicit criteria designed for this vulnerable population or to make the case for establishing a multidisciplinary team that includes a pharmacist.

Older adults with advanced CKD adhere to their treatment mainly because they understand the necessity to keep their health status stable. The complexity of medication regimens did not affect patients' adherence, which may point to patients' having the higher purpose of achieving the best possible health state. This knowledge should be used as a guide when objective measurements or results of a treatment point in opposite directions. Assessing medication adherence and complexity could raise awareness to changes in patients' lives, which might in turn lead to re-adjustment in medications and treatment goals. This awareness is towards personalized care and medication treatment, and it might lead to a better HRQoL.

12 SUGGESTIONS FOR FUTURE RESEARCH

“In my view, the lost art of listening and ignoring the patient as a human being is a quintessential failure of our health care.”

Bernard Lown

In this thesis, we tried to understand appropriateness, complexity and adherence in pharmaceutical therapy in older adults, but actually, it turned out that we highlighted more questions and missing knowledge than expected. All knowledge acquired up to this point can be seen as a starting point for more thorough further research.

Our findings showed a high prevalence of inappropriate prescriptions in all populations with advanced CKD. Currently available screening tools do not cover renal-specific medications and their side effects. Thus, the next step should be the development of an appropriate screening tool for this vulnerable patient group and to test it for effectiveness in longitudinal, randomized, clinical trials. Maybe through acknowledging and emphasising inappropriate medications, we can embrace the challenge of de-prescribing and shared-decision making.

Another study, maybe as a RCT, could compare different allocation groups (i.e., criteria, pharmacist and control) regarding medication complexity and adherence with follow-up times of 3, 6, 12 or 18 months.

The medication adherence of this older group was not assessed based on other important elements that influence adherence such as psychological factors. A qualitative study addressing important factors affecting medication adherence and exploring the differences between groups of young-old and old-old adults would contribute more knowledge. Especially, with older adults 80 years and older, there is a need to understand their goals in life, their understanding of their health condition, and their fears while suffering advanced CKD.

Another interesting research area would be to learn whether interventions from a pharmacotherapeutic viewpoint would have clinical relevance and a real impact on the lives of older adults. How can we sustain and improve quality of life? How can we acknowledge and support a shortening lifespan using medications without creating an endless list?

13 REFERENCES

1. World Health Organization. Proposed working definition of an older person in Africa for the MDS Project: World Health Organization; 2019 [Available from: <https://www.who.int/healthinfo/survey/ageingdefnolder/en/>].
2. Encyclopaedia Britannica. Old age 2019 [Available from: <https://www.britannica.com/science/old-age>].
3. Little W. Introduction to Sociology - 1st Canadian Edition. 2014. Available from: <https://opentextbc.ca/intr>.
4. Giannakouris K. Regional population projections EUROPOP2008: Most EU regions face older population profile i 2030. Eurostat: Statistics in focus. 2010;1/2010.
5. Spinewine A, Schmader KE, Barber N, Hughes C, Lapane KL, Swine C, et al. Appropriate prescribing in elderly people: how well can it be measured and optimised? *Lancet*. 2007;370(9582):173-84.
6. American Geriatrics Society 2015 Updated Beers Criteria for Potentially Inappropriate Medication Use in Older Adults. *J Am Geriatr Soc*. 2015;63(11):2227-46.
7. Ruggiero C, Dell'Aquila G, Gasperini B, Onder G, Lattanzio F, Volpato S, et al. Potentially inappropriate drug prescriptions and risk of hospitalization among older, Italian, nursing home residents: the ULISSE project. *Drugs Aging*. 2010;27(9):747-58.
8. O'Mahony D, Gallagher PF. Inappropriate prescribing in the older population: need for new criteria. *Age Ageing*. 2008;37(2):138-41.
9. Gallagher P, Barry P, O'Mahony D. Inappropriate prescribing in the elderly. *J Clin Pharm Ther*. 2007;32(2):113-21.
10. Bernell S, Howard SW. Use Your Words Carefully: What Is a Chronic Disease? *Front Public Health*. 2016;4(159).
11. Merriam Webster Dictionary. Chronic 2019 [Available from: <https://www.merriam-webster.com/dictionary/chronic>].
12. He Z, Bian J, Carretta HJ, Lee J, Hogan WR, Shenkman E, et al. Prevalence of Multiple Chronic Conditions Among Older Adults in Florida and the United States: Comparative Analysis of the OneFlorida Data Trust and National Inpatient Sample. *J Med Internet Res*. 2018;20(4):e137-e.
13. Wolff JL, Starfield B, Anderson G. Prevalence, expenditures, and complications of multiple chronic conditions in the elderly. *Arch Intern Med*. 2002;162(20):2269-76.
14. Coin A, Sergi G, Inelmen EM, Enzi G. Pathophysiology of body composition changes in elderly people. In: Mantovani G, Anker SD, Inui A, Morley JE, Fanelli FR, editors. *Cachexia and Wasting: A Modern Approach*. Milano: Springer; 2006.
15. Muravchick S. Anesthesia for the Elderly. In: Miller RD, editor. *Anesthesia*. 5th ed. Philadelphia: Churchill Livingstone; 2000. p. 2140-56.
16. Collins Dictionary. Volume of distribution 2019 [Available from: <https://www.collinsdictionary.com/dictionary/english/volume-of-distribution>].
17. Mangoni AA, Jackson SHD. Age-related changes in pharmacokinetics and pharmacodynamics: basic principles and practical applications. *Br J Clin Pharmacol*. 2004;57(1):6-14.
18. Bowie MW, Slattum PW. Pharmacodynamics in Older Adults: A Review. *Am J Geriatr Pharmacother*. 2007;5(3):263-303.
19. National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis*. 2002;39(2 Suppl 1):S1-266.

20. Agarwal R. Defining end-stage renal disease in clinical trials: a framework for adjudication. *Nephrol Dial Transplant*. 2016;31(6):864-7.
21. Sakula A. Pierre Adolphe Piorry (1794-1879): pioneer of percussion and pleximetry. *Thorax*. 1979;34(5):575-81.
22. National Institute for Health and Care Excellence (NICE). Chronic kidney disease in adults: assessment and management 2015 [Available from: <https://www.nice.org.uk/guidance/cg182/chapter/1-Recommendations#referral-criteria>].
23. Nolin TD, Unruh ML. Clinical relevance of impaired nonrenal drug clearance in ESRD. *Semin Dial*. 2010;23(5):482-5.
24. Coresh J, Selvin E, Stevens LA, Manzi J, Kusek JW, Eggers P, et al. Prevalence of chronic kidney disease in the United States. *JAMA*. 2007;298(17):2038-47.
25. Gansevoort RT, Correa-Rotter R, Hemmelgarn BR, Jafar TH, Heerspink HJL, Mann JF, et al. Chronic kidney disease and cardiovascular risk: epidemiology, mechanisms, and prevention. *Lancet*. 2013;382(9889):339-52.
26. Aitken GR, Roderick PJ, Fraser S, Mindell JS, Donoghue D, Day J, et al. Change in prevalence of chronic kidney disease in England over time: comparison of nationally representative cross-sectional surveys from 2003 to 2010. *BMJ Open*. 2014;4(9).
27. Hallan SI, Øvrehus MA, Romundstad S, Rifkin D, Langhammer A, Stevens PE, et al. Long-term trends in the prevalence of chronic kidney disease and the influence of cardiovascular risk factors in Norway. *Kidney Int*. 2016;90(3):665-73.
28. Hoerger TJ, Simpson SA, Yarnoff BO, Pavkov ME, Ríos Burrows N, Saydah SH, et al. The Future Burden of CKD in the United States: A Simulation Model for the CDC CKD Initiative. *Am J Kidney Dis*. 2015;65(3):403-11.
29. Kane PM, Vinen K, Murtagh FE. Palliative care for advanced renal disease: A summary of the evidence and future direction. *Palliat Med*. 2013;27(9):817-21.
30. Wolfe RA, Ashby VB, Milford EL, Ojo AO, Ettenger RE, Agodoa LY, et al. Comparison of mortality in all patients on dialysis, patients on dialysis awaiting transplantation, and recipients of a first cadaveric transplant. *N Engl J Med*. 1999;341(23):1725-30.
31. Russell JD, Beecroft ML, Ludwin D, Churchill DN. The quality of life in renal transplantation--a prospective study. *Transplantation*. 1992;54(4):656-60.
32. Dobbels F, De Bleser L, De Geest S, Fine RN. Quality of Life After Kidney Transplantation: The Bright Side of Life? *Adv Chronic Kidney Dis*. 2007;14(4):370-8.
33. Heldal K, Leivestad T, Hartmann A, Svendsen MV, Lien BH, Midtvedt K. Kidney transplantation in the elderly—the Norwegian experience. *Nephrol Dial Transplant*. 2007;23(3):1026-31.
34. World Health Organization. Traditional medicine: World Health Organization; 2019 [Available from: <https://www.who.int/traditional-complementary-integrative-medicine/about/en/>].
35. Qato DM, Wilder J, Schumm L, Gillet V, Alexander G. Changes in prescription and over-the-counter medication and dietary supplement use among older adults in the united states, 2005 vs 2011. *JAMA Intern Med*. 2016;176(4):473-82.
36. Berg C. Reseptregisteret 2013-2017 (The Norwegian Prescription Database 2013-2017). Oslo, Norge: Folkehelseinstituttet 2018.
37. Laroche M, Charmes J, Bouthier F, Merle L. Inappropriate Medications in the Elderly. *Clin Pharmacol Ther*. 2009;85(1):94-7.
38. Gorard DA. Escalating polypharmacy. *QJM*. 2006;99(11):797-800.

39. Haider SI, Johnell K, Thorslund M, Fastbom J. Trends in polypharmacy and potential drug-drug interactions across educational groups in elderly patients in Sweden for the period 1992 - 2002. *Int J Clin Pharmacol Ther*. 2007;45(12):643-53.
40. Barat I, Andreasen F, Damsgaard EM. Drug therapy in the elderly: what doctors believe and patients actually do. *Br J Clin Pharmacol*. 2001;51(6):615-22.
41. Viktil KK, Blix HS, Moger TA, Reikvam A. Polypharmacy as commonly defined is an indicator of limited value in the assessment of drug-related problems. *Br J Clin Pharmacol*. 2007;63(2):187-95.
42. Merriam Webster Dictionary. Polypharmacy 2019 [Available from: <https://www.merriam-webster.com/dictionary/polypharmacy>].
43. Masnoon N, Shakib S, Kalisch-Ellett L, Caughey GE. What is polypharmacy? A systematic review of definitions. *BMC Geriatr*. 2017;17(1):230.
44. Milton JC, Hill-Smith I, Jackson SHD. Prescribing for older people. *BMJ*. 2008;336(7644):606-9.
45. Steinman MA, Landefeld CS, Rosenthal GE, Berthenthal D, Sen S, Kaboli PJ. Polypharmacy and prescribing quality in older people. *J Am Geriatr Soc*. 2006;54(10):1516-23.
46. Murray MD, Kroenke K. Polypharmacy and medication adherence: small steps on a long road. *J Gen Intern Med*. 2001;16(2):137-9.
47. Kroenke K. Polypharmacy. Causes, consequences, and cure. *Am J Med*. 1985;79(2):149-52.
48. St Peter WL. Improving medication safety in chronic kidney disease patients on dialysis through medication reconciliation. *Adv Chronic Kidney Dis*. 2010;17(5):413-9.
49. Chiu YW, Teitelbaum I, Misra M, de Leon EM, Adzize T, Mehrotra R. Pill burden, adherence, hyperphosphatemia, and quality of life in maintenance dialysis patients. *Clin J Am Soc Nephrol*. 2009;4(6):1089-96.
50. Beers MH. Explicit criteria for determining potentially inappropriate medication use by the elderly. An update. *Arch Intern Med*. 1997;157(14):1531-6.
51. Beers MH, Ouslander JG, Rollingher I, Reuben DB, Brooks J, Beck JC. Explicit criteria for determining inappropriate medication use in nursing home residents. UCLA Division of Geriatric Medicine. *Arch Intern Med*. 1991;151(9):1825-32.
52. Hanlon JT, Schmader KE, Samsa GP, Weinberger M, Uttech KM, Lewis IK, et al. A method for assessing drug therapy appropriateness. *J Clin Epidemiol*. 1992;45(10):1045-51.
53. Hyttinen V, Jyrkka J, Valtonen H. A Systematic Review of the Impact of Potentially Inappropriate Medication on Health Care Utilization and Costs Among Older Adults. *Med Care*. 2016;54(10):950-64.
54. Wang KN, Bell JS, Chen EYH, Gilmartin-Thomas JFM, Ilomäki J. Medications and Prescribing Patterns as Factors Associated with Hospitalizations from Long-Term Care Facilities: A Systematic Review. *Drugs Aging*. 2018;35(5):423-57.
55. Hedna K, Hakkarainen KM, Gyllenstein H, Jönsson AK, Petzold M, Hägg S. Potentially inappropriate prescribing and adverse drug reactions in the elderly: a population-based study. *Eur J Clin Pharmacol*. 2015;71(12):1525-33.
56. Gallagher P, Ryan C, Byrne S, Kennedy J, O'Mahony D. STOPP (Screening Tool of Older Person's Prescriptions) and START (Screening Tool to Alert doctors to Right Treatment). Consensus validation. *Int J Clin Pharmacol Ther*. 2008;46(2):72-83.
57. Laroche ML, Charmes JP, Merle L. Potentially inappropriate medications in the elderly: a French consensus panel list. *Eur J Clin Pharmacol*. 2007;63(8):725-31.

58. McLeod PJ, Huang AR, Tamblyn RM, Gayton DC. Defining inappropriate practices in prescribing for elderly people: a national consensus panel. *CMAJ*. 1997;156(3):385-91.
59. Winit-Watjana W, Sakulrat P, Kespichayawattana J. Criteria for high-risk medication use in Thai older patients. *Arch Gerontol Geriatr*. 2008;47(1):35-51.
60. Rognstad S, Brekke M, Fetveit A, Spigset O, Wyller TB, Straand J. The Norwegian General Practice (NORGE) criteria for assessing potentially inappropriate prescriptions to elderly patients. A modified Delphi study. *Scand J Prim Health Care*. 2009;27(3):153-9.
61. Parker K, Aasebø W, Stavem K. Potentially inappropriate medications in elderly haemodialysis patients using the STOPP criteria. *Drugs - Real World Outcomes*. 2016;3(3):359-63.
62. Tesfaye WH, Castelino RL, Wimmer BC, Zaidi STR. Inappropriate prescribing in chronic kidney disease: A systematic review of prevalence, associated clinical outcomes and impact of interventions. *Int J Clin Pract*. 2017;71(7).
63. Kondo N, Nakamura F, Yamazaki S, Yamamoto Y, Akizawa T, Akiba T, et al. Prescription of potentially inappropriate medications to elderly hemodialysis patients: prevalence and predictors. *Nephrol Dial Transplant*. 2015;30(3):498-505.
64. Dörks M, Allers K, Schmiemann G, Herget-Rosenthal S, Hoffmann F. Inappropriate Medication in Non-Hospitalized Patients With Renal Insufficiency: A Systematic Review. *J Am Geriatr Soc*. 2017;65(4):853-62.
65. Kelley SO. Measurement of the Complexity of Medication Regimens of the Elderly [thesis]. Columbia, Mo: University of Missouri; 1988.
66. Wimmer BC, Bell JS, Fastbom J, Wiese MD, Johnell K. Medication Regimen Complexity and Polypharmacy as Factors Associated With All-Cause Mortality in Older People: A Population-Based Cohort Study. *Ann Pharmacother*. 2016;50(2):89-95.
67. Wimmer BC, Bell JS, Fastbom J, Wiese MD, Johnell K. Medication Regimen Complexity and Number of Medications as Factors Associated With Unplanned Hospitalizations in Older People: A Population-based Cohort Study. *J Gerontol A Biol Sci Med Sci*. 2016;71(6):831-7.
68. Colavecchia AC, Putney DR, Johnson ML, Aparasu RR. Discharge medication complexity and 30-day heart failure readmissions. *Res Social Adm Pharm*. 2017;13(4):857-63.
69. Schoonover H, Corbett CF, Weeks DL, Willson MN, Setter SM. Predicting potential postdischarge adverse drug events and 30-day unplanned hospital readmissions from medication regimen complexity. *J Patient Saf*. 2014;10(4):186-91.
70. World Health Organization Library. Adherence to long-term therapies:evidence for action. In: Library WHO, editor.: World Health Organization; 2003.
71. Haynes RB, Ackloo E, Sahota N, McDonald HP, Yao X. Interventions for enhancing medication adherence. *Cochrane Database Syst Rev*. 2008(2):Cd000011.
72. Morris LS, Schulz RM. Medication compliance: the patient's perspective. *Clin Ther*. 1993;15(3):593-606.
73. Royal Pharmaceutical Society of Great Britain. From compliance to concordance: achieving shared goals in medicine taking: Royal Pharmaceutical Society of Great Britain and Merck Sharp & Dohme; 1997.
74. Sørensen K, Van den Broucke S, Fullam J, Doyle G, Pelikan J, Slonska Z, et al. Health literacy and public health: A systematic review and integration of definitions and models. *J BMC Public Health*. 2012;12(1):80.

75. Taylor DM, Fraser SDS, Bradley JA, Bradley C, Draper H, Metcalfe W, et al. A Systematic Review of the Prevalence and Associations of Limited Health Literacy in CKD. *Clin J Am Soc Nephrol*. 2017;12(7):1070-84.
76. Alison Phillips L, Leventhal H, Leventhal EA. Assessing theoretical predictors of long-term medication adherence: Patients' treatment-related beliefs, experiential feedback and habit development. *Psychol Health*. 2013;28(10):1135-51.
77. Horne R, Weinman J. Patients' beliefs about prescribed medicines and their role in adherence to treatment in chronic physical illness. *J Psychosom Res*. 1999;47(6):555-67.
78. Blaschke TF, Osterberg L, Vrijens B, Urquhart J. Adherence to medications: insights arising from studies on the unreliable link between prescribed and actual drug dosing histories. *Annu Rev Pharmacol Toxicol*. 2012;52:275-301.
79. Osterberg L, Blaschke T. Adherence to medication. *N Engl J Med*. 2005;353(5):487-97.
80. Feinstein AR. On white-coat effects and the electronic monitoring of compliance. *Arch Intern Med*. 1990;150(7):1377-8.
81. Wroe AL. Intentional and Unintentional Nonadherence: A Study of Decision Making. *J Behav Med*. 2002;25(4):355-72.
82. Grymonpre RE, Didur CD, Montgomery PR, Sitar DS. Pill count, self-report, and pharmacy claims data to measure medication adherence in the elderly. *Ann Pharmacother*. 1998;32(7-8):749-54.
83. Marcum ZA, Gellad WF. Medication Adherence to Multi-Drug Regimens. *Clin Geriatr Med*. 2012;28(2):287-300.
84. Hughes CM. Medication non-adherence in the elderly: how big is the problem? *Drugs Aging*. 2004;21(12):793-811.
85. Vitolins MZ, Rand CS, Rapp SR, Ribisl PM, Sevick MA. Measuring Adherence to Behavioral and Medical Interventions. *Control Clin Trials*. 2000;21(5, Supplement 1):S188-S94.
86. Sturgess IK, Hughes CM, McElnay JC, King M. Identifying non-compliance of elderly patients: a comparison of three methods. *British Pharmaceutical Conference London: Pharmaceutical Journal*; 1998. p. 348-9.
87. Balkrishnan R. Predictors of medication adherence in the elderly. *Clin Ther*. 1998;20(4):764-71.
88. Weintraub M. Intelligent noncompliance with special emphasis on the elderly. *Contemp Pharm Pract*. 1981;4(1):8-11.
89. Ghimire S, Castelino RL, Lioufas NM, Peterson GM, Zaidi ST. Nonadherence to medication therapy in haemodialysis patients: a systematic review. *PLoS One*. 2015;10(12):e0144119.
90. Griva K, Lai AY, Lim HA, Yu Z, Foo MW, Newman SP. Non-adherence in patients on peritoneal dialysis: a systematic review. *PLoS One*. 2014;9(2):e89001.
91. Magacho EJC, Ribeiro LC, Chaoubah A, Bastos MG. Adherence to drug therapy in kidney disease. *Braz J Med Biol Res*. 2011;44:258-62.
92. Chironda G, Bhengu B. Ethical, legal and cultural implications in the management of chronic kidney disease (CKD) patients: a critical review of literature. *Trans R Soc S Afr*. 2016;71(2):137-43.
93. Mechta Nielsen T, Frøjk Juhl M, Feldt-Rasmussen B, Thomsen T. Adherence to medication in patients with chronic kidney disease: a systematic review of qualitative research. *Clin Kidney J*. 2018;11(4):513-27.

94. Fick DM, Cooper JW, Wade WE, Waller JL, Maclean JR, Beers MH. Updating the Beers criteria for potentially inappropriate medication use in older adults: results of a US consensus panel of experts. *Arch Intern Med.* 2003;163(22):2716-24.
95. Campanelli C. American Geriatrics Society updated Beers Criteria for potentially inappropriate medication use in older adults. *J Am Geriatr Soc.* 2012;60(4):616-31.
96. Fick D. Potentially inappropriate medication use in a Medicare managed care population: association with higher costs and utilization. *J Manag Care Pharm.* 2001;7(5):407-13.
97. Fick DM, Mion LC, Beers MH, J LW. Health outcomes associated with potentially inappropriate medication use in older adults. *Res Nurs Health.* 2008;31(1):42-51.
98. Hanlon JT, Fillenbaum GG, Kuchibhatla M, Artz MB, Boulton C, Gross CR, et al. Impact of inappropriate drug use on mortality and functional status in representative community dwelling elders. *Med Care Res Rev.* 2002;40(2):166-76.
99. Stockl KM, Le L, Zhang S, Harada AS. Clinical and economic outcomes associated with potentially inappropriate prescribing in the elderly. *Am J Manag Care.* 2010;16(1):e1-10.
100. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis.* 1987;40(5):373-83.
101. Beddhu S, Bruns FJ, Saul M, Seddon P, Zeidel ML. A simple comorbidity scale predicts clinical outcomes and costs in dialysis patients. *Am J Med.* 2000;108(8):609-13.
102. Gallagher P, Baeyens JP, Topinkova E, Madlova P, Cherubini A, Gasperini B, et al. Inter-rater reliability of STOPP (Screening Tool of Older Persons' Prescriptions) and START (Screening Tool to Alert doctors to Right Treatment) criteria amongst physicians in six European countries. *Age Ageing.* 2009;38(5):603-6.
103. Altman DG. *Practical Statistics for Medical Research.* London: Chapman & Hall; 1991.
104. George J, Phun YT, Bailey MJ, Kong DC, Stewart K. Development and validation of the medication regimen complexity index. *Ann Pharmacother.* 2004;38(9):1369-76.
105. Ghimire S, Peterson GM, Castolino RL, Jose MD, Zaidi ST. Medication Regimen Complexity and Adherence in Haemodialysis Patients: An Exploratory Study. *Am J Nephrol.* 2016;43(5):318-24.
106. Cardone KE, Manley HJ, Grabe DW, Meola S, Hoy CD, Bailie GR. Quantifying home medication regimen changes and quality of life in patients receiving nocturnal home hemodialysis. *Hemodial Int.* 2011;15(2):234-42.
107. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12(3):189-98.
108. Strobel C, Engedal K. Norsk revidert Mini Mental Status Evaluering (MMSE-NR) 2008 [Available from: <http://www.legeforeningen.no/geriatri>].
109. Viswanathan M, Golin CE, Jones CD, et al. Interventions to improve adherence to self-administered medications for chronic diseases in the united states: A systematic review. *Ann Intern Med.* 2012;157(11):785-95.
110. Morisky DE, Ang A, Krousel-Wood M, Ward HJ. Predictive validity of a medication adherence measure in an outpatient setting. *J Clin Hypertens.* 2008;10(5):348-54.
111. Morisky DE, Green LW, Levine DM. Concurrent and predictive validity of a self-reported measure of medication adherence. *Med Care.* 1986;24(1):67-74.
112. Morisky DE, DiMatteo MR. Improving the measurement of self-reported medication nonadherence: response to authors. *J Clin Epidemiol.* 2011;64(3):255-7.

113. Krousel-Wood M, Islam T, Webber LS, Re RN, Morisky DE, Muntner P. New medication adherence scale versus pharmacy fill rates in seniors with hypertension. *Am J Manag Care*. 2009;15(1):59-66.
114. Alkatheri AM, Alyousif SM, Alshabanah N, Albekairy AM, Alharbi S, Alhejaili FF, et al. Medication adherence among adult patients on hemodialysis. *Saudi J Kidney Dis Transpl*. 2014;25(4):762-8.
115. Kefale B, Tadesse Y, Alebachew M, Engidawork E. Management practice, and adherence and its contributing factors among patients with chronic kidney disease at Tikur Anbessa Specialized Hospital: A hospital-based cross-sectional study. *PLoS One*. 2018;13(7):e0200415.
116. Brant R. Assessing proportionality in the proportional odds model for ordinal logistic regression. *Biometrics*. 1990;46(4):1171-8.
117. Korevaar JC, Merkus MP, Jansen MA, Dekker FW, Boeschoten EW, Krediet RT. Validation of the KDQOL-SF: a dialysis-targeted health measure. *Qual Life Res*. 2002;11(5):437-47.
118. Barotfi S, Molnar MZ, Almasi C, Kovacs AZ, Remport A, Szeifert L, et al. Validation of the Kidney Disease Quality of Life-Short Form questionnaire in kidney transplant patients. *J Psychosom Res*. 2006;60(5):495-504.
119. Loge JH, Kaasa S, Hjermsstad MJ, Kvien TK. Translation and performance of the Norwegian SF-36 Health Survey in patients with rheumatoid arthritis. I. Data quality, scaling assumptions, reliability, and construct validity. *J Clin Epidemiol*. 1998;51(11):1069-76.
120. Hays RD, Kallich JD, Mapes DL, Coons SJ, Carter WB. Development of the kidney disease quality of life (KDQOL) instrument. *Qual Life Res*. 1994;3(5):329-38.
121. Ware JE, Kosinski M. SF-36 Physical & Mental Health Summary Scales: A Manual for Users of Version 1: QualityMetric; 2001.
122. O'Mahony D, O'Sullivan D, Byrne S, O'Connor MN, Ryan C, Gallagher P. STOPP/START criteria for potentially inappropriate prescribing in older people: version 2. *Age Ageing*. 2015;44(2):213-8.
123. Counter D, Millar JWT, McLay JS. Hospital readmissions, mortality and potentially inappropriate prescribing: a retrospective study of older adults discharged from hospital. *Br J Clin Pharmacol*. 2018;84(8):1757-63.
124. Parker K, Bull-Engelstad I, Benth JS, Aasebo W, von der Lippe N, Reier-Nilsen M, et al. Effectiveness of using STOPP/START criteria to identify potentially inappropriate medication in people aged ≥ 65 years with chronic kidney disease: a randomized clinical trial. *Eur J Clin Pharmacol*. 2019.
125. Hill-Taylor B, Sketris I, Hayden J, Byrne S, O'Sullivan D, Christie R. Application of the STOPP/START criteria: a systematic review of the prevalence of potentially inappropriate prescribing in older adults, and evidence of clinical, humanistic and economic impact. *J Clin Pharm Ther*. 2013;38(5):360-72.
126. Hill-Taylor B, Walsh KA, Stewart S, Hayden J, Byrne S, Sketris IS. Effectiveness of the STOPP/START (Screening Tool of Older Persons' potentially inappropriate Prescriptions/Screening Tool to Alert doctors to the Right Treatment) criteria: systematic review and meta-analysis of randomized controlled studies. *J Clin Pharm Ther*. 2016;41(2):158-69.
127. Cherubini A, Corsonello A, Lattanzio F. Underprescription of Beneficial Medicines in Older People. *Drugs Aging*. 2012;29(6):463-75.
128. Caruana EJ, Roman M, Hernández-Sánchez J, Solli P. Longitudinal studies. *J Thorac Dis*. 2015;7(11):E537-E40.

129. Dekkers OM, von Elm E, Algra A, Romijn JA, Vandenbroucke JP. How to assess the external validity of therapeutic trials: a conceptual approach. *Int J Epidemiol*. 2010;39(1):89-94.
130. Kennedy-Martin T, Curtis S, Faries D, Robinson S, Johnston J. A literature review on the representativeness of randomized controlled trial samples and implications for the external validity of trial results. *Trials*. 2015;16:495.
131. Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas*. 1960;20(1):37-46.
132. Ryan C, O'Mahony D, Byrne S. Application of STOPP and START criteria: interrater reliability among pharmacists. *Ann Pharmacother*. 2009;43(7):1239-44.
133. Blanco-Reina E, García-Merino MR, Ocaña-Riola R, Aguilar-Cano L, Valdellós J, Bellido-Estévez I, et al. Assessing Potentially Inappropriate Prescribing in Community-Dwelling Older Patients Using the Updated Version of STOPP-START Criteria: A Comparison of Profiles and Prevalences with Respect to the Original Version. *PLoS One*. 2016;11(12):e0167586.
134. Brown JD, Hutchison LC, Li C, Painter JT, Martin BC. Predictive Validity of the Beers and Screening Tool of Older Persons' Potentially Inappropriate Prescriptions (STOPP) Criteria to Detect Adverse Drug Events, Hospitalizations, and Emergency Department Visits in the United States. *J Am Geriatr Soc*. 2016;64(1):22-30.
135. Sanchez-Gili M, Toro-Chico P, Perez-Encinas M, Gomez-Pedrero AM, Portoles-Perez JM. [Pharmaceutical intervention on the therapeutic adherence in patients with chronic renal disease]. *Rev Calid Asist*. 2011;26(3):146-51.
136. Pednekar PP, Agh T, Malmenas M, Raval AD, Bennett BM, Borah BJ, et al. Methods for Measuring Multiple Medication Adherence: A Systematic Review-Report of the ISPOR Medication Adherence and Persistence Special Interest Group. *Value Health*. 2019;22(2):139-56.
137. Moon SJ, Lee WY, Hwang JS, Hong YP, Morisky DE. Accuracy of a screening tool for medication adherence: A systematic review and meta-analysis of the Morisky Medication Adherence Scale-8. *PLoS One*. 2017;12(11):e0187139.
138. Pezzotti P, Scalmana S, Mastromattei A, Di Lallo D, Group tPAW. The accuracy of the MMSE in detecting cognitive impairment when administered by general practitioners: A prospective observational study. *BMC Fam Pract*. 2008;9(1):29.
139. Berger I, Wu S, Masson P, Kelly PJ, Duthie FA, Whiteley W, et al. Cognition in chronic kidney disease: a systematic review and meta-analysis. *BMC Med*. 2016;14(1):206-.
140. Seifter JL, Samuels MA. Uremic encephalopathy and other brain disorders associated with renal failure. *Semin Neurol*. 2011;31(2):139-43.
141. Bugnicourt J-M, Godefroy O, Chillon J-M, Choukroun G, Massy ZA. Cognitive Disorders and Dementia in CKD: The Neglected Kidney-Brain Axis. *J Am Soc Nephrol*. 2013;24(3):353.
142. Watanabe K, Watanabe T, Nakayama M. Cerebro-renal interactions: Impact of uremic toxins on cognitive function. *Neurotoxicology*. 2014;44:184-93.
143. Murray AM, Pederson SL, Tupper DE, Hochhalter AK, Miller WA, Li Q, et al. Acute Variation in Cognitive Function in Hemodialysis Patients: A Cohort Study With Repeated Measures. *Am J Kidney Dis*. 2007;50(2):270-8.
144. Tamadon MR, Adibimehr AR, Ghorbani R. Cognitive Impairment in Dialysis and Non-Dialysis Patients Suffering from Chronic Renal Failure and Comparing Them with a Control Group. *Middle East J Rehabil Health Stud*. 2017;4(4):e12815.

145. Drew DA, Weiner DE. Chronic kidney disease and neurocognition. In: Alosco ML, Stern RA, editors. *The Oxford Handbook of Adult Cognitive Disorders*. New York, NY: Oxford University Press; 2019. p. 137.
146. Brysch EG, Cauthon KAB, Kalich BA, Sarbacker GB. Medication Regimen Complexity Index in the Elderly in an Outpatient Setting: A Literature Review. *Consult Pharm*. 2018;33(9):484-96.
147. Miettinen O. Confounding and effect-modification. *Am J Epidemiol*. 1974;100(5):350-3.
148. Pourhoseingholi MA, Baghestani AR, Vahedi M. How to control confounding effects by statistical analysis. *Gastroenterol Hepatol Bed Bench*. 2012;5(2):79-83.
149. Siegerink B, Dekker FW, Suttorp MM, Jager KJ, Zoccali C. Graphical presentation of confounding in directed acyclic graphs. *Nephrol Dial Transplant*. 2014;30(9):1418-23.
150. Merriam Webster Dictionary. Bias 2019 [Available from: <https://www.merriam-webster.com/dictionary/bias>].
151. Pannucci CJ, Wilkins EG. Identifying and avoiding bias in research. *Plast Reconstr Surg*. 2010;126(2):619-25.
152. Little RJ, D'Agostino R, Cohen ML, Dickersin K, Emerson SS, Farrar JT, et al. The prevention and treatment of missing data in clinical trials. *N Engl J Med*. 2012;367(14):1355-60.
153. Yeatts SD, Martin RH. What is Missing from my Missing Data Plan? *Stroke*. 2015;46(6):e130-e2.
154. Neovius M, Jacobson SH, Eriksson JK, Elinder C-G, Hylander B. Mortality in chronic kidney disease and renal replacement therapy: a population-based cohort study. *BMJ Open*. 2014;4(2).
155. Casals M, Girabent-Farrés M, Carrasco JL. Methodological quality and reporting of generalized linear mixed models in clinical medicine (2000-2012): a systematic review. *PLoS One*. 2014;9(11):e112653-e.
156. Twisk J, de Vente W, Apeldoorn A, de Boer M. Should we use logistic mixed model analysis for the effect estimation in a longitudinal RCT with a dichotomous outcome variable? *Epidemiol Biostat Public Health*. 2017;14(3).
157. Nedin Ranković G, Janković SM, Velićković Radovanović R, Jovic Z, Pesic G, Pavlovic S, et al. Potentially inappropriate prescribing of drugs in elderly patients on chronic hemodialysis treatment. *Clin Nephrol*. 2018;89(6):453-60.
158. Jones SA, Bhandari S. The prevalence of potentially inappropriate medication prescribing in elderly patients with chronic kidney disease. *Postgrad Med J*. 2013;89(1051):247-50.
159. Martins VdS, Mori ALPM, Dorea EL, Pinto GA, Hirata MH, Hirata FDC, et al. Exposure to potentially inappropriate medications in Brazilian elderly outpatients with metabolic diseases. *RBCF*. 2016;52:699-707.
160. Boland B, Guignard B, Dalleur O, Lang PO. Application of STOPP/START and Beers criteria: Compared analysis on identification and relevance of potentially inappropriate prescriptions. *Eur Geriatr Med*. 2016;7(5):416-23.
161. Fialova D, Topinkova E, Gambassi G, Finne-Soveri H, Jonsson PV, Carpenter I, et al. Potentially inappropriate medication use among elderly home care patients in Europe. *JAMA*. 2005;293(11):1348-58.
162. Lam MP, Cheung BMY. The use of STOPP/START criteria as a screening tool for assessing the appropriateness of medications in the elderly population. *Expert Rev Clin Pharmacol*. 2012;5(2):187-97.

-
163. Steinman MA, Rosenthal GE, Landefeld CS, Bertenthal D, Kaboli PJ. Agreement Between Drugs-to-Avoid Criteria and Expert Assessments of Problematic Prescribing. *Arch Intern Med.* 2009;169(14):1326-32.
 164. Neri L, Martini A, Andreucci VE, Gallieni M, Rey LA, Brancaccio D. Regimen complexity and prescription adherence in dialysis patients. *Am J Nephrol.* 2011;34(1):71-6.
 165. Pantuzza LL, Ceccato M, Silveira MR, Junqueira LMR, Reis AMM. Association between medication regimen complexity and pharmacotherapy adherence: a systematic review. *Eur J Clin Pharmacol.* 2017;73(11):1475-89.
 166. Correr CJ, Melchior AC, Fernandez-Llimos F, Pontarolo R. Effects of a pharmacotherapy follow-up in community pharmacies on type 2 diabetes patients in Brazil. *Int J Clin Pharm.* 2011;33(2):273-80.
 167. Negewo NA, Gibson PG, Wark PA, Simpson JL, McDonald VM. Treatment burden, clinical outcomes, and comorbidities in COPD: an examination of the utility of medication regimen complexity index in COPD. *Int J Chron Obstruct Pulmon Dis.* 2017;12:2929-42.
 168. Barnette DJ, Hanks C, Li W, Porter K. Patient-Level Medication Regimen Complexity in an Adolescent and Adult Population with Autism Spectrum Disorders. *Pharmacotherapy.* 2019;39(6):636-44.
 169. Hirsch JD, Metz KR, Hosokawa PW, Libby AM. Validation of a patient-level medication regimen complexity index as a possible tool to identify patients for medication therapy management intervention. *Pharmacotherapy.* 2014;34(8):826-35.
 170. Oosthuizen F, Dhooat E, Kazi S, Masondo B, Omarjee N, Sacoer Z, et al. Assessing the complexity of medicine regimens A pilot study. *Afr J Pharm Pharmacol.* 2011;5(16):1863-6.
 171. Advinha AM, de Oliveira-Martins S, Mateus V, Pajote SG, Lopes MJ. Medication regimen complexity in institutionalized elderly people in an aging society. *Int J Clin Pharm.* 2014;36(4):750-6.
 172. van Wijk CM, van Vliet KP, Kolk AM. Gender perspectives and quality of care: Towards appropriate and adequate health care for women. *Soc Sci Med.* 1996;43(5):707-20.
 173. Kestenbaum B, Sampson JN, Rudser KD, Patterson DJ, Seliger SL, Young B, et al. Serum phosphate levels and mortality risk among people with chronic kidney disease. *J Am Soc Nephrol.* 2005;16(2):520-8.
 174. Giachelli CM. The emerging role of phosphate in vascular calcification. *Kidney Int.* 2009;75(9):890-7.
 175. Tangkiatkumjai M, Walker DM, Praditpornsilpa K, Boardman H. Association between medication adherence and clinical outcomes in patients with chronic kidney disease: a prospective cohort study. *Clin Exp Nephrol.* 2017;21(3):504-12.
 176. Lau DT, Briesacher B, Mercaldo ND, Halpern L, Osterberg EC, Jarzebowski M, et al. Older Patients' Perceptions of Medication Importance and Worth: An Exploratory Study. *Drugs Aging.* 2008;25(12):1061-75.
 177. Park DC, Morrell RW, Frieske D, Kincaid D. Medication adherence behaviors in older adults: effects of external cognitive supports. *Psychol Aging.* 1992;7(2):252-6.
 178. Monane M, Bohn RL, Gurwitz JH, Glynn RJ, Avorn J. Noncompliance With Congestive Heart Failure Therapy in the Elderly. *Arch Intern Med.* 1994;154(4):433-7.
 179. Huang E, Segev DL, Rabb H. Kidney Transplantation in the Elderly. *Semin Nephrol.* 2009;29(6):621-35.

180. Gonzalez-Lopez A, Nava-Rebollo A, Chocarro-Martinez A, Andres-Martin B, Santana-Zapatero H, Herrera-Gomez F, et al. Potentially inappropriate prescribing in patients on dialysis using STOPP-START criteria. *Nefrología (English Edition)*. 2019;39(1):92-4.
181. Motter FR, Fritzen JS, Hilmer SN, Paniz EV, Paniz VMV. Potentially inappropriate medication in the elderly: a systematic review of validated explicit criteria. *Eur J Clin Pharmacol*. 2018;74(6):679-700.
182. Berger AK, Duval S, Krumholz HM. Aspirin, beta-blocker, and angiotensin-converting enzyme inhibitor therapy in patients with end-stage renal disease and an acute myocardial infarction. *J Am Coll Cardiol*. 2003;42(2):201-8.
183. Tonelli M, Bohm C, Pandeya S, Gill J, Levin A, Kiberd BA. Cardiac risk factors and the use of cardioprotective medications in patients with chronic renal insufficiency. *Am J Kidney Dis*. 2001;37(3):484-9.
184. Gowdak LH, Arantes RL, de Paula FJ, Krieger EM, De Lima JJ. Underuse of American College of Cardiology/American Heart Association Guidelines in hemodialysis patients. *Ren Fail*. 2007;29(5):559-65.
185. Wauters M, Elseviers M, Vaes B, Degryse J, Dalleur O, Vander Stichele R, et al. Too many, too few, or too unsafe? Impact of inappropriate prescribing on mortality, and hospitalization in a cohort of community-dwelling oldest old. *Br J Clin Pharmacol*. 2016;82(5):1382-92.
186. Anderson K, Stowasser D, Freeman C, Scott I. Prescriber barriers and enablers to minimising potentially inappropriate medications in adults: a systematic review and thematic synthesis. *BMJ Open*. 2014;4(12):e006544.
187. Heser K, Pohontsch NJ, Scherer M, Loffler A, Luck T, Riedel-Heller SG, et al. Perspective of elderly patients on chronic use of potentially inappropriate medication - Results of the qualitative CIM-TRIAD study. *PLoS One*. 2018;13(9):e0202068.
188. Karanicolas PJ, Farrokhyar F, Bhandari M. Practical tips for surgical research: blinding: who, what, when, why, how? *Can J Surg*. 2010;53(5):345-8.
189. Kaplan B, Mason NA, Shimp LA, Ascione FJ. Chronic Hemodialysis Patients. Part I: Characterization and Drug-Related Problems. *Ann Pharmacother*. 1994;28(3):316-9.
190. Gallagher PF, O'Connor MN, O'Mahony D. Prevention of potentially inappropriate prescribing for elderly patients: a randomized controlled trial using STOPP/START criteria. *Clin Pharmacol Ther*. 2011;89(6):845-54.
191. Department of Health, National Service Framework for Older People,. Medicines and older people: implementing medications-related aspects of the NSF for older people. In: Health. Do, editor. London 2001.
192. Marcum ZA, Hanlon JT, Murray MD. Improving medication adherence and health outcomes in older adults: an evidence-based review of randomized controlled trials. *Drugs Aging*. 2017;34(3):191-201.
193. Volume CI, Farris KB, Kassam R, Cox CE, Cave A. Pharmaceutical care research and education project: patient outcomes. *J Am Pharm Assoc (Wash)*. 2001;41(3):411-20.
194. Lam KD, Miao Y, Steinman MA. Cumulative changes in the use of long-term medications: a measure of prescribing complexity. *JAMA Intern Med*. 2013;173(16):1546-7.
195. Steinman MA, Low M, Balicer RD, Shadmi E. Impact of a nurse-based intervention on medication outcomes in vulnerable older adults. *BMC Geriatr*. 2018;18(1):207-.
196. Aziz VM, Hill N, Kumar S. Completed audit cycle to explore the use of the STOPP/START toolkit to optimise medication in psychiatric in-patients with dementia. *BJPsych Bull*. 2018;42(1):37-41.

14 LIST OF ERRATA

Page 66, reference 3, correction. Original text: ... *Sociaology* ... Corrected text: ... *Sociology*...

Page 66, reference 16, correction. Original text: ... *Dictionary C* ... Corrected text: ... *Collins Dictionary* ...

Page 66, reference 36, correction. Original text: *Sakshaug S.Legemiddelforbruket, i Norge 2013-2017 (Drug Consumption in Norway 2013-2017). Legemiddelstatistikk 2018*; Corrected text: *Berg C. Reseptregisteret 2013-2017(The Norwegian Prescription Database 2013-2017).Oslo,Norge: Folkehelseinstituttet 2018.*

Page 66+68, references 11+42, correction. Original text: ... *Dictionary MW* ... Corrected text: ... *Merriam Webster Dictionary* ...

Page 66, 67+69, references 1, 34+70, correction. Original text: ... *Organization WH*... Corrected text: ...*World Health Organization*...; correction. Original text: ... *an older peson* ... Corrected text: ... *an older person* ...

PAPERS

SHORT COMMUNICATION

Potentially Inappropriate Medications in Elderly Haemodialysis Patients Using the STOPP Criteria

Krystina Parker^{1,2} · Willy Aasebø¹ · Knut Stavem^{2,3,4}

Published online: 8 August 2016

© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract

Background Polypharmacy is commonly applied to elderly haemodialysis patients for treating terminal renal failure and multiple co-morbidities. Potentially inappropriate medications (PIMs) in multidrug regimens in geriatric populations can be identified using specially designed screening tools.

Objective The aims of this study were to estimate the prevalence of PIMs by applying the Screening Tool of Older Persons' Prescriptions (STOPP) criteria and the Beers criteria to elderly haemodialysis patients and to assess the association of some risk factors with the presence of PIMs.

Methods Fifty-one elderly haemodialysis patients participated; their median age was 74 (range 65–89) years, and 77 % of them were male. Demographic data, co-morbidity and medication lists were collected from the electronic medical records of the patients. The STOPP criteria were applied by two physicians independently to identify PIMs. The association of some risk factors with PIMs were assessed using Fisher's exact test.

Results The patients used a median of 13 (range 7–21) medications per day. The overall prevalence of PIMs using

the STOPP criteria was 63 %, and using the Beers criteria was 43 %. The most prevalent PIMs were proton-pump inhibitors. Benzodiazepines and first-generation antihistamines were related to side effects such as falls in the previous 3 months, and calcium-channel blockers were associated with chronic constipation. The number of PIMs was not significantly associated with number of medications, age, sex and co-morbidity.

Conclusions The STOPP criteria revealed a high prevalence of PIMs in a population of elderly patients receiving haemodialysis.

Key Points

Potentially inappropriate medications according to STOPP criteria were identified in 63 % of elderly haemodialysis patients and in 43 % using the Beers criteria.

Proton pump inhibitors were the most prevalent drugs according to STOPP criteria.

The number of potentially inappropriate medications was not significantly associated with number of medications, age, sex and co-morbidity.

✉ Krystina Parker
krystina.parker@medisin.uio.no

¹ Medical Division, Department of Nephrology, Akershus University Hospital, 1478 Lørenskog, Norway

² Institute of Clinical Medicine, University of Oslo, Oslo, Norway

³ Medical Division, Department of Pulmonary Medicine, Akershus University Hospital, Lørenskog, Norway

⁴ HØKH, Department of Health Services Research, Akershus University Hospital, Lørenskog, Norway

1 Background

Elderly haemodialysis patients are considered to be a vulnerable group due to the presence of renal failure with underlying co-morbidities that require the use of multiple

drugs. Drug efficacy in these patients is influenced by age-related changes, altered nutritional state and the haemodialysis treatment they receive. Haemodialysis alters the pharmacokinetics and pharmacodynamics of many drugs due to the presence of continuous changes in the fluid balance and uraemic toxin levels, and this may increase the risk of drug-to-drug interactions and adverse side effects in haemodialysis patients.

International guidelines for management of patients with renal failure point to the importance of dose reduction or discontinuation of certain drugs and provide cautionary notes for prescribing (Kidney Disease Improving Global Outcome [KDIGO] 2013) [1] or recommend regular review of medication lists (Kidney Disease Outcome Quality Initiative [KDOQI]) [2]. Several other approaches can be used to identify potentially inappropriate medications (PIMs), such as a targeted multidisciplinary team approach, consulting with pharmacists [3–5] or using validated screening tools [6–8]. Two recent studies used such a screening tool called the Beers criteria [9, 10], reporting a high prevalence of PIMs in elderly patients with chronic kidney disease and end-stage renal disease (ESRD). The Screening Tool of Older Persons' Prescriptions (STOPP) was used in two randomized trials involving geriatric populations, in which it contributed to a significant reduction of PIMs both at the time of discharge after acute hospitalization and up to 6 months after discharge [11], as well as reductions in the number of falls and costs [12]. Two European studies involving geriatric populations found that the number of PIMs was higher when using the STOPP criteria than when using the Beers criteria [11, 13]. However, to our knowledge the STOPP criteria have not been previously applied to elderly haemodialysis patients.

The aims of this study were to determine the prevalence of PIMs using the STOPP criteria and the Beers criteria in elderly haemodialysis patients and to assess the association of age, sex, number of medications and co-morbidity as risk factors for PIMs in this population.

2 Methods

2.1 Study Population and Data Collection

There were 102 patients treated in the dialysis centre of Akershus University Hospital in Norway between July and December 2012, and those aged ≥ 65 years were asked to participate in this study. Fifty-one patients were eligible for inclusion, and they all agreed to participate. The following information was collected from the electronic medical records of the patients: age, sex, number of prescribed medications, cause of kidney disease, severity of co-morbidity (classified using the Charlson Co-morbidity Index

[CCI]), time on haemodialysis and dialysis treatment quality index (quantified as urea clearance (Kt/V)). Supplementary information about some medication side effects was collected from patient interviews, such as falls during the previous 3 months, chronic constipation (for more than 3 months before starting haemodialysis) and current dizziness when at home.

2.2 Identification of Potentially Inappropriate Medications

PIMs were identified using the STOPP criteria and the updated Beers Criteria [6, 14]. The STOPP criteria were developed and validated through a Delphi consensus process by 18 experts in geriatric pharmacotherapy and include 65 indicators to identify important drug-to-drug and drug-to-disease interactions [6]. The STOPP criteria exhibit a high sensitivity in detecting PIMs and good inter-rater reliability [6, 15, 16]. Two physicians independently applied the STOPP and Beers criteria to review the medication lists of all of the patients included in the present study. In case of disagreement in identified PIMs between the two raters a consensus was achieved by open discussion.

We also collected information on the use of nephrology-specific drugs, which are not included in STOPP criteria, such as phosphate binders, Vitamin D analogues, active vitamin D and erythropoietin.

2.3 Statistical Analyses

The data were analysed using SPSS statistical software (version 20, IBM, SPSS, Chicago, IL, USA). Descriptive statistics are presented using mean $\pm$ SD or median (range) values. Fisher's exact test was used to compare the number of PIMs between groups dichotomized according to age, sex, CCI, time in dialysis and number of medications based on those equal to or above the median versus those below the median.

We applied a 5 % significance cut-off in two-sided tests. Inter-rater agreement between the two raters for assessments using the STOPP and Beers screening tools is presented with % agreement (proportion of cases for which the raters agree) and kappa coefficient for 2×2 tables. Strength of reliability coefficients was defined using the following nomenclature [17]: <0.20 poor, 0.21 – 0.40 fair, 0.41 – 0.60 moderate, 0.61 – 0.80 good, 0.81 – 1.00 very good.

3 Results

The clinical and demographic characteristics of the study population are presented in Table 1. The most common co-morbidities were hypertension, cardiovascular disease and

diabetes mellitus. The patients used a total of 652 medications and a median of 13 (range 7–21) different medications each day, and of these, a median of 5.2 (range 2–8) drugs

Table 1 Characteristics of the study population ($n = 51$)

Sex, male	39 (77)
Age, years, median (range)	74 (65–89)
Dialysis	
Time in haemodialysis, months, median (range)	8 (0–108)
Urea clearance, Kt/V , mean $\pm$ SD	1.47 $\pm$ 0.29
Arteriovenous fistula	29 (57)
Haemodialysis catheter	22 (43)
Diagnosis	
Nephrosclerosis	18 (35)
Post-renal disease ^a	8 (16)
Other ^b	8 (16)
Diabetic nephropathy	7 (14)
Unknown	7 (14)
Glomerulonephritis	3 (6)
Co-morbidities	
Charlson Co-morbidity Index, median (range)	5 (2–9)
Hypertension	47 (92)
Cardiovascular disease	28 (55)
Diabetes mellitus	21 (41)
Malignancies ^c	13 (25)
Atrial fibrillation	12 (24)
COPD ^d	11 (22)
Number of medications, median (range)	13 (7–21)

Data are given as number (%) except where stated otherwise

^a Post-renal disease = hydronephrosis, kidney-stone disease or retroperitoneal fibrosis

^b Other = kidney cancer, loss of kidney graft, amyloidosis, poisoning or anti-glomerular-basement-membrane nephritis

^c Malignancies = prostate cancer, colorectal cancer, testicular cancer, lung cancer, kidney cancer, lymphoma or myelomatosis

^d COPD = chronic obstructive pulmonary disease

Table 2 Potentially inappropriate medications identified by the Screening Tool of Older Persons' Prescriptions

Cardiovascular system	
Calcium-channel blocker with chronic constipation	7 (14)
Aspirin (>150 mg/day)	4 (8)
Gastrointestinal system	
Proton-pump inhibitor for peptic ulcer disease at full therapeutic dosage for >8 weeks	11 (22)
Loperamide	2 (4)
Central nervous system	
Long-term administration of long-acting benzodiazepines	6 (12)
First-generation antihistamine (prolonged use for >1 week)	6 (12)
Tricyclic antidepressants with an opiate or calcium-channel blocker	2 (4)
Opiates	2 (4)
Endocrine system	
Beta-blockers in those with diabetes mellitus and frequent hypoglycaemic episodes	1 (2)
Total number of potentially inappropriate medications	41

Data are given as number (%)

can be defined as nephrology-specific drugs. PIMs were identified in 32 patients (63 %) by using STOPP criteria. Table 2 lists the most prevalent inappropriate medications according to the STOPP criteria. Twenty-one patients (42 %) had one PIM and 11 patients (22 %) had two PIMs. Nineteen patients (37 %) reported falls during the previous 3 months, nine patients (18 %) reported chronic constipation before start of dialysis and three patients (6 %) reported dizziness. The most prevalent PIMs according to STOPP were proton pump inhibitors at full therapeutic dosage use ($n = 11$; 22 %), calcium-channel blockers ($n = 7$; 14 %) in patients with chronic constipation and benzodiazepines ($n = 6$; 12 %) in those prone to falls.

The number of PIMs did not differ significantly with age (≤ 74 vs. >74 years, $p = 0.39$), sex (male vs. female, $p = 0.74$), co-morbidity severity (CCI ≤ 5 vs. >5 , $p = 1.00$), time in haemodialysis (≤ 8 months vs. >8 months, $p = 0.78$) or number of medications (≤ 13 vs. >13 , $p = 0.56$).

Beers criteria defined PIMs in 22 patients (43 %). The most common PIM according to the Beers criteria was proton pump inhibitors (Table 3).

The kappa coefficient between raters for the STOPP criteria was 0.42 (0.24–0.56, 95 % CI) and for the Beers criteria kappa was 1.0. The % agreement was 0.78 for STOPP and 1.00 for Beers, respectively. The disagreements occurred within two of the STOPP criteria on the interpretation of adverse effects: history of falls and chronic constipation.

4 Discussion

The prevalence of PIMs based on the STOPP criteria was high (63 %) in this study of elderly Norwegian haemodialysis patients. No previous studies have presented

Table 3 Potentially inappropriate medications identified by the Beers criteria

Cardiovascular system	
Amiodarone	2 (4)
Gastrointestinal system	
Proton-pump inhibitors (over >8 weeks avoid, unless for high-risk patients, risk of <i>Clostridium difficile</i> infection, bone loss and fractures)	10 (20)
Metoclopramide (can cause extrapyramidal effects)	1 (2)
Central nervous system	
Selective serotonin reuptake inhibitors with a history of falls	4 (8)
Long-acting and short-acting benzodiazepines with a history of falls	3 (6)
Tricyclic antidepressants with a history of falls	2 (4)
Total number of potentially inappropriate medications	22

Data are given as number (%)

use of the STOPP criteria in a haemodialysis population. However, our findings are in line with two recent studies involving patients with chronic kidney disease that applied Beers criteria [9, 10]. In previous studies involving general geriatric populations the prevalence of PIMs ranged widely (21.4–79 %) when applying the STOPP criteria [18–22]. The variations between the reported prevalence of PIMs can probably be attributed to differences in study populations [6, 11, 13, 16, 20, 23–25].

The present study chose to apply the STOPP criteria because of their documented association with a reduction in PIMs in randomized studies [11, 12], and because the STOPP criteria reportedly identify more PIMs than the Beers criteria in European elderly patients [11, 13, 23]. In the present study, the prevalence of PIMs was higher with the STOPP criteria than with the Beers criteria, which is in line with previous reports [13, 26]. The inter-rater reliability of STOPP was lower than in previous reports [6, 15, 27]. However, these differences should be interpreted with caution because of differences in study designs and populations.

Previous studies involving elderly populations found that the number of PIMs was positively associated with the number of medications, age and number of co-morbidities [11, 23, 25]. In contrast, the present study found no associations between the number of PIMs and age, sex, co-morbidity severity, time on haemodialysis and number of medications. This difference between the studies may be due to differences in the case mix between the populations or the small number of participants in the present study.

This study had several limitations. The patients were relatively old and the number of patients was small, and so the results cannot be generalized to all patients with ESRD. The small sample size also limited the analysis of risk factors for PIMs, and multivariable analysis was not considered meaningful. The study did not include a comparison group. Finally, the STOPP criteria were not developed specifically for use in patients with ESRD and they do not evaluate several drugs commonly used by patients with

renal impairment, such as phosphate binders, erythropoietin, potassium binders and calcimimetics.

5 Conclusions

In conclusion, the prevalence of PIMs among elderly haemodialysis patients was high when applying the STOPP criteria and the Beers criteria, even though these criteria do not evaluate nephrology-specific drugs. In contrast, the STOPP criteria identified more PIMs compared with the Beers criteria in our study population. This finding supports the use of medication review to identify and avoid PIMs in haemodialysis patients, who are typically exposed to polypharmacy.

Compliance with Ethical Standards

Ethical Approval This study was approved by the ethics committee of Akershus University Hospital and was performed in accordance with the ethical standards of the Declaration of Helsinki. All participants gave their consent at the start of the study.

Funding No funding was received for the conduct of the study or the preparation of this manuscript.

Conflict of interest Krystina Parker, Willy Aasebø and Knut Stavem have no conflicts of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any noncommercial use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. KDIGO. Kidney disease improving global outcome. 2013. <http://kdigo.org/home/guidelines/>.

2. KKDOQI. 2015. <https://www.kidney.org/professionals/guidelines/hemodialysis2015>.
3. Kaur S, Mitchell G, Vitetta L, et al. Interventions that can reduce inappropriate prescribing in the elderly: a systematic review. *Drugs Aging*. 2009;26(12):1013–28.
4. Lutters M, Harbarth S, Janssens JP, et al. Effect of a comprehensive, multidisciplinary, educational program on the use of antibiotics in a geriatric university hospital. *J Am Geriatr Soc*. 2004;52(1):112–6.
5. St Peter WL. Management of polypharmacy in dialysis patients. *Semin Dial*. 2015;28(4):427–32.
6. Gallagher P, Ryan C, Byrne S, et al. STOPP (Screening Tool of Older Person's Prescriptions) and START (Screening Tool to Alert doctors to Right Treatment). Consensus validation. *Int J Clin Pharmacol Ther*. 2008;46(2):72–83.
7. Beers MH. Explicit criteria for determining potentially inappropriate medication use by the elderly. An update. *Arch Intern Med*. 1997;157(14):1531–6.
8. McLeod PJ, Huang AR, Tamblyn RM, et al. Defining inappropriate practices in prescribing for elderly people: a national consensus panel. *CMAJ*. 1997;156(3):385–91.
9. Kondo N, Nakamura F, Yamazaki S, et al. Prescription of potentially inappropriate medications to elderly hemodialysis patients: prevalence and predictors. *Nephrol Dial Transplant*. 2015;30(3):498–505.
10. Jones SA, Bhandari S. The prevalence of potentially inappropriate medication prescribing in elderly patients with chronic kidney disease. *Postgrad Med J*. 2013;89(1051):247–50.
11. Gallagher PF, O'Connor MN, O'Mahony D. Prevention of potentially inappropriate prescribing for elderly patients: a randomized controlled trial using STOPP/START criteria. *Clin Pharmacol Ther*. 2011;89(6):845–54.
12. Frankenthal D, Lerman Y, Kalendariev E, et al. Intervention with the screening tool of older persons potentially inappropriate prescriptions/screening tool to alert doctors to right treatment criteria in elderly residents of a chronic geriatric facility: a randomized clinical trial. *J Am Geriatr Soc*. 2014;62(9):1658–65.
13. Hamilton H, Gallagher P, Ryan C, et al. Potentially inappropriate medications defined by STOPP criteria and the risk of adverse drug events in older hospitalized patients. *Arch Intern Med*. 2011;171(11):1013–9.
14. American Geriatrics Society 2015 updated Beers criteria for potentially inappropriate medication use in older adults. *J Am Geriatr Soc*. 2015;63(11):2227–46.
15. Gallagher P, Baeyens JP, Topinkova E, et al. Inter-rater reliability of STOPP (Screening Tool of Older Persons' Prescriptions) and START (Screening Tool to Alert doctors to Right Treatment) criteria amongst physicians in six European countries. *Age Ageing*. 2009;38(5):603–6.
16. Ryan C, O'Mahony D, O'Donovan DO, et al. A comparison of the application of STOPP/START to patients' drug lists with and without clinical information. *Int J Clin Pharm*. 2013;35(2):230–5.
17. Altman DG. Practical statistics for medical research. London: Chapman & Hall; 1991.
18. Garcia-Gollarte F, Baleriola-Julvez J, Ferrero-Lopez I, et al. Inappropriate drug prescription at nursing home admission. *J Am Med Dir Assoc*. 2012;13(1):83.e9–15.
19. Ryan C, O'Mahony D, Kennedy J, et al. Potentially inappropriate prescribing in an Irish elderly population in primary care. *Br J Clin Pharmacol*. 2009;68(6):936–47.
20. Hill-Taylor B, Sketris I, Hayden J, et al. Application of the STOPP/START criteria: a systematic review of the prevalence of potentially inappropriate prescribing in older adults, and evidence of clinical, humanistic and economic impact. *J Clin Pharm Ther*. 2013;38(5):360–72.
21. Liu CL, Peng LN, Chen YT, et al. Potentially inappropriate prescribing (IP) for elderly medical inpatients in Taiwan: a hospital-based study. *Arch Gerontol Geriatr*. 2012;55(1):148–51.
22. Parsons C, Johnston S, Mathie E, et al. Potentially inappropriate prescribing in older people with dementia in care homes: a retrospective analysis. *Drugs Aging*. 2012;29(2):143–55.
23. O'Sullivan DP, O'Mahony D, Parsons C, et al. A prevalence study of potentially inappropriate prescribing in Irish long-term care residents. *Drugs Aging*. 2013;30(1):39–49.
24. Frankenthal D, Lerman Y, Lerman Y. The impact of hospitalization on potentially inappropriate prescribing in an acute medical geriatric division. *Int J Clin Pharm*. 2015;37(1):60–7.
25. Onatade R, Auyeung V, Scutt G, et al. Potentially inappropriate prescribing in patients on admission and discharge from an older peoples' unit of an acute UK hospital. *Drugs Aging*. 2013;30(9):729–37.
26. Gallagher P, Lang PO, Cherubini A, et al. Prevalence of potentially inappropriate prescribing in an acutely ill population of older patients admitted to six European hospitals. *Eur J Clin Pharmacol*. 2011;67(11):1175–88.
27. Ryan C, O'Mahony D, Byrne S. Application of STOPP and START criteria: interrater reliability among pharmacists. *Ann Pharmacother*. 2009;43(7):1239–44.

Effectiveness of using STOPP/START criteria to identify potentially inappropriate medication in people aged ≥ 65 years with chronic kidney disease: a randomized clinical trial

Krystina Parker^{1,2} · Ingrid Bull-Engelstad³ · Jūratė Šaltytė Benth^{2,4} · Willy Aasebø¹ · Nanna von der Lippe⁵ · Morten Reier-Nilsen³ · Ingrid Os^{2,5} · Knut Stavem^{2,4,6}

Received: 13 April 2019 / Accepted: 17 July 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Purpose Polypharmacy and inappropriate prescribing are common in elderly with chronic kidney disease (CKD). This study identified potentially inappropriate prescriptions (PIPs) and potential prescribing omissions (PPOs) using the Screening Tool of Older Persons' Prescriptions (STOPP) and the Screening Tool to Alert doctors to the Right Treatment (START) criteria in elderly with advanced CKD and determined the effect of a medication review on medication adherence and health-related quality of life (HRQoL).

Methods The intervention consisted of a medication review using STOPP/START criteria with a recommendation to a nephrologist or similar review without a recommendation. End points were prevalence of PIP and PPO, medication adherence, and HRQoL. Group differences in outcomes were assessed using a generalized linear mixed model. The trial was registered under www.clinicaltrials.gov (ID: NCT02424786).

Results We randomized 180 patients with advanced CKD (mean age 77 years, 23% female). The prevalence of PIPs and PPOs in the intervention group was 54% and 50%, respectively. The odds of PPOs were lower in the intervention than the control group (OR 0.42, 95% CI 0.19–0.92, $p = 0.032$), while there was no intergroup difference in the number of PIPs (OR 0.57, CI 0.27–1.20, $p = 0.14$). There was no difference in changes in medication adherence or HRQoL from baseline to 6 months between the groups.

Conclusions The intervention with the STOPP/START criteria identified a high prevalence of inappropriate medications in the elderly with advanced CKD and reduced the number of PPOs. However, there was no detectable impact of the intervention on medication adherence or HRQoL.

Keywords Polypharmacy · Chronic kidney disease · Medication adherence · Elderly · Inappropriate medication

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00228-019-02727-9>) contains supplementary material, which is available to authorized users.

✉ Krystina Parker
krystina.parker@medisin.uio.no

¹ Department of Nephrology, Medical Division, Akershus University Hospital, Lørenskog, Norway

² Faculty of Medicine, Institute of Clinical Medicine, University of Oslo, Oslo, Norway

³ Department of Nephrology, Medical Division, Vestre Viken HF, Drammen Hospital, Drammen, Norway

⁴ Department of Health Services Research Unit, Akershus University Hospital, Lørenskog, Norway

⁵ Department of Nephrology, Medical Division, Oslo University Hospital Ullevål, Oslo, Norway

⁶ Department of Pulmonary Medicine, Medical Division, Akershus University Hospital, Lørenskog, Norway

Introduction

Polypharmacy is common in patients with advanced chronic kidney disease (CKD). Patients with CKD at stages 2 to 5 take an average of 8 different medications, while dialysis patients typically take 10 to 12 different medications [1, 2]. Polypharmacy is associated with adverse drug events (ADEs) and inappropriate medications [3, 4], that is, medications whose risks outweigh their benefits. Inappropriate medications are associated with morbidity, mortality, ADEs, and higher costs [5, 6].

Specialized instruments such as the Screening Tool of Older Persons' Prescriptions (STOPP), the Screening Tool to Alert doctors to Right Treatment (START), and Beers Criteria may be used to detect potentially inappropriate medication (PIM) [7, 8]. The STOPP/START criteria are used worldwide, whereas the Beers Criteria are widely used in the USA,

Australia, and Asia [9]. The STOPP criteria aim to identify potentially inappropriate prescriptions (PIPs), whereas the START criteria identify potential prescribing omissions (PPOs). PIPs include any prescription without a clear clinical indication, dosage, and duration inappropriate for clinical use (overprescribing) or prescribing a medication with an adverse risk-benefit profile when safer alternatives are available (misprescribing). PPOs define any medication that is not prescribed despite having a clear clinical indication (underprescribing) [10].

A recent meta-analysis assessed the effectiveness of applying the STOPP/START criteria in general hospital facilities, nursing homes, or frail elderly included only four randomized trials [11]. Those authors concluded that the use of these criteria was associated with reductions in inappropriate prescriptions, falls, delirium episodes, hospital length of stay, and medication costs, but no improvement was observed in health-related quality of life (HRQoL) or mortality [11]. The effectiveness of using the STOPP/START criteria to detect inappropriate medications in older adults with advanced CKD has not been reported previously.

The present randomized clinical trial used the STOPP/START criteria to identify PIPs and PPOs in patients aged ≥ 65 years with CKD at stage 5 treated either conservatively or with dialysis. For the assessment of medication adherence and its change, we used the eight-item Morisky Medication Adherence Scale (MMAS-8), which has also been used in many previous studies [12]. HRQoL was assessed with the SF-12, which is commonly used in geriatric patients and patients with CKD [13, 14]. We hypothesized that the use of STOPP/START criteria would identify inappropriate medications and lead to improved medication adherence, and this would be associated with improved HRQoL. The primary objectives were to identify any differences in the numbers of PIPs and PPOs and in medication adherence between the intervention and control groups after 6 months of follow-up. The secondary objectives were to determine differences between the groups in the average number of medications and HRQoL scores over the 6-month follow-up.

Materials and methods

Study design and population

This clinical trial had a single-blind, multicenter, parallel-group randomized design. Patients were included from three nephrology centers (Akershus University Hospital; Oslo University Hospital, Ullevål; and Vestre Viken Hospital Trust, Drammen) from July 2015 to January 2017. All patients aged ≥ 65 years with CKD at stage 5 (estimated glomerular filtration rate < 15 ml/min/1.73 m², as calculated by the Chronic Kidney Disease Epidemiology Collaboration

equation) who were either treated conservatively or with peritoneal dialysis (PD) or hemodialysis (HD) were asked to participate. Patients in the three hospitals were identified from local registers of patients receiving HD or PD, or from the list of patients with CKD scheduled for outpatient visits. We excluded patients with a severe-to-moderate reduction in cognitive function (Mini Mental State Examination–Norwegian Revision (MMSE-NR) score of < 23 before or during the study) [15], severe hearing or visual impairment, or inadequate knowledge of the Norwegian language.

The National Committee for Medical and Health Research Ethics (South/East) and the Akershus University Hospital data protection officer approved the study. The study was conducted in accordance with the Declaration of Helsinki. The patients were given information both orally and in writing, and written consents were obtained. The trial was registered at www.clinicaltrials.gov (ID: NCT02424786).

Baseline data collection

All of the included patients participated in a semi-structured interview prior to randomization. The interviewer asked closed- and open-ended questions, and no audio recording for transcription was made during the interview. Information was collected on medications that included over-the-counter medication, administration mode, side effects, concomitant diseases, accidental falls, and specific symptoms such as vertigo, obstipation, pruritus, or dyspepsia. An accidental fall was defined as “any event in which a person inadvertently or intentionally comes to rest on the ground or another lower level” during the previous 3 months [16]. Falls that occurred after the HD sessions were considered as adverse events.

At the end of the interview, each patient was given questionnaires about HRQoL and medication adherence, which were to be answered at home and returned in postage-prepaid envelopes. Non-respondents received one reminder by telephone.

The Charlson Comorbidity Index (CCI) was calculated at baseline based on information in the medical record of a subject and supplementary information obtained during the interview. This index consists of an aggregate of 19 comorbidities that are weighted and summarized into a single number. The CCI is an accurate predictor of the 2-year mortality in patients with CKD [17, 18].

Randomization and intervention

All patients were randomly assigned at a ratio of 1:1 to the intervention and control groups using random numbers generated by a computer program, with the allocations stored in sealed numbered envelopes. The investigators and physicians were blinded to group allocation until the first interview was

completed, while all of the subjects remained blinded throughout the project.

The intervention consisted of a medication review by one investigator, a nephrologist (KP), who used the STOPP/START criteria to identify possible inappropriate medications. In case of an identified inappropriateness, a follow-up recommendation was written in the electronic medical record to inform the attending physician. These recommendations comprised simple statements explaining why this medication was identified as inappropriate. The attending nephrologists were free to decide whether or not they would comply with the recommendations and revise a patient's current medication list. For the control group, the same investigator performed an identical medication review using the STOPP/START criteria, but no notes were entered into the medical record.

Follow-up data collection

After 6 months, the same investigator carried out a second round of semi-structured interviews with all of the participants using similar questions as in the baseline interview about all medications, administration mode, possible side effects, and information about changes in medications. The same data as at baseline were registered, which also included new comorbidities and the number of hospitalization during the previous 6 months. Each participant was given the same questionnaires about HRQoL and medication adherence to be completed at home and returned by mail.

Outcome measurements

Screening tool to identify potentially inappropriate prescriptions and potential prescribing omissions

The STOPP/START version 2 criteria consist of 80 STOPP and 34 START criteria and have been validated using the Delphi consensus method [19]. The STOPP/START criteria are grouped according to organ systems (e.g., cardiovascular system and musculoskeletal system) to facilitate easy and rapid medication reviews. For each criterion, the tool contains a brief explanation of why a medication or a combination of medicines is considered appropriate or potentially inappropriate.

Medication adherence

The eight-item Morisky Medication Adherence Scale (MMAS-8) is a self-reported and validated questionnaire on the adherence to medication whose total score ranges from 0 (non-adherent) to 8 (adherent) [20–22]. Seven items have a “yes/no” response, and the eighth is scored from 1 to 4. For further analysis of data, medication adherence was

dichotomized into non-adherent (MMAS-8 score < 6) and adherent (MMAS-8 score $\geq$ 6) [23].

Health-related quality of life

We used the 12-item Short-Form Health Survey (SF-12) to assess the patients' HRQoL, which is part of the Kidney Disease Quality of Life Instrument. The SF-12 is self-administered and has been validated in various patient groups, including CKD [24]. The physical and mental health statuses were aggregated into two summary scores in this study: the physical component summary (PCS) score and the mental component summary (MCS) score [25].

Trial end points

The primary end points were the reduction in PIPs and PPOs and the improvement of medication adherence during the 6-month observation period. The secondary end points were the change in the number of medications and PCS and MCS scores.

Statistical analysis

Patient characteristics were described as number (percentage), mean $\pm$ SD, or median (minimum–maximum) values, as appropriate. Baseline characteristics were compared between respondents and non-respondents using the independent-samples *t* test for continuous variables or the χ^2 test or Fisher's exact test for categorical variables.

The differences between the intervention and control group at follow-up were assessed using a linear mixed model for the continuous outcomes (number of medications and PCS and MCS scores), while a generalized linear mixed model was applied to estimate the dichotomous outcomes (medication adherence: adherent versus non-adherent), PIPs according to STOPP criteria (none versus $\geq$ 1), and PPOs according to START criteria (none versus $\geq$ 1). The models contained fixed effects for measurement time point (baseline or 6-month follow-up) and the interaction between the time point and group (intervention or control). The interaction term in such a model quantifies differences between groups at follow-up adjusted for the baseline values. Random effects for patients were included. The models were estimated for cases with observations available at baseline. For sensitivity analyses, we conducted complete-case analysis using longitudinal analysis of covariance for the continuous outcomes (fixed effects for the group and baseline values) and binary logistic regression analysis for the dichotomous outcomes (fixed effects for the group).

Prior to the inclusion of patients, we estimated that to detect a difference of 0.5 standard deviations (SDs) in medication adherence score between the groups with a statistical power

of 80% at a significance level of 5%, the study would need a sample size of 63 subjects in each group.

The analyses were performed with STATA (version 15.1, StataCorp, College Station, TX, USA) or SAS (version 9.4, SAS Institute, Cary, NC, USA). All tests were two-sided, and results with p values below 0.05 were considered statistically significant.

Results

Study population

We recruited and randomized 180 patients from 319 eligible patients (Fig. 1). In total, 11% of the patients included at baseline were lost to follow-up due to death and 17% due to no response or incomplete questionnaires. This group was defined as non-respondents ($n = 50$). Respondents and non-respondents had the same baseline characteristics except for the latter having a lower MMSE-NR score ($p = 0.025$) and more hospitalizations during the follow-up period ($p = 0.014$).

The baseline characteristics did not differ between the allocated groups (Table 1).

Identification of potentially inappropriate medications

Among the 180 included patients, 265 inappropriate medications (PIPs and PPOs) according to the STOPP/START

criteria were found at baseline (Supplementary Table 1). The most common PIPs were proton-pump inhibitors, benzodiazepines, and first-generation antihistamines, while the most common PPOs were angiotensin-converting enzyme (ACE) inhibitor, statins, and vitamin D. The prevalence of PIPs at baseline was 54% in the intervention group and 55% in the control group; the corresponding prevalence rates of PPOs were 50% and 56%, respectively.

Outcomes of the intervention

Primary outcomes

The number of patients with one or more PIPs decreased in the intervention group whereas it remained almost the same in the control group (Supplementary Table 2). The probability of PIPs did not differ between the intervention and control groups at follow-up, whereas that of PPOs was lower in the intervention group than the control group (odds ratio (OR) = 0.42, 95% confidence interval (CI) = 0.19–0.92, $p = 0.032$) (Table 2). In the control group, we identified no severe PIPs or PPOs. After 6 months, there was no difference between the groups in medication adherence (Table 2).

Secondary outcomes

There was no significant intergroup difference in the average number of medications or HRQoL score at follow-up (Table 3).

Sensitivity analysis

The sensitivity analyses confirmed the above-mentioned results. There were no intergroup differences in the average number of medications, PCS and MCS scores, medication adherence, or PIPs. The probability of PPOs remained lower in the intervention group than the control group ($n = 161$; OR = 0.46, 95% CI = 0.24–0.87, $p = 0.017$).

Discussion

Using the STOPP/START criteria to screen medication in patients with advanced CKD revealed a high prevalence of PIMs and a reduction in PPOs during the 6-month follow-up period in this study. However, the intervention did not lead to improvements in the number of PIPs, medication adherence, average number of medications, or HRQoL scores. We are not aware of any previous study that has assessed the use of STOPP/START criteria in older people with advanced CKD.

The number of PPOs was the only variable that was reduced in the present study. This contrasts with previous reports of improvements in both PIPs and PPOs after similar

Fig. 1 Flow chart of study inclusion

Table 1 Characteristics of the populations ($n = 180$) in the two study groups

	Intervention (<i>n</i> = 92)		Control (<i>n</i> = 88)	
Sex, female	23	(25)	23	(26)
Age, years	76.0 ± 6.6		76.0 ± 7.6	
Geriatric features				
Mini-Mental State Examination Norwegian Revision score	28	(23–30)	28	(24–30)
Polypharmacy (> 5 drugs)	87	(95)	86	(98)
Living alone	31	(34)	36	(41)
≥ 1 fall within 3 months prior to inclusion	14	(15)	10	(11)
Symptoms				
Vertigo	25	(27)	20	(23)
Obstipation	22	(24)	22	(25)
Pruritus	46	(50)	32	(36)
Dyspepsia	13	(14)	8	(9)
Most frequent comorbidities				
Hypertension	77	(85)	67	(76)
Coronary disease	34	(39)	37	(42)
Malignancy	28	(31)	31	(35)
Diabetes mellitus	30	(34)	22	(25)
Atrial fibrillation	11	(12)	14	(16)
Charlson Comorbidity Index ≥4	61	(66)	62	(70)
Nephrological treatment				
Hemodialysis	46	(43)	41	(47)
Peritoneal dialysis	11	(12)	9	(10)
Conservative treatment	35	(38)	38	(43)

Data are number (percentage), mean $\pm$ SD, or median (minimum–maximum) values

interventions using the STOPP/START criteria [11]. Our literature search did not reveal other randomized trials evaluating the effect of using the STOPP/START criteria on clinical outcomes in older people with advanced CKD. The lower impact in the present study compared to previous studies may have been due to the implementation of suggested medication changes being left entirely to the judgment of the attending physician, differences in populations and settings, or the length of the follow-up period. A relatively short follow-up was chosen in this study because of an expected high mortality rate in our older cohort [26], and this was also used in previous randomized controlled trials [11].

The prevalence of PIMs in the present cohort as detected with the STOPP/START version 2 criteria was lower than that in a recent survey of HD patients using the same criteria [27], but similar to those in previous descriptive studies using the STOPP/START version 1 criteria [28, 29]. The differences in the prevalence of PIMs between studies may be caused by differences in the populations, medication practice or traditions, or the use of different versions of the STOPP/START criteria, which also makes interstudy comparisons more difficult.

The PIMs identified in the present study were almost identical to those reported in general geriatric or HD populations [27, 28, 30]. In both arms of the present study, more than 50% of the participants had omissions of recommended preventive medications or medications with documented therapeutic effects in advanced CKD. Such underprescribing is common in older people with or without CKD, such as in treatments for primary or secondary cardiovascular prevention with a beta-blocker after acute myocardial infarction, or statins and vitamin D in CKD [31–34]. This issue is complicated by current guidelines for lipid management in CKD stratifying statin treatment according to the stage of CKD [35].

The present study identified ACE inhibitor as the most common medication omission by applying the START criteria. Patients with CKD are regarded as a high-risk group for cardiovascular events; however, ACE inhibitors are often neglected in this patient group [36–38]. ACE inhibitors are cardio-protective, reduce vascular morbidity and mortality, and slow the progression of renal failure directly and indirectly by meticulous blood pressure control [36, 37]. However, the use of ACE inhibitors in advanced CKD is complicated by risks of side effects, such as hyperkalemia, metabolic acidosis,

Table 2 Differences in primary outcomes between the intervention group and the control group (reference) from a generalized linear mixed model

	<i>n</i>	Odds ratio ^a (95% CI)	<i>p</i>
Potentially inappropriate prescriptions according to STOPP criteria (none versus ≥ 1)	180	0.57 (0.27 to 1.20)	0.14
Potentially inappropriate omissions according to START criteria (none versus ≥ 1)	180	0.42 (0.19 to 0.92)	0.032
Medication adherence (adherent versus non-adherent)	157	1.17 (0.41 to 3.39)	0.77

^a Overall intervention effect over time, or difference between groups at follow-up, with no adjustment for baseline, *CI* confidence interval

and a possibility of further decline of GFR. Therefore, in some patients, nephrologists may justify omission of an ACE inhibitor.

Underprescribing may be attributed to an intention to avoid polypharmacy, lack of knowledge or limited evidence for use of the drugs, or because older people with advanced CKD are regularly excluded from clinical trials [39].

The intervention in the present study did not improve medication adherence. The adherence was high at both baseline and the 6-month follow-up, which supports previous reports of high medication adherence in older patients [40, 41]. This lack of response may therefore be explained by the high medication adherence at baseline, a ceiling effect of the instrument used for assessment or the definition used for high adherence excluding a large proportion of patients from further possible improvements. The findings are in line with those of a pharmacist-led intervention in community-dwelling older people [42].

The lack of improvement in HRQoL in the present study cannot be compared with similar interventions in patients with CKD, as this is the first study in advanced CKD [42]. However, the HRQoL scores of our population were low, and in accordance with those found in other patients with advanced CKD [43, 44].

Only a few previous trials have investigated the changes in the number of medications in cohorts of older people. Our finding of no effect on the average number of medications is in line with that of another randomized trial [45] but contrasts

with observations made in patients with dementia and the residents of nursing homes [9, 46].

The present study was a multicenter, randomized, clinical trial with older patients with advanced CKD and a high vulnerability due to polypharmacy and impaired kidney function. The population included in this study is likely to be representative of this patient population in Norway. As the study was randomized, it by design adjusted for both observable and non-observable covariates.

Some other challenges in this study should be noted. When calculating the sample size, we used the medication adherence score as the outcome; however, we had no figures from comparable studies in the literature about the size of effect to be expected. Furthermore, the ceiling effect of the instrument may also have limited the possible improvement, possibly contributing to an underestimation of the number of patients needed to detect a statistical significance. The study had a high proportion of missing data for the outcomes, and it would not be feasible to impute these missing values. We used a generalized linear mixed model to include all patients with available baseline data. Complete-case analyses were performed as sensitivity analyses to assess the robustness of the results and whether the assumptions made in the analyses were valid [47]. Non-respondents had lower MMSE-NR scores and higher hospitalization rates than the respondents, suggesting that the latter were healthier and less frail.

A medication review according to STOPP/START was performed in control patients, but not entered into the

Table 3 Differences in secondary outcomes between the intervention group and the control group (reference) from a linear mixed model

	<i>n</i>	Coefficient ^a (95% CI)	<i>p</i>
Number of medications	180	0.40 (− 0.29 to 1.09)	0.26
Quality of life, SF-12			
Physical component summary score	148	1.04 (− 1.89 to 3.98)	0.48
Mental component summary score	148	1.39 (− 1.27 to 4.06)	0.30

^a Regression coefficient represents overall intervention effect over time, or difference between groups at follow-up, with adjustment for baseline, *CI* confidence interval

medical records. If *severe inappropriateness* would have been detected among the controls, this might have represented an ethical dilemma. During the course of this study, however, we did not identify severe inappropriateness in this group. In case of severe inappropriateness, recommendations would have been written in the electronic medical record or the issue would have been discussed directly with the attending nephrologist, which we think would be in agreement with the general principles of the declaration of Helsinki.

The attending physicians who were responsible for implementing medication changes in the study did follow patients in both the intervention and control groups. It is therefore possible that there was a learning effect from the recommendations for medication changes in the intervention group, which may have benefited the control group, that is, a spill-over effect. This effect may have contributed to underestimations of the differences between the intervention and control groups.

The STOPP/START criteria were developed with a primary focus on the medication for older patients generally and not for specific disease populations. The findings obtained when applying the STOPP/START criteria therefore enable comparisons between different patient categories; however, it might not be feasible to apply the criteria in all settings. A limitation of the present study was that the validity and clinical relevance of these criteria in patients with advanced CKD have not yet been documented. Therefore, the decision of implementing the recommendations was left to the judgment of the attending nephrologist. In general, little is known about the clinical relevance of PIPs or PPOs, as detected using the STOPP/START criteria, at an individual level [48]. However, a recent study of the elderly with hip fracture showed that one in two PIM is clinically relevant at the individual level [49].

Each administered medication should ideally be appropriate, evidence-based, and safe, independent of the specific diseases or age of the patient. The use of a simple screening tool such as the STOPP/START criteria can facilitate a dialog between those involved in the prescribing, dispensing, and administering medications and the affected patients, and in theory, should help to improve the appropriateness of medication regimens. Finding feasible and effective ways to benefit from such processes and documenting their impact on medication adherence and ultimately the HRQoL of patients require further research.

In conclusion, in this study, an intervention based on the STOPP/START criteria detected PIMs and reduced the number of PPOs, but it did not lead to a reduction in the number of PIPs or improvements in medication adherence or HRQoL.

Acknowledgments The use of the Morisky Medication Adherence Scale (© MMAS) is protected by US copyright laws. Permission for use is required. A license agreement is available from Donald E. Morisky at dmorisky@gmail.com or MMAS Research LLC 14725 NE 20th ST. Bellevue, WA 98007.

Authors' contributions K.P. and K.S. designed the study with support from the other authors. K.P., M.R.N., I.B.E., W.A., and N.v.d.L. participated in screening, recruitment, and data collection. K.S., J.S.B., and K.P. analyzed the data. K.P. drafted the manuscript with support from K.S. All of the authors contributed to data interpretation, critically reviewed the manuscript, and approved its final version.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Informed consent Informed consent was obtained from all individual participants included in the study.

References

- Bailie GR, Eisele G, Liu L, Roys E, Kiser M, Finkelstein F, Wolfe R, Port F, Burrows-Hudson S, Saran R (2005) Patterns of medication use in the RRI-CKD study: focus on medications with cardiovascular effects. *Nephrol Dial Transplant* 20(6):1110–1115
- St Peter WL (2015) Management of polypharmacy in dialysis patients. *Semin Dial* 28(4):427–432
- Taché SV, Sönnichsen A, Ashcroft DM (2011) Prevalence of adverse drug events in ambulatory care: a systematic review. *Ann Pharmacother* 45(7–8):977–989
- Cadogan CA, Ryan C, Hughes CM (2016) Appropriate polypharmacy and medicine safety: when many is not too many. *Drug Saf* 39(2):109–116
- Spinewine A, Schmader KE, Barber N, Hughes C, Lapane KL, Swine C, Hanlon JT (2007) Appropriate prescribing in elderly people: how well can it be measured and optimised? *Lancet* 370(9582):173–184
- Hamilton H, Gallagher P, Ryan C, Byrne S, O'Mahony D (2011) Potentially inappropriate medications defined by STOPP criteria and the risk of adverse drug events in older hospitalized patients. *Arch Intern Med* 171(11):1013–1019
- Gallagher P, Ryan C, Byrne S, Kennedy J, O'Mahony D (2008) STOPP (Screening Tool of Older Person's Prescriptions) and START (Screening Tool to Alert doctors to Right Treatment). Consensus validation. *Int J Clin Pharmacol Ther* 46(2):72–83
- Beers MH, Ouslander JG, Rollingher I, Reuben DB, Brooks J, Beck JC (1991) Explicit criteria for determining inappropriate medication use in nursing home residents. UCLA Division of Geriatric Medicine. *Arch Intern Med* 151(9):1825–1832
- Curtin D, Gallagher PF, O'Mahony D (2019) Explicit criteria as clinical tools to minimize inappropriate medication use and its consequences. *Ther Adv Drug Saf* 10:2042098619829431
- O'Mahony D, Gallagher PF (2008) Inappropriate prescribing in the older population: need for new criteria. *Age Ageing* 37(2):138–141
- Hill-Taylor B, Walsh KA, Stewart S, Hayden J, Byrne S, Sketris IS (2016) Effectiveness of the STOPP/START (Screening Tool of Older Persons' potentially inappropriate Prescriptions/Screening Tool to Alert doctors to the Right Treatment) criteria: systematic

- review and meta-analysis of randomized controlled studies. *J Clin Pharm Ther* 41(2):158–169
12. Moon SJ, Lee WY, Hwang JS, Hong YP, Morisky DE (2017) Accuracy of a screening tool for medication adherence: a systematic review and meta-analysis of the Morisky Medication Adherence Scale-8. *PLoS One* 12(11):e0187139
 13. Wyld M, Morton RL, Hayen A, Howard K, Webster AC (2012) A systematic review and meta-analysis of utility-based quality of life in chronic kidney disease treatments. *PLoS Med* 9(9):e1001307
 14. (2015) American Geriatrics Society 2015 Updated Beers Criteria for Potentially Inappropriate Medication Use in Older Adults. *J Am Geriatr Soc* 63(11):2227–2246
 15. Strobel C, Engedal K (2008) Norsk revidert Mini Mental Status Evaluering (MMSE-NR). <http://www.legeforeningen.no/geriatri>
 16. World Health Organization (2018) Falls. <http://www.who.int/news-room/fact-sheets/detail/falls>. Accessed 3 June 2019
 17. Charlson ME, Pompei P, Ales KL, MacKenzie CR (1987) A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 40(5):373–383
 18. van Manen JG, Korevaar JC, Dekker FW, Boeschoten EW, Bossuyt PMM, Krediet RT (2002) How to adjust for comorbidity in survival studies in ESRD patients: a comparison of different indices. *Am J Kidney Dis* 40(1):82–89
 19. O'Mahony D, O'Sullivan D, Byrne S, O'Connor MN, Ryan C, Gallagher P (2015) STOPP/START criteria for potentially inappropriate prescribing in older people: version 2. *Age Ageing* 44(2): 213–218
 20. Berlowitz DR, Foy CG, Kazis LE, Bolin LP, Conroy MB, Fitzpatrick P, Gure TR, Kimmel PL, Kirchner K, Morisky DE, Newman J, Olney C, Oparil S, Pajewski NM, Powell J, Ramsey T, Simmons DL, Snyder J, Supiano MA, Weiner DE, Whittle J, Group SR (2017) Effect of intensive blood-pressure treatment on patient-reported outcomes. *N Engl J Med* 377(8):733–744
 21. Morisky DE, Ang A, Krousel-Wood M, Ward HJ (2008) Predictive validity of a medication adherence measure in an outpatient setting. *J Clin Hypertens* 10(5):348–354
 22. Morisky DE, DiMatteo MR (2011) Improving the measurement of self-reported medication nonadherence: response to authors. *J Clin Epidemiol* 64(3):255–257
 23. Yang A, Wang B, Zhu G, Jiao Z, Fang Y, Tang F, Ma C, Zhao Y, Cheng C, Zhong M (2014) Validation of Chinese version of the Morisky Medication Adherence Scale in patients with epilepsy. *Seizure* 23(4):295–299
 24. Korevaar JC, Merkus MP, Jansen MA, Dekker FW, Boeschoten EW, Krediet RT (2002) Validation of the KDQOL-SF: a dialysis-targeted health measure. *Qual Life Res* 11(5):437–447
 25. Hays RD, Kallish JD, Mapes DL, Coons SJ, Carter WB (1994) Development of the kidney disease quality of life (KDQOL) instrument. *Qual Life Res* 3(5):329–338
 26. System USRD (2017) United States Renal Data System. 2017 USRDS annual data report: epidemiology of kidney disease in the United States. Chapter 5: Mortality. National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD, 2017. https://www.usrds.org/2018/view/v2_05.aspx. Accessed 13 June 2018
 27. Gonzalez-Lopez A, Nava-Rebollo A, Chocarro-Martinez A, Andres-Martin B, Santana-Zapatero H, Herrera-Gomez F, Diego-Martin J, Escaja-Muga C, Grande-Villoria J (2019) Potentially inappropriate prescribing in patients on dialysis using STOPP-START criteria. *Nefrología (English Edition)* 39(1):92–94
 28. Parker K, Aasebø W, Stavem K (2016) Potentially inappropriate medications in elderly haemodialysis patients using the STOPP criteria. *Drugs - Real World Outcomes* 3(3):359–363
 29. Nedin Ranković G, Janković SM, Veličković Radovanović R, Jovic Z, Pesic G, Pavlovic S, Rankovic B, Rankovic J, Stokanovic D, Krtinic D (2018) Potentially inappropriate prescribing of drugs in elderly patients on chronic hemodialysis treatment. *Clin Nephrol* 89(6):453–460
 30. Frankenthal D, Lerman Y, Kalendariev E, Lerman Y (2014) Intervention with the screening tool of older persons potentially inappropriate prescriptions/screening tool to alert doctors to right treatment criteria in elderly residents of a chronic geriatric facility: a randomized clinical trial. *J Am Geriatr Soc* 62(9):1658–1665
 31. Soumerai SB, McLaughlin TJ, Spiegelman D, Hertzmark E, Thibault G, Goldman L (1997) Adverse outcomes of underuse of beta-blockers in elderly survivors of acute myocardial infarction. *JAMA* 277(2):115–121
 32. Gallagher P, Lang PO, Cherubini A, Topinkova E, Cruz-Jentoft A, Montero Errasquin B, Madlova P, Gasperini B, Baeyens H, Baeyens JP, Michel JP, O'Mahony D (2011) Prevalence of potentially inappropriate prescribing in an acutely ill population of older patients admitted to six European hospitals. *Eur J Clin Pharmacol* 67(11):1175–1188
 33. Navaneethan SD, Pansini F, Strippoli GF (2006) Statins in patients with chronic kidney disease: evidence from systematic reviews and randomized clinical trials. *PLoS Med* 3(5):e123
 34. Jean G, Souberbielle JC, Chazot C (2017) Vitamin D in chronic kidney disease and dialysis patients. *Nutrients* 9(4):E328
 35. Sarnak MJ, Bloom R, Muntner P, Rahman M, Saland JM, Wilson PWF, Fried L (2015) KDOQI US commentary on the 2013 KDIGO clinical practice guideline for lipid management in CKD. *Am J Kidney Dis* 65(3):354–366
 36. Gowdak LH, Arantes RL, de Paula FJ, Krieger EM, De Lima JJ (2007) Underuse of American College of Cardiology/American Heart Association Guidelines in hemodialysis patients. *Ren Fail* 29(5):559–565
 37. Tonelli M, Bohm C, Pandeya S, Gill J, Levin A, Kiberd BA (2001) Cardiac risk factors and the use of cardioprotective medications in patients with chronic renal insufficiency. *Am J Kidney Dis* 37(3): 484–489
 38. Berger AK, Duval S, Krumholz HM (2003) Aspirin, beta-blocker, and angiotensin-converting enzyme inhibitor therapy in patients with end-stage renal disease and an acute myocardial infarction. *J Am Coll Cardiol* 42(2):201–208
 39. Barry PJ, Gallagher P, Ryan C, O'Mahony D (2007) START (screening tool to alert doctors to the right treatment)—an evidence-based screening tool to detect prescribing omissions in elderly patients. *Age Ageing* 36(6):632–638
 40. Griva K, Lai AY, Lim HA, Yu Z, Foo MW, Newman SP (2014) Non-adherence in patients on peritoneal dialysis: a systematic review. *PLoS One* 9(2):e89001
 41. Ghimire S, Castelino RL, Lioufas NM, Peterson GM, Zaidi ST (2015) Nonadherence to medication therapy in haemodialysis patients: a systematic review. *PLoS One* 10(12):e0144119
 42. Marcum ZA, Hanlon JT, Murray MD (2017) Improving medication adherence and health outcomes in older adults: an evidence-based review of randomized controlled trials. *Drugs Aging* 34(3):191–201
 43. Brown EA, Johansson L, Farrington K, Gallagher H, Sensky T, Gordon F, Da Silva-Gane M, Beckett N, Hickson M (2010) Broadening Options for Long-term Dialysis in the Elderly (BOLDE): differences in quality of life on peritoneal dialysis compared to haemodialysis for older patients. *Nephrol Dial Transplant* 25(11):3755–3763
 44. Tsai HB, Chao CT, Chang RE, Hung KY, Group CS (2017) Conservative management and health-related quality of life in end-stage renal disease: a systematic review. *Clin Invest Med* 40(3):E127–E134
 45. Gallagher PF, O'Connor MN, O'Mahony D (2011) Prevention of potentially inappropriate prescribing for elderly patients: a randomized controlled trial using STOPP/START criteria. *Clin Pharmacol Ther* 89(6):845–854

46. Aziz VM, Hill N, Kumar S (2018) Completed audit cycle to explore the use of the STOPP/START toolkit to optimise medication in psychiatric in-patients with dementia. *BJPsych Bull* 42(1):37–41
47. Little RJ, D'Agostino R, Cohen ML, Dickersin K, Emerson SS, Farrar JT, Frangakis C, Hogan JW, Molenberghs G, Murphy SA, Neaton JD, Rotnitzky A, Scharfstein D, Shih WJ, Siegel JP, Stern H (2012) The prevention and treatment of missing data in clinical trials. *N Engl J Med* 367(14):1355–1360
48. Cooper JA, Cadogan CA, Patterson SM, Kerse N, Bradley MC, Ryan C, Hughes CM (2015) Interventions to improve the appropriate use of polypharmacy in older people: a Cochrane systematic review. *BMJ Open* 5(12):e009235
49. Lönnbro J, Wallerstedt SM (2017) Clinical relevance of the STOPP/START criteria in hip fracture patients. *Eur J Clin Pharmacol* 73(4): 499–505

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.