Roadmap towards a harmonized pan-European surveillance of obesity-related lifestyle behaviors and their determinants in children and adolescents

Running title: Harmonized pan-European surveillance in children and adolescents

5 Antje Hebestreit^{1*}, Barbara Thumann¹, Maike Wolters¹, Jens Bucksch², Inge Huybrechts³,
Joanna Inchley⁴, Cornelia Lange⁵, Nanna Lien⁶, Kristin Manz⁵, Nadia Slimani³, Hidde P. van
der Ploeg⁷, Wolfgang Ahrens^{1, 8} on behalf of the DEDIPAC Consortium

¹ Leibniz Institute for Prevention Research and Epidemiology – BIPS, Bremen, Germany

10 ² University of Education Heidelberg, Heidelberg, Germany

³ International Agency for Research on Cancer, Lyon, France

⁴ University of St Andrews, St Andrews, United Kingdom

⁵ Robert Koch Institute, Berlin, Germany

⁶ University of Oslo, Oslo, Norway

15 ⁷ VU University Medical Center, Amsterdam, The Netherlands

⁸ University of Bremen, Institute of Statistics, Bremen, Germany

***Address for correspondence:** Dr. Antje Hebestreit, Leibniz Institute for Prevention

20 Research and Epidemiology – BIPS, Achterstr. 30, D-28359 Bremen, Germany;

Tel. +49-421-218.56849, Fax +49-421-218.56821, hebestreit@leibniz-bips.de

Conflict of Interest

25 The authors declare that they have no competing interests.

Abstract

Objectives: To develop a roadmap towards a harmonized pan-European surveillance system for children and adolescents.

Methods: Representatives of five European surveillance systems and the German Health Interview and Examination Survey for Children and Adolescents contributed to the roadmap through a structured workshop in 2016.

Results: A conceptual framework for this roadmap was developed with seven action points (AP) guiding the successive cross-country harmonisation. First, key indicators of health behavior and their determinants in children and adolescents will be identified (AP1,2); short screening instruments will be developed and implemented to assess and monitor key indicators (AP3,4). In parallel, optional supplementary modules could be implemented to provide objective data (AP5). This would allow mutual calibration and improvement of existing instruments before their progressive replacement by more comparable measurement tools (AP6). Establishment of a competence platform is envisaged for guiding the harmonization process (AP7).

Conclusion: This approach builds on existing systems, provides comparable key health indicators across European regions, helps to assess temporal trends, and - once in place - will facilitate health reporting and monitoring of national and international health targets.

Keywords: surveillance, policy evaluation, health, diet, physical activity, sedentary behavior

Abbreviations

AP – Action Points; ASA24 - Automated Self-Administered 24-hour Recall; DEDIPAC - Determinants of Diet and Physical Activity; ECHI - European Core Health Indicators; EGHI - Expert Group on Health Information; EFSA - European Food Safety Authority; EHIS - European Health Interview Survey; FFQ – Food Frequency Questionnaire; HBSC – Healthy Behavior in School Children; HEPA PAT - Health-enhancing physical activity policy audit tool; INFORMAS - International Network for Food and Obesity/ non-communicable Diseases Research, Monitoring and Action Support; KiGGS - German Health Interview and Examination Survey for Children and Adolescents; NHANES - National Health and Nutrition Examination Survey; OECD - Organisation for Economic Co-operation and Development; SB – Sedentary behaviour; WHO – World Health Organization; WHO-COSI - WHO European Childhood Obesity Surveillance Initiative; 24HDR – 24-hour dietary recall; PA – Physical Activity

The Vienna Declaration on Nutrition and Non-communicable Diseases and the European child and adolescent health strategy 2015–2020 (World Health Organization 2014; World Health Organization Europe 2013), acknowledged the disease burden caused by unhealthy lifestyle patterns and its major health, societal and economic impacts in Europe.

65 To address these challenges a series of actions were adopted and member states from the World Health Organization (WHO) European region committed themselves to support nutrition and health related actions and surveillance systems across the life course (World Health Organization 2016). European member states have already recognised the need for harmonized public health surveillance systems to obtain comparable data across countries
70 and to align their policies, action plans and recommendations to combat unhealthy diets, physical inactivity and overweight within the European region (World Health Organization - Europe 2014). However, one major obstacle to comparing the prevalence of unhealthy behaviors and health outcomes, such as overweight and obesity, across countries is the lack of harmonized data based on objective methods and standardized protocols.

75 Within the Determinants of Diet and Physical Activity (DEDIPAC) Knowledge Hub (Brug et al. 2017) we conducted an inventory of existing (pan-) European surveillance systems (Bel-Serrat et al. 2017) including initiatives based on the following criteria: 1) national, regional (e.g. Nordic Monitoring System) and international initiatives (e.g. WHO European Childhood Obesity Surveillance Initiative (WHO-COSI)) collecting data on dietary,
80 physical activity (PA), sedentary behaviour (SB) and their determinants; and 2) surveillance systems currently on-going, or with at least one recently completed wave, carried out on a periodic basis. Most importantly, the inventory reported that only a few surveillance systems involve young populations. Internationally harmonized and comparable data on school-aged children and adolescents are currently provided by two international surveillance systems,
85 the WHO-COSI and the Health Behavior in School-aged Children: WHO Collaborative Cross-National survey (HBSC). Together, they cover the age groups 6 - 9.9 years (WHO-COSI) and 11, 13 and 15 years (HBSC), but the comparability between these systems is limited. In

addition, health data of younger children (<6 years) are not captured by current international surveillance systems despite that this developmental period is considered critical for later health outcomes.

We propose a roadmap towards a harmonized pan-European surveillance system for the monitoring of lifestyle behaviors and their main determinants in children and adolescents. The end goal is to maximize coverage and comparability of data and provide a robust evidence base to inform policy and related actions to improve young people's health across the European region.

Methods

Identification of existing surveillance systems covering children and adolescents

Based on the results of the DEDIPAC inventory (Bel-Serrat et al. 2017) we identified existing European surveillance systems that involve young populations and that either provide state-of-the-art instruments or that have already created a pan-European infrastructure potentially serving as a structural basis to successively develop a harmonized surveillance system. Harmonization in this respect refers to the process of minimizing differences in comparability of measures, variables and methods, so that outcomes are comparable. Further, standardization refers to the process of developing and implementing methodological or technical standards based on the consensus of all relevant parties to maximize comparability, interoperability, repeatability, or quality.

****INSERT TABLE 1 HERE****

Structured workshop

Five international surveillance systems were selected as suitable: WHO-COSI, HBSC, EU-Menu, the European Health Interview Survey (EHIS) and the Nordic Monitoring of Diet, Physical Activity and Overweight (Table 1). Additionally, the GloboDiet initiative was selected as an initiative that provides a standardized instrument suitable for assessing harmonized dietary data. According to the DEDIPAC inventory (Bel-Serrat et al. 2017), objectively measured PA and biomarkers were not analysed in international surveillance systems, however, PA data and sampling collection was conducted in several national monitoring surveys, such as the German Health Interview and Examination Survey for Children and Adolescents (KiGGS study); and so KiGGS was selected to serve as a model for the implementation of objective measurement methods and the collection of biosamples in population-based surveys (Kamtsiuris et al. 2007). Representatives of these surveillance systems contributed to the roadmap through a structured workshop in April 2016.

Results

Conceptual framework

The proposed conceptual framework for this roadmap focuses on children's and adolescents' health and shall guide the future development and implementation by a series of process-oriented action points (AP)(Figure).

****INSERT FIGURE HERE****

Figure 1: Conceptual framework for the roadmap towards the establishment of a standardized children and adolescents health surveillance system

First, we aim to develop and prioritise an agreed set of indicators at the individual- and setting-level according to the different areas of interest: dietary intake, dietary behavior,

PA, SB, health status (AP1) and their determinants (AP2). Out of those, key indicators with a high priority for measuring policy impact will be selected (e.g. consumption of soft drinks (Expert Group on Health Information (EGHI))). A set of core variables (or proxies) to assess the selected key indicators will be developed (e.g. consumption frequency of soft drinks) (AP3). Then, state-of-the-art methods/instruments to measure those variables (e.g. food frequency questionnaire, 'how often do you drink soft drinks per day?') will be identified and compiled in short screening instruments (screeners) which should be incorporated into existing surveillance systems (in the full sample)(AP4). This screening tool is a set of simple standard instruments/questions measuring variables that are needed to describe the most relevant indicators in the full surveillance sample. In parallel, a set of supplementary objective measurement methods (such as activity trackers, mHealth technology) may be implemented among sub-samples in these systems (AP5).

Before integration of new harmonized measurement instruments into existing surveillance systems can take place, their feasibility needs to be assessed in pilot studies. Subsequent methodological studies will allow calibration of existing instruments across surveys. Eventually, the gradual replacement of some of the current non-harmonized questions and measurements by new measurement modules would result in harmonization across existing systems (AP6). A methodological competence platform would be essential for coordinating and guiding the methodological studies that need to accompany the development, implementation and calibration of existing instruments in the long run (AP7). From early on the proposed approach will provide a core set of harmonized data for health reporting, benchmarking and monitoring of national and international health targets.

Discussion

Identification and prioritization of key indicators (AP1 and AP2)

Key indicators of health behaviors (AP1)

Surveillance systems need to assess indicators that are related to health targets to inform progress and performance of health policies. In 2012, a Joint Action of the European Commission and the member states resulted in a shortlist of 88 European Core Health Indicators (ECHI), classified by policy area, such as health determinants (National Institute for Public Health and the Environment 2012). Several criteria had to be met by the ECHI short list indicators (Expert Group on Health Information (EGHI)), e.g. indicators should support potential policy action (at the EU and Member State level); should build on existing indicator systems (e.g.OECD Health Data); should already be in use as widely as possible; and should focus on major public health problems (including health inequalities).and on the best potentials for effective policy action, both at the EU and Member State level.

As an example, indicators of dietary patterns like consumption frequency of sugar sweetened beverages (Malik et al. 2013), fast food (Rosenheck 2008) or fruits and vegetables (Wang et al. 2014) meet these requirements. These food items affect the quality of the diet and are associated with public health outcomes, such as overweight and obesity (Fernandez-Alvira et al. 2017). Further, eating patterns dominated by high energy-dense foods (Hebestreit et al. 2014) or breakfast consumption (Pearson et al. 2009) are often used for public health monitoring in children and adolescents (Currie et al. 2000).

In relation to PA, frequency, duration and intensity are typically measured across different domains (e.g. transport, leisure time, or organized PA) as well as assessment of overall PA level expressed in relation to meeting the current PA recommendations (Poitras et al. 2016). Objective assessment of PA and SB has higher validity and should preferably be used in parallel to supplement self-report assessments. However, most existing population-based surveillance systems are still solely dependent on self-report due to cost and practical

considerations. PA and SB have been reported to be associated with overweight/obesity, metabolic disorders and cardiovascular disease in children and adolescents (de Moraes et al. 2015). Screen time is often used as a proxy for total SB (McVeigh et al. 2016) but is only a partial indicator of time spent sitting and presents a number of methodological challenges (Atkin et al. 2012). Moreover, screen time involves exposure to marketing of unhealthy foods that may in itself be related to the same health outcomes as SB, mediated through food choice (Olafsdottir et al. 2014). Hence, objective assessment of SB (together with PA) would make an important addition to surveillance systems.

For the proposed roadmap, priority will be given to established indicators used by European member states and pan-European public health initiatives. For the selection of key indicators clear criteria will be applied, such as 1) evidence of association with health or health-related behaviors; 2) importance for health status; 3) usefulness for analyzing policy impact; and 4) usefulness for analyzing implementation facilitators and barriers, taking ECHI criteria also into account (see under AP1). The selection process itself will be accomplished in multiple steps. First, indicators will be identified after a literature review and will then be prioritized by experts during a workshop and a subsequent Delphi expert survey. Second, a preliminary set of indicators will then be mapped against available monitoring and surveillance data provided for example, by the WHO STEPSwise approach or the EHIS initiative. This work will be accomplished within the Policy Evaluation Network <https://www.jpi-pen.eu>, which runs until 2020.

Key indicators of determinants (AP2)

The ECHI shortlist includes 14 indicators of health determinants, including socio-economic, health status and intervention indicators (Table 2). During the last decades it became evident that individual-level determinants of dietary behavior are often overridden by environment-level (upstream) determinants in youth (Sleddens et al. 2015) (Table 3). Therefore, relevant determinants at the environmental level of schools (e.g. availability and

accessibility of un-/healthy foods/ drinks in schools and other settings (Buck et al. 2013)),
and family (e.g. number of family meals (Hebestreit et al. 2017)) should both be considered
215 in order to assess the impact of the policies at both levels and to assess their interactions.

With regard to PA, self-efficacy and perceived social support at the individual level
(Bauman et al. 2012) combined with indicators of well implemented physical education and
PA programs at the school level and with indicators of land use in the neighbourhood/
community may be useful to assess policy impact (Heath et al. 2012).

220 Knowledge on SB and its determinants in children and adolescents is currently
increasing: Apart from age – older adolescents are more sedentary – also gender, socio-
economic status of the family, weight status and environmental determinants are known to
be correlated to time spent in SB (Stierlin et al. 2015). As ECHI provides predominantly
screen-use as an indicator for SB, more evidence is required and determinants need to be
225 identified for surveillance purposes.

Additionally, we will consider indicators used by the International Network for Food
and Obesity/ non-communicable diseases Research, Monitoring and Action Support
(INFORMAS) that facilitate evaluating the current situation in European public policies
affecting the food policy environments. The INFORMAS approach comprises indicators such
230 as food provision, prices, promotion, labelling, retail and trade, leadership and governance
etc. We will further include indicators of the WHO Health-Enhancing physical Activity (HEPA)
Policy Audit Tool (PAT) which is a standardised tool designed to help researchers and policy
makers collect information on what policies exist across different sectors (e.g. organized
sport, school, transport, recreation) that directly or indirectly impact on PA (and sedentary
235 behaviour).

****INSERT TABLE 2 AND 3 HERE****

Development and evaluation of screeners and supplementary modules (AP3, AP4 and AP5)

Selection of a set of core variables (AP3)

Based on the selected and prioritised key indicators, a set of core variables (or proxies) will be identified that can be measured with a few screening questions (screeners).

To give an example, assessing fruits and vegetables intake can be measured as daily intake (piece/day) or as adherence to the 5-a-day recommendation (in %). Therefore the screener may include respective instruments to measure fruits and vegetables intake, e.g. a FFQ (as provided by EHIS), a 24HDR (GloboDiet), “How often do you eat fruit?” (HBSC survey 2013/2014) or “How often do you eat fruits and vegetables during a day, a week or a month?” (Nordic Monitoring System). Accordingly, the screeners will capture the most relevant aspects of health-related topics and allow ideally quantitative assessments; they should be self-sufficient for inclusion in different survey instruments and they should capture central components of key indicators; the selection of screeners for specific topics should be based on validity, reliability and the evidence regarding impact on health and health behaviour. A few studies such as HBSC already provide suitable screening tools for the collection of behavior- and health-related data that may be considered for the harmonization process (Brooks 2015).

Selection of instruments to assess key indicators (AP4)

Accepted and validated methodologies and definitions exist for a number of indicators, but the corresponding variables are not always comparable between systems. Dietary intake is mainly assessed using food frequency questionnaires (FFQ; without estimation of portion size), 24-h dietary recall (24HDR; ideally repeated recalls) (De Keyzer

et al. 2015) or food diary/ records (ideally repeated days recorded) (DIET@NET partnership
265 2017).

Multiple 24HDR (or dietary records of children) are considered as the least-biased
way to monitor food, nutrient and contaminant intake in populations. The traditional
interviewer- or self-administered 24HDR method is still the standardized assessment method
used in large-scale surveys, such as ASA24 (Automated Self-Administered 24-hour Recall;
270 <http://epi.grants.cancer.gov/asa24/>). In Europe, the interviewer-administered GloboDiet
24HDR software is used as a standardized system to assess dietary data in some European
countries. Also 24-hour urine voids are used to monitor certain components/ nutrients such
as sodium and iodine intake (biomonitoring surveys) (Brussaard et al. 2002).

Dietary behavior questionnaires have been validated less frequently compared to
275 dietary intake questionnaires, although questions are less complex. Dietary behavior (e.g.
breakfast consumption) may be measured using questionnaires or FFQs. Several short
FFQs focusing on a smaller number of key indicators have been developed for use in cross-
country studies such as WHO-COSI and HBSC. Validity of these FFQs was tested in both
HBSC and the Nordic Monitoring of Diet, Physical Activity and Overweight, while
280 reproducibility was only tested in HBSC (Bel-Serrat et al. 2017).

As FFQs create less respondent burden and are less resource intensive compared to
dietary records, 24HDR or biomarker use, FFQs have so far been a preferred method for
population-based surveys.

Official reports about PA levels in European children use data obtained through
285 questionnaires (Janssen et al. 2005). As questionnaires are imprecise in the assessment of
PA in children and adolescents and given the advances made in this area over the last
decades, objective methods should be favoured in large-scale studies or for surveillance
purposes (Basterfield et al. 2008). In contrast, various PA questionnaires have been used to
assess PA, but their validity is known to be limited (Helmerhorst et al. 2012). For example,
290 PA screeners used in HBSC have been shown to have acceptable psychometric properties

and validity (Health Behaviour in School-aged Children Study 2014). SB is most often assessed via questionnaires that mainly collect information on reading or screen-based behaviors, with some studies showing that these measures are reliable and valid (Bobakova et al. 2015). However, there is a lack of validated measures for total sitting time in children and adolescents.

Activity monitors and other new technological devices are promising objective methods for measuring PA and SB in surveillance surveys as they will facilitate the harmonization and integration of high quality measurements (Van Hecke et al. 2016; Verloigne et al. 2016). A major limitation that prevents the widespread use of activity monitors is the relatively high cost of the measurement devices as well as the laborious data management (Pedisic and Bauman 2015).. Furthermore, for the comparability of accelerometer data standardized proceedings in data collection and processing are essential (Pedisic and Bauman 2015). However, data management may become simpler and faster as the technology progresses and costs will presumably drop in the future as these devices are becoming the standard in current research. Additionally, suitable instruments (a) have to be valid and reliable for use in a cross-cultural context; (b) should overlap as much as possible with methods and instruments already used by current systems; (c) have to be easily applicable to provide robust estimates; and (d) should be affordable.

Selection of a set of supplementary modules (AP5)

The feasibility of replacing self-reports by objective measurements (e.g. activity monitors) and mHealth technology (World Health Organization 2011) should be evaluated for selected core variables. In parallel to the incorporation of screeners into existing surveillance systems (full sample), a set of supplementary objective measurement methods may be implemented (among sub-samples) in these systems, where feasible; this will enhance data quality and completeness. This may initially be accomplished in supplementary modules for smaller subgroups, while core variables may be used to assess temporal trends in the full

study population. Supplementary modules may provide quantitative data and could be used to validate self-reports or calibrate survey instruments. For dietary assessment, the European Food Safety Authority (EFSA) recommends dietary records for children aged 3-9 years and 24h-dietary recall interviews for children aged over 10 years, adolescents and adults for an application in subgroups (European Food Safety Authority 2014).

Objective measurements provide information on patterns and intensities, at least in subgroups; in combination with a PA diary, activity monitors allow the assessment of PA intensities and duration by domain (school-related, transport, leisure-time). To our knowledge, no pan-European surveillance initiative has used activity monitors so far but examples exist in national surveys such as the KiGGS study. In the second examination wave, KiGGS and the supplementary module Motorik-Modul (MoMo) used a physical activity questionnaire in addition to accelerometers (Actigraph GT3X+/wGT3X-BT) in a subsample. Accelerometer data sets for 6,720 respondents (11 to 29 years) have been collected. In the long run, KiGGS and MoMo will be able to investigate temporal trends across the total sample (Woll et al. 2017).

Implementation of screeners accompanied by methodological studies (AP6)

A set of screeners will be integrated into existing surveillance systems to provide comparable data between these systems for a limited number of variables only. Due to the strong selection criteria (validity, reliability and robustness of estimates in a cross-cultural context) the implementation of screeners facilitates quality assurance across the systems and over time. Screeners will be used to calibrate existing instruments with which they overlap across different surveillance systems. This calibration approach could be integrated in pilot studies (comparing data collected using new screening instruments with data from established instruments) and validation studies (comparing data collected using new screening instruments with data from established 'gold standard' methods, such as activity monitors for PA and SB). Since the screeners may be implemented by surveillance systems

without discarding existing instruments the system's internal integrity is secured and their ability to assess temporal trends is retained.

Then, in subsequent steps, more screeners or supplementary modules may be added and enable the gradual transition of existing surveillance systems towards harmonized and higher quality measures. Eventually, some of the original non-harmonized instruments may be replaced by new harmonized measurements without the loss of information on temporal trends if the calibration was successful.

Sustainable harmonization through a methodological competence platform (AP7)

The success of the harmonisation of current surveillance systems will depend on the willingness of existing systems to work together for the common aim of a pan-European system. Support may be greater if existing systems retain a certain degree of flexibility to provide data serving regional or national interests. Sustained support is needed for the selection, development, pretesting and validation of novel tools that are affordable, easy to use in different settings and that provide robust estimates. Existing surveillance systems may undertake a joint effort to establish a sustainable methodological competence platform for supporting and coordinating the necessary methodological developments. This most certainly will comprise information and communication technology, technical features and qualified personnel; the involvement or linkage with national and international surveillance systems will be desirable. Such a platform ought to offer instruments based on a modular system and enable the user to decide which tools they want to apply in their (national) surveillance system, e.g. from a basic questionnaire up to physical or even biochemical measurements. Such a methodological platform may facilitate (a) the development and validation of novel instruments and measurement modules; (b) the gradual replacement of the original measures with valid harmonized measurements; (c) the promotion of common standards for data management, data pooling, data access, ethics, training and quality management; and (d) the collaboration of surveillance systems.

In conclusion, the authors do not recommend the development of a novel pan-European surveillance system as several well-established systems are operational. Instead, we propose a stepwise harmonisation process building on these existing surveillance systems in order to facilitate increased comparability of data across surveys, age groups and countries. This approach would of course require additional resources to integrate these new measures into the existing surveillance systems, which might prove challenging as these systems are often financially stretched already.

The modular approach would allow to build a bridge between health reporting and research systems. This idea is already inherent to the proposed validation studies that would be conducted in subsamples. As already done in several surveillance systems such as National Health and Nutrition Examination Survey (NHANES) and KiGGS, longitudinal subsamples could become a fully integrated component to assess the causal relationship between behaviors, determinants and health outcomes and could further constitute a powerful pan-European cohort. Close concatenation with research may be a means to improve the quality and usefulness of the surveillance data and to optimise data use for the impact evaluation of health promoting policies.

Funding

This work is supported by the Joint Programming Initiative 'A Healthy Diet for a Healthy Life'. The funding agencies supporting this work are (in alphabetical order of participating country): Germany: Federal Ministry of Education and Research (BMBF); The Netherlands: The Netherlands Organisation for Health Research and Development (ZonMw); Norway: The Research Council of Norway, Division for Society and Health.

Availability of data and material

Not applicable

Authors' contributions

All authors were closely involved in the development of the conceptual framework and provided to the discussion section related to their field of expertise. All authors provided feedback and contributed to further versions. All authors read and approved the final manuscript.

Compliance with Ethical Standards

This article does not contain any studies with human participants or animals performed by any of the authors.

References

- Atkin AJ, Gorely T, Clemes SA et al. (2012) Methods of Measurement in epidemiology: sedentary Behaviour International journal of epidemiology 41:1460-1471 doi:10.1093/ije/dys118
- Basterfield L, Adamson AJ, Parkinson KN et al. (2008) Surveillance of physical activity in the UK is flawed: validation of the Health Survey for England Physical Activity Questionnaire Archives of disease in childhood 93:1054-1058 doi:10.1136/adc.2007.135905
- Bauman AE, Reis RS, Sallis JF et al. (2012) Correlates of physical activity: why are some people physically active and others not? Lancet 380:258-271 doi:10.1016/S0140-6736(12)60735-1
- Bel-Serrat S, Huybrechts I, Thumann BF et al. (2017) Inventory of surveillance systems assessing dietary, physical activity and sedentary behaviours in Europe: a DEDIPAC study Eur J Public Health 27:747-755 doi:10.1093/eurpub/ckx023
- Bobakova D, Hamrik Z, Badura P et al. (2015) Test-retest reliability of selected physical activity and sedentary behaviour HBSC items in the Czech Republic, Slovakia and Poland International Journal of Public Health 60:59-67
- Brooks F, Magnusson, J., Klemra, E., Chester, K., Spencer, N., and Smeeton, N. (2015) HBSC England National Report 2014. University of Hertfordshire, Hatfield, UK
- Brug J, van der Ploeg HP, Løyen A et al. (2017) Determinants of diet and physical activity (DEDIPAC): a summary of findings Int J Behav Nutr Phys Act 14:150 doi:10.1186/s12966-017-0609-5
- Brussaard JH, Lowik MR, Steingrimsdottir L et al. (2002) A European food consumption survey method--conclusions and recommendations Eur J Clin Nutr 56 Suppl 2:S89-94 doi:10.1038/sj.ejcn.1601432
- Buck C, Bornhorst C, Pohlabein H et al. (2013) Clustering of unhealthy food around German schools and its influence on dietary behavior in school children: a pilot study Int J Behav Nutr Phys Act 10:65
- Currie C, Hurrelmann K, Settertobulte W et al. (2000) Health and health behaviour among young people (Health Policy for Children and Adolescents, No.1). WHO Regional Office for Europe, Copenhagen
- De Keyser W, Bracke T, McNaughton SA et al. (2015) Cross-continental comparison of national food consumption survey methods--a narrative review Nutrients 7:3587-3620 doi:10.3390/nu7053587
- de Moraes AC, Carvalho HB, Siani A et al. (2015) Incidence of high blood pressure in children - effects of physical activity and sedentary behaviors: the IDEFICS study: High blood pressure, lifestyle and children International journal of cardiology 180:165-170 doi:10.1016/j.ijcard.2014.11.175

- DIET@NET partnership (2017) Nutritools. Medical Research Council. <https://www.nutritools.org/>. Accessed November 2017
- 440 European Food Safety Authority (2014) Guidance on the EU Menu methodology EFSA Journal 12:3944
doi:10.2903/j.efsa.2014.3944
- Expert Group on Health Information (EGHI) The European Core Health Indicators shortlist
http://ec.europa.eu/health/indicators/docs/echi_shortlist_by_policy_area_en.pdf. Accessed September 2016
- 445 Fernandez-Alvira JM, Bammann K, Eiben G et al. (2017) Prospective associations between dietary patterns and body composition changes in European children: the IDEFICS study Public Health Nutr:1-9
doi:10.1017/s1368980017002361
- Health Behaviour in School-aged Children Study (2014) Health Behaviour in School-aged Children (HBSC) Study Protocol: Background, Methodology and Mandatory items for the 2013/14 Survey. . St Andrews: CAHRU,
- 450 Heath GW, Parra DC, Sarmiento OL et al. (2012) Evidence-based intervention in physical activity: lessons from around the world Lancet 380:272-281 doi:10.1016/S0140-6736(12)60816-2
- Hebestreit A, Bornhorst C, Pala V et al. (2014) Dietary energy density in young children across Europe International Journal of Obesity 38:S124-S134 doi:10.1038/ijo.2014.143
- 455 Hebestreit A, Intemann T, Siani A et al. (2017) Dietary Patterns of European Children and Their Parents in Association with Family Food Environment: Results from the I.Family Study Nutrients 9
doi:10.3390/nu9020126
- Helmerhorst HJ, Brage S, Warren J et al. (2012) A systematic review of reliability and objective criterion-related validity of physical activity questionnaires Int J Behav Nutr Phys Act 9:103 doi:10.1186/1479-5868-9-103
- 460 Janssen I, Katzmarzyk PT, Boyce WF et al. (2005) Comparison of overweight and obesity prevalence in school-aged youth from 34 countries and their relationships with physical activity and dietary patterns Obesity reviews : an official journal of the International Association for the Study of Obesity 6:123-132
doi:10.1111/j.1467-789X.2005.00176.x
- 465 Kamtsiuris P, Lange M, Schaffrath Rosario A (2007) [The German Health Interview and Examination Survey for Children and Adolescents (KiGGS): sample design, response and nonresponse analysis] Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz 50:547-556 doi:10.1007/s00103-007-0215-9
- Malik VS, Pan A, Willett WC et al. (2013) Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis The American journal of clinical nutrition 98:1084-1102
doi:10.3945/ajcn.113.058362
- 470 McVeigh J, Smith A, Howie E et al. (2016) Trajectories of Television Watching from Childhood to Early Adulthood and Their Association with Body Composition and Mental Health Outcomes in Young Adults PloS one 11:e0152879 doi:10.1371/journal.pone.0152879
- 475 National Institute for Public Health and the Environment R (2012) ECHIM Echi Indicator Development and Documentation, Joint Action for ECHIM Final Report Part II. Bilthoven, The Netherlands
- Olafsdottir S, Berg C, Eiben G et al. (2014) Young children's screen activities, sweet drink consumption and anthropometry: results from a prospective European study Eur J Clin Nutr 68:223-228
- Pearson N, Biddle SJ, Gorely T (2009) Family correlates of breakfast consumption among children and adolescents. A systematic review Appetite 52:1-7
- 480 Pedisic Z, Bauman A (2015) Accelerometer-based measures in physical activity surveillance: current practices and issues Br J Sports Med 49:219-223 doi:10.1136/bjsports-2013-093407
- Poitras VJ, Gray CE, Borghese MM et al. (2016) Systematic review of the relationships between objectively measured physical activity and health indicators in school-aged children and youth Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme 41:S197-239
doi:10.1139/apnm-2015-0663
- 485 Rosenheck R (2008) Fast food consumption and increased caloric intake: a systematic review of a trajectory towards weight gain and obesity risk Obesity reviews : an official journal of the International Association for the Study of Obesity 9:535-547 doi:10.1111/j.1467-789X.2008.00477.x
- 490 Sleddens EF, Kroeze W, Kohl LF et al. (2015) Determinants of dietary behavior among youth: an umbrella review Int J Behav Nutr Phys Act 12:7 doi:10.1186/s12966-015-0164-x
- Stierlin AS, De Lepeleere S, Cardon G et al. (2015) A systematic review of determinants of sedentary behaviour in youth: a DEDIPAC-study Int J Behav Nutr Phys Act 12:133 doi:10.1186/s12966-015-0291-4
- Van Hecke L, Loyen A, Verloigne M et al. (2016) Variation in population levels of physical activity in European children and adolescents according to cross-European studies: a systematic literature review within DEDIPAC Int J Behav Nutr Phys Act 13:70 doi:10.1186/s12966-016-0396-4
- 495

- Verloigne M, Loyen A, Van Hecke L et al. (2016) Variation in population levels of sedentary time in European children and adolescents according to cross-European studies: a systematic literature review within DEDIPAC Int J Behav Nutr Phys Act 13:69 doi:10.1186/s12966-016-0395-5
- 500 Wang X, Ouyang Y, Liu J et al. (2014) Fruit and vegetable consumption and mortality from all causes, cardiovascular disease, and cancer: systematic review and dose-response meta-analysis of prospective cohort studies BMJ (Clinical research ed) 349:g4490 doi:10.1136/bmj.g4490
- Woll A, Albrecht C, Worth A (2017) Monitoring Motorik-Module (MoMo) – the KiGGS Wave 2 module to survey motor performance and physical activity Journal of Health Monitoring 2 doi:DOI
- 505 10.17886/RKI-GBE-2017-110
- World Health Organization - Europe (2014) WHO European Childhood Obesity Surveillance Initiative (COSI). <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/activities/monitoring-and-surveillance/who-european-childhood-obesity-surveillance-initiative-cosi>. Accessed November 2016
- 510 World Health Organization (2011) mHealth: New horizons for health through mobile technologies: second global survey on eHealth, Global Observatory for eHealth series - Volume 3 edn. World Health Organization,
- World Health Organization (2014) The European child and adolescent health strategy 2015–2020. Copenhagen, Denmark
- 515 World Health Organization (2016) Report of the commission on: ending childhood obesity. http://apps.who.int/iris/bitstream/10665/204176/1/9789241510066_eng.pdf?ua=1. Accessed June 2016
- World Health Organization Europe (2013) Vienna Declaration on Nutrition and Noncommunicable Diseases in the Context of Health 2020.

Table 1: Main pan-EU surveillance systems (adapted from (Bel-Serrat et al. 2017))

Survey system	Year of 1 st wave	No. of completed waves	No. of countries 1 st wave	No. of countries last/current wave	Target population	Sample size (year of wave)
Nordic Monitoring System	2011	2	5	5	7-12 years and 18-65 years	2,479 children; 9,153 adults (2011)
Healthy Behavior in School Children	1983/84	9	5	47	11, 13, 15 years	219,460 (2013/14)
WHO European Childhood Obesity Surveillance Initiative	2007/08	4	13	36	6-9 years	256,157 (2012/13)
European Health Interview Survey	2006-2009	2	17	30	> 15 years	210,000 (2014)
EU Menu	2011	6	2	16	3 months - 74 years	≥130 valid subjects per Sex and age class

Table 2: Health determinant indicators (and related ECHI numeration) of the European Core Health Indicators (Expert Group on Health Information (EGHI))

Health determinants indicators of the European Core Health Indicators	
Proximal health determinants	Distal health determinants
42. Body mass index	47. Hazardous alcohol consumption
43. Blood pressure	48. Use of illicit drugs
49. Consumption of fruit	53. Work-related health risks
50. Consumption of vegetables	54. Social support
52. Physical activity	55. Particulate matter exposure
46. Total alcohol consumption	
44. Regular smokers	
45. Pregnant women smoking	

Table 3: Priority list of indicators of the European Core Health Indicators (Expert Group on Health Information (EGHI))

Indicators of dietary intake	Indicators of physical activity	Indicators of sedentary behaviour
<ul style="list-style-type: none"> • Consumption of vegetables • Consumption of fruits • Consumption of soft drinks • Alcohol intake 	<ul style="list-style-type: none"> • Physical activity frequency, duration and intensity • Leisure time physical activity • Transport physical activity • Organised physical activity 	<ul style="list-style-type: none"> • Media use • Screen time • Computer use • Television time