Mitophagy and Neuroprotection

Guofeng Lou^{1,6}, Konstantinos Palikaras^{2,3,6}, Sofie Lautrup^{1,6}, Morten Scheibye-Knudsen⁴, Nektarios Tavernarakis^{2,3}, Evandro F. Fang^{1,5*}

¹Department of Clinical Molecular Biology, University of Oslo and Akershus University Hospital, 1478 Lørenskog, Norway

²Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology-Hellas, Greece

³Department of Basic Sciences, Faculty of Medicine, University of Crete, Heraklion, 70013, Crete, Greece

⁴Center for Healthy Aging, Department of Cellular and Molecular Medicine, University of Copenhagen, Copenhagen, Denmark.

⁵The Norwegian Centre on Healthy Ageing (NO-Age), Norway

⁶Co-first authors

*Correspondence:

e.f.fang@medisin.uio.no (E.F. Fang)

Abstract

Neurodegenerative diseases are strongly age-related and currently cannot be cured, with a surge of patient numbers in the coming decades in view of the emerging worldwide ageing population, bringing healthcare and socio-economic challenges. Effective therapies are urgently needed, and are dependent on new etiological mechanisms. In neurons, efficient clearance of damaged mitochondria, through the highly evolutionary conserved cellular process termed mitophagy, plays a fundamental role in mitochondrial and metabolic homeostasis, energy supply, neuronal survival and health. Conversely, defective mitophagy leads to accumulation of damaged mitochondria and cellular dysfunction, contributing to ageing and age-predisposed neurodegeneration. Here, we discuss the contribution of defective mitophagy in these diseases, and underlying molecular mechanisms, and highlight novel therapeutics based on new discovered mitophagy-inducing strategies.

Key words: ageing, Alzheimer's disease, amyotrophic lateral sclerosis, energy homeostasis, mitochondria, mitophagy, neurodegeneration, Parkinson's disease

Mitochondrial Maintenance Through Mitophagy: More Than the Homeostasis of Cellular Powerhouses and Metabolism

Mitochondria produce ATP and are indispensable for life and health in most eukaryotic organisms [1-5]. Decades of studies on cell fates (**apoptosis, necrosis, and autophagy**) (see Glossary) and diseases have greatly expanded our understanding of mitochondria as multi-faceted organelles displaying critical roles in metabolic homeostasis, regulation of Ca^{2+} and redox signalling, mitochondria-nuclear communications, and the arbitration of cell survival/stress resistance and death[4, 6, 7]. Mitochondrial dysfunction has been associated with a broad variety of human diseases. While mutations of nuclear or mitochondria-encoded mitochondrial proteins cause rare mitochondrial disorders[3], mitochondria-mediated ATP deprivation, oxidative stress and impaired cell signalling have been linked to the pathogenesis of prominent metabolic and neurodegenerative diseases (e.g., Alzheimer's disease/AD, Parkinson's disease/PD, and amyotrophic lateral sclerosis/ALS) [8-12] (see Clinician's Corner).

Cells have developed sophisticated, inter-connected regulatory pathways to maintain mitochondrial homeostasis through balanced mitochondrial biogenesis and clearance of damaged mitochondria. **Mitophagy**, a form of autophagy, mediates the removal of defective/superfluous mitochondria, and is the only known pathway through which whole mitochondria can be selectively removed (**Fig. 1**). Mitophagy is responsible for the basal mitochondrial turnover that eliminates dysfunctional mitochondria; however, mitophagy can also be induced under certain physiological conditions whereby healthy mitochondria are degraded. For example, during the maturation of erythrocytes, where mitochondria and other organelles are removed from the cell, and in the development of fertilized oocytes, when paternal mitochondria are eliminated [13, 14]. Additionally, mitophagy can be induced as a stress-response mechanism to inhibit mitochondria-dependent apoptosis. Current known mitophagy pathways are summarized here (**Fig. 2**) as well as in recent reviews [7, 9, 15].

In this review, we summarize the very recent progress of mitophagy in neuroprotection, covering topics from the changes of mitophagy in normal ageing and accelerated ageing conditions, as well as in the common neurodegenerative disorders, discuss

mitophagy induction as a therapeutic strategy as well as bring opinion on outstanding questions and future perspectives in this field.

Mitophagy and Healthy Longevity

Mitophagy and the Ageing Process

While ageing is the primary driver of the sporadic cases (which are dominant compared to familial ones) of neurodegenerative diseases, the underlying molecular mechanisms are elusive. Mitochondrial defects are accompanied by a plethora of age-related pathological conditions [3, 7, 9]. Impaired energy metabolism, which is characterized by accelerated mtDNA mutation rate, reduced electron transport chain (ETC) function, elevated reactive oxygen species (ROS) levels, defective cytoplasmic calcium buffering and enhanced release of pro-apoptotic factors, are a hallmark of ageing [16, 17].

During the last decades, multiple lines of experimental evidence in several model organisms, including yeast, flies, nematodes and mice, highlight the vital contribution of diminished mitochondrial activity in age-associated disorders [4, 7, 18]. Uncontrolled accumulation of dysfunctional organelles has been demonstrated in several types of cells and tissues during ageing [2, 19]. Cellular inability to discard impaired organelles, potentially due to the age-dependent decline of mitophagy or severe defects in macroautophagy, exacerbates energy homeostasis collapse through the accumulation of damaged mitochondrial population [20]. A recent study demonstrates that compromised mitochondrial function promotes a senescence response both in murine and human cells highlighting its association with the development and progression of ageing phenotypes [21]. Taken together, mitophagy might be a “gate-keeper” of energy metabolism sustaining cellular function, tissue physiology and organismal healthspan.

Several conserved longevity pathways, including low insulin/IGF-1 signaling, caloric restriction and moderate levels of mitochondrial dysfunction among others, converge on mitophagy stimulation to regulate lifespan. In *C. elegans*, mitophagy is induced in response to stress conditions eliminating damaged mitochondria and thereby promoting cellular homeostasis and viability [19]. Mitophagy deficiency mediates runaway accrual of dysfunctional organelles resulting in altered mitochondrial metabolism and increased susceptibility to stress and infection [18, 19, 22].

Experimental evidence supporting the vital role of mitophagy in the maintenance of energy metabolism comes from recent studies showing that mitochondrial removal is declined with age in mice and human cells [23, 24]. Interestingly, hippocampal neurons present approximately 70% reduction of mitophagic events in old mice. These results signify the crucial role of mitophagy in this particular brain region that is especially vulnerable to deterioration at early stages of AD. However, we should be extremely judicious in data extrapolation based on these experiments since these mt-Keima mice have early neurodegeneration and impaired circadian biology, compared with commonly used C57BL6/J mice [25]. Recent proof-of-concept studies demonstrate that mitophagy is blocked both in Tau and amyloid- β (A β)-based *C. elegans* and mouse models of AD, as well as in postmortem hippocampal brain tissues from AD patients, highlighting mitophagy defects as a common feature of AD pathogenesis [10, 26].

Taken together, mitochondrial metabolism and mitophagy hold a pivotal role in the maintenance of organismal homeostasis and ageing. Alterations in mitophagy balance result in the accumulation of defective mitochondria that contribute to the development and progression of age-associated pathologies [7]. Thus, the delineation of the molecular underpinnings and the identification of genetic and/or pharmacological interventions that influence mitophagy and mitochondrial function could be beneficial for cellular homeostasis and organismal healthspan.

Impaired Mitochondrial Clearance in Premature Aging Diseases With Neurodegeneration

Premature ageing diseases are extremely rare autosomal dominant genetic disorders in which clinical syndromes are manifested at a very early age resembling aspects of ageing [18, 27, 28]. Mutations of genes involved in DNA repair are the primary cause of premature ageing diseases, such as the mutations of genes encoded for xeroderma pigmentosum (XP) proteins, ataxia telangiectasia (AT)-mutated (ATM) and cockayne syndrome group A (CSA) or B (CSB) [18, 27, 28]. Recent studies indicate that mitochondrial dysfunction is also a major contributor of premature ageing in these pathological conditions. In XP group A (XPA), AT, and CSB, accumulation of nuclear DNA damage drives depletion of intracellular **NAD⁺** levels leading to the impairment of mitophagy and to subsequent accrual of defective mitochondria via reduction of the NAD⁺/sirtuins pathway [18, 28, 29]. Defective mitophagy was also reported in other

DNA repair-deficient-related diseases, such as Fanconi anaemia (FA) [30]. These data strongly suggest a role of 'nuclear DNA damage signalling to mitochondria' (N-M signalling) in mitophagy [4], and highlight an additive/synergistic role with genetic-oriented nuclear DNA damage to promote premature ageing, as well as related neurodegeneration phenotypes.

Defective Mitophagy in Neurodegenerative Diseases

Mitochondria are the major producers of ATP that is essential for the excitability and survival of neurons. Mitochondrial signalling pathways regulate synaptic signalling and related long-term modifications in neuronal structure and function [6, 9]. Thus, mitochondrial dysfunction as well as inefficient clearance of impaired mitochondria are detrimental to neurons and normal brain function [6, 9].

Defective Mitophagy in Alzheimer's disease

AD is the most common neurodegenerative disorder, resulting in severe memory loss and cognitive dysfunction. The principal pathological hallmarks of AD are intracellular neurofibrillary tangles (NFTs), mostly constituted by hyper-phosphorylated Tau protein (pTau), and extracellular A β plaques that derive from proteolytic processing of amyloid precursor protein (APP) [9, 10, 31]. Genetic and animal studies suggest that impairment of the autophagic/lysosomal/endosomal system contributes to accumulation of Tau- and A β -based **proteinopathies** [32, 33]. In addition to pTau and A β , mitochondrial dysfunction with accumulation of defective organelles are common features in brain tissues from AD patients as well as in experimental AD mouse models [9, 10, 34, 35]. Mounting evidence argues that alterations in mitochondrial dynamics, morphology, motility, and activity are linked to AD, and together with oxidative stress seem to be important early events in disease development [9, 10, 34, 35]. Since mitochondrial dysfunction is a hallmark of AD, and neuronal autophagy is also impaired in affected neurons, we have suggested a 'defective mitophagy in AD' theory [9]. Indeed, the basal level of mitophagy is dramatically reduced in post-mortem hippocampal tissues from AD patients, AD induced pluripotent stem cell (iPSC)-derived cortical neurons, as well as in the APP/PS1 and the 3xTg AD mouse models[10]. At the molecular level, the mitophagy machinery is hampered, evidenced by impaired activation of two autophagic/mitophagic initiator machinery proteins; ULK1 (low phosphorylated Serine 555 site) and TBK1 (low phosphorylated Serine 172 site)

[10]. Furthermore, lower basal as well as an impaired mitochondrial stress-induced mitophagy is evident in pan-neuronal pTau- or A β -expressing nematodes [10], indicating that Tau- and A β -based proteinopathies can affect the efficiency of mitophagic machinery. Possible additional mechanisms of defective mitophagy (accumulation of damaged mitochondria) in AD include: impaired Parkin-presenilin 1 (PS1)/ γ -secretase- Amyloid precursor protein intracellular domain (AICD)-PINK1 transcription axis [36], β -secretase-derived APP fragment/CTF β (C99)-induced endosomal-autophagic-lysosomal dysfunction [37], dysfunctional over-sized lysosomes[38], defective retrograde transport of damaged mitochondria to the soma for mitophagic degradation [39], and impaired mitochondrial stress response [40].

It remains to be determined whether the presence of impaired mitophagy is a consequence of Tau/A β proteinopathies, or whether inefficient mitophagy is an early event preceding and causing Tau/A β proteinopathies. Data from interventional studies, where mitophagy was reinstalled pharmacologically, favour the latter [10]. There are limitations to the use of transgenic Tau/A β animal models to address this puzzle. In AD mouse models, Tau/A β is already artificially overexpressed from the day of birth, a time-point several months ahead of disease pathologies. Studies using sporadic AD iPSC-derived neurons or other similar models are necessary to further dissect a confounding question: whether defective mitophagy plays a causative role or serving as a compensatory response in Tau/A β proteinopathies.

Defective mitophagy And Imbalanced Mitochondria-Derived Vesicles (Mdvs) Formation in Parkinson's disease

The symptoms of PD are caused by a loss of dopaminergic neurons in the substantia nigra pars compacta (SNpc) resulting in a decrease of dopamine (DA) levels in the striatum. Mitochondrial dysfunction and its related oxidative stress and inflammation are common features in PD patient brain samples, PD iPSC-derived neurons, and/or PD animal models [11, 12, 41, 42]. To note, pathological studies in the substantia nigra regions from 6 patients suggest that Complex I deficiency did not correlate with parkinsonism [43]. At the molecular level, mutations of the *PARK6* (encoding PINK1) and *PARK2* (encoding Parkin) genes cause <5% of PD cases (familial PD) [44, 45]. In addition to the well-known function in mitophagy, PINK1 and Parkin repress mitochondrial antigen presentation (MitAP). Experimental studies point to a significant

contribution of a hyper-activated immune system, provoked by inflammation, in PD progression [42]. A recent study shows that PINK1 and Parkin repress MitAP through inhibition of Sorting nexin 9 (Snx9)-dependent formation of mitochondria-derived vesicles (MDVs) [46].

Although extensive mechanistic studies of the PINK1/Parkin pathway have been performed, the role of PINK1/Parkin-dependent mitophagy *in vivo* remains elusive. Both *Prkn*^{-/-} and *Pink1*^{-/-} mice show no substantial PD-relevant phenotypes [47, 48]. In addition, basal mitophagy is comparable between PINK1 wild-type and *Pink1*^{-/-} mice [49]. This might be due to sufficient compensation of the loss of PINK1-dependent mitophagy by other PINK1-independent pathways under physiological conditions. To note, *PINK1* deletion in rhesus monkeys resulted in neurodegeneration; however, this phenotype was not associated with altered mitochondrial morphology in the neurons, suggesting the existence of none-mitophagy-regulatory role(s) of PINK1 [50]. PD patients show strong inflammatory phenotypes, with elevated levels of IL-1 β , IL-6, TNF- α , and IFN- γ [51]. Interestingly, a recent study showed that under acute (exhaustive exercise) or chronic (mutation of DNA Polymerase *PoIG*, named *mutator*) mitochondrial stress conditions, knock down of *Prkn* or *Pink1* in mice lead to an exacerbation of inflammation status via a **STING** dependent manner [51]. Additional studies link mitochondrial dysfunction and inflammation. Accumulation of damaged mitochondria can trigger inflammation via the **NLRP3 inflammasome** and/or the cGAS (a DNA sensor, also known as MB21D1)-cGAMP-STNG-TBK1-IRF3 pathway, linked to various human diseases, including pathological ageing and neurodegeneration [10, 51-53]. In cells with mitochondrial inner membrane permeabilization, mtDNA release induces activation of the cGAS-STING signalling [53]. Strikingly, loss of STING has been shown to dramatically rescue inflammation in both *Pink1*^{-/-}, *Prkn*^{-/-} and *Prkn*^{-/-}/*mutator* mouse models, and it even inhibited loss of dopaminergic neurons and motor defects observed in aged *Prkn*^{-/-}/*mutator* mice [51]. To note, the *in vivo* mitophagy levels were performed in the heart, but not in the dopaminergic neurons in the *Pink1*^{-/-} mt-Keima mice [51]. In summary, these *in vivo* data suggest that the PINK1/Parkin-dependent mitophagy may be dispensable at physiological conditions but is indeed necessary at stress/pathological conditions for the function and survival of PD-related dopaminergic neurons. More information between defective mitophagy and PD has been elegantly summarized in a recent review [54].

Emerging evidence from multiple studies suggests that the story is far more complex than previously appreciated with respect to PINK1 mutation, mitophagy status and PD[49, 50, 53, 55] as PINK1 and Parkin may have potential mitophagy-independent functions[50, 56]. Further understandings of PD aetiologies through the investigation of different mitophagy pathways, other non-canonical mitochondrial clearance pathways, and mitochondria-unrelated pathways are necessary.

Defective mitophagy in ALS and other neurodegenerative diseases

From the studies presented above, proteinopathies with mitochondrial dysfunction seem to be common features of neurodegenerative diseases. In addition to AD and PD, other neurodegenerative disorders, such as ALS, frontotemporal dementia (FTD) and Huntington's disease (HD) share similar features.

ALS is a fatal disease characterized by the selective degeneration of motor neurons. A hallmark of the disease, as in other neurodegenerative pathologies, is the accumulation of aggregated proteins within the affected neurons. A study using HeLa cells with protein overexpression suggests that inefficient turnover and aggregates of damaged mitochondria, may contribute to disease progression in ALS [57]. FTD affects cortical neurons and the basal ganglia, resulting in cognitive impairment, language deficiency and changes in social behaviour and conduct. Despite the differences in the types of neurons affected, both diseases show clinical and genetic overlap [58]. Many of the genes linked to ALS and FTD encode proteins involved in mitophagy/selective autophagy, including: OPTN, TBK1, p62, and receptor-interacting protein kinase 1 (PIPK1), among others [57-60], but how mutation of these genes contribute to the pathology are not fully understood.

HD is an autosomal-dominant neurodegenerative disorder caused by an expansion of the numbers of cytosine-adenine-guanine (CAG) trinucleotide repeats encoding a polyglutamine (polyQ) tract in the amino-terminal region of Huntingtin (Htt) protein. Htt is a large protein that can interact with and probably impair numerous other proteins. Interestingly, mutant Htt has also been proposed to affect the endosomal/autophagic system proteins[61]. Mitochondrial dysfunction and proteostasis alterations, particularly in the autophagic/endosomal system, including the mutation of *ATG7* [32],

impairment of GAPDH-mediated mitophagy by mutant Htt [62] are linked to the pathogenesis of HD. In summary, while evidence of impaired autophagy and pathological mitophagy in HD, determination of mitophagy status (upregulated/defective) in HD and more detailed molecular mechanisms are necessary.

Summarizing, it is likely that impaired mitophagy is a common phenomenon in many neurodegenerative pathologies (**Table 1**). To note, this opinion is based on the data dominantly from cell culture and laboratory animal models, clear cut post-mortem evidence from patients that demonstrates dysregulated mitophagy triggers multiple forms of neurodegeneration is scarce. In view of the high instability and diversity of mitophagy status in the human brain tissues, it is essential to combine large cohort of human brain samples obtained under tightly controlled conditions and state-of-the-art tissue processing methods, as recently performed [63], to further address this question.

Targeting Clearance of Damaged Mitochondria as Therapeutic Strategies

Maintenance of a healthy mitochondrial pool may serve as a robust approach to maintain healthy ageing and high quality of life at vulnerable old age. Since mitochondrial damage is a hallmark of several neurological pathologies, chemical substances that enhance the clearance of defective mitochondria are likely to have significant therapeutic benefit [6, 9, 17]. For diseases with genetic mutations, the best candidates for therapy remain small molecule drugs. Several robust mitophagy inducers have been discovered with promising benefits to the extension of healthspan and the protection of neurons in animal models and/or human cells. Among them are NAD⁺ precursors, urolithin A (UA), actinonin, spermidine [64, 65] and the FDA-approved drugs, rapamycin and metformin [66] (**Table 2**).

NAD⁺ precursors as mitophagy inducers

Major NAD⁺ precursors include nicotinamide, nicotinamide mononucleotide (NMN) and nicotinamide riboside (NR). Because of low NAD⁺ levels in tissues from laboratory AD animal models, NAD⁺ replenishment through supplementation with NAD⁺ precursors can inhibit A β and Tau pathology and reverse cognitive deficits in 3xTgAD and APP/PS1 models of AD via the NAD⁺-dependent SIRT1 and SIRT3, increased activities of PI3K-AKT, MAPK/ERK1/2, and increased expression of CREB [10, 67-70].

In addition to an intriguing benefit to AD, NAD⁺ replenishment also rescued mitochondrial defects and dopaminergic neuronal loss in iPSC and fly models of PD [41]. A more compressive summary of multi-faceted pathways between NAD⁺ and mitophagy is in a recent review [71]. Several 1-3 months clinical trials indicate that the NAD⁺ precursor NR is orally bioavailable and safe (1g to 2 g/day/adult) [72, 73], and it is important to see the benefits of NAD⁺ replenishment in clinical trials for AD patients.

Urolithin A as a mitophagy inducer

UA, an ellagitannins-derived metabolite, has been shown to induce muscular mitophagy in cultured cells and *C. elegans* [74], and neuronal mitophagy in both *C. elegans* and the mouse brain[10]. Indeed, UA and the antibiotic actinonin (AC) [23] were shown to trigger mitophagy in AD mouse and *C. elegans* models in a PINK1/Parkin/NIX-dependent manner [10]. Treatment with either compound lead to the amelioration of several AD pathological features including memory and learning deficits, A β -/Tau-related aggregation and neuroinflammation [10]. UA-induced mitophagy protects against neuroinflammation by decreasing the generation of pro-inflammatory cytokines, such as TNF- α and IL-6, and enhancing the levels of the anti-inflammatory IL-10 [10]. A recent study demonstrates that IL-10 triggers macroautophagy/mitophagy by diminishing **the mammalian target of rapamycin (mTOR)** through **AMPK** induction, inhibiting inflammasome activation and unresolved inflammation [75]. Results from a very recent stage I clinical trial of UA on 60 healthy elderly individuals, indicated that oral administration of UA (up to 1,000 mg/day) over a 4-week period was safe and was able to induce plasma acylcarnitines and skeletal muscle mitochondrial gene expression (clinicaltrials.gov NCT02655393) [76].

Spermidine as a mitophagy inducer

Polyamines, including spermidine, decrease with age in the human brain and animal models. Treatment with spermidine improves memory impairments and has life-prolonging effects in flies, nematodes, and mice [64, 77]. These beneficial effects depend on the expression of autophagy-related genes, hence the activity of autophagy [65]. In addition to autophagy, spermidine stimulates mitophagy in cultured cell lines, including human fibroblasts and cardiomyocytes among others [78]. Spermidine induces mitophagy by inhibition of mTOR and activation of AMPK [65, 79]. Additionally, the ATM-dependent PINK1/Parkin signalling also can be activated by spermidine[80].

However, the exact underlining mechanisms between spermidine and mitophagy activation remain elusive.

Rapamycin and metformin

Rapamycin and metformin are two FDA-approved mTOR inhibitors, which exhibit both anti-cancer and anti-ageing properties [81]. Rapamycin prevents mTOR by direct binding to and inhibition of mTOR complex 1 (mTORC1) [82]. Furthermore, rapamycin supplementation promotes the activation of AMPK in several model organisms, including flies, nematodes, and mice [66]. In turn, AMPK targets and phosphorylates mTOR leading to its subsequent inhibition. Similar to rapamycin, metformin, the most widely supplemented anti-diabetic drug, is shown to induce mitochondrial removal. Metformin inhibits mTOR indirectly via multiple signalling pathways that are poorly understood [83, 84]. The primary mode of action of metformin may include the inhibition of mitochondrial complex I, activation of SIRT1, AMPK, and Parkin-mediated mitophagy [84, 85]. Table 2 summarizes the health benefits, including neuroprotection, of different mitophagy-inducing compounds.

Concluding Remarks

Efficient mitochondrial quality control is pivotal for cellular and organismal homeostasis. Despite the extensive studies focusing on the molecular mechanisms that govern mitophagy, some outstanding questions remain. Firstly, it is necessary to further study the interconnected interactions between the different mitophagy pathways and how they coordinate the regulation of mitochondrial removal. Secondly, the balance between mitophagy, mitochondrial biogenesis, and the mitochondrial fusion-fission processes should be examined. Thirdly, untangling of the multi-faceted roles of mitophagy in cell survival and death is needed. Mitophagy stimulation might be both beneficial and detrimental for tissue homeostasis depending on cellular bioenergetics under certain pathophysiological conditions. Furthermore, status of mitophagy as well as its linkages to neurodegenerative diseases in large cohort of high quality post-mortem human brain tissues from different neurodegenerative disease cases needs to be carefully determined. We should be extremely judicious in data extrapolation. Lastly, studies investigating the clinical benefits of known mitophagy inducers and to further develop specific, robust, but low toxic mitophagy inducers are lacking.

Emerging evidence of the causative linkage between defective mitophagy in AD, PD, and other neurodegenerative pathologies, propels drug development targeting damaged mitochondria, mitophagy, and other mitochondrial elimination pathways. Significant progress on using mitophagy inducers (e.g., supplementation with NAD⁺ precursors) to inhibit AD/PD has been made in both *in vivo* animal studies and patient iPSC-based systems, and further clinical studies are in progress. Small compound-based mitophagy induction may also benefit healthy ageing and the reduction of neurodegeneration. Furthermore, trends of a close collaboration between academia and industry as well as the application of artificial intelligence in mechanistic studies and drug development may accelerate the efficient development of transformative molecules targeting on these currently incurable neurodegenerative diseases [86]. Importantly, life-style interventions, such as physical exercise and caloric restriction [87], represent promising candidates to counteract deterioration of mitochondrial metabolism and prevent neurodegeneration. We urge large population-based, randomized, double-blind, placebo-controlled clinical trials for promising mitophagy-inducing molecules on these neurodegenerative diseases.

Acknowledgements

The authors acknowledge the valuable work of the many investigators whose published articles they were unable to cite owing to space limitations. E.F.F. is supported by the HELSE Sør-ØST (#2017056), the Research Council of Norway (#262175 and #277813), and an Akershus University Hospital Strategic grant (#269901). K.P. is supported by an AXA Research Fund post-doctoral long-term fellowship and the Hellenic Foundation for Research and Innovation (HFRI) and the General Secretariat for Research and Technology (GSRT). N.T. is funded by grants from the European Research Council (ERC – GA695190 – MANNA), the European Commission Framework Programmes, and the Greek Ministry of Education.

Disclaimer Statement

E.F.F. has CRADA arrangement with ChromaDex and is a consultant for Aladdin Healthcare Technologies and Vancouver Dementia Prevention Centre.

416

417

Glossary

5' adenosine monophosphate-activated protein kinase (AMPK): An evolutionary conserved and fundamental metabolic regulator that senses changes in the intracellular energy. In response to energy deprivation, AMP, ADP, and the newly discovered fructose-1,6-bisphosphate (FBP), and aldolase activate AMPK for glucose and fatty acid uptake and oxidation.

Apoptosis: the process of programmed cell death occurring in multicellular organisms, with distinct morphological characteristics and energy-dependent biochemical mechanisms. Classical features of apoptotic cells include cell shrinkage, blebbing, and nuclear fragmentation.

Autophagy: Autophagy is a cellular self-clearance system which delivers unnecessary/damaged cytoplasmic substrates to lysosomes for degradation. Autophagy can be further classified into macroautophagy, microautophagy, and chaperon-mediated autophagy. Autophagy plays a fundamental role in lifespan and health, including neuronal development and survival.

Mammalian target of rapamycin (mTOR): is a kinase in the phosphatidylinositol 3-kinase-related kinase family, regulating a broad spectrum of cellular processes. Through the binding of two distinct protein complexes, mTOR can be further classified as mTOR complex 1 and mTOR complex 2. The major functions of mTOR are the regulations of growth, proliferation and motility of cells and body. mTOR inhibition activates autophagy.

Mitochondria-derived vesicles (MDV): While cells initiate mitophagy to eliminate severely damaged whole mitochondria, cells can also remove damaged cargos from a mildly defective mitochondrion, through the generation of MDVs.

Mitophagy: A sub-type of macro-autophagy by which the double membraned cup-shaped phagophore recognizes damaged mitochondria for lysosomal degradation. The first and most well-defined mitophagy pathway is the PINK1/Parkin-dependent pathway whereas mutations of the genes *PINK1* and *Parkin* link to PD.

Nicotinamide adenine dinucleotide (NAD⁺): it is an essential molecule (co-factor) for numerous proteins involved in cellular energy metabolism, healthspan and longevity. NAD⁺ depletion is a fundamental feature of ageing that may lead to broad chronic diseases, including neurodegenerative disorders.

Necrosis: is an acute cell injury resulting in both regulated and unregulated cell death. Classical features of necrosis include DNA damage and degradation, nucleus shrinks, plasma membrane rupture, and inflammatory responses.

NLRP3: with the full name of 'nucleotide-binding domain, leucine-rich-containing family, pyrin domain-containing-3' or 'Nod-like receptor protein 3'. It is an inflammasome involved in a wide range of autoimmune diseases, auto-inflammatory diseases, and neurodegenerative diseases, as it can sense inflammatory proteins and aggregated proteins, including A β .

Proteinopathies: A type of pathological feature with intracytoplasmic protein misfolding and aggregation. They are common characteristics of many late-onset neurodegenerative diseases, including AD, PD, and HD.

Stimulator of interferon genes (STING): also known as TMEM173. Activated by intracellular/extracellular pathogens, STING participates in innate immunity through the induction of type I interferon. Studies using PD animal models indicate STING activation is a driver of PD.

Text Boxes

Clinician's Corner

Dementia affects over 50 million individuals worldwide, with 60% of them are patients with AD. While there is still no conclusive link between Tau/A β proteinopathies and cognitive symptoms, it is generally believed that AD is caused by a mix of multiple factors, including ageing, genetic, environmental, and other non-genetic factors.

There are over 15 clinical trials for the NAD⁺ precursors NR and NMN on different diseases, including biological ageing, obesity, and neurodegenerative diseases. Several finished independent clinical studies show that NR is orally bioavailable and safe at a dose of 1-2 g/day/adult for 1-3 months.

Defective mitophagy is a likely a common feature of broad neurodegenerative diseases, at least based on the data from animal models of AD and PD and ALS. To

note, clear cut post-mortem evidence from patients that demonstrates dysregulated mitophagy actually triggers multiple forms of neurodegeneration is scarce. Large clinical trials on such drug candidates are necessary.

Figure Legends

Figure 1. An overview of the mitophagy pathway.

Mitophagy comprises initiation, membrane nucleation and phagophore formation, phagophore expansion, fusion with the lysosome and finally degradation. Under different mitophagy-induction conditions which normally activate AMPK and/or inhibit mTOR, resulting in the activation of ULK1 and following formation of the ULK1 initiation complex. Activated ULK1 phosphorylates Beclin1 (the mammalian orthologue of yeast Atg6) and activates VPS34 lipid kinase (a class III phosphatidylinositol 3-kinase/PI3K), leading to the formation of the PI3K III nucleation complex. AMBRA1 may also participate in these processes, including stabilization of ULK1 and binding to the PI3KIII complex. The PI3K III nucleation complex induces the synthesis of phosphatidylinositol-3-phosphate (PI3P) which creates platforms that serve to recruit specific effectors for membrane trafficking events. Typical PI3P-binding complex comprises zinc-finger FYVE domain-containing protein 1 (DFCP1) and WD repeat domain phosphoinositide-interacting proteins (WIPIs). The ATG12 conjugation system ('CS') then binds on DFCP1 and WIPIs. Together with the ATG12 'CS' and the LC3/GABARAPs conjugation system (only LC3 is shown for simplicity, LC3 'CS'), these processes enable the formation and elongation of the double-membrane phagophore. The ATG5-ATG12-ATG16L1 complex promotes conversion of LC3 to LC3-II, with is then conjugated with phosphatidylethanolamine (PE) to form LC3-II (similar to other Atg8 members). LC3-II is incorporated into the phagophore/autophagosome which enables autophagic recognition and engulfment of damaged mitochondria through binding on LC3-interacting motifs (LIRs) in the mitophagy receptors. Current known mitophagy receptors are listed. The mature mitophagosome can then fuse with the acidic lysosome through at least two known mechanisms as noted. Conditions which affect lysosome function are summarized.

Figure 2. A summary of different mitophagy pathways.

A. The PINK1-dependent and Parkin-(in)dependent mitophagy pathways. From upper clockwise: **(1):** For the PINK1- and Parkin-dependent mitophagy pathway, reduced mitochondrial membrane potential enables the stabilization of full-length PINK1 at OMM. Full-length auto-phosphorylated PINK1 phosphorylates ubiquitin, enabling the binding of Parkin to phospho-ubiquitin and the recruitment and binding to OMM receptors (Rs). Parkin, phosphorylated by PINK1, ubiquitinates several known 'Rs', including VDAC, mitofusins (MFN1 and MFN2), and the mammalian mitochondrial Rho (Miro). **(2):** Mitochondrial fission facilitates mitophagy since smaller mitochondria are easily engulfed and degraded. **(3):** PINK1 and Parkin arrest mitochondrial motility through degradation of Miro. **(4):** The PINK1-dependent, Parkin-independent mitophagy pathway. In addition to Parkin, alternative E3 ubiquitin ligases like SIAH1 and ARIH1 participate in PINK1-dependent, Parkin-independent mitophagy.

529 **B. Receptor-mediated mitophagy pathways.** Here, the receptor-mediated mitophagy pathways are
530 divided into three groups. **(1):** Group one comprises OMM proteins, including FUNDC1, NIX (BNIP3L),
531 BNIP3, FKBP8, and BCL3L13 (the mammalian orthologue of the yeast Atg32). **(2):** Group two contains
532 mitophagy receptors which are IMM components, including PHB2 and cardiolipin (Card.). **(3):** Group
533 three comprises autophagy receptors which are also involved in mitophagy including AMBRA1, NDP52,
534 TAXIBP1, NBR1, OPTN, and p62.
535 For references, see the main text.
536

Table 1. Proteins Involved in Mitophagy and Their Relations to Neurodegenerative Diseases.

Proteins	Functions in mitophagy	Relation to diseases
Specific receptors for mitophagy		
PINK1	Ser/Thr kinase; initiates mitophagy by phosphorylating components of mitophagy machinery including Ubiquitin and Parkin among others.	PD, AD, ALS
Parkin	E3 ubiquitin ligase: initiates mitophagy by ubiquitinating proteins on the OMM. PTEN-L inhibits PINK1/Parkin dependent mitophagy via dephosphorylation of PINK1-mediated pSer65Ub.	PD, AD, ALS
RIPK1	Binds on PGAM5 and PINK1 to induce mitophagy	ALS, Cancer
PGAM5	Involved in PINK1-dependent mitophagy.	PD, ALS
ARIH1/HHAR1	E3 ligase, RING-between-RING E3 ligase family: participates in PINK1-dependent Parkin-independent mitophagy in Parkin-deficient cancer cells.	Cancer, breast and lung cancer especially
NIX/BNIP3L	Mitochondrial receptor protein: essential in maturation of reticulocytes and hypoxia-induced mitophagy.	AD, PD
BNIP3	Involved in hypoxia-induced mitophagy; regulates OPA1 disassembling and DRP1 recruitment to mitochondria to facilitate mitophagy; stabilizes PINK1.	PD?
FUNDC1	OMM protein: initiates hypoxia-induced mitophagy, where phosphorylation of FUNDC1 by CK2 and Src is inhibited and dephosphorylation by PAGM5 is stimulated, hereby allowing FUNDC1 to bind to LC3 and recruit DRP1/DNM1L.	-
BCL2L13	OMM protein: involved in mitochondrial dynamics, Parkin-independent mitophagy and cargo binding.	-
FKBP8/(38)	OMM protein: involved in Parkin-independent mitophagy by binding to LC3A.	-
PHB2	IMM protein: interacts with LC3 following mitochondrial membrane rupture. Participates in Parkin-mediated mitophagy.	-
Cardiolipin	IMM phospholipid: involved in cargo labelling, via interaction with LC3 following mitochondrial membrane rupture.	-
Receptors for both macroautophagy and mitophagy		
AMBRA1	Working with PI3K III nucleation complex to initiate phagophore formation; Participates in PINK1/Parkin (in)dependent mitophagy.	Neural tube defects
ULK1	Ser/Thr kinase: Activation of AMPK or inhibition of mTOR induces composition of ULK1 initiation complex.	AD, development defects of axon
p62 (SQSTM1)	Autophagy/mitophagy receptor: involved in cargo sequestration; degrades sperm-derived mitochondria in fruit flies and mammals.	Motor neuron disease
NDP52	Autophagy receptor: involved in cargo sequestration.	-
OPTN	Involved in cargo sequestration, where it is being regulated by TBK1.	ALS
Beclin 1	Promotes formation of PI3KC3-C1 and regulates the lipid kinase.	PD
TBK1	Kinase: binds ubiquitin chains on mitochondrial proteins, then phosphorylates autophagy receptors and facilitates mitophagy/autophagy.	Motor neuron disease ALS

References: PINK1: [44, 45, 88]; Parkin: [44, 45, 89-92]; PTEN-L: [91]; RIPK1: [59, 60]; PGAM5: [59, 60]; ARIH1: [93]; NIX: [13, 94]; BNIP3: [95-99]; FUNDC1: [100-103]; BCL2L13: [104-108]; FKBP8: [109, 110]; PHB2: [111]; Cardiolipin: [112, 113]; LC3: [114]; GABARAP: [7, 114, 115]; AMBRA1: [116-118]; ULK1: [9, 98, 119-123]; p62: [124-126]; NDP52: [88]; OPTN: [127]; Beclin 1: [120]; and other references [7, 9, 128, 129].

546 **Table 2. Compounds targeting mitophagy and their known Health Effects.**

Compounds	Mechanistic linkage to mitophagy	Health benefits (More than mitophagy)
NR	Precursor of NAD ⁺ ; NAD ⁺ enhances expression of NIX (DCT-1); induces mitochondrial fission; enhances autophagy through sirtuin-dependent deacetylation of Atg5, Atg7, and Atg8; Under starvation, the NAD ⁺ /SIRT1 pathway deacetylates nuclear LC3 to facilitate its cytoplasmic redistribution.	Inhibits cognitive deficits, inflammation, and Tau pathology in AD; inhibits hearing loss; promotes healthspan and lifespan in mice and worms.
NMN	Precursor of NAD ⁺ ; NAD ⁺ enhances expression of NIX (DCT-1); induces mitochondrial fission; enhances autophagy through sirtuin-dependent deacetylation of Atg5, Atg7, and Atg8; Under starvation, the NAD ⁺ /SIRT1 pathway deacetylates nuclear LC3 to facilitate its cytoplasmic redistribution.	Inhibits cognitive deficits, inflammation, and Tau pathology in AD; inhibits hearing loss; promotes healthspan and lifespan in mice and worms.
NAM	Precursor of NAD ⁺ but higher level inhibits SIRT1 activity; affects SIRT1, SIRT3, PI3K-AKT, MAPK/ERK1/2, CREB	Reduces A β and Tau and inhibits cognitive deficits in 3xTgAD; Promotes healthspan not lifespan in mice
Urolithin A	Induces expression of mitophagy proteins, including full-length PINK1, Parkin, OPTN; p-ULK1, LC3B-II, Beclin1, Bcl2L13, AMBRA1, and FUNDC1 in SH-SY5Y cells; induces expression of full-length PINK1 in brain tissues of mice.	Inhibits A β and Tau pathology and reverses cognitive deficits in worm and mouse models of AD.
Actinonin	Induces expression of mitophagy proteins, including full-length PINK1, Parkin, OPTN; p-ULK1, LC3B-II, Beclin1, Bcl2L13, AMBRA1, and FUNDC1 in SH-SY5Y cells; induces expression of full-length PINK1 in brain tissues of mice.	Inhibits A β and Tau pathology and reverses cognitive deficits in worm and mouse models of AD.
Rapamycin	Induces macro-autophagy by direct binding and inhibition of mTOR; stimulates AMPK; extends lifespan in mice in an ULK1-dependent manner.	Promotes healthspan and lifespan in yeast, worms, flies, and mice; shows anti-cancer and anti-inflammation activities.
Spermidine	Induces mitophagy through multiple pathways, involving the ATM-PINK1-Parkin pathway, the Nrf2-SKN-1 pathway, and through activation of AMPK and inhibition of mTOR; inhibits EP300; induces BNIP3, CTSL, and ATGs.	Promotes healthspan and lifespan in yeast, worms, flies, and mice;
Metformin	Induces autophagy/mitophagy via SIRT1, IGF-1 and mTORC1, or via Parkin-mediated mitophagy.	Promotes healthspan and lifespan in worms and mice.

548
549
550
551
552
References: NR and NMN: [18, 28, 29, 70, 130, 131]; NAM: [68, 69]; Urolithin A: [10, 74]; Actinonin: [10, 23]; Rapamycin: [66, 81, 82]; Spermidine: [64, 65, 77, 80]; Metformin: [81, 83-85]; and other references[7, 9, 128, 129].

References

1. Lane, N. and Martin, W. (2010) The energetics of genome complexity. *Nature* 467 (7318), 929-34.
2. Wallace, D.C. (1999) Mitochondrial diseases in man and mouse. *Science* 283 (5407), 1482-8.
3. Scheibye-Knudsen, M. et al. (2015) Protecting the mitochondrial powerhouse. *Trends Cell Biol* 25 (3), 158-70.
4. Fang, E.F. et al. (2016) Nuclear DNA damage signalling to mitochondria in ageing. *Nat Rev Mol Cell Biol* 17 (5), 308-21.
5. Scheibye-Knudsen, M. et al. (2014) A high-fat diet and NAD(+) activate Sirt1 to rescue premature aging in cockayne syndrome. *Cell Metab* 20 (5), 840-55.
6. Mattson, M.P. et al. (2008) Mitochondria in neuroplasticity and neurological disorders. *Neuron* 60 (5), 748-66.
7. Palikaras, K. et al. (2018) Mechanisms of mitophagy in cellular homeostasis, physiology and pathology. *Nat Cell Biol* 20 (9), 1013-1022.
8. Wallace, D.C. (2012) Mitochondria and cancer. *Nat Rev Cancer* 12 (10), 685-98.
9. Kerr, J.S. et al. (2017) Mitophagy and Alzheimer's Disease: Cellular and Molecular Mechanisms. *Trends Neurosci* 40 (3), 151-166.
10. Fang, E.F. et al. (2019) Mitophagy inhibits amyloid- β and tau pathology and reverses cognitive deficits in models of Alzheimer's disease. *Nature Neuroscience* 22 (3), 401-412.
11. Lin, M.T. and Beal, M.F. (2006) Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature* 443 (7113), 787-95.
12. Bose, A. and Beal, M.F. (2019) Mitochondrial dysfunction and oxidative stress in induced pluripotent stem cell models of Parkinson's disease. *Eur J Neurosci* 49 (4), 525-532.
13. Sandoval, H. et al. (2008) Essential role for Nix in autophagic maturation of erythroid cells. *Nature* 454 (7201), 232-5.
14. Sato, M. and Sato, K. (2011) Degradation of paternal mitochondria by fertilization-triggered autophagy in *C. elegans* embryos. *Science* 334 (6059), 1141-4.
15. Fivenson, E.M. et al. (2017) Mitophagy in neurodegeneration and aging. *Neurochem Int* 109, 202-209.
16. Lopez-Otin, C. et al. (2013) The hallmarks of aging. *Cell* 153 (6), 1194-217.
17. Fang, E.F. et al. (2017) NAD(+) in Aging: Molecular Mechanisms and Translational Implications. *Trends in Molecular Medicine* 23 (10), 899-916.
18. Fang, E.F. et al. (2014) Defective mitophagy in XPA via PARP-1 hyperactivation and NAD(+)/SIRT1 reduction. *Cell* 157 (4), 882-896.
19. Palikaras, K. et al. (2015) Coordination of mitophagy and mitochondrial biogenesis during ageing in *C. elegans*. *Nature* 521 (7553), 525-8.
20. Palikaras, K. et al. (2017) Mitophagy and age-related pathologies: Development of new therapeutics by targeting mitochondrial turnover. *Pharmacol Ther* 178, 157-174.
21. Wiley, C.D. et al. (2016) Mitochondrial Dysfunction Induces Senescence with a Distinct Secretory Phenotype. *Cell Metabolism* 23 (2), 303-314.
22. Kirienko, N.V. et al. (2015) Mitophagy confers resistance to siderophore-mediated killing by *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 112 (6), 1821-6.
23. Sun, N. et al. (2015) Measuring In Vivo Mitophagy. *Mol Cell* 60 (4), 685-96.
24. Garcia-Prat, L. et al. (2016) Autophagy maintains stemness by preventing senescence. *Nature* 529 (7584), 37-42.
25. Pugh, P.L. et al. (2004) A behavioural characterisation of the FVB/N mouse strain. *Behav Brain Res* 155 (2), 283-9.
26. Cummins, N. et al. (2018) Disease-associated tau impairs mitophagy by inhibiting Parkin translocation to mitochondria. *EMBO J*.
27. Keijzers, G. et al. (2018) Monogenic Diseases of DNA Repair. *N Engl J Med* 378 (5), 491-492.
28. Scheibye-Knudsen, M. et al. (2014) A high-fat diet and NAD(+) activate Sirt1 to rescue premature aging in cockayne syndrome. *Cell Metab* 20 (5), 840-855.
29. Fang, E.F. et al. (2016) NAD(+) Replenishment Improves Lifespan and Healthspan in Ataxia Telangiectasia Models via Mitophagy and DNA Repair. *Cell Metab* 24 (4), 566-581.
30. Sumpter, R., Jr. et al. (2016) Fanconi Anemia Proteins Function in Mitophagy and Immunity. *Cell* 165 (4), 867-81.
31. Canter, R.G. et al. (2016) The road to restoring neural circuits for the treatment of Alzheimer's disease. *Nature* 539 (7628), 187-196.
32. Menzies, F.M. et al. (2015) Compromised autophagy and neurodegenerative diseases. *Nat Rev Neurosci* 16 (6), 345-57.
33. Nixon, R.A. (2013) The role of autophagy in neurodegenerative disease. *Nat Med* 19 (8), 983-97.

34. Swerdlow, R.H. et al. (2014) The Alzheimer's disease mitochondrial cascade hypothesis: progress and perspectives. *Biochim Biophys Acta* 1842 (8), 1219-31.
35. Cai, Q. and Tammineni, P. (2016) Alterations in Mitochondrial Quality Control in Alzheimer's Disease. *Front Cell Neurosci* 10, 24.
36. Goiran, T. et al. (2018) beta-Amyloid Precursor Protein Intracellular Domain Controls Mitochondrial Function by Modulating Phosphatase and Tensin Homolog-Induced Kinase 1 Transcription in Cells and in Alzheimer Mice Models. *Biol Psychiatry* 83 (5), 416-427.
37. Lauritzen, I. et al. (2016) Intraneuronal aggregation of the beta-CTF fragment of APP (C99) induces Abeta-independent lysosomal-autophagic pathology. *Acta Neuropathol* 132 (2), 257-76.
38. Martin-Maestro, P. et al. (2017) Mitophagy Failure in Fibroblasts and iPSC-Derived Neurons of Alzheimer's Disease-Associated Presenilin 1 Mutation. *Front Mol Neurosci* 10, 291.
39. Tammineni, P. et al. (2017) Impaired retrograde transport of axonal autophagosomes contributes to autophagic stress in Alzheimer's disease neurons. *Elife* 6.
40. Sorrentino, V. et al. (2017) Enhancing mitochondrial proteostasis reduces amyloid-beta proteotoxicity. *Nature*.
41. Schondorf, D.C. et al. (2018) The NAD⁺ Precursor Nicotinamide Riboside Rescues Mitochondrial Defects and Neuronal Loss in iPSC and Fly Models of Parkinson's Disease. *Cell Rep* 23 (10), 2976-2988.
42. Mosley, R.L. et al. (2012) Inflammation and adaptive immunity in Parkinson's disease. *Cold Spring Harb Perspect Med* 2 (1), a009381.
43. Palin, E.J. et al. (2013) Mesencephalic complex I deficiency does not correlate with parkinsonism in mitochondrial DNA maintenance disorders. *Brain* 136 (Pt 8), 2379-92.
44. Kitada, T. et al. (1998) Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature* 392 (6676), 605-8.
45. Valente, E.M. et al. (2004) Hereditary early-onset Parkinson's disease caused by mutations in PINK1. *Science* 304 (5674), 1158-60.
46. Matheoud, D. et al. (2016) Parkinson's Disease-Related Proteins PINK1 and Parkin Repress Mitochondrial Antigen Presentation. *Cell* 166 (2), 314-327.
47. Perez, F.A. and Palmiter, R.D. (2005) Parkin-deficient mice are not a robust model of parkinsonism. *Proc Natl Acad Sci U S A* 102 (6), 2174-9.
48. Kitada, T. et al. (2007) Impaired dopamine release and synaptic plasticity in the striatum of PINK1-deficient mice. *Proc Natl Acad Sci U S A* 104 (27), 11441-6.
49. McWilliams, T.G. et al. (2018) Basal Mitophagy Occurs Independently of PINK1 in Mouse Tissues of High Metabolic Demand. *Cell Metab* 27 (2), 439-449 e5.
50. Yang, W. et al. (2019) CRISPR/Cas9-mediated PINK1 deletion leads to neurodegeneration in rhesus monkeys. *Cell Res* 29 (4), 334-336.
51. Sliter, D.A. et al. (2018) Parkin and PINK1 mitigate STING-induced inflammation. *Nature* 561 (7722), 258-262.
52. West, A.P. et al. (2015) Mitochondrial DNA stress primes the antiviral innate immune response. *Nature* 520 (7548), 553-7.
53. Riley, J.S. et al. (2018) Mitochondrial inner membrane permeabilisation enables mtDNA release during apoptosis. *EMBO J* 37 (17).
54. Ryan, B.J. et al. (2015) Mitochondrial dysfunction and mitophagy in Parkinson's: from familial to sporadic disease. *Trends Biochem Sci* 40 (4), 200-10.
55. Marcassa, E. et al. (2018) Dual role of USP30 in controlling basal pexophagy and mitophagy. *EMBO Rep* 19 (7).
56. Yamada, T. et al. (2018) Mitochondrial Stasis Reveals p62-Mediated Ubiquitination in Parkin-Independent Mitophagy and Mitigates Nonalcoholic Fatty Liver Disease. *Cell Metab* 28 (4), 588-604 e5.
57. Moore, A.S. and Holzbaur, E.L. (2016) Dynamic recruitment and activation of ALS-associated TBK1 with its target optineurin are required for efficient mitophagy. *Proc Natl Acad Sci U S A* 113 (24), E3349-58.
58. Weishaupt, J.H. et al. (2016) Common Molecular Pathways in Amyotrophic Lateral Sclerosis and Frontotemporal Dementia. *Trends Mol Med* 22 (9), 769-783.
59. Ito, Y. et al. (2016) RIPK1 mediates axonal degeneration by promoting inflammation and necroptosis in ALS. *Science* 353 (6299), 603-8.
60. Hawk, M.A. et al. (2018) RIPK1-mediated induction of mitophagy compromises the viability of extracellular-matrix-detached cells. *Nat Cell Biol* 20 (3), 272-284.
61. Ross, C.A. and Tabrizi, S.J. (2011) Huntington's disease: from molecular pathogenesis to clinical treatment. *Lancet Neurol* 10 (1), 83-98.

62. Hwang, S. et al. (2015) Impaired GAPDH-induced mitophagy contributes to the pathology of Huntington's disease. *EMBO Mol Med* 7 (10), 1307-26.
63. Moreno-Jimenez, E.P. et al. (2019) Adult hippocampal neurogenesis is abundant in neurologically healthy subjects and drops sharply in patients with Alzheimer's disease. *Nat Med* 25 (4), 554-560.
64. Gupta, V.K. et al. (2013) Restoring polyamines protects from age-induced memory impairment in an autophagy-dependent manner. *Nat Neurosci* 16 (10), 1453-60.
65. Madeo, F. et al. (2018) Spermidine in health and disease. *Science* 359 (6374).
66. Ravikumar, B. et al. (2004) Inhibition of mTOR induces autophagy and reduces toxicity of polyglutamine expansions in fly and mouse models of Huntington disease. *Nat Genet* 36 (6), 585-95.
67. Hou, Y. et al. (2018) NAD(+) supplementation normalizes key Alzheimer's features and DNA damage responses in a new AD mouse model with introduced DNA repair deficiency. *Proc Natl Acad Sci U S A* 115 (8), E1876-E1885.
68. Liu, D. et al. (2013) Nicotinamide forestalls pathology and cognitive decline in Alzheimer mice: evidence for improved neuronal bioenergetics and autophagy procession. *Neurobiol Aging* 34 (6), 1564-80.
69. Turunc Bayrakdar, E. et al. (2014) Nicotinamide treatment reduces the levels of oxidative stress, apoptosis, and PARP-1 activity in Aβeta(1-42)-induced rat model of Alzheimer's disease. *Free Radic Res* 48 (2), 146-58.
70. Gong, B. et al. (2013) Nicotinamide riboside restores cognition through an upregulation of proliferator-activated receptor-gamma coactivator 1α regulated beta-secretase 1 degradation and mitochondrial gene expression in Alzheimer's mouse models. *Neurobiol Aging* 34 (6), 1581-8.
71. Fang, E.F. (2019) Mitophagy and NAD(+) inhibit Alzheimer disease. *Autophagy*, 1-3.
72. Trammell, S.A. et al. (2016) Nicotinamide riboside is uniquely and orally bioavailable in mice and humans. *Nat Commun* 7, 12948.
73. Dollerup, O.L. et al. (2018) A randomized placebo-controlled clinical trial of nicotinamide riboside in obese men: safety, insulin-sensitivity, and lipid-mobilizing effects. *Am J Clin Nutr* 108 (2), 343-353.
74. Ryu, D. et al. (2016) Urolithin A induces mitophagy and prolongs lifespan in *C. elegans* and increases muscle function in rodents. *Nat Med* 22 (8), 879-88.
75. Ip, W.K.E. et al. (2017) Anti-inflammatory effect of IL-10 mediated by metabolic reprogramming of macrophages. *Science* 356 (6337), 513-519.
76. Andreux, P.A. et al. (2019) The mitophagy activator urolithin A is safe and induces a molecular signature of improved mitochondrial and cellular health in humans. *Nature Metab* 1, 593-603.
77. Eisenberg, T. et al. (2009) Induction of autophagy by spermidine promotes longevity. *Nat Cell Biol* 11 (11), 1305-14.
78. Eisenberg, T. et al. (2016) Cardioprotection and lifespan extension by the natural polyamine spermidine. *Nat Med* 22 (12), 1428-1438.
79. Fan, J. et al. (2017) Spermidine coupled with exercise rescues skeletal muscle atrophy from D-gal-induced aging rats through enhanced autophagy and reduced apoptosis via AMPK-FOXO3a signal pathway. *Oncotarget* 8 (11), 17475-17490.
80. Qi, Y. et al. (2016) ATM mediates spermidine-induced mitophagy via PINK1 and Parkin regulation in human fibroblasts. *Sci Rep* 6, 24700.
81. Aliper, A. et al. (2017) Towards natural mimetics of metformin and rapamycin. *Aging (Albany NY)* 9 (11), 2245-2268.
82. Roth, G.S. and Ingram, D.K. (2016) Manipulation of health span and function by dietary caloric restriction mimetics. *Ann N Y Acad Sci* 1363, 5-10.
83. Howell, J.J. et al. (2017) Metformin Inhibits Hepatic mTORC1 Signaling via Dose-Dependent Mechanisms Involving AMPK and the TSC Complex. *Cell Metab* 25 (2), 463-471.
84. Zheng, Z. et al. (2012) Sirtuin 1-mediated cellular metabolic memory of high glucose via the LKB1/AMPK/ROS pathway and therapeutic effects of metformin. *Diabetes* 61 (1), 217-28.
85. Song, Y.M. et al. (2016) Metformin Restores Parkin-Mediated Mitophagy, Suppressed by Cytosolic p53. *Int J Mol Sci* 17 (1).
86. Bakula, D. et al. (2018) Aging and drug discovery. *Aging (Albany NY)* 10 (11), 3079-3088.
87. Mattson, M.P. et al. (2018) Intermittent metabolic switching, neuroplasticity and brain health. *Nat Rev Neurosci* 19 (2), 63-80.
88. Lazarou, M. et al. (2015) The ubiquitin kinase PINK1 recruits autophagy receptors to induce mitophagy. *Nature* 524 (7565), 309-314.
89. Geisler, S. et al. (2010) PINK1/Parkin-mediated mitophagy is dependent on VDAC1 and p62/SQSTM1. *Nat Cell Biol* 12 (2), 119-31.
90. Gegg, M.E. et al. (2010) Mitofusin 1 and mitofusin 2 are ubiquitinated in a PINK1/parkin-dependent manner upon induction of mitophagy. *Hum Mol Genet* 19 (24), 4861-70.

91. Wang, X. et al. (2011) PINK1 and Parkin target Miro for phosphorylation and degradation to arrest mitochondrial motility. *Cell* 147 (4), 893-906.
92. Gao, F. et al. (2015) The mitochondrial protein BNIP3L is the substrate of PARK2 and mediates mitophagy in PINK1/PARK2 pathway. *Hum Mol Genet* 24 (9), 2528-38.
93. Villa, E. et al. (2017) Parkin-Independent Mitophagy Controls Chemotherapeutic Response in Cancer Cells. *Cell Rep* 20 (12), 2846-2859.
94. Schwarten, M. et al. (2009) Nix directly binds to GABARAP: a possible crosstalk between apoptosis and autophagy. *Autophagy* 5 (5), 690-8.
95. Zhu, Y. et al. (2013) Modulation of serines 17 and 24 in the LC3-interacting region of Bnip3 determines pro-survival mitophagy versus apoptosis. *J Biol Chem* 288 (2), 1099-113.
96. Zhang, H. et al. (2008) Mitochondrial autophagy is an HIF-1-dependent adaptive metabolic response to hypoxia. *J Biol Chem* 283 (16), 10892-903.
97. Quinsay, M.N. et al. (2010) Bnip3 mediates permeabilization of mitochondria and release of cytochrome c via a novel mechanism. *J Mol Cell Cardiol* 48 (6), 1146-56.
98. Lee, Y. et al. (2011) Mitochondrial autophagy by Bnip3 involves Drp1-mediated mitochondrial fission and recruitment of Parkin in cardiac myocytes. *Am J Physiol Heart Circ Physiol* 301 (5), H1924-31.
99. Zhang, T. et al. (2016) BNIP3 Protein Suppresses PINK1 Kinase Proteolytic Cleavage to Promote Mitophagy. *J Biol Chem* 291 (41), 21616-21629.
100. Liu, L. et al. (2012) Mitochondrial outer-membrane protein FUNDC1 mediates hypoxia-induced mitophagy in mammalian cells. *Nat Cell Biol* 14 (2), 177-85.
101. Chen, G. et al. (2014) A regulatory signaling loop comprising the PGAM5 phosphatase and CK2 controls receptor-mediated mitophagy. *Mol Cell* 54 (3), 362-77.
102. Wu, W. et al. (2014) ULK1 translocates to mitochondria and phosphorylates FUNDC1 to regulate mitophagy. *EMBO Rep* 15 (5), 566-75.
103. Chen, M. et al. (2016) Mitophagy receptor FUNDC1 regulates mitochondrial dynamics and mitophagy. *Autophagy* 12 (4), 689-702.
104. Kanki, T. et al. (2009) Atg32 is a mitochondrial protein that confers selectivity during mitophagy. *Dev Cell* 17 (1), 98-109.
105. Okamoto, K. et al. (2009) Mitochondria-anchored receptor Atg32 mediates degradation of mitochondria via selective autophagy. *Dev Cell* 17 (1), 87-97.
106. Kanki, T. et al. (2013) Casein kinase 2 is essential for mitophagy. *Embo Reports* 14 (9), 788-794.
107. Wei, H.F. et al. (2015) Selective removal of mitochondria via mitophagy: distinct pathways for different mitochondrial stresses. *Biochimica Et Biophysica Acta-Molecular Cell Research* 1853 (10), 2784-2790.
108. Murakawa, T. et al. (2015) Bcl-2-like protein 13 is a mammalian Atg32 homologue that mediates mitophagy and mitochondrial fragmentation. *Nature Communications* 6.
109. Misaka, T. et al. (2018) FKBP8 protects the heart from hemodynamic stress by preventing the accumulation of misfolded proteins and endoplasmic reticulum-associated apoptosis in mice. *J Mol Cell Cardiol* 114, 93-104.
110. Bhujabal, Z. et al. (2017) FKBP8 recruits LC3A to mediate Parkin-independent mitophagy. *EMBO Rep* 18 (6), 947-961.
111. Wei, Y. et al. (2017) Prohibitin 2 Is an Inner Mitochondrial Membrane Mitophagy Receptor. *Cell* 168 (1-2), 224-238 e10.
112. Shen, Z. et al. (2017) Cardiolipin Regulates Mitophagy through the Protein Kinase C Pathway. *J Biol Chem* 292 (7), 2916-2923.
113. Chu, C.T. et al. (2013) Cardiolipin externalization to the outer mitochondrial membrane acts as an elimination signal for mitophagy in neuronal cells. *Nat Cell Biol* 15 (10), 1197-1205.
114. Shpilka, T. et al. (2011) Atg8: an autophagy-related ubiquitin-like protein family. *Genome Biol* 12 (7), 226.
115. Weidberg, H. et al. (2011) LC3 and GATE-16 N termini mediate membrane fusion processes required for autophagosome biogenesis. *Dev Cell* 20 (4), 444-54.
116. Fimia, G.M. et al. (2011) Unleashing the Ambra1-Becn1 complex from dynein chains: Ulk1 sets Ambra1 free to induce autophagy. *Autophagy* 7 (1), 115-7.
117. Di Rita, A. et al. (2018) HUWE1 E3 ligase promotes PINK1/PARKIN-independent mitophagy by regulating AMBRA1 activation via IKKalpha. *Nat Commun* 9 (1), 3755.
118. Van Humbeeck, C. et al. (2011) Parkin interacts with Ambra1 to induce mitophagy. *J Neurosci* 31 (28), 10249-61.
119. Fimia, G.M. et al. (2007) Ambra1 regulates autophagy and development of the nervous system. *Nature* 447 (7148), 1121-5.

120. Russell, R.C. et al. (2013) ULK1 induces autophagy by phosphorylating Beclin-1 and activating VPS34 lipid kinase. *Nat Cell Biol* 15 (7), 741-50.
121. Egan, D.F. et al. (2011) Phosphorylation of ULK1 (hATG1) by AMP-activated protein kinase connects energy sensing to mitophagy. *Science* 331 (6016), 456-61.
122. Cheong, H. et al. (2011) Ammonia-induced autophagy is independent of ULK1/ULK2 kinases. *Proc Natl Acad Sci U S A* 108 (27), 11121-6.
123. Wang, B. et al. (2018) The autophagy-inducing kinases, ULK1 and ULK2, regulate axon guidance in the developing mouse forebrain via a noncanonical pathway. *Autophagy* 14 (5), 796-811.
124. Pankiv, S. et al. (2007) p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of ubiquitinated protein aggregates by autophagy. *J Biol Chem* 282 (33), 24131-45.
125. Politi, Y. et al. (2014) Paternal mitochondrial destruction after fertilization is mediated by a common endocytic and autophagic pathway in *Drosophila*. *Dev Cell* 29 (3), 305-20.
126. Song, W.H. et al. (2016) Autophagy and ubiquitin-proteasome system contribute to sperm mitophagy after mammalian fertilization. *Proc Natl Acad Sci U S A* 113 (36), E5261-70.
127. Richter, B. et al. (2016) Phosphorylation of OPTN by TBK1 enhances its binding to Ub chains and promotes selective autophagy of damaged mitochondria. *Proc Natl Acad Sci U S A* 113 (15), 4039-44.
128. Rubinsztein, D.C. et al. (2011) Autophagy and aging. *Cell* 146 (5), 682-95.
129. Hansen, M. et al. (2018) Autophagy as a promoter of longevity: insights from model organisms. *Nat Rev Mol Cell Biol* 19 (9), 579-593.
130. Brown, K.D. et al. (2014) Activation of SIRT3 by the NAD(+) precursor nicotinamide riboside protects from noise-induced hearing loss. *Cell Metab* 20 (6), 1059-68.
131. Someya, S. et al. (2010) Sirt3 mediates reduction of oxidative damage and prevention of age-related hearing loss under caloric restriction. *Cell* 143 (5), 802-12.