Convergence rates for stochastic and degenerate-parabolic balance laws

Erlend Briseid Storrøsten

Dissertation presented for the degree of Philosophiae Doctor

Department of Mathematics
University of Oslo

October 2015

© Erlend Briseid Storrøsten, 2015

*Series of dissertations submitted to the
Faculty of Mathematics and Natural Sciences, University of Oslo
No. 1726*

ISSN 1501-7710

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

Acknowledgments

The writing of this thesis was carried out at the Department of Mathematics at the University of Oslo, and has been funded by the Research Council of Norway (NFR) through the Center of Mathematics for Applications (CMA). Let me express my gratitude towards CMA, NFR, the Department of Mathematics, and the University of Oslo for making the writing of this thesis possible.

Carrying out the project has truly been a long distance race, and I would not have been able to complete it without the encouragement and support of a great number of people. Let me first of all thank my supervisors Nils Henrik Risebro and Kenneth H. Karlsen. I am deeply grateful for everything I have learned. It is not only a matter of mathematics, it is also a matter of spirit. A heartfelt thanks to all my colleagues at the department, and in particular at CMA, for coffee breaks, discussions, beers and much more. A special thanks is given to Torstein K. Nilssen, Johanna Ridder and Håkon Andreas Hoel for discussing, reading and commenting on the introduction of the thesis.

Although, slightly less involved in the thesis, understandably, I would like to mention my family. Thanks for all your love and support, unconditionally there, always. Finally, let me thank Tonje. I could never have dreamed of anyone more supportive and understanding. Hopefully, life will be a little less work, and a little more of everything else in years to come.

Oslo, October 2015
Erlend Briseid Storrøsten

Contents

Acknowledgments	iii
Introduction	1
Overview of the thesis	1
Scalar conservation laws	4
Doubling-of-variables	6
The kinetic formulation	10
Discussion of main ideas and techniques	14
Part I: Finite difference approximations of degenerate convection-diffusion equations	14
Part II: Stochastic scalar conservation laws	19
References	29
Papers	31
Kenneth H. Karlsen, Nils Henrik Risebro and Erlend B. Storrøsten. $\mathbb{L}^1$ error estimates for difference approximations of degenerate convection-diffusion equations. <i>Math. Comp.</i> , 83(290):2717–2762, 2014.	35
Kenneth H. Karlsen, Nils Henrik Risebro and Erlend B. Storrøsten. On the convergence rate of finite difference methods for degenerate convection-diffusion equations in several space dimensions. <i>ESAIM:M2AN</i> Forthcoming.	83
Kenneth H. Karlsen and Erlend B. Storrøsten. On stochastic conservation laws and Malliavin calculus. <i>Submitted</i>	127
Kenneth H. Karlsen and Erlend B. Storrøsten. Analysis of a splitting method for stochastic balance laws. <i>Preprint</i>	183
Appendix	221
A.1 A note on the representation of nonlinear limits	221

A.2 A degenerate convection-diffusion equation arising in two phase flow in porous media.	234
References	236

Introduction

Overview of the thesis

This thesis consists of two parts, presented through four papers, henceforth referred to as paper I to IV.

The first part is concerned with finite difference approximations of the Cauchy problem

$$\partial_t u + \nabla \cdot f(u) = \Delta A(u), \quad u(0, x) = u_0(x), \quad (1)$$

where the flux function $f : \mathbb{R} \rightarrow \mathbb{R}^d$ and the diffusion function $A : \mathbb{R} \rightarrow \mathbb{R}$ are given nonlinear Lipschitz continuous functions. Furthermore we assume that $A' \geq 0$. The initial datum u_0 belongs to $L^\infty \cap L^1 \cap BV$. The above equation is referred to as a *degenerate convection-diffusion equation*, where “degenerate” refers to the fact that A is merely assumed to be nondecreasing. Problems of the form (1) arise in several important applications. Let me mention sedimentation processes [6, 7] and flow in porous media [15]. An example of the latter is given in Appendix A.2. Degeneracy is of importance due to the fact that when A' vanishes, solutions to (1) may develop discontinuities (shocks) in finite time. As a consequence (1) must be interpreted in the weak sense. Regarding weak solutions, a second condition is necessary to ensure that (1) is well posed, the so-called entropy condition.

To obtain a finite difference approximation of (1), the flux is discretized by a two-point monotone numerical flux and the diffusion term is approximated by a second order central difference. As an example, consider a semidiscrete scheme in one spatial dimension and suppose for simplicity that $f' \geq 0$. Let $\Delta x > 0$ and suppose $U = U_j(t)$ satisfies

$$\frac{d}{dt} U_j(t) + \frac{f(U_j) - f(U_{j-1})}{\Delta x} = \frac{A(U_{j+1}) - 2A(U_j) + A(U_{j-1}))}{\Delta x^2}, \quad U_j(0) = \int_{I_j} u_0(x) dx,$$

where $I_j = [(j - 1/2)\Delta x, (j + 1/2)\Delta x)$. Let $u_{\Delta x}$ be the piecewise constant interpolation, i.e., $u_{\Delta x}(t, x) = U_j(t)$ whenever $x \in I_j$. Let $T > 0$ be a fixed final time. The problem is to determine the largest possible number $r > 0$ such that

$$\|u_{\Delta x}(t, \cdot) - u(t, \cdot)\|_{L^1} \leq C\Delta x^r, \quad (0 \leq t \leq T), \quad (2)$$

for some constant $C > 0$ independent of $\Delta x > 0$. The number $r > 0$ is then referred to as the *rate of convergence* for the approximation $u_{\Delta x}$. The rate of convergence characterizes the speed of convergence, and as such it is a fundamental property regarding the efficiency of the numerical approximation. When $A' \geq \eta$ for some real number $\eta > 0$, solutions to (1)

are smooth and convergence takes place according to the formal order of the scheme. In the purely hyperbolic case, i.e., $A = 0$, Kuznecov showed that monotone finite difference schemes converge with rate $r = \frac{1}{2}$. Furthermore this is optimal [24]. In the general case of degenerate diffusion, the only previous result seems to be by Karlsen et al. in [21]. They established (2) with $r = \frac{1}{11}$ for a semidiscrete scheme in one spatial dimension.

In paper I, entitled “ L^1 error estimates for difference approximations of degenerate convection-diffusion equations”, the case of one spatial dimension is considered for semidiscrete, implicit and explicit finite difference schemes. Here (2) is established with $r = 1/3$, of course subject to certain constraints on the time stepping. The proofs are based on the method of “doubling of variables”, first introduced by Kruřkov for scalar conservation laws [23], and later extended to degenerate convection-diffusion equations by Carrillo [9]. Paper II is entitled “On the convergence rate of finite difference methods for degenerate convection-diffusion equations in several space dimensions”. In this paper, (2) is established for monotone semidiscrete schemes with $r = 2/(19 + d)$ where d is the spatial dimension. The result follows similarly for the implicit scheme. The proof in this paper is based on the so-called kinetic formulation developed by Lions, Perthame and Tadmor [28] and Chen and Perthame [11]. In the case $d = 1$, the rate $r = 1/3$ is retrieved by this technique as well.

The second part deals with scalar conservation laws perturbed by a random source. More precisely we consider the Cauchy problem

$$\partial_t u + \nabla \cdot f(u) = \sigma(x, u) \partial_t W(t), \quad u(0, x) = u_0(x), \quad (3)$$

where W is a Brownian motion. In fact, the model studied in Paper III allows for a slightly more general noise, that is, we let W be a cylindrical Wiener process. Both f and $u \mapsto \sigma(x, u)$ are deterministic Lipschitz continuous functions. Furthermore, we assume the existence of a constant $C > 0$ and a $\kappa \in (0, 1/2]$ such that

$$|\sigma(x_1, u) - \sigma(x_2, u)| \leq C |x_1 - x_2|^{\kappa + \frac{1}{2}} (1 + |u|), \quad (4)$$

for all $x_1, x_2 \in \mathbb{R}^d$ and $u \in \mathbb{R}$. Scalar conservation laws perturbed by a random noise has recently caught a lot of attention. In particular existence and uniqueness for (3) has previously been established by Feng and Nualart in [17] as well as Bauzet et al. in [2]. Both articles are based on a “doubling of variables” approach à la Kruřkov. Similar results has been obtained by Debussche and Vovelle in [12] based on the kinetic formulation. Let me also mention the related works [5, 22, 35, 18, 14, 19]. For an overview of some applications related to (3), see [3].

In paper III, entitled “On stochastic conservation laws and Malliavin calculus”, we propose a new notion of entropy solutions to (3) based on some tools from Malliavin calculus. The goal is to shed some light on certain aspects of both [17] and [2]. To be more precise, we propose a generalization of the Kruřkov entropy condition by allowing the Kruřkov constants to be Malliavin differentiable random variables. That is, we say that any predictable process $u = u(t, x, \omega)$ is an entropy solution to (3) if

$$E \left[\partial_t S(u - V) + \nabla \cdot Q(u, V) + S''(u - V) D_t V \sigma(x, u) - \frac{1}{2} S''(u - V) \sigma^2(x, u) \right] \leq 0,$$

in the sense of distributions for any convex $S \in C^2(\mathbb{R})$ and any Malliavin differentiable random variable V . Here $\partial_u Q(u, V) = S'(u - V)f'(u)$ and $D_t V$ denotes the Malliavin derivative of V evaluated at t . As in [2], existence of solutions to (3) is first established in the class of Young measure valued entropy solutions via the viscous approximation. Subsequently, as a consequence of the uniqueness argument, one may conclude that the Young measure valued entropy solution is in fact a regular entropy solution.

In paper IV, entitled “Analysis of a splitting method for stochastic balance laws”, we consider a time splitting procedure for the Cauchy problem (3). Let $T > 0$ be a fixed final time and $\Delta t = T/N$ for some $N \in \mathbb{N}$. Let $\mathcal{S}_{\text{CL}}$ denote the solution operator associated with the entropy solution of the deterministic Cauchy problem

$$\partial_t u + \nabla \cdot f(u) = 0, \quad u(0, x) = u^0(x).$$

That is, $\mathcal{S}_{\text{CL}}(t)u^0 := u(t)$. Similarly, we let $\mathcal{S}_{\text{SDE}}$ denote the solution operator to the stochastic differential equation $\partial_t u = \sigma(x, u) \partial_t W$. That is, if

$$u(t, x) = u(s, x) + \int_s^t \sigma(x, u(r, x)) dW(r), \quad x \in \mathbb{R},$$

then $\mathcal{S}_{\text{SDE}}(t, s)u(s) = u(t)$. The approximation $\{u^n\}_{n=0}^N$ is defined by

$$u^0 = u_0 \quad \text{and} \quad u^{n+1} = \mathcal{S}_{\text{SDE}}(t_{n+1}, t_n) \mathcal{S}_{\text{CL}}(\Delta t) u^n \quad \text{for } n = 0, \dots, N-1, \quad (5)$$

where $t_n = n\Delta t$. Let $u_{\Delta t}(t, x, \omega) = u^n(x, \omega)$ for $t \in [t_n, t_{n+1})$. Compactness of $\{u_{\Delta t}\}_{\Delta t > 0}$ is established in the sense of Young measures (Theorem 1, Appendix A.1). Under the assumption (4), it is shown that the $u_{\Delta t}$ converges to a Young measure valued entropy solution of (3), and hence to an entropy solution. A similar result has recently been established by Bauzet in [1] under the assumption that $\sigma(x, u) = \sigma(u)$ and $\sigma(0) = 0$. In the case of an x -dependent source term, solutions are no longer of bounded variation, and one has to rely on certain fractional BV estimates [10]. In the case that $\sigma(x, u) = \sigma(u)$ and $\sigma \in L^\infty$, we also establish a rate of convergence. That is, for any $t \in [0, T]$ we establish the existence of a constant $C > 0$ independent of Δt such that

$$E \left[\|u_{\Delta t}(t) - u(t)\|_{L^1_{\text{loc}}(\mathbb{R}^d)} \right] \leq C \Delta t^{1/3}, \quad u_0 \in BV, \quad (6)$$

where u is the entropy solution to (3). Operator splitting for equations of the form (3) has previously been considered in [20, 1, 4]. However, to the best of the authors knowledge, the convergence rate (6) is the first of its kind.

Both the degenerate convection-diffusion equations and the stochastic balance laws under consideration are perturbed versions of scalar conservation laws. As a consequence, the mathematical framework surrounding scalar conservation laws is of fundamental importance. In particular, the notion of entropy solutions and the Kruřkov technique of “doubling of variables” lies at the heart of the matter. Naturally, this is also the case for the alternative framework provided by the kinetic formulation. Therefore, I have chosen to include a brief, informal introduction to scalar conservation laws. The main part is dedicated to the Kruřkov theory of scalar conservation laws, emphasizing the need of an entropy criterion and establishing the technique referred to as “doubling of variables”. An outline of the kinetic approach, alongside a brief

comparison with the Kruřkov theory, is also given. There are several excellent introductions on the topic of scalar conservation laws, see for instance [33, 26, 20, 27]. Concerning the kinetic formulation see [32, 28]. Having presented the main topics in the theory of scalar conservation laws, the introduction continues with a discussion of the main ideas and techniques of the current thesis. In part two, the theory of Young measures plays an important role. I have thus chosen to include some notes on the topic in Appendix A.1.

Scalar conservation laws

Let $(t, x) \mapsto u(t, x)$ denote the density of a certain extensive quantity where t denotes time, and $x = (x_1, \dots, x_d)$ is the spatial variable. In mathematics, a *balance law* states that the rate of change of that quantity in any region $U \subset \mathbb{R}^d$ is given by the flux $(t, x) \mapsto f(t, x)$ through the boundary ∂U plus a given source $(t, x) \mapsto g(t, x)$. That is,

$$\frac{d}{dt} \int_U u(t, x) dx = - \int_{\partial U} f(t, x) \cdot \nu(x) dS(x) + \int_U g(t, x) dx, \quad (7)$$

where $\nu(x)$ denotes the outward pointing unit normal at a point x on the boundary of U and S denotes the surface measure. Alternatively, the production of the extensive quantity is balanced by the flux through the boundary in any given region. When nothing is produced, i.e., $g = 0$, the total amount remains constant, or conserved, and the above is referred to as a *conservation law*. Under the assumption that f and u are sufficiently smooth, an application of the divergence theorem yields

$$\partial_t u(t, x) + \nabla \cdot f(t, x) = g(t, x).$$

The most prominent example of such differential equations are the Euler equations, governing adiabatic (no heat transfer) and inviscid (frictionless) flow. The equations are a consequence of the conservation of mass, momenta and energy and in total constitutes what is referred to as a (hyperbolic) system of conservation laws. In this work we are concerned with the simpler case of a single scalar conservation law. While the theory for systems in general is rather incomplete, the theory for scalar conservation laws is fairly well understood.

For simplicity, we consider the one dimensional case. Suppose $f = f(u)$ where $f : \mathbb{R} \rightarrow \mathbb{R}$ is some nonlinear Lipschitz continuous function. Consider the Cauchy problem

$$\partial_t u + \partial_x f(u) = 0, \quad u(0, x) = u_0(x). \quad (8)$$

Let me state a few observations regarding solutions to (8).

(i) **Formation of shocks.** By the method of characteristics it is seen that even for smooth initial data the solution may become discontinuous after a finite time. This is referred to as the formation of shocks.

(ii) **Invertibility.** Consider the reversed problem

$$\partial_t \tilde{u} + \partial_x \tilde{f}(\tilde{u}) = 0, \quad \tilde{u}(0, x) = \tilde{u}^0(x), \quad (9)$$

where $\tilde{f} = -f^1$. Suppose there exists unique solutions to both (8) and (9). Let S and $\tilde{S}$ denote the solution operators of (8) and (9) respectively. That is, $S(t)u_0 = u(t)$ and $\tilde{S}(t)\tilde{u}^0 = \tilde{u}(t)$. For any $t_0 \geq 0$, $\tilde{u}(t, x) = u(t_0 - t, x)$ is a solution of (9) with $\tilde{u}^0 = u(t_0)$. It follows that $\tilde{S}(t)S(t) = \mathcal{I}$ where $\mathcal{I}$ denotes the identity operator. By symmetry $S(t)\tilde{S}(t) = \mathcal{I}$. Hence $\tilde{S}(t) = (S(t))^{-1}$.

As a consequence of (i) the equation must be considered in the weak sense. That is, if u satisfies

$$\int_0^\infty \int_{\mathbb{R}^d} u(t, x) \partial_t \varphi(t, x) + f(u(t, x)) \partial_x \varphi(t, x) dx dt + \int_{\mathbb{R}} u_0(x) \varphi(0, x) dx = 0$$

for any test function $\varphi \in C_c^\infty([0, \infty) \times \mathbb{R})$, we say that u is a *weak solution* of (8). Note that this is essentially a return to the more fundamental version (7), and so the weak solution should indeed be consistent with the underlying physics. Under the formation of a shock information is lost in the sense that we can find $u_1 \neq u_2$ such that $S(t)u_1 = S(t)u_2$. Now, it may be checked that (ii) is still valid when applied to weak solutions, but this implies a contradiction, yielding the fact that weak solutions of the reversed problem cannot be unique. Hence, weak solutions to (8) are in general not unique.

In (ii) we saw that reversing the direction of the flow provided a way to reverse the dynamics. Another illustration of the fact that the dynamics governed by (8) seems to be reversible is given by the following observation:

- (iii) **Reverseability.** Whenever f is symmetric, reversing the sign of u provides a reversal of the dynamics. That is, under the assumption that $f(u) = f(-u)$ it follows that $v(t, x) = -u(t_0 - t, x)$ is a solution of (8) with initial condition $-u(t_0)$.

The observation (iii) is valid for the inviscid burgers equation, $f(u) = \frac{1}{2}u^2$. Note that in this case, u models the momentum of a fluid. In fact, a similar observation holds true for the Euler equations as well. This is in contrast to the fact that the system which one seeks to model is the archetype of an irreversible processes. If one, after taking observations (i), (ii) and (iii) into account, is still inclined to apply (8) as a model, one is forced to separate the weak solutions representing the physics of the model from those that are mathematical artifacts.

Typically, in applications, equation (8) is an approximation of the problem:

$$\partial_t u^\varepsilon + \partial_x f(u^\varepsilon) = \varepsilon \partial_x^2 u^\varepsilon, \quad u^\varepsilon(0, x) = u_0(x), \quad (10)$$

where $\varepsilon > 0$ is some small number. In Burgers equation, the term $\varepsilon \partial_x^2 u^\varepsilon$ is naturally considered to represent the effect of viscosity, and so (10) is referred to as *the viscous approximation* of (8). The Cauchy problem (10) has a unique smooth solution. Furthermore, the system is irreversible, in contrast to (8) as observed in (ii) and (iii). From a physical perspective, this is easily understood as a consequence of the dissipation of mechanical energy into heat due to friction. To quantify the dissipation, let $S \in C^2(\mathbb{R})$ be a convex function, and let $Q' = S' f'$.

¹Changing the sign of f amounts to change the direction of the flow.

Multiplying (10) by $S'(u^\varepsilon)$ it follows by the chain-rule that

$$\begin{aligned}\partial_t S(u^\varepsilon) + \partial_x Q(u^\varepsilon) &= \varepsilon S'(u^\varepsilon) \partial_x^2 u^\varepsilon \\ &= \varepsilon \partial_x^2 S(u^\varepsilon) - \varepsilon S''(u^\varepsilon) (\partial_x u^\varepsilon)^2.\end{aligned}$$

Formally, assuming that both $u^\varepsilon(t, x) \rightarrow 0$ and $\partial_x u^\varepsilon(t, x) \rightarrow 0$ as $x \rightarrow \pm\infty$ sufficiently fast, we conclude that

$$\partial_t \int_{\mathbb{R}} S(u^\varepsilon(t, x)) dx = -\varepsilon \int_{\mathbb{R}} S''(u^\varepsilon(t, x)) (\partial_x u^\varepsilon(t, x))^2 dx \leq 0.$$

Note that the inequality is in fact strict for any nontrivial example and a suitably chosen S . Suppose there exist some u such that $u^\varepsilon \rightarrow u$ as $\varepsilon \downarrow 0$. What happens with this inequality as $\varepsilon \downarrow 0$? Wherever u is smooth, the term clearly vanishes as $\partial_x u$ remains bounded. At a shock, however, the term remains and strict inequality is preserved. That is, the limit u continues to dissipate “energy”. Inspired by this, we introduce the admissibility criterion

$$\int_0^\infty \int_{\mathbb{R}} S(u) \partial_t \varphi + Q(u) \partial_x \varphi dx dt + \int_{\mathbb{R}} S(u_0(x)) \varphi(0, x) dx \geq 0 \quad (11)$$

for any nonnegative $\varphi \in C_c^\infty([0, \infty) \times \mathbb{R})$. The pair (S, Q) is called an *entropy, entropy-flux pair*, and (11) is referred to as the *entropy inequality*. Now, one may check that in the presence of a strict inequality, i.e., after the formation of a shock, the reversed dynamics described in (ii) and (iii) does not satisfy the entropy inequality. If u satisfies the entropy inequality for any entropy, entropy-flux pair (S, Q) and any nonnegative φ , it is said to be an *entropy solution* of (8). Regarding integrability conditions, one generally consider a final time $T > 0$ and search for entropy solutions in the space $L^\infty(\Pi_T) \cap C([0, T]; L^1(\mathbb{R}))$ where $\Pi_T = [0, T] \times \mathbb{R}$.

Doubling-of-variables

Is it possible to conclude from the entropy inequality (11) that the entropy solution is unique? Let me give an outline of the “doubling of variables” argument given in [23]. At first, let me give the standard presentation of this argument. As is often the case, this is somehow the story told backwards. As a remedy, I’ll try to tell the story forward afterwards. Let me introduce the *Kruřkov entropies*

$$S(u) = |u - c|, \quad Q(u) = \text{sign}(u - c) (f(u) - f(c)), \quad c \in \mathbb{R}.$$

Any entropy solution u to (8) with initial condition u_0 satisfies

$$\int_0^\infty \int_{\mathbb{R}} |u - c| \partial_t \varphi + \text{sign}(u - c) (f(u) - f(c)) \partial_x \varphi dx dt + \int_{\mathbb{R}} |u_0(x) - c| \varphi(0, x) dx \geq 0, \quad (12)$$

for any $c \in \mathbb{R}$ and any nonnegative $\varphi \in C_c^\infty([0, \infty) \times \mathbb{R})$. Solutions characterized by this particular family of entropies are often referred to as *Kruřkov entropy solutions*. Let v be another Kruřkov entropy solution to (8) with initial condition v^0 . Now, we take u and v to be functions of different sets of variables, say $u = u(t, x)$ and $v = v(s, y)$. Let $\varphi = \varphi(t, x, s, y)$ be a

nonnegative compactly supported smooth function. Let $c = v(s, y)$ in (12). Integration in (s, y) yields

$$\begin{aligned} & \iiint_{\Pi^2} |u(t, x) - v(s, y)| \partial_t \varphi \, dx dt dy ds \\ & + \iiint_{\Pi^2} \text{sign}(u(t, x) - v(s, y)) (f(u(t, x)) - f(v(s, y))) \partial_x \varphi \, dx dt dy ds \\ & + \iint_{\Pi} \int_{\mathbb{R}} |u_0(x) - v(s, y)| \varphi(0, x, s, y) \, dx dy ds \geq 0, \end{aligned}$$

where $\Pi = [0, \infty) \times \mathbb{R}$. Similarly, as v is a Kruřkov entropy solution,

$$\begin{aligned} & \iiint_{\Pi^2} |v(s, y) - u(t, x)| \partial_s \varphi \, dx dt dy ds \\ & + \iiint_{\Pi^2} \text{sign}(v(s, y) - u(t, x)) (f(v(s, y)) - f(u(t, x))) \partial_y \varphi \, dx dt dy ds \\ & + \iint_{\Pi} \int_{\mathbb{R}} |v_0(y) - u(t, x)| \varphi(t, x, 0, y) \, dy dx dt \geq 0. \end{aligned}$$

Adding the two inequalities, we arrive at

$$\mathcal{T}_1 + \mathcal{T}_2 + \mathcal{T}_3 \geq 0, \quad (13)$$

where

$$\begin{aligned} \mathcal{T}_1 &:= \iiint_{\Pi^2} |u(t, x) - v(s, y)| (\partial_t + \partial_s) \varphi \, dx dt dy ds, \\ \mathcal{T}_2 &:= \iiint_{\Pi^2} \text{sign}(u(t, x) - v(s, y)) (f(u(t, x)) - f(v(s, y))) \partial_x \varphi \, dx dt dy ds \\ &+ \iiint_{\Pi^2} \text{sign}(v(s, y) - u(t, x)) (f(v(s, y)) - f(u(t, x))) \partial_y \varphi \, dx dt dy ds, \\ \mathcal{T}_3 &:= \iint_{\Pi} \int_{\mathbb{R}} |u_0(x) - v(s, y)| \varphi(0, x, s, y) \, dx dy ds \\ &+ \iint_{\Pi} \int_{\mathbb{R}} |v_0(y) - u(t, x)| \varphi(t, x, 0, y) \, dy dx dt. \end{aligned}$$

The next step is to pick φ in a clever way. Let J denote a symmetric mollifier on $\mathbb{R}$, i.e.,

$$J \geq 0, \quad \text{supp}(J) \subset (-1, 1), \quad \int J(\sigma) \, d\sigma = 1, \quad J(-\sigma) = J(\sigma). \quad (14)$$

Given $r > 0$, we let $J_r(\sigma) = \frac{1}{r} J(\frac{\sigma}{r})$. Take

$$\varphi(t, x, s, y) = \rho_\gamma \left(\frac{t+s}{2} \right) J_{r_0}(t-s) J_r(x-y), \quad \text{where } \rho_\gamma(t) = 1 - \int_0^t J_\gamma(\hat{t} - \tau) \, d\hat{t}. \quad (15)$$

Note that φ is not compactly supported, however one may check that all the integrals are indeed well defined. Next we want to send $r_0 \downarrow 0$. Note that

$$(\partial_t + \partial_s)\varphi(t, x, s, y) = \rho'_\gamma \left(\frac{t+s}{2} \right) J_{r_0}(t-s) J_r(x-y).$$

Hence, one may show that

$$\lim_{r_0 \downarrow 0} \mathcal{T}_1 = - \int_0^\infty \int_{\mathbb{R}} \int_{\mathbb{R}} |u(t, x) - v(t, y)| J_\gamma(t-\tau) J_r(x-y) dx dy dt.$$

Next we observe that

$$\text{sign}(u-v)(f(u) - f(v)) = \text{sign}(v-u)(f(v) - f(u)).$$

Furthermore $(\partial_x + \partial_y)\varphi = 0$. Hence $\mathcal{T}_2 = 0$. Concerning $\mathcal{T}_3$, one may show that

$$\lim_{r_0 \downarrow 0} \mathcal{T}_3 = \int_{\mathbb{R}} \int_{\mathbb{R}} |u_0(x) - v_0(y)| J_r(x-y) dx dy.$$

Upon sending $\gamma \downarrow 0$ it follows by (13) and the above limits that

$$\int_{\mathbb{R}} \int_{\mathbb{R}} |u(\tau, x) - v(\tau, y)| J_r(x-y) dx dy \leq \int_{\mathbb{R}} \int_{\mathbb{R}} |u_0(x) - v_0(y)| J_r(x-y) dx dy. \quad (16)$$

A first consequence of (16) is obtained by letting $r \downarrow 0$. This yields

$$\|u(t) - v(t)\|_{L^1} \leq \|u_0 - v_0\|_{L^1},$$

which is the L^1 -contraction property. Hence uniqueness of Kruřkov entropy solutions follows. A second consequence of (16) is the (fractional) BV regularity. Let $v = u$ in (17) and $x = y + z$. It follows that

$$\int_{\mathbb{R}} \|u(t, \cdot + z) - u(t, \cdot)\|_{L^1} J_r(z) dz \leq \int_{\mathbb{R}} \|u_0(\cdot + z) - u_0(\cdot)\|_{L^1} J_r(z) dz.$$

Next we try to tell the story forwards. The goal is to uncover the main “mechanics” of the proof as well as making it clear that the choice of the test function φ above is indeed rather obvious. To achieve this, the exposition is divided into two steps. First we assume that u, v are smooth entropy solutions of (8). Next, we see what happens when we try to rewrite this approach in its weak form. Let S be a symmetric convex function and take

$$\Psi(t, s) = \iint_{\mathbb{R} \times \mathbb{R}} S(u(t, x) - v(s, y)) J_r(x-y) dx dy.$$

We want to show that

$$\frac{d}{dt} \Psi(t, t) = \frac{d}{dt} \iint_{\mathbb{R} \times \mathbb{R}} S(u(t, x) - v(t, y)) J_r(x-y) dx dy \leq 0. \quad (17)$$

Given that u, v is smooth, the chain rule yields

$$\frac{d}{dt}\Psi(t, t) = [\partial_t \Psi(t, s) + \partial_s \Psi(t, s)]_{s=t}. \quad (18)$$

As u is a smooth entropy solution,

$$\partial_t S(u - c) + \partial_x Q(u, c) \leq 0, \quad \text{where } Q(u, c) = \int_c^u S'(z - c) f'(z) dz,$$

for any constant $c \in \mathbb{R}$. Hence

$$\begin{aligned} \partial_t \Psi(t, s) &= \iint_{\mathbb{R} \times \mathbb{R}} \partial_t S(u(t, x) - v(s, y)) J_r(x - y) dx dy \\ &\leq \iint_{\mathbb{R} \times \mathbb{R}} Q(u(t, x), v(s, y)) \partial_x J_r(x - y) dx dy. \end{aligned}$$

Similarly,

$$\partial_s \Psi(t, s) \leq \iint_{\mathbb{R} \times \mathbb{R}} Q(v(s, y), u(t, x)) \partial_y J_r(x - y) dx dy.$$

Suppose that Q is symmetric, i.e. $Q(u, v) = Q(v, u)$. Then it follows that

$$\partial_t \Psi(t, s) + \partial_s \Psi(t, s) \leq \iint_{\mathbb{R} \times \mathbb{R}} Q(u(t, x), v(s, y)) (\partial_x + \partial_y) J_r(x - y) dx dy = 0.$$

Hence, (17) follows given that Q is symmetric. In general, integration by parts yields

$$Q(u, v) = S'(u - v)(f(u) - f(v)) - \int_v^u S''(z - v)(f(z) - f(v)) dz.$$

The first term is symmetric, but the second is not. Letting $S \rightarrow |\cdot|$ the second term vanishes. This yields (17) with $S = |\cdot|$. How do we relax the assumption that u and v are smooth? That is, we want to apply the entropy inequality in its weak form (11). The essential step is to find a weak formulation of (18). Instead of $\frac{d}{dt}\Psi(t, t) \leq 0$, we want to show its weak equivalent

$$\int_0^\infty \Psi(t, t) \rho'(t) dt + \Psi(0, 0) \rho(0) \geq 0 \quad (19)$$

for any nonnegative $\rho \in C_c^\infty([0, \infty))$. This inequality yields (16) upon letting $\rho \rightarrow \mathbb{1}_{[0, \tau]}$. Observe that

$$\begin{aligned} \Psi(t, t) \rho'(t) &= \left[\Psi(t, s) (\partial_t + \partial_s) \rho \left(\frac{t + s}{2} \right) \right]_{s=t} \\ &= \lim_{r_0 \downarrow 0} \int_0^T \Psi(t, s) (\partial_t + \partial_s) \left(\rho \left(\frac{t + s}{2} \right) J_{r_0}(t - s) \right) ds. \end{aligned}$$

Hence, (19) follows if

$$\lim_{r_0 \downarrow 0} \iiint_{\Pi^2} S(u(t, x) - v(s, y)) (\partial_t + \partial_s) \varphi(t, x, s, y) dx dy ds dt$$

$$+ \int_{\mathbb{R}} \int_{\mathbb{R}} S(u_0(x) - v_0(y)) \rho(0) J_r(x - y) dx dy \geq 0,$$

where

$$\varphi(t, x, s, y) = \rho\left(\frac{t+s}{2}\right) J_{r_0}(t-s) J_r(x-y).$$

As we have already seen, this is exactly what we may conclude from the entropy inequalities (11) provided that Q is symmetric. Hence the choice $S = |\cdot|$.

The kinetic formulation

Let us once more turn back to the characterization of the entropy solution as the limit of the viscous approximation $\{u^\varepsilon\}_{\varepsilon>0}$ as $\varepsilon \downarrow 0$. Recall that after multiplying (10) by $S'(u^\varepsilon)$ we obtained

$$\partial_t S(u^\varepsilon) + \partial_x Q(u^\varepsilon) = \varepsilon \partial_x^2 S(u^\varepsilon) - \varepsilon S''(u^\varepsilon) (\partial_x u^\varepsilon)^2. \quad (20)$$

The key to the characterization of the limit was seen to be the fact that the square term on the right did not vanish as $\varepsilon \downarrow 0$. By choosing S to be convex, the term is negative, and this fact turns out to be sufficient for the characterization of the limit. The key point is that by studying the dynamics of nonlinear functions applied to the solution, we may single out the limit $u = \lim_{\varepsilon \downarrow 0} u^\varepsilon$ among other weak solutions to (8). The kinetic formulation provides an alternative description of these dynamics. Observe that

$$S(u) - S(0) = \int_{\mathbb{R}} S'(\xi) \chi(u; \xi) d\xi, \quad \text{where } \chi(u; \xi) = \begin{cases} 1, & \text{for } 0 < \xi < u, \\ -1, & \text{for } u < \xi < 0, \\ 0, & \text{otherwise.} \end{cases}$$

Applying this identity to (20), we arrive at

$$\begin{aligned} \partial_t \int_{\mathbb{R}} S'(\xi) \chi(u^\varepsilon; \xi) d\xi + \partial_x \int_{\mathbb{R}} S'(\xi) f'(\xi) \chi(u^\varepsilon; \xi) d\xi = \\ \varepsilon \partial_x^2 \int_{\mathbb{R}} S'(\xi) \chi(u^\varepsilon; \xi) d\xi - \varepsilon S''(u^\varepsilon) (\partial_x u^\varepsilon)^2. \end{aligned}$$

Furthermore, we note that

$$\varepsilon S''(u^\varepsilon) (\partial_x u^\varepsilon)^2 = \varepsilon \int_{\mathbb{R}} S''(\xi) \delta(u^\varepsilon - \xi) (\partial_x u^\varepsilon)^2 d\xi.$$

Assume $S(0) = 0$ such that $S(u_0) = \int_{\mathbb{R}} S'(\xi) \chi(u_0; \xi) d\xi$. Let $\varphi \otimes S'(t, x, \xi) = \varphi(t, x) S'(\xi)$. Then

$$\begin{aligned} \iiint_{\Pi \times \mathbb{R}} \chi(u^\varepsilon; \xi) \partial_t (\varphi \otimes S') + f'(\xi) \chi(u^\varepsilon; \xi) \partial_x (\varphi \otimes S') dx dt d\xi \\ + \iint_{\mathbb{R} \times \mathbb{R}} \chi(u_0; \xi) (\varphi \otimes S')(0, x, \xi) dx d\xi \end{aligned}$$

$$\begin{aligned}
&= -\varepsilon \iiint_{\Pi \times \mathbb{R}} \chi(u^\varepsilon; \xi) \partial_x^2(\varphi \otimes S') dx dt d\xi \\
&\quad + \varepsilon \iiint_{\Pi \times \mathbb{R}} \delta(u^\varepsilon - \xi) (\partial_x u^\varepsilon)^2 \partial_\xi(\varphi \otimes S') dx dt d\xi,
\end{aligned}$$

where $\Pi = [0, \infty) \times \mathbb{R}$ and $\varphi \in C_c^\infty(\Pi)$. Suppose there exists some function $u = u(t, x)$ and a measure $m = m(t, x, \xi)$ such that

$$\chi(u^\varepsilon; \cdot) \rightarrow \chi(u; \cdot) \quad \text{and} \quad \varepsilon \delta(u^\varepsilon - \cdot) (\partial_x u^\varepsilon)^2 \rightarrow m$$

as $\varepsilon \downarrow 0$. Then $\chi(u; \cdot)$ satisfies

$$\begin{cases} \partial_t \chi(u(t, x); \xi) + f'(\xi) \partial_x \chi(u(t, x); \xi) = \partial_\xi m(t, x, \xi), \\ \chi(u(0, x); \xi) = \chi(u_0(x); \xi), \end{cases} \quad (21)$$

in the sense of distributions. The crucial sign property, yielding the entropy inequality, is naturally expressed through the condition that the measure m is nonnegative. That is, u is an entropy solution of (8) given that u satisfies (21) for some nonnegative measure m [32, Theorem 3.2.1]. The measure m is often referred to as the *kinetic entropy defect measure*. Let me argue why the kinetic formulation may be considered as a weak formulation of the Kruřkov entropy condition. To this end, suppose $u \in L^\infty$ and let $S \in C_c^2(\mathbb{R})$ be convex on the region $|\xi| \leq \|u\|_\infty$. Replace c by ξ in (12), multiply by $S''(\xi)$ and integrate in ξ . This yields

$$\begin{aligned}
&\iiint_{\Pi \times \mathbb{R}} S''(\xi) |u - \xi| \partial_t \varphi + S''(\xi) \text{sign}(u - \xi) (f(u) - f(\xi)) \partial_x \varphi dx dt d\xi \\
&\quad + \iint_{\mathbb{R} \times \mathbb{R}} S''(\xi) |u_0(x) - \xi| \varphi(0, x) dx d\xi \geq 0.
\end{aligned}$$

Integration by parts yields

$$\begin{aligned}
&\iiint_{\Pi \times \mathbb{R}} \text{sign}(u - \xi) \partial_t(\varphi \otimes S') + \text{sign}(u - \xi) f'(\xi) \partial_x(\varphi \otimes S') dx dt d\xi \\
&\quad + \iint_{\mathbb{R} \times \mathbb{R}} \text{sign}(u_0(x) - \xi) (\varphi \otimes S')(0, x, \xi) dx d\xi \geq 0.
\end{aligned}$$

Next we note that

$$\text{sign}(u - \xi) = 2\chi(u; \xi) - \text{sign}(\xi). \quad (22)$$

Hence

$$\begin{aligned}
&2 \iiint_{\Pi \times \mathbb{R}} \chi(u; \xi) \partial_t(\varphi \otimes S') + \chi(u; \xi) f'(\xi) \partial_x(\varphi \otimes S') dx dt d\xi \\
&\quad + 2 \iint_{\mathbb{R} \times \mathbb{R}} \chi(u_0; \xi) (\varphi \otimes S')(0, x, \xi) dx d\xi - \underbrace{\iint_{\mathbb{R} \times \mathbb{R}} \text{sign}(\xi) (\varphi \otimes S')(0, x, \xi) dx d\xi}_{2S(0)\|\varphi(0, \cdot)\|_{L^1(\mathbb{R})}} \geq 0.
\end{aligned}$$

Assuming $S(0) = 0$, we recover the left hand side of the kinetic formulation. However, the explicit description of the dissipation is lost in place of a sign property relating to the convexity of S . There are many interesting aspects related to this new formulation. Among them is a second path to uniqueness of entropy solutions. Let me give an outline of the proof. The first observation is that for any $u, v \in \mathbb{R}$,

$$\int_{\mathbb{R}} Q(u, v; \xi) d\xi = |u - v|,$$

where

$$Q(u, v; \xi) = |\chi(u; \xi)| + |\chi(v; \xi)| - 2\chi(u; \xi)\chi(v; \xi).$$

This is a simple consequence of the equality

$$|\chi(u - v; \xi - v)| = (\chi(u; \xi) - \chi(v; \xi))^2.$$

Suppose u, v are entropy solutions of (8) with defect measures m_u and m_v respectively. Then

$$\partial_t |u - v| = \int_{\mathbb{R}} \text{sign}(\xi) \partial_t \chi(u; \xi) d\xi + \int_{\mathbb{R}} \text{sign}(\xi) \partial_t \chi(v; \xi) d\xi - 2 \int_{\mathbb{R}} \partial_t (\chi(u; \xi) \chi(v; \xi)) d\xi \quad (23)$$

Here we applied the fact that $|\chi(u; \xi)| = \text{sign}(\xi) \chi(u; \xi)$. Applying this and the fact that u, v satisfies (21),

$$\begin{aligned} \iint_{\mathbb{R} \times \mathbb{R}} \text{sign}(\xi) \partial_t \chi(u; \xi) dx d\xi &= \iint_{\mathbb{R} \times \mathbb{R}} \text{sign}(\xi) \partial_\xi m_u(t, x, \xi) dx d\xi, \\ \iint_{\mathbb{R} \times \mathbb{R}} \text{sign}(\xi) \partial_t \chi(v; \xi) dx d\xi &= \iint_{\mathbb{R} \times \mathbb{R}} \text{sign}(\xi) \partial_\xi m_v(t, x, \xi) dx d\xi. \end{aligned}$$

Concerning the product term, we multiply the equation for $\chi(u; \xi)$ by $\chi(v; \xi)$ and vice versa. Adding up yields

$$\partial_t (\chi(u; \xi) \chi(v; \xi)) + f'(\xi) \partial_x (\chi(u; \xi) \chi(v; \xi)) = \chi(v; \xi) \partial_\xi m_u + \chi(u; \xi) \partial_\xi m_v.$$

Integration in x and ξ yields

$$\iint_{\mathbb{R} \times \mathbb{R}} \partial_t (\chi(u; \xi) \chi(v; \xi)) dx d\xi = \iint_{\mathbb{R} \times \mathbb{R}} \chi(v; \xi) \partial_\xi m_u dx d\xi + \iint_{\mathbb{R} \times \mathbb{R}} \chi(u; \xi) \partial_\xi m_v dx d\xi.$$

By (23) we conclude that

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} |u - v| dx &= \iint_{\mathbb{R} \times \mathbb{R}} (\text{sign}(\xi) - 2\chi(v; \xi)) \partial_\xi m_u(t, x, \xi) dx d\xi \\ &\quad + \iint_{\mathbb{R} \times \mathbb{R}} (\text{sign}(\xi) - 2\chi(u; \xi)) \partial_\xi m_v(t, x, \xi) dx d\xi. \end{aligned}$$

By (22)

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} |u - v| \, dx &= -2 \iint_{\mathbb{R} \times \mathbb{R}} \delta(\xi - v) m_u(t, x, \xi) \, dx d\xi \\ &\quad - 2 \iint_{\mathbb{R} \times \mathbb{R}} \delta(\xi - u) m_v(t, x, \xi) \, dx d\xi \leq 0. \end{aligned}$$

A slightly different approach reveals some similarities with the Kruřkov technique. Applying (22) and Leibniz rule directly to (23), we arrive at

$$\partial_t |u - v| = \int_{\mathbb{R}} \text{sign}(\xi - v) \partial_t \chi(u; \xi) \, d\xi + \int_{\mathbb{R}} \text{sign}(\xi - u) \partial_t \chi(v; \xi) \, d\xi, \quad (24)$$

which is natural to compare with (18). In fact, a simple doubling argument combined with the basic property of the χ function yields (24) directly. To this end, observe that

$$\begin{aligned} \partial_t |u - v| &= \left[\partial_t |u(t) - v(s)| + \partial_s |v(s) - u(t)| \right]_{s=t} \\ &= \left[\partial_t \int_{\mathbb{R}} \text{sign}(\xi - v(s)) \chi(u(t); \xi) \, d\xi + \partial_s \int_{\mathbb{R}} \text{sign}(\xi - u(t)) \chi(v(s); \xi) \, d\xi \right]_{s=t}. \end{aligned}$$

As u, v satisfies (21), we conclude from (24)

$$\begin{aligned} \partial_t |u - v| &= - \int_{\mathbb{R}} f'(\xi) (\text{sign}(\xi - v) \partial_x \chi(u; \xi) + \text{sign}(\xi - u) \partial_x \chi(v; \xi)) \, d\xi \\ &\quad + \int_{\mathbb{R}} \text{sign}(\xi - v) \partial_\xi m_u \, d\xi + \int_{\mathbb{R}} \text{sign}(\xi - u) \partial_\xi m_v \, d\xi \\ &=: \mathcal{T}_1 + \mathcal{T}_2. \end{aligned}$$

Differentiating (22) we obtain $\partial_x \chi(u; \xi) = -\frac{1}{2} \partial_x \text{sign}(\xi - u)$. Hence it follows that

$$\mathcal{T}_1 = \frac{1}{2} \int_{\mathbb{R}} f'(\xi) \partial_x (\text{sign}(\xi - v) \text{sign}(\xi - u)) \, d\xi.$$

For $f \in C_0^1(\mathbb{R})$,

$$\frac{1}{2} \int_{\mathbb{R}} f'(\xi) \text{sign}(\xi - v) \text{sign}(\xi - u) \, d\xi = -\text{sign}(u - v) (f(u) - f(v)).$$

Hence we conclude that

$$\partial_t |u - v| + \partial_x (\text{sign}(u - v) (f(u) - f(v))) = \mathcal{T}_2.$$

Integration by parts yields $\mathcal{T}_2 \leq 0$ as above, and so the L^1 -contraction property follows upon integration in x .

Discussion of main ideas and techniques

The goal of the following section is to present the main ideas and techniques applied in the thesis. In doing so, certain choices are made. In particular, regarding Part II, questions of compactness and the application of Young measures are omitted.

Part I: Finite difference approximations of degenerate convection-diffusion equations

Let us return to the problem of finding a rate of convergence for finite difference approximations of degenerate convection-diffusion equations of the form (1). For the following discussion we restrict our attention to one spatial dimension. That is, we consider the Cauchy problem

$$\partial_t u + \partial_x f(u) = \partial_x^2 A(u), \quad u(0, x) = u_0(x), \quad (25)$$

where $A' \geq 0$. The accuracy of the approximation of a derivative by a finite difference is ultimately dependent on the regularity of the function. Consequently, regularity of solutions is expected to play a crucial role when considering convergence rates for finite difference approximations. In general, adding diffusion to the equation has an effect of smoothing the solution, and so one may expect that (25) should at least not be any more problematic than the conservation law (8). To gain some insight into the effect of degenerate diffusion terms, let us consider the degenerate elliptic problem

$$\frac{d^2}{dx^2} u^3(x) = 0 \quad \text{on } (-1, 1), \quad u(-1) = -1, u(1) = 1. \quad (26)$$

This amounts to study stationary solutions to (25) with $f = 0$ and $A(u) = u^3$. Letting $w = u^3$, it follows that $w''(x) = 0$ and $w(-1) = -1$ and $w(1) = 1$. Hence, $w(x) = x$ and so $u(x) = \text{sign}(x) \sqrt[3]{|x|}$ solves (26). Note however, $u'(x) \rightarrow \infty$ as $x \rightarrow 0$. In the case $A(u) = u^3$, A is mildly degenerate in the sense that $A'(u) = 0$ in a single point $u = 0$. In the case of strong degeneracy, i.e., $A'(u) = 0$ on an entire interval, solutions are in general discontinuous. Hence, one might expect that the degenerate diffusion term itself is responsible for certain jumps in the solution. For a concrete example of this, see [16]. Despite the lack of regularity of solutions for conservation laws, Kuznecov was able to show that finite difference schemes converge with rate $1/2$ [24]. To see how a degenerate diffusion term significantly complicates the situation, we start by taking a closer look at Carrillo's extension [9] of the Kruřkov theory to equations of the form (25). Consider the viscous approximation $\{u^\eta\}_{\eta>0}$ defined by

$$\partial_t u^\eta + \partial_x f(u^\eta) = \partial_x^2 A(u^\eta) + \eta \partial_x^2 u^\eta, \quad u^\eta(0, x) = u_0(x). \quad (27)$$

Multiply (27) by $\text{sign}(u - c)$. We have already seen how to treat the left hand side, so we restrict our attention to the diffusion terms. Regarding the first term on the right, for $A \in C^1(\mathbb{R})$,

$$\text{sign}(u^\eta - c) \partial_x A(u^\eta) = \text{sign}(A(u^\eta) - A(c)) \partial_x A(u^\eta). \quad (28)$$

This is a consequence of the fact that if $A'(u^\eta) = 0$, both sides are zero, and if $A'(u^\eta) > 0$, then $\text{sign}(u^\eta - c) = \text{sign}(A(u^\eta) - A(c))$. Consequently

$$\text{sign}(u^\eta - c) \partial_x^2 A(u^\eta) = \partial_x^2 |A(u) - A(c)| - \text{sign}'(A(u^\eta) - A(c)) (\partial_x A(u^\eta))^2.$$

The last term on the right yields

$$\eta \text{sign}(u^\eta - c) \partial_x^2 u^\eta = \eta \partial_x^2 |u^\eta - c| - \eta \text{sign}'(u^\eta - c) (\partial_x u^\eta)^2.$$

It follows that

$$\begin{aligned} \partial_t |u^\eta - c| + \partial_x (\text{sign}(u^\eta - c) (f(u^\eta) - f(c))) - \partial_x^2 |A(u) - A(c)| = \\ \eta \partial_x^2 |u^\eta - c| - \underbrace{\eta \text{sign}'(u^\eta - c) (\partial_x u^\eta)^2}_{\text{hyperbolic dissipation}} - \underbrace{\text{sign}'(A(u^\eta) - A(c)) (\partial_x A(u^\eta))^2}_{\text{parabolic dissipation}}. \end{aligned}$$

Upon sending $\eta \downarrow 0$ we obtain the entropy criterion

$$\partial_t |u - c| + \partial_x (\text{sign}(u - c) (f(u) - f(c))) - \partial_x^2 |A(u) - A(c)| \leq 0, \quad (29)$$

weakly on $[0, \infty) \times \mathbb{R}$. A fundamental fact regarding the uniqueness of entropy solutions is that the parabolic dissipation term is needed explicitly to arrive at the L^1 -contraction property. To see this, let v be another entropy solution of (25). Adding up the entropy conditions for $u = u(t, x)$ and $v = v(s, y)$ yields

$$(\partial_t + \partial_s) |u - v| + (\partial_x + \partial_y) (\text{sign}(u - v) (f(u) - f(v))) - (\partial_x^2 + \partial_y^2) |A(u) - A(v)| \leq 0. \quad (30)$$

However, $(\partial_x^2 + \partial_y^2)$ does not “reduce” to a single double derivative as $y \rightarrow x$. That is,

$$\partial_x^2 \Psi(x, x) = [(\partial_x + \partial_y)^2 \Psi(x, y)]_{x=y} = [(\partial_x^2 + 2\partial_x \partial_y + \partial_y^2) \Psi(x, y)]_{x=y}$$

for any smooth $\Psi = \Psi(x, y)$. Observe that

$$-2\partial_x \partial_y |A(u) - A(v)| = 2\text{sign}'(A(u) - A(v)) \partial_x A(u) \partial_y A(v). \quad (31)$$

Hence, to arrive at a “complete” double derivative on the left hand side, we need to add (31) to (30). At this point, it becomes clear that the explicit form of the parabolic dissipation term is needed. Replace (29) by

$$\begin{aligned} \partial_t |u - c| + \partial_x (\text{sign}(u - c) (f(u) - f(c))) - \partial_x^2 |A(u) - A(c)| \leq \\ - \text{sign}'(A(u) - A(c)) (\partial_x A(u))^2. \end{aligned}$$

Proceeding as above we arrive at

$$\begin{aligned} (\partial_t + \partial_s) |u - v| + (\partial_x + \partial_y) (\text{sign}(u - v) (f(u) - f(v))) \\ - (\partial_x^2 + 2\partial_x \partial_y + \partial_y^2) |A(u) - A(v)| \leq \mathcal{D}, \end{aligned}$$

where

$$\mathcal{D} = -\text{sign}'(A(u) - A(v)) \underbrace{\left((\partial_x A(u))^2 - 2\partial_x A(u)\partial_y A(v) + (\partial_y A(v))^2 \right)}_{(\partial_x A(u) - \partial_y A(v))^2} \leq 0.$$

To obtain a rate of convergence for finite difference approximations, we need to replicate the above argument as closely as possible. Since the fundamental difficulties stem from the diffusion term, we assume that $f = 0$ in the following discussion. As (28) does not have a natural analogue for finite differences, we instead work with $A^\eta(u) := A(u) + \eta u$. The goal is to find an error estimate that does not blow up in the limit $\eta \downarrow 0$. Let $\Delta x > 0$ and consider the semidiscrete finite difference scheme

$$\frac{d}{dt} U_j^\eta = \frac{A^\eta(U_{j-1}^\eta) - 2A^\eta(U_j^\eta) + A^\eta(U_{j+1}^\eta)}{\Delta x^2}, \quad U_j^\eta(0) = U_j^0, \quad (32)$$

for all $j \in \mathbb{Z}$. Let the piecewise constant approximation $u_{\Delta x}^\eta$ associated with $\{U_j^\eta\}$ be defined by

$$u_{\Delta x}^\eta(t, x) = U_j^\eta(t), \text{ whenever } x \in I_j := [(j - 1/2)\Delta x, (j + 1/2)\Delta x).$$

Let us introduce the difference quotients

$$D_+ u(x) = \frac{u(x + \Delta x) - u(x)}{\Delta x} \quad \text{and} \quad D_- u(x) = \frac{u(x) - u(x - \Delta x)}{\Delta x},$$

for any $u : \mathbb{R} \rightarrow \mathbb{R}$. In this notation (32) reads

$$\partial_t u_{\Delta x}^\eta = D_- D_+ A^\eta(u_{\Delta x}^\eta), \quad u_{\Delta x}^\eta(0, x) = u_{\Delta x}^0(x), \quad (33)$$

where $u_{\Delta x}^0(x) = U_j^0$ whenever $x \in I_j$. To simplify the notation, we suppress the η -dependence from the following calculations, and instead assume $A' > 0$. Following the above strategy, we multiply (33) by $\text{sign}(u_{\Delta x} - c)$. Since A is strictly increasing,

$$\partial_t |u_{\Delta x} - c| = \text{sign}(A(u_{\Delta x}) - A(c)) D_+ D_- A(u_{\Delta x}).$$

As a substitute for the chain-rule we introduce some new functions. By a misuse of notation, we let

$$\begin{aligned} D_+ |A(u_{\Delta x}) - A(c)| &= \text{sign}(A(\theta_{\Delta x}^+) - A(c)) D_+ A(u_{\Delta x}), \\ D_+ \text{sign}(A(u_{\Delta x}) - A(c)) &= \text{sign}'(A(\tau_{\Delta x}^+) - A(c)) D_+ A(u_{\Delta x}). \end{aligned}$$

Analogously we replace $\theta_{\Delta x}^+$ by $\theta_{\Delta x}^-$ when D_- replaces D_+ . The same applies to $\tau_{\Delta x}^-$. Note that the notation would be valid for continuously differentiable functions by the mean value theorem. In our case, the above may be taken as a definition. To be more specific, let $S_{\Delta x}$ denotes a translation, i.e. $S_{\Delta x} u(x) = u(x + \Delta x)$ for any $u = u(x)$. Then

$$\text{sign}(A(\theta_{\Delta x}^+) - A(c)) := \begin{cases} \frac{|A(S_{\Delta x} u_{\Delta x}) - A(c)| - |A(u_{\Delta x}) - A(c)|}{A(S_{\Delta x} u_{\Delta x}) - A(u_{\Delta x})}, & A(S_{\Delta x} u_{\Delta x}) \neq A(u_{\Delta x}), \\ 0 & \text{otherwise.} \end{cases}$$

Similar considerations apply to $\theta_{\Delta x}^-, \tau_{\Delta x}^+$ and $\tau_{\Delta x}^-$. By Leibniz rule for difference quotients,

$$D_-(\text{sign}(A(S_{\Delta x}u_{\Delta x}) - A(c)) D_+A(u_{\Delta x})) = \text{sign}'(A(\tau_{\Delta x}^+) - A(c)) (D_+A(u_{\Delta x}))^2 \\ + \text{sign}(A(u_{\Delta x}) - A(c)) D_-D_+A(u_{\Delta x}).$$

Similarly

$$D_+(\text{sign}(A(S_{-\Delta x}u_{\Delta x}) - A(c)) D_-A(u_{\Delta x})) = \text{sign}'(A(\tau_{\Delta x}^-) - A(c)) (D_-A(u_{\Delta x}))^2 \\ + \text{sign}(A(u_{\Delta x}) - A(c)) D_+D_-A(u_{\Delta x}).$$

Hence, we obtain

$$\text{sign}(A(u_{\Delta x}) - A(c)) D_-D_+A(u_{\Delta x}) = D_-D_+|A(u_{\Delta x}) - A(c)| + \mathcal{T}_1 \\ - \frac{1}{2}\text{sign}'(A(\tau_{\Delta x}^+) - A(c)) (D_+A(u_{\Delta x}))^2 \\ - \frac{1}{2}\text{sign}'(A(\tau_{\Delta x}^-) - A(c)) (D_-A(u_{\Delta x}))^2,$$

where

$$\mathcal{T}_1 = \frac{1}{2}D_- \left((\text{sign}(A(S_{\Delta x}u_{\Delta x}) - A(c)) - \text{sign}(A(\theta_{\Delta x}^+) - A(c))) D_+A(u_{\Delta x}) \right) \\ + \frac{1}{2}D_+ \left((\text{sign}(A(S_{-\Delta x}u_{\Delta x}) - A(c)) - \text{sign}(A(\theta_{\Delta x}^-) - A(c))) D_-A(u_{\Delta x}) \right).$$

We thus obtain the “entropy inequality”

$$\partial_t |u_{\Delta x} - c| - D_-D_+|A(u_{\Delta x}) - A(c)| = -\frac{1}{2}\text{sign}'(A(\tau_{\Delta x}^+) - A(c)) (D_+A(u_{\Delta x}))^2 \\ - \frac{1}{2}\text{sign}'(A(\tau_{\Delta x}^-) - A(c)) (D_-A(u_{\Delta x}))^2 + \mathcal{T}_1. \quad (34)$$

Let $u = u(s, y)$ denote the solution to (25) with $f = 0$. Then, as above,

$$\partial_s |u - c| - \partial_y^2 |A(u) - A(c)| = -\text{sign}'(A(u) - A(c)) (\partial_y A(u))^2. \quad (35)$$

Next we establish a replacement for (31) by exchanging the x -derivatives with finite differences. This yields

$$\partial_y(D_+ + D_-)|A(u_{\Delta x}) - A(u)| = -\text{sign}'(A(\tau_{\Delta x}^+) - A(u)) \partial_y A(u) D_+A(u_{\Delta x}) \\ - \text{sign}'(A(\tau_{\Delta x}^-) - A(u)) \partial_y A(u) D_-A(u_{\Delta x}). \quad (36)$$

Let $c = u(s, y)$ in (34) and $c = u_{\Delta x}(t, x)$ in (35). Add the resulting equations together with (36). Completing the squares we arrive at

$$(\partial_t + \partial_s) |u_{\Delta x} - u| - (D_-D_+ + \partial_y(D_+ + D_-) + \partial_y^2) |A(u_{\Delta x}) - A(u)|$$

$$\begin{aligned}
&= -\frac{1}{2} \text{sign}'(A(\tau_{\Delta x}^+) - A(c)) (D_+ A(u_{\Delta x}) - \partial_y A(u))^2 \\
&\quad - \frac{1}{2} \text{sign}'(A(\tau_{\Delta x}^-) - A(c)) (D_- A(u_{\Delta x}) - \partial_y A(u))^2 \\
&\quad + \mathcal{T}_1 + \mathcal{T}_2 \\
&\leq \mathcal{T}_1 + \mathcal{T}_2,
\end{aligned}$$

where

$$\begin{aligned}
\mathcal{T}_2 &= -\frac{1}{2} \left(\text{sign}'(A(u) - A(u_{\Delta x})) - \text{sign}'(A(u) - A(\tau_{\Delta x}^+)) \right) (\partial_y A(u))^2 \\
&\quad - \frac{1}{2} \left(\text{sign}'(A(u) - A(u_{\Delta x})) - \text{sign}'(A(u) - A(\tau_{\Delta x}^-)) \right) (\partial_y A(u))^2 \\
&=: \mathcal{T}_2^+ + \mathcal{T}_2^-.
\end{aligned}$$

Let

$$\mathcal{T}_3 = (D_- D_+ + \partial_y (D_+ + D_-) + \partial_y^2) |A(u_{\Delta x}) - A(u)|,$$

so that

$$(\partial_t + \partial_s) |u_{\Delta x} - u| \leq \mathcal{T}_1 + \mathcal{T}_2 + \mathcal{T}_3.$$

Let φ be the Kruřkov test function defined in (15). To obtain a rate of convergence we need to find upper bounds on

$$\iiint_{\Pi^2} \mathcal{T}_i(t, x, s, y) \varphi \, ds dy dt dx, \quad (i = 1, 2, 3),$$

in terms of $\Delta x, r, r_0, \eta$. Let me give a heuristic outline of how the term $\mathcal{T}_2$ is treated. In particular, let me argue why this term separates the case of spatial dimension $d = 1$ from the case $d > 1$. Consider $\mathcal{T}_2^+$, the term $\mathcal{T}_2^-$ is treated similarly. For the sake of simplicity, let us assume that $\tau_{\Delta x}^+$ is actually some function satisfying

$$\min\{S_{\Delta x} u_{\Delta x}, u_{\Delta x}\} \leq \tau_{\Delta x}^+ \leq \max\{S_{\Delta x} u_{\Delta x}, u_{\Delta x}\},$$

as would be the case if $\sigma \mapsto \text{sign}(\sigma)$ was continuously differentiable. Then

$$\int_{\mathbb{R}} |\tau_{\Delta x}^+ - u_{\Delta x}| \, dx \leq \int_{\mathbb{R}} |S_{\Delta x} u_{\Delta x} - u_{\Delta x}| \, dx \leq \|D_+ u_{\Delta x}\|_{L^1(\mathbb{R})} \Delta x.$$

Therefore it is natural to estimate the term by establishing a modulus of continuity for the parabolic dissipation term as a function of the Kruřkov constant. That is, we wish to establish a modulus of continuity for the mapping

$$c \mapsto \iint_{\Pi} \text{sign}'(A(u) - A(c)) (\partial_y A(u))^2 \varphi \, dy ds =: \zeta(c).$$

To do this, we return to (35). Thus

$$\zeta(c) = \iint_{\Pi} |u - c| \partial_s \varphi \, dy ds + \int_{\mathbb{R}} |u_0 - c| \varphi|_{s=0} \, dy + \iint_{\Pi} |A(u) - A(c)| \partial_y^2 \varphi \, dy ds.$$

It follows that

$$\begin{aligned} |\zeta(c_2) - \zeta(c_1)| &\leq \int_{c_1}^{c_2} |\zeta'(\xi)| \, d\xi \\ &\leq \int_{c_1}^{c_2} \left| \iint_{\Pi} \text{sign}(u - \xi) \partial_s \varphi \, dy ds \right| d\xi \\ &\quad + \int_{c_1}^{c_2} \left| \int_{\mathbb{R}} \text{sign}(u_0 - \xi) \varphi|_{s=0} \, dy \right| d\xi \\ &\quad + \int_{c_1}^{c_2} \left| \iint_{\Pi} \text{sign}(A(u) - A(\xi)) A'(\xi) \partial_y^2 \varphi \, dy ds \right| d\xi \\ &\leq \left(\|\partial_s \varphi\|_{L^1(\Pi_{s,y})} + \|\varphi|_{s=0}\|_{L^1(\mathbb{R}_y)} + \|A'\|_{\infty} \|\partial_y^2 \varphi\|_{\Pi_{s,y}} \right) |c_2 - c_1|. \end{aligned}$$

This is the essential step towards a rate of convergence in one spatial dimension. In several space dimensions, the error $\mathcal{T}_2$ splits into a sum of directional components. Therefore the $\mathcal{T}_2$ is not expressed through the continuity of parabolic dissipation. As a remedy we have to depend on the regularization of the Kruřkov entropy for the term $\mathcal{T}_2$ to vanish. That is, the regularization parameter will have to move slowly to zero compared with Δx in order for $\mathcal{T}_2$ to vanish in the limit $\Delta x \downarrow 0$. As a consequence, the rate is lowered considerably for dimension $d > 1$. A second observation regarding the estimation of $\mathcal{T}_2$ is, loosely speaking, that the kinetic formulation appears naturally. This intuition is based on the view that the kinetic formulation may be regarded as a weak form of the Kruřkov entropy condition, formally obtained by differentiating the entropy condition with respect to the Kruřkov constant.

Finally, let me mention a possible continuation of the work. As we see from the above discussion, the fundamental difficulties stems from the lack of a chainrule for finite differences. One might wonder if it is possible to construct schemes which enables a more accurate replication of the chain rule. Candidates for such approximations may be the higher order schemes considered in [29].

Part II: Stochastic scalar conservation laws

In the following discussion we restrict our attention to stochastic balance laws of the form

$$\partial_t u + \partial_x f(u) = \sigma(u) \partial_t W(t), \quad u(0, x) = u_0(x). \quad (37)$$

Here $W = W(t, \omega)$, $\omega \in \Omega$ is a Brownian motion on the probability space $(\Omega, \mathcal{F}, P)$ endowed with its natural filtration $\{\mathcal{F}_t\}_{t \geq 0}$. The coefficients $f = f(u)$ and $\sigma = \sigma(u)$ are given deterministic Lipschitz continuous functions. Regarding well-posedness, the following discussion is limited to the question of uniqueness, thus focusing on the extension of the “doubling of variables” technique. Along these lines, the discussion of the operator-splitting (5) is limited to the question of a rate of convergence. In what follows, computations are carried through in the

strong form. However, as we have seen, “doubling of variables” is suitable for weak formulations, and as such the below computations and formulations have natural weak counterparts.

Extension of the method of “doubling of variables” to stochastic balance laws

We have already seen how “doubling of variables” enables us to handle the flux term. So to arrive at the heart of the matter, let us instead consider how “doubling of variables” may be used to derive uniqueness of solutions to the stochastic differential equation

$$du(t) = \sigma(u(t)) dW(t), \quad u(0) = u_0. \quad (38)$$

Let u and v be solutions with initial condition u_0 and v_0 respectively. Let $S \in C^2(\mathbb{R})$ be a symmetric convex function. In the strong form, “doubling of variables” is based on the equality

$$\frac{d}{dt}\Psi(t, t) = [\partial_t \Psi(t, s) + \partial_s \Psi(t, s)]_{s=t}, \text{ where } \Psi(t, s) = S(u(t) - v(s)). \quad (39)$$

Furthermore, we recall that to handle equations of the form (37) we need to let $S \rightarrow |\cdot|$ due to the treatment of the flux term. To gain insight into what we should obtain from the right hand side of (39), let us compute the left hand side directly. By Itô’s formula

$$\begin{aligned} \frac{d}{dt}S(u(t) - v(t)) &= S'(u(t) - v(t))(\sigma(u(t)) - \sigma(v(t))) \dot{W}(t) \\ &\quad + \frac{1}{2}S''(u(t) - v(t))(\sigma(u(t)) - \sigma(v(t)))^2. \end{aligned} \quad (40)$$

Upon letting $S \rightarrow |\cdot|$ and taking the expectation, this yields

$$E[|u(t) - v(t)|] = E[|u_0 - v_0|],$$

from which uniqueness follows. Is it possible to obtain the right hand side of (40) by computing the right hand side of (39)? Following Kružkov, we compute

$$\begin{aligned} \frac{d}{dt}S(u(t) - c) &= S'(u(t) - c)\sigma(u(t))\dot{W}(t) + \frac{1}{2}S''(u(t) - c)(\sigma(u(t)))^2, \\ \frac{d}{ds}S(v(s) - c) &= S'(v(s) - c)\sigma(v(s))\dot{W}(s) + \frac{1}{2}S''(v(s) - c)(\sigma(v(s)))^2, \end{aligned}$$

for any constant $c \in \mathbb{R}$. Attempting to set $c = v(s)$ in the first equality, $c = u(t)$ in the second equality and inserting the resulting expressions into (39), we do not obtain the right hand side of (40). We are missing the term

$$-S''(u(t) - v(s))\sigma(u(t))\sigma(v(s)).$$

The problem stems from the fact that we cannot simply insert $c = v(s)$ in the Itô formula due to the fact that $v(s)$ is a random variable dependent on the Brownian motion up to time s . The same is true for the case $c = u(t)$. To see what happens when we insert a random variable instead of a constant, we take a closer look at the Itô formula. In order to do this, we need

some tools from Malliavin Calculus. Let us consider the Malliavin derivative as a directional derivative on the Wiener space $\Omega = C_0([0, T]) = \{\omega \in C([0, T]) : \omega(0) = 0\}$. This enables us to give some formal arguments, showing that the necessary relations are at least intuitively correct. For an introduction to the field, see for instance [31, 8, 13]. Let $H = L^2([0, T])$ with inner product $\langle h, g \rangle_H = \int_0^T h(t)g(t) dt$. Given a functional $F : \Omega \rightarrow \mathbb{R}$, we would like to define a derivative DF satisfying

$$\underbrace{\lim_{\varepsilon \downarrow 0} \frac{F(\omega + \varepsilon\gamma) - F(\omega)}{\varepsilon}}_{D_\gamma F(\omega)} = \langle DF(\omega), h \rangle_H, \quad \text{where} \quad \gamma(t) = \int_0^t h(s) ds,$$

for any $h \in H$. As a motivation for the above definition, let $F(\omega) = \int_0^T f(t) dW(t, \omega) = \int_0^T f(t) d\omega(t)$ for some $f \in H$. Then

$$\begin{aligned} D_\gamma F(\omega) &= \lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \left(\int_0^T f(s) d(w(s) + \varepsilon\gamma(s)) - \int_0^T f(s) dw(s) \right) \\ &= \int_0^T f(s) d\gamma(s) = \int_0^T f(s) h(s) ds. \end{aligned}$$

Hence $DF = f$. Regarding the domain of D , an extension along the same lines as the classical extension to weak derivatives is possible. However, for the sake of simplicity all random variables will be assumed to be in the domain of D . Let δ be the adjoint of D . That is, for a process $u = u(t, \omega)$ we let

$$E[\langle DF, u \rangle_H] = E[F\delta(u)] \text{ for all suitable } F = F(\omega).$$

It turns out that δ coincides with an extension of the Itô integral known as the Skorohod integral. For our purposes the following property is crucial:

$$F \int_0^T u(s) dW(s) = \underbrace{\int_0^T Fu(s) dW(s)}_{\text{Skorohod}} + \int_0^T D_s F u(s) ds, \quad (41)$$

where $D_s F$ denotes the Malliavin derivative of F evaluated at s . This is a simple consequence of the Leibniz rule for the Malliavin derivative. To this end, observe that for any suitable random variable $G = G(\omega)$,

$$\begin{aligned} E[GF\delta(u)] &= E[\langle D(GF), u \rangle_H] = E[\langle DG, Fu \rangle_H] + E[G \langle DF, u \rangle_H] \\ &= E[G\delta(Fu)] + E[G \langle DF, u \rangle_H]. \end{aligned}$$

Let us return to the Itô formula. Let V be Malliavin differentiable and $t_i^n := \frac{it}{2^n}$. By Taylor's formula,

$$S(u(t) - V) - S(u(0) - V) = \sum_{i=0}^{2^n-1} S'(u(t_i^n) - V)(u(t_{i+1}^n) - u(t_i^n))$$

$$\begin{aligned}
& + \frac{1}{2} \sum_{i=0}^{2^n-1} S''(\theta_i^n - V)(u(t_{i+1}^n) - u(t_i^n))^2 \\
& = \mathcal{J}_1^n + \mathcal{J}_2^n,
\end{aligned}$$

where θ_i^n is some intermediate point between $u(t_i^n)$ and $u(t_{i+1}^n)$. Now,

$$\mathcal{J}_2^n \rightarrow \frac{1}{2} \int_0^t S''(u(s) - V)(\sigma(u(s)))^2 ds$$

as $n \rightarrow \infty$. By (41)

$$\begin{aligned}
\mathcal{J}_1^n &= \sum_{i=0}^{2^n-1} S'(u(t_i^n) - V) \int_{t_i^n}^{t_{i+1}^n} \sigma(u(s)) dW(s) \\
&= \sum_{i=0}^{2^n-1} \int_{t_i^n}^{t_{i+1}^n} S'(u(t_i^n) - V) \sigma(u(s)) dW(s) \\
&\quad - \sum_{i=0}^{2^n-1} \int_{t_i^n}^{t_{i+1}^n} S''(u(t_i^n) - V) D_s V \sigma(u(s)) ds.
\end{aligned}$$

Here we applied the chain rule and the fact that $D_s u(t) = 0$ whenever $t < s$ to conclude that

$$D_s S'(u(t_i^n) - V) = -S''(u(t_i^n) - V) D_s V.$$

Hence

$$\mathcal{J}_1^n \rightarrow \int_0^t S'(u(s) - V) \sigma(u(s)) dW(s) - \int_0^t S''(u(s) - V) D_s V \sigma(u(s)) ds,$$

where the stochastic integral is interpreted as a Skorohod integral. With this Itô formula at hand, we define entropy solutions of (38) to be a Malliavin differentiable and adapted process u satisfying

$$E \left[\frac{d}{dt} S(u(t) - V) \right] = E \left[-S''(u(t) - V) D_t V \sigma(u(t)) + \frac{1}{2} S''(u(t) - V) (\sigma(u(t)))^2 \right], \quad (42)$$

for any Malliavin differentiable random variable V and any symmetric convex S in $C^2(\mathbb{R})$. Suppose $u = u(t)$ and $v = v(s)$ are entropy solutions to (38) with initial conditions u^0 and v^0 respectively. Applying the entropy condition (42) for both u and v it follows that

$$\begin{aligned}
& E [\partial_t \Psi(t, s) + \partial_s \Psi(t, s)] \\
&= E \left[S''(u(t) - v(s)) \underbrace{\left(\frac{1}{2} \sigma^2(u(t)) - D_t v(s) \sigma(u(t)) - D_s u(t) \sigma(v(s)) + \frac{1}{2} \sigma^2(v(s)) \right)}_{\mathcal{T}} \right].
\end{aligned}$$

Let us rewrite $\mathcal{T}$ according to

$$\mathcal{J} = \frac{1}{2} (\sigma(u(t)) - \sigma(v(s)))^2 + \underbrace{(\mathbb{1}_{t < s} \sigma(v(s)) - D_t v(s)) \sigma(u(t))}_{\text{zero for } s < t} + \underbrace{(\mathbb{1}_{s < t} \sigma(u(t)) - D_s u(t)) \sigma(v(s))}_{\text{zero for } t < s}.$$

Hence

$$E [\partial_t \Psi(t, s) + \partial_s \Psi(t, s)] = \mathcal{J}_1 + \mathcal{J}_2 + \mathcal{J}_3,$$

where

$$\begin{aligned}\mathcal{J}_1 &= \frac{1}{2} E \left[S''(u(t) - v(s)) (\sigma(u(t)) - \sigma(v(s)))^2 \right], \\ \mathcal{J}_2 &= E \left[S''(u(t) - v(s)) (\mathbb{1}_{t < s} \sigma(v(s)) - D_t v(s)) \sigma(u(t)) \right], \\ \mathcal{J}_3 &= E \left[S''(u(t) - v(s)) (\mathbb{1}_{s < t} \sigma(u(t)) - D_s u(t)) \sigma(v(s)) \right].\end{aligned}$$

Note that $\mathcal{J}_1$ is exactly what we are expecting from the direct calculation (40). Furthermore, it vanishes as $S \rightarrow |\cdot|$. Comparing with we (40), $\mathcal{J}_2$ and $\mathcal{J}_3$ should vanish as $s \rightarrow t$. In its weak form, the actual limit $s \rightarrow t$ is the essential limit

$$\lim_{r_0 \downarrow 0} \int_{\mathbb{R}} \mathcal{J}_i(t, s) J_{r_0}(s - t) ds, \quad i = 2, 3,$$

where J_{r_0} is a mollifier as defined in (14). An important observation is that the mollifier need not be symmetric. That is, we may assume $\text{supp}(J_{r_0}) \subset (0, r_0)$, which amounts to consider the essential limit as $s \downarrow t$. As $u(t)$ is $\mathcal{F}_t$ -adapted $D_s u(t) = 0$ whenever $t < s$. That is, if $u(t)$ is independent of the noise at time s , the derivative should be zero. Hence $\mathcal{J}_3(s, t) = 0$ whenever $s > t$ and so $\lim_{s \downarrow t} \mathcal{J}_3(s, t) = 0$. To see why $\lim_{s \downarrow t} \mathcal{J}_2(s, t) = 0$, we need expressions for $D_t v(s)$. Now,

$$\begin{aligned}D_\gamma v(s; \omega) &:= \lim_{\varepsilon \downarrow 0} \frac{v(s; \omega + \varepsilon \gamma) - v(s; \omega)}{\varepsilon} \\ &= D_\gamma v^0 + \lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \left(\int_0^s \sigma(v(r; \omega + \varepsilon \gamma)) d(\omega + \varepsilon \gamma)(r) - \int_0^s \sigma(r; \omega) d\omega(r) \right) \\ &= D_\gamma v^0 + \lim_{\varepsilon \downarrow 0} \left(\int_0^s \sigma(v(r; \omega + \varepsilon \gamma)) d\gamma(r) + \int_0^s \frac{\sigma(v(r; \omega + \varepsilon \gamma)) - \sigma(v(r; \omega))}{\varepsilon} d\omega(r) \right) \\ &= D_\gamma v^0 + \int_0^s \sigma(v(r; \omega)) d\gamma(r) + \int_0^s D_\gamma(\sigma(v(r; \omega))) d\omega(r).\end{aligned}$$

By the chain rule $D_\gamma(\sigma(v(r; \omega))) = \sigma'(v(r; \omega)) D_\gamma v(r; \omega)$. Hence the Malliavin derivative satisfies

$$\langle Dv(s), h \rangle_H = \langle Dv^0, h \rangle_H + \langle \mathbb{1}_{[0, s]} \sigma(v), h \rangle_H + \int_0^s \sigma'(v(r)) \langle Dv(r), h \rangle_H d\omega(r).$$

For $t < s$ the above equation yields

$$D_t v(s) = \sigma(v(t)) + \int_t^s \sigma'(v(r)) D_t v(r) dW(r).$$

Hence

$$\sigma(v(s)) - D_t v(s) = (\sigma(v(s)) - \sigma(v(t))) + \int_t^s \sigma'(v(r)) D_t v(r) dW(r),$$

from which we may conclude that $\lim_{s \downarrow t} \mathcal{T}_3(s, t) = 0$ essentially. Hence, we conclude

$$\begin{aligned} \partial_t E[S(u(t) - v(t))] &= \lim_{s \downarrow t} E[\partial_t \Psi(t, s) + \partial_s \Psi(t, s)] \\ &= \lim_{s \downarrow t} \frac{1}{2} E \left[S''(u(t) - v(s)) (\sigma(u(t)) - \sigma(v(s)))^2 \right]. \end{aligned}$$

Note that the latter term vanishes as $S \rightarrow |\cdot|$ independently of s and t . Let me point out a fundamental observation regarding the above technique: We need at least one of the solutions to be Malliavin differentiable. Which one depends on whether we take the limit $s \downarrow t$ or $s \uparrow t$. When we add the flux term, this causes a fundamental difficulty. Formally $D_t v(s)$ should satisfy

$$\partial_s D_t v + \nabla \cdot (f'(v) D_t v) = (\sigma'(v) D_t v) \partial_s W(s) \quad \text{for } s > t$$

and $D_t v(s) = \sigma(v(s))$. However, even in the deterministic setting ($\sigma' = 0$), the well-posedness of such equations is a subtle issue due to the discontinuity of $f'(v)$. To cope with the situation, one may instead replace one of the solutions by its viscous approximation. That is, we replace v by v^ε where

$$\partial_s v^\varepsilon + \nabla \cdot f(v^\varepsilon) = \sigma(v^\varepsilon) \partial_t W + \varepsilon \Delta v^\varepsilon, \quad v^\varepsilon(0, x) = v_0(x).$$

Letting $s \downarrow t$ before we let $\varepsilon \downarrow 0$ allows us to handle the term corresponding to $\mathcal{T}_2$.

Even though the kinetic formulation of (37) is not applied in any of the works of this thesis, it is the technique applied in several related works [12, 19]. Indeed, this technique seems more easily adapted to the stochastic setting. Let me therefore, for comparison, include a short outline of how the kinetic formulation may be applied to derive uniqueness of (38). As above, let u and v be solutions to (38). By Itô's formula

$$dS(u) = S'(u) \sigma(u) dW + \frac{1}{2} S''(u) \sigma^2(u) dt,$$

for any $S \in C^2(\mathbb{R})$. Applying the χ -function we derive the “kinetic formulation”

$$d\chi(u; \xi) = \delta(u - \xi) \sigma(u) dW - \frac{1}{2} \partial_\xi \delta(u - \xi) \sigma^2(u) dt,$$

in distributional sense. As in (23)

$$d|u - v| = \int_{\mathbb{R}} \underbrace{\text{sign}(\xi) d\chi(u; \xi) + \text{sign}(\xi) d\chi(v; \xi) - 2d(\chi(u; \xi) \chi(v; \xi))}_{dQ(u, v; \xi)} d\xi$$

By Leibniz rule

$$d(\chi(u; \xi)\chi(v; \xi)) = \chi(u; \xi)d\chi(v; \xi) + \chi(v; \xi)d\chi(u; \xi) + [\chi(u; \xi), \chi(v; \xi)] dt, \quad (43)$$

where $[\cdot, \cdot]$ denotes the cross-variation process. That is,

$$[\chi(u; \xi), \chi(v; \xi)] = \delta(u - \xi)\delta(v - \xi)\sigma(u)\sigma(v).$$

Hence, by (22)

$$\begin{aligned} dQ(u, v; \xi) &= \text{sign}(\xi - v) d\chi(u; \xi) + \text{sign}(\xi - u) d\chi(v; \xi) - 2\delta(u - \xi)\delta(v - \xi)\sigma^2(\xi) dt \\ &= \text{sign}(\xi - v) \delta(u - \xi)\sigma(u)dW - \frac{1}{2}\text{sign}(\xi - v) \partial_\xi \delta(u - \xi)\sigma^2(u)dt \\ &\quad + \text{sign}(\xi - u) \delta(v - \xi)\sigma(v)dW - \frac{1}{2}\text{sign}(\xi - u) \partial_\xi \delta(v - \xi)\sigma^2(v)dt \\ &\quad - 2\delta(u - \xi)\delta(v - \xi)\sigma(u)\sigma(v) dt \end{aligned}$$

Upon taking expectation and integrating in ξ

$$\begin{aligned} E[d|u - v|] &= -\frac{1}{2}E \left[\int_{\mathbb{R}} \text{sign}(\xi - v) \partial_\xi \delta(u - \xi)\sigma^2(u) dt d\xi \right] \\ &\quad - \frac{1}{2}E \left[\int_{\mathbb{R}} \text{sign}(\xi - u) \partial_\xi \delta(v - \xi)\sigma^2(v) dt d\xi \right] \\ &\quad - 2E \left[\int_{\mathbb{R}} \delta(u - \xi)\delta(v - \xi)\sigma(u)\sigma(v) dt d\xi \right] \end{aligned}$$

Integrating by parts and completing the square yields

$$E[d|u - v|] = E \left[\int_{\mathbb{R}} \underbrace{\delta(u - \xi)\delta(v - \xi)}_{0 \text{ for } u \neq v} \underbrace{(\sigma(u) - \sigma(v))^2}_{0 \text{ for } u = v} d\xi \right] = 0.$$

Loosely speaking, the sought cross term, which is obtained via the Malliavin derivative, appears directly as the cross-variation process via the application of Leibniz rule in (43).

A rate of convergence

We continue with an outline of the error estimate for the operator splitting established in Paper IV. Recall that $\mathcal{S}_{\text{CL}}$ denotes the solution operator associated with the entropy solution of the conservation law (8), while $\mathcal{S}_{\text{SDE}}$ denotes the solution operator associated with the solution of the stochastic differential equation $du = \sigma(u)dW$. Let $T > 0$ be some fixed final time. Divide $[0, T]$ into N parts of length Δt . The approximation $\{u^n\}_{n=0}^N$ is defined by

$$u^0 = u_0 \quad \text{and} \quad u^{n+1} = \mathcal{S}_{\text{SDE}}(t_{n+1}, t_n)\mathcal{S}_{\text{CL}}(\Delta t)u^n \quad \text{for } n = 0, \dots, N-1.$$

To obtain an error estimate, we adopt the above method of “doubling of variables” by replacing one of the solutions by the scheme $\{u^n\}$. However, to do this we need to pick a time-interpolant of $\{u^n\}_{n=0}^N$. Let me point out that choosing the right one is crucial. Indeed there are many

different choices, and the form of the error terms is dependent on the choice of interpolant. Inspired by [25] we consider the interpolant

$$\eta_{\Delta t}(t) = (\mathcal{S}_{\text{SDE}}(t, t_n) - \mathcal{I})\mathcal{S}_{\text{CL}}(\Delta t)u^n + \mathcal{S}_{\text{CL}}(t - t_n)u^n, \quad t_n \leq t \leq t_{n+1},$$

where $\mathcal{I}$ denotes the identity. Furthermore, to simplify the notation we let

$$\begin{aligned} u_{\Delta t}(t) &:= \mathcal{S}_{\text{SDE}}(t, t_n)\mathcal{S}_{\text{CL}}(\Delta t)u^n, & t_n < t \leq t_{n+1}, \\ v_{\Delta t}(t) &:= \mathcal{S}_{\text{CL}}(t - t_n)u^n, & t_n \leq t < t_{n+1}. \end{aligned}$$

It follows that

$$\eta_{\Delta t} = u_{\Delta t} + v_{\Delta t} - \mathcal{S}_{\text{CL}}(\Delta t)u^n.$$

Formally $du_{\Delta t} = \sigma(u_{\Delta t})dW$ and $dv_{\Delta t} + \partial_x f(v_{\Delta t})dt = 0$, and so

$$d\eta_{\Delta t} + \partial_x f(v_{\Delta t})dt = \sigma(u_{\Delta t})dW, \quad t_n < t < t_{n+1},$$

and $\eta_{\Delta t}(t_n) = u^n$. Hence, the error estimate is basically a question of continuous dependence with respect to the coefficients. Let $v = v(t, x)$ be a solution to (37). Formally Itô's formula yields

$$\begin{aligned} dS(\eta_{\Delta t} - v) &= -S'(\eta_{\Delta t} - v)\partial_x(f(v_{\Delta t}) - f(v))dt + S'(\eta_{\Delta t} - v)(\sigma(u_{\Delta t}) - \sigma(v))dW \\ &\quad + \frac{1}{2}S''(\eta_{\Delta t} - v)(\sigma(u_{\Delta t}) - \sigma(v))^2 dt \end{aligned}$$

In light of the above “doubling of variables” argument, we may expect that this is what we should obtain given that we are able to send $s \rightarrow t$ independently of the other parameters. Separating out the error terms, and taking expectation we obtain

$$E[dS(\eta_{\Delta t} - v)] = \mathcal{T}_1 + \cdots + \mathcal{T}_4,$$

where

$$\begin{aligned} \mathcal{T}_1 &:= -E[S'(\eta_{\Delta t} - v)\partial_x(f(\eta_{\Delta t}) - f(v))dt], \\ \mathcal{T}_2 &:= \frac{1}{2}E[S''(\eta_{\Delta t} - v)(\sigma(\eta_{\Delta t}) - \sigma(v))^2 dt], \\ \mathcal{T}_3 &:= E[S'(\eta_{\Delta t} - v)\partial_x(f(\eta_{\Delta t}) - f(v_{\Delta t}))dt], \\ \mathcal{T}_4 &:= E\left[S''(\eta_{\Delta t} - v)\left(\int_{\eta_{\Delta t}}^{u_{\Delta t}} (\sigma(z) - \sigma(v))\sigma'(z)dz\right)dt\right]. \end{aligned}$$

Here we applied the identity

$$(\sigma(u_{\Delta t}) - \sigma(v))^2 - (\sigma(\eta_{\Delta t}) - \sigma(v))^2 = 2 \int_{\eta_{\Delta t}}^{u_{\Delta t}} (\sigma(z) - \sigma(v))\sigma'(z)dz.$$

Although the above computations are formal, they turn out to be highly suggestive. To obtain a rate of convergence it is essential to know how fast the different terms vanish. To achieve

this requires detailed estimates. Let us anyhow point out some essential features regarding $\mathcal{T}_1, \dots, \mathcal{T}_4$. $\mathcal{T}_1$ is a familiar term from the theory of conservation laws. To handle it we need to integrate in x and let $S \rightarrow |\cdot|$. In fact, for smooth $\eta_{\Delta t}$ and $v_{\Delta t}$

$$\mathcal{T}_1 = \partial_x \left(S'(\eta_{\Delta t} - v)(f(\eta_{\Delta t}) - f(v)) \right) dt \quad \text{when } S' = \text{sign}(\cdot).$$

Similarly $\mathcal{T}_2$ vanishes as $S \rightarrow |\cdot|$. However, this is not the case for the term $\mathcal{T}_4$. It vanishes as $\Delta t \downarrow 0$, but only under the assumption that S'' is bounded. Hence, we need to balance the speed with which $S \rightarrow |\cdot|$ with respect to Δt . Note that $\mathcal{T}_4$ is a consequence of the “extra square term” in the Itô formula. As such the need to balance the speed with which $S \rightarrow |\cdot|$ is fundamentally a stochastic phenomena, eventually leading to a lower rate in the stochastic setting. Concerning $\mathcal{T}_3$ we note that

$$\eta_{\Delta t} - v_{\Delta t} = u_{\Delta t} - \mathcal{S}_{\text{CL}}(\Delta t)u^n = \int_{t_n}^t \sigma(u_{\Delta t}(s)) dW(s).$$

Consequently

$$\partial_x f(\eta_{\Delta t}) - \partial_x f(v_{\Delta t}) = (f'(\eta_{\Delta t}) - f'(v_{\Delta t})) \partial_x v_{\Delta t} + f'(\eta_{\Delta t}) \int_{t_n}^t \partial_x \sigma(u_{\Delta t}(s)) dW(s).$$

Furthermore

$$E [|(f'(\eta_{\Delta t}) - f'(v_{\Delta t})) \partial_x v_{\Delta t}|] \leq \|f'\|_{\text{Lip}} E \left[E \left[\left| \int_{t_n}^t \sigma(u_{\Delta t}(s)) dW(s) \right| \middle| \mathcal{F}_{t_n} \right] |\partial_x v_{\Delta t}| \right]$$

Hence the integral of $\mathcal{T}_3$ should vanish as $\Delta t \downarrow 0$ due to the fact that $v_{\Delta t}(t)$ is of bounded variation.

References

- [1] C. Bauzet. On a time-splitting method for a scalar conservation law with a multiplicative stochastic perturbation and numerical experiments. *J. Evol. Equ.*, 14(2):333–356, 2014.
- [2] C. Bauzet, G. Vallet, and P. Wittbold. The Cauchy problem for conservation laws with a multiplicative stochastic perturbation. *J. Hyperbolic Differ. Equ.*, 9(4):661–709, 2012.
- [3] J. Bec and K. Khanin. Burgers turbulence. *Phys. Rep.*, 447(1-2):1–66, 2007.
- [4] A. Bensoussan, R. Glowinski, and A. Răşcanu. Approximation of some stochastic differential equations by the splitting up method. *Appl. Math. Optim.*, 25(1):81–106, 1992.
- [5] I. H. Biswas, K. H. Karlsen, and A. K. Majee. Conservation laws driven by Lévy white noise. *J. Hyperbolic Differ. Equ.*, 12(3):581–654, 2015.
- [6] R. Bürger and W. L. Wendland. Sedimentation and suspension flows: historical perspective and some recent developments. *J. Engrg. Math.*, 41(2-3):101–116, 2001. Sedimentation and suspension flows: some recent contributions (Stuttgart, 1999).

- [7] M. C. Bustos, F. Concha, R. Bürger, and E. M. Tory. *Sedimentation and thickening*, volume 8 of *Mathematical Modelling: Theory and Applications*. Kluwer Academic Publishers, Dordrecht, 1999. Phenomenological foundation and mathematical theory.
- [8] R. A. Carmona and M. R. Tehranchi. *Interest rate models: an infinite dimensional stochastic analysis perspective*. Springer Finance. Springer-Verlag, Berlin, 2006.
- [9] J. Carrillo. Entropy solutions for nonlinear degenerate problems. *Arch. Ration. Mech. Anal.*, 147(4):269–361, 1999.
- [10] G. Chen, Q. Ding, and K. H. Karlsen. On nonlinear stochastic balance laws. *Arch. Ration. Mech. Anal.*, 204(3):707–743, 2012.
- [11] G. Chen and B. Perthame. Well-posedness for non-isotropic degenerate parabolic-hyperbolic equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 20(4):645–668, 2003.
- [12] A. Debussche and J. Vovelle. Scalar conservation laws with stochastic forcing. *J. Funct. Anal.*, 259(4):1014–1042, 2010.
- [13] G. Di Nunno, B. Øksendal, and F. Proske. *Malliavin calculus for Lévy processes with applications to finance*. Universitext. Springer-Verlag, Berlin, 2009.
- [14] Weinan E, K. Khanin, A. Mazel, and Ya. Sinai. Invariant measures for Burgers equation with stochastic forcing. *Ann. of Math. (2)*, 151(3):877–960, 2000.
- [15] M. S. Espedal and K. H. Karlsen. Numerical solution of reservoir flow models based on large time step operator splitting algorithms. In *Filtration in Porous Media and Industrial Application*, volume 1734 of *Lecture Notes in Mathematics*, pages 9–77. Springer Berlin Heidelberg, 2000.
- [16] S. Evje and K. H. Karlsen. Monotone difference approximations of BV solutions to degenerate convection-diffusion equations. *SIAM J. Numer. Anal.*, 37(6):1838–1860 (electronic), 2000.
- [17] J. Feng and D. Nualart. Stochastic scalar conservation laws. *J. Funct. Anal.*, 255(2):313–373, 2008.
- [18] P. K. Friz and B. Gess. Stochastic scalar conservation laws driven by rough paths. *Annales de l’Institut Henri Poincaré (C) Non Linear Analysis*, 2015.
- [19] M. Hofmanová. Degenerate parabolic stochastic partial differential equations. *Stochastic Process. Appl.*, 123(12):4294–4336, 2013.
- [20] H. Holden and N. H. Risebro. *Front tracking for hyperbolic conservation laws*, volume 152 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 2002.
- [21] K. H. Karlsen, U. Koley, and N. H. Risebro. An error estimate for the finite difference approximation to degenerate convection-diffusion equations. *Numer. Math.*, 121(2):367–395, 2012.

- [22] J. U. Kim. On a stochastic scalar conservation law. *Indiana Univ. Math. J.*, 52(1):227–256, 2003.
- [23] S. N. Kružkov. First order quasilinear equations with several independent variables. *Mat. Sb. (N.S.)*, 81 (123):228–255, 1970.
- [24] N. N. Kuznecov. The accuracy of certain approximate methods for the computation of weak solutions of a first order quasilinear equation. *Ž. Vyčisl. Mat. i Mat. Fiz.*, 16(6):1489–1502, 1627, 1976.
- [25] J. O. Langseth, A. Tveito, and R. Winther. On the convergence of operator splitting applied to conservation laws with source terms. *SIAM J. Numer. Anal.*, 33(3):843–863, 1996.
- [26] P. D. Lax. *Hyperbolic systems of conservation laws and the mathematical theory of shock waves*. Society for Industrial and Applied Mathematics, Philadelphia, Pa., 1973. Conference Board of the Mathematical Sciences Regional Conference Series in Applied Mathematics, No. 11.
- [27] R. J. LeVeque. *Finite volume methods for hyperbolic problems*. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 2002.
- [28] P.-L. Lions, B. Perthame, and E. Tadmor. A kinetic formulation of multidimensional scalar conservation laws and related equations. *J. Amer. Math. Soc.*, 7(1):169–191, 1994.
- [29] Y. Liu, C. Shu, and M. Zhang. High order finite difference WENO schemes for nonlinear degenerate parabolic equations. *SIAM J. Sci. Comput.*, 33(2):939–965, 2011.
- [30] J. C. Muccino, W. G. Gray, and L. A. Ferrand. Toward an improved understanding of multiphase flow in porous media. *Reviews of Geophysics*, 36(3):401–422, 1998.
- [31] D. Nualart. *The Malliavin calculus and related topics*. Probability and its Applications (New York). Springer-Verlag, Berlin, second edition, 2006.
- [32] B. Perthame. *Kinetic formulation of conservation laws*, volume 21 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2002.
- [33] D. Serre. *Systems of conservation laws. I*. Cambridge University Press, Cambridge, 1999. Hyperbolicity, entropies, shock waves, Translated from the 1996 French original by I. N. Sneddon.
- [34] A. Szymkiewicz. *Modelling Water Flow in Unsaturated Porous Media*. GeoPlanet: Earth and Planetary Sciences. Springer Berlin Heidelberg, Berlin, Heidelberg, 2013.
- [35] G. Vallet and P. Wittbold. On a stochastic first-order hyperbolic equation in a bounded domain. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 12(4):613–651, 2009.

Papers

- Paper I** **Kenneth H. Karlsen, Nils Henrik Risebro and Erlend B. Storrøsten**
 $\mathbb{L}^1$ error estimates for difference approximations of degenerate convection-diffusion equations. *Mathematics of Computation* **Volume 83, Number 290, November 2014**, Pages 2717-2762.
- Paper II** **Kenneth H. Karlsen, Nils Henrik Risebro and Erlend B. Storrøsten**
On the convergence rate of finite difference methods for degenerate convection-diffusion equations in several space dimensions.
ESAIM:M2AN Forthcoming.
- Paper III** **Kenneth H. Karlsen and Erlend B. Storrøsten**
On stochastic conservation laws and Malliavin calculus.
Submitted
- Paper IV** **Kenneth H. Karlsen and Erlend B. Storrøsten**
Analysis of a splitting method for stochastic balance laws
Preprint

L^1 ERROR ESTIMATES FOR DIFFERENCE APPROXIMATIONS OF DEGENERATE CONVECTION-DIFFUSION EQUATIONS

K. H. KARLSEN, N. H. RISEBRO, AND E. B. STORRØSTEN

ABSTRACT. We analyze monotone finite difference schemes for strongly degenerate convection-diffusion equations in one spatial dimension. These nonlinear equations are well-posed within a class of (discontinuous) entropy solutions. We prove that the L^1 error between the approximate and exact solutions is $\mathcal{O}(\Delta x^{1/3})$, where Δx is the spatial grid parameter. This result should be compared with the classical $\mathcal{O}(\Delta x^{1/2})$ error estimate for conservation laws (Kuznecov, 1976), and a recent estimate of $\mathcal{O}(\Delta x^{1/11})$ for degenerate convection-diffusion equations (Karlsen, Koley, Risebro 2012).

1. INTRODUCTION

Nonlinear convection-dominated flow problems arise in a range of applications, such as fluid dynamics, meteorology, transport of oil and gas in porous media, electro-magnetism, as well as in many other applications. As a consequence it has become a very important undertaking to construct robust, accurate, and efficient methods for the numerical approximation of such problems. Over the years a large number of stable (convergent) numerical methods have been developed for linear and nonlinear convection-diffusion equations in which the “diffusion part” is small, or even vanishing, relative to the “convection part” of the equation. There is a large literature on this topic, and we will provide a few relevant references later.

One central but exceedingly difficult issue relating to numerical methods for convection-diffusion equations, is the derivation of (a priori) error estimates that are robust in the singular limit as the diffusion coefficient vanishes, avoiding the exponential growth of error constants. This problem has been resolved only partly in special situations, such as for linear equations or in the completely degenerate case of no diffusion (scalar conservation laws). For general nonlinear equations containing both convection and (degenerate) diffusion terms this is a long standing open problem in numerical analysis.

This paper makes a small contribution to this general problem by deriving an error estimate for a class of simple difference schemes for nonlinear and strongly degenerate convection-diffusion problems of the form

$$(1.1) \quad \begin{cases} \partial_t u + \partial_x f(u) = \partial_x^2 A(u), & (x, t) \in \Pi_T, \\ u(0, x) = u^0(x), & x \in \mathbb{R}, \end{cases}$$

Received by the editor May 4, 2012 and, in revised form, January 25, 2013.

2010 *Mathematics Subject Classification.* Primary 65M06, 65M15; Secondary 35K65, 35L65.

Key words and phrases. Degenerate convection-diffusion equations, entropy conditions, finite difference schemes, error estimates.

where $\Pi_T = \mathbb{R} \times (0, T)$ for some fixed final time $T > 0$, and $u(x, t)$ is the scalar unknown function that is sought. The initial function $u_0 : \mathbb{R} \rightarrow \mathbb{R}$ is a given integrable and bounded function, while the convection flux $f : \mathbb{R} \rightarrow \mathbb{R}$ and the diffusion function $A : \mathbb{R} \rightarrow \mathbb{R}$ are given functions satisfying

$$f, A \text{ locally } C^1; A(0) = 0; A \text{ nondecreasing.}$$

The moniker *strongly degenerate* means that we allow $A'(u) = 0$ for all u in some interval $[\alpha, \beta] \subset \mathbb{R}$. Thus, the class of equations becomes very general, including purely hyperbolic equations (scalar conservation laws),

$$(1.2) \quad \partial_t u + \partial_x f(u) = 0,$$

as well as nondegenerate (uniformly parabolic) equations, such as the heat equation $\partial_t u = \partial_x^2 u$, and point-degenerate diffusion equations, such as the heat equation with a power-law nonlinearity: $\partial_t u = \partial_x(u^m \partial_x u)$, which is degenerate at $u = 0$.

Whenever the problem (1.1) is uniformly parabolic (i.e., $A' \geq \eta$ for some $\eta > 0$), it is well known that the problem admits a unique classical (smooth) solution. On the other hand, in the strongly degenerate case, (1.1) must be interpreted in the weak sense to account for possibly discontinuous (shock wave) solutions. Regarding weak solutions, it turns out that one needs an additional condition, the so-called *entropy condition*, to ensure that (1.1) is well-posed. More precisely, the following is known: For $u_0 \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$, there exists a unique solution $u \in C((0, T); L^1(\mathbb{R}^d))$, $u \in L^\infty(\Pi_T)$ of (1.1) such that $\partial_x A(u) \in L^2(\Pi_T)$ and for all convex functions $S : \mathbb{R} \rightarrow \mathbb{R}$ with $q'_S = f'S'$ and $r'_S = A'S'$,

$$(1.3) \quad \partial_t S(u) + \partial_x q_S(u) - \partial_x^2 r_S(u) \leq 0 \quad \text{in the weak sense on } [0, T) \times \mathbb{R}.$$

The satisfaction of these inequalities for all convex S is the entropy condition, and a weak solution satisfying the entropy condition is called an entropy solution. The well-posedness of entropy solutions is a famous result due to Kružkov [21] for conservation laws (1.2), and a more recent work by Carrillo [5] extends this to degenerate parabolic equations (1.1). These results are available in the multi-dimensional context, and we refer to [1, 10] for an overview of the relevant literature. For uniqueness of entropy solutions in the *BV* class; see [26, 28].

One traditional way of constructing entropy solutions is by the vanishing viscosity method, which starts off from classical solutions to the nondegenerate equation

$$\partial_t u_\eta + \partial_x f(u_\eta) = \partial_x^2 A(u_\eta) + \eta \partial_x^2 u_\eta, \quad \eta > 0,$$

and establishes the strong convergence of u_η as $\eta \rightarrow 0$ by deriving *BV* estimates that are independent of η ; see Vol'pert and Hudjaev [27].

Besides proving that u_η converges in the L^1 norm to the unique entropy solution u of (1.1), it is possible to prove the error estimate

$$(1.4) \quad \|u_\eta(\cdot, t) - u(\cdot, t)\|_{L^1} \leq C \sqrt{\eta}, \quad (\text{whenever } u_0 \in BV);$$

see [14] (cf. also [15]). The error bound (1.4) can also be obtained as a consequence of the more general continuous dependence estimate derived in [9]; see also [6, 18].

Herein we are interested in the much more difficult problem of deriving error estimates for numerical approximations of entropy solutions to convection-diffusion equations. Convergence results (without error estimates) have been obtained for finite difference schemes [12] (see also [13, 19]); finite volume schemes [16] (see also [2]); operator splitting methods [17]; and BGK approximations [3, 4], to mention just a few references. For a posteriori estimates for finite volume schemes, see [24].

To be concrete in what follows, let us for simplicity assume $f' \geq 0$ and consider the semi-discrete difference scheme

$$(1.5) \quad \frac{d}{dt}u_j(t) + \frac{f(u_j) - f(u_{j-1})}{\Delta x} = \frac{A(u_{j+1}) - 2A(u_j) + A(u_{j-1}))}{\Delta x^2},$$

where $u_j(t) \approx u(t, j\Delta x)$ and $\Delta x > 0$ is the spatial mesh size. Convergence of this scheme can be proved as in the works [12, 13], where explicit and implicit time discretizations are treated. Denote by $u_{\Delta x}(x, t)$ the piecewise constant interpolation of $\{u_j(t)\}_j$. The basic question we address in this paper is the following one: Does there exist a number $r \in (0, 1)$ and a constant C , independent of Δx , such that

$$(1.6) \quad \|u_{\Delta x}(\cdot, t) - u(\cdot, t)\|_{L^1} \leq C \Delta x^r,$$

where u is the unique entropy solution of (1.1). We refer to the number r as the rate of convergence.

In the purely hyperbolic case (1.2) ($A' \equiv 0$), the answer to this question is a classical result due to Kuznetsov [22], who proved that the rate of convergence is $1/2$ for viscous approximations as well as monotone difference schemes, and this is optimal for discontinuous solutions. The work of Kuznetsov [22] turned out to be extremely influential, and by now a large number of related works have been devoted to error estimation theory for conservation laws. We refer to [7] for an overview of the relevant results and literature.

Unfortunately, the situation is unclear in the degenerate parabolic case (1.1). Let us expose some reasons why adding a nonlinear diffusion term to (1.2) can make the error analysis significantly more difficult than in the streamlined Kuznetsov theory. First of all, it is well known that the purely hyperbolic difference scheme

$$(1.7) \quad \frac{d}{dt}u_j(t) + \frac{f(u_j) - f(u_{j-1}))}{\Delta x} = 0$$

has as a model equation, the second order viscous equation

$$\partial_t u + \partial_x f(u) = \frac{\Delta x}{2} \partial_x^2 f(u),$$

an equation that is compatible with the notion of entropy solution for (1.2). Indeed, an error estimate for this viscous equation is highly suggestive for what to expect for the upwind scheme (1.7) (this is of course what Kuznetsov proved). However, for convection-diffusion equations such as (1.1) the situation changes. The model equation for (1.5) is no longer second order but rather fourth order:

$$\partial_t u + \partial_x f(u) = \partial_x^2 A(u) + \frac{\Delta x}{2} \partial_x^2 f(u) - \frac{\Delta x^2}{12} \partial_x^4 A(u);$$

hence the error estimate (1.4) appears no longer so relevant for numerical schemes. Another added difficulty comes from the necessity to work with an explicit form of the parabolic dissipation term associated with (1.1). Indeed, in the analysis one needs to replace (1.3) by the following more precise entropy equation [5],

$$(1.8) \quad \begin{aligned} & \partial_t |u - c| + \partial_x (\text{sign}(u - c)(f(u) - f(c))) - \partial_x^2 |A(u) - A(c)| \\ & = -\text{sign}'(A(u) - A(c)) |\partial_x A(u)|^2, \quad c \in \mathbb{R}, \end{aligned}$$

which is formally obtained multiplying (1.1) by $\text{sign}(A(u) - A(c))$, assuming for the sake of this discussion that $A'(\cdot) > 0$. The term on the right-hand side is the parabolic dissipation term, which is a finite (signed) measure and thus very singular. To illustrate why the parabolic dissipation term is needed, let $u(y, s)$ and

$v(x, t)$ be two solutions satisfying (1.8). In the entropy equation for $u(y, s)$ one takes $c = v(x, t)$, while in the entropy equation for $v(x, t)$ one takes $c = u(y, s)$. Adding the two resulting equations yields

$$\begin{aligned} & (\partial_t + \partial_s) |u - v| + (\partial_x + \partial_y) (\text{sign}(u - v)(f(u) - f(v))) \\ & - (\partial_x^2 + \partial_y^2) |A(u) - A(v)| = -\text{sign}'(A(u) - A(v)) (|\partial_y A(u)|^2 + |\partial_x A(v)|^2), \end{aligned}$$

By adding $-2\partial_{xy}^2 |A(u) - A(v)|$ to both sides of this equation, noting that

$$-2\partial_{xy}^2 |A(u) - A(v)| = 2\text{sign}'(A(u) - A(v)) \partial_y A(u) \partial_x A(v),$$

we arrive at

$$\begin{aligned} & (\partial_t + \partial_s) |u - v| + (\partial_x + \partial_y) (\text{sign}(u - v)(f(u) - f(v))) \\ & - (\partial_x^2 - 2\partial_{xy}^2 + \partial_y^2) |A(u) - A(v)| \\ (1.9) \quad & = -\text{sign}'(A(u) - A(v)) (|\partial_y A(u)| - |\partial_x A(v)|)^2 \\ & \leq 0, \end{aligned}$$

from which the contraction property $\frac{d}{dt} \|u(\cdot, t) - v(\cdot, t)\|_{L^1} \leq 0$ follows [5]. Similarly, to obtain error estimates for numerical methods, it is necessary to derive a “discrete” version of (1.9) with v replaced by $u_{\Delta x}$. The main challenge is to suitably replicate at the discrete level the delicate balance between the two terms in (1.9) involving A ; the difficulty stems from the lack of a chain rule for finite differences.

Despite the mentioned difficulties, in this paper we will prove that there exists a constant C , independent of Δx , such that for any $t > 0$,

$$\|u_{\Delta x}(\cdot, t) - u(\cdot, t)\|_{L^1} \leq C \Delta x^{\frac{1}{3}}.$$

The only other work we are aware of that provides L^1 error estimates for numerical approximations of (1.1) is [20]; therein (1.6) is established with $r = \frac{1}{11}$; if A is a linear function, then the convergence rate is the usual one, namely $r = \frac{1}{2}$. In addition to the semi-discrete scheme (1.7), we will prove similar results for fully discrete (implicit and explicit) difference schemes.

Roughly speaking, the reason is two-fold for why we can significantly improve the result in [20]. First, we are herein able to provide a more faithful analog of (1.9) at the discrete level. Second, since $\text{sign}'(\cdot)$ is singular, one has to work with a Lipschitz continuous approximation $\text{sign}_\varepsilon(\cdot)$ of the sign function $\text{sign}(\cdot)$. The use of this approximation breaks the symmetry of the corresponding entropy fluxes, and introduces new error terms that depend on the parameter ε ; the process of “balancing” terms involving Δx and ε lowers the convergence rate (to $r = \frac{1}{11}$) [20]. In the present paper we are able to dispense with this balancing act. Indeed, we show that it is possible to send $\varepsilon \rightarrow 0$ independently of Δx .

The remaining part of this paper is organized as follows: In Section 2 we list some relevant a priori estimates satisfied by viscous approximations and entropy solutions, and provide a definition of entropy solutions. The semi-discrete difference scheme is defined and proved to be well-posed in Section 3. We also list several relevant a priori estimates. Section 4 is devoted to the proof of the error estimate. In Section 5 we show that the proof in Section 4 can be adapted to a fully discrete scheme that is implicit in the time variable. In fact, we go through all the steps of the proof and provide the details where there are considerable differences between the two cases. In Section 6 the explicit version of the scheme is treated. We end the paper with a few concluding remarks in Section 7.

2. PRELIMINARY MATERIAL

Set $A^\eta(u) := A(u) + \eta u$ for any fixed $\eta > 0$, and consider the uniformly parabolic problem

$$(2.1) \quad \begin{cases} u_t^\eta + f(u^\eta)_x = A^\eta(u^\eta)_{xx}, & (x, t) \in \Pi_T, \\ u^\eta(x, 0) = u^0(x), & x \in \mathbb{R}. \end{cases}$$

It is well known that (2.1) admits a unique classical (smooth) solution.

We collect some relevant (standard) a priori estimates in the next three lemmas.

Lemma 2.1. *Suppose $u^0 \in L^\infty(\mathbb{R}) \cap L^1(\mathbb{R}) \cap BV(\mathbb{R})$, and let u^η be the unique classical solution of (2.1). Then for any $t > 0$,*

$$\begin{aligned} \|u^\eta(\cdot, t)\|_{L^1(\mathbb{R})} &\leq \|u^0\|_{L^1(\mathbb{R})}, \\ \|u^\eta(\cdot, t)\|_{L^\infty(\mathbb{R})} &\leq \|u^0\|_{L^\infty(\mathbb{R})}, \\ |u^\eta(\cdot, t)|_{BV(\mathbb{R})} &\leq |u^0|_{BV(\mathbb{R})}. \end{aligned}$$

For a proof of the previous and next lemmas, see for example [27].

Lemma 2.2. *Suppose $u^0 \in L^\infty(\mathbb{R}) \cap L^1(\mathbb{R})$ and $f(u^0) - A(u^0)_x \in BV(\mathbb{R})$. Let u^η be the unique classical solution of (2.1). Then for any $t_1, t_2 > 0$,*

$$\|u^\eta(\cdot, t_2) - u^\eta(\cdot, t_1)\|_{L^1(\mathbb{R})} \leq |f(u^0) - A(u^0)_x|_{BV(\mathbb{R})} |t_2 - t_1|.$$

Regarding the following lemma, see [12, 25].

Lemma 2.3. *Suppose $u^0 \in L^\infty(\mathbb{R}) \cap L^1(\mathbb{R})$ and $f(u^0) - A(u^0)_x \in L^\infty(\mathbb{R}) \cap BV(\mathbb{R})$. Let u^η be the unique classical solution of (2.1). Then for any $t > 0$,*

$$\begin{aligned} \|f(u^\eta(\cdot, t)) - A(u^\eta(\cdot, t))_x\|_{L^\infty(\mathbb{R})} &\leq \|f(u^0) - A(u^0)_x\|_{L^\infty(\mathbb{R})}, \\ |f(u^\eta(\cdot, t)) - A(u^\eta(\cdot, t))_x|_{BV(\mathbb{R})} &\leq |f(u^0) - A(u^0)_x|_{BV(\mathbb{R})}. \end{aligned}$$

Note that $\|A(u^\eta)_x\|_{L_t^\infty(L_x^\infty)}$ and $\|A(u^\eta)_{xx}\|_{L_t^\infty(L_x^1)}$ are bounded independently of η provided that $A(u^0)_x$ is in $BV(\mathbb{R})$.

The results above imply that $\{u^\eta\}_{\eta>0}$ is relatively compact in $C([0, T]; L_{\text{loc}}^1(\mathbb{R}))$. If $u = \lim_{\eta \rightarrow 0} u^\eta$, then

$$\|u^\eta - u\|_{L^1(\Pi_T)} \leq C\eta^{1/2},$$

for some constant C which does not depend on η ; see [14]. Moreover, u is an entropy solution according to the following definition.

Definition 2.1. An entropy solution of the Cauchy problem (1.1) is a measurable function $u = u(x, t)$ satisfying:

- (D.1) $u \in L^\infty(\Pi_T) \cap C((0, T); L^1(\mathbb{R}))$.
- (D.2) $A(u) \in L^2((0, T); H^1(\mathbb{R}))$.
- (D.3) For all constants $c \in \mathbb{R}$ and test functions $0 \leq \varphi \in C_0^\infty(\mathbb{R} \times [0, T])$, the following entropy inequality holds:

$$\begin{aligned} \iint_{\Pi_T} |u - c| \varphi_t + \text{sign}(u - c) (f(u) - f(c)) \varphi_x + |A(u) - A(c)| \varphi_{xx} dx dt \\ + \int_{\mathbb{R}} |u_0 - c| \varphi(x, 0) dx \geq 0. \end{aligned}$$

The uniqueness of entropy solutions follows from the work [5]. Actually, in view of the above a priori estimates, the relevant functional class is $BV(\Pi_T)$, in which case we can replace (D.2) by the condition $A(u)_x \in L^\infty(\Pi_T)$. For a uniqueness result in the BV class, see [28].

3. DIFFERENCE SCHEME

We start by specifying the numerical flux to be used in the difference scheme.

Definition 3.1 (Numerical flux). We call a function $F \in C^1(\mathbb{R}^2)$ a two-point numerical flux for f if $F(u, u) = f(u)$ for $u \in \mathbb{R}$. If

$$\frac{\partial}{\partial u} F(u, v) \geq 0 \quad \text{and} \quad \frac{\partial}{\partial v} F(u, v) \leq 0$$

holds for all $u, v \in \mathbb{R}$, we call F monotone.

Let F_u and F_v denote the partial derivatives of F with respect to the first and second variable, respectively. We will also assume that F is Lipschitz continuous.

Let $\Delta x > 0$ and set $x_j = j\Delta x$ for $j \in \mathbb{Z}$, and define

$$D_\pm \sigma_j = \pm \frac{\sigma_{j\pm 1} - \sigma_j}{\Delta x},$$

for any sequence $\{\sigma_j\}$.

We may now define a semi-discrete approximation of the solution to (1.1) as the solution to the (infinite) system of ordinary differential equations

$$(3.1) \quad \begin{cases} \frac{d}{dt} u_j(t) + D_- F_{j+1/2} = D_- D_+ A(u_j), & t > 0, \\ u_j(0) = \frac{1}{\Delta x} \int_{I_j} u^0(x) dx, \end{cases} \quad j \in \mathbb{Z},$$

where $F_{j+1/2} = F(u_j, u_{j+1})$ is a numerical flux function and $I_j = (x_{j-1/2}, x_{j+1/2}]$.

The problem above can be viewed as an ordinary differential equation in the Banach space $\ell^1(\mathbb{Z})$ (see, e.g., [23]). To get bounds independent of Δx we define

$$\|\sigma\|_1 = \Delta x \sum_j |\sigma_j| \quad \text{and} \quad |\sigma|_{BV} = \sum_j |\sigma_{j+1} - \sigma_j| = \|D_+ \sigma\|_1.$$

If these are bounded we say that $\sigma = \{\sigma_j\}$ is in ℓ^1 and of bounded variation. Let $u(t) = \{u_j(t)\}_{j \in \mathbb{Z}}$, $u^0 = \{u_j(0)\}_{j \in \mathbb{Z}}$, and define the operator $\mathcal{A} : \ell^1 \rightarrow \ell^1$ by $(\mathcal{A}(u))_j := D_-(F(u_j, u_{j+1}) - D_+ A(u_j))$. Then (3.1) takes the following form:

$$\frac{du}{dt} + \mathcal{A}(u) = 0, \quad t > 0, \quad u(0) = u^0.$$

This problem has a unique continuously differentiable solution since $\mathcal{A}$ is Lipschitz continuous for each fixed $\Delta x > 0$. This solution defines a strongly continuous semigroup $\mathcal{S}(t)$ on ℓ^1 . If $\mathcal{S}$ also satisfies

$$\|\mathcal{S}(t)u - \mathcal{S}(t)v\|_1 \leq \|u - v\|_1 \quad \text{for} \quad u, v \in \ell^1,$$

we say that it is nonexpansive. The next lemma sums up some important properties of the solutions to (3.1) (for a proof see [11]).

Lemma 3.1. *Suppose that F is monotone. Then there exists a unique solution $u = \{u_j\}$ to (3.1) on $[0, T]$ with the following properties:*

- (a) $\|u(t)\|_1 \leq \|u^0\|_1$.
- (b) For every $j \in \mathbb{Z}$ and $t \in [0, T]$,

$$\inf_k \{u_k^0\} \leq u_j(t) \leq \sup_k \{u_k^0\}.$$

- (c) $|u(t)|_{BV} \leq |u^0|_{BV}$.
- (d) If $v = \{v_j\}$ is another solution with initial data v^0 , then

$$\|u(t) - v(t)\|_1 \leq \|u^0 - v^0\|_1.$$

Lemma 3.2. *If F is monotone, then*

$$(3.2) \quad \|F(u_j, u_{j+1}) - D_+ A(u_j)\|_{\ell^\infty} \leq \|F(u_j^0, u_{j+1}^0) - D_+ A(u_j^0)\|_{\ell^\infty},$$

$$(3.3) \quad |F(u_j, u_{j+1}) - D_+ A(u_j)|_{BV} \leq |F(u_j^0, u_{j+1}^0) - D_+ A(u_j^0)|_{BV}.$$

Furthermore, $t \mapsto \{u_j(t)\}_{j \in \mathbb{Z}}$ is ℓ^1 Lipschitz continuous.

Proof. The proof follows [12]. Let $v_j = \Delta x \sum_{k \leq j} \frac{du_k}{dt}$. Then v_j satisfies

$$(3.4) \quad v_j = \Delta x \sum_{k=-\infty}^j D_- (D_+ A(u_k) - F(u_k, u_{k+1})) = D_+ A(u_j) - F(u_j, u_{j+1}),$$

and we may define v_j for all $t \in [0, T]$. Note that $\{v_j(t)\}$ is in ℓ^1 for all t by Lemma 3.1. Differentiating (3.4) with respect to t we obtain

$$\begin{aligned} \frac{dv_j}{dt} &= \frac{1}{\Delta x} \left[a(u_{j+1}) \frac{du_{j+1}}{dt} - a(u_j) \frac{du_j}{dt} \right] \\ &\quad - F_u(u_j, u_{j+1}) \frac{du_j}{dt} - F_v(u_j, u_{j+1}) \frac{du_{j+1}}{dt}, \end{aligned}$$

where $a(u) = A'(u)$. Note that $D_- v_j = \frac{du_j}{dt}$ and $D_+ v_j = \frac{du_{j+1}}{dt}$. Therefore

$$(3.5) \quad \begin{aligned} \frac{dv_j}{dt} &= \left(\frac{1}{\Delta x} a(u_{j+1}) - F_v(u_j, u_{j+1}) \right) D_+ v_j \\ &\quad - \left(\frac{1}{\Delta x} a(u_j) + F_u(u_j, u_{j+1}) \right) D_- v_j. \end{aligned}$$

Assume $v_{j_0}(t_0)$ is a local maximum in j . Then $D_+ v_{j_0}(t_0) \leq 0$ and $D_- v_{j_0}(t_0) \geq 0$ so $\frac{dv_{j_0}}{dt}(t_0) \leq 0$ since F is monotone. Similarly, if $v_{j_0}(t_0)$ is a local minimum in j , then $\frac{dv_{j_0}}{dt}(t_0) \geq 0$. Then inequality (3.2) follows by the fact that $\{v_j(t)\} \in \ell^1$. Consider (3.3). We want to show that $\frac{d}{dt}(|v(t)|_{BV}) \leq 0$. Now,

$$\frac{d}{dt} \left(\sum_j |v_{j+1} - v_j| \right) = \sum_j \text{sign}(v_{j+1} - v_j) \frac{d}{dt} (v_{j+1} - v_j),$$

so we may use (3.5). Thus

$$\begin{aligned}
& \frac{d}{dt} |v(t)|_{BV} \\
&= \sum_j \left(\frac{1}{\Delta x} a(u_{j+2}) - F_v(u_{j+1}, u_{j+2}) \right) (D_+ v_{j+1}) \operatorname{sign}(v_{j+1} - v_j) \\
&\quad - \sum_j \left(\frac{1}{\Delta x} a(u_{j+1}) + F_u(u_{j+1}, u_{j+2}) \right) |D_+ v_j| \\
&\quad - \sum_j \left(\frac{1}{\Delta x} a(u_{j+1}) - F_v(u_j, u_{j+1}) \right) |D_+ v_j| \\
&\quad + \sum_j \left(\frac{1}{\Delta x} a(u_j) + F_u(u_j, u_{j+1}) \right) ((D_- v_j) \operatorname{sign}(v_{j+1} - v_j)) \\
&= \sum_j \left(\frac{1}{\Delta x} a(u_{j+1}) - F_v(u_j, u_{j+1}) \right) [(D_+ v_j) \operatorname{sign}(v_j - v_{j-1}) - |D_+ v_j|] \\
&\quad + \sum_j \left(\frac{1}{\Delta x} a(u_j) + F_u(u_j, u_{j+1}) \right) [(D_- v_j) \operatorname{sign}(v_{j+1} - v_j) - |D_- v_j|] \\
&\leq 0,
\end{aligned}$$

since $a(u) > 0$, $F_v \leq 0$, and $F_u \geq 0$. Given the preceding estimates, the ℓ^1 Lipschitz continuity is straightforward to prove. $\square$

It turns out that we need more conditions on F than mere monotonicity.

Definition 3.2. Given an entropy pair (ψ, q) and a numerical flux F , we define $Q \in C^1(\mathbb{R}^2)$ by

$$\begin{aligned}
Q(u, u) &= q(u), \\
\frac{\partial}{\partial v} Q(v, w) &= \psi'(v) \frac{\partial}{\partial v} F(v, w), \\
\frac{\partial}{\partial w} Q(v, w) &= \psi'(w) \frac{\partial}{\partial w} F(v, w).
\end{aligned}$$

We call Q a numerical entropy flux.

The next lemma gives a sufficient condition on the numerical flux to ensure that there exists a numerical entropy flux.

Lemma 3.3. *Given a two-point numerical flux F , assume that there exist C^1 functions F_1, F_2 such that*

$$(3.6) \quad F(u, v) = F_1(u) + F_2(v), \quad F'_1(u) + F'_2(u) = f'(u),$$

for all relevant u and v . Then there exists a numerical entropy flux Q for any entropy flux pair (ψ, q) .

Proof. Let (ψ, q) be an entropy pair. Then q has the form

$$q(u) = \int_c^u \psi'(z) f'(z) dz + C,$$

for some constant C . Define Q by

$$(3.7) \quad Q(u, v) = \int_c^u \psi'(z) F_1'(z) dz + \int_c^v \psi'(z) F_2'(z) dz + C.$$

It is easily verified that Q is a numerical entropy flux. $\square$

Let us list a few numerical flux functions to which Lemma 3.3 applies.

Example 3.1 (Engquist-Osher flux). Let

$$f'_+(s) = \max(f'(s), 0) \quad \text{and} \quad f'_-(s) = \min(f'(s), 0).$$

Then, in the terminology of Lemma 3.3, let $F(u, v) = F_1(u) + F_2(v)$ with

$$F_1(u) = f(0) + \int_0^u f'_+(s) ds \quad \text{and} \quad F_2(v) = \int_0^v f'_-(s) ds.$$

It is easily seen that the criteria given in Lemma 3.3 are satisfied, and F is also clearly monotone.

Example 3.2. Let $a, b \in \mathbb{R}$ and define

$$F_1(u) = af(u) + bu \quad \text{and} \quad F_2(v) = (1-a)f(v) - bv.$$

Note that $F(u, v) = F_1(u) + F_2(v)$ is monotone if

$$a \inf_u \{f'(u)\} \geq -b \quad \text{and} \quad (1-a) \sup_u \{f'(u)\} \leq b.$$

This example includes both the upwind scheme and the Lax-Friedrichs scheme.

From a more general point of view we may consider any flux splitting, that is, $f(u) = f^+(u) + f^-(u)$ with $(f^+)'(u) \geq 0$ and $(f^-)'(u) \leq 0$ for all $u \in \mathbb{R}$. Then the numerical flux

$$F(u, v) = f^+(u) + f^-(v)$$

satisfies the assumptions of Lemma 3.3. Note also that any convex combination of numerical flux functions which satisfy the hypothesis of Lemma 3.3, itself satisfies the assumptions of the lemma.

If (3.6) holds, then we have a representation of Q given by (3.7). It follows that

$$(3.8) \quad Q(u, v) = q(u) + \int_u^v \psi'(z) F_2'(z) dz.$$

Note that we may obtain another representation depending on F_1 by splitting up the first integral.

4. ERROR ESTIMATE

Let $\{u_j\}_{j \in \mathbb{Z}}$ be the solution to (3.1). We associate with it the piecewise constant function

$$(4.1) \quad u_{\Delta x}(x, t) = u_j(t) \quad \text{for } x \in I_j.$$

To derive the error estimate we need many of the uniform bounds from Sections 2 and 3. For these estimates to hold independently of Δx , we make the following assumptions on the initial data u^0 :

- (i) $u^0 \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R}) \cap BV(\mathbb{R})$.
- (ii) $A(u^0)_x \in BV(\mathbb{R})$.

We may now state the theorem.

Theorem 4.1. *Let u be the entropy solution to (1.1) and $\{u_j(t)\}_{j \in \mathbb{Z}}$ solve the semi-discrete difference scheme (3.1). If u^0 satisfies (i) and (ii) above, then for all sufficiently small Δx ,*

$$\|u_{\Delta x}(\cdot, t) - u(\cdot, t)\|_{L^1(\mathbb{R})} \leq \|u_{\Delta x}^0 - u^0\|_{L^1(\mathbb{R})} + C_T \Delta x^{\frac{1}{3}}, \quad t \in [0, T],$$

where the constant C_T depends on A , f , u^0 , and T , but not on Δx .

Let us define some of the functions we are going to work with. First, we will use the following approximation of the sign function:

$$\text{sign}_\varepsilon(\sigma) = \begin{cases} \sin(\frac{\pi\sigma}{2\varepsilon}) & \text{for } |\sigma| < \varepsilon, \\ \text{sign}(\sigma) & \text{otherwise,} \end{cases}$$

where $\varepsilon > 0$. Note that sign_ε is continuously differentiable and nondecreasing. We define

$$|u|_\varepsilon = \int_0^u \text{sign}_\varepsilon(z) dz.$$

Furthermore, we introduce an entropy pair $(\psi_\varepsilon, q_\varepsilon)$ defined by

$$\begin{aligned} \psi_\varepsilon(u, c) &= \int_c^u \text{sign}_\varepsilon(A(z) - A(c)) dz, \\ q_\varepsilon(u, c) &= \int_c^u \psi'_\varepsilon(z, c) f'(z) dz = \int_c^u \text{sign}_\varepsilon(A(z) - A(c)) f'(z) dz, \end{aligned}$$

where ψ'_ε is the derivative with respect to the first variable.

Lemma 4.1. *Suppose $A' > 0$. Let $u = u(y, s)$ be the classical solution of (1.1). Then for any constant $c \in \mathbb{R}$,*

$$\partial_s \psi_\varepsilon(u, c) + \partial_y q_\varepsilon(u, c) - \partial_y^2 |A(u) - A(c)|_\varepsilon = -\partial_y \psi'_\varepsilon(u, c) \partial_y A(u).$$

Proof. Multiply equation (1.1) by $\psi'_\varepsilon(u, c)$ to obtain

$$\partial_s \psi_\varepsilon(u, c) + \partial_y q_\varepsilon(u, c) = \psi'_\varepsilon(u, c) \partial_y^2 A(u).$$

The term on the right may be rewritten according to

$$\partial_y(\psi'_\varepsilon(u, c) \partial_y A(u)) = \partial_y \psi'_\varepsilon(u, c) \partial_y A(u) + \psi'_\varepsilon(u, c) \partial_y^2 A(u).$$

By the chain rule,

$$\partial_y(\psi'_\varepsilon(u, c) \partial_y A(u)) = \partial_y^2 |A(u) - A(c)|_\varepsilon.$$

Combining these equalities proves the lemma. $\square$

The next lemma is a simple identity taken from [8].

Lemma 4.2. *For any differentiable function g and all real numbers a, b, c ,*

$$\begin{aligned} \psi'_\varepsilon(a, c)(g(b) - g(a)) &= \int_c^b \psi'_\varepsilon(z, c) g'(z) dz - \int_c^a \psi'_\varepsilon(z, c) g'(z) dz \\ &\quad + \int_a^b \psi''_\varepsilon(z, c)(g(z) - g(b)) dz. \end{aligned}$$

Proof. Integration by parts yields

$$\psi'_\varepsilon(\zeta, c)(g(\zeta) - g(b)) = \int_c^\zeta \psi'_\varepsilon(z, c)g'(z) dz + \int_c^\zeta \psi''_\varepsilon(z, c)(g(z) - g(b)) dz,$$

for any $\zeta \in \mathbb{R}$. Take the two equations obtained by taking $\zeta = a$ and $\zeta = b$ and subtract one from the other. $\square$

Lemma 4.3. *Let u_j be the solution to (3.1). Then for all $c \in \mathbb{R}$,*

$$\begin{aligned} & \partial_t \psi_\varepsilon(u_j, c) + D_- Q^c(u_j, u_{j+1}) - D_- D_+ |A(u_j) - A(c)|_\varepsilon \\ & \leq -\frac{1}{(\Delta x)^2} \int_{u_{j+1}}^{u_j} \psi''_\varepsilon(z, c)(A(z) - A(u_{j+1})) dz \\ & \quad - \frac{1}{(\Delta x)^2} \int_{u_{j-1}}^{u_j} \psi''_\varepsilon(z, c)(A(z) - A(u_{j-1})) dz, \end{aligned}$$

where $Q^c(u, v) := Q_1^c(u) + Q_2^c(v)$,

$$Q_1^c(u) := \int_c^u \psi'_\varepsilon(z, c)F'_1(z) dz, \quad Q_2^c(v) := \int_c^v \psi'_\varepsilon(z, c)F'_2(z) dz,$$

for all real numbers u and v .

Proof. From (3.1) it follows that

$$\psi'_\varepsilon(u_j, c)\partial_t u_j + \psi'_\varepsilon(u_j, c)D_- F(u_j, u_{j+1}) = \psi'_\varepsilon(u_j, c)D_- D_+ A(u_j).$$

Note that

$$\psi'_\varepsilon(u_j, c)D_- F(u_j, u_{j+1}) = \psi'_\varepsilon(u_j, c)D_- F_1(u_j) + \psi'_\varepsilon(u_j, c)D_+ F_2(u_j),$$

and so we may apply Lemma 4.2. Let $g = F_1$. Then we obtain

$$\psi'_\varepsilon(u_j, c)D_- F_1(u_j) = D_- Q_1^c(u_j) - \frac{1}{\Delta x} \int_{u_j}^{u_{j-1}} \psi''_\varepsilon(z, c)(F_1(z) - F_1(u_{j-1})) dz.$$

Similarly, let $g = F_2$ to obtain

$$\psi'_\varepsilon(u_j, c)D_+ F_2(u_j) = D_+ Q_2^c(u_j) + \frac{1}{\Delta x} \int_{u_j}^{u_{j+1}} \psi''_\varepsilon(z, c)(F_2(z) - F_2(u_{j+1})) dz.$$

Finally, apply Lemma 4.2 twice with $g = A$. Adding the equations we obtain

$$\begin{aligned} & \psi'_\varepsilon(u_j, c)(A(u_{j-1}) - 2A(u_j) + A(u_{j+1})) \\ & = \int_{u_j}^{u_{j+1}} \psi'_\varepsilon(z, c)A'(z) dz + \int_{u_j}^{u_{j-1}} \psi'_\varepsilon(z, c)A'(z) dz \\ & \quad + \int_{u_j}^{u_{j+1}} \psi''_\varepsilon(z, c)(A(z) - A(u_{j+1})) dz \\ & \quad + \int_{u_j}^{u_{j-1}} \psi''_\varepsilon(z, c)(A(z) - A(u_{j-1})) dz. \end{aligned}$$

Note that

$$\begin{aligned} & \left[\int_{u_j}^{u_{j+1}} \psi'_\varepsilon(z, c) A'(z) dz + \int_{u_j}^{u_{j-1}} \psi'_\varepsilon(z, c) A'(z) dz \right] \\ &= \left[\int_{u_j}^{u_{j+1}} \frac{\partial}{\partial z} |A(z) - A(c)|_\varepsilon dz + \int_{u_j}^{u_{j-1}} \frac{\partial}{\partial z} |A(z) - A(c)|_\varepsilon dz \right] \\ &= [|A(u_{j-1}) - A(c)|_\varepsilon - 2|A(u_j) - A(c)|_\varepsilon + |A(u_{j+1}) - A(c)|_\varepsilon]. \end{aligned}$$

Combining the above computations we obtain

$$\partial_t \psi_\varepsilon(u_j, c) + D_- Q^c(u_j, u_{j+1}) - D_- D_+ |A(u_j) - A(u)|_\varepsilon = -E^c(u_{j-1}, u_j, u_{j+1}),$$

where

$$\begin{aligned} E^c(u_{j-1}, u_j, u_{j+1}) &= \frac{1}{\Delta x} \int_{u_{j-1}}^{u_j} \psi''_\varepsilon(z, c) (F_1(z) - F_1(u_{j-1})) dz \\ &\quad - \frac{1}{\Delta x} \int_{u_{j+1}}^{u_j} \psi''_\varepsilon(z, c) (F_2(z) - F_2(u_{j+1})) dz \\ &\quad + \frac{1}{(\Delta x)^2} \int_{u_{j+1}}^{u_j} \psi''_\varepsilon(z, c) (A(z) - A(u_{j+1})) dz \\ &\quad + \frac{1}{(\Delta x)^2} \int_{u_{j-1}}^{u_j} \psi''_\varepsilon(z, c) (A(z) - A(u_{j-1})) dz. \end{aligned}$$

The result follows from the monotonicity of F . $\square$

We shall need the next lemma, which deals with a mixed term, in order to carry out the “second order” doubling-of-the-variables argument.

Lemma 4.4. *Let $\{u_j\}$ be some sequence and u some differentiable function of y . Then*

$$\begin{aligned} & \left(\frac{1}{\Delta x} \int_{u_{j-1}}^{u_j} \psi''_\varepsilon(z, u) dz + \frac{1}{\Delta x} \int_{u_j}^{u_{j+1}} \psi''_\varepsilon(z, u) dz \right) \partial_y A(u) \\ &= -\partial_y (D_- + D_+) |A(u_j) - A(u)|_\varepsilon. \end{aligned}$$

Proof. Let a, b be fixed real numbers. Then

$$\begin{aligned} & \int_a^b \psi''_\varepsilon(z, u) dz A(u)_y \\ &= \int_a^b \text{sign}'_\varepsilon(A(z) - A(u)) A(u)_y A'(z) dz \\ &= -\frac{\partial}{\partial y} \left(\int_a^b \text{sign}_\varepsilon(A(z) - A(u)) A'(z) dz \right) \\ &= -\frac{\partial}{\partial y} (|A(b) - A(u)|_\varepsilon - |A(a) - A(u)|_\varepsilon). \end{aligned}$$

Let $a = u_{j-1}$, $b = u_j$ and $a = u_j$, $b = u_{j+1}$. Then add up the resulting equations and divide by Δx . $\square$

We are now in a position to carry out the doubling-of-the-variables argument.

Lemma 4.5. Suppose $A' > 0$. Let $u = u(y, s)$ be the classical solution to (1.1) and let $\{u_j\} = \{u_j(t)\}$ be the solution to (3.1). Then

$$\begin{aligned} & \partial_t \psi_\varepsilon(u_j, u) + \partial_s \psi_\varepsilon(u, u_j) + \partial_y q_\varepsilon(u, u_j) + D_- Q^u(u_j, u_{j+1}) \\ & - (\partial_y^2 + \partial_y(D_- + D_+) + D_- D_+) |A(u_j) - A(u)|_\varepsilon \leq -E_j^\varepsilon, \end{aligned}$$

where $E_j^\varepsilon := E^\varepsilon[u](u_{j-1}, u_j, u_{j+1})$ with

$$\begin{aligned} E^\varepsilon[u](u_{j-1}, u_j, u_{j+1}) &:= \frac{1}{(\Delta x)^2} \int_{u_{j+1}}^{u_j} \psi_\varepsilon''(z, u)(A(z) - A(u_{j+1})) dz \\ &+ \frac{1}{(\Delta x)^2} \int_{u_{j-1}}^{u_j} \psi_\varepsilon''(z, u)(A(z) - A(u_{j-1})) dz \\ &- \frac{1}{\Delta x} \int_{u_{j-1}}^{u_j} \psi_\varepsilon''(z, u) dz \partial_y A(u) \\ &- \frac{1}{\Delta x} \int_{u_j}^{u_{j+1}} \psi_\varepsilon''(z, u) dz \partial_y A(u) \\ &+ \partial_y \psi_\varepsilon'(u, u_j) \partial_y A(u). \end{aligned}$$

Proof. Let $c = u_j$ in Lemma 4.1 and $c = u$ in Lemma 4.3. Then add up the equations together with Lemma 4.4. $\square$

Remark 4.1. Note that E_j^ε is a function of y, s, t .

In what follows it will be necessary to work with the piecewise constant approximation defined in (4.1). To do this we introduce some new notation. Let the shift operator S_σ be defined for any $\varphi : \Pi_T \rightarrow \mathbb{R}$ by

$$S_\sigma \varphi(x, t) = \varphi(x + \sigma, t),$$

and let the difference quotient be defined by

$$D_\pm \varphi = \pm \frac{S_{\pm \Delta x} \varphi - \varphi}{\Delta x}.$$

Note that for any two functions u, v of x we have $D_+(uv) = S_{\Delta x} u D_+ v + (D_+ u)v$. If uv has compact support it follows that

$$\int_{\mathbb{R}} (D_+ u)v dx = - \int_{\mathbb{R}} u D_- v dx.$$

We will refer to these identities as the Leibniz rule for difference quotients and integration by parts for difference quotients. We will frequently integrate over the domain Π_T^2 . To avoid writing four integral signs we will in general write one for each domain Π_T and let $dX = dx dt dy ds$.

Lemma 4.6. Suppose $A' > 0$. Let $u_{\Delta x} = u_{\Delta x}(x, t)$ be defined by (4.1), and let $u = u(y, s)$ be the classical solution of (1.1). Let $\rho \in C_0^\infty(\mathbb{R})$ satisfy

$$\text{supp}(\rho) \subset [-1, 1], \quad \rho(-\sigma) = \rho(\sigma), \quad \rho(\sigma) \geq 0, \quad \int_{\mathbb{R}} \rho(\sigma) d\sigma = 1,$$

and set

$$\omega_r(x) = \frac{1}{r} \rho\left(\frac{x}{r}\right), \quad \rho_\alpha(\xi) = \frac{1}{\alpha} \rho\left(\frac{\xi}{\alpha}\right), \quad \rho_{r_0}(t) = \frac{1}{r_0} \rho\left(\frac{t}{r_0}\right),$$

for positive (small) r , α and r_0 . Let ν and τ be such that $0 < \nu < \tau < T$ and define

$$\psi^\alpha(t) := H_\alpha(t - \nu) - H_\alpha(t - \tau), \quad H_\alpha(t) = \int_{-\infty}^t \rho_\alpha(\xi) d\xi.$$

Let

$$\varphi(x, t, y, s) = \psi^\alpha(t) \omega_r(x - y) \rho_{r_0}(t - s).$$

To ensure $\varphi|_{t=0} \equiv 0$, $\varphi|_{s=0} \equiv 0$, we choose ν and τ such that $0 < r_0 < \min(\nu, T - \tau)$ and $0 < \alpha < \min(\nu - r_0, T - \tau - r_0)$. Then

$$\begin{aligned} (4.2) \quad & \iint_{\Pi_T^2} |u_{\Delta x} - u| \rho_\alpha(t - \nu) \omega_r \rho_{r_0} dX \\ & + \iint_{\Pi_T^2} \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)) (D_+ \varphi + \varphi_y) dX \\ & + \iint_{\Pi_T^2} \left(\int_{u_{\Delta x}}^{S_{\Delta x} u_{\Delta x}} \text{sign}(z - u) F'_2(z) dz \right) D_+ \varphi dX \\ & + \iint_{\Pi_T^2} |A(u_{\Delta x}) - A(u)| (D_- D_+ \varphi + (D_+ + D_-) \varphi_y + \varphi_{yy}) dX \\ & \geq \iint_{\Pi_T^2} |u_{\Delta x} - u| \rho_\alpha(t - \tau) \omega_r \rho_{r_0} dX + \liminf_{\varepsilon \downarrow 0} \iint_{\Pi_T^2} E_{\Delta x}^\varepsilon \varphi dX, \end{aligned}$$

where $E_{\Delta x}^\varepsilon(x, t, y, s) = E_j^\varepsilon(t, y, s)$ for $x \in I_j$.

Remark 4.2. Note that both

$$\varphi_x + \varphi_y = 0 \quad \text{and} \quad \varphi_{xx} + 2\varphi_{xy} + \varphi_{yy} = 0.$$

In equation (4.2) these expressions appear with difference quotients instead of x -derivatives. We expect that these equalities turn into good approximations as long as Δx tends relatively fast to zero compared to r . We will show that this is the case in what follows.

Proof. By Lemma 4.5 it follows that

$$\begin{aligned} & \partial_t \psi_\varepsilon(u_{\Delta x}, u) + \partial_s \psi_\varepsilon(u, u_{\Delta x}) + \partial_y q_\varepsilon(u, u_{\Delta x}) + D_- Q^u(u_{\Delta x}, S_{\Delta x} u_{\Delta x}) \\ & - (\partial_y^2 + \partial_y (D_- + D_+) + D_- D_+) |A(u_{\Delta x}) - A(u)|_\varepsilon \leq -E_{\Delta x}^\varepsilon, \end{aligned}$$

for all $(x, t, y, s) \in \Pi_T^2$. Let us multiply with φ and integrate over Π_T^2 . Using both ordinary integration by parts and integration by parts for difference quotients, we obtain

$$\begin{aligned} & \iint_{\Pi_T^2} \psi_\varepsilon(u_{\Delta x}, u) \varphi_t + \psi_\varepsilon(u, u_{\Delta x}) \varphi_s dX \\ & + \iint_{\Pi_T^2} q_\varepsilon(u, u_{\Delta x}) \varphi_y + Q^u(u_{\Delta x}, S_{\Delta x} u_{\Delta x}) D_+ \varphi dX \\ & + \iint_{\Pi_T^2} |A(u_{\Delta x}) - A(u)|_\varepsilon (\varphi_{yy} + (D_- + D_+) \varphi_y + D_- D_+ \varphi) dX \\ & \geq \iint_{\Pi_T^2} E_{\Delta x}^\varepsilon \varphi dX. \end{aligned}$$

We want to take the limit as $\varepsilon \downarrow 0$. Consider the first term on the left. By the dominated convergence theorem, for any $a, b \in \mathbb{R}$,

$$\lim_{\varepsilon \downarrow 0} \psi_\varepsilon(a, b) = \lim_{\varepsilon \downarrow 0} \int_b^a \text{sign}_\varepsilon(A(z) - A(b)) dz = |a - b|,$$

since $A' > 0$. It follows that

$$\lim_{\varepsilon \downarrow 0} \psi_\varepsilon(u_{\Delta x}, u) = \lim_{\varepsilon \downarrow 0} \psi_\varepsilon(u, u_{\Delta x}) = |u_{\Delta x} - u|.$$

Furthermore,

$$(\varphi_t + \varphi_s)(x, t, y, s) = (\rho_\alpha(t - \nu) - \rho_\alpha(t - \tau))\omega_r(x - y)\rho_{r_0}(t - s),$$

so by the dominated convergence theorem

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} \iint_{\Pi_T^2} \psi_\varepsilon(u_{\Delta x}, u) \varphi_t + \psi_\varepsilon(u, u_{\Delta x}) \varphi_s dX \\ = \iint_{\Pi_T^2} |u_{\Delta x} - u| \rho_\alpha(t - \nu) \omega_r \rho_{r_0} dX - \iint_{\Pi_T^2} |u_{\Delta x} - u| \rho_\alpha(t - \tau) \omega_r \rho_{r_0} dX. \end{aligned}$$

Consider the second term on the left. By (3.8) we obtain

$$Q^u(u_{\Delta x}, S_{\Delta x} u_{\Delta x}) = q_\varepsilon(u_{\Delta x}, u) + \int_{u_{\Delta x}}^{S_{\Delta x} u_{\Delta x}} \text{sign}_\varepsilon(A(z) - A(u)) F_2'(z) dz.$$

Since $A' > 0$,

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} q_\varepsilon(u_{\Delta x}, u) &= \lim_{\varepsilon \downarrow 0} \int_u^{u_{\Delta x}} \text{sign}_\varepsilon(A(z) - A(u_{\Delta x})) f'(z) dz \\ &= \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)). \end{aligned}$$

It follows that

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} Q^u(u_{\Delta x}, S_{\Delta x} u_{\Delta x}) &= \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)) \\ &\quad + \int_{u_{\Delta x}}^{S_{\Delta x} u_{\Delta x}} \text{sign}(z - u) F_2'(z) dz. \end{aligned}$$

As above

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} q_\varepsilon(u, u_{\Delta x}) &= \text{sign}(u - u_{\Delta x}) (f(u) - f(u_{\Delta x})) \\ &= \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)). \end{aligned}$$

Hence, again by the dominated convergence theorem,

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} \iint_{\Pi_T^2} q_\varepsilon(u, u_{\Delta x}) \varphi_y + Q^u(u_{\Delta x}, S_{\Delta x} u_{\Delta x}) D_+ \varphi dX \\ = \iint_{\Pi_T^2} \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)) (\varphi_y + D_+ \varphi) dX \\ + \iint_{\Pi_T^2} \left(\int_{u_{\Delta x}}^{S_{\Delta x} u_{\Delta x}} \text{sign}(z - u) F_2'(z) dz \right) D_+ \varphi dX. \quad \square \end{aligned}$$

Lemma 4.7. Let $E_{\Delta x}^\varepsilon$ and φ be defined in Lemma 4.6. Then

$$\liminf_{\varepsilon \downarrow 0} \iint_{\Pi_T^2} E_{\Delta x}^\varepsilon \varphi dX \geq \int_{\Pi_T} \liminf_{\varepsilon \downarrow 0} \left(\int_{\Pi_T} E_{\Delta x}^\varepsilon \varphi dy ds \right) dx dt.$$

Proof. Let

$$f_\varepsilon(x, t) := \int_{\Pi_T} E_{\Delta x}^\varepsilon \varphi \, dy ds$$

and

$$h_\varepsilon(x, t) := \int_{\Pi_T} \partial_y (D_- + D_+) |A(u_{\Delta x}) - A(u)|_\varepsilon \varphi \, dy ds.$$

Recall that $E_{\Delta x}^\varepsilon(x, t, y, s) = E_j^\varepsilon(t, y, s)$ for $x \in I_j$, where E_j^ε is defined in Lemma 4.5. Note that

$$\begin{aligned} E_j^\varepsilon &\geq -\frac{1}{\Delta x} \int_{u_{j-1}}^{u_j} \psi_\varepsilon''(z, u) \, dz \, \partial_y A(u) \\ &\quad - \frac{1}{\Delta x} \int_{u_j}^{u_{j+1}} \psi_\varepsilon''(z, u) \, dz \, \partial_y A(u), \end{aligned}$$

so by Lemma 4.4 it follows that $f_\varepsilon \geq h_\varepsilon$. Using integration by parts and the triangle inequality we obtain the bound

$$|h_\varepsilon| \leq (|D_- A(u_{\Delta x})| + |D_+ A(u_{\Delta x})|) \left(\int_{\Pi_T} |\varphi_y| \, dy ds \right) =: h.$$

It follows by Lemma 3.1 that h is an integrable nonnegative function such that $-h \leq f_\varepsilon$. By Fatou's lemma we obtain

$$\liminf_{\varepsilon \downarrow 0} \int_{\Pi_T} f_\varepsilon \, dx dt \geq \int_{\Pi_T} \liminf_{\varepsilon \downarrow 0} f_\varepsilon \, dx dt. \quad \square$$

Note that as $\varepsilon \downarrow 0$ the terms in E_j^ε concentrate on the domains specified by $u \in \text{int}(u_j, u_{j+1})$, $u \in \text{int}(u_{j-1}, u_j)$, or $u = u_j$. In order to analyze this limit we will need the following elementary lemma:

Lemma 4.8. *Let $\{u_j\}_{j \in \mathbb{Z}}$ be some sequence in $\mathbb{R}$ and let $A : \mathbb{R} \rightarrow \mathbb{R}$ be a strictly increasing continuously differentiable function. For any $u \in \mathbb{R}$ there exist sequences $\{\tau_j^\pm\}_{j \in \mathbb{Z}}$, $\{\theta_j^\pm\}_{j \in \mathbb{Z}}$ such that for each $j \in \mathbb{Z}$ both $\tau_j^\pm$ and $\theta_j^\pm$ are in $\text{int}(u_j, u_{j \pm 1})$ and*

$$\begin{aligned} D_\pm \text{sign}_\varepsilon(A(u_j) - A(u)) &= \text{sign}_\varepsilon'(A(\tau_j^\pm) - A(u)) D_\pm A(u_j), \\ D_\pm |A(u_j) - A(u)|_\varepsilon &= \text{sign}_\varepsilon(A(\theta_j^\pm) - A(u)) D_\pm A(u_j). \end{aligned}$$

If u is a differentiable function of y , then for each $j \in \mathbb{Z}$,

$$(4.3) \quad \text{sign}_\varepsilon'(A(\tau_j^\pm) - A(u)) A(u)_y = -(\text{sign}_\varepsilon(A(\theta_j^\pm) - A(u)))_y.$$

Both $\{\tau_j^\pm\}_{j \in \mathbb{Z}}$ and $\{\theta_j^\pm\}_{j \in \mathbb{Z}}$ depend on u and ε .

Proof. The first statement is a direct consequence of the mean value theorem. Consider (4.3). First note that $\tau_j^- = \tau_{j-1}^+$ and $\theta_j^- = \theta_{j-1}^+$, so it suffices to consider τ_j^+ and θ_j^+ . If $u_j = u_{j+1}$, then $\theta_j = \tau_j$ is independent of u and hence of y , so (4.3) follows by the chain rule. In general,

$$\begin{aligned} \text{sign}_\varepsilon'(A(\tau_j) - A(u)) A(u)_y D_+ A(u_j) &= D_+ \text{sign}_\varepsilon(A(u_j) - A(u)) A(u)_y \\ &= -D_+ (|A(u_j) - A(u)|_\varepsilon)_y \\ &= -\text{sign}_\varepsilon(A(\theta_j) - A(u))_y D_+ A(u_j). \end{aligned}$$

In the case $u_j \neq u_{j+1}$ we have $D_+ A(u_j) \neq 0$ and (4.3) follows. $\square$

The following result is concerned with the pointwise limit of $\text{sign}_\varepsilon(A(\theta_j^\pm) - A(u))$ as $\varepsilon \downarrow 0$. The explicit formula for this limit, which will be used later, shows that the limit is in fact a Lipschitz continuous function in the case that $A(u_j) \neq A(u_{j+1})$.

Lemma 4.9. *Let*

$$\text{sg}_{(a,b)}(\sigma) := \begin{cases} \frac{|a-\sigma| - |b-\sigma|}{b-a} & \text{if } a \neq b, \\ \text{sign}(a - \sigma) & \text{if } a = b, \sigma \neq b, \\ 0 & \text{if } a = b = \sigma, \end{cases}$$

for any real numbers a and b . Under the same assumptions as in Lemma 4.8,

$$\lim_{\varepsilon \downarrow 0} \text{sign}_\varepsilon(A(\theta_j^\pm) - A(u)) = -\text{sg}_{(A(u_j), A(u_{j+1}))}(A(u)).$$

Furthermore, if $a \neq b$, then

$$\text{sg}_{(a,b)}(\sigma) = \begin{cases} -1 & \text{if } \sigma \leq \min\{a, b\}, \\ \frac{2}{|a-b|}(\sigma - \frac{1}{2}(b+a)) & \text{if } \sigma \in \text{int}(a, b), \\ 1 & \text{if } \sigma \geq \max\{a, b\}. \end{cases}$$

Proof. To prove the first statement we consider the case of θ_j^+ . The same argument applies to θ_j^- . Recall the definition of θ_j^+ :

$$\text{sign}_\varepsilon(A(\theta_j^+) - A(u))(A(u_{j+1}) - A(u_j)) = |A(u_{j+1}) - A(u)|_\varepsilon - |A(u_j) - A(u)|_\varepsilon.$$

If $u_{j+1} = u_j$, then $\theta_j^+ = u_j$ for all u and ε , since $\theta_j^+ \in \text{int}(u_j, u_{j+1})$. Thus in this case

$$\lim_{\varepsilon \downarrow 0} \text{sign}_\varepsilon(A(\theta_j^+) - A(u)) = \begin{cases} 0 & \text{if } u = u_j, \\ \text{sign}(A(u_j) - A(u)) & \text{otherwise.} \end{cases}$$

Assume that $D_+A(u_j) \neq 0$. Then

$$\text{sign}_\varepsilon(A(\theta_j^+) - A(u)) = \frac{|A(u_{j+1}) - A(u)|_\varepsilon - |A(u_j) - A(u)|_\varepsilon}{A(u_{j+1}) - A(u_j)},$$

and the result follows by letting $\varepsilon \downarrow 0$. Let us prove the second statement. First observe that all expressions are symmetric in a and b , so we may assume that $a < b$. Under this assumption we have

$$\begin{aligned} (b-a)\text{sg}_{(a,b)}(\sigma) &= |a-\sigma| - |b-\sigma| \\ &= \text{sign}(a-\sigma)(a-\sigma) - \text{sign}(b-\sigma)(b-\sigma) \\ &= \begin{cases} \text{sign}(b-\sigma)(a-b) & \text{if } \sigma \notin (a, b), \\ 2\sigma - (b+a) & \text{if } \sigma \in (a, b). \end{cases} \end{aligned}$$

Dividing by $(b-a)$ concludes the proof. $\square$

Lemma 4.10. *Let $E_{\Delta x}^\varepsilon$ and φ be defined as in Lemma 4.6. For each $(x, t) \in \Pi_T$,*

$$\begin{aligned} & \liminf_{\varepsilon \downarrow 0} \int_{\Pi_T} E_{\Delta x}^\varepsilon \varphi \, dy \, ds \\ & \geq \int_{\Pi_T} D_- \left(D_+ \operatorname{sign}(A(u_{\Delta x}) - A(u)) \right. \\ & \quad \times \left. \left[\frac{1}{2} (A(u_{\Delta x}) + A(S_{\Delta x} u_{\Delta x})) - A(u) \right] \right) \varphi \, dy \, ds \\ & \quad + \liminf_{\varepsilon \downarrow 0} \frac{1}{2} \int_{\Pi_T} (\zeta^\varepsilon(u_{\Delta x}, \tau_{\Delta x}^-, u) + \zeta^\varepsilon(u_{\Delta x}, \tau_{\Delta x}^+, u)) (A(u)_y)^2 \varphi \, dy \, ds, \end{aligned}$$

where

$$\zeta^\varepsilon(a, b, c) := \operatorname{sign}'_\varepsilon(A(a) - A(c)) - \operatorname{sign}'_\varepsilon(A(b) - A(c)), \quad \forall a, b, c \in \mathbb{R}.$$

Proof. We split the proof into two claims.

Claim 1.

$$\begin{aligned} E_j^\varepsilon & \geq \frac{1}{2(\Delta x)^2} \int_{u_{j+1}}^{u_j} \zeta^\varepsilon(z, \tau_j^+, u) \partial_z (A(z) - A(u_{j+1}))^2 \, dz \\ & \quad + \frac{1}{2(\Delta x)^2} \int_{u_{j-1}}^{u_j} \zeta^\varepsilon(z, \tau_j^-, u) \partial_z (A(z) - A(u_{j-1}))^2 \, dz \\ & \quad + \frac{1}{2} (\zeta^\varepsilon(u_j, \tau_j^-, u) + \zeta^\varepsilon(u_j, \tau_j^+, u)) (A(u)_y)^2. \end{aligned}$$

Proof of Claim 1. Let

$$\begin{aligned} T^- & := \frac{1}{(\Delta x)^2} \int_{u_{j-1}}^{u_j} \psi_\varepsilon''(z, u) (A(z) - A(u_{j-1})) \, dz \\ & \quad - \frac{1}{\Delta x} \int_{u_{j-1}}^{u_j} \psi_\varepsilon''(z, u) \, dz A(u)_y + \frac{1}{2} \partial_y \psi_\varepsilon'(u, u_j) A(u)_y. \end{aligned}$$

We start by rewriting the first term as follows:

$$\begin{aligned} & \frac{1}{(\Delta x)^2} \int_{u_{j-1}}^{u_j} \psi_\varepsilon''(z, u) (A(z) - A(u_{j-1})) \, dz \\ & = \frac{1}{2(\Delta x)^2} \int_{u_{j-1}}^{u_j} \operatorname{sign}'_\varepsilon(A(z) - A(u)) \partial_z (A(z) - A(u_{j-1}))^2 \, dz \\ & = \frac{1}{2(\Delta x)^2} \int_{u_{j-1}}^{u_j} \operatorname{sign}'_\varepsilon(A(\tau_j^-) - A(u)) \partial_z (A(z) - A(u_{j-1}))^2 \, dz \\ & \quad + \frac{1}{2(\Delta x)^2} \int_{u_{j-1}}^{u_j} \zeta^\varepsilon(z, \tau_j^-, u) \partial_z (A(z) - A(u_{j-1}))^2 \, dz \\ & = \frac{1}{2} \operatorname{sign}'_\varepsilon(A(\tau_j^-) - A(u)) (D_- A(u_j))^2 \\ & \quad + \frac{1}{2(\Delta x)^2} \int_{u_{j-1}}^{u_j} \zeta^\varepsilon(z, \tau_j^-, u) \partial_z (A(z) - A(u_{j-1}))^2 \, dz. \end{aligned}$$

Concerning the second term in the definition of T^- , Lemma 4.8 gives

$$\begin{aligned} -\frac{1}{\Delta x} \int_{u_{j-1}}^{u_j} \psi''_\varepsilon(z, u) dz A(u)_y &= -D_- \text{sign}_\varepsilon(A(u_j) - A(u)) A(u)_y \\ &= -\text{sign}'_\varepsilon(A(\tau_j^-) - A(u)) D_- A(u_j) A(u)_y. \end{aligned}$$

For the last term we simply add and subtract to obtain

$$\begin{aligned} \frac{1}{2} \partial_y \psi'_\varepsilon(u, u_j) A(u)_y &= \frac{1}{2} \text{sign}'_\varepsilon(A(\tau_j^-) - A(u)) (A(u)_y)^2 \\ &\quad + \frac{1}{2} \zeta^\varepsilon(u_j, \tau_j^-, u) (A(u)_y)^2. \end{aligned}$$

Hence,

$$\begin{aligned} T^- &= \frac{1}{2} \text{sign}'_\varepsilon(A(\tau_j^-) - A(u)) (D_- A(u_j) - A(u)_y)^2 \\ &\quad + \frac{1}{2(\Delta x)^2} \int_{u_{j-1}}^{u_j} \zeta^\varepsilon(z, \tau_j^-, u) \partial_z (A(z) - A(u_{j-1}))^2 dz \\ &\quad + \frac{1}{2} \zeta^\varepsilon(u_j, \tau_j^-, u) (A(u)_y)^2. \end{aligned}$$

Define

$$\begin{aligned} T^+ &:= \frac{1}{(\Delta x)^2} \int_{u_{j+1}}^{u_j} \psi''_\varepsilon(z, u) (A(z) - A(u_{j+1})) dz \\ &\quad - \frac{1}{\Delta x} \int_{u_{j+1}}^{u_j} \psi''_\varepsilon(z, u) dz A(u)_y + \frac{1}{2} \partial_y \psi'_\varepsilon(u, u_j) A(u)_y. \end{aligned}$$

Using the same strategy as above we arrive at

$$\begin{aligned} T^+ &= \frac{1}{2} \text{sign}'_\varepsilon(A(\tau_j^+) - A(u)) (D_+ A(u_j) - A(u)_y)^2 \\ &\quad + \frac{1}{2(\Delta x)^2} \int_{u_{j+1}}^{u_j} \zeta^\varepsilon(z, \tau_j^+, u) \partial_z (A(z) - A(u_{j+1}))^2 dz \\ &\quad + \frac{1}{2} \zeta^\varepsilon(u_j, \tau_j^+, u) (A(u)_y)^2. \end{aligned}$$

Note that $E_j^\varepsilon = T^- + T^+$, so Claim 1 follows by removing the nonnegative terms on the righthand side. $\square$

Claim 2. Suppose that $x \in I_j$. Then

(4.4)

$$\begin{aligned} \liminf_{\varepsilon \downarrow 0} \frac{1}{2(\Delta x)^2} \int_{\Pi_T} \left[\int_{u_{j-1}}^{u_j} \zeta^\varepsilon(z, \tau_j^-, u) \frac{d}{dz} (A(z) - A(u_{j-1}))^2 dz \right. \\ \left. + \int_{u_{j+1}}^{u_j} \zeta^\varepsilon(z, \tau_j^+, u) \frac{d}{dz} (A(z) - A(u_{j+1}))^2 dz \right] \varphi dy ds \\ = \int_{\Pi_T} D_- \left(D_+ \text{sign}(A(u_j) - A(u)) \left[\frac{1}{2} (A(u_j) + A(u_{j+1})) - A(u) \right] \right) \varphi dy ds. \end{aligned}$$

Proof of Claim 2. Let

$$\begin{aligned} T_1^\varepsilon &:= \frac{1}{2(\Delta x)^2} \int_{u_{j-1}}^{u_j} \text{sign}'_\varepsilon(A(z) - A(u)) \frac{d}{dz} (A(z) - A(u_{j-1}))^2 dz, \\ K_1^\varepsilon &:= \frac{1}{2(\Delta x)^2} \int_{u_{j-1}}^{u_j} \text{sign}'_\varepsilon(A(\tau_j^-) - A(u)) \frac{d}{dz} (A(z) - A(u_{j-1}))^2 dz, \\ T_2^\varepsilon &:= \frac{1}{2(\Delta x)^2} \int_{u_{j+1}}^{u_j} \text{sign}'_\varepsilon(A(z) - A(u)) \frac{d}{dz} (A(z) - A(u_{j+1}))^2 dz, \\ K_2^\varepsilon &:= \frac{1}{2(\Delta x)^2} \int_{u_{j+1}}^{u_j} \text{sign}'_\varepsilon(A(\tau_j^+) - A(u)) \frac{d}{dz} (A(z) - A(u_{j+1}))^2 dz, \end{aligned}$$

and note that the left-hand side of (4.4) may be written as

$$\liminf_{\varepsilon \downarrow 0} \int_{\Pi_T} \left((T_1^\varepsilon - K_1^\varepsilon) + (T_2^\varepsilon - K_2^\varepsilon) \right) \varphi dy ds.$$

Let us rewrite T_1^ε as follows:

$$\begin{aligned} T_1^\varepsilon &= \frac{1}{(\Delta x)^2} \int_{u_{j-1}}^{u_j} \text{sign}'_\varepsilon(A(z) - A(u)) A'(z) (A(z) - A(u_{j-1})) dz \\ &= \frac{1}{(\Delta x)^2} \int_{u_{j-1}}^{u_j} \text{sign}'_\varepsilon(A(z) - A(u)) A'(z) (A(u) - A(u_{j-1})) dz \\ &\quad + \frac{1}{(\Delta x)^2} \int_{u_{j-1}}^{u_j} \text{sign}'_\varepsilon(A(z) - A(u)) A'(z) (A(z) - A(u)) dz \\ &= D_- \text{sign}_\varepsilon(A(u_j) - A(u)) \frac{(A(u) - A(u_{j-1}))}{\Delta x} + R_1^\varepsilon, \end{aligned}$$

where

$$\begin{aligned} R_1^\varepsilon &:= \frac{1}{\Delta x^2} \left[\text{sign}_\varepsilon(A(z) - A(u)) (A(z) - A(u)) \Big|_{z=u_{j-1}}^{z=u_j} \right. \\ &\quad \left. - \int_{u_{j-1}}^{u_j} \frac{d}{dz} |A(z) - A(u)|_\varepsilon dz \right]. \end{aligned}$$

Concerning K_1^ε , we apply Lemma 4.8 to obtain

$$\begin{aligned} K_1^\varepsilon &= \frac{1}{2(\Delta x)^2} \int_{u_{j-1}}^{u_j} \text{sign}'_\varepsilon(A(\tau_j^-) - A(u)) \partial_z (A(z) - A(u_{j-1}))^2 dz \\ &= \frac{1}{2} \text{sign}'_\varepsilon(A(\tau_j^-) - A(u)) (D_- A(u_j))^2 dz \\ &= \frac{1}{2} D_- \text{sign}_\varepsilon(A(u_j) - A(u)) D_- A(u_j). \end{aligned}$$

It now follows that

$$T_1^\varepsilon - K_1^\varepsilon = -\frac{1}{\Delta x} D_- \text{sign}_\varepsilon(A(u_j) - A(u)) \left[\frac{1}{2} (A(u_j) + A(u_{j-1})) - A(u) \right] + R_1^\varepsilon.$$

Performing the same type of computations as above yields

$$T_2^\varepsilon - K_2^\varepsilon = \frac{1}{\Delta x} D_+ \text{sign}_\varepsilon(A(u_j) - A(u)) \left[\frac{1}{2} (A(u_{j+1}) + A(u_j)) - A(u) \right] + R_2^\varepsilon,$$

where

$$R_2^\varepsilon := \frac{1}{\Delta x^2} \left[\int_{u_j}^{u_{j+1}} \frac{d}{dz} |A(z) - A(u)|_\varepsilon dz \right. \\ \left. - \operatorname{sign}_\varepsilon(A(z) - A(u))(A(z) - A(u)) \Big|_{z=u_j}^{z=u_{j+1}} \right].$$

Next, observe that

$$R_1^\varepsilon = \frac{1}{\Delta x^2} \left[\int_{z=u_{j-1}}^{z=u_j} \operatorname{sign}_\varepsilon(A(z) - A(u))(A(z) - A(u)) - |A(z) - A(u)|_\varepsilon \right],$$

so $\lim_{\varepsilon \downarrow 0} R_1^\varepsilon = 0$. The same considerations apply to R_2^ε so $\lim_{\varepsilon \downarrow 0} R_2^\varepsilon = 0$ also. Claim 2 follows from an application of the dominated convergence theorem. Finally, combining Claim 1 and Claim 2 finishes the proof. $\square$

4.1. Estimates. The purpose of this section is to find bounds on the “unwanted” terms in inequality (4.2) and Lemma 4.10. Throughout this section the notation is the one given in Lemma 4.6. We let C denote a generic constant. By constant it is meant that it does not depend on the “small” variables but it might depend on T and the initial data. For any set A , let $\mathbb{1}_A$ denote its characteristic function.

For future reference we collect some standard estimates in a lemma.

Lemma 4.11. *Let φ be defined in Lemma 4.6. Then*

$$\left| \frac{\partial^k}{\partial x^k} \varphi(x, t, y, s) \right| \leq \psi(t) \frac{\|\rho^{(k)}\|_{L^\infty}}{r^{k+1}} \mathbb{1}_{\{|x-y| \leq r\}}(x, y) \rho_{r_0}(t-s).$$

Recall that $S_\sigma \varphi(x, t, y, s) = \varphi(x + \sigma, t, y, s)$. If $|\sigma| \leq \Delta x$, then

$$\left| \frac{\partial^k}{\partial x^k} S_\sigma \varphi(x, t, y, s) \right| \leq \psi(t) \frac{\|\rho^{(k)}\|_{L^\infty}}{r^{k+1}} \mathbb{1}_{\{|x-y| \leq r+\Delta x\}}(x, y) \rho_{r_0}(t-s).$$

Considering the difference quotient applied to ω_r we have

$$|D_+ \omega_r(x - y)| \leq \frac{\|\rho'\|_{L^\infty}}{r^2} \mathbb{1}_{\{|x-y| \leq r+\Delta x\}}(x, y).$$

Proof. Note that

$$\frac{\partial^k}{\partial x^k} \omega_r(x) = \frac{1}{r^{k+1}} \rho^{(k)}\left(\frac{x}{r}\right).$$

Since $\operatorname{supp}(\rho) \subset [-1, 1]$ we have

$$\left| \frac{\partial^k}{\partial x^k} \omega_r(x) \right| \leq \frac{\|\rho^{(k)}\|_{L^\infty}}{r^{k+1}} \mathbb{1}_{\{|x| \leq r\}}(x),$$

which proves the first statement.

Consider the second statement. If $|x - y| \geq r + \Delta x$, then

$$|x + \sigma - y| \geq |x - y| - |\sigma| \geq r + \Delta x - \Delta x = r,$$

so it follows that $\mathbb{1}_{\{|x+\sigma-y| \leq r\}}(x, y) \leq \mathbb{1}_{\{|x-y| \leq r+\Delta x\}}(x, y)$; this proves the second statement.

To prove the last statement, recall that

$$D_+ \omega_r(x) = \frac{\omega_r(x + \Delta x) - \omega_r(x)}{\Delta x}.$$

If $|x| \geq r + \Delta x$, then $\omega_r(x + \Delta x) = \omega_r(x) = 0$, so $\text{supp}(D_+(\omega_r)) \subset [-r - \Delta x, r + \Delta x]$. By the mean value theorem and the fact that $\|\omega'_r\|_{L^\infty} = \|\rho'\|_{L^\infty} r^{-2}$ we get

$$|\omega_r(x + \Delta x) - \omega_r(x)| \leq \frac{\|\rho'\|_{L^\infty}}{r^2} \Delta x.$$

The last statement follows from this. $\square$

Estimate 4.1.

$$\left| \iint_{\Pi_T^2} \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)) (D_+\varphi + \varphi_y) dX \right| \leq C \frac{\Delta x}{r} \left(1 + \frac{\Delta x}{r} \right).$$

Proof. Let

$$\beta := \iint_{\Pi_T^2} \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)) (D_+\varphi + \varphi_y) dX.$$

First note that

$$D_+\varphi + \varphi_y = D_+\varphi - \varphi_x.$$

We claim that

$$(4.5) \quad (D_+\varphi - \varphi_x)(x, t, y, s) = \frac{1}{\Delta x} \int_0^{\Delta x} (\Delta x - \sigma) \varphi_{xx}(x + \sigma, t, y, s) d\sigma.$$

Hence

$$\beta = \frac{1}{\Delta x} \iint_{\Pi_T^2} \int_0^{\Delta x} \text{sign}_\varepsilon(A(u_{\Delta x}) - A(u)) (f(u_{\Delta x}) - f(u)) (\Delta x - \sigma) S_\sigma \varphi_{xx} d\sigma dX.$$

We can write

$$\begin{aligned} & \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)) (x, t, y, s) \\ &= \sum_j \underbrace{\text{sign}(u_j - u) (f(u_j) - f(u)) (t, y, s)}_{\Theta_j} \mathbb{1}_{\{I_j\}}(x). \end{aligned}$$

Using summation by parts we get

$$\begin{aligned} & \frac{1}{\Delta x} \int_{\mathbb{R}} \int_0^{\Delta x} \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)) (\Delta x - \sigma) S_\sigma \varphi_{xx} d\sigma dx \\ &= \frac{1}{\Delta x} \int_0^{\Delta x} \sum_j \Theta_j \int_{\mathbb{R}} \mathbb{1}_{\{I_j\}}(x) (\Delta x - \sigma) S_\sigma \varphi_{xx} dx d\sigma \\ &= \frac{1}{\Delta x} \int_0^{\Delta x} \sum_j \Theta_j \int_{I_j} \varphi_{xx}(x + \sigma, t, y, s) dx (\Delta x - \sigma) d\sigma \\ &= \int_0^{\Delta x} \sum_j \Theta_j (D_- S_\sigma \varphi_{x,j+1/2}) (\Delta x - \sigma) d\sigma \\ &= - \sum_j D_+ \Theta_j \int_0^{\Delta x} S_\sigma \varphi_{x,j+1/2} (\Delta x - \sigma) d\sigma, \end{aligned}$$

where $S_\sigma \varphi_{x,j+1/2} = \varphi_x(x_{j+1/2} + \sigma, t, y, s)$. By Lemma 4.11 we have

$$|\varphi_x(x + \sigma, t, y, s)| \leq C \frac{1}{r^2} \mathbb{1}_{\{|x-y| \leq r+\Delta x\}}(x, y) \rho_{r_0}(t-s).$$

Hence,

$$\left| \int_0^{\Delta x} S_\sigma \varphi_{x,j+1/2}(\Delta x - \sigma) d\sigma \right| \leq C \frac{\Delta x^2}{r^2} \mathbb{1}_{\{|x_{j+1/2}-y| \leq r+\Delta x\}}(y) \rho_{r_0}(t-s).$$

Now

$$|D_+ \Theta_j| \leq \|f\|_{\text{Lip}} |D_+ u_j|.$$

Therefore,

$$\begin{aligned} & \left| \frac{1}{\Delta x} \int_{\mathbb{R}} \int_0^{\Delta x} \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)) (\Delta x - \sigma) \varphi_{xx}^\sigma d\sigma dx \right| \\ & \leq \sum_j |D_+ \Theta_j| \left| \int_0^{\Delta x} S_\sigma \varphi_{x,j+1/2}(\Delta x - \sigma) d\sigma \right| \\ & \leq C \|f\|_{\text{Lip}} \sum_j |D_+ u_j| \frac{\Delta x^2}{r^2} \mathbb{1}_{\{|x_{j+1/2}-y| \leq r+\Delta x\}}(y) \rho_{r_0}(t-s). \end{aligned}$$

It follows by the above and Lemma 3.1 that

$$\begin{aligned} |\beta| & \leq C \Delta x^2 \frac{r + \Delta x}{r^2} \int_0^T \sum_j |D_+ u_j| dt \\ & = C \frac{r + \Delta x}{r^2} \int_{\Pi_T} |u_{\Delta x}(x + \Delta x, t) - u_{\Delta x}(x, t)| dx dt \\ & = CT \frac{1}{r} \left(1 + \frac{\Delta x}{r} \right) \Delta x |u_{\Delta x}^0|_{BV(\mathbb{R})}. \end{aligned}$$

This concludes the proof. $\square$

Estimate 4.2.

$$\begin{aligned} & \left| \iint_{\Pi_T^2} |A(u_{\Delta x}) - A(u)| (D_- D_+ \varphi + (D_+ + D_-) \varphi_y + \varphi_{yy}) dX \right| \\ & \leq C \frac{\Delta x^2}{r^3} \left(1 + \frac{\Delta x}{r} \right). \end{aligned}$$

Proof. Since $\varphi_{xx} + 2\varphi_{xy} + \varphi_{yy} = 0$ it follows that

$$D_- D_+ \varphi + (D_+ + D_-) \varphi_y + \varphi_{yy} = (D_- D_+ \varphi - \varphi_{xx}) + ((D_+ + D_-) \varphi - 2\varphi_x)_y.$$

Thus

$$\begin{aligned} & \iint_{\Pi_T^2} |A(u_{\Delta x}) - A(u)| (D_- D_+ \varphi + (D_+ + D_-) \varphi_y + \varphi_{yy}) dX \\ & = \iint_{\Pi_T^2} |A(u_{\Delta x}) - A(u)| (D_- D_+ \varphi - \varphi_{xx}) dX \\ & \quad + \iint_{\Pi_T^2} |A(u_{\Delta x}) - A(u)| ((D_+ + D_-) \varphi - 2\varphi_x)_y dX \\ & =: \zeta_1 + \zeta_2. \end{aligned}$$

Consider the term ζ_1 . We use the same strategy as in Estimate 4.1. Writing $\mu(\sigma) = \varphi(x + \sigma, t, y, s)$, a Taylor expansion gives

$$\mu(z) - \mu(0) = z\mu'(0) + \frac{1}{2}z^2\mu''(0) + \frac{1}{6}z^3\mu^{(3)}(0) - \frac{1}{6} \int_0^z (\sigma - z)^3 \mu^{(4)}(\sigma) d\sigma.$$

Using this, we get

$$\begin{aligned} & \mu(\Delta x) - 2\mu(0) + \mu(-\Delta x) - \Delta x^2 \mu''(0) \\ &= -\frac{1}{6} \int_0^{\Delta x} (\sigma - \Delta x)^3 \mu^{(4)}(\sigma) d\sigma + \frac{1}{6} \int_{-\Delta x}^0 (\sigma + \Delta x)^3 \mu^{(4)}(\sigma) d\sigma. \end{aligned}$$

It follows that

$$\begin{aligned} D_+ D_- \varphi - \varphi_{xx} &= -\frac{1}{6\Delta x^2} \int_0^{\Delta x} (\sigma - \Delta x)^3 \frac{\partial^4}{\partial x^4} \varphi(x + \sigma, t, y, s) d\sigma \\ &\quad + \frac{1}{6\Delta x^2} \int_{-\Delta x}^0 (\sigma + \Delta x)^3 \frac{\partial^4}{\partial x^4} \varphi(x + \sigma, t, y, s) d\sigma. \end{aligned}$$

Splitting ζ_1 according to this equality we get

$$\begin{aligned} \zeta_1 &= \iint_{\Pi_T^2} |A(u_{\Delta x}) - A(u)| (D_- D_+ \varphi - \varphi_{xx}) dX \\ &= -\frac{1}{6\Delta x^2} \iint_{\Pi_T^2} \int_0^{\Delta x} |A(u_{\Delta x}) - A(u)| (\sigma - \Delta x)^3 \frac{\partial^4}{\partial x^4} \varphi(x + \sigma, t, y, s) d\sigma dX \\ &\quad + \frac{1}{6\Delta x^2} \iint_{\Pi_T^2} \int_{-\Delta x}^0 |A(u_{\Delta x}) - A(u)| (\sigma + \Delta x)^3 \frac{\partial^4}{\partial x^4} \varphi(x + \sigma, t, y, s) d\sigma dX \\ &=: \zeta_{1,1} + \zeta_{1,2}. \end{aligned}$$

We also have that

$$|A(u_{\Delta x}) - A(u)|(x, t, y, s) = \sum_j \underbrace{|A(u_j) - A(u)|(t, y, s)}_{\Phi_j} \mathbb{1}_{\{I_j\}}(x).$$

Consider $\zeta_{1,1}$,

$$\begin{aligned} & -\int_0^{\Delta x} \int_{\mathbb{R}} |A(u_{\Delta x}) - A(u)| (\sigma - \Delta x)^3 \frac{\partial^4}{\partial x^4} S_\sigma \varphi dx d\sigma \\ &= -\sum_j |A(u_j) - A(u)|(t, y, s) \int_0^{\Delta x} (\sigma - \Delta x)^3 \int_{\mathbb{R}} \mathbb{1}_{\{I_j\}}(x) \frac{\partial^4}{\partial x^4} S_\sigma \varphi dx d\sigma \\ &= -\Delta x \int_0^{\Delta x} (\sigma - \Delta x)^3 \sum_j \Phi_j D_- \varphi_{xxx, j+1/2}^\sigma d\sigma \\ &= \Delta x \sum_j D_+ \Phi_j \int_0^{\Delta x} (\sigma - \Delta x)^3 S_\sigma \varphi_{xxx, j+1/2} d\sigma, \end{aligned}$$

where

$$S_\sigma \varphi_{xxx, j+1/2}(t, y, s) = \frac{\partial^3}{\partial x^3} \varphi(x_{j+1/2} + \sigma, t, y, s).$$

Now we use Lemma 4.11 to estimate this term as follows:

$$\begin{aligned} |\zeta_{1,1}| &= \left| \frac{1}{6\Delta x^2} \iint_{\Pi_T^2} \int_0^{\Delta x} |A(u_{\Delta x}) - A(u)| (\sigma - \Delta x)^3 \frac{\partial^4}{\partial x^4} \varphi(x + \sigma, t, y, s) d\sigma dX \right| \\ &= \left| \frac{1}{6\Delta x} \int_{\Pi_T} \int_0^T \sum_j D_+ \Phi_j \int_0^{\Delta x} (\sigma - \Delta x)^3 S_\sigma \varphi_{xxx,j+1/2} d\sigma dt dy ds \right| \\ &\leq C \frac{r + \Delta x}{\Delta x^2 r^4} \int_{\Pi_T} |D_+ A(u_{\Delta x})| \left(\int_0^{\Delta x} (\sigma - \Delta x)^3 d\sigma \right) dx dt \\ &\leq C \Delta x^2 \frac{r + \Delta x}{r^4} = C \frac{\Delta x^2}{r^3} \left(1 + \frac{\Delta x}{r} \right), \end{aligned}$$

where we have used that $|A(u_{\Delta x}(\cdot, t))|_{BV(\mathbb{R})}$ is bounded independently of $\Delta x, t, \eta$ by Lemma 3.1. The term $\zeta_{1,2}$ is estimated in a similar way.

Consider ζ_2 . Again, let $\mu(\sigma) = \varphi(x + \sigma, t, y, s)$. Then

$$(D_+ + D_-)\varphi - 2\varphi_x = \frac{1}{\Delta x} [\mu(\Delta x) - \mu(-\Delta x) - 2\Delta x \mu'(0)].$$

By a Taylor expansion

$$\mu(z) - \mu(0) = z\mu'(0) + \frac{1}{2}z^2\mu''(0) + \frac{1}{2} \int_0^z (\sigma - z)^2 \mu^{(3)}(\sigma) d\sigma.$$

Writing $z = \pm\Delta x$ and subtracting the corresponding equations we obtain

$$\begin{aligned} (D_+ + D_-)\varphi - 2\varphi_x &= \frac{1}{2\Delta x} \int_0^{\Delta x} (\sigma - \Delta x)^2 \frac{\partial^3}{\partial x^3} \varphi(x + \sigma, t, y, s) d\sigma \\ &\quad + \frac{1}{2\Delta x} \int_{-\Delta x}^0 (\sigma + \Delta x)^2 \frac{\partial^3}{\partial x^3} \varphi(x + \sigma, t, y, s) d\sigma. \end{aligned}$$

We may split ζ_2 into the two terms

$$\begin{aligned} \zeta_2 &= \frac{1}{2\Delta x} \iint_{\Pi_T^2} \int_0^{\Delta x} |A(u_{\Delta x}) - A(u)| (\sigma - \Delta x)^2 \frac{\partial^3}{\partial x^3} \frac{\partial}{\partial y} \varphi(x + \sigma, t, y, s) d\sigma dX \\ &\quad + \frac{1}{2\Delta x} \iint_{\Pi_T^2} \int_{-\Delta x}^0 |A(u_{\Delta x}) - A(u)| (\sigma + \Delta x)^2 \frac{\partial^3}{\partial x^3} \frac{\partial}{\partial y} \varphi(x + \sigma, t, y, s) d\sigma dX \\ &=: \zeta_{2,1} + \zeta_{2,2}. \end{aligned}$$

Performing integration by parts, $\zeta_{2,1}$ becomes

$$\frac{1}{2\Delta x} \iint_{\Pi_T^2} \int_0^{\Delta x} \text{sign}(A(u_{\Delta x}) - A(u)) A(u)_y (\sigma - \Delta x)^2 \frac{\partial^3}{\partial x^3} \varphi(x + \sigma, t, y, s) d\sigma dX.$$

Thus, by Lemma 4.11,

$$\begin{aligned} |\zeta_{2,1}| &\leq \frac{1}{2\Delta x} \iint_{\Pi_T^2} |A(u)_y| \left| \int_0^{\Delta x} (\sigma - \Delta x)^2 \frac{\partial^3}{\partial x^3} \varphi(x + \sigma, t, y, s) d\sigma \right| dX \\ &\leq TC \frac{r + \Delta x}{r^4 \Delta x} \left(\int_0^{\Delta x} (\sigma - \Delta x)^2 d\sigma \right) \int_{\Pi_T} |A(u)_y| dy ds \\ &\leq C \frac{\Delta x^2}{r^3} \left(1 + \frac{\Delta x}{r} \right), \end{aligned}$$

as $|A(u(\cdot, s))|_{BV(\mathbb{R})} \leq |A(u^0(\cdot))|_{BV(\mathbb{R})}$ for all s . The same estimate holds for $\zeta_{2,2}$. $\square$

Estimate 4.3.

$$\left| \iint_{\Pi_T^2} \left(\int_{u_{\Delta x}}^{S_{\Delta x} u_{\Delta x}} \operatorname{sign}(z - u) F_2'(z) dz \right) D_+ \varphi dX \right| \leq C \frac{\Delta x}{r} \left(1 + \frac{\Delta x}{r} \right).$$

Proof. By definition F_2' is bounded. Hence,

$$\left| \int_{u_j}^{u_{j+1}} \operatorname{sign}(z - u) F_2'(z) dz \right| \leq \|F_2\|_{\text{Lip}} \Delta x |D_+ u_j|.$$

Note that $|u_{\Delta x}(\cdot, t)|_{BV(\mathbb{R})}$ is bounded independently of $\Delta x, t, \eta$ by Lemma 3.1, so we may apply Lemma 4.11 to obtain the result. $\square$

Next, we consider the terms from Lemma 4.10.

Estimate 4.4.

$$\begin{aligned} & \iint_{\Pi_T^2} D_- \left(D_+ \operatorname{sign}(A(u_j) - A(u)) \left[\frac{1}{2}(A(u_j) + A(u_{j+1})) - A(u) \right] \right) \varphi dX \\ & \geq -C(1 + r + \Delta x) \frac{\Delta x}{r^2} \left(1 + \frac{\Delta x}{r} \right)^3. \end{aligned}$$

Proof. Let us first show that

$$(4.6) \quad \left| D_+ \operatorname{sign}(A(u_j) - A(u)) \left[\frac{1}{2}(A(u_j) + A(u_{j+1})) - A(u) \right] \right| \leq \Delta x D_+ \operatorname{sign}(A(u_j) - A(u)) D_+(A(u_j)).$$

First note that

$$\begin{aligned} & D_+ \operatorname{sign}(A(u_j) - A(u)) \\ & = \frac{2}{\Delta x} \operatorname{sign}(A(u_j) - A(u_{j+1})) \mathbb{1}_{\{A(u) \in \operatorname{int}(A(u_j), A(u_{j+1}))\}}, \end{aligned}$$

so the left-hand side of (4.6) is zero whenever $A(u) \notin \operatorname{int}(A(u_j), A(u_{j+1}))$. Second, if $c \in \operatorname{int}(a, b)$, then it follows that

$$\left| \frac{1}{2}(a + b) - c \right| = \frac{1}{2}(|b - c| + |a - c|) \leq |b - a|.$$

Since $z \mapsto \operatorname{sign}(A(z) - A(u))$ is increasing, the right-hand side is positive. This proves (4.6).

Performing integration by parts we obtain

$$\begin{aligned} & \left| \iint_{\Pi_T^2} D_- \left(D_+ \operatorname{sign}(A(u_j) - A(u)) \left[\frac{1}{2}(A(u_j) + A(u_{j+1})) - A(u) \right] \right) \varphi dX \right| \\ & \leq \iint_{\Pi_T^2} \left| D_+ \operatorname{sign}(A(u_j) - A(u)) \left[\frac{1}{2}(A(u_j) + A(u_{j+1})) - A(u) \right] \right| |D_+ \varphi| dX \\ & \leq \Delta x \iint_{\Pi_T^2} D_+ \operatorname{sign}(A(u_j) - A(u)) D_+(A(u_j)) |D_+ \varphi| dX. \end{aligned}$$

Using integration by parts for difference quotients and the Leibniz rule for difference quotients, we obtain

$$\begin{aligned}
 & \iint_{\Pi_T^2} D_+ \text{sign}_\varepsilon(A(u_{\Delta x}) - A(u)) D_+ A(u_{\Delta x}) |D_+ \varphi| \, dX \\
 &= - \iint_{\Pi_T^2} \text{sign}_\varepsilon(A(u_{\Delta x}) - A(u)) D_+ A(u_{\Delta x}) D_- |D_+ \varphi| \, dX \\
 &\quad - \iint_{\Pi_T^2} \text{sign}_\varepsilon(A(u_{\Delta x}) - A(u)) D_- D_+ A(u_{\Delta x}) |D_- \varphi| \, dX \\
 &=: \zeta_1 + \zeta_2.
 \end{aligned}$$

To estimate ζ_1 we first observe that $D_- |D_+ \varphi| \leq |D_+ D_- \varphi|$. Furthermore, when proving Estimate 4.2, we established that

$$\begin{aligned}
 & D_+ D_- \varphi(x, t, y, s) \\
 &= \varphi_{xx}(x, t, y, s) - \frac{1}{6\Delta x^2} \int_0^{\Delta x} (\sigma - \Delta x)^3 \frac{\partial^4}{\partial x^4} \varphi(x + \sigma, t, y, s) \, d\sigma \\
 &\quad + \frac{1}{6\Delta x^2} \int_{-\Delta x}^0 (\sigma + \Delta x)^3 \frac{\partial^4}{\partial x^4} \varphi(x + \sigma, t, y, s) \, d\sigma.
 \end{aligned}$$

By Lemma 4.11,

$$\begin{aligned}
 & \left| \int_0^{\pm \Delta x} (\sigma \mp \Delta x)^3 \frac{\partial^4}{\partial x^4} \varphi(x + \sigma, t, y, s) \, d\sigma \right| \\
 &\leq C \frac{(\Delta x)^4}{r^5} \mathbb{1}_{\{|x-y| \leq r+\Delta x\}}(x, y) \rho_{r_0}(t-s).
 \end{aligned}$$

Using Lemma 4.11 once more, the above implies that

$$\begin{aligned}
 \int_{\Pi_T} |D_+ D_- \varphi| \, dyds &\leq \int_{\Pi_T} |\varphi_{xx}| \, dyds + C \frac{\Delta x^2}{r^4} \left(1 + \frac{\Delta x}{r}\right) \\
 &\leq C \left(\frac{1}{r^2} + \frac{\Delta x^2}{r^4} \right) \left(1 + \frac{\Delta x}{r}\right).
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 |\zeta_1| &= \left| \iint_{\Pi_T^2} \text{sign}_\varepsilon(A(u_{\Delta x}) - A(u)) D_+ A(u_{\Delta x}) D_- |D_- \varphi| \, dX \right| \\
 &\leq \int_{\Pi_T} |D_+ A(u_{\Delta x})| \left(\int_{\Pi_T} |D_+ D_- \varphi| \, dyds \right) dxdt \\
 &\leq C \left(\frac{1}{r^2} + \frac{\Delta x^2}{r^4} \right) \left(1 + \frac{\Delta x}{r}\right) \int_{\Pi_T} |D_+ A(u_{\Delta x})| \, dxdt.
 \end{aligned}$$

Recall that $|A(u_{\Delta x}(\cdot, t))|_{BV(\mathbb{R})}$ is bounded independently of $\Delta x, t, \eta$ by Lemma 3.1.

Concerning ζ_2 we have

$$\begin{aligned} |\zeta_2| &= \left| \iint_{\Pi_T^2} \operatorname{sign}_\varepsilon(A(u_{\Delta x}) - A(u)) (D_- D_+ A(u_{\Delta x})) |D_- \varphi| dX \right| \\ &\leq \iint_{\Pi_T^2} |D_- D_+ A(u_{\Delta x})| |D_- \varphi| dX \\ &\leq C \frac{r + \Delta x}{r^2} \int_{\Pi_T} |D_- D_+ A(u_{\Delta x})| dx dt. \end{aligned}$$

Note that it follows from (3.1) and Lemma 3.2 that $\|D_- D_+ A(u_{\Delta x}(\cdot, t))\|_{L^1(\mathbb{R})}$ is bounded independently of $\Delta x, t, \eta$. Hence,

$$\begin{aligned} \Delta x \iint_{\Pi_T^2} D_+ \operatorname{sign}_\varepsilon(A(u_{\Delta x}) - A(u)) D_+ A(u_{\Delta x}) |D_+ \varphi| dX \\ \leq \Delta x (|\zeta_1| + |\zeta_2|) \\ \leq C(1 + r + \Delta x) \left(\frac{\Delta x}{r^2} + \frac{\Delta x^3}{r^4} \right) \left(1 + \frac{\Delta x}{r} \right) \\ \leq C(1 + r + \Delta x) \frac{\Delta x}{r^2} \left(1 + \frac{\Delta x}{r} \right)^3. \quad \square \end{aligned}$$

Estimate 4.5.

$$\begin{aligned} \int_{\Pi_T} \left(\liminf_{\varepsilon \downarrow 0} \int_{\Pi_T} \frac{1}{2} (\zeta^\varepsilon(u_{\Delta x}, \tau_{\Delta x}^-, u) + \zeta^\varepsilon(u_{\Delta x}, \tau_{\Delta x}^+, u)) (A(u)_y)^2 \varphi dy ds \right) dx dt \\ \geq -C \left(\frac{\Delta x}{r_0} + \frac{\Delta x}{r} + \frac{\Delta x}{r^2} \right). \end{aligned}$$

Proof. Set

$$\begin{aligned} \mathcal{R}_j^\varepsilon &:= (\zeta^\varepsilon(u_j, \tau_j^-, u) + \zeta^\varepsilon(u_j, \tau_j^+, u)) (A(u)_y)^2 \\ &= (\operatorname{sign}'_\varepsilon(A(u_j) - A(u)) - \operatorname{sign}'_\varepsilon(A(\tau_j^-) - A(u))) (A(u)_y)^2 \\ &\quad + (\operatorname{sign}'_\varepsilon(A(u_j) - A(u)) - \operatorname{sign}'_\varepsilon(A(\tau_j^+) - A(u))) (A(u)_y)^2, \end{aligned}$$

and $\mathcal{R}_{\Delta x}^\varepsilon(x, t, y, s) = \mathcal{R}_j^\varepsilon(y, t, s)$ for $x \in I_j$. Note that the term we want to estimate may be written as

$$\begin{aligned} \liminf_{\varepsilon \downarrow 0} \int_{\Pi_T} \mathcal{R}_{\Delta x}^\varepsilon(x, t, y, s) \varphi(x, t, y, s) dy ds \\ = \sum_j \liminf_{\varepsilon \downarrow 0} \left(\int_{\Pi_T} \mathcal{R}_j^\varepsilon(t, y, s) \varphi(x, t, y, s) dy ds \right) \mathbb{1}_{\{I_j\}}(x). \end{aligned}$$

Let us define an entropy function by

$$\begin{aligned} \partial_u \Psi_\varepsilon(u, u_{j-1}, u_j, u_{j+1}) \\ := \operatorname{sign}_\varepsilon(A(\theta_j^-) - A(u)) - 2 \operatorname{sign}_\varepsilon(A(u_j) - A(u)) + \operatorname{sign}_\varepsilon(A(\theta_j^+) - A(u)). \end{aligned}$$

Recall that $\theta_j^\pm = \theta_j^\pm(u)$, so the above function is not as explicit as it appears. However, by Lemma 4.9 we are able to obtain an explicit expression for the limit as $\varepsilon \rightarrow 0$. To simplify the notation we write $\Psi'_{\varepsilon,j}(u)$ for $\partial_u \Psi_\varepsilon(u, u_{j-1}, u_j, u_{j+1})$. Let us also define the entropy flux functions

$$\Xi'_{\varepsilon,j}(u) = \Psi'_{\varepsilon,j}(u) f'(u), \quad \Phi'_{\varepsilon,j}(u) = \Psi'_{\varepsilon,j}(u) A'(u).$$

That is, $(\Psi_{\varepsilon,j}, \Xi_{\varepsilon,j}, \Phi_{\varepsilon,j})$ is an entropy-entropy flux triple.

Multiplying equation (1.1) by $\Psi'_{\varepsilon,j}(u)$ yields

$$\Psi_{\varepsilon,j}(u)_s + \Xi_{\varepsilon,j}(u)_y = \Phi_{\varepsilon,j}(u)_{yy} - \partial_y \Psi'_{\varepsilon,j}(u) A(u)_y.$$

By Lemma 4.8 we see that

$$\begin{aligned} \partial_y \Psi'_{\varepsilon,j}(u) A(u)_y &= \left[\text{sign}_{\varepsilon}(A(\theta_j^-) - A(u))_y - 2 \text{sign}_{\varepsilon}(A(u_j) - A(u))_y \right. \\ &\quad \left. + \text{sign}_{\varepsilon}(A(\theta_j^+) - A(u))_y \right] A(u)_y \\ &= - \left[\text{sign}'_{\varepsilon}(A(\tau_j^-) - A(u)) - 2 \text{sign}'_{\varepsilon}(A(u_j) - A(u)) \right. \\ &\quad \left. + \text{sign}'_{\varepsilon}(A(\tau_j^+) - A(u)) \right] (A(u)_y)^2 \\ (4.7) \qquad \qquad \qquad &= \mathcal{R}_j^{\varepsilon}. \end{aligned}$$

It follows that we can write

$$\begin{aligned} (4.8) \quad \int_{\Pi_T} \mathcal{R}_j^{\varepsilon} \varphi \, dy ds &= \int_{\Pi_T} \Psi_{\varepsilon,j}(u) \varphi_s + \Xi_{\varepsilon,j}(u) \varphi_y + \Phi_{\varepsilon,j}(u) \varphi_{yy} \, dy ds \\ &=: T_1^{\varepsilon} + T_2^{\varepsilon} + T_3^{\varepsilon}. \end{aligned}$$

Let us consider the three terms separately.

By Lemma 4.9,

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} \Psi_{\varepsilon,j}(u) &= \lim_{\varepsilon \downarrow 0} \int_{u_j}^u \Psi'_{\varepsilon,j}(z) \, dz = \int_{u_j}^u \lim_{\varepsilon \downarrow 0} \Psi'_{\varepsilon,j}(z) \, dz \\ &= \int_{u_j}^u \text{sign}(A(z) - A(u_j)) - \text{sg}_{j-1}(A(z)) \, dz \\ &\quad + \int_{u_j}^u \text{sign}(A(z) - A(u_j)) - \text{sg}_j(A(z)) \, dz, \end{aligned}$$

where $\text{sg}_j(\sigma) := \text{sg}_{(A(u_j), A(u_{j+1}))}(\sigma)$. Again by Lemma 4.9, the mapping

$$z \mapsto \text{sign}(A(z) - A(u_j)) - \text{sg}_{j-1}(A(z))$$

has support in $\text{int}(u_j, u_{j-1})$. Similar considerations apply to the second term. Hence,

$$\begin{aligned} \left| \lim_{\varepsilon \downarrow 0} \Psi_{\varepsilon,j}(u) \right| &\leq \left| \int_{u_j}^{u_{j-1}} \text{sign}(A(z) - A(u_j)) - \text{sg}_{j-1}(A(z)) \, dz \right| \\ &\quad + \left| \int_{u_j}^{u_{j+1}} \text{sign}(A(z) - A(u_j)) - \text{sg}_j(A(z)) \, dz \right| \\ &\leq 2|u_j - u_{j-1}| + 2|u_{j+1} - u_j|. \end{aligned}$$

By the same type of reasoning we obtain the bound

$$\begin{aligned} \left| \lim_{\varepsilon \downarrow 0} \Xi_j^{\varepsilon}(u) \right| &\leq \left| \int_{u_j}^{u_{j-1}} [\text{sign}(A(z) - A(u_j)) - \text{sg}_{j-1}(A(z))] f'(z) \, dz \right| \\ &\quad + \left| \int_{u_j}^{u_{j+1}} [\text{sign}(A(z) - A(u_j)) - \text{sg}_j(A(z))] f'(z) \, dz \right| \\ &\leq 2 \|f'\|_{L^{\infty}} (|u_j - u_{j-1}| + |u_{j+1} - u_j|). \end{aligned}$$

Concerning Φ_j^ε we use substitution and the explicit expression given in Lemma 4.9. This leads to

$$\begin{aligned} \left| \lim_{\varepsilon \downarrow 0} \Phi_j^\varepsilon(u) \right| &= \left| \int_{u_j}^u (-\operatorname{sg}_{j-1}(A(z)) + 2 \operatorname{sign}(A(z) - A(u_j)) - \operatorname{sg}_j(A(z))) A'(z) dz \right| \\ &= \left| \int_{A(u_j)}^{A(u)} -\operatorname{sg}_{j-1}(\sigma) + 2 \operatorname{sign}(\sigma - A(u_j)) - \operatorname{sg}_j(\sigma) d\sigma \right| \\ &\leq \left| \int_{A(u_j)}^{A(u_{j-1})} \operatorname{sign}(\sigma - A(u_j)) - \operatorname{sg}_{j-1}(\sigma) d\sigma \right| \\ &\quad + \left| \int_{A(u_j)}^{A(u_{j+1})} \operatorname{sign}(\sigma - A(u_j)) - \operatorname{sg}_j(\sigma) d\sigma \right| \\ &\leq |A(u_j) - A(u_{j-1})| + |A(u_{j+1}) - A(u_j)|. \end{aligned}$$

Let us return to equation (4.8). By the dominated convergence theorem and the above computations we have

$$\begin{aligned} \left| \lim_{\varepsilon \downarrow 0} T_1^\varepsilon \right| &\leq \left\| \lim_{\varepsilon \downarrow 0} \Psi_j^\varepsilon \right\|_{L^\infty} \int_{\Pi_T} |\varphi_s| dy ds \leq C(|D_- u_j| + |D_+ u_j|), \\ \left| \lim_{\varepsilon \downarrow 0} T_2^\varepsilon \right| &\leq \left\| \lim_{\varepsilon \downarrow 0} \Xi_j^\varepsilon \right\|_{L^\infty} \int_{\Pi_T} |\varphi_y| dy ds \leq C \frac{\Delta x}{r} (|D_- u_j| + |D_+ u_j|), \\ \left| \lim_{\varepsilon \downarrow 0} T_3^\varepsilon \right| &\leq \left\| \lim_{\varepsilon \downarrow 0} \Phi_j^\varepsilon \right\|_{L^\infty} \int_{\Pi_T} |\varphi_{yy}| dy ds \leq C \frac{\Delta x}{r^2} (|D_- A(u_j)| + |D_+ A(u_j)|). \end{aligned}$$

Hence,

$$\begin{aligned} &\int_{\Pi_T} \sum_j \liminf_{\varepsilon \downarrow 0} \left(\int_{\Pi_T} \mathcal{R}_j^\varepsilon(t, y, s) \varphi(x, t, y, s) dy ds \right) \mathbb{1}_{\{I_j\}}(x) dx dt \\ &\geq -C \left(\frac{\Delta x}{r_0} + \frac{\Delta x}{r} \right) \int_{\Pi_T} |D_- u_{\Delta x}| + |D_+ u_{\Delta x}| dx dt \\ &\quad - C \frac{\Delta x}{r^2} \int_{\Pi_T} |D_- A(u_{\Delta x})| + |D_+ A(u_{\Delta x})| dx dt. \end{aligned}$$

The desired estimate now follows from the uniform bounds in Lemma 3.1. $\square$

4.2. Proof of Theorem 4.1. Let us now combine the previous results to conclude the proof of Theorem 4.1. We begin by stating a rather standard lemma.

Lemma 4.12. *Set*

$$\kappa(t) := \int_{\mathbb{R}} \int_{\Pi_T} |u_{\Delta x}(x, t) - u(y, s)| \omega_r(x - y) \rho_{r_0}(t - s) dy ds dx.$$

Let $t \geq r_0$, and denote by L_c the Lipschitz constant of $t \mapsto \|u(\cdot, t)\|_{L^1(\mathbb{R})}$. Then

$$\left| \kappa(t) - \|u_{\Delta x}(\cdot, t) - u(\cdot, t)\|_{L^1(\mathbb{R})} \right| \leq |u(\cdot, t)|_{BV(\mathbb{R})} r + L_c r_0.$$

Proof. By the reverse triangle inequality,

$$\begin{aligned} & \left| \kappa(t) - \|u_{\Delta x}(\cdot, t) - u(\cdot, t)\|_{L^1(\mathbb{R})} \right| \\ & \leq \int_{\mathbb{R}} \int_{\Pi_T} |u(y, s) - u(x, t)| \omega_r(x - y) \rho_{r_0}(t - s) dy ds dx \\ & \leq \int_0^T \left(\int_{\mathbb{R}} |u(y, s) - u(t, y)| dy \right) \rho_{r_0}(t - s) ds \\ & \quad + \int_{\mathbb{R}} \int_{\mathbb{R}} |u(t, y) - u(x, t)| \omega_r(x - y) dy dx \\ & \leq L_c r_0 + |u(\cdot, t)|_{BV(\mathbb{R})} r. \end{aligned} \quad \square$$

Proof of Theorem 4.1. Our starting point is Lemma 4.6. Let $A(\sigma) = \hat{A}(\sigma) + \eta\sigma$, where $\hat{A}$ is the original degenerate diffusion function. Let

$$\begin{aligned} \Xi &= \iint_{\Pi_T^2} \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)) (D_+ \varphi + \varphi_y) dX \\ & \quad + \iint_{\Pi_T^2} \left(\int_{u_{\Delta x}}^{S_{\Delta x} u_{\Delta x}} \text{sign}(z - u) F'_2(z) dz \right) D_+ \varphi dX \\ & \quad + \iint_{\Pi_T^2} |A(u_{\Delta x}) - A(u)| (D_- D_+ \varphi + (D_+ + D_-) \varphi_y + \varphi_{yy}) dX. \end{aligned}$$

By Estimate 4.1, Estimate 4.2, and Estimate 4.3, it follows that

$$(4.9) \quad |\Xi| \leq C \frac{\Delta x}{r} \left(1 + \frac{\Delta x}{r^2} \right) \left(1 + \frac{\Delta x}{r} \right) =: E_1.$$

Furthermore, by Lemma 4.7, Lemma 4.10, Estimate 4.4, and Estimate 4.5, it follows that

$$(4.10) \quad \liminf_{\varepsilon \downarrow 0} \iint_{\Pi_T^2} E_{\Delta x}^\varepsilon \varphi dX \geq -C(1 + r + \Delta x) \frac{\Delta x}{r^2} \left(1 + \frac{\Delta x}{r} \right)^3 - C \frac{\Delta x}{r_0} =: -E_2.$$

Applying the estimates (4.9) and (4.10), the inequality (4.2) becomes

$$\begin{aligned} & \iint_{\Pi_T^2} |u_{\Delta x} - u| \rho_\alpha(t - \tau) \omega_r(x - y) \rho_{r_0}(t - s) dX \\ & \leq \iint_{\Pi_T^2} |u_{\Delta x} - u| \rho_\alpha(t - \nu) \omega_r(x - y) \rho_{r_0}(t - s) dX + E_1 + E_2. \end{aligned}$$

Note that both E_1 and E_2 are independent of α . Thus, we can send α to zero, arriving at

$$\kappa(\tau) \leq \kappa(\nu) + E_1 + E_2,$$

where κ is defined as in Lemma 4.12.

By Lemma 4.12 it follows that

$$\begin{aligned} & \|u_{\Delta x}(\cdot, \tau) - u(\cdot, \tau)\|_{L^1(\mathbb{R})} \\ & \leq \|u_{\Delta x}(\cdot, \nu) - u(\cdot, \nu)\|_{L^1(\mathbb{R})} + 2 \left(L_c r_0 + |u^0|_{BV(\mathbb{R})} r \right) + E_1 + E_2. \end{aligned}$$

Recall that we had to pick $\nu > r_0$. Denote by L_d the L^1 Lipschitz constant of $t \mapsto u_{\Delta x}(\cdot, t)$. By the triangle inequality

$$\begin{aligned} & \|u_{\Delta x}(\cdot, \nu) - u(\cdot, \nu)\|_{L^1(\mathbb{R})} \\ & \leq \|u_{\Delta x}(\cdot, \nu) - u_{\Delta x}^0\|_{L^1(\mathbb{R})} + \|u_{\Delta x}^0 - u^0\|_{L^1(\mathbb{R})} + \|u^0 - u(\cdot, \nu)\|_{L^1(\mathbb{R})} \\ & \leq L_d \nu + \|u_{\Delta x}^0 - u^0\|_{L^1(\mathbb{R})} + L_c \nu. \end{aligned}$$

This means that

$$\begin{aligned} \|u_{\Delta x}(\cdot, \tau) - u(\cdot, \tau)\|_{L^1(\mathbb{R})} & \leq \|u_{\Delta x}^0 - u^0\|_{L^1(\mathbb{R})} + (L_c + L_d) \nu \\ & \quad + 2 \left(L_c r_0 + |u^0|_{BV(\mathbb{R})} r \right) + E_1 + E_2. \end{aligned}$$

Choose $r^3 = r_0^2 = \Delta x$ and $\nu = 2r_0$. Then there exists a constant C such that

$$\|u_{\Delta x}(\cdot, \tau) - u(\cdot, \tau)\|_{L^1(\mathbb{R})} \leq \|u_{\Delta x}^0 - u^0\| + C \Delta x^{\frac{1}{3}}.$$

Now recall that $A(\sigma) = \hat{A}(\sigma) + \eta\sigma$ and so we need to send η to zero to finish the proof. If u_η is the classical solution of the regularized equation and u is the entropy solution of the nonregularized equation, then it is well known that $u_\eta(\cdot, t) \rightarrow u(\cdot, t)$ in $L^1(\mathbb{R})$ as $\eta \rightarrow 0$ (see Section 2). Concerning the scheme, one may prove continuous dependence in ℓ^1 on η using Gronwall's inequality. Hence, we can also send η to zero in the scheme. This finishes the proof of Theorem 4.1. $\square$

5. IMPLICIT DIFFERENCE SCHEMES

In this section we show that the arguments presented in the previous sections carry through for implicit schemes. Fix a time step $\Delta t > 0$. We consider implicit difference schemes of the form

$$(5.1) \quad D_-^t u_j^n + D_- F(u_j^n, u_{j+1}^n) = D_- D_+ A(u_j^n) \quad n \geq 1, j \in \mathbb{Z},$$

where

$$D_-^t u_j^n = \frac{u_j^n - u_j^{n-1}}{\Delta t}.$$

Let $t_n = n\Delta t$ and $x_j = j\Delta x$. We define the grid cells

$$I_j^n = [x_{j-1/2}, x_{j+1/2}) \times (t_{n-1}, t_n] \quad \text{for } n \geq 0 \text{ and } j \in \mathbb{Z}.$$

The piecewise constant approximation is defined for all $(x, t) \in \mathbb{R} \times (-\Delta t, T]$ by

$$(5.2) \quad u_\Delta(x, t) = u_j^n \quad \text{for } (x, t) \in I_j^n.$$

The domain is chosen so that $D_-^t u_\Delta$ is defined for all $(x, t) \in \mathbb{R} \times (0, T)$. For the existence of a unique solution u_j^n to the nonlinear equation (5.1) and the convergence of u_Δ to an entropy solution; see [11].

We now state the main theorem.

Theorem 5.1. *Let u be the entropy solution to (1.1), and let u_Δ be defined via u_j^n by (5.2), where u_j^n solves (5.1). If u^0 satisfies the same assumptions as in Theorem 4.1, then for all sufficiently small Δx and Δt , and for all $n \in \mathbb{N}$ such that $t_n \in [0, T]$,*

$$\|u_\Delta(\cdot, t_n) - u(\cdot, t_n)\|_{L^1(\mathbb{R})} \leq \|u_\Delta^0 - u^0\|_{L^1(\mathbb{R})} + C \left(\Delta x^{1/3} + \Delta t^{1/2} \right),$$

where the constant C_T depends on u_0, A, f, T , but not on $\Delta x, \Delta t$.

To prove this theorem we will follow step-by-step the proof of Theorem 4.1 and present the details whenever there is a significant difference between the two cases.

Thanks to [11, Lemma 2.4], we have the following L^1 Lipschitz continuity result:

Lemma 5.1. *Let m and n be two nonnegative integers. Then*

$$\|u_{\Delta}(\cdot, t_n) - u_{\Delta}(\cdot, t_m)\|_{L^1(\mathbb{R})} \leq L_d |t_n - t_m|,$$

where $L_d = |F(u_j^0, u_{j+1}^0) - D_+ A(u_j^0)|_{BV}$.

Next, let us prove an implicit version of Lemma 4.3.

Lemma 5.2. *Let u_j^n be the solution to (5.1). Then for all $c \in \mathbb{R}$,*

$$\begin{aligned} & D_-^t \psi_{\varepsilon}(u_j^n, c) + D_- Q^c(u_j^n, u_{j+1}^n) - D_- D_+ |A(u_j^n) - A(c)|_{\varepsilon} \\ & \leq -\frac{1}{(\Delta x)^2} \int_{u_{j+1}^n}^{u_j^n} \psi_{\varepsilon}''(z, c) (A(z) - A(u_{j+1}^n)) dz \\ & \quad - \frac{1}{(\Delta x)^2} \int_{u_{j-1}^n}^{u_j^n} \psi_{\varepsilon}''(z, c) (A(z) - A(u_{j-1}^n)) dz, \end{aligned}$$

where $Q^c(u, v)$ is defined in Lemma 4.3.

Proof. From (5.1) it follows that

$$\psi_{\varepsilon}'(u_j^n, c) D_-^t u_j^n + \psi_{\varepsilon}'(u_j^n, c) D_- F(u_j^n, u_{j+1}^n) = \psi_{\varepsilon}'(u_j^n, c) D_- D_+ A(u_j^n).$$

Apply Lemma 4.2 with $g(\sigma) = \sigma$, $a = u_j^n$, and $b = u_j^{n-1}$ to obtain

$$\begin{aligned} \psi_{\varepsilon}'(u_j^n, c) D_-^t u_j^n &= D_-^t \psi_{\varepsilon}(u_j^n, c) - \frac{1}{\Delta t} \int_{u_j^n}^{u_j^{n-1}} \psi_{\varepsilon}''(z, c) (z - u_j^{n-1}) dz \\ &\geq D_-^t \psi_{\varepsilon}(u_j^n, c). \end{aligned}$$

The remaining part of the proof follows exactly as in the proof of Lemma 4.3. $\square$

Let us define the time shift operator

$$S_{\Delta t}^t \sigma(t) = \sigma(t + \Delta t),$$

for any function $\sigma = \sigma(t)$.

Lemma 5.3. *Suppose $A' > 0$. Let $u_{\Delta} = u_{\Delta}(x, t)$ be defined by (5.2), and let $u = u(y, s)$ be the classical solution of (1.1). Let $\psi(t) := \mathbb{1}_{[\nu, \tau)}(t)$ and define*

$$\varphi(x, t, y, s) = \psi(t) \omega_r(x - y) \rho_{r_0}(t - s),$$

where $\omega_r, \rho_{r_0}, \nu, \tau$ are chosen as in Lemma 4.6. Then

$$\begin{aligned}
& \iint_{\Pi_T^2} |u_\Delta - u| \delta_{\Delta t}^-(t - \nu) \omega_r \rho_{r_0} dX \\
& + \iint_{\Pi_T^2} |u_\Delta - u| S_{\Delta t}^t \psi \omega_r (D_+^t \rho_{r_0} - \partial_t \rho_{r_0}) dX \\
& + \Delta t \iint_{\Pi_T^2} |u_\Delta - u| D_+^t \psi \omega_r \partial_s \rho_{r_0} dX \\
& + \iint_{\Pi_T^2} \text{sign}(u_\Delta - u) (f(u_\Delta) - f(u)) (D_+ \varphi + \varphi_y) dX \\
& + \iint_{\Pi_T^2} \left(\int_{u_\Delta x}^{S_{\Delta x} u_\Delta} \text{sign}(z - u) F_2'(z) dz \right) D_+ \varphi dX \\
& + \iint_{\Pi_T^2} |A(u_\Delta) - A(u)| (D_- D_+ \varphi + (D_+ + D_-) \varphi_y + \varphi_{yy}) dX \\
& \geq \iint_{\Pi_T^2} |u_\Delta - u| \delta_{\Delta t}^-(t - \tau) \omega_r \rho_{r_0} dX + \liminf_{\varepsilon \downarrow 0} \iint_{\Pi_T^2} E_\Delta^\varepsilon \varphi dX,
\end{aligned}$$

where

$$\delta_{\Delta t}^-(t) = \frac{1}{\Delta t} \mathbb{1}_{\{[-\Delta t, 0)\}}(t),$$

and $E_\Delta^\varepsilon(x, t, y, s) = E^\varepsilon[u](u_{j-1}^n, u_j^n, u_{j+1}^n)(y, s)$ for $(x, t) \in I_j^n$.

Proof. As in Lemma 4.5, we obtain by Lemma 5.2 the inequality

$$\begin{aligned}
& D_-^t \psi_\varepsilon(u_j^n, u) + \partial_s \psi_\varepsilon(u, u_j^n) + \partial_y q_\varepsilon(u, u_j^n) + D_- Q^u(u_j^n, u_{j+1}^n) \\
& - (\partial_y^2 + \partial_y (D_- + D_+) + D_- D_+) |A(u_j^n) - A(u)|_\varepsilon \leq -E_{j,n}^\varepsilon,
\end{aligned}$$

where $E_{j,n}^\varepsilon := E^\varepsilon[u](u_{j-1}^n, u_j^n, u_{j+1}^n)$ is defined in Lemma 4.5. Let us multiply by φ and integrate over Π_T^2 . Integration by parts for difference quotients and ordinary integration by parts gives

$$\begin{aligned}
& \iint_{\Pi_T^2} \psi_\varepsilon(u_\Delta, u) D_+^t \varphi + \psi_\varepsilon(u, u_\Delta) \varphi_s dX \\
& + \iint_{\Pi_T^2} q_\varepsilon(u, u_\Delta) \varphi_y + Q^u(u_\Delta, S_{\Delta x} u_\Delta) D_+ \varphi dX \\
& + \iint_{\Pi_T^2} |A(u_\Delta) - A(u)|_\varepsilon (\varphi_{yy} + (D_- + D_+) \varphi_y + D_- D_+ \varphi) dX \\
& \geq \iint_{\Pi_T^2} E_\Delta^\varepsilon \varphi dX.
\end{aligned}$$

Consider the first term on the left. Let ε tend to zero as in the proof of Lemma 4.6. Using the Leibniz rule for difference quotients and adding and subtracting we obtain

$$D_+^t \varphi = S_{\Delta t}^t \psi \omega_r D_+^t \rho_{r_0} + D_+^t \psi \omega_r \rho_{r_0}.$$

Furthermore,

$$\varphi_s = -S_{\Delta t}^t \psi \omega_r \partial_t \rho_{r_0} + \Delta t D_+^t \psi \omega_r \partial_s \rho_{r_0}.$$

Hence,

$$\begin{aligned} & \iint_{\Pi_T^2} |u_\Delta - u| (D_+^t \varphi + \varphi_s) dX \\ &= \iint_{\Pi_T^2} |u_\Delta - u| S_{\Delta t}^t \psi \omega_r (D_+^t \rho_{r_0} - \partial_t \rho_{r_0}) dX \\ & \quad + \Delta t \iint_{\Pi_T^2} |u_\Delta - u| D_+^t \psi \omega_r \partial_s \rho_{r_0} dX + \iint_{\Pi_T^2} |u_\Delta - u| D_+^t \psi \omega_r \rho_{r_0} dX. \end{aligned}$$

Finally, we use that

$$(5.3) \quad D_+^t \psi = \delta_{\Delta t}^-(t - \nu) - \delta_{\Delta t}^-(t - \tau).$$

The lemma now follows, as in the proof of Lemma 4.6, by letting ε tend to zero. $\square$

Comparing the terms in Lemma 4.6 with the terms in Lemma 5.3 we recognize all but two terms.

Estimate 5.1.

$$\left| \iint_{\Pi_T^2} |u - u_\Delta| S_{\Delta t}^t \psi \omega_r (D_+^t \rho_{r_0} - \partial_t \rho_{r_0}) dX \right| \leq C \frac{\Delta t}{r_0} \left(1 + \frac{\Delta t}{r_0} \right).$$

Proof. To show this we use a Taylor expansion:

$$\begin{aligned} & \rho_{r_0}(t + \Delta t - s) - \rho_{r_0}(t - s) \\ &= \int_t^{t+\Delta t} \frac{\partial}{\partial z} \rho_{r_0}(z - s) dz \\ &= \frac{\partial}{\partial z} \Big|_{z=t} \rho_{r_0}(z - s) \Delta t - \int_t^{t+\Delta t} \frac{\partial^2}{\partial z^2} \rho_{r_0}(z - s) (z - (t + \Delta t)) dz. \end{aligned}$$

It follows that

$$(5.4) \quad D_+^t \rho_{r_0} - \partial_t \rho_{r_0} = -\frac{1}{\Delta t} \int_t^{t+\Delta t} \frac{\partial^2}{\partial z^2} \rho_{r_0}(z - s) (z - (t + \Delta t)) dz.$$

Integration by parts yields

$$\begin{aligned} & -\frac{1}{\Delta t} \int_t^{t+\Delta t} \int_0^T |u - u_\Delta|_\varepsilon \frac{\partial^2}{\partial z^2} \rho_{r_0}(z - s) (z - (t + \Delta t)) ds dz \\ &= \frac{1}{\Delta t} \int_t^{t+\Delta t} \int_0^T |u - u_\Delta|_\varepsilon \frac{\partial}{\partial s} \frac{\partial}{\partial z} \rho_{r_0}(z - s) (z - (t + \Delta t)) ds dz \\ &= -\frac{1}{\Delta t} \int_t^{t+\Delta t} \int_0^T \text{sign}_\varepsilon(u - u_\Delta) u_s \frac{\partial}{\partial z} \rho_{r_0}(z - s) (z - (t + \Delta t)) ds dz. \end{aligned}$$

Since

$$\frac{\partial}{\partial z} \rho_{r_0}(z - s) = \frac{1}{r_0} \frac{\partial}{\partial z} \rho \left(\frac{z - s}{r_0} \right) = \frac{1}{r_0^2} \rho' \left(\frac{z - s}{r_0} \right)$$

and ρ_{r_0} has support in $[-r_0, r_0]$, it follows that

$$\begin{aligned} & \frac{1}{\Delta t} \left| \int_t^{t+\Delta t} \int_0^T |u(y, s) - u_\Delta(x, t)|_\varepsilon \frac{\partial^2}{\partial z^2} \rho_{r_0}(z - s)(z - (t + \Delta t)) ds dz \right| \\ & \leq \frac{C}{r_0^2 \Delta t} \int_0^T |u_s(y, s)| \int_t^{t+\Delta t} \mathbb{1}_{\{|z-s| \leq r_0\}} |z - (t + \Delta t)| dz ds \\ & \leq C \frac{\Delta t}{r_0^2} \int_0^T |u_s| \mathbb{1}_{\{|t-s| \leq r_0 + \Delta t\}} ds. \end{aligned}$$

Multiply the above inequality by $\psi(t)\omega_r(x - y)$ and integrate in x, y, t . From the resulting inequality and (5.4), we arrive at the estimate:

$$\begin{aligned} & \left| \iint_{\Pi_T^2} |u - u_\Delta|_\varepsilon S_{\Delta t}^t \psi \omega_r (D_+^t \rho_{r_0} - \partial_t \rho_{r_0}) dX \right| \\ & \leq C \frac{\Delta t}{r_0^2} \iint_{\Pi_T^2} |u_s| S_{\Delta t}^t \psi \omega_r \mathbb{1}_{\{|t-s| \leq r_0 + \Delta t\}} dX \leq C \frac{\Delta t}{r_0} \left(1 + \frac{\Delta t}{r_0} \right) \|u_s\|_{L^1(\Pi_T)}. \end{aligned}$$

Since $\|u_s(\cdot, s)\|_{L^1(\mathbb{R})}$ is uniformly bounded on $[0, T]$, the estimate follows from the dominated convergence theorem. $\square$

Estimate 5.2.

$$\left| \Delta t \iint_{\Pi_T^2} |u - u_\Delta| D_+^t \psi \omega_r \partial_s \rho_{r_0} dX \right| \leq C \frac{\Delta t}{r_0}.$$

Proof. Integration by parts yields

$$\left| \iint_{\Pi_T^2} |u - u_\Delta| D_+^t \psi \omega_r \partial_s \rho_{r_0} dX \right| \leq \iint_{\Pi_T^2} |u_s| |D_+^t \psi| \omega_r \rho_{r_0} dX.$$

Because of (5.3) and since

$$\|\rho_{r_0}\|_{L^\infty} \leq \frac{\|\rho\|_{L^\infty}}{r_0},$$

it follows that

$$\iint_{\Pi_T^2} |u_s| |D_+^t \psi| \omega_r \rho_{r_0} dX \leq 2 \frac{\|\rho\|_{L^\infty}}{r_0} \|u_s\|_{L^1(\Pi_T)}. \quad \square$$

Proof of Theorem 5.1. We start out from Lemma 5.3 with $A(\sigma) = \hat{A}(\sigma) + \eta\sigma$, where $\hat{A}$ is the original degenerate diffusion function. By Estimate 5.1 and Estimate 5.2,

$$\begin{aligned} (5.5) \quad & \iint_{\Pi_T^2} |u - u_\Delta| S_{\Delta t}^t \psi \omega_r (D_+^t \rho_{r_0} - \partial_t \rho_{r_0}) dX \\ & + \Delta t \iint_{\Pi_T^2} |u - u_\Delta| D_+^t \psi \omega_r \partial_s \rho_{r_0} dX \leq C \frac{\Delta t}{r_0} \left(1 + \frac{\Delta t}{r_0} \right) =: E_3. \end{aligned}$$

Since all the estimates from Section 4.1 apply, we obtain

$$\begin{aligned} & \iint_{\Pi_T^2} |u_\Delta - u| \delta^{\Delta t}(t - \tau) \omega_r(x - y) \rho_{r_0}(t - s) dX \\ & \leq \iint_{\Pi_T^2} |u_{\Delta x} - u| \delta^{\Delta t}(t - \nu) \omega_r(x - y) \rho_{r_0}(t - s) dX + E_1 + E_2 + E_3, \end{aligned}$$

where E_1 and E_2 are defined respectively in (4.9) and (4.10).

Let us make the simplifying assumption that $\nu = t_m$ and $\tau = t_n$ for some $m, n \in \mathbb{N}$. Then the above inequality is rewritten as

$$\kappa(t_n) \leq \kappa(t_m) + E_1 + E_2 + E_3,$$

where

$$\kappa(t) = \int_{\mathbb{R}} \int_{\Pi_T} |u_{\Delta}(x, t) - u(y, s)| \omega_r(x - y) \rho_{r_0}(t - s) dy ds dx.$$

Applying Lemmas 4.12 and 5.1, and following the reasoning given in the semi-discrete case, we arrive at

$$\begin{aligned} & \|u_{\Delta}(\cdot, t_n) - u(\cdot, t_n)\|_{L^1(\mathbb{R})} \\ & \leq \|u_{\Delta}^0 - u^0\|_{L^1(\mathbb{R})} \\ & \quad + (L_c + L_d) t_m + 2 \left(L_c r_0 + |u^0|_{BV(\mathbb{R})} r \right) \\ & \quad + C(1 + r + \Delta x)^2 \left(1 + \frac{\Delta x}{r} \right)^3 \frac{\Delta x}{r^2} + C \frac{\Delta x}{r_0} + C \frac{\Delta t}{r_0} \left(1 + \frac{\Delta t}{r_0} \right) \\ & \leq \|u_{\Delta}^0 - u^0\|_{L^1(\mathbb{R})} + C \left(\frac{\Delta x}{r^2} + \frac{\Delta x + \Delta t}{r_0} + r + r_0 \right), \end{aligned}$$

where L_d is the constant in Lemma 5.1 and L_c is the constant from Lemma 4.12. Minimizing over r and r_0 , it is straightforward to see that for sufficiently small Δt , the minimum of the last term is dominated by

$$C \left(\Delta x^{1/3} + \Delta t^{1/2} \right).$$

This proves the theorem. $\square$

6. EXPLICIT DIFFERENCE SCHEMES

In this section we use the techniques developed in the previous section to provide a similar result concerning the explicit scheme. Fix a time step $\Delta t > 0$. We consider explicit schemes of the form

$$(6.1) \quad D_+^t u_j^n + D_- F(u_j^n, u_{j+1}^n) = D_- D_+ A(u_j^n) \quad n \geq 1, j \in \mathbb{Z},$$

where

$$D_+^t u_j^n = \frac{u_j^{n+1} - u_j^n}{\Delta t}.$$

The relevant a priori estimates and convergence to an entropy solution is proved in [12] under the hypothesis

$$(6.2) \quad 1 - \frac{\Delta t}{\Delta x} (F_1'(z) - F_2'(z)) - 2 \frac{\Delta t}{\Delta x^2} A'(w) \geq 0, \quad \forall (z, w) \in \mathbb{R}^2.$$

Let $t_n = n\Delta t$ and $x_j = j\Delta x$. We define the grid cells

$$I_j^n = [x_{j-1/2}, x_{j+1/2}) \times [t_n, t_{n+1}), \quad \text{for } n \geq 0 \text{ and } j \in \mathbb{Z}.$$

The piecewise constant approximation is defined for all $(x, t) \in \Pi_T$ by

$$(6.3) \quad u_{\Delta}(x, t) = u_j^n \text{ for } (x, t) \in I_j^n.$$

Theorem 6.1. *Let u be the entropy solution to (1.1), and let u_Δ be defined by (6.3) via u_j^n , where u_j^n solves (6.1). Suppose Δt and Δx are chosen such that (6.2) and the strengthened condition $\Delta t \leq C\Delta x^{8/3}$ hold. If u^0 satisfies the same assumptions as in Theorem 4.1, then for all sufficiently small Δx , and for all $n \in \mathbb{N}$ such that $t_n \in [0, T]$,*

$$\|u_\Delta(\cdot, t_n) - u(\cdot, t_n)\|_{L^1(\mathbb{R})} \leq \|u_\Delta^0 - u^0\|_{L^1(\mathbb{R})} + C\Delta x^{1/3},$$

where the constant C_T depends on A, f, u^0, T , but not on Δx .

We begin by proving the following lemma.

Lemma 6.1. *Let $\{u_j^n\}$ be the solution to (6.1). Suppose that*

$$(6.4) \quad 1 - \frac{\Delta t}{\Delta x} (F'_1(z) - F'_2(z)) \geq 0,$$

for all $z \in \mathbb{R}$. Then for all $j \in \mathbb{Z}$ and $n \in \mathbb{N}$,

$$\begin{aligned} & D_+^t \psi_\varepsilon(u_j^n, c) + D_- Q^c(u_j^n, u_{j+1}^n) - D_- D_+ |A(u_j^n) - A(c)|_\varepsilon \\ & \leq -\frac{1}{(\Delta x)^2} \int_{u_{j-1}^n}^{u_j^n} \psi_\varepsilon''(z, c) (A(z) - A(u_{j-1}^n)) dz \\ & \quad - \frac{1}{(\Delta x)^2} \int_{u_{j+1}^n}^{u_j^n} \psi_\varepsilon''(z, c) (A(z) - A(u_{j+1}^n)) dz \\ & \quad + \left(\int_{u_j^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c) dz \right) D_- D_+ A(u_j^n), \end{aligned}$$

where $Q^c(u, v)$ is defined by

$$Q^c(u, v) = \int_c^u \psi_\varepsilon'(z, c) F'_1(z) dz + \int_c^v \psi_\varepsilon'(z, c) F'_2(z) dz, \quad u, v \in \mathbb{R}.$$

Proof. We divide the proof into two steps.

Claim 1. Let $\{u_j^n\}$ be a solution to (6.1). Then

$$\begin{aligned} & D_+^t \psi_\varepsilon(u_j^n, c) + D_- Q^c(u_j^n, u_{j+1}^n) - D_- D_+ |A(u_j^n) - A(c)|_\varepsilon \\ & = -E^c(u_j^n, u_j^{n+1}, u_{j+1}^n, u_{j-1}^n), \end{aligned}$$

where

$$\begin{aligned} & E^c(u_j^n, u_j^{n+1}, u_{j+1}^n, u_{j-1}^n) \\ & = \frac{1}{\Delta t} \int_{u_j^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c) (z - u_j^{n+1}) dz \\ & \quad + \frac{1}{\Delta x} \int_{u_{j-1}^n}^{u_j^n} \psi_\varepsilon''(z, c) \left[(F_1(z) - F_1(u_{j-1}^n)) + \frac{1}{\Delta x} (A(z) - A(u_{j-1}^n)) \right] dz \\ & \quad + \frac{1}{\Delta x} \int_{u_j^n}^{u_{j+1}^n} \psi_\varepsilon''(z, c) \left[(F_2(z) - F_2(u_{j+1}^n)) - \frac{1}{\Delta x} (A(z) - A(u_{j+1}^n)) \right] dz. \end{aligned}$$

Proof of Claim 1. By definition (6.1) of $\{u_j^n\}$ it follows that

$$\psi_\varepsilon'(u_j^n, c) [D_+^t u_j^n + D_- F(u_j^n, u_{j+1}^n) - D_- D_+ A(u_j^n)] = 0.$$

Let $g(z) = z$ in Lemma 4.2. It follows that

$$\psi'_\varepsilon(u_j^n, c) D_+^t u_j^n = D_+^t \psi_\varepsilon(u_j^n, c) + \frac{1}{\Delta t} \int_{u_j^n}^{u_j^{n+1}} \psi''_\varepsilon(z, c) (z - u_j^{n+1}) dz.$$

The remaining terms can be treated as in Lemma 4.3. $\square$

Claim 2. Suppose (6.4) holds. Then

$$\begin{aligned} (6.5) \quad & \frac{1}{\Delta t} \int_{u_j^n}^{u_j^{n+1}} \psi''_\varepsilon(z, c) (z - u_j^{n+1}) dz \\ & + \frac{1}{\Delta x} \int_{u_{j-1}^n}^{u_j^n} \psi''_\varepsilon(z, c) (F_1(z) - F_1(u_{j-1}^n)) dz \\ & + \frac{1}{\Delta x} \int_{u_j^n}^{u_{j+1}^n} \psi''_\varepsilon(z, c) (F_2(z) - F_2(u_{j+1}^n)) dz \\ & \geq - \int_{u_j^n}^{u_j^{n+1}} \psi''_\varepsilon(z, c) D_- D_+ A(u_j^n) dz. \end{aligned}$$

Proof of Claim 2. Consider the first term on the left-hand side of (6.5). By definition, $u_j^{n+1} = u_j^n - \Delta t D_- F(u_j^n, u_{j+1}^n) + \Delta t D_- D_+ A(u_j^n)$, and so

$$\begin{aligned} & \frac{1}{\Delta t} \int_{u_j^n}^{u_j^{n+1}} \psi''_\varepsilon(z, c) (z - u_j^{n+1}) dz \\ & = \frac{1}{\Delta t} \int_{u_j^n}^{u_j^{n+1}} \psi''_\varepsilon(z, c) (z - u_j^n) dz + \int_{u_j^n}^{u_j^{n+1}} \psi''_\varepsilon(z, c) D_- F(u_j^n, u_{j+1}^n) dz \\ & \quad - \int_{u_j^n}^{u_j^{n+1}} \psi''_\varepsilon(z, c) D_- D_+ A(u_j^n) dz =: T_1 + T_2 + T_3. \end{aligned}$$

Note that T_1 is positive.

Let us split T_2 according to

$$D_- F(u_j^n, u_{j+1}^n) = \frac{1}{\Delta x} (F_1(u_j^n) - F_1(u_{j-1}^n)) + \frac{1}{\Delta x} (F_2(u_{j+1}^n) - F_2(u_j^n)),$$

and thus

$$\begin{aligned} T_2 & = \frac{1}{\Delta x} \int_{u_j^n}^{u_j^{n+1}} \psi''_\varepsilon(z, c) (F_1(u_j^n) - F_1(u_{j-1}^n)) dz \\ & \quad + \frac{1}{\Delta x} \int_{u_j^n}^{u_j^{n+1}} \psi''_\varepsilon(z, c) (F_2(u_{j+1}^n) - F_2(u_j^n)) dz. \end{aligned}$$

Now, let us split the two other terms appearing in equation (6.5):

$$\begin{aligned} S_1 &:= \frac{1}{\Delta x} \int_{u_{j-1}^n}^{u_j^n} \psi_\varepsilon''(z, c)(F_1(z) - F_1(u_{j-1}^n)) dz \\ &= \frac{1}{\Delta x} \int_{u_{j-1}^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c)(F_1(z) - F_1(u_{j-1}^n)) dz \\ &\quad - \frac{1}{\Delta x} \int_{u_j^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c)(F_1(z) - F_1(u_{j-1}^n)) dz \end{aligned}$$

and

$$\begin{aligned} S_2 &:= \frac{1}{\Delta x} \int_{u_j^n}^{u_{j+1}^n} \psi_\varepsilon''(z, c)(F_2(z) - F_2(u_{j+1}^n)) dz \\ &= -\frac{1}{\Delta x} \int_{u_{j+1}^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c)(F_2(z) - F_2(u_{j+1}^n)) dz \\ &\quad + \frac{1}{\Delta x} \int_{u_j^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c)(F_2(z) - F_2(u_{j+1}^n)) dz. \end{aligned}$$

Combining the above expressions we obtain

$$\begin{aligned} T_2 + S_1 + S_2 &= -\frac{1}{\Delta x} \int_{u_j^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c)(F_1(z) - F_1(u_j^n)) dz \\ &\quad + \frac{1}{\Delta x} \int_{u_j^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c)(F_2(z) - F_2(u_j^n)) dz \\ &\quad + \frac{1}{\Delta x} \int_{u_{j-1}^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c)(F_1(z) - F_1(u_{j-1}^n)) dz \\ &\quad - \frac{1}{\Delta x} \int_{u_{j+1}^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c)(F_2(z) - F_2(u_{j+1}^n)) dz. \end{aligned}$$

The two last terms on the right-hand side are positive as F is monotone. Let

$$H(z) = z - \frac{\Delta t}{\Delta x} (F_1(z) - F_2(z)).$$

Then, by assumption (6.4),

$$T_1 + T_2 + S_1 + S_2 \geq \frac{1}{\Delta t} \int_{u_j^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c) [H(z) - H(u_j^n)] dz \geq 0.$$

Adding T_3 to both sides proves Claim 2. □

By Claim 2,

$$\begin{aligned} E^c(u_j^n, u_j^{n+1}, u_{j+1}^n, u_{j-1}^n) \\ \geq \frac{1}{\Delta x^2} \int_{u_{j-1}^n}^{u_j^n} \psi_\varepsilon''(z, c)(A(z) - A(u_{j-1}^n)) dz \\ + \frac{1}{\Delta x^2} \int_{u_{j+1}^n}^{u_j^n} \psi_\varepsilon''(z, c)(A(z) - A(u_{j+1}^n)) dz \\ - \int_{u_j^n}^{u_j^{n+1}} \psi_\varepsilon''(z, c) D_- D_+ A(u_j^n) dz. \end{aligned}$$

Combining this with Claim 1 proves the lemma. $\square$

Lemma 6.2. *Suppose $A' > 0$, and (6.4) applies. Let $u_\Delta = u_\Delta(x, t)$ be defined by (6.3), and let $u = u(y, s)$ be the classical solution of (1.1). Set $\psi(t) := \mathbb{1}_{\{[\nu, \tau)\}}(t)$ and define*

$$\varphi(x, t, y, s) = \psi(t) \omega_r(x - y) \rho_{r_0}(t - s),$$

where $\omega_r, \rho_{r_0}, \nu, \tau$ are chosen as in Lemma 4.6. Then

$$\begin{aligned} & \iint_{\Pi_T^2} |u_\Delta - u| \delta_{\Delta t}^+(t - \nu) \omega_r \rho_{r_0} dX \\ & + \iint_{\Pi_T^2} |u_\Delta - u| S_{-\Delta t}^t \psi \omega_r (D_-^t \rho_{r_0} - \partial_t \rho_{r_0}) dX \\ & + \Delta t \iint_{\Pi_T^2} |u_\Delta - u| D_-^t \psi \omega_r \partial_s \rho_{r_0} dX \\ & + \iint_{\Pi_T^2} \text{sign}(u_\Delta - u) (f(u_\Delta) - f(u)) (D_+ \varphi + \varphi_y) dX \\ & + \iint_{\Pi_T^2} \left(\int_{u_\Delta x}^{S_{\Delta x} u_\Delta} \text{sign}(z - u) F_2'(z) dz \right) D_+ \varphi dX \\ & + \iint_{\Pi_T^2} |A(u_\Delta) - A(u)| (D_- D_+ \varphi + (D_+ + D_-) \varphi_y + \varphi_{yy}) dX \\ & \geq \iint_{\Pi_T^2} |u_\Delta - u| \delta_{\Delta t}^+(t - \tau) \omega_r \rho_{r_0} dX \\ & + \liminf_{\varepsilon \downarrow 0} \iint_{\Pi_T^2} E_\Delta^\varepsilon \varphi dX \\ & - \Delta t \iint_{\Pi_T^2} D_+^t \text{sign}_\varepsilon(A(u_\Delta) - A(u)) D_- D_+ A(u_\Delta) \varphi dX, \end{aligned}$$

where

$$\delta_{\Delta t}^+(t) = \frac{1}{\Delta t} \mathbb{1}_{\{[0, \Delta t)\}}(t),$$

and $E_\Delta^\varepsilon(x, t, y, s) = E^\varepsilon[u](u_{j-1}^n, u_j^n, u_{j+1}^n)(y, s)$ for $(x, t) \in I_j^n$.

Proof. By Lemma 6.1, we obtain as in Lemma 4.5 the following inequality:

$$\begin{aligned} & D_+^t \psi_\varepsilon(u_j^n, u) + \partial_s \psi_\varepsilon(u, u_j^n) + \partial_y q_\varepsilon(u, u_j^n) + D_- Q^u(u_j^n, u_{j+1}^n) \\ & - (\partial_y^2 + \partial_y(D_- + D_+) + D_- D_+) |A(u_j^n) - A(u)|_\varepsilon \\ & \leq -E_{j,n}^\varepsilon + \left(\int_{u_j^n}^{u_j^{n+1}} \psi_\varepsilon''(z, u) dz \right) D_- D_+ A(u_j^n), \end{aligned}$$

where $E_{j,n}^\varepsilon := E^\varepsilon[u](u_{j-1}^n, u_j^n, u_{j+1}^n)$ is defined in Lemma 4.5. Note that

$$\int_{u_j^n}^{u_j^{n+1}} \psi_\varepsilon''(z, u) dz = \Delta t D_+^t \text{sign}_\varepsilon(A(u_j^n) - A(u)).$$

Integration by parts for difference quotients and ordinary integration by parts gives

$$\begin{aligned} & \iint_{\Pi_T^2} \psi_\varepsilon(u_\Delta, u) D_-^t \varphi + \psi_\varepsilon(u, u_\Delta) \varphi_s dX \\ & + \iint_{\Pi_T^2} q_\varepsilon(u, u_\Delta) \varphi_y + Q^u(u_\Delta, S_{\Delta x} u_\Delta) D_+ \varphi dX \\ & + \iint_{\Pi_T^2} |A(u_\Delta) - A(u)|_\varepsilon (\varphi_{yy} + (D_- + D_+) \varphi_y + D_- D_+ \varphi) dX \\ & \geq \iint_{\Pi_T^2} E_\Delta^\varepsilon \varphi dX - \Delta t \iint_{\Pi_T^2} D_+^t \text{sign}_\varepsilon(A(u_{\Delta t}) - A(u)) D_- D_+ A(u_\Delta) \varphi dX. \end{aligned}$$

Consider the first term on the left-hand side. Let ε tend to zero as in the proof of Lemma 4.6. Using the Leibniz rule for difference quotients we obtain

$$D_-^t \varphi = S_{-\Delta t}^t \psi \omega_r D_-^t \rho_{r_0} + D_-^t \psi \omega_r \rho_{r_0}.$$

Recall that $\partial_s \rho_{r_0} = -\partial_t \rho_{r_0}$, so adding and subtracting gives

$$\varphi_s = -S_{-\Delta t}^t \psi \omega_r \partial_t \rho_{r_0} + \Delta t D_-^t \psi \omega_r \partial_s \rho_{r_0}.$$

Hence,

$$\begin{aligned} & \iint_{\Pi_T^2} |u_\Delta - u| (D_-^t \varphi + \varphi_s) dX \\ & = \iint_{\Pi_T^2} |u_\Delta - u| S_{-\Delta t}^t \psi \omega_r (D_-^t \rho_{r_0} - \partial_t \rho_{r_0}) dX \\ & \quad + \Delta t \iint_{\Pi_T^2} |u_\Delta - u| D_-^t \psi \omega_r \partial_s \rho_{r_0} dX + \iint_{\Pi_T^2} |u_\Delta - u| D_-^t \psi \omega_r \rho_{r_0} dX. \end{aligned}$$

Finally, we use that

$$(6.6) \quad D_-^t \psi = \delta_{\Delta t}^+(t - \nu) - \delta_{\Delta t}^+(t - \tau).$$

Concerning the second term on the left-hand side, we apply (3.8). The lemma now follows by sending ε to zero, as in the proof of Lemma 4.6. $\square$

As seen by comparing Lemmas 6.2 and 5.3, there is one new term. To estimate this term we will use a result from [12, p. 1853].

Lemma 6.3. *Let u_j^n be the solution to (6.1). Suppose the CFL condition (6.2) is satisfied. Then there exists a constant L such that*

$$\Delta x \sum_j |D_+ A(u_j^m) - D_+ A(u_j^n)| \leq L \sqrt{(m-n)\Delta t}, \quad \text{for all } m \geq n.$$

Estimate 6.1. Suppose (6.2) is satisfied. Then

$$\begin{aligned} & \left| \Delta t \iint_{\Pi_T^2} D_+^t \text{sign}_\varepsilon(A(u_\Delta) - A(u)) D_- D_+ A(u_\Delta) \varphi dX \right| \\ & \leq C \frac{\sqrt{\Delta t}}{\Delta x} + C \frac{\Delta t}{r_0} \left(1 + \frac{\Delta t}{r_0} \right) + C \Delta t. \end{aligned}$$

Proof. Integration by parts for difference quotients gives

$$\begin{aligned} & \Delta t \iint_{\Pi_T^2} D_+^t \text{sign}_\varepsilon(A(u_\Delta) - A(u)) D_- D_+ A(u_\Delta) \varphi dX \\ & = -\Delta t \iint_{\Pi_T^2} \text{sign}_\varepsilon(A(u_\Delta) - A(u)) S_{-\Delta t}^t D_- D_+ A(u_\Delta) D_-^t \varphi dX \\ & \quad - \Delta t \iint_{\Pi_T^2} \text{sign}_\varepsilon(A(u_\Delta) - A(u)) D_-^t D_- D_+ A(u_\Delta) \varphi dX =: T_1 + T_2, \end{aligned}$$

where we have used that

$$D_-^t (D_- D_+ A(u_\Delta) \varphi) = S_{-\Delta t}^t D_- D_+ A(u_\Delta) D_-^t \varphi + D_-^t D_- D_+ A(u_\Delta) \varphi.$$

Let us consider T_1 first. By the Leibniz rule for difference quotients,

$$\begin{aligned} T_1 & = \Delta t \iint_{\Pi_T^2} \text{sign}_\varepsilon(A(u_\Delta) - A(u)) S_{-\Delta t}^t D_- D_+ A(u_\Delta) S_{-\Delta t}^t \psi \omega_r D_-^t \rho_{r_0} dX \\ & \quad + \Delta t \iint_{\Pi_T^2} \text{sign}_\varepsilon(A(u_\Delta) - A(u)) S_{-\Delta t}^t D_- D_+ A(u_\Delta) D_-^t \psi \omega_r \rho_{r_0} dX \\ & =: T_{1,1} + T_{1,2}. \end{aligned}$$

Using equation (6.6),

$$|T_{1,2}| \leq \Delta t \int_{\Pi_T} |S_{-\Delta t}^t D_- D_+ A(u_\Delta)| (|\delta_{\Delta t}^+(t - \nu)| + |\delta_{\Delta t}^+(t - \tau)|) dx dt \leq C \Delta t,$$

as $\|D_- D_+ A(u_\Delta(\cdot, t))\|_{L^1(\mathbb{R})}$ is bounded independent of Δ and t ([13, Lemma 3.4]). Now, as in Lemma 4.11,

$$|D_-^t \rho_{r_0}| \leq \frac{\|\rho'\|_{L^\infty}}{r_0^2} \mathbb{1}_{\{|t-s| \leq r_0 + \Delta t\}}(t, s)$$

and, therefore,

$$|T_{1,1}| \leq C \Delta t \frac{r_0 + \Delta t}{r_0^2} \int_{\Pi_T} |S_{-\Delta t}^t D_- D_+ A(u_\Delta)| dx dt \leq C \frac{\Delta t}{r_0} \left(1 + \frac{\Delta t}{r_0} \right).$$

Next, we consider T_2 . By Lemma 6.3,

$$\begin{aligned} |T_2| &\leq \Delta t \int_{\nu}^{\tau} \int_{\mathbb{R}} |D_-^t D_- D_+ A(u_{\Delta})| \, dx dt \\ &\leq 2 \frac{\Delta t}{\Delta x} \int_{\nu}^{\tau} \|D_-^t D_+ A(u_{\Delta}(\cdot, t))\|_{L^1(\mathbb{R})} \, dt \\ &\leq 2TL \frac{\sqrt{\Delta t}}{\Delta x}. \end{aligned} \quad \square$$

Proof of Theorem 6.1. We start out from Lemma 6.2 with $A(\sigma) = \hat{A}(\sigma) + \eta\sigma$, where $\hat{A}$ is the original degenerate diffusion function. By Estimate 6.1,

$$\begin{aligned} &\left| \Delta t \iint_{\Pi_T^2} D_+^t \text{sign}_{\varepsilon}(A(u_{\Delta}) - A(u)) D_- D_+ A(u_{\Delta}) \varphi \, dX \right| \\ &\leq C \frac{\sqrt{\Delta t}}{\Delta x} + C \frac{\Delta t}{r_0} \left(1 + \frac{\Delta t}{r_0}\right) + C \Delta t =: E_4. \end{aligned}$$

Since all the estimates from Section 4.1 apply, we obtain

$$\begin{aligned} &\iint_{\Pi_T^2} |u_{\Delta} - u| \delta^{\Delta t}(t - \tau) \omega_r(x - y) \rho_{r_0}(t - s) \, dX \\ &\leq \iint_{\Pi_T^2} |u_{\Delta x} - u| \delta^{\Delta t}(t - \nu) \omega_r(x - y) \rho_{r_0}(t - s) \, dX \\ &\quad + E_1 + E_2 + E_3 + E_4, \end{aligned}$$

where E_1, E_2, E_3 are defined respectively in (4.9), (4.10), and (5.5). Let us make the assumption that $\nu = t_m$ and $\tau = t_n$ for some $m, n \in \mathbb{N}$. Then the above inequality takes the form

$$\kappa(t_n) \leq \kappa(t_m) + E_1 + E_2 + E_3 + E_4,$$

where

$$\kappa(t) := \int_{\mathbb{R}} \int_{\Pi_T} |u_{\Delta}(x, t) - u(y, s)| \omega_r(x - y) \rho_{r_0}(t - s) \, dy ds dx.$$

Applying Lemmas 4.12 and 5.1, and following the reasoning given in the semi-discrete case, we deduce

$$\begin{aligned} &\|u_{\Delta}(\cdot, t_n) - u(\cdot, t_n)\|_{L^1(\mathbb{R})} \\ &\leq \|u_{\Delta}^0 - u^0\|_{L^1(\mathbb{R})} + (L_c + L_d) t_m \\ &\quad + 2 \left(L_c r_0 + |u^0|_{BV(\mathbb{R})} r \right) + C(1 + r + \Delta x)^2 \left(1 + \frac{\Delta x}{r} \right)^3 \frac{\Delta x}{r^2} \\ &\quad + C \frac{\Delta x}{r_0} + C \frac{\Delta t}{r_0} \left(1 + r_0 + \frac{\Delta t}{r_0} \right) + C \frac{\sqrt{\Delta t}}{\Delta x} \\ &\leq \|u_{\Delta}^0 - u^0\|_{L^1(\mathbb{R})} + C \left(\frac{\Delta x}{r^2} + \frac{\Delta x + \Delta t}{r_0} + \frac{\sqrt{\Delta t}}{\Delta x} + r + r_0 \right), \end{aligned}$$

where L_d is the constant in Lemma 5.1 and L_c is the constant from Lemma 4.12. Let $r = r_0, \Delta x = r^3$ and $\Delta t = r^8$. It follows that

$$\|u_{\Delta}(\cdot, t_n) - u(\cdot, t_n)\|_{L^1(\mathbb{R})} \leq \|u_{\Delta}^0 - u^0\|_{L^1(\mathbb{R})} + C \Delta x^{1/3}.$$

Finally, we send $\eta \rightarrow 0$ to conclude the proof of the theorem. $\square$

7. CONCLUDING REMARKS

The added complexity of convection-diffusion equations versus conservation laws [22] arises as a result of the need to work with an explicit form of the parabolic dissipation term. This is reflected in the fact that the rate of convergence is lowered to $1/3$ (from $1/2$ for conservation laws) due to Estimate 4.4 and Estimate 4.5. The optimality of the $\frac{1}{3}$ rate is an open problem. Concerning Section 6 (explicit schemes), one may wonder if it is possible to remove the strengthened CFL condition $\Delta t \sim \Delta x^{8/3}$ (the usual one demands $\Delta t \sim \Delta x^2$). The difficulty is that the parabolic dissipation term is needed to balance the temporal error contribution as well as to carry out the doubling-of-the-variables argument, and this forces us to impose a stronger relation between Δt and Δx in order to appropriately control the temporal error contribution. We do not know if the condition $\Delta t \sim \Delta x^{8/3}$ is genuinely needed or is simply an artifact of our method of proof. Finally, we are currently investigating the multidimensional case. For the semi-discrete scheme the main challenge seems to be the adaptation of Estimate 4.5, or more precisely to produce a multidimensional analogue of (4.7). As an additional difficulty, Lemma 6.3 is not available in several space dimensions; see [12]. At the moment our multidimensional convergence rates are lower than in the one-dimensional case.

REFERENCES

- [1] Boris Andreianov and Noureddine Igbida, *On uniqueness techniques for degenerate convection-diffusion problems*, Int. J. Dyn. Syst. Differ. Equ. **4** (2012), no. 1-2, 3–34, DOI 10.1504/IJDSDE.2012.045992. MR2908633
- [2] B. Andreianov, M. Bendahmane, and K. H. Karlsen, *Discrete duality finite volume schemes for doubly nonlinear degenerate hyperbolic-parabolic equations*, J. Hyperbolic Differ. Equ. **7** (2010), no. 1, 1–67, DOI 10.1142/S0219891610002062. MR2646796 (2011d:65241)
- [3] D. Aregba-Driollet, R. Natalini, and S. Tang, *Explicit diffusive kinetic schemes for nonlinear degenerate parabolic systems*, Math. Comp. **73** (2004), no. 245, 63–94 (electronic), DOI 10.1090/S0025-5718-03-01549-7. MR2034111 (2004m:65136)
- [4] F. Bouchut, F. R. Guarguaglini, and R. Natalini, *Diffusive BGK approximations for nonlinear multidimensional parabolic equations*, Indiana Univ. Math. J. **49** (2000), no. 2, 723–749, DOI 10.1512/iumj.2000.49.1811. MR1793689 (2001k:35162)
- [5] José Carrillo, *Entropy solutions for nonlinear degenerate problems*, Arch. Ration. Mech. Anal. **147** (1999), no. 4, 269–361, DOI 10.1007/s002050050152. MR1709116 (2000m:35132)
- [6] Gui-Qiang Chen and Kenneth H. Karlsen, *Quasilinear anisotropic degenerate parabolic equations with time-space dependent diffusion coefficients*, Commun. Pure Appl. Anal. **4** (2005), no. 2, 241–266, DOI 10.3934/cpaa.2005.4.241. MR2149515 (2006h:35119)
- [7] Bernardo Cockburn, *Continuous dependence and error estimation for viscosity methods*, Acta Numer. **12** (2003), 127–180, DOI 10.1017/S0962492902000107. MR2249155 (2007e:35183)
- [8] Bernardo Cockburn and Pierre-Alain Gremaud, *A priori error estimates for numerical methods for scalar conservation laws. I. The general approach*, Math. Comp. **65** (1996), no. 214, 533–573, DOI 10.1090/S0025-5718-96-00701-6. MR1333308 (96g:65089)
- [9] B. Cockburn and G. Gripenberg, *Continuous dependence on the nonlinearities of solutions of degenerate parabolic equations*, J. Differential Equations **151** (1999), no. 2, 231–251, DOI 10.1006/jdeq.1998.3499. MR1669570 (2000a:35101)
- [10] Constantine M. Dafermos, *Hyperbolic conservation laws in continuum physics*, 3rd ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 325, Springer-Verlag, Berlin, 2010. MR2574377 (2011i:35150)
- [11] Steinar Evje and Kenneth H. Karlsen, *Degenerate convection-diffusion equations and implicit monotone difference schemes*, Hyperbolic problems: theory, numerics, applications, Vol. I (Zürich, 1998), Internat. Ser. Numer. Math., vol. 129, Birkhäuser, Basel, 1999, pp. 285–294. MR1717198 (2000k:65137)
- [12] Steinar Evje and Kenneth H. Karlsen, *Monotone difference approximations of BV solutions to degenerate convection-diffusion equations*, SIAM J. Numer. Anal. **37** (2000), no. 6, 1838–1860 (electronic), DOI 10.1137/S0036142998336138. MR1766850 (2001g:65110)

- [13] Steinar Evje and Kenneth Hvistendahl Karlsen, *Discrete approximations of BV solutions to doubly nonlinear degenerate parabolic equations*, Numer. Math. **86** (2000), no. 3, 377–417, DOI 10.1007/s002110000156. MR1785415 (2001g:65109)
- [14] Steinar Evje and Kenneth H. Karlsen, *An error estimate for viscous approximate solutions of degenerate parabolic equations*, J. Nonlinear Math. Phys. **9** (2002), no. 3, 262–281, DOI 10.2991/jnmp.2002.9.3.3. MR1916386 (2003f:35176)
- [15] Robert Eymard, Thierry Gallouët, and Raphaële Herbin, *Error estimate for approximate solutions of a nonlinear convection-diffusion problem*, Adv. Differential Equations **7** (2002), no. 4, 419–440. MR1869118 (2002h:35156)
- [16] Robert Eymard, Thierry Gallouët, Raphaële Herbin, and Anthony Michel, *Convergence of a finite volume scheme for nonlinear degenerate parabolic equations*, Numer. Math. **92** (2002), no. 1, 41–82, DOI 10.1007/s002110100342. MR1917365 (2003e:65159)
- [17] Helge Holden, Kenneth H. Karlsen, Knut-Andreas Lie, and Nils Henrik Risebro, *Splitting methods for partial differential equations with rough solutions*, Analysis and MATLAB programs. EMS Series of Lectures in Mathematics, European Mathematical Society (EMS), Zürich, 2010. MR2662342 (2011j:65002)
- [18] Kenneth Hvistendahl Karlsen and Nils Henrik Risebro, *On the uniqueness and stability of entropy solutions of nonlinear degenerate parabolic equations with rough coefficients*, Discrete Contin. Dyn. Syst. **9** (2003), no. 5, 1081–1104, DOI 10.3934/dcds.2003.9.1081. MR1974417 (2004h:35102)
- [19] Kenneth Hvistendahl Karlsen and Nils Henrik Risebro, *Convergence of finite difference schemes for viscous and inviscid conservation laws with rough coefficients*, M2AN Math. Model. Numer. Anal. **35** (2001), no. 2, 239–269, DOI 10.1051/m2an:2001114. MR1825698 (2002b:35138)
- [20] K. H. Karlsen, U. Koley, and N. H. Risebro, *An error estimate for the finite difference approximation to degenerate convection-diffusion equations*, Numer. Math. **121** (2012), no. 2, 367–395, DOI 10.1007/s00211-011-0433-9. MR2917166
- [21] S. N. Kružkov, *First order quasilinear equations with several independent variables*. (Russian), Mat. Sb. (N.S.) **81 (123)** (1970), 228–255. MR0267257 (42 #2159)
- [22] N. N. Kuznecov, *The accuracy of certain approximate methods for the computation of weak solutions of a first order quasilinear equation* (Russian), Ž. Vyčisl. Mat. i Mat. Fiz. **16** (1976), no. 6, 1489–1502, 1627. MR0483509 (58 #3510)
- [23] G. E. Ladas and V. Lakshmikantham, *Differential equations in abstract spaces*, Academic Press, New York, 1972. Mathematics in Science and Engineering, Vol. 85. MR0460832 (57 #824)
- [24] Mario Ohlberger, *A posteriori error estimates for vertex centered finite volume approximations of convection-diffusion-reaction equations*, M2AN Math. Model. Numer. Anal. **35** (2001), no. 2, 355–387, DOI 10.1051/m2an:2001119. MR1825703 (2002a:65142)
- [25] Tamir Tassa, *Regularity of weak solutions of the nonlinear Fokker-Planck equation*, Math. Res. Lett. **3** (1996), no. 4, 475–490. MR1406013 (98a:35079)
- [26] A. I. Vol’pert, *Spaces BV and quasilinear equations* (Russian), Mat. Sb. (N.S.) **73 (115)** (1967), 255–302. MR0216338 (35 #7172)
- [27] A. I. Vol’pert and S. I. Hudjaev, *The Cauchy problem for second order quasilinear degenerate parabolic equations* (Russian), Mat. Sb. (N.S.) **78 (120)** (1969), 374–396. MR0264232 (41 #8828)
- [28] Zhuo Qun Wu and Jing Xue Yin, *Some properties of functions in BV_x and their applications to the uniqueness of solutions for degenerate quasilinear parabolic equations*, Northeast. Math. J. **5** (1989), no. 4, 395–422. MR1053519 (91j:35153)

CENTER OF MATHEMATICS FOR APPLICATIONS (CMA), UNIVERSITY OF OSLO, P.O. BOX 1053, BLINDERN, N-0316 OSLO, NORWAY

E-mail address: kennethk@math.uio.no

CENTER OF MATHEMATICS FOR APPLICATIONS (CMA), UNIVERSITY OF OSLO, P.O. BOX 1053, BLINDERN, N-0316 OSLO, NORWAY

E-mail address: nilshr@math.uio.no

CENTER OF MATHEMATICS FOR APPLICATIONS (CMA), UNIVERSITY OF OSLO, P.O. BOX 1053, BLINDERN, N-0316 OSLO, NORWAY

E-mail address: erlenbs@math.uio.no

E-mail address: erlenbs@gmail.com

ON THE CONVERGENCE RATE OF FINITE DIFFERENCE METHODS FOR DEGENERATE CONVECTION-DIFFUSION EQUATIONS IN SEVERAL SPACE DIMENSIONS

K. H. KARLSEN, N. H. RISEBRO, AND E. B. STORRØSTEN

ABSTRACT. We analyze upwind difference methods for strongly degenerate convection-diffusion equations in several spatial dimensions. We prove that the local L^1 -error between the exact and numerical solutions is $\mathcal{O}(\Delta x^{2/(19+d)})$, where d is the spatial dimension and Δx is the grid size. The error estimate is robust with respect to vanishing diffusion effects. The proof makes effective use of specific kinetic formulations of the difference method and the convection-diffusion equation. This paper is a continuation of [25], in which the one-dimensional case was examined using the Kružkov-Carrillo entropy framework.

CONTENTS

1. Introduction	1
2. Viscosity approximations and entropy solutions	6
3. Difference method and main result	7
4. Kinetic formulations	8
4.1. Kinetic formulation of convection-diffusion equation	8
4.2. Kinetic formulation of discrete equations	9
4.3. Various regularizations	12
5. Proof of Theorem 3.1	15
5.1. Main error equation	15
5.2. Dissipative term	18
5.3. Bounding error terms	22
5.4. Concluding the proof of Theorem 3.1	32
Appendix A. Well-posedness of difference method	37
References	39

1. INTRODUCTION

The design of numerical methods for convection-diffusion problems is important for many applications in science and engineering. It is especially challenging to construct accurate methods for nonlinear problems in which the “diffusion part” is small or vanishing, relative to the “convection part” of the problem. Connected to this is the difficult problem of deriving error estimates for numerical methods that are robust in the singular limit as the diffusion coefficient vanishes, thereby avoiding the usual exponential growth of error constants.

In this paper we are interested in deriving error estimates for a class of finite difference methods for nonlinear, possibly strongly degenerate, convection-diffusion

Date: September 3, 2015.

2010 Mathematics Subject Classification. Primary: 65M06, 65M15; Secondary: 35K65, 35L65.

Key words and phrases. Degenerate convection-diffusion equations, entropy conditions, finite difference methods, error estimates.

problems of the form

$$(1.1) \quad \begin{cases} \partial_t u + \nabla \cdot f(u) = \Delta A(u), & (t, x) \in \Pi_T, \\ u(0, x) = u_0(x), & x \in \mathbb{R}^d, \end{cases}$$

where $\Pi_T = (0, T) \times \mathbb{R}^d$, $T > 0$, $d \geq 1$, and $u : \Pi_T \rightarrow \mathbb{R}$ is the unknown function that is sought. The initial datum u_0 is an integrable and bounded function, while the *flux* function $f : \mathbb{R} \rightarrow \mathbb{R}^d$ and the *diffusion* function $A : \mathbb{R} \rightarrow \mathbb{R}$ satisfy the standing assumptions

$$(1.2) \quad f, A \text{ locally } C^1; A(0) = 0; A \text{ is nondecreasing.}$$

By *strongly degenerate* it is meant that we allow for $A'(u) = 0$ for all u in some interval $[\alpha, \beta] \subset \mathbb{R}$. The resulting class of equations therefore contains parabolic and hyperbolic equations, as well as a mix thereof. In the nondegenerate (uniformly parabolic) case $A'(\cdot) > 0$, it is well known that (1.1) admits a unique classical solution. On the other hand, for strongly degenerate equations with discontinuous solutions, the well-posedness is ensured only in a class of weak solutions satisfying an entropy condition. The following result is known: For $u_0 \in L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$, there exists a unique solution $u \in C((0, T); L^1(\mathbb{R}^d))$, $u \in L^\infty(\Pi_T)$ of (1.1) such that $\partial_x A(u) \in L^2(\Pi_T)$ and for all convex functions S with $q'_S = S'f'$ and $r'_S = S'A'$,

$$\partial_t S(u) + \nabla \cdot q_S(u) - \Delta r_S(u) \leq 0, \quad \text{weakly on } [0, T) \times \mathbb{R}^d.$$

These inequalities are referred to as entropy inequalities and the corresponding solution is called an entropy solution.

For conservation laws ($A' \equiv 0$), the well-posedness of entropy solutions is a celebrated result due to Kruřkov [26]. Carrillo [8] extended this result to degenerate parabolic problems such as (1.1). For uniqueness of entropy solutions in the *BV* class, see [35, 36]. An alternative well-posedness theory, based on the so-called kinetic formulation, was developed by Lions, Perthame, and Tadmor [29] and Chen and Perthame [12]. We refer to [2, 16] for an overview of the relevant literature on hyperbolic and mixed hyperbolic-parabolic problems.

In this paper we derive error estimates for numerical approximations of entropy solutions to convection-diffusion equations. Convergence results, without error estimates, have been obtained for difference methods [18, 17, 24]; finite volume methods [21, 1]; splitting methods [22]; and BGK approximations [3, 5], to mention a few references. For a posteriori error estimates for finite volume methods, see [31].

We are herein interested in estimating the error committed by a class of monotone difference methods. The monotone methods make use of an upwind discretization of the convection term and a centred discretization of the parabolic term. For notational simplicity in the introduction, let us assume $f^{i,\prime}(\cdot) \geq 0$ and consider the prototype (semi-discrete) difference method

$$\frac{d}{dt} u_\alpha + \sum_{i=1}^d \frac{f^i(u_\alpha) - f^i(u_{\alpha-e_i})}{\Delta x} = \sum_{i=1}^d \frac{A(u_{\alpha+e_i}) - 2A(u_\alpha) + A(u_{\alpha-e_i}))}{\Delta x^2},$$

where $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{Z}^d$ is a multi-index, e_i is the i th unit vector in $\mathbb{R}^d$, and $\Delta x > 0$ is the spatial grid size. Although our methods are semi-discrete, i.e., not discretized in time, the results and proofs can be adjusted to cover some fully discrete methods, such as the implicit method analyzed in [18]. We refer to [25] for a discussion of this topic when $d = 1$.

Denote by $u_{\Delta x}$ the piecewise constant interpolant linked to u_α . The goal is to determine a number (convergence rate) $\gamma > 0$ such that

$$(1.3) \quad \|u_{\Delta x}(t, \cdot) - u(t, \cdot)\|_{L^1} \leq C \Delta x^\gamma, \quad (u_0 \in BV),$$

for some constant $C > 0$ independent of Δx and (the smallness of) A' .

In the purely hyperbolic case ($A' \equiv 0$), a prominent result due to Kuznetsov [28] says that γ is $1/2$ for monotone difference methods, as well as for the vanishing viscosity method. Influenced by [28], a number of works have further developed the “Kruřkov-Kuznetsov” error estimation theory for conservation laws. We refer to [6, 13] for an overview of the relevant results. Making use of the kinetic formulation, Perthame [33] provided an alternative route to error estimates.

With regard to convection-diffusion equations (1.1) with $A'(\cdot) \geq 0$, the subject of error estimates is significantly more difficult. It is only recently that there has been some progress. The simplest case is the vanishing viscosity method. Denote by u^η the solution of the uniformly parabolic equation

$$(1.4) \quad u_t^\eta + \nabla \cdot f(u^\eta) = \Delta A^\eta(u^\eta), \quad A^\eta(u) = A(u) + \eta u,$$

where $\eta > 0$ is a (small) viscosity parameter. We have the following error estimate for the viscosity approximation u^η :

$$(1.5) \quad \|u^\eta(\cdot, t) - u(\cdot, t)\|_{L^1} \leq C \sqrt{\eta}, \quad (u_0 \in BV),$$

where u is the entropy solution of (1.1). A “Kruřkov-Kuznetsov” type proof of this result is given in [19], see also [20] for a boundary value problem. The error bound (1.5) can also be seen as an outcome of continuous dependence estimates [13, 10] or the kinetic formulation [11, 30].

For conservation laws, the error estimate (1.5) for the viscous equation reveals what to expect for monotone difference methods [28]. This suggestive link breaks down for degenerate convection-diffusion equations (1.1), cf. [25], a fact that may foreshadow added difficulties coming from a second order operator. Indeed, for general A satisfying (1.2) and in one spatial dimension, the work [23] established (1.3) with $\gamma = 1/11$, a rate that was recently improved to $\gamma = 1/3$ in [25]. Although $\gamma = 1/3$ is the best available rate at the moment, its optimality is unknown and also far from the convergence rate $\gamma = 1/2$ known to be optimal for conservation laws. But in spite of that, with a linear diffusion function A , the convergence rate improves to $\gamma = 1/2$ [23, 25].

Apart from a result ($\gamma = 1/2$) for linear convection-diffusion equations [11], we are not aware of any results for multi-dimensional equations (1.1) with a degenerate, nonlinear diffusion part. In this paper we establish (1.3) with

$$(1.6) \quad \gamma = \frac{2}{19 + d} \quad (d \text{ is the spatial dimension}),$$

for general diffusion functions A obeying (1.2).

A technical aspect of the proof of (1.3) is that we are not applying the difference method directly to (1.1) but rather to (1.4). Denoting the corresponding numerical solution by $u_{\Delta x}^\eta$, we will prove that (1.3) holds with $u_{\Delta x}, u$ replaced by $u_{\Delta x}^\eta, u^\eta$, respectively, and that the error constant C is not depending on the parameter η . Our original claim (1.3) follows from this, since we have the error estimate (1.5).

To help motivate the technical arguments coming later, let us lay out a “high-level” overview of the analysis and some of the difficulties involved. As just alluded to, we will mostly work under the assumption $A' > 0$. As a consequence no information is lost upon working with $A(u)$ instead of u in the kinetic formulation (compare with the u -based formulation in [12]). Set $B = A^{-1}$ and define g by $g \circ A = f$. Then the solution u of (1.1) satisfies

$$(1.7) \quad B'(\zeta) \partial_t \chi_{A(u)} + g'(\zeta) \cdot \nabla \chi_{A(u)} - \Delta \chi_{A(u)} = \partial_\zeta m_{A(u)},$$

where

$$m_{A(u)} = m_{A(u)}(\zeta) = \delta(\zeta - A(u)) |\nabla A(u)|^2,$$

$$\chi_{A(u)} = \chi_{A(u)}(\zeta) = \begin{cases} 1 & \text{if } 0 < \zeta \leq A(u), \\ -1 & \text{if } A(u) \leq \zeta < 0, \\ 0 & \text{otherwise.} \end{cases}$$

This new formulation, although restricted to nondegenerate (isotropic) diffusion, allows for a simpler proof of the L^1 contraction property and thus the error estimate (1.3). More specifically, certain error terms linked to the regularization of the χ function [12] can be avoided, a fact that we use to our benefit.

Now we indicate how (1.7) leads to the L^1 contraction property. Let u and v be solutions to (1.1) with initial values u_0 and v_0 , respectively. Following [12, 33], we introduce the microscopic contraction functional

$$(1.8) \quad Q_{u,v}(\xi) = |\chi_u(\xi)| + |\chi_v(\xi)| - 2\chi_u(\xi)\chi_v(\xi).$$

Under the change of variable $\zeta = A(\xi)$,

$$|u - v| = \int_{\mathbb{R}} Q_{u,v}(\xi) d\xi = \int_{\mathbb{R}} B'(\zeta) Q_{A(u), A(v)}(\zeta) d\zeta,$$

and hence

$$(1.9) \quad \begin{aligned} \partial_t |u - v| &= \int_{\mathbb{R}} B'(\zeta) \partial_t Q_{A(u), A(v)}(\zeta) d\zeta \\ &= \int_{\mathbb{R}} B'(\zeta) \partial_t |\chi_{A(u)}(\zeta)| d\zeta + \int_{\mathbb{R}} B'(\zeta) \partial_t |\chi_{A(v)}(\zeta)| d\zeta \\ &\quad - 2 \int_{\mathbb{R}} B'(\zeta) \partial_t (\chi_{A(u)}(\zeta) \chi_{A(v)}(\zeta)) d\zeta. \end{aligned}$$

In view of (1.7), the chain rule yields

$$(1.10) \quad B'(\zeta) \partial_t |\chi_{A(u)}| + g'(\zeta) \cdot \nabla |\chi_{A(u)}| - \Delta |\chi_{A(u)}| = \text{sign}(\zeta) \partial_\zeta m_{A(u)},$$

with an analogous equation for v . Using the equations for $\chi_{A(u)}$, $\chi_{A(v)}$ and Leibniz's product rule, we easily check that

$$(1.11) \quad \begin{aligned} B'(\zeta) \partial_t (\chi_{A(u)} \chi_{A(v)}) + g'(\zeta) \cdot \nabla (\chi_{A(u)} \chi_{A(v)}) - \Delta (\chi_{A(u)} \chi_{A(v)}) \\ = \chi_{A(v)} \partial_\zeta m_{A(u)} + \chi_{A(u)} \partial_\zeta m_{A(v)} - 2 \nabla \chi_{A(u)} \cdot \nabla \chi_{A(v)}. \end{aligned}$$

Making use of (1.10) and (1.11) in (1.9) yields

$$(1.12) \quad \begin{aligned} \partial_t |u - v| &= - \int_{\mathbb{R}} g'(\zeta) \cdot \nabla Q_{A(u), A(v)}(\zeta) d\zeta + \int_{\mathbb{R}} \Delta Q_{A(u), A(v)}(\zeta) d\zeta \\ &\quad + \int_{\mathbb{R}} D(\zeta) d\zeta, \end{aligned}$$

where

$$\begin{aligned} D(\zeta) &= \left(\text{sign}(\zeta) - 2\chi_{A(v)}(\zeta) \right) \partial_\zeta m_{A(u)} + \left(\text{sign}(\zeta) - 2\chi_{A(u)}(\zeta) \right) \partial_\zeta m_{A(v)} \\ &\quad + 4 \nabla \chi_{A(u)}(\zeta) \cdot \nabla \chi_{A(v)}(\zeta) \\ &=: D_1(\zeta) + D_2(\zeta) + D_3(\zeta); \end{aligned}$$

the term $D(\cdot)$ accounts for the parabolic dissipation effects associated with u, v . Integrating (1.12) in x gives

$$\frac{d}{dt} \int |u(t, x) - v(t, x)| dx = \int \int_{\mathbb{R}} D(\zeta) d\zeta dx.$$

Although the computations have been formal up to this point, they are valid when interpreted in the sense of distributions. Moreover, as will be seen later, these computations can with some effort be replicated at the discrete level, i.e., when we replace the function v by the numerical solution $u_{\Delta x}$.

Clearly, the L^1 -contraction property follows if we can confirm that

$$(1.13) \quad \int_{\mathbb{R}} D(\zeta) d\zeta \leq 0.$$

This step is rather delicate and will ask for a regularization of the χ functions. Indeed, the hard part of the proof leading up to (1.3), (1.6) is related to this step. Let us for the moment ignore the regularization procedure, and continue with formal computations. Note that

$$\text{sign}(\zeta) - 2\chi_{A(v)}(\zeta) = \text{sign}(\zeta - A(v)),$$

and thus, after an integration by parts followed by an application of the chain rule,

$$\int_{\mathbb{R}} D_1(\zeta) d\zeta = -2 \int_{\mathbb{R}} \delta(\zeta - A(u)) \delta(\zeta - A(v)) |\nabla A(u)|^2 d\zeta.$$

Similarly,

$$\int_{\mathbb{R}} D_2(\zeta) d\zeta = -2 \int_{\mathbb{R}} \delta(\zeta - A(u)) \delta(\zeta - A(v)) |\nabla A(v)|^2 d\zeta.$$

Again by the chain rule,

$$D_3(\zeta) = 4\delta(\zeta - A(u))\delta(\zeta - A(v))\nabla A(u) \cdot \nabla A(v).$$

Combining these formal computations we finally arrive at (1.13):

$$\int_{\mathbb{R}} D(\zeta) d\zeta = -2 \int_{\mathbb{R}} \delta(\zeta - A(u))\delta(\zeta - A(v)) |\nabla A(u) - \nabla A(v)|^2 d\zeta \leq 0.$$

One crucial insight in [25] is that the convergence rate can be improved if one can send a certain parameter ε to zero independently of the grid size Δx , where ε controls the regularization of the Kruřkov entropies. In this paper the regularization of the entropies is replaced by the regularization of the χ functions, and as before we would like to send ε to zero independently of Δx (and other parameters). It turns out that in one spatial dimension we can do this, reaching the convergence rate $\gamma = 1/3$ as in [25]. In several dimensions we have not been able to carry out this “ $\varepsilon \rightarrow 0$ before other parameters” program.

A serious difficulty stems from the lack of a chain rule for finite differences, in combination with the highly nonlinear nature of the dissipation function $D(\cdot)$, resulting in a series of intricate error terms. A feature of the kinetic approach is that the crucial error term can be expressed via the parabolic dissipation measure. To be a bit more precise, at the continuous level, the convergence rate $\gamma = 1/3$ in the one-dimensional case depends decisively on the (weak) continuity of the map

$$(1.14) \quad c \mapsto \int_{\mathbb{R}} \delta(\zeta - c) m_{A(u)}(\zeta) d\zeta = \delta(c - A(u)) (\partial_x A(u))^2,$$

where u is the entropy solution and $m_{A(u)}$ is the parabolic dissipation measure. The continuity of this map follows from (1.7). Unfortunately, in several space dimensions the continuity becomes a subtle matter, since the parabolic dissipation measure splits into directional components,

$$m_{A(u)} = \sum_{i=1}^d m_{A(u)}^i, \quad m_{A(u)}^i = \delta(\zeta - A(u)) (\partial_{x_i} A(u))^2.$$

It appears difficult to claim from the kinetic equation (1.7) the continuity of the individual components

$$c \mapsto \int_{\mathbb{R}} \delta(\zeta - c) m_{A(u)}^i(\zeta) d\zeta = \delta(c - A(u)) (\partial_{x_i} A(u))^2, \quad i = 1, \dots, d.$$

Not being able to send the χ -regularisation parameter ε to zero, we must instead balance ε against the grid size Δx and a number of other parameters, at long last arriving at (1.3) with the convergence rate (1.6).

The optimality of (1.6) ($d > 1$), in the $L^\infty \cap BV$ class, is an open problem. It is informative to compare with recent results on viscosity solutions and error bounds for degenerate fully nonlinear elliptic and parabolic equations. We refer to Krylov [27], Barles and Jakobsen [4], and Caffarelli and Souganidis [7] for some recent works. For monotone approximations of fully nonlinear, first-order equations with Lipschitz solutions, Crandall and Lions [15] proved in 1984 the optimal L^∞ convergence rate $1/2$. However, finding a rate for degenerate second order equations remained an open problem. The first result is due to Krylov with the rate $1/27$. Later Barles and Jakobsen improved this to $1/5$, with a further improvement by Krylov to $1/2$ for equations with special structure. We remark that these results concern equations with convex nonlinearities. Caffarelli and Souganidis proved that there is an algebraic rate of convergence for a class of nonconvex equations. The convergence rate is not explicit but known to be some (small) positive number. Here we should point out that in our framework convexity plays no role; the error estimate applies to general nonlinearities.

The remaining part of this paper is organized as follows: In Section 2 we gather some relevant a priori estimates for nondegenerate convection-diffusion equations and state precisely the definition of an entropy solution. The difference method and the main result are presented in Section 3. In Section 4 we supply certain kinetic formulations of the convection-diffusion and difference equations. Section 5 is devoted to the proof of the main result, achieved through the derivation of an error equation based on the kinetic formulations, along with a lengthy series of estimates bounding “unwanted” terms in this equation. In Appendix A we collect results relating to well-posedness and a priori estimates for the difference method.

2. VISCOSITY APPROXIMATIONS AND ENTROPY SOLUTIONS

Let us define the viscosity approximations. Set $A^\eta(u) := A(u) + \eta u$ for any fixed $\eta > 0$, and consider the the uniformly parabolic problem

$$(2.1) \quad \begin{cases} u_t^\eta + \nabla \cdot f(u^\eta) = \Delta A^\eta(u^\eta), & (t, x) \in \Pi_T, \\ u^\eta(0, x) = u_0(x), & x \in \mathbb{R}^d. \end{cases}$$

It is well known that (2.1) admits a unique classical (smooth) solution. We collect some relevant (standard) estimates from [35].

Lemma 2.1. *Suppose $u_0 \in L^\infty(\mathbb{R}^d) \cap L^1(\mathbb{R}^d) \cap BV(\mathbb{R}^d)$, and let u^η be the unique classical solution of (2.1). Then for any $t > 0$,*

$$\begin{aligned} \|u^\eta(t, \cdot)\|_{L^1(\mathbb{R}^d)} &\leq \|u_0\|_{L^1(\mathbb{R}^d)}, \\ \|u^\eta(t, \cdot)\|_{L^\infty(\mathbb{R}^d)} &\leq \|u_0\|_{L^\infty(\mathbb{R}^d)}, \\ |u^\eta(t, \cdot)|_{BV(\mathbb{R}^d)} &\leq |u_0|_{BV(\mathbb{R}^d)}. \end{aligned}$$

Lemma 2.2. *Suppose $u_0 \in L^\infty(\mathbb{R}^d) \cap L^1(\mathbb{R}^d)$ and $\nabla \cdot (f(u_0) - \nabla A(u_0)) \in L^1(\mathbb{R}^d)$. Let u^η be the unique classical solution of (2.1). Then for any $t_1, t_2 > 0$,*

$$\|u^\eta(t_2, \cdot) - u^\eta(t_1, \cdot)\|_{L^1(\mathbb{R}^d)} \leq \|\nabla \cdot (f(u_0) - \nabla A(u_0))\|_{L^1(\mathbb{R}^d)} |t_2 - t_1|.$$

These results imply that the family $\{u^\eta\}_{\eta>0}$ is relatively compact in $C([0, T]; L^1_{loc}(\mathbb{R}^d))$. If $u = \lim_{\eta \rightarrow 0} u^\eta$, then

$$(2.2) \quad \|u^\eta - u\|_{L^1(\Pi_T)} \leq C\eta^{1/2},$$

for some constant C which does not depend on η , see, e.g., [19]. Moreover, u is an entropy solution according to the following definition:

Definition 2.1. An *entropy solution* of (1.1) is a measurable function $u = u(t, x)$ satisfying:

- (D.1) $u \in L^\infty([0, T]; L^1(\mathbb{R}^d)) \cap L^\infty(\Pi_T) \cap C((0, T); L^1(\mathbb{R}^d))$.
- (D.2) $A(u) \in L^2([0, T]; H^1(\mathbb{R}^d))$.
- (D.3) For all constants $c \in \mathbb{R}$ and all test functions $0 \leq \phi \in C_0^\infty(\mathbb{R}^d \times [0, T])$, the following entropy inequality holds:

$$\begin{aligned} \iint_{\Pi_T} |u - c| \partial_t \phi + \text{sign}(u - c) (f(u) - f(c)) \cdot \nabla \phi + |A(u) - A(c)| \Delta \phi \, dt dx \\ + \int_{\mathbb{R}^d} |u_0 - c| \phi(x, 0) \, dx \geq 0. \end{aligned}$$

The uniqueness of entropy solutions is proved in [8], see the introduction for additional references.

3. DIFFERENCE METHOD AND MAIN RESULT

Let $f = (f^1, \dots, f^d)$, and let Δx denote the mesh size. For simplicity we consider a uniform grid in $\mathbb{R}^d$ consisting of cubes with sides Δx . For a multi-index $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{Z}^d$, we let I_α denote the grid cell

$$I_\alpha = [x_{\alpha_1-1/2}, x_{\alpha_1+1/2}) \times \dots \times [x_{\alpha_d-1/2}, x_{\alpha_d+1/2}),$$

where $x_{j+1/2} = (j + 1/2)\Delta x$ for $j \in \mathbb{Z}$. Let $e_k \in \mathbb{Z}^d$ be the vector with value one in the k -th component and zero otherwise. Then we define the forward and backward discrete partial derivatives in the k -th direction as

$$D_\pm^k(\sigma_\alpha) = \pm \frac{\sigma_{\alpha \pm e_k} - \sigma_\alpha}{\Delta x} \quad k = 1, \dots, d.$$

Definition 3.1. (Numerical flux) We call a function $F \in C^1(\mathbb{R}^2)$ a *monotone two point numerical flux* for f , if $F(u, u) = f(u)$ and

$$\frac{\partial}{\partial u} F(u, v) \geq 0 \quad \text{and} \quad \frac{\partial}{\partial v} F(u, v) \leq 0$$

holds for all u and v . We say that the numerical flux *splits* whenever F can be written

$$F(u, v) = F_1(u) + F_2(v).$$

Note that $F_1' \geq 0$ and $F_2' \leq 0$ whenever F is monotone.

Let F^k be a numerical flux function corresponding to f^k for $k = 1, \dots, d$. The semi-discrete approximation of (1.1) is the solution of the equations

$$(3.1) \quad \begin{cases} \frac{d}{dt} u_\alpha + \sum_{i=1}^d D_-^i F^i(u_\alpha, u_{\alpha+e_i}) = \sum_{i=1}^d D_-^i D_+^i A(u_\alpha), & \alpha \in \mathbb{Z}^d, \, t \in (0, T), \\ u_\alpha(0) = u_{\alpha,0}, & \alpha \in \mathbb{Z}^d, \end{cases}$$

where $u_{\alpha,0} = \frac{1}{\Delta x^d} \int_{I_\alpha} u_0(x) \, dx$. See Appendix A, in particular Lemmas A.2 and A.3, regarding existence and solution properties to this infinite system of ODEs.

Define the piecewise constant (in x) function $u_{\Delta x}$ by

$$(3.2) \quad u_{\Delta x}(t, x) = u_\alpha(t) \text{ for } x \in I_\alpha.$$

Our main result is the following:

Theorem 3.1. *Suppose f and A satisfy (1.2) and the initial function u_0 is in $BV(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d) \cap L^1(\mathbb{R}^d)$. Let F^i be a monotone, Lipschitz, two point numerical flux corresponding to f^i that splits for $1 \leq i \leq d$. Let u be the entropy solution to (1.1) and $u_{\Delta x}$ be defined by (3.2), where u_α is the solution to (3.1).*

Then, for any positive R and T , there exists a constant C depending only on f , A , u_0 , R and T , such that

$$\|u_{\Delta x}(t) - u(t)\|_{L^1(B(0,R))} \leq C \Delta x^{\frac{2}{19+d}}, \quad t \in [0, T].$$

4. KINETIC FORMULATIONS

In this section we supply certain kinetic formulations of the continuous and discrete equations (1.1) and (3.1). As a preparation for the error estimate, we also regularize the kinetic equations by mollification. As explained in the introduction, due to the application of the viscous approximations in the proof of the error estimate, we assume $A' > 0$ for these intermediate results.

4.1. Kinetic formulation of convection-diffusion equation.

Lemma 4.1. *Assume that $A' > 0$ and set $B := A^{-1}$. Let u be the solution of (1.1). Define g by $g(A(z)) = f(z)$ for all $z \in \mathbb{R}$. Let $S \in C^2(\mathbb{R})$,*

$$\begin{aligned} \psi(u) &= \int_0^u S'(z) B'(z) dz, & \psi_A(u) &= \psi(A(u)), \\ q(u) &= \int_0^u S'(z) g'(z) dz, & q_A(u) &= q(A(u)), \end{aligned}$$

and $S_A(u) = S(A(u))$. Then

$$\partial_t \psi_A(u) + \nabla \cdot q_A(u) - \Delta S_A(u) = -S''_A(u) |\nabla A(u)|^2.$$

Proof. Multiplying (1.1) by $\psi'_A(u)$ gives

$$\partial_t \psi_A(u) + \psi'_A(u) \nabla \cdot f(u) = \psi'_A(u) \Delta A(u).$$

Using a change of variables $A(\sigma) = z$,

$$\begin{aligned} q'_A(u) &= \partial_u \left(\int_0^{A(u)} S'(z) g'(z) dz \right) \\ &= \partial_u \left(\int_0^u S'(A(\sigma)) g'(A(\sigma)) A'(\sigma) d\sigma \right) = S'(A(u)) f'(u). \end{aligned}$$

Hence

$$\psi'_A(u) \nabla \cdot f(u) = \nabla \cdot q_A(u).$$

Similarly we obtain $\psi'_A(u) = S'(A(u))$. Finally, observe that

$$\Delta S_A(u) = S''(A(u)) |\nabla A(u)|^2 + \psi'_A(u) \Delta A(u).$$

□

The above entropy equation can be rephrased in terms of the χ function. Recall that for any locally Lipschitz continuous $\Psi : \mathbb{R} \rightarrow \mathbb{R}$,

$$(4.1) \quad \Psi(u) - \Psi(0) = \int_{\mathbb{R}} \Psi'(\xi) \chi(u; \xi) d\xi, \quad (u \in \mathbb{R}).$$

The next lemma reveals the equation satisfied by $\chi(A(u); \zeta)$, where u solves (1.1), i.e., the kinetic formulation of the convection-diffusion equation.

Lemma 4.2. Assume that $A' > 0$ and set $B := A^{-1}$. Let u be the solution of (1.1). Define $\rho(t, x, \zeta) = \chi(A(u(t, x)); \zeta)$. Then

$$(4.2) \quad \begin{cases} B'(\zeta) \partial_t \rho + g'(\zeta) \cdot \nabla \rho - \Delta \rho = \partial_\zeta m & \text{in } \mathcal{D}'((0, T) \times \mathbb{R}^d \times \mathbb{R}), \\ \rho(0, x, \zeta) = \chi(A(u_0(x)); \zeta), & (x, \zeta) \in \mathbb{R}^d \times \mathbb{R}, \end{cases}$$

where

$$m(t, x, \zeta) = \delta(\zeta - A(u)) |\nabla A(u)|^2,$$

and g satisfies $g(A(z)) = f(z)$ for all $z \in \mathbb{R}$.

Proof. By Lemma 4.1 and (4.1),

$$\begin{aligned} \partial_t \int_{\mathbb{R}} S'(\zeta) B'(\zeta) \chi(A(u); \zeta) d\zeta + \nabla \cdot \int_{\mathbb{R}} S'(\zeta) g'(\zeta) \chi(A(u); \zeta) d\zeta \\ - \Delta \int_{\mathbb{R}} S'(\zeta) \chi(A(u); \zeta) d\zeta = \int_{\mathbb{R}} S'(\zeta) \partial_\zeta m(t, x, \zeta) d\zeta. \end{aligned}$$

□

4.2. Kinetic formulation of discrete equations. Stability/uniqueness analysis for differential equations often revolve around the chain rule. The chain rule breaks down for numerical methods, but for us the next lemma will act as a substitute.

Lemma 4.3. Let $S \in C^2(\mathbb{R})$ satisfy $S'(0) = 0$. For any $g \in C^1(\mathbb{R})$ and any real numbers a, b and c ,

$$\begin{aligned} S'(a)(g(b) - g(a)) = \int_0^b S'(z) g'(z) dz - \int_0^a S'(z) g'(z) dz \\ + \int_a^b S''(z)(g(z) - g(b)) dz. \end{aligned}$$

Proof. For any $\zeta \in \mathbb{R}$, integration by parts yields

$$S'(\zeta)(g(\zeta) - g(b)) = \int_0^\zeta S'(z) g'(z) dz + \int_0^\zeta S''(z)(g(z) - g(b)) dz.$$

Take the two equations obtained by setting ζ be equal to a and b and subtract one from the other. □

To make the discrete and continuous calculus notations similar, we introduce the discrete gradient

$$D_\pm \sigma = (D_\pm^1 \sigma, \dots, D_\pm^d \sigma), \quad \text{for any } \sigma : \mathbb{Z}^d \rightarrow \mathbb{R}.$$

The upcoming lemma contains the equation satisfied by $\chi(u_\alpha; \zeta)$, where u_α is the solution of the scheme (3.1).

Lemma 4.4. Suppose $A' \geq 0$. Let $\{u_\alpha\}_{\alpha \in \mathbb{Z}^d}$ be the solution to (3.1). Then $\rho_\alpha(t, \xi) := \chi(u_\alpha(t); \xi)$ satisfies

$$\begin{aligned} \partial_t \rho + (F'_1(\xi) \cdot D_- + F'_2(\xi) \cdot D_+) \rho - A'(\xi) D_- \cdot D_+ \rho = \partial_\xi (m_F + m_A), \\ \zeta_\alpha(0, \xi) = \chi(u_{\alpha,0}; \xi), \end{aligned}$$

in $\mathcal{D}'(\mathbb{R} \times [0, T])$ for each $\alpha \in \mathbb{Z}^d$, where

$$\begin{aligned} m_F = \sum_{i=1}^d \left((F_1^i(\xi) - F_1^i(u_{\alpha-e_i})) D_-^i \chi(u_\alpha; \xi) \right. \\ \left. + (F_2^i(\xi) - F_2^i(u_{\alpha+e_i})) D_+^i \chi(u_\alpha; \xi) \right) \end{aligned}$$

and

$$m_A = \sum_{i=1}^d \left(\frac{1}{\Delta x} (A(u_{\alpha+e_i}) - A(\xi)) D_+^i \chi(u_\alpha; \xi) + \frac{1}{\Delta x} (A(\xi) - A(u_{\alpha-e_i})) D_-^i \chi(u_\alpha; \xi) \right).$$

Proof. Since $\{u_\alpha\}$ is a solution of (3.1),

$$(4.3) \quad S'(u_\alpha(t)) \partial_t u_\alpha(t) + \sum_{i=1}^d S'(u_\alpha(t)) D_-^i F^i(u_\alpha(t), u_{\alpha+e_i}(t)) = \sum_{i=1}^d S'(u_\alpha(t)) D_-^i D_+^i A(u_\alpha(t)),$$

for all $t \in (0, T)$ and $\alpha \in \mathbb{Z}^d$. By the chain rule

$$S'(u_\alpha(t)) \partial_t u_\alpha(t) = \partial_t S(u_\alpha(t)).$$

Consider the flux term. For each i , we have that $F^i = F_2^i + F_1^i$, and therefore

$$S'(u_\alpha(t)) D_-^i F^i(u_\alpha(t), u_{\alpha+e_i}(t)) = S'(u_\alpha(t)) D_-^i F_1^i(u_\alpha) + S'(u_\alpha(t)) D_+^i F_2^i(u_\alpha).$$

By Lemma 4.3, with g equal to F_1^i and F_2^i , we obtain

$$\begin{aligned} S'(u_\alpha(t)) D_-^i F_1^i(u_\alpha) &= D_-^i Q_1^i(u_\alpha) - \frac{1}{\Delta x} \int_{u_\alpha}^{u_{\alpha-e_i}} S''(z) (F_1^i(z) - F_1^i(u_{\alpha-e_i})) dz, \\ S'(u_\alpha(t)) D_+^i F_2^i(u_\alpha) &= D_+^i Q_2^i(u_\alpha) + \frac{1}{\Delta x} \int_{u_\alpha}^{u_{\alpha+e_i}} S''(z) (F_2^i(z) - F_2^i(u_{\alpha+e_i})) dz, \end{aligned}$$

where

$$Q_j^i(u) := \int_0^u S'(z) (F_j^i)'(z) dz \quad \text{for } j = 1, 2.$$

Consider the term on the right-hand side of (4.3). Let

$$R(u) := \int_0^u S'(z) A'(z) dz.$$

Fix i and apply Lemma 4.3 with $g = A$, $a = u_\alpha$, $b = u_{\alpha-e_i}$, and $u_{\alpha+e_i}$. Adding the two equations yields

$$\begin{aligned} S'(u_\alpha) D_-^i D_+^i (A(u_\alpha)) &= D_-^i D_+^i R(u_\alpha) \\ &\quad + \frac{1}{\Delta x^2} \int_{u_\alpha}^{u_{\alpha+e_i}} S''(z) (A(z) - A(u_{\alpha+e_i})) dz \\ &\quad + \frac{1}{\Delta x^2} \int_{u_\alpha}^{u_{\alpha-e_i}} S''(z) (A(z) - A(u_{\alpha-e_i})) dz. \end{aligned}$$

Hence (4.3) turns into

$$\begin{aligned} \partial_t S(u_\alpha) + \sum_{i=1}^d (D_-^i Q_1^i(u_\alpha) + D_+^i Q_2^i(u_\alpha)) &- \sum_{i=1}^d D_-^i D_+^i R(u_\alpha) \\ &= \sum_{i=1}^d \frac{1}{\Delta x} \int_{u_\alpha}^{u_{\alpha-e_i}} S''(z) (F_1^i(z) - F_1^i(u_{\alpha-e_i})) dz \\ &\quad - \sum_{i=1}^d \frac{1}{\Delta x} \int_{u_\alpha}^{u_{\alpha+e_i}} S''(z) (F_2^i(z) - F_2^i(u_{\alpha+e_i})) dz \end{aligned}$$

$$\begin{aligned}
& + \sum_{i=1}^d \frac{1}{\Delta x^2} \int_{u_\alpha}^{u_{\alpha+e_i}} S''(z)(A(z) - A(u_{\alpha+e_i})) dz \\
& + \sum_{i=1}^d \frac{1}{\Delta x^2} \int_{u_\alpha}^{u_{\alpha-e_i}} S''(z)(A(z) - A(u_{\alpha-e_i})) dz.
\end{aligned}$$

By equation (4.1),

$$D_-^i Q_1^i(u_\alpha) + D_+^i Q_2^i(u_\alpha) = \int_{\mathbb{R}} S'(\xi)((F_1^i)'(\xi)D_-^i + (F_2^i)'(\xi)D_+^i)\chi(u_\alpha; \xi) d\xi.$$

Similarly,

$$D_-^i D_+^i R(u_\alpha) = \int_{\mathbb{R}} S'(\xi)A'(\xi)D_-^i D_+^i \chi(u_\alpha; \xi) d\xi.$$

Consider the right-hand side. For any $g \in C(\mathbb{R})$,

$$\int_a^b S''(z)(g(z) - g(b)) dz = \int_{\mathbb{R}} S''(\xi)(g(\xi) - g(b)) (\chi(b; \xi) - \chi(a; \xi)) d\xi.$$

Hence

$$\begin{aligned}
\frac{1}{\Delta x} \int_{u_\alpha}^{u_{\alpha-e_i}} (F_1^i(z) - F_1^i(u_{\alpha-e_i})) dz &= - \int_{\mathbb{R}} S''(\xi)(F_1^i(\xi) - F_1^i(u_{\alpha-e_i})) D_-^i \chi(u_\alpha; \xi) d\xi, \\
\frac{-1}{\Delta x} \int_{u_\alpha}^{u_{\alpha+e_i}} (F_2^i(z) - F_2^i(u_{\alpha+e_i})) dz &= - \int_{\mathbb{R}} S''(\xi)(F_2^i(\xi) - F_2^i(u_{\alpha+e_i})) D_+^i \chi(u_\alpha; \xi) d\xi.
\end{aligned}$$

Similarly,

$$\begin{aligned}
& \frac{1}{\Delta x^2} \int_{u_\alpha}^{u_{\alpha+e_i}} S''(z)(A(z) - A(u_{\alpha+e_i})) dz \\
&= \frac{-1}{\Delta x} \int_{\mathbb{R}} S''(\xi)(A(u_{\alpha+e_i}) - A(\xi)) D_+^i \chi(u_\alpha; \xi) d\xi, \\
& \frac{1}{\Delta x^2} \int_{u_\alpha}^{u_{\alpha-e_i}} S''(z)(A(z) - A(u_{\alpha-e_i})) dz \\
&= \frac{-1}{\Delta x} \int_{\mathbb{R}} S''(\xi)(A(\xi) - A(u_{\alpha-e_i})) D_-^i \chi(u_\alpha; \xi) d\xi.
\end{aligned}$$

The result follows. $\square$

For a function $u : \mathbb{R}^d \rightarrow \mathbb{R}$ we define the shift operator S_y by $S_y u(x) = u(x + y)$. Then the discrete derivatives may be expressed as

$$D_\pm^i u = \pm \frac{S_{\pm \Delta x_i} u - u}{\Delta x},$$

where $\Delta x_i = \Delta x e_i$.

Making a change of variable $\zeta = A(\xi)$, we can obtain an equation satisfied by $\chi(A(u_{\Delta x}); \zeta)$, where $u_{\Delta x}$ is the numerical solution (3.2), resulting in the “discrete” kinetic formulation to be utilized later.

Lemma 4.5. *Suppose $A' > 0$. Let $\{u_\alpha\}$ be the solution to (3.1) and define $u_{\Delta x}$ by (3.2). Let $G_j : \mathbb{R} \rightarrow \mathbb{R}^d$ satisfy $G_j(A(u)) = F_j(u) \forall u$, for $j = 1, 2$. Then $\rho^{\Delta x}(t, x, \zeta) = \chi(A(u_{\Delta x}(t, x)); \zeta)$ satisfies*

$$\begin{aligned}
& B'(\zeta) \partial_t \rho^{\Delta x} + (G_1'(\zeta) \cdot D_+ + G_2'(\zeta) \cdot D_-) \rho^{\Delta x} - D_- \cdot D_+ \rho^{\Delta x} = \partial_\zeta (n_A^{\Delta x} + n_G^{\Delta x}), \\
& \rho^{\Delta x}(0, \zeta) = \chi(A(u_{\Delta x}^0); \zeta),
\end{aligned}$$

in $\mathcal{D}'(\mathbb{R} \times \Pi_T)$, where

$$n_G^{\Delta x} = \sum_{i=1}^d (G_1^i(\zeta) - G_1^i(A(S_{-\Delta x_i} u_{\Delta x}))) D_-^i \chi(A(u_{\Delta x}); \zeta) \\ + \sum_{i=1}^d (G_2^i(\zeta) - G_2^i(A(S_{\Delta x_i} u_{\Delta x}))) D_+^i \chi(A(u_{\Delta x}); \zeta)$$

and

$$(4.4) \quad n_A^{\Delta x} = \sum_{i=1}^d \frac{1}{\Delta x} (A(S_{\Delta x_i} u_{\Delta x}) - \zeta) D_+^i \chi(A(u_{\Delta x}); \zeta) \\ + \sum_{i=1}^d \frac{1}{\Delta x} (\zeta - A(S_{-\Delta x_i} u_{\Delta x})) D_-^i \chi(A(u_{\Delta x}); \zeta).$$

Proof. Let $S \in C_c^\infty(\mathbb{R})$ and define $S_A(\xi) = S(A(\xi))$. By Lemma 4.4,

$$\partial_t \int_{\mathbb{R}} S_A(\xi) \chi(u_\alpha; \xi) d\xi + \int_{\mathbb{R}} S_A(\xi) (F_1'(\xi) \cdot D_- + F_2'(\xi) \cdot D_+) \chi(u_\alpha; \xi) d\xi \\ - \int_{\mathbb{R}} S_A(\xi) A'(\xi) D_- \cdot D_+ \chi(u_\alpha; \xi) d\xi = - \int_{\mathbb{R}} S_A'(\xi) (m_F + m_A) d\xi.$$

Let $\zeta = A(\xi)$ and note that $\chi(u_\alpha; \xi) = \chi(A(u_\alpha); A(\xi))$. The terms on the left-hand side are straightforward to verify. Next,

$$\int_{\mathbb{R}} S_A'(\xi) m_F(\xi) d\xi \\ = \sum_{i=1}^d \int_{\mathbb{R}} S'(A(\xi)) (G_1^i(A(\xi)) - G_1^i(A(u_{\alpha-e_i}))) D_-^i \chi(A(u_\alpha); A(\xi)) A'(\xi) d\xi \\ + \sum_{i=1}^d \int_{\mathbb{R}} S'(A(\xi)) (G_2^i(A(\xi)) - G_2^i(A(u_{\alpha+e_i}))) D_+^i \chi(A(u_\alpha); A(\xi)) A'(\xi) d\xi \\ = \sum_{i=1}^d \int_{\mathbb{R}} S'(\zeta) (G_1^i(\zeta) - G_1^i(A(u_{\alpha-e_i}))) D_-^i \chi(A(u_\alpha); \zeta) d\zeta \\ + \sum_{i=1}^d \int_{\mathbb{R}} S'(\zeta) (G_2^i(\zeta) - G_2^i(A(u_{\alpha+e_i}))) D_+^i \chi(A(u_\alpha); \zeta) d\zeta.$$

A similar computation shows the second equality involving $n_A^{\Delta x}$. $\square$

4.3. Various regularizations. In this section we study mollified versions of Lemma 4.2 and 4.5. Let us first introduce some notation. Let $J \in C_c^\infty(\mathbb{R})$ denote a function satisfying

$$\text{supp}(J) \subset [-1, 1], \quad \int_{\mathbb{R}} J(x) dx = 1 \text{ and } J(-x) = J(x)$$

for all $x \in \mathbb{R}$. That is, J is a symmetric mollifier on $\mathbb{R}$ with support in $[-1, 1]$. For any $\sigma > 0$ we let $J_\sigma(x) = \sigma^{-1} J(\sigma^{-1}x)$. For any $n \geq 1$, $J_\sigma^{\otimes n}$ is a symmetric mollifier on $\mathbb{R}^n$ with support in $[-\sigma, \sigma]^n$. In general the dimension of the argument will define n , so to simplify the notation we write J_σ instead of $J_\sigma^{\otimes n}$.

Let $\psi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a continuous function and $u, v \in L^1(\mathbb{R})$. Then we define

$$(\psi(u, v) \stackrel{(u,v)}{\star} f \otimes g)(x) := \int_{\mathbb{R}} \int_{\mathbb{R}} \psi(u(y_1), v(y_2)) f(x - y_1) g(x - y_2) dy_1 dy_2,$$

where $f, g \in L^1(\mathbb{R})$. Similarly, we let

$$\psi(u, v) \stackrel{(u)}{\star} f := \int_{\mathbb{R}} \psi(u(y), v(x)) f(x - y) dy.$$

This notation generalizes in an obvious way to functions of several variables.

We start by introducing regularizations of $\text{sign}(\cdot)$ and $\chi(u; \cdot)$

Lemma 4.6. *For $\varepsilon > 0$, define*

$$\text{sign}_\varepsilon(\xi) := 2 \int_0^\xi J_\varepsilon(\zeta) d\zeta, \quad \chi_\varepsilon(u; \xi) := \int_{\mathbb{R}} \chi(u; \zeta) J_\varepsilon(\xi - \zeta) d\zeta.$$

Then

- (i) *For each ξ , $u \mapsto \chi_\varepsilon(u; \xi) \in C^\infty(\mathbb{R})$ and $\partial_u \chi_\varepsilon(u; \xi) = J_\varepsilon(\xi - u)$.*
- (ii) *For all u and ξ*

$$\text{sign}_\varepsilon(\xi) - 2\chi_\varepsilon(u; \xi) = \text{sign}_\varepsilon(\xi - u).$$

- (iii) *For any u*

$$\int_{\mathbb{R}} |\chi_\varepsilon(u; \xi) - \chi(u; \xi)| d\xi \leq 4\varepsilon.$$

Proof. We first prove (i). Let $H'_\varepsilon(\sigma) = J_\varepsilon(\sigma)$. Since $J_\varepsilon(\xi - \zeta) = J_\varepsilon(\zeta - \xi)$,

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{1}{h} \int_{\mathbb{R}} (\chi(u + h; \zeta) - \chi(u; \zeta)) J_\varepsilon(\xi - \zeta) d\zeta \\ = \lim_{h \rightarrow 0} \frac{1}{h} (H_\varepsilon(u + h - \xi) - H_\varepsilon(u - \xi)) = J_\varepsilon(u - \xi). \end{aligned}$$

Next we prove (ii). Let $\sigma = \zeta - \xi$. By the symmetry of J_ε ,

$$\chi_\varepsilon(u; \xi) = \int_{\mathbb{R}} \chi(u; \sigma + \xi) J_\varepsilon(\sigma) d\sigma.$$

A calculation (or (5.24)) yields

$$\chi(u; \sigma + \xi) = \chi(u - \xi; \sigma) - \chi(-\xi; \sigma).$$

Note that $\chi(-\xi; \sigma) = -\chi(\xi; -\sigma)$. Hence

$$\chi_\varepsilon(u; \xi) = \int_{\mathbb{R}} (\chi(u - \xi; \zeta) + \chi(\xi; -\zeta)) J_\varepsilon(\zeta) d\zeta.$$

It follows that

$$\begin{aligned} \text{sign}_\varepsilon(\xi) - 2\chi_\varepsilon(u; \xi) &= -2 \int_{\mathbb{R}} \chi(u - \xi; \zeta) J_\varepsilon(\zeta) d\zeta + 2 \int_{\mathbb{R}} (\chi(\xi; \zeta) - \chi(\xi; -\zeta)) J_\varepsilon(\zeta) d\zeta \\ &=: \mathcal{T}_1 + \mathcal{T}_2. \end{aligned}$$

Since $(\chi(\xi; \zeta) - \chi(\xi; -\zeta))$ is antisymmetric in ζ and J_ε is symmetric it follows that $\mathcal{T}_2 = 0$. Now

$$\mathcal{T}_1 = -2 \int_0^{u-\xi} J_\varepsilon(\zeta) d\zeta = 2 \int_0^{\xi-u} J_\varepsilon(\zeta) d\zeta = \text{sign}_\varepsilon(\xi - u).$$

To prove (iii), note that

$$|\chi_\varepsilon(u; \xi) - \chi(u; \xi)| = 0 \text{ whenever } \xi \notin (-\varepsilon, \varepsilon) \cup (u - \varepsilon, u + \varepsilon).$$

□

For $\varepsilon > 0$ and $f \in C(\mathbb{R})$, let $R_\varepsilon^f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$(4.5) \quad \int_{\mathbb{R}} f(\sigma) \chi(u; \sigma) J_\varepsilon(\zeta - \sigma) d\sigma = R_\varepsilon^f(u, \zeta) + f(\zeta) \chi_\varepsilon(u; \zeta),$$

for all $u, \zeta \in \mathbb{R}$.

Now we are ready to provide “regularized” versions of Lemmas 4.2 and 4.5. As the mollification will take place on a slightly smaller region, we introduce the notation $\Pi_T^{r_0} := (r_0, T - r_0) \times \mathbb{R}^d$.

We start with the regularization of the kinetic formulation of the convection-diffusion equation.

Lemma 4.7. *Assume that $A' > 0$. Let u be the solution of (1.1) and define*

$$\rho_{\varepsilon, r, r_0} := \chi(A(u); \cdot) \star J_{r_0} \otimes J_r \otimes J_\varepsilon.$$

Then for $(t, x, \zeta) \in \Pi_T^{r_0} \times \mathbb{R}$, the function $\rho_{\varepsilon, r, r_0}$ satisfies

$$B'(\zeta) \partial_t \rho_{\varepsilon, r, r_0} + g'(\zeta) \cdot \nabla \rho_{\varepsilon, r, r_0} - \Delta \rho_{\varepsilon, r, r_0} + \partial_t R_{\varepsilon, r, r_0}^{B'} + \nabla \cdot R_{\varepsilon, r, r_0}^{g'} = \partial_\zeta n_{A, \varepsilon, r, r_0},$$

where

$$R_{\varepsilon, r, r_0}^f = R_\varepsilon^f(A(u), \zeta) \star J_r \otimes J_{r_0},$$

with R_ε^f defined by (4.5), and

$$n_{A, \varepsilon, r, r_0}(t, x, \zeta) = \left(J_\varepsilon(\zeta - A(u)) |\nabla A(u)|^2 \star J_{r_0} \otimes J_r \right)(t, x).$$

Proof. Starting off from Lemma 4.2, take the convolution of equation (4.2) with J_ε and apply (4.5). Finally, convolve the resulting equation with $J_r \otimes J_{r_0}$. $\square$

Next up is the regularization of the kinetic formulation of the discrete equations.

Lemma 4.8. *Under the same assumptions and with the same notation as in Lemma 4.5, define*

$$\rho_{\varepsilon, r, r_0}^{\Delta x} := \chi(A(u_{\Delta x}); \cdot) \star J_{r_0} \otimes J_r \otimes J_\varepsilon.$$

For $(t, x, \zeta) \in \Pi_T^{r_0} \times \mathbb{R}$, the function $\rho_{\varepsilon, r, r_0}^{\Delta x}$ satisfies

$$\begin{aligned} B'(\zeta) \partial_t \rho_{\varepsilon, r, r_0}^{\Delta x} + g'(\zeta) \cdot \nabla \rho_{\varepsilon, r, r_0}^{\Delta x} - \Delta \rho_{\varepsilon, r, r_0}^{\Delta x} + G_1'(\zeta) \cdot (D_+ - \nabla) \rho_{\varepsilon, r, r_0}^{\Delta x} \\ + G_2'(\zeta) \cdot (D_- - \nabla) \rho_{\varepsilon, r, r_0}^{\Delta x} + (\Delta - D_- \cdot D_+) \rho_{\varepsilon, r, r_0}^{\Delta x} + \partial_t R_{\varepsilon, r, r_0}^{B', \Delta x} \\ + D_+ \cdot R_{\varepsilon, r, r_0}^{G_1', \Delta x} + D_- \cdot R_{\varepsilon, r, r_0}^{G_2', \Delta x} = \partial_\zeta (n_{A, \varepsilon, r, r_0}^{\Delta x} + n_{G, \varepsilon, r, r_0}^{\Delta x}). \end{aligned}$$

Here, $R_{\varepsilon, r, r_0}^{f, \Delta x} = R_\varepsilon^f(A(u_{\Delta x}), \cdot) \star J_r \otimes J_{r_0}$ with R_ε^f coming from (4.5). Furthermore,

$$n_{A, \varepsilon, r, r_0}^{\Delta x} = n_A^{\Delta x} \star (J_\varepsilon \otimes J_r \otimes J_{r_0}) \text{ and } n_{G, \varepsilon, r, r_0}^{\Delta x} = n_G^{\Delta x} \star (J_\varepsilon \otimes J_r \otimes J_{r_0}).$$

Proof. In view of Lemma 4.5 and (4.5),

$$\begin{aligned} B'(\zeta) \partial_t \rho_\varepsilon^{\Delta x} + (G_1'(\zeta) \cdot D_+ + G_2'(\zeta) \cdot D_-) \rho_\varepsilon^{\Delta x} - D_- \cdot D_+ \rho_\varepsilon^{\Delta x} \\ + \partial_t R_\varepsilon^{B'}(u_{\Delta x}, \zeta) + D_+ \cdot R_\varepsilon^{G_1'}(u_{\Delta x}, \zeta) + D_- \cdot R_\varepsilon^{G_2'}(u_{\Delta x}, \zeta) = \partial_\zeta (n_{A, \varepsilon}^{\Delta x} + n_{G, \varepsilon}^{\Delta x}), \end{aligned}$$

where $\rho_\varepsilon^{\Delta x}(t, x, \zeta) = \chi_\varepsilon(A(u_{\Delta x}); \zeta)$ and $n_{A, \varepsilon}^{\Delta x} = n_A^{\Delta x} \star J_\varepsilon$ and $n_{G, \varepsilon}^{\Delta x} = n_G^{\Delta x} \star J_\varepsilon$. Take the convolution of the above equation with $J_r \otimes J_{r_0}$. Recall that $G_1' + G_2' = g'$ and add and subtract to obtain the result. $\square$

5. PROOF OF THEOREM 3.1

We are now ready to embark on the proof of the error estimate (Theorem 3.1). Instead of working directly with the microscopic contraction functional (1.8), we introduce a regularized version Q_ε of it. For $u, v, \xi \in \mathbb{R}$, define

$$(5.1) \quad Q_\varepsilon(u, v; \xi) := \text{sign}_\varepsilon(\xi) \chi_\varepsilon(u; \xi) + \text{sign}_\varepsilon(\xi) \chi_\varepsilon(v; \xi) - 2\chi_\varepsilon(u; \xi)\chi_\varepsilon(v; \xi),$$

where sign_ε and χ_ε are given in Lemma 4.6. One may show that

$$\int_{\mathbb{R}} (\chi_\varepsilon(u; \xi) - \chi_\varepsilon(v; \xi))^2 d\xi = \int_{\mathbb{R}} Q_\varepsilon(u, v; \xi) d\xi.$$

This equality is, however, not directly useful to us, since we will be working with functions like $\chi_\varepsilon(A(u); \xi)$ with $A(\cdot)$ nonlinear, but see the related Lemma 5.8.

5.1. Main error equation. We will use the kinetic formulations of the convection-diffusion equation and the difference method to derive a fundamental equation for the error quantity $Q_\varepsilon(A(u(t, x)), A(u_{\Delta x}(t, x)); \zeta)$ (properly regularized).

Lemma 5.1. *Assume that $A' > 0$. With the notation of Lemma 4.7 and 4.8, define*

$$Q_{\varepsilon, r, r_0}(\zeta) = Q_\varepsilon(A(u), A(u_{\Delta x}); \zeta) \stackrel{(u, u_{\Delta x})}{\star} (J_{r_0} \otimes J_r) \otimes (J_{r_0} \otimes J_r).$$

Then, for all $(t, x) \in \Pi_T^{r_0}$,

$$(5.2) \quad \int_{\mathbb{R}} B'(\zeta) \partial_t Q_{\varepsilon, r, r_0} d\zeta + \int_{\mathbb{R}} g'(\zeta) \cdot \nabla Q_{\varepsilon, r, r_0} d\zeta$$

$$(5.3) \quad = \int_{\mathbb{R}} \Delta Q_{\varepsilon, r, r_0} d\zeta + 2 \int_{\mathbb{R}} \nabla \rho_{\varepsilon, r, r_0} \cdot (2\nabla - (D_+ + D_-)) \rho_{\varepsilon, r, r_0}^{\Delta x} d\zeta$$

$$(5.4) \quad - \int_{\mathbb{R}} (\text{sign}_\varepsilon(\zeta - A(u_{\Delta x})) \star J_{r_0} \otimes J_r) \partial_t R_{\varepsilon, r, r_0}^{B'} d\zeta$$

$$(5.5) \quad - \int_{\mathbb{R}} (\text{sign}_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) \partial_t R_{\varepsilon, r, r_0}^{B', \Delta x} d\zeta$$

$$(5.6) \quad - \int_{\mathbb{R}} (\text{sign}_\varepsilon(\zeta - A(u_{\Delta x})) \star J_{r_0} \otimes J_r) \nabla \cdot R_{\varepsilon, r, r_0}^{g'} d\zeta$$

$$(5.7) \quad - \int_{\mathbb{R}} (\text{sign}_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) \left(D_+ \cdot R_{\varepsilon, r, r_0}^{G'_1, \Delta x} + D_- \cdot R_{\varepsilon, r, r_0}^{G'_2, \Delta x} \right) d\zeta$$

$$(5.8) \quad - \int_{\mathbb{R}} (\text{sign}_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) (G'_1(\zeta) \cdot (D_+ - \nabla)$$

$$(5.9) \quad + G'_2(\zeta) \cdot (D_- - \nabla)) \rho_{\varepsilon, r, r_0}^{\Delta x} d\zeta.$$

$$(5.10) \quad + \int_{\mathbb{R}} (\text{sign}_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) (\Delta - D_- \cdot D_+) \rho_{\varepsilon, r, r_0}^{\Delta x} d\zeta$$

$$(5.11) \quad - 2 \int_{\mathbb{R}} (J_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) n_{G, \varepsilon, r}^{\Delta x} d\zeta$$

$$(5.12) \quad - 2 \int_{\mathbb{R}} E_{\Delta x, \varepsilon, r, r_0}(\zeta) d\zeta,$$

where

$$(5.12) \quad \begin{aligned} E_{\Delta x, \varepsilon, r, r_0}(\zeta) = & -\nabla \rho_{\varepsilon, r, r_0} \cdot (D_+ + D_-) \rho_{\varepsilon, r, r_0}^{\Delta x} \\ & + (J_\varepsilon(\zeta - A(u_{\Delta x})) \star J_{r_0} \otimes J_r) n_{A, \varepsilon, r, r_0} \\ & + (J_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) n_{A, \varepsilon, r, r_0}^{\Delta x}. \end{aligned}$$

Proof. By definition,

$$Q_{\varepsilon, r, r_0}(t, x, \zeta) = \text{sign}_\varepsilon(\zeta) \rho_{\varepsilon, r, r_0}(t, x, \zeta) + \text{sign}_\varepsilon(\zeta) \rho_{\varepsilon, r, r_0}^{\Delta x}(t, x, \zeta)$$

$$- 2\rho_{\varepsilon,r,r_0}(t,x,\zeta)\rho_{\varepsilon,r,r_0}^{\Delta x}(t,x,\zeta).$$

Hence,

$$\begin{aligned}\partial_t \int_{\mathbb{R}} Q_{\varepsilon,r,r_0} B'(\zeta) d\zeta &= \int_{\mathbb{R}} \text{sign}_{\varepsilon}(\zeta) \partial_t(\rho_{\varepsilon,r,r_0} + \rho_{\varepsilon,r,r_0}^{\Delta x}) B'(\zeta) d\zeta \\ &\quad + \int_{\mathbb{R}} \partial_t(\rho_{\varepsilon,r,r_0} \rho_{\varepsilon,r,r_0}^{\Delta x}) B'(\zeta) d\zeta \\ &=: \mathcal{T}_1 + \mathcal{T}_2.\end{aligned}$$

By Lemmas 4.7 and 4.8

$$\begin{aligned}\mathcal{T}_1 &= - \underbrace{\int_{\mathbb{R}} \text{sign}_{\varepsilon}(\zeta) g'(\zeta) \cdot \nabla(\rho_{\varepsilon,r,r_0} + \rho_{\varepsilon,r,r_0}^{\Delta x}) d\zeta}_{\mathcal{T}_1^1} \\ &\quad + \underbrace{\int_{\mathbb{R}} \text{sign}_{\varepsilon}(\zeta) \Delta(\rho_{\varepsilon,r,r_0} + \rho_{\varepsilon,r,r_0}^{\Delta x}) d\zeta}_{\mathcal{T}_1^2} \\ &\quad - \underbrace{\int_{\mathbb{R}} \text{sign}_{\varepsilon}(\zeta) \partial_t(R_{\varepsilon,r,r_0}^{B'} + R_{\varepsilon,r,r_0}^{B',\Delta x}) d\zeta}_{\mathcal{T}_1^3} \\ &\quad - \underbrace{\int_{\mathbb{R}} \text{sign}_{\varepsilon}(\zeta) \left(\nabla \cdot R_{\varepsilon,r,r_0}^{g'} + D_+ \cdot R_{\varepsilon,r,r_0}^{G'_1,\Delta x} + D_- \cdot R_{\varepsilon,r,r_0}^{G'_2,\Delta x} \right) d\zeta}_{\mathcal{T}_1^4} \\ &\quad + \underbrace{\int_{\mathbb{R}} \text{sign}_{\varepsilon}(\zeta) (\partial_{\zeta} n_{A,\varepsilon,r,r_0} + \partial_{\zeta} n_{A,\varepsilon,r,r_0}^{\Delta x}) d\zeta}_{\mathcal{T}_1^5} \\ &\quad - \underbrace{\int_{\mathbb{R}} \text{sign}_{\varepsilon}(\zeta) (G'_1(\zeta) \cdot (D_+ - \nabla) + G'_2(\zeta) \cdot (D_- - \nabla)) \rho_{\varepsilon,r,r_0}^{\Delta x} d\zeta}_{\mathcal{T}_1^6} \\ &\quad + \underbrace{\int_{\mathbb{R}} \text{sign}_{\varepsilon}(\zeta) (\Delta - D_- \cdot D_+) \rho_{\varepsilon,r,r_0}^{\Delta x} d\zeta}_{\mathcal{T}_1^7} \\ &\quad + \underbrace{\int_{\mathbb{R}} \text{sign}_{\varepsilon}(\zeta) \partial_{\zeta} n_{G,\varepsilon,r,r_0}^{\Delta x} d\zeta}_{\mathcal{T}_1^8}.\end{aligned}$$

Similarly for $\mathcal{T}_2$ we obtain

$$\begin{aligned}\mathcal{T}_2 &= 2 \underbrace{\int_{\mathbb{R}} g'(\zeta) \cdot \nabla(\rho_{\varepsilon,r,r_0} \rho_{\varepsilon,r,r_0}^{\Delta x}) d\zeta}_{\mathcal{T}_2^1} \\ &\quad - 2 \underbrace{\int_{\mathbb{R}} \rho_{\varepsilon,r,r_0} \Delta \rho_{\varepsilon,r,r_0}^{\Delta x} + \Delta \rho_{\varepsilon,r,r_0} \rho_{\varepsilon,r,r_0}^{\Delta x} d\zeta}_{\mathcal{T}_2^2} \\ &\quad + 2 \underbrace{\int_{\mathbb{R}} \rho_{\varepsilon,r,r_0} \partial_t R_{\varepsilon,r,r_0}^{B',\Delta x} + \rho_{\varepsilon,r,r_0}^{\Delta x} \partial_t R_{\varepsilon,r,r_0}^{B'} d\zeta}_{\mathcal{T}_2^3}\end{aligned}$$

$$\begin{aligned}
& + 2 \underbrace{\int_{\mathbb{R}} \rho_{\varepsilon,r,r_0}^{\Delta x} \nabla \cdot R_{\varepsilon,r,r_0}^{g'} + \rho_{\varepsilon,r,r_0} D_+ \cdot R_{\varepsilon,r,r_0}^{G'_1, \Delta x} + \rho_{\varepsilon,r,r_0} D_- \cdot R_{\varepsilon,r,r_0}^{G'_2, \Delta x} d\zeta}_{\mathcal{T}_2^4} \\
& - 2 \underbrace{\int_{\mathbb{R}} \rho_{\varepsilon,r,r_0} \partial_{\zeta} n_{A,\varepsilon,r,r_0}^{\Delta x} + \rho_{\varepsilon,r,r_0}^{\Delta x} \partial_{\zeta} n_{A,\varepsilon,r,r_0} d\zeta}_{\mathcal{T}_2^5} \\
& + 2 \underbrace{\int_{\mathbb{R}} \rho_{\varepsilon,r,r_0} (G'_1(\zeta) \cdot (D_+ - \nabla) + G'_2(\zeta) \cdot (D_- - \nabla)) \rho_{\varepsilon,r,r_0}^{\Delta x} d\zeta}_{\mathcal{T}_2^6} \\
& - 2 \underbrace{\int_{\mathbb{R}} \rho_{\varepsilon,r,r_0} (\Delta - D_- \cdot D_+) \rho_{\varepsilon,r,r_0}^{\Delta x} d\zeta}_{\mathcal{T}_2^7} \\
& - 2 \underbrace{\int_{\mathbb{R}} \rho_{\varepsilon,r,r_0} \partial_{\zeta} n_{G,\varepsilon,r,r_0}^{\Delta x} d\zeta}_{\mathcal{T}_2^8}.
\end{aligned}$$

We compute $\mathcal{T}_1 + \mathcal{T}_2$ term by term, and thereby explain each of the terms (5.3)–(5.11) in the lemma. We start with

$$\mathcal{T}_1^1 + \mathcal{T}_2^1 = - \int_{\mathbb{R}} g'(\zeta) \cdot \nabla Q_{\varepsilon,r,r_0} d\zeta,$$

which gives the last term in (5.2).

To make the second derivative terms a complete derivative we need to add and subtract. Hence we may write

$$\begin{aligned}
\mathcal{T}_1^2 + \mathcal{T}_2^2 &= \Delta \int_{\mathbb{R}} Q_{\varepsilon,r,r_0} d\zeta + 4 \int_{\mathbb{R}} \nabla \rho_{\varepsilon,r,r_0} \cdot \nabla \rho_{\varepsilon,r,r_0}^{\Delta x} d\zeta \\
&= \Delta \int_{\mathbb{R}} Q_{\varepsilon,r,r_0} d\zeta + 2 \int_{\mathbb{R}} \nabla \rho_{\varepsilon,r,r_0} \cdot (D_+ + D_-) \rho_{\varepsilon,r,r_0}^{\Delta x} d\zeta \\
&\quad + 2 \int_{\mathbb{R}} \nabla \rho_{\varepsilon,r,r_0} (2\nabla - (D_+ + D_-)) \rho_{\varepsilon,r,r_0}^{\Delta x} d\zeta,
\end{aligned}$$

which explains (5.3) and the first term in $E_{\Delta x, \varepsilon, r, r_0}$.

By Lemma 4.6 it follows that

$$\begin{aligned}
\text{sign}_{\varepsilon}(\zeta) - 2\rho_{\varepsilon,r,r_0} &= \text{sign}_{\varepsilon}(\zeta - A(u)) \star J_{r_0} \otimes J_r, \\
\text{sign}_{\varepsilon}(\zeta) - 2\rho_{\varepsilon,r,r_0}^{\Delta x} &= \text{sign}_{\varepsilon}(\zeta - A(u_{\Delta x})) \star J_{r_0} \otimes J_r.
\end{aligned}$$

Hence,

$$\begin{aligned}
\mathcal{T}_1^3 + \mathcal{T}_2^3 &= - \int_{\mathbb{R}} (\text{sign}_{\varepsilon}(\zeta - A(u_{\Delta x})) \star J_{r_0} \otimes J_r) \partial_t R_{\varepsilon,r,r_0}^{B'} d\zeta \\
&\quad - \int_{\mathbb{R}} (\text{sign}_{\varepsilon}(\zeta - A(u)) \star J_{r_0} \otimes J_r) \partial_t R_{\varepsilon,r,r_0}^{B', \Delta x} d\zeta,
\end{aligned}$$

which explains (5.4) and (5.5). Similarly,

$$\begin{aligned}
\mathcal{T}_1^4 + \mathcal{T}_2^4 &= - \int_{\mathbb{R}} (\text{sign}_{\varepsilon}(\zeta - A(u_{\Delta x})) \star J_{r_0} \otimes J_r) \nabla \cdot R_{\varepsilon,r,r_0}^{g'} d\zeta \\
&\quad - \int_{\mathbb{R}} (\text{sign}_{\varepsilon}(\zeta - A(u)) \star J_{r_0} \otimes J_r) \left(D_+ \cdot R_{\varepsilon,r,r_0}^{G'_1, \Delta x} + D_- \cdot R_{\varepsilon,r,r_0}^{G'_2, \Delta x} \right) d\zeta,
\end{aligned}$$

which explains the presence of (5.6) and (5.7).

Performing integration by parts we obtain, using Lemma 4.6,

$$\begin{aligned}\mathcal{T}_1^5 + \mathcal{T}_2^5 &= \int_{\mathbb{R}} (\text{sign}_{\varepsilon}(\zeta - A(u_{\Delta x})) \star J_{r_0} \otimes J_r) \partial_{\zeta} n_{A,\varepsilon,r,r_0} \\ &\quad + (\text{sign}_{\varepsilon}(\zeta - A(u)) \star J_{r_0} \otimes J_r) \partial_{\zeta} n_{A,\varepsilon,r,r_0}^{\Delta x} d\zeta \\ &= -2 \int_{\mathbb{R}} (J_{\varepsilon}(\zeta - A(u_{\Delta x})) \star J_{r_0} \otimes J_r) n_{A,\varepsilon,r,r_0} \\ &\quad + (J_{\varepsilon}(\zeta - A(u)) \star J_{r_0} \otimes J_r) n_{A,\varepsilon,r,r_0}^{\Delta x} d\zeta,\end{aligned}$$

which explains the two last terms in $E_{\Delta x,\varepsilon,r,r_0}$.

Similarly,

$$\begin{aligned}\mathcal{T}_1^6 + \mathcal{T}_2^6 &= - \int_{\mathbb{R}} (\text{sign}_{\varepsilon}(\zeta - A(u)) \star J_{r_0} \otimes J_r) (G_1'(\zeta) \cdot (D_+ - \nabla) \\ &\quad + G_2'(\zeta) \cdot (D_- - \nabla)) \rho_{\varepsilon,r,r_0}^{\Delta x} d\zeta,\end{aligned}$$

and

$$\mathcal{T}_1^7 + \mathcal{T}_2^7 = \int_{\mathbb{R}} (\text{sign}_{\varepsilon}(\zeta - A(u)) \star J_{r_0} \otimes J_r) (\Delta - D_- \cdot D_+) \rho_{\varepsilon,r,r_0}^{\Delta x} d\zeta,$$

explaining the terms (5.8) and (5.9).

Finally, integration by parts yields

$$\mathcal{T}_1^8 + \mathcal{T}_2^8 = -2 \int_{\mathbb{R}} (J_{\varepsilon}(\zeta - A(u)) \star J_{r_0} \otimes J_r) n_{G,\varepsilon,r,r_0}^{\Delta x} d\zeta,$$

which is the term (5.10). $\square$

5.2. Dissipative term. In this subsection we are concerned with finding an upper bound on (5.12). In the continuous setting, this “dissipative” term is negative, cf. (1.13), which comes as a consequence of the chain rule of calculus. The following elementary lemma will help us contend with the lack of a discrete chain rule.

Lemma 5.2. *Let a and b be two real numbers. Then there exist real numbers $\tau = \tau_{\varepsilon}(a, b, \zeta)$ and $\theta = \theta_{\varepsilon}(a, b, \zeta)$ such that τ and θ are between a and b , and*

$$(5.13) \quad \int_{\mathbb{R}} J_{\varepsilon}(\zeta - \xi) (\chi(b; \xi) - \chi(a; \xi)) d\xi = J_{\varepsilon}(\zeta - \theta)(b - a),$$

$$(5.14) \quad \int_{\mathbb{R}} J_{\varepsilon}(\zeta - \xi) (b - \xi) (\chi(b; \xi) - \chi(a; \xi)) d\xi = \frac{1}{2} J_{\varepsilon}(\zeta - \tau)(b - a)^2.$$

Furthermore, whenever $a \neq b$:

(i)

$$\begin{aligned}J_{\varepsilon}(\zeta - \theta) &= \frac{1}{b - a} \int_a^b J_{\varepsilon}(\zeta - \xi) d\xi; \\ J_{\varepsilon}(\zeta - \tau) &= \frac{2}{(b - a)^2} \int_a^b J_{\varepsilon}(\zeta - \xi) (b - \xi) d\xi;\end{aligned}$$

(ii)

$$(J_{\varepsilon}(\zeta - \theta) - J_{\varepsilon}(\zeta - \tau))(b - a) = \frac{1}{b - a} \int_a^b J_{\varepsilon}(\zeta - \xi) (2\xi - (b + a)) d\xi;$$

(iii)

$$J_{\varepsilon}(\zeta - \tau) - J_{\varepsilon}(\zeta - a) = \frac{2}{(b - a)^2} \left(\int_a^b (J_{\varepsilon}(\zeta - \xi) - J_{\varepsilon}(\zeta - a)) (b - \xi) d\xi \right);$$

(iv)

$$J_\varepsilon(\zeta - \theta_\varepsilon(a, b, \zeta)) - J_\varepsilon(\zeta - \tau_\varepsilon(a, b, \zeta)) = \\ - (J_\varepsilon(\zeta - \theta_\varepsilon(b, a, \zeta)) - J_\varepsilon(\zeta - \tau_\varepsilon(b, a, \zeta))).$$

Proof. To prove (5.14), note that

$$\int_{\mathbb{R}} J_\varepsilon(\zeta - \xi)(b - \xi)(\chi(b; \xi) - \chi(a; \xi)) d\xi = \int_a^b J_\varepsilon(\zeta - \xi)(b - \xi) d\xi.$$

By the mean value theorem there exists a τ between a and b such that

$$\int_a^b J_\varepsilon(\zeta - \xi)(b - \xi) d\xi = J_\varepsilon(\zeta - \tau) \int_a^b (b - \xi) d\xi.$$

Equation (5.13) follows in a similar way. The proof of (i) is immediate. Let us prove (ii). By (i)

$$(J_\varepsilon(\zeta - \theta) - J_\varepsilon(\zeta - \tau))(b - a) = \int_a^b J_\varepsilon(\zeta - \xi) \left(1 - 2\frac{b - \xi}{b - a}\right) d\xi.$$

It remains to observe that

$$1 - 2\frac{b - \xi}{b - a} = \frac{2\xi - (b + a)}{b - a}.$$

To prove (iii), note that

$$J_\varepsilon(\zeta - a)(b - a)^2 = 2 \int_a^b J_\varepsilon(\zeta - a)(b - \xi) d\xi.$$

Hence (iii) follows by (i). To prove (iv), observe that the expression on the right-hand side of (ii) is symmetric in a and b . $\square$

The next result can be viewed as a discrete counterpart of the chain rule, enabling us to write the nonlinear term $n_A^{\Delta x}$, properly regularized, on a form that resembles a parabolic dissipation term like (1.14).

Lemma 5.3. *With the notation of Lemma 4.5, for each $1 \leq i \leq d$, let*

$$\tau_{\Delta x, i}^+ = \tau_\varepsilon(A(u_{\Delta x}), S_{\Delta x_i} A(u_{\Delta x}), \zeta), \quad \tau_{\Delta x, i}^- = \tau_\varepsilon(A(u_{\Delta x}), S_{-\Delta x_i} A(u_{\Delta x}), \zeta)$$

and

$$\theta_{\Delta x, i}^+ = \theta_\varepsilon(A(u_{\Delta x}), S_{\Delta x_i} A(u_{\Delta x}), \zeta), \quad \theta_{\Delta x, i}^- = \theta_\varepsilon(A(u_{\Delta x}), S_{-\Delta x_i} A(u_{\Delta x}), \zeta),$$

where $\tau_\varepsilon, \theta_\varepsilon$ is defined in Lemma 5.2. Then

(i)

$$n_A^{\Delta x} \star J_\varepsilon(t, x, \zeta) = \frac{1}{2} \sum_{i=1}^d J_\varepsilon(\zeta - \tau_{\Delta x, i}^+) (D_+^i A(u_{\Delta x}))^2 \\ + \frac{1}{2} \sum_{i=1}^d J_\varepsilon(\zeta - \tau_{\Delta x, i}^-) (D_-^i A(u_{\Delta x}))^2;$$

(ii) for $1 \leq i \leq d$,

$$D_+^i \chi_\varepsilon(A(u_{\Delta x}); \zeta) = J_\varepsilon(\zeta - \theta_{\Delta x, i}^+) D_+^i(A(u_{\Delta x})), \\ D_-^i \chi_\varepsilon(A(u_{\Delta x}); \zeta) = J_\varepsilon(\zeta - \theta_{\Delta x, i}^-) D_-^i(A(u_{\Delta x})).$$

Proof. By the definition (4.4) of $n_A^{\Delta x}$, recalling that S_y commutes with function evaluation,

$$\begin{aligned} & n_A^{\Delta x} \star J_\varepsilon(t, x, \zeta) \\ &= \sum_{i=1}^d \frac{1}{\Delta x^2} \int_{\mathbb{R}} J_\varepsilon(\zeta - \xi) (S_{\Delta x_i} A(u_{\Delta x}) - \xi) (\chi(S_{\Delta x_i} A(u_{\Delta x}); \xi) - \chi(A(u_{\Delta x}); \xi)) d\xi \\ &+ \sum_{i=1}^d \frac{1}{\Delta x^2} \int_{\mathbb{R}} J_\varepsilon(\zeta - \xi) (S_{-\Delta x_i} A(u_{\Delta x}) - \xi) (\chi(S_{-\Delta x_i} A(u_{\Delta x}); \xi) - \chi(A(u_{\Delta x}); \xi)) d\xi. \end{aligned}$$

Hence (i) follows by Lemma 5.2. To prove (ii) note that by Lemma 5.2,

$$\begin{aligned} & D_+^i \int_{\mathbb{R}} \chi(A(u_{\Delta x}); \xi) J_\varepsilon(\zeta - \xi) d\xi \\ &= \frac{1}{\Delta x} \int_{\mathbb{R}} J_\varepsilon(\zeta - \xi) (\chi(S_{\Delta x_i} A(u_{\Delta x}); \xi) - \chi(A(u_{\Delta x}); \xi)) d\xi \\ &= J_\varepsilon(\zeta - \theta_{\Delta x, i}^+) D_+^i(A(u_{\Delta x})). \end{aligned}$$

The same argument applies to $\theta_{\Delta x, i}^-$. $\square$

We have now come to the key result of this subsection, namely a lower bound on the discrete dissipation term (5.12).

Lemma 5.4. *Let $E_{\Delta x, \varepsilon, r, r_0}$ be defined in Lemma 5.1. Then*

$$E_{\Delta x, \varepsilon, r, r_0} \geq \sum_{k=1}^2 (R_k^+ + R_k^-) \text{ everywhere in } (r_0, T - r_0) \times \mathbb{R}^d \times \mathbb{R},$$

for all positive numbers Δx , ε , r , and r_0 , where

$$\begin{aligned} R_1^+(\zeta) &= \sum_{i=1}^d ((J_\varepsilon(\zeta - \tau_{\Delta x, i}^+) - J_\varepsilon(\zeta - \theta_{\Delta x, i}^+)) D_+^i A(u_{\Delta x}) \star J_{r_0} \otimes J_r) \partial_{x_i} \rho_{\varepsilon, r, r_0}, \\ R_1^-(\zeta) &= \sum_{i=1}^d ((J_\varepsilon(\zeta - \tau_{\Delta x, i}^-) - J_\varepsilon(\zeta - \theta_{\Delta x, i}^-)) D_-^i A(u_{\Delta x}) \star J_{r_0} \otimes J_r) \partial_{x_i} \rho_{\varepsilon, r, r_0}, \\ R_2^+(\zeta) &= \frac{1}{2} \sum_{i=1}^d \left[\left((J_\varepsilon(\zeta - A(u_{\Delta x})) - J_\varepsilon(\zeta - A(\tau_{\Delta x, i}^+))) \star J_{r_0} \otimes J_r \right) \right. \\ &\quad \left. \times (J_\varepsilon(\zeta - A(u)) (\partial_{x_i} A(u))^2 \star J_{r_0} \otimes J_r) \right], \\ R_2^-(\zeta) &= \frac{1}{2} \sum_{i=1}^d \left[\left((J_\varepsilon(\zeta - A(u_{\Delta x})) - J_\varepsilon(\zeta - A(\tau_{\Delta x, i}^-))) \star J_{r_0} \otimes J_r \right) \right. \\ &\quad \left. \times (J_\varepsilon(\zeta - A(u)) (\partial_{x_i} A(u))^2 \star J_{r_0} \otimes J_r) \right]. \end{aligned}$$

Proof. By Lemma 5.3,

$$\begin{aligned} & (J_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) n_{A, \varepsilon, r, r_0}^{\Delta x} \\ &= \frac{1}{2} \sum_{i=1}^d (J_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) (J_\varepsilon(\zeta - \tau_{\Delta x, i}^+) (D_+^i A(u_{\Delta x}))^2 \star J_{r_0} \otimes J_r) \\ &\quad + \frac{1}{2} \sum_{i=1}^d (J_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) (J_\varepsilon(\zeta - \tau_{\Delta x, i}^-) (D_-^i A(u_{\Delta x}))^2 \star J_{r_0} \otimes J_r) \\ &=: \mathcal{T}_1^+ + \mathcal{T}_1^-. \end{aligned}$$

Observe that

$$\partial_{x_i} \rho_{\varepsilon, r, r_0} = \partial_{x_i} (\chi_\varepsilon(A(u); \zeta) \star J_{r_0} \otimes J_r) = J_\varepsilon(\zeta - A(u)) \partial_{x_i} A(u) \star J_{r_0} \otimes J_r.$$

Using Lemma 5.3 once more gives

$$\begin{aligned} (D_+^i + D_-^i) \rho_{\varepsilon, r, r_0}^{\Delta x} &= (J_\varepsilon(\zeta - \theta_{\Delta x, i}^+) D_+^i A(u_{\Delta x}) \star J_{r_0} \otimes J_r \\ &\quad + J_\varepsilon(\zeta - \theta_{\Delta x, i}^-) D_-^i A(u_{\Delta x})) \star J_{r_0} \otimes J_r. \end{aligned}$$

Hence,

$$\begin{aligned} \nabla \rho_{\varepsilon, r, r_0} \cdot (D_+ + D_-) \rho_{\varepsilon, r, r_0}^{\Delta x} &= \sum_{i=1}^d (J_\varepsilon(\zeta - A(u)) \partial_{x_i} A(u) \star J_{r_0} \otimes J_r) (J_\varepsilon(\zeta - \theta_{\Delta x, i}^+) D_+^i A(u_{\Delta x}) \star J_{r_0} \otimes J_r) \\ &\quad + \sum_{i=1}^d (J_\varepsilon(\zeta - A(u)) \partial_{x_i} A(u) \star J_{r_0} \otimes J_r) (J_\varepsilon(\zeta - \theta_{\Delta x, i}^-) D_-^i A(u_{\Delta x})) \star J_{r_0} \otimes J_r. \end{aligned}$$

Adding and subtracting we obtain

$$-\nabla \rho_{\varepsilon, r, r_0} \cdot (D_+ + D_-) \rho_{\varepsilon, r, r_0}^{\Delta x} = \mathcal{T}_2^+ + \mathcal{T}_2^- + R_1^+ + R_1^-,$$

where

$$\begin{aligned} \mathcal{T}_2^+ &= - \sum_{i=1}^d (J_\varepsilon(\zeta - A(u)) \partial_{x_i} A(u) \star J_{r_0} \otimes J_r) (J_\varepsilon(\zeta - \tau_{\Delta x, i}^+) D_+^i A(u_{\Delta x}) \star J_{r_0} \otimes J_r), \\ \mathcal{T}_2^- &= - \sum_{i=1}^d (J_\varepsilon(\zeta - A(u)) \partial_{x_i} A(u) \star J_{r_0} \otimes J_r) (J_\varepsilon(\zeta - \tau_{\Delta x, i}^-) D_-^i A(u_{\Delta x})) \star J_{r_0} \otimes J_r. \end{aligned}$$

For each $1 \leq i \leq d$,

$$\begin{aligned} J_\varepsilon(\zeta - A(u_{\Delta x})) &= \frac{1}{2} \left(J_\varepsilon(\zeta - A(u_{\Delta x})) - J_\varepsilon(\zeta - A(\tau_{\Delta x, i}^+)) \right) \\ &\quad + \frac{1}{2} \left(J_\varepsilon(\zeta - A(u_{\Delta x})) - J_\varepsilon(\zeta - A(\tau_{\Delta x, i}^-)) \right) \\ &\quad + \frac{1}{2} \left(J_\varepsilon(\zeta - A(\tau_{\Delta x, i}^+)) + J_\varepsilon(\zeta - A(\tau_{\Delta x, i}^-)) \right). \end{aligned}$$

It follows that

$$(J_\varepsilon(\zeta - A(u_{\Delta x})) \star J_{r_0} \otimes J_r) n_{A, \varepsilon, r, r_0} = \mathcal{T}_3^+ + \mathcal{T}_3^- + R_2^+ + R_2^-,$$

where

$$\begin{aligned} \mathcal{T}_3^+ &= \frac{1}{2} \sum_{i=1}^d (J_\varepsilon(\zeta - A(\tau_{\Delta x, i}^+)) \star J_{r_0} \otimes J_r) (J_\varepsilon(\zeta - A(u)) (\partial_{x_i} A(u))^2 \star J_{r_0} \otimes J_r), \\ \mathcal{T}_3^- &= \frac{1}{2} \sum_{i=1}^d (J_\varepsilon(\zeta - A(\tau_{\Delta x, i}^-)) \star J_{r_0} \otimes J_r) (J_\varepsilon(\zeta - A(u)) (\partial_{x_i} A(u))^2 \star J_{r_0} \otimes J_r). \end{aligned}$$

Note that

$$E_{\Delta x, \varepsilon, r, r_0} = \sum_{k=1}^3 (\mathcal{T}_k^+ + \mathcal{T}_k^-) + \sum_{k=1}^2 (R_k^+ + R_k^-).$$

Now,

$$\begin{aligned} \sum_{k=1}^3 \mathcal{T}_k^+ &= \frac{1}{2} \sum_{i=1}^d (J_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) ((J_\varepsilon(\zeta - \tau_{\Delta x, i}^+) (D_+^i A(u_{\Delta x}))^2 \star J_{r_0} \otimes J_r) \\ &\quad - \sum_{i=1}^d (J_\varepsilon(\zeta - A(u)) \partial_{x_i} A(u) \star J_{r_0} \otimes J_r) (J_\varepsilon(\zeta - \tau_{\Delta x, i}^+) D_+^i A(u_{\Delta x}) \star J_{r_0} \otimes J_r) \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2} \sum_{i=1}^d (J_\varepsilon(\zeta - A(\tau_{\Delta x, i}^+)) \star J_{r_0} \otimes J_r) (J_\varepsilon(\zeta - A(u)) (\partial_{x_i} A(u))^2 \star J_{r_0} \otimes J_r) \\
& = \frac{1}{2} \sum_{i=1}^d J_\varepsilon(\zeta - A(u)) J_\varepsilon(\zeta - A(\tau_{\Delta x, i}^+)) (\partial_{x_i} A(u) - D_+^i A(u_{\Delta x}))^2 \\
& \quad \star^{(u, u_{\Delta x})} J_{r_0} \otimes J_r \otimes J_{r_0} \otimes J_r \\
& \geq 0.
\end{aligned}$$

The obtained inequality holds for all $(t, x, \zeta) \in (r_0, T - r_0) \times \mathbb{R}^d \times \mathbb{R}$. Similarly,

$$\sum_{i=1}^3 \mathcal{T}_i^-(t, x, \zeta) \geq 0 \text{ for all } (t, x, \zeta) \in (r_0, T - r_0) \times \mathbb{R}^d \times \mathbb{R}.$$

This concludes the proof of the lemma. $\square$

5.3. Bounding error terms. We are going to estimate a series of “unwanted” terms coming from Lemmas 5.1 and 5.4. To this end, we will need to gather three technical lemmas, the first one being a simple application of Young’s inequality for convolutions.

Lemma 5.5. *Let $\psi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a measurable function, and $u, v : \mathbb{R}^d \rightarrow \mathbb{R}$ be measurable functions satisfying*

$$|\psi(u(x_1), v(x_2))| \leq K_1(x_1) K_2(x_2) \quad (x_1, x_2 \in \mathbb{R}^d),$$

for some $K_1 \in L^p(\mathbb{R}^d)$, $1 \leq p \leq \infty$, and $K_2 \in L^1(\mathbb{R}^d)$. Then

$$\left\| \psi(u, v) \star^{(u, v)} f \otimes g \right\|_{L^1(\mathbb{R}^d)} \leq \|K_1\|_{L^p(\mathbb{R}^d)} \|K_2\|_{L^1(\mathbb{R}^d)} \|f\|_{L^q(\mathbb{R}^d)} \|g\|_{L^1(\mathbb{R}^d)},$$

for any $g \in L^1(\mathbb{R})$ and $f \in L^q(\mathbb{R})$ where $p^{-1} + q^{-1} = 1$.

If $\psi \in L^\infty(\mathbb{R}^2)$, then

$$(5.15) \quad \left\| \psi(u, v) \star^{(u, v)} f \otimes g \right\|_{L^\infty(\mathbb{R}^d)} \leq \|\psi\|_{L^\infty(\mathbb{R}^2)} \|f\|_{L^1(\mathbb{R}^d)} \|g\|_{L^1(\mathbb{R}^d)}.$$

Proof. Observe that

$$\begin{aligned}
& \left\| \psi(u, v) \star^{(u, v)} f \otimes g \right\|_{L^1(\mathbb{R}^d)} \\
& \leq \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d} K_1(y_1) K_2(y_2) |f(x - y_1)| |g(x - y_2)| dy_1 dy_2 dx \\
& = \|(K_1 \star |f|)(K_2 \star |g|)\|_{L^1(\mathbb{R}^d)}.
\end{aligned}$$

By Hölder’s inequality,

$$\|(K_1 \star |f|)(K_2 \star |g|)\|_{L^1(\mathbb{R}^d)} \leq \|K_1 \star |f|\|_{L^\infty(\mathbb{R}^d)} \|K_2 \star |g|\|_{L^1(\mathbb{R}^d)}.$$

By Young’s inequality for convolutions, $\|K_1 \star |f|\|_{L^\infty(\mathbb{R}^d)} \leq \|K_1\|_{L^p(\mathbb{R}^d)} \|f\|_{L^q(\mathbb{R}^d)}$ and $\|K_2 \star |g|\|_{L^1(\mathbb{R}^d)} \leq \|K_2\|_{L^1(\mathbb{R}^d)} \|g\|_{L^1(\mathbb{R}^d)}$. Equation (5.15) follows, since

$$\left| \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \psi(u(y_1), v(y_2)) f(x - y_1) g(x - y_2) dy_1 dy_2 \right| \leq \|\psi\|_{L^\infty(\mathbb{R}^2)} \|f\|_{L^1(\mathbb{R}^d)} \|g\|_{L^1(\mathbb{R}^d)}.$$

$\square$

The next lemma is at the heart of the matter, permitting us to estimate some terms involving convolutions against approximate delta functions.

Lemma 5.6. *For real numbers a and b , let $\tau = \tau_\varepsilon(a, b, \zeta)$ and $\theta = \theta_\varepsilon(a, b, \zeta)$ be as in Lemma 5.2.*

(i) For $f \in L^1_{\text{loc}}(\mathbb{R})$, define

$$\mathcal{T}_\varepsilon^1(f) = \int_{\mathbb{R}} (J_\varepsilon(\zeta - \tau) - J_\varepsilon(\zeta - a)) f(\zeta) d\zeta.$$

Then

$$|\mathcal{T}_\varepsilon^1(f)| \leq \left| \int_a^b |\partial_\xi(f \star J_\varepsilon)(\xi)| d\xi \right|.$$

(ii) For $f \in L^\infty(\mathbb{R})$, define

$$\mathcal{T}_\varepsilon^2(f) = \int_{\mathbb{R}} (J_\varepsilon(\zeta - \theta) - J_\varepsilon(\zeta - \tau))(b - a) f(\zeta) d\zeta.$$

Then

$$|\mathcal{T}_\varepsilon^2(f)| \leq \frac{1}{2} \|f\|_{L^\infty(\mathbb{R})} |b - a|.$$

(iii) Suppose $\{f_\varepsilon\}_{\varepsilon>0} \subset W^{1,1}_{\text{loc}}(\mathbb{R})$ and assume that there exists $f \in W^{1,1}_{\text{loc}}(\mathbb{R})$ such that $f_\varepsilon \rightarrow f$ in $W^{1,1}_{\text{loc}}(\mathbb{R})$. Then

$$\left| \lim_{\varepsilon \downarrow 0} \mathcal{T}_\varepsilon^1(f_\varepsilon) \right| \leq \left| \int_a^b |f'(\xi)| d\xi \right|.$$

Proof. Assume that $a < b$. By Lemma 5.2 and Fubini's Theorem,

$$\mathcal{T}_\varepsilon^1(f) = \frac{2}{(b-a)^2} \int_a^b ((f \star J_\varepsilon)(\xi) - (f \star J_\varepsilon)(a)) (b - \xi) d\xi.$$

Since $\partial_\xi(b - \xi)^2 = -2(b - \xi)$, integration by parts yields

$$(5.16) \quad \mathcal{T}_\varepsilon^1(f) = \int_a^b \partial_\xi(f \star J_\varepsilon)(\xi) \frac{(b - \xi)^2}{(b - a)^2} d\xi.$$

Then statement (i) follows, since

$$(5.17) \quad \frac{(b - \xi)^2}{(b - a)^2} \leq 1, \text{ whenever } a \leq \xi \leq b.$$

By Lemma 5.2,

$$\mathcal{T}_\varepsilon^2(f) = \frac{1}{b-a} \int_a^b (f \star J_\varepsilon)(\xi) (2\xi - (a+b)) d\xi.$$

As $\|f \star J_\varepsilon\|_{L^\infty(\mathbb{R})} \leq \|f\|_{L^\infty(\mathbb{R})}$, we may conclude that

$$\mathcal{T}_\varepsilon^2(f) \leq \frac{1}{b-a} \int_a^b \|f\|_{L^\infty(\mathbb{R})} |(2\xi - (a+b))| d\xi.$$

This implies statement (ii), because

$$\int_a^b |2\xi - (a+b)| d\xi = \frac{1}{2}(b-a)^2.$$

Finally, we establish statement (iii). By the triangle inequality and Young's inequality for convolutions,

$$\|f'_\varepsilon \star J_\varepsilon - f'\|_{L^1(V)} \leq \|f'_\varepsilon - f'\|_{L^1(V)} + \|f' \star J_\varepsilon - f'\|_{L^1(V)}$$

for any compact $V \subset \mathbb{R}$. Hence $(f'_\varepsilon \star J_\varepsilon) \rightarrow f'$ in $L^1_{\text{loc}}(\mathbb{R})$ as $\varepsilon \downarrow 0$. By (5.16) and (5.17) it follows that

$$\lim_{\varepsilon \downarrow 0} \mathcal{T}_\varepsilon^1(f_\varepsilon) = \lim_{\varepsilon \downarrow 0} \int_a^b (f'_\varepsilon \star J_\varepsilon)(\xi) \frac{(b - \xi)^2}{(b - a)^2} d\xi = \int_a^b f'(\xi) \frac{(b - \xi)^2}{(b - a)^2} d\xi.$$

The estimate follows thanks to (5.17). $\square$

We need one more lemma bounding some specific convolution integrals.

Lemma 5.7. *Suppose $f \in C(\mathbb{R})$ and let $R_\varepsilon^f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by (4.5). Then*

$$(5.18) \quad R_\varepsilon^f(u, \zeta) = \int_0^u (f(\sigma) - f(\zeta)) J_\varepsilon(\sigma - \zeta) d\sigma,$$

for all $u, \zeta \in \mathbb{R}$. Furthermore, if f is Lipschitz continuous, then

$$(5.19) \quad \int_{\mathbb{R}} |R_\varepsilon^f(u, \zeta)| d\zeta \leq \varepsilon \|f\|_{\text{Lip}} |u|.$$

Suppose $A' \geq \eta > 0$ and let g be defined by $g \circ A = f$. For $a, b \in \mathbb{R}$, let

$$Z(a, b) = \int_{\mathbb{R}} \text{sign}_\varepsilon(\zeta - a) R_\varepsilon^{g'}(b; \zeta) d\zeta.$$

Then

$$(5.20) \quad |Z(a, b)| \leq 4 \|f\|_{\text{Lip}} \frac{\varepsilon}{\eta}.$$

Proof. Observe that

$$\begin{aligned} & \int_{\mathbb{R}} f(\sigma) \chi(u; \sigma) J_\varepsilon(\zeta - \sigma) d\sigma \\ &= \underbrace{\int_{\mathbb{R}} (f(\sigma) - f(\zeta)) \chi(u; \sigma) J_\varepsilon(\zeta - \sigma) d\sigma}_{R_\varepsilon^f(u, \zeta)} + f(\zeta) \chi_\varepsilon(u; \zeta). \end{aligned}$$

Let $q'(\sigma) = (f(\sigma) - f(\zeta)) J_\varepsilon(\zeta - \sigma)$. Equation (5.18) follows, since

$$\int_{\mathbb{R}} (f(\sigma) - f(\zeta)) \chi(u; \sigma) J_\varepsilon(\zeta - \sigma) d\sigma = q(u) - q(0) = \int_0^u q'(\sigma) d\sigma.$$

To prove (5.19) observe that

$$\int_{\mathbb{R}} |R_\varepsilon^f(u, \zeta)| d\zeta \leq \int_{\mathbb{R}} |\chi(u; \sigma)| \left(\int_{\mathbb{R}} |f(\sigma) - f(\zeta)| J_\varepsilon(\zeta - \sigma) d\zeta \right) d\sigma.$$

The result follows as

$$\int_{\mathbb{R}} |f(\sigma) - f(\zeta)| J_\varepsilon(\zeta - \sigma) d\sigma \leq \|f\|_{\text{Lip}} \varepsilon.$$

Let us prove (5.20). Take

$$H_\varepsilon^{g'}(b; \zeta) = \int_0^\zeta R_\varepsilon^{g'}(b; \sigma) d\sigma.$$

Integration by parts yields

$$Z(a, b) = -2 \int_{\mathbb{R}} J_\varepsilon(\zeta - a) H_\varepsilon^{g'}(b; \zeta) d\zeta = -(H_\varepsilon^{g'}(b; \cdot) \star J_\varepsilon)(a).$$

By (5.18),

$$H_\varepsilon^{g'}(b; a) = \int_0^a \int_0^b (g'(\omega) - g'(\sigma)) J_\varepsilon(\omega - \sigma) d\omega d\sigma.$$

Due to the symmetry of J_ε ,

$$\begin{aligned} H_\varepsilon^{g'}(b; a) &= \int_0^a \int_0^b g'(\omega) J_\varepsilon(\omega - \sigma) d\omega d\sigma \\ &\quad - \int_0^a \int_0^b g'(\sigma) J_\varepsilon(\omega - \sigma) d\omega d\sigma \\ &= \int_0^a \int_0^b g'(\omega) J_\varepsilon(\omega - \sigma) d\omega d\sigma \end{aligned}$$

$$\begin{aligned}
& - \int_0^b \int_0^a g'(\omega) J_\varepsilon(\omega - \sigma) d\omega d\sigma \\
& = \int_{\mathbb{R}} g'(\omega) \chi(b; \omega) \left(\int_0^a J_\varepsilon(\omega - \sigma) d\sigma \right) d\omega \\
& \quad - \int_{\mathbb{R}} g'(\omega) \chi(a; \omega) \left(\int_0^b J_\varepsilon(\omega - \sigma) d\sigma \right) d\omega.
\end{aligned}$$

Note that

$$\int_0^a J_\varepsilon(\omega - \sigma) d\sigma = \int_{\mathbb{R}} \chi(a; \sigma) J_\varepsilon(\omega - \sigma) d\sigma = \chi_\varepsilon(a; \omega).$$

Hence,

$$H_\varepsilon^{g'}(b; a) = \int_{\mathbb{R}} g'(\omega) (\chi(b; \omega) \chi_\varepsilon(a; \omega) - \chi(a; \omega) \chi_\varepsilon(b; \omega)) d\omega.$$

Set

$$\lambda(a, b; \omega) := \chi(b; \omega) \chi_\varepsilon(a; \omega) - \chi(a; \omega) \chi_\varepsilon(b; \omega).$$

To find the support of $\lambda(a, b; \omega)$ we first observe that $\lambda(a, b; \omega) = -\lambda(b, a; \omega)$. This reduces the situation to the following cases:

$$\begin{cases} 0 \leq a \leq b : & |\lambda(a, b; \omega)| \leq \mathbb{1}_{|a-\omega| \leq \varepsilon}, \\ b \leq a \leq 0 : & |\lambda(a, b; \omega)| \leq \mathbb{1}_{|a-\omega| \leq \varepsilon}, \\ a \leq 0 \leq b : & |\lambda(a, b; \omega)| \leq \mathbb{1}_{|\omega| \leq \varepsilon}. \end{cases}$$

It thus follows that

$$|H_\varepsilon^{g'}(b, a)| \leq 2 \|g'\|_\infty \varepsilon.$$

Statement (5.20) follows as $g'(A(z))A'(z) = f'(z)$, which implies $\|g'\|_\infty \leq \|f\|_{\text{Lip}} \eta^{-1}$. $\square$

We have now the tools needed to start estimating the error terms in Lemmas 5.1 and 5.4, starting with those in Lemma 5.4.

Estimate 5.1. *Let $R_1^\pm$ be defined in Lemma 5.4. Then there exists a constant $C = C(d, J)$ such that*

$$\left\| \int_{\mathbb{R}} R_1^+(\zeta) + R_1^-(\zeta) d\zeta \right\|_{L^1(\Pi_T^{r_0})} \leq C \frac{\Delta x}{r^2} \left(1 + \frac{\Delta x}{r} \right) \|D_- A(u_{\Delta x})\|_{L^1(\Pi_T; \mathbb{R}^d)}.$$

Proof. Let us first make an observation regarding the similarity of these terms. By statement (iv) of Lemma 5.2, recalling also the definition of $\theta_{\Delta x, i}^\pm$ and $\tau_{\Delta x, i}^\pm$ in Lemma 5.3,

$$\begin{aligned}
& S_{\Delta x_i} \left(J_\varepsilon(\zeta - \tau_{\Delta x, i}^-) - J_\varepsilon(\zeta - \theta_{\Delta x, i}^-) \right) \\
& = J_\varepsilon(\zeta - \tau_\varepsilon(S_{\Delta x_i} A(u_{\Delta x}), A(u_{\Delta x}), \zeta)) - J_\varepsilon(\zeta - \theta_\varepsilon(S_{\Delta x_i} A(u_{\Delta x}), A(u_{\Delta x}), \zeta)) \\
& = - \left(J_\varepsilon(\zeta - \tau_{\Delta x, i}^+) - J_\varepsilon(\zeta - \theta_{\Delta x, i}^+) \right).
\end{aligned}$$

Recalling that $\rho_{\varepsilon, r, r_0} = \chi_\varepsilon(A(u); \cdot) \star J_{r_0} \otimes J_r$, which implies

$$\begin{aligned}
& R_1^+(\zeta) + R_1^-(\zeta) \\
& = - \sum_{i=1}^d S_{\Delta x_i} \left[\left(J_\varepsilon(\zeta - \tau_{\Delta x, i}^-) - J_\varepsilon(\zeta - \theta_{\Delta x, i}^-) \right) D_-^i(u_{\Delta x}) \star (J_{r_0} \otimes J_r) \right] \partial_{x_i} \rho_{\varepsilon, r, r_0} \\
& \quad + \sum_{i=1}^d \left[\left(J_\varepsilon(\zeta - \tau_{\Delta x, i}^+) - J_\varepsilon(\zeta - \theta_{\Delta x, i}^+) \right) D_-^i(u_{\Delta x}) \star (J_{r_0} \otimes J_r) \right] \partial_{x_i} \rho_{\varepsilon, r, r_0}
\end{aligned}$$

$$\begin{aligned}
&= - \sum_{i=1}^d (S_{\Delta x_i} - 1) \partial_{x_i} \rho_{\varepsilon, r, r_0} \\
&\quad \times \left[\left(J_\varepsilon(\zeta - \tau_{\Delta x, i}^-) - J_\varepsilon(\zeta - \theta_{\Delta x, i}^-) \right) D_-^i(u_{\Delta x}) \star (J_{r_0} \otimes J_r) \right] \\
&= -\Delta x \sum_{i=1}^d \left(J_\varepsilon(\zeta - \tau_{\Delta x, i}^-) - J_\varepsilon(\zeta - \theta_{\Delta x, i}^-) \right) D_-^i A(u_{\Delta x}) \chi_\varepsilon(A(u); \zeta) \\
&\quad \stackrel{(u, u_{\Delta x})}{\star} (J_{r_0} \otimes \partial_{x_i} J_r) \otimes (J_{r_0} \otimes D_+^i J_r).
\end{aligned}$$

By statement (ii) of Lemma 5.6,

$$\begin{aligned}
(5.21) \quad & \left| \int_{\mathbb{R}} R_1^+(\zeta) + R_1^-(\zeta) d\zeta \right| \\
& \leq \frac{\Delta x}{2} \sum_{i=1}^d \|\chi_\varepsilon(A(u); \cdot)\|_{L^\infty(\mathbb{R})} |D_-^i A(u_{\Delta x})| \stackrel{(u, u_{\Delta x})}{\star} |J_{r_0} \otimes \partial_{x_i} J_r \otimes J_{r_0} \otimes D_+^i J_r|.
\end{aligned}$$

By Lemma 5.5,

$$\begin{aligned}
\left\| \int_{\mathbb{R}} R_1^+(\zeta) + R_1^-(\zeta) d\zeta \right\|_{L^1(\Pi_T^{r_0})} &\leq \frac{\Delta x}{2} \sum_{i=1}^d \|D_-^i A(u_{\Delta x})\|_{L^1(\Pi_T)} \\
&\quad \times \|\partial_{x_i} J_r\|_{L^1(\mathbb{R}^d)} \|D_+^i J_r\|_{L^1(\mathbb{R}^d)}.
\end{aligned}$$

Recall that $\|\partial_{x_i} J_r\|_{L^1(\mathbb{R}^d)} \leq 2 \|J'\|_\infty r^{-1}$. Note that

$$\begin{aligned}
|D_+^i J_r(x)| &= \frac{1}{\Delta x} |J_r(x_i + \Delta x) - J_r(x_i)| \prod_{j \neq i} J_r(x_j) \\
&\leq \frac{1}{r^2} \|J'\|_\infty \mathbf{1}_{|x_i| \leq r + \Delta x} \prod_{j \neq i} J_r(x_j).
\end{aligned}$$

Hence

$$(5.22) \quad \|D_+^i J_r\|_{L^1(\mathbb{R}^d)} \leq \frac{1}{r^2} \|J'\|_\infty \int_{\mathbb{R}} \mathbf{1}_{|x_i| \leq r + \Delta x} dx_i = 2 \|J'\|_\infty \frac{1}{r} \left(1 + \frac{\Delta x}{r} \right)$$

The estimate follows from (5.22) and (5.21). $\square$

Estimate 5.2. Let $R_2^\pm$ be defined in Lemma 5.4. Then there exists a constant $C = C(d, J)$ such that

$$\begin{aligned}
&\left\| \int_{\mathbb{R}} R_2^+(\zeta) + R_2^-(\zeta) d\zeta \right\|_{L^1(\Pi_T^{r_0})} \\
&\leq C \frac{\Delta x}{\varepsilon^2 \sqrt{r_0 r^d}} \|D_+ A(u_{\Delta x})\|_{L^2(\Pi_T; \mathbb{R}^d)} \|\nabla A(u)\|_{L^2(\Pi_T; \mathbb{R}^d)}^2.
\end{aligned}$$

Proof. Let us consider R_2^+ . The term R_2^- is treated the same way. By Lemma 5.6,

$$\begin{aligned}
\left| \int_{\mathbb{R}} R_2^+(\zeta) d\zeta \right| &\leq \frac{1}{2} \sum_{i=1}^d \left| \int_{\mathbb{R}} \left(J_\varepsilon(\zeta - A(u_{\Delta x})) - J_\varepsilon(\zeta - A(\tau_{\Delta x, i}^+)) \right) J_\varepsilon(\zeta - A(u)) d\zeta \right| \\
&\quad \times (\partial_{x_i} A(u))^2 \stackrel{(u, u_{\Delta x})}{\star} (J_{r_0} \otimes J_r) \otimes (J_{r_0} \otimes J_r) \\
&\leq \frac{1}{2} \sum_{i=1}^d \left| \int_{A(u_{\Delta x})}^{S_{\Delta x_i} A(u_{\Delta x})} |\partial_\xi (J_\varepsilon(\cdot - A(u)) \star J_\varepsilon(\xi))| d\xi \right| \\
&\quad \times (\partial_{x_i} A(u))^2 \stackrel{(u, u_{\Delta x})}{\star} (J_{r_0} \otimes J_r) \otimes (J_{r_0} \otimes J_r).
\end{aligned}$$

By Young's inequality for convolutions,

$$\|J_\varepsilon(\cdot - A(u)) \star J'_\varepsilon\|_{L^\infty(\mathbb{R})} \leq \|J_\varepsilon(\cdot - A(u))\|_{L^\infty(\mathbb{R})} \|J'_\varepsilon\|_{L^1(\mathbb{R})} \leq \frac{2}{\varepsilon^2} \|J\|_\infty \|J'\|_\infty.$$

Hence,

$$\left| \int_{\mathbb{R}} R_2^+(\zeta) d\zeta \right| \leq \frac{\Delta x}{\varepsilon^2} \|J\|_\infty \|J'\|_\infty \sum_{i=1}^d |D_+^i A(u_{\Delta x})| (\partial_{x_i} A(u))^2 \stackrel{(u, u_{\Delta x})}{\star} (J_{r_0} \otimes J_r) \otimes (J_{r_0} \otimes J_r).$$

Applying Lemma 5.5, with $K_1(u) = |u|$, $K_2(v) = v^2$, $p = q = 2$, we get

$$\left\| \int_{\mathbb{R}} R_2^+(\zeta) d\zeta \right\|_{L^1(\Pi_T^{r_0})} \leq \frac{\Delta x}{\varepsilon^2} \|J\|_\infty \|J'\|_\infty \sum_{i=1}^d \|D_+^i A(u_{\Delta x})\|_{L^2(\Pi_T)} \|\partial_{x_i} A(u)\|_{L^2(\Pi_T)}^2 \times \|J_{r_0} \otimes J_r\|_{L^2(\mathbb{R} \times \mathbb{R}^d)} \|J_{r_0} \otimes J_r\|_{L^1(\mathbb{R} \times \mathbb{R}^d)}.$$

Now

$$\|J_{r_0} \otimes J_r\|_{L^2(\mathbb{R} \times \mathbb{R}^d)} = \|J_{r_0}\|_{L^2(\mathbb{R})} \prod_{i=1}^d \|J_r\|_{L^2(\mathbb{R})} = \frac{1}{\sqrt{r_0 r^d}} \|J\|_{L^2(\mathbb{R})}^{d+1}.$$

□

Estimate 5.3. Let $R_1^\pm$ be defined in Lemma 5.4 and suppose $d = 1$. Then there exists a constant $C = C(J)$ such that

$$\left\| \lim_{\varepsilon \downarrow 0} \int_{\mathbb{R}} R_2^+(\zeta) + R_2^-(\zeta) d\zeta \right\|_{L^1(\Pi_T^{r_0})} \leq C \left(\frac{\Delta x}{r} \|f\|_{\text{Lip}} + \frac{\Delta x}{r^2} \|A\|_{\text{Lip}} + \frac{\Delta x}{r_0} \right) \|D_+ u_{\Delta x}\|_{L^1(\Pi_T)}.$$

Proof. We consider R_2^+ . The R_2^- term can be treated similarly. Note that for $d = 1$,

$$\int_{\mathbb{R}} R_2^+(\zeta) d\zeta = \int_{\mathbb{R}} (J_\varepsilon(\zeta - A(u_{\Delta x})) - J_\varepsilon(\zeta - A(\tau_{\Delta x}^+))) n_{A, \varepsilon, r, r_0}(\zeta) d\zeta \stackrel{(u_{\Delta x})}{\star} J_{r_0} \otimes J_r,$$

where $n_{A, \varepsilon, r, r_0}$ is defined in Lemma 4.7. The map $\zeta \mapsto n_{A, \varepsilon, r, r_0}(t, x, \zeta)$ belongs to $W_{\text{loc}}^{1,1}(\mathbb{R})$ for each fixed $(t, x) \in (r_0, T - r_0) \times \mathbb{R}$. Due to Lemmas 4.7, 5.7, and 4.6,

$$\lim_{\varepsilon \rightarrow 0} \partial_\zeta n_{A, \varepsilon, r, r_0}(\zeta) = B'(\zeta) \partial_t \rho_{r, r_0}(\zeta) + g'(\zeta) \partial_x \rho_{r, r_0}(\zeta) - \partial_x^2 \rho_{r, r_0}(\zeta)$$

in $L^1(\mathbb{R})$ for each fixed (t, x) , where

$$\rho_{r, r_0}(\zeta) = \chi(A(u); \zeta) \star J_{r_0} \otimes J_r.$$

By statement (iii) of Lemma 5.6,

$$\begin{aligned} \left| \lim_{\varepsilon \downarrow 0} \int_{\mathbb{R}} R_2^+(\zeta) d\zeta \right| &\leq \left| \int_{A(u_{\Delta x})}^{S_{\Delta x} A(u_{\Delta x})} |B'(\zeta) \partial_t \rho_{r, r_0}(\zeta)| d\zeta \right| \stackrel{(u_{\Delta x})}{\star} J_{r_0} \otimes J_r \\ &\quad + \left| \int_{A(u_{\Delta x})}^{S_{\Delta x} A(u_{\Delta x})} |g'(\zeta) \partial_x \rho_{r, r_0}(\zeta)| d\zeta \right| \stackrel{(u_{\Delta x})}{\star} J_{r_0} \otimes J_r \\ &\quad + \left| \int_{A(u_{\Delta x})}^{S_{\Delta x} A(u_{\Delta x})} |\partial_x^2 \rho_{r, r_0}(\zeta)| d\zeta \right| \stackrel{(u_{\Delta x})}{\star} J_{r_0} \otimes J_r \\ &=: \mathcal{T}_1 + \mathcal{T}_2 + \mathcal{T}_3. \end{aligned}$$

We consider each term separately. Let $B(\zeta) = \xi$ or equivalently $A(\xi) = \zeta$. It follows that

$$\begin{aligned} \mathcal{T}_1 &\leq \left| \int_{A(u_{\Delta x})}^{S_{\Delta x} A(u_{\Delta x})} |B'(\zeta) \chi(A(u); \zeta)| d\zeta \right|^{(u, u_{\Delta x})} \star |\partial_t J_{r_0}| \otimes J_r \otimes J_{r_0} \otimes J_r \\ &= \left| \int_{u_{\Delta x}}^{S_{\Delta x} u_{\Delta x}} |\chi(u; \xi)| d\xi \right|^{(u, u_{\Delta x})} \star |\partial_t J_{r_0}| \otimes J_r \otimes J_{r_0} \otimes J_r \\ &\leq \Delta x |D_+ A(u_{\Delta x})|^{(u, u_{\Delta x})} \star |\partial_t J_{r_0}| \otimes J_r \otimes J_{r_0} \otimes J_r. \end{aligned}$$

By Lemma 5.5,

$$\|\mathcal{T}_1\|_{L^1(\Pi_T^{r_0})} \leq 2 \frac{\Delta x}{r_0} \|J'\|_{\infty} \|D_+ u_{\Delta x}\|_{L^1(\Pi_T)}.$$

Observe that $g'(A(\xi))A'(\xi) = f'(\xi)$ and $d\zeta = A'(\xi)d\xi$. Hence,

$$\begin{aligned} \mathcal{T}_2 &\leq \left| \int_{A(u_{\Delta x})}^{S_{\Delta x} A(u_{\Delta x})} |g'(\zeta) \chi(A(u); \zeta)| d\zeta \right|^{(u, u_{\Delta x})} \star J_{r_0} \otimes |\partial_x J_r| \otimes J_{r_0} \otimes J_r \\ &= \left| \int_{u_{\Delta x}}^{S_{\Delta x} u_{\Delta x}} |f'(\xi) \chi(u; \xi)| d\xi \right|^{(u, u_{\Delta x})} \star J_{r_0} \otimes |\partial_x J_r| \otimes J_{r_0} \otimes J_r \\ &\leq \Delta x \|f\|_{\text{Lip}} |D_+ u_{\Delta x}|^{(u, u_{\Delta x})} \star J_{r_0} \otimes |\partial_x J_r| \otimes J_{r_0} \otimes J_r. \end{aligned}$$

By Lemma 5.5,

$$\|\mathcal{T}_2\|_{L^1(\Pi_T^{r_0})} \leq 2 \|f\|_{\text{Lip}} \frac{\Delta x}{r} \|J'\|_{\infty} \|D_+ u_{\Delta x}\|_{L^1(\Pi_T)}.$$

Similarly,

$$\begin{aligned} \mathcal{T}_3 &\leq \left| \int_{A(u_{\Delta x})}^{S_{\Delta x} A(u_{\Delta x})} |\chi(A(u); \zeta)| d\zeta \right|^{(u, u_{\Delta x})} \star J_{r_0} \otimes |\partial_x^2 J_r| \otimes J_{r_0} \otimes J_r \\ &\leq \Delta x |D_+ A(u_{\Delta x})|^{(u, u_{\Delta x})} \star J_{r_0} \otimes |\partial_x^2 J_r| \otimes J_{r_0} \otimes J_r. \end{aligned}$$

By Lemma 5.5,

$$\|\mathcal{T}_3\|_{L^1(\Pi_T^{r_0})} \leq 2 \frac{\Delta x}{r^2} \|J''\|_{\infty} \|D_+ A(u_{\Delta x})\|_{L^1(\Pi_T)}.$$

□

Estimate 5.4. Let U be the second term in (5.3), Lemma 5.1, that is,

$$U = 2 \int_{\mathbb{R}} \nabla \rho_{\varepsilon, r, r_0} \cdot (2\nabla - (D_+ + D_-) \rho_{\varepsilon, r, r_0}^{\Delta x} d\zeta.$$

Then there exists a constant $C = C(d, J)$ such that

$$\|U\|_{L^1(\Pi_T^{r_0})} \leq C \frac{\Delta x^2}{r^3} \left(1 + \frac{\Delta x}{r}\right) \|A(u_{\Delta x})\|_{L^1([0, T]; BV(\mathbb{R}^d))}.$$

Remark 5.1. The BV norm may be replaced by the L^1 norm at the expense of an extra factor r^{-1} .

Proof. Clearly,

$$\begin{aligned} &\left\| \int_{\mathbb{R}} \nabla \rho_{\varepsilon, r, r_0} \cdot (2\nabla - (D_+ + D_-) \rho_{\varepsilon, r, r_0}^{\Delta x} d\zeta \right\|_{L^1(\Pi_T^{r_0})} \\ &\leq \|\nabla \rho_{\varepsilon, r, r_0}\|_{L^\infty(\Pi_T^{r_0} \times \mathbb{R}; \mathbb{R}_\infty^d)} \|(2\nabla - (D_+ + D_-) \rho_{\varepsilon, r, r_0}^{\Delta x})\|_{L^1(\Pi_T^{r_0} \times \mathbb{R}; \mathbb{R}_1^d)}. \end{aligned}$$

By Young's inequality for convolutions,

$$\|\partial_{x_i} \rho_{\varepsilon, r, r_0}\|_{L^\infty(\Pi_T^{r_0} \times \mathbb{R})} \leq \|\chi(A(u); \cdot)\|_{L^\infty(\Pi_T \times \mathbb{R})} \|J_\varepsilon \otimes \partial_{x_i} J_r \otimes J_{r_0}\|_{L^1(\mathbb{R} \times \mathbb{R}^d \times \mathbb{R})}$$

$$\leq \|\partial_{x_i} J_r\|_{L^1(\mathbb{R}^d)} \leq 2 \|J'\|_\infty r^{-1}.$$

We have

$$(2\partial_{x_i} - (D_+^i + D_-^i))\rho_{\varepsilon,r,r_0}^{\Delta x} = \chi(A(u_{\Delta x}); \cdot) \star J_\varepsilon \otimes (2\partial_{x_i} - (D_+^i + D_-^i))J_r \otimes J_{r_0}.$$

Using Taylor expansions with remainders,

$$\begin{aligned} ((D_+^i + D_-^i) - 2\partial_{x_i}) J_r(x) &= \frac{1}{2\Delta x} \int_0^{\Delta x} (z - \Delta x)^2 \partial_{x_i}^3 J_r(x_i + z) dz \prod_{j \neq i} J_r(x_j) \\ &\quad + \frac{1}{2\Delta x} \int_{-\Delta x}^0 (z + \Delta x)^2 \partial_{x_i}^3 J_r(x_i + z) dz \prod_{j \neq i} J_r(x_j) \\ &=: \partial_{x_i}(\varphi_1^i(x) + \varphi_2^i(x)), \end{aligned}$$

see for instance [25, p. 25]. Hence

$$(2\partial_{x_i} - (D_+^i + D_-^i))\rho_{\varepsilon,r,r_0}^{\Delta x} = \partial_{x_i}(\chi(A(u_{\Delta x}); \cdot) \star J_\varepsilon \otimes (\varphi_1^i + \varphi_2^i) \otimes J_{r_0}).$$

By Young's inequality for convolutions

$$\begin{aligned} \|(2\partial_{x_i} - (D_+^i + D_-^i))\rho_{\varepsilon,r,r_0}^{\Delta x}\|_{L^1(\Pi_T^{r_0} \times \mathbb{R})} &\leq \|\chi(A(u_{\Delta x}); \cdot)\|_{L^1([0,T] \times \mathbb{R}; BV(\mathbb{R}^d))} \\ &\quad \times \|\varphi_1^i + \varphi_2^i\|_{L^1(\mathbb{R}^d)}. \end{aligned}$$

Note that $\|\chi(A(u_{\Delta x}); \cdot)\|_{L^1([0,T] \times \mathbb{R}; BV(\mathbb{R}^d))} = \|A(u_{\Delta x})\|_{L^1([0,T]; BV(\mathbb{R}^d))}$. Now, as $\partial_{x_i} J_r(x_i + z) \leq r^{-3} \|J''\|_\infty \mathbf{1}_{|x_i + z| \leq r}$, it follows that

$$\begin{aligned} \|\varphi_1^i\|_{L^1(\mathbb{R}^d)} &= \frac{1}{2\Delta x} \int_{\mathbb{R}} \left| \int_0^{\Delta x} (z - \Delta x)^2 \partial_{x_i}^2 J_r(x_i + z) dz \right| dx_i \\ &\leq \frac{(r + \Delta x)}{\Delta x r^3} \|J''\|_\infty \int_0^{\Delta x} (z - \Delta x)^2 dz \\ &\leq \frac{1}{3} \|J''\|_\infty \frac{\Delta x^2}{r^2} \left(1 + \frac{\Delta x}{r}\right). \end{aligned}$$

The same estimate applies to φ_2^i . $\square$

Estimate 5.5. Let $\mathcal{T}$ be the term (5.9) from Lemma 5.1, that is,

$$\mathcal{T} = \int_R (\text{sign}_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r)(\Delta - D_- \cdot D_+) \rho_{\varepsilon,r,r_0}^{\Delta x} d\zeta.$$

Then there exists a constant $C = C(d, J)$ such that

$$\|\mathcal{T}\|_{L^1(\Pi_T^{r_0})} \leq C \frac{\Delta x^2}{r^3} \left(1 + \frac{\Delta x}{r}\right) \|A(u_{\Delta x})\|_{L^1([0,T]; BV(\mathbb{R}^d))}.$$

Remark 5.2. At the cost of an extra factor r^{-1} , the BV norm may be replaced by the L^1 norm.

Proof. First note that $|\text{sign}_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r| \leq 1$, so

$$|\mathcal{T}| \leq \|(\Delta - D_- \cdot D_+) \rho_{\varepsilon,r,r_0}^{\Delta x}\|_{L^1(\Pi_T \times \mathbb{R})}.$$

Now,

$$(\partial_{x_i}^2 - D_-^i D_+^i) \rho_{\varepsilon,r,r_0}^{\Delta x} = \chi(A(u_{\Delta x}); \cdot) \star J_\varepsilon \otimes (\partial_{x_i}^2 - D_-^i D_+^i) J_r \otimes J_{r_0}.$$

Using a Taylor expansion [25, p.24],

$$(\partial_{x_i}^2 - D_-^i D_+^i) J_r(x) = \frac{1}{6\Delta x^2} \int_0^{\Delta x} (z - \Delta x)^3 \partial_{x_i}^4 J_r(x_i + z) dz \prod_{j \neq i} J_r(x_j)$$

$$\begin{aligned}
& - \frac{1}{6\Delta x^2} \int_{-\Delta x}^0 (z + \Delta x)^3 \partial_{x_i}^4 J_r(x_i + z) dz \prod_{j \neq i} J_r(x_j) \\
& =: \partial_{x_i}(\varphi_1^i(x) + \varphi_2^i(x)).
\end{aligned}$$

Hence,

$$(\partial_{x_i}^2 - D_+^i D_+^i) \rho_{\varepsilon, r, r_0}^{\Delta x} = \partial_{x_i} (\chi(A(u_{\Delta x}); \cdot) \star J_\varepsilon \otimes (\varphi_1^i + \varphi_2^i) \otimes J_{r_0}).$$

By Young's inequality for convolutions,

$$\begin{aligned}
\|(\partial_{x_i}^2 - D_+^i D_+^i) \rho_{\varepsilon, r, r_0}^{\Delta x}\|_{L^1(\Pi_T \times \mathbb{R})} & \leq \|\chi(A(u_{\Delta x}); \cdot)\|_{L^1([0, T] \times \mathbb{R}; BV(\mathbb{R}^d))} \\
& \quad \times \|\varphi_1^i + \varphi_2^i\|_{L^1(\mathbb{R}^d)}.
\end{aligned}$$

It remains to estimate the L^1 norm of φ_1^i and φ_2^i :

$$\begin{aligned}
\|\varphi_1^i\|_{L^1(\mathbb{R}^d)} & = \frac{1}{6\Delta x^2} \int_{\mathbb{R}} \left| \int_0^{\Delta x} (z - \Delta x)^3 \partial_{x_i}^3 J_r(x_i + z) dz \right| dx_i \\
& \leq \frac{r + \Delta x}{3\Delta x^2 r^4} \|J^{(3)}\|_\infty \left| \int_0^{\Delta x} (z - \Delta x)^3 dz \right| \\
& = \frac{\|J^{(3)}\|_\infty}{12} \frac{\Delta x^2}{r^3} \left(1 + \frac{\Delta x}{r} \right).
\end{aligned}$$

A similar estimate applies to φ_2^i . □

Estimate 5.6. Let $\mathcal{T}_1$ and $\mathcal{T}_2$ be the terms from (5.8) in Lemma 5.1, that is,

$$\begin{aligned}
\mathcal{T}_1 & = \int_{\mathbb{R}} (\text{sign}_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) G'_1(\zeta) \cdot (D_+ - \nabla) \rho_{\varepsilon, r, r_0}^{\Delta x} d\zeta, \\
\mathcal{T}_2 & = \int_{\mathbb{R}} (\text{sign}_\varepsilon(\zeta - A(u)) \star J_{r_0} \otimes J_r) G'_2(\zeta) \cdot (D_- - \nabla) \rho_{\varepsilon, r, r_0}^{\Delta x} d\zeta,
\end{aligned}$$

where $G_k(A(u)) = F_k(u)$ for $j = 1, 2$. Then there exists a constant $C = C(d, J)$ such that

$$\|\mathcal{T}_k\|_{L^1(\Pi_T^{r_0})} \leq C \frac{\Delta x}{r} \left(1 + \frac{\Delta x}{r} \right) \|F'_k\|_{L^\infty(\mathbb{R}; \mathbb{R}^d)} \|A(u_{\Delta x})\|_{L^1([0, T]; BV(\mathbb{R}^d))}$$

for $k = 1, 2$.

Remark 5.3. Again the BV norm may be replaced by the L^1 norm at the cost of an extra factor r^{-1} .

Proof. Consider $\mathcal{T}_1$. We can change variables $\zeta = A(\xi)$, which yields

$$\mathcal{T}_1 = \int_{\mathbb{R}} (\text{sign}_\varepsilon(A(\xi) - A(u)) \star J_{r_0} \otimes J_r) F'_1(\zeta) \cdot (D_+ - \nabla) \rho_{\varepsilon, r, r_0}^{\Delta x} d\xi.$$

Then observe that

$$\|\mathcal{T}_1\|_{L^1(\Pi_T^{r_0})} \leq \|F'_1 \cdot (D_+ - \nabla) (\chi(A(u_{\Delta x}); \cdot) \star J_\varepsilon \otimes J_r \otimes J_{r_0})\|_{L^1(\Pi_T \times \mathbb{R})}.$$

We have

$$\begin{aligned}
& (D_+^i - \partial_{x_i}) (\chi(A(u_{\Delta x}); \cdot) \star J_\varepsilon \otimes J_r \otimes J_{r_0}) \\
& = \chi(A(u_{\Delta x}); \cdot) \star J_\varepsilon \otimes (D_+^i - \partial_{x_i}) J_r \otimes J_{r_0}.
\end{aligned}$$

By Taylor expansions,

$$\begin{aligned}
(D_+^i - \partial_{x_i}) J_r(x) & = \frac{1}{\Delta x} \int_0^{\Delta x} (\Delta x - z) \partial_{x_i}^2 J_r(x_i + z) dz \prod_{j \neq i} J_r(x_j) \\
& =: \partial_{x_i} \varphi(x).
\end{aligned}$$

By Young's inequality for convolutions,

$$\begin{aligned} & \left\| (F_1^i)' (D_+^i - \partial_{x_i}) (\chi(A(u_{\Delta x}); \cdot) \star J_\varepsilon \otimes J_r \otimes J_{r_0}) \right\|_{L^1(\Pi_T \times \mathbb{R})} \\ & \leq \left\| (F_1^i)' \right\|_{L^\infty(\mathbb{R})} \left\| \chi(A(u_{\Delta x}); \cdot) \right\|_{L^1([0, T] \times \mathbb{R}; BV(\mathbb{R}^d))} \left\| \varphi \right\|_{L^1(\mathbb{R}^d)}. \end{aligned}$$

It remains to estimate $\|\varphi\|_{L^1(\mathbb{R}^d)}$:

$$\begin{aligned} \|\varphi\|_{L^1(\mathbb{R}^d)} &= \frac{1}{\Delta x} \int_{\mathbb{R}} \left| \int_0^{\Delta x} (\Delta x - z) \partial_{x_i} J_r(x_i + z) dz \right| dx_i \\ &\leq \frac{1}{\Delta x} 2 \|J'\|_\infty \frac{r + \Delta x}{r^2} \int_0^{\Delta x} (\Delta x - z) dz \\ &= \|J'\|_\infty \frac{\Delta x}{r} \left(1 + \frac{\Delta x}{r} \right), \end{aligned}$$

from which the estimate of $\mathcal{T}_1$ follows. Similar arguments apply to $\mathcal{T}_2$. $\square$

Estimate 5.7. Consider the terms (5.4), (5.5), (5.6), and (5.7) from Lemma 5.1. Suppose $A' > \eta$ and set $B := A^{-1}$. Let

$$\begin{aligned} \mathcal{T}_1 &= \int_{\mathbb{R}} (\text{sign}_\varepsilon (\zeta - A(u_{\Delta x})) \star J_{r_0} \otimes J_r) \partial_t R_{\varepsilon, r, r_0}^{B'}(\zeta) d\zeta, \\ \mathcal{T}_2 &= \int_{\mathbb{R}} (\text{sign}_\varepsilon (\zeta - A(u)) \star J_{r_0} \otimes J_r) \partial_t R_{\varepsilon, r, r_0}^{B', \Delta x}(\zeta) d\zeta, \\ \mathcal{T}_3 &= \int_{\mathbb{R}} (\text{sign}_\varepsilon (\zeta - A(u_{\Delta x})) \star J_{r_0} \otimes J_r) \nabla \cdot R_{\varepsilon, r, r_0}^{g'}(\zeta) d\zeta, \\ \mathcal{T}_4 &= \int_{\mathbb{R}} (\text{sign}_\varepsilon (\zeta - A(u)) \star J_{r_0} \otimes J_r) \\ &\quad \times \left(D_+ \cdot R_{\varepsilon, r, r_0}^{G'_1, \Delta x}(\zeta) + D_- \cdot R_{\varepsilon, r, r_0}^{G'_2, \Delta x}(\zeta) \right) d\zeta, \end{aligned}$$

where

$$R_{\varepsilon, r, r_0}^f(\zeta) = R_\varepsilon^f(A(u), \zeta) \star J_{r_0} \otimes J_r \text{ and } R_{\varepsilon, r, r_0}^{f, \Delta x}(\zeta) = R_\varepsilon^f(A(u_{\Delta x}), \zeta) \star J_{r_0} \otimes J_r$$

for any function f , and R_ε^f is defined in equation (4.5). Then

$$\begin{aligned} \|\mathcal{T}_k\|_{L^\infty(\Pi_T^{r_0})} &\leq 8 \frac{\varepsilon}{\eta r_0} \|J'\|_\infty \text{ for } k = 1, 2, \\ \|\mathcal{T}_3\|_{L^\infty(\Pi_T^{r_0})} &\leq 8 \frac{\varepsilon}{\eta r} \|J'\|_\infty \sum_{i=1}^d \|f_i\|_{\text{Lip}}, \\ \|\mathcal{T}_4\|_{L^\infty(\Pi_T^{r_0})} &\leq 8 \frac{\varepsilon}{\eta r} \left(1 + \frac{\Delta x}{r} \right) \|J'\|_\infty \sum_{k=1}^2 \sum_{i=1}^d \|F_k^i\|_{\text{Lip}}. \end{aligned}$$

Proof. Consider $\mathcal{T}_1$. Moving the t derivative onto J_{r_0} , we have that

$$\mathcal{T}_1 = \int_{\mathbb{R}} \text{sign}_\varepsilon (\zeta - A(u_{\Delta x})) R_\varepsilon^{B'}(A(u), \zeta) d\zeta \stackrel{(u_{\Delta x}, u)}{\star} J_{r_0} \otimes J_r \otimes \partial_t J_{r_0} \otimes J_r.$$

By Lemma 5.5, equation (5.15), Lemma 5.7, and equation (5.20) with $f(z) = z$,

$$\|\mathcal{T}_1\|_{L^\infty(\Pi_T^{r_0})} \leq 4 \frac{\varepsilon}{\eta} \|J_{r_0} \otimes J_r\|_{L^1(\mathbb{R} \times \mathbb{R}^d)} \|\partial_t J_{r_0} \otimes J_r\|_{L^1(\mathbb{R} \times \mathbb{R}^d)} \leq 8 \frac{\varepsilon}{\eta r_0} \|J'\|_\infty.$$

The L^∞ bound on $\mathcal{T}_2$ follows similarly.

Let us consider $\mathcal{T}_3$:

$$\mathcal{T}_3 = \sum_{i=1}^d \int_{\mathbb{R}} \text{sign}_\varepsilon (\zeta - A(u_{\Delta x})) R_\varepsilon^{g'_i}(A(u), \zeta) d\zeta \stackrel{(u_{\Delta x}, u)}{\star} J_{r_0} \otimes J_r \otimes J_{r_0} \otimes \partial_{x_i} J_r.$$

By Lemma 5.5, equation (5.15), Lemma 5.7, and equation (5.20) with $f(z) = f_i(z)$,

$$\|\mathcal{T}_3\|_{L^\infty(\Pi_T^{r_0})} \leq 4 \frac{\varepsilon}{\eta} \sum_{i=1}^d \|f_i\|_{\text{Lip}} \|\partial_{x_i} J_r\|_{L^1(\mathbb{R}^d)} \leq 8 \frac{\varepsilon}{\eta r} \|J'\|_\infty \sum_{i=1}^d \|f_i\|_{\text{Lip}}.$$

The terms in $\mathcal{T}_4$ are estimated in the same way, but in view (5.22) we can utilize the bound

$$\|J_{r_0} \otimes D_\pm J_r\|_{L^1(\mathbb{R} \times \mathbb{R}^d)} \leq 2 \|J'\|_\infty \frac{1}{r} \left(1 + \frac{\Delta x}{r}\right).$$

□

5.4. Concluding the proof of Theorem 3.1. Recall that Q_ε , cf. (5.1), was introduced as an approximation to the contraction functional Q , cf. (1.8). Recall the basic property [12, 33]

$$(5.23) \quad |u - v| = \int_{\mathbb{R}} Q(u, v; \xi) d\xi, \quad u, v \in \mathbb{R}.$$

To argue for this relation, note that

$$Q(u, v; \xi) = |\chi(u; \xi)| + |\chi(v; \xi)| - 2\chi(u; \xi)\chi(v; \xi) = (\chi(u; \xi) - \chi(v; \xi))^2.$$

Next, observe that

$$(5.24) \quad \chi(u; \xi) - \chi(v; \xi) = \chi(u - v; \xi - v);$$

indeed, for any $S \in C_0^1(\mathbb{R})$,

$$\begin{aligned} \int_{\mathbb{R}} S'(\xi)(\chi(u; \xi) - \chi(v; \xi)) d\xi &= \int_u^v S'(\xi) d\xi = \int_0^{u-v} S'(\sigma + v) d\sigma \quad (\text{here } \sigma = \xi - v) \\ &= \int_{\mathbb{R}} S'(\sigma + v) \chi(u - v; \sigma) d\sigma \\ &= \int_{\mathbb{R}} S'(\xi) \chi(u - v; \xi - v) d\xi. \end{aligned}$$

Hence, the claim follows:

$$\int_{\mathbb{R}} (\chi(u; \xi) - \chi(v; \xi))^2 d\xi = \int_{\mathbb{R}} |\chi(u - v; \xi - v)| d\xi = |u - v|.$$

Let us quantify the approximation properties of Q_ε .

Lemma 5.8. *Let $A' \geq \eta > 0$, $B = A^{-1}$, and $f = g \circ A$. Define*

$$\begin{aligned} P &= \int_{\mathbb{R}} Q_\varepsilon(A(u), A(v); \zeta) B'(\zeta) d\zeta - |u - v|, \\ M &= \int_{\mathbb{R}} Q_\varepsilon(A(u), A(v); \zeta) g'(\zeta) d\zeta - \text{sign}(u - v) (f(u) - f(v)), \end{aligned}$$

and

$$N = \int_{\mathbb{R}} Q_\varepsilon(A(u), A(v); \zeta) d\zeta - |A(u) - A(v)|,$$

for any u and v , and where Q_ε is given by (5.1). Then

$$|P| \leq 16 \frac{\varepsilon}{\eta}, \quad |M| \leq 8 \frac{\varepsilon}{\eta}, \quad |N| \leq 8 \frac{\varepsilon}{\eta}.$$

Proof. Because $A' > 0$, $Q(u, v; \xi) = Q(A(u), A(v); A(\xi))$. Hence we can use (5.23) and a change of variables to obtain the identify

$$P = \int_{\mathbb{R}} (Q_\varepsilon(A(u), A(v); \zeta) - Q(A(u), A(v); \zeta)) B'(\zeta) d\zeta.$$

By definition of Q and the equality $|\chi(u; \xi)| = \text{sign}(\xi) \chi(u; \xi)$,

$$\begin{aligned} Q(A(u), A(v); \zeta) &= \text{sign}(\zeta) \chi(A(u); \zeta) + \text{sign}(\zeta) \chi(A(v); \zeta) \\ &\quad - 2\chi(A(u); \zeta)\chi(A(v); \zeta). \end{aligned}$$

Thus,

$$\begin{aligned} P &= \int_{\mathbb{R}} (\text{sign}_{\varepsilon}(\zeta) \chi_{\varepsilon}(A(u); \zeta) - \text{sign}(\zeta) \chi(A(u); \zeta)) B'(\zeta) d\zeta \\ &\quad + \int_{\mathbb{R}} (\text{sign}_{\varepsilon}(\zeta) \chi_{\varepsilon}(A(v); \zeta) - \text{sign}(\zeta) \chi(A(v); \zeta)) B'(\zeta) d\zeta \\ &\quad + 2 \int_{\mathbb{R}} (\chi(A(u); \zeta)\chi(A(v); \zeta) - \chi_{\varepsilon}(A(u); \zeta)\chi_{\varepsilon}(A(v); \zeta)) B'(\zeta) d\zeta \\ &=: P_1 + P_2 + P_3. \end{aligned}$$

Finding that the measure of the support of the integrand is bounded by 4ε for P_1 , P_2 , and P_3 , we conclude that

$$|P| \leq 16\varepsilon \|B'\|_{\infty},$$

and then the bound on P follows since $\|B'\|_{\infty} \leq \eta^{-1}$.

To prove the inequality for M , note that

$$\begin{aligned} &\text{sign}(u - v)(f(u) - f(v)) \\ &= \int_{\mathbb{R}} \text{sign}(u - v) (\chi(u; \zeta) - \chi(v; \zeta)) f'(\zeta) d\zeta \\ &= \int_{\mathbb{R}} |\chi(u; \zeta) - \chi(v; \zeta)| f'(\zeta) d\zeta \\ &= \int_{\mathbb{R}} [\text{sign}(\zeta) \chi(u; \zeta) + \text{sign}(\zeta) \chi(v; \zeta) - 2\chi(u; \zeta)\chi(v; \zeta)] f'(\zeta) d\zeta. \end{aligned}$$

Changing variables, we arrive at

$$\int_{\mathbb{R}} Q_{\varepsilon}(A(u), A(v); \zeta) g'(\zeta) d\zeta = \int_{\mathbb{R}} Q_{\varepsilon}(A(u), A(v); A(\zeta)) f'(\zeta) d\zeta,$$

and, since $\text{sign}(\zeta) \chi(w; \zeta) = \text{sign}(A(\zeta)) \chi(A(w); A(\zeta))$, we find that

$$\begin{aligned} |M| &\leq \int_{\mathbb{R}} \left| \text{sign}_{\varepsilon}(A(\zeta)) \chi_{\varepsilon}(A(u); A(\zeta)) - \text{sign}(A(\zeta)) \chi(A(u); A(\zeta)) \right| |f'(\zeta)| d\zeta \\ &\quad + \int_{\mathbb{R}} \left| \text{sign}_{\varepsilon}(A(\zeta)) \chi_{\varepsilon}(A(v); A(\zeta)) - \text{sign}(A(\zeta)) \chi(A(v); A(\zeta)) \right| |f'(\zeta)| d\zeta \\ &\quad + 2 \int_{\mathbb{R}} |\chi_{\varepsilon}(A(u); A(\zeta))\chi_{\varepsilon}(A(v); A(\zeta)) - \chi(A(u); A(\zeta))\chi(A(v); A(\zeta))| |f'(\zeta)| d\zeta. \end{aligned}$$

Each of the three integrands is bounded by 2 and has support where $|A(\zeta)| < \varepsilon$, i.e., where $|\zeta| \leq \varepsilon/\eta$, hence $|M| \leq 8\varepsilon/\eta$. The proof of the bound on $|N|$ is similar. $\square$

Concluding the proof of Theorem 3.1. We shall choose a positive test function $\phi \leq 1$, such that $|\nabla \phi|$ and $|\Delta \phi|$ are bounded by $C\phi$. This will be convenient when we estimate terms containing $\nabla \phi$ or $\Delta \phi$.

A test function with the necessary properties can be defined as follows, fix $R > d\sqrt{d}$ and define $\hat{\phi} : \mathbb{R}^d \rightarrow \mathbb{R}$ by

$$\hat{\phi}(x) = \begin{cases} 1 & \text{if } |x| \leq R + \sqrt{d}, \\ \exp((R + \sqrt{d} - |x|)/\sqrt{d}) & \text{otherwise.} \end{cases}$$

Define $\phi = \hat{\phi} \star J^{\otimes n}$ and note that $\phi(x) = 1$ for $x \in B(0, R)$. Note that $\hat{\phi}$ is weakly differentiable and satisfies

$$\partial_{x_i} \hat{\phi} = \begin{cases} -\frac{x_i}{\sqrt{d}|x|} \hat{\phi}(x) & |x| > R + \sqrt{d}, \\ 0 & |x| < R + \sqrt{d}. \end{cases}$$

It follows that $|\nabla \phi(x)| \leq \frac{1}{\sqrt{d}} \phi(x)$. In order to bound $\Delta \phi$ we first note that

$$\Delta \hat{\phi}(x) = \left(\frac{1}{d} - \frac{d-1}{\sqrt{d}|x|} \right) \hat{\phi}(x), \quad \text{for } |x| > R + \sqrt{d}.$$

Furthermore,

$$\frac{1}{d^2} \leq \left(\frac{1}{d} - \frac{d-1}{\sqrt{d}|x|} \right) \leq \frac{1}{d} \quad \text{for } |x| \geq d\sqrt{d}.$$

It follows that $|\Delta \hat{\phi}| \leq \frac{1}{d} \hat{\phi}(x)$ whenever $|x| > R + \sqrt{d}$. Hence

$$|\Delta \phi(x)| \leq \frac{1}{d} \phi(x) \quad \text{for } |x| > R + 2\sqrt{d}.$$

If $|x| \leq R + 2\sqrt{d}$ it follows by the lower bound $\phi(x) \geq e^{-2}$, that there exists a constant $C = C(d, J)$ such that $|\Delta \phi(x)| \leq C |\phi(x)|$.

The next lemma estimates how far $|u_{\Delta x} - u|$ is from its regularized counterpart

$$\int B'(\zeta) (\chi_\varepsilon(A(u_{\Delta x}); \zeta) - \chi_\varepsilon(A(u); \zeta))^2 d\zeta \stackrel{(u_{\Delta x}, u)}{\star} J_{r_0} \otimes J_r \otimes J_{r_0} \otimes J_r.$$

Lemma 5.9. *With the notation and assumptions of Lemma 5.1,*

$$(5.25) \quad \int_{\mathbb{R}^d} \left| \int_{\mathbb{R}} B'(\zeta) Q_{\varepsilon, r, r_0}(A(u_{\Delta x}), A(u); \zeta) d\zeta - |u_{\Delta x} - u| \phi dx \right| \leq C \left(r + r_0 + \|\phi\|_{L^1(\mathbb{R}^d)} \frac{\varepsilon}{\eta} \right),$$

$$(5.26) \quad \int_{r_0}^T \int_{\mathbb{R}^d} \left| \left[\int_{\mathbb{R}} g'(\zeta) Q_{\varepsilon, r, r_0}(A(u_{\Delta x}), A(u); \zeta) d\zeta - \text{sign}(u_{\Delta x} - u) (f(u) - f(u_{\Delta x})) \right] \cdot \nabla \phi \right| dx dt \leq CT \left(r + r_0 + \|\phi\|_{L^1(\mathbb{R}^d)} \frac{\varepsilon}{\eta} \right),$$

and

$$(5.27) \quad \int_{r_0}^T \int_{\mathbb{R}^d} \left| \left[\int_{\mathbb{R}} Q_{\varepsilon, r, r_0}(A(u_{\Delta x}), A(u); \zeta) d\zeta - |A(u_{\Delta x}) - A(u)| \right] \Delta \phi \right| dx dt \leq CT \left(r + r_0 + \|\phi\|_{L^1(\mathbb{R}^d)} \frac{\varepsilon}{\eta} \right),$$

where the constant C only depends on the initial data, A , and f .

Proof of Lemma 5.9. We establish (5.25) as follows:

$$\begin{aligned} & \int_{\mathbb{R}^d} \left| \int_{\mathbb{R}} B'(\zeta) Q_{\varepsilon, r, r_0}(A(u_{\Delta x}), A(u); \zeta) d\zeta - |u_{\Delta x} - u| \phi dx \right| \\ &= \int_{\mathbb{R}^d} \left| \int_0^T \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}} B'(\zeta) Q_\varepsilon(A(u_{\Delta x}(s, y)), A(u(s, y)); \zeta) d\zeta - |u_{\Delta x}(t, x) - u(t, x)| \right) J_{r_0}(t-s) J_r(x-y) dy ds \right| \phi dx \\ &\leq \int_{\mathbb{R}^d} \int_0^T \int_{\mathbb{R}^d} \left| \int_{\mathbb{R}} B'(\zeta) Q_\varepsilon(A(u_{\Delta x}(s, y)), A(u(s, y)); \zeta) d\zeta - |u_{\Delta x}(s, y) - u(s, y)| \right| J_{r_0}(t-s) J_r(x-y) dy ds \phi dx \end{aligned}$$

$$\begin{aligned}
& + \int_{\mathbb{R}^d} \int_0^T \int_{\mathbb{R}^d} \left[|u_{\Delta x}(t, x) - u_{\Delta x}(s, y)| + |u(t, x) - u(s, y)| \right] \\
& \quad \times J_{r_0}(t - s) J_r(x - y) dy ds dx \\
& \leq 16 \frac{\varepsilon}{\eta} \int_{\mathbb{R}^d} \phi dx + 2 \left(|u_0|_{BV(\mathbb{R}^d)} + |u_{\Delta x}(0, \cdot)|_{BV(\mathbb{R}^d)} \right) (r + r_0).
\end{aligned}$$

The bounds (5.26) and (5.27) are proved in the same way. $\square$

Writing the equation in Lemma 5.1 as

$$\int_{\mathbb{R}} B'(\zeta) \partial_t Q_{\varepsilon, r, r_0} d\zeta + \int_{\mathbb{R}} g'(\zeta) \nabla Q_{\varepsilon, r, r_0} d\zeta = \int_{\mathbb{R}} \Delta Q_{\varepsilon, r, r_0} d\zeta + \mathcal{E}_{\varepsilon, r, r_0}^{\Delta x},$$

we multiply by ϕ , integrate over $t \in [r_0, \tau]$ where $r_0 < \tau \leq T - r_0$, and integrate by parts in x , finally obtaining

$$\begin{aligned}
& \int_{\mathbb{R}^d} \int_{\mathbb{R}} B'(\zeta) Q_{\varepsilon, r, r_0} d\zeta \phi \Big|_{t=r_0}^{t=\tau} dx - \int_0^\tau \int_{\mathbb{R}^d} \int_{\mathbb{R}} g'(\zeta) Q_{\varepsilon, r, r_0} \cdot \nabla \phi d\zeta dx dt \\
& = \int_{r_0}^\tau \int_{\mathbb{R}^d} \int_{\mathbb{R}} Q_{\varepsilon, r, r_0} \Delta \phi d\zeta dx dt + \int_{r_0}^\tau \int_{\mathbb{R}^d} \mathcal{E}_{\varepsilon, r, r_0}^{\Delta x} \phi dx dt.
\end{aligned}$$

Combining this with Lemma 5.9 gives

$$\begin{aligned}
& \int_{\mathbb{R}^d} |u_{\Delta x} - u| \phi \Big|_{t=r_0}^{t=\tau} dx - \int_{r_0}^\tau \int_{\mathbb{R}^d} \text{sign}(u_{\Delta x} - u) (f(u_{\Delta x}) - f(u)) \cdot \nabla \phi dx dt \\
& \leq \int_{r_0}^\tau \int_{\mathbb{R}^d} |A(u_{\Delta x}) - A(u)| \Delta \phi dx dt + \int_{r_0}^\tau \int_{\mathbb{R}^d} \mathcal{E}_{\varepsilon, r, r_0}^{\Delta x} \phi dx dt \\
& \quad + C_T \left(r + r_0 + \frac{\varepsilon}{\eta} \right),
\end{aligned}$$

where C_T depends (linearly) on T . Using properties of ϕ , this can be rewritten as

$$\Lambda(\tau) - \Lambda(r_0) \leq C \int_{r_0}^\tau \Lambda(t) dt + \bar{\mathcal{E}}_{\varepsilon, r, r_0}^{\Delta x, \tau},$$

where

$$\begin{aligned}
\Lambda(t) &= \int_{\mathbb{R}^d} |u_{\Delta x}(t, x) - u(t, x)| \phi(x) dx, \\
\bar{\mathcal{E}}_{\varepsilon, r, r_0}^{\Delta x, \tau} &= \int_{r_0}^\tau \int_{\mathbb{R}^d} \mathcal{E}_{\varepsilon, r, r_0}^{\Delta x} \phi dx dt + C_T \left(r + r_0 + \frac{\varepsilon}{\eta} \right).
\end{aligned}$$

Gronwall's inequality then implies that

$$\Lambda(\tau) \leq \Lambda(r_0) + \tau e^{C\tau} \left(\Lambda(r_0) + \bar{\mathcal{E}}_{\varepsilon, r, r_0}^{\Delta x, \tau} \right).$$

Recall that u depends on η , and we now make this dependence explicit by writing u^η and Λ^η . Our aim is to estimate $u_{\Delta x} - u^0$. By (2.2),

$$\begin{aligned}
& \int_{B(0, R)} |u_{\Delta x}(\tau, \cdot) - u^0(\tau, \cdot)| dx dt - C\sqrt{\eta} \\
& \leq \Lambda^\eta(\tau) \\
& \leq C_T \|u_{\Delta x}(r_0, \cdot) - u^0(r_0, \cdot)\|_{L^1(\mathbb{R}^d)} + C_T \bar{\mathcal{E}}_{\varepsilon, r, r_0}^{\Delta x, \tau} \\
& \leq Cr_0 + \|u_{\Delta x}(0, \cdot) - u_0\|_{L^1(\mathbb{R}^d)} + C_T \bar{\mathcal{E}}_{\varepsilon, r, r_0}^{\Delta x, \tau}.
\end{aligned}$$

Next, we estimate the terms in the integral of $\mathcal{E}_{\varepsilon, r, r_0}^{\Delta x}$; these are the terms in (5.3)–(5.11). By Estimate 5.4,

$$(5.28) \quad \iint_{\Pi_T^{r_0}} |\text{second term in (5.3)}| \, dx dt \leq C \frac{\Delta x^2}{r^3} \left(1 + \frac{\Delta x}{r}\right),$$

where C depends on the initial data.

The integral of the terms (5.4)–(5.7) is bounded by Estimate 5.7:

$$\iint_{\Pi_T^{r_0}} |(5.4) + \dots + (5.7)| \, dx dt \leq C \frac{\varepsilon}{\eta} \left(\frac{1}{r_0} + \frac{1}{r} \left(1 + \frac{\Delta x}{r}\right) \right).$$

The integral of (5.8) is bounded by Estimate 5.6 as follows:

$$\iint_{\Pi_T^{r_0}} |(5.8)| \, dx dt \leq C \frac{\Delta x}{r} \left(1 + \frac{\Delta x}{r}\right).$$

The integral of (5.9) is bounded using Estimate 5.5:

$$\iint_{\Pi_T^{r_0}} |(5.9)| \, dx dt \leq C \frac{\Delta x^2}{r^3} \left(1 + \frac{\Delta x}{r}\right).$$

The term (5.11) is bounded using Estimates 5.1 and 5.2 (if $d > 1$):

$$(5.29a) \quad \iint_{\Pi_T^{r_0}} |(5.11)| \, dx dt \leq C \Delta x \left(\frac{1}{r^2} + \frac{1}{r} + \frac{1}{\varepsilon^2 \sqrt{r_0 r^d}} \right).$$

If $d = 1$, we can use Estimate 5.3 to achieve the better bound

$$(5.29b) \quad \iint_{\Pi_T^{r_0}} \left| \lim_{\varepsilon \rightarrow 0} (5.11) \right| \, dx dt \leq C \Delta x \left(\frac{1}{r^2} + \frac{1}{r} + \frac{1}{r_0} \right).$$

Finally, the term (5.10) is non-positive.

The fraction $\Delta x/r$ will turn out to be uniformly bounded (in fact vanishingly small), so we can overestimate it by a constant. Thus the bounds (5.28)–(5.29b) give the following estimate for $\bar{\mathcal{E}}_{\varepsilon, r, r_0}^{\Delta x, \tau}$:

$$\bar{\mathcal{E}}_{\varepsilon, r, r_0}^{\Delta x, \tau} \leq C_T \left(r + r_0 + \frac{\varepsilon}{\eta} + \frac{\varepsilon}{\eta r_0} + \frac{\varepsilon}{\eta r} + \frac{\Delta x}{r} + \frac{\Delta x}{r^2} + \frac{\Delta x}{\varepsilon^2 \sqrt{r_0 r^d}} \right).$$

If $u_0 \in BV(\mathbb{R}^d)$, $\|u_{\Delta x}(0, \cdot) - u_0\|_{L^1(\mathbb{R}^d)} \leq |u_0|_{BV(\mathbb{R}^d)} \Delta x$, so that

$$\begin{aligned} & \|u_{\Delta x}(\tau, \cdot) - u^0(\tau, \cdot)\|_{L^1(B(0, R))} \\ & \leq C_T \left(\Delta x + \sqrt{\eta} + r + r_0 + \frac{\varepsilon}{\eta} + \frac{\varepsilon}{\eta r_0} + \frac{\varepsilon}{\eta r} + \frac{\Delta x}{r} + \frac{\Delta x}{r^2} + \frac{\Delta x}{\varepsilon^2 \sqrt{r_0 r^d}} \right). \end{aligned}$$

Now we set $r = r_0 = \sqrt{\eta}$, $\varepsilon = r^4$; using that $r < 1$, the above simplifies to

$$\|u_{\Delta x}(\tau, \cdot) - u^0(\tau, \cdot)\|_{L^1(B(0, R))} \leq C_T \left(r + \frac{\Delta x}{r^{\frac{17+d}{2}}} \right).$$

Finally, minimizing with respect to r yields

$$\|u_{\Delta x}(\tau, \cdot) - u^0(\tau, \cdot)\|_{L^1(B(0, R))} \leq C_T \Delta x^{\frac{2}{19+d}}.$$

□

Remark 5.4. If $d = 1$, the above estimate gives a convergence rate of $1/10$, which is better than the rate reported in [23]. However, when $d = 1$, we can use (5.29b) instead of (5.29a). Then we have no terms with ε in the denominator, so we can send ε to zero in (5.2)–(5.11) before taking absolute values and integrating. Proceeding as above, i.e., setting $r = r_0 = \sqrt{\eta}$, this yields the bound

$$\|u_{\Delta x}(\tau, \cdot) - u^0(\tau, \cdot)\|_{L^1(B(0, R))} \leq C_T \left(r + \frac{\Delta x}{r^2} \right),$$

which gives the rate $1/3$ [25].

APPENDIX A. WELL-POSEDNESS OF DIFFERENCE METHOD

In this appendix we establish the well-posedness of the semi-discrete method. We also collect a series of a priori bounds.

Introduce

$$\|\sigma\|_1 = \Delta x^d \sum_{\alpha} |\sigma_{\alpha}| \quad \text{and} \quad |\sigma|_{BV} = \sum_{\alpha} \sum_{i=1}^d |\sigma_{\alpha+e_i} - \sigma_{\alpha}|.$$

If these quantities are bounded we say that $\sigma = \{\sigma_{\alpha}\}$ is in $\ell^1(\mathbb{Z}^d)$ and of bounded variation. Let $u(t) = \{u_{\alpha}(t)\}_{\alpha \in \mathbb{Z}^d}$ and $u_0 = \{u_{\alpha}(0)\}_{\alpha \in \mathbb{Z}^d}$ and define the operator $\mathcal{A} : \ell^1(\mathbb{Z}^d) \rightarrow \ell^1(\mathbb{Z}^d)$ by

$$(\mathcal{A}(u))_{\alpha} = \sum_{i=1}^d D_{-}^i [F^i(u_{\alpha}, u_{\alpha+e_i}) - D_{+}^i A(u_{\alpha})].$$

Then (3.1) may be considered as the Cauchy problem

$$\begin{cases} \frac{du}{dt} + \mathcal{A}(u) = 0, & t > 0, \\ u(0) = u_0. \end{cases}$$

This problem has a unique continuously differentiable solution for small t , since $\mathcal{A}$ is Lipschitz continuous for each $\Delta x > 0$. The solution defines a strongly continuous semigroup $\mathcal{S}(t)$ on ℓ^1 . We want to show that this semigroup is ℓ^1 contractive. This follows by the theory presented in [14], given that $\mathcal{A}$ is accretive, i.e.,

$$\sum_{\alpha} \text{sign}(u_{\alpha} - v_{\alpha}) (\mathcal{A}(u) - \mathcal{A}(v))_{\alpha} \geq 0.$$

for any u and v in $\ell^1(\mathbb{Z}^d)$ [32, 34].

Lemma A.1. *The operator $\mathcal{A} : \ell^1(\mathbb{Z}^d) \rightarrow \ell^1(\mathbb{Z}^d)$ is accretive.*

Proof. By definition

$$(\mathcal{A}(u) - \mathcal{A}(v))_{\alpha} = \sum_{i=1}^d D_{-}^i [F^i(u_{\alpha}, u_{\alpha+e_i}) - F^i(v_{\alpha}, v_{\alpha+e_i}) - D_{+}^i (A(u_{\alpha}) - A(v_{\alpha}))].$$

Let $\partial_1 F^i$ and $\partial_2 F^i$ denote the partial derivatives of F^i with respect to the first and second variable respectively. Since F^i is continuously differentiable there exist for each (α, i) some number $\tau_{\alpha, i}$ such that

$$F^i(u_{\alpha}, u_{\alpha+e_i}) - F^i(v_{\alpha}, u_{\alpha+e_i}) = \partial_1 F^i(\tau_{\alpha, i}, u_{\alpha+e_i})(u_{\alpha} - v_{\alpha})$$

and similarly a number $\theta_{\alpha, i}$ such that

$$F^i(v_{\alpha}, u_{\alpha+e_i}) - F^i(v_{\alpha}, v_{\alpha+e_i}) = \partial_2 F^i(v_{\alpha}, \theta_{\alpha, i})(u_{\alpha+e_i} - v_{\alpha+e_i}).$$

Let $w_{\alpha} = u_{\alpha} - v_{\alpha}$ then

$$F^i(u_{\alpha}, u_{\alpha+e_i}) - F^i(v_{\alpha}, v_{\alpha+e_i})$$

$$\begin{aligned}
&= F^i(u_\alpha, u_{\alpha+e_i}) - F^i(v_\alpha, u_{\alpha+e_i}) + F^i(v_\alpha, u_{\alpha+e_i}) - F^i(v_\alpha, v_{\alpha+e_i}) \\
&= \partial_1 F^i(\tau_{\alpha,i}, u_{\alpha+e_i}) w_\alpha + \partial_2 F^i(v_\alpha, \theta_{\alpha,i}) w_{\alpha+e_i}.
\end{aligned}$$

Let $A' = a$. Then there exist some ξ_α such that

$$A(u_\alpha) - A(v_\alpha) = a(\xi_\alpha) w_\alpha.$$

Using these expressions we obtain

$$\begin{aligned}
&\sum_{\alpha} \text{sign}(u_\alpha - v_\alpha) (\mathcal{A}(u) - \mathcal{A}(v))_\alpha \\
&= \sum_{\alpha} \sum_{i=1}^d \text{sign}(w_\alpha) D_-^i [\partial_1 F^i(\tau_{\alpha,i}, u_{\alpha+e_i}) w_\alpha + \partial_2 F^i(v_\alpha, \theta_{\alpha,i}) w_{\alpha+e_i}] \\
&\quad - \sum_{\alpha} \sum_{i=1}^d \text{sign}(w_\alpha) D_-^i D_+^i (a(\xi_\alpha) w_\alpha) := \mathcal{T}_1 - \mathcal{T}_2.
\end{aligned}
\tag{A.1}$$

Consider $\mathcal{T}_1$ first. Since

$$\begin{aligned}
D_-^i [\partial_1 F^i(\tau_{\alpha,i}, u_{\alpha+e_i}) w_\alpha + \partial_2 F^i(v_\alpha, \theta_{\alpha,i}) w_{\alpha+e_i}] &= \frac{1}{\Delta x} [\partial_1 F^i(\tau_{\alpha,i}, u_{\alpha+e_i}) w_\alpha \\
&\quad - \partial_1 F^i(\tau_{\alpha-e_i,i}, u_\alpha) w_{\alpha-e_i} + \partial_2 F^i(v_\alpha, \theta_{\alpha,i}) w_{\alpha+e_i} - \partial_2 F^i(v_{\alpha-e_i}, \theta_{\alpha-e_i,i}) w_\alpha],
\end{aligned}$$

it follows that

$$\begin{aligned}
\mathcal{T}_1 &= \frac{1}{\Delta x} \sum_{\alpha} \sum_{i=1}^d \left[\partial_1 F^i(\tau_{\alpha,i}, u_{\alpha+e_i}) |w_\alpha| - \partial_1 F^i(\tau_{\alpha-e_i,i}, u_\alpha) \text{sign}(w_\alpha) w_{\alpha-e_i} \right. \\
&\quad \left. + \partial_2 F^i(v_\alpha, \theta_{\alpha,i}) \text{sign}(w_\alpha) w_{\alpha+e_i} - \partial_2 F^i(v_{\alpha-e_i}, \theta_{\alpha-e_i,i}) |w_\alpha| \right] \\
&= \frac{1}{\Delta x} \sum_{i=1}^d \left[\sum_{\alpha} \partial_1 F^i(\tau_{\alpha,i}, u_{\alpha+e_i}) |w_\alpha| - \sum_{\alpha} \partial_1 F^i(\tau_{\alpha,i}, u_{\alpha+e_i}) \text{sign}(w_{\alpha+e_i}) w_\alpha \right. \\
&\quad \left. + \sum_{\alpha} \partial_2 F^i(v_\alpha, \theta_{\alpha,i}) \text{sign}(w_\alpha) w_{\alpha+e_i} - \sum_{\alpha} \partial_2 F^i(v_\alpha, \theta_{\alpha,i}) |w_{\alpha+e_i}| \right]
\end{aligned}$$

Since each F^i is monotone, it follows that $\mathcal{T}_1 \geq 0$. Considering $\mathcal{T}_2$, we have

$$\begin{aligned}
\mathcal{T}_2 &= \frac{1}{\Delta x^2} \sum_{i=1}^d \sum_{\alpha} \left[a(\xi_{\alpha+e_i}) \text{sign}(w_\alpha) w_{\alpha+e_i} \right. \\
&\quad \left. - 2a(\xi_\alpha) |w_\alpha| + a(\xi_{\alpha-e_i}) \text{sign}(w_\alpha) w_{\alpha-e_i} \right],
\end{aligned}$$

from which it follows that $\mathcal{T}_2 \leq 0$. $\square$

Lemma A.2. *Suppose F^i is monotone for each $1 \leq i \leq d$. For any positive T , there exists a unique solution $u = \{u_\alpha\}$ to (3.1) on $[0, T]$ with the properties:*

(i) $\|u(t)\|_1 \leq \|u_0\|_1$.

(ii) For every $\alpha \in \mathbb{Z}^d$ and $t \in [0, T]$,

$$\inf_{\beta} \{u_{\beta,0}\} \leq u_\alpha(t) \leq \sup_{\beta} \{u_{\beta,0}\}.$$

(iii) $|u(t)|_{BV} \leq |u_0|_{BV}$.

(iv) If $v = \{v_\alpha\}$ is a solution of the same problem with initial data v_0 , then

$$\|u(t) - v(t)\|_1 \leq \|u_0 - v_0\|_1.$$

Proof. Parts (i), (iii) and (iv) follows since $\mathcal{S}(t)$ is a contraction semigroup. Part (ii) follows from [9]. $\square$

Note that the ℓ^1 bound in [(i)] implies that $u_\alpha(t)$ exists for all t , and not only for t small.

Lemma A.3. *Suppose F^i is monotone for each $1 \leq i \leq d$. If u is a solution to (3.1) and $\mathcal{A}(u_0) \in \ell^1(\mathbb{Z}^d)$, then for each $h > 0$,*

$$\|u(t+h) - u(t)\|_{\ell^1} \leq \|\mathcal{A}(u_0)\|_{\ell^1} h.$$

Proof. Suppose that $\|u'(t)\| \leq C$. Then

$$\|u(t+h) - u(t)\| = \left\| \int_t^{t+h} u'(s) ds \right\| \leq \int_t^{t+h} \|u'(s)\| ds \leq Ch,$$

and so Lipschitz continuity would follow. We claim that

$$(A.2) \quad \frac{\partial}{\partial t} \|u'(t)\| \leq 0.$$

Indeed,

$$\begin{aligned} \frac{\partial}{\partial t} \|u'(t)\| &= \frac{\partial}{\partial t} \|\mathcal{A}(u(t))\| \\ &= \frac{\partial}{\partial t} \left[\Delta x^d \sum_{\alpha} \text{sign}(\mathcal{A}(u(t))_{\alpha}) \mathcal{A}(u(t))_{\alpha} \right] \\ &= \Delta x^d \sum_{\alpha} \text{sign}(\mathcal{A}(u(t))_{\alpha}) \partial_t \mathcal{A}(u(t))_{\alpha}, \end{aligned}$$

and

$$\begin{aligned} \partial_t \mathcal{A}(u(t))_{\alpha} &= \frac{\partial}{\partial t} \sum_{i=1}^d D_-^i [F^i(u_{\alpha}(t), u_{\alpha+e_i}(t)) - D_+^i \mathcal{A}(u(t))_{\alpha}] \\ &= \sum_{i=1}^d D_-^i [\partial_1 F^i(u_{\alpha}(t), u_{\alpha+e_i}(t)) u'_{\alpha}(t) + \partial_2 F^i(u_{\alpha}(t), u_{\alpha+e_i}(t)) u'_{\alpha+e_i}(t)] \\ &\quad - \sum_{i=1}^d D_-^i D_+^i a(u_{\alpha}(t)) u'_{\alpha}(t) \\ &= - \sum_{i=1}^d D_-^i [\partial_1 F^i(u_{\alpha}(t), u_{\alpha+e_i}(t)) \mathcal{A}(u(t))_{\alpha} + \partial_2 F^i(u_{\alpha}(t), u_{\alpha+e_i}(t)) \mathcal{A}(u(t))_{\alpha+e_i}] \\ &\quad + \sum_{i=1}^d D_-^i D_+^i a(u_{\alpha}(t)) \mathcal{A}(u(t))_{\alpha}. \end{aligned}$$

Considering the similarity between this computation and (A.1), it is seen that (A.2) holds. We conclude that $\|u'(t)\| \leq \|\mathcal{A}(u_0)\|$ and so the lemma follows. $\square$

REFERENCES

- [1] B. Andreianov, M. Bendahmane, and K. H. Karlsen. Discrete duality finite volume schemes for doubly nonlinear degenerate hyperbolic-parabolic equations. *J. Hyperbolic Differ. Equ.*, 7(1):1–67, 2010.
- [2] B. Andreianov and N. Igibida. On uniqueness techniques for degenerate convection-diffusion problems. *Int. J. Dyn. Syst. Differ. Equ.*, 4(1-2):3–34, 2012.
- [3] D. Aregba-Driollet, R. Natalini, and S. Tang. Explicit diffusive kinetic schemes for nonlinear degenerate parabolic systems. *Math. Comp.*, 73(245):63–94 (electronic), 2004.
- [4] G. Barles and E. R. Jakobsen. Error bounds for monotone approximation schemes for Hamilton-Jacobi-Bellman equations. *SIAM J. Numer. Anal.*, 43(2):540–558 (electronic), 2005.
- [5] F. Bouchut, F. R. Guarguaglini, and R. Natalini. Diffusive BGK approximations for nonlinear multidimensional parabolic equations. *Indiana Univ. Math. J.*, 49(2):723–749, 2000.

- [6] F. Bouchut and B. Perthame. Kružkov's estimates for scalar conservation laws revisited. *Trans. Amer. Math. Soc.*, 350(7):2847–2870, 1998.
- [7] L. A. Caffarelli and P. E. Souganidis. A rate of convergence for monotone finite difference approximations to fully nonlinear, uniformly elliptic PDEs. *Comm. Pure Appl. Math.*, 61(1):1–17, 2008.
- [8] J. Carrillo. Entropy solutions for nonlinear degenerate problems. *Arch. Ration. Mech. Anal.*, 147(4):269–361, 1999.
- [9] A. Chambolle and B. J. Lucier. Un principe du maximum pour des opérateurs monotones. *C. R. Acad. Sci. Paris Sér. I Math.*, 326(7):823–827, 1998.
- [10] G.-Q. Chen and K. H. Karlsen. Quasilinear anisotropic degenerate parabolic equations with time-space dependent diffusion coefficients. *Commun. Pure Appl. Anal.*, 4(2):241–266, 2005.
- [11] G.-Q. Chen and K. H. Karlsen. L^1 -framework for continuous dependence and error estimates for quasilinear anisotropic degenerate parabolic equations. *Trans. Amer. Math. Soc.*, 358(3):937–963, 2006.
- [12] G.-Q. Chen and B. Perthame. Well-posedness for non-isotropic degenerate parabolic-hyperbolic equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 20(4):645–668, 2003.
- [13] B. Cockburn. Continuous dependence and error estimation for viscosity methods. *Acta Numer.*, 12:127–180, 2003.
- [14] M. G. Crandall and T. M. Liggett. Generation of semi-groups of nonlinear transformations on general Banach spaces. *Amer. J. Math.*, 93:265–298, 1971.
- [15] M. G. Crandall and P.-L. Lions. Two approximations of solutions of Hamilton-Jacobi equations. *Math. Comp.*, 43(167):1–19, 1984.
- [16] C. M. Dafermos. *Hyperbolic conservation laws in continuum physics*, volume 325 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, third edition, 2010.
- [17] S. Evje and K. H. Karlsen. Discrete approximations of BV solutions to doubly nonlinear degenerate parabolic equations. *Numer. Math.*, 86(3):377–417, 2000.
- [18] S. Evje and K. H. Karlsen. Monotone difference approximations of BV solutions to degenerate convection-diffusion equations. *SIAM J. Numer. Anal.*, 37(6):1838–1860 (electronic), 2000.
- [19] S. Evje and K. H. Karlsen. An error estimate for viscous approximate solutions of degenerate parabolic equations. *J. Nonlinear Math. Phys.*, 9(3):262–281, 2002.
- [20] R. Eymard, T. Gallouët, and R. Herbin. Error estimate for approximate solutions of a nonlinear convection-diffusion problem. *Adv. Differential Equations*, 7(4):419–440, 2002.
- [21] R. Eymard, T. Gallouët, R. Herbin, and A. Michel. Convergence of a finite volume scheme for nonlinear degenerate parabolic equations. *Numer. Math.*, 92(1):41–82, 2002.
- [22] H. Holden, K. H. Karlsen, K.-A. Lie, and N. H. Risebro. *Splitting methods for partial differential equations with rough solutions*. EMS Series of Lectures in Mathematics. European Mathematical Society (EMS), Zürich, 2010. Analysis and MATLAB programs.
- [23] K. H. Karlsen, U. Koley, and N. H. Risebro. An error estimate for the finite difference approximation to degenerate convection-diffusion equations. *Numer. Math.*, 121(2):367–395, 2012.
- [24] K. H. Karlsen and N. H. Risebro. Convergence of finite difference schemes for viscous and inviscid conservation laws with rough coefficients. *M2AN Math. Model. Numer. Anal.*, 35(2):239–269, 2001.
- [25] K. H. Karlsen, N. H. Risebro, and E. B. Storrøsten. L^1 error estimates for difference approximations of degenerate convection-diffusion equations. *Math. Comp.*, 83(290):2717–2762, 2014.
- [26] S. N. Kružkov. First order quasilinear equations with several independent variables. *Mat. Sb. (N.S.)*, 81 (123):228–255, 1970.
- [27] N. V. Krylov. The rate of convergence of finite-difference approximations for Bellman equations with Lipschitz coefficients. *Appl. Math. Optim.*, 52(3):365–399, 2005.
- [28] N. N. Kuznecov. The accuracy of certain approximate methods for the computation of weak solutions of a first order quasilinear equation. *Ž. Vyčisl. Mat. i Mat. Fiz.*, 16(6):1489–1502, 1627, 1976.
- [29] P.-L. Lions, B. Perthame, and E. Tadmor. A kinetic formulation of multidimensional scalar conservation laws and related equations. *J. Amer. Math. Soc.*, 7(1):169–191, 1994.
- [30] C. Makridakis and B. Perthame. Optimal rate of convergence for anisotropic vanishing viscosity limit of a scalar balance law. *SIAM J. Math. Anal.*, 34(6):1300–1307, 2003.
- [31] M. Ohlberger. A posteriori error estimates for vertex centered finite volume approximations of convection-diffusion-reaction equations. *M2AN Math. Model. Numer. Anal.*, 35(2):355–387, 2001.
- [32] N. H. Pavel. *Differential equations, flow invariance and applications*, volume 113 of *Research Notes in Mathematics*. Pitman (Advanced Publishing Program), Boston, MA, 1984.

- [33] B. Perthame. Uniqueness and error estimates in first order quasilinear conservation laws via the kinetic entropy defect measure. *J. Math. Pures Appl. (9)*, 77(10):1055–1064, 1998.
- [34] K. Sato. On the generators of non-negative contraction semigroups in Banach lattices. *J. Math. Soc. Japan*, 20:423–436, 1968.
- [35] A. I. Vol'pert and S. I. Hudjaev. The Cauchy problem for second order quasilinear degenerate parabolic equations. *Mat. Sb. (N.S.)*, 78 (120):374–396, 1969.
- [36] Z. Q. Wu and J. X. Yin. Some properties of functions in BV_x and their applications to the uniqueness of solutions for degenerate quasilinear parabolic equations. *Northeast. Math. J.*, 5(4):395–422, 1989.

(Kenneth Hvistendahl Karlsen) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSLO, P.O. BOX 1053, BLINDERN, N-0316 OSLO, NORWAY.

E-mail address: `kennethk@math.uio.no`

(Nils Henrik Risebro) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSLO, P.O. BOX 1053, BLINDERN, N-0316 OSLO, NORWAY.

E-mail address: `nilshr@math.uio.no`

(Erlend Briseid Storrøsten) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSLO, P.O. BOX 1053, BLINDERN, N-0316 OSLO, NORWAY.

E-mail address: `erlenbs@math.uio.no`

ON STOCHASTIC CONSERVATION LAWS AND MALLIAVIN CALCULUS

K. H. KARLSEN AND E. B. STORRØSTEN

ABSTRACT. For stochastic conservation laws driven by a semilinear noise term, we propose a generalization of the Kružkov entropy condition by allowing the Kružkov constants to be Malliavin differentiable random variables. Existence and uniqueness results are provided. Our approach sheds some new light on the stochastic entropy conditions put forth by Feng and Nualart [17] and Bauzet, Vallet, and Wittbold [3], and in our view simplifies some of the proofs.

CONTENTS

1. Introduction	1
2. Entropy solutions	6
3. The viscous approximation	13
3.1. A priori estimates and well-posedness	13
3.2. Malliavin differentiability	23
4. Existence of entropy solutions	27
5. Uniqueness of entropy solutions	32
6. Appendix	43
6.1. Some “doubling of variables” tools	43
6.2. Weighted L^p spaces.	45
6.3. A version of Itô’s formula	47
6.4. The Lebesgue-Bochner space	49
6.5. Young measures	51
6.6. Weak compactness in L^1 .	51
References	52

1. INTRODUCTION

Stochastic partial differential equations (stochastic PDEs) arise in many fields, such as biology, physics, engineering, and economics, in which random phenomena play a crucial role. Complex systems always contain some element of uncertainty. Uncertainty may arise in the system parameters, initial and boundary conditions, and external forcing processes. Moreover, in many situations there is incomplete or partial understanding of the governing physical laws, and many models are therefore best formulated using stochastic PDEs.

Recently there has been an interest in studying the effect of stochastic forcing on nonlinear conservation laws [3, 4, 9, 14, 21, 20, 17, 36, 35], with particular emphasis on existence and uniqueness questions (well-posedness). Deterministic conservation laws possess discontinuous (shock) solutions, and a weak formulation coupled with

Date: August 18, 2015.

2010 Mathematics Subject Classification. Primary: 60H15, 35L60; Secondary: 35L65, 60G15.

Key words and phrases. Stochastic conservation law, entropy condition, Malliavin calculus, anticipating stochastic integral, existence, uniqueness.

an appropriate entropy condition is required to ensure the well-posedness [23, 26]. The question of well-posedness gets somewhat more difficult by adding a stochastic source term, due to the interaction between noise and nonlinearity.

In a different direction, we also mention the recent works [24, 25] by Lions, Perthame, and Souganidis on conservation laws with (rough) stochastic fluxes.

To be more precise, we are interested in stochastic conservation laws driven by Gaussian noise of the following form:

$$(1.1) \quad \begin{cases} du(t, x) + \nabla \cdot f(u(t, x))dt = \int_Z \sigma(x, u(t, x), z)W(dt, dz), & (t, x) \in \Pi_T, \\ u(0, x) = u^0(x), & x \in \mathbb{R}^d, \end{cases}$$

where $\Pi_T = (0, T) \times \mathbb{R}^d$, $T > 0$ is some fixed final time, $u^0 = u^0(x, \omega)$ is a given $\mathcal{F}_0$ -measurable random function, and the unknown $u = u(t, x, \omega)$ is a random (scalar) function. The flux function

$$(\mathcal{A}_f) \quad f : \mathbb{R} \rightarrow \mathbb{R}^d \text{ is assumed to be Lipschitz and } C^1(\mathbb{R}).$$

Concerning the source term, we let $Z \subseteq \mathbb{R}^n$ for some $n \geq 1$ and W be a space-time Gaussian white noise martingale random measure with respect to the filtration $\{\mathcal{F}_t\}_{0 \leq t \leq T}$ [39]. Its covariance measure is given by $dt \otimes d\mu$, where dt denotes Lebesgue measure on $[0, T]$ and μ is some deterministic σ -finite Borel measure on Z , that is, for measurable sets $A, B \subset Z$

$$E[W(t, A)W(t, B)] = t\mu(A \cap B).$$

The noise coefficient $\sigma : \mathbb{R}^d \times \mathbb{R} \times Z \rightarrow \mathbb{R}$ is a measurable function satisfying

$$(\mathcal{A}_\sigma) \quad \exists M \in L^2(Z) \text{ s.t. } \begin{cases} |\sigma(x, u, z) - \sigma(x, v, z)| \leq |u - v| M(z), \\ |\sigma(x, u, z)| \leq M(z)(1 + |u|), \end{cases}$$

for all $(x, z) \in \mathbb{R}^d \times Z$. Note that W induces a cylindrical Wiener process (with identity covariance operator) on $L^2(Z) = L^2(Z, \mathcal{B}(Z), \mu)$ which we also denote by W [30, § 7.1.2]. Let $G(u) : L^2(Z) \rightarrow L^2(\mathbb{R}^d, \phi)$ be defined by

$$(1.2) \quad G(u)h(x) = \int_Z \sigma(x, u(x), z)h(z) d\mu(z).$$

By $(\mathcal{A}_\sigma)$, G is a Lipschitz map from $L^2(\mathbb{R}^d, \phi)$ into $\mathcal{L}_2(L^2(Z), L^2(\mathbb{R}^d, \phi))$ with $\|G\|_{\text{Lip}} \leq \|M\|_{L^2(Z)}$, where $\mathcal{L}_2(L^2(Z), L^2(\mathbb{R}^d, \phi))$ denotes the space of Hilbert-Schmidt operators from $L^2(Z)$ into $L^2(\mathbb{R}^d, \phi)$, cf. Section 2. In this setting, (1.1) may be written as

$$du + \nabla \cdot f(u)dt = G(u)dW(t),$$

where the right-hand side is interpreted with respect to the cylindrical Wiener process [14]. In what follows, we will in general stick to the σ notation. We refer to [13] for a comparison of the stochastic integrals. The Malliavin calculus used later is developed with respect to the isonormal Gaussian process $W : H \rightarrow L^2(\Omega)$ defined by

$$(1.3) \quad W(h) = \int_0^T \int_Z h(s, z)W(ds, dz) = \int_0^T h(s)dW(s),$$

where H denotes the space $L^2([0, T] \times Z, \mathcal{B}([0, T]) \otimes \mathcal{B}(Z), dt \otimes d\mu)$. Concerning the notation and basic theory of Malliavin calculus we refer to [27].

When the noise term in (1.1) is additive (σ is independent of u), Kim [21] used Kružkov's entropy condition and proved the well-posedness of entropy solutions, see also Vallet and Wittbold [36]. When the noise term is additive, a change of variable turns (1.1) into a conservation law with random flux function and well-known "deterministic" techniques apply.

When the noise is multiplicative (i.e., σ depends on u), a simple adaptation of Kruřkov’s techniques fails to capture a specific “noise-noise” interaction term correlating two entropy solutions, and as a consequence they do not lead to the L^1 -contraction principle. This issue was resolved by Feng and Nualart [17], introducing an additional condition capturing the missing noise-noise interaction. These authors employ the Kruřkov entropy condition (on Itô form)

$$(1.4) \quad \begin{aligned} & \partial_t |u - c| + \partial_x [\text{sign}(u - c)(f(u) - f(c))] \\ & \leq \frac{1}{2} \text{sign}'(u - c) \sigma(u)^2 + \sigma(u) \partial_t W, \quad \forall c \in \mathbb{R}, \end{aligned}$$

which is understood in the distributional sense (and via an approximation of $\text{sign}(\cdot)$). Here, for the sake of simplicity, we take W to be an ordinary Brownian motion (Z is a point) and $d = 1$. The above family of inequalities, indexed over the “Kruřkov” constants c , is in [17] augmented with an additional condition related to certain substitution formulas [27, § 3.2.4], which enables the authors to recover the above mentioned interaction term and thus provide, for the first time, a general uniqueness result for stochastic conservation laws.

The additional condition proposed in [17] is rather technical and difficult to comprehend at first glance. Furthermore, the existence proof (passing to the limit in a sequence vanishing viscosity approximations) becomes increasingly difficult, with several added arguments revolving around fractional Sobolev spaces, estimates of the moments of increments, modulus of continuity of Itô processes, and the Garcia-Rodemich-Rumsey lemma.

Recently, Bauzet, Vallet, and Wittbold [3] provided a framework that uses the Kruřkov entropy inequalities (1.4) but bypasses the Feng-Nualart condition. Rather than comparing two entropy solutions directly, their uniqueness result compares the entropy solution against the vanishing viscosity solution, which is generated as the weak limit (as captured by the Young measure) of a sequence of solutions to stochastic parabolic equations with vanishing viscosity parameter. Although with this approach the existence proof becomes simple, many technical difficulties are added to the uniqueness proof. At this point, let us mention that Debussche and Vovelle [14] have provided an alternative well-posedness theory based on a kinetic formulation. The kinetic formulation avoids some of the difficulties alluded to above, thanks to the so-called entropy defect measure.

The purpose of our work is to propose a slight modification of the Kruřkov entropy condition (1.4) that will shed some new light on [17], and also [3]. To this end, we recall that a uniqueness proof for entropy solutions is based on a technique known as “doubling of variables”. Suppose that v is another entropy solution of (1.1) with initial condition v^0 . The key idea is to consider v as a function of a different set of variables, say $v = v(s, y)$, and then for each fixed $(s, y) \in \Pi_T$, take $c = v(s, y)$ in the entropy condition for u . In the case that u and v are stochastic fields, $v(s, y)$ is no longer a constant, but rather a random variable. Hence it seems natural to utilize an entropy condition in which the Kruřkov parameters c in (1.4) are random variables rather than constants.

Let us do an informal derivation of an entropy condition based on this idea. As above, we let W be an ordinary Brownian motion and $d = 1$. For each fixed $\varepsilon > 0$, suppose u^ε is a sufficiently regular solution of the stochastic parabolic equation

$$\partial_t u^\varepsilon + \partial_x f(u^\varepsilon) = \sigma(u^\varepsilon) \partial_t W + \varepsilon \partial_x^2 u^\varepsilon,$$

where the time derivative is understood in the sense of distributions. We apply the *anticipating* Itô formula (Theorem 6.7) to $|u^\varepsilon - V|$, with V being an arbitrary

Malliavin differentiable random variable. Taking expectations, we obtain

$$\begin{aligned} & E [\partial_t |u^\varepsilon - V| + \partial_x(\text{sign}(u^\varepsilon - V)(f(u^\varepsilon) - f(V)))] \\ & + E [\text{sign}'(u^\varepsilon - V) \sigma(u^\varepsilon) D_t V] - \frac{1}{2} E [\text{sign}'(u^\varepsilon - V) \sigma^2(u^\varepsilon)] \\ & = \varepsilon E [\text{sign}(u^\varepsilon - V) \partial_x^2 u^\varepsilon], \end{aligned}$$

where $D_t V$ is the Malliavin derivative of V at t . As

$$\varepsilon E [\text{sign}(u^\varepsilon - V) \partial_x^2 u^\varepsilon] \leq \varepsilon E [\partial_x^2 |u^\varepsilon - V|],$$

it follows that

$$\begin{aligned} & E [\partial_t |u^\varepsilon - V| + \partial_x(\text{sign}(u^\varepsilon - V)(f(u^\varepsilon) - f(V)))] \\ & + E [\text{sign}'(u^\varepsilon - V) \sigma(u^\varepsilon) D_t V] - \frac{1}{2} E [\text{sign}'(u^\varepsilon - V) \sigma^2(u^\varepsilon)] \\ & \leq \varepsilon E [\partial_x^2 |u^\varepsilon - V|]. \end{aligned}$$

Suppose $u^\varepsilon \rightarrow u$ in a suitable sense as $\varepsilon \downarrow 0$. Then the limit u ought to satisfy

$$\begin{aligned} & E [\partial_t |u - V| + \partial_x(\text{sign}(u - V)(f(u) - f(V)))] \\ (1.5) \quad & + E [\text{sign}'(u - V) \sigma(u) D_t V] - \frac{1}{2} E [\text{sign}'(u - V) \sigma^2(u)] \leq 0, \end{aligned}$$

which is the entropy condition that we propose should replace (1.4).

At least informally, it is easy to see why this entropy condition implies the L^1 contraction principle. Let $u = u(t, x)$ and $v = v(s, y)$ be two solutions satisfying the entropy condition (1.5). Suppose u, v are both Malliavin differentiable and spatially regular. The entropy condition for u yields

$$\begin{aligned} & E [\partial_t |u - v| + \partial_x(\text{sign}(u - v)(f(u) - f(v)))] \\ & + E [\text{sign}'(u - v) \sigma(u) D_t v] - \frac{1}{2} E [\text{sign}'(u - v) \sigma^2(u)] \leq 0. \end{aligned}$$

Similarly, the entropy condition of v yields

$$\begin{aligned} & E [\partial_s |v - u| + \partial_y(\text{sign}(v - u)(f(v) - f(u)))] \\ & + E [\text{sign}'(v - u) \sigma(v) D_s u] - \frac{1}{2} E [\text{sign}'(v - u) \sigma^2(v)] \leq 0. \end{aligned}$$

Suppose that $t > s$. Then $D_t v(s) = 0$ as v is adapted (to the underlying filtration). Adding the last two equations we obtain

$$\begin{aligned} & E [(\partial_t + \partial_s) |u - v| + (\partial_x + \partial_y)(\text{sign}(u - v)(f(u) - f(v)))] \\ & + E [\text{sign}'(u - v) \sigma(v) D_s u] - \frac{1}{2} E [\text{sign}'(u - v) (\sigma^2(u) + \sigma^2(v))] \leq 0. \end{aligned}$$

Completing the square yields

$$\begin{aligned} (1.6) \quad & E [(\partial_t + \partial_s) |u - v| + (\partial_x + \partial_y)(\text{sign}(u - v)(f(u) - f(v)))] \\ & + E [\text{sign}'(u - v) \sigma(v) (D_s u - \sigma(u))] - \underbrace{\frac{1}{2} E [\text{sign}'(u - v) (\sigma(u) - \sigma(v))^2]}_{=0} \leq 0. \end{aligned}$$

Next we write

$$D_s u(t) - \sigma(u(t)) = (D_s u(t) - \sigma(u(s))) + (\sigma(u(s)) - \sigma(u(t))),$$

and attempt to send $t \downarrow s$. The second term tends to zero almost everywhere. Formally, for fixed s , we observe that $D_s u(t)$ satisfies the initial value problem

$$(1.7) \quad \begin{cases} dw + \partial_x(f'(u)w) dt = (\sigma'(u)w) dW(t), & (t > s), \\ w(s) = \sigma(u(s)). \end{cases}$$

And so, concluding that

$$D_s u(t) \rightarrow \sigma(u(s)), \quad \text{as } t \downarrow s,$$

amounts to showing that the solution to (1.7) satisfies the initial condition (in some weak sense). Given this result, the L^1 contraction property follows from (1.6) in a standard way:

$$(1.8) \quad \frac{d}{dt} E \left[\|u(t) - v(t)\|_{L^1(\mathbb{R})} \right] \leq 0.$$

The above argument is hampered by an obstacle; namely, that the Malliavin differentiability of an entropy solution seems hard to establish. This can be seen related to the *discontinuous* coefficient $f'(u)$ in the stochastic continuity equation (1.7), making it difficult to establish the existence of a (properly defined) weak solution. In the deterministic context, continuity (and related transport) equations with low-regularity coefficients have been an active area of research in recent years, see for example [1, 5, 6] (and also [18] for a particular stochastic setting). Continuity equations arise in many applications, such as fluid mechanics. They also appear naturally when one linearizes a nonlinear conservation law $u_t + f(u)_x = 0$ into $w_t + (f'(u)w)_x = 0$, see references in [1, 5, 6]. The present work shows that stochastic continuity equations arise naturally as well, through linearisation (by the Malliavin derivative) of stochastic conservation laws driven by semilinear noise. However, the study of such equations is beyond the present paper and is left for future work.

As alluded to above, to make the L^1 contraction argument rigorous we would need to know that at least one of the two entropy solutions being compared, is Malliavin differentiable. To avoid this nontrivial issue, we shall employ a more indirect approach, motivated by [3], comparing one entropy solution against the solution of the viscous problem linked to the other entropy solution, relying on weak compactness in the space of Young measures for the viscous approximation. The Malliavin differentiability of the viscous solution is then established and its Malliavin derivative is shown to satisfy a linear stochastic parabolic equation, with an initial condition fulfilled in the weak sense. Given these results, the proof of the L^1 contraction property follows as outlined above.

Finally, we mention that the approach developed herein appears useful in the study of error estimates for numerical approximations of stochastic conservation laws, whenever the approximation is Malliavin differentiable. It seems to us that this Malliavin differentiability is indeed often available. Furthermore, the approach may be extended so as to cover strongly degenerate parabolic equations with Lévy noise, cf. [4], [34]. It also constitutes a starting point for developing a well-posedness theory for stochastic conservation laws with random, possibly anticipating initial data. Note however that this seems to depend on the Malliavin differentiability of the entropy solution (Lemma 5.3 is no longer applicable).

The remaining part of the paper is organized as follows: We present the solution framework and gather some preliminary results in Section 2. Well-posedness results for the viscous approximations are provided in Section 3. Furthermore, we establish the Malliavin differentiability of these approximations and show that the Malliavin derivative can be cast as the solution of a linear stochastic parabolic equation. The question of (weak) satisfaction of the initial condition is addressed. Sections 4 and

5 supply detailed proofs for the existence and uniqueness of Young measure-valued entropy solutions. Finally, some basic results are collected in Section 6.

2. ENTROPY SOLUTIONS

Under the assumption $\sigma(x, 0, z) = 0$, the ordinary L^p spaces ($2 \leq p < \infty$) constitute a natural choice for (1.1). Without this assumption, it turns out that a certain class of weighted L^p spaces are better suited. For non-negative ϕ we define

$$\|u\|_{p,\phi} := \left(\int_{\mathbb{R}^d} |u(x)|^p \phi(x) dx \right)^{1/p}.$$

The relevant weights, denoted by $\mathfrak{N}$, consist of non-zero $\phi \in C^1(\mathbb{R}^d) \cap L^1(\mathbb{R}^d)$ for which there is a constant C_ϕ such that $|\nabla \phi(x)| \leq C_\phi \phi(x)$. The weighted L^p -space associated with ϕ is denoted by $L^p(\mathbb{R}^d, \phi)$.

To see that $\mathfrak{N}$ is non-empty, consider $\phi_N(x) = (1 + |x|^2)^{-N}$ for $N \in \mathbb{N}$. Then we claim that $\phi_N \in \mathfrak{N}$ for all $N \geq d$. To this end, observe that

$$\nabla \phi_N(x) = -2N \frac{x}{1 + |x|^2} \phi_N(x),$$

so $|\nabla \phi_N(x)| \leq 2N \phi_N(x)$. Furthermore

$$\int_{\mathbb{R}^d} \phi_N(x) dx = \int_0^\infty \int_{\partial B(0,r)} \left(\frac{1}{1 + r^2} \right)^N dS(r) dr \leq d\alpha(d)\pi/2.$$

Another function belonging to $\mathfrak{N}$ is $\phi_\lambda(x) = \exp(-\lambda\sqrt{1 + |x|^2})$ for $\lambda > 0$ [38].

The fact that $\phi \in L^1(\mathbb{R}^d)$ yields $L^q(\mathbb{R}^d, \phi) \subset L^p(\mathbb{R}^d, \phi)$ for all $1 \leq p < q$. Indeed, with $r = q/p$ and $s = q/(q - p)$, observe that $r^{-1} + s^{-1} = 1$. Take $h = |u(t)|^p \phi^{1/r}$ and $g = \phi^{1/s}$ and apply Hölder's inequality to $\int |hg| dx$. This yields

$$(2.1) \quad \int_{\mathbb{R}^d} |u(x)|^p \phi dx \leq \left(\int_{\mathbb{R}^d} |u(x)|^q \phi(x) dx \right)^{p/q} \left(\int_{\mathbb{R}^d} \phi(x) dx \right)^{1-p/q},$$

or equivalently $\|u\|_{p,\phi} \leq \|u\|_{q,\phi} \|\phi\|_{L^1(\mathbb{R}^d)}^{1/p-1/q}$.

We shall also make use of the weighted L^∞ -norm

$$\|h\|_{\infty,\phi^{-1}} := \sup_{x \in \mathbb{R}^d} \{ |h(x)| \phi^{-1}(x) \}, \quad h \in C(\mathbb{R}^d).$$

Note that any compactly supported $h \in C(\mathbb{R}^d)$ is bounded in this norm, for $\phi \in \mathfrak{N}$. The norm is convenient due to the inequality $\|u\|_{p,\phi} \leq \|u\|_{p,\phi} \|h\|_{\infty,\phi^{-1}}$.

Denote by $\mathcal{E}$ the set of non-negative convex functions in $C^2(\mathbb{R})$ with $S(0) = 0$, S' bounded, and S'' compactly supported. Suppose $Q : \mathbb{R}^2 \rightarrow \mathbb{R}^d$ satisfies

$$\partial_1 Q(u, c) = S'(u - c) f'(u) \quad (u, c \in \mathbb{R}),$$

where $S \in \mathcal{E}$. Then we call $(S(\cdot - c), Q(\cdot, c))$ an *entropy/entropy-flux pair* (indexed over $c \in \mathbb{R}$). For short, we say that (S, Q) is in $\mathcal{E}$ if S is in $\mathcal{E}$. For $(S, Q) \in \mathcal{E}$, $\varphi \in C_c^\infty([0, T) \times \mathbb{R}^d)$, and $V \in \mathbb{D}^{1,2}$, we define the functional

$$\begin{aligned} \mathbf{Ent}[(S, Q), \varphi, V](u) &:= E \left[\int_{\mathbb{R}^d} S(u^0(x) - V) \varphi(0, x) dx \right] \\ &+ E \left[\iint_{\Pi_T} S(u - V) \partial_t \varphi + Q(u, V) \cdot \nabla \varphi dx dt \right] \\ &- E \left[\iint_{\Pi_T} \int_Z S''(u - V) \sigma(x, u, z) D_{t,z} V \varphi d\mu(z) dx dt \right] \\ &+ \frac{1}{2} E \left[\iint_{\Pi_T} \int_Z S''(u - V) \sigma(x, u, z)^2 \varphi d\mu(z) dx dt \right], \end{aligned}$$

where $D_{t,z}V$ is the Malliavin derivative of V at $(t, z) \in [0, T] \times Z$. We denote by $\mathbb{D}^{1,2}$ the space of Malliavin differentiable random variables in $L^2(\Omega)$ with Malliavin derivative in $L^2(\Omega; L^2([0, T] \times Z))$ [27, p. 27].

We claim that **Ent** is well-defined whenever $V \in \mathbb{D}^{1,2}$, $u \in L^2([0, T] \times \Omega; L^2(\mathbb{R}^d, \phi))$,

$$\|\varphi(t)\|_{\infty, \phi^{-1}}, \|\partial_t \varphi(t)\|_{\infty, \phi^{-1}}, \|\nabla \varphi(t)\|_{\infty, \phi^{-1}} \quad \text{are bounded on } [0, T]$$

(any $\varphi \in C_c^\infty([0, T] \times \mathbb{R}^d)$ meets these criteria). To this end, observe that the first three terms are bounded due to the Lipschitz condition on S . Indeed,

$$(2.2) \quad |Q(u, V)| = \left| \int_V^u S'(u - V) \partial f(z) dz \right| \leq \|S\|_{\text{Lip}} \|f\|_{\text{Lip}} |u - V|,$$

and so

$$(2.3) \quad \begin{aligned} \left| E \left[\iint_{\Pi_T} Q(u, V) \cdot \nabla \varphi dx dt \right] \right| &\leq \|S\|_{\text{Lip}} \|f\|_{\text{Lip}} E \left[\iint_{\Pi_T} (|u| + |V|) |\nabla \varphi| dx dt \right] \\ &\leq \|S\|_{\text{Lip}} \|f\|_{\text{Lip}} \int_0^T E \left[\|u(t)\|_{1, \phi} \right] \|\nabla \varphi(t)\|_{\infty, \phi^{-1}} + E[|V|] \|\nabla \varphi(t)\|_{L^1(\mathbb{R}^d)} dt, \end{aligned}$$

which is finite. The terms involving σ is most easily seen to be well defined as a consequence of the fact that the Hilbert Schmidt norm of $G(u)$ (cf. (1.2)) is bounded. To simplify the notation, we let $HS = \mathcal{L}_2(L^2(Z); L^2(\mathbb{R}^d, \phi))$. Due to assumption $(\mathcal{A}_\sigma)$ and Remark 2.1,

$$\begin{aligned} \|G(u)\|_{HS}^2 &= \int_{\mathbb{R}^d} \int_Z \sigma^2(x, u(x), z) \phi(x) d\mu(z) dx \\ &\leq 2 \|M\|_{L^2(Z)}^2 \int_{\mathbb{R}^d} (1 + |u(x)|^2) \phi(x) dx. \end{aligned}$$

Boundedness of the last term follows as

$$(2.4) \quad \begin{aligned} \left| E \left[\iint_{\Pi_T} \int_Z S''(u - V) \sigma(x, u, z)^2 \varphi d\mu(z) dx dt \right] \right| \\ \leq \|S''\|_\infty E \left[\int_0^T \|\varphi(t)\|_{\infty, \phi^{-1}} \|G(u(t))\|_{HS}^2 dt \right]. \end{aligned}$$

By (2.1), the sub-multiplicativity of the Hilbert Schmidt norm, and the Hölder inequality, it follows that

$$(2.5) \quad \begin{aligned} &\left| E \left[\iint_{\Pi_T} \int_Z S''(u - V) \sigma(x, u, z) D_{t,z} V \varphi d\mu(z) dx dt \right] \right| \\ &\leq \|S''\|_\infty E \left[\iint_{\Pi_T} \|\varphi(t)\|_{\infty, \phi^{-1}} \left| \int_Z \sigma(x, u, z) D_{t,z} V d\mu(z) \right| \phi dx dt \right] \\ &\leq \|S''\|_\infty \|\phi\|_{L^1(\mathbb{R}^d)}^{1/2} E \left[\int_0^T \|\varphi(t)\|_{\infty, \phi^{-1}} \|G(u(t)) D_t V\|_{2, \phi} dt \right] \\ &\leq \|S''\|_\infty \|\phi\|_{L^1(\mathbb{R}^d)}^{1/2} E \left[\int_0^T \|\varphi(t)\|_{\infty, \phi^{-1}}^2 \|G(u(t))\|_{HS}^2 dt \right]^{1/2} \\ &\quad \times \|DV\|_{L^2(\Omega; L^2([0, T] \times Z))} < \infty. \end{aligned}$$

Remark 2.1. Let A be an integral operator with kernel $a = a(x, z)$. In the above calculations, we applied the relation

$$\|A\|_{HS}^2 = \int_{\mathbb{R}^d} \int_Z a^2(x, z) \phi(x) d\mu(z) dx.$$

To see why this holds, recall that $\|A\|_{\text{HS}}^2 = \text{tr}(A^*A)$. Now

$$(A^*Ah)(z) = \int_{\mathbb{R}^d} a(x, z) \left(\int_Z a(x, z') h(z') d\mu(z') \right) \phi(x) dx.$$

Hence, A^*A is an integral operator with kernel $k \in L^2(Z^2)$ defined by

$$k(z, z') = \int_{\mathbb{R}^d} a(x, z) a(x, z') \phi(x) dx,$$

and so

$$\text{tr}(A^*A) = \int_Z k(z, z) d\mu(z) = \int_{\mathbb{R}^d} \int_Z a^2(x, z) \phi(x) d\mu(z) dx.$$

Let $\mathcal{P}$ denote the predictable σ -algebra on $[0, T] \times \Omega$ with respect to $\{\mathcal{F}_t\}$ [10, § 2.2]. In general we are working with equivalence classes of functions with respect to the measure $dt \otimes dP$. The equivalence class u is said to be *predictable* if it has a version $\tilde{u}$ that is $\mathcal{P}$ -measurable. In some of the arguments, to avoid picking versions, we consider the completion of $\mathcal{P}$ with respect to $dt \otimes dP$, denoted by $\mathcal{P}^*$. We recall that any jointly measurable and adapted process is $\mathcal{P}^*$ -measurable, see [10, Theorem 3.7].

Definition 2.1 (Entropy solution). An entropy solution $u = u(t, x; \omega)$ of (1.1) with initial condition $u^0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$ is a function satisfying:

- (i) u is a predictable process in $L^2([0, T] \times \Omega; L^2(\mathbb{R}^d, \phi))$.
- (ii) For any random variable $V \in \mathbb{D}^{1,2}$, any entropy/entropy-flux pair (S, Q) in $\mathcal{E}$, and all nonnegative test functions $\varphi \in C_c^\infty([0, T] \times \mathbb{R}^d)$,

$$\mathbf{Ent}[(S, Q), \varphi, V](u) \geq 0.$$

Here $L^2([0, T] \times \Omega; L^2(\mathbb{R}^d, \phi))$ is the Lebesgue-Bochner space, see Section 6.4.

Remark 2.2 (Weak solutions). By Theorem 5.1 and the proof of Theorem 4.1, the $(\varepsilon \downarrow 0)$ limit of the viscous approximation (3.1) is an entropy solution in the sense of Definition 2.1. It follows that any entropy solution is a weak solution. *Informally* this is a consequence of the integration by parts formula for Malliavin derivatives. To illustrate this, let $d = 1$, W be an ordinary Brownian motion, and suppose that u is a Malliavin differentiable and spatially regular entropy solution. We want to outline an informal argument showing that u is a (strong) solution of (1.1). Let

$$(u)_+ = \begin{cases} u & \text{for } u > 0, \\ 0 & \text{else,} \end{cases} \quad \text{and} \quad \text{sign}_+(u) = \begin{cases} 1 & \text{for } u > 0, \\ 0 & \text{else,} \end{cases}$$

so that $(u)'_+ = \text{sign}_+(u)$. The entropy inequality yields

$$\begin{aligned} & E [\partial_t(u - V)_+ + \partial_x(\text{sign}_+(u - V)(f(u) - f(V)))] \\ & + E [\text{sign}'_+(u - V) \sigma(u) D_t V] - \frac{1}{2} E [\text{sign}'_+(u - V) \sigma^2(u)] \leq 0. \end{aligned}$$

Note that

$$\sigma(u) D_t V - \frac{1}{2} \sigma^2(u) = -\sigma(u)(\sigma(u) - D_t V) + \frac{1}{2} \sigma^2(u).$$

By (1.7) and the chain rule, $D_t \text{sign}_+(u - V) = \text{sign}'_+(u - V)(\sigma(u) - D_t V)$. Hence,

$$\begin{aligned} & E [\partial_t(u - V)_+ + \partial_x(\text{sign}_+(u - V)(f(u) - f(V)))] \\ & - E [D_t \text{sign}_+(u - V) \sigma(u)] + \frac{1}{2} E [\text{sign}'_+(u - V) \sigma^2(u)] \leq 0 \end{aligned}$$

The integration by parts formula of Malliavin calculus yields

$$E [D_t \text{sign}_+(u - V) \sigma(u)] = E [\text{sign}_+(u - V) \sigma(u) \partial_t W],$$

and so

$$\begin{aligned} E \left[\partial_t(u - V)_+ + \partial_x(\text{sign}_+(u - V)(f(u) - f(V))) \right] \\ \leq E \left[\text{sign}_+(u - V) \sigma(u) \partial_t W \right]. \end{aligned}$$

Suppose there is a Malliavin differentiable random variable V satisfying

$$\begin{cases} u - V > 0 & \text{for } \omega \in A, \\ u - V < 0 & \text{else,} \end{cases}$$

for any $A \in \mathcal{F}_T$. Then

$$\partial_t u + \partial_x f(u) \leq \sigma(u) \partial_t W.$$

The opposite inequality follows by considering $(\cdot)^-$.

Let us fix some notation. For $n = 1, 2, \dots$, we will denote by J^n a non-negative, smooth function satisfying

$$\text{supp}(J^n) \subset B(0, 1), \quad \int_{\mathbb{R}^n} J^n(x) dx = 1, \quad \text{and} \quad J^n(x) = J^n(-x),$$

for all $x \in \mathbb{R}^n$. For any $r > 0$ we let $J_r^n(x) = \frac{1}{r^n} J^n(\frac{x}{r})$. For $n = 1$ we let $J_r^+(x) = J_r(x - r)$ and note that $\text{supp}(J_r^+) \subset (0, 2r)$. As the value of n is understood from the context, we will write $J = J^n$.

According to Theorem 4.1 and Theorem 5.1, if $u^0 \in L^p(\Omega; L^p(\mathbb{R}^d, \phi))$, then the entropy solution belongs to $L^p(\Omega \times [0, T]; L^p(\mathbb{R}^d, \phi))$ for any $2 \leq p < \infty$. As a consequence of the entropy inequality we obtain the following:

Proposition 2.1. *Let $2 \leq p < \infty$ and suppose $u^0 \in L^p(\Omega, \mathcal{F}_0, P; L^p(\mathbb{R}^d, \phi))$. If $u \in L^p([0, T] \times \Omega, L^p(\mathbb{R}^d, \phi))$ is an entropy solution of (1.1), then*

$$\text{ess sup}_{0 \leq t \leq T} \left\{ E \left[\|u(t)\|_{p, \phi}^p \right] \right\} < \infty.$$

Proof. Let

$$\varphi_\delta(t, x) = \left(1 - \int_0^t J_\delta(\sigma - \tau) d\sigma \right) \phi(x).$$

Let

$$S_R(u) = \begin{cases} R^p + pR^{p-1}(u - R) & \text{for } u \geq R, \\ |u|^p & \text{for } -R < u < R, \\ R^p - pR^{p-1}(u + R) & \text{for } u \leq -R. \end{cases}$$

Strictly speaking, S_R is not in $\mathcal{E}$, but this can be amended by a simple mollification step (which we ignore). Note that $S_R \rightarrow |\cdot|^p$ pointwise. Furthermore

$$|S'_R(u)| \leq p|u|^{p-1} \quad \text{and} \quad |S''_R(u)| \leq p(p-1)|u|^{p-2}.$$

Under these assumptions it is straightforward to supply estimates of the type (2.3) and (2.4). Hence we may apply the dominated convergence theorem to compute the limit $\lim_{\delta \downarrow 0} \mathbf{Ent}[(S_R, Q_R), \varphi_\delta, 0](u) \geq 0$. This yields

$$\begin{aligned} E \left[\int_{\mathbb{R}^d} S_R(u(\tau)) \phi(x) dx \right] &\leq E \left[\int_{\mathbb{R}^d} S_R(u^0(x)) \phi(x) dx \right] \\ &\quad + E \left[\int_0^\tau \int_{\mathbb{R}^d} Q_R(u, 0) \cdot \nabla \phi dx dt \right] \\ &\quad + \frac{1}{2} E \left[\int_0^\tau \int_{\mathbb{R}^d} \int_Z S''_R(u) \sigma(x, u, z)^2 \phi d\mu(z) dx dt \right], \end{aligned}$$

for almost all $\tau \in [0, T]$. Again, by the dominated convergence theorem, we may send $R \rightarrow \infty$. The result follows. $\square$

It is enough to consider smooth random variables in Definition 2.1, i.e., random variables of the form

$$V = f(W(h_1), \dots, W(h_n))$$

where $f \in C_c^\infty(\mathbb{R}^n)$, W is the isonormal Gaussian process defined by (1.3), and $h_1, \dots, h_n$ are in $H = L^2([0, T] \times Z)$, see [27, p. 25]. We denote the space of smooth random variables by $\mathcal{S}$.

Lemma 2.2. *Suppose $(\mathcal{A}_f)$ and $(\mathcal{A}_\sigma)$ are satisfied. Fix $u \in L^2([0, T] \times \Omega; L^2(\mathbb{R}^d))$, an entropy/entropy-flux pair $(S, Q) \in \mathcal{E}$, and $\varphi \in C_c^\infty([0, T] \times \mathbb{R}^d)$. Then*

$$V \mapsto \mathbf{Ent}[(S, Q), \varphi, V](u)$$

is continuous on $\mathbb{D}^{1,2}$ (in the strong topology).

Remark 2.3. It is not necessary that S'' is compactly supported in the upcoming proof (boundedness and continuity are enough).

Proof. Suppose that $V_n \rightarrow V$ in $\mathbb{D}^{1,2}$ as $n \rightarrow \infty$, and write

$$\begin{aligned} & \mathbf{Ent}[(S, Q), \varphi, V](u) - \mathbf{Ent}[(S, Q), \varphi, V_n](u) \\ &= E \left[\int_{\mathbb{R}^d} (S(u^0(x) - V) - S(u^0(x) - V_n)) \varphi(0, x) dx \right] \\ &+ E \left[\iint_{\Pi_T} (S(u - V) - S(u - V_n)) \partial_t \varphi dx dt \right] \\ &+ E \left[\iint_{\Pi_T} (Q(u, V) - Q(u, V_n)) \cdot \nabla \varphi dx dt \right] \\ &+ E \left[\iint_{\Pi_T} \int_Z (S''(u - V_n) D_{t,z} V_n - S''(u - V) D_{t,z} V) \sigma(x, u, z) \varphi d\mu(z) dx dt \right] \\ &+ \frac{1}{2} E \left[\iint_{\Pi_T} \int_Z (S''(u - V) - S''(u - V_n)) \sigma(x, u, z)^2 \varphi d\mu(z) dx dt \right] \\ &=: \mathcal{T}_1 + \mathcal{T}_2 + \mathcal{T}_3 + \mathcal{T}_4 + \mathcal{T}_5. \end{aligned}$$

We need to show that $\lim_{n \rightarrow \infty} \mathcal{T}_i(n) = 0$ for $1 \leq i \leq 5$.

First, note that $V_n \rightarrow V$ in $L^2(\Omega)$. Next,

$$|\mathcal{T}_1| \leq \|S\|_{\text{Lip}} E[|V - V_n|] \|\varphi(0)\|_{L^1(\mathbb{R})}.$$

Similarly,

$$|\mathcal{T}_2| \leq \|S\|_{\text{Lip}} E[|V - V_n|] \|\partial_t \varphi\|_{L^1(\Pi_T)}.$$

It follows as $V_n \rightarrow V$ in $L^2(\Omega)$ that $\mathcal{T}_1, \mathcal{T}_2 \rightarrow 0$ as $n \rightarrow \infty$.

Concerning $\mathcal{T}_3$, we first observe that for any $\zeta, \xi, \theta \in \mathbb{R}$,

$$\begin{aligned} |Q(\zeta, \xi) - Q(\zeta, \theta)| &= \left| \int_\xi^\zeta S'(z - \xi) \partial f(z) dz - \int_\theta^\zeta S'(z - \theta) \partial f(z) dz \right| \\ &\leq \left| \int_\xi^\zeta (S'(z - \xi) - S'(z - \theta)) \partial f(z) dz \right| + \left| \int_\xi^\theta S'(z - \theta) \partial f(z) dz \right|. \end{aligned}$$

Hence,

$$\begin{aligned} |\mathcal{T}_3| &\leq E \left[\iint_{\Pi_T} \left| \int_V^u (S'(z - V) - S'(z - V_n)) \partial f(z) dz \right| |\nabla \varphi| dx dt \right] \\ &+ E \left[\iint_{\Pi_T} \left| \int_V^{V_n} S'(z - V_n) \partial f(z) dz \right| |\nabla \varphi| dx dt \right] \\ &=: \mathcal{T}_3^1 + \mathcal{T}_3^2. \end{aligned}$$

We want to handle $\mathcal{T}_3^1$ using the dominated convergence theorem. The uniform bound is obtained as in (2.3). Since

$$|S'(z - \xi) - S'(z - \theta)| \leq \|S''\|_\infty |\xi - \theta| \text{ for all } z \in \mathbb{R},$$

it follows that

$$\psi(\theta) := \int_\xi^\zeta (S'(z - \xi) - S'(z - \theta)) \partial f(z) dz$$

is continuous at ξ and satisfies $\psi(\xi) = 0$. Suppose that $\{V_{n(j)}\}$ is a subsequence converging a.e. to V . By the dominated convergence theorem,

$$\begin{aligned} & \lim_{j \rightarrow \infty} \mathcal{T}_3^1(n(j)) \\ &= E \left[\iint_{\Pi_T} \lim_{j \rightarrow \infty} \left| \int_V^u (S'(z - V) - S'(z - V_{n(j)})) \partial f(z) dz \right| |\nabla \varphi| dx dt \right] = 0. \end{aligned}$$

By the above, any subsequence of $\{\mathcal{T}_3^1(n)\}_{n \geq 1}$ possesses a subsequence converging to 0, and so $\lim_{n \rightarrow \infty} \mathcal{T}_3^1(n) = 0$. Consider $\mathcal{T}_3^2$. Since

$$\mathcal{T}_3^2 \leq \|S\|_{\text{Lip}} \|f\|_{\text{Lip}} E[|V_n - V|] \|\nabla \varphi\|_{L^1(\Pi_T)},$$

it follows that $\lim_{n \rightarrow \infty} \mathcal{T}_3^2 = 0$.

Concerning the $\mathcal{T}_4$ -term, we first split it as follows:

$$\begin{aligned} \mathcal{T}_4 &= E \left[\iint_{\Pi_T} \int_Z S''(u - V_n) (D_{t,z} V_n - D_{t,z} V) \sigma(x, u, z) \varphi d\mu(z) dx dt \right] \\ &+ E \left[\iint_{\Pi_T} \int_Z (S''(u - V_n) - S''(u - V)) D_{t,z} V \sigma(x, u, z) \varphi d\mu(z) dx dt \right] \\ &= \mathcal{T}_4^1 + \mathcal{T}_4^2. \end{aligned}$$

By (2.5), $\lim_{n \rightarrow \infty} \mathcal{T}_4^1 = 0$. Owing to (2.5), the dominated convergence theorem implies $\lim_{n \rightarrow \infty} \mathcal{T}_4^2 = 0$. Finally, by (2.4) and the dominated convergence theorem, also $\lim_{n \rightarrow \infty} \mathcal{T}_5 = 0$. $\square$

For the proof of existence of solutions, it is convenient to introduce a weaker notion of entropy solution based on Young measures (see, e.g., [3, 15, 16, 28]). Instead of representing the solution as an element in $\mathcal{Y}(\Pi_T \times \Omega; \mathbb{R})$ we use the notion of entropy process proposed in [16] or equivalently the strong measure-valued solution proposed in [28]. Any probability measure ν on the real line may be represented by a measurable function $u : [0, 1] \rightarrow \mathbb{R} \cup \{\infty\}$ such that ν is the image of the Lebesgue measure $\mathcal{L}$ on $[0, 1]$ by u . In fact we may take (See [37, § 2.2.2])

$$(2.6) \quad u(\alpha) = \inf \{ \xi \in \mathbb{R} : \nu((-\infty, \xi]) > \alpha \}.$$

A Young measure $\nu \in \mathcal{Y}(\Pi_T \times \Omega; \mathbb{R})$ is thus represented by a (higher dimensional) function $u : \Pi_T \times [0, 1] \times \Omega \rightarrow \mathbb{R}$ where u satisfies $\nu_{t,x,\omega}(B) = \mathcal{L}(u(t, x, \cdot, \omega)^{-1}(B))$ for any measurable $B \subset \mathbb{R}$. The extension to Young measure-valued solutions is thus obtained through the embedding defined by

$$(2.7) \quad \Phi(u)(t, x, \alpha, \omega) = u(t, x, \omega).$$

Given a functional F we define the extension

$$\mathcal{Y}(F)(u) = \int_0^1 F(u(\alpha)) d\alpha,$$

so that $\mathcal{Y}(F) \circ \Phi = F$. For $1 \leq p < \infty$ we let

$$\|u\|_{p, \phi \otimes 1} = \left(\int_0^1 \int_{\mathbb{R}^d} |u(x, \alpha)|^p \phi(x) dx d\alpha \right)^{1/p}.$$

The associated space is denoted by $L^p(\mathbb{R}^d \times [0, 1], \phi)$.

Definition 2.2 (Young measure-valued entropy solution). A Young measure-valued entropy solution $u = u(t, x, \alpha; \omega)$ of (1.1), with initial condition u^0 belonging to $L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$, is a function satisfying:

- (i) u is a predictable process in $L^2([0, T] \times \Omega; L^2(\mathbb{R}^d \times [0, 1], \phi))$.
- (ii) For any random variable $V \in \mathbb{D}^{1,2}$, any entropy/entropy-flux pair (S, Q) in $\mathcal{E}$, and all nonnegative test functions $\varphi \in C_c^\infty([0, T] \times \mathbb{R}^d)$,

$$(2.8) \quad \mathcal{Y}(\mathbf{Ent}[(S, Q), \varphi, V])(u) \geq 0.$$

The next result is concerned with the essential continuity of the solutions at $t = 0$. A similar argument can be found in [7].

Lemma 2.3 (Initial condition). Suppose $(\mathcal{A}_f)$ and $(\mathcal{A}_\sigma)$ are satisfied, and that u^0 belongs to $L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$. Let u be a Young measure-valued entropy solution of (1.1) in the sense of Definition 2.2. Let $S : \mathbb{R} \rightarrow [0, \infty)$ be Lipschitz continuous and satisfy $S(0) = 0$. For any $\psi \in C_c^\infty(\mathbb{R}^d)$,

$$\mathcal{T}_{r_0} := E \left[\iint_{\Pi_T} \int_{[0,1]} S(u(t, x, \alpha) - u^0(x)) \psi(x) J_{r_0}^+(t) d\alpha dx dt \right] \rightarrow 0 \text{ as } r_0 \downarrow 0.$$

Remark 2.4. The proof does not depend on the differentiability of $J_{r_0}^+$. Hence the above limit may be replaced by

$$\lim_{\tau \downarrow 0} E \left[\frac{1}{\tau} \int_0^\tau \int_{\mathbb{R}^d} \int_{[0,1]} S(u(t, x, \alpha) - u^0(x)) \psi(x) d\alpha dx dt \right] = 0.$$

Proof. Let $S \in C^\infty(\mathbb{R})$ with bounded derivatives. Take

$$\varphi(t, x, y) = \xi_{r_0}(t) \psi(x) J_r(x - y) \text{ where } \xi_{r_0}(t) = 1 - \int_0^t J_{r_0}^+(s) ds.$$

Then let $V = u^0(y)$ in (2.8) and integrate in y . This implies

(2.9)

$$\begin{aligned} I &:= E \left[\int_{\mathbb{R}^d} \iint_{\Pi_T} \int_{[0,1]} S(u(t, x, \alpha) - u^0(y)) \psi(x) J_r(x - y) J_{r_0}^+(t) d\alpha dx dt dy \right] \\ &\leq E \left[\int_{\mathbb{R}^d} \iint_{\Pi_T} \int_{[0,1]} Q(u(t, x, \alpha), u^0(y)) \cdot \nabla_x \varphi d\alpha dx dt dy \right] \\ &\quad + E \left[\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} S(u^0(x) - u^0(y)) \varphi(0, x, y) dx dy \right] \\ &\quad + \frac{1}{2} E \left[\int_{\mathbb{R}^d} \iint_{\Pi_T} \int_{[0,1]} \int_Z S''(u(t, x, \alpha) - u^0(y)) \sigma(x, u, z)^2 \varphi d\mu(z) d\alpha dx dt dy \right] \\ &=: \mathcal{T}^1 + \mathcal{T}^2 + \mathcal{T}^3. \end{aligned}$$

Let us first observe that

$$\begin{aligned} I &= \mathcal{T}_{r_0} + E \left[\int_{\mathbb{R}^d} \iint_{\Pi_T} \int_{[0,1]} (S(u(t, x, \alpha) - u^0(y)) - S(u(t, x, \alpha) - u^0(x))) \right. \\ &\quad \left. \times \psi(x) J_r(x - y) J_{r_0}^+(t) d\alpha dx dt dy \right] =: \mathcal{T}_{r_0} + I^1. \end{aligned}$$

We want to take the limit $r_0 \downarrow 0$. First observe that we have the bound

$$|I^1| \leq \|S\|_{\text{Lip}} E \left[\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |u^0(x) - u^0(y)| \psi(x) J_r(x - y) dx dy \right] =: R,$$

which is independent of r_0 . Similarly, $|\mathcal{T}^2| \leq R$. Note that $\xi_{r_0} \rightarrow 0$ a.e. as $r_0 \downarrow 0$, so due to assumptions $(\mathcal{A}_f)$ and $(\mathcal{A}_\sigma)$, one may conclude by the dominated convergence theorem and estimates similar to those in (2.3) and (2.4) that

$$\lim_{r_0 \downarrow 0} \mathcal{T}^1 = \lim_{r_0 \downarrow 0} \mathcal{T}^3 = 0.$$

Thus, it follows by (2.9) that

$$\lim_{r_0 \downarrow 0} \mathcal{T}_{r_0} \leq 2R.$$

Since $r > 0$ was arbitrary, and $\lim_{r \downarrow 0} R = 0$, we have arrived at $\lim_{r_0 \downarrow 0} \mathcal{T}_{r_0} \leq 0$. The desired result follows, since we can approximate any Lipschitz function uniformly by smooth functions with bounded derivatives. $\square$

3. THE VISCOUS APPROXIMATION

For each fixed $\varepsilon > 0$, we denote by u^ε the solution of the regularized problem

$$(3.1) \quad \begin{cases} du^\varepsilon + \nabla \cdot f(u^\varepsilon) dt = \int_Z \sigma(x, u^\varepsilon, z) W(dt, dz) + \varepsilon \Delta u^\varepsilon dt, & (t, x) \in \Pi_T, \\ u^\varepsilon(0, x) = u^0(x), & x \in \mathbb{R}^d. \end{cases}$$

As in the deterministic case, the idea is to let $\varepsilon \rightarrow 0$ and obtain a solution to the stochastic conservation law (1.1). The entropy condition is meant to single out this limit as the only proper (weak) solution; the entropy solution. To show that this limit exists, a type of compactness argument is needed [3, 17, 9].

The existence of a unique solution to (3.1) may be found several places [3, 17]. In particular, the semi-group approach presented in [30, ch. 9] may be applied. The functional setting of [30] is that of a Hilbert space, and so the natural choice here is $L^2(\mathbb{R}^d, \phi)$ where $\phi \in \mathfrak{N}$. Due to the new functional setting, we have chosen to include proofs for some of the results relating to (3.1).

3.1. A priori estimates and well-posedness. Let S_ε be the semi-group generated by the heat kernel. That is $S_\varepsilon(t)u = \Phi_\varepsilon(t) \star u$ where

$$\Phi_\varepsilon(t, x) := \frac{1}{(4\varepsilon\pi t)^{d/2}} \exp\left(-\frac{|x|^2}{4\varepsilon t}\right).$$

Let $F(u) = \nabla \cdot f(u)$ and G be defined by (1.2). In this setting the key conditions [30, p. 142] for well posedness of (3.1) is given by:

- (F) $D(F)$ is dense in $L^2(\mathbb{R}^d, \phi)$ and there is a function $a : (0, \infty) \rightarrow (0, \infty)$ satisfying $\int_0^T a(t) dt < \infty$ for all $T < \infty$ such that, for all $t > 0$ and $u, v \in D(F)$,

$$\begin{aligned} \|S_\varepsilon(t)F(u)\|_{L^2(\mathbb{R}^d, \phi)} &\leq a(t) \left(1 + \|u\|_{L^2(\mathbb{R}^d, \phi)}\right), \\ \|S_\varepsilon(t)(F(u) - F(v))\|_{L^2(\mathbb{R}^d, \phi)} &\leq a(t) \|u - v\|_{L^2(\mathbb{R}^d, \phi)}. \end{aligned}$$

- (G) $D(G)$ is dense in $L^2(\mathbb{R}^d, \phi)$ and there is a function $b : (0, \infty) \rightarrow (0, \infty)$ satisfying $\int_0^T b^2(t) dt < \infty$ for all $T < \infty$ such that, for all $t > 0$ and $u, v \in D(G)$,

$$\begin{aligned} \|S_\varepsilon(t)G(u)\|_{\mathcal{L}_2(L^2(Z); L^2(\mathbb{R}^d, \phi))} &\leq b(t) \left(1 + \|u\|_{L^2(\mathbb{R}^d, \phi)}\right), \\ \|S_\varepsilon(t)(G(u) - G(v))\|_{\mathcal{L}_2(L^2(Z); L^2(\mathbb{R}^d, \phi))} &\leq b(t) \|u - v\|_{L^2(\mathbb{R}^d, \phi)}. \end{aligned}$$

Suppose $u^0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$. Under assumptions (F) and (G) we may conclude by [30, Theorem 9.15, Theorem 9.29] that there exists a unique predictable process $u^\varepsilon : [0, T] \times \Omega \rightarrow L^2(\mathbb{R}^d, \phi)$ such that

(i)

$$(3.2) \quad \sup_{0 \leq t \leq T} E \left[\|u^\varepsilon(t)\|_{2,\phi}^2 \right] < \infty.$$

(ii) For all $0 \leq t \leq T$

(3.3)

$$\begin{aligned} u^\varepsilon(t, x) &= \int_{\mathbb{R}^d} \Phi_\varepsilon(t, x - y) u^0(y) dy \\ &\quad - \int_0^t \int_{\mathbb{R}^d} \nabla_x \Phi_\varepsilon(t - s, x - y) \cdot f(u^\varepsilon(s, y)) dy ds \\ &\quad + \int_0^t \int_Z \int_{\mathbb{R}^d} \Phi_\varepsilon(t - s, x - y) \sigma(y, u^\varepsilon(s, y), z) dy W(ds, dz). \end{aligned}$$

(iii) u^ε is a weak solution of (3.1), i.e., for any test function $\varphi \in C_c^\infty(\mathbb{R}^d)$ and any pair of times t_0, t with $0 \leq t_0 \leq t \leq T$,

$$(3.4) \quad \begin{aligned} \int_{\mathbb{R}^d} u^\varepsilon(t) \varphi dx &= \int_{\mathbb{R}^d} u^\varepsilon(t_0) \varphi dx - \int_{t_0}^t \int_{\mathbb{R}^d} f(u^\varepsilon(s)) \cdot \nabla \varphi dx ds \\ &\quad + \int_{t_0}^t \int_{\mathbb{R}^d} \int_Z \sigma(x, u^\varepsilon(s, x), z) \varphi W(ds, dz) dx + \varepsilon \int_{t_0}^t \int_{\mathbb{R}^d} u^\varepsilon \Delta \varphi dx ds, \end{aligned}$$

 dP -almost surley.

To see that conditions (F) and (G) are satisfied we prove the following estimate:

Lemma 3.1. *Let $\phi \in \mathfrak{N}$ and $1 \leq p < \infty$. Suppose $C_\phi \sqrt{4\varepsilon t} \leq 1$. Let $v \in W^{1,p}(\mathbb{R}^d, \phi; \mathbb{R}^d)$ and $u \in L^p(\mathbb{R}^d, \phi)$. Whenever $C_\phi \sqrt{4\varepsilon t} \leq 1$,*

$$\begin{aligned} (i) \quad & \|\Phi_\varepsilon(t) \star u\|_{p,\phi} \leq \kappa_{1,d} \|u\|_{p,\phi}, \\ (ii) \quad & \|\Phi_\varepsilon(t) \star \nabla \cdot v\|_{p,\phi} \leq \frac{\kappa_{2,d}}{\sqrt{\varepsilon t}} \|v\|_{p,\phi}, \end{aligned}$$

where $\kappa_{1,d} = c_{d-1} \frac{d\alpha(d)}{\pi^{d/2}}, \kappa_{2,d} = c_d \frac{d\alpha(d)}{\pi^{d/2}}$, and

$$c_d = \int_0^\infty \zeta^d (1 + \zeta)^2 \exp(\zeta - \zeta^2) d\zeta.$$

The volume of the unit ball in $\mathbb{R}^d$ is denoted by $\alpha(d)$.

Before we give a proof let us see why (F) and (G) follows. Recall that we may assume $f(0) = 0$ without any loss of generality. By Lemma 3.1 and $(\mathcal{A}_f)$

$$\|S_\varepsilon(t)F(u)\|_{L^2(\mathbb{R}^d, \phi)} = \|\Phi_\varepsilon(t) \star \nabla \cdot f(u)\|_{L^2(\mathbb{R}^d, \phi)} \leq \underbrace{\frac{\kappa_{2,d}}{\sqrt{\varepsilon t}} \|f\|_{\text{Lip}}}_{a(t)} \|u\|_{2,\phi}.$$

It remains to observe that $\int_0^T \frac{1}{\sqrt{t}} dt = 2\sqrt{T} < \infty$. The second part of (F) follows similarly. Let us consider (G). First observe that

$$S_\varepsilon(t)G(u)h(x) = \int_Z \left(\int_{\mathbb{R}^d} \Phi_\varepsilon(t, x - y) \sigma(y, u(y), z) dy \right) h(z) d\mu(z).$$

Recall that $HS = \mathcal{L}_2(L^2(Z); L^2(\mathbb{R}^d, \phi))$. By Lemma 3.1 and $(\mathcal{A}_\sigma)$

$$\begin{aligned} \|S_\varepsilon(t)G(u)\|_{HS}^2 &= \int_Z \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \Phi_\varepsilon(t, x - y) \sigma(y, u(y), z) dy \right)^2 \phi(x) dx d\mu(z) \\ &= \int_Z \|\Phi_\varepsilon(t) \star \sigma(\cdot, u, z)\|_{2,\phi}^2 d\mu(z) \\ &\leq \kappa_{1,d}^2 \|M\|_{L^2(Z)}^2 (\|\phi\|_{L^1(\mathbb{R}^d)} + \|u\|_{2,\phi})^2. \end{aligned}$$

This yields the first part of condition (G). The second part follows similarly, in view of the Lipschitz assumption on σ .

Proof of Lemma 3.1. Consider (i). By Proposition 6.4,

$$\|\Phi_\varepsilon(t) \star u\|_{p,\phi} \leq \underbrace{\left(\int_{\mathbb{R}^d} |\Phi_\varepsilon(t, x)| (1 + w_{p,\phi}(|x|)) dx \right)}_{\|\Phi(t)\|} \|u\|_{p,\phi}.$$

where

$$w_{p,\phi}(r) = \frac{C_\phi}{p} r \left(1 + \frac{C_\phi}{p} r \right) \exp \left(\frac{C_\phi}{p} r \right).$$

We apply polar coordinates to compute $\|\Phi(t)\|$. This yields

$$\begin{aligned} \|\Phi(t)\| &= \int_0^\infty \int_{\partial B(0,r)} |\Phi_\varepsilon(t)| (r) (1 + w_{p,\phi}(r)) dS(r) dr \\ &= \frac{d\alpha(d)}{(4\varepsilon\pi t)^{d/2}} \int_0^\infty r^{d-1} \exp \left(\frac{C_\phi}{p} r - \frac{r^2}{4\varepsilon t} \right) \\ &\quad \times \left(\exp \left(-\frac{C_\phi}{p} r \right) + \frac{C_\phi}{p} r \left(1 + \frac{C_\phi}{p} r \right) \right) dr. \end{aligned}$$

To simplify, we note that

$$\exp \left(-\frac{C_\phi}{p} r \right) + \frac{C_\phi}{p} r \left(1 + \frac{C_\phi}{p} r \right) \leq \left(1 + \left(\frac{C_\phi}{p} r \right) \right)^2$$

Let $\zeta = r/\sqrt{4\varepsilon t}$. Provided $C_\phi\sqrt{4\varepsilon t} \leq 1$, it follows that

$$\frac{C_\phi}{p} r = \frac{C_\phi}{p} \sqrt{4\varepsilon t} \zeta \leq \zeta.$$

Inserting this we obtain

$$\|\Phi(t)\| \leq \frac{d\alpha(d)}{\pi^{d/2}} \int_0^\infty \zeta^{d-1} (1 + \zeta)^2 \exp(\zeta - \zeta^2) d\zeta$$

Estimate (ii) follows along the same lines. Integration by parts yields

$$\int_{\mathbb{R}^d} \Phi_\varepsilon(t, x - y) \nabla \cdot v(y) dy = \int_{\mathbb{R}^d} \nabla_x \Phi_\varepsilon(t, x - y) \cdot v(y) dy.$$

Hence,

$$|\Phi_\varepsilon(t) \star \nabla \cdot v| \leq |\nabla \Phi(t)| \star |v|.$$

By Proposition 6.4,

$$\|\nabla_x \Phi_\varepsilon(t) \star v\|_{L^p(\mathbb{R}^d)} \leq \underbrace{\left(\int_{\mathbb{R}^d} |\nabla \Phi_\varepsilon(t, x)| (1 + w_{p,\phi}(|x|)) dx \right)}_{\|\nabla \Phi(t)\|} \|v\|_{L^p(\mathbb{R}^d, \phi; \mathbb{R}^d)}.$$

Let $r = |x|$. Then

$$\|\nabla \Phi(t)\| = \int_0^\infty \underbrace{\int_{\partial B(0,r)} |\nabla \Phi_\varepsilon(t)| (r) (1 + w_{p,\phi}(r)) dS(r)}_{\Psi(r)} dr.$$

Now,

$$\nabla \Phi_\varepsilon(t, x) = -\frac{2\pi x}{(4\pi\varepsilon t)^{d/2+1}} \exp \left(-\frac{|x|^2}{4\varepsilon t} \right),$$

and so

$$\Psi(r) = \frac{d\alpha(d)}{2\varepsilon t \pi^{d/2}} \left(\frac{r}{\sqrt{4\varepsilon t}} \right)^d \exp \left(\frac{C_\phi}{p} r - \frac{r^2}{4\varepsilon t} \right) \left(\exp \left(-\frac{C_\phi}{p} r \right) + \frac{C_\phi}{p} r \left(1 + \frac{C_\phi}{p} r \right) \right).$$

Let $\zeta(r) = r/\sqrt{4\epsilon t}$ and suppose $C_\phi\sqrt{4\epsilon t} \leq 1$. Then

$$\begin{aligned} \int_0^\infty \left(\frac{r}{\sqrt{4\epsilon t}}\right)^d \exp\left(\frac{C_\phi}{p}r - \frac{r^2}{4\epsilon t}\right) \left(1 + \left(\frac{C_\phi}{p}r\right)\right)^2 dr \\ \leq \sqrt{4\epsilon t} \int_0^\infty \zeta^d(1 + \zeta)^2 \exp(\zeta - \zeta^2) d\zeta. \end{aligned}$$

This concludes the proof of the lemma. $\square$

The following two estimates constitute the reason why Lemma 3.1 is the key to the well-posedness of (3.1). As we will see, the relevant properties of u^ϵ follow rather easily with these estimates at hand.

Lemma 3.2. *Let $1 \leq p \leq \infty$ and $\phi \in \mathfrak{N}$. Suppose $v \in C([0, T]; W^{1,p}(\mathbb{R}^d, \phi; \mathbb{R}^d))$. Set*

$$\mathcal{T}[v](t, x) = \int_0^t \int_{\mathbb{R}^d} \Phi_\epsilon(t-s, x-y) (\nabla \cdot v(s, y)) dy ds.$$

Then, for any $1 \leq q < \infty$,

$$\|\mathcal{T}[v](t)\|_{L^p(\mathbb{R}^d, \phi)}^q \leq \kappa_{2,d}^q \left(2\sqrt{\frac{t}{\epsilon}}\right)^{q-1} \int_0^t \frac{1}{\sqrt{\epsilon(t-s)}} \|v(s)\|_{L^p(\mathbb{R}^d, \phi)}^q ds,$$

where $\kappa_{2,d} = c_d \frac{d\alpha(d)}{\pi^{d/2}}$ and c_d is defined in Lemma 3.1.

Proof. By Minkowski's integral inequality [31, p.271] and Lemma 3.1,

$$\begin{aligned} \|\mathcal{T}[v](t)\|_{p,\phi}^q &\leq \left(\int_0^t \|\Phi_\epsilon(t-s) \star (\nabla \cdot v(s, y))\|_{p,\phi} ds \right)^q \\ &\leq \left(\int_0^t \frac{\kappa_{2,d}}{\sqrt{\epsilon(t-s)}} \|v(s)\|_{p,\phi} ds \right)^q. \end{aligned}$$

If $q = 1$ we are done, so we may assume $1 < q < \infty$. Let r satisfy $1 = r^{-1} + q^{-1}$ and take

$$h(s) := \left(\frac{1}{\sqrt{\epsilon(t-s)}}\right)^{1/r} \quad \text{and} \quad g(s) := \left(\frac{1}{\sqrt{\epsilon(t-s)}}\right)^{1-1/r} \|v(s)\|_{p,\phi}.$$

By Hölder's inequality, $\|hg\|_{L^1([0,t])}^q \leq \|g\|_{L^q([0,t])}^q \|h\|_{L^r([0,t])}^q$, and so

$$\|\mathcal{T}[v](t)\|_{L^p(\mathbb{R}^d)}^q \leq (\kappa_{2,d} \|h\|_{L^r([0,t])})^q \int_0^t \frac{1}{\sqrt{\epsilon(t-s)}} \|v(s)\|_{L^p(\mathbb{R}^d; \mathbb{R}^d)}^q ds.$$

A simple computation yields

$$\|h\|_{L^r([0,t])}^q = \left(\int_0^t \frac{1}{\sqrt{\epsilon(t-s)}} ds \right)^{q/r} = \left(2\sqrt{\frac{t}{\epsilon}} \right)^{q-1}.$$

The result follows. $\square$

Lemma 3.3. *Let $2 \leq p < \infty$ and $\phi \in \mathfrak{N}$. Suppose $v : \Omega \times [0, T] \times Z \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a predictable process satisfying*

$$|v(s, x, z)| \leq K(s, x)M(z),$$

for $M \in L^2(Z)$ and a process $K \in L^2([0, T]; L^p(\Omega; L^p(\mathbb{R}^d, \phi)))$. Define

$$\mathcal{T}[v](t, x) = \int_0^t \int_Z \int_{\mathbb{R}^d} \Phi_\epsilon(t-s, x-y) v(s, y, z) dy W(dz, ds).$$

Then

$$E \left[\|\mathcal{T}[v](t)\|_{p,\phi}^p \right]^{1/p} \leq c_p^{1/p} \kappa_{1,d} \|M\|_{L^2(Z)} \left(\int_0^t E \left[\|K(s)\|_{p,\phi}^p \right]^{2/p} ds \right)^{1/2},$$

where c_p is the constant appearing in the Burkholder-Davis-Gundy inequality and $\kappa_{1,d} = c_{d-1} \frac{d\alpha(d)}{\pi^{d/2}}$, with c_d defined in Lemma 3.1.

Remark 3.1. To prove this result we use the Burkholder-Davis-Gundy inequality for $\mathbb{R}$ valued processes. Using Banach space valued versions [12, 33], one can derive more general estimates.

Proof. First note that

$$M(t, x) = \int_0^t \int_Z \int_{\mathbb{R}^d} \Phi_\varepsilon(\tau - s, x - y) v(s, y, z) dy W(dz, ds)$$

is a martingale on $[0, \tau]$, and so by the Burkholder-Davis-Gundy inequality [22],

$$E \left[|\mathcal{T}[v](t, x)|^p \right] \leq c_p E \left[\left(\int_0^t \int_Z |\Phi_\varepsilon(t - s) \star v(s, \cdot, z)(x)|^2 d\mu(z) ds \right)^{p/2} \right].$$

Upon integrating in space and applying Minkowski's inequality, it follows that

$$\begin{aligned} \|\mathcal{T}[v](t)\|_{p,\phi}^2 &\leq c_p^{2/p} E \left[\int_{\mathbb{R}^d} \left(\int_0^t \int_Z |\Phi_\varepsilon(t - s) \star v(s, \cdot, z)(x)|^2 d\mu(z) ds \right)^{p/2} \phi(x) dx \right]^{2/p} \\ &\leq c_p^{2/p} \int_0^t \int_Z \left(E \left[\int_{\mathbb{R}^d} |\Phi_\varepsilon(t - s) \star v(s, \cdot, z)(x)|^p \phi(x) dx \right] \right)^{2/p} d\mu(z) ds. \end{aligned}$$

By Lemma 3.1,

$$\|\mathcal{T}[v](t)\|_{p,\phi}^2 \leq c_p^{2/p} \kappa_{1,d}^2 \int_0^t \int_Z E \left[\|v(s, \cdot, z)\|_{p,\phi}^p \right]^{2/p} d\mu(z) ds.$$

By assumption,

$$\int_0^t \int_Z E \left[\|v(s, \cdot, z)\|_{p,\phi}^p \right]^{2/p} d\mu(z) ds \leq \|M\|_{L^2(Z)}^2 \int_0^t E \left[\|K(s)\|_{p,\phi}^p \right]^{2/p} ds.$$

□

For a Banach space E we denote by $\mathcal{X}_{\beta,q,E}$ the space of pathwise continuous predictable processes $u : [0, T] \times \Omega \rightarrow E$ normed by

$$(3.5) \quad \|u\|_{\beta,q,E} := \left(\sup_{t \in [0,T]} e^{-\beta t} E \left[\|u(t)\|_E^q \right] \right)^{1/q}.$$

The existence of a solution to (3.1) is obtained by the Banach fixed-point theorem, applied to the operator

$$\begin{aligned} \mathcal{S}(u)(t, x) &:= \int_{\mathbb{R}^d} \Phi_\varepsilon(t, x - y) u^0(y) dy \\ &\quad - \int_0^t \int_{\mathbb{R}^d} \nabla_x \Phi_\varepsilon(t - s, x - y) \cdot f(u(s, y)) dy ds \\ &\quad + \int_0^t \int_Z \int_{\mathbb{R}^d} \Phi_\varepsilon(t - s, x - y) \sigma(y, u(s, y), z) dy W(ds, dz) \end{aligned}$$

on the space $\mathcal{X}_{\beta,2,L^2(\mathbb{R}^d,\phi)}$, with $\beta \in \mathbb{R}$ sufficiently large. It follows that the sequence $\{u_n\}_{n \geq 1}$ defined inductively by $u_0 = 0$ and $u_{n+1} = \mathcal{S}(u_n)$ converges to u^ε in $\mathcal{X}_{\beta,2,L^2(\mathbb{R}^d,\phi)}$ as $n \rightarrow \infty$. By Lemmas 3.2 and 3.3 we are free to use the space $\mathcal{X}_{\beta,p,L^p(\mathbb{R}^d,\phi)}$ for any $2 \leq p < \infty$ in the fixed-point argument [17].

We can use Lemmas 3.2 and 3.3 to deduce a continuous dependence result. To do this, we need a measure of the distance between the coefficients. For the flux function f , the Lipschitz norm is a reasonable choice. Concerning the noise function σ , we introduce the norm $\|\sigma\|_{\text{Lip}} = \|M_\sigma\|_{L^2(Z)}$, where

$$M_\sigma(z) = \sup_{x \in \mathbb{R}^d} \left\{ \sup_{u \in \mathbb{R}} \frac{|\sigma(x, u, z)|}{1 + |u|} \right\} + \sup_{x \in \mathbb{R}^d} \left\{ \sup_{u \neq v} \frac{|\sigma(x, u, z) - \sigma(x, v, z)|}{|u - v|} \right\}.$$

Note that for any σ satisfying $(\mathcal{A}_\sigma)$, we have $\|\sigma\|_{\text{Lip}} < \infty$.

Proposition 3.4 (Continuous dependence). *Let $2 \leq p < \infty$ and $\phi \in \mathfrak{N}$. Let f_1, f_2 satisfy $(\mathcal{A}_f)$ and σ_1, σ_2 satisfy $(\mathcal{A}_\sigma)$. Suppose $u_1^0, u_2^0 \in L^p(\Omega, \mathcal{F}_0, P; L^p(\mathbb{R}^d, \phi))$. Let u_1^ε and u_2^ε denote the weak solutions of the corresponding problems (3.1) with $f = f_i, \sigma = \sigma_i$, and $u^0 = u_i^0$, for $i = 1, 2$. Then, for $\beta > 0$ sufficiently large there exists a constant $C = C(\beta, \varepsilon, T, f_1, \sigma_1)$ such that*

$$\begin{aligned} \|u_1^\varepsilon - u_2^\varepsilon\|_{\beta, p, L^p(\mathbb{R}^d, \phi)} &\leq C \left(E \left[\|u_1^0 - u_2^0\|_{p, \phi}^p \right] + \|f_1 - f_2\|_{\text{Lip}} \|u_1^\varepsilon\|_{\beta, p, L^p(\mathbb{R}^d, \phi)} \right. \\ &\quad \left. + \|\sigma_1 - \sigma_2\|_{\text{Lip}} \left(\|\phi\|_{L^1(\mathbb{R}^d)} + \|u_1^\varepsilon\|_{\beta, p, L^p(\mathbb{R}^d, \phi)} \right) \right), \end{aligned}$$

where the norm $\|\cdot\|_{\beta, p, L^p(\mathbb{R}^d, \phi)}$ is defined in (3.5).

Proof. By (3.3),

$$\begin{aligned} u_1^\varepsilon(t, x) - u_2^\varepsilon(t, x) &= \int_{\mathbb{R}^d} \Phi_\varepsilon(t, x - y)(u_1^0(y) - u_2^0(y)) dy \\ &\quad - \int_0^t \int_{\mathbb{R}^d} \nabla_x \Phi_\varepsilon(t - s, x - y) \cdot (f_1(u_1^\varepsilon(s, y)) - f_2(u_2^\varepsilon(s, y))) dy ds \\ &\quad + \int_0^t \int_Z \int_{\mathbb{R}^d} \Phi_\varepsilon(t - s, x - y)(\sigma_1(y, u_1^\varepsilon(s, y), z) - \sigma_2(y, u_2^\varepsilon(s, y), z)) dy W(ds, dz) \\ &= \mathcal{T}_1 + \mathcal{T}_2 + \mathcal{T}_3. \end{aligned}$$

By Lemma 3.1,

$$E \left[\|\mathcal{T}_1(t)\|_{p, \phi}^p \right] \leq \kappa_{1, d}^p E \left[\|u_1^0 - u_2^0\|_{p, \phi}^p \right].$$

where $\kappa_{1, d}$ is defined in Lemma 3.3. Hence

$$(3.6) \quad \|\mathcal{T}_1\|_{\beta, p, L^p(\mathbb{R}^d, \phi)} \leq \kappa_{1, d} E \left[\|u_1^0 - u_2^0\|_{p, \phi}^p \right]^{1/p}.$$

Consider $\mathcal{T}_2$. Note that

$$\begin{aligned} E \left[\|f_1(u_1^\varepsilon(s)) - f_2(u_2^\varepsilon(s))\|_{p, \phi}^p \right]^{1/p} &\leq \|f_1 - f_2\|_{\text{Lip}} E \left[\|u_1^\varepsilon(s)\|_{p, \phi}^p \right]^{1/p} \\ &\quad + \|f_2\|_{\text{Lip}} E \left[\|u_1^\varepsilon(s) - u_2^\varepsilon(s)\|_{p, \phi}^p \right]^{1/p}. \end{aligned}$$

By Lemma 3.2,

$$\begin{aligned} &E \left[\|\mathcal{T}_2(t)\|_{L^p(\mathbb{R}^d)}^p \right] \\ &\leq \kappa_{2, d}^p \left(4\sqrt{\frac{t}{\varepsilon}} \right)^{p-1} \|f_1 - f_2\|_{\text{Lip}}^p \int_0^t \frac{1}{\sqrt{\varepsilon(t-s)}} E \left[\|u_1^\varepsilon(s)\|_{p, \phi}^p \right] ds \\ &\quad + \kappa_{2, d}^p \left(4\sqrt{\frac{t}{\varepsilon}} \right)^{p-1} \|f_2\|_{\text{Lip}}^p \int_0^t \frac{1}{\sqrt{\varepsilon(t-s)}} E \left[\|u_1^\varepsilon(s) - u_2^\varepsilon(s)\|_{p, \phi}^p \right] ds. \end{aligned}$$

Multiplying by $e^{-\beta t}$ and taking the supremum yields

$$(3.7) \quad \|\mathcal{T}_2\|_{\beta,p,L^p(\mathbb{R}^d,\phi)} \leq \delta_{\beta,1} \|f_1 - f_2\|_{\text{Lip}} \|u_1^\varepsilon\|_{\beta,p,L^p(\mathbb{R}^d,\phi)} \\ + \delta_{\beta,1} \|f_2\|_{\text{Lip}} \|u_1^\varepsilon - u_2^\varepsilon\|_{\beta,p,L^p(\mathbb{R}^d,\phi)},$$

where

$$\delta_{\beta,1} = \kappa_{2,d} \sup_{t \in [0,T]} \left(4\sqrt{\frac{t}{\varepsilon}} \right)^{1-1/p} \left(\int_0^t \frac{e^{-\beta(t-s)}}{\sqrt{\varepsilon(t-s)}} ds \right)^{1/p}.$$

Consider $\mathcal{T}_3$. First, observe that

$$|\sigma_1(y, u_1^\varepsilon, z) - \sigma_2(y, u_2^\varepsilon, z)| \leq M_{\sigma_1}(z) |u_1^\varepsilon - u_2^\varepsilon| + M_{\sigma_1 - \sigma_2}(z)(1 + |u_1^\varepsilon|).$$

Due to a simple extension of Lemma 3.3,

$$E \left[\|\mathcal{T}_3(t)\|_{L^p(\mathbb{R}^d)}^p \right]^{1/p} \leq c_p^{1/p} \kappa_{1,d} \|\sigma_1\|_{\text{Lip}} \left(\int_0^t E \left[\|u_1^\varepsilon(s) - u_2^\varepsilon(s)\|_{p,\phi}^p \right]^{2/p} ds \right)^{1/2} \\ + c_p^{1/p} \kappa_{1,d} \|\sigma_1 - \sigma_2\|_{\text{Lip}} \left(\int_0^t E \left[\|1 + |u_1^\varepsilon(s)|\|_{p,\phi}^p \right]^{2/p} ds \right)^{1/2}.$$

Multiplication by $e^{-\beta t/p}$ and then taking the supremum yields

$$(3.8) \quad \|\mathcal{T}_3\|_{\beta,p,L^p(\mathbb{R}^d,\phi)} \leq \delta_{\beta,2} \|\sigma_1\|_{\text{Lip}} \|u_1^\varepsilon - u_2^\varepsilon\|_{\beta,p,L^p(\mathbb{R}^d,\phi)} \\ + \delta_{\beta,2} \|\sigma_1 - \sigma_2\|_{\text{Lip}} (\|\phi\|_{L^1(\mathbb{R}^d)} + \|u_1^\varepsilon\|_{\beta,p,L^p(\mathbb{R}^d,\phi)}),$$

where

$$\delta_{\beta,2} = c_p^{1/p} \kappa_{1,d} \sup_{t \in [0,T]} \left(\int_0^t e^{-\beta 2(t-s)/p} ds \right)^{1/2} \leq c_p^{1/p} \sqrt{\frac{p}{2\beta}}.$$

Here we used that $\|1\|_{\beta,p,L^p(\mathbb{R}^d,\phi)} = \|\phi\|_{L^1(\mathbb{R}^d)}$. Combine (3.6), (3.7), and (3.8), and note that $\delta_{\beta,i} \rightarrow 0$ as $\beta \rightarrow \infty$ for $i = 1, 2$. This concludes the proof. $\square$

In order to apply Itô's formula to the process $t \mapsto u^\varepsilon(t, x)$ we need to know that the weak (mild) solution u^ε of (3.1) is in fact a strong solution. The following result provides the existence of weak derivatives.

Proposition 3.5. *Fix $\phi \in \mathfrak{N}$ and a multiindex α . Make the following assumptions:*

- (i) *The flux-function f belongs to $C^{|\alpha|}(\mathbb{R}; \mathbb{R}^d)$ with all bounded derivatives.*
- (ii) *For each fixed $z \in Z$, $(x, u) \mapsto \sigma(x, u, z)$ belongs to $C^{|\alpha|}(\mathbb{R}^d \times \mathbb{R})$ and for each $0 < \zeta \leq \alpha$ and $0 \leq n \leq |\alpha|$ there exists $M_{\zeta,n} \in L^2(Z)$ such that*

$$\begin{cases} \partial_1^\zeta \partial_2^n \sigma(x, u, z) \leq M_{\zeta,n}(z), & 1 \leq n \leq |\alpha|, \\ \partial_1^\zeta \sigma(x, u, z) \leq M_{\zeta,0}(z)(1 + |u|). \end{cases}$$

- (iii) *The initial function u^0 satisfies for all $\zeta \leq \alpha$,*

$$E \left[\|\partial^\zeta u^0\|_{p,\phi}^p \right] < \infty \quad (2 \leq p < \infty).$$

Let u^ε be the weak solution of (3.1). For any $\zeta \leq \alpha$, there exists a predictable process

$$(t, x, \omega) \mapsto \partial_x^\zeta u^\varepsilon(t, x, \omega) \text{ in } L^p([0, T] \times \Omega; L^p(\mathbb{R}^d, \phi))$$

such that for all $\varphi \in C_c^\infty(\Pi_T)$,

$$\iint_{\Pi_T} \partial_x^\zeta u^\varepsilon \varphi dx dt = (-1)^{|\zeta|} \iint_{\Pi_T} u^\varepsilon \partial_x^\zeta \varphi dx dt, \quad dP\text{-almost surely.}$$

To prove Proposition 3.5 we apply

Lemma 3.6. *Let $\sigma \in C^\infty(\mathbb{R}^d \times \mathbb{R})$ and suppose $u \in C^\infty(\mathbb{R}^d)$. For any multiindex α , let $\partial_x^\alpha := \prod_k \partial_{x_k}^{\alpha_k}$. Then*

$$\partial_x^\alpha \sigma(x, u(x)) = \sum_{\zeta \leq \alpha} \sum_{\gamma \in \pi(\zeta)} C_{\gamma, \alpha} \partial_1^{\alpha - \zeta} \partial_2^{|\gamma|} \sigma(x, u(x)) \prod_{i=1}^{|\gamma|} \partial_x^{\gamma^i} u(x).$$

Here $\pi(\zeta)$ denote all partitions of ζ , i.e., all multiindices $\gamma = \{\gamma^i\}_{i \geq 1}$ such that $\sum \gamma^i = \zeta$. Furthermore, $|\gamma|$ denotes the number of terms in the partition γ .

Remark 3.2. Whenever $\zeta \neq 0$ we assume that the terms γ^i in the partition γ satisfies $\gamma^i \neq 0$. If $\zeta = 0$ we let $\gamma = \gamma^1 = 0$ and by convention let $|\gamma| = 0$.

Proof. One may prove by induction and the chain-rule that

$$\partial_x^\alpha \sigma(x, u(x)) = [(\partial_z + \partial_y)^\alpha \sigma(y, u(z))]_{y=x, z=x}.$$

By the binomial theorem,

$$(\partial_z + \partial_y)^\alpha \sigma(y, u(z)) = \sum_{\zeta \leq \alpha} \binom{\alpha}{\zeta} \partial_y^{\alpha - \zeta} \partial_z^\zeta \sigma(y, u(z)).$$

Thanks to [19, Propositions 1 and 2], it follows that

$$\partial_z^\zeta \sigma(y, u(z)) = \sum_{\gamma \in \pi(\zeta)} M_\gamma \partial_u^{|\gamma|} \sigma(y, u(z)) \prod_i \partial_z^{\gamma^i} u(z),$$

where M_γ is a constant. The result follows by combining the above identities. $\square$

Proof of Proposition 3.5. We divide the proof into two steps.

Step 1 (uniform estimates on $\{u^n\}_{n \geq 1}$). For all $\zeta < \alpha$, suppose

$$(3.9) \quad \sup_{0 \leq s \leq T} E \left[\|\partial^\zeta u^n(s)\|_{p, \phi}^p \right] \leq C_{\zeta, p} \quad (2 \leq p < \infty).$$

We claim that there exists a constant $C \geq 0$, independent of β and n , and a number $\delta_\beta \geq 0$ such that

$$(3.10) \quad \|\partial^\alpha u^{n+1}\|_{\beta, p, L^p(\mathbb{R}^d, \phi)} \leq C + \delta_\beta \|\partial^\alpha u^n\|_{\beta, p, L^p(\mathbb{R}^d, \phi)},$$

where $\delta_\beta < 1$ for some $\beta > 0$, and $\|\cdot\|_{\beta, p, L^p(\mathbb{R}^d, \phi)}$ is defined in (3.5). Given (3.10), it follows that

$$\|\partial^\alpha u^n\|_{\beta, p, L^p(\mathbb{R}^d, \phi)} \leq C \sum_{k=0}^{n-1} \delta_\beta^k \leq \frac{C}{1 - \delta_\beta},$$

and we are done.

To establish (3.10), observe that the weak derivative satisfies

$$\begin{aligned} \partial_x^\alpha u^{n+1}(t, x) &= \int_{\mathbb{R}^d} \Phi_\varepsilon(t, x - y) \partial_y^\alpha u^0(y) dy \\ &\quad - \int_0^t \int_{\mathbb{R}^d} \nabla_x \Phi_\varepsilon(t - s, x - y) \cdot \partial_y^\alpha f(u^n(s, y)) dy ds \\ &\quad + \int_0^t \int_Z \int_{\mathbb{R}^d} \Phi_\varepsilon(t - s, x - y) \partial_y^\alpha \sigma(y, u^n(s, y), z) dy W(ds, dz) \\ &=: \mathcal{T}_1(t, x) + \mathcal{T}_2(t, x) + \mathcal{T}_3(t, x). \end{aligned}$$

To justify this, multiply by a test function and apply the Fubini theorem [39, p.297].

By the triangle inequality we may estimate each term separately.

Consider $\mathcal{T}_1$. By Lemma 3.1,

$$E \left[\|\mathcal{T}_1(t)\|_{p, \phi}^p \right] \leq \kappa_{1, d}^p E \left[\|\partial^\alpha u^0\|_{p, \phi}^p \right],$$

where $\kappa_{1,d}$ is defined in Lemma 3.3. By assumption (iii) it follows that there exists a constant C such that

$$(3.11) \quad \|\mathcal{T}_1\|_{\beta,p,L^p(\mathbb{R}^d,\phi)} \leq C.$$

Consider $\mathcal{T}_2$. By Lemma 3.2,

$$\begin{aligned} \|\mathcal{T}_2\|_{\beta,p,L^p(\mathbb{R}^d,\phi)}^p &= \sup_{t \in [0,T]} e^{-\beta t} E \left[\|\mathcal{T}_2(t)\|_{p,\phi}^p \right] \\ &\leq \kappa_{2,d}^p \left(2\sqrt{\frac{T}{\varepsilon}} \right)^{p-1} \sup_{t \in [0,T]} \int_0^t \frac{e^{-\beta(t-s)}}{\sqrt{\varepsilon(t-s)}} e^{-\beta s} E \left[\|\partial^\alpha f(u^n(s))\|_{p,\phi}^p \right] ds \\ &\leq \kappa_{2,d}^p \left(2\sqrt{\frac{T}{\varepsilon}} \right)^{p-1} \left(\int_0^T \frac{e^{-\beta(t-s)}}{\sqrt{\varepsilon(t-s)}} ds \right) \|\partial^\alpha f(u^n(s))\|_{\beta,p,L^p(\mathbb{R}^d,\phi)}^p. \end{aligned}$$

By Lemma 3.6, the triangle inequality, and the generalized Hölder's inequality,

$$\begin{aligned} E \left[\|\partial^\alpha f(u^n(s))\|_{p,\phi}^p \right]^{1/p} &\leq \sum_{\gamma \in \pi(\alpha)} C_{\gamma,\alpha} \|\partial^{|\gamma|} f\|_\infty E \left[\left\| \prod_{i=1}^{|\gamma|} \partial^{\gamma^i} u^n(s) \right\|_{p,\phi}^p \right]^{1/p} \\ &\leq \sum_{\gamma \in \pi(\alpha)} C_{\gamma,\alpha} \|\partial^{|\gamma|} f\|_\infty \prod_{i=1}^{|\gamma|} E \left[\left\| \partial^{\gamma^i} u^n(s) \right\|_{q_i,\phi}^{q_i} \right]^{1/q_i}, \end{aligned}$$

where $q_i = |\alpha|p/|\gamma_i|$. By assumption there exists a constant C , independent of n , such that

$$E \left[\left\| \partial^{\gamma^i} u^n(s) \right\|_{q_i,\phi}^{q_i} \right] \leq C,$$

for all terms where $\gamma^i < \alpha$. Since $C_{\alpha,\alpha} = 1$, there is another constant C such that

$$E \left[\|\partial^\alpha f(u^n(s))\|_{p,\phi}^p \right]^{1/p} \leq C + \|f'\|_\infty E \left[\|\partial^\alpha u^n(s)\|_{p,\phi}^p \right]^{1/p},$$

for all $n \geq 1$. Multiply by $e^{-\beta t/p}$ and take the supremum to obtain

$$\|\partial^\alpha f(u^n)\|_{\beta,p,L^p(\mathbb{R}^d,\phi)} \leq C + \|f'\|_\infty \|\partial^\alpha u^n\|_{\beta,p,L^p(\mathbb{R}^d,\phi)}.$$

It follows that

$$(3.12) \quad \begin{aligned} \|\mathcal{T}_2\|_{\beta,p,L^p(\mathbb{R}^d,\phi)} &\leq c_d \left(2\sqrt{\frac{T}{\varepsilon}} \right)^{1-1/p} \left(\int_0^T \frac{e^{-\beta(t-s)}}{\sqrt{\varepsilon(t-s)}} ds \right)^{1/p} \\ &\quad \times \left(C + \|f'\|_\infty \|\partial^\alpha u^n\|_{\beta,p,L^p(\mathbb{R}^d,\phi)} \right). \end{aligned}$$

Consider $\mathcal{T}_3$. By Lemma 3.6

$$\begin{aligned} \mathcal{T}_3(t,x) &= \sum_{\zeta \leq \alpha} \sum_{\gamma \in \pi(\zeta)} C_{\gamma,\alpha} \int_0^t \int_Z \int_{\mathbb{R}^d} \Phi_\varepsilon(t-s, x-y) \\ &\quad \times \partial_1^{\alpha-\zeta} \partial_2^{|\gamma|} \sigma(y, u^n(s,y), z) \left(\prod_{i=1}^{|\gamma|} \partial_y^{\gamma^i} u^n(s,y) \right) dy W(ds, dz) \\ &= \mathcal{T}_3^0(t,x) + \mathcal{T}_3^1(t,x), \end{aligned}$$

where $\mathcal{T}_3^0$ contains the term with $\zeta = 0$. By Lemma 3.3, assumption (ii), and the generalised Hölder inequality,

$$E \left[\left\| \mathcal{T}_3^1(t) \right\|_{L^p(\mathbb{R}^d)}^p \right]^{1/p} \leq \sum_{0 < \zeta \leq \alpha} \sum_{\gamma \in \pi(\zeta)} C_{\gamma, \alpha} c_p^{1/p} \kappa_{1,d} \left\| M_{\alpha - \zeta, |\gamma|} \right\|_{L^2(Z)} \\ \times \left(\int_0^t \prod_{i=1}^{|\gamma|} E \left[\left\| \partial^{\gamma^i} u^n(s) \right\|_{L^{q_i}(\mathbb{R}^d)}^{q_i} \right]^{2/q_i} ds \right)^{1/2},$$

where $q_i = |\zeta|p/|\gamma_i|$. The term $\mathcal{T}_3^0$ is estimated similarly by applying the second case of assumption (ii). It follows from (3.9) that there exists a constant C such that

$$E \left[\left\| \mathcal{T}_3(t) \right\|_{p, \phi}^p \right]^{1/p} \leq C + c_p^{1/p} \kappa_{1,d} \left\| M_{0,1} \right\|_{L^2(Z)} \left(\int_0^t E \left[\left\| \partial^\alpha u^n(s) \right\|_{p, \phi}^p \right]^{2/p} ds \right)^{1/2}.$$

Multiplying by $(e^{-\beta t})^{1/p}$ and taking the supremum yields

(3.13)

$$\left\| \mathcal{T}_3 \right\|_{\beta, p, L^p(\mathbb{R}^d, \phi)} \leq C + c_p^{1/p} \kappa_{1,d} \left\| M_{0,1} \right\|_{L^2(Z)} \\ \times \sup_{t \in [0, T]} \left(\int_0^t e^{-2\beta(t-s)/p} \left(e^{-\beta s} E \left[\left\| \partial^\alpha u^n(s) \right\|_{p, \phi}^p \right] \right)^{2/p} ds \right)^{1/2} \\ \leq C + c_p^{1/p} \kappa_{1,d} \left\| M_{0,1} \right\|_{L^2(Z)} \sqrt{\frac{p}{2\beta}} \left\| \partial^\alpha u^n \right\|_{\beta, p, L^p(\mathbb{R}^d, \phi)}.$$

Combining (3.11), (3.12), and (3.13) we obtain inequality (3.10), where

$$\delta_\beta = \kappa_{2,d} \left(2\sqrt{\frac{T}{\varepsilon}} \right)^{1-1/p} \left(\int_0^T \frac{e^{-\beta(t-s)}}{\sqrt{\varepsilon(t-s)}} ds \right)^{1/p} \|f'\|_\infty \\ + c_p^{1/p} \kappa_{1,d} \left\| M_{0,1} \right\|_{L^2(Z)} \sqrt{\frac{p}{2\beta}}.$$

It is clear that $\delta_\beta \rightarrow 0$ as $\beta \rightarrow \infty$ and so (3.10) follows. By induction, estimate (3.9) holds for all $\zeta \leq \alpha$.

Step 2 (convergence of u^n). Fix $\zeta \leq \alpha$. We apply Theorem 6.9 to the family $\{\partial^\zeta u^n\}_{n \geq 1}$ on the space

$$(X, \mathcal{A}, \mu) = (\Omega \times \Pi_T, \mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d), dP \otimes dt \otimes \phi(x)dx).$$

By means of (3.9),

$$\sup_{n \geq 1} \left\{ E \left[\iint_{\Pi_T} |\partial^\zeta u^n(t, x)|^2 \phi(x) dx dt \right] \right\} < \infty.$$

Hence, $\{\partial^\zeta u^n\}_{n \geq 1}$ has a Young measure limit $\nu^\zeta \in \mathcal{Y}(\Omega \times \Pi_T)$. Next, define $\partial^\zeta u^\varepsilon(t, x, \omega) := \int_{\mathbb{R}} d\nu_{t, x, \omega}^\zeta$. By definition, the limit has a $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d)$ measurable version. Furthermore, $\partial^\zeta u^\varepsilon \in L^p(\Omega \times [0, T]; L^p(\mathbb{R}^d, \phi))$, cf. proof of Theorem 4.1 and Lemma 6.8. Let us show that $\partial^\zeta u^\varepsilon$ is the weak derivative of u^ε . To this end, observe that

$$\iint_{\Pi_T} u^n(t, x) \partial^\zeta \varphi dx dt = (-1)^{|\zeta|} \iint_{\Pi_T} \partial^\zeta u^n(t, x) \varphi dx dt,$$

for any $\varphi \in C_c^\infty(\mathbb{R}^d)$. By Lemma 6.11(ii) and Theorem 6.10, there is a subsequence $\{n(j)\}_{j \geq 1}$ such that for any $A \in \mathcal{F}$,

$$\begin{aligned} \lim_{j \rightarrow \infty} E \left[\mathbb{1}_A \iint_{\Pi_T} \partial^\zeta u^{n(j)} \varphi \, dx dt \right] &= E \left[\iint_{\Pi_T} \int_{\mathbb{R}} \varphi(t, x) \mathbb{1}_A(\omega) \, d\nu_{t,x,\omega}^\zeta(\xi) \, dx dt \right] \\ &= E \left[\mathbb{1}_A \iint_{\Pi_T} \partial^\zeta u^\varepsilon \varphi \, dx dt \right]. \end{aligned}$$

As $u^n \rightarrow u^\varepsilon$ in $\mathcal{X}_{\beta,2,L^2(\mathbb{R}^d,\phi)}$, it follows that

$$\lim_{n \rightarrow \infty} E \left[\mathbb{1}_A \iint_{\Pi_T} u^n \partial^\zeta \varphi \, dx dt \right] = E \left[\mathbb{1}_A \iint_{\Pi_T} u^\varepsilon \partial^\zeta \varphi \, dx dt \right].$$

The conclude the proof. $\square$

3.2. Malliavin differentiability. In this subsection we will establish the Malliavin differentiability of the viscous approximation. Furthermore, we observe that the Malliavin derivative satisfies a linear parabolic equation. This equation is then applied to show that $D_{r,z} u^\varepsilon(t, x) \rightarrow \sigma(x, u^\varepsilon(r, x), z)$ as $t \downarrow r$ in a weak sense (Lemma 3.8); a property which is crucial in the proof of uniqueness.

Proposition 3.7 (Malliavin derivative of viscous approximation). *Suppose $(\mathcal{A}_f)$ and $(\mathcal{A}_\sigma)$ are satisfied. Fix $\phi \in \mathfrak{N}$ and $u^0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$. Let u^ε be the solution of (3.1). Then u^ε belongs to $\mathbb{D}^{1,2}(L^2([0, T]; L^2(\mathbb{R}^d, \phi)))$ and*

$$(3.14) \quad \text{ess sup}_{0 \leq t \leq T} \|u^\varepsilon(t)\|_{\mathbb{D}^{1,2}(L^2(\mathbb{R}^d, \phi))} < \infty.$$

Furthermore, for $dr \otimes d\mu$ -a.a. (r, z) , the $L^2(\mathbb{R}^d, \phi)$ -valued process $\{D_{r,z} u^\varepsilon(t)\}_{t > r}$ is a predictable weak solution of

$$(3.15) \quad \begin{cases} dw + \nabla \cdot (f'(u^\varepsilon)w) \, dt = \int_Z \partial_2 \sigma(x, u^\varepsilon, z') w \, W(dt, dz') + \varepsilon \Delta w \, dt, & t \in [r, T], \\ w(r, x, z) = \sigma(x, u^\varepsilon(r, x), z), \end{cases}$$

while $D_{r,z} u^\varepsilon(t) = 0$ if $r > t$. Furthermore

$$(3.16) \quad \text{ess sup}_{r \in [0, T]} \left\{ \sup_{t \in [0, T]} E \|D_r u^\varepsilon(t)\|_{L^2(Z; L^2(\mathbb{R}^d, \phi))}^2 \right\} < \infty.$$

Remark 3.3. Let $w_{r,z}(t, x) = D_{r,z} u^\varepsilon(t, x)$, $t > r$. Estimate (3.16) may be seen as a consequence of the Grönwall-type estimate

$$E \left[\|w_{r,z}(t)\|_{L^2(\mathbb{R}^d, \phi)}^2 \right] \leq \left(1 + C e^{C(t-r)} \right) E \left[\|w_{r,z}(r)\|_{L^2(\mathbb{R}^d, \phi)}^2 \right],$$

for $t \geq r$. From the perspective of a uniqueness result (see Lemma 3.8), it is of interest to know whether one can derive such estimates independent of ε .

Proof. We divide the proof into two steps.

Step 1 (uniform bounds). Consider the Picard approximation $\{u^n\}_{n \geq 1}$ of u^ε . We want to prove that

$$(3.17) \quad \sup_{0 \leq t \leq T} \|u^n(t)\|_{\mathbb{D}^{1,2}(L^2(\mathbb{R}^d, \phi))} \leq C, \quad \text{for all } n \geq 1.$$

By [27, Proposition 1.2.4, Proposition 1.3.8, Proposition 1.2.8],

(3.18)

$$\begin{aligned}
& D_{r,z}u^{n+1}(t, x) \\
&= \int_{\mathbb{R}^d} \Phi_\varepsilon(t-r, x-y) \sigma(y, u^n(r, y), z) dy \\
&\quad - \int_r^t \int_{\mathbb{R}^d} \nabla_x \Phi_\varepsilon(t-s, x-y) \cdot f'(u^n(s, y)) D_{r,z}u^n(s, y) dy ds \\
&\quad + \int_r^t \int_Z \int_{\mathbb{R}^d} \Phi_\varepsilon(t-s, x-y) \partial_2 \sigma(y, u^n(s, y), z') D_{r,z}u^n(s, y) dy W(ds, dz') \\
&=: \mathcal{T}_1^n + \mathcal{T}_2^n + \mathcal{T}_3^n,
\end{aligned}$$

for all $r \in (0, t]$. Whenever $r > t$, $D_r u^{n+1}(t) = 0$ since u^{n+1} is adapted, see [27, Corollary 1.2.1]. Hence, given (3.17), we conclude that $\{D_{r,z}u^\varepsilon\}_{t \geq r}$ is a mild solution of (3.15) for almost all $(r, z) \in (0, t] \times Z$. Next, we apply [30, Theorem 9.15] to infer that the mild solution is also a weak solution.

Let us prove (3.17). Recall that

$$\|u(t)\|_{\mathbb{D}^{1,2}(L^2(\mathbb{R}^d, \phi))}^2 = E \left[\|u(t)\|_{2, \phi}^2 \right] + E \left[\|Du(t)\|_{H \otimes L^2(\mathbb{R}^d, \phi)}^2 \right],$$

where H denotes the space $L^2([0, T] \times Z)$. To obtain a uniform bound we want to establish the existence of a constant C such that

$$(3.19) \quad \|Du^{n+1}\|_{\beta, 2, H \otimes L^2(\mathbb{R}^d, \phi)} \leq C + \delta_\beta \|Du^n\|_{\beta, 2, H \otimes L^2(\mathbb{R}^d, \phi)} \quad \text{where } \delta_\beta < 1,$$

for some $\beta > 0$. Under these assumptions it follows that

$$\|Du^n\|_{\beta, 2, H \otimes L^2(\mathbb{R}^d, \phi)} \leq C \sum_{k=0}^{n-1} \delta_\beta^k \leq \frac{C}{1 - \delta_\beta},$$

and (3.17) follows. To establish (3.19) we proceed by (3.18) to estimate each term separately.

Consider $\mathcal{T}_1^n$. By Lemma 3.1 and assumption $(\mathcal{A}_\sigma)$,

$$\begin{aligned}
\|\mathcal{T}_1^n(r, t)\|_{L^2(Z; L^2(\mathbb{R}^d, \phi))}^2 &= \int_Z \|\Phi_\varepsilon(t-r) \star \sigma(\cdot, u^n(r), z)\|_{2, \phi}^2 d\mu(z) \\
&\leq \kappa_{1,d}^2 \|M\|_{L^2(Z)}^2 (\|\phi\|_{L^1(\mathbb{R}^d)} + \|u^n(r)\|_{2, \phi})^2,
\end{aligned}$$

for each $0 \leq r < t$. It follows that

(3.20)

$$\begin{aligned}
& \|\mathcal{T}_1^n\|_{\beta, 2, H \otimes L^2(\mathbb{R}^d, \phi)} \\
(3.21) \quad & \leq \kappa_{1,d} \|M\|_{L^2(Z)} \left(\sup_{0 \leq t \leq T} e^{-\beta t} E \left[\int_0^t (\|\phi\|_{L^1(\mathbb{R}^d)} + \|u^n(r)\|_{2, \phi})^2 dr \right] \right)^{1/2} \leq C.
\end{aligned}$$

Let us consider $\mathcal{T}_2^n$. By Lemma 3.2,

$$\begin{aligned}
\|\mathcal{T}_2^n(r, t)\|_{L^2(Z; L^2(\mathbb{R}^d, \phi))}^2 &= \int_Z \left\| \int_r^t \nabla \Phi_\varepsilon(t-s) \star f'(u^n(s)) D_{r,z}u^n(s) ds \right\|_{2, \phi}^2 d\mu(z) \\
&\leq 2\kappa_{2,d}^2 \|f'\|_\infty^2 \sqrt{\frac{t-r}{\varepsilon}} \int_r^t \frac{1}{\sqrt{\varepsilon(t-s)}} \int_Z \|D_{r,z}u^n(s)\|_{2, \phi}^2 d\mu(z) ds.
\end{aligned}$$

Multiplication by $e^{-\beta t}$ and integration in r yields

$$\begin{aligned} e^{-\beta t} E \left[\|\mathcal{T}_2^n(t)\|_{H \otimes L^2(\mathbb{R}^d, \phi)}^2 \right] \\ \leq 2\kappa_{2,d}^2 \|f'\|_\infty^2 \sqrt{\frac{t}{\varepsilon}} \int_0^t \frac{e^{-\beta(t-s)}}{\sqrt{\varepsilon(t-s)}} e^{-\beta s} E \left[\|Du^n(s)\|_{H \otimes L^2(\mathbb{R}^d, \phi)}^2 \right] ds. \end{aligned}$$

It follows that

$$(3.22) \quad \|\mathcal{T}_2^n\|_{\beta, 2, H \otimes L^2(\mathbb{R}^d, \phi)} \leq \kappa_{2,d} \|f'\|_\infty \left(2\sqrt{\frac{T}{\varepsilon}} \int_0^T \frac{e^{-\beta(T-s)}}{\sqrt{\varepsilon(T-s)}} ds \right)^{1/2} \times \|Du^n(s)\|_{\beta, 2, H \otimes L^2(\mathbb{R}^d, \phi)}.$$

Consider $\mathcal{T}_3^n$. Due to $(\mathcal{A}_\sigma)$,

$$|\partial_2 \sigma(x, u, z') D_{r,z} u^n(s, y)| \leq M(z') |D_{r,z} u^n(s, y)|.$$

By Lemma 3.3,

$$\begin{aligned} E \left[\|\mathcal{T}_3^n(r, t)\|_{L^2(Z; L^2(\mathbb{R}^d, \phi))}^2 \right] \\ \leq c_2 \kappa_{1,d}^2 \|M\|_{L^2(Z)}^2 \int_r^t E \left[\|D_r u^n(s)\|_{L^2(Z; L^2(\mathbb{R}^d, \phi))}^2 \right] ds. \end{aligned}$$

Integrate in r and multiply by $e^{-\beta t}$ to obtain

$$\begin{aligned} e^{-\beta t} E \left[\|\mathcal{T}_3^n(t)\|_{H \otimes L^2(\mathbb{R}^d, \phi)}^2 \right] \\ \leq c_2 \kappa_{1,d}^2 \|M\|_{L^2(Z)}^2 \int_0^t e^{-\beta(t-s)} e^{-\beta s} E \left[\|Du^n(s)\|_{H \otimes L^2(\mathbb{R}^d, \phi)}^2 \right] ds. \end{aligned}$$

Hence

$$(3.23) \quad \|\mathcal{T}_3^n\|_{\beta, 2, H \otimes L^2(\mathbb{R}^d, \phi)} \leq c_2^{1/2} \kappa_{1,d} \|M\|_{L^2(Z)} \frac{1}{\sqrt{\beta}} \|Du^n\|_{\beta, 2, H \otimes L^2(\mathbb{R}^d, \phi)}.$$

Combining (3.20), (3.22), and (3.23) yields (3.19) with

$$\delta_\beta = \kappa_{2,d} \|f'\|_\infty \left(2\sqrt{\frac{T}{\varepsilon}} \int_0^T \frac{e^{-\beta(T-s)}}{\sqrt{\varepsilon(T-s)}} ds \right)^{1/2} + c_2^{1/2} \kappa_{1,d} \|M\|_{L^2(Z)} \frac{1}{\sqrt{\beta}},$$

where $\kappa_{2,d}$ is the constant from Lemma 3.2, while c_2 and $\kappa_{1,d}$ are the constants from Lemma 3.3. Note that $\delta_\beta \downarrow 0$ as $\beta \rightarrow \infty$. Leaving out the integration in r throughout Step 1, we deduce the estimate

$$(3.24) \quad \|D_r u^n\|_{\beta, 2, L^2(Z) \otimes L^2(\mathbb{R}^d, \phi)} \leq \frac{C}{1 - \delta_\beta}.$$

Step 2 (convergence). Let E denote the space $L^2([0, T]; L^2(\mathbb{R}^d, \phi))$ and recall that $H = L^2(Z \times [0, T])$. Consider $\{u_n\}_{n \geq 1}$ as a sequence in $\mathbb{D}^{1,2}(E)$. By (3.17) and the Hilbert space valued version of [27, Lemma 1.2.3] (see [8, Lemma 5.2]), it follows that u^ε belongs to $\mathbb{D}^{1,2}(E)$ and that $Du^n \rightharpoonup Du^\varepsilon$ in $L^2(\Omega; H \otimes E)$, i.e., for any $h \in H$, $\varphi \in E$, and $V \in L^2(\Omega)$,

$$E [\langle Du^n, h \otimes \varphi \rangle_{H \otimes E} V] \rightarrow E [\langle Du^\varepsilon, h \otimes \varphi \rangle_{H \otimes E} V].$$

It follows that the map

$$(t, \omega) \mapsto Du^\varepsilon(t, \omega) \in L^2(H \otimes L^2(\mathbb{R}^d, \phi))$$

is $\mathscr{P}^*$ -measurable. Note that Lemma 6.8 extends to cover this case so that $(t, \omega) \mapsto D_{r,z} u^\varepsilon(t, \omega)$ is $\mathscr{P}^*$ -measurable $dr \otimes d\mu$ almost all $(r, z) \in [0, T] \times Z$.

For each fixed $t \in [0, T]$, we can conclude by (3.17) as above that $u^\varepsilon(t) \in \mathbb{D}^{1,2}(L^2(\mathbb{R}^d, \phi))$, where $Du^n(t) \rightharpoonup Du^\varepsilon(t)$ along some subsequence. Besides, this

limit agrees dt -almost everywhere with the evaluation of the limit taken in $\mathbb{D}^{1,2}(E)$. This follows by definition for smooth Hilbert space valued random variables and may be extended to the general case by approximation. The weak lower semi-continuity of the norm yields (3.14). Similarly, we may apply (3.24) and the Banach-Alaoglu theorem to extract a weakly convergent subsequence in the space $L^\infty([0, T]; \mathcal{X}_{\beta, 2, L^2(Z) \otimes L^2(\mathbb{R}^d, \phi)})$. This yields the bound (3.16).

As above, $u^\varepsilon(t, x) \in \mathbb{D}^{1,2}$ for $dt \otimes dx$ -almost all (t, x) , and for such (t, x) we have $D_{r,z}u^\varepsilon(t, x) = Du^\varepsilon(t, x, r, z)$ for $d\mu \otimes dr$ almost all (r, z) where $Du^\varepsilon(t, x, r, z)$ denotes the evaluation of the limit taken in $\mathbb{D}^{1,2}(E)$. Taking the Malliavin derivative of (3.3) (as above on u^{n+1}) it follows that $t \mapsto D_{r,z}u^\varepsilon(t)$ is a mild solution of (3.15) for $dr \otimes d\mu$ almost all (r, z) . To conclude that it is a weak solution we can easily verify conditions (F) and (G), with $F(w, t) = \nabla \cdot (f'(u^\varepsilon(t))w)$ and

$$G(w, t)h(x) = \int_Z \partial_2 \sigma(x, u^\varepsilon(t, x), z)w(x)h(z) d\mu(z).$$

□

The next result concerns the limit of $D_{r,z}u^\varepsilon(t, x)$ as $t \downarrow r$. In view of Lemma 3.7, this is a question about the satisfaction of the initial condition for (3.15).

Lemma 3.8. *Let $\phi \in C_c^\infty(\mathbb{R}^d)$ be non-negative. In the setting of Proposition 3.7, for $\Psi \in L^2(\Omega \times Z; L^2(\mathbb{R}^d, \phi))$, set*

$$\mathcal{T}_{r_0}(\Psi) := E \left[\iiint_{Z \times \Pi_T} (D_{r,z}u^\varepsilon(t, x) - \sigma(x, u^\varepsilon(r, x), z)) J_{r_0}^+(t - r) \Psi \phi dt dx d\mu(z) \right].$$

Then there exists a constant C independent of r_0 such that

$$(3.25) \quad |\mathcal{T}_{r_0}(\Psi)| \leq CE \left[\|\Psi\|_{L^2(Z; L^2(\mathbb{R}^d, \phi))}^2 \right]^{1/2}.$$

and $\lim_{r_0 \downarrow 0} \mathcal{T}_{r_0}(\Psi) = 0$ for dr -almost all $r \in [0, T]$.

Proof. Note that $\mathcal{T}_{r_0} = \mathcal{T}_{r_0}^1 - \mathcal{T}_{r_0}^2$, and consider each term separately. By Hölder's inequality,

$$\begin{aligned} |\mathcal{T}_{r_0}^1| &= \left| \int_0^T E \left[\int_Z \int_{\mathbb{R}^d} \Psi(x, z) D_{r,z}u^\varepsilon(t, x) \phi(x) d\mu(z) dx \right] J_{r_0}^+(t - r) dt \right| \\ &\leq \text{ess sup}_{t \in [0, T]} E \left[\int_Z \int_{\mathbb{R}^d} \Psi(x, z) D_{r,z}u^\varepsilon(t, x) \phi(x) d\mu(z) dx \right] \\ &\leq E \left[\|\Psi\|_{L^2(Z; L^2(\mathbb{R}^d, \phi))}^2 \right]^{1/2} \text{ess sup}_{t \in [0, T]} \left\{ E \left[\|D_r u^\varepsilon(t)\|_{L^2(Z; L^2(\mathbb{R}^d, \phi))}^2 \right]^{1/2} \right\}. \end{aligned}$$

Furthermore, due to $(\mathcal{A}_\sigma)$,

$$\begin{aligned} \mathcal{T}_{r_0}^2 &= E \left[\int_Z \int_{\mathbb{R}^d} \Psi(x, z) \sigma(x, u^\varepsilon(r, x), z) \phi(x) dx d\mu(z) \right] \\ &\leq \|M\|_{L^2(Z)} E \left[\|\Psi\|_{L^2(Z; L^2(\mathbb{R}^d, \phi))}^2 \right]^{1/2} E \left[\|1 + |u^\varepsilon(r)|\|_{L^2(\mathbb{R}^d, \phi)}^2 \right]^{1/2}. \end{aligned}$$

The uniform bound (3.25) follows by (3.16) and (3.2). Note that $\Psi \mapsto \mathcal{T}_{r_0}(\Psi)$ is a linear functional on $L^2(\Omega \times Z; L^2(\mathbb{R}^d, \phi))$, for each $r_0 > 0$. By (3.25) the family $\{\mathcal{T}_{r_0}\}_{r_0 > 0}$ is uniformly continuous. Hence, by approximation, it suffices to prove the lemma for Ψ smooth in x with bounded derivatives. Let

$$\varphi(t, x, z) = \Psi(x, z) \phi(x) \xi_{r_0, r}(t), \quad \xi_{r_0, r}(t) = 1 - \int_0^t J_{r_0}^+(\sigma - r) d\sigma.$$

By Proposition 3.7,

$$\begin{aligned}
0 &= \int_{\mathbb{R}^d} \sigma(x, u^\varepsilon(r, x), z) \varphi(r, x, z) dx \\
&+ \int_r^T \int_{\mathbb{R}^d} D_{r,z} u^\varepsilon(t, x) \partial_t \varphi(t, x, z) dx dt \\
&+ \int_r^T \int_{\mathbb{R}^d} f'(u^\varepsilon(t, x)) D_{r,z} u^\varepsilon(t, x) \cdot \nabla \varphi(t, x, z) dx dt \\
&+ \varepsilon \int_r^T \int_{\mathbb{R}^d} D_{r,z} u^\varepsilon(t, x) \Delta \varphi(t, x, z) dx dt \\
&+ \int_r^T \int_Z \int_{\mathbb{R}^d} \partial_2 \sigma(x, u^\varepsilon(t, x), z') D_{r,z} u^\varepsilon(t, x) \varphi(t, x, z) dx W(dz', dt),
\end{aligned}$$

$dr \otimes d\mu \otimes dP$ -almost all (r, z, ω) . Note that

$$\partial_t \varphi(t, x, z) = -\Psi(x, z) \phi(x) J_{r_0}^+(t - r).$$

Taking expectations and integrating in z we obtain

$$\begin{aligned}
\mathcal{T}_{r_0}(\Psi) &= E \left[\int_Z \int_r^T \int_{\mathbb{R}^d} f'(u^\varepsilon(t, x)) D_{r,z} u^\varepsilon(t, x) \cdot \nabla(\Psi \phi) \xi_{r_0}(t, r) dx dt d\mu(z) \right] \\
&+ \varepsilon E \left[\int_Z \int_r^T \int_{\mathbb{R}^d} D_{r,z} u^\varepsilon(t, x) \Delta(\Psi \phi) \xi_{r_0}(t, r) dx dt d\mu(z) \right].
\end{aligned}$$

As $\lim_{r_0 \downarrow 0} \xi_{r_0, r}(t) = 0$ for all $t > r$, it follows by the dominated convergence theorem that $\lim_{r_0 \downarrow 0} \mathcal{T}_{r_0}(\Psi) = 0$. $\square$

4. EXISTENCE OF ENTROPY SOLUTIONS

We will now prove the existence entropy solutions, as defined in Section 2.

Theorem 4.1. *Fix $\phi \in \mathfrak{N}$ and $2 \leq p < \infty$. Suppose $u^0 \in L^p(\Omega, \mathcal{F}_0, P; L^p(\mathbb{R}^d, \phi))$, and $(\mathcal{A}_f)$ and $(\mathcal{A}_\sigma)$ hold. There exists a Young measure-valued entropy solution u of (1.1) in the sense of Definition 2.2. Moreover, $u \in L^p(\Omega \times [0, T]; L^p(\mathbb{R}^d \times [0, 1], \phi))$.*

The bounds in Proposition 3.5 blow up as $\varepsilon \downarrow 0$. Below we establish bounds that are independent of the regularization parameter $\varepsilon > 0$.

Lemma 4.2 (Uniform bounds). *Suppose $(\mathcal{A}_f)$ and $(\mathcal{A}_\sigma)$ hold, and u^0 belongs to $L^p(\Omega, \mathcal{F}_0, P; L^p(\mathbb{R}^d, \phi))$ for some even number $p \geq 2$ and $\phi \in \mathfrak{N}$. Then there exists a constant C , depending on u^0, f, σ, p, T and ϕ but not on ε , such that*

$$(4.1) \quad E \left[\|u^\varepsilon(t)\|_{p, \phi}^p \right] \leq C, \quad t \in [0, T].$$

Proof. Assume that u^0, f, σ satisfy the assumptions of Proposition 3.5 for $|\alpha| \leq 2$. The general result follows by approximation, in view of Proposition 3.4. Let us for the moment also assume $\phi \in \mathfrak{N} \cap C^\infty$, so that there is a constant $C_{\phi, 2}$ satisfying $|\Delta \phi| \leq C_{\phi, 2} \phi$. By Proposition 3.5, u^ε is a strong solution of (3.1). Hence we may apply Itô's formula to the function $S(u) = |u|^p$, cf. Step 2 in the upcoming proof

of Theorem 4.1. After multiplying by ϕ and integrating the result in x ,

$$\begin{aligned} \|u^\varepsilon(t)\|_{p,\phi}^p &= \|u^0\|_{p,\phi}^p \\ &\quad - p \int_0^t \int_{\mathbb{R}^d} |u^\varepsilon(s,x)|^{p-1} \operatorname{sign}(u^\varepsilon(s,x)) (\nabla \cdot f(u^\varepsilon(s,x)) - \varepsilon \Delta u^\varepsilon(s,x)) \phi(x) dx ds \\ &\quad + p \int_0^t \int_Z \int_{\mathbb{R}^d} |u^\varepsilon(s,x)|^{p-1} \operatorname{sign}(u^\varepsilon(s,x)) \sigma(x, u^\varepsilon(s,x), z) \phi(x) dx W(ds, dz) \\ &\quad + \frac{1}{2} p(p-1) \int_0^t \int_{\mathbb{R}^d} |u^\varepsilon(s,x)|^{p-2} \int_Z \sigma^2(x, u^\varepsilon(s,x), z) \phi(x) d\mu(z) dx ds. \end{aligned}$$

Let $q(u) = p \int_0^u |z|^{p-1} \operatorname{sign}(z) \partial f(z) dz$ and note that by $(\mathcal{A}_f)$,

$$(4.2) \quad |q(u)| = \left| \int_0^u p |z|^{p-1} \operatorname{sign}(z) \partial f(z) dz \right| \leq \|f\|_{\operatorname{Lip}} |u|^p.$$

It follows that $q(u^\varepsilon(t))\phi \in L^1(\Omega; L^1(\mathbb{R}^d; \mathbb{R}^d))$ for $0 \leq t \leq T$.

By the chain rule and integration by parts,

$$\begin{aligned} \mathcal{T}_1 &:= \int_0^t \int_{\mathbb{R}^d} p |u^\varepsilon(s,x)|^{p-1} \operatorname{sign}(u^\varepsilon(s,x)) \nabla \cdot f(u^\varepsilon(s,x)) \phi(x) dx ds \\ &= \int_0^t \int_{\mathbb{R}^d} \partial q(u^\varepsilon(s,x)) \cdot \nabla u^\varepsilon(s,x) \phi(x) dx ds \\ &= - \int_0^t \int_{\mathbb{R}^d} q(u^\varepsilon(s,x)) \cdot \nabla \phi(x) dx ds. \end{aligned}$$

By (4.2) and the fact that $\phi \in \mathfrak{N}$,

$$|\mathcal{T}_1| \leq C_\phi \|f\|_{\operatorname{Lip}} \int_0^t \|u^\varepsilon(s)\|_{p,\phi}^p ds.$$

Again by the chain rule and integration by parts,

$$\begin{aligned} \mathcal{T}_2 &:= \varepsilon \int_0^t \int_{\mathbb{R}^d} p |u^\varepsilon(s,x)|^{p-1} \operatorname{sign}(u^\varepsilon(s,x)) \Delta u^\varepsilon(s,x) \phi(x) dx ds \\ &= -\varepsilon p(p-1) \int_0^t \int_{\mathbb{R}^d} |u^\varepsilon(s,x)|^{p-2} |\nabla u^\varepsilon(s,x)|^2 \phi(x) dx ds \\ &\quad - \varepsilon \int_0^t \int_{\mathbb{R}^d} p |u^\varepsilon(s,x)|^{p-1} \operatorname{sign}(u^\varepsilon(s,x)) \nabla u^\varepsilon(s,x) \cdot \nabla \phi(x) dx ds \\ &= -\varepsilon p(p-1) \int_0^t \int_{\mathbb{R}^d} |u^\varepsilon(s,x)|^{p-2} |\nabla u^\varepsilon(s,x)|^2 \phi(x) dx ds \\ &\quad + \varepsilon \int_0^t \int_{\mathbb{R}^d} |u^\varepsilon(s,x)|^p \Delta \phi(x) dx ds. \end{aligned}$$

Using the assumption on ϕ ,

$$|\mathcal{T}_2| \leq C_{\phi,2\varepsilon} \int_0^t \|u^\varepsilon(s)\|_{p,\phi}^p ds.$$

Finally, by assumption $(\mathcal{A}_\sigma)$,

$$\begin{aligned} \mathcal{T}_3 &:= \frac{1}{2} p(p-1) \int_0^t \int_{\mathbb{R}^d} |u^\varepsilon(s,x)|^{p-2} \int_Z \sigma^2(x, u^\varepsilon(s,x), z) \phi(x) d\mu(z) dx ds \\ &\leq \frac{1}{2} p(p-1) \|M\|_{L^2(Z)}^2 \int_0^t \int_{\mathbb{R}^d} |u^\varepsilon(s,x)|^{p-2} (1 + |u^\varepsilon(s,x)|)^2 \phi(x) dx ds \\ &\leq p(p-1) \|M\|_{L^2(Z)}^2 \left(\int_0^t \|u^\varepsilon(s)\|_{p-2,\phi}^{p-2} ds + \int_0^t \|u^\varepsilon(s)\|_{p,\phi}^p ds \right). \end{aligned}$$

After taking expectations and summarizing our findings, we arrive at

$$\begin{aligned} E \left[\|u^\varepsilon(t)\|_{p,\phi}^p \right] &\leq E \left[\|u^0\|_{p,\phi}^p \right] + \underbrace{p(p-1) \|M\|_{L^2(Z)}^2 \int_0^t E \left[\|u^\varepsilon(s)\|_{p-2,\phi}^{p-2} \right] ds}_{C_2} \\ &\quad + \underbrace{\left(C_\phi \|f\|_{\text{Lip}} + \varepsilon C_{\phi,2} + p(p-1) \|M\|_{L^2(Z)}^2 \right) \int_0^t E \left[\|u^\varepsilon(s)\|_{p,\phi}^p \right] ds}_{C_1}, \end{aligned}$$

and hence, appealing to Grönwall's inequality,

$$E \left[\|u^\varepsilon(t)\|_{p,\phi}^p \right] \leq C_2 (1 + C_1 t e^{C_1 t}).$$

Observe that the result holds for p if it also holds for $p-2$, as long as u^0 belongs to $L^p(\Omega; L^p(\mathbb{R}^d, \phi))$. For this reason, (4.1) follows by induction. In view of Lemma 6.5, it takes an easy approximation argument to extend to the case $\phi \in \mathfrak{N}$. $\square$

Remark 4.1. In the forgoing proof, it is certainly possible to apply the Burkholder-Davis-Gundy inequality, resulting in the improvement

$$E \left[\sup_{0 \leq t \leq T} \|u^\varepsilon(t)\|_{L^p(\mathbb{R}^d, \phi)}^p \right] \leq C,$$

for some constant C independent of ε .

Proof of Theorem 4.1. We divide the proof into two main steps.

Step 1 (convergence). We apply Theorem 6.9 to the viscous approximation $\{u^\varepsilon\}_{\varepsilon>0}$ on the measure space

$$(X, \mathcal{A}, \mu) = (\Omega \times \Pi_T, \mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d), dP \otimes dt \otimes \phi(x)dx).$$

By Lemma 4.2,

$$\sup_{\varepsilon>0} \left\{ E \left[\iint_{\Pi_T} |u^\varepsilon(t, x)|^2 \phi(x) dx dt \right] \right\} < \infty,$$

so we may take $\zeta(\xi) = \xi^2$. It follows that there exists a Young measure $\nu = \nu_{t,x,\omega}$ such that for any Carathéodory function $\psi = \psi(u, t, x, \omega)$ satisfying

$$\psi(u^\varepsilon(\cdot), \cdot) \rightharpoonup \bar{\psi}(\cdot) \text{ in } L^1(\Omega \times \Pi_T),$$

we have

$$\bar{\psi}(t, x, \omega) = \int_{\mathbb{R}} \psi(\xi, t, x, \omega) d\nu_{t,x,\omega}(\xi) = \int_0^1 \psi(u(t, x, \alpha, \omega), t, x, \omega) d\alpha.$$

Here $u(t, x, \cdot, \omega)$ is defined through (2.6), i.e.,

$$u(t, x, \alpha, \omega) = \inf \{ \xi \in \mathbb{R} : \nu_{t,x,\omega}((-\infty, \xi]) > \alpha \}.$$

We want to show that the limit u is measurable, i.e. that it has a version $\tilde{u}$ such that for any $\beta \in \mathbb{R}$,

$$B_\beta = \{(t, x, \alpha, \omega) : \tilde{u}(t, x, \alpha, \omega) \geq \beta\} \in \mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d) \otimes \mathcal{B}([0, 1]).$$

Note that

$$\tilde{u}(t, x, \alpha, \omega) \geq \beta \Leftrightarrow \inf_{\xi \in \mathbb{R}} \{ \nu_{t,x,\omega}((-\infty, \xi]) > \alpha \} \geq \beta \Leftrightarrow \nu_{t,x,\omega}((-\infty, \beta]) \leq \alpha.$$

By definition of the Young measure we pick a version (not relabeled) such that, the mapping $(t, x, \omega) \mapsto \nu_{t,x,\omega}((-\infty, \beta])$ is $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable. Furthermore, if it was finitely valued it would be clear that B_β is in the product topology, i.e., $B_\beta \in \mathcal{F} \otimes \mathcal{B}(\Pi_T) \otimes \mathcal{B}([0, 1])$. Hence, the result follows upon approximation by simple functions [11, Example 5.3.1].

Let us show that $u \in L^p([0, T] \times \Omega; L^p(\mathbb{R}^d \times [0, 1], \phi))$. That, is

$$(4.3) \quad E \left[\iint_{\Pi_T} \int_0^1 |u(t, x, \alpha)|^p \phi(x) d\alpha dx dt \right] < \infty.$$

Let $\psi \in C_c^\infty(\mathbb{R})$ be supported in $(-1, 1)$ and satisfy $0 \leq \psi \leq 1, \psi(0) = 1$. Take $\psi_R(u) = \psi(u/R)$. It follows that $\lim_{R \rightarrow \pm\infty} \psi_R(u) = 1$ for each $u \in \mathbb{R}$. Moreover, take $S_R(u) = |u|^p \psi_R(u)$, and note that $S_R(u) \uparrow |u|^p$ for all $u \in \mathbb{R}$. Since S_R is compactly supported it follows that $\{S_R(u^\varepsilon)\}_{\varepsilon>0}$ is uniformly integrable on X . By Theorem 6.10, there is a subsequence $\varepsilon_j \downarrow 0$ such that

$$\begin{aligned} E \left[\iint_{\Pi_T} \int_0^1 S_R(u(t, x, \alpha)) \phi(x) d\alpha dx dt \right] \\ = \lim_{j \rightarrow \infty} E \left[\iint_{\Pi_T} S_R(u^{\varepsilon_j}(t, x)) \phi(x) dx dt \right] \\ \leq \limsup_{\varepsilon \downarrow 0} E \left[\iint_{\Pi_T} |u^\varepsilon(t, x)|^p \phi(x) dx dt \right] \leq C \quad (\text{by Lemma 4.2}), \end{aligned}$$

for some constant C independent of R . The claim (4.3) follows upon letting $R \rightarrow \infty$, applying the monotone convergence theorem.

Step 2 (entropy condition). Let us for the moment assume that f, σ, u^0 satisfy the assumptions of Proposition 3.5 for all multiindices $|\alpha| \leq 2$. Fix an entropy/entropy-flux pair (S, Q) in $\mathcal{E}$, a nonnegative test function $\varphi \in C_c^\infty([0, T] \times \mathbb{R})$, and a random variable $V \in \mathcal{S}$. We want to apply the anticipating Itô formula (Theorem 6.7), for fixed $x \in \mathbb{R}^d$, to $X_t = u^\varepsilon(t, x)$ and $F(X, V, t) = S(X - V)\varphi(t, x)$.

By Proposition 3.5 u^ε is a strong solution of (3.1).

$$\begin{aligned} u^\varepsilon(t, x) = u^0(x) + \int_0^t \varepsilon \Delta u^\varepsilon(s, x) - \nabla \cdot f(u^\varepsilon(s, x)) ds \\ + \int_0^t \int_Z \sigma(x, u^\varepsilon(s, x), z) W(ds, dz), \quad dx \otimes dP\text{-almost surely.} \end{aligned}$$

To this end, consider the weak form (3.4), integrate by parts (cf. Proposition 3.5), and use a (separating) countable subset $\{\varphi_n\}_{n \geq 1} \subset C_c^\infty(\mathbb{R}^d)$ of test functions. The anticipating Itô formula yields, after taking expectations and integrating in x ,

$$\begin{aligned} (4.4) \quad 0 = E \left[\int_{\mathbb{R}^d} S(u^0 - V) \varphi(0) dx \right] \\ + E \left[\int_0^T \int_{\mathbb{R}^d} S(u^\varepsilon(t) - V) \partial_t \varphi(t) dx dt \right] \\ - E \left[\int_0^T \int_{\mathbb{R}^d} \nabla \cdot f(u^\varepsilon(t)) S'(u^\varepsilon(t) - V) \varphi(t) dx dt \right] \\ + E \left[\varepsilon \int_0^T \int_{\mathbb{R}^d} \Delta u^\varepsilon(t) (S'(u^\varepsilon(t) - V) \varphi(t)) dx dt \right] \\ - E \left[\int_0^T \int_{\mathbb{R}^d} \int_Z S''(u^\varepsilon(t) - V) \varphi(t) \sigma(x, u^\varepsilon(t), z) D_{t,z} V d\mu(z) dx dt \right] \\ + \frac{1}{2} E \left[\int_0^T \int_{\mathbb{R}^d} \int_Z S''(u^\varepsilon(t) - V) \varphi(t) \sigma(x, u^\varepsilon(t), z)^2 d\mu(z) dx dt \right], \end{aligned}$$

where $D_{t,z}V$ is the Malliavin derivative of V at (t, z) . By the chain rule and integration by parts, it follows from (4.4) that

$$\begin{aligned}
& E \left[\int_0^T \int_{\mathbb{R}^d} \underbrace{S(u^\varepsilon(t, x) - V) \partial_t \varphi(t, x)}_{\psi_1(u^\varepsilon, \cdot)} + \underbrace{Q(u^\varepsilon(t, x), V) \cdot \nabla \varphi(t, x)}_{\psi_2(u^\varepsilon, \cdot)} dx dt \right] \\
& + E \left[\int_{\mathbb{R}^d} S(u^0(x) - V) \varphi(0, x) dx \right] \\
(4.5) \quad & \geq E \left[\int_0^T \int_{\mathbb{R}^d} \underbrace{S''(u^\varepsilon(t, x) - V) \int_Z \sigma(x, u^\varepsilon(t, x), z) D_{t,z} V d\mu(z) \varphi(t, x)}_{\psi_3(u^\varepsilon, \cdot)} dx dt \right] \\
& - \frac{1}{2} E \left[\int_0^T \int_{\mathbb{R}^d} \underbrace{S''(u^\varepsilon(t, x) - V) \int_Z \sigma^2(x, u^\varepsilon(t, x), z) d\mu(z) \varphi(t, x)}_{\psi_4(u^\varepsilon, \cdot)} dx dt \right] \\
& - \varepsilon E \left[\int_0^T \int_{\mathbb{R}^d} S(u^\varepsilon(t, x) - V) \Delta \varphi(t, x) dx dt \right].
\end{aligned}$$

At this point we may apply Proposition 3.4 to relax the assumptions on f, σ, u^0 to the ones listed in Theorem 4.1, leaving the details to the reader.

Obviously,

$$\lim_{\varepsilon \downarrow 0} E \left[\varepsilon \int_0^T \int_{\mathbb{R}^d} S(u^\varepsilon(t, x) - V) \Delta \varphi(t, x) dx dt \right] = 0.$$

For the remaining terms, it suffices by Step 1 and the upcoming Theorem 6.10 to show that $\{\psi_i(u^\varepsilon, \cdot) \phi^{-1}\}_{\varepsilon > 0}$ is uniformly integrable ($i = 1, 2, 3, 4$).

Regarding ψ_1 , by Lemma 6.11(ii), we must show that

$$(4.6) \quad \sup_{\varepsilon > 0} E \left[\iint_{\Pi_T} |\psi_1(u^\varepsilon(t, x), t, x) \phi^{-1}(x)|^2 \phi(x) dx dt \right] < \infty.$$

As S is in $\mathcal{E}$ and $\varphi \in C_c^\infty(\Pi_T)$,

$$\begin{aligned}
|\psi_1(u^\varepsilon(t, x), t, x) \phi^{-1}(x)| &= |S(u^\varepsilon(t, x) - V) \partial_t \varphi(t, x) \phi^{-1}(x)|^2 \\
&\leq 2 \|S\|_{\text{Lip}}^2 \|\partial_t \varphi(t)\|_{\infty, \phi^{-1}} (|u^\varepsilon(t, x)|^2 + |V|^2).
\end{aligned}$$

So (4.6) follows by Lemma 4.2. The term involving ψ_2 follows along the same lines.

Consider the term involving the Malliavin derivative, namely ψ_3 . As above, $\{\psi_3(u^\varepsilon, \cdot) \phi^{-1}\}_{\varepsilon > 0}$ is uniformly integrable provided

$$\sup_{\varepsilon > 0} E \left[\iint_{\Pi_T} |\psi_3(u^\varepsilon(t, x), t, x) \phi^{-1}(x)|^2 \phi(x) dx dt \right] < \infty.$$

By $(\mathcal{A}_\sigma)$,

$$\left| \int_Z \sigma(x, u^\varepsilon(t, x), z) D_{t,z} V d\mu(z) \right|^2 \leq \|M\|_{L^2(Z)}^2 \|D_t V\|_{L^2(Z)}^2 (1 + |u^\varepsilon(t, x)|)^2.$$

Recall that V is uniformly bounded and also that $\text{supp}(S'') \subset (-R, R)$ for some $R < \infty$. Hence,

$$S''(u^\varepsilon - V)(1 + |u^\varepsilon|) \leq \|S''\|_\infty (1 + R + \|V\|_\infty).$$

Consequently,

$$\begin{aligned} & E \left[\iint_{\Pi_T} |\psi_3(u^\varepsilon(t, x), t, x) \phi^{-1}(x)|^2 \phi(x) dx dt \right] \\ & \leq \|S''\|_\infty^2 \|M\|_{L^2(Z)}^2 (1+R+\|V\|_\infty)^2 \|\varphi\|_{\infty, \phi^{-1}}^2 E \left[\int_0^T \|D_t V\|_{L^2(Z)}^2 dt \right] \|\phi\|_{L^1(\mathbb{R}^d)}. \end{aligned}$$

Consider the ψ_4 -term. By $(\mathcal{A}_\sigma)$

$$S''(u^\varepsilon - V) \int_Z \sigma^2(x, u^\varepsilon, z) d\mu(z) \leq \|S''\|_\infty \|M\|_{L^2(Z)}^2 (1 + R + \|V\|_\infty)^2.$$

Hence

$$\begin{aligned} & E \left[\iint_{\Pi_T} |\psi_4(u^\varepsilon(t, x), t, x) \phi^{-1}(x)|^2 \phi(x) dx dt \right] \\ & \leq \|S''\|_\infty^2 \|M\|_{L^2(Z)}^4 (1 + R + \|V\|_\infty)^4 \|\varphi\|_{\infty, \phi^{-1}}^2 \iint_{\Pi_T} \phi(x) dx dt. \end{aligned}$$

Summarizing, upon sending $\varepsilon \downarrow 0$ along a subsequence, it follows that

$$\mathcal{Y}(\mathbf{Ent}[(S, Q), \varphi, V])(u) \geq 0,$$

where u is the process defined in Step 1. Finally, the result follows for general $V \in \mathbb{D}^{1,2}$ by the density of $\mathcal{S} \subset \mathbb{D}^{1,2}$ and Lemma 2.2. $\square$

5. UNIQUENESS OF ENTROPY SOLUTIONS

To prove the uniqueness of Young measure-valued entropy solutions, we need an additional assumption on σ : there exists $M \in L^2(Z)$ and $\kappa > 0$ such that

$$(\mathcal{A}_{\sigma,1}) \quad |\sigma(x, u, z) - \sigma(y, u, z)| \leq M(z) |x - y|^{\kappa+1/2} (1 + |u|),$$

for $x, y \in \mathbb{R}^d$ and $u \in \mathbb{R}$. Actually, it suffices that the criterion is satisfied locally, i.e., for each compact $K \subset \mathbb{R}^d \times \mathbb{R}^d$ there exists $M = M_K$ such that $(\mathcal{A}_{\sigma,1})$ is satisfied for all $(x, y) \in K$.

Theorem 5.1. *Fix $\phi \in \mathfrak{N}$, and suppose $u^0 \in L^2(\Omega, P, \mathcal{F}_0; L^2(\mathbb{R}^d, \phi))$. Under assumptions $(\mathcal{A}_f)$, $(\mathcal{A}_\sigma)$, and $(\mathcal{A}_{\sigma,1})$ there exists but one Young measure-valued entropy solution to (1.1) in the sense of Definition 2.2. Moreover, this solution is an entropy solution in the sense of Definition 2.1.*

The proof is found at the end of this section. As discussed in the introduction, due to the lack of Malliavin differentiability at the hyperbolic level, the uniqueness argument will invoke the viscous approximations and their limit taken in the weak sense of Young measures.

Retracing the proof of Theorem 5.1, making some small modifications, we obtain the following spatial regularity result:

Proposition 5.2 (Spatial regularity). *Fix $\phi \in \mathfrak{N}$, and suppose u^0 belongs to $L^2(\Omega, P, \mathcal{F}_0; L^2(\mathbb{R}^d, \phi))$. Under assumptions $(\mathcal{A}_f)$, $(\mathcal{A}_\sigma)$, and $(\mathcal{A}_{\sigma,1})$ the entropy solution u to (1.1) satisfies*

$$\begin{aligned} & E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u(t, x+z) - u(t, x-z)| \phi(x) J_r(z) dx dz \right] \\ & \leq CE \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u^0(x+z) - u^0(x-z)| \phi(x) J_r(z) dx dz \right] + \mathcal{O}(r^\kappa), \end{aligned}$$

where the constant C depends only on $C_\phi, \|f\|_{\text{Lip}}, T$, and κ is the exponent from assumption $(\mathcal{A}_{\sigma,1})$.

See [14, 9] for similar results, and how to turn this result into a fractional BV estimate. The proof of Proposition 5.2 is found at the very end of the section.

The next lemma contains the “entropy condition” at the parabolic level, which will be utilized later in the uniqueness proof.

Lemma 5.3. *For each fixed $\varepsilon > 0$, let u^ε be the solution of (3.1). Suppose $V \in L^2(\Omega)$ is $\mathcal{F}_s$ -measurable for some $s \in (0, T)$, and $0 \leq \varphi \in C_c^\infty([0, T] \times \mathbb{R}^d)$ with $\text{supp}(\varphi) \subset (s, T) \times \mathbb{R}^d$. Then*

$$\begin{aligned} & E \left[\iint_{\Pi_T} S(u^\varepsilon - V) \partial_t \varphi + Q(u^\varepsilon, V) \cdot \nabla \varphi \, dx dt d\alpha \right] \\ & \geq -\frac{1}{2} E \left[\iint_{\Pi_T} \int_Z S''(u^\varepsilon - V) \sigma(x, u^\varepsilon, z)^2 \varphi(t, x) \, d\mu(z) \, dx dt d\alpha \right] \\ & \quad - \varepsilon E \left[\iint_{\Pi_T} S(u^\varepsilon(t) - V) \Delta \varphi(t) \, dx dt \right], \end{aligned}$$

for any entropy/entropy-flux pair (S, Q) in $\mathcal{E}$.

Proof. Consider (4.5). Note that for any $V \in \mathcal{S}_b$ that is $\mathcal{F}_s$ -measurable,

$$E \left[\iint_{\Pi_T} \int_Z S''(u^\varepsilon(t) - V) \sigma(x, u^\varepsilon(t), z) D_{t,z} V \varphi(t) \, d\mu(z) \, dx dt \right] = 0,$$

thanks to [27, Proposition 1.2.8]. The general result follows by approximation as in Lemma 2.2. $\square$

The following “doubling of variables” lemma is at the heart of the matter. To some extent it may be instructive to compare its proof with the rather involved computations in [17, Lemma 3.2] and [3, Section 4.1].

Lemma 5.4. *Suppose $(\mathcal{A}_f), (\mathcal{A}_\sigma)$ hold. Fix $\phi \in \mathfrak{N}$, and let $\{u^\varepsilon\}_{\varepsilon>0}$ be a sequence of viscous approximations with initial condition $u^0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$. Let v be a Young measure-valued entropy solution in the sense of Definition 2.1 with initial condition $v^0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$.*

For any $0 < \gamma < \frac{1}{2}T$ take $t_0 \in [0, T - 2\gamma]$ and define

$$\xi_{\gamma, t_0}(t) := 1 - \int_0^t J_\gamma^+(s - t_0) \, ds.$$

Let $\psi \in C_c^\infty(\mathbb{R}^d)$ be non-negative and define

$$\varphi(t, x, s, y) = \frac{1}{2^d} \psi\left(\frac{x+y}{2}\right) J_r\left(\frac{x-y}{2}\right) \xi_{\gamma, t_0}(t) J_{r_0}^+(t-s).$$

Let S_δ be a function satisfying

$$S'_\delta(\sigma) = 2 \int_0^\sigma J_\delta(z) \, dz, \quad S_\delta(0) = 0.$$

Furthermore, define

$$Q_\delta(u, c) = \int_c^u S'_\delta(z - c) f'(z) \, dz,$$

and note that the pair (S_δ, Q_δ) belongs to $\mathcal{E}$.

Then

$$(5.1) \quad L \geq R + F + \mathcal{T}_1 + \mathcal{T}_2 + \mathcal{T}_3,$$

where

$$\begin{aligned}
L &= E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} S_\delta(v^0(y) - u^\varepsilon(t, x)) \varphi(t, x, 0, y) dy dx dt \right], \\
R &= -E \left[\iiint_{\Pi_L^2} \int_{[0,1]} S_\delta(v - u^\varepsilon) (\partial_s + \partial_t) \varphi d\beta dX \right], \\
F &= -E \left[\iiint_{\Pi_L^2} \int_{[0,1]} Q_\delta(u^\varepsilon, v) \cdot \nabla_x \varphi + Q_\delta(v, u^\varepsilon) \cdot \nabla_y \varphi d\beta dX \right], \\
\mathcal{T}_1 &= -\frac{1}{2} E \left[\iiint_{\Pi_L^2} \int_{[0,1]} \int_Z S''_\delta(v - u^\varepsilon) (\sigma(y, v, z) - \sigma(x, u^\varepsilon, z))^2 \varphi d\mu(z) d\beta dX \right], \\
\mathcal{T}_2 &= E \left[\iiint_{\Pi_L^2} \int_{[0,1]} \int_Z S''_\delta(v - u^\varepsilon) (D_{s,z} u^\varepsilon - \sigma(x, u^\varepsilon, z)) \sigma(y, v, z) \varphi d\mu(z) d\beta dX \right], \\
\mathcal{T}_3 &= -\varepsilon E \left[\iiint_{\Pi_L^2} \int_{[0,1]} S_\delta(u^\varepsilon - v) \Delta_x \varphi d\beta dX \right],
\end{aligned}$$

where $dX = dx dt dy ds$.

Remark 5.1. In [17, Section 4.6] the authors prove existence of a *strong* entropy solution. The additional condition attached to the notion of strong solution stems from the difficulties in sending $\varepsilon \downarrow 0$ before $r_0 \downarrow 0$. In our setting, the existence of a strong entropy solution amounts to showing that we can send $r_0 \downarrow 0$ and $\varepsilon \downarrow 0$ simultaneously in such a way that $\lim_{(\varepsilon, t) \downarrow (0, s)} \mathcal{T}_2 = 0$. This requires a careful study of how the continuity properties of (3.15) depends on ε , cf. Lemma 3.8. We do not proceed along this path in this paper, instead we let $r_0 \downarrow 0$ before $\varepsilon \downarrow 0$ as in [3].

Proof. Recall that $\text{supp}(J_{r_0}^+) \subset (0, 2r_0)$, so $J_{r_0}^+(t - s)$ is zero whenever $s \geq t$. Applying Lemma 5.3 with $V = v(s, y, \beta)$ and integrating in y, s, β , we obtain

$$\begin{aligned}
& E \left[\iiint_{\Pi_T^2} \int_{[0,1]} S_\delta(u^\varepsilon - v) \partial_t \varphi + Q_\delta(u^\varepsilon, v) \cdot \nabla \varphi d\beta dX \right] \\
(5.2) \quad & \geq -\frac{1}{2} E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \int_Z S''_\delta(u^\varepsilon - v) \sigma(x, u^\varepsilon, z)^2 \varphi d\mu(z) d\beta dX \right] \\
& \quad - \varepsilon E \left[\iiint_{\Pi_T^2} \int_{[0,1]} S_\delta(u^\varepsilon - v) \Delta_x \varphi d\beta dX \right].
\end{aligned}$$

Similarly, in the entropy inequality for $v = v(s, y, \beta)$ we take $V = u^\varepsilon(t, x)$ and integrate in t, x , resulting in

$$\begin{aligned}
& E \left[\iiint_{\Pi_T^2} \int_{[0,1]} S_\delta(v - u^\varepsilon) \partial_s \varphi + Q_\delta(v, u^\varepsilon) \cdot \nabla_y \varphi d\beta dX \right] \\
& + E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} S_\delta(v^0(y) - u^\varepsilon(t, x)) \varphi(t, x, 0, y) dy dx dt \right] \\
(5.3) \quad & \geq E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \int_Z S''_\delta(v - u^\varepsilon) D_{s,z} u^\varepsilon \sigma(y, v, z) \varphi d\mu(z) d\beta dX \right] \\
& \quad - \frac{1}{2} E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \int_Z S''_\delta(v - u^\varepsilon) \sigma(y, v, z)^2 \varphi d\mu(z) d\beta dX \right].
\end{aligned}$$

The result follows by adding (5.2) and (5.3). $\square$

Proposition 5.5 (Kato inequality). *Fix $\phi \in \mathfrak{N}$. Suppose $(\mathcal{A}_f)$, $(\mathcal{A}_\sigma)$, and $(\mathcal{A}_{\sigma,1})$ hold. Let u be the Young measure-valued limit of the viscous approximations $\{u^\varepsilon\}_{\varepsilon>0}$ with initial condition $u^0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$, constructed in Theorem 4.1. Let v be a Young measure-valued entropy solution in the sense of Definition 2.1 with initial condition $v^0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$. Then, for almost all $t_0 \in (0, T)$ and any non-negative $\psi \in C_c^\infty(\mathbb{R}^d)$,*

$$(5.4) \quad \begin{aligned} & E \left[\int_{\mathbb{R}^d} \iint_{[0,1]^2} |u(t_0, x, \alpha) - v(t_0, x, \beta)| \psi(x) d\alpha d\beta dx \right] \\ & \leq E \left[\int_{\mathbb{R}^d} |u^0(x) - v^0(x)| \psi(x) dx \right] \\ & + E \left[\int_0^{t_0} \int_{\mathbb{R}^d} \iint_{[0,1]^2} \text{sign}(u(t, x, \alpha) - v(t, x, \beta)) \right. \\ & \quad \left. \times (f(u(t, x, \alpha)) - f(v(t, x, \beta))) \cdot \nabla \psi(x) d\beta d\alpha dx dt \right]. \end{aligned}$$

Proof. Starting off from (5.1), we send r_0 and ε to zero (in that order). Next, we send (δ, r) to $(0, 0)$ simultaneously. In view of Limits 3 and 4, we let $\delta(r) = r^{1+\eta}$ with $0 < \eta < 2\kappa - 1$. Finally, we send $\gamma \downarrow 0$. We arrive at the Kato inequality (5.4) thanks to the upcoming Limits 1–6. $\square$

Remark 5.2. Later we will make repeated use of two elementary identities. Set

$$\xi_r(x) := \frac{1}{2^d} \int_{\mathbb{R}^d} \psi\left(\frac{x+y}{2}\right) J_r\left(\frac{x-y}{2}\right) dy.$$

Then note that $\xi_r = \psi \star J_r$. Indeed, making the change of variable $z = (x+y)/2$, it follows that $(x-y)/2 = x-z$ and $dy = 2^d dz$. Next, consider the change of variables

$$\Phi(x, y) = \left(\frac{x+y}{2}, \frac{x-y}{2} \right) = (\tilde{x}, z).$$

By the change of variables formula

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} g(\tilde{x}, z) d\tilde{x} dz = \iint_{\mathbb{R}^d \times \mathbb{R}^d} g(\Phi(x, y)) |\det(\partial\Phi(x, y))| dx dy,$$

for any measurable function $g(\cdot, \cdot)$. A computation yields $|\det(\partial\Phi(x, y))| = 1/2^d$. It follows that

$$\begin{aligned} & \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \underbrace{h(x, y) \psi\left(\frac{x+y}{2}\right) J_r\left(\frac{x-y}{2}\right)}_{g(\Phi(x, y))} dx dy \\ & = \iint_{\mathbb{R}^d \times \mathbb{R}^d} \underbrace{h(\tilde{x} + z, \tilde{x} - z) \psi(\tilde{x}) J_r(z)}_{g(\tilde{x}, z)} d\tilde{x} dz, \end{aligned}$$

for any measurable function $h(\cdot, \cdot)$. Most of the time we drop the tilde and write x instead of $\tilde{x}$.

Limit 1. With L defined in Lemma 5.4,

$$\lim_{r_0 \downarrow 0} L = E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(v^0(x-z) - u^0(x+z)) \psi(x) J_r(z) dx dz \right].$$

If $\delta = \delta(r)$ is a nondecreasing function satisfying $\delta(r) \downarrow 0$ as $r \downarrow 0$, then

$$\lim_{\gamma, (\delta, r), \varepsilon, r_0 \downarrow 0} L = E \left[\|v^0 - u^0\|_{1, \psi} \right].$$

Proof. Note that

$$(5.5) \quad |S_\delta(b) - S_\delta(a)| = \left| \int_a^b S'_\delta(z) dz \right| \leq |b - a|.$$

Hence, due to Remark 5.2,

$$\begin{aligned} & \left| L - E \left[\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} S_\delta(v^0(y) - u^0(x)) \frac{1}{2^d} \psi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) dx dy \right] \right| \\ & \leq E \left[\iint_{\Pi_T} |u^0(x) - u^\varepsilon(t, x)| J_{r_0}^+(t) (\psi \star J_r)(x) dx dt \right]. \end{aligned}$$

Arguing as in Lemma 2.3 for the viscous approximation, it follows that

$$\begin{aligned} \lim_{r_0 \downarrow 0} L &= E \left[\frac{1}{2^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} S_\delta(v^0(y) - u^0(x)) \psi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) dx dy \right] \\ &= E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(v^0(x-z) - u^0(x+z)) \psi(x) J_r(z) dx dz \right]. \end{aligned}$$

This proves the first limit. The second limit follows by the dominated convergence theorem and Lemma 6.1. $\square$

Remark 5.3. To establish Limits 2 and 3 we need to send $\varepsilon \downarrow 0$ in terms of the form

$$E \left[\iiint_{\Pi_T \times \mathbb{R}^d \times [0,1]} \Psi(u^\varepsilon(t, x, \omega), t, x, y, \beta, \omega) \underbrace{\frac{1}{2^d} \phi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) d\beta dy dx dt}_{d\eta_{\phi,r}} \right],$$

where Ψ is continuous in the first variable. Essentially we proceed as in the proof of Theorem 4.1, but now the underlying measure space is $\Pi_T \times \mathbb{R}^d \times [0, 1] \times \Omega$ instead of $\Pi_T \times \Omega$. By Lemma 4.2 and Remark 5.2,

$$\begin{aligned} & \sup_{\varepsilon > 0} \left\{ E \left[\iiint_{\Pi_T \times \mathbb{R}^d \times [0,1]} |u^\varepsilon(t, x)|^2 d\eta_{\phi,r} \right] \right\} \\ & = \sup_{\varepsilon > 0} \left\{ E \left[\int_0^T \|u^\varepsilon(t)\|_{2, \phi \star J_r}^2 dt \right] \right\} < \infty. \end{aligned}$$

By Theorem 6.9, there exists $\nu \in \mathcal{Y}(\Pi_T \times \mathbb{R}^d \times [0, 1] \times \Omega)$ such that whenever $\Psi(u^\varepsilon, \cdot) \rightharpoonup \bar{\Psi}$ along some subsequence in $L^1(\Pi_T \times \mathbb{R}^d \times [0, 1] \times \Omega, d\eta_{\phi,r} \otimes dP)$,

$$\bar{\Psi} = \int_{\mathbb{R}} \Psi(\xi, t, x, y, \beta, \omega) d\nu_{t,x,\omega}(\xi) = \int_0^1 \Psi(u(t, x, \alpha, \omega), t, x, y, \beta, \omega) d\alpha,$$

where u is defined through (2.6). The fact that $\nu_{t,x,y,\beta,\omega} = \nu_{t,x,\omega}$ comes out since the limit is independent of y, β when Ψ is independent of y, β . For measurability considerations, see Step 1 in proof of Theorem 4.1.

Limit 2. With R defined in Lemma 5.4,

$$\lim_{\gamma, \varepsilon, r_0 \downarrow 0} R = E \left[\iiint_{\mathbb{R}^d \times \mathbb{R}^d \times [0,1]^2} S_\delta(v(t_0, x-z, \beta) - u(t_0, x+z, \alpha)) \psi(x) J_r(z) d\alpha d\beta dx dz \right],$$

for dt -a.a. $t_0 \in [0, T]$. If $\delta = \delta(r)$ is a nondecreasing function satisfying $\delta(r) \downarrow 0$ as $r \downarrow 0$, then

$$\lim_{\gamma, (\delta, r), \varepsilon, r_0 \downarrow 0} R = E \left[\int_{\mathbb{R}^d} \iint_{[0,1]^2} |v(t_0, x, \beta) - u(t_0, x, \alpha)| \psi(x) d\alpha d\beta dx \right],$$

for dt -a.a. $t_0 \in [0, T]$.

Proof. Since

$$(\partial_s + \partial_t)\varphi(t, x, s, y) = -\frac{1}{2^d}\psi\left(\frac{x+y}{2}\right)J_r\left(\frac{x-y}{2}\right)J_\gamma^+(t-t_0)J_{r_0}^+(t-s),$$

it follows that

$$R = E \left[\frac{1}{2^d} \iiint_{\Pi_T^2} \int_{[0,1]} S_\delta(v - u^\varepsilon) \psi\left(\frac{x+y}{2}\right) J_r\left(\frac{x-y}{2}\right) \times J_\gamma^+(t-t_0) J_{r_0}^+(t-s) d\beta dX \right].$$

Thanks to

$$|S_\delta(v - u^\varepsilon)| \leq |v| + |u^\varepsilon|,$$

we can apply the dominated convergence theorem and Lemma 6.2, resulting in

$$\lim_{r_0 \downarrow 0} R = E \left[\iiint_{\Pi_T \times \mathbb{R}^d \times [0,1]} \underbrace{S_\delta(v(t, y, \beta) - u^\varepsilon(t, x)) J_\gamma^+(t-t_0) (\psi \phi^{-1})\left(\frac{x+y}{2}\right)}_{\Psi(u^\varepsilon, \cdot)} \times \frac{1}{2^d} \phi\left(\frac{x+y}{2}\right) J_r\left(\frac{x-y}{2}\right) d\beta dy dx dt \right].$$

By Lemma 6.11(ii), $\{\Psi_\varepsilon(u^\varepsilon, \cdot)\}$ is uniformly integrable, and so, cf. Theorem 6.10, we can extract a weakly convergent subsequence. By Remark 5.3 and 5.2

$$\begin{aligned} \lim_{\varepsilon, r_0 \downarrow 0} R &= E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} \iint_{[0,1]^2} S_\delta(v(t, y, \beta) - u(t, x, \alpha)) J_\gamma^+(t-t_0) \right. \\ &\quad \left. \times \frac{1}{2^d} \psi\left(\frac{x+y}{2}\right) J_r\left(\frac{x-y}{2}\right) d\alpha d\beta dy dx dt \right] \\ &= E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} \iint_{[0,1]^2} S_\delta(v(t, x-z, \beta) - u(t, x+z, \alpha)) \right. \\ &\quad \left. \times J_\gamma^+(t-t_0) \psi(x) J_r(z) d\alpha d\beta dz dx dt \right] \end{aligned}$$

Note that

$$|S_\delta(a-b) - S_\delta(c-d)| \leq |b-d| + |a-c|, \quad a, b, c, d \in \mathbb{R}.$$

Applying this inequality and Lemma 6.2, we can send $\gamma \downarrow 0$ to obtain the first inequality. To send $(\delta, r) \downarrow (0, 0)$ we apply the dominated convergence theorem and Lemma 6.1, yielding

$$\lim_{r, \varepsilon, r_0 \downarrow 0} R = E \left[\iint_{\Pi_T} \iint_{[0,1]^2} |v(t, x, \beta) - u(t, x, \alpha)| \psi(x) J_\gamma^+(t-t_0) d\alpha d\beta dx dt \right].$$

To send $\gamma \downarrow 0$ we apply Lemma 6.2. This provides the second limit. $\square$

Limit 3. With F defined in Lemma 5.4,

$$(5.6) \quad \lim_{\gamma, \varepsilon, r_0 \downarrow 0} F_1 = E \left[\int_0^{t_0} \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times [0,1]^2} S'_\delta(u(t, x+z, \alpha) - v(t, x-z, \beta)) \right. \\ \left. \times (f(u(t, x+z, \alpha)) - f(v(t, x-z, \beta))) \cdot \nabla \psi(x) J_r(z) d\alpha d\beta dx dz dt \right] \\ + \mathcal{O} \left(\delta + \frac{\delta}{r} \right).$$

If $\delta : [0, \infty) \rightarrow [0, \infty)$ satisfy $\lim_{r \rightarrow 0} \frac{\delta(r)}{r} = 0$, then

$$(5.7) \quad \lim_{\gamma, r, \varepsilon, r_0 \downarrow 0} F = -E \left[\int_0^{t_0} \int_{\mathbb{R}^d} \iint_{[0,1]^2} \text{sign}(u(t, x, \alpha) - v(t, x, \beta)) \right. \\ \left. \times (f(u(t, x, \alpha)) - f(v(t, x, \beta))) \cdot \nabla \psi(x) d\alpha d\beta dx dt \right].$$

Proof. Using integration by parts,

$$Q_\delta(u^\varepsilon, v) = S'_\delta(u^\varepsilon - v)(f(u^\varepsilon) - f(v)) - \int_v^{u^\varepsilon} S''_\delta(z - v)(f(z) - f(v)) dz$$

and

$$Q_\delta(v, u^\varepsilon) = S'_\delta(v - u^\varepsilon)(f(v) - f(u^\varepsilon)) - \int_{u^\varepsilon}^v S''_\delta(z - u^\varepsilon)(f(z) - f(u^\varepsilon)) dz.$$

Due to the symmetry of S_δ ,

$$F = -E \left[\iiint_{\Pi_T^2} \int_{[0,1]} Q_\delta(u^\varepsilon, v) \cdot \nabla_x \varphi + Q_\delta(v, u^\varepsilon) \cdot \nabla_y \varphi d\beta dX \right] \\ = -E \left[\iiint_{\Pi_T^2} \int_{[0,1]} S'_\delta(u^\varepsilon - v)(f(u^\varepsilon) - f(v)) \cdot (\nabla_x + \nabla_y) \varphi d\beta dX \right] \\ + E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \left(\int_v^{u^\varepsilon} S''_\delta(z - v)(f(z) - f(v)) dz \right) \cdot \nabla_x \varphi d\beta dX \right] \\ + E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \left(\int_{u^\varepsilon}^v S''_\delta(z - u^\varepsilon)(f(z) - f(u^\varepsilon)) dz \right) \cdot \nabla_y \varphi d\beta dX \right] \\ = -F_1 + F_2 + F_3,$$

where $dX = dx dt dy ds$ as in Lemma 5.4. Note that

$$(5.8) \quad \left| \int_v^u S''_\delta(z - v)(f(z) - f(v)) dz \right| \leq \|f\|_{\text{Lip}} \delta, \quad u, v \in \mathbb{R}.$$

To see this, recall that $S''_\delta(\sigma) = 2J_\delta(\sigma)$. By $(\mathcal{A}_f)$,

$$\left| \int_v^u S''_\delta(z - v)(f(z) - f(v)) dz \right| \leq 2 \|f\|_{\text{Lip}} \text{sign}(u - v) \int_v^u J_\delta(z - v) |z - v| dz,$$

and letting $\xi = |z - v|/\delta$,

$$\text{sign}(u - v) \int_v^u J_\delta(z - v) |z - v| dz = \delta \int_0^{\delta^{-1}|u-v|} J(\xi) \xi d\xi \leq \frac{\delta}{2}.$$

In view of (5.8), it is clear that

$$F_2 \leq \|f\|_{\text{Lip}} \delta \iiint_{\Pi_T^2} |\nabla_x \varphi| dX.$$

A computation shows $\|\nabla\varphi\|_{L^1(\Pi_T^2)} \leq C(1+r^{-1})$, for some constant C depending only on J, T, ψ . Consequently,

$$F_2 \leq C \|f\|_{\text{Lip}} \delta \left(1 + \frac{1}{r}\right).$$

The same type of estimate applies to F_3 .

Let us consider F_1 . Observe that

$$(\nabla_x + \nabla_y)\varphi(t, x, s, y) = \frac{1}{2^d} \nabla\psi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) \xi_{\gamma, t_0}(t) J_{r_0}^+(t-s).$$

For $\delta > 0$, define

$$\mathcal{F}_\delta(a, b) := S'_\delta(a-b)(f(a) - f(b)), \quad a, b \in \mathbb{R},$$

and note that $(t, b) \mapsto \mathcal{F}_\delta(u^\varepsilon(t, x), b)$ obeys the hypotheses of Lemma 6.2. By the dominated convergence theorem and Lemma 6.2,

$$\begin{aligned} \lim_{r_0 \downarrow 0} F_1 = E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} \int_{[0,1]} \underbrace{\mathcal{F}_\delta(u^\varepsilon(t, x), v(t, y, \beta)) \cdot \zeta \left(\frac{x+y}{2} \right) \xi_{\gamma, t_0}(t)}_{\Psi(u^\varepsilon, \cdot)} \right. \\ \left. \times \frac{1}{2^d} \phi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) d\beta dy dx dt \right], \end{aligned}$$

where $\zeta(x) = \phi^{-1}(x) \nabla\psi(x)$. The uniform integrability of $\{\Psi(u^\varepsilon, \cdot)\}_{\varepsilon > 0}$ follows thanks to Lemma 6.11(ii). Indeed, $|\zeta| \leq C_\phi$ and $|\mathcal{F}_\delta(u^\varepsilon, v)| \leq \|f\|_{\text{Lip}} |u^\varepsilon - v|$, so

$$|\Psi(u^\varepsilon, \cdot)|^2 \leq 2C_\phi^2 \|f\|_{\text{Lip}}^2 (|u^\varepsilon|^2 + |v|^2).$$

By Theorem 6.10 and Remark 5.3

$$\begin{aligned} \lim_{\varepsilon, r_0 \downarrow 0} F_1 = E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} \iint_{[0,1]^2} \mathcal{F}_\delta(u(t, x, \alpha), v(t, y, \beta)) \right. \\ \left. \cdot \frac{1}{2^d} \nabla\psi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) \xi_{\gamma, t_0}(t) d\alpha d\beta dy dx dt \right], \end{aligned}$$

along a subsequence. Sending $\gamma \downarrow 0$, applying Remark 5.2, yields (5.6).

Next we want to prove (5.7). To send $r \downarrow 0$ we apply Lemma 6.1. It is easily verified that condition (i) and (iii) are satisfied with $F_\delta = \mathcal{F}_\delta$. Consider condition (ii). Since $S'_\delta(-\sigma) = -S'_\delta(\sigma)$ for all $\sigma \in \mathbb{R}$, it follows that

$$\begin{aligned} \mathcal{F}_\delta(a, b) - \mathcal{F}_\delta(a, c) &= S'_\delta(b-a)(f(b) - f(a)) - S'_\delta(c-a)(f(c) - f(a)) \\ &= \int_b^c \partial_z(S'_\delta(z-a)(f(z) - f(a))) dz \\ &= \int_b^c S''_\delta(z-a)(f(z) - f(a)) dz + \int_b^c S'_\delta(z-a)f'(z) dz, \end{aligned}$$

for $a, b, c \in \mathbb{R}$. By (5.8),

$$|\mathcal{F}_\delta(a, b) - \mathcal{F}_\delta(a, c)| \leq \underbrace{\left| \int_b^c S''_\delta(z-a)(f(z) - f(a)) dz \right|}_{\leq \|f\|_{\text{Lip}} 2\delta} + \underbrace{\left| \int_b^c S'_\delta(z-a)f'(z) dz \right|}_{\leq \|f\|_{\text{Lip}} |b-c|}.$$

This and the symmetry of $\mathcal{F}_\delta$, i.e., $\mathcal{F}_\delta(a, b) = \mathcal{F}_\delta(b, a)$ for $a, b \in \mathbb{R}$, yields condition (ii). Hence, by Lemma 6.1,

$$\lim_{(\delta, r), \varepsilon, r_0 \downarrow 0} F_1 = E \left[\iint_{\Pi_T} \iint_{[0,1]^2} \text{sign}(u(t, x, \alpha) - v(t, x, \beta)) \right. \\ \left. \times (f(u(t, x, \alpha)) - f(v(t, x, \beta))) \cdot \nabla \psi(x) \xi_{\gamma, t_0}(t) d\beta d\alpha dx dt \right].$$

At long last, Limit (5.7) follows by sending $\gamma \downarrow 0$. $\square$

Limit 4. Suppose assumptions $(\mathcal{A}_{\sigma,1})$ and $(\mathcal{A}_\sigma)$ hold. With $\mathcal{T}_1$ defined in Lemma 5.4,

$$\mathcal{T}_1 = \mathcal{O} \left(\frac{r^{2\kappa+1}}{\delta} + \delta \right).$$

Proof. By assumption $(\mathcal{A}_{\sigma,1})$ and $(\mathcal{A}_\sigma)$,

$$|\sigma(y, v, z) - \sigma(x, u^\varepsilon, z)| \leq M(z) |y - x|^\kappa (1 + |u^\varepsilon|) + M(z) |v - u^\varepsilon|.$$

and thus

$$|\mathcal{T}_1| = \frac{1}{2} E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \int_Z S''_\delta(v - u^\varepsilon) (\sigma(y, v, z) - \sigma(x, u^\varepsilon, z))^2 \varphi d\mu(z) d\beta dX \right] \\ \leq \|M\|_{L^2(Z)}^2 E \left[\iiint_{\Pi_T^2} \int_{[0,1]} S''_\delta(v - u^\varepsilon) |x - y|^{2\kappa+1} (1 + |u^\varepsilon|)^2 \varphi d\beta dX \right] \\ + \|M\|_{L^2(Z)}^2 E \left[\iiint_{\Pi_T^2} \int_{[0,1]} S''_\delta(v - u^\varepsilon) |v - u^\varepsilon|^2 \varphi d\beta dX \right] \\ =: \mathcal{T}_1^1 + \mathcal{T}_1^2.$$

Since $J_r(\frac{x-y}{2}) = 0$ whenever $|x - y| \geq 2r$,

$$\mathcal{T}_1^1 \leq 4 \|M_K\|_{L^2(Z)}^2 \|J\|_\infty \frac{r^{2\kappa+1}}{\delta} E \left[\iiint_{\Pi_T^2} (1 + |u^\varepsilon|)^2 \varphi dX \right].$$

Moreover, as

$$E \left[\iiint_{\Pi_T^2} (1 + |u^\varepsilon|)^2 \varphi dX \right] \leq \int_0^T E \left[\|1 + u^\varepsilon(t)\|_{2, \psi \star J_r}^2 \right] dt,$$

there is a constant $C > 0$, independent of $r_0, \varepsilon, \delta, \gamma, r$, such that $\mathcal{T}_1^1 \leq Cr^{2\kappa+1}\delta^{-1}$. Regarding the second term $\mathcal{T}_1^2$, observe that

$$S''_\delta(v - u^\varepsilon) |v - u^\varepsilon|^2 = J_\delta(v - u^\varepsilon) |v - u^\varepsilon|^2 \leq 2 \|J\|_\infty \delta.$$

Hence, $\mathcal{T}_1^2 \leq \delta 2 \|J\|_\infty \|M\|_{L^2(Z)}^2 \|\varphi\|_{L^1(\Pi_T^2)}$. $\square$

Let us consider the term involving the Malliavin derivative.

Limit 5. With $\mathcal{T}_2$ defined in Lemma 5.4,

$$\lim_{r_0 \downarrow 0} \mathcal{T}_2 = 0.$$

Proof. Let us split $\mathcal{T}_2$ as follows:

$$\begin{aligned}
\mathcal{T}_2 = & E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \int_Z S''_\delta(v - u^\varepsilon(s, x)) \left(D_{s,z} u^\varepsilon(t, x) - \sigma(x, u^\varepsilon(s, x), z) \right) \right. \\
& \quad \left. \times \sigma(y, v, z) \varphi d\mu(z) d\beta dX \right] \\
& + E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \int_Z \left(S''_\delta(v - u^\varepsilon(t, x)) - S''_\delta(v - u^\varepsilon(s, x)) \right) \right. \\
& \quad \left. \times D_{s,z} u^\varepsilon(t, x) \sigma(y, v, z) \varphi d\mu(z) d\beta dX \right] \\
& + E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \int_Z S''_\delta(v - u^\varepsilon(s, x)) \left(\sigma(x, u^\varepsilon(s, x), z) - \sigma(x, u^\varepsilon(t, x), z) \right) \right. \\
& \quad \left. \times \sigma(y, v, z) \varphi d\mu(z) d\beta dX \right] \\
& + E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \int_Z \left(S''_\delta(v - u^\varepsilon(s, x)) - S''_\delta(v - u^\varepsilon(t, x)) \right) \right. \\
& \quad \left. \times \sigma(x, u^\varepsilon, z) \sigma(y, v, z) \varphi d\mu(z) d\beta dX \right] \\
= & \mathcal{T}_2^1 + \mathcal{T}_2^2 + \mathcal{T}_2^3 + \mathcal{T}_2^4.
\end{aligned}$$

Consider $\mathcal{T}_2^1$. We want to apply Lemma 3.8 for fixed (s, y, β) with

$$\begin{aligned}
\Psi_{s,y,\beta}(x, z) &= S''_\delta(v - u^\varepsilon(s, x)) \sigma(y, v, z), \\
\phi_y(x) &= \frac{1}{2^d} \psi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right).
\end{aligned}$$

Then

$$\mathcal{T}_2^1 = \iint_{\Pi_T} \int_0^1 \mathcal{T}_{r_0}(\Psi_{s,y,\beta}) d\beta ds dy.$$

By means of Lemma 3.8, $\lim_{r_0 \downarrow 0} \mathcal{T}_{r_0}(\Psi_{s,y,\beta}) = 0$ $ds dy d\beta$ -a.e., and so $\lim_{r_0 \downarrow 0} \mathcal{T}_2^1 = 0$ by the dominated convergence theorem. To this end, in view of (3.25), there exists a constant C , not depending on r_0 , such that

$$\begin{aligned}
|\mathcal{T}_{r_0}(\Psi_{s,y,\beta})|^2 &\leq C^2 E \left[\iint_{Z \times \mathbb{R}^d} |\Psi_{s,y,\beta}(x, z)|^2 \phi_y(x) dx d\mu(z) \right] \\
&\leq C^2 \|S''_\delta\|_\infty^2 E \left[\int_Z |\sigma(y, v, z)|^2 (\psi \star J_r)(y) d\mu(z) \right] \\
&\leq C^2 \|S''_\delta\|_\infty^2 \|M\|_{L^2(Z)}^2 E [(1 + |v|)^2 (\psi \star J_r)(y)].
\end{aligned}$$

Due to the compact support of $\psi \star J_r$, we see that $|\mathcal{T}_{r_0}(\Psi_{s,y,\beta})|$ is dominated by an integrable function.

Let us consider $\mathcal{T}_2^2$. Note that

$$\begin{aligned}
& |S''_\delta(v - u^\varepsilon(t, x)) - S''_\delta(v - u^\varepsilon(s, x))| \\
& \leq \underbrace{\max \left\{ 2 \|S''_\delta\|_\infty, \|S''_\delta\|_{\text{Lip}} |u^\varepsilon(t, x) - u^\varepsilon(s, x)| \right\}}_{\Psi}.
\end{aligned}$$

By Hölder's inequality,

$$\begin{aligned} \mathcal{T}_2^2 &\leq E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \int_Z \Psi^2(s, t, x) |\sigma(y, v, z)|^2 \varphi d\mu(z) d\beta dX \right]^{1/2} \\ &\quad \times E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \int_Z |D_{s,z} u^\varepsilon(t, x)|^2 \varphi d\mu(z) d\beta dX \right]^{1/2} \\ &=: F_1 \times F_2. \end{aligned}$$

By the uniform boundedness of Ψ we can apply the dominated convergence theorem and Lemma 6.2, to conclude that $\lim_{r_0 \downarrow 0} F_1 = 0$. It remains to show that $|F_2| \leq C$, with C independent of $r_0 > 0$. We deduce easily

$$\begin{aligned} F_2^2 &= \int_0^T \int_0^T E \left[\|D_s u^\varepsilon(t)\|_{L^2(Z; L^2(\mathbb{R}^d, \psi \star J_r))}^2 \right] J_{r_0}^+(t-s) \xi_{\gamma, t_0}(t) ds dt \\ &\leq \int_0^T \sup_{t \in [0, T]} \left\{ E \left[\|D_s u^\varepsilon(t)\|_{L^2(Z; L^2(\mathbb{R}^d, \psi \star J_r))}^2 \right] \right\} ds, \end{aligned}$$

and so $|F_2|$ is uniformly bounded by (3.16).

Consider $\mathcal{T}_2^3$. By Hölder's inequality and $(\mathcal{A}_\sigma)$,

$$\begin{aligned} |\mathcal{T}_2^3| &\leq \|S_\delta''\|_\infty \|M\|_{L^2(Z)} E \left[\iiint_{\Pi_T^2} \int_{[0,1]} \int_Z |\sigma(y, v, z)|^2 \varphi d\mu(z) d\beta dX \right]^{1/2} \\ &\quad \times E \left[\iiint_{\mathbb{R}^d \times [0, T]^2} |u^\varepsilon(s, x) - u^\varepsilon(t, x)|^2 (\psi \star J_r)(x) J_{r_0}^+(t-s) dx dt ds \right]^{1/2} \end{aligned}$$

By the dominated convergence theorem and Lemma 6.2, $\lim_{r_0 \downarrow 0} \mathcal{T}_2^3 = 0$.

The term $\mathcal{T}_2^4$ is treated in the same manner as $\mathcal{T}_2^2$, resulting in $\lim_{r_0 \downarrow 0} \mathcal{T}_2^4 = 0$. $\square$

Limit 6. With $\mathcal{T}_3$ defined in Lemma 5.4,

$$\mathcal{T}_3 = \mathcal{O}(\varepsilon).$$

Proof. Note that

$$|S_\delta(u^\varepsilon - v)\Delta_x \varphi| \leq (|u^\varepsilon| + |v|) |\Delta_x \varphi|.$$

Using this inequality, it follows from Lemma 4.2 that

$$E \left[\iiint_{\Pi_T^2} S_\delta(u^\varepsilon - v)\Delta_x \varphi d\beta dX \right] \leq C,$$

for some constant $C > 0$ independent of ε and r_0 . $\square$

Having established Proposition 5.5, the proof of Theorem 5.1 follows easily.

Proof of Theorem 5.1. In the setting of Proposition 5.5, suppose $u^0 = v^0$. Let $\{\phi_R\}_{R>1}$ be as in Lemma 6.6 and take $\psi = \phi_R$ in (5.4). Exploiting that ϕ belongs to $\mathfrak{N}$, sending $R \rightarrow \infty$ yields

$$\eta(t_0) \leq C_\phi \|f\|_{\text{Lip}} \int_0^{t_0} \eta(t) dt,$$

where

$$\eta(t) = E \left[\iiint_{\mathbb{R}^d \times [0, 1]^2} |u(t, x, \alpha) - v(t, x, \beta)| \phi(x) d\beta d\alpha dx \right].$$

An application of Grönwall's inequality gives $\eta(t_0) = 0$ for dt -a.a. $t_0 \in [0, T]$. Hence, any Young measure-valued entropy solution of (1.1) (in the sense of Definition 2.2)

with initial value u^0 , coincides with the one obtained in Theorem 4.1. Furthermore, $u(t, x, \alpha) = \hat{u}(t, x) := \int u(t, x, \alpha) d\alpha$ for $dt dx d\alpha dP$ -a.a. (t, x, α, ω) . It follows that $\hat{u}$ is an entropy solution (in the sense of definition 2.1). $\square$

Proof of Proposition 5.2. Let $\{\phi_R\}_{R>1}$ be as in Lemma 6.6, and start off from Lemma 5.4 with $\psi = \phi_R$ and $v^0 = u^0$. We then compute the limits $r_0 \downarrow 0$, $\varepsilon \downarrow 0$, and $\gamma \downarrow 0$ (in that order). Recall that by Theorem 5.1, $v = u$ with $u = \lim_{\varepsilon \downarrow 0} u^\varepsilon$. Furthermore, u is a solution according to Definition 2.1. Due to Limits 1–6 we arrive at the inequality

$$\begin{aligned}
 (5.9) \quad & E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(u^0(x-z) - u^0(x+z)) \phi_R(x) J_r(z) dx dz \right] \\
 & \geq E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(u(t_0, x-z) - u(t_0, x+z)) \phi_R(x) J_r(z) dx dz \right] \\
 & \quad + E \left[\int_0^{t_0} \iint_{\mathbb{R}^d \times \mathbb{R}^d} S'_\delta(u(t, x+z) - u(t, x-z)) \right. \\
 & \quad \quad \times (f(u(t, x+z)) - f(u(t, x-z))) \cdot \nabla \phi_R(x) J_r(z) dx dz dt \left. \right] \\
 & \quad + \mathcal{O} \left(\delta + \frac{\delta}{r} + \frac{r^{2\kappa+1}}{\delta} \right),
 \end{aligned}$$

where $\mathcal{O}(\cdot)$ is independent of R , cf. Limits 3 and 4 and Lemmas 6.5 and 6.6.

Note that

$$|S_\delta(\sigma) - |\sigma|| \leq \delta, \quad \forall \sigma \in \mathbb{R},$$

and $|\nabla \phi| \leq C_\phi \phi$. With the help of Lemma 6.6, we can now send $R \rightarrow \infty$ in (5.9), obtaining

$$\eta(t_0) \leq \eta(0) + C_\phi \|f\|_{\text{Lip}} \int_0^{t_0} \eta(t) dt + \mathcal{O} \left(\delta + \frac{\delta}{r} + \frac{r^{2\kappa+1}}{\delta} \right),$$

where

$$\eta(t) = E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u(t, x-z) - u(t, x+z)| \phi(x) J_r(z) dx dz \right].$$

By Grönwall's inequality,

$$\eta(t) \leq \left(1 + C_\phi \|f\|_{\text{Lip}} t e^{C_\phi \|f\|_{\text{Lip}} t} \right) \left(\eta(0) + \mathcal{O} \left(\delta + \frac{\delta}{r} + \frac{r^{2\kappa+1}}{\delta} \right) \right).$$

Prescribing $\delta = r^{\kappa+1}$ concludes the proof. $\square$

6. APPENDIX

6.1. Some “doubling of variables” tools.

Lemma 6.1. *Suppose $u, v \in L^1_{\text{loc}}(\mathbb{R}^d)$ and $\{F_\delta\}_{\delta>0}$ satisfy:*

- (i) *There is $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ such that $F_\delta \rightarrow F$ pointwise as $\delta \downarrow 0$.*
- (ii) *There exists a constant $C > 0$ such that*

$$|F_\delta(a, b) - F_\delta(c, d)| \leq C(|a - c| + |b - d| + \delta),$$

for all $a, b, c, d \in \mathbb{R}$ and all $\delta > 0$.

(iii) *There is a constant $C > 0$ such that*

$$|F_\delta(a, a)| \leq C(1 + |a|) \text{ for all } \delta > 0.$$

Fix $\psi \in C_c(\mathbb{R}^d)$. Suppose that $\delta : [0, \infty) \rightarrow [0, \infty)$ satisfies $\delta(r) \downarrow 0$ as $r \downarrow 0$. Set

$$\begin{aligned} \mathcal{T}_r := & \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} F_{\delta(r)}(u(x), v(y)) \frac{1}{2^d} \psi\left(\frac{x+y}{2}\right) J_r\left(\frac{x-y}{2}\right) dy dx \\ & - \int_{\mathbb{R}^d} F(u(x), v(x)) \psi(x) dx. \end{aligned}$$

Then $\mathcal{T}_r \rightarrow 0$ as $r \downarrow 0$.

Proof. Due to Remark 5.2,

$$\begin{aligned} \mathcal{T}_r = & \int_{\mathbb{R}^d} \underbrace{\int_{\mathbb{R}^d} F_{\delta(r)}(u(x+z), v(x-z)) \psi(x) dx}_{g_\delta(z)} J_r(z) dz \\ & - \underbrace{\int_{\mathbb{R}^d} F(u(x), v(x)) \psi(x) dx}_{g(0)}. \end{aligned}$$

Suppose for the moment that given a number $\varepsilon > 0$, there exists two numbers $\eta = \eta(\varepsilon) > 0$ and $\delta = \delta_0(\varepsilon) > 0$ such that

$$(6.1) \quad |g_\delta(z) - g(0)| \leq \varepsilon, \text{ whenever } |z| \leq \eta \text{ and } \delta < \delta_0.$$

The change of variables $z = r\zeta$ yields

$$|\mathcal{T}_r| \leq \int_{\mathbb{R}^d} |g_\delta(z) - g(0)| J_r(z) dz = \int_{\mathbb{R}^d} |g_\delta(r\zeta) - g(0)| J(\zeta) d\zeta.$$

Fix $\varepsilon > 0$, and pick η, δ_0 as dictated by (6.1). Let $r_0 > 0$ satisfy $r_0 \leq \eta$ and $\delta(r_0) \leq \delta_0$. It follows by (6.1) that $|\mathcal{T}_{r_0}| \leq \varepsilon$. Hence, $\mathcal{T}_{r_0} \downarrow 0$ as $r \downarrow 0$.

Let us now prove (6.1). By assumption (ii),

$$\begin{aligned} |g_\delta(z) - g(0)| & \leq C \int_{\mathbb{R}^d} |u(x+z) - u(x)| \psi(x) dx + C \int_{\mathbb{R}^d} |v(x-z) - v(x)| \psi(x) dx \\ & \quad + \int_{\mathbb{R}^d} |F_\delta(u(x), v(x)) - F(u(x), v(x))| \psi(x) dx + C\delta \|\psi\|_{L^1(\mathbb{R}^d)}. \end{aligned}$$

Because of assumptions (i) and (iii), we can apply the dominated convergence theorem to conclude that

$$\lim_{\delta \downarrow 0} \int_{\mathbb{R}^d} |F_\delta(u(x), v(x)) - F(u(x), v(x))| \psi(x) dx = 0.$$

It remains to show that

$$(6.2) \quad \lim_{z \rightarrow 0} \int_{\mathbb{R}^d} |u(x+z) - u(x)| \psi(x) dx = 0.$$

The term involving v follows by the same argument. Pick a compact $K \subset \mathbb{R}^d$ such that $\bigcup_{|z| \leq 1} \text{supp}(\psi(\cdot + z)) \subset K$. Fix $\varepsilon > 0$. By the density of continuous functions in $L^1(K)$, we can find $w \in C(K)$ such that $\|w - u\|_{L^1(K)} \leq \varepsilon$. Then

$$\int_{\mathbb{R}^d} |u(x+z) - u(x)| \psi(x) dx \leq 2 \|\psi\|_\infty \varepsilon + \int_{\mathbb{R}^d} |w(x+z) - w(x)| \psi(x) dx,$$

for any $|z| \leq 1$. Next we send $z \rightarrow 0$. The claim (6.2) follows by the dominated convergence theorem and the arbitrariness of $\varepsilon > 0$. $\square$

Lemma 6.2. *Let $v \in L^p([0, T])$, $1 \leq p < \infty$. Moreover, Let $F : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be measurable in the first variable and Lipschitz in the second variable,*

$$|F(s, a) - F(s, b)| \leq C |a - b|, \quad \forall a, b \in \mathbb{R}, \forall s \in [0, T],$$

for some constant $C > 0$. Set

$$\mathcal{T}_{r_0}(s) = \left(\int_0^T |F(s, v(t)) - F(s, v(s))|^p J_{r_0}^+(t-s) dt \right)^{1/p}.$$

Then $\mathcal{T}_{r_0}(s) \rightarrow 0$ ds -a.e. as $r_0 \downarrow 0$.

Proof. We can write $v = v_1^n + v_2^n$ with v_1^n continuous and $\|v_2^n\|_{L^p([0, T])} \leq 1/n$. This is possible since the continuous functions are dense in $L^p([0, T])$. Assuming $s \in [0, T - 2r_0]$, an application of the triangle inequality gives

$$\begin{aligned} |\mathcal{T}_{r_0}(s)| &\leq C \left(\int_0^T |v(t) - v(s)|^p J_{r_0}^+(t-s) dt \right)^{1/p} \\ &\leq C \left(\int_0^T |v_1^n(t) - v_1^n(s)|^p J_{r_0}^+(t-s) dt \right)^{1/p} \\ &\quad + (|v_2^n|^p \star J_{r_0}(s))^{1/p} + |v_2^n(s)|. \end{aligned}$$

Sending $r_0 \downarrow 0$, it follows that $\lim_{r_0 \downarrow 0} |\mathcal{T}_{r_0}(s)| \leq 2|v_2^n(s)|$ for ds -a.a. $s \in [0, T]$. Since $v_2^n \rightarrow 0$ in $L^p([0, T])$, it has a subsequence that converges ds -a.e., and this concludes the proof. $\square$

6.2. Weighted L^p spaces. First we make some elementary observations regarding functions in $\mathfrak{N}$ (see Section 2 for the definition of $\mathfrak{N}$).

Lemma 6.3. *Suppose $\phi \in \mathfrak{N}$ and $0 < p < \infty$. Then, for $x, z \in \mathbb{R}^d$,*

$$\left| \phi^{1/p}(x+z) - \phi^{1/p}(x) \right| \leq w_{p,\phi}(|z|) \phi^{1/p}(x),$$

where

$$w_{p,\phi}(r) = \frac{C_\phi}{p} r \left(1 + \frac{C_\phi}{p} r e^{C_\phi r/p} \right),$$

which is defined for all $r \geq 0$. As a consequence it follows that if $\phi(x_0) = 0$ for some $x_0 \in \mathbb{R}^d$, then $\phi \equiv 0$ (and by definition $\phi \notin \mathfrak{N}$).

Proof. Set $g(\lambda) = \phi^{1/p}(x + \lambda z)$. Then

$$g'(\lambda) = \frac{1}{p} \phi^{1/p-1}(x + \lambda z) (\nabla \phi(x + \lambda z) \cdot z).$$

Since $\phi \in \mathfrak{N}$, it follows that $|g'(\lambda)| \leq \frac{C_\phi}{p} g(\lambda) |z|$. Hence

$$g(\lambda) \leq g(0) + \frac{C_\phi}{p} |z| \int_0^\lambda g(\xi) d\xi.$$

By Grönwall's inequality,

$$g(\lambda) \leq g(0) \left(1 + \frac{C_\phi}{p} |z| \lambda e^{C_\phi |z| \lambda/p} \right).$$

Hence,

$$|g(1) - g(0)| \leq \frac{C_\phi}{p} |z| g(0) \left(1 + \frac{C_\phi}{p} |z| e^{C_\phi |z|/p} \right).$$

This concludes the proof. $\square$

Next, we consider an adaption of Young's inequality for convolutions.

Proposition 6.4. Fix $\phi \in \mathfrak{N}$. Suppose $f \in C_c(\mathbb{R}^d)$, and $g \in L^p(\mathbb{R}^d, \phi)$ for some finite $p \geq 1$. Then

$$\|f \star g\|_{L^p(\mathbb{R}^d, \phi)} \leq \left(\int_{\mathbb{R}^d} |f(x)| (1 + w_{p,\phi}(|x|)) dx \right) \|g\|_{L^p(\mathbb{R}^d, \phi)}.$$

where $w_{p,\phi}$ is defined in Lemma 6.3.

Proof. First observe that

$$\begin{aligned} \|f \star g\|_{L^p(\mathbb{R}^d, \phi)}^p &= \int_{\mathbb{R}^d} \left| \int_{\mathbb{R}^d} f(x-y)g(y) dy \right|^p \phi(x) dx \\ &\leq \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} |f(x-y)| \left(\frac{\phi(x)}{\phi(y)} \right)^{1/p} |g(y)| \phi^{1/p}(y) dy \right)^p dx. \end{aligned}$$

By Lemma 6.3(iii),

$$\left(\frac{\phi(x)}{\phi(y)} \right)^{1/p} \leq \frac{1}{\phi^{1/p}(y)} \left(\phi^{1/p}(y) + \left| \phi^{1/p}(x) - \phi^{1/p}(y) \right| \right) \leq (1 + w_{p,\phi}(|x-y|)).$$

Set

$$\zeta(x) := |f(x)| (1 + w_{p,\phi}(|x|)), \quad \xi(x) := |g(x)| \phi^{1/p}(x).$$

Then, by Young's inequality for convolutions,

$$\|f \star g\|_{L^p(\mathbb{R}^d, \phi)} \leq \|\zeta \star \xi\|_{L^p(\mathbb{R}^d)} \leq \|\zeta\|_{L^1(\mathbb{R}^d)} \|\xi\|_{L^p(\mathbb{R}^d)}.$$

□

Lemma 6.5. Fix $\phi \in \mathfrak{N}$, and let $w_{p,\phi}$ be defined in Lemma 6.3. Then

- (i) If $\varphi \in L^1(\mathbb{R}^d)$ is nonnegative, then $\phi \star \varphi \in \mathfrak{N}$ with $C_{\phi \star \varphi} = C_\phi$.
- (ii) Let $\varphi \in L^1(\mathbb{R}^d)$ have support in $B(0, 1)$ and $\|\varphi\|_{L^1(\mathbb{R}^d)} = 1$. Take $\varphi_\delta(z) = \frac{1}{\delta^d} \varphi(\frac{z}{\delta})$. Then, for any $u \in L^p(\mathbb{R}^d, \phi)$,

$$\left| \|u\|_{p,\phi}^p - \|u\|_{p,\phi \star \varphi_\delta}^p \right| \leq w_{1,\phi}(\delta) \|\varphi\|_{L^1(\mathbb{R}^d)} \|u\|_{p,\phi}^p.$$

- (iii) Let $\varphi \in C_c^2(\mathbb{R}^d)$. Then

$$|\Delta(\phi \star \varphi)(x)| \leq C_\phi \left(\int_{\mathbb{R}^d} |\nabla \varphi(z)| (1 + w_{1,\phi}(|z|)) dz \right) \phi(x).$$

If φ has support in $B(0, 1)$ this yields

$$|\Delta(\phi \star \varphi_\delta)(x)| \leq \frac{1}{\delta} (1 + w_{1,\phi}(|\delta|)) C_\phi \|\nabla \varphi\|_{L^1(\mathbb{R}^d)} \phi(x).$$

Proof. Consider (i). By Young's inequality for convolutions,

$$\|\phi \star \varphi\|_{L^1(\mathbb{R}^d)} \leq \|\phi\|_{L^1(\mathbb{R}^d)} \|\varphi\|_{L^1(\mathbb{R}^d)}.$$

Furthermore, $\phi \star \varphi \in C^1(\mathbb{R}^d)$ and

$$|\nabla(\phi \star \varphi)(x)| = \left| \int_{\mathbb{R}^d} \varphi(y) \nabla \phi(x-y) dy \right| \leq C_\phi (\phi \star \varphi)(x).$$

Consider (ii). By Lemma 6.3,

$$\begin{aligned} \left| \|u\|_{p,\phi}^p - \|u\|_{p,\phi \star \varphi_\delta}^p \right| &= \left| \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |u(x)|^p (\phi(x-z) - \phi(x)) \varphi_\delta(z) dz dx \right| \\ &\leq \|u\|_{p,\phi}^p \int_{\mathbb{R}^d} w_{1,\phi}(|z|) \varphi_\delta(z) dz. \end{aligned}$$

This proves (ii). Consider (iii). Integration by parts yields

$$\begin{aligned} |\Delta(\phi \star \varphi)(x)| &= \left| \int_{\mathbb{R}^d} \nabla \varphi(x-y) \cdot \nabla \phi(y) dy \right| \\ &\leq C_\phi \int_{\mathbb{R}^d} |\nabla \varphi(x-y)| \phi(y) dy. \end{aligned}$$

By Lemma 6.3,

$$\int_{\mathbb{R}^d} |\nabla \varphi(x-y)| \phi(y) dy \leq \left(\int_{\mathbb{R}^d} |\nabla \varphi(x-y)| (1 + w_{1,\phi}(|x-y|)) dy \right) \phi(x).$$

□

Lemma 6.6. *Let $\phi \in \mathfrak{N}$. Then there exists $\{\phi_R\}_{R>1} \subset C_c^\infty(\mathbb{R}^d)$ such that*

- (i) $\phi_R \rightarrow \phi$ and $\nabla \phi_R \rightarrow \nabla \phi$ pointwise in $\mathbb{R}^d$ as $R \rightarrow \infty$,
- (ii) $\exists$ a constant C independent of $R > 1$ such that

$$\max \left\{ \|\phi_R\|_{\infty, \phi^{-1}}, \|\nabla \phi_R\|_{\infty, \phi^{-1}} \right\} \leq C.$$

Proof. Modulo a mollification step, we may assume $\phi \in C^\infty$. Let $\zeta \in C_c^\infty(\mathbb{R}^d)$ satisfy $0 \leq \zeta \leq 1$, $\zeta(0) = 1$. Let $\phi_R(x) := \phi(x)\zeta(R^{-1}x)$. Then

$$\nabla \phi_R(x) = \nabla \phi(x)\zeta(R^{-1}x) + R^{-1}\phi(x)\nabla \zeta(R^{-1}x).$$

Hence (i) follows. Clearly, $\|\phi_R\|_{\infty, \phi^{-1}} = \sup_x \{|\phi_R(x)|\phi^{-1}(x)\} = \|\zeta\|_\infty$. Furthermore,

$$|\nabla \phi_R(x)| \leq (C_\phi \zeta(R^{-1}x) + R^{-1}|\nabla \zeta(R^{-1}x)|) \phi(x).$$

Hence, $\|\nabla \phi_R\|_{\infty, \phi^{-1}} \leq C_\phi + R^{-1}\|\nabla \zeta\|_\infty$. □

6.3. A version of Itô's formula. Here we establish the particular anticipating Itô formula applied in the proof of Theorem 4.1.

Theorem 6.7. *Let*

$$X(t) = X_0 + \int_0^t \int_Z u(s, z) W(dz, ds) + \int_0^t v(s) ds,$$

where $u : [0, T] \times Z \times \Omega \rightarrow \mathbb{R}$ and $v : [0, T] \times \Omega \rightarrow \mathbb{R}$ are jointly measurable and $\{\mathcal{F}_t\}$ -adapted processes, satisfying

$$(6.3) \quad E \left[\left(\int_0^T \int_Z u^2(s, z) d\mu(z) ds \right)^2 \right] < \infty, \quad E \left[\int_0^T v^2(s) ds \right] < \infty.$$

Let $F : \mathbb{R}^2 \times [0, T] \rightarrow \mathbb{R}$ be twice continuously differentiable. Suppose there exists a constant $C > 0$ such that for all $(\zeta, \lambda, t) \in \mathbb{R}^2 \times [0, T]$,

$$\begin{aligned} |F(\zeta, \lambda, t)|, |\partial_3 F(\zeta, \lambda, t)| &\leq C(1 + |\zeta| + |\lambda|), \\ |\partial_1 F(\zeta, \lambda, t)|, |\partial_{1,2}^2 F(\zeta, \lambda, t)|, |\partial_1^2 F(\zeta, \lambda, t)| &\leq C. \end{aligned}$$

Let $V \in \mathcal{S}$. Then $s \mapsto \partial_1 F(X(s), V, s)u(s)$ is Skorohod integrable, and

$$\begin{aligned}
F(X(t), V, t) &= F(X_0, V, 0) \\
&+ \int_0^t \partial_3 F(X(s), V, s) ds \\
&+ \int_0^t \int_Z \partial_1 F(X(s), V, s) u(s, z) W(dz, ds) \\
&+ \int_0^t \partial_1 F(X(s), V, s) v(s) ds \\
&+ \int_0^t \int_Z \partial_{1,2}^2 F(X(s), V, s) D_{s,z} V u(s, z) d\mu(z) ds \\
&+ \frac{1}{2} \int_0^t \int_Z \partial_1^2 F(X(s), V, s) u^2(s, z) d\mu(z) ds, \quad dP\text{-almost surely.}
\end{aligned}$$

Proof. The proof follows [27, Theorem 3.2.2 and Proposition 1.2.5]. We give an outline and some details where there are considerable differences. Furthermore, we leave out the t variable F as this is a standard modification.

Set $t_i^n = \frac{it}{2^n}$, $0 \leq i \leq 2^n$. By Taylor's formula,

$$\begin{aligned}
F(X(t), V) &= F(X_0, V) + \underbrace{\sum_{i=0}^{2^n-1} \partial_1 F(X(t_i^n), V) (X(t_{i+1}^n) - X(t_i^n))}_{\mathcal{T}_n^1} \\
&+ \underbrace{\frac{1}{2} \sum_{i=0}^{2^n-1} \partial_1^2 F(\bar{X}_i, V) (X(t_{i+1}^n) - X(t_i^n))^2}_{\mathcal{T}_n^2},
\end{aligned}$$

where $\bar{X}_i$ denotes a random intermediate point between $X(t_i^n)$ and $X(t_{i+1}^n)$. As in the proof of [27, Proposition 1.2.5],

$$\mathcal{T}_n^2 \rightarrow \frac{1}{2} \int_0^t \int_Z \partial_1^2 F(X(s), V) u^2(s, z) d\mu(z) ds, \quad \text{in } L^1(\Omega) \text{ as } n \rightarrow \infty.$$

Note that

$$\begin{aligned}
\mathcal{T}_n^1 &= \underbrace{\sum_{i=0}^{2^n-1} \partial_1 F(X(t_i^n), V) \int_{t_i^n}^{t_{i+1}^n} \int_Z u(s, z) W(dz, ds)}_{\mathcal{T}_n^{1,1}} \\
&+ \underbrace{\sum_{i=0}^{2^n-1} \partial_1 F(X(t_i^n), V) \int_{t_i^n}^{t_{i+1}^n} v(s) ds}_{\mathcal{T}_n^{1,2}}.
\end{aligned}$$

Clearly,

$$\mathcal{T}_n^{1,2} \rightarrow \int_0^t \partial_1 F(X(s), V) v(s) ds, \quad \text{in } L^1(\Omega) \text{ as } n \rightarrow \infty.$$

Consider $\mathcal{T}_n^{1,1}$. By [27, Proposition 1.3.5], $s \mapsto \partial_1 F(X(t_i^n), V)u(s)$ is Skorohod integrable on $[t_i^n, t_{i+1}^n]$ and

$$\begin{aligned} \mathcal{T}_n^{1,1} = & \underbrace{\sum_{i=0}^{2^n-1} \int_{t_i^n}^{t_{i+1}^n} \int_Z \partial_1 F(X(t_i^n), V)u(s, z) W(dz, ds)}_{\mathcal{T}_n^{1,1,1}} \\ & + \underbrace{\sum_{i=0}^{2^n-1} \int_{t_i^n}^{t_{i+1}^n} \int_Z \partial_{1,2}^2 F(X(t_i^n), V)D_{s,z}Vu(s, z) d\mu(z)ds}_{\mathcal{T}_n^{1,1,2}}. \end{aligned}$$

As before

$$\mathcal{T}_n^{1,1,2} \rightarrow \int_0^t \int_Z \partial_{1,2}^2 F(X(s), V)D_{s,z}Vu(s, z) d\mu(z)ds, \quad \text{in } L^1(\Omega) \text{ as } n \rightarrow \infty.$$

Consider $\mathcal{T}_n^{1,1,1}$. Let

$$\zeta_n(s, z) = \sum_{i=0}^{2^n-1} \partial_{1,2}^2 F(X(t_i^n), V) \mathbb{1}_{[t_i^n, t_{i+1}^n)}(s) D_{s,z}Vu(s, z),$$

and note that ζ_n is Skorohod integrable on $[0, t]$. We need to show the following:

- (i) There exists $\zeta \in L^2(\Omega; H)$ such that $\zeta_n \rightarrow \zeta$ in $L^2(\Omega; H)$.
- (ii) There exists a $G \in L^2(\Omega)$ such that for each $U \in \mathcal{S}$

$$E \left[\int_0^t \int_Z \zeta_n(s, z) W(dz, ds) U \right] \rightarrow E[G U].$$

Then we may conclude by [27, Proposition 1.3.6] that ζ is Skorohod integrable and $\int_0^t \zeta(s) dW(s) = G$. The result then follows. Consider (i). Let

$$\zeta(s, z) = \partial_{1,2}^2 F(X(s), V) D_{s,z}Vu(s, z).$$

Then

$$E \left[\int_0^t \int_Z |\zeta_n(s) - \zeta(s)|^2 d\mu(z)ds \right] \leq E \left[H_n \int_0^t \int_Z |D_{s,z}Vu(s, z)|^2 d\mu(z)ds \right],$$

where

$$H_n = \sup_{|t_i^n - s| \leq t2^{-n}} \left\{ \left| \partial_{1,2}^2 F(X(t_i^n), V) - \partial_{1,2}^2 F(X(s), V) \right|^2 \right\}.$$

Hence, (i) follows by the dominated convergence theorem. Consider (ii). The existence of a random variable G follows by the convergence of the other terms. This also yields the weak convergence. It remains to check that $G \in L^2(\Omega)$. This is a consequence of assumptions (6.3). $\square$

6.4. The Lebesgue-Bochner space. Let $(X, \mathcal{A}, \mu)$ be a σ -finite measure space and E a Banach space. In the previous sections $X = [0, T] \times \Omega$, $\mu = dt \otimes dP$, E is typically $L^p(\mathbb{R}^d, \phi)$ for some $1 \leq p < \infty$, and $\mathcal{A}$ is the predictable σ -algebra $\mathcal{P}$. A function $u : X \rightarrow E$ is *strongly μ -measurable* if there exists a sequence of μ -simple functions $\{u_n\}_{n \geq 1}$ such that $u_n \rightarrow u$ μ -almost everywhere. By a *μ -simple function* $s : X \rightarrow E$ we mean a function of the form

$$s(\zeta) = \sum_{k=1}^N \mathbb{1}_{A_k}(\zeta) x_k, \quad \zeta \in X,$$

where $x_k \in E$ and $A_k \in \mathcal{A}$ satisfy $\mu(A_k) < \infty$ for all $1 \leq k \leq N$. The Lebesgue-Bochner space $L^p(X, \mathcal{A}, \mu; E)$ is the linear space of μ -equivalence classes of strongly measurable functions $u : X \rightarrow E$ satisfying

$$\int_X \|u(\xi)\|_E^p d\mu(\xi) < \infty.$$

A map $u : X \rightarrow E$ is *weakly μ -measurable* if the map $\xi \mapsto \langle u(\xi), \varphi^* \rangle$ has a μ -version which is $\mathcal{A}$ -measurable for each φ^* in the dual space E^* . By the Pettis measurability theorem [32, Theorem 1.11], strong μ -measurability is equivalent to weak μ -measurability, whenever E is separable.

For $u \in L^1(X, \mathcal{A}, \mu; L^1(\mathbb{R}^d, \phi))$, it is convenient to know that $\zeta \mapsto u(\zeta)(x)$ has a μ -version which is $\mathcal{A}$ -measurable for almost all x . In fact this is crucial to the manipulations performed in the previous sections. The following result verifies that this is indeed the case.

Lemma 6.8. *Let $(X, \mathcal{A}, \mu)$ be a σ -finite measure space and $\phi \in \mathfrak{N}$. Let*

$$\Psi : L^1(X \times \mathbb{R}^d, \mathcal{A} \otimes \mathcal{B}(\mathbb{R}^d), d\mu \otimes d\phi) \rightarrow L^1(X, \mathcal{A}, \mu; L^1(\mathbb{R}^d, \phi))$$

be defined by $\Psi(u)(\xi) = u(\xi, \cdot)$. Then Ψ is an isometric isomorphism.

Remark 6.1. The measure space $(X \times \mathbb{R}^d, \mathcal{A} \otimes \mathcal{B}(\mathbb{R}^d), d\mu \otimes d\phi)$ is not necessarily complete. Strictly speaking we should rather consider its completion. What this ensures is that every representative is measurable with respect to the complete σ -algebra. A remedy is to define $L^1(X \times \mathbb{R}^d, \mathcal{A} \otimes \mathcal{B}(\mathbb{R}^d), d\mu \otimes d\phi)$ by asking that any element u has a $d\mu \otimes d\phi$ -version $\tilde{u}$ which is $\mathcal{A} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable. Now, $\tilde{u}(\cdot, x)$ is $\mathcal{A}$ -measurable, and so for $d\phi$ -almost all x , $u(\cdot, x)$ has a μ -version which is $\mathcal{A}$ -measurable.

Proof. Let us first check that $\Psi(u) \in L^1(X; L^1(\mathbb{R}^d, \phi))$. By the Pettis measurability theorem [32, Theorem 1.11], strong μ -measurability follows due to the separability of $L^1(\mathbb{R}^d, \phi)$ if $\Psi(u)$ is weakly μ -measurable. That is, for any $\varphi \in L^\infty(\mathbb{R}^d, \phi)$, the map

$$\xi \mapsto \int_{\mathbb{R}^d} \varphi(x) u(\xi, x) \phi(x) dx,$$

has a μ -version which is $\mathcal{A}$ -measurable. This is a consequence of Fubini's theorem [11, Proposition 5.2.2]. The fact that Ψ is an isometry is obvious. It remains to prove that Ψ is surjective. Let $v \in L^1(X; L^1(\mathbb{R}^d, \phi))$. By definition there exists a sequence $\{v_n\}_{n \geq 1}$ of simple functions such that $v_n \rightarrow v$ μ -almost everywhere. Set

$$v_n(\xi) = \sum_{k=1}^{N_n} \mathbf{1}_{A_{k,n}}(\xi) f_{k,n}, \quad u_n(\xi, x) = v_n(\xi)(x),$$

where $A_{k,n} \in \mathcal{A}$, $f_{k,n} \in L^1(\mathbb{R}^d, \phi)$. Note that u_n is $\mathcal{A} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable, and $\Psi(u_n) = v_n$. By the Lebesgue dominated convergence theorem, $v_n \rightarrow v$ in $L^1(X, L^1(\mathbb{R}^d, \phi))$ [32, Proposition 1.16]. By the isometry property, $\{u_n\}_{n \geq 1}$ is Cauchy, and so by completeness there exists u such that

$$u_n \rightarrow u \text{ in } L^1(X \times \mathbb{R}^d, \mathcal{A} \otimes \mathcal{B}(\mathbb{R}^d), d\mu \otimes d\phi).$$

Since

$$\begin{aligned} \int_X \|v - \Psi(u)\|_{1,\phi} d\mu &= \lim_{n \rightarrow \infty} \int_X \|v_n - \Psi(u)\|_{1,\phi} d\mu \\ &= \lim_{n \rightarrow \infty} \iint_{X \times \mathbb{R}^d} |u_n(\xi, x) - u(\xi, x)| d\mu \otimes d\phi(\xi, x) = 0, \end{aligned}$$

it follows that $\Psi(u) = v$. □

6.5. Young measures. The purpose of this subsection is to provide a reference for some results concerning Young measures and their application as generalized limits. Let $(X, \mathcal{A}, \mu)$ be a σ -finite measure space, and $\mathcal{P}(\mathbb{R})$ denote the set of probability measures on $\mathbb{R}$. In the previous sections X is typically $\Pi_T \times \Omega$. A *Young measure* from X into $\mathbb{R}$ is a function $\nu : X \rightarrow \mathcal{P}(\mathbb{R})$ such that $x \mapsto \nu_x(B)$ is $\mathcal{A}$ -measurable for every Borel measurable set $B \subset \mathbb{R}$. We denote by $\mathcal{Y}(X, \mathcal{A}, \mu; \mathbb{R})$, or simply $\mathcal{Y}(X; \mathbb{R})$ if the measure space is understood, the set of all Young measures from X into $\mathbb{R}$. The following theorem is proved in [29, Theorem 6.2] in the case that $X \subset \mathbb{R}^n$ and μ is the Lebesgue measure:

Theorem 6.9. *Let $(X, \mathcal{A}, \mu)$ be a σ -finite measure space. Let $\zeta : [0, \infty) \rightarrow [0, \infty]$ be a continuous, nondecreasing function satisfying $\lim_{\xi \rightarrow \infty} \zeta(\xi) = \infty$ and $\{u^n\}_{n \geq 1}$ a sequence of measurable functions such that*

$$\sup_n \int_X \zeta(|u^n|) d\mu(x) < \infty.$$

Then there exist a subsequence $\{u^{n_j}\}_{j \geq 1}$ and $\nu \in \mathcal{Y}(X, \mathcal{A}, \mu; \mathbb{R})$ such that for any Carathéodory function $\psi : \mathbb{R} \times X \rightarrow \mathbb{R}$ with $\psi(u^{n_j}(\cdot), \cdot) \rightharpoonup \bar{\psi}$ in $L^1(X)$, we have

$$\bar{\psi}(x) = \int_{\mathbb{R}} \psi(\xi, x) d\nu_x(\xi).$$

The proof is based on the embedding of $\mathcal{Y}(X; \mathbb{R})$ into $L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$. Here $\mathcal{M}(\mathbb{R})$ denotes the space of Radon measures on $\mathbb{R}$. The crucial observation is that $(L^1(X, C_0(\mathbb{R})))^*$ is isometrically isomorphic to $L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$ also in the case that $(X, \mathcal{A}, \mu)$ is an abstract σ -finite measure space. It is relatively straightforward to go through the proof and extend to this more general case [26, Theorem 2.11]. Note however that the use of weighted L^p spaces allows us to stick with the version for finite measure spaces.

6.6. Weak compactness in L^1 . To apply Theorem 6.9 one must first be able to extract from $\{\psi(\cdot, u^n(\cdot))\}_{n \geq 1}$ a weakly convergent subsequence in $L^1(X)$. The key result is the Dunford-Pettis Theorem.

Definition 6.1. Let $\mathcal{K} \subset L^1(X, \mathcal{A}, \mu)$.

(i) $\mathcal{K}$ is *uniformly integrable* if for any $\varepsilon > 0$ there exists $c_0(\varepsilon)$ such that

$$\sup_{f \in \mathcal{K}} \int_{|f| \geq c} |f| d\mu < \varepsilon, \quad \text{whenever } c > c_0(\varepsilon).$$

(ii) $\mathcal{K}$ has *uniform tail* if for any $\varepsilon > 0$ there exists $E \in \mathcal{A}$ with $\mu(E) < \infty$ such that

$$\sup_{f \in \mathcal{K}} \int_{X \setminus E} |f| d\mu < \varepsilon.$$

If $\mathcal{K}$ satisfies both (i) and (ii) it is said to be *equiintegrable*.

Remark 6.2. Note that (ii) is void when μ is finite. As a consequence uniform integrability and equiintegrability are equivalent for finite measure spaces.

Theorem 6.10 (Dunford-Pettis). *Let $(X, \mathcal{A}, \mu)$ be a σ -finite measure space. A subset $\mathcal{K}$ of $L^1(X)$ is relatively weakly sequentially compact if and only if it is equiintegrable.*

By the Eberlain-Šmulian theorem [40], in the weak topology of a Banach space, relative weak compactness is equivalent with relative sequentially weak compactness. There are a couple of well known reformulations of uniform integrability.

Lemma 6.11. *Suppose $\mathcal{K} \subset L^1(X)$ is bounded. Then $\mathcal{K}$ is uniformly integrable if and only if:*

(i) For any $\varepsilon > 0$ there exists $\delta(\varepsilon) > 0$ such that

$$\sup_{f \in \mathcal{K}} \int_E |f| d\mu < \varepsilon, \quad \text{whenever } \mu(E) < \delta(\varepsilon).$$

(ii) There is an increasing function $\Psi : [0, \infty) \rightarrow [0, \infty)$ such that $\Psi(\zeta)/\zeta \rightarrow \infty$ as $\zeta \rightarrow \infty$ and

$$\sup_{f \in \mathcal{K}} \int_X \Psi(|f(x)|) d\mu(x) < \infty.$$

Remark 6.3. Suppose there exists $g \in L^1(X)$ such that $|f| \leq g$ for all $f \in \mathcal{K}$. Then

$$\sup_{f \in \mathcal{K}} \int_E |f| d\mu \leq \int_E g d\mu, \quad \forall E \in \mathcal{A}.$$

Since $\{g\} \subset L^1(X)$ is uniformly integrable, it follows by Lemma 6.11(i) that $\mathcal{K}$ is uniformly integrable.

REFERENCES

- [1] L. Ambrosio and G. Crippa. Continuity equations and ODE flows with non-smooth velocity. *Proc. Roy. Soc. Edinburgh Sect. A*, 144(6):1191–1244, 2014.
- [2] E. J. Balder. Lectures on Young measure theory and its applications in economics. *Rend. Istit. Mat. Univ. Trieste*, 31(suppl. 1):1–69, 2000. Workshop on Measure Theory and Real Analysis (Italian) (Grado, 1997).
- [3] C. Bauzet, G. Vallet, and P. Wittbold. The Cauchy problem for conservation laws with a multiplicative stochastic perturbation. *J. Hyperbolic Differ. Equ.*, 9(4):661–709, 2012.
- [4] I. H. Biswas, K. H. Karlsen, and A. K. Majee. Conservation laws driven by Lévy white noise. *J. Hyperbolic Differ. Equ.*, to appear.
- [5] F. Bouchut and F. James. One-dimensional transport equations with discontinuous coefficients. *Nonlinear Anal.*, 32(7):891–933, 1998.
- [6] F. Bouchut, F. James, and S. Mancini. Uniqueness and weak stability for multi-dimensional transport equations with one-sided Lipschitz coefficient. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 4(1):1–25, 2005.
- [7] C. Cancès and T. Gallouët. On the time continuity of entropy solutions. *J. Evol. Equ.*, 11(1):43–55, 2011.
- [8] R. A. Carmona and M. R. Tehranchi. *Interest rate models: an infinite dimensional stochastic analysis perspective*. Springer Finance. Springer-Verlag, Berlin, 2006.
- [9] G.-Q. Chen, Q. Ding, and K. H. Karlsen. On nonlinear stochastic balance laws. *Arch. Ration. Mech. Anal.*, 204(3):707–743, 2012.
- [10] K. L. Chung and R. J. Williams. *Introduction to stochastic integration*. Modern Birkhäuser Classics. Birkhäuser/Springer, New York, second edition, 2014.
- [11] D. L. Cohn. *Measure theory*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser/Springer, New York, second edition, 2013.
- [12] S. Cox and M. Veraar. Vector-valued decoupling and the Burkholder-Davis-Gundy inequality. *Illinois J. Math.*, 55(1):343–375 (2012), 2011.
- [13] R. C. Dalang and L. Quer-Sardanyons. Stochastic integrals for spde’s: a comparison. *Expo. Math.*, 29(1):67–109, 2011.
- [14] A. Debussche and J. Vovelle. Scalar conservation laws with stochastic forcing. *J. Funct. Anal.*, 259(4):1014–1042, 2010.
- [15] R. J. DiPerna. Measure-valued solutions to conservation laws. *Arch. Rational Mech. Anal.*, 88(3):223–270, 1985.
- [16] R. Eymard, T. Gallouët, and R. Herbin. Existence and uniqueness of the entropy solution to a nonlinear hyperbolic equation. *Chinese Ann. Math. Ser. B*, 16(1):1–14, 1995.
- [17] J. Feng and D. Nualart. Stochastic scalar conservation laws. *J. Funct. Anal.*, 255(2):313–373, 2008.
- [18] F. Flandoli, M. Gubinelli, and E. Priola. Well-posedness of the transport equation by stochastic perturbation. *Invent. Math.*, 180(1):1–53, 2010.
- [19] M. Hardy. Combinatorics of partial derivatives. *Electron. J. Combin.*, 13(1):Research Paper 1, 13 pp. (electronic), 2006.
- [20] H. Holden and N. H. Risebro. Conservation laws with random source. *Appl. Math. Optim.*, 36(2) (1997) 229–241.

- [21] J. U. Kim. On a stochastic scalar conservation law. *Indiana Univ. Math. J.* **52** (1) (2003) 227–256.
- [22] D. Khoshnevisan. A primer on stochastic partial differential equations. In *A minicourse on stochastic partial differential equations*, volume 1962 of *Lecture Notes in Math.*, pages 1–38. Springer, Berlin, 2009.
- [23] S. N. Kružkov. First order quasilinear equations with several independent variables. *Mat. Sb. (N.S.)*, 81 (123):228–255, 1970.
- [24] P.-L. Lions, B. Perthame, and P. Souganidis. Scalar conservation laws with rough (stochastic) fluxes. *Stoch. Partial Differ. Equ. Anal. Comput.*, 1(4):664–686, 2013.
- [25] P.-L. Lions, B. Perthame, and P. E. Souganidis. Scalar conservation laws with rough (stochastic) fluxes: the spatially dependent case. *Stoch. Partial Differ. Equ. Anal. Comput.*, 2(4):517–538, 2014.
- [26] J. Málek, J. Nečas, M. Rokyta, and M. Růžička. Weak and measure-valued solutions to evolutionary PDEs, volume 13 of *Applied Mathematics and Mathematical Computation*. Chapman & Hall, London, 1996.
- [27] D. Nualart. The Malliavin calculus and related topics. *Probability and its Applications (New York)*. Springer-Verlag, Berlin, second edition, 2006.
- [28] E. Yu. Panov. On measure-valued solutions of the Cauchy problem for a first-order quasilinear equation. *Izv. Ross. Akad. Nauk Ser. Mat.*, 60(2):107–148, 1996.
- [29] P. Pedregal. Parametrized measures and variational principles. *Progress in Nonlinear Differential Equations and their Applications*, 30. Birkhäuser Verlag, Basel, 1997.
- [30] S. Peszat and J. Zabczyk. Stochastic partial differential equations with Lévy noise, volume 113 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2007.
- [31] E. M. Stein. Singular integrals and differentiability properties of functions. *Princeton Mathematical Series*, No. 30. Princeton University Press, Princeton, N.J., 1970.
- [32] J. M. A. M. van Neerven. Stochastic Evolution Equations. *ISEM Lecture Notes*, 2007/08.
- [33] J. M. A. M. van Neerven, M. C. Veraar, and L. Weis. Stochastic integration in UMD Banach spaces. *Ann. Probab.*, 35(4):1438–1478, 2007.
- [34] Giulia Di Nunno, Bernt Øksendal, and Frank Proske. *Malliavin calculus for Lévy processes with applications to finance*. Universitext. Springer-Verlag, Berlin, 2009.
- [35] G. Vallet. Dirichlet problem for a nonlinear conservation law. *Rev. Mat. Complut.*, 13(1):231–250, 2000.
- [36] G. Vallet and P. Wittbold. On a stochastic first-order hyperbolic equation in a bounded domain. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 12(4):613–651, 2009.
- [37] C. Villani. Topics in optimal transportation, volume 58 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2003.
- [38] A. I. Vol’pert and S. I. Hudjaev. The Cauchy problem for second order quasilinear degenerate parabolic equations. *Mat. Sb. (N.S.)*, 78 (120):374–396, 1969.
- [39] J. B. Walsh. An introduction to stochastic partial differential equations. In *École d’été de probabilités de Saint-Flour, XIV—1984*, volume 1180 of *Lecture Notes in Math.*, pages 265–439. Springer, Berlin, 1986.
- [40] R. Whitley. An elementary proof of the Eberlein-Šmulian theorem. *Math. Ann.*, 172:116–118, 1967.

(Kenneth Hvistendahl Karlsen)

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF OSLO

P.O. BOX 1053, BLINDERN

N-0316 OSLO, NORWAY

E-mail address: kennethk@math.uio.no

(Erlend Briseid Storrøsten)

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF OSLO

P.O. BOX 1053, BLINDERN

N-0316 OSLO, NORWAY

E-mail address: erlenbs@math.uio.no

ANALYSIS OF A SPLITTING METHOD FOR STOCHASTIC BALANCE LAWS

K. H. KARLSEN AND E. B. STORRØSTEN

ABSTRACT. We analyze a semi-discrete splitting method for conservation laws driven by a semilinear noise term. Making use of fractional BV type estimates, we show that the splitting method produces a compact sequence of approximate solutions converging to the exact solution, as the time step Δt tends to zero. Under the assumption of a homogenous noise function, and thus the availability of BV estimates, we provide an error estimate. Bringing into play a generalization of Kružkov's entropy condition, permitting the "Kružkov constants" to be Malliavin differentiable random variables, we establish an L^1 convergence rate of order $\frac{1}{3}$ in Δt .

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. Operator splitting	5
4. A priori estimates	6
5. Convergence	18
6. Error estimate	21
7. Appendix	31
References	37

1. INTRODUCTION

Recently there have been many works studying the effect of stochastic forcing on scalar conservation laws [3, 4, 5, 7, 12, 11, 8, 25, 24], with emphasis on existence, uniqueness, and stability questions. Deterministic conservation laws exhibit shocks (discontinuous solutions), and a weak formulation coupled with an appropriate entropy condition is required to ensure the well-posedness [15]. The question of uniqueness gets somewhat more difficult by adding a stochastic source term, due to the interaction between noise and nonlinearity. In a different direction, we have the pathwise theory in [17, 18] for conservation laws with (rough) stochastic fluxes.

In this paper we are interested in the convergence of approximate solutions to conservation laws driven by a multiplicative Wiener noise term, i.e., stochastic balance laws of the form

$$du + \operatorname{div} f(u) dt = \sigma(x, u) dB, \quad (t, x) \in \Pi_T, \quad (1.1)$$

Date: October 29, 2015.

2010 Mathematics Subject Classification. Primary: 60H15, 35L60; Secondary: 35L65, 60G15.

Key words and phrases. Stochastic conservation law, entropy condition, Malliavin calculus, numerical method, operator splitting, convergence, error estimate.

Acknowledgment: We are grateful to Nils Henrik Risebro for valuable comments and suggestions.

with initial data:

$$u(0, x) = u_0(x), \quad x \in \mathbb{R}^d. \quad (1.2)$$

We denote by ∇ and $\operatorname{div} = \nabla \cdot$ the spatial gradient and divergence, respectively. Moreover, $\Pi_T = \mathbb{R}^d \times (0, T)$ for some fixed final time $T > 0$, and $u(x, t)$ is the scalar unknown function that is sought. The random force in (1.1) is driven by a Wiener process $B = B(t) = B(t, \omega)$, $\omega \in \Omega$, over a stochastic basis $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$, where P is a probability measure, $\mathcal{F}$ is a σ -algebra, and $\{\mathcal{F}_t\}_{t \geq 0}$ is a right-continuous filtration on $(\Omega, \mathcal{F})$ such that $\mathcal{F}_0$ contains all the P -negligible subsets.

The convection flux $f : \mathbb{R} \rightarrow \mathbb{R}^d$ satisfies

$$f \text{ is Lipschitz continuous on } \mathbb{R}. \quad (\mathcal{A}_f)$$

Furthermore, we will sometimes make use of the assumption

$$f'' \text{ is uniformly bounded on } \mathbb{R}. \quad (\mathcal{A}_{f,1})$$

The noise coefficient $\sigma : \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}$ is assumed to satisfy

$$\|\sigma\|_{\text{Lip}} = \sup_{x \in \mathbb{R}^d} \sup_{u \neq v} \left\{ \frac{|\sigma(x, u) - \sigma(x, v)|}{|u - v|} \right\} < \infty, \quad |\sigma(\cdot, 0)| \in L^\infty(\mathbb{R}^d). \quad (\mathcal{A}_\sigma)$$

This yields the bounds

$$\begin{aligned} |\sigma(x, u) - \sigma(x, v)| &\leq \|\sigma\|_{\text{Lip}} |u - v|, \\ |\sigma(x, u)| &\leq \max \left\{ \|\sigma\|_{\text{Lip}}, \|\sigma(\cdot, 0)\|_{L^\infty(\mathbb{R}^d)} \right\} (1 + |u|). \end{aligned}$$

Furthermore, we will often assume that there is a constant M_σ such that

$$|\sigma(x, u) - \sigma(y, u)| \leq M_\sigma |x - y|^{\kappa_x + 1/2} (1 + |u|), \quad \kappa_x \in (0, 1/2]. \quad (\mathcal{A}_{\sigma,1})$$

A general difficulty affecting convergence/error analysis is related to the time discretization and the interplay between noise and nonlinearity. Up to now there are only a few studies investigating this problem. Holden and Risebro [9] study a one-dimensional equation with bounded initial data and a compactly supported, homogeneous noise function $\sigma = \sigma(u)$, ensuring L^∞ -bounds on the solution. An operator splitting method is used to construct approximate solutions, and it is shown that a subsequence of these approximations converges to a (possible non-unique) weak solution. Recently this work was generalized to stochastic entropy solutions and extended to the multi-dimensional case by Bauzet [1]. Kröker and Rohde [13] analyze semi-discrete (time continuous) finite volume methods. They use the L^p version of the compensated compactness method to prove convergence to a stochastic entropy solution for one-dimensional equations, with non-homogeneous noise function $\sigma = \sigma(x, u)$. Bauzet, Charrier, and Galloüet [2] analyze fully discrete finite volume methods for multi-dimensional equations, with homogeneous noise function $\sigma = \sigma(x, u)$. Their proof relies on weak BV (energy) estimates and a uniqueness result for measure-valued stochastic entropy solutions.

In this paper, as in [9, 1], we will investigate the semi-discrete splitting method for calculating approximations to stochastic entropy solutions of (1.1). Roughly speaking, this method is based on “splitting off” the effect of the stochastic source $\sigma(x, u) dB$. This Godunov-type operator splitting can be used to extend sophisticated numerical methods for deterministic conservation laws to stochastic balance laws. Generally speaking, the tag “operator splitting” refers to the well-known idea of constructing numerical methods for complicated partial differential equations by reducing them to a progression of simpler equations, each of which can be solved by some tailor-made numerical method. The operator splitting approach is described in a large number of articles and books. We do not survey the literature here, referring the reader instead to the bibliography in [10]. The main focus of the book

[10] is on convergence results, within classes of discontinuous functions, for general splitting algorithms for deterministic nonlinear partial differential equations.

Compared to the earlier results of Holden-Risebro and Bauzet [1, 9], the main contributions of the present paper are twofold. First, we establish convergence of the splitting approximations to a stochastic entropy solution in the case of non-homogenous noise functions $\sigma(x, u)$. Whenever σ has a dependency on the spatial position x , BV estimates are no longer available and the approach resorted to in [9, 1] does not apply. Following an idea laid out in [5], and independently in [7], we derive a fractional BV_x estimate, which, via an interpolation argument, is turned into a temporal equicontinuity estimate. These a priori estimates, along with Young measures and an earlier uniqueness result, are used to show that splitting approximations converge to a stochastic entropy solution. Second, for homogenous noise functions $\sigma = \sigma(u)$, we exhibit an L^1 error estimate of the form $O\left(\Delta t^{\frac{1}{3}}\right)$. Except for the expected convergence rate for the vanishing viscosity method [5], this appears to be the first error estimate derived for approximate solutions to stochastic conservation laws. The rate $\frac{1}{3}$ should be compared to the first order convergence rate available for conservation laws with deterministic source [16]. Our proof relies on BV estimates and a generalization of the Kruřkov entropy condition, allowing for the “Kruřkov constants” to be Malliavin differentiable random variables, which was put forward in the recent work [11].

The remaining part of this paper is organized as follows: Section 2 collects some preliminarily material along with the relevant notion of (stochastic entropy) solution. The operator splitting method is defined precisely in Section 3. A series of a priori estimates are derived in Section 4, which are subsequently used in Section 5 to prove convergence towards a stochastic entropy solution. Section 6 is devoted to the proof of the error estimate. Section 7 is an appendix collecting some definitions and useful results used elsewhere the paper.

2. PRELIMINARIES

Denote by $\mathfrak{N}$ all $\phi \in C^1(\mathbb{R}^d) \cap L^1(\mathbb{R}^d)$ for which there exists a constant C such that $|\nabla \phi| \leq C\phi$. Let

$$C_\phi = \inf \{C \mid |\nabla \phi| \leq C\phi\}.$$

For $\phi \in \mathfrak{N}$ we associate the weighted L^p -norm $\|\cdot\|_{p,\phi}$ defined by

$$\|u\|_{p,\phi} := \left(\int_{\mathbb{R}^d} |u(x)|^p \phi(x) dx \right)^{1/p}.$$

The weighted L^p -space associated with ϕ is denoted by $L^p(\mathbb{R}^d, \phi)$. The results we use for functions in $\mathfrak{N}$ are collected in Section 7.2.

Before we define the notion of entropy solution we need the notion of entropy pairs. Let us denote by $\mathcal{E}$ the set of non-negative convex functions in $C^2(\mathbb{R})$ such that S' is bounded, and S'' is compactly supported. A pair of functions (S, Q) is called an entropy pair if $S : \mathbb{R} \rightarrow \mathbb{R}$ is C^2 and $Q = (Q_1, \dots, Q_d) : \mathbb{R} \mapsto \mathbb{R}^d$ satisfies $Q' = S'f'$. An entropy pair (S, Q) is said to belong to $\mathcal{E}$ if S belongs to $\mathcal{E}$.

Let $\mathcal{P}$ denote the predictable σ -algebra on $[0, T] \times \Omega$ with respect to $\{\mathcal{F}_t\}$ [6, § 2.2]. In general we are working with equivalence classes of functions with respect to the measure $dt \otimes dP$. The equivalence class u is said to be *predictable* if it has a version $\tilde{u}$ that is $\mathcal{P}$ -measurable. Equivalently we could ask of any representative to be $\mathcal{P}^*$ measurable, where $\mathcal{P}^*$ is the completion of $\mathcal{P}$ with respect to $dt \otimes dP$. Note that any (jointly) measurable and adapted process is $\mathcal{P}^*$ -measurable [6, Theorem 3.7].

We use the notion of entropy solution defined in [11].

Definition 2.1. Let $\phi \in \mathfrak{N}$. An entropy solution u of (1.1)–(1.2) where $u^0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$ is a stochastic process

$$u = \{u(t) = u(t, x) = u(t, x; \omega)\}_{t \in [0, T]}$$

satisfying the following conditions:

- (i) u is a predictable process in $L^2([0, T] \times \Omega; L^2(\mathbb{R}^d, \phi))$.
- (ii) For any random variable $V \in \mathbb{D}^{1,2}$, any entropy, entropy-flux pair $(S, Q) \in \mathcal{E}$

$$\begin{aligned} E \left[\iint_{\Pi_T} S(u - V) \partial_t \varphi + Q(u, V) \cdot \nabla \varphi \, dx dt + \int_{\mathbb{R}^d} S(u_0(x) - V) \varphi(0, x) \, dx \right] \\ - E \left[\iint_{\Pi_T} S''(u - V) \sigma(x, u) D_t V \varphi \, dx dt \right] \\ + \frac{1}{2} E \left[\iint_{\Pi_T} S''(u - V) \sigma(x, u)^2 \varphi \, dx dt \right] \geq 0, \end{aligned}$$

for all non-negative $\varphi \in C_c^\infty([0, T] \times \mathbb{R}^d)$.

Here $L^2([0, T] \times \Omega; L^2(\mathbb{R}^d, \phi))$ denotes the Lebesgue-Bochner space and $D_t V$ denotes the Malliavin derivative of V evaluated at time t . The Malliavin calculus is developed with respect to the Gaussian process induced by the Wiener process $\{B(t)\}_{t \geq 0}$ [21]. $\mathbb{D}^{1,2}$ consists of all Malliavin differentiable $V \in L^2(\Omega, \mathcal{F}_T, P)$ such that $DV \in L^2(\Omega \times [0, T])$. The existence and uniqueness of such entropy solutions to (1.1)–(1.2) is established under assumption $(\mathcal{A}_f)$, $(\mathcal{A}_\sigma)$ and $(\mathcal{A}_{\sigma,1})$ in [11]. Let us also mention that whenever $u^0 \in L^p(\Omega; L^p(\mathbb{R}^d, \phi))$ for $2 \leq p < \infty$ it follows that

$$\text{ess sup}_{0 \leq t \leq T} \left\{ E \left[\|u(t)\|_{p, \phi}^p \right] \right\} < \infty.$$

Under the additional assumption $(\mathcal{A}_{\sigma,1})$, by [11, Proposition 5.2] the entropy solution u satisfies

$$\begin{aligned} E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u(t, x + z) - u(t, x - z)| J_r(z) \phi(x) \, dx \right] \\ \leq CE \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u_0(x + z) - u_0(x - z)| J_r(z) \phi(x) \, dx \right] + \mathcal{O}(r^{\kappa_x}) \end{aligned}$$

where κ_x is given in $(\mathcal{A}_{\sigma,1})$ and J_r is defined below (2.1). Whenever $\sigma(x, u) = \sigma(u)$ the last term on the right vanishes, i.e., $\mathcal{O} = 0$.

Let us fix some notation that will be applied throughout along with some basic observations. Let $\{J_\delta\}_{\delta > 0}$ be a sequence of symmetric mollifiers on $\mathbb{R}^d$, i.e.,

$$J_\delta(x) = \frac{1}{\delta^d} J^d\left(\frac{x}{\delta}\right), \quad (2.1)$$

where J^d is a nonnegative, smooth, symmetric function satisfying $\text{supp}(J^d) \subset B(0, 1)$ and $\int J = 1$. In general the dimension d is understood by the dimension of the argument and we write $J^d = J$. Whenever $d = 1$ we let $J^+(x) = J(x - 1)$ so that $\text{supp}(J^+) \subset (0, 2)$.

Next, we define a C^2 -approximation of the absolute value function. Let S_δ satisfy

$$S'_\delta(\sigma) = 2 \int_0^\sigma J_\delta(z) \, dz \text{ and } S_\delta(0) = 0. \quad (2.2)$$

Then

$$|r| - \delta \leq S_\delta(r) \leq |r|, \quad |S''_\delta(r)| \leq \frac{2}{\delta} \|J\|_\infty \mathbf{1}_{|r| < \delta}, \quad (2.3)$$

Given the entropy S_δ , we define the entropy flux $Q_\delta(u, v) : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ by

$$Q_\delta(u, v) = \int_v^u S'_\delta(\xi - v) f'(\xi) d\xi, \quad u, v \in \mathbb{R}. \quad (2.4)$$

It satisfies

$$|\partial_u(Q_\delta(u, v) - Q_\delta(v, u))| \leq \|f''\|_{L^\infty} \delta, \quad (2.5)$$

and

$$|Q_\delta(u, v)| \leq \|f\|_{\text{Lip}} S_\delta(u - v). \quad (2.6)$$

3. OPERATOR SPLITTING

We will now describe the basic operator splitting method for (1.1), that is, we will obtain an approximate solution through a composition of the solution operator to a deterministic conservation law and to a stochastic differential equation.

Let $\mathcal{S}_{\text{CL}}(t)$ be the solution operator that maps an initial function $v_0(x)$ to the unique entropy solution of the deterministic conservation law

$$\partial_t v + \text{div } f(v) = 0, \quad v(0, x) = v_0(x), \quad (3.1)$$

That is, if $v(t) := \mathcal{S}_{\text{CL}}(t)v_0$ then v is the unique entropy solution of (3.1). In fact, more precisely, for each $\tau \in [0, T]$

$$\begin{aligned} & \int_{\mathbb{R}^d} |v^0(x) - c| \varphi(0, x) dx - \int_{\mathbb{R}^d} |v(\tau) - c| \varphi(\tau, x) dx \\ & + \int_0^\tau \int_{\mathbb{R}^d} |v - c| \partial_t \varphi + \text{sign}(v - c) (f(v) - f(c)) \cdot \nabla \varphi dx dt \geq 0 \end{aligned}$$

for all $c \in \mathbb{R}$ and all non-negative $\varphi \in C_c^\infty([0, T] \times \mathbb{R})$. Let $v(t) := \mathcal{S}_{\text{CL}}(t)v_0$ and $u(t) := \mathcal{S}_{\text{CL}}(t)u_0$ where $u_0, v_0 \in L^1(\mathbb{R}^d, \phi)$ and $\phi \in \mathfrak{N}$. Then, for any $t \in [0, T]$

$$\|v(t) - u(t)\|_{1, \phi} \leq (1 + C_\phi \|f\|_{\text{Lip}} t e^{C_\phi \|f\|_{\text{Lip}} t}) \|u_0 - v_0\|_{1, \phi}.$$

Suppose $u^0 \in L^1(\Omega, \mathcal{F}_0, P; L^1(\mathbb{R}^d, \phi))$. By the decomposition of the map $\Omega \ni \omega \mapsto u_0(\omega) \mapsto \mathcal{S}_{\text{CL}}(t)u_0(\omega)$, it follows that $\mathcal{S}_{\text{CL}}(t)u_0$ is $\mathcal{F}_0$ measurable as a an element in $L^1(\mathbb{R}^d, \phi)$ [20, § 3.3].

Similarly, for $t > s$ we let $\mathcal{S}_{\text{SDE}}(t, s)$ denote the two-paramater semigroup defined by $\mathcal{S}_{\text{SDE}}(t, s)w^s = w(t)$ where w is the strong solution of

$$w(t, x) = w^s(x) + \int_s^t \sigma(x, w(r, x)) dB(r).$$

Suppose $w^s, v^s \in L^2(\Omega, \mathcal{F}_s, P; L^2(\mathbb{R}^d, \phi))$ and let $w(t) = \mathcal{S}_{\text{SDE}}(t, s)w^0$ and $v(t) = \mathcal{S}_{\text{SDE}}(t, s)v^s$. Then

$$E[\|w(t) - v(t)\|_{1, \phi}] = E[\|w^s - v^s\|_{1, \phi}]. \quad (3.2)$$

To this end, consider the quantity $S_\delta(w(t, x) - v(t, x))$. An application of Itô's formula yields

$$\begin{aligned} S_\delta(w(t, x) - v(t, x)) &= S_\delta(w^s(x) - v^s(x)) \\ &+ \int_s^t S'_\delta(w(r, x) - v(r, x))(\sigma(x, w(r, x)) - \sigma(x, v(r, x))) dB(r) \\ &+ \frac{1}{2} \int_s^t S''_\delta(w(r, x) - v(r, x))(\sigma(x, w(r, x)) - \sigma(x, v(r, x)))^2 dt. \end{aligned}$$

Multiply by $\phi(x)$, take the expectation, integrate in x and send $\delta \downarrow 0$. The result follows due to assumption $(\mathcal{A}_\sigma)$. Due to (3.2) it is seen that

$$\mathcal{S}_{\text{SDE}}(\cdot, s) : L^1(\Omega, \mathcal{F}_s, P; L^1(\mathbb{R}^d, \phi)) \rightarrow L^1([s, T] \times \Omega, \mathcal{P}_{[s, T]}, dt \otimes dP; L^1(\mathbb{R}^d, \phi)),$$

where $\mathcal{P}_{[s,T]}$ denotes the predictable σ -algebra relative to $\{\mathcal{F}_t\}_{s \leq t \leq T}$ on $[s, T] \times \Omega$.

Fix $N \in \mathbb{N}$ and take $\Delta t = T/N$. let $t_n = n\Delta t$. Set $u^0 := u_0$ and define

$$u^{n+1}(x; \omega) = [\mathcal{S}_{\text{SDE}}(t_{n+1}, t_n; \omega) \circ \mathcal{S}_{\text{CL}}(\Delta t)] u^n(x; \omega), \quad (3.3)$$

for $n = 0, 1, \dots, N-1$.

To investigate the convergence of the semi-discrete splitting algorithm (3.3), we need to work with functions that are not only defined for each $t_n = n\Delta t$, but in the entire interval $[0, T]$. To this end, we introduce two different “time-interpolants” $u_{\Delta t}(t) = u_{\Delta t}(t, x; \omega)$, $v_{\Delta t}(t) = v_{\Delta t}(t, x; \omega)$ defined by

$$u_{\Delta t}(t) = \mathcal{S}_{\text{SDE}}(t, t_n) \circ \mathcal{S}_{\text{CL}}(\Delta t) u^n, \quad t \in (t_n, t_{n+1}], \quad (3.4)$$

and

$$v_{\Delta t}(t) = \mathcal{S}_{\text{CL}}(t - t_n) u^n, \quad t \in [t_n, t_{n+1}) \quad (3.5)$$

for $n = 0, \dots, N-1$. Furthermore we let $u_{\Delta t}((t_n)+) = v_{\Delta t}((t_{n+1})-) = \mathcal{S}_{\text{CL}}(\Delta t) u^n$.

4. A PRIORI ESTIMATES

To establish the convergence of $\{u_{\Delta t}\}_{\Delta t > 0}$ to a stochastic entropy solution we will need a series of a priori estimates. These are also crucial when deriving the error estimate. The following result explains the introduction of the weight functions $\mathfrak{N}$.

Proposition 4.1 (Local L^p estimates). *Suppose $(\mathcal{A}_f)$ and $(\mathcal{A}_\sigma)$ are satisfied. Let $2 \leq p < \infty$ and $M \geq \|f\|_{\text{Lip}}$. For $t \in (0, T)$, $R > 0$ let $\Gamma(t) = \max\{0, R - Mt\}$. Suppose $\phi \in C^1(\mathbb{R})$ is nonnegative and satisfies $|\nabla \phi| \leq C_\phi \phi$. Then there exists constants C_1 and C_2 depending only on p, σ, f and C_ϕ such that*

$$\begin{aligned} E \left[\int_{B(0, \Gamma(t_n))} |u^n(x)|^p \phi(x) dx \right] &\leq e^{C_1 t_n} E \left[\int_{B(0, R)} |u_0(x)|^p \phi(x) dx \right] \\ &\quad + C_2 t_n e^{C_1 t_n} \int_{B(0, R)} \phi(x) dx. \end{aligned} \quad (4.1)$$

In the case that $\sigma(x, 0) = 0$ we have $C_2 = 0$.

Proof of Proposition 4.1. Deterministic step. We want to prove the following: Let $1 \leq p < \infty$, $v^0 \in L^p_{\text{loc}}(\mathbb{R}^d)$ and $v(t) = \mathcal{S}_{\text{CL}}(t) v^0$. Then

$$\int_{B(0, \Gamma(t))} |v(t, x)|^p \phi(x) dx \leq e^{\|f\|_{\text{Lip}} C_\phi t} \int_{B(0, R)} |v^0(x)|^p \phi(x) dx. \quad (4.2)$$

We might as well assume $\Gamma(t) > 0$. As v is an entropy solution of (3.1) it follows that

$$\iint_{\Pi_T} \int_{\mathbb{R}^d} S(v(t, x)) \partial_t \varphi + Q(v(t, x)) \cdot \nabla_x \varphi dx dt + \int_{\mathbb{R}} S(v^0(x)) \varphi(0, x) dx \geq 0$$

for all convex entropy, entropy-flux pairs (S, Q) and all non-negative $\varphi \in C_c^\infty([0, T] \times \mathbb{R}^d)$. Let $\varphi(t, x) = \psi_\delta(t) H_\delta(\Gamma(t), |x|) \phi(x)$ where

$$\psi_\delta(t) = 1 - \int_0^t J_\delta(\zeta - \tau) d\zeta \quad \text{and} \quad H_\delta(L, r) = \int_{-\delta}^L J_\delta(\zeta - r) d\zeta.$$

Then

$$\begin{aligned} \partial_t \varphi(t, x) &= -J_\delta(t - \tau) H_\delta(\Gamma(t), |x|) \phi(x) - M \psi_\delta(t) J_\delta(\Gamma(t) - |x|) \phi(x), \\ \nabla \varphi(t, x) &= -\psi_\delta(t) J_\delta(\Gamma(t) - |x|) \frac{x}{|x|} \phi(x) + \psi_\delta(t) H_\delta(\Gamma(t), |x|) \nabla \phi(x). \end{aligned}$$

Hence

$$\begin{aligned}
 \int_{\mathbb{R}} S(v^0(x)) H_{\delta}(R, |x|) \phi(x) dx &\geq \iint_{\Pi_T} S(v(t, x)) J_{\delta}(t - \tau) H_{\delta}(\Gamma(t), |x|) \phi(x) dx dt \\
 &\quad + \underbrace{\iint_{\Pi_T} \left(Q(v(t, x)) \cdot \frac{x}{|x|} + MS(v(t, x)) \right) \psi_{\delta}(t) J_{\delta}(\Gamma(t) - |x|) \phi(x) dx dt}_{\mathcal{T}^1} \\
 &\quad - \underbrace{\iint_{\Pi_T} Q(v(t, x)) \psi_{\delta}(t) H_{\delta}(\Gamma(t), |x|) \cdot \nabla \phi(x) dx dt}_{\mathcal{T}^2}.
 \end{aligned} \tag{4.3}$$

Suppose $S'(0) = S(0) = 0$. Then

$$|Q(v)| = \left| \int_0^v S'(z) f'(z) dz \right| \leq \|f\|_{\text{Lip}} S(v).$$

It follows as $M \geq \|f\|_{\text{Lip}}$ that $\mathcal{T}^1 \geq 0$. Due to the assumption on ϕ

$$|\mathcal{T}_2| \leq \|f\|_{\text{Lip}} C_{\phi} \iint_{\Pi_T} S(v) \psi_{\delta}(t) H_{\delta}(R, |x|) \phi(x) dx dt.$$

Let $\delta, \delta \downarrow 0$. Inequality (4.3) takes the following form

$$X(\tau) \leq X(0) + \|f\|_{\text{Lip}} C_{\phi} \int_0^{\tau} X(r) dr,$$

where

$$X(t) = \int_{B(0, \Gamma(t))} S(v(t, x)) \phi(x) dx.$$

Equation (4.2) follows by taking $S(v) = |v|^p$ and applying Grönwall's inequality.

Stochastic step. We want to prove the following: Let $2 \leq p < \infty$. Suppose $w(s) \in L^p(\Omega; L^p_{\text{loc}}(\mathbb{R}))$ and take $w(t) = \mathcal{S}_{\text{SDE}}(t, s)w(s)$. For any $R > 0$ there exists constants C_3 and C_2 depending only on p and σ such that

$$\begin{aligned}
 E \left[\int_B |w(t, x)|^p \phi(x) dx \right] &\leq e^{C_3(t-s)} \left(E \left[\int_B |w(s, x)|^p \phi(x) dx \right] \right. \\
 &\quad \left. + C_2(t-s) \int_B \phi(x) dx \right). \tag{4.4}
 \end{aligned}$$

If $\sigma(x, 0) = 0$ then we may pick $C_2 = 0$.

By Ito's lemma,

$$d\eta(w) = \frac{1}{2} \eta''(w) \sigma(x, w)^2 dt + \eta'(w) \sigma(x, w) dB,$$

for any $\eta \in C^2$. Without loss of generality, we can assume $p = 2, 4, 6, \dots$. Taking $\eta(u) = |u|^p$ multiplying by ϕ and integrating over $B = B(0, R)$ yields

$$\begin{aligned}
 E \left[\int_B |w(t, x)|^p \phi(x) dx \right] &- E \left[\int_B |w(s, x)|^p \phi(x) dx \right] \\
 &\leq \frac{p(p-1)}{2} \int_s^t E \left[\int_B w(r, x)^{p-2} \sigma(x, w(r, x))^2 \phi(x) dx \right] dr.
 \end{aligned}$$

Recall that $\sigma(x, w) \leq |\sigma(x, 0)| + \|\sigma\|_{\text{Lip}} |w|$. Hence, according to assumption $(\mathcal{A}_{\sigma})$

$$\begin{aligned}
 T_1 &:= \frac{p(p-1)}{2} E \left[\int_B w(r, x)^{p-2} \sigma(x, w(r, x))^2 \phi(x) dx \right] \\
 &\leq p(p-1) \left(\|\sigma(\cdot, 0)\|_{\infty}^2 E \left[\int_B |w(r, x)|^{p-2} \phi(x) dx \right] \right.
 \end{aligned}$$

$$+ \|\sigma\|_{\text{Lip}}^2 E \left[\int_B |w(r, x)|^p \phi(x) dx \right].$$

Applying Hölder's inequality and then Young's inequality yields

$$\int_B |w(r, x)|^{p-2} \phi(x) dx \leq \frac{p-2}{p} \int_B |w(r, x)|^p \phi(x) dx + \frac{2}{p} \int_B \phi(x) dx.$$

Hence

$$\begin{aligned} T_1 \leq & \underbrace{(p-1) \left((p-2) \|\sigma(\cdot, 0)\|_\infty + p \|\sigma\|_{\text{Lip}}^2 \right)}_{C_3} E \left[\int_B |w(r, x)|^p \phi(x) dx \right] \\ & + \underbrace{2(p-1) \|\sigma(\cdot, 0)\|_\infty^2}_{C_4} \int_B \phi(x) dx. \end{aligned}$$

It follows that

$$\begin{aligned} E \left[\int_B |w(t, x)|^p \phi(x) dx \right] & \leq E \left[\int_B |w(s, x)|^p \phi(x) dx \right] \\ & + C_3 \int_s^t E \left[\int_B |w(r, x)|^p \phi(x) dx \right] dr + C_4 \left(\int_B \phi(x) dx \right) (t-s). \end{aligned}$$

This inequality is of the following general form:

$$X(t) \leq X(s) + \int_s^t K(r) X(r) dr + \int_s^t H(r) dr. \quad (4.5)$$

Appealing to Gronwall's inequality yields

$$X(t) \leq \exp \left[\int_s^t K(r) dr \right] X(s) + \int_s^t \exp \left[\int_r^t K(u) du \right] H(r) dr. \quad (4.6)$$

Identifying $K = C_3$ and $H = C_4 \|\phi\|_{L^1(B)}$, it follows that

$$\begin{aligned} E \left[\int_B |w(t, x)|^p \phi(x) dx \right] & \leq e^{C_3(t-s)} E \left[\int_B |w(s, x)|^p \phi(x) dx \right] \\ & + C_4 \|\phi\|_{L^1(B)} \int_s^t e^{C_3(t-r)} dr. \end{aligned}$$

Integration by parts yields

$$\begin{aligned} \int_s^t e^{C_3(t-r)} dr & = C_3 \int_s^t e^{C_3(t-r)} (t-r) dr + e^{C_3(t-s)} (t-s) \\ & \leq \left(\frac{C_3}{2} (t-s) + 1 \right) e^{C_3(t-s)} (t-s). \end{aligned}$$

Equation (4.4) follows. In the case that $\sigma(x, 0) = 0$ we observe that $C_4 = 0$ and hence $C_2 = 0$.

Inductive step. Let P_n be the statement that inequality (4.1) is true. Note that P_0 is trivially true. We want to show that P_n implies P_{n+1} . By (3.3) $u^{n+1} = \mathcal{S}_{\text{SDE}}(t_{n+1}, t_n) \mathcal{S}_{\text{CL}}(\Delta t) u^n$. Recall that $v_{\Delta t}((t_{n+1})-) = \mathcal{S}_{\text{CL}}(\Delta t) u^n$. By (4.2)

$$E \left[\int_{B(0, \Gamma(t_{n+1}))} |v_{\Delta t}((t_{n+1})-, x)|^p \phi(x) dx \right] \leq e^{\|f\|_{\text{Lip}} C_\phi \Delta t} E \left[\int_{B(0, \Gamma(t_n))} |u^n(x)|^p \phi(x) dx \right].$$

Since $u^{n+1} = \mathcal{S}_{\text{SDE}}(t_{n+1}, t_n) v_{\Delta t}((t_{n+1})-)$ it follows by (4.4)

$$E \left[\int_{B(0, \Gamma(t_{n+1}))} |u^{n+1}(x)|^p \phi(x) dx \right] \leq$$

$$e^{C_3 \Delta t} \left(E \left[\int_{B(0, \Gamma(t_{n+1}))} |v_{\Delta t}((t_{n+1})-, x)|^p \phi(x) dx \right] + C_2 \int_{B(0, \Gamma(t_{n+1}))} \phi(x) dx \Delta t \right).$$

Combining the two yields

$$\begin{aligned} E \left[\int_{B(0, \Gamma(t_{n+1}))} |u^{n+1}(x)|^p \phi(x) dx \right] \\ \leq e^{C_3 \Delta t} \left(e^{\|f\|_{\text{Lip}} C_\phi \Delta t} E \left[\int_{B(0, \Gamma(t_n))} |u^n(x)|^p \phi(x) dx \right] \right. \\ \left. + C_2 \Delta t \int_{B(0, \Gamma(t_{n+1}))} \phi(x) dx \right) \\ \leq e^{C_1 \Delta t} \left(E \left[\int_{B(0, \Gamma(t_n))} |u^n(x)|^p \phi(x) dx \right] \right. \\ \left. + C_2 \Delta t \int_{B(0, R)} \phi(x) dx \right) \end{aligned}$$

where $C_1 = \|f\|_{\text{Lip}} C_\phi + C_3$. Inserting the induction hypothesis finishes the proof of (4.1). $\square$

Corollary 4.2. *Let $u_{\Delta t}$ and $v_{\Delta t}$ be defined by (3.4) and (3.5) respectively. Suppose u_0 belongs to $L^q(\Omega; L^q(\mathbb{R}^d, \phi))$ for some $2 \leq q < \infty$ and some $\phi \in \mathfrak{N}$. Then, for each $1 \leq p \leq q$ there exists a finite constant C independent of Δt (but dependent on T, p, ϕ, f, σ and u_0) such that*

$$\max \left\{ E \left[\|u_{\Delta t}(t)\|_{p, \phi}^p \right], E \left[\|v_{\Delta t}(t)\|_{p, \phi}^p \right] \right\} \leq C, \quad t \in [0, T].$$

Proof. It suffices to prove the result for $p = q$. To this end suppose $1 \leq p < q$ and $w \in L^q(\mathbb{R}^d, \phi)$. Let $r = q/p$ and $s = q/(q-p)$ and observe that $r^{-1} + s^{-1} = 1$. Take $f = |u|^p \phi^{1/r}$ and $g = \phi^{1/s}$ and apply Hölder's inequality. This yields

$$\int_{\mathbb{R}^d} |w(x)|^p \phi(x) dx \leq \left(\int_{\mathbb{R}^d} |w(x)|^q \phi(x) dx \right)^{p/q} \left(\int_{\mathbb{R}^d} \phi(x) dx \right)^{1-p/q}. \quad (4.7)$$

Consider the case $p = q$. By Proposition 4.1 it follows that there exists a constant $C > 0$ depending only on q, f, σ, u_0, T and ϕ such that

$$E \left[\|u^n\|_{q, \phi}^q \right] \leq C$$

for all $0 \leq n \leq N$. By (4.2)

$$E \left[\|\mathcal{S}_{\text{CL}}(t - t_n) u^n\|_{q, \phi}^q \right] \leq e^{\|f\|_{\text{Lip}} C_\phi (t - t_n)} E \left[\|u^n\|_{q, \phi}^q \right].$$

This finishes the proof for $v_{\Delta t}$. For $u_{\Delta t}$ the result follows by (4.4). $\square$

The following result should be compared with [11, Proposition 5.2]. See also [5, § 6].

Proposition 4.3 (fractional BV estimates). *Suppose $(\mathcal{A}_f)$, $(\mathcal{A}_{f,1})$, $(\mathcal{A}_\sigma)$ and $(\mathcal{A}_{\sigma,1})$ are satisfied. Let $\phi \in \mathfrak{N}$. Suppose $u^0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$. Let $u_{\Delta t}$ and $v_{\Delta t}$ be defined by (3.4) and (3.5) respectively. Then there exists a constant $C_{T, \phi}$, independent of Δt , such that*

$$E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u_{\Delta t}(t, x+z) - u_{\Delta t}(t, x-z)| J_r(z) \phi(x) dx dz \right]$$

$$\leq C_{T,\phi} E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u_0(x+z) - u_0(x-z)| J_r(z) \phi(x) dx dz \right] + C_{T,\phi} r^{\kappa_x},$$

for any $t \in (0, T)$. Here $\kappa_x \in (0, 1/2]$ defined in $(\mathcal{A}_{\sigma,1})$. If $\sigma(x, u) = \sigma(u)$ the last term on the right vanishes. The same holds with $u_{\Delta t}$ replaced by $v_{\Delta t}$.

Before we continue, let me state some convenient change-of-variables identities. For $h = h(\cdot, \cdot)$

$$\begin{aligned} \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} h(x, y) \phi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) dx dy \\ = \iint_{\mathbb{R}^d \times \mathbb{R}^d} h(\tilde{x} + z, \tilde{x} - z) \phi(\tilde{x}) J_r(z) d\tilde{x} dz. \end{aligned} \quad (4.8)$$

This follows by the change of variables $(\tilde{x}, z) = (\frac{x+y}{2}, \frac{x-y}{2})$. Next,

$$\frac{1}{2^d} \int_{\mathbb{R}^d} \phi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) dy = \phi \star J_r(x). \quad (4.9)$$

This follows by taking $z = (x+y)/2$ and observing that $dy = 2^d dz$.

Proof of Proposition 4.3. Given a stochastic process $u = u(t) = u(t, x; \omega)$, we introduce the quantity

$$\mathcal{D}_r^u(t) := E \left[\frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} |u(t, x) - u(t, y)| J_r \left(\frac{x-y}{2} \right) \phi \left(\frac{x+y}{2} \right) dx dy \right].$$

Actually, at first we are not going to work with this quantity but rather

$$\mathcal{D}_{r,\delta}^u(t) := E \left[\frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(u(t, x) - u(t, y)) J_r \left(\frac{x-y}{2} \right) \phi \left(\frac{x+y}{2} \right) dx dy \right],$$

where the regularized entropy S_δ is defined in (2.2). In view of (2.3), and (4.9)

$$|\mathcal{D}_r^u(t) - \mathcal{D}_{r,\delta}^u(t)| \leq \|\phi\|_{L^1(\mathbb{R}^d)} \delta, \quad t > 0. \quad (4.10)$$

1. Deterministic step. Let $v(t, x)$ be the unique entropy solution of the conservation law (3.1). We want to prove the following claim: There exist a constant C_1 depending only on J and C_ϕ such that for all $0 < r \leq 1$

$$\mathcal{D}_{r,\delta}^v(t) \leq e^{C_\phi \|f\|_{\text{Lip}} t} \left(\mathcal{D}_{r,\delta}^v(0) + C_1 \|f''\|_\infty E \left[\|v_0\|_{1,\phi} \right] t \left(\frac{\delta}{r} + \delta \right) \right). \quad (4.11)$$

Let S_δ and Q_δ be the functions defined in (2.2) and (2.4), respectively. Using the entropy inequalities and Kružkov's method of doubling the variables, it follows

in a standard way that for $t > 0$

$$\begin{aligned}
 & \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(v(t, x) - v(t, y)) J_r\left(\frac{x-y}{2}\right) \phi\left(\frac{x+y}{2}\right) dx dy \\
 & \quad - \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(v_0(x) - v_0(y)) J_r\left(\frac{x-y}{2}\right) \phi\left(\frac{x+y}{2}\right) dx dy \\
 & \leq \frac{1}{2^d} \int_0^t \iint_{\mathbb{R}^d \times \mathbb{R}^d} Q_\delta(v(s, x), v(s, y)) \cdot \nabla \phi\left(\frac{x+y}{2}\right) J_r\left(\frac{x-y}{2}\right) dx dy ds \\
 & \quad + \frac{1}{2^d} \int_0^t \iint_{\mathbb{R}^d \times \mathbb{R}^d} (Q_\delta(v(s, y), v(s, x)) - Q_\delta(v(s, x), v(s, y))) \\
 & \quad \cdot \nabla_y \left(\phi\left(\frac{x+y}{2}\right) J_r\left(\frac{x-y}{2}\right) \right) dx dy ds \\
 & =: \mathcal{T}_{\text{CL}}^1 + \mathcal{T}_{\text{CL}}^2.
 \end{aligned}$$

By (2.6),

$$|\mathcal{T}_{\text{CL}}^1| \leq C_\phi \|f\|_{\text{Lip}} \frac{1}{2^d} \int_0^t \iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(v(s, x) - v(s, y)) J_r\left(\frac{x-y}{2}\right) \phi\left(\frac{x+y}{2}\right) dx dy ds.$$

Consider $\mathcal{T}_{\text{CL}}^2$. Thanks to (2.5),

$$|Q_\delta(v, u) - Q_\delta(u, v)| = \left| \int_v^u \partial_\xi (Q_\delta(\xi, v) - Q_\delta(v, \xi)) d\xi \right| \leq \|f''\|_\infty |u - v| \delta,$$

so that

$$\begin{aligned}
 |\mathcal{T}_{\text{CL}}^2| & \leq \frac{\|f''\|_\infty}{2} \delta \frac{1}{2^d} \int_0^t \iint_{\mathbb{R}^d \times \mathbb{R}^d} |v(s, x) - v(s, y)| \left| \nabla J_r\left(\frac{x-y}{2}\right) \right| \phi\left(\frac{x+y}{2}\right) dx dy ds \\
 & \quad + \frac{\|f''\|_\infty}{2} \delta \frac{1}{2^d} \int_0^t \iint_{\mathbb{R}^d \times \mathbb{R}^d} |v(s, x) - v(s, y)| J_r\left(\frac{x-y}{2}\right) \left| \nabla \phi\left(\frac{x+y}{2}\right) \right| dx dy ds \\
 & = \mathcal{T}_{\text{CL}}^{2,1} + \mathcal{T}_{\text{CL}}^{2,2}.
 \end{aligned}$$

Consider $\mathcal{T}_{\text{CL}}^{2,1}$. Let $\varphi_r(z) = \|\nabla J\|_1^{-1} \frac{1}{r^d} |\nabla J(\frac{z}{r})|$ so that

$$\left| \nabla J_r\left(\frac{x-y}{2}\right) \right| = \|\nabla J\|_1 \frac{1}{r} \varphi_r\left(\frac{x-y}{2}\right).$$

By the triangle inequality and (4.9)

$$\begin{aligned}
 & \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} |v(s, x) - v(s, y)| \left| \nabla J_r\left(\frac{x-y}{2}\right) \right| \phi\left(\frac{x+y}{2}\right) dx dy \\
 & \leq \|\nabla J\|_1 \frac{2}{r} \int_{\mathbb{R}^d} |v(s, x)| (\phi \star \varphi_r)(x) dx = \|\nabla J\|_1 \frac{2}{r} \|v(s)\|_{1, \phi \star \varphi_r}.
 \end{aligned}$$

Consider $\mathcal{T}_{\text{CL}}^{2,2}$. Similarly, as $\phi \in \mathfrak{N}$

$$\begin{aligned}
 & \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} |v(s, x) - v(s, y)| J_r\left(\frac{x-y}{2}\right) \left| \nabla \phi\left(\frac{x+y}{2}\right) \right| dx dy \\
 & \leq 2C_\phi \int_{\mathbb{R}^d} |v(s, x)| (\phi \star J_r)(x) dx = 2C_\phi \|v(s)\|_{1, \phi \star J_r}.
 \end{aligned}$$

By Lemma 7.3 there is a constant C_2 depending only on J and C_ϕ such that for all $0 \leq r \leq 1$

$$\max \left\{ \|v(s)\|_{1, \phi \star \varphi_r}, \|v(s)\|_{1, \phi \star J_r} \right\} \leq \|v(s)\|_{1, \phi} (1 + rC_2).$$

Hence

$$|\mathcal{T}_{\text{CL}}^2| \leq \|f''\|_\infty (1 + rC_2) \left(\int_0^t \|v(s)\|_{1, \phi} ds \right) \left(\|\nabla J\|_1 \frac{1}{r} + C_\phi \right) \delta$$

Due to (4.2) it follows that $\|v(s)\|_{1, \phi} \leq e^{\|f\|_{\text{Lip}} C_\phi s} \|v_0\|_{1, \phi}$. Summarizing, we have established the inequality

$$\begin{aligned} & \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(v(t, x) - v(t, y)) J_r\left(\frac{x-y}{2}\right) \phi\left(\frac{x+y}{2}\right) dx dy \\ & \quad - \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(v_0(x) - v_0(y)) J_r\left(\frac{x-y}{2}\right) \phi\left(\frac{x+y}{2}\right) dx dy \\ & \leq \underbrace{C_\phi \|f\|_{\text{Lip}}}_K \int_0^t \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(v(s, x) - v(s, y)) J_r\left(\frac{x-y}{2}\right) \phi\left(\frac{x+y}{2}\right) dx dy ds \\ & \quad + \underbrace{\int_0^t C_1 \|f''\|_\infty \|v_0\|_{1, \phi} e^{\|f\|_{\text{Lip}} C_\phi s} \left(\frac{\delta}{r} + \delta \right) ds}_{H(s)}. \end{aligned} \quad (4.12)$$

This inequality is of the form (4.5). By Gronwall's inequality (4.6),

$$\begin{aligned} & \iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(v(t, x) - v(t, y)) J_r\left(\frac{x-y}{2}\right) \phi\left(\frac{x+y}{2}\right) dx dy \\ & \leq e^{C_\phi \|f\|_{\text{Lip}} t} \left(\iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(v_0(x) - v_0(y)) J_r\left(\frac{x-y}{2}\right) \phi\left(\frac{x+y}{2}\right) dx dy \right. \\ & \quad \left. + C_1 \|f''\|_\infty \|v_0\|_{1, \phi} t \left(\frac{\delta}{r} + \delta \right) \right). \end{aligned}$$

This proves the claim (4.11)

2. *Stochastic step.* Let $w(t) = \mathcal{S}_{\text{SDE}}(t, s)w(s)$. We will now derive an estimate for w similar to (4.12). In fact, we want to prove the following claim: There exists constants C_1 and C_2 depending only on J, σ and ϕ such that

$$\mathcal{D}_{r, \delta}^w(t) \leq \mathcal{D}_{r, \delta}^w(s) + C_1 \frac{r^{2\kappa+1}}{\delta} \int_s^t E \left[\|1 + |w(\zeta)|\|_{2, \phi}^2 \right] d\zeta + C_2(t-s)\delta, \quad (4.13)$$

for all $0 \leq r \leq 1$. If $M_\sigma = 0$ then $C_1 = 0$.

To this end observe that $w(t, x) - w(t, y)$ solves the stochastic differential equation

$$d(w(t, x) - w(t, y)) = (\sigma(x, w(t, x)) - \sigma(y, w(t, y))) dB(t).$$

Applying Ito's formula to $S_\delta(w(t, x) - w(t, y))$ gives

$$\begin{aligned} & dS_\delta(w(t, x) - w(t, y)) \\ & = \frac{1}{2} S_\delta''(w(t, x) - w(t, y)) (\sigma(x, w(t, x)) - \sigma(y, w(t, y)))^2 dt \\ & \quad + S_\delta'(w(t, x) - w(t, y)) (\sigma(x, w(t, x)) - \sigma(y, w(t, y))) dB(t), \end{aligned}$$

and by integrating against the test function $J_r(\frac{x-y}{2})\phi(\frac{x+y}{2})$, we arrive at

$$\begin{aligned}
 & \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(w(t, x) - w(t, y)) J_r(\frac{x-y}{2}) \phi(\frac{x+y}{2}) dx dy \\
 & \quad - \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta(w(s, x) - w(s, y)) J_r(\frac{x-y}{2}) \phi(\frac{x+y}{2}) dx dy \\
 & = \int_s^t \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \frac{1}{2} S_\delta''(w(\zeta, x) - w(\zeta, y)) \\
 & \quad \times (\sigma(x, w(\zeta, x)) - \sigma(y, w(\zeta, y)))^2 J_r(\frac{x-y}{2}) \phi(\frac{x+y}{2}) dx dy d\zeta \\
 & \quad + \int_s^t \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} S_\delta'(w(\zeta, x) - w(\zeta, y)) (\sigma(x, w(\zeta, x)) - \sigma(y, w(\zeta, y))) dx dy dB(\zeta) \\
 & = \mathcal{T}_{\text{SDE}}^1 + \mathcal{T}_{\text{SDE}}^2,
 \end{aligned}$$

where the $\mathcal{T}_{\text{SDE}}^2$ -term has zero expectation. Note that

$$(\sigma(x, u) - \sigma(y, v))^2 \leq 2(\sigma(x, u) - \sigma(x, v))^2 + 2(\sigma(x, v) - \sigma(y, v))^2$$

for any $u, v \in \mathbb{R}$. We estimate the $\mathcal{T}_{\text{SDE}}^1$ -term as follows:

$$\begin{aligned}
 E[|\mathcal{T}_{\text{SDE}}^1|] & \leq 2E \left[\int_s^t \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} J_\delta(w(\zeta, x) - w(\zeta, y)) (\sigma(x, w(\zeta, x)) - \sigma(y, w(\zeta, x)))^2 \right. \\
 & \quad \times J_r(\frac{x-y}{2}) \phi(\frac{x+y}{2}) dx dy d\zeta \Big] \\
 & \quad + 2E \left[\int_s^t \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} J_\delta(w(\zeta, x) - w(\zeta, y)) (\sigma(y, w(\zeta, x)) - \sigma(y, w(\zeta, y)))^2 \right. \\
 & \quad \times J_r(\frac{x-y}{2}) \phi(\frac{x+y}{2}) dx dy d\zeta \Big] =: S_1 + S_2.
 \end{aligned}$$

Regarding S_1 , recall that $|J_\delta| \leq \|J\|_\infty / \delta$. By $(\mathcal{A}_{\sigma,1})$

$$\begin{aligned}
 |S_1| & \leq \|J\|_\infty \frac{2}{\delta} E \left[\int_s^t \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} |\sigma(x, w(\zeta, x)) - \sigma(y, w(\zeta, x))|^2 J_r(\frac{x-y}{2}) \phi(\frac{x+y}{2}) dx dy ds \right] \\
 & \leq \|J\|_\infty M_\sigma^2 \frac{2}{\delta} E \left[\int_s^t \frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} |x - y|^{2\kappa_x + 1} (1 + |w(\zeta, x)|)^2 J_r(\frac{x-y}{2}) \phi(\frac{x+y}{2}) dx dy ds \right] \\
 & \leq 2\|J\|_\infty M_\sigma^2 \frac{r^{2\kappa_x + 1}}{\delta} \int_s^t E \left[\|1 + |w(\zeta)|\|_{2, \phi \star J_r}^2 \right] d\zeta.
 \end{aligned}$$

By Lemma 7.3 there exist a constant C depending on C_ϕ only such that

$$\|1 + |w(\zeta)|\|_{2, \phi \star J_r}^2 \leq \|1 + |w(\zeta)|\|_{2, \phi}^2 (1 + rC_1)$$

for all $0 \leq r \leq 1$. It follows that

$$|S_1| \leq 2\|J\|_\infty M_\sigma^2 (1 + rC_1) \int_s^t E \left[\|1 + |w(\zeta)|\|_{2, \phi}^2 \right] d\zeta \frac{r^{2\kappa_x + 1}}{\delta}$$

Consider S_2 . Due to assumption $(\mathcal{A}_\sigma)$

$$J_\delta(w(\zeta, x) - w(\zeta, y)) (\sigma(y, w(\zeta, x)) - \sigma(y, w(\zeta, y)))^2 \leq \|\sigma\|_{\text{Lip}}^2 \|J\|_\infty \delta.$$

Hence

$$|S_2| \leq 2 \|\sigma\|_{\text{Lip}}^2 \|J\|_\infty \|\phi\|_{L^1(\mathbb{R}^d)} (t - s) \delta$$

This proves (4.13)

3. Inductive step. For u^n be defined by (3.3) and $u_{\Delta t}$ by (3.4), let P_n be the following claim: There exists constants C_1, C_2 and C_3 depending only on J, ϕ, σ such that for all $0 < r \leq 1$

$$\begin{aligned} \mathcal{D}_{r,\delta}^{u^n} &\leq e^{C_\phi \|f\|_{\text{Lip}} t_n} \left(\mathcal{D}_{r,\delta}^{u_0} + C_3 \|f''\|_\infty \left(\Delta t \sum_{k=0}^{n-1} E \left[\|u^k\|_{1,\phi} \right] \right) \left(\frac{\delta}{r} + \delta \right) \right. \\ &\quad \left. + C_1 \frac{r^{2\kappa_x+1}}{\delta} \int_0^{t_n} E \left[\|1 + |u_{\Delta t}(t)|\|_{2,\phi}^2 \right] dt + C_2 t_n \delta \right). \end{aligned} \quad (4.14)$$

If $M_\sigma = 0$ then $C_1 = 0$. Note that P_0 is trivially true. Assume that P_n is true, we want to show that P_{n+1} is holds. Recall that $u^{n+1} = \mathcal{S}_{\text{SDE}}(t_{n+1}, t_n) \mathcal{S}_{\text{CL}}(\Delta t) u^n$. Let $w^n = \mathcal{S}_{\text{CL}}(\Delta t) u^n$ and note that $\mathcal{S}_{\text{SDE}}(t, t_n) w^n = u_{\Delta t}(t)$ for $t_n \leq t < t_{n+1}$. As $1 \leq e^{C_\phi \|f\|_{\text{Lip}} \Delta t}$, (4.13) yields

$$\mathcal{D}_{r,\delta}^{u^{n+1}} \leq \mathcal{D}_{r,\delta}^{w^n} + e^{C_\phi \|f\|_{\text{Lip}} \Delta t} \left(C_1 \frac{r^{2\kappa_x+1}}{\delta} \int_{t_n}^{t_{n+1}} E \left[\|1 + |u_{\Delta t}(t)|\|_{2,\phi}^2 \right] dt + C_2 \Delta t \delta \right).$$

By (4.11)

$$\mathcal{D}_{r,\delta}^{w^n} \leq e^{C_\phi \|f\|_{\text{Lip}} \Delta t} \left(\mathcal{D}_{r,\delta}^{u^n} + C_3 \|f''\|_\infty E \left[\|u^n\|_{1,\phi} \right] \Delta t \left(\frac{\delta}{r} + \delta \right) \right).$$

Hence

$$\begin{aligned} \mathcal{D}_{r,\delta}^{u^{n+1}} &\leq e^{C_\phi \|f\|_{\text{Lip}} \Delta t} \left(\mathcal{D}_{r,\delta}^{u^n} + C_3 \|f''\|_\infty E \left[\|u^n\|_{1,\phi} \right] \Delta t \left(\frac{\delta}{r} + \delta \right) \right. \\ &\quad \left. + C_1 \frac{r^{2\kappa_x+1}}{\delta} \int_{t_n}^{t_{n+1}} E \left[\|1 + |u_{\Delta t}(t)|\|_{2,\phi}^2 \right] dt + C_2 \Delta t \delta \right). \end{aligned}$$

Inserting the hypothesis P_n yields P_{n+1} .

4. Concluding the proof. Consider (4.14). By Corollary 4.2

$$\mathcal{D}_{r,\delta}^{u^n} \leq e^{C t_n} \mathcal{D}_{r,\delta}^{u_0} + C t_n e^{C t_n} \left(\frac{\delta}{r} + \delta + \frac{r^{2\kappa_x+1}}{\delta} \right),$$

Thanks to (4.10), this translates into

$$\mathcal{D}_r^{u^n} \leq C_T \mathcal{D}_r^{u_0} + C_T \left(\frac{\delta}{r} + \delta + \frac{r^{2\kappa_x+1}}{\delta} + \|\phi\|_{L^1(\mathbb{R}^d)} \delta \right), \quad 0 \leq n \leq N.$$

We may reason via (4.13) to obtain

$$\mathcal{D}_r^{u_{\Delta t}}(t) \leq C_T \mathcal{D}_r^{u_0} + C_T \left(\frac{\delta}{r} + \delta + \frac{r^{2\kappa_x+1}}{\delta} + \|\phi\|_{L^1(\mathbb{R}^d)} \delta \right), \quad t \in [0, T].$$

Note that the same holds true if we replace $u_{\Delta t}$ by $v_{\Delta t}$ due to (4.11). Viewing $r > 0$ as fixed, we can choose $\delta = \mathcal{O}(r^{\kappa+1})$ to arrive at the bound

$$\mathcal{D}_r^{u_{\Delta t}}(t) \leq C_T \mathcal{D}_r^{u_0} + C_T r^{\kappa_x}.$$

The result follows by (4.8). In the case that $M_\sigma = 0$ we have

$$\mathcal{D}_r^{u_{\Delta t}}(t) \leq C_T \mathcal{D}_r^{u_0} + C_T \left(\frac{\delta}{r} + \delta + \|\phi\|_{L^1(\mathbb{R}^d)} \delta \right), \quad t \in (0, T).$$

and we may let $\delta \downarrow 0$ independent of r . $\square$

Next we consider time continuity. Recall that the interpolants $u_{\Delta t}, v_{\Delta t}$ are discontinuous at $t_n = n\Delta t, n = 1, \dots, N-1$. Hence, the result must somehow quantify the size of the jumps as $\Delta t \downarrow 0$.

Proposition 4.4 (L^1 time continuity). *Assume that $(\mathcal{A}_f), (\mathcal{A}_{f,1}), (\mathcal{A}_\sigma)$ and $(\mathcal{A}_{\sigma,1})$ are satisfied. Let $\phi \in \mathfrak{N}$ and $u^0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$ satisfy*

$$E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u_0(x+z) - u_0(x-z)| |J_r(z)\phi(x)| dx dz \right] = \mathcal{O}(r^{\kappa_0}), \quad (4.15)$$

for some $0 < \kappa_0 \leq 1$. Let

$$\kappa := \begin{cases} \min \{\kappa_0, \kappa_x\} & \text{if } \sigma = \sigma(x, u), \\ \kappa_0 & \text{if } \sigma = \sigma(u). \end{cases}$$

Let $u_{\Delta t}$ and $v_{\Delta t}$ be defined by (3.4) and (3.5) respectively. Then:

- (i) Suppose $0 < \tau_1 < \tau_2 \leq T$ satisfies $\tau_1 \in (t_k, t_{k+1}]$ and $\tau_2 \in (t_l, t_{l+1}]$. Then there exists a finite constant $C_{T,\phi}$, independent of Δt , such that

$$E \left[\int_{\mathbb{R}^d} |u_{\Delta t}(\tau_2, x) - u_{\Delta t}(\tau_1, x)| \phi(x) dx \right] \leq C_{T,\phi} \left(|(l-k)\Delta t|^{\frac{\kappa}{\kappa+1}} + \sqrt{\tau_2 - \tau_1} \right).$$

- (ii) Suppose $0 \leq \tau_1 \leq \tau_2 < T$ satisfies $\tau_1 \in [t_k, t_{k+1})$ and $\tau_2 \in [t_l, t_{l+1})$. Then there exists a finite constant $C_{T,\phi}$, independent of Δt , such that

$$E \left[\int_{\mathbb{R}^d} |v_{\Delta t}(\tau_2, x) - v_{\Delta t}(\tau_1, x)| \phi(x) dx \right] \leq C_{T,\phi} \left(\sqrt{(l-k)\Delta t} + |\tau_2 - \tau_1|^{\frac{\kappa}{\kappa+1}} \right)$$

Proof. We shall first quantify weak continuity in the mean of $t \mapsto u_{\Delta t}(t), t \mapsto v_{\Delta t}(t)$, and then—via Proposition 4.3 along the lines of [14, 15]—turn this into fractional L^1 time continuity in the mean. The reason for first exhibiting a weak estimate is that the deterministic step does not produce functions that are Lipschitz continuous in time, thereby preventing a direct “inductive argument”, see [14].

1. *Weak estimate.* Let $t_n = n\Delta t$. Suppose $0 < \tau_1 \leq \tau_2 \leq T$ satisfies $\tau_1 \in (t_k, t_{k+1}]$ and $\tau_2 \in (t_l, t_{l+1}]$. Fix any $\varphi \in L^\infty(\Omega; C^1(\mathbb{R}^d))$ and $\phi \in \mathfrak{N}$. We claim that there exists a constant $C > 0$ independent of Δt and φ such that

$$\begin{aligned} E \left[\int_{\mathbb{R}^d} \varphi(x) \left(u_{\Delta t}(\tau_2, x) - u_{\Delta t}(\tau_1, x) \right) \phi(x) dx \right] \\ \leq C \left((\|\nabla \varphi\|_{L^\infty} + \|\varphi\|_{L^\infty}) (l-k)\Delta t + \|\varphi\|_{L^\infty} \sqrt{\tau_2 - \tau_1} \right). \end{aligned} \quad (4.16)$$

Let $\psi = \varphi\phi$. To this end we continue as follows:

$$\begin{aligned} \mathcal{T} &= E \left[\int_{\mathbb{R}^d} \psi(x) \left(u_{\Delta t}(\tau_2, x) - u_{\Delta t}(\tau_1, x) \right) dx \right] \\ &= E \left[\int_{\mathbb{R}^d} \psi(x) \left(u_{\Delta t}(\tau_2, x) - u_{\Delta t}((t_l)+, x) \right) dx \right] \\ &\quad + E \left[\int_{\mathbb{R}^d} \psi(x) \left(u_{\Delta t}(t_{k+1}, x) - u_{\Delta t}(\tau_1, x) \right) dx \right] \\ &\quad + E \left[\int_{\mathbb{R}^d} \psi(x) \left(u_{\Delta t}((t_l)+, x) - u_{\Delta t}(t_{k+1}, x) \right) dx \right] \\ &=: \mathcal{T}_1 + \mathcal{T}_2 + \mathcal{T}_3. \end{aligned}$$

Recall that $u_{\Delta t}((t_n)+) = v_{\Delta t}((t_{n+1})-) = \mathcal{S}_{\text{CL}}(\Delta t)u^n$. Regarding the last term

$$u_{\Delta t}((t_l)+, x) - u_{\Delta t}(t_{k+1}, x) = v_{\Delta t}((t_{l+1})-, x) - v_{\Delta t}(t_l, x)$$

$$+ \sum_{n=k+1}^{l-1} u_{\Delta t}(t_{n+1}, x) - u_{\Delta t}(t_n, x)$$

Furthermore we note that

$$\begin{aligned} u_{\Delta t}(t_{n+1}, x) - u_{\Delta t}(t_n, x) &= (u_{\Delta t}(t_{n+1}, x) - u_{\Delta t}((t_n)+, x)) \\ &\quad + (v_{\Delta t}((t_{n+1})-, x) - v_{\Delta t}(t_n, x)). \end{aligned}$$

This yields

$$\begin{aligned} \mathcal{T}_3 &= E \left[\sum_{n=k+1}^{l-1} \int_{\mathbb{R}^d} \psi(x) \left(u_{\Delta t}(t_{n+1}, x) - u_{\Delta t}((t_n)+, x) \right) dx \right] \\ &\quad + E \left[\sum_{n=k+1}^l \int_{\mathbb{R}^d} \psi(x) \left(v_{\Delta t}((t_{n+1})-, x) - v_{\Delta t}(t_n, x) \right) dx \right] \\ &= E \left[\int_{\mathbb{R}^d} \psi(x) \int_{t_{k+1}}^{t_{l-1}} \sigma(x, u_{\Delta t}(t, x)) dB(t) dx \right] \\ &\quad + E \left[\sum_{n=k+1}^l \int_{\mathbb{R}^d} \psi(x) \left(\mathcal{S}_{\text{CL}}(\Delta t) u^n(x) - u^n(x) \right) dx \right] \end{aligned}$$

It follows that

$$\mathcal{T} = \mathcal{T}^{\text{CL}} + \mathcal{T}^{\text{SDE}}$$

where

$$\begin{aligned} \mathcal{T}^{\text{CL}} &:= E \left[\sum_{n=k+1}^l \int_{\mathbb{R}^d} \psi(x) \left(\mathcal{S}_{\text{CL}}(\Delta t) u^n(x) - u^n(x) \right) dx \right], \\ \mathcal{T}^{\text{SDE}} &:= E \left[\int_{\mathbb{R}^d} \psi(x) \int_{\tau_1}^{\tau_2} \sigma(x, u_{\Delta t}(t, x)) dB(t) dx \right]. \end{aligned}$$

Note that $\mathcal{T}^{\text{CL}} = 0$ if $k = l$. Let $v(t, x)$ be the entropy solution of the conservation law (3.1). Since v is a weak solution it follows for almost all $0 \leq s \leq t$

$$\begin{aligned} \left| \int_{\mathbb{R}^d} (v(t, x) - v(s, x)) \psi(x) dx \right| &\leq \left| \int_s^t \int_{\mathbb{R}^d} f(v(r, x)) \cdot \nabla \psi(x) dx dr \right| \\ &\leq \|f\|_{\text{Lip}} \int_s^t \int_{\mathbb{R}^d} |v(r, x)| |\nabla \varphi(x) \phi(x) + \varphi(x) \nabla \phi(x)| dx dr \\ &\leq \|f\|_{\text{Lip}} (\|\nabla \varphi\|_{L^\infty} + C_\phi \|\varphi\|_{L^\infty}) \left(\int_s^t \|v(r)\|_{1, \phi} dr \right). \end{aligned}$$

Hence we arrive at

$$|\mathcal{T}^{\text{CL}}| \leq \|f\|_{\text{Lip}} (\|\nabla \varphi\|_{L^\infty} + C_\phi \|\varphi\|_{L^\infty}) \left(\int_{t_{k+1}}^{t_{l+1}} \|v_{\Delta t}(t)\|_{1, \phi} dt \right).$$

By (4.7), Jensen's inequality and the Itô isometry

$$\begin{aligned} |\mathcal{T}^{\text{SDE}}| &\leq \|\varphi\|_{L^\infty} \int_{\mathbb{R}^d} E \left[\left| \int_{\tau_1}^{\tau_2} \sigma(x, u_{\Delta t}(t, x)) dB(t) \right|^2 \right] \phi(x) dx \\ &\leq \|\varphi\|_{L^\infty} \|\phi\|_{L^1(\mathbb{R}^d)}^{1/2} \left(\int_{\mathbb{R}^d} E \left[\left| \int_{\tau_1}^{\tau_2} \sigma(x, u_{\Delta t}(t, x)) dB(t) \right|^2 \right] \phi(x) dx \right)^{1/2} \\ &\leq \|\varphi\|_{L^\infty} \|\phi\|_{L^1(\mathbb{R}^d)}^{1/2} \left(\int_{\mathbb{R}^d} E \left[\int_{\tau_1}^{\tau_2} \sigma^2(x, u_{\Delta t}(t, x)) dt \right] \phi(x) dx \right)^{1/2} \end{aligned}$$

$$= \|\varphi\|_{L^\infty} \|\phi\|_{L^1(\mathbb{R}^d)}^{1/2} \left(\int_{\tau_1}^{\tau_2} E \left[\|\sigma(\cdot, u_{\Delta t}(t, \cdot))\|_{2, \phi}^2 \right] dt \right)^{1/2}$$

Due to $(\mathcal{A}_\sigma)$ there is a constant C such that $|\sigma(x, u_{\Delta t})| \leq C(1 + |u_{\Delta t}|)$. Appealing to Corollary 4.2 the above estimates implies the existence of a constant C independent of Δt and φ such that

$$|\mathcal{T}| \leq C \left((\|\nabla \varphi\|_{L^\infty} + \|\varphi\|_{L^\infty})(k - l)\Delta t + \|\varphi\|_{L^\infty} \sqrt{\tau_2 - \tau_1} \right),$$

which yields (4.16).

Let us consider $v_{\Delta t}$. Suppose $0 \leq \tau_1 \leq \tau_2 < T$ where $\tau_1 \in [t_k, t_{k+1})$ and $\tau_2 \in [t_l, t_{l+1})$. We claim that there exists a constant $C > 0$ independent of Δt and φ such that

$$\begin{aligned} E \left[\int_{\mathbb{R}^d} \varphi(x) \left(v_{\Delta t}(\tau_2, x) - v_{\Delta t}(\tau_1, x) \right) \phi(x) dx \right] \\ \leq C \left((\|\nabla \varphi\|_{L^\infty} + \|\varphi\|_{L^\infty}) |\tau_2 - \tau_1| + \|\varphi\|_{L^\infty} \sqrt{(l - k)\Delta t} \right). \end{aligned} \quad (4.17)$$

Note that

$$\begin{aligned} v_{\Delta t}(\tau_2, x) - v_{\Delta t}(\tau_1, x) &= v_{\Delta t}(\tau_2, x) - v_{\Delta t}(t_l, x) \\ &\quad + \sum_{n=k+1}^l v_{\Delta t}(t_n, x) - v_{\Delta t}((t_n)-, x) \\ &\quad + \sum_{n=k+1}^{l-1} v_{\Delta t}((t_{n+1})-, x) - v_{\Delta t}(t_n, x) \\ &\quad + v_{\Delta t}((t_{k+1})-, x) - v_{\Delta t}(\tau_1, x). \end{aligned}$$

This yields

$$E \left[\int_{\mathbb{R}^d} \psi(x) \left(v_{\Delta t}(\tau_2, x) - v_{\Delta t}(\tau_1, x) \right) dx \right] = \mathcal{T}^{\text{CL}} + \mathcal{T}^{\text{SDE}}$$

where

$$\begin{aligned} \mathcal{T}^{\text{CL}} &= E \left[\int_{\mathbb{R}^d} \psi(x) \left(\mathcal{S}_{\text{CL}}(\tau_2 - t_l) u^l(x) - u^l(x) \right) dx \right] \\ &\quad + \sum_{n=k+1}^{l-1} E \left[\int_{\mathbb{R}^d} \psi(x) \left(\mathcal{S}_{\text{CL}}(\Delta t) u^n(x) - u^n(x) \right) dx \right] \\ &\quad + E \left[\int_{\mathbb{R}^d} \psi(x) \left(\mathcal{S}_{\text{CL}}(\Delta t) u^k - \mathcal{S}_{\text{CL}}(\tau_1 - t_k) u^k \right) dx \right], \end{aligned}$$

and

$$\begin{aligned} \mathcal{T}^{\text{SDE}} &= \sum_{n=k+1}^l E \left[\int_{\mathbb{R}^d} \psi(x) \int_{t_{n-1}}^{t_n} \sigma(x, u_{\Delta t}(t, x)) dB(t) dx \right] \\ &= E \left[\int_{\mathbb{R}^d} \psi(x) \int_{t_k}^{t_l} \sigma(x, u_{\Delta t}(t, x)) dB(t) dx \right] \end{aligned}$$

Appealing to the above estimates yields (4.17).

2. Strong estimate. Let $d(x) = u_{\Delta t}(\tau_2) - u_{\Delta t}(\tau_1)$, define $\beta(x) = \text{sign}(d(x))$ and $\beta_\delta = \beta \star J_{2\delta}$. By the triangle inequality

$$\begin{aligned} E \left[\int_{\mathbb{R}^d} |u_{\Delta t}(\tau_2, x) - u_{\Delta t}(\tau_1, x)| \phi(x) dx \right] \\ \leq \left| E \left[\int_{\mathbb{R}^d} \beta_\delta(x) d(x) \phi(x) dx \right] \right| + E \left[\int_{\mathbb{R}^d} ||d(x)| - \beta_\delta(x) d(x)| \phi(x) dx \right] \end{aligned}$$

$$=: \mathcal{T}_1 + \mathcal{T}_2$$

Note that β_δ is in $L^\infty(\Omega; C^1(\mathbb{R}^d))$ for each $\delta > 0$ and $\|\nabla \beta_\delta\|_{L^\infty} = \mathcal{O}(1/\delta)$. By (4.16)

$$\mathcal{T}_1 = \mathcal{O}\left(\frac{1}{\delta}(l-k)\Delta t + \sqrt{\tau_2 - \tau_1}\right).$$

Consider $\mathcal{T}_2$. Following an argument found in [15, Lemma 1]

$$\begin{aligned} ||d(x)| - \beta_\delta(x)d(x)| &\leq \int_{\mathbb{R}^d} ||d(x)| - d(x)\text{sign}(d(y))| J_{2\delta}(x-y) dy \\ &\leq 2 \int_{\mathbb{R}^d} |d(x) - d(y)| J_{2\delta}(x-y) dy. \end{aligned}$$

Upon adding and subtracting and making the change of variables $2\tilde{x} = x + y$ and $2z = x - y$ it follows (after relabeling $\tilde{x}$ by x)

$$\begin{aligned} \int_{\mathbb{R}^d} ||d(x)| - \beta_\delta(x)d(x)| \phi(x) dx &\leq 2 \iint_{\mathbb{R}^d \times \mathbb{R}^d} |d(x+z) - d(x-z)| \\ &\quad \times J_\delta(z) |\phi(x+z) - \phi(x)| dz dx \\ &\quad + 2 \iint_{\mathbb{R}^d \times \mathbb{R}^d} |d(x+z) - d(x-z)| J_\delta(z) \phi(x) dz dx \\ &=: \mathcal{T}_2^2 + \mathcal{T}_2^1 \end{aligned}$$

Consider $\mathcal{T}_2^2$. By Lemma 7.2

$$|\phi(x+z) - \phi(x)| \leq w_{1,\phi}(|z|)\phi(x).$$

Hence by the symmetry of J_δ and the triangle inequality

$$\begin{aligned} |\mathcal{T}_2^2| &\leq 4 \iint_{\mathbb{R}^d \times \mathbb{R}^d} |d(x-z)| J_\delta(z) w_{1,\phi}(|z|) \phi(x) dz dx \\ &\leq 4w_{1,\phi}(\delta) \iint_{\mathbb{R}^d \times \mathbb{R}^d} |d(y)| J_\delta(x-y) \phi(x) dy dx \\ &= 4w_{1,\phi}(\delta) \|u_{\Delta t}(\tau_2) - u_{\Delta t}(\tau_1)\|_{1,\phi \star J_\delta} \end{aligned}$$

By Lemma 7.3 and Corollary 4.2 $E[|\mathcal{T}_2^2|] = \mathcal{O}(\delta)$. By Proposition 4.3 it follows due to assumption (4.15) that $E[|\mathcal{T}_2^1|] = \mathcal{O}(\delta^\kappa)$. Consequently

$$\mathcal{T}_1 + \mathcal{T}_2 = \mathcal{O}\left(\frac{1}{\delta}(l-k)\Delta t + \sqrt{\tau_2 - \tau_1} + \delta^\kappa\right)$$

Choosing $\delta = ((l-k)\Delta t)^{1/(\kappa+1)}$ concludes the proof of (i). The result (ii) follows analogously due to (4.17). $\square$

5. CONVERGENCE

Having established estimates uniform in Δt we are prepared to study the limiting behavior of $u_{\Delta t}$ and $v_{\Delta t}$ as $\Delta t \downarrow 0$. We will apply the framework of Young measures, cf. Section 7.4.

We start by establishing an approximate entropy inequality for the operator splitting.

Lemma 5.1. *Suppose $u_0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$, $\phi \in \mathfrak{N}$. Let $u_{\Delta t}$ and $v_{\Delta t}$ be defined by (3.4) and (3.5) respectively. For any $(S, Q) \in \mathcal{E}$, any $V \in \mathbb{D}^{1,2}$ and any non-negative $\varphi \in C_c^\infty([0, T] \times \mathbb{R}^d)$*

$$E\left[\int_{\mathbb{R}^d} S(u^0(x) - V)\varphi(0, x) dx\right]$$

$$\begin{aligned}
 & + E \left[\iint_{\Pi_T} S(u_{\Delta t}(t, x) - V) \partial_t \varphi(t, x) + Q(v_{\Delta t}(t, x), V) \cdot \nabla \varphi(t, x) \, dx dt \right] \\
 & - E \left[\iint_{\Pi_T} S''(u_{\Delta t}(t, x) - V) D_t V \sigma(u_{\Delta t}(t, x), x) \varphi(t, x) \, dx dt \right] \\
 & + E \left[\frac{1}{2} \iint_{\Pi_T} S''(u_{\Delta t}(t, x) - V) \sigma^2(u_{\Delta t}(t, x), x) \varphi(t, x) \, dx dt \right] \\
 & + \sum_{n=0}^{N-1} E \left[\int_{t_n}^{t_{n+1}} \int_{\mathbb{R}^d} (S(v_{\Delta t}(t, x) - V) - S(v_{\Delta t}((t_{n+1})-, x) - V)) \partial_t \varphi(t, x) \, dx dt \right] \geq 0.
 \end{aligned}$$

Proof. By definition $v_{\Delta t}$ satisfies:

$$\begin{aligned}
 & \int_{\mathbb{R}^d} S(u^n(x) - V) \varphi(t_n, x) \, dx - \int_{\mathbb{R}^d} S(v_{\Delta t}((t_{n+1})-, x) - V) \varphi(t_{n+1}, x) \, dx \\
 & + \int_{t_n}^{t_{n+1}} \int_{\mathbb{R}^d} S(v_{\Delta t}(t, x) - V) \partial_t \varphi(t, x) \, dx \, dt \\
 & + \int_{t_n}^{t_{n+1}} \int_{\mathbb{R}^d} Q(v_{\Delta t}(t, x), V) \cdot \nabla \varphi(t, x) \, dx \, dt \geq 0.
 \end{aligned}$$

By [21, Theorem 3.2.2],

$$\begin{aligned}
 \int_{\mathbb{R}^d} S(u^{n+1}(x) - V) \varphi(t_{n+1}, x) \, dx & = \int_{\mathbb{R}^d} S(u_{\Delta t}((t_n)+, x) - V) \varphi(t_n, x) \, dx \\
 & + \int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} S(u_{\Delta t}(t, x) - V) \partial_t \varphi(t, x) \, dt \, dx \\
 & + \int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} S'(u_{\Delta t}(t, x) - V) \sigma(u_{\Delta t}(t, x), x) \varphi(t, x) \, dB(t) \, dx \\
 & - \int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} S''(u_{\Delta t}(t, x) - V) D_t V \sigma(u_{\Delta t}(t, x), x) \varphi(t, x) \, dt \, dx \\
 & + \frac{1}{2} \int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} S''(u_{\Delta t}(t, x) - V) \sigma^2(u_{\Delta t}(t, x), x) \varphi(t, x) \, dt \, dx
 \end{aligned}$$

where the stochastic integral is a Skorohod integral. Note that

$$\begin{aligned}
 & \int_{\mathbb{R}^d} S(u_{\Delta t}((t_n)+, x) - V) \varphi(t_n, x) \, dx - \int_{\mathbb{R}^d} S(v_{\Delta t}((t_{n+1})-, x) - V) \varphi(t_{n+1}, x) \, dx \\
 & = - \int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} S(v_{\Delta t}((t_{n+1})-, x) - V) \partial_t \varphi(t, x) \, dt \, dx
 \end{aligned}$$

Adding the two equations and taking expectations yields

$$\begin{aligned}
& E \left[\int_{\mathbb{R}^d} S(u^n(x) - V) \varphi(t_n, x) dx \right] - E \left[\int_{\mathbb{R}^d} S(u^{n+1}(x) - V) \varphi(t_{n+1}, x) dx \right] \\
& + E \left[\int_{t_n}^{t_{n+1}} \int_{\mathbb{R}^d} (S(v_{\Delta t}(t, x) - V) - S(v_{\Delta t}((t_{n+1})-, x) - V)) \partial_t \varphi(t, x) dx dt \right] \\
& + E \left[\int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} S(u_{\Delta t}(t, x) - V) \partial_t \varphi(t, x) dt dx \right] \\
& + E \left[\int_{t_n}^{t_{n+1}} \int_{\mathbb{R}^d} Q(v_{\Delta t}(t, x), V) \cdot \nabla \varphi(t, x) dx dt \right] \\
& - E \left[\int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} S''(u_{\Delta t}(t, x) - V) D_t V \sigma(u_{\Delta t}(t, x), x) \varphi(t, x) dt dx \right] \\
& + E \left[\frac{1}{2} \int_{\mathbb{R}^d} \int_{t_n}^{t_{n+1}} S''(u_{\Delta t}(t, x) - V) \sigma^2(u_{\Delta t}(t, x), x) \varphi(t, x) dt dx \right] \geq 0.
\end{aligned}$$

Next we sum over $n = 0, 1, \dots, N-1$. The result follows. $\square$

Theorem 5.1. *Suppose $(\mathcal{A}_f)$, $(\mathcal{A}_{f,1})$, $(\mathcal{A}_\sigma)$ and $(\mathcal{A}_{\sigma,1})$ are satisfied. Let $\phi \in \mathfrak{N}$ and $2 \leq p < \infty$. Suppose $u_0 \in L^p(\Omega, \mathcal{F}_0, P; L^p(\mathbb{R}^d, \phi))$ satisfy (4.15). Let $u_{\Delta t}$ and $v_{\Delta t}$ be defined by (3.4) and (3.5) respectively. Then there exists a subsequence $\{\Delta t_j\}$ and a predictable $u \in L^p([0, T] \times \Omega; L^p(\mathbb{R}^d \times [0, 1], \phi))$ such that both $u_{\Delta t_j} \rightarrow u$ and $v_{\Delta t_j} \rightarrow u$ in the following sense:*

For any Carathodory function $\Psi : \mathbb{R} \times \Pi_T \times \Omega \rightarrow \mathbb{R}$ such that $\Psi(u_{\Delta t_j}, \cdot) \rightharpoonup \bar{\Psi}$ (respectively $\Psi(v_{\Delta t_j}, \cdot) \rightharpoonup \bar{\Psi}$) in $L^1(\Pi_T \times \Omega, d\phi \otimes dt \otimes dP)$ it follows that

$$\bar{\Psi}(t, x, \omega) = \int_0^1 \Psi(u(t, x, \alpha, \omega), t, x, \omega) d\alpha. \quad (5.1)$$

The process $\tilde{u} = \int_0^1 u d\alpha$ is an entropy solution in the sense of Definition 2.1.

Proof. Step 1: Existence of limits. Let us consider $u_{\Delta t}$, the exact same considerations apply to $v_{\Delta t}$. This follows exactly as in the proof of [11, Theorem 4.1], Step 1. Let us anyway give an outline of the argument. We apply Theorem 7.1 to $\{u_{\Delta t}\}$ on the measure space

$$(X, \mathcal{A}, \mu) = (\Omega \times \Pi_T, \mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d), dP \otimes dt \otimes d\phi).$$

By Corollary 4.2

$$\sup_{\Delta t > 0} \left\{ E \left[\iint_{\Pi_T} |u_{\Delta t}|^2 \phi(x) dx dt \right] \right\} < \infty.$$

Hence there exists a Young measure $\nu = \nu_{t,x,\omega}$ such that for any Carathodory function Ψ satisfying $\Psi(u_{\Delta t_j}, \cdot) \rightharpoonup \bar{\Psi}$ in $L^1(\Pi_T \times \Omega, d\phi \otimes dt \otimes dP)$ it follows that

$$\bar{\Psi}(t, x, \omega) = \int_{\mathbb{R}} \Psi(\xi, t, x, \omega) d\nu_{t,x,\omega}(\xi).$$

Let

$$u(t, x, \alpha, \omega) := \inf \{ \xi \in \mathbb{R} : \nu_{t,x,\omega}(-\infty, \xi] > \alpha \}.$$

The equality (5.1) follows by the fact that $\mathcal{L} \circ u^{-1}(t, x, \cdot, \omega) = \nu_{t,x,\omega}$ where $\mathcal{L}$ denotes the Lebesgue measure on $[0, 1]$. For predictability, and the fact that $u \in L^p([0, T] \times \Omega; L^p(\mathbb{R}^d \times [0, 1], \phi))$ see the proof of [11, Theorem 4.1].

Step 2: Independence of interpolation. Let us denote by v the limit obtained by considering $\{v_{\Delta t}\}$ instead of $\{u_{\Delta t}\}$ in Step 1. We want to show that $v = u$. By [22,

Lemma 6.3] this holds true if

$$\mathcal{T}(\Delta t) := E \left[\iint_{\Pi_T} |u_{\Delta t}(t, x) - v_{\Delta t}(t, x)| \phi(x) dt dx \right] \rightarrow 0 \text{ as } \Delta t \downarrow 0. \quad (5.2)$$

To this end observe that

$$\begin{aligned} \mathcal{T} &\leq E \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \int_{\mathbb{R}^d} |u_{\Delta t}(t, x) - u_{\Delta t}((t_n)+, x)| \phi(x) dt dx \right] \\ &\quad + E \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \int_{\mathbb{R}^d} |v_{\Delta t}((t_{n+1})-, x) - v_{\Delta t}(t, x)| \phi(x) dt dx \right] \\ &= \mathcal{T}_1 + \mathcal{T}_2. \end{aligned}$$

By Proposition 4.4(i) it follows that

$$\mathcal{T}_1 \leq C_{T,\psi} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \sqrt{t - t_n} dt = \frac{2}{3} C_{T,\psi} T \sqrt{\Delta t}.$$

By Proposition 4.4(ii) it follows that

$$\mathcal{T}_2 \leq C_{T,\phi} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} (t_{n+1} - t)^r dt \leq C_{T,\phi} T \Delta t^r$$

where $r = \frac{\kappa}{\kappa+1} > 0$ and κ is the exponent from Proposition 4.4. This proves (5.2).

Step 3: Entropy inequality. We need to prove that u is a Young measure-valued entropy solution in the sense of [11, Definition 2.2]. The result then follows by [11, Theorem 5.1]. Let S, V and φ be as in Lemma 5.1 and define

$$\mathcal{T}_{\Delta t} = \sum_{n=0}^{N-1} E \left[\int_{t_n}^{t_{n+1}} \int_{\mathbb{R}^d} (S(v_{\Delta t}(t, x) - V) - S(v_{\Delta t}((t_{n+1})-, x) - V) \partial_t \varphi(t, x) dx dt \right].$$

We want to show that $\mathcal{T}_{\Delta t} \rightarrow 0$ as $\Delta t \downarrow 0$. By Proposition 4.4

$$\begin{aligned} |\mathcal{T}_{\Delta t}| &\leq \|S\|_{\text{Lip}} \sup_{t \in [0, T]} \left\{ \|\partial_t \varphi\|_{\infty, \phi^{-1}} \right\} \\ &\quad \times \sum_{n=0}^{N-1} E \left[\int_{t_n}^{t_{n+1}} \int_{\mathbb{R}^d} |v_{\Delta t}(t, x) - v_{\Delta t}((t_{n+1})-, x)| \phi(x) dx dt \right] \\ &\leq \|S\|_{\text{Lip}} \sup_{t \in [0, T]} \left\{ \|\partial_t \varphi\|_{\infty, \phi^{-1}} \right\} C_{T,\phi} T \Delta t^r, \end{aligned}$$

as in the proof of Step 2. Concerning the remaining terms of Lemma 5.1, the limit $\Delta t \downarrow 0$ is treated exactly as in [11, Proof of Theorem 4.1, Step 2]. It follows that u is a Young measure-valued entropy solution. $\square$

6. ERROR ESTIMATE

We now restrict our attention to the case

$$\sigma(x, u) = \sigma(u), \quad \sigma \in L^\infty. \quad (\mathcal{A}_{\sigma,2})$$

We may now state the main result of this section.

Theorem 6.1. *Let $\phi \in \mathfrak{N}$. Suppose $(\mathcal{A}_f)$, $(\mathcal{A}_{f,1})$, $(\mathcal{A}_\sigma)$ and $(\mathcal{A}_{\sigma,2})$ are satisfied. Suppose furthermore that $u_0 \in L^2(\Omega, \mathcal{F}_0, P; L^2(\mathbb{R}^d, \phi))$ satisfies (4.15) with $\kappa_0 = 1$. Let u be an entropy solution of (1.1) – (1.2) according to Definition 2.1 and let $u_{\Delta t}$ be defined by (3.4). Then there exists a constant C , independent of Δt , but dependent on σ, f and T such that*

$$E \left[\|u(\tau) - u_{\Delta t}(\tau)\|_{1,\phi} \right] \leq C \Delta t^{\frac{1}{3}}.$$

for all $\tau \in [0, T]$.

The proof is split into several parts, and at long last the pieces are collected at the end of this section. To achieve this, inspired by [16], we consider yet another time interpolant. Let

$$\eta_{\Delta t}(t) := \underbrace{(\mathcal{S}_{\text{SDE}}(t, t_n) - \mathcal{I})\mathcal{S}_{\text{CL}}(\Delta t)u^n}_{u_{\Delta t}(t) - \mathcal{S}_{\text{CL}}(\Delta t)u^n} + \underbrace{\mathcal{S}_{\text{CL}}(t - t_n)u^n}_{v_{\Delta t}(t)}, \quad t \in [t_n, t_{n+1}].$$

Hopefully $\eta_{\Delta t}$ satisfies an approximation to (1.1). Formally we have that

$$\partial_t \eta_{\Delta t} + \operatorname{div} f(v_{\Delta t}) = \sigma(u_{\Delta t}) \partial_t W. \quad (6.1)$$

This line of reasoning is made rigorous by the approximate entropy inequality expressed in Proposition 6.1, cf. [16, Lemma 3.1]. Formally (6.1) expresses that the error terms are in general a perturbation of the coefficients f and σ . Due to regularity issues we choose to consider an approximation of $\eta_{\Delta t}$. Given $\{v^n = v^n(x)\}_{n=0}^{N-1}$ we let

$$\psi(t) = (\mathcal{S}_{\text{SDE}}(t, t_n) - \mathcal{I})v^n \text{ and } w(t) = \mathcal{S}_{\text{CL}}(t - t_n)u^n, \quad t \in [t_n, t_{n+1}]. \quad (6.2)$$

Note that $\eta_{\Delta t} = \psi + w$ whenever $v^n = \mathcal{S}_{\text{CL}}(\Delta t)u^n$ for $n = 0, \dots, N-1$. However, due to the lack of differentiability of $\mathcal{S}_{\text{CL}}(\Delta t)u^n$, we let v^n approach $\mathcal{S}_{\text{CL}}(\Delta t)u^n$ at a later stage.

Proposition 6.1. *Suppose $(\mathcal{A}_f)$ and $(\mathcal{A}_\sigma)$ are satisfied. Let $\tilde{\eta} = \psi + w$ where ψ and w are defined in (6.2). Then, for all $\phi \in C_c^\infty([t_n, t_{n+1}] \times \mathbb{R}^d)$ and all entropy, entropy-flux pairs $(S, Q) \in \mathcal{E}$*

$$\begin{aligned} & E \left[\int_{\mathbb{R}^d} S(\tilde{\eta}(t_n, x) - V) \phi(t_n, x) dx \right] \\ & - E \left[\int_{\mathbb{R}^d} S(\tilde{\eta}((t_{n+1})-, x) - V) \phi(t_{n+1}, x) dx \right] \\ & + E \left[\iint_{\Pi_n} S(\tilde{\eta} - V) \partial_t \phi + Q(\tilde{\eta}, V) \cdot \nabla \phi dx dt \right] \\ & + E \left[\iint_{\Pi_n} \int_V^{\tilde{\eta}} S'(z - V) (f'(z - \psi) - f'(z)) dz \cdot \nabla \phi dx dt \right] \\ & + E \left[\iint_{\Pi_n} \int_V^{\tilde{\eta}} S''(z - V) f'(z - \psi) dz \cdot \nabla \psi \phi dx dt \right] \\ & - E \left[\iint_{\Pi_n} S''(\tilde{\eta} - V) D_t V \sigma(\psi + v^n) \phi(t, x) dx dt \right] \\ & + \frac{1}{2} E \left[\iint_{\Pi_n} S''(\tilde{\eta} - V) \sigma^2(\psi + v^n) \phi(t, x) dx dt \right] \geq 0 \end{aligned}$$

where $\Pi_n = [t_n, t_{n+1}] \times \mathbb{R}$.

The proof of Proposition 6.1 is deferred to Section 7.1. To make sure that we work with Malliavin differentiable quantities, we substitute the entropy solution u with the viscous approximation u^ε , defined by

$$du^\varepsilon + \nabla \cdot f(u^\varepsilon) dt = \sigma(u^\varepsilon) dB(t) + \varepsilon \Delta u^\varepsilon dt, \quad u^\varepsilon(0) = u_0,$$

and then let $\varepsilon \downarrow 0$ at a later stage. See [11, § 3] regarding properties of u^ε . As a consequence of [11, Theorem 5.1] and [22, Proposition 6.12] it follows that $u^\varepsilon \rightarrow u$ in $L^1([0, T] \times \Omega; L^1(\mathbb{R}^d, \phi))$ as $\varepsilon \downarrow 0$. In fact, under the assumptions of Theorem 6.1 $u^\varepsilon \rightarrow u$ with rate $1/2$ [5, Theorem 5.2]. We may now proceed to the doubling-of-the-variables argument.

Lemma 6.2. *Let $\phi \in \mathfrak{N}$. Let $u^\varepsilon = u^\varepsilon(s, y)$ be the viscous approximation of (1.1). Let $v(t, x) = v^n(x)$ for $t \in [t_n, t)$ and $\psi = \psi(t, x), w = w(t, x), \tilde{\eta} = \tilde{\eta}(t, x)$ be defined in Proposition 6.1. For any $0 < \gamma < \frac{1}{2}T$ take $t_0 \in [0, T - 2\gamma]$ and define*

$$\xi_\gamma(t) = 1 - \int_0^t J_\gamma^+(s - t_0) ds.$$

Furthermore, let

$$\varphi(t, x, s, y) = \frac{1}{2^d} \phi\left(\frac{x+y}{2}\right) J_r\left(\frac{x-y}{2}\right) J_{r_0}^+(s-t) \xi_\gamma(t),$$

and S_δ be defined by (2.2). Then

$$L - R + F + \mathcal{T}_1 + \mathcal{T}_2 + \mathcal{T}_3 + \mathcal{T}_4 + \mathcal{T}_5 + \mathcal{T}_6 + \mathcal{T}_7 \geq 0,$$

where

$$\begin{aligned} L &= E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} S_\delta(\tilde{\eta}(0, x) - u^\varepsilon(s, y)) \varphi(0, x, s, y) dx ds dy \right], \\ R &= -E \left[\iiint_{\Pi_T^2} S_\delta(\tilde{\eta} - u^\varepsilon) (\partial_t + \partial_s) \varphi dX \right], \\ F &= E \left[\iiint_{\Pi_T^2} Q(u^\varepsilon, \tilde{\eta}) \cdot \nabla_y \varphi + Q(\tilde{\eta}, u^\varepsilon) \cdot \nabla_x \varphi dX \right], \\ \mathcal{T}_1 &= \frac{1}{2} E \left[\iiint_{\Pi_T^2} S_\delta''(u^\varepsilon - \tilde{\eta}) (\sigma(u^\varepsilon) - \sigma(\tilde{\eta}))^2 \varphi dX \right], \\ \mathcal{T}_2 &= E \left[\iiint_{\Pi_T^2} S_\delta''(u^\varepsilon - \tilde{\eta}) (\sigma(u^\varepsilon) - D_t u^\varepsilon) \sigma(\psi + v) \varphi dX \right], \\ \mathcal{T}_3 &= E \left[\iiint_{\Pi_T^2} S_\delta''(u^\varepsilon - \tilde{\eta}) \left(\int_{\tilde{\eta}}^{\psi+v} (\sigma(z) - \sigma(u^\varepsilon)) \sigma'(z) dz \right) \varphi dX \right], \\ \mathcal{T}_4 &= E \left[\iiint_{\Pi_T^2} \int_{u^\varepsilon}^{\tilde{\eta}} S'(z - u^\varepsilon) (f'(z - \psi) - f'(z)) dz \cdot \nabla_x \varphi dX \right] \\ &\quad + E \left[\iiint_{\Pi_T^2} \int_{u^\varepsilon}^{\tilde{\eta}} S''(z - u^\varepsilon) f'(z - \psi) dz \cdot \nabla_x \psi \varphi dX \right], \\ \mathcal{T}_5 &= \varepsilon E \left[\iiint_{\Pi_T^2} S_\delta(u^\varepsilon - \tilde{\eta}) \Delta_y \varphi dX \right], \\ \mathcal{T}_6 &= - \sum_{n=0}^{N-1} E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} \left(S_\delta(\tilde{\eta}((t_{n+1}), x) - u^\varepsilon(s, y)) \right. \right. \\ &\quad \left. \left. - S_\delta(\tilde{\eta}((t_{n+1})-, x) - u^\varepsilon(s, y)) \right) \varphi(t_{n+1}, x, s, y) dx ds dy \right] \end{aligned}$$

Proof. Let us first assume $\phi \in C_c^\infty(\mathbb{R}^d)$, the result for $\phi \in \mathfrak{N}$ follows by approximation. Setting $V = \tilde{\eta}(t, x)$ in [11, Lemma 5.3] we get

$$\begin{aligned} &E \left[\iiint_{\Pi_T^2} S_\delta(u^\varepsilon - \tilde{\eta}) \partial_s \varphi + Q(u^\varepsilon, \tilde{\eta}) \cdot \nabla_y \varphi dX \right] \\ &\quad + \frac{1}{2} E \left[\iiint_{\Pi_T^2} S_\delta''(u^\varepsilon - \tilde{\eta}) \sigma^2(u^\varepsilon) \varphi dX \right] + \varepsilon E \left[\iiint_{\Pi_T^2} S_\delta(u^\varepsilon - \tilde{\eta}) \Delta_y \varphi dX \right] \geq 0. \end{aligned}$$

Take $V = u^\varepsilon(s, y)$ in Proposition 6.1, integrate in (s, y) over Π_T and sum over $n = 0, \dots, N-1$. This yields

$$E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} S_\delta(\tilde{\eta}(0, x) - u^\varepsilon(s, y)) \varphi(0, x, s, y) dx ds dy \right]$$

$$\begin{aligned}
& + E \left[\iiint_{\Pi_T^2} S_\delta(\tilde{\eta} - u^\varepsilon) \partial_t \varphi + Q(\tilde{\eta}, u^\varepsilon) \cdot \nabla_x \varphi dX \right] \\
& + E \left[\iiint_{\Pi_T^2} \int_{u^\varepsilon}^{\tilde{\eta}} S'(z - u^\varepsilon) (f'(z - \psi) - f'(z)) dz \cdot \nabla_x \varphi dX \right] \\
& + E \left[\iiint_{\Pi_T^2} \int_{u^\varepsilon}^{\tilde{\eta}} S''(z - u^\varepsilon) f'(z - \psi) dz \cdot \nabla_x \psi \varphi dX \right] \\
& - E \left[\iiint_{\Pi_T^2} S_\delta''(\tilde{\eta} - u^\varepsilon) D_t u^\varepsilon \sigma(\psi + v) \varphi dX \right] \\
& + \frac{1}{2} E \left[\iiint_{\Pi_T^2} S_\delta''(\tilde{\eta} - u^\varepsilon) \sigma^2(\psi + v) \varphi dX \right] \\
& - \sum_{n=0}^{N-1} E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} \left(S_\delta(\tilde{\eta}((t_{n+1}), x) - u^\varepsilon(s, y)) \right. \right. \\
& \quad \left. \left. - S_\delta(\tilde{\eta}((t_{n+1})-, x) - u^\varepsilon(s, y)) \right) \varphi(t_{n+1}, x, s, y) dx ds dy \right] \geq 0
\end{aligned}$$

Note that

$$\begin{aligned}
& \frac{1}{2} \sigma^2(u^\varepsilon) - D_t u^\varepsilon \sigma(\psi + v) + \frac{1}{2} \sigma^2(\psi + v) \\
& = \frac{1}{2} (\sigma(\psi + v) - \sigma(u^\varepsilon))^2 + (\sigma(u^\varepsilon) - D_t u^\varepsilon) \sigma(\psi + v) \\
& = \frac{1}{2} (\sigma(\tilde{\eta}) - \sigma(u^\varepsilon))^2 + \int_{\tilde{\eta}}^{\psi+v} (\sigma(z) - \sigma(u^\varepsilon)) \sigma'(z) dz \\
& \quad + (\sigma(u^\varepsilon) - D_t u^\varepsilon) \sigma(\psi + v).
\end{aligned}$$

The result follows upon adding the two inequalities. $\square$

Next follows a series of estimates of the terms appearing in Lemma 6.2. The underlying assumptions are the ones made in Theorem 6.1. In general we let C denote a generic constant, meaning that it is independent of the “small” parameters $\Delta t, r, r_0, \gamma, \varepsilon$ and δ . Similar considerations apply to the Landau notation $\mathcal{O}$.

Estimate 6.1. *Let L be defined in Lemma 6.2. Then*

$$\lim_{r_0 \downarrow 0} L = E[\|u_0 - u^0\|_{1,\phi}] + \mathcal{O}(\delta + r).$$

Proof. Recall that

$$L = E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} S_\delta(\tilde{\eta}(0, x) - u^\varepsilon(s, y)) \varphi(0, x, s, y) dx ds dy \right].$$

By (2.3)

$$|S_\delta(\tilde{\eta}(0, x) - u^\varepsilon(s, y)) - \tilde{\eta}(0, x) - u^\varepsilon(s, y)| \leq \delta.$$

By the inverse triangle inequality

$$\begin{aligned}
& ||\tilde{\eta}(0, x) - u^\varepsilon(s, y)| - |\tilde{\eta}(0, x) - u_0(y)|| \leq |u^\varepsilon(s, y) - u_0(y)|, \\
& ||\tilde{\eta}(0, x) - u_0(y)| - |\tilde{\eta}(0, x) - u_0(x)|| \leq |u_0(y) - u_0(x)|.
\end{aligned}$$

Hence, adding and subtracting, noting that $\tilde{\eta}(0) = u^0$ it follows by the triangle inequality that

$$\begin{aligned}
& |S_\delta(\tilde{\eta}(0, x) - u^\varepsilon(s, y)) - |u^0(x) - u_0(x)|| \\
& \leq \delta + |u^\varepsilon(s, y) - u_0(y)| + |u_0(y) - u_0(x)|
\end{aligned}$$

By (4.9)

$$\begin{aligned} & \left| L - E \left[\|u^0 - u_0\|_{1, \phi \star J_r} \right] \right| \\ & \leq \delta \|\phi\|_{L^1(\mathbb{R}^d)} + \underbrace{\int_0^T E \left[\|u^\varepsilon(s) - u_0\|_{1, \phi \star J_r} \right] J_{r_0}^+(s) ds}_{\mathcal{Z}_1} \\ & \quad + \underbrace{E \left[\frac{1}{2^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} |u_0(y) - u_0(x)| \phi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) dx dy \right]}_{\mathcal{Z}_2}. \end{aligned}$$

By [11, Lemma 2.3] we have that $\mathcal{Z}_1 \rightarrow 0$ as $r_0 \rightarrow 0$. Regarding $\mathcal{Z}_2$ we apply (4.8). Due to the assumption (4.15) with $\kappa_0 = 1$

$$\mathcal{Z}_2 = E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u_0(x+z) - u_0(x-z)| \phi(x) J_r(z) dx dz \right] = \mathcal{O}(r).$$

Finally we apply Lemma 7.3 to conclude that

$$\left| E \left[\|u^0 - u_0\|_{1, \phi \star J_r} - \|u^0 - u_0\|_{1, \phi} \right] \right| = \mathcal{O}(r).$$

□

Estimate 6.2. *Let R be defined in Lemma 6.2. Then*

$$\lim_{\varepsilon, r_0 \downarrow 0} R = E \left[\int_0^T \|\tilde{\eta}(t) - u(t)\|_{1, \phi} J_\gamma^+(t - t_0) dt \right] + \mathcal{O}(\delta + r).$$

Proof. Differentiating it follows that

$$R = E \left[\iiint_{\Pi_T^2} S_\delta(\tilde{\eta}(t, x) - u^\varepsilon(s, y)) \frac{1}{2^d} \phi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) J_{r_0}(s-t) J_\gamma^+(t-t_0) dX \right].$$

Adding and subtracting we obtain

$$\begin{aligned} |S_\delta(\tilde{\eta}(t, x) - u^\varepsilon(s, y)) - |\tilde{\eta}(t, x) - u^\varepsilon(t, x)|| & \leq \\ & \delta + |u^\varepsilon(s, y) - u^\varepsilon(t, y)| + |u^\varepsilon(t, y) - u^\varepsilon(t, x)|. \end{aligned}$$

This yields

$$\begin{aligned} & \left| R - E \left[\int_0^T \|\tilde{\eta}(t) - u^\varepsilon(t)\|_{1, \phi \star J_r} J_\gamma^+(t - t_0) dt \right] \right| \\ & \leq \delta \|\phi\|_{L^1(\mathbb{R}^d)} + \underbrace{E \left[\iint_{[0, T]^2} \|u^\varepsilon(s) - u^\varepsilon(t)\|_{\phi \star J_r} J_{r_0}^+(s-t) J_\gamma^+(t-t_0) ds dt \right]}_{\mathcal{Z}_1} \\ & \quad + \underbrace{E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} |u^\varepsilon(t, y) - u^\varepsilon(t, x)| \frac{1}{2^d} \phi \left(\frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) J_\gamma^+(t-t_0) dx dy dt \right]}_{\mathcal{Z}_2}. \end{aligned}$$

By Lemma 7.4 it follows that $\lim_{r_0 \downarrow 0} \mathcal{Z}_1 = 0$. Next we apply the fact that $u^\varepsilon \rightarrow u$ in $L^1([0, T] \times \Omega; L^1(\mathbb{R}^d, \phi))$ and (4.8) to conclude that

$$\lim_{\varepsilon, r_0 \downarrow 0} \mathcal{Z}_2 = \int_0^T E \left[\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u(t, x+z) - u(t, x-z)| \phi(x) J_r(z) dx dz \right] J_\gamma^+(t-t_0) dt.$$

It follows by [11, Proposition 5.2] and the assumption (4.15) with $\kappa_0 = 1$ that $|\lim_{\varepsilon, r_0 \downarrow 0} \mathcal{Z}_2| = \mathcal{O}(r)$. The result then follows by Lemma 7.3. □

Estimate 6.3. *Let F be defined in Lemma 6.2. Then*

$$\lim_{\varepsilon, r_0 \downarrow 0} F \leq C_\phi \|f\|_{\text{Lip}} E \left[\int_0^T \|u(t) - \tilde{\eta}(t)\|_{1,\phi} \xi_\gamma(t) dt \right] + \mathcal{O} \left(\delta \left(1 + \frac{1}{r} \right) + r \right).$$

Proof. We first observe that

$$F = F_1 + F_2 + F_3 \tag{6.3}$$

where

$$\begin{aligned} F_1 &= E \left[\iiint_{\Pi_T^2} S'_\delta(u^\varepsilon - \tilde{\eta})(f(u^\varepsilon) - f(\tilde{\eta}))(\nabla_x + \nabla_y)\varphi dX \right], \\ F_2 &= -E \left[\iiint_{\Pi_T^2} \int_{u^\varepsilon}^{\tilde{\eta}} S''_\delta(z - u^\varepsilon)(f(z) - f(u^\varepsilon)) dz \cdot \nabla_x \varphi dX \right], \\ F_3 &= -E \left[\iiint_{\Pi_T^2} \int_{\tilde{\eta}}^{u^\varepsilon} S''_\delta(z - \tilde{\eta})(f(z) - f(\tilde{\eta})) dz \cdot \nabla_y \varphi dX \right]. \end{aligned}$$

To this end recall that

$$F = E \left[\iiint_{\Pi_T^2} Q_\delta(u^\varepsilon, \tilde{\eta}) \cdot \nabla_y \varphi + Q_\delta(\tilde{\eta}, u^\varepsilon) \cdot \nabla_x \varphi dX \right].$$

Integration by parts yields

$$\begin{aligned} Q_\delta(u^\varepsilon, \tilde{\eta}) &= S'_\delta(u^\varepsilon - \tilde{\eta})(f(u^\varepsilon) - f(\tilde{\eta})) - \int_{\tilde{\eta}}^{u^\varepsilon} S''_\delta(z - \tilde{\eta})(f(z) - f(\tilde{\eta})) dz, \\ Q_\delta(\tilde{\eta}, u^\varepsilon) &= S'_\delta(\tilde{\eta} - u^\varepsilon)(f(\tilde{\eta}) - f(u^\varepsilon)) - \int_{u^\varepsilon}^{\tilde{\eta}} S''_\delta(z - u^\varepsilon)(f(z) - f(u^\varepsilon)) dz. \end{aligned}$$

This yields (6.3). Next we want to show that

$$|F_2| + |F_3| = \mathcal{O} \left(\delta \left(1 + \frac{1}{r} \right) \right). \tag{6.4}$$

Consider F_2 . The term F_3 is estimated analogously. Note that

$$\left| \int_{u^\varepsilon}^{\tilde{\eta}} S''_\delta(z - u^\varepsilon)(f(z) - f(u^\varepsilon)) dz \right| \leq \|f\|_{\text{Lip}} \delta.$$

Hence it follows that

$$|F_3| \leq \|f\|_{\text{Lip}} \delta E \left[\iiint_{\Pi_T^2} |\nabla_x \varphi| dX \right].$$

A straight forward computation yields the existence of a constant $C > 0$ such that

$$\iiint_{\Pi_T^2} |\nabla_x \varphi| dX \leq C \left(1 + \frac{1}{r} \right).$$

This proves (6.4). Next we claim that

$$\lim_{\varepsilon, r_0 \downarrow 0} F_1 \leq C_\phi \|f\|_{\text{Lip}} E \left[\int_0^T \|u(t) - \tilde{\eta}(t)\|_{1,\phi * J_r} \xi_\gamma(t) dt \right] + \mathcal{O}(\delta + r). \tag{6.5}$$

Let

$$\mathcal{F}_\delta(a, b) = S'_\delta(a - b)(f(a) - f(b)).$$

Then [11, Limit 3]

$$|\mathcal{F}_\delta(b, a) - \mathcal{F}_\delta(c, a)| \leq 2 \|f\|_{\text{Lip}} \delta + \|f\|_{\text{Lip}} |b - c|.$$

It follows that

$$|\mathcal{F}_\delta(u^\varepsilon(s, y), \tilde{\eta}(t, x)) - \mathcal{F}_\delta(u^\varepsilon(t, x), \tilde{\eta}(t, x))|$$

$$\leq \|f\|_{\text{Lip}} (2\delta + |u^\varepsilon(s, y) - u^\varepsilon(t, y)| + |u^\varepsilon(t, y) - u^\varepsilon(t, x)|).$$

Hence

$$\begin{aligned} & \left| F_1 - E \left[\iint_{\Pi_T} \mathcal{F}_\delta(u^\varepsilon(t, x), \tilde{\eta}(t, x)) \cdot (\nabla \phi * J_r)(x) \xi_\gamma(t) dx dt \right] \right| \\ & \leq C_\phi \|f\|_{\text{Lip}} E \left[\iint_{[0, T]^2} \|u^\varepsilon(s) - u^\varepsilon(t)\|_{1, \phi * J_r} J_{r_0}^+(s - t) \xi_\gamma(t) ds dt \right] \\ & + C_\phi \|f\|_{\text{Lip}} E \left[\int_0^T \iint_{\mathbb{R}^d \times \mathbb{R}^d} |u^\varepsilon(t, x + z) - u^\varepsilon(t, x - z)| |\phi(x)| J_r(z) \xi_\gamma(t) dx dz dt \right] \\ & + 2\delta \|f\|_{\text{Lip}} T \|\nabla \phi\|_{L^1(\mathbb{R}^d)}, \end{aligned}$$

where we have made a change of variables as in Estimate 6.2. Following the same reasoning as in Estimate 6.2 it follows that

$$\left| \lim_{\varepsilon, r_0 \downarrow 0} F_1 - E \left[\iint_{\Pi_T} \mathcal{F}_\delta(u(t, x), \tilde{\eta}(t, x)) \cdot (\nabla \phi * J_r)(x) \xi_\gamma(t) dx dt \right] \right| = \mathcal{O}(\delta + r).$$

The inequality (6.5) follows by the fact that $\mathcal{F}_\delta(a, b) \leq \|f\|_{\text{Lip}} |a - b|$, $|\nabla \phi| \leq C_\phi \phi$. The result follows by combining the above steps. $\square$

Estimate 6.4. Let $\mathcal{T}_1$ be defined in Lemma 6.2. Then

$$|\mathcal{T}_1| \leq C\delta$$

Proof. Recall that

$$\mathcal{T}_1 = \frac{1}{2} E \left[\iiint_{\Pi_T^2} S_\delta''(u^\varepsilon - \tilde{\eta})(\sigma(u^\varepsilon) - \sigma(\tilde{\eta}))^2 \varphi dX \right].$$

Recall that $S_\delta'' = 2J_\delta$. Hence

$$S_\delta''(u^\varepsilon - \tilde{\eta})(\sigma(u^\varepsilon) - \sigma(\tilde{\eta}))^2 \leq 2 \|\sigma\|_{\text{Lip}}^2 J_\delta(u^\varepsilon - \tilde{\eta}) |u^\varepsilon - \tilde{\eta}|^2 \leq 2 \|\sigma\|_{\text{Lip}}^2 \|J\|_\infty \delta.$$

Due to (4.9) and Young's inequality for convolutions

$$\iiint_{\Pi_T^2} \varphi dX = \left(\int_0^T \int_0^T J_{r_0}^+(s - t) \xi_\gamma(t) ds dt \right) \left(\int_{\mathbb{R}^d} \phi * J_r(x) dx \right) \leq T \|\phi\|_{L^1(\mathbb{R}^d)}.$$

The result follows. $\square$

Estimate 6.5. Let $\mathcal{T}_2$ be defined in Lemma 6.2. Then

$$\lim_{r_0 \downarrow 0} \mathcal{T}_2 = 0$$

Proof. This follows exactly as in [11, Limit 5]. However, letting $\sigma \in L^\infty$ simplifies the analysis, and allows for $\phi \in \mathfrak{N}$ instead of $C_c^\infty(\mathbb{R}^d)$. $\square$

Estimate 6.6. Let $\mathcal{T}_3$ be defined in Lemma 6.2. Then

$$|\mathcal{T}_3| \leq C \frac{1}{\delta} E \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \|v^n - w(t)\|_{\phi * J_r} dt \right]$$

Proof. Recall that

$$\mathcal{T}_3 = E \left[\iiint_{\Pi_T^2} S_\delta''(u^\varepsilon - \tilde{\eta}) \int_{\tilde{\eta}}^{\psi+v} (\sigma(z) - \sigma(u^\varepsilon)) \sigma'(z) dz \varphi dX \right].$$

Now, as $\tilde{\eta} = \psi + w$

$$\left| \int_{\tilde{\eta}}^{\psi+v} (\sigma(z) - \sigma(u^\varepsilon)) \sigma'(z) dz \right| \leq 2 \|\sigma\|_\infty \|\sigma\|_{\text{Lip}} |v - w|.$$

Recall that $v(t) = v^n$ for $t \in [t_n, t_{n+1})$. The result follows by (2.3) and (4.9). $\square$

Estimate 6.7. *Let $\mathcal{T}_4$ be defined in Lemma 6.2. Then*

$$|\mathcal{T}_4| \leq C\sqrt{\Delta t} \left(1 + E \left[\int_0^T \|\nabla v(t)\|_{1,\phi \star J_r} dt \right] \right).$$

Proof. Recall that

$$\begin{aligned} \mathcal{T}_4 &= E \left[\iiint_{\Pi_T^2} \int_{u^\varepsilon}^{\tilde{\eta}} S'_\delta(z - u^\varepsilon) (f'(z - \psi) - f'(z)) dz \cdot \nabla_x \varphi dX \right] \\ &\quad + E \left[\iiint_{\Pi_T^2} \int_{u^\varepsilon}^{\tilde{\eta}} S''_\delta(z - u^\varepsilon) f'(z - \psi) dz \cdot \nabla_x \psi \varphi dX \right]. \end{aligned}$$

Assume that w is smooth. The general result follows by approximation. Integration by parts and the chain rule yields

$$\begin{aligned} &E \left[\iiint_{\Pi_T^2} \int_{u^\varepsilon}^{\tilde{\eta}} S'_\delta(z - u^\varepsilon) (f'(z - \psi) - f'(z)) dz \cdot \nabla_x \varphi dX \right] \\ &= -E \left[\iiint_{\Pi_T^2} S'_\delta(\tilde{\eta} - u^\varepsilon) (f'(w) - f'(\tilde{\eta})) \cdot \nabla_x \tilde{\eta} \varphi dX \right] \\ &\quad + E \left[\iiint_{\Pi_T^2} \int_{u^\varepsilon}^{\tilde{\eta}} S'_\delta(z - u^\varepsilon) f''(z - \psi) dz \cdot \nabla_x \psi \varphi dX \right]. \end{aligned}$$

Next we observe that

$$\int_{u^\varepsilon}^{\tilde{\eta}} S'_\delta(z - u^\varepsilon) f''(z - \psi) dz = - \int_{u^\varepsilon}^{\tilde{\eta}} S''_\delta(z - u^\varepsilon) f'(z - \psi) dz + S'_\delta(\tilde{\eta} - u^\varepsilon) f'(w).$$

Combining the above we arrive at

$$\begin{aligned} \mathcal{T}_4 &= E \left[\underbrace{\iiint_{\Pi_T^2} S'_\delta(\tilde{\eta} - u^\varepsilon) f'(\tilde{\eta}) \cdot \nabla_x \psi \varphi dX}_{\mathcal{Z}_1} \right] \\ &\quad + E \left[\underbrace{\iiint_{\Pi_T^2} S'_\delta(\tilde{\eta} - u^\varepsilon) (f'(\tilde{\eta}) - f'(w)) \cdot \nabla_x w \varphi dX}_{\mathcal{Z}_2} \right]. \end{aligned}$$

Concider $\mathcal{Z}_2$. Since $w(t)$ is $\mathcal{F}_{t_n}$ -measurable for all $t \in [t_n, t_{n+1})$

$$\begin{aligned} |\mathcal{Z}_2| &\leq E \left[\iiint_{\Pi_T^2} |f'(\tilde{\eta}) - f'(w)| |\nabla_x w| \varphi dX \right] \\ &\leq \|f'\|_{\text{Lip}} \sum_{n=0}^{N-1} \iiint_{\Pi_T \times \Pi_n} E \left[E[|\psi| | \mathcal{F}_{t_n}] |\nabla_x w| \right] \varphi dX. \end{aligned}$$

By definition

$$\psi(t, x) = \int_{t_n}^t \sigma(\psi(r, x) + v^n(x)) dB(r), \quad t_n \leq t < t_{n+1}. \quad (6.6)$$

By the Burkholder-Davies-Gundy inequality there exists a constant $c_1 > 0$ such that

$$E \left[|\psi| | \mathcal{F}_{t_n} \right] \leq c_1 E \left[\left(\int_{t_n}^t \sigma^2(\psi(r, x) + v^n(x)) dr \right)^{1/2} \middle| \mathcal{F}_{t_n} \right] \leq c_1 \|\sigma\|_\infty \sqrt{t - t_n}.$$

It follows by Proposition 4.3

$$|\mathcal{Z}_2| \leq c_1 \|\sigma\|_\infty \|f'\|_{\text{Lip}} \sqrt{\Delta t} E \left[\int_0^T \|\nabla_x w(t)\|_{1,\phi \star J_r} dt \right] \leq C\sqrt{\Delta t}$$

Consider $\mathcal{Z}_1$. By (4.9)

$$|\mathcal{Z}_1| \leq \|f\|_{\text{Lip}} E \left[\iiint_{\Pi_T^2} |\nabla_x \psi| \varphi dX \right] \leq \|f\|_{\text{Lip}} E \left[\iint_{\Pi_T} |\nabla_x \psi| (\phi \star J_r) dx dt \right].$$

Differentiating (6.6) it follows that

$$\nabla_x \psi(t, x) = \int_{t_n}^t \sigma'(\psi(r, x) + v^n(x)) (\nabla_x \psi(r, x) + \nabla v^n(x)) dB(r), \quad t_n \leq t < t_{n+1}.$$

By Lemma 6.3 below there exists a constant $C > 0$, depending only on σ , such that

$$E \left[|\nabla_x \psi(t, x)| \right] \leq C \sqrt{t - t_n} E \left[|\nabla v^n(x)| \right], \quad t_n \leq t < t_{n+1}.$$

It follows that

$$|\mathcal{Z}_1| \leq C \left(E \left[\int_0^T \|\nabla v(t)\|_{1, \phi \star J_r} dt \right] \right) \sqrt{\Delta t}.$$

□

Lemma 6.3. *Let $1 \leq p < \infty$. Suppose $h : [t_n, t_{n+1}] \times \Omega \rightarrow \mathbb{R}$ is jointly measurable, $\mathcal{F}_t$ -adapted and*

$$P \left[\int_{t_n}^t |h(s)|^2 ds < \infty \right] = 1.$$

Suppose $X(t_n) \in L^p(\Omega, \mathcal{F}_{t_n}, P)$ and let the process $X : [t_n, t_{n+1}] \times \Omega \rightarrow \mathbb{R}$ satisfy

$$X(t) = X(t_n) + \int_{t_n}^t h(s) dB(s), \quad t \in [t_n, t_{n+1}].$$

Suppose there exists a constant K and $Y \in L^p(\Omega, \mathcal{F}_{t_n}, P)$ such that

$$|h(t; \omega)| \leq Y(\omega) + K |X(t)| \quad (6.7)$$

for all $t \in [t_n, t_{n+1}]$. Then for all $t \in [t_n, t_{n+1}]$, $\beta > p(c_p^{1/p} K)^2/2$

$$E \left[\sup_{t_n \leq s \leq t} |X(s)|^p \right]^{1/p} \leq C(\beta) e^{\beta(t-t_n)} \left(E \left[|X(t_n)|^p \right]^{1/p} + c_p^{1/p} \sqrt{t - t_n} E \left[|Y|^p \right]^{1/p} \right),$$

where $C(\beta) = (1 - c_p^{1/p} K \sqrt{p/2\beta})^{-1}$ and c_p is the constant from the Burkholder-Davies-Gundy inequality.

Proof. Let

$$\|X\|_{\beta, p, \tau} := \left(\sup_{t_n \leq t \leq \tau} e^{-\beta(t-t_n)} E \left[|X(t)|^p \right] \right)^{1/p}$$

By the triangle inequality

$$\begin{aligned} \left(e^{-\beta(t-t_n)} E \left[|X(t)|^p \right] \right)^{1/p} &\leq \left(e^{-\beta(t-t_n)} E \left[\left| \int_{t_n}^t h(s) dB(s) \right|^p \right] \right)^{1/p} \\ &\quad + \left(e^{-\beta(t-t_n)} E \left[|X(t_n)|^p \right] \right)^{1/p}. \end{aligned} \quad (6.8)$$

By the Burkholder-Davies-Gundy inequality

$$E \left[\left| \int_{t_n}^t h(s) dB(s) \right|^p \right] \leq c_p E \left[\left(\int_{t_n}^t h^2(s) ds \right)^{p/2} \right]$$

By (6.7) and Minkowski's inequality

$$T(t) = \left(e^{-\beta(t-t_n)} E \left[\left(\int_{t_n}^t h^2(s) ds \right)^{p/2} \right] \right)^{1/p}$$

$$\begin{aligned}
&\leq E \left[\left(e^{-2\beta(t-t_n)/p} \int_{t_n}^t (Y + K_2 |X(s)|)^2 ds \right)^{p/2} \right]^{1/p} \\
&\leq e^{-\beta(t-t_n)/p} \sqrt{t-t_n} E[|Y|^p]^{1/p} \\
&\quad + K_2 \left(e^{-2\beta(t-t_n)/p} E \left[\left(\int_{t_n}^t |X(s)|^2 ds \right)^{p/2} \right]^{2/p} \right)^{1/2}.
\end{aligned}$$

Due to Minkowski's integral inequality

$$\begin{aligned}
e^{-2\beta(t-t_n)/p} E \left[\left(\int_{t_n}^t |X(s)|^2 ds \right)^{p/2} \right]^{2/p} &\leq e^{-2\beta(t-t_n)/p} \int_{t_n}^t \|X(s)\|_{L^p(\Omega)}^2 ds \\
&\leq \|X\|_{\beta,p,t}^2 \int_{t_n}^t e^{-2\beta(t-s)/p} ds \\
&= \|X\|_{\beta,p,t}^2 \frac{p}{2\beta} \left(1 - e^{-2\beta(t-t_n)/p} \right),
\end{aligned}$$

where we have used that

$$e^{-\beta(t-t_n)/p} \|X(s)\|_{L^p(\Omega)} = \left(e^{-\beta(s-t_n)} E[|X(s)|^p] \right)^{1/p} e^{-\beta(t-s)/p}.$$

It follows that

$$T(t) \leq e^{-\beta(t-t_n)/p} \sqrt{t-t_n} E[|Y|^p]^{1/p} + K_2 \|X\|_{\beta,p,t} \sqrt{\frac{p}{2\beta}} \quad (6.9)$$

Apply estimate (6.9) in (6.8) and take the supremum over $t_n \leq t \leq \tau$. It follows that

$$\|X\|_{\beta,p,\tau} \leq c_p^{1/p} K_2 \sqrt{\frac{p}{2\beta}} \|X\|_{\beta,p,\tau} + c_p^{1/p} \sqrt{\tau-t_n} E[|Y|^p]^{1/p} + E[|X(t_n)|^p]^{1/p}.$$

Choosing β sufficiently large it follows that $c_p^{1/p} K_2 \sqrt{p/2\beta} < 1$. Hence

$$\|X\|_{\beta,p,\tau} \leq \frac{1}{1 - c_p^{1/p} K_2 \sqrt{p/2\beta}} \left(c_p^{1/p} \sqrt{\tau-t_n} E[|Y|^p]^{1/p} + E[|X(t_n)|^p]^{1/p} \right).$$

The result follows upon multiplication by $e^{\beta(\tau-t_n)/p}$ as

$$e^{\beta(\tau-t_n)/p} \|X\|_{\beta,p,\tau} = \left(\sup_{t_n \leq t \leq \tau} e^{\beta(\tau-t)} E[|X(s)|^p] \right)^{1/p} \geq \sup_{t_n \leq t \leq \tau} E[|X(t)|^p]^{1/p}.$$

□

Estimate 6.8. Let $\mathcal{T}_6$ be defined in Lemma 6.2. Then

$$\mathcal{T}_6 = \mathcal{O}(\varepsilon).$$

Proof. This follows as in [11, Limit 6].

□

Estimate 6.9. Let $\mathcal{T}_6$ be defined in Lemma 6.2. Then

$$|\mathcal{T}_6| \leq \sum_{n=0}^{N-1} E \left[\|\mathcal{S}_{CL}(\Delta t) u^n - v^n\|_{1,\phi \star J_r} \right].$$

Proof. Recall that

$$\begin{aligned}
\mathcal{T}_6 = - \sum_{n=0}^{N-1} E \left[\iint_{\Pi_T} \int_{\mathbb{R}^d} \left(S_\delta(\tilde{\eta}(t_{n+1}, x) - u^\varepsilon(s, y)) \right. \right. \\
\left. \left. - S_\delta(\tilde{\eta}((t_{n+1})-, x) - u^\varepsilon(s, y)) \right) \varphi(t_{n+1}, x, s, y) dx ds dy \right].
\end{aligned}$$

First, we note that $|S_\delta(b) - S_\delta(a)| \leq |b - a|$. This and (4.9) yields

$$|\mathcal{T}_6| \leq \sum_{n=0}^{N-1} E \left[\|\tilde{\eta}(t_{n+1}) - \tilde{\eta}((t_{n+1})-)\|_{1,\phi \star J_r} \right].$$

Now,

$$\tilde{\eta}(t_{n+1}) - \tilde{\eta}((t_{n+1})-) = \mathcal{S}_{\text{SDE}}(t_{n+1}, t_n)(\mathcal{S}_{\text{CL}}(\Delta t)u^n - v^n) + \mathcal{S}_{\text{CL}}(\Delta t)u^n - v^n.$$

The result follows by (3.2). $\square$

Proof of Theorem 6.1. Consider Lemma 6.2. Send $r_0 \downarrow 0, \varepsilon \downarrow 0$ and $\gamma \downarrow 0$ in that order. Next we let $v^n \rightarrow \mathcal{S}_{\text{CL}}(\Delta t)u^n$ in $L^1(\Omega, L^1(\mathbb{R}^d, \phi))$. Due to the L^1 -Lipschitz continuity of $\mathcal{S}_{\text{CL}}$ in time it follows by Estimate 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8 and 6.9 that

$$\begin{aligned} E \left[\|u_0 - u^0\|_{1,\phi} \right] + C_\phi \|f\|_{\text{Lip}} \int_0^{t_0} E \left[\|\eta_{\Delta t}(t) - u(t)\|_{1,\phi} \right] dt \\ + \mathcal{O} \left(\delta + r + \sqrt{\Delta t} + \frac{\delta}{r} + \frac{\Delta t}{\delta} \right) \geq E \left[\|\eta_{\Delta t}(t_0) - u(t_0)\|_{1,\phi} \right]. \end{aligned}$$

Next, let $u^0 = u_0$, apply Grönwall's inequality and let $\delta = \Delta t^{2/3}$ and $r = \Delta t^{1/3}$. This proves Theorem 6.1. $\square$

7. APPENDIX

7.1. Proof of Proposition 6.1. The proof of Proposition 6.1 is based on the following Lemma:

Lemma 7.1. *Suppose $u, v \in L^2(\Omega, P, \mathcal{F}_{t_n}; L^2(\mathbb{R}^d))$, v smooth. Let*

$$\psi(t) = (\mathcal{S}_{\text{SDE}}(t, t_n) - \mathcal{I})v \quad \text{and} \quad w(t) = \mathcal{S}_{\text{CL}}(t - t_n)u,$$

for $t \in [t_n, t_{n+1}]$. Then, for any $(S, Q) \in \mathcal{E}$, $\varphi \in C_c^\infty(\Pi_n^2)$ and any $V \in \mathbb{D}^{1,2}$

$$R - L + \mathcal{T}_1 + \mathcal{T}_2 - \mathcal{T}_3 + \mathcal{T}_4 \geq 0$$

where

$$\begin{aligned} L &= E \left[\iint_{\Pi_n} \int_{\mathbb{R}^d} S(w(t_{n+1}, x) + \psi(s, y) - V) \varphi(t_{n+1}, x, s, y) dx dy ds \right] \\ &\quad + E \left[\iint_{\Pi_n} \int_{\mathbb{R}^d} S(w(t, x) + \psi(t_{n+1}, y) - V) \varphi(t, x, t_{n+1}, y) dy dx dt \right], \\ R &= E \left[\iint_{\Pi_n} \int_{\mathbb{R}^d} S(w(t_n, x) + \psi(s, y) - V) \varphi(t_n, x, s, y) dx dy ds \right] \\ &\quad + E \left[\iint_{\Pi_n} \int_{\mathbb{R}^d} S(w(t, x) + \psi(t_n, y) - V) \varphi(t, x, t_n, y) dy dx dt \right], \\ \mathcal{T}_1 &= E \left[\iiint_{\Pi_n^2} S(w(t, x) + \psi(s, y) - V) (\partial_t + \partial_s) \varphi dX \right], \\ \mathcal{T}_2 &= E \left[\iiint_{\Pi_n^2} Q(w(t, x), V - \psi(s, y)) \cdot \nabla_x \varphi dX \right], \\ \mathcal{T}_3 &= E \left[\iiint_{\Pi_n^2} S''(w(t, x) + \psi(s, y) - V) D_s V \sigma(\psi(s, y) + v(y)) \varphi dX \right], \\ \mathcal{T}_4 &= \frac{1}{2} E \left[\iiint_{\Pi_n^2} S''(w(t, x) + \psi(s, y) - V) \sigma^2(\psi(s, y) + v(y)) \varphi dX \right], \end{aligned}$$

where $\Pi_n = [t_n, t_{n+1}] \times \mathbb{R}^d$.

Proof of Lemma 7.1. The entropy inequality yields

$$\begin{aligned} \int_{\mathbb{R}^d} S(w(t_n, x) - c) \varphi(t_n, x, s, y) - S(w(t_{n+1}, x) - c) \varphi(t_{n+1}, x, s, y) dx \\ + \iint_{\Pi_n} S(w - c) \partial_t \varphi + Q(w, c) \cdot \nabla_x \varphi dt dx \geq 0, \end{aligned} \quad (7.1)$$

for all $c \in \mathbb{R}$ and all $s, y \in \Pi_n$. Let $c = V - \psi(s, y)$ in (7.1), integrate in s and y and take expectation to obtain

$$\begin{aligned} E \left[\iint_{\Pi_n} \int_{\mathbb{R}^d} S(w(t_n, x) + \psi(s, y) - V) \varphi(t_n, x, s, y) dx ds dy \right] \\ - E \left[\iint_{\Pi_n} \int_{\mathbb{R}^d} S(w(t_{n+1}, x) + \psi(s, y) - V) \varphi(t_{n+1}, x, s, y) dx ds dy \right] \\ + E \left[\iiint_{\Pi_n^2} S(w + \psi - V) \partial_t \varphi + Q(w, V - \psi) \cdot \nabla_x \varphi dX \right] \geq 0. \end{aligned} \quad (7.2)$$

Note that $w(t)$ is $\mathcal{F}_{t_n}$ adapted for all $t \in [t_n, t_{n+1}]$. To see what equation ψ satisfies, let $\zeta(t) = \mathcal{S}_{\text{SDE}}(t, t_n)v$. By definition

$$\zeta(t) = v + \int_{t_n}^t \sigma(\zeta(r)) dB(r).$$

Since $\psi(t) = \zeta(t) - v$ it follows that

$$\psi(t) = \int_{t_n}^t \sigma(\psi(r) + v) dB(r), \quad (t \in [t_n, t_{n+1}]) \quad (7.3)$$

Fix $t, x \in \Pi_n$ and $y \in \mathbb{R}$ and let

$$X(s) := w(t, x) + \psi(s, y) - V \text{ and } Y(s) := S(X(s)) \varphi(t, x, s, y), \quad s \in [t_n, t_{n+1}].$$

By (7.3)

$$X(s) = w(t, x) - V + \int_{t_n}^s \sigma(\psi(r, y) + v(y)) dB(r).$$

By [21, Theorem 3.2.2]

$$\begin{aligned} Y(t_{n+1}) &= Y(t_n) \\ &+ \int_{t_n}^{t_{n+1}} S(X(s)) \partial_s \varphi ds \\ &+ \int_{t_n}^{t_{n+1}} S'(X(s)) \sigma(\psi(s) + v) \varphi dB(s) \\ &+ \frac{1}{2} \int_{t_n}^{t_{n+1}} S''(X(s)) \sigma^2(\psi(s) + v) \varphi ds \\ &- \int_{t_n}^{t_{n+1}} S''(X(s)) D_s V \sigma(\psi(s) + v) \varphi ds. \end{aligned}$$

where the stochastic integral is interpreted as a Skorohod integral. Upon integrating in t, x, y and taking expectation

$$\begin{aligned}
 & E \left[\iint_{\Pi_n} \int_{\mathbb{R}^d} S(w(t, x) + \psi(t_n, y) - V) \varphi(t, x, t_n, y) dy dt dx \right] \\
 & - E \left[\iint_{\Pi_n} \int_{\mathbb{R}^d} S(w(t, x) + \psi(t_{n+1}, y) - V) \varphi(t, x, t_{n+1}, y) dy dt dx \right] \\
 & + E \left[\iiint_{\Pi_n^2} S(w(t, x) + \psi(s, y) - V) \partial_s \varphi dX \right] \\
 & + \frac{1}{2} E \left[\iiint_{\Pi_n^2} S''(w(t, x) + \psi(s, y) - V) (\sigma(\psi(s, y) + v(y)))^2 \varphi dX \right] \\
 & - E \left[\iiint_{\Pi_n^2} S''(w(t, x) + \psi(s, y) - V) D_s V \sigma(\psi(s, y) + v(y)) \varphi dX \right] = 0.
 \end{aligned} \tag{7.4}$$

The result follows by adding (7.2) and (7.4). $\square$

Proof of Proposition 6.1. Consider Lemma 7.1. Let

$$\varphi(t, x, s, y) := \frac{1}{2^d} \phi \left(\frac{t+s}{2}, \frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) J_{r_0}(t-s). \tag{7.5}$$

Then send r_0, r to zero in that order. The result follows by Limit 1,2,3,4 and 5. $\square$

Limit 1. Let L and R be the terms defined in Lemma 7.1 and φ be defined by (7.5). Then

$$\begin{aligned}
 \lim_{r, r_0 \downarrow 0} L(r, r_0) &= E \left[\int_{\mathbb{R}^d} S(w(t_{n+1}, x) + \psi(t_{n+1}, x) - V) \phi(t_{n+1}, x) dx \right], \\
 \lim_{r, r_0 \downarrow 0} R(r, r_0) &= E \left[\int_{\mathbb{R}^d} S(w(t_n, x) + \psi(t_n, x) - V) \phi(t_n, x) dx \right].
 \end{aligned}$$

Proof. Consider

$$L_1 = E \left[\iint_{\Pi_n} \int_{\mathbb{R}^d} S(w(t_{n+1}, x) + \psi(s, y) - V) \varphi(t_{n+1}, x, s, y) dx dy ds \right].$$

The same reasoning holds true for the remaining terms. As a consequence of the dominated convergence theorem and Lemma 7.4 we may conclude that

$$\begin{aligned}
 \lim_{r_0 \downarrow 0} L_1 &= \frac{1}{2} E \left[\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} S(w(t_{n+1}, x) + \psi(t_{n+1}, y) - V) \right. \\
 & \quad \left. \times \frac{1}{2^d} \phi \left(t_{n+1}, \frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) dx dy \right].
 \end{aligned}$$

Note that continuity yields the fact that it holds for all t_{n+1} . Next, we apply Lemma 7.4 to conclude that

$$\lim_{r, r_0 \downarrow 0} L_1 = \frac{1}{2} E \left[\int_{\mathbb{R}^d} S(w(t_{n+1}, x) + \psi(t_{n+1}, x) - V) \phi(t_{n+1}, x) dx \right].$$

$\square$

Limit 2. Let $\mathcal{T}_1$ be defined in Lemma 7.1 and φ in (7.5). Then

$$\lim_{r, r_0 \downarrow 0} \mathcal{T}_1 = E \left[\iint_{\Pi_n} S(u(t, x) - V) \partial_t \phi(t, x) dx dt \right].$$

Proof. Recall that

$$\mathcal{T}_1 = E \left[\iiint_{\Pi_n^2} S(w(t, x) + \psi(s, y) - V) (\partial_t + \partial_s) \varphi \, dX \right].$$

Observe that

$$(\partial_t + \partial_s) \varphi(t, x, s, y) = \frac{1}{2^d} \partial_1 \phi \left(\frac{t+s}{2}, \frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) J_{r_0}(t-s).$$

The result follows by the Dominated Convergence Theorem and Lemma 7.4 as in the proof of Limit 1. $\square$

Limit 3. Let $\mathcal{T}_2$ be defined in Lemma 7.1 and φ in (7.5). Then

$$\begin{aligned} \lim_{r, r_0 \downarrow 0} \mathcal{T}_2 &= E \left[\iint_{\Pi_n} Q(w + \psi, V) \cdot \nabla \phi \, dx dt \right] \\ &\quad + E \left[\iint_{\Pi_n} \left(\int_V^{w+\psi} S'(z - V) (f'(z - \psi) - f'(z)) \, dz \right) \cdot \nabla \phi \, dx dt \right] \\ &\quad + E \left[\iint_{\Pi_n} \left(\int_V^{w+\psi} S''(z - V) f'(z - \psi) \, dz \right) \cdot \nabla \psi \phi \, dx dt \right], \end{aligned}$$

where $w = w(t, x)$, $\psi = \psi(t, x)$ and $\phi = \phi(t, x)$.

Proof. Recall that

$$\mathcal{T}_2 = E \left[\iiint_{\Pi_n^2} Q(w(t, x), V - \psi(s, y)) \cdot \nabla_x \varphi \, dX \right].$$

First observe that

$$\begin{aligned} \nabla_x \varphi(t, x, s, y) &= \frac{1}{2^{d+1}} \partial_2 \phi \left(\frac{t+s}{2}, \frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) J_{r_0}(t-s) \\ &\quad - \frac{1}{2^d} \phi \left(\frac{t+s}{2}, \frac{x+y}{2} \right) \nabla_y J_r \left(\frac{x-y}{2} \right) J_{r_0}(t-s). \end{aligned}$$

Using integration by parts it follows that

$$\begin{aligned} &E \left[\iiint_{\Pi_n^2} Q(w(t, x), V - \psi(s, y)) \nabla_x \varphi \, dx dt dy ds \right] \\ &= E \left[\iiint_{\Pi_n^2} Q(w(t, x), V - \psi(s, y)) \right. \\ &\quad \cdot \left. \frac{1}{2^d} \partial_2 \phi \left(\frac{t+s}{2}, \frac{x+y}{2} \right) J_r \left(\frac{x-y}{2} \right) J_{r_0}(t-s) \, dx dt dy ds \right] \\ &\quad + E \left[\iiint_{\Pi_n^2} \nabla_y \cdot Q(w(t, x), V - \psi(s, y)) \varphi(t, x, s, y) \, dx dt dy ds \right] \\ &=: \mathcal{T}_2^1 + \mathcal{T}_2^2. \end{aligned}$$

As above

$$\lim_{r, r_0 \downarrow 0} \mathcal{T}_2^1 = E \left[\iint_{\Pi_n} Q(w(t, x), V - \psi(t, x)) \cdot \nabla \phi(t, x) \, dx dt \right].$$

Finally we apply the identity

$$Q(w, V - \psi) = Q(w + \psi, V) + \int_V^{w+\psi} S'(z - V) (f'(z - \psi) - f'(z)) \, dz.$$

Consider $\mathcal{T}_2^2$. By the chain rule it follows that

$$\mathcal{T}_2^2 = -E \left[\iiint_{\Pi_n^2} \partial_2 Q(w(t, x), V - \psi(s, y)) \cdot \nabla_y \psi(s, y) \varphi(t, x, s, y) dx dt dy ds \right].$$

Letting $r_0 \downarrow 0$ and $r \downarrow 0$ we arrive at

$$\lim_{r, r_0 \downarrow 0} \mathcal{T}_2^2 = -E \left[\iint_{\Pi_n} \partial_2 Q(w(t, x), V - \psi(t, x)) \cdot \nabla_x \psi(t, x) \phi(t, x) dx dt \right].$$

Finally, note that

$$\begin{aligned} \partial_2 Q(w, V - \psi) &= - \int_{V-\psi}^w S''(z - V + \psi) f'(z) dz \\ &= - \int_V^{w+\psi} S''(z - V) f'(z - \psi) dz. \end{aligned}$$

□

Limit 4. Let $\mathcal{T}_3$ be defined in Lemma 7.1 and φ by (7.5). Then

$$\lim_{r, r_0 \downarrow 0} \mathcal{T}_3 = E \left[\iint_{\Pi_n} S''(w(t, x) + \psi(t, x) - V) D_t V \sigma(\psi(t, x) + v(x)) \phi(t, x) dx dt \right].$$

Proof. Recall that

$$\mathcal{T}_3 = E \left[\iiint_{\Pi_n^2} S''(w(t, x) + \psi(s, y) - V) D_s V \sigma(\psi(s, y) + v(y)) \varphi dX \right].$$

It is straight forward to apply the dominated convergence theroem and Lemma 7.4 to this term. □

Limit 5. Let $\mathcal{T}_4$ be defined in Lemma 7.1 and φ by (7.5). Then

$$\lim_{r, r_0 \downarrow 0} \mathcal{T}_4 = \frac{1}{2} E \left[\iint_{\Pi_n} S''(w(t, x) + \psi(t, x) - V) \sigma^2(\psi(t, x) + v(x)) \phi(t, x) dx dt \right].$$

Proof. The proof is similar to the above. □

7.2. Weighted L^p spaces. Here we state some elementary results regarding functions in $\mathfrak{N}$. For proofs, see [11].

Lemma 7.2. Let $\phi \in \mathfrak{N}$ and $0 < p < \infty$. Then

$$\left| \phi^{1/p}(x+z) - \phi^{1/p}(x) \right| \leq w_{p,\phi}(|z|) \phi^{1/p}(x),$$

where

$$w_{p,\phi}(r) = \frac{C_\phi}{p} r \left(1 + \frac{C_\phi}{p} r e^{C_\phi r/p} \right)$$

defined for all $r \geq 0$. As a consequence it follows that if $\phi(x_0) = 0$ for some $x_0 \in \mathbb{R}^d$, then $\phi = 0$ (and by definition $\phi \notin \mathfrak{N}$).

Lemma 7.3. Let $\phi \in \mathfrak{N}$. Then

- (i) If $\varphi \in L^1(\mathbb{R}^d)$ is nonnegative, then $\phi \star \varphi \in \mathfrak{N}$ with $C_{\phi \star \varphi} = C_\phi$.
- (ii) Let $\varphi \in L^1(\mathbb{R}^d)$ have support in $B(0, 1)$ and $\|\varphi\|_{L^1(\mathbb{R}^d)} = 1$. Take $\varphi_\delta(z) = \frac{1}{\delta^d} \varphi(\frac{z}{\delta})$. Then, for any $u \in L^p(\mathbb{R}^d, \phi)$

$$\left| \|u\|_{p,\phi}^p - \|u\|_{p,\phi \star \varphi_\delta}^p \right| \leq w_{1,\phi}(\delta) \|u\|_{p,\phi}^p.$$

7.3. A “Doubling of variables” tool. The following result follows along the lines of [23, Lemma 2.7.2]. See also [11, § 6].

Lemma 7.4. *Suppose $u, v \in L^1_{\text{loc}}(\mathbb{R}^d)$ and F is Lipschitz on $\mathbb{R}^2$. Fix $\psi \in C_c(\mathbb{R}^d)$ and let*

$$\begin{aligned} \mathcal{T}_r := & \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} F(u(x), v(y)) \frac{1}{2^d} \psi\left(\frac{x+y}{2}\right) J_r\left(\frac{x-y}{2}\right) dy dx \\ & - \int_{\mathbb{R}^d} F(u(x), v(x)) \psi(x) dx, \end{aligned}$$

where J_r is defined in (2.1). Then $\mathcal{T}_r \rightarrow 0$ as $r \downarrow 0$.

Similarly, let $G : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be measurable in the first variable and Lipschitz continuous in the second variable. Let $w \in L^1([0, T])$. Set

$$\mathcal{T}_{r_0}(s) = \int_0^T |G(s, w(t)) - G(s, w(s))| J_{r_0}(t-s) dt.$$

Then $\mathcal{T}_{r_0}(s) \rightarrow 0$ Lebesgue-a.e. as $r_0 \downarrow 0$. The result does not depend on the symmetry of J .

7.4. Young measures. The purpose of this section is to provide a reference for some results concerning Young measures and their applications as generalized limits. Let $(X, \mathcal{A}, \mu)$ be a σ -finite measure space, and $\mathcal{P}(\mathbb{R})$ denote the set of probability measures on $\mathbb{R}$. In the previous sections X is typically $\Pi_T \times \Omega$. A *Young measure* from X into $\mathbb{R}$ is a function $\nu : X \rightarrow \mathcal{P}(\mathbb{R})$ such that $x \mapsto \nu_x(B)$ is $\mathcal{A}$ -measurable for every Borel measurable set $B \subset \mathbb{R}$. We denote by $\mathcal{Y}(X, \mathcal{A}, \mu; \mathbb{R})$, or simply $\mathcal{Y}(X; \mathbb{R})$ if the measure space is understood, the set of all Young measures from X into $\mathbb{R}$. The following theorem is proved in [22, Theorem 6.2] in the case that $X \subset \mathbb{R}^n$ and μ is the Lebesgue measure:

Theorem 7.1. *Let $(X, \mathcal{A}, \mu)$ be a σ -finite measure space. Let $\zeta : [0, \infty) \rightarrow [0, \infty]$ be a continuous, non decreasing function satisfying $\lim_{\xi \rightarrow \infty} \zeta(\xi) = \infty$ and $\{u^n\}_{n \geq 1}$ a sequence of measurable functions such that*

$$\sup_n \int_X \zeta(|u^n|) d\mu(x) < \infty.$$

Then there exist a subsequence $\{u^{n_j}\}_{j \geq 1}$ and a $\nu \in \mathcal{Y}(X, \mathcal{A}, \mu; \mathbb{R})$ such that for any Carathéodory function $\psi : \mathbb{R} \times X \rightarrow \mathbb{R}$ with $\psi(u^{n_j}(\cdot), \cdot) \rightharpoonup \bar{\psi}$ in $L^1(X)$ we have

$$\bar{\psi}(x) = \int_{\mathbb{R}} \psi(\xi, x) d\nu_x(\xi).$$

The proof is based on the embedding of $\mathcal{Y}(X; \mathbb{R})$ into $L^\infty_{w*}(X, \mathcal{M}(\mathbb{R}))$. Here $\mathcal{M}(\mathbb{R})$ denotes the space of Radon measures on $\mathbb{R}$. The crucial observation is that $(L^1(X, C_0(\mathbb{R})))^*$ is isometrically isomorphic to $L^\infty_{w*}(X, \mathcal{M}(\mathbb{R}))$ also in the case that $(X, \mathcal{A}, \mu)$ is an abstract σ -finite measure space. It is relatively straight forward to go through the proof and extend to the more general case [19, Theorem 2.11]. Note however that due to the application of the weighted L^p spaces it suffices, in our case, with the version for finite measure spaces.

7.5. Weak compactness in L^1 . To apply Theorem 7.1 it is necessary to determine whether or not $\{\psi(\cdot, u^n(\cdot))\}_{n \geq 1}$ has a weakly convergent subsequence in $L^1(X)$. The key result is known as the Dunford-Pettis Theorem.

Definition 7.1. Let $\mathcal{K} \subset L^1(X, \mathcal{A}, \mu)$.

(i) $\mathcal{K}$ is *uniformly integrable* if for any $\varepsilon > 0$ there exists a $c_0(\varepsilon)$ such that

$$\sup_{f \in \mathcal{K}} \int_{|f| \geq c} |f| d\mu \leq \varepsilon \text{ whenever } c \geq c_0(\varepsilon).$$

- (ii) $\mathcal{K}$ has *uniform tail* if for any $\varepsilon > 0$ there exists a $E \in \mathcal{A}$ with $\mu(E) < \infty$ such that

$$\sup_{f \in \mathcal{K}} \int_{X \setminus E} |f| d\mu \leq \varepsilon.$$

If $\mathcal{K}$ satisfies both (i) and (ii) it is said to be *equiintegrable*.

Remark 7.5. Note that (ii) is void when μ is finite.

Theorem 7.2 (Dunford-Pettis). *Let $(X, \mathcal{A}, \mu)$ be a σ -finite measure space. A subset $\mathcal{K}$ of $L^1(X)$ is relatively weakly sequentially compact if and only if it is equiintegrable.*

Recall that by the Eberlein-Šmulian Theorem, in the weak topology of a Banach space, relative weak compactness is equivalent with relative sequentially weak compactness [26]. There are a couple of well known reformulations of uniform integrability.

Lemma 7.6. *Suppose $\mathcal{K} \subset L^1(X)$ is bounded. Then $\mathcal{K}$ is uniformly integrable if and only if:*

- (i) *For any $\varepsilon > 0$ there exists a $\delta(\varepsilon) > 0$ such that*

$$\sup_{f \in \mathcal{K}} \int_E |f| d\mu \leq \varepsilon \text{ whenever } \mu(E) \leq \delta(\varepsilon).$$

- (ii) *There exists an increasing function $\Psi : [0, \infty) \rightarrow [0, \infty)$ such that $\Psi(\zeta)/\zeta \rightarrow \infty$ as $\zeta \rightarrow \infty$ satisfying*

$$\sup_{f \in \mathcal{K}} \int_X \Psi(|f(x)|) d\mu(x) < \infty.$$

REFERENCES

- [1] C. Bauzet. Time-splitting approximation of the cauchy problem for a stochastic conservation law. *Mathematics and Computers in Simulation*, 118:73–86, 2015.
- [2] C. Bauzet, J. Charrier, and T. Gallouët. Convergence of flux-splitting finite volume schemes for hyperbolic scalar conservation laws with a multiplicative stochastic perturbation. Preprint, 2014.
- [3] C. Bauzet, G. Vallet, and P. Wittbold. The Cauchy problem for conservation laws with a multiplicative stochastic perturbation. *J. Hyperbolic Differ. Equ.*, 9(4):661–709, 2012.
- [4] I. H. Biswas, K. H. Karlsen, and A. K. Majee. Conservation laws driven by Lévy white noise. *J. Hyperbolic Differ. Equ.*, 12(3):581–654, 2015.
- [5] G.-Q. Chen, Q. Ding, and K. H. Karlsen. On nonlinear stochastic balance laws. *Arch. Ration. Mech. Anal.*, 204(3):707–743, 2012.
- [6] K. L. Chung and R. J. Williams. *Introduction to stochastic integration*. Modern Birkhäuser Classics. Birkhäuser/Springer, New York, second edition, 2014.
- [7] A. Debussche and J. Vovelle. Scalar conservation laws with stochastic forcing. *J. Funct. Anal.*, 259(4):1014–1042, 2010.
- [8] J. Feng and D. Nualart. Stochastic scalar conservation laws. *J. Funct. Anal.*, 255(2):313–373, 2008.
- [9] H. Holden and N. H. Risebro. Conservation laws with random source. *Appl. Math. Optim.*, 36(2) (1997) 229–241.
- [10] H. Holden, K. H. Karlsen, K.-A. Lie, and N. H. Risebro. *Splitting Methods for Partial Differential Equations with Rough Solutions: Analysis and MATLAB programs*. EMS Series of Lectures in Mathematics. European Mathematical Society (EMS), Zürich, 2010.
- [11] K. H. Karlsen and E. B. Storrøsten. On stochastic conservation laws and Malliavin calculus. Preprint, [arXiv:1507.05518](https://arxiv.org/abs/1507.05518) [math.AP].
- [12] J. U. Kim. On a stochastic scalar conservation law. *Indiana Univ. Math. J.* 52 (1) (2003) 227–256.
- [13] I. Kröker and C. Rohde. Finite volume schemes for hyperbolic balance laws with multiplicative noise. *Appl. Numer. Math.*, 62(4):441–456, 2012.
- [14] S. N. Kružkov. Results on the nature of the continuity of solutions of parabolic equations, and certain applications thereof. *Mat. Zametki*, 6:97–108, 1969.

- [15] S. N. Kružkov. First order quasilinear equations with several independent variables. *Mat. Sb. (N.S.)*, 81 (123):228–255, 1970.
- [16] J. O. Langseth, A. Tveito, and R. Winther. On the convergence of operator splitting applied to conservation laws with source terms. *SIAM Journal on Numerical Analysis*, 33(3):pp. 843–863, 1996.
- [17] P.-L. Lions, B. Perthame, and P. Souganidis. Scalar conservation laws with rough (stochastic) fluxes. *Stoch. Partial Differ. Equ. Anal. Comput.*, 1(4):664–686, 2013.
- [18] P.-L. Lions, B. Perthame, and P. E. Souganidis. Scalar conservation laws with rough (stochastic) fluxes: the spatially dependent case. *Stoch. Partial Differ. Equ. Anal. Comput.*, 2(4):517–538, 2014.
- [19] J. Málek, J. Nečas, M. Rokyta, and M. Růžička. *Weak and measure-valued solutions to evolutionary PDEs*, volume 13 of *Applied Mathematics and Mathematical Computation*. Chapman & Hall, London, 1996.
- [20] S. Mishra and Ch. Schwab. Sparse tensor multi-level Monte Carlo finite volume methods for hyperbolic conservation laws with random initial data. *Math. Comp.*, 81(280):1979–2018, 2012.
- [21] David Nualart. *The Malliavin calculus and related topics*. Probability and its Applications (New York). Springer-Verlag, Berlin, second edition, 2006.
- [22] Pablo Pedregal. *Parametrized measures and variational principles*. Progress in Nonlinear Differential Equations and their Applications, 30. Birkhäuser Verlag, Basel, 1997.
- [23] Denis Serre. *Systems of conservation laws. 1*. Cambridge University Press, Cambridge, 1999. Hyperbolicity, entropies, shock waves, Translated from the 1996 French original by I. N. Sneddon.
- [24] G. Vallet. Dirichlet problem for a nonlinear conservation law. *Rev. Mat. Complut.*, 13(1):231–250, 2000.
- [25] G. Vallet and P. Wittbold. On a stochastic first-order hyperbolic equation in a bounded domain. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 12(4):613–651, 2009.
- [26] Robert Whitley. An elementary proof of the Eberlein-Šmulian theorem. *Math. Ann.*, 172:116–118, 1967.

(Kenneth H. Karlsen)

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF OSLO
P.O. BOX 1053, BLINDERN
N-0316 OSLO, NORWAY
E-mail address: kennethk@math.uio.no

(Erlend Briseid Storrøsten)

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF OSLO
P.O. BOX 1053, BLINDERN
N-0316 OSLO, NORWAY
E-mail address: erlenbs@math.uio.no

Appendix

A.1 A note on the representation of nonlinear limits

Suppose $\{u^n\}_{n \geq 1}$ is a sequence of real valued functions on a measure space $(X, \mathcal{A}, \mu)$. We seek to answer the following question:

Question 1. Suppose there exists some class of nonlinear functions $\psi : X \times \mathbb{R} \rightarrow \mathbb{R}$ such that $\psi(\cdot, u^n) \rightharpoonup \bar{\psi}$. Under which assumptions does there exist some limiting object ν , depending on $\{u^n\}_{n \geq 1}$ only, and a mapping $\mathcal{F}$ such that $\bar{\psi} = \mathcal{F}(\psi, \nu)$.

Assuming that $\{u^n\}_{n \geq 1}$ is compact in $L^1(X)$, we could subtract a strongly convergent subsequence $\{u^{n_j}\}_{j \geq 1}$ and a limit $u \in L^1(X)$ such that $u^{n_j} \rightarrow u$ in $L^1(X)$. In this case, it is in general true that $\bar{\psi} = \psi(u)$. To be more specific, suppose $\psi(x, u^n) \rightharpoonup \bar{\psi}$ in $L^1(X)$. As $u^{n_j} \rightarrow u$ strongly in $L^1(X)$ there exists a subsequence, not relabeled, such that $u^{n_j} \rightarrow u$ μ -almost everywhere. For any $\varphi \in L^\infty(X)$,

$$\int_X \bar{\psi} \varphi d\mu = \lim_{j \rightarrow \infty} \int_X \psi(x, u^{n_j}(x)) \varphi d\mu = \int_X \psi(x, u(x)) \varphi d\mu,$$

given that we may apply the dominated convergence theorem. In fact, by Vitali's convergence theorem it suffices that $\{\psi(\cdot, u^{n_j})\}_{j \geq 1}$ is uniformly integrable for the second equality to hold. It follows that $\bar{\psi} = \psi(\cdot, u)$. Often, strong compactness of $\{u^n\}_{n \geq 1}$ is not available. As a remedy one might want to try weak convergence instead. However, weak convergence of a subsequence $u^{n_j} \rightharpoonup u$ is not sufficient to conclude that $\bar{\psi} = \psi(u)$. As a counterexample, take $X = [0, 2\pi]$ and $u^n(x) = \sin(nx)$. Then $\sin(nx) \rightharpoonup 0$ in $L^1([0, 2\pi])$, but $\int_0^{2\pi} |\sin(nx)| dx = 4$ implying that $|\sin(nx)| \not\rightarrow 0$. So how do we handle Question 1 when strong compactness is unavailable? Let me sketch two slightly different approaches. Assume that $\psi(x, \cdot) \in C_0^1(\mathbb{R})$. Then

$$\psi(x, u^n(x)) = - \int_{\mathbb{R}} \partial_\xi \psi(x, \xi) \underbrace{\mathbb{1}_{[u^n(x), \infty)}(\xi)}_{f^n(x, \xi)} d\xi.$$

By weak* compactness of $\{f^n\}_{n \geq 1}$ in $L^\infty(X \times \mathbb{R})$ we may subtract a weakly convergent subsequence $f^{n_j} \rightharpoonup f$. A possible solution is given by

$$\bar{\psi}(x) = - \int_{\mathbb{R}} \partial_\xi \psi(x, \xi) f(x, \xi) d\xi = \int_{\mathbb{R}} \psi(x, \xi) \partial_\xi f(x, \xi) d\xi,$$

provided the limit f is differentiable in some sense. A second representation is given by

$$\psi(x, u^n(x)) = \int_{\mathbb{R}} \psi(x, \xi) \underbrace{\delta(\xi - u^n(x))}_{\nu_x^n(\xi)} d\xi.$$

Suppose that we may establish compactness of $\{\nu^n\}_{n \geq 1}$ so that we may subtract a weakly convergent subsequence $\nu^{n_j} \rightharpoonup \nu$. This yields the presentation

$$\bar{\psi}(x) = \int_{\mathbb{R}} \psi(x, \xi) \nu_x(\xi) d\xi.$$

Of course, if this is to make any sense $\nu = \partial_\xi f$. In these notes both these approaches are considered, and both are shown to yield Theorem 1. Since the first approach outlined above is associated with the kinetic formulation of scalar conservation laws, we refer to this as the kinetic approach. The second will be referred to as the canonical approach, as this is the more standard functional analytic one. As the applicability of Theorem 1 is depending on the weak compactness of $\{\psi(\cdot, u^n)\}_{n \geq 1}$ in $L^1(X)$, a section on this topic is also included.

The kinetic approach

Let $(X, \mathcal{A}, \mu)$ be a σ -finite measure space. Let $\mathcal{P}(\mathbb{R})$ denote the set of Borel regular probability measures on $\mathbb{R}$. A *Young measure* from X into $\mathbb{R}$ is a function $\nu : X \rightarrow \mathcal{P}(\mathbb{R})$ such that $x \mapsto \nu(x)(B)$ is $\mathcal{A}$ -measurable for every Borel measurable set $B \subset \mathbb{R}$. We denote by $\mathcal{Y}(X, \mathcal{A}, \mu; \mathbb{R})$ or $\mathcal{Y}(X; \mathbb{R})$ the set of all Young measures from X into $\mathbb{R}$. Let $\nu \in \mathcal{Y}(X, \mathcal{A}, \mu; \mathbb{R})$ and let $f(x, \xi) := \nu_x((-\infty, \xi])$. Then one may check that f is $\mathcal{A} \otimes \mathcal{B}(\mathbb{R})$ measurable and for any $\varphi \in L^1(X, C(\mathbb{R})) \cap L^1(X, W^{1,1}(\mathbb{R}))$

$$\int_X \int_{\mathbb{R}} f(x, \xi) \partial_\xi \varphi(x, \xi) d\xi d\mu(x) = - \int_X \int_{\mathbb{R}} \varphi(x, \xi) d\nu_x(\xi) d\mu(x).$$

What we need is rather a converse. That is, we need a characterization of the functions f for which there exists a Young measure ν such that $f(x, \xi) := \nu_x((-\infty, \xi])$. At the heart of this result is of course the Riesz-Markov Theorem associating any linear functional on $C_c(\mathbb{R})$ to a Radon measure on $\mathbb{R}$. Recall that a Borel regular measure is Radon if it is finite on compact sets.

Lemma 1. *Let $f : X \rightarrow L^\infty(\mathbb{R})$ be weakly measurable. That is*

$$x \mapsto \int_{\mathbb{R}} f(x, \xi) g(\xi) d\xi$$

is measurable for each $g \in C_c(\mathbb{R})$. Let $\mathcal{G} = \{g \in C_c^1(\mathbb{R}) \mid |g| \leq 1\}$. For $g \in C_c^1(\mathbb{R})$ and $A \in \mathcal{A}$ with $\mu(A) < \infty$ we let

$$\mathcal{S}_A(g) := - \int_A \int_{\mathbb{R}} f(x, \xi) g'(\xi) d\xi d\mu(x).$$

Then, for all $A \in \mathcal{A}$ such that $\mu(A) < \infty$

$$\sup_{g \in \mathcal{G}} \mathcal{S}_A(g) = \mu(A) \quad \text{and} \quad \inf_{g \geq 0} \mathcal{S}_A(g) \geq 0 \quad (44)$$

if and only if there exists a $\nu \in \mathcal{Y}(X; \mathbb{R})$ such that

$$\int_X \int_{\mathbb{R}} f(x, \xi) \partial_{\xi} \varphi(x, \xi) d\xi d\mu(x) = - \int_X \int_{\mathbb{R}} \varphi(x, \xi) d\nu_x(\xi) d\mu(x) \quad (45)$$

for all $\varphi \in L^1(X, W^{1,1}(\mathbb{R})) \cap L^1(X, C(\mathbb{R}))$.

Remark 1. The condition (44) are weak formulations of the conditions

$$\int_{\mathbb{R}} \partial_{\xi} f(x, \xi) d\xi = 1 \quad \text{and} \quad \partial_{\xi} f(x, \xi) \geq 0.$$

Proof. Suppose that there exists a $\nu \in \mathcal{Y}(X; \mathbb{R})$ satisfying (45). Then, for any $\varphi(x, \xi) = \mathbb{1}_A(x)g(\xi)$ where A and g is as above we obtain by (45)

$$\mathcal{S}_A(g) = - \int_A \int_{\mathbb{R}} f(x, \xi) g'(\xi) d\xi d\mu(x) = \int_A \int_{\mathbb{R}} g(\xi) d\nu_x(\xi) d\mu(x).$$

It follows immediately that $0 \leq \mathcal{S}_A(g)$ whenever $g \geq 0$ and $\mathcal{S}_A(g) \leq \mu(A)$ for all $g \in \mathcal{G}$. For the opposite inequality let $g_R \in C_c^1(\mathbb{R})$ satisfy $g_R(\xi) = 1$ for $\xi \in [-R, R]$ and $0 \leq g_R \leq 1$. By Fatou's lemma and the fact that $\nu_x([-R, R]) \rightarrow 1$ as $R \rightarrow \infty$ for μ -a.a. $x \in X$ we obtain

$$\liminf_{R \rightarrow \infty} \mathcal{S}_A(g_R) = \liminf_{R \rightarrow \infty} \int_A \int_{\mathbb{R}} g_R(\xi) d\nu_x(\xi) d\mu(x) \geq \mu(A).$$

It follows that $\sup_{g \in \mathcal{G}} \mathcal{S}_A(g) \geq \mu(A)$.

Let us consider the converse. We first claim that for μ -almost all $x \in X$

$$\int_{\mathbb{R}} f(x, \xi) g'(\xi) d\xi \leq 0 \quad \text{for all nonnegative } g \in C_c^1(\mathbb{R}). \quad (46)$$

By assumption $\mathcal{S}_A(g) \geq 0$ for all $A \in \mathcal{A}$, $\mu(A) < \infty$. It follows that there exists a null set N_g such that

$$\int_{\mathbb{R}} f(x, \xi) g'(\xi) d\xi \leq 0 \quad \text{for all } x \in X \setminus N_g.$$

The result does follow by the separability of $C_c^1(\mathbb{R})$ and the continuity of the above term with respect to convergence in $C^1(\mathbb{R})$. Next we want to prove the following equality

$$\mu(A) = - \int_A \inf_{g \in \mathcal{G}} \left\{ \int_{\mathbb{R}} f(x, \xi) g'(\xi) d\xi \right\} d\mu(x). \quad (47)$$

Let $\{g_j\}_{j \geq 1}$ be a dense sequence in $\mathcal{G}$. Let $s_j \in C_c^1(\mathbb{R})$ be nonnegative and satisfy

$$\bigvee_{1 \leq k \leq j} g_k \leq s_j \leq 1, \quad \text{and} \quad s_j \leq s_{j+1}$$

for all $j \geq 1$. Here $\vee$ denotes the pointwise maximum over the given functions. Now, for any $1 \leq k \leq j$ we have by (46)

$$\int_{\mathbb{R}} f(x, \xi) g'_k(\xi) d\xi \geq \int_{\mathbb{R}} f(x, \xi) s'_j(\xi) d\xi,$$

and so, whenever $f(x, \cdot)$ belongs to $L^\infty(\mathbb{R})$ we have

$$\inf_{g \in \mathcal{G}} \int_{\mathbb{R}} f(x, \xi) g'(\xi) d\xi = \inf_{k \geq 1} \int_{\mathbb{R}} f(x, \xi) g'_k(\xi) d\xi \geq \lim_{j \rightarrow \infty} \int_{\mathbb{R}} f(x, \xi) s'_j(\xi) d\xi.$$

The opposite inequality follows as $s_j \in \mathcal{G}$ for all $j \geq 1$. Similarly

$$\sup_{g \in \mathcal{G}} \mathcal{S}_A(g) = - \lim_{j \rightarrow \infty} \int_A \int_{\mathbb{R}} f(x, \xi) s'_j(\xi) d\xi d\mu(x)$$

for all $A \in \mathcal{A}$, $\mu(A) < \infty$. Hence, due to the assumption $\sup_{g \in \mathcal{G}} \mathcal{S}_A(g) = \mu(A)$ and the monotone convergence theorem

$$\begin{aligned} \mu(A) &= - \lim_{j \rightarrow \infty} \int_A \int_{\mathbb{R}} f(x, \xi) s'_j(\xi) d\xi d\mu(x) \\ &= - \int_A \lim_{j \rightarrow \infty} \int_{\mathbb{R}} f(x, \xi) s'_j(\xi) d\xi d\mu(x) \\ &= - \int_A \inf_{g \in \mathcal{G}} \int_{\mathbb{R}} f(x, \xi) g'(\xi) d\xi d\mu(x). \end{aligned}$$

This proves (47). It follows that

$$\sup_{g \in \mathcal{G}} \int_{\mathbb{R}} f(x, \xi) g'(\xi) d\xi = 1 \text{ } \mu\text{-a.a } x \in X.$$

Combined with (46) we may conclude by theorem [4, § 1.8, Theorem 1 and Corollary 1] that for μ -a.a. $x \in X$ there exists a $\nu_x \in \mathcal{P}(\mathbb{R})$ such that

$$\int_{\mathbb{R}} f(x, \xi) g'(\xi) d\mu(x) = - \int_{\mathbb{R}} g(\xi) d\nu_x(\xi) \text{ for all } g \in C_c^1(\mathbb{R}).$$

The (weak) measurability follows immediately, and so $\{\nu_x\}$ is in $\mathcal{Y}(X; \mathbb{R})$. The extension of the integration by parts formula to more general test functions is left to the reader. $\square$

Combining Lemma 1 with Alaoglu's theorem applied to $L^\infty(X \times \mathbb{R})$ we arrive at the following result:

Lemma 2. *Let $\zeta : [0, \infty) \rightarrow [0, \infty]$ be a continuous, non decreasing function satisfying $\lim_{\xi \rightarrow \infty} \zeta(\xi) = \infty$ and $\{u^n\}_{n \geq 1}$ a sequence of measurable functions such that*

$$\sup_n \int_A \zeta(|u^n|) d\mu(x) < \infty$$

for all $A \in \mathcal{A}$, $\mu(A) < \infty$. Then there exists a $\nu \in \mathcal{Y}(X; \mathbb{R})$, and a subsequence $\{u^{n(j)}\}_{j \geq 1}$

such that

$$\lim_{j \rightarrow \infty} \int_X \varphi(x, u^{n(j)}(x)) d\mu(x) = \int_X \int_{\mathbb{R}} \varphi(x, \xi) d\nu_x(\xi) d\mu(x)$$

for all $\varphi \in L^1(X, C_0(\mathbb{R}))$.

Proof. Let $f^n(x, \xi) := \mathbb{1}_{\xi \geq u^n(x)}$. As $\sup_n \left\{ \|f^n\|_{L^\infty(X \times \mathbb{R})} \right\} \leq 1$ there exists a $f \in L^\infty(X \times \mathbb{R})$ and some sub-sequence still labeled $\{f^n\}$ such that

$$\lim_{n \rightarrow \infty} \int_X \int_{\mathbb{R}} f^n(x, \xi) \varphi(x, \xi) d\xi d\mu(x) = \int_X \int_{\mathbb{R}} f(x, \xi) \varphi(x, \xi) d\xi d\mu(x)$$

for all $\varphi \in L^1(X \times \mathbb{R})$. Let $\varphi \in L^1(X, C_c^1(\mathbb{R})) \cap L^1(X \times \mathbb{R})$. Then

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_X \varphi(x, u^n(x)) d\mu(x) &= \lim_{n \rightarrow \infty} - \int_X \int_{\mathbb{R}} f^n(x, \xi) \partial_\xi \varphi(x, \xi) d\xi d\mu(x) \\ &= - \int_X \int_{\mathbb{R}} f(x, \xi) \partial_\xi \varphi(x, \xi) d\xi d\mu(x). \end{aligned}$$

Next we want to check that f satisfies the conditions of Lemma 1. Let $\varphi(x, \xi) = \mathbb{1}_A(x)g(\xi)$ where $g \in C_c^1(\mathbb{R})$ satisfies $|g| \leq 1$ and $A \in \mathcal{A}$, $\mu(A) < \infty$. Then

$$- \int_A \int_{\mathbb{R}} f(x, \xi) g'(\xi) d\xi d\mu(x) = \lim_{n \rightarrow \infty} \int_A g(u^n(x)) d\mu(x). \quad (48)$$

Let $g_k(\xi) \in \mathcal{G}$ satisfy $g_k(\xi) = 1$ for $\xi \in [-k, k]$. Then

$$\begin{aligned} \int_A g_k(u^n(x)) d\mu(x) &= \mu(\{|u^n(x)| \leq k\} \cap A) \\ &\quad + \int_{A \cap \{|u^n(x)| > k\}} g_k(u^n(x)) d\mu(x) =: T_{n,k}^1 + T_{n,k}^2. \end{aligned}$$

Now, since ζ is non-decreasing

$$\zeta(k) \sup_n \mu(A \cap \{|u^n(x)| > k\}) \leq \sup_n \int_A \zeta(|u^n(x)|) d\mu(x) < C_A,$$

and so $\sup_n \mu(A \cap \{|u^n(x)| > k\}) \rightarrow 0$ as $k \rightarrow \infty$. Hence

$$|\mu(A) - T_{n,k}^1| = \mu(A \cap \{|u^n(x)| > k\}) \leq \frac{C_A}{\zeta(k)}.$$

Similarly $T_{n,k}^2 \leq C_A/\zeta(k)$. Hence

$$\left| \int_A g_k(u^n(x)) d\mu(x) - \mu(A) \right| \leq 2 \frac{C_A}{\zeta(k)}.$$

It follows that $\sup_{g \in \mathcal{G}} \mathcal{S}_A(g) = \mu(A)$. Next we want to prove that $\mathcal{S}_A(g) \geq 0$ whenever $g \geq 0$. This follows immediately from (48). Hence, by Lemma 1 there exists a Young measure

$\nu \in \mathcal{Y}(X; \mathbb{R})$ such that

$$\lim_{n \rightarrow \infty} \int_X \varphi(x, u^n(x)) d\mu(x) = \int_X \int_{\mathbb{R}} \varphi(x, \xi) d\nu_x(\xi) d\mu(x).$$

It follows by approximation that the same is true for $\varphi \in L^1(X, C_0(\mathbb{R}))$. $\square$

Theorem 1. *Let $\zeta : [0, \infty) \rightarrow [0, \infty]$ be a continuous, non decreasing function satisfying $\lim_{\xi \rightarrow \infty} \zeta(\xi) = \infty$ and $\{u^n\}_{n \geq 1}$ a sequence of measurable functions such that*

$$\sup_{n \geq 1} \int_X \zeta(|u^n|) d\mu(x) < \infty.$$

Then there exist a sub-sequence $\{u^{n(j)}\}_{j \geq 1}$ and an associated Young measure $\nu \in \mathcal{Y}(X, \mathcal{A}, \mu; \mathbb{R})$ such that for any Carathéodory function $\psi : X \times \mathbb{R} \rightarrow \mathbb{R}$ satisfying $\psi(\cdot, u^n(\cdot)) \rightharpoonup \bar{\psi}$ in $L^1(X)$

$$\bar{\psi}(x) = \int_{\mathbb{R}} \psi(x, \xi) d\nu_x(\xi).$$

Proof. This is just the extension of the result in Lemma 2 and it follows exactly as in [8, Theorem 6.2]. $\square$

The canonical approach.

Let us denote the set of signed Radon measures on $\mathbb{R}$ by $\mathcal{M}(\mathbb{R})$. These are normed by the total variation norm

$$\|\nu\|_{\mathcal{M}} = \sup \{ |\langle \nu, g \rangle| \mid g \in C_0(\mathbb{R}), |g| \leq 1 \}$$

where $\langle \nu, g \rangle$ denotes the dual pairing between $\mathcal{M}(\mathbb{R})$ and $C_0(\mathbb{R})$. We say that $\nu : X \rightarrow \mathcal{M}(\mathbb{R})$ is weak*-measurable if

$$x \mapsto \langle \nu_x, g \rangle \text{ is measurable for all } g \in C_0(\mathbb{R}).$$

Consider the vector space of all weak*-measurable, bounded maps $\nu : X \rightarrow \mathcal{M}(\mathbb{R})$. The space $L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$ is defined as the quotient space induced by the equivalence relation $\nu^1 \sim \nu^2$ if $\langle \nu^1, g \rangle = \langle \nu^2, g \rangle$ in $L^\infty(X)$ for all $g \in C_0(\mathbb{R})$. Due to the separability of $C_0(\mathbb{R})$ this is the same as saying that there exists a null set $N \subset X$ such that $\nu_x^1 = \nu_x^2$ in $\mathcal{M}(\mathbb{R})$ for all $x \in X \setminus N$. Let us define a norm on $L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$ by

$$\sup \left\{ \|\langle \nu, g \rangle\|_{L^\infty(X)} : g \in C_0(\mathbb{R}), |g| \leq 1 \right\}.$$

A consequence of the separability of $C_0(\mathbb{R})$ is that the map $x \mapsto \|\nu_x\|_{\mathcal{M}(\mathbb{R})}$ is measurable and

$$\|\nu\|_{L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))} = \text{ess sup}_{x \in X} \left\{ \|\nu_x\|_{\mathcal{M}(\mathbb{R})} \right\}.$$

For any $\nu \in \mathcal{Y}(X; \mathbb{R})$ we have $\|\nu_x\|_{\mathcal{M}(\mathbb{R})} = 1$ for μ -a.a. $x \in X$. Hence $\mathcal{Y}(X; \mathbb{R})$ is a subset of the closed unit ball in $L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$. Suppose that the space $(L^1(X, C_0(\mathbb{R})))^*$ is

isometrically isomorphic to $L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$. Due to Alaoglu's theorem $\mathcal{Y}(X; \mathbb{R})$ is relatively compact in $L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$. It follows that for any sequence $\{\nu^n\}_{n \geq 1} \subset \mathcal{Y}(X; \mathbb{R})$ there exist a sub-sequence $\{\nu^{n(j)}\}$ and a limit $\nu \in L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$ such that

$$\lim_{j \rightarrow \infty} \int_X \langle \nu_x^{n(j)}, \varphi(x) \rangle d\mu(x) = \int_X \langle \nu_x, \varphi(x) \rangle d\mu(x)$$

for all $\varphi \in L^1(X, C_0(\mathbb{R}))$. Furthermore, since the unit ball is closed in the weak* topology $\|\nu_x\|_{\mathcal{M}(\mathbb{R})} \leq 1$ for μ -a.a. $x \in X$. Hence, due to the identification $(L^1(X, C_0(\mathbb{R})))^* \simeq L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$ we obtain a representation of the limit. In the case that $X \subset \mathbb{R}^d$ and μ is the Lebesgue measure the result is proved in [6, Theorem 2.11]. Despite the fact that the result should be rather standard it has not been easy to find it in the literature. Anyway, the proof turns out to be rather straight forward except possibly the need of a lift [5] on $M^\infty(X)$, the space of measurable bounded functions on X .

Theorem 2. *Let $(X, \mathcal{A}, \mu)$ be a complete, σ -finite measure space. Then $(L^1(X, C_0(\mathbb{R})))^*$ is isometrically isomorphic to $L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$.*

Remark 2. Note that the below proof is not dependent on the separability of $C_0(\mathbb{R})$. However, one may discard the assumption of completeness of the measure space $(X, \mathcal{A}, \mu)$, by applying the separability and just picking representatives in place of using a lift.

Proof. For each $\nu \in L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$ we define $\Phi_\nu \in (L^1(X, C_0(\mathbb{R})))^*$ by

$$\Phi_\nu(\varphi) = \int_X \langle \nu_x, \varphi(x) \rangle d\mu(x).$$

Note that

$$|\Phi_\nu(\varphi)| \leq \|\nu\|_{L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))} \|\varphi\|_{L^1(X, C_0(\mathbb{R}))}$$

so Φ_ν is indeed an element of $(L^1(X, C_0(\mathbb{R})))^*$. Furthermore, note that it is well defined on $L^1(X) \otimes_\pi C_0(\mathbb{R})$, so it is well defined on $L^1(X) \hat{\otimes}_\pi C_0(\mathbb{R}) \simeq L^1(X, C_0(\mathbb{R}))$.

Injectivity: Suppose that $\Phi_\nu = 0$. We want to prove that $\langle \nu, g \rangle = 0$ in $L^\infty(X)$ for all $g \in C_0(\mathbb{R})$. Fix $g \in C_0(\mathbb{R})$. Since $\Phi_\nu = 0$ it follows that

$$\int_X h(x) \langle \nu_x, g \rangle d\mu(x) = 0 \text{ for all } h \in L^1(X).$$

Let $A = \{x \in X : \langle \nu_x, g \rangle \geq 0\}$ and note that $A \in \mathcal{A}$. Then by letting $h = 1_B$ where $\mu(B) < \infty$ it follows that

$$0 = \int_{A \cap B} \langle \nu_x, g \rangle d\mu(x) + \int_{B \setminus A} \langle \nu_x, g \rangle d\mu(x)$$

It follows that $x \mapsto \langle \nu_x, g \rangle = 0$ locally μ -almost anywhere. Assuming that $(X, \mathcal{A}, \mu)$ is σ -finite we may conclude that $\langle \nu, g \rangle = 0$ in $L^\infty(X)$.

Surjectivity: Let $\Phi \in (L^1(X, C_0(\mathbb{R})))^*$. We want to prove the existence of some $\nu \in L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$ such that $\Phi = \Phi_\nu$. Fix $g \in C_0(\mathbb{R})$. Let $\Lambda_g : L^1(X) \rightarrow \mathbb{R}$ be defined by $\Lambda_g(h) = \Phi(h \otimes g)$.

Note that

$$|\Lambda_g(h)| = |\Phi(h \otimes g)| \leq \|\Phi\| \|h\|_{L^1(X)} \|g\|_\infty,$$

so $\Lambda_g \in (L^1(X))^*$. By the Riesz Representation Theorem [1, Theorem 8.14] $(X, \mathcal{A}, \mu)$ is σ -finite) there exists a $u_g \in L^\infty(X)$ with $\|u_g\|_{L^\infty(X)} = \|\Phi\| \|g\|_\infty$ such that

$$\Phi(h \otimes g) = \int_X h(x)u_g(x) d\mu(x) \text{ for all } h \in L^1(X).$$

Note that the map $u : C_0(\mathbb{R}) \rightarrow L^\infty(X)$ given by $u(g) = u_g$ is linear and bounded. Due to [5, § IV, Theorem 3] $((X, \mathcal{A}, \mu)$ is complete and σ -finite) there exists a lifting $\rho : L^\infty(X) \rightarrow M^\infty(X)$. Here $M^\infty(X)$ is the set of all bounded measurable functions from X to $\mathbb{R}$. For any $x \in X$ we consider the map $\Lambda_x : C_0(\mathbb{R}) \rightarrow \mathbb{R}$ defined by $\Lambda_x(g) = \rho(u_g)(x)$. It immediately follows that Λ_x is linear. Due to the properties of the lift $|\rho(u_g)(x)| \leq \|u_g\|_{L^\infty(X)}$ for all $x \in X$. Hence

$$|\Lambda_x(g)| = |\rho(u_g)(x)| \leq \|u_g\|_{L^\infty(X)} = \|\Phi\| \|g\|_\infty \text{ for all } g \in C_0(\mathbb{R}).$$

It follows that $\Lambda_x \in (C_0(\mathbb{R}))^*$. By [4, § 1.8, Theorem 1] there exists a radon measure ν_x for each $x \in X$ with $\|\nu_x\|_{\mathcal{M}(\mathbb{R})} \leq \|\Phi\|$ such that $\Lambda_x(g) = \langle \nu_x, g \rangle$ for all $g \in C_0(\mathbb{R})$. Since $\langle \nu_x, g \rangle = \rho(u_g)(x)$ it follows that $x \mapsto \langle \nu_x, g \rangle$ is measurable. Hence (taking the equivalence class) $\nu \in L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))$. It follows that

$$\begin{aligned} \Phi(h \otimes g) &= \int_X h(x)u_g(x) d\mu(x) = \int_X h(x)\rho(u_g(x)) d\mu(x) \\ &= \int_X h(x) \langle \nu_x, g \rangle d\mu(x) = \Phi_\nu(h \otimes g). \end{aligned}$$

Since $L^1(X, C_0(\mathbb{R})) \simeq L^1(X) \hat{\otimes}_\pi C_0(\mathbb{R})$ [9, Example 2.19] it follows that $\Phi = \Phi_\nu$. This proves the surjectivity. Finally note that by the above construction

$$\|\Phi_\nu\| = \|\Phi\| \geq \sup_{x \in X} \|\nu_x\|_{\mathcal{M}(\mathbb{R})} = \|\nu\|_{L_{w*}^\infty(X, \mathcal{M}(\mathbb{R}))},$$

so $\nu \mapsto \Phi_\nu$ is indeed an isometry. □

Weak convergence in L^1 .

In order to apply Theorem 1 it is necessary to show that $\{\psi(\cdot, u^n(\cdot))\}_{n \geq 1}$ has a weakly convergent subsequence. That is, we need to show weak compactness of $\{\psi(\cdot, u^n(\cdot))\}_{n \geq 1}$ in $L^1(X)$. The key property to weak convergence in L^1 is uniform integrability. The main references for this section is [3] and [2].

Definition 1. A subset $\mathcal{K}$ of $L^1(X)$ is called *uniformly integrable* if

$$\sup_{f \in \mathcal{K}} \int_{|f| \geq c} |f| d\mu \rightarrow 0 \text{ as } c \rightarrow \infty.$$

Uniform integrability is of interest due to the following result [3, § IV.8, Corollary 11]:

Theorem 3. *If a set $\mathcal{K}$ in $L^1(X, \mathcal{A}, \mu)$ is weakly sequentially compact then it is uniformly integrable. If $\mu(X) < \infty$ then uniform integrability of $\mathcal{K}$ is sufficient for $\mathcal{K}$ to be weakly*

sequentially compact.

Recall that by the Eberlain-Šmulian Theorem, in the weak topology of a Banach space, relative weak compactness is equivalent with relative sequentially weak compactness [11]. We will not be satisfied with just considering the case $\mu(X) < \infty$, so another condition turns out to be necessary.

Definition 2. A set $\mathcal{K}$ in $L^1(X)$ has a *uniform tail* if for any $\varepsilon > 0$ there exists an $E \in \mathcal{A}$ with $\mu(E) < \infty$ such that

$$\sup_{f \in \mathcal{K}} \int_{X \setminus E} |f| d\mu \leq \varepsilon.$$

It turns out that uniform integrability may be formulated in many different ways. Let me restate Theorem 3 exactly as in [3, § IV.8, Corollary 11]

Theorem 3 (Restated). If a set $\mathcal{K}$ in $L^1(X, \mathcal{A}, \mu)$ is weakly sequentially compact then

$$\lim_{\mu(E) \rightarrow 0} \int_E f d\mu = 0 \quad (49)$$

uniformly for f in $\mathcal{K}$. If $\mu(X) < \infty$ then conversely this condition is sufficient for a bounded set $\mathcal{K}$ to be weakly sequentially compact.

The two following results makes it clear that the two versions of Theorem 3 are indeed the same. Let us first compare the condition (49) with the notion of uniform integrability in Definition 1.

Proposition 1. Let $\mathcal{K} \subset L^1(X)$ be bounded. Then $\mathcal{K}$ is uniformly integrable if and only if given any $\varepsilon > 0$ there exists a $\delta(\varepsilon) > 0$ such that

$$\sup_{f \in \mathcal{K}} \int_E |f| d\mu \leq \varepsilon \text{ whenever } \mu(E) \leq \delta(\varepsilon).$$

Proof. Suppose that $\mathcal{K}$ is uniformly integrable. For any $E \in \mathcal{A}, c \geq 0$ and $f \in \mathcal{K}$

$$\int_E |f| d\mu = \int_{E \cap \{|f| \geq c\}} |f| d\mu + \int_{E \cap \{|f| < c\}} |f| d\mu. \quad (50)$$

Let $\varepsilon > 0$ and chose $c_0(\varepsilon/2)$ such that

$$\int_{|f| \geq c_0} |f| d\mu \leq \varepsilon/2 \text{ for all } f \in \mathcal{K}.$$

Next, pick $\delta(\varepsilon)$ such that $\delta c_0 \leq \varepsilon/2$. For any $f \in \mathcal{K}$ we obtain by (50) that

$$\int_E |f| d\mu = \frac{\varepsilon}{2} + \mu(E)c_0 \leq \varepsilon$$

for any $E \in \mathcal{A}$ satisfying $\mu(E) \leq \delta$. Consider the converse. Pick $g \in \mathcal{K}$. Then

$$\mu(\{|g| \geq c\}) = \int_{|g| \geq c} 1 d\mu \leq \frac{1}{c} \sup_{f \in \mathcal{K}} \int_{|f| \geq c} |f| d\mu \leq \frac{1}{c} \sup_{f \in \mathcal{K}} \|f\|_{L^1(X)}.$$

Hence, given $\delta > 0$ there exists a $c_0(\delta) > 0$ such that $\mu(\{|g| \geq c\}) \leq \delta$ for all $c \geq c_0$ and all $g \in \mathcal{K}$. By assumption, given $\varepsilon > 0$ there exists a $\delta(\varepsilon)$ such that

$$\sup_{f \in \mathcal{K}} \int_{\{|g| \geq c\}} |f| \, d\mu \leq \varepsilon \text{ whenever } \mu(\{|g| \geq c\}) \leq \delta.$$

Hence, for all $g \in \mathcal{K}$

$$\int_{|g| \geq c} |g| \, d\mu \leq \sup_{f \in \mathcal{K}} \int_{\{|g| \geq c\}} |f| \, d\mu \leq \varepsilon \text{ when } c \geq c_0(\varepsilon).$$

□

Lemma 3. Suppose $\mu(X) < \infty$. Let $\mathcal{K} \subset L^1(X)$. If $\mathcal{K}$ is uniformly integrable, then it is bounded.

Proof. Note that

$$\int |f| \, d\mu = \int_{\{|f| \geq c\}} |f| \, d\mu + \int_{\{|f| < c\}} |f| \, d\mu.$$

According to the definition of uniform integrability we may pick $c_0(1)$ such that

$$\int_{|f| \geq c_0} |f| \, d\mu \leq 1 \text{ for all } f \in \mathcal{K}.$$

Hence $\|f\|_{L^1(X)} \leq 1 + \mu(X)c_0$ for all $f \in \mathcal{K}$.

□

Let $1 < p < \infty$. Due to reflexivity, boundedness suffices for weak compactness in $L^p(X)$. In the case that $\mu(X) < \infty$ we have $L^p(X) \subset L^1(X)$ and so $(L^1(X))^* \subset (L^p(X))^*$. It follows that weak convergence in $L^p(X)$ implies weak convergence in $L^1(X)$. So by Theorem 3 uniform boundedness in $L^p(X)$ imply uniform integrability. The connection is rather straight forward:

Lemma 4. $\mathcal{K} \subset L^1(X)$ is uniformly integrable if there exists an increasing function $\Psi : [0, \infty) \rightarrow [0, \infty)$ such that $\Psi(\zeta)/\zeta \rightarrow \infty$ as $\zeta \rightarrow \infty$ and a $M \geq 0$

$$\int_X \Psi(|f(x)|) \, d\mu(x) \leq M \text{ for all } f \in \mathcal{K}$$

Remark 3. This is the de la Vallée Poussin criterion. The converse demands a more elaborate proof.

Proof. Let $\varepsilon > 0$. Choose $\zeta_0(\varepsilon)$ so that $\Psi(\zeta)/\zeta \geq M/\varepsilon$ for all $\zeta \geq \zeta_0$. This implies that

$$\left(\frac{\varepsilon}{M}\right) \frac{\Psi(|f|)}{|f|} \geq 1 \text{ whenever } |f| \geq \zeta_0.$$

Hence, whenever $\zeta \geq \zeta_0$

$$\int_{|f| \geq \zeta} |f| \, d\mu \leq \frac{\varepsilon}{M} \int_{|f| \geq \zeta} \Psi(|f|) \, d\mu \leq \varepsilon.$$

□

Let us consider the sufficiency part of Theorem 3. According to J. Diestel: “The converse is not so easy, nor should it be. There are bigger fish to fry.” [2, p. 47]. So we are apparently dealing with the easy part. The compactness stems as usual from Alaoglu’s theorem. Hence we need to consider $L^1(X)$ as a subspace of $(L^1(X))^{**}$. By the Riesz Representation theorem [1] it follows that $(L^1(X))^*$ is isometrically isomorphic to $L^\infty(X)$ whenever $(X, \mathcal{A}, \mu)$ is σ -finite. So what is $(L^\infty(X))^*$? Let us denote by $\mathbf{ba}(X, \mathcal{A}, \mu)$ all finitely additive signed finite measures on $\mathcal{A}$ which are absolutely continuous with respect to μ . We embed $\mathbf{ba}(X, \mathcal{A}, \mu)$ with the norm

$$\|\nu\|_{\mathbf{ba}(X)} = \sup_{\pi \in \Pi_0} \sum_{A \in \pi} |\nu(A)|$$

where Π_0 consists of all finite partitions of X in $\mathcal{A}$. This is the total variation norm. It makes sense to integrate functions in $L^\infty(X)$ with respect to any $\nu \in \mathbf{ba}(X)$. The procedure is straight forward. First, for a simple function $g = \sum_{i=1}^n \alpha_i \mathbb{1}_{A_i}$ we define

$$\int g d\nu = \sum_{i=1}^n \alpha_i \nu(A_i).$$

Since ν is absolutely continuous with respect to μ it follows that

$$\left| \int g d\nu \right| \leq \sum_{i=1}^n |\alpha_i| |\nu(A_i)| \leq \|g\|_{L^\infty(X)} \|\nu\|_{\mathbf{ba}(X)}. \quad (51)$$

As the simple functions are dense in $L^\infty(X)$ [1, Lemma 2.11] we may extend the integral due to the above inequality.

Remark 4. Note that the lack of σ -additivity for the measure ν destroys the strong properties of convergence for the integral associated with regular σ -additive measures (such as The Dominated Convergence Theorem).

Proposition 2. The map $\Phi : \mathbf{ba}(X, \mathcal{A}, \mu) \rightarrow (L^\infty(X, \mathcal{A}, \mu))^*$ defined by

$$\Phi(\nu)(f) = \int_X f d\nu \quad (\nu \in \mathbf{ba}(X), f \in L^\infty(X))$$

is an isometric isomorphism.

Proof. First remark that Φ is well defined and linear. Let us show that it is an isometry. It follows by (51) that $\|\Phi(\nu)\|_{(L^\infty(X))^*} \leq \|\nu\|_{\mathbf{ba}(X)}$. For any $\pi \in \Pi_0$ let $g_\pi = \sum_{A \in \pi} \text{sign}(\nu(A)) \mathbb{1}_A$. Then $\|g_\pi\|_{L^\infty(X)} \leq 1$ and

$$\sum_{A \in \pi} |\nu(A)| = \sum_{A \in \pi} \text{sign}(\nu(A)) \nu(A) = \int g_\pi d\nu.$$

Taking the supremum over all $\pi \in \Pi_0$ we obtain $\|\Phi(\nu)\|_{(L^\infty(X))^*} \geq \|\nu\|_{\mathbf{ba}(X)}$. It remains to show surjectivity. Suppose $\Lambda \in (L^\infty(X))^*$. For any $A \in \mathcal{A}$, let $\nu(A) := \Lambda(\mathbb{1}_A)$. It is straight

forward to verify that ν is a finitely additive measure that is absolutely continuous with respect to μ . Let g_π be as above. Then

$$\sum_{A \in \pi} |\nu(A)| = \sum_{A \in \pi} \text{sign}(\nu(A)) \Lambda(\mathbb{1}_A) = \Lambda(g_\pi).$$

Hence $\|\nu\|_{\mathbf{ba}(X)} \leq \|\Lambda\|_{(L^\infty(X))^*}$ and $\nu \in \mathbf{ba}(X, \mathcal{A}, \mu)$. Since

$$\int f d\nu = \Lambda(f) \text{ for all simple } f \in L^\infty(X)$$

it follows by density that $\Phi(\nu) = \Lambda$. □

We are now in a position to prove the sufficiency part of Theorem 3.

Proposition 3. Let $(X, \mathcal{A}, \mu)$ be a σ -finite measure space. Suppose $\mathcal{K} \subset L^1(X, \mathcal{A}, \mu)$ is bounded, uniformly integrable and has a uniform tail. Then $\mathcal{K}$ is relatively (sequentially) compact.

Remark 5. In the below proof we apply the following characterization of σ -additivity for finite measures: A finitely additive finite measure ν is σ -additive if and only if for any nondecreasing sequence $\{A_n\}_{n \geq 1} \subset \mathcal{A}$

$$\nu\left(\bigcup_{n \geq 1} A_n\right) = \lim_{n \rightarrow \infty} \nu(A_n).$$

Proof. Let $\{f_n\}_{n \geq 1} \subset \mathcal{K}$. We want to show that it has a weakly convergent subsequence. For any $f \in L^1(X)$ the measure ν_f defined by

$$\nu_f(A) := \int_A f d\mu \quad (A \in \mathcal{A})$$

belongs to $\mathbf{ba}(X, \mathcal{A}, \mu)$. Furthermore, $\|f\|_{L^1(X)} = \|\nu_f\|_{\mathbf{ba}(X)}$. By Proposition 2 and Alaoglu's theorem it follows that there exists a $\nu \in \mathbf{ba}(X)$ and a subsequence $\{\nu_{f_j}\}$ of $\{\nu_{f_n}\}_{n \geq 1}$ such that $\nu_{f_j} \xrightarrow{*} \nu$ as $j \rightarrow \infty$. That is,

$$\int g f_j d\mu \rightarrow \int g d\nu \text{ for all } g \in L^\infty(X).$$

We want to show that $\nu = \nu_f$ for some $f \in L^1(X)$. By the Radon-Nikodým Theorem it follows that $\nu = \nu_f$ for some measurable function f if ν is a σ -additive measure which is absolutely continuous with respect to μ . We already know that ν is absolutely continuous with respect to μ . Let us show that ν is σ -additive. That is, we need to show that for any increasing sequence $\{A_k\}_{k \geq 1} \subset \mathcal{A}$ we have

$$\nu(A) = \lim_{k \rightarrow \infty} \nu(A_k) \text{ where } A := \bigcup_{k \geq 1} A_k.$$

Let $\varepsilon > 0$. We show that there exists a $k_0(\varepsilon) \geq 1$ such that

$$|\nu(A) - \nu(A_k)| \leq \varepsilon \text{ for all } k \geq k_0(\varepsilon).$$

By the triangle inequality

$$|\nu(A) - \nu(A_k)| \leq |\nu(A) - \nu_{f_j}(A)| + |\nu_{f_j}(A) - \nu_{f_j}(A_k)| + |\nu_{f_j}(A_k) - \nu(A_k)|. \quad (52)$$

For any $E \in \mathcal{A}$

$$|\nu_{f_j}(A) - \nu_{f_j}(A_k)| \leq \int_{A \setminus A_k} |f_j| d\mu \leq \int_{E \cap (A \setminus A_k)} |f_j| d\mu + \int_{X \setminus E} |f_j| d\mu.$$

Since $\mathcal{K}$ has a uniform tail it follows that there exists an E_0 with $\mu(E_0) < \infty$ such that

$$\sup_{j \geq 1} \int_{X \setminus E_0} |f_j| d\mu \leq \frac{\varepsilon}{2}.$$

Due to uniform integrability there exists a $\delta(\varepsilon/2)$ such that

$$\sup_{f \in \mathcal{K}} \int_E |f| d\mu \leq \varepsilon/2 \text{ whenever } \mu(E) \leq \delta.$$

Observe that

$$\mu(E_0 \cap (A \setminus A_k)) = \mu((E_0 \cap A) \setminus (E_0 \cap A_k)) = \mu(E_0 \cap A) - \mu(E_0 \cap A_k).$$

Hence it follows by the σ -additivity of μ that there exists a $k_0(\delta)$ such that

$$\mu(E_0 \cap (A \setminus A_k)) \leq \delta \text{ whenever } k \geq k_0(\delta).$$

Hence, for all $j \geq 1$

$$\int_{E_0 \cap (A \setminus A_k)} |f_j| d\mu \leq \varepsilon/2 \text{ whenever } k \geq k_0(\delta).$$

It follows that

$$|\nu_{f_j}(A) - \nu_{f_j}(A_k)| \leq \varepsilon \text{ whenever } k \geq k_0$$

Letting $j \rightarrow \infty$ in (52) proves the claim. To show that f belongs to $L^1(X)$ we need to show that ν has finite total variation. By the boundedness of $\mathcal{K}$ it follows that there exists some $M \geq 0$ such that

$$\|\nu_{f_j}\|_{\mathbf{ba}(X)} = \|f_j\|_{L^1(X)} \leq M \text{ for all } j \geq 1.$$

Since the closed ball $B(0, M) \subset \mathbf{ba}(X)$ is weak* closed it follows that $\|\nu\|_{\mathbf{ba}(X)} \leq M$. $\square$

A.2 A degenerate convection-diffusion equation arising in two phase flow in porous media.

Equations of the form (1) arise in connection with flow in porous media. As an example, we consider a model describing the flow of oil and water in a porous medium. For a discussion of the adequacy and assumptions underlying this model see for instance [10, 7]. Let s_o and s_w denote the saturation of oil and water respectively. That is, the fraction of the pore space occupied by the given fluid. By definition it follows that $s_w + s_o = 1$. In general one of the fluids will be preferentially attracted by the surface of the solid skeleton (the porous medium). As a consequence, the pressure of the two fluids at equilibrium is discontinuous across the interface of the two fluids. The pressure drop across the interface are known as the capillary pressure p_{cow} . That is, $p_{cow} = p_o - p_w$. A common assumption is that p_{cow} is a non-decreasing function of the water saturation. That is, $p_{cow} = p_{cow}(s_w)$ and $p'_{cow} \geq 0$. Let q_o and q_w denote the flux of oil and water respectively. Under the assumption that gravitational forces may be neglected (horizontal flow), Darcy's law yields

$$q_w = -\lambda_w(s_w)\nabla p_w, \quad q_o = -\lambda_o(s_w)\nabla p_o,$$

where λ_o and λ_w is the phase mobility of oil and water respectively. Conservation of oil and water yields

$$\partial_t s_w + \nabla \cdot q_w = 0, \quad \text{and} \quad \partial_t s_o + \nabla \cdot q_o = 0. \quad (53)$$

Let the auxiliary pressure p_c satisfy $\nabla p_c = \frac{\lambda_w}{\lambda_T} \nabla p_{cow}$ where $\lambda_T = \lambda_o + \lambda_w$ and define $p = p_o - p_c$. We want to find equations for the unknowns p and s_w . Now, q_w may be expressed through ∇p and ∇p_{cow} by the relation

$$\begin{aligned} q_w &= -\lambda_w \nabla p_w = -\lambda_w \nabla (p + p_c - p_{cow}) = -\lambda_w \nabla p - \lambda_w \left(\frac{\lambda_w}{\lambda_T} - 1 \right) \nabla p_{cow} \\ &= \lambda_w \nabla p - \frac{\lambda_w \lambda_o}{\lambda_T} \nabla p_{cow}. \end{aligned}$$

Inserting this into (53) for s_w we arrive at

$$\partial_t s + \nabla \cdot (\lambda_w(s) \nabla p) = \nabla \cdot \left(\frac{\lambda_w(s) \lambda_o(s)}{\lambda_T(s)} \nabla p_{cow}(s) \right),$$

where $s = s_w$. This is known as the saturation equation. Next we need an equation for the pressure p . First observe that

$$\begin{aligned} q_T &= q_o + q_w = -\lambda_o \nabla p_o - \lambda_w \nabla p_w = -\lambda_o \nabla p_o - \lambda_w \nabla (p_o - p_{cow}) \\ &= -\lambda_T \nabla p_o + \lambda_w \nabla p_{cow} = -\lambda_T \nabla (p_o - p_c) = -\lambda_T \nabla p. \end{aligned}$$

As $s_w + s_o = 1$ it follows by (53) that $\nabla \cdot q_T = 0$. Hence we obtain the pressure equation

$$\nabla \cdot (\lambda_T \nabla p) = 0.$$

In the one dimensional case we may conclude that $\lambda_T \nabla p$ is constant, say $\lambda_T \partial_x p = 1$. Regarding permeability functions, simple power-type relations are often postulated. That is, for instance $\lambda_w(s) = s^2$ and $\lambda_o(s) = (1 - s)^2$. This yields the equation

$$\partial_t s + \partial_x \left(\underbrace{\frac{s^2}{s^2 + (1 - s)^2}}_{f(s)} \right) = \partial_x \left(\underbrace{\frac{s^2(1 - s)^2}{s^2 + (1 - s)^2} p'_{cow}(s)}_{A'(s)} \partial_x s \right)$$

which is of the form (1).

References

- [1] Robert G. Bartle. *The elements of integration and Lebesgue measure*. Wiley Classics Library. John Wiley & Sons Inc., New York, 1995. Containing a corrected reprint of the 1966 original [1t The elements of integration, Wiley, New York; MR0200398 (34 #293)], A Wiley-Interscience Publication.
- [2] Joe Diestel. Uniform integrability: an introduction. *Rend. Istit. Mat. Univ. Trieste*, 23(1):41–80 (1993), 1991. School on Measure Theory and Real Analysis (Grado, 1991).
- [3] Nelson Dunford and Jacob T. Schwartz. *Linear Operators. I. General Theory*. With the assistance of W. G. Bade and R. G. Bartle. Pure and Applied Mathematics, Vol. 7. Interscience Publishers, Inc., New York; Interscience Publishers, Ltd., London, 1958.
- [4] Lawrence C. Evans and Ronald F. Gariepy. *Measure theory and fine properties of functions*. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1992.
- [5] A. Ionescu Tulcea and C. Ionescu Tulcea. *Topics in the theory of lifting*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 48. Springer-Verlag New York Inc., New York, 1969.
- [6] J. Málek, J. Nečas, M. Rokyta, and M. Ružička. *Weak and measure-valued solutions to evolutionary PDEs*, volume 13 of *Applied Mathematics and Mathematical Computation*. Chapman & Hall, London, 1996.
- [7] Julia C. Muccino, William G. Gray, and Lin A. Ferrand. Toward an improved understanding of multiphase flow in porous media. *Reviews of Geophysics*, 36(3):401–422, 1998.
- [8] Pablo Pedregal. *Parametrized measures and variational principles*. Progress in Nonlinear Differential Equations and their Applications, 30. Birkhäuser Verlag, Basel, 1997.
- [9] Raymond A. Ryan. *Introduction to tensor products of Banach spaces*. Springer Monographs in Mathematics. Springer-Verlag London Ltd., London, 2002.
- [10] Adam. Szymkiewicz and SpringerLink (Online service). *Modelling Water Flow in Unsaturated Porous Media*. GeoPlanet: Earth and Planetary Sciences,. Springer Berlin Heidelberg :, Berlin, Heidelberg :, 2013.

- [11] Robert Whitley. An elementary proof of the Eberlein-Šmulian theorem. *Math. Ann.*, 172:116–118, 1967.