HMGB1 and IL-33 kinetics in trauma and hemorrhagic shock

A prospective study in trauma patients

William Ottestad

Department of Anesthesiology, Division of Emergencies and Critical Care
Oslo University Hospital

Institute of Clinical Medicine, Faculty of Medicine, University of Oslo
2019

© William Ottestad, 2020

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-640-9

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

In memory of Endre Gustavsen

*“Logic will get you from A to B. Imagination will take you everywhere”
Albert Einstein*

Contents

Acknowledgements	9
List of papers	11
Abbreviations	13
General introduction	15
Synopsis	15
Inflammation	16
The immune system and the danger theory	17
Innate immune sensors – recognizing danger	19
DAMPs and alarmins	20
Genomic storm	21
Trauma epidemiology	22
Systemic inflammatory response syndrome	23
Post-injury multiple organ failure	24
Pathophysiology of remote organ failure	25
Iatrogenic factors contributing to multiple organ failure	27
HMGB1	27
Historical background	27
Structure and intracellular function	28
Release mechanisms and cytoplasmic shuttling	29
Cell death modes and HMGB1 release	30
Necrosis	31
Apoptosis	31
Necroptosis	31
NETosis	32
Pyroptosis	32
HMGB1 post-translational modifications	33
HMGB1 receptors	35
TLR4	35
RAGE	36
CXCL12	37
Sequential and pleiotropic effects of HMGB in sterile inflammation	38
HMGB1 in disease	40
Sepsis	40
Trauma and hemorrhagic shock	40

HMGB1 and outcome in human trauma patients	42
Ischemia-reperfusion injury	42
Acetaminophen drug-induced liver injury	43
Chronic inflammatory disease and cancer	44
HMGB1 resolution of inflammation and tissue repair	44
HMGB1 antagonism	45
IL-33	46
Background	46
Structure, expression pattern and intracellular function	47
IL-33 processing and release	48
The IL-33 receptor complex	49
Extracellular function	49
Post-translational modifications	50
IL-33 in disease	50
Aims and hypotheses	52
Primary objectives	52
Secondary objectives	52
Hypotheses	52
Methodological considerations	53
Study population and epidemiology	53
Study design	54
Sampling and processing	54
Detection systems	55
ELISA	55
Bead-based immunoassay	57
Western blotting	58
Project database and OUH trauma registry	59
Outcome variables	59
Explanatory variables	60
Statistical considerations	62
Ethical aspects	63
Summary of results	64
Paper I	64
Paper II	64
Paper III	65

General Discussion	67
HMGB1 kinetics in trauma patients	67
First order kinetics – potential decay mechanisms	67
Biphasic release and correlation to outcome	68
Second wave HMGB1 – cellular source	69
HMGB1 detection – pre-analytical conditions and sample material	70
IL-33 kinetics in trauma and shock	73
IL-33 detection	74
IL-33 source	75
Concluding remarks	76
Limitations	77
Future perspectives	78
Appendix	79
References	82

Acknowledgements

The present work is based on clinical samples from trauma patients admitted to Oslo University Hospital (OUH) Ullevål during 2011–2018. My deepest gratitude goes to our patients and their relatives, who chose to contribute during their darkest hours.

Torsten Eken, my principal supervisor, words cannot express the gratitude I feel for having you as my supervisor. Your presence and enthusiasm has been outstanding. This project has been strenuous due to some unexpected challenges, but when it comes to hard work, you lead by example. I admire your diligence, discipline and honesty. Your support and guidance has been exceptional. It has been a privilege being your student, as a teacher in science you are truly a giant.

Ulf Kongsgaard, my co-supervisor, you have been with us since the inception of this project with your warm support and expert advice. In times of great difficulty you have been a true friend and a gentle moderator.

Special thanks to Ingrid Nygren Rognes. Undoubtedly, this project would not have been possible without your support. We have spent many long days and late hours together at Ullevål recruiting patients to the project, at times it was hard to endure, but your smile, laughter and uplifting spirit brought us through.

Ulf Andersson, you are a superstar in the HMGB1 field, and it has been an honor to have you with us on this project. You have contributed substantially in the preparation of our main manuscript and have been instrumental to the interpretation of data. Your knowledge of HMGB1 biology is exceptional, and your enthusiasm for our project has been truly motivational. Thank you for your almost weekly updates on news from the HMGB1 field, and for providing me with a constant flow of exciting papers.

Peter Lundbäck, thank you for taking upon you the grim task of teaching an anesthesiologist the essentials of molecular biology, and for your patience and kind support throughout this project. You have contributed heavily with ideas, experiments and interpretation of data. We share a passion for whisky, and our next project should dive into the world of Scotch and Bourbon!

Signe Søvik, I want to express my sincere appreciation for all those great scientific discussions, and thank you for bringing me and Torsten safe back down to earth when our enthusiasm skyrocketed us into orbit.

This research project has relied heavily on my co-authors who have contributed with experiments, ideas and manuscript preparation. Special thanks go to Olav Sundes and Guttorm Haraldsen for excellent collaboration and invaluable contribution to Paper III.

I am grateful to Ivan Jonassen Rimstad, Astrid Arnesen Hug, Linda Soilammi, Erlend Skaga, Cassandra Frisvoll, Per Christian Weiseth, and Tora Torrissen for help with the recruitment of patients, acquisition and pre-analytical handling of samples, and to Anne-Marie Siebke Trøseid and Jens Petter Berg for help with the optimization of pre-analytical conditions and for laboratory facilities. I am also grateful to the Department of Medical Biochemistry at OUH for providing routine laboratory analyses and to Ronny Kristiansen for extracting them in electronic form. Special thanks to Julie Katrine Lindstad and Frank Sætre for help with enzyme-linked immunosorbent assay analyses. I am grateful to Jo Røislien for his authoritative advice regarding data interpretation and statistics. Helena Erlandsson Harris, thank you for valuable discussions, and for your kind help with Paper II. HMGB1 analyses for Paper I have been performed at the Department of Immunology, OUH under the leadership of Tom Eirik Mollnes and with help and supervision from Søren E. Pischke. The study was financially supported by OUH, the University of Oslo, South-Eastern Norway Regional Health Authority, and The Research Council of Norway.

This project would not have been possible without clinical data from the OUH Trauma Registry and for the relentless and expedient work by registrars Morten Hestnes and Hans Johansson. I would also like to thank my great colleagues and the helicopter crews at the OUH Division of Pre-Hospital Services for their continuous interest in this project and help in the acquisition of prehospital samples. Last but not least, I am grateful to the many doctors, nurses and clinical laboratory technicians who have allowed us and our project to intrude into their work with critically injured patients.

Finally I would like to express my deepest love and appreciation for my family. I could not have endured this without your unconditional love and support.

William Ottestad

Oslo, December 2019

List of papers

The thesis is based on the following papers, which will be referred to by their Roman numerals:

I. Biphasic Release of the Alarmin High Mobility Group Box 1 Protein Early After Trauma Predicts Poor Clinical Outcome

William Ottestad, Ingrid N. Rognes, Soeren E. Pischke, Tom E. Mollnes, Ulf Andersson, Torsten Eken

Crit Care Med. 2019 Aug;47(8):e614-e622.

II. HMGB1 concentration measurements in trauma patients: assessment of pre-analytical conditions and sample material

William Ottestad, Ingrid N. Rognes, Erlend Skaga, Cassandra Frisvoll, Guttorm Haraldsen, Torsten Eken, Peter Lundbäck

Mol Med. 2019 Dec 31;26(1):5.

III. Rapid systemic surge of IL-33 after severe human trauma

Olav Sundnes*, William Ottestad*, Camilla Schjalm, Peter Lundbäck, Lars la Cour Poulsen, Tom Eirik Mollnes, Guttorm Haraldsen, Torsten Eken

* equal contribution by these authors

Submitted manuscript

Abbreviations

APAP	N-acetyl-para-aminophenol (Acetaminophen/Paracetamol)
ATP	Adenosine triphosphate
AUC	Area under the curve
CRP	C-reactive protein
DAMP	Damage-associated molecular pattern
DNA	Deoxyribonucleic acid
ELISA	Enzyme-linked immunosorbent assay
Fc	Crystallizable fragment, the tail region of an antibody
HAT	Histone acetyl transferase
HDAC	Histone deacetylase
HMGB1	High mobility group box protein 1
HUVEC	Human umbilical vein endothelial cell
ILC	Innate lymphoid cells
Ig	Immunoglobulin
IL	Interleukin
IL-1RA	Interleukin-1 receptor antagonist
IRAK	Interleukin-1 receptor-associated kinase
ISS	injury severity score
JAK	Janus kinase
LPS	lipopolysaccharide
mAb	Monoclonal antibody
MBL	Mannose-binding lectin
mRNA	messenger RNA
NADPH	Nicotinamide adenine dinucleotide phosphate
NET	Neutrophil extracellular trap

NF- κ B	Nuclear factor kappa B
NISS	New injury severity score
NK cells	Natural killer cells
NLR	Nucleotide-binding oligomerization domain-like receptors
NLS	Nuclear localization sequence
pAb	Polyclonal antibody
PAMP	Pathogen associated molecular pattern
PRR	Pattern recognition receptor
RAGE	Receptor for advanced glycation end products
REC	Regional Committee for Medical and Health Research Ethics
RNA	Ribonucleic acid
ROS	Reactive oxygen species
SDS	Sodium dodecyl sulfate
SIRS	Systemic inflammatory response syndrome
SLE	Systemic lupus erythematosus
ST2	suppression of tumorigenicity 2
Th	T helper
TIR	Toll/interleukin-1 receptor
TNF	Tumor necrosis factor
TRIF	TIR-domain-containing adapter-inducing interferon- β

General introduction

Synopsis

Trauma is the main cause of death for people under 40 years, with a considerable death toll even if patients arrive in hospital alive. Hemostatic resuscitation and damage-control surgery has reduced early deaths, however secondary complications, including respiratory distress syndrome, multiple organ failure and nosocomial infections remain significant causes of morbidity and mortality. These events have been linked to unbridled systemic inflammation and are the result of innate immune system activation.

A group of proteins referred to as alarmins are thought to be the most proximal mediators of the inflammatory response to injury. Alarmins are constitutively expressed proteins that harbor important housekeeping functions within the cell, however once released to the extracellular space from damaged or stressed cells they function as a “mayday signal” that initiates the innate immune response. Several alarmins have been described, among them high mobility group box protein 1 (HMGB1), interleukin 1 α (IL-1 α), interleukin 33 (IL-33), heat-shock proteins and S100 proteins.

This research effort focused in particular on HMGB1. HMGB1 has been subject to intense research for its role in sterile inflammation, and is considered to be an archetype alarmin. The causal relationship between HMGB1 and systemic inflammation is well established in numerous in vitro and in vivo models, and antibody-based HMGB1 antagonism in rodent models of trauma attenuates systemic inflammation and dramatically increases survival. Additionally we investigated the role of the novel alarmin IL-33 in human trauma due to its expression in barrier tissues and vascular endothelium. At the time of the inception of this project, detailed HMGB1 and IL-33 kinetics in human trauma was largely unexplored.

Our main goal was to generate high resolution observational data, providing solid statistical relationships between the alarmins HMGB1 and IL-33 and outcome in terms of organ failure and survival after trauma. The research groups forming the backbone of this project at Oslo University Hospital (OUS), the University of Oslo (UiO), and the Karolinska Institute (KI) have studied HMGB1 kinetics after trauma in a unique series of 1094 blood samples from 136 patients with a broad spectrum of injuries and outcomes. We coupled this information to data from the Oslo University Hospital Trauma Registry and found that levels of HMGB1 were highly correlated with adverse outcome.

HMGB1 release after trauma exhibited a biphasic pattern. Surprisingly, a delayed release of HMGB1 3–6 hours after injury predicted outcome, and thus we identified a potential clinically accessible time window for targeting HMGB1 after trauma in humans, opening up for exciting future perspectives. In cooperation with the Department of Immunology at Oslo University Hospital, we explored IL-33 kinetics in the same series of blood samples, and observed a rapid and transient release of IL-33 in a subset of critically injured patients. Finally, we found that the protein soluble suppression of tumorigenicity receptor 2 (sST2), the decoy receptor of IL-33, exhibited an inverse kinetic pattern to IL-33.

There is currently no standard operating procedure for pre-analytical handling of clinical HMGB1 samples, despite that pre-analytical conditions account for a substantial part of the overall error rate in laboratory testing. Several studies have explored HMGB1 concentrations in trauma patients utilizing standard Enzyme-linked immunosorbent assay (ELISA), however there is a considerable variation in reported concentrations and kinetics even in comparable patient populations. We hypothesized that choice of sample material or differences in pre-analytical conditions could account for some of these reported differences. Therefore a methodological study assessing the impact of pre-analytical conditions and sample material was performed. We found that HMGB1 concentrations in arterial and venous samples cannot be directly compared, and concentration values in plasma and serum must be compared with caution due to wide limits of agreement. Although HMGB1 levels in clinical samples from trauma patients are fairly stable, strict adherence to a pre-analytical protocol is advisable in order to protect sample integrity.

Inflammation

Inflammation has increasingly been recognized as an important causative or contributing factor in a wide spectrum of diseases, including cardiovascular disease, cancer, and Alzheimer. The purpose of the inflammatory response is to eliminate the inducing agent, remove damaged tissue, initiate repair and restore tissue function. (1) However, in the case of severe injury or infection, a dysregulated immune response can inflict collateral damage in remote organs not initially affected by the insult.

The cardinal signs of acute inflammation; calor (heat), rubor (redness), tumor (swelling), dolor (pain) and functio laesa (loss of function) were recognized in ancient

times and first documented by Aulus Cornelius Celsus more than 2000 years ago.

Inflammation is a tightly controlled, multicellular and highly complex response to tissue injury or altered homeostasis involving both surveillance, defense and repair mechanisms.

(1) Tissue injury can be a result of different noxious stimuli such as mechanical trauma, pathogens, heat, foreign substances, toxins, and radiation. The inflammatory response is initiated by tissue resident sensor cells also known as sentinel or guardian cells.

Macrophages and mast cells are among the most abundant sentinel cells. These cells reside in all tissues, they are not constitutively activated but respond rapidly to endogenous or exogenous stimuli of noxious character. Once activated, they secrete cytokines and chemokines that exhibit both auto-, para- and endocrine functions. Blood-borne leukocytes are recruited in large numbers to the site of injury, and the accumulation of leukocytes in the affected tissues is considered a cornerstone of inflammation.

Neutrophils are the most abundant type of blood-borne leukocytes, and they can ingest and kill a range of invading pathogens without the need for an adaptive immune response.

(2) The cardinal signs of inflammation; heat, redness, swelling and pain are a result of increased blood flow, endothelial leakage, and leukocyte extravasation. An inflammatory exudate stimulates nociceptors, and the pain serves to alert the host of an altered state. Cytokines, among them IL-1 and IL-6, act as pyrogens increasing the thermoregulatory set point in the hypothalamus, and pain and fever contribute to sickness behavior, a set of adaptive behavioral changes.

The immune system and the danger theory

Innate immunity is non-adaptive and non-specific, and represents the first line defense against infection or other noxious stimuli resulting in tissue injury. Key components of the innate immune system are evolutionary conserved and present in lower organisms. The innate immune system also includes epithelial and endothelial barriers, and the complement system. Innate immune cells recognize evolutionary conserved proteins of both exogenous and endogenous origin, and can respond to sterile or infectious stimuli that cause cellular injury or death. If barriers are breached, tissue resident innate immune cells initiate and propagate the immune response. (3)

In contrast to the innate immune system, the adaptive system recognizes pathogens in an antigen specific manner, and possesses an almost infinite memory function, serving to

protect us from re-infections. The innate and adaptive systems are not completely separate entities, and appropriate adaptive responses depend on crosstalk with innate immune cells. (4) The memory function has been considered to be exclusive to the adaptive system, however this dogma has recently been challenged. Epigenetic modifications are proposed to enhance the expression of innate immune genes in the event of reinfection, and this has been coined “innate memory”. (5)

In 1949 Burnet and Fenner introduced the *self/non-self (SNS) model* of immunity. (6) This model is based on the fundamental and simple idea that the immune system discriminates between self (defined early in life) and non-self, tolerating self and attacking non-self. According to this theory the commander of the immune response is the B lymphocyte which produces antibodies when it recognizes any non-self structure. This theory was supported by the seminal works of Medawar et al. on acquired immune tolerance. (7) They found that adult mice would accept foreign skin grafts if they had been injected with cells from the donors early in life. (8) However, the SNS model failed to answer a number of fundamental questions, like; What happens when self changes in pregnancy, puberty, and aging? Why do we fail to reject tumors that express mutated proteins? Accordingly, the SNS model was constantly refined in order to incorporate incompatible data. It was later discovered that the B lymphocyte needs help from a CD4+ T helper cell (signal 2) in order to become activated. (9) Additionally, the T helper cell also required a co-stimulatory signal from an antigen-presenting cell (APC) to be activated. (10) These observations posed a major problem for the SNS model, because APCs are not antigen specific, consequently, one has to hypothesize that yet another layer exists which not only allows non-self antigens to activate immunity.

In 1986 Charles Janeway presented a solution to this problem and launched the *infectious non-self* model. (11) He proposed that APCs are quiescent until activated via germ line encoded pattern recognition receptors (PRRs) that recognize evolutionary conserved pathogen-associated molecular patterns (PAMPs). PAMPs are a diverse set of microbial molecules, e.g. lipopolysaccharide (LPS) and viral RNA, and APCs recognize these as *infectious non-self*. However, a brisk immune response is clearly evident after severe tissue injury or shock in void of any non-self stimuli, and the SNS and *infectious non-self* models fails to explain this important observation. (12)

In 1994 Polly Matzinger proposed the Danger model: *“this viewpoint outlines a model of immunity based on the idea that the immune system is more concerned with entities that do damage than with those that are foreign”*. (13,14) The Danger model represented a paradigm shift in the understanding of the immune system and how discrimination at the effector stage of the immune response does not depend on non-self stimuli. The Danger model proposes that damage is the key to triggering an immune response, and this response is initiated by endogenous danger signals from injured or stressed cells. The organ tissues themselves are the key players in the immune response, and APCs are activated by danger signals from injured cells exposed to pathogens, toxins or mechanical trauma. Sentinel cells constantly sample their environment for danger signals, and without their activation the T helper cells cannot initiate an adaptive response. The last decade a large body of experimental data has emerged that supports this model of immunity, and a growing number of endogenous danger signals have been described in sepsis and trauma. (15-17) These endogenous danger signals have received various names including alarmins and damage-associated molecular patterns (DAMPs). (18) Endogenous DAMPs can be recognized by the same PRRs that recognize microbial PAMPs and consequently the downstream immune response is propagated through common pathways, eliciting a stereotype response. (19)

Innate immune sensors – recognizing danger

There are several families of PRRs; membrane bound, cytosolic and soluble receptors. Acute phase proteins such as C-reactive protein (CRP) and mannose-binding lectin (MBL) are soluble PRRs that recognize pathogen carbohydrates, promoting phagocytosis and complement activation. Membrane bound PRRs include toll-like receptors (TLRs) and C-type lectin receptors. (19) TLRs were first discovered in *Drosophila*, and in humans constitute a family of 10 highly conserved transmembrane spanning proteins. (20) Professional phagocytic cells express a diversity of TLRs, and TLRs are instrumental in the activation of sentinel cells. TLRs activate several signalling pathways, among which NF- κ B is the most distinctive, resulting in up-regulation of immune stimulatory cell surface proteins and pro-inflammatory cytokines. (21) The nucleotide-binding oligomerization domain-like receptors (NLRs) constitute a family of intracellularly expressed sensor proteins that become activated by cytosolic PAMPs and DAMPs. Activation of NLRs may lead to assembly of the inflammasome, a multi-protein complex

that has the capacity to activate caspase-1. (22) TLRs and NLRs are promiscuous receptors and they can recognize both endogenous and exogenous ligands.

DAMPs and alarmins

“Damp is neither wet nor dry and conjures up unpleasant moisture that is noisome or nagging. Things that are damp are usually cool, but a DAMP jettisoned from a damaged cell ignites a local conflagration, inducing the calor, rubor, tumor, and dolor that are the hallmarks of inflammation.” David S Pisetsky

The endogenous danger signals released by tissues or innate cells in the face of injury or stress have received a variety of different names, including; damage or danger associated molecular patterns (DAMPs), alarmins, hyppos and cell-death associated molecular patterns (CDAMPs). (18,23) These terms are often used synonymously, however the nomenclature is confusing and the use of these terms is inconsistent in the literature. The term DAMP is especially ambiguous, and denotes a large group of different danger molecules including multi-molecular structures such as microparticles and neutrophil extracellular traps (NETs). (23) These extracellular organelles contain many different immunostimulatory molecules, additionally the term DAMP does not fully capture the activity of these structures. DAMPs are more than patterns to be recognized and the name does not address that many of these structures and molecules are effectors critical to host defense (e.g. NETs trap and kill invading bacteria). Many consider DAMPs as the endogenous analog to PAMPs, however some use the term DAMP for all danger signals, both of endogenous origin and for microbial derived signals (PAMPs). (18) Matzinger introduced the term “hyppo” which stands for hydrophobic protein parts. (14) Hyppos are common motifs in PAMPs and in many endogenous molecules recognized by TLRs. The term alarmin was introduced by J. Oppenheim and co-workers in 2004, (24) and the following criteria to identify and classify alarmins were proposed: (25,26)

- Alarmins are constitutively available and are rapidly released following non-programmed cell death, but are not released by apoptotic cells
- Upon activation, immune cells can actively secrete alarmins by unconventional secretory mechanisms, activating inflammation-related pathways such as NF- κ B
- Alarmins recruit and activate receptor-expressing cells of the innate immune system and enhance both innate and adaptive immune responses

- Alarmins restore homeostasis by promoting tissue repair

The classic dichotomy between programmed and non-programmed cell death has recently been blurred by the discovery of programmed necrosis, termed necroptosis. (27) Alarmins are rapidly released during necroptosis, and the first criterion needs to be updated accordingly. (28) The distinction between an alarmin and a cytokine can be difficult to make. Alarmins are constitutively expressed in many cell types, they have important homeostatic roles intracellularly, and when released to the extracellular compartment they acquire a new identity to serve as potent cytokines or chemokines. (25) Contrary to alarmins, cytokines need to be induced by danger signals, and they are usually not chemotactic. (17) Honoring the above criteria the term alarmin will be used throughout the thesis.

Genomic storm

Severe trauma represents extreme stress on the cellular level. In a recent study of 167 patients, trauma altered the expression of > 80% of the leukocyte transcriptome over the first 28 days after injury. (29) In total 16,820 genes were either up or down regulated, and compared to healthy controls, a twofold change in expression was seen in 5136 genes, with more genes down regulated than up regulated. The greatest change in gene expression was seen < 12 hours after injury. Interestingly, injury severity measured by injury severity score (ISS) did not affect gene expression, and base deficit as a surrogate for physiological stress had a marginal effect with expression changes in 8 genes. In this study they were not able to identify unique expression patterns associated with the development of remote organ failure or infection when comparing complicated vs. non-complicated recovery in patients. There was no qualitative difference in gene expression pattern between the groups, the same clusters of genes were up and down regulated, however magnitude of change and duration varied significantly. In the complicated recovery group, expression was still not returned to baseline within 28 d, in contrast the uncomplicated recovery group returned to baseline within 7–14 days. There was no transcriptional evidence of a “second hit” later in the clinical course. Transcriptional activation of anti-inflammatory or immune suppressive genes occurred early and simultaneously with activation of innate immunity and pro-inflammatory pathways. The current paradigm of a sequential compensatory anti-inflammatory response syndrome as a

downstream response to the initial pro-inflammatory response was not consistent with the transcriptional pattern in blood leucocytes after injury in this study.

The genomic response to trauma, burns and endotoxemia was compared. Surprisingly, low-dose bacterial endotoxin in healthy subjects elicited a high degree of similar gene expression pattern in the blood leukocyte transcriptome compared to the expression pattern following trauma. There was an 88% overlap in gene expression pattern with respect to magnitude and direction, however duration of the response was shorter in endotoxemia. The genomic response to burns and trauma demonstrated an impressive 98% overlap. The common response was thought to be the result of endotoxin and alarmin mediated activation of the TLR4 receptor and downstream transcription factors.

This study provides genomic evidence for the danger model, and it seems like the innate immune response to different stressors is uniform and fundamental. Invading pathogens, mechanical trauma, hypoxia, radiation, thermal injury and chemical insults all lead to cellular injury, the stressors are different, however the immune response is apparently the same. (30)

Trauma epidemiology

Trauma is one of the main causes of premature death, in Norway and internationally. Worldwide, more people in the age group 0 – 40 years die from trauma than from all other diseases. (31) Approximately 5 million people died from trauma in the year 2000, or 13,700 per day. In Norway, 2,310 people died from trauma related to accidents or violence in the year 2018, excluding drowning and intoxication (statistikkbank.fhi.no/dar/).

In 1983, Scientific American published a review on trauma epidemiology in the United States written by Donald D. Trunkey. (32) This paper became one of the most influential in the history of traumatology, and fuelled the recognition of trauma as a noteworthy disease with its own epidemiology, pathophysiology, management and prognosis. Trunkey introduced the trimodal distribution of trauma deaths based upon the time interval from injury to death: Immediate deaths account for 50–60% of all trauma deaths. These patients die immediately from exsanguination, massive brain injury or airway compromise, and they are declared dead on scene, during transport or shortly after arrival to the hospital. The proportion of immediate deaths has remained largely unchanged over

time and across different trauma systems, and reflects the devastating nature of the injury patterns in these patients. (33,34) A second peak of deaths occurs within hours after arrival to the hospital. The definition of these “early” deaths varies between studies, and some authors include deaths occurring within 48 hours after arrival to trauma center. The majority of these patients die from brain injury or uncontrolled hemorrhage, despite that hemorrhage as a cause of death has decreased during the last decade. Early deaths account for 25–30% of all trauma deaths, and the proportion of deaths in this group is unchanged or has increased compared to historical controls(34,35) A contribution to this may be that a modern approach to prehospital care has assured a “scoop and run” strategy with rapid transfer to hospital, and it is possible that more patients with unsurvivable injuries are brought to hospital alive. The third peak in the trimodal distribution of trauma deaths occurs days or weeks after injury. In Trunkey and Baker’s seminal work from the 1970s these late deaths accounted for 20% of all trauma deaths, and they reported that 80% of late deaths were due to sepsis or multiple organ failure. (36) Late mortality has decreased dramatically in the last decade as a result of better intensive care. (34,37,38) In a study from 2005 of 678 trauma deaths in Dallas County USA, late deaths accounted for 9% of all trauma deaths, and the authors claim that we are moving towards a bimodal distribution of deaths. (35) A retrospective study of 260 trauma deaths from a Level 1 trauma center in Norway in the period 1996 – 2004 confirmed that over 50% of trauma victims died within the first hour after injury. (39) There was no distinct second peak, however a third peak of late deaths occurred beyond 96 h post-injury and accounted for 10% of all deaths. The risk of death from non-traumatic causes among trauma patients remains elevated for months to years after discharge from hospital, however the underlying causes for this are poorly studied and remain speculative. (40)

Systemic inflammatory response syndrome

In the case of severe injury a dysregulated and metastatic immune response can inflict collateral damage in remote organs not initially affected by the insult. (41,42) Several studies have documented a close association between systemic inflammation and increased incidence of multiple organ failure, sepsis and mortality. (43-46)

Systemic inflammatory response syndrome (SIRS) was first defined in 1991 and was the result of a sepsis-definition consensus conference charged with the task of developing a set of clinical criteria for the early identification of sepsis. (47) Four SIRS criteria were

defined: tachycardia (heart rate > 90 beats/min), tachypnea or hyperventilation (respiratory rate > 20 breaths/min or PaCO₂ < 32 mmHg), fever or hypothermia (temperature > 38 or < 36 °C), and leukocytosis or leukopenia (white blood cells > 12,000/mm³, < 4,000/mm³ or the presence of > 10% immature neutrophils). Patients who met two or more of these criteria fulfilled the definition of SIRS, and sepsis was defined as confirmed infection leading to the onset of SIRS. The systemic inflammatory response is not unique to sepsis as nearly all patients admitted to the intensive care unit meet the SIRS definition, including patients with trauma, burn injuries, pancreatitis and cardiogenic shock. (41) Thus, the SIRS criteria are highly sensitive but not specific, and the positive predictive value is low as many patients that fulfill these criteria never develop infection or adverse outcomes. (48) Therefore, in the third international consensus definition for sepsis, the SIRS criteria were abandoned and substituted with the Sequential Organ Failure Assessment (SOFA) score. (49) The predictive ability of the SOFA score for in-hospital mortality is superior to that of the SIRS definition in patients with suspected sepsis, (50) The SOFA score is increasingly applied as a tool to define organ dysfunction or failure in diverse settings, including trauma and shock. (51,52)

Organ dysfunction or failure in patients with critical illness cannot be considered a dichotomous event that is either present or absent, since progression from normal function to failure is a continuum of physiologic derangements. A cut-off level of any variable or scoring system used to assess organ function will be somewhat arbitrary and one can argue that a reasonable approach will strive to establish a dose-response relationship between any predictor variable of interest and organ dysfunction. (53)

Post-injury multiple organ failure

Multiple organ failure was first described as a downstream event to trauma, shock and major surgery in the 1970s by Baue and Eiseman. (54,55) Prevalence and temporal trends of multiple organ failure vary widely across different trauma systems. (37,38,40,56-58) In a retrospective study of 260 consecutive trauma autopsies in Norway (single center 1996–2004), multiple organ failure accounted for only 8% of the total mortality. (39) Nevertheless, several reports from different trauma systems have confirmed that patients progressing to multiple organ failure endure three to four times the number of days on mechanical ventilation compared to patients without organ failure. The absolute number of patients dying from multiple organ failure is modest, however the total burden on the

patient and the health care system is severe, and this condition continues to be resource intensive, morbid and lethal. (37,56)

Multiple scoring systems have been developed in the attempt to define multiple organ failure in epidemiological studies, but no single definition has been accepted as a gold standard. (59) Three different scoring systems are commonly used in the context of trauma: Denver Post-injury Multiple Organ Failure Score, Marshall Multiple Organ Dysfunction Score, and the previously mentioned SOFA score. Historically the Denver and Marshall scores are most commonly applied in trauma research, however recently the SOFA score has been used in a number of studies. (59) These scoring systems share several variables, and all include $\text{PaO}_2/\text{FiO}_2$ ratio, creatine and bilirubin measurements. Denver and SOFA include vasoactive drug doses, while Marshall uses pressure-adjusted heart rate, which is calculated as the product of the heart rate and the ratio of central venous pressure to mean arterial pressure. Platelet counts and Glasgow Coma Scale score are included in Marshall and SOFA, but not in the Denver score. Comparisons between studies using different scoring systems should be done with caution, since they are sufficiently different to generate apparently different organ dysfunction patterns, incidences and mortality predictions even in the same population. (52,59)

Pathophysiology of remote organ failure

Systemic inflammation is commonly associated with severe injury, and the clinical hallmarks are similar to those observed during sepsis. (12) Translocation of bacteria from the gut or as a result of broken epithelial barriers can conceivably contribute to systemic inflammation, but it is now generally accepted that SIRS following trauma can occur without microbial stimuli. (12,14,60,61)

The host response to severe injury involves hemostatic, inflammatory, endocrine and neurological systems, a concert of pathways that ultimately affect tissues far away from the site of injury (Figure 1). (41) Loss of endothelial integrity is considered a cornerstone in the pathophysiology of shock and remote organ failure. (41,62-64) The endothelial glycocalyx (glykys = sweet, kalyx = coat) is increasingly being recognized as the key to vascular homeostasis. (65) It constitutes a layer of membrane bound proteoglycans and glycoproteins on the luminal surface of the vasculature, which regulates vascular permeability and tone, facilitates migration of immune cells through expression of adhesion molecules, and contributes to hemostasis. (66) Traumatic injury leads to

Figure 1. Tissue injury and hemorrhage can lead to systemic inflammatory response syndrome (SIRS), which, if persistent, may cause remote organ failure. Activation of innate immune cells and neuroendocrine pathways enables humoral and cellular factors to damage tissues far from the site of the original injury. Damage-associated molecular patterns (DAMPs), cytokines, reactive oxygen species and catecholamines compromise endothelial integrity leading to increased vascular permeability and tissue edema. Modified from Lord et al. with permission. (41)

glycocalyx shedding or fragmentation by several mechanisms. (62,63,67) Activated neutrophils release proteolytic enzymes and promote synthesis of reactive oxygen species (ROS) leading to glycocalyx fragmentation. (68) Additionally, the neuro-endocrine response to tissue injury, pain and hypoxia releases a massive catecholamine surge further compromising the integrity of the glycocalyx layer. (69) Extensive glycocalyx shedding may ultimately lead to increased vascular permeability and tissue edema, diminishing oxygen delivery at the cellular level and contributing to remote organ failure.

Recently, a new paradigm has been proposed that explains organ failure as a transient and adaptive response to severe stress, leading to a state of hibernation. (70,71) Several experimental sepsis models and studies of patients with critical illness have documented mitochondrial dysfunction in shock-induced organ failure, however it is uncertain whether this is causative or epiphenomenal. (72-75) Post-mortem histology of heart and kidney samples from sepsis patients have revealed extensive mitochondrial injury, however minimal cell death. (76-78) There are several plausible explanations to how inflammation can cause mitochondrial dysfunction or injury. Inflammation induces NF- κ B mediated down regulation of mitochondrial proteins, and this is documented in volunteers receiving endotoxin and in sepsis patients. (79,80) Further, the exaggerated inflammatory

state is associated with increased oxidative stress, and excessive amounts of ROS and nitric oxide (NO) directly inhibit mitochondrial respiration. (81)

These two models should not be considered incompatible, a synergistic model incorporating elements from both theories is possible. Whether oxygen delivery or oxygen consumption is the limiting factor is probably highly dynamic, and varies between patients, organs, vascular beds and along the clinical course.

Iatrogenic factors contributing to multiple organ failure

Hemostatic resuscitation and damage control surgery are strategies designed to limit bleeding and restore organ perfusion, however these strategies themselves inevitably introduce several iatrogenic factors that might eventually contribute to remote organ failure. The application of massive transfusion protocols exposes the trauma patients to immunogenic factors, carrying the risk of transfusion associated acute lung injury, infections, allergic and hemolytic reactions. (82,83) The widespread use of vasopressors in the intensive care setting can lead to gut ischemia, endothelial dysfunction, hyperglycemia, cardiomyopathy and down-regulation of adaptive immunity. (84) Crystalloid and colloid fluids increase interstitial edema and reduce oxygen delivery at the cellular level, and fluid overload is associated with poor outcome. (85) The use of 100% oxygen during initial resuscitation leads to hyperoxia and the generation of reactive oxygen species (ROS), causing alveolar and microvascular damage. (86,87) Additionally, mechanical ventilation exerts mechanical stress on the alveoli and is a well recognized contributing factor to the development of acute respiratory distress syndrome (ARDS). (88)

HMGB1

Historical background

In 1973 a new group of chromatin associated proteins was discovered, and these non-histone proteins were named “high mobility group” proteins (HMG) due to their high mobility in gel electrophoresis. (89) The HMG protein family was further subdivided into three different families named HMGA, HMGB and HMGN, based on their structural properties and function. (90) The HMGB family contains four members; HMGB1, 2, 3 and 4, which all have HMG boxes that bind DNA with little sequence specificity. (90)

Initially HMGB1 was thought to reside only in the nucleus, participating in regulation of gene expression, however in 1979 Bustin et al. showed that HMGB1 can be localized in the cytoplasm implicating functions outside the nucleus. (91) A secreted protein named amphoterin involved in neurite outgrowth was later identified to be identical with HMGB1. (92) In 1999 HMGB1 was identified as a pro inflammatory cytokine and a late mediator in sepsis. (93) Delayed administration (24 h after insult) of anti-HMGB1 antibodies was shown to reverse established and lethal sepsis in a mouse model, sparking off a new era of research in inflammatory disease. (94) Administration of HMGB1 antagonists improved survival and reduced morbidity also in animal models of sterile inflammation in trauma, shock, and ischemia-reperfusion injury. The proinflammatory role of HMGB1 was later questioned when highly purified recombinant HMGB1 failed to induce cytokine responses in macrophages and neutrophils. (95-97) In 2012 mass-spectrometric analyses revealed that recombinant HMGB1 cytokine function was dependent on the redox state of the molecule, explaining the divergent in vitro data reported by different labs. (98) HMGB1 is now considered to be a prototypical alarmin, and a substantial body of evidence supports its crucial role as a mediator of systemic inflammation in a number of diseases, and has initiated a new field of translational medicine for targeting HMGB1 in disease. (99,100)

Structure and intracellular function

HMGB1 is a 25-kDa nuclear protein ubiquitously expressed in most eukaryotic cells. The amino acid sequence is highly conserved through evolution with a 99% sequence homology between mammals. HMGB1 is composed of 214 residues in a distinctive three-domain structure containing two DNA-binding domains (box A and box B) and an acidic, repetitive tail (Figure 2). (101,102) HMG box domains are present in many proteins and contain three alpha helices that fold up in a V-shaped structure that can bind to the minor groove of DNA, resulting in an allosteric transition which can promote binding of other proteins. (90) HMGB1 has a DNA chaperone like function, however the binding to DNA is transient compared to histones. HMGB1 plays an instrumental role in gene transcription. (103) HMGB1 can bind directly to nucleosomes inducing the loosening of wrapped DNA and thus facilitating chromatin remodeling. The HMGB1-nucleosome interaction is extremely dynamic, and a single HMGB1 molecule interacts with a new nucleosome every second. (104) HMGB1 binds to general transcription

factors such as TATA binding protein (TBP) and specific transcription factors including steroid nuclear receptors, HOX, p53 and NF- κ B subunits. (90,105) HMGB1 is also involved in maintenance of genome integrity, DNA replication and repair. (106,107) HMGB1 deficient mice die within a few hours from birth, underpinning the critical role of HMGB1 in normal development. (108) Additionally, HMGB1 contains two non-classical nuclear localization sequences (NLS). These lysine-rich sequences reside within box A (NLS1) and between box B and the C-terminal tail (NLS2). Concomitant acetylation of NLS1 and NLS2 promotes HMGB1 relocation to the cytosol (Figure 2). (109) Three redox-sensitive cysteine residues (C23, C45, and C106) regulate HMGB1s cytokine inducing capacity by modulating receptor usage outside the cell. (98,110)

Figure 2. The structure of HMGB1, indicating its two DNA binding domains (the A and B boxes) and the acidic C-terminal tail. The biological actions of HMGB1 are diverse and dependent on the redox status of its three inherent cysteine residues (C23, C45, and C106). Intracellular shuttling of HMGB1 between the cytoplasm and nucleus is regulated by the acetylation of lysine residues in the NLS. The sequence of human HMGB1 is included, with color coding of amino acid sequences corresponding to the functional domains. HMGB1 is highly conserved; the only difference between human and mouse/rat is the substitution of two amino acids. Abbreviations: NLS, nuclear localization signal domains; TLR4, Toll-like receptor 4; RAGE, receptor for advanced glycation end products. From Harris et al. with permission. (99)

Release mechanisms and cytoplasmic shuttling

Innate immune cells can actively secrete HMGB1 in response to inflammatory stimuli in the event of infection or tissue injury. Monocytes, macrophages, dendritic cells and natural killer cells have all been shown to actively secrete HMGB1 in response to cellular activation. (109,111-113) However it is been increasingly recognized that non-immune parenchymal cells also participate in active HMGB1 secretion. Hepatocytes are the main source of actively secreted HMGB1 during oxidative stress in the context of ischemia-reperfusion (I/R) injury in the liver. (114) HMGB1 secretion is also documented in neurons, epithelial and endothelial cells, platelets and smooth muscle cells. (92,115-118)

HMGB1 lacks a secretory peptide signal and cannot be exported through the endoplasmic reticulum–Golgi system, the classical export route for extracellular proteins. Under normal physiological conditions HMGB1 constantly shuttles between the nucleus and the cytoplasm with the equilibrium shifted towards a predominant nuclear localization. Hyperacetylation of the NLS regions block re-entry of HMGB1 into the nucleus, and shifts the equilibrium towards cytosolic accumulation. (109) Histone acetyltransferases (HATs) and histone deacetylases (HDACs) are two families of enzymes that participate in epigenetic control through acetylation and deacetylation of histones and other intracellular proteins. HATs and HDACs dynamically regulate the acetylation of HMGB1. (109,114)

Several endogenous and exogenous signals might trigger HMGB1 translocation and release. In vitro, HDAC activity is suppressed in hepatocytes exposed to hypoxia. In vivo, liver I/R injury and hypoxia has been shown to induce early release of acetylated HMGB1. (114) Binding of IL-1, TNF α , LPS or HMGB1 to surface receptors on monocytic cells activate the NF- κ B and MAP kinase (MAPK) pathways. Phosphorylated MAPKs activate HATs or inhibit HDACs, and the balance is shifted in favor of acetylation, and HMGB1 accumulates in the cytosol and is packed into secretory lysosomes. (109) Exocytosis and HMGB1 release from LPS activated monocytes is dependent on lysophosphatidylcholine (LPC) a bioactive lipid which is generated at the site of inflammation in a delayed manner, supporting the evidence that HMGB1 is a late mediator of sepsis(119) Recent evidence demonstrates that the inflammasome also mediates HMGB1 release by inducing an inflammatory type of cell death called pyroptosis (see below), which thus represents another mode of active HMGB1 release. (120-122) The complete mechanisms behind HMGB1 release are diverse and still remain a largely unexplored area of HMGB1 biology.

Cell death modes and HMGB1 release

Although macrophages and monocytes can actively secrete HMGB1 via secretory lysosomes, (111) cell death-dependent release of HMGB1 is conceivably an important source of extracellular HMGB1 in sterile inflammation. The immune system is continuously exposed to dying cells during normal cell turnover, injury and infection. Mechanisms have evolved that discriminate between cell death events signaling potential

threats and physiological programmed cell death that should be tolerated. (27) I.e., the immune system determines whether cell death is immunogenic or tolerogenic.

Necrosis

Necrosis is non-programmed cell death caused by irreversible external injury. Loss of cell membrane integrity is the hallmark of necrosis, leading to uncontrolled release of intracellular proteins and organelles which initiate an inflammatory response. Several noxious stimuli can lead to necrosis, such as infection, hypoxia, mechanical trauma, radiation, and heat. HMGB1 is bound loosely to chromatin and passively leaked from dying cells during tissue injury when the integrity of membranes are lost. (104,123) Extracellular HMGB1 serves as a marker of necrosis, however the classic dichotomy between programmed and non-programmed cell death is blurred by the discovery of several pathways leading to highly regulated and genetically controlled necrosis. (27)

Apoptosis

Apoptotic cell death is the classical form of programmed cell death and leads to a highly controlled disintegration of the cell, playing an essential role in tissue development and homeostasis. (124) Apoptosis and necrosis can be viewed as the two extremes of a continuum. Apoptotic cells retain their HMGB1 tightly bound to chromatin until cleared by macrophages, and apoptosis is largely a non-immunogenic process. (104) Apoptotic cells that escape prompt clearance by phagocytic cells undergo secondary necrosis and HMGB1 can be released to a minor extent. (104) Apoptosis is mediated by a cascade of intracellular cysteine proteases called caspases, and this process is highly energy (ATP) dependent. (124,125)

Necroptosis

During energy depletion or the presence of viral caspase inhibitors, the cell can induce a caspase independent pathway leading to a programmed form of necrosis called necroptosis. The protein kinase receptor-interacting protein 3 (RIP3) is a molecular switch between apoptosis and necroptosis. (126) TNF α stimulation leads to the formation of an RIP3–RIP1 protein complex called the necrosome. (127) The necrosome activates the pro-necrotic protein mixed lineage kinase domain like pseudokinase (MLKL) via phosphorylation, allowing to MLKL insert into and permeabilize plasma membranes and organelles. Necroptosis involves a regulated loss of membrane integrity and release of

several DAMPs including HMGB1. Human cytomegalovirus and several other viruses express caspase inhibitors that arrest the apoptotic pathway, and necroptosis might have evolved as a redundant pathway in cases where cells are unable to induce apoptosis. (128,129) Oxidative stress leading to mitochondria-derived ROS facilitates necroptosis, (130) and necroptosis is a dominant pathway during ischemic conditions like myocardial infarction, stroke and ischemia-reperfusion injury. (131-133) In vitro and in vivo models of erythrocyte transfusions have been shown to induce necroptosis of lung endothelial cells. (134)

NETosis

Neutrophils are the most abundant leukocytes in mammals and the first line of cellular defense against microbes. Neutrophils eliminate microbes through several mechanisms, among them phagocytosis, ROS production, and the release of neutrophil extracellular traps (NETs). (135) NETs are composed of externalized DNA and nuclear chromatin forming a complex meshwork of fibers plastered with granule-derived antimicrobial peptides and enzymes. NETs trap and kill invading microorganisms independent of phagocytosis. NETosis is a type of programmed cell death in activated neutrophils leading to release of NETs and ultimately plasma membrane rupture. (136) NETosis can be induced by pro inflammatory stimuli such as LPS, ROS, TNF α , IL-8 and HMGB1. HMGB1 promotes NET formation, (137) and NET formation can contribute to perpetuation of sterile inflammation and tissue injury, and plays a role in the pathogenesis of systemic lupus erythematosus (SLE), thrombosis and I/R injury. (138-140)

Pyroptosis

Pyroptosis is a caspase-dependent proinflammatory programmed cell death, classically executed by macrophages and monocytes in response to infection and sterile injury. (141) Pyroptosis features rapid plasma membrane rupture, DNA fragmentation and release of pro-inflammatory intracellular content. The complex molecular machinery of pyroptosis is still being elucidated, however pyroptosis is induced in response to sensing of cytosolic PAMPs or DAMPs. (121,141) The nucleotide-binding oligomerization domain-like receptors (NLRs) are intracellularly expressed sensor proteins and part of the larger family of pattern recognition receptors. NLRs sense cytosolic PAMPs and DAMPs and upon activation they induce formation of the inflammasome, a large multi-protein complex. (22) There are several different inflammasomes named after their NLR component, where

the NLRP3 inflammasome is the best characterized. The inflammasome mediates caspase-1 activation which in turn induces proteolytic cleavage and activation of pro-IL-1 β and pro-IL-18, and ultimately pyroptosis if caspase-1 is sufficiently triggered. Pyroptosis was initially thought to result from caspase-1-dependent formation of plasma membrane pores that dissipate cellular ionic gradients, (142) however increasing evidence suggests a crucial role of caspase-11 (caspase-4 in humans) mediated pyroptosis in endotoxemia. (143-147) Pyroptosis leads to rapid release of cytosolic content and subsequently several DAMPs, and is considered a major source of HMGB1 release in sepsis and endotoxemia. Additionally HMGB1 triggers macrophage/monocyte pyroptosis through a novel mechanism of RAGE mediated HMGB1 endocytosis. (141) Recent evidence suggests that hemorrhagic shock augments inflammasome activation and facilitates pyroptosis in lung vascular endothelium. (148)

HMGB1 post-translational modifications

HMGB1 is subject to extensive post-translational modifications, including reversible and terminal cysteine oxidation, acetylation, methylation, and phosphorylation. (149) Specific post-translational modifications influence cellular localization, release mechanisms, and receptor binding ability (Figure 3). (150) High-resolution mass spectroscopy have identified different HMGB1 isoforms, and the study of the functional properties of these isoforms has been instrumental for the understanding of HMGB1 biology.

The biological actions of HMGB1 are diverse and dependent on the redox status of its three inherent cysteine residues (C23, C45, and C106). (98) The thiol side chains of cysteine residues are susceptible to oxidation that may give rise to intra-/intermolecular disulfide bridges. Terminal oxidation renders the thiol side chain as irreversible sulphonic acid (SO₃H). The reducing environment inside the cell keeps HMGB1 in its reduced form with the thiol cysteine side chains conserved, and this redox isoform is denoted “all-thiol” or “fully reduced HMGB1”. Fully reduced HMGB1 is a potent chemotactic mediator. (110) However, direct cytokine inducing effect of HMGB1 requires a disulfide bond between C23 and C45, concomitantly C106 needs to remain in its reduced (thiol) form. (98,110) This cytokine inducing isoform is denoted “disulfide HMGB1”. HMGB1 with any of the cysteines terminally oxidized (“sulfonyl HMGB1”) does not have any known pro-inflammatory functions, (98) however one might speculate that sulfonyl HMGB1 could be involved in resolution of inflammation and tissue repair. (151) The terminal

Figure 3. In normal conditions, HMGB1 is mainly localized in the nucleus as the non-acetylated fully reduced (all-thiol) isoform. Upon tissue injury, this isoform leaks out from dead cells and can be easily converted to disulfide HMGB1. Leukocytes are recruited to the site of injury and secrete acetylated HMGB1, which may be partially oxidized before and/or after release to the extracellular space. Reductants or oxidizers present in the extracellular environment may also alter the redox state of HMGB1. From Venereau et al. with permission (100)

oxidation of HMGB1 during apoptosis is a key feature required for immune tolerance induction by apoptotic cells. During apoptosis, caspase-dependent mechanisms initiate HMGB1 oxidation through mitochondrial ROS production. (152) The cysteine redox status is a functional switch for HMGB1 inflammatory effects, and terminal oxidation might be a mechanism to limit the inflammatory response both in magnitude and duration.

Acetylation is one of the most common post-translational modifications of proteins and is crucial for the regulation and function of many proteins. HMGB1 contains 43 acetylation-susceptible lysines, potentially generating a vast number of acetylation variants. (109) Conserved key lysine stretches reside within the NLS regions, and acetylation of all lysines in the NLS regions is termed hyperacetylation. Hyperacetylation of the NLS regions shifts the HMGB1 localization equilibrium towards the cytosol and

promotes extracellular release in activated cells. (109) Monomethylation of a lysine in position 43 (K43) induces HMGB1 release from neutrophils and represents the only cell-type specific post-translational modification reported. (153) However, monomethylation of K43 occurs concomitantly with NLS hyperacetylation and its impact on HMGB1 translocation, release and function remains unknown.

Analysis of HMGB1 isoforms can give important information on cell death mode and release mechanism. In general the acetylation state can distinguish passively from actively released HMGB1, since hyperacetylation of HMGB1 shifts the equilibrium towards cytosolic accumulation. (109) HMGB1 in a resting cell is predominantly in its fully reduced form, due to the reducing intracellular environment, and consequently non-acetylated fully reduced HMGB1 is mainly a biomarker of necrosis (Figure 3). HMGB1 released by pyroptosis and necroptosis is hyperacetylated and predominantly disulfide form, however terminally oxidized forms may be released as well. (122)

HMGB1 receptors

HMGB1 acts as an innate immune sensor and signal amplifier of other DAMPs, however this has not been sufficiently studied in vivo. (154) HMGB1 has been reported to operate as a promiscuous molecule and to interact with as many as 15 distinct receptors, but this view can be challenged (155,156) Only two receptor systems, TLR4 and RAGE, have so far been extensively studied in carefully defined molecular systems and confirmed in a substantial number of studies to act as specific HMGB1 receptors. (157) Additionally, HMGB1 has a remarkable capacity to form complexes with a number of other molecules, including LPS, haptoglobin, nucleic acids, nucleosomes, histones and additional danger molecules to function in a synergistic mode, amplifying inflammation. (154,158-163)

TLR4

Toll-like receptors (TLRs) are a family of evolutionary conserved pattern-recognition receptors playing a critical role in the host's immune response to infection and tissue damage. HMGB1 has been reported to bind to TLR2, TLR4 and TLR9, which all share the ability to recognize both endogenous and exogenous ligands. (164-166) Increasing evidence indicates that TLR4 is the main functional TLR receptor system for HMGB1. HMGB1-induced activation of TLR4 requires, in similarity to LPS, the extracellular adaptor protein MD-2. HMGB1 binding to the TLR4/MD-2 complex demands HMGB1

to be in its disulfide form (C23-C45) with C106 in its thiol form. (167,168) The HMGB1-TLR4 activation triggers signaling through MYD88- and TRIF-dependent pathways subsequently inducing NF- κ B nuclear translocation, and production and release of TNF α , IL-1 β and IL-6. In vitro, extracellular HMGB1 is incapable of activating NF- κ B, the hallmark of HMGB1 signaling, when TLR4 is absent or functionally blocked. (167,168)

RAGE

The receptor for advanced glycation end products (RAGE) is a receptor of the immunoglobulin superfamily that recognizes multiple ligands including HMGB1, amyloid- β protein, S100 proteins and advanced glycation end products (AGEs). (169) RAGE was the first identified HMGB1 receptor, discovered in the context of neurite outgrowth in the fetal brain. Later discoveries have identified several HMGB1/RAGE-dependent processes mediating inflammation including pyroptosis and subsequent remote organ failure in trauma and endotoxemia. (141,148,170)

RAGE can interact with all HMGB1 redox isoforms, but the disulfide isoform binds with considerably higher affinity. (118) HMGB1/RAGE interaction induces nuclear NF- κ B translocation and transcription of chemokines, cytokines and genes involved in adhesion/migration. (171,172) However, the HMGB1/RAGE-mediated cytokine formation may not reflect direct RAGE-activated intracellular signal cascades, but may rather depend on later downstream mechanisms. This is supported by the observation that TLR4 gene-deficient macrophages with maintained RAGE expression do not produce cytokines when activated by highly purified HMGB1 of any redox isoform. (168)

It was recently demonstrated that HMGB1 binding to RAGE mediates dynamin-dependent endocytosis of HMGB1 in macrophages triggering pyroptosis. (141) Fluorochrome-labeled HMGB1 added to macrophage cultures was promptly endocytosed via RAGE and could be visualized in endosomes within 15 min and later translocated into lysosomes. The end result was inflammasome activation, pyroptosis and subsequent release of proinflammatory mediators.

RAGE can bind HMGB1 complexed with several different DAMPs and PAMPs, most noticeably LPS. (147) It has recently been demonstrated that HMGB1 delivers LPS into the cytosol of macrophages and endothelial cells through RAGE-mediated internalization. (147) Once internalized, HMGB1 acts like a detergent under the acidic conditions in lysosomes, where HMGB1 disrupts the lysosomal membrane enabling LPS to avoid

degradation, escape to the cytosol, and activate its intracellular receptor caspase-11 (caspase-4/5 in humans) with subsequent inflammasome activation and pyroptosis.

The HMGB1/RAGE axis has been shown to mediate ischemic brain injury, remote lung injury and gut barrier dysfunction in models of hemorrhagic shock and brain injury. (173-175) RAGE-deficient mice are protected from gut barrier dysfunction and pathological bacterial translocation induced by hemorrhagic shock. (173) RAGE-mediated endocytosis of HMGB1 facilitates pyroptosis in both macrophages and lung endothelial cells. (141,148) In a model of hemorrhagic shock, HMGB1 primed lung endothelial cells for pyroptosis in a RAGE-dependent manner. Binding of HMGB1 to RAGE initiated endothelial cell endocytosis of HMGB1, which in turn triggered lysosome rupture and pyroptosis. (148) This is a novel mechanism for remote organ injury after trauma and shock. Additionally the HMGB1/RAGE axis promotes neutrophil migration and triggers neutrophil-mediated injury amplification following necrosis in the liver. (170)

CXCL12

HMGB1 forms a complex with CXCL12, an abundant chemokine broadly expressed by endothelial cells. The HMGB1-CXCL12 complex binds to the CXCR4 receptor, inducing migration of leukocytes and fibroblasts. (176) HMGB1 acting in complex with CXCR12 via CXCR4 requires HMGB1 to be in its fully reduced form. (176) HMGB1 signaling through RAGE increases CXCL12 transcription, assuring sustained and increased CXCL12 production, which is used autocrinally. (177) The chemoattractant activity of HMGB1 is impaired in RAGE knockout mice, and inhibited by anti-RAGE antibodies, but RAGE is only necessary for CXCL12 secretion, while the chemotactic activity by CXCR4 is RAGE-dispensable. (176,178,179) In a model of muscle injury in mice abundant leucocyte recruitment and cell migration was demonstrated as early as 6 hours after injury, and CXCR4 blockage with its specific antagonist AMD3100 abolished chemotaxis completely. (176,177)

In summary, several in vivo models strongly indicate that TLR4 and RAGE receptors are the two main functional HMGB1 receptor systems in sterile inflammation. Cytokine production is mediated mainly by TLR4, and RAGE promotes pyroptosis, chemotaxis and amplification of the immune response (Figure 4).

Figure 4. HMGB1 redox isoform determines receptor usage. Fully reduced (all-thiol) HMGB1 (atHMGB1) forms a complex with CXCL12 that signals via CXCR4. Disulfide HMGB1 (dsHMGB1) signals via TLR4. atHMGB1, dsHMGB1 and sulfonyl HMGB1 (oxHMGB1) all bind to RAGE. From Andersson et al. with permission. (157)

Sequential and pleiotropic effects of HMGB in sterile inflammation

Genetic deletion of HMGB1 or neutralization of extracellular HMGB1 markedly reduces necrotic cell induced TNF α production, underpinning the key role of HMGB1 as a mediator of sterile inflammation. (104,168) The role of HMGB1 in sterile inflammation is context sensitive and highly dynamic along the temporal course. (149) Based on the current knowledge a simplified model of sterile inflammation might be proposed: mechanical trauma to any tissue, e.g. muscle, will lead to rapid release of fully reduced non-acetylated HMGB1 into the extracellular space. (110,149) Fully reduced HMGB1 forms heterocomplexes with CXCL12, which is always present in low concentrations in the extracellular fluid. HMGB1 activation of RAGE on neighboring cells will lead to increased CXCL12 transcription, and subsequent CXCL12 secretion. The HMGB1-CXCL12 heterocomplexes will activate CXCR4 receptors in the sub-nanomolar concentration range. The CXCR4 receptor is expressed by both endothelial cells and leukocytes. (176) Leukocytes are efficiently recruited from the microcirculation by a direct effect and indirectly by activation of endothelial cells. (171) Fully reduced HMGB1 will be partly oxidized to disulfide HMGB1 in the oxidative environment outside the cell. (110,180) Disulfide HMGB1 and other DAMPs present in the extracellular space will activate migrating leucocytes through the TLR4 receptor, inducing active secretion of

acetylated disulfide HMGB1 and cytokines (Figure 5). (110) A vicious cycle may thus evolve, where the HMGB1/RAGE axis triggers neutrophil-mediated injury amplification following necrosis. (170) Disulfide HMGB1 may also conceivably spill over to the systemic circulation and trigger platelets and immune cells in remote organs.

Hemorrhagic shock and hypoxic conditions have both been demonstrated to facilitate pyroptosis in distant tissues, thereby promoting remote organ failure. (114,148,173) The oxidation of fully reduced HMGB1 to disulfide HMGB1 is facilitated by leucocyte ROS production in the traumatized tissues. Eventually HMGB1 will be terminally oxidized to its sulfonyl form, conceivably contributing to resolution of the pro-inflammatory response. (151)

Figure 5. HMGB1 functions during muscle injury. Fully reduced (all-thiol) HMGB1 is released by necrotic muscle cells and promotes leukocyte recruitment after forming a heterocomplex with CXCL12. Disulfide HMGB1, originating from infiltrating leukocytes, activates them to release proinflammatory cytokines/chemokines, but cannot bind CXCL12 and has no chemoattractant activity. Reactive oxygen species (ROS) produced by leukocytes induce the terminal oxidation of HMGB1 cysteines to sulfonates (SO₃⁻), and HMGB1 gets inactivated. From Venereau et al. with permission. (177)

HMGB1 in disease

HMGB1 has emerged as a key player in both acute and chronic inflammation.

Experimental evidence supports its role in the pathogenesis of a multitude of diseases, ranging from sepsis and trauma to cancer and rheumatoid arthritis. (100) HMGB1 is a late mediator in infectious diseases, in contrast to conditions associated with substantial necrosis or cell damage such as trauma and shock. (123) The temporal release pattern of HMGB1 in different conditions is dependent on cellular source and release mechanism.

Sepsis

In 1999 HMGB1 was identified as a late mediator of endotoxin lethality in mice, and subsequently this was reproduced in Gram-negative experimental sepsis caused by cecal ligation and puncture, (CLP). (93,94) Pro-inflammatory cytokines like $\text{TNF}\alpha$ and IL-1 peaked within 1–2 hours after induction of experimental sepsis, in contrast to HMGB1 which peaked at 16–32 hours. HMGB1 antibodies given before or at induction of experimental sepsis did not prevent death. Remarkably and clinically highly relevant, HMGB1 antibodies administered in a delayed manner and as late as 24 hours after cecal perforation protected against sepsis lethality. (94) This was at the time of the experiments a truly unparalleled result in preclinical models of sepsis for several reasons: therapy was initiated when animals were morbidly ill, and delayed immunotherapy administered more than 8 hours after CLP had never been effective for any other targeted therapy. (181-183) Improved survival has been confirmed in other models of experimental sepsis targeting HMGB1 with specific antibodies. (120,184,185) Additionally, in similar models, HMGB1 antibodies reduced post-sepsis cognitive impairment, splenomegaly, gut barrier dysfunction, anemia, and systemic inflammation. (185-189) Systemic injection of highly purified recombinant disulfide HMGB1 in healthy animals produces a sepsis-like syndrome and ultimately death. (93,166) Increased HMGB1 serum levels have been documented in sepsis patients, and the levels are higher in non-survivors. (93,190-192)

Trauma and hemorrhagic shock

Experimental work in mice has unambiguously demonstrated a central mechanistic role for HMGB1/TLR4-mediated inflammation in trauma, shock, and ischemia-reperfusion injury (Table 1). (116,193) Systemic inflammation and epithelial barrier disruption following trauma require HMGB1. In mice, HMGB1 antagonists attenuate systemic inflammation and prevent remote organ injury in several models of trauma and

Model	Species	KO	Agent	Timing (h:m)	Therapeutic effects	Ref.
Crush injury hind limbs	Rat	-	pAb	0:05 6:00	Survival at 7 days post injury 30% → 65% Significantly decreased serum IL-6 and TNF α levels 6 h post injury. Remote lung injury with interstitial edema, and intra alveolar hemorrhage was observed in the control group, these changes were attenuated in the anti-HMGB1 treatment group.	Shimazaki et al. 2012 (197)
Pseudofracture	Mouse	-	mAb	0:00 24:00	T-cell proliferation and TH1 cytokine response was markedly depressed after trauma. Anti-HMGB1 antibody ameliorates the impaired T-cell response and immune-suppression after trauma.	Ruan et al. 2015 (199)
Closed mid-shaft femur fracture	Mouse	TLR4	pAb	0:00	Decreased systemic inflammation evaluated by IL-6 and IL-10 levels. Decreased hepatic and gut epithelial inflammation and damage. TLR ko mice did not receive any additional benefit from anti-HMGB1 therapy.	Levy et al. 2007 (193)
Femur fracture and hemorrhagic shock	Mouse	TLR4; intestinal epithelial HMGB1	pAb	-1:00	Reduced remote lung injury manifested histologically. TLR4 deficient mice and mice lacking HMGB1 in the intestinal epithelium were protected from lung injury.	Sodhi et al. 2015 (116)
Femur fracture and hemorrhagic shock	Rat	-	mAb	0:00	No effect on overall survival. Decreased serum IL-6 and TNF α levels 3 h post injury. Markedly attenuated endothelial damage and traumatic coagulopathy evaluated by thrombelastography and serum levels of activated Protein C and Syndecan-1	Xu et al. 2016 (196)
Hemorrhagic shock	Mouse	-	pAb	1:00	Decreased pulmonary levels of IL-1 β and IL-6 and complete prevention of hemorrhage induced acute lung injury.	Kim et al. 2005 (194)
Hemorrhagic shock	Mouse	-	pAb	2:30	Survival at 24 hours post hemorrhage 46% → 90%. Decreased systemic inflammation assessed by serum IL-6 and IL-10 levels. Ileal mucosal hyper-permeability and bacterial translocation substantially reduced	Yang et al. 2006 (174)
Hemorrhagic shock	Rat	-	mAb	1:30	Attenuated cardiac depression, decreased serum CK-MB/LDH levels, and decreased shock induced release of IL-6, IL-1 β and TNF α . Electron microscopy demonstrated severe morphological changes in cardiocytes and these changes were less severe in the anti-HMGB1 treatment group.	Zhou et al. 2015 (195)
Traumatic brain injury	Rat, mouse	RAGE	mAb	6:00	Reduced blood-brain barrier leakage by 88%, and leakage restricted to primary lesion site, in contrast to the control group where leakage occurred in a global manner. Motor function was substantially impaired in the control group at 6 and 24 h after injury, however the anti HMGB1 treated rats had similar motor function to sham control rats. RAGE KO mice exhibited less brain edema and anti-HMGB1 mAb did not offer any additional protection	Okuma et al. 2012 (198)

Table 1. Murine models of trauma and shock, with specific HMGB1 antagonists and reported therapeutic effects. Administration of antibodies (polyclonal, pAb, or monoclonal, mAb) was done either before injury, at the time of the insult, or after injury according to hours and minutes (h:m) in the timing column. In some of the experiments knockout (ko) animals were used in order to elucidate the receptor pathway in play.

hemorrhagic shock, among them traumatic brain injury and crush injury (Table 1).

(116,174,193-199) In these models systemic inflammation and release of IL-1 β , IL-6 and TNF α seems to be a downstream event to HMGB1/TLR4 signaling. Further, HMGB1

activation of RAGE in lung tissue after hemorrhagic shock, results in neutrophil infiltration and amplification of the inflammatory response. (170)

HMGB1 and outcome in human trauma patients

At the time of inception of this project only 3 studies had documented HMGB1 serum or plasma levels after trauma in humans. The first study was done in a small population of blunt trauma victims with hemorrhagic shock defined as admission base deficit (BD, defined as negative base excess > 6 mmol/L or SBP < 90 mmHg. HMGB1 levels were increased within 6 h of injury, and HMGB1 serum concentration was sustained for the first 24 h post-injury. (174) In this study blood samples were obtained at two time points within 24 hours after admission and no correlations to outcome was reported. In 2009 Peltz et al. reported serial measurements of HMGB1 early after trauma. Peltz et al. found markedly elevated plasma concentrations within 1 h of injury, however peak plasma HMGB1 levels occurred 2–6 h after trauma, and HMGB1 remained elevated above healthy volunteers through 136 h post injury. In this study no clear relationship was found between HMGB1 levels and ISS, transfusion requirements or BD at admission. (200) In contrast, Cohen et al. in their 2009 paper found elevated plasma levels of HMGB1 within 30 min after trauma, that correlated with severity of injury (ISS), tissue hypoperfusion (BD), traumatic coagulopathy, and levels of inflammatory mediators and complement activation. HMGB1 correlated with the development of SIRS and multiple organ failure (MOF), with significantly higher levels of HMGB1 among non-survivors. (201) The peak HMGB1 plasma concentrations in these studies varies considerably with respect to absolute concentrations, temporal aspect, and correlation to outcome.

Ischemia-reperfusion injury

Ischemia-reperfusion injury is seen in trauma, organ transplants, myocardial infarction and stroke. Tissue reperfusion after a prolonged period of ischemia triggers a number of molecular cascades that ultimately lead to cell death by necrotic, necroptotic and apoptotic pathways. Evidence from experimental models both in vitro and in vivo strongly suggests a central role of HMGB1 in the propagation of warm I/R injury in liver, kidney, heart and brain. (202-204) Hypoxia is a key initiating event in I/R injury, and aberrant HMGB1 expression, cytosolic accumulation and extracellular release increase significantly in hepatocytes exposed to hypoxia in vitro. (114) Further, HMGB1 expression is up-regulated in the liver and kidney in vivo after warm I/R injury. (114,203)

Delayed administration (immediately after reperfusion) of neutralizing antibodies to HMGB1 protects against liver and kidney I/R injury, evaluated by histology and serum liver transaminases and creatinine respectively. (202,203) Administration of recombinant disulfide HMGB1 magnifies I/R injury in liver and kidney, while TLR4 knockout mice are protected from I/R injury with no additional protective effects by HMGB1 antagonism.

The brain is profoundly sensitive to ischemia, and reperfusion of ischemic regions of the brain compromises the blood-brain barrier, however the integrity of the non-ischemic hemisphere is commonly affected as well. (205) This metastatic effect is propagated by the systemic release of HMGB1 from the site of infarction, and loss of endothelial integrity is a downstream event to global inflammation in the brain. (205) In an experimental model of 2-h occlusion of the middle cerebral artery in rats, delayed administration of anti-HMGB1 monoclonal antibody (mAb) led to a 75% reduction of infarction volume 48 h after reperfusion. (206) This dramatic reduction of infarction size markedly and consistently improved neurological deficits. HMGB1 levels are elevated in cerebrospinal fluid as early as 2 h after reperfusion of the brain. (205) The mechanism leading to loss of blood-brain barrier integrity is uncertain, however extracellular HMGB1 increases expression of IL-1 β , IL-6 and TNF α in monocytic cells which produce large amounts of ROS, threatening endothelial integrity. (207,208) HMGB1 blockade effectively protects the integrity of the blood-brain barrier both functionally and structurally, attenuating vasogenic edema. (198,205,209) Cellular edema in the brain is suggested to be a result of HMGB1/TLR4 signaling. TLR4 expression in microglial cells exhibits a temporal and spatial overlap with the development of cerebral edema. Microglia cells release IL-6 in response to HMGB1/TLR4 signaling, and astrocytes increase the expression of their water channel aquaporin-4 (AQP4) in response to IL-6. (210)

Acetaminophen drug-induced liver injury

Acetaminophen-induced liver injury is considered a highly HMGB1-dependent condition and is studied extensively in both patients and animals. (211) The initial injury caused by a hepatotoxic intracellular metabolite induces necrotic hepatocyte death followed by release of non-acetylated, fully-reduced HMGB1. This initial hepatocyte-induced HMGB1 release causes leukocyte influx into the liver and leukocyte activation, that may

be lethal due to TLR4-mediated fulminant inflammation. (167) Treatment with HMGB1-specific antagonists in acetaminophen-overdosed mice is highly successful and provides an extended therapeutic time window compared to standard therapy. (167,212)

Chronic inflammatory disease and cancer

Several studies have identified HMGB1 as a mediator of autoimmune disease such as systemic lupus erythematosus (SLE) and rheumatoid arthritis (RA). (99) HMGB1 is over-expressed in many different cancers, among them colorectal cancer, breast cancer, melanoma and mesothelioma. (213,214) HMGB1 is suggested to propagate and sustain a pro-inflammatory microenvironment in tumors, promoting angiogenesis and tumor growth. (215)

HMGB1 resolution of inflammation and tissue repair

Sterile Inflammation is often perceived as a harmful process, however we should not overlook the regenerative properties of the inflammatory response to injury. (216) HMGB1 is described to have regenerative properties in diverse tissues, and blocking HMGB1 might inflict collateral damage by interfering with physiological repair and regeneration of tissues. (151,217,218) The regenerative properties of fully reduced HMGB1 have been confirmed in several in vivo models of tissue injury, including skin, muscle and heart. (219-224) Fully reduced HMGB1 is a potent chemotactic and pro-angiogenic factor, (225) and recruits not only immune cells but also stem cells, epithelial and endothelial cells to the site of injury. At the site of injury HMGB1 facilitates proliferation of both stem cells and tissue resident cells. (226-228) HMGB1 stimulates endothelial cells and macrophages to release pro-angiogenic factors, and absence of HMGB1 in leukocytes has been shown to impair muscle regeneration due to defective angiogenesis. (229) Intramuscular administration of HMGB1 enhances skeletal muscle regeneration and vascularization, (222) and topical treatment with fully reduced HMGB1 accelerated wound healing in diabetic mice. (220) In a mouse model of myocardial infarction local administration of HMGB1 induced myocardial regeneration and restored cardiac function, (224,228,230) whereas HMGB1 blockade increased myocardial scarring and led to hypertrophy of the non-infarcted area. (223)

HMGB1 antagonism

Initial experimental studies on HMGB1 antagonism in sepsis and trauma were done with polyclonal antibodies, however due to their immunogenic nature they are not suitable for clinical applications. (93,184,231) The generation of effective monoclonal antibodies (mAb) targeting highly conserved proteins like HMGB1 is inherently difficult due to the low immunogenicity of such proteins. (232) However the monoclonal antibody 2G7 binds to an epitope within the A box and has proven effective in several disease models ranging from sepsis and trauma to arthritis and transplantation. (185,233)

Immunogenicity of xenogeneic antibodies can negatively affect clinical efficacy in human disease, but the application of recombinant DNA techniques has successfully generated human chimeric antibodies for clinical use in the treatment of autoimmune disease and cancer. A humanized chimeric 2G7 antibody (h2G7) has been developed and recent evidence confirms that the h2G7 antibody displayed similar specificity and affinity compared to its murine analog. The h2G7 proved highly effective in an in vivo model of acetaminophen drug induced liver injury, and accordingly this antibody represents a possible future tool in human disease. (212)

Affibody molecules are a class of engineered proteins acting as antibody mimetics. Affibody molecules are significantly smaller than antibodies, and conceivably they have favourable bio-distribution properties during trauma and shock. Affibody molecules specific for HMGB1 have been developed, and proven effective in a highly HMGB1-dependent disease model in mice. The Affibody molecules can be designed in different versions with biological half-life in vivo ranging from 20 min to several days. (234)

Another strategy for limiting HMGB1-driven inflammation is to interfere with HMGB1 binding to its receptors. The first indication of the feasibility of this approach came with the observation that injecting mice with a purified fragment of HMGB1, the box A domain, protected against lethality during CLP. (94) The mechanism of action is not fully understood, but since box A is a fragment of HMGB1, it might bind to HMGB1 receptors without activating them or acting as a weak agonist. It has recently been demonstrated that box A binds to RAGE and blocks the RAGE-mediated endocytosis of HMGB1 and HMGB1-complexes. (235)

A tetrameric peptide (P5779) derived from HMGB1 selectively inhibits the interaction between disulfide HMGB1 and its adaptor MD-2, blocking activation of the TLR4

receptor and the release of pro-inflammatory cytokines. (167) P5779 ameliorates APAP-mediated toxicity, I/R injury, and sepsis mortality in vivo. (167) Most importantly it does not interfere with lipopolysaccharide (LPS)-induced cytokine induction, avoiding a compromised defense against Gram-negative infections.

There are several small molecules that bind to HMGB1, blocking cytokine release and other downstream effects. Interestingly several of these molecules are derived from natural and medical herbs, among them glycyrrhizin and salicylic acid. Glycyrrhizin is the sweet tasting compound of the licorice plant *Glycyrrhiza glabra*, and in Japan it is commonly used for the treatment of hepatitis. Glycyrrhizin has been demonstrated to have protective effects in vivo in models of sepsis, brain injury and ischemia/reperfusion injury. (100) Ethyl pyruvate and ketamine have been shown to interfere with HMGB1 release mechanisms. Ketamine is a widely used intravenous anesthetic in trauma and shock patients. Ketamine inhibits LPS-induced HMGB1 release in vitro and in vivo, and attenuates sepsis induced acute lung injury. (236-238)

HMGB1 antagonism has proven successful in several preclinical models of trauma and shock, ameliorating disease severity and promoting survival. (155,239) Specific targeting of extracellular HMGB1 seems like the most rational approach for several reasons: HMGB1 has important intracellular functions critical for cell homeostasis, and HMGB1 interacts with several receptor systems which it shares with numerous other ligands. The pleiotropic role of HMGB1 in sterile inflammation is regulated by extensive post-translational modifications along the temporal course of the disease process, however antibodies blocking specific HMGB1 isoforms are still not available. In order to define a therapeutic time window the half-life and kinetics for HMGB1 and its isoforms need to be explored in patients.

IL-33

Background

Originally, IL-33 was given the name DSV27 and was cloned in a study examining differentially expressed genes in a canine model of cerebral vasospasm. (240) The function of nuclear IL-33 has been studied in human endothelial cells in which it was first described in 2003 by Baekkevold et al. (241) Later IL-33 was identified as a member of

the IL-1 superfamily of cytokines, and a ligand for the orphan receptor ST2 (also known as IL1RL1). (242) IL-33 is considered a novel alarmin with a large body of evidence supporting its role in the pathogenesis of asthma and allergic disease. (243,244) Recently there have been exciting discoveries regarding IL-33 as a potential alarmin in trauma and hemorrhagic shock (see below). (245)

Structure, expression pattern and intracellular function

The human IL-33 gene is located on chromosome 9. Full-length IL-33 (30 kDa) is produced from two transcripts which encode one protein. (246) A biologically active splice variant lacking exon 3 has been described in a human cell line of hepatic carcinoma. (247) Full-length IL-33 consists of two highly conserved N- and C-terminal regions. The N-terminal chromatin-binding domain (amino acid 1-65) of IL33 defines its normal nuclear localization, while the extracellular activities reside in its C-terminal cytokine domain (amino acid 112-270). The cytokine domain contains 12 beta-strands constituting a beta trefoil indispensable for receptor activation, a characteristic feature of all IL-1 family members. (248) In contrast to HMGB1, IL-33 is thought to repress gene expression by facilitating chromatin compaction, (249) however IL-33 housekeeping functions remain obscure. Recent evidence demonstrates that knockdown of nuclear IL-33 does not induce changes in the endothelial cell proteome, suggesting that the nuclear localization is a mechanism of regulation for extracellular cytokine activity through nuclear sequestration. (250)

There are wide differences in IL-33 expression patterns across different species. (251,252) In humans, IL-33 is constitutively expressed in quiescent, non-proliferating endothelial cells and in bronchial and gut epithelial cells. (252-254) In contrast, constitutive IL-33 production is generally not observed in mouse endothelial cells. In mice, IL-33 is constitutively expressed in epithelial cells including keratinocytes, in opposition to IL-33 expression in human skin that is found to be weak or absent, but strongly induced by interferon gamma. (251) The widely different IL-33 expression patterns in mice and humans challenge the translational value of mechanistic studies done in murine models.

IL-33 processing and release

When barriers are breached and cells destroyed, nuclear IL-33 is leaked to the extracellular space and acts as a potent alarmin. Several cytokines in the IL-1 family must undergo enzymatic cleavage by caspase-1 in order to become a functional cytokine. In contrast, IL-33 is biologically active in its 30 kDa full length form, (255,256) and caspase-1 cleavage results in attenuation or inactivation of IL-33. (256) Elastase and cathepsin G which are secreted by neutrophils, macrophages and mast cells at sites of inflammation can cleave IL-33 into N-terminally truncated forms that have up to 30-fold higher biological activity than full length IL-33 (Figure 6). (257) Due to the low potency of full length IL-33, limited proteolysis of the N-terminus is an efficient mechanism to amplify the inflammatory response at the site of inflammation. In contrast to HMGB1, oxidation outside the cell leads to the formation of two disulfide bridges that inhibits IL-33 binding to its receptor, quickly terminating the cytokine activity. (258) In the context of apoptosis nuclear IL-33 is cleaved by caspase-3 and caspase-7 to yield

Figure 6. After cell injury or necrosis, IL-33 is released as a full-length protein that can be cleaved extracellularly by proteases from cells such as neutrophils or mast cells, generating highly active mature forms that exhibit 10–30-fold higher activity. From Caroyl et al. with permission (334)

biologically inactive fragments. (259) IL-33 lacks a signal peptide for conventional secretion, and active release may depend on packaging in non-canonical secretory vesicles. (260,261) IL-33 up-regulation and translocation to cytoplasmic vesicles in an ATP dependent fashion have been documented in both fibroblasts and myocytes in response to non-lethal mechanical stress both in vitro and in vivo. (260,262) In vitro, inducible expression and active secretion have been reported in macrophages, dendritic cells and mast cells in response to LPS and pro-inflammatory cytokines. (252) The exact mechanism of IL-33 translocation from nucleus to cytoplasm is not known, however deletion of the N-terminal nuclear domain prevents IL-33 from entering the nucleus and transgenic mice lacking the N-terminal domain have high systemic IL-33 levels, developing severe Th2-cell biased systemic inflammation (see below). (263)

The IL-33 receptor complex

The IL-33 receptor complex consists of two transmembrane molecules, ST2 (also known as IL-1R4) and IL-1R3 (also known as IL1RAcP). (242,264) IL-33 binds to ST2 inducing a conformational change of ST2 that allows recruitment of accessory receptor IL-1RAcP. (265) The adaptor protein myeloid differentiation primary response protein 88 (MyD88) and IL-1R-associated kinase (IRAK) family members are recruited to this transmembrane complex via protein-protein interactions, ultimately activating NF- κ B, MAPK and JNK pathways. (265-267) ST2 also exists in a splice variant that lacks its membrane-anchoring domain, resulting in a secreted/soluble ST2 variant (sST2). (268) Expression of sST2 is up-regulated as a response to increased extracellular IL-33 levels and sST2 acts as a decoy receptor. (269) Increased serum levels of sST2 is reported in several disease states, and sST2 is a promising prognostic marker in acute coronary syndrome, and heart failure. (270-274) In sepsis elevated serum sST2 correlates with cardiac dysfunction, disease severity and mortality. (275,276)

Extracellular function

IL-33 plays an important role in Th2-cell associated immunology. IL-33 activates naive T cells and promotes their maturation toward a Th2 phenotype leading to the release of Th2-type cytokines including IL-4, IL-5 and IL-13. (242) Administration of recombinant IL-33 to mice leads to eosinophilia, splenomegaly, increased IgE and massive histological changes in the lungs and intestine. (242) The Th2 biased cytokine response has been

considered the hallmark of IL-33 signaling, favoring the recruitment of eosinophils and basophils to the site of acute inflammation. However, IL-33 also has the ability to induce Th2-type cytokines in mice that lack an adaptive immune system, (277) thus having no Th2 cells. This observation laid the ground for the intriguing discovery of innate production of Th2 cytokines by innate lymphoid cells (ILCs). (278-280) Group 2 innate lymphoid cells (ILC2) can be activated to produce IL-5 and IL-13 in response to IL-33. In a recent study, IL-33 mediated activation of ILC2 was a key driver of lung inflammation after injury and shock. (245) In addition to its Th2 type responses, IL-33 targets multiple immune cells, including dendritic cells, macrophages and mast cells, resulting in the induction of pro-inflammatory cytokines such as IL-1 β , IL-6 and TNF α , as well as chemokine synthesis. (281)

Post-translational modifications

Mature human IL-33 displays four free cysteine residues susceptible for oxidation. A recent study utilizing mass spectrometry revealed a 4-Da mass reduction in recombinant human IL-33 after it was subjected to oxidation, consistent with formation of two disulfide bridges. (258) Spectroscopy demonstrated a change in both secondary and tertiary structure leading to an extensive conformational change, terminating IL-33/ST2 interaction. Nuclear magnetic resonance (NMR) analysis confirmed a significant change in protein conformation between IL-33 and disulfide bonded IL-33 (DSB IL-33). The oxidation of IL-33 to DSB IL-33 was confirmed both in vitro and in vivo. In a culture media of human sera at 37 °C a 50% conversion from IL-33 to DSB IL-33 was evident within 1–2 h and > 90% conversion by 4 h. In vivo IL-33 is primarily released in its fully reduced form, and rapidly oxidized to DSB IL-33. This has been demonstrated in both IL-33 humanized mice, in human lung explants, and in sputum samples from asthma patients. (258)

IL-33 in disease

The involvement of IL-33 in Th2-related diseases like asthma and allergy is well documented. Recent genome-wide association studies have proposed that *IL33* and *IL1RL1* are susceptibility genes for asthma onset without being located on the same chromosome. (244,282) Furthermore, levels of IL-33 and sST2 are increased in lung tissue, bronchoalveolar lavage fluid (BAL) and sera from asthma patients and in

experimental models of allergic lung disease. (271,283,284) Cigarette smoke induces increased IL-33 expression in epithelial cells, and increased ST2 expression in macrophages and NK cells, leading to pronounced TNF α , IL-12 and interferon secretion. (285) It has been speculated that this mechanism can contribute to exacerbated inflammation and disease progress in Chronic Obstructive Pulmonary Disease patients.

The role of IL-33 in infection is context sensitive, and IL-33 can exert both deleterious and protective effects upon parasitic or viral infection. Injection of recombinant IL-33 has been shown to enhance bacterial clearance, attenuate organ damage and reduce mortality in CLP-induced sepsis in mice. (286,287) IL-33 prevents TLR4-induced down-regulation of CXCR2 in both mouse and human neutrophils, subsequently facilitating chemotaxis and thereby enhancing neutrophil influx to the site of infection. (287) The potentially protective effects of IL-33 in sepsis models is supported indirectly by the observation that sepsis survivors have lower serum concentrations of sST2. (276)

At the time of inception of this study no data existed on IL-33 in human trauma. However a recent study in a cohort of severely injured blunt trauma patients (n = 472, mean ISS 20) documented elevated IL-33 plasma levels at admission and a delayed increase in plasma sST2 levels. (245) Relative to IL-33, peak levels in sST2 were delayed by 12–24 h. The early peak in IL-33 correlated with increased systemic levels of Th2-type cytokines and the development of nosocomial infection and pulmonary dysfunction. However the polarization of native T cells is a slow process and the early appearance of Th2-type cytokines in the circulation after injury suggest that other pathways contribute to the rapid Th2-type response in trauma. In a model of blunt trauma and shock in mice, ILC2 cell counts and IL-5 expression increased rapidly within 6 h after trauma, and were dependent on IL-33 release. (245) IL-33 knockout mice proved ILC2 expression and activation as well as neutrophil IL-5 up-regulation to be IL-33 dependent. The increase in ILC2 counts and IL-5 expression was associated with severe lung injury, and both IL-33 deletion, ILC2 deficiency and IL-5 neutralization prevented trauma induced lung injury. In summary, IL-33 released after trauma and shock increased ILC2 production of IL-5 that subsequently stimulated IL-5 production by neutrophils in the lung.

Aims and hypotheses

Primary objectives

Investigate in individual trauma patients the time course of plasma concentrations of HMGB1 and IL-33, and its relationship to injury, treatment, and outcome.

Secondary objectives

1. Describe in detail serial measurements of plasma concentration of HMGB1 and IL-33
2. Relate these findings to anatomic injury, physiological derangement, acute treatment, and outcome through coupling to the Oslo University Hospital Trauma Registry and other clinical data
3. In statistical models quantify to what extent our findings predict outcome, relative to other known variables
4. Identify factors that may be used for therapeutic or diagnostic purposes, monitoring, and prediction of outcome in trauma patients

Hypotheses

The following hypotheses were stated before start of the project:

- HMGB1 concentration as a function of time after trauma is causally related to and therefore predicted by the extent of anatomic injury and physiological derangement
- HMGB1 contributes to systemic inflammation, remote organ failure and death after trauma, and its concentration as a function of time can therefore be used to predict outcome
- HMGB1 concentration changes rapidly, and repeated measurements with high time resolution are needed to fully understand the innate immune response
- Previous conflicting results regarding HMGB1 concentration and its time course after injury can be explained by different sampling times

After achieving our preliminary results, we have added to this:

- Previous conflicting results regarding HMGB1 concentration and its time course after injury can be explained by differences in pre-analytical conditions or sample material

- IL-33 will be released during severe trauma and shock due to its expression in vascular endothelium and barrier tissues

Methodological considerations

In this section I will discuss the rationale for study design, selected methods and its limitations. For comprehensive details regarding study population and technical aspects of HMGB1 and IL-33 analyses, I refer to the respective papers.

Study population and epidemiology

All investigations have been conducted on trauma patients who were admitted to Oslo University Hospital (OUH) Ullevål from January 2011 through February 2017. The hospital is equivalent to a Level 1 trauma centre, and it is the trauma referral centre for South-Eastern Norway, i.e., for 2.8 million people. Approximately 1,800 trauma patients are admitted annually and all are included in the hospital-based trauma registry, which currently contains more than 26,000 patients. Approximately 40% are severely injured, defined as ISS $\geq$ 16, and overall mortality is approximately 6%. Hospital performance is monitored with standard and novel survival prediction models and a recently developed stratified statistical process control technique, and OUH Ullevål performs favorably. (288) We aimed to study a population with a broad spectrum of injuries and outcomes in order to establish a potential dose-response relationship between injury severity, physiological derangement, HMGB1 and IL-33 release, and outcome.

This study was prospective and descriptive in nature, and recruitment of patients was done by convenience. All adult patients who met criteria for trauma team activation were eligible for enrollment. Patients less than 18 years old, burn injuries, and pregnant women were not included. Patients were followed until 10 days, discharge from ICU, or death, whichever event came first. Reference HMGB1 and IL-33 values were obtained from 20 healthy volunteers. Patients with burn injuries would have been an interesting study objective, however the majority of patients with severe burn injuries are rapidly transferred to the Burn Unit, Haukeland University Hospital, Bergen. Pregnancy is associated with substantial changes in physiology, and elevated HMGB1 levels have been documented in high risk pregnancies. To avoid this confounding factor we did not include pregnant women in our study. (289)

Study design

At the time of inception of this study the kinetics of HMGB1 and IL-33 release in trauma patients was unknown. Previous studies relied primarily on admission samples taken at variable time from injury or daily samples in the intensive care unit. We hypothesized that the HMGB1 and IL-33 concentrations change rapidly after trauma, and that analysis of a single sample and correlation to outcome would be confounded by variable time from injury to sampling. In the case of rapidly changing kinetics, concentration at admission or peak values will therefore potentially fail to reflect the true effect on outcome. The notion that effect on outcome is an effect of admission or peak value is not biologically plausible, yet a very common approach in medical research in general. (290)

We wanted to explore the effects of HMGB1 and IL-33 over time utilizing area under the concentration curves as a summary measure. Peltz et al. (200) found that plasma HMGB1 was markedly elevated within an hour after injury, peaked at 2–6 h after injury, and remained elevated through 136 h. In their study, time resolution during the first 24 h was 5–12 h. Based on this we expected the peak HMGB1 value to be found somewhere between 2–6 h. In order to better describe the initial HMGB1 release after trauma, we decided to obtain blood samples on arrival to OUH Ullevål and thereafter every 2 h the first 8 h, followed by daily samples until the patient left the ICU or until 10 days after injury. We also acquired prehospital blood samples, i.e., as close to the time of injury as possible, from some of the patients arriving by helicopter. Our main study population is comprised of 136 patients, and samples from 1094 time points are analysed. After the initial investigation on HMGB1 kinetics we decided to use the same approach exploring the potential role of the novel alarmin IL-33 in human trauma.

Sampling and processing

Inflammatory molecules have been reported to be sensitive to in vitro activation or degradation after sampling, (291) and strict control of pre-analytical conditions should be incorporated in protocols to ensure valid data. Sampling, handling and processing of sample material was performed by personnel dedicated specifically to this project, this was done to assure adherence to protocol and minimize variability in sampling routines. The sample material consisted of four test tubes for plasma, serum, coagulation analyses and RNA preservation. The plasma and serum tubes were put on ice slush immediately after sampling and centrifuged within 15 min after sampling. The supernatant was

transferred to three sterile polypropylene tubes and transferred to a -80°C freezer. Long-term storage was at the regional biobank facility operated by South-Eastern Norway Regional Health Authority.

Immediately after injury the primary source of HMGB1 or IL-33 release is likely to be mechanically injured cells. It is reasonable to assume that venous blood drained from areas of traumatized tissue will have a higher HMGB1 or IL-33 concentration than mixed venous blood. Conversely, venous blood drained from extremities with ongoing intravenous transfusion could theoretically dilute HMGB1 or IL-33 in the sample. Additionally, the common application of venous stasis during sampling of peripheral venous blood might induce hemolysis, potentially interfering with our analyses. Mixed venous blood will probably reflect the total HMGB1 or IL-33 release most accurately, however central venous lines are seldom established early after trauma. Therefore, arterial blood samples were preferred whenever possible to avoid the potential confounding factors associated with venous sampling. Potential HMGB1 or IL-33 modification or elimination in the pulmonary circulation could not be accounted for, however arterial blood reflects the HMGB1 and IL-33 signal delivered to the end organs.

Detection systems

ELISA

An immunoassay is an antibody-based biochemical method that measures the presence or concentration of a ligand, commonly a protein. Enzyme-linked immunosorbent assay (ELISA) was first introduced in the early 1970s and is the gold standard method for quantitative detection of analytes such as peptides, proteins, and antibodies. ELISA is a plate-based technique that can be used in a high-throughput manner to selectively detect and quantify low concentrations of analytes and is rapid and cost effective. The first step in an ELISA is the immobilization of antigens or antibodies to a solid surface. This step is also referred to as coating and is based upon passive adsorption, which depends on hydrophobic bonds and/or hydrophilic interactions. Polystyrene plates are most commonly used but are hydrophobic by nature, hence hydrophilic groups are introduced on the surface to improve the binding capacity of hydrophilic molecules including antibodies. ELISAs are divided into “direct”, “indirect” or “sandwich” ELISAs. In the direct and indirect ELISA the analyte is directly bound (unspecific binding) to the solid surface and detection depends only on one antigen-specific antibody, while the sandwich

ELISA requires two antigen-specific antibodies with distinct epitopes to allow parallel binding of two antibodies. In the sandwich ELISA, the solid surface is coated with a capture antibody and non-specific binding sites are blocked. When sample is added the soluble antigen binds to the capture antibody and the solid surface is washed to remove unbound non-antigen molecules and thus reduces the complexity of the sample. Subsequent detection of the captured antigen by an additional enzyme-linked antibody, known as detection antibody, confers a second layer of specificity. The detection antibody is conjugated with an enzyme, most commonly horseradish peroxidase. Antigen is stuck between two antibodies, hence the name “sandwich” ELISA.

Monoclonal antibodies are highly specific recognizing only one epitope, thus reducing the probability of cross-reactivity or unspecific binding. However, small changes in the epitope or conformational change might render a monoclonal antibody unable to detect the target protein. Development of monoclonal antibodies with high affinity against HMGB1 has proven difficult, and commercial ELISAs are based on polyclonal capture antibodies of avian origin (chicken). Polyclonal antibodies have higher overall affinity against the antigen due to recognition of multiple epitopes, however this comes with the cost of higher potential for cross reactivity and unspecific binding. In the HMGB1 sandwich ELISA (IBL/Shino-Test) applied in Paper I and II, a polyclonal antibody specific for HMGB1 has been pre-coated onto the wells of a polystyrene multi-well plate. Samples are added to the wells, HMGB1 binds to the immobilized antibody, and the plate is washed to remove unbound antigen. A peroxidase (POD) conjugated anti-HMGB1,2 monoclonal antibody is applied, and chromogenic solution is added to the antigen-antibody complex. POD catalyzes oxidation of the (near) colorless chromogenic substrate 3,3',5,5'-Tetramethylbenzidine (TMB) to a blue colored product with peak absorbances at 370 nm and 650 nm. Oxidation of TMB is a continuous reaction, and high concentrations of oxidized TMB suffer from poor solubility. A sulfuric acid stop-solution is therefore added to inactivate the reaction. Quantification is assessed as spectrophotometric absorbance at 450 nm, and absolute concentration values can be determined by comparing to standard curves with known concentrations of recombinant porcine HMGB1. Human HMGB1 (UniProtKB P09429) and porcine HMGB1 (UniProtKB P12682) have 99% sequence homology with only two residues that differ (Q142H and D207E). Nevertheless, absolute values need to be interpreted with caution especially as recombinant and endogenous proteins might differ with respect to post-translational

modifications and structure, thus making them bind with different affinity to detection antibodies. Additionally, the enzyme-mediated amplification of a signal is not always linear and can thus skew results. In general, ELISAs achieve protein quantification with high specificity and sensitivity. Several commercial ELISAs for HMGB1 exists, however the IBL kit designed by Shino-Test corporation is regarded as a robust assay by many experts in the field, and is widely used and cited in many publications. (292) Cross-reactivity with HMGB2 has been documented with several polyclonal antibodies, but is reported to be < 2% in the Shino-Test kit. (293).

Bead-based immunoassay

Building upon the “sandwich” ELISA concept, other antibody-based platforms have been developed, also multiplex alternatives. The bead-based immunoassay used to detect IL-33 in Paper III is one such multiplex assay and utilizes antibody-conjugated magnetic microspheres (“beads”) from Luminex Corporation. These beads have been internally dyed with red and infrared fluorophores of differing intensities, allowing differentiation of one bead from another. Each bead has a unique spectral identity, and this allows analysis of up to fifty analytes in one single panel. Individual bead populations are covalently conjugated with a capture antibody for one specific analyte, and many different detection reactions can be carried out simultaneously in a very small sample volume. After an analyte from a test sample is captured on the bead surface, a biotinylated detection antibody is introduced, followed by addition of a streptavidin–phycoerythrin conjugate. Phycoerythrin serves as a fluorescent reporter molecule, and its fluorescent intensity is proportional to the amount of detection antibody–analyte complexes present.

Different technologies exist for detection and quantification of analytes and are either based on laser and flow cytometry or light-emitting diodes (LED) and imaging. We applied the Bio-Plex Magpix system which utilizes LED based optics to excite the fluorophores involved, and a magnet to immobilize a monolayer of magnetic beads. Green LED light (525 nm) is used to excite the phycoerythrin (detection signal) and red LED light (634 nm) for the beads (analyte identity). Emission peaks of phycoerythrin and the beads are identified and quantified with a charge-coupled device (CCD) imager. Bead-based immunoassays have several advantages over conventional ELISA, as it allows for detection of multiple proteins simultaneously and generally has a broader dynamic range. In the bead-based assay, non-specific binding is reduced since the immobilized capture

antibody sits on a smaller surface area, requiring less antibody and smaller sample volume. (294) However multiplexing introduces a potential problem with cross-reactivities and subsequently reduced signal to noise ratio, often termed “the matrix effect”. (295) In order to minimize artifacts from multiplexing, varying degrees of sample dilution is mandatory and in the case of complex matrices like serum or plasma a fourfold dilution (1:4) is usually applied.

Western blotting

Western blotting is another immunoassay technique commonly used to identify and measure relative amounts of specific proteins in tissue lysates, supernatants or serum/plasma. Western blot provides separation of proteins based on molecular weight. Proteins are first denatured and reduced by heating the sample in a buffer supplemented with sodium dodecyl sulfate (SDS), reducing agent and bromophenol blue (tracking dye). SDS is a denaturing detergent that binds proteins via hydrophobic interactions, and SDS coating of proteins disrupts the tertiary structures in folded proteins and renders them linear. SDS binds rather uniformly to linear proteins and gives all proteins a uniform negative charge which masks the intrinsic charges of amino-acid side chains. Thus, charge becomes proportional to molecular weight so that electrophoretic separation is not biased by differences in electrical charges or structural differences. The prepared sample is applied to the wells of an SDS polyacrylamide electrophoresis gel, an electric field is applied to the gel for a defined time period, and the SDS-treated proteins will move toward the positive anode at different rates depending on their molecular weight.

In order to make the proteins accessible to antibody detection, they are transferred (“electro-blotted”) onto a membrane (commonly nitrocellulose). The membrane is subsequently treated with a non-fat dry milk solution in order to block areas where no proteins have been transferred. The membrane is probed with an antibody specific to the protein of interest (primary antibody), and then washed to remove unbound primary antibody. An enzyme conjugated secondary antibody targets the primary antibody, and is used to amplify the signal and to cleave a chemiluminescent agent, producing luminescence signal directly proportional to the amount of protein detected by the primary antibody.

Cross-reactivity of the antibodies with other proteins in the mixture can be distinguished by the expected molecular weight of the protein of interest, granting

western blot a high test specificity. However, a caveat is that post-translational modifications or proteolytic cleavage can alter the apparent size of the protein of interest in different samples. Additionally, proteins show variability in the amount of bound SDS, altering migration speed during gel electrophoresis. (296) Test sensitivity is mostly dependent on the antibodies, and validation of an antibody in an ELISA assay does not necessarily translate well for the use in western blotting or vice versa. Western blot involves a denaturing step, thus altering native epitopes, and one must use antibodies that recognize linear epitopes. Further, high protein content in complex matrices like serum and plasma can interfere with protein migration in the electrophoretic gel and thus limit the sample volume that can be loaded, however an enrichment step can enhance both specificity and sensitivity of the western blot in such cases.

Project database and OUH trauma registry

We have developed a relational database where all analysis results are stored corresponding to time after injury. The analysis results are also linked to electronically acquired routine laboratory tests from the hospital stay and to clinical data obtained from the OUH Trauma Registry, which is an internal hospital quality registry where all trauma patients are described formally in great detail. Available information includes demographical data, injury mechanism, classification and scoring of all anatomic injuries, physiological derangement, information on outcome, as well as times for injury, transport, and treatment. We have supplemented information about treatment in the trauma registry with data collected manually from the intensive care units (ICUs) at OUH Ullevål.

Outcome variables

Lung failure is the most frequent organ failure after trauma. (42,56,297) Pulmonary contusion, ISS > 25, and massive transfusion are independent risk factors for the development of ARDS after blunt trauma(298-300) Respiratory failure as a single cause of death is rare, however ARDS is strongly associated with multiple organ failure. (301,302) Trauma patients with multiple organ failure require longer time on ventilator, and they endure longer ICU stays. (56) Ventilator free days (VFD), defined as days alive and off ventilator during the first 30 days after trauma, was used as primary outcome measure. There is a close correlation between duration of ventilation and mortality, and VFD is considered as a combined measure of mortality and morbidity. (303) The main

focus in our study was remote organ failure as a downstream effect of unbridled inflammation. Therefore, we did not include mortality as an endpoint since the majority of the patients that died did so within the first 24 h after admission, reflecting the devastating nature of the injury patterns in these patients. The early deaths of these patients can be accounted for by injury severity and pattern, and the extent of the innate immune response is less likely to play a substantial role in the pathophysiology.

Secondary outcome measures, focusing on early signs of remote organ injury were maximal serum creatinine, alanine aminotransferase (ALT), aspartate aminotransferase (AST), and bilirubin within 48 h after injury, and body weight-adjusted total noradrenaline dose administered from 8 h after admission through the ICU day (24-h period from 7 am) starting no later than 48 h after admission.

Systemic inflammation leads to vascular leakage, reduced vasomotor tone, and the need for vasopressors or inotropic drugs to avoid hypotension. We reasoned that the accumulated vasopressor dose could be considered as a surrogate marker for the severity of proceeding systemic inflammation. Noradrenaline was the predominant drug of choice to ensure adequate systemic vascular resistance and subsequently blood pressure in our trauma patients during the ICU course.

Explanatory variables

Several models for prediction of mortality after trauma have been developed. The US Trauma and Injury Severity Score (TRISS) and the UK Trauma Audit and Research Network (TARN) probability of survival model are the most widely used models to monitor institutional performance and quality of trauma care. (304) Both models incorporate measures of anatomic injury, physiological derangement, age, and mechanism of injury, which are all well-founded predictors of outcome in trauma. The TRISS model has been criticized for dichotomizing age and both TARN and TRISS for not taking into account pre-injury co-morbidity. Recently a Norwegian survival model in trauma (NORMIT) has been developed in order to resolve some of the problems with TRISS and TARN. (288,305)

Injury severity. The Injury Severity Score (ISS) was developed as a tool to predict survival after trauma. (306) ISS is an anatomical scoring system where each injury is assigned an Abbreviated Injury Scale (AIS) code allocated to one of six body regions.

Each AIS code has a severity score, which is an ordinal value from 1 to 6 describing the range from minor to unsurvivable injuries. The 3 most severely injured body regions have their single maximal score squared, and their sum is called the ISS. The ISS is consequently scaled from 1 to 75, however the 44 possible ISS values are not uniformly distributed. New injury severity score (NISS) is a modification of ISS and is defined as the sum of squares of the AIS scores of each of a patient's three most severe injuries, regardless of body region. When only the 3 most severe injuries are taken into account, multiple injuries might not be well reflected in the score of an individual patient. Many different injury patterns can yield the same ISS/NISS score, and injuries to different body regions are not weighted. (307) An ISS/NISS value resulting from injuries to the head will conceivably represent a higher mortality than numerically similar values generated by injuries to the extremities, however this is not reflected in the score. Excluding patients with major head trauma conceivably reduces the NISS score as a source of unjustified heterogeneity.

AIS score triplets that produce identical ISS/NISS scores have been proven to have markedly different mortality rates. (307) NISS is shown to be a better predictor of mortality than ISS, however there is conflicting data on this issue. (308,309) Taking into account the 3 most severe injuries, we reasoned that the NISS would perform better than ISS to reflect the total burden of trauma. Additionally, NISS is shown to perform better than ISS in predicting MOF. (308) ISS and NISS do not exhibit a linear relationship to mortality, and both scoring systems are markedly non-monotonic, displaying large spikes and drops in mortality for successive values. In one large study the mortality dropped from 38% to 8% when the NISS score increased from 25 to 29. (307)

In our multiple regression analysis regarding HMGB1, we have chosen to use NISS as a surrogate marker for the total burden of trauma despite its non-linear and non-monotonic nature. We acknowledge the limitations in applying NISS as a tool to adjust for injury severity in our regression models, however no other scoring systems are easily accessible. The universal and frequent use of the ISS/NISS scoring system makes it possible to compare results across studies and different trauma systems.

Base excess. Base excess (BE) is defined as the amount of strong acid that must be added to each liter of fully oxygenated blood in order to return the pH to 7.4 at a temperature of 37 °C and pCO₂ of 5.3 kPa. The predominant base contributing to base

excess in the blood is bicarbonate. During progressive hypovolemic shock, metabolic acids accumulate in the blood stream as a result of anaerobic metabolism. Metabolic acids most notably lactic acid deplete bicarbonate. A negative base excess, also termed a base deficit (BD), is a useful surrogate marker of cellular hypoxia.

Admission BD is considered a robust predictor of outcome in trauma, and this has been demonstrated in different trauma populations and in different trauma systems over a timespan of 25 years. (310) BD has been found to predict outcomes in terms of transfusion requirements, cytokine response, MOF, and survival. (311-314) A critical threshold exists at or slightly above a BD of 6, where the probability of MOF and death rise exponentially. This threshold is widely used in trauma studies to dichotomise patients into groups with poor or good prognosis. (310,311) Lactate is another metabolic correlate of oxygen debt and severity of shock, (311) Few studies have compared lactate and BD as predictors of outcome in trauma, and data are conflicting. (314,315) Lactate may be elevated after excessive alcohol intake, administration of catecholamines and during decreased hepatic clearance, and these factors may confound the assessment of shock severity utilizing blood lactate as a metabolic correlate of shock, (311,316) and we therefore chose BE over lactate as a surrogate marker of shock.

Age. Age is a predictor for survival after trauma in the TRISS, TARN and NORMIT models. (288) We therefore included age as predictor for outcome in our statistical models. Age was included as a continuous variable.

Mechanism of injury and gender. In a retrospective study of 5363 trauma patients admitted to OUH Ullevål in the time period 2000–2006, mechanism of injury and gender did not contribute as independent predictors of outcome. (305) However, mechanism of injury is included in the TRISS model, and studies examining the effects of gender on outcome in trauma have yielded conflicting results, (317) with a reported possible interaction between age and gender. (304) We therefore chose to include these variables in our statistical modeling.

Statistical considerations

Data collection was designed as serial measurements in individual subjects, and our statistical analyses take advantage of this. In all aspects we strived to establish continuous dose–response relationships between HMGB1 concentrations and other variables, in

contrast to previous publications which have often converted continuous data into two or more categories. Multivariable analysis techniques were utilized to evaluate individual and simultaneous effects of several putative factors taken together.

Ethical aspects

Permission for the project was granted from the Regional Committee for Medical and Health Research Ethics (REC) and from the institutional Privacy Ombudsman for Research. The main ethical consideration was the recruitment of and blood sampling from individuals who were not able to consent. We therefore collaborated closely with the REC to achieve acceptable handling of this issue, which was balanced against the negative consequences arising for future patients if no research can be done in severely injured patients. During the first 24 h after admission a total volume of 128 ml blood was withdrawn from an individual trauma patient. In non-bleeding patients this represents a small fraction of the total blood volume, and in patients subject to massive transfusion this is minor compared to the large volume shift during bleeding and subsequent transfusion. It was concluded that blood sampling from patients not able to consent was acceptable if next of kin consented for them, and the patients themselves granted informed consent as soon as they were able. Consent from next of kin was sufficient in patients permanently unable to consent. For patients who were unable to consent until after they left OUH Ullevål, it was deemed sufficient to send information about the project and their right to withdraw. All patients who were unable to consent were checked against the Norwegian national biological research reservation registry before final inclusion. It turned out that only two of the originally enrolled patients did not want to participate, and consequently all sample material was destructed. Background, practical and theoretical aspects of the project have been continuously presented to and discussed with patients and their next of kin, and their interest in the project and eagerness to contribute has been overwhelming.

Summary of results

Paper I

We investigated HMGB1 concentration kinetics in serial plasma samples from 136 trauma patients with a broad spectrum of injuries and outcomes. Extracellular HMGB1 release in trauma patients demonstrated a biphasic pattern. An immediate phase with high concentrations of HMGB1 declining exponentially with a half-life of 26 min was seen in all patients. In the most severely injured patients the immediate decay phase was blunted by a second wave of HMGB1 release peaking 3–6 h after injury. Multivariable analyses showed that admission HMGB1 concentration was predicted by the degree of anatomical injury (NISS) alone. Second wave HMGB1 (area under the concentration curve [AUC] 3–6 h post-injury after subtraction of the exponentially declining immediate phase) was predicted by degree of physiological derangement (admission BE) and admission HMGB1 concentration. The total area under an exponential decay curve is proportional to the initial concentration. Consequently, immediate phase HMGB1 release in terms of both admission value and AUC lacked prognostic significance. However, the second wave HMGB1 3–6 h post-injury was a strong and highly significant predictor of outcome, rendering insignificant the effects of anatomical injury and physiological derangement on all outcome variables except maximum creatinine value.

Paper II

To the best of our knowledge, this is the first study to compare HMGB1 levels in simultaneously acquired arterial and venous samples from trauma patients. We found a close correlation between arterial and venous HMGB1 concentrations, surprisingly however arterial concentrations were found to be consistently lower than venous.

Previous reports have raised the concern that serum components may interfere with the detection of HMGB1 by ELISA. We therefore compared HMGB1 levels in simultaneously acquired serum and plasma samples from trauma patients. Additionally, we evaluated the suitability of western blot analysis as an alternative to ELISA in our patient population. The kinetic pattern of HMGB1 in parallel plasma and serum samples from the same individual was strikingly similar, however it displayed wide limits of agreement. Standard western blot analysis employing two diverse detection antibodies

failed to detect HMGB1 at clinically relevant concentrations in plasma and serum samples.

Adherence to strict protocol regarding pre-analytical conditions was a major priority throughout this research project, and we evaluated the impact of storing samples at room temperature for a variable time period prior to centrifugation, and of freeze / thaw cycles and long term storage at -80°C . HMGB1 concentrations were stable for approximately 3–6 h at room temperature but thereafter increased exponentially. In accordance with previous reports we confirmed that HMGB1 plasma levels are fairly stable in samples subjected to long time storage and freeze / thaw cycles.

Paper III

We explored IL-33 concentration kinetics in the same series of plasma samples employed in Paper I. In the population of 136 patients, 28 patients were IL-33 positive, and IL-33 was primarily detected in a small subset of severely injured patients (median ISS 42, $n = 24$). Peak IL-33 plasma levels were detected upon admission and declined below detection limit within 8 h after injury in all patients. IL-33 was not detected later in the clinical course despite several patients suffering from severe post-trauma complications. IL-33 decline obeyed first order kinetics with a calculated half life of 1.4 h. IL-33 positive patients had more widespread tissue injuries compared to IL-33 negative patients, calculated as both NISS and number of ISS body regions affected. IL-33 positive patients displayed increased incidence of early coagulopathy defined as $\text{INR} > 1.2$ and $\text{aPTT} > 35$ seconds, and had higher transfusion requirements.

Concentration kinetics of the IL-33 decoy receptor sST2 were measured in a subset of IL-33 positive and negative patients. We confirmed previous reports that sST2 is abundantly expressed in healthy controls, however a massive increase in plasma sST2 concentration was observed starting from around 2 h after injury and peaking within 32 h after injury, with higher peak concentrations seen in IL-33 positive patients.

In summary, we show a rapid but transient release of IL-33 after severe trauma and delayed release of the decoy receptor sST2 displaying an inverse kinetic relationship to IL-33. The early release of IL-33 after trauma with peak levels 0.5–2.5 h after injury indicates passive release of nuclear IL-33 rather than de novo synthesis, strengthening the notion that IL-33 has the potential to fulfill the role as an alarmin in human trauma and

shock. Conversely, the delayed release of the sST2 receptor peaking approximately 12-24 h after injury suggests de novo synthesis of this protein.

General Discussion

HMGB1 kinetics in trauma patients

In Paper I we explore high resolution HMGB1 kinetics in individual trauma patients. Peltz et al. were the first to explore the temporal pattern of HMGB1 early after trauma, and they reported peak levels 2 – 6 h post injury, which represented a striking 10 fold increase from their initial sample taken within 1 h after injury. (200) We report an inverse temporal pattern with peak levels at admission and a rapid decline in plasma concentration within 6 h after injury. Additionally, Peltz et al. reported HMGB1 plasma levels reaching almost 100 fold higher concentrations than what we have documented in a comparable trauma population.

Sustained HMGB1 levels at 24 h and up to several days after trauma have been reported in 3 previous studies. (174,200,318) Peltz et al. reported a median level of almost 200 ng/ml 24–136 h post injury, and this stands in sharp contrast to our findings, with the majority of patients having a HMGB1 plasma level < 12 ng/ml during this time interval. In a small population of blunt chest trauma patients (mean ISS 17) HMGB1 plasma concentrations increased gradually day 1–7 after trauma, reaching 74 ng/ml at day 7 after injury, (174) and similar results were found in another study of blunt chest trauma. (318) The temporal pattern of HMGB1 is remarkably consistent in our heterogeneous population of trauma patients, and sustained levels were not seen in any patients. Differences in kinetics and concentration values reported in these studies might be attributed to pre-analytical conditions or the use of different detection systems. This will be discussed below.

A recent study by Namas et al. reported HMGB1 plasma kinetics within 24 h after injury which are in accordance with the findings in our study, however they did not report absolute concentration values or time from injury. (45)

First order kinetics – potential decay mechanisms

The first-order decay, with a HMGB1 concentration half-life of 26 min observed in the immediate phase was consistent across the whole population, despite differences in injury severity and admission BE. Additionally, several patients underwent damage control surgery and aggressive resuscitation without any obvious change in decay rate. This is counterintuitive, since trauma patients receive various amounts of fluids, drugs and

transfusions one could expect this to impact decay rate unevenly. The mechanism behind the rapid first order decay of HMGB1 concentration during the immediate phase is a matter of speculation, and several mechanisms can be responsible.

It is well documented that HMGB1 is a sticky protein, and interaction with other proteins of intracellular or extracellular origin could mask HMGB1 detection by ELISA. (319) HMGB1 is reported to bind to haptoglobin, extracellular DNA and histones, all are proteins that could potentially be in abundance in the bloodstream after trauma. (157,320) Rapid proteolytic cleavage is another plausible explanation, but HMGB1 clearance in serum by proteases has been shown to be much slower than what we have reported in our study. (180) HMGB1 binding to soluble RAGE is another possible explanation, but soluble RAGE levels are reported to peak at approximately 2000 pg/mL early after severe trauma in humans. (321,322)

Biphasic release and correlation to outcome

The key discovery in our study is the identification of a second wave HMGB1 occurring in certain patients with severe tissue injury and physiological derangement. This is a distinctly novel finding, and raises important questions regarding cellular source, release mechanism and isoforms.

It is tempting to speculate that the different relationships of admission HMGB1 and second wave to outcome might be caused by different HMGB1 isoforms. (98) Non-acetylated fully reduced HMGB1 is constitutively present in the nucleus of all cells during homeostasis and will inevitably be released due to mechanical trauma or necrosis. Consequently, it is reasonable to expect that the immediate HMGB1 release after injury constitutes mainly fully reduced HMGB1. (150) This isoform mediates chemotaxis but not cytokine induction, which is compatible with admission HMGB1's lack of correlation to outcome in our multivariable statistical models. Cohen et al. had reported a strong bivariable correlation between admission HMGB1 value and outcome by the means of organ failure and ventilator free days. (201) In view of these findings it was somewhat surprising that admission HMGB1 in our population was not a significant predictor to any outcome variable. We believe that the correlation between admission HMGB1 and outcome in bivariable analysis reflects early HMGB1 release as being a proxy of tissue injury severity rather than being a causal factor.

Second-wave HMGB1 was the dominant predictor for our primary outcome measure, rendering all other putative variables insignificant except for a small contribution from age. The results from our statistical modeling suggest that second wave HMGB1 is a product of injury severity and shock, acting as mediator of the combined effects of anatomical injury and physiological derangement. The second wave could represent passive release due to ongoing cell death, or active secretion due to cellular stress, or a combination of the two. Isoform analyses providing acetylation status would be required in order to distinguish whether the second wave is a result of active secretion or passive release. Despite being highly speculative, I propose that the strong correlation of second-wave AUC₃₋₆ to outcome is not simply reflecting prolonged release of HMGB1 in general terms, but rather a switch in HMGB1 redox state from reduced to disulfide form. Disulfide HMGB1 is a TLR4 and RAGE ligand resulting in sterile inflammation and injury amplification. (118,167,170)

HMGB1 will be rapidly oxidized outside the cell due to the oxidative environment, and consequently one would expect initially released fully reduced HMGB1 to be oxidized to disulfide HMGB1. The common use of 100% oxygen during resuscitation in severe trauma and shock would further facilitate the formation of disulfide bridges. In vitro oxidation of fully reduced HMGB1 in human serum was studied recently by in situ protein NMR spectroscopy. Fully reduced HMGB1 was rapidly oxidized to disulfide HMGB1 with first order kinetics and a half-life of 17 min. (180)

Second wave HMGB1 – cellular source

Neutrophils and monocytes migrate into traumatized tissue where they become activated by the pro-inflammatory environment, (177) and the second wave release documented in our study is in agreement with the expected kinetics for active HMGB1 secretion by monocytes. (119) However, it is well documented that also non-immune parenchymal cells such as hepatocytes, neurons, endothelial cells and smooth muscle cells contribute to active HMGB1 secretion. (92,114,116,117)

A few experimental models have addressed specific organs as source of HMGB1 release in trauma and shock. In a model of bilateral femur fracture and hemorrhagic shock in mice, acute remote lung injury was mediated by release of HMGB1 from the gut following activation of intestinal epithelial TLR4 receptors. (116) Mice in which HMGB1 was genetically selectively deleted from the intestinal epithelium had lower systemic

HMGB1 levels and were protected from acute lung injury. HMGB1 isoforms were not investigated. The same group had previously documented increased HMGB1 levels as early as 2 h after hemorrhagic shock and resuscitation in mice in the serum, lung and liver, but not in the kidney. (323) Pulmonary endothelial cells are activated at a very early stage of lung inflammation and play a critical role in the development of acute lung injury after trauma and shock. (148,324) An experimental model of hemorrhagic shock in mice have demonstrated NLRP3 inflammasome activation in lung endothelial cells, promoting pyroptosis. (325) Pulmonary endothelial cells undergoing pyroptosis may consequently be a source of second-wave HMGB1.

In the context of hemorrhagic shock, decreased perfusion gives rise to hypoxia and concomitant cellular stress in peripheral tissues. Interestingly, hyperacetylated HMGB1 has been detected in serum of mice subjected to warm hepatic I/R injury as early as 1 hour after reperfusion. (114) Hypoxia was believed to be the triggering event, and hyperacetylation of HMGB1 was also induced in hepatocytes exposed to hypoxia in vitro (1% O₂, 5% CO₂ and 94% N₂). Hyperacetylated HMGB1 increased in the hepatocyte cell culture supernatants as early as 3 h after the hypoxic exposure. (114) Negative BE was a strong predictor of second-wave plasma HMGB1 level in our trauma population, and we speculate that hypoxia might be a trigger for active HMGB1 secretion contributing to the second wave also in our trauma patients. Taken together these experimental models underpin the importance of HMGB1 release from liver, lung and gut epithelial cells in the pathogenesis of organ failure after trauma and shock.

Recent evidence has also revealed that resting platelets contain substantial amounts of HMGB1. The intracellular HMGB1 is translocated to the cell surface upon platelet activation. (326,327) Platelets are highly activated in trauma and shock and could conceivably be an important source of HMGB1.

HMGB1 detection – pre-analytical conditions and sample material

“HMGB1 is one of the pillars of the immune system, so whenever you try use the immune system to create reagents against HMGB1 you are asking the immune system to make antibodies against one of the main cogs in the immune machinery.” Marco Bianchi

At the time of inception of this research project, 3 previous studies by Yang, Cohen and Peltz respectively had investigated HMGB1 in trauma patients. (174,200,201) The studies

by Cohen and Peltz both relied on plasma samples and standard ELISA technique, and the study by Yang et al. utilized serum samples and quantitative western blot. In order to be able to compare our results with previous work in the field, we chose to use the HMGB1 ELISA kit by IBL, utilized by both Cohen and Peltz. Several commercial ELISAs for HMGB1 exists, however the IBL kit identical to that originally designed by Shino-Test corporation is regarded as a the gold standard by most experts in the field. Additionally, its validity had been tested on both plasma and serum samples. (292).

In the context of trauma and hemorrhagic shock, there is an abundance of intracellular proteins in the plasma due to necrosis and loss of endothelial integrity and a massive activation of several humoral cascade systems. Consequently it's reasonable to anticipate increased noise or interference with antibody detection in any immunoassay. A recent study of patients with hematological disorders reported that HMGB1 concentrations measured with ELISA were about 10-fold lower in serum than in plasma. (328) In contrast, in two different studies of healthy individuals, serum yielded higher HMGB1 concentrations than EDTA plasma. (292,329) However, similar studies from trauma patients were lacking when we started this research project and we reasoned that interference or noise with the ELISA assay could conceivably depend on disease state and sample material. In our trauma samples, HMGB1 serum concentrations were found to be both higher and lower compared with plasma concentrations, with no specific pattern. However, we found a strong correlation between concentration measurements in plasma and serum, consequently the comparison of kinetic profiles in serum and plasma samples should be valid despite wide limits of agreement. Thus, we believe that serum and plasma samples cannot be directly compared but that they are equally valid in the study of HMGB1 kinetics in trauma patients. We were somewhat surprised that arterial concentrations were found to be consistently lower than venous. However, we relied primarily on plasma from arterial samples, in particular from patients with severe and critical injuries since they routinely get an arterial line. The first blood sample after admission is also routinely obtained through arterial puncture in our trauma centre.

HMGB1's propensity to form complexes with other soluble proteins raises an important question regarding to which extent the ELISA detects complexed HMGB1. Interference caused by IgG and IgM antibodies has been documented in serum and plasma from Rheumatoid arthritis and SLE patients and in healthy donors. (319,330) Incubation of recombinant human HMGB1 with IL-1 β , LPS or specific antibodies

significantly reduces the amount of HMGB1 detected by the IBL ELISA. Interestingly, interference can be largely reversed with perchloric acid (PCA) treatment of the sample. (331) PCA can dissociate protein-protein complexes due to its acidic properties. This approach has been tested on serum samples from critically ill sepsis patients, and PCA treatment of the sample prior to ELISA increased HMGB1 detection significantly. (331) In a novel approach utilizing electrophoretic mobility shift, a highly sensitive and specific HMGB1 assay has been developed. (332) The assay relies on detection of high affinity complexes formed between HMGB1 and radio-labeled hemicatenated DNA loops. Unfortunately, the assay is complex, has a low throughput and does not resolve the detection of HMGB1 isoforms.

Western blotting is a laborious and semi-quantitative method, however the sodium dodecyl sulphate-mediated denaturation step theoretically enables the method to detect HMGB1 complexed with other proteins due to dissociation of these complexes. Indeed, western blot has been reported to be more sensitive than ELISA when applied on serum and plasma samples from patients with Rheumatoid arthritis and SLE. (319,330) However, in our hands western blotting was not able to detect clinically relevant HMGB1 levels in plasma or serum from trauma patients. To determine the lowest level of detection by western blot, we diluted recombinant HMGB1 in normal serum and plasma, and were only able to detect HMGB1 bands at concentrations equal to or above 1000 ng/mL, 10–50 times the maximum level measured by ELISA in our clinical samples.

Monoclonal antibodies are often less effective for detecting proteins in denatured state or altered conformation compared to polyclonal antibodies that detect multiple epitopes. However, the application of a polyclonal antibody (Abcam cat no. 18256) did not increase sensitivity with our protocol. In the seminal work by Wang et al., the monoclonal antibody 2G7 was applied successfully in western blotting of serum samples from sepsis patients with a detection limit of 50 pg. (93) We cannot explain these discrepancies, however enrichment approaches such as sample concentrators with semipermeable membranes with specific molecular weight cut-offs can be used to enhance sensitivity. (329)

Time from sampling to centrifugation and storage temperature are crucial aspects in order to protect sample integrity, yet are often overlooked or not reported in clinical research. (333) A study by Lehner et al. reported that storage of blood samples at room

temperature (25 °C) often led to higher HMGB1 values in sera after 6 h, however with a considerable individual variability. (292) In line with this we report that HMGB1 plasma levels increase exponentially after 6 h in blood samples that are stored in room temperature (25 °C) prior to centrifugation. The results were consistent, however notably HMGB1 concentration from the two patients with highest initial value did not change during the 24 h storage. These samples were among the earliest samples after injury, and one might speculate that the stability of the sample is redox sensitive and therefore may also vary according to predominant isoform in the sample.

In a previous study, HMGB1 concentrations in serum samples stored in room temperature (25 °C) for up to 7 days after centrifugation or subjected to freeze/thaw cycles were remarkably stable. (292) In accordance with this we found a strong linear correlation between concentrations in originally analyzed plasma and plasma subjected to long-term storage and freeze/thaw cycles. The wide limits of agreement in our study may probably in part be caused by inter-assay variability, which was avoided in the study by Lehner et al.

Ongoing degradation in the sample due to protease activity or interaction with other proteins may depend on disease state, and may also vary according to predominant isoform. The redox state is shown to change quickly under in vitro conditions, (180) and since isoform analyses might be instrumental to future HMGB1 research, handling of samples should follow strict protocol, especially prior to centrifugation. HMGB1 detection and stability in a sample is context sensitive, and therefore methodological studies should be performed for all disease states where HMGB1 is investigated as a potential biomarker.

IL-33 kinetics in trauma and shock

The role of the alarmin IL-33 and its receptor sST2 is poorly understood in trauma and hemorrhagic shock. Experimental work on IL-33 has mainly focused on its role in allergic and airway disease, (334) and most studies have been performed in rodents. In contrast to HMGB1, IL-33 tissue expression patterns differ markedly between mouse and humans, therefore the degree to which results from mechanistic studies in mice translate to human disease is uncertain.

In Paper III we explore systemic IL-33 levels in human trauma. Our results differ markedly from the recent work by Xu et al. (245) We found peak IL-33 levels at admission, corresponding to 0:29–2:27 h after injury, and then a rapid decrease following first order kinetics with a calculated half-life of 1.4 h. Xu et al. did not report time from injury, however peak values were also found in their admission samples, and the authors speculate that this coincides approximately with 6 h post injury. Interestingly, Xu et al. reported sustained IL-33 levels for several days after injury in the majority of patients, in contrast to our results where IL-33 was never detected beyond 12 h post injury. These divergent results cannot be explained by differences in injury severity as IL-33 was detected in all trauma patients by Xu et al. (average ISS 20), as opposed to the dichotomic distribution in our trauma population where most patients was IL-33 negative and a small group of severely injured patients (mean ISS 42) was IL-33 positive. Additionally, we did not detect IL-33 in healthy controls opposed to an average baseline of 23 pg/ml in the study by Xu et al., suggesting differences in either sensitivity or background noise in the immunoassays applied in our respective studies. In agreement with Xu et al. we documented a delayed release of sST2 peaking approximately 12–24 h after injury in a small subset of patients.

IL-33 detection

Poor agreement or total absence of relation between results have been shown between commercially available ELISAs. (335,336) Constitutive levels of IL-33 in healthy controls have been determined in a number of studies and show a wide range from undetectable or low (1–10 pg/mL) (337,338) to several hundred picograms or even nanograms pr mL. (339-342) In a recent study, an ELISA validated for sera was unable to detect IL-33 in healthy controls, despite showing high sensitivity and specificity in recombinant IL-33 spike-in serum samples. (336) It might be speculated that the IL-33 positive controls and the sustained levels in trauma patients in the Xu et al. study represents a problem with background noise in the Millipore assay. Conversely, the low or absent levels in our study could be due to poor sensitivity of the BioRad assay.

Poor sensitivity has been shown in several IL-33 ELISAs due to interaction with factors in serum, possibly globulins, albumin and sST2 that interfere with IL-33 quantification in a dose dependent manner. (336) In a recent study of IL-33 regulation, recombinant IL-33 was found to be rapidly oxidized to its disulfide form after exposure to

serum or cell culture media (> 90% conversion by 4 h). (258) Two different ELISA assays, including the Millipore assay used by Xu et al., were validated in this study. They were found to primarily detect DSB IL-33, and the DSB IL-33 signal was not affected by the presence of ST2. However, the reduced IL-33 signal diminished by the presence of ST2 in a dose dependent manner, and the authors conclude that these assays detect “free” IL-33, either non-ST2 bound or DSB IL-33 which is non-ST2 binding. To the best of our knowledge similar validation studies are not done on the Bio-Rad assay used in our study. The Bio-Rad assay has antibodies directed towards an unspecified epitope on the c-terminal part of IL-33, and IL-33 cleavage by neutrophils or mast cell proteases or extracellular oxidation to DSB IL-33 could conceivably affect antibody detection. In the light of this, we cannot rule out that our Bio-Rad assay detects only a fraction of the bioavailable IL-33

Differences in pre-analytical conditions between studies could introduce major confounding factors since differences in storage temperature and time to centrifugation might conceivably affect redox state of IL-33 in plasma and serum samples. Additionally, quantification of IL-33 could conceivably be affected by the predominant IL-33 isoform measured versus the IL-33 used as reference standard. The IL-33 concentration measurements in our study may thus be compromised by the obvious problems with validation of IL-33 ELISA systems, and the lack of specific IL-33 isoform detection. Our supplementary experiments using immunoprecipitation and western blotting could not detect IL-33 in normal plasma or in patients with high IL-33 plasma concentration as determined by ELISA, even when reducing the background noise by cleaning out a large fraction of albumin and immunoglobulin from samples. The sustained levels seen days after injury in the Xu study are likely to be DSB IL33, since reduced IL-33 is rapidly oxidized or bound to sST2. Conversely, the earlier samples in our study might be comprised of reduced IL-33 not yet bound to sST2.

IL-33 source

We consider it likely that the source of systemically released IL-33 is the vascular endothelium, which in humans shows strong constitutive expression of IL-33. (254,334) Endothelial cells as primary source is in accordance with the low IL-33 levels detected, as these cells represent a small tissue volume. In contrast, HMGB1 can be released from all nucleated cells which represents a much larger tissue volume.

In a study of patients undergoing liver resection, surgery resulted in a 5-fold increase in post-operative serum IL-33 compared with pre-operative levels. (343) In the same study, IL-33 protein expression was up-regulated and IL-33 released by liver sinusoidal endothelial cells as early as 1 hour after 60 min of warm ischemia in mice. In light of these data the liver could potentially be an import source of IL-33 in human trauma, which coincides with the observation that the majority of our IL-33 positive patients (88%) had injuries in the abdominal region. Interestingly, patients with major isolated head injury were all IL-33 negative, supporting that that IL-33 is not expressed strongly in human brain despite abundant expression in mice. (334)

CD45+ hematopoietic cells, such as macrophages and mast cells, have been proposed to represent a major cellular source of IL-33, however convincing evidence of inducible protein expression in leukocytes has not yet been provided. (334) In our study we did not detect IL-33 beyond 8 h after injury, suggesting that systemic release of IL-33 in trauma is related to acute cellular injury. Adding to this, microarray data of the circulating leukocyte transcriptome after severe trauma proved no increase in IL-33 expression, (29) however it is unclear whether the systemic circulation serves as an accurate mirror of the pro-inflammatory environment in the peripheral tissues. In conclusion, early release (peak levels 0:29–2:27 h after injury) and the association with injury severity, fits well with the notion that IL-33 is constitutively available and rapidly released due to necrosis.

Concluding remarks

Overpowering systemic inflammation with ensuing multiple organ failure is a common and deadly feature after severe trauma, as in sepsis and other critical conditions. In this thesis, new knowledge has been provided regarding high resolution kinetics of the prototypical alarmin HMGB1 and the novel alarmin IL-33 in trauma and shock.

- We present the first report on high time resolution HMGB1 kinetics in individual patients with time locked to the injury event. This approach revealed two separate HMGB1 release phases, with very different predictive ability for outcome.
- There is a close correlation between HMGB1 concentrations in plasma and serum samples and in arterial vs. venous samples, however agreement is poor. Arterial concentrations were found to be consistently lower than venous.

- HMGB1 is relatively stable in plasma and serum during long time storage and freeze thaw cycles. Adherence to strict protocols regarding pre-analytical conditions is advisable as HMGB1 levels increase exponentially after 3–6 h if blood samples are stored in room temperature prior to centrifugation.
- IL-33 is released early after trauma in severely injured patients, and there is a delayed compensatory release of the sST2 decoy receptor. Based on previous work and on our data, the cellular source of systemic IL-33 is likely to be the vascular endothelium. Biological interpretation of IL-33 studies in trauma and shock is severely compromised by the lack of validated ELISAs that can distinguish IL-33 isoforms.

The innate immune response to cellular injury and stress is fundamental and uniform. Consequently, the biological mechanisms we describe may be relevant in a large number of diseases, including sepsis, burn injury, surgical stress, pancreatitis, and ischemia-reperfusion injury. Our primary interest is the causal mechanisms of systemic inflammation in humans; the reason why we have chosen trauma as a model is simply that the exact moment of the triggering event is known. Although our primary intent was not to search for biomarkers, our manuscript may provide a generic recipe for evaluation of biomarkers. Scientists often study the concentrations of molecules at the time the patient arrives in hospital. Our study demonstrates that this may be completely misleading in highly dynamic situations.

Limitations

Several limitations for this work have been mentioned in the specific papers. The study has a relatively small size and is a single center study. Analyses regarding mechanism of injury are uncertain due to the low number of penetrating injuries. NISS was used as a summary measure of anatomical injury, but it lacks a direct relationship to the total volume of damaged tissue. The relationship between NISS and subsequent immunological activation is therefore complex. Since the focus in the present study was inflammation and remote organ failure, patients with major head trauma had to be excluded from the analysis of our major outcome measure; ventilator free days. Patients with severe head trauma spend substantial time on ventilator support due to the head injury per se, independent of inflammation or remote organ failure. Ventilator free days can be considered a combined measure of morbidity and mortality, but how this translates to mortality is uncertain, and can vary between different countries and hospitals. Our

multivariable models were based on linear relationships despite biological systems being inherently nonlinear. Non-linear statistical methods could therefore conceivably generate more optimal models.

Future perspectives

HMGB1 is a promising biomarker in diverse diseases. The predictive value of second wave of HMGB1 in our trauma cohort we believe is a result of HMGB1 being a key player in the pathogenesis of unbridled systemic inflammation, and not only a proxy for injury and shock severity. In order to translate this knowledge to the clinic there is an urgent need to develop antibodies that enable isoform-specific ELISAs that can facilitate high throughput and expedient analyses.

In the context of trauma we believe a biomarker has limited value since treatment options are mainly organ supportive, and intensive care is largely guided by clinical surrogates. Targeted therapy does not currently exist, and we would like to draw attention to the potential therapeutic implications of our findings. HMGB1 antagonistic therapy has proven successful in a wide range of animal models of sepsis, trauma and hemorrhagic shock. (239) The kinetics of HMGB1 release identified in our study indicate a clinically accessible time window for early administration of HMGB1 antagonists after severe trauma in humans. This could potentially attenuate the development of devastating systemic inflammation and secondary remote organ failure.

IL-33 is a novel alarmin in trauma, however mechanistic studies in murine models have limited translational value due to marked differences in expression pattern between species. Interpretation of IL-33 studies in humans is severely hampered by the lack of validated ELISAs and consequently divergent results. In order for the field to move forward there is an urgent need for high throughput and reliable immunoassays that can detect different IL-33 isoforms.

Appendix

In the autumn of 2017, the University of Liverpool launched an internal investigation of research misconduct involving Daniel J. Antoine at the proteomics facility. Daniel Antoine became a superstar in the HMGB1 research community when he and colleagues published quantitative mass spectrometry (MS) data on HMGB1 isoforms in liver disease. (344) In what was considered groundbreaking work, Antoine et al. concluded that acetylated disulfide HMGB1 outperformed all other biomarkers, including total HMGB1 levels, in predicting clinical outcome for acetaminophen-intoxicated patients. This paper sparked off an intensive research effort into the role of HMGB1 isoforms in diverse disease states. Antoine and coworkers had apparently developed a robust MS assay that could reliably quantify HMGB1 isoforms with respect to both redox and acetylation state in a single sample. Researchers from all over the world reached out to Antoine in order to get their clinical or experimental samples analyzed with respect to HMGB1 isoform profile. This generated dozens of HMGB1 papers published in highly influential journals. Antoine is a co-author of 103 papers on HMGB1 biology, however 28 of these are reviews containing no primary data (Ulf Andersson personal communication). In the ongoing investigation several of Antoine's papers have been revealed to contain some form of manipulated data. There is speculation that Antoine has provided spiked samples for MS analysis and that in several instances he has generated false data in the void of actual analyses.

In the spring of 2016 we sent samples from our trauma patients to Liverpool for MS analysis. We hypothesized that the immediate phase of HMGB1 release in our samples primarily consisted of non-acetylated fully reduced HMGB1, and the second wave primarily of acetylated disulfide HMGB1. We discussed our idea with Antoine, who was keen to perform MS analysis on a subset of our samples. He provided isoform data from 91 samples taken from 16 patients, which correlated with our ELISA data and apparently confirmed our hypothesis. We therefore wrote a paper based on both ELISA and MS data describing HMGB1 isoform kinetics in trauma patients. The paper was accepted for publication by Critical Care Medicine (CCM) in late March 2018. While the paper was in page proof stage it came to our attention that Antoine was the subject of an internal investigation due to concerns about the reproducibility and integrity of data from one of his research projects. We immediately contacted CCM's chief editor in order to freeze

publication of our paper as a precautionary measure. We then performed a statistical analysis of the data set that Antoine had provided to us, in an effort to assess the validity of the data. The data set consisted of 91 HMGB1 concentrations given with one decimal digit. We found that the distribution of the last digits was highly unlikely: none of them were “0”, two were “9”, whereas 23 were “1” (Pearson’s chi-squared test $p < 10^{-5}$, compared to flat distribution of last digit). The dataset was sent to the investigators in Liverpool, and they confirmed that it had been fabricated by Antoine in the void of any actual analysis. We were deeply shocked by this discovery, and it represented a major setback for our work. Luckily, the publication of our paper could be stopped, and we later submitted a revised version which was accepted by Critical Care Medicine.

The implications and extent of Antoine’s misconduct are currently not fully known, but all his HMGB1 isoform analyses should be regarded highly suspicious until proven otherwise. To date (December 2019) 10 of his papers are recommended for retraction and many more retractions are anticipated. It should however be noted that Antoine was never involved in any fundamentally basic discoveries regarding HMGB1 biology. His scientific misconduct concerns quantitative aspects of assessments of acetylated or redox-modified HMGB1 isoforms in clinical or preclinical samples. It does not alter our general understanding of HMGB1 biology regarding modifications (acetylation, phosphorylation, methylation, redox changes) or the functional consequences or receptor usage caused by these conversions.

References

1. Medzhitov R. Origin and physiological roles of inflammation. *Nature*. 2008 Jul 24;454(7203):428–35.
2. Nathan C. Neutrophils and immunity: challenges and opportunities. *Nat Rev Immunol*. 2006 Mar;6(3):173–82.
3. Barton GM. A calculated response: control of inflammation by the innate immune system. *J Clin Invest*. 2008 Feb;118(2):413–20.
4. Jain A, Pasare C. Innate Control of Adaptive Immunity: Beyond the Three-Signal Paradigm. *J Immunol*. 2017 May;198(10):3791–800.
5. Netea MG, Latz E, Mills KHG, O'Neill LAJ. Innate immune memory: a paradigm shift in understanding host defense. *Nat Immunol*. 2015 Jul;16(7):675–9.
6. Burnet F, Fenner F. *The Production of Antibodies*. Second edition. Melbourne; Macmillan; 1949.
7. Ribatti D. Peter Brian Medawar and the discovery of acquired immunological tolerance. *Immunol Lett*. 2015 Oct 1;167(2):63–6.
8. Billingham RE, Brent L, Medawar PB. Actively acquired tolerance of foreign cells. *Nature*. 1953 Oct;172(4379):603–6.
9. Bretscher P, Cohn M. A theory of self-nonsel self discrimination. *Science*. 1970 Sep;169(3950):1042–9.
10. Lafferty KJ, Cunningham AJ. A new analysis of allogeneic interactions. *Aust J Exp Biol Med Sci*. 1975 Feb;53(1):27–42.
11. Janeway CAJ. Approaching the asymptote? Evolution and revolution in immunology. *Cold Spring Harb Symp Quant Biol*. 1989;54 Pt 1:1–13.
12. Manson J, Thiemermann C, Brohi K. Trauma alarmins as activators of damage-induced inflammation. *Br J Surg*. 2011 Dec 22;99 Suppl 1(S1):12–20.
13. Matzinger P. Tolerance, danger, and the extended family. *Annu Rev Immunol*. 1994;12:991–1045.
14. Matzinger P. The danger model: a renewed sense of self. *Science*. 2002 Apr 12;296(5566):301–5.
15. Hirsiger S, Simmen H-P, Werner CML, Wanner GA, Rittirsch D. Danger Signals Activating the Immune Response after Trauma. *Mediators Inflamm*. 2012;2012(2):1–10.
16. Yang H, Wang H, Ju Z, Ragab AA, Lundbäck P, Long W, et al. Circulating mitochondrial DAMPs cause inflammatory responses to injury. *Nature*. 2015 Jan 12;464(7285):104–7.
17. Rider P, Voronov E, Dinarello CA, Apte RN, Cohen I. Alarmins: Feel the Stress. *J Immunol*. 2017 Feb 6;198(4):1395–402.
18. Bianchi ME. DAMPs, PAMPs and alarmins: all we need to know about danger. *J Leukoc Biol*. 2006 Oct 10;81(1):1–5.

19. Takeuchi O, Akira S. Pattern recognition receptors and inflammation. *Cell*. 2010 Mar 19;140(6):805–20.
20. Lemaitre B, Nicolas E, Michaut L, Reichhart JM, Hoffmann JA. The dorsoventral regulatory gene cassette *spatzle/Toll/cactus* controls the potent antifungal response in *Drosophila* adults. *Cell*. 1996 Sep;86(6):973–83.
21. Zarembek KA, Godowski PJ. Tissue expression of human Toll-like receptors and differential regulation of Toll-like receptor mRNAs in leukocytes in response to microbes, their products, and cytokines. *J Immunol*. 2002 Jan;168(2):554–61.
22. Shaw PJ, Lamkanfi M, Kanneganti T-D. NOD-like receptor (NLR) signaling beyond the inflammasome. *Eur J Immunol*. 2010 Mar;40(3):624–7.
23. Pisetsky DS. The effectors of innate immunity: DAMPs, DAMEs, or DIMEs? *Arthritis Res Ther*. 2013;15(5):123.
24. Yang D, Oppenheim JJ. Antimicrobial proteins act as “alarmins” in joint immune defense. *Arthritis Rheum*. 2004 Nov;50(11):3401–3.
25. Yang D, la Rosa de G, Tewary P, Oppenheim JJ. Alarmins link neutrophils and dendritic cells. *Trends Immunol*. 2009 Nov;30(11):531–7.
26. Harris HE, Raucchi A. Alarmin(g) news about danger: workshop on innate danger signals and HMGB1. *EMBO Rep*. 2006 Aug;7(8):774–8.
27. Vanden Berghe T, Linkermann A, Jouan-Lanhouet S, Walczak H, Vandenabeele P. Regulated necrosis: the expanding network of non-apoptotic cell death pathways. *Nat Rev Mol Cell Biol*. 2014 Feb;15(2):135–47.
28. Kaczmarek A, Vandenabeele P, Krysko DV. Necroptosis: the release of damage-associated molecular patterns and its physiological relevance. *Immunity*. 2013 Feb;38(2):209–23.
29. Xiao W, Mindrinos MN, Seok J, Cuschieri J, Cuenca AG, Gao H, et al. A genomic storm in critically injured humans. *J Exp Med*. 2011 Dec 19;208(13):2581–90.
30. Raymond SL, Holden DC, Mira JC, Stortz JA, Loftus TJ, Mohr AM, et al. Microbial recognition and danger signals in sepsis and trauma. *Biochim Biophys Acta Mol Basis Dis*. 2017 Oct;1863(10 Pt B):2564–73.
31. Vos T, Barber RM, Bell B, Bertozzi-Villa A. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2015 Aug 22;386(9995):743–800.
32. Trunkey DD. Trauma. Accidental and intentional injuries account for more years of life lost in the U.S. than cancer and heart disease. Among the prescribed remedies are improved preventive efforts, speedier surgery and further research. *Sci Am*. 1983 Aug;249(2):28–35.
33. Sobrino J, Shafi S. Timing and causes of death after injuries. *Proc (Bayl Univ Med Cent)*. 2013 Apr;26(2):120–3.
34. Pfeifer R, Tarkin IS, Rocos B, Pape H-C. Patterns of mortality and causes of death in polytrauma patients—Has anything changed? *Injury*. 2009 Sep;40(9):907–11.

35. Gunst M, Ghaemmaghami V, Gruszecki A, Urban J, Frankel H, Shafi S. Changing epidemiology of trauma deaths leads to a bimodal distribution. *Proc (Bayl Univ Med Cent)*. 2010 Oct;23(4):349–54.
36. Baker CC, Oppenheimer L, Stephens B, Lewis FR, Trunkey DD. Epidemiology of trauma deaths. *Am J Surg*. 1980 Jul;140(1):144–50.
37. Dewar DC, Tarrant SM, King KL, Balogh ZJ. Changes in the epidemiology and prediction of multiple-organ failure after injury. *J Trauma Acute Care Surg*. 2013 Mar;74(3):774–9.
38. Fröhlich M, Lefering R, Probst C, Paffrath T, Schneider MM, Maegele M, et al. Epidemiology and risk factors of multiple-organ failure after multiple trauma: an analysis of 31,154 patients from the TraumaRegister DGU. *J Trauma Acute Care Surg*. 2014 Apr;76(4):921–7.
39. Soreide K, Krüger AJ, Vårdal AL, Ellingsen CL, Søreide E, Lossius HM. Epidemiology and Contemporary Patterns of Trauma Deaths: Changing Place, Similar Pace, Older Face. *World J Surg*. 2007 Sep 25;31(11):2092–103.
40. Probst C, Zelle BA, Sittaro NA, Lohse R, Krettek C, Pape HC. Late death after multiple severe trauma: when does it occur and what are the causes? *J Trauma*. 2009 Apr;66(4):1212–7.
41. Lord JM, Midwinter MJ, Chen Y-F, Belli A, Brohi K, Kovacs EJ, et al. The systemic immune response to trauma: an overview of pathophysiology and treatment. *Lancet*. 2014 Oct 18;384(9952):1455–65.
42. Dewar D, Moore FA, Moore EE, Balogh Z. Postinjury multiple organ failure. *Injury*. 2009 Sep;40(9):912–8.
43. Napolitano LM, Ferrer T, McCarter RJ, Scalea TM. Systemic inflammatory response syndrome score at admission independently predicts mortality and length of stay in trauma patients. *J Trauma*. 2000 Oct;49(4):647–52.
44. Jastrow KM, Gonzalez EA, McGuire MF, Suliburk JW, Kozar RA, Iyengar S, et al. Early Cytokine Production Risk Stratifies Trauma Patients for Multiple Organ Failure. *J Am Coll Surg*. 2009 Sep 1;209(3):320–31.
45. Namas RA, Vodovotz Y, Almahmoud K, Abdul-Malak O, Zaaqoq A, Namas R, et al. Temporal Patterns of Circulating Inflammation Biomarker Networks Differentiate Susceptibility to Nosocomial Infection Following Blunt Trauma in Humans. *Ann Surg*. 2016 Jan;263(1):191–8.
46. Maier B, Lefering R, Lehnert M, Laurer HL, Steudel WI, Neugebauer EA, et al. Early versus late onset of multiple organ failure is associated with differing patterns of plasma cytokine biomarker expression and outcome after severe trauma. *Shock*. 2007 Dec;28(6):668–74.
47. Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest*. 1992 Jun;101(6):1644–55.
48. Churpek MM, Zdravetz FJ, Winslow C, Howell MD, Edelson DP. Incidence and Prognostic Value of the Systemic Inflammatory Response Syndrome and

- Organ Dysfunctions in Ward Patients. *Am J Respir Crit Care Med*. 2015 Oct 15;192(8):958–64.
49. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016 Feb 23;315(8):801–10.
 50. Seymour CW, Liu VX, Iwashyna TJ, Brunkhorst FM, Rea TD, Scherag A, et al. Assessment of Clinical Criteria for Sepsis: For the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016 Feb 23;315(8):762–74.
 51. Minne L, Abu-Hanna A, de Jonge E. Evaluation of SOFA-based models for predicting mortality in the ICU: A systematic review. *Crit Care*. 2008;12(6):R161.
 52. Dewar DC, White A, Attia J, Tarrant SM, King KL, Balogh ZJ. Comparison of postinjury multiple-organ failure scoring systems. *J Trauma Acute Care Surg*. 2014 Oct;77(4):624–9.
 53. Royston P, Altman DG, Sauerbrei W. Dichotomizing continuous predictors in multiple regression: a bad idea. *Stat Med*. 2006 Jan 15;25(1):127–41
 54. Baue AE. Multiple, progressive, or sequential systems failure. A syndrome of the 1970s. *Arch Surg*. 1975 Jul;110(7):779–81.
 55. Eiseman B, Beart R, Norton L. Multiple organ failure. *Surg Gynecol Obstet*. 1977 Mar;144(3):323–6.
 56. Sauaia A, Moore EE, Johnson JL, Chin TL, Banerjee A, Sperry JL, et al. Temporal trends of postinjury multiple-organ failure. *J Trauma Acute Care Surg*. 2014 Mar;76(3):582–93.
 57. Minei JP, Cuschieri J, Sperry J, Moore EE. The changing pattern and implications of multiple organ failure after blunt injury with hemorrhagic shock. *Crit Care Med*. 2012 Apr;40(4):1129–1135.
 58. de Kneegt C, Meylaerts SAG, Leenen LPH. Applicability of the trimodal distribution of trauma deaths in a Level I trauma centre in the Netherlands with a population of mainly blunt trauma. *Injury*. 2008 Sep;39(9):993–1000.
 59. Hutchings L, Watkinson P, Young JD, Willett K. Defining multiple organ failure after major trauma: A comparison of the Denver, Sequential Organ Failure Assessment, and Marshall scoring systems. *J Trauma Acute Care Surg*. 2017 Mar;82(3):534–41.
 60. Moore FA, Moore EE, Poggetti R, McAnena OJ, Peterson VM, Abernathy CM, et al. Gut bacterial translocation via the portal vein: a clinical perspective with major torso trauma. *J Trauma*. 1991 May;31(5):629–36
 61. Deitch EA. Role of the gut lymphatic system in multiple organ failure. *Curr Opin Crit Care*. 2001 Apr;7(2):92–8.
 62. Tuma M, Canestrini S, Alwahab Z, Marshall J. Trauma and Endothelial Glycocalyx. *Shock*. 2016 Oct;46(4):352–7.

63. Rahbar E, Cardenas JC, Baimukanova G, Usadi B, Bruhn R, Pati S, et al. Endothelial glycocalyx shedding and vascular permeability in severely injured trauma patients. *J Transl Med*. 2015 Apr 12;13(1):127.
64. Chignalia AZ, Yetimakman F, Christiaans SC, Unal S, Bayrakci B, Wagener BM, et al. The Glycocalyx and trauma. *Shock*. 2016 Apr;45(4):338–48.
65. Schött U, Solomon C, Fries D, Bentzer P. The endothelial glycocalyx and its disruption, protection and regeneration: a narrative review. *Scand J Trauma Resusc Emerg Med*. 2016 Apr 12;24:48
66. Ostrowski SR, Johansson PI. Endothelial glycocalyx degradation induces endogenous heparinization in patients with severe injury and early traumatic coagulopathy. *J Trauma Acute Care Surg*. 2012 Jul;73(1):60–6.
67. Johansson PI, Stensballe J, Rasmussen LS, Ostrowski SR. A High Admission Syndecan-1 Level, A Marker of Endothelial Glycocalyx Degradation, Is Associated With Inflammation, Protein C Depletion, Fibrinolysis, and Increased Mortality in Trauma Patients. *Ann Surg*. 2011 Aug;254(2):194–200.
68. van Golen RF, Reiniers MJ, Vrisekoop N, Zuurbier CJ, Olthof PB, van Rheeën J, et al. The Mechanisms and Physiological Relevance of Glycocalyx Degradation in Hepatic Ischemia/Reperfusion Injury. *Antioxid Redox Signal*. 2014 Sep;21(7):1098–118.
69. Johansson PI, Stensballe J, Rasmussen LS, Ostrowski SR. High circulating adrenaline levels at admission predict increased mortality after trauma. *J Trauma Acute Care Surg*. 2012 Feb;72(2):428–36.
70. Mongardon N, Dyson A, Singer M. Is MOF an outcome parameter or a transient, adaptive state in critical illness? *Curr Opin Crit Care*. 2009 Oct;15(5):431–6.
71. Singer M, De Santis V, Vitale D, Jeffcoate W. Multiorgan failure is an adaptive, endocrine-mediated, metabolic response to overwhelming systemic inflammation. *Lancet*. 2004 Aug;364(9433):545–8.
72. Fullerton J, Singer M. Organ Failure in the ICU: Cellular Alterations. *Semin Respir Crit Care Med*. 2011 Oct 11;32(05):581–6.
73. Brealey D, Brand M, Hargreaves I, Heales S, Land J, Smolenski R, et al. Association between mitochondrial dysfunction and severity and outcome of septic shock. *Lancet*. 2002 Jul 20;360(9328):219–23.
74. Fredriksson K, Hammarqvist F, Strigard K, Hulténby K, Ljungqvist O, Wernerman J, et al. Derangements in mitochondrial metabolism in intercostal and leg muscle of critically ill patients with sepsis-induced multiple organ failure. *Am J Physiol Endocrinol Metab*. 2006 Nov;291(5):E1044–50
75. Singer M. The role of mitochondrial dysfunction in sepsis-induced multi-organ failure. *Virulence*. 2013 Oct 23;5(1):66–72.
76. Takasu O, Gaut JP, Watanabe E, To K, Fagley RE, Sato B, et al. Mechanisms of Cardiac and Renal Dysfunction in Patients Dying of Sepsis. *Am J Respir Crit Care Med*. 2013 Mar;187(5):509–17.
77. Noble JS, MacKirdy FN, Donaldson SI, Howie JC. Renal and respiratory failure in Scottish ICUs. *Anaesthesia*. 2001 Feb;56(2):124–9.

78. Hotchkiss RS, Swanson PE, Freeman BD, Tinsley KW, Cobb JP, Matuschak GM, et al. Apoptotic cell death in patients with sepsis, shock, and multiple organ dysfunction. *Crit Care Med*. 1999 Jul;27(7):1230–51.
79. Calvano SE, Xiao W, Richards DR, Felciano RM, Baker HV, Cho RJ, et al. A network-based analysis of systemic inflammation in humans. *Nature*. 2005 Aug 31;437(7061):1032–7.
80. Carré JE, Orban J-C, Re L, Felsmann K, Iffert W, Bauer M, et al. Survival in Critical Illness Is Associated with Early Activation of Mitochondrial Biogenesis. *Am J Respir Crit Care Med*. 2010 Sep 15;182(6):745–51.
81. Larsen FJ, Schiffer TA, Weitzberg E, Lundberg JO. Regulation of mitochondrial function and energetics by reactive nitrogen oxides. *Free Radic Biol Med*. 2012 Nov 15;53(10):1919–28.
82. Vlaar APJ, Juffermans NP. Transfusion-related acute lung injury: a clinical review. *Lancet*. 2013 Sep;382(9896):984–94.
83. Bolton-Maggs PHB, Cohen H. Serious Hazards of Transfusion (SHOT) haemovigilance and progress is improving transfusion safety. *Br J Haematol*. 2013 Sep 14;163(3):303–14.
84. Andreis DT, Singer M. Catecholamines for inflammatory shock: a Jekyll-and-Hyde conundrum. *Intensive Care Med*. 2016 Sep;42(9):1387–97.
85. Malbrain MLNG, Marik PE, Witters I, Cordemans C, Kirkpatrick AW, Roberts DJ, et al. Fluid overload, de-resuscitation, and outcomes in critically ill or injured patients: a systematic review with suggestions for clinical practice. *Anaesthesiol Intensive Ther*. 2014 Nov 28;46(5):361–80.
86. Vincent J-L, Taccone FS, He X. Harmful Effects of Hyperoxia in Postcardiac Arrest, Sepsis, Traumatic Brain Injury, or Stroke: The Importance of Individualized Oxygen Therapy in Critically Ill Patients. *Can Respir J*. 2017;2017:2834956.
87. Pannu S. Too Much Oxygen: Hyperoxia and Oxygen Management in Mechanically Ventilated Patients. *Semin Respir Crit Care Med*. 2016 Feb;37(1):16–22
88. Henderson WR, Chen L, Amato MBP, Brochard LJ. Fifty Years of Research in ARDS. *Respiratory Mechanics in Acute Respiratory Distress Syndrome*. *Am J Respir Crit Care Med*. 2017 Oct 1;196(7):822–33
89. Goodwin GH, Sanders C, Johns EW. A new group of chromatin-associated proteins with a high content of acidic and basic amino acids. *Eur J Biochem*. 1973 Sep 21;38(1):14–9.
90. Bianchi ME, Agresti A. HMG proteins: dynamic players in gene regulation and differentiation. *Curr Opin Genet Dev*. 2005 Oct;15(5):496–506.
91. Bustin M, Neihart NK. Antibodies against chromosomal HMG proteins stain the cytoplasm of mammalian cells. *Cell*. 1979 Jan;16(1):181–9.
92. Merenmies J, Pihlaskari R, Laitinen J. 30-kDa heparin-binding protein of brain (amphoterin) involved in neurite outgrowth. Amino acid sequence and

- localization in the filopodia of the advancing plasma. *J Biol Chem*. 1991 Sep 5; 266 (25) 16722-9.
93. Wang H. HMG-1 as a Late Mediator of Endotoxin Lethality in Mice. *Science*. 1999 Jul 9;285(5425):248–51.
 94. Yang H, Ochani M, Li J, Qiang X, Tanovic M, Harris HE, et al. Reversing established sepsis with antagonists of endogenous high-mobility group box 1. *Proc Natl Acad Sci USA*. 2004 Jan 6;101(1):296–301.
 95. Rouhiainen A, Tumova S, Valmu L. Analysis of proinflammatory activity of highly purified eukaryotic recombinant HMGB1 (amphoterin). *J Leukoc Biol*. 2007 Jan;81(1):49-58
 96. Tsan M-F. Heat shock proteins and high mobility group box 1 protein lack cytokine function. *J Leukoc Biol*. 2011 Jun;89(6):847–53.
 97. Bianchi ME. HMGB1 loves company. *J Leukoc Biol*. 2009 Sep;86(3):573-6.
 98. Yang H, Lundbäck P, Ottosson L, Erlandsson-Harris H, Venereau E, Bianchi ME, et al. Redox modification of cysteine residues regulates the cytokine activity of high mobility group box-1 (HMGB1). *Mol Med*. 2012 Mar 30;18:250–9.
 99. Harris HE, Andersson U, Pisetsky DS. HMGB1: a multifunctional alarmin driving autoimmune and inflammatory disease. *Nat Rev Rheumatol*. 2012 Jan 31;8(4):195–202.
 100. Venereau E, De Leo F, Mezzapelle R, Careccia G, Musco G, Bianchi ME. HMGB1 as biomarker and drug target. *Pharmacol Res*. 2016 Sep;111:534–44.
 101. Hardman CH, Broadhurst RW, Raine AR, Grasser KD, Thomas JO, Laue ED. Structure of the A-domain of HMG1 and its interaction with DNA as studied by heteronuclear three- and four-dimensional NMR spectroscopy. *Biochemistry*. 1995 Dec;34(51):16596–607.
 102. Weir HM, Kraulis PJ, Hill CS, Raine AR. Structure of the HMG box motif in the B-domain of HMG1. *EMBO J*. 1993 Apr;12(4):1311-9.
 103. Agresti A, Bianchi ME. HMGB proteins and gene expression. *Curr Opin Genet Dev*. 2003 Apr;13(2):170-8.
 104. Scaffidi P, Misteli T, Bianchi ME. Release of chromatin protein HMGB1 by necrotic cells triggers inflammation. *Nature*. 2002 Jul 11;418(6894):191-5.
 105. Banerjee S, Kundu TK. The acidic C-terminal domain and A-box of HMGB-1 regulates p53-mediated transcription. *Nucleic Acids Res*. 2003 Jun 15;31(12):3236-47.
 106. Reeves R. Nuclear functions of the HMG proteins. *Biochim Biophys Acta*. 2010 Jan;1799(1-2):3–14.
 107. Reeves R, Adair JE. Role of high mobility group (HMG) chromatin proteins in DNA repair. *DNA Repair*. 2005 Jul 28;4(8):926-38
 108. Calogero S, Grassi F, Aguzzi A, Voigtländer T, Ferrier P, Ferrari S, et al. The lack of chromosomal protein Hmg1 does not disrupt cell growth but causes lethal hypoglycaemia in newborn mice. *Nat Genet*. 1999 Jul;22(3):276–80.

109. Bonaldi T, Talamo F, Scaffidi P, Ferrera D, Porto A, Bachi A, et al. Monocytic cells hyperacetylate chromatin protein HMGB1 to redirect it towards secretion. *EMBO J*. 2003 Oct 15;22(20):5551–60.
110. Venereau E, Casalgrandi M, Schiraldi M. Mutually exclusive redox forms of HMGB1 promote cell recruitment or proinflammatory cytokine release. *J Exp Med*. 2012 Aug 27;209(9):1519–28.
111. Wahamoa H, Vallerskog T, Qin S, Lunderius C, LaRosa G, Andersson U, et al. HMGB1-secreting capacity of multiple cell lineages revealed by a novel HMGB1 ELISPOT assay. *J Leukoc Biol*. 2007 Jan;81(1):129–36.
112. Zetterstrom CK, Bergman T, Rynnel-Dagoo B, Erlandsson, Andersson U, et al. High mobility group box chromosomal protein 1 (HMGB1) is an antibacterial factor produced by the human adenoid. *Pediatr Res*. 2002 Aug;52(2):148–54.
113. Semino C, Angelini G, Poggi A, Rubartelli A. NK/iDC interaction results in IL-18 secretion by DCs at the synaptic cleft followed by NK cell activation and release of the DC maturation factor HMGB1. *Blood*. 2005 Jul 15;106(2):609–16.
114. Evankovich J, Cho SW, Zhang R, Cardinal J, Dhupar R, Zhang L, et al. High mobility group box 1 release from hepatocytes during ischemia and reperfusion injury is mediated by decreased histone deacetylase activity. *J Biol Chem*. 2010 Dec 17;285(51):39888–97.
115. Liu S, Stolz DB, Sappington PL, Macias CA, Killeen ME, Tenhunen JJ, et al. HMGB1 is secreted by immunostimulated enterocytes and contributes to cytomix-induced hyperpermeability of Caco-2 monolayers. *Am J Physiol Cell Physiol*. 2006 Apr;290(4):C990–9.
116. Sodhi CP, Jia H, Yamaguchi Y, Lu P, Good M, Egan C, et al. Intestinal Epithelial TLR-4 Activation Is Required for the Development of Acute Lung Injury after Trauma/Hemorrhagic Shock via the Release of HMGB1 from the Gut. *J Immunol*. 2015 May 15;194(10):4931–9.
117. Porto A, Palumbo R, Pieroni M, Aprigliano G, Chiesa R. Smooth muscle cells in human atherosclerotic plaques secrete and proliferate in response to high mobility group box 1 protein. *FASEB J*. 2006 Dec;20(14):2565–6.
118. Stark K, Philippi V, Stockhausen S, Busse J. Disulfide HMGB1 derived from platelets coordinates venous thrombosis in mice. *Blood*. 2016 Nov 17;128(20):2435–2449.
119. Gardella S, Andrei C, Ferrera D, Lotti LV, Torrisi MR, Bianchi ME, et al. The nuclear protein HMGB1 is secreted by monocytes via a non-classical, vesicle-mediated secretory pathway. *EMBO Rep*. 2002 Oct;3(10):995–1001.
120. Lamkanfi M, Sarkar A, Vande Walle L, Vitari AC, Amer AO, Wewers MD, et al. Inflammasome-dependent release of the alarmin HMGB1 in endotoxemia. *J Immunol*. 2010 Oct 1;185(7):4385–92.
121. Lu B, Nakamura T, Inouye K, Li J, Tang Y, Lundbäck P, et al. Novel role of PKR in inflammasome activation and HMGB1 release. *Nature*. 2012 Aug 30;488(7413):670–4.

122. Nyström S, Antoine DJ, Lundbäck P, Lock JG, Nita AF, Höglstrand K, et al. TLR activation regulates damage-associated molecular pattern isoforms released during pyroptosis. *EMBO J*. 2013 Jan 9;32(1):86–99.
123. Raucci A, Palumbo R, Bianchi ME. HMGB1: a signal of necrosis. *Autoimmunity*. 2007 Jun;40(4):285–9.
124. Zamaraeva MV, Sabirov RZ, Maeno E. Cells die with increased cytosolic ATP during apoptosis: a bioluminescence study with intracellular luciferase. *Cell Death Differ*. 2005 Nov;12(11):1390–7.
125. Richter C, Schweizer M, Cossarizza A, Franceschi C. Control of apoptosis by the cellular ATP level. *FEBS Lett*. 1996 Jan 8;378(2):107–10.
126. Newton K, Dugger DL, Wickliffe KE, Kapoor N, et al. Activity of protein kinase RIPK3 determines whether cells die by necroptosis or apoptosis. *Science*. 2014 Mar 21;343(6177):1357–60.
127. Zhang J, Yang Y, He W, Sun L. Necrosome core machinery: MLKL. *Cell Mol Life Sci*. 2016 Jun;73(11-12):2153–63.
128. Galluzzi L, Kepp O, Morselli E, Vitale I, Senovilla L, Pinti M, et al. Viral strategies for the evasion of immunogenic cell death. *J Intern Med*. 2010 May; 267(5):526–42.
129. Han J, Zhong CQ, Zhang DW. Programmed necrosis: backup to and competitor with apoptosis in the immune system. *Nat Immunol*. 2011 Nov 16;12(12):1143–9.
130. Takemoto K, Hatano E, Iwaisako K, Takeiri M, Noma N, Ohmae S, et al. Necrostatin-1 protects against reactive oxygen species (ROS)-induced hepatotoxicity in acetaminophen-induced acute liver failure. *FEBS Open Bio*. 2014 Sep 6;4:777–87.
131. Zhang T, Zhang Y, Cui M, Jin L, Wang Y, Lv F, et al. CaMKII is a RIP3 substrate mediating ischemia- and oxidative stress-induced myocardial necroptosis. *Nature Med*. 2016 Feb;22(2):175–82.
132. Degterev A, Huang Z, Boyce M, Li Y, Jagtap P, Mizushima N, et al. Chemical inhibitor of nonapoptotic cell death with therapeutic potential for ischemic brain injury. *Nat Chem Biol*. 2005 Jul;1(2):112–9.
133. Liang X, Chen Y, Zhang L, Jiang F, Wang W, Ye Z, et al. Necroptosis, a novel form of caspase-independent cell death, contributes to renal epithelial cell damage in an ATP-depleted renal ischemia model. *Mol Med Rep*. 2014 Aug; 10(2):719–24.
134. Qing DY, Conegliano D, Shashaty MGS, Seo J, Reilly JP, Worthen GS, et al. Red blood cells induce necroptosis of lung endothelial cells and increase susceptibility to lung inflammation. *Am J Respir Crit Care Med*. 2014 Dec 1;190(11):1243–54.
135. Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, et al. Neutrophil extracellular traps kill bacteria. *Science*. 2004 Mar;303(5663):1532–5.

136. Yipp BG, Petri B, Salina D, Jenne CN, Scott BNV, Zbytnuik LD, et al. Infection-induced NETosis is a dynamic process involving neutrophil multitasking in vivo. *Nature Med.* 2012 Sep;18(9):1386–93.
137. Tadie J-M, Bae H-B, Jiang S, Park DW, Bell CP, Yang H, et al. HMGB1 promotes neutrophil extracellular trap formation through interactions with Toll-like receptor 4. *Am J Physiol Lung Cell Mol Physiol.* 2013 Mar 1;304(5):L342–9.
138. Fuchs TA, Brill A, Duerschmied D. Extracellular DNA traps promote thrombosis. *Proc Natl Acad Sci USA.* 2010 Sep 7;107(36):15880-5.
139. Garcia-Romo GS, Caielli S, Vega B, Connolly J, Allantaz F, Xu Z, et al. Netting neutrophils are major inducers of type I IFN production in pediatric systemic lupus erythematosus. *Sci Transl Med.* 2011 Mar 9;3(73):73ra20.
140. Savchenko AS, Borissoff JI, Martinod K, De Meyer SF. VWF-mediated leukocyte recruitment with chromatin decondensation by PAD4 increases myocardial ischemia/reperfusion injury in mice. *Blood.* 2014 Jan 2;123(1):141-8.
141. Xu J, Jiang Y, WANG J, Shi X, Liu Q, Liu Z, et al. Macrophage endocytosis of high-mobility group box 1 triggers pyroptosis. *Cell Death Differ.* 2014 Aug; 21(8):1229–39.
142. Fink SL, Cookson BT. Caspase-1-dependent pore formation during pyroptosis leads to osmotic lysis of infected host macrophages. *Cell Microbiol.* 2006 Nov; 8(11):1812–25.
143. Yang D, He Y, Muñoz-Planillo R, Liu Q, Núñez G. Caspase-11 Requires the Pannexin-1 Channel and the Purinergic P2X7 Pore to Mediate Pyroptosis and Endotoxic Shock. *Immunity.* 2015 Nov 17;43(5):923–32.
144. de Gassart A, Martinon F. Pyroptosis: Caspase-11 Unlocks the Gates of Death. *Immunity.* 2015 Nov 17;43(5):835–7.
145. Kayagaki N, Warming S, Lamkanfi M, Vande Walle L, Louie S, Dong J, et al. Non-canonical inflammasome activation targets caspase-11. *Nature.* 2011 Oct 16;479(7371):117–21.
146. Knodler LA, Crowley SM, Sham HP, Yang H. Noncanonical inflammasome activation of caspase-4/caspase-11 mediates epithelial defenses against enteric bacterial pathogens. *Cell host microbe.* 2014 Aug 13;16(2):249–256.
147. Deng M, Tang Y, Li W, Wang X, Zhang R, Zhang X, et al. The Endotoxin Delivery Protein HMGB1 Mediates Caspase-11-Dependent Lethality in Sepsis. *Immunity.* 2018 Oct 16;49(4):740–53.e7
148. Yang J, Zhao Y, Zhang P, Li Y, Yang Y, Yang Y, et al. Hemorrhagic shock primes for lung vascular endothelial cell pyroptosis: role in pulmonary inflammation following LPS. *Cell Death Dis.* 2016 Sep 8;7(9):e2363.
149. Yang H, Antoine DJ, Andersson U. The many faces of HMGB1: molecular structure-functional activity in inflammation, apoptosis, and chemotaxis. *J Leukoc Biol.* 2013 Jun;93(6):865-73.

150. Tang Y, Zhao X, Antoine D, Xiao X, Wang H, Andersson U, et al. Regulation of Posttranslational Modifications of HMGB1 During Immune Responses. *Antioxid Redox Signal*. 2016 Apr 20;24(12):620–34.
151. Venereau E, Ceriotti C, Bianchi ME. DAMPs from Cell Death to New Life. *Front Immunol*. 2015;6:422.
152. Kazama H, Ricci JE, Herndon JM, Hoppe G, Green DR. Induction of immunological tolerance by apoptotic cells requires caspase-dependent oxidation of high-mobility group box-1 protein. *Immunity*. 2008 Jul 18;29(1):21–32.
153. Ito I, Fukazawa J, Yoshida M. Post-translational methylation of high mobility group box 1 (HMGB1) causes its cytoplasmic localization in neutrophils. *J Biol Chem*. 2007 Jun 1;282(22):16336–44.
154. Hreggvidsdóttir HS, Ostberg T, Wahamaa H, Schierbeck H, Aveberger A-C, Klevenvall L, et al. The alarmin HMGB1 acts in synergy with endogenous and exogenous danger signals to promote inflammation. *J Leukoc Biol*. 2009 Sep; 86(3):655–62.
155. Andersson U, Tracey KJ. HMGB1 is a therapeutic target for sterile inflammation and infection. *Annu Rev Immunol*. 2011;29:139–62.
156. Kang R, Chen R, Zhang Q, Hou W, Wu S, Cao L. HMGB1 in health and disease. *Mol aspects Med*. 2014 Dec;40:1-116.
157. Andersson U, Yang H, Harris H. High-mobility group box 1 protein (HMGB1) operates as an alarmin outside as well as inside cells. *Semin Immunol*. 2018 Aug; 38:40-48.
158. Hreggvidsdóttir HS, Lundberg AM, Aveberger A-C, Klevenvall L, Andersson U, Harris HE. High mobility group box protein 1 (HMGB1)-partner molecule complexes enhance cytokine production by signaling through the partner molecule receptor. *Mol Med*. 2012 Mar 27;18:224–30.
159. Wahamaa H, Schierbeck H. High mobility group box protein 1 in complex with lipopolysaccharide or IL-1 promotes an increased inflammatory phenotype in synovial fibroblasts. *Arthritis Res Ther*. 2011 Aug 26;13(4):R136.
160. Leclerc P, Wahamaa H, Idborg H, Jakobsson PJ, HARRIS HE, Korotkova M. IL-1 β /HMGB1 complexes promote The PGE2 biosynthesis pathway in synovial fibroblasts. *Scand J Immunol*. 2013 May;77(5):350–60.
161. Tian J, Avalos AM, Mao SY, Chen B, Senthil K. Toll-like receptor 9–dependent activation by DNA-containing immune complexes is mediated by HMGB1 and RAGE. *Nat Immunol*. 2007 May;8(5)487–96.
162. Yanai H, Ban T, Wang Z, Choi MK, Kawamura T, Negishi H, et al. HMGB proteins function as universal sentinels for nucleic-acid-mediated innate immune responses. *Nature*. 2009 Nov 5;462(7269):99–103.
163. Youn JH, Oh YJ, Kim ES, Choi JE. High mobility group box 1 protein binding to lipopolysaccharide facilitates transfer of lipopolysaccharide to CD14 and enhances lipopolysaccharide-mediated TNF- α in human monocytes. *J Immunol*. 2008 Apr 1;180(7):5067–74.

164. Yu M, Wang H, Ding A, Golenbock DT, Latz E, Czura CJ, et al. HMGB1 signals through toll-like receptor (TLR) 4 and TLR2. *Shock*. 2006 Aug;26(2):174–9.
165. Ivanov S, Dragoi A-M, Wang X, Dallacosta C, Louten J, Musco G, et al. A novel role for HMGB1 in TLR9-mediated inflammatory responses to CpG-DNA. *Blood*. 2007 Sep 15;110(6):1970–81.
166. van Zoelen MAD, Yang H, Florquin S, Meijers JCM, Akira S, Arnold B, et al. Role of toll-like receptors 2 and 4, and the receptor for advanced glycation end products in high-mobility group box 1-induced inflammation in vivo. *Shock*. 2009 Mar;31(3):280–4.
167. Yang H, Wang H, Ju Z, Ragab AA, Lundbäck P, Long W, et al. MD-2 is required for disulfide HMGB1-dependent TLR4 signaling. *J Exp Med*. 2015 Jan 12;212(1):5–14.
168. Yang H, Hreggvidsdottir HS, Palmblad K, Wang H, Ochani M, Li J, et al. A critical cysteine is required for HMGB1 binding to Toll-like receptor 4 and activation of macrophage cytokine release. *Proc Natl Acad Sci USA*. 2010 Jun 29;107(26):11942–7.
169. Chuah YK, Basir R, Talib H, Tie TH, et al. Receptor for advanced glycation end products and its involvement in inflammatory diseases. *Int J Inflamm*. 2013;2013:403460
170. Huebener P, Pradere J-P, Hernandez C, Gwak G-Y, Caviglia JM, Mu X, et al. The HMGB1/RAGE axis triggers neutrophil-mediated injury amplification following necrosis. *J Clin Invest*. 2015 Feb;125(2):539–50.
171. Fiuza C, Bustin M, Talwar S, Tropea M, Gerstenberger E, Shelhamer JH, et al. Inflammation-promoting activity of HMGB1 on human microvascular endothelial cells. *Blood*. 2003 Apr 1;101(7):2652–60.
172. Penzo M, Molteni R, Suda T, et al. Inhibitor of NF- κ B kinases α and β are both essential for high mobility group box 1-mediated chemotaxis. *J Immunol*. 2010 Apr 15;184(8):4497–509.
173. Raman KG, Sappington PL, Yang R, Levy RM, Prince JM, Liu S, et al. The role of RAGE in the pathogenesis of intestinal barrier dysfunction after hemorrhagic shock. *Am J Physiol Gastrointest Liver Physiol*. 2006 Oct;291(4):G556–65.
174. Yang R, Harada T, Mollen KP, Prince JM, Levy RM, Englert JA, et al. Anti-HMGB1 neutralizing antibody ameliorates gut barrier dysfunction and improves survival after hemorrhagic shock. *Mol Med*. 2006 Apr;12(4-6):105–14.
175. Gao TL, Yuan XT, Yang D, Dai HL, et al. Expression of HMGB1 and RAGE in rat and human brains after traumatic brain injury. *J Trauma Acute Care Surg*. 2012 Mar;72(3):643–9.
176. Schiraldi M, Raucci A, Muñoz LM, Livoti E, et al. HMGB1 promotes recruitment of inflammatory cells to damaged tissues by forming a complex with CXCL12 and signaling via CXCR4. *J Exp Med*. 2012 Mar 12;209(3):551–63.
177. Venereau E, Schiraldi M, Uguccioni M, Bianchi ME. HMGB1 and leukocyte migration during trauma and sterile inflammation. *Mol Immunol*. 2013 Aug; 55(1):76–82.

178. Degryse B, Bonaldi T, Scaffidi P, Müller S, Resnati M, Sanvito F, et al. The high mobility group (HMG) boxes of the nuclear protein HMGB1 induce chemotaxis and cytoskeleton reorganization in rat smooth muscle cells. *J Cell Biol.* 2001 Mar 19;152(6):1197–206.
179. Orlova VV, Choi EY, Xie C, Chavakis E, Bierhaus A, Ihanus E, et al. A novel pathway of HMGB1-mediated inflammatory cell recruitment that requires Mac-1-integrin. *EMBO J.* 2007 Feb 21;26(4):1129–39.
180. Zandarashvili L, Sahu D, Lee K, Lee YS, Singh P, Rajarathnam K, et al. Real-time Kinetics of High-mobility Group Box 1 (HMGB1) Oxidation in Extracellular Fluids Studied by in Situ Protein NMR Spectroscopy. *J Biol Chem.* 2013 Apr 26;288(17):11621–7.
181. Calandra T, Echtenacher B, Roy DL, Pugin J, Metz CN, Hultner L, et al. Protection from septic shock by neutralization of macrophage migration inhibitory factor. *Nature Med.* 2000 Feb;6(2):164–70.
182. Vanden Berghe T, Demon D, Bogaert P, Vandendriessche B, Goethals A, Depuydt B, et al. Simultaneous targeting of IL-1 and IL-18 is required for protection against inflammatory and septic shock. *Am J Respir Crit Care Med.* 2014 Feb 1;189(3):282–91.
183. Eskandari MK, Bolgos G, Miller C, Nguyen DT, DeForge LE, Remick DG. Anti-tumor necrosis factor antibody therapy fails to prevent lethality after cecal ligation and puncture or endotoxemia. *J Immunol.* 1992 May 1;148(9):2724–30.
184. Suda K, Kitagawa Y, Ozawa S, Saikawa Y, Ueda M, Ebina M, et al. Anti-high-mobility group box chromosomal protein 1 antibodies improve survival of rats with sepsis. *World J Surg.* 2006 Sep;30(9):1755–62.
185. Qin S, Wang H, Yuan R, Li H, Ochani M, Ochani K, et al. Role of HMGB1 in apoptosis-mediated sepsis lethality. *J Exp Med.* 2006 Jul 10;203(7):1637–42.
186. Chavan SS, Huerta PT, Robbiati S, Valdés-Ferrer SI, Ochani M, Dancho M, et al. HMGB1 mediates cognitive impairment in sepsis survivors. *Mol Med.* 2012 Sep 7;18:930–7.
187. Ferrer SV, Ballina MR, Olofsson PS, et al. HMGB1 mediates splenomegaly and expansion of splenic CD11b+ Ly-6C (high) inflammatory monocytes in murine sepsis survivors. *J Intern Med.* 2013 Oct;274(4):381–90.
188. Yang R, Miki K, Oksala N, Nakao A, Lindgren L, Killeen ME, et al. Bile high-mobility group box 1 contributes to gut barrier dysfunction in experimental endotoxemia. *Am J Physiol Regul Integr Comp Physiol.* 2009 Jul 21;297(2):R362–9.
189. Valdés-Ferrer SI, Papoin J, Dancho ME, Olofsson P, Li J, Lipton JM, et al. HMGB1 mediates anemia of inflammation in murine sepsis survivors. *Mol Med.* 2016 May;21(1):951–958.
190. Angus DC, Yang L, Kong L, Kellum JA, Delude RL, Tracey KJ, et al. Circulating high-mobility group box 1 (HMGB1) concentrations are elevated in both uncomplicated pneumonia and pneumonia with severe sepsis. *Crit Care Med.* 2007 Apr;35(4):1061–7.

191. Karlsson S, Pettilä V, Tenhunen J, Laru-Sompa R, Hynninen M, Ruokonen E. HMGB1 as a predictor of organ dysfunction and outcome in patients with severe sepsis. *Intensive Care Med.* 2008 Jun;34(6):1046–53.
192. Sund n-Cullberg J, Norrby-Teglund A, Rouhiainen A, Rauvala H, Herman G, Tracey KJ, et al. Persistent elevation of high mobility group box-1 protein (HMGB1) in patients with severe sepsis and septic shock. *Crit Care Med.* 2005 Mar;33(3):564–73.
193. Levy RM, Mollen KP, Prince JM, Kaczorowski DJ, Vallabhaneni R, Liu S, et al. Systemic inflammation and remote organ injury following trauma require HMGB1. *Am J Physiol Regul Integr Comp Physiol.* 2007 Oct;293(4):R1538–44.
194. Kim JY, Park JS, Strassheim D, Douglas I, Diaz del Valle F, Asehnoune K, et al. HMGB1 contributes to the development of acute lung injury after hemorrhage. *Am J Physiol Lung Cell Mol Physiol.* 2005 May;288(5):L958–65.
195. Zhou Y, Li Y, Mu T. HMGB1 Neutralizing Antibody Attenuates Cardiac Injury and Apoptosis Induced by Hemorrhagic Shock/Resuscitation in Rats. *Biol Pharm Bull.* 2015;38(8):1150–60.
196. Xu L, Zhao K, Shen X, Fan X-X, Ding K, Liu R-M, et al. Blockade of Extracellular High-Mobility Group Box 1 Attenuates Systemic Inflammation and Coagulation Abnormalities in Rats with Acute Traumatic Coagulopathy. *Med Sci Monit.* 2016 Jul 20;22:2561–70.
197. Shimazaki J, Matsumoto N, Ogura H, Muroya T, Kuwagata Y, Nakagawa J, et al. Systemic involvement of high-mobility group box 1 protein and therapeutic effect of anti-high-mobility group box 1 protein antibody in a rat model of crush injury. *Shock.* 2012 Jun;37(6):634–8.
198. Okuma Y, Liu K, Wake H, Zhang J, Maruo T, Date I, et al. Anti-high mobility group box-1 antibody therapy for traumatic brain injury. *Ann Neurol.* 2012 Sep; 72(3):373–84.
199. Ruan X, Darwiche SS, Cai C, Scott MJ, Pape H-C, Billiar TR. Anti-HMGB1 monoclonal antibody ameliorates immunosuppression after peripheral tissue trauma: attenuated T-lymphocyte response and increased splenic CD11b (+) Gr-1 (+) myeloid-derived suppressor cells require HMGB1. *Mediators Inflamm.* 2015 Jan 29;2015:458626.
200. Peltz ED, Moore EE, Eckels PC, Damle SS, Tsuruta Y, Johnson JL, et al. HMGB1 is markedly elevated within 6 hours of mechanical trauma in humans. *Shock.* 2009 Jul;32(1):17–22.
201. Cohen MJ, Brohi K, Calfee CS, Rahn P, Chesebro BB, Christiaans SC, et al. Early release of high mobility group box nuclear protein 1 after severe trauma in humans: role of injury severity and tissue hypoperfusion. *Crit Care.* 2009;13(6):R174.
202. Tsung A, Sahai R, Tanaka H, Nakao A, Fink MP, Lotze MT, et al. The nuclear factor HMGB1 mediates hepatic injury after murine liver ischemia-reperfusion. *J Exp Med.* 2005 Apr 4;201(7):1135–43.

203. Wu H, Ma J, Wang P, Corpuz TM, Panchapakesan U, Wyburn KR, et al. HMGB1 contributes to kidney ischemia reperfusion injury. *J Am Soc Nephrol*. 2010 Nov; 21(11):1878–90.
204. Andrassy M, Volz HC, Igwe JC, Funke B, Eichberger SN, Kaya Z, et al. High-mobility group box-1 in ischemia-reperfusion injury of the heart. *Circulation*. 2008 Jun 24;117(25):3216–26.
205. Zhang J, Takahashi HK, Liu K, Wake H, Liu R, Maruo T, et al. Anti-high mobility group box-1 monoclonal antibody protects the blood-brain barrier from ischemia-induced disruption in rats. *Stroke*. 2011 May;42(5):1420–8.
206. Liu K, Mori S, Takahashi HK, Tomono Y, Wake H, Kanke T, et al. Anti-high mobility group box 1 monoclonal antibody ameliorates brain infarction induced by transient ischemia in rats. *FASEB J*. 2007 Dec;21(14):3904–16.
207. Zheng YJ, Xu WP, Ding G, Gao YH, Wang HR, Pan SM. Expression of HMGB1 in septic serum induces vascular endothelial hyperpermeability. *Mol Med Rep*. 2016 Jan;13(1):513–21.
208. Heo JH, Han SW, Lee SK. Free radicals as triggers of brain edema formation after stroke. *Free Radic Biol Med*. 2005 Jul 1;39(1):51–70.
209. Haruma J, Teshigawara K, Hishikawa T, Wang D, Liu K, Wake H, et al. Anti-high mobility group box-1 (HMGB1) antibody attenuates delayed cerebral vasospasm and brain injury after subarachnoid hemorrhage in rats. *Sci rep*. 2016 Nov 24;6:37755.
210. Laird MD, Shields JS, Sukumari-Ramesh S, Kimbler DE, Fessler RD, Shakir B, et al. High mobility group box protein-1 promotes cerebral edema after traumatic brain injury via activation of toll-like receptor 4. *Glia*. 2014 Jan;62(1):26–38.
211. Lea JD, Clarke JI, McGuire N, Antoine DJ. Redox-Dependent HMGB1 Isoforms as Pivotal Co-Ordinators of Drug-Induced Liver Injury: Mechanistic Biomarkers and Therapeutic Targets. *Antioxid Redox Signal*. 2016 Apr 20;24(12):652–65.
212. Lundbäck P, Lea JD, Sowinska A, Ottosson L, Furst CM, Steen J, et al. A novel high mobility group box 1 neutralizing chimeric antibody attenuates drug-induced liver injury and postinjury inflammation in mice. *Hepatology*. 2016 Nov; 64(5):1699–710.
213. Ellerman JE, Brown CK, de Vera M, Zeh HJ, Billiar T, Rubartelli A, et al. Masquerader: high mobility group box-1 and cancer. *Clin Cancer Res*. 2007 May 15;13(10):2836–48.
214. Jube S, Rivera ZS, Bianchi ME, Powers A, Wang E. Cancer cell secretion of the DAMP protein HMGB1 supports progression in malignant mesothelioma. *Cancer Res*. 2012 Jul 1;72(13):3290–301.
215. Zeh HJ, Lotze MT. Addicted to death: invasive cancer and the immune response to unscheduled cell death. *J Immunother*. 2005 Jan;28(1):1–9.
216. Karin M, Clevers H. Reparative inflammation takes charge of tissue regeneration. *Nature*. 2016 Jan 21;529(7586):307–15.
217. Bianchi ME, Crippa MP, Manfredi AA, Mezzapelle R, Rovere-Querini P, Venereau E. High-mobility group box 1 protein orchestrates responses to tissue

- damage via inflammation, innate and adaptive immunity, and tissue repair. *Immunol Rev.* 2017 Oct 13;280(1):74–82.
218. Tirone M, Tran NL, Ceriotti C, Gorzanelli A, Canepari M, Bottinelli R, et al. High mobility group box 1 orchestrates tissue regeneration via CXCR4. *J Exp Med.* 2018 Jan 2;215(1):303–18.
 219. Dardenne AD, Wulff BC, Wilgus TA. The alarmin HMGB-1 influences healing outcomes in fetal skin wounds. *Wound Repair Regen.* 2013 Mar;21(2):282–91.
 220. Straino S, Di Carlo A, Mangoni A, De Mori R. High-mobility group box 1 protein in human and murine skin: involvement in wound healing. *J Invest Dermatol.* 2008 Jun;128(6):1545–53.
 221. Dormoy-Raclet V, Cammas A, Celona B, Lian XJ, van der Giessen K, Zivojnovic M, et al. HuR and miR-1192 regulate myogenesis by modulating the translation of HMGB1 mRNA. *Nat Commun.* 2013;4:2388.
 222. De Mori R, Straino S, Di Carlo A, Mangoni A, Pompilio G, Palumbo R, et al. Multiple effects of high mobility group box protein 1 in skeletal muscle regeneration. *Arterioscler Thromb Vasc Biol.* 2007 Nov;27(11):2377–83.
 223. Kohno T, Anzai T, Naito K, Miyasho T, Okamoto M, Yokota H, et al. Role of high-mobility group box 1 protein in post-infarction healing process and left ventricular remodelling. *Cardiovasc Res.* 2009 Feb 15;81(3):565–73.
 224. Kitahara T, Takeishi Y, Harada M, Niizeki T, Suzuki S, Sasaki T, et al. High-mobility group box 1 restores cardiac function after myocardial infarction in transgenic mice. *Cardiovasc Res.* 2008 Oct 1;80(1):40–6.
 225. Yang S, Xu L, Yang T, Wang F. High-mobility group box-1 and its role in angiogenesis. *J Leukoc Biol.* 2014 Apr;95(4):563–74.
 226. Tamai K, Yamazaki T, Chino T, Ishii M. PDGFR α -positive cells in bone marrow are mobilized by high mobility group box 1 (HMGB1) to regenerate injured epithelia. *Proc Natl Acad Sci USA.* 2011 Apr 19;108(16):6609–14.
 227. Rossini A, Zacheo A, Mocini D, Totta P, Facchiano A, Castoldi R, et al. HMGB1-stimulated human primary cardiac fibroblasts exert a paracrine action on human and murine cardiac stem cells. *J Mol Cell Cardiol.* 2008 Apr;44(4):683–93.
 228. Limana F, Germani A, Zacheo A, Kajstura J, Di Carlo A, Borsellino G, et al. Exogenous high-mobility group box 1 protein induces myocardial regeneration after infarction via enhanced cardiac C-kit⁺ cell proliferation and differentiation. *Circ Res.* 2005 Oct 14;97(8):e73–83.
 229. Campana L, Santarella F, Esposito A, Maugeri N, Rigamonti E, Monno A, et al. Leukocyte HMGB1 is required for vessel remodeling in regenerating muscles. *J Immunol.* 2014 Jun 1;192(11):5257–64.
 230. Takahashi K, Fukushima S, Yamahara K, Yashiro K, Shintani Y, Coppen SR, et al. Modulated inflammation by injection of high-mobility group box 1 recovers post-infarction chronically failing heart. *Circulation.* 2008 Sep 30;118(14 Suppl):S106–14.
 231. Tsung A, Tohme S, Billiar TR. High-mobility group box-1 in sterile inflammation. *J Intern Med.* 2014 Nov;276(5):425–43.

232. Sinclair NRS. B cell/antibody tolerance to our own antigens. *Front Biosci.* 2004 Sep;9:3019–28.
233. Schierbeck H, Lundbäck P, Palmblad K, Klevenvall L, Erlandsson-Harris H, Andersson U, et al. Monoclonal anti-HMGB1 (high mobility group box chromosomal protein 1) antibody protection in two experimental arthritis models. *Mol Med.* 2011 Sep;17(9-10):1039–44.
234. Löfblom J, Feldwisch J, Tolmachev V, Carlsson J, Ståhl S, Frejd FY. Affibody molecules: engineered proteins for therapeutic, diagnostic and biotechnological applications. *FEBS Letters.* 2010 Jun 18;584(12):2670–80.
235. Yang H, Liu H, Zeng Q, Imperato GH, Addorisio ME, Li J, et al. Inhibition of HMGB1/RAGE-mediated endocytosis by HMGB1 antagonist box A, anti-HMGB1 antibodies, and cholinergic agonists suppresses inflammation. *Mol Med.* 2019 Apr 11;25(1):13.
236. Li K, Yang J, Han X. Ketamine attenuates sepsis-induced acute lung injury via regulation of HMGB1-RAGE pathways. *Int Immunopharmacol.* 2016 May; 34:114–28.
237. Qin M-Z, Gu Q-H, Tao J, Song X-Y, Gan G-S, Luo Z-B, et al. Ketamine effect on HMGB1 and TLR4 expression in rats with acute lung injury. *Int J Clin Exp Pathol.* 2015;8(10):12943–8.
238. Zhang Z, Zhang L, Zhou C, Wu H. Ketamine inhibits LPS-induced HGMB1 release in vitro and in vivo. *Int Immunopharmacol.* 2014 Nov;23(1):14–26.
239. Andersson U, Yang H, Harris H. Extracellular HMGB1 as a therapeutic target in inflammatory diseases. *Expert Opin Ther Targets.* 2018 Mar;22(3):263–77
240. Onda H, Kasuya H, Takakura K, Hori T, Imaizumi T, Takeuchi T, et al. Identification of genes differentially expressed in canine vasospastic cerebral arteries after subarachnoid hemorrhage. *J Cereb Blood Flow Metab.* 1999 Nov; 19(11):1279–88.
241. Baekkevold ES, Roussigné M, Yamanaka T, Johansen F-E, Jahnsen FL, Amalric F, et al. Molecular characterization of NF-HEV, a nuclear factor preferentially expressed in human high endothelial venules. *Am J Pathol.* 2003 Jul;163(1):69–79.
242. Schmitz J, Owyang A, Oldham E, Song Y, Murphy E, McClanahan TK, et al. IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines. *Immunity* 2005 Nov;23(5):479–90.
243. Saluja R, Ketelaar ME, Hawro T, Church MK, Maurer M, Nawijn MC. The role of the IL-33/IL-1RL1 axis in mast cell and basophil activation in allergic disorders. *Mol Immunol.* 2015 Jan;63(1):80–5.
244. Moffatt MF, Gut IG, Demenais F, Strachan DP, Bouzigon E, Heath S, et al. A large-scale, consortium-based genomewide association study of asthma. *N Engl J Med.* 2010 Sep 23;363(13):1211–21.
245. Xu J, Guardado J, Hoffman R, Xu H, Namas R, Vodovotz Y, et al. IL33-mediated ILC2 activation and neutrophil IL5 production in the lung response after severe

- trauma: A reverse translation study from a human cohort to a mouse trauma model. *PLoS Med.* 2017 Jul;14(7):e1002365.
246. Sundlisæter E, Edelmann RJ, Hol J, Sponheim J, Kchler AM, Weiss M, et al. The Alarmin IL-33 Is a Notch Target in Quiescent Endothelial Cells. *Am J Pathol.* 2012 Sep 1;181(3):1099–111.
 247. Hong J, Bae S, Jhun H, Lee S, Choi J, Kang T, et al. Identification of constitutively active interleukin 33 (IL-33) splice variant. *J Biol Chem.* 2011 Jun 3;286(22):20078–86.
 248. Lingel A, Weiss TM, Niebuhr M, Pan B, Appleton BA, Wiesmann C, et al. Structure of IL-33 and Its Interaction with the ST2 and IL-1RAcP Receptors—Insight into Heterotrimeric IL-1 Signaling Complexes. *Structure.* 2009 Oct; 17(10):1398–410.
 249. Roussel L, Erard M, Cayrol C, Girard J-P. Molecular mimicry between IL-33 and KSHV for attachment to chromatin through the H2A–H2B acidic pocket. *EMBO Rep.* 2008 Aug 8;9(10):1006–12.
 250. Gautier V, Cayrol C, Farache D, Roga S, Monsarrat B, Burlet-Schiltz O, et al. Extracellular IL-33 cytokine, but not endogenous nuclear IL-33, regulates protein expression in endothelial cells. *Sci Rep.* 2016 Oct 3;6:34255.
 251. Sundnes O, Pietka W, Loos T, Sponheim J, Rankin AL, Pflanz S, et al. Epidermal Expression and Regulation of Interleukin-33 during Homeostasis and Inflammation: Strong Species Differences. *J Invest Dermatol.* 2015 Jan; 135(7):1771–80.
 252. Martin NT, Martin MU. Interleukin 33 is a guardian of barriers and a local alarmin. *Nat Immunol.* 2016 Feb;17(2):122–31.
 253. Kchler AM, Pollheimer J, Balogh J, Sponheim J, Manley L, Sorensen DR, et al. Nuclear Interleukin-33 Is Generally Expressed in Resting Endothelium but Rapidly Lost upon Angiogenic or Proinflammatory Activation. *Am J Pathol.* 2008 Oct;173(4):1229–42.
 254. Moussion C, Ortega N, Girard J-P. The IL-1-Like Cytokine IL-33 Is Constitutively Expressed in the Nucleus of Endothelial Cells and Epithelial Cells In Vivo: A Novel “Alarmin”. *PLoS ONE.* 2008 Oct 6;3(10):e3331.
 255. Talabot-Ayer D, Lamacchia C, Gabay C, Palmer G. Interleukin-33 is biologically active independently of caspase-1 cleavage. *J Biol Chem.* 2009 Jul 17;284(29): 19420–6.
 256. Cayrol C, Girard J-P. The IL-1-like cytokine IL-33 is inactivated after maturation by caspase-1. *Proc Natl Acad Sci USA.* 2009 Jun 2;106(22):9021–6.
 257. Lefranais E, Roga S, Gautier V, Gonzalez-de-Peredo A, Monsarrat B, Girard J-P, et al. IL-33 is processed into mature bioactive forms by neutrophil elastase and cathepsin G. *Proc Natl Acad Sci USA.* 2012 Jan 31;109(5):1673–8.
 258. Scott IC, Majithiya JB, Rapley L, Kemp BP, England E, Rees DG, et al. Oxidation of the alarmin IL-33 regulates ST2-dependent inflammation. *Nat Commun.* 2015 Sep 4;6:8327.

259. LUthi AU, Cullen SP, McNeela EA, Duriez PJ, Afonina IS, Sheridan C, et al. Suppression of interleukin-33 bioactivity through proteolysis by apoptotic caspases. *Immunity*. 2009 Jul 17;31(1):84–98.
260. Kakkar R, Hei H, Dobner S, Lee RT. Interleukin 33 as a Mechanically Responsive Cytokine Secreted by Living Cells. *J Biol Chem*. 2012 Feb 24;287(9):6941–8.
261. Zhao W, Hu Z. The enigmatic processing and secretion of interleukin-33. *Cell Mol Immunol*. 2010;7(4):260–2.
262. Chen W-Y, Hong J, Gannon J, Kakkar R, Lee RT. Myocardial pressure overload induces systemic inflammation through endothelial cell IL-33. *Proc Natl Acad Sci USA*. 2015 Jun 9;112(23):7249–54.
263. Bessa J, Meyer CA, de Vera Mudry MC, Schlicht S, Smith SH, Iglesias A, et al. Altered subcellular localization of IL-33 leads to non-resolving lethal inflammation. *J Autoimmun*. 2014 Dec;55(C):33–41.
264. Chackerian AA, Oldham ER, Murphy EE, Schmitz J, Pflanz S, Kastelein RA. IL-1 Receptor Accessory Protein and ST2 Comprise the IL-33 Receptor Complex. *J Immunol*. 2007 Aug 3;179(4):2551–5.
265. Liu X, Hammel M, He Y, Tainer JA, Jeng U-S, Zhang L, et al. Structural insights into the interaction of IL-33 with its receptors. *Proc Natl Acad Sci USA*. 2013 Sep 10;110(37):14918–23.
266. Martin MU. Special aspects of interleukin-33 and the IL-33 receptor complex. *Semin Immunol*. 2013 Dec 15;25(6):449–57.
267. Ali S, Huber M, Kollewe C, Bischoff SC, Falk W, Martin MU. IL-1 receptor accessory protein is essential for IL-33-induced activation of T lymphocytes and mast cells. *Proc Natl Acad Sci USA*. 2007 Nov 20;104(47):18660–5.
268. Yanagisawa K, Takagi T, Tsukamoto T, Tetsuka T, et al. Presence of a Novel Primary Response Gene St2l, encoding a Product Highly Similar to the Interleukin-1 Receptor Type-1. *FEBS Lett*. 1993 Feb 22;318(1):83–7.
269. Hayakawa H, Hayakawa M, Kume A, Tominaga S-I. Soluble ST2 blocks interleukin-33 signaling in allergic airway inflammation. *J Biol Chem*. 2007 Sep 7;282(36):26369–80.
270. Tajima S, Oshikawa K, Tominaga S-I, Sugiyama Y. The increase in serum soluble ST2 protein upon acute exacerbation of idiopathic pulmonary fibrosis. *Chest*. 2003 Oct;124(4):1206–14.
271. Oshikawa K, Kuroiwa K, Tago K, Iwahana H, YANAGISAWA K, Ohno S, et al. Elevated soluble ST2 protein levels in sera of patients with asthma with an acute exacerbation. *Am J Respir Crit Care Med*. 2001 Jul 15;164(2):277–81.
272. Shimpo M. Serum Levels of the Interleukin-1 Receptor Family Member ST2 Predict Mortality and Clinical Outcome in Acute Myocardial Infarction. *Circulation*. 2004 May 11;109(18):2186–90.
273. Wang Y-P, Wang J-H, Wang X-L, Liu J-Y, Jiang F-Y, Huang X-L, et al. Roles of ST2, IL-33 and BNP in predicting major adverse cardiovascular events in acute

- myocardial infarction after percutaneous coronary intervention. *J Cell Mol Med*. 2017 Jun 17;21(11):2677–84.
274. Parikh RH, Seliger SL, Christenson R, Gottdiener JS, Psaty BM, deFilippi CR. Soluble ST2 for Prediction of Heart Failure and Cardiovascular Death in an Elderly, Community-Dwelling Population. *J Am Heart Assoc*. 2016 Aug 1;5(8):e003188.
 275. Hoogerwerf JJ, Tanck MWT, van Zoelen MAD, Wittebole X, Laterre P-F, van der Poll T. Soluble ST2 plasma concentrations predict mortality in severe sepsis. *Intensive Care Med*. 2010 Feb 12;36(4):630–7.
 276. Hur M, Kim H, Kim HJ, Yang HS, Magrini L, Marino R, et al. Soluble ST2 Has a Prognostic Role in Patients With Suspected Sepsis. *Ann Lab Med*. 2015 Nov; 35(6):570–7
 277. Kondo Y, Yoshimoto T, Yasuda K, Futatsugi-Yumikura S, Morimoto M, Hayashi N, et al. Administration of IL-33 induces airway hyperresponsiveness and goblet cell hyperplasia in the lungs in the absence of adaptive immune system. *Int Immunol*. 2008 Jun 1;20(6):791–800.
 278. Neill DR, Wong SH, Bellosi A, Flynn RJ, Daly M, Langford TKA, et al. Nuocytes represent a new innate effector leukocyte that mediates type-2 immunity. *Nature*. 2010 Mar 3;464(7293):1367–70.
 279. Price AE, Liang H-E, Sullivan BM, Reinhardt RL, Eisley CJ, Erle DJ, et al. Systemically dispersed innate IL-13-expressing cells in type 2 immunity. *Proc Natl Acad Sci USA*. 2010;107(25):11489–94.
 280. Walker JA, McKenzie AN. Development and function of group 2 innate lymphoid cells. *Curr Opin Immunol*. 2013 Apr 1;25(2):148–55.
 281. Lunderius-Andersson C, Enoksson M, Nilsson G. Mast Cells Respond to Cell Injury through the Recognition of IL-33. *Front Immunol*. 2012 Apr 19;3:82.
 282. Bønnelykke K, Sleiman P, Nielsen K, Kreiner-Møller E, Mercader JM, Belgrave D, et al. A genome-wide association study identifies CDHR3 as a susceptibility locus for early childhood asthma with severe exacerbations. *Nat Genet*. 2014 Jan; 46(1):51–5.
 283. Préfontaine D, Lajoie-Kadoch S, Foley S, Audusseau S, Olivenstein R, Halayko AJ, et al. Increased expression of IL-33 in severe asthma: evidence of expression by airway smooth muscle cells. *J Immunol*. 2009 Oct 15;183(8):5094–103.
 284. Kurowska-Stolarska M, Stolarski B, Kewin P, Murphy G, Corrigan CJ, Ying S, et al. IL-33 Amplifies the Polarization of Alternatively Activated Macrophages That Contribute to Airway Inflammation. *J Immunol*. 2009 Nov 4;183(10):6469–77.
 285. Kearley J, Silver JS, Sanden C, Liu Z, Berlin AA, White N, et al. Cigarette Smoke Silences Innate Lymphoid Cell Function and Facilitates an Exacerbated Type I Interleukin-33-Dependent Response to Infection. *Immunity*. 2015 Mar 17;42(3):566–79.
 286. Li S, Zhu F-X, Zhao X-J, An Y-Z. The immunoprotective activity of interleukin-33 in mouse model of cecal ligation and puncture-induced sepsis. *Immunol Lett*. 2016 Jan;169:1–7.

287. Alves-Filho JC, nego FSO, Souto FO, Freitas A, Verri WA, Auxiliadora-Martins M, et al. Interleukin-33 attenuates sepsis by enhancing neutrophil influx to the site of infection. *Nat Med*. 2010 Jun;16(6):708–12.
288. Skaga NO, Eken T, Søvik S. Validating performance of TRISS, TARN and NORMIT survival prediction models in a Norwegian trauma population. *Acta Anaesthesiol Scand*. 2018 Feb;62(2):253–66.
289. Nadeau-Vallée M, Obari D, Palacios J, Brien M-È, Duval C, Chemtob S, et al. Sterile inflammation and pregnancy complications: a review. *Reproduction*. 2016 Dec;152(6):R277–92.
290. Matthews JN, Altman DG, Campbell MJ, Royston P. Analysis of serial measurements in medical research. *BMJ*. 1990 Jan 27;300(6719):230–5.
291. Skogstrand K, Ekelund CK, Thorsen P, Vogel I, Jacobsson B, Nørgaard-Pedersen B, et al. Effects of blood sample handling procedures on measurable inflammatory markers in plasma, serum and dried blood spot samples. *J Immunol Methods*. 2008 Jul 20;336(1):78–84.
292. Lehner J, Wittwer C, Fersching D, Siegele B, Holdenrieder S, Stoetzer OJ. Methodological and preanalytical evaluation of an HMGB1 immunoassay. *Anticancer Res*. 2012 May;32(5):2059–62.
293. Davies JE, Apta BHR, Harper MT. Cross-reactivity of anti-HMGB1 antibodies for HMGB2. *J Immunol Methods*. 2018 May;456:72–76
294. Elshal MF, McCoy JP. Multiplex bead array assays: performance evaluation and comparison of sensitivity to ELISA. *Methods*. 2006 Apr;38(4):317–23.
295. Warade J. Retrospective Approach to Evaluate Interferences in Immunoassay. *EJIFCC*. 2017 Oct;28(3):224–32.
296. Rath A, Glibowicka M, Nadeau VG, Chen G, Deber CM. Detergent binding explains anomalous SDS-PAGE migration of membrane proteins. *Proc Natl Acad Sci USA*. 2009 Feb 10;106(6):1760–5.
297. Bakowitz M, Bruns B, McCunn M. Acute lung injury and the acute respiratory distress syndrome in the injured patient. *Scand J Trauma Resusc Emerg Med*. 2012 Aug 10;20:54.
298. Miller PR, Croce MA, Kilgo PD, Scott J, Fabian TC. Acute respiratory distress syndrome in blunt trauma: identification of independent risk factors. *Am Surg*. 2002 Oct;68(10):845–50.
299. Navarrete-Navarro P, Rivera-Fernández R, Rincón-Ferrari MD, García-Delgado M, Muñoz A, Jiménez JM, et al. Early markers of acute respiratory distress syndrome development in severe trauma patients. *J Crit Care*. 2006 Sep;21(3):253–8.
300. Watkins TR, Nathens AB, Cooke CR, Psaty BM, Maier RV, Cuschieri J, et al. Acute respiratory distress syndrome after trauma. *Crit Care Med*. 2012 Aug;40(8):2295–303.
301. Del Sorbo L, Slutsky AS. Acute respiratory distress syndrome and multiple organ failure. *Curr Opin Crit Care*. 2011 Feb;17(1):1–6.

302. Vincent J-L, Zambon M. Why Do Patients Who Have Acute Lung Injury/Acute Respiratory Distress Syndrome Die from Multiple Organ Dysfunction Syndrome? Implications for Management. *Clin Chest Med*. 2006 Dec;27(4):725–31.
303. Schoenfeld DA, Bernard GR, ARDS Network. Statistical evaluation of ventilator-free days as an efficacy measure in clinical trials of treatments for acute respiratory distress syndrome. *Crit Care Med*. 2002 Aug;30(8):1772–7.
304. Bouamra O, Wrotchford A, Hollis S, Vail A, Woodford M, Lecky F. A New Approach to Outcome Prediction in Trauma: A Comparison With the TRISS Model. *J Trauma*. 2006 Sep;61(3):701–10.
305. Jones JM, Skaga NO, Søvik S, Lossius HM, Eken T. Norwegian survival prediction model in trauma: modelling effects of anatomic injury, acute physiology, age, and co-morbidity. *Acta Anaesthesiol Scand*. 2014 Jan 20;58(3):303–15.
306. Baker SP, O'Neill B, Haddon WJ, Long WB. The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma*. 1974 Mar;14(3):187–96.
307. Kilgo PD, Meredith JW, Hensberry R, Osler TM. A note on the disjointed nature of the injury severity score. *J Trauma*. 2004 Sep;57(3):479–85.
308. Balogh Z, Offner PJ, Moore EE, Biffi WL. NISS predicts postinjury multiple organ failure better than the ISS. *J Trauma*. 2000 Apr;48(4):624–7.
309. Meredith JW, Evans G, Kilgo PD, MacKenzie E, Osler T, McGwin G, et al. A comparison of the abilities of nine scoring algorithms in predicting mortality. *J Trauma*. 2002 Oct;53(4):621–8.
310. Ibrahim I, Chor WP, Chue KM, Tan CS, Tan HL, Siddiqui FJ, et al. Is arterial base deficit still a useful prognostic marker in trauma? A systematic review. *Am J Emerg Med*. 2016 Mar;34(3):626–35.
311. Rixen D, Siegel JH. Bench-to-bedside review: oxygen debt and its metabolic correlates as quantifiers of the severity of hemorrhagic and post-traumatic shock. *Crit Care*. 2005 Oct 5;9(5):441–53.
312. Ouellet J-F, Roberts DJ, Tiruta C, Kirkpatrick AW, Mercado M, Trottier V, et al. Admission base deficit and lactate levels in Canadian patients with blunt trauma: are they useful markers of mortality? *J Trauma Acute Care Surg*. 2012 Jun;72(6):1532–5.
313. Kroezen F, Bijlsma TS, Liem MSL, Meeuwis JD, Leenen LPH. Base Deficit-Based Predictive Modeling of Outcome in Trauma Patients Admitted to Intensive Care Units in Dutch Trauma Centers. *J Trauma*. 2007 Oct;63(4):908–13.
314. Callaway DW, Shapiro NI, Donnino MW, Baker C, Rosen CL. Serum Lactate and Base Deficit as Predictors of Mortality in Normotensive Elderly Blunt Trauma Patients. *J Trauma*. 2009 Apr;66(4):1040–4.
315. Raux M, Le Manach Y, Gauss T, Baumgarten R, Hamada S, Harrois A, et al. Comparison of the Prognostic Significance of Initial Blood Lactate and Base Deficit in Trauma Patients. *Anesthesiology*. 2017 Mar;126(3):522–33.

316. Gjedsted J, Buhl M, Nielsen S, Schmitz O, Vestergaard ET, Tønnesen E, et al. Effects of adrenaline on lactate, glucose, lipid and protein metabolism in the placebo controlled bilaterally perfused human leg. *Acta Physiol*. 2011 Aug; 202(4):641–8.
317. Joestl J, Lang NW, Kleiner A, Platzer P, Aldrian S. The Importance of Sex Differences on Outcome after Major Trauma: Clinical Outcome in Women Versus Men. *J Clin Med*. 2019 Aug;8(8):1263.
318. Wang X-W, Karki A, Zhao X-J, Xiang X-Y, Lu Z-Q. High plasma levels of high mobility group box 1 is associated with the risk of sepsis in severe blunt chest trauma patients: a prospective cohort study. *J Cardiothorac Surg*. 2014 Aug 2;9:133.
319. Urbonaviciute V, Fürnrohr BG, Weber C, Haslbeck M, Wilhelm S, Herrmann M, et al. Factors masking HMGB1 in human serum and plasma. *J Leukoc Biol*. 2007 Jan;81(1):67–74.
320. Yang H, Wang H, Levine YA, Gunasekaran MK, Wang Y, Addorisio M, et al. Identification of CD163 as an antiinflammatory receptor for HMGB1-haptoglobin complexes. *JCI Insight*. 2016 May 19;1(7):e85375
321. Joly P, Marshall JC, Tessier PA, Massé C, Page N, Frenette AJ, et al. S100A8/A9 and sRAGE kinetic after polytrauma; an explorative observational study. *Scand J Trauma Resusc Emerg Med*. 2017 Nov 25;25(1):114.
322. Cohen MJ, Carles M, Brohi K, Calfee CS, Rahn P, Call MS, et al. Early Release of Soluble Receptor for Advanced Glycation Endproducts After Severe Trauma in Humans. *J Trauma*. 2010 Jun;68(6):1273–8.
323. Fan J, LI Y, Levy RM, Fan JJ, Hackam DJ, Vodovotz Y, et al. Hemorrhagic Shock Induces NAD(P)H Oxidase Activation in Neutrophils: Role of HMGB1-TLR4 Signaling. *J Immunol*. 2007 May 2;178(10):6573–80.
324. Xu P, Wen Z, Shi X, LI Y, Fan L, Xiang M, et al. Hemorrhagic Shock Augments Nlrp3 Inflammasome Activation in the Lung through Impaired Pyrin Induction. *J Immunol*. 2013 May 3;190(10):5247–55.
325. Xiang M, Shi X, LI Y, Xu J, Yin L, Xiao G, et al. Hemorrhagic Shock Activation of NLRP3 Inflammasome in Lung Endothelial Cells. *J Immunol*. 2011 Oct 19;187(9):4809–17.
326. Maugeri N, Campana L, Gavina M, Covino C, De Metrio M, Panciroli C, et al. Activated platelets present high mobility group box 1 to neutrophils, inducing autophagy and promoting the extrusion of neutrophil extracellular traps. *J Thromb Haemost*. 2014 Nov 14;12(12):2074–88.
327. Maugeri N, Franchini S, Campana L, Baldini M, Ramirez GA, Sabbadini MG, et al. Circulating platelets as a source of the damage-associated molecular pattern HMGB1 in patients with systemic sclerosis. *Autoimmunity*. 2012 Sep 17;45(8):584–7.
328. Basso D, Padoan A, Laufer T, Aneloni V, Moz S, Schroers H, et al. Relevance of pre-analytical blood management on the emerging cardiovascular protein biomarkers TWEAK and HMGB1 and on miRNA serum and plasma profiling. *Clin Biochem*. 2017 Mar;50(4-5):186–93.

329. Weng L, Guo L, Vachani A, Mesaros C, Blair IA. Quantification of Serum High Mobility Group Box 1 by Liquid Chromatography/High-Resolution Mass Spectrometry: Implications for Its Role in Immunity, Inflammation, and Cancer. *Anal Chem*. 2018 Jun 19;90(12):7552–60.
330. Abdulahad DA, Westra J, Bijzet J, Limburg PC, Kallenberg CGM, Bijl M. High mobility group box 1 (HMGB1) and anti-HMGB1 antibodies and their relation to disease characteristics in systemic lupus erythematosus. *Arthritis Res Ther*. 2011 May 6;13(3):R71.
331. Barnay-Verdier SP, Gaillard C, Messmer M, Borde C, Gibot S, Maréchal V. PCA-ELISA: A sensitive method to quantify free and masked forms of HMGB1. *Cytokine*. 2011 Jul 1;55(1):4–7.
332. Gaillard C, Borde C, Gozlan J, Maréchal V, Strauss F. A High-Sensitivity Method for Detection and Measurement of HMGB1 Protein Concentration by High-Affinity Binding to DNA Hemicatenanes. *PLoS ONE*. 2008 Aug 6;3(8):e2855.
333. Moore HM, Kelly A, Jewell SD, McShane LM, Clark DP, Greenspan R, et al. Biospecimen Reporting for Improved Study Quality. *Biopreserv Biobank*. 2011 Apr;9(1):57–70.
334. Cayrol C, Girard J-P. Interleukin-33 (IL-33): A nuclear cytokine from the IL-1 family. *Immunol Rev*. 2017 Dec 16;281(1):154–68.
335. Wu Q, Chaplin H, Ambrose N, Sen D, Leandro MJ, Wing C, et al. Juvenile arthritis disease activity score is a better reflector of active disease than the disease activity score 28 in adults with polyarticular juvenile idiopathic arthritis. *Ann Rheum Dis*. 2016 Feb 10;75(3):635–6.
336. Ketelaar ME, Nawijn MC, Shaw DE, Koppelman GH, Sayers I. The challenge of measuring IL-33 in serum using commercial ELISA: lessons from asthma. *Clin Exp Allergy*. 2016 May 26;46(6):884–7.
337. Dhillon OS, Narayan HK, Quinn PA, Squire IB, Davis JE, et al. Interleukin 33 and ST2 in non-ST-elevation myocardial infarction: Comparison with Global Registry of Acute Coronary Events Risk Scoring and NT-proBNP. *Am Heart J*. 2011 Jun 1;161(6):1163–70.
338. Thierry A, Giraud S, Robin A, Barra A, Bridoux F, Ameteau V, et al. The Alarmin Concept Applied to Human Renal Transplantation: Evidence for a Differential Implication of HMGB1 and IL-33. *PLoS ONE*. 2014 Feb 20;9(2):e88742.
339. Barbosa IG, Rodrigues DH, Rocha NP, da Cunha Sousa LF, Vieira ELM, Simões-e-Silva AC, et al. Plasma levels of alarmin IL-33 are unchanged in autism spectrum disorder: a preliminary study. *J Neuroimmunol*. 2015 Jan 15;278:69–72.
340. Yanaba K, Yoshizaki A, Asano Y, Kadono T, Sato S. Serum IL-33 levels are raised in patients with systemic sclerosis: association with extent of skin sclerosis and severity of pulmonary fibrosis. *Clin Rheumatol*. 2011 Jan 19;30(6):825–30.
341. Pastorelli L, Garg RR, Hoang SB, Spina L, Mattioli B, Scarpa M, et al. Epithelial-derived IL-33 and its receptor ST2 are dysregulated in ulcerative colitis and in experimental Th1/Th2 driven enteritis. *Proc Natl Acad Sci USA*. 2010 Apr 27;107(17):8017–22.

- 342. Chen T, Jia R-Z, Guo Z-P, Cao N, Li M-M, Jiao X-Y. Elevated serum interleukin-33 levels in patients with Henoch-Schönlein purpura. *Arch Dermatol Res*. 2012 Jul 27;305(2):173–7.
- 343. Yazdani HO, Chen H-W, Tohme S, Tai S, van der Windt DJ, Loughran P, et al. IL-33 exacerbates liver sterile inflammation by amplifying neutrophil extracellular trap formation. *J Hepatol*. 2017 Sep 21;68:130–9
- 344. Antoine DJ, Jenkins RE, Dear JW, Williams DP, McGill MR, Sharpe MR, et al. Molecular forms of HMGB1 and keratin-18 as mechanistic biomarkers for mode of cell death and prognosis during clinical acetaminophen hepatotoxicity. *J Hepatol*. 2012 May 1;56(5):1070–9.

Paper II

RESEARCH ARTICLE

Open Access

HMGB1 concentration measurements in trauma patients: assessment of pre-analytical conditions and sample material

William Ottestad^{1,2*} , Ingrid N. Rognes², Erlend Skaga², Cassandra Frisvoll³, Guttorm Haraldsen^{4,5}, Torsten Eken^{1,2†} and Peter Lundbäck^{4,5†}

Abstract

Background: HMGB1 is a mediator of systemic inflammation in sepsis and trauma, and a promising biomarker in many diseases. There is currently no standard operating procedure for pre-analytical handling of HMGB1 samples, despite that pre-analytical conditions account for a substantial part of the overall error rate in laboratory testing. We hypothesized that the considerable variations in reported HMGB1 concentrations and kinetics in trauma patients could be partly explained by differences in pre-analytical conditions and choice of sample material.

Methods: Trauma patients ($n = 21$) admitted to a Norwegian Level I trauma center were prospectively included. Blood was drawn in K₂EDTA coated tubes and serum tubes. The effects of delayed centrifugation were evaluated in samples stored at room temperature for 15 min, 3, 6, 12, and 24 h respectively. Plasma samples subjected to long-term storage in -80°C and to repeated freeze/thaw cycles were compared with previously analyzed samples. HMGB1 concentrations in simultaneously acquired arterial and venous samples were also compared. HMGB1 was assessed by standard ELISA technique, additionally we investigated the suitability of western blot in both serum and plasma samples.

Results: Arterial HMGB1 concentrations were consistently lower than venous concentrations in simultaneously obtained samples (arterial = $0.60 \times$ venous; 95% CI 0.30–0.90). Concentrations in plasma and serum showed a strong linear correlation, however wide limits of agreement. Storage of blood samples at room temperature prior to centrifugation resulted in an exponential increase in plasma concentrations after ≈ 6 h. HMGB1 concentrations were fairly stable in centrifuged plasma samples subjected to long-term storage and freeze/thaw cycles. We were not able to detect HMGB1 in either serum or plasma from our trauma patients using western blotting.

Conclusions: Arterial and venous HMGB1 concentrations cannot be directly compared, and concentration values in plasma and serum must be compared with caution due to wide limits of agreement. Although HMGB1 levels in clinical samples from trauma patients are fairly stable, strict adherence to a pre-analytical protocol is advisable in order to protect sample integrity. Surprisingly, we were unable to detect HMGB1 utilizing standard western blot analysis.

Keywords: HMGB1 protein, Pre-analytical phase, Wounds and injuries, Enzyme-linked immunosorbent assay, Blotting, western

* Correspondence: williamottestad@gmail.com

†Torsten Eken and Peter Lundbäck contributed equally to this work.

¹Department of Anaesthesiology, Oslo University Hospital, PO Box 4956 Nydalen, NO-0424 Oslo, Norway

²Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway

Full list of author information is available at the end of the article

© The Author(s). 2019 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Background

High-mobility group box 1 protein (HMGB1) is an evolutionary conserved DNA-binding protein present in all nucleated eukaryotic cells and in platelets. Under physiological conditions, HMGB1 is located predominantly in the nucleus where it acts as a nucleosomal stabilizer and a transcriptional regulator. (Bianchi and Agresti 2005) Outside the cell, HMGB1 acquires a new identity to serve as a powerful mediator of inflammation. (Yang et al. 2013) It is passively released or actively secreted in a multitude of pathological conditions including sepsis, trauma, cancer, and auto-immune diseases, and in many cases a predictable correlation to disease activity is evident. (Harris et al. 2012; Ottestad et al. 2019) The biological actions of HMGB1 are remarkable in their diversity, caused by a marked propensity for post-translational modifications. (Yang et al. 2012)

Several studies have explored HMGB1 concentrations in trauma patients utilizing standard ELISA techniques, however there is a considerable variation in reported concentrations and kinetics even in comparable patient populations. (Ottestad et al. 2019; Peltz et al. 2009; Cohen et al. 2009; Yang et al. 2006; Namas et al. 2016) Although pre-analytical conditions have been reported to account for 46 to 68% of the overall error rate in laboratory testing in general, (Carraro et al. 2012; Plebani 2012) there are no recommended operating procedures for clinical HMGB1 analysis that are widely applied. (Moore et al. 2011) We have therefore explored some crucial aspects for measurement of HMGB1 concentrations in trauma patients.

First, analyses of inflammatory molecules are often biased by sample degradation or ex vivo sample activation. This is especially true for constitutively expressed inflammatory markers. (Skogstrand et al. 2008; Flower et al. 2000) Release of HMGB1 from dying blood cells could also potentially affect total levels in blood samples over time. Further complicating the issue, HMGB1 stability in a given sample could conceivably depend on disease type and state, potentially invalidating experience gained from studies in other disease categories. We have therefore evaluated effects on measured HMGB1 concentrations in samples from trauma patients stored at room temperature for variable time periods prior to centrifugation and freezing, and subjected to long-term storage at -80°C with several freeze/thaw cycles after centrifugation.

Second, immunoassays are prone to interference by endogenous factors that alter antibody binding or the measurable concentration of the analyte. (Tate and Ward 2004) Cross-reactivity, unsuspected protein-protein bindings, and lipemia are all known to cause interference, and these factors can depend on type of sample material. (Tate and Ward 2004; Clerico et al. 2018) In line with this, both

serum and plasma components that interfere with HMGB1 detection by ELISA systems have been reported, yielding different results in serum vs. plasma depending on disease state. (Urbonaviciute et al. 2007; Basso et al. 2017; Lehner et al. 2012) We have therefore compared HMGB1 levels measured by ELISA in simultaneously acquired serum and plasma samples from patients early after trauma. The suitability of western blot (WB) analysis as an alternative to ELISA was also addressed, as western blotting is less prone to interference by protein-protein interactions due to a denaturing step.

Third, previous studies of HMGB1 in trauma patients have either utilized venous samples only, (Cohen et al. 2009) mainly arterial samples, (Ottestad et al. 2019) an unspecified mixture of arterial and venous samples, (Namas et al. 2016) or not reported the source of their material. (Peltz et al. 2009; Yang et al. 2006) It could be argued that arterial samples may be preferable in order to obtain blood that is not draining from any particular injured body part, (Ottestad et al. 2019) however we are not aware of any study comparing HMGB1 concentration in arterial vs. venous blood after trauma. We have therefore also investigated HMGB1 concentrations in plasma derived from simultaneously acquired arterial and venous samples in trauma patients.

Methods

Recruitment of trauma patients was performed January 2011 – November 2018 at Oslo University Hospital Ullevål, a Norwegian Level I trauma center. (Ottestad et al. 2019) Patients were recruited by convenience. All patients ≥ 18 years who met criteria for trauma team activation were eligible for enrollment. Patients with burn injuries and pregnant women were excluded.

Arterial or venous blood was drawn in K_2EDTA coated tubes and serum tubes (Vacuette 454209 and 367977, Greiner Bio-One International GmbH, Austria) immediately after admission. The tubes were put directly in ice slush after 8–10 inversions, and after 15 min centrifuged at 2500 g at 4°C for 15 min. Aliquots of the supernatants were immediately transferred to sterile polypropylene tubes (NUNC CryoTubes 479–6843; VWR International AS, Oslo, Norway) and stored at -80°C .

In a recent study of HMGB1 kinetics in trauma patients (Ottestad et al. 2019) we have reported HMGB1 plasma concentrations as a function of time after injury from 136 of the patients. In the current study, we have compared previously reported plasma concentration measurements from several time points in 5 of these patients to reanalyzed plasma and to serum samples that were not originally analyzed, but subjected to long-term storage at -80°C .

Simultaneous arterial and venous blood samples were acquired in 6 additional patients, four of them both at

admission and at approximately 4 h after injury. The latter time point was selected to embrace the previously described “second wave” of HMGB1 release after trauma. (Ottestad et al. 2019) In another 10 patients, admission arterial samples were drawn in five 2 ml EDTA tubes which were stored at room temperature ($\approx 25^{\circ}\text{C}$) for 15 min, 3, 6, 12, and 24 h respectively before centrifugation and freezing. An overview of patients and samples in each group is displayed in Table 1.

HMGB1 concentrations were obtained by standard ELISA technique according to manufacturer instruction (HMGB1 ELISA Kit II; ST51011; Shino-Test Corporation, Tokyo, Japan). Serum samples and plasma samples subjected to freeze/thaw and long-term storage were analyzed with a single ELISA kit, however compared with original plasma analyses performed with ELISA kits from a different production batch for our previous study. Analyses of simultaneously acquired arterial and venous samples were done within a single ELISA kit in order to rule out inter-assay variability, in quadruplicate with results reported as median concentrations.

Western blotting was performed with 1 μL plasma or serum diluted in 9 μL H_2O + 3 μL Laemmli sample buffer (100 mM Tris-HCl pH 6.8, 45% glycerol, 5% sodium dodecyl sulfate, 12.5% 2-mercaptoethanol, and 0.25% bromophenol blue), and incubated at 97°C for 10 min. Recombinant human HMGB1 (HMGBiotech HM100,

Milano, Italy) was used as a positive control in western blot analyses at different spike-in concentrations in normal venous plasma and serum from one healthy subject. SDS-PAGE electrophoresis was performed at 200 V for 30 min using mini-protean TGX gels 4–20% (Bio-Rad Laboratories AB, Oslo, Norway). The gel was transferred to nitrocellulose membrane (Trans-Blot[®] Turbo[™] Mini Nitrocellulose 0.2 μm , BioRad Laboratories AB, Oslo, Norway). Membranes were blocked with 5% milk powder in Tris-Buffered Saline and Tween20 TBS-T for 45 min at room temperature, and rinsed once in TBS-T. The membranes were incubated with primary anti-HMGB1 antibodies (Abcam 18256, Cambridge, UK, or m2G7, kind gift from Professor Helena Erlandsson Harris) (Lundbäck et al. 2016) diluted to 1 $\mu\text{g}/\text{mL}$ in 1% milk/TBS-T over night at 4°C . Gels were then washed 4×5 min in TBS-T and incubated with HRP-conjugated anti-rabbit antibody (Jackson ImmunoResearch 711-035-152, Immunolab, Oslo, Norway) or HRP-conjugated anti-mouse antibody (Jackson ImmunoResearch 711-035-151, Immunolab, Oslo, Norway) for 2 h at room temperature, diluted 1:20,000 in 1% milk/TBS-T, followed by extensive washing with TBS-T.

Data analysis was undertaken using JMP 11.2.1 and 13.2.0 (SAS Institute, Cary, NC). A two-tailed $p \leq 0.05$ was chosen to represent statistical significance. Group comparisons were performed with the Wilcoxon rank-

Table 1 Characteristics of the three study populations

Characteristics	Sample Storage patients (n = 10)	Serum vs. Plasma patients (n = 5)	Arterial vs. Venous patients (n = 6)
Demographics			
Sex (male: female)	7: 3	3: 2	6: 0
Age (years)	48 (29–68; 19–78); 10	54 (30–55; 23–55); 5	35 (19–52; 17–57); 6
Pre-injury ASA PS (ASA I: II: III)	5: 3: 2	3: 1: 1	5: 0: 1
Injuries			
Mechanism of injury (blunt: penetrating)	10: 0	4: 1	5: 1
NISS	34 (22–43; 12–75); 10	41 (22–53; 9–57); 5	41 (7–59; 1–66); 6
ISS	28 (20–32; 5–48); 10	24 (15–46; 9–57); 5	36 (4–49; 1–50); 6
Admission BE (mmol/L)	−4.1 (−7.3 – −1.0; −8.3 – 0.1); 10	−3.4 (−5.6 – −2.0; −6.2 – −1.6); 4	−2.3 (−6.2–0.3; −8.7 – 3.7); 6
HMGB1 analyses			
Admission HMGB1 (ng/mL) ^a	6.57 (3.01–29.9; 1.68–155); 10	50.8 (1.88–91.2; 1.54–97.9); 5	13.2 (8.62–17.7; 7.31–20.4); 6
Time from injury to first sample (hours) ^b	3:07 (1:03–5:44; 0:39–7:26); 9	0:45 (0:31–2:41; 0:29–4:06); 5	1:41 (0:49–2:49; 0:17–3:50); 6
Hospital treatment			
ICU length of stay (days)	6 (2–11; 1–17); 10	7 (3–9; 2–10); 5	6 (2–11; 2–12); 6
Ventilator treatment (y: n)	5: 5	2: 3	5: 1
Outcome			
Dead at 30 days (y: n)	1: 9	1: 4	1: 5
Ventilator-free days	29 (11–30; 0–30); 10	30 (0–30; 0–30); 5	23 (17–29; 0–30); 6

Numbers are given as median (quartiles; range) and number of patients if not otherwise specified. NISS New Injury Severity Score, ISS Injury Severity Score, BE Base Excess, ICU intensive care unit, Ventilator-free days, days alive and off ventilator during the first 30 days after trauma. ^aFor “Arterial vs. Venous” group, as median concentration from first arterial sample. ^bTime from injury missing for one patient in “Sample Storage” group

sum test, or with the Wilcoxon signed-rank test for paired measurements. Agreement between methods was assessed with the Bland–Altman method for repeated measures. (Myles 2007; Bland and Altman 2007) Linear mixed model analysis with patient as random effect was utilised to assess correlation between original plasma, serum, and reanalyzed plasma, relations between arterial and venous HMGB1 concentration in simultaneous samples, and HMGB1 concentration change during storage. The reported coefficient of multiple determination (R^2) estimates the proportion of variation in the response that can be attributed to the model rather than to random error.

Results

Comparisons between HMGB1 concentrations in samples analyzed early, subjected to long-term storage or stored for a variable time in room temperature before centrifugation and freezing, in plasma vs. serum, and in arterial vs. venous blood were done in three separate groups of trauma patients. Characteristics of the study populations are shown in Table 1.

Effects of storage in room temperature before centrifugation and freezing were evaluated in admission samples from 10 trauma patients (Table 1; Fig. 1). There was no significant difference between HMGB1 concentrations after 15 min whether the samples were cooled in ice slush or left in room temperature ($p = 0.16$, Wilcoxon signed-rank test). Concentrations were stable for approximately 3–6 h, but thereafter increased exponentially

Fig. 1 Semilogarithmic plot of HMGB1 concentrations as a function of storage time in room temperature before centrifugation and freezing. Individual patients are color coded

in samples from eight of the ten patients (Fig. 1). Surprisingly, HMGB1 concentrations from the two patients with highest initial value did not change during the 24 h storage period (linear mixed model; $p = 0.60$).

Effects of long-term storage and freeze/thaw cycles on HMGB1 plasma concentration were assessed through reanalysis of 24 plasma samples from five trauma patients (Table 1). Following the initial ELISA-based HMGB1 analyses, each sample was subjected to > 2 years of -80°C storage and 3 freeze/thaw cycles. HMGB1 concentrations in original and reanalyzed plasma showed a strong linear correlation ($R^2 = 0.986$, $p < 0.0001$), however wide limits of agreement (Fig. 2a; mean difference -1.2 ng/mL, Limits of Agreement [LoA] -21.7 to 19.3).

Correspondence between HMGB1 serum concentrations and previously analyzed plasma determined by ELISA analysis was assessed in 22 sample pairs from the same patients (Table 1). HMGB1 concentrations in plasma and serum showed a strong linear correlation ($R^2 = 0.83$, $p < 0.0001$), however wide limits of agreement (Fig. 2b; mean difference -1.7 ng/mL, LoA -25.5 to 22.1 ng/mL).

Fig. 2 Bland–Altman scatter plot of difference between previously analyzed plasma and reanalyzed plasma (a) and previously analyzed plasma and serum (b) against the average of the two measurements. Samples from individual patients are color coded

Western blot analyses were performed on simultaneously obtained serum and plasma admission samples from two patients, with ELISA plasma HMGB1 concentrations of 50.8 and 97.9 ng/mL respectively. These are among the highest HMGB1 concentrations reported in our previous study (population median at admission 3.74 ng/mL, range 0.31–223 ng/mL, $n = 135$). (Ottestad et al. 2019) We were unable to detect HMGB1 in any of the samples, utilizing either the polyclonal rabbit antibody or the monoclonal mouse antibody (data not shown). To determine the lowest level of detection by western blot, we diluted recombinant HMGB1 in normal serum and plasma, and were only able to detect HMGB1 bands at spike-in concentrations equal to or above 1000 ng/mL.

Simultaneously acquired arterial and venous samples were obtained from 6 patients at admission (median 1:41 h after injury, range 0:17–3:50 h, Table 1; T01 in Fig. 3). Samples were also obtained from 4 of the patients at a median of 4:12 h after injury (range 3:56–4:16 h; T02 in Fig. 3). Median absolute time difference between simultaneously acquired samples was 3 min (range 1–4 min); arterial blood was sampled before venous in 6 of the 10 sample pairs. Arterial concentrations were consistently lower than venous

(Fig. 3; $p = 0.01$, Wilcoxon signed-rank test), and there was no significant effect of whether the arterial sample was obtained before or after the venous sample ($p = 0.83$, Wilcoxon rank-sum test). Effects of venous HMGB1 concentration on arterial HMGB1 concentration were assessed in a mixed model with patient as random effect. Arterial HMGB1 concentration was $0.60 \times$ venous HMGB1 concentration (95% CI 0.30–0.90, $R^2 = 0.68$, $p = 0.002$).

Discussion

We are not aware of any previously published reports comparing venous and arterial blood in clinical samples with respect to systemic HMGB1 levels. Surprisingly, arterial concentrations were 40% lower than venous concentrations, and it is tempting to speculate whether the pulmonary circulation might be a site for HMGB1 clearance. Alternatively, factors in arterial blood may interfere with HMGB1 ELISA detection. HMGB1 undergoes rapid cysteine redox changes, (Zandarashvili et al. 2013) and since ambient oxygen pressure increases when venous blood reaches the pulmonary circulation, HMGB1 oxidation to disulfide or sulfonyl isoforms could potentially alter protein-protein interactions and consequently detection by ELISA. The fraction of free vs. complexed HMGB1 could conceivably also be affected by redox state, as is the case with the alarmin IL33. (Scott et al. 2015)

HMGB1 analyses are often performed on plasma or serum samples that have been subjected to variable duration of storage and sometimes repeated freeze/thaw cycles. Previous studies in non-trauma patients have reported that HMGB1 concentrations in serum samples stored in room temperature for up to 7 days after centrifugation or subjected to freeze/thaw cycles were remarkably stable. (Lehner et al. 2012; Wang et al. 2015) In accordance with this we found a strong linear correlation between concentrations in originally analyzed plasma and plasma subjected to long-term storage and freeze/thaw cycles, however comparison of absolute concentration values was not straightforward due to wide limits of agreement. Despite HMGB1 being fairly stable during storage in room temperature, we found an exponential increase in HMGB1 levels after 3–6 h when blood samples from trauma patients were stored at room temperature before centrifugation and freezing. Surprisingly, HMGB1 concentrations from the two samples with the highest initial HMGB1 value did not exhibit an exponential increase during storage. Neutrophils and thrombocytes have been reported as major vehicles for HMGB1, both cell types are easily activated and short lived, and consequently potential sources of ex-vivo release of HMGB1 in blood samples. Contribution from other cell types can also not be excluded. Although highly speculative, ex-vivo release of HMGB1 might

conceivably be limited upon depletion due to extravasation to injured tissues or clot formation following traumatic coagulopathy. Kinetic analysis of more cell type specific markers may be required to explain this observation.

Based on these findings, we speculate that differences in previously reported HMGB1 concentrations in comparable populations of trauma patients (Ottestad et al. 2019; Cohen et al. 2009; Yang et al. 2006; Namas et al. 2016) might be caused by variability in pre-analytical conditions. Peltz et al. (2009) published the first paper to document elevated HMGB1 plasma concentrations after trauma in humans, however their HMGB1 concentrations were 100-fold higher than HMGB1 levels reported in similar studies with comparable populations of trauma patients. (Ottestad et al. 2019; Cohen et al. 2009; Yang et al. 2006; Namas et al. 2016) Notably, in contrast to the other authors, Peltz et al. utilized samples from EDTA tubes previously drawn for clinical indications and processed within 24 h. Storage temperature and time from sampling to centrifugation were not reported, and their results must therefore be interpreted with caution.

HMGB1 concentrations in parallel plasma and serum samples showed strong linear correlation, but wide limits of agreement. Consequently, we recommend that absolute concentration values from plasma and serum samples should be compared with caution, at least in the context of trauma.

The application of immunoassays in complex matrices such as plasma and serum is challenged by interference, (Tate and Ward 2004; Clerico et al. 2018) also known as “matrix effects”. Polyreactive antibodies or unsuspected protein–protein interactions may interfere with antibody detection, decreasing both sensitivity and specificity of the assay and consequently challenging reliable HMGB1 detection in clinical samples. Previous reports have suggested that factors in serum and plasma interfere with HMGB1 detection in ELISA systems. (Urbonaviciute et al. 2007; Bianchi 2009; Abdulahad et al. 2011) HMGB1 is a “sticky” protein that interacts with a multitude of molecules, including proteins, DNA and lipids. In trauma there is an abundance of intracellular proteins leaked from necrotic cells into the circulating plasma, and proteins binding to HMGB1 could potentially negatively impact HMGB1 detection in the ELISA system. ELISA depends on an antibody capture step in order to decrease the complexity of the sample and increase signal-to-noise ratio. Thus, test sensitivity will be affected by sample refinement and type of capture antibody used.

In western blotting, under reducing and denaturing conditions, interfering proteins or other factors will not stay complexed to HMGB1 unless they are covalently bound. Consequently, western blotting could potentially allow for a more unbiased detection of HMGB1.

However, we were unable to detect HMGB1 in either serum or plasma from our trauma patients, using two diverse primary antibodies that recognize surface-exposed epitopes. The mAb has a specificity to A-box residue 53–63, and was generated by immunization with recombinant full-length HMGB1. (Qin et al. 2006) The polyclonal Ab has specificity to residue 151–185, including parts of the B-box and linker and both antibodies have been described to be suitable as primary western blot antibodies. However, antibody epitopes may be inaccessible because of sample-specific post-translational modifications (e.g. glycosylation). HMGB1 bands were only detected at spike-in concentrations of 10–50 times the maximum level measured by ELISA in our trauma patients. High protein content in most clinical samples limits the possible sample-loading amount, as expected we experienced problems with unspecific binding and interference with migration at higher serum or plasma volumes than 1 µL/lane. We avoided the use of immunoprecipitation or other enrichment strategies in order to ensure an unbiased approach.

Our study has several limitations. It was a single-center study, and sample size was small and limited to trauma patients only. Reanalyzed plasma and serum samples were compared with plasma analyzed with ELISA assays from a different production batch, and part of the disagreement between original plasma and reanalyzed plasma or original plasma and serum might be explained by inter-assay imprecision which has been reported to be up to 14% for the HMGB1 immunoassay that was used. (Lehner et al. 2012)

Conclusions

HMGB1 concentration measurements are sensitive to pre-analytical conditions and choice of sample material. Consequently, standard operating procedures for clinical HMGB1 research should be established so that robust conclusions can be drawn when comparing results from different studies. The observation that arterial HMGB1 concentrations were consistently lower than venous concentrations in simultaneously obtained samples is novel and raises the question whether the pulmonary circulation is a site for HMGB1 clearance. Absolute concentration values in plasma and serum must be compared with caution due to wide limits of agreement. In accordance with earlier studies we confirmed that HMGB1 plasma levels are fairly stable in samples subjected to long-term storage and freeze / thaw cycles, however storage in room temperature before centrifugation resulted in exponential increase in HMGB1 concentrations after 3–6 h. The high correlation between concentration measurements before and after long-term storage allows studies of kinetic profiles within subjects despite wide limits of agreement. Somewhat unexpectedly, standard western

blot analysis failed to detect HMGB1 at clinically relevant levels in plasma and serum samples from our trauma patients.

Abbreviations

DNA: Deoxyribonucleic acid; EDTA: Ethylenediaminetetraacetate; ELISA: Enzyme-linked immunosorbent assay; HMGB1: High mobility group box 1 protein; HRP: Horseradish peroxidase

Acknowledgements

We thank Julie Katrine Lindstad for help with project design and ELISA analyses. We are grateful to Ivan Jonassen Rimstad, Astrid Arnesen Hug, Linda Soilammi, and Tomas Drægni for help with recruitment of patients and acquisition of samples, and to Anne-Marie Siebke Trøseid and Jens Petter Berg for help with optimization of pre-analytical conditions and for lab facilities. We thank the Oslo University Hospital Trauma Registry and registrars Morten Hestnes and Hans Johansson for providing clinical data. Special thanks to Frank Sætre for help with ELISA analyses, and to Helena Erlandsson Harris for providing us with anti-HMGB1 antibodies (m2G7) for our western blot analysis.

Authors' contributions

TE, WO and PL conceived the study and were responsible for overall design; WO, INR, TE, ES, and CF collected the data. INR, WO and PL performed the ELISA and WB analyses. TE and WO designed the statistical analyses and did the initial data interpretation; TE, WO, GH and PL finalized analyses and interpretation. WO and PL wrote the first version of the manuscript, and all authors critically revised and approved the final version.

Funding

This study was financially supported by Oslo University Hospital, the University of Oslo, and South-Eastern Norway Regional Health Authority.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

All parts of the study were approved by the Regional committee for medical and health research ethics (2010/2014 REK Sør-Øst D), in accordance with the Declaration of Helsinki. Patients were enrolled on admission, and written informed consent was obtained as soon as possible. A temporary written consent from the closest relative was obtained when patients were unable to consent. All material was destroyed if patients or relatives did not consent. The Regional committee for medical and health research ethics approved the use of biological material from patients who died before a fully informed consent could be obtained, and the use of written information with the possibility for withdrawal for patients who were transferred to other hospitals or discharged before consent could be obtained. All those patients were checked against the Norwegian national biological research reservation registry before final inclusion.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Anaesthesiology, Oslo University Hospital, PO Box 4956 Nydalen, NO-0424 Oslo, Norway. ²Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway. ³Faculty of Medicine, University of Oslo, Oslo, Norway. ⁴K.G. Jebsen Inflammation Research Centre, Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway. ⁵Department of Pathology, Oslo University Hospital, Oslo, Norway.

Received: 18 September 2019 Accepted: 18 December 2019

References

- Abdulahad DA, Westra J, Bijzet J, Limburg PC, Kallenberg CGM, Bijl M. High mobility group box 1 (HMGB1) and anti-HMGB1 antibodies and their relation to disease characteristics in systemic lupus erythematosus. *Arthritis Res Ther*. 2011;13(3):R71.
- Basso D, Padoan A, Laufer T, Aneloni V, Moz S, Schroers H, et al. Relevance of pre-analytical blood management on the emerging cardiovascular protein biomarkers TWEAK and HMGB1 and on miRNA serum and plasma profiling. *Clin Biochem*. 2017;50(4–5):186–93.
- Bianchi ME. HMGB1 loves company. *J Leukoc Biol*. 2009;86(3):573–6.
- Bianchi ME, Agresti A. HMG proteins: dynamic players in gene regulation and differentiation. *Curr Opin Genet Dev*. 2005;15(5):496–506.
- Bland JM, Altman DG. Agreement between methods of measurement with multiple observations per individual. *J Biopharm Stat*. 2007;17(4):571–82.
- Carraro P, Zago T, Plebani M. Exploring the initial steps of the testing process: frequency and nature of pre-preanalytic errors. *Clin Chem*. 2012;58(3):638–42.
- Clerico A, Belloni L, Carrozza C, Correale M, Dittadi R, Dotti C, et al. A black swan in clinical laboratory practice: the analytical error due to interferences in immunoassay methods. *Clin Chem Lab Med*. 2018;56(3):397–402.
- Cohen MJ, Brohi K, Calfee CS, Rahn P, Chesebro BB, Christiaans SC, et al. Early release of high mobility group box nuclear protein 1 after severe trauma in humans: role of injury severity and tissue hypoperfusion. *Crit Care*. 2009;13(6):R174.
- Flower L, Ahuja RH, Humphries SE, Mohamed-Ali V. Effects of sample handling on the stability of interleukin 6, tumour necrosis factor-alpha and leptin. *Cytokine*. 2000;12(11):1712–6.
- Harris HE, Andersson U, Pisetsky DS. HMGB1: a multifunctional alarmin driving autoimmune and inflammatory disease. *Nat Rev Rheumatol*. 2012;8(4):195–202.
- Lehner J, Wittwer C, Fersching D, Siegele B, Holdenrieder S, Stoetzer OJ. Methodological and preanalytical evaluation of an HMGB1 immunoassay. *Anticancer Res*. 2012;32(5):2059–62.
- Lundbäck P, Lea JD, Sowinska A, Ottosson L, Furst CM, Steen J, et al. A novel high mobility group box 1 neutralizing chimeric antibody attenuates drug-induced liver injury and postinjury inflammation in mice. *Hepatology*. 2016;64(5):1699–710.
- Moore HM, Kelly A, Jewell SD, McShane LM, Clark DP, Greenspan R, et al. Biospecimen reporting for improved study quality. *Biopreserv Biobank*. 2011;9(1):57–70.
- Myles PS, Cui J. Using the Bland-Altman method to measure agreement with repeated measures. *Br J Anaesth*. 2007;99(3):309–11.
- Namas RA, Vodovotz Y, Almahmoud K, Abdul-Malak O, Zaaqoq A, Namas R, et al. Temporal patterns of circulating inflammation biomarker networks differentiate susceptibility to nosocomial infection following blunt trauma in humans. *Ann Surg*. 2016;263(1):191–8.
- Ottestad W, Rognes IN, Pischke SE, Mollnes TE, Andersson U, Eken T. Biphasic release of the alarmin High Mobility Group Box 1 protein early after trauma predicts poor clinical outcome. *Crit Care Med*. 2019;47(8):e614–e622.
- Peltz ED, Moore EE, Eckels PC, Damle SS, Tsuruta Y, Johnson JL, et al. HMGB1 is markedly elevated within 6 hours of mechanical trauma in humans. *Shock*. 2009;32(1):17–22.
- Plebani M. Quality indicators to detect pre-analytical errors in laboratory testing. *Clin Biochem Rev*. 2012;33(3):85–8.
- Qin S, Wang H, Yuan R, Li H, Ochani M, Ochani K, et al. Role of HMGB1 in apoptosis-mediated sepsis lethality. *J Exp Med*. 2006;203(7):1637–42.
- Scott IC, Majithiya JB, Rapley L, Kemp BP, England E, Rees DG, et al. Oxidation of the alarmin IL-33 regulates ST2-dependent inflammation. *Nat Commun*. 2015;6:1–10 Nature Publishing Group.
- Skogstrand K, Ekelund CK, Thorsen P, Vogel I, Jacobsson B, Nørgaard-Pedersen B, et al. Effects of blood sample handling procedures on measurable inflammatory markers in plasma, serum and dried blood spot samples. *J Immunol Methods*. 2008;336(1):78–84.
- Tate J, Ward G. Interferences in immunoassay. *Clin Biochem Rev*. 2004;25(2):105–20.
- Urbancic V, Füllrohr BG, Weber C, Haslbeck M, Wilhelm S, Herrmann M, et al. Factors masking HMGB1 in human serum and plasma. *J Leukoc Biol*. 2007;81(1):67–74.
- Wang J, Zhu HH, Xue JH, Wu SS, Chen Z. Effects of storage conditions on the stability of serum CD163, NGAL, HMGB1 and MIP2. *Int J Clin Exp Pathol*. 2015;8(4):4099–105.

- Yang H, Antoine DJ, Andersson U. The many faces of HMGB1: molecular structure-functional activity in inflammation, apoptosis, and chemotaxis. *J Leukoc Biol.* 2013;93(6):865–73.
- Yang H, Lundbäck P, Ottosson L, Erlandsson-Harris H, Venereau E, Bianchi ME, et al. Redox modification of cysteine residues regulates the cytokine activity of high mobility group box-1 (HMGB1). *Mol Med.* 2012 Mar 30; 18:250–9.
- Yang R, Harada T, Mollen KP, Prince JM, Levy RM, Englert JA, et al. Anti-HMGB1 neutralizing antibody ameliorates gut barrier dysfunction and improves survival after hemorrhagic shock. *Mol Med.* 2006;12(4–6):105–14.
- Zandarashvili L, Sahu D, Lee K, Lee YS, Singh P, Rajarathnam K, et al. Real-time kinetics of high-mobility group box 1 (HMGB1) oxidation in extracellular fluids studied by in situ protein NMR spectroscopy. *J Biol Chem.* 2013;288(17):11621–7.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions