**Regional Structure and Basin Development of the mid-
Norwegian Volcanic Passive Margin: Anatomy of a
Polyphased Rifted System**

Dmitry Zastrozhnov

© Dmitry Zastrozhnov , 2020

*Series of dissertations submitted to the
Faculty of Mathematics and Natural Sciences, University of Oslo
No. 2269*

ISSN 1501-7710

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

“A geophysicist is a person who passes as an exacting expert on the basis of being able to turn out with prolific fortitude infinite strings of incomprehensible formulae calculated with micrometric precision from vague assumptions, which are based on debatable figures taken from inconclusive experiments, carried out with instruments of problematic accuracy by persons of doubtful reliability and questionable morality for the avowed purpose of annoying and confounding a hopeless chimerical group of fanatics known as geologists who are themselves the lunatic fringe surrounding the hard working mining operator”

-Quarterly News Bulletin of the Geological Society of South Africa, 1964

Preface

This PhD thesis entitled “*Regional Structure and Basin Development of the mid-Norwegian Volcanic Passive Margin: Anatomy of a Polyphased Rifted System*” has been submitted to the Department of Geosciences at the University of Oslo in accordance with the requirements for the degree of Philosophiae Doctor (PhD). The work presented herein was primarily carried out at the Centre for Earth Evolution and Dynamics (CEED) in the University of Oslo as a part of *OMNIS Project (Offshore Mid-Norway: Integrated Margin and Basin Studies)*. Candidate continued his research activity on this topic in Volcanic Basin Petroleum Research AS (VBPR) to further develop this thesis using a flow of new data. The seismic, magnetic, and gravity data presented in this study were kindly provided by TGS (Asker, Norway). Thus, the thesis is a product of long-term close partnership between UiO-CEED, VBPR and TGS. The principal supervisor for this work was Professor Jan Inge Faleide (UiO) with co-supervision from Sverre Planke (UiO and VBPR) and close collaboration with Laurent Gernigon (Geological Survey of Norway, Trondheim).

This thesis focuses on the structure, tectonostratigraphic development and petroleum prospectivity of the mid-Norwegian volcanic passive margin and is organized in two parts. The first part includes research motivation for this study, scientific background, regional geological framework, a summary of the research articles produced during this PhD work, concluding remarks as well as recommendations for further research projects. The second part comprises of published research articles authored by the candidate.

Dmitry Zastrozhnov

Oslo, 3 February 2020

To my granddad Stanislav, who is my role model both in life and profession of geologist

Summary

This PhD thesis presents a multidisciplinary study of the mid-Norwegian volcanic passive margin. A large dataset consisting of vintage and recently acquired 2D and 3D seismic surveys, potential field data, exploration wells, as well as updated regional biolithostratigraphic schemes has been jointly interpreted along the mid-Norwegian margin.

This PhD thesis shows that the long basin development of the mid-Norwegian margin was polyphased and was partly controlled by pre-existing deep inherited crustal blocks. After the regional NE Atlantic Late Jurassic-earliest Cretaceous rifting event, an extensional mode shifted from a stretching to a thinning, which was concentrated in the central basin province and accommodated in a deep crustal level along large detachment faults at least until the mid-Albian. At that time the main depocentres focused principally in the Møre Basin. Since the Cenomanian, the main deposition was concentrated in the Vøring Basin and was associated with the sequential migration of rifting axis towards the line of future breakup. The migration of rift axis is explained by strain hardening due to lithospheric cooling.

The regional mapping of the mid-Norwegian margin performed in the frame of this PhD work helped to revise previously defined structural elements (e.g. Hel Graben, Nyk High, Någrind Syncline, Utgard High) as well as identify and describe new structural elements mainly along the outer Vøring and Møre basins (e.g. Kolga High, Møre Marginal Plateau, Ervik Ridge, Grimm High, Hermod Basin). The outer Vøring and Møre basins have been interpreted to be underlain by a relatively thick continental crust below elevated marginal plateaus. The existence of such elevated crustal marginal plateaus along the outer mid-Norwegian margin better explains (1) the observed structural geometries (e.g. shallowing of the BCU towards the outer domain), (2) tectonostratigraphic evolution of the margin (e.g. discrete extensional events and crustal boudinage), (3) low magnetization of the crust, (4) subaerial emplacement of breakup-related lavas and geochemical signatures of mantle melts interaction with upper crust, (5) long time lag (100-110 Myr) between the necking event in the Late Jurassic-earliest Cretaceous and lithospheric breakup in the earliest Eocene.

Along the outer mid-Norwegian margin, the interpreted shallow crustal blocks possibly contain potential pre-Cretaceous plays. This may bring back exploration interest to this underexplored part of the mid-Norwegian margin. The tilted/rotated fault blocks here can form structural traps typical for the Haltenbanken area. For instance, it was shown that pre-Cretaceous rocks in the outer Møre Basin are located in the oil and gas maturation window in the present-day temperature field. Despite being buried under volcanic sequences, these potential pre-

Cretaceous rocks can be a potential source for intra-volcanic reservoirs similar to Rosebank discovery in the West of Shetland Basin and/or Lower to Upper Cretaceous-Paleocene reservoirs east of the volcanic domain (e.g. Tulipan discovery).

The insights of this PhD thesis will be both useful for generic understanding of passive margins and volcanic margins in particular, as well as for ongoing and future regional studies along the mid-Norwegian margin. This PhD thesis urges the importance of multidisciplinary approach utilizing different datasets and observations at various scales, while studying such complex first-order geological structures as passive margins.

Acknowledgements

I am deeply grateful to my main supervisor **Jan Inge Faleide** for inviting me to Oslo and involving me in OMNIS Project. I thank you for your mentoring, guidance, and patience during my work on this PhD thesis. I will be forever in debt to my co-supervisor **Sverre Planke** for providing me an opportunity to join Volcanic Basin Petroleum Research (VBPR), when the things did not look promising for me, and eventually put a bow on this PhD dissertation.

One of the cornerstone moments during this PhD time was getting to know **Laurent Gernigon** during a warm spring day of the year 2014 at EGU in Vienna. Since then, you became my closest collaborator, scientific inspirator and very good friend, to who I can call at any time of the day and discuss everything related to science. Even though you were not an official supervisor, your mentorship and contribution were invaluable for me. I hope sincerely that the completion of this PhD project will not terminate our positive collaboration.

My special thanks go to **Iakov Gogin**, my old good friend from student days, who, by lucky happenstance, turned out to work with the biostratigraphy of the NE Atlantic margins and kindly provided appropriate materials which I was long and unsuccessfully trying to get from other sources. My deepest appreciation goes to **Mansour Abdelmalak**, my officemate both at UiO and VBPR, for his help and support at early days of my PhD time. **Sergei Medvedev's** support and encouragement were always invaluable for me. You have such versatile personality: geologist, physicist, mathematician, politician, family relationships expert, philologist, singer, ice-hockey expert, Chelyabinsk-citizen; what always makes our regular get-togethers in Piano Bar (a bar with no piano actually) a great fun.

I would like to offer my special thanks to **Dani Schmid** for his grant assistance with modelling. **Michel Heeremans** is thanked for always being supportive and helpful, when it comes to software, seismic projects and data. I owe an important debt to **Trine-Lise Gørbitz** for her help with administrative stuff, which allowed me to stay in Norway and continue working on this PhD project. I wish to thank my dear colleagues from VBPR gang (**Ellen, Dwarika, Nina, Stephane, Benjamin, Ben, Romain**) for the nice and friendly working environment. TGS Company and particularly **Reidun Myklebust** are greatly acknowledged for providing geophysical data for this work.

I ought to transfer my special thanks to the beautiful city of Saint-Petersburg. My lovely friends and colleagues, **Bor'ka "Lysy" Petrushkov, Gleb "Frantsuz" Lipenkov, Dima "Khas" Khasis**, are always with me during my ups and downs, and distance has never been a problem to maintain our all-weather friendship. My former boss from VSEGEI, **Vasily Proskurnin**, is

greatly acknowledged for allowing me to go to Norway and his legendary and ironic (or not?) send-off speech (*“You can go whistle for going to Norway”*) still heartens me. All my other colleagues from “Taimyr Crew” and “Khopyor Crew” of VSEGEI are thanked for good days in the field. I also want to thank my master thesis supervisors, **Andrei Khudoley** and **Vladimir Verzhbitskiy**, for introducing me to the amazing world of science and structural geology in particular.

It would be unfair not to acknowledge my good dorm friends who made my daily life in Oslo funnier and easier, except some hard Saturday mornings. **Serioga, Alondra, Rashik, Zhenyochek, Vadya, Kolyamba, Mayka, Almirka, Andryucha, Tanyusha, Przemek**, you guys rock! If we wrote the screenplay from all our stories, it would be the greatest teen comedy ever. My teammates from OSI Futsal are thanked for the nice practices, tournaments and Friday beers we had together, which helped me to release stress and keep my body and spirit up. My very special thanks go to **J.** for making my life sweeter and bitter.

I want to express my deepest gratitude and love to my parents, **Olga** and **Andrei**, for their support, belief and motivation. Being a son of geologists is sometimes hard, but I am really happy to follow your steps. I would like to thank my grandfather **Stanislav**, a great person and geologist (yeah, too), who always listens to my complaints about work and gives invaluable advices and support. I love you very much! My big love and thanks go to my grandma, **Lyudmila**, who fortunately does not ask a lot of questions about my PhD (apparently because she is not a geologist), but always motivates me with her *“You know what you have to do”* at the end of our phone calls. Finally, I want to thank my little son **Romka**, who is actually about to become a teenager, while waiting the completion of my PhD, for his existence, smiles, hugs, laughter and love. You are sunshine of my life!

List of articles

Main publications

Article 1 – Published in Tectonics

Zastrozhnov, D.^{1,2}, Gernigon, L.³, Gogin, I.⁴, Abdelmalak, M. M.^{1,2}, Planke, S.^{1,2}, Faleide, J. I.¹, Eide, S.⁵ & Myklebust, R.⁶ (2018). Cretaceous-Paleocene Evolution and Crustal Structure of the Northern Vøring Margin (Offshore Mid-Norway): Results from Integrated Geological and Geophysical Study. *Tectonics*, 37(2), 497-528.

¹Centre for Earth Evolution and Dynamics, University of Oslo, Oslo, Norway

²Volcanic Basin Petroleum Research (VBPR), Oslo Science Park, Oslo, Norway

³Geological Survey of Norway (NGU), Trondheim, Norway

⁴RPS Ichron, Northwich, UK

⁵First Geo AS, Skøyen, Norway

⁶TGS, Asker, Norway

Article 2 – Published in Tectonophysics

Theissen-Krah, S.^{1,2}, Zastrozhnov, D.^{2,3}, Abdelmalak, M. M.^{2,3}, Schmid, D. W.⁴, Faleide, J. I.², & Gernigon, L.⁵ (2017). Tectonic Evolution and Extension at the Møre Margin–Offshore mid-Norway. *Tectonophysics*, 721, 227-238.

¹GEOMAR Helmholtz Centre for Ocean Research Kiel, Wischhofstr. 1-3, 24148 Kiel, Germany

²Centre for Earth Evolution and Dynamics (CEED), University of Oslo, Norway

³Volcanic Basin Petroleum Research (VBPR), Oslo Science Park, 0349 Oslo, Norway

⁴Physics of Geological Processes (PGP), University of Oslo, Norway

⁵First Geo AS, Skøyen, Norway

⁶Geological Survey of Norway (NGU), Leiv Erikssons vei, 39, Trondheim, Norway

Article 3 – Accepted in Marine and Petroleum Geology

Zastrozhnov, D.^{1,2}, Gernigon, L.³, Gogin, I.⁴, Planke S.^{1,2}, Abdelmalak, M.M.^{1,2}, Polteau, S.^{5,6}, Faleide, J.I.¹, Manton, B.² & Myklebust, R.⁷ (2020). Regional Structure and Polyphased Cretaceous-Paleocene Rift and Basin Development of the mid-Norwegian Volcanic Passive Margin. *Marine and Petroleum Geology*, 115, 104269.

¹Centre for Earth Evolution and Dynamics (CEED), University of Oslo, Norway

²*Volcanic Basin Petroleum Research (VBPR), Oslo Science Park, Oslo, Norway*

³*Geological Survey of Norway (NGU), Trondheim, Norway*

⁴*Ichron Ltd, Northwich, Cheshire, UK*

⁵*SurfExGeo, Oslo, Norway*

⁶*Institute for Energy Technology (IFE), Kjeller, Norway*

⁷*TGS, Asker, Norway*

Article 4 – Published in Geological Society, London, Special Publications

Millett, J. M.^{1,2}, Manton, B. M.¹, Zastrozhnov, D.^{1,3}, Planke, S.^{1,3}, Maharjan, D.¹, Bellwald, B.¹, Gernigon, L.⁴, Faleide, J.I.³, Jolley, D.W.², Walker, F.², Abdelmalak, M.M.¹, Jerram, D.A.^{3,5}, Myklebust, R.⁶, Kjølhamar, B.E.⁶, Halliday, J.⁷ & Birch-Hawkins, A.⁷. (2020). Basin Structure and Prospectivity of the NE Atlantic Volcanic Rifted Margin: Cross-border examples from the Faroe-Shetland, Møre and Southern Vøring Basins. *Geological Society, London, Special Publications*, 495.

¹*VBPR, Oslo, Norway*

²*Department of Geology & Petroleum Geology, University of Aberdeen, UK*

³*CEED, University of Oslo, Norway*

⁴*Geological Survey of Norway (NGU), Trondheim*

⁵*DougalEARTH Ltd, Solihull, UK*

⁶*TGS, Asker, Norway*

⁷*TGS, Surbiton, UK*

List of additional publications building up to main publications

Appendix A – Published in Tectonics

Abdelmalak, M.M.^{1,2}, Faleide, J.I.¹, Planke, S.^{1,2}, Gernigon, L.³, Zastrozhnov, D.^{1,2}, Shephard G.E.¹ & Myklebust, R.⁴ (2017). The T-Reflection and the Deep Crustal Structure of the Vøring Margin Offshore Mid-Norway. *Tectonics*, 36, 2497–2523.

¹*Centre for Earth Evolution and Dynamics (CEED), University of Oslo, Oslo, Norway*

²*Volcanic Basin Petroleum Research (VBPR), Oslo Science Park, Oslo, Norway*

³*Geological Survey of Norway (NGU), Trondheim, Norway*

⁴*TGS, Asker, Norway*

Appendix B – Published in Journal of Geophysical Research: Solid Earth

Abdelmalak, M. M.¹, Planke, S.^{1,2}, Faleide, J.I.¹, Jerram, D.A.^{1,3,4}, Zastrozhnov, D.¹, Eide, S.⁵ & Myklebust, R.⁶. (2016). The Development of Volcanic Sequences at Rifted Margins: New Insights from the Structure and Morphology of the Vøring Escarpment, mid-Norwegian Margin. *Journal of Geophysical Research: Solid Earth*, 121, 5212–5236.

¹*Centre for Earth Evolution and Dynamics, University of Oslo, Oslo, Norway*

²*Volcanic Basin Petroleum Research, Oslo Research Park, Oslo, Norway*

³*DougalEARTH Ltd., Solihull, UK*

⁴*Earth, Environmental and Biological Sciences, Queensland University of Technology, Brisbane, Queensland, Australia*

⁵*First Geo AS, Fornebu, Norway*

⁶*TGS, Asker, Norway*

Appendix C – an oral and poster talk presented at the 34th Nordic Geological Winter Meeting, January 8th-10th 2020, Oslo, Norway

Gernigon, L.¹, Zastrozhnov, D.^{2,3}, Planke, S.^{2,3}, Abdelmalak, M.M.³, Faleide, J.I.³, Maharjan, D.², Manton, B.², Myklebust, R.⁴. (2020). A New Nomenclature Map of the mid-Norwegian Margin: Status Update. *The 34th Nordic Geological Winter Meeting*, January 8th-10th 2020, Oslo, Norway.

¹*Geophysics Group, Geological Survey of Norway (NGU), Trondheim, Norway*

²*Volcanic Basin Petroleum Research (VBPR), Oslo, Norway*

³*Centre for Earth Evolution and Dynamics (CEED), Department of Geosciences, University of Oslo, Norway*

⁴*TGS, Asker, Norway*

Table of Contents

Preface	v
Summary	ix
Acknowledgements	xi
List of articles	xiii
1. Research Motivation: Towards a New Understanding of the Structure and Evolution of the mid-Norwegian Margin: Why Go Back to This Region?	21
2. Scientific Background	35
2.1. Mechanisms of rifting and their impact on rift basin architecture and development	35
2.2. Magma-poor vs magma-rich margins	38
3. Regional Geological Framework of the mid-Norwegian margin	45
4. PhD Publications. Synthesis.	49
5. Concluding remarks	55
6. Future work	56
References	59
Paper I	73
• <i>Cretaceous-Paleocene Evolution and Crustal Structure of the Northern Vøring Margin (Offshore Mid-Norway): Results from Integrated Geological and Geophysical Study</i>	
Paper II	107
• <i>Tectonic Evolution and Extension at the Møre Margin–Offshore mid-Norway</i>	
Paper III	121
• <i>Regional Structure and Polyphased Cretaceous-Paleocene Rift and Basin Development of the mid-Norwegian Volcanic Passive Margin</i>	
Paper IV	155
• <i>Basin Structure and Prospectivity of the NE Atlantic Volcanic Rifted Margin: Cross-border examples from the Faroe-Shetland, Møre and Southern Vøring Basins</i>	
Appendix A	197
• <i>The T-Reflection and the Deep Crustal Structure of the Vøring Margin Offshore Mid-Norway</i>	
Appendix B	227
• <i>The Development of Volcanic Sequences at Rifted Margins: New Insights from the Structure and Morphology of the Vøring Escarpment, mid-Norwegian Margin</i>	
Appendix C	255
• <i>A New Nomenclature Map of the mid-Norwegian Margin: Status Update</i>	

PART I

1. Research Motivation: Towards a New Understanding of the Structure and Evolution of the mid-Norwegian Margin: Why Go Back to This Region?

Passive margins are well spread in the world (Figure 1) and their importance is dictated by their industrial significance as they host almost 70% of discovered conventional hydrocarbon reserves (Mann et al., 2003; Levell et al., 2010). From a theoretical point of view, they are key elements in Wilson Cycles and an important component of the global geodynamic evolution (Wilson, 1966; Burke et al., 1976; Bradley, 2008). Thus, these structures are among the most scientifically attractive first-order geological structures on the Earth. The last four decades are characterized by technological advances and extensive data acquisition along passive margins and therefore a need for data analysis, interpolation, and compilation rises. This PhD thesis utilizes recently acquired data to refine the understanding of passive margins formation and structure using the example of the mid-Norwegian margin.

Figure 1. Worldwide distribution of different types of passive margins (modified from Berndt et al., 2019).

The formation of passive margins can be divided in two major stages: continental and oceanic rifting (drifting). The transition from continental to oceanic rifting, the continental breakup, represents a fundamental and complex process in the evolution of passive margins. Our planet offers examples of passive margins at different stages, from initial rifting to drifting and further fossilization in orogenic belts. The African and Baikal rift systems represent a few

modern examples of the initial stage of continental rifting, while mature stages of passive margin evolution are well presented worldwide (Figure 1; e.g. Atlantic, Arctic and Antarctic margins; Bradley, 2008).

The variety of world-wide examples of passive margins allows us to study controlling mechanisms and deformation styles at all stages of margins evolution. Drifting or post-breakup stage is usually characterized by thermal subsidence, mild basin inversion, modest sedimentation, and gravity-related re-adjustments while major tectonic activity shifts away from passive margins and concentrates in the oceanic domain. In contrast, during the pre-breakup stage passive margins are subjected to active stretching of the continental crust, and associated formation of sedimentary basins. The continental breakup can be associated with voluminous magmatism as is the case of volcanic (magma-rich) passive margins, or it can be nearly amagmatic as exemplified by magma-poor passive margins. Even though the phases of margin evolution are well understood, and different types of passive margins are present worldwide (Figure 1), the relationships between pre-breakup development of passive margins and processes responsible for eventual continental breakup often remain poorly understood and controversial. The mid-Norwegian margin is one of best areas to study these processes. It has a long and sometime polemic study history of mechanisms and structural styles of continental rifting and breakup-related magmatism. Newly acquired data ask for interpretation which may refine our understanding and resolve existing debates.

The mid-Norwegian volcanic rifted margin was subjected to extensive geological and geophysical studies since late 1960s (Blystad et al., 1995; Brekke, 2000; Faleide et al., 2010 and references therein). The start of exploration activity started in 1980s and concentrated mainly in the shallow shelf part (Halten and Dønna terraces and Trøndelag Platform). The first phase of exploration led to pioneer regional mapping of the basin structures and the first understanding of the pre- and post-breakup tectonic development of the mid-Norwegian margin (Gabrielsen et al., 1984; Bøen et al., 1984; Brekke & Riis, 1987). Despite a reasonable understanding of the proximal province, the mid-Norwegian margin was still poorly explored in its distal and outer parts. In the late 1970s and early 1980s, the results from Deep Sea Drilling Project (DSDP) and Oceanic Drilling Program (ODP) on the outer Vøring Margin and sparsely acquired 2D seismic and refraction data (Hinz et al., 1987; Mutter & Zehnder, 1988; Zehnder et al., 1990; Planke et al., 1991) contributed to major step in the primary understanding of volcanic margin formation (e.g. Eldholm et al., 1989; Skogseid & Eldholm, 1989).

Summarizing the existing data and performed studies, the Norwegian Petroleum Directorate (NPD) published a structural map of the Norwegian Sea Region in 1995 (Figure 2a)

(Blystad et al., 1995) where main structural elements were named and described. The Blystad et al. (1995) map became a regional standard which is still widely used and cited by both academic and exploration communities. However, at the time of map compilation, the structure of the deep shelf provinces (Møre and Vøring basins) of the mid-Norwegian margin was still poorly known in terms of pre-breakup development due to limited exploration activity in these provinces. Industry and academia focused on research in the deep basin settings and acquired additional seismic and well data only in late 1990s. The interpretations of this dataset resulted in numerous publications on the tectonostratigraphic development of the Møre and Vøring Basins (e.g. Lundin & Dore, 1997; Bjørnseth et al., 1997; Swiecicki et al., 1998; Ren et al., 1998, 2003; Brekke, 2000; Skogseid et al., 2000; Færseth & Lien, 2002; Gernigon et al., 2003, 2004). However, most of the study area was still lacking dense regional coverage of good-quality seismic and well data to provide reliable age for solid tectonostratigraphic framework and detailed basin architecture. Also, the old conventional seismic could not image the deep parts of the basin that extended up to 7-8 s twt.

Since late 1990s, one of the main directions of the mid-Norwegian margin research was the geological and geophysical understanding of the deep offshore Møre and Vøring basins (Figure 2a-b). The important point was to properly determine the crustal architecture, the geometry and evolution of the complex mid-Norwegian rifted system since the collapse of the Caledonian orogen (Walker et al., 1997; Ren et al., 1998; Osmundsen et al., 2002; Mosar, 2000, 2003; Mjelde et al., 2002, 2003, 2005, 2008, 2009; Gomez et al., 2004; Gernigon et al., 2004; 2006; Scheck-Wenderoth et al., 2007; Reynisson et al., 2010). More recently, the results of these studies led to a heated discussion on to what extent the mid-Norwegian volcanic rifted passive margin and volcanic margins in general can be compared to an Iberian type of magma-poor margins (Peron-Pinvidic et al., 2013; Nirrengarten et al., 2014; Peron-Pinvidic & Osmundsen, 2016, 2018; Gernigon et al., 2015, 2019; Osmundsen & Peron-Pinvidic, 2018; Theissen-Krah et al., 2017; Zastrozhnov et al., 2018; Guan et al., 2019). A key issue in this dispute is the interpretation of the sub-sedimentary and sub-volcanic structures of the deep Møre and Vøring basins, particularly the nature of the high-velocity ($V_p > 7.0$ km/s) lower crustal body (LCB) and associated deep seismic reflections (e.g. T-Reflection, Gernigon et al., 2003; Abdelmalak et al., 2017). One group of researchers interpreted these features as an exhumed serpentinized mantle domain in large parts of the deep Møre and Vøring basin provinces (Reynisson et al., 2010; Lundin & Doré, 2011; Rüpke et al., 2013; Peron-Pinvidic & Osmundsen, 2016, 2018) implying structural similarities with the magma-poor margins. Peron-Pinvidic & Osmundsen (2016) suggested, for example, that the onset of volcanism in the distal

margin was preceded by a phase of mantle exhumation and denudation and that the volume of magmatism between magma-poor and volcanic margin were not so significantly different. The other group of researchers doubted that the crust underneath the Møre and Vøring basins was so drastically thinned and suggested instead that the high-velocity lower crust underneath the basin may simply be the remnants of the exhumed old inherited crust later intruded by deep sills and magmatic cumulates (Nirrengarten et al., 2014; Abdelmalak et al., 2017; Gernigon et al., 2015, 2019; Petersen and Schiffer, 2016; Theissen-Krah et al., 2017; Zastrozhnov et al., 2018).

Another point of concern is whether the mid-Norwegian margin was involved in either several and discrete extension and cooling events during the Cretaceous-Paleocene (Lundin & Doré, 1997; Doré et al., 1999; Reemst and Cloetingh, 2000; van Wijk and Cloetingh, 2002; Gernigon et al., 2003, 2004, 2015, 2019; Zastrozhnov et al., 2018), or in a single phase of continuous lithospheric deformation leading to breakup which fits an Iberian-type model (Peron-Pinvidic & Osmundsen, 2018). Resolving this debate may lead to better understanding the thermal evolution of the mid-Norwegian margin and potential in application of classic basin evolution models.

For instance, the important question is whether the classic basin evolution model of McKenzie (1978) of uniform and instantaneous pure shear extension followed by thermal subsidence is applicable for the formation and subsidence history of the large and thick sedimentary Møre and Vøring basins. The McKenzie model is an important reference mechanism for first-order tectonic predictions in rifted margins (Allen & Allen, 2013). However, being generally favorable for rift basins such as the North Sea (Wood & Barton, 1983; Marsden et al., 1990; Nadin & Kusznir, 1995), proximal margins and sediment-starved passive margins such as the Galicia Margin in the Central Atlantic (Moullade et al., 1988), this model requires significant modifications for the complex tectonic settings such as the volcanic rifted mid-Norwegian margin. The formation of the large and thick sedimentary Møre and Vøring basins is associated with distinct periods of rifting and accelerated subsidence, which, together with voluminous breakup-related magmatism, cannot be reasonably explained by the theoretical subsidence predictions of the McKenzie model.

Figure 2a. NPD structural elements map of the mid-Norwegian margin (modified from Blystad et al., 1995, volcanic cover is removed).

Figure 2b. An updated structural elements map of the mid-Norwegian margin (pre-breakup structural elements) (Zastrozhnov et al., 2020; Gernigon et al., in prep.) compiled from the results of this PhD thesis.

In an exploration context, the understanding of the crustal nature, thermal and tectonic evolution, and sedimentary infill history of the Møre and Vøring basins, has major implications for petroleum systems. Recent interpretations of the pre-breakup development of the mid-Norwegian margin inspired by the Iberian model assumed thin or nearly absent pre-Cretaceous sedimentary sequences resting at greater depths on top of the exhumed serpentinitized mantle in large parts of the outer Møre and Vøring basins (Bergslien & Syvertsen, 2015; Peron-Pinvidic & Osmundsen, 2016). Therefore, such a model questions and limits the presence of regionally prolific Jurassic play in the outer basins (Bergslien & Syvertsen, 2015). In contrast, a model with moderately thinned crust in the outer Møre and Vøring basins still leaves a possibility for the prospective pre-Cretaceous sequence to be present at a relatively shallow depth within oil and gas maturation window and therefore may facilitate future exploration activities in the area (Theissen-Krah et al., 2017; Millet et al., 2020).

Over the last 15 years, a significant data accumulation including new and reprocessed 2D seismic reflection and refraction lines, large 3D seismic surveys, regional interpolated 3D volume (J-Cube MN), potential field data, new exploration wells, seabed sampling (Figure 3), and revised biostratigraphic schemes enabled an improved understanding of the mid-Norwegian margin. Particularly new and reprocessed seismic data achieved a better imaging of intra-volcanic sequences and sub-basalt basin geometries in the outer domain of the mid-Norwegian margin (Figure 4a-b), and deep basin structures in the central parts of the Møre and Vøring basins (Figure 5a-b).

Figure 3. Regional geological and geophysical dataset used in this PhD thesis. Grey lines - vintage and new reflection seismic (data courtesy of TGS), red polygons - 3D seismic surveys, brown polygon - J-Cube MN 3D volume (data courtesy of TGS), black lines - published refraction data, black dots - available well data, yellow dots - seabed sampling (data courtesy of VBPR).

The main goal of this PhD thesis was to compile, integrate and interpret this comprehensive geological and geophysical dataset to improve the understanding of processes controlling the formation and development of the deep Cretaceous-Paleocene basins of the mid-

Norwegian margin. During this integrated study 9 regional seismic horizons have been mapped out within the entire mid-Norwegian margin using 2D seismic and incoming brand-new 3D surveys, and have been calibrated with new wells, revised biostratigraphic charts and potential field data (Figures 6 and 7). The interpretations were further tested and validated by 2D potential field and geodynamic modelling. More specifically, this PhD thesis aims to:

- revise and identify structural elements in the distal and outer basins, which were poorly imaged and discussed in previous regional publications;
- establish an updated tectonostratigraphic framework for the volcanic rifted mid-Norwegian margin with particular emphasis on the Møre and Vøring basins;
- study crustal and basin architecture of the mid-Norwegian margin;
- understand the role of structural inheritance and pre-existing basin topography in the formation of the large sedimentary basins;
- discuss and review existing crustal nature and evolution scenarios for the interpreted sequence of tectonic events in the basins;
- evaluate validity of applying magma-poor Iberian type of passive margins scenario for the mid-Norwegian case;
- propose a conceptual model for the pre-breakup development of the Møre and Vøring basins;
- utilize all new observations and interpretations for reappraisal of the prospectivity of the yet underexplored Møre and Vøring basins.

Figure 4. An example of sub-basalt and intra-basalt imaging improvements in the outer Vøring Basin. (a) – original 2D seismic line, (b) – reprocessed seismic line. Data courtesy of TGS.

Figure 5. An example of deep basin imaging improvements in the Møre Basin. (a) Vintage GMNR line (1994). (b) AMS 3D survey (2019). Data courtesy of TGS.

Finally, an improved mapping of the mid-Norwegian margin presented in this PhD thesis led to the compilation of an updated nomenclature map of the mid-Norwegian margin in collaboration with Volcanic Basin Petroleum Research (VBPR), Geological Survey of Norway (NGU) and TGS (Figure 2b; Appendix C). This new regional map brings a great addition and update to Blystad et al. (1995) landmark map (Figure 2a) and will help both academia and industry geologists to be introduced to the up-to-date regional framework of the mid-Norwegian margin.

Figure 6. (a) – Interpreted regional Base Cretaceous Unconformity surface (in twt) showing basin configuration of the mid-Norwegian margin and its adjacency. (b) – An example of the integrated seismic-gravity-magnetic interpretation, where the regional BCU horizon is jointly interpreted and calibrated with potential field data.

Figure 7. Regional seismic grids interpreted during this PhD work for the entire mid-Norwegian margin.

2. Scientific Background

2.1. Mechanisms of rifting and their impact on rift basin architecture and development

The mechanism of rifting and subsequent formation of passive margins are generally attributed to stretching and thinning of the continental lithosphere (e.g. McKenzie, 1978). Rifts are the centres of crustal extension, which can be aborted or instead lead to breakup, oceanic spreading and the formation of passive margins.

The instantaneous and uniform extension of the lithosphere (McKenzie, 1978) is considered as the simplest reference model for the rift formation and development. This model implies symmetrical and coaxial stretching, and therefore is compliant with the condition of pure shear (Figure 8a). The upper crust is affected by brittle faulting, whereas the lower crust is extended in ductile manner. The maximum thermal gradient is observed in the centre of the rift and is related to passive upwelling of hot asthenosphere, which is required to maintain isostatic equilibrium. The tectonic subsidence in rift basins, which is an isostatic response to the stretching of the continental lithosphere, comprises of two components: an initial fault-controlled (syn-rift) subsidence and post-rift thermal subsidence. The initial fault-controlled subsidence is instantaneous and is dependent on the initial crustal to lithospheric thickness ratios and the stretching factor β , which is the ratio of the initial to the final crustal/lithospheric thickness. The following post-rift thermal subsidence is a result of thermal relaxation (cooling) of the lithosphere to its pre-stretching conditions and depends only on the applied stretching factor (β). It decreases exponentially with time as a consequence of a decrease of heat flow.

While being favorable for first-order predictions of crustal thicknesses and subsidence histories in many rift basins, the McKenzie model assumptions do not always predict variations in continental margin architecture (e.g. conjugate asymmetries), modes of crustal thinning, accelerated subsidence and syn-rift uplift (e.g. Lister et al., 1986). To explain structural asymmetries, which often occur in rift basins and across conjugate margins, Wernicke (1981, 1985) proposed a model of continental extension by simple shear along large crustal detachments. In such a simple shear model, the continental lithosphere extends asymmetrically, and the zone of brittle crustal stretching is laterally offset from the zone of ductile lower crustal or subcrustal stretching along the detachment (Figure 8b). Therefore, the maximum thermal gradient is also offset from the centre of the rift basin in the simple shear model. Consequently, it has significant implications for uplift and subsidence patterns in rift basins. In this case, crustal stretching produces the fault-controlled syn-rift subsidence in the middle of the rift basin, while the lower crustal and subcrustal stretching causes asthenospheric upwelling and

tectonic uplift within the proximal part of the rift basin (Figure 8b). Therefore, subsequent thermal relaxation and post-rift subsidence is spatially separated from the initial fault-controlled subsidence and occur above the sub-crustal stretching zone.

Figure 8. Idealized models for crustal extension. (a) - pure shear model of McKenzie (1978), (b) - simple shear model of Wernicke (1981, 1985), (c) - hybrid model (e.g. Coward, 1986): simple shear in the crust and pure shear in sub-crustal lithosphere. Modified from Fossen (2016) and Lister et al. (1991).

Lister et al. (1986) developed a concept of simple shear lithospheric extension along major detachment faults and proposed the existence of so-called upper-plate and lower-plate passive margins (Figure 9). The asymmetrical shape of conjugate passive margins, according to Lister et al. (1986), is controlled by the dipping direction of the master detachment faults either towards the ocean or the continent. The upper-plate passive margin geometry is controlled by a continent-ward dipping detachment fault, which comprises crust devoid of structures in its hanging wall and is characterized by a narrow continental shelf. The lower-

plate passive margins are characterized by a wide continental shelf and are controlled by seaward dipping detachment faults (Figure 9). The lower-plate margins comprise exhumed deep crystalline rocks, which are overlain by tilted blocks with fragments of the upper plate. Lister et al. (1986) also highlighted a second-order structuring of passive margins where more than one detachment system act, allowing the formation of high-standing undeformed regions. In this case, a so-called continental ribbon is bounded by oppositely dipping detachments and a marginal plateau is bounded by two detachments zones with the same dip (Figure 10).

Figure 9. Detachment-fault model of passive margins with lower-plate and upper plate characteristics. Modified from Lister et al. (1986).

Figure 10. Paired detachment systems model of Lister et al. (1986) explaining possible variations in the crustal architecture of passive margins.

However, compared to the pure shear model, the large-scale simple shear model cannot explain the thermal subsidence spatially superimposed on the initial fault-controlled syn-rift subsidence, which is the case for many rift basins worldwide (e.g. North Sea). To explain this involving detachment faults, hybrid models were proposed (Coward, 1986; Kusznir et al., 1991), where the upper crust is deformed by simple shear along major detachments, while the

ductile lower crust and subcrustal lithospheric mantle are extended coaxially by pure shear (Figure 8c).

Generally, pure shear, simple shear and/or combined simple-pure shear models are the main models of rift basin development. However, to explain a variety of geological and geophysical observations accumulated over last three decades due to extensive seismic acquisition and drilling within the passive margins worldwide, some modifications to these models were provided including: a) depth-dependent stretching, where stretching factors vary in different lithospheric levels (Rowley & Sahagian, 1986; Huisman & Beaumont, 2002, 2008); b) magmatic activity influence caused by melting, igneous underplating or dynamic support from a hot asthenosphere (White & McKenzie 1989; Skogseid et al., 2000; Geoffroy et al., 2015); c) lithospheric phase transitions caused by strong changes in the pressure-temperature conditions (e.g. Podladchikov et al., 1994; Baird et al. 1995; Kaus et al., 2005; Hartz et al., 2017)

2.2. Magma-poor vs magma-rich margins

Passive margins can be generally divided into magma-poor and magma-rich types (e.g. Geoffroy, 2005; Reston, 2009; Franke, 2013; Peron-Pinvidic et al., 2013). This classification does not only concern the deformation mode and magmatic budget during continental breakup, but also reflects important variations in terms of structural styles and subsidence histories. While having similar structural styles in the proximal parts, magma-poor and magma-rich margins display contrasting differences in the distal and outer parts (e.g. Franke, 2013; Geoffroy et al., 2015; Gernigon et al., 2019).

Magma-poor margins are generally characterized by extreme rates of crustal thinning, which progressively increases towards the oceanic domain and promotes the presence of a broad zone of partially serpentinized and exhumed mantle along top-basement detachment faults in the distal and outer domains (e.g. Boillot & Froitzheim, 2001; Perez-Gussinye et al., 2003; Reston, 2007, 2009; Reston & Manatschal, 2011; Peron-Pinvidic et al., 2013) (Figure 11a). The magma-poor margins generally have low to moderate sediment accumulation rates, deep to ultra-deep-water continental breakup and very small to moderate volumes of magmatism (e.g. Galicia Margin: Manatschal & Bernoulli, 1999; Mohn et al., 2015; Goban Spur, North of Biscay: de Graciansky et al., 1985).

The magma-rich margins represent the majority of passive continental margins (Menzies et al., 2002; Skogseid, 2001; Franke, 2013) and are generally characterized by sub-

aerial continental breakup, which results into formation of thick volcanic wedges of seaward-dipping reflectors (SDRs) along the proto-breakup axis (Hinz, 1981; Mutter et al., 1982; Eldholm et al., 1989), extensive emplacement of mafic sills and dikes in the crust and sedimentary basins (Planke et al., 2005) and the presence of high V_p (>7.0 km/s) and high-density lower crustal bodies (LCB) (e.g. Mjelde et al., 2007, 2009, 2016; Breivik et al., 2014; Abdelmalak et al. 2017) (Figure 11b). The sedimentation rates in sag-basins within the magma-rich margins are usually much higher compared to magma-poor margins (e.g. NE Atlantic vs Iberian-Biscay margins: Straume et al., 2019). The magmatism in magma-rich margins are often associated with large igneous provinces (LIPs) (e.g. Eldholm & Grue, 1994; Menzies et al., 2002) and occurs in relatively short geological time interval of 1-3 Myr (Franke, 2013; Abdelmalak et al., 2016a). The origin and composition of LCBs is debated and is commonly interpreted to represent either breakup-related mafic underplating (e.g. White & McKenzie, 1989; Eldholm et al., 1989; Skogseid et al., 1992), old inherited high-grade crust (Gernigon et al., 2003; Ebbing et al., 2006; Mjelde et al., 2016; Petersen and Schiffer, 2016; Zastrozhnov et al., 2018), highly intruded lower crust (White et al., 2008; Abdelmalak et al., 2017) or serpentized mantle (Ren et al., 1998; Reynisson et al., 2010; Péron-Pinvidic & Osmundsen, 2016). Such a discussion on the origin of LCBs in the magma-rich margins is particularly due to poor seismic imaging of deep basin structure in the outer domain below a thick volcanic pile and non-uniqueness of petrophysical values of the crust obtained from potential field modelling exercises and refraction experiments (e.g. Saltus & Blakely, 2011; Peron-Pinvidic et al., 2016).

Figure 11. Two archetypal end-members of passive margins. (a) – Schematic crustal section along magma-poor “Iberian type” passive margin, which is characterized by progressive thinning of continental crust and exhumation and serpentinization of underlying lithospheric mantle, (b) – Schematic crustal section along the outer part of the “mid-Norwegian type” of magma-rich volcanic margins, which is characterized by voluminous extrusive and intrusive magmatism and presence of enigmatic high V_p (>7 km/s) lower crustal body (LCB). Modified from Gernigon et al. (2006).

A significant reappraisal of the passive margins allowed Peron-Pinvidic et al. (2013) and Peron-Pinvidic & Manatschal (2019) to postulate global first-order similarities both between magma-poor and magma-rich margins. As a corner-stone model they implied the structure and development of the well-studied and drilled Newfoundland-Iberian conjugate margin, which is traditionally referred as “archetypal magma-poor margins” (Peron-Pinvidic et al., 2013). They propose that each passive margin goes through the same sequence of deformations starting from (1) the stretching in a proximal domain, followed by (2) extreme crustal thinning and mantle exhumation in the distal domain, and (3) subsequent magmatism and oceanization in the outer and oceanic domains respectively. These phases assume a general continuum of crustal and lithospheric deformations which migrate successively oceanward until continental breakup is achieved. Local structural and evolution variations between passive margins or their segments, according to Peron-Pinvidic et al. (2013), can be explained by the

crustal inheritance influence, extension rates, thermal and compositional lithospheric states and rheology. In their model, the volcanic rifted mid-Norwegian margin resembles the Iberian type of passive margins (Figure 12a) displaying similar crustal architecture, and sequence and rates of deformational events, with only contrasting difference in the voluminous magmatic phase in the outer domain in the mid-Norwegian margin (Peron-Pinvidic & Osmundsen, 2016, 2018).

Figure 12. Two recent crustal models proposed for the mid-Norwegian margin. (a) – “Iberian type” model suggesting similarities with the magma-poor margins, where the distal and outer domains represent zone of exhumed/serpentinized mantle with no continental crust preserved. The deep T-reflection in this case represents a top of exhumed and serpentinized lithospheric mantle. Modified from Peron-Pinvidic et al. (2016, 2018). (b) – “Marginal Plateau” model proposed in this PhD thesis (Zastrozhnov et al., 2020). The distal domain is characterized by thinned crust with local mantle serpentinization windows, while the outer domain represents a marginal plateau, which is an elevated crustal block. The deep T-reflection in this case represents a top of inherited high Vp lower crustal body mixed with breakup-related mafic and ultra-mafic material.

However, such a uniform model for the development of passive margins was extensively questioned, especially for the case of polyphased and volcanic (magma-rich) passive margins (e.g. Geoffroy et al., 2015; Davis & Lavier, 2017; Guan et al., 2019) and the mid-Norwegian margin in particular (Gernigon et al., 2015, 2019; Theissen-Krah et al., 2017; Zastrozhnov et al., 2018, 2020). Geoffroy et al. (2015) summarized observations from the South and North Atlantic volcanic margins and concluded that sharp transition from continental rifting to oceanic crust formation does not require mantle exhumation and appears to be a fundamental feature of volcanic passive margins. Geoffroy et al. (2015) argued that extreme lithospheric extension is not the one and only prerequisite for continental breakup. This can simply be achieved by magma support (e.g. Buck & Karner, 2004) expressed by voluminous mafic diking into the crust which therefore may localize the deformation and increase rapidly the lithosphere (e.g. Callot et al., 2001; Yamasaki & Gernigon, 2009). In their numerical modelling, Davis & Lavier (2017) also concluded that a thick continental crust preserved in the outer domain may be a favorable factor for the formation of volcanic passive margins. Moreover, subaerial breakup conditions in the outer domains of the magma-rich margins should be supported by the presence of relatively elevated continental block there, which is not the case for deep-water outer domains of Iberian-type margins. Geochemical study of breakup-related volcanics in the mid-Norwegian margin showed that magma mixing between asthenospheric and upper crustal melts must have occurred prior to their emplacement in the Vøring and Møre Marginal highs (Meyer et al., 2007, 2009; Abdelmalak et al., 2016a; Bakke, 2017). Also it was shown that the LCBs and associated deep crustal reflections (*T*-Reflection, *UH*-Reflection) in the outer and distal domains of the mid-Norwegian margin were affected by faulting long before breakup meaning that they were present before significant thinning of the crust and the initiation of magmatism (Gernigon et al., 2003, 2004; Zastrozhnov et al., 2018, 2020). While the *T*-Reflection coincides with high gravity signal, its regional magnetic signature in the mid-Norwegian margin is markedly low compared to exhumed mantle ridges in the Iberian margin (Zastrozhnov et al., 2020). Combination of these observations does not support breakup-related magmatic underplating as a primary model for the LCB emplacement and also cast doubts on serpentinized mantle origin of the LCB and therefore support its primarily crustal origin.

The final architecture and timing of rift systems of magma-poor and magma-rich margins may also reflect first-order orogenic inheritance. As such, Chenin et al. (2015) showed that distribution of magma-rich margins in the North Atlantic coincide with the Caledonian orogenic lithosphere, while magma-poor margins tend to develop over the Variscan orogenic lithosphere. Chenin et al. (2015) also documented a significant time lag between necking and

breakup in the NE Atlantic magma-rich margins, which in the order of 80-100 Myr or more, while for the NE Atlantic magma-poor margins this deformation period does not exceed 45 Myr. This notable difference in the pre-breakup deformation duration between magma-poor and magma-rich margins, however, is not reported in the South Atlantic passive margins (Mohriak et al., 2008; Mohriak & Leroy, 2013).

3. Regional Geological Framework of the mid-Norwegian margin

The mid-Norwegian margin consists of three large segments: Møre, Vøring and Lofoten-Vesterålen, which are about 400-500 km each and separated respectively by the Jan Mayen Corridor and Bivrost Lineament (Figure 2b). These segments developed along the mid-Norwegian passive margin display significant contrasts in structural style on both lithospheric and basin scales, timing of basin subsidence and basin infill history.

The Møre Margin has a narrow pre-Cretaceous platform domain (Møre Platform) which to the west passes into the wide and deep Cretaceous Møre Basin. To the west towards the continent-ocean ‘boundary’ (COB), the Møre Basin is confined by the wide and shallow pre-Cretaceous Møre Marginal Plateau (Gernigon et al., 2015; Theissen-Krah et al., 2017; Manton et al., 2018; Millet et al., 2020), which is partly covered by the breakup-related volcanics of the Møre Marginal High (Figure 2b).

Compared to the Møre Margin, the Vøring Margin has a wide proximal domain (Trøndelag Platform), which is bounded to the west by the Halten and Dønna terraces. Further west the wide distal domain is formed by the large and deep Cretaceous Vøring Basin, which is dissected by an elongated boundary fault system of the Fles Fault Complex. The outer part of the Vøring Basin extends to COB and represents a series of Late Cretaceous-Paleocene structural highs (Gjallar Ridge, Nyk High, Grimm High, Skoll High) separated by sub-basins (Hel Graben, Vigrid and Någrind Synclines, Fenris Graben).

The Lofoten-Vesterålen Margin is characterized by narrow and elevated Cretaceous sedimentary basins, bounded by basement highs (Figure 2b). The thickness of continental crust in the Lofoten-Vesterålen Margin appears to be much larger than in the greatly extended Møre and Vøring basins further south (Faleide et al., 2008; Tsikalas et al., 2012; Maystrenko et al., 2017a, 2017b).

The formation of the mid-Norwegian margin together with other NE Atlantic adjacent and conjugate continental margins is the result of the continental breakup between Baltica and Laurentia in the early Eocene (Faleide et al., 2008; Abdelmalak et al., 2016a; Gernigon et al., 2019). The pre-breakup rifting development of the mid-Norwegian margin is characterized by a series of extensional events developed after the collapse of the Scandinavian Caledonides in the Late Devonian-Early Carboniferous and before the late Paleocene-earliest Eocene formation of a distal volcanic margin (Doré et al., 1999; Brekke et al., 2001; Faleide et al., 2008; Tsikalas et al., 2012; Gernigon et al., 2019).

The late Paleozoic-early Mesozoic rifting history of the mid-Norwegian margin was controlled by the pre-existing Caledonian grain. It is expressed by apparent structural

coincidence between rift localization and orientation, and ancient Caledonian axes (Doré et al., 1997; Schiffer et al., 2019). It is believed that the first rifting phase was in the mid-Carboniferous time, but it remains poorly constrained in the mid-Norwegian margin and was suggested based on structural observations onshore East Greenland (e.g. Stemmerik et al., 1993; Stemmerik, 2000). The seismic evidence of subsequent Permo-Triassic extension is well documented in the Trøndelag Platform and was interpreted to culminate in the Early Triassic (Müller et al. 2005).

The Middle to Late Triassic and earliest Jurassic represent a post-rift phase with relative tectonic quiescence, when the area experienced a prominent paleogeographic change from marine to coastal and lacustrine sedimentary environments (Brekke et al., 2000; Müller et al. 2005). This quiescent period was followed by intermittent and moderate Early and Middle Jurassic stretching events (Blystad et al., 1995; Martinius et al. 2001; Ravnås et al., 2014). The Lower-Middle Jurassic strata (mainly sandstones) reflect shallow marine deposition prior to the onset of the next major rift phase.

The Late Jurassic–earliest Cretaceous rifting event is related to northward propagation of Atlantic rifting and reflects major extension and crustal stretching in the NE Atlantic (Lundin & Doré, 2011; Stoker et al., 2017). The stretching of the crust and block faulting created accommodation space where syn-rift deposits were accumulating. During this period the development of large and deep Cretaceous sedimentary basins in the distal part of the mid-Norwegian margin (e.g. the Møre and Vøring basins) was initiated (Blystad et al., 1995; Brekke, 2000; Zastrozhnov et al., 2020). Continuous stretching led to the localization of the central basin high system (e.g. Ervik Ridge, Vigra High, Slettringen Ridge; Figure 2b), activation of border faults (e.g. Klakk Fault Complex), and initial structural individualization of marginal plateaus in the outer basin domains (Zastrozhnov et al., 2020). Since the Barremian and up to the mid-Albian the extension mode shifted from stretching to severe thinning which was localized and accommodated by large detachment faults along the central basin high system. In the Early Cretaceous the outer portion of the mid-Norwegian margin including the marginal plateaus and isolated basement highs (e.g. Rån Ridge, Vigrid Syncline, Nyk High) were affected by local extensional episodes (e.g. Gernigon et al., 2003; Zastrozhnov et al., 2018; 2020), which, however, were accompanied by a moderate stretching of the crust. Since the mid-Albian up to recent the central Møre Basin was regionally subjected to thermal cooling and passive sedimentation without any significant tectonism. The thermal cooling in the central Vøring Basin lasted until the mid-Cenomanian/Turonian, when it was followed by renewed

extensional phase affecting mainly the Rås and Træna basins, and locally the Gjallar Ridge (Zastrozhnov et al., 2018; 2020).

Since the mid-Campanian and up to the final breakup and onset of sea floor spreading in the earliest Eocene, the main deformation axis started to progressively migrate from the central Vøring Basin towards the line of breakup, which resulted in the formation of such structural elements as the Nyk High, the Hel Graben, the Hel Terrace and the Grimm High and associated sub-basins. The Late Cretaceous-Paleocene structural development of the outer Vøring Basin has been reconsidered and attributed to extensional processes (Zastrozhnov et al., 2018; 2020), rather than compressional buckling (Brekke, 2000; Lundin et al., 2013). In the outer Møre Margin and western termination of the Jan Mayen Corridor, the presence of the Late Cretaceous-Paleocene extension is questionable, but if present, it was likely localized and expressed by block faulting in the Kolga High and in the sub-basalt Bylgja Ridge.

The Late Cretaceous-Paleocene extension was followed by the continental breakup and associated extrusive and intrusive magmatism in the mid-Norwegian margin. The final breakup was diachronous and initiated in the Thanetian at 57-58 Ma in the Møre segment and later propagated to the outer Vøring Margin in the early Ypresian at 56-55 Ma (Gernigon et al., 2019; Zastrozhnov et al., 2020). The breakup-related magmatism is characterized by emplacement of sub-aerial and submarine volcanic facies, whose lateral distribution and thickness variations were strongly controlled by the pre-existing basin topography and paleobathymetry, and movements along the initiated oceanic fracture zones (Abdelmalak et al., 2016b). The cause of breakup-related magmatism in the NE Atlantic is still debated and is usually attributed to rift-related processes and/or alternative sub-lithospheric processes (White, 1992; Skogseid et al., 2000; Meyer et al., 2007; Foulger et al., 2005; Foulger & Anderson, 2005).

The drifting (post-breakup) stage of the development of the mid-Norwegian margin is characterized by thermal cooling and subsidence of the sedimentary basins (Faleide et al., 2008). However, a series of mid-Cenozoic compressional events affected the Vøring Basin and Jan Mayen Corridor in the mid-Eocene, Oligocene, mid-Miocene and Pleistocene (Lundin & Doré, 2002; Doré et al., 2008). It is expressed by basin inversion structures (Helland-Hansen Arch, Modgunn Arch, Naglfar Dome etc.) and local reactivation of structural highs along the Fles Fault Complex (e.g. Vimur and Utgard highs) and in the proximal domain (Nordland Ridge). These compressional events could be the results of ridge push and coeval spreading reorganization in the oceanic domain, and their notable absence in the Møre Basin was explained by strain partitioning along the Jan Mayen Fracture Zone (Doré et al., 2008).

The Quaternary evolution of the mid-Norwegian margin is closely linked to the northern hemisphere glaciation where large and thick glacial depocentres expanded over the Norwegian Sea (Hjelstuen et al., 2004). This period was also associated with neotectonic activity including large landslides (Laberg & Vorren, 2000; Dahlgren et al., 2002; Haflidason et al., 2004; Bellwald et al., 2019) and vertical movements of Scandinavia caused by isostatic response to ice sheets during glacial phases (Mangerud et al., 2011; Fjeldskaar & Amantov, 2017; Medvedev et al., 2019).

4. PhD Publications. Synthesis.

This article-based thesis contains two first-authored publications, one second-authored paper and one third-authored paper. One of the first-authored papers was published in *Tectonics* in 2018 (Zastrozhnov et al., 2018 – Paper 1) and the second has been recently published in *Marine and Petroleum Geology* (Zastrozhnov et al., 2020 – Paper 3). The second-authored paper (Theissen-Krah et al., 2017 – Paper 2) was published in *Tectonophysics* in 2017. The third-authored paper has been recently published in *Geological Society of London Special Publication* (Millet et al., 2020– Paper 4).

The seismic mapping results presented in the main papers of this PhD were also integrated in two fifth-authored papers, which are included in the Appendix A and B. The first paper was published in *Tectonics* in 2017 (Abdelmalak et al., 2017 – Appendix A) and the second paper was published in *Journal of Geophysical Research: Solid Earth* in 2016 (Abdelmalak et al., 2016b – Appendix B).

The concepts of all the papers were developed within the same project. All together these papers highlight the importance of multidisciplinary integrated study of the mid-Norwegian margin by combining observations from the crustal and lithospheric scale to sedimentary/volcanic and prospect levels. These papers mutually contribute to each other and form a basis for an updated regional tectonostratigraphic framework of the mid-Norwegian margin and the Møre and Vøring basins in particular.

Paper I “Cretaceous-Paleocene Evolution and Crustal Structure of the Northern Vøring Paper (Offshore Mid-Norway): Results from Integrated Geological and Geophysical Study” investigates the northern Vøring segment of the mid-Norwegian margin. A multidisciplinary study including detailed interpretation of new and reprocessed 2D seismic data, a complete biostratigraphic re-evaluation of released borehole data and potential field modelling of a representative regional transect was performed to constrain the tectonostratigraphic framework and crustal architecture of this yet understudied segment of the mid-Norwegian margin. The seismic interpretation results show evidence for Barremian-Aptian (?) faulting activity in the Någrind Syncline. The present-day configuration of the Någrind Syncline was explained by extensional processes enhanced by a Late Cretaceous-Paleocene footwall uplift of the adjacent Nyk High. The possible mid-Cenomanian-Santonian extension in the Træna Basin was assumed to explain high tectonic subsidence rates obtained from backstripping analysis. The seismic interpretation supports the presence of biostratigraphically questionable thick Paleocene strata in the Hel Graben. Both structural observations and results from potential field modelling do not support the presence of the wide serpentized mantle

domain in the study area. The high-velocity and high-density LCBs and associated deep crustal reflections were interpreted to represent old high-grade Caledonian crust. The specific crustal configuration and basin architecture in the northern Vøring margin was explained by the crustal boudinage model where the main sub-basins underlain by significantly thinned crust were developed in between rigid and thick crustal blocks (“buffers”). The narrow serpentinitized mantle ‘windows’, if present, were likely localized below these sub-basins.

Paper II “Tectonic Evolution and Extension at the Møre Margin - Offshore mid-Norway” presents the study of the Møre segment of the mid-Norwegian margin. An extensive compilation of new and reprocessed 2D seismic data was used to better understand the development of the Møre Margin. The potential field and forward basin modelling of a representative crustal transect across the central Møre Margin was performed to reconstruct the crustal architecture and basin development and to estimate stretching factors for each regional rifting phase. The results support the presence of the elevated pre-Cretaceous domain in the outer Møre Basin, which is referred as the Møre Marginal Plateau. This study also shows that the main deposition in the study area took place in the Early Cretaceous and was concentrated in the Møre Basin along the flanks of deep-seated structural highs. Since the Cenomanian the main subsidence phases and associated depocentres migrated northwards to the Vøring Basin. The results of the study also show that the observed and modelled crust are too thick to favor the outer portion of the LCB to be of serpentinitized mantle origin (e.g. a zone of exhumed continental mantle). In turn, the distal LCB was interpreted to be related to magmatic intrusion into the inherited lower crust. Finally, the stretching factors along the transect obtained from different approaches (forward modelling, backstripping and crustal thinning) were compared and discussed.

Paper III “Regional Structure and Polyphased Cretaceous-Paleocene Rift and Basin Development of the mid-Norwegian Volcanic Passive Margin” presents a regional study of the large Cretaceous-Paleocene sedimentary provinces of the Møre and Vøring basins. The paper uses an extensive dataset of new and reprocessed 2D/3D seismic surveys and new and re-evaluated well biostratigraphy to investigate crustal and basin architecture of the Møre and Vøring basins and examine 4D sequential evolution of the Cretaceous-Paleocene sedimentary infill. Several new structural elements have been identified and described in the central and outer Møre and Vøring basins including the Ervik Ridge, Bylgja Ridge, Kolga High, Hevring High and Dufa High (Figure 2b). Previously defined structural elements in the outer Vøring Basin (e.g. Rån Ridge, Vigrid Syncline and Gjallar Ridge) have been revised. Distal elevated marginal plateaus defined at the BCU level were suggested to present in the outer Møre and

Vøring basins. The paper shows that the Early Cretaceous to Paleocene evolution of Møre and Vøring basins is associated with episodic phases of extensional events separated by intermediate cooling phases. This study also shows that the Vøring and Møre basins, in details, have different structural and sedimentary evolution. Regional Late Jurassic-earliest Cretaceous rifting was proposed to continue until mid-Albian time, when it was mainly concentrated and accommodated along the central high system in the Møre and Vøring basins. The high sedimentation and subsidence rates in Cenomanian-Santonian in the Vøring Basin are explained by additional thinning event likely accommodated in the lower crust. Final Late Cretaceous-Paleocene rifting affected mainly the outer Vøring Basin but is also proposed for the outer Jan Mayen Corridor (Kolga High) and the outer Møre Basin (Bylgja Ridge). The overall configuration of the Møre and Vøring basins is explained in the concept of Lister et al. (1986, 1991) where large-scale detachment faults facilitate individualization of distal elevated continental blocks (marginal plateaus). The presence of such elevated crustal areas may explain emplacement of sub-aerial breakup-related volcanics, which contain traces of significant crustal contamination. Therefore, the applicability of Iberian-type margin scenario for the pre-breakup development of the mid-Norwegian margin was strongly questioned.

Paper IV “Basin Structure and Prospectivity of the NE Atlantic Volcanic Rifted Margin: Cross-border examples from the Faeroe-Shetland, Møre and Southern Vøring Basins” discusses and summarizes new seismic mapping results done for the Faeroe-Shetland, Møre and southern Vøring basins in terms of basin structure and hydrocarbon prospectivity and cross-border comparisons between the UK and Norwegian sectors. The work mainly focuses on new large 3D surveys alongside extensive 2D surveys recently acquired in the mid-Norwegian margin and Faeroe-Shetland Basin. New structures in the yet underexplored areas within the Møre and southern Vøring basins were described and compared with regionally proven hydrocarbon plays. Three major untested basin-margin structural highs (the Rån Ridge, the Kolga High and the Bylgja Ridge) have been interpreted to comprise potential pre-Cretaceous rocks within drillable depths. In the outer Møre Basin the presence of the sub-volcanic elevated pre-Cretaceous crustal domain within the present-day oil and gas maturation window could facilitate hydrocarbon generation and subsequent migration into intra-volcanic reservoirs. Such conditions were clearly demonstrated in the notable Rosebank discovery in the Faeroe-Shetland Basin. Detailed regional mapping of the Base and Top Basalt horizons shows the presence of areas where the basalt thickness is only several hundred meters or even down to zero meters (e.g. the Kolga High) and therefore highlights targets with reduced volcanic-related drilling risks. The Upper Cretaceous-Paleocene play has the main exploration interest both in the mid-

Norwegian margin and Faroe-Shetland Basin comprising turbidite systems of reservoir quality with new potential structures identified within the Helland Hansen and Modgunn arches. Furthermore, high-amplitude saucer-shaped clastic intrusions and clear brightening of glacial sands in the Neogene-Quaternary interval mapped out in the Møre and southern Vøring basins may provide additional exploration interest for the shallow targets, which are proven plays in the adjacent Northern North Sea.

The paper in the Appendix A “The T-Reflection and the Deep Crustal Structure of the Vøring Margin, Offshore mid-Norway” focuses on the regional distribution and morphological features of the so-called deep *T*-Reflection in the Vøring Margin and subsequently discusses possible explanations of its origin. Integrated seismic-gravity-magnetic interpretation of the extensive geophysical dataset helped to identify different morphologies of the *T*-Reflection including single and multiple reflections, smooth, rough, faulted and domal shapes. The distinct correlation between gravity highs and positive geometries of the *T*-Reflection supports that the reflection represents a high-impedance boundary associated with a high-density and high-velocity body. Indeed, the *T*-Reflection in over 50% of the outer Vøring Margin corresponds to the top of the lower crustal body defined as the top of the 7+ km/s velocity layer in refraction data. Notably, there is no significant magnetic anomaly associated with the *T*-Reflection. The seismic mapping results also show the spatial correlation between *T*-Reflection, magma intrusions and fault planes along basement highs. The results of the study are in favor of the complex origin of the *T*-Reflection representing a mixture of mafic and ultra-mafic cumulates and sills and high-grade inherited lower crust.

The paper in the Appendix B “The Development of Volcanic Sequences at Rifted Margins: New Insights from the Structure and Morphology of the Vøring Escarpment, mid-Norwegian Margin” represents a detailed analysis of complex morphologies of breakup-related sub-aerial and shallow marine volcanic sequences along the Vøring Escarpment. The detailed 2D seismic interpretation shows that the Vøring Escarpment is distinctively segmented depending on (1) the sea level change; (2) volumes of magma supply; (3) pre-existing sub-basalt topography; (4) influence of the adjacent Jan Mayen Fracture Zone; or the mutual impact of all the factors. The results of the study highlight the importance of integrated interpretation of volcanic facies, sub-basalt topography and regional tectonostratigraphy in order to understand the margin development just prior to and during the onset of the continental breakup in the earliest Eocene.

Finally, in the *Appendix C* the two sheets of the updated structural and magmatic elements map of the mid-Norwegian margin, which was presented during the Nordic

Geological Winter Meeting in 2020, are included. This map is based on the mapping results of this PhD thesis and is now together with the explanatory text in preparations for submission to *Norwegian Journal of Geology*.

5. Concluding remarks

An integrated regional interpretation of the new large geological and geophysical dataset presented in this PhD thesis allowed to reinvestigate the pre-breakup development of the entire mid-Norwegian volcanic rifted passive margin (with focus on the Møre and Vøring basins) from crustal to sedimentary/volcanic and prospect levels. A better imaging of deep basins and sub-basalt domains allowed to identify new and revise previously known structural elements. The updated tectonostratigraphic framework for the mid-Norwegian margin is based on the most recent high-quality dataset available offshore Norway. An integrated interpretation of seismic data calibrated with published refraction data and tested by potential field and forward basin modelling augmented understanding of the crustal and basin architecture of the Møre and Vøring basins as a part of the mid-Norwegian passive margin.

This PhD thesis questions the assumptions of the presence of a wide zone of serpentinized mantle right beneath the large parts of the Møre and Vøring basins. The data analysis confirms the primarily crustal nature of the enigmatic high-velocity and high-density lower crustal body and associated *T*-Reflection, which was interpreted as exhumed high-grade Caledonian crust later mixed with breakup-related mafic and ultra-mafic magmatic material. Altogether the results of this study suggest that the outer portion of the mid-Norwegian margin is underlain by continental blocks (>5-8 km of thickness) often observed on top of the LCB (Figure 12b).

This PhD thesis suggests irrelevance of the applicability of the Iberian margin model for the mid-Norwegian margin. The mid-Norwegian margin does not show a single extension phase from the strong necking event in the Late Jurassic-earliest Cretaceous to the late Paleocene-early Eocene lithospheric breakup, but instead experienced a series of episodic and localized Cretaceous to Paleocene extensional events which were separated by intermediate cooling/relaxation. The time lag between the initial necking event and eventual continental breakup in the mid-Norwegian margin is about 100-110 Myr (Figure 12), which is far more than observed in the North Atlantic magma-poor margins (35-45 Myr). If the Iberian model accepted for the mid-Norwegian margin, the breakup conditions would have been achieved around Aptian-Albian time somewhere in the central part of the Møre and Vøring basins. Instead, the rifting was locally aborted at this time (e.g. the Møre Basin), the lithosphere started to cool down and strengthen, and the renewed phases of extension started to migrate laterally and further jumped towards the future COB (Figure 12b), which extends diagonally across the pre-existing rift system. The emplacement of sub-aerial and shallow marine breakup-related volcanics with apparent traces of upper crustal contamination disfavor the Iberian scenario,

where the breakup usually took place in the deep-water domain with no continental crust preserved on top of the mantle.

The conceptual model of the first-order configuration of the mid-Norwegian margin presented in this PhD thesis (Figure 12b) resembles a modified simple-shear model of Lister et al. (1986, 1991), where several large-scale detachments control the formation of rift valleys in the central basins and elevated marginal plateaus in the outer domains. Additionally, isolated ribbons of continental crust (e.g. basement highs) can affect second-order configuration of the margin similar to crustal boudinage mode (e.g. Gartrell, 1997) as it was shown for the Northern Vøring case (Zastrow et al., 2018 – *Paper I*). The Lister et al. (1986, 1991) model agrees with observations and results presented in this PhD thesis and therefore better explains crustal and basin architecture, sedimentary infill, and tectonomagmatic evolution of the mid-Norwegian margin.

6. Future work

As is always the case in science, new studies usually do not only answer the questions, but raise a series of new questions. This thesis is not an exception and results in several important issues:

(1) The mechanisms of deformations in the lower crust and lithospheric mantle remain unclear and require further investigations. An integrated study which combines new 3D regional lithospheric models (e.g. Maystrenko et al., 2017a; 2017b), new geological observations and realistic 2D-3D numerical modelling is a must to understand what are the triggering forces for lower and upper crustal deformations and how did they affect sedimentary and tectonic processes in the basins. Seismic interpretation and regional mapping presented in this PhD thesis provide all observations required by modellers to test various scenarios.

(2) The newly defined Kolga High, Bara High and Bylgja Ridge are located in the outer sub-volcanic basin domain and are associated with prominent gravity and magnetic anomalies and the presence of deep crustal reflections like the *T*-Reflection. A detailed seismic interpretation of large 3D seismic surveys covering these structures accompanied by potential field modelling will help to better assess the crustal architecture and evolution of the outer basin province of the mid-Norwegian margin.

(3) The Fles Fault Complex (FFC) is a major basin boundary in the Vøring segment. It dissects almost 12 km of the Cretaceous-Cenozoic section and is segmented along strike by old transfer zones into N-S and NE-SW parts. A series of compressional events in the Cenozoic led to the local reactivation of the fault system into reverse sense which could govern the growth

of the large Helland-Hansen Arch. An additional study will aim to analyze fault geometries/morphologies and amplitudes, timing of main fault reactivations and controlling factors of localization and nucleation of new faults. The results of this study will be important both for the understanding the origin and formation of large and long-lived intra-basin fault systems and regional tectonostratigraphic framework of the mid-Norwegian margin.

(4) The tectonomagmatic development of the outer Vøring and Møre margins also remains unclear. New well data and available 2D/3D seismic data will help to restore tectonomagmatic and breakup development of this important part of the mid-Norwegian margin. This can be achieved by detailed seismostratigraphic mapping of lower and upper vent complexes, volcanic sequence boundaries, Cenozoic horizons, faults and compressional structures. Such study will result in better understanding of the dynamic of the breakup propagation and the diachronous volcanic development of the mid-Norwegian margin recently suggested (Gernigon et al., 2019). Detailed mapping is necessary to evaluate properly the subtle ages and rates of the Cenozoic magmatic activity in the different marginal segments. Eventually, understanding of how the sequential development of breakup-related magmatism led to PETM (Paleocene-Eocene Thermal Maximum) conditions is also a fundamental task in climate science and current environmental issues.

(5) It is important to integrate results of this PhD thesis into the conjugate margin context to assess the geology and petroleum systems across the entire NE Atlantic province. The existing plate reconstruction models allow to restore confidently the positions of Greenland and Norway back to the breakup time at 56 Ma, however earlier Paleozoic-Mesozoic plate configurations need to be revised and updated. Careful estimations of paleostress orientations and stretching factors for major rifting events in each margin segment will help to quantify amount of extension and therefore to integrate these results into existing plate reconstruction models.

(6) Challenges also remain in both sub-basalt and intra-basalt imaging. In particular seismic attenuation and multiples have proven problematic to confidently interpret sub-basalt and intra basalt geological features. Petrophysical properties of volcanic facies are now being extensively studied based on well data and onshore analogues. This, together with integrated detailed volcanological and sub-volcanic basin interpretations performed in the frame of this PhD thesis, can improve seismic velocity models to make them more realistic and workable, and strengthen sub-basalt imaging.

(7) Fossil passive margins sandwiched in orogenic belts onshore are good objects to study the crustal nature and development of deep parts of passive margins, which are now

uplifted and exposed within orogenic nappes. Compared to well-studied fossil magma-poor margins (e.g. Mohn et al., 2012; Masini et al., 2013, 2014; Beltrando et al., 2014; Clerc et al., 2016), the fossil onshore magma-rich margins are rare and remain poorly studied. It was suggested that rigid and dense volcanic outer domains and/or COT of magma-rich margins are likely to be lost through subduction in convergence settings (Reston & Manatschal, 2011). However, a good example of COT-domain of a pre-Caledonian fossil magma-rich passive margin is located in the Scandinavian Mountains and was recently subjected to detailed petrological, geochemical and structural studies (Abdelmalak et al., 2015; Jakob et al., 2019; Kjøl et al., 2019). A continued study, which combines new incoming data and observations both from fossil onshore and offshore magma-rich margins, is necessary to better understand processes related to the formation of magma-rich margins.

References

- Abdelmalak, M. M., Andersen, T. B., Planke, S., Faleide, J. I., Corfu, F., Tegner, C., ... & Myklebust, R. (2015). The ocean-continent transition in the mid-Norwegian margin: Insight from seismic data and an onshore Caledonian field analogue. *Geology*, 43(11), 1011-1014.
- Abdelmalak, M. M., Meyer, R., Planke, S., Faleide, J. I., Gernigon, L., Frieling, J., ... & Said, A. (2016a). Pre-breakup magmatism on the Vøring Margin: Insight from new sub-basalt imaging and results from Ocean Drilling Program Hole 642E. *Tectonophysics*, 675, 258-274.
- Abdelmalak, M. M., Planke, S., Faleide, J. I., Jerram, D. A., Zastrozhnov, D., Eide, S., & Myklebust, R. (2016b). The development of volcanic sequences at rifted margins: New insights from the structure and morphology of the Vøring Escarpment, mid-Norwegian Margin. *Journal of Geophysical Research: Solid Earth*, 121(7), 5212-5236.
- Abdelmalak, M. M., Faleide, J. I., Planke, S., Gernigon, L., Zastrozhnov, D., Shephard, G. E., & Myklebust, R. (2017). The T-Reflection and the Deep Crustal Structure of the Vøring Margin, Offshore mid-Norway. *Tectonics*, 36(11), 2497-2523.
- Allen, P. A. & Allen, J. R. (2013), *Basin Analysis: Principles and Application to Petroleum Play Assessment*, pp. 642, John Wiley.
- Baird, D. J., Knapp, J. H., Steer, D. N., Brown, L. D., & Nelson, K. D. (1995). Upper-mantle reflectivity beneath the Williston basin, phase-change Moho, and the origin of intracratonic basins. *Geology*, 23(5), 431-434.
- Bakke, V. N. (2017). *Volcanic rocks at the Møre Marginal High: geochemistry, petrogenesis and emplacement mechanisms* (Master's thesis, The University of Bergen).
- Bellwald, B., Urlaub, M., Hjelstuen, B. O., Sejrup, H. P., Sørensen, M. B., Forsberg, C. F., & Vanneste, M. (2019). NE Atlantic continental slope stability from a numerical modeling perspective. *Quaternary Science Reviews*, 203, 248-265.
- Beltrando, M., Manatschal, G., Mohn, G., Dal Piaz, G. V., Brovarone, A. V., & Masini, E. (2014). Recognizing remnants of magma-poor rifted margins in high-pressure orogenic belts: The Alpine case study. *Earth-Science Reviews*, 131, 88-115.
- Bergslie, D. & Syvertsen, S.E. (2015). Challenges in Outer High Exploration – Norwegian Berndt, C., Planke, S., Teagle, D., Huismans, R., Torsvik, T., Frieling, J., ... & Coxall, H. (2019). Northeast Atlantic breakup volcanism and consequences for Paleogene climate change-MagellanPlus Workshop report. Scientific Drilling, 26, 69-85. *Exploration Revived*, 16th March 2015, Bergen, Norway.
- Berndt, C., Planke, S., Teagle, D., Huismans, R., Torsvik, T., Frieling, J., ... & Coxall, H. (2019). Northeast Atlantic breakup volcanism and consequences for Paleogene climate change-MagellanPlus Workshop report. *Scientific Drilling*, 26, 69-85.
- Bjørnseth, H. M., Grant, S. M., Hansen, E. K., Hossack, J. R., Roberts, D. G., & Thompson, M. (1997). Structural evolution of the Vøring Basin, Norway, during the Late Cretaceous and Palaeogene. *Journal of the Geological Society*, 154(3), 559-563.

- Blystad, P., H. Brekke, R. B. Færseth, B. T. Larsen, J. Skogseid, and B. Tørudbakken. (1995). Structural elements of the Norwegian continental shelf Part II: the Norwegian Sea Region. *NPD-Bulletin*, The Norwegian Petroleum Directorate, 8.
- Boillot, G., & Froitzheim, N. (2001). Non-volcanic rifted margins, continental break-up and the onset of sea-floor spreading: some outstanding questions. *Geological Society, London, Special Publications*, 187(1), 9-30.
- Bradley, D. C. (2008). Passive margins through earth history. *Earth-Science Reviews*, 91(1-4), 1-26.
- Breivik, A., Faleide, J. I., Mjelde, R., Flueh, E., & Murai, Y. (2014). Magmatic development of the outer Vøring margin from seismic data. *Journal of geophysical research: solid earth*, 119(9), 6733-6755.
- Brekke, H., & Riis, F. (1987). Tectonics and basin evolution of the Norwegian shelf between 62 N and 72 N. *Norsk geologisk tidsskrift*, 67, 295-322.
- Brekke, H. (2000). The tectonic evolution of the Norwegian Sea Continental Margin with emphasis on the Vøring and Møre Basins. *Geological Society, London, Special Publications*, 167(1), 327-378.
- Brekke, H., Sjulstad, H. I., Magnus, C., & Williams, R. W. (2001). Sedimentary environments offshore Norway—an overview. In *Norwegian Petroleum Society Special Publications* (Vol. 10, pp. 7-37). Elsevier.
- Buck, W. R., & Karner, G. D. (2004). Consequences of asthenospheric variability on continental rifting. *Rheology and deformation of the lithosphere at continental margins*, 62, 1-30.
- Burke, K., Dewey, J. F., & Kidd, W. S. F. (1976). Dominance of horizontal movements, arc and microcontinental collisions during the later permobile regime. *The early history of the Earth*, 113, 129.
- Bøen, F., Eggen, S., & Vollset, J. (1984). Structures and basins of the margin from 62 to 69 N and their development. In *Petroleum Geology of the North European Margin* (pp. 253-270). Springer, Dordrecht.
- Callot, J. P., Grigné, C., Geoffroy, L., & Brun, J. P. (2001). Development of volcanic passive margins: Two-dimensional laboratory models. *Tectonics*, 20(1), 148-159.
- Chenin, P., Manatschal, G., Lavier, L. L., & Erratt, D. (2015). Assessing the impact of orogenic inheritance on the architecture, timing and magmatic budget of the North Atlantic rift system: a mapping approach. *Journal of the Geological Society*, 172(6), 711-720.
- Clerc, C., Lagabrielle, Y., Labaume, P., Ringenbach, J. C., Vauchez, A., Nalpas, T., ... & Fourcade, S. (2016). Basement–Cover decoupling and progressive exhumation of metamorphic sediments at hot rifted margin. Insights from the Northeastern Pyrenean analog. *Tectonophysics*, 686, 82-97.
- Coward, M. P. (1986). Heterogeneous stretching, simple shear and basin development. *Earth and Planetary Science Letters*, 80(3-4), 325-336.

- Dahlgren, K. T., Vorren, T. O., & Laberg, J. S. (2002). Late Quaternary glacial development of the mid-Norwegian margin—65 to 68 N. *Marine and Petroleum Geology*, 19(9), 1089-1113.
- Davis, J. K., & Lavier, L. L. (2017). Influences on the development of volcanic and magma-poor morphologies during passive continental rifting. *Geosphere*, 13(5), 1524-1540.
- de Graciansky, P. C., Poag, C. W., CUNNINGHAM JR, R. O. B. E. R. T., Loubere, P., Masson, D. G., Mazzullo, J. M., ... & Sigal, J. (1985). The Goban Spur transect: Geologic evolution of a sediment-starved passive continental margin. *Geological Society of America Bulletin*, 96(1), 58-76.
- Doré, A. G., Lundin, E. R., Fichler, C., & Olesen, O. (1997). Patterns of basement structure and reactivation along the NE Atlantic margin. *Journal of the Geological Society*, 154(1), 85-92.
- Doré, A. G., Lundin, E. R., Jensen, L. N., Birkeland, Ø., Eliassen, P. E., & Fichler, C. (1999). Principal tectonic events in the evolution of the northwest European Atlantic margin. In *Geological society, London, petroleum geology conference series* (Vol. 5, No. 1, pp. 41-61). Geological Society of London.
- Doré, A. G., Lundin, E. R., Kusznir, N. J., & Pascal, C. (2008). Potential mechanisms for the genesis of Cenozoic domal structures on the NE Atlantic margin: pros, cons and some new ideas. *Geological Society, London, Special Publications*, 306(1), 1-26.
- Ebbing, J., Lundin, E., Olesen, O., & Hansen, E. K. (2006). The mid-Norwegian margin: a discussion of crustal lineaments, mafic intrusions, and remnants of the Caledonian root by 3D density modelling and structural interpretation. *Journal of the Geological Society*, 163(1), 47-59.
- Eldholm, O., Thiede, J. and Taylor E. (1989). Evolution of the Vøring Volcanic Margin. In *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by O. Eldholm, J. Thiede, and E. Taylor, pp. 1033–1065, Ocean Drilling Program, College Station, Tex.
- Eldholm, O. and Grue, K. (1994). North-Atlantic Volcanic Margins - Dimensions and Production-Rates. *Journal of Geophysical Research-Solid Earth*, 99(B2): 2955-2968.
- Faleide, J. I., Tsikalas, F., Breivik, A. J., Mjelde, R., Ritzmann, O., Engen, Ø., Wilson, J., and Eldholm, O. (2008). Structure and evolution of the continental margin off Norway and the Barents Sea. *Episodes*, 31(No. 1), 82-91.
- Fjeldskaar, W., & Amantov, A. (2018). Effects of glaciations on sedimentary basins. *Journal of Geodynamics*, 118, 66-81.
- Fossen, H. (2016). *Structural geology*. Cambridge University Press.
- Foulger, G. R., Natland, J. H., & Anderson, D. L. (2005). A source for Icelandic magmas in remelted Iapetus crust. *Journal of Volcanology and Geothermal Research*, 141(1-2), 23-44.
- Foulger, G. R., & Anderson, D. L. (2005). A cool model for the Iceland hotspot. *Journal of Volcanology and Geothermal Research*, 141(1-2), 1-22.

- Franke, D. (2013). Rifting, lithosphere breakup and volcanism: Comparison of magma-poor and volcanic rifted margins. *Marine and Petroleum Geology*, 43, 63-87.
- Færseth, R. B., & Lien, T. (2002). Cretaceous evolution in the Norwegian Sea—a period characterized by tectonic quiescence. *Marine and Petroleum Geology*, 19(8), 1005-1027.
- Gabrielsen, R. H., Færseth, R., Hamar, G., & Rønnevik, H. (1984). Nomenclature of the main structural features on the Norwegian Continental Shelf north of the 62nd parallel. In *Petroleum Geology of the North European Margin* (pp. 41-60). Springer, Dordrecht.
- Gartrell, A. P. (1997). Evolution of rift basins and low-angle detachments in multilayer analog models. *Geology*, 25(7), 615-618.
- Geoffroy, L. (2005). Volcanic passive margins. *Comptes Rendus Geoscience*, 337(16), 1395-1408.
- Geoffroy, L., Burov, E. B., & Werner, P. (2015). Volcanic passive margins: another way to break up continents. *Scientific Reports*, 5, 14828.
- Gernigon, L., Ringenbach, J. C., Planke, S., Le Gall, B., & Jonquet-Kolstø, H. (2003). Extension, crustal structure and magmatism at the outer Vøring Basin, Norwegian margin. *Journal of the Geological Society*, 160(2), 197-208.
- Gernigon, L., Ringenbach, J.-C., Planke, S., Le Gall, B. (2004). Deep structures and breakup along volcanic rifted margins: insights from integrated studies along the outer Vøring Basin (Norway). *Marine and Petroleum Geology*, 21 (3), 363–372.
- Gernigon, L., Lucazeau, F., Brigaud, F., Ringenbach, J.-C., Planke, S., Le Gall, B. (2006). A moderate melting model for the Vøring margin (Norway) based on structural observations and a thermo-kinematical modelling: Implication for the meaning of the lower crustal bodies. *Tectonophysics*, 412 (3), 255–278.
- Gernigon, L., Blischke, A., Nasuti, A., & Sand, M. (2015). Conjugate volcanic rifted margins, seafloor spreading, and microcontinent: Insights from new high-resolution aeromagnetic surveys in the Norway Basin. *Tectonics*, 34(5), 907-933.
- Gernigon, L., Franke, D., Geoffroy, L., Schiffer, C., Foulger, G. R., & Stoker, M. (2019). Crustal fragmentation, magmatism, and the diachronous opening of the Norwegian-Greenland Sea. *Earth-Science Reviews*.
- Gernigon L., Zastrozhnov D., Planke S., Abdelmalak M.M., Maharjan D., Manton B., Faleide J.I. and Myklebust R. (in prep.). A new structural and magmatic elements map of the mid-Norwegian margin. *Volcanic Basin Petroleum Research/Geological Survey of Norway (NGU)*, Oslo/Trondheim, Norway.
- Gomez, M., Verges, J., Fernandez, M., Torne, M., Ayala, C., Wheeler, W., & Karpuz, R. (2004). Extensional geometry of the Mid Norwegian Margin before Early Tertiary continental breakup. *Marine and Petroleum Geology*, 21(2), 177-194.
- Guan, H., Geoffroy, L., Gernigon, L., Chauvet, F., Grigné, C., & Werner, P. (2019). Magmatic ocean-continent transitions. *Marine and Petroleum Geology*, 104, 438-450.

- Haflidason, H., Sejrup, H. P., Nygård, A., Mienert, J., Bryn, P., Lien, R., ... & Masson, D. (2004). The Storegga Slide: architecture, geometry and slide development. *Marine geology*, 213(1-4), 201-234.
- Hartz, E. H., Medvedev, S., & Schmid, D. W. (2017). Development of sedimentary basins: differential stretching, phase transitions, shear heating and tectonic pressure. *Basin Research*, 29(5), 591-604.
- Hinz, K. (1981). Wedges of very thick oceanward dipping layers beneath passive continental margins. The origin and paleoenvironmental significance. *Geologisches Jahrbuch E*, 22, 3-28.
- Hinz, K., Mutter, J. C., Zehnder, C. M., & NGT Study Group. (1987). Symmetric conjugation of continent-ocean boundary structures along the Norwegian and East Greenland margins. *Marine and Petroleum Geology*, 4(3), 166-187.
- Hjelstuen, B. O., Sejrup, H. P., Haflidason, H., Berg, K., & Bryn, P. (2004). Neogene and Quaternary depositional environments on the Norwegian continental margin, 62 N–68 N. *Marine geology*, 213(1-4), 257-276.
- Huisman, R. S., & Beaumont, C. (2002). Asymmetric lithospheric extension: The role of frictional plastic strain softening inferred from numerical experiments. *Geology*, 30(3), 211-214.
- Huisman, R. S., & Beaumont, C. (2008). Complex rifted continental margins explained by dynamical models of depth-dependent lithospheric extension. *Geology*, 36(2), 163-166.
- Jakob, J., Andersen, T. B., & Kjøll, H. J. (2019). A review and reinterpretation of the architecture of the South and South-Central Scandinavian Caledonides—A magma-poor to magma-rich transition and the significance of the reactivation of rift inherited structures. *Earth-science reviews*.
- Kaus, B. J., Connolly, J. A., Podladchikov, Y. Y., & Schmalholz, S. M. (2005). Effect of mineral phase transitions on sedimentary basin subsidence and uplift. *Earth and Planetary Science Letters*, 233(1-2), 213-228.
- Kjøll, H. J., Andersen, T. B., Corfu, F., Labrousse, L., Tegner, C., Abdelmalak, M. M., & Planke, S. (2019). Timing of break-up and thermal evolution of a pre-Caledonian Neoproterozoic exhumed magma-rich rifted margin. *Tectonics*.
- Kusznir, N. J., Marsden, G., & Egan, S. S. (1991). A flexural-cantilever simple-shear/pure-shear model of continental lithosphere extension: applications to the Jeanne d'Arc Basin, Grand Banks and Viking Graben, North Sea. *Geological Society, London, Special Publications*, 56(1), 41-60.
- Laberg, J. S., & Vorren, T. O. (2000). The Trænadjupet Slide, offshore Norway—morphology, evacuation and triggering mechanisms. *Marine Geology*, 171(1-4), 95-114.
- Levell, B., Argent, J., Doré, A. G., & Fraser, S. (2010). Passive margins: overview. In *Geological Society, London, Petroleum Geology Conference series* (Vol. 7, No. 1, pp. 823-830). Geological Society of London.
- Lister, G. S., Etheridge, M. A., & Symonds, P. A. (1986). Detachment faulting and the evolution of passive continental margins. *Geology*, 14(3), 246-250.

- Lister, G. S., Etheridge, M. A., & Symonds, P. A. (1991). Detachment models for the formation of passive continental margins. *Tectonics*, 10(5), 1038-1064.
- Lundin, E. R., & Doré, A. G. (1997). A tectonic model for the Norwegian passive margin with implications for the NE Atlantic: Early Cretaceous to break-up. *Journal of the Geological Society*, 154(3), 545-550.
- Lundin, E., & Doré, A. G. (2002). Mid-Cenozoic post-breakup deformation in the 'passive' margins bordering the Norwegian–Greenland Sea. *Marine and Petroleum Geology*, 19(1), 79-93.
- Lundin, E. R., & Doré, A. G. (2011). Hyperextension, serpentinization, and weakening: A new paradigm for rifted margin compressional deformation. *Geology*, 39(4), 347-350.
- Lundin, E. R., Doré, A. G., Rønning, K., & Kyrkjebø, R. (2013). Repeated inversion and collapse in the Late Cretaceous–Cenozoic northern Vøring Basin, offshore Norway. *Petroleum Geoscience*, 19(4), 329-341.
- Manatschal, G., & Bernoulli, D. (1999). Architecture and tectonic evolution of nonvolcanic margins: Present-day Galicia and ancient Adria. *Tectonics*, 18(6), 1099-1119.
- Mangerud, J., Gyllencreutz, R., Lohne, Ø., & Svendsen, J. I. (2011). Glacial history of Norway. In *Developments in Quaternary Sciences* (Vol. 15, pp. 279-298). Elsevier.
- Mann, P., Gahagan, L. & Gordon, M. B. (2003). Tectonic Setting of the World's Giant Oil and Gas Fields. American Association of Petroleum Geologists, Tulsa, OK, Memoirs, 78, 15–105
- Manton, B., Planke, S., Zastrozhnov, D., Abdelmalak, M. M., Millett, J., Maharjan, D., ... & Kjølhamar, B. E. (2018). Pre-Cretaceous prospectivity of the outer Møre and Vøring basins constrained by new 3D seismic data. In *80th EAGE Conference and Exhibition 2018*.
- Marsden, G., Yielding, G., Roberts, A. M., & Kusznir, N. J. (1990). Application of a flexural cantilever simple-shear/pure shear model of continental lithosphere extension to the formation of the northern North Sea basin. *Tectonic evolution of the North Sea rifts*, 241-261.
- Martinius, A. W., Kaas, I., Helgesen, G., Kj, J. M., & Leith, D. A. (2001). Sedimentology of the heterolithic and tide-dominated Tilje Formation (Early Jurassic, Halten Terrace, offshore mid-Norway). In *Norwegian Petroleum Society Special Publications* (Vol. 10, pp. 103-144). Elsevier.
- Masini, E., Manatschal, G., & Mohn, G. (2013). The Alpine Tethys rifted margins: Reconciling old and new ideas to understand the stratigraphic architecture of magma-poor rifted margins. *Sedimentology*, 60(1), 174-196.
- Masini, E., Manatschal, G., Tugend, J., Mohn, G., & Flament, J. M. (2014). The tectono-sedimentary evolution of a hyper-extended rift basin: the example of the Arzacq–Mauléon rift system (Western Pyrenees, SW France). *International Journal of Earth Sciences*, 103(6), 1569-1596.
- Maystrenko, Y. P., Olesen, O., Gernigon, L., & Gradmann, S. (2017a). Deep structure of the Lofoten-Vesterålen segment of the Mid-Norwegian continental margin and adjacent areas

derived from 3-D density modeling. *Journal of Geophysical Research: Solid Earth*, 122(2), 1402-1433.

Maystrenko, Y. P., Gernigon, L., Nasuti, A., & Olesen, O. (2017b). Deep structure of the Mid-Norwegian continental margin (the Vøring and Møre basins) according to 3-D density and magnetic modelling. *Geophysical Journal International*, 212(3), 1696-1721.

McKenzie, D. (1978). Some remarks on the development of sedimentary basins. *Earth and Planetary science letters*, 40(1), 25-32.

Medvedev, S., Hartz, E. H., Schmid, D. W., Zakariassen, E., & Varhaug, P. (2019). Influence of glaciations on North Sea petroleum systems. *Geological Society, London, Special Publications*, 494, SP494-2018.

Menzies, M. A., Klemperer, S. L., Ebinger, C. J., & Baker, J. (2002). Characteristics of volcanic rifted margins. *Special Papers-Geological Society of America*, 1-14.

Meyer, R., Van Wijk, J., & Gernigon, L. (2007). The North Atlantic Igneous Province: A review of models for its formation. *Geological Society of America Special Papers*, 430, 525-552.

Meyer, R., Hertogen, J., Pedersen, R. B., Viereck-Götte, L., & Abratis, M. (2009). Interaction of mantle derived melts with crust during the emplacement of the Vøring Plateau, NE Atlantic. *Marine Geology*, 261(1-4), 3-16.

Millett, J. M., Manton, B. M., Zastrozhnov, D., Planke, S., Maharjan, D., Bellwald, B., ... & Abdelmalak, M. M. (2020). Basin Structure and Prospectivity of the NE Atlantic Volcanic Rifted Margin: Cross-border examples from the Faroe-Shetland, Møre and Southern Vøring Basins. *Geological Society, London, Special Publications*, 495.

Mjelde, R., Kasahara, J., Shimamura, H., Kamimura, A., Kanazawa, T., Kodaira, S., ... & Shiobara, H. (2002). Lower crustal seismic velocity-anomalies; magmatic underplating or serpentinized peridotite? Evidence from the Vøring Margin, NE Atlantic. *Marine Geophysical Researches*, 23(2), 169-183.

Mjelde, R., Shimamura, H., Kanazawa, T., Kodaira, S., Raum, T., & Shiobara, H. (2003). Crustal lineaments, distribution of lower crustal intrusives and structural evolution of the Vøring Margin, NE Atlantic; new insight from wide-angle seismic models. *Tectonophysics*, 369(3-4), 199-218.

Mjelde, R., Raum, T., Breivik, A., Shimamura, H., Murai, Y., Takanami, T., & Faleide, J. I. (2005). Crustal structure of the Vøring Margin, NE Atlantic: a review of geological implications based on recent OBS data. In *Geological Society, London, Petroleum Geology Conference Series* (Vol. 6, No. 1, pp. 803-813). Geological Society of London.

Mjelde, R., Raum, T., Murai, Y., & Takanami, T. (2007). Continent-ocean-transitions: review, and a new tectono-magmatic model of the Vøring Plateau, NE Atlantic. *Journal of Geodynamics*, 43(3), 374-392.

Mjelde, R., Breivik, A. J., Raum, T., Mittelstaedt, E., Ito, G., & Faleide, J. I. (2008). Magmatic and tectonic evolution of the North Atlantic. *Journal of the Geological Society*, 165(1), 31-42.

- Mjelde, R., Faleide, J. I., Breivik, A. J., & Raum, T. (2009). Lower crustal composition and crustal lineaments on the Vøring Margin, NE Atlantic: a review. *Tectonophysics*, 472(1-4), 183-193.
- Mjelde, R., Kvarven, T., Faleide, J. I., & Thybo, H. (2016). Lower crustal high-velocity bodies along North Atlantic passive margins, and their link to Caledonian suture zone eclogites and Early Cenozoic magmatism. *Tectonophysics*, 670, 16-29.
- Mohn, G., Manatschal, G., Beltrando, M., Masini, E., & Kuszniir, N. (2012). Necking of continental crust in magma-poor rifted margins: Evidence from the fossil Alpine Tethys margins. *Tectonics*, 31(1).
- Mohn, G., Karner, G. D., Manatschal, G., & Johnson, C. A. (2015). Structural and stratigraphic evolution of the Iberia–Newfoundland hyper-extended rifted margin: a quantitative modelling approach. *Geological Society, London, Special Publications*, 413(1), 53-89.
- Mohriak, W., Nemčok, M., & Enciso, G. (2008). South Atlantic divergent margin evolution: rift-border uplift and salt tectonics in the basins of SE Brazil. *Geological Society, London, Special Publications*, 294(1), 365-398.
- Mohriak, W. U., & Leroy, S. (2013). Architecture of rifted continental margins and break-up evolution: insights from the South Atlantic, North Atlantic and Red Sea–Gulf of Aden conjugate margins. *Geological Society, London, Special Publications*, 369(1), 497-535.
- Mosar, J. (2000). Depth of extensional faulting on the Mid-Norway Atlantic passive margin. In *Norges geologiske undersøkelse Bulletin* (Vol. 437, p. 33).
- Mosar, J. (2003). Scandinavia's North Atlantic passive margin. *Journal of Geophysical Research: Solid Earth*, 108(B8).
- Moullade, M., Brunet, M.F. & Boillot, G. (1988). Subsidence and deepening of the Galicia Margin: the paleoenvironmental control. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 103, pp. 733-740). Ocean Drilling Program College Station, TX.
- Mutter, J. C., Talwani, M., & Stoffa, P. L. (1982). Origin of seaward-dipping reflectors in oceanic crust off the Norwegian margin by “subaerial sea-floor spreading”. *Geology*, 10(7), 353-357.
- Mutter, J. C., & Zehnder, C. M. (1988). Deep crustal structure and magmatic processes: the inception of seafloor spreading in the Norwegian-Greenland Sea. *Geological Society, London, Special Publications*, 39(1), 35-48.
- Müller, R., Ngstuen, J. P., Eide, F., & Lie, H. (2005). Late Permian to Triassic basin infill history and palaeogeography of the Mid-Norwegian shelf—East Greenland region. In *Norwegian Petroleum Society Special Publications* (Vol. 12, pp. 165-189). Elsevier.
- Nadin, P. A., & Kuszniir, N. J. (1995). Palaeocene uplift and Eocene subsidence in the northern North Sea Basin from 2D forward and reverse stratigraphic modelling. *Journal of the Geological Society*, 152(5), 833-848.

- Nirrengarten, M., Gernigon, L., & Manatschal, G. (2014). Lower crustal bodies in the Møre volcanic rifted margin: Geophysical determination and geological implications. *Tectonophysics*, 636, 143-157.
- Osmundsen, P. T., Sommaruga, A., Skilbrei, J. R., & Olesen, O. (2002). Deep structure of the Mid Norway rifted margin. *Norwegian Journal of Geology/Norsk Geologisk Forening*, 82(4).
- Osmundsen, P. T., & Péron-Pinvidic, G. (2018). Crustal-scale fault interaction at rifted margins and the formation of domain-bounding breakaway complexes: Insights from offshore Norway. *Tectonics*, 37(3), 935-964.
- Perez-Gussinye, M., Ranero, C.R., Reston, T.J. and Sawyer, D. (2003). Mechanisms of extension at nonvolcanic margins: Evidence from the Galicia interior basin, west of Iberia. *Journal of Geophysical Research: Solid Earth*, 108(B5).
- Peron-Pinvidic, G., Manatschal, G., & Osmundsen, P. T. (2013). Structural comparison of archetypal Atlantic rifted margins: A review of observations and concepts. *Marine and Petroleum Geology*, 43, 21-47.
- Peron-Pinvidic, G., & Osmundsen, P. T. (2016). Architecture of the distal and outer domains of the Mid-Norwegian rifted margin: Insights from the Rån-Gjallar ridges system. *Marine and Petroleum Geology*, 77, 280-299.
- Peron-Pinvidic, G., Osmundsen, P. T., & Ebbing, J. (2016). Mismatch of geophysical datasets in distal rifted margin studies. *Terra Nova*, 28(5), 340-347.
- Peron-Pinvidic, G., & Osmundsen, P. T. (2018). The Mid Norwegian-NE Greenland conjugate margins: Rifting evolution, margin segmentation, and breakup. *Marine and Petroleum Geology*, 98, 162-184.
- Peron-Pinvidic, G., & Manatschal, G. (2019). Rifted margins: state of the art and future challenges. *Frontiers in Earth Science*, 7, 218.
- Petersen, K. D., & Schiffer, C. (2016). Wilson cycle passive margins: Control of orogenic inheritance on continental breakup. *Gondwana Research*, 39, 131-144.
- Planke, S., Skogseid, J., & Eldholm, O. (1991). Crustal structure off Norway, 62 to 70 north. *Tectonophysics*, 189(1-4), 91-107.
- Planke, S., Rasmussen, T., Rey, S. S., & Myklebust, R. (2005). Seismic characteristics and distribution of volcanic intrusions and hydrothermal vent complexes in the Vøring and Møre basins. In *Geological Society, London, Petroleum Geology Conference series* (Vol. 6, No. 1, pp. 833-844). Geological Society of London.
- Podladchikov, Y. Y., Poliakov, A. N. B., & Yuen, D. A. (1994). The effect of lithospheric phase transitions on subsidence of extending continental lithosphere. *Earth and Planetary Science Letters*, 124(1-4), 95-103.
- Ravnås, R., Berge, K., Campbell, H., Harvey, C., & Norton, M. J. (2014). Halten Terrace Lower and Middle Jurassic inter-rift megasequence analysis: megasequence structure, sedimentary architecture and controlling parameters. *From Depositional Systems to Sedimentary Successions on the Norwegian Continental Margin*, 46(46), 215-252.

- Reemst, P., & Cloetingh, S. (2000). Polyphase rift evolution of the Vøring margin (mid-Norway): Constraints from forward tectonostratigraphic modeling. *Tectonics*, 19(2), 225-240.
- Ren, S., Skogseid, J., & Eldholm, O. (1998). Late Cretaceous-Paleocene extension on the Vøring volcanic margin. *Marine Geophysical Researches*, 20(4), 343-369.
- Ren, S., Faleide, J. I., Eldholm, O., Skogseid, J., & Gradstein, F. (2003). Late Cretaceous–Paleocene tectonic development of the NW Vøring basin. *Marine and Petroleum Geology*, 20(2), 177-206.
- Reston, T. J. (2007). The formation of non-volcanic rifted margins by the progressive extension of the lithosphere: the example of the West Iberian margin. *Geological Society, London, Special Publications*, 282(1), 77-110.
- Reston, T. J. (2009). The structure, evolution and symmetry of the magma-poor rifted margins of the North and Central Atlantic: A synthesis. *Tectonophysics*, 468(1-4), 6-27.
- Reston, T., & Manatschal, G. (2011). Rifted margins: Building blocks of later collision. In *Arc-continent collision* (pp. 3-21). Springer, Berlin, Heidelberg.
- Reynisson, R. F., Ebbing, J., Lundin, E., & Osmundsen, P. T. (2010). Properties and distribution of lower crustal bodies on the mid-Norwegian margin. In *Geological Society, London, Petroleum Geology Conference Series* (Vol. 7, No. 1, pp. 843-854). Geological Society of London.
- Rowley, D. B., & Sahagian, D. (1986). Depth-dependent stretching: A different approach. *Geology*, 14(1), 32-35.
- Rüpke, L. H., Schmid, D. W., Perez-Gussinye, M., & Hartz, E. (2013). Interrelation between rifting, faulting, sedimentation, and mantle serpentinization during continental margin formation—including examples from the Norwegian Sea. *Geochemistry, Geophysics, Geosystems*, 14(10), 4351-4369.
- Saltus, R. W., & Blakely, R. J. (2011). Unique geologic insights from “non-unique” gravity and magnetic interpretation. *GSA Today*, 21(12), 4-11.
- Scheck-Wenderoth, M., Raum, T., Faleide, J. I., Mjelde, R., & Horsfield, B. (2007). The transition from the continent to the ocean: a deeper view on the Norwegian margin. *Journal of the Geological Society*, 164(4), 855-868.
- Schiffer, C., Doré, A. G., Foulger, G. R., Franke, D., Geoffroy, L., Gernigon, L., ... & Peace, A. (2019). Structural inheritance in the North Atlantic. *Earth-Science Reviews*, 102975.
- Skogseid, J., Eldholm, O., (1989. Vøring plateau continental margin: Seismic interpretation, stratigraphy and vertical movements. In *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by O. Eldholm, J. Thiede, and E. Taylor, pp. 993–1030, Ocean Drilling Program, College Station, Tex.
- Skogseid, J., Pedersen, T., Eldholm, O., & Larsen, B. T. (1992). Tectonism and magmatism during NE Atlantic continental break-up: the Vøring Margin. *Geological Society, London, Special Publications*, 68(1), 305-320.

- Skogseid, J., Planke, S., Faleide, J. I., Pedersen, T., Eldholm, O., & Neverdal, F. (2000). NE Atlantic continental rifting and volcanic margin formation. *Geological Society, London, Special Publications*, 167(1), 295-326.
- Skogseid, J. (2001). Volcanic margins: geodynamic and exploration aspects. *Marine and Petroleum Geology*, 18(4), 457-461.
- Stemmerik, L., Christiansen, F., Piasecki, S., Jordt, B., Marcussen, C., & Nøhr-Hansen, H. (1993). Depositional history and petroleum geology of the Carboniferous to Cretaceous sediments in the northern part of East Greenland. In *Norwegian Petroleum Society Special Publications* (Vol. 2, pp. 67-87). Elsevier.
- Stemmerik, L. (2000). Late Palaeozoic evolution of the North Atlantic margin of Pangea. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 161(1-2), 95-126.
- Stoker, M. S., Stewart, M. A., Shannon, P. M., Bjerager, M., Nielsen, T., Blischke, A., ... & Ólavsdóttir, J. (2017). An overview of the Upper Palaeozoic–Mesozoic stratigraphy of the NE Atlantic region. *Geological Society, London, Special Publications*, 447(1), 11-68.
- Straume, E. O., Gaina, C., Medvedev, S., Hochmuth, K., Gohl, K., Whittaker, J. M., ... & Hopper, J. R. (2019). GlobSed: Updated total sediment thickness in the world's oceans. *Geochemistry, Geophysics, Geosystems*, 20(4), 1756-1772.
- Swiecicki, T., Gibbs, P. B., Farrow, G. E., & Coward, M. P. (1998). A tectonostratigraphic framework for the Mid-Norway region. *Marine and Petroleum Geology*, 15(3), 245-276.
- Theissen-Krah, S., Zastrozhnov, D., Abdelmalak, M. M., Schmid, D. W., Faleide, J. I., & Gernigon, L. (2017). Tectonic evolution and extension at the Møre Margin–Offshore mid-Norway. *Tectonophysics*, 721, 227-238.
- Tsikalas, F., Faleide, J. I., Eldholm, O., & Blaich, O. A. (2012). The NE Atlantic conjugate margins. *Regional geology and tectonics: Phanerozoic passive margins, cratonic basins and global tectonic maps*, 1, 140-201.
- Van Wijk, J. W., & Cloetingh, S. A. P. L. (2002). Basin migration caused by slow lithospheric extension. *Earth and Planetary Science Letters*, 198(3-4), 275-288.
- Walker, I. M., Berry, K. A., Bruce, J. R., Bystøl, L., & Snow, J. H. (1997). Structural modelling of regional depth profiles in the Vøring Basin: implications for the structural and stratigraphic development of the Norwegian passive margin. *Journal of the Geological Society*, 154(3), 537-544.
- Wernicke, B. (1981). Low-angle normal faults in the Basin and Range Province: nappe tectonics in an extending orogen. *Nature*, 291(5817), 645.
- Wernicke, B. (1985). Uniform-sense normal simple shear of the continental lithosphere. *Canadian Journal of Earth Sciences*, 22(1), 108-125.
- White, R., & McKenzie, D. (1989). Magmatism at rift zones: the generation of volcanic continental margins and flood basalts. *Journal of Geophysical Research: Solid Earth*, 94(B6), 7685-7729.

- White, R. S. (1992). Crustal structure and magmatism of North Atlantic continental margins. *Journal of the Geological Society*, 149(5), 841-854.
- White, R. S., Smith, L. K., Roberts, A. W., Christie, P. A. F., & Kuszniir, N. J. (2008). Lower-crustal intrusion on the North Atlantic continental margin. *Nature*, 452(7186), 460.
- Wilson, J. T. (1966). Did the Atlantic close and then re-open? *Nature*, v.211, p.676-681.
- Wood, R., & Barton, P. (1983). Crustal thinning and subsidence in the North Sea. *Nature*, 302(5904), 134.
- Yamasaki, T., & Gernigon, L. (2009). Styles of lithospheric extension controlled by underplated mafic bodies. *Tectonophysics*, 468(1-4), 169-184.
- Zastrozhnov, D., Gernigon, L., Gogin, I., Abdelmalak, M. M., Planke, S., Faleide, J. I., ... & Myklebust, R. (2018). Cretaceous-Paleocene Evolution and Crustal Structure of the Northern Vøring Margin (Offshore Mid-Norway): Results from Integrated Geological and Geophysical Study. *Tectonics*, 37(2), 497-528.
- Zastrozhnov, D., Gernigon, L., Gogin, I., Planke S., Abdelmalak, M.M., Polteau, S., ... & Myklebust, R. (2020). Regional structure and polyphased Cretaceous-Paleocene rift and basin development of the mid-Norwegian volcanic passive margin. *Marine and Petroleum Geology*.
- Zehnder, C. M., Mutter, J. C., & Buhl, P. (1990). Deep seismic and geochemical constraints on the nature of rift-induced magmatism during breakup of the North Atlantic. *Tectonophysics*, 173(1-4), 545-565.

PART II

Paper I

Cretaceous-Paleocene Evolution and Crustal Structure of the Northern Vøring Margin (Offshore Mid-Norway): Results from Integrated Geological and Geophysical Study

By

Dmitry Zastrozhnov, Laurent Gernigon, Iakov Gogin, Mansour M. Abdelmalak, Sverre Planke, Jan Inge Faleide, Sigurd Eide and Reidun Myklebust

Tectonics, 37(2), 497-528

RESEARCH ARTICLE

10.1002/2017TC004655

Key Points:

- We present a new tectonic scenario for the Cretaceous-Paleocene development of the volcanic rifted mid-Norwegian Margin and the northern Vøring segment in particular
- Crustal architecture and tectonic development of the volcanic rifted mid-Norwegian Margin significantly differs from the Iberian type margins
- The development of sag basins is strongly related to inherited crustal blocks ("buffers")

Correspondence to:

D. Zastrozhnov,
zastrozhe@gmail.com

Citation:

Zastrozhnov, D., Gernigon, L., Gogin, I., Abdelmalak, M. M., Planke, S., Faleide, J. I., ... Myklebust, R. (2018). Cretaceous-Paleocene evolution and crustal structure of the northern Vøring Margin (offshore mid-Norway): Results from integrated geological and geophysical study. *Tectonics*, 37. <https://doi.org/10.1002/2017TC004655>

Received 11 MAY 2017

Accepted 11 JAN 2018

Accepted article online 15 JAN 2018

Cretaceous-Paleocene Evolution and Crustal Structure of the Northern Vøring Margin (Offshore Mid-Norway): Results from Integrated Geological and Geophysical Study

D. Zastrozhnov^{1,2} , L. Gernigon³, I. Gogin⁴, M. M. Abdelmalak^{1,2} , S. Planke^{1,2}, J. I. Faleide¹ , S. Eide⁵ , and R. Myklebust⁶
¹Centre for Earth Evolution and Dynamics, University of Oslo, Oslo, Norway, ²Volcanic Basin Petroleum Research (VBPR), Oslo Science Park, Oslo, Norway, ³Geological Survey of Norway (NGU), Trondheim, Norway, ⁴RPS Ichron, Northwich, UK, ⁵First Geo AS, Skøyen, Norway, ⁶TGS, Asker, Norway

Abstract We present results of a multidisciplinary study of the northern segment of the Vøring volcanic rifted margin, offshore mid-Norway. This segment represents a transitional margin domain that is less investigated compared to the adjacent segments of the margin. In order to understand the geological evolution of the study area, we performed an integrated interpretation of an extensive geological and geophysical data set. This data set includes recently acquired and reprocessed 2-D reflection seismic, published refraction data and potential field data, as well as new borehole data. Two-dimensional potential field modeling was performed to better assess the crustal architecture and evolution of the northern Vøring Margin. We then consider how crustal-scale structures and processes affected the basin formation. The outer and distal northern Vøring Margin represents a series of deep Cretaceous (Træna Basin and Någrind Syncline) and Cretaceous-Paleocene (Hel Graben) sag subbasins underlain by a significantly thinned continental crust. These subbasins developed in between structural highs (Utgard, Nyk, and Grimm Highs), which are underlain by a thicker crust and interpreted as a series of rigid continental blocks ("buffers"). In addition to the regional Late Jurassic-Early Cretaceous rifting events, we found structural evidence of local Neocomian and mid-Cretaceous extensional reactivation affecting the northern segment of the Vøring Basin. During the mid-Late Cretaceous-Paleocene, the extensional axis within the Vøring Basin province migrated sequentially northwestward to the present-day continent-ocean "boundary". We also show fundamental differences between the volcanic rifted mid-Norwegian Margin and nonvolcanic (Iberian-type) margins and how preexisting structures events can shape the evolution and architecture of the margin.

1. Introduction

The processes through which continental rifting develops into oceanic rifting is a major challenge in the Earth Sciences. The relationship between continental rifting and magmatism and the transition from rifting to spreading (e.g., the breakup stage) are still poorly understood and controversial. Important questions usually concern the following: (1) how much continental extension and thinning of the lithosphere predates the initiation of oceanic rifting (e.g., Brune et al., 2016; Huisman & Beaumont, 2011; Lavie & Manatschal, 2006; Van Avendonk et al., 2009), (2) how the rifting and breakup extension are physically and rheologically accommodated by faulting/shearing and/or igneous additions (e.g., Brune et al., 2012; Buck, 2006; Ebinger & Casey, 2001; Rosenbaum et al., 2010; Yamasaki & Gernigon, 2009), (3) the role of inheritance in rift and drift development (e.g., Harry & Bowling, 1999; Petersen & Schiffer, 2016), and (4) the fundamental magmatic and tectonic differences between "magma-poor" and volcanic rifted margins (Franke, 2013; Geoffroy et al., 2015).

In this context, we explore the crustal structure and tectonostratigraphic development of the Vøring Margin, which is a part of the mid-Norwegian Margin (Figure 1). The Vøring Margin is long recognized to be a classic example of a volcanic rifted margin. It is characterized by a long period of rifting, distal basaltic seaward dipping reflectors (SDRs), extensive distribution of mafic dikes and sills in the continental crust and sedimentary basins, and the presence of high-velocity ($V_p > 7.0$ km/s) lower crustal bodies (Abdelmalak et al., 2015, 2017; Eldholm et al., 1989; Mjelde et al., 2016; Planke et al., 1991; Skogseid et al., 2000).

Figure 1. (a) Updated nomenclature map of the northern mid-Norwegian Margin with prebreakup structural elements. The fault pattern is mainly based on this study and modified from Blystad et al. (1995) and Gernigon et al. (2003). Locations of seismic close-ups are shown in purple solid lines. Extent of isochore maps is defined by black rectangle. (b) Geological and geophysical data set used in this study: Red circles = expended spread profiles (centers), solid black lines = ocean bottom seismometers profiles, solid light gray lines = reflection seismic, and yellow circles = main wells used in this study.

The northern part of the Vøring Margin represents a specific transitional structural domain bounded by two prominent transfer zones—the Surt and Bivrost Lineaments (Figure 1a). However, this particular segment of the Vøring Margin was poorly documented and less studied in terms of prebreakup tectonic evolution and crustal architecture. The main period of data acquisition and interpretation that resulted in the development of the tectonostratigraphic framework of this segment occurred two decades ago. This geophysical and geological material has been reviewed and summarized in the context of regional tectonics (Blystad et al., 1995; Brekke, 2000; Lundin et al., 2013; Mjelde et al., 1998; Mosar, 2000; Mosar et al., 2002; Osmundsen et al., 2002; Planke et al., 1991; Skogseid et al., 1992; Tsikalas et al., 2008). Only a few semi-regional case studies focus on and document the Late Cretaceous-Paleocene rifting event preceding the breakup (Fjellanger et al., 2005; Gernigon et al., 2003; Imber et al., 2005; Mogensen et al., 2000; Ren et al., 2003, 1998). Other studies have used published data sets and materials to underpin potential field modeling exercises (Ebbing et al., 2006; Kusznir et al., 2005; Maystrenko et al., 2017; Scheck-Wenderoth et al., 2007; Theissen-Krah & Rupke, 2010; Walker et al., 1997; Wangen et al., 2011).

Various controversial scenarios have been proposed to explain the dynamic and depositional evolution of the Vøring Margin and its associated basins. The most recent studies have suggested some similarities with the Iberian (magma-poor) type of passive margins (Lundin & Doré, 2011; Péron-Pinvidic & Osmundsen, 2016; Reynisson et al., 2010; Rüpké et al., 2013) implying the presence of relatively wide, serpentinized/exhumed mantle domains in the central, distal, and volcanic parts of the margin. In contrast, other publications consider fundamental differences with magma-poor margin (Iberian type) highlighting the role of crustal inheritance and rift duration on passive margin development and showing that there are no clear prerequisites and clear evidence to support a broad zone of mantle exhumation/serpentinization within the mid-Norwegian Margin (Abdelmalak et al., 2017; Gernigon et al., 2003, 2004, 2015; Maystrenko et al., 2017; Nirrengarten et al., 2014; Petersen & Schiffer, 2016; Theissen-Krah et al., 2017). One of the debate concerns the origin and geological interpretation of the strong crustal reflections underlying mostly Cretaceous highs and basins in the distal and outer domains of the margin. This is a key aspect to consider in defining a model of the development of

the mid-Norwegian Margin and the Vøring segment in particular. A better combination of multidisciplinary geological data sets (refraction and reflection seismic, gravity and magnetic, biostratigraphy/lithostratigraphy, structural geology, tectonic and potential field modeling, etc.) is therefore necessary to validate both sedimentary and crustal interpretations (see Saltus & Blakely, 2011, for a discussion).

In this contribution, we use new and reprocessed seismic data sets, combined with recently released and revised well biostratigraphy, in order to refine the basin architecture and the Cretaceous-Paleocene tectonic evolution of the northern Vøring Margin. Over the last decade, a new generation of long-offset, high-quality seismic reflection profiles covering the entire mid-Norwegian Margin has permitted an improvement in the imaging of the deep basin structures and allowed better regional seismic interpretation. Our seismic reflection interpretation is compared with available ocean bottom seismometers (OBS) data and expanded spread profiles (ESPs) and combined with a potential field modeling along a representative NW-SE oriented crustal transect across the northern Vøring Margin. This multidisciplinary approach allows us to evaluate the properties of the deep structures beneath the sedimentary cover and test hypotheses about the style of crustal deformation and sedimentation in a poly rifted volcanic margin setting. We finally propose and discuss a crustal and tectonic scenario for the rifted margin evolution.

2. Regional Setting

The present structure of the Vøring Margin off mid-Norway is a result of several post-Caledonian extensional episodes culminating with the complete separation between Norway and Greenland in the earliest Eocene (Brekke, 2000; Faleide et al., 2008; Talwani & Eldholm, 1977). The Vøring Margin can be divided into a series of second-order structures (basins, highs, grabens, etc.) (Figure 1a) formed during Late Jurassic-Early Cretaceous and Late Cretaceous-Paleocene rifting (Blystad et al., 1995). Additional extensional events took place in mid-Cretaceous time (Bjørnseth et al., 1997; Gernigon et al., 2003; Lundin & Doré, 1997; Pascoe et al., 1999; Roberts et al., 2009) but is still a matter of debate because some authors believe that the mid-Cretaceous period was dominantly a quiet tectonic period (Færseth, 2012; Færseth & Lien, 2002). The Late Cretaceous-Paleocene rifting phase, which culminated in continental breakup near the Paleocene-Eocene transition (56 Ma), was associated with extensive and voluminous magmatic activity (Eldholm et al., 2002). This large magmatic event is traditionally correlated with the arrival of the Icelandic mantle “plume” (Skogseid et al., 2000) and/or independent sublithospheric processes (Foulger, 2010; Meyer et al., 2007; Simon et al., 2009; van Wijk et al., 2001).

The postbreakup evolution of the Vøring Margin is characterized mainly by thermal cooling and regional subsidence of the sedimentary basins (Brekke, 2000; Faleide et al., 2008). A mid-Cenozoic compressional event is also expressed by a number of inversion structures (domes/arches, reverse faults, etc.), which are likely the result of a compressive stress dominantly induced by ridge push (Doré et al., 2008; Vågnes et al., 1998). The final development of the margin is closely linked with Northern Hemisphere glaciation events, when large Plio-Pleistocene depocenters expanded over the area, increasing the regional subsidence and tilt of the margin (Faleide et al., 2008; Hjelstuen et al., 1999; Rise et al., 2005).

The northern Vøring Margin segment, which forms the focus of this study, is located west of the Nordland region onshore Norway (Figure 1). The Trøndelag Platform, which defines the proximal part of the margin segment, is a large and shallow platform containing post-Caledonian Late Paleozoic to Late Mesozoic basins (Figures 1 and 4a). The distal and outer parts of the Vøring Basin province are characterized by NE-SW oriented structural basin highs (e.g., the Utgard High and the Nyk High) and deep subbasins (e.g., the Traena Basin, the Någrind Syncline, and the Hel Graben) (Figures 1 and 4a). The northern Vøring Basin is bounded by the Surt and Bivrost Lineaments defined between the adjacent central Vøring Margin segment and the Lofoten Margin to the north (Figure 1). The transition to the outer margin domain is strongly affected by Late Cretaceous-Paleocene extensional deformation within the so-called Grimm-Rym-Nyk structural “ring” which resulted in the formation of the deep Paleocene Hel Graben and the newly defined Hermod Basin (Figure 1). The outer domain is outlined by the Paleocene Hel Terrace and Grimm High; the latter is covered by relatively thin breakup-related volcanic sequences and is interpreted as a deep-seated structural high (Abdelmalak, Planke, et al., 2016).

3. Data Set and Methods

We interpreted a large set of high-quality seismic reflection lines acquired and processed by TGS (Figure 1b). The seismic data are time migrated with a positive amplitude at the seabed and are of good to excellent

Figure 2. Potential field data for the northern Vøring Margin. (a) The 50 km high-pass-filtered Bouguer data. (b) The 50 km high-pass-filtered magnetic data. COB = continent-ocean “boundary”, DT = Donna Terrace, FG = Fenris Graben, GH = Grimm High, GS = Gleipne Saddle, GrH = Grønøy High, HB = Helgeland Basin, HG = Hel Graben, HT = Halten Terrace, HmB = Hermod Basin, LR = Lofoten Ridge, NGR = North Gjallar Ridge, NH = Nyk High, NS = Någrind Syncline, JH = Jennega High, RB = Rås Basin, RH = Røst High, RR = Rån Ridge, RbB = Ribban Basin, RsB = Røst Basin, SGR = South Gjallar Ridge, SH = Sandflesa High, SkH = Skoll High, TB = Træna Basin, UH = Utgard High, UR = Utrøst Ridge, VB = Vestfjorden Basin, VDT = Valflesa Deep Terrace, VS = Vigrid Syncline. Data courtesy of TGS.

quality, although some deterioration occurs below sill intrusions. The potential field data were provided and processed by TGS. Well formation tops were obtained from the Norwegian Petroleum Directorate (NPD) (NPD factpages, 2017) and the Norwegian Offshore Stratigraphic Lexicon (NORLEX) group (Gradstein et al., 2010). The raw results of biostratigraphic analyses and wireline logs for well 6608/2-1S have been provided by the NPD but have been reevaluated by RPS Ichron (see Appendix A and Figure A1).

The seismic data set has been jointly interpreted together with potential field data (Figure 2) and published crustal-scale velocity data (Figure 1b) derived from ocean bottom seismometers (OBS) and expanded spread profiles (EPS) surveys (Digranes et al., 1998; Mjelde et al., 1998; Planke et al., 1991).

3.1. Seismic Interpretation

The regional seismic interpretation has been carried out using IHS Kingdom software. The line spacing of the final data set (Figure 1b) ranges from 0.2 to 2 km, which was sufficient enough to confidently tie the different seismic units mapped in this study. We interpreted a set of regional Cretaceous-early Cenozoic seismic horizons over the entire northern Vøring Margin (Figure 3). These units were correlated with the well database. The final regional horizons include the following: the Base Cretaceous Unconformity (BCU), mid-Albian (?) (MA), mid-Cenomanian (MC), Top Turonian (TT), intra mid-Campanian (IMC), the Base Tertiary Unconformity (BT), and Top Paleocene (TP) (Figure 3). Additionally, several pre-Cretaceous reflections were picked on the Trøndelag Platform, while Top Santonian (TS) and Base Late Pliocene (BLP) have been interpreted locally. We constructed a number of time-thickness (isochore) maps (Figure 11) in order to highlight the regional configuration of the main depocenters, their migration through time and space, and their relationship with the deep crustal architecture of the margin.

The crustal structure for the proximal part (Trøndelag Platform) was obtained from the reflection data where the interpreted top crystalline basement (TB) marker can be reliably identified over the area and tied with wells. To further constrain the crustal structure of the deep Vøring Basin, we compared the seismic interpretation with the top basement grid from a 3-D lithospheric model (previously published by Scheck-Wenderoth and Maystrenko, 2011). In the areas where the top basement grid crosses confidently interpreted sedimentary successions, we moved it down to a depth level consistent with the seismic observation since the new and reprocessed long-offset seismic data allow us now to more confidently identify deeper strong midcrustal reflections in the sedimentary basin. Most likely deeper than 10–11 s two-way travel time (TWT), the Moho is barely recognized on the seismic reflection data; as a result, we used OBS and ESP data and potential field modeling results to identify and map the Moho.

The 50 km high-pass-filtered Bouguer gravity anomaly map processed by TGS (Figure 2a) reveals a complex pattern of high- and low-amplitude anomalies along the margin. High-pass filtering of the data makes it possible to dissociate gravity anomalies caused by shallow basement structures from density variations in the deepest crust and/or the upper mantle. Cutoff wavelengths of 50 km emphasize density contrasts with source depth usually deeper than 10 km.

The 50 km high-pass-filtered magnetic data (Figure 2b) are based on the Mid-Norway Aeromagnetic Compilation by TGS, including the most recent ship track and aeromagnetic surveys in the Vøring Margin. Within the continental domain, these data allowed identification of the main deep-sourced magnetic anomalies. It provides information about the deep continental basement rocks expected below the nonmagnetic sedimentary cover. The 50 km high-pass-filtered potential field data were included in the seismic project, scaled to two-way travel time, converted to pseudohorizons, and displayed on the seismic lines to facilitate the joint interpretation of the seismic, gravity, and magnetic data.

Potential field modeling is a useful technique to evaluate and check further basement depth, nature, and properties. To illustrate the crustal configuration of the northern Vøring Margin, a 2-D potential field modeling has been carried out along a characteristic NW-SE section striking from the Helgeland Basin on the Trøndelag Platform up to the continent-ocean "boundary" (Figure 4). This section illustrates the main crustal configuration interpreted beneath the main structural highs and subbasins described in this study. The forward modeling has been carried out using the commercial software GM-SYS integrated within the GEOSOFT Oasis Montaj software. The modeling itself is based on the conventional and original method of Talwani (1973), where sets of irregular polygons with different physical properties along a 2-D transect create magnetic and gravity signals that are then compared with the observed potential field values. The geometry and properties of the polygons are changed accordingly until the best fit between the observed and calculated signal is achieved.

The 2-D margin transect was first depth converted using hiQbe™ Mid-Norway, a regional high-quality seismic velocity cube covering the entire mid-Norwegian Margin (<http://www.first-geo.com/products/hiqbe/norway>). The velocity cube has been compiled from 536 seismic stacking velocity data sets, including the stacking velocities along the transect, employing proprietary and industry standard geostatistical methods. The

Figure 4. Crustal transect across the northern Vøring Margin derived from (a) seismic observations and (b) potential field modeling (see Figure 1a for location). (c) Similar section at 1:1 scale with structural features discussed in the text. In the northern Vøring Margin we propose a model of crustal boudinage, where sag basins with thin and very thin crust are separated by rigid continental buffers. LC = lower crust, LCB = lower crustal body, MC = middle crust, ULCB = ultrahigh-density lower crustal body. Note the differences between the interpreted and modeled transect in the central Hel Graben, where we have to remove pre-Cretaceous sediments in order to get better fit with the gravity.

stacking velocities are calibrated to 230 check shots from wells, compensating for delta anisotropy—a parameter describing the near-vertical *P* wave propagation (Thomsen, 1986). At depths below well control, the delta anisotropy cannot be calculated directly from the well check shot. Calibration was then achieved by extrapolating the delta anisotropy downward in such a way that the hiQbe™ velocities are comparable to those derived from seismic refraction modeling of OBS data (e.g., Ni Dheasúna et al., 2012).

The software TecMod 2-D was also used to estimate total and tectonic subsidence rates for a series of rifting episodes in the Træna Basin calculated with forward modeling and classic backstripping approach. Kinematic, isostatic, and thermal principles of these approaches and their comparison are described in detail by Rüpke et al. (2008, 2010), Theissen-Krah and Rüpke (2010), and Clark et al. (2014).

4. Results

4.1. Geological Structure, Seismic Interpretation, and Correlation With Potential Field Data Along the Transect

Transect in Figure 4a illustrates the main basin and structural configuration of the northern Vøring Margin. The 342 km long transect strikes NW-SE from the Ylvingen Fault Zone at the Helgeland Basin on the Trøndelag Platform across the Vøring Basin and terminates close to the expected continent-ocean “boundary”.

The Trøndelag Platform defines the proximal and shallower part of the margin segment and shows Devonian-Early Carboniferous? to Jurassic basins (Blystad et al., 1995; Osmundsen et al., 2002) bounded to the east by the Caledonian basement that crops out in the onshore part. The structure of the thick Late Paleozoic basin (up to 6 km) is dominantly controlled by fault systems (Figure 4), which were possibly initiated during post-Caledonian orogenic collapse and then sequentially reactivated during Carboniferous?-Permian rifting episodes (e.g., Doré et al., 1999). To the west, this Late Paleozoic basin extends to a prominent basement horst lying at the northern prolongation of the Nordland Ridge (Figures 1a and 4a). Due to high density, this basement horst coincides with a pronounced gravity anomaly observed at that level (Figures 2b and 4b). We were able to tie a top basement marker to well 6609/7-1, where this horst structure was drilled and basement rocks were reached at a shallow level of 1,912 m (NPD factpages, 2017). The recovered rocks predominantly consist of pre-Cambrian quartzite and other low-medium-grade metamorphic rocks with evident sedimentary origin (Slagstad et al., 2011).

Later Mesozoic faulting activity in the area generated second-order half-grabens, grabens, and horst structures. Triassic to Jurassic sediments covered the deep Late Paleozoic basin and the basement horst and were subsequently peneplained in the Nordland Ridge at the BCU level. The superimposed Helgeland Basin (Figure 4a) observed in the inner platform contains a mainly Early Cretaceous sedimentary fill (Blystad et al., 1995; Brekke, 2000).

Our seismic interpretation shows basinward dipping fault systems at the northwestern flank of the Nordland Ridge/Grønøy High (Figure 5). Those faults controlled the Cretaceous evolution of the narrow and deep Træna Basin, which was poorly imaged by the previous seismic data sets. Compared to surroundings, the Træna Basin is characterized by NE trending negative gravity and magnetic anomalies (Figure 2). Despite a uniformly deep configuration of the basin along its strike, we observed a subtle increase of the magnetic and gravity signal (Figure 2) toward the Lofoten-Vesterålen Margin, which coincides with a strong seismic reflection underlying the deep-seated pre-Cretaceous fault blocks (Figure 5). The transition to the adjacent deep Rås Basin is marked by prominent north trending, relatively strong, positive gravity and magnetic anomalies (Figures 2 and 5) lying in the prolongation of the Surt Lineament, which is interpreted as an old inherited basement structure (Blystad et al., 1995). Our interpretation highlights a deep basin configuration (BCU at 7–8 s in the axial part; Figure 5). Regional mapping and our revised calibration suggest thin Lower Cretaceous successions (about 1 s TWT) underlying thick Coniacian to mid-Campanian sequences (Figure 5).

The northwestern flank of the Træna Basin is defined by the Utgard High. This prominent basement high, which is characterized by a strong NE trending gravity anomaly, can be divided into two segments (Figure 2a). The gravity anomaly is clearly connected to a prominent high-amplitude reflection observed in the deep part of the high (Figure 6). We have to note here that the observed reflection represents some local deep features and we were not able to establish any direct and spatial connection with the notable and similar deep *T*-Reflection defined and described farther west in the outer part of the Vøring Basin (see Abdelmalak

Farther to the northwest, the Utgard High flanks the Någrind Syncline, a 10 km thick deep Cretaceous sag basin better imaged now with the new seismic data (Figure 7). The Surt Lineament separates the Någrind Syncline from the Vigrid Syncline in the southwest, whereas its northern continuation is limited by the Bivrost Lineament (Figure 1a). Like the Træna Basin, the Någrind Syncline also corresponds to strong negative gravity and magnetic anomalies (Figure 2). The tectonic history of the Någrind Syncline is strongly related to the Late Jurassic to Paleocene development of the adjacent highs (the Nyk and Utgard Highs) and the

Figure 6. Structure of the Utgard High and configuration of the ULCB. Southern segment: (a) uninterpreted and (b) interpreted seismic sections. Northern segment: (c) uninterpreted and (d) interpreted seismic sections. White arrows on the uninterpreted sections point at the *UH-Reflection* interpreted as the top of the ULCB. Note the significant increase of the gravity and magnetic signals (G50 and M50, respectively) in the northern segment of the high. The common legend for the sedimentary part is shown in Figure 4. The abbreviations for the horizons are given in section 3.1. The location of the seismic line is shown in Figure 1a. Data courtesy of TGS.

structural evolution of the flanking/bounding transfer zones (e.g., the Surt and Bivrost Lineaments) (Blystad et al., 1995; Brekke, 2000). We found evidence that the Jurassic-early Cretaceous rifting did not terminate here at BCU level (Berriasian, according to Dalland et al., 1988) but prolonged into the Neocomian (Barremian-Aptian?) time. The early and late synrift wedges of Jurassic-Early Cretaceous and possible Neocomian age respectively were controlled by normal faulting on the flank of a deep horst lying below the Nyk High (Figures 4a and 7).

The nearly NE trending positive gravity and magnetic trend in the Nyk High correlates well with a shallow basement high observed at that level (Figure 2). The anomalies curve in an almost southern direction within the Surt Lineament mimicking the deep structural configuration. The dominant and shallower structural style of the Nyk High at the present day represents the uplifted and eroded footwall of a rifting system, which initiated in Late Cretaceous time and was active during most of the Paleocene (Gernigon et al., 2003; Mogensen et al., 2000; Ren et al., 2003). Rift shoulder uplift of the Nyk High was concomitant with a significant subsidence of the adjacent Ngrind Syncline, which is not particularly affected by classic rotated fault blocks. The ongoing extension during the Paleocene promoted severe collapse of the western flank of the Nyk High, thus creating significant accommodation space resulting in preservation of a thick Paleocene succession observed and drilled (e.g., well 6706/6-1) in the Hel Graben (Figure 4a). Later Paleocene-early Eocene minor sinistral transpressional reactivation complicated the fault pattern within the Nyk High (Imber et al., 2005).

There is still an ongoing debate on the Late Cretaceous-Paleocene development of the Hel Graben. The more recent interpretations rely on the controversial stratigraphic analysis of well 6706/6-1. The NPD claims that there is at least 1,400 m of Paleocene deposits in the well (NPD factpages, 2017) with no penetration within Upper Cretaceous sediments expected at that level in predrilling interpretation (Blystad et al., 1995; Ren et al., 2003). In contrast, the NORLEX team (Gradstein et al., 2010, and well report at <http://www.nhm.uio.no/norlex>) interprets the same section as comprising Upper Cretaceous Shetland Group sediments, suggesting only 200 m of Paleocene sediments in the well; this interpretation implies that the Hel Graben is dominantly a Late Cretaceous rather than Paleocene synrift feature (e.g., Fjellanger et al., 2005). Our own seismic tie

Figure 7. Seismic line across the Nâgrind-Nyk segment of the study area: (a) uninterpreted and (b) interpreted. Here we can recognize Jurassic and Neocomian synrift wedges developed on the eastern flank of the Nyk High. The deep and old synrift sequences are controlled by an east dipping faults. Maastrichtian-Paleocene rifting in the area was associated with west dipping faulting, the uplift of the Nyk High, and significant subsidence of the Nâgrind Syncline, where a large depocenter was formed. The common legend for the sedimentary part is shown in Figure 4. The abbreviations for the horizons are given in section 3.1. The location of the seismic line is shown in Figure 1a. Data courtesy of TGS.

from the adjacent Vigrid Syncline into the Hel Graben across the Rym Fault Zone (Figure 8) does support the presence of thick Paleocene sediments in the Hel Graben. However, our interpretation of the deepest parts of the structure is less confident due to poor imaging caused by extensive distribution of magmatic intrusives, making our assessment of the pre-Paleocene structure more speculative. However, seismic imaging locally allows to estimate the relative depth of the BCU (Figure 8).

The Hel Graben is characterized by a rounded negative gravity anomaly (Figure 2a). It laterally passes into the Paleocene Hermod Basin interpreted as a separate subbasin in this study (Figures 1a and 8). The limit between the two subbasins is characterized by a deep-seated high (Figure 8). The gravity pattern shows that the Hermod Basin may extend northeast toward the outer Lofoten Margin (Figure 2a) and possibly merge with the Røst Basin as recently suggested by Maystrenko et al. (2017). The 3-D potential field modeling of Maystrenko et al. (2017) suggests that thick Late Cretaceous-Paleocene sequences (undifferentiated) may likely exist underneath the Lofoten lava inner flows.

Figure 8. Seismic tie from the Vigrid Syncline to the Hel Graben revealing the deep configuration of the basin formed underneath thick Paleocene successions: (a) uninterpreted and (b) interpreted lines. The Hel Graben and the adjacent Hermod Basin are structurally divided by a deep-seated structural high. The common legend for the sedimentary part is shown in Figure 4. The abbreviations for the horizons are given in section 3.1. The location of the seismic line is shown in Figure 1a. The 50 km high-pass-filtered Bouguer gravity (G50) and magnetic (M50) anomalies are shown. Data courtesy of TGS.

The northern boundary of the Hel Graben is defined by the Hel Terrace. Together with the Hermod Basin, this is also a newly defined structural element. It is characterized by a series of southward dipping Late Cretaceous-Paleocene growth faults (Figure 9a) possibly decoupled and developing over overpressured mobile shales widely distributed in Lower to mid-Cretaceous intervals in the deep Vøring Basin (Brekke et al., 1999; Dalland et al., 1988; Vergara et al., 2001). Farther northeast, the northern flank of the Hel Terrace is characterized by a graben-like feature with clear synrift Paleocene seismic wedges postdating the main Late Cretaceous-Paleocene extensional phase leading to the formation of the Hel Graben (Figure 9b). The new seismic data also suggest that the footwall of the Hel Terrace was exposed and peneplained during this period. Notably, we discovered evidence of earlier pulses of magmatism (intra-Paleocene) in the Hermod Basin compared to the main early Eocene volcanic event recognized in the Vøring Marginal High. This early phase of magmatism is witnessed by older vent complexes that developed in the Paleocene successions and genetically linked with deeper sill intrusions (Figure 9b).

The subbasalt Grimm High (Figures 4 and 10) has been defined based on a previous integrated seismic-gravity-magnetic interpretation of the volcanic margin (Abdelmalak, Planke, et al., 2016). It coincides with a gravity high that correlates with a positive magnetic anomaly, suggesting a shallow basement beneath the basalt (Figures 2 and 10). The anomalies also correlate well with a structural high interpreted at the top and base basalt levels at the Vøring Marginal High (Abdelmalak, Planke, et al., 2016). The Grimm High has a thin

Figure 9. Seismic lines across the Hel Terrace and the Hel Graben. (a) Uninterpreted and (b) interpreted lines: central part of the Hel Terrace with Paleocene rift structures (growth faults) that are controlled at depth by probable lower-mid-Cretaceous mobile shales. Later in the Paleocene, the structure was mildly reactivated in strike-slip regime. The increase of the gravity signal toward the Vøring Marginal High could correspond to the shallowing of the BCU level and presence of pre-Cretaceous high/terrace below volcanics. Eastern part of the Hel Terrace: (c) uninterpreted and (d) interpreted lines. Late Cretaceous synrift sequences could develop in the central part of the structure and then superimposed by thick Paleocene sediments related to the collapse of the Hel Graben. Later in the Paleocene, the northern flank of the Hel Terrace was affected by rifting and associated rift-shoulder uplift and erosion. This event was preceded by minor sill intrusions expressed by a venting developed in Paleocene sequences. Major magmatic activity occurred at the Paleocene-Eocene transition with extensive sill intrusions (shown in purple) and development of magmatic sequences at the Vøring Marginal High (inner flows). The 50 km high-pass-filtered Bouguer gravity (G50) and magnetic (M50) anomalies are shown for Figure 9a only. The common legend for the sedimentary part is shown in Figure 4. The abbreviations for the horizons are given in section 3.1. The location of the seismic lines is shown in Figure 1a. Data courtesy of TGS.

Figure 10. A part of the studied depth-converted transect (see Figure 4a) adjacent to continent-ocean boundary showing subbasalt configuration of the Grimm High. Tilted fault blocks below volcanics could develop within Cretaceous sequences on the analogue with the North Gjallar Ridge (see Gernigon et al., 2003). At about 15 km deep and faulted crustal reflection (*T*-Reflection) is observed. The Base Tertiary and Base Cretaceous Unconformities are shown in dashed green and dotted red, respectively. Data courtesy of TGS.

basaltic cover (Abdelmalak, Planke, et al., 2016), reflecting its relatively high elevation when subaerial volcanic sequences were deposited in the earliest Eocene. Notably, the gravity high observed at that level also fits a high-amplitude reflection observed at 15 km in the deep part of the Grimm High (Figures 4a, 4b, and 10). The deep reflection coincides spatially with the deep and faulted *T*-Reflection described and defined in the outer Vøring Basin (Abdelmalak et al., 2017; Gernigon et al., 2003, 2004). At the level of the Grimm High, the new seismic data revealed tilted fault block system below the volcanics (Figure 10), with a structural style similar to that observed in the adjacent North Gjallar Ridge (Gernigon et al., 2003, 2004; Ren et al., 2003). The upper synrift feature could represent the equivalent of the Campanian-Maastrichtian Springar Formation penetrated by well 6704/12-1 in the North Gjallar Ridge (Fjellanger et al., 2005; Gernigon et al., 2003).

Figure 11. Isochore maps of seismic sequences K10 through P10 showing general distribution and migration of the basin depocenters in time and space. Present structural outline of the main structural highs is shown in transparent white. The red line shows a landward limit of basalt flows Extent of the maps is shown in Figure 1a. GrH = Grønøy High, HB = Helgeland Basin, HG = Hel Graben, HmB = Hermod Basin, HT = Hel Terrace, NGR = North Gjallar Ridge, NH = Nyk High, NS = Någrind Syncline, RFZ = Rym Fault Zone, TB = Træna Basin, UH = Utgard High, VS = Vigrid Syncline.

4.2. Regional Development of the Main Cretaceous-Paleocene Tectonostratigraphic Units

Having described the present margin geometry, we here infer its evolution based on different isochore maps (Figure 11). We organize this from oldest to youngest, focusing on thickness trends and onlap patterns around major subbasins and flanking structural highs.

4.2.1. Unit K10 (Valanginian to Mid-Albian)

The Valanginian to mid-Albian sequence K10 defined in this study is rather reinterpreted as a synthinning sequence and not a postrift as suggested in previous publications (Færseth & Lien, 2002; Vergara et al., 2001). The isolated rounded depocenter in the Någrind Syncline reflects the Neocomian (Barremian-Aptian?) faulting/rifting activity observed on the eastern flank of the Nyk High (Figures 7 and 11a). However, no prominent and concomitant depocenter is identified within the Træna Basin (Figure 11a). The strata of the K10 sequences onlap the deep terraces imaged in the southern part of the Træna Basin and along the northwestern flank of the Nordland Ridge, which was already an elevated feature at that time (Figure 11a). The uplift of the Nordland Ridge was also associated with lateral subsidence of and sedimentation within the Helgeland Basin, which tends to deepen in the northeastern direction (Figure 11a). As previously interpreted (Blystad et al., 1995; Brekke, 2000), the Lower Cretaceous succession thins toward the Utgard High, suggesting that a local and central horst configuration was already initiated at the early stage of the Vøring Basin formation. Slight thinning and local absence of the K10 sequence on top of the Utgard High is confirmed by our seismic observations (Figures 6 and 11a).

Nevertheless, rather thick Lower Cretaceous sediments are also confirmed by well 6608/2-1S in the northern Utgard High. This reflects a complex structural topography at the early stages of the high formation.

The absence and thinning of the Valanginian to mid-Albian K10 sequence in the southern and central Nyk High suggests NE striking basement/BCU highs, which were already emerged at that time. Several minor depocenters are observed at the transition between the North Træna Basin and the adjacent Lofoten-Vesterålen Margin (Figure 11a).

4.2.2. Unit K20 (Mid-Albian to Mid-Cenomanian).

Similar depositional pattern can be deduced from the mid-Albian to mid-Cenomanian isochore map (unit K20, Figure 11b). Clear erosion is seen on the Nordland Ridge and partly on the deep terraces mapped in the southeastern part of the Træna Basin (Figure 11b). According to the sediment thickness distribution, the Træna Basin was still characterized by a slow sedimentation rate during the K20 deposition. Thinning of the sequence K20 is also interpreted toward the Utgard High (Figure 11b). However, locally poor quality of seismic imaging there makes such interpretation a bit uncertain. We identified several scattered depocenters blanketing the Någrind Syncline and its transition to the Vigrid Syncline and partly the Nyk High (Figure 11b). We also observe significant thickening of the K20 at the transition between the Någrind Syncline and the Lofoten-Vesterålen Margin.

4.2.3. Unit K30 (Mid-Cenomanian-Early Coniacian)

Deposition of the mid-Cenomanian-Early Coniacian unit K30 mainly occurred in the Træna Basin, which defined a deep (up to 1.7 s TWT), prominent and NE-SW elongated (100 × 25 km), fault-controlled depocenter that deepened toward the southern Rås Basin (Figure 11c). The depocenter-related normal faults tended to develop over older fault systems observed on both flanks of the Træna Basin (Figure 4a). Both seismic observations and isochores indicate that the synthinning subsidence shifted northeastward from the Rås Basin. During the same period, a large and relatively thick rounded depocenter (55 × 45 km; up to 1–1.5 s TWT thick) developed in the adjacent Någrind Syncline. Thinning of the K30 unit is observed on the SE flank of the North Gjallar Ridge and in the Utgard Ridge, indicating their relatively high structural position during this period (Figure 11c).

4.2.4. Unit K40 (Coniacian to Mid-Campanian)

Rapid subsidence and accommodation of a large, NW-SE elongated (150 × 30 km) thick depocenter (up to 2.9 s TWT or 6 km thick) during the Coniacian to mid-Campanian K40 period (Figure 11d) concentrated in the Træna Basin. Based on seismic observations, we suggest that fault-driven tectonic subsidence ceased around Santonian and then the Træna Basin was passively infilled during the early to mid-Campanian. Thinning and erosion of the K40 unit on both flanks of the Utgard High confirm its shallow bathymetric configuration at that period (Figure 11d). Rather high sediment accumulation rates persisted along the axis of the Någrind Syncline with the deposition of the Nise Formation sourced from NE Greenland (Morton & Grant, 1998). In the Campanian, the North Gjallar Ridge was still in a relative high position and shallow marine setting as illustrated by the absence and/or pinch out of the K40 unit on its SE flank as confirmed by well 6705/10-1 (Asterix). Furthermore, thinning of the unit K40 toward the Hel Terrace (Figure 11d) also indicates a shallower structure in that domain.

4.2.5. Unit K50 (Mid-Campanian to Base Tertiary)

A dramatic depocenter shift of the unit K50 occurred from the mid-Campanian to the beginning of the Tertiary (e.g., the Base Tertiary Unconformity). The main sedimentation axis moved northwestward from the Træna Basin to the Någrind Syncline (Figure 11e). The depocenter shape mimics the preexisting basin configuration developing in between deep-seated structural highs (Figure 11e). In the SW Någrind Syncline and its transition to the Vigrid Syncline, the depocenter is seemingly not fault controlled. However, subsidence and higher sedimentation in the NE segment of the Någrind Syncline could be enhanced by additional reactivation of the SE dipping faults locally observed on the flank of the adjacent Nyk High (Figure 4a). The sequence K50 is thinned significantly along the Nyk High and the North Gjallar Ridge, which were both uplifted during this period. Smaller thicknesses of K50 are also observed on the Utgard High. Notably, the Hel Terrace was still in a shallower position with K50 onlaps onto its southern flank, suggesting a structural evolution quite similar to the adjacent North Gjallar Ridge.

4.2.6. Unit P10 (Paleocene)

Another marked change in sedimentation pattern happened during deposition of the sequence P10 in the Paleocene, when the depocenter suddenly migrated northwestward toward the Hel Graben and the

Table 1
Density and Magnetic Properties of the Units Modeled in the Present Study

Layers	Density ($\text{kg} \cdot \text{m}^{-3}$)	Susceptibility (SI units)	Remanence (A/m)
Water	1,450	0	0
Mid-Pliocene to Holocene	1,800–1,900	0	0
Mid-Miocene to Mid-Pliocene	2,000	0	0
Eocene to Mid-Miocene	2,050	0	0
Basalts/SDR	2,550	0.01	-1 (Upper); +1 (Lower)
Paleocene	2,225	0	0
Intra Mid-Campanian to Maastrichtian	2,400	0	0
Lower Campanian to Intra Mid-Campanian	2,400–2,525 (deep sag)	0	0
Mid-Coniacian to Upper Santonian	2,525–2,575 (deep sag)	0	0
Mid-Cenomanian to Lower Coniacian	2,525–2,600 (deep sag)	0	0
Mid-Albian to Mid-Cenomanian	2,550–2,675 (deep sag)	0	0
Valanginian to Mid-Albian	2,550–2,700 (deep sag)	0	0
Pre-Cretaceous (Und)	2,650–2,775 (deep sag)	0	0
Callovian to Berriasian	2,500	0	0
Toarcian to Bathonian	2,550	0	0
Hettangian to Pliensbachian	2,559	0	0
Upper Triassic	2,575	0	0
Mid-Triassic	2,585	0	0
Mid-Permian to Lower Triassic	2,600	0	0
Upper Devonian? -Permian	2,700	0	0
Upper crust (approximately Caledonian nappes)	2,750	0.01	0
Middle crust	2,850	0.03	0
Lower crust	2,950	0.005	0
High-density lower crust (LCB)	3,100	0.005	0
Ultrahigh density lower crust (ULCB)	3,200	0.015	0
Mantle	3,300 (Lower)–3,250 (Upper)	0	0

Hermod Basin (Figure 11f). In the central part of the outer Vøring Basin, faulting migrated progressively toward the volcanic domain growing at the same period (Gernigon et al., 2004). In the Hel Graben and the Hermod Basin deposition and subsidence were largely controlled by a collapse of the NW flank of the Nyk High and facilitated by a slip along the Rym Fault Zone. Additional evidence for later Paleocene faulting/rifting is seen in the northern part of the basin (e.g., the Hel Terrace) (Figures 9c and 9d).

Minor depocenters in the Vigrid/Någrind syncline axis and near the Lofoten-Vesterålen Margin (Figures 7 and 11f) are not fault controlled; instead, they are likely interpreted as sedimentary bodies (e.g., fans) sourced from the neighboring eroded highs. Erosion of the sequence P10 in the Utgard High is possibly the result of late stages of subsequent Early Tertiary strike-slip reactivation along the Fles Fault Complex as previously noted by Blystad et al. (1995).

4.3. Potential Field Modeling of the Regional Transect and Crustal Configuration

The 2-D forward potential field modeling of the northern Vøring transect (Figures 4a and 4b) was carried out to better constrain the crustal architecture and basement properties underlying the complex depocenter system previously described in this study. The initial crustal parameters used in the potential field modeling are shown in Table 1 (see also Appendix B).

While the Moho configuration in the Trøndelag Platform is constrained by several ESPs (Planke et al., 1991), the depth of top basement was entirely obtained from the new seismic reflection data, where the expected top basement surface represents a well-defined seismic marker. The total crustal thickness decreases down to 10 km in the axial part of the platform, where we have to presume attenuated upper to lower crust and exhumation of the high density, lower crustal body ($3,100 \text{ kg/m}^3$) at 15 km depth; an interpretation consistent with the high velocities (around 7.1 km/s) and depth estimations was taken from the available ESP data (Planke et al., 1991).

The transition zone to the Træna Basin corresponds to an abrupt thinning of the continental crust over less than 40 km from thickness in 18 km at the Nordland Ridge to 4–5 km in the central axis of the Traena Basin

(Figure 4b). As suggested by Maystrenko et al. (2017), the first necking of the rift system is located in this position. The Moho shallows from 25 km to 20 km in the Utgard-Træna area (Mjelde et al., 1998; Planke et al., 1991).

The northern Utgard High forms the most prominent magnetic and gravity signals along the profile (Figure 4b). In order to fit the prominent gravity signature and surrounding refraction data, we assumed a thick (10–12 km), ultrahigh-density body (ULCB) ($3,200 \text{ kg/m}^3$) from Moho level to relatively shallow position (8–10 km) in the lower and upper parts of the crust. Based on our modeling results, we have to assume 1 km thick midcrustal lenses, which are most likely preserved on both flanks of the structure and almost in contact with the meta-sedimentary prerift and synrift rocks. According to our seismic observations, the top of the ULCB deduced from potential field modeling correlates well with the *UH*-Reflection (Figures 4b and 6) implying a significant impedance (density/velocity) contrast at that level. The basement below the Træna Basin necessitates massive and very high density and magnetic rocks (Figure 4b). Our modeling results do not support the presence and/or preservation of deeper Mesozoic sedimentary rocks in depth. Accordingly, we suppose that the thin and ultrahigh-density lower continental crust may directly underlie the basin possibly coinciding with the strong reflection observed within pre-Cretaceous fault blocks (Figure 5).

A minimum crustal thickness of 5–6 km in the deep-sag Nâgrind Syncline is expected at the connection to the NW flank of the Utgard High (Figure 4b). The top of the basement is reliably defined in reflection seismic and correlative with refraction data lying at 13 km at the cross point with OBS line 7-96 (Mjelde et al., 1998). The basement shallows to 7.5 km toward the Nyk High and forms a prominent paleohigh with total crustal thickness in 13–15 km also confirmed by OBS line 7-96 (Mjelde et al., 1998) and ESP data (Planke et al., 1991). Lens-shaped bodies of middle, lower crust and a dense LCB (1 to 3 km thick apiece) are required there to fit the observed potential field data.

The axial part of the Hel Graben is not crossed by any refraction profiles (Figures 2b and 4b). We modeled that the Moho rises to 18 km depth, which corresponds to a drastic thinning of the crust reduced to 3–4 km thick in the central part of the Hel Graben. Since we believed that the BCU was confidently interpreted at the position of the transect, we made structural corrections for the pre-Cretaceous configuration in the central Hel Graben (Figure 4a). Based on our modeling, we had to agree on extremely thin or no pre-Cretaceous sediments below this unconformity in order to get a better fit with the observed gravity (Figure 4b). As a result, the Cretaceous sediments most likely lie directly on top of a high-density lower crust also expected in the central part of the Hel Graben (Figure 4b). This crustal configuration also implies that upper to lower crust is most likely substantially delaminated or absent there. The presence of pre-Cretaceous sediments is not totally excluded, but if present, these metamorphosed sequences are extremely compacted and/or highly deformed. With realistic magnetic susceptibility values in the basement, we have not been able to reach a solid fit with the observed magnetic anomalies at the Hel Graben (Figure 4b). This discrepancy could be explained by the magnetic influence of the large shallow sill complex observed in the Hel Graben (e.g., Berndt et al., 2000) but is not considered in our study.

Presence of the deep-seated continental basement block underneath the Vøring Marginal High within the subbasalt Grimm High (Abdelmalak, Planke, et al., 2016) is in agreement with our modeling results (Figure 4b). A crustal fragment with a thickness of more than 10 km is expected there. The Moho beneath the Grimm High is also relatively well constrained at 25 km from previous OBS and ESP data (Mjelde et al., 1997, 1998; Planke et al., 1991) and then shallows to less than 20 km toward the continent-ocean “boundary”. The prolongation of the *T*-Reflection expected at that level partly corresponds to the boundary between the lower crust and the LCB bodies modeled in the distal part of the transect (Figures 4a and 4b). To fit the gravity, our model also suggests that up to 3 km of pre-Cretaceous strata could be preserved above the continental basement expected underneath the basalt.

5. Discussion

5.1. Origin and Geophysical Properties of the Lower Crustal Bodies (LCBs)

5.1.1. Origin of the Utgard Ultrahigh-Density Body (ULCB)

Below the Utgard High our modeling suggests the presence of an up to 12 km thick, ultrahigh-density (3200 kg/m^3) lower crustal body (Figure 4b), the top of which is located almost at the level of the *UH*-Reflection. The shallowing of the *UH*-Reflection in the central and northern parts of the Utgard High fits

Figure 12. Cartoon illustrating Cretaceous-Paleocene evolution of the northern Vøring Margin.

well with the main magnetic and gravity signals (Figures 6c and 6d). The nature of these potential field anomalies has been extensively discussed. Earlier models (Planke et al., 1991; Skogseid et al., 1992) speculated on the crustal origin of the anomaly, implying that it most likely represents an old, Caledonian, high-density basement block with Mesozoic or pre-Mesozoic age of emplacement. Our observation shows (Figures 6c and 6d) that the *UH*-Reflection corresponds to the top of the local LCB and has been affected by a series of Mesozoic (and older?) faulting events. Consequently, we also considered that the Utgard High LCB (or part of it) is most likely an inherited basement structure.

Mjelde et al. (2013, 2016) also proposed that the basement here partly consists of lower crustal eclogites with a density of up to $3,500 \text{ kg/m}^3$. These eclogites may be related to a sliver of the Caledonian suture zone (e.g., Petersen & Schiffer, 2016). In a recent 3-D lithospheric model of the Lofoten-Vesterålen margin and adjacent parts of the Vøring Margin, a thick high-density lower crustal layer of Caledonian origin, with high densities varying from $2,985 \text{ kg/m}^3$ to $3,070 \text{ kg/m}^3$, has also been considered beneath the Utgard High (Maystrenko et al., 2017).

Reynisson et al. (2010), based on their regional 3-D potential field modeling, interpreted part of the LCB within the Utgard High as heavily (and synrift) serpentinized mantle, assuming very thin crust underneath the high. Our modeling and observations suggest, however, that the Utgard High was an old crustal horst partly shielded from drastic thinning (Figures 4 and 12), compared to the surrounding subbasins (e.g., Træna Basin and Nāgrind Basin). The clear correlation with the central horst system acting as a thick tectonic buffer since the early stage of rifting in the northern Vøring Margin suggests that the high-density rocks were most likely part of the prerift basement even before the onset of drastic thinning of the crust. Wangen et al. (2011) also got unrealistic rates of extension in their tectonic modeling run with a shallow and thin crust at the position of the Utgard High. Therefore, we suggest that the nature of the anomaly is better regarded as a remnant of old Caledonian or Precambrian high-density crustal blocks. Speculative serpentinization within the Utgard High is unlikely for us. This system of crustal blocks may extend toward the Lofoten-Vesterålen Margin and merge with either the Lofoten Ridge or the Røst High, which are also expressed as prominent, elongated, positive gravity anomalies. It is likely that the anomaly could also be amplified by significant intrusion of breakup-related, high-velocity, dolerite sills ($>7.0 \text{ km/s}$) as suggested by Berndt et al. (2000) and Neumann et al. (2013). The density expected at the level of the ULCB is so high ($3,200 \text{ kg/cm}^3$) and so massive that we disregarded the possibility of preserved deep sediments at that level. This is in contradiction with recent alternative assumptions from Osmundsen et al. (2016). Shallow magnetic source estimations

(Figure 4b) also support the presence of magnetic (basement) rock at 10–15 km and accordingly cannot easily explain the presence of Mesozoic sediments at that level.

5.1.2. Outer LCB

To fit both the observed gravity and the high velocities also deduced from ESP and OBS data, our modeling also requires a high-density LCB ($3,100 \text{ kg/cm}^3$) in the outer part of the northern Vøring Basin and below the Grimm High (Figure 4b). This body is typically characterized by V_p velocities between 7.1 and 7.7 km/s (Mjelde et al., 2003). The nature of the outer LCB located along continent-ocean boundary has been widely discussed by many authors in the past. Earlier schemes suggest that LCB in the distal part of the mid-Norwegian Margin represented mafic crust underplated during continental breakup (e.g., White & McKenzie, 1989). However, later studies promoted a variety of alternatives on the origin and properties of LCB, ranging from high-velocity intrusions into the lower crust (White et al., 2008, 2010), heavily serpentinized/exhumed mantle (Osmundsen & Ebbing, 2008; Péron-Pinvidic & Osmundsen, 2016; Ren et al., 1998; Reynisson et al., 2010), and/or a retrograde/high-grade Caledonian lower crust (Abdelmalak et al., 2017; Ebbing et al., 2006; Gernigon et al., 2004; Mjelde et al., 2016; Petersen & Schiffer, 2016).

According to our modeling results (Figure 4b) and new observations (Figure 10), the upper boundary of the outer LCB coincides partly with the subbasalt prolongation of the T-Reflection located at approximately 15 km depth. This high-amplitude reflection also correlates with a strong gravity signal within the Grimm High (Figure 2a). Similar high-amplitude reflection has been earlier described and mapped within the South and North Gjallar Ridges (Gernigon et al., 2003, 2004), where it is also interpreted as a part of an old (possibly Caledonian) inherited piece of lower crust (Ebbing et al., 2006; Gernigon et al., 2004). The T-Reflection influenced the basin evolution long before the onset of the breakup magmatism and as early as the initiation of Late Cretaceous–Paleocene rifting in the mid-Late Campanian (Gernigon et al., 2003, 2004). Furthermore, in the recent study by Abdelmalak et al. (2017), the T-Reflection in the Hel Graben was also interpreted as the top of LCB, where it coincides with lower crustal sill intrusions and detachment faults. An alternative interpretation of the prebreakup emplacement of this LCB may also be explained in terms of serpentinization model in the North Gjallar Ridge (Ren et al., 1998). However, seawater penetration into the mantle could not be easily explained if we consider the thick (8–9 km) prebreakup sedimentary package and the associated decoupled fault system overlying the deep LCB (Figure 10). Thus, in our model we agree that an inherited crustal origin of the LCB is most likely for this distal part of the margin.

5.2. Basin Evolution and Crustal Deformation

5.2.1. Træna Basin Revisited: Too Narrow and Deep To Be Just a Postrift Cretaceous Sag Basin?

Very few papers have described the Træna Basin in detail (Blystad et al., 1995; Gomez et al., 2004; Osmundsen et al., 2002). According to Osmundsen et al. (2002), the Træna Basin represents a large rollover structure controlled by an east dipping detachment fault along the southeast side of the Utgard High, which influenced the subsidence of the basin and the formation of two large Lower and Upper Cretaceous depocenters. In contrast, Gomez et al. (2004) proposed that the Træna Basin was rather controlled by a large west dipping detachment fault defining the NW flank of the Nordland Ridge, whereas the NW dipping flexure of the Nordland Ridge was explained by Late Cretaceous, postrift thermal subsidence. We do not see any evidence for a significant west dipping detachment faulting at the BCU or deeper level along the flank of the Nordland Ridge and its transition to the Træna Basin. The Træna Basin could be initiated as a half-graben on the flank of the Utgard High in the Jurassic–Early Cretaceous, with the Nordland Ridge being uplifted and peneplained (Blystad et al., 1995; Brekke & Riis, 1987) as a flexural response during this rifting event, as later proposed by Osmundsen et al. (2002). According to our observations, the subsequent formation of the elongate, narrow, and extremely thick Upper Cretaceous depocenter of the Træna Basin is fault controlled on both sides of the basin but is not only structurally limited to the southeastern side of the Utgard High. Therefore, the existence of an active and persistent west dipping detachment through the entire thinning history of the Træna Basin is questionable.

The conjugate fault systems observed on both sides of the Træna Basin (Figure 4a) are sealed by Santonian strata and then were moderately reactivated mainly along the Utgard High as a continuation of the Fles Fault Complex in latest Cretaceous–Paleocene times. One could explain this faulting as a simple result of differential compaction during a long period of postrift thermal sagging of the basin (Færseth & Lien, 2002). However, the backstripped tectonic subsidence curve (Figure 13) calculated for the central part of the Træna Basin shows

Figure 13. Total and tectonic subsidence curves calculated by different methods (forward modeling and backstripping) on the axial part of the Træna Basin. Extreme subsidence rates obtained for mid-Cenomanian-Santonian time are typical for strike-slip related basins and also can be attributed to rift basins associated with steep-dipping faults. Subsidence curves were computed using the TecMod 2-D software.

an abrupt transition to relatively rapid subsidence magnitude (~2.5 km) starting in the mid-Cenomanian (~95 Ma) and ceasing at Santonian (~85 Ma). Such rapid subsidence (Figure 13), the atypical high rates of sedimentation during this geologically short time period (10 Myr), and the narrow basin configuration (~20 km wide) cannot be easily explained by a McKenzie-like thermal subsidence model (McKenzie, 1978) invoking uniform and instantaneous pure shear extension. Narrow, small basins characterized by accelerated and short-lived tectonic subsidence (2–4 km in 10 Myr) typically form in pull-apart settings, where transtensional strike-slip regime is involved (Christie-Blick & Biddle, 1985; Xie & Heller, 2009). Typical strike-slip basins (e.g., Dead Sea Rift (ten Brink & Ben-Avraham, 1989)) tend to lengthen rather than widen during ongoing extension. Transtensional regimes also favor much faster and efficient thinning of the crust, which seems to be the case in the Træna Basin according to the very thin crust expected there (Figure 4b). Notably, Early to Late Cretaceous oblique extension and dextral strike slip have been described in the adjacent Lofoten-Vesterålen Margin (Bergh et al., 2007).

The observed fault systems controlling the mid-Cenomanian-Santonian subsidence in the Træna Basin show relatively steep configuration (Figure 4a). However, direct evidence for both large-scale strike-slip motions (e.g., negative flower structures) is not well imaged along the basin. Furthermore, the backstripped tectonic subsidence rate in 2.5 km is not uncommon for rift basins (Allen & Allen, 2013). The formation of steep-dipping faults in a narrow rift basin could explain ongoing rapid subsidence even after rift opening had slowed, as is shown in the Rio Grande Rift (van Wijk et al., 2018). In this case, local transtensional reactivation or extension by slip on steep-dipping faults could explain the mid-Cenomanian-Santonian structural evolution of the narrow, deep Træna Basin. Unfortunately, our observations and modeling results cannot fully discriminate between the two scenarios. Nonetheless, a local extensional component during the Cenomanian-Santonian thinning of the Træna Basin could be concomitant with the uplift and tilting of the Trøndelag Platform (Brekke, 2000), the Nordland Ridge (Fjellanger et al., 2005), the Lofoten Margin (Henstra et al., 2016), and the Utgard High. Our biostratigraphic study of well 6608/2-15 in the northern Utgard High (see appendix and Figure A1 for details) reveals the presence of a significant Turonian hiatus that could be a result of vertical movements linked with this possible thinning event. Mid-Coniacian extensional reactivation along the flanks of the Træna Basin was also assumed in the regional paleobathymetric model by Roberts et al. (2009). However, the authors did not provide any structural examples for that.

In previous publications, the Træna Basin was characterized by a thick (2 s TWT) Lower Cretaceous succession and the Base Cretaceous Unconformity lying at approximately 6–7 s TWT (Blystad et al., 1995), shallowing toward the Lofoten-Vesterålen Margin. In previous study, Osmundsen et al. (2002) interpreted the BCU at a deeper level (8 s TWT) but also presumed a thick Lower Cretaceous succession (up to 3 s thick TWT). Our interpretation and regional seismic tie consider thinner Lower Cretaceous sediments in the Træna Basin (Figures 4a, 5, and 11a). The absence of thick Lower Cretaceous depocenters in the Træna Basin is also in

contrast to Vergara et al. (2001), who proposed regional sediment passive infill of inherited rift-related Jurassic lows during the Berriasian to Albian. On the one hand, the Træna Basin could have been a sediment-starved unfilled rift system at this time. However, relatively thick Lower Cretaceous strata are also preserved on top of the Utgard High “crustal buffer” (Figure 6d), which was located in a shallower position at that stage (Figure 11a). Accordingly, we speculate that much thicker Early Cretaceous sediments would have most likely coexisted in the adjacent subbasin. To explain such an ambiguity, we propose a crustal scenario where pre-Cretaceous and partly pre-mid-Albian successions were most likely deposited in the proto-Træna Basin but were subsequently drastically stretched during the Late Jurassic-Cretaceous extensional events (Figure 12). Such a mechanism was described onshore in the inverted Northeastern Pyrenean rifted margin (Clerc et al., 2015, 2016), where similar synrift basins developed over greatly thinned continental crust with basin floors representing locally exhumed continental lower crust. In such drastic thinning environments, Clerc et al. (2016) show that both prerift and synrift sediments can be affected by synmetamorphic ductile deformation and transported/delaminated together with the extended and highly thinned continental crust. This field analogue could most likely apply to the Træna Basin, where the potential field modeling also supports a drastic thinning of the crust just underneath the Cretaceous sediments (Figures 4b and 12). The thinned continental crust in the Træna Basin predominantly consists of ultrahigh dense material of possibly Caledonian and/or older origin. This crustal body extends and thickens northwestward in the prolongation of the seismic and structural grain observed in the deep basement part of the Utgard High. We propose that the upper-middle crust with prerift and synrift sediments were deformed and largely squeezed and delaminated in the central part of the Træna Basin, during the exhumation of an old lower crust which are almost in contact with the synsag sediments. Due to the great depth of the sediments observed in the Træna Basin, metamorphism and associated ductile deformation is a reasonable hypothesis to propose. It may also explain the difficulty to image any consistent seismic packages underneath the BCU. After Late Jurassic-Early Cretaceous rifting, another active phase of crustal thinning renewed in the Træna Basin and adjacent areas during the mid-Cenomanian-Santonian. This could explain the sudden increase of the subsidence rate suggested by Figure 13.

5.2.2. Deformation Patterns in the Nyk-Någrind Segment and the Hel Graben

Thanks to the new TGS data, we see clear evidence for a separate Neocomian (Barremian-Aptian?) rifting between the Någrind Syncline and the Nyk High (Figures 7 and 11a). This event succeeds the Jurassic-earliest Cretaceous extensional phase particularly well recognized and constrained in the platform area (Blystad et al., 1995; Brekke, 2000). Up to the mid-Cenomanian, postrift mainly concentrated within the basin low created during the Late Jurassic-Neocomian (Barremian-Aptian?) extension. From the mid-Cenomanian to Campanian, both increasing subsidence and deposition shifted southeastward (Figures 11b and 11c). The depocenters appear to be partly controlled by a west dipping detachment fault along the adjacent flank of the Utgard High (Figures 4a and 4b). This observation is partly in agreement with Skogseid et al. (1992), who speculated earlier on a similar development of the Utgard High and its surrounding basins. The middle crust and possible pre-Cretaceous prerift/synrift sediments in the adjunction to the Utgard High are drastically stretched along the detachment, with the total crustal thickness of 5–6 km (Figure 4b) constrained by our 2-D potential field modeling and also in agreement with Maystrenko et al. (2017).

Previous studies (Brekke, 2000; Lundin et al., 2013; Lundin & Doré, 2011) suggest that the folded geometry of the Någrind Syncline is not a result of a thermal subsidence following the Jurassic-Early Cretaceous rifting phase (Doré et al., 1999) but the consequence of a large-scale lithospheric folding caused by a modest Maastrichtian regional compression (Bjørnseth et al., 1997; Lundin et al., 2013). Parallel-bedded strata with relatively uniform thickness below the syncline could support this interpretation (Bjørnseth et al., 1997). However, we argue that after initial subsidence caused by a series of Late Jurassic to Neocomian extensional events, the Någrind Syncline had a “steer’s head” basin geometry with slightly increased sedimentation in its central part and sediment thinning toward the adjacent Nyk and Utgard Highs (Figure 11). Such a geometry is typical for basins that experienced thermal subsidence after a stretching episode (White & McKenzie, 1988). Subsequent reshaping of the Någrind Syncline occurred in mid-Campanian-Maastrichtian time, when the rift axis jumped toward the Hel Graben (Figures 11e and 11f). During this renewed activity, the northwest flank of the Nyk High started to rise and fault as a rift shoulder. The uplift of the Nyk High (Figures 4a, 5, 11e, 11f, and 12) generated a concomitant tectonic flexure and rapid subsidence in the Någrind Syncline leading to the formation of a deep depocenter (Figure 11e). We, therefore, prefer to attribute the present-day geometry

of the Någrind Syncline to extensional processes rather than a compressional buckling as proposed by Lundin et al. (2013).

Furthermore, in the outer Vøring Basin clear extension dominated in the North and South Gjallar Ridges since the mid-Campanian (Gernigon et al., 2003; Ren et al., 2003). This late rifting event is associated with the formation of clear and prominent synrift wedges (Spingar Formation) and normal faults partly cut by the Base Tertiary unconformity (Gernigon et al., 2004). We also precise that the presence of a shallowing Turonian (Lysing Formation) across the North Gjallar Ridge and the Fenris Graben (Osmundsen et al., 2002) is not any longer valid since well 6704/12-1 confirms that these synrift sediments are dominantly Campanian-Paleocene in age. In contrast to Lundin et al. (2013), we also believe that the main extension phase in the Gjallar Ridge initiated in the mid-Campanian and not Paleocene, where most of the deformation already migrated toward the volcanic province. In the adjacent Vigrid Syncline, as suggested by Kjennerud and Vergara (2005), an increased sedimentation in the mid-Campanian-Maastrichtian could also be a response to the onset of this rifting phase in the outer Vøring Basin.

Our study shows that the subbasaltic Grimm High resembles the North Gjallar Ridge. In addition to the faulting shape of the T-Reflection, we also define about 8–10 km of possible pre-Cenozoic rocks capping the reflection (Figure 10), although poorer seismic imaging below volcanics does not allow a more confident interpretation. Similar structural styles (i.e., tilted fault blocks within Cretaceous strata) and nearly synchronous vertical movements within the Grimm High area and the North Gjallar Ridge during Turonian to Maastrichtian times (Figures 3, 11d, and 11e) suggest that they might be parts of the common structural domain at this time.

The nature of the sudden Paleocene collapse of the Hel Graben is still debatable (Lundin et al., 2013). It is clearly associated with the Nyk High hanging wall slipping toward the graben center, but the crustal mechanism responsible for the structure formation remains unclear. Lundin et al. (2002) proposed that the Hel Graben was formed by a Paleocene caldera eruption possibly coeval with igneous activity associated with the extensive intrusion of high-velocity sills, which thus might have taken place prior to breakup time (Berndt et al., 2000). Evidence for older intra-Paleocene sill intrusions is observed in the Hel Graben (Figure 9b) and could support this interpretation, but the size and origin of such a putative caldera remain difficult to explain. Another model (Gernigon et al., 2003) assumed that breakup-related magmatism and thinning could provoke lateral flow of a ductile lower crust below the rift zone. This model could similarly explain both Paleocene uplift of the Nyk High and the collapse of the Hel Graben. Later, Lundin et al. (2013) proposed several other possibilities including similar crustal delamination and magma withdrawal during chamber deflation (e.g., Pinel & Jaupart, 2005).

Our seismic interpretation and modeling results consider a highly thinned crust (<5 km) below the Cretaceous-Paleocene Hel Graben (Figures 4b and 4c). We can only restore and constrain evidence of Late Cretaceous to Late Paleocene rifting/faulting in the Hel Graben and surrounding structures. However, it is possible that the deepest and older Cretaceous sediments rest directly upon the LCB (Figures 4b and 4c), while middle-upper crust and pre-Cretaceous prerift/synrift deposits (if they still exist) were totally delaminated and/or boudined in the similar mode previously described in the Traena Basin (Figure 12). Such crustal removal could be largely controlled by a detachment faulting system observed along the western flank of the Nyk High and possibly on the eastern flank of the Grimm High/Hel Terrace lying on the conjugate side of the expected Paleocene rift/thinning axis (Figures 4b and 4c). Together with the breakup-related magmatism, this thinning event could have promoted a collapse of the Hel Graben and uplift of the Nyk High as proposed in the model of Gernigon et al. (2003). Depositional patterns and the rapid Paleocene subsidence rates may indicate that the main phase of thinning took place during the Paleocene rifting phase restricted to the Hel Graben and possibly underneath the basalt in the central outer part of the Vøring Margin (e.g., the Fenris Graben). Poorer seismic imaging at greater depth in the area makes hard to estimate the amount of crustal thinning in the earlier rifting phases. The possible presence of pre-Cretaceous sediments at the Grimm and Nyk highs (Figures 4a and 4b) allows us to speculate that the pre-Cretaceous basin on top of the thinned crust could also coexist within the Hel Graben, although appropriate sediments could be highly metamorphosed and delaminated at least during Late Cretaceous-Paleocene rifting and likely during earlier extensional phases.

Compared to the Hel Graben, the crustal blocks preserved on the flanks (e.g., the Grimm High and Nyk High) show evidence of thicker continental crust (Figure 4b), less affected by the Paleocene thinning compared to the adjacent Hel Graben. We interpret this specific crustal configuration in terms of boudinage of the crust (Figure 12) (see Gartrell, 1997) which appears to be a key process in many other rifted margins worldwide (Clerc et al., 2017). The Nyk and Grimm Highs most likely represent thick crustal swell or boudin in between the subbasin (Hel Graben), where drastic thinning deformations were localized (Figure 12).

It is also important to note that the Hel Graben experienced a different tectonic history compared to the other segments of the outer Vøring Basin. Increasing faulting events and drastic Paleocene sedimentation in the Hel Graben (Figure 11f) are synchronous to uplift and progressive decrease of the normal faulting observed farther south in the central segment of the outer Vøring Basin (e.g., North and South Gjallar Ridges). We propose that this difference is possibly linked with the localization and migration of the deformation during the Paleocene. The crustal thinning estimation suggests that in the Paleocene, the Hel Graben was almost close to the breakup stage if we assume a crustal thickness of less than 4 km (Figures 4b and 4c). However, the breakup finally occurred 65 km farther west. It suggests that the Paleocene rift axis that almost dissected the Hel Graben failed to propagate farther south and jumped west of the Grimm High during the earliest Eocene breakup. The final jump is possibly explained by the onset of magmatism (Figure 12), likely leading to extra weakening and rift localization in the western branch of this complex overlapping system. Similar processes of rift migration/abortion triggered by magmatism and underplating are described in Yamasaki and Gernigon (2010).

5.3. Regional and Generic Implications

5.3.1. Early (Neocomian) and Mid-Cretaceous Extension in Its Regional Context

We may identify evidence for significant Cretaceous tectonic activity in the northern Vøring Margin. While Late Jurassic-Early Cretaceous and Late Cretaceous-Paleocene extensional phases are widely accepted and confirmed by structural observations and drilling, Neocomian and mid-Cretaceous (Aptian to nearly Campanian) rifting events are often neglected, rather ambiguous, and often interpreted as a passive sedimentary infill during postrift thermal subsidence (Færseth, 2012; Færseth & Lien, 2002; Skogseid et al., 2000). Recent drilling results from the Vøring Basin; for example, well 6603/5-1 at the South Gjallar Ridge and well 6608/2-15 at the Utgard High (NPD factpages, 2017) proved much thicker Lower and Upper Cretaceous sediments than previously believed (Blystad et al., 1995; Skogseid et al., 2000; Swiecicki et al., 1998). Such misinterpretation in the proposed stratigraphy is primarily the results of previous underestimation of the sedimentation rates throughout the Cretaceous period and consequently may explain the minimization of the mid-Cretaceous extension (Færseth & Lien, 2002). However, previous evidence for prominent early-to mid-Cretaceous extensional tectonism has been reported in the Rån and Gjallar Ridges in the outer Vøring Basin (Gernigon et al., 2003), the Lofoten-Vesterålen Margin (Henstra et al., 2016; Tsikalas et al., 2001, 2005), the Någrind Syncline, and the Træna Basin (Figures 7 and 12; this study). Aptian-Cenomanian extension at the southern end of the Nordland Ridge was also suggested by Pascoe et al. (1999). It is not impossible that this extensional deformation may have propagated north, along the flank of the Nordland Ridge, triggering mid-Cenomanian-Santonian extensional reactivation in the Træna Basin. Activation of the Nordland Ridge as a sand source in the Cenomanian and Turonian-Coniacian (Lysing Formation) was also suggested by Doré et al. (1999) and Fjellanger et al. (2005). Furthermore, coeval tectonic reactivation was observed along the onshore continuation of the Møre-Trøndelag Fault Complex (Sømme & Jackson, 2013). Nearly coeval apatite fission track ages (since 100 Ma) of fault reactivation and block uplift of kilometers scale within the Møre-Trøndelag Fault Complex have been reported from the Norwegian mainland (Redfield et al., 2005). Beyond that, in the wider North Atlantic realm, we have well-documented evidence of Early to mid-Cretaceous extension, including Aptian rifting in the SW Barents Sea (Faleide et al., 1993; Serck et al., 2017), mid-Albian faulting activity in the NE Greenland (Whitham et al., 1999), and Berriasian to Cenomanian synrift deposition in the Rockall Trough (Corfield et al., 1999). The prominent Late Mesozoic rifting phases in the Faeroe-Shetland Basin likely occurred in the Neocomian, Aptio-Albian, Cenomanian-Santonian, and Campanian-Maastrichtian (Dean et al., 1999), whereas minor Turonian-Coniacian fault reactivation in the region promoted coeval sand deposition identical to the Lysing Formation in the Norwegian Sea (Grant et al., 1999). All above mentioned structural observations make it likely that the entire mid-Norwegian Margin and the northern Vøring Margin segment, in particular, were subjected to similar early to mid-Cretaceous events in the regional context contrary to the quiescence

interpretation of Færseth and Lien (2002). These events were possibly not associated with a prominent and continuous rifting axis in the mid-Norwegian Margin but represented scattered, isolated, and migrating rifting centers that were selectively reactivated along basement highs and old mobile zones (Lofoten Ridge, Nordland Ridge, Nyk High, Surt Lineament, Jan Mayen Corridor, etc.).

5.3.2. On the Role of Crustal Inheritance in the Margin Development and the Presence of Serpentinized Mantle in the Vøring Basin

As mentioned above, recent tectonic models suggested the presence of a partially to heavily serpentinized mantle in the Vøring Basin (Péron-Pinvidic et al., 2013; Péron-Pinvidic & Osmundsen, 2016; Reynisson et al., 2010; Rüpké et al., 2013). Extremely thinned crust in the Hel Graben and the Træna Basin (Figure 4b) could favor such tectonic scenario there. Pérez-Gussinyé and Reston (2001) suggested that serpentinization can be likely achieved in the settings with thinning (beta) factors above 3 and a crustal thickness less than 8 km. However, in addition, we have to consider the timing and duration of the main thinning phases, decompacted sediment thicknesses during these phases, and the development of basin-scale faults in order to understand environments that could promote or block water passage to the mantle and lead to a partial serpentinization. All these parameters should be then estimated, integrated, and tested within different modeling approaches to allow testing of various hypotheses. However, our tectonic model also supports the early presence of inherited continental crust and related LCB over most of the highs observed in the Vøring Basin (Figure 12). It is unlikely that a broad zone of exhumed mantle affected the entire Vøring Basin, and the presence of exhumed mantle underneath the basalt cannot explain the shallow environments suggested by new seismic interpretation and petrology derived from the ODP drilling in the Vøring Marginal High (Abdelmalak, Meyer, et al., 2016; Abdelmalak, Planke, et al., 2016). If serpentinization occurs, it would eventually be restricted to local windows in the deepest part of the subsag basins involved in this boudined and mullioned type of superextension (Figure 12). Other observations from worldwide rifted margins (Clerc et al., 2017) and also various numerical models also support the presence and persistence of inherited crust underneath similarly large sag basins (Brune et al., 2014; Petersen & Schiffer, 2016).

The presence of older crustal grains and crustal weakness zones has a significant influence on basin development and architecture. While analogue and numerical models deal mostly with homogenous and uniform mode of extension in the rifted margins (Lavie & Manatschal, 2006), crustal inheritance and late magmatism could explain complex geometries and deformation migration and localization discussed in this paper. The inherited crustal blocks and associated structures/fabrics initiate and control the onset and development of thinning deformations of the continental crust (Abdelmalak et al., 2017). This “boudinage style of deformation” is similar to what has been previously described and proposed by Lister et al. (1986), Gartrell (1997), or Clerc et al. (2017). For the Northern Vøring and Lofoten-Vesterålen Margins, such a tectonic scenario was proposed almost 30 years ago by Anderson (1986) who, based on gravity anomalies analysis, suggested the existence of large-scale boudinage of the continental crust which makes up the margin. Furthermore, these weakness zones could be associated with major lithosphere structural discontinuities (e.g., sutures and mega-shear zones). Their orientation and deformational pattern could give us better constraints in paleoreconstructions and better understand their role in the margin segmentation. Similar tectonic regimes occurred in the development of other rifted margins (Chian et al., 2001; Clerc et al., 2017; Franke et al., 2014; Funck et al., 2008; Gernigon et al., 2014; Sun et al., 2009). Therefore, we caution researchers to apply “classical” hyperextended Iberian scenarios in complex tectonic settings such as the mid-Norwegian Margin. In the mid-Norwegian Margin larger and thicker continental crustal masses with rather complex petrophysical properties and spatial distribution are involved. Its rift development is also polyphase and shows a much longer period of episodic rifting event compared to the Iberian rifted margin types.

6. Conclusions

1. We investigated and interpreted new seismic data in the northern Vøring Margin poorly constrained by deep seismic data in the past. Combined with regional map production and potential field modeling, we propose a new tectonic scenario for the mid-Norwegian Margin.
2. We argue that the northern Vøring Margin and the entire mid-Norwegian Margin were tectonically active during mid-Cretaceous in contrast to Færseth and Lien (2002). We found seismic evidence of Barremian-Aptian (?) faulting activity in the Någrind Syncline and mid-Cenomanian-Santonian extensional

reactivation in the Træna Basin. This may indicate a nonuniform and nonlinear distribution of rifting centers in the Early to mid-Cretaceous within the Vøring Margin likely reflecting selective tectonic reactivations along preexisting mobile zones.

3. The Træna Basin is a narrow basin with a thin crust (4–5 km) which initiated since the Late Jurassic as a half-graben developed on the flank of the Utgard High. Possible later mid-Cenomanian-Santonian extensional reactivation significantly increased the basin subsidence and the thinning of the underlying continental crust. Presynrift and partly synrift sediments together with upper-middle crustal rocks at the basin floor might have been significantly deformed and delaminated/compacted during these extensional phases
4. According to our observations and modeling results, the crust underneath the Utgard High likely represents ultrahigh-density body ($3,200 \text{ kg/m}^3$) of Caledonian or older (Precambrian) origin. The top of this body corresponds to the strong *UH*-Reflection. Detachment faults developed on the flanks of this rigid continental block and controlled the prethinning and synthinning Mesozoic evolution of the adjacent basins (e.g., the Nâgrind Syncline and the Træna Basin).
5. We see no prerequisites to explain the configuration of the Nâgrind Syncline as a result of compressional buckling as previously suggested by Lundin et al. (2013). Our analysis shows that the Nâgrind Syncline has been formed by a migrating sequence of extensional and postextensional events and associated vertical movements from the Early Cretaceous to the Early Eocene.
6. The Hel Graben is a deep Cretaceous-Paleocene basin that developed in between two rigid block ("continental buffers"). The thinning deformation in the Hel Graben was controlled by detachments on the flanks of these blocks (the Grimm High and the Nyk High) with a culmination in the Paleocene that led to a collapse of the structure. We suggest very thin upper and middle crustal rocks and prerift and synrift sediments in the axial part of the Hel Graben. They are most likely highly deformed and metamorphosed along detachments. Extremely thinned crust (3–4 km) indicates that the Hel Graben was very close to the breakup stage in the Paleocene until the rifting axes eventually jumped in the late Paleocene-earliest Eocene toward the present continent-ocean "boundary". We also found evidence of intra-Paleocene intrusive magmatism preceding the main magmatic event during the earliest Eocene continental breakup.
7. The *T*-Reflection observed underneath the North Gjallar Ridge extends toward the Grimm High and might be of the same origin, that is, old inherited Caledonian lower crust. These structures could be part of a common structural trend before the Paleocene extensional phase.
8. We do not expect wide zones of mantle exhumation and serpentinization in the Vøring Basin; however, local areas of serpentinization could eventually develop in deep subbasins with significantly thinned crust (Træna Basin and Hel Graben). Nonetheless, in our opinion, the mid-Norwegian Margin represents a unique type of a volcanic rifted margin, where large inherited crustal blocks define a long and polyphase development and a complex deformation distribution. Combined with the presence of SDRs along the continent-ocean transition, it has very little in common with the Iberian nonvolcanic rifted margin types.

Appendix A: Stratigraphic Comment on Well 6608/2-1S, Utgard High

This appendix provides a detailed stratigraphic comment on the selected Cretaceous intervals and log pattern interpretation of well 6608/2-1S. The well 6608/2-1S has been analyzed by RPS Ichron. The raw results of biostratigraphic analyses and wireline logs have been provided by the Norwegian Petroleum Directorate (NPD) via the Diskos National Data Repository. After review and integration, these data formed the basis of stratigraphic interpretation.

A1. Early Cretaceous Intervals

The Early Cretaceous interval is dated in both the main hole and ST2 bypass where the sidewall core samples have been analyzed providing stronger age confidence (Figure A1).

The top of the Langebarn Formation is defined with the use of the change in the log pattern and is supported with the presence of *Caudammina ovula crassa* foraminifera (Figure A1). The Albian interval of the formation is characterized with the sparse palynological assemblage, consistent with the interpretation, with rare presence of *Batioladinium micropodum*, *Muderongia asymmetrica*, and taxa of a broad Early Cretaceous range. The presence of *Aptea polymorpha* is considered to be characteristic of the Albian period in the northern latitudes (Nøhr-Hansen, 1993).

Figure A1. Well 6608/2-1S and ST2 bypass. Biostratigraphy summary chart of Barremian to Coniacian intervals. Formation names after NORLEX (Gradstein et al., 2010).

The Late Aptian section of the well is characterized by the prominent gamma log pattern (Figure A1), suggesting the paleoenvironments with the decreased level of oxygen typical for this age. The presence of *Cerbia tabulata* and *Aptea eisenackii* and increase in *A. polymorpha* confirm the dating for the interval (Costa & Davey, 1992). The presence of *Vesperopsis mayi* is typical for the northern latitudes (Nøhr-Hansen, 1993) and tentatively suggest stressed environment (Harding & Allen, 1995).

The Early Aptian is confirmed with the presence of the red stained foraminifera and LAD of *Ctenidodinium elegantulum*, *Sirmiodinium grossii*, and *Pseudoceratium pelliferum* (Costa & Davey, 1992). The pronounce spike increase in gamma (Figure A1), together with abundant influx of bisaccate pollen grains and sapropelic organic in core, confirms the presence of OAE1 sediments, representing Selli event (Ainsworth et al., 2000). This interval is an equivalent to Fischschiefer bed in North Sea. The presence of *Subtilisphaera perlucida* is additional evidence of anoxic environment and confirms the age interpretation.

The presence of Barremian sediments within the section is supported with the downward change of the log pattern (Figure A1) and is confirmed with the LAD of *Aptea anphrissa* and *Nelchinopsis kostromiensis*, together with the base of abundant presence of *P. cretaceum* and overall assemblage change (Costa & Davey, 1992).

A2. Coniacian-Albian Interval

The Lysing Formation, typically representing Turonian-Late Coniacian strata, is not prominently expressed in the log profile and has not been analyzed for biostratigraphy (Figure A1). The underlying interval is characterized by the rich assemblage indicative of Turonian age, present in two samples (Figure A1). The cooccurrence of acme event of *Palaeohystrichophora infusorioides* together with abundance of

Cauveridinium membraniphorum is indicative of latest Turonian (Costa & Davey, 1992; Olde et al., 2014). The cooccurring *Stephodium coronatum* and *Apteodinium maculatum* serve as important markers as their LAD are observed at top Turonian boundary (Costa & Davey, 1992), specifically for North Træna and Møre Basins (Radmacher et al., 2015; Smelror et al., 1994). The cooccurring *Heterosphaeridium difficile* is known to have its FAD within the Turonian (Costa & Davey, 1992), specifically in the North Træna Basin (Radmacher et al., 2015). The influx of *Surculosphaeridium longifurcatum*, observed in the sample below, is known to be characteristic of mid-Turonian strata (Costa & Davey, 1992; Radmacher et al., 2015).

Due to its lower thickness in comparison to the known sections of the basin, the Turonian interval of the well is considered to be either condensed or possessing a stratigraphic unconformity. Since the studied assemblages possess strong Middle-Late Turonian marker events and lack any Early Turonian events, this negative evidence is tentatively suggesting the presence of unconformity with the Early and partly Middle Turonian intervals missing from the section.

The underlying part of the Blållange Formation is characterized by extensively expanded section with its age defined in two samples from the upper part of the studied interval (Figure A1). The assemblage in these samples is represented by the abundant presence of *Subtilisphaera kalaatii* and other peredinioid cysts such as *Palaeoperidinium cretaceum* and *Palaeohystrichophora infusorioides*, together with increase of common *Chlamydomphorella nyei* and single occurrences of *Rhombodella paucispina* and *Ovoidinium scabrosum*. These events suggest the correlation to the *Palaeohystrichophora infusorioides*-*Palaeohystrichophora palaeoinfusa* Interval Zone in the sense of Radmacher et al. (2014) and are indicative of Middle-Late Cenomanian.

The presence of the assemblages dominated with peridinioid cysts, specifically *Palaeoperidinium cretaceum* and *Subtilisphaera kalaatii*, in Cenomanian strata is characteristic for North Træna Basin (Radmacher et al., 2015), southwestern Barents Sea (Radmacher et al., 2014), and East Greenland (Nøhr-Hansen, 1993, 2012).

Appendix B: A Comment on the Petrophysical Properties Used in the 2-D Potential Field Modeling of the Northern Vøring Transect

All potential field studies combine a priori constraints such as physical property data, geological mapping, or seismic interpretations to limit the infinite number of possible theoretical solutions (Saltus & Blakely, 2011). Some of the modeling assumptions derived mostly (1) from local onshore-offshore correlations of some gravity and magnetic anomalies, (2) the nature of the rocks expected, (3) the correlation with V_p values from refraction/OBS data, and (4) the conclusion of the potential field modeling itself.

Rock property data from onshore (see NGU database) (e.g., Olesen et al., 2010) and local offshore density logs also provide local constraints, but mostly limited to the upper part of the sedimentary basin and/or the proximal part of the margin. We assume that the density in the deepest part of the basin does not exceed an average value of $2,750 \text{ kg/m}^3$, which also fits conventional V_p /density function. This threshold value was also voluntarily chosen to reach the maximum of crustal thinning we could expect along the transect.

Sedimentary rocks are also almost nonmagnetic (e.g., Olesen et al., 2010), and we assumed, accordingly, that most of the magnetic sources source from the crystalline basement or the basalt in the outer part of the margin.

Review of the onshore measurements (see Olesen et al., 2010) shows a large variation of the rock properties. However, all rocks from the upper part of the crust (e.g., the Caledonian nappes) have, in average, lower density and lower magnetic susceptibilities ($<0.01\text{--}0.02 \text{ SI}$) compared to the pre-Cambrian crust showing higher density and much higher susceptibilities ($>0.03 \text{ SI}$). Previous modelings have shown that a simple crustal model with an upper crust dominated by rocks equivalent of the Caledonian nappes and a magnetic middle crust associated with pre-Cambrian types of rocks could easily explain the observed gravity and magnetic signature of the mid-Norwegian Margin (e.g., Ebbing et al., 2006; Gernigon et al., 2015; Maystrenko et al., 2017; Olesen et al., 2010; Reynisson et al., 2010). The lower to the upper crust with V_p velocities higher than 7 km/s are most likely characterized by high-density material as confirmed by our modeling (e.g., the prominent gravity anomaly of the Utgard High). In general, the crustal densities are also realistic and remain in the range of average crustal values measured at specific depth on the Earth (Christensen & Mooney, 1995; Herzberg et al., 1983).

In addition to the forward modeling, we also consider Werner independent deconvolution techniques to estimate the potential field sources directly from measured anomalies without appeal to specific assumptions about the source distributions. When finally combined with seismic data, the final potential field model fits all observations and geophysical data available (e.g., seismic, gravity, and magnetic). Therefore, if a model produces results that agree with data, we will regard it as successful until new data and understanding allow us to modify or even reject modeling results. The parameters in Table 1 just show the best parameters required to fit all the data and observations. Without extra constrain, a sensitivity analysis will only result in a minor variation of the relative crustal properties involved.

Acknowledgments

Funding for this work came from OMNIS Project: Offshore Mid-Norwegian: Integrated Margin and Basin Studies (project: 210429/E30) funded by the Norwegian Research Council through its Centre of Excellence CEED (Centre for Earth Evolution and Dynamics). The seismic, magnetic, and gravity data presented in this study were provided by TGS. RPS Ichron is acknowledged for performing biostratigraphic analysis of well 6608/2-15. Seismic interpretation was done using IHS Kingdom software. Grid interpolations and map compilations were established using Geosoft Oasis Montaj and ArcGIS softwares. We would like to acknowledge First Geo AS and TGS for the use of hiQbe™ regional velocity cube employed in depth conversion. We thank GeoModelling Solutions for providing an academic license for TecMod 2-D software. The seismic and well data used in this study are available through the Diskos National Data Repository (NDR). Well formation tops are also available at the NPD factpages (www.npd.no) and the NORLEX team website (www.nhm2.uio.no/norlex/). We are grateful to Jolante van Wijk (New Mexico Tech), Daniel Schmid (PGP, University of Oslo), and Sergei Medvedev (CEED, University of Oslo) for their help and fruitful discussions that improved the paper, and we thank Ben Manton (VBPR) for the linguistic remarks. We also would like to thank Christopher Jackson, Philip Ball, and the Editor Nathan Niemi for their constructive and helpful reviews.

References

- Abdelmalak, M. M., Andersen, T. B., Planke, S., Faleide, J. I., Corfu, F., Tegner, C., ... Myklebust, R. (2015). The ocean-continent transition in the mid-Norwegian margin: Insight from seismic data and an onshore Caledonian field analogue. *Geology*, 43(11), 1011–1014. <https://doi.org/10.1130/G37086.1>
- Abdelmalak, M. M., Faleide, J. I., Planke, S., Gernigon, L., Zastrozhnov, D., Shephard, G. E., & Myklebust, R. (2017). The T-Reflection and the deep crustal structure of the Vøring Margin offshore mid-Norway. *Tectonics*, 36, 2497–2523. <https://doi.org/10.1002/2017TC004617>
- Abdelmalak, M. M., Meyer, R., Planke, S., Faleide, J. I., Gernigon, L., Frieling, J., ... Myklebust, R. (2016). Pre-breakup magmatism on the Vøring Margin: Insight from new sub-basalt imaging and results from Ocean Drilling Program Hole 642E. *Tectonophysics*, 675, 258–274. <https://doi.org/10.1016/j.tecto.2016.02.037>
- Abdelmalak, M. M., Planke, S., Faleide, J. I., Jerram, D. A., Zastrozhnov, D., Eide, S., & Myklebust, R. (2016). The development of volcanic sequences at rifted margins: New insights from the structure and morphology of the Vøring Escarpment, mid-Norwegian Margin. *Journal of Geophysical Research: Solid Earth*, 121, 5212–5236. <https://doi.org/10.1002/2015JB012788>
- Ainsworth, N. R., Riley, L. A., & Gallagher, L. T. (2000). An Early Cretaceous lithostratigraphic and biostratigraphic framework for the Britannia Field reservoir (Late Barremian–Late Aptian), UK North Sea. *Petroleum Geoscience*, 6(4), 345–367. <https://doi.org/10.1144/petgeo.6.4.345>
- Allen, P. A., & Allen, J. R. (2013). *Basin analysis: Principles and application to petroleum play assessment* (p. 642). Chichester: John Wiley.
- Anderson, J. A. (1986). Arctic geodynamics: The opening of the Arctic Ocean. *Advances in Space Research*, 6(9), 79–83. [https://doi.org/10.1016/0273-1177\(86\)90356-X](https://doi.org/10.1016/0273-1177(86)90356-X)
- Bergh, S. G., Eig, K., Klovjan, O. S., Henningsen, T., Olesen, O., & Hansen, J. A. (2007). The Lofoten-Vesterålen continental margin: A multiphase Mesozoic–Palaeogene rifted shelf as shown by offshore-onshore brittle fault-fracture analysis. *Norwegian Journal of Geology*, 87, 29–58.
- Berndt, C., Skogly, O. P., Planke, S., Eldholm, O., & Mjelde, R. (2000). High-velocity breakup-related sills in the Vøring Basin, off Norway. *Journal of Geophysical Research*, 105, 28, 443–28, 454. <https://doi.org/10.1029/2000JB900217>
- Bjørnseth, H. M., Grant, S. M., Hansen, E. K., Hossack, J. R., Roberts, D. G., & Thompson, M. (1997). Structural evolution of the Vøring Basin, Norway, during the Late Cretaceous and Paleogene. *Journal of the Geological Society (London)*, 154(3), 559–563. <https://doi.org/10.1144/gsjgs.154.3.0559>
- Blystad, P., Brekke, H., Færseth, R. B., Larsen, B. T., Skogseid, J., & Tørudbakken, B. (1995). Structural elements of the Norwegian continental shelf. Part II: The Norwegian Sea region, *NPD-Bulletin*, The Norwegian Petroleum Directorate, 8.
- Brekke, H. (2000). The tectonic evolution of the Norwegian Sea continental margin with emphasis on the Vøring and Møre Basins. *Geological Society, London, Special Publications*, 167(1), 327–378. <https://doi.org/10.1144/gsl.sp.2000.167.01.13>
- Brekke, H., Dahlgren, S., Nyland, B., & Magnus, C. (1999). The prospectivity of the Vøring and Møre Basins on the Norwegian Sea continental margin. *Geological Society, London, Petroleum Geology Conference series*, 5, 261–274. <https://doi.org/10.1144/0050261>
- Brekke, H., & Riis, F. (1987). Tectonics and basin evolution of the Norwegian shelf between 62°N and 72°N. *Norsk Geologisk Tidsskrift*, 67, 295–322.
- Brune, S., Heine, C., Perez-Gussinye, M., & Sobolev, S. V. (2014). Rift migration explains continental margin asymmetry and crustal hyper-extension. *Nature Communications*, 5. <https://doi.org/10.1038/ncomms5014>
- Brune, S., Popov, A. A., & Sobolev, S. V. (2012). Modeling suggests that oblique extension facilitates rifting and continental break-up. *Journal of Geophysical Research*, 117, B08402. <https://doi.org/10.1029/2011JB008860>
- Brune, S., Williams, S. E., Butterworth, N. P., & Müller, R. D. (2016). Abrupt plate accelerations shape rifted continental margins. *Nature*, 536(7615), 201–204. <https://doi.org/10.1038/nature18319>
- Buck, W. R. (2006). The role of magma in the development of the Afro-Arabian rift system. In G. Yirgu, C. J. Ebinger, & P. K. H. Maguire (Eds.), *The Afar Volcanic Province within the East African Rift System*, Geological Society, London, Special Publications (Vol. 259, pp. 43–54). <https://doi.org/10.1144/GSL.SP.2006.259.01.05>
- Chian, D., Reid, I. D., & Jackson, H. R. (2001). Crustal structure beneath Orphan Basin and implications for nonvolcanic continental rifting. *Journal of Geophysical Research*, 106(B6), 10,923–10,940. <https://doi.org/10.1029/2000JB900422>
- Christensen, N. I., & Mooney, W. D. (1995). Seismic velocity structure and composition of the continental crust: A global view. *Journal of Geophysical Research*, 100(B6), 9761–9788. <https://doi.org/10.1029/95JB00259>
- Christie-Blick, N., & Biddle, K. T. (1985). Deformation and basin formation along strike-slip faults. *SEPM Special Publication*, 37, 1–34. <https://doi.org/10.2110/pec.85.37.0001>
- Clark, S. A., Glørstad-Clark, E., Faleide, J. I., Schmid, D., Hartz, E. H., & Fjeldskaar, W. (2014). Southwest Barents Sea rift basin evolution: Comparing results from backstripping and time-forward modelling. *Basin Research*, 26(4), 550–566. <https://doi.org/10.1111/bre.12039>
- Clerc, C., Lagabrielle, Y., Labaume, P., Ringenbach, J.-C., Vauchez, A., Nalpas, T., ... Fourcade, S. (2016). Basement–cover decoupling and progressive exhumation of metamorphic sediments at hot rifted margin. Insights from the Northeastern Pyrenean analog. *Tectonophysics*, 686, 82–97. <https://doi.org/10.1016/j.tecto.2016.07.022>
- Clerc, C., Lahfid, A., Monié, P., Lagabrielle, Y., Chopin, C., Poujol, M., ... de St Blanquat, M. (2015). High-temperature metamorphism during extreme thinning of the continental crust: Areappraisal of the north Pyrenean paleo-passive margin. *Solid Earth*, 6(2), 643–668. <https://doi.org/10.5194/se-6-643-2015>
- Clerc, C., Ringenbach, J.-C., Jolivet, L., & Ballard, J.-F. (2017). Rifted margins: Ductile deformation, boudinage, continentward-dipping normal faults and the role of the weak lower crust. *Gondwana Research*. <https://doi.org/10.1016/j.gr.2017.04.030>
- Cohen, K. M., Finney, S. C., Gibbard, P. L., & Fan, J.-X. (2013). The ICS international chronostratigraphic chart. *Episodes*, 36(3), 199–204.

- Corfield, S., Murphy, N., & Parker, S. (1999). The structural and stratigraphic framework of the Irish Rockall Trough. *Geological Society, London, Petroleum Geology Conference Series*, 5, 407–420. <https://doi.org/10.1144/0050407>
- Costa, L. I., & Davey, R. J. (1992). Dinoflagellate cysts of the Cretaceous system. In A. J. Powell (Ed.), *A stratigraphic index of dinoflagellate cysts* (pp. 99–131). London: Chapman & Hall. https://doi.org/10.1007/978-94-011-2386-0_3
- Dalland, A., Worsley, D., & Ofstad, K. (1988). A lithostratigraphic scheme for the Mesozoic and Cenozoic succession offshore mid- and northern Norway. *NPD Bulletin*, 4, 63.
- Dean, K., McLachlan, K., & Chambers, A. (1999). Rifting and the development of the Faeroe–Shetland Basin. *Geological Society, London, Petroleum Geology Conference Series*, 5, 533–544. <https://doi.org/10.1144/0050533>
- Digranes, P., Mjelde, R., Kodaira, S., Shimamura, H., Kanazawa, T., Shiobara, H., & Berg, E. W. (1998). A regional shear-wave velocity model in the central Vøring Basin, N. Norway, using three-component ocean bottom seismographs. *Tectonophysics*, 293(3–4), 157–174. [https://doi.org/10.1016/S0040-1951\(98\)00093-6](https://doi.org/10.1016/S0040-1951(98)00093-6)
- Doré, A. G., Lundin, E. R., Jensen, L. N., Birkland, Ø., Eliassen, P. E., & Fichler, C. (1999). Principal tectonic events in the evolution of the northwest European Atlantic margin. *Geological Society, London, Petroleum Geology Conference Series*, 5, 41–61. <https://doi.org/10.1144/0050041>
- Doré, A. G., Lundin, E. R., Kusznir, N. J., & Pascal, C. (2008). Potential mechanisms for the genesis of Cenozoic domal structures on the NE Atlantic margin: Pros and cons and some new ideas. In H. Johnson, et al. (Eds.), *The nature and origin of compression in passive margins*, *Geological Society, London, Special Publications* (Vol. 306, pp. 1–26). <https://doi.org/10.1144/SP306.1>
- Ebbing, J., Lundin, E., Olesen, O., & Hansen, E. K. (2006). The mid-Norwegian margin: A discussion of crustal lineaments, mafic intrusions, and remnants of the Caledonian root by 3D density modelling and structural interpretation. *Journal of the Geological Society, London*, 163(1), 47–59. <https://doi.org/10.1144/0016-764905-029>
- Ebinger, C., & Casey, M. (2001). Continental breakup in magmatic provinces: An Ethiopian example. *Geology*, 29(6), 527–530. [https://doi.org/10.1130/0091-7613\(2001\)029<0527:CBIMPA>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<0527:CBIMPA>2.0.CO;2)
- Eldholm, O., Thiede, J., & Taylor, E. (1989). Evolution of the Vøring volcanic margin. In O. Eldholm, J. Thiede, & E. Taylor (Eds.), *Proceedings of the Ocean Drilling Program, scientific results* (pp. 1033–1065). College Station, TX: Ocean Drilling Program.
- Eldholm, O., Tsikalas, F., & Faleide, J. I. (2002). Continental margin off Norway 62–75°N: Palaeogene tectono-magmatic segmentation and sedimentation. *Geological Society of London, Special Publication*, 197(1), 39–68. <https://doi.org/10.1144/gsl.sp.2002.197.01.03>
- Færseth, R. B. (2012). Structural development of the continental shelf offshore Lofoten–Vesterålen, northern Norway. *Norwegian Journal of Geology*, 92, 19–40.
- Færseth, R. B., & Lien, T. (2002). Cretaceous evolution in the Norwegian Sea—A period characterized by tectonic quiescence. *Marine and Petroleum Geology*, 19(8), 1005–1027. [https://doi.org/10.1016/S0264-8172\(02\)00112-5](https://doi.org/10.1016/S0264-8172(02)00112-5)
- Faleide, J. I., Tsikalas, F., Breivik, A. J., Mjelde, R., Ritzmann, O., Engen, Ø., ... Eldholm, O. (2008). Structure and evolution of the continental margin off Norway and the Barents Sea. *Episodes*, 31(1), 82–91.
- Faleide, J. I., Vågnes, E., & Gudlaugsson, S. T. (1993). Late Mesozoic–Cenozoic evolution of the southwestern Barents Sea in a regional rift-shear tectonic setting. *Marine and Petroleum Geology*, 10(3), 186–214. [https://doi.org/10.1016/0264-8172\(93\)90104-Z](https://doi.org/10.1016/0264-8172(93)90104-Z)
- Fjellanger, E., Surlyk, F., Wamsteeker, L. C., & Midtun, T. (2005). Upper Cretaceous basin-floor fans in the Vøring Basin, mid-Norway shelf. In B. Wandås, et al. (Eds.), *Onshore-offshore relationships on the North Atlantic Margin, Norwegian Petroleum Society Special Publications* (Vol. 12, pp. 135–164). [https://doi.org/10.1016/S0928-8937\(05\)80047-5](https://doi.org/10.1016/S0928-8937(05)80047-5)
- Foulger, G. R. (2010). *Plates vs plumes: A geological controversy*. Oxford: Wiley-Blackwell. <https://doi.org/10.1002/9781444324860>
- Franke, D. (2013). Rifting, lithosphere breakup and volcanism: Comparison of magma-poor and volcanic rifted margins. *Marine and Petroleum Geology*, 43, 63–87. <https://doi.org/10.1016/j.marpetgeo.2012.11.003>
- Franke, D., Savva, D., Pubellier, M., Steuer, S., Mouly, B., Auzietre, J. L., ... Chamot-Rooke, N. (2014). The final rifting evolution in the South China Sea. *Marine and Petroleum Geology*, 58, 704–720. <https://doi.org/10.1016/j.marpetgeo.2013.11.020>
- Funck, T., Andersen, M. S., Neish, J. K., & Dahl-Jensen, T. (2008). A refraction seismic transect from the Faroe Islands to the Hatton-Rockall Basin. *Journal of Geophysical Research*, 113, B12405. <https://doi.org/10.1029/2008JB005675>
- Gartrell, A. P. (1997). Evolution of rift basins and low-angle detachments in multilayer analog models. *Geology*, 25(7), 615–618. [https://doi.org/10.1130/0091-7613\(1997\)025<0615:EORBAL>2.3.CO;2](https://doi.org/10.1130/0091-7613(1997)025<0615:EORBAL>2.3.CO;2)
- Geoffroy, L., Burov, E. B., & Werner, P. (2015). Volcanic passive margins: Another way to break up continents. *Scientific Reports*, 5. <https://doi.org/10.1038/srep14828>
- Gernigon, L., Blischke, A., Nasuti, A., & Sand, M. (2015). Conjugate volcanic rifted margins, seafloor spreading, and microcontinent: Insights from new high-resolution aeromagnetic surveys in the Norway Basin. *Tectonics*, 34, 907–933. <https://doi.org/10.1002/2014TC003717>
- Gernigon, L., Brönnner, M., Roberts, D., Olesen, O., Nasuti, A., & Yamasaki, T. (2014). Crustal and basin evolution of the southwestern Barents Sea: From Caledonian orogeny to continental breakup. *Tectonics*, 33, 347–373. <https://doi.org/10.1002/2013TC003439>
- Gernigon, L., Ringenbach, J.-C., Planke, S., & Jonquet-Kolsto, H. (2003). Extension, crustal structure and magmatism at the outer Vøring Basin, Norwegian Margin. *Journal of Geological Society*, 160(2), 197–208. <https://doi.org/10.1144/0016-764902-055>
- Gernigon, L., Ringenbach, J.-C., Planke, S., & Le Gall, B. (2004). Deep structures and breakup along volcanic rifted margins: Insights from integrated studies along the outer Vøring Basin (Norway). *Marine and Petroleum Geology*, 21(3), 363–372. <https://doi.org/10.1016/j.marpetgeo.2004.01.005>
- Gomez, M., Verges, J., Fernandez, M., Torne, M., Ayala, C., Wheeler, W., & Karpuz, R. (2004). Extensional geometry of the Mid-Norwegian Margin before Early Tertiary continental breakup. *Marine and Petroleum Geology*, 21(2), 177–194. <https://doi.org/10.1016/j.marpetgeo.2003.11.017>
- Gradstein, F. M., Anthonissen, E., Brunstad, H., Charnock, M., Hammer, Ø., Hellem, T., & Lervik, K. S. (2010). Norwegian Offshore Stratigraphic Lexicon (NORLEX). *Newsletters on Stratigraphy*, 44(1), 73–86. <https://doi.org/10.1127/0078-0421/2010/0005>
- Grant, N., Bouma, A., & McIntyre, A. (1999). The Turonian play in the Faeroe–Shetland Basin. *Geological Society, London, Petroleum Geology Conference Series*, 5, 661–673. <https://doi.org/10.1144/0050661>
- Harding, I., & Allen, R. (1995). Dinocysts and the palaeoenvironmental interpretation of non-marine sediments: An example from the Wealden of the Isle of Wight (Lower Cretaceous, southern England). *Cretaceous Research*, 16(6), 727–743. <https://doi.org/10.1006/cres.1995.1046>
- Harry, D. L., & Bowling, J. C. (1999). Inhibiting magmatism on nonvolcanic rifted margins. *Geology*, 27(10), 895–898. [https://doi.org/10.1130/0091-7613\(1999\)027<0895:IMONRM>2.3.CO;2](https://doi.org/10.1130/0091-7613(1999)027<0895:IMONRM>2.3.CO;2)
- Henstra, G. A., Gawthorpe, R. L., Helland-Hansen, W., Ravnås, R., & Rotevatn, A. (2016). Depositional systems in multiphase rifts: Seismic case study from the Lofoten margin, Norway. *Basin Research*, 29(4), 447–469. <https://doi.org/10.1111/bre.12183>
- Herzberg, C. T., Fyfe, W. S., & Carr, M. J. (1983). Density constraints on the formation of the continental Moho and crust. *Contributions to Mineralogy and Petrology*, 84(1), 1–5. <https://doi.org/10.1007/BF01132324>

- Hjelstuen, B. O., Eldholm, O., & Skogseid, J. (1999). Cenozoic evolution of the northern Vøring Margin. *Geological Society of America Bulletin*, 111(12), 1,792–1,807. [https://doi.org/10.1130/0016-7606\(1999\)111<1792:CEOTNV>2.3.CO;2](https://doi.org/10.1130/0016-7606(1999)111<1792:CEOTNV>2.3.CO;2)
- Huisman, R. S., & Beaumont, C. (2011). Depth-dependent extension, two-stage breakup and cratonic underplating at rifted margins. *Nature*, 473(7,345), 74–78. <https://doi.org/10.1038/nature09988>
- Imber, J., Holdsworth, R. E., McCaffrey, K. J. W., Wilson, R. W., Jones, R. R., England, R. W., & Gjeldvik, G. (2005). Early Tertiary sinistral transpression and fault reactivation in the western Vøring Basin, Norwegian Sea: Implications for hydrocarbon exploration and pre-breakup deformation in ocean margin basins. *AAPG Bulletin*, 89(8), 1043–1069. <https://doi.org/10.1306/02240504043>
- Kjennerud, T., & Vergara, L. (2005). Cretaceous to Palaeogene 3D palaeobathymetry and sedimentation in the Vøring Basin, Norwegian Sea. In A. G. Doré & B. A. Vining (Eds.), *Petroleum geology: Northwest Europe and global perspectives—Proceedings of the 6th Petroleum Geology Conference* (pp. 815–831). Geological Society, London.
- Kusznir, N. J., Hunsdale, R., Roberts, A. M., & iSIMM Team (2005). Timing and magnitude of depth-dependent lithosphere stretching on the S. Lofoten and N. Vøring continental margins offshore mid-Norway: Implications for subsidence and hydrocarbon maturation at volcanic rifted margins. In A. G. Doré & B. A. Vining (Eds.), *Petroleum geology: Northwest Europe and global perspectives, Proceedings of the 6th Petroleum Geology Conference* (pp. 767–783). Geological Society, London.
- Lavier, L. L., & Manatschal, G. (2006). A mechanism to thin the continental lithosphere at magma-poor margins. *Nature*, 440(7,082), 324–328. <https://doi.org/10.1038/nature04608>
- Lister, G. S., Etheridge, M. A., & Symonds, P. A. (1986). Detachment faulting and the evolution of passive continental margins. *Geology*, 14(3), 246–250. [https://doi.org/10.1130/0091-7613\(1986\)14<246:DFATEO>2.0.CO;2](https://doi.org/10.1130/0091-7613(1986)14<246:DFATEO>2.0.CO;2)
- Lundin, E. R., & Doré, A. G. (1997). A tectonic model for the Norwegian passive margin with implications for the NE Atlantic: Early Cretaceous to break-up. *Geological Society London Journal*, 154(3), 545–550. <https://doi.org/10.1144/gsjgs.154.3.0545>
- Lundin, E. R., & Doré, A. G. (2011). Hyperextension, serpentinization, and weakening: A new paradigm for rifted margin compressional deformation. *Geology*, 39(4), 347–350. <https://doi.org/10.1130/G31499.1>
- Lundin, E. R., Doré, A. G., Rønning, K., & Kyrkjebø, R. (2013). Repeated inversion and collapse in the Late Cretaceous–Cenozoic northern Vøring Basin, offshore Norway. *Petroleum Geoscience*, 19(4), 329–341. <https://doi.org/10.1144/petgeo2012-022>
- Lundin, E. R., Rønning, K., Doré, A. G., & Olesen, O. (2002). Hel Graben, Vøring Basin, Norway—A possible major cauldron? Paper presented at 25th Nordic Geological Winter Meeting, 6–9 January, Reykjavik, Abstract Volume, 132.
- Maystrenko, Y. P., Olesen, O., Gernigon, L., & Gradmann, S. (2017). Deep structure of the Lofoten–Vesterålen segment of the mid-Norwegian continental margin and adjacent areas derived from 3-D density modeling. *Journal of Geophysical Research: Solid Earth*, 122, 1402–1433. <https://doi.org/10.1002/2016JB013443>
- McKenzie, D. (1978). Some remarks on the development of sedimentary basins. *Earth and Planetary Science Letters*, 40(1), 25–32. [https://doi.org/10.1016/0012-821X\(78\)90071-7](https://doi.org/10.1016/0012-821X(78)90071-7)
- Meyer, R., van Wijk, J., & Gernigon, L. (2007). The North Atlantic Igneous Province: A review of models for its formation. In G. R. Foulger & D. M. Jurdy (Eds.), *Plates, plumes and planetary processes, Geological Society of America Special Paper* (Vol. 430, pp. 525–552). [https://doi.org/10.1130/2007.2430\(26\)](https://doi.org/10.1130/2007.2430(26))
- Mjelde, R., Digranes, P., Shimamura, H., Shiobara, H., Kodaira, S., Brekke, S. H., ... Thorbjørnsen, T. (1998). Crustal structure of the northern part of the Vøring Basin, mid-Norway margin, from wide-angle seismic and gravity data. *Tectonophysics*, 293(3–4), 175–205. [https://doi.org/10.1016/S0040-1951\(98\)00090-0](https://doi.org/10.1016/S0040-1951(98)00090-0)
- Mjelde, R., Goncharov, A., & Müller, R. D. (2013). The Moho: Boundary above upper mantle peridotites or lower crustal eclogites? A global review and new interpretations for passive margins. *Tectonophysics*, 609, 636–650. <https://doi.org/10.1016/j.tecto.2012.03.001>
- Mjelde, R., Kodaira, S., Shimamura, H., Kanazawa, T., Shiobara, H., Berg, E. W., & Riise, O. (1997). Crustal structure of the central part of the Vøring Basin, mid-Norway margin, from ocean bottom seismographs. *Tectonophysics*, 277, 235–257. [https://doi.org/10.1016/S0040-1951\(97\)00028-0](https://doi.org/10.1016/S0040-1951(97)00028-0)
- Mjelde, R., Kvarven, T., Faleide, J. I., & Thybo, H. (2016). Lower crustal high-velocity bodies along North Atlantic passive margins, and their link to Caledonian suture zone eclogites and Early Cenozoic magmatism. *Tectonophysics*, 670, 16–29. <https://doi.org/10.1016/j.tecto.2015.11.021>
- Mjelde, R., Raum, T., Digranes, P., Shimamura, H., Shiobara, H., & Kodaira, S. (2003). V_p/V_s ratio along the Vøring Margin, NE Atlantic, derived from OBS data: Implications on lithology and stress field. *Tectonophysics*, 369(3–4), 175–197. [https://doi.org/10.1016/S0040-1951\(03\)00198-7](https://doi.org/10.1016/S0040-1951(03)00198-7)
- Mogensen, T. E., Nyby, R., Karpuz, R., & Haremo, P. (2000). Late Cretaceous and Tertiary structural evolution of the northeastern part of the Vøring Basin, Norwegian Sea. In A. Nøttvedt et al. (Eds.), *Dynamics of the Norwegian margin: Geological Society, London, Special Publication* (Vol. 167, pp. 379–396). <https://doi.org/10.1144/GSL.SP.2000.167.01.14>
- Morton, A. C., & Grant, S. (1998). Cretaceous depositional systems in the Norwegian Sea: Heavy mineral constraints. *AAPG Bulletin*, 82(2), 274–290.
- Mosar, J. (2000). Depth of extensional faulting on the Mid-Norway Atlantic passive margin. *Norgesgeologiske undersøkelse Bulletin*, 437, 33–41.
- Mosar, J., Eide, E. A., Osmundsen, P. T., Sommaruga, A., & Torsvik, T. (2002). Greenland–Norway separation: A geodynamic model for the North Atlantic. *Norwegian Journal of Geology*, 82, 281–298.
- Neumann, E.-R., Svensen, H., Tegner, C., Planke, S., Thirlwall, M., & Jarvis, K. E. (2013). Sill and lava geochemistry of the mid-Norway and NE Greenland conjugate margins. *Geochemistry, Geophysics, Geosystems*, 14, 3666–3690. <https://doi.org/10.1002/ggge.20224>
- Ní Dheasúna, M., M. Ballesteros, J. Leven (2012). A regional velocity cube for depth conversion on the Northwest Shelf, *PESA News Resources*, August/September, 2012, 40–43. Retrieved from www.pnronline.com.au/article.aspx?p=1&id=810
- Nirrengarten, M., Gernigon, L., & Manatschal, G. (2014). Lower crustal bodies in the Møre volcanic rifted margin: Geophysical determination and geological implications. *Tectonophysics*, 636, 143–157. <https://doi.org/10.1016/j.tecto.2014.08.004>
- Nøhr-Hansen, H. (1993). Dinoflagellate cyst stratigraphy of the Barremian to Albian, Lower Cretaceous, North-East Greenland. *Bulletin. Grønlands Geologiske Undersøgelse*, 166, 1–171.
- Nøhr-Hansen, H. (2012). Palynostratigraphy of the Cretaceous–lower Palaeogene sedimentary succession in the Kangerlussuaq Basin, southern East Greenland. *Review of Palaeobotany and Palynology*, 178, 59–90. <https://doi.org/10.1016/j.revpalbo.2012.03.009>
- Norwegian Petroleum Directorate (NPD) (2017). Factpages. Retrieved from <http://factpages.npd.no/factpages/>, last access April 2017.
- Olde, K., Jarvis, I., Pearce, M. M., Uličný, D., Tocher, B., Trabucchi-Alexandre, J., & Gröcke, D. (2014). A revised northern European Turonian (Upper Cretaceous) dinoflagellate cyst biostratigraphy: Integrating palynology and carbon isotope events. *Review of Palaeobotany and Palynology*, 213, 1–16.
- Olesen, O., Brønner, M., Ebbing, J., Gellein, J., Gernigon, L., Koziel, J., ... Usov, S. (2010). New aeromagnetic and gravity compilations from Norway and adjacent areas—Methods and applications. In B. A. Vining & S. C. Pickering (Eds.), *Petroleum geology: From mature basins*

- to new frontiers, *proceedings of the 7th Petroleum Geology Conference* (pp. 559–586). London: Geological Society. <https://doi.org/10.1144/0070559>
- Osmundsen, P. T., & Ebbing, J. (2008). Style of extension offshore mid-Norway and implications for mechanisms of crustal thinning at passive margins. *Tectonics*, 27, TC6016. <https://doi.org/10.1029/2007TC002242>
- Osmundsen, P. T., Péron-Pinvidic, G., Ebbing, J., Erratt, D., Fjellanger, E., Bergslien, D., & Syvertsen, S. E. (2016). Extension, hyperextension and mantle exhumation offshore Norway: A discussion based on 6 crustal transects. *Norwegian Journal of Geology*, 96, 343–372. <https://doi.org/10.17850/njg96-4-05>
- Osmundsen, P. T., Sommaruga, A., Skilbrei, J. R., & Olesen, O. (2002). Deep structure of the mid-Norway rifted margin. *Norwegian Journal of Geology*, 82, 205–224.
- Pascoe, R., Hooper, R., Storhaug, K., & Harper, H. (1999). Evolution of extensional styles at the southern termination of the Nordland Ridge, Mid-Norway: A response to variations in coupling above Triassic salt. *Geological Society, London, Petroleum Geology Conference Series*, 5, 83–90. <https://doi.org/10.1144/0050083>
- Pérez-Gussinyé, M., & Reston, T. J. (2001). Rheological evolution during extension at nonvolcanic rifted margins: Onset of serpentinization and development of detachments leading to continental breakup. *Journal of Geophysical Research*, 106, 3961–3975. <https://doi.org/10.1029/2000JB900325>
- Péron-Pinvidic, G., Manatschal, G., & Osmundsen, P. T. (2013). Structural comparison of archetypal Atlantic rifted margins: A review of observations and concepts. *Marine and Petroleum Geology*, 43, 21–47. <https://doi.org/10.1016/j.marpetgeo.2013.02.002>
- Péron-Pinvidic, G., & Osmundsen, P. T. (2016). Architecture of the distal and outer domains of the mid-Norwegian rifted margin: Insights from the Rån-Gjallar ridges system. *Marine and Petroleum Geology*, 77, 280–299. <https://doi.org/10.1016/j.marpetgeo.2016.06.014>
- Petersen, K. D., & Schiffer, C. (2016). Wilson cycle passive margins: Control of orogenic inheritance on continental breakup. *Gondwana Research*, 39, 131–144. <https://doi.org/10.1016/j.gr.2016.06.012>
- Pinel, V., & Jaupart, C. (2005). Caldera formation by magma withdrawal from a reservoir beneath a volcanic edifice. *Earth and Planetary Science Letters*, 230(3–4), 273–287. <https://doi.org/10.1016/j.epsl.2004.11.016>
- Planke, S., Skogseid, J., & Eldholm, O. (1991). Crustal structure off Norway, 62–70°N. *Tectonophysics*, 189(1–4), 91–107. [https://doi.org/10.1016/0040-1951\(91\)90489-F](https://doi.org/10.1016/0040-1951(91)90489-F)
- Radmacher, W., Mangerud, G., & Tyska, J. (2015). Dinoflagellate cyst biostratigraphy of Upper Cretaceous strata from two wells in the Norwegian Sea. *Review of Palaeobotany and Palynology*, 216, 18–32. <https://doi.org/10.1016/j.revpalbo.2014.12.007>
- Radmacher, W., Tyska, J., Mangerud, G., & Pearce, M. A. (2014). Dinoflagellate cyst biostratigraphy of the Late Albian to Early Maastrichtian in the southwestern Barents Sea. *Marine and Petroleum Geology*, 57, 109–121. <https://doi.org/10.1016/j.marpetgeo.2014.04.008>
- Redfield, T. F., Braathen, A., Gabrielsen, R. H., Osmundsen, P. T., Torsvik, T., & Andriessen, P. A. M. (2005). Late Mesozoic to Early Cenozoic components of vertical separation across the Møre Trøndelag Fault Complex, Norway. *Tectonophysics*, 395(3–4), 233–249. <https://doi.org/10.1016/j.tecto.2004.09.012>
- Ren, S., Faleide, J. I., Eldholm, O., Skogseid, J., & Gradstein, F. (2003). Late Cretaceous–Paleocene development of the NW Vøring Basin. *Marine and Petroleum Geology*, 20(2), 177–206. [https://doi.org/10.1016/S0264-8172\(03\)00005-9](https://doi.org/10.1016/S0264-8172(03)00005-9)
- Ren, S., Skogseid, J., & Eldholm, O. (1998). Late Cretaceous–Paleocene extension on the Vøring volcanic margin. *Marine Geophysical Researches*, 20(4), 343–369. <https://doi.org/10.1023/A:1004554217069>
- Reynisson, R. F., Ebbing, J., Lundin, E., & Osmundsen, P. T. (2010). Properties and distribution of lower crustal bodies on the mid-Norwegian margin. *Geological Society, London, Petroleum Geology Conference Series*, 7, 843–854. <https://doi.org/10.1144/0070843>
- Rise, L., Ottesen, D., Berg, K., & Lundin, E. (2005). Large-scale development of the mid-Norwegian margin during the last 3 million years. *Marine and Petroleum Geology*, 22(1–2), 33–44. <https://doi.org/10.1016/j.marpetgeo.2004.10.010>
- Roberts, A. M., Corfield, R. I., Kusznir, N. J., Matthews, S. J., Hansen, E.-K., & Hooper, R. J. (2009). Mapping palaeostructure and palaeobathymetry along the Norwegian Atlantic continental margin, Møre and Vøring basins. *Petroleum Geoscience*, 15(1), 27–43. <https://doi.org/10.1144/1354-079309-804>
- Rosenbaum, G., Regenauer-Lieb, K., & Weinberg, R. F. (2010). Interaction between mantle and crustal detachments: A nonlinear system controlling lithospheric extension. *Journal of Geophysical Research*, 115, B11412. <https://doi.org/10.1029/2009JB006696>
- Rüpke, L. H., Schmalholz, S. M., Schmid, D. W., & Podladchikov, Y. Y. (2008). Automated thermotectonostratigraphic basin reconstruction: Viking Graben case study. *AAPG Bulletin*, 92(3), 309–326. <https://doi.org/10.1306/11140707009>
- Rüpke, L. H., Schmid, D. W., Hartz, E. H., & Martinsen, B. (2010). Basin modelling of a transform margin setting: Structural, thermal and hydrocarbon evolution of the Tano Basin, Ghana. *Petroleum Geoscience*, 16(3), 283–298. <https://doi.org/10.1144/1354-079309-905>
- Rüpke, L. H., Schmid, D. W., Perez-Gussinye, M., & Hartz, E. (2013). Interrelation between rifting, faulting, sedimentation, and mantle serpentinization during continental margin formation—including examples from the Norwegian Sea. *Geochemistry, Geophysics, Geosystems*, 14, 4351–4369. <https://doi.org/10.1002/ggge.20268>
- Saltus, R. W., & Blakely, R. J. (2011). Unique geologic insights from “non-unique” gravity and magnetic interpretation. *GSA Today*, 21(12), 4–10. <https://doi.org/10.1130/G136A.1>
- Scheck-Wenderoth, M., & Maystrenko, Y. P. (2011). 3D lithospheric-scale structural model of the Norwegian continental margin (the Vøring and Møre basins). *Scientific Technical Report STR11/02–Data*, GFZ Potsdam. <https://doi.org/10.2312/GFZ.b103-11027>
- Scheck-Wenderoth, M., Raum, T., Faleide, J. I., Mjelde, R., & Horsfield, B. (2007). The transition from the continent to the ocean: A deeper view on the Norwegian margin. *Journal of the Geological Society of London*, 164(4), 855–868. <https://doi.org/10.1144/0016-76492006-131>
- Serck, C. S., Faleide, J. I., Braathen, A., Kjøllhamar, B., & Escalona, A. (2017). Jurassic to Early Cretaceous basin configuration (s) in the Fjerdjupet subbasin, SW Barents Sea. *Marine and Petroleum Geology*, 86, 874–891. <https://doi.org/10.1016/j.marpetgeo.2017.06.044>
- Simon, K., Huismans, R. S., & Beaumont, C. (2009). Dynamical modelling of lithospheric extension and small-scale convection: Implications for magmatism during the formation of volcanic rifted margins. *Geophysical Journal International*, 176(1), 327–350. <https://doi.org/10.1111/j.1365-246X.2008.03891.x>
- Skogseid, J., Eldholm, O., & Larsen, V. B. (1992). Vøring Basin: Subsidence and tectonic evolution. In R. M. Larsen, et al. (Eds.), *Structural and tectonic modeling and its application to petroleum geology*, Norwegian Petroleum Society, Special Publications (Vol. 1, pp. 55–82). Elsevier, Amsterdam.
- Skogseid, J., Planke, S., Faleide, J. I., Pedersen, T., Eldholm, O., & Neverdal, F. (2000). NE Atlantic continental rifting and volcanic margin formation. In A. Nøttvedt et al. (Eds.), *Dynamics of the Norwegian margin*, Geological Society, London, Special Publication (Vol. 167, pp. 295–326). Geological Society, London. <https://doi.org/10.1144/GSL.SP.2000.167.01.12>
- Slagstad, T., Davidsen, B., & Daly, J. S. (2011). Age and composition of crystalline basement rocks on the Norwegian continental margin: Offshore extension and continuity of the Caledonian–Appalachian orogenic belt. *Journal of the Geological Society, London*, 168(5), 1167–1185. <https://doi.org/10.1144/0016-76492010-136>

- Smelror, M., Jacobsen, T., Rise, L., Skarbø, O., Verdenius, J. G., & Vigran, J. O. (1994). Jurassic to Cretaceous stratigraphy of shallow cores on the Møre Basin Margin, Mid-Norway. *Norsk Geologisk Tidsskrift*, 74, 89–107.
- Somme, T. O., & Jackson, C. A. L. (2013). Source-to-sink analysis of ancient sedimentary systems using a subsurface case study from the Møre–Trøndelag area of southern Norway: Part 2—Sediment dispersal and forcing mechanisms. *Basin Research*, 25, 512–531. <https://doi.org/10.1111/bre.12014>
- Sun, Z., Zhou, D., Wu, S. M., Zhong, Z. H., Myra, K., Jiang, J. Q., & Fan, H. (2009). Patterns and dynamics of rifting on passive continental margin from shelf to slope of the northern South China Sea: Evidence from 3D analogue modeling. *Journal of Earth Science China*, 20(1), 136–146. <https://doi.org/10.1007/s12583-009-0011-6>
- Swiecicki, T., Gibbs, P. B., Farrow, G. E., & Coward, M. P. (1998). A tectonostratigraphic framework for the mid-Norway region. *Marine and Petroleum Geology*, 15(3), 245–276. [https://doi.org/10.1016/S0264-8172\(97\)00029-9](https://doi.org/10.1016/S0264-8172(97)00029-9)
- Talwani, M. (1973). Computer usage in the computation of gravity anomalies. *Methods in Computational Physics*, 13, 343–389.
- Talwani, M., & Eldholm, O. (1977). Evolution of the Norwegian-Greenland Sea. *Bulletin Geological Society of America*, 88(7), 969–999. [https://doi.org/10.1130/0016-7606\(1977\)88<969:EOTNS>2.0.CO;2](https://doi.org/10.1130/0016-7606(1977)88<969:EOTNS>2.0.CO;2)
- ten Brink, U. S., & Ben-Avraham, Z. (1989). The anatomy of a pull-apart basin: Seismic reflection observations of the Dead Sea Basin. *Tectonics*, 8(2), 333–350. <https://doi.org/10.1029/TC008i002p00333>
- Theissen-Krah, S., & Rüpke, L. H. (2010). Feedbacks of sedimentation on crustal heat flow: New insights from the Vøring Basin, Norwegian Sea. *Basin Research*, 22, 976–990. <https://doi.org/10.1111/j.1365-2117.2009.00437>
- Theissen-Krah, S., Zastrozhnov, D., Abdelmalak, M. M., Schmid, D. W., Faleide, J. I., & Gernigon, L. (2017). Tectonic evolution and extension at the Møre Margin—Offshore mid-Norway. *Tectonophysics*, 721, 227–238. <https://doi.org/10.1016/j.tecto.2017.09.009>
- Thomsen, L. (1986). Weak elastic anisotropy. *Geophysics*, 51(10), 1954–1966. <https://doi.org/10.1190/1.1442051>
- Tsikalas, F., Faleide, J. I., & Eldholm, O. (2001). Lateral variations in tectono-magmatic style along the Lofoten-Vesterålen margin off Norway. *Marine and Petroleum Geology*, 18(7), 807–832. [https://doi.org/10.1016/S0264-8172\(01\)00030-7](https://doi.org/10.1016/S0264-8172(01)00030-7)
- Tsikalas, F., Faleide, J. I., Eldholm, O., & Wilson, J. (2005). Late Mesozoic–Cenozoic structural and stratigraphic correlations between the conjugate mid-Norway and NE Greenland continental margins. In A. G. Doré & B. A. Vining (Eds.), *Petroleum geology: Northwest Europe and global perspectives—Proceedings of the 6th Petroleum Geology Conference, Geological Society, London* (pp. 785–801). <https://doi.org/10.1144/0060785>
- Tsikalas, F., Faleide, J. I., & Kuznir, N. J. (2008). Along-strike variations in rifted margin crustal architecture and lithosphere thinning between northern Vøring and Lofoten margin segments off mid-Norway. *Tectonophysics*, 458(1–4), 68–81. <https://doi.org/10.1016/j.tecto.2008.03.001>
- Våagnes, E., Gabrielsen, R. H., & Haremo, P. (1998). Late Cretaceous–Cenozoic intraplate contractional deformation at the Norwegian continental shelf: Timing, magnitude and regional implications. *Tectonophysics*, 300(1–4), 29–46. [https://doi.org/10.1016/S0040-1951\(98\)00232-7](https://doi.org/10.1016/S0040-1951(98)00232-7)
- Van Avendonk, H. J. A., Lavie, L. L., Shillington, D. J., & Manatschal, G. (2009). Extension of continental crust at the margin of the eastern Grand Banks, Newfoundland. *Tectonophysics*, 468(1–4), 131–148. <https://doi.org/10.1016/j.tecto.2008.05.030>
- Van Wijk, J., Koning, D., Axen, G., Coblenz, D., Gragg, E., & Sion, B. (2018). Tectonic subsidence, geoid analysis, and the Miocene–Pliocene unconformity in the Rio Grande rift, southwestern United States: Implications for mantle upwelling as a driving force for rift opening. *Geosphere*. <https://doi.org/10.1130/GES01522.1>
- Van Wijk, J. W., Huismans, R. S., Ter Voorde, M., & Cloetingh, S. A. P. L. (2001). Melt generation at volcanic continental margins: No need for a mantle plume? *Geophysical Research Letters*, 28(20), 3995–3998. <https://doi.org/10.1029/2000GL012848>
- Vergara, L., Wreglesworth, I., Trayfoot, M., & Richardsen, G. (2001). The distribution of Cretaceous and Paleocene deep-water reservoirs in the Norwegian Sea basins. *Petroleum Geoscience*, 7(4), 395–408. <https://doi.org/10.1144/petgeo.7.4.395>
- Walker, I. M., Berry, K. A., Bruce, J. R., Bystøl, L., & Snow, J. H. (1997). Structural modelling of regional depth profiles in the Vøring Basin: Implications for the structural and stratigraphic development of the Norwegian passive margin. *Journal of the Geological Society, London*, 154(3), 537–544. <https://doi.org/10.1144/gsjgs.154.3.0537>
- Wangen, M., Mjelde, R., & Faleide, J. I. (2011). The extension of the Vøring margin (NE Atlantic) in case of different degrees of magmatic underplating. *Basin Research*, 23(1), 83–100. <https://doi.org/10.1111/j.1365-2117.2010.00467.x>
- White, N. J., & McKenzie, D. P. (1988). Formation of the “Steer’s Head” geometry of sedimentary basins by differential stretching of the crust and mantle. *Geology*, 16(3), 250–253. [https://doi.org/10.1130/0091-7613\(1988\)016<0250:FOTSSH>2.3.CO;2](https://doi.org/10.1130/0091-7613(1988)016<0250:FOTSSH>2.3.CO;2)
- White, R. S., Eccles, J. D., & Roberts, A. W. (2010). Constraints on volcanism, igneous intrusion and stretching on the Rockall–Faroe continental margin. *Petroleum Geology Conference Series*, 7, 831–842. <https://doi.org/10.1144/0070831>
- White, R. S., & McKenzie, D. P. (1989). Magmatism at rift zones: The generation of volcanic continental margins and flood basalts. *Journal of Geophysical Research*, 94(B6), 7685–7729. <https://doi.org/10.1029/JB094iB06p07685>
- White, R. S., Smith, L. K., Roberts, A. W., Christie, P. A. F., Kuznir, N. J., & iSIMM Team (2008). Lower-crustal intrusion on the North Atlantic continental margin. *Nature*, 452(7186), 460–464. <https://doi.org/10.1038/nature06687>
- Whitham, A. G., Price, S. P., Koriani, A. M., & Kelly, S. R. A. (1999). Cretaceous (post-Valanginian) sedimentation and rift events in NE Greenland (71–77°N). *Geological Society, London, Petroleum Geology Conference Series*, 5, 325–336. <https://doi.org/10.1144/0050325>
- Xie, X., & Heller, P. L. (2009). Plate tectonics and basin subsidence history. *GSA Bulletin*, 121, 55–64. <https://doi.org/10.1130/B26398.1>
- Yamasaki, T., & Gernigon, L. (2009). Styles of lithospheric extension controlled by underplated mafic bodies. *Tectonophysics*, 468(1–4), 169–184. <https://doi.org/10.1016/j.tecto.2008.04.024>
- Yamasaki, T., & Gernigon, L. (2010). Redistribution of the lithosphere deformation by the emplacement of underplated mafic bodies: Implications for microcontinent formation. *Journal of the Geological Society of London*, 167(5), 961–971. <https://doi.org/10.1144/0016-76492010-027>

Paper II

Tectonic Evolution and Extension at the Møre Margin–Offshore mid-Norway

By

*Sonja Theissen-Krah, Dmitry Zastrozhnov, Mansour M. Abdelmalak, Daniel W. Schmid,
Jan Inge Faleide & Laurent Gernigon*

Tectonophysics, 721, 227-238

II

Tectonic evolution and extension at the Møre Margin – Offshore mid-Norway

S. Theissen-Krah^{a,b,*}, D. Zastrozhnov^{b,c}, M.M. Abdelmalak^{b,c}, D.W. Schmid^d, J.I. Faleide^b, L. Gernigon^e

^a GEOMAR Helmholtz Centre for Ocean Research Kiel, Wischhofstr. 1-3, 24148 Kiel, Germany

^b Centre for Earth Evolution and Dynamics (CEED), University of Oslo, Norway

^c Volcanic Basin Petroleum Research (VBPR), Oslo Science Park, 0349 Oslo, Norway

^d Physics of Geological Processes (PGP), University of Oslo, Norway

^e Geological Survey of Norway (NGU), Leiv Erikssons vei, 39, Trondheim, Norway

ARTICLE INFO

Keywords:

Basin modelling
Stratigraphy reconstruction
Møre continental margin
Extension
Stretching factors

ABSTRACT

Lithospheric stretching is the key process in forming extensional sedimentary basins at passive rifted margins. This study explores the stretching factors, resulting extension, and structural evolution of the Møre segment on the Mid-Norwegian continental margin. Based on the interpretation of new and reprocessed high-quality seismic, we present updated structural maps of the Møre margin that show very thick post-rift sediments in the central Møre Basin and extensive sill intrusion into the Cretaceous sediments. A major shift in subsidence and deposition occurred during mid-Cretaceous. One transect across the Møre continental margin from the Slørebotn Subbasin to the continent-ocean boundary is reconstructed using the basin modelling software TecMod. We test different initial crustal configurations and rifting events and compare our structural reconstruction results to stretching factors derived both from crustal thinning and the classical backstripping/decompaction approach. Seismic interpretation in combination with structural reconstruction modelling does not support the lower crustal bodies as exhumed and serpentinised mantle. Our extension estimate along this transect is $\sim 188 \pm 28$ km for initial crustal thickness varying between 30 and 40 km.

1. Introduction

The Møre Margin is a segment of the Mid-Norwegian passive continental margin and is located between 62° and 65°N and 0° and 6°E (Fig. 1). It is separated from the Vøring Margin by the Jan Mayen Corridor in the north and by the north-westward extension of the Tampen Spur from the Magnus Basin in the south. The Møre continental margin comprises of the deep sedimentary Møre Basin, which is bounded to the Møre Marginal High by the Faroe-Shetland-Escarpment to the west and the Møre Trøndelag Fault Complex to the east. The Møre Margin is divided into several sub-basins and highs (Fig. 1) and includes the Møre Trøndelag Fault Complex and the Slørebotn Subbasin in the east.

The Mid-Norwegian margin has been studied extensively for hydrocarbon exploration and consequently large amounts of seismic data have been acquired. Many studies that address similar stretching and extension estimates focused on the Vøring Margin (e.g. Gernigon et al., 2003; Reemst and Cloetingh, 2000; Theissen and Rüpke, 2010; Wangen and Faleide, 2008; Wangen et al., 2011). However few studies address

the Møre continental margin segment specifically (Gabrielsen et al., 1999; Gomez et al., 2004; Skogseid et al., 2000). This study focuses on stretching factors and extension along a seismic profile crossing the entire Møre continental margin. We further discuss the implication of different modelling approaches, seismic interpretation, rifting events, and initially assumed crustal configuration. The modelling tool TecMod and the classical backstripping/decompaction method are used to reconstruct the stratigraphy and calculate stretching and extension. The stretching factor (beta) is compared to estimates from crustal thinning, where beta is estimated by dividing the assumed initial crustal thickness prior to rifting by the today's thickness.

2. Geological setting

The Mid-Norwegian margin developed through several rift episodes since the collapse of the Caledonides in the Devonian. The final rift phase during the Upper Cretaceous eventually led to breakup between Greenland and Eurasia in the Eocene (Talwani and Eldholm, 1972).

Several early rift phases during Late Palaeozoic-Early Mesozoic have

* Corresponding author at: GEOMAR Helmholtz Centre for Ocean Research Kiel, Wischhofstr. 1-3, 24148 Kiel, Germany.

E-mail address: stheissen@geomar.de (S. Theissen-Krah).

Fig. 1. a) Bathymetry and topography map showing the study area. The red lines indicate the seismic lines used for the structural reconstruction; the red frame marks the location of (b). b) Close up of the study area including main structural features in the Møre Margin. Seismic profiles used for the reconstruction study in red. Black dots indicate well locations in the area and the wells projected onto the key profile are marked in red. GH = Gossa High, MMH = Møre Marginal High, OH = Ona High, MTFC = Møre-Trøndelag Fault Complex, SSB = Sjørebotn Sub-basin, VH = Vigra High, VMH = Vøring Marginal High. Interpretation of volcanic facies units are taken from [Abdelmalak et al. \(2015, 2016b\)](#). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Fig. 2. Top (a–c) row shows time-structure maps of the Cretaceous units. Time-thickness maps are plotted at the bottom (d–f). The blue and red lines mark the continent ocean boundary (COB) and the extent of the volcanics, respectively. The COB is defined as seaward termination of the K-reflector (Abdelmalak et al., 2016a). The thick red line shows the location of the key profile. EH = Ervik High; FH = Frøya High; GH = Grip High; GNU = Gnausen High; GSH = Gossa High; HD = Haysule Dome; HHA = Helland-Hansen Arch; HT = Halten Terrace; ID = Isak Dome; KH = Kinn High; KRH = Kiran High; MA = Modgunn Arch; MB = Marulk Basin; MR = Manet Ridge; OH = Ona High; OLD = Ormen Lange Dome; RB = Rås Basin; SMA = Southern Modgunn Arch; SR = Slettringen Ridge; SS = Slorebotn Subbasin; VH = Vigra High. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

been suggested for the Norwegian-Greenland rift system (Grunnaleite and Gabrielsen, 1995). The earliest rift episode that can be observed in the seismic sedimentary strata at the Mid-Norwegian margin is Permian-Triassic in age. Evidence for this event has been described at the Trøndelag Platform and the Halten Terrace (Tsikalas et al., 2012). During this early rift phase Caledonian basement trends were re-activated and rifting was oriented ENE-WSW (Gomez et al., 2004).

The major rift event during which the crust was highly thinned took place during Late Jurassic–Early Cretaceous and was mainly non-magmatic (Brekke, 2000) with minor magmatic activity suggested by evidence for Cretaceous seamounts and the presence of pyroclastic rocks (Lundin and Dore, 1997) in the Vøring Basin. Both tectonic and thermal subsidence during the Cretaceous led to the accumulation of up to 8 km of sediments in local depocenters in the Vøring and Møre basins (Scheck-Wenderoth et al., 2007). The extension direction during Late Jurassic–Early Cretaceous shifted to a NW-SE direction and the final opening occurred almost perpendicular to the present day coastline (Doré et al., 1999; Mosar et al., 2002). A separate rift event during the mid-Cretaceous is debated; Grunnaleite and Gabrielsen (1995) suggest a mid-Cretaceous rift event, while other studies conclude that no stretching occurred during the mid-Cretaceous (Faereth and Lien, 2002).

A last stretching event during Late Cretaceous–Palaeocene is constrained by seismic data from the Vøring margin and it has been

suggested that the main rift event was initiated during Late Maastrichtian (Ren et al., 1998) or mid-Campanian time (Gernigon et al., 2003; Ren et al., 2003). The Campanian–Palaeocene rifting is less pronounced in seismic data from the Møre Margin, but evidence for brittle deformation is observed close to the Tertiary lava flows. Final lithospheric separation was accompanied by voluminous magmatism resulting in numerous sill intrusions, lava flows and characteristic seaward dipping reflectors (SDR), that cover large areas towards and over the continent-ocean boundary (COB) (Berndt et al., 2001; Planke et al., 2005; Ren et al., 2003 and Fig. 1). Breakup was initiated during the Palaeocene–Eocene transition (C24r, (Gernigon et al., 2012)).

Lower crustal bodies (LCBs) with high P-wave velocities (> 7 km/s) are characteristic for the outer parts of the Mid-Norwegian margin. LCBs occur below or are part of the lower crust and are observed beneath the marginal highs and the volcanic provinces. They continue to the east below the stretched and thinned crust of the deep Vøring and Møre basins. The LCBs have been imaged in most of the parts of the Møre Basin and are described to be 2–5 km thick with thickening beneath the intrabasinal highs (Faleide et al., 2008; Kvarven et al., 2014; Lundin and Dore, 2011; Mjelde et al., 2009). Traditionally the LCBs in the outer part of the Møre and Vøring margins have been interpreted as breakup related magmatic underplating (Mjelde et al., 2009; Olafsson et al., 1992; White and McKenzie, 1989), but alternative magmatic and non-magmatic origins (e.g. a post Caledonian metamorphic core

complex, a sill complex intruded into the lower crust or serpentinised mantle rocks) have been suggested and discussed (Dore et al., 1997; Gernigon et al., 2006; Gernigon et al., 2004; Osmundsen and Ebbing, 2008; Reynisson et al., 2010). The concept of mantle serpentinisation during continental rifting is based on the observations from the non-volcanic Iberian margin (Perez-Gussinye et al., 2001) and similar processes have been suggested for the inner parts of the LCB at the Norwegian margin (Lundin and Dore, 2011). Prerequisite for serpentinisation is a complete embrittlement of the crust and the presence of fluids. Rüpke et al. (2013) showed that despite the thick sedimentary cover that lead to higher temperatures in the lower crust compared to basin with thin sediment thicknesses, serpentinisation is possible above a critical stretching factor. The critical stretching factor depends on sedimentation rate and rift duration, but is generally high, e.g. above 6 for an average sedimentation rate of 0.2 mm/yr. The nature and origin of the LCB and in particular the question if serpentinisation can effectively occur at volcanic (and rifted) passive margins, and to what extent a comparison between volcanic and non-volcanic margins is valid (Peron-Pinvidic et al., 2013) or not (Geoffroy et al., 2015; Gernigon et al., 2015), is an ongoing debate.

3. Seismic interpretation

The study area comprises approximately 600 new and reprocessed seismic reflection and refraction data. The key seismic reflection lines cross the Møre Basin in northwest-southeast direction approximately perpendicular to the line of breakup. The main tectonic features are well imaged by the seismic data set, including several intra-basinal highs (Grip High, Vigra High, Ona High and Giske High), and the Slørebotn Subbasin. The southern extensions of the Modgunn Arch and Ormen Lange Dome are imaged in the northern part of the study area

(Figs. 1b, 2, and 3).

Structures and sedimentary layering are well recognised in the south-eastern part of the Møre Basin, whereas interpretation of the north-western part is hindered by numerous sill intrusions and lava flows close to the COB. The sedimentary units were tied to wells available in the Møre and southern Vøring area (Fig. 1b). Data of the deeper parts of the Møre Basin are limited and only 5 wells are drilled through Late Cretaceous or older formations, two of which are located in the Slørebotn Subbasin. The depth to the basement is therefore difficult to constrain in some parts of the area, and earlier studies define the depth of the Møre Basin to be the base Cretaceous level (Blystad et al., 1995). The new and reprocessed data set allows better interpretation of the depth to the basement in the proximal part of the margin. In the distal domain we used the lithosphere model from Scheck-Wenderoth and Maystrenko (2008) for the interpretation of the top basement.

Most of the seismic sections show the existence of thick sedimentary strata below the base Cretaceous reflection down to the recorded depth of 10 s TWT. The ages of this succession remain unknown since the base Cretaceous has never been drilled in the deep Møre Basin, but it is likely that these sediments correlate to the pre-Cretaceous sequences that have been drilled on the platforms and in the Slørebotn Subbasin.

A strong reflector within the Cretaceous sequence interpreted to be of intra-Cenomanian age can be followed on all seismic profiles in the study area. The depth to this reflector is about 5.5 to 6 s TWT in the central part of the basin (Fig. 2). Another prominent reflector interpreted as the base Cretaceous unconformity (BCU), defines Jurassic-Cretaceous syn-rift to post-rift transition reaches depth to 8 s TWT in the deep basins. This reflection is however not as distinct on all profiles as the intra-Cenomanian and different interpretations are possible (Grunnleite and Gabrielsen, 1995). In addition to the intra-Cretaceous

Fig. 3. a) Seismic reflection of the Møre transect (red line in Fig. 1b). The red area in the western part indicates the breakup lava flows. b) Geological interpretation of a). The question mark and dashed lines show the locations where the interpretation of the pre mid-Cretaceous strata are difficult and uncertain due to the widespread sill intrusions. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

and the base Cretaceous marker, the reflection corresponding to the Top Cretaceous level has been identified on all profiles, which allows us to estimate the thickness of the Cretaceous sequence in the entire Møre Basin (Fig. 2).

The present-day relief of the BCU is shown in Fig. 2a. The BCU is deepest, up to 8 s TWT on the western flank of the major structural highs and faults in the central basin (e.g. Vigra High, Grip High, Slettringen Ridge). The BCU shallows again towards the COB, which is also expressed in the Møre Marginal Plateau, a shallow platform, which can be interpreted as a northern continuation of the Faroe Platform. As suggested by Gernigon et al. (2015) this shallow platform (Møre Marginal Plateau) could represent a continental and marginal plateau that existed prior to continental breakup. In this case the plateau can be directly related to the prolongation of the Jan Mayen Microcontinent (JMMC) which was initially part of the Møre Basin between the Faroe Platform and the outer Vøring Basin (Gaina et al., 2009; Gernigon et al., 2012). From potential field modelling and the resulting pre-drift configuration, Gernigon et al. (2015) suggested that Mesozoic and possibly Palaeozoic sediments could be present beneath the outer Møre Margin and underneath a large part of the Møre Basin.

The mid-Cenomanian map represents the basin configuration after ~40 m.y. of syn-rift. The depth to this reflector increases from the Slørebotn Subbasin towards the central Møre Basin with maximum depth of 6–6.5 s TWT west of the Vigra and Grip highs, and shallows again towards the Møre Marginal High. Mid-Cenomanian reflector is absent on the Manet Ridge, Frøya High and in the western part of the Halten Terrace. The deepening of this reflector west of Vigra and Grip high, and the Slettringen Ridge is comparable to the BCU map (Fig. 2a). In contrast to the configuration at the base Cretaceous, the deepening of the Mid-Cenomanian reflector extends further to the west in the central parts and towards the COB. This subsidence suggests that minor faulting occurred during that time, probably due to reactivation of the old fault systems.

The Top Cretaceous (Fig. 2c) map shows a different pattern with a rather uniform depth except the prominent low in central basin and along the margin (Fig. 2b). Thick lava flows covering the western margin and post-breakup thermal subsidence is likely to have caused the deepening of the Top Cretaceous layer at the margin close to the COB. In the northern part (Møre-Vøring transition) the Top Cretaceous level seems to thin slightly towards the COB. Glacial loading and the corresponding high sedimentation rates have probably caused further subsidence seen in the central part.

The Late Cretaceous sediments are affected by the formation of domes during the Mid-Miocene compression (Grunnaleite and Gabrielsen, 1995; Lundin and Dore, 2002) and are distinct features on the Mid-Cenomanian (Ormen Lange dome, Isaak dome, Helland Hansen Arch) and Top Cretaceous time-structure maps (Fig. 2b,c).

Fig. 2d–f show the cumulative sediment thickness of the Cretaceous units and the evolution of the depocenters in the Early and Late Cretaceous. Several depocenters, up to 6 s TWT thicknesses, trending NNW-SSE are observed along the western flanks of major faults. Two prominent depocenters with up to 6.3 s thickness are present, one in the southern part of the central Møre Basin and a second depocenter that is located in the northern part at the transition to the Vøring Basin (southern Rås Basin). The comparison of the thickness distribution of the upper and Lower Cretaceous reveals a major shift in depositional environment during the mid-Cretaceous. The depocenters that were initiated during the main rift event in the Jurassic/Cretaceous are filled with thick Lower Cretaceous sediments (Fig. 2e). Very thin or absent Lower Cretaceous strata (e.g. Vigra, Ona, and Frøya High) is probably caused by non-deposition and/or erosion due to footwall uplift and rotation of the central and continental rafts preserved in the deep part of the Cretaceous Møre Basins.

The thickness distribution of the Upper Cretaceous (Fig. 2d) shows a different pattern with nearly uniform sediment thickness in the central Møre Basin. Subsidence and significantly higher sedimentation occur in

the northern part towards the transition to the Vøring Basin, which suggests different tectonic environments in the Møre Basin during the Late Cretaceous with only minor tectonic activity at the Møre margin.

Using average interval velocities proposed by Scheck-Wenderoth et al. (2007) for layer depth conversion allows thickness and depth estimates for the different sedimentary units. The Cenozoic succession is around 2 km thick in the Slørebotn Subbasin and increases to a maximum depth of over 4 km west of Vigra High. The base of the Upper Cretaceous reaches its maximum with 7.6 km in the sag basin between the Vigra and Grip highs. In average the central Møre Basin consists of 3 to > 4 km of Upper Cretaceous sedimentary strata.

Our data agree with previous interpretations that the Møre Basin is a deep sedimentary basin with thick Cretaceous sequence (e.g. Gomez et al., 2004). Particularly in the depocenters on either side of Vigra High thick sedimentary wedges are imaged down to 10 s TWT (corresponding to depths of 15–16 km). Additionally the new seismic data suggest, that the Møre Margin also contains thicker layers of possibly pre-Cretaceous strata and not shallow basement or an outer zone of exhumed continental mantle as suggested by (Peron-Pinvidic et al., 2013).

3.1. Modelling

We use the basin modelling software TecMod (Rüpke et al., 2008) for the reconstruction of the Møre transect. TecMod couples a forward model to an inverse modelling algorithm to fit a specific basin stratigraphy. The forward model is based on pure-shear kinematics and resolves for basin scale, e.g. sedimentation and compaction, and lithosphere processes, such as rifting, heat transfer, and flexural isostasy on a Lagrangian finite-element mesh. The advantage of this reconstruction is that in each time step, new sediment packages are deposited to be part of the computational domain for the following time steps. The sediments are therefore included in the thermal calculations to account for thermal blanketing effects which may lead to considerable differences in heat flow and subsidence history compared to the results of method that decouple both thermal and structural processes (Theissen and Rüpke, 2010; Wangen, 1995). During the inversion the stretching factors and palaeo-water depths values are iteratively updated until the forward modelling results match the observed input stratigraphy. During the fitting process only the thickness and the location of sediment packages are used to update the forward parameters and well data such as present-day temperature, vitrinite reflectance data and estimates of palaeo-water depth can therefore be used to verify the corresponding model results. Further details of the automated basin reconstruction method are described in Rüpke et al. (2008, 2010).

3.2. Møre transect

Our key profile across the Møre margin consists of a new TGS seismic line (Fig. 3a) and line M-M' (MB-6-92) in Blystad et al. (1995) to the southeast. Fig. 3b shows our geological interpretation of the Møre profile. The seismic line LOS 99-002 overlaps with this transect for ~25 km and runs from the Møre Marginal High to the extinct Aegir Ridge (Fig. 4). The deep crustal interpretation for this part of the profile is taken from Breivik et al. (2006).

The transect runs from the continental shelf to the Møre Marginal High and images the sedimentary strata in the Møre Basin and the typical lava flows and SDR approaching the continent-ocean boundary in the northeast. This profile can be subdivided into structural highs separating the sedimentary depocenters from southeast to northwest, these are: Giske High, Ona High and Vigra High. In addition to the main highs the new data show narrow and deep sub-basins, each of around 10 km width and up to 14 km deep between the Vigra and Ona highs. Small basins bounded by rotated fault blocks can be identified in the northeast whereas the normal faults dip mainly towards northwest. Several sill intrusions within the Lower Cretaceous sequence are present

Fig. 4. a) Seismic line LOS 99-002 with interpretation of basement from (Breivik et al., 2006). b) Combined transect including sediments overlying oceanic crust. This transect is the stratigraphy input for the second part of the reconstruction study.

between km 100–150 (Fig. 3a). The eastern part the Møre Basin displays a basin shape with sedimentary sequences dipping to the east, whereas in the western part the Upper Cretaceous and Cenozoic sequences run nearly horizontal towards the Møre Marginal High.

In contrast to previous interpretations (Faereth and Lien, 2002; Gomez et al., 2004) which describe no evidence for tectonic activity in the Cretaceous, we observe normal faults in the western part of the transect that cut at least through the lowermost Cretaceous sediments, suggesting rifting or tectonic activity during Early Cretaceous time. As mentioned above large areas approaching the COB are covered with lava flows and sill intrusions, which makes interpretation of the western part of the profile difficult (0–70 km). Our interpretation in this part agrees well with published interpretation mainly based on wide-angle seismic data (Faleide et al., 2008; Kvarven et al., 2014; Nirrengarten et al., 2014).

3.2.1. Model setup

Two sets of models are investigated. The first set focuses on the structural evolution of the Møre Basin (Fig. 3), while in the second the whole transect to the Aegir Ridge (Fig. 4) is reconstructed to investigate the impact of the breakup and spreading in the modelling.

Commonly used pre-rift crustal thicknesses for the Norwegian margin are between 30 and 40 km (Gernigon et al., 2006; Rüpke et al., 2013; Theissen and Rüpke, 2010; Wangen and Faleide, 2008). The choice of the initial crustal thickness influences the stretching factors necessary to create a sedimentary basin of a given depth and we therefore investigate the effect of different initial crustal thicknesses, i.e. 30 to 40 km. All input parameters and material properties are listed in Table 1. Additionally we introduce initial crustal heterogeneities and lower crustal bodies at various locations to fit the observed gravity and OBS data. The details of these models are described in the gravity section below. We assume the following three rift phases for our reference setup:

(1) Permian/Early Triassic rifting from 270 to 250 Ma to account for

the early stretching episodes, prior to the main phase during
(2) Late Jurassic/earliest Cretaceous (170–145 Ma).
(3) A final rift phase starting in Late Cretaceous (80 Ma) until breakup at 55 Ma.

We also test a longer second rift phase extending until Aptian time (115 Ma), which is observed at the central and outer Vøring Margin. Due to the number of ages applied to the sedimentary layers in our stratigraphy this alternative scenario leads to a Jurassic/Cretaceous rift phase from 170 to 115 Ma.

The temperature is fixed to 5 °C at the seafloor and 1300 °C at the boundary between lithosphere and asthenosphere. The lithosphere-asthenosphere boundary is considered to be the isostatic compensation depth and is at 125 km in all model setups. Radiogenic heat production in the crystalline crust exponentially decreases (e-fold length = 20 km) with depth from 2 $\mu\text{W}/\text{m}^3$ in the upper crust and 0.18 $\mu\text{W}/\text{m}^3$ in the lower crust. The effective sediment conductivities are computed as the geometric average between matrix and pore fluid conductivities.

Serpentinisation can potentially occur in all our model setups. The requirement is that high stretching leads to a complete embrittlement of the crust (Rüpke et al., 2013), which allows us to test if the LCB observed along the Møre profile could be serpentinised mantle and how the initial configuration influences the possibility of serpentine formation.

The second setup is the extended profile shown in Fig. 4 that runs to the Aegir Ridge and the reconstruction focus on the extension of the Norwegian part of the Norway-Greenland rift system. Material properties, rift phases and initial crustal thicknesses described above are the same in both setups. Breakup at 55 Ma is included and spreading in the Norway Basin is active until 28 Ma and produces 9 km thick oceanic crust. All models assume flexural isostasy with a necking depth of 15 km and an effective elastic thickness of 5 km.

3.2.2. Gravity modelling

Gravity modelling is performed to test different interpretation and

Table 1

Material properties of the crust, mantle, and stratigraphic layers.

		Matrix density (kg m ⁻³)	Thermal expansion (K ⁻¹)	Radiogenic heat (W m ⁻³)	Conductivity (W m ⁻¹ K ⁻¹)	Heat capacity (J kg ⁻¹ K ⁻¹)	Surface porosity (%)	Compaction length (km ⁻¹)
Mantle	Peridotite	3340	3.2×10^{-5}	0	3.5	1000		
Lower Crust	Diabase	2850	2.4×10^{-5}	1.8×10^{-7}	2.6	800		
Upper Crust	Granite	2750	2.4×10^{-5}	2×10^{-6}	2.6	800		
Quaternary	Sandstone	2650		7×10^{-7}	3.95	855	41	0.31
Neogene	Sandstone	2700		7×10^{-7}	3.95	855	41	0.31
Palaeogene	Shale	2700		2×10^{-6}	1.64	860	70	0.83
Upper Cretaceous	Organic rich shale	2610		2×10^{-6}	1.54	880	70	0.83
Lower Cretaceous	Shale	2700		2×10^{-6}	1.64	860	70	0.83
Upper Jurassic	Sandstone	2720		7×10^{-7}	3.95	855	41	0.31
Lower Jurassic	Marl	2700		1×10^{-6}	2	850	50	0.5
Triassic	Siltstone	2710		1×10^{-6}	2.01	940	55	0.51
Permian	Sandy shale	2700		1.71×10^{-6}	1.84	860	65	0.83
Inner LCB		3300	2.4×10^{-5}	1.8×10^{-7}	2.6	800		
Outer LCB/magmatic underplate		3100	2.4×10^{-5}	0	2.6	1000		

possible locations of the known LCBs that are used as input parameter for the reconstruction modelling. TecMod's gravity modelling is based on the computation of the gravitational acceleration due to a 2-dimensional body described by Talwani et al. (1959) and the reformulated algorithm by Won and Bevis (1987).

TecMod's finite-element mesh is used to calculate the gravity anomalies. This method allows for computing the changes in gravity anomaly along a reconstructed profile, which can then be compared to measured gravity anomaly changes. We compare observed free-air gravity anomaly data provided by TGS, which are comparable to the Word Gravity Map (WGM2012) that is freely available from the Bureau Gravimétrique International (BGI) <http://bgi.omp.obs-mip.fr/> (The International Gravimetric Bureau, 2012) with our modelling results for the profile across the Møre Basin (Fig. 3).

The gravity comparison consists of three steps of modelling: (1) the profile (Fig. 3b) is reconstructed with TecMod as described above to compute the necessary stretching factors to create the Møre Basin. This setup does not include any initial heterogeneities or lower crustal bodies. (2) The actual gravity comparison is performed with TecMod's forward modelling scheme due to the faster computation compared to the reconstruction. Stretching factors resulting from step 1 are used as input parameters. In this step we introduce high-density bodies to mimic the lower crustal body. The initial heterogeneities together with the LCBs give a close match with the observed gravity anomalies. In the last step (3), we run again the full basin reconstruction including the LCB and the heterogeneous initial crust to verify the results from the initial reconstruction. We test different densities and locations for an inner and outer LCB based on our interpretation and suggested in various publications (e.g. Kvarven et al., 2014; Nirrengarten et al., 2014). The density is 3300 kg/m³ and 3100 kg/m³ for the inner and outer LCB, respectively. With this assumption we do not specify the nature of the LCBs, but the LCBs are treated as crustal heterogeneities. In an alternative setup the outer LCB is treated as magmatic underplate coming into the model at 55 Ma. Additionally, it is assumed that the tectonic evolution at the Norwegian margin before mid-Permian (the starting age in our model) led to a heterogeneous crust where the initial thicknesses of the upper and lower crust vary horizontally. It should be noted that the heterogeneities and the LCBs do not change the total initial crustal thickness; it changes the density distribution throughout the crust due to varying proportions of upper and lower crust, and the implemented higher density LCBs/underplate. The total initial thickness is varied between 30 and 40 km.

The best fit of the gravity data is achieved with an inner LCB between 565 and 615 km with a maximum thickness of 4.5 km was needed to fit the gravity observations in the south-eastern part of the transect (Fig. 5). The outer LCB or alternatively magmatic underplating

occurs from the north-western start of the profile to the intrabasinal low between Vigra and Ona High. A thicker magmatic underplate is needed to fit the gravity data, compared to the outer LCB as part of the lower crust.

4. Modelling results

4.1. Stratigraphy, well comparison and crustal configuration

All reconstructions were run for 30 iterations and the input stratigraphy is fitted well for all models. To calibrate the results we compare computed temperature and maturity with available measurements from wells (Fig. 6). Vitrinite reflectance data (%Ro, Sweeney and Burnham, 1990) as indicator for maturity is sensitive to the temperature-time evolution and therefore well suited to verify the thermal solution of the reconstruction. For four wells temperature and vitrinite reflectance data were available: well 6302/6-1 and well 6403/10-1 in the Møre Basin are located ca. 30 south and 24 km north of the transect, respectively. The other two boreholes are also ca. 30 km north of the profile and have been drilled in the Slørebotn Subbasin, which can explain the local mismatch in temperature data. Nevertheless the overall computed temperature and maturity data match the present-day measurements well (Fig. 6), which confirms that the modelled thermal evolution (temperature and heat flow through time) provide valid solutions and it constrains the thermal parameter choice (radiogenic heat production, sediment conductivities).

In addition to the stratigraphy comparison we compare the computed crustal configuration with results inferred from our seismic data and potential field modelling (Faleide et al., 2008; Kvarven et al., 2014; Nirrengarten et al., 2014). At the end of the reconstruction the crust in the central basin is considerably thinned. The best fit is achieved with initially 32–35 km thick crust, also regarding the depth of the Moho, which lies in 20 km depth beneath the sag basins in the center and shallows to ~17 km towards the north-western end of the profile (Fig. 5).

Although the reconstruction only aims at matching the basin stratigraphy by pure shear deformation without considering faults in the crust, the resulting crustal thicknesses and Moho depth are comparable to interpretations from potential field modelling and wide-angle seismic data. The discrepancies in LCB thickness are also due to the modelling focus on sedimentary basin reconstruction, but as shown below, the consideration of such heterogeneities is important for the calculation of stretching factors and therefore the estimation of extension at continental margins.

Fig. 5. Crustal configuration after basin reconstruction and gravity comparison. a) The modelled (black line) free-air gravity anomaly fits well the measured data (red line). b) Resulting crustal thickness and stratigraphy of the reconstructed transect with total initial crustal thickness of 32 km. The locations of the inner and outer LCBs are constrained by gravity comparison and implemented as initial heterogeneities that change the density distribution of the crust. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Fig. 6. Basin temperature and comparison of observed and modelled data for the four wells projected onto the transect (Fig. 5). The top left figure (a) shows the modelled basin temperature and the period boundaries (black lines) from Quaternary to Triassic. The left plot of each well (b–e) shows observed (dots) and modelled (solid line) vitrinite reflectance (Easy Ro%). Temperature comparison is shown in the right panels. Modelled values are displayed as black line and observed data as diamonds.

Fig. 7. a) Total stretching and stretching factors for each rift phase along the transect for a setup with 35 km initial crustal thickness. Most thinning occurred prior to the final rifting which accounts for ~35% of total stretching. Highest stretching occurs at the depocenters. b) Comparison of total stretching factor from structural reconstruction for 30 (black line), 32 (red lines) and 40 km (grey line) initial crustal thickness. Higher stretching factors are needed if a LCB is considered in the reconstruction modelling. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

4.2. Stretching factors and extension

The second (breakup) setup is used to investigate stretching factors between Norway and the Aegir Ridge. The stretching factors calculated by the stratigraphy reconstruction are shown in Fig. 7. Fig. 7a shows the cumulative crustal stretching factor and stretching factors for each rift phases for an initial crustal thickness of 35 km. High stretching occurs at the deepest depocenters, e.g. on either side of the Vigra High with a maximum stretching factor $\beta = 3.3$ in the center of the basin after three periods of rifting. The highest thinning factors are computed in the north-western part at the breakup location, but it should be noted that the depth of the depocenter here is not well resolved due to the overlying volcanics. Stretching decreases towards the southeast with beta factors below 2. Major thinning took place during the first two rift episodes between Triassic and Early Cretaceous. The final rift phase that eventually led to breakup accounts only for ~35% of lithospheric stretching.

Assuming an initial crustal thickness of only 30 km the average β -factor is higher (3.9) with maximum values of 5 in the deep sub-basins. Lower values are computed for a thicker initial crustal thickness of 40 km, 2.2 and 2.7 for average and maximum stretching, respectively (Fig. 7b). Including LCBs as magmatic underplate or initial heterogeneities requires higher stretching factors compared to models that do not include any heterogeneities in the crust (Fig. 7b).

The computed stretching factors required to fit the input stratigraphy of all setups are lower than the critical value to favour brittle deformation and hence serpentinisation.

In the setup that includes the extended Cretaceous rift phase yield the same total stretching factors, only the distribution is different. Compared to the models where stretching ceased in the Early Cretaceous, these setups show slightly less stretching during the last and final rift event, but higher beta factors during the Jurassic/

Cretaceous (170–115 Ma) rift phase.

Crustal extension can be calculated by integrating crustal thinning (γ) along the profile length, where crustal thinning is defined by $\gamma = 1 - \frac{1}{\beta}$. The resulting extension is 188 ± 28 km (215 for 30 km crust and 160 for 40 km crust) for the models with an initial crustal thickness of 30–40 km.

5. Discussion

The results from new and reprocessed seismic data allow for detailed mapping of the Cretaceous and Cenozoic sedimentary successions in the Møre Basin. The major rift event during Jurassic/Early Cretaceous created several deep depocenters along the Møre Margin that are filled with thick Cretaceous and underlain by pre-Cretaceous sediments. The clear shallowing of the Base Cretaceous unconformity towards the Møre Marginal High and the shallow platform (Møre Marginal Plateau) highlighted by the new seismic data suggest that the crust is moderately thinned and that crustal fragments are still preserved and probably too thick to favour both mantle exhumation and serpentinisation to occur, which is in agreement with recent results from potential field modelling (Gernigon et al., 2015; Nirrengarten et al., 2014). Our stratigraphy reconstruction of the Møre transect supports this interpretation with a final crustal thicknesses of minimum 5–8 km (> 10 km in the vicinity of the inner LCB) and no serpentine formation. This result is in contrast to the suggestion that the inner LCB could represent serpentinised mantle (Lundin and Dore, 2011; Rüpke et al., 2013) because our seismic interpretation and stratigraphy calibration is different compared to those studies. The location of the outer LCB seems to correlate to the location of sill intrusions (Abdelmalak et al., under review) and we therefore prefer an interpretation for the outer LCB as magmatic underplate or magmatic intrusion into the preserved lower crust.

The thickness distribution of the Upper and Lower Cretaceous units (Fig. 3e,f) reveal that subsidence and sediment deposition shifted northwards towards the outer Vøring Margin after Cenomanian. The Møre Basin experienced subsidence and sedimentation mainly during the Early Cretaceous and less tectonic activity in Late Cretaceous, comparable to the observation by Scheck-Wenderoth et al. (2007), who suggested that fault activity is reduced in the Møre Basin during Late Cretaceous.

A comparison with previously published stretching factors and extension estimates is difficult because most studies focus only on post middle Jurassic rifting. Gomez et al. (2004) and Skogseid et al. (2000) estimated stretching and extension for the Møre margin by crustal thinning. Gomez et al. (2004) used only the upper crust for reconstruction and their results highly depend on the assumed pre-Cretaceous crustal thickness, i.e. the crustal thickness after earlier stretching episodes and it requires proper identification of the Base Cretaceous sediments and basement rocks. In the preferred model of Gomez et al. (2004) the pre-Cretaceous rift crustal thickness is already thinned to 20 km during earlier stretching episodes. Both studies suggest an extension of ca. 140–150 km since the middle Jurassic, which is comparable to the results of this study. However the distribution of extension is different, the final rift phase in our study only accounts for ca. 35%, which corresponds to ca. 60 km, whereas Skogseid et al. (2000) suggested 87 km for Maastrichtian/Palaeocene rifting and in an earlier study calculated an extension of 90 to 111 km during Late Cretaceous/Palaeocene (Skogseid, 1994).

Roberts et al. (2009) proposed increasing stretching factors from 1 (unstretched) at the Mid-Norwegian margin to up to 4 at the Møre Marginal High for breakup related stretching during Palaeocene from flexural backstripping. They assumed an average $\beta = 1.4$ for the Base Cretaceous. A study (Rüpke et al., 2013) using the same modelling approach, but different seismic data suggests very high stretching ($\beta > 8$) for the deep sub-basins for the rift phase(s) prior to breakup

Fig. 8. A comparison of stretching factors calculated with different methods for 35 km initial crust is shown. The comparison is calculated for the central Møre Basin only without breakup. The classical backstripping method results in lower stretching factors compared to the reconstruction due to the underestimation of sedimentation. See text for discussion.

related stretching. Our cumulative β (maximum 5 for 30 km crust) for these rift events is lower because we assume slightly shallower basins and apply the onset of the last rift phase during mid-Late Cretaceous and therefore do not need to create all the accommodation space for the Cretaceous sediments already during the Jurassic rift phase.

The determination of stretching factors and therefore also the estimated extension is not only dependent on the accuracy of geological interpretation, e.g. stratigraphy, crustal thickness or the assumed initial crustal thickness, but also highly affected by the method used to determine the stretching factors.

Fig. 8 shows a comparison of total stretching factors for the Møre transect calculated with three different methods for an initial crustal thickness of 35 km: the classical backstripping and decompaction approach (Steckler and Watts, 1978), stretching estimated by crustal thinning, and the computed β -factors from the reconstruction modelling. In the backstripping approach individual sediment packages are consecutively decompacted to recover the subsidence history (Steckler and Watts, 1978) and the stretching factors can be calculated from the resulting tectonic subsidence (Stewart et al., 2000):

$$\beta^{-1} = 1 - \frac{S_{\text{tec}}(\rho_m - \rho_w)}{t_{c0}(\rho_m - \rho_w)}, \quad (1)$$

where S_{tec} are total tectonic subsidence, t_{c0} the initial crustal thickness prior to rifting, and ρ_m , ρ_w , and ρ_c are densities of the mantle, water, and crust, respectively. Stretching factors derived from the subsidence analyses are usually lower and are likely to be underestimated because stretching/thinning of the sediments is not included (Theissen and Rüpke, 2010; Wangen and Faleide, 2008). In our case the calculated sediment thicknesses are up to 50–60% lower for certain sediment packages (e.g. between 115 and 145 Ma) if the sediments are only backstripped and decompacted but not thinned during the formation of the basin, compared to the reconstruction modelling where thinning of the sediments is included. Crustal stretching inferred by crustal thinning is poorly constrained because only the initial thickness is compared to the crustal thickness observed today and both values itself are subjected to large uncertainties. This method does not include any fault reconstruction or area balancing and often gives large discrepancies if compared to other methods (Ranero and Perez-Gussinye, 2010). The stretching factors are higher compared to the reconstruction modelling due to the thicker crust at the end of the reconstruction compared to what has been observed. Despite the discrepancies in crustal configuration and thickness of the LCB's between the our modelling results and the initial seismic interpretation the influence of implementing LCBs is seen in the computed stretching factors with higher stretching required in models that include crustal heterogeneities.

By applying an initial crustal thickness of 32–35 km thickness, compared to the even thicker (~40 km) crust observed close to the Norwegian mainland (Kvarven et al., 2014), we assume that the crust has already been partly thinned during the episodes of stretching prior to the Permian/Early Triassic rift phase. Despite all research, also driven by the hydrocarbon industry the mechanism that leads to highly thinned crust is not fully understood. Osmundsen and Ebbing (2008) suggested that the crust is thinned along large low-angle detachment faults that thin the crust from initial thicknesses of up to 40 km to < 10 km. This mechanism is compared to the 'thinning mode' (Lavie and

Manatschal, 2006) derived from the Iberian margin. Although we do not see large basin flank detachment fault in our profiles across the Møre Basin it is possible that low-angle normal faults play an important role in thinning the crust but are in our case overprinted by the thick sedimentary cover and sill complexes. This is also in agreement with Lavie and Manatschal (2006) who describe that the thinning mode often affects parts of the margin that are usually buried under thick sediments. However, we also find a comparison with the Iberian margin difficult because the major rift phase during Late Jurassic/Early Cretaceous, responsible for the highly thinned crust, most likely aborted during mid-Cretaceous and was not a continuous process leading to the Eocene breakup. An alternative model to stretch the crust has been suggested by Ranero and Perez-Gussinye (2010) who showed that very high thinning of the crust can be explained by normal faulting if the faults are active sequentially in time.

In plate reconstruction studies extension is quantified by the overlap of the continent-ocean boundary and has been estimated to be ~400 km between Greenland and the Mid-Norwegian margin, which corresponds to an average stretching factor of at least 2 (Torsvik and Cocks, 2005; Torsvik et al., 2001). Using recently published rift velocity (Brune et al., 2016) the total extension between Greenland and Norway at the location of our transect of ~440 km between 200 Ma and breakup. Our model data fit with the proposed stretching estimates derived from plate reconstruction studies (Brune et al., 2016; Torsvik et al., 2001), if we assume that about half the extension between the Norway and East Greenland takes place at the Møre Margin and the other half is distributed between the Jan Mayen Microcontinent and the conjugate East Greenland Margin.

Very high stretching factors of about 4–5 (maybe locally higher) characterize the deepest sub-basin. A more precise estimation of stretching and extension derived from stratigraphy reconstruction is only possible if sub-volcanic imaging/interpretation is improved and if deeper well data allow for better age constraints.

6. Conclusions

New and reprocessed seismic data from the Norwegian passive margin allow for better interpretation of the seismic stratigraphy in the central Møre Basin. To derive stretching and extension that occurred during several episodes of rifting we have reconstructed the Møre Basin using different methods. Gravity modelling supports two LCBs along the transect and suggests the existence of heterogeneous thickness of the upper and lower crust prior to the first applied stretching event during Permian/lower Triassic.

The reconstruction study fits well the present-day stratigraphy, temperature, and maturity data derived from boreholes. The key points of our study are:

- The central Møre Basin is a deep basin with thick post-rift sedimentary strata following the main rift episode during Late Jurassic/Early Cretaceous. The sedimentary infill has locally maximum depth of 10 s TWT (corresponding to 15–16 km). Cretaceous depocenters are concentrated along the western flank of the structural highs. A major shift of subsidence and deposition to the north towards the Møre-Vøring transition occurred during mid-Cretaceous.

- The estimation of extension highly depends on the method. Compared to the reconstruction modelling, stretching evaluated by crustal thinning gives very high values, while the classical backstripping approach yields lower beta factors due to the underestimation of sedimentation. Up to ~60% more sedimentary infill is necessary if the sediments are also stretched during reconstruction of the present-day stratigraphy. This leads to higher stretching factors compared to the decompaction/backstripping calculations.
- Our result does not support the interpretation of the LCBs as serpentinised mantle along most of the reconstructed profile. The observed and modelled crustal remains too thick for complete mantle exhumation and high rate of serpentinisation required to explain the P-waves values of the LCB. We prefer a model where the outer LCB is related to magmatic intrusion into the pre-existing inherited lower crust and/or magmatic underplating.
- The extension along the reconstructed profile across the Møre Basin is estimated to 188 ± 28 km. It is comparable to estimations from plate reconstruction studies. The main thinning phase occurred before the final and separate rift episode leading to breakup.

Acknowledgment

This work is part of the project OMNIS (Offshore Mid-Norway: Integrated Margin and Basin Studies, 210429/E30) funded by the Norwegian Research Council through its Centre of Excellence (project 223272, Centre for Earth Evolution and Dynamics (CEED)). We thank R. Myklebust from TGS for providing the seismic data set and First Geo and TGS for providing the depth converted profile. We further thank L. Rüpke for his suggestions and comments on the reconstruction modelling, and Per Terje Osmundsen and Yuriy Maystrenko for their constructive and helpful reviews.

References

- Abdelmalak, M.M., Andersen, T.B., Planke, S., Faleide, J.I., Corfu, F., Tegner, C., Shephard, G.E., Zastrozhnov, D., Myklebust, R., 2015. The ocean-continent transition in the mid-Norwegian margin: insight from seismic data and an onshore Caledonian field analogue. *Geology* 43, 1011–1014.
- Abdelmalak, M.M., Meyer, R., Planke, S., Faleide, J.I., Gernigon, L., Frieling, J., Sluijs, A., Reichart, G.J., Zastrozhnov, D., Theissen-Krah, S., Said, A., Myklebust, R., 2016a. Pre-breakup magmatism on the Vøring margin: insight from new sub-basalt imaging and results from Ocean Drilling program hole 642E. *Tectonophysics* 675, 258–274.
- Abdelmalak, M.M., Planke, S., Faleide, J.I., Jerram, D.A., Zastrozhnov, D., Eide, S., Myklebust, R., 2016b. The development of volcanic sequences at rifted margins: new insights from the structure and morphology of the Vøring Escarpment, mid-Norwegian Margin. *J. Geophys. Res. Solid Earth* 121, 5212–5236.
- Abdelmalak, M.M., Faleide, J.I., Planke, S., Gernigon, L., Zastrozhnov, D., Shephard, G.E., Myklebust, R., 2017. The T-reflection and the deep crustal structure of the Vøring Margin: offshore Mid-Norway. *Tectonics* (under review).
- Berndt, C., Planke, S., Alvestad, E., Tsikalas, F., Rasmussen, T., 2001. Seismic volcanotratigraphy of the Norwegian Margin: constraints on tectonomagmatic break-up processes. *J. Geol. Soc.* 158, 413–426.
- Blystad, P., Brekke, H., Faerseth, R.B., Larsen, B.T., Skogseid, J., Tørudbakke, B., 1995. Structural elements of the Norwegian Continental Shelf. Part II: the Norwegian Sea region. *Norw. Petrol. Direct. Bull.* 8.
- Breivik, A.J., Mjelde, R., Faleide, J.I., Murai, Y., 2006. Rates of continental breakup magmatism and seafloor spreading in the Norway Basin-Iceland plume interaction. *J. Geophys. Res. Solid Earth* 111.
- Brekke, H., 2000. The tectonic evolution of the Norwegian Sea Continental Margin with emphasis on the Vøring and More Basins. In: Nottvedt, A., Olausen, S., Torudbakken, B., Gabrielsen, R.H., Brekke, H., Birkeland, O., Skogseid, J. (Eds.), *Dynamics of the Norwegian Margin*. Geological Society, London, Special Publications 167, pp. 327–378.
- Brune, S., Williams, S.E., Butterworth, N.P., Müller, R.D., 2016. Abrupt plate accelerations shape rifted continental margins. *Nature* 1–4 (advance online publication).
- Dore, A.G., Lundin, E.R., Fichler, C., Olesen, O., 1997. Patterns of basement structure and reactivation along the NE Atlantic margin. *J. Geol. Soc.* 154, 85–92.
- Doré, A.G., Lundin, E.R., Jensen, L.B., Birkeland, O., Eliassen, P.E., Fichler, C., 1999. Principal tectonic events in the evolution of the northwest European Atlantic margin. In: Fleet, A.J., Boldy, S.A.R. (Eds.), *Petroleum Geology of Northwest Europe: Proceedings of the 5th Conference*. Geological Society, London, pp. 41–61.
- Faerseth, R.B., Lien, T., 2002. Cretaceous evolution in the Norwegian Sea - a period characterized by tectonic quiescence. *Mar. Pet. Geol.* 19, 1005–1027.
- Faleide, J.I., Tsikalas, F., Breivik, A.J., Mjelde, R., Ritzmann, O., Engen, O., Wilson, J., Eldholm, O., 2008. Structure and evolution of the continental margin off Norway and Barents Sea. *Episodes* 31, 82–91.
- Gabrielsen, R.H., Odinsen, T., Grunnaleite, I., 1999. Structuring of the Northern Viking Graben and the More Basin; the influence of basement structural grain, and the particular role of the More-Trondelag Fault Complex. *Mar. Pet. Geol.* 16, 443–465.
- Gaina, C., Gernigon, L., Ball, P., 2009. Palaeocene-Recent plate boundaries in the NE Atlantic and the formation of the Jan Mayen microcontinent. *J. Geol. Soc.* 166, 601–616.
- Geoffroy, L., Burov, E.B., Werner, P., 2015. Volcanic passive margins: another way to break up continents. *Sci Rep* 5, 14828.
- Gernigon, L., Ringenbach, J.C., Planke, S., Le Gall, B., Jonquet-Kolsto, H., 2003. Extension, crustal structure and magmatism at the outer Vøring Basin, Norwegian margin. *J. Geol. Soc.* 160, 197–208.
- Gernigon, L., Ringenbach, J.C., Planke, S., Le Gall, B., 2004. Deep structures and breakup along volcanic rifted margins: insights from integrated studies along the outer Vøring Basin (Norway). *Mar. Pet. Geol.* 21, 363–372.
- Gernigon, L., Lucazeau, F., Brigaud, F., Ringenbach, J.C., Planke, S., Le Gall, B., 2006. A moderate melting model for the Vøring margin (Norway) based on structural observations and a thermo-kinematical modelling: implication for the meaning of the lower crustal bodies. *Tectonophysics* 412, 255–278.
- Gernigon, L., Gaina, C., Olesen, O., Ball, P.J., Péron-Pinvidic, G., Yamasaki, T., 2012. The Norway Basin revisited: from continental breakup to spreading ridge extinction. *Mar. Pet. Geol.* 35, 1–19.
- Gernigon, L., Blischke, A., Nasuti, A., Sand, M., 2015. Conjugate volcanic rifted margins, sea-floor spreading and microcontinent: insights from new high-resolution aeromagnetic surveys in the Norway Basin. *Tectonics* 907–933.
- Gomez, M., Verges, J., Fernandez, M., Torne, M., Ayala, C., Wheeler, W., Karpuz, R., 2004. Extensional geometry of the mid Norwegian Margin before Early Tertiary continental breakup. *Mar. Pet. Geol.* 21, 177–194.
- Grunnaleite, I., Gabrielsen, R.H., 1995. Structure of the More Basin, mid-Norway continental margin. *Tectonophysics* 252, 221–251.
- Kvarven, T., Ebbing, J., Mjelde, R., Faleide, J.I., Libak, A., Thybo, H., Flueh, E.R., Murai, Y., 2014. Crustal structure across the Møre margin, mid-Norway, from wide-angle seismic and gravity data. *Tectonophysics* 626, 21–40.
- Lavier, L.L., Manatschal, G., 2006. A mechanism to thin the continental lithosphere at magma-poor margins. *Nature* 440, 324–328.
- Lundin, E.R., Dore, A.G., 1997. A tectonic model for the Norwegian passive margin with implications for the NE Atlantic: Early Cretaceous to break-up. *J. Geol. Soc.* 154, 545–550.
- Lundin, E., Dore, A.G., 2002. Mid-Cenozoic post-breakup deformation in the 'passive' margins bordering the Norwegian-Greenland Sea. *Mar. Pet. Geol.* 19, 79–93.
- Lundin, E.R., Dore, A.G., 2011. Hyperextension, serpentinization, and weakening: a new paradigm for rifted margin compressional deformation. *Geology* 39, 347–350.
- Mjelde, R., Raum, T., Kandilarov, A., Murai, Y., Takanami, T., 2009. Crustal structure and evolution of the outer More Margin, NE Atlantic. *Tectonophysics* 468, 224–243.
- Mosar, J., Eide, E.A., Osmundsen, P.T., Sommaruga, A., Torsvik, T.H., 2002. Greenland - Norway separation: a geodynamic model for the North Atlantic. *Nor. J. Geol.* 82, 281–298.
- Nirrengarten, M., Gernigon, L., Manatschal, G., 2014. Lower crustal bodies in the More volcanic rifted margin: geophysical determination and geological implications. *Tectonophysics* 636, 143–157.
- Olafsson, I., Sundvor, E., Eldholm, O., Grue, K., 1992. Møre Margin: crustal structure from analysis of Expanded Spread Profiles. *Mar. Geophys. Res.* 14, 137–162.
- Osmundsen, P.T., Ebbing, J., 2008. Styles of extension offshore mid-Norway and implications for mechanisms of crustal thinning at passive margins. *Tectonics* 27.
- Perez-Gussinye, M., Reston, T.J., Phipps Morgan, J., 2001. Serpentinization and magmatism during extension at non-volcanic margins: the effect of initial lithospheric structure. *Geol. Soc. Lond., Spec. Publ.* 187, 551–576.
- Peron-Pinvidic, G., Manatschal, G., Osmundsen, P.T., 2013. Structural comparison of archetypal Atlantic rifted margins: a review of observations and concepts. *Mar. Pet. Geol.* 43, 21–47.
- Planke, S., T., R., Rey, S.S., Myklebust, R., 2005. Seismic characteristics and distribution of volcanic intrusions and hydrothermal vent complexes in the Vøring and Møre basins. In: *Geological Society, London, Petroleum Geology Conference Series* 6, pp. 833–844.
- Ranero, C.R., Perez-Gussinye, M., 2010. Sequential faulting explains the asymmetry and extension discrepancy of conjugate margins. *Nature* 468 (294-U180).
- Reemst, P., Cloetingh, S., 2000. Polyphase rift evolution of the Vøring margin (mid-Norway): constraints from forward tectonostratigraphic modeling. *Tectonics* 19, 225–240.
- Ren, S.C., Skogseid, J., Eldholm, O., 1998. Late Cretaceous-Paleocene extension on the Vøring Volcanic Margin. *Mar. Geophys. Res.* 20, 343–369.
- Ren, S.C., Faleide, J.I., Eldholm, O., Skogseid, J., Gradstein, F., 2003. Late Cretaceous-Paleocene tectonic development of the NW Vøring Basin. *Mar. Pet. Geol.* 20, 177–206.
- Reynisson, R.F., Ebbing, J., Lundin, E., Osmundsen, P.T., 2010. Properties and Distribution of Lower Crustal Bodies on the Mid-Norwegian Margin. pp. 843–854.
- Roberts, A.M., Corfield, R.I., Kusznir, N.J., Matthews, S.J., Hansen, E.K., Hooper, R.J., 2009. Mapping palaeostructure and palaeobathymetry along the Norwegian Atlantic continental margin: More and Vøring basins. *Pet. Geosci.* 15, 27–43.
- Rüpke, L.H., Schmalholz, S.M., Schmid, D.W., Podladchikov, Y.Y., 2008. Automated thermotectonostratigraphic basin reconstruction: Viking Graben case study. *AAPG Bull.* 92, 309–326.
- Rüpke, L.H., Schmid, D.W., Hartz, E.H., Martinsen, B., 2010. Basin modelling of a transform margin setting: structural, thermal and hydrocarbon evolution of the Tano Basin, Ghana. *Pet. Geosci.* 16, 283–298.
- Rüpke, L.H., Schmid, D.W., Perez-Gussinye, M., Hartz, E., 2013. Interrelation between

- rifting, faulting, sedimentation, and mantle serpentinization during continental margin formation—including examples from the Norwegian Sea. *Geochem. Geophys. Geosyst.* 4351–4369.
- Scheck-Wenderoth, M., Maystrenko, Y., 2008. How warm are passive continental margins? A 3-D lithosphere-scale study from the Norwegian margin. *Geology* 36, 419–422.
- Scheck-Wenderoth, M., Raum, T., Faleide, J.I., Wjelde, R., Horsfield, B., 2007. The transition from the continent to the ocean: a deeper view on the Norwegian margin. *J. Geol. Soc.* 164, 855–868.
- Skogseid, J., 1994. Dimensions of the Late Cretaceous Paleocene Northeast Atlantic rift derived from Cenozoic subsidence. *Tectonophysics* 240, 225–247.
- Skogseid, J., Planke, S., Faleide, J.I., Pedersen, T., Eldholm, O., Neverdal, F., 2000. NE Atlantic continental rifting and volcanic margin formation. In: Nottvedt, A., Larsen, B.T., Gabrielsen, R.H., Olausen, S., Torudbakken, B., Skogseid, J., Brekke, H., Birkeland, O. (Eds.), *Dynamics of the Norwegian Margin*. Geological Society, London, Special Publications 167. pp. 295–326.
- Steckler, M.S., Watts, A.B., 1978. Subsidence of Atlantic-type continental-margin off New-York. *Earth Planet. Sci. Lett.* 41, 1–13.
- Stewart, J., Watts, A.B., Bagguley, J.G., 2000. Three-dimensional subsidence analysis and gravity modelling of the continental margin offshore Namibia. *Geophys. J. Int.* 141, 724–746.
- Sweeney, J.J., Burnham, A.K., 1990. Evaluation of a simple model of vitrinite reflectance based on chemical kinetics. *AAPG Bull. Am. Assoc. Pet. Geol.* 74, 1559–1570.
- Talwani, M., Eldholm, O., 1972. Continental margin off Norway: a geophysical study. *Geol. Soc. Am. Bull.* 83, 3575–3606.
- Talwani, M., Worzel, J.L., Landisman, M., 1959. Rapid gravity computations for 2-dimensional bodies with application to the Mendocino submarine fracture zone. *J. Geophys. Res.* 64, 49–59.
- The International Gravimetric Bureau, 2012. *IAG Geodesist's Handbook - Journal of Geodesy*. Springer, pp. 787–974.
- Theissen, S., Rüpke, L.H., 2010. Feedbacks of sedimentation on crustal heat flow: new insights from the Voring Basin, Norwegian Sea. *Basin Res.* 22, 976–990.
- Torsvik, T.H., Cocks, L.R.M., 2005. Norway in space and time: a centennial cavalcade. *Nor. J. Geol.* 85, 73–86.
- Torsvik, T.H., Van der Voo, R., Meert, J.G., Mosar, J., Walderhaug, H.J., 2001. Reconstructions of the continents around the North Atlantic at about the 60th parallel. *Earth Planet. Sci. Lett.* 187, 55–69.
- Tsikalas, F., Faleide, J.I., Eldholm, O., Antonio Blaich, O., 2012. 5 - The NE Atlantic Conjugate Margins, Regional Geology and Tectonics: Phanerozoic Passive Margins, Cratonic Basins and Global Tectonic Maps. Elsevier, Boston, pp. 140–201.
- Wangen, M., 1995. The blanketing effect in sedimentary basins. *Basin Res.* 7, 283–298.
- Wangen, M., Faleide, J.I., 2008. Estimation of crustal thinning by accounting for stretching and thinning of the sedimentary basin — an example from the Voring margin, NE Atlantic. *Tectonophysics* 457, 224–238.
- Wangen, M., Mjelde, R., Faleide, J.I., 2011. Modelling thermal transients from magmatic underplating—an example from the Voring margin (NE Atlantic). *Comput. Geosci.* 15, 771–788.
- White, R., McKenzie, D., 1989. Magmatism at rift zones: the generation of volcanic continental margins and flood basalts. *J. Geophys. Res. Solid Earth* 94, 7685–7729.
- Won, I.J., Bevis, M., 1987. Computing the gravitational and magnetic-anomalies due to a polygon - algorithms and Fortran subroutines. *Geophysics* 52, 232–238.

Paper III

Regional Structure and Polyphased Cretaceous-Paleocene Rift and Basin Development of the mid-Norwegian Volcanic Passive Margin

By

Dmitry Zastrozhnov, Laurent Gernigon, Iakov Gogin, Sverre Planke, Mansour M. Abdelmalak, Stéphane Polteau, Jan Inge Faleide, Ben Manton & Reidun Myklebust

Marine and Petroleum Geology, 115, 104269

III

Research paper

Regional structure and polyphased Cretaceous-Paleocene rift and basin development of the mid-Norwegian volcanic passive margin

Dmitry Zastrozhnov^{a,b,*}, Laurent Gernigon^c, Iakov Gogin^{d,f}, Sverre Planke^{a,b},
Mansour M. Abdelmalak^{a,b}, Stéphane Polteau^{b,g,h}, Jan Inge Faleide^a, Ben Manton^b,
Reidun Myklebust^e

^a Centre for Earth Evolution and Dynamics (CEED), University of Oslo, Norway

^b Volcanic Basin Petroleum Research (VBPR), Oslo Science Park, Oslo, Norway

^c Geological Survey of Norway (NGU), Trondheim, Norway

^d Ichtron Ltd, Northwich, Cheshire, UK

^e TGS, Asker, Norway

^f CGG Robertson, Llanrhos, UK

^g SurfExGeo AS, Oslo, Norway

^h Institute for Energy Technology (IFE), Kjeller, Norway

ARTICLE INFO

Keywords:

NE Atlantic
Passive margin
Mid-Norwegian margin
Møre and Vøring basins
Basin evolution
Hyperextension
Seismic interpretation

ABSTRACT

The Møre and Vøring basins of the mid-Norwegian volcanic passive margin are characterized by thick accumulations of Cretaceous to Paleocene sedimentary strata. They were formed during a series of Late Mesozoic–Early Cenozoic extensional events and represent vast underexplored areas with a limited number of wells. Recently, a new generation of long-offset 2D seismic reflection lines and 3D seismic data, together with new well data, has permitted a significant improvement in the regional understanding of the Møre and Vøring basins. This has enabled much better imaging of the deep Cretaceous subbasins and sub-basalt structures. In light of this significant data improvement, we performed a regional tectonostratigraphic synthesis of the pre-breakup development of the Møre and Vøring basins. We have interpreted eight regional Cretaceous and Paleocene horizons and constructed a series of structural and thickness maps. The new interpretations allow us to examine the sequential evolution of the Cretaceous to Paleocene sedimentary infill and to discuss its relationship to the deep crustal structures and regional tectonic events. We conclude that the long and polyphased development of the Møre and Vøring basins is partly controlled by deep-seated structural highs. We show that active deposition in the Early Cretaceous was mainly focused in the Møre Basin, while the main Cenomanian and subsequent Late Cretaceous–Paleocene depocentres developed principally in the Vøring Basin and migrated sequentially west towards the present continent–ocean boundary. We argue that the outer Møre and Vøring basins are likely underlain by a relatively thick continental crust compared to the inner part of the regional sag basin. In this setting our observations do not support evidence for a large zone of exhumed upper mantle, which has previously been proposed to have formed before magmatism and breakup.

1. Introduction

The transition from rifting to drifting is a fundamental geodynamic process within Earth Sciences (Wilson, 1966). Important issues include the mechanisms of sedimentary basin formation and the modes of lithospheric thinning that predate oceanic rifting (e.g. Lavier and Manatschal, 2006; Van Avendonk et al., 2009; Reston, 2009; Huismans and Beaumont, 2011; Brune et al., 2014). Generic models of rifting have difficulty explaining the diversity of continental margins worldwide. In

particular, under scrutiny are the impacts of different deformation modes, structural inheritance and magmatic additions during rifting and breakup (e.g. Chenin et al., 2015; Ligi et al., 2018; Simon et al., 2009; Petersen and Schiffer, 2016; Guan et al., 2019; Nirrengarten et al., 2020; Schiffer et al., 2019).

The mid-Norwegian margin is a well-studied example of a volcanic passive margin that illustrates the complexity of rifting, sedimentary basin development and breakup processes (Gaina et al., 2009; Gernigon et al., 2019). The formation of the mid-Norwegian margin is the result

* Corresponding author. Centre for Earth Evolution and Dynamics (CEED), University of Oslo, Norway.

E-mail address: zastrozhe@gmail.com (D. Zastrozhnov).

<https://doi.org/10.1016/j.marpetgeo.2020.104269>

Received 11 September 2019; Received in revised form 28 January 2020; Accepted 29 January 2020

Available online 05 February 2020

0264-8172/ © 2020 Elsevier Ltd. All rights reserved.

Fig. 1. An updated nomenclature map of the mid-Norwegian margin with pre-breakup structural elements. The fault pattern is mainly based on this study and modified from Blystad et al. (1995), Gernigon et al. (2003) and Zastrozhnov et al. (2018). Regional transects A-H (Fig. 8) are shown in light purple solid lines. Locations of seismic close-ups are shown in white solid lines. The ocean-continent boundary (OCB) is defined as the inner edge of the true oceanic crust. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

of a long history of Paleozoic and Mesozoic rifting in the NE Atlantic region (Doré et al., 1999; Tsikalas et al., 2012) that culminated with the rupture of the lithosphere in the Early Cenozoic (Talwani and Eldholm, 1977; Skogseid and Eldholm, 1989). The Møre and Vøring rifted margin segments of the mid-Norwegian margin are prolongations of the North

Sea and the Faroe-Shetland margin and preserve a series of N and NE-trending deep Cretaceous basins, flanked by paleo-highs, terraces and shallow platforms (Blystad et al., 1995; Brekke, 2000) (Fig. 1).

A characteristic feature of the rift-to-drift transition of the mid-Norwegian margin is its widespread volcanism that is part of the North

Atlantic Igneous Province (Saunders et al., 1997; Meyer et al., 2007; Abdelmalak et al., 2016a, 2016b; Millett et al., 2020). The volcanism characterizes it as a volcanic passive margin. The main characteristics of volcanic passive margins are: (1) thick volcanic wedges of seaward dipping reflector sequences (SDRs) emplaced along the proto-breakup axes (Hinz, 1981; Mutter et al., 1982; Eldholm et al., 1989; Berndt et al., 2001; Planke et al., 2017); (2) massive emplacement of sill and dyke intrusions in the sedimentary basins (Planke et al., 2005); and (3) presence of a high- V_p and high density lower crustal bodies (LCB) (Breivik et al., 2014; Mjelde et al., 2002, 2016; Abdelmalak et al., 2017).

Despite widespread consensus on these geophysical observations, an understanding of the rifted structure and evolution up to the onset of magmatic breakup remains incomplete and debated in the outer and distal parts of the mid-Norwegian margin, which are the Møre and Vøring basins. A particular point of contention is whether the final phase of extension of the mid-Norwegian margin evolved similarly to magma-poor margins in continuous extension from necking to breakup (Péron-Pinvidic et al., 2013; Péron-Pinvidic and Osmundsen, 2018), or if the mid-Norwegian margin evolved in a fundamentally different polyphased manner consisting of discrete pulses of extension and final magmatic weakening (Zastrow et al., 2018; Gernigon et al., 2019; Guan et al., 2019). In this context, the regional and geophysical understanding of the geometry of deep offshore sedimentary basins of the mid-Norwegian margin is important to constrain, so as to properly evaluate and determine the 3D structure and 4D evolution of the complex rifted system, and to what extent the mid-Norwegian margin can be compared to Iberian magma-poor type margins.

We use new and reprocessed long-offset 2D seismic reflection profiles and large 3D surveys covering the entire study area. In addition, we re-evaluated the biostratigraphy of released wells drilled in this under-explored area in order to calibrate and perform our regional seismic-to-well ties. This new dataset allows us to better visualize and confidently interpret the deep structures lying beneath the thick Cretaceous Møre and Vøring basins, although some of the deep structures remain poorly imaged below the breakup-related basalts. In turn, we better constrain the main phases of tectono-stratigraphic development and the structural geometries of the Møre and Vøring basins since the initial Late Jurassic-earliest Cretaceous rifting event until the onset of continental breakup. We therefore re-evaluate the timing of uplift of the structural highs and establish a series of new isopach maps to illustrate the spatio-temporal evolution of the main depocentres, associated with the sag-basins and structural highs. The new seismic data also reveal new structures within distal volcanic provinces. Together with extensive seismic interpretation, an updated chronostratigraphic event chart is presented in terms of the tectonostratigraphic evolution of the entire rifted margin. We find that the mid-Norwegian margin formed in a polyphased manner and represents a specific type of passive margin, where basin distribution and development are largely controlled by pre-existing crustal blocks and where the outer domain is underlain by a relatively thick continental crust.

2. Geological setting

The mid-Norwegian margin experienced several post-Caledonian extensional episodes, which ultimately resulted in the complete separation between Greenland and Norway in the earliest Eocene (Talwani and Eldholm, 1977; Brekke, 2000; Faleide et al., 2008). The large and thick Møre and Vøring basins of the mid-Norwegian margin formed during the Late Jurassic-earliest Cretaceous and Late Cretaceous-Paleocene rifting phases (Blystad et al., 1995; Brekke, 2000; Faleide et al., 2008). One additional extensional event was suggested for the mid-Cretaceous (Bjørnseth et al., 1997; Lundin and Doré, 1997; Pascoe et al., 1999; Gernigon et al., 2003; Roberts et al., 2009; Zastrow et al., 2018), although this period has also been considered tectonically quiet (Færseth and Lien, 2002; Lien, 2005; Færseth, 2012).

After the final rifting episode in the Late Cretaceous-Paleocene, continental breakup initiated and was accompanied with extensive and voluminous intrusive and extrusive magmatic activity (Eldholm et al., 2002; Planke et al., 2005; Abdelmalak et al., 2016a, 2016b). The submarine and subaerial volcanic sequences partially cover the Late Cretaceous-Paleocene and older structures which are likely to be present in the outer Møre and Vøring basins (Fig. 1).

The Møre Basin is limited to the east by the narrow pre-Cretaceous Møre Platform (Fig. 1). The southern limit of the Møre Basin is located where the NE-SW trending Møre-Trøndelag Fault Complex truncates the sub-meridional trends of the northern North Sea (Brekke, 2000). To the west, towards the ocean-continent 'boundary' (OCB), which we define here as the inner edge of the true oceanic crust, the Møre Basin is bordered by the wide and shallow, pre-Cretaceous Møre Marginal Plateau (Gernigon et al., 2015; Theissen-Krah et al., 2017; Manton et al., 2018; Millett et al., 2020). To the north, the transition to the Vøring Basin is defined by the Jan Mayen Corridor, which is interpreted to be a complex long-lived inherited crustal transfer system that indirectly influenced the segmentation of the oceanic domain (Gernigon et al., 2012, 2015; Nirrengarten et al., 2014). The Møre Basin is characterized by high rates of subsidence and sedimentation during the Early Cretaceous with less tectonic activity in the Late Cretaceous compared to the outer Vøring Basin. The major shift of subsidence and deposition to the north towards the Vøring Basin occurred during the Cenomanian (Theissen-Krah et al., 2017).

The Vøring Basin is limited to the east by the large Trøndelag Platform and the Halten and Dønna terraces (Fig. 1). The outer portion of the Vøring Basin most likely extends towards the proto-oceanic domain to the west. It is characterized by a series of structural highs (Gjallar Ridge, Nyk High, Grimm High, Skoll High) and associated subbasins (Hel Graben and Fenris Graben) formed during the Late Cretaceous-Paleocene rifting phase preceding the onset of breakup (Gernigon et al., 2003; Ren et al., 2003; Zastrow et al., 2018). The Bivrost Lineament separates the Vøring Basin from the narrow and elevated Lofoten-Vesterålen Margin to the north (Brekke, 2000; Maystrenko et al., 2017a; Tsikalas et al., 2019). The Fles Fault Complex, which entirely dissects the Cretaceous-Paleocene sedimentary package, is a NE-SW trending, elongated boundary fault system that splits the Vøring Basin into western and eastern parts (Blystad et al., 1995; Brekke, 2000).

3. Dataset and methods

3.1. Seismic data and interpretation

In this study, we used over 150,000 km of new and reprocessed vintage 2D seismic reflection lines acquired with a line spacing ranging from 0.2 to 2 km covering the entire mid-Norwegian margin (260,000 km²) (Fig. 2). This large dataset was time migrated and displays a positive amplitude at the seabed. Overall, the data are of good to excellent quality, allowing us to tie confidently seismic stratigraphic units.

In addition, we also fine-tuned our interpretation based on two recent large 3D seismic surveys acquired by TGS in 2017–2018 (Fig. 2). The Atlantic Margin North (AMN17) seismic survey covers c. 16,000 km² in the southern Vøring Basin. The Atlantic Margin South (AMS17-18) seismic data covers c. 31,000 km² in the Møre Basin.

Together with both 2D and 3D seismic data, we use an interpolated 3D seismic volume (e.g. Whiteside et al., 2013), called J-Cube MN, compiled by TGS from a dense grid of 2D and 3D seismic data on the mid-Norwegian margin. The advantage of using J-Cube includes the ability to better constrain basin-wide seismic-to-well ties by creating arbitrary and composite lines in any direction. The interpretation workflow is the same as for conventional 3D seismic volumes. However, the level of imaging quality of J-Cube is heavily dependent on the line spacing, and hence the quality varies locally.

Fig. 2. Geological and geophysical dataset used in this study: black dots – available exploration wells; solid light grey lines – 2D reflection seismic; polygons AMN17 and AMS17-18 – recently acquired large TGS 3D cubes; J-Cube MN – regional interpolated 3D volume. Bathymetry Data – Gebco 2014 grid merged with J-Cube seabed grid.

The regional seismic interpretation presented in this paper focuses on stratigraphic intervals from the Early Cretaceous to Near Top Paleocene (Fig. 3). The seismic interpretation was carried out using the IHS Kingdom suite, DUG Insight and Petrel software. Eight regional seismic horizons were picked and tied to the available well dataset and mapped out across the entire study area. These horizons are the Base Cretaceous Unconformity (BCU), mid-Albian (MA), mid-Cenomanian (MC), Near Top Turonian (NTT), intra-mid-Campanian (IMC), the Base Paleogene Unconformity (BPU), Near Top Paleocene (NTP) and Base Basalt (BP) (Figs. 3 and 4). A few additional Cenozoic horizons including Top Tare (TT) and Base Naust (BN) horizons were mapped locally along the profiles. The final grids of the interpreted regional horizons represent a merge from all 2D and 3D seismic datasets available in this study (Fig. 2).

For integration purposes, we used the potential field data as pseudo-horizons, which were loaded into the seismic projects by scaling the values to fit the two-way travel time of the seismic data. We used 50 km high-pass-filtered Bouguer gravity and magnetic data (Fig. 5), both of which were processed and filtered by TGS. This integrated seismic-gravity-magnetic approach (e.g. Abdelmalak et al., 2016b, 2017; Zastrozhnov et al., 2018) helps us to better constrain the crustal configuration below the thick sedimentary cover, and further identify poorly imaged deep-seated structural basement highs below breakup basalts and sill intrusions. High-pass filtering of the gravity data (Fig. 5a) enables the differentiation of regional gravity anomalies caused by shallow basement structures from density variations in the deepest crust and the upper mantle. Cut-off wavelengths of 50 km usually emphasize density contrasts with source depths of < 10 km. With the exception of the volcanic domains, the 50 km high-pass filtered magnetic data (Fig. 5b) provides information on deep-sourced anomalies often related to basement rocks which floor non-magnetic sedimentary basins. In the volcanic province, the magnetic data correlates very well with the different volcanic facies identified within the

seismic data.

3.2. Well data and correlation principles

All key wells used for the Møre and Vøring basins (Fig. 6) were reviewed and interpreted based on the data available in the Diskos National Data Repository. The established marker events were interpreted with the integration of recent zonation schemes for the mid-Norwegian shelf, Barents Sea, and onshore northeast Greenland. The local marker events may not provide a key to regional ties or precise dating but indicate events and surfaces of correlative value. For the rest of the available wells, we followed the lithostratigraphic formation tops as defined by the NORLEX group (Gradstein et al., 2010; <http://nhm2.uio.no/norlex>).

The stratigraphic interpretation of the key wells (see Fig. 6 and Appendix 1) is based on the identification of correlatable biostratigraphic zones together with log interpretations. The calibration of the biostratigraphic data against the wireline logs, lithostratigraphy and seismic data helped to constrain and link together the assemblage alterations to chronostratigraphic events. The different biostratigraphic zones are defined using the downhole occurrence of key marker taxa and changes in assemblage composition. Locally, reworked taxa were identified with the interpreted clastic sediments, and contaminated caved assemblages were identified based on taxa distribution patterns and the geological history of the basins.

The biostratigraphic zonation of Radmacher et al. (2015) was used as the reference regional biostratigraphic scheme, because it is the most recent link between the Norwegian and the Barents seas taxa. In addition, we used the zonation of Duxbury (2001) as calibration for Early Cretaceous events on the Norwegian Continental Shelf and North Sea basins, and of Radmacher et al. (2014) for tying the Late Cretaceous basins close to the Barents Sea. We further combined the zonations of Nøhr-Hansen (1993, 1996, 2012) for understanding the correspondence

Fig. 3. Tectonostratigraphic summary and segment map of the study area. Time scale after Cohen et al. (2013). Expansion of abbreviations is given in Appendix 2.

between palynological assemblages of the Barremian to Paleocene within the Greenland-Norwegian Seaway. We also used the zonation of Powell (1992) for Northwest Europe to constrain the overall Early Cenozoic biostratigraphic framework.

4. Results

4.1. Characteristics of the main regional seismic horizons

4.1.1. Base Cretaceous Unconformity (BCU)

The BCU (Fig. 7) forms the floor of the Møre and Vøring basins and the isochron map illustrates the first order configuration of the mid-Mesozoic rift system which was affected by regional extension during the Late Jurassic-earliest Cretaceous period. The BCU shows great regional variability that reflects the different structural position of the main basin elements as well as regional variations influenced by basin-scale tectonic, thermal and/or isostatic process and global sea level variation.

The BCU mostly represents a composite surface (e.g. unconformity complex) rather than a single erosional event (e.g. Kyrkjebø et al., 2004). The BCU is mostly defined as a clear onlap surface on the terraces as well as on the deep-seated structural highs in the basin province (Brekke, 2000, Fig. 4a). This structural criterion was the main factor in identifying BCU. In general, the reflection is mostly soft (negative amplitude), but varies depending on the character of underlying seismic facies (Fig. 4a). The BCU is difficult to interpret along deep basin axes where the surface appears as a paraconformity. Locally, the BCU is inconsistent between 2D seismic lines due to edge effects and poor migration along fault/detachment complexes, terraces, and deep-seated structural highs. The western limit of the mapped BCU horizon is located where extrusive and intrusive volcanic sequences are entirely masking the pre-breakup geology. Underlying pre-Cretaceous seismic facies vary from coherent to incoherent reflections. Overlying Lower to Upper Cretaceous successions represent coherent to semi-coherent and incoherent reflections of moderate to high amplitudes (Fig. 4a).

Numerous wells penetrated the BCU on the terraces (eg. on the Halten and Donna terraces) and platform domains, but no wells reached the BCU in the central and outer Vøring and Møre basins. Although wells 6603/5-1 S (Dalsnuten) on the South Gjallar Ridge (Fig. 6) and well 6608/2-1 S (Sverdrup) in the northern Utgard High targeted deep pre-Cretaceous sequences (> 5000 m b.s.l.), both were abandoned in Early to Late Cretaceous sequences, thus indicating that the BCU is much deeper than previously thought in interpretations pre-dating the drilling (Blystad et al., 1995; Swiecicki et al., 1998; Brekke, 2000; Skogseid et al., 2000).

4.1.2. Mid-Albian horizon (MA)

The mid-Albian seismic horizon is a strong hard reflection (positive amplitude) sandwiched between thick soft reflections (Fig. 4b). In general, the mid-Albian horizon is poorly imaged within domes, deep basins (e.g. the Rås and Træna basins, Vigrid Syncline), and below sill intrusions and basalts. The mid-Albian horizon terminates at the edge of the proximal terraces. The western extent of the mapped mid-Albian horizon is again limited by the overlying volcanic sequences. The overlying seismic facies are of high amplitude, mostly coherent to semi-coherent, continuous to semi-continuous reflections (Fig. 4b).

Numerous wells penetrated the mid-Albian unit along the eastern flank of the Vøring Basin in the Halten and Dønna terraces, but only one well (6608/2-1 S) reached this level in the basin province on the western flank of the Træna Basin in the northern Utgard High (Zastrozhnov et al., 2018). However, we cannot firmly assign the age of this horizon using these wells since common onlaps, truncations, and hiatuses suggest that the Lower Cretaceous section is condensed on the flanks of the terraces and structural highs, thus resulting in a poor age estimation for the top of the unit. The mid-Albian age of this horizon is instead based on well 6205/3-1 R, which was drilled in the central part of the

Fig. 4. Typical manifestation of the main regional seismic horizons mapped in this study: (a) – Base Cretaceous Unconformity (BCU), (b) – mid-Albian (MA), (c) – mid-Cenomanian (MC), (d) – Near Top Turonian (NTT), (e) – intra-mid-Campanian (IMC), (f) – Base Paleogene Unconformity (BPU), (g) – Near Top Paleocene (NTP). Seismic examples are from AMN17 and AMS17-18 3D cubes. Data courtesy of TGS.

Slørebotn Subbasin (Fig. 6). The Lower Cretaceous succession in this basin is well imaged on the seismic data, appears to be complete, and is less affected by tectonic and stratigraphic hiatuses. In well 6603/5-1 S on the South Gjallar Ridge (Dalsnuten prospect), the Norwegian Petroleum Directorate indicates an Early Cretaceous (Aptian) age for the deepest Cretaceous interval drilled in the outer Vøring Basin. However, our biostratigraphic re-evaluation of the drilling results (Fig. 6) shows that the well rather terminates at Cenomanian level. Another important biostratigraphic age estimation comes from the North Træna Basin in the Lofoten-Vesterålen Margin, where this horizon was reached by shallow boreholes 6710/03-U-01 and 6711/04-U-01 (e.g. Hansen et al., 1992; Henstra et al., 2016).

4.1.3. Mid-Cenomanian horizon (MC)

The mid-Cenomanian seismic horizon is a distinct hard reflection (Fig. 4c), also poorly imaged within the domes, highly faulted structural highs (South and North Gjallar ridges) and volcanic areas. In the deep

subbasins of the Møre Basin (Slørebotn, Runde, Vigra and Grip subbasins), this horizon represents locally a strong and double reflection similar to the typical appearance of the mid-Albian horizon (Fig. 4c). Overlying seismic reflections are variable from semi-coherent to incoherent facies of moderate to low amplitude in the deepest basinal area.

Numerous wells reached the Cenomanian interval on the Halten and Dønna terraces but are not always applicable for the basin-wide correlations due to their marginal setting. In the Vøring Basin only two wells are reported to reach Cenomanian level: well 6603/5-1 S (Dalsnuten) in the South Gjallar Ridge (Fig. 6) and well 6608/2-1 S (Sverdrup) on the Northern Utgard High (Zastrozhnov et al., 2018). However, due to complex faulting geometries within these structural highs and generally poor seismic imaging, this horizon cannot be confidently tied away from the wells into the deep basin setting. Like the mid-Albian, the age of the horizon was interpreted principally on the biostratigraphy results of well 6205/3-1 R in the Slørebotn Subbasin

Fig. 5. Potential field data for the mid-Norwegian margin. Transparent white polygons with dashed outline are main structural highs. (a) 50-km high-pass filtered Bouguer data. (b) 50-km high pass filtered magnetic data. Data courtesy of TGS.

(Fig. 6) and shallow boreholes in the Lofoten-Vesterålen Margin.

4.1.4. Near Top Turonian horizon (NTT)

The Near Top Turonian reflection is a distinct and mostly soft reflection that is sandwiched between relatively thick hard reflections (Fig. 4d). Seismic facies above are usually faulted, display moderate amplitudes that are semi-coherent to incoherent. Seismic imaging is degraded mainly within the Cenozoic domes (Helland-Hansen Arch, Ormen Lange Dome) and below the volcanic sequences.

Numerous wells intersect the Turonian interval on the terraces, but only a few wells reached this level in the basin province. Here, we used wells 6504/5-1 S, 6605/8-1, 6605/10-1 and 6403/6-1 in central basin settings to set a well-constrained seismic tie (Fig. 6). The Near Top Turonian horizon fits well with the top of the Lysing Formation. However, the biostratigraphic assemblages could not pin-point the top Turonian boundary, which is often complicated by the diachronous late Turonian-early Coniacian development of the Lysing sands along the mid-Norwegian margin (Gradstein et al., 2010, NORLEX; Fig. 6; see also Appendix 1).

4.1.5. Intra-mid-Campanian horizon (IMC)

The intra-mid-Campanian horizon is a distinct reflection when compared to the relatively incoherent unit above, although its character is highly variable, ranging from a strong hard to strong soft reflection (Fig. 4e). The intra-mid-Campanian horizon, where present and well defined, is generally the first coherent reflection below the Base Paleogene Unconformity. The intra-mid-Campanian horizon is usually highly faulted, and the seismic imaging of the unit is rather complex within the domes and below sill intrusions.

This horizon correlates well to the top of the Nise Formation in the outer Vøring Basin (Figs. 3 and 6), which has been reached by several wells in the Vøring and Møre basins (e.g. 6704/12-1; 6605/8-1; 6403/6-1, 6505/10-1, 6405/7-1). The intra-mid-Campanian age of this horizon is very well constrained and is expected to mark the onset of the

last rifting episode preceding the breakup of the mid-Norwegian margin (Kittelsen et al., 1999; Gernigon et al., 2003; Ren et al., 2003; Fjellanger et al., 2005; Gradstein et al., 2010, NORLEX; NPD, 2019).

4.1.6. Base Paleogene Unconformity (BPU)

The Base Paleogene Unconformity, or top Cretaceous horizon, is defined by a regional unconformity, which mostly appears as a clear and hard reflection (Fig. 4f). However, this reflection cannot be traced confidently to the west into the volcanic domain but can be interpreted tentatively below thin basalt flows. Seismic facies above the BPU have coherent reflections in the Paleocene succession.

This horizon represents an erosional surface along the flanks of the Møre and Vøring basins (Brekke, 2000; Gjelberg et al., 2005). However, our biostratigraphy analysis also suggests a continuous stratigraphic development throughout the Late Cretaceous and early Paleocene in the central and deepest parts of the Vøring (6705/10-1, 6603/12-1, and 6505/10-1; Fig. 6), and Møre basins (6302/6-1 and 6403/6-1; Fig. 6; see Appendix 1). This is in agreement with the interpretations of Hjelstuen et al. (1999) and Gjelberg et al. (2005).

4.1.7. Near Top Paleocene (NTP)

The Near Top Paleocene horizon represents a hard high-amplitude reflection with a thick soft below (Fig. 4g). This horizon often appears as an onlap surface along the flanks of the Mimir High, Ygg High, Gjallar Ridge, Nyk High, and Hel Graben (Naglfar Dome). Overlying seismic facies are relatively continuous, parallel to sub-parallel and of high amplitude (Fig. 4g).

Our regional seismic-tie shows that the Near Top Paleocene corresponds to the HV1 horizon described at the well 6302/6-1 (Tulipan) in the Møre Basin (Kjoberg et al., 2017; Schmiedel et al., 2017). HV1 is a paleosurface showing numerous craters of uppermost hydrothermal vent complexes emplaced at the Paleocene-Eocene boundary. The horizon merges into the Inner Flows facies in the volcanic domain in the outer Vøring Basin (cf. Hjelstuen et al., 1999; and Planke et al., 2005),

Fig. 6. Regional correlation of the Cretaceous-Paleocene succession across the key wells in the Vøring and Møre basins. Correlations lines are the main seismic horizons: BCU - Base Cretaceous Unconformity, MA - mid-Albian, MC - mid-Cenomanian, NTT - Near Top Turonian, IMC - intra-mid-Campanian, BPU - Base Paleogene Unconformity, NTP - Near Top Paleocene. Blg - Blållange Formation, spg - Springar Formation, tg - tang Formation.

but extends over the Inner Flows in the southwestern Vøring Basin (cf. Hjelstuen et al., 1999), and onlaps onto the Faeroe-Shetland Escarpment in the outer Møre Basin.

The age of the horizon is not very well-constrained. According to the NPD nomenclature (Dalland et al., 1988), the top of the Paleocene corresponds to the top of the Tang Formation. However, based on the detailed biostratigraphic re-evaluation of the Tulipan well (6302/6-1), Kjoberg et al. (2017), showed that this horizon corresponds to the top of a “transition zone” between the Tang and Tare formations, which they estimated to be of earliest Eocene in age. Eventually, given these stratigraphic uncertainties, we name this horizon “Near Top Paleocene”. In order to illustrate the structural and basin configuration of the volcanic domain prior to breakup time, we merge the Near Top Paleocene and Base Basalt horizons northwest of the volcanic escarpments.

4.1.8. Base Basalt horizon (BB)

The Base Basalt horizon represents the base of the volcanic sequence, and as such is restricted to the outer Møre and Vøring basins. The Base Basalt horizon is characterized by a soft reflection with moderate amplitude (Fig. 4h) thus interpreted to be the downward transition from high-velocity/high-density hyaloclastites into pre-breakup sedimentary sequences. Interpretation of this horizon was commonly challenging and in some cases was aided by more subjectively picking of the horizon above the first occurrence of clear saucer-shaped sills emplaced into sub-basalt sediments. The nature of the Base Basalt surface is defined by the volcanic seismic facies above, ranging from submarine (Lower Series and Inner Flows), to coastal (Lava Delta) and subaerial (Landward Flows and Inner SDR) (e.g. Abdelmalak et al., 2016b).

Fig. 7. Time-structure map of the Base Cretaceous Unconformity (BCU). Landward extent of late Paleocene-early Eocene lavas is shown in solid red line. Note that the interpretation below the lavas is less confident due to poorer seismic imaging. Expansion of abbreviations used on the map is given in [Appendix 2](#). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

4.2. Structure of the Møre and Vøring Basins

In this section, we describe the configuration of the Møre and Vøring basins using a series of time-structure maps based on the integration of our BCU, Base Paleogene and Near Top Paleocene horizons, as well as regional transects and seismic close-ups. The updated mapping allowed us to define and image better a set of new structural

elements along the mid-Norwegian rift system, which is crucial to describe and present in order to form a basis for the updated regional tectonostratigraphic framework. The geometry of the BCU mostly reflects the regional architecture of the Vøring and Møre basins affected by Late Jurassic-earliest Cretaceous rifting. The geometries of the Base Paleogene Unconformity and Near Top Paleocene time-structure maps are used to characterize the configuration of the outer Vøring Basin,

(caption on next page)

Fig. 8. Regional time-transects across the Møre and Vøring basins. Location of the transects is shown on Fig. 1. Transects B-H are from conventional 2D seismic survey, while transect A is extracted from the J-Cube MN volume. Main detachment faults are shown in thick dashed red line. 50-km high-pass filtered Bouguer and 50-km high pass filtered magnetic data are displayed on each transect as pseudo-horizons in thick solid brown line and thick solid blue line respectively. Data courtesy of TGS. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

which was mostly affected by a distinct Late Cretaceous-Paleocene renewed rifting episode (Gernigon et al., 2003; Ren et al., 2003; Zastrozhnov et al., 2018).

4.2.1. Møre Basin

As shown in Fig. 7, in the Møre margin segment the BCU deepens from 2 to 2.5 s on the western flank of the Møre Platform down to 7.5–8 s in the Runde Subbasin. The western part of the Runde Subbasin is bounded by a NE-SW trending ridge, informally called the Ervik Ridge. The Ervik Ridge is a new 125 × 20 km NE-SW elongated structural element defined by a shallowing of the BCU level up to 4.5–5 s associated with positive gravity and magnetic anomalies (Figs. 1, 5, 7, 8a and 9). The Ervik Basin, which is located further to the west towards the present OCB, forms the basin on the NW flank of the Ervik Ridge (Figs. 1, 7, 8a and 9). We interpreted the Ervik Basin as a major supra-detachment basin controlled by west dipping low-angle faults active up to Albian time (Figs. 8a and 9).

The outer Møre Basin exhibits a shallow BCU level defined as the Møre Marginal Plateau (Figs. 1, 7 and 8a-d). The Møre Marginal Plateau extends from the Jan Mayen Corridor in the north down to the Brendan's Dome in the south. Further south in the UK sector, the Møre Marginal Plateau likely merges with the terraces on the eastern flank of the Faroe-Shetland Basin (Millett et al., 2020). The eastern flank of the Møre Marginal Plateau represents a large east-dipping monocline that is locally dissected by a set of west-dipping fault systems similar to those at the Rån Ridge in the southwestern Vøring Basin (Figs. 1, 7 and 8a-d). The faults are delimiting relatively small NE-SW oriented rotated fault blocks (e.g. the Bara and Hrann highs; see Figs. 1, 7 and 8b-c) and half grabens defined at the BCU level. Both the Bara and Hrann highs formed at the northeastern flank of the Møre Marginal Plateau and are observed at the edge of the Inner Flows.

Seismic interpretation within the shallow Møre Marginal Plateau is challenging due to the development of the Early Cenozoic lava flows and sill intrusions. Nonetheless, potential field data shows the presence of a prominent positive gravity anomaly along the western flank of the Møre Marginal Plateau (Figs. 5a and 8a-d). The new AMS17-18 3D data at this setting supports the presence of tilted fault blocks with internal layered units below the shallow BCU (Fig. 9). We refer and name this new sub-basalt structure as the Bylgja Ridge. The NE-SW strike of the Bylgja Ridge coincides with the strike of the Faroe-Shetland Escarpment, which developed just above the Bylgja Ridge. A deep-seated crustal reflection beneath the Bylgja Ridge follows the structural geometry of the ridge (Fig. 9). This deep crustal reflection has a similar form to the gravity anomaly likely contributing to its occurrence (Fig. 9). The western flank of the Bylgja Ridge is associated with a prominent west-dipping normal fault system that controlled the accommodation space of the adjacent sub-basalt Bylgja Basin (Fig. 1). These faults likely terminate in the Late Cretaceous-Paleocene sequence (Fig. 9).

Further north towards the Jan Mayen Corridor, the BCU defines NE-SW trending structural highs (the Kiran and Vigra highs) and adjoining subbasins (the East and West Kiran subbasins, and Vigra subbasins) in the Møre Basin (Figs. 1 and 7). These structural highs also correlate with strong positive gravity and magnetic anomalies (Fig. 5).

4.2.2. Jan Mayen Corridor and southern Vøring Basin: regional transfer domain

The most prominent manifestation of the Jan Mayen Corridor is observed at the BCU level as the sudden change of structural trend at the northern termination of the Vigra High, switching from a NE-SW

trend typical for the Møre Basin to a nearly N-S direction in the southern Vøring Basin (Figs. 1 and 7). The structural trends are defined by the Grip High and Slettringen Ridge, where the BCU shallows up to 4.5 s and 5 s respectively. Both structural highs correlate to positive gravity and magnetic anomalies (Figs. 5a and 8d-e), while the strike of the Jan Mayen Corridor is expressed by a gravity low (Fig. 5). These structures are adjacent to prominent subbasins, including the Holmen, Grip, Golma, and Slettringen subbasins, all characterized by deeper BCU levels (Fig. 7). Notably, the Holmen Subbasin and a small subbasin in between the Grip High and Slettringen Ridge have NE-SW trends typical for the Møre and central Vøring basins (Figs. 1 and 7). This trend is also prominent on the Halten Terrace (Fig. 1) and is expressed by a number of small graben structures (e.g. the Grinda Graben).

On the BCU map, a NW-SE lateral ramp between the Vigra and Grip highs is evident along the Jan Mayen Corridor (Fig. 7). This transfer zone seems to extend to the east and merges with a NW-SE segment of the Klakk Fault Complex at the southern termination of the Frøya High. Also, the northern part of the Slettringen Ridge strikes to the NE-SW repeating a dominant structural trend in the Møre and central Vøring basins and is visually segmented by a NW-SE dextral lateral ramp from a N-S trending part of the ridge to the south. This transfer zone coincides with the location of the Hevring Lineament newly defined in this study (Figs. 1, 5 and 7). This lineament extends to the east towards the Halten Terrace, where the Klakk Fault Complex is segmented at the southern termination of the Sklinna Ridge (Fig. 7). In between these transfer zones both the Slettringen and Golma subbasins have an apparent rhomboidal shape as depicted on the BCU map (Fig. 7).

Towards the north, we interpret the Hevring High (Figs. 1 and 7) as a horst structure characterized by a west-dipping detachment fault on its western flank. This fault controlled the creation of accommodation space for the Early to mid-Cretaceous Hevring Subbasin (Figs. 1, 7, 8e and 11). The eastern flank of the Hevring High corresponds to the east-dipping faults of the Fles Fault Complex (Figs. 1 and 9).

In the outer southern Vøring Basin, the rapid shallowing of the BCU is observed at the Kolga High (up to 3.8–4 s), which coincides with the western termination of the Jan Mayen Corridor and northern prolongation of the Møre Marginal Plateau (Figs. 1 and 7). The Kolga High is also a newly defined structural element characterized by a positive gravity anomaly (Figs. 5a, 8d and 10). Generally, the pattern of the Bouguer gravity anomalies corresponds to the geometry of a deep crustal reflection beneath the high (Fig. 10). The Kolga High has a very thin, even locally absent volcanic cover (Manton et al., 2018; Millett et al., 2020), and our seismic interpretation indicates that its structure is defined by a series of decoupled tilted fault blocks that developed both in pre-Cretaceous and Late (?) Cretaceous sequences (Fig. 10). The deepest structures of the Kolga High are reminiscent of those at the Rån Ridge located further north along the same regional trend (Figs. 1 and 7).

The Rån Ridge represents the most prominent BCU high in the southern Vøring Basin (Gernigon et al., 2003) and is the northern termination of the Jan Mayen Corridor (Fig. 7). It is characterized by a positive gravity anomaly (Figs. 5a, 8e and 11). The structure of the Rån Ridge is defined by fault blocks and detachment faults, which are capped by the BCU and likely detached along evaporite layers (of potentially Triassic age) (Fig. 11) or along similar intra-crustal weakness zones. Local Early to mid-Cretaceous normal faults on the western flank of the Rån Ridge control the accommodation space for the Rån Basin (Gernigon et al., 2003; Abdelmalak et al., 2017). The western flank of the Rån Basin is delimited by the small Dufa High (Figs. 1, 7 and 8e). The thin basalt flows covering the Dufa High did not degrade

Fig. 8. (continued)

significantly the resolution of the seismic data, allowing interpretation of a likely pre-Cretaceous tilted fault block below the BCU (Fig. 8e). Similar to the Rån Ridge, the Dufa High also corresponds to a clear positive gravity anomaly (Figs. 5a and 8e).

4.2.3. Central and northern Vøring basins

The change of the nearly meridional structural trends typical for the southern Vøring Basin to the dominant NE-SW direction in the central and northern Vøring Basin occurs just north of the Hevring High and Rån Lineament (Figs. 1 and 7). In the central and northern Vøring Basin, the BCU is much deeper than in the other segments of the mid-Norwegian margin and reaches a maximum depth at around 8–9 s which is observed in the Træna and Rås basins (Figs. 7 and 8f-h). There, the western flank of the basins is marked by a prominent chain of deep-seated NE-SW trending structural highs along the Fles Fault Complex, corresponding to the newly defined Hevring, Unn and Vimur highs, and the well-studied Utgard High (Figs. 1, 7, 8g-h and 11). All the highs are characterized by distinct gravity anomalies (Figs. 5a, 8g-h and 11) suggesting the presence of underlying basement highs. The average BCU depth along these highs is at approximately 6 s, with a minimum

depth of 4.5 s at the Utgard High.

Further to the northeast along the Fles Fault Complex, we interpret the Unn High, which is a small deep-seated horst structure (Fig. 1). Finally, the Vimur High is a deep-seated footwall block of the SW-dipping Fles Fault Complex (Fig. 8g).

Towards the northern Vøring Basin along the Surt Lineament, the northern Vimur High changes its strike to the E-W direction forming a lateral ramp in connection to the Utgard High (Figs. 1 and 7). Both southern parts of the Utgard and Nyk highs attached to the Surt Lineament are N-S oriented defining the rhomboidal geometry of the southern Någrind Syncline (Fig. 7). The wide and deep Vigrid Syncline forms the eastern flank of the entire Gjallar Ridge reaching maximum dimensions (Figs. 1, 7 and 8f-g) in between the North Gjallar Ridge and Vimur High.

4.2.4. Outer Vøring Basin

The BCU shallows towards the outer Vøring Basin, with areas above 6 s defining the NE-SW trending South and North Gjallar ridges and the previously described structural elements called the Skoll and Grimm highs (Figs. 7 and 8g; e.g. Abdelmalak et al., 2016b; Planke et al., 2017;

Fig. 9. A seismic line extracted from AMS17-18 3D volume showing configuration of the southwestern part of the Møre Basin and its transition to the Northern North Sea: (a) uninterpreted and (b) interpreted. Notably, the Near Top Paleocene horizon overlies the volcanic Inner Flows facies and onlaps onto the volcanic escarpment. White arrows on the uninterpreted profile show deep crustal reflection (T-Reflection?) in the outer Møre Basin below the Bylgja Ridge, whereas blue arrows highlight possible configuration of the BCU which caps deep-seated fault blocks with tilted reflections and syn-rift wedges. Note how gravity anomaly pattern mimics the configuration of the deep structures (BCU, deep reflections). Horizon abbreviation expansions are given in Fig. 4 and in section 3.1. The common legend for the main tectonostratigraphic units is shown on Fig. 8. G50 - 50-km high-pass filtered Bouguer anomaly pseudo-horizon; M50 - 50-km high pass filtered magnetic data pseudo-horizon. Location of the seismic line is shown on Fig. 1. Data courtesy of TGS. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

Fig. 10. A seismic line extracted from AMN17 3D volume showing configuration of the southern Vøring Basin and Jan Mayen Corridor: (a) uninterpreted and (b) interpreted. Our seismic tie from the basin part towards the Kolga High shows that it is primarily the Late Cretaceous-Paleocene structural high underlain by a deep-seated pre-Cretaceous block and lower (?) crustal dome expressed by the T-Reflection. Note how gravity anomaly pattern mimics the configuration of the deep structures (BCU, deep reflections). Horizon abbreviation expansions are given in Fig. 4 and in section 3.1. G50 - 50-km high-pass filtered Bouguer anomaly pseudo-horizon; M50 - 50-km high pass filtered magnetic data pseudo-horizon. The common legend for the main tectonostratigraphic units is shown on Fig. 8. White arrows on the uninterpreted profile show deep crustal reflection (T-Reflection?) in the southern Vøring Basin. Location of the seismic line is shown on Fig. 1. Location of the seismic line is shown on Fig. 1. Data courtesy of TGS.

Zastrozhnov et al., 2018).

The relatively shallow BCU highs below the North and South Gjallar ridges are clearly segmented at the Gleipne Saddle (Fig. 7). The North and South Gjallar ridges consist of a NE-SW trending system of rotated fault blocks eroded at the Base Paleogene level (Fig. 8f–g and 12–15) (e.g. Blystad et al., 1995; Gernigon et al., 2003; Ren et al., 2003). Notably in both structures along their southeastern flanks, we observe low-angle faulting, which affects deeper stratigraphic levels and is sealed by the mid-Albian level (Figs. 13 and 14). Furthermore, we observe Cenomanian-Turonian syn-extensional wedges at the western flank of the North Gjallar Ridge (Fig. 15). Both the southern and northern segments of the Gjallar Ridge correlate with a relatively low amplitude magnetic signature, but pronounced gravity anomalies (Figs. 5, 8f–g, 13 and 14), which are interpreted to originate mostly from the deep crustal T-Reflection. This deep, lower crustal and inherited structure controlled the observed segmentation of the outer Vøring Basin before the onset of breakup related magmatism (Gernigon et al., 2003, 2004; Abdelmalak et al., 2017).

In the northwestern part of the Vøring Basin, the Nyk High represents an uplifted and eroded footwall fault block of the collapsed Late Cretaceous-Paleocene Hel Graben (e.g. Lundin et al., 2013;

Zastrozhnov et al., 2018). The northern flank of the Hel Graben is defined by the Hel Terrace (Figs. 1 and 12), which is mostly characterized by a series of southward dipping growth faults developed in Upper Cretaceous-Paleocene strata and possibly detached within over-pressured Cretaceous mobile shales (Zastrozhnov et al., 2018).

Towards the OCB, the Skoll and Grimm highs correspond to elevated areas at the Base Basalt/Near Top Paleocene level (Fig. 16). The BCU configuration of the Skoll High remains uncertain due to poor sub-basalt seismic imaging (Figs. 7 and 8g). The sub-basalt Grimm High corresponds to a prominent positive gravity anomaly (Fig. 5a) which was previously attributed to a relatively shallow BCU high based on integrated seismic-gravity-magnetic interpretation (Abdelmalak et al., 2016b; Zastrozhnov et al., 2018).

4.3. Sediment thickness distribution

A series of updated isochore (time-thickness) maps (Figs. 17–19) shows the main subbasin trends and distribution, and inter-relationships with adjacent structural highs and terraces. The isochore maps are particularly useful to illustrate the tectonostratigraphic evolution of the Møre and Vøring basins from the Early Cretaceous to Paleocene.

Fig. 11. A seismic line extracted from AMN17 3D volume showing configuration of the Rån Ridge and adjacent subbasins in the southern Vøring Basin: (a) uninterpreted and (b) interpreted. Main detachment faults in the Rån Ridge and Hevring High could develop along the Permian-Triassic evaporites as proposed by Gernigon et al. (2003). Note how gravity anomaly pattern mimics the configuration of the deep structures (BCU, deep reflections). Horizon abbreviation expansions are given in Fig. 4 and in section 3.1. G50 - 50-km high-pass filtered Bouguer anomaly pseudo-horizon; M50 - 50-km high pass filtered magnetic data pseudo-horizon. The common legend for the main tectonostratigraphic units is shown on Fig. 8. Location of the seismic line is shown on Fig. 1. Data courtesy of TGS.

4.3.1. BCU to mid-Albian (unit K10)

The K10 thickness map (Fig. 17a) shows the depositional trends from the Base Cretaceous to the mid-Albian time. The maximum sediment thickness focused in the Møre and southern Vøring basins. Significant minimization of thicknesses occurred to the north of the Rån Lineament (Fig. 17a). Four individual depocentres formed in the NE-SW trending Runde and Slørebotn subbasins, the round-shaped East Kiran Subbasin, and the N-S trending Golma Subbasin. The main depocentres are structurally controlled by dominantly west-dipping detachment faults observed on the flanks of the adjacent deep-seated structural highs and main breakaways (Fig. 8a–e and 17a). Above these highs, the K10 sequence significantly thins or is absent. In addition, prominent thinning and pinch-out of the K10 sequence is also observed towards the Møre Marginal Plateau and along the Møre-Trøndelag Fault Complex, both of which were standing in a high position during this period.

4.3.2. Mid-Albian to mid-Cenomanian (unit K20)

During the mid-Albian to mid-Cenomanian period, the main depocentre of the unit K20 was located in the Rås Basin (Fig. 17b). This depocentre is confined to the northwest by the structural highs along the Fles Fault Complex, where the unit K20 is thinner. Its southwestern termination is confined by the Rån Lineament (Fig. 17b). In the outer

Vøring Basin, additional local depocentres of the unit K20 developed along the southeastern flank of the South and North Gjallar ridges (Fig. 17b). Thicker deposition is also observed in the adjacent Rån Basin and is controlled by a major west-dipping fault/detachment system partly decoupled from the basement structures (Fig. 8e; e.g. Gernigon et al., 2003, 2019). The Rån Ridge is characterized by a very thin package or even an absence of the K20 on its top. In the southern Vøring Basin and almost the entire Møre Basin, K20 shows a nearly uniform thickness (Fig. 17b) except at the basin flanks, above the deep-seated structural highs and along the Møre-Trøndelag Fault Complex where it gradually becomes thinner and locally pinches-out (Fig. 17b).

4.3.3. Mid-Cenomanian to Near Top Turonian (unit K30)

The unit K30 displays a nearly uniform thickness in the Møre Basin, with slight thinning and ultimate pinch-out towards the western and eastern basin flanks (Fig. 18a). The main depocentres of the unit K30 mostly developed in the Rås Basin and the Hevring Subbasin and appear to be wider (100 km wide) and less focused compared to the older K10 and K20 depocentres (width lower than 50 km). Thick deposits of the unit K30 are also observed at the southwestern Træna Basin (Fig. 18a). The unit K30 significantly thins along the northwestern flank of the Træna Basin (which is the Utgard High), whereas it thickens over the

structural highs along the Fles Fault Complex and further towards the Vigrid Syncline and Hevring Subbasin. To the west, unit K30 thins and pinches-out over the Rån and Gjallar ridges. The increased thickness of K30 is interpreted in the southern Vøring Basin over the Slettringen Ridge and adjacent areas (Fig. 18a).

The unit K40 was deposited between the Near Top Turonian and intra-mid-Campanian horizons. The main depocentre of the unit is observed in the Vøring Basin. The unit K40 shows a localized maximum thickness in the Rås and Træna basins (Fig. 18b), where it is confined in between the Rån and Bivrost lineaments (Fig. 18b). The thickest part of this large depocentre increases northeast towards the Træna Basin. An

Fig. 13. A seismic line showing configuration of the South Gjallar Ridge and adjacent basins: (a) uninterpreted and (b) interpreted. White arrows on the uninterpreted profile show deep crustal T-Reflection, whereas blue arrows highlight possible configuration of the BCU which floors a deep basin in the Vigrid Syncline and slightly shallows towards the South Gjallar Ridge. Note how gravity anomaly pattern mimics the configuration of the deep structures (BCU, T-Reflection). Note a prominent intra-Paleocene lava flow package which is developed in the Vigrid Syncline and reflects earlier stages of the breakup-related magmatism along the mid-Norwegian margin. Horizon abbreviation expansions are given in Fig. 4 and in section 3.1. G50 - 50-km high-pass filtered Bouguer anomaly pseudo-horizon; M50 - 50-km high pass filtered magnetic data pseudo-horizon. The common legend for the main tectonostratigraphic units is shown on Fig. 8. Location of the seismic line is shown on Fig. 1. Data courtesy of TGS. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

important secondary depocentre is also located in the Någrind Syncline between the Utgard and Nyk highs. The K40 unit thins along the Fles Fault Complex, Utgard High and Gjallar ridges. Compared to the Vøring Basin, the K40 unit deposition in the Møre Basin is nearly uniform with slight thinning and pinch-out towards the basin flanks (Fig. 18b).

4.3.5. Intra-mid-Campanian to Base Paleogene (unit K50)

The mid-Campanian-Maastrichtian main sedimentation axis moved north-westward from the Træna and Rås basins to the Någrind Syncline, where the main depocentre of the unit K50 is located (Fig. 19a). The main depocentre merges with a thinner one located in the Vigrid Syncline and Gleipne Saddle. Another distinct depocentre with thinner unit K50 developed in the Hel Graben, which is separated from the Vigrid Syncline by the Rym Fault Zone (Fig. 19a). The unit K50 also thins towards the Nyk High, the North and South Gjallar ridges, along the Fles Fault Complex and the Utgard High. Like the K40 unit, the southern Vøring and Møre basins are characterized by a relatively uniform thickness showing a gradual thinning towards the basin flanks (Fig. 19a).

4.3.6. Paleocene (unit P10)

A marked change in sedimentation patterns happened during the deposition of the unit P10 in the Paleocene (Fig. 19b). From the Late Cretaceous to Paleocene, the main depocentres migrated further north-westward toward the Hermod Basin, Hel Graben and likely to the sub-basalt Fenris Graben. The southwestern termination of the Hel Graben depocentre is clearly controlled by the Surt Lineament (Fig. 19b). The P10 unit is very thin and nearly absent along the entire Gjallar Ridge, the Nyk and Utgard highs. It was partly eroded during this period. The southern Vøring Basin is characterized by a small and nearly uniform thickness of the unit P10. In the Møre Basin prominent depocentres developed both on the western and eastern basin flanks, while in the central part of the basin the thickness of the P10 unit is small and does not show any significant fluctuations (Fig. 19b).

5. Discussion

5.1. Implications for the Møre and Vøring Basins evolution

This section aims to discuss and compare our observations in the context of the published regional models of the development of the

Fig. 14. A seismic line showing configuration of the North Gjallar Ridge and adjacent basins: (a) uninterpreted and (b) interpreted. White arrows on the uninterpreted profile show deep crustal T-Reflection, whereas blue arrows highlight possible configuration of the BCU which floors a deep basin in the Vigrid Syncline and shallows towards the North Gjallar Ridge. Note how gravity anomaly pattern mimics the configuration of the deep structures (BCU, T-Reflection). Horizon abbreviation expansions are given in Fig. 4 and in section 3.1. G50 - 50-km high-pass filtered Bouguer anomaly pseudo-horizon; M50 - 50-km high pass filtered magnetic data pseudo-horizon. The common legend for the main tectonostratigraphic units is shown on Fig. 8. Location of the seismic line is shown on Fig. 1. Data courtesy of TGS. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

Fig. 15. A seismic example showing structural style of the North Gjallar Ridge. The syn-extensional growth strata and rollover anticlines are observed in the Cenomanian-Turonian interval on the western flank of the North Gjallar Ridge. Horizon abbreviation expansions are given in Fig. 4 and in section 3.1. Location of the seismic line is shown on Fig. 1. Data courtesy of TGS.

Fig. 16. Time-structure map of the Near Top Paleocene horizon merged with the nearly concomitant Base Basalt horizon in the outer mid-Norwegian margin. Merge location is a white solid line. Landward extent of late Paleocene-early Eocene lavas is shown in solid red line. Expansion of abbreviations used on the map is given in [Appendix 2](#). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

Møre and Vøring basins, which are summarized on [Fig. 20](#). Our detailed tectonostratigraphic framework is summarized on [Fig. 3](#).

5.1.1. Early Cretaceous stage: transition from syn-rift to post-rift or basin-wide extension?

The Early Cretaceous stage in the Møre and Vøring basins was preceded by a regional basin-wide Late Jurassic-earliest Cretaceous

ripping event. This event resulted in the formation of the deep Møre and Vøring basins which became distinct from the platforms and terraces ([Blystad et al., 1995](#); [Brekke, 2000](#); [Faleide et al., 2008](#)). Subbasins formed during this stage are characterized by large sediment thicknesses, especially in the Møre and southern Vøring basins, and their distribution is apparently controlled both by large west and east-dipping detachments on the bounding flanks of the major structural highs

Fig. 17. Time-thickness maps of seismic sequences K10 (a) and K20 (b) showing general distribution and migration of the basin depocentres in time and space. Present structural outline of the main structural highs is shown in dashed black line. Landward extent of late Paleocene-early Eocene lavas is shown in solid red line. Expansion of abbreviations used on the map is given in [Appendix 2](#). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

and terraces (Figs. 8 and 17a).

In the outer Møre Basin, the new imaging also supports the presence of an isolated elevated domain (the Møre Marginal Plateau). The distribution of depocentres and lateral pinch-outs of the Cretaceous sequences show that the distal sub-platform defined as the Møre Marginal Plateau was standing in a higher position and did not experience severe subsidence during most of the Cretaceous when compared to the central Møre Basin (Figs. 17–19). Likewise, a wide and shallow continental block also coexisted in the outer Vøring Basin at that time before its subsequent dislocation in the Late Cretaceous-Paleocene. The remnants of the dislocated marginal crustal block in the outer Vøring Basin are now represented as deep terraces (Fig. 1) and were mapped out there based on the proposed BCU configuration (Figs. 7, 13 and 14). The distribution of the K10 unit also suggests that major subsidence/sedimentation focused in the central Møre and southern Vøring basins during the Early Cretaceous (Fig. 17a). These domains coincide with the main thinning of the crust observed between the edge of the main terraces (e.g. the crustal necking) and the structural highs observed in the central axis of the Møre and southern Vøring basins (Kvarven et al., 2016; Maystrenko et al., 2017b).

We suggest that the major extensional event did not cease around the Late Jurassic-earliest Cretaceous transition in the Møre and Vøring basins, as was proposed earlier (Blystad et al., 1995; Brekke, 2000; Færseth and Lien, 2002; Lien, 2005). The Late Jurassic-earliest Cretaceous phase was characterized by intense regional stretching of the upper crust, whereas since the earliest Cretaceous and up to the mid-Albian the extension shifted from stretching to a thinning mode that was localized in the Møre and Vøring basins along the central basin high system (Figs. 21a and 22). The thinning mode typically shows very little or even no evidence for upper crustal extension and faulting in the

seismic sections (e.g. Lavier and Manatschal, 2006). During the thinning mode the extension is transferred to lower levels of the crust and is accommodated along the deep-seated shear/detachment zones localized along strong crustal bodies (Péron-Pinvidic and Manatschal, 2010), which seems to be the case for the central Møre and Vøring basins (Fig. 8). The existence of large basin-floor detachments in the Møre and Vøring basins was also earlier discussed by Osmundsen and Ebbing (2008) based on the analysis of sparse old 2D long-offset seismic lines. In this context, the definition of the BCU may not reflect the transition from syn-rift to post-rift stage of subsidence during the Jurassic–Cretaceous basin development (e.g. Færseth and Lien, 2002), but a shift in crustal deformation regime from a regional stretching type of deformation to a severe thinning regime. The idea of the Early Cretaceous extensional phase in the Møre and Vøring basins presented in this study is partly aligned with the regional models of Doré et al. (1999), Tsikalas et al. (2005), Lundin et al. (2013) and Péron-Pinvidic and Osmundsen (2018) (Fig. 20).

Concurrently, a renewed phase of Early Cretaceous (Neocomian) upper crustal stretching occurred locally in the outer Vøring Basin (Figs. 13, 14 and 21a), where normal faulting with significant displacement affected the western flank of the Rån Ridge (Gernigon et al., 2003), the proto-Gjallar Ridge and Vigrid Syncline, and the eastern flank of the proto-Nyk High (Zastrozhnov et al., 2018). It may reflect mild pulses of regional Early Cretaceous extensional reactivation documented in the NE Atlantic within the Faroe-Shetland Basin (Dean et al., 1999; Stoker, 2016; Schöpfer and Hinsch, 2019), NE Greenland (Whitham et al., 1999), the Lofoten-Vesterålen Margin (Henstra et al., 2016; Tsikalas et al., 2001, 2005, 2019) and the southwestern Barents Sea (Faleide et al., 1993). In the regional context, it is tempting to speculate that the propagation of the Faroe-Shetland Early Cretaceous

Fig. 18. Time-thickness maps of seismic sequences K10 (a) and K20 (b) showing general distribution and migration of the basin depocentres in time and space. Present structural outline of the main structural highs is shown in dashed black line. Landward extent of late Paleocene-early Eocene lavas is shown in solid red line. Expansion of abbreviations used on the map is given in [Appendix 2](#). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

extensional branch could also have potentially affected the proto-Jan Mayen Microplate Complex located between the outer Vøring and the Faroe Plateau before breakup (Fig. 21a). It is not also excluded that the hypothetical northern extensional systems from the Danmarkshavn and Thetis basins, and the documented one from the Lofoten-Vesterålen Margin (Henstra et al., 2016) could propagate to the south and overlap within the northern Vøring Basin extensional systems (Fig. 21a).

5.1.2. Cenomanian-Santonian stage: thermal sagging in the Møre Basin and increased subsidence and structurally controlled deposition in the central Vøring Basin

The mid-Cretaceous period (post-Albian) has often been interpreted as a period of relative tectonic quiescence (Færseth and Lien, 2002; Lien, 2005, Fig. 20). Færseth and Lien (2002) assumed that a regional increase of “post-rift” sedimentation took place both in the Møre and Vøring basins during most of the Early Cretaceous and during the Cenomanian-Santonian. Our study shows, however, that the subsidence rate is extremely variable between the Møre and Vøring Basin. The subsidence rate increased in the central Vøring Basin and is moderate in large parts of the Møre Basin (Fig. 18a–b). We interpret this difference in terms of tectonic regime variation. After the mid-Albian, the main thinning phase seemed to have failed in the Møre Basin and the whole central Møre Basin likely switched to a classic post-rift thermal subsidence mode (Fig. 21b) reflected by aggradational infill of the post-Albian sequences (Fig. 8a–h). Likewise, in the Vigrid Syncline, the minor mid-Albian-mid-Cenomanian depocentres followed local zones of Early Cretaceous stretching and reflect a transition to a thermal subsidence mode in that part of the basin.

Active sedimentation and subsidence in the Cenomanian to Santonian took place in the Rån and Træna basins and is restricted in

between flanking deep-seated highs and terraces (Fig. 18a–b). According to Færseth and Lien (2002) and Lien (2005), increased Cenomanian-Santonian subsidence in the Møre and Vøring basins is the result of thermal contraction of the lithosphere and an increased sediment input. These authors explain this sudden sediment load as a result of smoothing of the basin topography by the Cenomanian time, leading to subsequent redistribution of sediment fairways from the hinterland to the deep basins. An isostatic response due to the sediment load caused flexure and subsequent regional uplift and erosion of the basin flanks of the NE Greenland and mid-Norwegian margins (Lien, 2005). Both emerged areas were provenance regions for a sand-rich deep-water fan succession of the Lysing Member (Brekke et al., 2001; Fjellanger et al., 2005). This uplift and erosion of the basin flanks formed a pronounced Turonian/Coniacian unconformity (Brekke et al., 2001; Sømme and Jackson, 2013).

However, based on our regional mapping, the magnitude of the tectonic subsidence at that time in the Træna Basin is estimated to be in the order of 0.2–0.4 km/Myr (Zastrow et al., 2018). Such a localized and accelerated subsidence rate is too high to be explained by a classical thermal relaxation process (McKenzie, 1978), but is typical for rift or strike slip-basin regimes (Allen and Allen, 2013). Moreover, the short wavelengths of the deep and restricted Cenomanian-Santonian depocentres in the Rån and Træna basins (~30–50 km; Fig. 18) is difficult to reconcile with a large-scale topographic influence on the sedimentation rate and/or a large-scale flexure or cooling of the lithosphere without involving processes assuming a continental lithosphere instability. We propose that the thick Cenomanian-Santonian sedimentation is the result of a renewed, but local thinning episode in the Vøring Basin, which could be attributed to ductile boudinage of the underlying attenuated continental crust (e.g. Zastrow et al., 2018).

Fig. 19. Time-thickness maps of seismic sequences K50 (a) and P10 (b) showing general distribution and migration of the basin depocentres in time and space. Near Top Paleocene surface is merged with the Base Basalt horizon in the outer mid-Norwegian margin. Merge location is shown in solid white line on figure (b). Present structural outline of the main structural highs is shown in dashed black line. Landward extent of Late Paleocene-Early Eocene lavas is shown in solid red line. Expansion of abbreviations used on the map is given in [Appendix 2](#). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

Chronostratigraphy			Doré et al. (1999)	Brekke (2000)	Færseth & Lien (2002)	Tsikalas et al. (2005)	Lundin et al. (2013)	Peron-Pinvidic & Osmundsen (2018)	This study
Paleogene	Eocene	L Ypresian 56.0	Magmatism	Magmatism	Not Discussed	Magmatism	Magmatism	Magmatism	Magmatism
		L Thanetian 59.2	Uplift and Extension	Uplift and Transpression	Rifting	Rifting	Extension	Breakup/ Oceanization	Extension
		M Selandian 61.6					Compression		
		E Danian 66.0							
Cretaceous	Late	Maastrichtian 72.1	Extension	Rifting	Regional Subsidence	Subsidence	Not Discussed	Exhumation	Thinning
		Campanian 83.6							
		Santonian 86.3		Tilting of Basin Margins					
		Coniacian 89.8							
		Turonian 93.9							
		Cenomanian 100.5							
	Early	Albian 113.0	Not Discussed	Subsidence Minor Faulting	Thermal Subsidence	Rifting	Not Discussed	Hyperextension	Thinning
		Aptian 125.0							
		Barremian 129.4							
		Hauterivian 132.9	Extension						
		Valanginian 139.8							
		Berriasian 145.0							
J	Late	Tithonian 152.1	Extension	Main Rifting Phase	Rifting		Hyperextension	Thinning	Stretching

Fig. 20. Diagram comparing previously published tectonostratigraphic models of the Møre and Vøring basins and the model presented in this study. Note, that only major tectonic events are shown. The detailed tectonostratigraphic model, which is based on the results of this study, is shown on [Fig. 3](#).

Fig. 21. Schematic map of the distribution of main tectonic regimes acting in the mid-Norway-Greenland conjugate margins in the mid-Cretaceous (a) and in the mid-Cenomanian-Turonian (b) times. Expansion of abbreviations used on the map is given in Appendix 2.

and subsequently ductile lateral escape of the lower crust (Fig. 22). A flow through ductile lower crust away from the rift centre in the Rås and Træna basins towards the future breakup axis in the outer Vøring Basin may further explain the observed relatively quick (~10 Myr) rift jump (see [Tetreault and Buitier, 2018](#)). Such “depth-dependent” thinning mechanism may also explain the absence of prominent Cenomanian-Santonian syn-rift wedges and accompanying strong upper crustal faulting in the Rås and Træna basins.

In addition, the western flank of the Gjallar Ridge (Fenris Graben) experienced renewed phase of minor extensional activity in the Cenomanian-Turonian (Figs. 15 and 21b). The concomitant footwall uplift and thinning of the Turonian strata mark the first evidence of the

rise of the Gjallar Ridge (Figs. 13, 14 and 18a). This extensional system could propagate from the Thetis Basin in the north and extend to the Rån Basin/Dufa High in the south (Fig. 21b), where minor faulting affected Cenomanian-Turonian strata (Fig. 8e).

5.1.3. Late Cretaceous-Paleocene stage: towards break-up

Since the mid-Campanian, the main rift zone migrated towards the distal part of the margin and affected mostly the outer Vøring Basin. Meanwhile, the central parts of the Vøring and Møre basins were in the thermal subsidence mode (Fig. 23a), except local extensional reactivation along the Fles Fault Complex (cf. [Brekke, 2000](#)). Normal faulting developed in the Nyk High, the Grimm High ([Zastrozhnov](#)

Fig. 22. Schematic section illustrating configuration and development of the mid-Norwegian margin by the initial seafloor spreading time in the early Eocene. The LCB represents a mixture of exhumed high-grade Caledonian crust and breakup-related magmatic material.

Fig. 23. Schematic map of the distribution of main tectonic regimes acting in the Mid-Norway-Greenland conjugate margins in the end Cretaceous (a) and in the end Paleocene (b) times. Expansion of abbreviations used on the map is given in [Appendix 2](#).

et al., 2018), South and North Gjallar ridges, the western part of the Rån Basin. Here, dominantly west-dipping normal faults are present, partly cut by the Base Paleogene Unconformity (Fig. 8f–g, 12, 13, 14 and 15). At the same time, the western flank of the Hel Graben (Hel Terrace) was affected by faults with the opposite polarity (Ren et al., 2003; Zastrozhnov et al., 2018). The development of the main deep mid-Campanian-Maastrichtian depocentre in the Nâgrind Syncline (Fig. 19a) was a result of tectonic flexure and rapid subsidence due to the rising of the adjacent Nyk High as a rift shoulder (Zastrozhnov et al., 2018). Similarly, the footwall uplifts in the Gjallar Ridge system enhanced the subsidence and higher sedimentation of the adjacent and poorly faulted Vigrid Syncline (Fig. 19a). Based on these observations, we interpret the Late Cretaceous-Paleocene subsidence in the Vigrid and Nâgrind synclines as a result of extensional processes rather than a compressional inversion as proposed by Lundin et al. (2013) (Fig. 20). The Surt Lineament at that time acted as a high-relief accommodation zone (Ren et al., 2003) offsetting the rift zones between the Gjallar Ridge and the Nyk High/Hel Graben (Fig. 23a). The Surt Lineament also reflects a change in the deformation style separating a poorly faulted tectonic sag associated with a drastic thinning of underlying crust in the Hel Graben and a moderate stretching phase in the uplifted Gjallar Ridge characterized by numerous normal faults, syn rift wedges and possibly thicker crust.

The deformation pattern within the sub-basalt Grimm High to the north of the Surt Lineament is not easy to interpret. It could either form a common and connected extensional domain with the Gjallar Ridge system further south (Zastrozhnov et al., 2018), or alternatively could reflect propagation of the extensional system from the Thetis Basin in the north (Fig. 23a), which was already offset from the southern extensional branches of the Gjallar ridges by the Surt Lineament.

Later in the Paleocene in the central outer Vøring Basin, the main rift zone progressively migrated further to the OCB and localized in

the Fenris Graben (Fig. 23b). At the same time, the increased faulting activity on both flanks of the Hel Graben led to the collapse of the structure which was accompanied by high sedimentation rates (Fig. 19b). Zastrozhnov et al. (2018) suggested that the Paleocene rift axis in the Hel Graben failed to propagate further south across the Surt Lineament, and consequently jumped west of the Grimm High during the earliest Eocene at the onset of breakup in the Vøring Margin. This final rift jump was further supported by the onset of the magmatism along the nascent Vøring volcanic margin, which led to the additional weakening of the crust (Zastrozhnov et al., 2018).

Due to the volcanic cover, the interpretation of the deformational styles and time constraints in the outer part of the Møre and southern Vøring basins is somewhat uncertain. Lundin and Doré (2018) suggested the absence of Paleocene extension along the outer Møre Basin involving orthogonal transform regimes in the later breakup. However, in the regional context, lithospheric rupture without preceding Cretaceous-Paleocene extension is unlikely in the outer part of the Møre Basin. Like the outer Vøring Basin, a large part of the Møre Marginal Plateau and possibly the proto-Jan Mayen Microplate Complex was likely affected by significant Paleocene extension prior to breakup, but until now this large sub-basaltic domain (100–150 km) remained poorly imaged. Based on the new 3D seismic data, we interpret the Kolga High as a structural high primarily developed due to Late Cretaceous-Paleocene faulting (Fig. 10). We also expect occurrences of Late Cretaceous-Paleocene rifting at the western flank of the Bylgja Ridge (Bylgja Basin) to explain the rotated faulted structure observed just beneath the lava flows (Figs. 8a and 9). In such a scenario, the southern Late Cretaceous-Paleocene extensional branch could terminate at the Jan Mayen Corridor (Fig. 23b), which offsets it from the extensional branch in the outer Vøring Basin along the proto-Vøring transform margin. The proto-Vøring transform margin could act then as an accommodation zone, facilitating sediment transfer from NE Greenland to the adjacent

basins in the mid-Norwegian margin (e.g. Brekke et al., 2001).

Our seismic observations also support assumptions of Gernigon et al. (2019) about diachronous breakup of the mid-Norwegian margin, suggesting that the breakup and related magmatism along the Møre Margin initiated somewhat earlier in the Thanetian, compared to the earliest Eocene breakup in the Vøring Margin (Figs. 9 and 21b). The interpretation of the diachronous breakup-related magmatism is supported by the stratigraphic position of the Near Top Paleocene horizon over and within the Inner Flows facies in the Møre and Vøring respectively, as well as by the diachronous development of the vent complexes, where the lowermost complex observed in the upper Paleocene (Thanetian?) strata (e.g. Planke et al., 2005). Earlier pulses of sill intrusions are also described in the Hel Graben (Zastrozhnov et al., 2018).

5.2. Deep basin configuration and the role of crustal architecture and structural inheritance on the basin development: the need for a new model?

The configuration and nature of the crust flooring the Møre and Vøring basins is a matter of a recent scientific discussion. Several studies have suggested structural and crustal similarities with Iberian-type magma-poor passive margins, implying that large parts of the Møre and Vøring basins are underlain by serpentized/exhumed mantle domain (Reynisson et al., 2010; Lundin and Doré, 2011; Rüpke et al., 2013; Péron-Pinvidic et al., 2013; Péron-Pinvidic and Osmundsen, 2016, 2018; Osmundsen et al., 2016; Osmundsen and Péron-Pinvidic, 2018). Alternative studies advocate that there is no clear geological or geophysical evidence to support a broad zone of mantle serpentization/exhumation along the mid-Norwegian margin (Mjelde et al., 2002; Gernigon et al., 2004, 2006, 2015, 2019; Ebbing et al., 2006; Myer et al., 2013; Nirrengarten et al., 2014; Theissen-Krah et al., 2017; Abdelmalak et al., 2017; Zastrozhnov et al., 2018; Guan et al., 2019). Here we discuss our new results in the context of the aforementioned scenarios and how structural and lithospheric inheritance affects the distribution of deformation and sedimentation in the Møre and Vøring basins.

5.2.1. Crustal architecture of the outer Møre and Vøring basins

The crustal architecture of the outer Møre and Vøring basins is a key topic to understand the crustal and lithospheric setting before the onset of magmatism leading to the formation of volcanic margin and breakup. Our new data and structural interpretation support a clear shallowing of the BCU towards the OCB in the Møre Basin (Figs. 7, 8a-c and 9), where it is expressed by a wide sub-basalt crustal block called the Møre Marginal Plateau.

According to nomenclature of Lister et al. (1986), “a marginal plateau” represents a relatively undeformed, high-standing region bounded by large detachments faults. Although imaging of large detachments zones is complicated below volcanics, we believe that such “marginal plateau” was present along the outer Møre margin before the formation of the volcanic margin (Fig. 8a–b, e.g. Theissen-Krah et al., 2017). The later formation of the SDR sequence is proposed to be controlled by continent-ward dipping detachment faults (e.g. Geoffroy et al., 2015). This detachment fault system could act earlier together with an ocean-ward detachment fault system in the central basin province, which led to the individualization of the marginal plateau (Fig. 22).

Our structural observations also suggest that pre-Cretaceous sediments and relatively thicker and high-standing crustal blocks could be present within the Møre Marginal Plateau and on several deep crustal rafts of the Møre and Vøring basins (Fig. 8a–c and 9) (e.g. Nirrengarten et al., 2014; Gernigon et al., 2015; Theissen-Krah et al., 2017). The Møre Marginal Plateau likely represents an intermediate and relatively thick crustal and lithospheric domain in between the failed mid-Mesozoic inner rift system and the distal volcanic margin emplaced in a weaker lithospheric region (Fig. 22) (Gernigon et al., 2019; Guan et al.,

2019). Refraction data interpretation of Kvarven et al. (2016) corroborates the presence of elevated and thicker continental crust domain in the outer part of the Møre Basin. The presence of such shallow and buoyant crustal blocks may also explain the late and wide emplacement of the subaerial basalt in the Møre segment of the mid-Norwegian margin. A similar (continuous?) marginal plateau may have extended in the outer Vøring Basin before the mid-Late Cretaceous. However, compared to the Møre Marginal Plateau, the shallow Cretaceous domain, expected in the outer Vøring Basin, was further dislocated during the mid to Late Cretaceous–Paleocene rifting episode(s) (Fig. 8g–h, 13, 14 and 15). Based on structural modelling and refraction data, the existence of a wide continental block bounded from the west by a continent-ward dipping detachment fault system below the Vøring Marginal High was also earlier proposed by Walker et al. (1997) and Mjelde et al. (2003).

In the outer Vøring Basin the nature of the underlying crust has been extensively discussed in terms of the deep crustal reflection usually referred as the T-Reflection (Gernigon et al., 2003; Ebbing et al., 2006; Abdelmalak et al., 2017; Zastrozhnov et al., 2018). The T-Reflection usually coincides with the upper part of the high-velocity (7+ km/s) outer lower crustal body (LCB). In earlier models, the nature of the outer LCB has been mostly interpreted and discussed in terms of magmatic underplating concurrent to continental breakup (e.g., White and McKenzie, 1989). On the other hand, several amagmatic and partly magmatic models for the nature of the outer LCB in the Vøring Basin have been alternatively advocated including: 1) high velocity intrusions into the lower crust (White et al., 2008, 2010), 2) heavily serpentized/exhumed/denudated mantle (Ren et al., 1998; Osmundsen and Ebbing, 2008; Reynisson et al., 2010; Péron-Pinvidic and Osmundsen, 2016), and 3) retrograde/high-grade Caledonian lower crust (Gernigon et al., 2003; Ebbing et al., 2006; Mjelde et al., 2016; Petersen and Schiffer, 2016; Zastrozhnov et al., 2018.) possibly intruded by late magmatic sills and cumulates (Abdelmalak et al., 2017).

It was shown that the deformation of the T-Reflection in the outer Vøring Basin is decoupled from younger Late Cretaceous–Paleocene faulting (Gernigon et al., 2003; Zastrozhnov et al., 2018). This earlier deformation of the T-Reflection (Figs. 8e, 11, 13 and 14) implies that the main basement faults and high-velocity blocks were already present before final thinning of the crust and before the onset of major magmatism. Interpretation of a thin salt layer and local rollers (Triassic?) in the Rån Ridge also suggest that the T-Reflection already influenced the structural style of the ridge before and during the Cretaceous (Fig. 11). Accordingly, it is likely that recent interpretation of controversial intra-crustal/serpentized mantle detachment may instead simply represent a salt-related decollement system and not necessarily a crust/mantle coupled system. Moreover, if the T-Reflection represented mantle domes exhumed below the structural highs in the outer Vøring Basin, higher magnetic susceptibility and remanence induced by high degree of mantle serpentization would be expected there (e.g. Oufi et al., 2002), but the lack of prominent magnetic signature show that this is not the case there (Figs. 5b, 8e–h, 11, 13 and 14). According to the Iberian-type scenario (e.g. Péron-Pinvidic and Osmundsen, 2018), serpentization should occur during most of the Cretaceous period which is characterized by several polarity changes of the magnetic field. In this case we should expect higher regional values or at least a striped pattern of the magnetic signal. However, the entire Møre–Vøring basin province has generally low magnetic values (below or close to 0 nT) except in the central highs complex area where the thicker continental crust corresponds to a few positive magnetic anomalies. Gernigon et al. (2015) also showed that the thinnest part of the crust can be simply explained by a contact between the sediment and low magnetic lower crust and/or not serpentized mantle with no remanence and almost no susceptibility. Although the thermal effect of the proposed magmatic underplating in the outer mid-Norwegian margin could locally demagnetize the crust at the level of the T-Reflection just prior to breakup according to the models with large final volumes of magma (Gernigon

et al., 2006), the paleotemperatures from the wells in the outer Vøring Basin are better explained by numerical calculations without magmatic underplating (Fjeldskaar et al., 2009). Finally, according to Zastrozhnov et al. (2018), serpentinized mantle, if present, could be restricted locally within the subbasins in the northern Vøring Basin, an area that experienced extreme thinning rates throughout Mesozoic and Paleocene (Fig. 21) (i.e. Rås Basin, Træna Basin, Hel Graben).

Furthermore, the mid-Norwegian margin exhibits a significant time lag of almost 100 Myr between the necking event in the Late Jurassic-earliest Cretaceous and further lithospheric breakup in the late Paleocene-early Eocene. In contrast, the time lag between the necking and lithospheric breakup along the NE Atlantic magma-poor margins is markedly lower and is about 35–45 Myr (Chenin et al., 2015). So, if the Iberian magma-poor model worked along the mid-Norwegian margin, the breakup conditions would have been likely achieved by the Aptian-Albian time along the axial part of the Møre and Vøring basins. While along the magma-poor margins the necking to lithospheric breakup deformation period represents a continuous and regional thinning phase, along the mid-Norwegian margin the stretching/thinning phases were episodic (Figs. 3 and 20) and migrated both laterally along the margin and towards the future breakup axis (Figs. 21 and 23).

Along the mid-Norwegian margin, the volcanism and breakup developed when the environment was shallow marine to subaerial (e.g. Abdelmalak et al., 2016b; Millett et al., 2020). In contrast, continental breakup along the Iberian magma-poor margin occurred in deep-water conditions (Mohn et al., 2015). Mohn et al. (2015) estimated paleo-water depth during a final rifting stage along the Galicia segment of the Iberian margin to be in the range of 1–3 km. Thus, to explain the rapid uplift of heavily altered serpentinized mantle in the distal part of the mid-Norwegian margin to facilitate shallow marine and subaerial emplacement of basalts an additional exterior force would be required. Ren et al. (1998) favored the Iberian model for the crustal configuration of the outer Vøring Basin and suggested late Paleocene uplift as much as 1.2 km before breakup due to tectonomagmatic factors including underplating. Indeed, the uplift of this magnitude could be compatible with subaerial environment during late stages of rifting and initial seafloor spreading. However, later paleotemperature analyses in the outer Vøring Basin (Fjeldskaar et al., 2009) did not favor underplating of a magmatic body emplaced during the Early Cenozoic. Moreover, geochemical study of breakup-related volcanics showed that magma mixing between asthenospheric and upper crustal melts must have occurred prior to their emplacement within the Vøring and Møre Marginal highs (Meyer et al., 2009; Abdelmalak et al., 2016a; Bakke, 2017). Also, in their numerical modelling, Davis and Lavie (2017) concluded that a thick continental crust preserved in the outer domain is a favorable factor for the formation of volcanic passive margins.

In summary, taking all the above-mentioned observations and assumptions into account, we find the applicability of the Iberian model to the mid-Norwegian margin problematic, and favor the scenario where the distal and outer Møre and Vøring basins cannot easily represent a large zone of exhumed continental mantle before breakup. Most likely the volcanic margin formed on top of a distal marginal plateau, where a significant thickness of continental crust (> 5–8 km) was still present (Fig. 22).

5.2.2. On the role of structural and lithospheric inheritance

Both structural and lithospheric inheritance played important roles in the along-strike variations in architecture, segmentation and Cretaceous-Paleocene evolution and sedimentary infill of the Møre and Vøring basins (Zastrozhnov et al., 2018; Gernigon et al., 2019). From the Late Jurassic to mid-Albian time, active syn-thinning deposition is well-documented in the Møre and Vøring basins along pre-existing NE-SW trending rift valleys and were controlled by large detachment faults developed on the flanks of the bounding structural highs (Fig. 17a). The main tectonic sag basins developed between the proximal platforms and the western marginal plateaus. Later Cenomanian-Santonian and

Maastrichtian-Paleocene extensional events, which developed in the Vøring Basin, subsequently migrated sequentially towards the future OCB and still the main rifting-related depocentres were controlled by pre-existing deep-seated structural highs acting as crustal buffers (Figs. 18 and 19).

We suggest that the mid-Norwegian margin represents a specific type of passive rifted margin where large crustal blocks control the formation and regional distribution of the thick sedimentary sag-basins (Fig. 22). This specific configuration of the passive margin resembles the “crustal boudinage” model Gartrell (1997) and was already proposed for the Northern Vøring Margin case (Zastrozhnov et al., 2018) and has been suggested for many other rifted margins worldwide (Clerc et al., 2017; Jolivet et al., 2018). Here, the migration of the main deformational axis could be explained by the strengthening of lithosphere (Fig. 22) resulting from the intermediate thermal relaxation/subsidence phase (Van Wijk and Cloetingh, 2002; Gernigon et al., 2019). Van Wijk and Cloetingh (2002) considered ~20–60 Myr as the timing of thermal re-equilibration of the mid-Norwegian margin, which fits well with the observations in the Møre segment. However, according to our interpretation, the sharp and local relocation of the extensional axis in the Rås and Træna basins further west to the outer Vøring Basin could occur within just ~10 Myr and we suggest this enhancement could be the consequence of the potential lower crustal flow (Fig. 22).

The structure and depocentres distribution described in the results (Figs. 17–19) show that the NW-SE trending Jan Mayen Corridor, Rån, Surt and Bivrost lineaments define a long-lived and along-strike segmentation of the mid-Norwegian margin still effective during most of the Cretaceous period. The meaning and origin of these lineaments have been widely attributed to the Caledonian and/or pre-Caledonian shear zones (Blystad et al., 1995; Doré et al., 1997; Skilbrei and Olesen, 2005; Mjelde et al., 2009; Schiffer et al., 2019), or density contrasts within the upper mantle and at crustal levels (Maystrenko et al., 2017a, 2017b). The NE-SW structural trend appears to be dominant in the Møre and Vøring basins, except for the N-S trend typical for the southern Vøring Basin and adjacent Halten Terrace (Figs. 1 and 7). The rhomboidal structural geometries typical for this domain are constrained by the Jan Mayen Corridor and Rån Lineament (Fig. 7). Doré et al. (1999) interpreted the N-S trend as a result of E-W Jurassic extension which was then superimposed by NW-SE Early Cretaceous extension. However, Skilbrei and Olesen (2005) indicated that the internal grain of the Halten Terrace is also dissected by NE-SW trending subbasins formed before the Cretaceous (Figs. 1 and 7). Our results show that the NE-SW orientation of the structures is also present in the southern Vøring Basin (Figs. 1 and 7).

Together with Færseth and Lien (2002) we also argue for a consistent NW-SE Late Jurassic-earliest Cretaceous extension which re-activated the pre-existing N-S basement grain in the Halten Terrace and adjacent southern Vøring Basin in an oblique manner. This may have influenced the segmentation of the Møre and Vøring basins along NW-SE striking, extension parallel, transfer/weakness zones corresponding to the Jan Mayen Corridor and the Hevring and Rån lineaments. Further north, the Surt Lineament controlled the distribution of the extensional axis during the Late Cretaceous-Paleocene (Fig. 23). The rhomboidal geometries defined at the BCU level along the Surt Lineament support its manifestation as a transfer zone at least since the beginning of the Cretaceous (Fig. 7). The Bivrost Lineament has also been interpreted as a crustal transfer zone that influenced sedimentary basin segmentation in between the northern Vøring Basin and the Lofoten-Vesterålen Margin (Maystrenko et al., 2017a).

6. Conclusions

This contribution illustrates the structure and Cretaceous-Paleocene tectonostratigraphic evolution of the rift system between Norway and Greenland. Based on new 2D/3D data and a biostratigraphic re-evaluation of the exploration wells available, we have mapped the

entire margin and identified new structures in the distal part of the mid-Norwegian margin. We conclude that:

- The evolution of the mid-Norwegian margin from the Early Cretaceous to Paleocene exhibits episodic phases of extension and intermediate cooling events that preconditioned the margin lithosphere prior to the volcanic margin formation.
- After a significant Late Jurassic-earliest Cretaceous rifting event, a deformation mode shifted from a stretching to a thinning which was accommodated along large basin-floor detachments in the central basin domain until the mid-Albian. Also at least until the mid-Albian a prominent extensional branch developed in the outer domain of the Vøring Basin, where it is observed in the Rån Basin, Vigrid Syncline/Gjallar Ridge and on the eastern flank of the Nyk High (Någrind Syncline).
- An additional local Cenomanian-Santonian rifting event is proposed in the Vøring Basin to explain the atypical sedimentation rate within the Træna and Rås basins. This event proves that the tectonic evolution of rifted margins can vary significantly in space from one margin segment (e.g. the Møre Margin) to another margin segment (e.g. the Vøring Margin) separated by a regional crustal transfer system (e.g. the Jan Mayen Corridor).
- The final Late Cretaceous-Paleocene rift is commonly offset from the dominant mid-Mesozoic or older thinning axis of the mid-Norwegian margin. Strain hardening due to Mesozoic lithospheric cooling as well as additional enhancements from lateral lower crustal flow may explain the migration of depocentres and/or jumps of the rift axes over time. This may also have partly influenced the location of the final rift and volcanic margin together with the diachronous breakup configuration.
- The volcanic margin seems to have developed preferentially either on thick continental ribbons or moderately thinned continental crust. This interpretation is supported by the presence of the distal marginal plateaus highlighted by a shallow BCU and thicker crystalline crust which are constrained by the most recent 2D and 3D seismic surveys. The progressive thinning of the Cretaceous sequences towards the distal basin margins suggests that the marginal plateaus remained in a relatively high position during most of the Cretaceous period. A difference is, however, observed in the outer Vøring Basin, where a renewed phase of Late Cretaceous-Paleocene extension dislocated the predicted pre-existing distal plateau. This interpretation contrasts with a previous interpretation of the distal domain of the mid-Norwegian margin, in which a wide zone of exhumed mantle was proposed to fit a classical magma-poor Iberian margin scenario.
- Our regional reinterpretation of the Møre and Vøring basins shows that the long and polyphased rift system preceding the formation of the volcanic domain along the mid-Norwegian margin is markedly different to sediment-starved, magma-poor passive rifted margins. The mid-Norwegian margin has a different crustal architecture, a different basin evolution, higher sediment rates as well as a significant magmatic budget associated with the final lithospheric rupture between Eurasia and Greenland.

Author contribution

Dmitry Zastrozhnov: Conceptualization: Lead; Investigation: Lead; Validation: Lead; Visualization: Lead; Writing: Original Draft. Laurent Gernigon: Conceptualization: Supporting; Investigation: Supporting; Supervision: Lead; Validation: Supporting; Writing: review and editing. Iakov Gogin: Investigation: Supporting; Visualization: Supporting; Sverre Planke: Funding acquisition: Lead; Supervision: Supporting; Data Curation: Lead; Validation: Supporting; Writing: review and editing. Mansour M. Abdelmalak: Data Curation: Supporting; Visualization: Supporting; Writing: review and editing. Stephane Polteau: Writing: review and editing; Visualization: Supporting;

Supervision: Supporting. Jan Inge Faleide: Funding acquisition: Lead; Project administration: Lead; Writing: review and editing. Ben Manton: Visualization: Supporting; Writing – review and editing. Reidun Myklebust: Data Curation: Supporting; Visualization: Supporting.

Acknowledgments

The results of this work came from OMNIS Project, which received funding from the Research Council of Norway through its Centres of Excellence funding scheme, project number 223272. The seismic, magnetic, and gravity data presented in this study were provided by TGS. RPS Ichron is acknowledged for performing biostratigraphic analysis of key regional wells. Seismic interpretation was done using Kingdom Suite, Petrel and DUG InSight software. Grid interpolations and map compilations were established using Petrel, Geosoft Oasis Montaj and ArcGIS softwares. The seismic and raw well data used in this study are available through the Diskos National Data Repository (NDR). Well formation tops are available at the NPD factpages (www.npd.no) and the NORLEX group website (www.nhm2.uio.no/norlex/). Reviews from Tim Minshall and Kamaldeen Omosanya greatly helped us to improve the final manuscript.

Appendix 1. Stratigraphic comment on the main tectonostratigraphic units

BCU – mid-Albian

The Early Cretaceous interval of the section is not widely represented in the Vøring and Møre basins due to the reason that most of the wells in the area targeted Late Cretaceous and Tertiary structures. The Langebarn Formation in this area is holding a transitional position and both the Greenland-Barents Sea and North Sea/Norwegian shelf areas have their influence on the assemblage composition. The well 6610/3-1R provides an example of confidently dated Hauterivian-Barremian interval with such typical for the Norwegian shelf and the North Sea indicative taxa as *Cassiculosphaeridia magna*, *Hystrichosphaeridium arborispinum*, *Kleithrisphaeridium corrugatum* and *Batioladinium longicornutum* (Duxbury, 2001). On the other hand, the well 6608/2-1S (and ST2 bypass) represents much more expanded Early Cretaceous section where the presence of typical Greenland-Barents Sea events becomes more prominent. The Aptian interval of this well is possessing such typical northern latitudes event as an influx of *Vesperopsis mayi*, and the Albian section is characterized by the presence of *Pseudoceratium nudum*, *Batioladinium micropodum* and *Muderongia assymetrica* which also links to northern basins. The early Aptian OAE1 Selli event is prominently expressed in this well with the Gamma sharp increase on the log, and palynologically marked with abundant bisaccate pollens and sapropelic organics.

Mid-Albian – mid-Cenomanian

The early to middle Cenomanian section is well characterized biostratigraphically in number of wells and extensively expressed in the studied basin, reaching 1200 m thickness in the well 6603/5-1S. The consistent and abundant presence of *Sindridinium* genus representatives allows reliable intra-basinal correlation, tying the Vøring Basin to the Greenland, Barents Sea and North Sea reference sections (Nøhr-Hansen et al., 2017). The presence of abundant to superabundant peridinioid cysts in this particular well, but also in the basin overall (wells 6609/6-1, 6610/3-1R) is indicative for typical northern latitudes Cenomanian anoxic/high productivity environment with domination of the eutrophic taxa representatives (Pavlishina, 2017; Olde et al., 2014). The presence of representatives of *Rhombodella paucispina* in the basin is also providing a useful link between typical Cenomanian of the North Sea (Gradstein et al., 1999) and Greenland (Nøhr-Hansen et al., 2017). Otherwise, the presence of such taxa as *Litosphaeridium siphoniphorum*

and *Epelidosphaeridia spinosa* is evidence of typical European Cenomanian strata (Costa and Davey, 1992).

Mid-Cenomanian – Near Top Turonian

The late Cenomanian interval is represented in the studied area with very low thickness sections while the typical Cenomanian-Turonian boundary OAE2 event is not expressed in the basin. The Turonian section is widely presented in the basin, reaching more than 600 m in the well 6505/10-1 (Fig. 6) but lacks indicative marker events due to enriched carbonated lithology. The best representation of the interval is observed in the north-eastern wells where the relatively condensed sections possess a characteristic assemblage. The lower part of the section in the wells 6608/2-1S and 6609/6-1 is represented with early-middle Turonian influx of *Surculosphaeridium longifurcatum*, while the upper part of Turonian is well characterized with the presence of *Stephodinium coronatum*, *Aptodinium maculatum* and increased presence of *Palaeohystrichophora infusorioides* (Costa and Davey, 1992; Radmacher et al., 2015).

Top Turonian – mid-Campanian

The interval of the section begins with the final stage of Blålänge Formation sedimentation represented with Lysing Member sands accumulation. The events of the sand deposition show substantial diachrony, spanning Turonian-Coniacian boundary (Fig. 6). The oldest phase of the event is represented in the well 6505/10-1 where the typical top Turonian marker events such as presence of *Stephodinium coronatum* and increase in *Aptodinium maculatum* and *Palaeohystrichophora infusorioides* constantly appear in the 200 m mudstone succession above the top of the Lysing sand. In the well 6610/3-1R we can observe the Turonian/Coniacian boundary within the sandstone interval, supported with occurrence of *Stephodinium coronatum* and *Aptodinium maculatum*. The more typical case for the area is represented in the well 6706/11-1 where the early Coniacian acme of *Heterosphaeridium difficile* is recorded just above the sand while the top Turonian events of *Stephodinium coronatum* and *Palaeohystrichophora infusorioides* are observed just below the sand, supporting early Coniacian age for the sandstone deposition. Further sand development can be observed in the well 6605/8-1, where the basal early Coniacian acme of *Heterosphaeridium difficile* is observed between two phases of sand deposition, while the presence of *Spiniferites porosus* and *Psaligonyaulax deflandrei* just above the sand marks the top early Coniacian boundary.

The Kvitnos Formation is well biostratigraphically characterized in the Vøring Basin area, in particular in the well 6605/8-2, with its late Coniacian section being characterized with acme of *Chatangiella spinosa* and early Santonian interval possessing such markers as *Surculosphaeridium longifurcatum* and rare to common *Heterosphaeridium difficile*. The only sandstone deposition event observed for this interval is the presence of thin Tumler Member in the well 6706/11-1.

The major episode of sand deposition in the basin happened alongside with Nise Formation sedimentation, forming the late Santonian-Campanian Spekkhogger Member. These major sand units are greatly developed in the north of the basin, on the Gjallar Ridge and Nyk High and are unofficially referred to as Delfin sands. This sandstone deposition is observed in two stages, subdivided with early Campanian events of *Alterbidinium ioanidessii*, influx of *Chatangiella niiga* and acme of *Trithyrodinium suspectum*, generally occurring between the intervals of the sand deposition. The top part of this sandstone is characterized with typical Campanian events (Radmacher et al., 2015; Costa and Davey, 1992) such as increase and acme of *Heterosphaeridium heteracanthum*, common presence of *Trithyrodinium suspectum*, and occurrence of *Callaiosphaeridium asymmetricum* in the well 6706/11-1.

Mid-Campanian – Base Paleogene

The Springar Formation is well developed in the western part of the Vøring Basin nearby the Gjallar Ridge. The sand deposition in the area was active during early Maastrichtian and late Maastrichtian and is represented with Hvithval Member. The well 6604/2-1 is representing two episodes of sand deposition: (1) the lower Maastrichtian part of the Member is characterized with increase and acme events of *Alterbidinium acutulum* and *Chatangiella biapertura*; (2) upper Maastrichtian section is characterized with typical for this age occurrence and acme of *Palynodinium grallator*, *Trithyrodinium evittii* and *Impagidinium septentrionalis*, while the bottom part of this interval can be indicated with the increase of *Cerodinium diebelii*.

Base Paleogene - Top Paleocene

The Paleocene interval is represented with the Tang Formation in the section. The Danian part of the interval is expanded in the well 6604/10-1 where it is characterized with the influx of *Senoniasphaeria inornata*, followed by influx of *Palaeocystodinium bulliforme* and the presence of *Alisocysta reticulata* (Powell, 1992; Nøhr-Hansen, 2012). The close well 6604/2-1 is possessing also *Spongodinium delitiense* and *Spiniferites “magnifica”* at this level. The Selandian part of the interval is dated in the well 6604/10-1 with the increased presence of *Isabelidium viborgense*.

The Thanetian interval is well characterized in the basin and is represented in the well 6604/10-1. The basal part of the interval is marked with the increased presence of *Glaphyrocysta ordinata* group. The interval is characterized with the presence of numerous marker events including the presence and acme of *Alisocysta margarita*, *Alisocysta gippingensis*, *Spiniferites “membranispina”* and *Leptodinium “obscurum”* (Heilmann-Clausen, 1985; Powell, 1992). The top of Paleocene section is marked with the presence of *Palaeoperidinium pyrporum* and *Alisocysta margarita*.

A thick studied interval in the well 6706/6-1 from first returns to TD possesses mixed assemblage of Paleocene and Late Cretaceous taxa. The Tertiary assemblage is represented with Selandian-Danian marker taxa such as abundant *Alisocysta margarita* and *Palaeocystodinium bulliforme*. The Cretaceous assemblage is represented with early Maastrichtian to late-early Campanian succession of marker events. Although the succession of events may imply the Cretaceous age of the strata contaminated with cavings, we accept official NPD (NPD, 2019) viewpoint to assign the section to Paleocene until the further drilling in the area clarifies the stratigraphic uncertainty.

Appendix 2. Expansion of abbreviations used on the map figures

AH – Aukra High, AS – Aukra Subbasin, BH – Bara High, BL – Bivrost Lineament, BR – Bylgja Ridge, CMB – Central Møre Basin, CVB – Central Vøring Basin; DB – Danmarkshavn Basin, DH – Dufa High, DR – Danmarkshavn Ridge, EB – Ervik Basin, EKB – East Kiran Subbasin, ER – Ervik Ridge, FH – Frøya High, FG – Fenris Graben, GH – Grimm High, GIH – Golma High, GnH – Gnausen High, GR – Gjallar Ridge, GpB – Grip Subbasin, GpH – Grip High, GrH – Grønøy High, GS – Gleipne Saddle, GsB – Golma Subbasin, GsH – Gossa High, HB – Hermod Basin, HG – Hel Graben, HH – Hevring High, HL – Hevring Lineament, HIIH – Holmen High, HrH – Hrann High, HsB – Holmen Subbasin, HT – Hel Terrace, HvB – Hevring Subbasin, IMB – Inner Møre Basin, KH – Kinn High, KIH – Kolga High, KrH – Kiran High, JMC – Jan Mayen Corridor, JMMC – Jan Mayen Microcontinent Complex, LR – Lofoten Ridge, LVM – Lofoten-Vesterålen, MdS – Mardoll Subbasin, MgB – Magnus Basin, MH – Mimir High, MMP – Møre Marginal Plateau, MR – Manet Ridge, MS – Marmæle Spur, MTFC – Møre-Trøndelag Fault Complex, ND – Naglfar Dome, NGR – North Gjallar Ridge, NH – Nyk High, NSR – North Sklinna Ridge, NS – Någrind Syncline, NTB – North Træna Basin, NVB – Northern Vøring Basin, OCB – Ocean-continent boundary, OH – Ona

High, OLD – Ormen Lange Dome, OMB – Outer Møre Basin, OS – Orta Subbasin, OSVB – Outer Southern Vøring Basin, OVB – Outer Vøring Basin, RB – Rås Basin, RbB – Ribban Basin, RdB – Runde Subbasin, RH – Røst High, RL – Rån Lineament, RnB – Rån Basin, RR – Rån Ridge, RsB – Røst Basin, SGR – South Gjallar Ridge, Sfh – Sandflesa High, SH – Skoll High, SIB – Slørebotn Subbasin, SL – Surt Lineament, SR – Slettringen Ridge, SsB – Slettringen Subbasin, SSR – South Sklinna Ridge, StH – Steinen High, SVB – Southern Vøring Basin, TB – Træna Basin, TtB – Thetis Basin, TS – Tampen Spur, UH – Unn High, UtH – Utgard High, UR – Utrøst Ridge, VB – Vestfjorden Basin, VD – Vema Dome, VH – Vigra High, VfH – Valflesa High, VmH – Vimur High, VS – Vigrid Syncline, VsB – Vigra Subbasin, WKB – West Kiran Subbasin, YH – Ygg High.

References

- Abdelmalak, M.M., Meyer, R., Planke, S., Faleide, J.I., Gernigon, L., Frieling, J., Sluijs, A., Reichart, G.-J., Zastrozhnov, D., Theissen-Krah, S., Said, A., Myklebust, R., 2016a. Pre-breakup magmatism on the Vøring margin: Insight from new sub-basalt imaging and results from ocean drilling program hole 642E. *Tectonophysics* 675, 258–274. <https://doi.org/10.1016/j.tecto.2016.02.037>.
- Abdelmalak, M.M., Planke, S., Faleide, J.I., Jerram, D.A., Zastrozhnov, D., Eide, S., Myklebust, R., 2016b. The development of volcanic sequences at rifted margins: new insights from the structure and morphology of the Vøring Escarpment, mid-Norwegian Margin. *J. Geophys. Res. Solid Earth* 121, 5212–5236. <https://doi.org/10.1002/2015JB012788>.
- Abdelmalak, M.M., Faleide, J.I., Planke, S., Gernigon, L., Zastrozhnov, D., Shephard, G.E., Myklebust, R., 2017. The T-Reflection and the deep crustal structure of the Vøring Margin offshore Mid-Norway. *Tectonics* 36, 2497–2523. <https://doi.org/10.1002/2017TC004617>.
- Allen, P.A., Allen, J.R., 2013. *Basin Analysis: Principles and Application to Petroleum Play Assessment*. John Wiley, pp. 642.
- Bakke, V.N., 2017. *Volcanic Rocks at the Møre Marginal High: Geochemistry, Petrogenesis and Emplacement Mechanisms*. Master's thesis. The University of Bergen.
- Berndt, C., Planke, S., Alvestad, E., Tsikalas, F., Rasmussen, T., 2001. Seismic volcanotratigraphy of the Norwegian Margin: constraints on tectonomagmatic break-up processes. *J. Geol. Soc.* 158, 413–426.
- Blystad, P., Brekke, H., Færseth, R.B., Larsen, B.T., Skogseid, J., Tørrudbakken, B., 1995. Structural elements of the Norwegian continental shelf Part II: the Norwegian Sea Region, NPD-Bulletin. The Norwegian Petroleum Directorate 8.
- Bjørnseth, H.M., Grant, S.M., Hansen, E.K., Hossack, J.R., Roberts, D.G., Thompson, M., 1997. Structural evolution of the Vøring basin, Norway, during the late cretaceous and Paleogene. *J. Geol. Soc. (Lond.)* 154 (3), 559–563. <https://doi.org/10.1144/gsjgs.154.3.0559>.
- Breivik, A., Faleide, J.I., Mjelde, R., Flueh, E., Murai, Y., 2014. Magmatic development of the outer Vøring margin from seismic data. *Journal of Geophysical Research-Solid Earth* 119 (9), 6733–6755.
- Brekke, H., 2000. The tectonic evolution of the Norwegian sea continental margin with emphasis on the Vøring and Møre basins. Geological Society, London, Special Publications 167 (1), 327–378. <https://doi.org/10.1144/gsl.sp.2000.167.01.13>.
- Brekke, H., Sjulstad, H.I., Magnus, C., Williams, R.W., 2001. Sedimentary environments offshore Norway—an overview. In: In: Martinsen, O.J., Dreyer, T. (Eds.), *Sedimentary Environments Offshore Norway—Palaeozoic to Recent*, vol. 10. NPF Special Publication, pp. 7–37.
- Brune, S., Heine, C., Perez-Gussinye, M., Sobolev, S.V., 2014. Rift migration explains continental margin asymmetry and crustal hyper-extension. *Nat. Commun.* 5. <https://doi.org/10.1038/ncomms5014>.
- Chenin, P., Manatschal, G., Lavier, L.L., Errat, D., 2015. Assessing the impact of orogenic inheritance on the architecture, timing and magmatic budget of the North Atlantic rift system: a mapping approach. *J. Geol. Soc.* 172 (6), 711–720. <https://doi.org/10.1144/jgs2014-139>.
- Clerc, C., Ringenbach, J.-C., Jolivet, L., Ballard, J.-F., 2017. Rifted margins: ductile deformation, boudinage, continentward-dipping normal faults and the role of the weak lower crust. *Gondwana Res.* <https://doi.org/10.1016/j.jgr.2017.04.030>.
- Cohen, K.M., Finney, S.C., Gibbard, P.L., Fan, J.-X., 2013. The ICS international chronostratigraphic chart. *Episodes* 36 (3), 199–204.
- Costa, L.I., Davey, R.J., 1992. Dinoflagellate cysts of the cretaceous system. In: Powell, A.J. (Ed.), *A Stratigraphic Index of Dinoflagellate Cysts*. Chapman and Hall, London.
- Dalland, A., Worsley, D., Ofstad, K., 1988. A lithostratigraphic scheme for the Mesozoic and Cenozoic succession offshore mid- and northern Norway. *NPD Bulletin* 4, 63.
- Davis, J.K., Lavier, L.L., 2017. Influences on the development of volcanic and magma-poor morphologies during passive continental rifting. *Geosphere* 13 (5), 1524–1540.
- Dean, K., McLachlan, K., Chambers, A., 1999. Rifting and the development of the Faeroe-Shetland Basin. Geological Society, London, Petroleum Geology Conference series 5, 533–544. <https://doi.org/10.1144/0050533>.
- Doré, A.G., Lundin, E.R., Fichler, C., Olesen, O., 1997. Patterns of basement structure and reactivation along the NE Atlantic margin. *J. Geol. Soc.* 154, 85–92.
- Doré, A.G., Lundin, E.R., Jensen, L.N., Birkland, Ø., Eliassen, P.E., Fichler, C., 1999. Principal tectonic events in the evolution of the northwest European Atlantic margin. Geological Society, London, Petroleum Geology Conference series 5, 41–61. <https://doi.org/10.1144/0050041>.
- Duxbury, S., 2001. A palynological zonation scheme for the lower cretaceous - United Kingdom sector, central North Sea. *N. Jb. Geol. Pal. Abh.* 219, 95–137.
- Ebbing, J., Lundin, E., Olesen, O., Hansen, E.K., 2006. The mid-Norwegian margin: a discussion of crustal lineaments, mafic intrusions, and remnants of the Caledonian root by 3D density modelling and structural interpretation. *J. Geol. Soc.* 163, 47–59. <https://doi.org/10.1144/0016-764905-029>. London.
- Eldholm, O., Thiede, J., Taylor, E., 1989. Evolution of the Vøring volcanic margin. In: Eldholm, O., Thiede, J., Taylor, E. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results. Ocean Drilling Program, College Station, Tex.*, pp. 1033–1065.
- Eldholm, O., Tsikalas, F., Faleide, J.I., 2002. Continental margin off Norway 62–75°N: palaeogene tectono-magmatic segmentation and sedimentation. *Geol. Soc. London Spec. Publ.* 197 (1), 39–68. <https://doi.org/10.1144/gsl.sp.2002.197.01.03>.
- Faleide, J.I., Vågnes, E., Gudlaugsson, S.T., 1993. Late Mesozoic-Cenozoic evolution of the southwestern Barents Sea in a regional rift-shear tectonic setting. *Mar. Petrol. Geol.* 10, 186–214. [https://doi.org/10.1016/0264-8172\(93\)90104-Z](https://doi.org/10.1016/0264-8172(93)90104-Z).
- Faleide, J.I., Tsikalas, F., Breivik, A.J., Mjelde, R., Ritzmann, O., Engen, Ø., Wilson, J., Eldholm, O., 2008. Structure and evolution of the continental margin off Norway and the Barents Sea. *Episodes* 31 (No. 1), 82–91.
- Fjeldskaar, W., Grunnaleite, I., Zweigel, J., Mjelde, R., Faleide, J.I., Wilson, J., 2009. Modelled palaeo-temperature on Vøring, offshore mid-Norway - the effect of the Lower Crustal Body. *Tectonophysics* 474 (3–4), 544–558.
- Fjellanger, E., Surlyk, F., Wamsteeker, L.C., Midtun, T., 2005. Upper Cretaceous basin-floor fans in the Vøring basin, mid Norway shelf. In: In: Wandås, B., Nystuen, J.P., Eide, E., Gradstein, F.M. (Eds.), *Onshore-Offshore Relationships on the North Atlantic Margin*, vol. 12. Norwegian Petroleum Society Special Publications, pp. 135–164. [https://doi.org/10.1016/S0928-8937\(05\)80047-5](https://doi.org/10.1016/S0928-8937(05)80047-5).
- Færseth, R.B., Lien, T., 2002. Cretaceous evolution in the Norwegian Sea – a period characterized by tectonic quiescence. *Mar. Petrol. Geol.* 19, 1005–1027. [https://doi.org/10.1016/S0264-8172\(02\)00112-5](https://doi.org/10.1016/S0264-8172(02)00112-5).
- Færseth, R.B., 2012. Structural development of the continental shelf offshore Lofoten-Vesterålen, northern Norway. *Norw. J. Geol.* 92, 19–40.
- Gaina, C., Gernigon, L., Ball, P., 2009. Paleocene-recent plate boundaries in the NE Atlantic and the formation of the Jan Mayen microcontinent. *J. Geol. Soc.* 166 (4), 601–616.
- Geoffroy, L., Burov, E., Werner, P., 2015. Volcanic passive margins: another way to break up continents. *Sci. Rep.* 5, 14828.
- Gernigon, L., Ringenbach, J.-C., Planke, S., Jonquet-Kolsto, H., 2003. Extension, crustal structure and magmatism at the outer Vøring basin, Norwegian margin. *J. Geol. Soc.* 160, 197–208. <https://doi.org/10.1144/0016-764902-055>. London.
- Gernigon, L., Ringenbach, J.-C., Planke, S., Le Gall, B., 2004. Deep structures and breakup along volcanic rifted margins: insights from integrated studies along the outer Vøring Basin (Norway). *Mar. Petrol. Geol.* 21 (3), 363–372.
- Gernigon, L., Lucaczeu, F., Brigaud, F., Ringenbach, J.-C., Planke, S., Le Gall, B., 2006. A moderate melting model for the Vøring margin (Norway) based on structural observations and a thermo-kinematical modelling: implication for the meaning of the lower crustal bodies. *Tectonophysics* 412 (3), 255–278.
- Gernigon, L., Gaina, C., Olesen, O., Ball, P., Péron-Pinvidic, G., Yamasaki, T., 2012. The Norway Basin revisited: from continental breakup to spreading ridge extinction. *Mar. Petrol. Geol.* 35 (1), 1–19.
- Gernigon, L., Blischke, A., Nasuti, A., Sand, M., 2015. Conjugate volcanic rifted margins, seafloor spreading, and microcontinent: insights from new high-resolution aeromagnetic surveys in the Norway Basin. *Tectonics* 34, 907–933. <https://doi.org/10.1002/2014TC003717>.
- Gernigon, L., Franke, D., Geoffroy, L., Schiffer, C., Foulger, G.R., Stoker, M., 2019. Crustal fragmentation, magmatism, and the diachronous opening of the Norwegian-Greenland Sea. *Earth Sci. Rev.* <https://doi.org/10.1016/j.earscirev.2019.04.011>.
- Gjelberg, J.G., Martinsen, O.J., Charnock, M., Møller, N., Antonsen, P., 2005. The reservoir development of the late maastichtian - early Paleocene omen Lange gas field, Møre basin, mid-Norwegian shelf. In: Dore, A.G., Vining, B.A. (Eds.), *Petroleum Geology: North-West Europe and Global Perspectives - Proceedings of the 6th Petroleum Geology Conference*. Geological Society, London, pp. 1165–1184.
- Gradstein, F.M., Kaminski, M.A., Agterberg, F.P., 1999. Micropaleontology and paleoceanography of the cretaceous seaway between Norway and Greenland. In: Gradstein, F.M., Van der Zwaan, G. (Eds.), *Ordering the Fossil Record - Challenges in Stratigraphy and Paleontology*, Symp. Volume 75th Birthday Prof. C. W. Drooger, Spec. Publ. Earth Science Review. Elsevier Publ. Co., Amsterdam, pp. 75.
- Gradstein, F.M., Antonissen, E., Brunstad, H., Charnock, M., Hammer, Ø., Hellem, T., Lervik, K.S., 2010. Norwegian offshore stratigraphic lexicon (NORLEX). *News. Stratigr.* 44, 73–86.
- Guan, H., Geoffroy, L., Gernigon, L., Chauvet, F., Grigné, C., Werner, P., 2019. Magmatic ocean-continent transitions. *Mar. Petrol. Geol.* <https://doi.org/10.1016/j.marpetgeo.2019.04.003>.
- Hansen, J.W., Bakke, S., Fanavoll, S., 1992. *Shallow Drilling Nordland VI and VII 1992*. report 23. IKU.
- Henstra, G.A., Gawthorpe, R.L., Helland-Hansen, W., Ravnås, R., Rotevatn, A., 2016. Depositional systems in multiphase rifts: seismic case study from the Lofoten margin, Norway. *Basin Res.* 1–23. <https://doi.org/10.1111/bre.12183>.
- Hinz, K., 1981. A hypothesis on terrestrial catastrophes. Wedges of very thick seaward dipping layers beneath passive continental margins. Their origin and paleoenvironmental significance. *Geol. Jahrb.* E22, 3–28.
- Hjelstuen, B.O., Eldholm, O., Skogseid, J., 1999. Cenozoic evolution of the northern Vøring Margin. *Geol. Soc. Am. Bull.* 111, 1792–1807. [https://doi.org/10.1130/0016-7606\(1999\)111<1792:CEOTNV>2.3.CO;2](https://doi.org/10.1130/0016-7606(1999)111<1792:CEOTNV>2.3.CO;2).
- Huismans, R.S., Beaumont, C., 2011. Depth-dependent extension, two-stage breakup and cratonic underplating at rifted margins. *Nature* 473. <https://doi.org/10.1038/nature09988>.

- Jolivet, L., Menant, A., Clerc, C., Sternai, P., Bellahsen, N., Leroy, S., Pik, R., Stab, M., Faccenna, C., Gorini, G., 2018. Extensional crustal tectonics and crust-mantle coupling, a view from the geological record. *Earth Sci. Rev.* 185, 1187–1209.
- Kittelsen, J.E., Olsen, R.R., Marten, R.F., Hansen, E.K., Hollingsworth, R.R., 1999. The first deepwater well in Norway and its implications for the Upper Cretaceous Play, Vøring Basin. Geological Society, London, Petroleum Geology Conference series 5, 275–280. <https://doi.org/10.1144/0050275>.
- Kjoberg, S., Schmiedel, T., Planke, S., Svensen, H., Millett, J., Jerram, D.A., Galland, O., Lecomte, I., Schofield, N., Haug, Ø.T., Heslem, A., 2017. 3D structure and formation of hydrothermal vent complexes at the Paleocene-Eocene transition, the Møre Basin. mid-Norwegian Margin: Interpretation 5 <https://doi.org/10.1190/int-2016-0159.1>. this issue.
- Kvarven, T., Mjelde, R., Hjelstuen, B.O., Faleide, J.I., Thybo, H., Flueh, E.R., Murai, Y., 2016. Crustal composition of the more margin and compilation of a conjugate Atlantic margin transect. *Tectonophysics* 666, 144–157.
- Kyrkjebø, R., Gabrielsen, R.H., Faleide, J.I., 2004. Unconformities related to the Jurassic–Cretaceous synrift–post-rift transition of the northern North Sea. *J. Geol. Soc.* 161 (1), 1–17.
- Lavier, L.L., Manatschal, G., 2006. A mechanism to thin the continental lithosphere at magma-poor margins. *Nature* 440 (7082), 324–328. <https://doi.org/10.1038/nature04608>.
- Lien, T., 2005. From rifting to drifting: effects on the development of deep-water hydrocarbon reservoirs in a passive margin setting. *Norwegian Sea. Nor. J. Geol./Nor. Geol. Foren.* 85 (4).
- Ligi, M., Bonatti, E., Bosworth, W., Cai, Y., Cipriani, A., Palmiotto, C., Ronca, S., Seyler, M., 2018. Birth of an ocean in the Red Sea: oceanic-type basaltic melt intrusions precede continental rupture. *Gondwana Res.* 54, 150–160.
- Lister, G.S., Etheridge, M.A., Symonds, P.A., 1986. Detachment faulting and the evolution of passive continental margins. *Geology* 14, 246–250. [https://doi.org/10.1130/0091-7613\(1986\)14<246:DFATEO>2.0.CO;2](https://doi.org/10.1130/0091-7613(1986)14<246:DFATEO>2.0.CO;2).
- Lundin, E.R., Doré, A.G., 1997. A tectonic model for the Norwegian passive margin with implications for the NE Atlantic: early Cretaceous to break-up. *Geological Society London Journal* 154, 545–550. <https://doi.org/10.1144/gsjgs.154.3.0545>.
- Lundin, E.R., Doré, A.G., 2011. Hyperextension, serpentinization, and weakening: a new paradigm for rifted margin compressional deformation. *Geology* 39, 347–350. <https://doi.org/10.1130/G31499.1>.
- Lundin, E.R., Doré, A.G., Rønning, K., Kyrkjebø, R., 2013. Repeated inversion and collapse in the Late Cretaceous–Cenozoic northern Vøring Basin, offshore Norway. *Petrol. Geosci.* 19, 329–341. <https://doi.org/10.1144/petgeo2012-022>.
- Lundin, E., Doré, A., 2018. Non-Wilsonian Break-Up Predisposed by Transforms: Examples from the North Atlantic and Arctic, vol. 470 Geological Society, London, Special Publications (SP470). 6).
- Manton, B., Planke, S., Zastrozhnov, D., Abdelmalak, M.M., Millett, J., Maharjan, D., Polteau, S., Faleide, J.I., Gernigon, L., Myklebust, R., 2018. Pre-Cretaceous prospectivity of the outer Møre and Vøring basins constrained by new 3D seismic data. In: 80th EAGE Conference and Exhibition 2018.
- Maystrenko, Y.P., Olesen, O., Gernigon, L., Gradmann, S., 2017a. Deep structure of the Lofoten–Vesterålen segment of the Mid-Norwegian continental margin and adjacent areas derived from 3-D density modeling. *J. Geophys. Res. Solid Earth* 122. <https://doi.org/10.1002/2016JB013443>.
- Maystrenko, Y.P., Gernigon, L., Nasuti, A., Olesen, O., 2017b. Deep structure of the Mid-Norwegian continental margin (the Vøring and Møre basins) according to 3-D density and magnetic modelling. *Geophys. J. Int.* 212 (3), 1696–1721.
- Meyer, R., van Wijk, J., Gernigon, L., 2007. The North Atlantic igneous province: a review of models for its formation. In: In: Foulger, G.R., Jurdy, D.M. (Eds.), *Plates, Plumes and Planetary Processes*, vol. 430. Geological Society of America Special Paper, pp. 525–552. [https://doi.org/10.1130/2007.2430\(26](https://doi.org/10.1130/2007.2430(26).
- Meyer, R., Hertogen, J., Pedersen, R.B., Viereck-Götte, L., Abratis, M., 2009. Interaction of mantle derived melts with crust during the emplacement of the Vøring Plateau, NE Atlantic. *Mar. Geol.* 261 (1–4), 3–16.
- McKenzie, D., 1978. Some remarks on the development of sedimentary basins. *Earth Planet Sci. Lett.* 40, 25–32. [https://doi.org/10.1016/0012-821X\(78\)90071-7](https://doi.org/10.1016/0012-821X(78)90071-7).
- Millett, J.M., Manton, B.M., Zastrozhnov, D., Planke, S., Maharjan, D., Bellwald, B., Abdelmalak, M.M., 2020. Basin Structure and Prospectivity of the NE Atlantic Volcanic Rifted Margin: Cross-Border Examples from the Faroe-Shetland, Møre and Southern Vøring Basins. Geological Society, London, Special Publications, pp. 495.
- Mjelde, R., Kasahara, J., Shimamura, H., Kamimura, A., Kanazawa, T., Kodaira, S., Raum, T., Shiobara, H., 2002. Lower crustal seismic velocity anomalies; magmatic underplating or serpentinized peridotite? Evidence from the Vøring Margin, NE Atlantic. *Mar. Geophys. Res.* 23 (2), 169–183.
- Mjelde, R., Faleide, J.I., Breivik, A.J., Raum, T., 2009. Lower crustal composition and crustal lineaments on the Vøring margin, NE Atlantic: a review. *Tectonophysics* 472 (1–4), 183–193.
- Mjelde, R., Kvarven, T., Faleide, J.I., Thybo, H., 2016. Lower crustal high-velocity bodies along North Atlantic passive margins, and their link to Caledonian suture zone eclogites and Early Cenozoic magmatism. *Tectonophysics* 670, 16–29. <https://doi.org/10.1016/j.tecto.2015.11.021>.
- Mjelde, R., Shimamura, H., Kanazawa, T., Kodaira, S., Raum, T., Shiobara, H., 2003. Crustal lineaments, distribution of lower crustal intrusives and structural evolution of the Vøring Margin, NE Atlantic; new insight from wide-angle seismic models. *Tectonophysics* 369 (3–4), 199–218.
- Mohn, G., Karner, G.D., Manatschal, G., Johnson, C.A., 2015. Structural and stratigraphic evolution of the Iberia–Newfoundland hyper-extended rifted margin: a quantitative modelling approach. *Geological Society, London, Special Publications* 413 (1), 53–89.
- Mutter, J.C., Talwani, M., Stoffa, P.L., 1982. Origin of seaward-dipping reflectors in oceanic crust off the Norwegian margin by “subaerial sea-floor spreading”. *Geology* 10 (7), 353–357.
- Myer, D., Constable, S., Key, K., 2013. Magnetotelluric evidence for layered mafic intrusions beneath the Vøring and Exmouth rifted margins. *Phys. Earth Planet. In.* 220, 1–10.
- Nirrengarten, M., Gernigon, L., Manatschal, G., 2014. Lower crustal bodies in the Møre volcanic rifted margin: geophysical determination and geological implications. *Tectonophysics* 636, 143–157. <https://doi.org/10.1016/j.tecto.2014.08.004>.
- Norwegian Petroleum Directorate (NPD), 2019. Factpages. available at: <http://factpages.npd.no/factpages/> (last access: September 2019).
- Nirrengarten, M., Mohn, G., Schito, A., Corrado, S., Gutiérrez-García, L., Bowden, S., Despinois, F., February 2020. The thermal imprint of continental breakup during the formation of the South China Sea. *Earth Planet Sci. Lett.* 531, 115972.
- Nøhr-Hansen, H., 1993. Dinoflagellate cyst stratigraphy of the barremian to albian, lower cretaceous, north-east Greenland. *Bull. Grøn. Geol. Unders.* 166, 1–171.
- Nøhr-Hansen, H., 1996. Upper Cretaceous dinoflagellate cyst stratigraphy, onshore West Greenland. *Bull. - Grøn. Geol. Undersøgelser* 170, 104.
- Nøhr-Hansen, H., Costa, L.I., Pearce, M.A., Alsen, P., 2017. New Albian to Cenomanian (Cretaceous) Dinoflagellate Cyst Taxa of Ovoidinoid Affinities from East Greenland, the Barents Sea and England. *Palynology*, pp. 1–26.
- Nøhr-Hansen, H., 2012. Palynostratigraphy of the cretaceous–lower palaeogene sedimentary succession in the kangerlussuaq basin, southern east Greenland. *Rev. Palaeobot. Palynol.* 178, 59–90.
- Olde, K., Jarvis, I., Pearce, M., Uličný, D., Tocher, B., Trabuco-Alexandre, J., Gröcke, D., 2014. A revised northern European Turonian (Upper Cretaceous) dinoflagellate cyst biostratigraphy: integrating palynology and carbon isotope events. *Rev. Palaeobot. Palynol.* 213, 1–16.
- Osmundsen, P.T., Ebbing, J., 2008. Style of extension offshore mid-Norway and implications for mechanisms of crustal thinning at passive margins. *Tectonics* 27, 1–12. <https://doi.org/10.1029/2007TC002242>.
- Osmundsen, P.T., Péron-Pinvidic, G., Ebbing, J., Erratt, D., Fjellanger, E., Bergslien, D., Syvertsen, S.E., 2016. Extension, hyperextension and mantle exhumation offshore Norway: a discussion based on 6 crustal transects. *Norw. J. Geol.* 96, 343–372. <https://doi.org/10.17850/njg96-4-05>.
- Osmundsen, P.T., Péron-Pinvidic, G., 2018. Crustal-scale fault interaction at rifted margins and the formation of domain-bounding breakaway complexes: insights from offshore Norway. *Tectonics* 37 (3), 935–964. <https://doi.org/10.1002/2017TC004792>.
- Pascoe, R., Hooper, R., Storhaug, K., Harper, H., 1999. Evolution of extensional styles at the southern termination of the Nordland Ridge, Mid-Norway: a response to variations in coupling above Triassic salt. In: *Geological Society, London, Petroleum Geology Conference Series*, vol. 5. pp. 83–90. <https://doi.org/10.1144/0050083>.
- Pavlishina, P., 2017. Palynostratigraphy and palaeoenvironments around the Albian–Cenomanian boundary interval (OAE1d), North Bulgaria. *Sci. China Earth Sci.* 60, 71–79.
- Péron-Pinvidic, G., Manatschal, G., 2010. From microcontinents to extensional allochthons: witnesses of how continents rift and break apart? *Petrol. Geosci.* 16 (3), 189–197.
- Péron-Pinvidic, G., Manatschal, G., Osmundsen, P.T., 2013. Structural comparison of archetypal Atlantic rifted margins: a review of observations and concepts. *Mar. Petrol. Geol.* 43, 21–47. <https://doi.org/10.1016/j.marpetgeo.2013.02.002>.
- Péron-Pinvidic, G., Osmundsen, P.T., 2016. Architecture of the distal and outer domains of the Mid-Norwegian rifted margin: insights from the Rån-Gjallar ridges system. *Mar. Petrol. Geol.* 77, 280–299. <https://doi.org/10.1016/j.marpetgeo.2016.06.014>.
- Péron-Pinvidic, G., Osmundsen, P.T., 2018. The Mid Norwegian-NE Greenland conjugate margins: rifting evolution, margin segmentation, and breakup. *Mar. Petrol. Geol.* 98, 162–184.
- Petersen, K.D., Schiffer, C., 2016. Wilson cycle passive margins: control of orogenic inheritance on continental breakup. *Gondwana Res.* 39, 131–144. <https://doi.org/10.1016/j.jgr.2016.06.012>.
- Planke, S., Rasmussen, T., Rey, S., Myklebust, R., 2005. Seismic characteristics and distribution of volcanic intrusions and hydrothermal vent complexes in the Vøring and Møre basins. In: *Geological Society, London, Petroleum Geology Conference Series. Geological Society of London*, pp. 833–844.
- Planke, S., Millett, J.M., Maharjan, D., Jerram, D.A., Abdelmalak, M.M., Groth, A., Hoffmann, J., Berndt, C., Myklebust, R., 2017. Igneous seismic geomorphology of buried lava fields and coastal escarpments on the Vøring volcanic rifted margin. *Interpretation* 5 (3), 1–42.
- Powell, J.A., 1992. Dinoflagellate cysts of the tertiary system. In: Powell, A.J. (Ed.), *A Stratigraphic Index of Dinoflagellate Cysts*. Chapman and Hall, London.
- Radmacher, W., Tyszkaj, J., Mangerud, G., Pearce, M.A., 2014. Dinoflagellate cyst biostratigraphy of the late albian to early mastrichtian in the southwestern Barents sea. *Mar. Petrol. Geol.* 57, 109–121.
- Radmacher, W., Mangerud, G., Tyszkaj, J., 2015. Dinoflagellate cyst biostratigraphy of Upper Cretaceous strata from two wells in the Norwegian Sea. *Rev. Palaeobot. Palynol.* 216, 18–32.
- Ren, S., Skogseid, J., Eldholm, O., 1998. Late Cretaceous–Paleocene extension on the Vøring volcanic margin. *Mar. Geophys. Res.* 20 (4), 343–369. <https://doi.org/10.1023/A:1004554217069>.
- Ren, S., Faleide, J.I., Eldholm, O., Skogseid, J., Gradstein, F., 2003. Late cretaceous-paleocene development of the NW Vøring basin. *Mar. Petrol. Geol.* 20, 177–206. [https://doi.org/10.1016/S0264-8172\(03\)00005-9](https://doi.org/10.1016/S0264-8172(03)00005-9).
- Reynisson, R.F., Ebbing, J., Lundin, E., Osmundsen, P.T., 2010. Properties and distribution of lower crustal bodies on the mid-Norwegian margin. *Geological Society, London, Petroleum Geology Conference series* 7, 843–854. <https://doi.org/10.1144/0070843>.

- Roberts, A.M., Corfield, R.I., Kusznir, N.J., Matthews, S.J., Hansen, E.-K., Hooper, R.J., 2009. Mapping palaeostructure and palaeobathymetry along the Norwegian Atlantic continental margin, Møre and Vøring basins. *Petrol. Geosci.* 15, 27–43. <https://doi.org/10.1144/1354-079309-804>.
- Reston, T.J., 2009. The structure, evolution and symmetry of the magma-poor rifted margins of the North and Central Atlantic: a synthesis. *Tectonophysics* 468 (1–4), 6–27.
- Rüpke, L.H., Schmid, D.W., Perez-Gussinye, M., Hartz, E., 2013. Interrelation between rifting, faulting, sedimentation, and mantle serpentinization during continental margin formation-including examples from the Norwegian Sea. *Geochim. Geophys. Geosyst.* 14, 4351–4369. <https://doi.org/10.1002/ggge.20268>.
- Saunders, A.D., Fitton, J.G., Kerr, A.C., Norry, M.J., Kent, R.W., 1997. The North Atlantic igneous provinces. In: Mahoney, J.J., Coffin, M.F. (Eds.), *Large Igneous Provinces: Continental, Oceanic, and Planetary Flood Volcanism*. American Geophysical Union, Geophysical Monograph, pp. 45–93.
- Schiffer, C., Doré, A.G., Foulger, G.R., Franke, D., Geoffroy, L., Gernigon, L., Peace, A., 2019. Structural inheritance in the north Atlantic. *Earth Sci. Rev.* 102975.
- Schmiedel, T., Kjøberg, S., Planke, S., Magee, C., Galland, O., Schofield, N., Jackson, C.A.-L., Jerram, D.A., 2017. Mechanisms of overburden deformation associated with the emplacement of the Tulipan sill, mid-Norwegian margin. *Interpretation* 5, SK23–SK38.
- Schöpfer, K., Hinsch, R., 2019. Late Palaeozoic – Mesozoic tectonostratigraphic development of the eastern Faroe-Shetland Basin: new insights from high-resolution 3D seismic and well data. *Mar. Petrol. Geol.* 109, 494–518. <https://doi.org/10.1016/j.marpetgeo.2019.04.007>.
- Simon, K., Huisman, R.S., Beaumont, C., 2009. Dynamical modelling of lithospheric extension and small-scale convection: implications for magmatism during the formation of volcanic rifted margins. *Geophys. J. Int.* 176 (1), 327–350.
- Skilbrei, J.R., Olesen, O., 2005. Deep structure of the Mid-Norwegian shelf and onshore-offshore correlations: Insight from potential field data. In: Wandås, B., Nystuen, J.P., Eide, E.A. (Eds.), *Onshore-offshore Relationships on the North Atlantic Margin*, vol. 12. Norwegian Petroleum Society (NPF) Special Publication, pp. 43–67.
- Skogseid, J., Eldholm, O., 1989. Vøring plateau continental margin: seismic interpretation, stratigraphy and vertical movements. In: Eldholm, O., Thiede, J., Taylor, E. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results, College Station, TX (Ocean Drilling Pro)*. Proceedings of the Ocean Drilling Program, Scientific Results, vol. 104. Ocean Drilling Program, College Station, TX, pp. 993–1030.
- Skogseid, J., Planke, S., Faleide, J.I., Pedersen, T., Eldholm, O., Neverdal, F., 2000. NE Atlantic continental rifting and volcanic margin formation. In: Nøttvedt, A. (Ed.), *Dynamics of the Norwegian Margin*, vol. 167. Geological Society (London) Special Publication, pp. 295–326. <https://doi.org/10.1144/GSL.SP.2000.167.01.12>.
- Stoker, M.S., 2016. Cretaceous tectonostratigraphy of the faroe-shetland region. *Scot. J. Geol.* 52 (1), 19–41.
- Swiecicki, T., Gibbs, P.B., Farrow, G.E., Coward, M.P., 1998. A tectonostratigraphic framework for the mid-Norway region. *Mar. Petrol. Geol.* 15 (3), 245–276. [https://doi.org/10.1016/S0264-8172\(97\)00029-9](https://doi.org/10.1016/S0264-8172(97)00029-9).
- Sømme, T.O., Jackson, C.A.L., 2013. Source-to-sink analysis of ancient sedimentary systems using a subsurface case study from the Møre-Trøndelag area of southern Norway: Part 2 – sediment dispersal and forcing mechanisms. *Basin Res.* 25, 512–531. <https://doi.org/10.1111/bre.12014>.
- Talwani, M., Eldholm, O., 1977. Evolution of the Norwegian-Greenland sea. *Geol. Soc. Am. Bull.* 88, 969–999.
- Tetreault, J.L., Buiter, S.J.H., 2018. The influence of extension rate and crustal rheology on the evolution of passive margins from rifting to break-up. *Tectonophysics* 746, 155–172. <https://doi.org/10.1016/j.tecto.2017.08.029>.
- Theissen-Krah, S., Zastrozhnov, D., Abdelmalak, M.M., Schmid, D.W., Faleide, J.I., Gernigon, L., 2017. Tectonic evolution and extension at the Møre Margin – offshore mid-Norway. *Tectonophysics* 721, 227–238. <https://doi.org/10.1016/j.tecto.2017.09.009>.
- Tsikalas, F., Faleide, J.I., Eldholm, O., 2001. Lateral variations in tectono-magmatic style along the Lofoten-Vesterålen margin off Norway. *Mar. Petrol. Geol.* 18, 807–832. [https://doi.org/10.1016/S0264-8172\(01\)00030-7](https://doi.org/10.1016/S0264-8172(01)00030-7).
- Tsikalas, F., Faleide, J.I., Eldholm, O., Wilson, J., 2005. Conjugate mid-Norway and NE Greenland continental margins. In: Doré, A.G., Vining, B.A. (Eds.), *Petroleum Geology: North-West Europe and Global Perspectives—Proceedings of the 6th Petroleum Geology Conference*, pp. 785–801.
- Tsikalas, F., Faleide, J.I., Eldholm, O., Blaich, O.A., 2012. The NE Atlantic conjugate margins. In: Roberts, D.G., Bally, A.W. (Eds.), *Phanerozoic Passive Margins, Cratonic Basins and Global Tectonic Maps*, pp. 141–201.
- Tsikalas, F., Faleide, J.I., Kalač, A., 2019. New insights into the Cretaceous-Cenozoic tectono-stratigraphic evolution of the southern Lofoten margin, offshore Norway. *Mar. Petrol. Geol.* 110, 832–855. <https://doi.org/10.1016/j.marpetgeo.2019.07.025>.
- Van Avendonk, H.J.A., Lavie, L.L., Shillington, D.J., Manatschal, G., 2009. Extension of continental crust at the margin of the eastern Grand Banks, Newfoundland. *Tectonophysics* 468, 131–148. <https://doi.org/10.1016/j.tecto.2008.05.030>.
- Van Wijk, J.W., Cloetingh, S.A.P.L., 2002. Basin migration caused by slow lithospheric extension. *Earth Planet Sci. Lett.* 198 (3–4), 275–288.
- Walker, I.M., Berry, K.A., Bruce, J.R., Bystøl, L., Snow, J.H., 1997. Structural modelling of regional depth profiles in the Vøring Basin, implications for the structural and stratigraphic development of the Norwegian passive margin. *J. Geol. Soc. London* 154, 537–544.
- White, R.S., McKenzie, D.P., 1989. Magmatism at rift zones: the generation of volcanic continental margins and flood basalts. *J. Geophys. Res.* 94, 7685–7729. <https://doi.org/10.1029/JB094iB06p07685>.
- White, R.S., Smith, L.K., Roberts, A.W., Christie, P.A.F., Kusznir, N.J., and the rest of the iSIMM Team, 2008. Lower-crustal intrusion on the North Atlantic continental margin. *Nature* 452, 460–465. <https://doi.org/10.1038/nature06687>.
- White, R.S., Eccles, J.D., Roberts, A.W., 2010. Constraints on volcanism, igneous intrusion and stretching on the Rockall–Faroe continental margin. In: *Petroleum Geology Conference Series*, vol. 7. pp. 831–842. <https://doi.org/10.1144/0070831>.
- Whiteside, W., Wang, B., Bondeson, H., Li, Z., 2013. 3D Imaging from 2D Seismic Data, an Enhanced Methodology, vol. 2013 SEG Technical Program Expanded Abstracts. <https://doi.org/10.1190/segam2013-1148.1>.
- Whitham, A.G., Price, S.P., Koriani, A.M., Kelly, S.R.A., 1999. Cretaceous (post-Valanginian) sedimentation and rift events in NE Greenland (71–77°N). *Geological Society, London, Petroleum Geology Conference series* 5, 325–336. <https://doi.org/10.1144/0050325>.
- Wilson, J.T., 1966. Did the Atlantic close and then re-open? *Nature* 211, 676–681.
- Zastrozhnov, D., Gernigon, L., Gogin, I., Abdelmalak, M.M., Planke, S., Faleide, J.I., Eide, S., Myklebust, R., 2018. Cretaceous-paleocene evolution and crustal structure of the northern Vøring margin (offshore mid-Norway): results from integrated geological and geophysical study. *Tectonics* 37 (2), 497–528.
- Oufi, O., Cannat, M., Horen, H., 2002. Magnetic properties of variably serpentinised abyssal peridotites. *Journal of Geophysical Research-Solid Earth* 107 (B5). <https://doi.org/10.1029/2001JB000549>.
- Heilmann-Clausen, C. (1985). Dinoflagellate stratigraphy of uppermost Danian to Ypresian in the Viborg 1 borehole, central Jylland, Denmark.

Appendix A

The T-Reflection and the Deep Crustal Structure of the Vøring Margin Offshore Mid-Norway

By

Mansour M. Abdelmalak, Jan Inge Faleide, Sverre Planke, Laurent Gernigon, Dmitry Zastrozhnov, Grace E. Shephard & Reidun Myklebust

Tectonics, 36, 2497–2523

RESEARCH ARTICLE

10.1002/2017TC004617

Key Points:

- We address the relationships between the lower crustal *T*-Reflection and the high-velocity and high-density Lower Crustal Body (LCB) on the Vøring Margin using integrated multidisciplinary geophysical data sets
- The *T*-Reflection represents a high impedance boundary associated with intruded sills with different shapes and characters
- The LCB is made of a mixture of pre- and syn-breakup mafic and ultramafic rocks associated with older metamorphic rocks such as granulites and eclogites

Correspondence to:

M. M. Abdelmalak,
m.m.abdelmalak@geo.uio.no;
abdelmalak@vbpr.no;
abdelmalak_mansour@yahoo.fr

Citation:

Abdelmalak, M. M., Faleide, J. I., Planke, S., Gernigon, L., Zastrozhnov, D., Shephard, G. E., & Myklebust, R. (2017). The *T*-Reflection and the deep crustal structure of the Vøring Margin, offshore mid-Norway. *Tectonics*, 36, 2497–2523. <https://doi.org/10.1002/2017TC004617>

Received 12 APR 2017

Accepted 7 OCT 2017

Accepted article online 17 OCT 2017

Published online 14 NOV 2017

The *T*-Reflection and the Deep Crustal Structure of the Vøring Margin, Offshore mid-Norway

M. M. Abdelmalak^{1,2}, J. I. Faleide¹, S. Planke^{1,2}, L. Gernigon³, D. Zastrozhnov^{1,2}, G. E. Shephard¹, and R. Myklebust⁴

¹Centre for Earth Evolution and Dynamics (CEED), University of Oslo, Oslo, Norway, ²Volcanic Basin Petroleum Research (VBPR), Oslo Science Park, Oslo, Norway, ³Geological Survey of Norway (NGU), Trondheim, Norway, ⁴TGS, Asker, Norway

Abstract Seismic reflection data along volcanic passive margins frequently provide imaging of strong and laterally continuous reflections in the middle and lower crust. We have completed a detailed 2-D seismic interpretation of the deep crustal structure of the Vøring Margin, offshore mid-Norway, where high-quality seismic data allow the identification of high-amplitude reflections, locally referred to as the *T*-Reflection. Using a dense seismic grid, we have mapped the geometry of the *T*-Reflection in order to compare it with filtered Bouguer gravity anomalies and seismic refraction data. The *T*-Reflection is identified between 7 and 10 s. Sometimes it consists of one single smooth reflection. However, it is frequently associated with a set of rough multiple reflections displaying discontinuous segments with varying geometries, amplitudes, and contact relationships. The *T*-Reflection seems to be connected to deep sill networks and is locally identified at the continuation of basement high structures or terminates over fractures and faults. The *T*-Reflection presents a low magnetic signal. The spatial correlation between the filtered positive Bouguer gravity anomalies and the deep dome-shaped reflections indicates that the latter represent a high-impedance boundary contrast associated with a high-density and high-velocity body. In ~50% of the outer Vøring Margin, the depth of the mapped *T*-Reflection is found to correspond to the depth of the top of the Lower Crustal Body (LCB), which is characterized by high *P* wave velocities (>7 km/s). We present a tectonic scenario, where a large part of the deep crustal structure is composed of preserved upper continental crustal blocks and middle to lower crustal lenses of inherited high-grade metamorphic rocks. Deep intrusions into the faulted crustal blocks are responsible for the rough character of the *T*-Reflection, whereas intrusions into the ductile lower crust and detachment faults are likely responsible for its smoother character. Deep magma intrusions can be responsible for regional metamorphic processes leading to an increasing velocity of the lower crust to more than 7 km/s. The result is a heterogeneous LCB that likely represents a complex mixture of pre- to syn-breakup mafic and ultramafic rocks (cumulates and sills) and old metamorphic rocks such as granulites and eclogites. An increasing degree of melting toward the breakup axis is responsible for an increasing proportion of cumulates and sill intrusions in the lower crust.

1. Introduction

The increased availability of high-quality seismic and potential field data, combined with deep sea drilling and onshore analogue studies, has shown that extensional processes can lead to complex crustal configurations, depending on lithospheric composition, thermal structure and heat transfer, far field stresses, extension rate and duration, mantle flow, and structural inheritance (Clerc et al., 2017; England, 1983; Koptev et al., 2015; Kusznir & Park, 1987; Lavecchia et al., 2017; Reston, 2007; Rosenbaum et al., 2008; Ziegler & Cloetingh, 2004). Many studies have led to a range of conceptual and numerical models of plate breakup at magma-poor margins (Brune et al., 2017; Huisman & Beaumont, 2003; Lavie & Manatschal, 2006; Pérez-Gussinyé, 2012). However, more than 50% of passive margins worldwide are interpreted to be magma-rich (Coffin & Eldholm, 1994; Menzies et al., 2002; Skogseid, 2001). These are characterized by massive occurrence of mafic extrusive and intrusive rocks, emplaced before and during plate breakup (e.g., Abdelmalak, 2010; Abdelmalak et al., 2012; Eldholm & Grue, 1994; Geoffroy, 2005), and playing a major role in determining the evolution and the deep structure of magma-rich margins. Seismic reflection data along volcanic passive margins frequently provide imaging of strong and laterally continuous crustal reflections in the middle and lower crusts (e.g., Clerc et al., 2015; Kusznir et al., 2015).

In the NE Atlantic (Figure 1A), wide-angle seismic surveying across most of the conjugate volcanic margins revealed high-velocity layers at the base of the crust referred to as Lower Crustal Bodies (LCBs; Holbrook

Figure 1. (A) Onshore and offshore distribution of the basalt flows in the North Atlantic Igneous Province (Abdelmalak et al., 2014). The Seaward Dipping Reflectors (SDR) locations and different refraction profiles used in this study are indicated (see the text for data references). (B) 3-D reconstruction at C21 (~47 Ma) of the crustal structure across the Vøring Margin (Faleide et al., 2008) and its conjugate Greenland margin in the Thetis Basin (Schlindwein & Jokat, 1999; Voss & Jokat, 2007). The magnetic anomalies are defined using released magnetic data EMAG2 (Maus et al., 2009). The continent–ocean boundary (COB) is indicated for the conjugate margins. Dashed black lines indicate shear zones. NGR: North Gjallar Ridge; TR: T-Reflection (dashed red line). AR: Aegir Ridge; BK: Bløseville Kyst; BVP: Boreas Volcanic Province; DB: Danmarkshavn Basin; EGFZ: East Greenland Fault Zone; FB: Foster Basin; FVP: Foster Volcanic Province; HB: Harstad Basin; HwH: Hold with Hope; JL: Jamson Land; JMFZ: Jan Mayen Fracture Zone; JMMC: Jan Mayen Micro-Continent; JMR: Jan Mayen Ridge; S: Shannon; TB: Thetis Basin; TØ: Traill Ø; TØIC: Traill Ø Igneous Complex; VVP: Vestbakken Volcanic Province; WJMFZ: West Jan Mayen Fracture Zone.

et al., 2001; Klingelhöfer et al., 2005; Mjelde et al., 2007; Mjelde, Raum, et al., 2009; Mjelde, Raum, Myhren, et al., 2005; Roberts et al., 2009; Schiffer et al., 2016; Voss & Jokat, 2007; Voss et al., 2009; Figure 1B). LCBs are characterized by P wave velocities (V_p) of 7.1 to 7.7 km/s and V_p/V_s ratios ranging between 1.8 and 1.9 (Mjelde, Raum, et al., 2003; O'Reilly et al., 1996). The nature and origin of the LCBs are still controversial and probably more complex than hitherto thought (e.g., Ebbing et al., 2006; Gernigon et al., 2004; Mjelde et al., 2016). Early studies have proposed that the anomalously high velocities in the LCBs represent massive mafic magmatic underplating emplaced during the final stage of rifting and continental breakup. In this context, LCBs could be explained by magma showing high concentrations of MgO, which in turn may reflect high asthenospheric temperatures and/or compositional inhomogeneities in the asthenospheric source (e.g., R. S. White & McKenzie, 1989).

Lower Crustal Bodies are often located along the continent–ocean transition (COT; Eldholm et al., 2000) but can extend continentward, outside the identified volcanic province (Kvarven et al., 2014; Mjelde et al., 2016; Nirrengarten et al., 2014). In recent years, discussion about the interpretation of the LCBs focuses on its nature in terms of magmatic versus inherited and/or serpentinized material (Ebbing et al., 2006; Fichler et al., 2011; Lundin & Doré, 2011; Mjelde et al., 2002; Reynisson et al., 2010; Wangen et al., 2011). Given the strong overlap of the geophysical values attributed to continental crust, as well as mafic and ultramafic rocks, the geophysical modeling of LCBs alone does not easily permit the preference of one interpretation over the other, but rather, the observed correlation with structure allows the discussion of scenarios and realistic hypotheses (Gernigon et al., 2004; Ren et al., 1998; Schiffer et al., 2016). In such conditions, the architecture of rifted margins is still the subject of debate in order to explain its structural, stratigraphic, and magmatic evolution.

On the other hand, the interactions between magmatism, deep structures, and extensional processes are also not completely understood. Generally, transport of mafic magma through the upper brittle crust is widely viewed as being dominated by dykes (Lister & Kerr, 1991; Rubin, 1995), which are often invoked as the main feeders for flood basalt sequences (Abdelmalak et al., 2015; Coffin & Eldholm, 1994; Ernst et al., 1995; Klausen & Larsen, 2002). In contrast to dykes, bedding-parallel sills are widely regarded as having only a limited role in vertical magma transport. How deep sill complexes along volcanic margin are associated with LCBs, or not, is a challenging problem. To address these issues, it is common to supplement the reflection-based seismic interpretation with velocity and density models derived from refraction and potential field studies (e.g., Corseri et al., 2017).

High-amplitude deep crustal reflections, located at depths >6 – 7 s, have been imaged in the outer Vøring Basin. Reflections display a domal shape below the South and North Gjallar ridges and have been interpreted as the top of a core complex of melted crust triggered by underplating and labelled “mid-crustal convex-up reflector” (Lundin & Doré, 1997) or “MCD” for “middle crustal dome” (Ren et al., 1998) in the North Gjallar Ridge. In later studies, these reflections have been regionally mapped along the outer Vøring Basin and named the T -Reflection (Gernigon et al., 2001). In the North and South Gjallar ridges, Gernigon et al. (2003) further discussed the definition and structural interpretation of the T -Reflection and suggested that it coincides with the top of the continental part of the LCB geophysically modeled at approximately the same depth based on ocean bottom seismometer (OBS) data (Mjelde, Faleide, et al., 2009; Rouzo et al., 2006; Figure 1). The part of the LCB imaged beneath the North Gjallar Ridge appears to have been emplaced before the main volcanic event and influenced the faulting and distal development of the sedimentary basin at least 10–15 Ma before the breakup and the main magmatic event (Gernigon et al., 2004).

The aim of this study was to determine the nature and origin of the deep crust along the conjugate volcanic margins of the Norwegian–Greenland Sea. Despite expansive coverage of seismic data along the mid-Norwegian Margin (Figure 3B), several questions about the deep structure of rifted margins remain unresolved: (1) What is the origin of the T -Reflection? (2) What are the relationships between the T -Reflection, basin deformation, continental breakup, and the LCB? (3) What is the origin of the LCB in the continental and distal parts of the rifted margin? (4) Are LCBs fully representative of breakup-related igneous rocks? To answer these questions, we have investigated the structure, nature, and possible origins of the T -Reflection based on the use of new and combined geophysical data sets (reflection and refraction seismic, gravity, and magnetic data).

2. Geological Setting

The conjugate volcanic rifted margins along the Norwegian–Greenland Sea are parts of the North Atlantic large igneous province formed during the continental breakup in the Paleogene (Eldholm et al., 2000;

Figure 2. Simplified map of the mid-Norwegian margin showing the main tectonic units, fault zones (using our new seismic interpretation and modified from Blystad et al., 1995; and Gernigon et al., 2003), and volcanic seismic facies units (using our new seismic interpretation and modified from Berndt et al., 2001). FG: Fenris Graben; GH: Grimm High; HG/ND: Hel Graben/Nagfar Dome; NR: Norland Ridge; NS: Någrind Syncline; RB: Rån Basin; RR: Rån Ridge; SDRs: Seward dipping reflectors; SH: Skoll High; TB: Træna Basin; UH: Utgard High; VS: Vigrid Syncline; YH: Ygg High.

Meyer et al., 2007; Saunders et al., 1997; Figure 1). The onset of continental breakup marked the culmination of a ~300 Ma period of extension and basin formation subsequent to the Caledonian orogeny (Doré et al., 1999; Skogseid et al., 2000; Tsikalas et al., 2008; Ziegler, 1988). The main extensional events occurred in Late Paleozoic–Triassic, Late Jurassic–Early/Middle Cretaceous, and Late Cretaceous–Paleocene times (Brekke, 2000; Eldholm & Grue, 1994; Gernigon et al., 2004; Lundin & Doré, 2005). Through the Paleozoic and Mesozoic, lithospheric thinning resulted in the formation of large sedimentary basins controlled by normal faults (e.g., Faleide et al., 2008; Tsikalas et al., 2012). Subsidence was especially important during the Cretaceous, giving rise to the accumulation of up to 8 km of sediments in local depocenters in the Vøring and Møre basins (Blystad et al., 1995; Brekke, 2000; Lien, 2005; Scheck-Wenderoth et al., 2007).

On the Vøring Margin, the Late Jurassic–Early/mid-Cretaceous extension and the Late Cretaceous extension formed large-scale rift blocks and structural highs (Early Cretaceous ridges and highs: Rån Ridge, Utgard High; Late Cretaceous–Early Paleocene ridges and highs: North and South Gjallar ridges, Nyk High, Ygg High, Grimm High, and Skoll High) separated by subbasins such as the Træna Basin, the Någrind Syncline, the Rån Basin, the Rås Basin, the Vigrid Syncline, and the Fenris Graben (Abdelmalak, Planke, et al., 2016; Figure 2). During the Late Cretaceous–Paleocene, the locus of maximum extension migrated NW toward the zone of the future continental separation (Skogseid et al., 2000). The final continental breakup recorded in the Vøring Marginal High occurred at early Eocene (~56–55 Ma according to Gradstein et al., 2012, time-scale) and resulted in voluminous extrusive and intrusive igneous activities within the adjacent

Figure 3. (A) Seismic reflection data used in this study. (B) Bouguer residual gravity anomaly, 50 km high-pass filtered. White shaded areas correspond to structural highs. Black lines indicate faults. Gravity data courtesy of TGS. FFC: Fles Fault Complex; FG: Fenris Graben; GH: Grimm High; GS: Gleipne Saddle; HeH: Hevring High; HG/ND: Hel Graben/Nagfar Dome; NH: Nyk High; NGR: North Gjallar Ridge; NR: Norland Ridge; NS: Någrind Syncline; MH: Mimir High; MMH: Møre Marginal High; RB: Rån Basin; RR: Rån Ridge; SGR: South Gjallar Ridge; SH: Skoll High; SR: Slettringen Ridge; TB: Træna Basin; UH: Utgard High; UnH: Unn High; VMH: Vøring Marginal High; VS: Vigrid Syncline; YH: Ygg High.

sedimentary basins and preexisting continental crust (Breivik et al., 2014; Eldholm & Grue, 1994; Hinz, 1981; Mjelde et al., 2007; Mutter et al., 1982; Planke et al., 2005; R. S. White & McKenzie, 1989; Figure 2).

On the NE Greenland side, extrusive volcanism is also exposed in several places onshore (e.g., Brooks, 2011; Geissler et al., 2016; Larsen et al., 2014; Tegner et al., 1998; e.g., Shannon Island, Blosseville Kyst area, Hold with Hope; Figure 1A). Offshore, a thin volcanic complex, extending approximately 110 km, is mapped along the Thetis Margin. Seaward Dipping Reflectors (SDRs) are observed east of this volcanic complex where they are located approximately along the continent–ocean boundary (Abdelmalak, Planke, et al., 2016). The SDRs pinch out to the south against the Foster Volcanic Province, which is bounded by the Traill Ø Igneous Complex to the south and the Foster Basin to the north (Figure 1A). The Foster Volcanic Province consists of thick volcanic flows draped by Eocene and younger sediments (Reynolds et al., 2017). However, it has been classified as transitional crust by Voss and Jokat (2007). Nonetheless, we consider the Foster Volcanic Province as an oceanic plateau. Further south, the Traill Ø Igneous Complex defines a trend parallel to the West Jan Mayen Fracture Zone. The formation of this igneous complex is linked to the breakup between East Greenland and the Jan Mayen Micro-Continent during the Late Eocene–Oligocene time. North of the East Greenland Fracture Zone, the Boreas Volcanic Province is considered to be the conjugate to the Vestbakken Volcanic Province in the SW Barents Sea (Figure 1A).

3. Data and Methods

3.1. Data

This work is based on the interpretation of a dense grid of more than 1,200 regional 2-D seismic lines with a total length of more than 150,000 km and with a spacing ranging between 0.2 and 2 km (Figure 3A). This data set includes recently acquired as well as reprocessed high-quality seismic reflection data that provide improved imaging of the deeper parts of the margin. The new data include, notably, 10,000 km of the MNR-11 seismic survey acquired in 2011 by TGS and Fugro. Our database also comprises previous long-offset seismic surveys including in particular the VMT-95, VBT-94, and the GMNR-94 surveys, which were recorded with 11 to 14 s (two-way travel time).

In addition, more than 40 refraction profiles from the Møre, Vøring, and Lofoten-Vesterålen margins have been used in this study (e.g., Breivik et al., 2009, 2014; Mjelde et al., 1996, 2007; Mjelde, Raum, et al., 2009; Mjelde, Raum, Breivik, et al., 2005; Raum et al., 2002, 2006; Figure 3A). Conjugate refraction profiles from NE Greenland (e.g., Hermann, 2013; Schlindwein & Jokat, 1999; Schmidt-Aursch & Jokat, 2005; Voss & Jokat, 2007; Voss et al., 2009; Weigel et al., 1995) and Jan Mayen (e.g., Breivik et al., 2012) margins have also been considered (Figure 1A). These data have been used to compare the different velocity models of the crust and the difference between the reflection and refraction seismic data.

Ship-borne gravity and magnetic data have been acquired along most of the seismic profiles of the mid-Norwegian margin. The data have been processed and gridded by TGS. The data have been combined with regional public grids of satellite gravity (Sandwell & Smith, 2009), magnetic compilations (Maus et al., 2009; Verhoef et al., 1996), and bathymetry (GEBCO centenary edition). The gravity data have been Bouguer corrected using a correction density of $2,200 \text{ kg/m}^3$. Both the gravity and magnetic grids have been high-pass filtered with cutoff wavelengths of 50, 100, 200, and 400 km.

3.2. Methods

The regional seismic interpretation was carried out using the SMT Kingdom Suite software (www.ihs.com). We mapped the *T*-Reflection horizon in the outer Vøring Margin based on its strong amplitude characteristics. The seismic line density was high enough to laterally correlate the deep crustal reflections and to generate grids.

Bouguer-corrected gravity maps are useful for geological interpretations as the effect of water depth variations is mainly removed (e.g., Berndt, 2002). The Bouguer anomaly map mainly reflects variations in crustal thickness. A high-pass filtered Bouguer anomaly map with a cutoff wavelength of 50 km emphasizes the deep basin and midcrustal density variations, with typical source depths of less than 10 km. The anomaly pattern (e.g., wavelength and amplitude) provides information about fault blocks and configuration of subbasins (e.g., Berndt, 2002; Figure 3B). Magnetic anomaly maps may reveal large tectonic trends and define oceanic crust with characteristic sea floor spreading anomalies. High-pass filtered magnetic anomaly maps usually highlight middle and upper crustal variations. The wavelength and amplitude of anomalies provide information about depth to source and magnetic properties. The high-amplitude positive magnetic anomalies on the shelf often correspond to basement ridges or shallow igneous rocks, whereas negative anomalies are generally associated with deep sedimentary strata. High-pass filtered potential field data were scaled to two-way travel time, converted to pseudohorizons, and loaded into the workstation to facilitate the joint interpretation of seismic and potential field data (e.g., Planke et al., 2015).

To improve the 2-D deep crustal interpretation, several key refraction profiles were time-converted (to two-way travel time) and loaded into the workstation. This approach allowed us to display the crustal and the velocity structure along the seismic reflection data for comparison. The grid generated from seismic horizon picking (the *T*-Reflection grid) was depth converted with HiQbeTM (<http://www.first-geo.com/products/hiqbe/norway>), a regional seismic velocity cube covering the mid-Norwegian margin and compiled from 536 seismic stacking velocity data sets and 230 check shots from wells and vertical seismic profile data. The stacking velocities used for the depth conversion are constraining the velocity structure of the sedimentary basin where reflections are imaged. At greater depths, a delta-anisotropy function was used for depth conversion (e.g., Thomsen, 1986). The stacking velocities are, however, lower than the OBS velocities because of differences in the frequency content and the paths of wave propagation (Thomsen, 1986). The HiQbeTM velocity cube intends to replicate the velocity for true vertical wave propagation, while OBS data are modeled with a great horizontal component often without taking into concern the anisotropic effect (horizontal being greater than vertical velocity, in general). Consequently, the HiQbeTM is presented with average velocities (from sea surface down to target depth), while OBS data are mostly presented as interval velocities (layer-specific velocities) where interval velocities are often higher (~10–15%) than the average velocities. Combined with uncertainties related to the different modeling techniques, additional errors could be driven by uncertainties in the OBS velocity forward modeling. These considerations will lead to obvious uncertainties and differences when comparing refraction and depth-converted reflection data. However, by comparing the stacking velocities and OBS velocity in selected 1-D profiles in the Vøring Margin, we found comparable results in most of the cases.

4. Results

4.1. The Vøring Margin Structure

The 50 km high-pass filtered Bouguer gravity anomaly map reveals a complex pattern of positive and negative amplitude anomalies along the Vøring Margin (Figure 3B). In the Vøring Basin, several elongated negative gravity anomalies, striking NE–SW, are identified and correspond to the Hel Graben, Fenris Graben, Vigrid Syncline, Någrind Syncline, Træna Basin, and Rås Basin. These different subbasins are separated by NE–SW elongated positive anomalies, corresponding to the Nyk High, the North and South Gjallar ridges, and the Utgard High. The Rån Ridge approximately coincides with an ENE–WSW positive anomaly. High-amplitude rounded positive gravity anomalies are distributed along the Vøring Escarpment and correspond to positive elongated NE-trending magnetic anomalies (Abdelmalak, Planke, et al., 2016; Planke et al., 2017). Oceanward of the Vøring Escarpment, the gravity anomalies likely represent intra- and subbasalt structural highs. These anomalies correspond to Skoll High, Grimm High, and Ygg High, as also observed on seismic data. Toward the SW Vøring Basin, the gravity anomalies change orientation from NE–SE (Unn High) to approximately N–S (Slettringen Ridge; Figure 3B).

4.2. Reflection Data: The T-Reflection

From reprocessed and high-quality seismic data in the Vøring Margin, a more accurate interpretation from the shallow parts to the deeper parts of the margin, down to the *T*-Reflection, is now possible. Examples of the reprocessed seismic data, shown in Figure 4, illustrate the main elements discussed in this paper. For the shallow parts, the base Tertiary unconformity represents a regional erosional surface onlapped by Paleogene sediments. The base Cretaceous unconformity (BCU) is robustly mapped landward of the basalt. The BCU is generally considered to record the rift climax activity of the major Late Jurassic–Early Cretaceous extensional phase and is often used as a key regional marker to constrain the spatial and temporal history of the Late Jurassic–Early Cretaceous rift system (Blystad et al., 1995). Below the basalt, the seismic imaging is of poor quality but confirms locally that volcanic basins do exist underneath the lava flows. The BCU is deeper below the Rån Basin and the Vigrid Syncline and shallows below the South Gjallar and Rån ridges. Several low-angle faults characterizing the Cretaceous and the pre-Cretaceous sediments are assumed to be of at least Early Campanian–Early Paleocene age (Gernigon et al., 2003; Ren et al., 2003). Jurassic to Early/middle Cretaceous faulting was previously observed on the Rån Ridge area where the BCU and older syn-rift sequences could be clearly identified near the basalts (Gernigon et al., 2001, 2003). Abundant Paleogene sill intrusions are also identified within the Cretaceous and pre-Cretaceous sequences.

For the deeper part, the outer Vøring Basin displays a well-defined high-amplitude and strong reflection with variable continuity and coherence referred to as the *T*-Reflection (Gernigon et al., 2003, 2004). This high-amplitude deep crustal reflection is observed at depths ranging between 7 and 10 s (Figures 4 and 5). In the seismic data, the *T*-Reflection is sometimes smooth and is primarily identified on a single event (Figures 5A and 5B). Sometimes the *T*-Reflection is associated with a series of reflections (Figure 5D). It is therefore discontinuous and displays distinct steep and flat segments with varying geometries, amplitude, and contact relationships (Figures 5C and 5D).

In this paper we discuss the *T*-Reflection as a single characteristic feature, regardless of its shape and number of cycles. Our seismic interpretation reveals that the *T*-Reflection is mainly identified in the outer Vøring Basin and extends between the Vøring Escarpment and Fles Fault Complex and Utgard High (Figure 6A). West of the Vøring Escarpment, the *T*-Reflection is difficult to map beneath the thick volcanic extrusives that give rise to scattering of the seismic energy. The *T*-Reflection is clearly mapped in the North Gjallar Ridge where the high-amplitude reflection is rather continuous, shows a smooth shape, and can be mapped laterally over 50 km (Figure 6B). In the southwest Vøring Basin, the *T*-Reflection can be identified over more than 150 km along the Rån Basin, Rån Ridge, and Vigrid Syncline (Figure 6B).

Beneath the northwestern parts of the North Gjallar, South Gjallar, and Rån ridges, the *T*-Reflection exhibits domal shapes of more than 20 km in diameter, with its shallowest part between 7 and 8 s. Compared to the first map of the *T*-Reflection (Gernigon et al., 2003), our new and denser seismic data set permits the identification of additional subtle variations across the Rån and the South Gjallar ridges. Less prominent *T*-Reflection domes (Figure 6A) are now observed below the so-called Hevring High, Unn High, and

Figure 4. (A) Interpreted seismic section (in two-way travel time) showing the smooth *T*-Reflection along the South Gjallar Ridge, the Rån Basin, and the Rån Ridge. (B) Interpreted seismic section (in two-way travel time) showing the discontinuous *T*-Reflection along the Rån Ridge. The 7+ km/s (considered as top LCB) and the 8+ km/s (the seismic Moho) layers are extracted from refraction data and displayed on the seismic reflection data. Sill intrusions are indicated by thin red lines. The 50 km high-pass filtered gravity (B50) and magnetic (M50) data are plotted above the seismic profiles. BCU: base Cretaceous unconformity; FFC: Fles Fault Complex; IM: intra Miocene; TC: top Cretaceous; TO: top Oligocene; TR: *T*-Reflection. See Figure 6 for line location. Data courtesy from TGS.

Slettringen Ridge (Figure 3B). The *T*-Reflection deepens below the Gleipne Saddle, the Vigrid Syncline, and the Hel Graben where the top is mapped around 9–10 s (Figure 6A).

The *T*-Reflection near the Rån Ridge is more ambiguous and consists of multiple reflections and shows a rough character (Figure 6B). As mentioned by Gernigon et al. (2001), the *T*-Reflection in this area could be correlated with high-velocity basement locally intruded by sills; however, its morphology is characterized by a contrasting and ambiguous faulted geometry (Figure 4B) that differs from the smooth character observed below the South and North Gjallar ridges (Figure 6B). Most of the overlying faults on the Rån Ridge seem to have been active during the Early Cretaceous until the Cenomanian where they terminate and were covered by thick Upper Cretaceous sediments (Figure 4B). Older fault activities that affected the pre-Cretaceous sedimentary succession (Triassic–Jurassic?) are also expected (e.g., Gernigon et al., 2003).

Figure 5. Different shapes for the *T*-Reflection. (A, B) Single reflection and smooth shape. (C, D) Multiple reflections and rough shape. Black arrows indicate the *T*-Reflection. See Figure 6 for lines location.

Locally, the *T*-Reflection seems to be connected to deep sill networks (Figure 7A), with evidence of venting (Figure 7B). Close to Slettringen Ridge, the *T*-Reflection presents a rough shape and may represent the continuation of a preexisting basement high (Figure 7C). The *T*-Reflection often disappears or terminates over fractures and faults (Figure 7D).

By draping the 50 km filtered Bouguer anomaly over the *T*-Reflection topography (Figure 6C), a good correlation is found. Generally, the positive gravity signature corresponds to the different *T*-Reflection domes and highs, whereas the negative filtered Bouguer anomalies correspond to the deeper levels of the *T*-Reflection. The well-defined correlation between the *T*-Reflection and the gravity anomaly confirms that the *T*-Reflection represents a high-impedance boundary associated with a high-density body. No magnetic anomalies are associated with the *T*-Reflection domes on the Vøring Margin (Figure 4). The seismic interpretation also reveals that there is a good correlation between the basin structure (subbasins, highs, and ridges) defined at Cretaceous level and the *T*-Reflection geometry.

4.3. Refraction Data: The LCB Spatial Distribution

From the seismic refraction data (Figure 3A), we compiled maps of the seismic Moho depth (Figure 8A) and the depth (Figure 8B) and thickness of the LCB (Figure 8C). Accordingly, we picked the velocity depth values along individual profiles and created isovelocity maps. The Moho is characterized by isovelocities higher than 8.0–8.3 km/s. The Moho depth along the coast of mainland Norway is tied to Fennoscandia profiles (Breivik et al., 2011; Kinck et al., 1993; Kvarven et al., 2014; Stratford et al., 2009). On the Norwegian mainland, the Moho is about 35 km deep; it becomes slightly shallower below the Vøring and Møre basins (25–27 km), deepens again below the Vøring Marginal High, and rises dramatically toward the oceanic domain (9–15 km) (e.g., Planke et al., 1991). The Moho topography shows an apparent sinistral offset along the Jan Mayen Fracture Zone. The Moho depth increases below the Vøring Spur where thick igneous crust is identified (up to 15 km) and is interpreted to be the result of excess melting along the east Jan Mayen Fracture Zone

Figure 6. (A) Time-structure map (in two-way travel time) of the *T*-Reflection grid in the Vøring Margin. (B) Shape and character of the *T*-Reflection. (C) 3-D view of the 50 km high-pass filtered Bouguer anomaly draped above the topography of the *T*-Reflection. Thin black lines indicate faults.

during Early–mid-Eocene time (Gernigon et al., 2009), or alternatively during a younger Miocene magmatic event (Breivik et al., 2014).

The top LCB (Figure 8B) is generally defined as the top of the 7+ km/s velocity layer. The top of the LCB is deeper in the Vøring and Møre basins (depths ranging between 18 and 22 km) and becomes shallower toward the oceanic crust (depths ranging between 10 and 12 km). In the oceanic Lofoten Basin, the top LCB is shallower (between 10 and 16 km) compared to the Vøring and Møre margins. For the LCB thickness map (Figure 8C), we considered V_p velocities between 7.1 and 7.7 km/s as criteria to calculate the LCB thickness. We extended our mapping into the oceanic domain, and we defined the outer limit of the LCB where “normal” oceanic crust (6–8 km thickness) was clearly identified at the location of magnetic chrons C22–C23. The thickness of the LCB varies considerably, from 0 to about 8–9 km, which might be caused by variations in the prebreakup structure and/or by the late “magmatic underplating” process that affected the preexisting crust.

On the Vøring Margin, the thickness of the LCB changes along and across the strike of the seismic profiles. Rounded and elongated shaped features, with thicknesses ranging between 6 and 8 km, characterize the Vøring Basin. On the Vøring Marginal High, local increase of LCB thickness to more than 6 km could be correlated with location of the thicker part of the Seaward Dipping Reflectors. Near the Vøring Transform Margin, the LCB is thinner (less than 2 km) and shows a limited extent. On the Møre Margin, our mapping shows a more regular distribution of the LCB thickness. The thicker part (~5 km) is situated beneath the Møre Marginal High and could be correlated with the extent of the Seaward Dipping Reflectors. The LCB thickness decreases oceanward until disappearing west of magnetic chron C23. A local anomaly below the Vigra High is noticed where the LCB reaches a thickness of 4 km. The LCB thickness decreases toward the limit of sill intrusion in the sedimentary basin (Figure 8C). On the Lofoten–Vesterålen Margin, northeast of the Bivrost Lineament, the LCB has a very limited extent. The thicker part (~4 km) is situated along the continent–ocean boundary, and a rapid decrease in thickness is noticed in the oceanic domain toward magnetic chron C23.

Figure 7. *T*-Reflection and relationship to deep basin structure. (A) Possible connection between *T*-Reflection and the overlying sill network. (B) Possible venting event associated with *T*-Reflection. White shaded area indicates possible venting event. (C) Continuation of the *T*-Reflection at the location of the basement high. (D) The *T*-Reflection terminates along a fault. Black arrows indicate the *T*-Reflection, and red arrows indicate the sill intrusions. We added a transparency around the *T*-Reflection to highlight it in the seismic profile. See Figure 6 for line location.

5. Discussion

5.1. Comparison Between the *T*-Reflection and the LCB Using Seismic Data

The generation of long-offset and high-quality seismic reflection profiles acquired along most of the rifted margins leads to the unraveling of an unexpected variety of structures (e.g., Clerc et al., 2017) and provides unprecedented access to the processes occurring in the middle and lower continental crusts. On the Vøring volcanic margin, offshore mid-Norway, seismic reflection data reveal the existence of strong and laterally continuous reflections referred to as the *T*-Reflection. It has been suggested that the *T*-Reflection on the North Gjallar Ridge matches with the top of the continental part of the LCB (Gernigon et al., 2004); thus, understanding the relationship between the *T*-Reflection and the LCB is of first-order importance to address the deep structure of the margin.

5.1.1. 2-D Comparison

From the time-converted refraction profiles, we have attempted a direct comparison between the seismic reflection and refraction data as illustrated in Figure 9A. The refraction profile 11-96 was shot in a SW-NE direction along the Fenris Graben. The different velocities modeled along the refraction profile are indicated (e.g., Mjelde, Shimamura, et al., 2003). The LCB is indicated by velocities ranging between 7.2 and 7.5 km/s. On the seismic reflection line covering the central part of the refraction profile 11-96, the basaltic layer contributes to an imprecise image of the deeper parts of the sedimentary basin and crustal structure (Figure 9B). The modeled *P* wave velocities allow the identification of different crustal structures (top crystalline basement, LCB, and Moho). Along the refraction profile 11-96, the *T*-Reflection is located at different level (with depths ranging between 8 and 10 s) and displaying different *P* wave velocities. Below the Gleipne Saddle, the *T*-Reflection is deeper than the top of the LCB and shows a level close to the Moho defined by refraction modeling (Figure 9A). Along the Fenris Graben, the *T*-Reflection is located within the LCB layer but often fits

Figure 8. (A) Seismic Moho depth. (B) Top of the Lower Crustal Body (LCB). (C) Thickness map of the LCB. The different maps are compiled and interpolated using all available seismic refraction data. The extrusive extent corresponds to the breakup related volcanics defined from the volcanostratigraphic volcanic facies units. MMH: Møre Marginal High; VMH: Vøring Marginal High; VTM: Vøring Transform Margin; VS: Vøring Spur.

the top of the LCB or is close to it. Along the Rån Ridge, the *T*-Reflection is well defined and can be mapped continuously. On the refraction profile 6-00 (Raum et al., 2006; Figure 9C), the *T*-Reflection is associated with velocities ranging between 5 and 6 km/s. However, *P* wave velocity of 7+ km/s have been modeled at the level of the *T*-Reflection, along the same transect (Rouzo et al., 2006), showing already that OBS modeling can produce a large range of uncertainties.

Figure 9. (A) Time-converted refraction profile 11-96 (Mjelde, Shimamura, et al., 2003) overlain by (B) the coincident seismic profile. (C) T-Reflection plotted above the refraction profile 6-00 (Raum et al., 2006). The 50 km high-pass filtered gravity (B50) and magnetic (M50) data are plotted above the seismic profiles. TR: T-Reflection (red line). See Figure 6 for lines location.

Figure 10. (A) Depth-converted T-Reflection. (B) Difference map between top LCB and depth-converted T-Reflection. The interval difference showing values ranging between -2500 m and $+2500$ m (yellow color) correspond to the area where T-Reflection and top LCB match. A positive (blue color) or negative (red, orange, and green colors) difference indicates that the T-Reflection is deeper or shallower than top LCB, respectively.

Figure 11. Deep sill network in the Vøring Margin. The 50 km high-pass filtered gravity (B50) and magnetic (M50) data are plotted above the seismic profiles. Different pipe structures are identified and interpreted as hydrothermal vent complexes and seepages. See Figure 6 for location.

5.1.2. 3-D Comparison

The depth-converted seismic data shows that the *T*-Reflection depth ranges between 11 and 19 km in the Vøring Basin (Figure 10A). The *T*-Reflection depth ranges between 11 and 13 km above the North and South Gjallar ridges and the Rån Ridge. A deeper *T*-Reflection, between 13 and 15 km, is interpreted below the Hevring High and the Unn High and deepens in Vigrid Syncline and Hel Graben/Nagfar Dome to 15–19 km.

The difference map between the top LCB and the depth-converted *T*-Reflection is also shown in Figure 10B. Locally, the differences are quite significant and can be explained by uncertainties in the different techniques used in depth conversion (see section 3.2). Nonetheless, we consider an error of ± 2500 m when comparing the top LCB compiled from refraction data and the depth-converted *T*-Reflection using stacking velocities (Figure 10B). In this context, the interval with values ranging between -2500 m and $+2500$ m corresponds to the area where the *T*-Reflection and the top LCB likely match. This area corresponds approximately to 50% of the area where the *T*-Reflection is mapped on the Vøring Margin. In the Vigrid Syncline, the Hel Graben, and the Gleipne Saddle, the mapped *T*-Reflection is generally deeper than the top LCB based on the refraction data. This difference could possibly be related to depth conversion issues especially in the deepest part of the sedimentary basin and basement where the stacking velocities are more uncertain. In the North and South Gjallar ridges, the *T*-Reflection depth mostly corresponds to the top LCB. The *T*-Reflection is shallower than the top LCB in the southwestern corner of the Vøring Basin. In this area, which has been affected by strike-slip tectonic movement, the LCB is thin or absent (Berndt, Mjelde, et al., 2001). Furthermore, high V_p velocities on the Rån Ridge have been interpreted as indication of the existence of a deep seated eclogite body (Figure 9C; e.g., Raum et al., 2006). However, we note that an alternative model suggests that the *T*-Reflection coincides with velocities of up to 7 km/s (Rouzo et al., 2006).

5.2. Spatial Correlation Between Sill Intrusions, *T*-Reflection, and LCB

On the basis of our new observations at the deeper level of seismic profiles, the distribution of deep sill networks becomes clearer. An example of a complex sill network in the Vøring Basin is illustrated in Figure 11. Sills are easily identified by their characteristic high-amplitude seismic response, saucer shape, discordant

reflection geometry, and abrupt terminations (e.g., Planke et al., 2005). They are also associated with hydrothermal vent complexes that were mainly formed by explosive eruptions of gases, liquids, and sediments during sills emplacement (e.g., Svensen et al., 2004). Sills intruded at deeper crustal levels are larger than those identified at shallower levels in terms of both thickness and lateral extent (Figure 11). A characteristic feature is that most of the sills are vertically connected via magmatic junctions (e.g., Cartwright & Hansen, 2006). This observation indicates that individual sills are often interconnected and part of larger sill complexes (e.g., Hansen et al., 2004). In the North Gjallar Ridge, the sill network can be traced vertically from Paleocene sediments down to the T-Reflection level inferred to represent the top of the Lower Crustal Body. Wilson and Wheeler (2002) previously inferred a connection between the LCB and the overlying sill complex noting that magma “appears to move from sill to sill via a complex system of narrow, dyke-like feeders.”. Fractures and faults may play an important role in magma transfer as it is easier for magma to use a zone of weakness. In Figure 11 an interesting feature can be observed, the sill network appears to terminate against the Surt Lineament and use this weakness zone as a conduit to shallower levels.

In the Vøring Basin, the sedimentary sequences above the LCB are strongly intruded by sills (e.g., Brekke, 2000; Planke et al., 2005), and the concentration of sills appears to be highest in the central part of the basin where the LCB is thicker (e.g., Mjelde, Faleide, et al., 2009). Sills are generally found at their deepest position and lowest stratigraphic levels in the eastern part of the Vøring Basin (e.g., Planke et al., 2005) and typically step up toward the west to their highest levels close to the Vøring Escarpment. However, subsill imaging remains difficult partly due to transmission loss across high-impedance contrast boundaries. Complicating factors, such as offset-dependent tuning, complex 3-D geometries of sills, velocity increases in the metamorphic aureole, and structural variations in the overburden, commonly make it difficult to obtain good resolution criteria (Planke et al., 2005, 2015). It is often possible to interpret several sill intrusions, but the continuity of deeper sills is commonly lost when the thickness and/or number of the overlying sills increases. In addition, high-amplitude seismic events below the uppermost sill are commonly not primary events but rather peg-leg multiples or converted waves (e.g., Berndt et al., 2000).

In the Vøring Basin, exploration well 6607/5-2 at the Utgard High penetrated two microgabbro sills intruded into Upper Cretaceous mudstone and sandstone lithologies (Neumann et al., 2013). Sonic logs from sills show an average velocity of 7 km/s (Planke et al., 2005, 2015). Multichannel seismic reflection profiles in the Hel Graben reveal a sill complex at approximately 5 km depth associated with exceptionally high, 7.4 km/s seismic wide-angle velocities (Berndt et al., 2000). Such velocities are within the velocity range that characterizes the LCB in volcanic passive margins. Interestingly, in the mid-Norwegian margin, the sill extent correlates very well the LCB extent (Figure 8C).

5.3. The Nature of the LCB in the Continental Domain

Unraveling the nature of the LCB is crucial to understand the deep structure and tectonic evolution of the volcanic margins and its implications in terms of crustal thinning, heat flow, and vertical motion. Depending on alternative interpretations, the magmatic rock volume of the North Atlantic large igneous province may be significantly less than previously thought, considering that 60–80% of the volume of this large igneous province has been estimated to reside in the LCB (Eldholm & Grue, 1994; R. S. White et al., 1987).

Refraction data along the Vøring Margin indicates large variations in thickness and velocity within the LCB, which have been proposed to relate to the distribution of mantle melts via feeder dykes (Mjelde et al., 2002). The velocities of this layer vary between 7.0 and 7.7 km/s and have often been interpreted as a function of differences in the magma composition due to inhomogeneities in the asthenospheric source or as a function of mafic differentiation inside the LCB (Mjelde, Raum, Breivik, et al., 2005). The variability in LCB dimensions and velocities demand a consideration of properties in terms of geodynamic setting. First, similarity in LCB velocity ranges does not necessarily imply the same emplacement process and composition. Secondly, temperature–pressure relations may, in some settings, allow secondary metamorphic processes during or subsequent to emplacement.

Geochemical analyses of the Utgard High sills demonstrate that the LCB can be explained as a heterogeneous mixture of cumulates associated with the opening related magmatism and less dense rocks such as old continental basement (Neumann et al., 2013). In this context, the “cumulate” could be defined as a rock that forms via partial crystallization of a melt, after which the remaining melt is removed from the system.

Because igneous cumulates commonly contain more MgO than their parental liquids and because rocks with higher MgO have a higher seismic velocity, cumulate lower crust formed by fractional crystallization should have the highest velocity. This provides an upper limit on the deviation of lower crustal velocity from bulk crustal velocity (e.g., Korenaga et al., 2002).

Wangen et al. (2011) also concluded that a scenario involving a LCB constituting solely of underplated material would require an unrealistic amount of extension. A scenario where underplating or lower crustal intrusion (magmatic addition) accounts for maximum half the LCB is more likely. Berndt et al. (2000) noted that the geophysical expression of the high-velocity mafic lower crust is complex and may alternatively be interpreted as a preexisting crust injected by scattered high-velocity sills. A model involving a mixture of rocks with contrasting physical properties is in agreement with the large variation in V_p (7.1–7.8 km/s) and relatively low V_p/V_s ratios (1.7–1.85) documented within the LCB by Mjelde et al. (2002).

P wave velocities in low-grade crystalline rocks are found to increase from ~6.0 km/s in granite (felsic) to ~6.9 in gabbro (mafic). The V_p/V_s ratio increases correspondingly from ~1.70 to ~1.85 (e.g., Holbrook et al., 1992). Increasing the P wave velocity in gabbro to the range observed in the Vøring Basin can be achieved by (1) increasing the temperature of the melt and thereby the MgO content or (2) increasing the metamorphic grade to granulite facies (Hurich et al., 2001). The observed densities, V_p , and V_p/V_s ratios are compatible with both hypotheses.

Lower crustal P wave velocities in the range of 7.2–7.6 km/s are also compatible with a crustal layer comprising partly serpentinized peridotites, the existence of which has been postulated in the deeper parts of the Møre Basin (Osmundsen & Ebbing, 2008; Reynisson et al., 2010). Lundin and Doré (2011) proposed that the anomalously wide LCB could have two origins: an inner part made up of partially serpentinized mantle related to Early Cretaceous hyperextension and an outer narrower part of intruded lower crust that overprinted the margin edge during breakup. These authors argued that partial serpentinization (10–30%) can readily account for the P wave velocities of the LCB beneath the Cretaceous basin (e.g., Escartín et al., 2001). Partial serpentinization probably takes place during or soon after hyperextension (Skelton et al., 2005), and a LCB formed in this manner would explain a temporal and spatial relationship to the crustal thinning and overlying basin fill.

However, the low V_p/V_s ratios observed on the Vøring Marginal High and in the northern Vøring Basin could be inconsistent with serpentinized peridotites (Mjelde et al., 2002; Mjelde, Raum, et al., 2003). Despite uncertainties about V_p/V_s estimates and interpretations, the authors also argued against the serpentinized peridotite hypothesis in the central and southern Vøring Basin. The existence of clear Moho reflections, S wave anisotropy, the absence of P wave anisotropy, and low stretching factors do not support the presence of serpentinized mantle. The same conclusion was drawn by Gernigon et al. (2004), primarily based on arguments related to the stretching factors, fault decoupling within the sedimentary formation, and interrogation about the fluid budget. Nirrengarten et al. (2014) and Theissen-Krah et al. (2017) also argued against a serpentinization origin of the LCB in the Møre Margin from 2-D potential field modeling results. Gernigon et al. (2015) suggested that too much continental crust is likely preserved on top of the distal LCB to fit a broad zone of exhumed serpentinized mantle in the outer part of the Møre Margin. A comparison with the distal parts of magma-poor margin (Iberian type) was not easily supported by the data and do not necessarily fit with the long period of rifting and noncontinuum of deformation observed in the mid-Norwegian margin (Gernigon et al., 2015). The correlation between the landward sill extent and the LCB extent is an evidence for a magmatic component in the deeper part of the margin. A serpentinized mantle could not generate sufficient melt to account for the highly intruded sedimentary basin.

Despite limitations, discussion and numerous arguments mentioned above, Peron-Pinvidic and Osmundsen (2016) suggested the presence of a broad zone of exhumed serpentinized mantle directly underneath the lava flows and SDR in the Vøring and Møre marginal highs and presented a magma-poor and ultra-slow spreading scenario to explain the development of the volcanic margin. However, velocities compiled in our study support the presence of thick crystalline basement crust preserved on top of the LCB underneath the Landward Flows and adjacent SDR seismic facies units. The petrology and geochemistry results of the lava flows from ODP Hole 642E also show evidence of continental contamination (Abdelmalak, Meyer, et al., 2016; Meyer et al., 2009) that cannot easily be explained by the presence of a shallow serpentinized mantle lying just underneath the subaerial lava flows. In a similar tectonic setting (Uruguay volcanic margin),

Figure 12. Velocity structure of the continent–ocean transition in the Vøring Margin plotted above the seismic reflection line HV-3-96. The seismic velocity structure is determined using ocean bottom seismometer (OBS) profiles crossing the line (profiles 3-96 and 4-96 (e.g., Mjelde, Shimamura, et al., 2003)) and tied to the Oceanic Drilling Program (ODP) Hole 642E. The total field magnetic and the free air gravity data are plotted above the seismic profiles. Velocity depth profiles for the continental (P1), the continent–ocean transition (P2), and the oceanic (P3) domains are plotted. For each profile the crustal structure is indicated (see section 5.3 for more details and Figure 3A for profile locations).

Clerc et al. (2015) showed evidence that the continental lower crust is most likely preserved underneath the SDR and adjacent basins and also confirm that volcanic margin and magma-poor margins show striking differences (e.g., Clerc et al., 2017; Franke, 2013; Geoffroy et al., 2015).

5.4. The Nature of LCB in the Continent–Ocean Transition and Oceanic Domains

A seismic transect across the continent–ocean transition in the Vøring Margin is shown in Figure 12. On the seismic reflection profile, the top basalt reflection is easily identifiable due to the high-impedance contrast between the postbreakup sediments and the volcanics. The top of the basaltic sequences is an unconformity and correlative conformity between the postbreakup sediments and the underlying basaltic rocks and represents the continentward continuation of the top oceanic basement reflection further west (Abdelmalak, Planke, et al., 2016; Berndt, Planke, et al., 2001; Planke et al., 2000). The SDR wedge is characterized by divergent arcuate reflections with increasing dip in the deeper part.

The P wave velocity model shows a narrow continent–ocean transition, followed by thick igneous crust that tapers off toward normal oceanic crust. On the continental side, very close to the SDR wedge, the top of the crystalline basement is a surface obtained by interpolation of the depth at which OBS-derived P wave velocities change from values <5.5 km/s to values >6 km/s, typical of continental crystalline rocks (e.g., Mjelde, Raum, Myhren, et al., 2005; Raum et al., 2002). The crystalline basement is overlain by an intruded sedimentary layer with P wave velocities between 4.2 and ~ 5.5 km/s. Below the crystalline basement near the base of the preserved continental crust, a LCB is identified ($V_p > 7.0$ km/s).

On the oceanic side, postbreakup sediments lie directly above the oceanic basement. From the velocity model, oceanic layer 2 is characterized by P wave velocities of ~ 4 to 6 km/s and is interpreted as pillow lavas (layer 2A, V_p 4–5 km/s) underlain by a sheeted dyke layer (layer 2B, V_p : 5–6 km/s; e.g., Christeson et al., 2007). Oceanic layer 3A is characterized by P wave velocities ranging between 6 and 6.7 km/s to 6.9 km/s (e.g., R. S. White et al., 1992). Studies of the oceanic crust demonstrate that the lithology of this middle oceanic layer can vary, depending on the spreading rate when it was formed and on how much deformation affected this layer after that (Dijkstra & Cawood, 2004; Mitchell, 2001). Generally, this layer represents a mixture of mafic intrusive rocks and sheeted dykes in unknown proportion (e.g., W. M. White & Klein, 2014). The lowest oceanic layer, 3B, is characterized by P wave velocities of 6.9–7.6 km/s (e.g., R. S. White et al., 1992). Because of these high P wave velocities and a high V_p/V_s ratio, Raum et al. (2006) interpreted this layer as gabbroic intrusive rocks. At depth, the model is limited by the crust–mantle boundary (Moho), defined when the P wave velocities increase from values of 7.6 km/s to values of 7.9–8 km/s in the oceanic domain and 8.0–8.3 km/s in the continental domain (e.g., Mjelde, Raum, Myhren, et al., 2005).

Igneous crustal thickness of 10 km has been inferred for the Møre Margin (Breivik et al., 2006) and Lofoten Margin (Kodaira et al., 1995). Thicker igneous crust has been proposed in the Vøring Marginal High by Breivik et al. (2009). A reduction in igneous crustal and LCB thicknesses in the oceanic domain and the related melt production shortly after breakup may reflect a changing temperature structure of the residual mantle (Eldholm & Grue, 1994). This is attributed to heat advecting from the system (e.g., R. S. White, 1988) and a change to steady-state sea floor spreading.

A ~ 40 to 50 km wide COT zone separates the crystalline basement and the oceanic layers 3A and 3B (Figure 12). The transition from continental to oceanic crust is characterized by lateral velocity changes at midcrustal levels near the inner edge of the SDR (e.g., Breivik et al., 2014; Mjelde, Raum, Myhren, et al., 2005). Below the SDR sequences, vertical and inclined reflections are identified and interpreted as the feeder dyke system (Abdelmalak et al., 2015). It has been suggested that this transitional crust resembles oceanic layer 3A (Eldholm & Grue, 1994). It may consist of dykes near the transition with the extrusive volcanics and could represent a gabbroic complex in the lower regions (Zehnder et al., 1990). Layer 3B could be assigned as a lateral continuation of the LCB since these two layers are characterized by similar P wave velocities (e.g., Raum et al., 2006).

In the Vøring Margin at the COT, highly intruded lower crust below the SDR wedge was observed in several profiles (Figure 13) and has also been identified in the Faroe Margin in the COT zone below the SDRs (R. S. White et al., 2008; R. S. White & Smith, 2009). Variations in seismic velocity in the LCB may reflect variable percentages of intruded igneous rock into preexisting continental crust. Such highly intruded lower crust has been proposed for accretion models of igneous crust in the context of fast spreading ridges (e.g., Korenaga et al., 2002). Nevertheless, calculated half-spreading rates in the Møre and Vøring Margins during the breakup magmatism time indicate a moderate to slow seafloor spreading process (Breivik et al., 2006, 2009; Gernigon et al., 2015).

5.5. The Relationship Between the T -Reflection and the LCB

In the absence of deep drilling, calibration of the lithological composition of rocks at depth very often comes from potential field and seismic refraction studies. Magnetic susceptibility, density, and velocity values (and gradients) are regularly used as conclusive of sedimentary, crystalline, magmatic, or mantle rocks. However, it is also well known that different lithologies, with distinct structural, thermal, and geochemical histories, can correspond to similar sets of geophysical parameters (e.g., Mjelde, Faleide, et al., 2009; Saltus & Blakely, 2011).

In the Vøring Margin, the strong correlation between the high-amplitude T -Reflection and the gravity signal (Figure 6C) indicates that the T -Reflection represents a high-impedance boundary, for example, between

Figure 13. (A) Seismic example along the Vøring Margin showing a highly intruded lower crust below the SDR wedge. The interpretation of the profile is indicated. The 50 km high-pass filtered gravity (B50) and magnetic (M50) data are plotted above the seismic profiles. See Figure 3A for the profile location. (B) Close-up of the highly intruded lower crust below the SDR.

sedimentary rocks and high-density basement rocks such as mid to lower crust or serpentinized mantle (Ren et al., 1998), crustal and underplated gabbroic material (Mjelde et al., 1997, 2002), and/or between upper crustal rocks and high-grade metamorphic rocks such as granulites and eclogites material (Gernigon et al., 2004).

The absence of a magnetic anomaly associated with the dome shape of the *T*-Reflections does not favor a mafic or ultramafic origin (Gernigon et al., 2003, 2004, 2006). From the overall characteristics mentioned above, Gernigon et al. (2006) concluded that the continental part of the LCB observed beneath the outer Vøring Basin may be partly, or fully, attributed to high-pressure granulite/eclogite inherited lower crustal material that is known to display both high *P* wave velocities (7.2–8.5 km/s) and high densities (2.8–3.6 kg/km³). Raum et al. (2006) and Mjelde, Faleide, et al. (2009) interpreted the lower crust in the southwestern corner of the Vøring Basin as in situ eclogites based on anomalously high *P* wave velocities (8.4 km/s). Such Caledonian eclogite bodies are well documented from western Norway onshore (e.g., Andersen & Jamtveit, 1990), have been inferred from wide-angle seismic data in the lower crust of the mid-Norwegian margin (Kvarven et al., 2014, 2016; Mjelde et al., 2013, 2016; Olafsson et al., 1992), and are also documented offshore East Greenland (Schiffer et al., 2015). High-grade metamorphic rocks are interpreted to have been formed by metamorphism of lower crustal gabbros during the Caledonian orogeny (e.g., Corfu et al., 2014; Kvarven et al., 2014).

All these observations provide a consistent model for preferentially interpreting the *T*-Reflection as the top of an inherited old crystalline, high *V_p* basement crust, likely to have been exhumed and preserved in the deep part of the rifted margin. Recent modeling suggests that the deep root of the Caledonides can be exhumed

Figure 14. Schematic illustration showing the tectonic evolution of the Vøring Margin before (redrawn after Gartrell, 1997) and during the breakup magmatic stage (see section 5.5 for more details). The crustal structure is characterized by the occurrence of poorly thinned crustal boudins separated by thinned and sheared inter-boudins covered by thick syn-tectonic sediments.

and lie directly underneath the sedimentary basin formed by subsequent rifting of the crust between Norway and Greenland (Petersen & Schiffer, 2016). The dome shape features represented by T-Reflection could be the result of an impedance contrast between granulite/eclogitic material, and crystalline rocks and/or metasediments that can be differently compacted and faulted on both sides of a preexisting dome during the deformation of the outer Vøring Basin.

We favor a tectonic scenario (Figure 14) where a large part of the crust observed underneath the pre- and syn-rift sediments could represent both preserved upper continental crustal blocks and middle to lower crustal lenses of inherited and intruded, high-grade metamorphic rocks (e.g., Gernigon et al., 2004). The softening of the crust favored by the high heat flow caused by the igneous rocks may lead to a ductile shearing of the lower crust during extension. As a result, boudinage appears as a recurrent deformation process accounting for the thinning of the continental crust at variable scale (e.g., Clerc et al., 2017; Gartrell, 1997). Consequently, the crustal structure is characterized by the occurrence of rigid crustal blocks, interpreted as thick and poorly thinned crustal blocks, separated by inter-boudins characterized by thinned and sheared crust covered by thick syn-tectonic sediments (Figure 14).

It is well known that the mid-Norwegian margin experienced a prolonged history of extension and basin formation since the collapse of the Caledonian orogeny (e.g., Faleide et al., 2008). The Late Jurassic–Early Cretaceous extension and the Late Cretaceous/Early Paleocene extension are the most important events that led to major fault activity and deep detachments, which in turn control the crustal deformation and subsidence of the margin. Extension formed large-scale rotated rift blocks or structural highs trending NE–SW parallel to weakness zones, probably inherited from the Caledonian crustal configuration. The deep detachments that flattened out at the lower crustal level influenced the development of the sedimentary basin since the drastic thinning phase in Late Jurassic–Early Cretaceous.

Continental breakup between Norway and Greenland occurred in association with a large-scale magmatic activity and volcanism (Eldholm et al., 1989; Skogseid & Eldholm, 1989). Magma intrusion into the lower crust enhanced the metamorphic processes that led to an increasing velocity of the lower crust. The final result was a heterogeneous body made of a mixture of pre- and syn-breakup, and possibly post-breakup, mafic and

Figure 15. (A) LCB thickness map for the NE Atlantic conjugate margins restored to C21 (~47 Ma) in a fixed Eurasia reference frame using the finite rotations from Gaina et al. (2009). The refraction lines used to compile this map are shown in Figure 1. Examples of deep strong reflections in (B) the NE Greenland margin, (C) the Jan Mayen Ridge, and (D) the SW Barents Sea. Black arrows indicate the deep strong reflection similar to the *T*-Reflection in Norway. The profile location is indicated in Figure 15A. B100 and B200 are the high-pass filtered gravity data with a cutoff of 100 and 200 km, respectively. Seismic interpretation comes from our unpublished data.

ultramafic rocks (cumulates and sills) associated with old metamorphic rocks such as granulites and eclogites. Additionally, the lithological boundary probably also served as favored conduits for igneous intrusions, resulting in the sill-like appearance for parts of the reflections and which may explain the varying reflectivity pattern and the abrupt endings of many of the reflections. Magma may also have intruded the low-angle faults and detachments and top basement contact expected at the level of the *T*-Reflection. The magmatic intrusions may increase the reflectivity of the contacts that appear as very bright reflections on some profiles. This could also be responsible for the local discontinuity of the *T*-Reflection. An increasing melting degree toward the breakup axis is responsible for increasing rate of cumulates and sill intrusions into the lower crust at the COT.

5.6. Lateral Variation Along the Conjugate Margins

To complete the whole picture of the deep structures of the mid-Norwegian margin, we also investigated the NE Greenland conjugate margin (Figure 15). High-quality seismic reflection lines from the NE Greenland margin are sparse but show high-amplitude deep crustal reflections at depths ranging between 7 and 9 s (Figure 15B). These reflections, similar to the *T*-Reflection discussed here, were also mapped beneath the Danmarkshavn Ridge and in the eastern part of the Thetis Basin (Figure 15A). They are discontinuous, show a rough character, and fit with the high-pass filtered Bouguer gravity anomaly. Similar to the *T*-Reflection, the conjugate reflections also represent a high-impedance boundary associated with a high-density body. A strong deep crustal high-amplitude reflection is also found in the Jan Mayen Ridge at depths ranging between 7 and 8 s (Figure 15C). A double midcrustal dome-shaped feature with high-amplitude reflections (named D1 and D2) is found at depths ranging between 7 and 10 s in the Harstad Basin situated in the SW Barents Sea (Figure 15D). The high-pass filtered Bouguer anomaly correlates very well with the doming of

these two deep crustal reflections, which also have no obvious magnetic signature. The main density contrast is likely across D2, whereas D1 is possibly an intra-Jurassic reflection (Figure 15D).

Compared to the mid-Norwegian Margin, the density of refraction lines along the NE Greenland and the Jan Mayen Micro-Continent is lower, and therefore, the shape and the variation of the LCB are not as well constrained. However, seismic refraction data reveal the existence of LCB along the NE Greenland margins (Hermann, 2013; Mandler & Jokat, 1998; Schlindwein & Jokat, 2000; Schmidt-Aursch & Jokat, 2005; Voss & Jokat, 2007; Voss et al., 2009; Weigel et al., 1995). From the available refraction data (Figure 3A), we also compiled the LCB thickness map along the NE Greenland margin (Figure 15A) and find it thicker than at the mid-Norwegian margin. The thicker part (12–15 km) of the LCB is situated below the Foster Volcanic Province, and it becomes thinner below the Jameson Land where sill intrusions but no volcanic extrusives are found. Along the Thetis Margin, the thickness of the LCB does not exceed 4 km. Similarly to the mid-Norwegian margin, the LCB thickness decreases at magnetic chrons C23–C22 and merges into oceanic layer 3. A high-velocity lower crust, with V_p of more than 7 km/s, is also identified in the eastern margin of the Jan Mayen Micro-Continent (Breivik et al., 2012).

Offshore NE Greenland, two igneous sill complexes are identified in the Thetis Basin and the Danmarkshavn Basin (e.g., Reynolds et al., 2017). The sill complex of the Danmarkshavn Basin is continuous landward until the Jameson Land area (Figure 15A). If we consider a link between the igneous sill complexes and the extent of the LCB, we expect that the latter should be present below the Thetis and the Danmarkshavn basins. Such an assumption could only be verified by new refraction data. We estimated the total extent of the LCB and its volume for the Vøring–Lofoten and NE Greenland conjugate margins. We considered a fully magmatic LCB, which is therefore an overestimation (see discussion above) and thus represents a maximum of melt production. For the NE Greenland area, the LCB covers an area of $\sim 136,000 \text{ km}^2$ and indicates a maximum volume of $0.56 \times 10^6 \text{ km}^3$. On the conjugate Vøring–Lofoten margins, the LCB covers a similar area of $\sim 140,000 \text{ km}^2$ and indicates a maximum volume of $0.50 \times 10^6 \text{ km}^3$, which is arguably surprising if we consider the asymmetry of the NE Atlantic margins.

6. Conclusions

In this paper we investigated the deep structure of the Vøring Margin offshore mid-Norway using an integrated multidisciplinary geophysical data sets, including reflection and refraction seismic, gravity, and magnetic data. This detailed analysis has resulted in the following conclusions:

1. New seismic data permit a refined mapping of the *T*-Reflection in the outer Vøring Margin. The *T*-Reflection is identified between 7 and 10 s and sometimes consists of one single smooth reflection, such as in the North and South Gjallar ridges. However, it is frequently associated with a set of rough multiple reflections and displays discontinuous steep and flat segments with varying geometries, amplitudes, and contact relationships. The *T*-Reflection seems to be connected to deep sill networks and locally located at the continuation of basement high structures or terminates over fractures and faults.
2. The positive correlation between the geometries of the 50 km high-pass filtered Bouguer anomaly and the *T*-Reflection indicates that the latter represents a high-impedance boundary associated with a high-density body. However, no significant magnetic anomaly is associated with the *T*-Reflection.
3. A comparison between the *T*-Reflection and the top LCB depths, deduced from independent seismic reflection and refraction results, indicates that these two features are found at the same depth (within the uncertainties) in $\sim 50\%$ of the outer Vøring Margin areas.
4. We present a tectonic scenario where a large part of the deep crustal structure is composed of preserved upper continental basement and middle to lower crustal lenses of inherited and intruded high-grade metamorphic rocks. Intrusions into faulted crustal blocks are responsible for the rough character of the *T*-Reflection whereas sill intrusion into the preexisting lower crust and detachment faults could explain the smooth character of the *T*-Reflection.
5. Magma intrusion into the lower crust might be responsible for metamorphic processes that may have led to an increased velocity of the lower crust. The result is a heterogeneous LCB made of a mixture of pre- and syn-breakup mafic and ultramafic rocks (sills and cumulates) associated with older metamorphic rocks such as granulites and eclogites. The positive correlation between the extent, continuity, and lateral thickness variation of the LCB and sill intrusion in the basin is an argument in favor of a breakup magmatic

origin of the LCB. An increased degree of melting toward the breakup axis is likely responsible for an oceanward increase in volumes of cumulates and sill intrusions into the lower crust.

6. Similar domal shapes and high-amplitude reflections associated with high gravity and low magnetic anomalies have been identified on other NE Atlantic margins. However, further investigations are probably required to understand their nature and their possible correlation with the mid-Norwegian margin T-Reflection.

Acknowledgments

Funding for this work came from the OMNIS Project: Offshore Mid-Norway: Integrated Margin and Basin Studies (project 210429) funded by the Research Council of Norway through its Centres of Excellence funding scheme, project 223272 CEED (Centre for Earth Evolution and Dynamics). The seismic, magnetic, and gravity data presented in this study were provided by TGS. We would like to acknowledge First Geo AS and TGS for the use of Hiqbe™ regional velocity cube employed in depth conversion of the seismic profiles. Seismic interpretation was done using IHS Kingdom software. Grid interpolations and map compilations were established using Geosoft Oasis Montaj and ArcGis softwares. GES acknowledges VISTA—a basic research program in collaboration between The Norwegian Academy of Science and Letters, and Statoil (project 6268, “DEFMOD”). We thank Camille Clerc, Christian Schiffer, the associated Editor, and the Editor Claudio Faccenna for useful comments and suggestions that improved the paper.

References

- Abdelmalak, M. M. (2010). Transition Spatio-temporelle entre rift sédimentaire et marge passive volcanique: l'exemple de la Baie de Baffin, Centre Ouest Groenland, (Spatio-temporal transition between a sedimentary basin to a volcanic passive margin: The Baffin Bay case example, central west Greenland) (pp. 266). [Available at Université du Maine (France) Retrieved from <http://cyberdoc.univ-lemans.fr/theses/2010/2010LEMA1030.pdf>, Le Mans].
- Abdelmalak, M. M., Andersen, T. B., Planke, S., Faleide, J. I., Corfu, F., Tegner, C., ... Myklebust, R. (2015). The ocean–continent transition in the mid-Norwegian margin: Insight from seismic data and an onshore Caledonian field analogue. *Geology*, 43, 1011–1014. <https://doi.org/10.1130/G37086.1>
- Abdelmalak, M. M., Faleide, J. I., Planke, S., Theissen-Krah, S., Zastrozhnov, D., Breivik, A. J., ... Myklebust, R. (2014). Breakup magmatism style on the North Atlantic Igneous Province: Insight from Mid-Norwegian volcanic margin. Paper presented at EGU General Assembly Conference Abstracts, Vienna.
- Abdelmalak, M. M., Geoffroy, L., Angelier, J., Bonin, B., Callot, J. P., Gélard, J. P., & Aubourg, C. (2012). Stress fields acting during lithosphere breakup above a melting mantle: A case example in West Greenland. *Tectonophysics*, 581, 132–143. <https://doi.org/10.1016/j.tecto.2011.11.020>
- Abdelmalak, M. M., Meyer, R., Planke, S., Faleide, J. I., Gernigon, L., Frieling, J., ... Myklebust, R. (2016). Pre-breakup magmatism on the Vøring Margin: Insight from new sub-basalt imaging and results from Ocean Drilling Program Hole 642E. *Tectonophysics*, 675, 258–274. <https://doi.org/10.1016/j.tecto.2016.02.037>
- Abdelmalak, M. M., Planke, S., Faleide, J. I., Jerram, D. A., Zastrozhnov, D., Eide, S., & Myklebust, R. (2016). The development of volcanic sequences at rifted margins: New insights from the structure and morphology of the Vøring Escarpment, mid-Norwegian margin. *Journal of Geophysical Research: Solid Earth*, 121, 5212–5236. <https://doi.org/10.1002/2015JB012788>
- Andersen, T. B., & Jamtveit, B. (1990). Uplift of deep crust during orogenic extensional collapse: A model based on field studies in the Sogn-Sunnfjord Region of western Norway. *Tectonics*, 9(5), 1097–1111. <https://doi.org/10.1029/TC0091005p01097>
- Berndt, C. (2002). Residual Bouguer satellite gravity anomalies reveal basement grain and structural elements of the Vøring Margin, off Norway. *Norwegian Journal of Geology*, 82, 31–36.
- Berndt, C., Mjelde, R., Planke, S., Shimamura, H., & Faleide, J. I. (2001). Controls on the tectono-magmatic evolution of a volcanic transform margin: The Vøring Transform Margin, NE-Atlantic. *Marine Geophysical Researches*, 22(3), 133–152.
- Berndt, C., Planke, S., Alvestad, E., Tsikalas, F., & Rasmussen, T. (2001). Seismic volcanostratigraphy of the Norwegian margin: Constraints on tectonomagmatic break-up processes. *Journal of the Geological Society*, 158(3), 413–426. <https://doi.org/10.1144/jgs.158.3.413>
- Berndt, C., Skogly, O. P., Planke, S., Eldholm, O., & Mjelde, R. (2000). High-velocity breakup-related sills in the Vøring Basin, off Norway. *Journal of Geophysical Research*, 105, 28,443–428,454.
- Blystad, P., Brekke, H., Færseth, R. B., Larsen, B. T., Skogseid, J., & Tørudbakken, B. (1995). Structural elements of the Norwegian continental shelf Part II: The Norwegian sea region, *NPD-Bulletin, The Norwegian Petroleum Directorate*, 8.
- Breivik, A., Faleide, J. I., Mjelde, R., Flueh, E., & Murai, Y. (2014). Magmatic development of the outer Vøring Margin from seismic data. *Journal of Geophysical Research: Solid Earth*, 119, 6733–6755. <https://doi.org/10.1002/2014JB011040>
- Breivik, A. J., Faleide, J. I., Mjelde, R., & Flueh, E. R. (2009). Magma productivity and early seafloor spreading rate correlation on the northern Vøring Margin, Norway—Constraints on mantle melting. *Tectonophysics*, 468(1–4), 206–223. <https://doi.org/10.1016/j.tecto.2008.09.020>
- Breivik, A. J., Mjelde, R., Faleide, J. I., & Murai, Y. (2006). Rates of continental breakup magmatism and seafloor spreading in the Norway Basin–Iceland plume interaction. *Journal of Geophysical Research*, 111, B07102. <https://doi.org/10.1029/2005JB004004>
- Breivik, A. J., Mjelde, R., Faleide, J. I., & Murai, Y. (2012). The eastern JanMayen microcontinent volcanic margin. *Geophysical Journal International*, 188(3), 798–818.
- Breivik, A. J., Mjelde, R., Raum, T., Faleide, J. I., Murai, Y., & Flueh, E. R. (2011). Crustal structure beneath the Trøndelag Platform and adjacent areas of the mid-Norwegian margin, as derived from wide-angle seismic and potential field data. *Norwegian Journal of Geology*, 90, 141–161.
- Brekke, H. (2000). The tectonic evolution of the Norwegian Sea Continental Margin with emphasis on the Vøring and Møre Basins. *Geological Society, London, Special Publications*, 167(1), 327–378. <https://doi.org/10.1144/gsl.sp.2000.167.01.13>
- Brooks, C. K. (2011). The East Greenland rifted volcanic margin. *Geological Survey of Denmark and Greenland Bulletin*, 24, 96.
- Brune, S., Heine, C., Clift, P. D., & Pérez-Gussinyé, M. (2017). Rifted margin architecture and crustal rheology: Reviewing Iberia–Newfoundland, Central South Atlantic, and South China Sea. *Marine and Petroleum Geology*, 79, 257–281. <https://doi.org/10.1016/j.marpetgeo.2016.10.018>
- Cartwright, J., & Hansen, D. M. (2006). Magma transport through the crust via interconnected sill complexes. *Geology*, 34(11), 929–932. <https://doi.org/10.1130/g22758a.1>
- Christeson, G. L., McIntosh, K. D., & Karson, J. A. (2007). Inconsistent correlation of seismic layer 2a and lava layer thickness in oceanic crust. *Nature*, 445(7126), 418–421. http://www.nature.com/nature/journal/v445/n7126/supinfo/nature05517_S1.html
- Clerc, C., Jolivet, L., & Ringenbach, J. C. (2015). Ductile extensional shear zones in the lower crust of a passive margin. *Earth and Planetary Science Letters*, 431, 1–7. <https://doi.org/10.1016/j.epsl.2015.08.038>
- Clerc, C., Ringenbach, J.-C., Jolivet, L., & Ballard, J.-F. (2017). Rifted margins: Ductile deformation, boudinage, continentward-dipping normal faults and the role of the weak lower crust. *Gondwana Research*. <https://doi.org/10.1016/j.gr.2017.04.030>
- Coffin, M. F., & Eldholm, O. (1994). Large igneous provinces: Crustal structure, dimensions and external consequence. *Reviews of Geophysics*, 32, 1–36.
- Corfu, F., Andersen, T. B., & Gasser, D. (2014). The Scandinavian Caledonides: Main features, conceptual advances and critical questions. *Geological Society, London, Special Publications*, 390(1), 9–43. <https://doi.org/10.1144/sp390.25>

- Corseri, R., Senger, K., Selway, K., Abdelmalak, M. M., Planke, S., & Jerram, D. (2017). Magnetotelluric evidence for massive sulphide mineralization in intruded sediments of the outer Vøring Basin, mid-Norway. *Tectonophysics*, 706–707, 196–205. <https://doi.org/10.1016/j.tecto.2017.04.011>
- Dijkstra, A. H., & Cawood, P. A. (2004). Base-up growth of ocean crust by multiple phases of magmatism: Field evidence from Macquarie Island. *Journal of the Geological Society*, 161(5), 739–742. <https://doi.org/10.1144/0016-764904-033>
- Doré, A. G., Lundin, E. R., Jensen, L. N., Birkland, Ø., Eliassen, P. E., & Fichler, C. (1999). Principal tectonic events in the evolution of the northwest European Atlantic margin. *Geological Society, London, Petroleum Geology Conference Series*, 5, 41–61. <https://doi.org/10.1144/0050041>
- Ebbing, J., Lundin, E., Olesen, O., & Hansen, E. K. (2006). The mid-Norwegian margin: A discussion of crustal lineaments, mafic intrusions, and remnants of the Caledonian root by 3D density modelling and structural interpretation. *Journal of the Geological Society, London*, 163, 47–59.
- Eldholm, O., Gladchenko, T. P., Skogseid, J., & Planke, S. (2000). Atlantic volcanic margins: A comparative study. In A. E. A. Nottvedt (Ed.), *Dynamics of the Norwegian margin, Special Publications* (pp. 411–428). London: Geological Society.
- Eldholm, O., & Grue, K. (1994). North Atlantic volcanic margins: Dimensions and production rates. *Journal of Geophysical Research*, 99, 2955–2968.
- Eldholm, O., Thiede, J., & Taylor, E. (1989). Evolution of the Vøring Volcanic Margin. In O. Eldholm, et al. (Eds.), *Proceedings of the ODP, Scientific Results 104* (pp. 1033–1065). College Station, TX: Ocean Drilling Program.
- England, P. C. (1983). Constraints on extension of continental lithosphere. *Journal of Geophysical Research*, 88, 1145–1152.
- Ernst, R. E., Buchan, K. L., & Palmer, H. C. (1995). Giant dyke swarms: Characteristics, distribution and geotectonic applications. In G. Baer & A. Heimann (Eds.), *Physics and chemistry of dykes* (pp. 3–21). Rotterdam: Balkema.
- Escartin, J., Hirth, G., & Evans, B. (2001). Strength of slightly serpentinized peridotites: Implications for the tectonics of oceanic lithosphere. *Geology*, 29(11), 1023–1026. [https://doi.org/10.04.106/0091-7613\(2001\)029%3C1023:sossp%3E2.0.co;2](https://doi.org/10.04.106/0091-7613(2001)029%3C1023:sossp%3E2.0.co;2)
- Faleide, J. I., Tsikalas, F., Breivik, A. J., Mjelde, R., Ritzmann, O., Engen, Ø., ... Eldholm, O. (2008). Structure and evolution of the continental margin off Norway and the Barents Sea. *Episodes*, 31(1), 82–91.
- Fichler, C., Odinsen, T., Rueslåtten, H., Olesen, O., Vindstad, J. E., & Wienecke, S. (2011). Crustal inhomogeneities in the northern North Sea from potential field modeling: Inherited structure and serpentinites? *Tectonophysics*, 510(1), 172–185. <https://doi.org/10.1016/j.tecto.2011.06.026>
- Franke, D. (2013). Rifting, lithosphere breakup and volcanism: Comparison of magma-poor and volcanic rifted margins. *Marine and Petroleum Geology*, 43, 63–87. <https://doi.org/10.1016/j.marpetgeo.2012.11.003>
- Gaina, C., Gernigon, L., & Ball, P. (2009). Palaeocene–Recent plate boundaries in NE Atlantic and formation of the Jan Mayen microcontinent. *Journal of the Geological Society of London*, 166, 601–616.
- Gartrell, A. P. (1997). Evolution of rift basins and low-angle detachments in multilayer analog models. *Geology*, 25(7), 615–618.
- Geissler, W. H., Gaina, C., Hopper, J. R., Funck, T., Blischke, A., Arting, U., ... Abdelmalak, M. M. (2016). Seismic volcanostratigraphy of the NE Greenland continental margin. *Geological Society, London, Special Publications*, 447. <https://doi.org/10.1144/sp447.11>
- Geoffroy, L. (2005). Volcanic passive margins. *Comptes Rendus Géosciences*, 337, 1395–1408.
- Geoffroy, L., Burov, E. B., & Werner, P. (2015). Volcanic passive margins: Another way to break up continents. *Scientific Reports*, 5, 14828. <https://doi.org/10.1038/srep14828>
- Gernigon, L., Blischke, A., Nasuti, A., & Sand, M. (2015). Conjugate volcanic rifted margins, sea-floor spreading and microcontinent: Insights from new high-resolution aeromagnetic surveys in the Norway Basin. *Tectonics*, 34, 907–933. <https://doi.org/10.1002/2014TC003717>
- Gernigon, L., Lucazeau, F., Brigaud, F., Ringenbach, J.-C., Planke, S., & Le Gall, B. (2006). A moderate melting model for the Vøring Margin (Norway) based on structural observations and a thermo-kinematical modelling: Implication for the meaning of the lower crustal bodies. *Tectonophysics*, 412(3–4), 255–278.
- Gernigon, L., Olesen, O., Ebbing, J., Wienecke, S., Gaina, C., Mogaard, J. O., ... Myklebust, R. (2009). Geophysical insights and early spreading history in the vicinity of the Jan Mayen Fracture Zone, Norwegian–Greenland Sea. *Tectonophysics*, 468(1–4), 185–205.
- Gernigon, L., Ringenbach, J. C., Planke, S., Jonquet-Kolstø, E., Ballard, J. F., & Le Gall, B. (2001). Rifting and segmentation along the outer Vøring Basin, North Atlantic margin, Norway. In *American Association of Petroleum Geologists (AAPG) annual meeting* (pp. 1–7). Denver, CO.
- Gernigon, L., Ringenbach, J.-C., Planke, S., & Jonquet-Kolsto, H. (2003). Extension, crustal structure and magmatism at the outer Vøring Basin, Norwegian margin. *Journal of the Geological Society of London*, 160, 197–208.
- Gernigon, L., Ringenbach, J.-C., Planke, S., & Le Gall, B. (2004). Deep structures and breakup along volcanic rifted margins: Insights from integrated studies along the outer Vøring Basin (Norway). *Marine and Petroleum Geology*, 21(3), 363–372.
- Gradstein, F. M., Ogg, J. G., Schmitz, M. D., & Ogg, G. M. (2012). The Geologic time scale 2012.
- Hansen, D. M., Cartwright, J. A., & Thomas, D. (2004). 3D seismic analysis of the geometry of igneous sills and sill junction relationships. *Geological Society, London, Memoirs*, 29(1), 199–208. <https://doi.org/10.1144/gsl.mem.2004.029.01.19>
- Hermann, T. (2013). The northeast Greenland Margin tectonic evolution (pp. 170). vorgelegt dem Rat der Chemisch-Geowissenschaftlichen Fakultät der Friedrich-Schiller-Universität Jena.
- Hinz, K. (1981). Hypothesis on terrestrial catastrophes: Wedges of very thick oceanward dipping layers beneath passive margins—Their origin and palaeoenvironment significance. *Geologisches Jahrbuch*, 22, 345–363.
- Holbrook, W. S., Larsen, H. C., Korenaga, J., Dahl-Jensen, T., Reid, I. D., Kelemen, P. B., ... Detrick, R. S. (2001). Mantle thermal structure and active upwelling during continental breakup in the North Atlantic. *Earth and Planetary Science Letters*, 190, 251–266.
- Holbrook, W. S., Mooney, W. D., & Christensen, N. J. (1992). The seismic velocity structure of the deep continental crust. In R. A. D. M. Fountain & R. W. Kay (Eds.), *Continental lower crust, development in geotectonics* (pp. 1–43). Amsterdam: Elsevier.
- Huisman, S. R., & Beaumont, C. (2003). Symmetric and asymmetric lithospheric extension: Relative effects of frictional-plastic and viscous strain softening. *Journal of Geophysical Research*, 108(B10), 2496. <https://doi.org/10.1029/2002JB002026>
- Hurich, C. A., Deemer, S. J., Indares, A., & Salisbury, M. (2001). Compositional and metamorphic controls on velocity and reflectivity in the continental crust: An example from the Grenville Province of eastern Québec. *Journal of Geophysical Research*, 106(B1), 665–682. <https://doi.org/10.1029/2000JB900244>
- Kinck, J. J., Husebye, E. S., & Larsson, F. R. (1993). The Moho depth distribution in Fennoscandia and the regional tectonic evolution from Archean to Permian times. *Precambrian Research*, 64(1–4), 23–51. [https://doi.org/10.1016/0301-9268\(93\)90067-C](https://doi.org/10.1016/0301-9268(93)90067-C)
- Klausen, M. B., & Larsen, H. C. (2002). East Greenland coast-parallel dike swarm and its role in continental breakup. In M. A. Menzies, S. L. Klemperer, C. J. Ebinger, & J. Baker (Eds.), *Volcanic rifted margins, Geological Society of America Special Paper* (pp. 133–158). Boulder, CO: Geological Society of America.

- Klingelhöfer, F., Edwards, R. A., Hobbs, R. W., & England, R. W. (2005). Crustal structure of the NE Rockall Trough from wide-angle seismic data modeling. *Journal of Geophysical Research*, 110, B11105. <https://doi.org/10.1029/2005JB003763>
- Kodaira, S., Goldschmidt-Rokita, A., Hartmann, J. M., Hirschleber, H. B., Iwasaki, T., Kanazawa, T., ... Shimamura, H. (1995). Crustal structure of the Lofoten continental margin, off northern Norway, from ocean-bottom seismographic studies. *Geophysical Journal International*, 121(3), 907–924. <https://doi.org/10.1111/j.1365-246X.1995.tb06447.x>
- Koptev, A., Calais, E., Burov, E., Leroy, S., & Gerya, T. (2015). Dual continental rift systems generated by plume–lithosphere interaction. *Nature Geoscience*, 8(5), 388–392. <https://doi.org/10.1038/ngeo2401>
- Korenaga, J., Kelemen, P. B., & Holbrook, W. S. (2002). Methods for resolving the origin of large igneous provinces from crustal seismology. *Journal of Geophysical Research*, 107(B9), 2178. <https://doi.org/10.1029/2001JB001030>
- Kusznir, N. J., & Park, J. K. (1987). The extensional strength of the continental lithosphere: Its dependence on geothermal gradient, and crustal composition and thickness. In M. P. Coward, J. F. Dewey, & P. L. Hancock (Eds.), *Continental extensional tectonics, Geological Society Special Publication* (Vol. 28, pp. 35–52). London: The Geological Society.
- Kusznir, N. J., Roberts, A., & Bellingham, P. (2015). Deep seismic reflectivity at volcanic margins: Reflections from the petrological Moho or from within the mantle? Paper presented at EGU General Assembly, Vienna.
- Kvarven, T., Ebbing, J., Mjelde, R., Faleide, J. I., Libak, A., Thybo, H., ... Murai, Y. (2014). Crustal structure across the Møre Margin, mid-Norway, from wide-angle seismic and gravity data. *Tectonophysics*, 626, 21–40. <https://doi.org/10.1016/j.tecto.2014.03.021>
- Kvarven, T., Mjelde, R., Hjelstuen, B. O., Faleide, J. I., Thybo, H., Flueh, E. R., & Murai, Y. (2016). Crustal composition of the Møre Margin and compilation of a conjugate Atlantic margin transect. *Tectonophysics*, 666, 144–157. <https://doi.org/10.1016/j.tecto.2015.11.002>
- Larsen, L. M., Pedersen, A. K., Tegner, C., & Duncan, R. A. (2014). Eocene to Miocene igneous activity in NE Greenland: Northward younging of magmatism along the East Greenland Margin. *Journal of the Geological Society*, 171(4), 539–553. <https://doi.org/10.1144/jgs2013-118>
- Lavecchia, A., Thieulot, C., Beekman, F., Cloetingh, S., & Clark, S. (2017). Lithosphere erosion and continental breakup: Interaction of extension, plume upwelling and melting. *Earth and Planetary Science Letters*, 467, 89–98. <https://doi.org/10.1016/j.epsl.2017.03.028>
- Lavier, L. L., & Manatschal, G. (2006). A mechanism to thin the continental lithosphere at magma-poor margins. *Nature*, 440, 607–610.
- Lien, T. (2005). From rifting to drifting: Effects on the development of deep-water hydrocarbon reservoirs in a passive margin setting, Norwegian sea, *Norsk Geologisk Tidsskrift volume 85 2005 number 4 side 319-332*.
- Lister, J. R., & Kerr, R. C. (1991). Fluid-mechanical models of crack propagation and their application to magma transport in dykes. *Journal of Geophysical Research*, 96, 10,049–10,077.
- Lundin, E. R., & Doré, A. G. (1997). A tectonic model for the Norwegian passive margin with implications for the NE Atlantic: Early Cretaceous to break-up. *Journal of the Geological Society*, 154(3), 545–550. <https://doi.org/10.1144/gsjgs.154.3.0545>
- Lundin, E. R., & Doré, A. G. (2005). NE Atlantic break-up: A re-examination of the Iceland mantle plume model and the Atlantic–Arctic linkage. *Geological Society, London, Petroleum Geology Conference Series*, 6, 739–754. <https://doi.org/10.1144/0060739>
- Lundin, E. R., & Doré, A. G. (2011). Hyperextension, serpentinization, and weakening: A new paradigm for rifted margin compressional deformation. *Geology*, 39(4), 347–350. <https://doi.org/10.1130/g31499.1>
- Mandler, H. A. F., & Jokat, W. (1998). The crustal structure of central East Greenland: Results from combined land–sea seismic refraction experiments. *Geophysical Journal International*, 135(1), 63–76. <https://doi.org/10.1046/j.1365-246X.1998.00586.x>
- Maus, S., Barckhausen, U., Berkenbosch, H., Bournas, N., Brozena, J., Childers, V., ... Caratori Tontini, F. (2009). EMAG2: A 2-arc min resolution Earth Magnetic Anomaly Grid compiled from satellite, airborne, and marine magnetic measurements. *Geochemistry, Geophysics, Geosystems*, 10, Q08005. <https://doi.org/10.1029/2009GC002471>
- Menzies, M. A., Klemperer, S. L., Ebinger, C. J., & Baker, J. (2002). Characteristics of volcanic rifted margins. In M. A. Menzies, et al. (Eds.), *Volcanic rifted margins* (Vol. 362, pp. 1–14). Boulder, CO: Geological Society of America.
- Meyer, R., Hertogen, J., Pedersen, R. B., Viereck-Götte, L., & Abratis, M. (2009). Interaction of mantle derived melts with crust during the emplacement of the Vøring Plateau, NE Atlantic. *Marine Geology*, 261(1–4), 3–16. <https://doi.org/10.1016/j.margeo.2009.02.007>
- Meyer, R., van Wijk, J., & Gernigon, L. (2007). The North Atlantic Igneous Province: A review of models for its formation. *Geological Society of America Special Papers*, 430, 525–552. [https://doi.org/10.1130/2007.2430\(26\)](https://doi.org/10.1130/2007.2430(26))
- Mitchell, N. C. (2001). Random sequences of lithologies exposed on the Mid-Atlantic Ridge. *Journal of Geophysical Research*, 106(B11), 26,365–26,378. <https://doi.org/10.1029/2001JB000241>
- Mjelde, R., Faleide, J. I., Breivik, A. J., & Raum, T. (2009). Lower crustal composition and crustal lineaments on the Vøring Margin, NE Atlantic: A review. *Tectonophysics*, 472(1–4), 183–193. <https://doi.org/10.1016/j.tecto.2008.04.018>
- Mjelde, R., Goncharov, A., & Müller, R. D. (2013). The Moho: Boundary above upper mantle peridotites or lower crustal eclogites? A global review and new interpretations for passive margins. *Tectonophysics*, 609, 636–650. <https://doi.org/10.1016/j.tecto.2012.03.001>
- Mjelde, R., Kasahara, J., Shimamura, H., Kamimura, A., Kanazawa, T., Kodaira, S., ... Shiobara, H. (2002). Lower crustal seismic velocity-anomalies; magmatic underplating or serpentinized peridotite? Evidence from the Vøring Margin, NE Atlantic. *Marine Geophysical Researches*, 23(2), 169–183. <https://doi.org/10.1023/a:1022480304527>
- Mjelde, R., Kodaira, S., Hassan, R. K., Goldschmidt-Rokita, A., Tomita, N., Sellevoll, M. A., ... Kanazawa, T. (1996). The continent/ocean transition of the Lofoten volcanic margin, N. Norway. *Journal of Geodynamics*, 22(3–4), 189–206. [https://doi.org/10.1016/0264-3707\(96\)00016-6](https://doi.org/10.1016/0264-3707(96)00016-6)
- Mjelde, R., Kodaira, S., Shimamura, H., Kanazawa, T., Shiobara, H., Berg, E. W., & Riise, O. (1997). Crustal structure of the central part of the Vøring Basin, mid-Norway margin, from ocean bottom seismographs. *Tectonophysics*, 277(4), 235–257. [https://doi.org/10.1016/S0040-1951\(97\)00028-0](https://doi.org/10.1016/S0040-1951(97)00028-0)
- Mjelde, R., Kvarven, T., Faleide, J. I., & Thybo, H. (2016). Lower crustal high-velocity bodies along North Atlantic passive margins, and their link to Caledonian suture zone eclogites and Early Cenozoic magmatism. *Tectonophysics*, 670, 16–29. <https://doi.org/10.1016/j.tecto.2015.11.021>
- Mjelde, R., Raum, T., Breivik, A., Shimamura, H., Murai, Y., Takanami, T., & Faleide, J. I. (2005). Crustal structure of the Vøring Margin, NE Atlantic: A review of geological implications based on recent OBS data. *Geological Society, London, Petroleum Geology Conference Series*, 6, 803–813. <https://doi.org/10.1144/0060803>
- Mjelde, R., Raum, T., Digranes, P., Shimamura, H., Shiobara, H., & Kodaira, S. (2003). V_p/V_s ratio along the Vøring Margin, NE Atlantic, derived from OBS data: Implications on lithology and stress field. *Tectonophysics*, 369(3–4), 175–197. [https://doi.org/10.1016/S0040-1951\(03\)00198-7](https://doi.org/10.1016/S0040-1951(03)00198-7)
- Mjelde, R., Raum, T., Kandilarov, A., Murai, Y., & Takanami, T. (2009). Crustal structure and evolution of the outer Møre Margin, NE Atlantic. *Tectonophysics*, 468(1–4), 224–243. <https://doi.org/10.1016/j.tecto.2008.06.003>
- Mjelde, R., Raum, T., Murai, Y., & Takanami, T. (2007). Continent–ocean-transitions: Review, and a new tectono-magmatic model of the Vøring Plateau, NE Atlantic. *Journal of Geodynamics*, 43(3), 374–392.
- Mjelde, R., Raum, T., Myhren, B., Shimamura, H., Murai, Y., Takanami, T., ... Næss, U. (2005). Continent–ocean transition on the Vøring Plateau, NE Atlantic, derived from densely sampled ocean bottom seismometer data. *Journal of Geophysical Research*, 110, B05101. <https://doi.org/10.1029/2004JB003026>

- Mjelde, R., Shimamura, H., Kanazawa, T., Kodaira, S., Raum, T., & Shiobara, H. (2003). Crustal lineaments, distribution of lower crustal intrusives and structural evolution of the Vøring Margin, NE Atlantic; new insight from wide-angle seismic models. *Tectonophysics*, 369(3–4), 199–218. [https://doi.org/10.1016/S0040-1951\(03\)00199-9](https://doi.org/10.1016/S0040-1951(03)00199-9)
- Mutter, J. C., Talwani, M., & Stoffa, P. L. (1982). Origin of seaward-dipping reflectors in oceanic crust off the Norwegian margin by “subaerial sea-floor spreading”. *Geology*, 10, 353–357.
- Neumann, E.-R., Svensen, H., Tegner, C., Planke, S., Thirlwall, M., & Jarvis, K. E. (2013). Sill and lava geochemistry of the mid-Norway and NE Greenland conjugate margins. *Geochemistry, Geophysics, Geosystems*, 14, 3666–3690. <https://doi.org/10.1002/ggge.20224>
- Nirrengarten, M., Gernigon, L., & Manatschal, G. (2014). Lower crustal bodies in the Møre volcanic rifted margin: Geophysical determination and geological implications. *Tectonophysics*, 636, 143–157. <https://doi.org/10.1016/j.tecto.2014.08.004>
- Olafsson, I., Sundvor, E., Eldholm, O., & Grue, K. (1992). Møre Margin: Crustal structure from analysis of expanded spread profiles. *Marine Geophysical Researches*, 14(2), 137–162. <https://doi.org/10.1007/bf01204284>
- O'Reilly, B. M., Hauser, F., Jacob, A. W. B., & Shannon, P. M. (1996). The lithosphere below the Rockall Trough: Wide-angle seismic evidence for extensive serpentinisation. *Tectonophysics*, 255(1–2), 1–23. [https://doi.org/10.1016/0040-1951\(95\)00149-2](https://doi.org/10.1016/0040-1951(95)00149-2)
- Osmundsen, P. T., & Ebbing, J. (2008). Style of extension offshore mid-Norway and implications for mechanisms of crustal thinning at passive margins. *Tectonics*, 27, TC6016. <https://doi.org/10.1029/2007TC002242>
- Pérez-Gussinyé, M. (2012). A tectonic model for hyperextension at magma-poor rifted margins: An example from the West Iberia–Newfoundland conjugate margins. *Geological Society, London, Special Publications*, 369. <https://doi.org/10.1144/sp369.19>
- Peron-Pinvidic, G., & Osmundsen, P. T. (2016). Architecture of the distal and outer domains of the Mid-Norwegian rifted margin: Insights from the Rån–Gjallar ridges system. *Marine and Petroleum Geology*, 77, 280–299. <https://doi.org/10.1016/j.marpetgeo.2016.06.014>
- Petersen, K. D., & Schiffer, C. (2016). Wilson cycle passive margins: Control of orogenic inheritance on continental breakup. *Gondwana Research*, 39, 131–144. <https://doi.org/10.1016/j.gr.2016.06.012>
- Planke, S., Millett, J. M., Maharjan, D., Jerram, D. A., Abdelmalak, M. M., Groth, A., ... Myklebust, R. (2017). Igneous seismic geomorphology of buried lava fields and coastal escarpments on the Vøring volcanic rifted margin. *Interpretation*, 5(3), SK161–SK177. <https://doi.org/10.1190/int-2016-0164.1>
- Planke, S., Rasmussen, T., Rey, T., & Myklebust, R. (2005). Seismic characteristics and distribution of volcanic intrusions and hydrothermal vent complexes in the Vøring and Møre basins. In A. G. Doré & B. A. Vining (Eds.), *Petroleum geology: North-west Europe and global perspectives—Proceedings of the 6th Petroleum Geology Conference* (pp. 833–844). London: Geological Society.
- Planke, S., Skogseid, J., & Eldholm, O. (1991). Crustal structure off Norway, 62° to 70° north. *Tectonophysics*, 189(1–4), 91–107. [https://doi.org/10.1016/0040-1951\(91\)90489-F](https://doi.org/10.1016/0040-1951(91)90489-F)
- Planke, S., Svensen, H., Myklebust, R., Bannister, S., Manton, B., & Lorenz, L. (2015). *Geophysics and remote sensing* (pp. 1–16). Springer Berlin. https://doi.org/10.1007/11157_2014_6
- Planke, S., Symonds, P. A., Avelstad, E., & Skogseid, J. (2000). Seismic volcanostratigraphy of large-volume basaltic extrusive complexes on rifted margins. *Journal of Geophysical Research*, 105, 19,333–19,351.
- Raum, T., Mjelde, R., Digraanes, P., Shimamura, H., Shiobara, H., Kodaira, S., ... Thorbjørnsen, T. (2002). Crustal structure of the southern part of the Vøring Basin, mid-Norway margin, from wide-angle seismic and gravity data. *Tectonophysics*, 355(1–4), 99–126. [https://doi.org/10.1016/S0040-1951\(02\)00136-1](https://doi.org/10.1016/S0040-1951(02)00136-1)
- Raum, T., Mjelde, R., Shimamura, H., Murai, Y., Bråstein, E., Karpuz, R. M., ... Kolstø, H. J. (2006). Crustal structure and evolution of the southern Vøring Basin and Vøring Transform Margin, NE Atlantic. *Tectonophysics*, 415(1–4), 167–202. <https://doi.org/10.1016/j.tecto.2005.12.008>
- Ren, S., Faleide, J. I., Eldholm, O., Skogseid, J., & Gradstein, F. (2003). Late Cretaceous–Paleocene tectonic development of the NW Vøring Basin. *Marine and Petroleum Geology*, 20(2), 177–206. [https://doi.org/10.1016/S0264-8172\(03\)00005-9](https://doi.org/10.1016/S0264-8172(03)00005-9)
- Ren, S., Skogseid, J., & Eldholm, O. (1998). Late Cretaceous–Paleocene extension on the Vøring volcanic margin. *Marine Geophysical Researches*, 20(4), 343–369. <https://doi.org/10.1023/a:1004554217069>
- Reston, T. J. (2007). Extension discrepancy at North Atlantic nonvolcanic rifted margins: Depth-dependent stretching or unrecognized faulting? *Geology*, 35(04), 367–370.
- Reynisson, R. F., Ebbing, J., Lundin, E., & Osmundsen, P. T. (2010). Properties and distribution of lower crustal bodies on the mid-Norwegian margin. *Geological Society, London, Petroleum Geology Conference Series*, 7, 843–854. <https://doi.org/10.1144/0070843>
- Reynolds, P., Planke, S., Millett, J. M., Jerram, D. A., Trulsvik, M., Schofield, N., & Myklebust, R. (2017). Hydrothermal vent complexes offshore northeast Greenland: A potential role in driving the PETM. *Earth and Planetary Science Letters*, 467, 72–78. <https://doi.org/10.1016/j.epsl.2017.03.031>
- Roberts, A. W., White, R. S., & Christie, P. A. F. (2009). Imaging igneous rocks on the North Atlantic rifted continental margin. *Geophysical Journal International*, 179(2), 1024–1038. <https://doi.org/10.1111/j.1365-246X.2009.04306.x>
- Rosenbaum, G., Weinberg, R. F., & Regenauer-Lieb, K. (2008). The geodynamics of lithospheric extension. *Tectonophysics*, 458(1–4), 1–8.
- Rouzo, S., Klingelhöfer, F., Jonquet-Kolstø, H., Karpuz, R., Kravik, K., Mjelde, R., ... Géli, L. (2006). 2-D and 3-D modelling of wide-angle seismic data: An example from the Vøring volcanic passive margin. *Marine Geophysical Researches*, 27(3), 181–199. <https://doi.org/10.1007/s11001-006-0001-3>
- Rubin, A. M. (1995). Propagation of magma-filled cracks. *Annual Review of Earth and Planetary Sciences*, 23(1), 287–336. <https://doi.org/10.1146/annurev.ea.23.050195.001443>
- Saltus, R. W., & Blakely, R. J. (2011). Unique geologic insights from “non-unique” gravity and magnetic interpretation. *GSA Today*, 21, 4–10.
- Sandwell, D. T., & Smith, W. H. F. (2009). Global marine gravity from retracked Geosat and ERS-1 altimetry: Ridge segmentation versus spreading rate. *Journal of Geophysical Research*, 114, B01411. <https://doi.org/10.1029/2008JB006008>
- Saunders, A. D., Fitton, J. G., Kerr, A. C., Norry, M. J., & Kent, R. W. (1997). The North Atlantic igneous province. In J. J. Mahoney & M. F. Coffin (Eds.), *Large igneous provinces: Continental, oceanic, and planetary flood volcanism* (pp. 45–93). Washington, DC: American Geophysical Union.
- Scheck-Wenderoth, M., Raum, T., Faleide, J. I., Mjelde, R., & Horsfield, B. (2007). The transition from the continent to the ocean: A deeper view on the Norwegian margin. *Journal of the Geological Society*, 164(4), 855–868. <https://doi.org/10.1144/0016-76492006-131>
- Schiffer, C., Balling, N., Ebbing, J., Jacobsen, B. H., & Nielsen, S. B. (2016). Geophysical-petrological modelling of the East Greenland Caledonides—Isostatic support from crust and upper mantle. *Tectonophysics*, 692(Part A), 44–57. <https://doi.org/10.1016/j.tecto.2016.06.023>
- Schiffer, C., Jacobsen, B. H., Balling, N., Ebbing, J., & Nielsen, S. B. (2015). The East Greenland Caledonides—Teleseismic signature, gravity and isostasy. *Geophysical Journal International*, 203(2), 1400–1418. <https://doi.org/10.1093/gji/ggv373>
- Schindlwein, V., & Jokat, W. (1999). Structure and evolution of the continental crust of northern East Greenland from integrated geophysical studies. *Journal of Geophysical Research*, 104, 15,227–15,245.

- Schindwein, V., & Jokat, W. (2000). Post-collisional extension of the East Greenland Caledonides: A geophysical perspective. *Geophysical Journal International*, 140(3), 559–567. <https://doi.org/10.1046/j.1365-246X.2000.00036.x>
- Schmidt-Aursch, M. C., & Jokat, W. (2005). The crustal structure of central East Greenland—I: From the Caledonian orogen to the Tertiary igneous province. *Geophysical Journal International*, 160(2), 736–752. <https://doi.org/10.1111/j.1365-246X.2005.02514.x>
- Skelton, A., Whitmarsh, R., Arghe, F., Crill, P., & Koyi, H. (2005). Constraining the rate and extent of mantle serpentinization from seismic and petrological data: Implications for chemosynthesis and tectonic processes. *Geofluids*, 5(3), 153–164. <https://doi.org/10.1111/j.1468-8123.2005.00111.x>
- Skogseid, J. (2001). Volcanic margins: Geodynamic and exploration aspects. *Marine and Petroleum Geology*, 18, 457–461.
- Skogseid, J., & Eldholm, O. (1989). Vøring Plateau continental margin: Seismic interpretation, stratigraphy, and vertical movements. In O. Eldholm, et al. (Eds.), *Proceedings of the ODP, Scientific Results*, 104, (pp. 993–1030). College Station, TX: Ocean Drilling Program.
- Skogseid, J., Planke, S., Faleide, J. I., Pedersen, T., Eldholm, O., & Neverdal, F. (2000). NE Atlantic continental rifting and volcanic margin formation. In A. E. A. Nottvedt (Ed.), *Dynamics of the Norwegian margin* (pp. 295–326). London: Geological Society, London, Special Publications.
- Stratford, W., Thybo, H., Faleide, J. I., Olesen, O., & Tryggvason, A. (2009). New Moho map for onshore southern Norway. *Geophysical Journal International*, 178(3), 1755–1765. <https://doi.org/10.1111/j.1365-246X.2009.04240.x>
- Svensen, H., Planke, S., Malthes-Sorensen, A., Jamtveit, B., Myklebust, R., Rasmussen Eidem, T., & Rey, S. S. (2004). Release of methane from a volcanic basin as a mechanism for initial Eocene global warming. *Nature*, 429(6991), 542–545.
- Tegner, C., Leshner, C. E., Larsen, L. M., & Watt, W. S. (1998). Evidence from the rare-earth-element record of mantle melting for cooling of the Tertiary Iceland plume. *Nature*, 395(6702), 591–594. http://www.nature.com/nature/journal/v395/n6702/supinfo/395591a0_S1.html
- Theissen-Krah, S., Zastrozhnov, D., Abdelmalak, M. M., Schmid, D. W., Faleide, J. I., & Gernigon, L. (2017). Tectonic evolution and extension at the Møre Margin—Offshore mid-Norway. *Tectonophysics*, 721, 227–238.
- Thomsen, L. (1986). Weak elastic anisotropy. *Geophysics*, 51(10), 1954–1966. <https://doi.org/10.1190/1.1442051>
- Tsikalas, F., Faleide, J. I., Eldholm, O., & Antonio Blach, O. (2012). 5 - The NE Atlantic conjugate margins. In D. G. R. W. Bally (Ed.), *Regional geology and tectonics: Phanerozoic passive margins, cratonic basins and global tectonic maps* (pp. 140–201). Boston, MA: Elsevier. <https://doi.org/10.1016/B978-0-444-56357-6.00004-4>
- Tsikalas, F., Faleide, J. I., & Kuznir, N. J. (2008). Along-strike variations in rifted margin crustal architecture and lithosphere thinning between northern Vøring and Lofoten margin segments off mid-Norway. *Tectonophysics*, 458, 68–81.
- Verhoef, J., MacNab, R., Roest, W., Arkani-Hamed, J., & Members of the Project Team (1996). Magnetic anomalies of the Arctic and North Atlantic Oceans and adjacent land areas (GAMM-AA5: Gridded Aeromagnetic and Marine Magnetics of the North Atlantic and Arctic, 5km) Rep. 3125, Geological Survey of Canada, Open File Report 3125a.
- Voss, M., & Jokat, W. (2007). Continent–ocean transition and voluminous magmatic underplating derived from P-wave velocity modelling of the East Greenland continental margin. *Geophysical Journal International*, 170(580–604).
- Voss, M., Schmidt-Aursch, M. C., & Wilfried, J. (2009). Variations in magmatic processes along the East Greenland volcanic margin. *Geophysical Journal International*, 177, 755–782.
- Wangen, M., Mjelde, R., & Faleide, J. I. (2011). The extension of the Vøring Margin (NE Atlantic) in case of different degrees of magmatic underplating. *Basin Research*, 23(1), 83–100. <https://doi.org/10.1111/j.1365-2117.2010.00467.x>
- Weigel, W., Flüß, E. R., Miller, H., Butzke, A., Dehghani, G. A., Gebhardt, V., ... Groekort Study Group (1995). Investigations of the East Greenland continental margin between 70° and 72°N by deep seismic sounding and gravity studies. *Marine Geophysical Researches*, 17(2), 167–199. <https://doi.org/10.1007/bf01203425>
- White, R. S. (1988). A hot-spot model for early Tertiary volcanism on the N Atlantic. In A. C. Morton & L. M. Parson (Eds.), *Early Tertiary volcanism and the opening of the NE Atlantic*, Geological Society Special Publication (pp. 3–13).
- White, R. S., McKenzie, D., & O’Nions, R. K. (1992). Oceanic crustal thickness from seismic measurements and rare earth element inversions. *Journal of Geophysical Research*, 97(B13), 19,683–19,715. <https://doi.org/10.1029/92JB01749>
- White, R. S., & McKenzie, D. P. (1989). Magmatism at rift zones: The generation of volcanic continental margins and flood basalts. *Journal of Geophysical Research*, 94, 7685–7729.
- White, R. S., & Smith, L. K. (2009). Crustal structure of the Hatton and the conjugate East Greenland rifted volcanic margins, NE Atlantic. *Journal of Geophysical Research*, 114, B02305. <https://doi.org/10.1029/2008JB005856>
- White, R. S., Smith, L. K., Roberts, A. W., Christie, P. A. F., Kusznir, N. J., & the rest of the iSIMM Team (2008). Lower-crustal intrusion on the North Atlantic continental margin. *Nature*, 452, 460–465.
- White, R. S., Spence, G. D., Fowler, S. R., McKenzie, D. P., Westbrook, G. K., & Bowen, A. N. (1987). Magmatism at rifted continental margins. *Nature*, 330, 439–444.
- White, W. M., & Klein, E. M. (2014). Composition of the oceanic crust. In *Treatise on geochemistry* (2nd ed., pp. 457–496). Amsterdam, Netherlands: Elsevier.
- Wilson, M., & Wheeler, W. (2002). Magma migration through the continental crust: 3-D seismic and thermo-mechanical constraints on sites of crustal contamination. In *Eos Transactions, American Geophysical Union* (pp. F1407). Washington, DC.
- Zehnder, M. C., Mutter, J., & Buhl, P. (1990). Deep seismic and geochemical constraints on the nature of rift-induced magmatism during breakup of the North Atlantic, Seismic probing of continents and their margins. *Tectonophysics*, 173(1), 545–565. [https://doi.org/10.1016/0040-1951\(90\)90245-4](https://doi.org/10.1016/0040-1951(90)90245-4)
- Ziegler, P. A. (1988). *Evolution of the Arctic–North Atlantic and the Western Tethys* (Vol. 43). American Association of Petroleum Geologist Memoir.
- Ziegler, P. A., & Cloetingh, S. (2004). Dynamic processes controlling evolution of rifted basins. *Earth Science Reviews*, 64, 1–50.

Appendix B

The Development of Volcanic Sequences at Rifted Margins: New Insights from the Structure and Morphology of the Vøring Escarpment, mid-Norwegian Margin.

By

Mansour M. Abdelmalak, Sverre Planke, Jan Inge Faleide, Dougal A. Jerram, Dmitry Zastrozhnov, Sigurd Eide & Reidun Myklebust

Journal of Geophysical Research: Solid Earth, 121, 5212–5236

RESEARCH ARTICLE

10.1002/2015JB012788

Key Points:

- Volcanostratigraphic mapping of the lava delta system during the formation of the Vøring Escarpment
- Different control factors leading to the escarpment segmentation with different segment morphology
- Preservation of subbasalt basins and structural highs in the Vøring Marginal High

Correspondence to:

M. M. Abdelmalak,
m.m.abdelmalak@geo.uio.no;
abdelmalak_mansour@yahoo.fr

Citation:

Abdelmalak, M. M., S. Planke, J. I. Faleide, D. A. Jerram, D. Zastrozhnov, S. Eide, and R. Myklebust (2016), The development of volcanic sequences at rifted margins: New insights from the structure and morphology of the Vøring Escarpment, mid-Norwegian Margin, *J. Geophys. Res. Solid Earth*, 121, 5212–5236, doi:10.1002/2015JB012788.

Received 31 DEC 2015

Accepted 12 JUN 2016

Accepted article online 15 JUN 2016

Published online 4 JUL 2016

The development of volcanic sequences at rifted margins: New insights from the structure and morphology of the Vøring Escarpment, mid-Norwegian Margin

M. M. Abdelmalak¹, S. Planke^{1,2}, J. I. Faleide¹, D. A. Jerram^{1,3,4}, D. Zastrozhnov¹, S. Eide⁵, and R. Myklebust⁶

¹Centre for Earth Evolution and Dynamics, University of Oslo, Oslo, Norway, ²Volcanic Basin Petroleum Research, Oslo Research Park, Oslo, Norway, ³DougalEARTH Ltd., Solihull, UK, ⁴Earth, Environmental and Biological Sciences, Queensland University of Technology, Brisbane, Queensland, Australia, ⁵First Geo AS, Fornebu, Norway, ⁶TGS, Asker, Norway

Abstract On the Vøring Margin offshore mid-Norway, Paleogene continental breakup was characterized by the extrusion of large volumes of flood basalts erupted in different depositional environments. The transition from subaerial to submarine emplacement environment is marked by the formation of the Vøring Escarpment which records the early encroachment of flood basalt into the basin and the buildup of a lava delta system. The increased availability of new and reprocessed high-quality seismic data allows a more detailed characterization of the along-strike and across-strike continuity and variability of the different volcanic seismic facies units. Detailed seismic interpretation shows that the ~350 km long NE-SW trending Vøring Escarpment is a prominent feature along the Vøring Margin with a height ranging between 200 and 1600 m. Structurally, the Vøring Escarpment is segmented along strike into five segments (E1–E5) with different controlling factors leading to variation in accommodation space. Relative sea level change and magma supply are the major controlling factors for segments E2 and E4 which are characterized by a well-developed lava delta system and significant escarpment height. Tectonic movements along the Jan Mayen Fracture Zone resulted in second-order segmentation of the E1 segment into pseudoescarpments with a very thin lava delta system and limited escarpment height. Segments E3 and E5, situated along the flanks of Cretaceous/Paleocene highs, are controlled by the structural highs, which were possibly reactivated during breakup time. Our mapping results provide crucial information about the paleogeography and yield important information regarding the paleo-water depth and depocenter locations prior to and during the breakup of the Vøring Margin.

1. Introduction

This paper presents new results from a detailed seismic interpretation of the Vøring Margin offshore mid-Norway, concentrating on the ~350 km long NE-SW aligned Vøring Escarpment. Historically it has been difficult to acquire good reflection seismic records from sequences containing major lava flows. This has changed with the increased availability of high-quality data and improved processing techniques. Together with reinterpretation of old data on volcanic rifted margins, newer seismic data have revealed that breakup magmatism is common and widespread, although its intensity and internal character may vary significantly both along margins and between margin segments.

Volcanic successions display a variety of reflection configurations in the subsurface. These are indicative of the depositional environment during flood volcanism (e.g., subaerial versus subaqueous) and the subsequent mass transport of material within the margin [e.g., Planke *et al.*, 2000; Berndt *et al.*, 2001b; Wright *et al.*, 2012; Watton *et al.*, 2013]. Recent advances in processing seismic data [e.g., Gallagher and Dromgoole, 2007; Soubaras and Dowle, 2010; Woodburn *et al.*, 2014] have greatly improved the imaging of geometries within volcanic deposits as well as in the sediments below the basalts. This allows better constraints and interpretation of the breakup-related igneous rocks, the paleogeography, and the rift topography before and during the breakup. Furthermore, the understanding of the temporal evolution of breakup volcanism has been improved significantly by several deep-sea drilling surveys: DSDP Leg 38 (Deep Sea Drilling Project) and ODP Leg 104 (Oceanic Drilling Program) on the Vøring Margin [Eldholm *et al.*, 1987, 1989; Planke, 1994; Planke and Eldholm, 1994] and ODP Legs 152 [Larsen and Saunders, 1998; Planke and Cambray, 1998] and 163 [Larsen *et al.*, 1999] on the Southeast Greenland Margin (Figure 1).

Figure 1. Offshore distribution of volcanic facies units in the Vøring Margin and regional reconstruction at ~52 Ma (inset; based on *Gaia et al.* [2009]) showing onshore and offshore volcanic rocks in the North Atlantic Igneous Province (NAIP). The DSDP/ODP Leg locations are indicated in the general reconstruction: a, ODP Leg 152; b, ODP Leg 163; c, ODP Leg 104; d, DSDP Leg 38. The profile A-A' (in two-way traveltimes) shows an example of 2-D mapping of the different seismic facies units on the Vøring Margin. The 2-D seismic profiles used in this study are indicated in the map. BL: Bivrost Lineament; COB: continent-ocean boundary; FG: Fenris Graben; SH: Skoll High; GH: Grimm High; GS: Gleipne Saddle; JMFZ: Jan Mayen Fracture Zone, LSF: Lower Series Flows; NGR: North Gjallar Ridge; SDR: Seaward Dipping Reflector; SGR: South Gjallar Ridge; MH: Mimir High; YH: Ygg High.

Multichannel seismic data have allowed the definition and characterization of the seismic “volcanostratigraphy” concept based on their shape, reflection pattern, and boundary reflections [e.g., *Planke et al.*, 2000; *Berndt et al.*, 2001b]. On the Norwegian Margin, several volcanic seismic facies units have been identified: (1) Landward Flows, (2) Lava Delta, (3) Inner Flows, (4) inner Seaward Dipping Reflectors (inner SDR), (5) Outer High, and (6) outer Seaward Dipping Reflectors (outer SDR) (see Table 1 and Figure 1). Such facies succession represents a typical volcanic rifted margin sequence and describes the evolution of the breakup extrusive complex landward of, and/or very close to, the first magnetic seafloor spreading anomalies.

Escarpment structure and lava delta systems that characterize the volcanic margins worldwide are identified in seismic profiles and in onshore analog margins such as in West Greenland [*Pedersen et al.*, 1998; *Japsen et al.*, 2005], Iceland [e.g., *Watton et al.*, 2013], and Antarctica [*Smellie et al.*, 2011]. By definition the escarpment is a steep slope or long cliff formed by erosion or by vertical movements along a fault. The term is often used interchangeably with scarp but is more accurately associated with cliffs produced

Table 1. Description of the Volcanic Seismic Facies Units Based On *Planke and Alvestad* [1999], *Planke et al.* [1999, 2000], *Berndt et al.* [2001b], *Rey et al.* [2008], *Nelson et al.* [2009b], *Jerram et al.* [2009], and *Abdelmalak et al.* [2016]

Seismic Facies Unit	Shape	Reflections Characteristics		Volcanic Facies	Depositional Environment
		Boundaries	Internal		
Inner Flows	Sheet	Top: high amplitude, disrupted, rough being onlapped or concordant; base: negative polarity often obscured	Chaotic sheet-like body of very disrupted or hummocky reflections	Massive and fragmented flows, volcanoclastics and hyaloclastics	Shallow marine deposited in broad basin
Lava Delta	Bank	Top: high amplitude or reflection truncation; base: reflection truncation or termination	Progradational reflection configuration	Massive and fragmented basalts and volcanoclastics	Coastal
Landward Flows	Sheet	Top: high amplitude, smooth being onlapped or concordant; base: low amplitude, disrupted	Parallel to subparallel. High-amplitude disrupted	Flood basalts	Subaerial or shallow marine flood basalts deposited on a plain or in broad basin
Inner SDR	Wedge	Top: intermediate to high amplitude, smooth with pseudoescape; base: seldom defined	Divergent arcuate or sometimes a divergent-planar pattern	Flood basalts	Subaerial flood basalts deposited in subsiding structure
Outer High	Mound	Top: high amplitude, disrupted and rough; base: not visible	Chaotic	Hyaloclastic flows and volcanoclastics	Shallow marine environment
Outer SDR	Wedge	Top: intermediate to high amplitude, smooth with pseudoescape; base: seldom defined	Divergent arcuate internal reflectors, lower amplitude than the Inner SDR	Flood basalts mixed with pillow basalts sediments and sills	Deep marine depositional environment
Lava Flows	Sheet	Top: high amplitude with pseudoescape; base: not visible	Chaotic	Flood basalts	Subaerial to shallow submarine depositional environment
Lower Series Flows	Sheet	Top: high amplitude, strong and continuously negative in polarity reflection; base: seldom defined	Wavy to continuous subparallel reflections with an internal disrupted and hummocky shape	Flood basalts mixed with volcanoclastic sediments, hyaloclastics, and sills	Subaqueous flows extruding and intruding into wet sediments

by depositional/erosional processes rather than those produced by faulting. The escarpment marks the transition from subaerial to submarine environment and records the early encroachment of the flood basalt into the basin and the buildup of a lava delta system (Figure 2). The transition from one environment to another provides a means to reconstruct the relative position of the shoreline and the water depth at the time of the delta progradation [e.g., *Jerram et al.*, 2009; *Wright et al.*, 2012] and is therefore an important structure that can shed light on the paleogeography and topography during the development of the volcanic margin.

When the escarpment morphology is controlled by relative sea level changes and/or magma supply, coastal and subaqueous erosion are enhanced and the escarpment is defined as the height of the delta front corresponding to the height between the Inner Flows and the Landward Flows (Figure 2). Tectonic movement along faults may considerably influence the morphology and structure of the escarpment, emphasizing the polygenetic nature of some of these landforms. In this case a large fault scarp could result, and greatly increasing the height between the Landward Flows and the Inner Flows.

On the Vøring Margin, the final continental breakup resulted in voluminous igneous activity generating both extrusive and intrusive rocks in the adjacent sedimentary basin and preexisting continental crust (profile A-A' in Figure 1) [Planke et al., 2005; Scheck-Wenderoth et al., 2007; Breivik et al., 2014]. Large volumes of flood basalts erupted in submarine to subaerial settings [e.g., Berndt et al., 2001b] and created the Vøring Escarpment. The identification of rift topography before/during breakup is improved by the determination of the accommodation space which gives better constraints on the paleogeography of the area during the buildup of the lava delta system. It yields valuable information regarding the paleo-sea level and the location of depocenters directly prior to the volcanism. Furthermore, the paleo-water depths, which are indicated by the delta front heights, provide important input parameters in basin and petroleum system modeling. Therefore it is imperative that the volcanic extrusive should be carefully investigated to understand their emplacement and influence on the margin structure. Until now, no detailed investigations have been made on the Vøring Escarpment. The main reasons have been the lack of good quality seismic data and poor data

Figure 2. (a) Uninterpreted and (b) interpreted seismic sections highlighting key seismic facies units of the lava delta system. (c) Schematic representation (not to scale) of a developing lava-fed delta. TB: top basalt, BB: base basalt. Data courtesy of TGS.

coverage across the outer part of the volcanic margins because all the prospects are being focused on the sedimentary basin far from the extrusive volcanic rocks. In recent years, some prospects located in volcanic systems have been successful, such as the Rosenbank discovery in the North Atlantic [Schofield and Jolley, 2013] and the Neuquén Basin in Argentina [Rodríguez Monreal et al., 2009]. Such new hydrocarbon discoveries make the volcanic margin a new target and new frontier [e.g., Jerram, 2015].

Recently, the increased density of seismic data on the Vøring Margin and improvement in seismic imaging now permit more detailed and accurate studying and characterizing of the along-strike and across-strike continuity and variability of the different volcanic seismic units. We are thus able to present the structure and the morphology of the Vøring Escarpment using a dense grid of high-quality seismic data combined with gravity and magnetic surveys. Investigating in detail the reflection geometries of the Vøring Escarpment has allowed us to improve the constraints on how flood basalt develops in its transition from a subaerial to a submarine environment. This in turn enabled us to reconstruct the development and evolution of the Vøring Escarpment and to address the different factors that controlled the buildup of this escarpment. Our results provide more information about how the paleoshoreline evolved at the time of volcanism and give

more constraints on the subbasalt geology, accommodation space, paleo–water depth, and volcanotectonic processes occurring during the formation of the escarpment. In a broader context, our results will improve the understanding of escarpment development in volcanic margins.

2. Context of the Vøring Margin Development and Breakup

The Vøring rifted margin, part of the mid-Norwegian Margin, is a ~350 km NE-SW trending area bounded by the Jan Mayen Fracture Zone to the southwest and the Bivrost Lineament to the northeast [Blystad *et al.*, 1995]. The Vøring Margin comprises the Trøndelag Platform, the Vøring Basin, and the Vøring Marginal High (Figure 1) and is the consequence of a significant Late Jurassic to Early Cretaceous crustal thinning, with episodic subsidence leading to a very thick Cretaceous depocenter [Blystad *et al.*, 1995; Skogseid *et al.*, 2000; Scheck-Wenderoth *et al.*, 2007]. This extremely thick Cretaceous sedimentary sequences (~10 km) mostly concealed the geometry of the deeper synrift sequences in the Vøring Basin [Faleide *et al.*, 2008]. The thinned crust at the Vøring Margin was, again, the locus of a new phase of extensional deformation in the latest Cretaceous to Early Paleocene. This prebreakup rifting event has been particularly documented in the outer Vøring Basin [Eldholm *et al.*, 2002; Gernigon *et al.*, 2003; Ren *et al.*, 2003; Gernigon *et al.*, 2004]. The Late Jurassic–Early Cretaceous extension and the Late Cretaceous extension formed large-scale rift blocks or structural highs (Early Cretaceous ridges and highs: Rån Ridge and Utgard High; Late Cretaceous–Early Paleocene ridges and highs: North/South Gjallar Ridge, Nyk High, Mimir High, Ygg High, Grimm High, and Skoll High) (Figure 1).

In the Vøring Margin, the magmatic-tectonic processes that led to the final breakup are restricted to a 100–150 km wide region of the outer Vøring Basin and the Vøring Marginal High presently situated to the east of the first oceanic magnetic chrons (Figure 1). The volcanic sequences emplaced during the onset of the breakup, partially masked seismically the Late Cretaceous–Early Eocene and older sedimentary strata and continental structures of the outer Vøring Margin.

3. Data and Methods

3.1. Data

This work is based on the interpretation of a dense grid of more than 500 regional 2-D seismic lines with a total length of more than 80,000 km and covering the entire area of the Vøring Margin with a line spacing ranging between 0.2 and 2 km. This exceptional data set includes recently acquired as well as reprocessed high-quality seismic reflection data that provide improved imaging of the deeper parts of the margin. The line density is high enough to correlate laterally the different facies units along the mid-Norwegian Margin. Some of the seismic lines we have used on the Vøring Margin are shown in Figure 1. Our seismic interpretation was carried out within the framework of a project covering the entire mid-Norwegian Margin. Seismic lines were depth converted with HiQbe™ (<http://www.first-geo.com/products/hiqbe/norway>), a regional high-quality seismic velocity cube covering the mid-Norwegian Margin compiled from 536 seismic stacking velocity data sets and 230 check shots from wells and vertical seismic profile data [First Geo AS and TGS, 2015].

In addition to seismic profile data, we used available public-domain bathymetry (e.g., GEBCO, the GEBCO_08 Grid, version 20100927, "<http://www.gebco.net>"), satellite gravity [Sandwell and Smith, 2009], released magnetic data (EMAG2) [Maus *et al.*, 2009], and local aeromagnetic data reprocessed by Geological Survey of Norway and TGS.

High-pass-filtered potential field data (100 km) were compiled for most of the studied area to better understand the intermediate to deep structures of the Vøring Margin. The high-pass-filtered gravity and magnetic data were converted to pseudohorizons and loaded into the seismic interpretation project to facilitate an integrated seismic-gravity-magnetic interpretation. The 100 km high-pass-filtered Bouguer anomaly makes it possible to remove the regional gravity anomalies caused by density variations in the deeper crust and the upper mantle [e.g., Berndt, 2002]. A cutoff wavelength of 100 km emphasizes density contrasts with source depth <10 km. Thus, these data provide important structural and lithological information about basins. Magnetic data are most useful for determining the presence of trends and depths of mafic volcanic rocks and metamorphic terrains. The 100 km high-pass-filtered magnetic anomalies are the best way to detect variations in continental crystalline crustal terranes and sea floor-spreading anomalies.

3.2. Methods: Mapping the Volcanic Facies Units of the Vøring Margin

In this study we use the concept of seismic volcanostratigraphy for the interpretation of large volcanic constructions [e.g., Planke et al., 2000; Berndt et al., 2001b]. Our dense grid of high-quality seismic data allows us to refine the mapping and to study the variability in space and time of the extrusive rocks. Moreover, our new observations allow a better definition and interpretation of the genetic processes and the evolution of the different seismic facies units. First of all, we interpret the volcanic succession as one seismic sequence, i.e., a depositional unit of genetically related strata bounded by unconformities or correlative conformities. This step is called the seismic sequence analysis. The second step consists of the mapping of seismic facies units within the volcanic sequence. Other igneous-related feature like sill intrusion and associated hydrothermal vent complexes are included in the mapping [e.g., Svensen et al., 2004; Hansen et al., 2005; Planke et al., 2005, 2015]. Finally, we interpret the volcanic nature of the mapped volcanic seismic facies units taking into account well ties, field analogs, wave propagation phenomena such as internal multiples, and seismic processing. It is frequently possible to associate objectively identified seismic facies units with typical volcanic deposits, which in turn may be indicative of distinct emplacement environments [Planke et al., 2000; Berndt et al., 2001b; Jerram, 2002; Jerram et al., 2009].

3.2.1. Seismic Sequence Analysis

The regional extent of the extrusive breakup complex is interpreted on seismic reflection data primarily on the basis of its top reflection. The top of the basaltic sequences is a top unconformity/conformity between the postbreakup sediments and the underlying volcanic rocks (predominantly basalt). The top lava also represents the continentward continuation of the top oceanic basement reflection further west. The top basalt reflection is easily identifiable due to the high impedance contrast between the postbreakup sediments and the volcanic rocks. It is often smooth but may locally be irregular and faulted and sometimes shows evidence of pseudoescarpments [Planke et al., 2000; Berndt et al., 2001b].

The basal sequence boundary is frequently difficult or impossible to identify. This is often due to imaging problems related to the seismic properties of the overlying basaltic constructions. The high impedance contrasts at the top and/or within the lava pile give rise to a scattering of seismic energy. In addition, the lower boundary of the volcanic extrusive complexes is often poorly defined [e.g., Planke et al., 2015]. The base of the basalts can also consist of a transition and composite zone of sequentially deposited extrusive volcanic rocks mixed with subsequently intruded sills and volcanoclastic sediments. Locally, basaltic sequences lie over the intruded sedimentary basin. Then, we assume that the base of the basalts should be above the sill intrusions identified on seismic profiles (e.g., profile A-A' in Figure 1).

3.2.2. Volcanic Seismic Facies Units

A seismic facies unit is a part of the volcanic sequence and which has a uniform internal seismic character. It must be mappable, i.e., it must differ from the adjacent units, extend laterally to the neighboring seismic profiles, and must have a uniform overall shape. Here we focus on the seismic facies units that document the transition from subaerial (Landward Flows) to subaqueous (Lava Delta and Inner Flows) environments (Figure 2). The seismic characteristics of these facies unit are defined below and summarized in Table 1.

The *Landward Flows* are characterized by parallel to subparallel high-amplitude reflections. This seismic facies unit is commonly identified on seismic profiles landward and below the inner SDR. The top reflection is a strong, fairly smooth event. The external shape is sheet-like, whereas internal reflections are disrupted or hummocky and subparallel. The basal boundary may be identified as a negative in polarity reflection on high-quality multichannel seismic reflection data. Deeper reflections are discontinuous low-amplitude events. The unit frequently terminates at a regional escarpment or merges with prograding reflections of the Lava Delta facies unit (Figures 2a and 2b).

The *Inner Flows* form a sheet-like body of very disrupted or hummocky reflections. The top reflection is a high-amplitude, disrupted event. Internal reflection configuration is chaotic. The base is a weak negative-polarity reflection but is often difficult to identify. Overlying reflections are dominantly subparallel, whereas deeper reflections are frequently masked (Figures 2a and 2b). The unit is up to 60 km wide on the mid-Norwegian Margin and is typically several hundred meters thick.

The *Lava Delta* facies unit is characterized by a progradational reflection configuration. The upper boundary is determined by reflection terminations or a change in reflection geometry from fairly flat-laying to disrupted, arcuate events (Figures 2a and 2b). The lower boundary is interpreted as a surface connecting the lower

terminations of the prograding reflections but is often more difficult to identify. The Lava Delta unit has not been drilled on the Norwegian Margin, but a combined interpretation of well-defined reflection configuration compared with modern and ancient analogs suggests that this unit can be interpreted as the foresets of a lava delta.

Where the delta foresets are well imaged, it is possible to determine the transition from the Landward Flows (subaerial) to the Lava Delta (subaqueous) facies units (Figure 2). This transition from one environment to another provides a means to reconstruct the relative position of the shoreline and the water depth at the time of the delta migration [e.g., *Jerram et al.*, 2009; *Wright et al.*, 2012]. The lava flowing into the sea undergoes quenching and fragmentation into hyaloclastic breccias which were rapidly deposited downslope under gravitational processes to form inclined foresets and create the escarpment (Figure 2c).

As with the top of the Lava Delta providing information on the paleoshoreline through time, the thickness of the delta front and Inner Flows packages provides a very good proxy for paleo-water depth. It should be noted that the best estimate for the water depth can be made at the escarpment front as this is the most clearly imaged part of the seismic sections.

Generally, the Lava Delta facies unit displays similarities with siliciclastic delta systems, by filling the available accommodation space and by reacting to changes in relative sea level, variation in supply of material, and the vertical tectonic movements (subsidence or uplift) (Figure 2c). Delta growth is through deposition of new lava flows and hyaloclastic breccias, with successive phases of volcanism producing a stacking pattern geometry characteristic of the lava delta systems and resulting in the progradation of the shoreline. The geometry of the stacking depends on the dominant factor at the time of the deposition (i.e., the relative sea level changes, lava supply, and the tectonic vertical movements) (Figure 2c), making it possible to reconstruct the depositional environment and interpret the Lava Delta facies unit within a sequence stratigraphic framework [e.g., *Kjørboe*, 1999].

4. Results

4.1. The Vøring Marginal High Structure

The filtered magnetic anomaly map shows that the Vøring Escarpment marks a significant change in magnetic field between the Vøring Marginal High and the outer Vøring Basin to the east (Figure 3a). Basinward, the magnetic field is characterized by low-amplitude magnetic anomalies related to thick sedimentary units in the Vøring Basin in areas with little or no extrusive volcanic rocks. Oceanward of the escarpment, the Vøring Marginal High is characterized by elongated high-amplitude magnetic anomalies (Figure 3a).

The filtered Bouguer anomaly map reveals a complex pattern of high-amplitude and low-amplitude anomalies along the outer Vøring Margin (Figure 3b). The Bouguer effect is calculated as if the water layer is replaced by rock having the same density as nearby sediments at the same depth. The map therefore reflects basin and crustal density variations more accurately than free-air high-pass-filtered data. In the map, the North and South Gjallar ridges are characterized by high-amplitude positive gravity anomalies (Figure 3b). The Hel Graben, the Rån Basin, and the Fenris Graben are characterized by circular negative gravity anomalies. High-amplitude circular positive gravity anomalies are distributed along the Vøring Escarpment and correspond to positive elongated NE trending magnetic anomalies. One negative gravity anomaly corresponds to the high-amplitude magnetic anomaly parallel to the Vøring Escarpment. The gravity anomalies likely represent subbasalt structures, basins, and/or basement highs. On the seismic data, this anomaly corresponds to the new high that we named the Skoll High. Two other highs are also defined along the escarpment and named the Grimm High and the Ygg High.

4.2. The Vøring Escarpment Structure

The Vøring Escarpment is a ~350 km NE-SW trending feature along the entire Vøring Margin which characterizes the top basalt. It is situated between 66°5'N–68°15'N and 1°55'E–8°20'E. Using a dense grid of high-quality seismic data, we refined the structural mapping of the Vøring Escarpment. First of all, we were able to map top and base basalt on the Vøring Margin. The depth to the top of the basalts varies considerably (Figure 3c). The top basalt is deeper toward the Jan Mayen Fracture Zone (JMFZ) and in the Inner Flows domain. Generally, the top basalt is shallower on the Vøring Marginal High (VMH) especially above the Skoll High and the Grimm High (Figure 3c).

Figure 3. (a) The 100 km high-pass-filtered magnetic data. (b) The 100 km high-pass-filtered Bouguer anomaly. (c) Top basalt depth (in km). (d) Basalt thickness map. (e) Top basalt slope. We used the average velocity of 4 km/s in basalt to calculate the basalt thickness [e.g., Planke, 1994; Jerram *et al.*, 2009]. The foot of the scarp is indicated. FG: Fenris Graben; SH: Skoll High; GH: Grimm High; HG: Hel Graben; JMZF: Jan Mayen Fracture Zone; ND: Naglfar Dome; NGR: North Gjallar Ridge; SGR: South Gjallar Ridge; VMH Vøring marginal high; YH: Ygg high. Magnetic and gravity data courtesy of TGS.

Using our top and base basalt interpretation, we have established the basalt thickness by applying an average velocity of 4 km/s in basalt [e.g., Planke, 1994; Jerram *et al.*, 2009]. Assuming such a constant value, the basalt thickness increases from less than hundred meters in the Inner Flows to more than 6 km in the SDR domain (Figure 3d) where our mapping reveal separate depocenters within the extent of the SDR. In addition, thin basalts are identified above the Skoll High and in the Jan Mayen Fracture Zone areas. Top basalt does not show significant variation in slope although in some places it can exceed 10° (Figure 3e). Some structural highs mapped in the Fenris Graben on the basis of seismic data also show relatively steep slopes (see Figure 1). These NE-SW highs are parallel to the North Gjallar Ridge and its characteristic fault system. Interestingly, the Vøring Escarpment strikes parallel to the breakup axis in the Vøring Margin and to the different highs mapped in the Fenris Graben. The Skoll High represents a deep-seated topographic high

Figure 4. (a) Lava Delta (LD) thickness map along the Vøring Margin. (b) Landward Flows (LF) thickness map along the Vøring Margin. (c) Vøring Escarpment height. Segments E1 to E5 are identified. (d) Cross section along the Vøring Escarpment showing the height variation. The gray area indicates the average height of the escarpments. An average velocity of 4 km/s in basalt is used to calculate the lava delta thickness, the landward flows thickness, and the Vøring Escarpment height [e.g., Planke, 1994; Jerram *et al.*, 2009].

draped with lava. The base of the slope of the Skoll High is rather steep. This transition also defines the foot of the scarp (Figure 3).

On the Vøring Margin, we have mapped the extent of the Lava Delta facies unit using the progradational shape identified on the seismic profile. The Lava Delta facies unit covers an area of more than 4000 km², and its width varies along strike (Figure 1). We calculated the Lava Delta and the Landward Flows thicknesses in the area of the Lava Delta extent (Figures 4a and 4b). The Lava Delta thickness map (Figure 4a), the Landward Flows thickness map (Figure 4b), and the Vøring Escarpment height (Figure 4c), show that at a first-order observation the Vøring Escarpment is divided into five segments (E1 to E5) differing from each other in escarpment height, Lava Delta, and Landward Flows thicknesses (Figure 4).

By considering the delta front height as a criterion to define the escarpment height, the relief of the Vøring Escarpment varies between 200 m and 1600 m (Figures 4c and 4d). The E1 segment is characterized by a less developed escarpment relief with a height ranging between 200 and 500 m with an average of 400 m (Figure 4d). This area is also characterized by thin development of the Lava Delta (less than 500 m) and the Landward Flows (also less 500 m) (Figure 4). The lava delta system in the E1 segment is limited in extent. The E1 is located near to the Jan Mayen Fracture Zone and shows several pseudoescarpments. In the transitional area between the E1 and E2 segments, the Ygg High was identified on the basis of new seismic and potential field data. The transition between E1 and E2 is also characterized by a sharp increase of the escarpment height from 500 m to 1000 m (Figure 4d).

Figure 5. (a) Uninterpreted and (b) interpreted seismic sections (in two-way traveltime) highlighting the key facies units of the lava delta system of escarpment segment E2. (c) Schematic illustration of the depth-converted seismic section showing the accommodation space and the position of the paleoshoreline. The paleo-water depth is indicated: ~ 1.5 km. See Figures 1 and 4c for location. Data courtesy of TGS.

The E2 segment characterizes the South Gjallar Ridge area. This segment shows higher escarpment relief ranging between 800 and 1600 m (average height of 1300 m) and a well-developed lava delta system with a thick Lava Delta (ranging between 700 m and 1000 m) and thick Landward Flows (ranging between 500 m and 1500 m) (Figure 4). The transition between E2 and E3 is abrupt, and the escarpment height decreases from 800 m to 400 m (Figure 4d). In the E3 segment, situated above the Skoll High, we mapped a thin Lava Delta thickness (ranging between 300 m and 500 m) and thin Landward Flows (less than 500 m) (Figure 4). In the E4 segment, the escarpment relief is well developed with a height ranging between 800 and 1600 (Figure 4c) with an average height of 1200 m (Figure 4d). In this escarpment segment, situated at the North Gjallar Ridge, the lava delta system is well developed with thickness ranging between 700 m and 1500 m, and the Landward Flows are also thick (ranging between 500 m and 2000 m) (Figure 4). The transition between E4 and E5 is more gentle. The escarpment segment E5 is characterized by low height ranging between 200 and 500 m (average height of ~ 450 m) and a less developed lava delta system. The Grimm High was identified on the basis of new seismic profiles and potential field data. In this segment we mapped a thin Lava Delta thickness (less than 500 m) and thin Landward Flows (less than 500 m) (Figure 4).

Figure 6. (a) Uninterpreted and (b) interpreted seismic sections (in two-way traveltime) highlighting the key facies units of the lava delta system of the escarpment segment E4. (c) Schematic illustration of the depth-converted seismic section showing the accommodation space and the position of the paleoshoreline. The paleo-water depth is indicated: ~1.3 km. See Figures 1 and 4c for location. Data courtesy of TGS.

4.3. The Vøring Escarpment Morphology

We investigated in more detail the Vøring Escarpment morphology and the paleo-water depth for the different segments using seismic reflection data. Example of the representative lava delta system of segments E2 and E4 are shown in Figures 5 and 6, respectively. The Landward Flows, the Inner Flows, and the Lava Delta are well imaged in these seismic sections. High-amplitude reflections with saucer shape and abrupt terminations are interpreted as sill intrusions (Figures 5a, 5b, 6a, and 6b). The initial stacking pattern is progradational into the basin filling the accommodation space. The progradational Lava Delta is a good indication of the starting and end positions of the paleoshoreline (Figures 5c and 6c). The point where the delta deposits pinch out on the landward side marks the location of the paleoshoreline where the delta began to form. With sufficient supply of volcanic material the paleoshoreline moved basinward with the prograding volcanic sequences. The crest of the escarpment marks the end position of the paleoshoreline (Figures 5c and 6b). With a rapid and voluminous supply of volcanics into the basin, the accommodation space available at the point of entry of the volcanics into the sea is filled. This is seen clearly by the development of major prograding foresets imaged on seismic data (Figures 5a, 5b, 6a, and 6b). Slumped blocks with disrupted internal reflection configuration are commonly identified at the foot of the escarpment (Figure 7). With the

Figure 7. Seismic section (in two-way traveltimes) showing a zoom on slumped blocks along escarpment segment E4 indicated by disrupted internal reflection configuration. Data courtesy of TGS.

progradation of the Lava Delta, these slumped blocks will be covered by the supply of new material into the basin. Some of these slumped blocks show more than 4 km long and over 0.5 km height (Figure 6c). A step-back delta system identified in seismic profile (Figure 5b) developed as a result of rising sea level and/or decrease in magma supply into the basin. A paleo-water depth ranging between 1.3 and 1.5 km characterizes the segment E2 and E4 (Figures 5c and 6c).

In the E1 segment, the Vøring Escarpment is second-order segmented into several pseudoescarpments due to the tectonic effect of the Jan Mayen Fracture Zone. A seismic reflection example is shown in Figure 8. Multiple Lava Deltas and escarpments are identified in this section (Figures 8a and 8b). The Lava Delta shows a limited thickness. In this section two development stages are observed. The first stage is characterized by development of a normal lava delta system where the Inner Flows and the Lava Delta units are identified. This first system was then rotated, uplifted, and subsequently covered by a second lava delta system (Figure 8c). Vertical movements along the fracture zone can also play a role in the second-order segmentation of the E1 segment. The paleo-water depth along this segment is limited and does not exceed 0.5 km.

Representative examples of segments E3 and E5 are shown in Figures 9 and 10. In the E3 segment overlying the Skoll High, the lava delta system is limited in extent. Only the delta front defines the escarpment height even though it has a much larger scarp (Figure 9). In the Fenris Graben along the scarp (see Figure 1), the seismic image consists mainly of packages of inclined reflections that can be interpreted as sliding and slumping flows over an elevated topography (Figures 9a and 9b). This resulted in an increase of the accommodation space in the area (Figure 9c) with paleo-water depths of more than 1.5 km. The identification of slumped blocks in the Fenris Graben could indicate that this structure was tectonically active during the lava progradation.

Similar to the E3 segment, the E5 segment developed above a deep-seated high, the Grimm High, but in contrast to the E3 segment, no elevated topographic feature is identified. In this segment a paleo-water depth of 0.8 km has been measured (Figure 10c). Some hydrothermal vent complexes are identified in the sedimentary basin (Figures 10a and 10b). Again, similar to the E3 segment, the lava delta system is less developed. In these two segments (E3 and E5) the Lava Delta and Landward Flows are thinner than in the E2 and E4 segments (Figures 9c and 10c). The stacking pattern is progradational into the basin, filling the accommodation space and allowing the identification of the starting and end positions of the paleoshoreline (Figures 9c and 10c). Delta front collapses were also common during the development of the lava delta system; an example is shown in Figure 11. The initial stacking pattern is progradational into the basin filling the accommodation space. Sometimes, the unconsolidated delta front subsided causing fractures to propagate

Figure 8. (a) Uninterpreted and (b) interpreted seismic sections (in two-way traveltime) crossing the E1 escarpment segment. Multiple lava deltas and escarpments are identified in this section. (c) Schematic illustration of the depth-converted seismic section showing two lava delta systems. The first system is rotated and partly covered by a second lava delta system. The paleo-water depth for the second system is indicated: ~ 0.5 km. BB: base basalt, IF: inner flow, LD: lava delta, TB: top basalt. See Figures 1 and 4c for location. Data courtesy of TGS.

through the delta front (Figure 11c). Such fractures will have induced large-scale delta collapse, sliding/slumping, and gravity-driven debris avalanches into the basin. From the seismic profile, the Inner Flows are seen to be characterized by the presence of highs that appear to be the result of gas release in the area. Surface seepages are clearly identified and are related to the development of these highs (Figures 11a and 11b). The location of the highs and related gas seepages are shown in Figure 1. Generally, these highs are elongated and strike parallel to the fault system of the North Gjallar Ridge.

5. Discussion

5.1. Lava Delta System Component

As exploration in volcanic rifted margins increases, more seismic data shot over lava cover and/or, where extensive intrusions are present within the basin, have become more available [e.g., Planke *et al.*, 2005;

Figure 9. (a) Uninterpreted and (b) interpreted seismic sections (in two-way traveltime) highlighting the key facies units of the lava delta system of Vøring Escarpment segment E3. (c) Schematic illustration of the depth-converted seismic section showing the accommodation space and the position of the paleoshoreline. The paleo-water depth is indicated: ~1.8 km. See Figures 1 and 4c for location. Data courtesy of TGS.

Jerram et al., 2009; Woodburn et al., 2014; Schofield et al., 2015b). While significant developments, especially in seismic processing, have improved the quality of offshore studies and thus our understanding of these margins, work on the volcanic facies from outcrop analogs has also been instrumental for improving the interpretation of the subsurface geophysical data. Volcanic facies mapping in flood basalts has enabled the key geometries to be identified [e.g., Jerram, 2002], and their application to onshore sequences helps us to determine how to link subseismic-scale observations onshore with how they may be expressed offshore [e.g., Jerram et al., 2009; Schofield et al., 2015a]. It is becoming clear that the internal architecture of flood basalt sequences is markedly heterogeneous in terms of rock properties [e.g., Jerram et al., 2009; Nelson et al., 2009a]. These variations in rock properties are related to key volcanic facies types (e.g., tabular flows and hyaloclastites) which can be defined in outcrop and where wells have intersected volcanic rocks [e.g., Nelson et al., 2009b; Watton et al., 2014; Nelson et al., 2015]. Furthermore, for the onset of flood volcanism,

Figure 10. (a) Uninterpreted and (b) interpreted seismic sections (in two-way traveltime) highlighting the key facies units of the lava delta system of escarpment segment E5. (c) Schematic illustration of the depth-converted seismic section showing the accommodation space and the position of the paleoshoreline. The paleo-water depth is indicated: ~0.8 km. See Figures 1 and 4c for location. Data courtesy of TGS.

facies mapping of onshore analogs has demonstrated a wide variety of volcanic successions that vary both spatially and temporally [Jerram *et al.*, 2009, 2015a, 2015b]. We therefore have the potential to map out facies, understand their internal properties, and to better define how they might be manifested in the seismic sections.

Good onshore analog examples of lava-fed deltas and thick hyaloclastite sequences have been recognized in outcrops in Greenland [Pedersen *et al.*, 1998; Japsen *et al.*, 2005; Larsen and Pedersen, 2009] (Figure 9a), Iceland [e.g., Watton *et al.*, 2013] (Figure 9b), and Antarctica [Smellie *et al.*, 2011]. Furthermore, such volcanoclastic sequences where lavas interact with wet environments are becoming an increasingly recognized factor at the onset of flood volcanism in many flood basalt provinces and associated volcanic rifted margins [e.g., Ross *et al.*, 2005; Jerram *et al.*, 2015a, 2015b]. In the West Greenland area the lava-fed delta developed in a lacustrine setting [e.g., Japsen *et al.*, 2005; Dam *et al.*, 2009; Abdelmalak, 2010; Abdelmalak *et al.*, 2012a, 2012b], as erupted lava flows dammed indigenous drainage systems and infilled the accommodation space in lake bodies. Large wedge-shaped bodies of hyaloclastite breccia are overlain by thick and extensive horizontal lava flows and interbedded with thin sedimentary units. In Iceland, examples where lava flows enter the sea and emergent volcanic systems have been studied to better understand the facies and development of hyaloclastite sequences [Watton *et al.*, 2013]. Modern examples of lava-fed deltas presently being formed are found in Hawaii, where multiple lava flows enter the sea forming extensive hyaloclastite deposits (Figure 12c). The Hawaii examples have also been sampled by a Hawaiian Scientific Drilling Project well which intersected the subaerial to subaqueous transition which was revealed by facies variations within the hyaloclastites [e.g., Watton *et al.*, 2014].

In the Vøring Margin, the offshore succession of the lava delta system, which records the transition from subaerial to subaqueous environment, was interpreted using the character and geometry of the seismic reflections, combined with onshore exposures from key flood basalt sequences, such as the North Atlantic

Figure 11. (a) Uninterpreted and (b) interpreted seismic sections (in two-way traveltime) showing delta front collapse. The inner flows are characterized by the presence of highs or domes that appear to be the result of gas release in the area. These domes postdate the volcanism since they affect the geometry of younger sediments. (c) Schematic illustration of the depth-converted seismic section. The paleo-water depth is indicated ~ 1.2 km. See Figures 1 and 4c for location. Data courtesy of TGS.

Igneous Province [Planke and Alvestad, 1999; Planke et al., 1999; Jerram, 2002; Jerram and Widdowson, 2005; Nelson et al., 2009b]. Figure 13 shows a 3-D illustration of the schematic outline of the volcanic processes and associated deposits found around the escarpment area, with relation to the seismic volcanic facies seen.

The Landward Flows seismic facies unit is characterized by parallel to subparallel high-amplitude reflections. Individual high-amplitude reflections may be picked over tens of kilometers. The lateral extent of the reflections suggests that the volcanic rocks in these parts of the lava sequence may be of tabular type facies. Landward Flows facies units have been drilled at a number of margins, e.g., Western Australia [Symonds et al., 1998; Rey et al., 2008] and the northeast Atlantic (e.g., ODP Leg 104 Site 642 and ODP Leg 152 Site 917 near the boundary of the inner SDR and Landward Flows). The drilled interval consists of subaerially erupted and emplaced flood basalts, sometimes with thin interbasalt sediment layers. This facies unit may be considered to be akin to the trap like tabular flows [e.g., Jerram et al., 2009]. Because subaerial flood basalts

Figure 12. (a) Onshore analog hyaloclastite breccia in West Greenland (see Figure 1 for location). (b) Example of analog hyaloclastite foresets in Iceland. (c) Modern example of lava entering the sea on Hawaii. Figure 12a, Abdelmalak; Figures 12b and 12c, Jerram.

may flow for several kilometers, volcanoes and fissural flows are most likely the origin of the Landward Flows. Onshore analogs include the Hawaiian lava flows, which are predominantly emplaced as pāhoehoe flows and can travel tens of kilometers from source to shoreline [e.g., Heliker *et al.*, 1998; Heliker and Mattox, 2003].

The Lava Delta facies unit is situated below and basinward of the Landward Flows facies unit. In seismic profiles, this facies unit shows a prograding reflection configuration. The Lava Delta is interpreted to be

Figure 13. Schematic outline of the volcanic processes and associated deposits found around the escarpment area. The respective seismic volcanic facies (e.g., Inner Flows and Lava Delta) are also labeled. The sequence is characterized by a mixture of subaerial and subaqueous eruptions and transitions from subaerial to subaqueous environment. Lava Delta prograde out from the paleoshoreline and deposit reworked volcanic debris/turbidite flows into the basin.

deposited in a subaqueous predominantly coastal depositional environment. In places the Landward Flows unit merges with the Lava Delta facies unit and forms basinward dipping reflections. This facies is interpreted to be composed of hyaloclastite fans or aprons that record the flow of lava directly into the offshore basin. The magma supply and the progradation of the delta foreset into the basin will favor sliding and slumping of the unconsolidated hyaloclastite along the escarpment.

The Inner Flows seismic unit is identified landward and sometimes below the Lava Delta and the Landward Flows. The Inner Flows unit is a sheet-like body of highly disrupted showing hummocky reflections with a chaotic internal reflection configuration. The Inner Flows are deposited in a broad basin in a shallow marine depositional environment. This seismic unit consists of a mixture of massive and fragmented basalt, volcanoclastic rocks, pillow lavas, and shallow intrusions. The main sources of the Inner Flows are fissures along the region of the continental breakup, as this unit is interpreted to have formed simultaneously with Landward Flows and Lava Delta units.

5.2. Buildup of the Lava Delta System

The reconstruction of the emplacement history of continental flood basalt around Vøring Escarpment is based on the key volcanic facies and seismic reflection relationships in the Vøring Margin (this study) and from a worldwide integration of lava delta system interpretation. The seismic reflection configuration appears to have been deposited sequentially with a stacking pattern revealing a variation in the accommodation space relative to sea level change, volcanic material supply, and tectonic movements. As in conventional delta systems, the height of the delta front may be a proxy for water depth at the time of delta deposition [e.g., *Kjørboe, 1999; Spitzer et al., 2008*]. The dominant control is thought to be relative sea level change. However, in the case of high magma supply, the Lava Delta is capable of continuous progradation despite fluctuation in sea level and accommodation.

Figures 14a–14d show a schematic representation of the lava delta system development relative to sea level and magma supply. This model can be used to explain the development of the “step-back deltas.” The initial stacking pattern is progradational into the basin during the initial high stand time (highstand system tract or HST). Deposition was likely controlled by the volumes of erupted lava entering the basin and filling the accommodation space (Figure 14a). During volcanic activity, the lava flows prograde into the basin and define the extent of progradation of the lava-fed delta. If the supply is high enough, progradation will occur throughout the entire sea level cycle (Figure 11b). The deposition of volcanic rocks can also affect accommodation, with an increase in subsidence causing aggradation. Coastal erosion and slumping are enhanced during the period of falling sea level (lowstand system tract or LST) (Figure 14c). The unconsolidated delta front subsides causing fractures to propagate through the delta front. Such fractures will induce large-scale delta collapse, sliding/slumping and gravity-driven debris avalanches into the basin (Figure 14c). The stacking pattern changes to retrogradation during a sea level rise (transgressive systems tract or TST) and/or decreased magma supply. Any relative rise in sea level floods the lava flows and moves the shoreline significantly. These different factors will favor the development of a step-back delta (Figure 14d), where the next lava sequences prograde out and fill the newly developed accommodation space [e.g., *Wright et al., 2012*]. During this period subaqueous erosion of the lava delta system is enhanced.

Structural highs could considerably affect the lava delta system buildup (Figure 14e). Generally, the Landward Flows and Lava Delta units are thinner, since there is less accommodation space above these highs, which sometime can present pronounced topographic features. Magma supply will drape the highs and progradation of the lava flows above the structural high starts (Figure 14e). Slumping and sliding are common along the flank of the structural high and could be related to the progradation of the lava flows and/or in response to vertical movement. Strike-slip movements and vertical movements along fracture zones will result in a second-order segmentation of the escarpment into several pseudoescarpments with different lava delta systems (Figure 14f).

5.3. Escarpments Along the North Atlantic Margin

The breakup volcanic sequences in the mid-Norwegian Margin include two prominent escarpments: the Vøring Escarpment on the Vøring Margin and the Faeroe-Shetland Escarpment on the Møre and United Kingdom margins (Figure 15a). The Vøring Escarpment presents a ~350 km prominent feature along the

Figure 14. Idealized development of a lava delta system in response to different controlling factors. Figures 14a–14d show development in response to relative sea level and/or magma supply. (a and b) Progradation of the volcanic sequences occurs rapidly during magmatic events and fills the accommodation space (highstand system tract or HST). (c) During periods of low volcanic output and where the relative sea level falls (lowstand system tract or LST), slumping is enhanced. (d) Where relative sea level rises (transgressive systems tract or TST), step-back deltas can occur. (e) Structural high effect. (f) Tectonic effect of fracture zone on the lava delta system.

Vøring Margin and separates the Vøring Marginal High to the west from the Vøring Basin to the east. The Faeroe-Shetland Escarpment extends from the region east of the Faeroe Islands to the Jan Mayen Fracture Zone between 61°20'N–65°N and 4°W–3°15'E and bounds the Møre Marginal High to the west from the Møre Basin to the east. Good examples of the development of the lava delta system along the Faeroe-Shetland Escarpment have been reported by Wright *et al.* [2012] and can be seen to show similar internal characteristics to those reported in this study. Both escarpments are situated at the boundary between Inner Flows and Landward Flows and are absent elsewhere (Figure 15a). North of the intersection with the Bivrost Lineament, the Vøring Escarpment does not continue into the Lofoten Margin. There the escarpment seems to represent a volcanic buildup or simply a flow front. From seismic data on the NE Greenland conjugate margin, we identified similar escarpment (the Thetis Escarpment) facing to the west (Figure 15b). The NE Greenland escarpment presents a limited extent compared to the Vøring and Faeroe-Shetland examples. Future detailed mapping of the Greenland Margin volcanic stratigraphy may improve the characterization of this escarpment.

5.4. Escarpment Along the Vøring Margin

The nature and origin of the Vøring Escarpment has been debated for decades. Talwani and Eldholm [1972] described it as a first-order geological feature related to the continent ocean boundary. Furthermore, they

Figure 15. (a) Offshore distribution of the volcanic facies seismic units in the mid-Norwegian Margin updated from *Berndt et al.* [2001b]. The Vøring Escarpment and the Faeroe-Shetland Escarpment are indicated in the map. (b) Offshore distribution of the volcanic facies seismic units in the NE Greenland Margin. The Thetis Escarpment is indicated on the map. See Figure 1 for location.

noted that the escarpment marks significant changes in a number of geological and geophysical parameters. In particular, the magnetic field becomes quiet on the landward side, gravity highs are located on the seaward side, and there is a pronounced difference in the velocity-depth functions on either side of the escarpment. Although the detailed expression of the escarpment varies locally, *Hagevang et al.* [1983] demonstrated that it is associated with a continuous sharp negative magnetic anomaly on the Vøring Marginal High. There was also little agreement about the relative movement along the escarpment. *Caston* [1976] proposed that it has acted as a normal fault throughout the Tertiary. *Skogseid and Eldholm* [1987] suggested that the escarpment may represent a reactivated Late Jurassic-Early Cretaceous fault. *Blystad et al.* [1995] suggested that the Vøring Escarpment may have developed as a fault scarp in the Maastrichtian-Paleocene rifting episodes and was subsequently covered by flood basalts during the Early Eocene breakup. Similar to the Vøring Escarpment, the structural expression of the Faeroe-Shetland escarpment varies along strike. To the south, it locally reveals a lobate edge, apparently defining the front of the volcanic flows [e.g., *Kjørboe*, 1999; *Wright et al.*, 2012]. North of about 63°N, along the Møre Margin, the escarpment is well expressed, fairly linear, and may reflect an interrelationship between a volcanic edge and a deeper structural setting [e.g., *Blystad et al.*, 1995].

Table 2. Summary of the Different Controlling Factors of the Vøring Escarpment^a

Control factors	Lava delta system morphology	LD extent	Escarpment segment	Escarpment height	LD thickness	LF thickness
Relative sea level Lava supply		Well extended	E2	800 to 1600 m Average~ 1300 m	700 m to 1000 m	500 m to 1500 m
			E4	800 to 1600 m Average~ 1200 m	700 m to 1500 m	500 m to 2000 m
High		Limited extent	E3	500 to 800 m Average~ 600 m	300 m to 500 m	Less than 500 m
			E5	200 to 500 m Average~ 450 m	Less than 500 m	Less than 500 m
JMFBZ		Limited extent	E1	200 to 500 m Average~ 400 m	Less than 500 m	Less than 500 m

^aRelated segment characteristic and morphology is indicated for each controlling factor. LD: Lava Delta; LF: Landward Flows. LD: Lava Delta; LF: Landward Flows.

Smythe *et al.* [1983] proposed that the Faeroe-Shetland Escarpment was a flow edge separating subaerially erupted basalt on top of the Mesozoic sediments from a restricted shallow shelf, thus defining a paleoshoreline. Berndt *et al.* [2001b] suggested that the continuity and undulation of both the Faeroe-Shetland Escarpment and the Vøring Escarpment make it very unlikely that the escarpments are dominantly fault controlled as suggested by Skogseid and Eldholm [1989] and Brekke *et al.* [1999]. The fact that the escarpments exist exclusively between the Landward Flows and Inner Flows suggests that they are rather the result of a generic margin forming process as suggested by Smythe *et al.* [1983] for the Faeroe-Shetland Escarpment. Planke and Alvestad [1999] interpreted the prograding reflections underneath the Landward Flows to result from the foresets of a Gilbert-type Lava Delta terminating with the escarpment. Such a setting is similar to those of Lava Deltas and bench collapse at the Kilauea Volcano, Hawaii [Moore *et al.*, 1973]. An origin of this type implies that the escarpments develop during or after the emplacement of the Landward Flows and submergence of the marginal high to below sea level. Detailed mapping of the southern part of the Faeroe-Shetland Escarpment has led to similar interpretation for that escarpment [Kjørboe, 1999; Wright *et al.*, 2012].

In this study, the structural expression of the Vøring Escarpment is shown to vary along strike (Figures 3 and 4) with five segments defined showing different morphologies and different heights (Table 2). This first-order segmentation is due to the different factors that control the Vøring Escarpment buildup. Segments 2 and 4, which display the higher escarpment topography, are characterized by a well-developed lava delta system with thicker Lava Delta and Landward Flows compared to the other segments. Relative sea level change and the magma supply are the main factors that control the establishment of these two segments. During the rise of sea level and/or decreased magma supply a step-back delta can occur. The lava delta system of the segments E3 and E5 and part of the segment E1 developed above Skoll High, Grimm High, and Ygg High, respectively. Generally, the lava delta system shows a limited extent with thinner Lava Delta and Landward Flows thicknesses compared with the E2 and E4 segments. These structural highs seem to control the morphology of these escarpment segments. In segment E3, sliding and slumping indicates that this structure was tectonically active during the lava progradation. Finally, the escarpment segment E1 is characterized by a second-order segmentation, mainly due to the tectonomagmatic control of the Jan Mayen Fracture Zone [e.g., Berndt *et al.*, 2001a]. Multiple pseudoescarpments and lava delta systems with limited extent and thin flows characterize this segment. Despite these different control factors, the Vøring Escarpment strikes NE-SW parallel to the breakup axis and to the Late Cretaceous- Early Paleocene Gjallar Ridge and faults. The outline of the escarpment structure is clearly seen in the potential field data (Figures 3a and 3b). The escarpment marks significant changes in the magnetic field which becomes quiet on the landward side. On the filtered Bouguer anomaly map, the different high-amplitude and low-amplitude anomalies along the Vøring Escarpment correlate with the segments identified from the seismic data.

5.5. Subbasalt Structure

It is well known that lava delta systems are constructed by explosive and nonexplosive interaction of magma and water by basalt quenching and fragmentation when subaerially erupted lavas flow into the sea [e.g., Planke *et al.*, 2000; Watton *et al.*, 2013]. This process has been observed in detail in the Kilauea volcano, Hawaii, where subaerial lavas were emplaced on top of shallow marine, foreset-bedded hyaloclastites [e.g., Moore *et al.*, 1973; Watton *et al.*, 2014]. Lava Deltas can commonly develop in emergent volcanic systems and where local or regionally elevated areas funnel lava flows toward coastlines [e.g., Watton *et al.*, 2013], and by analogy, large parts of the Vøring Margin were elevated or emerging when the volcanics were emplaced. Nevertheless, on the Vøring Marginal High, the nature of the crust below extrusive rocks remains unclear since the seismic signal is often obscured by the volcanic cover. Several strong reflections are identified and interpreted as a sill complex beneath the lava flows. Such sill intrusions are typically characterized by high-amplitude, abrupt termination and sometimes saucer-shaped geometries [e.g., Planke *et al.*, 2005, 2015]. This observation implies the presence and preservation of subbasalt basins in the Vøring Marginal High [Abdelmalak *et al.*, 2015]. This interpretation is in good accordance with the resistivity structure within the Vøring Marginal High, showing low resistivity below the volcanic unit and suggesting the presence of kilometer-scale thick subbasalt sedimentary sequences [Planke *et al.*, 2015]. Basalt thickness mapping illustrates separate depocenters in the SDR that could indicate complicated rift geometry in the outer Vøring Margin and significant local variations in the accommodation space possibly controlled by tectonism during SDR growth [see Abdelmalak *et al.*, 2016].

ODP Hole 642E, located near the apex of the SDR on the Vøring Marginal High, is situated ~25 km west of the Vøring Escarpment. The volcanic units sampled at Hole 642E consist of peraluminous, cordierite bearing peperitic basaltic andesitic to dacitic flows interbedded with thick volcanosedimentary deposits [Meyer *et al.*, 2009; Abdelmalak *et al.*, 2016]. The peraluminous geochemistry combined with available C, Sr, Nd, and Pb isotopes data point toward an upper crustal anatectic source of these magmas with a significant contribution of organic carbon-rich pelagic sedimentary material [Abdelmalak *et al.*, 2016]. As drilling stopped within the basaltic andesitic flow unit, little is known about the geology prior to the eruption of these crustal anatectic melts.

On the outer Vøring Basin, the Late Cretaceous-Paleocene topography was characterized by low-angle normal faulting, subsidence, synrift sedimentation, and local uplift [Gernigon *et al.*, 2003; Ren *et al.*, 2003]. From the Upper Cretaceous to the upper Paleocene (Danian to Early Thanetian: 65–59 Ma) a regional uplift event is recorded both in the Vøring Basin and the entire NE Atlantic [Brekke *et al.*, 2001]. This uplift has been correlated with upwelling of hot mantle that was responsible for a dramatic shallowing of the outer Vøring Basin and the emergence and local erosion of the preexisting Cretaceous highs (e.g., Gjallar Ridge) [Brekke *et al.*, 2001; Ren *et al.*, 2003]. Vertical movement could also have affected the escarpment height and led to a variation in the paleo-water depth and depocenter locations during the breakup magmatism event. However, no biostratigraphic data are available that could confirm the paleo-water depth in the Outer Vøring Margin since no wells have been drilled in the area.

The identification in this study of subbasins on the Vøring Marginal High that may contain pelagic sedimentary rocks rich in organic carbon [Abdelmalak *et al.*, 2016] could enhance the area for petroleum exploration. Gas seepage/release along the Fenris Graben could (Figures 1 and 11) also indicate the hydrocarbon potential of Vøring Marginal High. Such gas releases in the area are probably caused by the increased maturation of organic matter as a result of extensive sill intrusion in the sedimentary basin below the volcanic rocks.

6. Conclusions

The increased amount of new and reprocessed high-quality seismic data permit a more accurate characterization of the along-strike and across-strike continuity and the variability of the different volcanic seismic facies units along the Vøring Margin. Using good quality imaging, we have mapped in detail the ~350 km long NE-SW trending Vøring Escarpment which is a prominent structure along the Vøring Margin with a height ranging between 200 and 1600 m. This detailed analysis has resulted in the following conclusions:

1. The Vøring Escarpment is divided along strike into five main segments. This segmentation is due to three main control factors: the sea level change/magma supply, structural highs, and the JMFZ.
2. The well-developed Vøring Escarpment morphology along the segments E2 and E4 was mainly controlled by relative sea level change and magma supply during the lava delta progradation and filling of the basin.
3. In segments E3 and E5, the escarpment morphology is a result of a combination of the front of a volcanic buildup and a partly faulted boundary of differential subsidence between the Cretaceous/Tertiary structural highs and the basin area to the east.
4. Segment E1 is characterized by pseudoescarpments, mainly identified around the Jan Mayen Fracture Zone which is the main controlling factor of this segment. Vertical movements along this fracture zone could have played an important role in forming the escarpment morphology in this area.
5. The recognition of the interplay between lava supply, relative sea level, the development of key volcanic facies, and their correct identification in offshore sequences provides insights into the breakup process at volcanic rifted margins. The continuing development and movement of the paleoshoreline during the buildup of volcanism and its dynamic interplay along large segments of the margin (as has been shown here) has profound implications for how basins and depocenters develop during the onset and main rifting phases along such continental margins.
6. Integrated seismic, gravity, and magnetic interpretation indicate complicated rift geometry in the outer Vøring Margin with the existence of several subbasins and structural highs. The identification of sedimentary rocks rich in organic carbon in the outer Vøring Margin as well as gas seepages along the Vøring Escarpment could enhance the area for petroleum exploration.

Acknowledgments

Funding for this work came from OMNIS Project: Offshore Mid-Norwegian: Integrated Margin and Basin Studies, (project 210429/E30) funded by the Norwegian Research Council. The seismic, magnetic, and gravity data presented in this study were provided by TGS. We would like to acknowledge First Geo AS and TGS for the use of Hqbe™ regional velocity cube employed in depth conversion of the seismic profiles. Dougal Jerram and Sverre Planke are supported by the Research Council of Norway through its Centre of Excellence funding scheme, project 223272 (CEED). Seismic interpretation was done using IHS: Kingdom suite software. Grid interpolations and map compilations were established using Geosoft Oasis Montaj and ArcGis softwares. We thank Nick Schofield, Christian Berndt, John R. Hopper, the Editor Uri ten Brink, and an anonymous Associated Editor for useful comments that improved the paper. We thank Adrian Read for reading and revising our corrected manuscript.

References

- Abdelmalak, M. M. (2010), Transition spatio-temporelle entre rift sédimentaire et marge passive volcanique: L'exemple de la Baie de Baffin, Centre Ouest Groenland, (Spatio-temporal transition between a sedimentary basin to a volcanic passive margin: The Baffin Bay case example, Central West Greenland), p. 266, Le Mans. [Available at Université du Maine, France] online: <http://cyberdoc.univ-lemans.fr/theses/2010/2010LEMA1030.pdf>]
- Abdelmalak, M. M., C. Aubourg, L. Geoffroy, and F. Laggoun-Deffarge (2012a), A new oil-window indicator? The magnetic assemblage of claystones from the Baffin Bay volcanic margin, *AAPG Bull.*, 96(2), 205–215.
- Abdelmalak, M. M., L. Geoffroy, J. Angelier, B. Bonin, J. P. Callot, J. P. Gélard, and C. Aubourg (2012b), Stress fields acting during lithosphere breakup above a melting mantle: A case example in West Greenland, *Tectonophysics*, 581, 132–143, doi:10.1016/j.tecto.2011.11.020.
- Abdelmalak, M. M., T. B. Andersen, S. Planke, J. I. Faleide, F. Corfu, C. Tegner, G. E. Shephard, D. Zastrozhnov, and R. Myklebust (2015), The ocean-continent transition in the mid-Norwegian Margin: Insight from seismic data and an onshore Caledonian field analogue, *Geology*, 43, 1011–1014, doi:10.1130/G37086.1.
- Abdelmalak, M. M., et al. (2016), Pre-breakup magmatism on the Vøring Margin: Insight from new sub-basalt imaging and results from Ocean Drilling Program Hole 642E, *Tectonophysics*, 675, 258–274, doi:10.1016/j.tecto.2016.02.037.
- Berndt, C. (2002), Residual Bouguer satellite gravity anomalies reveal basement grain and structural elements of the Vøring Margin, off Norway, *Nor. J. Geol.*, 82, 31–36.
- Berndt, C., R. Mjelde, S. Planke, H. Shimamura, and J. I. Faleide (2001a), Controls on the tectono-magmatic evolution of a volcanic transform margin: The Vøring Transform Margin, NE-Atlantic, *Mar. Geophys. Res.*, 22(3), 133–152.
- Berndt, C., S. Planke, E. Alvestad, F. Tsikalas, and T. Rasmussen (2001b), Seismic volcanostratigraphy of the Norwegian Margin: Constraints on tectonomagmatic break-up processes, *J. Geol. Soc.*, 158(3), 413–426, doi:10.1144/jgs.158.3.413.
- Blystad, P., H. Brekke, R. B. Færseth, B. T. Larsen, J. Skogseid, and B. Tørudbakken (1995), Structural elements of the Norwegian continental shelf Part II: The Norwegian Sea Region, *NPD-Bulletin, The Norwegian Petroleum Directorate.*, 8.
- Breivik, A., J. I. Faleide, R. Mjelde, E. Flueh, and Y. Murai (2014), Magmatic development of the outer Vøring margin from seismic data, *J. Geophys. Res. Solid Earth*, 119, 6733–6755, doi:10.1002/2014JB011040.
- Brekke, H., S. Dahlgren, B. Nyland, and C. Magnus (1999), The prospectivity of the Vøring and Møre basins on the Norwegian Sea continental margin, *Geol. Soc. Lond. Petrol. Geol. Conf. Ser.*, 5, 261–274, doi:10.1144/0050261.
- Brekke, H., H. I. Sjulstad, C. Magnus, and R. W. Williams (2001), Sedimentary environments offshore Norway—An overview, in *Norwegian Petroleum Society Special Publications*, edited by J. M. Ole and D. Tom, pp. 7–37, Elsevier, Amsterdam, doi:10.1016/S0928-8937(01)80006-0.
- Caston, V. N. D. (1976), Tertiary sediments of the Vøring Plateau, Norwegian Sea, recovered by Leg 38 Of the Deep Sea Drilling Project, in *Init. Rep. DSDP*, 38, edited by M. Talwani et al., pp. 761–782, U.S. Govt. Print. Off., Washington.
- Dam, G., G. K. Pedersen, M. Sonderholm, H. Midtgaard, L. M. Larsen, H. Nohr-Hansen, and A. K. Pedersen (2009), Lithostratigraphy of the Cretaceous-Pleocene Nuussuaq Group, Nuussuaq Basin, West Greenland, *Geol. Surv. Denmark Greenland Bull.*, 19, 171.
- Eldholm, O., J. Thiede, E. Taylor, and S. S. Party (1987), Summary and preliminary conclusions, ODP Leg 104, in *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by O. Eldholm et al., pp. 751–771, Ocean Drilling Program, College Station, Tex.
- Eldholm, O., J. Thiede, and E. Taylor (1989), Evolution of the Vøring Volcanic Margin, in *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by O. Eldholm, J. Thiede, and E. Taylor, pp. 1033–1065, Ocean Drilling Program, College Station, Tex.
- Eldholm, O., F. Tsikalas, and J. I. Faleide (2002), Continental margin off Norway 62–75°N: Palaeogene tectono-magmatic segmentation and sedimentation, *Geol. Soc. London Spec. Publ.*, 197(1), 39–68, doi:10.1144/gsl.sp.2002.197.01.03.
- Faleide, J. I., F. Tsikalas, A. J. Breivik, R. Mjelde, O. Ritzmann, Ø. Engen, J. Wilson, and O. Eldholm (2008), Structure and evolution of the continental margin off Norway and the Barents Sea, *Episodes*, 31(1), 82–91.
- First Geo AS and TGS (2015), Regional high quality velocity Cube for mid Norway, Norwegian sector *Rep.*, 25 pp.

- Gaina, C., L. Gernigon, and P. Ball (2009), Palaeocene-recent plate boundaries in NE Atlantic and formation of the Jan Mayen microcontinent, *J. Geol. Soc. London*, *166*, 601–616.
- Gallagher, J. W., and P. W. Dromgoole (2007), Exploring below the basalt, offshore Faroes: A case history of sub-basalt imaging, *Petrol. Geosci.*, *13*(3), 213–225, doi:10.1144/1354-079306-711.
- Gernigon, L., J.-C. Ringenbach, S. Planke, and H. Jonquet-Kolsto (2003), Extension, crustal structure and magmatism at the outer Voring Basin, Norwegian Margin, *J. Geol. Soc. London*, *160*, 197–208.
- Gernigon, L., J.-C. Ringenbach, S. Planke, and B. Le Gall (2004), Deep structures and breakup along volcanic rifted margins: Insights from integrated studies along the outer Voring Basin (Norway), *Mar. Pet. Geol.*, *21*(3), 363–372.
- Hagevang, T., O. Eldholm, and I. Aalstad (1983), Pre-23 magnetic anomalies between Jan Mayen and Greenland-Senja fracture zones in the Norwegian sea, *Mar. Geophys. Res.*, *5*(4), 345–363, doi:10.1007/BF00305388.
- Hansen, J. P. V., J. A. Cartwright, M. Huuse, and O. R. Clausen (2005), 3D seismic expression of fluid migration and mud remobilization on the Gjallar Ridge, offshore mid-Norway, *Basin Res.*, *17*(1), 123–139, doi:10.1111/j.1365-2117.2005.00257.x.
- Heliker, C., and T. N. Mattox (2003), The first two decades of the Pu' u 'Ō'ō-Kūpaianaha eruption: Chronology and selected bibliography, *U.S. Geol. Surv. Prof. Pap.*, 1–28.
- Heliker, C. C., M. T. Mangan, T. N. Mattox, J. P. Kauahikaua, and R. T. Helz (1998), The character of long-term eruptions: Inferences from episodes 50–53 of the Pu' u 'Ō'ō-Kūpaianaha eruption of Kilauea Volcano, *Bull. Volcanol.*, *59*(6), 381–393, doi:10.1007/s004450050198.
- Japsen, P., P. F. Green, and J. A. Chalmers (2005), Separation of Palaeogene and Neogene uplift on Nuussuaq, West Greenland, *J. Geol. Soc. London*, *162*, 299–314.
- Jerram, D. A. (2002), Volcanology and facies architecture of flood basalts, *Geol. Soc. Am. Spec. Pap.*, *362*, 119–132, doi:10.1130/0-8137-2362-0.119.
- Jerram, D. A. (2015), Hot rocks and oil: Are volcanic margins the new frontier?, *Exploration & Production*.
- Jerram, D. A., and M. Widdowson (2005), The anatomy of continental flood basalt provinces: Geological constraints on the processes and products of flood volcanism, *Lithos*, *79*(3–4), 385–405, doi:10.1016/j.lithos.2004.09.009.
- Jerram, D. A., R. T. Single, R. W. Hobbs, and C. E. Nelson (2009), Understanding the offshore flood basalt sequence using onshore volcanic facies analogues: An example from the Faroe-Shetland basin, *Geol. Mag.*, *146*(3), 353–367, doi:10.1017/s0016756809005974.
- Jerram, D. A., H. H. Svensen, S. Planke, A. G. Polozov, and T. H. Torsvik (2015a), The onset of flood volcanism in the north-western part of the Siberian Traps: Explosive volcanism versus effusive lava flows, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, doi:10.1016/j.palaeo.2015.04.022.
- Jerram, D. A., M. Widdowson, P. B. Wignall, Y. Sun, X. Lai, D. P. G. Bond, and T. H. Torsvik (2015b), Submarine palaeoenvironments during Emeishan flood basalt volcanism, SW China: Implications for plume–lithosphere interaction during the Capitanian, Middle Permian ('end Guadalupian') extinction event, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, doi:10.1016/j.palaeo.2015.06.009.
- Kjørboe, L. (1999), Stratigraphic relationships of the Lower Tertiary of the Faeroe Basalt Plateau and the Faeroe-Shetland Basin, *Geol. Soc. London Petrol. Geol. Conf. Ser.*, *5*, 559–571, doi:10.1144/00505559.
- Larsen, H. C., and A. D. Saunders (1998), Tectonism and volcanism at the southeast Greenland rifted margin: A record of plume impact and later continental rupture, in *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by A. D. Saunders, H. C. Larsen, and S. W. J. Wise, pp. 503–533, Ocean Drilling Program, College Station, Tex.
- Larsen, L. M., and A. K. Pedersen (2009), Petrology of the Paleocene picrites and flood basalts on Disko and Nuussuaq, West Greenland, *J. Petrol.*, *50*(9), 1667–1711.
- Larsen, L. M., R. Waagstein, A. K. Pedersen, and M. Storey (1999), Trans-Atlantic correlation of the Palaeogene volcanic successions in the Faeroe Islands and East Greenland, *J. Geol. Soc.*, *156*(6), 1081–1095, doi:10.1144/gsjgs.156.6.1081.
- Maus, S., et al. (2009), EMAG2: A 2-arc min resolution Earth Magnetic Anomaly Grid compiled from satellite, airborne, and marine magnetic measurements, *Geochem. Geophys. Geosyst.*, *10*, Q08005, doi:10.1029/2009GC002471.
- Meyer, R., J. Hertogen, R. B. Pedersen, L. Viereck-Götte, and M. Abratis (2009), Interaction of mantle derived melts with crust during the emplacement of the Voring Plateau, N.E. Atlantic, *Mar. Geol.*, *261*(1–4), 3–16, doi:10.1016/j.margeo.2009.02.007.
- Moore, J. G., R. L. Phillips, R. W. Grigg, D. W. Peterson, and D. A. Swanson (1973), Flow of lava into the sea, 1969–1971, Kilauea Volcano, Hawaii, *Geol. Soc. Am. Bull.*, *84*(2), 537–546, doi:10.1130/0016-7606(1973)84<537:folits>2.0.co;2.
- Nelson, C. E., D. A. Jerram, and R. W. Hobbs (2009a), Flood basalt facies from borehole data: Implications for prospectivity and volcanology in volcanic rifted margins, *Petrol. Geosci.*, *15*(4), 313–324, doi:10.1144/1354-079309-842.
- Nelson, C. E., D. A. Jerram, R. T. Single, and R. W. Hobbs (2009b), Understanding the facies architecture of flood basalts and volcanic rifted margins and its effect on geophysical properties, in *Faroe Islands Exploration Conference: Proceedings of the 2nd Conference, Annales Societatis Scientiarum Faroensis, Supplementum*, edited by T. Varming and H. Ziska, pp. 83–103.
- Nelson, C. E., D. A. Jerram, J. A. P. Clayburn, A. M. Halton, and J. Roberge (2015), Eocene volcanism in offshore southern Baffin Bay, *Mar. Pet. Geol.*, *67*, 678–691, doi:10.1016/j.marpetgeo.2015.06.002.
- Pedersen, G. K., L. M. Larsen, A. K. Pedersen, and B. F. Hjortkjaer (1998), The syn-volcanic Naajaat lake, Paleocene of West Greenland, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, *140*, 271–287.
- Planke, S. (1994), Geophysical response of flood basalts from analysis of wire line logs: Ocean Drilling Program Site 642, Voring volcanic margin, *J. Geophys. Res.*, *99*, 9279–9296, doi:10.1029/94JB00496.
- Planke, S., and E. Alvestad (1999), Seismic volcanostratigraphy of the extrusive breakup complexes in the northeast Atlantic: Implications from ODP/DSDP drilling, in *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by H. C. Larsen et al., pp. 3–16, Ocean Drilling Program, College Station, Tex.
- Planke, S., and H. Cambray (1998), Seismic properties of flood basalt from the 917A downhole data, Southeast Greenland volcanic margin, in *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by A. D. Saunders, H. C. Larsen, and S. W. Wise Jr., pp. 453–462, Ocean Drilling Program, College Station, Tex.
- Planke, S., and O. Eldholm (1994), Seismic response and construction of seaward dipping wedges of flood basalts: Voring volcanic margin, *J. Geophys. Res.*, *99*(B5), 9263–9278, doi:10.1029/94JB00468.
- Planke, S., E. Alvestad, and O. Eldholm (1999), Seismic characteristics of basaltic extrusive and intrusive rocks, *Leading Edge*, *18*, 342–348.
- Planke, S., P. A. Symonds, E. Avelstad, and J. Skogseid (2000), Seismic volcanostratigraphy of large-volume basaltic extrusive complexes on rifted margins, *J. Geophys. Res.*, *105*, 19,333–19,351, doi:10.1029/1999JB900005.
- Planke, S., T. Rasmussen, T. Rey, and R. Myklebust (2005), Seismic characteristics and distribution of volcanic intrusions and hydrothermal vent complexes in the Voring and More basins, in *Petroleum Geology: North-West Europe and Global Perspectives—Proceedings of the 6th Petroleum Geology Conference*, edited by A. G. Doré and B. A. Vining, pp. 833–844, Geol. Soc., London.
- Planke, S., H. Svensen, R. Myklebust, S. Bannister, B. Manton, and L. Lorenz (2015), *Geophysics and Remote Sensing*, pp. 1–16, Springer, Berlin Heidelberg, doi:10.1007/11157_2014_6.

- Ren, S., J. I. Faleide, O. Eldholm, J. Skogseid, and F. Gradstein (2003), Late Cretaceous–Paleocene tectonic development of the NW Vøring Basin, *Mar. Pet. Geol.*, *20*(2), 177–206, doi:10.1016/S0264-8172(03)00005-9.
- Rey, S. S., S. Planke, P. A. Symonds, and J. I. Faleide (2008), Seismic volcanostratigraphy of the Gascoyne margin, Western Australia, *J. Volcanol. Geotherm. Res.*, *172*(1–2), 112–131, doi:10.1016/j.jvolgeores.2006.11.013.
- Rodriguez Monreal, F., H. J. Villar, R. Baudino, D. Delpino, and S. Zencich (2009), Modeling an atypical petroleum system: A case study of hydrocarbon generation, migration and accumulation related to igneous intrusions in the Neuquen Basin, Argentina, *Mar. Pet. Geol.*, *26*(4), 590–605, doi:10.1016/j.marpetgeo.2009.01.005.
- Ross, P. S., I. Uktins Peate, M. K. McClintock, Y. G. Xu, I. P. Skilling, J. D. L. White, and B. F. Houghton (2005), Mafic volcanoclastic deposits in flood basalt provinces: A review, *J. Volcanol. Geotherm. Res.*, *145*(3–4), 281–314, doi:10.1016/j.jvolgeores.2005.02.003.
- Sandwell, D. T., and W. H. F. Smith (2009), Global marine gravity from retracked Geosat and ERS-1 altimetry: Ridge segmentation versus spreading rate, *J. Geophys. Res.*, *114*, B01411, doi:10.1029/2008JB006008.
- Scheck-Wenderoth, M., T. Raum, J. I. Faleide, R. Mjelde, and B. Horsfield (2007), The transition from the continent to the ocean: A deeper view on the Norwegian Margin, *J. Geol. Soc.*, *164*(4), 855–868, doi:10.1144/0016-76492006-131.
- Schofield, N., and D. W. Jolley (2013), Development of intra-basaltic lava-field drainage systems within the Faroe-Shetland Basin, *Petrol. Geosci.*, *19*(3), 273–288, doi:10.1144/petgeo2012-061.
- Schofield, N., et al. (2015a), Regional magma plumbing and emplacement mechanisms of the Faroe-Shetland Sill Complex: Implications for magma transport and petroleum systems within sedimentary basins, *Basin Res.*, doi:10.1111/bre.12164.
- Schofield, N., D. A. Jerram, S. Holford, N. Mark, A. Hartley, J. Howell, D. Muirhead, P. Green, D. Hutton, and C. Stevenson (2015b), Sills in sedimentary basins and petroleum systems, in *Advances in Volcanology*, pp. 1–22, Springer, Berlin, Heidelberg, doi:10.1007/11157_2015_17.
- Skogseid, J., and O. Eldholm (1987), Early Cenozoic crust at the Norwegian continental margin and the conjugate Jan Mayen Ridge, *J. Geophys. Res.*, *92*(B11), 11,471–11,491, doi:10.1029/JB092iB11p11471.
- Skogseid, J., and O. Eldholm (1989), Vøring plateau continental margin: Seismic interpretation stratigraphy, and vertical movements, in *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by O. Eldholm, J. Thiede, and E. Taylor, pp. 993–1030, Ocean Drilling Program, College Station.
- Skogseid, J., S. Planke, J. I. Faleide, T. Pedersen, O. Eldholm, and F. Neverdal (2000), NE Atlantic continental rifting and volcanic margin formation, in *Dynamics of the Norwegian Margin*, edited by A. Nottvedt et al., pp. 295–326, Geol. Soc. London Spec. Publ., London.
- Smellie, J. L., S. Rocchi, and P. Armienti (2011), Late Miocene volcanic sequences in northern Victoria Land, Antarctica: Products of glaciovolcanic eruptions under different thermal regimes, *Bull. Volcanol.*, *73*(1), 1–25, doi:10.1007/s00445-010-0399-y.
- Smythe, D. K., J. A. Chalmers, A. G. Skucet, A. Dobinson, and A. S. Mould (1983), Early opening history of the North Atlantic—I. Structure and origin of the Faeroe-Shetland Escarpment, *Geophys. J. R. Astron. Soc.*, *72*(2), 373–398, doi:10.1111/j.1365-246X.1983.tb03791.x.
- Soubaras, R., and R. Dowle (2010), Variable-depth streamer—A broadband marine solution, *First Break*, *28*, 89–96.
- Spitzer, R., R. S. White, P. A. F. Christie, and iSSIMM Team (2008), Seismic characterization of basalt flows from the Faroes margin and the Faroe-Shetland basin, *Geophys. Prospect.*, *56*(1), 21–31, doi:10.1111/j.1365-2478.2007.00666.x.
- Svensen, H., S. Planke, A. Mørche-Sørensen, B. Jamtveit, R. Myklebust, T. Rasmussen Eidem, and S. S. Rey (2004), Release of methane from a volcanic basin as a mechanism for initial Eocene global warming, *Nature*, *429*(6991), 542–545.
- Symonds, P. A., S. Planke, Ø. Frey, and J. Skogseid (1998), Volcanic development of the Western Australian continental margin and its implications for basin development, in *The Sedimentary Basins of Western Australia 2: Proc. of Pet. Expl. Soc. of Australia Symp., Perth*, edited by P. G. Purcell and R. R. Purcell, pp. 33–54.
- Talwani, M., and O. Eldholm (1972), Continental margin off Norway: A geophysical study, *Geol. Soc. Am. Bull.*, *83*(12), 3575–3606, doi:10.1130/0016-7606(1972)83[3575:cmonag]2.0.co;2.
- Watton, T. J., D. A. Jerram, T. Thordarson, and R. J. Davies (2013), Three-dimensional lithofacies variations in hyaloclastite deposits, *J. Volcanol. Geotherm. Res.*, *250*, 19–33, doi:10.1016/j.jvolgeores.2012.10.011.
- Watton, T. J., K. A. Wright, D. A. Jerram, and R. Brown (2014), The petrophysical and petrographical properties of hyaloclastite deposits: Implications for petroleum exploration, *AAPG Bull.*, *98*(3), 449–463.
- Woodburn, N., A. Hardwick, H. Masoomzadeh, and T. Travis (2014), Improved signal processing for sub-basalt imaging, *Geol. Soc. London Spec. Publ.*, *397*(1), 163–171, doi:10.1144/sp397.6.
- Wright, K. A., R. J. Davies, D. A. Jerram, J. Morris, and R. Fletcher (2012), Application of seismic and sequence stratigraphic concepts to a lava-fed delta system in the Faroe-Shetland Basin, UK and Faroes, *Basin Res.*, *24*(1), 91–106, doi:10.1111/j.1365-2117.2011.00513.x.

Appendix C

A new nomenclature map of the mid-Norwegian margin: status update

By

Laurent Gernigon, Dmitry Zastrozhnov, Sverre Planke, Mohamed Mansour Abdelmalak, Jan Inge Faleide, Dwarika Maharjan, Ben Manton, Reidun Myklebust

The 34th Nordic Geological Winter Meeting, January 8th-10th 2020, Oslo, Norway.

A new nomenclature map of the mid-Norwegian margin: status update

Laurent Gernigon¹, Dmitry Zastrozhnov^{2,3}, Sverre Planke^{2,3}, Mohamed Mansour Abdelmalak³, Jan Inge Faleide³, Dwarika Maharjan², Ben Manton², Reidun Myklebust⁴

¹ Geophysics Group, Geological Survey of Norway (NGU), Trondheim, Norway, Laurent.Gernigon@NGU.NO

² Volcanic Basin Petroleum Research (VBPR), Oslo, Norway, dmitry@vbpr.no

³ Centre for Earth Evolution and Dynamics (CEED), Department of Geosciences, University of Oslo, Norway

⁴ TGS, Asker, Norway

Since the last official NPD nomenclature map of the mid-Norwegian margin (MNM) was published in 1995 (Blystad et al., 1995), a great amount of data including new 2D and 3D seismic surveys, refraction data, potential field data, seabed sampling, exploration wells, revised biostratigraphic schemes have been acquired. This has resulted in improved mapping of the volcanic rifted margin, especially in its deepest part and in the outer (volcanic) province. Within different scientific projects we perform a comprehensive interpretation of this new regional geological and geophysical dataset. This led to the compilation of new nomenclature map of the mid-Norwegian margin, where we unified our structural observations. In order to avoid map visualization complexities and to reflect basin configuration at different stages of the MNM development, we present two separate map sheets showing: 1) the pre-breakup structural elements and 2) syn- to post-breakup elements. The pre-breakup map sheet documents the first order rift architecture of the MNM before the onset of volcanism and includes new and revised structural elements on the platform, terraces and deep Møre and Vøring basin province and related fault systems. The syn- and post-breakup map sheet provides the most up-to-date regional mapping of the different volcanic facies recognized in the outer province. The second map sheet also includes sills and associated hydrothermal vent complexes, magnetic anomaly chrons and oceanic fracture zones, basin inversion structures, slides and glacial depocentres. After the publication, the digital version of the map will be freely available at NGU and VBPR websites and will include different layers in standard GIS formats (e.g. shapefiles) and a ready-to-use GIS project including embedded files, specific color codes, templates and detailed attribute tables. We believe that our new regional map brings a great addition and update to Blystad's landmark map and will help both academic and exploration communities to be introduced to the up-to-date regional framework of the MNM.