Title Page

Title of paper: Increased endothelial activation in patients with mixed connective tissue disease

Authors and affiliations: Mona Skjelland^{1,2}, Mahtab Zamani^{1,2}, Siri Hetlevik³, Vibke Lilleby³, Bente Halvorsen^{2,4}, Karolina Skagen¹

¹Department of Neurology, Oslo University Hospital Rikshospitalet, Oslo Norway

² Institute of Clinical Medicine, University of Oslo, Oslo, Norway

³Department of Rheumatology, Oslo University Hospital Rikshospitalet, Oslo, Norway

⁴ Research Institute of Internal Medicine, Oslo University Hospital, Rikshospitalet, Oslo, Norway

Abstract

Objective: Juvenile-onset mixed connective tissue disease (JMCTD) is a chronic inflammatory disease. We have previously demonstrated preclinical atherosclerosis in these patients, now exploring this further by assessing markers of endothelial dysfunction.

Methods: Thirty-three patients with JMCTD and 33 age-and sex-matched controls were included. sICAM-1, IL-6 and vWF were assayed from blood taken at the time of carotid ultrasound.

Results: Our major findings were: (i) Levels of sICAM-1 ($p<0.001$), IL-6 ($p=0.004$) and vWF ($p=0.001$) were higher, whereas (ii) HDL cholesterol (<0.01) and ApoA1 ($p<0.01$) were lower in the patient group compared to controls.

Conclusions: Patients with JMCTD had significantly increased levels of markers of endothelial dysfunction.

1. Introduction

Mixed connective tissue disease (MCTD) is a chronic inflammatory, systemic rheumatic, autoimmune disease where up to 23 % of patients present in childhood(1, 2). Patients with chronic inflammatory rheumatic disease manifest endothelial dysfunction, and have an increased risk of premature cardiovascular disease (CVD) such as stroke and myocardial infarction(3, 4). We have recently documented pre-clinical atherosclerosis with increased carotid intima-media thickness for patients with juvenile mixed connective tissue disease (JMCTD) independent of traditional risk factors for cardiovascular disease(5) . The relative contribution of systemic inflammation and disease-specific factors of rheumatologic disease in the development of atherosclerotic vascular disease in this patient group remains unclear. The majority of the reported increased cardiovascular risk is from adult patient populations making it difficult to explore the atherogenic mechanisms in this patient group, and a younger, juvenile population might therefore be better suited. The long-term risk for patients with JMCTD disease is uncertain(1, 2, 6) and the connection between endothelial cell damage and atherosclerosis in JMCTD, has not been described previously.

Endothelial cell dysfunction (ECD) is a hallmark for both initiation and maintenance of atherosclerosis resulting in detectable lesions of atherosclerotic cardiovascular disease (CVD), possibly serving as a marker for future risk of cardiovascular events (7). Dysfunctional endothelial cells express intercellular adhesion molecules on their surface allowing migration and recruitment of circulating leukocytes and accumulation of low density lipoprotein (LDL) across the vascular endothelium into the intimal space(8). High density lipoprotein (HDL) has been reported to have anti-atherogenic effect by countering this process with reverse cholesterol transport (RCT). Membrane bound intercellular adhesion molecule-1 (ICAM-1) is cleaved and released to form soluble ICAM-1 (sICAM-1) into the bloodstream where it acts as a marker of ICAM-1 and is upregulated during autoimmune activation(9). Concentration of sICAM-1 in serum/plasma has been associated with cardiovascular disease in several prospective epidemiological studies of mature adults (10, 11), and associations have been found between sICAM-1 and cardiovascular mortality in both healthy individuals (11, 12) and populations at high risk including pulmonary fibrosis and SLE (11, 13). Previous studies examining the relationship between sICAM-1 and subclinical atherosclerosis have produced mixed results with the positive association of sICAM-1 to preclinical atherosclerosis attenuated after adjustment for traditional cardiovascular risk factors (14, 15).

In this juvenile patient group with sub-clinical atherosclerosis we therefore aimed to assess endothelial activation in this group, assuming that increased endothelial activation would most likely be attributable to the inflammatory burden of the disease itself rather than cardiovascular risk factors.

2. Materials and Methods

2.1 Patients and control subjects

Data on the patients and controls included in this study as well as methods used for patient and control identification is reported previously(16). Patients with symptom onset before the age 18 years were included. Thirty-three patients had serum collected and represented the total sample available for analyzing markers of endothelial dysfunction. "Controls were randomly selected from the Norwegian population register. Controls with history of autoimmune disease necessitating immunosuppressive medication, heart disease, and/or lung disease (other than mild asthma) were excluded".

The study was approved by the Regional Ethics Committee for Medical Research (Id 2012/1721) and written informed consent was obtained from patients and controls <18 years old. For patients and controls younger than 16, consent was obtained from parents as well(17).

2.2 Carotid ultrasound and IMT measurement

Patients and controls were examined with color duplex US of the carotid arteries by an experienced ultrasonographer, who was blinded for patients' and controls' clinical and laboratory data. The examination was carried out in the supine position with the head angled at 45° toward the contralateral side. IMT measurements were synchronized with the QRS-complex on the echocardiogram and made in each carotid artery at the peak of the R wave where the lumen was widest. IMT was defined as the distance between the lumen-adventitia borders of the vessel, identified as a double-line pattern in a longitudinal image. IMT was measured in the carotid arteries on both sides of the neck in 3 segments: (1) in the far wall of the common carotid artery, 1 cm proximal to the bifurcation, over an area of 1 cm and in 3 different projections: lateral, posterior and anterior; (2) in the far wall of the carotid bifurcations over an area of 1 cm in the lateral position; (3) in the far wall of the proximal internal carotid arteries immediately distal to the bifurcation over an area of 1 cm in the lateral projection. We used the averaged value of the 10 carotid IMT measurements in each subject for further analyses. The carotid US examinations were carried out according to the American Society of Echocardiography guidelines²¹ using a Toshiba/Canon medical systems Aplio 500.

2.3 Clinical and blood sample assessment

All patients and controls underwent clinical examination at Oslo University Hospital (OUH) from March 2013 to June 2015. Details on physical examination as well as assessment of disease activity and clinical remission is reported previously (16). Fasting blood samples were analysed by routine methods at OUH and included; low density lipoprotein (LDL), high density lipoprotein (HDL), triglycerides (TG), apolipoprotein A1 (ApoA1), apolipoprotein b (ApoB), fasting glucose, glycosylated haemoglobin (HbA1c). Serum samples were separated from whole blood and were kept at -80°C until biochemical analysis. Serum levels of sICAM, was measured by Meso Scale Diagnostics V-plex. Rockville, USA and IL-6 was ELISA (Bio-Techne, Minneapolis, USA) and von Willenbrand

factor (vWF) quantified by in house analysis using antibody obtained from R&D Systems, Abingdon, UK

Cholesterol efflux capacity was measured as previously described (18, 19) Cholesterol efflux PMA differentiated THP-1 macrophages were lipid loaded by incubation with ox LDL (20 µg/ml) in regular growth medium plus 0.5 ICi/l (18.5 mBq/l) ¹⁴C-Cholesterol (American Radiolabel Chemicals, Saint Louis, MO), dissolved in ethanol, was added. After 24 h, radiolabeled media were removed, and the foam cells were washed twice with 0.2% bovine serum albumin (BSA) (wt:v) in RPMI. Thereafter, the cells were incubated in RPMI 1640 with patient or control sera (2.5%, v/v) as cholesterol acceptors. After 34 h, the cell medium was collected and the cells were harvested in 0.2 mol/L NaOH. The radioactivity was measured by liquid scintillation counting using TRI-CARB 2300 TR Scintillation Counter (Packard, Waltham, MA). Data are presented as fractional (%) cholesterol efflux calculated as $\text{dpm (media)}/\text{dpm (media + cell-associated)} \times 100$. Percentages efflux was calculated using the equation; $[\text{DPM}_{\text{medium}}/(\text{DPM}_{\text{cell+medium}})] \times 100$.

2.4 Statistical analysis

Categorical variables were compared with the Mann-Whitney u test (two groups) and Student's t-test/paired samples t-test was used for comparison of normally distributed data. Patients and controls were compared using paired testing. The Chi square test was used for analyzing categorical data. All calculations were performed with SPSS for Windows statistical software (Version 21.0; SPSS Inc, Chicago, IL). The correlation of well-known risk factors for cardiovascular disease to the markers of endothelial dysfunction was assessed on primary analysis. The factors found significant were included in the regression analysis.

3. Results

3.1 Baseline characteristics of patients and controls

Thirty-three patients had blood tests at the same time as carotid ultrasound and were all included for analysis of markers of endothelial dysfunction. Patient characteristics are shown in Table 1. Twenty-eight patients were female. The mean age of diagnosis in the patient group was 13 years, whereas mean duration of disease was 16.9 years. In the control group there were 29 female patients (88%) and the mean age was 28.7 years (SD 9.4).

Table 1. Patient Characteristics.

Characteristics	Patients with JMCTD, n=33
-----------------	---------------------------

Female gender*	28 (84.8)
Age at inclusion, years	27.9 (9.7)
Age at diagnosis, years	13.3 (3.6)
Disease duration	16.9 (10.1)
PGA at inclusion, cm VAS**	17 (4-58)
Current medication at inclusion*	12 (36)
Steroids per os	4 (12)
Azathioprine	10 (30)
Methotrexate	19 (57)
Hydroxychloroquine	1 (3)
Mycophenolate Mofetil	1 (3)
Rituximab	

The values are given as mean (SD) or *number (percentage) or **median (range). PGA: Physician global assessment.

Cardiovascular risk factors for patients and controls are shown in Table 2. Patients had significantly lower levels of HDL cholesterol compared with controls (1.3 vs 2.8 mmol/l, $p < 0.01$), as well as lower levels of Apolipoprotein A at 1.4 vs 1.69 mmol/L, $p < 0.01$). Patients had higher levels of both triglycerides and ESR compared with controls (1.1 ± 0.8 vs 0.8 ± 0.3 mmol/l, $p = 0.13$) and (11.4 ± 8.6 vs 5.8 ± 3.9 , $p < 0.01$), respectively. Intima media thickness remained significantly higher in patients compared with healthy controls (0.58 ± 0.08 vs 0.53 ± 0.07 , $p = 0.025$).

Table 2. Cardiovascular risk factors in patients with juvenile onset MCTD and controls.

Variables	Patients, n=33	Controls, n=33	P
BMI, kg/m ²	22.9 $\pm$ 3.9	23.5 $\pm$ 3.1	0.44
Cigarette smoking, yes n (%)	6 (18.2)	6 (18.2)	0.70
Diabetes, n (%)	1 (3.0)	0	0.32
Systolic blood pressure, mmHg	109 $\pm$ 13.9	115 $\pm$ 14.7	0.12
Diastolic blood pressure, mmHg	63 $\pm$ 16.9	68 $\pm$ 9.4	0.19
HDL, mmol/L	1.3 $\pm$ 0.4	2.8 $\pm$ 0.9	<0.001
LDL, mmol/L	2.5 $\pm$ 0.8	2.8 $\pm$ 0.9	0.14
Triglycerides, mmol/L	1.1 $\pm$ 0.8	0.8 $\pm$ 0.3	0.13
ESR mm	11.4 $\pm$ 8.6	5.8 $\pm$ 3.9	0.003
HbA1c, %	5.3 $\pm$ 0.2	5.4 $\pm$ 0.9	0.56
CRP, mg/L*	2.3 $\pm$ 3.2	1.4 $\pm$ 1.9	0.19
Apolipoprotein A1	1.42 $\pm$ 0.29	1.69 $\pm$ 0.27	0.001
Apolipoprotein B	0.79 $\pm$ 0.24	0.85 $\pm$ 0.22	0.36

Values are mean $\pm$ SD or *median (range). BMI: body mass index; HDL: high-density lipoprotein; LDL: low-density lipoprotein; HbA1c: glycosylated hemoglobin; CRP: C-reactive protein; ESR: erythrocyte sedimentation rate; IMT: intima media thickness. °Smoke occasionally or every day.

3.2. Cholesterol efflux capacity.

Even though within normal reference values, both HDL and ApoA1 were significantly reduced within patients as compared to controls. The degree of lipid accumulation in the progression of atherosclerosis is dependent on the efficacy of cholesterol efflux involving high-density lipoprotein (HDL)-related mechanisms. The comparatively low levels of HDL in patients poses the question whether they had reduced atherogenic protection causing the observed increase in atherosclerosis. To explore this, we assessed the cholesterol efflux capacity exhibiting the functional properties of serum as surrogate marker of HDL's efflux capacity. We, however, found the efflux to be similar in the two groups with mean efflux capacity (%) of 5.5 ± 1.3 SD in patients compared to 5.5 ± 0.9 SD in controls.

3.3 Increased serum levels of sICAM, IL-6 and vWF in patients compared with controls

Next, we assessed whether the patient's low HDL could cause altered endothelial function. To study this we measured 3 classic endothelial markers (sICAM-1, IL-6 and vWF). On paired analysis assessing serum levels we found higher levels of sICAM (256.2 ± 107.4 vs 111.3 ± 40.5 , $p < 0.001$), vWF (251.1 ± 165.4 vs 167.0 ± 68.5 , $p = 0.01$) and IL-6 (38.3 ± 48.6 vs 17.9 ± 23.9 , $p = 0.04$) in patients compared to controls respectively. On regression analysis with adjustment for factors significant on primary analysis (ESR, Apolipoprotein A and HDL) sICAM-1 ($p < 0.001$) and HDL ($p < 0.001$) remained significantly higher in patients compared with controls.

Figure 1.

Fig 1. Box plot showing levels of IL-6, s-ICAM and vWf in patients and healthy controls. The bottom and the top boxes represent the first and third quartile. Horizontal lines in boxes represent the median value and the whiskers the range limit. Median levels of IL-6, s-ICAM and vWf were significantly higher in patients compared to controls. ** $p < 0.001$ and * $p < 0.05$.

4. Discussion

This study investigated circulating markers of endothelial dysfunction in patients with juvenile MCTD known to have pre-clinical atherosclerosis. The main findings were that patients had (i) increased levels of sICAM-1, vWF and IL-6, and (ii) lower levels of HDL and ApoA1 compared to healthy age- and sex-matched controls. Elevated sICAM-1 in patients with negligible presence of traditional cardiovascular risk factors, indicates that the driver of endothelial dysfunction and atherosclerosis in this group might be related to the inflammatory burden associated with the underlying disease itself.

Chronic inflammation plays a key role in the development of atherosclerotic CVD and chronic inflammatory diseases such as MCTD, Rheumatoid arthritis (RA) and systemic lupus erythematosus (SLE) are known to be associated with an increased risk of premature CVD. CVD in patients with adult connective tissue disease, however, has also been associated with an increased prevalence of traditional risk factors and a juvenile population might therefore be better suited to explore the mechanisms involved in the development of premature atherosclerosis in this disease.

IL-6 and vWF were increased in patients compared to controls (although this difference was attenuated when adjusting for erythrocyte sedimentation rate (ESR)), likely reflecting the general inflammatory state of the patients. The increased levels of sICAM-1 in these patients with documented preclinical atherosclerosis are in keeping with previous studies reporting sICAM-1 to be associated with both IMT (20) and change in IMT(21). Adhesion molecules are key players in the development of atherosclerosis by facilitating monocyte recruitment to the endothelium and migration across the endothelial surface into the intima wall layer. In fact, previous reports have indicated associations between ICAM-1 and stroke with ICAM-1 expression found up-regulated in vascular endothelium of patients with ischemic stroke(22).

In this current study, high-density lipoprotein (HDL) and ApoA1 were higher in controls compared to patients indicating a possible loss of vascular protection from HDL in the patient group. Cholesterol efflux, however, demonstrated patients to have HDL function similar to control group in its removal of radiolabeled cholesterol from lipid laden macrophages. Even though the efflux capacity was normal, it is tempting to speculate that the reduced HDL level causes reduced protection of endothelium and subsequent activation driven by the inflammatory burden in the patients. Further studies investigating the mechanisms leading to increased subclinical atherosclerosis in this patient group is warranted. Despite the number of patients included in this study being comparable to other published reports on markers of ECD and cardiovascular risk in patients with connective tissue disease, it still remains a small population. The strengths of this study are the relatively long duration of follow up from time of diagnosis/disease onset of a well-defined cohort of patients “..results compared with paired controls, allowing for the presentation of novel data on markers of endothelial dysfunction and pre-clinical atherosclerosis in JMCDT.

Acknowledgment

The author thanks Ellen Lund Sagen for technical assistance. The study is partly funded by Norwegian Research Council RC grant 144245

References

1. Kotajima L, Aotsuka S, Sumiya M, Yokohari R, Tojo T, Kasukawa R. Clinical features of patients with juvenile onset mixed connective tissue disease: Analysis of data collected in a nationwide collaborative study in Japan. *J Rheumatol* 1996;23:1088-94.
2. Burdt MA, Hoffman RW, Deutscher SL, Wang GS, Johnson JC, Sharp GC. Long-term outcome in mixed connective tissue disease: Longitudinal clinical and serologic findings. *Arthritis Rheum* 1999;42:899-909.
3. Bjornadal L, Yin L, Granath F, Klareskog L, Ekbom A. Cardiovascular disease a hazard despite improved prognosis in patients with systemic lupus erythematosus: Results from a Swedish population based study 1964-95. *The Journal of rheumatology* 2004;31:713-9.
4. Symmons DP, Gabriel SE. Epidemiology of CVD in rheumatic disease, with a focus on RA and SLE. *Nature reviews Rheumatology* 2011;7:399-408.
5. Skagen K, Hetlevik SO, Zamani M, Lilleby V, Skjelland M. Preclinical carotid atherosclerosis in patients with juvenile-onset mixed connective tissue disease. *J Stroke Cerebrovasc Dis* 2019.
6. Mier RJ, Shishov M, Higgins GC, Rennebohm RM, Wortmann DW, Jerath R, et al. Pediatric-onset mixed connective tissue disease. *Rheum Dis Clin North Am* 2005;31:483-96, vii.
7. Stary HC. Natural history and histological classification of atherosclerotic lesions: An update. *Arterioscler Thromb Vasc Biol* 2000;20:1177-8.
8. Libby P, Ridker PM, Maseri A. Inflammation and atherosclerosis. *Circulation* 2002;105:1135-43.
9. Witkowska AM, Borawska MH. Soluble intercellular adhesion molecule-1 (sICAM-1): An overview. *Eur Cytokine Netw* 2004;15:91-8.
10. Ridker PM, Hennekens CH, Roitman-Johnson B, Stampfer MJ, Allen J. Plasma concentration of soluble intercellular adhesion molecule 1 and risks of future myocardial infarction in apparently healthy men. *Lancet* 1998;351:88-92.
11. Malik I, Danesh J, Whincup P, Bhatia V, Papacosta O, Walker M, et al. Soluble adhesion molecules and prediction of coronary heart disease: A prospective study and meta-analysis. *Lancet* 2001;358:971-6.
12. Rohde LE, Hennekens CH, Ridker PM. Cross-sectional study of soluble intercellular adhesion molecule-1 and cardiovascular risk factors in apparently healthy men. *Arterioscler Thromb Vasc Biol* 1999;19:1595-9.
13. Gross MD, Bielinski SJ, Suarez-Lopez JR, Reiner AP, Bailey K, Thyagarajan B, et al. Circulating soluble intercellular adhesion molecule 1 and subclinical atherosclerosis: The coronary artery risk development in young adults study. *Clin Chem* 2012;58:411-20.
14. Tang W, Pankow JS, Carr JJ, Tracy RP, Bielinski SJ, North KE, et al. Association of sICAM-1 and MCP-1 with coronary artery calcification in families enriched for coronary heart disease or hypertension: The NHLBI Family Heart Study. *BMC Cardiovasc Disord* 2007;7:30.
15. Rohatgi A, Owens AW, Khera A, Ayers CR, Banks K, Das SR, et al. Differential associations between soluble cellular adhesion molecules and atherosclerosis in the Dallas Heart Study: A distinct role for soluble endothelial cell-selective adhesion molecule. *Arterioscler Thromb Vasc Biol* 2009;29:1684-90.
16. Hetlevik SO, Flato B, Rygg M, Nordal EB, Brunborg C, Hetland H, et al. Long-term outcome in juvenile-onset mixed connective tissue disease: A nationwide Norwegian study. *Ann Rheum Dis* 2017;76:159-65.
17. Hetlevik SO, Flato B, Aalokken TM, Lund MB, Reiser S, Mynarek GK, et al. Pulmonary manifestations and progression of lung disease in juvenile-onset mixed connective tissue disease. *J Rheumatol* 2019;46:93-100.
18. Halvorsen B, Holm S, Yndestad A, Scholz H, Sagen EL, Nebb H, et al. Interleukin-10 increases reverse cholesterol transport in macrophages through its bidirectional interaction with liver X receptor alpha. *Biochem Biophys Res Commun* 2014;450:1525-30.

19. Macpherson ME, Halvorsen B, Yndestad A, Ueland T, Mollnes TE, Berge RK, et al. Impaired hdl function amplifies systemic inflammation in common variable immunodeficiency. *Sci Rep* 2019;9:9427.
20. Papagianni A, Dovas S, Bantis C, Belechri AM, Kalovoulos M, Dimitriadis C, et al. Carotid atherosclerosis and endothelial cell adhesion molecules as predictors of long-term outcome in chronic hemodialysis patients. *Am J Nephrol* 2008;28:265-74.
21. Kondo K, Kitagawa K, Nagai Y, Yamagami H, Hashimoto H, Hougaku H, et al. Associations of soluble intercellular adhesion molecule-1 with carotid atherosclerosis progression. *Atherosclerosis* 2005;179:155-60.
22. Deddens LH, van Tilborg GA, van der Toorn A, van der Marel K, Paulis LE, van Bloois L, et al. Mri of icam-1 upregulation after stroke: The importance of choosing the appropriate target-specific particulate contrast agent. *Mol Imaging Biol* 2013;15:411-22.