Title:

An empirical study of factors associated with height-for-age z-scores of children aged 6–23 months in northwest Rwanda: the role of care practices related to child feeding and health.

Authors:

Theogene Dusingizimana^{1,2 *}, Janet L Weber¹, Thiagarajah Ramilan³, Per Ole Iversen^{4,5,6} and Louise Brough¹

¹ School of Food and Advanced Technology, Massey University, Tennent Drive, Palmerston North 4442, New Zealand.

² Department of Food Science and Technology, College of Agriculture, Animal Sciences and Veterinary Medicine, University of Rwanda, P.O. Box 210 Musanze, Rwanda.

³ School of Agriculture and Environment, Massey University, Tennent Drive, Palmerston North 4442, New Zealand.

⁴ Department of Nutrition, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway.

⁵ Department of Hematology, Oslo University Hospital, Oslo, Norway.

⁶ Division of Human Nutrition, Faculty of Medicine and Health Sciences, Stellenbosch University, Tygerberg, South Africa.

Corresponding author:

Theogene Dusingizimana

¹ School of Food and Advanced Technology, Massey University, Tennent Drive, Palmerston North 4442, New Zealand

² Department of Food Science and Technology, College of Agriculture, Animal Sciences and Veterinary Medicine, University of Rwanda, P.O. Box 210 Musanze, Rwanda

Email address: T.Dusingizimana@massey.ac.nz / dusingizetho@yahoo.fr

This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114520004961

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Short title:

Determinants of child nutritional status in Rwanda.

Introduction

Stunting, or low height-for-age (HAZ) ⁽¹⁾, also referred to as growth retardation, is the most common form of undernutrition among children under 5 years ⁽²⁾. With an estimated 149 million (22%) of children affected ⁽³⁾, stunting has been identified as a global public health priority. Stunting is associated with an increased risk of morbidity and mortality, poor cognitive and physical development, with significant educational and economic consequences ⁽⁴⁾ as well as an increased risk of non-communicable diseases in adulthood ^(4, 5). The World Health Assembly has set targets to reduce stunting by 40% between 2010 and 2025 ⁽⁶⁾; however, at the current rate of reduction, several countries, especially low- and middle-income countries (LMIC) are unlikely to achieve the targets ⁽²⁾.

For many years, policies to reduce child undernutrition in LMIC, particularly in sub-Saharan Africa, have often focused on improving household income and food security ^(7, 8). However, evidence shows that increasing household income and food security, though necessary, is not sufficient for improved child nutritional status ⁽⁹⁾. Research also shows that interventions to improve child nutritional status through improved household income and/or food security depend largely on child care practices within the household ^(10, 11). Several studies showed that better care practices related to feeding and preventive health practices can significantly improve child nutritional status, even in households with limited economic and food resources ⁽¹²⁻¹⁴⁾.

In Rwanda, 35% of children younger than 5 years were stunted in 2018 ⁽¹⁵⁾. Data on prevalence of stunting in Rwanda show a national downward trend (48.3% in 2000 to 35% in 2018); however, the reduction has not been consistent across the country. For example, in 14

out 30 districts of Rwanda, 40%–54% of children under 5 years are stunted ⁽¹⁵⁾. Based on the World Health Organization (WHO) thresholds ⁽¹⁶⁾, these rates of stunting are of very high public health significance and require urgent actions. Thus, research is needed to understand the factors underlying the persistence of child stunting in Rwanda. Given the significant disparity in stunting prevalence, a greater understanding of subnational level factors is needed to inform the design of more targeted programmes and policies to effectively address child stunting in Rwanda.

Studies ^(17, 18) have documented gaps in care practices related to child feeding practices in Rwanda. For example, the number of children aged 6–23 months who met the minimum acceptable diet remained unchanged (17%) between 2010 and 2018 ⁽¹⁵⁾. Increasing evidence also suggests that poor care practices, including sub-optimal child feeding practices are potentially contributing to the high rates of child undernutrition in Rwanda ^(15, 19, 20). In addition, studies indicate inadequate health practices such as low utilization of child growth monitoring for screening of child nutritional status ⁽²¹⁾. However, research is limited on the influence of child feeding patterns on child nutrition status. A few studies examining the relationship between infant and young child feeding (IYCF) practices and child nutrition status in Rwanda have assessed single indicators of feeding practices and produced mixed results. Uwiringiyimana *et al.* ⁽²²⁾ found that exclusive breastfeeding was significantly positively associated with HAZ in children aged 5–30 months ($n= 138$). Conversely, Matsiko ⁽²³⁾ found that achieving minimum dietary diversity, minimum meal frequency and acceptable diet were not significantly associated with HAZ in children aged 6–12 months ($n= 192$). Given that child feeding has several dimensions (e.g. the type, quality and variety of foods, the quantity of foods, responsive feeding, etc.), studies examining the association between single feeding practices and child nutrition status may not provide a comprehensive picture of the association between overall child feeding patterns and nutritional outcomes ⁽²⁴⁾. Ruel &

Menon ⁽²⁵⁾ proposed a novel approach of assessing the relationship between child feeding patterns and nutritional status using an index that combined various dimensions of child feeding. Drawing from Ruel and Menon's approach, Condo *et al.* ⁽²⁶⁾ examined the association between a feeding index and nutritional status in Rwanda. The authors found no association between the feeding index and child nutritional status; however, their findings may not apply to all children because the study focused on HIV-infected children whose caregivers' feeding practices may differ from those of caregivers with healthy children.

The objectives of the current study were to identify the factors associated with child stunting in Rwanda, including the relationship between indices of child feeding and health practices and child stunting. Our assumption was that better feeding and health practices are positively associated with better child HAZ, and that mothers of non-stunted children have better child feeding and health practices than those of stunted children. This study contributes to the knowledge on the determinants of child stunting in Rwanda and extends the literature on the usefulness of indices of child feeding and health practices in relation to child nutritional status in the Rwandan context.

Methods

Settings

The study was conducted in Rutsiro district, northwest of Rwanda, 150 km from the capital city, Kigali. The inhabitants of the district are predominantly rural and engaged in subsistence agriculture ⁽¹⁵⁾. About 50% of the population in Rutsiro are food insecure, and the prevalence of stunting increased from 46% in 2015 to 54% in 2018 ⁽¹⁵⁾. The public health system of the district consists of a network of 17 health centres and one district hospital ⁽²⁷⁾.

Design and sample size

This was a cross-sectional study involving mother-child pairs. The sample size was calculated using the Cochran formula for estimation of a single proportion ⁽²⁸⁾. Although our focus was not to estimate the proportion of stunted children, this would allow us to analyse the determinants of child nutritional status with a 95% confidence of having an anticipated prevalence of stunting.

$$n = Z^2 pq/d^2$$

where n is the sample size, Z is 1.96, p is the proportion of stunting (46% at the time of the survey ⁽²⁹⁾), q is 1-p, and d is the relative desired precision of 5%. The calculated sample size (n= 382) was increased by 5% to account for contingencies such as recording error, resulting in a final sample of 400. Because of the variability in geographical landscape, it was assumed that child feeding and/or health practices might differ across the district. Therefore, the district was divided into three zones based on the main roads connecting Rutsiro district to its neighboring districts. Health centres were used as an entry point to facilitate access to the community, and community health workers (CHWs) who work under supervisions of the health centers assisted in identifying potential participants. Three health centres (9 in total across the district) were purposively selected in each zone to maximize geographic distribution. Within each health centre's catchment area, two administrative entities (i.e. cells) were selected. The cell containing the health centre was automatically selected, and the next was randomly selected among distant cells. In each cell, one village was randomly selected, and a list of all potentially eligible mothers with children aged 6–23 months was compiled using growth monitoring records obtained from CHWs, and children were randomly selected from the lists. Eligibility criteria were: (1) child aged 6 to 23 months; (2) child was apparently healthy; and (3) being in the lowest socioeconomic categories. All

participants gave oral informed consent prior to data collection. The survey was undertaken from September 2018 to January 2019.

Ethical approval

All study guidelines were in accordance with the guidelines laid down in the Declaration of Helsinki. The study procedures were also approved by the Massey University Human Ethics Committee (reference: SOA 17/67) and the Institutional Review Board of the University of Rwanda's College of Medicine and Health Sciences (reference: 003/CMHS IRB/2017).

Permission to collect data was also obtained from the District public health office. Oral informed consent was obtained from all participants, after the purpose of the study was explained. Each participant received 5 soap bars (~\$US 1) as a recognition for their participation.

Face-to-face interviews were conducted with mothers, and data were collected using a structured questionnaire digitally preprogrammed on the CommCare platform⁽³⁰⁾. The data was captured on a hand-held tablet. Information on socio-demographic characteristics, maternal and child characteristics were collected. Mothers also reported infant birth weight, child morbidities in previous 4 weeks: diarrhoea (defined as ≥ 3 watery or loose stools per day), upper respiratory infections (coughing, runny nose, or wheezing), fever, and other illnesses. In addition, household socioeconomic characteristics such as size, number of children under 5 years, ownership of agricultural land and other household assets, and source of drinking water were reported. Altitude and location of the household premises were recorded using a handheld Global Positioning System (Tremble Juno SB Handheld).

Measures

Anthropometry

Anthropometric measurements were taken following standard procedures⁽³¹⁾. Child height was measured to the nearest 1 mm in a recumbent position using a UNICEF designed length board. The height of mothers was measured using a wall-mounted portable stadiometer to the nearest 0.1 cm. The weight of both mother and child was measured to the nearest 100 g using an electronic scale (SECA Model 874, Hamburg, Germany).

Feeding and health practices

Breastfeeding and complementary feeding practices, health practices (antenatal care visits during the last pregnancy, attendance at growth monitoring, use of deworming tablets, child immunization status, use of vitamin A supplements) were reported by the mother. Birth certificates and immunization records were used to verify information on birth weight and immunization status, respectively.

Infant and child feeding index

We adapted the methodology used by Ruel & Menon⁽²⁵⁾ and others⁽³²⁻³⁴⁾ to construct an infant and child feeding index (ICFI). The ICFI consisted of five child feeding practice components: current breastfeeding (yes/no), duration of exclusive breastfeeding (i.e. whether the mother exclusively breastfed for the first 6 months), dietary diversity indicator, feeding frequency, and responsiveness during feeding. Dietary diversity and meal frequency were calculated using 24 h dietary recall data⁽³⁵⁾. The feeding practice components and the scoring system used to create the ICFI for the different age-groups are described in the **Supplemental Table 1**. Briefly, a score of 1 or 2 was assigned for a positive practice, and a score of 0 was assigned for a potentially negative practice. The practices were judged positive or negative

based on child feeding recommendations ⁽³⁶⁾ as well as evidence about their potential benefits or risks ⁽³⁷⁾.

For children aged 6–8 months and 9–11 months, a score of 2 was assigned for mothers whose children were still breastfeeding. For children aged 12–23 months, a score of 1 was assigned for mothers whose children were still breastfeeding. For all children, a score of 0 was assigned for mothers whose children were not breastfeeding. A score of 1 was also assigned for mothers who reported exclusive breastfeeding for the first 6 months, and a score of 0 was given to those who introduced complementary foods to their children prior to 6 months.

A dietary diversity indicator was created using information collected on the number of food groups consumed by the child in the previous 24 hours. Following the guidelines for young child feeding ⁽³⁵⁾, seven food groups, adapted from the Rwanda's Demographic and Health Survey ⁽²⁹⁾, were considered: grains/roots/tubers, legumes/nuts, milk or dairy products, flesh foods (meat/fish/poultry), eggs, vitamin A fruits and vegetables, and other fruits and vegetables. A score of 1 was given for each food group consumed, resulting in a dietary diversity score ranging from 0 to 7 which was divided into three categories: low, medium, and high. For all age groups, low category included children who had not received any foods. For both 6–8 months and 9–11 months age groups, a child was classified into medium category if a child received 1–2 food groups, or into high category if a child received 3 or more food groups. For children 12–23 months, a medium category comprised of children who had received 2–3 food groups, whereas a high category comprised those receiving ≥ 4 food groups.

Feeding frequency scoring criteria used information collected on the number of meals received by the child in the past 24 hours. This indicator reflects the WHO recommended

minimum meal frequency for breast-fed and non-breast-fed children (i.e. breast-fed children aged 6–8 months and 9–23 months should receive solid- or semi-solid or soft foods at least 2 times a day and 3 times a day, respectively). Thin/watery porridges and trivial snacks were not counted ⁽³⁵⁾.

Indicator of responsive feeding was based on current young child feeding recommendation ⁽³⁶⁾, and it was informed by a previous study ⁽³⁸⁾ conducted in the same population. It was defined as caregiver's self-report of encouraging food intake of children who refuse to eat due to loss of appetite or during diarrhoea ⁽¹⁰⁾. A score of 1 was assigned for an action to encourage food intake, and a score of 0 if the action potentially restricted/discouraged child's food intake.

The ICFI was created by adding up the scores obtained from all the component practices. For each age group, the minimum ICFI score was 0 and maximum possible ICFI score was 9. The scores were ranked and tertiles created to form three categories: low, medium and high. Grouping the ICFI score into tertiles, although arbitrary, was done to allow comparison with other studies that have examined the relationship between ICFI and child HAZ.

Health practices index

A methodology used by Ruel *et al.* ⁽¹³⁾ was adapted to develop a health practices index (HPI). Five health practices were considered: attendance at growth monitoring in the past month, the number of antenatal care visits when pregnant with the study child, child diarrhoea treatment methods, vitamin A supplementation (whether the child received vitamin A supplements in the previous 6 months) and immunization status (whether the child received all age-specific vaccines). The scoring system for the different health practices is described in the

Supplemental Table 2.

Household hunger level

Household hunger level was assessed and categorized using a validated cross-cultural household hunger scale (HHS) ⁽³⁹⁾. Following the HHS measurement guide, a 6-point household hunger score was generated and categorized into three levels of household hunger: little or no hunger (score: 0–1 score); moderate hunger (score: 2–3), and severe hunger (score: 4–6). The moderate and severe hunger level categories were combined to form one category, i.e. moderate/severe, as there were few households (4.2%, n=16) in the severe hunger category.

Household wealth index

A household wealth index – a proxy measure of household socioeconomic status – was constructed using a principal component analysis (PCA) ⁽⁴⁰⁾, with the following 12 variables: access to agricultural land, purchase of commercial fertilizer in the previous cropping season, the quality of housing (type of floor, wall), degree of household crowding, i.e. the number of persons sleeping in a room (1 if ≤ 3 persons/room and 0 if ≥ 4 persons/room), source of lighting (none, petrol lamp or solar/electricity), ownership of durable assets (radio, thermos, mattress, umbrella), ownership of health insurance for all household members, and a measure of animal ownership expressed in tropical livestock units (TLU). The TLU is a metric combining multiple species of livestock into a weighted measure representing total body weight and potentially market value ⁽⁴¹⁾. The TLU variable was skewed and, therefore, 3 categories were created. All variables were thus categorical, and were ranked in ascending order, i.e. from the worst to the best. The scores of first two extracted component of the PCA were used as a measure of household wealth index and included as covariates in multivariable regression analyses.

Statistical analysis

The WHO Anthro software version 3.2.2 was used to calculate height-for-age z-scores (HAZ) ⁽⁴²⁾. A child was classified as stunted if his/her HAZ was 2 standard deviations (SD) below the median of the WHO reference population ⁽⁴³⁾. A binary indicator of stunting (stunted/non-stunted) was used to allow comparison between care practices and other characteristics of mothers of non-stunted and those of stunted children. Continuous variables were summarized using means and SD or medians and interquartile ranges (IQR) for non-normally distributed variables. Categorical variables were summarized using frequencies and percentages. Mann-Whitney test or independent *t* test (continuous variables) and Chi-square test (categorical variables) were used to test differences in care practices and other characteristics between non-stunted and stunted children. Multiple linear regression analyses were used to examine the relationship between various factors and child HAZ (dependent variable). To identify variables to include in the regression models, we first represented hypothetical assumptions about the causal relationships between various factors and child HAZ using DAGitty software ⁽⁴⁴⁾. The software was used to construct directed acyclic graphs (DAGs) which specify the relationships among factors based on *a priori* evidence, theoretical knowledge and researcher's subject matter expertise ⁽⁴⁵⁾, while enabling identifying covariates to adjust for in multiple regression models ⁽⁴⁵⁾. Although we aimed to examine association between various factors and child HAZ, ICFI and HPI were particularly of interest because feeding and health practices are modifiable factors often targeted by interventions to prevent stunting. Therefore, two different regression models containing ICFI and HPI as main independent variables were fitted. The first model (model 1) examined association between ICFI and child HAZ, adjusting for all other covariates at child-, maternal-, and household-level included in the model. The second model (model 2) examined association between HPI and child HAZ, adjusting for the same covariates as in model 1. Since feeding practices are linked to child food intake, which is a direct determinant of child HAZ (besides illnesses), we also adjusted

for ICFI in model 2. Because of age-related variability in child feeding practices ⁽³⁶⁾, we ran separate regression analyses: for all children, and for those aged 6–11 months and 12–23 months. All assumptions were assessed for all models. Multicollinearity was checked using the variance inflation factor (VIF) of 10 for all explanatory variables ⁽⁴⁶⁾. Homoscedasticity was assessed by visual inspection of a plot of standardized residuals versus standardized predicted values. Variables with $P < 0.05$ were considered statistically significant. All analyses were performed using SPSS version 25 (IBM, Armonk, NY, USA).

Results

Of the 400 child-mother pairs recruited, twenty-one children (5%) were excluded from the analysis due to preterm birth or lack of health cards to verify birthweight information. Thus, the descriptive analysis was performed on 379 children. Of these, 35 mothers (9%) had no height measurements, whereas 10 (2.5%) had no data on education. The final sample considered for regression analyses was thus 334 (~84%) mother-child pairs. The median age was significantly lower ($P < 0.001$) for non-stunted children than for stunted children (**Table 1**). Mothers of non-stunted children were older and taller than those of stunted children. Approximately 60% of the mothers had either never attended school or did not complete primary education; 16% had some secondary education, and 65% did not achieve the recommended number of ≥ 4 antenatal care visits. Forty-three percent of households had experienced moderate to severe hunger in the past 4 weeks.

Nutritional status

The mean (SD) HAZ was -1.69 (SD 1.3). Thirty-eight percent of the children were stunted (Table 3) and the stunting prevalence in older children was more than twice that in younger children (47% vs 21%, $P < 0.001$). There was no significant difference in the prevalence of stunting between males and females or between household wealth index categories.

Child feeding practices

Almost all children (96%) were still breastfed at the time of the survey. However, over one third of the mothers (39%) introduced complementary foods to their children before the recommended age (i.e. 6 months) (**Table 2**). Approximately half (54%) of children were fed at the recommended minimum frequency; however, the diversity of foods consumed was low. On average, children consumed 3 food groups, and only 36% met the WHO recommended minimum dietary diversity (≥ 4 food groups) (**Supplemental Table 3**). As compared to children aged 12–23 months, children in the age group of 6–11 months were less likely to achieve the recommended minimum dietary diversity (23% vs 42%, $P=0.001$) (data not shown). Nearly all children (93%) consumed staples (grains, roots or tubers), whereas the consumption of animal source foods was low (27%) (**Supplemental Table 3**). The mean ICFI score for all children was 5.8 (SD 1.6.) Non-stunted children had significantly higher ICFI score than stunted children (6.0 vs. 5.4, $P=0.001$) (Table 2).

Child morbidity and health practices

Morbidity among children during the previous 4 weeks was high: 47% had experienced diarrhoea, 78% had symptoms of respiratory infections, 43% had fever and 11% had other illness (**Supplemental Table 3**). No significant difference was observed in the prevalence of morbidity between non-stunted and stunted children. However, results from sub-analyses (data not shown) showed that, in younger children (6–11 months), stunted children were more likely to have diarrhoea than non-stunted (64% vs 42%, $P=0.05$). About half (46%) of the mothers whose children had suffered from diarrhoea reported either never seeking treatment or using home remedies to treat their child diarrhoea. The prevalence of stunting was higher among children whose mothers did not seek treatment or used home remedies for child diarrhoea than among children whose mothers sought diarrhoea treatment from a health centre or health worker (58% vs 48%, $P=0.057$). The mean HPI was 4.5 (SD 1.0), with no significant difference between non-stunted and stunted children ($P=0.58$) (**Table 3**).

Association between ICFI, HPI and child nutritional status

Results from bivariate analyses showed that ICFI was associated with child nutritional status. We found a significant difference of 0.34 in mean HAZ between children in the highest compared to the lowest ICFI tertiles ($P < 0.001$) (data not shown). Additionally, the prevalence of stunting was significantly lower among children whose mothers were in the high ICFI tertile compared to those in the lowest tertile (27% vs 50%, $P = 0.001$) (**Table 2**). In bivariate analysis, the association between HPI and HAZ was not statistically significant ($P = 0.10$) (data not shown). The prevalence of stunting was also not as expected; it was significantly higher among children whose mothers were in the medium HPI tertile as compared to low and high tertiles (Table 3). However, when the two lowest tertiles (low and medium) were combined and compared with the high tertile, the prevalence of stunting was significantly lower among children in the highest tertile compared to the lowest tertile (33.5% vs. 43.8 %, $P = 0.04$).

Results from multiple linear regression analyses are shown in **Table 4**. There was a significant and positive association between ICFI and HAZ, after controlling for potential confounders at child-, maternal-, and household-level. We found a mean difference of 0.14 HAZ between children of mothers in the high ICFI tertile compared to those in the low ICFI tertile ($P = 0.039$) (**Table 4, model 1**). When analysis was stratified by age, the association between ICFI and HAZ remain significant ($\beta = 0.16$, $P = 0.025$) only in older children (12–23 months) (**Supplemental Table 4**). The association between HPI and HAZ was not statistically significant, neither when all age groups were combined (**Table 4, model 2**) nor in stratified analysis (**Supplemental Table 5**). We also analysed individual components of ICFI and HPI, while controlling for variables as in ICFI or HPI models (data not shown). Among the components of ICFI, a significant and positive association was observed between breastfeeding and child HAZ ($\beta = 0.13$, $P = 0.011$). The association between dietary diversity

indicator and child HAZ also approached statistical significance ($\beta = 0.17$, $P = 0.087$). None of the components of HPI were significantly associated with HAZ.

In both model 1 and model 2, child's birth weight and maternal height were positively associated with HAZ. By contrast, child's age and sex (male), and altitude were negatively associated with HAZ. In models including all children (6–23 months), respiratory infections tended to be significant and negatively associated with HAZ, but no association was observed between diarrhoea and HAZ. When analysed by age group, we found a statistically significant and negative association between both diarrhoea and respiratory infections and HAZ only in younger children (6–11 months) (**Supplemental Tables 4 & 5**).

Discussion

Findings from the current study indicate that the burden of stunting among children aged 6–23 months in Rutsiro district is high (39%). National estimates show that the prevalence of stunting increases from 18% among children 6–8 months to 21% among those aged 9–11 months to 49% among those aged 18–23 months⁽²⁹⁾. Our results showed a similar trend, and are comparable to those observed in other studies showing that the odds of stunting increases with age⁽⁴⁸⁾.

In the present study, we found a feeding pattern that is similar to what has been observed in other studies conducted in Rwanda. For example, 39% of the mothers reported introducing complementary foods to their children before 6 months. In a cross-sectional study⁽²²⁾ and a longitudinal study⁽²³⁾ conducted in northern and southern provinces of Rwanda, respectively, researchers found that only 50% of children were exclusively breastfed for the first 6 months. These results suggest that national estimates of exclusive breastfeeding prevalence (87%)⁽²⁹⁾ may be an overestimation in rural Rwanda. In addition, the diet of children was less diverse, with majority of children (64%) consuming less than the recommended 4 food groups, and the consumption of animal sources foods was extremely

low. Other studies conducted in rural Rwanda have reported that 65–70% of children younger than 24 months do not meet the WHO recommended dietary diversity ^(22, 29). Although monotonous diets are correlated with household poverty in LMIC ⁽⁴⁹⁾, several studies conducted in Rwanda revealed other important factors underlying poor child feeding practices, including maternal beliefs and food restrictions, especially during child illness ^(18, 38).

Our finding of a positive association between ICFI and child HAZ is consistent with other studies that have investigated the relationship between a summary of child feeding practices and child HAZ in rural ^(32, 34, 50) and urban populations ^(13, 51). However, our result contrasts those of Ntab *et al.* ⁽⁵²⁾ who found no association between ICFI and child HAZ in rural Senegalese children ($n = 500$, age 12–42 months). These inconsistent results are probably due to differences in feeding practices. For example, contrary to our observation, breastfed children in the Senegalese study had lower mean HAZ than non-breastfed children because mothers prolonged breastfeeding for stunted children. Our findings of an association between ICFI and child HAZ only in older children is consistent with those of Ruel & Menon ⁽²⁵⁾ who found a positive association between ICFI and HAZ in older children (12–36 months). These authors suggested that the cumulative effect of improved care practices on child health may increase over time, and therefore, the effect of better child feeding practices may become more apparent in older children than in younger children ⁽²⁵⁾.

In the current study, two components of ICFI, i.e. breastfeeding and dietary diversity indicators, were associated with child HAZ. Other studies in Rwanda ⁽²²⁾ and Kenya ⁽⁵³⁾ have reported positive associations between breastfeeding and linear growth among children < 24 months. However, our results regarding breastfeeding should be interpreted with caution because only a small number (5%) of children were not breastfeeding. Our finding of a positive association between dietary diversity and child HAZ is also supported by findings

from several countries in Africa (including Rwanda), Asia and Latin America ^(54, 55). This result underscore the use of the dietary diversity as a better proxy of feeding practices and suggests that interventions to diversify children's diet have potential to improve child nutritional status of young children in Rwanda. However, studies found that complementary foods in Rwanda may be source of mycotoxins ^(56, 57). Consumption of mycotoxins, particularly aflatoxins, through complementary foods was found to be associated with child stunting in children (9–60 months) ⁽⁵⁸⁾. This could be one of the reasons why the relationship of ICFI with child HAZ was not as strong as might be expected, but further investigation is needed in this area. Thus, to maximize the benefits of dietary diversification on children's nutritional status, interventions should focus not only on improving the diversity of complementary foods, but also the safety of these foods.

The lack of association between HPI and child HAZ in the current study contrasts findings of Liu *et al.*, ⁽⁵⁹⁾ who reported a significant and positive association between a health practices index (measured by access to prenatal care, timing and frequency of prenatal care visits) and child HAZ in a multi-country/pooled analysis of longitudinal data from four LMICs. The discrepancy between our results and those of Liu *et al.* may be due to methodological differences. Our health index included five aspects of health practices, whereas that of Liu *et al.* included only indicators related to antenatal care. Additionally, some components (e.g. growth monitoring and antenatal care) used to construct the HPI serve as an entry point to nutrition services (e.g. counselling on IYCF practices and micronutrient supplementation), therefore, the effect of HPI in the current study on child nutritional status may be much more dependent on the quality of services provided to mothers. It was found in India and Bangladesh that growth monitoring programmes had little or no effect on child nutritional status because of failure to identify growth faltering among children or ineffective nutritional counselling to mothers ⁽⁶⁰⁾.

Research shows that health promoting practices tend to cluster. That is, a mother who, for example, exclusively breastfeeds is more likely to engage in other positive practices ⁽²⁴⁾. So, it has been suggested that summary indices may be useful to comprehensively assess the cumulative effect of various dimensions of child feeding and/or health practices on child nutritional status, while accounting for the interrelationships between these practices ⁽⁵¹⁾. Thus, conceptually, one would expect the effect size (i.e. adjusted mean HAZ difference between high and low categories) of the index on child HAZ to be higher than that of single practices. However, in the present analysis, only a few components (i.e. breastfeeding and dietary diversity) were associated with child HAZ, and the effect size of each of these components on child HAZ was comparable to that of the ICFI. Moreover, the addition or removal of the ICFI from the model did not significantly change the adjusted R-squared (data not shown). These results suggest that, at least in this sample, these indices did not perform better than individual practices. Although indices have some advantages, such as summarizing information on child feeding or health practices, they can also mask the individual practices they include ⁽³⁴⁾. In addition, while composite indices have previously been used to examine associations between feeding/health practices and child nutritional status in various settings, there still no consensus on methodology to construct these indices. Therefore, further research is needed to validate these indices and to address their limitations. Such research could, for example, examine the performance of ICFI constructed by including other aspects of child feeding such as hygiene practices during food preparations, the types or amounts of foods as opposed to food groups.

Among covariates, child age, sex (male) and altitude were negatively associated with HAZ. These factors have been found to be associated with child stunting in Rwanda ^(61, 62) and other sub-Saharan African countries ⁽⁶³⁾. In addition, maternal height and infant birth weight were positively associated with child HAZ. This relationship has been described in

previous studies showing that both maternal height and child birthweight reflect the prenatal environment and strong determinants of subsequent child nutritional status^(64, 65). The association between maternal height and child HAZ may be explained by two mechanisms: shorter mothers have small uterine volume which constrains fetal growth⁽⁶⁶⁾, and/or placental transport mechanisms that limit provision of nutrients to the fetus and growth⁽⁶⁷⁾. It is also suggested that this association could reflect shared genetic factors and/or common environmental factors that affect a mother during her early childhood and subsequently the growth of her offspring⁽⁶⁸⁾. However, studies indicated that, in early childhood, the effect of environmental factors on child linear growth is stronger than that of genetic factors, especially in poorer regions where adverse environmental conditions, such as poor quality of diet and diseases, may not permit full expression of height potential⁽⁶⁸⁻⁷⁰⁾. A significant negative association was observed between child morbidities (diarrhoea and respiratory infection) and child HAZ, but only in younger children. This finding corroborates the previous aforementioned longitudinal study⁽²³⁾ that also found a significant negative association between days with diarrhoea or respiratory infection with decreased child linear growth among children aged < 12 months.

Strengths and limitations

One of the strengths of the current study is the use of summary indices to comprehensively examine the relationship between a variety of feeding and health practices and child HAZ. There are some limitations to the current study. First, the cross-sectional nature of the current study means that our analysis cannot establish causal relationships. A longitudinal study design is recommended to somewhat address this limitation. Second, the study was conducted in one district only and was limited to the lowest socio-economic groups, which limits the generalizability of our findings. Third, most of the information on feeding and health practices was obtained from mothers' recall which is prone to recall bias and social

desirability. Fourth, although we used the standard 24 h recall method to assess child feeding practices, dietary information may not represent the usual diet of children because certain foods may be infrequently consumed. For example, the consumption of animal source foods may be rare whereas some fruits may be available only in certain seasons. Lastly, even though we controlled for several covariates, we cannot rule out the influence of unmeasured variables, such as hygiene and sanitation which are known to influence child nutritional status, particularly in resource-limited settings ⁽⁵³⁾.

Conclusion

Findings from the current study indicated that stunting among children 6–23 months in Rutsiro district remains of high public health concern, and that child HAZ is associated with both pre- and postnatal factors. The results on the relationship between ICFI, HPI and child HAZ provided a mixed picture, making the overall conclusion regarding the role of care practices less clear cut. On one hand, the finding of a positive association between ICFI and child HAZ supports our assumption that better feeding practices are positively associated with better child HAZ in this setting. On the other hand, however, the fact that only a few components of the ICFI were significantly associated with child HAZ (findings also observed in other studies), and given the lack of a significant association between HPI and child HAZ, the findings of the current study raise questions on the usefulness of the composite indices, and call for further research to validate such indices. Such research could help elucidate the strengths and weaknesses of using indices versus individual components. In the context of the current study, breastfeeding and child dietary diversity appear to have potential to improve child HAZ.

References

1. de Onis M (2006) WHO Child Growth Standards based on length/height, weight and age. *Acta Paediatr* 95, 76-85.
2. Development Initiatives (2018) 2018 Global Nutrition Report: Shining a light to spur action on nutrition. Bristol: Development Initiatives.
3. United Nations Children's Fund (UNICEF), World Health Organization International Bank for Reconstruction and Development/The World Bank (2020) Levels and trends in child malnutrition: Key Findings of the 2020 Edition of the Joint Child Malnutrition Estimates. Geneva: World Health Organization.
4. Victora CG, Adair L, Fall C *et al.* (2008) Maternal and child undernutrition: consequences for adult health and human capital. *Lancet* 371, 340-357.
5. Gluckman PD, Hanson MA Beedle AS (2007) Early life events and their consequences for later disease: a life history and evolutionary perspective. *Am J Hum Biol* 19, 1-19.
6. World Health Organization (2014) Global Nutrition Targets 2025: stunting policy brief (WHO/NMH/NHD/14.3). Geneva: World Health Organization.
7. Amugsi DA, Mittelmark MB, Lartey A *et al.* (2014) Influence of childcare practices on nutritional status of Ghanaian children: a regression analysis of the Ghana Demographic and Health Surveys. *BMJ Open* 4, e005340.
8. Kennedy E & Haddad L (1992) Food security and nutrition, 1971–91: Lessons learned and future priorities. *Food Policy* 17, 2-6.
9. Engle PL, Bentley M Pelto G (2007) The role of care in nutrition programmes: current research and a research agenda. *Proc Nutr Soc* 59, 25-35.

10. Engle PL, Lhostika L Armstrong H (1997) The Care Initiative: Assessment, Analysis and Action to Improve Care for Nutrition. New York: United Nations Children's Fund.
11. Kennedy E & Peters P (1992) Household food security and child nutrition: the interaction of income and gender of household head. *World Dev* 20, 1077-1085.
12. Brown LV, Zeitlin MF, Peterson KE *et al.* (1992) Evaluation of the impact of weaning food messages on infant feeding practices and child growth in rural Bangladesh. *Am J Clin Nutr* 56, 994-1003.
13. Ruel MT, Levin CE, Armar-Klemesu M *et al.* (1999) Good care practices can mitigate the negative effects of poverty and low maternal schooling on children's nutritional status: Evidence from Accra. *World Dev* 27, 1993-2009.
14. Zeitlin M (1991) Nutritional resilience in a hostile environment: positive deviance in child nutrition. *Nutr Rev* 49, 259-268.
15. National Institute of Statistics of Rwanda (2018) Comprehensive Food Security and Vulnerability Analysis 2018. <https://www.wfp.org/publications/rwanda-comprehensive-food-security-vulnerability-analysis-december-2018> (accessed July 2020)
16. de Onis M, Borghi E, Arimond M *et al.* (2019) Prevalence thresholds for wasting, overweight and stunting in children under 5 years. *Public Health Nutr* 22, 175-179.
17. Ministry of Health [Rwanda] & United Nations Children's Fund (UNICEF) (2014) *Knowledge, Attitudes and Practices Assessment on Early Nurturing of Children Report*. Kigali: Ministry of Health & UNICEF Rwanda.
18. Lee J, Dusingizimana T Umutoni F (2016) Understanding consumer demand for nutritious food in Nyanza district, Rwanda.

<https://www.usaid.gov/sites/default/files/documents/1860/GAIN%20Summary%20Focused%20Ethnographic%20Survey%20Report.pdf> (accessed June 2020)

19. Lung'aho M, Birachi E, Butare L *et al.* (2015) Rwanda nutrition, markets and gender analysis 2015: An integrated approach towards alleviating malnutrition among vulnerable populations in Rwanda. Nairobi: Government of Rwanda/International Center for Tropical Agriculture (CIAT).
20. Ministry of Health (2014) Rwanda National Food and Nutrition Policy. Kigali: Ministry of Health, Government of Rwanda.
21. Ngirabega JD, Leonard W, Munyanshongore C *et al.* (2010) Utilization of community based growth monitoring services by eligible children in rural Rwanda. *Rwanda Med J* 68, 40-47.
22. Uwiringiyimana V, Ocké MC, Amer S *et al.* (2019) Predictors of stunting with particular focus on complementary feeding practices: A cross-sectional study in the northern province of Rwanda. *Nutrition* 60, 11-18.
23. Matsiko E (2019) Exploring linear growth retardation in Rwandan children : Ecological and biological factors. PhD Thesis, Wageningen University.
24. Arimond M & Ruel MT (2001) *Assessing care: progress towards the measurement of selected childcare and feeding practices, and implications for programs. Discussion Paper 119*. Washington, DC: International Food Policy Research Institute.
25. Ruel MT & Menon P (2002) Child feeding practices are associated with child nutritional status in Latin America: innovative uses of the demographic and health surveys. *J Nutr* 132, 1180-1187.
26. Condo JU, Gage A, Mock N *et al.* (2015) Sex differences in nutritional status of HIV-exposed children in Rwanda: a longitudinal study. *Trop Med Int Health* 20, 17-23.

27. National Institute of Statistics of Rwanda (2012) 2012 Rwanda fourth population and housing census. District profile: Rutsiro. Kigali: National Institute of Statistics of Rwanda.
28. Cochran WG (1977) *Sampling techniques*, 3rd ed ed, *Wiley series in probability and mathematical statistics*. New York: John Wiley & Sons Inc.
29. National Institute of Statistics of Rwanda (NISR) [Rwanda], Ministry of Health (MOH) [Rwanda] ICF International (2015) Rwanda Demographic and Health Survey 2014-15. Rockville, Maryland: NISR, MOH and ICF International.
30. Dimagi CommCare. <https://www.dimagi.com/commcare/> (accessed January 2019)
31. Cogill B (2003) Anthropometric indicators measurement guide. Washington, DC: Food and Nutrition Technical Assistance (FANTA) Project, FHI 360.
32. Jones AD (2015) The production diversity of subsistence farms in the Bolivian Andes is associated with the quality of child feeding practices as measured by a validated summary feeding index. *Public Health Nutr* 18, 329-342.
33. Arimond M & Ruel M (2002) Progress in developing an infant and child feeding index: An example using the Ethiopia demographic health survey 2000. Washington DC: International Food Policy Research Institute.
34. Sawadogo PS, Martin-Prevel Y, Savy M *et al.* (2006) An infant and child feeding index is associated with the nutritional status of 6-to 23-month-old children in rural Burkina Faso. *J Nutr* 136, 656-663.
35. World Health Organization (2010) Indicators for assessing infant and young child feeding practices: part 2: Measurement. Geneva: WHO.
36. Dewey KG (2003) Guiding principles for complementary feeding of the breastfed child. Washington, DC: Pan American Health Organization/World Health Organization.

37. Brown K, Dewey K Allen L (1998) Complementary feeding of young children in developing countries: A review of current scientific knowledge. Geneva: WHO.
38. Dusingizimana T, Weber JL, Ramilan T *et al.* A qualitative analysis of infant and young child feeding practices in rural Rwanda. *Public Health Nutr.* doi: 10.1017/S1368980020001081.
39. Ballard T, Coates J, Swindale A *et al.* (2011) *Household hunger scale: indicator definition and measurement guide*. Washington DC: Food and Nutrition Technical Assistance II Project, FHI 360.
40. Vyas S & Kumaranayake L (2006) Constructing socio-economic status indices: how to use principal components analysis. *Health Policy Plan* 21, 459-468.
41. Mosites EM, Rabinowitz PM, Thumbi SM *et al.* (2015) The relationship between livestock ownership and child stunting in three countries in Eastern Africa using national survey data. *PLoS One* 10, e0136686.
42. World Health Organization (2011) WHO Anthro for personal computers, version 3.2.2: Software for assessing growth and development of the world's children. Geneva: World Health Organization.
43. World Health Organization (2006) *Multicentre Growth Reference Study Group. WHO Child Growth Standards: Length/Height-for-Age, Weight-for-Age, Weight-for-Length, Weight-for-Height and Body Mass Index-for-Age: Methods and Development*. Geneva: World Health Organization.
44. Textor J, Hardt J Knüppel S (2011) DAGitty: a graphical tool for analyzing causal diagrams. *Epidemiology* 22, 745.
45. Greenland S, Pearl J Robins JM (1999) Causal diagrams for epidemiologic research. *Epidemiology* 10, 37-48.

46. James G, Witten D, Hastie T *et al.* (2013) *An introduction to statistical learning*. vol. 112. New York: Springer.
47. World Health Organization & United Nations Children's Fund (2015) Joint Monitoring Programme (JMP) for water supply and sanitation. 2015.
48. Binagwaho A, Rukundo A, Powers S *et al.* (2020) Trends in burden and risk factors associated with childhood stunting in Rwanda from 2000 to 2015: policy and program implications. *BMC Public Health* 20, 83.
49. Onyango AW (2003) Dietary diversity, child nutrition and health in contemporary African communities. *Comp Biochem Physiol A Mol Integr Physiol* 136, 61-69.
50. Qu P, Mi B, Wang D *et al.* (2017) Association between the infant and child feeding index (ICFI) and nutritional status of 6- to 35-month-old children in rural western China. *PLoS One* 12, e0171984.
51. Moursi MM, Trèche S, Martin-Prével Y *et al.* (2008) Association of a summary index of child feeding with diet quality and growth of 6–23 months children in urban Madagascar. *Eur J Clin Nutr* 63, 718-724.
52. Ntab B, Simondon KB, Milet J *et al.* (2005) A young child feeding index is not associated with either height-for-age or height velocity in rural Senegalese children. *J Nutr* 135, 457-464.
53. Onyango AW, Esrey SA, Kramer MS (1999) Continued breastfeeding and child growth in the second year of life: a prospective cohort study in western Kenya. *The Lancet* 354, 2041-2045.
54. Arimond M & Ruel MT (2004) Dietary Diversity Is Associated with Child Nutritional Status: Evidence from 11 Demographic and Health Surveys. *J Nutr* 134, 2579-2585.

55. Menon P, Bamezai A, Subandoro A *et al.* (2015) Age-appropriate infant and young child feeding practices are associated with child nutrition in India: insights from nationally representative data. *Matern Child Nutr* 11, 73-87.
56. Grosshagauer S, Milani P, Kraemer K *et al.* (2020) Inadequacy of nutrients and contaminants found in porridge-type complementary foods in Rwanda. *Matern Child Nutr* 16, e12856.
57. Matsiko F, Kanyange C, Ingabire G *et al.* (2017) Detection and quantification of aflatoxin in cassava and maize flour sold in Kigali open markets, Rwanda. *Int Food Res J* 24, 459–464.
58. Gong Y, Hounsa A, Egal S *et al.* (2004) Postweaning exposure to aflatoxin results in impaired child growth: a longitudinal study in Benin, West Africa. *Environ Health Perspect* 112, 1334-1338.
59. Liu X, Behrman JR, Stein AD *et al.* (2017) Prenatal care and child growth and schooling in four low- and medium-income countries. *PLoS One* 12, e0171299.
60. Ashworth A, Shrimpton R, Jamil K (2008) Growth monitoring and promotion: review of evidence of impact. *Matern Child Nutr* 4 86-117.
61. Habyarimana F, Zewotir T, Ramroop S *et al.* (2016) Spatial distribution of determinants of malnutrition of children under five years in Rwanda: Simultaneous measurement of three anthropometric indices. *J Hum Ecol* 54, 138-149.
62. Uwiringiyimana V, Veldkamp A, Amer S (2019) Stunting spatial pattern in Rwanda: An examination of the demographic, socio-economic and environmental determinants. *Geospat Health* 14.
63. Wamani H, Åström AN, Peterson S *et al.* (2007) Boys are more stunted than girls in Sub-Saharan Africa: a meta-analysis of 16 demographic and health surveys. *BMC Pediatr* 7, 17.

64. Young MF, Nguyen PH, Gonzalez Casanova I *et al.* (2018) Role of maternal preconception nutrition on offspring growth and risk of stunting across the first 1000 days in Vietnam: A prospective cohort study. *PLoS One* 13, e0203201.
65. Schmidt MK, Muslimatun S, West CE *et al.* (2002) Nutritional status and linear growth of Indonesian infants in west Java are determined more by prenatal environment than by postnatal factors. *J Nutr* 132, 2202-2207.
66. Zhang X, Cnattingius S, Platt RW *et al.* (2007) Are Babies Born to Short, Primiparous, or Thin Mothers “Normally” or “Abnormally” Small? *J Pediatr* 150, 603-607.e603.
67. Price KC & Coe C (2000) Maternal constraint on fetal growth patterns in the rhesus monkey (*Macaca mulatta*): the intergenerational link between mothers and daughters. *Hum Reprod* 15 2, 452-457.
68. Hernandez-Diaz S, Peterson K, Dixit S *et al.* (1999) Association of maternal short stature with stunting in Mexican children: common genes vs common environment. *Eur J Clin Nutr* 53, 938-945.
69. Jelenkovic ASund RHur YM *et al.* (2016) Genetic and environmental influences on height from infancy to early adulthood: An individual-based pooled analysis of 45 twin cohorts. *Sci Rep* 6, 28496.
70. Liu Q, Yu C, Gao W *et al.* (2015) Genetic and Environmental Effects on Weight, Height, and BMI Under 18 Years in a Chinese Population-Based Twin Sample. *Twin Res Hum Genet* 18, 571-580.

Table 1. Child-, maternal-, and household-level characteristics, by child nutritional status¹
(Frequencies and percentages; means and standard deviations; medians and interquartile ranges (IQR))

Characteristics	Full sample		Non-stunted		Stunted		P
	n	%	n	%	n	%	
Number of children	379	100	233	61.5	146	38.5	
<i>Child-level</i>							
Age (months)							<0.001
Median	15.1		13.3		17.2		
IQR	10.9, 19.4		9.7, 18.5		13.5, 20.8		
Age group							<0.001
6–11	120	31.7	95	79.2	25	20.8	
12–23	259	68.3	138	53.3	121	46.7	
Sex							0.13
Males	184	48.5	106	57.6	78	42.4	
Females	195	51.5	127	65.1	68	34.9	
Birth weight (kg)							<0.001
Median	3.2		3.3		2.8		
IQR	3.0, 3.6		3.0, 3.6		3.0, 3.5		
HAZ							<0.001
Mean	−1.69		−0.97		−2.83		
SD	1.3		0.9		0.8		
<i>Maternal-level</i>							
Age (years)							0.02
Median	28		30		27		
IQR	24, 34		25, 35		23, 33		
Height (cm)							<0.001
Mean	156		157.3		153.8		
SD	5.9		5.5		5.7		
Marital status							0.06
Married	304	80.0	194	64.0	110	36.0	
Unmarried/widowed	75	20.0	39	52.0	36	48.0	
Number of ANC visits							0.58
< 4	248	65	150	60.0	98	40.0	
≥ 4	131	35	83	63.0	48	37.0	
Education level							0.54
None/incomplete primary	219	59.3	132	60.3	87	39.7	
Completed primary	88	23.8	53	60.2	35	39.8	
Some secondary	62	16.8	20	67.7	42	32.3	
<i>Household-level</i>							
Household size							0.22
Median	4		5		4		
IQR	3, 6		3, 6		3, 5		
Number of children < 5 years							0.42
Median	1		1		1		
IQR	1, 2		1, 2		1, 2		
Altitude (masl)							0.003
Median	1977.2		1913		2023.5		

	IQR	1729.0, 2198.0	1726.5, 2137.5	1760.3, 2229.8		
Wealth index						0.11
Lowest tertile	126	33.2	78	61.9	48	38.1
Middle tertile	126	33.2	69	54.8	57	45.2
Upper tertile	127	33.5	86	67.7	41	32.3
Hunger level						0.10
Little/no hunger	215	56.7	140	65.1	75	34.9
Moderate/severe	164	43.3	93	56.7	71	43.3
Source of drinking water [§]						0.92
Improved	222	58.6	136	58.4	86	38.7
Unimproved	157	41.4	97	41.6	60	38.2
Zone						0.19
Zone 1	75	24.4	43	22.3	32	27.8
Zone 2	81	26.3	47	24.4	34	29.6
Zone 3	152	49.4	103	53.4	49	42.6

IQR, interquartile range (1st and 3rd quartiles); HAZ, height-for-age z-score; SD, standard deviation; ANC, antenatal care; masl, metres above sea level.

¹ Data are missing for maternal education (n= 10) and maternal height (n= 36).

[§] Source of drinking water defined according to UNICEF/WHO ⁽⁴⁷⁾: improved= public taps, standpipes, tube wells, boreholes, protected well and springs, rainwater; unimproved= unprotected well and spring, surface water.

Table 2. Distribution of the ICFI and its components by child nutritional status (Frequencies and percentages; range; means and standard deviations)

	Full sample		Non-stunted		Stunted		P
	n	%	n	%	n	%	
Current breastfeeding							<0.001
Yes	362	95.5	230	63.5	132	36.5	
No	17	4.5	3	17.6	14	82.4	
Child was exclusively breastfed to 6 months							0.20
Yes	231	60.9	148	64.1	56	35.9	
No	148	39.1	85	57.4	63	42.6	
Dietary diversity score [†]							0.043
Low	28	7.4	18	64.3	10	35.7	
Medium	172	45.4	94	54.7	78	45.3	
High	179	47.2	121	67.3	58	32.4	
Meal frequency score [‡]							0.58
Low	43	11.3	28	65.1	15	34.9	
Medium	195	61.5	115	59.0	80	41.0	
High	141	37.2	90	63.8	51	36.2	
Responsive feeding when child refuses foods							0.50
Yes	262	69.1	164	62.6	98	37.4	
No	117	30.9	69	59.0	48	41.0	
Food restriction during diarrhoea							0.11
Yes	175	46.2	100	57.1	75	42.9	
No	204	53.8	133	65.2	71	34.8	
ICFI score							
Range		1–9		1–9		1–9	
Mean		5.7		6.0		5.4	<0.001
SD		1.6		1.6		1.6	
ICFI tertiles							0.001
Low (1–4)	82	21.6	41	50.0	41	50.0	
Medium (5–6)	165	43.5	95	57.6	70	42.2	
High (7–9)	132	34.8	97	73.5	35	26.5	

ICFI, infant and child feeding index; SD, standard deviation;

[†] For age 12–23 months, medium= 2–3 food groups/d and high= 4 or more food groups/d.[‡] For age 12–23 months, medium= 1–2 meals/d and high= 4+ meals/d.

Table 3. Distribution of the HPI and its components by nutritional status of children aged 6 – 23 months, Rutsiro district, Rwanda, September 2018 to January 2019.
(Frequencies and percentages; range; means and standard deviations)

	Full sample		Non-stunted		Stunted		P
	n	%	n	%	n	%	
Attendance at growth monitoring (past month)							0.67
Yes	318	83.9	197	61.9	121	38.1	
No	61	16.1	36	59.0	25	41.0	
Antenatal care							0.66
Poor (0–1)	148	39.1	85	57.4	63	42.6	
Average (2–3 visits)	231	60.9	148	64.1	56	35.9	
Good (≥ 4 visits)							
Diarrhoea treatment method							0.06
No treatment/home remedy	174	45.9	98	56.3	76	43.7	
Health center/health worker	205	54.1	135	65.9	70	34.1	
Child received vitamin A supplements (past 6 months)							0.07
Yes	362	95.5	219	60.5	143	39.5	
No	17	4.5	14	82.4	3	17.6	
Child fully immunized							0.26
Yes	357	94.2	217	60.8	140	39.2	
No	22	5.8	16	72.7	6	27.3	
HPI score							
Range		1–6		2–6		1–6	
Mean		4.5		4.5		4.5	0.58
SD		1.0		1.0		1.0	
HPI tertiles							0.043
Low (score: 1–3)	58	15.3	37	63.8	21	36.2	
Medium (score: 4)	127	33.5	67	52.8	60	47.2	
High (score: 5–6)	194	51.2	129	66.5	65	33.5	

HPI, health practices index; SD, standard deviation.

Table 4. Factors associated with HAZ of children aged 6–23 months in Rutsiro district, Rwanda, September 2018–January 2019 ¹
(Standardized coefficients and 95% confidence intervals)

	Model 1			Model 2		
	β	95% CI	P	β	95% CI	P
ICFI terciles (Ref.= Low)						
Medium	0.07	−0.13, 0.47	0.27	0.06	−0.16, 0.45	0.35
High	0.14	0.02, 0.72	0.039	0.13	−0.01, 0.72	0.06
HPI (Ref.= Low)						
Medium	–	–	–	0.03	−0.29, 0.42	0.71
High	–	–	–	0.09	−0.13, 0.57	0.22
Age (months)	−0.29	−0.10, −0.05	< 0.001	−0.30	−0.10, −0.05	< 0.001
Birth weight (kg)	0.22	0.29, 0.72	< 0.001	0.22	0.29, 0.72	< 0.001
Sex (Ref.= female)						
Male	−0.12	−0.53, −0.08	0.008	−0.13	−0.54, −0.9	0.007
Diarrhoea – yes	−0.04	0.32, 0.14	0.45	−0.03	−0.30, 0.17	0.58
Respiratory infection – yes	−0.09	−0.51, 0.04	0.09	−0.09	−0.53, 0.02	0.07
Maternal height (cm)	0.28	0.04, 0.08	<0.001	0.28	0.04, 0.08	< 0.001
Maternal education (Ref. = none/incomplete primary)						
Complete primary	0.07	−0.08, 0.49	0.16	0.08	−0.07, 0.50	0.13
Some secondary	0.08	−0.03, 0.63	0.08	0.09	−0.05, 0.61	0.10
Altitude (1000 m above sea level)	−0.15	−1.06, 0.25	0.002	−0.14	−1.00, −0.18	0.005
Household hunger (Ref. = little/no hunger)						
Mild/severe	−0.05	−0.37, 0.13	0.34	−0.05	−0.37, 0.14	0.37
Wealth index						
1 st Factor	−0.01	−0.14, 0.11	0.82	−0.02	−0.16, 0.10	0.65
2 nd Factor	−0.03	−0.16, 0.09	0.58	−0.04	−0.17, 0.08	0.49
Adjusted R ²		0.31			0.31	

HAZ, height-for-age z-score; β , standardized coefficients; CI, confidence intervals; ICFI, infant and child feeding index; Ref., reference category; HPI, health practices index.

¹ Model 1 included ICFI as the main independent variable. Model 2 included HPI as the main independent variable. Both models included the same covariates, except that model 2 also included ICFI. (–) means that HPI was not included in model 1.