UiO : **University of Oslo**

Ulrik Bo Rufus Enstad

Heisenberg modules and Balian–Low theorems

Applications of operator algebras to Gabor analysis

Thesis submitted for the degree of Philosophiae Doctor

Department of Mathematics

Faculty of Mathematics and Natural Sciences

2020

© **Ulrik Bo Rufus Enstad, 2020**

*Series of dissertations submitted to the
Faculty of Mathematics and Natural Sciences, University of Oslo
No. 2366*

ISSN 1501-7710

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

Preface

This thesis is submitted in partial fulfillment of the requirements for the degree of *Philosophiae Doctor* at the University of Oslo. The research presented here was conducted under the supervision of professor Nadia Larsen and professor Franz Luef.

The thesis is a collection of four papers, presented in chronological order of writing. The papers are preceded by an introductory chapter that relates them together and provides background information and motivation for the work.

Acknowledgements

I would like to begin by thanking my supervisor Nadia Larsen. She has been immensely supportive and encouraging during my time as a PhD student. I want to thank her for all our frequent, inspiring discussions, for encouraging me to go to numerous conferences and for making it possible for me to stay abroad in Boulder.

I am grateful to my co-supervisor Franz Luef for his support and all the insightful discussions we have had in Trondheim and elsewhere. I want to thank him for introducing me to time-frequency analysis during my master's studies, when some of the ideas in this thesis began to develop.

I want to express my gratitude to Judith Packer for her hospitality and for all the enlightening discussions we had during my stay in Boulder. I also want to thank Shen Lu, Alex Munday, Andrew Stocker, Colton Grainger and of course the Ekrems for making the stay so pleasant.

Everyday life as a PhD student at UiO would not have been the same without the Operator Algebras group. I want to thank everyone who has been in the group during my time at UiO for maintaining such a good atmosphere at work during our daily lunch and coffee breaks.

I also want to thank Petter Nyland for being a great travel buddy, my co-authors Are Austad, Mads Jakobsen and Tron Omland for rewarding collaborations, and Martin Helsø for his assistance with L^AT_EX.

I want to thank my parents Christine and Syver and my siblings Tita and Truls for being an awesome family. Finally, I want to thank Luca for always being there for me and being the best.

• **Ulrik Bo Rufus Enstad**
Oslo, May 2020

List of Papers

Paper I

Ulrik Enstad, Mads S. Jakobsen and Franz Luef “Time-frequency analysis on the adeles over the rationals”. In: *Comptes Rendus Mathématique*. Vol. 357, no. 2 (2019), pp. 188–199. DOI: 10.1016/j.crma.2018.12.004.

Paper II

Are Austad and Ulrik Enstad “Heisenberg modules as function spaces”. In: *Journal of Fourier Analysis and Applications*. Vol. 26, no. 2 (2020). DOI: 10.1007/s00041-020-09729-7.

Paper III

Ulrik Enstad “The Balian–Low theorem for locally compact abelian groups and vector bundles”. In: *Journal de Mathématiques Pures et Appliquées*. (In press) (2019) DOI: 10.1016/j.matpur.2019.12.005.

Paper IV

Ulrik Enstad, Mads S. Jakobsen, Franz Luef and Tron Omland “Deformations of Gabor frames on the adeles and other locally compact abelian groups”. Preprint, arXiv:2001.07080.

Contents

Preface	i
List of Papers	iii
Contents	v
1 Introduction	1
1.1 From the uncertainty principle to the Balian–Low theorem	1
1.2 Time-frequency analysis on locally compact abelian groups	5
1.3 Operator algebras	10
1.4 Twisted group C^* -algebras and noncommutative tori . . .	11
1.5 Vector bundles and projective modules	13
1.6 Heisenberg modules	16
1.7 Summary of Papers	21
References	22
Papers	28
I Time-frequency analysis on the adeles over the rationals	29
I.1 Introduction	29
I.2 The p -adic numbers and adeles over the rationals	30
I.3 Feichtinger’s algebra for p -adic groups and the adeles . . .	34
I.4 Gabor frames	37
References	45
II Heisenberg modules as function spaces	51
II.1 Introduction	51
II.2 Preliminaries	55
II.3 Results	62
References	77
III The Balian–Low theorem for locally compact abelian groups and vector bundles	81
III.1 Introduction	81
III.2 Hilbert C^* -modules and their frames	85
III.3 Time-frequency analysis and Heisenberg modules	91
III.4 Vector bundles and quasiperiodic functions	95
III.5 Connecting Heisenberg modules and vector bundles	100
III.6 The group $\mathbb{R} \times \mathbb{Q}_p$	106
III.A Fundamental domains	113

III.B	Cutoff functions in the Feichtinger algebra	115
	References	118
IV	Deformations of Gabor frames on the adeles and other locally compact abelian groups	123
IV.1	Introduction	123
IV.2	Locally compact abelian groups and their automorphisms .	129
IV.3	Time-frequency analysis	135
IV.4	Deformation results	139
IV.5	The Balian–Low theorem	146
IV.6	Gabor analysis on the adeles of a global field	151
	References	163

Chapter 1

Introduction

The scope of this thesis lies in the interplay between the fields of operator algebras and time-frequency analysis. The main theme is the Balian–Low theorem (or lack thereof) on locally compact abelian groups, and its connection to Heisenberg modules. In particular, we initiate time-frequency analysis on the group of adeles over the rationals and show that the Balian–Low theorem holds in this setting. All the papers in the thesis are related to this main theme.

The classical Balian–Low theorem in time-frequency analysis on the real line can be seen as a manifestation of the uncertainty principle. In the following introduction, we begin with a survey of the uncertainty principle and time-frequency analysis, ending at the Balian–Low theorem. We then introduce C^* -algebras and their Hilbert modules, in particular Heisenberg modules, and explain the link to time-frequency analysis. Finally, we formulate some of the main achievements based on this link, and we outline how the link will be exploited in the papers that constitute the thesis.

1.1 From the uncertainty principle to the Balian–Low theorem

The uncertainty principle is one of the fundamental principles of quantum mechanics, discovered by Werner Heisenberg in 1927 [Hei27]. It asserts that one cannot determine the position and momentum of a particle simultaneously up to arbitrary precision: The more precisely one knows the position, the less precisely one can know the momentum, and vice versa.

Matrix mechanics, the first mathematically rigorous formulation of quantum mechanics, was described in a series of papers by Heisenberg, Born and Jordan [BHJ26; BJ25; Hei25]. In their formalism, the position and momentum of a particle moving along a line were represented by infinite matrices X and P satisfying the commutation relation

$$[X, P] = XP - PX = i\hbar I \tag{1.1}$$

where $\hbar$ denotes the reduced Planck constant and I is the identity matrix. A year later, Schrödinger published four papers (see e.g. [Sch26]) on what became known as the wave mechanics formalism of quantum mechanics. Wave mechanics was formulated in terms of wave functions of real variables that acted as probability densities associated to states. It was von Neumann who gave the first mathematically rigorous proof that the matrix mechanics of Heisenberg, Born and Jordan and the wave mechanics of Schrödinger are equivalent: In a series of papers published in 1927 (e.g. [vNe27]), he invented the concept of a

1. Introduction

Hilbert space, and showed that matrix mechanics and wave mechanics can be formulated in the context of the Hilbert spaces $\ell^2(\mathbb{N})$ and $L^2(\mathbb{R})$, respectively.

Using von Neumann's formalism, the position X and the momentum P of a particle moving along a line can be represented by the unbounded self-adjoint operators on $L^2(\mathbb{R})$ given by

$$(Xf)(t) = tf(t), \quad (Pf)(t) = -i\hbar f'(t) \quad (1.2)$$

for $f \in L^2(\mathbb{R})$. These two operators satisfy Heisenberg's commutation relation (1.1). One defines the expectation $\langle X \rangle_\psi$ and uncertainty $\Delta_\psi X$ of an observable X in the state $\psi \in L^2(\mathbb{R})$ via the following formulas:

$$\begin{aligned} \langle X \rangle_\psi &= \langle X\psi, \psi \rangle, \\ \Delta_\psi X &= \langle (X - \langle X \rangle_\psi I)^2 \rangle_\psi^{1/2}. \end{aligned}$$

A quantitative version of the uncertainty principle proved by Earle Hesse Kennard [Ken27] in 1927 then states that

$$(\Delta_\psi X)(\Delta_\psi P) \geq \frac{\hbar}{2}. \quad (1.3)$$

The inequality (1.3) can be derived from (1.1) using the Cauchy–Schwarz inequality.

Mathematically, the uncertainty principle can be seen as a statement about the Fourier transform on the real line. Since the momentum operator is the position operator conjugated by the Fourier transform, one can write (1.3) as

$$\left(\int_{\mathbb{R}} (t-a)^2 |f(t)|^2 dt \right)^{1/2} \left(\int_{\mathbb{R}} (\xi-b)^2 |\widehat{f}(\xi)|^2 d\xi \right)^{1/2} \geq \frac{\|f\|_2^2}{4\pi} \quad (1.4)$$

for any $f \in L^2(\mathbb{R})$ and any $a, b \in \mathbb{R}$. This expresses the fact that a signal represented by the function f cannot be arbitrarily well-localized in both time and frequency.

In signal and harmonic analysis, time-frequency analysis comprises those techniques that study signals simultaneously in their time and frequency domains. More mathematically, it is the field of harmonic analysis where functions and operators are studied using the structure of time-frequency shifts. It is therefore to be expected that the uncertainty principle manifests itself in the field. Time-frequency analysis has its origins in quantum mechanics, e.g. in the work of Wigner [Wig32].

The basic operation in time-frequency analysis is the time-frequency shift. The time shift operator T_x and frequency shift M_ω ($x, \omega \in \mathbb{R}$) are operators on $L^2(\mathbb{R})$ given by

$$(T_x f)(t) = f(t-x), \quad (M_\omega f)(t) = e^{2\pi i \omega t} f(t) \quad (1.5)$$

for $f \in L^2(\mathbb{R})$, $t \in \mathbb{R}$. They satisfy the commutation relation

$$M_\omega T_x = e^{2\pi i x \omega} T_x M_\omega, \quad x, \omega \in \mathbb{R}. \quad (1.6)$$

A time-frequency shift is defined by $\pi(x, \omega) = M_\omega T_x$ for $x, \omega \in \mathbb{R}$. Using time-frequency shifts, one can define the *short-time Fourier transform*. The exposition that follows is inspired by Gröchenig’s book [Grö01]. The idea behind the short-time Fourier transform is that one wants to find the frequency content of a signal $f \in L^2(\mathbb{R})$ that varies over time (such as a piece of music) around a certain time x . If one takes the usual Fourier transform of the whole signal, this gives the frequency content of the whole signal rather than at the specific time x . To circumvent this, one picks a function g localized around 0, multiplies f by $T_x g$, the shift of g by x , and takes the Fourier transform of the resulting product. Formally, the short-time Fourier transform of f with respect to the window g is the function $V_g f: \mathbb{R}^2 \rightarrow \mathbb{C}$ given by

$$V_g f(x, \omega) = \widehat{f T_x g}(\omega) = \int_{\mathbb{R}} f(t) e^{-2\pi i \omega t} \overline{g(t-x)} dt = \langle f, \pi(x, \omega) g \rangle \quad (1.7)$$

for $(x, \omega) \in \mathbb{R}^2$. Similarly to the ordinary Fourier transform, the short-time Fourier transform extends to an isometry $V_g: L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}^2)$ for nonzero $g \in L^2(\mathbb{R})$. The short-time Fourier transform also has orthogonality relations and a reconstruction formula. In particular, one can write any $f \in L^2(\mathbb{R})$ as

$$f = \int_{\mathbb{R}^2} V_g f(z) \pi(z) h dz \quad (1.8)$$

for any $g, h \in L^2(\mathbb{R})$ with $\langle g, h \rangle = 1$, where the integral is understood in a weak sense.

The properties of the short-time Fourier transform V_g are reflected by the chosen window g . Thinking of f as a piece of music, one can reason as follows: For $V_g f$ to accurately measure the starting point and duration of each of the sounds that make up f , the window g has to be well-localized. Similarly, for $V_g f$ to accurately measure the pitch of each sound of f , $\widehat{g}$ has to be well-localized. But as we have seen, the uncertainty principle implies that g cannot be arbitrarily well-localized both in time and frequency. Therefore, one typically picks a window g for which both g and $\widehat{g}$ have a certain degree of localization, such as the Gaussian.

In the recovery formula (1.8), one needs to know all the values $V_g f(z)$, $z \in \mathbb{R}^2$, in order to reconstruct f . It would be more feasible if one could reconstruct f from a set of samples of $V_g f$. Thus, the question arises whether one can obtain a discretization of (1.8) to a lattice Δ in $\mathbb{R}^2$, say $\Delta = \alpha\mathbb{Z} \times \beta\mathbb{Z}$ for $\alpha, \beta > 0$. This would give us expressions of the form

$$f = \sum_{z \in \Delta} \langle f, \pi(z) g \rangle \pi(z) h \quad (1.9)$$

for every $f \in L^2(\mathbb{R})$. In the language of frame theory, (1.9) holds for every $f \in L^2(\mathbb{R})$ if and only if the sequence of time-frequency translates of g over Δ

$$\mathcal{G}(g, \Delta) = \{\pi(z) g : z \in \Delta\} \quad (1.10)$$

1. Introduction

is a *frame* for the Hilbert space $L^2(\mathbb{R})$. Abstractly, a sequence $(e_j)_{j \in J}$ in a Hilbert space H is called a *frame* if there exist constants $A, B > 0$ such that

$$A\|f\|^2 \leq \sum_{j \in J} |\langle f, e_j \rangle|^2 \leq B\|f\|^2 \quad (1.11)$$

for every $f \in H$. The frame property (1.11) for $(e_j)_j$ is equivalent to the existence of a *dual frame* $(f_j)_{j \in J}$, which is a sequence for which an analogue (1.9) holds for all $f \in H$.

The set $\mathcal{G}(g, \Delta)$ is called the *Gabor system* over Δ generated by the *window* g , and we call it a *Gabor frame* if it forms a frame for $L^2(\mathbb{R})$. As such, a Gabor frame is precisely what one gets after a successful discretization of the short-time Fourier transform. Gabor systems have their name from the Hungarian physicist and electrical engineer Dennis Gabor, who won the Nobel prize in physics for his invention of holography. His 1946 paper “Theory of communication” is highly influential and one of the starting points of time-frequency analysis. Here, one finds the (not entirely mathematically precise) claim that one can write every function $f \in L^2(\mathbb{R})$ in the form

$$f(t) = 2^{1/4} \sum_{k, l \in \mathbb{Z}} c_{k, l} e^{2\pi i l t} e^{-\pi(t-k)^2} \quad (1.12)$$

for suitable scalar coefficients $c_{k, l} \in \mathbb{C}$. In the language of time-frequency analysis, Gabor’s claim concerns the completeness properties of the Gabor system $\mathcal{G}(\phi, \mathbb{Z} \times \mathbb{Z})$, where ϕ denotes the Gaussian $\phi(t) = 2^{1/4} e^{-\pi t^2}$. A. J. E. M. Janssen, one of the pioneers of mathematical time-frequency analysis, investigated Gabor’s claim in [Jan81]. He showed that one can find convergent expansions as in (1.12) for every $f \in L^2(\mathbb{R})$, but only in the sense of tempered distributions and not in the L^2 -norm. In particular, $\mathcal{G}(\phi, \mathbb{Z} \times \mathbb{Z})$ is not a Gabor frame!

The fact that the Gaussian does not give a Gabor frame over the lattice $\mathbb{Z} \times \mathbb{Z}$ can be understood as a consequence of the *Balian–Low theorem*, one of the manifestations of the uncertainty principle in time-frequency analysis. The Balian–Low theorem is an improvement of another theorem, the *density theorem*, which we will now explain: Given the lattice $\alpha\mathbb{Z} \times \beta\mathbb{Z}$, the product $\alpha\beta$ is called the *volume* of the lattice and measures how far apart the lattice points are spread in $\mathbb{R}^2$. The smaller the volume is, the more densely packed in $\mathbb{R}^2$ the lattice is. The density theorem states that for every $g \in L^2(\mathbb{R})$, for $\mathcal{G}(g, \Delta)$ to be a Gabor frame, it is necessary that the volume of Δ is less than or equal to 1. This reflects the intuition that in order to recover a signal from discrete samples of its short-time Fourier transform, one cannot sample too sparsely in the time-frequency plane. The Balian–Low theorem is a strengthening of the density theorem for well-localized functions, and comes in many variants. The first version that was formulated independently by Roger Balian [Bal81] and Francis Low [Low85] states the following:

Theorem 1.1.1. *Suppose $g \in L^2(\mathbb{R})$ is such that $\mathcal{G}(g, \alpha\mathbb{Z} \times \beta\mathbb{Z})$ is a Gabor frame and both of the integrals*

$$\int_{\mathbb{R}} t^2 |g(t)|^2 dt \quad \text{and} \quad \int_{\mathbb{R}} \omega^2 |\widehat{g}(\omega)|^2 d\omega \quad (1.13)$$

converge. Then $\alpha\beta < 1$.

The proofs of Balian and Low both employed the Zak transform, but contained a technical gap that was later closed by Daubechies in [Dau90]. The first proof without gaps was given by Guy Battle in [Bat88] and is based directly on the Heisenberg commutation relation (1.1), which shows that the Balian–Low theorem is intimately related to the uncertainty principle. Indeed, the requirement in Theorem 1.1.1 is equivalent to g being in the domain of both the position and momentum operators from quantum mechanics.

There are several variations of the Balian–Low with different hypotheses on the window g but with the same conclusion. One of them involves a function space in time-frequency analysis called the *Feichtinger algebra* $S_0(\mathbb{R})$ [Fei81]. It consists of those functions $f \in L^2(\mathbb{R})$ with absolutely integrable short-time Fourier transform:

$$S_0(\mathbb{R}) = \{f \in L^2(\mathbb{R}) : \int_{\mathbb{R}^2} |V_f f(z)| dz < \infty\}.$$

The function space $S_0(\mathbb{R})$ is an example of a modulation space, and it is a Fourier invariant space of continuous functions containing the Schwartz space. It can be shown that if $g \in S_0(\mathbb{R})$, then the integrals in (1.13) converge. Consequently, we get the following weaker version of Theorem 1.1.1:

Theorem 1.1.2. *Suppose $\mathcal{G}(g, \alpha\mathbb{Z} \times \beta\mathbb{Z})$ is a Gabor frame and $g \in S_0(\mathbb{R})$. Then $\alpha\beta < 1$.*

The advantage with the above variant of the Balian–Low theorem is that its statement can be easily generalized to the setting of a locally compact abelian group, which we will now detail.

1.2 Time-frequency analysis on locally compact abelian groups

Time-frequency analysis can be readily done on a locally compact abelian (LCA) group G . We write the group operation multiplicatively, and denote by $\widehat{G}$ the Pontryagin dual of G . In this setting, the time and frequency shift operators are defined as follows: Given $x \in G$ and $\omega \in \widehat{G}$,

$$(T_x \xi)(t) = \xi(x^{-1}t), \quad (M_\omega \xi)(t) = \omega(t)\xi(t) \quad (1.14)$$

for $\xi \in L^2(G)$, $t \in G$. One then sets

$$\pi(x, \omega) = M_\omega T_x, \quad (x, \omega) \in G \times \widehat{G}. \quad (1.15)$$

1. Introduction

Thus, in the locally compact abelian case, $\mathbb{R}$ is interchanged with G and $\mathbb{R}^2 \cong \mathbb{R} \times \widehat{\mathbb{R}}$ is interchanged with $G \times \widehat{G}$. We have the following commutation relation

$$M_\omega T_x = \omega(x) T_x M_\omega, \quad (x, \omega) \in G \times \widehat{G}$$

from which one can deduce

$$\pi(x, \omega)\pi(y, \tau) = \overline{\tau(x)}\pi(xy, \omega\tau), \quad (x, \omega), (y, \tau) \in G \times \widehat{G}. \quad (1.16)$$

Recall that a (strongly continuous) unitary representation of a group G on a Hilbert space H is a homomorphism $G \rightarrow \mathcal{U}(H)$, where $\mathcal{U}$ denotes the unitary operators on H , such that $x \mapsto U(x)\xi$ is continuous for every $\xi \in H$. The maps $x \mapsto T_x$ and $\omega \mapsto M_\omega$ are both unitary representations of G and $\widehat{G}$ on $L^2(G)$ respectively. However, because of the relation (1.16), $\pi: G \times \widehat{G} \rightarrow \mathcal{U}(H)$ is not a unitary representation on the nose, but rather a *projective* unitary representation.

In general, a projective representation of a group G on a Hilbert space H is a strongly continuous map $U: G \rightarrow \mathcal{U}(H)$ such that

$$U(x)U(y) = \sigma(x, y)U(xy) \quad (1.17)$$

for some complex scalar $\sigma(x, y)$ of modulus 1 and all $x, y \in G$. We also require $U(1) = I$. It can be shown from these conditions that σ must satisfy the following relations:

$$\sigma(x, y)\sigma(xy, z) = \sigma(x, yz)\sigma(y, z), \quad x, y, z \in G, \quad (1.18)$$

$$\sigma(1, 1) = 1. \quad (1.19)$$

The function $\sigma: G \times G \rightarrow \mathbb{T}$ will in general only be measurable and is called a 2-cocycle. The representation U is called σ -projective.

When one views the time-frequency shifts π as a projective representation of $G \times \widehat{G}$, it is often called the *Weyl–Heisenberg representation*. It can alternatively be viewed as an ordinary representation of an associated Heisenberg group [Grö01], although this is a viewpoint we will not adopt here. The 2-cocycle c associated to the Weyl–Heisenberg representation is continuous and is given by

$$c((x, \omega), (y, \tau)) = \overline{\tau(x)}, \quad (x, \omega), (y, \tau) \in G \times \widehat{G}.$$

Gabor systems over lattices Δ in $G \times \widehat{G}$ with windows $g \in L^2(G)$ are defined just as in the real case:

$$\mathcal{G}(g, \Delta) = \{\pi(z)g : z \in \Delta\}.$$

Note that this is the orbit of g under the representation π . The system is called a Gabor frame if it is a frame for $L^2(G)$, i.e. there exist $C, D > 0$ such that

$$C\|f\|_2^2 \leq \sum_{z \in \Delta} |\langle f, \pi(z)g \rangle|^2 \leq D\|f\|_2^2$$

for all $f \in L^2(G)$. If $\mathcal{G}(g, \Delta)$ is a Gabor frame, then $\overline{\text{span} \mathcal{G}(g, \Delta)} = L^2(G)$. Or, in terms of representation theory, g is a cyclic vector for the restricted representation $\pi|_{\Delta}: \Delta \rightarrow \mathcal{U}(L^2(G))$. In that sense, finding Gabor frames over a lattice Δ can be thought of as finding “well-behaved” cyclic vectors for the representation $\pi|_{\Delta}$, or *frame vectors* as they have been called in the literature [HL00].

A vector f in $L^2(G)$ is called integrable with respect to π if its matrix coefficient is absolutely integrable, i.e.

$$\int_{G \times \widehat{G}} |\langle f, \pi(z)f \rangle| dz < \infty.$$

Thus, the Feichtinger algebra $S_0(G)$ consists precisely of the integrable vectors of π . By fixing any $g \in S_0(G)$, one defines a norm on $S_0(G)$ via

$$\|f\|_{S_0(G),g} = \int_{G \times \widehat{G}} |\langle f, \pi(z)g \rangle| dz$$

for $f \in S_0(G)$. The Feichtinger algebra is a Banach algebra with respect to this norm and both pointwise multiplication or convolution, and choosing a different g yields an equivalent norm [Jak18].

Many theorems of Gabor analysis on the real line, such as the density theorem [JL16], carry over to the locally compact abelian setting:

Theorem 1.2.1. *Let Δ be a lattice in $G \times \widehat{G}$. If there exists $g \in L^2(G)$ such that $\mathcal{G}(g, \Delta)$ is a Gabor frame, then $\text{vol}(\Delta) \leq 1$.*

Let us discuss Gabor analysis on concrete groups other than $\mathbb{R}$ and $\mathbb{R}^n$. The groups $\mathbb{Z}$, $\mathbb{T}$ and $\mathbb{Z}/(m)$ appear naturally when sampling or periodizing Gabor frames on $\mathbb{R}$ [Jan97; JL19]. In general, it follows from the structure theory of locally compact abelian groups [HR63] that a LCA group G can be written as

$$G \cong \mathbb{R}^n \times H$$

where H is a LCA group containing a compact open subgroup. Gröchenig shows in [Grö98] that Gabor analysis on groups containing a compact open subgroup is very different from that of $\mathbb{R}^n$. For instance, there is no Balian–Low theorem for the Feichtinger algebra on discrete or compact groups. This is intuitive for e.g. compact groups, as the notion of good localization in time using integrals is trivially satisfied because the Haar measure is finite. Similarly, on discrete groups, good localization in frequency becomes trivially satisfied since the dual group is compact. As an explicit example, let $G = \mathbb{Z}$, let $\Delta = \mathbb{Z} \times \{0\} \subseteq \mathbb{Z} \times \mathbb{T} \cong \mathbb{Z} \times \widehat{\mathbb{Z}}$ and let $g = \delta_0$. Then $g \in S_0(\mathbb{Z}) = \ell^1(\mathbb{Z})$, and $\mathcal{G}(g, \Delta)$ is the standard orthonormal basis $\{\delta_k : k \in \mathbb{Z}\}$ for $\ell^2(\mathbb{Z})$.

Also, groups containing a compact open subgroup may fail to have lattices: For instance, the group $\mathbb{Q}_p$ of p -adic numbers has a compact open subgroup, namely the p -adic integers $\mathbb{Z}_p$. However, the only discrete subgroup of $\mathbb{Q}_p$ is the trivial one, so in particular, $\mathbb{Q}_p$ admits no lattices.

1. Introduction

This does not mean products $\mathbb{R}^n \times H$ where H contains a compact open subgroup are not interesting. For instance, the group $\mathbb{R} \times \mathbb{Q}_p$ contains in a nontrivial way the subgroup $\mathbb{Z}[1/p] = \{a/p^k : a, k \in \mathbb{Z}\}$ as a lattice, namely through the embedding $q \mapsto (q, q) \in \mathbb{R} \times \mathbb{Q}_p$. The group $\mathbb{R} \times \mathbb{Q}_p$ can be viewed as a truncated version of the *adeles*, a locally compact ring appearing in algebraic number theory [RV99]. In general, one can associate an adele group to any global field. For the rational numbers, the associated adele group is the restricted product

$$\mathbb{A}_{\mathbb{Q}} = \prod_p^{\mathbb{Z}_p} \mathbb{Q}_p = \left\{ (x_p)_p \in \prod_p \mathbb{Q}_p : x_p \in \mathbb{Z}_p \text{ for all but finitely many } p \right\} \quad (1.20)$$

where the product is taken over all primes p . The rational numbers $\mathbb{Q}$ embed as a lattice into the adeles via the map $q \mapsto (q, q, \dots)$. In Paper I, we initiate Gabor analysis on both $\mathbb{R} \times \mathbb{Q}_p$ and the adele group, and show that there are natural examples of Gabor frames over variations of the above lattices embedded into the time-frequency plane. We also give an explicit description of the Feichtinger algebra $S_0(\mathbb{A}_{\mathbb{Q}})$ in this setting. In Paper IV, we introduce the S -adeles $\mathbb{A}_{\mathbb{Q}, S}$, which generalize both $\mathbb{R} \times \mathbb{Q}_p$ and $\mathbb{A}_{\mathbb{Q}}$. Here, one takes the restricted product in (1.20), but over a fixed set S of primes. Similarly to $\mathbb{R} \times \mathbb{Q}_p$ and $\mathbb{A}_{\mathbb{Q}}$, the ring $\mathbb{Z}[S^{-1}]$ embeds as a lattice into $\mathbb{A}_{\mathbb{Q}, S}$ via the diagonal map.

Returning to the Balian–Low theorem, we can now ask whether there is an analogue for the adeles. After all, the group of adeles is neither a compact nor discrete group, and contains $\mathbb{R}$ as a subgroup. More generally we can ask, using the Feichtinger algebra, for which LCA groups there is a Balian–Low theorem. That is, solve the following problem:

Problem 1.2.2. *Determine which locally compact abelian groups G have the following property: For every lattice Δ in $G \times \widehat{G}$, the existence of a Gabor frame $\mathcal{G}(g, \Delta)$ with $g \in S_0(G)$ implies that the volume of Δ (i.e. the measure of a fundamental domain of Δ in $G \times \widehat{G}$) is strictly less than one.*

One can get a more specific instance of Problem 1.2.2 by considering lattices in $G \times \widehat{G}$ of the form $\Delta = \Lambda \times \Lambda^{\perp}$ for Λ a lattice in G . Here, the *annihilator* $\Lambda^{\perp}$ of Λ is the group of all characters of G which are constantly equal to 1 on Λ . Because of the formula $\text{vol}(\Lambda) \text{vol}(\Lambda^{\perp}) = 1$, such lattices in $G \times \widehat{G}$ always have volume 1. We then get the following:

Problem 1.2.3. *Determine which locally compact abelian groups G have the following property: For every lattice Λ in G , there are no Gabor frames $\mathcal{G}(g, \Lambda \times \Lambda^{\perp})$ with $g \in S_0(G)$.*

The advantage of working with so-called critical lattices $\Delta = \Lambda \times \Lambda^{\perp}$ as in Problem 1.2.3 is that one can employ the Zak transform for locally compact abelian groups. This transform is used in time-frequency analysis to diagonalize the frame operator corresponding to Gabor systems with rational density. However, it was first introduced by Gelfand [Gel50] for a problem in

differential equations, and was later generalized by Weil to locally compact abelian groups [Wei64]. We will introduce the Zak transform in Paper III.

The Zak transform on $\mathbb{R}$ was used by Balan in [Bal01] to show that the Balian–Low theorem for $S_0(\mathbb{R})$ can be seen as a consequence of the nontriviality of a certain vector bundle of rank 1 over $\mathbb{T}^2$. What the Zak transform does is that it realizes windows of Gabor systems as continuous sections of this vector bundle. Consequently, the sections must have a zero, which implies that the initial window cannot give a Gabor frame. A natural question is whether there is an analogous vector bundle in the setting of Problem 1.2.3, and whether its nontriviality determines whether Problem 1.2.3 holds for the given group. The goal of Paper III is to show that this is the case, and furthermore that these vector bundles show up naturally as Heisenberg modules. We also prove that there is a Balian–Low theorem for the group $\mathbb{R} \times \mathbb{Q}_p$ in Paper III by exhibiting the nontriviality of the associated vector bundle.

The Zak transform on locally compact abelian groups was used by Kaniuth and Kutyniok [KK98] to study Problem 1.2.3, where they showed the following:

Theorem 1.2.4. *Let G be a second countable, compactly generated LCA group with noncompact identity component. Then for every $g \in S_0(G)$ and every lattice Λ in G , $\mathcal{G}(g, \Lambda \times \Lambda^\perp)$ is not a Gabor frame.*

The assumption that G is second countable is technical and is used to guarantee the surjectivity of the Zak transform. The assumption that G is compactly generated is more essential for their proof, because they use structure theory to decompose their group as $G = \mathbb{R}^a \times \mathbb{Z}^b \times K$, where K is a compact abelian group. Among the assumptions in Theorem 1.2.4, the only assumption that obviously cannot be removed is the condition that the identity component of G is noncompact. This is exactly what excludes e.g. compact or discrete groups, where Theorem 1.2.4 does not hold.

The question then becomes whether one can remove the assumptions that G is second countable and compactly generated from Theorem 1.2.4. In other words, whether noncompactness of the identity component characterizes groups that have a Balian–Low theorem for lattices of the form $\Lambda \times \Lambda^\perp$. In fact, one might even ask whether the more general Problem 1.2.2 holds for exactly those groups with noncompact identity component. Since the group of S -adeles has noncompact identity component, namely $\mathbb{R}$, this group would then satisfy a Balian–Low theorem for the Feichtinger algebra. Investigating this matter is one of the goals of Paper IV, where we characterize LCA groups G for which Problem 1.2.2 holds as exactly the ones with noncompact identity component. The proof that noncompactness of the identity component of G implies that there is a Balian–Low theorem is based on a technique in [AFK14]. Here they use deformation theory for Gabor frames on $L^2(\mathbb{R}^n)$ [FK04] to arrive at a contradiction: If $\mathcal{G}(g, \Delta)$ is a Gabor frame with $g \in S_0(\mathbb{R}^n)$ and $\text{vol}(\Delta) = 1$, then the lattice can be deformed slightly so that $\text{vol}(\Delta) > 1$ and the frame property is preserved. This contradicts the density theorem Theorem 1.2.1, so we conclude that $\mathcal{G}(g, \Delta)$ is not a Gabor frame. Note that the assumption that $g \in S_0(\mathbb{R}^n)$ is essential for such deformation results. Using the Braconnier topology, we prove

a similar deformation theory for Gabor frames on LCA groups, and show that it implies the validity of Problem 1.2.2 for LCA groups with noncompact identity component.

Conversely, if G has compact identity component, then there exists for every lattice Λ in G a Gabor orthonormal basis of the form $\mathcal{G}(g, \Lambda \times \Lambda^\perp)$ for some $g \in S_0(G)$. We prove this in Paper IV, and the proof relies on results from the theory of totally disconnected groups.

1.3 Operator algebras

The field of operator algebras has its origin in the work of Murray, von Neumann, Gelfand and Naimark. In his work on the mathematical foundations of quantum mechanics, von Neumann gave the formal definition of an abstract Hilbert space. This set the stage for a series of papers written by Murray and von Neumann in the 1930's on "rings of operators" [MN36], which today are known as *von Neumann algebras*. The most basic example of a von Neumann algebra is the algebra $\mathcal{B}(H)$ of bounded linear operators on a Hilbert space H . This algebra is equipped with an involution given by taking the adjoint of an operator. More generally, a von Neumann algebra on H is a subalgebra of $\mathcal{B}(H)$ that is closed under taking adjoints (a $*$ -algebra) and closed under taking weak limits of operators. The commutant of a von Neumann algebra on $A \subseteq \mathcal{B}(H)$ is the set A' of all operators that commutes with every operator in A . Murray and von Neumann proved that a von Neumann algebra A can alternatively be described as a $*$ -algebra of operators on a Hilbert space which is equal to its own double commutant, i.e. $A = (A')'$. This theorem became known as the bicommutant theorem, and is one of the cornerstones of the theory of von Neumann algebras.

In 1943, the Soviet mathematicians Gelfand and Naimark wrote a paper [GN43] on a certain class of Banach algebras A equipped with an involution $a \mapsto a^*$ satisfying the identity

$$\|a^*a\| = \|a\|^2$$

for all $a \in A$. This class of Banach $*$ -algebras are today known as (unital) C^* -algebras, and the above identity is known as the C^* -identity. Gelfand and Naimark proved two fundamental results about C^* -algebras.

Theorem 1.3.1. *Every commutative unital C^* -algebra A is isomorphic to the C^* -algebra $C(X)$ of continuous functions on a (canonically determined) compact Hausdorff space X :*

$$A \cong C(X).$$

In fact, the functor sending a space X to its C^ -algebra of continuous functions is a contravariant equivalence of categories between the category of compact Hausdorff spaces and the category of commutative unital C^* -algebras.*

Theorem 1.3.2. *Every C^* -algebra A is isomorphic to a C^* -subalgebra of $\mathcal{B}(H)$ for some Hilbert space H . In other words, there exists a faithful representation*

$$\pi: A \rightarrow \mathcal{B}(H).$$

Both of these theorems have tremendous technical consequences that makes the theory of C^* -algebras rich. A consequence of (1.3.1) is the *continuous functional calculus*, which allows one to prove e.g. that positive elements of C^* -algebras always have square roots. This is done by looking at the C^* -subalgebra generated by the positive element a , an algebra which necessarily must be commutative, and then viewing a as a continuous function on its spectrum using Theorem 1.3.2. More generally, non-unital C^* -algebras may be realized as continuous functions vanishing at infinity on locally compact Hausdorff spaces. Gelfand and Naimark's second theorem (Theorem 1.3.2) shows that all C^* -algebras can be realized as algebras of operators on a Hilbert space. This theorem is what makes C^* -algebras considered part of operator algebras, in contrast to Banach algebras. In particular, every von Neumann algebra is a C^* -algebra, since the norm topology is stronger than the weak topology.

Gelfand and Naimark's first theorem also has conceptual content. It suggests that a noncommutative C^* -algebra can be thought of somehow as a “noncommutative” space. This idea turns out to be fruitful, and is the basis of the noncommutative geometry developed by Alain Connes, in which notions such as de Rham cohomology are extended from topological spaces to operator algebras [Con94]. Another great success in operator algebras which is based on Gelfand and Naimark's first theorem is *operator K -theory*, which generalizes the K -theory of locally compact Hausdorff spaces. Together with some other data, it has recently been shown that K -theory completely classifies a large class of C^* -algebras in the sense that two C^* -algebras from this class are isomorphic if and only if their K -theory are isomorphic. This is the *Elliott classification program* initiated by George A. Elliott in [Ell76].

1.4 Twisted group C^* -algebras and noncommutative tori

Some of the most versatile examples of C^* -algebras are the noncommutative tori. In order to define them, let $\Theta = (\theta_{ij})_{i,j=1}^n$ be a skew symmetric real $n \times n$ matrix. A noncommutative n -torus A_Θ is a C^* -algebra universally generated by n unitary elements $u_1, \dots, u_n$ satisfying the commutation relations

$$u_i u_j = e^{2\pi i \theta_{ij}} u_j u_i, \quad 1 \leq i, j \leq n. \quad (1.21)$$

Here, universally generated is meant in the sense that if B is any other C^* -algebra with unitary elements $v_1, \dots, v_n$ satisfying the same relations as in (1.21), then there exists a unique $*$ -homomorphism $\phi: A_\Theta \rightarrow B$ with $\phi(u_i) = v_i$ for each $1 \leq i \leq n$. The elements in A_Θ of the form

$$\sum_{k_1, \dots, k_n \in \mathbb{Z}} c_{k_1, \dots, k_n} u_1^{k_1} \cdots u_n^{k_n}, \quad \text{where } (c_{k_1, \dots, k_n}) \in \ell^1(\mathbb{Z}^n) \quad (1.22)$$

form a natural Banach $*$ -subalgebra of A_Θ .

The name “noncommutative torus” comes from the fact that the C^* -algebra of continuous functions on the n -torus $\mathbb{T}^n$ is universally generated by n commuting

1. Introduction

unitary elements, i.e. unitaries satisfying (1.21) with $\theta_{i,j} = 0$ for all $1 \leq i, j \leq n$. If $\mathbb{T}^n = \mathbb{R}^n / \mathbb{Z}^n$, then these unitaries are given by

$$u_j(t_1, \dots, t_n) = e^{2\pi i t_j}, \quad 1 \leq j \leq n.$$

The sums from (1.22) then correspond precisely to Fourier expansions of continuous functions on $\mathbb{T}^n$. Thus, one can view the C^* -algebras A_Θ as twisted versions of $C(\mathbb{T}^n)$.

In the representation theory of groups, the representation theory of an appropriate algebra often captures the representation theory of a group. For instance, if G is a finite group, then the representations of G are in one-to-one correspondence with modules over the complex group algebra $\mathbb{C}G$. For general locally compact groups G , a similar role is played by the (full) group C^* -algebra $C^*(G)$ of G . Similarly, if σ is a 2-cocycle on G , the σ -projective representation theory of G is captured by the σ -twisted group C^* -algebra $C^*(G, \sigma)$ of G . More precisely, there is a bijection between unitary σ -projective representations of G on a Hilbert space H and nondegenerate representations of $C^*(G, \sigma)$ on H . The twisted group C^* -algebra can be constructed as follows: For our purposes, we assume that G is discrete. Consider the Banach algebra $\ell^1(G)$, but with *twisted convolution*:

$$(a * b)(x) = \sum_{y \in G} \sigma(y, y^{-1}x) a(y) b(y^{-1}x), \quad a, b \in \ell^1(G), x \in G.$$

The resulting Banach algebra is denoted by $\ell^1(G, \sigma)$. It also comes equipped with an involution turning it into a Banach $*$ -algebra:

$$a^*(x) = \overline{\sigma(x, x^{-1}) a(x^{-1})}, \quad x \in G.$$

Now given any Banach $*$ -algebra A , there is an associated *universal C^* -algebra* in which A embeds as a dense $*$ -subalgebra. For $\ell^1(G, \sigma)$ this universal C^* -algebra is precisely $C^*(G, \sigma)$. It satisfies the following universal property: It is universally generated by unitary elements $(u_x)_{x \in G}$ that satisfy the relations

$$u_x u_y = \sigma(x, y) u_{xy}, \quad x, y \in G. \quad (1.23)$$

Given a σ -projective unitary representation $U: G \rightarrow \mathcal{U}(H)$, it extends to a $*$ -representation $\bar{U}: \ell^1(G, \sigma) \rightarrow \mathcal{B}(H)$ via the following formula:

$$\bar{U}(a) = \sum_{x \in G} a(x) U(x), \quad a \in \ell^1(G, \sigma). \quad (1.24)$$

This representation further extends to a representation of the C^* -algebra $C^*(G, \sigma)$.

Now if the 2-cocycle σ is constantly equal to 1, one recovers ordinary unitary representations and the untwisted full group C^* -algebra $C^*(G)$. If the discrete group G is abelian, then convolution becomes a commutative operation, so $C^*(G)$ is a commutative unital C^* -algebra. It follows from Theorem 1.3.1 that it

should be isomorphic to the C^* -algebra of continuous functions on some compact Hausdorff space. It turns out that the Fourier transform recovers this space: As usual, the Fourier transform

$$\mathcal{F}: \ell^1(G) \rightarrow C(\widehat{G})$$

is an injective $*$ -homomorphism of Banach $*$ -algebras with dense range. One can then extend the Fourier transform to an isomorphism of C^* -algebras $C^*(G) \rightarrow C(\widehat{G})$. Thus, the Pontryagin dual $\widehat{G}$ is the compact space we were looking for. This motivates the viewpoint that $C^*(G, \sigma)$ can be thought of as a noncommutative or twisted analogue of the space $\widehat{G}$, if one does not distinguish the latter from its algebra of continuous functions. In particular, the C^* -algebras $C^*(\mathbb{Z}^n, \sigma)$ should be noncommutative analogues of $\widehat{\mathbb{Z}^n} \cong \mathbb{T}^n$, that is, noncommutative tori. And this definition of noncommutative tori turns out to be equivalent to the one we have already given: For $G = \mathbb{Z}^n$, every 2-cocycle is (up to cohomology) of the form $\sigma_\Theta(k, l) = e^{\pi i \langle k, \Theta l \rangle / 2}$ for some skew-symmetric real matrix Θ [Ell84]. If one sets $u_i = \delta_{e_i}$ where e_i is the i -th standard basis vector of $\mathbb{Z}^n$, one deduces the relations (1.21) from (1.23).

1.5 Vector bundles and projective modules

A vector bundle over a compact space X is, roughly speaking, a family of finite dimensional vector spaces continuously parametrized by X . More precisely, a (complex) vector bundle E over X of rank k consists of a continuous surjection $\pi: E \rightarrow X$ and the structure of a k -dimensional complex vector space on each fiber $E_x = \pi^{-1}(\{x\})$ such that the following holds: For all $x \in X$, there exists a neighbourhood U of x and a homeomorphism $h: \pi^{-1}(U) \rightarrow U \times \mathbb{C}^k$ such that $\pi(h^{-1}(x, v)) = x$, and such that the restriction $h|_{E_x}: E_x \rightarrow \{x\} \times \mathbb{C}^k$ is a linear isomorphism. In geometry, one studies smooth vector bundles over manifolds, with the tangent and cotangent bundles of a manifold being important examples.

Vector bundles over a space X carry important topological information about X . One way to collect this information into a mathematical object is to form the monoid $V(X)$ consisting of isomorphism classes of vector bundles over X , where the binary operation is the *Whitney sum* of vector bundles: Given two bundles E and F over X , their Whitney sum $E \oplus F$ has the fiber $E_x \oplus F_x$ over each point $x \in X$. The construction $X \mapsto V(X)$ is functorial and preserves e.g. inverse limits. From any monoid V one can construct the associated *Grothendieck group*, which consists of formal differences of elements from V . The Grothendieck group of $V(X)$ is precisely the K -theory group $K^0(X)$.

K -theory was first developed by Alexander Grothendieck for sheaves over algebraic varieties in 1957 [BS58], and then for topological spaces by Michael Atiyah and Friedrich Hirzebruch in 1959 [AH59]. It was the insight of Serre [Ser55] and Swan [Swa62] that there is an analogy between vector bundles and projective modules, which we will now outline: If A is a ring, then a left A -module $\mathcal{E}$ is said to be (*finitely generated*) *projective* if there exists another left A -module $\mathcal{F}$ such that $\mathcal{E} \oplus \mathcal{F} \cong A^k$ for some $k \in \mathbb{N}$. Using direct sums of modules, one

1. Introduction

can form a monoid $V(A)$ of isomorphism classes of projective modules over A . Furthermore, $K_0(A)$ is the Gröthendieck group of $V(A)$.

A (*continuous*) *section* of a vector bundle $\pi: E \rightarrow X$ over a compact space X is a continuous function $\xi: X \rightarrow E$ such that $\pi \circ \xi = \text{id}_X$. The set $\Gamma(E)$ of all continuous sections of E forms a projective module over $C(X)$ with respect to pointwise operations. The Serre–Swan theorem [Ser55; Swa62] states the following:

Theorem 1.5.1. *Let X be a compact Hausdorff space. Then for every projective $C(X)$ -module $\mathcal{E}$, there exists a (unique up to isomorphism) vector bundle $E \rightarrow X$ such that*

$$\mathcal{E} \cong \Gamma(E).$$

In fact, the functor from the category of vector bundles over X to the category of projective modules over $C(X)$ sending a bundle E to its $C(X)$ -module of sections is an equivalence of categories.

The Serre–Swan theorem has important implications in noncommutative geometry: It allows one to think of projective modules over a noncommutative C^* -algebra A as “noncommutative” vector bundles over the noncommutative space A . A consequence of the Serre–Swan theorem is also that $K_0(C(X)) \cong K^0(X)$, which shows that K -theory for C^* -algebras is a generalization of K -theory for topological spaces.

In geometry, vector bundles $E \rightarrow X$ are often equipped with a natural extra structure, namely an *inner product*: This is a collection of inner products $\langle \cdot, \cdot \rangle_x$ on each fiber E_x that varies continuously with x . For instance, if $TM \rightarrow M$ is the tangent bundle of a smooth manifold M , then an inner product on TM that varies smoothly with $x \in M$ is precisely a Riemannian metric. If one starts with an inner product on E , one gets an inner product-like structure on the $C(X)$ -module $\Gamma(E)$ defined as follows: Given $\xi, \eta \in \Gamma(E)$, the function $\bullet\langle \xi, \eta \rangle$ on X is given by

$$\bullet\langle \xi, \eta \rangle(x) = \langle \xi(x), \eta(x) \rangle_x, \quad x \in X.$$

By continuity of the inner product, this defines a map $\Gamma(E) \times \Gamma(E) \rightarrow C(X)$. This “ $C(X)$ -valued inner product” enjoys many of the properties that an ordinary $\mathbb{C}$ -valued inner product has. Thus we are lead naturally to the following notion.

Let A be a unital C^* -algebra. A *left Hilbert A -module* $\mathcal{E}$ is a left A -module that is equipped with an A -valued inner product $\bullet\langle \cdot, \cdot \rangle: \mathcal{E} \times \mathcal{E} \rightarrow A$ that satisfies the following axioms:

- (i) $\bullet\langle a\xi + b\eta, \gamma \rangle = a\bullet\langle \xi, \gamma \rangle + b\bullet\langle \eta, \gamma \rangle$ for $a, b \in A$ and $\xi, \eta, \gamma \in \mathcal{E}$.
- (ii) $\bullet\langle \xi, \eta \rangle^* = \bullet\langle \eta, \xi \rangle$ for $\xi, \eta \in \mathcal{E}$.
- (iii) $\bullet\langle \xi, \xi \rangle \geq 0$, with $\bullet\langle \xi, \xi \rangle = 0$ if and only if $\xi = 0$.
- (iv) $\mathcal{E}$ is complete with respect to the norm $\|\xi\|_{\mathcal{E}} = \|\bullet\langle \xi, \xi \rangle\|^{1/2}$.

A version of the Serre–Swan theorem then states that finitely generated Hilbert C^* -modules over $C(X)$ correspond precisely to Hermitian vector bundles over X , i.e. vector bundles equipped with an inner product.

The correct notion of a morphism of Hilbert A -modules $\mathcal{E}$ and $\mathcal{F}$ is an *adjointable* map $T: \mathcal{E} \rightarrow \mathcal{F}$, i.e. an A -linear map which possesses an adjoint $T^*: \mathcal{F} \rightarrow \mathcal{E}$ in the sense that

$$\bullet \langle T\xi, \eta \rangle = \bullet \langle \xi, T^*\eta \rangle, \quad \xi \in \mathcal{E}, \eta \in \mathcal{F}.$$

In contrast to the Hilbert space case, not all bounded linear maps have adjoints.

Hilbert C^* -modules were first introduced in the commutative case by Kaplansky in 1953 [Kap53]. For general C^* -algebras they were first studied by William Paschke [Pas73] and then by Marc Rieffel in his seminal works on induced representations [Rie74a] and Morita equivalence [Rie74b] for operator algebras. In [Rie74a], Rieffel showed that Mackey’s theory of induced representations of locally compact groups can be generalized to the setting of C^* -algebras.

In algebra, two rings are said to be *Morita equivalent* if their left (or equivalently right) module categories are equivalent as categories. Morita’s theorem [Mor58] characterizes Morita equivalence of rings R and S in terms of an R - S -bimodule satisfying certain properties. Rieffel’s notion of Morita equivalence for C^* -algebras is today an essential part of operator algebras, and is based on Hilbert C^* -bimodules. C^* -algebras A and B are Morita equivalent if there exists an *imprimitivity A - B -bimodule*, i.e. an A - B -bimodule $\mathcal{E}$ which is a left Hilbert A -module and a right Hilbert B -module that satisfies suitable compatibility conditions. We denote the inner product on $\mathcal{E}$ as a left Hilbert A -module by $\bullet \langle \cdot, \cdot \rangle$ and the inner product on $\mathcal{E}$ as a right Hilbert B -module by $\langle \cdot, \cdot \rangle_\bullet$. Among the axioms of an imprimitivity bimodule is

$$\bullet \langle \xi, \eta \rangle \gamma = \xi \langle \eta, \gamma \rangle_\bullet, \quad \xi, \eta, \gamma \in \mathcal{E}, \quad (1.25)$$

known as the associativity condition. One also requires that the ideal $\text{span}\{\bullet \langle \xi, \eta \rangle : \xi, \eta \in \mathcal{E}\}$ is dense in A , and similarly for B . As in the case of rings, a Morita equivalence between C^* -algebras A and B implements an isomorphism $B \cong \mathcal{K}(\mathcal{E})$, where $\mathcal{K}(\mathcal{E})$ is a C^* -algebra of operators on the left module $\mathcal{E}$ known as (*generalized*) *compact operators*. They are the closure of the span of the “rank one” adjointable operators $\Theta_{\eta, \gamma}: \mathcal{E} \rightarrow \mathcal{E}$ given by

$$\Theta_{\eta, \gamma} \xi = \bullet \langle \xi, \eta \rangle \gamma, \quad \xi, \eta, \gamma \in \mathcal{E}.$$

Note that Θ is the conventional notation for rank one operators and should not be confused with the skew-symmetric matrix defining a noncommutative torus. Following Rieffel [Rie81, Theorem 2.1], If B is unital with identity 1_B , then the dense ideal consisting of the span of the elements $\langle \xi, \eta \rangle_\bullet$ for $\xi, \eta \in \mathcal{E}$ intersects the open set of invertible elements of B , hence it is all of B . Consequently one can find elements $\eta_1, \dots, \eta_k, \gamma_1, \dots, \gamma_k \in \mathcal{E}$ such that

$$\sum_{j=1}^k \langle \eta_j, \gamma_j \rangle_\bullet = 1_B. \quad (1.26)$$

1. Introduction

The isomorphism $\mathcal{K}(\mathcal{E}) \cong B$ takes $\Theta_{\eta, \gamma}$ to $\langle \eta, \gamma \rangle_\bullet$, which implies that

$$\sum_{j=1}^k \Theta_{\eta_j, \gamma_j} = I, \quad (1.27)$$

where I is the identity operator on $\mathcal{E}$. Consequently the sets $\{\eta_1, \dots, \eta_k\}$ and $\{\gamma_1, \dots, \gamma_k\}$ are both algebraic generating sets for $\mathcal{E}$ (as a left A -module).

The operator $\Theta = \sum_{j=1}^k \Theta_{\eta_j, \gamma_j}$ is a generalization of an operator appearing in frame theory on Hilbert spaces known as a *frame-type* operator. The fact that such an operator is equal to the identity operator expresses exactly the fact that two given sequences are dual frames to each other. In 2002, Frank and Larson [FL02] extended the theory of frames for Hilbert spaces to the setting of Hilbert C^* -modules. Formally, a sequence $(\eta_j)_{j \in J}$ in a left Hilbert A -module $\mathcal{E}$ is a *frame* for $\mathcal{E}$ if there exist constants $C, D > 0$ such that

$$C \bullet \langle \xi, \xi \rangle \leq \sum_{j \in J} \bullet \langle \xi, \eta_j \rangle \bullet \langle \xi, \eta_j \rangle^* \leq D \bullet \langle \xi, \xi \rangle \quad (1.28)$$

for every $\xi \in \mathcal{E}$. The inequalities have to be interpreted in terms of positivity in A . Comparing to (1.11), one sees that (1.28) is a natural generalization of frames for Hilbert spaces. As in the Hilbert space case, $(\eta_j)_j$ is a frame if and only if the associated frame operator $\Theta: \mathcal{E} \rightarrow \mathcal{E}$ given by

$$\Theta \xi = \sum_{j \in J} \bullet \langle \xi, \eta_j \rangle \eta_j$$

is invertible.

1.6 Heisenberg modules

We have now come to the objects that connect operator algebras with Gabor frames, namely Heisenberg modules. Besides their link to Gabor frames, Heisenberg modules have been used numerous times in operator algebras, see e.g. [Boc99; Bön+18; Ech+10; LP13]. Heisenberg modules were introduced by Rieffel in his work on projective modules over higher-dimensional noncommutative tori [Rie88]. The motivation for the construction of Heisenberg modules is to produce Hilbert C^* -modules over a given twisted group C^* -algebra of an abelian discrete group.

Let Δ be an abelian discrete group with 2-cocycle σ . Suppose one can find an embedding of Δ as a lattice inside $G \times \widehat{G}$ for a locally compact abelian group G in such a way that σ is cohomologous to $c|_\Delta$, where c denotes the 2-cocycle of the Weyl–Heisenberg representation associated to $G \times \widehat{G}$ from Section 1.2. It then follows that $C^*(\Delta, \sigma) \cong C^*(\Delta, c)$, so we can focus on constructing projective modules over $C^*(\Delta, c)$.

What Rieffel then demonstrates is that there is a very natural way to construct a Hilbert C^* -module over $C^*(\Delta, c)$ in this setting. The key ingredient is the

Weyl–Heisenberg representation $\pi|_{\Delta}$ restricted to the lattice Δ , which extends to a representation of $C^*(\Delta, c)$ (as in Section 1.4) which is *faithful*. Rieffel starts with the Schwartz–Bruhat space of functions, but one can equally well use the Feichtinger algebra $S_0(G)$, as shown by Luef in [Lue09]. Both of these spaces satisfy the same essential properties which we formulate for $S_0(G)$. First, $S_0(G)$ is invariant under time-frequency shifts, in fact $\|\pi(z)f\|_{S_0} = \|f\|_{S_0}$ for every $f \in S_0(G)$. Thus, the twisted convolution algebra $\ell^1(\Delta, c)$ acts on $S_0(G)$ via

$$a \cdot f = \bar{\pi}(a)f = \sum_{z \in \Delta} a(z)\pi(z)\xi, \quad a \in \ell^1(\Delta, c),$$

where $\bar{\pi}$ denotes the extension of the Weyl–Heisenberg representation to $\ell^1(\Delta, c)$, see (1.24). Furthermore, for $f, g \in S_0(G)$, the short-time Fourier transform $V_g f$ is in $S_0(G \times \widehat{G})$. From general properties of $S_0(G)$ [Jak18], it follows that the restriction of $V_g f$ to Δ is in $S_0(\Delta) = \ell^1(\Delta)$, and consequently the function $\bullet \langle f, g \rangle$ on Δ given by

$$\bullet \langle f, g \rangle(z) = \langle f, \pi(z)g \rangle = V_g f(z), \quad z \in \Delta,$$

is in $\ell^1(\Delta)$. One must then check that $S_0(G)$ with this left structure satisfies the axioms of a left *pre-inner product* $\ell^1(\Delta, c)$ -module. These are routine calculations except for the positivity of $\bullet \langle f, f \rangle$ which is a consequence of the Poisson summation formula. One can then complete $S_0(G)$ into a left Hilbert C^* -module $\mathcal{E}_{\Delta}$ over $C^*(\Delta, c)$, called a *Heisenberg module*. In a similar fashion, the C^* -algebra $C^*(\Delta^{\circ}, \bar{c})$ acts on $\mathcal{E}_{\Delta}$ on the right, where Δ° is the adjoint lattice of Δ . In this way, $\mathcal{E}_{\Delta}$ becomes an imprimitivity module. When written out, the associativity condition (1.25) of this imprimitivity module becomes

$$\sum_{z \in \Delta} \langle f, \pi(z)g \rangle \overline{\langle h, \pi(z)k \rangle} = \frac{1}{\text{vol}(\Delta)} \sum_{z \in \Delta^{\circ}} \langle k, \pi(z)g \rangle \overline{\langle h, \pi(z)f \rangle}, \quad f, g, h, k \in S_0(G)$$

after taking the inner product of each side with a fourth function. In time-frequency analysis, the above identity is known as the *fundamental identity of Gabor analysis* and was proved by Janssen and Tolomieri–Orr [Jan95; TO95] independently from (and 7 years after) Rieffel’s work.

As we show in Paper II, one can recover the Hilbert space $L^2(G)$ from $\mathcal{E}_{\Delta}$ by means of localization [Lan95]: By using the canonical faithful tracial state tr on $C^*(\Delta, c)$, the following defines a complex-valued inner product on $\mathcal{E}_{\Delta}$:

$$\langle f, g \rangle := \text{tr}(\bullet \langle f, g \rangle).$$

In fact, for $f, g \in S_0(G) \subseteq L^2(G)$, this is the usual L^2 -inner product. Thus, the completion of $\mathcal{E}_{\Delta}$ with respect to the norm coming from this inner product can be naturally identified with $L^2(G)$ in such a way that the inclusions $S_0(G) \subseteq \mathcal{E}_{\Delta} \subseteq L^2(G)$ are preserved. Furthermore, this localization procedure extends adjointable operators on $\mathcal{E}_{\Delta}$ to bounded operators on $L^2(G)$. As shown in Paper II, the rank one operator $\Theta_{g,h}$ of $\mathcal{E}_{\Delta}$ extends exactly to the Gabor frame-type operator $S_{g,h}$ given by

$$S_{g,h}f = \sum_{z \in \Delta} \langle f, \pi(z)g \rangle \pi(z)h, \quad f \in L^2(G).$$

1. Introduction

This further extends to finite sums of rank one operators in $\mathcal{E}_\Delta$. Using the link between the frame property and invertibility of the associated frame operator, one arrives at the following result:

Theorem 1.6.1. *Let G be a locally compact abelian group, and let Δ be a lattice in $G \times \widehat{G}$. Then a finite subset $\{g_1, \dots, g_k\} \subseteq \mathcal{E}_\Delta$ is a module frame (that is, an algebraic generating set) for $\mathcal{E}_\Delta$ if and only if*

$$\mathcal{G}(g_1, \dots, g_k; \Delta) := \mathcal{G}(g_1, \Delta) \cup \dots \cup \mathcal{G}(g_k, \Delta)$$

is a frame for $L^2(G)$.

The above result is among the main theorems of the paper of Luef [Lue09], where it is proved for windows in $S_0(G)$ and $G = \mathbb{R}^n$. It was generalized to the locally compact abelian case by Jakobsen and Luef in [JL18].

A union of Gabor systems over the same lattice as in Theorem 1.6.1 is called a *multiwindow Gabor system*. The above theorem shows that the search for multiwindow Gabor frames with windows in $\mathcal{E}_\Delta$ is equivalent to the search for finite generating sets of $\mathcal{E}_\Delta$. A finite generating set for $\mathcal{E}_\Delta$ can always be perturbed to lie in $S_0(G)$. Thus, since $\mathcal{E}_\Delta$ is a finitely generated module, the following is an immediate consequence as pointed out by Luef [Lue09]:

Corollary 1.6.2. *There exists a multiwindow Gabor frame $\mathcal{G}(g_1, \dots, g_k; \Delta)$ with $g_1, \dots, g_k \in S_0(G)$ for any given lattice Δ in $G \times \widehat{G}$.*

This existence result for multiwindow Gabor frames was not known before [Lue09] and is one of the first examples of how the connection to Heisenberg modules has given new results in Gabor analysis.

We have seen that the density theorem (Theorem 1.2.1) states that for Gabor frames to exist over a lattice Δ , it is necessary that $\text{vol}(\Delta) \leq 1$. One can then ask whether the converse holds, that is, if Δ is a lattice in $G \times \widehat{G}$, does there exist $g \in L^2(G)$ such that $\mathcal{G}(g, \Delta)$ is a Gabor frame? This was solved by Bekka [Bek04] using von Neumann algebraic techniques, but an important calculation was already done by Rieffel in [Rie88], which we briefly outline: Representing $C^*(\Delta, c)$ faithfully on $L^2(G)$ using the integrated form of the Weyl–Heisenberg representation, one can form the von Neumann algebra

$$W^*(\Delta, c) = \overline{\pi(C^*(\Delta, c))}'' = \{\pi(z) : z \in \Delta\}''.$$

Similarly, one forms $W^*(\Delta^\circ, \bar{c})$ from the right C^* -algebra. These are both finite von Neumann algebras with faithful normal traces, and they are commutants of each other on $L^2(G)$. The center valued von Neumann dimension $\text{cdim}_{W^*(\Delta, c)} L^2(G)$ (or coupling operator) of the left Hilbert $W^*(\Delta, c)$ -module $L^2(G)$ is calculated to be exactly $\text{vol}(\Delta)I$. As shown by Bekka, a Hilbert M -module H is cyclic, i.e. there exists $\xi \in H$ such that $\overline{M}\xi = H$, if and only if $\text{cdim}_M H \leq I$. Furthermore, the existence of a cyclic vector in the above setting is shown to be equivalent to the existence of a frame vector. Thus, one arrives at the following theorem:

Theorem 1.6.3. *Let Δ be a lattice in $G \times \widehat{G}$. Then there exists $g \in L^2(G)$ such that $\mathcal{G}(g, \Delta)$ is a Gabor frame if and only if $\text{vol}(\Delta) \leq 1$.*

Note that Bekka only proves the above for $G = \mathbb{R}^n$, but the argument generalizes to the locally compact abelian setting.

The above von Neumann algebraic considerations only guarantee the existence of windows in $L^2(G)$. If we want well-localized windows, e.g. in $S_0(G)$, the situation becomes more intricate. As we have seen for $G = \mathbb{R}^n$, the Balian–Low theorem is an obstruction to the existence of well-localized windows when $\text{vol}(\Delta) = 1$. It is an open problem whether the converse of the Balian–Low theorem holds: That is, if $\text{vol}(\Delta) < 1$ implies the existence of a function $g \in S_0(\mathbb{R}^n)$ for which $\mathcal{G}(g, \Delta)$ is a Gabor frame. We will now show how Jakobsen and Luef gave a partial solution to this problem by using Rieffel’s results in [Rie88]. First of all, by the link to Heisenberg modules, we can use the following proposition to reformulate the problem:

Proposition 1.6.4. *The left $C^*(\Delta, c)$ -module $\mathcal{E}_\Delta$ is singly generated if and only if there exists $g \in S_0(G)$ such that $\mathcal{G}(g, \Delta)$ is a Gabor frame.*

For $G = \mathbb{R}^n$, every lattice Δ in $\mathbb{R}^n \times \widehat{\mathbb{R}^n} = \mathbb{R}^{2n}$ is of the form $M\mathbb{Z}^{2n}$ for some real, invertible $n \times n$ matrix M . Thus, $C^*(\Delta, c) = C^*(M\mathbb{Z}^{2n}, c)$ is a twisted group C^* -algebra on $M\mathbb{Z}^{2n} \cong \mathbb{Z}^{2n}$, hence a noncommutative $2n$ -torus. One can find the corresponding Θ -matrix as follows: The $2n$ unitaries that generate $C^*(M\mathbb{Z}^{2n}, c)$ are $\delta_{Me_1}, \dots, \delta_{Me_{2n}}$, and the commutation relations between these are

$$\delta_{Me_i} \delta_{Me_j} = e^{2\pi i \langle Me_i, JMe_j \rangle} \delta_{Me_j} \delta_{Me_i}, \quad 1 \leq i, j \leq 2n,$$

where J denotes the standard $2n \times 2n$ symplectic matrix

$$J = \begin{pmatrix} 0_n & I_n \\ -I_n & 0_n \end{pmatrix}.$$

Thus, the Θ matrix such that $C^*(\Delta, c) \cong A_\Theta$ is given by $\Theta = M^t J M$.

Some of the main results of Rieffel in his work on projective modules over noncommutative tori [Rie88] are the following:

Theorem 1.6.5. *Let A_Θ be an n -dimensional noncommutative torus where Θ contains at least one irrational number. Let tr denote the canonical faithful tracial state of A_Θ . Then the following hold:*

- (i) $K_0(A_\Theta)$ has cancellation. That is, suppose $\mathcal{E}$ and $\mathcal{F}$ are projective A_Θ -modules such that their classes $[\mathcal{E}]$ and $[\mathcal{F}]$ in K -theory are equal, i.e. there exists some projective module $\mathcal{G}$ such that $\mathcal{E} \oplus \mathcal{G} \cong \mathcal{F} \oplus \mathcal{G}$. Then one can conclude that $\mathcal{E} \cong \mathcal{F}$.
- (ii) The positive cone of $K_0(A_\Theta)$ is determined by the trace. That is, if $x \in K_0(A_\Theta)$ has $K_0(\text{tr})(x) > 0$ where $K_0(\text{tr})$ denotes the homomorphism $K_0(A_\Theta) \rightarrow \mathbb{C}$ induced by tr , then x is in the positive cone of $K_0(A_\Theta)$, i.e. $x = [\mathcal{E}]$ for some projective module $\mathcal{E}$.

1. Introduction

We get the following immediate corollary:

Corollary 1.6.6. *Let A_Θ be a noncommutative torus where Θ contains at least one irrational number. If $\mathcal{E}$ is a projective module over A_Θ such that $0 < K_0(\text{tr})([\mathcal{E}]) < 1$, then $\mathcal{E}$ is singly generated.*

Proof. Set $x = [A_\Theta] - [\mathcal{E}] \in K_0(A_\Theta)$. Since $0 < K_0(\text{tr})([\mathcal{E}]) < 1$, we also have that $0 < K_0(\text{tr})(x) < 1$. By Theorem 1.6.5, it follows that there exists some projective module $\mathcal{F}$ such that $x = [\mathcal{F}]$. Consequently $[\mathcal{E}] + [\mathcal{F}] = [A_\Theta]$, so again by Theorem 1.6.5, we have that $\mathcal{E} \oplus \mathcal{F} \cong A_\Theta$ as A_Θ -modules. But then by picking the projection of A_Θ onto $\mathcal{E}$, we obtain a projection $p \in A_\Theta$ such that $\mathcal{E} \cong pA_\Theta$. Hence $\mathcal{E}$ is singly generated. ■

Assume now that a lattice $\Delta = M\mathbb{Z}^{2n}$ is precisely such that the corresponding matrix $\Theta = M^t J M$ contains at least one irrational entry. Then by Corollary 1.6.6, $\mathcal{E}_\Delta$ is singly generated if $0 < K_0(\text{tr})(\mathcal{E}_\Delta) < 1$. But as shown by Rieffel in [Rie88], the trace of the Heisenberg module $\mathcal{E}_\Delta$ is precisely $\text{vol}(\Delta)$. Thus, using Proposition 1.6.4, Jakobsen and Luef arrived at the following:

Theorem 1.6.7. *Let $\Delta = M\mathbb{Z}^{2n}$ be a lattice in $\mathbb{R}^{2n}$ such that the matrix $\Theta = M^t J M$ has at least one irrational entry. Suppose $\text{vol}(\Delta) < 1$. Then there exists $g \in S_0(\mathbb{R}^n)$ such that $\mathcal{G}(g, \Delta)$ is a Gabor frame.*

The case when the lattice is given by a rational matrix remains unsolved.

We now return to the setting of a locally compact abelian group G and revisit the Balian–Low theorem. In the specialized setting of Problem 1.2.3, we looked at lattices in $G \times \hat{G}$ of the form $\Delta = \Lambda \times \Lambda^\perp$ for a lattice Λ in G . Let us first look at the left coefficient algebra $C^*(\Delta, c)$. For $(x, \omega), (y, \tau) \in \Lambda \times \Lambda^\perp$, we have that

$$c((x, \omega), (y, \tau)) = \overline{\tau(x)} = 1,$$

hence the twisted group C^* -algebra $C^*(\Delta, c)$ is just the group C^* -algebra $C^*(\Delta)$. As we have seen, this commutative C^* -algebra is isomorphic to the continuous functions on the Pontryagin dual of Δ , which we can identify as follows:

$$X_{G, \Lambda} := \hat{\Delta} \cong \hat{\Lambda} \times \hat{\Lambda}^\perp \cong (\hat{G}/\Lambda^\perp) \times (G/\Lambda) \cong (G/\Lambda) \times (\hat{G}/\Lambda^\perp). \quad (1.29)$$

Furthermore, by the Serre–Swan theorem Theorem 1.5.1, the Heisenberg module $\mathcal{E}_\Delta$ must now correspond to a vector bundle E over the space $X_{G, \Lambda}$. Paper III is dedicated to finding a description of this bundle, and showing that it generalizes the bundle discussed by Balan in [Bal01]. The map that implements the isomorphism $\mathcal{E}_\Delta \rightarrow \Gamma(E)$ turns out to be the already mentioned Zak transform. The bundle E is a line bundle, so a generator for $\Gamma(E)$ is the same as a nonvanishing section, which then trivializes the bundle. Thus, $\mathcal{E}_\Delta$ is singly generated, i.e. there exists a Gabor frame $\mathcal{G}(g, \Delta)$ with $g \in S_0(G)$, if and only if the bundle E is trivial. The validity of a Balian–Low theorem for the lattice $\Lambda \times \Lambda^\perp$ then amounts to the nontriviality of the associated bundle E . We prove in Paper III explicitly the nontriviality of the vector bundle for $G = \mathbb{R} \times \mathbb{Q}_p$ and Λ the lattice $\mathbb{Z}[1/p]$. This in turn gives us a Balian–Low theorem in that setting.

However, in Paper IV we generalize this Balian–Low theorem using completely different techniques to all groups with noncompact identity component and *all* lattices in $G \times \widehat{G}$ with volume 1, not just those of the form $\Lambda \times \Lambda^\perp$.

1.7 Summary of Papers

Paper I This paper introduces the rational adele group and the group $\mathbb{R} \times \mathbb{Q}_p$ as natural settings for Gabor analysis. This comes from the fact that both of these groups admit lattices, namely $\mathbb{Q}$ and $\mathbb{Z}[1/p]$ respectively. We give a characterization of the Feichtinger algebra for these groups, and prove the existence of certain Gabor frames over variations of the mentioned lattices.

Paper II In this paper, we use localization of Hilbert C^* -modules to recover $L^2(G)$ from the Heisenberg module associated to a lattice Δ in the time-frequency plane of G . This gives a continuous embedding of the Heisenberg module into $L^2(G)$ which is needed to formulate some of the results in Paper III. In addition, we obtain a new proof of the description of generators of Heisenberg modules as multiwindow Gabor frames that extends to windows in the Heisenberg module and not just $S_0(G)$.

Paper III This paper is devoted to the study of the Balian–Low theorem in the setting of LCA groups and its connection to Heisenberg modules. We show how Balan’s bundle E from [Bal01] shows up naturally when considering Heisenberg modules, and that the nontriviality of E can be explained through the Hilbert C^* -module isomorphism of $\Gamma(E)$ with the appropriate Heisenberg module. We consider this situation for general LCA groups, and connect the validity of the Balian–Low theorem for LCA groups to the nontriviality of associated vector bundles. In particular, we show that the triviality of these bundles explain why there is no Balian–Low theorem for compact or discrete groups. We also revisit the setting of Paper I, namely the truncated adele group $\mathbb{R} \times \mathbb{Q}_p$, and show that in this case the associated vector bundle is nontrivial: Thereby providing a Balian–Low theorem for this group.

Paper IV In this paper, we prove deformation results for Gabor frames on general locally compact abelian groups with respect to the Braconnier topology. Using the Braconnier modular function, we prove that the Balian–Low theorem for $S_0(G)$ holds for groups with noncompact identity component. We remark that this is for *all* lattices of volume 1 in $G \times \widehat{G}$, not just those of the form $\Lambda \times \Lambda^\perp$. In particular, this implies Balian–Low theorems on the S -adeles, which generalizes one of the main theorems of Paper III. We are also able to prove a converse of the Balian–Low theorem for groups with compact identity component, and demonstrate that topologically, the adeles constructed from a global function field are of this form.

References

- [AFK14] Ascensi, G., Feichtinger, H. G., and Kaiblinger, N. “Dilation of the Weyl symbol and Balian-Low theorem”. In: *Trans. Amer. Math. Soc.* Vol. 366, no. 7 (2014), pp. 3865–3880.
- [AH59] Atiyah, M. F. and Hirzebruch, F. “Riemann-Roch theorems for differentiable manifolds”. In: *Bull. Amer. Math. Soc.* Vol. 65 (1959), pp. 276–281.
- [Bal01] Balan, R. “Topological obstructions to localization results”. In: *SPIE Wavelets Applications in Signal and Image Processing VIII* (2001), pp. 184–192.
- [Bal81] Balian, R. “Un principe d’incertitude fort en théorie du signal ou en mécanique quantique”. In: *C. R. Acad. Sci. Paris Sér. II Méc. Phys. Chim. Sci. Univers Sci. Terre* vol. 292, no. 20 (1981), pp. 1357–1362.
- [Bat88] Battle, G. “Heisenberg proof of the Balian-Low theorem”. In: *Lett. Math. Phys.* Vol. 15, no. 2 (1988), pp. 175–177.
- [Bek04] Bekka, B. “Square integrable representations, von Neumann algebras and an application to Gabor analysis”. In: *J. Fourier Anal. Appl.* Vol. 10, no. 4 (2004), pp. 325–349.
- [BHJ26] Born, M., Heisenberg, W., and Jordan, P. “Zur Quantenmechanik. II.” In: *Zeitschrift für Physik* vol. 35, no. 8 (Aug. 1926), pp. 557–615.
- [BJ25] Born, M. and Jordan, P. “Zur Quantenmechanik”. In: *Zeitschrift für Physik* vol. 34, no. 1 (Dec. 1925), pp. 858–888.
- [Boc99] Boca, F. P. “Projections in rotation algebras and theta functions”. In: *Comm. Math. Phys.* Vol. 202, no. 2 (1999), pp. 325–357.
- [Bön+18] Bönicke, C. et al. “Isomorphism and Morita equivalence classes for crossed products of irrational rotation algebras by cyclic subgroups of $SL_2(\mathbb{Z})$ ”. In: *J. Funct. Anal.* Vol. 275, no. 11 (2018), pp. 3208–3243.
- [BS58] Borel, A. and Serre, J.-P. “Le théorème de Riemann-Roch”. fr. In: *Bulletin de la Société Mathématique de France* vol. 86 (1958), pp. 97–136.
- [Con94] Connes, A. *Noncommutative geometry*. Academic Press, Inc., San Diego, CA, 1994, pp. xiv+661.
- [Dau90] Daubechies, I. “The wavelet transform, time-frequency localization and signal analysis”. In: *IEEE Trans. Inform. Theory* vol. 36, no. 5 (1990), pp. 961–1005.
- [Ech+10] Echterhoff, S. et al. “The structure of crossed products of irrational rotation algebras by finite subgroups of $SL_2(\mathbb{Z})$ ”. In: *J. Reine Angew. Math.* Vol. 639 (2010), pp. 173–221.
- [Ell76] Elliott, G. A. “On the classification of inductive limits of sequences of semisimple finite-dimensional algebras”. In: *J. Algebra* vol. 38, no. 1 (1976), pp. 29–44.

-
- [Ell84] Elliott, G. A. “On the K -theory of the C^* -algebra generated by a projective representation of a torsion-free discrete abelian group”. In: *Operator algebras and group representations, Vol. I (Neptun, 1980)*. Vol. 17. Monogr. Stud. Math. Pitman, Boston, MA, 1984, pp. 157–184.
- [Fei81] Feichtinger, H. G. “On a new Segal algebra”. English. In: *Monatsh. Math.* Vol. 92 (1981), pp. 269–289.
- [FK04] Feichtinger, H. G. and Kaiblinger, N. “Varying the time-frequency lattice of Gabor frames”. In: *Trans. Amer. Math. Soc.* Vol. 356, no. 5 (2004), pp. 2001–2023.
- [FL02] Frank, M. and Larson, D. R. “Frames in Hilbert C^* -modules and C^* -algebras”. In: *J. Operator Theory* vol. 48, no. 2 (2002), pp. 273–314.
- [Gel50] Gelfand, I. M. “Expansion in characteristic functions of an equation with periodic coefficients”. In: *Doklady Akad. Nauk SSSR (N.S.)* Vol. 73 (1950), pp. 1117–1120.
- [GN43] Gelfand, I. and Neumark, M. “On the imbedding of normed rings into the ring of operators in Hilbert space”. In: *Rec. Math. [Mat. Sbornik] N.S.* Vol. 12(54) (1943), pp. 197–213.
- [Grö01] Gröchenig, K. *Foundations of time-frequency analysis*. Applied and Numerical Harmonic Analysis. Birkhäuser Boston, Inc., Boston, MA, 2001, pp. xvi+359.
- [Grö98] Gröchenig, K. “Aspects of Gabor analysis on locally compact abelian groups”. In: *Gabor analysis and algorithms*. Appl. Numer. Harmon. Anal. Birkhäuser Boston, Boston, MA, 1998, pp. 211–231.
- [Hei25] Heisenberg, W. “Über quantentheoretische Umdeutung kinematischer und mechanischer Beziehungen.” In: *Zeitschrift für Physik* vol. 33, no. 1 (Dec. 1925), pp. 879–893.
- [Hei27] Heisenberg, W. “Über den anschaulichen Inhalt der quantentheoretischen Kinematik und Mechanik”. In: *Zeitschrift für Physik* vol. 43, no. 3 (Mar. 1927), pp. 172–198.
- [HL00] Han, D. and Larson, D. R. “Frames, bases and group representations”. In: *Mem. Amer. Math. Soc.* Vol. 147, no. 697 (2000), pp. x+94.
- [HR63] Hewitt, E. and Ross, K. A. *Abstract harmonic analysis. Vol. I: Structure of topological groups. Integration theory, group representations*. Die Grundlehren der mathematischen Wissenschaften, Bd. 115. Academic Press, Inc., Publishers, New York; Springer-Verlag, Berlin-Göttingen-Heidelberg, 1963, pp. viii+519.
- [Jak18] Jakobsen, M. S. “On a (no longer) new Segal algebra: a review of the Feichtinger algebra”. In: *J. Fourier Anal. Appl.* Vol. 24, no. 6 (2018), pp. 1579–1660.

- [Jan81] Janssen, A. J. E. M. “Gabor representation of generalized functions”. In: *J. Math. Anal. Appl.* Vol. 83, no. 2 (1981), pp. 377–394.
- [Jan95] Janssen, A. J. E. M. “Duality and biorthogonality for Weyl-Heisenberg frames”. In: *J. Fourier Anal. Appl.* Vol. 1, no. 4 (1995), pp. 403–436.
- [Jan97] Janssen, A. J. E. M. “From continuous to discrete Weyl-Heisenberg frames through sampling”. In: vol. 3. 5. Dedicated to the memory of Richard J. Duffin. 1997, pp. 583–596.
- [JL16] Jakobsen, M. S. and Lemvig, J. “Density and duality theorems for regular Gabor frames”. In: *J. Funct. Anal.* Vol. 270, no. 1 (2016), pp. 229–263.
- [JL18] Jakobsen, M. S. and Luef, F. “Duality of Gabor Frames and Heisenberg Modules”. preprint, arXiv.org:1806.05616. 2018.
- [JL19] Jakobsen, M. S. and Luef, F. “Sampling and periodization of generators of Heisenberg modules”. In: *Int. J. Math.* Vol. 30, no. 10 (2019), p. 1950001.
- [Kap53] Kaplansky, I. “Modules over operator algebras”. In: *Amer. J. Math.* Vol. 75 (1953), pp. 839–858.
- [Ken27] Kennard, E. H. “Zur Quantenmechanik einfacher Bewegungstypen”. In: *Zeitschrift für Physik* vol. 44, no. 4 (Apr. 1927), pp. 326–352.
- [KK98] Kaniuth, E. and Kutyniok, G. “Zeros of the Zak transform on locally compact abelian groups”. In: *Proc. Amer. Math. Soc.* Vol. 126, no. 12 (1998), pp. 3561–3569.
- [Lan95] Lance, E. C. *Hilbert C^* -modules*. Vol. 210. London Mathematical Society Lecture Note Series. A toolkit for operator algebraists. Cambridge University Press, Cambridge, 1995, pp. x+130.
- [Low85] Low, F. “Complete sets of wave packets”. In: *A Passion for Physics: Essays in Honor of Geoffrey Chew*. World Scientific, 1985, pp. 17–22.
- [LP13] Latrémolière, F. and Packer, J. A. “Noncommutative solenoids and their projective modules”. In: *Commutative and noncommutative harmonic analysis and applications*. Vol. 603. Contemp. Math. Amer. Math. Soc., Providence, RI, 2013, pp. 35–53.
- [Lue09] Luef, F. “Projective modules over noncommutative tori are multi-window Gabor frames for modulation spaces”. In: *J. Funct. Anal.* Vol. 257, no. 6 (2009), pp. 1921–1946.
- [MN36] Murray, F. J. and Neumann, J. von. “On rings of operators”. In: *Ann. of Math. (2)* vol. 37, no. 1 (1936), pp. 116–229.
- [Mor58] Morita, K. “Duality for modules and its applications to the theory of rings with minimum condition”. In: *Sci. Rep. Tokyo Kyoiku Daigaku Sect. A* vol. 6 (1958), pp. 83–142.

-
- [Pas73] Paschke, W. L. “Inner product modules over B^* -algebras”. In: *Trans. Amer. Math. Soc.* Vol. 182 (1973), pp. 443–468.
 - [Rie74a] Rieffel, M. A. “Induced representations of C^* -algebras”. In: *Advances in Math.* Vol. 13 (1974), pp. 176–257.
 - [Rie74b] Rieffel, M. A. “Morita equivalence for C^* -algebras and W^* -algebras”. In: *J. Pure Appl. Algebra* vol. 5 (1974), pp. 51–96.
 - [Rie81] Rieffel, M. A. “ C^* -algebras associated with irrational rotations”. In: *Pacific J. Math.* Vol. 93, no. 2 (1981), pp. 415–429.
 - [Rie88] Rieffel, M. A. “Projective modules over higher-dimensional noncommutative tori”. In: *Canad. J. Math.* Vol. 40, no. 2 (1988), pp. 257–338.
 - [RV99] Ramakrishnan, D. and Valenza, R. J. *Fourier Analysis on Number Fields*. New York: Springer, 1999.
 - [Sch26] Schrödinger, E. “Quantisierung als Eigenwertproblem”. In: *Annalen der Physik* vol. 385, no. 13 (1926), pp. 437–490.
 - [Ser55] Serre, J.-P. “Faisceaux algébriques cohérents”. In: *Ann. of Math. (2)* vol. 61 (1955), pp. 197–278.
 - [Swa62] Swan, R. G. “Vector bundles and projective modules”. In: *Trans. Amer. Math. Soc.* Vol. 105 (1962), pp. 264–277.
 - [TO95] Tolimieri, R. and Orr, R. S. “Poisson summation, the ambiguity function, and the theory of Weyl-Heisenberg frames”. In: *J. Fourier Anal. Appl.* Vol. 1, no. 3 (1995), pp. 233–247.
 - [vNe27] von Neumann, J. “Mathematische Begründung der Quantenmechanik.” German. In: *Nachr. Ges. Wiss. Göttingen, Math.-Phys. Kl.* Vol. 1927 (1927), pp. 1–57.
 - [Wei64] Weil, A. “Sur certains groupes d’opérateurs unitaires”. In: *Acta Math.* Vol. 111 (1964), pp. 143–211.
 - [Wig32] Wigner, E. “On the Quantum Correction For Thermodynamic Equilibrium”. In: *Phys. Rev.* Vol. 40 (5 June 1932), pp. 749–759.

Papers

Time-frequency analysis on the adeles over the rationals

Ulrik Enstad, Mads S. Jakobsen, Franz Luef

Published in *Comptes Rendus Mathématique*, 2019, volume 357, issue 2, pp. 188–199. DOI: 10.1016/j.crma.2018.12.004.

Abstract

We show that the construction of Gabor frames in $L^2(\mathbb{R})$ with generators in $S_0(\mathbb{R})$ and with respect to time-frequency shifts from a rectangular lattice $\alpha\mathbb{Z} \times \beta\mathbb{Z}$ is equivalent to the construction of certain Gabor frames for L^2 over the adeles over the rationals and the group $\mathbb{R} \times \mathbb{Q}_p$. Furthermore, we detail the connection between the construction of Gabor frames on the adeles and on $\mathbb{R} \times \mathbb{Q}_p$ with the construction of certain Heisenberg modules.

I.1 Introduction

The theory of Gabor systems and their frame properties is available on any locally compact abelian (LCA) group [JL16]. However, the construction of explicit examples of Gabor frames with time-frequency shifts from lattices (discrete and co-compact subgroups) is mostly restricted to the *elementary* LCA groups $\mathbb{R}$, $\mathbb{Z}$, $\mathbb{T}$ and $\mathbb{Z}/d\mathbb{Z}$, $d \in \mathbb{N}$. While the group of the p -adic numbers $\mathbb{Q}_p$ does not contain any lattice, the group $\mathbb{R} \times \mathbb{Q}_p$ and also the adèle group over the rationals, $\mathbb{A}_{\mathbb{Q}}$, do contain discrete and co-compact subgroups. This makes these groups eligible for Gabor analysis.

Other efforts to do time-frequency and time-scale (wavelet) analysis on local fields include [AES10; ASS18; BB18; KS10; LJ07; Sha16a; Sha16b] and [SS09].

We mention that [CG15] provides a method of constructing Gabor frames on any group with time-frequency shifts from lattices. In case of the p -adics or other groups that do not contain lattices other methods of building somewhat structured families of functions with the use of *quasi-lattices* have been suggested in [GS07].

In this paper we combine established theory on the p -adic numbers, the adeles, Gabor analysis and the theory of the Feichtinger algebra and the modulation spaces. Our main result (Theorem I.4.2) shows that the construction of (dual) Gabor frames for $L^2(\mathbb{R})$ with generators in the Feichtinger algebra $S_0(\mathbb{R})$ is equivalent to both (I) the construction of certain (dual) Gabor frames for $L^2(\mathbb{A}_{\mathbb{Q}})$

with generators in $S_0(\mathbb{A}_{\mathbb{Q}})$ and (II) the construction of certain (dual) Gabor frames for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ with generators in $S_0(\mathbb{R} \times \mathbb{Q}_p)$.

In Section I.2 we describe the groups $\mathbb{Q}_p$ and $\mathbb{A}_{\mathbb{Q}}$. The *Feichtinger algebra* on the p -adic groups $\mathbb{Z}_p$ and $\mathbb{Q}_p$, the adeles $\mathbb{A}_{\mathbb{Q}}$, and $\mathbb{R} \times \mathbb{Q}_p$ is described in Section I.3. Finally, in Section I.4 we state and prove our main result. In Section I.4.1 we connect the construction of dual Gabor frames in $L^2(\mathbb{A}_{\mathbb{Q}})$ and $L^2(\mathbb{R} \times \mathbb{Q}_p)$ with the construction of idempotent elements in twisted group C^* -algebras of the groups $\mathbb{A}_{\mathbb{Q}}$ and $\mathbb{R} \times \mathbb{Q}_p$ as explained in [JL18]. In this way we answer the question raised in [LP13, Remark 5.8]. This work is related to the investigation of noncommutative solenoids by Latrémolière and Packer [LP13; LP15; LP17] and the work of Larsen and Li [LL12].

We also mention that after submitting the preprint for this paper, we have discovered a simultaneous generalization of the main result to Gabor frames over $L^2(\mathbb{R} \times \Omega_a)$, where Ω_a denotes the group of a -adic integers [HN98, p. 184], [HR63, Section 10]. These groups generalize both the p -adic numbers and the finite adeles over the rationals, but do not appear often in the literature. An example of an application of these groups in operator algebras can be found in [KOQ14].

I.2 The p -adic numbers and adeles over the rationals

In this section we give a brief exposition of the field of p -adic numbers and the adèle group $\mathbb{A}_{\mathbb{Q}}$ over the rational numbers. Given a prime number p , the *p -adic absolute value* on $\mathbb{Q}$ is defined by

$$|x|_p = p^{-k}$$

where $x = p^k(a/b)$ and p divides neither a nor b . One also sets $|0|_p = 0$. The p -adic absolute value satisfies a strengthened version of the triangle inequality (the ultrametric triangle inequality), namely

$$|x + y|_p \leq \max\{|x|_p, |y|_p\}. \quad (\text{I.1})$$

The completion of $\mathbb{Q}$ with respect to the metric $d_p(x, y) = |x - y|_p$ is a field denoted by $\mathbb{Q}_p$ and its elements are called *p -adic numbers*. The topology inherited from the metric makes $\mathbb{Q}_p$ into a locally compact Hausdorff space. In particular the p -adic numbers $\mathbb{Q}_p$ form a (non-compact) locally compact abelian group with respect to the topology induced by the above metric and under addition. One can show that every p -adic number x has a *p -adic expansion* of the form

$$x = \sum_{k=-\infty}^{\infty} a_k p^k,$$

where $a_k \in \{0, \dots, p-1\}$ for each k and there exists some $n \in \mathbb{Z}$ such that $a_k = 0$ for all $k < n$. The sequence $(a_k)_{k \in \mathbb{Z}}$ in this expansion is unique.

The p -adic integers. The closed unit ball in $\mathbb{Q}_p$ is denoted by $\mathbb{Z}_p$ and its elements are called p -adic integers. Because of (I.1) and the multiplicativity of $|\cdot|_p$, $\mathbb{Z}_p$ is a subring of $\mathbb{Q}_p$. In terms of p -adic expansions, a p -adic number $x = \sum_{k \in \mathbb{Z}} a_k p^k$ is a p -adic integer if and only if $a_k = 0$ for $k < 0$. The map $\{0, \dots, p-1\}^{\mathbb{N}} \rightarrow \mathbb{Z}_p$ given by $(a_k)_k \mapsto \sum_k a_k p^k$ is a homeomorphism, which shows that $\mathbb{Z}_p$ has the topology of a Cantor set. In particular, $\mathbb{Z}_p$ is a compact subgroup of $\mathbb{Q}_p$. But $\mathbb{Z}_p$ is also open in $\mathbb{Q}_p$. Indeed, if $x \in \mathbb{Z}_p$, then using (I.1) one shows that the open ball $B_{1/2}(x) = \{y \in \mathbb{Q}_p : |y - x|_p < 1/2\}$ is contained in $\mathbb{Z}_p$.

We take the Haar measure $\mu_{\mathbb{Q}_p}$ on $\mathbb{Q}_p$ so that $\mu_{\mathbb{Q}_p}(\mathbb{Z}_p) = 1$. The Haar measure on $\mathbb{Z}_p$ is the one on $\mathbb{Q}_p$ restricted to $\mathbb{Z}_p$.

The dual groups of $\mathbb{Q}_p$ and $\mathbb{Z}_p$. Denote by $\mathbb{Z}[1/p]$ the subring of $\mathbb{Q}$ consisting of rational numbers of the form a/p^k where $k, a \in \mathbb{Z}$. Then $\mathbb{Q}_p = \mathbb{Z}_p + \mathbb{Z}[1/p]$ and $\mathbb{Z}_p \cap \mathbb{Z}[1/p] = \mathbb{Z}$, so that

$$\mathbb{Q}_p / \mathbb{Z}_p = \frac{\mathbb{Z}_p + \mathbb{Z}[1/p]}{\mathbb{Z}_p} \cong \frac{\mathbb{Z}[1/p]}{\mathbb{Z}_p \cap \mathbb{Z}[1/p]} = \mathbb{Z}[1/p] / \mathbb{Z}$$

as abelian groups. Denote the quotient map $\mathbb{Q}_p \rightarrow \mathbb{Z}[1/p] / \mathbb{Z}$ by $x \mapsto \{x\}_p$. In terms of p -adic expansions, we have $\left\{ \sum_{k \in \mathbb{Z}} a_k p^k \right\}_p = \sum_{k=-\infty}^{-1} a_k p^k + \mathbb{Z}$ (observe that for any p -adic number x only finitely many of the a_k are nonzero). Every character $\omega \in \hat{\mathbb{Q}}_p$ is of the form

$$\omega : \mathbb{Q}_p \rightarrow \mathbb{C}, \quad \omega(x) = e^{2\pi i \{xy\}_p}, \quad x \in \mathbb{Q}_p,$$

for some $y \in \mathbb{Q}_p$. In fact, the map $\mathbb{Q}_p \rightarrow \hat{\mathbb{Q}}_p$ given by mapping y to the ω defined above is a topological isomorphism. Hence the Pontryagin dual of $\mathbb{Q}_p$, $\hat{\mathbb{Q}}_p$, can be identified with $\mathbb{Q}_p$ itself. We will use this particular identification for the rest of the paper.

Recall that the *annihilator* of a closed subgroup H of a locally compact abelian group G is defined by

$$H^\perp = \{\omega \in \hat{G} : \omega(h) = 1 \text{ for all } h \in H\}.$$

In general we have topological isomorphisms $\hat{H} \cong \widehat{\hat{G}/H^\perp}$ and $\widehat{\widehat{G}/H} \cong H^\perp$. Under the identification of $\hat{\mathbb{Q}}_p$ with $\mathbb{Q}_p$, the annihilator $\mathbb{Z}_p^\perp$ of the p -adic integers $\mathbb{Z}_p$ is identified with $\mathbb{Z}_p$ itself. Hence the dual group $\hat{\mathbb{Z}}_p$ is isomorphic to $\hat{\mathbb{Q}}_p / \mathbb{Z}_p^\perp \cong \mathbb{Q}_p / \mathbb{Z}_p \cong \mathbb{Z}[1/p] / \mathbb{Z}$. There is another interesting description of $\hat{\mathbb{Z}}_p$, namely it is isomorphic to the Prüfer p -group $\mathbb{Z}(p^\infty)$ which is the subgroup of $\mathbb{T}$ consisting of all p^n -th roots of unity as n ranges over all $n = 0, 1, 2, \dots$,

$$\mathbb{Z}(p^\infty) = \{z \in \mathbb{C} : z = e^{2\pi i k p^{-n}}, \quad k = 1, \dots, p^n, \quad n \in \mathbb{N}_0\}. \quad (\text{I.2})$$

Observe that any $z \in \mathbb{Z}(p^\infty)$ can be uniquely written as $z = \exp(2\pi i \sum_{k=1}^{\infty} a_k p^{-k})$ where each $a_k \in \{0, 1, \dots, p-1\}$ and only finitely

I. Time-frequency analysis on the adeles over the rationals

many a_k are non-zero. This allows us to identify every element $z \in \mathbb{Z}(p^\infty)$ with a unique p -adic number, also to be denoted by z , given by $z = \sum_{k=1}^{\infty} a_k p^{-k}$. Every element $z \in \mathbb{Z}(p^\infty)$ defines a character $\omega_z \in \widehat{\mathbb{Z}}_p$ via

$$\omega_z : \mathbb{Z}_p \rightarrow \mathbb{C}, \quad \omega_z(x) = e^{2\pi i \{xz\}_p}, \quad x \in \mathbb{Z}_p. \quad (\text{I.3})$$

Conversely, every element in $\widehat{\mathbb{Z}}_p$ is given in this way for some $z \in \mathbb{Z}(p^\infty)$.

Restricted products. Let $(G_i)_{i \in I}$ be a collection of locally compact abelian groups for some index set I . If for each $i \in I$ the group H_i is a compact and open subgroup of G_i , then the *restricted product* of the groups $(G_i)_{i \in I}$ (with respect to the H_i) is defined to be

$$G = \prod_{i \in I}^{H_i} G_i = \left\{ (x_i)_{i \in I} \in \prod_{i \in I} G_i : x_i \in H_i \text{ for all but finitely many } i \in I \right\}.$$

The correct topology in order to turn G into a locally compact group is the one with a basis of open sets of the form $\prod_{i \in I} U_i$, where each U_i is open in G_i and $U_i = H_i$ for all but finitely many $i \in I$. This is called the *restricted product topology*. If each G_i is locally compact and each H_i is compact, then the restricted product is a locally compact group [RV99, Proposition 5-1(i)]. Moreover, every character $\omega \in \widehat{G}$ is of the form

$$\omega((x_i)_{i \in I}) = \prod_{i \in I} \omega_i(x_i),$$

where $(\omega_i)_{i \in I}$ is an element of the restricted product $\prod_{i \in I}^{H_i^\perp} \widehat{G}_i$, [RV99, Theorem 5-4]. This gives us an identification

$$\left(\prod_{i \in I}^{H_i} G_i \right)^\wedge \cong \prod_{i \in I}^{H_i^\perp} \widehat{G}_i.$$

If for each $i \in I$ the Haar measure on G_i is μ_i , then the product measure $\mu = \prod_{i \in I} \mu_i$ gives a Haar measure on the restricted product of the G_i . Typically the Haar measure on each G_i is normalized so that $\mu_i(H_i) = 1$.

The adeles. The group of *finite adeles* is the restricted product

$$\mathbb{A}_{\mathbb{Q}, \text{fin}} = \prod_{p \in \mathbb{P}}^{\mathbb{Z}_p} \mathbb{Q}_p.$$

As $\widehat{\mathbb{Q}}_p \cong \mathbb{Q}_p$ and $\mathbb{Z}_p^\perp \cong \mathbb{Z}_p$, it follows from the previous paragraph that $\mathbb{A}_{\mathbb{Q}, \text{fin}}$ is a self-dual locally compact abelian group. The group of *adeles* is

$$\mathbb{A}_{\mathbb{Q}} = \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}}.$$

It is also a self dual locally compact abelian group. We write elements of the adeles as $(x_\infty, (x_p)_p)$ where $x_\infty \in \mathbb{R}$ and $(x_p)_p \in \mathbb{A}_{\mathbb{Q}, \text{fin}}$. The Haar measure on $\mathbb{A}_{\mathbb{Q}}$ is the product measure of the Lebesgue measures on $\mathbb{R}$ and the measure on all the p -adics $\mathbb{Q}_p$ (normalized for each p so that $\mu_{\mathbb{Q}_p}(\mathbb{Z}_p) = 1$).

Every element $y = (y_\infty, (y_p)_p) \in \mathbb{A}_{\mathbb{Q}}$ defines a character $\omega_y \in \widehat{\mathbb{A}_{\mathbb{Q}}}$ via

$$\omega_y : \mathbb{A}_{\mathbb{Q}} \rightarrow \mathbb{T}, \quad x = (x_\infty, (x_p)_p) \mapsto e^{2\pi i x_\infty y_\infty} \cdot \prod_{p \in \mathbb{P}} e^{-2\pi i \{x_p y_p\}_p}. \quad (\text{I.4})$$

Moreover, every character on $\mathbb{A}_{\mathbb{Q}}$ is given in such a way by some $y \in \mathbb{A}_{\mathbb{Q}}$. The minus in the exponential for the p -adics is not necessary for this identification. It is however required for the following neat fact: we will see in a moment that the rationals $\mathbb{Q}$ can be embedded into $\mathbb{A}_{\mathbb{Q}}$ as a discrete and co-compact subgroup. By the identification of $\mathbb{A}_{\mathbb{Q}}$ with $\widehat{\mathbb{A}_{\mathbb{Q}}}$ above one has $\mathbb{Q}^\perp = \mathbb{Q}$. The identification in (I.4) is the same that is used in [Dei10, Satz 5.4.2]. In [MP05, Section 4.3.7] the identification of $\mathbb{A}_{\mathbb{Q}}$ with $\widehat{\mathbb{A}_{\mathbb{Q}}}$ is done with the minus in the exponential for the real part.

Lattices in the adeles. A subgroup H of an abelian topological group G is called a *lattice* if it is discrete and the quotient group G/H is compact. A *fundamental domain* for H in G is a Borel set $B \subseteq G$ such that every $x \in G$ can be uniquely written as $x = b + h$ where $b \in B$ and $h \in H$. Fundamental domains always exist for lattices in abelian groups [KK98, Lemma 2]. The *size* or *covolume* of a lattice H in G , denoted by $s(H)$, is the measure $\mu(B)$ of any fundamental domain B for H in G .

Naturally, there is an abundance of lattices in $\mathbb{R}$ (they are all of the form $H = \alpha\mathbb{Z}$, $\alpha \in \mathbb{R} \setminus \{0\}$). The p -adic numbers contain *no* lattices (the only co-compact subgroup in $\mathbb{Q}_p$ is $\mathbb{Q}_p$ itself, and the only discrete subgroup of $\mathbb{Q}_p$ is the trivial subgroup $\{0\}$). However, both $\mathbb{A}_{\mathbb{Q}}$ and $\mathbb{R} \times \mathbb{Q}_p$ do contain lattices.

Proposition I.2.1. *For any $\alpha \in \mathbb{R} \setminus \{0\}$ the mapping*

$$\varphi_\alpha : \mathbb{Q} \rightarrow \mathbb{A}_{\mathbb{Q}}, \quad \varphi_\alpha(q) = (\alpha q, (q)_p)$$

embeds $\mathbb{Q}$ as a lattice into $\mathbb{A}_{\mathbb{Q}}$. The set $B_\alpha = [0, |\alpha|) \times \prod_{p \in \mathbb{P}} \mathbb{Z}_p$ is a fundamental domain for $\varphi_\alpha(\mathbb{Q})$ in $\mathbb{A}_{\mathbb{Q}}$ and $s(\varphi_\alpha(\mathbb{Q})) = |\alpha|$. Moreover, under the identification of $\mathbb{A}_{\mathbb{Q}}$ with $\widehat{\mathbb{A}_{\mathbb{Q}}}$ as in (I.4), the group $\varphi_\alpha(\mathbb{Q})^\perp$ can be identified with $\varphi_{1/\alpha}(\mathbb{Q})$.

Proof. As described in, e.g., [RV99, Theorem 5-11], [Dei10, Satz 5.2.1], and [MP05, Section 4.3.7], $\varphi_1(\mathbb{Q})$ is a lattice in $\mathbb{A}_{\mathbb{Q}}$ with fundamental domain $[0, 1) \times \prod_{p \in \mathbb{P}} \mathbb{Z}_p$ and $s(\varphi_1(\mathbb{Q})) = 1$. Moreover, it is also shown that $\varphi_1(\mathbb{Q})^\perp = \varphi_1(\mathbb{Q})$. The embedding of $\mathbb{Q}$ into $\mathbb{A}_{\mathbb{Q}}$ via φ_α for $\alpha \neq 1$ corresponds to an application of the topological group automorphism $x \mapsto \alpha \cdot x$, $x \in \mathbb{R}$ on the real component of $\mathbb{A}_{\mathbb{Q}}$. It is immediate that the desired results hold. $\blacksquare$

Lattices in $\mathbb{R} \times \mathbb{Q}_p$. In a similar fashion to the construction given in equation (I.4), every $y = (y_\infty, y_p) \in \mathbb{R} \times \mathbb{Q}_p$ defines a character $\omega_y \in \widehat{\mathbb{R}} \times \widehat{\mathbb{Q}}_p$ via

$$\omega_y : \mathbb{R} \times \mathbb{Q}_p, x = (x_\infty, x_p) \mapsto e^{2\pi i(x_\infty y_\infty - \{x_p y_p\}_p)}. \quad (\text{I.5})$$

One can show that every character in $\widehat{\mathbb{R}} \times \widehat{\mathbb{Q}}_p$ is given as in (I.5) for some $y \in \mathbb{R} \times \mathbb{Q}_p$.

As for the adeles, there is an abundance of lattices in $\mathbb{R} \times \mathbb{Q}_p$. This is well-known and can be found in, e.g., [LP15].

Proposition I.2.2. *Let p be a prime number. For any $\alpha \in \mathbb{R} \setminus \{0\}$ the mapping*

$$\psi_\alpha : \mathbb{Z}[1/p] \rightarrow \mathbb{R} \times \mathbb{Q}_p, \psi_\alpha(q) = (\alpha q, q)$$

embeds $\mathbb{Z}[1/p]$ as a lattice into $\mathbb{R} \times \mathbb{Q}_p$. The set $B_\alpha = [0, |\alpha|) \times \mathbb{Z}_p$ is a fundamental domain for $\psi_\alpha(\mathbb{Z}[1/p])$ in $\mathbb{R} \times \mathbb{Q}_p$ and $s(\psi_\alpha(\mathbb{Z}[1/p])) = |\alpha|$. Moreover, under the identification of $\mathbb{R} \times \mathbb{Q}_p$ with $\widehat{\mathbb{R}} \times \widehat{\mathbb{Q}}_p$ as in (I.5), the group $\psi_\alpha(\mathbb{Z}[1/p])^\perp$ can be identified with $\psi_{1/\alpha}(\mathbb{Z}[1/p])$.

The proof is very similar in nature to that of Proposition I.2.1 and it is therefore omitted.

I.3 Feichtinger's algebra for p -adic groups and the adeles

For any locally compact abelian group the *Feichtinger algebra* S_0 [Fei81; Jak18; Los80] (sometimes denoted by $\mathbf{M}^1$) is a Banach space of functions that behaves very much like the Schwartz-Bruhat space $\mathcal{S}$ (in fact, $\mathcal{S}$ is a dense subspace of S_0 , see [Fei81, Theorem 9]). For example, S_0 is invariant under the Fourier transform and the Poisson formula holds pointwise. In this section we describe S_0 for the following concrete groups: the real line $\mathbb{R}$, the p -adic integers $\mathbb{Z}_p$, the p -adic numbers $\mathbb{Q}_p$, the group $\mathbb{R} \times \mathbb{Q}_p$, the finite adeles $\mathbb{A}_{\mathbb{Q}, \text{fin}}$, and the adeles $\mathbb{A}_{\mathbb{Q}}$.

The description below of S_0 on $\widehat{\mathbb{Q}}_p$ and $\mathbb{A}_{\mathbb{Q}}$ makes it clear that S_0 is a far larger and interesting space of functions than the Schwartz-Bruhat space on these groups. The latter consists “only” of the collection of all locally constant functions with compact support.

S_0 on the real line, $\mathbb{R}$. The Feichtinger algebra on the real line, $S_0(\mathbb{R})$, is described in detail in, e.g., [FZ98] and [Grö01]. Here we only mention the following definition. If we let g be the Gaussian function $g(x) = e^{-x^2}$, $x \in \mathbb{R}$, then

$$S_0(\mathbb{R}) = \left\{ f \in L^1(\mathbb{R}) : \int_{\mathbb{R}} \|f * E_\omega g\|_1 d\omega < \infty \right\}.$$

Here $*$ is the usual convolution of functions and $E_\omega : f(x) \mapsto e^{2\pi i \omega x} f(x)$, $x \in \mathbb{R}$ is the modulation operator. The S_0 -norm of f is given by $\|f\|_{S_0(\mathbb{R})} = \int_{\mathbb{R}} \|f * E_\omega g\|_1 d\omega$.

S_0 on the p -adic integers, $\mathbb{Z}_p$. Since $\mathbb{Z}_p$ is a compact group, it follows from, e.g., [Fei81, Remark 3] or [Jak18, Lemma 4.11], that $S_0(\mathbb{Z}_p)$ consists exactly of all continuous functions on $\mathbb{Z}_p$ that have an absolutely summable Fourier series. With the identification of the Prüfer p -group $\mathbb{Z}(p^\infty)$ as the dual group of $\mathbb{Z}_p$ (see (I.2) and (I.3)) we can describe $S_0(\mathbb{Z}_p)$ as follows:

$$S_0(\mathbb{Z}_p) = \left\{ f \in \mathbf{C}(\mathbb{Z}_p) : f(x) = \sum_{z \in \mathbb{Z}(p^\infty)} c(z) e^{2\pi i \{xz\}_p}, x \in \mathbb{Z}_p, c \in \ell^1(\mathbb{Z}(p^\infty)) \right\}.$$

Moreover, the norm on $S_0(\mathbb{Z}_p)$ is defined by $\|f\|_{S_0(\mathbb{Z}_p)} = \|c\|_1$, where f and c are related as above.

S_0 on the p -adic numbers, $\mathbb{Q}_p$. The p -adic numbers $\mathbb{Q}_p$ contain the p -adic integers $\mathbb{Z}_p$ as a compact open subgroup. A set of coset representatives Y_p of $\mathbb{Q}_p/\mathbb{Z}_p$ is given by

$$Y_p = \{y \in \mathbb{Q}_p : y = \sum_{j=1}^{\infty} c_j p^{-j}, c_j \in \{0, 1, p-1\}, c_j = 0 \text{ for all but finitely many } j\}. \quad (\text{I.6})$$

It follows by, e.g., [Fei81, Lemma 8(ii)], [Jak18, Theorem 7.7], or [Rei89, §2.9] that $S_0(\mathbb{Q}_p)$ consists exactly of all continuous functions f for which their restrictions to each of the cosets of $\mathbb{Q}_p/\mathbb{Z}_p$, the collection of functions

$$\{f_y : \mathbb{Z}_p \rightarrow \mathbb{C}, x \mapsto f(x+y), x \in \mathbb{Z}_p\}_{y \in Y_p},$$

belongs to $S_0(\mathbb{Z}_p)$ and such that $\|f\|_{S_0(\mathbb{Q}_p)} := \sum_{y \in Y_p} \|f_y\|_{S_0(\mathbb{Z}_p)} < \infty$.

The characterization of $S_0(\mathbb{Q}_p)$ in terms of the functions with absolutely convergent Fourier series allows us to describe $S_0(\mathbb{Q}_p)$ as those functions that are exactly of the form

$$f(x) = \sum_{z \in \mathbb{Z}(p^\infty)} c(y, z) e^{2\pi i \{(x-y)z\}_p} \text{ for all } x \in y + \mathbb{Z}_p \quad (\text{I.7})$$

and for all $y \in Y_p$, and where $c \in \ell^1(Y_p \times \mathbb{Z}(p^\infty))$. Moreover, the norm on $S_0(\mathbb{Q}_p)$ is equivalently defined by $\|f\|_{S_0(\mathbb{Q}_p)} = \|c\|_{\ell^1(Y_p \times \mathbb{Z}(p^\infty))}$, where f and c are related as in (I.7).

S_0 on $\mathbb{R} \times \mathbb{Q}_p$. It follows from, e.g., [Fei81, Theorem 7] or [Jak18, Theorem 7.4], that the functions in $S_0(\mathbb{R} \times \mathbb{Q}_p)$ are exactly those of the form

$$f = \sum_{j \in \mathbb{N}} f_j^{(\mathbb{R})} \otimes f_j^{(\mathbb{Q}_p)} \text{ where } f_j^{(\mathbb{R})} \in S_0(\mathbb{R}), f_j^{(\mathbb{Q}_p)} \in S_0(\mathbb{Q}_p) \quad (\text{I.8})$$

for all $j \in \mathbb{N}$ and such that $\sum_{j \in \mathbb{N}} \|f_j^{(\mathbb{R})}\|_{S_0(\mathbb{R})} \|f_j^{(\mathbb{Q}_p)}\|_{S_0(\mathbb{Q}_p)} < \infty$. The norm on $S_0(\mathbb{R} \times \mathbb{Q}_p)$ is given by

$$\|f\|_{S_0(\mathbb{R} \times \mathbb{Q}_p)} = \inf \left\{ \sum_{j \in \mathbb{N}} \|f_j^{(\mathbb{R})}\|_{S_0(\mathbb{R})} \|f_j^{(\mathbb{Q}_p)}\|_{S_0(\mathbb{Q}_p)} \right\},$$

I. Time-frequency analysis on the adeles over the rationals

where the functions f , $\{f_j^{(\mathbb{R})}\}_{j \in \mathbb{N}}$ and $\{f_j^{(\mathbb{Q}_p)}\}_{j \in \mathbb{N}}$ are related as in (I.8) and the infimum is taken over all possible representations of f as in (I.8).

S_0 on $\mathbb{A}_{\mathbb{Q}, \text{fin}}$. The finite adeles $\mathbb{A}_{\mathbb{Q}, \text{fin}}$ contain $H = \prod_{p \in \mathbb{P}} \mathbb{Z}_p$ as a compact open subgroup. A set of coset representatives Y of $\mathbb{A}_{\mathbb{Q}, \text{fin}}/H$ is given by

$$Y = \{(y_2, y_3, \dots) \in \prod_{p \in \mathbb{P}} Y_p : \text{where } Y_p \text{ is as in (I.6) and } y_p = 0 \text{ for all but finitely many } p\}.$$

By [Fei81, Lemma 8(ii)], [Jak18, Theorem 7.7], or [Rei89, §2.9] the Banach space $S_0(\mathbb{A}_{\mathbb{Q}, \text{fin}})$ consists exactly of all continuous functions f on $\mathbb{A}_{\mathbb{Q}, \text{fin}}$ for which their restriction to each of the cosets of $\mathbb{A}_{\mathbb{Q}, \text{fin}}/H$, the functions

$$\{f_y : H \mapsto \mathbb{C}, x \mapsto f(x+y), x \in H\}_{y \in Y},$$

belong to $S_0(H)$ and such that

$$\|f\|_{S_0(\mathbb{A}_{\mathbb{Q}, \text{fin}})} = \sum_{y \in Y} \|f_y\|_{S_0(H)} < \infty. \quad (\text{I.9})$$

Here $S_0(H)$ is the Banach space of continuous functions over the compact group $H = \prod_{p \in \mathbb{P}} \mathbb{Z}_p$ with absolutely convergent Fourier series.

Another characterization of $S_0(\mathbb{A}_{\mathbb{Q}, \text{fin}})$ is given in [Rei89, §9.4] in the following way: any function f in $S_0(\mathbb{A}_{\mathbb{Q}, \text{fin}})$ is exactly of the form

$$f = \sum_{j \in \mathbb{N}} \bigotimes_{p \in \mathbb{P}} f_j^{(\mathbb{Q}_p)} = \sum_{j \in \mathbb{N}} f_j^{(\mathbb{Q}_2)} \otimes f_j^{(\mathbb{Q}_3)} \otimes \dots, \text{ where } f_j^{(\mathbb{Q}_p)} \in S_0(\mathbb{Q}_p), p \in \mathbb{P}, j \in \mathbb{N}, \quad (\text{I.10})$$

and for each $j \in \mathbb{N}$ *only finitely many* of the functions $f_j^{(\mathbb{Q}_p)}$, $p \in \mathbb{P}$ are *not equal* to $\mathbf{1}_{\mathbb{Z}_p}$, and such that

$$\sum_{j \in \mathbb{N}} \prod_{p \in \mathbb{P}} \|f_j^{(\mathbb{Q}_p)}\|_{S_0(\mathbb{Q}_p)} < \infty.$$

The norm

$$\|f\|_{S_0(\mathbb{A}_{\mathbb{Q}, \text{fin}})} = \inf \left\{ \sum_{j \in \mathbb{N}} \prod_{p \in \mathbb{P}} \|f_j^{(\mathbb{Q}_p)}\|_{S_0(\mathbb{Q}_p)} \right\},$$

where the infimum is taken over all possible representation of f as in (I.10), is a norm on $S_0(\mathbb{A}_{\mathbb{Q}, \text{fin}})$ that is equivalent to the norm in (I.9).

S_0 on the adele group, $\mathbb{A}_{\mathbb{Q}}$. By definition $\mathbb{A}_{\mathbb{Q}} = \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}}$. It follows from [Fei81, Theorem 7] or [Jak18, Theorem 7.4] that a function f belongs to $S_0(\mathbb{A}_{\mathbb{Q}})$ if and only if

$$f = \sum_{j \in \mathbb{N}} \bigotimes_{p \in \{\infty\} \cup \mathbb{P}} f_j^{(\mathbb{Q}_p)} = \sum_{j \in \mathbb{N}} f_j^{(\mathbb{R})} \otimes f_j^{(\mathbb{Q}_2)} \otimes f_j^{(\mathbb{Q}_3)} \otimes \dots, \quad (\text{I.11})$$

$(\mathbb{Q}_\infty = \mathbb{R})$ where for each $j \in \mathbb{N}$ only finitely many of the functions $f_j^{(\mathbb{Q}_p)}$, $p \in \mathbb{P}$ are not equal to $\mathbb{1}_{\mathbb{Z}_p}$, and such that

$$\sum_{j \in \mathbb{N}} \prod_{p \in \{\infty\} \cup \mathbb{P}} \|f_j^{(\mathbb{Q}_p)}\|_{S_0(\mathbb{Q}_p)} < \infty.$$

Moreover, the $S_0(\mathbb{A}_\mathbb{Q})$ -norm is given by

$$\|f\|_{S_0(\mathbb{A}_\mathbb{Q})} = \inf \left\{ \sum_{j \in \mathbb{N}} \prod_{p \in \{\infty\} \cup \mathbb{P}} \|f_j^{(\mathbb{Q}_p)}\|_{S_0(\mathbb{Q}_p)} \right\},$$

where the infimum is taken over all possible representations of f as in (I.11).

I.4 Gabor frames

In this section we describe how one can construct Gabor frames for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ and $L^2(\mathbb{A}_\mathbb{Q})$ from existing Gabor frames for $L^2(\mathbb{R})$. The theory of Gabor frames is well understood, see e.g., the books [Chr16; Grö01] and the recent paper [JL16] that develops the theory of Gabor frames for general LCA groups. We give a very brief account of Gabor frames for general LCA groups before we state our main result, Theorem I.4.2.

The theory of Gabor frames for general LCA groups. For a moment let G be a general locally compact abelian group. We denote the dual group by $\widehat{G}$. The action that an element $\omega \in \widehat{G}$ has on $x \in G$ is written as $\omega(x)$. For any $x \in G$ and $\omega \in \widehat{G}$ we define the *translation* T_x and the *modulation* operator E_ω as follows:

$$T_x f(t) = f(t - x), \quad E_\omega f(t) = \omega(t) f(t), \quad t \in G.$$

The translation and modulation operators are unitary operators on $L^2(G)$ and isometries on $S_0(G)$. For convenience we define the *time-frequency* shift operator for any $\lambda = (x, \omega) \in G \times \widehat{G}$ to be

$$\pi(\lambda) = \pi(x, \omega) = E_\omega T_x.$$

Let Λ be a lattice (a discrete and co-compact subgroup) of the time-frequency domain $G \times \widehat{G}$ and let g be a function in $L^2(G)$. We let $\langle \cdot, \cdot \rangle$ denote the L^2 -inner product with the linearity in the first entry. The collection of functions $\{\pi(\lambda)g\}_{\lambda \in \Lambda} \subset L^2(G)$ is a *Gabor system*. Such a system is a *frame* for $L^2(G)$ if there exist constants $A, B > 0$ such that

$$A \|f\|_2^2 \leq \sum_{\lambda \in \Lambda} |\langle f, \pi(\lambda)g \rangle|^2 \leq B \|f\|_2^2 \quad \text{for all } f \in L^2(G).$$

Equivalently, the associated Gabor frame operator

$$S_{g, \Lambda} : L^2(G) \rightarrow L^2(G), \quad S_{g, \Lambda} f = \sum_{\lambda \in \Lambda} \langle f, \pi(\lambda)g \rangle \pi(\lambda)g$$

is well-defined, linear, bounded and invertible. The usefulness of (Gabor) frames lies in the following. If $g \in L^2(G)$ and the lattice Λ in $G \times \widehat{G}$ are such that $\{\pi(\lambda)g\}_{\lambda \in \Lambda}$ is a frame for $L^2(G)$, then there exists a (in general non-unique) function $h \in L^2(G)$ such that

$$f = \sum_{\lambda \in \Lambda} \langle f, \pi(\lambda)g \rangle \pi(\lambda)h \quad \text{for all } f \in L^2(G). \quad (\text{I.12})$$

If g and h are such that (I.12) holds, then they are called a dual pair of Gabor frame generators. For a given Gabor frame $\{\pi(\lambda)g\}_{\lambda \in \Lambda}$ the canonical choice of the function h such that (I.12) holds is the *canonical dual generator* $h = S_{g,\Lambda}^{-1}g$. It is a celebrated result of Gabor analysis that if $g \in S_0(G)$ generates a Gabor frame, then the canonical dual generator also belongs to $S_0(G)$ [GL04].

For our purposes we mention only the following result of Gabor analysis:

Lemma I.4.1 ([JL16, Theorem 6.1]). *Let G be an LCA group and let Λ be a lattice of $G \times \widehat{G}$. Two functions $g, h \in S_0(G)$ are a dual pair of Gabor frame generators for $L^2(G)$ with respect to time-frequency shifts from Λ (i.e., (I.12) holds) if and only if*

$$\langle h, \pi(\lambda^\circ)g \rangle = \text{vol}(\Lambda) \delta_{\lambda^\circ, 0} \quad \text{for all } \lambda^\circ \in \Lambda^\circ,$$

where Λ° is the adjoint lattice of Λ ,

$$\Lambda^\circ = \{\lambda^\circ \in G \times \widehat{G} : \pi(\lambda)\pi(\lambda^\circ) = \pi(\lambda^\circ)\pi(\lambda) \text{ for all } \lambda \in \Lambda\}.$$

Gabor systems in $L^2(\mathbb{R})$. Recall that for every $\omega \in \mathbb{R}$ the modulation operator E_ω is given by $E_\omega f(t) = e^{2\pi i \omega t} f(t)$. A Gabor system in $L^2(\mathbb{R})$ generated by a function $g \in L^2(\mathbb{R})$ with time-frequency shifts from the lattice $\Lambda = \alpha\mathbb{Z} \times \beta\mathbb{Z}$, $\alpha, \beta > 0$ is thus of the form

$$\{\pi(\lambda)g\}_{\lambda \in \alpha\mathbb{Z} \times \beta\mathbb{Z}} = \{E_{m\beta} T_{n\alpha} g\}_{m,n \in \mathbb{Z}} = \{t \mapsto e^{2\pi i m \beta t} g(t - n\alpha)\}_{m,n \in \mathbb{Z}}.$$

Celebrated results in time-frequency analysis include the following: (A) the Gaussian function $g(x) = e^{-\pi x^2}$ [Lyu92; Sei92] and all totally positive functions [GRS18; GS13] generate a Gabor frame for $L^2(\mathbb{R})$ whenever α and β are such that $\alpha\beta < 1$. (B) the values of α, β and γ such that the Gabor system $\{E_{m\beta} T_{n\alpha} \mathbf{1}_{[0,\gamma]}\}_{m,n \in \mathbb{Z}}$ is a frame for $L^2(\mathbb{R})$ is much more difficult to describe [DS16; Jan03].

Gabor systems in $L^2(\mathbb{R} \times \mathbb{Q}_p)$. For every $\omega = (\omega_\infty, \omega_p) \in \mathbb{R} \times \mathbb{Q}_p$ the modulation operator $E_\omega \equiv E_{\omega_\infty, \omega_p}$ on functions f over $\mathbb{R} \times \mathbb{Q}_p$ is defined by

$$E_\omega f(t_\infty, t_p) \equiv E_{\omega_\infty, \omega_p} f(t_\infty, t_p) = e^{2\pi i(\omega_\infty t_\infty - \{\omega_p t_p\}_p)} f(t_\infty, t_p), \quad (t_\infty, t_p) \in \mathbb{R} \times \mathbb{Q}_p.$$

A Gabor system generated by a function $g \in L^2(\mathbb{R} \times \mathbb{Q}_p)$ and the lattice

$$\Lambda = \psi_\alpha(\mathbb{Z}[1/p]) \times \psi_\beta(\mathbb{Z}[1/p]) = \{(\alpha q, q, \beta r, r) : q, r \in \mathbb{Z}[1/p]\}, \quad \alpha, \beta > 0$$

is thus of the form

$$\{\pi(\lambda)g\}_{\lambda \in \Lambda} = \{(t_\infty, t_p) \mapsto e^{2\pi i(\beta r t_\infty - \{r t_p\}_p)} g(t_\infty - \alpha q, t_p - q)\}_{q,r \in \mathbb{Z}[1/p]}.$$

Gabor systems in $L^2(\mathbb{A}_{\mathbb{Q}})$. For every $\omega = (\omega_{\infty}, (\omega_p)_p) \in \mathbb{A}_{\mathbb{Q}}$ the modulation operator $E_{\omega} \equiv E_{\omega_{\infty}, (\omega_p)_p}$ on functions f over $\mathbb{A}_{\mathbb{Q}}$ is defined by

$$E_{\omega_{\infty}, (\omega_p)_p} f(t_{\infty}, (t_p)_p) = e^{2\pi i \omega_{\infty} t_{\infty}} \prod_{p \in \mathbb{P}} e^{-2\pi i \{\omega_p t_p\}_p} f(t_{\infty}, (t_p)_p), \quad (t_{\infty}, (t_p)_p) \in \mathbb{A}_{\mathbb{Q}}.$$

A Gabor system generated by a function $g \in L^2(\mathbb{A}_{\mathbb{Q}})$ and a lattice

$$\Lambda = \varphi_{\alpha}(\mathbb{Q}) \times \varphi_{\beta}(\mathbb{Q}) = \{(\alpha q, (q)_p, \beta r, (r)_p) : q, r \in \mathbb{Q}\}, \quad \alpha, \beta > 0$$

is thus of the form

$$\{\pi(\lambda)g\}_{\lambda \in \Lambda} = \left\{ (t_{\infty}, (t_p)_p) \mapsto e^{2\pi i \beta r t_{\infty}} \prod_{p \in \mathbb{P}} e^{-2\pi i \{r t_p\}_p} g(t_{\infty} - \alpha q, (t_p - q)_p) \right\}_{q, r \in \mathbb{Q}}.$$

Gabor frames in $L^2(\mathbb{R} \times \mathbb{Q}_p)$ and $L^2(\mathbb{A}_{\mathbb{Q}})$. The following result describes that the construction of a Gabor frame in $L^2(\mathbb{R})$ implies that certain functions generate Gabor frames for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ and $L^2(\mathbb{A}_{\mathbb{Q}})$.

Theorem I.4.2. *Let $\alpha, \beta > 0$. For any two functions $g^{(\mathbb{R})}$ and $h^{(\mathbb{R})}$ in $S_0(\mathbb{R})$ the following statements are equivalent.*

- (i) $g^{(\mathbb{R})}$ and $h^{(\mathbb{R})}$ generate dual Gabor frames for $L^2(\mathbb{R})$ with respect to time-frequency shifts from the lattice $\alpha\mathbb{Z} \times \beta\mathbb{Z}$.
- (ii) For any $p \in \mathbb{P}$ the two functions $g = g^{(\mathbb{R})} \otimes \mathbf{1}_{\mathbb{Z}_p}$ and $h = h^{(\mathbb{R})} \otimes \mathbf{1}_{\mathbb{Z}_p}$ in $S_0(\mathbb{R} \times \mathbb{Q}_p)$ generate dual Gabor frames for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ with respect to the lattice

$$\Lambda = \psi_{\alpha}(\mathbb{Z}[1/p]) \times \psi_{\beta}(\mathbb{Z}[1/p]) = \{(\alpha q, q, \beta r, r) : q, r \in \mathbb{Z}[1/p]\} \subset \mathbb{R} \times \mathbb{Q}_p \times \mathbb{R} \times \mathbb{Q}_p.$$

- (iii) The two functions g and h in $S_0(\mathbb{A}_{\mathbb{Q}})$, defined by

$$g = g^{(\mathbb{R})} \otimes \mathbf{1}_{\mathbb{Z}_2} \otimes \mathbf{1}_{\mathbb{Z}_3} \otimes \dots \quad \text{and} \quad h = h^{(\mathbb{R})} \otimes \mathbf{1}_{\mathbb{Z}_2} \otimes \mathbf{1}_{\mathbb{Z}_3} \otimes \dots,$$

generate dual Gabor frames for $L^2(\mathbb{A}_{\mathbb{Q}})$ with respect to the lattice

$$\Lambda = \varphi_{\alpha}(\mathbb{Q}) \times \varphi_{\beta}(\mathbb{Q}) = \left\{ (\alpha q, (q)_p, \beta r, (r)_p) : q, r \in \mathbb{Q} \right\} \subset \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}} \times \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}}.$$

Corollary I.4.3. *For any $g^{(\mathbb{R})} \in S_0(\mathbb{R})$ and $\alpha, \beta > 0$ the following statements are equivalent.*

- (i) The function $g^{(\mathbb{R})}$ generates a Gabor frame for $L^2(\mathbb{R})$ with respect to the lattice $\alpha\mathbb{Z} \times \beta\mathbb{Z}$.
- (ii) For any $p \in \mathbb{P}$ the function $g = g^{(\mathbb{R})} \otimes \mathbf{1}_{\mathbb{Z}_p}$ generates a Gabor frame for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ with respect to the lattice

$$\Lambda = \psi_{\alpha}(\mathbb{Z}[1/p]) \times \psi_{\beta}(\mathbb{Z}[1/p]) = \{(\alpha q, q, \beta r, r) : q, r \in \mathbb{Z}[1/p]\} \subset \mathbb{R} \times \mathbb{Q}_p \times \mathbb{R} \times \mathbb{Q}_p.$$

(iii) The function $g \in S_0(\mathbb{A}_{\mathbb{Q}})$ defined by

$$g = g^{(\mathbb{R})} \otimes \mathbb{1}_{\mathbb{Z}_2} \otimes \mathbb{1}_{\mathbb{Z}_3} \otimes \dots,$$

generates a Gabor frame for $L^2(\mathbb{A}_{\mathbb{Q}})$ with respect to the lattice

$$\begin{aligned} \Lambda &= \varphi_{\alpha}(\mathbb{Q}) \times \varphi_{\beta}(\mathbb{Q}) \\ &= \{(\alpha q, (q)_p, \beta r, (r)_p) : q, r \in \mathbb{Q}\} \subset \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}} \times \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}}. \end{aligned}$$

Proof of Theorem I.4.2. It follows from the description of S_0 in Section I.3 that the functions g and h in ii and iii belong to $S_0(\mathbb{R} \times \mathbb{Q}_p)$ and $S_0(\mathbb{A}_{\mathbb{Q}})$, respectively. We only proof the equivalence between i and iii as the proof of the equivalence between i and ii is almost identical. Observe that $s(\Lambda) = \alpha\beta$ and that the adjoint lattice to the lattice Λ in iii is the discrete and co-compact subgroup of $\mathbb{A}_{\mathbb{Q}} \times \mathbb{A}_{\mathbb{Q}}$ given by

$$\begin{aligned} \Lambda^{\circ} &= \varphi_{1/\beta}(\mathbb{Q}) \times \varphi_{1/\alpha}(\mathbb{Q}) \\ &= \{(\beta^{-1}q, (q)_p, \alpha^{-1}r, (r)_p) : q, r \in \mathbb{Q}\} \subset \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}} \times \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}}. \end{aligned}$$

By Lemma I.4.1 the two functions g and h generate dual Gabor frames for $L^2(\mathbb{A}_{\mathbb{Q}})$ if and only if they satisfy

$$\langle h, \pi(\lambda^{\circ})g \rangle = \alpha\beta \delta_{\lambda^{\circ}, 0} \quad \text{for all } \lambda^{\circ} \in \Lambda^{\circ}. \quad (\text{I.13})$$

The tensor product form of g and h implies that (I.13) takes the form

$$\langle h^{(\mathbb{R})}, E_{\alpha^{-1}r} T_{\beta^{-1}q} g^{(\mathbb{R})} \rangle \prod_{p \in \mathbb{P}} \langle \mathbb{1}_{\mathbb{Z}_p}, E_r T_q \mathbb{1}_{\mathbb{Z}_p} \rangle = \alpha\beta \delta_{(q,r), (0,0)} \quad \text{for all } q, r \in \mathbb{Q}. \quad (\text{I.14})$$

Observe that a fraction $q \in \mathbb{Q}$ belongs to $\mathbb{Z}_p$ for all $p \in \mathbb{P}$ if and only if $q \in \mathbb{Z}$. Hence, if $q \in \mathbb{Q} \setminus \mathbb{Z}$, then, for some $p \in \mathbb{P}$ the support of the function $\mathbb{1}_{\mathbb{Z}_p}$ and $T_q \mathbb{1}_{\mathbb{Z}_p}$ is disjoint. This implies that (I.14) is satisfied for all $q \in \mathbb{Q} \setminus \mathbb{Z}$. Since modulation E_r is turned into translation T_r by the Fourier transform on $\mathbb{Q}_p$, it follows from Parsevals identity that also, for any given $r \in \mathbb{Q} \setminus \mathbb{Z}$ the inner-product

$$\langle \mathbb{1}_{\mathbb{Z}_p}, E_r \mathbb{1}_{\mathbb{Z}_p} \rangle_{L^2(\mathbb{Q}_p)} = 0$$

for some $p \in \mathbb{P}$. These two observations imply that we have verified (I.14) for all $q, r \in \mathbb{Q} \setminus \mathbb{Z}$. It remains to show that (I.14) holds for all $q, r \in \mathbb{Z}$. If $q, r \in \mathbb{Z}$, then $\langle \mathbb{1}_{\mathbb{Z}_p}, E_r T_q \mathbb{1}_{\mathbb{Z}_p} \rangle = 1$ for all $p \in \mathbb{P}$ (we have normalized the Haar measure on each $\mathbb{Q}_p$ such that $\int_{\mathbb{Q}_p} \mathbb{1}_{\mathbb{Z}_p} = 1$). This implies that we only need to verify

$$\langle h^{(\mathbb{R})}, E_{\alpha^{-1}r} T_{\beta^{-1}q} g^{(\mathbb{R})} \rangle = \alpha\beta \delta_{(q,r), (0,0)} \quad \text{for all } q, r \in \mathbb{Z}. \quad (\text{I.15})$$

Lemma I.4.1 states that (I.15) is satisfied if and only if the two Gabor system $\{E_{m\beta} T_{n\alpha} g^{(\mathbb{R})}\}_{m,n \in \mathbb{Z}}$ and $\{E_{m\beta} T_{n\alpha} h^{(\mathbb{R})}\}_{m,n \in \mathbb{Z}}$ are dual Gabor frames, which is i. ■

A Balian-Low theorem. The classical Balian-Low theorem for the Feichtinger algebra on the real numbers states that if $g \in S_0(\mathbb{R})$ and $\alpha\beta = 1$, then g cannot generate a Gabor frame for $L^2(\mathbb{R})$ over the lattice $\alpha\mathbb{Z} \times \beta\mathbb{Z} \subseteq \mathbb{R} \times \mathbb{R}$. It is a natural question to ask whether the following general statement for locally compact abelian groups holds: Suppose $g \in S_0(G)$ and Λ is a lattice in $G \times \widehat{G}$ of the form $\Gamma \times \Gamma^\perp$ with Γ a lattice in G . Then g does not generate a Gabor frame for $L^2(G)$ over Λ ?

It turns out that this does not hold for general LCA groups G and lattices Γ , see [Grö98]. However, it has been shown to hold for all lattices in second countable compactly generated LCA groups with noncompact component of the identity [KK98]. The groups $\mathbb{A}_{\mathbb{Q}}$ and $\mathbb{R} \times \mathbb{Q}_p$ are both second countable with noncompact component of the identity, but not compactly generated, so the result in [KK98] does not cover these groups. Note that the lattice $\Gamma = \varphi_\alpha(\mathbb{Q})$ in $\mathbb{A}_{\mathbb{Q}}$ has annihilator $\Gamma^\perp = \varphi_{1/\alpha}(\mathbb{Q})$. Thus, if $\alpha\beta = 1$, then $\varphi_\alpha(\mathbb{Q}) \times \varphi_\beta(\mathbb{Q}) \subseteq \mathbb{A}_{\mathbb{Q}} \times \mathbb{A}_{\mathbb{Q}}$ is a lattice of the form $\Gamma \times \Gamma^\perp$ (and similarly for the lattice $\psi_\alpha(\mathbb{Z}[1/p]) \times \psi_\beta(\mathbb{Z}[1/p]) \subseteq (\mathbb{R} \times \mathbb{Q}_p) \times (\mathbb{R} \times \mathbb{Q}_p)$).

Combining Corollary I.4.3 and the classical Balian-Low theorem for the Feichtinger algebra on $\mathbb{R}$, we obtain the following restricted Balian-Low type theorem for the groups $\mathbb{A}_{\mathbb{Q}}$ and $\mathbb{R} \times \mathbb{Q}_p$:

Proposition I.4.4. *Let $g \in S_0(\mathbb{R})$, and let $\alpha\beta = 1$. Then the following hold:*

- (i) *The function $g \otimes \mathbb{1}_{\mathbb{Z}_2} \otimes \mathbb{1}_{\mathbb{Z}_3} \otimes \cdots \in S_0(\mathbb{A}_{\mathbb{Q}})$ does not generate a Gabor frame for $L^2(\mathbb{A}_{\mathbb{Q}})$ over the lattice $\varphi_\alpha(\mathbb{Q}) \times \varphi_\beta(\mathbb{Q})$ in $\mathbb{A}_{\mathbb{Q}} \times \mathbb{A}_{\mathbb{Q}}$.*
- (ii) *The function $g \otimes \mathbb{1}_{\mathbb{Z}_p} \in S_0(\mathbb{R} \times \mathbb{Q}_p)$ does not generate a Gabor frame for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ over the lattice $\psi_\alpha(\mathbb{Z}[1/p]) \times \psi_\beta(\mathbb{Z}[1/p])$ in $(\mathbb{R} \times \mathbb{Q}_p) \times (\mathbb{R} \times \mathbb{Q}_p)$.*

The above result only holds for functions in $S_0(\mathbb{A}_{\mathbb{Q}})$ of the restricted form $g \otimes \mathbb{1}_{\mathbb{Z}_2} \otimes \mathbb{1}_{\mathbb{Z}_3} \otimes \cdots$ with $g \in S_0(\mathbb{R})$ (and analogously for $S_0(\mathbb{R} \times \mathbb{Q}_p)$). An interesting question is whether one can obtain the same conclusion for all elements of $S_0(\mathbb{A}_{\mathbb{Q}})$, and the first author is presently working on this.

Modulation spaces. Modulation spaces were invented by Feichtinger in the early 80s and can be defined on any locally compact abelian group, see e.g., [Fei03; Fei06; Fei83] and [Grö01].

It is well-known that the modulation spaces can be described using Gabor frames that are constructed with windows in S_0 [Grö01]. This characterization of the modulation spaces and the construction of the Gabor frames for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ and $L^2(\mathbb{A}_{\mathbb{Q}})$ in Theorem I.4.2 lead to the following.

Lemma I.4.5. *Let $g \in S_0(\mathbb{A}_{\mathbb{Q}})$ and Λ be as in Theorem I.4.2(iii) such that $\{\pi(\lambda)g\}_{\lambda \in \Lambda}$ is a Gabor frame for $L^2(\mathbb{A}_{\mathbb{Q}})$. The modulation space $M^{s,t}(\mathbb{A}_{\mathbb{Q}})$, $s, t \in [1, \infty]$ consists exactly of all elements $\sigma \in S_{0\text{prime}}(\mathbb{A}_{\mathbb{Q}})$ such that*

$$\|\sigma\|_{M^{s,t}(\mathbb{A}_{\mathbb{Q}})} := \left(\sum_{r \in \mathbb{Q}} \left(\sum_{q \in \mathbb{Q}} |\sigma(E_{\beta r, (r)_p} T_{\alpha q, (q)_p} g)|^s \right)^{t/s} \right)^{1/t},$$

with the obvious modification if s or t equal ∞ .

In a similar way the modulation spaces on $\mathbb{R} \times \mathbb{Q}_p$ can be defined using the Gabor frames constructed as in Theorem I.4.2(ii).

It is well-known that $S_0 \cong M^{1,1}$, $L^2 \cong M^{2,2}$, and that $S_{0\text{prime}} = M^{\infty,\infty}$.

In recent years the modulation spaces have been used successfully as spaces of symbols in the theory of pseudo-differential operators. For example, the space $M^{\infty,1}$ coincides with the *Sjöstrand class*. Among many we refer to, e.g., [Grö06; GS07].

I.4.1 Heisenberg modules

As described in [JL18], the construction of Gabor frames for $L^2(G)$ with time-frequency shifts from a closed subgroup $\Lambda \subset G \times \widehat{G}$ (where G is any locally compact abelian group and $\widehat{G}$ its dual group) is equivalent to the construction of certain projections in the twisted group C^* -algebra $C^*(\Lambda, c)$ where c denotes the cocycle coming from the Heisenberg representation [Rie88]. We state briefly some of the theory of [JL18] for the case of the Gabor frames for $L^2(\mathbb{A}_{\mathbb{Q}})$ and $L^2(\mathbb{R} \times \mathbb{Q}_p)$ constructed in Theorem I.4.2. This is the first example of a singly-generated Heisenberg module beyond the case of elementary locally compact abelian groups and it is not covered by the recent results in [JL18]. The equivalence bimodule that we use here is a suitable completion of the Feichtinger algebra on the respective groups. In contrast, the theory presented in [Rie88] and [LP13; LP15] use completions of the Schwartz–Bruhat space $\mathcal{S}(G)$ and $C_c(G)$, respectively, to construct equivalence bimodules between twisted group C^* -algebras of lattices in $G \times \widehat{G}$.

Heisenberg modules over $\mathbb{R} \times \mathbb{Q}_p$. For $\alpha, \beta > 0$ we define the following two Banach algebras:

$$\begin{aligned} \mathcal{A} &= \left\{ \mathbf{a} \in \mathcal{B}(L^2(\mathbb{R} \times \mathbb{Q}_p)) : \mathbf{a} = \sum_{q,r \in \mathbb{Z}[1/p]} a(q,r) E_{\beta r, r} T_{\alpha q, q}, a \in \ell^1(\mathbb{Z}[1/p]^2) \right\}, \\ \mathcal{B} &= \left\{ \mathbf{b} \in \mathcal{B}(L^2(\mathbb{R} \times \mathbb{Q}_p)) : \mathbf{b} = \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Z}[1/p]} b(q,r) (E_{\alpha^{-1}r, r} T_{\beta^{-1}q, q})^*, b \in \ell^1(\mathbb{Z}[1/p]^2) \right\}. \end{aligned}$$

Indeed, the norms $\|\mathbf{a}\|_{\mathcal{A}} = \|a\|_1$, $\|\mathbf{b}\|_{\mathcal{B}} = \|b\|_1$ (where $\mathbf{a}, a, \mathbf{b}$ and b are related as above) turn $\mathcal{A}$ and $\mathcal{B}$ into involutive Banach algebras with respect to composition of operators and where the involution is the L^2 -adjoint.

Observe that $\mathcal{A}$ and $\mathcal{B}$ are *not* generated by finitely many unitaries as is the case of the noncommutative 2-torus studied in relation to the construction of Gabor frames in $L^2(\mathbb{R})$ [Lue11].

Elements in $\mathcal{A}$ and $\mathcal{B}$ act on functions in $L^2(\mathbb{R} \times \mathbb{Q}_p)$ from the left and the right, respectively, by

$$\mathbf{a} \cdot f := \sum_{q,r \in \mathbb{Z}[1/p]} a(q,r) E_{\beta r, r} T_{\alpha q, q} f, \quad f \in L^2(\mathbb{R} \times \mathbb{Q}_p), \quad \mathbf{a} \in \mathcal{A},$$

$$f \cdot \mathbf{b} := \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Z}[1/p]} b(q,r) (E_{\alpha^{-1}r,r} T_{\beta^{-1}q,q})^* f, \quad f \in L^2(\mathbb{R} \times \mathbb{Q}_p), \quad \mathbf{b} \in \mathcal{B}.$$

We define an $\mathcal{A}$ - and $\mathcal{B}$ -valued inner product in the following way:

$$\begin{aligned} \bullet \langle \cdot, \cdot \rangle &: S_0(\mathbb{R} \times \mathbb{Q}_p) \times S_0(\mathbb{R} \times \mathbb{Q}_p) \rightarrow \mathcal{A}, \\ \bullet \langle f, g \rangle &= \sum_{q,r \in \mathbb{Z}[1/p]} \langle f, E_{\beta r,r} T_{\alpha q,q} g \rangle E_{\beta r,r} T_{\alpha q,q}, \\ \langle \cdot, \cdot \rangle_\bullet &: S_0(\mathbb{R} \times \mathbb{Q}_p) \times S_0(\mathbb{R} \times \mathbb{Q}_p) \rightarrow \mathcal{B}, \\ \langle f, g \rangle_\bullet &= \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Z}[1/p]} \langle g, (E_{\alpha^{-1}r,r} T_{\beta^{-1}q,q})^* f \rangle (E_{\alpha^{-1}r,r} T_{\beta^{-1}q,q})^*. \end{aligned}$$

One can show that

$$\bullet \langle f, g \rangle \cdot h = f \cdot \langle g, h \rangle_\bullet \quad \text{for all } f, g, h \in S_0(\mathbb{R} \times \mathbb{A}_{\mathbb{Q}}).$$

Denote by A and B the C^* -closures of $\mathcal{A}$ and $\mathcal{B}$ inside $\mathcal{B}(L^2(\mathbb{R} \times \mathbb{Q}_p))$, respectively. The actions and algebra-valued inner products defined give $S_0(\mathbb{R} \times \mathbb{Q}_p)$ the structure of a pre-imprimitivity $\mathcal{A}$ - $\mathcal{B}$ -bimodule. It can thus be completed into an imprimitivity A - B -bimodule, a *Heisenberg module* in the sense of Rieffel [Rie88], which sets up a Morita equivalence between A and B . It is worth noting that in this case, A is a twisted group C^* -algebra on the group $\mathbb{Z}[1/p] \times \mathbb{Z}[1/p]$. These have been termed *noncommutative solenoids* by F. Latrémolière and J. Packer and are studied in [LP13; LP15; LP17], where they prove that B is also a noncommutative solenoid.

Proposition I.4.6 ([JL18, Theorem 3.14]). *Let g, h be two functions in $S_0(\mathbb{R} \times \mathbb{Q}_p)$ and consider the lattices*

$$\Lambda = \{(\alpha q, q, \beta r, r) : q, r \in \mathbb{Z}[1/p]\} \quad \text{and} \quad \Lambda^\circ = \{(\beta^{-1}q, q, \alpha^{-1}r, r) : q, r \in \mathbb{Z}[1/p]\}$$

in $(\mathbb{R} \times \mathbb{Q}_p)^2$ with $\alpha, \beta > 0$ as in Theorem I.4.2. The following statements are equivalent.

- (i) $f = \bullet \langle f, g \rangle \cdot h$ for all $f \in S_0(\mathbb{R} \times \mathbb{Q}_p)$.
- (ii) $\langle g, h \rangle_\bullet$ is the identity operator on $L^2(\mathbb{R} \times \mathbb{Q}_p)$.
- (iii) g and h generate dual Gabor frames with respect to Λ for $L^2(\mathbb{R} \times \mathbb{Q}_p)$.
- (iv) $\bullet \langle g, h \rangle$ is an idempotent operator from $L^2(\mathbb{R} \times \mathbb{Q}_p)$ onto $V := \overline{\text{span}}\{\pi(\lambda^\circ)g\}_{\lambda^\circ \in \Lambda^\circ}$.
- (v) $f = g \cdot \langle h, f \rangle_\bullet$ for all $f \in S_0(\mathbb{R} \times \mathbb{Q}_p) \cap V$.

We close with a result on projections in $\mathcal{A}$ which follows from Theorem I.4.2 by choosing for $g^{(\mathbb{R})}$ the Gaussian $g_0^{(\mathbb{R})}(t) = e^{-\pi t^2}$.

I. Time-frequency analysis on the adeles over the rationals

Proposition I.4.7. *Let $g \in S_0(\mathbb{R} \times \mathbb{Q}_p)$ be the function defined by $g = g_0^{(\mathbb{R})} \otimes \mathbb{1}_{\mathbb{Z}_p}$, where $g_0^{(\mathbb{R})}$ is the Gaussian, and consider the lattice*

$$\Lambda = \varphi_\alpha(\mathbb{Z}[1/p]) \times \varphi_\beta(\mathbb{Z}[1/p]) = \{(\alpha q, q, \beta r, r) : q, r \in \mathbb{Z}[1/p]\} \subset \mathbb{R} \times \mathbb{Q}_p \times \mathbb{R} \times \mathbb{Q}_p.$$

Then $\bullet \langle S_{g,\Lambda}^{-1/2} g, S_{g,\Lambda}^{-1/2} g \rangle$ is a projection in $\mathcal{A}$ if and only if $\alpha\beta < 1$.

Proof. As shown in [Lue11] the construction of projections of the form $\bullet \langle g, g \rangle$ is equivalent to the construction of tight Gabor frames. Since $S_{g,\Lambda}^{-1/2} g$ generates the canonical tight Gabor frame the result follows from the result of Lyubarskii-Seip that $\{\pi(\alpha k, \beta l) g_0^{(\mathbb{R})}\}_{k,l \in \mathbb{Z}}$ is a Gabor frame if and only if $\alpha\beta < 1$. ■

Heisenberg modules over $\mathbb{A}_{\mathbb{Q}}$. For $\alpha, \beta > 0$ we define the following two Banach algebras:

$$\begin{aligned} \mathcal{A} &= \{ \mathbf{a} \in B(L^2(\mathbb{A}_{\mathbb{Q}})) : \mathbf{a} = \sum_{q,r \in \mathbb{Q}} a(q,r) E_{\beta r, (r)_p} T_{\alpha q, (q)_p}, a \in \ell^1(\mathbb{Q}^2) \}, \\ \mathcal{B} &= \{ \mathbf{b} \in B(L^2(\mathbb{A}_{\mathbb{Q}})) : \mathbf{b} = \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Q}} b(q,r) (E_{\alpha^{-1}r, (r)_p} T_{\beta^{-1}q, (q)_p})^*, b \in \ell^1(\mathbb{Q}^2) \}. \end{aligned}$$

The norms $\|\mathbf{a}\|_{\mathcal{A}} = \|a\|_1$, $\|\mathbf{b}\|_{\mathcal{B}} = \|b\|_1$ turn $\mathcal{A}$ and $\mathcal{B}$ into involutive Banach algebras with respect to composition of operators and where the involution is the L^2 -adjoint. As in the case of $\mathbb{R} \times \mathbb{Q}_p$ described before, $\mathcal{A}$ and $\mathcal{B}$ are not generated by finitely many unitaries. Elements in $\mathcal{A}$ and $\mathcal{B}$ act on functions in $L^2(\mathbb{A}_{\mathbb{Q}})$ from the left and the right, respectively, by

$$\begin{aligned} \mathbf{a} \cdot f &:= \sum_{q,r \in \mathbb{Q}} a(q,r) E_{\beta r, (r)_p} T_{\alpha q, (q)_p} f, \quad f \in L^2(\mathbb{A}_{\mathbb{Q}}), \quad \mathbf{a} \in \mathcal{A}, \\ f \cdot \mathbf{b} &:= \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Q}} b(q,r) (E_{\alpha^{-1}r, (r)_p} T_{\beta^{-1}q, (q)_p})^* f, \quad f \in L^2(\mathbb{A}_{\mathbb{Q}}), \quad \mathbf{b} \in \mathcal{B}. \end{aligned}$$

We define an $\mathcal{A}$ - and $\mathcal{B}$ -valued inner product in the following way:

$$\begin{aligned} \bullet \langle \cdot, \cdot \rangle &: S_0(\mathbb{A}_{\mathbb{Q}}) \times S_0(\mathbb{A}_{\mathbb{Q}}) \rightarrow \mathcal{A}, \\ \bullet \langle f, g \rangle &= \sum_{q,r \in \mathbb{Q}} \langle f, E_{\beta r, (r)_p} T_{\alpha q, (q)_p} g \rangle E_{\beta r, (r)_p} T_{\alpha q, (q)_p}, \\ \langle \cdot, \cdot \rangle_{\bullet} &: S_0(\mathbb{A}_{\mathbb{Q}}) \times S_0(\mathbb{A}_{\mathbb{Q}}) \rightarrow \mathcal{B}, \\ \langle f, g \rangle_{\bullet} &= \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Q}} \langle g, (E_{\alpha^{-1}r, (r)_p} T_{\beta^{-1}q, (q)_p})^* f \rangle (E_{\alpha^{-1}r, (r)_p} T_{\beta^{-1}q, (q)_p})^*. \end{aligned}$$

One can show that

$$\bullet \langle f, g \rangle \cdot h = f \cdot \langle g, h \rangle_{\bullet} \quad \text{for all } f, g, h \in S_0(\mathbb{A}_{\mathbb{Q}}).$$

In this case, as in the case with the group $\mathbb{R} \times \mathbb{Q}_p$, we obtain from $S_0(\mathbb{A}_{\mathbb{Q}})$ an imprimitivity A - B -bimodule, where A and B denote the C^* -closures of $\mathcal{A}$ and $\mathcal{B}$ in $B(L^2(\mathbb{A}_{\mathbb{Q}}))$, respectively.

Proposition I.4.8 ([JL18, Theorem 3.14]). *Let g, h be two functions in $S_0(\mathbb{A}_{\mathbb{Q}})$ and consider the lattices*

$$\Lambda = \{(\alpha q, (q)_p, \beta r, (r)_p) : q, r \in \mathbb{Q}\} \quad \text{and} \quad \Lambda^\circ = \{(\beta^{-1}q, (q)_p, \alpha^{-1}r, (r)_p) : q, r \in \mathbb{Q}\}$$

in $\mathbb{A}_{\mathbb{Q}}^2$ with $\alpha, \beta > 0$ as in Theorem I.4.2. The following statements are equivalent.

- (i) $f = \bullet \langle f, g \rangle \cdot h$ for all $f \in S_0(\mathbb{A}_{\mathbb{Q}})$.
- (ii) $\langle g, h \rangle_\bullet$ is the identity operator on $L^2(\mathbb{A}_{\mathbb{Q}})$.
- (iii) g and h generate dual Gabor frames with respect to Λ for $L^2(\mathbb{A}_{\mathbb{Q}})$.
- (iv) $\bullet \langle g, h \rangle$ is an idempotent operator from $L^2(\mathbb{A}_{\mathbb{Q}})$ onto $V := \overline{\text{span}}\{\pi(\lambda^\circ)g\}_{\lambda^\circ \in \Lambda^\circ}$.
- (v) $f = g \cdot \langle h, f \rangle_\bullet$ for all $f \in S_0(\mathbb{A}_{\mathbb{Q}}) \cap V$.

We close with a result on projections in $\mathcal{A}$ which follows from Theorem I.4.2 by choosing for $g^{(\mathbb{R})}$ the Gaussian $g_0^{(\mathbb{R})}(t) = e^{-\pi t^2}$. The proof is analogous to the one of Proposition I.4.7.

Proposition I.4.9. *Let $g \in S_0(\mathbb{A}_{\mathbb{Q}})$ be the function defined by*

$$g = g_0^{(\mathbb{R})} \otimes \mathbf{1}_{\mathbb{Z}_2} \otimes \mathbf{1}_{\mathbb{Z}_3} \otimes \dots,$$

where $g_0^{(\mathbb{R})}$ denotes the Gaussian, and consider the lattice

$$\Lambda = \varphi_\alpha(\mathbb{Q}) \times \varphi_\beta(\mathbb{Q}) = \{(\alpha q, (q)_p, \beta r, (r)_p) : q, r \in \mathbb{Q}\} \subset \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}} \times \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}}.$$

Then $\bullet \langle S_{g, \Lambda}^{-1/2} g, S_{g, \Lambda}^{-1/2} g \rangle$ is a projection in $\mathcal{A}$ if and only if $\alpha\beta < 1$.

Acknowledgments

The work of M.S.J. was carried out during the tenure of the ERCIM “Alain Bensoussan” Fellowship Programme at NTNU. This work was finished while M.S.J. and F.L. were visiting NuHAG at the Faculty of Mathematics at the University of Vienna. We are thankful for their hospitality. The first author thanks Nadia Larsen and Tron Omland for valuable discussions about the groups $\mathbb{A}_{\mathbb{Q}}$ and $\mathbb{R} \times \mathbb{Q}_p$ and for pointing out that these groups admit lattices.

References

- [AES10] Albeverio, S. A., Evdokimov, S., and Skopina, M. “ p -adic multiresolution analysis and wavelet frames”. English. In: *J. Fourier Anal. Appl.* Vol. 16, no. 5 (2010), pp. 693–714.

- [ASS18] Ahmad, O., Shah, F. A., and Sheikh, N. A. “Gabor frames on non-Archimedean fields”. In: *Int. J. Geom. Methods Mod. Phys.* Vol. 15, no. 5 (2018), pp. 1850079, 17.
- [BB18] Benedetto, J. and Benedetto, R. “Frames of translates for number-theoretic groups”. preprint, arXiv:1804.07783. 2018.
- [CG15] Christensen, O. and Goh, S. S. “Fourier-like frames on locally compact abelian groups”. In: *J. Approx. Theory* vol. 192, no. 0 (2015), pp. 82–101.
- [Chr16] Christensen, O. *An Introduction to Frames and Riesz Bases*. Second. Applied and Numerical Harmonic Analysis. Birkhäuser Basel, 2016, pp. xxv+704.
- [Dei10] Deitmar, A. *Automorphe Formen*. Springer, Heidelberg, 2010, pp. viii+250.
- [DS16] Dai, X.-R. and Sun, Q. “The *abc*-problem for Gabor systems”. In: *Mem. Amer. Math. Soc.* Vol. 244, no. 1152 (2016), pp. ix+99.
- [Fei03] Feichtinger, H. G. “Modulation spaces of locally compact Abelian groups”. In: *Proc. Internat. Conf. on Wavelets and Applications*. Ed. by Radha, R., Krishna, M., and Thangavelu, S. Chennai, January 2002: New Delhi Allied Publishers, 2003, pp. 1–56.
- [Fei06] Feichtinger, H. G. “Modulation spaces: looking back and ahead”. In: *Sampl. Theory Signal Image Process.* Vol. 5, no. 2 (2006), pp. 109–140.
- [Fei81] Feichtinger, H. G. “On a new Segal algebra”. English. In: *Monatsh. Math.* Vol. 92 (1981), pp. 269–289.
- [Fei83] Feichtinger, H. G. *Modulation spaces on locally compact Abelian groups*. Tech. rep. University of Vienna, 1983, p. 52.
- [FZ98] Feichtinger, H. G. and Zimmermann, G. “A Banach space of test functions for Gabor analysis”. English. In: *Gabor Analysis and Algorithms: Theory and Applications*. Ed. by Feichtinger, H. G. and Strohmer, T. Applied and Numerical Harmonic Analysis. Birkhäuser Boston, 1998, pp. 123–170.
- [GL04] Gröchenig, K. and Leinert, M. “Wiener’s lemma for twisted convolution and Gabor frames”. In: *J. Amer. Math. Soc.* Vol. 17, no. 1 (2004), pp. 1–18.
- [Grö01] Gröchenig, K. *Foundations of time-frequency analysis*. Applied and Numerical Harmonic Analysis. Birkhäuser Boston, Inc., Boston, MA, 2001, pp. xvi+359.
- [Grö06] Gröchenig, K. “A pedestrian’s approach to pseudodifferential operators”. In: *Harmonic Analysis and Applications*. Ed. by Heil, C. Vol. in Honor of John J. Benedetto’s 65th Birthday. Appl. Numer. Harmon. Anal. Boston, MA: Birkhäuser Boston, 2006, pp. 139–169.

-
- [Grö98] Gröchenig, K. “Aspects of Gabor analysis on locally compact abelian groups”. In: *Gabor analysis and algorithms*. Appl. Numer. Harmon. Anal. Birkhäuser Boston, Boston, MA, 1998, pp. 211–231.
 - [GRS18] Gröchenig, K., Romero, J. L., and Stöckler, J. “Sampling theorems for shift-invariant spaces, Gabor frames, and totally positive functions”. In: *Invent. Math.* Vol. 211, no. 3 (2018), pp. 1119–1148.
 - [GS07] Gröchenig, K. and Strohmer, T. “Pseudodifferential operators on locally compact abelian groups and Sjöstrand’s symbol class”. In: *J. Reine Angew. Math.* Vol. 613 (2007), pp. 121–146.
 - [GS13] Gröchenig, K. and Stöckler, J. “Gabor frames and totally positive functions”. In: *Duke Math. J.* Vol. 162, no. 6 (2013), pp. 1003–1031.
 - [HN98] Havin, V. P. and Nikol’skij, N. (*Commutative Harmonic Analysis II. Group Methods in Commutative Harmonic Analysis. Transl. from the Russian*. Berlin: Springer, 1998.
 - [HR63] Hewitt, E. and Ross, K. A. *Abstract harmonic analysis. Vol. I: Structure of topological groups. Integration theory, group representations*. Die Grundlehren der mathematischen Wissenschaften, Bd. 115. Academic Press, Inc., Publishers, New York; Springer-Verlag, Berlin-Göttingen-Heidelberg, 1963, pp. viii+519.
 - [Jak18] Jakobsen, M. S. “On a (no longer) New Segal Algebra: A Review of the Feichtinger Algebra”. In: *J. Fourier Anal. Appl.* (2018), pp. 1–82.
 - [Jan03] Janssen, A. J. E. M. “Zak transforms with few zeros and the tie”. In: *Advances in Gabor Analysis*. Ed. by Feichtinger, H. G. and Strohmer, T. Applied and Numerical Harmonic Analysis. Basel: Birkhäuser, 2003, pp. 31–70.
 - [JL16] Jakobsen, M. S. and Lemvig, J. “Density and duality theorems for regular Gabor frames”. In: *J. Funct. Anal.* Vol. 270, no. 1 (2016), pp. 229–263.
 - [JL18] Jakobsen, M. S. and Luef, F. “Duality of Gabor Frames and Heisenberg Modules”. preprint, arXiv.org:1806.05616. 2018.
 - [KK98] Kutyniok, G. and Kaniuth, E. “Zeros of the Zak transform on locally compact abelian groups”. In: *Proc. Amer. Math. Soc.* Vol. 126, no. 12 (1998), pp. 3561–3569.
 - [KOQ14] Kaliszewski, S., Omland, T., and Quigg, J. “Cuntz-Li algebras from a -adic numbers”. In: *Rev. Roumaine Math. Pures Appl.* Vol. 59, no. 3 (2014), pp. 331–370.
 - [KS10] King, E. and Skopina, M. A. “Quincunx multiresolution analysis for $L^2(\mathbf{Q}_2^2)$ ”. In: *p -Adic Numbers Ultrametric Anal. Appl.* Vol. 2, no. 3 (2010), pp. 222–231.
 - [LJ07] Li, D. and Jiang, H. “Basic results of Gabor frame on local fields”. In: *Chin. Ann. Math. Ser. B* vol. 28, no. 2 (2007), pp. 165–176.

- [LL12] Larsen, N. and Li, X. “The 2-adic ring C^* -algebra of the integers and its representations”. In: *J. Funct. Anal.* Vol. 262, no. 4 (2012), pp. 1392–1426.
- [Los80] Losert, V. “A characterization of the minimal strongly character invariant Segal algebra”. In: *Ann. Inst. Fourier (Grenoble)* vol. 30 (1980), pp. 129–139.
- [LP13] Latrémolière, F. and Packer, J. A. “Noncommutative solenoids and their projective modules”. In: *Commutative and noncommutative harmonic analysis and applications*. Vol. 603. Contemp. Math. Amer. Math. Soc., Providence, RI, 2013, pp. 35–53.
- [LP15] Latrémolière, F. and Packer, J. A. “Explicit construction of equivalence bimodules between noncommutative solenoids”. In: *Trends in harmonic analysis and its applications*. Vol. 650. Contemp. Math. Amer. Math. Soc., Providence, RI, 2015, pp. 111–140.
- [LP17] Latremoliere, F. and Packer, J. “Noncommutative solenoids and the Gromov-Hausdorff propinquity”. In: *Proc. Amer. Math. Soc.* Vol. 145 (2017), pp. 2043–2057.
- [Lue11] Luef, F. “Projections in noncommutative tori and Gabor frames”. In: *Proc. Amer. Math. Soc.* Vol. 139, no. 2 (2011), pp. 571–582.
- [Lyu92] Lyubarskii, Y. I. “Frames in the Bargmann space of entire functions.” In: *Entire and Subharmonic Functions*. Vol. 11. Adv. Sov. Math. Providence, RI: American Mathematical Society (AMS), 1992, pp. 167–180.
- [MP05] Manin, Y. and Panchishkin, A. *Introduction to modern number theory*. Second. Vol. 49. Encyclopaedia of Mathematical Sciences. Springer-Verlag, Berlin, 2005, pp. xvi+514.
- [Rei89] Reiter, H. *Metaplectic Groups and Segal Algebras*. Lect. Notes in Mathematics. Berlin: Springer, 1989.
- [Rie88] Rieffel, M. A. “Projective modules over higher-dimensional noncommutative tori”. In: *Canad. J. Math.* Vol. 40, no. 2 (1988), pp. 257–338.
- [RV99] Ramakrishnan, D. and Valenza, R. J. *Fourier Analysis on Number Fields*. New York: Springer, 1999.
- [Sei92] Seip, K. “Density theorems for sampling and interpolation in the Bargmann-Fock space. I.” In: *J. Reine Angew. Math.* Vol. 429 (1992), pp. 91–106.
- [Sha16a] Shah, F. “Periodic wavelet frames on local fields of positive characteristic”. In: *Numer. Funct. Anal. Optim.* Vol. 37, no. 5 (2016), pp. 603–627.
- [Sha16b] Shah, F. A. “Gabor frames on local fields of positive characteristic”. In: *Tbilisi Math. J.* Vol. 9, no. 2 (2016), pp. 129–139.

- [SS09] Shelkovich, V. and Skopina, M. “ p -adic Haar multiresolution analysis and pseudo-differential operators”. In: *J. Fourier Anal. Appl.* Vol. 15, no. 3 (2009), pp. 366–393.

Paper II

Heisenberg modules as function spaces

Are Austad, Ulrik Enstad

Published in *Journal of Fourier Analysis and Applications*, 2020, volume 26, issue 2. DOI: 10.1016/s00041-020-09729-7.

Abstract

Let Δ be a closed, cocompact subgroup of $G \times \widehat{G}$, where G is a second countable, locally compact abelian group. Using localization of Hilbert C^* -modules, we show that the Heisenberg module $\mathcal{E}_\Delta(G)$ over the twisted group C^* -algebra $C^*(\Delta, c)$ due to Rieffel can be continuously and densely embedded into the Hilbert space $L^2(G)$. This allows us to characterize a finite set of generators for $\mathcal{E}_\Delta(G)$ as exactly the generators of multi-window (continuous) Gabor frames over Δ , a result which was previously known only for a dense subspace of $\mathcal{E}_\Delta(G)$. We show that $\mathcal{E}_\Delta(G)$ as a function space satisfies two properties that make it eligible for time-frequency analysis: Its elements satisfy the fundamental identity of Gabor analysis if Δ is a lattice, and their associated frame operators corresponding to Δ are bounded.

II.1 Introduction

Gabor analysis concerns sets of time-frequency shifts of functions. The field has its roots in a paper by the electrical engineer and physicist Dennis Gabor [Gab46]. In this paper, the author made the claim that one could obtain basis-like representations of functions in $L^2(\mathbb{R})$ in terms of the set $\{e^{2\pi i l x} \phi(x - k) : k, l \in \mathbb{Z}\}$, where ϕ denotes a Gaussian. Today, one of the central problems of the field remains understanding the spanning and basis-like properties of sets of the form $\{e^{2\pi i \beta l x} \eta(x - \alpha k) : k, l \in \mathbb{Z}\}$ for a given $\eta \in L^2(\mathbb{R})$ and $\alpha, \beta > 0$.

Although Gabor analysis is usually carried out for functions of one or several real variables, the nature of time-frequency shifts makes it possible to generalize many aspects of the theory to the setting of a locally compact abelian group G [Grö98]. In this setting, elements of G represent time, while elements of the Pontryagin dual $\widehat{G}$ represent frequency. If $\eta \in L^2(G)$, then a time-frequency shift of η is a function of the form $\pi(x, \omega)\eta(t) = \omega(t)\eta(t - x)$ for $t, x \in G$ and $\omega \in \widehat{G}$. A Gabor system with generator η will in general be any collection of

II. Heisenberg modules as function spaces

time-frequency shifts of η . In this paper, we will allow continuous Gabor systems over any closed subgroup Δ of the time-frequency plane $G \times \widehat{G}$, which will be of the form $(\pi(z)\eta)_{z \in \Delta}$. We say that such a system forms a Gabor frame if it is a continuous frame for $L^2(G)$, which means that there exist $C, D > 0$ such that

$$C\|\xi\|_2^2 \leq \int_{\Delta} |\langle \xi, \pi(z)\eta \rangle|^2 dz \leq D\|\xi\|_2^2$$

for every $\xi \in L^2(G)$. Here, we integrate with respect to a fixed Haar measure on Δ . More generally, if $\eta_1, \dots, \eta_k \in L^2(G)$, one calls $(\pi(z)\eta_j)_{z \in \Delta, 1 \leq j \leq k}$ a multi-window Gabor frame if there exist $C, D > 0$ such that

$$C\|\xi\|_2^2 \leq \sum_{j=1}^k \int_{\Delta} |\langle \xi, \pi(z)\eta_j \rangle|^2 dz \leq D\|\xi\|_2^2$$

for all $\xi \in L^2(G)$. If Δ is a discrete subgroup of $G \times \widehat{G}$, one recovers the usual notion of a (discrete) regular Gabor frame. Here, regular means that the discrete subset Δ of $G \times \widehat{G}$ has the structure of a subgroup. A basic fact of Gabor frame theory is that $(\pi(z)\eta)_{z \in \Delta}$ is a Gabor frame if and only if the associated frame operator $S_{\eta}: L^2(G) \rightarrow L^2(G)$ is invertible. The operator is given weakly by

$$S_{\eta}\xi = \int_{\Delta} \langle \xi, \pi(z)\eta \rangle \pi(z)\eta dz$$

for $\xi \in L^2(G)$.

In [JL18; Lue09; Lue11], Luef and later Jakobsen and Luef discovered that the duality theory of regular Gabor frames is closely related to a class of imprimitivity bimodules constructed by Rieffel [Rie88]. These imprimitivity bimodules are known as *Heisenberg modules*. In general, a Hilbert C^* -module over a C^* -algebra A can be thought of as a generalized Hilbert space where the field of scalars $\mathbb{C}$ is replaced with A , and where the inner product takes values in A rather than $\mathbb{C}$. Hilbert C^* -modules were introduced by Kaplansky in [Kap53], and have since become essential in many parts of operator algebras and noncommutative geometry [Con94]. An imprimitivity A - B -bimodule is both a left Hilbert C^* -module over A and a right Hilbert C^* -module over B , with compatibility conditions on the left and right structures. If there exists an imprimitivity A - B -bimodule, then the C^* -algebras A and B are called Morita equivalent, a notion first described by Rieffel in [Rie74a; Rie74b]. Morita equivalent C^* -algebras share many important properties, such as representation theory and ideal structure.

For a closed subgroup Δ of $G \times \widehat{G}$, the Heisenberg module $\mathcal{E}_{\Delta}(G)$ can be constructed as a norm completion of the Feichtinger algebra $S_0(G)$ [Lue09]. The latter is an important space of functions in time-frequency analysis [Fei81]. The Heisenberg module implements the Morita equivalence between the twisted group C^* -algebras $C^*(\Delta, c)$ and $C^*(\Delta^{\circ}, \bar{c})$. Here, Δ° denotes the adjoint subgroup of

Δ , which consists of all points $w \in G \times \widehat{G}$ for which $\pi(w)$ commutes with $\pi(z)$ for every $z \in \Delta$. Readers familiar with Gabor analysis know that the adjoint subgroup plays a central role in results such as the fundamental identity of Gabor analysis, and this result can indeed be inferred directly from the structure of the Heisenberg modules. An important class of examples come from when $G = \mathbb{R}^n$ and Δ is a lattice in $G \times \widehat{G} \cong \mathbb{R}^{2n}$, in which case the twisted group C^* -algebras $C^*(\Delta, c)$ and $C^*(\Delta^\circ, \bar{c})$ are both noncommutative $2n$ -tori. Indeed, these examples were the original motivation for the construction of Heisenberg modules in [Rie88]. However, the construction has also been applied in other contexts, such as in the construction of finitely generated projective modules over noncommutative solenoids [EJL19; LP13; LP15].

For a general left Hilbert C^* -module $\mathcal{E}$ over a C^* -algebra A , one defines rank-one operators in analogy with the Hilbert space case. Specifically, if $\eta, \gamma \in \mathcal{E}$, the rank-one operator $\Theta_{\eta, \gamma}: \mathcal{E} \rightarrow \mathcal{E}$ is given by

$$\Theta_{\eta, \gamma} \xi = \bullet \langle \xi, \eta \rangle \gamma$$

for $\xi \in \mathcal{E}$. Here, $\bullet \langle \cdot, \cdot \rangle$ denotes the A -valued inner product on $\mathcal{E}$. A central observation in [Lue09] is that for $\eta \in S_0(G)$, the rank-one operator $\Theta_{\eta, \eta}$ associated to the Heisenberg module $\mathcal{E}_\Delta(G)$ agrees with the Gabor frame operator S_η on a dense subspace of $\mathcal{E}_\Delta(G)$, namely the Feichtinger algebra $S_0(G)$. This observation has an important consequence: It allows a finite generating set of the Heisenberg module coming from the dense subspace $S_0(G)$ to be characterized exactly as the generators of a multi-window Gabor frame over Δ [JL18, p. 14]. Moreover, such a finite generating set exists (that is, $\mathcal{E}_\Delta(G)$ is finitely generated) if and only if Δ is cocompact in $G \times \widehat{G}$ [JL18, Theorem 3.9]. However, since $\mathcal{E}_\Delta(G)$ is an abstract completion of $S_0(G)$, its elements can a priori not be interpreted as functions in any sense. Therefore, it is not straightforward to obtain a similar characterization for generators of $\mathcal{E}_\Delta(G)$ not necessarily in $S_0(G)$.

Nonetheless, it was recently remarked in [AL18] that $\mathcal{E}_\Delta(G)$ can be continuously embedded into $L^2(G)$. In the present paper, we elaborate on this embedding, and show how it arises naturally from the notion of localization of Hilbert C^* -modules as discussed in [Lan95]. The important extra structure on the Heisenberg module when localizing is a faithful trace on the C^* -algebra $C^*(\Delta, c)$. In the case that Δ is a lattice in $G \times \widehat{G}$, we use the canonical tracial state on $C^*(\Delta, c)$ (see e.g. [BO18, p. 951]). If Δ is only cocompact, we have to work a bit more, see Proposition II.3.1. It was already observed in [Lue09] that this trace plays an important role when connecting Heisenberg modules and Gabor frames. However, the consequence that the trace makes it possible to embed $\mathcal{E}_\Delta(G)$ continuously into $L^2(G)$ was first observed in [AL18].

Furthermore, in the language of localization, the rank-one operator $\Theta_{\eta, \eta}$ for $\eta \in \mathcal{E}_\Delta(G)$ extends uniquely to a bounded linear operator on $L^2(G)$, and we show in this paper that the extension is exactly the Gabor frame operator S_η (Theorem II.3.15). As a consequence, we generalize the equivalence between generators of Heisenberg modules and generators of multi-window Gabor frames to the case when the generators belong to $\mathcal{E}_\Delta(G)$ (Theorem II.3.16). We summarize some of our main results in the following.

Theorem A (cf. Proposition II.3.12, Theorem II.3.15, Theorem II.3.16). *Let G be a second countable, locally compact abelian group, and let Δ be a closed, cocompact subgroup of $G \times \widehat{G}$. Denote by $B_\Delta(G)$ the subspace of $L^2(G)$ consisting of those $\eta \in L^2(G)$ for which $(\pi(z)\eta)_{z \in \Delta}$ is a Bessel family for $L^2(G)$, that is,*

$$\int_{\Delta} |\langle \xi, \pi(z)\eta \rangle|^2 dz < \infty$$

for every $\xi \in L^2(G)$. This is a Banach space with respect to the norm

$$\|\eta\|_{B_\Delta(G)} = \|S_\eta\|^{1/2} = \sup_{\|\xi\|_2=1} \left(\int_{\Delta} |\langle \xi, \pi(z)\eta \rangle|^2 dz \right)^{1/2}.$$

The following hold:

- (i) *The Heisenberg module $\mathcal{E}_\Delta(G)$ has a concrete description as the completion of $S_0(G)$ in the Banach space $B_\Delta(G)$. The actions are given in Proposition II.3.12.*
- (ii) *For $\eta \in \mathcal{E}_\Delta(G)$, the Heisenberg module rank-one operator $\Theta_\eta: \mathcal{E}_\Delta(G) \rightarrow \mathcal{E}_\Delta(G)$ extends to the Gabor frame operator $S_\eta: L^2(G) \rightarrow L^2(G)$.*
- (iii) *Let $\eta_1, \dots, \eta_k \in \mathcal{E}_\Delta(G)$. Then $\{\eta_1, \dots, \eta_k\}$ is a generating set for $\mathcal{E}_\Delta(G)$ as a left $C^*(\Delta, c)$ -module if and only if $(\pi(z)\eta_j)_{z \in \Delta, 1 \leq j \leq k}$ is a multi-window Gabor frame for $L^2(G)$.*

Part iii of Theorem A gives a complete description of finite generating sets of the Heisenberg modules due to Rieffel, showing that they are the generators of a multi-window Gabor frame. Conversely, multi-window Gabor frames over Δ with generators in $\mathcal{E}_\Delta(G)$ give rise to finite generating sets for $\mathcal{E}_\Delta(G)$.

Note also that part i of Theorem A implies that $(\pi(z)\eta)_{z \in \Delta}$ is a Bessel family for $L^2(G)$ whenever $\eta \in \mathcal{E}_\Delta(G)$. Consequently, the Gabor analysis operator $C_\eta: L^2(G) \rightarrow L^2(\Delta)$, synthesis operator $D_\eta: L^2(\Delta) \rightarrow L^2(G)$, and frame operator $S_\eta: L^2(G) \rightarrow L^2(G)$ associated to η over Δ are all bounded linear operators. This is an attractive property of $\mathcal{E}_\Delta(G)$ as a function space in time-frequency analysis, at least when focusing on the subgroup Δ . We also show that elements of the Heisenberg module satisfy the fundamental identity of Gabor analysis over the subgroup Δ when it is a lattice (Proposition II.3.18).

We also comment on the assumption in Theorem A that Δ is cocompact. This is necessary for our localization techniques to work, see Proposition II.3.1. However, as shown in [JL16, Theorem 5.1], the existence of a multi-window Gabor frame over Δ implies that the quotient $(G \times \widehat{G})/\Delta$ is compact, i.e. Δ is a cocompact subgroup of $G \times \widehat{G}$. The assumption is therefore mild.

The paper is structured as follows: In Section II.2, we cover the necessary background material on frames in Hilbert C^* -modules, continuous Gabor frames and Heisenberg modules. In Section II.3, we introduce the notion of the localization of a Hilbert C^* -module with respect to a (possibly unbounded) trace on the coefficient algebra, and compute the localization of the Heisenberg module. We then give applications to Gabor analysis.

II.2 Preliminaries

II.2.1 Frames in Hilbert C^* -modules

In the interest of brevity, we will assume basic knowledge about C^* -algebras, Hilbert C^* -modules, imprimitivity bimodules and adjointable operators between such modules. We mention [Lan95; RW98] as references. Instead, we dedicate this section to introduce module frames.

The A -valued inner product of a left Hilbert A -module will in general be denoted by $\bullet\langle \cdot, \cdot \rangle$, while the A -valued inner product of a right Hilbert A -module will be denoted by $\langle \cdot, \cdot \rangle_\bullet$. We often refer to A as the *coefficient algebra* of $\mathcal{E}$. If $\mathcal{E}$ and $\mathcal{F}$ are left Hilbert A -modules, we use $\mathcal{L}_A(\mathcal{E}, \mathcal{F})$ to denote the Banach space of adjointable operators $\mathcal{E} \rightarrow \mathcal{F}$, or just $\mathcal{L}(\mathcal{E}, \mathcal{F})$ when there is no chance of confusion. As is standard, we write $\mathcal{L}(\mathcal{E}) = \mathcal{L}_A(\mathcal{E})$ for the C^* -algebra $\mathcal{L}_A(\mathcal{E}, \mathcal{E})$, and $\mathcal{K}(\mathcal{E}) = \mathcal{K}_A(\mathcal{E})$ for the (generalized) compact operators on $\mathcal{E}$.

For an (at most) countable index set J , we denote by $\ell^2(J, A)$ the left Hilbert A -module of all sequences $(a_j)_{j \in J}$ in A for which the sum $\sum_{j \in J} a_j a_j^*$ converges in A , with A -valued inner product

$$\bullet\langle (a_j)_{j \in J}, (b_j)_{j \in J} \rangle = \sum_{j \in J} a_j b_j^*.$$

There is an analogous way to make $\ell^2(J, A)$ into a *right* Hilbert A -module, by replacing $a_j b_j^*$ with $a_j^* b_j$ in the definition. We will work with left modules throughout this section, but obvious modifications can be made for the case of right modules as well.

We now define module frames in Hilbert A -modules, introduced in [FL02] in the case where A is unital. For a treatment of the possibly non-unital case, see [AB17].

Definition II.2.1. Let A be a C^* -algebra and $\mathcal{E}$ be a left Hilbert A -module. Furthermore, let J be some countable index set and let $(\eta_j)_{j \in J}$ be a sequence in $\mathcal{E}$. We say $(\eta_j)_{j \in J}$ is a *module frame* for $\mathcal{E}$ if there exist constants $C, D > 0$ such that

$$C \bullet\langle \xi, \xi \rangle \leq \sum_{j \in J} \bullet\langle \xi, \eta_j \rangle \bullet\langle \eta_j, \xi \rangle \leq D \bullet\langle \xi, \xi \rangle \quad (\text{II.1})$$

for all $\xi \in \mathcal{E}$, and the middle sum converges in norm. The constants C and D are called lower and upper frame bounds, respectively.

Remark II.2.2. If $A = \mathbb{C}$ in the above definition then $\mathcal{E}$ is a Hilbert space, and we recover the definition of frames in Hilbert spaces due to Duffin and Schaeffer [DS52].

Remark II.2.3. We will never treat frames over different index sets simultaneously, so to ease notation we will sometimes leave the index set implied.

Let $(\eta_j)_{j \in J}$ be a sequence in $\mathcal{E}$ that satisfies the upper frame bound condition in Definition II.2.1 but not necessarily the lower frame bound condition. Such a sequence is called a *Bessel sequence* and every constant $D > 0$ for which (II.1)

II. Heisenberg modules as function spaces

is true is called a *Bessel bound* for $(\eta_j)_{j \in J}$. To a Bessel sequence $(\eta_j)_{j \in J}$ we associate the *module analysis operator* $\Phi = \Phi_{(\eta_j)_j} : \mathcal{E} \rightarrow \ell^2(J, A)$ given by

$$\Phi\xi = (\bullet\langle\xi, \eta_j\rangle)_{j \in J} \quad (\text{II.2})$$

for $\xi \in \mathcal{E}$. It is an adjointable A -linear operator, and its adjoint $\Psi = \Psi_{(\eta_j)_j}$ is known as the *module synthesis operator*, and is given by

$$\Psi((a_j)_j) = \sum_{j \in J} a_j \cdot \eta_j, \quad (\text{II.3})$$

for $(a_j)_j \in \ell^2(J, A)$. Now let $(\gamma_j)_{j \in J}$ be another Bessel sequence. We then define the *module frame-like operator* $\Theta \in \mathcal{L}_A(\mathcal{E})$ by $\Theta = \Theta_{(\eta_j)_j, (\gamma_j)_j} := \Psi_{(\gamma_j)_j} \Phi_{(\eta_j)_j}$. That is, for all $\xi \in \mathcal{E}$ we have

$$\Theta\xi = \sum_{j \in J} \bullet\langle\xi, \eta_j\rangle \cdot \gamma_j. \quad (\text{II.4})$$

In case $(\eta_j)_j = (\gamma_j)_j$ we write $\Theta_{(\eta_j)_j} := \Theta_{(\eta_j)_j, (\eta_j)_j}$ and call it the *module frame operator* (associated to $(\eta_j)_j$). Since $\Theta_{(\eta_j)_j} = \Phi_{(\eta_j)_j}^* \Phi_{(\eta_j)_j}$, we see that $\Theta_{(\eta_j)_j}$ is always a positive operator.

A special case of the above situation is when we consider a sequence (η) consisting of a single element $\eta \in \mathcal{E}$, i.e. $|J| = 1$. It follows by the Cauchy-Schwarz inequality for Hilbert C^* -modules that such a sequence is automatically a Bessel sequence. We write $\Phi_\eta = \Phi_{(\eta)}$, $\Psi_\eta = \Psi_{(\eta)}$, $\Theta_{\eta, \gamma} = \Theta_{(\eta), (\gamma)}$ for another sequence (γ) where $\gamma \in \mathcal{E}$, and $\Theta_\eta = \Theta_{(\eta)}$. Note that in this case, $\Phi_\eta \in \mathcal{L}_A(\mathcal{E}, A)$, $\Psi_\eta \in \mathcal{L}_A(A, \mathcal{E})$ and $\Theta_{\eta, \gamma} \in \mathcal{L}_A(\mathcal{E}, \mathcal{E})$ are given by

$$\begin{aligned} \Phi_\eta\xi &= \bullet\langle\xi, \eta\rangle \\ \Psi_\eta a &= a \cdot \eta \\ \Theta_{\eta, \gamma}\xi &= \bullet\langle\xi, \eta\rangle \cdot \gamma \end{aligned}$$

for $\xi \in \mathcal{E}$, $a \in A$. Also, for a finite Bessel sequence $(\eta_1, \dots, \eta_k)$, we have that $\Phi_{(\eta_j)_{j=1}^k} = \sum_{j=1}^k \Phi_{\eta_j}$, and similar equalities for the synthesis and frame-like operators. The operator $\Theta_{\eta, \gamma}$ is often called a *rank-one operator*, and we have the following proposition, which is immediate by [RW98, Lemma 2.30, Proposition 3.8].

Proposition II.2.4. *Let η be an element of a full left Hilbert A -module $\mathcal{E}$. Then*

$$\|\eta\|_{\mathcal{E}} = \|\Theta_\eta\|_{\mathcal{L}_A(\mathcal{E})}.$$

More generally, if $\mathcal{E}$ is an imprimitivity A - B -bimodule, then

$$\|\bullet\langle\xi, \eta\rangle\|_A = \|\langle\eta, \xi\rangle_\bullet\|_B$$

for every $\xi, \eta \in \mathcal{E}$. Hence, the norm of $\mathcal{E}$ as a left Hilbert A -module coincides with the norm of $\mathcal{E}$ as a right Hilbert B -module.

The frame property of a Bessel sequence $(\eta_j)_{j \in J}$ can be characterized in terms of the invertibility of the associated frame operator $\Theta_{(\eta_j)_j}$. For a proof, see [AB17, Theorem 1.2].

Proposition II.2.5. *Let $(\eta_j)_{j \in J}$ be a Bessel sequence in $\mathcal{E}$. Then the frame operator $\Theta_{(\eta_j)_j}$ associated to $(\eta_j)_j$ is invertible if and only if $(\eta_j)_j$ is a module frame for $\mathcal{E}$.*

The following proposition shows that finite module frames are nothing more than (algebraic) generating sets, and conversely.

Proposition II.2.6. *Let $\mathcal{E}$ be a left Hilbert A -module, and let $\eta_1, \dots, \eta_k \in \mathcal{E}$. Then $(\eta_1, \dots, \eta_k)$ is a module frame for $\mathcal{E}$ if and only if it is a generating set for $\mathcal{E}$, i.e. for every $\xi \in \mathcal{E}$ there exist coefficients $a_1, \dots, a_k \in A$ such that*

$$\xi = \sum_{j=1}^k a_j \cdot \eta_j.$$

Proof. Let Θ be the module frame operator corresponding to $(\eta_j)_j$. If $(\eta_j)_j$ is a frame for $\mathcal{E}$, then by [AB17, Theorem 1.2] one has the expansion $\xi = \sum_{j=1}^k \langle \xi, \Theta^{-1} \eta_j \rangle \cdot \eta_j$ for every $\xi \in \mathcal{E}$. This shows that $(\eta_j)_j$ is a generating set for $\mathcal{E}$.

We now prove the converse. Denote by $\Phi : \mathcal{E} \rightarrow A^k$ the map $\Phi \xi = (\langle \xi, \eta_j \rangle)_{j=1}^k$. This is an adjointable A -module map, with $\Phi^*(a_j)_{j=1}^k = \sum_{j=1}^k a_j \eta_j$. By assumption Φ^* is a surjection. [Lan95, Theorem 3.2] then gives that the image of Φ is a complementable submodule of A^k . The usual Hilbert space argument then gives that $\Phi^* \Phi : \mathcal{E} \rightarrow \mathcal{E}$ is invertible, and it follows from Proposition II.2.5 that $(\eta_1, \dots, \eta_k)$ is a module frame for $\mathcal{E}$. ■

II.2.2 Gabor analysis on locally compact abelian groups

For the rest of the paper (unless stated otherwise), G will denote a second countable, locally compact abelian group with group operation written additively and with identity $0 \in G$, and Δ will denote a closed subgroup of the time-frequency plane $G \times \widehat{G}$. We fix a Haar measure on G and equip $\widehat{G}$ with the dual measure [Fol95, Theorem 4.21]. Furthermore, we pick a Haar measure on Δ , and let $(G \times \widehat{G})/\Delta$ have the unique measure such that Weil's formula holds [JL16, equation (2.4)]. We can then associate to Δ the quantity $s(\Delta) = \mu((G \times \widehat{G})/\Delta)$, known as the *size of Δ* [JL16, p. 235]. Here μ denotes the chosen Haar measure. The size of Δ is finite precisely when $(G \times \widehat{G})/\Delta$ is compact, that is, Δ is cocompact in $G \times \widehat{G}$.

Given $x \in G$ and $\omega \in \widehat{G}$, we define the translation operator T_x and modulation operator M_ω on $L^2(G)$ by

$$(T_x \xi)(t) = \xi(t - x), \quad (M_\omega \xi)(t) = \omega(t) \xi(t)$$

for $\xi \in L^2(G)$ and $t \in G$. The translation and modulation operators are unitary linear operators on $L^2(G)$. Moreover, a time-frequency shift is an operator of the form $\pi(x, \omega) = M_\omega T_x$ for $x \in G$ and $\omega \in \widehat{G}$.

The *adjoint subgroup* of Δ , denoted by Δ° , is the closed subgroup of $G \times \widehat{G}$ given by

$$\Delta^\circ = \{w \in G \times \widehat{G} : \pi(z)\pi(w) = \pi(w)\pi(z) \text{ for all } z \in \Delta\}.$$

We use the identification of Δ° with $((G \times \widehat{G})/\Delta)^\circ$ in [JL16, p. 234] to pick the dual measure on Δ° corresponding to the measure on $(G \times \widehat{G})/\Delta$ induced from the chosen measure on Δ . If Δ is cocompact in $G \times \widehat{G}$, then Δ° is discrete, and the induced measure on Δ° will be the counting measure scaled by the constant $s(\Delta)^{-1}$ [JL18, equation (13)].

We consider the two following important examples:

Example II.2.7. Suppose Δ is a lattice in $G \times \widehat{G}$, namely a discrete, cocompact subgroup of $G \times \widehat{G}$. Then Δ° is also a lattice in $G \times \widehat{G}$ [Rie88, Lemma 3.1]. In this situation, we will usually equip Δ with the counting measure. The size of Δ is then the measure of any fundamental domain for Δ in $G \times \widehat{G}$ [JL16, Remark 1]. Since Δ in particular is cocompact, the measure on Δ° will not be the counting measure in general, but rather the counting measure scaled by $s(\Delta)^{-1}$.

Example II.2.8. Let $\Delta = G \times \widehat{G}$. Δ is then cocompact in $G \times \widehat{G}$, since $(G \times \widehat{G})/\Delta$ is trivial. The natural choice of measure on Δ in this situation is the product measure coming from the chosen measure on G and the dual measure on $\widehat{G}$. The induced measure on $\Delta^\circ = \{0\}$ is then the normalized measure assigning the value 1 to $\{0\}$.

II.2.3 Gabor frames.

We will need a continuous version of Gabor frames, and so we cannot treat our Gabor frames as a special case of Definition II.2.1. However, note the similarities between the definitions and results given here and in Section II.2.1.

Given $\eta \in L^2(G)$, the family $\mathcal{G}(\eta; \Delta) = (\pi(z)\eta)_{z \in \Delta}$ is called a *Gabor system* over Δ with generator η . More generally, given $\eta_1, \dots, \eta_k \in L^2(G)$, the family $\mathcal{G}(\eta_1, \dots, \eta_k; \Delta) = (\pi(z)\eta_j)_{z \in \Delta, 1 \leq j \leq k}$ is called a *multi-window Gabor system* over Δ with generators $\eta_1, \dots, \eta_k$.

The multi-window Gabor system $\mathcal{G}(\eta_1, \dots, \eta_k; \Delta)$ is called a *multi-window Gabor frame* if it is a (*continuous*) *frame* [AAG93; JL16; Kai94] for $L^2(G)$ in the sense that both of the following hold:

- (i) The family $\mathcal{G}(\eta_1, \dots, \eta_k; \Delta)$ is weakly measurable, that is, for every $\xi \in L^2(G)$ and each $1 \leq j \leq k$, the map $z \mapsto \langle \xi, \pi(z)\eta_j \rangle$ is measurable.
- (ii) There exist positive constants $C, D > 0$ such that for all $\xi \in L^2(G)$ we have that

$$C\|\xi\|_2^2 \leq \sum_{j=1}^k \int_{\Delta} |\langle \xi, \pi(z)\eta_j \rangle|^2 dz \leq D\|\xi\|_2^2. \quad (\text{II.5})$$

The numbers C and D are called lower and upper frame bounds respectively. We may also refer to the upper frame bound as a *Bessel bound* in analogy with Section II.2. If the family $\mathcal{G}(\eta_1, \dots, \eta_k; \Delta)$ is weakly measurable and has an upper frame bound but not necessarily a lower frame bound, we call it a *Bessel family*. A (single-window) Gabor system which is a frame is called a *Gabor frame*.

The *analysis operator* associated to a Bessel family $(\pi(z)\eta)_{z \in \Delta}$ is the bounded linear operator $C_\eta: L^2(G) \rightarrow L^2(\Delta)$ given by

$$C_\eta \xi = (\langle \xi, \pi(z)\eta \rangle)_{z \in \Delta} \quad (\text{II.6})$$

for $\xi \in L^2(G)$. Its adjoint $D_\eta: L^2(\Delta) \rightarrow L^2(G)$ is called the *synthesis operator* and is given weakly by

$$D_\eta(c_z)_{z \in \Delta} = \int_{\Delta} c_z \pi(z)\eta \, dz \quad (\text{II.7})$$

for $(c_z)_{z \in \Delta} \in L^2(\Delta)$. The *frame-like operator* associated to two Bessel families $\mathcal{G}(\eta; \Delta)$ and $\mathcal{G}(\gamma; \Delta)$ is the operator $S_{\eta, \gamma} = D_\gamma C_\eta$ which is given weakly by

$$S_{\eta, \gamma} \xi = \int_{\Delta} \langle \xi, \pi(z)\eta \rangle \pi(z)\gamma \, dz \quad (\text{II.8})$$

for $\xi \in L^2(G)$. In particular, the *frame operator* associated to the Bessel family $\mathcal{G}(\eta; \Delta)$ is the operator $S_\eta := S_{\eta, \eta}$. This is a positive operator.

If $\mathcal{G}(\eta_1, \dots, \eta_k; \Delta)$ is a multi-window Gabor Bessel family, then its analysis, synthesis and frame operators are given respectively by $C = \sum_{j=1}^k C_{\eta_j}$, $D = \sum_{j=1}^k D_{\eta_j}$ and $S = \sum_{j=1}^k S_{\eta_j}$.

Note how the following proposition is analogous to Proposition II.2.5. The result is well-known in frame theory.

Proposition II.2.9. *Let $\eta_1, \dots, \eta_k \in L^2(G)$ be such that $\mathcal{G}(\eta_1, \dots, \eta_k; \Delta)$ is a Bessel family for $L^2(G)$. Then $\mathcal{G}(\eta_1, \dots, \eta_k; \Delta)$ is a multi-window Gabor frame if and only if the associated frame operator $S = \sum_{j=1}^k S_{\eta_j}$ is invertible on $L^2(G)$.*

The *Feichtinger algebra* $S_0(G)$ is the set of $\xi \in L^2(G)$ for which

$$\int_{G \times \widehat{G}} |\langle \xi, \pi(z)\xi \rangle| \, dz < \infty. \quad (\text{II.9})$$

See [Jak18] for a comprehensive introduction to $S_0(G)$. For us, the Feichtinger algebra will play a key role in the construction of Heisenberg modules as in [Lue09], see Proposition II.2.12. Note that in the original paper [Rie88], the Schwartz-Bruhat space $\mathcal{S}(G)$ was used instead. The Schwartz-Bruhat space has a more technical definition. Although it will not be important to us, we mention that the Feichtinger algebra has a natural Banach space structure [Fei81, Theorem 1].

Proposition II.2.10. *The following properties hold for the Feichtinger algebra:*

II. Heisenberg modules as function spaces

- (i) If $\eta \in S_0(G)$, then $\mathcal{G}(\eta; \Delta)$ is a Bessel family for $L^2(G)$.
- (ii) If G is discrete, then $S_0(G) = \ell^1(G)$.

For a proof of these results, see [JL16, Corollary A.5] and [Jak18, Lemma 4.11].

II.2.4 Twisted group C^* -algebras and Heisenberg modules

For the moment, let Δ be a general second countable, locally compact abelian group. A (normalized) continuous 2-cocycle on Δ is a continuous map $c: \Delta \times \Delta \rightarrow \mathbb{T}$ that satisfies the following two identities:

- (i) For every $z_1, z_2, z_3 \in \Delta$ we have that

$$c(z_1, z_2)c(z_1 + z_2, z_3) = c(z_1, z_2 + z_3)c(z_2, z_3). \quad (\text{II.10})$$

- (ii) If 0 denotes the identity element of Δ , then

$$c(0, 0) = 1. \quad (\text{II.11})$$

Note that if c is a continuous 2-cocycle, then its pointwise complex conjugate $\bar{c}$ is a continuous 2-cocycle as well.

Given a continuous 2-cocycle c on Δ , one can equip the Feichtinger algebra $S_0(\Delta)$ with a multiplication and involution as follows: For $a, b \in S_0(\Delta)$ and $z \in \Delta$, one defines

$$a * b(z) = \int_{\Delta} c(w, z - w) a(w) b(z - w) \, dw \quad (\text{II.12})$$

$$a^*(z) = \overline{c(z, -z) a(-z)}. \quad (\text{II.13})$$

The C^* -enveloping algebra of $S_0(\Delta, c)$ is called the c -twisted group C^* -algebra of Δ and is denoted by $C^*(\Delta, c)$. Note that this definition is equivalent to the usual definition of $C^*(\Delta, c)$ as the C^* -enveloping algebra of $L^1(\Delta, c)$, as $S_0(\Delta, c)$ is dense in $L^1(\Delta, c)$ and the L^1 -norm dominates the universal C^* -norm on $L^1(\Delta, c)$.

Let H be a Hilbert space, and denote by $\mathcal{U}(H)$ the unitary operators on H . A map $\pi: \Delta \rightarrow \mathcal{U}(H)$ is called a c -projective unitary representation of Δ on H if the following two properties hold:

- (i) π is strongly continuous, i.e. for every $\xi \in H$, the map $\Delta \rightarrow H, z \mapsto \pi(z)\xi$ is continuous.
- (ii) For every $z, w \in \Delta$, we have that

$$\pi(z)\pi(w) = c(z, w)\pi(z + w). \quad (\text{II.14})$$

The twisted group C^* -algebra $C^*(\Delta, c)$ captures the c -projective unitary representation theory of Δ in the following sense: For every c -projective unitary representation $\pi: \Delta \rightarrow \mathcal{U}(H)$ on a Hilbert space H , there is a nondegenerate $*$ -representation $\bar{\pi}: C^*(\Delta, c) \rightarrow \mathcal{L}(H)$ which for $a \in L^1(\Delta, c)$ is given weakly by

$$\bar{\pi}(a) = \int_{\Delta} a(z) \pi(z) \, dz. \quad (\text{II.15})$$

The above representation is called the *integrated* representation of π . Conversely, if $\Pi: C^*(\Delta, c) \rightarrow \mathcal{L}(H)$ is any nondegenerate $*$ -representation of $C^*(\Delta, c)$ on H , then there exists a unique c -projective unitary representation $\pi: \Delta \rightarrow \mathcal{U}(H)$ such that $\bar{\pi} = \Pi$. This correspondence can be seen as a consequence of e.g. [PR89, Proposition 2.7].

Note also that if π is a c -projective unitary representation, then π^* defined by $\pi^*(z) = \pi(z)^*$ is $\bar{c}$ -projective. This follows from taking the adjoint of both sides of (II.14) (it is essential that we are working with abelian groups in this situation).

When Δ is discrete, we have by Proposition II.2.10 ii that $S_0(\Delta, c) \cong \ell^1(\Delta, c)$. If we equip Δ with the counting measure, there is a canonical tracial state on $C^*(\Delta, c)$ [BO18, p. 951]. On the dense $*$ -subalgebra $\ell^1(\Delta, c)$, it is given by

$$\text{tr}(a) = a(0) \quad (\text{II.16})$$

for $a \in \ell^1(\Delta, c)$.

We now return to the situation where G is a second countable, locally compact abelian group, and Δ is a closed subgroup of $G \times \widehat{G}$. The map $c: \Delta \times \Delta \rightarrow \mathbb{T}$ given by

$$c((x, \omega), (y, \tau)) = \overline{\tau(x)} \quad (\text{II.17})$$

for $(x, \omega), (y, \tau) \in \Delta$ is a continuous 2-cocycle on Δ called the *Heisenberg 2-cocycle* [Rie88, p. 263]. Moreover, the time-frequency shifts $\pi(x, \omega) = M_{\omega} T_x$ define a c -projective unitary representation of $G \times \widehat{G}$ on $L^2(G)$, and so we have that

$$\pi(x, \omega) \pi(y, \tau) = \overline{\tau(x)} \pi(x + y, \omega \tau).$$

This representation is often called the *Heisenberg representation*. Restricting to the closed subgroup Δ of $G \times \widehat{G}$, we obtain a c -projective unitary representation of Δ on $L^2(G)$. We denote the restriction by π_{Δ} . This representation then induces a $*$ -representation of $C^*(\Delta, c)$ on $L^2(G)$, which we also (by slight abuse of notation) denote by π_{Δ} . We have the following result, see [Rie88, Proposition 2.2].

Proposition II.2.11. *The integrated representation $\pi_{\Delta}: C^*(\Delta, c) \rightarrow \mathcal{L}(L^2(G))$ is faithful, i.e. $\pi_{\Delta}(a) = 0$ implies $a = 0$ for all $a \in C^*(\Delta, c)$.*

In the following proposition, we give the definition of Heisenberg modules. For a proof, see the proof of [JL18, Theorem 3.4] or Rieffel's arguments from [Rie88] which are similar.

Proposition II.2.12. *Let G be a locally compact abelian group, and let Δ be a closed subgroup of $G \times \widehat{G}$, both with chosen Haar measures. Equip Δ° with the Haar measure determined as in Section II.2.2. The Heisenberg module $\mathcal{E}_\Delta(G)$ is an imprimitivity $C^*(\Delta, c)$ - $C^*(\Delta^\circ, \bar{c})$ -module obtained as a completion of the Feichtinger algebra $S_0(G)$. The actions and inner products are given densely as follows:*

- (i) *If $a \in S_0(\Delta, c)$, $b \in S_0(\Delta^\circ, \bar{c})$ and $\xi \in S_0(G)$, then $a \cdot \xi, \xi \cdot b \in S_0(G)$, with*

$$a \cdot \xi = \int_{\Delta} a(z) \pi(z) \xi \, dz, \quad \xi \cdot b = \int_{\Delta^\circ} b(w) \pi(w)^* \xi \, dw. \quad (\text{II.18})$$

- (ii) *If $\xi, \eta \in S_0(G)$, then $\bullet\langle \xi, \eta \rangle \in S_0(\Delta, c)$ and $\langle \xi, \eta \rangle_\bullet \in S_0(\Delta^\circ, \bar{c})$, with*

$$\bullet\langle \xi, \eta \rangle(z) = \langle \xi, \pi(z)\eta \rangle, \quad \langle \xi, \eta \rangle_\bullet(w) = \langle \pi(w)\eta, \xi \rangle \quad (\text{II.19})$$

for $z \in \Delta$ and $w \in \Delta^\circ$.

We can rewrite the left and right actions of Proposition II.2.12 as follows: Since $\pi: G \times \widehat{G} \rightarrow \mathcal{L}(L^2(G))$ is a c -projective unitary representation, it follows that π^* is $\bar{c}$ -projective. We restrict π and π^* to Δ and Δ° respectively and obtain the representations π_Δ and $\pi_{\Delta^\circ}^*$. Passing to the integrated representations, we obtain $*$ -representations of $C^*(\Delta, c)$ and $C^*(\Delta^\circ, \bar{c})$ which we also denote by π_Δ and $\pi_{\Delta^\circ}^*$ respectively. We can then write the left and right module actions given in (II.18) as

$$a \cdot \xi = \pi_\Delta(a)\xi, \quad \xi \cdot b = \pi_{\Delta^\circ}^*(b)\xi \quad (\text{II.20})$$

for $\xi \in S_0(G)$, $a \in S_0(\Delta, c)$ and $b \in S_0(\Delta^\circ, \bar{c})$.

II.3 Results

II.3.1 Localization of Hilbert C^* -modules.

We will use localization of Hilbert C^* -modules with respect to positive linear functionals as defined in [Lan95, p. 7]. Localization is a technique reminiscent of the GNS construction. It uses a positive linear functional on the coefficient algebra of a Hilbert C^* -module to embed the module continuously into a Hilbert space. The authors are not aware of many uses of localization in the literature, but an example is found in [KL12]. We will focus exclusively on the case of faithful traces, but we will need a version for (possibly) unbounded traces, which we develop after reviewing the case of finite faithful traces.

Let $\text{tr}: A \rightarrow \mathbb{C}$ denote a finite trace on A , i.e. a positive linear functional on A that satisfies $\text{tr}(a^*a) = \text{tr}(aa^*)$ for all $a \in A$. Assume also that tr is faithful, that is, $\text{tr}(a^*a) = 0$ implies $a = 0$ for all $a \in A$. If $\mathcal{E}$ is a left Hilbert A -module, it is easily verified that

$$\langle \xi, \eta \rangle_{\text{tr}} = \text{tr}(\bullet\langle \xi, \eta \rangle) \quad (\text{II.21})$$

for $\xi, \eta \in \mathcal{E}$ defines a ($\mathbb{C}$ -valued) inner product on $\mathcal{E}$, and we denote the Hilbert space completion of $\mathcal{E}$ in the norm $\|\cdot\|_{H_{\mathcal{E}}}$ coming from $\langle \cdot, \cdot \rangle_{\text{tr}}$ by $H_{\mathcal{E}}$. For $\xi \in \mathcal{E}$, the chain of inequalities

$$\|\xi\|_{H_{\mathcal{E}}}^2 = \text{tr}(\bullet \langle \xi, \xi \rangle) \leq \|\text{tr}\| \|\bullet \langle \xi, \xi \rangle\|_A = \|\text{tr}\| \|\xi\|_{\mathcal{E}}^2$$

shows that the embedding $\mathcal{E} \hookrightarrow H_{\mathcal{E}}$ is continuous. Moreover, if tr is a state, that is, $\|\text{tr}\| = 1$, then the embedding is norm-decreasing. The Hilbert space $H_{\mathcal{E}}$ is called the *localization* of $\mathcal{E}$ with respect to tr .

If $\mathcal{E}$ and $\mathcal{F}$ are left Hilbert A -modules, we obtain localizations $H_{\mathcal{E}}$ and $H_{\mathcal{F}}$ with respect to tr . Let $T: \mathcal{E} \rightarrow \mathcal{F}$ be an adjointable linear operator. Then in particular, T is a bounded linear operator when viewing the Hilbert C^* -modules as Banach spaces, and we denote its norm by $\|T\|$. For all $\xi \in \mathcal{E}$ we have that $\bullet \langle T\xi, T\xi \rangle \leq \|T\|^2 \bullet \langle \xi, \xi \rangle$ [RW98, Corollary 2.22]. Applying tr on both sides, we obtain

$$\|T\xi\|_{H_{\mathcal{F}}}^2 \leq \|T\|^2 \|\xi\|_{H_{\mathcal{E}}}^2, \quad (\text{II.22})$$

which shows that T extends to a bounded linear operator of Hilbert spaces $\bar{T}: H_{\mathcal{E}} \rightarrow H_{\mathcal{F}}$. If $\|\bar{T}\|_h$ denotes the norm of $\bar{T}$ as a Hilbert space operator, then (II.22) also shows that $\|\bar{T}\|_h \leq \|T\|$. Hence we have a norm-decreasing (hence continuous) inclusion of Banach spaces $\mathcal{L}(\mathcal{E}, \mathcal{F}) \rightarrow \mathcal{L}(H_{\mathcal{E}}, H_{\mathcal{F}})$. If $\mathcal{E} = \mathcal{F}$, then more is true: We obtain an injective $*$ -homomorphism of C^* -algebras [Lan95, p. 58] $\mathcal{L}(\mathcal{E}) \rightarrow \mathcal{L}(H_{\mathcal{E}})$. Since injective $*$ -homomorphisms of C^* -algebras are necessarily isometries [Mur90, Theorem 3.1.5], we deduce that $\mathcal{L}(\mathcal{E}) \rightarrow \mathcal{L}(H_{\mathcal{E}})$ is an isometry. Hence in this case we have

$$\|\bar{T}\|_h = \|T\| \quad (\text{II.23})$$

for all $T \in \mathcal{L}(\mathcal{E})$.

We can define the localization of a *right* Hilbert A -module $\mathcal{E}$ at a faithful trace tr similarly, except in this situation we have to set the inner product to be $\langle \xi, \eta \rangle_{\text{tr}} = \text{tr}(\langle \eta, \xi \rangle \bullet)$ for $\xi, \eta \in \mathcal{E}$ to get linearity in the first argument instead of the second. Just as with left modules, we obtain a Hilbert space $H_{\mathcal{E}}$ together with an injective bounded linear map $\mathcal{E} \hookrightarrow H_{\mathcal{E}}$.

In the following, we develop a version of localization with respect to a possibly unbounded trace that works for our purposes. Denote by A_+ the positive elements of the C^* -algebra A . By a *weight* on A , we will mean a function $\phi: A_+ \rightarrow [0, \infty]$ that satisfies $\phi(a + b) = \phi(a) + \phi(b)$ for all $a, b \in A_+$, $\phi(\lambda a) = \lambda \phi(a)$ for all $a \in A_+$ and $\lambda > 0$, and $\phi(0) = 0$. The weight ϕ is *lower semi-continuous* if whenever $(a_\alpha)_\alpha$ is a net in A_+ converging to a , then $\phi(a) \leq \liminf_\alpha \phi(a_\alpha)$. A weight ϕ on A is a *trace* if $\phi(a^*a) = \phi(aa^*)$ for all $a \in A$, and is *faithful* if $\phi(a) = 0$ implies $a = 0$ for every $a \in A_+$.

For a weight ϕ on A , let $A_+^\phi = \{a \in A_+ : \phi(a) < \infty\}$. The weight ϕ is called *densely defined* if A_+^ϕ is dense in A_+ (in the norm topology). Moreover, let $A^\phi = \text{span } A_+^\phi$. By [Ped79, Lemma 5.1.2], ϕ has a unique extension to a positive linear functional on A^ϕ , and ϕ is densely defined if and only if A^ϕ is dense in A .

II. Heisenberg modules as function spaces

A weight ϕ on A is called *finite* if $A_+^\phi = A_+$. In that case, ϕ extends uniquely to a positive linear functional on $A^\phi = \text{span } A_+^\phi = \text{span } A_+ = A$, and so we obtain a positive linear functional on the whole of A . Conversely, any positive linear functional on A restricts to a finite weight on A_+ . If A is a unital C^* -algebra, then ϕ is finite if and only if $1 \in A_+^\phi$ if and only if ϕ is densely defined.

Now let $\mathcal{E}$ be a left Hilbert A -module, and tr a (possibly unbounded) trace on A . There are two problems with localizing $\mathcal{E}$ with respect to A : The first one is that $\text{tr}(\bullet\langle\xi, \eta\rangle)$ might not be finite for $\xi, \eta \in \mathcal{E}$, which means that we do not get a well-defined inner product by setting $\langle\xi, \eta\rangle = \text{tr}(\bullet\langle\xi, \eta\rangle)$. The other problem is that we might not get a continuous embedding $\mathcal{E} \rightarrow H_{\mathcal{E}}$ even if the inner product is well-defined. However, the following set-up is sufficient for our purposes, and solves the aforementioned problems. The essential ingredient in the proof is a result due to Combes and Zettl [CZ83].

Proposition II.3.1. *Let A and B be C^* -algebras, and suppose tr_B is a faithful finite trace on B . Then the following hold:*

- (i) *If $\mathcal{E}$ is an imprimitivity A - B -bimodule, then there exists a unique lower semi-continuous trace tr_A such that*

$$\text{tr}_A(\bullet\langle\xi, \xi\rangle) = \text{tr}_B(\langle\xi, \xi\rangle\bullet) \quad (\text{II.24})$$

for all $\xi \in \mathcal{E}$. Moreover, tr_A is faithful and densely defined, with $\text{span}\{\bullet\langle\xi, \eta\rangle : \xi, \eta \in \mathcal{E}\} \subseteq A^{\text{tr}_A}$, and setting

$$\langle\xi, \eta\rangle_{\text{tr}_A} = \text{tr}_A(\bullet\langle\xi, \eta\rangle) \quad (\text{II.25})$$

for $\xi, \eta \in \mathcal{E}$ defines an inner product on $\mathcal{E}$, with $\langle\xi, \eta\rangle_{\text{tr}_A} = \langle\xi, \eta\rangle_{\text{tr}_B}$ for all $\xi, \eta \in \mathcal{E}$. Consequently, the Hilbert space obtained by completing $\mathcal{E}$ in the norm $\|\xi\|' = \text{tr}_A(\bullet\langle\xi, \xi\rangle)^{1/2}$ is just the localization of $\mathcal{E}$ with respect to tr_B .

- (ii) *If $\mathcal{E}$ and $\mathcal{F}$ are imprimitivity A - B -bimodules, then every adjointable A -linear operator $\mathcal{E} \rightarrow \mathcal{F}$ has a unique extension to a bounded linear operator $H_{\mathcal{E}} \rightarrow H_{\mathcal{F}}$. Furthermore, the map $\mathcal{L}_A(\mathcal{E}, \mathcal{F}) \rightarrow \mathcal{L}(H_{\mathcal{E}}, H_{\mathcal{F}})$ given by sending T to its unique extension is a norm-decreasing linear map of Banach spaces. Finally, if $\mathcal{E} = \mathcal{F}$, the map $\mathcal{L}_A(\mathcal{E}) \rightarrow \mathcal{L}(H_{\mathcal{E}})$ is an isometric $*$ -homomorphism of C^* -algebras.*

Proof. Suppose $\mathcal{E}$ is an imprimitivity A - B -bimodule. By [CZ83, Proposition 2.2], there is a unique lower semi-continuous trace tr_A on A such that the relation in equation (II.24) holds for all $\xi \in \mathcal{E}$. Since tr_B is finite, it is densely defined, and so tr_A is densely defined by the same proposition. The same goes for faithfulness. Since $\text{tr}_A(\bullet\langle\xi, \xi\rangle) = \text{tr}_B(\langle\xi, \xi\rangle\bullet) < \infty$, we have that $\text{span}\{\bullet\langle\xi, \xi\rangle : \xi \in \mathcal{E}\} \subseteq \text{span } A_+^{\text{tr}_A} = A^{\text{tr}_A}$. By the polarization identity for Hilbert C^* -modules, elements of the form $\bullet\langle\xi, \eta\rangle$ are in $\text{span}\{\bullet\langle\xi, \xi\rangle : \xi \in \mathcal{E}\}$, and so the unique extension of tr_A to a positive linear functional on A^{tr_A} is defined on all elements of the form $\bullet\langle\xi, \eta\rangle$ with $\xi, \eta \in \mathcal{E}$. Thus, in this situation the inner product proposed in (II.25) is well-defined. Again by the polarization identity,

the relation in (II.24) implies that $\text{tr}_A(\bullet\langle\xi, \eta\rangle) = \text{tr}_B(\langle\eta, \xi\rangle\bullet)$ for all $\xi, \eta \in \mathcal{E}$, and so $\langle\xi, \eta\rangle_{\text{tr}_A} = \langle\xi, \eta\rangle_{\text{tr}_B}$.

If $T \in \mathcal{L}(\mathcal{E}, \mathcal{F})$, then we have that $\bullet\langle T\xi, T\xi\rangle \leq \|T\|\bullet\langle\xi, \xi\rangle$ for every $\xi \in \mathcal{E}$. Taking the trace tr_A , we obtain that $\|T\xi\|_{H_{\mathcal{E}}} \leq \|T\|\|\xi\|_{H_{\mathcal{F}}}$, just as in the discussion of localization with respect to finite traces. This shows that T extends to a bounded linear map $H_{\mathcal{E}} \rightarrow H_{\mathcal{F}}$, and that the inclusion $\mathcal{L}(\mathcal{E}, \mathcal{F}) \rightarrow \mathcal{L}(H_{\mathcal{E}}, H_{\mathcal{F}})$ is norm-decreasing. In particular, if $\mathcal{E} = \mathcal{F}$, it becomes an isometric $*$ -homomorphism of C^* -algebras. $\blacksquare$

We will refer to the localization of $\mathcal{E}$ with respect to tr_B in Proposition II.3.1 above also as the localization of $\mathcal{E}$ with respect to tr_A .

Remark II.3.2. If both A and B are unital in Proposition II.3.1, then tr_A , being a densely defined trace on a unital C^* -algebra, has to be finite. In that case, we can localize $\mathcal{E}$ as a left A -module with respect to tr_A in the usual fashion, and then Proposition II.3.1 tells us that the localization is exactly the same as when done with respect to tr_B .

II.3.2 Localization of the twisted group C^* -algebra

The following proposition shows that for a discrete group Δ with a 2-cocycle c , the localization of $C^*(\Delta, c)$ as a left Hilbert module over itself with respect to the canonical trace can be identified in a natural way with $\ell^2(\Delta)$.

Proposition II.3.3. *Let Δ be a discrete group equipped with the counting measure and a 2-cocycle c . Denote by H the localization of $C^*(\Delta, c)$ as a left module over itself with respect to its canonical faithful tracial state. Then H can be identified with $\ell^2(\Delta)$ in such a way that the following diagram of inclusions commutes:*

$$\begin{array}{ccc} \ell^1(\Delta) & \hookrightarrow & C^*(\Delta, c) \\ \downarrow & & \downarrow \\ \ell^2(\Delta) & \xrightarrow{\cong} & H \end{array}$$

Moreover, the inclusion map $C^*(\Delta, c) \rightarrow \ell^2(\Delta)$ is norm-decreasing, that is, for all $a \in C^*(\Delta, c)$ we have that

$$\|a\|_{\ell^2(\Delta)} \leq \|a\|_{C^*(\Delta, c)}.$$

Proof. We have that $C^*(\Delta, c)$ is dense in H in the Hilbert space norm on H , and that $\ell^1(\Delta)$ is dense in $C^*(\Delta, c)$ in the C^* -norm on $C^*(\Delta, c)$. Since the C^* -norm on $C^*(\Delta, c)$ dominates the Hilbert space norm of H , we get that $\ell^1(\Delta)$ is also dense in H in the Hilbert space norm. Moreover $\ell^1(\Delta)$ is also dense in $\ell^2(\Delta)$ in the ℓ^2 -norm.

Denote by $\langle \cdot, \cdot \rangle$ the inner product on $\ell^2(\Delta)$. The $C^*(\Delta, c)$ -valued inner product on $C^*(\Delta, c)$ as a left Hilbert C^* -module over itself is given by $\bullet\langle a, b \rangle = ab^*$

II. Heisenberg modules as function spaces

for $a, b \in C^*(\Delta, c)$, and so the inner product with respect to tr is given by $\langle a, b \rangle_{\text{tr}} = \text{tr}(ab^*)$. If $a, b \in \ell^1(\Delta, c)$, then

$$\begin{aligned} \langle a, b \rangle_{\text{tr}} &= \text{tr}(ab^*) = (ab^*)(0) = \sum_{z \in \Delta} c(w, 0 - w) a(w) b^*(0 - w) \\ &= \sum_{z \in \Delta} c(w, -w) a(w) \overline{c(-w, w) b(w)} = \sum_{z \in \Delta} a(w) \overline{b(w)} = \langle a, b \rangle. \end{aligned}$$

This shows that $\langle \cdot, \cdot \rangle_{\text{tr}}$ and $\langle \cdot, \cdot \rangle$ agree on the subspace $\ell^1(\Delta, c)$ which is dense in both of the Hilbert spaces as argued. It follows that H can be identified with $\ell^2(\Delta)$ in such a way that the inclusions of $\ell^1(\Delta)$ into $\ell^2(\Delta)$ and $C^*(\Delta, c)$ are preserved. Moreover, since tr is a state, we have that the inclusion $C^*(\Delta, c) \hookrightarrow \ell^2(G)$ is norm-decreasing. $\blacksquare$

Remark II.3.4. In the sequel the following situation will be relevant: Let Δ be a discrete group, and denote by μ the counting measure on Δ . Let $k > 0$ be a constant. Then we can consider the C^* -algebra $C^*(\Delta, c)$ defined with respect to the measure $k\mu$ rather than μ , and so all sums involved in formulas for convolutions and norms will have a factor of k in front. In this situation there is still a faithful trace tr on $C^*(\Delta, c)$ given by $\text{tr}(a) = a(0)$ for $a \in \ell^1(\Delta, c)$. However, note that this is not a state when $k \neq 1$. Indeed, the multiplicative identity of $C^*(\Delta, c)$ is $k^{-1}\delta_0$ rather than δ_0 , and so

$$\text{tr}(1) = \text{tr}(k^{-1}\delta_0) = k^{-1}\delta_0(0) = k^{-1}.$$

If we rescale tr by k , we obtain a state.

II.3.3 Localization of the Heisenberg module

We will need a trace on the left C^* -algebra $A = C^*(\Delta, c)$ of the Heisenberg module in Proposition II.2.12. When Δ is a lattice in $G \times \widehat{G}$, we will just consider the canonical faithful trace tr_A on $C^*(\Delta, c)$. Note that by Proposition II.3.1 and Remark II.3.2, there exists a finite faithful trace on the right C^* -algebra $B = C^*(\Delta^\circ, \bar{c})$ such that $\text{tr}_A(\bullet \langle \xi, \eta \rangle) = \text{tr}_B(\langle \eta, \xi \rangle_\bullet)$ for all $\xi, \eta \in \mathcal{E}_\Delta(G)$. If $\xi, \eta \in S_0(G)$, then

$$\langle \xi, \eta \rangle = \langle \xi, \pi(0)\eta \rangle = \bullet \langle \xi, \eta \rangle (0) = \text{tr}_A(\bullet \langle \xi, \eta \rangle) = \text{tr}_B(\langle \eta, \xi \rangle_\bullet).$$

But there is a canonical trace tr'_B on B such that $\text{tr}'_B(b) = b(0)$ whenever $b \in \ell^1(\Delta^\circ, \bar{c})$. Since $\text{tr}'_B(\langle \eta, \xi \rangle_\bullet) = \langle \xi, \eta \rangle_\bullet(0) = \langle \pi(0)\xi, \eta \rangle = \langle \xi, \eta \rangle$, this shows that tr_B and tr'_B agree on $\text{span}\{\langle \xi, \eta \rangle_\bullet : \xi, \eta \in S_0(G)\}$. Since the latter is dense in B , we conclude that $\text{tr}_B = \text{tr}'_B$. Note however by Remark II.3.4 that the faithful trace tr_B which satisfies (II.25) is not a state unless $s(\Delta) = 1$.

In the case when Δ is only cocompact and not necessarily discrete, Δ° is discrete, and we obtain a (possibly unbounded) trace on $C^*(\Delta, c)$ by the following proposition. Note that we use the measures as chosen in the beginning of this section, and that B is equipped with the canonical trace that is not a state in general.

Proposition II.3.5. *Let G be a second countable, locally compact abelian group, and let Δ be a closed, cocompact subgroup of $G \times \widehat{G}$. Let $A = C^*(\Delta, c)$ and $B = C^*(\Delta^\circ, \bar{c})$. Denote by tr_B the canonical faithful trace on B as in Remark II.3.4. Then the induced trace tr_A on A via the Heisenberg module $\mathcal{E}_\Delta(G)$ as in Proposition II.3.1 is given by*

$$\text{tr}_A(\bullet \langle \xi, \eta \rangle) = \langle \xi, \eta \rangle$$

for $\xi, \eta \in S_0(G)$. In particular, if Δ is a lattice in $G \times \widehat{G}$, then tr_A is the canonical faithful tracial state on $C^*(\Delta, c)$.

Proof. By Proposition II.3.1, the induced trace tr_A satisfies

$$\text{tr}_A(\bullet \langle \xi, \eta \rangle) = \text{tr}_B(\langle \eta, \xi \rangle_\bullet)$$

for all $\xi, \eta \in \mathcal{E}_\Delta(G)$. If $\xi, \eta \in S_0(G)$, then $\langle \eta, \xi \rangle_\bullet \in S_0(\Delta^\circ, \bar{c}) = \ell^1(\Delta^\circ, \bar{c})$ by Proposition II.2.12 and Proposition II.2.10 ii, and so

$$\text{tr}_B(\langle \eta, \xi \rangle_\bullet) = \langle \eta, \xi \rangle_\bullet(0) = \langle \pi(0)\xi, \eta \rangle = \langle \xi, \eta \rangle.$$

If Δ is a lattice, then A is the twisted group C^* -algebra of a discrete group, and in this case we know that the canonical faithful tracial state tr on $C^*(\Delta, c)$ is given by $\text{tr}(a) = a(0)$ for $a \in \ell^1(\Delta, c) = S_0(\Delta, c)$. In particular, $\text{tr}(\bullet \langle \xi, \eta \rangle) = \langle \xi, \eta \rangle$. By fullness of $\mathcal{E}$ as a left Hilbert A -module, it follows that tr and tr_A agree on a dense subspace of A , hence on all of A . This shows that tr_A is indeed the faithful canonical tracial state on A . $\blacksquare$

Based on the above proposition, we make the following convention for the rest of the paper:

Convention II.3.6. *We fix a second countable, locally compact abelian group G , and a closed, cocompact subgroup Δ of $G \times \widehat{G}$. We fix Haar measures on G and Δ . If Δ is a lattice in $G \times \widehat{G}$, we assume the counting measure on Δ . From these measures, we obtain measures on $\widehat{G}$, $G \times \widehat{G}$ and Δ° as in Section II.2.2. Note that the measure on Δ° will be the counting measure scaled by a factor of $s(\Delta)^{-1}$. Let $A = C^*(\Delta, c)$ and $B = C^*(\Delta^\circ, \bar{c})$, so that the Heisenberg module $\mathcal{E}_\Delta(G)$ is an imprimitivity A - B -bimodule. We assume the canonical faithful trace tr_B on B given by $\text{tr}_B(b) = b(0)$ for $b \in \ell^1(\Delta^\circ, c)$. We equip A with the possibly unbounded trace tr_A induced from tr_B as in Proposition II.3.5. In particular, if Δ is a lattice, then tr_A is the canonical faithful tracial state on A .*

In the following proposition, we compute the localization of the Heisenberg module associated to a cocompact subgroup $\Delta \subseteq G \times \widehat{G}$.

Proposition II.3.7. *Let G denote a second countable locally compact abelian group, and let Δ be a closed, cocompact subgroup of $G \times \widehat{G}$. Then the localization H of the Heisenberg module $\mathcal{E}_\Delta(G)$ with respect to either of the traces on $C^*(\Delta, c)$ and $C^*(\Delta^\circ, \bar{c})$ can be identified with $L^2(G)$ in such a way that the diagram of inclusions commutes:*

$$\begin{array}{ccc}
 S_0(G) & \hookrightarrow & \mathcal{E}_\Delta(G) \\
 \downarrow & & \downarrow \\
 L^2(G) & \xrightarrow{\cong} & H
 \end{array}$$

Thus, the Heisenberg module can be continuously embedded into $L^2(G)$, with

$$\|\eta\|_2 \leq s(\Delta)^{1/2} \|\eta\|_{\mathcal{E}_\Delta(G)} \quad (\text{II.26})$$

for all $\eta \in \mathcal{E}_\Delta(G)$. In particular, if $(\eta_n)_n$ is a sequence in $\mathcal{E}_\Delta(G)$ that converges to an element $\eta \in \mathcal{E}_\Delta(G)$ in the $\mathcal{E}_\Delta(G)$ -norm, then $(\eta_n)_n$ also converges to η in the $L^2(G)$ -norm.

Proof. Let $\xi, \eta \in S_0(G)$. Then $\bullet\langle\xi, \eta\rangle \in S_0(\Delta, c)$ by Proposition II.2.12, and so by (II.19) and Proposition II.3.5 we obtain

$$\langle\xi, \eta\rangle_{\text{tr}_A} = \text{tr}_A(\bullet\langle\xi, \eta\rangle) = \bullet\langle\xi, \eta\rangle(0) = \langle\xi, \pi(0)\eta\rangle = \langle\xi, \eta\rangle.$$

This shows that $\langle\cdot, \cdot\rangle_{\text{tr}}$ and $\langle\cdot, \cdot\rangle$ agree on the dense subspace $S_0(G)$ of H . Hence, the localization H can be identified with $L^2(G)$ in such a way that the above diagram commutes. Moreover, since $\|\text{tr}_B\| = \text{tr}_B(1_B) = s(\Delta)$, see Remark II.3.4, we have

$$\|\eta\|_2^2 = \langle\eta, \eta\rangle = \text{tr}_B(\langle\eta, \eta\rangle_\bullet) \leq \|\text{tr}_B\| \|\langle\eta, \eta\rangle_\bullet\| = s(\Delta) \|\eta\|_{\mathcal{E}}^2.$$

This implies (II.26). ■

Proposition II.3.7 embeds the Heisenberg module as a dense subspace of $L^2(G)$, and allows us to think of $\mathcal{E}_\Delta(G)$ as a function space.

II.3.4 Applications to Gabor analysis

In light of Proposition II.3.7 and Proposition II.3.1, it follows that every adjointable $C^*(\Delta, c)$ -module operator $\mathcal{E}_\Delta(G) \rightarrow \mathcal{E}_\Delta(G)$ has a unique extension to a bounded linear map $L^2(G) \rightarrow L^2(G)$. The following lemma states that when $\eta, \gamma \in S_0(G)$, the extension of the adjointable operator $\Theta_{\eta, \gamma}$ on $\mathcal{E}_\Delta(G)$ to a bounded linear operator on $L^2(G)$ is equal to $S_{\eta, \gamma}$. This will be generalized to functions $\eta, \gamma \in \mathcal{E}_\Delta(G)$ in Theorem II.3.15. The lemma was observed in [Lue09] in the case of $G = \mathbb{R}^d$, but without using the language of localization. It was also covered in greater generality in [JL18, Theorem 3.14].

Lemma II.3.8. *Let $\eta, \gamma \in S_0(G)$. Then the module frame-like operator $\Theta_{\eta, \gamma}: \mathcal{E}_\Delta(G) \rightarrow \mathcal{E}_\Delta(G)$ extends to the Gabor frame-like operator $S_{\eta, \gamma}: L^2(G) \rightarrow L^2(G)$.*

Proof. Suppose $\eta, \gamma \in S_0(G)$. To begin with, let $\xi \in S_0(G)$. Then by Proposition II.2.12, $\bullet\langle\xi, \eta\rangle \in S_0(\Delta, c)$, and consequently $\bullet\langle\xi, \eta\rangle \cdot \gamma \in S_0(G)$. Moreover, equations (II.19) and (II.18) give that

$$\Theta_{\eta, \gamma} \xi = \bullet\langle\xi, \eta\rangle \gamma = \int_{\Delta} \bullet\langle\xi, \eta\rangle(z) \pi(z) \gamma \, dz = \int_{\Delta} \langle\xi, \pi(z) \eta\rangle \pi(z) \gamma \, dz = S_{\eta, \gamma} \xi.$$

Now let $\xi \in \mathcal{E}_{\Delta}(G)$, and suppose $(\xi_n)_n$ is a sequence in $S_0(G)$ that converges to ξ in the $\mathcal{E}_{\Delta}(G)$ -norm. Then by continuity, $\Theta_{\eta, \gamma} \xi = \lim_n \Theta_{\eta, \gamma} \xi_n$ in the $\mathcal{E}_{\Delta}(G)$ -norm. By Proposition II.3.7, the sequence $(\xi_n)_n$ also converges to ξ in the $L^2(G)$ -norm, and so by continuity, $S_{\eta, \gamma} \xi = \lim_n S_{\eta, \gamma} \xi_n$ in the $L^2(G)$ -norm. From what we already proved for functions in $S_0(G)$, we obtain that $\Theta_{\eta, \gamma} \xi = S_{\eta, \gamma} \xi$ (as elements of $L^2(G)$).

But this shows that $S_{\eta, \gamma}|_{\mathcal{E}_{\Delta}(G)} = \Theta_{\eta, \gamma}$, and since the extension of $S_{\eta, \gamma}|_{\mathcal{E}_{\Delta}(G)}$ to $L^2(G)$ is $S_{\eta, \gamma}$, we conclude that the extension of $\Theta_{\eta, \gamma}$ to $L^2(G)$ is $S_{\eta, \gamma}$. ■

The following lemma was also noted in [JL18, Lemma 3.6]. We give a different proof here which uses localization.

Lemma II.3.9. *Let $\eta \in S_0(G)$. Then the Heisenberg module norm of η can be expressed in the following ways:*

$$\|\eta\|_{\mathcal{E}_{\Delta}(G)} = \|C_{\eta}\| \quad (\text{II.27})$$

$$= \|S_{\eta}\|^{1/2} \quad (\text{II.28})$$

$$= \sup_{\|\xi\|_2=1} \left(\int_{\Delta} |\langle\xi, \pi(z) \eta\rangle|^2 \, dz \right)^{1/2} \quad (\text{II.29})$$

$$= \inf\{D^{1/2} : D \text{ is a Bessel bound for } \mathcal{G}(\eta; \Delta)\}. \quad (\text{II.30})$$

Proof. By Proposition II.2.4, the Heisenberg module norm of η is given by $\|\eta\|_{\mathcal{E}_{\Delta}(G)} = \|\Theta_{\eta}\|_{\mathcal{L}(\mathcal{E}_{\Delta}(G))}^{1/2}$. Since $\eta \in S_0(G)$, we get from Lemma II.3.8 and (II.23) that

$$\|\eta\|_{\mathcal{E}_{\Delta}(G)} = \|S_{\eta}\|_{\mathcal{L}(L^2(G))}^{1/2}.$$

Now from the equality $S_{\eta} = C_{\eta}^* C_{\eta}$ it follows that $\|S_{\eta}\|^{1/2} = \|C_{\eta}\|$. This takes care of (II.27) and (II.28). The expressions in (II.29) and (II.30) are well-known for the operator norm $\|C_{\eta}\|$. ■

We are now ready to prove the first of our main results:

Theorem II.3.10. *Let G be a second countable, locally compact abelian group, and let Δ be a closed, cocompact subgroup of $G \times \hat{G}$. If $\eta \in \mathcal{E}_{\Delta}(G)$, then $\mathcal{G}(\eta; \Delta)$ is a Bessel family for $L^2(G)$. That is, there exists a $D > 0$ such that*

$$\int_{\Delta} |\langle\xi, \pi(z) \eta\rangle|^2 \, dz \leq D \|\xi\|_2^2$$

for all $\xi \in L^2(G)$. Consequently, the analysis, synthesis and frame-like operators $C_{\eta} : L^2(G) \rightarrow L^2(\Delta)$, $D_{\eta} : L^2(\Delta) \rightarrow L^2(G)$, $S_{\eta, \gamma} : L^2(G) \rightarrow L^2(G)$ are all well-defined, bounded linear operators for $\eta, \gamma \in \mathcal{E}_{\Delta}(G)$.

II. Heisenberg modules as function spaces

Proof. Let $\eta \in \mathcal{E}_\Delta(G)$, and let $(\eta_n)_n$ be a sequence in $S_0(G)$ with

$$\lim_{n \rightarrow \infty} \|\eta - \eta_n\|_{\mathcal{E}_\Delta(G)} = 0.$$

Since $\eta_n \in S_0(G)$ for all n , $\mathcal{G}(\eta, \Delta)$ is a Bessel family for all n by Proposition II.2.10. Denote by D_n the optimal Bessel bound of $\mathcal{G}(\eta; \Delta)$ for each n , which by (II.30) in Lemma II.3.9 is equal to $\|\eta_n\|_{\mathcal{E}_\Delta(G)}^2$. Since $(\eta_n)_n$ is convergent in the Heisenberg module norm, it follows that $(\|\eta_n\|_{\mathcal{E}_\Delta(G)})_{n=1}^\infty$ is bounded, and so $(D_n)_{n=1}^\infty$ is bounded, by D say. We then have that

$$\int_\Delta |\langle \xi, \pi(z)\eta_n \rangle|^2 dz \leq D_n \|\xi\|_2^2 \leq D \|\xi\|_2^2$$

for every $\xi \in L^2(G)$ and every $n \in \mathbb{N}$. Since $(\eta_n)_n \rightarrow \eta$ in $\mathcal{E}_\Delta(G)$, we have from Proposition II.3.7 that $(\eta_n)_n \rightarrow \eta$ in $L^2(G)$ as well. Hence, continuity of the inner product gives for each $z \in \Delta$ and each $\xi \in L^2(G)$ that

$$\lim_{n \rightarrow \infty} |\langle \xi, \pi(z)\eta_n \rangle|^2 = |\langle \xi, \pi(z)\eta \rangle|^2.$$

By Fatou's lemma, we obtain for every $\xi \in L^2(G)$ that

$$\int_\Delta |\langle \xi, \pi(z)\eta \rangle|^2 dz \leq \liminf_{n \rightarrow \infty} \int_\Delta |\langle \xi, \pi(z)\eta_n \rangle|^2 dz \leq D \|\xi\|^2.$$

This proves that $\mathcal{G}(\eta; \Delta)$ is a Bessel family. ■

We are now able to extend the description of the Heisenberg module norm given in Lemma II.3.9 for functions in $S_0(G)$ to all of $\mathcal{E}_\Delta(G)$.

Proposition II.3.11. *Let $\eta \in \mathcal{E}_\Delta(G)$. Then the module norm of η can be expressed in the following ways:*

$$\|\eta\|_{\mathcal{E}_\Delta(G)} = \|C_\eta\| \tag{II.31}$$

$$= \|S_\eta\|^{1/2} \tag{II.32}$$

$$= \sup_{\|\xi\|=1} \left(\int_\Delta |\langle \xi, \pi(z)\eta \rangle|^2 dz \right)^{1/2} \tag{II.33}$$

$$= \inf\{D^{1/2} : D \text{ is a Bessel bound for } \mathcal{G}(\eta; \Delta)\}. \tag{II.34}$$

Proof. Let $\eta \in \mathcal{E}_\Delta(G)$. We will show that $\|\eta\|_{\mathcal{E}_\Delta(G)} = \|C_\eta\|$. Once this is shown, the rest of the expressions for $\|\eta\|_{\mathcal{E}_\Delta(G)}$ follow just as in the proof of Lemma II.3.9.

Let $(\eta_n)_{n=1}^\infty$ be a sequence in $S_0(G)$ such that

$$\lim_{n \rightarrow \infty} \|\eta - \eta_n\|_{\mathcal{E}_\Delta(G)} = 0.$$

Then $(\eta_n)_n$ is a Cauchy sequence in the Heisenberg module norm, and so for every $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that for all $m, n \geq N$ we have that

$$\|\eta_m - \eta_n\|_{\mathcal{E}_\Delta(G)} < \epsilon.$$

Since $\eta_n \in S_0(G)$ for all $n \in \mathbb{N}$ and $S_0(G)$ is a subspace of $L^2(G)$, we have that $\eta_m - \eta_n \in S_0(G)$ for all $m, n \in \mathbb{N}$, and so by Lemma II.3.9, we can write

$$\|\eta_m - \eta_n\|_{\mathcal{E}_\Delta(G)} = \|C_{\eta_m - \eta_n}\| = \|C_{\eta_m} - C_{\eta_n}\|.$$

But then by the above, we obtain that the sequence of operators $(C_{\eta_n})_{n=1}^\infty$ is Cauchy in $\mathcal{L}(L^2(G), L^2(\Delta))$, and so by completeness, there exists $T \in \mathcal{L}(L^2(G), L^2(\Delta))$ such that

$$\lim_{n \rightarrow \infty} \|T - C_{\eta_n}\| = 0.$$

Now fix $\xi \in L^2(G)$. Then we have that

$$\lim_{n \rightarrow \infty} \|T\xi - C_{\eta_n}\xi\|_2 = 0.$$

It is well-known that this implies the existence of a subsequence $(C_{\eta_{n_k}}\xi)_{k=1}^\infty$ that converges pointwise almost everywhere to $T\xi$ (see for instance [Rud87, Theorem 3.12]). However, since $(\eta_n)_n$ converges to η in the $L^2(G)$ -norm by Proposition II.3.7, we have that

$$\lim_{n \rightarrow \infty} C_{\eta_n}\xi(z) = \lim_{n \rightarrow \infty} \langle \xi, \pi(z)\eta_n \rangle = \langle \xi, \pi(z)\eta \rangle = C_\eta\xi(z)$$

for every $z \in \Delta$. Hence $(C_{\eta_n}\xi)_n$ converges pointwise to $C_\eta\xi$, and it follows that $(C_{\eta_{n_k}}\xi)_k$ converges pointwise to $C_\eta\xi$ as well. This shows that $C_\eta\xi = T\xi$ almost everywhere, and so they represent the same element in $L^2(\Delta)$. Since ξ was arbitrary, it follows that $C_\eta = T$, and so we have that

$$\lim_{n \rightarrow \infty} \|C_\eta - C_{\eta_n}\| = 0.$$

This implies that

$$\|\eta\|_{\mathcal{E}_\Delta(G)} = \lim_{n \rightarrow \infty} \|\eta_n\|_{\mathcal{E}_\Delta(G)} = \lim_{n \rightarrow \infty} \|C_{\eta_n}\| = \|C_\eta\|.$$

■

Let $B_\Delta(G)$ denote the set of $\eta \in L^2(G)$ such that $\mathcal{G}(\eta; \Delta)$ is a Bessel family for $L^2(G)$. Then $B_\Delta(G)$ is a Banach space when equipped with the norm

$$\|\eta\|_{B_\Delta(G)} = \|C_\eta\| = \inf\{D^{1/2} : D \text{ is a Bessel bound for } \mathcal{G}(\eta; \Delta)\}. \quad (\text{II.35})$$

By Proposition II.2.12, the Heisenberg module $\mathcal{E}_\Delta(G)$ is the completion of $S_0(G)$ with respect to the Heisenberg module norm. But by using our embedding of $\mathcal{E}_\Delta(G)$ into $L^2(G)$ in Proposition II.3.7 and the expression of the Heisenberg module norm provided in Proposition II.3.11, we obtain a concrete description of $\mathcal{E}_\Delta(G)$ as a subspace of $L^2(G)$. In the following proposition, we use the notation from (II.20).

Proposition II.3.12. *Let G be a second countable, locally compact abelian group, and let Δ be a closed, cocompact subgroup of $G \times \widehat{G}$. Then the Heisenberg module $\mathcal{E}_\Delta(G)$ is the completion of $S_0(G)$ in $B_\Delta(G)$. The bimodule structure can be described as follows: Let $a \in C^*(\Delta, c)$, $b \in C^*(\Delta^\circ, \bar{c})$ and $\xi \in \mathcal{E}_\Delta(G)$. Then*

$$a \cdot \xi = \pi_\Delta(a)\xi \quad (\text{II.36})$$

$$\xi \cdot b = \pi_{\Delta^\circ}^*(b)\xi \quad (\text{II.37})$$

Proof. By Proposition II.2.12, we know that $\mathcal{E}_\Delta(G)$ is the completion of $S_0(G)$ with respect to the Heisenberg module norm. By Proposition II.3.7, we know that $\mathcal{E}_\Delta(G)$ is continuously embedded into $L^2(G)$ in a way that respects the embedding of $S_0(G)$ into $L^2(G)$. By Proposition II.3.11, we have a description of the Heisenberg module norm as $\|\eta\|_{\mathcal{E}_\Delta(G)} = \|\eta\|_{B_\Delta(G)}$. It follows that $\mathcal{E}_\Delta(G)$ is the completion of $S_0(G)$ with respect to the norm of $B_\Delta(G)$.

To see that (II.36) holds, let $a \in C^*(\Delta, c)$ and $\xi \in \mathcal{E}_\Delta(G)$. Let $(a_n)_n$ be a sequence in $S_0(\Delta, c)$ such that $\lim_{n \rightarrow \infty} a_n = a$ in $C^*(\Delta, c)$. Let $(\xi_n)_n$ be a sequence in $S_0(G)$ such that $\lim_{n \rightarrow \infty} \xi_n = \xi$ in $\mathcal{E}_\Delta(G)$. Then by continuity of the left action of $C^*(\Delta, c)$ on $\mathcal{E}_\Delta(G)$, we have that

$$a \cdot \xi = \left(\lim_n a_n \right) \cdot \left(\lim_n \xi_n \right) = \lim_n (a_n \cdot \xi_n) = \lim_n \pi(a_n)\xi_n$$

in $\mathcal{E}_\Delta(G)$. The last equality follows from the description of $a \cdot \xi$ for $a \in S_0(\Delta, c)$ and $\xi \in S_0(G)$ as $\pi(a)\xi$ (see Proposition II.2.12). Since $\xi_n \rightarrow \xi$ in the Heisenberg module norm, we have that $\xi_n \rightarrow \xi$ in the $L^2(G)$ -norm. Also, since $\pi(a_n) \rightarrow \pi(a)$ in the operator norm, we have that $\pi(a_n)\xi_n \rightarrow \pi(a)\xi$ in the $L^2(G)$ -norm. Hence, interchanging the $\mathcal{E}_\Delta(G)$ -limit in the equation above with an $L^2(G)$ -limit, we obtain that $a \cdot \xi = \pi_\Delta(a)\xi$.

The argument for (II.37) is similar, as for $b \in S_0(\Delta^\circ, \bar{c})$ and $\xi \in S_0(G)$, the definition of $\xi \cdot b$ in Proposition II.2.12 is equal to $\pi_{\Delta^\circ}^*(b)\xi$. A similar approximation argument to the one above shows that $\xi \cdot b = \pi_{\Delta^\circ}^*(b)\xi$ also holds for $b \in C^*(\Delta^\circ, \bar{c})$ and $\xi \in \mathcal{E}_\Delta(G)$. ■

Example II.3.13. If one sets $\Delta = G \times \widehat{G}$, the Heisenberg module $\mathcal{E}_\Delta(G)$ is all of $L^2(G)$. To see this, note that $\Delta^\circ = \{0\}$. Thus, we have the identification $C^*(\Delta^\circ, \bar{c}) \cong \mathbb{C}$, where a sequence $a \in C^*(\Delta^\circ, \bar{c}) = \mathbb{C}\Delta^\circ$ is identified with its value $a(0)$ at 0. In this situation, the Heisenberg module $\mathcal{E}_\Delta(G)$ is a $C^*(\Delta, c)$ - $\mathbb{C}$ -imprimitivity bimodule. But then $\mathcal{E}_\Delta(G)$ is a right Hilbert C^* -module over $\mathbb{C}$, so it must be a Hilbert space (with linearity in the second argument of the inner product). The right action is given by

$$\xi \cdot b = \sum_{w \in \Delta^\circ} b(w)\pi(w)^*\xi = b(0)\xi,$$

which under the identification $C^*(\Delta^\circ, \bar{c}) \cong \mathbb{C}$ becomes $\xi \cdot \lambda = \xi\lambda$ for $\xi \in L^2(G)$ and $\lambda \in \mathbb{C}$, i.e. ordinary scalar multiplication. Furthermore, the inner product at the value 0 is given by $\langle \xi, \eta \rangle_\bullet(0) = \langle \pi(0)\eta, \xi \rangle = \langle \eta, \xi \rangle$, i.e. the right inner product is just the conjugate of the ordinary $L^2(G)$ -inner product.

It follows immediately from Proposition II.2.4 that the Heisenberg module norm in this case is just the $L^2(G)$ -norm, and so $\mathcal{E}_\Delta(G) = B_\Delta(G) = L^2(G)$. The statement $B_\Delta(G) = L^2(G)$ when Δ is the whole time-frequency plane is well known. Indeed, in this case the analysis operator C_η is the short-time Fourier transform. This is a bounded operator $L^2(G) \rightarrow L^2(G \times \widehat{G})$ for all $\eta \in L^2(G)$, and is invertible for any $\eta \neq 0$, see [Grö98, Theorem 6.2.1].

Example II.3.14. Suppose G is a discrete group, and that Δ is a cocompact subgroup of $G \times \widehat{G}$ (which must then be a lattice). Then $S_0(G) = \ell^1(G)$ (Proposition II.2.10, ii), and so the Heisenberg module satisfies $\ell^1(G) \subseteq \mathcal{E}_\Delta(G) \subseteq \ell^2(G)$. In particular, if G is finite, then $\mathcal{E}_\Delta(G) = \ell^1(G) = \ell^2(G) = \mathbb{C}G \cong \mathbb{C}^{|G|}$.

The following theorem extends Lemma II.3.8 and is one of our main results.

Theorem II.3.15. *Let G be a second countable, locally compact abelian group, and let Δ be a closed, cocompact subgroup of $G \times \widehat{G}$. Let $\eta, \gamma \in \mathcal{E}_\Delta(G)$. Then the module frame-like operator $\Theta_{\eta, \gamma}: \mathcal{E}_\Delta(G) \rightarrow \mathcal{E}_\Delta(G)$ extends via localization to the Gabor frame-like operator $S_{\eta, \gamma}: L^2(G) \rightarrow L^2(G)$.*

Proof. Let $(\eta_n)_{n=1}^\infty$ and $(\gamma_n)_{n=1}^\infty$ be sequences in $S_0(G)$ that converge towards η and γ respectively in the Heisenberg module norm. Let $\xi \in \mathcal{E}_\Delta(G)$. Then $(\Theta_{\eta_n, \gamma_n} \xi)_n$ converges towards $\Theta_{\eta, \gamma} \xi$ in the Heisenberg module norm. By Lemma II.3.8, we have that $\Theta_{\eta_n, \gamma_n} \xi = S_{\eta_n, \gamma_n} \xi$ for each n , and since convergence in the Heisenberg module norm implies convergence in the $L^2(G)$ -norm, we have that

$$\lim_{n \rightarrow \infty} \|S_{\eta_n, \gamma_n} \xi_n - \Theta_{\eta, \gamma} \xi\|_2 = 0. \quad (\text{II.38})$$

By Proposition II.3.11 and the identity $C_{\eta - \eta_n} = C_\eta - C_{\eta_n}$, the sequences of operators $(C_{\eta_n})_n$ and $(C_{\gamma_n}^*)_n$ converge in the operator norm to C_η and C_γ^* respectively, and so $(S_{\eta_n, \gamma_n})_n$ converges in the operator norm towards $S_{\eta, \gamma}$. It follows that the sequence $(S_{\eta_n, \gamma_n} \xi)_n$ converges to $S_{\eta, \gamma} \xi$ in the L^2 -norm. But then by (II.38), we have that $\Theta_{\eta, \gamma} \xi = S_{\eta, \gamma} \xi$. This shows that the restriction of $S_{\eta, \gamma}$ to $\mathcal{E}_\Delta(G)$ is equal to $\Theta_{\eta, \gamma}$, and so the unique extension of $\Theta_{\eta, \gamma}$ to a bounded linear operator on $L^2(G)$ must be $S_{\eta, \gamma}$. ■

We now arrive at another one of our main results. The following result was previously only known for generators in $S_0(G)$ [JL16; Lue09]. It states that finite module frames for $\mathcal{E}_\Delta(G)$ are exactly the generators of multi-window Gabor frames for $L^2(G)$, where the generators are allowed to come from $\mathcal{E}_\Delta(G)$. This gives a complete description of generators of Heisenberg modules in terms of multi-window Gabor frames.

Theorem II.3.16. *Let G be a second countable, locally compact abelian group, let Δ be a closed, cocompact subgroup of $G \times \widehat{G}$, and let $\eta_1, \dots, \eta_k$ be elements of the Heisenberg module $\mathcal{E}_\Delta(G)$. Then the following are equivalent:*

- (i) *The set $\{\eta_1, \dots, \eta_k\}$ generates $\mathcal{E}_\Delta(G)$ as a left $C^*(\Delta, c)$ -module. That is, for all $\xi \in \mathcal{E}_\Delta(G)$ there exist $a_1, \dots, a_k \in C^*(\Delta, c)$ such that*

$$\xi = \sum_{j=1}^k a_j \cdot \eta_j.$$

- (ii) *The system*

$$\mathcal{G}(\eta_1, \dots, \eta_k; \Delta) = \{\pi(z)\eta_j : z \in \Delta, 1 \leq j \leq k\}$$

is a multi-window Gabor frame for $L^2(G)$.

Proof. By Proposition II.2.6, the set $\{\eta_1, \dots, \eta_k\}$ is a generating set for $\mathcal{E}_\Delta(G)$ if and only if the sequence $(\eta_1, \dots, \eta_k)$ is a module frame for $\mathcal{E}_\Delta(G)$. By Proposition II.2.5, this happens if and only if $\Theta = \Theta_{(\eta_j)_{j=1}^k} = \sum_{j=1}^k \Theta_{\eta_j}$ is invertible as an element of $\mathcal{L}_{C^*(\Delta, c)}(\mathcal{E}_\Delta(G))$. By Theorem II.3.15 and linearity of the localization map $\mathcal{L}_{C^*(\Delta, c)}(\mathcal{E}_\Delta(G)) \hookrightarrow \mathcal{L}(L^2(G))$, this operator extends via localization to the Gabor multi-window frame operator $S = S_{(\eta_j)_{j=1}^k} = \sum_{j=1}^k S_{\eta_j}$ on $L^2(G)$. Since the localization map $\mathcal{L}(\mathcal{E}_\Delta(G)) \hookrightarrow \mathcal{L}(L^2(G))$ is an inclusion of unital C^* -subalgebras, it follows by inverse closedness [Mur90, Theorem 2.1.11] that Θ is invertible in $\mathcal{L}(\mathcal{E}_\Delta(G))$ if and only if S is invertible in $\mathcal{L}(L^2(G))$. But by Proposition II.2.9, the latter happens if and only if $\mathcal{G}(\eta_1, \dots, \eta_k; \Delta)$ is a frame for $L^2(G)$. $\blacksquare$

II.3.5 The fundamental identity of Gabor analysis

So far we have considered a closed subgroup Δ of $G \times \widehat{G}$, and from this we built the Heisenberg module $\mathcal{E}_\Delta(G)$, which is a $C^*(\Delta, c)$ - $C^*(\Delta^\circ, \bar{c})$ -imprimitivity bimodule. We focused specifically on the case when Δ is cocompact, since this implies Δ° is discrete and hence $C^*(\Delta^\circ, \bar{c})$ is unital. By [JL16, p. 5], Δ° is identical to the annihilator $\Delta^\perp$ (also defined in the same article) up to a measure-preserving change of coordinates, and it is also the case that $(\Delta^\perp)^\perp = \Delta$ by [DE14, Proposition 3.6.1]. Hence $(\Delta^\circ)^\circ = \Delta$. Imposing the restriction that Δ° also be cocompact, which implies that both Δ and Δ° are lattices, we could build $\mathcal{E}_{\Delta^\circ}(G)$ and ask how it relates to $\mathcal{E}_\Delta(G)$. The following proposition shows that the relationship is just about as good as we could hope for.

Proposition II.3.17. *Let Δ be a lattice in $G \times \widehat{G}$. Then $\mathcal{E}_\Delta(G) = \mathcal{E}_{\Delta^\circ}(G)$ as subspaces of $L^2(G)$, and $\|\eta\|_{\mathcal{E}_{\Delta^\circ}(G)} = s(\Delta)^{-1/2} \|\eta\|_{\mathcal{E}_\Delta(G)}$ for all $\eta \in \mathcal{E}_\Delta(G)$.*

Proof. Note first that since Δ is a lattice, so is Δ° . In particular, Δ° is a cocompact subgroup, so all the results in this section for Δ apply just as well for Δ° . Hence, by Proposition II.2.12 and Proposition II.3.12, one obtains the Heisenberg module $\mathcal{E}_{\Delta^\circ}(G)$ as a completion of $S_0(G)$ in $B_{\Delta^\circ}(G)$, which is a $C^*(\Delta^\circ, c)$ - $C^*(\Delta, \bar{c})$ -imprimitivity bimodule. Note that in the construction of $\mathcal{E}_{\Delta^\circ}(G)$ we put on Δ° the counting measure, and on Δ the counting measure

scaled with $s(\Delta^\circ)^{-1}$ as per Convention II.3.6. Denote the left inner product on $\mathcal{E}_{\Delta^\circ}(G)$ by $\bullet\langle\cdot, \cdot\rangle'$ and the right inner product by $\langle\cdot, \cdot\rangle'_\bullet$.

Let $\eta \in S_0(G)$. Denote by $\pi_{\Delta^\circ}^*$ the C^* -algebra representation as in the discussion following Proposition II.2.12. Denote by π_{Δ° the representation π_Δ in the same discussion, but with Δ replaced with Δ° . In other words, $\pi_{\Delta^\circ}^*$ is a representation of $C^*(\Delta^\circ, \bar{c})$ on $L^2(G)$, while π_{Δ° is a representation of $C^*(\Delta^\circ, c)$ on $L^2(G)$. Keeping in mind the right measures, we have that

$$\begin{aligned}\pi_{\Delta^\circ}^*(\langle\eta, \eta\rangle_\bullet) &= s(\Delta)^{-1} \sum_{w \in \Delta^\circ} \langle\pi(w)\eta, \eta\rangle \pi(w)^* \\ \pi_{\Delta^\circ}(\bullet\langle\eta, \eta\rangle') &= \sum_{w \in \Delta^\circ} \langle\eta, \pi(w)\eta\rangle \pi(w).\end{aligned}$$

From the above we see that

$$(\pi_{\Delta^\circ}^*(\langle\eta, \eta\rangle_\bullet))^* = s(\Delta)^{-1} \pi_{\Delta^\circ}(\bullet\langle\eta, \eta\rangle'). \quad (\text{II.39})$$

Using the faithfulness of the integrated representations (Proposition II.2.11), we obtain for all $\eta \in S_0(G)$ that

$$\begin{aligned}\|\eta\|_{\mathcal{E}_{\Delta^\circ}(G)}^2 &= \|\bullet\langle\eta, \eta\rangle'\|_{C^*(\Delta^\circ, c)} \\ &= \|\pi_{\Delta^\circ}(\bullet\langle\eta, \eta\rangle')\| && \text{by faithfulness of } \pi_{\Delta^\circ} \\ &= \|\pi_{\Delta^\circ}(\bullet\langle\eta, \eta\rangle')^*\| \\ &= \|s(\Delta)^{-1} \pi_{\Delta^\circ}^*(\langle\eta, \eta\rangle_\bullet)\| && \text{by (II.39)} \\ &= s(\Delta)^{-1} \|\langle\eta, \eta\rangle_\bullet\|_{C^*(\Delta^\circ, \bar{c})} && \text{by faithfulness of } \pi_{\Delta^\circ}^* \\ &= s(\Delta)^{-1} \|\bullet\langle\eta, \eta\rangle\|_{C^*(\Delta, c)} && \text{by Proposition II.2.4} \\ &= s(\Delta)^{-1} \|\eta\|_{\mathcal{E}_\Delta(G)}^2.\end{aligned}$$

By the above, we have that a sequence $(\eta_n)_n$ in $S_0(G)$ is Cauchy in the $\mathcal{E}_\Delta(G)$ -norm if and only if it is Cauchy in the $\mathcal{E}_{\Delta^\circ}(G)$ -norm. Thus, the sequence has a limit in $\mathcal{E}_\Delta(G)$ -norm if and only if it has a limit in $\mathcal{E}_{\Delta^\circ}(G)$ -norm. Since both of these norms dominate the $L^2(G)$ -norm, it follows if $(\eta_n)_n$ is Cauchy in either of the norms, the limit in either of the norms give the same element in $L^2(G)$. It follows that $\mathcal{E}_{\Delta^\circ}(G) = \mathcal{E}_\Delta(G)$, with $\|\eta\|_{\mathcal{E}_{\Delta^\circ}(G)} = s(\Delta)^{1/2} \|\eta\|_{\mathcal{E}_\Delta(G)}$ for all $\eta \in \mathcal{E}_\Delta(G)$. ■

Finally, we show that the fundamental identity of Gabor analysis (or FIGA) [FL06, Theorem 4.5] holds when all involved functions are in $\mathcal{E}_\Delta(G)$ when Δ is a lattice. In the following we let $S_{\eta, \gamma}^\Delta$ be the operator of (II.8), and let $S_{\eta, \gamma}^{\Delta^\circ}$ denote the operator of (II.8) but with Δ° instead of Δ . It is already known that FIGA holds for functions in $S_0(G)$ even when Δ is just a closed subgroup of the time-frequency plane $G \times \widehat{G}$ [JL16, Corollary 6.3]. However, in the proof of Proposition II.3.18 below we shall need to apply Theorem II.3.15 to frame operators with respect to Δ and with respect to Δ° . This then requires that both

II. Heisenberg modules as function spaces

Δ and Δ° are cocompact, hence they are both lattices. With these restrictions, FIGA is the statement

$$\sum_{z \in \Delta} \langle \eta, \pi(z)\gamma \rangle \langle \pi(z)\xi, \psi \rangle = \frac{1}{s(\Delta)} \sum_{z^\circ \in \Delta^\circ} \langle \xi, \pi(z^\circ)\gamma \rangle \langle \pi(z^\circ)\eta, \psi \rangle \quad (\text{II.40})$$

for $\eta, \gamma, \xi, \psi \in S_0(G)$. In short form it is just the statement

$$S_{\gamma, \xi}^\Delta \eta = s(\Delta)^{-1} S_{\gamma, \eta}^{\Delta^\circ} \xi \quad (\text{II.41})$$

for $\eta, \gamma, \xi \in S_0(G)$. With the restriction that Δ is a lattice in $G \times \widehat{G}$, the following proposition extends the range for the FIGA (for the particular lattice Δ) to functions in $\mathcal{E}_\Delta(G)$.

Proposition II.3.18. *Let G be a second countable, locally compact abelian group, and let Δ be a lattice in $G \times \widehat{G}$. Then (II.41) holds for $\eta, \gamma, \xi \in \mathcal{E}_\Delta(G)$.*

Proof. Let $(\eta_n)_n, (\gamma_n)_n$ and $(\xi_n)_n$ be sequences in $S_0(G)$ that converge to η, γ and ξ , respectively, in the $\mathcal{E}_\Delta(G)$ -norm. By Proposition II.3.17, the same is true in $\mathcal{E}_{\Delta^\circ}(G)$ -norm. Then, since the fundamental identity of Gabor analysis applies for functions in $S_0(G)$ by [JL16, Corollary 6.3], we have

$$S_{\gamma_n, \xi_n}^\Delta \eta_n = s(\Delta)^{-1} S_{\gamma_n, \eta_n}^{\Delta^\circ} \xi_n$$

for all $n \in \mathbb{N}$. By Theorem II.3.15 we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|S_{\gamma, \xi}^\Delta \eta - S_{\gamma_n, \xi_n}^\Delta \eta_n\|_{\mathcal{E}_\Delta(G)} &= 0 \\ \lim_{n \rightarrow \infty} \|S_{\gamma, \eta}^{\Delta^\circ} \xi - S_{\gamma_n, \eta_n}^{\Delta^\circ} \xi_n\|_{\mathcal{E}_{\Delta^\circ}(G)} &= 0. \end{aligned}$$

Since convergence in the Heisenberg module norm implies convergence in the $L^2(G)$ -norm, we conclude that the following equality holds in $L^2(G)$, were the limits are taken in $L^2(G)$:

$$S_{\gamma, \xi}^\Delta \eta = \lim_{n \rightarrow \infty} S_{\gamma_n, \xi_n}^\Delta \eta_n = \lim_{n \rightarrow \infty} s(\Delta)^{-1} S_{\gamma_n, \eta_n}^{\Delta^\circ} \xi_n = s(\Delta)^{-1} S_{\gamma, \eta}^{\Delta^\circ} \xi.$$

■

Remark II.3.19. As already mentioned, the FIGA holds for functions in $S_0(G)$ even when Δ is only a closed subgroup for $G \times \widehat{G}$. The techniques in this paper are based on localization of A - B -imprimitivity bimodules, which requires that we have a finite trace on at least one of the algebras A and B . Therefore the assumption that Δ is a lattice in $G \times \widehat{G}$ is necessary for our approach to the FIGA. There might be another technique that allows for an extension of the FIGA to $\mathcal{E}_\Delta(G)$ for Δ only a closed subgroup of $G \times \widehat{G}$ that the authors are not aware of. We remark again that for the existence of Gabor frames over a closed subgroup Δ of $G \times \widehat{G}$, it is necessary that Δ is cocompact in $G \times \widehat{G}$, which is the setting for most of the results in the paper.

Acknowledgements. The authors would like to thank Nadia Larsen, Franz Luef and Luca Gazdag for giving feedback on a draft of the paper. The second author would like to thank Erik Bédos for helpful discussions. The authors are also indebted to the referees for their invaluable feedback on the first draft of the paper, and to one of the referees for suggesting a simpler proof of Proposition II.2.6, which we have included.

References

- [AAG93] Ali, S. T., Antoine, J.-P., and Gazeau, J.-P. “Continuous frames in Hilbert space”. In: *Ann. Physics* vol. 222, no. 1 (1993), pp. 1–37.
- [AB17] Arambašić, L. and Bakić, D. “Frames and outer frames for Hilbert C^* -modules”. In: *Linear Multilinear Algebra* vol. 65, no. 2 (2017), pp. 381–431.
- [AL18] Austad, A. and Luef, F. “Modulation Spaces as a Smooth Structure in Noncommutative Geometry”. preprint, arXiv.org:1809.11063. 2018.
- [BO18] Bédos, E. and Omland, T. “On reduced twisted group C^* -algebras that are simple and/or have a unique trace”. In: *J. Noncommut. Geom.* Vol. 12, no. 3 (2018), pp. 947–996.
- [Con94] Connes, A. *Noncommutative geometry*. Academic Press, Inc., San Diego, CA, 1994, pp. xiv+661.
- [CZ83] Combes, F. and Zettl, H. “Order structures, traces and weights on Morita equivalent C^* -algebras”. In: *Math. Ann.* Vol. 265, no. 1 (1983), pp. 67–81.
- [DE14] Deitmar, A. and Eichterhoff, S. *Principles of harmonic analysis*. Second. Universitext. Springer, Cham, 2014, pp. xiv+332.
- [DS52] Duffin, R. J. and Schaeffer, A. C. “A class of nonharmonic Fourier series”. In: *Trans. Amer. Math. Soc.* Vol. 72 (1952), pp. 341–366.
- [EJL19] Enstad, U. B. R., Jakobsen, M. S., and Luef, F. “Time-frequency analysis on the adèles over the rationals”. In: *C. R. Math. Acad. Sci. Paris* vol. 357, no. 2 (2019), pp. 188–199.
- [Fei81] Feichtinger, H. G. “On a new Segal algebra”. In: *Monatsh. Math.* Vol. 92, no. 4 (1981), pp. 269–289.
- [FL02] Frank, M. and Larson, D. R. “Frames in Hilbert C^* -modules and C^* -algebras”. In: *J. Operator Theory* vol. 48, no. 2 (2002), pp. 273–314.
- [FL06] Feichtinger, H. G. and Luef, F. “Wiener amalgam spaces for the fundamental identity of Gabor analysis”. In: *Collect. Math.* Vol. 57 (2006), pp. 233–253.
- [Fol95] Folland, G. B. *A course in abstract harmonic analysis*. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1995, pp. x+276.

- [Gab46] Gabor, D. “Theory of communication”. In: *Journal of the Institution of Electrical Engineers* (1946), pp. 429–457.
- [Grö98] Gröchenig, K. “Aspects of Gabor analysis on locally compact abelian groups”. In: *Gabor analysis and algorithms*. Appl. Numer. Harmon. Anal. Birkhäuser Boston, Boston, MA, 1998, pp. 211–231.
- [Jak18] Jakobsen, M. S. “On a (no longer) new Segal algebra: a review of the Feichtinger algebra”. In: *J. Fourier Anal. Appl.* Vol. 24, no. 6 (2018), pp. 1579–1660.
- [JL16] Jakobsen, M. S. and Lemvig, J. “Density and duality theorems for regular Gabor frames”. In: *J. Funct. Anal.* Vol. 270, no. 1 (2016), pp. 229–263.
- [JL18] Jakobsen, M. S. and Luef, F. “Duality of Gabor Frames and Heisenberg Modules”. preprint, arXiv.org:1806.05616. 2018.
- [Kai94] Kaiser, G. *A friendly guide to wavelets*. Birkhäuser Boston, Inc., Boston, MA, 1994, pp. xviii+300.
- [Kap53] Kaplansky, I. “Modules over operator algebras”. In: *Amer. J. Math.* Vol. 75 (1953), pp. 839–858.
- [KL12] KaaD, J. and Lesch, M. “A local global principle for regular operators in Hilbert C^* -modules”. In: *J. Funct. Anal.* Vol. 262, no. 10 (2012), pp. 4540–4569.
- [Lan95] Lance, E. C. *Hilbert C^* -modules*. Vol. 210. London Mathematical Society Lecture Note Series. A toolkit for operator algebraists. Cambridge University Press, Cambridge, 1995, pp. x+130.
- [LP13] Latrémolière, F. and Packer, J. A. “Noncommutative solenoids and their projective modules”. In: *Commutative and noncommutative harmonic analysis and applications*. Vol. 603. Contemp. Math. Amer. Math. Soc., Providence, RI, 2013, pp. 35–53.
- [LP15] Latrémolière, F. and Packer, J. A. “Explicit construction of equivalence bimodules between noncommutative solenoids”. In: *Trends in harmonic analysis and its applications*. Vol. 650. Contemp. Math. Amer. Math. Soc., Providence, RI, 2015, pp. 111–140.
- [Lue09] Luef, F. “Projective modules over noncommutative tori are multi-window Gabor frames for modulation spaces”. In: *J. Funct. Anal.* Vol. 257, no. 6 (2009), pp. 1921–1946.
- [Lue11] Luef, F. “Projections in noncommutative tori and Gabor frames”. In: *Proc. Amer. Math. Soc.* Vol. 139, no. 2 (2011), pp. 571–582.
- [Mur90] Murphy, G. J. *C^* -algebras and operator theory*. Academic Press, Inc., Boston, MA, 1990, pp. x+286.
- [Ped79] Pedersen, G. K. *C^* -algebras and their automorphism groups*. Vol. 14. London Mathematical Society Monographs. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London-New York, 1979, pp. ix+416.

- [PR89] Packer, J. A. and Raeburn, I. “Twisted crossed products of C^* -algebras”. In: *Math. Proc. Cambridge Philos. Soc.* Vol. 106, no. 2 (1989), pp. 293–311.
- [Rie74a] Rieffel, M. A. “Induced representations of C^* -algebras”. In: *Advances in Math.* Vol. 13 (1974), pp. 176–257.
- [Rie74b] Rieffel, M. A. “Morita equivalence for C^* -algebras and W^* -algebras”. In: *J. Pure Appl. Algebra* vol. 5 (1974), pp. 51–96.
- [Rie88] Rieffel, M. A. “Projective modules over higher-dimensional noncommutative tori”. In: *Canad. J. Math.* Vol. 40, no. 2 (1988), pp. 257–338.
- [Rud87] Rudin, W. *Real and complex analysis*. Third. McGraw-Hill Book Co., New York, 1987, pp. xiv+416.
- [RW98] Raeburn, I. and Williams, D. P. *Morita equivalence and continuous-trace C^* -algebras*. Vol. 60. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1998, pp. xiv+327.

The Balian–Low theorem for locally compact abelian groups and vector bundles

Ulrik Enstad

To appear in *Journal de Mathématiques Pures et Appliquées*.

DOI: 10.1016/j.matpur.2019.12.005.

Abstract

Let Λ be a lattice in a second countable, locally compact abelian group G with annihilator $\Lambda^\perp \subseteq \widehat{G}$. We investigate the validity of the following statement: For every η in the Feichtinger algebra $S_0(G)$, the Gabor system $\{M_\tau T_\lambda \eta\}_{\lambda \in \Lambda, \tau \in \Lambda^\perp}$ is not a frame for $L^2(G)$. When Λ is a lattice in $G = \mathbb{R}$, this statement is a variant of the Balian–Low theorem. Extending a result of R. Balan, we show that whether the statement generalizes to (G, Λ) is equivalent to the nontriviality of a certain vector bundle over the compact space $(G/\Lambda) \times (\widehat{G}/\Lambda^\perp)$. We prove this equivalence using Heisenberg modules. More specifically, we show that the Zak transform can be viewed as an isomorphism of certain Hilbert C^* -modules. As an application, we prove a Balian–Low theorem in the new context of the group $\mathbb{R} \times \mathbb{Q}_p$, where $\mathbb{Q}_p$ denotes the p -adic numbers.

III.1 Introduction

In his work on projective modules over noncommutative tori [Rie88], M. Rieffel introduced a class of Hilbert C^* -modules known as Heisenberg modules. These modules establish the Morita equivalence of twisted group C^* -algebras associated to a lattice Δ in the time-frequency plane of a locally compact abelian (LCA) group G . Heisenberg modules have been applied numerous times in operator algebras and noncommutative geometry, see for example [Boc99; CL18; LM16; LP13; LP15].

It was shown by F. Luef in [Lue09; Lue11] that the Morita equivalence of Heisenberg modules over noncommutative tori is closely related to the duality theory of regular Gabor frames. These connections were recently generalized to the setting of LCA groups [JL18]. Gabor frames are the objects of study in Gabor analysis, which can be considered a subfield of time-frequency analysis.

The central problem of Gabor analysis is the recovery of signals from a discrete set of time-frequency translates of fixed functions in $L^2(\mathbb{R}^n)$. While Gabor analysis is usually carried out in $\mathbb{R}^n$, much of the framework can be generalized to the setting of a LCA group G as follows: If (x, ω) is an element of the time-frequency plane $G \times \widehat{G}$, the time-frequency shift operator $\pi(x, \omega)$ acts on functions $\xi \in L^2(G)$ via

$$\pi(x, \omega)\xi(t) = \omega(t)\xi(x^{-1}t)$$

for $t \in G$. By picking a lattice Δ in the time-frequency plane $G \times \widehat{G}$ and a finite set of functions $\eta_1, \dots, \eta_k \in L^2(G)$, one forms the associated *multiwindow Gabor system* as follows:

$$\mathcal{G}(\eta_1, \dots, \eta_k; \Delta) = \{\pi(z)\eta_j : z \in \Delta, 1 \leq j \leq k\}.$$

To allow for stable reconstruction of functions in $L^2(G)$ from a multiwindow Gabor system, one requires the frame property due to Duffin and Schaeffer [DS52] to be satisfied. That is, if there exist constants $K, L > 0$ such that

$$K\|\xi\|_2^2 \leq \sum_{j=1}^k \sum_{z \in \Delta} |\langle \xi, \pi(z)\eta_j \rangle|^2 \leq L\|\xi\|_2^2$$

for all $\xi \in L^2(G)$, one calls the multiwindow Gabor system $\mathcal{G}(\eta_1, \dots, \eta_k; \Delta)$ a *multiwindow Gabor frame*. In particular, a singlewindow Gabor system, or just Gabor system for short, is called a *Gabor frame* if it forms a frame for $L^2(G)$.

One of the main observations of [JL18; Lue09] is that if the windows $\eta_1, \dots, \eta_k$ of a multiwindow Gabor frame over the lattice $\Delta \subseteq G \times \widehat{G}$ are well-localized, they can be interpreted as a set of generators for the Heisenberg module $\mathcal{E}_\Delta(G)$ constructed from Δ . By well-localized, we mean that the generators are all elements of the Feichtinger algebra $S_0(G)$, a Banach space of test functions that is fundamental to time-frequency analysis. Since Heisenberg modules are finitely generated, an immediate consequence is the existence of a multiwindow Gabor frame $\mathcal{G}(\eta_1, \dots, \eta_k; \Delta)$ with $\eta_j \in S_0(G)$, $1 \leq j \leq k$, for any given lattice Δ in $G \times \widehat{G}$.

A longstanding problem in Gabor analysis on $G = \mathbb{R}^n$ is whether one can find a (singlewindow) Gabor frame $\mathcal{G}(\eta; \Delta)$ with a well-localized window $\eta \in S_0(\mathbb{R}^n)$ over a given lattice Δ in the time-frequency plane $\mathbb{R}^n \times \widehat{\mathbb{R}^n} \cong \mathbb{R}^{2n}$. One can ask the same question for a lattice Δ in the time-frequency plane $G \times \widehat{G}$ of a locally compact abelian group G : Does there exist an $\eta \in S_0(G)$ for which $\mathcal{G}(\eta, \Delta)$ is a Gabor frame? Or, in terms of Heisenberg modules, is the Heisenberg module $\mathcal{E}_\Delta(G)$ singly generated? A basic restriction on the lattice is provided by one of the density theorems [Hei07]: It is necessary that $\text{vol}(\Delta) \leq 1$ for Gabor frames $\mathcal{G}(\eta, \Delta)$ with $\eta \in S_0(G)$ to exist [JL16, Theorem 5.6]. But in the group $G = \mathbb{R}^n$, more is true:

Theorem III.1.1. *Let Δ be a lattice in the time-frequency plane $\mathbb{R}^n \times \widehat{\mathbb{R}^n}$ of $\mathbb{R}^n$. If there exists a function η in the Feichtinger algebra $S_0(\mathbb{R}^n)$ for which $\mathcal{G}(\eta, \Delta)$ is a Gabor frame for $L^2(\mathbb{R}^n)$, then $\text{vol}(\Delta) < 1$.*

The above result is an example of a Balian-Low theorem (BLT), as it is a non-existence result for well-localized Gabor frames at the so-called *critical density* $\text{vol}(\Delta) = 1$. The original Balian-Low theorem is due to R. Balian [Bal81] and F. Low [Low85] and concerns lattices of the form $\Delta = \alpha\mathbb{Z} \times \beta\mathbb{Z}$ for $\alpha, \beta > 0$ in $\mathbb{R} \times \widehat{\mathbb{R}}$. It also uses a slightly more general notion of time-frequency localization that does not involve the Feichtinger algebra, but which is particular to $G = \mathbb{R}$. In [Bat88], a proof of the original formulation of the Balian-Low theorem is deduced from the uncertainty principle, and a relation to noncommutative geometry is demonstrated in [Lue18]. The amalgam Balian-Low theorem is another early version of the BLT that employs Wiener amalgam spaces [BHW95; Hei90]. Versions of the Balian-Low theorem for more general lattices in $\mathbb{R}^n$ have since been proved, even for discrete sets Δ without any lattice structure [AFK14]. Theorem III.1.1 can also be deduced from the perturbation results of H. Feichtinger and N. Kaiblinger [FK04]. The converse of Theorem III.1.1 remains open for general lattices, but was proved for the class of non-rational lattices in [JL18]. The proof uses Heisenberg modules and K -theory of noncommutative tori.

It is no coincidence that the setting of Theorem III.1.1 is the group $G = \mathbb{R}^n$. Indeed, Theorem III.1.1 is easily shown to fail if one replaces $\mathbb{R}^n$ with an arbitrary LCA group G [Grö98]. For instance, it fails for discrete or compact groups (see Proposition III.5.7). One might then ask if there is a way to characterize the groups G for which Theorem III.1.1 holds. This will be the setup in the present paper, but we restrict ourselves to the case where the lattice Δ takes the form $\Lambda \times \Lambda^\perp$, where Λ is a lattice in G and $\Lambda^\perp$ is the annihilator of Λ in $\widehat{G}$ (III.10). These lattices always have volume 1. We will consider the following statement in the setting (G, Λ) , where G is a second countable LCA group and Λ is a lattice in G .

Statement III.1.2. *For all $\eta \in S_0(G)$, the Gabor system*

$$\mathcal{G}(\eta, \Lambda \times \Lambda^\perp) = \{\pi(\lambda, \tau)\eta : \lambda \in \Lambda, \tau \in \Lambda^\perp\}$$

is not a frame for $L^2(G)$.

Note that it is a consequence of Theorem III.1.1 that Statement III.1.2 holds true for any lattice Λ in $G = \mathbb{R}^n$.

The advantage of the formulation of Statement III.1.2 is that the *Zak transform* can be employed. Originally introduced by I. Gelfand [Gel50], the Zak transform was later generalized by A. Weil to the case of locally compact abelian groups [Wei64]. It takes its name from the physicist J. Zak who discovered it independently [Zak67]. Given a lattice Λ in the LCA group G , the Zak transform of a complex-valued function ξ on G is the function $Z_{G, \Lambda}\xi : G \times \widehat{G} \rightarrow \mathbb{C}$ given by

$$Z_{G, \Lambda}\xi(x, \omega) = \sum_{\lambda \in \Lambda} \xi(x\lambda)\omega(\lambda)$$

for $(x, \omega) \in G \times \widehat{G}$. The above defines a continuous function if e.g. $\xi \in S_0(G)$.

The first proofs of the amalgam version of the BLT [BHW95; Hei90] employed the Zak transform, and used the fact that it diagonalizes the frame operator associated to the Gabor system $\mathcal{G}(\eta, \Lambda \times \Lambda^\perp)$. In [KK98], E. Kaniuth and G. Kutyniok used the Zak transform to show that the Balian–Low statement in Statement III.1.2 holds for all lattices in compactly generated, second countable, locally compact abelian groups with noncompact identity component. It is an open problem whether the hypothesis that G is compactly generated can be dropped from their theorem.

One of the main points we want to make in this paper is that the Zak transform has a natural interpretation as an isomorphism of Hilbert C^* -modules. When $\Delta = \Lambda \times \Lambda^\perp$, the associated Heisenberg module $\mathcal{E}_\Delta(G)$ becomes a Hilbert C^* -module over the (un-twisted) group C^* -algebra of Δ . Using the Fourier transform, this algebra can be identified as the continuous functions on the Pontryagin dual $X := \widehat{\Delta} \cong (G/\Lambda) \times (\widehat{G}/\Lambda^\perp)$. By the Serre–Swan theorem, the projective module $\mathcal{E}_\Delta(G)$ must be isomorphic to a vector bundle over X . The role of the Zak transform in this respect is to identify the vector bundle in question. More precisely, we prove that there exists a complex vector bundle $E_{G,\Lambda}$ for which the following holds:

Theorem B (cf. Theorem III.5.1 / Proposition III.4.5). *Let Λ be a lattice in the second countable LCA group G , and let Δ be the lattice $\Lambda \times \Lambda^\perp$ in $G \times \widehat{G}$. Then the Zak transform implements an isomorphism of Hilbert C^* -modules*

$$Z_{G,\Lambda} : \mathcal{E}_\Delta(G) \rightarrow \Gamma(E_{G,\Lambda}).$$

Here, $\mathcal{E}_\Delta(G)$ is the Heisenberg module associated to Δ , and $\Gamma(E_{G,\Lambda})$ is the module of continuous sections of the complex line bundle $E_{G,\Lambda}$ constructed in Section III.4.2.

Now Statement III.1.2 is equivalent to the nonexistence of a single generator for the Heisenberg module $\mathcal{E}_\Delta(G)$. We show in Section III.2.3 that the $C(X)$ -module $\Gamma(E)$ of continuous sections of a vector bundle E over a compact Hausdorff space is singly generated if and only if E is a trivial bundle. Consequently, we can formulate Statement III.1.2 in terms of the vector bundle $E_{G,\Lambda}$ from Theorem B as follows:

Theorem C (cf. Theorem III.5.4). *Let Λ be a lattice in a second countable, locally compact abelian group G . Then the following are equivalent:*

- (i) $E_{G,\Lambda}$ is nontrivial.
- (ii) The Balian–Low statement (Statement III.1.2) holds in this setting. That is, whenever $\eta \in S_0(G)$, then the Gabor system

$$\mathcal{G}(\eta, \Lambda \times \Lambda^\perp) = \{\pi(\lambda, \tau)\eta : \lambda \in \Lambda, \tau \in \Lambda^\perp\}$$

is not a frame for $L^2(G)$.

The above theorem builds upon an idea of R. Balan connecting Gabor superframes to vector bundles over the 2-torus $\mathbb{T}^2$ [Bal01]. A special case of his

result is that the amalgam version of the Balian–Low theorem is a consequence of the nontriviality of a certain line bundle over $\mathbb{T}^2$. If $G = \mathbb{R}$ and $\Lambda = \alpha\mathbb{Z}$ in Theorem C, then the base space $X_{G,\Lambda}$ is homeomorphic to $\mathbb{T}^2$, and we will indeed show that $E_{G,\Lambda}$ in this case is closely related to Balan’s bundle (cf. Example III.4.6). Thus, Theorem C can be viewed as an extension of this special case of Balan’s result to general second countable LCA groups.

We end this paper by applying Theorem C to prove that the Balian–Low statement (Statement III.1.2) holds in a new setting: We set G to be the truncated adèle group $\mathbb{R} \times \mathbb{Q}_p$ where $\mathbb{Q}_p$ denotes the p -adic numbers. It is a well-known fact from number theory that the group of p -adic rationals $\Lambda = \mathbb{Z}[1/p] = \{a/p^k : a, k \in \mathbb{Z}\}$ embeds as a lattice in G . We then have the following:

Theorem D (cf. Theorem III.6.5). *Let G be the group $\mathbb{R} \times \mathbb{Q}_p$, and let Λ be the lattice $\mathbb{Z}[1/p]$ embedded into G as in Section III.6. Then the Balian–Low statement (Statement III.1.2) holds for (G, Λ) : That is, whenever $\eta \in S_0(\mathbb{R} \times \mathbb{Q}_p)$, the Gabor system*

$$\mathcal{G}(\eta, \Lambda \times \Lambda^\perp) = \{(s, x) \mapsto e^{2\pi i r s} e^{-2\pi i \{rx\}_p} \eta(s - q, x - q) : q, r \in \mathbb{Z}[1/p]\}$$

is not a frame for $L^2(\mathbb{R} \times \mathbb{Q}_p)$.

The above theorem is the first Balian–Low theorem in the context of a LCA group which is not compactly generated. Hence, it is not covered by the result of Kaniuth and Kutyniok [KK98]. Number-theoretic groups such as $\mathbb{R} \times \mathbb{Q}_p$ and the full adèles over the rationals have not been explored much in Gabor analysis so far. In [EJL19], examples of Gabor frames in these groups are constructed, and a mild Balian–Low type theorem, namely [EJL19, Proposition 4.4], is also proved. However, the result only holds for functions in $S_0(\mathbb{R} \times \mathbb{Q}_p)$ of a very specific form, and Theorem D is a generalization to all generators in $S_0(\mathbb{R} \times \mathbb{Q}_p)$.

The text is structured as follows: In Section III.2, we define Hilbert C^* -modules and their frames, and we discuss modules of sections of vector bundles. In Section III.3, we introduce Gabor analysis on locally compact abelian groups and Heisenberg modules. In Section III.4, we introduce the Zak transform, quasiperiodic functions and the vector bundle $E_{G,\Lambda}$ from Theorem B. In Section III.5, we show that the Zak transform gives an isomorphism of Hilbert C^* -modules, and prove theorems Theorem B and Theorem C. Then in Section III.6, we prove a Balian–Low theorem for the group $\mathbb{R} \times \mathbb{Q}_p$, namely Theorem D. In the appendix, we have collected some basic results that are needed but do not constitute a part of the spirit of the main text.

III.2 Hilbert C^* -modules and their frames

III.2.1 Hilbert C^* -modules

In this section we define Hilbert C^* -modules, see e.g. [RW98]. Let A be a unital C^* -algebra. A Hilbert C^* -module over A is, roughly speaking, a “Hilbert space”

with an inner product taking values in the C^* -algebra A rather than the complex numbers. More precisely, a *left Hilbert A -module* $\mathcal{E}$ is a left A -module that is equipped with an A -valued inner product $\bullet\langle \cdot, \cdot \rangle : \mathcal{E} \times \mathcal{E} \rightarrow A$ that satisfies the following axioms:

- (i) $\bullet\langle a\xi + b\eta, \gamma \rangle = a\bullet\langle \xi, \gamma \rangle + b\bullet\langle \eta, \gamma \rangle$ for $a, b \in A$ and $\xi, \eta, \gamma \in \mathcal{E}$.
- (ii) $\bullet\langle \xi, \eta \rangle^* = \bullet\langle \eta, \xi \rangle$ for $\xi, \eta \in \mathcal{E}$.
- (iii) $\bullet\langle \xi, \xi \rangle \geq 0$, with $\bullet\langle \xi, \xi \rangle = 0$ if and only if $\xi = 0$.
- (iv) $\mathcal{E}$ is complete with respect to the norm $\|\xi\|_{\mathcal{E}} = \|\bullet\langle \xi, \xi \rangle\|^{1/2}$.

Note that we use the dot in the notation for the A -valued inner product to distinguish it from $\mathbb{C}$ -valued inner products.

From the definition above, $\mathcal{E}$ has a complex vector space structure given by $\lambda\xi := (\lambda 1_A)\xi$ where $\lambda \in \mathbb{C}$, $\xi \in \mathcal{E}$ and 1_A is the multiplicative identity of A .

If $A = \mathbb{C}$, one recovers the definition of a (complex) Hilbert space.

Let $\mathcal{E}$ and $\mathcal{F}$ be left Hilbert A -modules, and let $T : \mathcal{E} \rightarrow \mathcal{F}$ be a bounded, A -linear map. Then T is called *adjointable* if there exists an A -linear map $S : \mathcal{F} \rightarrow \mathcal{E}$ such that

$$\bullet\langle T\xi, \eta \rangle = \bullet\langle \xi, S\eta \rangle$$

for all $\xi \in \mathcal{E}$ and $\eta \in \mathcal{F}$. The map S is then uniquely determined and is referred to the *adjoint* of S . It is denoted by $S = T^*$. If T is adjointable, then T is invertible if and only if its adjoint is invertible, and in that case we have the relationship $(T^*)^{-1} = (T^{-1})^*$. The set of adjointable operators $\mathcal{E} \rightarrow \mathcal{E}$ is a C^* -algebra and is denoted by $\mathcal{L}_A(\mathcal{E})$ or just $\mathcal{L}(\mathcal{E})$.

A bounded, A -linear map $T : \mathcal{E} \rightarrow \mathcal{F}$ is called *inner product preserving* if

$$\bullet\langle T\xi, T\eta \rangle = \bullet\langle \xi, \eta \rangle$$

for all $\xi, \eta \in \mathcal{E}$. An inner product preserving map is automatically injective, as it becomes an isometry in the sense of metric spaces. An inner product preserving map is not necessarily adjointable, see [RW98, Example 2.19]. However, if it is surjective, hence invertible, then it is also adjointable, and the adjoint is given by the inverse.

We say that two left Hilbert A -modules $\mathcal{E}$ and $\mathcal{F}$ are *isomorphic* if there exists a surjective inner product preserving (hence invertible and adjointable) A -linear map $T : \mathcal{E} \rightarrow \mathcal{F}$.

Let J be a countable index set. Denote by $\ell^2(J, A)$ the set of all sequences $(a_j)_{j \in J}$ for which the sum $\sum_{j \in J} a_j a_j^*$ converges unconditionally in A (this is the same as the direct sum $\bigoplus_{j \in J} A$, see [Lan95, p. 6]). This set forms a left Hilbert A -module with respect to pointwise left multiplication and the inner product

$$\bullet\langle (a_j)_j, (b_j)_j \rangle = \sum_{j \in J} a_j b_j^*$$

for $(a_j)_j, (b_j)_j \in \ell^2(J, A)$. If $J = \{1, \dots, k\}$ for some $k \in \mathbb{N}$, one obtains the left Hilbert A -module A^k , the *free* A -module of rank k . We represent the elements of A^k as row vectors. If M is a $k \times k$ matrix with entries in A , then *right multiplication* by M defines an A -linear map $A^n \rightarrow A^n$.

A left Hilbert A -module $\mathcal{E}$ is called *countably generated* if there exists a countable set $S \subseteq \mathcal{E}$ such that $\{a \cdot \xi : a \in A, \xi \in S\}$ is dense in $\mathcal{E}$. If $A = \mathbb{C}$ so that $\mathcal{E}$ is a Hilbert space, then being countably generated as an A -module is the same as being separable as a Hilbert space. The module $\mathcal{E}$ is *finitely generated* if there exists a finite generating set S as above. This is equivalent to the existence of a finite set $\{\eta_1, \dots, \eta_k\}$ in $\mathcal{E}$ with the following property: For every $\xi \in \mathcal{E}$, there exist elements $a_1, \dots, a_k \in A$ such that

$$\xi = a_1 \eta_1 + \dots + a_k \eta_k.$$

The set $\{\eta_1, \dots, \eta_k\}$ is called a generating set for $\mathcal{E}$. Moreover, $\mathcal{E}$ is *finitely generated projective* if it is isomorphic to a module of the form $A^k P$, where P is a $k \times k$ projection matrix with coefficients in A .

III.2.2 Module frames

A *module frame* [FL02] in a left Hilbert A -module $\mathcal{E}$ is a sequence $(\eta_j)_{j \in J}$ in $\mathcal{E}$ for which there exists $K, L > 0$ such that the following double inequality holds for all $\xi \in \mathcal{E}$:

$$K \cdot \langle \xi, \xi \rangle \leq \sum_{j \in J} \bullet \langle \xi, \eta_j \rangle \bullet \langle \xi, \eta_j \rangle^* \leq L \cdot \langle \xi, \xi \rangle. \quad (\text{III.1})$$

Here, the inequalities are to be interpreted with respect to the order structure on the self-adjoint elements of the C^* -algebra A .

Note that if A is the C^* -algebra of complex numbers, then $\mathcal{E}$ is a Hilbert space and the above double inequality reduces to

$$K \|\xi\|^2 \leq \sum_{j \in J} |\langle \xi, \eta_j \rangle|^2 \leq L \|\xi\|^2 \quad (\text{III.2})$$

for all $\xi \in \mathcal{H}$.

We call the numbers K and L *lower* and *upper frame bounds*, respectively. A frame is called *tight* if one can choose frame bounds $K = L$, and *normalized tight* if one can choose $K = L = 1$.

A sequence $(\eta_j)_{j \in J}$ that satisfies the upper frame bound condition but not necessarily the lower, is called a *Bessel sequence*. An upper frame bound for a Bessel sequence is called a *Bessel bound* for the sequence. If $(\eta_j)_j$ is a Bessel sequence in $\mathcal{E}$, one defines the operators $C: \mathcal{E} \rightarrow \ell^2(J, A)$ and $D: \ell^2(J, A) \rightarrow \mathcal{E}$ by

$$C\xi = (\bullet \langle \xi, \eta_j \rangle)_{j \in J} \quad (\text{III.3})$$

$$D(a_j)_j = \sum_{j \in J} a_j \eta_j \quad (\text{III.4})$$

III. The BLT for LCA groups and vector bundles

for $\xi \in \mathcal{E}$ and $(a_j)_j \in \ell^2(J, A)$. These are adjointable A -linear operators, and $C^* = D$. They go under different names in the literature. In [FL02], C is called the *frame transform*, while in Gabor analysis in general, they are called the *analysis* and *synthesis* operator, respectively.

One also defines $S = DC$ and calls it the *frame operator* corresponding to $(\eta_j)_j$. It is given by

$$S\xi = \sum_{j \in J} \bullet \langle \xi, \eta_j \rangle \eta_j \quad (\text{III.5})$$

for $\xi \in \mathcal{E}$.

The following gives an important characterization of the frame property of a Bessel sequence. It can be seen by noting that $\bullet \langle S\xi, \xi \rangle = \sum_{j \in J} \bullet \langle \xi, \eta_j \rangle \bullet \langle \xi, \eta_j \rangle^*$ and using the fact that $T_1 \leq T_2$ if and only if $\bullet \langle T_1 \xi, \xi \rangle \leq \bullet \langle T_2 \xi, \xi \rangle$ for all $\xi \in \mathcal{E}$, where $T_1, T_2 \in \mathcal{L}(\mathcal{E})$ are self-adjoint [RW98, Lemma 2.28].

Proposition III.2.1. *Let $\mathcal{E}$ be a left Hilbert A -module over a C^* -algebra A . Let $(\eta_j)_{j \in J}$ be a Bessel sequence in $\mathcal{E}$. Then the following are equivalent:*

- (i) $(\eta_j)_j$ is a frame for $\mathcal{E}$.
- (ii) The frame operator S corresponding to $(\eta_j)_j$ as in (III.5) is invertible in $\mathcal{L}(\mathcal{E})$.

Using the Cauchy–Schwarz inequality for Hilbert C^* -modules, one can show that for a finite sequence, there is always an upper frame bound. Thus, to show that a finite sequence is a module frame, one only needs to show that there exists a lower frame bound. In fact, we have the following characterizations of finite module frames.

Proposition III.2.2. *Let A be a unital C^* -algebra, and let $\mathcal{E}$ be a left Hilbert A -module. Let $\eta_1, \dots, \eta_k \in \mathcal{E}$. Then the following are equivalent:*

- (i) The set $\{\eta_1, \dots, \eta_k\}$ is a generating set for $\mathcal{E}$.
- (ii) The sequence $(\eta_1, \dots, \eta_k)$ is a module frame for $\mathcal{E}$.
- (iii) The frame operator S corresponding to $(\eta_1, \dots, \eta_k)$ as in (III.5) is invertible in $\mathcal{L}(\mathcal{E})$.

Moreover, if $(\eta_1, \dots, \eta_k)$ is a module frame, then the sequence $(\tilde{\eta}_1, \dots, \tilde{\eta}_k)$ where $\tilde{\eta}_j = S^{-1/2} \eta_j$ for $1 \leq j \leq k$ is a normalized tight module frame for $\mathcal{E}$. Hence, $\mathcal{E}$ being finitely generated is equivalent to the existence of a normalized tight module frame in $\mathcal{E}$.

Proof. The equivalence of (i) and (ii) is proved in e.g. [FL02, Proposition 5.9]. The equivalence of (ii) and (iii) is Proposition III.2.1. The last observation follows from writing out the identity

$$\bullet \langle \xi, \xi \rangle = \bullet \langle S^{-1/2} S S^{-1/2} \xi, \xi \rangle$$

for $\xi \in \mathcal{E}$, where S is the frame operator corresponding to the sequence $(\eta_j)_j$. ■

An important application of normalized tight frames is the following, see [Rie10, Proposition 7.2] for a reference.

Proposition III.2.3. *Let A be a unital C^* -algebra. Then every finitely generated left Hilbert A -module is projective. In fact, suppose $\{\eta_1, \dots, \eta_k\}$ is a normalized tight frame for the Hilbert A -module $\mathcal{E}$ (which exists by Proposition III.2.2). Then the $k \times k$ matrix*

$$P = \begin{pmatrix} \bullet\langle\eta_1, \eta_1\rangle & \cdots & \bullet\langle\eta_1, \eta_k\rangle \\ \vdots & \ddots & \vdots \\ \bullet\langle\eta_k, \eta_1\rangle & \cdots & \bullet\langle\eta_k, \eta_k\rangle \end{pmatrix}$$

is a projection in $M_k(A)$, and $\mathcal{E} \cong A^k P$.

Note especially the first sentence of Proposition III.2.3, namely that all finitely generated Hilbert C^* -modules are projective. One can show that this also holds more generally for countably generated projective C^* -modules, but this result will not be important to us.

III.2.3 Modules of sections of a vector bundle

In this section we look at finitely generated Hilbert C^* -modules over commutative C^* -algebras. If A is commutative, then by Gelfand duality, A is $*$ -isomorphic to the C^* -algebra $C(X)$ of continuous functions on a uniquely determined compact Hausdorff space X .

A way to obtain Hilbert C^* -modules over $C(X)$ is from Hermitian vector bundles over X . If $\pi: E \rightarrow X$ is a Hermitian vector bundle, we denote by $\Gamma(E)$ the set of continuous sections of E , i.e. all continuous functions $s: X \rightarrow E$ such that $\pi \circ s = \text{id}_X$. The set $\Gamma(E)$ has the structure of a finitely generated left Hilbert $C(X)$ -module with the following left action and $C(X)$ -valued inner product:

$$(f \cdot s)(x) = f(x)s(x), \quad (\text{III.6})$$

$$\bullet\langle s, t \rangle(x) = \langle s(x), t(x) \rangle_x. \quad (\text{III.7})$$

Here, $f \in C(X)$, $s, t \in \Gamma(E)$, $x \in X$ and $\langle \cdot, \cdot \rangle_x$ denotes the inner product on the fiber E_x of E . We also use the notation $\|v\|_x^2 = \langle v, v \rangle_x$.

That $\Gamma(E)$ is finitely generated comes from the fact that for every vector bundle E over a compact Hausdorff space X , there exists a vector bundle F over X such that $E \oplus F$ is isomorphic to a trivial bundle [Swa62, Corollary 5].

The remarkable fact about finitely generated projective modules over $C(X)$ is that they all come from vector bundles over X . This is known as the Serre–Swan theorem. The version that we state here is slightly modified to include the inner product structures found on Hermitian vector bundles and Hilbert C^* -modules.

Proposition III.2.4 (Serre–Swan theorem). *Suppose $\mathcal{E}$ is a finitely generated left Hilbert $C(X)$ -module. Then there exists a unique Hermitian vector bundle*

III. The BLT for LCA groups and vector bundles

$E \rightarrow X$ such that $\Gamma(E) \cong \mathcal{E}$ as left Hilbert $C(X)$ -modules. If $\mathcal{E}$ is represented as $C(X)^k P$ for a projection P in $M_k(C(X))$, then E can be constructed as the subbundle

$$E = \{(x, v) \in X \times \mathbb{C}^k : vP(x) = v\}$$

of the trivial bundle of rank k .

Proof. Note that by Proposition III.2.3, $\mathcal{E}$ is projective. By the Serre–Swan theorem for finitely generated projective $C(X)$ -modules [Swa62], there exists a unique complex vector bundle $E \rightarrow X$ such that $\Gamma(E) \cong \mathcal{E}$ as left $C(X)$ -modules. The description of E as a subbundle of $X \times \mathbb{C}^k$ also follows from [Swa62]. We suppress the isomorphism from the notation and think of $\mathcal{E}$ as being equal to $\Gamma(E)$.

We define an inner product on the fibers of E as follows: Let $x \in X$, $v, w \in E_x$, and pick sections $s, t \in \Gamma(E)$ such that $s(x) = v$ and $t(x) = w$. Set $\langle v, w \rangle_x = \bullet \langle s, t \rangle(x)$. This is independent of the choice of sections, and will give us the required inner product on E . ■

The next proposition describes generating sets of $\Gamma(E)$ as a left Hilbert $C(X)$ -module, where $E \rightarrow X$ is a Hermitian vector bundle. Recall that a line bundle is a vector bundle where the dimension of the fibers is equal to 1.

Proposition III.2.5. *Let E be a Hermitian vector bundle over a compact Hausdorff space X . Let $s_1, \dots, s_k \in \Gamma(E)$. The following are equivalent:*

- (i) *The set $\{s_1, \dots, s_k\}$ is a generating set for $\Gamma(E)$ as a left $C(X)$ -module.*
- (ii) *For every $x \in X$, we have that*

$$\text{span}\{s_1(x), \dots, s_k(x)\} = E_x.$$

Proof. The frame operator corresponding to the Bessel sequence $(s_j)_{j=1}^k$ in $\Gamma(E)$ is given by

$$Ss = \sum_{j=1}^k \bullet \langle s, s_j \rangle s_j \quad (\text{III.8})$$

for $s \in \Gamma(E)$. This is a $C(X)$ -linear operator $\Gamma(E) \rightarrow \Gamma(E)$, so by [Swa62, Theorem 1] there exists a unique vector bundle homomorphism $\theta : E \rightarrow E$ such that

$$(Ss)(x) = \theta_x(s(x))$$

for all $s \in \Gamma(E)$ and $x \in X$, where θ_x denotes the restriction of θ to E_x . For each $x \in X$, let $S_x : E_x \rightarrow E_x$ denote the frame operator of the sequence $\{s_1(x), \dots, s_k(x)\}$ in the Hilbert space E_x . By (III.8), we have that

$$\theta_x(s(x)) = \sum_{j=1}^k \bullet \langle s, s_j \rangle s_j(x) = \sum_{j=1}^k \langle s(x), s_j(x) \rangle_x s_j(x) = S_x(s(x)) \quad (\text{III.9})$$

for all $s \in \Gamma(E)$ and $x \in X$. Given $v \in E_x$, then letting $s \in \Gamma(E)$ be such that $s(x) = v$, we obtain that $\theta_x(v) = S_x(v)$. Hence, by (III.9), $\theta_x = S_x$ for all $x \in X$.

By Proposition III.2.2, $\{s_1, \dots, s_k\}$ is a generating set for $\Gamma(E)$ if and only if the frame operator S is invertible in $\Gamma(E)$. But this happens if and only if θ is invertible as a map of vector bundles. This is equivalent to each θ_x being invertible for every $x \in X$ by [MS74, Lemma 2.3]. This is equivalent to $\{s_1(x), \dots, s_k(x)\}$ being a generating set for E_x for each $x \in X$, using Proposition III.2.2. This finishes the proof. $\blacksquare$

We immediately obtain the following corollary:

Corollary III.2.6. *Suppose E is a line bundle, and that $s_1, \dots, s_k \in \Gamma(E)$. Then $\{s_1, \dots, s_k\}$ is a generating set for $\Gamma(E)$ if and only if for every $x \in X$, there exists $j \in \{1, \dots, k\}$ such that $s_j(x) \neq 0$. In particular, if $s \in \Gamma(E)$, then $\{s\}$ is a (single) generator for $\Gamma(E)$ if and only if s is nonvanishing on X . Consequently, the existence of a single generator for $\Gamma(E)$ is equivalent to the triviality of E .*

Note that the notion of a module frame in $\Gamma(E)$ is different from the common notions of local and global frames of sections of vector bundles, see e.g. [Lee13, p. 257]. In particular, a global frame for a vector bundle is a basis at every fiber, not just a spanning set.

III.3 Time-frequency analysis and Heisenberg modules

III.3.1 Time-frequency shifts and twisted group C^* -algebras

For the rest of the paper, unless otherwise stated, G will denote a second countable, locally compact abelian group. Denote by $\widehat{G}$ the Pontryagin dual of G . Given a closed subgroup H of G , the *annihilator* of H , denoted by $H^\perp$, is the set

$$H^\perp = \{\omega \in \widehat{G} : \omega(x) = 1 \text{ for all } x \in H\}. \quad (\text{III.10})$$

This will always be a closed subgroup of $\widehat{G}$. We have natural isomorphisms $\widehat{H} \cong \widehat{G}/H^\perp$ and $\widehat{G/H} \cong H^\perp$.

We call a subgroup H of G *cocompact* if the quotient G/H is compact. A subgroup Λ of G that is both discrete and cocompact is called a *lattice* in G . If Λ is a lattice in G , then $\Lambda^\perp$ is a lattice in $\widehat{G}$ [Rie88, Lemma 3.1].

Given $x \in G$ and $\omega \in \widehat{G}$, the unitary linear operators T_x and M_ω on $L^2(G)$ given by

$$T_x \xi(t) = \xi(x^{-1}t) \quad M_\omega \xi(t) = \omega(t) \xi(t) \quad (\text{III.11})$$

for $\xi \in L^2(G)$, $t \in G$, are called *time shift by x* and *frequency shift by ω* , respectively. These two operators satisfy the following commutation relation:

$$M_\omega T_x = \omega(x) T_x M_\omega. \quad (\text{III.12})$$

III. The BLT for LCA groups and vector bundles

A *time-frequency shift* is a combined operator of the form $\pi(x, \omega) = M_\omega T_x$. From (III.12), it follows that

$$\pi(x, \omega)\pi(y, \tau) = \overline{\tau(x)}\pi(xy, \omega\tau) \quad (\text{III.13})$$

for $(x, \omega), (y, \tau) \in G \times \widehat{G}$. The function $c: (G \times \widehat{G}) \times (G \times \widehat{G}) \rightarrow \mathbb{T}$ given by

$$c((x, \omega), (y, \tau)) = \overline{\tau(x)} \quad (\text{III.14})$$

satisfies the identities

$$\begin{aligned} c(z_1, z_2)c(z_1 z_2, z_3) &= c(z_1, z_2 z_3)c(z_2, z_3) \\ c(1, 1) &= 1 \end{aligned}$$

for all $z_1, z_2, z_3 \in G \times \widehat{G}$ and with 1 denoting the identity element of $G \times \widehat{G}$. Thus, when restricting c to any lattice Δ in $G \times \widehat{G}$, we obtain a 2-cocycle on Δ . Restricting also π to Δ , we obtain a *c-projective unitary representation* of the group Δ on the Hilbert space $L^2(G)$ [Wil07, Chapter D3].

Whenever we have a 2-cocycle on a discrete group, we can associate to it the corresponding *twisted group C^* -algebra*. In our case, we have the twisted group C^* -algebra $C^*(\Delta, c)$, which is the C^* -enveloping algebra of the Banach $*$ -algebra $\ell^1(\Delta, c)$. As a Banach space, $\ell^1(\Delta, c)$ is $\ell^1(\Delta)$, but the multiplication is *c-twisted convolution* and the involution is *c-twisted involution*. These operations are defined as follows, where $a, b \in \ell^1(\Delta)$ and $z \in \Delta$:

$$\begin{aligned} (a * b)(z) &= \sum_{w \in \Delta} c(w, w^{-1}z) a(w) b(w^{-1}z) \\ a^*(z) &= \overline{c(z, z^{-1}) a(z^{-1})}. \end{aligned}$$

This C^* -algebra captures the representation theory of Δ in the following way: The *c-projective unitary representations* of Δ are in 1–1 correspondence with nondegenerate $*$ -representations of $C^*(\Delta, c)$. In particular, our *c-projective representation* $\pi|_\Delta$ lifts to a representation $C^*(\Delta, c) \rightarrow \mathcal{B}(L^2(G))$, which we by abuse of notation also denote by π . On the dense $*$ -subalgebra $\ell^1(\Delta, c)$, π is given by

$$\pi(a) = \sum_{z \in \Delta} a(z) \pi(z) \quad (\text{III.15})$$

for $a \in \ell^1(\Delta, c)$. This representation is faithful [Rie88, Proposition 2.2].

III.3.2 Gabor frames

As before, let G be a second countable, locally compact abelian group. Let Δ be a lattice in $G \times \widehat{G}$, and let $\eta \in L^2(G)$. The *Gabor system generated by η over Δ* is the set

$$\mathcal{G}(\eta, \Delta) = \{\pi(z)\eta : z \in \Delta\}. \quad (\text{III.16})$$

If this set is a frame for the Hilbert space $L^2(G)$ as in (III.1), we call it a *Gabor frame* over Δ with generator η . That is, $\mathcal{G}(\eta, \Delta)$ is a Gabor frame if there exist $K, L > 0$ such that

$$K\|\xi\|_2^2 \leq \sum_{z \in \Delta} |\langle \xi, \pi(z)\eta \rangle|^2 \leq L\|\xi\|_2^2$$

for all $\xi \in L^2(G)$.

Generalizing, if $\eta_1, \dots, \eta_k \in L^2(G)$, then the *multiwindow Gabor system* generated by the $\eta_1, \dots, \eta_k$ over Δ is the set

$$\mathcal{G}(\eta_1, \dots, \eta_k; \Delta) := \{\pi(z)\eta_j : z \in \Delta, 1 \leq j \leq k\}. \quad (\text{III.17})$$

If this set is a frame for $L^2(G)$, then we call it a *multiwindow Gabor frame* over Δ with generators $\eta_1, \dots, \eta_k$.

Remark III.3.1. It is possible to consider Gabor systems over closed subgroups Δ of $G \times \widehat{G}$ that are not discrete. This leads to the notion of continuous Gabor systems and frames, see [JL16]. In our case, we will eventually consider Δ to be of the form $\Lambda \times \Lambda^\perp$, with Λ a closed subgroup of G . It turns out that for frames to exist over such a closed subgroup of $G \times \widehat{G}$, Λ must be a lattice in G [JL16, Corollary 5.8], so we will not lose anything interesting by assuming that Δ is discrete.

The *Feichtinger algebra* is the set $S_0(G)$ of all functions $\xi \in L^2(G)$ for which

$$\int_{G \times \widehat{G}} |\langle \xi, \pi(z)\xi \rangle| \, dz < \infty. \quad (\text{III.18})$$

The Feichtinger algebra has many different descriptions, see [Jak18]. Functions in $S_0(G)$ are continuous.

III.3.3 Heisenberg modules

As usual, let G denote a second countable, locally compact abelian group. Let Δ be a lattice in the time-frequency plane $G \times \widehat{G}$.

We now proceed to review the construction by Rieffel in [Rie88] of the modules that have been termed *Heisenberg modules*. We follow the approach by Jakobsen and Luef in [JL18] and use the Feichtinger algebra $S_0(G)$ instead of the Schwartz–Bruhat space $\mathcal{S}(G)$ from [Rie88]. Although the Heisenberg module has a natural structure as an imprimitivity bimodule, only the left module structure will be important to us.

By the proof of [JL18, Theorem 3.4], we have the following:

Proposition III.3.2. *Let G be a second countable, locally compact abelian group, and let Δ be a lattice in $G \times \widehat{G}$. Then the Feichtinger algebra $S_0(G)$ can be completed into a full finitely generated left Hilbert $C^*(\Delta, c)$ -module which we denote by $\mathcal{E}_\Delta(G)$. If $a \in \ell^1(\Delta, c)$ and $\xi \in S_0(G)$, then $a \cdot \xi \in S_0(G)$ and is given by*

$$a \cdot \xi = \sum_{z \in \Delta} a(z)\pi(z)\xi. \quad (\text{III.19})$$

If $\xi, \eta \in S_0(G)$, then $\bullet\langle\xi, \eta\rangle \in \ell^1(\Delta, c)$ and is given by

$$\bullet\langle\xi, \eta\rangle(z) = \langle\xi, \pi(z)\eta\rangle \quad (\text{III.20})$$

for $z \in \Delta$.

The module $\mathcal{E}_\Delta(G)$ is called a *Heisenberg module*. Note that $\mathcal{E}_\Delta(G)$ is finitely generated in our case since Δ is assumed to be a lattice in $G \times \widehat{G}$ (see the proof of [Rie88, Proposition 3.3] and [JL18, Theorem 3.9]).

The following is proved in [AE19, Proposition 3.2, Proposition 3.6, Theorem 3.11], and gives a characterization of generators of Heisenberg modules as exactly multiwindow Gabor frames as in (III.17).

Proposition III.3.3. *Let Δ be a lattice in $G \times \widehat{G}$. Then the following hold:*

- (i) *There is a continuous embedding of the Heisenberg module $\mathcal{E}_\Delta(G)$ into $L^2(G)$ that makes the following diagram commute:*

$$\begin{array}{ccc} S_0(G) & \longrightarrow & \mathcal{E}_\Delta(G) \\ & \searrow & \downarrow \\ & & L^2(G) \end{array}$$

Under this embedding, we have that

$$\|\eta\|_2 \leq \|\eta\|_{\mathcal{E}_\Delta(G)} \quad (\text{III.21})$$

for every $\eta \in \mathcal{E}_\Delta(G)$.

- (ii) *The norm on $\mathcal{E}_\Delta(G)$ is given by*

$$\|\eta\|_{\mathcal{E}_\Delta(G)} = \sup_{\|\xi\|_2=1} \left(\sum_{z \in \Delta} |\langle\xi, \pi(z)\eta\rangle|^2 \right)^{1/2}$$

for $\eta \in \mathcal{E}_\Delta(G)$. Thus, $\mathcal{E}_\Delta(G)$ can be described as the completion of $S_0(G)$ in the Banach space consisting of those $\eta \in L^2(G)$ for which $\mathcal{G}(\eta, \Delta)$ is a Bessel sequence for $L^2(G)$.

- (iii) *A subset $\{\eta_1, \dots, \eta_k\}$ of $\mathcal{E}_\Delta(G)$ is a generating set for $\mathcal{E}_\Delta(G)$ if and only if $\mathcal{G}(\eta_1, \dots, \eta_k; \Delta)$ is a multiwindow Gabor frame for $L^2(G)$ (see (III.16)).*

Remark III.3.4. Note that in (III.21), there is no appearance of the constant $s(\Delta)^{1/2}$ as in [AE19, Proposition 3.2]. The reason for this is that in the case of a lattice, both traces in the setting of [AE19, Convention 3.1] are finite. Thus, one can consider the trace tr_A instead of tr_B when proving the inequality of the $L^2(G)$ -norm and the Heisenberg module norm. Since tr_A is a state in this setting, we have $\|\text{tr}_A\| = 1$, so we obtain (III.21).

We are now ready to obtain a useful reformulation of Statement III.1.2 in terms of Heisenberg modules:

Proposition III.3.5. *Let $\Delta = \Lambda \times \Lambda^\perp$ for a lattice Λ in G . Then the following are equivalent:*

- (i) *The Heisenberg module $\mathcal{E}_\Delta(G)$ is not singly generated.*
- (ii) *For every $\eta \in \mathcal{E}_\Delta(G)$, the Gabor system $\mathcal{G}(\eta, \Delta)$ is not a frame for $L^2(G)$.*
- (iii) *Statement III.1.2 holds for (G, Λ) . That is, for every $\eta \in S_0(G)$, the Gabor system $\mathcal{G}(\eta, \Delta)$ is not a frame for $L^2(G)$.*

Proof. The equivalence of (i) and (ii) follows immediately from Proposition III.3.3 (iii), and (ii) implies (iii) since $S_0(G) \subseteq \mathcal{E}_\Delta(G)$. It remains to prove that (iii) implies (i).

First, note the following: If $\{\eta_1, \dots, \eta_k\}$ is a module frame for $\mathcal{E}_\Delta(G)$, then using the same procedure as in the proof of [Rie88, Proposition 3.7], one obtains a new module frame $\{\eta'_1, \dots, \eta'_k\}$ for $\mathcal{E}_\Delta(G)$ where $\eta'_j \in S_0(G)$ for $1 \leq j \leq k$. The argument works out because $\ell^1(\Delta, c)$, like the Schwartz–Bruhat space, is spectrally invariant in $C^*(\Delta, c)$. As pointed out in [Lue09, Theorem 2.2], this is a consequence of [GL04, Theorem 3.1] in the case of $G = \mathbb{R}^n$. However, as pointed out in [GL04, p. 16], the proofs generalize in a straightforward manner to the locally compact abelian case.

Thus, the existence of a one-element module frame (i.e. a single generator) $\eta \in \mathcal{E}_\Delta(G)$ implies the existence of a one-element module frame (i.e. single generator) $\eta' \in S_0(G)$. Thus, if no single generators can be found in $S_0(G)$, no single generators can be found in $\mathcal{E}_\Delta(G)$, which finishes the proof. ■

III.4 Vector bundles and quasiperiodic functions

III.4.1 The Zak transform

Throughout this section, we assume that G is a second countable, locally compact abelian group, and that Λ is a lattice in G .

The *Zak transform* [Gel50; Wei64] of a function $\xi \in L^2(G)$ with respect to Λ is the function $Z\xi = Z_{G,\Lambda}\xi: G \times \widehat{G} \rightarrow \mathbb{C}$ given by

$$Z\xi(x, \omega) = \sum_{\lambda \in \Lambda} \xi(x\lambda) \omega(\lambda) \quad (\text{III.22})$$

for $(x, \omega) \in G \times \widehat{G}$. By [KK98, Lemma 3], the Zak transform of a function $\xi \in L^2(G)$ is defined almost everywhere on $G \times \widehat{G}$.

The following result is proved for $G = \mathbb{R}^n$ in [Grö01, Lemma 8.2.1 (c)]. While the author believes the result for general locally compact abelian groups to be well-known, he was not able to find a reference for it, so it is proved in the appendix, see Proposition III.B.3.

Proposition III.4.1. *Let Λ be a lattice in a second countable, locally compact abelian group G . If $\xi \in S_0(G)$, then $Z_{G,\Lambda}\xi$ is continuous.*

Now note that for $\xi \in L^2(G)$, $(x, \omega) \in G \times \widehat{G}$ and $(\lambda, \tau) \in \Lambda \times \Lambda^\perp$, we have the following:

$$\begin{aligned} Z\xi(x\lambda, \omega\tau) &= \sum_{\lambda' \in \Lambda} \xi(x\lambda\lambda')(\omega\tau)(\lambda') \\ &= \sum_{\lambda' \in \Lambda} \xi(x\lambda')\omega(\lambda'\lambda^{-1}) \\ &= \overline{\omega(\lambda)}Z\xi(x, \omega). \end{aligned}$$

Thus, if $\xi \in S_0(G)$, then by Proposition III.4.1 and the above calculation, $Z\xi$ is an element of the following function space:

Definition III.4.2. Denote by $\mathcal{QP}_{G,\Lambda}$ the complex vector space of continuous functions $F: G \times \widehat{G} \rightarrow \mathbb{C}$ satisfying the relation

$$F(x\lambda, \omega\tau) = \overline{\omega(\lambda)}F(x, \omega) \quad (\text{III.23})$$

for all $(x, \omega) \in G \times \widehat{G}$ and $(\lambda, \tau) \in \Lambda \times \Lambda^\perp$.

The relation in (III.23) is referred to as a *quasiperiodicity relation* in the literature [KK98], so we will call elements of $\mathcal{QP}_{G,\Lambda}$ continuous, quasiperiodic functions on $G \times \widehat{G}$.

Define $X_{G,\Lambda}$ to be the space

$$X_{G,\Lambda} = (G/\Lambda) \times (\widehat{G}/\Lambda^\perp). \quad (\text{III.24})$$

Since Λ is a lattice in G , this is a compact space. The linear space $\mathcal{QP}_{G,\Lambda}$ becomes a left $C(X_{G,\Lambda})$ -module with the pointwise action

$$(f \cdot F)(x, \omega) = f([x], [\omega])F(x, \omega) \quad (\text{III.25})$$

for $(x, \omega) \in G \times \widehat{G}$, $f \in C(X_{G,\Lambda})$ and $F \in \mathcal{QP}_{G,\Lambda}$. Moreover, we define a $C(X_{G,\Lambda})$ -valued inner product on $\mathcal{QP}_{G,\Lambda}$ by

$$\bullet \langle F, G \rangle ([x], [\omega]) = F(x, \omega) \overline{G(x, \omega)} \quad (\text{III.26})$$

for $(x, \omega) \in G \times \widehat{G}$ and $F, G \in \mathcal{QP}_{G,\Lambda}$. The above structure is easily seen to satisfy the first three axioms in the definition of a Hilbert $C(X_{G,\Lambda})$ -module from Section III.2. The norm induced by the $C(X_{G,\Lambda})$ -valued inner product is just the supremum norm, so any Cauchy sequence $(F_n)_{n \in \mathbb{N}}$ of continuous, quasiperiodic functions will have a uniform limit $F \in C_b(G \times \widehat{G})$. Continuity then gives that the limit is also quasiperiodic, showing that $\mathcal{QP}_{G,\Lambda}$ is in fact a Hilbert C^* -module over $C(X_{G,\Lambda})$.

Our aim is to show that $\mathcal{QP}_{G,\Lambda}$ is finitely generated as a $C(X_{G,\Lambda})$ -module, so that it corresponds to a Hermitian vector bundle over $X_{G,\Lambda}$ by Proposition III.2.4. To do this, we begin by characterizing finite module frames of $\mathcal{QP}_{G,\Lambda}$:

Proposition III.4.3. *Let $\{G_1, \dots, G_k\}$ be a subset of $\text{QP}_{G,\Lambda}$. Then the following hold:*

- (i) *$\{G_1, \dots, G_k\}$ is a module frame in $\text{QP}_{G,\Lambda}$ if and only if for every $(x, \omega) \in G \times \widehat{G}$, there exists $1 \leq j \leq k$ such that $G_j(x, \omega) \neq 0$.*
- (ii) *$\{G_1, \dots, G_k\}$ is a normalized tight module frame in $\text{QP}_{G,\Lambda}$ if and only if*

$$\sum_{j=1}^k |G_j(x, \omega)|^2 = 1$$

for every $(x, \omega) \in G \times \widehat{G}$.

In particular, a single generator $\{G\}$ for $\text{QP}_{G,\Lambda}$ is just a nonvanishing continuous quasiperiodic function on $G \times \widehat{G}$.

Proof. Note that the module frame condition from (III.1) on $\{G_1, \dots, G_k\}$ translates into the existence of $K, L > 0$ such that

$$K|F(x, \omega)|^2 \leq \sum_{j=1}^k F(x, \omega) \overline{G_j(x, \omega)} F(x, \omega) G_j(x, \omega) \leq L|F(x, \omega)|^2$$

for all $F \in \text{QP}_{G,\Lambda}$ and all $(x, \omega) \in G \times \widehat{G}$. Hence, cancelling $|F(x, \omega)|^2$, we obtain the condition

$$K \leq \sum_{j=1}^k |G_j(x, \omega)|^2 \leq L$$

for all $(x, \omega) \in G \times \widehat{G}$. Since each $|G_j|^2$ can be seen as a continuous function on the compact space $X_{G,\Lambda}$, it is clear that the upper frame condition automatically holds by the extreme value theorem. For the same reason, the lower frame condition is equivalent to the expression $\sum_{j=1}^k |G_j(x, \omega)|^2$ being nonvanishing, which is equivalent to (i). Now (ii) follows directly by considering $K = L = 1$ in this situation. $\blacksquare$

The following proposition shows that $\text{QP}_{G,\Lambda}$ is finitely generated. In fact, one can always find a module frame consisting of images of elements in $S_0(G)$ under the Zak transform. Recall that T_x denotes a time shift as in (III.11).

Proposition III.4.4. *Let U be an open neighbourhood of 1 in G such that the collection $\{\lambda U : \lambda \in \Lambda\}$ is pairwise disjoint, let $K \subseteq U$ be a nonempty compact set, and let ξ a function in $S_0(G)$ satisfying $\xi|_K = 1$ and $\text{supp}(\xi) \subseteq U$ (the existence of such a function is guaranteed by Proposition III.B.2). Then there exist $x_1, \dots, x_k \in G$ such that*

$$\{Z_{G,\Lambda}(T_{x_1}\xi), \dots, Z_{G,\Lambda}(T_{x_k}\xi)\}$$

is a module frame for $\text{QP}_{G,\Lambda}$. Consequently, $\text{QP}_{G,\Lambda}$ is a finitely generated left Hilbert $C(X_{G,\Lambda})$ -module.

Proof. Let $(x, \omega) \in G \times \widehat{G}$. Then xU is an open neighbourhood of x , and the sets $\{\lambda xU : \lambda \in \Lambda\}$ are pairwise disjoint. Moreover, if $y \in xK$ and $\lambda \in \Lambda$ satisfy $\xi(x^{-1}y\lambda) \neq 0$, then $x^{-1}y\lambda \in \text{supp}(\xi) \subseteq U$. Thus $x^{-1}y \in U \cap (\lambda^{-1}U)$ which forces $\lambda = 1$. Consequently, we have that

$$Z(T_x\xi)(y, \omega) = \sum_{\lambda \in \Lambda} \xi(x^{-1}y\lambda)\omega(\lambda) = \xi(x^{-1}y) = 1. \quad (\text{III.27})$$

By (III.23) and (III.27), we have that that

$$Z(T_x\xi)(y\lambda', \omega) = \overline{\omega(\lambda')}Z(T_x\xi)(y, \omega) = \overline{\omega(\lambda')}$$

whenever $y \in xK$ and $\lambda' \in \Lambda$. This shows that the function $Z(T_x\xi)$ is nonvanishing on $xK^\circ\Lambda \subseteq xK\Lambda$. Now $\{xK^\circ : x \in G\}$ is an open cover of G , and Λ is cocompact in G , so we can find a finite cover of G of the form $\{x_jK^\circ\Lambda : 1 \leq j \leq k\}$ for $x_1, \dots, x_k \in G$. For each $1 \leq j \leq k$, $F_j := Z_{G,\Lambda}(T_{x_j}\xi)$ is nonvanishing on $x_jK^\circ\Lambda$. By Proposition III.4.1, each F_j is in $\text{QP}_{G,\Lambda}$. Thus, by Proposition III.4.3, the set $\{F_1, \dots, F_k\}$ is a module frame for $\text{QP}_{G,\Lambda}$.

Since a module frame is the same as a generating set by Proposition III.2.1, $\text{QP}_{G,\Lambda}$ is finitely generated as a left $C(X_{G,\Lambda})$ -module. $\blacksquare$

III.4.2 The line bundle $E_{G,\Lambda}$

In this section, we construct the line bundle mentioned in Theorem B and Theorem C in the introduction.

We begin with the left Hilbert $C(X_{G,\Lambda})$ -module $\text{QP}_{G,\Lambda}$ defined in Definition III.4.2, which was shown to be finitely generated in Proposition III.4.4. By Proposition III.2.4, there exists a Hermitian vector bundle $E \rightarrow X_{G,\Lambda}$ such that $\Gamma(E) \cong \text{QP}_{G,\Lambda}$. In fact, letting $\{G_1, \dots, G_k\}$ be a normalized tight frame for $\text{QP}_{G,\Lambda}$, we have that $\text{QP}_{G,\Lambda} \cong C(X_{G,\Lambda})^k P$, where

$$P = \begin{pmatrix} \bullet \langle G_1, G_1 \rangle & \cdots & \bullet \langle G_1, G_k \rangle \\ \vdots & \ddots & \vdots \\ \bullet \langle G_k, G_1 \rangle & \cdots & \bullet \langle G_k, G_k \rangle \end{pmatrix} = \begin{pmatrix} G_1 \overline{G_2} & \cdots & G_1 \overline{G_k} \\ \vdots & \ddots & \vdots \\ G_k \overline{G_1} & \cdots & G_k \overline{G_k} \end{pmatrix}.$$

It then follows from Proposition III.2.4 that

$$E \cong \{([x], [\omega], v) \in X_{G,\Lambda} \times \mathbb{C}^k : vP([x], [\omega]) = v\}.$$

The condition $vP([x], [\omega]) = v$ for $([x], [\omega]) \in X_{G,\Lambda}$ and $v = (v_j)_{j=1}^k \in \mathbb{C}^k$ translates into

$$v_j = \sum_{i=1}^k G_i(x, \omega) \overline{G_j(x, \omega)} v_i = \overline{G_j(x, \omega)} \sum_{i=1}^k G_i(x, \omega) v_i$$

for each $1 \leq j \leq k$. Thus, $v_j / \overline{G_j(x, \omega)}$ is constant in j , so we can write v as a complex multiple of the vector $(\overline{G_j(x, \omega)})_{j=1}^k$. It follows that we can describe E as

$$E \cong \{([x], [\omega], (z \overline{G_j(x, \omega)})_{j=1}^k) : x \in G, \omega \in \widehat{G}, z \in \mathbb{C}\}.$$

This shows that E is in fact a line bundle. Now we want to obtain a description of E independent of the chosen module frame. Note that the map $G \times \widehat{G} \times \mathbb{C} \rightarrow E_{G,\Lambda}$ given by

$$(x, \omega, z) \mapsto ([x], [\omega], (z \overline{G_j(x, \omega)})_{j=1}^k)$$

is continuous and surjective, and that (x, ω, z) and (x', ω', z') has the same image under this map if and only if there exist $\lambda \in \Lambda$ and $\tau \in \Lambda^\perp$ such that

$$\begin{aligned} x' &= x\lambda, \\ \omega' &= \omega\tau, \\ z' &= \overline{\omega(\lambda)}z. \end{aligned}$$

Thus, we get a continuous bijection

$$E_{G,\Lambda} := (G \times \widehat{G} \times \mathbb{C}) / \sim \rightarrow E$$

where $\sim$ is the equivalence relation defined above. If we transfer the vector bundle structure from E to $E_{G,\Lambda}$ via this map, then the projection $\pi: E_{G,\Lambda} \rightarrow X_{G,\Lambda}$ is given by $\pi([x, \omega, z]) = ([x], [\omega])$ for $[x, \omega, z] \in \tilde{E}_{G,\Lambda}$. The vector space structure on the fiber

$$\pi^{-1}([x], [\omega]) = \{[x, \omega, z] : z \in \mathbb{C}\}$$

is given by

$$[x, \omega, z] + [x, \omega, w] = [x, \omega, z + w], \quad (\text{III.28})$$

$$\mu[x, \omega, z] = [x, \omega, \mu z] \quad (\text{III.29})$$

for $x \in G$, $\omega \in \widehat{G}$ and $z, w, \mu \in \mathbb{C}$. Furthermore, the inner product on the same fiber is given by

$$\langle [x, \omega, z], [x, \omega, w] \rangle_{([x], [\omega])} = z\overline{w}. \quad (\text{III.30})$$

Since the map $E_{G,\Lambda} \rightarrow E$ restricts to a linear isomorphism on each fiber, it is a vector bundle isomorphism by [MS74, Lemma 2.3]. We have thus obtained a description of E independent on a chosen module frame. We summarize our results in the following proposition:

Proposition III.4.5. *Let $E_{G,\Lambda}$ denote the quotient of $G \times \widehat{G} \times \mathbb{C}$ by the equivalence relation given by $(x, \omega, z) \sim (x', \omega', z')$ if and only if there exist $\lambda \in \Lambda$ and $\tau \in \Lambda^\perp$ such that*

$$\begin{aligned} x' &= x\lambda, \\ \omega' &= \omega\tau, \\ z' &= \overline{\omega(\lambda)}z. \end{aligned}$$

Then $E_{G,\Lambda}$ has the structure of a Hermitian line bundle with operations given in (III.28), (III.29) and (III.30), and the map $\Psi: \text{QP}_{G,\Lambda} \rightarrow \Gamma(E_{G,\Lambda})$ given by

$$\Psi(F)([x], [\omega]) = [x, \omega, F(x, \omega)]$$

is an isomorphism of left Hilbert $C(X_{G,\Lambda})$ -modules.

Example III.4.6. We compare the bundle $E_{G,\Lambda}$ in the case of $G = \mathbb{R}$ and $\Lambda = \mathbb{Z}$ to the bundle ξ in [Bal01] in the case of $p = 1$. Then $1 = \alpha\beta = p/q = 1$ so that $q = 1$, and thus we can set $r_0 = 1$ and $n_0 = 0$ as well (see equation (21)). The matrix $E(t)$ used to define ξ in Equations (32) and (33) in [Bal01] becomes the 1×1 matrix $E(t) = e^{-2\pi it}$, so the Equations (32) and (33) in [Bal01] reduce to the relations that define $E_{\mathbb{R},\mathbb{Z}}$, modulo a sign. This shows that ξ is the dual bundle of $E_{\mathbb{R},\mathbb{Z}}$. However, since the Picard group of $\mathbb{T}^2$ (the abelian group of line bundles under tensor product where inversion is given by taking dual bundles) is isomorphic to $\mathbb{Z}$, a line bundle over $\mathbb{T}^2$ is nontrivial if and only if its dual bundle is nontrivial. Hence the nontriviality of ξ is equivalent to the nontriviality of $E_{\mathbb{R},\mathbb{Z}}$, and the nontriviality of the former bundle was demonstrated in [Bal01].

III.5 Connecting Heisenberg modules and vector bundles

III.5.1 The Zak transform as an isomorphism of Hilbert C^* -modules

We now combine the Heisenberg modules from Section III.3.3 and the setting of Statement III.1.2. We consider the Heisenberg module $\mathcal{E}_\Delta(G)$ over the twisted group C^* -algebra $C^*(\Delta, c)$, where the lattice $\Delta \subseteq G \times \widehat{G}$ is of the form $\Delta = \Lambda \times \Lambda^\perp$ for a lattice Λ in G . The 2-cocycle c from (III.14) restricted to Δ is given by

$$c((\lambda, \tau), (\lambda', \tau')) = \overline{\tau'(\lambda)} = 1$$

since $\lambda \in \Lambda$ and $\tau \in \Lambda^\perp$. Thus, the 2-cocycle is constantly equal to 1 on $\Lambda \times \Lambda^\perp$ and the twisted group C^* -algebra $C^*(\Lambda \times \Lambda^\perp, c)$ becomes the un-twisted group C^* -algebra $C^*(\Lambda \times \Lambda^\perp)$.

The group $\Lambda \times \Lambda^\perp$ is abelian, and its Pontryagin dual can be identified with $X_{G,\Lambda}$ from Section III.4.2 as follows:

$$\begin{aligned} \widehat{\Lambda \times \Lambda^\perp} &\cong (G \times \widehat{G}) / ((\Lambda \times \Lambda^\perp)^\perp) && \text{using the identity } \widehat{H} \cong \widehat{G}/H^\perp \\ &\cong (\widehat{G}/\Lambda^\perp) \times (G/\Lambda) \\ &\cong (G/\Lambda) \times (\widehat{G}/\Lambda^\perp) \\ &= X_{G,\Lambda}. \end{aligned}$$

Note that we flipped the two spaces in the last homeomorphism. Thus, we get $C^*(\Lambda \times \Lambda^\perp) \cong C(X_{G,\Lambda})$ via the Fourier transform, see [Wil07, Proposition 3.1]. We will use the Fourier transform $\phi: C^*(\Lambda \times \Lambda^\perp) \rightarrow C(X_{G,\Lambda})$ given by

$$\phi(a)([x], [\omega]) = \sum_{\lambda \in \Lambda, \tau \in \Lambda^\perp} a(\lambda, \tau) \omega(\lambda) \tau(x) \quad (\text{III.31})$$

for $a \in \ell^1(\Lambda \times \Lambda^\perp)$, $x \in G$, $\omega \in \widehat{G}$, to identify the two C^* -algebras.

Now the Heisenberg module $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$ is a finitely generated Hilbert C^* -module over $C^*(\Lambda \times \Lambda^\perp) \cong C(X_{G,\Lambda})$ by Proposition III.3.2. By the Serre–Swan

theorem (Proposition III.2.4), we know that $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$ must be isomorphic to the continuous sections of a Hermitian vector bundle over $X_{G,\Lambda}$. We now prove that the bundle in question is $E_{G,\Lambda}$ from Section III.4.2, and that the precise isomorphism is implemented by the Zak transform from Section III.4.1:

Theorem III.5.1. *Let Λ be a lattice in a second countable, locally compact abelian group G . Consider the Heisenberg module $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$ over $C^*(\Lambda \times \Lambda^\perp)$, with $S_0(G)$ as a dense $\ell^1(\Lambda \times \Lambda^\perp)$ -submodule. Let ϕ denote the Fourier transform from (III.31). For $a \in \ell^1(\Lambda \times \Lambda^\perp)$ and $\xi, \eta \in S_0(G)$, we have that*

$$Z(a \cdot \xi) = \phi(a) \cdot Z(\xi), \quad (\text{III.32})$$

$$\phi(\bullet \langle \xi, \eta \rangle) = \bullet \langle Z\xi, Z\eta \rangle. \quad (\text{III.33})$$

Using ϕ to identify $C^*(\Lambda \times \Lambda^\perp)$ with $C(X_{G,\Lambda})$, the Zak transform given in (III.22) is an isomorphism of left Hilbert modules

$$Z_{G,\Lambda} : \mathcal{E}_{\Lambda \times \Lambda^\perp}(G) \rightarrow \text{QP}_{G,\Lambda}.$$

Proof. First, Proposition III.4.1 ensures that Z maps $S_0(G)$ into $\text{QP}_{G,\Lambda}$. To show (III.32), we begin by letting $a = \delta_{(\lambda_0, \tau_0)} \in \ell^1(\Lambda \times \Lambda^\perp)$ for $(\lambda_0, \tau_0) \in \Lambda \times \Lambda^\perp$ and $\xi \in S_0(G)$. Then ϕ maps $\delta_{(\lambda_0, \tau_0)}$ to the function on $(G/\Lambda) \times (\widehat{G}/\Lambda^\perp)$ given by $([x], [\omega]) \mapsto \omega(\lambda_0)\tau_0(x)$. Therefore, letting $x \in G$ and $\omega \in \widehat{G}$, we obtain

$$\begin{aligned} Z(\delta_{(\lambda_0, \tau_0)} \cdot \xi)(x, \omega) &= \sum_{\lambda \in \Lambda} \pi(\lambda_0, \tau_0) \xi(x\lambda) \omega(\lambda) && \text{by (III.19)} \\ &= \sum_{\lambda \in \Lambda} \xi(x\lambda\lambda_0^{-1}) \tau_0(x\lambda) \omega(\lambda) \\ &= \sum_{\lambda \in \Lambda} \xi(x\lambda) \tau_0(x\lambda\lambda_0) \omega(\lambda\lambda_0) && \text{via } \lambda \mapsto \lambda\lambda_0 \\ &= \omega(\lambda_0) \tau_0(x) \sum_{\lambda \in \Lambda} \xi(x\lambda) \omega(\lambda) \\ &= (\phi(\delta_{(\lambda_0, \tau_0)}) \cdot (Z\xi))(x, \omega). \end{aligned}$$

By linearity and continuity, this proves (III.32) for all $a \in \ell^1(\Lambda \times \Lambda^\perp)$.

We move on to proving (III.33). Let $\xi, \eta \in S_0(G)$. Denote by $\mathcal{F}$ the usual Fourier transform $S_0(G) \rightarrow S_0(\widehat{G})$. Since the Poisson summation formula holds for functions in $S_0(G)$ [Jak18, Theorem 5.7 (iii)] and $S_0(G)$ is an algebra under pointwise multiplication, we have that the Poisson summation formula holds for the function $t \mapsto \xi(tx)\overline{\eta(\lambda^{-1}tx)}$, with $x \in G$, $\omega \in \widehat{G}$. We do the following calculation, where the Poisson summation formula is applied in the fifth equality:

$$\begin{aligned} \phi(\bullet \langle \xi, \eta \rangle)([x], [\omega]) &= \sum_{\lambda \in \Lambda, \tau \in \Lambda^\perp} \langle \xi, \pi(\lambda, \tau)\eta \rangle \omega(\lambda) \tau(x) \\ &= \sum_{\lambda \in \Lambda, \tau \in \Lambda^\perp} \int_G \xi(t) \overline{\eta(\lambda^{-1}t) \tau(t)} \omega(\lambda) \tau(x) \, dt \end{aligned}$$

$$\begin{aligned}
 &= \sum_{\lambda \in \Lambda, \tau \in \Lambda^\perp} \int_G \xi(tx) \overline{\eta(\lambda^{-1}tx) \tau(t)} \omega(\lambda) \, dt \quad (\text{via } t \mapsto tx) \\
 &= \sum_{\lambda \in \Lambda} \sum_{\tau \in \Lambda^\perp} \mathcal{F}[t \mapsto \xi(tx) \overline{\eta(\lambda^{-1}tx)}](\tau) \omega(\lambda) \\
 &= \sum_{\lambda \in \Lambda} \sum_{\lambda' \in \Lambda} \xi(\lambda'x) \overline{\eta(\lambda^{-1}\lambda'x)} \omega(\lambda) \quad (\text{by Poisson summation}) \\
 &= \sum_{\lambda, \lambda' \in \Lambda} \xi(x\lambda') \overline{\eta(x\lambda)} \omega(\lambda) \omega(\lambda') \quad (\text{via } \lambda \mapsto \lambda\lambda') \\
 &= Z\xi(x, \omega) \overline{Z\eta(x, \omega)}.
 \end{aligned}$$

This proves (III.33).

We have now established that the Zak transform is linear and inner product preserving (hence an isometry) on $S_0(G)$ and preserves the action of $\ell^1(\Lambda \times \Lambda^\perp)$ on $S_0(G)$.

Now take $\xi \in \mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$, and let $(\xi_n)_n$ be a sequence in $S_0(G)$ such that $\|\xi - \xi_n\|_{\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)} \rightarrow 0$. Since Z is an isometry on $S_0(G)$, we have that

$$\|Z\xi_m - Z\xi_n\|_\infty = \|Z(\xi_m - \xi_n)\|_\infty = \|\xi_m - \xi_n\|_{\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)}$$

for $m, n \in \mathbb{N}$. Since $(\xi_n)_n$ converges in the $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$ -norm, we have that $(Z\xi_n)_n$ converges uniformly to a function $F \in \mathbf{QP}_{G, \Lambda}$.

By [KK98, Remark 4], the Zak transform can be viewed as a unitary map from $L^2(G)$ to $L^2(B \times B')$, where B and B' are any fundamental domains for Λ in G and $\Lambda^\perp$ in $\widehat{G}$, respectively. Hence we have, using (III.21), that

$$\|Z\xi - Z\xi_n\|_{L^2(B \times B')} = \|\xi - \xi_n\|_{L^2(G)} \leq \|\xi - \xi_n\|_{\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)} \rightarrow 0,$$

so $Z\xi_n \rightarrow Z\xi$ in the L^2 -norm on $B \times B'$. But we also have that

$$\|Z\xi_n - F\|_{L^2(B \times B')} \leq \|Z\xi_n - F\|_{L^\infty(B \times B')} \rightarrow 0$$

which means that $Z\xi = F$ on $B \times B'$. By quasiperiodicity, they must be equal on the whole of $G \times \widehat{G}$. We have thus shown that the Zak transform maps $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$ into $\mathbf{QP}_{G, \Lambda}$, and by continuity it becomes an inner product preserving $C^*(\Delta, c)$ -linear map $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G) \rightarrow \mathbf{QP}_{G, \Lambda}$.

It remains to show that Z is surjective. Since Z is an isometry, it has closed range, so it suffices to show that $Z(S_0(G))$ is dense in $\mathbf{QP}_{G, \Lambda}$. We show this in the following lemma: ■

Lemma III.5.2. *The image of $S_0(G)$ under the Zak transform is dense in $\mathbf{QP}_{G, \Lambda}$.*

Proof. By Proposition III.2.2, we can find a module frame $\{G_1, \dots, G_k\}$ for $\mathbf{QP}_{G, \Lambda}$ where $G_j = Z\xi_j$ for functions $\xi_j \in S_0(G)$, $1 \leq j \leq k$.

Now suppose $F \in \mathbf{QP}_{G, \Lambda}$. Then since $\{G_1, \dots, G_k\}$ is a module frame, it is a generating set for $\mathbf{QP}_{G, \Lambda}$ by Proposition III.2.2. Thus, there exist functions

$f_1, \dots, f_k \in C(X_{G,\Lambda})$ such that $F = \sum_j f_j \cdot G_j$. Now since $\ell^1(\Lambda \times \Lambda^\perp)$ is dense in $C^*(\Lambda \times \Lambda^\perp)$ and $\phi(C^*(\Lambda \times \Lambda^\perp)) = C(X_{G,\Lambda})$, we can find sequences $(a_{i,j})_{i=1}^\infty$ in $\ell^1(\Lambda \times \Lambda^\perp)$ for each $1 \leq j \leq k$ such that $\lim_i \phi(a_{i,j}) = f_j$ in the sup-norm. By (III.32), we then have that

$$F = \lim_i \sum_j \phi(a_{i,j}) \cdot G_j = \lim_i Z \left(\sum_j a_{i,j} \cdot \xi_j \right).$$

By Proposition III.3.2 (III.19), we have that $\sum_j a_{i,j} \cdot \xi_j$ is in $S_0(G)$. this shows that F is in the closure of $Z(S_0(G))$, which finishes the proof. ■

As a consequence, we obtain the following description of the Heisenberg module $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$ in terms of the Zak transform.

Proposition III.5.3. *The Heisenberg module $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$ consists exactly of the functions $\xi \in L^2(G)$ for which $Z_{G,\Lambda}\xi$ is continuous.*

Proof. Let $\xi \in L^2(G)$. Then $Z\xi$ is continuous if and only if $Z\xi \in \text{QP}_{G,\Lambda}$. Since the Zak transform maps $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$ bijectively onto $\text{QP}_{G,\Lambda}$ by Theorem III.5.1, it follows that $\xi \in \mathcal{E}$ if and only if $Z\xi$ is continuous. ■

We are now in position to prove one of our main results:

Theorem III.5.4. *Let G be a second countable, locally compact abelian group, and let Λ be a lattice in G . Then the following are equivalent:*

- (i) *The vector bundle $E_{G,\Lambda}$ is nontrivial.*
- (ii) *The Heisenberg module $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$ is not singly generated.*
- (iii) *Every continuous map $F: G \times \widehat{G} \rightarrow \mathbb{C}$ satisfying*

$$F(x\lambda, \omega\tau) = \overline{\omega(\lambda)}F(x, \omega)$$

for all $(x, \omega) \in G \times \widehat{G}$ and $(\lambda, \tau) \in \Lambda \times \Lambda^\perp$, must have a zero.

- (iv) *Whenever $\xi \in L^2(G)$ is such that $Z_{G,\Lambda}\xi$ is continuous, then $Z_{G,\Lambda}\xi$ has a zero.*
- (v) *Whenever $\xi \in L^2(G)$ is such that $Z_{G,\Lambda}\xi$ is continuous, then $\mathcal{G}(\xi, \Lambda \times \Lambda^\perp)$ is not a Gabor frame for $L^2(G)$.*
- (vi) *Statement III.1.2 holds for (G, Λ) . That is, whenever $\xi \in S_0(G)$, then $\mathcal{G}(\xi, \Lambda \times \Lambda^\perp)$ is not a Gabor frame for $L^2(G)$.*

Proof. By Proposition III.4.5 and Theorem III.5.1, the three modules $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$, $\text{QP}_{G,\Lambda}$ and $\Gamma(E_{G,\Lambda})$ are all isomorphic, so if one of them is singly generated, then all of them are.

By Corollary III.2.6, (i) is equivalent to the statement that $\Gamma(E_{G,\Lambda})$ is not singly generated, which is again equivalent to (ii).

By Proposition III.4.3, (iii) is equivalent to the statement that $\text{QP}_{G,\Lambda}$ is not singly generated. By Proposition III.3.3 (iii) and Proposition III.5.3, (v) is equivalent to saying that there are no single generators for $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$. Since every $F \in \text{QP}_{G,\Lambda}$ is of the form $Z\xi$ for some $\xi \in \mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$, (iii) is equivalent to (iv).

Finally, (v) and (vi) are equivalent to (ii) by Proposition III.3.5, again using the description of $\mathcal{E}_{\Lambda \times \Lambda^\perp}(G)$ in Proposition III.5.3. ■

It is already known from [KK98, Remark 6] that (iii), (iv) and (v) in Theorem III.5.4 are equivalent. The proofs we present here however, are new.

III.5.2 Determining (non)triviality of $E_{G,\Lambda}$ for some examples

Now that we have proved our main result, Theorem III.5.4, we turn to determine the triviality or nontriviality of the bundle $E_{G,\Lambda}$ for specific (G, Λ) . By Theorem III.5.4, this will then give the validity or nonvalidity of the Balian–Low theorem for $S_0(G)$ over the lattice $\Lambda \times \Lambda^\perp$ (Statement III.1.2) in these settings.

We begin with the following result, which describes a kind of functoriality for the construction $(G, \Lambda) \mapsto E_{G,\Lambda}$.

Proposition III.5.5. *Let G and H be second countable, locally compact abelian groups. Let Λ be a lattice in G and let Γ be a lattice in H . Suppose $\phi: G \rightarrow H$ is a topological group isomorphism such that $\phi(\Lambda) = \Gamma$. Then the spaces $X_{G,\Lambda}$ and $X_{H,\Gamma}$ are homeomorphic, and the vector bundles $E_{G,\Lambda}$ and $E_{H,\Gamma}$ are isomorphic.*

Proof. Define $h: X_{G,\Lambda} \rightarrow X_{H,\Gamma}$ by $h([x], [\omega]) = ([\phi(x)], [\omega \circ \phi^{-1}])$. Then it is straightforward to show that h is a continuous bijection, hence a homeomorphism because the spaces are compact Hausdorff. Furthermore, define $\psi: E_{G,\Lambda} \rightarrow E_{H,\Gamma}$ by $\psi([x, \omega, z]) = [\phi(x), \omega \circ \phi^{-1}, z]$. This map is well-defined, as for $x \in G$, $\omega \in \hat{G}$, $\lambda \in \Lambda$, $\tau \in \Lambda^\perp$ and $z \in \mathbb{C}$, we have that

$$\begin{aligned} \psi([x\lambda, \omega\tau, \overline{\omega(\lambda)}z]) &= [\phi(x\lambda), (\omega\tau) \circ \phi^{-1}, \overline{\omega(\lambda)}z] \\ &= [\phi(x)\phi(\lambda), (\omega \circ \phi^{-1})(\tau \circ \phi^{-1}), \overline{\omega(\lambda)}z] \\ &= [\phi(x), \omega \circ \phi^{-1}, \omega \circ \phi^{-1}(\phi(\lambda))\overline{\omega(\lambda)}z] \\ &= [\phi(x), \omega \circ \phi^{-1}, z] \\ &= \psi([x, \omega, z]). \end{aligned}$$

Moreover, it is continuous and restricts to an isomorphism on each fiber, so it is an isomorphism of vector bundles. ■

In the following example we look at the vector bundles associated to the groups $G = \mathbb{R}^n$ for $n \in \mathbb{N}$.

Example III.5.6. Let $G = \mathbb{R}^n$. Any lattice in $\mathbb{R}^n$ is of the form $A\mathbb{Z}^n$ for an invertible real $n \times n$ matrix A . Since multiplication by A is a topological isomorphism on $\mathbb{R}^n$ that maps $\mathbb{Z}^n$ to $A\mathbb{Z}^n$, we have by Proposition III.5.5 that

all the vector bundles $E_{\mathbb{R}^n, \Lambda}$ are isomorphic regardless of the chosen lattice Λ . We will therefore concentrate on the simplest choice $\Lambda = \mathbb{Z}^n$.

The base space of the vector bundle $E = E_{\mathbb{R}^n, \mathbb{Z}^n}$ becomes $(\mathbb{R}^n/\mathbb{Z}^n) \times (\mathbb{R}^n/\mathbb{Z}^n) \cong \mathbb{T}^{2n}$. The fact that E is a nontrivial vector bundle in this case is a consequence of the usual amalgam Balian–Low theorem in higher dimensions. The proof found in e.g. [Grö01, p. 164 Lemma 8.4.2] assumes that there is a non-vanishing continuous quasiperiodic function $F: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{C}$ and arrives at a contradiction. We will outline an alternate proof using Chern–Weil theory for vector bundles, see [MS74].

The base space $\mathbb{T}^{2n}$ has the structure of a smooth manifold, and a frame for the vector fields on $\mathbb{T}^{2n}$ is given by $\{\partial_{x_j}, \partial_{\omega_j}\}_{j=1}^n$ where we view a coordinate in $\mathbb{T}^{2n}$ as $(x_1, \dots, x_n, \omega_1, \dots, \omega_n)$.

The vector bundle E can be shown to be a smooth vector bundle over $\mathbb{T}^{2n}$. The smooth sections of E can be identified with *smooth* quasiperiodic functions $F: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{C}$.

A calculation shows that defining

$$\begin{aligned}\nabla_{\partial_{x_j}} F(x, \omega) &= \frac{\partial F}{\partial x_j}(x, \omega), \\ \nabla_{\partial_{\omega_j}} F(x, \omega) &= 2\pi i x_j F(x, \omega) + \frac{\partial F}{\partial \omega_j}(x, \omega)\end{aligned}$$

for $(x, \omega) \in G \times \widehat{G}$, and extending linearly, defines a connection ∇ on E . We have that

$$(\nabla_{\partial_{x_j}} \nabla_{\partial_{\omega_k}} - \nabla_{\partial_{\omega_k}} \nabla_{\partial_{x_j}}) F(x, \omega) = \begin{cases} 2\pi i F(x, \omega) & \text{for } j = k \\ 0 & \text{for } j \neq k \end{cases}$$

for $1 \leq j, k \leq n$ and $(x, \omega) \in G \times \widehat{G}$. Hence the curvature F^∇ associated to ∇ is given by

$$F^\nabla = 2\pi i \sum_{j=1}^n dx_j \wedge d\omega_j.$$

It follows that the first Chern class of E is given by

$$c_1(E) = \frac{i}{2\pi} F^\nabla = - \sum_{j=1}^n dx_j \wedge d\omega_j.$$

This is a nontrivial element of the second cohomology group $H^2(\mathbb{T}^{2n}, \mathbb{Z}) \cong \mathbb{Z}^{n(2n-1)}$ which is generated by all wedges of any two 1-forms $dx_j, d\omega_k$, $j, k = 1, \dots, n$. Since the line bundle has nontrivial first Chern class, it follows that the bundle is nontrivial.

Proposition III.5.7. *Let G be a second countable abelian group which is either compact or discrete, and let Λ be any lattice in G . Then the vector bundle $E_{G, \Lambda}$ is trivial.*

III. The BLT for LCA groups and vector bundles

Proof. Suppose that G is compact. Then Λ is a compact and discrete subgroup of G , so it must be finite, say of order r . Since finite abelian groups are self-dual, $\widehat{G}/\Lambda^\perp \cong \widehat{\Lambda}$ must be finite of order r as well. Thus, the base space $X_{G,\Lambda}$ of the vector bundle $E = E_{G,\Lambda}$ is homeomorphic to a disjoint union of r copies of G/Λ :

$$X_{G,\Lambda} = (G/\Lambda) \times (\widehat{G}/\Lambda^\perp) = \coprod_{k=1}^r (G/\Lambda) \times \{\omega_k\}.$$

Here, $\omega_1, \dots, \omega_r$ are coset representatives for $\Lambda^\perp$ in $\widehat{G}$. Now, the vector bundle E is trivial if the restrictions of E to each component $(G/\Lambda) \times \{\omega_k\}$ is trivial. The restrictions are of the form

$$E_k := \pi^{-1}((G/\Lambda) \times \{\omega_k\}) = \{[x, \omega_k, z] : x \in G, z \in \mathbb{C}\}.$$

Define a map $\phi: E_k \rightarrow (G/\Lambda) \times \mathbb{C}$ from E_k to the trivial bundle over G/Λ of rank 1 by

$$\phi([x, \omega_k, z]) = ([x], \omega_k(x)z).$$

This map is well-defined, as it maps $[x\lambda, \omega_k\tau, \overline{\omega_k(\lambda)}z]$ to $([x\lambda], \omega_k(x\lambda)\overline{\omega_k(\lambda)}z) = ([x], \omega_k(x)z)$ for $(x, \omega) \in G \times \widehat{G}$, $(\lambda, \tau) \in \Lambda \times \Lambda^\perp$ and $z \in \mathbb{C}$. It is clear that it is continuous and restricts to a linear isomorphism on each fiber, hence it is an isomorphism of vector bundles. This proves the assertion that E is trivial.

If G is discrete, then G/Λ is both compact and discrete, hence finite. The argument above applies, with the roles of (G, Λ) and $(\widehat{G}, \Lambda^\perp)$ interchanged. ■

At this point, we would like to point out that even though one does not have a Balian–Low theorem for the Feichtinger algebra in compact or discrete groups, this does not mean there does not exist Balian–Low phenomena at all in these groups. In fact, it is shown in [NO19] that there exists a Balian–Low theorem in finite cyclic groups.

The following result follows directly from combining the main result of [KK98] with Theorem III.5.4.

Proposition III.5.8. *Let G be a second countable, locally compact abelian group that is compactly generated and has noncompact connected component of the identity. Then for any lattice Λ in G , the vector bundle $E_{G,\Lambda}$ is nontrivial.*

III.6 The group $\mathbb{R} \times \mathbb{Q}_p$

In this section, we will show that when $G = \mathbb{R} \times \mathbb{Q}_p$ and Λ is a certain lattice in G , then the associated bundle $E_{G,\Lambda}$ is nontrivial. By Theorem III.5.4, this will prove Statement III.1.2 for this setting, thereby giving us a Balian–Low theorem in this new setting.

Recall that the locally compact field $\mathbb{Q}_p$ of p -adic numbers is the completion of $\mathbb{Q}$ with respect to the absolute value $|x|_p = p^{-k}$ where $x = p^k(a/b)$ for $a, b, k \in \mathbb{Z}$,

$p \nmid a$, $p \nmid b$ and $|0|_p = 0$. Every p -adic number x can be expressed uniquely as a converging series

$$x = \sum_{k=K}^{\infty} x_k p^k$$

where K is a (possibly negative) integer and $x_k \in \{0, \dots, p-1\}$ for every integer $k \geq K$. The ring of p -adic integers $\mathbb{Z}_p$ is the compact subgroup of p -adic numbers x for which $|x|_p \leq 1$, or, equivalently, the series of the form $\sum_{k=0}^{\infty} x_k p^k$.

The group $\mathbb{Q}_p$ is self-dual via the pairing $\mathbb{Q}_p \times \mathbb{Q}_p \rightarrow \mathbb{T}$, $(x, y) \mapsto e^{2\pi i \{xy\}_p}$, where

$$\left\{ \sum_{k=K}^{\infty} x_k p^k \right\}_p = \sum_{k=K}^{-1} x_k p^k.$$

We identify the group $\widehat{\mathbb{R} \times \mathbb{Q}_p}$ with itself via $((s, x), (t, y)) \mapsto e^{2\pi i s t} e^{-2\pi i \{xy\}_p}$ (note the sign). A remarkable fact is that

$$\mathbb{Z}[1/p] = \{a/p^k : a, k \in \mathbb{Z}\}$$

can be embedded diagonally into $\mathbb{R} \times \mathbb{Q}_p$ as a lattice. In other words, the set $\mathbb{Z}[1/p]^{\Delta} = \{(q, q) : q \in \mathbb{Z}[1/p]\}$ is a lattice in $\mathbb{R} \times \mathbb{Q}_p$ (analogous to [RV99, Theorem 5.11]). Under the identification of the Pontryagin dual of $\mathbb{R} \times \mathbb{Q}_p$ above, the annihilator of $\mathbb{Z}[1/p]^{\Delta}$ is identified with $\mathbb{Z}[1/p]$. Thus we have that

$$\widehat{\mathbb{Z}[1/p]} \cong \widehat{\mathbb{Z}[1/p]^{\Delta}} \cong \frac{\mathbb{R} \times \mathbb{Q}_p}{(\mathbb{Z}[1/p]^{\Delta})^{\perp}} \cong \frac{\mathbb{R} \times \mathbb{Q}_p}{\mathbb{Z}[1/p]^{\Delta}}. \quad (\text{III.34})$$

Now $\mathbb{Z}[1/p]$ can be realized as the direct limit of the sequence

$$\mathbb{Z} \xrightarrow{p} \mathbb{Z} \xrightarrow{p} \mathbb{Z} \xrightarrow{p} \dots$$

where each map is multiplication by p . We index this sequence by the natural numbers including zero. The injection of the k -th copy of $\mathbb{Z}$ into $\mathbb{Z}[1/p]$ is given by $a \mapsto a/p^k$. Taking Pontryagin duals, we obtain that $\widehat{\mathbb{Z}[1/p]}$ is isomorphic to the inverse limit of the sequence

$$\dots \xrightarrow{\widehat{p}} \widehat{\mathbb{Z}} \xrightarrow{\widehat{p}} \widehat{\mathbb{Z}} \xrightarrow{\widehat{p}} \widehat{\mathbb{Z}}.$$

Using the usual identification $\widehat{\mathbb{Z}} \cong \mathbb{R}/\mathbb{Z}$, we obtain that $\widehat{\mathbb{Z}[1/p]}$ is isomorphic to the inverse limit of

$$\dots \xrightarrow{p} \mathbb{R}/\mathbb{Z} \xrightarrow{p} \mathbb{R}/\mathbb{Z} \xrightarrow{p} \mathbb{R}/\mathbb{Z}$$

where each map is multiplication by p . This space is known as the p -solenoid and is denoted by $\mathcal{S}_p$. It is an example of a compact Hausdorff space that is connected, but not path-connected.

Denote by $\pi_k: \widehat{\mathbb{Z}[1/p]} \rightarrow \mathbb{R}/\mathbb{Z}$ the projection down to the k -th copy of $\mathbb{R}/\mathbb{Z}$. It is given by mapping a character $\omega \in \widehat{\mathbb{Z}[1/p]}$ to the class $[t] \in \mathbb{R}/\mathbb{Z}$ such that $\omega(a/p^k) = e^{2\pi i a t}$ for all $a, k \in \mathbb{Z}$.

III. The BLT for LCA groups and vector bundles

Now using (III.34), ω is obtained from a class $[s, x] \in (\mathbb{R} \times \mathbb{Q}_p)/\mathbb{Z}[1/p]^\Delta$, i.e. it is given by $\omega(q) = e^{2\pi i q s} e^{-2\pi i \{q x\}_p}$ for $q \in \mathbb{Z}[1/p]$. Letting $\pi_k(\omega) = [t]$, we have that

$$e^{2\pi i a t} = \omega(a/p^k) = e^{2\pi i a s/p^k} e^{-2\pi i \{a x/p^k\}_p}$$

for all $a, k \in \mathbb{Z}$, so that $[t] = [p^{-k}s - \{p^{-k}x\}_p]$. We thus have the following proposition:

Proposition III.6.1. *The p -solenoid $\mathcal{S}_p$ given by the inverse limit of the sequence*

$$\dots \xrightarrow{p} \mathbb{R}/\mathbb{Z} \xrightarrow{p} \mathbb{R}/\mathbb{Z} \xrightarrow{p} \mathbb{R}/\mathbb{Z}$$

is isomorphic to the quotient group $(\mathbb{R} \times \mathbb{Q}_p)/\mathbb{Z}[1/p]^\Delta$. The projection π_k from $(\mathbb{R} \times \mathbb{Q}_p)/\mathbb{Z}[1/p]^\Delta$ down to the k -th factor of $\mathbb{R}/\mathbb{Z}$ is given by

$$\pi_k([s, x]) = [p^{-k}s - \{p^{-k}x\}_p]$$

for $(s, x) \in \mathbb{R} \times \mathbb{Q}_p$.

We now investigate the line bundle $\tilde{E} = E_{G, \Lambda}$ in the case of $G = \mathbb{R} \times \mathbb{Q}_p$ and $\Lambda = \mathbb{Z}[1/p]^\Delta$. The base space becomes

$$X_{G, \Lambda} = \frac{\mathbb{R} \times \mathbb{Q}_p}{\mathbb{Z}[1/p]^\Delta} \times \frac{\widehat{\mathbb{R} \times \mathbb{Q}_p}}{(\mathbb{Z}[1/p]^\Delta)^\perp} \cong \left(\frac{\mathbb{R} \times \mathbb{Q}_p}{\mathbb{Z}[1/p]^\Delta} \right)^2$$

which is homeomorphic to $\mathcal{S}_p^2$ by Proposition III.6.1. Thus, the vector bundle $\tilde{E}$ is a line bundle over $\mathcal{S}_p^2$, and by Proposition III.6.1, $\mathcal{S}_p^2$ can be realized as the inverse limit of the sequence

$$\dots \rightarrow (\mathbb{R}/\mathbb{Z})^2 \xrightarrow{(\cdot, p)} (\mathbb{R}/\mathbb{Z})^2 \xrightarrow{(\cdot, p)} (\mathbb{R}/\mathbb{Z})^2.$$

Here $(\cdot, p)([s], [t]) = ([ps], [pt])$ for $([s], [t]) \in (\mathbb{R}/\mathbb{Z})^2$, and the projection $\pi_k^2: \mathcal{S}_p^2 \rightarrow (\mathbb{R}/\mathbb{Z})^2$ given by

$$\pi_k^2([s, x], [t, y]) = ([p^{-k}s - \{p^{-k}x\}_p], [p^{-k}t - \{p^{-k}y\}_p]) \quad (\text{III.35})$$

for $([s, x], [t, y]) \in \mathcal{S}_p^2$.

Key to understanding vector bundles over inverse limits of spaces is the following result. It can be seen by combining [Bla98, Example 5.2.4] with the correspondence between vector bundles and finitely generated projective modules, as well as the continuity of the functor $C(\cdot)$ from compact Hausdorff spaces to unital C^* -algebras:

Proposition III.6.2. *Suppose X is the inverse limit of the sequence of compact Hausdorff spaces*

$$\dots \xrightarrow{f_3} X_3 \xrightarrow{f_2} X_2 \xrightarrow{f_1} X_1.$$

Then the monoid $\text{Vect}(X)$ of isomorphism classes of vector bundles over X (under direct sum) is isomorphic to the direct limit of the monoids

$$\text{Vect}(X_1) \xrightarrow{f_1^*} \text{Vect}(X_2) \xrightarrow{f_2^*} \text{Vect}(X_3) \xrightarrow{f_3^*} \dots,$$

where the induced maps are given by pullback of vector bundles.

The identity in $\text{Vect}(X)$ is the class of the unique trivial bundle of rank 0. Removing this class from $\text{Vect}(X)$, i.e. considering only bundles of positive rank, we obtain a semigroup that we will denote by $\text{Vect}^+(X)$. The functor Vect^+ preserves inverse limits in the same sense that Vect does in Proposition III.6.2.

Denote by $\mathbb{Z}^+$ the set of positive integers. By [Rie83, Theorem 3.9] and [PR04, Proposition 4.1], the semigroup $\text{Vect}^+(\mathbb{T}^2)$ is isomorphic to $\mathbb{Z}^+ \times \mathbb{Z}$, where (q, a) represents the vector bundle $E_{q,a}$ over $\mathbb{T}^2$ of rank q and Chern class $-a \in H^2(\mathbb{T}^2, \mathbb{Z}) \cong \mathbb{Z}$. The vector bundle $E = E_{\mathbb{R}, \mathbb{Z}}$ corresponds to $(q, a) = (1, -1)$ by Example III.5.6. It is easily shown that $(\cdot p, \cdot p)^*(E_{q,a}) \cong E_{q, p^2 a}$. Hence, the induced map of $(\cdot p, \cdot p): (\mathbb{R}/\mathbb{Z})^2 \rightarrow (\mathbb{R}/\mathbb{Z})^2$ on the level of vector bundles is given by $(q, a) \mapsto (q, p^2 a)$. This gives us that $\text{Vect}^+(\mathcal{S}_p^2)$ is the direct limit of the sequence of monoids

$$\mathbb{Z}^+ \times \mathbb{Z} \xrightarrow{(\cdot 1, \cdot p^2)} \mathbb{Z}^+ \times \mathbb{Z} \xrightarrow{(\cdot 1, \cdot p^2)} \mathbb{Z}^+ \times \mathbb{Z} \xrightarrow{(\cdot 1, \cdot p^2)} \dots$$

In other words, $\text{Vect}^+(\mathcal{S}_p^2) \cong \mathbb{Z}^+ \times \mathbb{Z}[1/p]$. It follows that we obtain the following classification of vector bundles over $\mathcal{S}_p^2$:

Proposition III.6.3. *Let $E_{q,a}$ denote the vector bundle of rank q and Chern class $-a$ over $\mathbb{T}^2$. Then every vector bundle over $\mathcal{S}_p^2$ of positive rank is of the form $(\pi_k^2)^*(E_{q,a})$ for some $k, q \in \mathbb{Z}^+$ and $a \in \mathbb{Z}$, where π_k^2 is the map in (III.35). Moreover, $(\pi_k^2)^*(E_{q,a}) \cong (\pi_{k'}^2)^*(E_{q',a'})$ if and only if $q = q'$ and $a/p^{2k} = a'/p^{2k'}$. Thus, $(\pi_k^2)^*(E_{q,a})$ is trivial if and only if $a = 0$.*

The next proposition, namely Proposition III.6.4, determines the bundle $E_{\mathbb{R} \times \mathbb{Q}_p, \mathbb{Z}[1/p]^\Delta} \rightarrow \mathcal{S}_p^2$ among the bundles in Proposition III.6.3. Before the proof, we present the functoriality of the section functor.

To begin with, let A and B be unital C^* -algebras and let $\phi: A \rightarrow B$ be a $*$ -homomorphism. If $\mathcal{E}$ is a left finitely generated projective A -module, there is a way to construct a left finitely generated projective B -module $\phi_*(\mathcal{E})$ known as *extension of scalars* [GVF01, p. 60]. In terms of projection matrices, extension of scalars can be described as follows: If $\mathcal{E}$ is represented by the projection $P = (p_{i,j})_{i,j=1}^k \in M_k(A)$, i.e. $\mathcal{E} \cong A^k P$, then we have that

$$\phi_*(A^k P) \cong B^k \phi_*(P). \quad (\text{III.36})$$

Here, $\phi_*(P)$ is the matrix $(\phi(p_{i,j}))_{i,j=1}^k \in M_k(B)$.

Now suppose we are in the following situation: We have a continuous map $f: X \rightarrow Y$ of compact Hausdorff spaces. We then get an induced $*$ -homomorphism $C(f): C(Y) \rightarrow C(X)$ given by precomposition by f . We can consider the extension of scalars along this $*$ -homomorphism. Let $E \rightarrow Y$ be a vector bundle over Y . Then we can form the pullback $f^*(E)$ of E along f , which is a vector bundle over X . Now $\Gamma(E)$ is a left finitely generated projective $C(Y)$ -module, while $\Gamma(f^*(E))$ is a finitely generated projective $C(X)$ -module. The section functor is functorial in the sense that if one performs extension of scalars on $\Gamma(E)$ using the induced map $C(f)$, then the resulting $C(Y)$ -module is isomorphic to $\Gamma(f^*(E))$. That is, we have that

$$C(f)_*(\Gamma(E)) \cong \Gamma(f^*(E)). \quad (\text{III.37})$$

For a proof of (III.37), see [GVF01, Proposition 2.12].

The proof of the following proposition will employ fundamental domains, which are introduced in Section III.A.

Proposition III.6.4. *Let E denote the line bundle $E_{\mathbb{R},\mathbb{Z}}$ over $(\mathbb{R}/\mathbb{Z})^2$ and let $\tilde{E}$ denote the line bundle $E_{\mathbb{R} \times \mathbb{Q}_p, \mathbb{Z}[1/p]^\Delta}$ over $\mathcal{S}_p^2$, both defined as in Section III.4.2. Then $\tilde{E}$ is isomorphic to the pullback of E along $\pi_0^2: \mathcal{S}_p^2 \rightarrow (\mathbb{R}/\mathbb{Z})^2$, where π_0^2 is given as in (III.35).*

Proof. Our proof goes as follows: Instead of showing directly that $\tilde{E} \cong (\pi_0^2)^*(E)$, we can instead show (by Proposition III.2.4) that $\Gamma(\tilde{E}) \cong \Gamma((\pi_0^2)^*(E))$ as $C(\mathcal{S}_p^2)$ -modules.

We consider the continuous map $\pi_0^2: \mathcal{S}_p^2 \rightarrow (\mathbb{R}/\mathbb{Z})^2$, which induces the map

$$C(\pi_0^2): C((\mathbb{R}/\mathbb{Z})^2) \rightarrow C(\mathcal{S}_p^2)$$

given by precomposition by π_0^2 .

We have that $\Gamma(\tilde{E}) \cong \text{QP}_{\tilde{G}, \tilde{\Lambda}}$ by Proposition III.4.5. We also have that

$$\begin{aligned} \Gamma((\pi_0^2)^*(E)) &\cong C(\pi_0^2)_*(\Gamma(E)) && \text{by (III.37)} \\ &\cong C(\pi_0^2)_*(\text{QP}_{\mathbb{R},\mathbb{Z}}) && \text{by Proposition III.4.5.} \end{aligned}$$

The goal is thus to show that $\text{QP}_{\tilde{G}, \tilde{\Lambda}}$ and $C(\pi_0^2)_*(\text{QP}_{\mathbb{R},\mathbb{Z}})$ are isomorphic as $C(\mathcal{S}_p^2)$ -modules. To show this, it suffices to show that they can be represented by the same projection matrix over $C(\mathcal{S}_p^2)$, i.e. that $\text{QP}_{\tilde{G}, \tilde{\Lambda}} \cong A^k P \cong C(\pi_0^2)_*(\text{QP}_{\mathbb{R},\mathbb{Z}})$ for some $k \in \mathbb{N}$ and $P \in M_k(C(\mathcal{S}_p^2))$. This will be our approach.

We begin by constructing normalized tight module frames for the modules $\text{QP}_{\tilde{G}, \tilde{\Lambda}}$ and $C(\pi_0^2)_*(\text{QP}_{\mathbb{R},\mathbb{Z}})$, using fundamental domains from Section III.A.

Set $\tilde{G} = \mathbb{R} \times \mathbb{Q}_p$ and $\tilde{\Lambda} = \mathbb{Z}[1/p]^\Delta$. A fundamental domain for $\tilde{\Lambda}$ in $\tilde{G}$ is given by $\tilde{B} = [0, 1) \times \mathbb{Z}_p$. To see this, note that if $(s, x) \in \mathbb{R} \times \mathbb{Q}_p$, then

$$(s, x) = (\{x\}_p + \lfloor s - \{x\}_p \rfloor, \{x\}_p + \lfloor s - \{x\}_p \rfloor) + (\text{fr}(s - \{x\}_p), x - \{x\}_p - \lfloor s - \{x\}_p \rfloor) \quad (\text{III.38})$$

where $\lfloor \cdot \rfloor$ denotes the floor function $\mathbb{R} \rightarrow \mathbb{Z}$ and fr denotes the fractional part of a real number, i.e. $\text{fr}(s) = s - \lfloor s \rfloor$. The first pair in (III.38) is in $\mathbb{Z}[1/p]^\Delta$ while the second pair is in $[0, 1) \times \mathbb{Z}_p$. Thus, the generalized floor function $\lfloor \cdot \rfloor_{\tilde{B}}: \mathbb{R} \times \mathbb{Q}_p \rightarrow \mathbb{Z}[1/p]$ from (III.41) associated to the fundamental domain $\tilde{B} = [0, 1) \times \mathbb{Z}_p$ is given by

$$\lfloor (s, x) \rfloor_{\tilde{B}} = \{x\}_p + \lfloor s - \{x\}_p \rfloor \quad (\text{III.39})$$

for $(s, x) \in \mathbb{R} \times \mathbb{Q}_p$. Now since $\mathbb{Z}_p$ is clopen in $\mathbb{Q}_p$ we have that

$$\partial([0, 1) \times \mathbb{Z}_p) = \partial([0, 1)) \times \overline{\mathbb{Z}_p} \cup \overline{[0, 1)} \times \partial(\mathbb{Z}_p) = \{0, 1\} \times \mathbb{Z}_p. \quad (\text{III.40})$$

By Proposition III.A.2, $\lfloor \cdot \rfloor_{\tilde{B}}$ is discontinuous at (s, x) if and only if $(s, x) \in \partial([0, 1) \times \mathbb{Z}_p) = \{0, 1\} \times \mathbb{Z}_p$. From (III.39), this happens if and

only if $s - \{x\}_p \in \mathbb{Z}$.

By Example III.A.4, we can find a continuous, Λ -periodic function $f: \mathbb{R} \rightarrow \mathbb{C}$ with $f(0) = 0$ and

$$|f(s)|^2 + |f(s + 1/2)|^2 = 1$$

for all $s \in \mathbb{R}$. Consequently, the set $\{G_1, G_2\}$ where

$$\begin{aligned} G_1(s, t) &= f(s)J_{[0,1)}(s, t) = f(s)e^{-2\pi i \lfloor s \rfloor t}, \\ G_2(s, t) &= f(s + 1/2)J_{[0,1)}(s + 1/2, t) = e^{-2\pi i \lfloor s + 1/2 \rfloor t}, \end{aligned}$$

is a normalized tight module frame for $\mathcal{Q}\mathcal{P}_{\mathbb{R}, \mathbb{Z}}$ as in Proposition III.A.3. By Proposition III.2.3, the matrix $P \in M_2(C(\mathbb{T}^2))$ given by

$$P([s], [t]) = \begin{pmatrix} |G_1(s, t)|^2 & G_1(s, t)\overline{G_2(s, t)} \\ G_2(s, t)\overline{G_1(s, t)} & |G_2(s, t)|^2 \end{pmatrix}$$

represents $\mathcal{Q}\mathcal{P}_{\mathbb{R}, \mathbb{Z}}$, i.e. $\mathcal{Q}\mathcal{P}_{\mathbb{R}, \mathbb{Z}} \cong C(\mathbb{T}^2)^2 P$. Now it follows from (III.36) that the module $(\pi_0^2)_*(\mathcal{Q}\mathcal{P}_{\mathbb{R}, \mathbb{Z}}) = (\pi_0^2)_*(C(\mathbb{T}^2)^2 P)$ is represented by the projection matrix $Q = (\pi_0^2)_*(P) = P \circ \pi_0^2$, which is given by $Q = (Q_{i,j})_{i,j=1}^2$ where

$$\begin{aligned} Q_{1,1}([s, x], [t, y]) &= |G_1(s - \{x\}_p, t - \{y\}_p)|^2, \\ Q_{1,2}([s, x], [t, y]) &= G_1(s - \{x\}_p, t - \{y\}_p)\overline{G_2(s - \{x\}_p, t - \{y\}_p)}, \\ Q_{2,1}([s, x], [t, y]) &= G_2(s - \{x\}_p, t - \{y\}_p)\overline{G_1(s - \{x\}_p, t - \{y\}_p)}, \\ Q_{2,2}([s, x], [t, y]) &= |G_2(s - \{x\}_p, t - \{y\}_p)|^2. \end{aligned}$$

Next, note that with $x_1 = (0, 0)$ and $x_2 = (-1/2, 0)$, the sets $x_1 + \tilde{B}^\circ + \tilde{\Lambda} = (0, 1) \times \mathbb{Z}_p + \mathbb{Z}[1/p]^\Delta$ and $x_2 + \tilde{B}^\circ + \tilde{\Lambda} = (-1/2, 1/2) \times \mathbb{Z}_p + \mathbb{Z}[1/p]^\Delta$ form an open cover of $\mathbb{R} \times \mathbb{Q}_p$ as in Proposition III.A.3. Now define $\tilde{f}: \mathbb{R} \times \mathbb{Q}_p \rightarrow \mathbb{C}$ by $\tilde{f}(s, x) = f(s - \{x\}_p)$. Then $\tilde{f}$ is a continuous, $\mathbb{Z}[1/p]^\Delta$ -periodic function and

$$|\tilde{f}(s, x)|^2 + |\tilde{f}(s + 1/2, x)|^2 = |f(s - \{x\}_p)|^2 + |f(s - \{x\}_p + 1/2)|^2 = 1$$

for all $(s, x) \in \mathbb{R} \times \mathbb{Q}_p$. Moreover, if $(k, x) \in \partial([0, 1) \times \mathbb{Z}_p) = \{0, 1\} \times \mathbb{Z}_p$ then

$$\tilde{f}(k, x) = f(k - \{x\}_p) = f(k) = 0,$$

since f is zero on integers. This shows that $\tilde{f}$ satisfies the requirements in Proposition III.A.3 with respect to the fundamental domain $\tilde{B}$ and x_1, x_2 . Consequently, with

$$\begin{aligned} \tilde{G}_1(s, x, t, y) &= \tilde{f}(s, x)J_{\tilde{B}}(s, x, t, y) \\ \tilde{G}_2(s, x, t, y) &= \tilde{f}(s + 1/2, x)J_{\tilde{B}}(s + 1/2, x, t, y), \end{aligned}$$

III. The BLT for LCA groups and vector bundles

we have that $\{\tilde{G}_1, \tilde{G}_2\}$ is a normalized tight frame for $\text{QP}_{\tilde{G}, \tilde{\Lambda}}$. By Proposition III.2.3, it follows that the matrix $\tilde{P} = (\tilde{P}_{i,j})_{i,j} \in M_2(C(\mathcal{S}_p^2))$ given by

$$\tilde{P}([s, x], [t, y]) = \begin{pmatrix} |\tilde{G}_1(s, x, t, y)|^2 & \tilde{G}_1(s, x, t, y) \tilde{G}_2(s, x, t, y) \\ \tilde{G}_2(s, x, t, y) \tilde{G}_1(s, x, t, y) & |\tilde{G}_2(s, x, t, y)|^2 \end{pmatrix}$$

represents the module $\text{QP}_{\tilde{G}, \tilde{\Lambda}}$, i.e. $\text{QP}_{\tilde{G}, \tilde{\Lambda}} \cong C(\mathcal{S}_p^2)^2 \tilde{P}$. We now compute $\tilde{P}$ and show that it is equal to the matrix Q . First, note that

$$|\tilde{G}_1(s, x, t, y)|^2 = |\tilde{f}(s, x)|^2 = |f(s - \{x\}_p)|^2 = |G_1(s - \{x\}_p, t - \{y\}_p)|^2$$

and similarly, $|\tilde{G}_2(s, x, t, y)|^2 = |G_2(s - \{x\}_p, t - \{y\}_p)|^2$. Thus, $\tilde{P}_{1,1} = Q_{1,1}$ and $\tilde{P}_{2,2} = Q_{2,2}$. Next, note that using (III.39), $J_{\tilde{B}}$ is given by

$$J_{\tilde{B}}(s, x, t, y) = e^{-2\pi i(\{x\}_p + \lfloor s - \{x\}_p \rfloor)t} e^{2\pi i(\{x\}_p + \lfloor s - \{x\}_p \rfloor)y}_p$$

for $(s, x), (t, y) \in \mathbb{R} \times \mathbb{Q}_p$. Thus, calculating $J_{\tilde{B}}(s, x, t, y) \overline{J_{\tilde{B}}(s + 1/2, x, t, y)}$, the exponentials involving $\{x\}_p$ cancel and we are left with

$$\begin{aligned} J_{\tilde{B}}(s, x, t, y) \overline{J_{\tilde{B}}(s + 1/2, x, t, y)} &= e^{-2\pi i(\lfloor s - \{x\}_p \rfloor - \lfloor s + 1/2 - \{x\}_p \rfloor)t} \\ &\quad \times e^{2\pi i(\lfloor s - \{x\}_p \rfloor - \lfloor s + 1/2 - \{x\}_p \rfloor)y}_p \\ &= e^{-2\pi i(\lfloor s + \{x\}_p \rfloor - \lfloor s + 1/2 - \{x\}_p \rfloor)(t - \{y\}_p)}. \end{aligned}$$

It follows that

$$\begin{aligned} \tilde{P}_{1,2}([s, x], [t, y]) &= \tilde{G}_1(s, x, t, y) \overline{\tilde{G}_2(s, x, t, y)} \\ &= \tilde{f}(s, x) \overline{\tilde{f}(s + 1/2, x)} J_{\tilde{B}}(s, x, t, y) \overline{J_{\tilde{B}}(s + 1/2, x, t, y)} \\ &= f(s - \{x\}_p) \overline{f(s + 1/2 - \{x\}_p)} e^{-2\pi i(\lfloor s + \{x\}_p \rfloor - \lfloor s + 1/2 - \{x\}_p \rfloor)(t - \{y\}_p)} \\ &= f(\pi_0(s, x)) \overline{f(\pi_0(s + 1/2, x))} J_B(s - \{x\}_p, t - \{y\}_p) \\ &\quad \times \overline{J_B(s + 1/2 - \{x\}_p, t - \{y\}_p)} \\ &= G_1(s - \{x\}_p, t - \{y\}_p) \overline{G_2(s - \{x\}_p, t - \{y\}_p)} \\ &= Q_{1,2}([s, x], [t, y]). \end{aligned}$$

This implies that

$$\tilde{P}_{2,1} = \overline{\tilde{P}_{1,2}} = \overline{Q_{1,2}} = Q_{2,1}.$$

We have now shown that $\tilde{P}_{i,j} = Q_{i,j}$ for all $i, j = 1, 2$. Thus $Q = \tilde{P}$, which finishes the proof. $\blacksquare$

With Proposition III.6.4 proved, our final main result is a straightforward corollary:

Theorem III.6.5. *Let $G = \mathbb{R} \times \mathbb{Q}_p$ and let $\Lambda = \mathbb{Z}[1/p]^\Delta$. Then the line bundle $\tilde{E} = E_{G, \Lambda}$ over $\mathcal{S}_p^2$ is nontrivial. Hence Statement III.1.2 holds for (G, Λ) , that is, if $\eta \in S_0(\mathbb{R} \times \mathbb{Q}_p)$, then the Gabor system*

$$\mathcal{G}(\eta, \Lambda \times \Lambda^\perp) = \{(s, x) \mapsto e^{2\pi i r s} e^{-2\pi i \{r x\}_p} \eta(s - q, x - q) : q, r \in \mathbb{Z}[1/p]\}$$

is not a frame for $L^2(\mathbb{R} \times \mathbb{Q}_p)$.

Proof. By Proposition III.6.4, $\tilde{E} \cong \pi_0^*(E_{1,1})$. By Proposition III.6.3, $\tilde{E}$ is then nontrivial, since otherwise it would be of the form $(\pi_0^2)^*(E_{q,0})$ for some $q \in \mathbb{N}$. By Theorem III.5.4, the nontriviality of $\tilde{E}$ implies Statement III.1.2 for (G, Λ) . ■

Appendix III.A Fundamental domains

In this appendix, we collect some results on fundamental domains and the construction of specific module frames for $\mathcal{Q}\mathcal{P}_{G,\Lambda}$. They are used in the proof of Proposition III.6.4 and in the proof of Proposition III.B.3.

Let Λ be a lattice in a second countable LCA group G . A measurable set $B \subseteq G$ is called a *fundamental domain* (or *Borel section*) [Mac52] for Λ in G if the collection $\{\lambda B : \lambda \in \Lambda\}$ forms a partition of G . Equivalently, every $x \in G$ can be written uniquely as $x = b\lambda$ where $b \in B$ and $\lambda \in \Lambda$.

Proposition III.A.1. *Let Λ be a lattice in a second countable, locally compact abelian group G . Then there exists a relatively compact fundamental domain B for Λ in G with nonempty interior.*

Proof. Let μ denote a Haar measure on G . Since G is both locally compact and second countable, it follows that G is σ -compact. By [Mar91, p. 35, paragraph (0.40) 1)], there exists a relatively compact fundamental domain B for Λ in G with $\mu(\partial B) = 0$.

We claim that B cannot be nowhere dense. Suppose otherwise, i.e. $(\overline{B})^\circ = \emptyset$. Then $(\overline{\lambda B})^\circ = \lambda(\overline{B})^\circ = \emptyset$ for every $\lambda \in \Lambda$. Since Λ is discrete in G and G is second countable, Λ has to be countable. The identity

$$G = \bigcup_{\lambda \in \Lambda} \lambda B$$

then expresses G , a locally compact Hausdorff space, as a countable union of nowhere dense subsets. This contradicts the Baire category theorem, so we must conclude that $(\overline{B})^\circ \neq \emptyset$.

Let U be a nonempty open subset of $\overline{B}$. Now suppose for a contradiction that $B^\circ = \emptyset$. Then $\overline{B} = B^\circ \cup \partial B = \partial B$, so $\mu(U) \leq \mu(\overline{B}) = \mu(\partial B) = 0$, which contradicts the properties of Haar measure. Hence $B^\circ \neq \emptyset$. ■

Let B be a fundamental domain for the lattice Λ in G . We define the function $\lfloor \cdot \rfloor_B : G \rightarrow \Lambda$ by letting $\lfloor x \rfloor_B$ be the unique $\lambda \in \Lambda$ such that $x = b\lambda$ for some (unique) $b \in B$. We think of this as a generalized floor function.

Note that we have

$$\lfloor x\lambda \rfloor_B = \lfloor x \rfloor_B \lambda \tag{III.41}$$

for all $x \in G$ and $\lambda \in \Lambda$.

Proposition III.A.2. *The function $\lfloor \cdot \rfloor_B$ is locally constant on $B^\circ \Lambda$ and discontinuous on $(B^\circ \Lambda)^c = (\partial B)\Lambda$.*

Proof. We begin by showing that $\lfloor \cdot \rfloor_B$ is locally constant on $B^\circ \Lambda$. Suppose $x \in B^\circ \Lambda$. Then $x = b\lambda$ for $b \in B$, $\lambda \in \Lambda$, and there exists an open set $U \subseteq B$ with $b \in U$. Then $x \in \lambda U$. If $y \in \lambda U$, say $y = b'\lambda$ with $b' \in U$, then $\lfloor y \rfloor_B = \lambda = \lfloor x \rfloor_B$, which proves that $\lfloor \cdot \rfloor_B$ is constant on the set λU .

Next, we show that if $\lfloor \cdot \rfloor_B$ is continuous at $x \in G$, then $x \in B^\circ \Lambda$. Suppose $\lfloor \cdot \rfloor_B$ is continuous at $x = b\lambda$ for $b \in B$, $\lambda \in \Lambda$. Let V be a neighbourhood of λ such that $V \cap \Lambda = \{\lambda\}$. By continuity, there exists a neighbourhood U of x for which $\lfloor \cdot \rfloor_B(U) \subseteq V$. But since $\lfloor \cdot \rfloor_B$ maps into Λ , we have that $\lfloor y \rfloor_B = \lambda$ for all $y \in U$. Hence $\lfloor z \rfloor_B = 1$ for all $z \in \lambda^{-1}U$, i.e. $\lambda^{-1}U \subseteq B$. Since $b \in \lambda^{-1}U$ we then have that $b \in B^\circ$. Thus $x \in B^\circ \Lambda$. ■

We describe how to obtain a specific module frame for $\text{QP}_{G,\Lambda}$ from certain fundamental domains. This is inspired by [PR04, p. 457]. Given a fundamental domain B for Λ in G , define the function $J_B: G \times \widehat{G} \rightarrow \mathbb{C}$ by

$$J_B(x, \omega) = \overline{\omega(\lfloor x \rfloor_B)}. \quad (\text{III.42})$$

Note that for $(x, \omega) \in G \times \widehat{G}$ and $(\lambda, \tau) \in \Lambda \times \Lambda^\perp$, we have that

$$J_B(x\lambda, \omega\tau) = \overline{(\omega\tau)(\lfloor x\lambda \rfloor_B)} = \overline{\omega(\lfloor x \rfloor_B \lambda)} = \overline{\omega(\lambda)} J_B(x, \omega).$$

Hence J_B satisfies the quasiperiodicity relation (III.23). However, it is not necessarily continuous everywhere on $G \times \widehat{G}$, so it is not an element of $\text{QP}_{G,\Lambda}$ in general. But by Proposition III.A.2, J_B is continuous on $B^\circ \Lambda \times \widehat{G}$. Thus, if $f: G \rightarrow \mathbb{C}$ is a continuous, Λ -periodic function with $f|_{\partial B} = 0$, then the function $G: G \times \widehat{G} \rightarrow \mathbb{C}$ given by

$$G(x, \omega) = f(x) J_B(x, \omega) \quad (\text{III.43})$$

for $(x, \omega) \in G \times \widehat{G}$ is continuous and quasiperiodic, hence an element of $\text{QP}_{G,\Lambda}$. We will use these functions in the following proposition to construct module frames for $\text{QP}_{G,\Lambda}$:

Proposition III.A.3. *Let B be a relatively compact fundamental domain with nonempty interior and let $x_1, \dots, x_k \in G$ be such that $\{x_j B^\circ \Lambda\}_{j=1}^k$ covers G . Let $f \in C(G)$ be a Λ -periodic function with $f|_{\partial B} = 0$ that satisfies*

$$\sum_{j=1}^k |f(x_j^{-1}x)|^2 = 1 \quad (\text{III.44})$$

for all $x \in G$. Set

$$G_j(x, \omega) = f(x_j^{-1}x) J_B(x_j^{-1}x, \omega). \quad (\text{III.45})$$

Then is a normalized tight module frame for $\text{QP}_{G,\Lambda}$.

Proof. As described in the discussion preceding the proposition, the function $f(x) J_B(x, \omega)$ is continuous on the whole of $G \times \widehat{G}$ and hence an element of

$\text{QP}_{G,\Lambda}$. By translation, the functions $f(x_j^{-1}x)J_B(x_j^{-1}x, \omega)$ are also elements of $\text{QP}_{G,\Lambda}$ for each $1 \leq j \leq k$. Finally, we have that

$$\sum_{j=1}^k |G_j(x, \omega)|^2 = \sum_{j=1}^k |f(x_j^{-1}x)|^2 = 1$$

for all $x \in G$. Hence, by Proposition III.4.3, the set $\{G_1, \dots, G_k\}$ is a normalized tight module frame for $\text{QP}_{G,\Lambda}$. $\blacksquare$

Example III.A.4. In this example we show how wavelet filter functions give us module frames as in Proposition III.A.3 in the case $G = \mathbb{R}$, $\Lambda = \mathbb{Z}$. The example closely follows the discussion in [PR04, p. 457].

Let $m: \mathbb{R} \rightarrow \mathbb{C}$ be a continuous wavelet filter function for dilation by 2 (see for instance [Hei11, Section 12.6]), i.e. it is $\mathbb{Z}$ -periodic and satisfies

$$|m(s)|^2 + |m(s + 1/2)|^2 = 1 \quad (\text{III.46})$$

for all $s \in \mathbb{R}$. Suppose also that m is normalized so that $m(0) = 1$. Then by (III.46) we have that $m(1/2) = 0$, which gives $m(k + 1/2) = 0$ for all $k \in \mathbb{Z}$ by periodicity. Let B be the fundamental domain $[0, 1)$ for $\mathbb{Z}$ in $\mathbb{R}$, and set $x_1 = 0$, $x_2 = -1/2$. We see that f given by $f(s) = m(s - 1/2)$ is continuous, vanishes on $\mathbb{Z} = (\partial B)\Lambda$ and satisfies (III.44), so by Proposition III.A.3, the functions

$$\begin{aligned} G_1(s, t) &= f(s)J_{[0,1)}(s, t), \\ G_2(s, t) &= f(s + 1/2)J_{1/2+[0,1)}(s, t) \end{aligned}$$

form a normalized tight frame for $\text{QP}_{\mathbb{R}, \mathbb{Z}}$.

Note that the functions $J_{\beta+[0,1)}$ are exactly the functions $j_{1,1,\beta}$ in [PR04] (and indeed this inspired the notation J_B) and that $\text{QP}_{G,\Lambda}$ is isomorphic to $X(1, 1)$, only that the latter is considered a right module instead of a left module.

Appendix III.B Cutoff functions in the Feichtinger algebra

In this appendix, we have collected some technical results concerning $S_0(G)$ for which the author was unable to find a reference.

The following lemma is used to prove Proposition III.B.2.

Lemma III.B.1. *The following hold in a locally compact abelian group:*

- (i) *If $K \subseteq U$ where K is compact and U is open, then there exists a neighbourhood N of the identity such that $KN \subseteq U$.*
- (ii) *If $K \subseteq V \subseteq \bar{V} \subseteq U$ where K is compact and U, V are open and U is contained in compact set, then there exists a neighbourhood N of the identity such that $K \subseteq \cap_{x \in N} (xV)$ and $\bar{V}N \subseteq U$.*

Proof. We begin by proving i. For every $x \in K$, we can find a neighbourhood N_x of the identity such that $x \in xN_x^2 \subseteq U$ [HR79, p. 18, Theorem (4.5)]. Then we have that the union $\bigcup_{x \in K} (xN_x)$ covers K , so there exists a finite subcovering given by $x_1, \dots, x_n \in K$. Let $N = \bigcap_{k=1}^n N_{x_k}$, which is a nonempty neighbourhood of the identity. If $x \in K$ and $y \in N$, then $x = x_k z$ for some k and $z \in N_{x_k}$. But then $zy \in N_{x_k}^2$, so $xy = x_k zy \in x_k N_{x_k}^2 \subseteq U$. This shows that $KN \subseteq U$.

We now prove ii. Denote by L the compact set that covers U . Since $\bar{V}$ is contained in L , it is compact. Using i, we can find an open neighbourhood N_1 of the identity such that $\bar{V}N_1 \subseteq U$. Taking relative complements in L , we obtain $L \setminus V \subseteq L \setminus K$. Now $L \setminus V$ is closed and contained in L , hence compact, and $L \setminus K$ is open. Thus we can apply i again and obtain an open neighbourhood N_2 of the identity such that $(L \setminus V)N_2 \subseteq L \setminus K$. But then

$$L \setminus ((L \setminus V)N_2) = L \setminus \left(\bigcup_{x \in N_2} x(L \setminus V) \right) = \bigcap_{x \in N_2} L \setminus (x(L \setminus V)) = \bigcap_{x \in N_2} xV$$

so $K \subseteq \bigcap_{x \in N_2} xV$. Setting $N = N_1 \cap N_2$, we see that this set satisfies both $\bar{V}N \subseteq U$ and $K \subseteq \bigcap_{x \in N} (xV)$, so the proof is finished. ■

Proposition III.B.2. *Let K be a compact subset of G , and let U be an open subset of G such that $K \subseteq U$. Then there exists a function $f \in S_0(G)$ such that $f|_K = 1$ and $\text{supp}(f) \subseteq U$.*

Proof. If $K = \emptyset$ then one can just set $f = 0$, so suppose that $K \neq \emptyset$. Since all locally compact groups are normal as topological spaces [HR79, p. 76, Theorem (8.13)], we can find an open set V such that $K \subseteq V \subseteq \bar{V} \subseteq U$, and since K is compact, we can assume that $\bar{V}$ compact. Now without loss of generality, we can assume that U is contained in a compact set.

Let N be an open neighbourhood as in part ii of Lemma III.B.1, i.e. we have both $K \subseteq \bigcap_{t \in N} tV$ and $N\bar{V} \subseteq U$. Since $K \neq \emptyset$, N is necessarily nonempty. Let $\phi \in C_c(G)$ satisfy $\int_G \phi(x) \, dx = 1$ and $\text{supp}(\phi) \subseteq N$. Define $f = \phi * \chi_V$. Then $f \in S_0(G)$ by [Jak18, Lemma 4.2 (iii)]. Now since $\text{supp}(\phi) \subseteq N$, we have that

$$f(x) = \int_G \phi(t) \chi_V(t^{-1}x) \, dt = \int_N \phi(t) \chi_V(t^{-1}x) \, dt.$$

If $x \in K$ then $x \in \bigcap_{t \in N} tV$, so $t^{-1}x \in V$ for all $t \in N$. Thus, $\chi_V(t^{-1}x) = 1$ for all $t \in N$ and hence

$$f(x) = \int_N \phi(t) \, dt = 1.$$

If $x \notin U$ then $x \notin \bar{V}N$ which means $t^{-1}x \notin \bar{V}$ for all $t \in N$, so $\chi_V(t^{-1}x) = 0$. This gives $f(x) = 0$. While this does not necessarily give $x \notin \text{supp}(f)$, we can pick another open set W of G with $K \subseteq W \subseteq \bar{W} \subseteq U$ and choose an $f \in S_0(G)$ as above with $f|_K = 1$ and $f(x) = 0$ whenever $x \notin W$. In that case, $\text{supp}(f) \subseteq \bar{W} \subseteq U$, which gives us what we wanted. ■

The proof of the next result is based on the proof given in [Grö01, Lemma 8.2.1 c)]. The proof employs the Wiener algebra $W(G)$ [Fei77]. Specifically, we will use that $S_0(G) \subseteq W(G)$, which can be seen by combining $W(\mathcal{FL}^1, L^1) = S_0(G)$, cf. [Fei81b, Remark 6], with the inclusion $W(\mathcal{FL}^1, L^1) \subseteq W(L^\infty, L^1)$, cf. [Fei81a, Lemma 1.2(iv)] and $W(G) = W(L^\infty, L^1)$.

Proposition III.B.3. *Let Λ be a lattice in a second countable, locally compact abelian group. If $\xi \in S_0(G)$, then $Z_{G,\Lambda}\xi$ is continuous. Hence, the Zak transform maps $S_0(G)$ into $\text{QP}_{G,\Lambda}$.*

Proof. Suppose $\xi \in S_0(G)$, so that $\xi \in W(G)$. Let B be a fundamental domain for Λ in G with nonempty interior as in Proposition III.A.1. Because of quasiperiodicity, it is enough to prove continuity of $Z\xi$ on $B \times \widehat{G}$, so let $(x, \omega) \in B \times \widehat{G}$ and suppose $(x_\alpha, \omega_\alpha)_\alpha$ is a net in $B \times \widehat{G}$ that converges towards (x, ω) . Let $\epsilon > 0$. There exists an α_1 such that whenever $\alpha \geq \alpha_1$, then $x_\alpha \in B^\circ \subseteq B$.

Since $\|\xi\|_{W(G)}$ is finite, there exists a finite subset F of Λ such that

$$\sum_{\lambda \in \Lambda \setminus F} \sup_{x \in B} |\xi(x\lambda)| < \frac{\epsilon}{2}.$$

Since ξ and ω are continuous, the function $(x, \omega) \mapsto \sum_{\lambda \in F} \xi(x\lambda)\omega(\lambda)$ is continuous, so there exists an α_2 such that we have that

$$\left| \sum_{\lambda \in F} \xi(x_\alpha \lambda) \omega_\alpha(\lambda) - \sum_{\lambda \in F} \xi(x\lambda) \omega(\lambda) \right| < \frac{\epsilon}{2}$$

for $\alpha \geq \alpha_2$. Combining all our observations, we see that if $\alpha \geq \alpha_1$ and $\alpha \geq \alpha_2$ then

$$\begin{aligned} |Z\xi(x_\alpha, \omega_\alpha) - Z\xi(x, \omega)| &= \left| \sum_{\lambda \in \Lambda} \xi(x_\alpha \lambda) \omega_\alpha(\lambda) - \sum_{\lambda \in \Lambda} \xi(x\lambda) \omega(\lambda) \right| \\ &\leq \frac{\epsilon}{2} + \left| \sum_{\lambda \in \Lambda \setminus F} \xi(x_\alpha \lambda) \omega_\alpha(\lambda) - \sum_{\lambda \in \Lambda \setminus F} \xi(x\lambda) \omega(\lambda) \right| \\ &\leq \frac{\epsilon}{2} + \sum_{\lambda \in \Lambda \setminus F} |\xi(x_\alpha \lambda)| + \sum_{\lambda \in \Lambda \setminus F} |\xi(x\lambda)| \\ &\leq \frac{\epsilon}{2} + 2 \sum_{\lambda \in \Lambda \setminus F} \sup_{x \in B} |\xi(x\lambda)| \\ &\leq \frac{\epsilon}{2} + 2 \cdot \frac{\epsilon}{4} \\ &= \epsilon. \end{aligned}$$

This finishes the proof. ■

Acknowledgements. The author is indebted to Nadia Larsen, Franz Luef, Judith Packer and Leonard Huang for valuable conversations. The author is grateful for the hospitality of Judith Packer, Carla Farsi and the rest of the operator algebras group at the University of Colorado, Boulder during his stay there in 2018–2019, during which a lot of the paper was written. The stay was supported by the Fulbright Foundation and the Trond Mohn Foundation. The author also wants to thank Mads S. Jakobsen for pointing out helpful references, and Andreas Andersson, Are Austad, Martin Helsø and Petter Nyland for giving feedback on drafts of the paper. Finally, the author wants to thank the referee for valuable feedback, in particular for pointing out how to better structure Section III.4, leading to shorter proofs and less dependence on fundamental domains.

References

- [AE19] Austad, A. and Enstad, U. “Heisenberg modules as function spaces”. Preprint, arXiv:1904.10826. 2019.
- [AFK14] Ascensi, G., Feichtinger, H. G., and Kaiblinger, N. “Dilation of the Weyl symbol and Balian-Low theorem”. In: *Trans. Amer. Math. Soc.* Vol. 366, no. 7 (2014), pp. 3865–3880.
- [Bal01] Balan, R. “Topological obstructions to localization results”. In: *SPIE Wavelets Applications in Signal and Image Processing VIII* (2001), pp. 184–192.
- [Bal81] Balian, R. “Un principe d’incertitude fort en théorie du signal ou en mécanique quantique”. In: *C. R. Acad. Sci. Paris Sér. II Méc. Phys. Chim. Sci. Univers Sci. Terre* vol. 292, no. 20 (1981), pp. 1357–1362.
- [Bat88] Battle, G. “Heisenberg proof of the Balian-Low theorem”. In: *Lett. Math. Phys.* Vol. 15, no. 2 (1988), pp. 175–177.
- [BHW95] Benedetto, J. J., Heil, C., and Walnut, D. F. “Differentiation and the Balian-Low theorem”. In: *J. Fourier Anal. Appl.* Vol. 1, no. 4 (1995), pp. 355–402.
- [Bla98] Blackadar, B. *K-theory for operator algebras*. Second. Vol. 5. Mathematical Sciences Research Institute Publications. Cambridge University Press, Cambridge, 1998, pp. xx+300.
- [Boc99] Boca, F. P. “Projections in rotation algebras and theta functions”. In: *Comm. Math. Phys.* Vol. 202, no. 2 (1999), pp. 325–357.
- [CL18] Chakraborty, S. and Luef, F. “Duality of Gabor Frames and Heisenberg Modules”. Preprint, arXiv:1705.11194. 2018.
- [DS52] Duffin, R. J. and Schaeffer, A. C. “A class of nonharmonic Fourier series”. In: *Trans. Amer. Math. Soc.* Vol. 72 (1952), pp. 341–366.
- [EJL19] Enstad, U. B. R., Jakobsen, M. S., and Luef, F. “Time-frequency analysis on the adèles over the rationals”. In: *Comptes Rendus Mathématique* vol. 357 (2019), pp. 188–199.

-
- [Fei77] Feichtinger, H. G. “A characterization of Wiener’s algebra on locally compact groups”. In: *Arch. Math. (Basel)* vol. 29, no. 2 (1977), pp. 136–140.
 - [Fei81a] Feichtinger, H. G. “Banach spaces of distributions of Wiener’s type and interpolation”. In: *Functional analysis and approximation (Oberwolfach, 1980)*. Vol. 60. Internat. Ser. Numer. Math. Birkhäuser, Basel-Boston, Mass., 1981, pp. 153–165.
 - [Fei81b] Feichtinger, H. G. “On a new Segal algebra”. In: *Monatsh. Math.* Vol. 92, no. 4 (1981), pp. 269–289.
 - [FK04] Feichtinger, H. G. and Kaiblinger, N. “Varying the time-frequency lattice of Gabor frames”. In: *Trans. Amer. Math. Soc.* Vol. 356, no. 5 (2004), pp. 2001–2023.
 - [FL02] Frank, M. and Larson, D. R. “Frames in Hilbert C^* -modules and C^* -algebras”. In: *J. Operator Theory* vol. 48, no. 2 (2002), pp. 273–314.
 - [Gel50] Gelfand, I. M. “Expansion in characteristic functions of an equation with periodic coefficients”. In: *Doklady Akad. Nauk SSSR (N.S.)* Vol. 73 (1950), pp. 1117–1120.
 - [GL04] Gröchenig, K. and Leinert, M. “Wiener’s lemma for twisted convolution and Gabor frames”. In: *J. Amer. Math. Soc.* Vol. 17, no. 1 (2004), pp. 1–18.
 - [Grö01] Gröchenig, K. *Foundations of time-frequency analysis*. Applied and Numerical Harmonic Analysis. Birkhäuser Boston, Inc., Boston, MA, 2001, pp. xvi+359.
 - [Grö98] Gröchenig, K. “Aspects of Gabor analysis on locally compact abelian groups”. In: *Gabor analysis and algorithms*. Appl. Numer. Harmon. Anal. Birkhäuser Boston, Boston, MA, 1998, pp. 211–231.
 - [GVF01] Gracia-Bondía, J. M., Várilly, J. C., and Figueroa, H. *Elements of noncommutative geometry*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Boston, Inc., Boston, MA, 2001, pp. xviii+685.
 - [Hei07] Heil, C. “History and evolution of the density theorem for Gabor frames”. In: *J. Fourier Anal. Appl.* Vol. 13, no. 2 (2007), pp. 113–166.
 - [Hei11] Heil, C. *A basis theory primer*. Expanded. Applied and Numerical Harmonic Analysis. Birkhäuser/Springer, New York, 2011, pp. xxvi+534.
 - [Hei90] Heil, C. E. *Wiener amalgam spaces in generalized harmonic analysis and wavelet theory*. Thesis (Ph.D.)—University of Maryland, College Park. ProQuest LLC, Ann Arbor, MI, 1990, p. 314.

- [HR79] Hewitt, E. and Ross, K. A. *Abstract harmonic analysis. Vol. I.* Second. Vol. 115. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin-New York, 1979, pp. ix+519.
- [Jak18] Jakobsen, M. S. “On a (No Longer) New Segal Algebra: A Review of the Feichtinger Algebra”. In: *J. Fourier Anal. Appl.* Vol. 24, no. 6 (2018), pp. 1579–1660.
- [JL16] Jakobsen, M. S. and Lemvig, J. “Density and duality theorems for regular Gabor frames”. In: *J. Funct. Anal.* Vol. 270, no. 1 (2016), pp. 229–263.
- [JL18] Jakobsen, M. S. and Luef, F. “Duality of Gabor Frames and Heisenberg Modules”. Preprint, arXiv.org:1806.05616. 2018.
- [KK98] Kaniuth, E. and Kutyniok, G. “Zeros of the Zak transform on locally compact abelian groups”. In: *Proc. Amer. Math. Soc.* Vol. 126, no. 12 (1998), pp. 3561–3569.
- [Lan95] Lance, E. C. *Hilbert C^* -modules*. Vol. 210. London Mathematical Society Lecture Note Series. A toolkit for operator algebraists. Cambridge University Press, Cambridge, 1995, pp. x+130.
- [Lee13] Lee, J. M. *Introduction to smooth manifolds*. Second. Vol. 218. Graduate Texts in Mathematics. Springer, New York, 2013, pp. xvi+708.
- [LM16] Lesch, M. and Moscovici, H. “Modular curvature and Morita equivalence”. In: *Geom. Funct. Anal.* Vol. 26, no. 3 (2016), pp. 818–873.
- [Low85] Low, F. “Complete sets of wave packets”. In: *A Passion for Physics: Essays in Honor of Geoffrey Chew*. World Scientific, 1985, pp. 17–22.
- [LP13] Latrémolière, F. and Packer, J. A. “Noncommutative solenoids and their projective modules”. In: *Commutative and noncommutative harmonic analysis and applications*. Vol. 603. Contemp. Math. Amer. Math. Soc., Providence, RI, 2013, pp. 35–53.
- [LP15] Latrémolière, F. and Packer, J. A. “Explicit construction of equivalence bimodules between noncommutative solenoids”. In: *Trends in harmonic analysis and its applications*. Vol. 650. Contemp. Math. Amer. Math. Soc., Providence, RI, 2015, pp. 111–140.
- [Lue09] Luef, F. “Projective modules over noncommutative tori are multi-window Gabor frames for modulation spaces”. In: *J. Funct. Anal.* Vol. 257, no. 6 (2009), pp. 1921–1946.
- [Lue11] Luef, F. “Projections in noncommutative tori and Gabor frames”. In: *Proc. Amer. Math. Soc.* Vol. 139, no. 2 (2011), pp. 571–582.
- [Lue18] Luef, F. “The Balian-Low theorem and noncommutative tori”. In: *Expo. Math.* Vol. 36, no. 2 (2018), pp. 221–227.
- [Mac52] Mackey, G. W. “Induced representations of locally compact groups. I”. In: *Ann. of Math. (2)* vol. 55 (1952), pp. 101–139.

-
- [Mar91] Margulis, G. A. *Discrete subgroups of semisimple Lie groups*. Vol. 17. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)* [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1991, pp. x+388.
 - [MS74] Milnor, J. W. and Stasheff, J. D. *Characteristic classes*. *Annals of Mathematics Studies*, No. 76. Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1974, pp. vii+331.
 - [NO19] Nitzan, S. and Olsen, J.-F. “Balian–Low type theorems in finite dimensions”. In: *Mathematische Annalen* vol. 373, no. 1 (Feb. 2019), pp. 643–677.
 - [PR04] Packer, J. A. and Rieffel, M. A. “Projective multi-resolution analyses for $L^2(\mathbb{R}^2)$ ”. In: *J. Fourier Anal. Appl.* Vol. 10, no. 5 (2004), pp. 439–464.
 - [Rie10] Rieffel, M. A. “Vector bundles and Gromov-Hausdorff distance”. In: *J. K-Theory* vol. 5, no. 1 (2010), pp. 39–103.
 - [Rie83] Rieffel, M. A. “The cancellation theorem for projective modules over irrational rotation C^* -algebras”. In: *Proc. London Math. Soc. (3)* vol. 47, no. 2 (1983), pp. 285–302.
 - [Rie88] Rieffel, M. A. “Projective modules over higher-dimensional noncommutative tori”. In: *Canad. J. Math.* Vol. 40, no. 2 (1988), pp. 257–338.
 - [RV99] Ramakrishnan, D. and Valenza, R. J. *Fourier analysis on number fields*. Vol. 186. *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1999, pp. xxii+350.
 - [RW98] Raeburn, I. and Williams, D. P. *Morita equivalence and continuous-trace C^* -algebras*. Vol. 60. *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1998, pp. xiv+327.
 - [Swa62] Swan, R. G. “Vector bundles and projective modules”. In: *Trans. Amer. Math. Soc.* Vol. 105 (1962), pp. 264–277.
 - [Wei64] Weil, A. “Sur certains groupes d’opérateurs unitaires”. In: *Acta Math.* Vol. 111 (1964), pp. 143–211.
 - [Wil07] Williams, D. P. *Crossed products of C^* -algebras*. Vol. 134. *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2007, pp. xvi+528.
 - [Zak67] Zak, J. “Finite Translations in Solid-State Physics”. In: *Phys. Rev. Lett.* Vol. 19 (24 Dec. 1967), pp. 1385–1387.