Dietary variations and specializations of killer whales (*Orcinus orca*) in Norway

Eve Jourdain

A dissertation presented for the degree
of Doctor Philosophiae (Dr Philos)
2020

Norwegian Orca Survey, Andenes

UiO

Department of Biosciences
Faculty of Mathematics and Natural Sciences
University of Oslo

© Eve Jourdain, 2020

*Series of dissertations submitted to the Department of Biosciences Faculty of Mathematics
and Natural Sciences, University of Oslo No. 2321*

ISSN 1501-7710

All rights reserved. No part of this publication should be reproduced, transmitted or used, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.

Photo cover: Eve Jourdain

Print production: Reprosentralen, University of Oslo.

ACKNOWLEDGEMENTS

Conducting this research has been the dream of a lifetime and I would like to thank everyone who contributed to make it a reality. My dear colleague and friend **Dag Vongraven** has played a tremendous role by believing in my determination, by giving me his endless support at all times and by contributing his dataset to my research. This research wouldn't have been possible without the fantastic faith and support from my friend **Todd Robeck** who believed in this project and never missed a chance to help with funding opportunities. I wish to thank my friend and colleague **Tiu Similä** for her support and for sharing her knowledge and material.

I would like to acknowledge my amazing partner **Richard Karoliussen**. While I had no plan to settle down in northern Norway, our love story led to me building this research which we committed to conduct together as partners both in life and in the field. He has earned as much credit on this project and I owe him a huge part of this success.

I wish to thank **Clare Andvik** for her creativity, hard work and determination and for being such a great team member. **Katrine Borgå** played an important role in facilitating the laboratory analyses and helping me with the Dr Philos.

I am grateful to **Marten Bril** and other people at **Sea Safari Andenes** for reporting killer whale sightings throughout the years and putting efforts into collecting identification photographs.

I wish to thank my family and friends for their constant support and for handling my tiresome passion for killer whales and my tedious need to constantly achieve dreams and goals over the years.

I wish to thank my friends and former colleagues at **Marineland, Antibes** and **Sea World, San Diego** for giving me the chance to experience a life with daily close contact with killer whales some years ago. These years gave me a life-time experience which greatly inspired my wish and determination to study killer whales in the wild and act for their protection.

This research was greatly inspired by killer whale studies conducted by pioneers in the field in British Columbia, Canada.

This study would have never been possible without the generous grants from the **Sea World and Busch Gardens Conservation fund, Sea World and Parks Entertainment** and **Spare Bank-1 Nord-Norge** to Norwegian Orca Survey. **Regatta Norway** and **Steiner** provided sponsorship with great gear in the field.

Table of Contents

SUMMARY	9
LIST OF PAPERS INCLUDED IN THE THESIS	11
I- INTRODUCTION	12
A- DIETARY PATTERNS: CAUSES AND CONSEQUENCES.....	12
B- DIETARY VARIATIONS IN CETACEANS	12
C- CASE STUDY - KILLER WHALES IN NORWAY	13
D- DIETARY SPECIALIZATIONS IN KILLER WHALES.....	16
II- OBJECTIVES.....	22
III- MATERIALS AND METHODS	24
A- STUDY AREA.....	24
B- SURVEYS	25
C- DATA COLLECTION AND PROCESSING	25
IV- RESULTS AND DISCUSSION.....	28
A- OBSERVATIONAL EVIDENCE	28
B- EVIDENCE FROM DIETARY MARKERS	35
C- POSSIBLE DRIVING FORCES OF DIETARY DIVERSITY.....	41
D- IMPLICATIONS OF DIETARY VARIATIONS/SPECIALIZATIONS	43
CLOSING REMARKS	46
REFERENCES.....	47
AUTHORSHIP CONTRIBUTIONS	60

SUMMARY

Killer whales (*Orcinus orca*) are found in all the world's oceans and even though a generalist predator, local populations tend to adopt remarkably specialized diets. Foraging specializations shape movement, behavior and sociality and thus have substantial conservation implications in this species. In Norway, killer whales have long been known to associate with the Norwegian Spring Spawning stock of Atlantic herring (*Clupea harengus*) as primary food source. Over the years, indications of additional prey types have arisen but, due to research efforts largely focused on wintering herring grounds, dietary patterns have remained poorly understood. This thesis investigates the incidence of inter-individual/group dietary variations and possible foraging specializations in killer whales in northern Norway.

Killer whales were photo-identified and their occurrence monitored throughout the year in 2013-2018 in northern Norway. For each encounter, systematic efforts aimed at collecting data on individuals' identity, behavior, group size and group composition. Whenever possible, target prey was identified either visually or through molecular identification of prey fragments retrieved from sub-surface feeding events. Biopsy samples of photo-identified killer whales were also collected in 2017-2018 for further analysis of skin nitrogen and carbon isotopic ratios in skin, total mercury levels in skin and polychlorinated biphenyls (PCBs) in blubber as dietary descriptors. Contaminant burdens were explored for the potential conservation implications of dietary patterns. Combining multiple data types collected in all seasons and over multiple years, this multi-disciplinary long-term study was the very first of its kind in Norway and allowed for new insights into the foraging ecology of its killer whale population.

Killer whale occurrence peaked in the winter months of high herring abundance, confirming the herring as a major food source. While most identified individuals left the study area after the herring initiated its migration southwards, ~75 herring-feeding killer whales returned to the study area each spring (March-April). The lumpfish (*Cyclopterus lumpus*) was identified as a seasonal target prey for these whales. Another ~15 killer whales, rarely or not seen at herring wintering grounds, were encountered in multiple years from May through October and consistently fed on harbor seal (*Phoca vitulina*) and gray seal (*Halichoerus grypus*).

Recent distributional shifts in wintering herring has promoted the new appearance of humpback whales (*Megaptera novaeangliae*) which fed alongside killer whales during this study, often on common prey patches at herring wintering grounds. Whilst killer whales appeared as the catalyst species in all mixed-species feeding aggregations, observational evidence pointed towards humpback whales depredating herring balls that had been herded by killer whales. These findings highlighted the dynamic nature of ecological interactions in varying environmental contexts and how competition costs may fluctuate. These processes may further influence dietary patterns.

Skin isotopic values of killer whales known to have been feeding on herring, lumpfish or seals at time of sampling revealed distinct isotopic niches with clusters that coincided with predatory

field observations. Skin isotopic values of lumpfish-eaters supported seasonal specialization on lumpfish. While these whales were known to be feeding on herring from the winter months in the same years, this constitutes the first evidence for killer whales seasonally switching between alternative prey resources and foraging behaviors in Norway.

Elevated nitrogen values for seal-eating individuals, regardless of sampling season and also at times of high fish prey availability, supported episodic field observations to be representative of these whales' long-term feeding habits. Predation on pinnipeds occurred throughout the year and persisted overtime, and up to several decades. Behavioral characteristics revealed a certain level of specialization to efficiently forage upon pinnipeds for these whales. However, isotopic values and field observations confirmed seal-eating individuals to also include fish prey in their diet. Estimates of relative dietary proportions further suggested an ecological gradient within this killer whale population, with individuals differing in the proportional contribution of different prey items to their diet.

Concentration profiles of PCBs and mercury in blubber and skin samples, respectively supported inter-individual variations in long-term dietary habits. As expected for individuals feeding at higher trophic levels, seal-eating killer whales showed PCB blubber concentrations three times higher and mercury skin levels twice as high as fish-eaters. These findings translated to a higher risk for health effects, with potential consequences at the population level. Whilst killer whales in Norway were previously wrongly assigned to a low risk category of pollution effects based on tissue samples collected from fish-eating individuals, this thesis calls for urgent revision of the status of this killer whale population, accounting for ecological differentiation among social groups.

This study contributes to our current understanding of foraging specializations in killer whales. By providing findings from a newly investigated region, this work broadens the possibilities for comparative studies, necessary to assess whether dietary specialization is a characteristic of the species and to identify underlying mechanisms and implications. This thesis brings the first conclusive evidence for intra-population variations in foraging specializations in the North Atlantic and encourages dietary studies in other adjacent regions.

LIST OF PAPERS INCLUDED IN THE THESIS

Paper I.

Jourdain, E., and Vongraven, D. (2017). Humpback whale (*Megaptera novaeangliae*) and killer whale (*Orcinus orca*) feeding aggregations for foraging on herring (*Clupea harengus*) in northern Norway. *Mammalian Biology*, 86, 27-32. doi: 10.1016/j.mambio.2017.03.006.

Paper II.

Jourdain, E., Karoliussen, R., de Vos, J., Zakharov, S. E., and Tougard, C. (2019). Killer whales (*Orcinus orca*) feeding on lumpfish (*Cyclopterus lumpus*) in northern Norway. *Marine Mammal Science*. doi: 10.1111/mms.12618.

Paper III.

Jourdain, E., Vongraven, D., Bisther, A., and Karoliussen, R. (2017). First longitudinal study of seal-feeding killer whales (*Orcinus orca*) in Norwegian coastal waters. *PloS one*, 12(6), e0180099. doi: 10.1371/journal.pone.0180099.

Paper IV.

Jourdain, E., Andvik, C., Karoliussen, R., Ruus, A., Vongraven, D., and Borgå, K. (2020). Isotopic niche differs between seal and fish-eating killer whales (*Orcinus orca*) in northern Norway. *Ecology and Evolution*, 00: 1-13. doi://10.1002/ece3.6182.

Paper V.

Andvik, C., **Jourdain, E.**, Ruus, A., Lyche, J. L., Karoliussen, R., and Borgå, K. At risk: Preying on seals pushes killer whales from Norway above pollution effect thresholds. *Scientific Reports*, 10:11888. doi://10.1038/s41598-020-68659-y.

I- INTRODUCTION

A- Dietary patterns: causes and consequences

Conspecific individuals have long been treated as equivalent in the study of species' or populations' diets. However, there has been growing evidence for individual variations in the use of a common pool of available resources across a wide range of taxa (e.g. fish: Holbrook & Schmitt 1992, Warburton et al. 1998, mammals: Estes et al. 2003, birds: Ceia & Ramos 2015). As such, dietary variation may arise from phenotypic variation (Price 1987, Hori 1993) or a diverse array of physiological (Afik & Karasov 1995), ecological and behavioral mechanisms that may promote individual specialization or generalization as a consumer's foraging strategy (Warburton et al. 1998, Estes et al. 2003, Svanbäck & Bolnick 2005, Svanbäck & Bolnick 2007, Tinker et al. 2008, Araújo et al. 2011). Prey specialization, also referred to as ecological or foraging specialization, occurs when there is behavioral specialization to only use a limited fraction of available resources (Bolnick et al. 2003, Araújo et al. 2011). It may arise when individuals benefit from enhanced foraging efficiency, owing to increased experience with feeding conditions over non-specialists (Tinker et al. 2008), and if foraging upon a range of taxonomically diverse prey types requires different, sometimes complex, behaviors for efficient use (Cunningham & Hughes 1984). Thus, a generalist population may be a mixture of true generalist or specialized individuals. Through increased variation in individuals' fitness and disruptive selection, individuality in foraging behavior has a broad range of implications for both population/community dynamics and evolutionary processes and should therefore be monitored (see Bolnick et al. 2003).

B- Dietary variations in cetaceans

In cetaceans, dietary specializations have received particular attention for several reasons: they commonly occur at the individual or group level (e.g. bird-associated foraging by minke whales *Balaenoptera acutorostrata*: Hoelzel et al. 1989, beach hunting by bottlenose dolphins *Tursiops sp*: Sargeant et al. 2005) or population level (e.g. lobtail and bubble net feeding in humpback whales *Megaptera novaeangliae*: Weinrich et al. 1992), they are maintained over long periods of time through cultural transmission (Rendell & Whitehead 2001, Kopps et al. 2014) and have been identified as potential drivers of population genetic structure, through cultural behavioral differentiation, in a barrierless environment (Hoelzel 1998, Kopps et al. 2014, Louis et al. 2014). As highly mobile, large body-sized predators that occupy top trophic

positions and with high daily prey requirements, cetacean prey-specialists may substantially impact prey populations and communities through top-down effects (e.g. Williams et al. 2004). The interest in assessing their role in marine ecosystems has therefore substantially added motivation to conducting dietary studies in cetaceans. Whilst underlying mechanisms have been poorly understood so far, assessing dietary patterns of extant cetacean populations and across regions are warranted for further comparative studies and to gain a better understanding of associated causes and consequences.

Due to ecological characteristics and substantial implications detailed hereafter, dietary variations are of particular importance to the study and management of killer whales (*Orcinus orca*). The species may also be considered a particularly suitable candidate for dietary studies for several reasons. They are reliably identifiable at the individual level (Bigg 1982), there has been a wide range of methods developed for the investigation of their diet and feeding ecology (observation of foraging behavior: e.g. Baird & Dill 1995, direct observations of surface predation, analysis of stomach contents from beached carcasses: e.g. Ford et al. 1998, collection of prey fragments for molecular identification of prey species: Ford & Ellis 2006, chemical analysis of skin and blubber biopsy samples collected from live individuals: e.g. Krahn et al. 2007) and there is a large body of knowledge available on their ecology and behavior worldwide, highly beneficial in interpreting results.

C- Case study - Killer whales in Norway

a- Background studies

Killer whales occur all along the Norwegian coast, in both nearshore and offshore waters and have long been known to forage on the Norwegian Spring Spawning stock of the Atlantic herring (*Clupea harengus*), as supported by stomach contents and occurrence patterns (Grieg 1906, Collett 1912, Jonsgård & Lyshoel 1970, Christensen 1982, Christensen & Øritsland 1982, Christensen 1988, Øien 1988). Because they were perceived as competitors to fishery industries, killer whales were culled by Norwegian whalers with a total of 2,435 catches in 1938-1981 across the North Atlantic, including 80% from the Norwegian Sea (Christensen 1982, Øien 1988). In 1982, and upon recommendations from the International Whaling Commission, the Norwegian killer whale catch ended (Christensen 1988, Øien 1988). The commission further recommended that photographic-based population assessments, as first

initiated in the eastern North Pacific (Bigg 1982), be carried out. In 1983-1985, a pilot study initiated off the coast of Møre and Lofoten marked the beginning of the long-term photographic identification study of killer whales in Norway for the purpose of studying this population's life history and ecology (Lyrholm 1984, 1987).

Surveys were conducted each year at herring wintering grounds in a fjord system (Vestfjord, Tysfjord, Ofotfjord) of the Lofoten region, northern Norway, in the period 1983-2005 (Lyrholm 1984, 1987, Bisther & Vongraven 1995, Similä et al. 1996, Kuningas et al. 2014), and in spring at the herring spawning grounds located off Møre from 1987 to 1996 (Bisther & Vongraven 1995, 2001). Between 1986 and 2003, up to 700 killer whales were present at herring wintering grounds (Similä et al. 1996, Kuningas et al. 2014, Vongraven & Bisther 2014). Surface and underwater observations of feeding behaviors revealed specialized techniques, including the highly cooperative carousel feeding in which killer whales coordinate their swimming by means of acoustic signaling to better herd the herring in a tight ball, close to the surface, and use the underside of their flukes to debilitate the prey (Similä & Ugarte 1993, Similä 1997, Simon et al. 2007a). Subsurface and travel feeding were described as additional, apparently less cooperative, feeding techniques (Similä 1997). As a highly adaptive predator, killer whales were also seen in association with herring purse-seiners, directly benefitting from compromised or dead herring falling from the nets during fishing operations (Van Opzeeland et al. 2005).

While behavioral observations confirmed killer whales as efficient herring predators, the following converging lines of evidence suggested potential year-round specialization. As such, re-sightings of photographically identified individuals at matching time of year and location, over consecutive years, confirmed killer whale site fidelity relative to seasonal herring abundance (Similä et al. 1996). Additionally, some individuals were shown to be seasonally moving between herring spawning and wintering grounds, as documented by identification photographs (Lyrholm 1988) and satellite tracking of seven satellite-tagged individuals in 2000-2001 (Similä & Stenersen 2004). Behavioral plasticity relative to varying distribution, abundance and schooling patterns that characterize the different herring life-stages i.e., wintering, spawning and feeding (Holst et al. 2004) confirmed the killer whales' ability to exploit their herring prey throughout the year (Samarra & Miller 2015).

Nitrogen stable isotopic ratios measured for killer whales sampled at herring wintering grounds ($n=17$ in 2005, 2007) and feeding grounds ($n=2$ in 2001, 2002) showed little variance, in support of a herring-specialized diet (Foote et al. 2012). Yet, enriched ^{15}N values for some sampled individuals were indicative of predation on higher trophic level prey for these whales and pointed towards possible inter-individual dietary variations. A lack of predation history for these whales however, precluded any meaningful interpretation of these apparent fine-scale dietary variations. In fact, historical predation records on squid, cephalopods, cod (*Gadus morhua*), porpoises, seals and bottlenose whales (*Hyperoodon ampullatus*) had already suggested dietary diversity for killer whales in Norway several decades ago (Collett 1912, Jonsgård 1968, Jonsgård & Lyshoel 1970, Christensen 1982). However, with 20 years (1986-2006) of field efforts largely focused on herring grounds, it was unlikely that the importance of other prey resources would be revealed.

b- Evidence for dietary variations

With recent expansion of research efforts to include all seasons, observations brought new evidence for ecological variations and documented new prey types for killer whales in Norway. A very first investigation of predation on pinnipeds compiled seven predatory events from central and northern coasts of Norway between 1988 and 2011 (Vongraven & Bisther 2014). Photographic identification of individual killer whales involved, also known as herring-eaters from field observations recorded in the same years, confirmed some social groups to adopt a mixed diet i.e. fish and mammal prey, as suggested by (Foote et al. 2009). As three of the seven predation events involved the same individual killer whales, and despite a large population size for this region (Kuningas et al. 2014), the authors suggested that only a small number of specialized killer whale groups may be feeding upon pinniped prey in Norway (Vongraven & Bisther 2014). A single account of predation on harbor porpoise (*Phocoena phocoena*), for which individual killer whales could not be photo-identified, further added to the list of mammalian prey in Norwegian coastal waters (Cosentino 2015).

New fish prey resources were also identified. A six month-study conducted in May-November 2011 revealed two killer whale groups apparently seasonally foraging on Atlantic salmon (*Salmo salar*) in the Lofoten region (Vester & Hammerschmidt 2013). Fishery surveys conducted offshore in the Norwegian Sea in July-August of 2007-2008 also revealed killer whale occurrence in association with the northeastern Atlantic mackerel (*Scomber scombrus*)

schools (Nøttestad et al. 2014). Greater availability i.e. schools distributed in upper waters throughout the day and night, and higher nutritional values for the mackerel compared to the herring, suggest it as a seasonally more profitable fish prey and could explain why killer whales may forage preferentially on mackerel in the summer months offshore (Nøttestad et al. 2014). Because mackerel-foraging killer whales were not photographically identified, it is unknown whether they are the same individuals as the ones identified feeding on herring in the winter months in northern Norwegian fjords (Similä & Ugarte 1993, Similä et al. 1996, Similä 1997, Van Opzeeland et al. 2005).

c- Knowledge gaps

In contrast with the initial assumption that the preferred prey of killer whales was herring in all seasons in Norway, the actual existence of ecological variations within this population has appeared increasingly clear. However, the lack of identification data and/or only brief periods of data collection for these recent studies has precluded any assessment of dietary differences and specializations among killer whale individuals/groups. A more accurate description of this population is needed, accounting for heterogeneity in patterns of resource use, as well as the investigation of further implications this may have for management and conservation.

D- Dietary specializations in killer whales

a- Diet and ecotypes

The killer whale occupies the top trophic position throughout its cosmopolitan range and, as a species, is considered a generalist predator with a wide range of ~140 prey species including cetaceans, pinnipeds, bony and cartilaginous fish, seabirds, squids, octopuses and sea turtles (Ford 2009). However, where killer whale densities peak in productive high latitude waters, local populations tend to adopt remarkably specialized diets, only selectively foraging on a narrow range of available prey. This was first documented in coastal waters of the eastern North Pacific where three killer whale forms, or ecotypes, show divergent ecological specializations despite overlapping distributions. In this region, so-called Resident killer whales appear to exclusively feed on teleost fishes, and primarily on salmonids, and squid; Transient killer whales selectively forage on marine mammal prey; and Offshore killer whales appear to specialize on shark and other high trophic level fish prey (Bigg et al. 1985, Bigg et al. 1987, Baird & Dill 1995, Ford et al. 1998, Ford & Ellis 2006, Ford et al. 2011). Forty-five

years of research that engaged field observations, analysis of stomach contents of beached carcasses and dietary markers in biopsy tissue samples supported fundamentally distinct, non-overlapping diets for these ecotypes (Ford et al. 1998, Saulitis et al. 2000, Herman et al. 2005, Krahn et al. 2007). Likely due to the specific requirements entitled to efficiently encounter and capture taxonomically and behaviorally divergent prey types, the three ecotypes differ in habitat use patterns (Baird & Dill 1995, Nichol & Shackleton 1996, Saulitis et al. 2000, Ford & Ellis 2006, Matkin et al. 2007, Dahlheim et al. 2008), social organization (Bigg et al. 1987, Baird & Dill 1996, Baird & Whitehead 2000), and acoustic and foraging behaviors (Baird & Dill 1995, Barrett-Lennard et al. 1996, Saulitis et al. 2000, Deecke et al. 2005).

Similar ecological specializations have been described in several regions worldwide. Two non-associating Resident and Transient forms, with similar characteristics as described above, have been documented in the Russian Far East (Burdin et al. 2004, Filatova et al. 2015). In Antarctic waters, five ecotypes, at least partially sympatric, are also currently recognized. Type A killer whales occur in open waters and appear to mainly feed on minke whales (*Balaenoptera bonaerensis*; Pitman & Ensor 2003). Type B1, also called Pack-ice killer whales, selectively forage on Weddell seals (*Leptonychotes weddellii*) among loose pack ice throughout its circumpolar range (Pitman & Durban 2012). Smaller type B2, also called Gerlache Strait killer whales, forages mainly on penguins in the Antarctic Peninsula (Pitman & Durban 2012). Type C, also known as Ross Sea killer whales, appear as fish-specialists in the deep pack ice (Pitman & Ensor 2003, Pitman et al. 2018). The most distinctive killer whale morphotype Type D, for which only six observations in deep oceanic waters were recorded in 2004-2010, adopt a subantarctic distribution and possibly a fish-based diet (Pitman et al. 2011). In the North Atlantic, two partially sympatric killer whale ecotypes have been suggested (Foote et al. 2009). Ecotype 1 includes fish-eating killer whales of Norway, Iceland and the UK region that associate with regional stocks of Atlantic herring and/or Northeastern Atlantic mackerel, but with individuality in the proportions of dietary components, including high trophic level pinniped prey (Foote et al. 2010, Vongraven & Bisther 2014). In contrast, ecotype 2 appears as a prey-specialist mainly feeding on baleen whales (Foote et al. 2009).

b- Underlying mechanisms of dietary patterns

Foraging specialization documented for most well-studied killer whale populations worldwide suggests it as a possible characteristic of the species. However, individual specialization may

depend on habitat productivity (Baird et al. 2006, Araújo et al. 2011), likely favored under stable conditions of prey abundance. Conversely, a diversified diet, with ability to switch among alternative prey types, may be more profitable under conditions of frequently fluctuating prey populations (Araújo et al. 2011). Keeping with this theory, killer whales occurring in tropical low productivity waters seem to be feeding on a much broader array of prey types compared to the high latitude killer whale populations mentioned above (Baird et al. 2006, Weir et al. 2010, Bolanos-Jimenez et al. 2014). However, long-term ecological data is still lacking for tropical regions and have thus not allowed for consistency of individuals' dietary habits to be tested. Therefore, group-specific dietary specializations cannot be excluded. It should be noted however that populations of true generalist killer whales i.e. with individuals feeding on multiple prey types (fish, birds and/or mammals), were documented from high latitude productive regions in both hemispheres (Crozet Archipelago: Guinet 1992, South Africa: Best et al. 2010, Iceland/Scotland: Foote et al. 2010, South America: Capella et al. 2014, Norway: Vongraven & Bisther 2014, Marion Island: Reisinger et al. 2016). In fact, theoretical modelling shows specialization can be advantageous in the short term, but may reduce resilience and increase vulnerability in the long term (Whitehead & Ford 2018).

Ecological interactions possibly constitute another determinant of dietary patterns in killer whales (Baird et al. 1992). By reducing availability of preferred resources, intraspecific competition may promote behavioral diversification and individual specialization (Svanbäck & Bolnick 2005, Svanbäck & Bolnick 2007, Tinker et al. 2008, Kobler et al. 2009). Interspecific competition could have both positive and negative effects on individual specialization (Svanbäck & Bolnick 2005, Araújo et al. 2011).

c- Implications of dietary specialization

i- Contamination patterns

Persistent and biomagnifying organohalogen contaminants such as polychlorinated biphenyls (PCBs) and organochlorine pesticides (e.g. dichlor-diphenyl-trichloroethane: DDTs) bioaccumulate in lipid tissues of top marine predators, and killer whales were listed among the most contaminated cetaceans in the world (Ross et al. 2000, Wolkers et al. 2007, Desforges et al. 2018). Because accumulation patterns of these pollutants are related to dietary habits and trophic level of individuals, resource-specialized sympatric ecotypes/populations/groups may be exposed to different contaminant levels and associated toxicological risks, e.g.

immunotoxicity and endocrine disruption (Ross et al. 2000, Ylitalo et al. 2001, Jepson et al. 2016, Pedro et al. 2017). For instance, marine mammal eating killer whales in the eastern North Pacific had PCB blubber concentrations six times higher than that of the sympatric Northern Resident community of fish-eating killer whales (Ross et al. 2000). Multiple synergistic stressors e.g. food depletion may further have cumulative/additive effects and impact populations in various ways. Exposure and vulnerability to contaminant concentrations should be monitored and considered in management actions.

ii- Population dynamics

Prey availability was shown to be a determinant of killer whales' survival and reproduction, ultimately driving population dynamics (Ward et al. 2009, Ford et al. 2010). Populations adopting divergent foraging strategies may follow different, or even contrasted, demographic trends owing to fluctuations in specific preferred prey populations. In the eastern North Pacific for instance, substantial differences in abundance and population trends were detected between the two sympatric Southern Resident and West Coast Transient populations (and ecotypes) since the mid-1990s (COSEWIC 2008). Because of different conservation needs, the two populations have been managed separately and assigned disparate conservation status (Southern Resident population: Endangered vs West Coast Transient population: Threatened, COSEWIC 2008).

Divergent demographic trends may also exist within genetically cohesive populations as a result of inter-group variations in feeding behaviors, ultimately influencing the dynamic of the population as a whole. In Crozet archipelago, killer whale matriline that depredate toothfish from longline fisheries showed higher survival, fecundity rates and overall growth than the non-interacting matriline (Tixier et al. 2015, Tixier et al. 2017). Artificial food provisioning from the longlines can greatly complement individuals' energy intake and enhance fitness and fecundity of depredating cetaceans. Similar observations of group-specific demographic trends were reported for killer whale groups interacting, or not, with drop-line tuna fisheries in the Strait of Gibraltar (Esteban et al. 2016). Hence, there is a critical need to account for heterogeneity in cultural foraging habits, even for killer whale groups occurring in sympatry.

iii- Population structure

Foraging specializations in killer whales appear to be culturally transmitted through generations within matrilineal social units (Riesch et al. 2012). Their apprenticeship may represent a substantial investment, especially for complex hunting techniques, further balanced by the benefits of enhanced prey searching and handling efficiencies through gained experience and familiarity with specific feeding behaviors (e.g. Guinet 1991). As a result, individuals dispersing to a social group that forages on a different prey type, using other specialized cooperative feeding behaviors, would likely risk a reduction in foraging efficiency and ultimately, a reduced fitness. This likely promotes social and reproductive isolation of the differentially resource-specialized groups instead. Cultural differentiation has been suggested as a possible mechanism shaping population genetic structure and facilitating the formation of separate ecotypes in sympatry (Hoelzel et al. 2007, Pilot et al. 2010, Moura et al. 2014, Foote et al. 2016).

Concurrent with ecological specializations, a mosaic of killer whale populations has been identified worldwide (Morin et al. 2010, Foote & Morin 2016). Both nuclear and mitochondrial DNA markers supported genetic differentiation of sympatric killer whale ecotypes in the eastern North Pacific (Hoelzel et al. 1998, Hoelzel et al. 2007, Morin et al. 2010, Pilot et al. 2010, Foote et al. 2016), in the Russian Far East (Filatova et al. 2015) and in the Antarctic (Leduc et al. 2008, Morin et al. 2010). Genetic structure of Northeastern Atlantic killer whale populations also coincides with apparent association to Atlantic herring, Atlantic mackerel and tuna stocks as primary food sources (Sigurjónsson et al. 1988, Similä et al. 1996, Guinet et al. 2007, Foote et al. 2011). Distribution of preferred prey, and inter-group differences in foraging strategies and ranging patterns were further identified as key factors in maintaining fine-scale genetic structure within ecotypes (eastern North Pacific: Parsons et al. 2013) and within a single population (Iceland: Tavares et al. 2018). Monitoring foraging habits may greatly assist in identifying demographically and evolutionary independent units, and thus should be considered an integral part of the conservation and management plans in killer whales.

iv- Resilience to changing habitats

A broad range of prey resources and habitats used within a population may imply a better buffering capacity against anthropogenic impacts and changing habitats as individuals/groups are likely to respond differently (Tinker et al. 2008). As such, identifying and preserving the

distinct ecological units i.e. ecological diversity within populations, is a fundamental key goal towards achieving a greater adaptive potential of marine species. In the light of changing marine ecosystems as promoted by global warming, assessing intra-population ecological variations also constitutes an important first step towards evaluating behavioral plasticity and resilience of killer whales to shifting habitats and prey resources (Kovacs & Lydersen 2008, Ferguson et al. 2010).

d- Knowledge gaps

A fascinating mosaic of ecological specializations in killer whales worldwide has emerged over many years of field studies. Long term ecological data have provided unequivocal support for foraging strategies shaping much of killer whales' biology and therefore triggering multiple implications. As a result, whilst management strategies are usually assigned at the population or regional level, it has been increasingly recognized that intra-population heterogeneity in foraging behaviors should be accounted for in the study of killer whales (COSEWIC 2008, Tixier et al. 2017). Although dietary patterns have been well documented for some populations, dietary studies are still warranted in most regions including across the North Atlantic. This thesis contributes to filling some of these knowledge gaps by providing new findings about dietary patterns of killer whales in Norway, and subsequent conservation implications.

II- OBJECTIVES

The overall objective of this thesis was to investigate the foraging ecology of killer whales in northern Norwegian fjords. The study was designed as a long-term multi-disciplinary approach to explore dietary variations, and to identify potential foraging specializations at the individual or group level, within this killer whale population. This work was based on continuous data collection maintained throughout the year, for six consecutive years in 2013-2018.

Sighting histories, predation records, behavioral observations and skin and blubber biopsy samples of photo-identified killer whales were collected. Individuals' occurrence patterns i.e. seasonality and site-fidelity, feeding habits and behavioral adaptations to target prey were investigated and compared over multiple years. Building upon killer whale identification datasets assembled by colleagues in Norway from 1987 through 1993 allowed for sighting and predation histories to span up to 30 years for some whales. Isotopic ratios and contaminant profiles were used as integrated dietary descriptors to infer on the persistence of foraging behaviors recorded from field observations and thus, to infer on individuals' long-term dietary habits and level of specialization. More specifically, the objectives of each paper were as follows.

Paper I investigated 1) killer whale occurrence at herring wintering grounds, newly located in coastal and offshore waters off Vesterålen from 2010 (Huse et al. 2010); and assessed 2) the scope and nature of mixed killer whales-humpback whales aggregations for feeding on herring schools. The study was based on observational data collected off Andøya during the three winter seasons 2013-2016.

Objectives of **Paper II** were 1) to explore seasonality and residency patterns of killer whales photo-identified in March-April off Andøya; 2) to identify the target prey seasonally eaten by these whales; and 3) to investigate the potential occurrence of behavioral adaptations to target prey, as an indication of seasonal foraging specialization. Data used in this study were collected over the five spring (March-April) seasons 2014-2018.

Aims of **Paper III** were to 1) review all available records of killer whale predation on pinnipeds in Norwegian coastal waters for the period 1988-2016; 2) to identify individual killer whales specifically involved in these predation events and explore consistency in feeding habits over

time; and 3) to uncover behavioral and social traits potentially adaptive of foraging upon mammalian prey as indicative of foraging specialization for these whales. Data used in this study were from multiple sources, including research field observations collected as part of this thesis in 2014-2016 and by colleagues in 1987-1993, as well as opportunistic imagery from wildlife enthusiasts leading to a dataset spanning from 1988 to 2016 for this study.

Dietary patterns identified in Papers I-III encouraged further research on specialization levels featuring integrated dietary descriptors, which led to **Paper IV**. In this study, nitrogen and carbon isotopic values in skin of photo-identified killer whales known from Papers I-III were used to 1) assess whether episodic predation records were reflective of consistent feeding behavior; and 2) to explore actual segregation in dietary habits among individuals/groups.

In line with Paper IV, contaminant profiles were used as additional dietary descriptors in **Paper V** to further investigate the validity of the dietary patterns identified in Papers I-IV. Specific objectives of this study were to 1) quantify and compare the PCB and mercury levels in blubber and skin, respectively of fish- and seal-eating killer whales; and 2) to explore the implications of dietary variations for management and conservation of killer whales in Norway.

Biopsy skin and blubber samples used in both Papers IV and V were collected in northern Norway in 2017-2018.

III- MATERIALS AND METHODS

A- Study area

The large majority of the data used in the studies covered in this thesis was collected around the northern tip of Andøya, centered on Andenes (69°19'N, 16°07'E), located in Vesterålen, northern Norway (Figure 1). Running to the east of Andøya, Andfjord is ~ 50 km long, reaches maximum depths of 500 m and is open in the north towards the Norwegian Sea. West of Andøya is the open sea where the Bleik Canyon, located 15 km offshore, stretches 40 to 50 km long and reaches 3,000 m depths. The canyon generates upwellings (Blindheim 1985), making the area particularly productive and rich in marine life (Skjoldal 2004). Andfjord was part of the wintering grounds of the Norwegian Spring Spawning herring from 2010 (Huse et al. 2010). However, over the course of this study, the herring stock has occupied diverse wintering areas (ICES 2018), encouraging seasonal fieldwork off Vengsøya (69°50'N, 18°32'E) in November 2015 and 2016, and off Skjervøya (70° 2'N, 20°58'E) in 2017 and 2018, both located in Troms County in northern Norway (Figure 1). Several interconnected fjords, characterized by a wide range of depths, were surveyed in both areas.

Figure 1. Locations of the four main study areas where fieldwork was conducted as part of this thesis in 2013-2018 (A, B, C). In addition, an historical dataset used in parts of this thesis was based on fieldwork conducted in region D in 1987-1993 (Bisther & Vongraven 1994, Bisther & Vongraven 2001).

B- Surveys

In 2013-2014, data collection was conducted from a whale-watch 36-foot-long rigid hull inflatable boat operated from Andenes. From 2015 onwards, surveys were conducted from a 20-foot-long aluminum vessel dedicated to the project. Opportunistic surveys were conducted in all seasons in response to sightings reported by locals, following land-based searches or directly as boat-based searches when killer whale occurrence peaked e.g. in the winter months of high herring abundance. Data collection was initiated upon encounter with a group of killer whales, defined as a grouping of individuals in apparent association and acting in a coordinated manner within the visual range of an observer (as per Baird & Dill 1995).

C- Data collection and processing

a- Photo-identification

Killer whales were approached and systematically photographed using a digital reflex Canon 5D camera and a mounted lens of 70-200 mm in focal length. Following photo-identification protocols developed by Bigg (1982), individuals were identified based on the size and shape of the dorsal fin, presence of nicks in the dorsal fin, and pigmentation and scarring patterns of the adjacent gray saddle patch (Figure 2). Killer whales in each encounter were catalogued using the best quality identification photograph available for each individual and assigning it a unique identification number.

b- Behavioral and predation records

Upon encounter with a group of killer whales, geographic location (GPS, Garmin) was recorded. Group size estimate, spatial cohesion of the group and behavioral state (Table 1) were systematically recorded at 5-min intervals for most encounters. Various clues including speed bursts, directional chases, fast and sharp turns, head-nodding movements and scavenging birds were used to identify feeding events of surfacing whales (Table 1, Ford et al. 1998), which were recorded *ad libitum* (Mann 1999). Collection of tissue fragments from prey killed and consumed by photo-identified killer whales was attempted whenever possible using a long-handled fine-mesh net, and for further molecular identification of prey species (Figure 2). An unmanned aircraft DJI Phantom III or IV carrying a camera was used from 2015 onwards to capture aerial images of prey capture events of photo-identified individuals, allowing for visual identification of target prey.

Table 1. Description of killer whale behavioral states as recorded throughout the study period (adapted from Ford 1989).

Behavioral state	Description
Foraging	Erratic swimming in varying directions, sudden lunges and high-speed swimming, presence of prey fragments and seabirds
Traveling	Consistent movement with all group members, in a tight group formation, following the same course at the same speed, while displaying synchronous dives and surfacings
Milling	Low speed movement with individuals engaged in frequent changes in direction leading to no directional movement by the group
Resting	Low level activity with individuals being stationary lined-up and floating motionless at the surface while displaying coordinated breathings
Socializing	High energy activity in which individuals may physically interact with one another and/or perform aerial displays such as leaps, tail slaps, and spy-hops (defined as vertical half-rise of the body out of the water)

c- Biopsy samples

Biopsy samples of photo-identified killer whales were collected using an ARTS darting system (Restech, Norway) and 25 x 9 mm or 40 x 9 mm stainless steel tips in 2017 and using an injection gun (Pneu-Dart Inc, USA) and 25 x 7 mm stainless steel tips in 2018. The region directly posterior to the dorsal fin of adult and sub-adult killer whales was the target area for sampling (Figure 2). Resulting skin and blubber samples were used for analysis of nitrogen and carbon stable isotopic values in skin. In addition, a total of 34 PCBs, 15 organochlorine pesticides, 18 brominated flame retardants, 16 hydroxylated metabolites of PCBs, and polybrominated diphenylethers (PBDEs) were determined in blubber samples of 31 killer whales. Mercury was determined in skin samples from 38 individuals.

d- Database

A catalogue of identified killer whales was held updated throughout the study period (Jourdain & Karoliussen 2018). In addition, available data for all catalogued killer whales were compiled into a database (Figure 2). Re-sightings of individuals over time and across locations were used to investigate individuals' patterns of seasonality and residency. Behavioral and predatory records allowed for identification of dietary patterns and comparison among individuals.

Dietary markers from biopsy samples were used to further assess the recurrence of dietary habits recorded from field observations.

e- Historical datasets

A photo-identification study conducted in November-March at herring spawning grounds off the coast of Møre in 1987-1993 resulted in a catalogue of 605 individual killer whales (Bisther & Vongraven 1994, Bisther & Vongraven 2001). This catalogue, along with individuals' sighting and behavioral histories, was available and used in the investigation of killer whales' long-term feeding habits.

Figure 2. Summary of data types collected throughout the study period (2013-2018) for photo-identified killer whales. All photographs by Eve Jourdain, Norwegian Orca Survey.

IV- RESULTS AND DISCUSSION

Findings are presented as lines of ecological evidence that support dietary variations and specializations, with results relating to several papers, rather than listed according to each paper.

A- Observational evidence

a- *Seasonality and target prey*

i- Herring foraging

Killer whale occurrence peaked from November through January at herring wintering grounds (Paper I) where 930 individuals were photo-identified from 2013-2018 (see Jourdain & Karoliussen 2018 for full identification catalogue). Behavioral observations and prey evidence confirmed foraging on herring during these winter months (Paper I), consistent with previous knowledge (Similä & Ugarte 1993, Similä et al. 1996, Similä 1997).

A compelling finding in this study was the appearance of humpback whales at current herring wintering grounds, never recorded in 17 years (1986 – 2003) of research at former herring wintering grounds (Paper I). In 2013-2016, 47 instances of killer whales and humpback whales feeding on common herring patches were recorded (Paper I). More specifically, killer whales that had cooperatively herded a herring school were often joined by one to three humpback whales lunging through the fish ball. Killer whales had initiated feeding in almost all (94%) mixed-species feeding aggregations, and often terminated feeding after humpback whale(s) had lunged (Paper I). These results suggested the observed feeding aggregations to be more beneficial to humpback whales and with possible costs to killer whales. Whilst interactions between killer whales and humpback whales reported worldwide have been largely agonistic (see Pitman et al. 2017 for a review), Paper I documents a new type of interaction between the two species.

The herring departed their wintering grounds in January-February, also leading to the sudden departure of killer whales (Paper I). Some killer whale groups may be following the herring migration (Lyrholm 1988, Similä et al. 1996, Similä & Stenersen 2004). However, as the herring undertakes its 800-1,000 km southward migration to spawning grounds, it undergoes substantial fat metabolism and fat content drops from ~ 18-22% when wintering to ~ 8-10% after spawning (Slotte 1999). Such changes in nutritional value, combined with lower densities

found at spawning and feeding grounds due to looser schools (Nøttestad et al. 1996), make the herring a less profitable prey throughout its different life stages (i.e. wintering, spawning, feeding; Holst et al. 2004). This could be promoting killer whale behavioral diversification and subsequent dietary variations, especially if other prey resources were available in a predictable way.

ii- Lumpfish foraging

From the 930 killer whales catalogued in the study area (Jourdain & Karoliussen 2018), 75 individuals returned to Andfjord each year in March-April between 2014 and 2018 (Paper II). Individuals remained in the area for up to 40 days, and foraging was recorded as the most frequently observed behavioral state (Paper II). Molecular or visual identification of prey remains revealed the lumpfish (*Cyclopterus lumpus*) as seasonal target prey for these whales when occurring in Andfjord in spring, representative of adult and subadult killer whales of both sexes (Paper II). Whilst most (81%) individuals were known herring-eaters from the winter months, these results brought the first conclusive evidence for killer whales in Norway to be supplementing their diet with, or to be switching to alternative prey resources, at times of lower herring availability/profitability.

The lumpfish is widely distributed throughout the Norwegian Sea (Holst 1993). While remaining at abyssal depths most of the year, it migrates to coastal breeding grounds where spawning occurs from April to July (Davenport 1985, Goulet et al. 1986, Mitamura et al. 2012). Fat content of lumpfish ranges from >8% in gravid females and up to 20% for males during spawning (Davenport & Kjørsvik 1986), equivalent to that of over-wintering herring (Slotte 1999). While lumpfish was not known as prey for killer whales in Norway prior to this study, it represented the entire stomach content of 30 killer whales harvested off West Greenland in 2003 (Laidre et al. 2006), confirming its nutritional qualities in killer whale diet. In Norway, seasonal specialization on temporarily abundant lumpfish prey may be highly profitable at times of herring scarcity. As lumpfish occupy spawning grounds in multiple regions along the Norwegian coast (Davenport 1985, Holst 1993), it could represent a significant prey resource for killer whales at the population level.

iii- Foraging on pinniped prey

Research datasets collected as part of this thesis in 2013-2016 and by colleagues in 1987-1993, as well as additional opportunistic imagery (and associated metadata) captured by wildlife

photographers in 2010-2016, documented a total of 23 events of killer whale predation on harbor seal or gray seal in 1988-2016 (Paper III). Photographic identification was possible for 14 of these predation events and included 13 killer whales from four social groups. Three of the killer whales observed feeding on seals in 1988-1992 off the Møre coast (Vongraven & Bisther 2014) were also observed in 2011-2016 off northern Norway, again preying on seals (Paper III, Figure 3). Given that there were 930 killer whales identified in the study area (Jourdain & Karoliussen 2018), and if all individuals were foraging on seals and had equal chance of being encountered, it would be unlikely to repeatedly observe the same individuals engaging in pinniped-foraging. Instead, these observations suggested that only a few killer whale groups may include pinniped prey in their diet.

Figure 3. **a)** Identification photograph of adult female killer whale *KI-03* collected off Orskjera, Møre as she was chasing a harbor seal on March 27, 1988 (in Vongraven & Bisther 2014); **b)** Identification photograph of *KI-03* collected off Andenes on June 27, 2015 as **c)** she was preying upon a harbor seal pup; **d)** Prey remains collected immediately after predation confirmed effective prey consumption (Paper III). This figure illustrates how field observations i.e. identification data and prey evidence were compiled in the investigation of individuals' dietary habits for this study. Photographs by (a) Dag Vongraven and (b), (c), (d) Richard Karoliussen.

A total of 37 encounters (including 14 with observed predation) with these seal-eating killer whales in the period 1988-2016 provided insights into specific occurrence patterns (Paper III). Seal-eating killer whales occurred in near-shore shallow areas, typical pinniped habitat (Bjørge 1991, Haug et al. 1994, Nilssen et al. 2010), with sightings concentrated in breeding areas for both gray and harbor seals along the Norwegian coast (Paper III). Timing of the observations,

with patterns consistent across years, further suggested killer whale movements reflective of local peaks in pinniped pupping and weaning (Paper III). For example, some of the seal-eating killer whales traveled from Vesterålen, where they were seen in May-September, northward to Troms in late October (Paper III), following the cline in timing of the gray seal pupping periods, which is delayed in the north (Haug et al. 1994).

Three of the seal-eating killer whales were observed feeding on herring at spawning grounds off Møre coast in 1991, suggesting a mixed diet including both fish and mammalian prey, or a prey switching (Vongraven & Bisther 2014). None of these whales were encountered at former herring wintering grounds in Tysfjord in 1986-2003 (Kuningas et al. 2014) or at herring wintering grounds in Andfjord-Troms in 2010-2016, where 584 and 656 individual killer whales were identified, respectively (Paper III). However, after Paper III was published, these same three individuals were encountered and observed feeding on herring at wintering grounds in Skjervøy in November 2018, confirming a mixed diet maintained over a lifetime for these whales.

iv- Other prey types

During the course of this study, field observations also documented killer whale predation on harbor porpoise, Atlantic salmon, Atlantic mackerel and black guillemot (*Cepphus grylle*). However, with only single records for these prey types, it was not useful for identification of individual-specific dietary patterns.

b- Variations in social organization

The size of killer whale groups varied throughout the study period, likely as a result of different foraging tactics used to efficiently search for and capture various target prey. Killer whales foraging on lumpfish in Andfjord in spring, also known as herring-eaters from winter observations, occurred in smaller groups in spring (median=8) than in winter (median=14) (Paper II, Figure 4). Smaller groups could be maximizing energetic gain when foraging upon a scattered prey resource, and could therefore be adaptive to foraging on solitary lumpfish (Davenport 1985) as opposed to foraging upon highly concentrated wintering herring (Nøttestad et al. 2002). Group size adjustment for fish prey concentration was also reported for killer whales feeding on mackerel in the summer months (Nøttestad et al. 2014).

Figure 4. Distribution of group size adopted by killer whales known as seal-eaters (Paper III) and fish-eaters as they seasonally forage on lumpfish in spring and herring in the winter (Paper II). Data distribution is displayed as second and third quartiles (box plot), median (horizontal crossing line) and mean (star symbol). Figure is based on data from Papers II and III.

Seal-eating killer whales, mainly encountered from May through October, also adopted a smaller group size (median=5) than their summer fish-eating counterparts (median=12) sharing the same waters (Paper III, Figure 4). Small groups were also reported for mammal-eating killer whales in the eastern North Pacific (Baird & Dill 1996), in Shetland, UK (Deecke et al. 2011), in Argentina (Iñiguez et al. 2005) and in the Crozet archipelago (Guinet 1992), suggesting a converging trait resulting from similar feeding ecology and common behavioral adaptations to efficiently hunt pinniped prey. A small group size was suggested to be most energetically efficient by ensuring stealthy and cooperative prey-searching and handling, whilst optimizing sharing of a single medium-sized pinniped prey (Baird & Dill 1996).

Investigating association patterns of seal-eating killer whales revealed a two-level social structure (Paper III, Figure 5). The first level consisted of highly stable basic units of three-to-five individuals. Regular, but temporary, associations of these units made up the second structural level (Paper III). By mostly keeping the basic units apart, seal-eating killer whales

may achieve a group size that enhances foraging efficiency, whilst temporary associations may allow for socializing and skill teaching (Baird & Dill 1996). Long-lasting social associations, the raw material for cultural transmission in killer whale societies (Riesch et al. 2012), may have facilitated the extreme persistence of dietary habits over a life-time for some of the seal-eating killer whales (Paper III).

Figure 5. Sociogram of seal-eating killer whales observed in at least four encounter days over the course of this study. Occurrence and thickness of links show association patterns between pairs of individuals, revealing permanent basic units that may temporary associate (Paper III).

c- Variations in foraging behavior

Herring-feeding killer whales were observed in tight group formations when cooperatively herding herring schools close to the surface (Paper I, Figure 6a). At lumpfish spawning grounds in spring, spatial cohesion of killer whale groups varied with behavior i.e. foraging, traveling, milling, resting, socializing, with individuals typically scattering over several hundred meters and up to 1 km when foraging (Paper II). Individuals moved in the same direction but surfaced independently and did not appear to cooperatively steer or capture lumpfish prey. Instead,

aerial drone imagery revealed individual prey capture (Figure 6b), with occasional prey sharing between adult females and their offspring(s) (Paper II). These observations, comparable to killer whales foraging on loose schools of salmonid prey (Ford & Ellis 2006, Vester & Hammerschmidt 2013, Ford & Ellis 2014), indicated behavioral adaptation to seasonally forage upon lumpfish.

Figure 6. Aerial photographs showing **a)** killer whales cooperatively herding a herring school off Andenes in January 2016 (Paper I) as opposed to **b)** individual foraging typically observed when killer whales fed upon lumpfish in spring in the same location (Paper II). These photographs highlight drone imagery as efficient means to study killer whale feeding behavior. Photographs by Richard Karoliussen.

Fish-eating killer whales remained in their core seasonal feeding areas for up to several weeks at a time, likely as a result of concentration of their migratory fish prey (Papers I, II). In contrast, seal-eating killer whales never stayed for long in a given location. Instead, they were constantly on the move, typically transiting from one seal colony to another (Paper III). Because apparently relying on stealth to capture pinnipeds, killer whales likely maximize their foraging efficiency by traveling elsewhere once their presence has been detected by potential prey. Similar contrasting occurrence patterns between fish and mammal-eating killer whales were initially highlighted in the eastern North Pacific (Ford & Ellis 2014, Ford 2019).

Seal-eating killer whales appeared to be hunting in near silence in the study area (Paper III). This observation was consistent with the acoustic behavior described for mammal-eating killer whales elsewhere, and likely represents an adaptation to avoid being detected by their acoustically sensitive marine mammal prey (Barrett-Lennard et al. 1996, Deecke et al. 2005, Deecke et al. 2011). In contrast, killer whales carousel-feeding on herring produce high rates of echolocation clicks and pulsed calls as means to both coordinate their attacks and promote schooling of their prey (Similä & Ugarte 1993, Simon et al. 2007a, Simon et al. 2007b).

Overall, killer whale groups encountered throughout the year in the study area differed considerably in their patterns of seasonal occurrence and behavior, suggesting adaptations to efficiently encounter and forage on the different target prey identified (Papers I-III). Most strikingly, behavioral features of seal-eating killer whales were in contrast with that of the sympatric fish-eaters (Papers I-III), but identical to that of mammal-eating killer whales elsewhere (Baird & Dill 1995, Baird & Dill 1996, Barrett-Lennard et al. 1996, Ford et al. 1998, Deecke et al. 2005, Deecke et al. 2011, Ford & Ellis 2014). Behavioral observations and predation records consistent over multiple years suggested inter-individual differences in dietary habits and indicated killer whale groups to be specialists of different prey types.

B- Evidence from dietary markers

As predatory history only provided a ‘snapshot’ of individuals’ feeding habits, inferences on the level of prey specialization were limited from Papers I-III. Integrated dietary markers (Papers IV, V) in skin and blubber biopsy samples of photo-identified killer whales, combined with time-series of field observations for these whales (Papers I-III), were used to further investigate consistency in dietary habits.

a- Stable isotopic values

The isotopic composition of a predator’s tissue reflects that of its prey resources in a predictable way (Newsome et al. 2010). Due to a greater retention of the heavier ^{15}N isotope than the lighter ^{14}N isotope in the production of nitrogenous waste, the nitrogen ratio of ^{15}N to ^{14}N ($\delta^{15}\text{N}$) shows a stepwise enrichment from prey to predator, and is therefore indicative of relative trophic position (DeNiro & Epstein 1978). The carbon ratio of ^{13}C to ^{12}C ($\delta^{13}\text{C}$) primarily reflects sources of carbon origin and thus, allows for discrimination between feeding areas in marine habitats, i.e. coastal/offshore vs benthic/pelagic (Hobson et al. 1994).

Skin $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values were measured for 38 killer whales, including individuals known as herring-, lumpfish- and seal-eaters from Papers I-III, biopsy sampled in all seasons in 2017-2018. Solely based on $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values, a Gaussian mixture model successfully assigned the 38 samples to three clusters (Paper IV, Figure 7) that largely aligned with predatory records available for these whales (Papers I-III).

Figure 7. Isospace of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values and niches as measured from killer whale skin samples (Paper IV). Solid lines represent the standard ellipses corrected for sample size and encompass 40% of the data, while dashed lines represent the convex hull area including the entire dataset for each cluster. Prey values are also plotted for comparison.

The low variance in isotopic values for both herring and lumpfish-eating killer whales led to a narrow isotopic niche for both groups and supported consistency in diet among individuals within each group (Paper IV, Figure 7). These results were in support of apparent seasonal specialization on herring in the winter months (Similä & Ugarte 1993, Similä et al. 1996, Similä 1997) and on spawning lumpfish in Andfjord in spring (Paper II). Whilst similar $\delta^{15}\text{N}$ values for herring and lumpfish-eaters indicated feeding at comparable trophic levels, the two dietary groups differed in $\delta^{13}\text{C}$ values (up to 2.5‰ difference, Paper IV, Figure 7). Such an increase in $\delta^{13}\text{C}$ values in spring could be due to temporary specialization on lumpfish prey, which is a semi-pelagic fish in a coastal habitat, as opposed to pelagic herring that spends most of its life offshore (Paper II).

Lumpfish-eaters (i.e. foraging upon lumpfish at time of sampling), sampled in April, were known to be feeding on herring in the months of November-January (Paper II). Segregation in isotopic niche between herring- and lumpfish-eaters (Paper IV), and differences in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values between lumpfish-eaters sampled in spring and herring-eaters sampled in winter,

suggested a possible diet switch from herring in November-February to lumpfish in March-April for these whales. However, inclusion of both herring and lumpfish, or even of other unidentified fish prey, cannot be ruled out.

Seal-eating killer whales showed significantly elevated $\delta^{15}\text{N}$ values regardless of sampling season (i.e. winter, spring, summer), compared to herring- and lumpfish-eaters (Paper IV, Figure 7). These $\delta^{15}\text{N}$ values confirmed predatory field observations of seal-eating killer whales (Paper III) to be reflective of consistent feeding habits, rather than opportunistic events. Distinct non-overlapping isotopic niches that characterized fish- and seal-eating killer whales further indicated dietary segregation into individuals/groups that include pinniped prey to their diet, and others that appear as fish-specialists (Paper IV).

Consistently elevated $\delta^{15}\text{N}$ values for seal-eating killer whales, even when sampled at seasonal herring and lumpfish grounds, supported these whales to be feeding on pinnipeds also when fish prey is abundant (Paper IV). However, seal-eating killer whales had lower mean $\delta^{15}\text{N}$ values than that expected under a pinniped-specialized diet ($\delta^{15}\text{N}_{\text{expected}} = 15.5\text{‰}$ vs $\delta^{15}\text{N}_{\text{measured}} = 12.6\text{‰}$) and also showed a wide isotopic niche (Figure 7), indicative of a diverse diet (Paper IV). These results supported a mixed diet with inclusion of both fish and mammalian prey for seal-eating killer whales, consistent with field records (Vongraven & Bisther 2014, Paper IV). Mixing models further estimated fish prey to represent an equal-to-secondary dietary component and group-specific variations in proportional contributions of the two prey groups (Paper IV).

b- Mercury levels

Although a natural element, mercury levels have increased in the environment due to anthropogenic activities, and is currently recognized as a global pollutant (Dietz et al. 2009). Methyl mercury has been shown to bioaccumulate and biomagnify through food webs, leading to highest concentrations in high trophic level marine species (Campbell et al. 2005). In this study, seal-eating killer whales had twice as high skin mercury levels as the two fish-eating groups herring-eaters and lumpfish-eaters (Paper V, Figure 8). Such a difference in concentrations was likely a result of higher mercury levels in pinnipeds that may constitute a substantial portion of the diet of seal-eating killer whales, compared to lower mercury levels expected in low trophic level fish species (Campbell et al. 2005). Hence, these results were in

support of the predation records in Papers I-III, and the isotopic segregation into fish and mammal-eating groups (Figure 7) in Paper IV, indicating that they are reflective of true long-term foraging habits and specializations for these individuals.

Figure 8. Total mercury levels in skin (µg/g dry weight) in seal-eating and fish-eating killer whales biopsy sampled in 2017-2018 in all seasons in northern Norway (Paper V). Horizontal line represents the median, whiskers represent the lower (first) and upper (fourth) quartiles, dots mark individual killer whales and points outside the whiskers/box are outliers.

c- Organohalogen contaminants (OHC)

Concentration patterns of organohalogen contaminants (OHC) such as PCBs, DTTs and PBDEs constitute valuable chemical tracers for differentiating killer whale prey-specialists, owing to differences in trophic level and relative OHC concentrations of their respective prey (Herman et al. 2005, Krahn et al. 2007, Krahn et al. 2008). Because adult reproductive females transfer their contaminant burdens to their calf, they may not be representative of OHCs levels of the population (Ross et al. 2000, Ylitalo et al. 2001, Haraguchi et al. 2009, Krahn et al. 2009, Pedro et al. 2017). Adult males, which never offload their contaminants and thus become increasingly contaminated as they grow older, appear better suited to fully apprehend differences in OHC concentrations among prey-specialists.

In this thesis, herring- and lumpfish-eating killer whales had similar levels of OHCs in their blubber (Paper V), in support of foraging at similar trophic level fish prey (Paper IV). However,

seal-eating killer whales showed substantially higher OHC levels than fish-eaters (Paper V). More specifically, adult male seal-eaters had three times higher Σ PCB levels (Figure 9), and twice as high Σ DDTs, Σ CHLs and Σ PBDEs levels (Paper V, Table 2) in their blubber than fish-eating killer whales frequenting the same waters.

Atlantic herring only have low concentrations of contaminants i.e. Σ PCBs: 17.4-25.5 ng/g wet weight (Wolkers et al. 2007) compared to harbor seal i.e. Σ PCBs: 2,246-11,320 ng/g lipid weight (Ruus et al. 1999) and gray seal i.e. Σ PCBs: 2,972-10,103 ng/g lipid weight (Ruus et al. 1999) in Norway. Therefore, higher OHCs levels measured in seal-eating killer whales (Paper V) are likely a result of increased exposure from biomagnification due to foraging upon highly contaminated pinniped prey (Ross et al. 2000, Ylitalo et al. 2001, Herman et al. 2005, Krahn et al. 2007) compared to lumpfish- and herring-eaters, which do not appear to include mammalian prey in their diet (Paper IV). Thus, these results confirmed pinniped-foraging as a long-term dietary habit for killer whales identified as seal-eaters from Papers III and IV.

Figure 9. Lipid-normalized Σ PCB levels ($\mu\text{g/g}$) in blubber of seal- and fish-eating killer whales sampled in 2017-2018 in all seasons in northern Norway (Paper V). Horizontal line represents the median, whiskers represent the lower (first) and upper (fourth) quartiles, dots mark individual killer whales and points outside the

PCBs and DDTs contributed ~86% of the total OHC burden for all three dietary groups (lumpfish-, herring- and seal-eaters), but relative abundance of the differently chlorinated PCB congeners differed markedly between seal- and fish-eating killer whales (Paper V). The seal-eating individuals had higher relative proportions of hexachloro- and heptachloro-congeners, likely due to increased presence of these congeners in marine mammals relative to fish (Hoekstra et al. 2003). These results were consistent with PCB patterns found for marine mammal- vs fish-eating killer whales of the eastern North Pacific (Herman et al. 2005) and of Greenland (Pedro et al. 2017), further supporting pinniped-foraging as a persistent dietary habit for seal-eating killer whales in Norway (Papers III, IV).

There was a five-fold difference between median Σ PCB levels in adult male fish- and seal-eating killer whales sampled for this thesis (Paper V). This is a much smaller distinction than the 20-fold difference in median Σ PCB levels measured between the (adult males) sympatric fish-eating (Northern Residents: 10.5 $\mu\text{g/g}$ lipid weight) and mammal-eating (Transients: 206 $\mu\text{g/g}$ lipid weight) ecotypes of the eastern North Pacific (Buckman et al. 2011), known to be strict prey specialists (Ford et al. 1998, Ford & Ellis 2006, Ford & Ellis 2014, Ford 2019). This is likely attributable to the fact that seal-eating killer whales in Norway adopt a mixed diet including both fish and mammal prey (Vongraven & Bisther 2014, Paper IV), and therefore, consume less of the most contaminated pinniped prey than the exclusive mammal-eating killer whales in the Pacific.

Table 2. Lipid-normalized concentrations ($\mu\text{g/g}$) of organohalogen contaminants in seal- and fish-eating adult male killer whales biopsy sampled in 2017-2018 in northern Norway (Paper V).

OHC group		Seal-eaters	Fish-eaters
ΣDDTs	mean $\pm$ SD	30.9 $\pm$ 16.7	18.5 $\pm$ 24.0
	range	(14.4–54.2)	(3.04–100)
ΣCHLs	mean $\pm$ SD	10.3 $\pm$ 3.75	5.12 $\pm$ 5.54
	range	(5.54–14.4)	(1.52–23.4)
ΣPBDEs	mean $\pm$ SD	1.61 $\pm$ 0.39	0.630 $\pm$ 0.440
	range	(1.27–2.13)	(0.258–1.79)

C- Possible driving forces of dietary diversity

Historical records of multiple prey types (Collett 1912, Jonsgård & Lyshoel 1970, Christensen 1982) suggest that inter-individual/group dietary variations may have always characterized this population. Alternatively, behavioral diversification could have intensified as a result of one or a combination of several of the following factors.

a- Collapse of the Norwegian Spring Spawning herring

With improvement of the fishing methods and over-exploitation throughout the 1950s, the Norwegian Spring Spawning stock of herring had collapsed by the end of the 1960s, reducing the spawning biomass from 10 million tons (Mt) in 1957 to 0.28 Mt in 1968 (Dragesund & Ulltang 1975, Dragesund et al. 1980). A thinner fatness layer was recorded for killer whales caught in 1951-1968, suggesting a negative effect of herring depletion on the condition of herring-feeding killer whales (Christensen 1982). Therefore, as herring became scarce, the realized benefit of specializing on herring prey may have been lost (Svanbäck & Bolnick 2005), forcing killer whales to diversify their diet as an adaptive response (Tinker et al. 2008, Randa et al. 2009).

b- Past culling

The past culling of killer whales in Norway, as a management action to preserve the remaining herring, may have played an additional role in the niche diversification process. In 1969-1970, 368 killer whales were culled over a 100 x 50 km area within a few weeks (Øien 1988). Entire groups were never culled, and catch records showed a general lack of small individuals and a skewed sex ratio of 2M:1F (Vongraven & Bisther 1995). Due to differential roles played by key members in cetacean societies and cultural transmission of prey specializations within matrilineal social groups in killer whales, feeding strategies may have been lost and foraging success compromised if key members were culled (Williams & Lusseau 2006, Brent et al. 2015, Natrass et al. 2019). This could have promoted the appearance of new feeding behaviors in this population (Papers II, III).

c- Shifts in prey populations

After suffering a severe historical decline, both the harbor seal (Bjørge et al. 2007, Nilssen et al. 2010) and the gray seal (Øigård et al. 2012, but see Nilssen 2017) have shown population growth in the recent decades. The gray seal has also notably expanded its range, with breeding

areas now found from Trøndelag through Finnmark counties (Haug et al. 1994, Bjørge et al. 2002). Such increased availability of lipid-rich pinnipeds residents in Norwegian coastal waters has undoubtedly promoted seals as a valuable food source to killer whales in this region, especially after the herring stock collapsed in the late 1960s, and likely facilitated dietary diversification for some groups (Paper III).

Major shifts in prey resources owing to warming climate and ocean in the North Atlantic over the last three decades (ICES 2018) may also have created new foraging opportunities and promoted dietary diversification in killer whales in Norway. Owing to fluctuations in oceanographic conditions, lumpfish abundance in the Barents Sea, with fish spawning along the coast of northern Norway, varied considerably over the last decades. Specifically, the biomass was lowest in the 1980s, increased throughout the 1990s, and was highest in the 2000s (Eriksen et al. 2014). It remains unknown whether killer whales in Norway have only recently started feeding on lumpfish (Paper II). Regardless, the increase in lumpfish biomass again since 2012 in the Norwegian Sea, reaching 150 million fish in 2019 (IMR 2020), may have promoted or intensified this fish as food source for killer whales in this region.

The northeastern Atlantic mackerel has expanded its distributional range northwards and has spent a longer time feeding in the Norwegian Sea in the summer (Utne et al. 2012, Nøttestad et al. 2015b). Increasing from 2 Mt in 2007 to 9 Mt in 2014 (Nøttestad et al. 2015b), the mackerel has become more abundant than the Norwegian Spring Spawning herring, which decreased from 12 Mt in 2009 to 5 Mt in 2014 (ICES 2013). Whilst killer whale presence has been increasing in the summer months in the Norwegian Sea (2009-2012, Nøttestad et al. 2015a), field observations further confirmed selective foraging upon mackerel in July and August (Nøttestad et al. 2014).

d- Ecological interactions

As forager density increases on a common pool of resources, prey availability decreases. Competition may therefore promote individuals' behavioral diversification to efficiently forage upon new exclusive prey types, reduce competition and ultimately enhance individuals' fitness (Svanbäck & Bolnick 2005, Svanbäck & Bolnick 2007, Tinker et al. 2008, Kobler et al. 2009). As such, fluctuations in prey populations, owing to inter- and intraspecific competition or changing environmental conditions, may play a substantial role in mediating temporal

dynamics of dietary variation and specializations (Svanbäck & Bolnick 2005, Araújo et al. 2011). The recent appearance of humpback whales at herring wintering grounds illustrates how changing ecological conditions, e.g. changes in herring distribution, may lead to variations in forager occurrence and competition forces (Paper I). The dynamics and complexity of ecological interactions that occur within and among species likely play a substantial role in shaping individuals' and populations' patterns of resource use and feeding behaviors.

D- Implications of dietary variations/specializations

a- Variations in individuals' fitness

The Arctic Monitoring and Assessment Program concluded that killer whales in Norway were at a lower risk from PCB-mediated health effects than other Arctic killer whale populations (AMAP 2018). This was based upon Σ PCB blubber concentrations from male herring-feeding killer whales sampled in 2002 (geometric mean= 27 $\mu\text{g/g}$ lw in Wolkers et al. 2007). However, Σ PCB levels in male seal-eating killer whales reported in this thesis (geometric mean= 46 $\mu\text{g/g}$ lipid weight, Paper V, Figure 9), were nearly double these levels and were more comparable to that measured in mammal-eating killer whales harvested in Greenland (Σ PCB arithmetic mean=65.1 mg/kg in Pedro et al. 2017), for which PCB-mediated effects were the highest (AMAP 2018).

All seal-eating killer whales in Paper V had Σ PCB blubber concentrations that exceeded thresholds for high-to-highest risks of immune and hormone system health effects (AMAP 2018), compared to only 46% of the fish-eaters (Paper V). The Σ PCB levels of seal-eating killer whales furthermore, exceeded the Σ PCB threshold of 41 $\mu\text{g/g}$ lipid weight, at which there could be risks of reproductive impairment with occlusions in the uterus (Jepson et al. 2016). As such, mean Σ PCB levels of female seal-eaters (48 $\mu\text{g/g}$ lipid weight) and female fish-eaters (3.6 $\mu\text{g/g}$ lipid weight) in Paper V could be translated to approximately 45% and 80% of calves surviving, respectively (Hall et al. 2018). Seal-eating killer whales were also exposed to higher health risks owing to higher mercury levels than fish-eaters (Paper V). Overall, these results indicate that variations in survival and breeding performance among killer whale groups adopting different diets could be expected, and therefore that ecological groups should be managed separately in Norway.

b- Population dynamics

A recent study modelled killer whale population growth around the world, accounting for population-specific PCB blubber concentrations and mediated health effects (Desforges et al. 2018). Using PCB levels from Wolkers et al. (2007), Norwegian killer whales were assigned to the least contaminated group. The results predicted little-to-no effect of PCB pollution on population growth, and even a 50% growth within the next 100 years. However, higher PCB burdens in seal-eating compared to fish-eating killer whales in Norway (Paper V) suggest higher PCB-mediated health effects, possibly leading to lower survival and reproduction rates, for seal-eaters in Norway. A decline of these whales could have further impact on the growth of the population as a whole.

c- Population structure

Strong genetic differentiation was reported between the two sympatric ecotypes that specialize on either fish or marine mammal prey in the Northeast Pacific (Hoelzel et al. 1998, Hoelzel et al. 2007, Morin et al. 2010, Parsons et al. 2013, Moura et al. 2014, Foote et al. 2016). In the North Atlantic, there is also evidence for a link between genetic divergence and ecological specialization on a large-scale (Foote et al. 2011) but patterns are less clear in the different regions, largely due to a lack of dietary studies so far. Results in the present thesis revealed ecological divergence between fish- and seal-eating killer whales occurring in sympatry in Norway (Papers I-IV), suggesting behavioral mechanisms that could lead to possible fine-scale genetic differentiation as shown in Icelandic waters (Tavares et al. 2018).

Chemical tracers used in this study revealed ecological differentiation between killer whale groups i.e. fish- vs seal-eaters (Papers IV, V), based on consistent differences in trophic ecology, and leading to different levels of exposure and vulnerability to OHC concentrations (Paper V). In light of these results, an integrated approach coupling ecological and molecular markers seems critical to accurately identify both genetic and ecological independent management units.

d- Resilience to changing habitats

A wide range of prey types for killer whales in Norway (Similä et al. 1996, Vester & Hammerschmidt 2013, Nøttestad et al. 2014, Vongraven & Bisther 2014, Cosentino 2015, Papers I-III) suggest this population could be somewhat buffered against changing marine

ecosystems, as individuals/groups are likely to respond to shifting prey populations in different ways. However, highly specialized groups could be severely impacted by changes in the abundance of their preferred prey (Ward et al. 2009, Ford et al. 2010). Consequently, the proportion of prey-specialists, and the scope of dependence on preferred prey vs foraging flexibility may greatly determine the resilience of this killer whale population to rapidly changing marine ecosystems in the Arctic.

CLOSING REMARKS

This thesis has showed that killer whales in Norway constitute a generalist population characterized by inter-individual dietary variations and a mixture of individual specialists of various prey types. Results have revealed that fish-specialists may respond to seasonal fluctuations in prey availability by switching between specialized foraging behaviors for efficient use of alternative fish prey resources. Other individuals seem to specialize, to various extents, on pinniped prey over a lifetime. However, there was no indication of killer whales feeding exclusively on marine mammal prey in Norway. These findings, along with previous knowledge of a mixed diet for killer whales in Greenland (Pedro et al. 2017, Bourque et al. 2018) and Iceland (Samarra et al. 2017) suggest that killer whales in the North Atlantic may not be as ecologically segregated as the eastern North Pacific prey-specialized killer whale communities.

As large sized predators with high daily caloric requirements, killer whales could be significantly impacting Norwegian pinniped (and other prey) populations through top-down effects (Williams et al. 2004). This thesis provides key information to assess killer whale predatory impact in this region.

Pollution levels and associated health risks differed substantially between fish- and seal-eating killer whales. A higher OHC burden in seal-eaters translated to a higher risk for health effects, with potential consequences at the population level. Such new knowledge points out the urgent need for a new assessment of the status of this killer whale population, accounting for ecological differentiation among social groups and possibly, considering fish- and seal-eaters as distinct ecological management units.

This study supports, once again, individuality in feeding behavior as an apparent characteristic of the species and underlines the importance of accounting for intra-population heterogeneity in dietary habits for effective management and conservation. Hence, the findings underline the importance of collecting individual-focused data, on the long-term, to study killer whales. This work, built upon six years of data collection on individual killer whales in northern Norway, has a lot of potential for further studies.

REFERENCES

- Afik D, Karasov WH (1995) The trade-offs between digestion rate and efficiency in warblers and their ecological implications. *Ecology* 76: 2247-2257. doi:10.2307/1941699.
- AMAP (2018) Amap assessment 2018: Biological effects of contaminants on arctic wildlife and fish: Arctic monitoring and assessment programme (amap).
- Araújo MS, Bolnick DI, Layman CA (2011) The ecological causes of individual specialisation. *Ecology Letters* 14: 948-958. doi:10.1111/j.1461-0248.2011.01662.x.
- Baird RW, Abrams PA, Dill LM (1992) Possible indirect interactions between transient and resident killer whales - implications for the evolution of foraging specializations in the genus *orcinus*. *Oecologia* 89: 125-132. doi:10.1007/bf00319024.
- Baird RW, Dill LM (1995) Occurrence and behaviour of transient killer whales: Seasonal and pod-specific variability, foraging behaviour, and prey handling. *Canadian Journal of Zoology* 73: 1300-1311. doi:10.1139/z95-154.
- Baird RW, Dill LM (1996) Ecological and social determinants of group size in transient killer whales. *Behavioral Ecology* 7: 408-416. doi:10.1093/beheco/7.4.408.
- Baird RW, McSweeney DJ, Bane C, Barlow J, Salden DR, La'Ren KA, LeDuc RG, Webster DL (2006) Killer whales in hawaiian waters: Information on population identity and feeding habits1. *Pacific Science* 60: 523-530. doi:10.1353/psc.2006.0024.
- Baird RW, Whitehead H (2000) Social organization of mammal-eating killer whales: Group stability and dispersal patterns. *Canadian Journal of Zoology* 78: 2096-2105. doi:10.1139/z00-155.
- Barrett-Lennard LG, Ford JKB, Heise KA (1996) The mixed blessing of echolocation: Differences in sonar use by fish-eating and mammal-eating killer whales. *Animal Behaviour* 51: 553-565. doi:10.1006/anbe.1996.0059.
- Best P, Meyer M, Lockyer C (2010) Killer whales in south african waters—a review of their biology. *African Journal of Marine Science* 32: 171-186. doi:10.2989/1814232X.2010.501544.
- Bigg M (1982) An assessment of killer whale *orcinus orca* stocks off vancouver island, british columbia. *Report of the International Whaling Commission* 32: 655-666.
- Bigg MA, Ellis GM, Ford JKB, Balcomb KC (1987) *Killer whales: A study of their identification, genealogy and natural history in british columbia and washington state*. Phantom Press, Nanaimo.
- Bigg MA, Ford JKB, Ellis GM. (1985) Two sympatric forms of killer whales off british columbia and washington *Sixth Biennial Conference on the Biology of Marine Mammals*, Vancouver, Canada.
- Bisther A, Vongraven D (1994) *Killer whales off the mid-norwegian coast*. Final Report, NFFR-PROJECT 4001-319.016, Amsterdam.

- Bisther A, Vongraven D (1995) Studies of social ecology of norwegian killer whales (*orcinus orca*). In: Blix AS, Walløe L, Ulltang Ø (eds) *Whales, seals, fish and man*, 169-176. Elsevier Science.
- Bisther A, Vongraven D (2001) Killer whales feeding on both mammals and fish: A transient, resident or opportunistic type? *Proceedings of the Poster presentation at the 14th Biennial Conference on the Biology of Marine Mammals*.
- Bjørge A (1991) Status of the harbor seal *phoca vitulina* in norway. *Biological Conservation* 58: 229-238. doi:10.1016/0006-3207(91)90121-o.
- Bjørge A, Øien N, Fagerheim KA (2007) Abundance of harbour seals (*phoca vitulina*) in norway based on aerial surveys and photographic documentation of hauled-out seals during the moulting season, 1996 to 1999. *Aquatic Mammals* 33: 269-275. doi:10.1578/AM.33.3.2007.269
- Bjørge A, Øien N, Hartvedt S, Bøthun G, Bekkby T (2002) Dispersal and bycatch mortality in gray, halichoerus grypus, and harbor, phoca vitulina, seals tagged at the norwegian coast. *Marine Mammal Science* 18: 963-976. doi:10.1111/j.1748-7692.2002.tb01085.x.
- Blindheim J (1985) Ecological features of the norwegian sea, Sixth conference of the Comité Arctique International, Fairbanks, Alaska: 366-401.
- Bolanos-Jimenez J, Mignucci-Giannoni AA, Blumenthal J, Bogomolni A, Casas JJ, Henriquez A *et al.* (2014) Distribution, feeding habits and morphology of killer whales *orcinus orca* in the caribbean sea. *Mammal Review* 44: 177-189. doi:10.1111/mam.12021.
- Bolnick DI, Svanbäck R, Fordyce JA, Yang LH, Davis JM, Darrin Hulsey C, Forister ML (2003) The ecology of individuals: Incidence and implications of individual specialization. *The American Naturalist* 161: 1-28. doi:10.1086/343878.
- Bourque J, Dietz R, Sonne C, St Leger J, Iverson S, Rosing-Asvid A, Hansen M, McKinney MA (2018) Feeding habits of a new arctic predator: Insight from full-depth blubber fatty acid signatures of greenland, faroe islands, denmark, and managed-care killer whales *orcinus orca*. *Marine Ecology Progress Series* 603: 1-12. doi:10.3354/meps12723
- Brent LNJ, Franks DW, Foster EA, Balcomb KC, Cant MA, Croft DP (2015) Ecological knowledge, leadership, and the evolution of menopause in killer whales. *Current Biology* 25: 746-750. doi:10.1016/j.cub.2015.01.037.
- Buckman AH, Veldhoen N, Ellis G, Ford JKB, Helbing CC, Ross PS (2011) Pcb-associated changes in mrna expression in killer whales (*orcinus orca*) from the ne pacific ocean. *Environmental Science & Technology* 45: 10194-10202. doi:10.1021/es201541j.
- Burdin AM, Hoyt E, Sato H, Tarasyan K, Filatova OA (2004) *Resident and transient-type killer whales, orcinus orca, in southeast kamchatka, russia*. Paper SC/56/SM15 presented to the IWC Scientific Committee.
- Campbell LM, Norstrom RJ, Hobson KA, Muir DCG, Backus S, Fisk AT (2005) Mercury and other trace elements in a pelagic arctic marine food web (northwater polynya, baffin bay). *Science of the Total Environment* 351-352: 247-263. doi:10.1016/j.scitotenv.2005.02.043.

- Capella JJ, Abramson JZ, Vilina YA, Gibbons J (2014) Observations of killer whales (*orcinus orca*) in the fjords of chilean patagonia. *Polar Biology* 37: 1533-1539. doi:10.1007/s00300-014-1535-5.
- Ceia FR, Ramos JA (2015) Individual specialization in the foraging and feeding strategies of seabirds: A review. *Marine Biology* 162: 1923-1938. doi:10.1007/s00227-015-2735-4.
- Christensen I (1982) Killer whales in norwegian coastal waters. *Report of the International Whaling Commission* 32: 633-641.
- Christensen I (1988) Distribution, movements and abundance of killer whales (*orcinus orca*) in norwegian coastal waters, 1982-1987, based on questionnaire surveys. *Rit Fiskideildar* 11: 79-88.
- Christensen I, Øritsland T (1982) *Census of killer whales on the coast of norway in february 1982*. Paper SC/34/SM4 presented to the IWC Scientific Committee.
- Collett R (1912) *Orcinus orca* (lin.) 1758. In: R. Collett e (ed) *Norges pattedyr: Cetacea. Aschehoug, christiania*, 707-716.
- Cosentino AM (2015) First record of norwegian killer whales attacking and feeding on a harbour porpoise. *Marine Biodiversity Records* 8. doi:10.1017/S1755267215000895.
- COSEWIC (2008) *Cosewic assessment and update status report on the killer whale orcinus orca, southern resident population, northern resident population, west coast transient population, offshore population and northwest atlantic / eastern arctic population, in canada. Committee on the status of endangered wildlife in canada. Ottawa*.
- Cunningham P, Hughes R (1984) Learning of predatory skills by shorecrabs *carcinus maenas* feeding on mussels and dogwhelks. *Marine Ecology Progress Series* 16: 1-26.
- Dahlheim ME, Schulman-Janiger A, Black N, Ternullo R, Ellifrit D, Balcomb KC (2008) Eastern temperate north pacific offshore killer whales (*orcinus orca*): Occurrence, movements, and insights into feeding ecology. *Marine Mammal Science* 24: 719-729. doi:10.1111/j.1748-7692.2008.00206.x.
- Davenport J (1985) *Synopsis of biological data on the lumpsucker, cyclopterus lumpus (linnaeus, 1758)*. FAO Fisheries, Rome, Italy: Food and Agriculture Organization of the United Nations.
- Davenport J, Kjørsvik E (1986) Buoyancy in the lumpsucker cyclopterus lumpus. *Journal of the Marine Biological Association of the United Kingdom* 66: 159-174. doi:10.1017/S0025315400039722.
- Deecke VB, Ford JKB, Slater PJB (2005) The vocal behaviour of mammal-eating killer whales: Communicating with costly calls. *Animal Behaviour* 69: 395-405. doi:10.1016/j.anbehav.2004.04.014.
- Deecke VB, Nykanen M, Foote AD, Janik VM (2011) Vocal behaviour and feeding ecology of killer whales *orcinus orca* around shetland, uk. *Aquatic Biology* 13: 79-88. doi:10.3354/ab00353.
- DeNiro MJ, Epstein S (1978) Influence of diet on the distribution of carbon isotopes in animals. *Geochimica et Cosmochimica Acta* 42: 495-506. doi:10.1016/0016-7037(78)90199-0.

- Desforges J-P, Hall A, McConnell B, Rosing-Asvid A, Barber JL, Brownlow A *et al.* (2018) Predicting global killer whale population collapse from pcb pollution. *Science* 361: 1373-1376. doi:10.1126/science.aat1953.
- Dietz R, Outridge PM, Hobson KA (2009) Anthropogenic contributions to mercury levels in present-day arctic animals—a review. *Science of the Total Environment* 407: 6120-6131. doi:10.1016/j.scitotenv.2009.08.036.
- Dragesund O, Hamre J, Ulltang Ø (1980) Biology and population dynamics of the norwegian spring-spawning herring. *Rapports et Procès-verbaux des Réunions Conseil International pour l'Exploration de la Mer* 177: 43-71.
- Dragesund O, Ulltang Ø (1975) Stock size fluctuations and rate of exploitation of the norwegian spring-spawning herring, 1950-1974. *FiskDir. Skr. Ser. HavUnders.* 16: 315-337.
- Eriksen E, Durif CMF, Prozorkevich D (2014) Lumpfish (*cyclopterus lumpus*) in the barents sea: Development of biomass and abundance indices, and spatial distribution. *Ices Journal of Marine Science* 71: 2398-2402. doi:10.1093/icesjms/fsu059.
- Esteban R, Verborgh P, Gauffier P, Giménez J, Guinet C, de Stephanis R (2016) Dynamics of killer whale, bluefin tuna and human fisheries in the strait of gibraltar. *Biological Conservation* 194: 31-38. doi:10.1016/j.biocon.2015.11.031.
- Estes J, Riedman M, Staedler M, Tinker M, Lyon B (2003) Individual variation in prey selection by sea otters: Patterns, causes and implications. *Journal of Animal Ecology* 72: 144-155. doi:10.1046/j.1365-2656.2003.00690.x.
- Ferguson SH, Higdon JW, Chmelnitsky EG (2010) The rise of killer whales as a major arctic predator. In: Ferguson SH, Loseto LL, Mallory ML (eds) *A little less arctic: Top predators in the world's largest northern inland sea, hudson bay*, 117-136. Springer Netherlands, Dordrecht.
- Filatova OA, Borisova EA, Shpak OV, Meschersky IG, Tiunov AV, Goncharov AA, Fedutin ID, Burdin AM (2015) Reproductively isolated ecotypes of killer whales *orcinus orca* in the seas of the russian far east. *Biology Bulletin* 42: 674-681. doi:10.1134/s1062359015070043.
- Foote A, Vijay N, Avila-Arcos M, Baird R, Durban J, Morin P *et al.* (2016) Genome-culture coevolution promotes rapid divergence in the killer whale. *Nature Communications*. doi:10.1101/040295.
- Foote AD, Morin PA (2016) Genome-wide snp data suggest complex ancestry of sympatric north pacific killer whale ecotypes. *Heredity*: 1-10. doi:10.1038/hdy.2016.54.
- Foote AD, Newton J, Piortney SB, Willerslev E, Gilbert MTP (2009) Ecological, morphological and genetic divergence of sympatric north atlantic killer whale populations. *Molecular Ecology* 18: 5207-5217. doi:10.1111/j.1365-294X.2009.04407.x.
- Foote AD, Similä T, Vikingsson G, Stevick PT (2010) Movement, site fidelity and connectivity in a top marine predator, the killer whale. *Evolutionary Ecology* 24: 803-814. doi:10.1007/s10682-009-9337-x.

- Foote AD, Vester H, Vikingsson GA, Newton J (2012) Dietary variation within and between populations of northeast atlantic killer whales, *orcinus orca*, inferred from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analyses. *Marine Mammal Science* 28: 472-485. doi:10.1111/j.1748-7692.2012.00563.x.
- Foote AD, Vilstrup JT, de Stephanis R, Verborgh P, Nielsen SCA, Deaville R *et al.* (2011) Genetic differentiation among north atlantic killer whale populations. *Molecular Ecology* 20: 629-641. doi:10.1111/j.1365-294X.2010.04957.x.
- Ford JK (2019) Killer whales: Behavior, social organization, and ecology of the oceans' apex predators. In: Würsig B (ed) *Ethology and behavioral ecology of odontocetes*, 239-259. Springer Nature, Switzerland.
- Ford JKB (1989) Acoustic behavior of resident killer whales (*orcinus orca*) off vancouver island, british columbia. *Canadian Journal of Zoology* 67: 727-745.
- Ford JKB (2009) Killer whales *orcinus orca*. In: Perrin WF, Würsig B, Thewissen JGM (eds) *The encyclopedia of marine mammals*, 650-657. Academic Press, San Diego, USA.
- Ford JKB, Ellis GM (2006) Selective foraging by fish-eating killer whales *orcinus orca* in british columbia. *Marine Ecology Progress Series* 316: 185-199. doi:10.3354/meps316185.
- Ford JKB, Ellis GM (2014) You are what you eat: Foraging specializations and their influence on the social organization and behavior of killer whales. In: Yamagiwa J, Karczmarski L (eds) *Primates and cetaceans: Field research and conservation of complex mammalian societies*, 75-98. Springer, Tokyo, Japan.
- Ford JKB, Ellis GM, Barrett-Lennard LG, Morton AB, Palm RS, Balcomb KC (1998) Dietary specialization in two sympatric populations of killer whales (*orcinus orca*) in coastal british columbia and adjacent waters. *Canadian Journal of Zoology* 76: 1456-1471. doi:10.1139/cjz-76-8-1456.
- Ford JKB, Ellis GM, Matkin CO, Wetklo MH, Barrett-Lennard LG, Withler RE (2011) Shark predation and tooth wear in a population of northeastern pacific killer whales. *Aquatic Biology* 11: 213-224. doi:10.3354/ab00307.
- Ford JKB, Ellis GM, Olesiuk PF, Balcomb KC (2010) Linking killer whale survival and prey abundance: Food limitation in the oceans' apex predator? *Biology Letters* 6: 139-142. doi:10.1098/rsbl.2009.0468.
- Goulet D, Green JM, Shears TH (1986) Courtship, spawning, and parental care behavior of the lumpfish, *cyclopterus lumpus* l., in newfoundland. *Canadian Journal of Zoology* 64: 1320-1325.
- Grieg JA (1906) Nogle notiser fra et spaekhuggerstaeng ved bildöströmmen i januar 1904. *Bergens Museums Årbok*: 1-28.
- Guinet C (1991) Intentional stranding apprenticeship and social play in killer whales *orcinus orca*. *Canadian Journal of Zoology* 69: 2712-2716.
- Guinet C (1992) Predation behavior of killer whales (*orcinus orca*) around crozet islands. *Canadian Journal of Zoology* 70: 1656-1667.

- Guinet C, Domenici P, de Stephanis R, Barrett-Lennard L, Ford JKB, Verborgh P (2007) Killer whale predation on bluefin tuna: Exploring the hypothesis of the endurance-exhaustion technique. *Marine Ecology Progress Series* 347: 111-119. doi:10.3354/meps07035.
- Hall AJ, McConnell BJ, Schwacke LH, Ylitalo GM, Williams R, Rowles TK (2018) Predicting the effects of polychlorinated biphenyls on cetacean populations through impacts on immunity and calf survival. *Environmental Pollution* 233: 407-418. doi:10.1016/j.envpol.2017.10.074.
- Haraguchi K, Hisamichi Y, Endo T (2009) Accumulation and mother-to-calf transfer of anthropogenic and natural organohalogenes in killer whales (*orcinus orca*) stranded on the pacific coast of japan. *Science of the Total Environment* 407: 2853-2859. doi:10.1016/j.scitotenv.2009.01.003.
- Haug T, Henriksen G, Kondakov A, Mishin V, Nilssen KT, Røv N (1994) The status of grey seals halichoerus grypus in north norway and on the murman coast, russia. *Biological Conservation* 70: 59-67.
- Herman DP, Burrows DG, Wade PR, Durban JW, Matkin CO, LeDuc RG, Barrett-Lennard LG, Krahn MM (2005) Feeding ecology of eastern north pacific killer whales *orcinus orca* from fatty acid, stable isotope, and organochlorine analyses of blubber biopsies. *Marine Ecology Progress Series* 302: 275-291. doi:10.3354/meps302275.
- Hobson KA, Piatt JF, Pitocchelli J (1994) Using stable isotopes to determine seabird trophic relationships. *Journal of Animal Ecology* 63: 786-798. doi:10.2307/5256.
- Hoekstra PF, O'Hara TM, Fisk AT, Borgå K, Solomon KR, Muir DCG (2003) Trophic transfer of persistent organochlorine contaminants (ocs) within an arctic marine food web from the southern beaufort–chukchi seas. *Environmental Pollution* 124: 509-522. doi:10.1016/S0269-7491(02)00482-7.
- Hoelzel AR (1998) Genetic structure of cetacean populations in sympatry, parapatry, and mixed assemblages: Implications for conservation policy. *Journal of Heredity* 89: 451-458. doi:10.1093/jhered/89.5.451.
- Hoelzel AR, Dahlheim M, Stern SJ (1998) Low genetic variation among killer whales (*orcinus orca*) in the eastern north pacific and genetic differentiation between foraging specialists. *Journal of Heredity* 89: 121-128. doi:10.1093/jhered/89.2.121.
- Hoelzel AR, Dorsey EM, Stern J (1989) The foraging specializations of individual minke whales. *Animal Behaviour* 38: 786-794.
- Hoelzel AR, Hey J, Dahlheim ME, Nicholson C, Burkanov V, Black N (2007) Evolution of population structure in a highly social top predator, the killer whale. *Molecular Biology and Evolution* 24: 1407-1415. doi:10.1093/molbev/msm063.
- Holbrook SJ, Schmitt RJ (1992) Causes and consequences of dietary specialization in surfperches: Patch choice and intraspecific competition. *Ecology* 73: 402-412. doi:10.2307/1940748.
- Holst JC (1993) Observations on the distribution of lumpsucker (*cyclopterus lumpus*, l.) in the norwegian sea. *Fisheries Research* 17: 369-372. doi:10.1016/0165-7836(93)90136-u.

- Holst JC, Røttingen I, Melle W (2004) The herring. In: HR S (ed) *The norwegian sea ecosystem*, 203-226. Tapir Academic Press, Trondheim.
- Hori M (1993) Frequency-dependent natural selection in the handedness of scale-eating cichlid fish. *Science* 260: 216-219. doi:10.1126/science.260.5105.216.
- Huse G, Fernö A, Holst JC (2010) Establishment of new wintering areas in herring co-occurs with peaks in the 'first time/repeat spawner' ratio. *Marine Ecology Progress Series* 409: 189-198. doi:10.3354/meps08620.
- ICES (2013) *Report of the working group on widely distributed stocks (wgwide)*, 27 august - 2 september 2013, ices headquarters. Copenhagen, Denmark.
- ICES (2018) *Report of the working group on widely distributed stocks (wgwide)*, 28 august- 3 september 2018. Torshavn, Faroe Islands.
- IMR (2020) *Rognkjeks/rognkall*. <https://www.hi.no/hi/temasider/arter/rognkjeks-rognkall>
- Iñiguez MA, Tossenberger VP, Gasparrou C. (2005) Socioecology of killer whales (*orcinus orca*) in northern patagonia, argentina, 9 pp, Unpublished paper to the IWC Scientific Committee, Ulsan, Korea, June 2005. (SC/57/SM5).
- Jepson PD, Deaville R, Barber JL, Aguilar A, Borrell A, Murphy S *et al.* (2016) Pcb pollution continues to impact populations of orcas and other dolphins in european waters. *Scientific Reports* 6: 18573. doi:10.1038/srep18573.
- Jonsgård Å (1968) A note on the attacking behaviour of the killer whale (*orcinus orca*). Nor. Hvalfangstmus. *Norsk Hvalfangst-Tidende* 57: 84-85.
- Jonsgård Å, Lyshoel PB (1970) A contribution to the knowledge of the biology of the killer whale *orcinus orca* l. *Nytt Magasin for Zoologi* 18: 41-48.
- Jourdain E, Karoliussen R. (2018) Identification catalogue of norwegian killer whales: 2007-2018. Figshare, doi: 10.6084/m9.figshare.4205226.
- Kobler A, Klefoth T, Mehner T, Arlinghaus R (2009) Coexistence of behavioural types in an aquatic top predator: A response to resource limitation? *Oecologia* 161: 837-847. doi:10.1007/s00442-009-1415-9.
- Kopps AM, Ackermann CY, Sherwin WB, Allen SJ, Bejder L, Krutzen M (2014) Cultural transmission of tool use combined with habitat specializations leads to fine-scale genetic structure in bottlenose dolphins. *Proceedings of the Royal Society B-Biological Sciences* 281: 1782. doi:10.1098/rspb.2013.3245.
- Kovacs KM, Lydersen C (2008) Climate change impacts on seals and whales in the north atlantic and adjacent shelf areas. *Science Progress* 91: 117-150. doi:10.3184/003685008X324010.
- Krahn MM, Bradley Hanson M, Schorr GS, Emmons CK, Burrows DG, Bolton JL, Baird RW, Ylitalo GM (2009) Effects of age, sex and reproductive status on persistent organic pollutant concentrations in "southern resident" killer whales. *Marine Pollution Bulletin* 58: 1522-1529. doi:10.1016/j.marpolbul.2009.05.014.

- Krahn MM, Herman DP, Matkin CO, Durban JW, Barrett-Lennard L, Burrows DG *et al.* (2007) Use of chemical tracers in assessing the diet and foraging regions of eastern north pacific killer whales. *Marine Environmental Research* 63: 91-114. doi:10.1016/j.marenvres.2006.07.002.
- Krahn MM, Pitman RL, Burrows DG, Herman DP, Pearce RW (2008) Use of chemical tracers to assess diet and persistent organic pollutants in antarctic type c killer whales. *Marine Mammal Science* 24: 643-663. doi:10.1111/j.1748-7692.2008.00213.x.
- Kuningas S, Similä T, Hammond PS (2014) Population size, survival and reproductive rates of northern norwegian killer whales (*orcinus orca*) in 1986-2003. *Journal of the Marine Biological Association of the United Kingdom* 94: 1277-1291. doi:10.1017/s0025315413000933.
- Laidre KL, Heide-Jorgensen MP, Orr JR (2006) Reactions of narwhals, *monodon monoceros*, to killer whale, *orcinus orca*, attacks in the eastern canadian arctic. *Canadian Field-Naturalist* 120: 457-465. doi:10.22621/cfn.v120i4.355.
- Leduc RG, Robertson KM, Pitman RL (2008) Mitochondrial sequence divergence among antarctic killer whale ecotypes is consistent with multiple species. *Biology Letters* 4: 426-429. doi:10.1098/rsbl.2008.0168.
- Louis M, Fontaine MC, Spitz J, Schlund E, Dabin W, Deaville R *et al.* (2014) Ecological opportunities and specializations shaped genetic divergence in a highly mobile marine top predator. *Proceedings of the Royal Society B: Biological Sciences* 281: 20141558. doi:10.1098/rspb.2014.1558.
- Lyrholm T. (1984) Observations of killer whales, *orcinus orca*, during a photodocumentation survey off the møre coast, norway, 7 pp, Unpublished paper to the IWC Scientific Committee, 1984 (SC/36/SM21).
- Lyrholm T. (1987) Photoidentification of killer whales, *orcinus orca*, off the coast of norway - preliminary results 1983-1986, 14 pp, Unpublished paper to the IWC Scientific Committee, 1987 (SC/39/SM19).
- Lyrholm T (1988) Photoidentification of individual killer whales, *orcinus orca*, off the coast of norway, 1983-1986. *Rit Fiskideildar* 11: 89-94.
- Mann J (1999) Behavioral sampling methods for cetaceans: A review and critique. *Marine Mammal Science* 15: 102-122. doi:10.1111/j.1748-7692.1999.tb00784.x.
- Matkin CO, Barrett-Lennard LG, Yurk H, Ellifrit D, Trites AW (2007) Ecotypic variation and predatory behavior among killer whales (*orcinus orca*) off the eastern aleutian islands, alaska. *Fishery Bulletin* 105: 74-87.
- Mitamura H, Thorstad EB, Uglem I, Bjorn PA, Okland F, Naesje TF, Dempster T, Arai N (2012) Movements of lump sucker females in a northern norwegian fjord during the spawning season. *Environmental Biology of Fishes* 93: 475-481. doi:10.1007/s10641-011-9942-8.
- Morin PA, Archer FI, Foote AD, Vilstrup J, Allen EE, Wade P *et al.* (2010) Complete mitochondrial genome phylogeographic analysis of killer whales (*orcinus orca*) indicates multiple species. *Genome Research* 20: 908-916. doi:10.1101/gr.102954.109.

- Moura AE, Kenny JG, Chaudhuri R, Hughes MA, Welch AJ, Reisinger RR *et al.* (2014) Population genomics of the killer whale indicates ecotype evolution in sympatry involving both selection and drift. *Molecular Ecology* 23: 5179-5192. doi:10.1111/mec.12929.
- Nattrass S, Croft DP, Ellis S, Cant MA, Weiss MN, Wright BM *et al.* (2019) Postreproductive killer whale grandmothers improve the survival of their grandoffspring 116: 26669-26673. doi:10.1073/pnas.1903844116.
- Newsome SD, Clementz MT, Koch PL (2010) Using stable isotope biogeochemistry to study marine mammal ecology. *Marine Mammal Science* 26: 509-572. doi:10.1111/j.1748-7692.2009.00354.x.
- Nichol LM, Shackleton DM (1996) Seasonal movements and foraging behavior of northern resident killer whales (*orcinus orca*) in relation to the inshore distribution of salmon (*oncorhynchus* spp.) in british columbia. *Canadian Journal of Zoology* 74: 983-991. doi:10.1139/z96-111.
- Nilssen KT, Skavberg NE, Poltermann M, Haug T, Härkönen T, Henriksen G (2010) Status of harbour seals (*phoca vitulina*) in mainland norway. *NAMMCO Scientific Publications* 8: 61-69. doi:10.7557/3.2672.
- Nilssen KTB, A. (2017) *Selhavert og steinkobbe*. Havforskningsrapporten 2017. Fisken og Havet, særnummer 1–2017.
- Nøttestad L, Aksland M, Beltestad A, Fernö A, Johannessen A, Arve Misund O (1996) Schooling dynamics of norwegian spring spawning herring (*clupea harengus* l.) in a coastal spawning area. *Sarsia* 80: 277-284. doi:10.1080/00364827.1996.10413601.
- Nøttestad L, Ferno A, Axelsen BE (2002) Digging in the deep: Killer whales' advanced hunting tactic. *Polar Biology* 25: 939-941. doi:10.1007/s00300-002-0437-0.
- Nøttestad L, Krafft BA, Anthonypillai V, Bernasconi M, Langård L, Mørk HL, Fernö A (2015a) Recent changes in distribution and relative abundance of cetaceans in the norwegian sea and their relationship with potential prey. *Frontiers in Ecology and Evolution* 2. doi:10.3389/fevo.2014.00083.
- Nøttestad L, Sivle LD, Krafft BA, Langard L, Anthonypillai V, Bernasconi M, Langoy H, Ferno A (2014) Prey selection of offshore killer whales *orcinus orca* in the northeast atlantic in late summer: Spatial associations with mackerel. *Marine Ecology Progress Series* 499: 275-283. doi:10.3354/meps10638.
- Nøttestad L, Utne KR, Óskarsson GJ, Jónsson SP, Jacobsen JA, Tangen Ø *et al.* (2015b) Quantifying changes in abundance, biomass, and spatial distribution of northeast atlantic mackerel (*scomber scombrus*) in the nordic seas from 2007 to 2014 73: 359-373. doi:10.1093/icesjms/fsv218.
- Øien N (1988) The distribution of killer whales (*orcinus orca*) in the north atlantic based on norwegian catches, 1938-1981, and incidental sightings, 1967-1987. *Rit Fiskideildar* 11: 65-78.
- Øigård TA, Frie AK, Nilssen KT, Hammill MO (2012) Modelling the abundance of grey seals (*halichoerus grypus*) along the norwegian coast. *Ices Journal of Marine Science* 69: 1436-1447. doi:10.1093/icesjms/fss103.

- Parsons KM, Durban JW, Burdin AM, Burkanov VN, Pitman RL, Barlow J *et al.* (2013) Geographic patterns of genetic differentiation among killer whales in the northern north pacific. *Journal of Heredity* 104: 737-754. doi:10.1093/jhered/est037.
- Pedro S, Boba C, Dietz R, Sonne C, Rosing-Asvid A, Hansen M, Provatas A, McKinney MA (2017) Blubber-depth distribution and bioaccumulation of pcbs and organochlorine pesticides in arctic-invading killer whales. *Science of the Total Environment* 601–602: 237-246. doi:10.1016/j.scitotenv.2017.05.193.
- Pilot M, Dahlheim ME, Hoelzel AR (2010) Social cohesion among kin, gene flow without dispersal and the evolution of population genetic structure in the killer whale (*orcinus orca*). *Journal of Evolutionary Biology* 23: 20-31. doi:10.1111/j.1420-9101.2009.01887.x.
- Pitman RL, Deecke VB, Gabriele CM, Srinivasan M, Black N, Denkinger J *et al.* (2017) Humpback whales interfering when mammal-eating killer whales attack other species: Mobbing behavior and interspecific altruism? *Marine Mammal Science* 33: 7-58. doi:10.1111/mms.12343.
- Pitman RL, Durban JW (2012) Cooperative hunting behavior, prey selectivity and prey handling by pack ice killer whales (*orcinus orca*), type b, in antarctic peninsula waters. *Marine Mammal Science* 28: 16-36. doi:10.1111/j.1748-7692.2010.00453.x.
- Pitman RL, Durban JW, Greenfelder M, Guinet C, Jorgensen M, Olson PA, Plana J, Tixier P, Towers JR (2011) Observations of a distinctive morphotype of killer whale (*orcinus orca*), type d, from subantarctic waters. *Polar Biology* 34: 303-306. doi:10.1007/s00300-010-0871-3.
- Pitman RL, Ensor P (2003) Three forms of killer whales in antarctic waters. *Journal of Cetacean Research and Management* 5: 131-139.
- Pitman RL, Fearnbach H, Durban JW (2018) Abundance and population status of ross sea killer whales (*orcinus orca*, type c) in mcmurdo sound, antarctica: Evidence for impact by commercial fishing? *Polar Biology*. doi:10.1007/s00300-017-2239-4.
- Price T (1987) Diet variation in a population of darwin's finches. *Ecology* 68: 1015-1028. doi:10.2307/1938373.
- Randa LA, Cooper DM, Meserve PL, Yunker JA (2009) Prey switching of sympatric canids in response to variable prey abundance. *Journal of Mammalogy* 90: 594-603. doi:10.1644/08-MAMM-A-092R1.1.
- Reisinger RR, Grocke DR, Lubcker N, McClymont EL, Hoelzel AR, de Bruyn PJN (2016) Variation in the diet of killer whales *orcinus orca* at marion island, southern ocean. *Marine Ecology Progress Series* 549: 263-274. doi:10.3354/meps11676.
- Rendell L, Whitehead H (2001) Culture in whales and dolphins. *Behavioral and Brain Sciences* 24: 309-382. doi:10.1017/s0140525x0100396x.
- Riesch R, Barrett-Lennard LG, Ellis GM, Ford JKB, Deecke VB (2012) Cultural traditions and the evolution of reproductive isolation: Ecological speciation in killer whales? *Biological Journal of the Linnean Society* 106: 1-17. doi:10.1111/j.1095-8312.2012.01872.x.

- Ross PS, Ellis GM, Ikonomou MG, Barrett-Lennard LG, Addison RF (2000) High pcb concentrations in free-ranging pacific killer whales, *orcinus orca*: Effects of age, sex and dietary preference. *Marine Pollution Bulletin* 40: 504-515.
- Ruus A, Ugland KI, Espeland O, Skaare JU (1999) Organochlorine contaminants in a local marine food chain from jarfjord, northern norway. *Marine Environmental Research* 48: 131-146. doi:10.1016/S0141-1136(99)00037-9.
- Samarra FIP, Miller PJO (2015) Prey-induced behavioural plasticity of herring-eating killer whales. *Marine Biology* 162: 809-821. doi:10.1007/s00227-015-2626-8.
- Samarra FIP, Vighi M, Aguilar A, Vikingsson GA (2017) Intra-population variation in isotopic niche in herring-eating killer whales off iceland. *Marine Ecology Progress Series* 564: 199-210. doi:10.3354/meps11998.
- Sargeant BL, Mann J, Berggren P, Krutzen M (2005) Specialization and development of beach hunting, a rare foraging behavior, by wild bottlenose dolphins (*tursiops* sp.). *Canadian Journal of Zoology* 83: 1400-1410. doi:10.1139/Z05-136.
- Saulitis E, Matkin C, Barrett-Lennard L, Heise K, Ellis G (2000) Foraging strategies of sympatric killer whale (*orcinus orca*) populations in prince william sound, alaska. *Marine Mammal Science* 16: 94-109. doi:10.1111/j.1748-7692.2000.tb00906.x.
- Sigurjónsson J, Lyrholm T, Leatherwood S, Jonsson E, Vikingsson G (1988) Photoidentification of killer whales, *orcinus orca*, off iceland, 1981 through 1986. *Rit Fiskideildar* 11: 99-114.
- Similä T (1997) Sonar observations of killer whales (*orcinus orca*) feeding on herring schools. *Aquatic Mammals* 23: 119-126.
- Similä T, Holst JC, Christensen I (1996) Occurrence and diet of killer whales in northern norway: Seasonal patterns relative to the distribution and abundance of norwegian spring-spawning herring. *Canadian Journal of Fisheries and Aquatic Sciences* 53: 769-779.
- Similä T, Stenersen J (2004) *Norwegian killer whales*. Tringa forlag, Oslo, Norway.
- Similä T, Ugarte F (1993) Surface and underwater observations of cooperatively feeding killer whales in northern norway. *Canadian Journal of Zoology* 71: 1494-1499.
- Simon M, McGregor PK, Ugarte F (2007a) The relationship between the acoustic behaviour and surface activity of killer whales (*orcinus orca*) that feed on herring (*clupea harengus*). *Acta Ethologica* 10: 47-53. doi:10.1007/s10211-007-0029-7.
- Simon M, Wahlberg M, Miller LA (2007b) Echolocation clicks from killer whales (*orcinus orca*) feeding on herring (*clupea harengus*). *Journal of the Acoustical Society of America* 121: 749-752. doi:10.1121/1.2404922.
- Skjoldal HR (2004) *The norwegian sea ecosystem*. Tapir Academic Press, Trondheim.
- Slotte A (1999) Differential utilisation of energy during wintering and spawning migration in norwegian spring-spawning herring. *Journal of Fish Biology* 54: 338-355. doi:10.1111/j.1095-8649.1999.tb00834.x.

- Svanbäck R, Bolnick DI (2005) Intraspecific competition affects the strength of individual specialization: An optimal diet theory method. *Evolutionary Ecology Research* 7: 993-1012.
- Svanbäck R, Bolnick DI (2007) Intraspecific competition drives increased resource use diversity within a natural population. *Proceedings of the Royal Society B: Biological Sciences* 274: 839-844. doi:10.1098/rspb.2006.0198.
- Tavares SB, Samarra FIP, Pascoal S, Graves JA, Miller PJO (2018) Killer whales (*orcinus orca*) in iceland show weak genetic structure among diverse isotopic signatures and observed movement patterns. *Ecology and Evolution* 0. doi:10.1002/ece3.4646.
- Tinker MT, Bentall G, Estes JA (2008) Food limitation leads to behavioral diversification and dietary specialization in sea otters. *Proceedings of the National Academy of Sciences* 105: 560. doi:10.1073/pnas.0709263105.
- Tixier P, Authier M, Gasco N, Guinet C (2015) Influence of artificial food provisioning from fisheries on killer whale reproductive output. *Animal Conservation* 18: 207-218. doi:10.1111/acv.12161.
- Tixier P, Barbraud C, Pardo D, Gasco N, Duhamel G, Guinet C (2017) Demographic consequences of fisheries interaction within a killer whale (*orcinus orca*) population. *Marine Biology* 164: 170. doi:10.1007/s00227-017-3195-9.
- Utne KR, Huse G, Ottersen G, Holst JC, Zabavnikov V, Jacobsen JA, Óskarsson GJ, Nøttestad LJMBR (2012) Horizontal distribution and overlap of planktivorous fish stocks in the norwegian sea during summers 1995–2006 8: 420-441. doi:10.1080/17451000.2011.640937.
- Van Opzeeland IC, Corkeron PJ, Leyssen T, Similä T, Van Parijs SM (2005) Acoustic behaviour of norwegian killer whales, *orcinus orca*, during carousel and seiner foraging on spring-spawning herring. *Aquatic Mammals* 31: 110-119. doi:10.1578/AM.31.1.2005.110.
- Vester H, Hammerschmidt K (2013) First record of killer whales (*orcinus orca*) feeding on atlantic salmon (*salmo salar*) in northern norway suggest a multi-prey feeding type. *Marine Biodiversity Records* 6: 1-5. doi:10.1017/S1755267212001030.
- Vongraven D, Bisther A (1995) Possible effects of previous catch on the present population of norwegian killer whales (*orcinus orca*). In: Blix AS, Walløe L, Ulltang Ø (eds) *Whales, seals, fish and man*, 177-179. Elsevier Science, Tromsø, Norway.
- Vongraven D, Bisther A (2014) Prey switching by killer whales in the north-east atlantic: Observational evidence and experimental insights. *Journal of the Marine Biological Association of the United Kingdom* 94: 1357-1365. doi:10.1017/s0025315413001707.
- Warburton K, Retif S, Hume D (1998) Generalists as sequential specialists: Diets and prey switching in juvenile silver perch. *Environmental Biology of Fishes* 51: 445-454. doi:10.1023/A:1007489100809.
- Ward EJ, Holmes EE, Balcomb KC (2009) Quantifying the effects of prey abundance on killer whale reproduction. *Journal of Applied Ecology* 46: 632-640. doi:10.1111/j.1365-2664.2009.01647.x.

- Weinrich MT, Schilling MR, Belt CR (1992) Evidence for acquisition of a novel feeding behavior - lobtail feeding in humpback whales *megaptera novaeangliae*. *Animal Behaviour* 44: 1059-1072. doi:10.1016/s0003-3472(05)80318-5.
- Weir CR, Collins T, Carvalho I, Rosenbaum HC (2010) Killer whales (*orcinus orca*) in angolan and gulf of guinea waters, tropical west africa. *Journal of the Marine Biological Association of the United Kingdom* 90: 1601-1611. doi:10.1017/s002531541000072x.
- Whitehead H, Ford JKB (2018) Consequences of culturally-driven ecological specialization: Killer whales and beyond. *Journal of Theoretical Biology* 456: 279-294. doi:10.1016/j.jtbi.2018.08.015.
- Williams R, Lusseau D (2006) A killer whale social network is vulnerable to targeted removals. *Biology Letters* 2: 497-500. doi:10.1098/rsbl.2006.0510.
- Williams TM, Estes JA, Doak DF, Springer AM (2004) Killer appetites: Assessing the role of predators in ecological communities. *Ecology* 85: 3373-3384. doi:10.1890/03-0696.
- Wolkers H, Corkeron PJ, van Parijs SM, Similä T, van Bavel B (2007) Accumulation and transfer of contaminants in killer whales (*orcinus orca*) from norway: Indications for contaminant metabolism. *Environmental Toxicology and Chemistry* 26: 1582-1590.
- Ylitalo GM, Matkin CO, Buzitis J, Krahn MM, Jones LL, Rowles T, Stein JE (2001) Influence of life-history parameters on organochlorine concentrations in free-ranging killer whales (*orcinus orca*) from prince william sound, ak. *Science of the Total Environment* 281: 183-203. doi:10.1016/S0048-9697(01)00846-4.

AUTHORSHIP CONTRIBUTIONS

Eve Jourdain is the sole author of the summary part of the thesis “Dietary variations and specializations in killer whales (*Orcinus orca*) in Norway”.

Contributions to Papers I-V are as follows:

Paper I:

Eve Jourdain: First author and corresponding author

EJ conceptualized the study. **EJ** acquired the funds. **EJ** collected field observations. **EJ** and DV analyzed the data. **EJ** wrote the paper. DV commented on earlier drafts of the manuscripts.

Paper II:

Eve Jourdain: First author and corresponding author

EJ conceptualized the study. **EJ** acquired the funds. **EJ**, RK, JDV and SEZ collected field observations and tissue samples. CT conducted the laboratory analyses. **EJ** analyzed the data. **EJ** wrote the paper. CT commented on earlier drafts of the manuscript.

Paper III:

Eve Jourdain: First author and corresponding author

EJ conceptualized the study. **EJ** acquired the funds. **EJ**, DV, AB and RK collected data. **EJ** analyzed the data. **EJ** and DV wrote the paper. AB commented on earlier drafts of the manuscript.

Paper IV:

Eve Jourdain: First author and corresponding author

EJ, CA, RK and DV conceptualized the study. **EJ** acquired the funds. RK, **EJ** and CA collected samples/data. CA conducted the laboratory analyses. **EJ** and CA analyzed the data. **EJ** wrote the paper. CA, DV, AR and KB commented on earlier drafts of the manuscript.

Paper V:

Eve Jourdain: Second author and co-supervisor on CA's MSc project

CA, **EJ**, AR, JLL, and KB designed the study. **EJ** acquired the funds. RK, **EJ** and CA collected samples/data. CA and JLL conducted the laboratory analyses. CA analyzed the data. CA wrote the paper. **EJ**, AR, JLL and KB commented on earlier drafts of the manuscript.

PAPER I

Humpback whale (*Megaptera novaeangliae*) and killer whale (*Orcinus orca*)
feeding aggregations for foraging on herring (*Clupea harengus*)
in northern Norway.

Jourdain, E., and Vongraven, D. (2017).

Mammalian Biology, 86: 27-32.

doi: 10.1016/j.mambio.2017.03.006.

Original investigation

Humpback whale (*Megaptera novaeangliae*) and killer whale (*Orcinus orca*) feeding aggregations for foraging on herring (*Clupea harengus*) in Northern NorwayE. Jourdain^{a,*}, D. Vongraven^b^a Norwegian Orca Survey – Haugesveien 30, N-8480 Andenes, Norway^b Norwegian Polar Institute – Fram Center, N-9296 Tromsø, Norway

ARTICLE INFO

Article history:

Received 16 July 2016

Accepted 30 March 2017

Handled by Nelika K. Hughes

Available online 1 April 2017

Keywords:

Humpback whale

Megaptera novaeangliae

Killer whale

Orcinus orca

Feeding behavior

Mixed-species aggregation

Interaction

Interspecific

ABSTRACT

Aggregations of predators on food patches have been documented for both terrestrial and marine animals. Here, we documented for the first time, and investigated, non-predatory aggregations occurring between humpback whales (*Megaptera novaeangliae*) and killer whales (*Orcinus orca*) while feeding on wintering Norwegian spring spawning herring (*Clupea harengus*) in Andfjord, northern Norway. Observational data were collected during 109 opportunistic surveys through three seasons 2013–2016. Killer whales were observed feeding on 59 occasions, with one to three humpback whales involved in 47 of these feeding events (79.7%), and there was an increased probability of finding feeding humpback whales when feeding killer whales also were observed. With killer whales identified as the initiating species in 94.4% of the feeding aggregations for which the first species was known, and with humpback whales joining and feeding on the fish ball afterwards, we suggest that humpback whales may benefit more from these aggregations than the opposite.

© 2017 Deutsche Gesellschaft für Säugetierkunde. Published by Elsevier GmbH. All rights reserved.

Introduction

The ocean is a dynamic environment where irregularly allocated resources define the distribution of organisms. Food patches, defined as dense concentrations of homogenous resources, often promote spatial and temporal aggregations of predators through two main facilitative mechanisms. First, a concentration of foragers on food patches may subsequently enhance discovery of unpredictable prey by neighbors in search of food, a passive process called local enhancement (Bruno et al., 2003; Grunbaum and Veit, 2003; Poysa, 1992; Silverman et al., 2004). Second, predators may actively associate with hetero-specifics if easier prey capture and increased foraging efficiency are provided (Bearzi, 2006; Berce, 2010; Goyert et al., 2014; Lett et al., 2014). Higher predator densities may also encourage schooling behavior of fish prey, promoting possible mutual benefits among foragers (Lett et al., 2014; Similä and Ugarte, 1993). Predator interactions remain advantageous as

long as prey density outweighs competition, and otherwise become negative (competitive) when resources diminish (Poysa, 1992).

Mixed-species feeding aggregations in the marine environment have been reported in many species including seabirds, pinnipeds, cetaceans and fish (e.g. Au and Pitman, 1988; Bearzi et al., 2006; Brager, 1998; Camphuysen and Webb, 1999; Goyert et al., 2014; Pitman and Ballance, 1992; Vaughn et al., 2007). As a common factor of all studies, some species appear to act as catalysts, initiating the feeding process, with other species arriving at a later stage when prey have been found or concentrated and predation eased, and are called opportunists (Camphuysen and Webb, 1999).

Off the coast of northern Norway, killer whales (*Orcinus orca*) occur throughout the year with a peak of observations during October–January when the Norwegian spring spawning (NSS) herring (*Clupea harengus*) over-winters in coastal waters (Similä et al., 1996). From 1986 through 2005, when herring wintered in the Tysfjord – Vestfjord system, studies highlighted killer whales as efficient herring predators with specialized feeding techniques including carousel feeding (Similä, 1997b; Similä and Ugarte, 1993). In carousel feeding, killer whales herd the fish into a tight ball close to the surface and capture individual herring after debilitating them with underwater tail slaps (Domenici et al., 2000; Similä, 1997a; Similä and Ugarte, 1993; Simon et al., 2006). Killer

* Corresponding author.

E-mail addresses: evejourdain@yahoo.fr (E. Jourdain), dag.vongraven@npolar.no (D. Vongraven).

whales were shown to adjust group size in relation to herring school size (Nøttestad et al., 2002), but feeding aggregations with other cetacean predators were never reported over the course of these long-term studies.

Humpback whales (*Megaptera novaeangliae*) use the Norwegian Sea as a migration corridor between their summer feeding areas, mainly located around Jan Mayen and Bear Island, and their low latitude breeding ground (Øien, 2007). Since 2010, the wintering ground of the NSS herring has been located in off-shore and coastal waters between 69°N and 73°N (Huse et al., 2010; Norges Sildesalgslag, 2010) where humpback whales, never before observed on the former herring wintering ground, have been present in large numbers and observed feeding on the same herring schools as killer whales (personal unpublished observations). Although adult humpback whales have previously been observed opportunistically feeding alongside another cetacean species, the Indo-Pacific bottlenose dolphin (*Tursiops aduncus*; Stockin and Burgess, 2005), interactions between humpback and killer whales reported to date have often been agonistic, and almost exclusively involved predation (killer whales towards humpback whales) or defense (e.g., humpbacks being attacked by killer whales or defending other prey species; see Pitman et al. (2017) for a review). This paper reports on the first observations of feeding aggregations formed by killer whales and humpback whales.

We investigated the scope of feeding aggregations formed by humpback and killer whales on the wintering ground of the NSS herring, and further examined if one of the species could be identified as initiating the feeding events. More specifically, using observations collected over three winter seasons (December–February 2013–2016), we tested if observing

feeding humpback whales was related to the occurrence of feeding killer whales, thus suggesting that in these mixed-species feeding aggregations killer whales act as catalysts while humpbacks are opportunists. Possible mechanisms involved in the formation of these feeding aggregations and potential competition are discussed. To our knowledge, this is the first report on humpback whales and killer whales feeding on common prey patches, therefore providing a valuable contribution to our current understanding of humpback whale–killer whale interactions.

Material and methods

Study area

Observations were made around the northern tip of Andøya, centered on Andenes (69°19'N, 16°08'E), Vesterålen, northern Norway (Fig. 1). The NSS herring stock, which has occupied diverse wintering areas over the last decades, started over-wintering in both coastal and offshore waters between 69 and 73°N in 2002 (Huse et al., 2010). Andfjord, a fjord running to the east of Andøya and open in the north towards the Norwegian Sea, has been part of the current wintering ground of the NSS herring. All the feeding events presented in this study occurred in near-shore waters where depths were all less than 250 m and commonly less than 100 m.

Data collection

Observations during the winter seasons 2013–2014 and 2014–2015 were collected during whale watching trips operated from Andenes on board a 36-foot-long rigid hull inflatable boat.

Fig. 1. The map shows the study area where all the field trips were conducted during the three seasons. Each symbol represents the exact location, based on GPS coordinates, of feeding events involving humpback and/or killer whales.

Table 1

Definition of relevant behavioral states and additional cues considered as indicators of ongoing feeding events involving killer whales and/or humpback whales (based on Similä and Ugarte, 1993 and personal observations).

Behavioral indicators and additional visual cues	Description
<i>Killer whales</i>	
No swimming pattern	No clear heading or speed
Diving	Back strongly arched due to vertical orientation of the body required to return to the underwater feeding spot
Dynamic state	High energy and speed, burst swims
<i>Humpback whales</i>	
Fluke-up	Humpback whales often show the fluke before lunging due to vertical orientation when heading to the underwater feeding spot
Surface-lunging	Humpback whales surface with mouth open and ventral throat pleats expanded while engulfing prey and expelling water through baleen plates
<i>Additional cues</i>	
Bird activity	Birds flying over the feeding whales, going down to the surface to fetch prey remains
Floating prey	Dead or stunned herring floating on the feeding spot
Jumping herring	The herring may leap through the sea surface in an attempt to escape underwater predator activity

During 2015–2016, observations were made during opportunistic research surveys conducted on board an 18-foot-long aluminum research vessel from Andenes. The field season lasted from the beginning of December, when the first whales were sighted, to early February, when herring started migrating to the spawning grounds further south and the whales abruptly deserted the area. Trips were conducted in sea state 4 or less on the Beaufort scale. Cetaceans were visually located through a land-based survey from Andenes lighthouse when visibility allowed, or by actively searching from the boat. On both platforms (whale-watching and research vessel), humpback and killer whales were equally searched for. If several cetacean individuals or groups were detected in the area, priority was given to observing feeding cetaceans, since watching humpback and killer whales feeding on herring is a major focus for winter tourism off Andenes. To ensure consistent sampling, this approach was maintained during 2015–2016 on board the research vessel. On both platforms, data collection was conducted as follows: when whales were encountered, species and GPS position were recorded. Based on pre-defined behavioral states and visual cues used as indicators of feeding state, we recorded all feeding events involving killer whales and/or humpback whales (Table 1). An *ad libitum* sampling method was chosen, appropriate for recording discrete predation events (Mann, 1999). When killer whales and humpback whales were found within an estimated 100 m distance for at least five minutes (according to and modified from Bearzi, 2006), the two species were recorded as occurring together. When simultaneously displaying feeding behaviors, and if at least one of the additional cues were observed (see Table 1), the two species were considered to be feeding on the same food patch. Whenever possible, we recorded which species initiated the feeding, and if one species joined the other after feeding had already commenced. If we arrived onsite after feeding by both species had already begun, we classified the species that initiated feeding as ‘missing data’. If humpback whales and killer whales were seen in the same area before initiating feeding, we recorded that both species initiated feeding. Since the study was conducted north of the Arctic circle, the lack of light limited observations to a maximum of one to five hours throughout the observation season. The whale-watching vessel stayed on the field as long as light enabled, and thus sampling ended for the same reasons on both platforms, namely when the group was lost or if sea/weather/light conditions became too poor to continue. No avoidance behavior was noticed for feeding whales, and thus boat effect was not considered.

Statistical analyses

We used the survey cruise as the sample unit, where presence and absence of each whale species was recorded. We summarized

presence and absence as a count for each of the three observation seasons and conducted a chi-square analysis to test the null hypothesis that co-occurrence of both species was independent across observational seasons. We then used count data from surveys in which humpback and killer whales were observed together (i.e., within 100 m of each other) to assess if the incidence of feeding humpback whales was dependent on the occurrence of feeding killer whales. For this final analysis, we used a chi-square exact procedure that estimated the test *p*-value via a Monte Carlo simulation with 5000 replicates. We accepted statistical significance for all tests at $\alpha = 0.05$. All statistical analyses were done using the R software (R Core Team, 2013) and data on feeding locations were plotted in ArcMap.

Results

Data were collected from 17 December 2013 through 6 February 2014 (season 1), from 2 December 2014 through 4 February 2015 (season 2) and from 30 December 2015 through 18 February 2016 (season 3), with a total of 109 opportunistic surveys. Each trip on sea lasted for an average of 2 h (range = 0.4–4.25 h, $n = 109$). We encountered killer whales and/or humpback whales on 100 trips. Time spent with whales averaged 1.7 h per trip (range = 0.25–3.9 h) (Table 2).

Humpback and killer whales were simultaneously observed on 67 of the 100 trips with whales encountered (Table 2). We recorded killer whales feeding on 59 occasions, with humpback whales being involved in 47 (79.7%) of these feedings (Figs. 1 and 2). On all 47 mixed-species feeding aggregations, typically 10–25 killer whales were cooperatively herding the herring close to the surface when, some minutes later, one to three humpback whales appeared on the feeding spot, lunging one to four times through the bait-ball. Of the 47 mixed-species feeding aggregations, the species initiating feeding was known for 36 of these, and killer whales were identified as the species initiating the feeding on 34 (94.4%) occasions. On two

Table 2

Summary of the survey effort during winters 2013–2014 (season 1), 2014–2015 (season 2) and 2015–2016 (season 3) off Andenes. Encounters with humpback whales (*Mn*) and/or killer whales (*Oo*) are also shown.

	Season 1	Season 2	Season 3	Total
Time on sea (hours)	85.6	68.1	63.5	217.2
Time with whales (hours)	67.8	46.1	54.3	168.2
Total number of trips	40	41	28	109
Trips with only <i>Oo</i> encountered	8	13	9	30
Trips with only <i>Mn</i> encountered	1	1	1	3
Trips with both <i>Mn</i> and <i>Oo</i> encountered	30	20	17	67
Trips with neither species encountered	1	7	1	9

Table 3

Summary of all observed humpback whale-killer whale (Mn-Oo) feeding aggregations, along with the species that started the feeding events over the three winters of observation.

	Season 1	Season 2	Season 3	Total
Trips with Mn-Oo feeding aggregations	22	15	10	47
Occasions where Oo started feeding	16	11	7	34
Occasions where Mn started feeding	0	0	0	0
Occasions where both Oo and Mn started feeding	2	0	0	2
Missing data	4	4	3	11
Trips with Oo feeding (single species)	0	8	4	12
Trips with Mn feeding (single species)	1	0	0	1

Fig. 2. Photograph showing two humpback whales surface-lunging through a herring ball being herded by a pod of killer whales.

occasions, both species appeared to initiate the feeding simultaneously. There was no instance recorded where humpback whales initiated a feeding bout and killer whales joined later (Table 3). On all feeding bouts, observation or collection of forage fish on the sea surface enabled confirmation of herring as the prey species. No predatory interactions (killer whales attacking humpback whales) or humpback whales avoiding/defending themselves against killer whales were recorded in our observations.

Proportions of observations of the two species did not vary between seasons ($\chi^2 = 0.007$, $df = 2$, $p = 0.99$), so data from all seasons were pooled. The pooled data are shown in Table 4, where counts of all observations of both species, feeding or not, from all trips are accounted for.

The chi-square exact test of count data from cruises where humpback whales and killer whales were detected together indicated that the likelihood of observing humpback whales feeding increased when killer whales were present and feeding ($\chi^2 = 43.25$, $df = 1$, $p < 0.001$).

Discussion

Despite reports of feeding aggregations involving multiple species from a broad range of taxa, feeding interactions between killer whales and humpback whales have not been reported previously. Here, we have documented 47 multi-species feeding

aggregations recorded over three observation seasons. Typically, killer whales actively feeding on herring were joined by one to three humpback whales abruptly showing up on the feeding spot, racing towards the fish ball and surface lunging, confirming effective predation.

Similar to other *Balaenopteridae*, humpback whales are filter feeders and support their large body size using lunge feeding, both at depths and near the surface, to engulf large amounts of prey (Goldbogen et al., 2008; Hazen et al., 2009). Adapting feeding strategies to prey availability and environmental conditions, humpback whales display a large repertoire of feeding behaviors, including surface lunging (Friedlaender et al., 2009; Hain et al., 1995), bottom feeding with associated prey flushing (Hain et al., 1995), bubble net feeding (D'Vincent et al., 1985; Hain et al., 1982; Jurasz and Jurasz, 1979; Wiley et al., 2011) and lobtail feeding (Weinrich et al., 1992). As such, humpback whales appear as one of the most versatile baleen whales in terms of feeding techniques (Friedlaender et al., 2009; Hain et al., 1982). Evidence for opportunistic feeding has also been documented by Stockin and Burgess (2005), who reported on an adult humpback whale joining and feeding on a fish-ball cooperatively herded below the surface by Indo-Pacific bottlenose dolphins, while the whale was migrating along the coast of Australia. This previous study supports our observations of humpback whales taking advantage of bait balls herded by another cetacean species. Indeed, in most of our feeding observations, killer whales had herded the herring into a compact ball below the surface when humpback whales reached the feeding event. Because studies elsewhere showed that surface feeding humpback whales maximize their foraging efficiency by preferentially lunge-feeding on denser and shallower prey patches of vertical configuration (Friedlaender et al., 2009; Hazen et al., 2009; Sharpe, 2001), we suggest that humpback whales aggregating with foraging killer whales can directly access a highly rewarding shallow prey patch, optimally concentrated, without the costs associated with prey location and herding.

The systematic occurrence over multiple years of a large mass of wintering herring in relatively confined fjords represents an important predictable food source for both killer whales and humpback whales. The NSS herring is known to form dense layers up to several kilometers in length when wintering in fjords, being highly concentrated over an extensive area that would largely enable predators to forage independently (Foote et al., 1996). Instead, we found feeding aggregations composed of humpback and killer whales that happened more often than expected by chance. Because killer whales

Table 4

Contingency table summarizing all 109 trips. When both humpback (Mn) and killer whales (Oo) were observed on the same trip, observation counts are given for the two species seen "Together" (T; within 100 m distance for at least 5 min, see Methods) and "Apart" (A).

	Oo observed feeding	Oo observed, but not feeding	Oo not observed
Mn observed feeding	T: 47 A: 0	0	1
Mn observed, but not feeding	T: 1 A: 0	T: 5 A: 14	2
Mn not observed	11	19	9

were the initiating species in 94% of the observed feeding events (when this parameter could be positively assessed) and because humpback whales aggregated at a later stage, we hypothesise that humpback whales may benefit more from these interactions than killer whales. Our observations thus suggest killer whales as the catalyst species, whose feeding behavior is detected by opportunist humpback whales that subsequently aggregate on food patches.

Killer whales are acoustically active when foraging on herring, producing high rates of calls and echolocation clicks (Simon et al., 2007a, 2007b). Therefore, we assume that humpback whales can hear vocal herring-feeding killer whales, attracting them towards the feeding spot. This was supported by an observation of two resting humpback whales that suddenly started travelling, and that we followed for about 1 km before reaching a pod of carousel feeding killer whales. The humpback whales clearly joined the feeding event as they surface-lunged twice through the herring ball. Evidence for humpback whales being capable of long-distance detection and response to killer whales' vocalizations has been documented by Pitman et al. (2017) and Curé et al. (2015). The latter study also suggested that humpback whales perceive some killer whale vocalizations as a threat, as humpback whales displayed marked avoidance behavior, an anti-predator response, upon hearing vocalizations of killer whales recorded during mammal-killing events. However, humpback whales' reaction to fish-eating killer whale vocalizations was not tested. It has been shown that harbor seals are able to discriminate between fish feeding and marine mammal feeding killer whales through learning and habituation, and it is likely that humpback whales have also habituated to sounds produced by fish feeding killer whales in the study area (Deecke et al., 2002). On the other hand, recent studies suggest that killer whales rarely prey on adult baleen whales, but instead direct their attacks on calves in breeding grounds or on their first migration to feeding grounds (McCordic et al., 2014; Mehta et al., 2007; Pitman et al., 2015). Humpback whale calves were not observed in our study area and killer whales are specialized on feeding on herring during the winter months. In such an ecological context, humpback whales may not perceive killer whales as predators, promoting the occurrence of observed feeding aggregations as a novel non-predatory type of interaction between the two species.

In the previous wintering grounds of NSS herring (Tysfjord, Ofotfjord and Vestfjord) where killer whales were present, humpback whales were not observed during 20 years of fieldwork (1986–2006, Tiu Similä pers. comm). The current, more offshore wintering grounds of the herring stock have brought the two species together where humpback whales, as a highly adaptive species, appear to actively aggregate with foraging killer whales for direct access to prey, with presumably reduced costs. Although competition was not investigated in this study, some feeding events were terminated almost instantly after humpback whale lunging was observed, which suggests potential competition costs for killer whales, which may greatly vary with the effort put into the prey location and herding processes. Nøttestad et al. (2002) observed killer whales diving to 160–180 m depth, where the herring remain during the daytime, and lifting dense schools to the surface by coordinated actions. Because such foraging behavior in deep waters demands a lot of energy, losing the herring ball to opportunistic humpback whales could represent a significant cost to killer whales. Our observations were made in areas where depths mostly spanned 50–100 meters. If the herring schools that are foraged upon come from small schools that remain in shallow waters during the day (Nøttestad, 1998), then any induced costs to killer whales may be lower than presumed, as they can easily move to another shallow prey patch. However, it cannot be ruled out that the killer whales had strategically pushed the herring towards these shallows as a feeding strategy (Nøttestad, 2002). If they subsequently lose the herded bait-ball, significant costs would be induced.

Humpback whales migrate annually between high-latitude feeding grounds and tropical breeding areas, where they mainly rely on their fat stores (Clapham and Mead, 1999). As such, predictably available food patches, with reduced foraging costs, in either the departure region or along the migration route may constitute a valuable additional prey intake (Stevick et al., 2003). We do not claim that humpback whales rely on the killer whales' skills for efficient herring-feeding. However, we suggest that aggregating with feeding killer whales may have become an emerging foraging strategy for humpback whales in northern Norwegian fjords during the winter months.

Acknowledgements

Sea Safari Andenes AS and Lofoten Opplevelser AS were the research platforms during the two first seasons of this study. Fieldwork on the third season was funded by grants from Sea World & Busch Garden Conservation Fund and Spare Bank-1 Norge. Todd Atwood provided critical statistical advice. Data collection was facilitated thanks to R. Karoliussen's great navigation skills.

References

- Au, D.W., Pitman, R.L., 1988. Seabird relationships with tropical tunas and dolphins. In: Burger, J. (Ed.), *Seabirds and Other Marine Vertebrates*. Columbia University Press, New York, pp. 174–212.
- Bearzi, G., Politi, E., Agazzi, S., Azzellino, A., 2006. Prey depletion caused by overfishing and the decline of marine megafauna in Eastern Ionian Sea coastal waters (Central Mediterranean). *Biol. Conserv.* 127, 373–382.
- Bearzi, M., 2006. California sea lions use dolphins to locate food. *J. Mammal.* 87, 606–617.
- Berec, L., 2010. Impacts of foraging facilitation among predators on predator-prey dynamics. *Bull. Math. Biol.* 72, 94–121.
- Brager, S., 1998. Feeding associations between white-fronted terns and Hector's dolphins in New Zealand. *Condor* 100, 560–562.
- Bruno, J.F., Stachowicz, J.J., Bertness, M.D., 2003. Inclusion of facilitation into ecological theory. *Trends Ecol. Evol.* 18, 119–125.
- Camphuysen, K., Webb, A., 1999. Multi-species feeding associations in North Sea seabirds: jointly exploiting a patchy environment. *Ardea* 87, 177–198.
- Clapham, P.J., Mead, J.G., 1999. Megaptera novaeangliae. *Mamm. Species* 604, 1–9.
- Curé, C., Sivle, L.D., Visser, F., Wensveen, P.J., Isojunno, S., Harris, C.M., Kvadsheim, P.H., Lam, F.-P.A., Miller, P.J.O., 2015. Predator sound playbacks reveal strong avoidance responses in a fight strategist baleen whale. *Mar. Ecol. Prog. Ser.* 526, 267–282.
- D'Vincent, C.G., Nilson, R.M., Hanna, R.E., 1985. Vocalization and Coordinated Feeding Behavior of the Humpback Whale Megaptera Novaeangliae in Southeastern Alaska, USA. *Scientific Reports of the Whales Research Institute, Tokyo* 41–48.
- Deecke, V.B., Slater, P.J.B., Ford, J.K.B., 2002. Selective habituation shapes acoustic predator recognition in harbour seals. *Nature* 420, 171–173.
- Domenici, P., Batty, R.S., Similä, T., Ogam, E., 2000. Killer whales (*Orcinus orca*) feeding on schooling herring (*Clupea harengus*) using underwater tail-slaps: kinematic analyses of field observations. *J. Exp. Biol.* 203, 283–294.
- Foote, K.G., Ostrowski, M., Røttingen, I., Slotte, A., 1996. Abundance estimation of Norwegian spring spawning herring wintering in the Vestfjord system, December 1996. *Copenhagen* 1996 p. 8.
- Friedlaender, A.S., Hazen, E.L., Nowacek, D.P., Halpin, P.N., Ware, C., Weinrich, M.T., Hurst, T., Wiley, D., 2009. Diel changes in humpback whale Megaptera novaeangliae feeding behavior in response to sand lance Ammodytes spp. behavior and distribution. *Mar. Ecol. Prog. Ser.* 395, 91–100.
- Goldbogen, J.A., Calambokidis, J., Croll, D.A., Harvey, J.T., Newton, K.M., Oleson, E.M., Schorr, G., Shadwick, R.E., 2008. Foraging behavior of humpback whales: kinematic and respiratory patterns suggest a high cost for a lunge. *J. Exp. Biol.* 211, 3712–3719.
- Goyert, H.F., Manne, L.L., Veit, R.R., 2014. Facilitative interactions among the pelagic community of temperate migratory terns, tunas and dolphins. *Oikos* 123, 1400–1408.
- Grunbaum, D., Veit, R.R., 2003. Black-browed albatrosses foraging on Antarctic krill: density-dependence through local enhancement? *Ecology* 84, 3265–3275.
- Hain, J.H.W., Carter, G.R., Kraus, S.D., Mayo, C.A., Winn, H.E., 1982. Feeding behavior of the humpback whale, Megaptera novaeangliae, in the Western North Atlantic. *Fish. Bull.* 80, 259–268.
- Hain, J.H.W., Ellis, S.L., Kenney, R.D., Clapham, P.J., Gray, B.K., Weinrich, M.T., Babb, I.G., 1995. Apparent bottom feeding by humpback whales in Stellwagen Bank. *Mar. Mamm. Sci.* 11, 464–479.
- Hazen, E.L., Friedlaender, A.S., Thompson, M.A., Ware, C.R., Weinrich, M.T., Halpin, P.N., Wiley, D.N., 2009. Fine-scale prey aggregations and foraging ecology of humpback whales Megaptera novaeangliae. *Mar. Ecol. Prog. Ser.* 395, 75–89.

- Huse, G., Fernö, A., Holst, J.C., 2010. Establishment of new wintering areas in herring co-occurs with peaks in the 'first time/repeat spawner' ratio. *Mar. Ecol. Prog. Ser.* 409, 189–198.
- Jurasz, C.M., Jurasz, V.P., 1979. Feeding modes of the humpback whale, *Megaptera novaeangliae*, in southeast Alaska. *Sci. Rep. Whales Res. Inst. Tokyo* 31, 69–83.
- Lett, C., Semeria, M., Thiebault, A., Tremblay, Y., 2014. Effects of successive predator attacks on prey aggregations. *Theor. Ecol.* 7, 239–252.
- Mann, J., 1999. Behavioral sampling methods for cetaceans: a review and critique. *Mar. Mamm. Sci.* 15, 102–122.
- McCordic, J.A., Todd, S.K., Stevick, P.T., 2014. Differential rates of killer whale attacks on humpback whales in the North Atlantic as determined by scarification. *J. Mar. Biol. Assoc. U. K.* 94, 1311–1315.
- Mehta, A.V., Allen, J.M., Constantine, R., Garrigue, C., Jann, B., Jenner, C., Marx, M.K., Matkin, C.O., Mattila, D.K., Minton, G., Mizroch, S.A., Olavarria, C., Robbins, J., Russell, K.G., Seton, R.E., Steiger, G.H., Vikingsson, G.A., Wade, P.R., Witteveen, B.H., Clapham, P.J., 2007. Baleen whales are not important as prey for killer whales *Orcinus orca* in high-latitude regions. *Mar. Ecol. Prog. Ser.* 348, 297–307.
- Nøttestad, L., Fernö, A., Axelsen, B.E., 2002. Digging in the deep: killer whales' advanced hunting tactic. *Polar Biol.* 25, 939–941.
- Nøttestad, L., 1998. Forholdet mellom sild og spekkhogger: Kampen for tilværelsen. *Naturen* 2, 78–81.
- Nøttestad, L., 2002. Killer Whales (*Orcinus Orca* L.) and Saithe (*Pollachius Virens* L.) Trap Herring (*Clupea Harengus* L.) in Shallow Water by Taking Advantage of Steep Bottom Topography. International Council for the Exploration of the Sea, Hillerød, Denmark, p. 11.
- Norges Sildesalgslag, 2010. Norwegian Spring-spawning Herring Fisheries, Fisheries Statistics Data Set. Norges Sildesalgslag, Bergen, Norway.
- Øien, N., 2007. Hval i Norskehavet. In: Skogen, M., Gjøsæter, H., Toresen, R., Robberstad, Y. (Eds.), *Havets ressurser og miljø*. Institute of Marine Research (In Norwegian).
- Pitman, R.L., Ballance, L.T., 1992. Parkinson petrel distribution and foraging ecology in the Eastern Pacific – aspects of an exclusive feeding relationship with dolphins. *Condor* 94, 825–835.
- Pitman, R.L., Totterdell, J.A., Fearnbach, H., Ballance, L.T., Durban, J.W., Kemps, H., 2015. Whale killers: prevalence and ecological implications of killer whale predation on humpback whale calves off Western Australia. *Mar. Mamm. Sci.* 31, 629–657.
- Pitman, R.L., Deecke, V.B., Gabriele, C.M., Srinivasan, M., Black, N., Denking, J., Durban, J.W., Mathews, E.A., Matkin, D.R., Neilson, J.L., Schulman-Janiger, A., Shearwater, D., Stap, P., Ternullo, R., 2017. Humpback whales interfering when mammal-eating killer whales attack other species: mobbing behavior and interspecific altruism? *Mar. Mamm. Sci.* 33, 7–58.
- Poysa, H., 1992. Group foraging in patchy environments – the importance of coarse level local enhancement. *Ornis Scand.* 23, 159–166.
- R Core Team, 2013. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>.
- Sharpe, F.A., 2001. Social Foraging of the Southeast Alaskan Humpback Whale *Megaptera novaeangliae*. Department of Biological Sciences, Simon Fraser University, p. 141.
- Silverman, E.D., Veit, R.R., Nevitt, G.A., 2004. Nearest neighbors as foraging cues: information transfer in a patchy environment. *Mar. Ecol. Prog. Ser.* 277, 25–36.
- Similä, T., Ugarte, F., 1993. Surface and underwater observations of cooperatively feeding killer whales in northern Norway. *Can. J. Zool.* 71, 1494–1499.
- Similä, T., Holst, J.C., Christensen, I., 1996. Occurrence and diet of killer whales in Northern Norway: seasonal patterns relative to the distribution and abundance of Norwegian spring-spawning herring. *Can. J. Fish. Aquat. Sci.* 53, 769–779.
- Similä, T., 1997a. Behavioral Ecology of Killer Whales in Northern Norway. Norwegian College of Fishery Science, University of Tromsø.
- Similä, T., 1997b. Sonar observations of killer whales (*Orcinus orca*) feeding on herring schools. *Aquat. Mamm.* 23, 119–126.
- Simon, M., Ugarte, F., Wahlberg, M., Miller, L.A., 2006. Icelandic killer whales *Orcinus orca* use a pulsed call suitable for manipulating the schooling behaviour of herring *Clupea harengus*. *Bioacoustics* 16, 57–74.
- Simon, M., McGregor, P.K., Ugarte, F., 2007a. The relationship between the acoustic behaviour and surface activity of killer whales (*Orcinus orca*) that feed on herring (*Clupea harengus*). *Acta Ethol.* 10, 47–53.
- Simon, M., Wahlberg, M., Miller, L.A., 2007b. Echolocation clicks from killer whales (*Orcinus orca*) feeding on herring (*Clupea harengus*) (L.). *J. Acoust. Soc. Am.* 121, 749–752.
- Stevick, P.T., Allen, J., Berube, M., Clapham, P.J., Katona, S.K., Larsen, F., Lien, J., Mattila, D.K., Palsboll, P.J., Robbins, J., Sigurjonsson, J., Smith, T.D., Oien, N., Hammond, P.S., 2003. Segregation of migration by feeding ground origin in North Atlantic humpback whales (*Megaptera novaeangliae*). *J. Zool.* 259, 231–237.
- Stockin, K.A., Burgess, E.A., 2005. Opportunistic feeding of an adult humpback whale (*Megaptera novaeangliae*) migrating along the coast of southeastern Queensland, Australia. *Aquat. Mammals* 31, 120–123.
- Vaughn, R.L., Shelton, D.E., Timm, L.L., Watson, L.A., Wuersig, B., 2007. Dusky dolphin (*Lagenorhynchus obscurus*) feeding tactics and multi-species associations. *N. Z. J. Mar. Freshw. Res.* 41, 391–400.
- Weinrich, M.T., Schilling, M.R., Belt, C.R., 1992. Evidence for acquisition of a novel feeding behavior – lobtail feeding in humpback whales *Megaptera novaeangliae*. *Anim. Behav.* 44, 1059–1072.
- Wiley, D., Ware, C., Bocconcelli, A., Cholewiak, D., Friedlaender, A., Thompson, M., Weinrich, M., 2011. Underwater components of humpback whale bubble-net feeding behaviour. *Behaviour* 148, 575–602.

PAPER II

Killer whales (*Orcinus orca*) feeding on lumpfish (*Cyclopterus lumpus*)
in northern Norway.

Jourdain, E., Karoliussen, R., de Vos, J., Zakharov, S. E., and Tougard, C. (2019).

Marine Mammal Science, 00: 1-14.

doi: 10.1111/mms.12618.

Killer whales (*Orcinus orca*) feeding on lumpfish (*Cyclopterus lumpus*) in northern Norway

Eve Jourdain¹ | Richard Karoliussen¹ | Jacques de Vos¹ |
Stanislav E. Zakharov¹ | Christelle Tougard²

¹Norwegian Orca Survey, Andenes, Norway

²ISEM, Université de Montpellier, CNRS, EPHE, IRD, Montpellier, France

Correspondence

Eve Jourdain, Norwegian Orca Survey,
Haugnesveien 30, 8480 Andenes, Norway.
Email: post@norwegianorcasurvey.no

Funding information

Sea World and Busch Gardens Conservation
Fund; Sea World and Parks Entertainment

Abstract

Killer whales (*Orcinus orca*) in Norwegian waters have long been known to rely on Atlantic herring (*Clupea harengus*) as a main prey resource. However, research almost exclusively conducted at seasonal herring grounds may have biased studies away from detecting other potentially significant prey species. Since 2013, dedicated research efforts have focused on monitoring killer whale occurrence and foraging ecology throughout the year in northern Norway. This study presents results on site-fidelity of photographically identified individuals, predation records and behavioral patterns from five spring seasons (March–April) in 2014–2018 in Andfjord, northern Norway. A minimum number of 75 adult and subadult killer whales (out of a catalog of 971 individuals) returned seasonally to the study area for foraging and residency for up to six weeks. Lumpfish (or lumpsucker, *Cyclopterus lumpus*) was the only type of prey identified (based on molecular or visual identification) on 22 predation events from 2016 ($n = 4$), 2017 ($n = 2$) and 2018 ($n = 16$). Spatial group cohesion observed when foraging was a potential adaptation for efficiently hunting this prey species. These whales were also encountered at herring wintering grounds the same years, but with different group sizes. Such behavioral adaptations suggested intraannual switching between prey resources and foraging strategies.

KEYWORDS

feeding behavior, foraging, foraging strategy, killer whale, *Orcinus orca*, seasonality, site-fidelity

1 | INTRODUCTION

The killer whale (*Orcinus orca*) is found in all the world's oceans and as a species is generalist, with a wide range of prey documented including fish, marine mammals, cephalopods, reptiles and seabirds (Ford, 2009). However, populations have been shown to be prey specialists. For example, in the temperate coastal northeastern Pacific, killer whales feed exclusively on either marine mammals or fish, (Baird & Dill, 1995; Ford & Ellis, 2006; Ford et al., 1998). Such prey specialization, including underlying foraging behavior and knowledge about particular prolific feeding areas, are culturally transmitted in the form of learned behavior within groups and persist over time (Ford et al., 1998; Riesch et al., 2012). Resulting ecological and cultural divergences among social groups promote the formation of sympatric socially and genetically isolated ecotypes (Foote et al., 2016; Ford & Ellis, 2014). Divergent foraging strategies, by creating differences in occurrence and movement patterns as well as social organization, may further play a role in population structuring (Hoelzel et al., 2007; Morin et al., 2010; Parsons et al., 2013).

In the North Atlantic, two ecotypes of killer whales were proposed based on dietary specialization, Type 1 and Type 2, with nitrogen stable isotope values and tooth wear used as proxies to indicate key prey choice for each type (Foote, Newton, Piortney, Willerslev, & Gilbert, 2009; Foote, Vester, Vikingsson, & Newton, 2012). Type 1 refers to killer whales from Iceland, Norway, and the North Sea, which rely on the Atlantic herring (*Clupea harengus*) and/or the Atlantic mackerel (*Scomber scombrus*) as their main food source(s). In contrast, Type 2 primarily feeds on other cetaceans. However, this simplistic classification seems unlikely to encapsulate the broad-scale diversity of the diet of North Atlantic killer whales. In fact, there is strong evidence for interindividual variation in the relative proportions of prey types consumed within Type 1 killer whales. For example, some may also prey on pinnipeds, as evidenced by field observations (Foote, Similä, Vikingsson, & Stevick, 2010; Jourdain, Vongraven, Bisther, & Karoliussen, 2017; Samarra et al., 2018; Vongraven & Bisther, 2014). As such, different feeding specializations may exist within populations. Recent studies using ecological tracers were further supportive of interindividual variation in dietary habits among regions (e.g., Faroe Islands/Denmark vs. Greenland; Bourque et al., 2018), within regions (Samarra, Vighi, Aguilar, & Vikingsson, 2017), and within social groups (in Greenland, see Bourque et al., 2018). However, the lack of killer whale identification data and predation records for most regions has impeded our understanding of fine-scale seasonal and/or group-specific prey specializations so far. The potential occurrence of variable foraging strategies and dependence on main and alternative prey resources remains to be investigated further in the North Atlantic.

Between 1986 and 2005, Norwegian killer whale studies were mainly focused on wintering grounds of the Norwegian Spring Spawning (NSS) stock of the Atlantic herring. At that time these grounds were located in a fjord system (Vestfjord-Ofotfjord-Tysfjord) in the Lofoten region, northern Norway (68°13'N, 15°37'E). Research efforts were focused on photographic identification and investigation of feeding behavior, occurrence patterns, social ecology, and acoustics (Domenici, Batty, Similä, & Ogam, 2000; Similä, 1997; Similä & Ugarte, 1993; Similä, Holst, & Christensen, 1996; Van Opzeeland, Corkeron, Leyssen, Similä, & Van Parijs, 2005; Van Parijs, Leyssen, & Similä, 2004). These studies highlighted killer whales as efficient herring predators using specialized feeding strategies (e.g., Similä & Ugarte, 1993). Photo-identification surveys conducted on both wintering and spawning grounds of the NSS herring, combined with satellite tracking of six individual whales tagged in 2000 and 2001, further suggested that some groups may follow the herring's movements for most of the year (Bisther & Vongraven, 1995; Stenersen & Similä, 2004; Similä et al., 1996), though effective herring foraging offshore remained largely inconclusive. As a result, and due to lack of evidence for additional prey items, herring has been regarded as the main prey for killer whales in Norwegian waters. This conclusion was further supported by a lack of intrapopulation variation in stable isotope values from skin samples collected from whales in the NSS wintering grounds (Foote et al., 2012).

However, focusing research efforts on seasonal herring grounds may have biased studies away from detecting other potentially significant prey species. In fact, recent studies conducted outside seasonal herring grounds documented predation on Atlantic salmon (*Salmo salar*; Vester & Hammerschmidt, 2013), Atlantic mackerel (Nøttestad et al., 2014), harbor porpoise (*Phocoena phocoena*; Cosentino, 2015), and on pinnipeds (Jourdain et al., 2017; Vongraven &

Bisther, 2014), suggesting intrapopulation ecological variation. Combining sighting and predation records of individuals over time would help in assessing prey specializations of killer whales in this region.

Longitudinal photo-identification studies, as initiated by colleagues three decades ago, have been reestablished in 2013 at current herring wintering (November–February) grounds (see Huse, Fernö, & Holst, 2010; Jourdain and Vongraven, 2017). From mid-January to mid-February, the herring migrate to the main spawning area located 800–1,000 km south, off the central coast of Norway (Dragesund, Johannessen, & Ulltang, 1997), leading to the departure of the winter aggregating killer whales. Research efforts yet maintained throughout the rest of the year in the study area aimed at collecting photo-identification data and predation records.

Here, we report on a subset of known herring-feeding killer whales showing site-fidelity in a northern Norwegian fjord in spring (March–April), after the herring has departed, for the period 2014–2018. Behavior and predation records were collected over multiple years in order to assess foraging as a predominant function for seasonal use of the area in spring, and to identify the target prey. Group cohesion was investigated with reference to the focal prey and group sizes were compared to winter observations for insights into behavioral adaptations and therefore, indication of seasonal prey switching and specialization for these whales.

2 | MATERIALS AND METHODS

2.1 | Data collection

Data were collected in March–April of 2015–2018 east of Andøya (Andfjord, 69°19'N, 16°08'E; Figure 1), in Vesterålen region, northern Norway. Because killer whale presence was not known to be regular in the study area at this specific time of year, surveys were largely opportunistic, and effort varied considerably across years. In 2015, surveys were always conducted in response to sightings being reported by locals. In the following years, and based on the 2015 sighting records, land-based watches searched for killer whales on good weather days, as did opportunistic boat surveys, in addition to relying on sighting reports. Supplementary photographs of individual killer whales within the study area with associated date and time data were provided by local wildlife photographers for April–March of 2014–2017, and those of high enough quality were accepted for use in the analyses.

Surveys were conducted on a 20-ft-long (6 m) aluminum research vessel in sea conditions of Beaufort scale 4 or less. We defined an encounter as a sighting of a group of killer whales. A group was defined as all individuals observed in apparent association, acting in a coordinated manner during the observation period and which were within the visual range of the observers (as per Baird & Dill, 1995). For each encounter, the geographic position (GPS Garmin) was recorded and the whales were systematically photographed using a digital reflex camera and a mounted lens of 70–200 mm in focal length. Predominant behavioral pattern, group size estimate and cohesion index (see below) were also recorded at 5-min intervals for the majority of surveys. Surveys terminated when the group was lost from sight, traveled away from the study site, or if light and/or sea conditions deteriorated.

Behavior was divided into five categories (adapted from Ford, 1989): (1) *Foraging*—erratic swimming in varying directions, sudden lunges and high-speed swimming, presence of prey fragments and seabirds; (2) *Traveling*—consistent movement with all group members, in a tight group formation, following the same course at the same speed, while displaying synchronous dives and surfacings; (3) *Milling*—low speed movement with individuals engaged in frequent changes in direction leading to no directional movement by the group; (4) *Resting*—low level activity with individuals being stationary lined-up and floating motionless at the surface while displaying coordinated breathings; (5) *Socializing*—high energy activity in which individuals may physically interact with one another and/or perform aerial displays such as leaps, tail slaps, and spy-hops (defined as vertical half-rise of the body out of the water). Activities displayed by the majority of the whales were considered as the main behavioral pattern for the group.

Spatial cohesion of the group, often variable throughout encounters, was characterized using a cohesion index (CI) (adapted from Daura-Jorge, Wedekin, Piacentini, & Simões-Lopes, 2005). Within a group, individuals scattered over distances (visually assessed) greater than 200 m were classified as low cohesion (CI = 1), individuals occupying a

FIGURE 1 Location of the study area in northern Norway. Box A corresponds to Andfjord where the data presented in this study were collected. Box B locates the wintering herring grounds where the whales identified in this study were encountered in November–February in 2015–2018 while feeding on herring. Spatial distribution of killer whale encounters during March–April in 2014–2018, with corresponding behavioral patterns are plotted. Depth contours (m), as well as GPS tracks (blue plots) of the opportunistic surveys conducted in 2018 are also shown.

60–200 m area were classified as intermediate cohesion ($CI = 2$) and individuals aggregated within 50 m were classified as high cohesion ($CI = 3$). Smaller subgroups, often composed of adult female(s) with offspring(s), were considered as part of the main group. Classifications were made by the same observer throughout the study, reducing potential observer-related bias.

From 2016 onwards, collection of tissue fragments from prey killed and consumed by killer whales was also attempted. Based on the methods developed by Ford et al. (1998), we used speed bursts, directional chases, fast and sharp turns, and head-nodding movements of surfacing whales along with scavenging seabirds as clues to feeding events. Upon detection of such events, the research vessel slowly approached the focal whale, stopped in the fluke print (glassy area visible at the surface as the whale dives and its tail creates an upwelling, as defined by Ford & Ellis, 2006) and observers assessed potential presence of prey remains in the water column. Using a long-handled (3 m) fine-mesh net, one observer onboard collected any accessible prey fragments. Because collection was challenged by fast-sinking prey parts and common rough sea state, a professional free-diver assisted in collection of prey remains in April 2018. Upon collection, fragments were placed in a plastic bag and stored in a cooling box while in the field. Ashore, fragments were stored in 1.5 ml vials containing 95% ethanol at -20°C until analysis.

Whenever a focal whale was observed feeding at close range to the vessel, photographs were also taken in an attempt to visually-identify target prey species. An unmanned aircraft DJI Phantom IV carrying a DJI 1/2.3" CMOS camera was used on three days in 2018 to capture still images and video footage of prey capture events.

2.2 | Data processing

Killer whales were individually identified based on location and number of notches in the leading and/or trailing edge of the dorsal fin, along with pigmentation pattern of and natural markings occurring on the saddle patch (Bigg, 1982). Only individuals displaying one or several unique features were considered as distinctive and effectively cataloged. Calves and other individuals lacking persistent identifying markings were excluded. Individuals with distinctly taller dorsal fins were classified as "Male" (Olesiuk, Bigg, & Ellis, 1990), adult female-sized individuals repeatedly seen with one calf or older offspring(s) swimming in echelon position throughout the study period were qualified as "Female" and individuals for which sex was uncertain were qualified as "Unknown." Photographs were scored for quality ($0 < Q < 2$) based upon sharpness, contrast and angle. To minimize misidentification, only excellent quality ($Q2$) photographic records of distinctive cataloged killer whales were considered in this study. Both left- and right-side identification photos were used for ID, but only whales cataloged from the left side were retained for analysis.

2.3 | Data analysis

2.3.1 | Sighting and behavioral patterns

To assess site-fidelity patterns, individual sighting records were compared across days and years. Despite sampling biases due to uneven effort throughout seasons and across years, we considered minimum residency to be informative of individuals' site-fidelity periods within seasons. Therefore, the number of days elapsed from the first to the last sighting within each season was calculated for each individual, as compared to the total length of survey period for each given season (as per Weinrich, Martin, Griffiths, Bove, & Schilling, 1997). Mean residency values were calculated across individuals per season, excluding individuals not seen, and averaged for the entire study period.

To investigate seasonal usage of the study area, relative predominance of observed behavioral states was calculated. Further seasonal behavioral characteristics, assumed to be adaptive to the target prey, were tested as follows. A chi-squared test was used to test the relationship between behavior and cohesion index (CI). Due to low occurrence, records of resting and socializing behaviors were not included in the test. Records of group size collected for this study were compared to group sizes recorded at herring wintering grounds for these same whales in 2015–2018 using a Mann–Whitney–Wilcoxon test. Statistical tests were performed using R software (R Core Team, 2016).

2.3.2 | Molecular and visual prey identification

Molecular analyses were conducted on prey tissue samples for identification of target species. Total DNA from 11 samples was extracted from ethanol-preserved samples following standard procedures (Sambrook, Fritsch, & Maniatis, 1989). A fragment of the cytochrome c oxidase I gene (COI) was PCR-amplified using the GoTaq G2 Hot Start Colorless Master Mix (Promega) and the primers Pros1Fwd and Pros2Rev (Sparks & Smith, 2004) with an annealing step at 46°C. Direct sequencing was carried out in both directions at the technical facilities of the Genotyping and Sequencing Platform of the Institut des Sciences de l'Evolution de Montpellier (ISEM). Sequences were then aligned in MEGA v7 (Kumar, Stecher, & Tamura, 2016) and blasted (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) against sequences from the GenBank database (<https://www.ncbi.nlm.nih.gov/genbank/>). Photographs and video footages of predation events were scanned for visual identification of prey species.

3 | RESULTS

3.1 | Effort

From 2015 to 2018, 42 days at sea in March–April resulted in 123.2 hr spent with killer whales and 14,170 photographs taken over 51 encounters from a dedicated research vessel. Photographic contributions from wildlife photographers led to 11 additional encounter days for the period 2014–2017. Behavior and CI were recorded for 62 hr from 48 encounters. Molecular ($n = 11$) and photographic ($n = 11$) evidence enabled positive identification of prey species for 22 predation events from 11 encounter days in 2016–2018. For the period 2014–2018, photographs resulted in 276 identifications of 75 distinctively marked individuals (Figure 2). The number of identified individuals per season varied from a minimum of six (in 2014) to a maximum of 42 (in 2018), with a mean of 27 (standard deviation $SD = 13.3$). Poor weather conditions in 2015 and 2017 resulted in lower numbers of encounter days and identifications for these years. Sampling effort is summarized in Table 1.

FIGURE 2 Cumulative number of identified individual killer whales in Andfjord in March–April plotted against cumulative identifications. Each data point represents a survey. The 1:1 plot is shown for reference. Dashed lines separate the five calendar years (2014–2018).

TABLE 1 Sampling effort for the period March–April in 2014–2018 in Andfjord.

	2014	2015	2016	2017	2018	Total
Dedicated effort by researchers						
Effort days	0	7	11	6	20	44
Encounter days	0	7	11	4	20	42
Killer whale encounters	0	7	15	4	25	51
Hours spent with whales	0	11.7	22.8	8.1	80.6	123.2
Hours of behavioral sampling	0	6.0	13.7	4.3	38.0	62.0
Photographs taken	0	2,480	5,080	990	5,620	14,170
Predation events for which prey remains were collected	0	0	2	0	9	11
Predation events for which prey were visually identified	0	0	2	2	7	11
Additional photographic contributions						
Encounter days	1	0	3	7	0	11
Photographs provided	300	0	127	1,280	0	1,707
Total						
Encounter days	1	7	14	11	20	53
Identifications	6	41	70	40	119	276
Distinct individuals	6	29	33	25	42	—

3.2 | Characterization of individual killer whales

The 75 identified individuals represent 7.7% of the 971 killer whales identified in the study area (Jourdain & Karoliussen, 2018). Of these, 34 (45.3%) were adult males, 21 (28%) were adult females and 20 (26.7%) were of unknown sex; 61 (81.3%) were encountered at herring wintering grounds on one or multiple years between 2013 and 2018 (unpublished data). Social associations appeared highly variable across encounters, and thus we considered individuals independently. Plotting the cumulative number of newly identified individuals in the period March–April from 2014 to 2018 revealed a steadily decreasing identification rate, suggesting that most distinctive whales using Andfjord in spring had been identified during the study (Figure 2).

3.3 | Sighting frequencies

Of 75 individuals, 20 (26.7%) were sighted only once, while 55 (73.3%) were sighted during two or more encounters in March–April from 2014 to 2018. Of these, 40 were seen in multiple years (range = 2–4; Figure 3). Averaging all five spring seasons, individuals were sighted during a mean of 1.8 seasons ($SD = 0.9$, range = 1–4) and on 3.7 d ($SD = 2.7$, range = 1–12). For individuals sighted on a minimum of two encounter days within each spring season (Figure 3), mean (minimum) residency varied from 8.3 days (in 2015) to 16 days (in 2018) leading to a mean of 11.3 days ($SD = 6.2$, range = 2–39; Table 2) throughout the study period.

3.4 | Behavioral patterns

Foraging (70.3%) and traveling (23.9%) represented the most frequently observed behavioral states, while milling, resting, and socializing accounted for less than 6% of the observations. Group cohesion varied with behavior ($\chi^2 = 3,940$, $df = 4$, $p < .001$). Individuals adopted a tighter group formation when traveling as opposed to scattered over several hundred meters and up to 1 km when foraging. Groups encountered were significantly smaller (Mann–

FIGURE 3 (a) Distribution of individual sighting frequencies within season (March–April), averaged over the entire study period 2014–2018; (b) Distribution of individual sighting frequencies across years, that is, number of seasons (March–April) in which individuals were photographed from 2014 to 2018.

TABLE 2 Summary of the parameters used to assess minimum residency of killer whales in Andfjord during March–April in 2014–2018. Calculated residency should be considered as minimum values due to opportunistic data collection.

	2014	2015	2016	2017	2018
Number of encounter days	1	7	14	11	20
First-to-last encounter period (days)	—	19	32	42	44
Individuals resighted	—	9	21	8	26
Residency mean $\pm$ SD (min–max)	—	8.3 $\pm$ 3.7 (2–14)	10.6 $\pm$ 4.8 (3–20)	10.3 $\pm$ 6.2 (4–16)	16 $\pm$ 10.3 (4–39)

Whitney–Wilcoxon: $W = 297$, $p = .001$) in spring (median = 8, range = 1–30, $n = 49$) than in winter (median = 14, range = 1–40, $n = 23$; Figure 4).

3.5 | Prey identification

Prey fragments from all 11 predation events sampled were identified to be from lumpfish (*Cyclopterus lumpus*; Table 3). The alignment of the COI fragment was 587 nucleotides long. From the 11 sequences obtained for molecular identification, all BLAST queries resulted in the identification of the lumpfish with 99% to 100% of similarity with the sequence JN311799 (Knebelsberger et al., 2014). The new sequences were deposited in GenBank under the accession numbers MF688049 and MH922998–MH923009.

Photo and aerial video material enabled unambiguous identification of the lumpfish as the consumed prey for seven and four additional predation events, respectively (Figure 5, Table 3).

Thirteen individual killer whales were confirmed feeding on lumpfish from these 22 predation events, of which five were adult males, three were adult females and five were of unknown sex including three subadults (Table 3). Five individuals (NKW-0004, NKW-0572, NKW-0712, NKW-0715, and Y093) fed on lumpfish on multiple occasions (range = 2–4) during the same encounter. Adult male NKW-0004 was further confirmed to be feeding on lumpfish over two different days, five days apart, in April 2018.

FIGURE 4 Distribution of group size adopted by the lumpfish-feeding killer whales when encountered at the lumpfish spawning grounds ($n = 49$) vs. herring wintering grounds ($n = 23$) in 2015–2018. Data distribution is displayed as second and third quartiles (box plot), median (vertical crossing line) and mean (star symbol) values for spring and winter seasons.

TABLE 3 Summary of predation events for which evidence was collected for identification of the target prey. All resulted in identification of the lumpfish as prey species.

Events	Date	Evidence	ID
1, 2	April 30, 2016	Prey fragments	NKW-0712
3, 4	April 30, 2016	Photographs	NKW-0712
5, 6	April 12, 2017	Photograph	NKW-0572
7	April 5, 2018	Prey fragments	NKW-0004
8–10	April 9, 2018	Prey fragments	NKW-0004
11	April 9, 2018	Photographs	NKW-0912
12	April 11, 2018	Photographs	NKW-0079
13	April 12, 2018	Prey fragments	NKW-0910
14	April 13, 2018	Aerial footage	Unidentified subadult
15	April 13, 2018	Aerial footage	NKW-0998
16	April 14, 2018	Aerial footage	NKW-0908
17	April 15, 2018	Aerial footage	Unidentified subadult
18, 19	April 22, 2018	Prey fragments	NKW-0715
20	April 22, 2018	Prey fragments	NKW-0716
21	April 24, 2018	Prey fragments	Y093
22	April 24, 2018	Photographs	Y093

4 | DISCUSSION

We have shown that a subset of killer whales seasonally visited Andfjord in March–April of 2014–2018 and remained in the area for several days and up to several weeks. Most of these whales were also encountered at herring wintering grounds in the same years. This further supports the idea that killer whales make adaptive seasonal movements in response to spatial and temporal changes in prey availability in this region (Similä et al., 1996). Marked seasonality and predominance of foraging in spring suggests the use of a seasonally reliable food source. All 22 predation events where the prey was identified revealed lumpfish to be the target prey species in March and April.

The lumpfish is widely distributed throughout the Norwegian Sea (Holst, 1993). It remains offshore often at abyssal depths most of the year and migrates to coastal breeding areas from April to July (Davenport, 1985). Due to dorsal muscles allowing only limited locomotory abilities (Davenport & Kjørsvik, 1986), the lumpfish travels at low swimming speeds (i.e., 0.72 km/hr in Mitamura et al., 2012). Spawning male and female lumpfish are about 28–30 and 40–45 cm in length, respectively (Albert et al., 2002). Spent females may complete spawning over a period of a few weeks and then return to the open ocean, or alternatively may visit adjacent fjords to lay eggs at multiple locations (Goulet, Green, & Shears, 1986; Mitamura et al., 2012). Males then guard the nest until the eggs hatch 6–10 weeks later (Davenport 1985). These movement patterns suggest seasonal increased availability of the lumpfish nearshore in spring, as supported by exclusive seasonal fishery from April through mid-June for this species (Bertelsen, 1994).

To the best of our knowledge, the lumpfish has not hitherto been reported as a prey species of Norwegian killer whales, despite previous analysis of 95 stomach contents (Christensen, 1982) and observational studies (e.g., Similä & Ugarte, 1993; Similä et al., 1996). However, the lumpfish was listed as prey of killer whales off West Greenland, where stomachs of 30 harvested killer whales contained only lumpfish (Laidre, Heide-Jørgensen, & Orr, 2006), and off Iceland from sporadic predation records (Samarra et al., 2018). Our results list the lumpfish as prey for killer whales in an additional region of the North Atlantic. Lumpfish repeatedly recorded as prey within single season ($n = 16$ records in 2018) and across years (2016–2018; Figure 5, Table 3) in this study suggests consistency of this seasonal dietary pattern, likely in response to the increased availability of lumpfish when migrating horizontally towards spawning areas.

In spring, not only were encountered killer whales and predation records consistent across seasons, but observed foraging behavior was in striking contrast to cooperative herring feeding strategies observed from November to February in the same region (Jourdain & Vongraven, 2017). When feeding on herring, killer whales aggregate in a tight group formation for coordinated school-herding, prior to slapping the fish with the underside of their flukes and then individually consuming stunned fish (Similä, 1997; Similä & Ugarte, 1993). Similar hunting behavior of killer whales feeding on schooling mackerel in the Norwegian Sea in July–August has also been observed (Nøttestad et al., 2014). In contrast, killer whales foraging on solitary lumpfish (Davenport, 1985) were spread over several hundred meters and aerial footage supported individual rather than group-based foraging. This behavior appears more similar to that of killer whales feeding on loose patches of Atlantic salmon in the Lofoten region, Norway (Vester & Hammerschmidt, 2013). This is in further support of killer whales adapting feeding strategies to behavioral traits of target prey (e.g., Baird & Dill, 1995; Samarra & Miller, 2015; Similä & Ugarte, 1993). Lumpfish-feeding killer whales also adopted smaller group sizes in spring than when observed at herring wintering grounds (Figure 4). By maximizing energetic gain, smaller groups could be adaptive to foraging on a scattered prey resource as opposed to highly concentrated wintering herring, where a more cooperative foraging strategy is required (Nøttestad & Axelsen, 1999; Nøttestad, Ferno, & Axelsen, 2002). Our results bring the first robust evidence, featuring individual records of site fidelity, behavior, and predation over several years, for seasonal adaptation and switching between alternative prey resources in Norwegian waters.

The NSS herring is found in most dense aggregations on its wintering grounds, compared to lower densities due to looser schools on spawning and feeding grounds (Nøttestad et al., 1996). The NSS herring is also subject to important changes in fat content throughout its life cycle with the highest rate of energy depletion occurring during the

FIGURE 5 Sample of photographs used as evidence supportive of killer whale predation on lumpfish in Andfjord. All photos were taken in April 2016–2018; (a) Lumpfish observed below the surface prior to capture and consumption by adult male killer whale NKW-0572; (b) Adult male killer whale NKW-0712 catching a lumpfish; (c) Adult female-sized killer whale feeding on the carcass of a lumpfish; (d) A subadult killer whale consumed the inner part of a lumpfish before discarding the carcass.

spawning migration. As such, herring fat content is about 22% when entering the wintering area in September and is reduced to eight to 10% after spawning in February–April (Slotte, 1999). As herring both loses fat content and becomes more dispersed in spring, lumpfish with fat content ranging from >8% for gravid females to 20% for males (Davenport & Kjørsvik, 1986) enters the coastal waters to spawn. Some killer whale groups may have diversified their diet and be able to switch to foraging on this alternative, seasonally abundant and apparently relatively nutritious, prey.

Our results showed that at least 75 different killer whales have adapted to the seasonality of spawning lumpfish in Andfjord, returning year after year for temporary residence and foraging (Figures 2 and 3, Table 2). This represents 7.7% of the total number of marked individuals identified in the study area between 2007 and 2018 (Jourdain & Karoliussen, 2018). This number should be considered as a baseline minimum since it does not include unmarked individuals and calves also observed feeding on lumpfish during this study. Additionally, because this study was entirely based on opportunistic efforts and covered a relatively small area, and because killer whales may temporarily travel away from feeding sites or display group-specific preferences in habitat use, individuals or entire groups may have been undetected (Kuningas, Similä, & Hammond, 2014). While the lumpfish is a common species along the Norwegian coast with spawning grounds found in multiple regions (Davenport, 1985; Holst, 1993) it could very well be part of the diet of killer whales elsewhere and this feeding behavior could be important at the population level.

Although the NSS herring has long been assumed to be the main prey of killer whales in Norway, recent year-round data collection efforts uncovered persistent dietary specializations on pinnipeds (Jourdain et al., 2017) and on lumpfish (present study) in northern Norway. This highlights the likelihood that killer whale research conducted on a seasonal basis can miss seasonal specialized feeding behaviors and underestimating important prey resources. Our results further emphasize the importance of individual-focused studies, allowing for the detection of interindividual variations in dietary habits and prey specializations.

ACKNOWLEDGMENTS

This research was supported by grants from the Sea World and Busch Gardens Conservation fund and from Sea World and Parks Entertainment. CT thanks the team of the Genotyping and Sequencing platform shared with the LabEx CeMEB (Centre Méditerranéen de l'Environnement et de la Biodiversité, Montpellier, France) for their technical help. Photographs contributed by Northern Explorers, Marten Bril (Sea Safari Andenes AS), Camilla Ilmoni, Helge Hellesund and Daniele Zanoni (Whale Safari AS) enabled identifications of individuals that were included in this study. We thank Dag Vongraven for mapping the study area. Comments and suggestions from three anonymous reviewers substantially improved the manuscript.

ORCID

Eve Jourdain <https://orcid.org/0000-0003-2619-7799>

REFERENCES

- Albert, O. T., Torstensen, E., Bertelsen, B., Jonsson, S. T., Pettersen, I. H., & Holst, J. C. (2002). Age-reading of lumpsucker (*Cyclopterus lumpus*) otoliths: Dissection, interpretation and comparison with length frequencies. *Fisheries Research*, 55, 239–252.
- Baird, R. W., & Dill, L. M. (1995). Occurrence and behaviour of transient killer whales: seasonal and pod-specific variability, foraging behaviour, and prey handling. *Canadian Journal of Zoology*, 73, 1300–1311.
- Bertelsen, B. (1994). *Bestandsutvikling og bestandsstruktur hos rognkjeks og rognkall (Cyclopterus lumpus L.) i Sifjorden i Troms, og maskeseleksjon og fangsteffektivitet ved garn-og rusefiske etter arten* [Population structure and trajectory of lumpsucker (*Cyclopterus lumpus* L.) in Sifjorden in Troms, mask selectivity and catch efficiency in nets and traps for this species]. Fiskerikandidatoppgave [Master's thesis], Norwegian College of Fishery Science, University of Tromsø, Tromsø, Norway.
- Bigg, M. (1982). An assessment of killer whale *Orcinus orca* stocks off Vancouver Island, British Columbia. *Report of the International Whaling Commission*, 32, 655–666.
- Bisther, A., & Vongraven, D. (1995). Studies of social ecology of Norwegian killer whales (*Orcinus orca*). In A. S. Blix, L. Walløe, & Ø. Ulltang (Eds.), *Whales, seals, fish and man* (pp. 169–176). Amsterdam, The Netherlands: Elsevier Science.
- Bourque, J., Dietz, R., Sonne, C., St. Leger, J., Iverson, S., Rosing-Asvid, A., ... McKinney, M. A. (2018). Feeding habits of a new Arctic predator: Insight from full-depth blubber fatty acid signatures of Greenland, Faroe Islands, Denmark, and managed-care killer whales *Orcinus orca*. *Marine Ecology Progress Series*, 603, 1–12.
- Christensen, I. (1982). Killer whales in Norwegian coastal waters. *Report of the International Whaling Commission*, 32, 633–641.
- Cosentino, A. M. (2015). First record of Norwegian killer whales attacking and feeding on a harbour porpoise. *Marine Biodiversity Records*, 8, E108.
- Daura-Jorge, F. G., Wedekin, L. L., Piacentini, V. D. Q., & Simões-Lopes, P. C. (2005). Seasonal and daily patterns of group size, cohesion and activity of the estuarine dolphin, *Sotalia guianensis* (P. J. van Bénédén) (Cetacea, Delphinidae), in southern Brazil. *Revista Brasileira de Zoologia*, 22, 1014–1021.
- Davenport, J. (1985). *Synopsis of biological data on the lumpsucker, Cyclopterus lumpus (Linnaeus, 1758)*. (FAO Fisheries Synopsis No. 147). Rome, Italy: Food and Agriculture Organization of the United Nations.
- Davenport, J., & Kjørsvik, E. (1986). Buoyancy in the lumpsucker *Cyclopterus lumpus*. *Journal of the Marine Biological Association of the UK*, 66, 159–174.
- Domenici, P., Batty, R. S., Similä, T., & Ogam, E. (2000). Killer whales (*Orcinus orca*) feeding on schooling herring (*Clupea harengus*) using underwater tail-slaps: Kinematic analyses of field observations. *Journal of Experimental Biology*, 203, 283–294.
- Dragesund, O., Johannessen, A., & Ulltang, Ø. (1997). Variation in migration and abundance of Norwegian Spring Spawning herring (*Clupea harengus* L.). *Sarsia*, 82, 97–105.
- Foote, A. D., Newton, J., Piortney, S. B., Willerslev, E., & Gilbert, M. T. P. (2009). Ecological, morphological and genetic divergence of sympatric North Atlantic killer whale populations. *Molecular Ecology*, 18, 5207–5217.
- Foote, A. D., Similä, T., Vikingsson, G., & Stevick, P. T. (2010). Movement, site fidelity and connectivity in a top marine predator, the killer whale. *Evolutionary Ecology*, 24, 803–814.
- Foote, A. D., Vester, H., Vikingsson, G. A., & Newton, J. (2012). Dietary variation within and between populations of Northeast Atlantic killer whales, *Orcinus orca*, inferred from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analyses. *Marine Mammal Science*, 28, E472–E485.

- Foote, A. D., Vijay, N., Avila-Arcos, M. C., Baird, R. W., Durban, J. W., Fumagalli, M., ... Wolf, J. B. W. (2016). Genome-culture coevolution promotes rapid divergence of killer whale ecotypes. *Nature Communications*, 7, Article number, 11693.
- Ford, J. K. B. (1989). Acoustic behavior of resident killer whales (*Orcinus orca*) off Vancouver Island, British Columbia. *Canadian Journal of Zoology*, 67, 727–745.
- Ford, J. K. B. (2009). Killer whales *Orcinus orca*. In W. F. Perrin, B. Würsig, & J. G. M. Thewissen (Eds.), *The encyclopedia of marine mammals* (pp. 650–657). San Diego, CA: Academic Press.
- Ford, J. K. B., & Ellis, G. M. (2006). Selective foraging by fish-eating killer whales *Orcinus orca* in British Columbia. *Marine Ecology Progress Series*, 316, 185–199.
- Ford, J. K. B., & Ellis, G. M. (2014). You are what you eat: Foraging specializations and their influence on the social organization and behavior of killer whales. In J. Yamagiwa & L. Karczmarski (Eds.), *Primates and cetaceans: field research and conservation of complex mammalian societies* (pp. 75–98). Tokyo, Japan: Springer.
- Ford, J. K. B., Ellis, G. M., Barrett-Lennard, L. G., Morton, A. B., Palm, R. S., & Balcomb, K. C. (1998). Dietary specialization in two sympatric populations of killer whales (*Orcinus orca*) in Coastal British Columbia and adjacent waters. *Canadian Journal of Zoology*, 76, 1456–1471.
- Goulet, D., Green, J. M., & Shears, T. H. (1986). Courtship, spawning, and parental care behavior of the lumpfish, *Cyclopterus lumpus* L., in Newfoundland. *Canadian Journal of Zoology*, 64, 1320–1325.
- Hoelzel, A. R., Hey, J., Dahlheim, M. E., Nicholson, C., Burkanov, V., & Black, N. (2007). Evolution of population structure in a highly social top predator, the killer whale. *Molecular Biology and Evolution*, 24, 1407–1415.
- Holst, J. C. (1993). Observations on the distribution of lumpsucker (*Cyclopterus lumpus*, L.) in the Norwegian Sea. *Fisheries Research*, 17, 369–372.
- Huse, G., Fernö, A., & Holst, J. C. (2010). Establishment of new wintering areas in herring co-occurs with peaks in the 'first time/repeat spawner' ratio. *Marine Ecology Progress Series*, 409, 189–198.
- Jourdain, E. & Karoliussen, R. (2018). *Identification catalogue of Norwegian killer whales: 2007–2018*. Figshare. doi: <https://doi.org/10.6084/m9.figshare.4205226>
- Jourdain, E., & Vongraven, D. (2017). Humpback whale (*Megaptera novaeangliae*) and killer whale (*Orcinus orca*) feeding aggregations for foraging on herring (*Clupea harengus*) in Northern Norway. *Mammalian Biology - Zeitschrift für Säugetierkunde*, 86, 27–32.
- Jourdain, E., Vongraven, D., Bisther, A., & Karoliussen, R. (2017). First longitudinal study of seal-feeding killer whales (*Orcinus orca*) in Norwegian coastal waters. *PLoS One*, 12, e0180099.
- Knebelsberger, T., Landi, M., Neumann, H., Kloppmann, M., Sell, A. F., Campbell, P. D., ... Costa, F. O. (2014). A reliable DNA barcode reference library for the identification of the North European shelf fish fauna. *Molecular Ecology Resources*, 14, 1060–1071.
- Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33, 1870–1874.
- Kuningas, S., Similä, T., & Hammond, P. S. (2014). Population size, survival and reproductive rates of northern Norwegian killer whales (*Orcinus orca*) in 1986–2003. *Journal of the Marine Biological Association of the UK*, 94, 1277–1291.
- Laidre, K. L., Heide-Jørgensen, M. P., & Orr, J. R. (2006). Reactions of narwhals, *Monodon monoceros*, to killer whale, *Orcinus orca*, attacks in the Eastern Canadian Arctic. *Canadian Field-Naturalist*, 120, 457–465.
- Mitamura, H., Thorstad, E. B., Uglem, I., Bjørn, P. A., Økland, F., Næsje, T. F., ... Arai, N. (2012). Movements of lumpsucker females in a northern Norwegian fjord during the spawning season. *Environmental Biology of Fishes*, 93, 475–481.
- Morin, P. A., Archer, F. I., Foote, A. D., Foote, A. D., Vilstrup, J., Allen, E. E., ... Harkins, T. (2010). Complete mitochondrial genome phylogeographic analysis of killer whales (*Orcinus orca*) indicates multiple species. *Genome Research*, 20, 908–916.
- Nøttestad, L., & Axelsen, B. E. (1999). Herring schooling manoeuvres in response to killer whale attacks. *Canadian Journal of Zoology*, 77, 1540–1546.
- Nøttestad, L., Aksland, M., Beltestad, A., Fernö, A., Johannessen, A., & Arve Misund, O. (1996). Schooling dynamics of Norwegian Spring Spawning herring (*Clupea harengus* L.) in a coastal spawning area. *Sarsia*, 80, 277–284.
- Nøttestad, L., Ferno, A., & Axelsen, B. E. (2002). Digging in the deep: Killer whales' advanced hunting tactic. *Polar Biology*, 25, 939–941.
- Nøttestad, L., Sivle, L. D., Krafft, B. A., Langård, L., Anthonypillai, V., Bernasconi, M., ... Fernö, A. (2014). Prey selection of off-shore killer whales *Orcinus orca* in the Northeast Atlantic in late summer: Spatial associations with mackerel. *Marine Ecology Progress Series*, 499, 275–283.
- Olesiuk, P. F., Bigg, M. A., & Ellis, G. M. (1990). Life history and population dynamics of resident killer whales *Orcinus orca* in the coastal waters of British Columbia and Washington State. *Report of the International Whaling Commission*, 12, 209–243.
- Parsons, K. M., Durban, J. W., Burdin, A. M., Burkanov, V. N., Pitman, R. L., Barlow, J., ... Wade, P. R. (2013). Geographic patterns of genetic differentiation among killer whales in the Northern North Pacific. *Journal of Heredity*, 104, 737–754.

- R Core Team. (2016). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <http://www.R-project.org>
- Riesch, R., Barrett-Lennard, L. G., Ellis, G. M., Ford, J. K. B., & Deecke, V. B. (2012). Cultural traditions and the evolution of reproductive isolation: ecological speciation in killer whales? *Biological Journal of the Linnean Society*, 106, 1–17.
- Samarra, F. I. P., & Miller, P. J. O. (2015). Prey-induced behavioural plasticity of herring-eating killer whales. *Marine Biology*, 162, 809–821.
- Samarra, F. I. P., Basso, M., Beesau, J., Elíasdóttir, M. Ó., Gunnarsson, K., Mrusczok, M.-T., ... Víkingsson, G. A. (2018). Prey of killer whales (*Orcinus orca*) in Iceland. *PLoS ONE*, 13(12), e0207287.
- Samarra, F. I. P., Vighi, M., Aguilar, A., & Víkingsson, G. A. (2017). Intra-population variation in isotopic niche in herring-eating killer whales off Iceland. *Marine Ecology Progress Series*, 564, 199–210.
- Sambrook, J., Fritsch, E. F., & Maniatis, T. (1989). *Molecular cloning: A laboratory manual*. New York, NY: Cold Spring Harbor Laboratory Press.
- Similä, T. (1997). Sonar observations of killer whales (*Orcinus orca*) feeding on herring schools. *Aquatic Mammals*, 23, 119–126.
- Similä, T., & Ugarte, F. (1993). Surface and underwater observations of cooperatively feeding killer whales in northern Norway. *Canadian Journal of Zoology*, 71, 1494–1499.
- Similä, T., Holst, J. C., & Christensen, I. (1996). Occurrence and diet of killer whales in northern Norway: Seasonal patterns relative to the distribution and abundance of Norwegian Spring-Spawning Herring. *Canadian Journal of Fisheries and Aquatic Sciences*, 53, 769–779.
- Slotte, A. (1999). Differential utilisation of energy during wintering and spawning migration in Norwegian Spring-Spawning herring. *Journal of Fish Biology*, 54, 338–355.
- Sparks, J. S., & Smith, W. L. (2004). Phylogeny and biogeography of cichlid fishes (Teleostei: Perciformes: Cichlidae). *Cladistics*, 20, 501–517.
- Stenersen, J., & Similä, T. (2004). *Norwegian killer whales*. Oslo, Norway: Tringa Forlag.
- Van Opzeeland, I. C., Corkeron, P. J., Leyssen, T., Similä, T., & Van Parijs, S. M. (2005). Acoustic behaviour of Norwegian killer whales, *Orcinus orca*, during carousel and seiner foraging on spring-spawning herring. *Aquatic Mammals*, 31, 110–119.
- Van Parijs, S. M., Leyssen, T., & Similä, T. (2004). Sounds produced by Norwegian killer whales, *Orcinus orca*, during capture. *Journal of the Acoustical Society of America*, 116, 557–560.
- Vester, H., & Hammerschmidt, K. (2013). First record of killer whales (*Orcinus orca*) feeding on Atlantic salmon (*Salmo salar*) in northern Norway suggest a multi-prey feeding type. *Marine Biodiversity Records*, 6, 1–5.
- Vongraven, D., & Bisther, A. (2014). Prey switching by killer whales in the north-east Atlantic: Observational evidence and experimental insights. *Journal of the Marine Biological Association of the United Kingdom*, 94, 1357–1365.
- Weinrich, M., Martin, M., Griffiths, R., Bove, J., & Schilling, M. (1997). A shift in distribution of humpback whales, *Megaptera novaeangliae*, in response to prey in the southern Gulf of Maine. *Fishery Bulletin*, 95, 826–836.

How to cite this article: Jourdain E, Karoliussen R, de Vos J, Zakharov SE, Tougard C. Killer whales (*Orcinus orca*) feeding on lumpfish (*Cyclopterus lumpus*) in northern Norway. *Mar Mam Sci*. 2019;1–14. <https://doi.org/10.1111/mms.12618>

PAPER III

First longitudinal study of seal-feeding killer whales (*Orcinus orca*)
in Norwegian coastal waters.

Jourdain, E., Vongraven, D., Bisther, A., and Karoliussen, R. (2017).

PloS One, 12(6): e0180099.

doi: 10.1371/journal.pone.0180099.

RESEARCH ARTICLE

First longitudinal study of seal-feeding killer whales (*Orcinus orca*) in Norwegian coastal waters

Eve Jourdain^{1☯*}, Dag Vongraven^{2☯}, Anna Bisther^{3‡}, Richard Karoliussen^{1‡}

1 Norwegian Orca Survey, Andenes, Norway, **2** Norwegian Polar Institute, Tromsø, Norway, **3** Reportagebörsen, Gothenburg, Sweden

☯ These authors contributed equally to this work.

‡ These authors also contributed equally to this work.

* eve@norwegianorcasurvey.no

Abstract

Killer whales (*Orcinus orca*) have been documented preying on either fish or marine mammals in several regions, suggesting that this odontocete species has the ability to specialize on different types of prey. Off Norway, killer whales have been shown to rely on the Atlantic herring (*Clupea harengus*) as a main prey resource. Infrequent observations have revealed seals as an additional component of their diet, yet the extent of predation on marine mammals has remained largely unknown. Here, we present the findings of 29 years of photographic and observational data on seal-feeding killer whale groups identified in Norwegian coastal waters. Four groups have been observed preying and feeding on seals over several years, taking both harbor (*Phoca vitulina*) and grey (*Halichoerus grypus*) seals. These stable groups are shown to adopt small group sizes, were typically observed in near-shore areas and were not encountered on herring wintering grounds. Behavioral and social traits adopted by these groups are similar to those of pinniped-feeding killer whales from other regions. The potential ecological reasons and the extent of such prey specializations are discussed.

OPEN ACCESS

Citation: Jourdain E, Vongraven D, Bisther A, Karoliussen R (2017) First longitudinal study of seal-feeding killer whales (*Orcinus orca*) in Norwegian coastal waters. PLoS ONE 12(6): e0180099. <https://doi.org/10.1371/journal.pone.0180099>

Editor: Yan Robert-Coudert, Centre National de la Recherche Scientifique, FRANCE

Received: April 10, 2017

Accepted: June 9, 2017

Published: June 30, 2017

Copyright: © 2017 Jourdain et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data from Figshare are now publicly available at https://figshare.com/articles/Data_from_First_longitudinal_study_of_seal-feeding_killer_whales_Orcinus_orca_in_Norwegian_coastal_waters/5039280 (DOI: [10.6084/m9.figshare.5039280](https://doi.org/10.6084/m9.figshare.5039280)).

Funding: Fieldwork conducted from 1988 through 1993 was partly funded by the Marine Mammal Program of the National Research Council (http://www.forskningsradet.no/en/Home_page/1177315753906) to DV and AB. From 2014

Introduction

Killer whales (*Orcinus orca*) are cosmopolitan apex predators, preying on over 140 prey species including bony and cartilaginous fish, squids, mammals, reptiles and birds throughout their cosmopolite range [1, 2]. Despite a broad diet, some local populations feed on a narrow range of prey, adapting feeding strategies to prey type and availability [3–6].

In coastal waters of the eastern North Pacific, the most extensive studies have been focused on two sympatric ecotypes of killer whales, differing in morphology, pigmentation, acoustics, social behavior, group size, diet, movement patterns and genetics [4, 7–13]. The so-called resident type appears to be exclusively fish-feeding, preferentially preying on salmon species, whereas the transient type preys exclusively on marine-mammals [4, 10, 12, 14].

More recently, similar ecological specializations have been identified in other regions; off southeastern Alaska [5], Russian Far East [15] and in Antarctica [6, 16, 17]. Interestingly, killer

through 2016, fieldwork was financed by grants from Spare-Bank 1 Nord-Norge (<https://www.sparebank1.no/bank/privat.html>) to RK and EJ, Sea World and Busch Gardens Conservation fund (<https://swbg-conservationfund.org>) to RK and EJ, and Sea World and Parks Entertainment (<https://seaworldparks.com>) to EJ and RK. One of the authors is also employed, at the time of writing, by a commercial company being Reportagebørsen. The two commercial funding organizations only provided financial support to the study, and Reportagebørsen only provides AB with support in the form of salary. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: Part of the funding was from a commercial source, being Sea World and Busch Gardens Conservation fund and Sea World and Parks Entertainment. AB is also employed (at the time of writing) by commercial company Reportagebørsen. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

whales thought to largely or exclusively specialize on pinniped-prey seem to adopt identical behavioral and social adaptations as described for transient killer whales [18–20]. A convergence of plastic traits has been suggested as an adaptation to the type of prey being hunted [21–23].

In the North Atlantic, similar variations in killer whale feeding behavior have been suggested but never sufficiently documented, and a possible ecotype delineation is less clear [24, 25]. Based on analyses of nitrogen stable isotopes and tooth wear in samples from museum and stranded specimen, Foote and colleagues [26] suggested the presence of a generalist Type 1 and a marine-mammal specialist Type 2, the two being partly sympatric in some overlapping regions of their respective ranges. Killer whales ranging off Norway, Iceland and in the North Sea were proposed to be the Type 1, having the Atlantic herring (*Clupea harengus*) as a main prey resource but with individual and/or group variations in the proportions of prey items consumed [26]. An ecological gradient was suggested within this Type 1, with some groups persistently feeding on higher trophic level prey.

Off Norway, research conducted over the past three decades has largely focused on areas where killer whales congregate seasonally in large numbers owing to abundance of the Norwegian Spring Spawning (NSS) stock of the Atlantic herring [27, 28]. Seasonal movement pattern, site fidelity of identified individuals as well as complex coordinated feeding behaviors highlighted killer whales as herring specialists [27–29]. Sporadic predation observations also suggested additional types of prey including seals and harbor porpoise that support individuals feeding at higher trophic levels [24, 30, 31]. However, because these opportunistic observations often lack identification of individuals, the potential ecological specializations of marine mammal eating killer whales in Norwegian waters has remained largely undocumented.

In this study, we document identified killer whales that have been persistently feeding on seals for a minimum of 30 years in Norwegian coastal waters. To determine if these whales displayed similar adaptations to pinniped-feeding killer whales in other regions, and as a first step towards assessing the scope of prey specialization, we looked into specific occurrence patterns, foraging behavior and social structure. Using photo-identification and predation data collected from 1988 through 2016, the specific aims of the study were to 1) highlight apparent prey specialization of seal-eating killer whales and 2) provide the first substantial baseline information about killer whale predation on pinnipeds in Norwegian waters, while contributing data on behavioral characteristics of seal-feeding killer whales from a newly investigated region.

Materials and methods

Data collection

The data set used for this study consisted of records of photo-identified individual killer whales and observations of predation events from different periods and various areas (Fig 1) as follows:

1. Long-term killer whale study initiated in 2013 in northern Norway [34]:
Records of predation events involving identified killer whales were collected year round off Andenes, northern Norway (approximately 69°19'N, 16°08'E; Fig 1), as part of a study on the foraging ecology of killer whales. Field efforts began after a sighting report from a network of local people including fishermen, all-seasons whale-watching companies and local inhabitants. Data collection was also conducted seasonally further north, off Troms (69°50'N, 18°30'E), in October–November 2014, 2015 and 2016. On each encounter, information including date, time, location, group size estimate and group composition was

Fig 1. Important prey areas and observations of seal-feeding killer whale groups. The left panel (A) shows seasonal herring grounds since 1990 while the middle panel (B) shows areas of abundance of coastal seals since 1990 [32]. The right panel (C) shows locations of K and KI whales sightings since 1988. (The map was made in ESRI ArcGIS software [33] using public domain map data @ naturalearthdata.com).

<https://doi.org/10.1371/journal.pone.0180099.g001>

logged. Using a DSLR camera equipped with a telephoto lens, individual killer whales were systematically photographed throughout the encounters. On a few occasions, highly distinctive individuals were identified from direct observations in the field. Behavior and predation events were recorded using *Ad libitum* sampling methods [35]. Surfacing whales displaying speed bursts, sudden changes in direction and circling behaviors often indicated ongoing predation incidents that were documented with photographs and/or video recordings. Whenever possible, prey species was visually identified on the field or through photographs. When the whales had finished feeding and moved on, the feeding spot was approached and prey remains collected using a fine-mesh dip net to confirm prey consumption. Encounters ended when weather, light and/or sea conditions became too poor to continue or if visual contact with the whales was lost.

2. Photo-identification study based on central Norwegian coast 1987–1993 [24]:
A photo-identification study was initiated in 1987 in areas covering the herring spawning ground that was mainly located off the Møre coast, and approximately 100 nm of coastline north from this area (between latitudes 62°45'N–64°05'N and longitudes 6°–9°15'E, Fig 1). From 1987 through 1993, the collecting of ID photographs was opportunistic and conducted from November through March, resulting in a catalogue of 605 individuals [24]. All photographs were taken with analogue cameras, fitted with 300 mm telephoto lenses and predation events were recorded *Ad libitum* [35]. Information including date, time, location and group size estimate was noted during each encounter. Additional fieldwork was conducted around harbor and grey seal pupping colonies in June and September–October, respectively. Field studies were conducted in Møre areas in the years 1987–1993 sporadically during the months November through March. In the period 1990–1993 field studies during October through December were carried out in the Tysfjord area (68°16'N, 15°52'E) for four consecutive years.
3. Photo-identification study based in northern Norway 1986–2003 [36]:
A catalogue of 585 individual killer whales photographically identified in northern Norway, as part of a long-term killer whale study, was available for review. This catalogue consisted of photographs taken in northern Norway in the former NSS herring wintering ground in

Tysfjord-Ofotfjord-Vestfjord from 1986 to 2003 (work of Similä and colleagues mentioned in [36]).

4. Opportunistic imagery from other sources:

One video containing unidentifiable whales preying on seals was collected from a wildlife photographer in 1988 at Orskjera, off the Møre coast (63°7'N, 7°15'E). Substantial amounts of high resolution opportunistic photographs, taken from 2010 through 2016 and accompanied by metadata on date and location, were also provided by wildlife photographers. Photographs were taken in all seasons in coastal waters between latitudes 62° and 72° (Fig 1). Six videos showing killer whales hunting seal prey, including two accompanied by high resolution photographs, were also available for review.

Data analysis

All accounts of killer whales preying on seals were recorded, including both successful and failed predation events. Following photo-identification protocols developed by Bigg [7], individual killer whales specifically involved in these predation events were identified using the shape, pigmentation and scarring patterns of the dorsal fin and saddle patch. Photographs of both left and right sides were used for individual identification. Best photographs for each individual in each encounter was selected and rated for quality (Q), ranging from 0 (lowest) to 2 (highest) based on sharpness, contrast and angle of the dorsal fin in relation to the camera. All further sightings of individual killer whales identified as seal-feeding were recorded, including information on date and location. Resulting sighting histories, including only identifications of quality Q1 and Q2, were further used to map temporal and spatial occurrence patterns.

A group of whales was defined as all individual killer whales visible within the range of the observer and displaying coordinated activities (as per [11, 22, 37]). To assess group size, only encounters for which all individuals were believed to be identified were used. Group sizes adopted by seal-feeding vs fish-feeding killer whales were compared using a Mann-Whitney-Wilcoxon test performed in R [38].

Association patterns between individuals and persistence of groups were analyzed utilizing SOCPROG 2.6, a software specifically designed to study social organization of animal populations [39]. Association indices were calculated for all pairs of individuals using the Simple-Ratio Index, with association defined as presence in the same group within a sampling period (i.e an encounter on a certain date) and with all group members presumed identified [40]. From opportunistic photographs, group membership could be determined by looking at individuals captured within frames, or by pooling frames within certain time intervals [21]. Unreliably marked calves were excluded from these analyses and only Q2 identifications were accounted for. In order to reduce bias associated with small sample sizes, restrictions were also set to account exclusively for individuals observed in at least four sampling periods (as per [41]). The resulting association matrix was summarized in a sociogram with nodes representing individuals and edges indicating incidence and strength of relationship between each pair of whales [42]. Permuting associations within samples enabled testing for potential preferred associations among individuals [43, 44]. If the standard deviation (SD) calculated on the pairwise associations was significantly greater for the real association indices than the random data set, the null hypothesis of individuals associating at random was rejected. To further investigate temporal patterns in detected associations, the lagged association rate was calculated [45, 46]. This analysis calculates the probability for two animals to be associated again after some time, given that they have been associated before. The null association rate, which is the expected value of the lagged association rate if individuals associate at random, was plotted for comparison.

Results

A total of 23 predation events of killer whales on seal prey in Norwegian coastal waters was documented from 15,000 photographs taken over 300 days of research effort between March 1988 and June 1992 [24], and 158,000 photographs, six videos and field notes taken over 468 days between July 2010 and December 2016. Nineteen of these predation events resulted in kills and consumption of the prey, and four were unsuccessful chases. Positive identification of individual killer whales involved was possible for 14 of these 23 predation events, and these whales were further re-sighted on 23 encounters, with no observed predation. Both young and adult individuals of grey (*Halichoerus grypus*) and harbor seals (*Phoca vitulina*) have been observed as part of the whales' chosen pinniped prey.

Identification of seal-eating killer whales

Five killer whales, including so-called KI-03, KI-05 and KI-06, were first identified chasing harbor seals in March 1988 off the central coast of Norway. Three more individuals were identified in this group while foraging in a grey seal pupping colony in October 1990, before being identified as part of a larger group of 11 whales in March 1991. KI-03, KI-05 and KI-06 were re-sighted on several occasions between May 2011 and August 2016, in northern Norway (Fig 2). To ensure consistence with the longitudinal study, all individual whales encountered with KI-03, KI-05 and KI-06 on at least one occasion between 2010 and 2016 were given similar alphanumeric codes and considered as members of the KI group.

KI whales were observed chasing or preying on pinnipeds on eight occasions (Fig 3). They were also observed twice feeding on herring at herring spawning grounds in 1991 off the Møre region, in association with other known herring-feeding killer whales, before travelling to a seal haul-out where they were observed for several hours unsuccessfully chasing harbor seals.

First identified in July 2010, individuals K1, K2, K3, K4 and K5, hereafter referred to as the K group, constitute the second assemblage of killer whales identified as seal-feeding.

Fig 2. Photographs of seal-feeding killer whales KI-03, KI-05 and KI-06. Individuals in upper photographs, observed off central coast of Norway in 1988, match individuals visible on the lower row of photographs, taken off Andenes in June 2015.

<https://doi.org/10.1371/journal.pone.0180099.g002>

Fig 3. Photograph of KI-03 preying upon a harbor seal pup in June 2015. The photograph (A) illustrates the predation observation and the inset (B) confirms effective prey consumption.

<https://doi.org/10.1371/journal.pone.0180099.g003>

K whales were observed chasing or preying on seals on six occasions and were never observed feeding on fish during the course of this study.

Predation observations for KI and K whales are summarized in [Table 1](#). There was no evidence for interaction between KI and K whales in this study.

Table 1. Summary of recorded predation events on seals where involved individual killer whales could be identified.

Date	Location	Predation type	Minimum whales identified	Seal species	Evidence
27 Mar 1988	Orskjera, Møre	Chase	KI03, KI05, KI06	Harbor seal	Video
15 Oct 1990	Froan, Trøndelag	Blubber feeding	KI03	Undetermined	Photographs
10 Mar 1991	Ona, Møre	Chase	KI03, KI05, KI06	Harbor seal	Photographs
27 Jun 2013	Stø, Nordland	Kill	KI02, KI05, KI06, KI08, KI09, KI10	Undetermined	Photographs
15 Jul 2013	Andenes, Nordland	Kill	K1, K2, K3, K4, K5	Harbor seal	Photographs
27 Jul 2013	Andenes, Nordland	Kill	KI01, KI03, KI07	Harbour seal	Photographs and video
30 May 2014	Andenes, Nordland	Kill	K1	Undetermined	Photographs
31 May 2014	Andenes, Nordland	Kill	K1, K2, K3, K4, K5	Undetermined	Photographs and video
14 Sept 2014	Vestfjord, Nordland	Chase	K1, K2, K3, K4, K5	Harbour seal	Video
27 Jun 2015	Andenes, Nordland	Kill (3)	KI01, KI02, KI03, KI04, KI05, KI06, KI07, KI08, KI09, KI10	Harbour seal (1), grey seal (1), undetermined (1)	Photographs
19 Aug 2016	Senja, Troms	Kill (2)	K1, K2, K3, K4, K5	Harbour seal (1) and undetermined (1)	Photographs and video

For encounters that occurred in the period 1988–1991, only identifications of whales re-sighted after 2010 are indicated.

<https://doi.org/10.1371/journal.pone.0180099.t001>

Table 2. Sighting histories of KI and K whales.

Group	1988	1989	1990	1991	1992	2010	2011	2012	2013	2014	2015	2016	Total
KI	1	0	1	3	1	0	1	1	4	2	2	3	19
K	0	0	0	0	0	1	0	0	2	5	3	4	15

The number of encounters with KI and K whales is given for each year during the periods 1988–1992 and 2010–2016.

<https://doi.org/10.1371/journal.pone.0180099.t002>

Occurrence pattern

From 1988 through 1992 off the central coast of Norway, KI whales were encountered on six observation days during the months of March (5) and October (1). Between 2010 and 2016 off northern Norway, these whales were observed on 13 days from May through August (Table 2 and Fig 4). K whales were also encountered on 15 occasions between 2010 and 2016, with sightings from May through October (Table 2 and Fig 4). Observations covered a maximum range of 800 km between re-sightings of KI whales and 350 km for K whales.

KI and K whales were not identified on the former herring wintering grounds in Tysfjord, where 584 individual killer whales were identified from 1986 through 2003 (work of Similä and colleagues, mentioned in [36]). They were also not encountered on the recently established herring wintering ground off Troms and Andfjord regions where 656 individuals were identified during November–February and, despite substantial effort, between 2010 and 2016 (Fig 4; [34]).

All encounters with KI and K groups documented in this study occurred within 300 meters from the shoreline or around reefs and seal haul-outs.

Fig 4. Observations of seal-feeding killer whales in relation to effort from 2010 through 2016. The plot (A) shows the cumulative sighting distribution of KI (dark bars) and K (light bars) whales while the plot (B) reports on the cumulative number of effort days spent in the field for each month, from 2010 through 2016 in northern Norway.

<https://doi.org/10.1371/journal.pone.0180099.g004>

Fig 5. Sociogram of seal-feeding killer whales observed in at least four sampling periods. Occurrence and thickness of links show association patterns between pairs of individuals, revealing the two assemblages KI and K.

<https://doi.org/10.1371/journal.pone.0180099.g005>

Group size and social organization

Mean group size for seal-feeding killer whales, accounting for both KI and K assemblages, was 5 individuals ($N = 27$, range = 3–11, median = 5). Fish-feeding killer whale groups encountered during spring-summers 2014, 2015 and 2016 averaged a group size of 13 individuals ($N = 30$, range = 6–25, median = 12), comparable to findings of Similä and colleagues [27]. The Mann-Whitney-Wilcoxon test supported a statistical difference between group size adopted by seal vs fish-feeding killer whales ($W = 782.5$, $p < 0.001$).

Sixty-eight identifications of 13 individuals over 17 sampling periods (= days) indicated two levels of association. First, individuals associated in 100% of observations enabled four apparently cohesive units to be identified. A second social level, involving temporary associations between cohesive units further indicated two distinct larger assemblages each represented by KI and K whales (Fig 5). The SD of calculated pairwise association indices was significantly higher for the real data set than for the random data ($SD_{\text{Real}} = 0.40031$, $SD_{\text{Random}} = 0.37726$, $p = 0.0010$), enabling us to reject the null hypothesis that individuals associate at random. Temporal analyses revealed stable associations over long periods of time and above the null association rate (Fig 6).

Discussion

We have documented a minimum of 13 killer whales, belonging to four stable units, and two larger assemblages (KI and K), that appear to regularly feed on seals in Norwegian coastal

Fig 6. Plots of lagged association rates. The blue plot represents our real data set while the red plot shows the expected lagged association rates if individuals associated at random.

<https://doi.org/10.1371/journal.pone.0180099.g006>

waters. Repeated observations over time of identified individuals preying on seals support persistent preference for pinniped prey over several years for K whales, and for nearly three decades for KI-03, KI-05 and KI-06. A comparison of KI and K whales' behavioral traits to the social structure, occurrence pattern and foraging behavior described for pinniped-feeding killer whales in other regions [9–11, 18, 22, 47, 48] provide further insights into the degree of prey specialization for these whales.

Observations of KI and K whales were widely distributed along the Norwegian coast, with all sightings concentrated in important breeding areas for harbor and grey seals. Major breeding sites for both pinniped species can be found in Froan (64°05'N, 09°15'E), central Norway and in the northern part of the recorded range of this study [49, 50]. With one notable exception discussed below, all encounters with KI and K whales occurred in near shore shallow areas, typical pinniped habitat [49, 50] where the whales were observed following reefs and shoreline contours, or circling haul-outs with long dives, erratic swimming patterns and no real travel direction. This 'haul-out and near shore foraging' behavior has been described for transient killer whales hunting pinnipeds in the Northeastern Pacific [9, 10], at Subantarctic Marion Island and Possession Island, Crozet [47, 48]. Even though seals are non-migratory, thus offering a constant and predictable prey resource throughout their range along the Norwegian coast, it may be more rewarding for killer whales to forage around haul-outs in the breeding season where seal pups and females regularly enter and exit the water. Pacific transient killer whales appear to adapt their range, following latitudinal clines in timing of pupping [10]. Similarly, observations in northern Norway suggested a likelihood for KI and K whales to be observed near haul-outs at times coinciding with pupping periods of harbor seals (June–July, [51]) and grey seals (September–October, [50]) in this region. The timing of observations also suggests a north-south movement pattern for these whales, which coincides with local peaks in pupping and weaning of pinnipeds. More specifically, K whales travelled from the

Nordland region where they were observed from May through September to the Troms region where sightings were recorded in late October. This same pattern was repeated in 2014, 2015 and 2016, and observations followed the cline of the grey seal pupping period, which is delayed further north and occurs as late as November–December northwards in the Troms and Finnmark regions [52].

Further evidence supporting prey specialization in KI and K whales is a group size significantly smaller than the group size of herring-feeding killer whales. Living in small groups has been suggested as a strategy to maximize energy intake for pinniped-feeding killer whales. Small group sizes seem to guarantee stealthy and cooperative prey-searching and handling, whilst optimizing prey sharing [11]. In our dataset, small groups of seal-feeding killer whales were also long lasting, as revealed by temporal analyses of social organization. This is consistent with long-term cohesiveness of basic social units found to be a characteristic of the species, regardless of the prey-type hunted (e.g. [8, 22, 37, 53]). Stable and long-lasting social groups undoubtedly provide the raw material needed for social transmission of feeding behaviors and persistence of dietary habits over the years [54]. Additionally, temporary loose associations between stable units may be necessary for social learning of hunting skills and prey handling, and may explain the larger groups that were observed on a few occasions [10, 22]. On 27 June 2015, the three groups of KI whales were observed cooperatively hunting and sharing three seals over a period of six hours. Two juveniles under three years of age and a calf of the year were recorded in the group, and social play was observed when the whales handled a harbor seal pup for 40 minutes before killing it.

Two acoustic recordings available for K whales whilst handling seals at a close distance revealed no audible sounds, but sporadic trains of echolocation clicks from the whales. This is consistent with findings for marine mammal feeding killer whales from other regions [18, 55, 56], and is in sharp contrast to the highly vocal herring-feeding killer whales [57].

KI and K whales seem to have adopted strikingly similar adaptations to other pinniped-feeding killer whales. KI whales were observed feeding on herring in the spawning grounds in 1991 where they were also seen preying on seals. Despite massive study efforts and hundreds of identifications, KI and K groups were not identified on former or current herring wintering grounds during 1986–2003 and 2013–2016, respectively [34]. This could suggest that herring does not represent a main prey resource to these whales. However, because killer whale groups may display variations in habitat usage and preferences, and because data collection so far was largely opportunistic and covering a limited area, individuals might be undetected and not available for sampling in surveyed areas. Thus, absence of KI and K whales from the herring wintering grounds remain inconclusive.

How long these two assemblages have shown this type of prey specialization, and the origins of their distinctive diet remain largely unknown, but some major factors could be at least partially explanatory. The NSS stock of herring, the main prey base for killer whales in Norwegian coastal waters, collapsed in 1970 [58, 59]. How killer whales coped with prey depletion remains unknown, however they would likely have the phenotypic plasticity to adapt to changing environments as suggested by the wide variety of prey items and elaborated feeding techniques documented throughout their range [6, 60, 61]. Whilst prey switching appears as a potential adaptive response to decreasing prey supplies for numerous predator species [62–64], it is possible that some killer whale groups had diversified their diet and developed specific foraging strategies for new prey types in response to the drastic decrease in herring biomass [65]. Von-graven and Bisther [30] discussed this as a potential explanation for the observation of KI whales observed feeding in association with known herring feeding-killer whales in the spawning areas in March 1991, before splitting off into a smaller group and travelling a few miles to inshore waters where they started chasing harbor seals. The past culling of the killer whale

population, organized off the central Norwegian coast as a management action to preserve the remaining herring, may also have played a role in the niche diversification process [66, 67]. In 1969 and 1970, 167 and 201 killer whales were culled, respectively, over a 100 x 50 km area and within a few weeks. Entire groups were apparently never culled, and catch records showed a general lack of small individuals and a skewed sex ratio of 2M:1F [67]. Due to differential roles played by key members in cetacean societies and social transmission of matrilineal-based prey specializations in killer whales, feeding strategies may have been lost in culled groups, jeopardizing feeding success if key members were removed, and promoting potential changes in diet [68, 69].

This study revealed two assemblages of killer whales that persistently fed on seal prey and displayed parallel adaptations to that of pinniped-feeding killer whales from other regions, suggesting a certain degree of prey specialization. It remains, however, unknown as to whether KI and K whales are seasonal prey-specialists, switching between different prey types, or if they specialize on mammals year-round. Despite the observations of KI and K whales fitting well with the distribution of pinniped prey over the years, all observations have been recorded within coastal areas covered by the annual migration of the NSS herring. Thus, further analyses of dietary markers in biopsy samples are warranted, and this will constitute a research priority in order to conclude on the degree of prey specialization of these whales.

Acknowledgments

We wish to thank Mellemværfet in Kristiansund and its residents for hospitality and accommodation during the fieldwork period 1988–1993. Tiu Similä gave us access to the identification catalogue from her long-term killer whale study. We are very grateful to photographers Nils Aukan, Ingar Øien, Camilla Ilmoni, Marten Bril, Inaki Aizpurua Quiroga, Laura Riem, Anders Frantzen, Camilla Figenschou, Simen Olsen, Steve Baines, Piet van den Bemd and Isabelle Dupré whose material contributed to building the sighting history of the seal-feeding killer whale groups in this study. We also thank Sea Safari Andenes, Arctic Whale Tours and Stø Safari for their involvement in this study by reporting killer whale sightings.

Author Contributions

Conceptualization: Eve Jourdain, Dag Vongraven.

Funding acquisition: Eve Jourdain, Dag Vongraven, Anna Bisther, Richard Karoliussen.

Investigation: Eve Jourdain, Dag Vongraven, Anna Bisther, Richard Karoliussen.

Methodology: Eve Jourdain, Dag Vongraven, Anna Bisther, Richard Karoliussen.

Project administration: Eve Jourdain, Dag Vongraven.

Resources: Eve Jourdain, Dag Vongraven, Richard Karoliussen.

Visualization: Eve Jourdain, Dag Vongraven.

Writing – original draft: Eve Jourdain, Dag Vongraven.

Writing – review & editing: Eve Jourdain, Dag Vongraven, Anna Bisther, Richard Karoliussen.

References

1. Forney KA, Wade PR. Worldwide distribution and abundance of killer whales. In: Estes JA, Demaster DP, Doak DF, Williams TM, Brownell RL, Jr., editors. *Whales, whaling and ocean ecosystems* 2006. p. 145–62.

2. Ford JKB. Killer whales *Orcinus orca*. In: Perrin WF, Würsig B, Thewissen JGM, editors. The encyclopedia of marine mammals. 2nd ed. San Diego, USA: Academic Press; 2009. p. 650–7.
3. Nichol LM, Shackleton DM. Seasonal movements and foraging behavior of northern resident killer whales (*Orcinus orca*) in relation to the inshore distribution of salmon (*Oncorhynchus* spp.) in British Columbia. *Canadian Journal of Zoology*. 1996; 74(6):983–91.
4. Ford JKB, Ellis GM, Barrett-Lennard LG, Morton AB, Palm RS, Balcomb KC. Dietary specialization in two sympatric populations of killer whales (*Orcinus orca*) in coastal British Columbia and adjacent waters. *Canadian Journal of Zoology*. 1998; 76(8):1456–71.
5. Saulitis E, Matkin C, Barrett-Lennard L, Heise K, Ellis G. Foraging strategies of sympatric killer whale (*Orcinus orca*) populations in Prince William Sound, Alaska. *Marine Mammal Science*. 2000; 16(1):94–109.
6. de Bruyn PJN, Tosh CA, Terauds A. Killer whale ecotypes: is there a global model? *Biological Reviews*. 2013; 88(1):62–80. <https://doi.org/10.1111/j.1469-185X.2012.00239.x> PMID: 22882545
7. Bigg M. An assessment of killer whale *Orcinus orca* stocks off Vancouver Island, British Columbia. Report of the International Whaling Commission. 1982; 32:655–66.
8. Bigg MA, Olesiuk PF, Ellis GM, Ford JKB, Balcomb KC. Social organization and genealogy of resident killer whales *Orcinus orca* in the coastal waters of British Columbia and Washington State. Report of the International Whaling Commission. 1990; 12:383–405.
9. Morton AB. A quantitative comparison of the behavior of resident and transient forms of the killer whale off the central British Columbia coast. Reports of the International Whaling Commission Special Issue. 1990:245–8.
10. Baird RW, Dill LM. Occurrence and behaviour of transient killer whales: Seasonal and pod-specific variability, foraging behaviour, and prey handling. *Canadian Journal of Zoology*. 1995; 73(7):1300–11.
11. Baird RW, Dill LM. Ecological and social determinants of group size in transient killer whales. *Behavioral Ecology*. 1996; 7(4):408–16.
12. Ford JKB, Ellis GM. You are what you eat: foraging specializations and their influence on the social organization and behavior of killer whales. In: Yamagiwa J, Karczmarski L, editors. *Primates and cetaceans: field research and conservation of complex mammalian societies*. Springer; 2014. p. 75–98.
13. Hoelzel AR, Dover GA. Genetic differentiation between sympatric killer whale populations. *Heredity*. 1991; 66:191–5.
14. Baird RW, Abrams PA, Dill LM. Possible indirect interactions between transient and resident killer whales: Implications for the evolution of foraging specializations in the genus *Orcinus*. *Oecologia*. 1992; 89:125–32. <https://doi.org/10.1007/BF00319024> PMID: 28313404
15. Filatova OA, Borisova EA, Shpak OV, Meschersky IG, Tiunov AV, Goncharov AA, et al. Reproductively isolated ecotypes of killer whales *Orcinus orca* in the seas of the Russian Far East. *Biology Bulletin*. 2015; 42(7):674–81. <https://doi.org/10.1134/s1062359015070043>
16. Pitman RL, Ensor P. Three forms of killer whales in Antarctic waters. *Journal of Cetacean Research and Management*. 2003; 5:131–9.
17. Krahn MM, Pitman RL, Burrows DG, Herman DP, Pearce RW. Use of chemical tracers to assess diet and persistent organic pollutants in Antarctic Type C killer whales. *Marine Mammal Science*. 2008; 24(3):643–63.
18. Deecke VB, Nykanen M, Foote AD, Janik VM. Vocal behaviour and feeding ecology of killer whales *Orcinus orca* around Shetland, UK. *Aquatic Biology*. 2011; 13(1):79–U186.
19. Dahlheim ME, White PA. Ecological aspects of transient killer whales *Orcinus orca* as predators in southeastern Alaska. *Wildlife Biology*. 2010; 16(3):308–22. <https://doi.org/10.2981/09-075>
20. Pitman RL, Durban JW. Cooperative hunting behavior, prey selectivity and prey handling by pack ice killer whales (*Orcinus orca*), type B, in Antarctic Peninsula waters. *Marine Mammal Science*. 2012; 28(1):16–36. <https://doi.org/10.1111/j.1748-7692.2010.00453.x>
21. Beck S, Kuningas S, Esteban R, Foote AD. The influence of ecology on sociality in the killer whale (*Orcinus orca*). *Behavioral Ecology*. 2012; 23(2):246–53.
22. Reisinger RR, Beukes C, Hoelzel AR, de Bruyn PJN. Kinship and association in a highly social apex predator population, killer whales at Marion Island. *Behavioral Ecology*. 2017; 28(3):750–9. <https://doi.org/10.1093/beheco/arx034>
23. Tavares SB, Samarra FIP, Miller PJO. A multilevel society of herring-eating killer whales indicates adaptation to prey characteristics. *Behavioral Ecology*. 2017; 28(2):500–14. <https://doi.org/10.1093/beheco/arw179>

24. Bisther A, Vongraven D, editors. Killer whales feeding on both mammals and fish: A transient, resident or opportunistic type? Poster presentation at the 14th Biennial Conference on the Biology of Marine Mammals; 2001; Vancouver, British Columbia, Canada 2001.
25. Samarra FIP, Foote AD. Seasonal movements of killer whales between Iceland and Scotland. *Aquatic Biology*. 2015; 24(1):75–9. <https://doi.org/10.3354/ab00637>
26. Foote AD, Newton J, Piethney SB, Willerslev E, Gilbert MTP. Ecological, morphological and genetic divergence of sympatric North Atlantic killer whale populations. *Molecular Ecology*. 2009; 18(24):5207–17. PMID: [20050301](#)
27. Similä T, Holst JC, Christensen I. Occurrence and diet of killer whales in northern Norway: Seasonal patterns relative to the distribution and abundance of Norwegian spring-spawning herring. *Canadian Journal of Fisheries and Aquatic Sciences*. 1996; 53(4):769–79.
28. Bisther A, Vongraven D. Studies of social ecology of Norwegian killer whales (*Orcinus orca*). In: Blix AS, Walløe L, Ulltang Ø, editors. *Whales, seals, fish and man*: Elsevier Science; 1995. p. 169–76.
29. Similä T, Ugarte F. Surface and underwater observations of cooperatively feeding killer whales in northern Norway. *Canadian Journal of Zoology*. 1993; 71:1494–9.
30. Vongraven D, Bisther A. Prey switching by killer whales in the north-east Atlantic: observational evidence and experimental insights. *Journal of the Marine Biological Association of the United Kingdom*. 2014; 94(6):1357–65.
31. Cosentino AM. First record of Norwegian killer whales attacking and feeding on a harbour porpoise. *Marine Biodiversity Records*. 2015; 8:null-null. <https://doi.org/10.1017/S1755267215000895>
32. Institute of Marine Research. Havforskningsrapporten 2016 (in Norwegian). *Fisken og havet*. 2016; Special Issue 1:203 pp.
33. ESRI. ArcGIS Desktop: Release 10.5. Redlands, CA: Environmental Systems Research Institute; 2016.
34. Jourdain E, Karoliussen R. The Norwegian Orca ID-catalogue. 2016. 10.6084/m9.figshare.4205226.
35. Altmann J. Observational study of behavior: Sampling methods. *Behaviour*. 1974; 49:227–67. PMID: [4597405](#)
36. Kuningas S, Similä T, Hammond PS. Population size, survival and reproductive rates of northern Norwegian killer whales (*Orcinus orca*) in 1986–2003. *Journal of the Marine Biological Association of the United Kingdom*. 2014; 94(6):1277–91. <https://doi.org/10.1017/s0025315413000933>
37. Baird RW, Whitehead H. Social organization of mammal-eating killer whales: group stability and dispersal patterns. *Canadian Journal of Zoology*. 2000; 78(12):2096–105.
38. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>. 2013.
39. Whitehead H. SOCPROG programs: analysing animal social structures. *Behavioral Ecology and Sociobiology*. 2009; 63(5):765–78. <https://doi.org/10.1007/s00265-008-0697-y>
40. Ginsberg JR, Young TP. Measuring association between individuals or groups in behavioural studies. *Animal Behaviour*. 1992; 44:377–9.
41. Ottensmeyer CA, Whitehead H. Behavioural evidence for social units in long-finned pilot whales. *Canadian Journal of Zoology*. 2003; 81:1327–38.
42. Whitehead H, Dufault S. Techniques for Analyzing Vertebrate Social Structure Using Identified Individuals: Review and Recommendations. In: Peter J.B. Slater JSRCTS, Timothy JR, editors. *Advances in the Study of Behavior*. Volume 28: Academic Press; 1999. p. 33–74.
43. Bejder L, Fletcher D, Brager S. A method for testing association patterns of social animals. *Animal Behaviour*. 1998; 56:719–25. <https://doi.org/10.1006/anbe.1998.0802> PMID: [9784222](#)
44. Whitehead H. Testing association patterns of social animals. *Animal Behaviour*. 1999; 57:F26–F9. <https://doi.org/10.1006/anbe.1999.1099> PMID: [10373270](#)
45. Whitehead H. Selection of models of lagged identification rates and lagged association rates using AIC and QAIC. *Communications in Statistics-Simulation and Computation*. 2007; 36(6):1233–46.
46. Whitehead H. Investigating structure and temporal scale in social organizations using identified individuals. *Behavioral Ecology*. 1995; 6(2):199–208. <https://doi.org/10.1093/beheco/6.2.199>
47. Reisinger RR, De Bruyn PJN, Tosh CA, Oosthuizen WC, Mufanadzo NT, Bester MN. Prey and seasonal abundance of killer whales at sub-Antarctic Marion Island. *African Journal of Marine Science*. 2011; 33(1):99–105.
48. Guinet C. Intentional stranding apprenticeship and social play in killer whales *Orcinus orca*. *Canadian Journal of Zoology*. 1991; 69:2712–6.

49. Nilssen KT, Skavberg NE, Poltermann M, Haug T, Härkönen T, Henriksen G. Status of harbour seals (*Phoca vitulina*) in mainland Norway. NAMMCO Scientific Publications. 2010; 8:61–9.
50. Nilssen KT, Haug T. Status of grey seals (*Halichoerus grypus*) in Norway. NAMMCO Scientific Publications. 2007; 6:23–31.
51. Bjørge A. Status of the harbor seal *Phoca vitulina* in Norway. Biological Conservation. 1991; 58(2):229–38. [https://doi.org/10.1016/0006-3207\(91\)90121-o](https://doi.org/10.1016/0006-3207(91)90121-o)
52. Haug T, Henriksen G, Kondakov A, Mishin V, Nilssen KT, Røv N. The status of grey seals *Halichoerus grypus* in North Norway and on the Murman Coast, Russia. Biological Conservation. 1994; 70:59–67.
53. Esteban PR. Abundancia, estructura social y parámetros de historia natural de las orcas (*Orcinus orca*) del Estrecho de Gibraltar University of Cadiz, Spain; 2008.
54. Riesch R, Barrett-Lennard LG, Ellis GM, Ford JKB, Deecke VB. Cultural traditions and the evolution of reproductive isolation: ecological speciation in killer whales? Biological Journal of the Linnean Society. 2012; 106:1–17.
55. Barrett-Lennard LG, Ford JKB, Heise KA. The mixed blessing of echolocation: Differences in sonar use by fish-eating and mammal-eating killer whales. Animal Behaviour. 1996; 51(3):553–65.
56. Guinet C. Comportement de chasse des orques *Orcinus orca* autour des îles Crozet. Canadian Journal of Zoology. 1992; 70:1656–67.
57. Simon M, McGregor PK, Ugarte F. The relationship between the acoustic behaviour and surface activity of killer whales (*Orcinus orca*) that feed on herring (*Clupea harengus*). Acta Ethologica. 2007; 10(2):47–53.
58. Dragesund O, Johannessen A, Ulltang Ø. Variation in migration and abundance of Norwegian spring spawning herring (*Clupea harengus* L.). Sarsia. 1997; 82:97–105.
59. Dragesund O, Hamre J, Ulltang Ø. Biology and population dynamics of the Norwegian spring-spawning herring. Rapports et Procès-verbaux des Réunions Conseil International pour l'Exploration de la Mer. 1980; 177(43–71).
60. Samarra FIP, Miller PJO. Prey-induced behavioural plasticity of herring-eating killer whales. Marine Biology. 2015; 162(4):809–21. <https://doi.org/10.1007/s00227-015-2626-8>
61. Whitman DW, Agrawal AA. What is phenotypic plasticity and why is it important? In: Whitman DW, Ananthakrishnan TN, editors. Phenotypic plasticity in insects. Enfield: Science Publishers; 2009. p. 1–63.
62. Panzacchi M, Linnell JDC, Odden J, Odden M, Andersen R. When a generalist becomes a specialist: patterns of red fox predation on roe deer fawns under contrasting conditions. Canadian Journal of Zoology-Revues Canadienne De Zoologie. 2008; 86(2):116–26. <https://doi.org/10.1139/z07-120>
63. Randa LA, Cooper DM, Meserve PL, Yunker JA. Prey switching of sympatric canids in response to variable prey abundance. Journal of Mammalogy. 2009; 90(3):594–603. <https://doi.org/10.1644/08-mamm-a-092r1.1>
64. Tinker MT, Bentall G, Estes JA. Food limitation leads to behavioral diversification and dietary specialization in sea otters. Proceedings of the National Academy of Sciences of the United States of America. 2008; 105(2):560–5. <https://doi.org/10.1073/pnas.0709263105> PMID: 18195370
65. Piltz SH, Porter MA, Maini PK. Prey Switching with a Linear Preference Trade-Off. SIAM Journal on Applied Dynamical Systems. 2014; 13(2):658–82. <https://doi.org/10.1137/130910920>
66. Marine Research Institute. Catch of killer whales in the North Atlantic 1938–1981. 2016.
67. Vongraven D, Bisther A. Possible effects of previous catch on the present population of Norwegian killer whales (*Orcinus orca*). In: Blix AS, Walløe L, Ulltang Ø, editors. Whales, seals, fish and man: Elsevier Science; 1995. p. 177–9.
68. Williams R, Lusseau D. A killer whale social network is vulnerable to targeted removals. Biology Letters. 2006; 2(4):497–500. <https://doi.org/10.1098/rsbl.2006.0510> PMID: 17148272
69. Whitehead H, Rendell L. Movements, habitat use and feeding success of cultural clans of South Pacific sperm whales. Journal of Animal Ecology. 2004; 73(1):190–6.

PAPER IV

Isotopic niche differs between seal and fish-eating
killer whales (*Orcinus orca*) in northern Norway.

Jourdain, E., Andvik, C., Karoliussen, R., Ruus, A., Vongraven, D., & Borgå, K. (2020).

Ecology and Evolution. 00: 1-13.

doi://10.1002/ece3.6182.

Isotopic niche differs between seal and fish-eating killer whales (*Orcinus orca*) in northern Norway

Eve Jourdain¹ | Clare Andvik² | Richard Karoliussen¹ | Anders Ruus^{2,3} |
Dag Vongraven⁴ | Katrine Borgå²

¹Norwegian Orca Survey, Andenes, Norway

²Department of Biosciences, University of Oslo, Oslo, Norway

³Norwegian Institute for Water Research, Oslo, Norway

⁴Norwegian Polar Institute, Tromsø, Norway

Correspondence

Eve Jourdain, Norwegian Orca Survey,
Daniel Olaisensvei 6, N-8480 Andenes,
Norway.
Email: post@norwegianorcasurvey.no

Funding information

Seaworld and Parks Entertainment;
SeaWorld and Busch Gardens Conservation
Fund

Abstract

Ecological diversity has been reported for killer whales (*Orcinus orca*) throughout the North Atlantic but patterns of prey specialization have remained poorly understood. We quantify interindividual dietary variations in killer whales ($n = 38$) sampled throughout the year in 2017–2018 in northern Norway using stable isotopic nitrogen ($\delta^{15}\text{N}$: $^{15}\text{N}/^{14}\text{N}$) and carbon ($\delta^{13}\text{C}$: $^{13}\text{C}/^{12}\text{C}$) ratios. A Gaussian mixture model assigned sampled individuals to three differentiated clusters, characterized by disparate nonoverlapping isotopic niches, that were consistent with predatory field observations: seal-eaters, herring-eaters, and lumpfish-eaters. Seal-eaters showed higher $\delta^{15}\text{N}$ values (mean $\pm$ SD: $12.6 \pm 0.3\text{‰}$, range = 12.3–13.2‰, $n = 10$) compared to herring-eaters (mean $\pm$ SD: $11.7 \pm 0.2\text{‰}$, range = 11.4–11.9‰, $n = 19$) and lumpfish-eaters (mean $\pm$ SD: $11.6 \pm 0.2\text{‰}$, range = 11.3–11.9, $n = 9$). Elevated $\delta^{15}\text{N}$ values for seal-eaters, regardless of sampling season, confirmed feeding at high trophic levels throughout the year. However, a wide isotopic niche and low measured $\delta^{15}\text{N}$ values in the seal-eaters, compared to that of whales that would eat solely seals ($\delta_{\text{N-measured}} = 12.6$ vs. $\delta_{\text{N-expected}} = 15.5$), indicated a diverse diet that includes both fish and mammal prey. A narrow niche for killer whales sampled at herring and lumpfish seasonal grounds supported seasonal prey specialization reflective of local peaks in prey abundance for the two fish-eating groups. Our results, thus, show differences in prey specialization within this killer whale population in Norway and that the episodic observations of killer whales feeding on prey other than fish are a consistent behavior, as reflected in different isotopic niches between seal and fish-eating individuals.

KEYWORDS

ecological niche, ecological variation, prey specialization, stable isotopes, trophic level

1 | INTRODUCTION

The killer whale (*Orcinus orca*) is an apex predator found in all the world's oceans (Forney & Wade, 2006). Although a generalist

species, studies have revealed discrete prey specializations of sympatric populations in some regions (e.g., Ford et al., 1998; Saulitis, Matkin, Barrett-Lennard, Heise, & Ellis, 2000) with foraging behaviors apparently culturally transmitted through generations within

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd

matrilineal social units (Ford & Ellis, 2014; Riesch, Barrett-Lennard, Ellis, Ford, & Deecke, 2012). Through cultural divergence and social isolation of specialized groups, prey specialization may influence population structure (Hoelzel et al., 2007; Riesch et al., 2012), and eventually facilitate ecotype formation (Foote et al., 2016). Due to different patterns of resource use, killer whale populations may be differentially impacted by human activities. In the coastal waters of the northeastern Pacific, so-called resident killer whales rely on salmonids as a main food source while transient killer whales appear to feed exclusively on marine mammal prey (e.g., Ford & Ellis, 2014; Ford et al., 1998). The resident and transient killer whale communities constitute distinct populations and ecotypes (Morin et al., 2010). Divergent demographic trends for these populations since the mid-1990s have resulted in disparate conservation status and management strategies (COSEWIC, 2008). This is an example of why understanding interindividual diet variations may be important in the management of this species.

In the North Atlantic, at least two types of killer whales differing in morphology, tooth wear and nitrogen isotopic values have been suggested (Foote, Newton, Piertney, Willerslev, & Gilbert, 2009). A far-ranging generalist so-called Type 1 includes herring-feeding killer whales off Norway and Iceland, but with interindividual variation in the dietary proportions of contributing prey items, including high trophic level prey (Foote et al., 2009). This suggestion was supported by a variation in intrapopulation ecological niche in Iceland (Samarra et al., 2018; Samarra, Vighi, Aguilar, & Vikingsson, 2017), and field observations of a subset of individuals switching between pinniped and fish prey from both Norway (Vongraven & Bisther, 2014) and Iceland (Foote, Similä, Vikingsson, & Stevick, 2010). In contrast, larger Type 2 killer whales appear to specialize on cetacean prey (Foote et al., 2009). However, current classification in two types may be oversimplistic considering the large diversity of ecological/dietary patterns across the North Atlantic (see Jourdain, Ugarte, et al., 2019 for a review). Foraging strategies of North Atlantic killer whales at the group and individual levels remain poorly understood.

Killer whales in Norway have historically been thought to specialize on herring (*Clupea harengus*) and to mainly associate with the most abundant Norwegian Spring Spawning (NSS) stock (Similä, Holst, & Christensen, 1996), as supported by concurrent observations. Herring made up almost the entire stomach contents of killer whales caught prior to 1980 (Christensen, 1982), and killer whales typically occur in large seasonal aggregations at herring wintering grounds (Bisther & Vongraven, 1995; Similä et al., 1996), where they display remarkably specialized feeding behaviors (Domenici, Batty, Similä, & Ogam, 2000; Similä & Ugarte, 1993). However, as killer whale studies have primarily been conducted at herring wintering grounds until recently, other prey utilized in other areas were unlikely to be identified. Research efforts extended to other seasons and regions in Norwegian waters have documented new prey species, for example, the Atlantic mackerel (*Scomber scombrus*, Nøttestad et al., 2014), the Atlantic salmon (*Salmo salar*, Vester & Hammerschmidt, 2013), the harbor porpoise (*Phocoena phocoena*, Cosentino, 2015), and the gray (*Halichoerus grypus*) and harbor (*Phoca vitulina*) seals

(Jourdain, Vongraven, Bisther, & Karoliussen, 2017; Vongraven & Bisther, 2014). However, the lack of identification data and/or only brief periods of data collection for these studies precluded any assessment of dietary differences and specializations among killer whale individuals/groups.

Efforts began in 2013 to investigate feeding habits at the individual level, combining predation records of photo-identified individuals, behavioral observations, and tissue samples collected throughout multiple years (Jourdain, Karoliussen, Vos, Zakharov, & Tougaard, 2019; Jourdain et al., 2017, this study). Results revealed that some social groups specialize to some extent on pinnipeds (Jourdain et al., 2017) and that some herring-eating individuals seasonally switch to feeding on locally abundant lumpfish (*Cyclopterus lumpus*) in spring (Jourdain, Karoliussen, et al., 2019). Social interactions among sympatric killer whales adopting distinct foraging behaviors have not been investigated to date. Field observations provide only a snapshot of observable feeding bouts; combining observational records with time-integrated dietary markers would assist in assessing interindividual variations in diet, as well as persistency of prey specialization over longer periods of time.

Because the isotopic composition of a predator's tissue reflects that of its prey resources in a predictable/quantifiable manner (DeNiro & Epstein, 1978), stable isotopes have been commonly used in dietary studies (see Newsome, Clementz, & Koch, 2010 for a review). Due to a greater retention of the heavier ^{15}N isotope than the lighter ^{14}N isotope in the production of nitrogenous waste, the nitrogen ratio of ^{15}N to ^{14}N ($\delta^{15}\text{N}$) shows a stepwise enrichment from food source to consumer and is therefore indicative of relative trophic position (DeNiro & Epstein, 1978; Hobson & Clark, 1992). The carbon ratio of ^{13}C to ^{12}C ($\delta^{13}\text{C}$), primarily reflects variable carbon origins of inshore/benthic versus pelagic/offshore sources, therefore allowing for discrimination between feeding locations in marine systems (Hobson, Piatt, & Pitocchelli, 1994). Typically, inshore/benthic ecosystems are characterized by higher $\delta^{13}\text{C}$ values than pelagic/offshore systems. Isotopic variance within a population, characterized by variance in the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of its individuals, can be used as a measure of diet variation, referred to as niche width (Bearhop, Adams, Waldron, Fuller, & Macleod, 2004). Typically, consumers that feed on a wide range of food sources will display larger variations in isotopic signatures and thus a wider niche than prey specialists that feed on a narrow range of prey. Similarly, consumers that feed in multiple locations will show greater variations in $\delta^{13}\text{C}$, that is, a wider niche (Bearhop et al., 2004).

In the study of killer whales, high variability in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values has proved effective in assessing inter and intrapopulation dietary variations and preferences (Durban, Fearnbach, Burrows, Ylitalo, & Pitman, 2017; Herman et al., 2005; Krahn et al., 2007; Reisinger et al., 2016; Samarra et al., 2017; Tixier et al., 2019). Isotopic profiles confirmed dietary segregation between sympatric resource specialists in the northeastern Pacific (Herman et al., 2005) and around the Antarctic Peninsula (Durban et al., 2017) but also revealed generalist killer whale populations adopting a mixed diet including both fish and mammal prey (Reisinger et al., 2016; Tixier

et al., 2019). In these studies, a priori knowledge of sampled individuals/populations through previous field observations has been highly beneficial for meaningful interpretation of dietary patterns (Newsome et al., 2010).

In this study, we use $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values from killer whale skin samples collected throughout the year in northern Norway. The aims were to (a) measure interindividual variations in dietary habits by comparing isotopic profiles, that is, trophic level and niche width, of fish and seal-eating killer whales; (b) estimate the contribution of pinniped prey to the diet of seal-eating killer whales. Results are discussed in light of predation records available for the sampled whales to assess consistency in individuals' dietary habits and further evaluate the degree of prey specialization for these whales.

2 | MATERIALS AND METHODS

2.1 | Sample collection

Killer whale biopsy samples were collected in August and November 2017 and from April through July 2018 in northern Norway. In November, samples were collected in Kvænangen, Skjervøy (Figure 1a). This fjord was part of the herring wintering grounds (ICES, 2018), and killer whales were observed foraging on herring at time of sampling. The rest of the year, samples were collected off Andøya (Figure 1b), where killer whales were encountered throughout the year and are known to seasonally feed on various prey types

including herring (Jourdain & Vongraven, 2017), pinnipeds (Jourdain et al., 2017), and lumpfish (Jourdain, Karoliussen, et al., 2019).

Killer whale biopsies were sampled using an ARTS darting system (Restech) and 25×9 mm or 40×9 mm stainless steel tips in 2017, and with an injection gun (Pneu-Dart Inc) and 25×7 mm tips in 2018. The biopsy tips were sterilized with boiling water and 95% ethanol and placed in clean plastic bags before use. Killer whales were sampled when travelling or feeding. The region directly posterior to the dorsal fin of adult and subadult killer whales was the target area for sampling. For each sampled individual, identification photographs were taken. Biopsy darts containing skin and blubber were retrieved, stored in a clean plastic bag and placed in a cooling box while at sea. Onshore, skin and blubber layers were sliced apart and stored separately at -20°C until analysis. Isotopic values measured in killer whale skin are expected to represent the individuals' diet in the four to six weeks prior to sampling. This is based on controlled diet experiments that estimated half-time turnover rates for bottlenose dolphin (*Tursiops truncatus*) skin to be 24 ± 8 d for carbon and 48 ± 19 d for nitrogen (Giménez, Ramírez, Almunia, Forero, & Stephanis, 2016).

Estimating the diet of consumers requires isotopic prey values. Because isotopic values may vary greatly both in space and time, it is important that both prey and killer whales are sampled within matching geographic areas and time periods (Phillips et al., 2014). Herring muscle ($n = 4$) was collected in November 2017 in Kvænangen, Skjervøy, lumpfish muscle ($n = 5$) was collected in March–April 2017 in Andfjord while conducting focal studies of feeding killer whales

FIGURE 1 Locations of the 38 killer whale biopsy samples collected in northern Norway in 2017–2018. Region (a) corresponds to Kvænangen, Skjervøy, the herring wintering grounds where killer whales were sampled in November 2017. Region (b) corresponds to Andøya, where whales were sampled at the lumpfish spring spawning grounds and throughout the summer in 2018. Blue dots ($n = 22$) correspond to individual whales sampled during November and assigned to sampling season group *Winter*. Gray dots ($n = 10$) correspond to whales sampled in April–May and assigned to sampling season group *Spring*. Red dots ($n = 6$) correspond to whales sampled in June–August and assigned to sampling season group *Summer*. Herring muscle ($n = 4$) was obtained in region (a). Lumpfish muscle ($n = 5$) and seal muscle ($n = 1$) were obtained from region (b)

(Jourdain, Karoliussen, et al., 2019). Muscle from a dead stranded harbor seal was sampled in November 2017 in Andenes.

2.2 | Data processing

Sampled killer whales were identified using nicks, shape and size of the dorsal fin, and scarring and pigmentation patterns of the saddle patch (Bigg, 1982). Individuals were matched to an existing catalogue of 971 killer whales identified between 2007 and 2018 in northern Norway (Jourdain & Karoliussen, 2018). Classification of sex was done as per Bigg (1982). Records of predation on seals collected in 2013–2018 were used to assign sampled killer whales a priori to one of the two diet groups. Individuals with a history of predation on seals were assigned to the group *Seal-eaters*, while individuals with no such history were classified as *Fish-eaters*. In addition, individuals were assigned a group reflecting season at sampling. Group *Winter* included individuals sampled at herring wintering grounds in November, group *Spring* contained individuals sampled at lumpfish spawning grounds in April–May and group *Summer* included the whales sampled from June through August in Andfjord (Figure 1).

2.3 | Stable isotope analysis

Skin samples from killer whales, and muscle from lumpfish, herring, and seal were freeze-dried and ground individually with an agate mortar and pestle to a fine powder. An aliquot was rinsed three times in a 2:1 chloroform: methanol solution to remove lipids, following the method developed by Folch, Lees, and Stanley (1957) and modified by Elliott, Roth, and Crook (2017). An aliquot from the bulk tissue was not treated with any chloroform: methanol solution. A duplicate analysis was run on non-lipid-extracted and lipid-extracted values, in accordance with recommendations (Lesage et al., 2010; Ryan et al., 2012). $\delta^{13}\text{C}$ values were determined from lipid-extracted samples to control for the low $\delta^{13}\text{C}$ found in the lipid fraction of an organism that can lead to bias (DeNiro & Epstein, 1978; Tarroux et al., 2010; Yurkowski, Hussey, Semeniuk, Ferguson, & Fisk, 2015). $\delta^{15}\text{N}$ values were determined from nonlipid-extracted samples due to the unpredictable changes in $\delta^{15}\text{N}$ values in fish muscle and cetacean skin following lipid extraction (Lesage et al., 2010; Ryan et al., 2012). The powdered sample (1 mg $\pm$ 5%) was weighed into a tin capsule. The $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ ratios were measured simultaneously using an Elemental Analyzer (EA) IsoLink Isotope Ratio Mass Spectrometer (IRMS) System, consisting of a Flash EA and a DeltaV IRMS (Thermo Scientific, Germany). All analyses were conducted at the Stable Isotope Laboratory at the University of Oslo. The quality of the analysis was assured by the incorporation into each run two internal reference materials, JGLUT (L-glutamic acid, $\delta^{13}\text{C}$ = -13.43‰ , $\delta^{15}\text{N}$ = -4.34‰ , Fisher Scientific) and POPPGLY (glycine, $\delta^{13}\text{C}$ = -36.58‰ , $\delta^{15}\text{N}$ = 11.25‰ , Fisher Scientific).

$\delta^{13}\text{C}$ was calibrated to the Vienna Pee Dee Belemnite (VPDB) scale using LSVEC (lithium carbonate, $\delta^{13}\text{C}$ = -46.6‰) and NBS-19 (calcium carbonate, $\delta^{13}\text{C}$ = 1.95‰) (both obtained from the International Atomic Energy Agency, Austria). $\delta^{15}\text{N}$ was calibrated to the AIR scale using USGS40 (L-glutamic acid, $\delta^{15}\text{N}$ = -4.52‰) and USGS41 (L-glutamic acid, $\delta^{15}\text{N}$ = 47.57‰) (both obtained from the United States Geological Survey). Analytical precision based on repeated analyses of quality assurance material JALA (alanine, $\delta^{13}\text{C}$ = -20.62‰ , $\delta^{15}\text{N}$ = -3.16‰ , Fisher Scientific) indicated measurement errors of $0.09 \pm 0.01\text{‰}$ for $\delta^{15}\text{N}$ and $0.06 \pm 0.02\text{‰}$ for $\delta^{13}\text{C}$.

2.4 | Statistical analysis

All statistics were performed in R v.3.4.1. for Mac OS X (R Development Core Team, 2017). The α level was set to $p = .05$. To test simultaneously the effect of independent variables sex, diet group (*Seal-eaters* vs. *Fish-eaters*), and sampling season group (*Winter*, *Spring*, and *Summer*) on isotopic ratios, a multiple linear regression model was fitted to the data measured for $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$. Diagnostic plots were used to investigate normality of the residuals, while Levene's test run in the package *car* (Fox & Weisberg, 2011) was used to validate homogeneity of variance in residuals. Significant explanatory variables were identified by conducting a forward model selection in the package *stats*, from the null model to the full global model featuring all variables. A lower Akaike Information Criterion (AIC) was used for model selection, and a higher adjusted R-squared as an indication of the fit of each model.

A mixture model-based clustering analysis using the *mclust* package was conducted on the $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values to estimate the most likely number of clusters and the probability of individuals belonging to each cluster (Scrucca, Fop, Murphy, & Raftery, 2016). The Bayesian Information Criterion (BIC) was used to select the best model. Individual assignment to clusters was compared to field observations for validation.

The isotopic niche, referring to the isospace delineated by $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values, of resulting clusters was estimated using calculated convex hull areas (encompassing all data points) and bivariate ellipses in the package *SIBER* (Jackson, Inger, Parnell, & Bearhop, 2011). The Standard Ellipse Area (SEA) is a measure of the standard deviation for bivariate data. SEA corrected for small sample size (SEA_c), containing 40% of the data regardless the sample size, enabled robust comparison between clusters. Bayesian Standard Ellipse Areas (SEA_b) were generated using 10^6 posterior draws for each cluster and used to statistically compare niche width between clusters (Jackson et al., 2011).

We used stable isotope mixing models (Parnell et al., 2013) in the package *simmr* (Parnell, 2016) to estimate relative contributions of herring, lumpfish, and seal prey to the diet of seal-eating killer whales at the cluster level. All three prey groups were confirmed to be part of the diet of seal-eating killer whales from field observations (see Table 1 and Discussion). In a Bayesian framework, and accounting

TABLE 1 Summary of the 38 killer whale biopsies sampled in 2017–2018 in northern Norway and used in this study

Whale ID	Sex	Age	Cluster	Diet at sampling	Sampling month	$\delta^{15}\text{N}$	$\delta^{13}\text{C}$
(a priori) Fish-eating individuals							
NKW-063	Male	Adult	Herring-eater	Herring	November	11.7	-19.5
NKW-153	Male	Adult	Herring-eater	Herring	November	11.7	-19.6
NKW-165	Male	Adult	Herring-eater	Herring	November	11.7	-19.4
NKW-180	Male	Adult	Herring-eater	Herring	November	11.8	-19.6
NKW-183	Male	Adult	Herring-eater	Herring	November	11.9	-19.4
NKW-421	Male	Adult	Herring-eater	Herring	November	11.7	-19.0
NKW-506	Male	Adult	Herring-eater	Herring	November	11.7	-19.6
NKW-511	Female	Adult	Herring-eater	Herring	November	11.4	-19.7
NKW-862	Male	Adult	Herring-eater	Herring	November	11.9	-18.9
NKW-867	Male	Adult	Herring-eater	Herring	November	11.5	-19.2
NKW-918	Male	Adult	Herring-eater	Herring	November	11.5	-19.4
NKW-919	Male	Adult	Herring-eater	Herring	November	11.8	-19.8
NKW-920	Male	Adult	Herring-eater	Herring	November	11.6	-19.6
NKW-922	Male	Adult	Herring-eater	Herring	November	11.7	-19.1
NKW-923	Male	Adult	Herring-eater	Herring	November	11.6	-18.9
NKW-924	Male	Adult	Seal-eater	Herring	November	12.6	-18.3
NKW-937	Male	Adult	Herring-eater	Herring	November	11.9	-19.5
UNK 1	N/A	Subadult	Herring-eater	Herring	November	11.4	-19.2
NKW-004	Male	Adult	Lumpfish-eater	Lumpfish	April	11.8	-18.1
NKW-079	Male	Adult	Lumpfish-eater	Lumpfish	April	11.5	-17.7
NKW-348	Male	Adult	Herring-eater	Lumpfish	April	11.5	-19.2
NKW-537	Female	Adult	Lumpfish-eater	Lumpfish	April	11.5	-18.4
NKW-572	Male	Adult	Lumpfish-eater	Lumpfish	April	11.5	-18.3
NKW-714	Male	Subadult	Lumpfish-eater	Lumpfish	April	11.9	-18.3
NKW-908	N/A	N/A	Lumpfish-eater	Lumpfish	April	11.8	-18.2
NKW-978	N/A	N/A	Lumpfish-eater	Lumpfish	April	11.3	-18.3
Y137	N/A	Subadult	Lumpfish-eater	Lumpfish	April	11.5	-18.1
NKW-598	Male	Adult	Lumpfish-eater	N/A	July	11.3	-18.6
(a priori) Seal-eating individuals							
NKW-785	Female	Adult	Herring-eater	Herring	November	11.5	-19.2

(Continues)

TABLE 1 (Continued)

Whale ID	Sex	Age	Cluster	Diet at sampling	Sampling month	$\delta^{15}\text{N}$	$\delta^{13}\text{C}$
K1	Female	Adult	Seal-eater	Lumpfish	May	12.6	-18.5
K101	Male	Adult	Seal-eater	N/A	November	12.4	-19.4
K103	Female	Adult	Seal-eater	N/A	November	12.3	-20.0
K105	Male	Adult	Seal-eater	N/A	November	12.8	-19.2
K106	Female	Adult	Seal-eater	Pinnipeds	August	13.2	-18.9
K107	Male	Adult	Seal-eater	Pinnipeds	June	12.3	-18.0
NKW-096	Female	Adult	Seal-eater	Pinnipeds	August	12.5	-19.1
NKW-823	N/A	N/A	Seal-eater	Pinnipeds	August	12.9	-19.4
NKW-921	Male	Adult	Seal-eater	Pinnipeds	August	12.5	-18.5

Note: Individuals were listed a priori as fish or seal-eaters from field observations. Unique identification (ID) codes are given (Jourdain & Karoliussen, 2018), as well as sex and age for each individual when known. Clustering of individuals (Cluster 1: seal-eaters, Cluster 2: herring-eaters, or Cluster 3: lumpfish-eaters), as determined from the maximum likelihood Gaussian mixture model, is indicated. Individuals sampled in November belong to sampling group Winter, individuals sampled in April–May belong to sampling group Spring and individuals sampled from June through August belong to sampling group Summer. Isotopic values refer to lipid-extracted $\delta^{13}\text{C}$ and nonlipid-extracted $\delta^{15}\text{N}$.

for the putative diet components of a consumer and uncertainties in both food source and consumer isotopic values, models estimate the distributions of possible diets. Killer whale $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ skin values were input as the consumer data and mean ($\pm$ SD) isotopic values of herring, lumpfish, and seal muscle were used as sources (see Results). To set uncertainty around the single data point available for seal prey, we used published uncertainty data ($SD_{\delta^{13}\text{C}} = 0.2$, $SD_{\delta^{15}\text{N}} = 0.3$) on isotopic muscle values from stranded harbor seals ($n = 9$) that were collected over a short period of time and from a single region (Hobson, Sease, Merrick, & Piatt, 1997). Concentration dependencies and corrections for trophic discrimination, only optional when running mixing models with *simmr*, were not computed. Due to the large variation in both $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values (see Results) among seal-eating killer whales, mixing models were also run at the individual level using the same segment values.

Expected $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values were also calculated for herring, lumpfish, and seal-eaters if they were prey specialists using the diet-to-tissue skin discrimination factors $1.57 \pm 0.52\text{‰}$ and $1.01 \pm 0.37\text{‰}$ for nitrogen and carbon, respectively, as estimated for bottlenose dolphins by Giménez et al. (2016). The following equations were used, as per Herman et al. (2005):

$$\delta_{\text{N-expected}} = \sum_{i=1}^n (\text{Diet } i * \delta^{15}\text{N } i) + 1.57$$

$$\delta_{\text{C-expected}} = \sum_{i=1}^n (\text{Diet } i * \delta^{13}\text{C } i) + 1.01$$

where n is the number of different prey species consumed, Diet i is the proportion of each prey species consumed, and $\delta^{15}\text{N } i$ and $\delta^{13}\text{C } i$ are the measured isotopic ratio values of the i th prey species being herring, lumpfish, or seal.

3 | RESULTS

Thirty-eight individual killer whales were biopsy sampled from 16 encounter days in 2017–2018 in northern Norway (Figure 1, Table 1). Due to dynamic social associations observed throughout the years and challenges to accurately identify stable units, individuals were considered independently except for one case discussed below (see Discussion). For the period 2013–2018, the sampled individuals were encountered throughout the year on one to 23 observation days (mean = 7.8; $SD = 5.3$). Resulting predatory observations confirmed that some of the individuals sampled at matching times and locations adopted different diets (see Jourdain, Karoliussen, et al., 2019; Jourdain et al., 2017; Table 1).

3.1 | Isotopic values and clustering

Killer whale skin $\delta^{15}\text{N}$ values ranged between 11.3‰ and 13.2‰ (mean $\pm$ SD : $11.9 \pm 0.5\text{‰}$, $n = 38$) and $\delta^{13}\text{C}$ values ranged between -20.0‰ and -17.7‰ ($-19.0 \pm 0.6\text{‰}$, $n = 38$), resulting in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ spanning a range of 1.9‰ and 2.3‰, respectively

(Table 1). Herring muscle $\delta^{15}\text{N}$ values ranged between 9.6‰ and 10.4‰ ($10.0 \pm 0.3\text{‰}$, $n = 4$) and $\delta^{13}\text{C}$ values ranged between -21.6‰ and -20.6‰ ($-21.0 \pm 0.5\text{‰}$, $n = 4$). Lumpfish muscle $\delta^{15}\text{N}$ values ranged between 11.0‰ and 12.3‰ ($11.5 \pm 0.5\text{‰}$, $n = 5$) and $\delta^{13}\text{C}$ values ranged between -20.6‰ and -20.3‰ ($-20.4 \pm 0.1\text{‰}$, $n = 5$). The one seal muscle sample had a $\delta^{15}\text{N}$ value of 13.9‰ and a $\delta^{13}\text{C}$ value of -19.9‰.

Multiple linear regression fitted to $\delta^{15}\text{N}$ values, and testing sex, sampling season, and a priori diet groups showed diet group as the only significant explanatory variable (adjusted $R^2 = 0.57$, $F_{5,32} = 10.96$, $p < .001$). Whales previously recorded preying on seals had higher $\delta^{15}\text{N}$ values than whales never recorded feeding on seal prey (*Seal-eaters* vs. *Fish-eaters*: coefficient estimate = 0.86, $t = 5.20$, and $p < .001$; Figure 2a). The multiple linear

regression fitted to $\delta^{13}\text{C}$ showed sampling season as the only significant explanatory variable (adjusted $R^2 = 0.57$, $F_{5,32} = 11$, $p < .001$; Figure 2b). Whales sampled at lumpfish spawning grounds in April and May had the highest $\delta^{13}\text{C}$ values (coefficient estimate = -18.31, $t = -147$, $p < .001$), followed by the whales sampled in the summer (*Spring* vs. *Summer*: coefficient estimate = -0.46, $t = -2.36$, $p = .30$). The whales sampled at herring wintering grounds in November showed the lowest $\delta^{13}\text{C}$ values (*Spring* vs. *Winter*: coefficient estimate = -1.02, $t = -6.86$, $p < .001$).

Based solely on the $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values in killer whale skin, the maximum likelihood Gaussian mixture model showed the most likely number of clusters to be three. *Cluster 1* was distinguished by higher $\delta^{15}\text{N}$ values (mean $\pm$ SD: $12.6 \pm 0.3\text{‰}$, range = 12.3–13.2‰, $n = 10$) than both *Cluster 2* (mean $\pm$ SD: $11.7 \pm 0.2\text{‰}$, range = 11.4–11.9‰,

FIGURE 2 (a) Boxplot of the $\delta^{15}\text{N}$ values (in ‰) measured in skin samples of killer whales known as *Fish-eaters* (i.e., no history of predation on seals, $n = 28$) and *Seal-eaters* (i.e., history of predation on seal prey, $n = 10$) from field observations; (b) Boxplot of the $\delta^{13}\text{C}$ values (in ‰) for each of the three sampling groups *Winter*, *Spring*, and *Summer*. For both plots, box represents second and third quartiles, horizontal line corresponds to the median, and whiskers represent the first and fourth quartiles. Data points are represented as dots, and dots outside the box and whiskers are outliers

FIGURE 3 Isospace of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values and niches as measured from killer whale ($n = 38$) skin samples. Seal-eaters ($n = 10$) are represented as black unfilled circles, herring-eaters ($n = 19$) as red, and lumpfish-eaters ($n = 9$) as dark blue. Solid lines represent the standard ellipses corrected for sample size (SEA_c) and encompassing 40% of the data, while dashed lines represent the convex hull area including the entire dataset for each cluster. Note the absence of overlap between clusters. Filled circles indicate $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values from muscle samples of prey ($n = 4$ herring; $n = 5$ lumpfish; $n = 1$ harbor seal)

FIGURE 4 Outputs from Bayesian isotopic mixing models showing proportional estimated contributions (mean, 25% and 75% percentiles) of herring, lumpfish, and harbor seal to the diet of seal-eating killer whales (*Cluster 1*, $n = 10$) biopsy sampled in northern Norway

$n = 19$) and *Cluster 3* (mean $\pm$ SD: $11.6 \pm 0.2\text{‰}$, range = $11.3\text{--}11.9$, $n = 9$). *Cluster 1* showed a wide variation in $\delta^{13}\text{C}$ with values intermediate to *Cluster 2* and *Cluster 3* (Table 1, Figure 3). Except for the three individuals NKW-348, NKW-785, and NKW-924 assigned to unexpected clusters (see Table 1), assignment of individuals to clusters coincided largely with predatory field observations (Jourdain, Karoliussen, et al., 2019; Jourdain & Vongraven, 2017; Jourdain et al., 2017; Table 1; Figure 3). *Clusters 1*: seal-eaters, *Cluster 2*: herring-eaters, and *Cluster 3*: lumpfish-eaters were therefore used for the isotopic analyses hereafter.

Average $\delta^{15}\text{N}$ values measured in individuals from *Cluster 1*: seal-eaters were approximately 1‰ higher than in fish-eating individuals from *Cluster 2*: herring-eaters and *Cluster 3*: lumpfish-eaters combined (mean $\pm$ SD: $11.6 \pm 0.2\text{‰}$, range = $11.3\text{--}11.9$, $n = 28$).

3.2 | Isotopic niche and diet composition

Niche width based on isotopic signatures for herring and lumpfish-eating killer whales appeared narrow and not significantly different as supported by a similar SEA_B ($p = .14$; Figure 3). SEA_B for seal-eaters indicated a wider niche than both clusters ($p < .001$) as supported by a wide convex hull (Figure 3). Total absence of SEA_C overlap among groups suggested that the three dietary clusters occupy distinct niches (Figure 3).

Bayesian stable isotope mixing models estimated mean relative contributions of the three prey groups to the diet of seal-eating killer whales to be: herring (mean $\pm$ SD) = 0.16 ± 0.08 , lumpfish = 0.26 ± 0.12 , and harbor seal = 0.58 ± 0.06 (Figure 4). Mixing models run separately for each seal-eating individual further suggested a large variation in proportional contribution of harbor seal compared to herring and lumpfish. For seven of the 10 seal-eating killer whales, harbor seal was the highest dietary contributor although fish prey also appeared as a significant food source (Figure 5). Mean contribution of harbor seal ranged from 17% (2.5%–97.5% quantiles: 2%–36%,

FIGURE 5 Individual variation in the mean proportional contribution of harbor seal, herring, and lumpfish to the diet of seal-eating killer whales (*Cluster 1*, $n = 10$) biopsy sampled in northern Norway, as estimated from Bayesian isotopic mixing models

TABLE 2 Calculated expected $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ skin values for killer whales feeding exclusively on herring, lumpfish, and seal prey as compared to mean true (measured) values for each dietary cluster

	Expected $\delta^{15}\text{N}$; $\delta^{13}\text{C}$ (in ‰)	True values $\delta^{15}\text{N}$; $\delta^{13}\text{C}$ (in ‰)
<i>Cluster 1</i> : seal-eaters	15.5; -18.9	12.6; -18.9
<i>Cluster 2</i> : herring-eaters	11.6; -20.0	11.7; -19.3
<i>Cluster 3</i> : lumpfish-eaters	13.0; -19.4	11.6; -18.2

whale KI07) to 93% (85%–98%, whale KI06, Figure 5). Expected values for potential prey specialists feeding on herring, lumpfish, and seals are shown in Table 2.

4 | DISCUSSION

Recent killer whale studies in Norway suggested multiple prey resources (Cosentino, 2015; Jourdain, Karoliussen, et al., 2019; Jourdain et al., 2017; Nøttestad et al., 2014; Vester & Hammerschmidt, 2013; Vongraven & Bisther, 2014) as opposed to initial observations of killer whales being herring specialists in this region (Christensen, 1982; Similä et al., 1996). Our results further support a generalist population characterized by interindividual dietary variations. Low variation in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ skin values for killer whales sampled at herring and lumpfish grounds supported previous field observations of seasonal specialization on abundant fish prey (Jourdain, Karoliussen, et al., 2019; Similä et al., 1996; Similä & Ugarte, 1993). Higher $\delta^{15}\text{N}$ skin values measured throughout the year for individuals repeatedly observed feeding on seals in 1988–2016 (Jourdain et al., 2017; Vongraven & Bisther, 2014) supported persistent feeding at higher trophic level and marked preference for pinniped prey for these whales, though a wide isotopic niche indicated a diversified diet.

Herring and lumpfish-eating killer whales had similar $\delta^{15}\text{N}$ values, indicative of feeding at comparable trophic levels. Low variation in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values in herring-eaters (*Cluster 2*) resulted in the narrowest niche of the three clusters, implying consistency in diet among individuals and apparent prey specialization in the winter months of high herring abundance. This is consistent with Foote, Vester, Vikingsson, and Newton (2012) who found only little variation in isotopic values (mean $\pm$ SD: 11.8 ± 0.5 , range = 10.9–12.9, $n = 20$) in killer whales sampled at herring wintering grounds in northern Norway in 2005 and 2007. Lumpfish-eating killer whales (*Cluster 3*), sampled in the first half of April in Andfjord, were shown to seasonally feed on lumpfish, based on predatory observations and molecular identification of target prey (Jourdain, Karoliussen, et al., 2019). These whales showed a slightly wider isotopic niche, possibly indicative of a gradual seasonal switch in diet, from herring to lumpfish. As the wintering herring initiates its progressive migration southwards to spawning grounds at the end of January in the study area (Røttingen, 1990), the lumpfish migration from offshore feeding areas to inland spawning grounds starts in February (Eriksen, Durif, & Prozorkevich, 2014). Such overlap in time and space between the two fish species could promote temporary inclusion of both fish prey in killer whales' diet. Indeed, sampled lumpfish-eating killer whales were confirmed to be feeding on herring from winter encounters between 2015 and 2018 (Jourdain, Karoliussen, et al., 2019). In any case, seasonal specialization on lumpfish is likely reflecting a seasonal local peak in prey abundance rather than true dietary preference.

A few individuals sampled at herring grounds (i.e., whales KI01, KI03, KI05, and NKW-924) and lumpfish grounds (i.e., whale K1) showed higher $\delta^{15}\text{N}$ skin values, indicative of feeding on higher trophic level prey. These whales were assigned to *Cluster 1*: seal-eaters

for which distinctive enrichment in ^{15}N correlated with apparent inclusion of seals in their diet. This was confirmed from field observations for nine of the 10 whales in this cluster (see Jourdain et al., 2017). Seal-eaters showed $\delta^{15}\text{N}$ skin values in average 1.0‰, and up to 1.9‰, higher compared to fish-eaters, regardless of sampling season. These values are consistent with approximately one trophic level, assuming an estimated discrimination factor of $\sim 1.57\text{‰}$ for nitrogen for bottlenose dolphin (Giménez et al., 2016). This falls within or exceeds the range of values discriminative of fish versus marine mammal-eating killer whales in Iceland ($\Delta\text{N} = 1.32\text{‰}$ in Samarra et al., 2017), British Columbia/Washington State ($\Delta\text{N} = 0.7\text{‰}$, in Herman et al., 2005), Eastern Aleutian Islands ($\Delta\text{N} = 1.2\text{‰}$ in Herman et al., 2005), and Antarctica ($\Delta\text{N} = 0.9\text{‰}$ in Durban et al., 2017). Disparate isotopic profiles among individuals sampled at matching times and locations in our study indicate co-occurrence of individuals that include, and others that do not include, pinnipeds to their diet. Elevated $\delta^{15}\text{N}$ values found in three of the 20 killer whales sampled at herring wintering grounds by Foote et al. (2012) also suggested such dietary structuring.

Combined with individual predation records collected over years and up to several decades (see Jourdain et al., 2017; Vongraven & Bisther, 2014), our results confirm persistent dietary preference rather than opportunistic feeding on pinnipeds for these whales. Consistently elevated $\delta^{15}\text{N}$ values for all ten seal-eaters (*Cluster 1*) sampled in all seasons further indicated that predation on seals (or other high trophic level prey) occurred throughout the year, even at times of high abundance of fish prey. This was supported by strikingly similar skin $\delta^{15}\text{N}$ values measured for individuals KI01, KI03, and KI07. All three individuals were shown to constitute a stable long-lasting social group that hunts and feeds cooperatively (Jourdain et al., 2017), and therefore, comparable diet and isotopic profiles are expected for these whales. Although sampled in different seasons for this study (see Table 1), year-round homogeneous $\delta^{15}\text{N}$ values support consistent feeding habits for this group regardless of the time of the year.

A wider isotopic niche found for seal-eating killer whales compared to fish-eaters implies a more diversified diet. This was supported by the following observations of seal-eating individuals also feeding on fish prey. Adult males NKW-924 and KI01 were photographed while scavenging around herring-purse seiners in November 2017 and 2018 (EJ, unpublished data), respectively, confirming effective inclusion of herring to their diets. Whales KI03, KI05, and KI06 were also observed carousel feeding on herring off Ona, Møre coast, Norway in 1991, immediately prior to travelling near-shore to chase harbor seals (Vongraven & Bisther, 2014). Female K1 was observed feeding on lumpfish at time of sampling in May 2018 before resuming nearshore foraging as previously described for seal-eating groups (Jourdain et al., 2017). Expected $\delta^{15}\text{N}$ values calculated for seal-eaters if they were prey specialists were in further support of a mixed diet (Table 2).

When computing herring, lumpfish, and harbor seal as putative prey of seal-eating killer whales (*Cluster 1*), isotope mixing models suggested that seal prey made up 46 to 68% of these whales'

diet (2.5 and 97.5% quartiles at the cluster level), therefore suggesting fish prey as an equal or secondary food source (Figure 4). However, these results should only be considered preliminary and interpreted with caution due to small sample size for both killer whale and prey values, and due to the limitations and assumptions involved in the use of these models. Mixing models are very sensitive to the assumption that all potential prey sources are computed (Bond & Diamond, 2011; Parnell et al., 2013; Phillips et al., 2014), which can be confirmed if isotopic values of the consumer fall within the mixing polygon of the connecting food sources once corrected for trophic fractionation (Phillips & Koch, 2002). This was not the case here (see Figure 3), implying missing prey sources. Also, if there was large temporal and/or individual variation in diet composition; that is, not all seal-eating individuals in *Cluster 1* feeding on all three prey types, and not at time of sampling, computing herring, lumpfish, and seal prey could be unsuitable (Phillips et al., 2014; Phillips & Koch, 2002). Another caveat in our models resides in the use of variation values from harbor seals that were not collected in our study area. Further efforts in monitoring food sources to better obtain a priori knowledge on individuals' diet, and collecting prey samples, are warranted. More complete datasets and inclusion of appropriate trophic discrimination factors would allow for more accurate diet reconstruction (Phillips et al., 2014) of seal-eating killer whales in Norway and should be a priority for future research.

Nevertheless, the large variation observed in estimated dietary contributions among seal-eating killer whales is consistent with the wide isotopic niche that characterizes this cluster (Figures 3–5). Importantly, there might be a gradient of dietary patterns among Norwegian killer whales ranging from fish specialists (*Clusters 2 and 3*: herring and lumpfish-eaters) on one end to individuals that feed most extensively on high trophic level prey (*Cluster 1*: seal-eaters) on the other end. Contrasting field observations and isotopic values for whale NKW-785 support this theory. This female was observed feeding on pinnipeds on one encounter day in June 2017 (EJ unpublished data) and feeding on herring at time of sampling in November of the same year. Owing to low $\delta^{15}\text{N}$ skin values, this whale was assigned to the cluster herring-eaters by the Gaussian mixture model, bringing evidence that not all killer whales known as seal-eaters from field observations forage on pinnipeds throughout the year. Instead, some individuals may seasonally switch between specialized feeding behaviors or may only opportunistically feed on seals.

Variations in $\delta^{13}\text{C}$ among the three sampling groups support variable foraging areas in relation to seasonal prey movement and/or foraging on prey that utilize different habitats. The lowest $\delta^{13}\text{C}$ values measured for herring-eaters (*Cluster 2*) sampled in November at herring wintering grounds coincide with killer whales just returning from their summer offshore distribution (Nøttestad et al., 2014, 2015), following the migration of the herring to the coastal wintering grounds in early fall. Lower $\delta^{13}\text{C}$ values could also be a result of specializing on herring, which is a pelagic fish spending most of its life offshore (Dragesund, Hamre, & Ulltang,

1980). The highest $\delta^{13}\text{C}$ values measured in killer whales sampled at lumpfish spawning grounds in spring (*Cluster 3*) were indicative of a coastal habitat for these whales. This is consistent with a winter spent in fjords foraging on wintering herring but could also be a result of temporarily specializing on the lumpfish which is a semipelagic fish (Davenport, 1985). Intermediate $\delta^{13}\text{C}$ values for whales sampled from June through August could indicate intergroup variations in foraging areas due to a mosaic of prey resources (Jourdain et al., 2017; Nøttestad et al., 2014; Similä et al., 1996) relied upon at this time of the year.

Expected values of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ in skin of killer whales that would exclusively feed on seal prey indicate that pinniped-eating individuals sampled in this study are not prey specialists. Vongraven and Bisther (2014) suggested that the near total collapse of the NSS herring in 1970 caused by overfishing, could have forced a herring-dependent population of killer whales to switch to other prey types including pinnipeds. Phenotypic plasticity and the ability to learn and culturally transmit new hunting techniques may have facilitated such a switch (Riesch et al. 2012; Samarra & Miller, 2015). Further resource specialization may increase foraging efficiency if experienced foragers benefit from enhanced searching and handling abilities of better selected cost-effective prey (see Bolnick et al., 2003). Under consistent environmental conditions, and if prey specialists indeed experienced a greater fitness than generalists, the level of prey specialization could increase over time regardless of target prey, as shown in other species (Annett & Pierotti, 1999; Golet, Kuletz, Roby, & Irons, 2000).

Despite a small sample size, our study captured a diversity of dietary patterns largely consistent with field observations. Results highlight dietary structuring and differences in prey specialization within this killer whale population which could reflect either seasonal localized food abundance, individuals' true dietary preferences, or both. Resampling of individuals over time and throughout the year would assist in assessing intraindividual dietary variations. While links between diet, genetics and social structure remain to be investigated in this region, our observations confirm that at least seasonal range overlap occurs among killer whale groups adopting distinct diets.

ACKNOWLEDGMENTS

This research was supported by grants from the Sea World and Busch Gardens Conservation fund and Sea World and Parks Entertainment. Biopsy sampling was conducted under the permit FOTS-ID 10176 issued by Mattilsynet (report nr. 2016/179856) to Norwegian Orca Survey.

CONFLICT OF INTERESTS

Authors have no competing interests to declare.

AUTHOR CONTRIBUTIONS

E.J., C.A., R.K., and D.V. conceptualized the study. E.J., C.A., and R.K. collected samples. E.J., C.A., A.R., and K.B. conceptualized the analyses. C.A. conducted laboratory work. E.J. and C.A. analyzed

data. E.J., C.A., and D.V. wrote the paper. All authors approved the final manuscript.

DATA AVAILABILITY STATEMENT

All isotopic values used in this study are available from Figshare Digital Repository: <https://doi.org/10.6084/m9.figshare.11901933.v1>

ORCID

Eve Jourdain <https://orcid.org/0000-0003-2619-7799>

Clare Andvik <https://orcid.org/0000-0002-1870-7106>

Anders Ruus <https://orcid.org/0000-0002-4374-7871>

Dag Vongraven <https://orcid.org/0000-0002-0505-7757>

Katrine Borgå <https://orcid.org/0000-0002-8103-3263>

REFERENCES

- Annett, C. A., & Pierotti, R. J. E. (1999). Long-term reproductive output in Western Gulls: Consequences of alternate tactics in diet choice. *Ecology*, 80, 288–297. [https://doi.org/10.1890/0012-9658\(1999\)080\[0288:LTROIW\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[0288:LTROIW]2.0.CO;2)
- Bearhop, S., Adams, C. E., Waldron, S., Fuller, R. A., & Macleod, H. (2004). Determining trophic niche width: A novel approach using stable isotope analysis. *Journal of Animal Ecology*, 73, 1007–1012. <https://doi.org/10.1111/j.0021-8790.2004.00861.x>
- Bigg, M. (1982). An assessment of killer whale *Orcinus orca* stocks off Vancouver Island, British Columbia. *Report of the International Whaling Commission*, 32, 655–666.
- Bisther, A., & Vongraven, D. (1995). Studies of social ecology of Norwegian killer whales (*Orcinus orca*). In: A. S. Blix, L. Walløe, & Ø. Ulltang (Eds.) *Whales, seals, fish and man*. Amsterdam: Elsevier Science.
- Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Darrin Hulsey, C., & Forister, M. L. (2003). The Ecology of Individuals: Incidence and Implications of Individual Specialization. *The American Naturalist*, 161, 1–28.
- Bond, A. L., & Diamond, A. W. (2011). Recent Bayesian stable-isotope mixing models are highly sensitive to variation in discrimination factors. *Journal of Ecological Applications*, 21, 1017–1023. <https://doi.org/10.1890/09-2409.1>
- Christensen, I. (1982). Killer whales in Norwegian coastal waters. *Report of the International Whaling Commission*, 32, 633–641.
- Cosentino, A. M. (2015). First record of Norwegian killer whales attacking and feeding on a harbour porpoise. *Marine Biodiversity Records*, 8, e108.
- COSEWIC. (2008). *COSEWIC assessment and update status report on the killer whale Orcinus orca, Southern Resident population, Northern Resident population, West Coast Transient population, Offshore population and Northwest Atlantic / Eastern Arctic population, in Canada*. Ottawa: Committee on the Status of Endangered Wildlife in Canada.
- Davenport, J. (1985). *Synopsis of biological data on the lumpsucker, Cyclopterus lumpus (Linnaeus, 1758)*. Rome, Italy: Food and Agriculture Organization of the United Nations., FAO Fisheries.
- Deniro, M. J., & Epstein, S. (1978). Influence of diet on the distribution of carbon isotopes in animals. *Geochimica et Cosmochimica Acta*, 42, 495–506.
- Domenici, P., Batty, R. S., Similä, T., & Ogam, E. (2000). Killer whales (*Orcinus orca*) feeding on schooling herring (*Clupea harengus*) using underwater tail-slaps: Kinematic analyses of field observations. *Journal of Experimental Biology*, 203, 283–294.
- Dragesund, O., Hamre, J., & Ulltang, Ø. (1980). Biology and population dynamics of the Norwegian spring-spawning herring. *Rapports Et Procès-Verbaux Des Réunions, Conseil International pour l'Exploration de la Mer*, 177, 43–71.
- Durban, J. W., Fearnbach, H., Burrows, D. G., Ylitalo, G. M., & Pitman, R. L. (2017). Morphological and ecological evidence for two sympatric forms of type B killer whale around the Antarctic Peninsula. *Polar Biology*, 40, 231–236.
- Elliott, K. H., Roth, J. D., & Crook, K. (2017). Lipid extraction techniques for stable isotope analysis and ecological assays. In: S. K. Bhattacharya (Ed.) *Lipidomics: Methods and Protocols*. New York, NY: Humana Press.
- Eriksen, E., Durif, C. M. F., & Prozorkevich, D. (2014). Lumpfish (*Cyclopterus lumpus*) in the Barents Sea: Development of biomass and abundance indices, and spatial distribution. *ICES Journal of Marine Science*, 71, 2398–2402.
- Folch, J., Lees, M., & Stanley, S. (1957). A simple method for the isolation and purification of total lipides from animal tissues. *The Journal of Biological Chemistry*, 226, 497–509.
- Foote, A. D., Newton, J., Piortney, S. B., Willerslev, E., & Gilbert, M. T. P. (2009). Ecological, morphological and genetic divergence of sympatric North Atlantic killer whale populations. *Molecular Ecology*, 18, 5207–5217. <https://doi.org/10.1111/j.1365-294x.2009.04407.x>
- Foote, A. D., Similä, T., Vikingsson, G., & Stevick, P. T. (2010). Movement, site fidelity and connectivity in a top marine predator, the killer whale. *Evolutionary Ecology*, 24, 803–814. <https://doi.org/10.1007/s10682-009-9337-x>
- Foote, A. D., Vester, H., Vikingsson, G. A., & Newton, J. (2012). Dietary variation within and between populations of Northeast Atlantic killer whales, *Orcinus orca*, inferred from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analyses. *Marine Mammal Science*, 28, 472–485.
- Foote, A. D., Vijay, N., Avila-Arcos, M. C., Baird, R. W., Durban, J. W., Fumagalli, M., ... Wolf, J. B. W. (2016). Genome-culture coevolution promotes rapid divergence of killer whale ecotypes. *Nature Communications*, 7.
- Ford, J. K. B., & Ellis, G. M. (2014). You are what you eat: Foraging specializations and their influence on the social organization and behavior of killer whales. In: J. Yamagiwa, & L. Karczmarski (Eds.), *Primates and cetaceans: Field research and conservation of complex mammalian societies*. Tokyo, Japan: Springer.
- Ford, J. K. B., Ellis, G. M., Barrett-Lennard, L. G., Morton, A. B., Palm, R. S., & Balcomb, K. C. (1998). Dietary specialization in two sympatric populations of killer whales (*Orcinus orca*) in Coastal British Columbia and adjacent waters. *Canadian Journal of Zoology*, 76, 1456–1471. <https://doi.org/10.1139/cjz-76-8-1456>
- Forney, K. A., & Wade, P. R. (2006). Worldwide distribution and abundance of killer whales. In: J. A. Estes, D. P. Demaster, D. F. Doak, T. M. Williams, R. L. Brownell Jr. (Eds.) *Whales, whaling and ocean ecosystems*. Berkeley, CA: University of California Press.
- Fox, J., & Weisberg, S. (2011). *An {R} companion to applied regression* (2nd ed.). Retrieved from <http://socserv.socsci.mcmaster.ca/jfox/Books/Companion>
- Giménez, J., Ramírez, F., Almunia, J., Forero, M. G., & de Stephanis, R. (2016). From the pool to the sea: Applicable isotope turnover rates and diet to skin discrimination factors for bottlenose dolphins (*Tursiops truncatus*). *Journal of Experimental Marine Biology and Ecology*, 475, 54–61. <https://doi.org/10.1016/j.jembe.2015.11.001>
- Golet, G. H., Kuletz, K. J., Roby, D. D., & Irons, D. B. J. T. A. (2000). Adult prey choice affects chick growth and reproductive success in pigeon guillemots. *The Auk*, 117, 82–91. <https://doi.org/10.1093/auk/117.1.82>
- Herman, D. P., Burrows, D. G., Wade, P. R., Durban, J. W., Matkin, C. O., Leduc, R. G., ... Krahn, M. M. (2005). Feeding ecology of Eastern North Pacific killer whales *Orcinus orca* from fatty acid, stable isotope, and organochlorine analyses of blubber biopsies. *Marine Ecology Progress Series*, 302, 275–291. <https://doi.org/10.3354/meps302275>

- Hobson, K. A., & Clark, R. G. (1992). Assessing avian diets using stable isotopes II: Factors influencing diet-tissue fractionation. *The Condor: Ornithological Applications*, 94, 189–197. <https://doi.org/10.2307/1368808>
- Hobson, K. A., Piatt, J. F., & Pitocchelli, J. (1994). Using stable isotopes to determine seabird trophic relationships. *Journal of Animal Ecology*, 63, 786–798. <https://doi.org/10.2307/5256>
- Hobson, K. A., Sease, J. L., Merrick, R. L., & Piatt, J. F. (1997). Investigating trophic relationships of pinnipeds in Alaska and Washington using stable isotope ratios of nitrogen and carbon. *Marine Mammal Science*, 13, 114–132. <https://doi.org/10.1111/j.1748-7692.1997.tb00615.x>
- Hoelzel, A. R., Hey, J., Dahlheim, M. E., Nicholson, C., Burkanov, V., & Black, N. (2007). Evolution of population structure in a highly social top predator, the killer whale. *Molecular Biology and Evolution*, 24, 1407–1415. <https://doi.org/10.1093/molbev/msm063>
- ICES. (2018). *Report of the Working Group on Widely Distributed Stocks (WGWIDE)*, 28 August– 3 September 2018, Faroe Islands, Torshavn.
- Jackson, A. L., Inger, R., Parnell, A. C., & Bearhop, S. (2011). Comparing isotopic niche widths among and within communities: SIBER – Stable Isotope Bayesian Ellipses in R. *Journal of Animal Ecology*, 80, 595–602.
- Jourdain, E., & Karoliussen, R. (2018). Identification catalogue of Norwegian killer whales: 2007–2018. Figshare. <https://doi.org/10.6084/m9.figshare.4205226>
- Jourdain, E., Karoliussen, R., de Vos, J., Zakharov, S. E., & Tougaard, C. (2019). Killer whales (*Orcinus orca*) feeding on lumpfish (*Cyclopterus lumpus*) in northern Norway. *Marine Mammal Science*, 1–14.
- Jourdain, E., Ugarte, F., Vikingsson, G. A., Samarra, F. I. P., Ferguson, S. H., Lawson, J., ... Desportes, G. (2019). North Atlantic killer whale *Orcinus orca* populations: A review of current knowledge and threats to conservation. *Mammal Review*, 49, 384–400. <https://doi.org/10.1111/mam.12168>
- Jourdain, E., & Vongraven, D. (2017). Humpback whale (*Megaptera novaeangliae*) and killer whale (*Orcinus orca*) feeding aggregations for foraging on herring (*Clupea harengus*) in Northern Norway. *Mammalian Biology - Zeitschrift für Säugetierkunde*, 86, 27–32.
- Jourdain, E., Vongraven, D., Bisther, A., & Karoliussen, R. (2017). First longitudinal study of seal-feeding killer whales (*Orcinus orca*) in Norwegian coastal waters. *PLoS ONE*, 12, e0180099.
- Krahn, M. M., Herman, D. P., Matkin, C. O., Durban, J. W., Barrett-Lennard, L., Burrows, D. G., ... Wade, P. R. (2007). Use of chemical tracers in assessing the diet and foraging regions of Eastern North Pacific killer whales. *Marine Environmental Research*, 63, 91–114.
- Lesage, V., Morin, Y., Rioux, É., Pomerleau, C., Ferguson, S. H., & Pelletier, É. (2010). Stable isotopes and trace elements as indicators of diet and habitat use in cetaceans: Predicting errors related to preservation, lipid extraction, and lipid normalization. *Marine Ecology Progress Series*, 419, 249–265.
- Morin, P. A., Archer, F. I., Foote, A. D., Vilstrup, J., Allen, E. E., Wade, P., ... Harkins, T. (2010). Complete mitochondrial genome phylogeographic analysis of killer whales (*Orcinus orca*) indicates multiple species. *Genome Research*, 20, 908–916.
- Newsome, S. D., Clementz, M. T., & Koch, P. L. (2010). Using stable isotope biogeochemistry to study marine mammal ecology. *Marine Mammal Science*, 26, 509–572.
- Nøttestad, L., Krafft, B. A., Anthonypillai, V., Bernasconi, M., Langård, L., Mørk, H. L., & Fernö, A. (2015). Recent changes in distribution and relative abundance of cetaceans in the Norwegian Sea and their relationship with potential prey. *Frontiers in Ecology and Evolution*, 2. <https://doi.org/10.3389/fevo.2014.00083>
- Nøttestad, L., Sivle, L. D., Krafft, B. A., Langard, L., Anthonypillai, V., Bernasconi, M., ... Ferno, A. (2014). Prey selection of offshore killer whales *Orcinus orca* in the Northeast Atlantic in late summer: Spatial associations with mackerel. *Marine Ecology Progress Series*, 499, 275–283. <https://doi.org/10.3354/meps10638>
- Parnell, A. (2016). *simmr*-package a stable isotope mixing model. Version 0.3.
- Parnell, A. C., Phillips, D. L., Bearhop, S., Semmens, B. X., Ward, E. J., Moore, J. W., ... Inger, R. (2013). Bayesian stable isotope mixing models. *Environmetrics*, 24, 387–399.
- Phillips, D. L., Inger, R., Bearhop, S., Jackson, A. L., Moore, J. W., Parnell, A. C., ... Ward, E. J. (2014). Best practices for use of stable isotope mixing models in food-web studies. *Canadian Journal of Zoology*, 92, 823–835. <https://doi.org/10.1139/cjz-2014-0127>
- Phillips, D. L., & Koch, P. L. (2002). Incorporating concentration dependence in stable isotope mixing models. *Oecologia*, 130, 114–125.
- R Development Core Team. (2017). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Reisinger, R. R., Grocke, D. R., Lubcker, N., McClymont, E. L., Hoelzel, A. R., & de Bruyn, P. J. N. (2016). Variation in the Diet of Killer Whales *Orcinus orca* at Marion Island, Southern Ocean. *Marine Ecology Progress Series*, 549, 263–274. <https://doi.org/10.3354/meps11676>
- Riesch, R., Barrett-Lennard, L. G., Ellis, G. M., Ford, J. K. B., & Deecke, V. B. (2012). Cultural traditions and the evolution of reproductive isolation: Ecological speciation in killer whales? *Biological Journal of the Linnean Society*, 106, 1–17.
- Røttingen, I. (1990). A review of variability in the distribution and abundance of Norwegian spring spawning herring and Barents Sea capelin. *Polar Research*, 8, 33–42.
- Ryan, C., McHugh, B., Trueman, C. N., Harrod, C., Berrow, S. D., & O'Connor, I. (2012). Accounting for the effects of lipids in stable isotope ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values) analysis of skin and blubber of balaenopterid whales. *Rapid Communications in Mass Spectrometry*, 26, 2745–2754. <https://doi.org/10.1002/rcm.6394>
- Samarra, F. I. P., Bassoi, M., Beesau, J., Eliasdottir, M. O., Gunnarsson, K., Mruscok, M. T., ... Vikingsson, G. A. (2018). Prey of killer whales (*Orcinus orca*) in Iceland. *PLoS ONE*, 13, e0207287.
- Samarra, F. I. P., & Miller, P. J. O. (2015). Prey-induced behavioural plasticity of herring-eating killer whales. *Marine Biology*, 162, 809–821. <https://doi.org/10.1007/s00227-015-2626-8>
- Samarra, F. I. P., Vighi, M., Aguilar, A., & Vikingsson, G. A. (2017). Intra-population variation in isotopic niche in herring-eating killer whales off Iceland. *Marine Ecology Progress Series*, 564, 199–210. <https://doi.org/10.3354/meps11998>
- Saulitis, E., Matkin, C., Barrett-Lennard, L., Heise, K., & Ellis, G. (2000). Foraging strategies of sympatric killer whale (*Orcinus orca*) populations in Prince William Sound, Alaska. *Marine Mammal Science*, 16, 94–109. <https://doi.org/10.1111/j.1748-7692.2000.tb00906.x>
- Scrucca, L., Fop, M., Murphy, T. B., & Raftery, A. E. (2016). mclust 5: Clustering, classification and density estimation using Gaussian finite mixture models. *The R Journal*, 8, 289–317. <https://doi.org/10.32614/RJ-2016-021>
- Similä, T., Holst, J. C., & Christensen, I. (1996). Occurrence and diet of killer whales in Northern Norway: Seasonal patterns relative to the distribution and abundance of Norwegian Spring-Spawning Herring. *Canadian Journal of Fisheries and Aquatic Sciences*, 53, 769–779. <https://doi.org/10.1139/f95-253>
- Similä, T., & Ugarte, F. (1993). Surface and underwater observations of cooperatively feeding killer whales in Northern Norway. *Canadian Journal of Zoology*, 71, 1494–1499. <https://doi.org/10.1139/z93-210>
- Tarroux, A., Ehrich, D., Lecomte, N., Jardine, T. D., Bêty, J., & Berteaux, D. (2010). Sensitivity of stable isotope mixing models to variation in isotopic ratios: Evaluating consequences of lipid extraction. *Methods in Ecology and Evolution*, 1, 231–241.
- Tixier, P., Giménez, J., Reisinger, R. R., Méndez-Fernandez, P., Arnould, J. P., Cherel, Y., & Guinet, C. (2019). Importance of toothfish in the diet of generalist subantarctic killer whales: Implications for fisheries

- interactions. *Marine Ecology Progress Series*, 613, 197–210. <https://doi.org/10.3354/meps12894>
- Vester, H., & Hammerschmidt, K. (2013). First record of killer whales (*Orcinus orca*) feeding on Atlantic salmon (*Salmo salar*) in Northern Norway suggest a multi-prey feeding type. *Marine Biodiversity Records*, 6, 1–5. <https://doi.org/10.1017/S1755267212001030>
- Vongraven, D., & Bisther, A. (2014). Prey switching by killer whales in the North-East Atlantic: Observational evidence and experimental insights. *Journal of the Marine Biological Association of the United Kingdom*, 94, 1357–1365. <https://doi.org/10.1017/s0025315413001707>
- Yurkowski, D. J., Hussey, N. E., Semeniuk, C., Ferguson, S. H., & Fisk, A. T. J. P. B. (2015). Effects of lipid extraction and the utility of lipid

normalization models on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in Arctic marine mammal tissues. *Polar Biology*, 38, 131–143. <https://doi.org/10.1007/s00300-014-1571-1>

How to cite this article: Jourdain E, Andvik C, Karoliussen R, Ruus A, Vongraven D, Borgå K. Isotopic niche differs between seal and fish-eating killer whales (*Orcinus orca*) in northern Norway. *Ecol Evol*. 2020;00:1–13. <https://doi.org/10.1002/ece3.6182>

PAPER V

At risk: Preying on seals pushes killer whales (*Orcinus orca*)
from Norway above pollution effects thresholds.

Andvik, C., Jourdain, E., Ruus, A., Lyche, J. L., Karoliussen, R., and Borgå, K.

Scientific Reports. 10:11888.

doi:10.1038/s41598-020-68659-y.

OPEN

Preying on seals pushes killer whales from Norway above pollution effects thresholds

Clare Andvik¹, Eve Jourdain², Anders Ruus^{1,3}, Jan L. Lyche⁴, Richard Karoliussen² & Katrine Borgå¹✉

Killer whales (*Orcinus orca*) are at risk from high levels of biomagnifying pollutants, such as polychlorinated biphenyls (PCBs) and mercury (Hg). Previous toxicological risk assessments for the Norwegian killer whale population have assumed fish as the primary prey source, and assessed the population as below established effect thresholds. However, some individuals have recently been identified to also feed on seals. This study is the first to quantify levels of pollutants in seal-eating killer whales from northern Norway, and to measure Hg levels in the skin of killer whales worldwide. We found higher levels of all pollutants in seal-eating than fish-eating killer whales, including the emerging brominated flame retardants pentabromoethylbenzene (PBEB), pentabromotoluene (PBT) and hexabromobenzene (HBB). Sum polychlorinated biphenyls (Σ PCBs) in the blubber of seal-eaters ($n=7$, geometric mean = 46 $\mu\text{g/g l.w.}$) were four times higher than fish-eaters ($n=24$, geometric mean = 11 $\mu\text{g/g l.w.}$), which pushed all seal-eating individuals above multiple thresholds for health effects. Total Hg levels in skin of seal-eaters ($n=10$, arithmetic mean = 3.7 $\mu\text{g/g d.w.}$) were twice as high as in fish-eaters ($n=28$, arithmetic mean = 1.8 $\mu\text{g/g d.w.}$). Our results indicate that by feeding on higher trophic prey, the Norwegian killer whale population is at higher risk of health effects from pollution than previously assumed.

Pollutants, such as polychlorinated biphenyls (PCBs) and mercury (Hg), have the potential to cause harmful effects in wildlife due to their persistent, bioaccumulative and toxic properties^{1,2}. PCBs are a type of organohalogen contaminant (OHC): anthropogenically produced pollutants that include a range of pesticides and industrial products. Hg is a naturally occurring element, however the concentrations measured in wildlife have increased by 3–4 orders of magnitude over the last 150 years due to anthropogenic activities³.

The killer whale (*Orcinus orca*) is a species particularly vulnerable to biomagnifying pollutants due to its position as an apex predator, long life span, and thick blubber layer in which lipophilic OHCs can accumulate⁴. High recorded tissue levels of PCBs and Hg led to the killer whale being named the most exposed species by the Arctic Monitoring and Assessment Programme (AMAP)⁵, and over 50% of killer whale populations worldwide have been predicted to collapse within the next 100 years due to PCB exposure⁶. The levels of pollutants in a killer whale population, and thus its risk of health effects, are higher in populations feeding at high trophic levels due to increased exposure from biomagnification in prey^{6,7}. For example, in the northeast Pacific, marine-mammal eating killer whale populations have higher levels of PCBs than fish-eating populations from the same geographic area^{8,9}. Variations in Hg levels according to dietary preference have not yet been recorded in killer whales, although higher Hg levels would be expected in groups feeding on marine mammal prey, as Hg also biomagnifies in food webs^{10,11}.

Killer whales frequenting the coast of Norway have long been assumed to feed primarily on Atlantic herring (*Clupea harengus*)^{12,13}. However, field observations and stable isotope dietary descriptors have recently confirmed that some individuals switch from fish to marine mammal prey, such as seals, on a long term and multi-seasonal basis^{14,15}. Toxicological risk assessments of the Norwegian killer whale population have been based on the PCB levels of nine, fish-eating killer whales sampled from one area of northern Norway in 2002^{5,6,16}. Hg levels have not been measured to date. With recent knowledge of the varied dietary preferences in the Norwegian killer

¹Department of Biosciences, University of Oslo, 0316 Oslo, Norway. ²Norwegian Orca Survey, Daniel Olaisensvei 6, 8480 Andenes, Norway. ³Norwegian Institute of Water Research, Gaustadalléen 21, 0349 Oslo, Norway. ⁴Department of Food Safety and Infection Biology, Norwegian University of Life Sciences, Campus Adamstuen, P.O. Box 5003, 1432 Ås, Norway. ✉email: katrine.borga@ibv.uio.no

Figure 1. ΣPCB levels (µg/g lipid weight) in blubber of seal-eating ($n=7$) and fish-eating ($n=24$) killer whales from Norway. **(a)** Levels in dietary groups. Horizontal lines represent the median, whiskers represent the lower (first) and upper (fourth) quartiles, dots represent individual whales, points outside the whiskers and box are outliers. **(b)** Levels in each individual whale, ordered on the x-axis in decreasing levels of ΣPCBs within dietary groups. The red horizontal lines (10 µg/g l.w. and 100 µg/g l.w.) are, respectively, the thresholds for a risk, and high risk, of health effects⁷.

whale population, along with changing environmental exposure, a quantification of the PCB and Hg levels of both fish and seal-eating killer whales is necessary for more accurate risk assessments.

The aim of this study was to quantify and compare the PCB levels in blubber, and total Hg levels in skin, of seal-eating and fish-eating killer whales in northern Norway. The assignment of whales to dietary groups was based on stable isotope analysis and observational data presented in a previous study¹⁴. Our results are relevant in conducting a more accurate risk assessment of contaminant-mediated health effects in this population, accounting for heterogeneity in foraging habits.

Results and discussion

PCB levels. We found higher sum of PCBs (ΣPCBs) levels in seal-eating killer whales than fish-eaters (Welch's t-test, $t=4.47$, $p=0.0006$; Fig. 1a, Table 1). The geometric mean ΣPCBs for adult male seal-eaters ($n=4$; 46 µg/g l.w.) was almost double the levels previously recorded in adult male fish-eating killer whales from Norway in 2002 ($n=8$; 27 µg/g l.w.)¹⁶. These lower levels from 2002 have been used to conclude that killer whales from Norway are at less risk from PCB-mediated health effects than other Arctic killer whale populations⁷. We found higher median ΣPCBs in adult male seal-eating killer whales from our study (51 µg/g l.w.), which were comparable to levels in an adult male killer whale harvested from Greenland with seal remains in the stomach (65 µg/g l.w.), suggesting equitable risk¹⁷.

PCBs are associated with immune suppression in marine mammals^{18,19}, and all the seal-eating killer whales from our study had ΣPCBs in blubber exceeding the 10 µg/g l.w. threshold for a risk of immune and hormone system health effects, determined from laboratory rat studies⁷, in contrast to only 46% of the fish-eating whales (Fig. 1b). Two of the seal-eating killer whales were even above the higher threshold for an elevated risk of health effects (100 µg/g l.w.). All seal-eating killer whales in our study (compared to 54% of fish-eaters) exceeded an additional threshold established using immune system endpoints in feeding experiments on harbour seals (*Phoca vitulina*), of 9 µg/g l.w.^{20,21}. The higher levels of PCBs in seal-eaters thus indicate potential weaknesses in the ability to fight diseases, which can have implications for survival and population growth^{22,23}.

PCBs also affect reproduction and offspring mortality in wildlife^{23,24}. In a recent study, the effects of PCBs on population growth was assessed via influences on calf survival and immune suppression by placing killer whale populations into exposure groups based on the ΣPCB levels in adult female killer whales⁶. The Norwegian killer whale population was placed in the lowest exposure group, 1, based on an estimated ΣPCB value in adult females of 10 µg/g l.w., extrapolated down from measured levels in adult males^{6,16}. Populations in exposure group 1 were

OHC group	Seal-eaters			Fish-eaters		
	All ^f (n = 7)	Adult male (n = 4)	Adult female (n = 2)	All ^f (n = 24)	Adult male (n = 17)	Adult female (n = 2)
ΣPCBs^a						
μ ± SD	56.0 ± 37.8	58.9 ± 35.9	64.6	16.7 ± 18.3	20.7 ± 20.4	4.07
Geometric mean	45.7	50.7	48.4	10.9	14.2	3.66
Median (range)	48.2 (21.8–108)	53.0 (22.0–108)	64.6 (21.8–107)	9.52 (2.3–77.1)	12.0 (4.03–77.1)	4.07 (2.3–5.85)
ΣDDTs^b						
μ ± SD	28.3 ± 21.1	30.9 ± 16.7	32.9	15.3 ± 21.4	19.3 ± 20.4	2.49
Geometric mean	21.4	27.7	19.9	8.59	11.4	2.20
Median (range)	26.8 (6.67–59.2)	27.5 (14.4–54.2)	32.9 (6.67–59.2)	7.79 (1.33–100)	8.96 (3.04–100)	2.49 (1.33–3.65)
ΣCHLs^c						
μ ± SD	9.78 ± 5.59	10.3 ± 3.75	11.6	4.35 ± 5.01	5.31 ± 5.65	1.17
Geometric mean	8.41	9.69	9.13	2.98	3.80	0.952
Median (range)	9.46 (4.06–18.9)	10.6 (5.54–14.4)	11.7 (4.41–18.9)	2.67 (0.489–23.4)	3.06 (1.52–23.4)	1.17 (0.489–1.85)
HCB						
μ ± SD	0.523 ± 0.467	0.389 ± 0.309	0.893	0.205 ± 0.0903	0.229 ± 0.0934	0.105
Geometric mean	0.401	0.324	0.650	0.18	0.216	0.0973
Median (range)	0.290 (0.220–0.146)	0.242 (0.220–0.852)	0.873 (0.290–1.45)	0.178 (0.0649–0.519)	0.194 (0.126–0.519)	0.105 (0.0649–0.146)
β-HCH						
μ ± SD	0.116 ± 0.090	0.0974 ± 0.0692	0.179	0.0470 ± 0.0369	0.0546 ± 0.0935	0.0186
Geometric mean	0.0929	0.0827	0.145	0.0378	0.0456	0.0146
Median (range)	0.0739 (0.0426–0.284)	0.074 (0.0426–0.199)	0.179 (0.0739–0.284)	0.0311 (<LOD–0.193)	0.0513 (<LOD–0.193)	0.0186 (<LOD–0.030)
Mirex						
μ ± SD	0.295 ± 0.148	0.256 ± 0.179	0.400	0.0700 ± 0.0777	0.0842 ± 0.0879	0.0384
Geometric mean	0.224	0.166	0.398	0.0425	0.0522	0.383
Median (range)	0.347 (<LOD–0.446)	0.283 (<LOD–0.437)	0.400 (0.354–0.446)	0.0449 (<LOD–0.350)	0.0478 (<LOD–0.304)	0.0384 (<LOD–0.0417)
ΣPBDEs^d						
μ ± SD	1.57 ± 0.73	1.61 ± 0.39	1.67	0.554 ± 0.415	0.491 ± 0.444	0.218
Geometric mean	1.41	1.58	1.22	0.447	0.553	0.180
Median (range)	1.38 (0.529–2.81)	1.51 (1.27–2.13)	1.67 (0.529–2.81)	0.466 (0.0960–1.79)	0.491 (0.258–1.79)	0.218 (0.0960–0.339)
HBB						
μ ± SD	0.009 ± 0.003	0.009 ± 0.004	0.010	0.003 ± 0.003	0.003 ± 0.004	0.002
Geometric mean	0.009	0.009	0.010	0.002	0.00	0.001
Median (range)	0.009 (0.006–0.014)	0.009 (0.006–0.014)	0.010 (0.010–0.010)	(0.0006–0.014)	0.002 (0.0006–0.014)	0.002 (0.001–0.002)
4'-OH-BDE49						
μ ± SD	0.008 ± 0.006	0.007 ± 0.004	0.010	0.003 ± 0.003	0.004 ± 0.004	0.0006
Geometric mean	0.006	0.006	0.007	0.001	0.002	0.0005
Median (range)	0.006 (0.002–0.018)	0.007 (0.002–0.011)	0.010 (0.003–0.018)	0.002 (< LOD–0.014)	0.003 (< LOD–0.014)	0.0006 (< LOD–0.0009)
ΣOHCs^e						
μ ± SD	96.6 ± 65.0	103 ± 56.2	112	37.3 ± 44.6	46.3 ± 50.2	8.11
Geometric mean	78.6	91.0	80.6	23.5	30.7	7.17
Median (range)	90.1 (34.1–191)	93.9 (43.6–178)	112 (34.1–191)	20.5 (4.31–202)	24.5 (9.10–202)	8.11 (4.31–11.9)

Table 1. Summary of lipid-normalised concentrations (μg/g) of organohalogen contaminants in sampled killer whales (n = 31). LOD Limit of detection. ^aSum of PCB-28, -66, -74, -87, -99, -101, -105, -110, -114, -118, -128, -137, -138, -141, -149, -151, -153, -156, -157, 170, -180, -183, -187, -189, 194, -196, -206, -209. ^bSum of p,p'-DDT, p,p'-DDD and p,p'-DDE. ^cSum of oxychlordane, *trans*-chlordane, *cis*-chlordane, *trans*-nonachlor and *cis*-nonachlor. ^dSum of BDE-28, -47, -99, -100, -153, -154. ^eSum of all organohalogen contaminants detected in more than 65% of whale samples. ^fIncluding whales of unknown sex, subadults, and of unknown age: summary statistics for these whales can be found in Supplementary Table S3.

modelled to double in size within the next 100 years, if model assumptions are accepted. In our study, however, the two adult seal-eating female killer whales had ΣPCB values of 22 μg/g l.w. and 107 μg/g l.w., which would place them in exposure groups 2 and 5, respectively. Populations in exposure group 2 are estimated to have modest population growth, albeit at a reduced rate, whereas populations in group 5 are estimated to be at high risk of population collapse within the next 100 years⁶. The geometric mean ΣPCB levels of all the seal-eating killer whales (46 μg/g l.w.), furthermore, exceeded the ΣPCB threshold of 41 μg/g l.w., at which reproductive impairment with

Figure 2. Total mercury levels in fish-eating ($n=28$) and seal-eating ($n=10$) killer whales from Norway. (a) Levels in skin ($\mu\text{g/g d.w.}$). Horizontal lines represent the median, whiskers represent the lower (first) and upper (fourth) quartiles, dots represent individual whales, points outside the whiskers and box are outliers. (b) Levels in liver ($\mu\text{g/g w.w.}$), estimated from skin using equation $\ln(Hg_{liver}) = 1.6124 \times \ln(Hg_{skin}) + 2.0346$ based on bottlenose dolphins (*Tursiops truncatus*)⁵⁸, and ordered on the x-axis in decreasing levels of total mercury within dietary groups. The red horizontal line ($16 \mu\text{g/g w.w.}$) is the threshold for a low risk of health effects in marine mammals, and ranges to $64 \mu\text{g/g w.w.}$ Levels below the threshold of $16 \mu\text{g/g w.w.}$ are classified as no risk of mercury-mediated health effects in marine mammals⁷.

occlusions in the uterus and increased glucocorticoid production in the adrenals was observed in ringed seals (*Pusa hispida*)^{21,24}. Killer whales from Norway are observed with viable offspring, and have an estimated calving rate of 0.197²⁵. However, this is based on killer whales studied on herring overwintering grounds, which are thus likely fish-eaters. Although we have a small sample size for adult female killer whales (just two individuals in each dietary group, with a large range), the higher levels in seal-eating killer whales may result in increased calf mortality. The difference in the geometric mean ΣPCB levels in female seal-eaters ($48 \mu\text{g/g l.w.}$) and female fish-eaters ($3.6 \mu\text{g/g l.w.}$) is translatable to approximately 45% and 80% of calves surviving, respectively²³. Median ΣPCB levels in female seal-eaters ($67 \mu\text{g/g l.w.}$) are furthermore comparable to the median ΣPCB levels in female, tuna-eating, killer whales from the Strait of Gibraltar ($51 \mu\text{g/g l.w.}$), which has one of the lowest reproductive rates for killer whales globally²¹. Lower reproductive rates in the seal-eating killer whales are relevant for the wider ecosystem, as seal-eating killer whales occupy a differentiated ecological niche than the fish-eaters, that could be lost with localised extinctions¹⁴.

Mercury. Seal-eating killer whales from our study had higher total Hg levels in skin than fish-eaters (arithmetic mean seal-eaters = $3.7 \mu\text{g/g d.w.}$ and fish-eaters = $1.8 \mu\text{g/g d.w.}$; Welch's t-test, $t=4.89$, $p=0.0008$; Fig. 2a, Table 2). The estimated hepatic total Hg levels for both seal-eaters (arithmetic mean: $13 \mu\text{g/g w.w.}$) and fish-eaters (arithmetic mean: $3.6 \mu\text{g/g w.w.}$) were below the threshold for liver abnormalities in bottlenose dolphins (*Tursiops truncatus*) ($61 \mu\text{g/g w.w.}$)²⁶, and below the threshold for a low risk of health effects in marine mammals ($16 \mu\text{g/g w.w.}$)⁷, based on toxic hepatitis, uremia and renal failure observed in harp seals (*Pagophilus groenlandicus*)²⁷. However, four of the seal-eaters (40%) were above the $16 \mu\text{g/g w.w.}$ threshold, compared to none of the fish-eaters, still indicating a higher risk of health effects for the seal-eating group (Fig. 2b). This is the first indication of mercury levels in killer whales from Norway to date. It is also the first study worldwide to measure Hg levels in the skin of killer whales, thus allowing for repeatable and minimally invasive biomonitoring. Whilst levels appear low, the extrapolation from liver to skin in our study has uncertainties, and the skin levels are comparable to at-risk populations of bottlenose dolphins from the English Channel²⁸ and Mediterranean²⁹. There was a positive correlation between the total Hg levels measured in the skin and the total OHC levels measured in the blubber of the whales (Spearman's rank correlation, $S=2,572$, $p=0.006$, $\rho=0.48$).

Total mercury levels	Seal-eaters			Fish-eaters		
	All ^a (n = 10)	Adult male (n = 5)	Adult female (n = 4)	All ^a (n = 28)	Adult male (n = 20)	Adult female (n = 3)
Measured in skin (µg/g d.w.)						
µ ± SD	3.73 ± 1.26	3.52 ± 1.66	3.97 ± 0.95	1.76 ± 0.32	1.78 ± 0.27	3.19 ± 1.49
Geometric mean	3.47	3.11	3.87	1.73	1.76	2.98
Median (range)	3.95 (1.19–5.44)	3.71 (1.19–5.44)	4.18 (2.62–4.88)	1.79 (1.01–2.40)	1.83 (1.29–2.25)	2.62 (2.08–4.88)
Estimated in liver (µg/g w.w.)^b						
µ ± SD	12.5 ± 5.62	11.6 ± 7.1	12.5 ± 4.47	3.57 ± 1.05	3.68 ± 0.987	3.70 ± 0.531
Geometric mean	10.5	8.77	11.8	3.41	3.54	3.68
Median (range)	14.2 (1.78–18.5)	14.3 (1.78–18.5)	13.2 (6.48–17.1)	3.47 (1.49–5.65)	3.61 (1.57–5.65)	3.411 (3.38–4.31)

Table 2. Summary of total mercury levels in sampled killer whales (n = 38). ^aIncluding whales of unknown sex, subadults, and of unknown age; summary statistics for these whales found in Supplementary Table S4.

^bEstimated using equation $\ln(Hg_{liver}) = 1.6124 \times \ln(Hg_{skin}) + 2.0346$ based on bottlenose dolphins (*Tursiops truncatus*)⁵⁸.

Figure 3. Principal component analysis (PCA) biplot based on organohalogen contaminant levels in blubber of killer whales from Norway. Response loadings are represented as black arrows, and passive explanatory variables are added as orange symbols representing the centroids of the variable (age, sex, dietary group). “S” refers to the centroid for seal-eaters (n = 7) and “F” for fish-eaters (n = 24). The centroid for males (n = 22), females (n = 4) and unknown sex (n = 5) are represented as ♂, ♀ and U, respectively. “A” refers to the centroid for the adult whales (n = 25), “SA” for subadults (n = 3), and “?” if the age is unknown (n = 3). The percentage of the total variation explained by PC1 and PC2 are given in brackets on each axis.

Other organohalogen contaminants. Levels of other organohalogen contaminants (OHCs), including the pesticide dichlorodiphenyltrichloroethane (DDT) and the industrial chemicals polybrominated diphenyl ethers (PBDEs) were also all higher in the seal-eating killer whales than the fish-eaters (RDA, $F_{1,28} = 10.53$, $p = 0.001$; Fig. 3). Threshold levels for possible toxic effects in marine mammals have not been established for these pollutants. However, many have been linked to oestrogenic, reproductive and endocrine effects in wildlife^{30,31}. The Stockholm Convention on Persistent Organic Pollutants has banned many of these pollutants³², and whilst downward trends have been observed in the environment³³, levels are still high and the geometric mean ΣPCB (14 µg/g l.w.), ΣDDT (11 µg/g l.w.) and ΣPBDE (0.55 µg/g l.w.) levels in adult male fish-eating killer whales sampled for this study were comparable to geometric mean levels in blubber of male killer whales sampled in Norway in 2002 (ΣPCB: 27 µg/g l.w., p,p'-DDE: 11 µg/g l.w., ΣPBDE: 0.48 µg/g l.w.)¹⁶. We also screened for four emerging brominated contaminants in the tissues of both seal and fish-eating sampled killer whales. Whilst levels of 3-dibromopropyl-2,4,6-tribromophenyl ether (DPTE) were below the limit of detection in all whales, pentabromoethylbenzene (PBEB) was found in one whale (a seal-eater), pentabromotoluene (PBT) was

found in 58% of the sampled whales (86% of the seal-eaters and 50% of the fish-eaters) and hexabromobenzene (HBB) was found in all individuals. HBB, included in the PCA, was correlated to the other organohalogen contaminants (Fig. 3). The presence of these emerging contaminants in top predators is a concern, as production is unregulated and the effects on marine mammals unknown.

ΣPCBs and ΣDDTs dominated the total OHC load for both killer whale dietary clusters, together contributing 87% and 86% of the total OHC burden for seal-eaters and fish-eaters, respectively (Table 1). PCB congeners – 99, – 138, – 153 and – 180 accounted for 67% (for seal-eaters) and 58% (for fish-eaters) of the total ΣPCB load, with PCB-153 contributing the most. The ΣDDTs were dominated by p,p'-DDE, contributing 90% of the total ΣDDTs for the seal-eaters and 92% for fish-eaters. The least contributing were the BFRs (1.5% for seal-eaters and 1.3% for fish-eaters of total OHC load) and OH-metabolites (0.01% for seal-eaters and 0.007% for fish-eaters of total OHC load).

Influence of sex/age class on pollutant levels. Previous studies have shown age-dependent increases for both OHC and total Hg in killer whales due to age accumulation^{34,35}, and lower OHC levels in reproductive females due to maternal offload^{17,36–38}. These factors were not possible to be tested in the present study due to a small sample size for adult female individuals in each dietary group (Tables 1, 2), the undetermined sex of five whales, and no possibility to obtain the precise age of the whales. The influence of these factors on the pollutant levels will be a focus for further research as our knowledge of the sampled whales, and their population, increases.

Implications for dietary ecology. The higher OHC and total Hg levels in seal-eaters indicate a long-term preference for higher trophic prey, in accordance with field observations and stable isotope analysis^{14,15}. The PCB patterns further support long-term preferences for marine mammals, with the seal-eating killer whales having a higher proportion of hexaCBs, heptaCBs, and higher CBs than the fish-eating killer whales (Welch's t-test, $t_{28} = 5.38$, $p = 0.001$; Supplementary Fig. S1). Higher-chlorinated PCBs are often found in marine mammals relative to fish^{39,40}, and higher proportions of hexaCBs and heptaCBs have been found in marine mammal-eating killer whales compared to fish-eating whales in the Northeast Pacific^{8,9}, and in seal-eating whales in Greenland compared to fish-eating whales in the Faroe Islands¹⁷.

There is a 20-fold difference in median ΣPCB levels between two killer whale populations inhabiting the same area of the northeast Pacific, one of which solely eats marine mammals and the other fish^{9,41}. This is a larger difference than the five-fold difference between median ΣPCB levels in fish and seal-eating killer whales in our study, and thus may indicate support for previous studies proposing a mixed diet for the seal-eating whales, in which fish prey is an equal or secondary food source to seal prey¹⁴. It is theorised that some killer whales in Norway diversified to eating seals as a response to the collapse of the herring fisheries in the 1970s¹⁵. Future changes in prey availability could mean that more killer whales switch to eating seal and/or incorporate a higher proportion of seals in the diet, with subsequent health and population effects. Our results illustrate that differentiations in dietary preferences need to be accounted for in effective risk assessment, management and conservation of the environment.

Methods

Sampling. Killer whale biopsy samples of skin and blubber from 38 individuals were collected year-round from August 2017 to July 2018 in northern Norway. All whales were sampled according to relevant guidelines and regulations, and conducted under the permit FOTS-ID 10176 issued by Mattilsynet (the Norwegian Food Safety Authority, report nr. 2016/179856). Details of seasonal sampling locations, stable isotope dietary descriptors and classification of sampled individuals are described in a previous study¹⁴. In the current study, total Hg was analysed in skin from all individuals ($n = 38$), whereas organohalogen contaminants (OHC) was analysed in blubber of 31 individuals due to insufficient blubber for the remaining 7 individuals.

OHC analysis. OHC analysis was conducted at the Laboratory of Environmental Toxicology at the Norwegian University of Life Sciences, Oslo, Norway. We analysed a total of 83 OHCs: 49 organochlorines (OCs), including 34 PCBs and 15 organochlorine pesticides (OCPs), 18 brominated flame retardants (BFRs), including newer and unregulated compounds, and 16 hydroxylated metabolites (OH-metabolites) of PCBs and polybrominated diphenylethers (PBDEs). A full list of analysed compounds can be found in Supplementary Table S1.

We analysed OCs and BFRs using a multicomponent method, first described in 1978⁴², and since modified for a range of compounds and biological matrices^{43–46}. The analysis of the OH-metabolites was conducted according to previously published methods^{47,48}. An outline of the method is described in the Supplementary Information. Reported concentrations were blank corrected based on the average concentration detected within blank samples. The limit of detection (LOD) was defined as three times the average noise in chromatograms, and ranged from 0.40 to 11.10 ng/g w.w. for OCs, 0.012 to 0.362 ng/g w.w. for BFRs and 0.013 to 0.040 ng/g w.w. for OH-metabolites (see Supplementary Table S2). Internal reference materials for OCs and BFRs (contaminated seal blubber, MTref01) and OH-metabolites (contaminated seal blood, MTref03) were also extracted in conjunction with sample material to assess method performance. Internal standard recoveries are listed in Supplementary Table S2.

Hg analysis. We analysed total Hg by atomic absorption spectrometry at the University of Oslo, using a Direct Mercury Analyser (DMA-80, Milestone Srl, Soirsole, Italy). Killer whale skin samples were freeze dried in a Leybold-Heraeus GT2 freeze dryer with a Leybold Vakuum GmbH vacuum pump (Leybold, Cologne, Germany) and then homogenised to a fine powder using an agate pestle and mortar. Approximately 0.002 g of killer whale skin were analysed in parallel with sample blanks and certified reference material (DORM-4, fish protein;

DOLT-5, dogfish liver, National Research Council, Ottawa, Canada). If enough material, samples were analysed in duplicates to ensure precision of measurements and the arithmetic mean value used. Average recoveries of the certified reference materials were within 10% of the reported values. The detection limit of the instrument was 0.05 ng mercury.

Data treatment. We included OHC compounds found in levels above the instrument's LOD in a minimum of 65% of the individual whale samples for statistical analysis (see Supplementary Table S1, Supporting Information for pollutants excluded). For individual concentrations below the LOD, we imputed left-censored data by replacing missing values with a random number between 0 and the LOD assuming a beta distribution ($\alpha = 5$, $\beta = 1$) to retain the pattern of the dataset. In total, 95 values below the LOD were replaced, representing 6.52% of the OHC dataset. All total Hg samples were above the LOD.

We defined the Σ PCBs as the sum of all 28 PCB congeners detected in more than 65% of the whale samples (PCB-28, -66, -74, -87, -99, -101, 105, -110, -114, -118, -128, -137, -138, -141, -149, -151, -153, -156-, 157, 170, -180, -183, -187, -189, 194, -196, -206, -209). The definition for Σ PCBs varies within killer whale literature, with some studies analysing only a few core PCB congeners³⁵, some all 209 of the possible congeners³⁶, and others not providing a definition (e.g. for thresholds for possible health effects³⁷). There will therefore inevitably be some errors in comparisons. However, since the Σ PCBs in killer whales is dominated by a few commonly reported congeners, typically PCB-153 and -138^{16,37}, it is unlikely that the inclusion of other minor constituents will have a major influence on the total load. PCBs were further grouped according to the number of chlorine substitutions per molecule, i.e. homologue group to compare the pattern of PCBs. Σ DDTs was defined as the sum of p,p'-DDT, p,p'-DDD and p,p'-DDE, the Σ PBDEs as the sum of BDE-28, -47, -99, -100, -153 and -154 and the sum of chlor-danes (Σ CHLs) as the sum of oxychlordane, *trans*-chlordane, *cis*-chlordane, *trans*-nonachlor and *cis*-nonachlor.

Statistical analyses. Statistical analyses were performed using R v. 3.4.1⁴⁹. The significance level was set to $\alpha = 0.05$, except in cases where the value was adjusted due to multiple testing, and was two-tailed. In addition to visual inspection, normality was tested using the Shapiro–Wilk's test⁵⁰ and homogeneity of variance by Levene's test⁵¹ using the R package *car*⁵².

Whale dietary groups. The dietary groups used in this study are based on a previous study, which used stable isotope values inputted into a Gaussian mixture model to assign sampled individuals to two fish-eating groups: *Herring-eaters* and *Lumpfish-eaters* and one mammal-eating group *Seal-eaters*¹⁴. The three dietary groups were characterised by disparate, non-overlapping isotopic niches that were consistent with predatory field observations. The seal-eating group was defined by higher $\delta^{15}\text{N}$ values than the two fish-eating groups.

We found that the herring and lumpfish-eating killer whales did not differ in either their OHC levels (Tukey's HSD: $p = 0.49$) or total Hg levels (pairwise Welch's t-test: $p = 0.67$). In this study, we thus combined the dietary groups *Herring-eaters* and *Lumpfish-eaters* into the group *Fish-eaters*, to enable easier comparison to the seal-eating killer whales.

We then used Welch's t-test to compare the Σ PCB levels in the seal-eating and fish-eating dietary groups (using a log10 transformation), and to compare the total Hg levels in the skin between the two dietary groups.

OHC dataset. We used multivariate analysis to compare and visualise the differences in all the OHCs between the dietary groups, age and sex classes using the *vegan* package in R⁵³. Principle Component Analysis (PCA) was used to visualise the main structure of the data: reducing the dimensions to two new, uncorrelated, latent variables termed principle components 1 and 2 (PC1 and PC2). We log-10 transformed contaminant levels to ensure normality and homogeneity of variance, and the presence of any influential outliers were checked by the Cook's distance test. Redundancy Analysis (RDA) was used to extract and summarise the variation in the OHC levels constrained, and thereby explained, by a set of explanatory variables⁵⁴. Significant associations between response variables and the explanatory variables were identified by an RDA based forward model selection, followed by a Monte Carlo forward permutation test (1,000 unrestricted permutations). The samples' scores along PC1 were subject to one-way Analysis of Variance (ANOVA) followed by Tukey's honestly significant difference post hoc test (Tukey's HSD) to analyse differences between the three dietary groups. PC1 scores were also used to evaluate correlation to total Hg levels in the skin using a Spearman's rank correlation test. Absolute concentrations were subject to PCA with lipid % as a covariate, after checking its significance using RDA, as lipid normalising data in inferential statistics can often lead to misleading conclusions⁵⁵.

We lipid-normalised OHC values when comparing levels to threshold values for toxicity or other killer whale populations, and used the geometric mean as the average for each dietary group to reflect the log normal distribution of the data. In accordance with convention, efforts were made to only compare adult males with other worldwide populations, as reproductive female whales are known to transfer a substantial portion of their OHC burden to their calves^{35,36,38}. In any case of comparison, similar metrics were compared (i.e. arithmetic mean, geometric mean, median) and all variables kept similar (i.e. sex, age, biopsy/stranded animals). We make the assumption in this study that the killer whales sampled in 2002 in Norway were fish-eaters for the following reasons: firstly, the whales were sampled on herring overwintering grounds, feeding on herring, and photographs were taken of five of the eight adults sampled and were identified as herring-eaters from previous field observations¹⁶. Secondly, the PCB pattern in the sampled whales showed 76% of Σ PCBs higher chlorinated congeners (hexaCBs or higher), which is more similar to the fish-eaters from our study (80% higher chlorinated congeners) than the seal-eaters (87% higher chlorinated congeners). Thirdly, the upper 95% confidence range of all pollutants reported in the 2002 killer whales falls below both the geometric and arithmetic mean values for seal-eaters from this study.

Total Hg dataset. The normal distribution of the data within each dietary group meant we used the arithmetic mean as an average. The three dietary groups (*Herring-eaters*, *Lumpfish-eaters* and *Seal-eaters*) were compared using a pairwise Welch's t-test with a Benjamini–Hochberg False Discovery Rate correction to adjust for multiple testing. Because we found no difference between the *Herring-eaters* and *Lumpfish-eaters* ($p = 0.67$), we combined these two groups to a group called “*Fish-eaters*” for easier comparison with the seal-eaters. The total Hg levels in the skin of the two groups, *Fish-eaters* and *Seal-eaters* were compared using Welch's t-test.

There is a strong positive correlation between Hg levels in the skin and liver in toothed whales, and this can be used to compare Hg levels measured in skin with hepatic toxicity threshold values^{56–58}. To extrapolate to liver from skin in our samples, we chose an equation based on a model using concentrations in the liver (Hg_{liver} $\mu\text{g/g}$ w.w) and skin (Hg_{skin} $\mu\text{g/g}$ w.w) of bottlenose dolphins (*Tursiops truncatus*) (Eq. 1)⁵⁸. We converted dry weight to wet weight using the water content for each individual whale measured during freeze drying.

$$\ln(Hg_{liver}) = 1.6124 \times \ln(Hg_{skin}) + 2.0346 \quad (1)$$

When comparing Hg concentrations to other worldwide populations, both male and female whales were included. This was due to a lack of information of sex in one of the populations for comparisons and because killer whales are unlikely to pass on Hg burdens to calves^{5,59}.

Data availability

All data is available in the main text or the Supplementary information.

Received: 15 November 2019; Accepted: 25 June 2020

Published online: 17 July 2020

References

- Letcher, R. J. *et al.* Exposure and effects assessment of persistent organohalogen contaminants in Arctic wildlife and fish. *Sci. Total Environ.* **408**, 2995–3043 (2010).
- Dietz, R. *et al.* What are the toxicological effects of mercury in Arctic biota?. *Sci. Total Environ.* **443**, 775–790 (2013).
- Dietz, R. *et al.* Three decades (1983–2010) of contaminant trends in East Greenland polar bears (*Ursus maritimus*). Part 1: Legacy organochlorine contaminants. *Environ. Int.* **59**, 485–493 (2013).
- Tanabe, S. PCB problems in the future: Foresight from current knowledge. *Environ. Pollut.* **50**, 5–28 (1988).
- AMAP. Arctic Monitoring and Assessment Programme: AMAP Assessment Report-Biological effects of contaminants on Arctic wildlife and fish, Tromsø, Norway (2018).
- Desforges, J.-P. *et al.* Predicting global killer whale population collapse from PCB pollution. *Science* **361**, 1373–1376 (2018).
- Dietz, R. *et al.* Current state of knowledge on biological effects from contaminants on arctic wildlife and fish. *Sci. Total Environ.* **696**, 133792 (2019).
- Herman, D. P. *et al.* Feeding ecology of Eastern North Pacific killer whales *Orcinus orca* from fatty acid, stable isotope, and organochlorine analyses of blubber biopsies. *Mar. Ecol. Prog. Ser.* **302**, 275–291 (2005).
- Buckman, A. H. *et al.* PCB-associated changes in mRNA expression in killer whales (*Orcinus orca*) from the NE Pacific ocean. *Environ. Sci. Technol.* **45**, 10194–10202 (2011).
- Ruus, A. *et al.* Methylmercury biomagnification in an Arctic pelagic food web. *Environ. Toxicol. Chem.* **34**, 2636–2643 (2015).
- Tokar, E. J., Boyd, W. A., Freedman, J. H. & Waalks, M. P. Toxic effects of metals. In *Casarett & Doull's Essentials of Toxicology* (eds Klaasen, C. D. & Watkins, J. B., III.) 347–359 (McGraw-Hill Education, New York, 2015).
- Bisther, A. & Vongraven, D. Studies of the social ecology of Norwegian killer whales (*Orcinus orca*). In *Whales, Seals, Fish and Man* (eds Blix, A. S. *et al.*) 169–176 (Elsevier Science, Amsterdam, 1995). [https://doi.org/10.1016/S0163-6995\(06\)80020-X](https://doi.org/10.1016/S0163-6995(06)80020-X).
- Similä, T., Holst, J. C. & Christensen, I. Occurrence and diet of killer whales in northern Norway: Seasonal patterns relative to the distribution and abundance of Norwegian spring-spawning herring. *Can. J. Fish. Aquat. Sci.* **53**, 769–779 (1996).
- Jourdain, E. *et al.* Isotopic niche differs between seal and fish-eating killer whales (*Orcinus orca*) in northern Norway. *Ecol. Evol.* <https://doi.org/10.1002/ece3.6182> (2020).
- Jourdain, E., Vongraven, D., Bisther, A. & Karoliussen, R. First longitudinal study of seal-feeding killer whales (*Orcinus orca*) in Norwegian coastal waters. *PLoS ONE* **12**, 1–14 (2017).
- Wolkers, H., Corkeron, P. J., Van Parijs, S. M., Simila, T. & Van Bavel, B. Accumulation and transfer of contaminants in killer whales (*Orcinus orca*) from Norway: Indications for contaminant metabolism. *Environ. Toxicol. Chem.* **26**, 1582–1590 (2007).
- Pedro, S. *et al.* Blubber-depth distribution and bioaccumulation of PCBs and organochlorine pesticides in Arctic-invading killer whales. *Sci. Total Environ.* **601–602**, 237–246 (2017).
- Desforges, J.-P. *et al.* Immunotoxic effects of environmental pollutants in marine mammals. *Environ. Int.* **86**, 126–139 (2016).
- Mori, C., Morsey, B., Levin, M., Nambiar, P. R. & De Guise, S. Immunomodulatory effects of in vitro exposure to organochlorines on T-cell proliferation in marine mammals and mice. *J. Toxicol. Environ. Health Part A* **69**, 283–302 (2006).
- Kannan, K., Blankenship, A. L., Jones, P. D. & Giesy, J. P. Toxicity reference values for the toxic effects of polychlorinated biphenyls to aquatic mammals. *Hum. Ecol. Risk Assess. Int. J.* **6**, 181–201 (2000).
- Jepson, P. D. *et al.* PCB pollution continues to impact populations of orcas and other dolphins in European waters. *Sci. Rep.* **6**, 1–17 (2016).
- Luster, M. *et al.* Risk assessment in immunotoxicology. II. Relationships between immune and host resistance tests. *Fundam. Appl. Toxicol.* **21**, 71–82 (1993).
- Hall, A. J. *et al.* Predicting the effects of polychlorinated biphenyls on cetacean populations through impacts on immunity and calf survival. *Environ. Pollut.* **233**, 407–418 (2018).
- Helle, E., Olsson, M. & Jensen, S. DDT and PCB levels and reproduction in ringed seal from the Bothnian Bay. *AMBIO A J. Hum. Environ.* **5**, 188–189 (1976).
- Kuningas, S., Similä, T. & Hammond, P. S. Population size, survival and reproductive rates of northern Norwegian killer whales (*Orcinus orca*) in 1986–2003. *J. Mar. Biol. Assoc. U.K.* **94**, 1277–1291 (2014).
- Rawson, A. J., Patton, G. W., Hofmann, S., Pietra, G. G. & Johns, L. Liver abnormalities associated with chronic mercury accumulation in stranded Atlantic bottlenose dolphins. *Ecotoxicol. Environ. Saf.* **25**, 41–47 (1993).
- Ronald, K., Tessaro, S. V., Uthe, J. F., Freeman, H. C. & Frank, R. Methylmercury poisoning in the harp seal (*Pagophilus groenlandicus*). *Sci. Total Environ.* **8**, 1–11 (1977).
- Zanuttini, C. *et al.* High pollutant exposure of bottlenose dolphins from the largest European community in the English Channel. *Sci. Rep.* **9**, 1–10 (2019).

29. Frodello, J. P., Romeo, M. & Viale, D. Distribution of mercury in the organs and tissues of five toothed-whale species of the Mediterranean. *Environ. Pollut.* **108**, 447–452 (2000).
30. Wójtowicz, A. K., Kajta, M. & Gregoraszczuk, E. L. DDT- and DDE-induced disruption of ovarian steroidogenesis in prepubertal porcine ovarian follicles: A possible interaction with the main steroidogenic enzymes and estrogen receptor beta. *J. Physiol. Pharmacol.* **58**, 873–885 (2007).
31. Hamers, T. *et al.* In vitro profiling of the endocrine-disrupting potency of brominated flame retardants. *Toxicol. Sci.* **92**, 157–173 (2006).
32. United Nations Environment Programme. *Stockholm convention on persistent organic pollutants (POPs)*. Geneva, Switzerland: United Nations Environment Programme. (2001). https://www.wipo.int/edocs/lexdocs/treaties/en/unep-pop/trt_unep_pop_2.pdf.
33. Rigét, F. *et al.* Temporal trends of persistent organic pollutants in Arctic marine and freshwater biota. *Sci. Total Environ.* **649**, 99–110 (2019).
34. Endo, T., Kimura, O., Hisamichi, Y., Minoshima, Y. & Haraguchi, K. Age-dependent accumulation of heavy metals in a pod of killer whales (*Orcinus orca*) stranded in the northern area of Japan. *Chemosphere* **67**, 51–59 (2006).
35. Krahn, M. M. *et al.* Effects of age, sex and reproductive status on persistent organic pollutant concentrations in 'Southern Resident' killer whales. *Mar. Pollut. Bull.* **58**, 1522–1529 (2009).
36. Ross, P. S., Ellis, G. M., Ikononou, M. G., Barrett-Lennard, L. G. & Addison, R. F. High PCB concentrations in free-ranging Pacific killer whales (*Orcinus orca*): Effects of age, sex and dietary preference. *Mar. Pollut. Bull.* **40**, 504–515 (2000).
37. Ylitalo, G. M. *et al.* Influence of life-history parameters on organochlorine concentrations in free-ranging killer whales (*Orcinus orca*) from Prince William Sound, AK. *Sci. Total Environ.* **281**, 183–203 (2001).
38. Haraguchi, K., Hisamichi, Y. & Endo, T. Accumulation and mother-to-calf transfer of anthropogenic and natural organohalogenes in killer whales (*Orcinus orca*) stranded on the Pacific coast of Japan. *Sci. Total Environ.* **407**, 2853–2859 (2009).
39. Hoekstra, P. F. *et al.* Trophic transfer of persistent organochlorine contaminants (OCs) within an Arctic marine food web from the southern Beaufort-Chukchi Seas. *Environ. Pollut.* **124**, 509–522 (2003).
40. Sobek, A. *et al.* A comparison of PCB bioaccumulation factors between an Arctic and a temperate marine food web. *Sci. Total Environ.* **408**, 2753–2760 (2010).
41. Ford, J. K. B. *et al.* Dietary specialization in two sympatric populations of killer whales (*Orcinus orca*) in coastal British Columbia and adjacent waters. *Can. J. Zool.* **76**, 1456–1471 (1998).
42. Brevik, E. M. Gas chromatographic method for the determination of organochlorine pesticides in human milk. *Bull. Environ. Contam. Toxicol.* **19**, 281–286 (1978).
43. Bernhoft, A. & Skaare, J. U. Levels of selected individual polychlorinated biphenyls in different tissues of harbour seals (*Phoca vitulina*) from the southern coast of Norway. *Environ. Pollut.* **86**, 99–107 (1994).
44. Andersen, M. *et al.* Geographic variation of PCB congeners in polar bears (*Ursus maritimus*) from Svalbard east to the Chukchi Sea. *Polar Biol.* **24**, 231–238 (2001).
45. Polder, A. *et al.* Levels and patterns of persistent organic pollutants (POPs) in tilapia (*Oreochromis* sp.) from four different lakes in Tanzania: Geographical differences and implications for human health. *Sci. Total Environ.* **488–489**, 252–260 (2014).
46. Polder, A. *et al.* Spatial and temporal changes of chlorinated pesticides, PCBs, dioxins (PCDDs/PCDFs) and brominated flame retardants in human breast milk from Northern Russia. *Sci. Total Environ.* **391**, 41–54 (2008).
47. Løken, K. B., Lie, E., Lundanes, E. & Skåre, J. U. Extension of a multicomponent method to include the determination of OH-PCBs and OH-PBDEs in biological matrices. *Organohalogen Compd.* **68**, 2430–2433 (2006).
48. Gabrielsen, K. M. *et al.* Levels and patterns of hydroxylated polychlorinated biphenyls (OH-PCBs) and their associations with thyroid hormones in hooded seal (*Cystophora cristata*) mother–pup pairs. *Aquat. Toxicol.* **105**, 482–491 (2011).
49. R Development Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna. <https://www.R-project.org/>. (2019).
50. Shapiro, S. S. & Wilk, M. B. An analysis of variance test for normality (complete samples). *Biometrika* **52**, 591–611 (1965).
51. Levene, H. A robust approximate confidence-interval for components of variance. *Ann. Math. Stat.* **31**, 534–535 (1960).
52. Fox, J. & Weisberg, S. *An R Companion to Applied Regression* (Sage, Thousand Oaks, 2011).
53. Oksanen, J. *et al.* *vegan: Community Ecology Package*. R package version 2.5-4. (2019).
54. Palmer, M. W., McGlenn, D. J., Westerberg, L. & Milberg, P. Indices for detecting differences in species composition: Some simplifications of RDA and CCA. *Ecology* **89**, 1769–1771 (2008).
55. Hebert, C. E. & Keenleyside, K. A. To normalize or not to normalize? Fat is the question. *Environ. Toxicol. Chem.* **14**, 801–807 (1995).
56. Aubail, A. *et al.* Use of skin and blubber tissues of small cetaceans to assess the trace element content of internal organs. *Mar. Pollut. Bull.* **76**, 158–169 (2013).
57. Cáceres-Saez, I., Goodall, R. N. P., Dellabianca, N. A., Cappozzo, H. L. & Guevara, S. R. The skin of Commerson's dolphins (*Cephalorhynchus commersonii*) as a biomonitor of mercury and selenium in Subantarctic waters. *Chemosphere* **138**, 735–743 (2015).
58. Stavros, H.-C.W. *et al.* Correlation and toxicological inference of trace elements in tissues from stranded and free-ranging bottlenose dolphins (*Tursiops truncatus*). *Chemosphere* **82**, 1649–1661 (2011).
59. Endo, T. *et al.* Distribution of total mercury, methyl mercury and selenium in pod of killer whales (*Orcinus orca*) stranded in the northern area of Japan: Comparison of mature females with calves. *Environ. Pollut.* **144**, 145–150 (2006).

Acknowledgements

We thank Malin Røyset Aarønes, Helene Thorstensen and Nils Christian Stenseth for helpful comments on the manuscript, as well as those from two anonymous reviewers. Research was supported by grants from the Sea World and Busch Gardens Conservation fund and Sea World and Parks Entertainment to Norwegian Orca Survey. C.A. was supported by a student stipend from Tekna.

Author contributions

C.A., E.J., A.R., J.L.L., and K.B. designed the study. C.A. conducted laboratory work. C.A. generated figures and performed all data analyses. E.J. and R.K. acquired funds. E.J., R.K., and C.A. obtained the samples for analysis. All authors were involved in interpretation of results and writing the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-020-68659-y>.

Correspondence and requests for materials should be addressed to K.B.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020