Acute Lisfranc Injuries

Epidemiology, diagnostic challenges and treatment

Thesis by

Are Haukåen Stødle

© Are Haukåen Stødle, 2021

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-871-7

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

Populærvitenskapelig sammendrag (norsk)

Akutte Lisfranc skader

- Epidemiologi, diagnostiske utfordringer og behandling

Lisfranc skader er skader i mellomfoten som kan bestå av både leddbånd- og bruddskader.

Skadene kan være utfordrende å diagnostisere og dersom en ustabil Lisfranc skade overses kan dette medføre betydelige plager for pasientene. En større oppmerksomhet rundt disse skadene og mer avanserte metoder i diagnostiseringen av dem har gjort at flere Lisfranc skader oppdages.

I sin doktoravhandling har Are Haukåen Stødle vist at forekomsten av Lisfranc skader er betydelig høyere enn tidligere rapportert og at de fleste kan relateres til lav-energi skademekanismer. Lisfranc skader overses relativt ofte på vanlige røntgenbilder, men avdekkes ved CT undersøkelse. Ved undersøkelse av forekomsten var over halvparten av skadene var stabile, det vil si at man ikke klarte å fremprovosere feilstilling i leddene i mellomfoten. De stabile skadene ble behandlet med gips og avlastning i 6 uker og hadde et utmerket sluttresultat. Hvilken behandling som er best for de ustabile skadene der det er feilstilling i leddene i mellomfoten, er omdiskutert. Stødle og kollegaer har sammenlignet to operasjonsmetoder for behandling av de ustabile Lisfranc skadene. Den ene gruppen fikk avstivet ledd i mellomfoten permanent mens den andre gruppen fikk kun midlertidig avstivet leddet ved hjelp av plate og skruer. Det var ingen tydelig forskjell i funksjonelt resultat mellom

gruppene. Gruppen med midlertidig avstiving hadde en mer anatomisk stilling i mellomfoten etter kirurgi, men hadde også en høy forekomst av radiologisk påvist slitasje i leddet som ble midlertidig avstivet.

Table of contents

Populærvitenskapelig sammendrag (norsk)	2
Acknowledgements.....	6
Abbreviations.....	8
List of papers.....	10
Summary	11
Introduction.....	13
Epidemiology.....	13
Anatomy and biomechanics of the Lisfranc joint.....	14
Mechanisms of injury.....	16
Diagnosis.....	16
Clinical signs.....	16
Imaging.....	17
Evaluation of stability.....	22
Classifications.....	23
Treatment.....	26
Nonoperative treatment.....	27
Operative treatment.....	28
Aim of thesis.....	32
Materials and Methods.....	33
Summary of results.....	39
Discussion.....	42

Methods.....	42
General discussion.....	46
Conclusions.....	52
Suggestion for future research.....	53
References.....	54
Appendix.....	
AOFAS midfoot score.....	
VAS pain score.....	
SF-36.....	
Paper 1.....	
Paper 2.....	
Paper 3.....	

Acknowledgements

The work presented in this thesis was carried out at Oslo University Hospital between 2011 and 2020. I am grateful for the financial support provided by Sophies Minde Foundation and Oslo University Hospital. This enabled this PhD project.

I would like to express my gratitude to the Division Orthopedic Surgery, Oslo University Hospital for the opportunity to conduct my research in an inspiring and stimulating environment. I am appreciative of the support from all my colleagues both with regard to research and every day clinical practice.

Throughout this project I have had the support and guidance of both Elisabeth Ellingsen Husebye, MD, PhD, and Kjetil Hvaal, MD, PhD. Elisabeth has been my main supervisor, always looking out for me and supporting me. Thank you for your hard work, quick responses and for keeping me on track. Kjetil has taught me just about everything I know in foot and ankle surgery and opened my eyes to this fascinating part of medicine. Thank you for always being supportive and always having time for a discussion.

During this project I have also had the pleasure to work with Professor Jan Erik Madsen. Thank you for your scientific guidance, insightful and always quick responses.

I would also like to thank my other co-authors:

Martine Enger, MD, it is a pleasure to work with you and your colleagues at Oslo A&E Department.

Helga Brøgger, MD, the support from you and the Department of Radiology and Nuclear Medicine has been crucial.

Throughout this period, I have worked alongside Marius Molund, MD, PhD and Fredrik Nilsen, MD, who have worked with their own PhD projects. We have become good friends and shared experiences and frustrations along the way.

To my good friend Gilbert Buru Moatshe, MD, PhD, who started orthopedic residency together with me in Haugesund. Thank you for your critical feedback on my manuscripts.

My parents, Anne and John Stødle, thanks for always being supportive and giving me all the opportunities in the world.

Finally, I would like to thank the most important people in my life; my wife and best friend Ida and our two children Erle and Eskil Syver. You are the best thing that ever happened to me!

Abbreviations

AOFAS	American Orthopaedic Foot and Ankle Society
AP	Antero-posterior
BP group	Bridge plating group
CI	Confidence interval
CT	Computerized tomography
C1-C2 distance	First cuneiform - second cuneiform distance
C1-M2 distance	First cuneiform - second metatarsal distance
IQR	Interquartile range
MOXFQ score	Manchester-Oxford Foot Questionnaire score
MRI	Magnetic resonance imaging
M1-M2	First metatarsal – second metatarsal distance
NRS	Numeric rating scale
OR	Odds ratio
OA	Osteoarthritis
ORIF	Open reduction and internal fixation
PA group	Primary arthrodesis group
RCT	Randomized controlled trial
SD	Standard deviation

SF-36	36 Item Short Form Health Survey
TMT	Tarsometatarsal
VAS	Visual analogue scale
VAS-FA	Visual analogue scale foot and ankle

List of papers

Paper 1

Lisfranc injuries: Incidence, mechanisms of injury and predictors of instability.

Stødle AH, Hvaal KH, Enger M, Brøgger H, Madsen JE, Ellingsen Husebye E. Foot Ankle Surg. 2020 Jul;26(5):535-540. doi: 10.1016/j.fas.2019.06.002. Epub 2019 Jun 20.PMID: 31257042

Paper 2

Outcome after nonoperative treatment of stable Lisfranc injuries.

A prospective cohort study.

Stødle AH, Hvaal KH, Brøgger HM, Madsen JE, Husebye EE. Foot
(Submitted)

Paper 3

***Temporary Bridge Plating vs Primary Arthrodesis of the First Tarsometatarsal Joint
in Lisfranc Injuries: Randomized Controlled Trial.***

Stødle AH, Hvaal KH, Brøgger HM, Madsen JE, Husebye EE. Foot Ankle Int. 2020 Aug;41(8):901-910. doi: 10.1177/1071100720925815. Epub 2020 Jun 5.PMID: 32501109

Summary

Lisfranc injuries are ligamentous and/or bony injuries to the midfoot. More advanced diagnostic methods as well as higher awareness of these injuries, seem to have increased the incidence compared to numbers previously reported. The Lisfranc injuries consist of a wide spectrum of injuries, ranging from severe fracture-dislocations to stable, non-displaced injuries. To differentiate the stable from the unstable injuries without obvious displacement can be challenging. Although there is agreement that the stable injuries should be treated nonoperatively, the literature reporting on the outcome after stable Lisfranc injuries is sparse. Unstable Lisfranc injuries should be treated with anatomic reduction and stable fixation. The best method of fixation, however, is debated.

The aims of the thesis were to obtain a better understanding of Lisfranc injuries by studying the incidence and mechanisms of injury, as well as the outcome following nonoperative treatment of stable injuries and operative treatment of unstable injuries.

The first study was a cross-sectional study focusing on the incidence and mechanisms of injury as well as trying to determine predictors of instability. The incidence of Lisfranc injuries detected was higher than previously reported. Low-energy trauma was the most common mechanism of injury. Factors that increased the chance of having an unstable injury were intraarticular fractures in the two lateral tarsometatarsal joints, female gender and a shorter second tarsometatarsal joint height. The second study was a prospective cohort study examining the outcome after nonoperatively treated stable Lisfranc injuries. An excellent

outcome was reported at the final follow-up at median 55 months after injury. The last study was a randomized controlled trial comparing arthrodesis to temporary bridge plate fixation of the first tarsometatarsal joint in unstable Lisfranc injuries. Both groups had good outcome scores and no method was found to be superior to the other with regard to the primary outcome score which was the American Orthopedic Foot and Ankle Score. The first metatarsal was better aligned in the group treated with temporary bridge plate fixation, but this group also had a high rate of radiologically detected osteoarthritis.

Introduction

Epidemiology

Injuries to the tarsometatarsal joint line is often referred to as Lisfranc injuries after the French surgeon and gynecologist Jacques Lisfranc de St. Martin (1787-1847), who described an amputation of the foot through the tarsometatarsal (TMT) joint line in 1815.¹

In modern literature Lisfranc injuries are not limited to the tarsometatarsal joint line, but also include the joints and ligamentous structures of the intercuneiform and naviculocuneiform joints.^{2,3} Lisfranc injuries are reported to be relatively uncommon and can lead to severe disability, especially if they are missed or treated inadequately.^{2,4-6} Hardcastle et al. reported an incidence of 1/60 000 person per year in 1982, previously Aitken et al. found 16 Lisfranc injuries during a 15 years period in their hospital treating 5500 fractures a year.^{7,8} English found that only 0.2 percent of the fractures treated at his hospital were Lisfranc injuries.⁹ These historical papers focus primarily on injuries with severe fracture-dislocations and not the more subtle Lisfranc injuries. Up to 24 % of the Lisfranc injuries are reported being missed on primary radiographs.¹⁰⁻¹² The increased availability of more advanced radiological modalities such as magnetic resonance imaging (MRI), computerized tomography (CT) scans, weightbearing radiographs and stress fluoroscopy in recent years, as well as an increased awareness of Lisfranc injuries, is probably the reason why the incidence of these injuries seems to be higher than previously reported.^{3,13-18} In a recent study by Ponkilainen et al., they reported an

incidence of Lisfranc injuries of 9.2/100 000 person-years when retrospectively looking at all CT scans obtained after foot injuries during a 5 years period.¹⁹

Anatomy and biomechanics of the Lisfranc joint

The anatomy of the foot is complex as there are 26 bones and 33 joints. The foot should be able to cushion the body, adapt to the surface, transfer the load from the body through the ankle and foot and provide a leverage for propulsion.²⁰ To be able to do this the joints in the foot have different functions, some joints have a greater range of motion thereby making the foot able to adapt to the surface, while other joints have a more restricted motion to give the acquired stability during propulsion and thereby make the load transfer possible.^{21–23} In addition, the flexibility of the foot change during the gait cycle, as the hindfoot inverts the transverse joints become divergent making the foot more rigid.²⁰ In the TMT joint line the medial three joints have little motion and provides stability of the medial and middle part of the foot during gait.²⁴ The lateral two TMT joints on the other hand have more inherent motion and play an important role in the foots ability to adapt to the surface.^{20,24} A fusion of these two joints have been shown to give a less favorable result.²⁵ Therefore, it seems that in the central and medial part of the midfoot stability is most important, while in the lateral part of the midfoot motion is more important than rigidity. Posttraumatic osteoarthritis (OA) is also better tolerated in the lateral two TMT joints than in the two central TMT joints where symptomatic posttraumatic osteoarthritis has been shown to be more common.^{26,27}

The primary stability of the midfoot is provided by both osseous and ligamentous structures. The cuneiforms and metatarsals are wedged shaped and form a transverse arch in the midfoot. In addition, the second metatarsal is recessed proximal in the mortise between the medial and lateral cuneiform, forming a “keystone”.²⁸ Strong ligamentous structures that transverse the TMT joints, especially the Lisfranc ligament which run from the medial cuneiform to the base of the second metatarsal, play an important part in stabilizing the midfoot. The Lisfranc ligament consist of three parts; the interosseous ligament running from the medial cuneiform to the base of the second metatarsal, which is the strongest, the plantar ligament running from the medial cuneiform to the plantar aspect of the second and third metatarsal bases and finally, the dorsal ligament running from the dorsolateral aspect of the medial cuneiform to dorsomedial on the base of the second metatarsal, which is the weakest.^{29–31} There are intermetatarsal ligaments between the base of the four lateral metatarsals, but no ligament between the base of the first and second metatarsal base, making the Lisfranc ligament the primary stabilizer between the first and second column.³¹ Injuries to these structures can cause instability and may lead to displacement of the TMT joints.^{16,29,32–34} Kaar et al. showed in their biomechanical study that sectioning of both the interosseous (Lisfranc) ligament between the medial cuneiform and base of the second metatarsal, and plantar ligament between the medial cuneiform and base of the second and third metatarsals, were required to produce transverse instability.³⁴ Instability was defined as 2 mm or more displacement between the base of the second metatarsal and the medial cuneiform and between the middle cuneiform and the second metatarsal. To produce longitudinal instability i.e., widening of the space between the

medial and middle cuneiform, the interosseous (Lisfranc) ligament and the interosseus ligament between the medial and middle cuneiform had to be sectioned.

Secondary stabilizers such as the plantar fascia, tendons (tibialis anterior and posterior, and peroneus longus) and intrinsic muscles, also contribute to the stability of the Lisfranc joint complex.³

Mechanisms of injury

The mechanisms of injury are often divided into high-energy or low-energy injuries. The high-energy injuries are seen with mechanisms of injury such as motor vehicle accidents, fall from heights or crush injuries and have previously been shown to account for the majority of the injuries.^{2,7,14,35–38} In more recent studies, however, high-energy injuries only accounted for 36.5–40 % of the cases.^{19,39} The low-energy injuries can occur during athletic activities or just by stumbling or falling during walking.^{19,39} These injuries are often caused by an axial or rotational force on a fixed foot in equinus (indirect trauma) causing disruption of the ligaments and/or fractures in the TMT region.^{2,39,40} The lateral TMT joints have been reported to be less frequently involved in low-energy injuries.³⁹

Diagnosis

Clinical signs

Lisfranc injuries can be challenging to detect, especially the subtle injuries and injuries in polytrauma patients can easily be missed. Swelling and pain in the midfoot are common clinical findings together with inability to weightbear. Palpation of the midfoot and manipulation of the

midfoot using adduction/pronation usually induces pain.⁴¹ If a plantar ecchymosis is present one should have a high suspicion of Lisfranc injury.⁴²

Imaging

Most patients with a suspected injury to their midfoot will have an initial non-weightbearing radiograph of their injured foot. Up to 24 % of the injuries have been reported to be missed on the plain radiographs.^{10,12,19} The severe Lisfranc fracture-dislocations can easily be detected on plain radiographs, but the minimally displaced injuries can be challenging or almost impossible to detect. To detect Lisfranc injuries on radiographs one should be familiar with the normal findings on radiographs of the midfoot. The medial boarder of the middle cuneiform should line up with the medial boarder of the second metatarsal on the AP view. On the oblique view the lateral boarder of the lateral cuneiform should line up with the lateral boarder of the third metatarsal, while the medial boarder of the cuboid lines up with the medial boarder of the fourth metatarsal. On the lateral view there should be no step-off at the TMT joints. ^{43,44}

Figure 1: Normal AP, oblique and lateral radiograph of right foot. a) The medial boarder of the middle cuneiform should line up with the medial boarder of the second metatarsal base on the AP radiograph. b) The medial boarder of the cuboid should line up with the medial boarder of the fourth metatarsal base on the oblique radiograph. c) There should be no step-off in the tarsometatarsal joints on the lateral radiograph.

When evaluating displacement in the more subtle Lisfranc injuries, three main distances have been reported measured; the distance between the base of the first and second metatarsal (M1-M2), the distance between the medial cuneiform and the base of the second metatarsal (C1-M2), and finally the distance between the medial and middle cuneiform (C1-C2). Even though the M1-M2 distance is most commonly used there is no consensus with regards to a definition of this distance.⁴⁵ The average M1-M2 distance has in non-injured feet been reported to be from 1.3 mm to 2.5 mm in non-weightbearing AP radiographs and up to 2.9 mm in weightbearing radiographs, although a distance up to 5.0 mm can be seen in normal feet. As the distance vary between individuals weightbearing radiographs of the contralateral side should be obtained for comparison, the difference should be less than 2 mm.^{6,16,41,45-48}

The C1-M2 distance is the interval where the interosseous Lisfranc ligaments is located as it originates from the lateral aspect of the medial cuneiform and inserts on the lateral and plantar aspect of the second metatarsal. On weightbearing AP radiographs the C1-M2 distance in normal feet has been reported to be up to 3.5 mm and compared to the contralateral side the difference should be less than 2 mm in stable Lisfranc injuries.^{46,49} To detect intercuneiform instability the C1-C2 distance should be compared to the contralateral side, an increased distance of 2 mm compared to the non-injured side is considered an abnormal finding.⁴⁶

Figure 2: Lisfranc injury to the left foot. An increased distance in the C1-M2 interval (red line) is revealed on weightbearing radiograph compared to the non-injured right side.

Although the distances mentioned above are commonly used to evaluate the stability of Lisfranc injuries, good weightbearing radiographs can be difficult to obtain due to the complex anatomic configuration of the midfoot. Overlapping of the tarsal and metatarsal bones can make the interpretation of the radiographs challenging.^{22,43,45,50} An avulsion fracture of the origin of the Lisfranc ligament from the lateral side of the medial cuneiform or from the

insertion on the medial aspect of the second metatarsal base (“fleck sign”) is indicative of a Lisfranc injury.³ Contrary to previous belief that the “fleck sign” was a sign of instability, it has not been shown to increase the risk of subsequent displacement in nonoperative treatment of minimally displaced Lisfranc injuries.^{51,52}

In a cadaver study by Lu et al. evaluating the ability of plain radiographs and CT scans to detect sequential displacement of the second metatarsal, none of the 1 mm dorsolateral displacements were visualized on radiographs and two thirds of the 2 mm displacements were not detected on radiographs, while all of the 1 and 2 mm displacements were detected on the CT scans. Both radiographs and CT scans were able to detect the 3 mm displacements. Detecting displacement in the TMT joints of less than 2-3 mm therefore seems inadequate using (standard/plain) radiographs.^{17,22} Ebraheim and Lu also showed in their study that a 3 mm dorsolateral displacement of the second metatarsal in relation to the middle cuneiform reduced the contact area by 38.6 %, while a 15 % reduction was seen where the second metatarsal was displaced 1 mm.⁵³ Minor displacements, not detected on radiographs, can thereby significantly reduce the contact area in the TMT joints.

Due to the limited ability of radiographs to detect minor joint displacements in the midfoot there is an increasing use of CT scans in diagnosing Lisfranc injuries. CT scans are useful in detecting subtle Lisfranc injuries as well as in differentiating between different fracture patterns and planning the surgery.^{11,14,17,54} Conventional CT scans have the drawback of being non-dynamic examinations and therefore cannot show displacement that only occurs during

weightbearing. As weightbearing CT scans become more available they are expected to make the detection of occult instability in Lisfranc injuries less challenging.⁵⁵

Evaluation of stability

To detect instability in the subtle, non-displaced Lisfranc injuries, both weightbearing radiographs and stress-tests under fluoroscopy have been advocated.^{5,16,33,36,40,45,56} Comparing the injured foot to the contralateral side is essential to detect subtle displacements.^{16,40,44} These examinations can be painful for the patients in the acute setting, but 7 to 14 days after injury most patient will tolerate testing of the stability.⁵⁶ The proponents for the weightbearing radiographs argue that weightbearing radiographs are more objective and less examiner dependent than the stress-test under fluoroscopy, while the proponents for the stress-test argue that many patients are not able to weightbear adequately enough to detect occult instability and the stress-test where the examiner perform the abduction and pronation of the foot are more likely to detect instability.^{4,13,16,33,34,41,45,48,57} Coss et al. and Kaar et al. have showed in a cadaveric studies that stress radiographs were better in detecting a transected Lisfranc ligament than simulated weightbearing.^{33,34} In both weightbearing radiographs and stress-tests the injured foot should be compared to the non-injured foot.^{16,28,40} If a patient is not able to weightbear and stress-test under fluoroscopy is not possible due to pain 7-14 days after the injury, a stress-test while the patient is under anesthesia can be performed. One of the challenges using the stress-test under fluoroscopy, is lack of standardization and subjectiveness of the examination.⁵⁷ Also there is no agreement of a threshold value for when the test is positive.

In a retrospective study on 21 subtle Lisfranc injuries by Raikin et al., MRI was shown to classify 90 % of subtle Lisfranc injuries correctly into stable or unstable injuries.¹³ They found that a rupture of the plantar ligament running from the medial cuneiform to the base of the second and third metatarsal had a sensitivity of 94 % and a specificity of 75 % and was the strongest predictor of instability in acute Lisfranc injuries. They used manual stress radiographs with the patient under anesthesia and intraoperative findings as their reference standard. Kitsukawa et al. showed in their study that in most cases where the plantar ligament was ruptured, the interosseous and dorsal ligament were ruptured as well.⁵⁸

Both conventional CT scans and MRIs are non-dynamic examinations, hence displacement occurring only during weightbearing will not be detected. Weightbearing CT scans are becoming more available and in the future this will most likely be the gold standard for detecting instability in the subtle Lisfranc injuries.^{55,59} With regard to follow-up, patients with injuries evaluated as stable should be followed up with weightbearing radiographs to detect any missed instability.^{41,51,56}

Classifications

In the literature at least fifteen different classification systems have been proposed for classifying Lisfranc injuries. They vary between focusing on the trauma mechanism, direction of displacement of the TMT joints or involved columns.⁶⁰

In 1909 Quènu and Küss first described injuries to the TMT joints based on the direction of metatarsal displacement, the classification has been modified by Hardcastle and later by Myerson.^{2,7} These classifications are based on joint incongruity and direction of dislocation and

are widely used by authors when describing Lisfranc fracture-dislocations, but are of limited use when it comes to guiding treatment and establishing prognosis, nor can they be used to classify the more subtle Lisfranc injuries without obvious displacement on plain radiographs.^{15,16,37,39,56}

In 2001 Chiodo and Myerson presented a classification for Lisfranc fracture-dislocations based on the columnar theory which focus on the three mechanical columns of the midfoot.⁵⁶ The medial column includes the first TMT and medial naviculocuneiform joint and allows about 3.5 mm of motion in the sagittal plane.²⁴ The middle column is rigid and only allow minimal of sagittal motion. This column includes the second and third TMT joint as well as the articulations between the middle and lateral cuneiform bones and the navicular bone. At last the lateral column includes the fourth and fifth TMT joints, these joints have considerably more motion, approximately 13 mm in the sagittal plane.^{3,24} This classification can potentially help plan the treatment as each column is thought to function as one unit and also probably has a better prognostic value as subtle joint incongruity seems to be better tolerated in the lateral column than in the middle column.²⁶ Lau et al. also showed in their study an association between the functional outcome and the number of columns injured.³⁷

Figure 3: The functional columns of the foot. The medial column (cyan) consists of the first naviculocuneiform and first tarsometatarsal joint. The middle column (green) consists of the second and third naviculocuneiform joints as well as the second and third tarsometatarsal joints. The lateral column (purple) consists of the fourth and fifth tarsometatarsal joints.

For the subtle Lisfranc injuries, Nunley and Vertullo in 2002 published a classification based on weightbearing radiographs and bone scintigrams.¹⁶ Weightbearing radiographs of the injured foot were compared to the non-injured side. A 2 mm increased diastasis in the interval between the base of the first and second metatarsal (M1-M2) compared to the non-injured side, was considered diagnostic for a displaced Lisfranc injury. If the weightbearing radiographs did not show any increased diastasis, a bone scintigram was obtained to confirm the diagnosis of a stable Lisfranc injury. Grade 1 injury was an injury without any displacement and positive

bone scintigram. Grade 2 injury had an increased diastasis of 2-5 mm between the first and second metatarsal bases, but no loss of arch height on the lateral weightbearing radiograph. Grade 3 had increased diastasis of 2-5 mm and loss of arch height. The grade 1 injuries, the stable midfoot injuries, were recommended treated nonoperatively, while the unstable grade 2 and 3 injuries were recommended treated operatively.

Schepers and Rammelt published a new classification system in 2018 based on the column theory of the midfoot, but did not base the classification on displacement of the TMT joints, thereby enabling classification of the more subtle, nondisplaced injuries as well.⁶⁰ They focus on the number of involved columns (medial, central and lateral) and combine this with the severity of joint involvement: no involvement (including extraarticular fractures), ligamentous injury (with or without avulsion fractures), simple intraarticular fractures or comminuted fractures of > 50 % of the joint surface. To classify the injuries and decide treatment strategy the authors recommend using the diagnostic tools needed (conventional radiographs, weightbearing radiographs, stress images, CT scans and/or MRI). Although this new classification system seems promising, the inter- and intra-observer reliability and correlation to outcome has yet to be determined.

Treatment

Lisfranc injuries consists of a wide spectrum of injuries ranging from stable, non-displaced injuries to severe fracture dislocations with soft tissue compromise. Historically, the main focus has been on the treatment of the fracture-dislocations, but during the last decades the subtle

injuries have gained attention. There still is no classification system proven good enough to properly guide the treatment and predicting the final outcome.^{60,61} It seems clear that due to the wide variety in these injuries they cannot all be treated in the same manner.⁶² The goal of the treatment should be to achieve a pain free, plantigrade and stable foot, preferably without development of post traumatic OA and the need for subsequent surgery. Regarding the treatment there are controversies to which fixation technique to use or when to perform a primary arthrodesis.^{18,62-64}

Nonoperative treatment

The stable, nondisplaced Lisfranc injuries are recommended treated nonoperatively, but only few studies report the outcome of nonoperative treatment after having verified the stability.^{4,16,51,65} The nonoperative treatment in these studies vary between weightbearing as tolerated in a below-knee-orthosis to immobilization in a below-the-knee cast for 6 weeks.^{4,16,51,65} Nunley and Vertullo reported excellent results in all (7) patients with a stable Lisfranc injury treated nonoperatively in their study.¹⁶ In their study looking at the rate of secondary displacement after conservative treatment of minimally displaced Lisfranc injuries, Chen et al. reported a mean American Orthopaedic Foot and Ankle Society (AOFAS) midfoot score of 78.0 (68.6-87.4) and a Manchester-Oxford Foot Questionnaire (MOXFQ) score of 24.8 (11.1-38.5) in the 12 patients where the injury remained stable at a mean follow-up of 54 months.⁵¹ The stability of the injuries were evaluated using weightbearing radiographs. Interestingly they reported the median time to detection of displacement to be 18 (range 2-141) days. Crates et al. also looked at the outcome of nonoperative treatment of subtle Lisfranc

injuries where the stability were evaluated by weightbearing radiographs without further specifying the timing of these.⁶⁵ Of the 36 patients included in the study, only 16 were successfully treated nonoperatively, while 20 required surgery after failed nonoperative treatment. The failure of conservative treatment was based on the judgement of the senior author and no subsequent radiographical findings are mentioned, thus the results of the study are difficult to interpret.

Ponkilainen et al. reported on nonoperative treatment of non-displaced Lisfranc injuries determined by CT scans.⁶⁶ The patients were evaluated at a median of 3.8 years after injury, 65% of the patients scored more than 80 points in the pain and function subscales of the visual analogue scale foot and ankle (VAS-FA). 78 % of the patients scored above 80 points overall. Although the results in this study were good, 7 % of the patients scored under 60 points overall. The patients were not tested for any occult instability which might have led to a poorer result in that subgroup of the patients.

Operative treatment

In Lisfranc injuries with instability or displacement operative treatment is recommended.^{4,15,16,41,48} Several studies have showed that the most important factors in achieving a good outcome is anatomic reduction and stable fixation.^{2,35,36,67,68} Anatomic reduction is most reliably achieved by open reduction.^{18,35,36,69} In the patients presenting with severe fracture-dislocations and compromised soft tissue, the dislocated joints should be reduced and temporary fixation can be achieved either using an external fixator or percutaneous pin fixation.⁷⁰

Closed reduction and cast immobilization of dislocated Lisfranc injuries have shown a high rate of unsatisfactory midfoot alignment at follow-up.^{2,71} Although a few older studies reports excellent functional outcome despite persistent joint displacement in patients treated with closed reduction and cast immobilization, maintenance of anatomic reduction is now widely recognized as one of the most important factors in achieving good functional outcome after Lisfranc injuries. Persistent displacement of the TMT joints tend to lead to a poor functional outcome.^{2,8,71,72} Fixation of the TMT joints using Kirschner wires improve the radiological and functional outcome after displaced Lisfranc injuries, compared to closed reduction and cast immobilization.^{2,71} Although Kirschner wire fixation improved the outcome after displaced Lisfranc injuries, a higher rate of redisplacement occur when using Kirschner wires compared to screw fixation.^{36,73} Transarticular screw fixation has in a biomechanical study been show to provide better stability of the TMT joints.⁷⁴

Since the introduction of transarticular screw fixation by Arntz et al. this treatment has been the gold standard in treating unstable Lisfranc injuries.^{18,36,63} This fixation technique is used in the medial and central part of the midfoot. When the lateral TMT joints are involved most authors recommend treating these joints with Kirschner wire fixation as these joints are more mobile than the central and medial joints.¹⁸ Good outcome scores and a low rate of redisplacement has been reported when using transarticular screw fixation of Lisfranc injuries.^{25,35,36,63,75}

While the Kirschner wires transfixing the lateral TMT joints usually are removed within 6-8 weeks, there seems to be no consensus with regard to removal of the screws as some authors

recommend removal after 4 months while others leave the hardware in permanently if the patients has no complaints.^{15,41,75}

Despite the good outcome scores reported after reduction and fixation of unstable Lisfranc injuries there is a high rate of posttraumatic OA.^{18,25,35,76} Whether this is mainly due to the primary trauma or part of this can be contributed to fixing the joints using transarticular screws is unknown. In recent years there has been a trend towards using a bridge plating technique in treating Lisfranc injuries aiming at inflicting less damage to the joint surface. This technique avoids passing screws through the joint, possibly reducing the incidence of posttraumatic OA. Alberta et al. showed that a 3.5 mm screw cause a destruction of 3.6 % of the joint surface when passed through the second TMT joint.⁷⁷ They also reported a similar ability of dorsal plates and transarticular screws to resist TMT joint displacement during weightbearing load. When using the bridge plating technique most authors recommend removing the plates after 4-6 months.^{67,78-81} Kirzner et al. compared the outcome of transarticular screw fixation to the bridge plating technique in a retrospective study and reported an significant better MOXFQ score and AOFAS score in the plate group (MOXFQ: 38.1 versus 25.6, $p<.001$ and AOFAS:71.0 versus 82.5, $p<.001$).⁷⁸ This was also confirmed by Hu et al. in their prospective comparative study reporting an AOFAS score of 83.1 in the bridge plate group and 78.5 in the transarticular screw group ($p<.01$).⁸⁰ A reduction of posttraumatic OA when using the bridge plating technique has not been show.

Although open reduction and internal fixation has been proven to yield good results in Lisfranc injuries, Kuo et al. reported poorer results in the subgroup of patients suffering from purely ligamentous Lisfranc injuries, despite adequate reduction and screw fixation.³⁵ This led to the

randomized controlled trial by Ly and Coetzee comparing primary arthrodesis of the medial two or three TMT joints to open reduction and internal fixation (ORIF) using transarticular screw fixation in the primary ligamentous Lisfranc injuries.⁶³ At 2-years follow-up they reported a significant better AOFAS score in the primary arthrodesis group compared to the group treated with ORIF (mean AOFAS 88.0 versus 68.6, $p<.005$). The good outcome scores using primary arthrodesis reported by Ly and Coetzee have also been confirmed by other studies comparing primary arthrodesis to ORIF.^{25,75,82}

Rammelt et al. compared primary ORIF using Kirschner wires or screws to secondary corrective arthrodesis and showed a significant better AOFAS score in the primary ORIF group (81.4 vs 71.8, $p=.031$), implying that treating these injuries in the acute setting is preferable to addressing any sequelae presenting at a later time.⁶⁹

Aim of thesis

The aim of this Ph.D. thesis was to obtain a better understanding of Lisfranc injuries by studying the incidence and mechanism of injury, as well as the outcome after nonoperative treatment of the stable injuries and operative treatment of the unstable injuries.

Specific aims of the papers

Paper 1: What is the incidence of Lisfranc injuries and which mechanisms of injury are the most common? Are there any radiological predictors of instability?

Paper 2: What is the outcome after stable Lisfranc injuries?

Paper 3: Is extraarticular bridge plating of an unstable Lisfranc injury better than primary arthrodesis?

Materials and Methods

Paper 1:

Lisfranc injuries: Incidence, mechanisms of injury and predictors of instability

Design: Cross-sectional study

Materials: 89 patients presenting with Lisfranc at Oslo University Hospital and Oslo Accident and Emergency Department were registered during a 1-year period. 84 patients were included in the study.

Methods: Lisfranc injuries were defined as injury to the TMT joints with avulsion fractures, intra-articular fractures and/or displacement in the TMT joints. The diagnosis was verified radiologically in all patients. Demographic data and mechanisms of injury were registered. In the non-displaced injuries stability was tested using a stress test under fluoroscopy or if this was undecisive weightbearing radiographs were obtained. The injury patterns were assessed on radiographs and CT scans. A treatment algorithm was used (Figure 4). The CT scans were also used to measure the mortise depth and the height of the second TMT joint, these findings were correlated to the stability of the injuries (Figure 5).

Figure 4: Treatment algorithm for Lisfranc injuries.

Figure 5: CT scans showing method for measuring second tarsometatarsal joint (TMT-2) height (a) and Lisfranc mortise depth (b). The red line in figure a indicate the plane for evaluating the Lisfranc mortise depth. In figure b the medial mortise depth is marked A and the lateral mortise depth is marked B.

Statistical analysis: Normal distributed data were presented using mean and standard deviation. The independent sample t-test was used to compare the means between groups. For categorical variables the odds ratio (OR) and Pearson Chi-square test were used. Logistic regression was used to assess the relation between the stability and fracture pattern, as well as the relation between stability and Lisfranc mortise depth. As the radiographs and CT scans were evaluated by three of the authors, the interrater reliability was also calculated using the interclass correlation coefficient.

Ethical considerations:

Prior to inclusion the Regional Ethics Committee approved the study (2014/853/REK) and the patients signed an informed consent form.

Paper 2:

Outcome after nonoperative treatment of stable Lisfranc injuries.

Design: Prospective cohort study

Materials: 26 patients with stable Lisfranc injuries were included. The patients were also included in study 1.

Methods: All the patients had nondisplaced Lisfranc injuries verified with CT scans. The stability was confirmed with a stress test under fluoroscopy. Weightbearing radiographs of both feet was obtained at follow-ups up to 1 year.

Outcome parameters: The main outcome score was the AOFAS midfoot score and the secondary outcome scores were the Visual Analog Scale (VAS) for pain at rest and during

walking, 36-Item Short Form Health Survey (SF-36) and radiological OA on follow-up radiographs.

The AOFAS score consist of three main categories (pain, function and alignment).⁸³ The best score being 100 and the worst possible score 0 (Appendix 1).

To evaluate the pain experienced by the patients the VAS pain scale was used. The patients marked the experienced pain at rest and during walking on a scale for 0 (no pain) to 10 (worst possible pain, Appendix 2). The SF-36 was used to evaluate the patient's health and well-being. The survey consists of 36 items which again contributes to eight health domains (physical functioning, role limitations due to physical problems, bodily pain, general health, vitality, social functioning, role limitations due to emotional problems and mental health). From these eight domains two summary scores can be calculated (physical component summary and mental component summary, Appendix 3) ^{84,85} To verify the stability and detect any development of OA, radiographs were obtained at follow-ups up to one year after trauma.

Statistical analyses: Normally distributed data were presented as means and standard deviations (SD). For statistical analyses the one sample t-test was used. Non-parametric data were presented as median and interquartile range (IQR), for statistical analysis the Wilcoxon signed rank test was used.

Ethical considerations:

The study was approved by the Regional Ethic Committee (2014/849/REK) and the patients signed an informed consent form prior to inclusion.

Paper 3:

Temporary bridge plating vs primary arthrodesis of the first tarsometatarsal joint in Lisfranc injuries.

Design: Randomized controlled trial (superiority study).

Materials: 48 patients with Lisfranc injuries were included between October 2011 and August 2015. Instability of the medial 3 TMT joints were confirmed radiologically.

Figure 6: CONSORT flowchart of the trial enrollment and analysis.

Methods: Demographic data and mechanism of injury was registered. Preoperative CT scans were obtained in all patients. The patients were randomized to either primary arthrodesis (PA group = 24 patients) or temporary bridge plate fixation (BP group = 24 patients) of the first TMT joint. The instability of the second and third TMT joint was treated with primary arthrodesis in

all patients. A dual incision technique using dorsomedial and dorsolateral incision was used and anatomic reduction was sought before final fixation. The patients were immobilized in a cast 6-8 weeks postoperative and then started weightbearing as tolerated in a Walker boot which was discontinued at 12 weeks. The bridge plate over the first TMT joint in the BP group was removed after 4-5 months. Postoperative CT scans and radiographs, as well as weightbearing radiographs at 6-8 weeks, 3, 12 and 24 months were obtained.

Figure 7: The two surgical techniques used in the study. A) primary arthrodesis of the three medial TMT joints. B) Temporary bridge plating of the first TMT joint and primary arthrodesis of the second and third TMT joint. In this patient the arthrodesis of the third TMT joint was fixated using a plate due to comminution of the joint. C) Radiograph after removal of the temporary bridge plate in b).

Outcome parameters: The main outcome parameter was the AOFAS midfoot score, while the secondary outcome scores were the VAS pain score, SF-36, pedobarography and complications. In addition, postoperative reduction was evaluated on CT scans. At the 2-years follow-up the

alignment of the foot was assessed on weightbearing radiographs as well as secondary displacement and development of posttraumatic OA was registered.

Statistical analysis: A power analysis for a superiority study was conducted prior to the inclusion of patients. Normally distributed data were presents as means and standard deviations. The t-test for independent samples was used to detect statistical significance. Non-parametric data were presented as medians and IQR, these data were analyzed using the Mann-Whitney U test. Categorical data were analyzed using the Fischer exact test. To calculate correlation between variables the Spearman rank correlation was used.

Ethical considerations: The study was approved by the Regional Ethics Committee (2011/1582/REK) and the hospitals data protection officer. All participants sign an informed consent form prior to inclusion. The study was registered in Clinical.Trials.gov (ID: NCT01448941).

Summary of results

Paper 1:

Lisfranc injuries: Incidence, mechanisms of injury and predictors of instability

In this study we found the incidence of Lisfranc injuries in the catchment area of Oslo University Hospital to be 14/100 000 person-years. The incidence of stable injuries was higher than of the unstable injuries (8/100 000 versus 6/100 000 person-years). Unstable injuries were more common in women than in men (OR = 3.1 95% CI= 1.3-7.6, p=.016). The single most common mechanism of injury was fall from own height/ twisting injury of the foot, accounting for 31%. Only 31% of the injuries were caused by high-energy trauma, while 20% were sports related

injuries. 28% of the primary non-weightbearing radiographs were unable to detect any injury. Joint displacement, as a sign of instability, was most likely to be detected on CT scans (17/84) or on stress-test under fluoroscopy (14/67). With regard to the fracture pattern evaluated on the CT scans, only intraarticular fractures of the fourth and/or fifth TMT joint was related to an increased chance of instability (OR = 6.0, 95% CI = 1.6 – 21.5). The stable Lisfranc injuries had a deeper second TMT joint than the unstable ones ($p=.04$). Men also had a deeper second TMT joint compared to women ($p=.001$).

Paper 2:

Outcome after nonoperative treatment of stable Lisfranc injuries.

In this study we prospectively examined the outcome after nondisplaced, stable Lisfranc injuries in 26 patients. The patients included in this study were the patients with stable Lisfranc injuries from paper 1 that were compliant to the conservative treatment and available for follow-up at Oslo University Hospital. The median follow-up was 55 (IQR 53 -60) months. All injuries remained stable on follow-up radiographs up to 1 year. The median AOFAS midfoot score was 89 (IQR 84 – 97) at one year and improved further to 100 (IQR 90 – 100) at the final follow-up ($p= .005$), consistent with an excellent outcome. This was mirrored by a VAS score of 0 (IQR (0 – 0) at the final follow-up. The SF-36 score at 1 year was equivalent to the score of the Norwegian general population, indicating a return to the same quality of life as the general population.

Radiological OA was only detected in 1 patient at the 1-year follow-up, no patients had clinical signs of OA. All patients included in this study had intraarticular fractures or avulsion fractures

in the TMT joint line on CT scans, but as many as 9/26 injuries was not detected on initial non-weightbearing radiographs.

Paper 3:

Temporary bridge plating vs primary arthrodesis of the first tarsometatarsal joint in Lisfranc injuries.

In this randomized controlled study (RCT) were not able to detect any statistically significant difference between the PA and BP groups at 2-years follow-up with regard to mean AOFAS midfoot score, 89 (SD 9) versus 85 (SD 15), median VAS pain score 0 (IQR 0 – 2.0) versus 0.9 (IQR 0-3.0) or in the different subgroups of the SF-36. The Meary's angle (talus-first metatarsal angle) in the AP weightbearing radiographs at 2-years follow-up showed better alignment in the BP group compared to the PA group ($p = .04$). The pedobarography showed a reduced peak pressure under the fifth metatarsal in the PA group ($p = .047$). The reduced fifth metatarsal peak pressure in the PA group was correlated to the lateral Meary's (talus-first metatarsal) angle ($r_s = -0.65$, $p = .002$). At the 2-years follow-up 11/24 patients in the BP group had radiological signs of OA on the radiographs. We registered more complications in the BP group than in the PA group, but this finding was not statistically significant.

Discussion

Methods

Paper 1

To determine the incidence of Lisfranc injuries in the Oslo University Hospital catchment area we registered all Lisfranc injuries presenting at our institution during a 1-year period (1 September 2014 to 31 August 2015) and only the patients living in the catchment area were included in the incidence calculation (Figure 8). Any patient presenting at other institutions or not seeking medical treatment would thereby not have been detected. This could potentially underestimate the incidence.

Figure 8: Patients presenting with Lisfranc injuries at Oslo University hospital during a 1-year period. * one patient with bilateral unstable Lisfranc injuries.

All injuries were evaluated using CT scans (except one who had an MRI), confirming injuries to the Lisfranc joint complex. Any pure ligamentous sprains without avulsion fractures or displacement could have been missed. A larger patient population would of course have increased the strength of the paper.

The mechanisms of injury were registered at presentation thereby reducing any recall bias. The mechanisms of injury are often classified as high- or low-energy injuries. We defined low-energy injuries as fall from own height, twisting injury of the foot, bike accidents, falling downstairs and kicking into an object. High-energy injuries included motor vehicle accidents, fall > 3 meters and crush injuries. Sports related injuries were registered as a separate category.

To differentiate between the stable and unstable nondisplaced injuries we examined these feet with stress-test under fluoroscopy 7-14 days after the trauma. There is no consensus regarding which method is the best in detecting instability of Lisfranc injuries, some authors prefer testing the stability using stress test under fluoroscopy, while others advocate obtaining weightbearing radiographs.^{4-6,13,16,33,34,36,40,45,56,86} As weightbearing radiographs were not obtained in most of the nondisplaced injuries, we were not able to compare the ability of the two tests to detected instability.

Several classifications have been proposed for Lisfranc injuries, most of these have limitations when it comes to guiding treatment and predicting the outcome. We chose to categorize the fracture patterns using the columnar theory (medial, middle and lateral column) as proposed by

Chiodo and Myerson, and later by Schepers and Rammelt, in addition to differentiating the injuries into stable and unstable injuries.^{56,60}

Paper 2

To evaluate the outcome after stable Lisfranc injuries in paper 2, we included patients with stable Lisfranc injuries from paper 1 that were available for follow-up at Oslo University Hospital and were compliant with the conservative treatment protocol. To ensure the stability of the injury, weightbearing radiographs were obtained at follow-ups up to 1 year as Chen et al. showed that subsequent displacement can be detected on weightbearing radiographs even after 12 weeks.⁵¹ The main outcome score in both paper 2 and 3 was the AOFAS midfoot score as this is the score most commonly used in evaluating the outcome after Lisfranc injuries.⁸⁷ The AOFAS score does, however, have several drawbacks. It combines patient reported items with physical examination, which can lead to inter-observer bias. Further, the AOFAS score cannot be calculated if an answer is missing, few response categories are available for each item, and several responses are formulated in a non-specific way. The score has a ceiling effect and consequently a limited ability to differentiate between patients with a good outcome scores, which was the case in study 2.⁸⁸ Ponkilainen et al. did, on the other hand, show that the AOFAS midfoot score does have a high convergent validity and an acceptable internal consistency.⁸⁸ The developers of the AOFAS scoring system have recently discouraged the use of the AOFAS score as the main outcome measure.⁸⁹ In addition to the AOFAS midfoot score the SF-36 was used, which is an extensively validated patient-reported survey of patient health.

At the final follow-up (median 55 months), 8 of the patients were only available for phone interview. The AOFAS score was thereby conducted by phone and the patients reported the physical examination items themselves. This may have induced bias to this part of the AOFAS score, although the patients reported an excellent functional outcome and no gait abnormalities or malalignment were detected on previous follow-ups. The VAS pain score also poses a problem when a phone interview is conducted, therefore the numeric rating scale (NRS) for pain was used as the NRS pain score is highly correlated to the VAS pain score.⁹⁰

Paper 3

This paper is a RCT conducted as a superiority trial aiming to show that the joint sparing procedure using temporary bridge plating technique was superior to primary arthrodesis of the first TMT joint in Lisfranc injuries. The decision to only compare the two different treatments in the first TMT joint and not all 3 medial TMT joints was based on the knowledge that the medial TMT joint is more mobile than the 2 central TMT joints, hence preserving this joint might therefore be more beneficial than preserving the central ones.²⁴ Painful OA is also more common in the second and third TMT joint than in the medial and lateral columns after Lisfranc injuries.²⁶

The AOFAS midfoot score was also in this study the main outcome parameter, as it has been the score most widely used when evaluating Lisfranc injuries. The AOFAS score has in recent years been criticized for having several flaws, as mentioned above, and is now longer recommended being used as a primary outcome measure.⁸⁸ In future studies we recommend

using other validated foot and ankle specific patient reported outcome measures as the primary outcome parameter.

When conducting an RCT the preferred method to reduce detection bias would be blinding of both the patients and the examiner at the follow-ups. As the patients in the BP group needed an additional surgery to remove the plate, patient-blinding was not feasible. Blinding of the examiner at the follow-ups was not done.

Several papers have confirmed that an anatomical reduction and stable fixation are two most important factors in obtaining a favorable result after Lisfranc injuries and Ebraheim et al. showed that displacement less than 3 mm in the TMT joint line was often not detected on radiographs, but easily detected on CT scans.^{2,35,36,53,67,68} To evaluate the postoperative reduction we therefore obtained postoperative CT scans of all patients.

Concern has been raised regarding the radiation exposure caused by CT scans. A CT scans of the ankle and foot exposes the patient to an effective dose of radiation of 0.07 mSv which is equivalent to 0.88 chest radiograph.⁹¹

General discussion

Paper 1

The incidence of 14/100 000 person-years that we found, was higher than reported in previous papers. Historically, an incidence of 1/60 000 has been reported.⁷ More recently Ponkilainen et al. reported an incidence of 9.2/100 000 person-years based on reviewing CT scans obtained after foot trauma, which is more in concordance with the findings we present.¹⁹ The high

incidence of Lisfranc injuries in our study compared to older studies can be explained by an increased focus on and awareness of the injury, as well as the use of CT scans which also detect the nondisplaced injuries.

Low-energy trauma such as falling from own height or a twisting injury of the foot, was the most common injury mechanism accounting for 44 % of the injuries. High-energy injuries, which often is reported to be the most common cause of Lisfranc injuries, only accounted for 32 % of the injuries.^{2,7,35-37} Our finding of low-energy trauma to be the most common cause of Lisfranc injuries is consistent with the findings of both Ponkilainen et al. and Renninger et al.^{19,39}

In our study, stable injuries were more common (55%) than unstable Lisfranc injuries (45%). To our knowledge the distribution between stable and unstable injuries has not been reported previously. The distribution between the genders were equal, but women were more likely to have an unstable injury than men ($p = .016$). This finding was unexpected as most authors report Lisfranc injuries to be more common in men, although Crates et al. and Komenda et al. reported an equal distribution between the genders.^{2,5,7,26,36,65,92}

When comparing the Lisfranc mortise between men and women, the latter had a shallower Lisfranc mortise, including a lower 2nd TMT joint height, which is in concordance with the findings of Yu-Kai et al.⁴⁷ This probably renders the Lisfranc joint less stable than in women than in men. The 2nd TMT joint height was also lower in the unstable injuries compared to the stable injuries. The medial mortise dept was not significant different ($p = 0.072$) between the stable and unstable group. A larger study population would maybe have changed this. The strength of the interosseous part of the Lisfranc ligament might be greater in patients with a deeper

Lisfranc mortise and a larger TMT-2 height, as one would expect this to facilitate a larger footprint for the origin and insertion of the ligament.

The fracture patterns of the of the Lisfranc injuries were determined on CT scans according to the columnar theory (divided into no fracture, avulsion fractures, intraarticular fractures and extraarticular fractures). Only intraarticular fractures in the lateral column increased the chance of the Lisfranc injury to be unstable (OR= 6.0, $p = .007$). Although the “fleck-sign” (avulsion fracture of the Lisfranc ligament) previously has been interpreted as a sign of instability we did not find the fleck-sign to increase the chance of instability, as also pointed out by Chen et al.^{2,51,52}

Paper 2

In this paper we studied the outcome after nonoperative treatment of the stable Lisfranc injuries. In general, there is an agreement of treating stable Lisfranc injuries nonoperatively, but only a few papers focus on the outcome after these less severe injuries.^{4,16,51,65} The few studies that report on these patients do report good outcome scores, but have included few patients (maximum 12) with confirmed stable injuries on follow-up radiographs.^{16,51} This study confirms that one can expect an excellent outcome after nondisplaced Lisfranc injuries that are verified to be stable using stress-test under fluoroscopy and weightbearing radiographs on follow-ups. One important finding was that the outcome score continued to improve even after 1 year, indicating that these patients can have a long rehabilitation time. Another interesting finding was that posttraumatic OA does not seem to be a frequent complication to stable Lisfranc

injuries at mid-term follow-up, unlike the high incidence of OA reported after unstable injuries.^{18,25,35,76} We only detected radiological signs of OA in 1 patient at the 1-year follow-up and no patients had clinical signs of AO at the final follow-up (median 55 months).

Lisfranc injuries are known to be easily overlooked on radiographs.^{10–12} The cases where there are no displacement in the TMT joints the injuries are even harder to detect.¹¹ This article did, not surprisingly, confirm that these injuries are easily missed on radiographs as 9/23 were not detected on initial radiographs. CT scans are therefore essential in detecting these injuries.

Paper 3

Preserving joints and mobility, thereby preserving the best possible functionality, is often the goal when treating orthopaedic injuries. In Lisfranc injuries, this goal must be weighed against the risk of developing an unstable midfoot not being able to provide the required stability needed during gait. Transarticular screw fixation has been the “gold standard” when treating unstable Lisfranc injuries.³⁵ Ly and Coetzee showed in their RCT published in 2006 better outcome when treating primarily ligamentous Lisfranc injuries with primary arthrodesis compared to transarticular screw fixation.⁶³ During the last decade several papers have been published on using bridge plating to stabilize the joints without inflicting further damage to the joints as transarticular screw fixation inevitably does.^{67,77–79,81} In paper 3 the joint preserving technique of temporary bridge plating the first TMT joint was compared to primary arthrodesis of the joint. The aim was to show that the joint preserving treatment was superior to the joint sacrificing treatment. Neither the AOFAS score, VAS pain score nor the SF-36 showed superiority of bridge plating technique compared to primary arthrodesis at 2-years follow-up.

However, we did confirm the findings of a high rate of mild posttraumatic OA after bridge plating seen in previous studies.^{67,79} A higher rate of complications were detected in the BP group compared to the PA group, although this was not statistically significant ($p=.348$). The additional surgery required in the BP group to remove the plate might be one of the reasons for the higher complication rate. In this study the mechanisms of injury, in the majority of the patients, were low-energy or sports related as the patients sustaining high-energy injuries often were polytrauma patients and thereby excluded. This is one of the weaknesses of the study as low-energy injuries are known to have better results than the injuries sustained during high-energy trauma.^{37,39,81}

Kirzner et al. has previously shown bridge plating to give better postoperative reduction compared to transarticular screw fixation, but this was not confirmed by Lau et al. in their study comparing bridge plating to transarticular screws or a combination of the two techniques^{67,78} We found a better postoperative reduction in the BP group when examining the alignment of the first metatarsal (AP Meary's angle). Pedobarographic examination revealed a statistically significant reduced peak pressure under the fifth metatarsal in the PA group both compared to the non-injured foot and compared to the BP group. Although the lateral Meary's angle in the PA group was not significantly reduced compared to the BP group, a slight reduction of the lateral Meary's angle together with shortening of the medial column occurring when performing an arthrodesis, can elevate the head of the first metatarsal. This can induce a pronation of the foot and thereby reduce the 5th metatarsal peak pressure.

Loss of reduction occurred in 11 % of the patients after treatment with bridge plating in the study by Kirzner et al. where the plates were removed at a minimum of 6 months.⁷⁸ We did not

see loss of reduction in our patients. This can possibly be explained by the fact that we fused the 2nd and 3rd TMT joints also in the BP group, whereas all unstable medial and central TMT joints were plated in Kirzners study.

Conclusions

- The incidence of all Lisfranc injuries was found to be 14/100 000 person-years, whereas the incidence of unstable injuries was 6/100 000 person-years. This is a higher incidence than previously reported.
- Low-energy trauma was the most common mechanism of injury and only 1/3 of the cases were caused by high-energy trauma.
- The distribution between genders were equal, but there was a higher proportion of unstable injuries in females.
- Intraarticular fractures in the lateral TMT joints increased the risk of having an unstable Lisfranc injury.
- Nonoperative treatment of nondisplaced, stable Lisfranc injuries yield excellent outcome. The rehabilitation period can be prolonged as improvement in the AOFAS score was detected even after 1 year.
- Posttraumatic OA does not seem to be at major problem after stable Lisfranc injuries.
- The outcome after temporary bridge plating of the first TMT joint in unstable Lisfranc injuries was not shown to be superior to primary arthrodesis.
- The alignment of the first metatarsal was better in the group treated with temporary bridge plating compared to primary arthrodesis.
- There was a high incidence of radiologically detected OA in the temporary bridge plate group (11/24) at 2 years.

Suggestion for future research

One of the most challenging issues when treating Lisfranc injuries can be to determine the stability of the injury, as the stability often dictates the treatment to follow. To date there is no consensus on the best way to determine stability in the nondisplaced injuries. Both weightbearing radiographs and stress-test under fluoroscopy have their proponents. The tests have been compared in cadaver studies, but not in a clinical setting. A study comparing the results of the two methods and following the patients to detect any missed instability could therefore help us determining which method is the best to detect occult instability in the Lisfranc joint complex.

Weightbearing CT scans is also expected to revolutionize the work-up of patients who has sustained a nondisplaced Lisfranc injury. Several papers have already been published on cadaver feet^{55,59,93} Investigating the displacement of subtle Lisfranc injuries on weightbearing CT scan is expected to make the evaluation of stability more accurate and to be a useful tool in the decision making to which injuries require surgery.

At last, Lisfranc injuries are a wide variety of injuries and all unstable injuries should not be treated the same way. Some injuries can benefit by only being treated by one homerun screw or a suture button device, while others are best treated by primary arthrodesis of the medial three TMT joints. Further studies are needed to evaluate these different treatment options.

References

1. Desmond EA, Chou LB. Current Concepts Review : Lisfranc Injuries. *Foot Ankle Int.* 2006;27(8):653-660. doi:10.1177/107110070602700819
2. Myerson MS, Fisher RT, Burgess AR, Kenzora JE. Fracture dislocations of the tarsometatarsal joints: end results correlated with pathology and treatment. *Foot Ankle.* 1986;6:225-242. doi:10.1177/107110078600600504
3. Myerson MS. The diagnosis and treatment of injury to the tarsometatarsal joint complex. *J Bone Joint Surg Br.* 1999;81(5):756-763. doi:10.1302/0301-620X.81B5.10369
4. Curtis MJ, Myerson M, Szura B. Tarsometatarsal joint injuries in the athlete. *Am J Sports Med.* 1993;21(4):497-502. doi:10.1177/036354659302100403
5. Goossens M, De Stoop N. Lisfranc's fracture-dislocations: etiology, radiology, and results of treatment. A review of 20 cases. *Clin Orthop Relat Res.* 1983;(176):154-162.
6. Faciszewski T, Burks RT, Manaster BJ. Subtle injuries of the Lisfranc joint. *J Bone Jt Surg.* 1990;72(10):1519-1522. doi:10.1016/j.pcad.2015.11.006
7. Hardcastle P, Reschauer R, Kutscha-Lissberg E, Schoffmann W. Injuries to the tarsometatarsal joint. Incidence, classification and treatment. *J Bone Jt Surg.* 1982;64-B(3):349-356.
8. Aitken A, Poulson D. Dislocations of the tarsometatarsal joint. *J Bone Jt Surg.* 1963;45-A(2):246-260.
9. English T. Dislocations of the metatarsal bone and adjacent toe. *J Bone Jt Surg.* 1964;46B(4):700-704.
10. Haapamaki V V, Kiuru MJ, Koskinen SK. Ankle and foot injuries: analysis of MDCT findings.

AJR Am J Roentgenol. 2004;183:615-622. doi:10.2214/ajr.183.3.1830615

11. Ponkilainen VT, Partio N, Salonen EE, et al. Inter- and intraobserver reliability of non-weight-bearing foot radiographs compared with CT in Lisfranc injuries. *Arch Orthop Trauma Surg.* March 2020. doi:10.1007/s00402-020-03391-w
12. Sherief TI, Mucci B, Greiss M. Lisfranc injury: how frequently does it get missed? And how can we improve? *Injury.* 2007;38(7):856-860. doi:10.1016/j.injury.2006.10.002
13. Raikin SM, Elias I, Dheer S, Besser MP, Morrison WB, Zoga AC. Prediction of midfoot instability in the subtle Lisfranc injury. Comparison of magnetic resonance imaging with intraoperative findings. *J Bone Joint Surg Am.* 2009;91(4):892-899. doi:10.2106/JBJS.H.01075
14. Haapamaki V, Kiuru M, Koskinen S. Lisfranc Fracture-dislocation in Patients with Multiple Trauma: Diagnosis with Multidetector Computed Tomography. *Foot Ankle Int.* 2004;25(9):614-619. doi:10.1177/107110070402500903
15. Watson TS, Shurnas PS, Denker J. Treatment of Lisfranc joint injury: current concepts. *J Am Acad Orthop Surg.* 2010;18(12):718-728.
16. Nunley JA, Vertullo CJ. Classification , Investigation , and Management of Midfoot Sprains Lisfranc Injuries in the Athlete. *Am J Sports Med.* 2002;30(6):871-878.
17. Lu J, Ebraheim N a, Skie M, Porshinsky B, Yeasting R a. Radiographic and computed tomographic evaluation of Lisfranc dislocation: a cadaver study. *Foot ankle Int.* 1997;18(6):351-355.
18. Stavlas P, Roberts CS, Xypnitos FN, Giannoudis P V. The role of reduction and internal fixation of Lisfranc fracture-dislocations: A systematic review of the literature. *Int Orthop.*

2010;34(8):1083-1091. doi:10.1007/s00264-010-1101-x

19. Ponkilainen VT, Laine HJ, Mäenpää HM, Mattila VM, Haapasalo HH. Incidence and Characteristics of Midfoot Injuries. *Foot Ankle Int.* 2019;40(1):105-112.
doi:10.1177/1071100718799741
20. CHAN CW, RUDINS A. Foot Biomechanics During Walking and Running. *Mayo Clin Proc.* 1994;69(5):448-461. doi:10.1016/S0025-6196(12)61642-5
21. DeOrio M, Erickson M, Usuelli FG, Easley M. Lisfranc injuries in sport. *Foot Ankle Clin.* 2009;14(2):169-186. doi:10.1016/j.fcl.2009.03.008
22. Eleftheriou KI, Rosenfeld PF, Calder JDF. Lisfranc injuries: an update. *Knee Surg Sports Traumatol Arthrosc.* 2013;21(6):1434-1446. doi:10.1007/s00167-013-2491-2
23. Pearce CJ, Calder JD. Surgical anatomy of the midfoot. *Knee Surgery, Sport Traumatol Arthrosc.* 2010;18(5):581-586. doi:10.1007/s00167-010-1101-9
24. Ouzounian TJ, Shereff MJ. In vitro determination of midfoot motion. *Foot Ankle.* 1989;10(3):140-146. doi:10.1177/107110078901000305
25. Mulier T, Reynders P, Dereymaeker G, Broos P. Severe Lisfrancs injuries: primary arthrodesis or ORIF? *Foot ankle Int / Am Orthop Foot Ankle Soc [and] Swiss Foot Ankle Soc.* 2002;23(10):902-905.
26. Komenda GA, Myerson MS, Biddinger KR. Results of arthrodesis of the tarsometatarsal joints after traumatic injury. *J Bone Jt Surg.* 1996;78(11):1665-1676.
27. Sangeorzan BJ, Veith RG, Hansen ST. Salvage of Lisfranc's tarsometatarsal joint by arthrodesis. *Foot Ankle.* 1990;10:193-200. doi:10.1177/107110079001000401
28. Weatherford BM, Anderson JG, Bohay DR. Management of Tarsometatarsal Joint

- Injuries. *J Am Acad Orthop Surg*. 2017;25(7):469-479. doi:10.5435/JAAOS-D-15-00556
29. Solan MC, Moorman CT, Miyamoto RG, Jasper LE, Belkoff SM. Ligamentous restraints of the second tarsometatarsal joint: a biomechanical evaluation. *Foot ankle Int*. 2001;22(8):637-641. doi:10.1177/107110070102200804
 30. de Palma L, Santucci a, Sabetta SP, Rapali S. Anatomy of the Lisfranc joint complex. *Foot Ankle Int*. 1997;18(6):356-364. <http://www.ncbi.nlm.nih.gov/pubmed/9208295>.
 31. Panchbhavi VK, Molina D, Villarreal J, Curry MC, Andersen CR. Three-dimensional, digital, and gross anatomy of the Lisfranc ligament. *Foot Ankle Int*. 2013;34(6):876-880. doi:10.1177/1071100713477635
 32. Panchbhavi VK, Andersen CR, Vallurupalli S, Yang J. A minimally disruptive model and three-dimensional evaluation of Lisfranc joint diastasis. *J Bone Joint Surg Am*. 2008;90(12):2707-2713. doi:10.2106/JBJS.G.01420
 33. Coss LHS, USNR M, Manos LRE, et al. Abduction Stress and AP Weightbearing Radiography of Purely Ligamentous Injury in the Tarsometatarsal Joint. *Foot Ankle Int*. 1998;19(8):537-541. doi:10.1177/107110079801900806
 34. Kaar S, Femino J, Morag Y. Lisfranc joint displacement following sequential ligament sectioning. *J Bone Jt Surg - Ser A*. 2007;89(10):2225-2232. doi:10.2106/JBJS.F.00958
 35. Kuo RS, Tejwani NC, Digiovanni CW, et al. Outcome after open reduction and internal fixation of Lisfranc joint injuries. *J Bone Jt Surg - Ser A*. 2000;82(11):1609-1618. doi:10.2106/00004623-200011000-00015
 36. Arntz CT, Veith RG, Hansen ST, et al. Fractures and fracture-dislocations of the tarsometatarsal joint. *J Bone Joint Surg Am*. 1988;70(2):173-181. doi:10.1097/00005373-

197403000-00001

37. Lau SC, Guest C, Hall M, Tacey M, Joseph S, Oppy A. Do columns or sagittal displacement matter in the assessment and management of Lisfranc fracture dislocation? An alternate approach to classification of the Lisfranc injury. *Injury*. 2017;48(7):1689-1695.
doi:10.1016/j.injury.2017.03.046
38. Vuori JP, Aro HT. Lisfranc joint injuries: trauma mechanisms and associated injuries. *J Trauma*. 1993;35:40-45. doi:10.1097/00005373-199307000-00007
39. Renninger CH, Cochran G, Tompane T, Bellamy J, Kuhn K. Injury Characteristics of Low-Energy Lisfranc Injuries Compared With High-Energy Injuries. *Foot Ankle Int*. 2017;38(9):964-969. doi:10.1177/1071100717709575
40. Shapiro MS, Wascher DC, Finerman G a. Rupture of Lisfranc's ligament in athletes. *Am J Sports Med*. 1992;22(5):687-691. doi:10.1177/036354659402200518
41. Myerson MS, Cerrato R. Current management of tarsometatarsal injuries in the athlete. *J Bone Jt Surg Am*. 2008;(Nov;90(11)):2522-2533.
<https://pubmed.ncbi.nlm.nih.gov/18978422/>. Accessed November 6, 2020.
42. Ross G, Cronin R, Hauzenblas J, Juliano P. *Plantar Ecchymosis Sign: A Clinical Aid to Diagnosis of Occult Lisfranc Tarsometatarsal Injuries*. *Journal of orthopaedic trauma* 10, 119-122 (1996). doi:10.1097/00005131-199602000-00008
43. Norfray JF, Geline RA, Steinberg RI, Galinski AW, Gilula LA. Subtleties of Lisfranc fracture-dislocations. *Am J Roentgenol*. 1981;137(6):1151-1156. doi:10.2214/ajr.137.6.1151
44. Foster SC, Foster RR. Lisfranc's tarsometatarsal fracture dislocation. *Radiology*. 1976;120(1):79-83. doi:10.1148/120.1.79

45. Sripanich Y, Weinberg MW, Krähenbühl N, et al. Imaging in Lisfranc injury: a systematic literature review. *Skeletal Radiol.* 2001;49(1):31-53. doi:10.1007/s00256-019-03282-1
46. Seo D, Lee H, Lee KW, Lee SK, Kim S. Nonweightbearing Radiographs in Patients With a Subtle Lisfranc Injury. *Foot ankle Int.* 2017;38(10):1120-1125.
doi:10.1177/1071100717717220
47. Yu-Kai Y, Shiu-Bii L. Anatomic Parameters of the Lisfranc Joint Complex in a Radiographic and Cadaveric Comparison. *J Foot Ankle Surg.* 2015;54(5):883-887.
doi:10.1053/j.jfas.2015.02.021
48. Shakked RJ. Lisfranc injury in the athlete. *JBJS Rev.* 2017;5(9):e4.
doi:10.2106/JBJS.RVW.17.00025
49. Potter HG, Deland JT, Gusmer PB, Carson E, Warren RF. Magnetic resonance imaging of the Lisfranc ligament of the foot. *Foot Ankle Int.* 1998;19(7):438-446.
<http://www.ncbi.nlm.nih.gov/pubmed/9694121>.
50. Preidler KW, Peicha G, Lajtai G, et al. Conventional radiography, CT, and MR imaging in patients with hyperflexion injuries of the foot: Diagnostic accuracy in the detection of bony and ligamentous changes. *Am J Roentgenol.* 1999;173(6):1673-1677.
doi:10.2214/ajr.173.6.10584818
51. Chen P, Ng N, Snowden G, Mackenzie SP, Nicholson JA, Amin AK. Rates of Displacement and Patient-Reported Outcomes Following Conservative Treatment of Minimally Displaced Lisfranc Injury. *Foot Ankle Int.* 2020;41(4):387-391.
doi:10.1177/1071100719895482
52. Seybold JD, Coetzee JC. Lisfranc Injuries: When to Observe, Fix, or Fuse. *Clin Sports Med.*

- 2015;34(4):705-723. doi:10.1016/j.csm.2015.06.006
53. Ebraheim N a, Yang H, Lu J, Biyani A. Computer evaluation of second tarsometatarsal joint dislocation. *Foot Ankle Int.* 1996;17(11):685-689.
 54. Lau S, Bozin M, Thillainadesan T. Lisfranc fracture dislocation: A review of a commonly missed injury of the midfoot. *Emerg Med J.* 2017. doi:10.1136/emmermed-2015-205317
 55. Sripanich Y, Weinberg M, Krähenbühl N, Rungprai C, Saltzman CL, Barg A. Change in the First Cuneiform–Second Metatarsal Distance After Simulated Ligamentous Lisfranc Injury Evaluated by Weightbearing CT Scans. *Foot Ankle Int.* 2020. doi:10.1177/1071100720938331
 56. Chiodo CP, Myerson MS. Developments and advances in the diagnosis and treatment of injuries to the tarsometatarsal joint. *Orthop Clin North Am.* 2001;32(1):11-20. doi:10.1016/S0030-5898(05)70190-5
 57. Naguib S, Meyr AJ. Reliability, Surgeon Preferences, and Eye-Tracking Assessment of the Stress Examination of the Tarsometatarsal (Lisfranc) Joint Complex. *J Foot Ankle Surg.* 2019;58(1):93-96. doi:10.1053/j.jfas.2018.08.015
 58. Kitsukawa K, Hirano T, Niki H, Tachizawa N, Nakajima Y, Hirata K. MR Imaging Evaluation of the Lisfranc Ligament in Cadaveric Feet and Patients with Acute to Chronic Lisfranc Injury. *Foot Ankle Int.* 2015;36(12):1483-1492. doi:10.1177/1071100715596746
 59. Sripanich Y, Weinberg MW, Krähenbühl N, Rungprai C, Saltzman CL, Barg A. Reliability of measurements assessing the Lisfranc joint using weightbearing computed tomography imaging. *Arch Orthop Trauma Surg.* 2020. doi:10.1007/s00402-020-03477-5
 60. Schepers T, Rammelt S. Classifying the Lisfranc injury: Literature overview and a new

classification. *Fuss und Sprunggelenk*. 2018;16(3):151-159.

doi:10.1016/j.fuspru.2018.07.003

61. Calder J. Results of isolated Lisfranc injuries and the effect of compensation claims. *J Bone Jt* 2004;527-530. doi:10.1302/0301-620X.86B4.13761
62. Coetzee JC. Making Sense of Lisfranc Injuries. *Foot Ankle Clin*. 2008;13(4):695-704.
doi:10.1016/j.fcl.2008.07.001
63. Ly T V., Coetzee JC. Treatment of Primarily Ligamentous Lisfranc Joint Injuries: Primary Arthrodesis Compared with Open Reduction and Internal Fixation<sbt
aid="1080838">A Prospective, Randomized Study</sbt> *J Bone Jt Surg*. 2006;88(3):514. doi:10.2106/JBJS.E.00228
64. Weatherford BM, Bohay DR, Anderson JG. Open Reduction and Internal Fixation Versus Primary Arthrodesis for Lisfranc Injuries. *Foot Ankle Clin*. 2017;22(1):1-14.
doi:10.1016/j.fcl.2016.09.002
65. Crates JM, Barber FA, Sanders EJ. Subtle Lisfranc Subluxation: Results of Operative and Nonoperative Treatment. *J Foot Ankle Surg*. 2015;54(3):350-355.
doi:10.1053/j.jfas.2014.07.015
66. Ponkilainen VT, Partio N, Salonen EE, et al. Outcomes after nonoperatively treated non-displaced Lisfranc injury: a retrospective case series of 55 patients. *Arch Orthop Trauma Surg*. 2020. doi:10.1007/s00402-020-03599-w
67. Lau S, Howells N, Millar M, De Villiers D, Joseph S, Oppy A. Plates, Screws, or Combination? Radiologic Outcomes After Lisfranc Fracture Dislocation. *J Foot Ankle Surg*. 2016. doi:10.1053/j.jfas.2016.03.002

68. Richter M, Thermann H, Huefner T, Schmidt U, Krettek C. Aetiology, treatment and outcome in Lisfranc joint dislocations and fracture dislocations. *Foot Ankle Surg.* 2002;8(1):21-32. doi:10.1046/j.1460-9584.2002.00294.x
69. Rammelt S, Schneiders W, Schikore H, Holch M, Heineck J, Zwipp H. Primary open reduction and fixation compared with delayed corrective arthrodesis in the treatment of tarsometatarsal (Lisfranc) fracture dislocations. *J Bone Joint Surg Br.* 2008;Nov 2008(90,11):1499.
70. Benirschke SK, Meinberg E, Anderson SA, Jones CB, Cole PA. Fractures and Dislocations of the Midfoot: Lisfranc and Chopart Injuries. *J Bone Jt Surg.* 2012;94-A(14):1326-1337. doi:10.1053/j.jfas.2010.08.005
71. Wilppula E. Tarsometatarsal fracture-dislocation. Late results in 26 patients. *Acta Orthop Scand.* 1973;44(3):335-345.
72. Brunet JA, Wiley JJ. The late results of tarsometatarsal joint injuries. *J Bone Jt Surg - Ser B.* 1987;69(3):437-440. doi:10.1302/0301-620x.69b3.3108261
73. Schepers T, Oprel PP, Van Lieshout EMM. Influence of Approach and Implant on Reduction Accuracy and Stability in Lisfranc Fracture-Dislocation at the Tarsometatarsal Joint. *Foot Ankle Int.* 2013;34(5):705-710. doi:10.1177/1071100712468581
74. Lee CA, Birkedal JP, Dickerson EA, Vieta PA, Webb LX, Teasdall RD. Stabilization of Lisfranc joint injuries: A biomechanical study. *Foot Ankle Int.* 2004;25(5):365-370. doi:10.1177/107110070402500515
75. Henning JA, Jones CB, Sietsema DL, Bohay DR, Anderson JG. Open reduction internal fixation versus primary arthrodesis for lisfranc injuries: A prospective randomized study.

- Foot Ankle Int.* 2009;30(10):913-922. doi:10.3113/FAI.2009.0913
76. Pereira, Cléber de Jesus; Espinosa EG. Evaluation of the Surgical Treatment of Lisfranc Joint Fracture-Dislocation. *Acta Orthop Bras.* 2008;16:93-97.
77. Alberta FG, Aronow MS, Barrero M, Diaz-Doran V, Sullivan RJ, Adams DJ. Ligamentous Lisfranc joint injuries: a biomechanical comparison of dorsal plate and transarticular screw fixation. *Foot Ankle Int.* 2005;26(6):462-473.
<http://www.ncbi.nlm.nih.gov/pubmed/15960913>. Accessed March 11, 2014.
78. Kirzner N, Zotov P, Goldbloom D, Curry H, Bedi H. Dorsal bridge plating or transarticular screws for Lisfranc fracture dislocations. *Bone Joint J.* 2018. doi:10.1302/0301-620X.100B4.BJJ-2017-0899.R2
79. Stern RE, Assal M. Dorsal Multiple Plating Without Routine Transarticular Screws for Fixation of Lisfranc Injury. *Orthopedics.* 2014;37(12):815-819. doi:10.3928/01477447-20141124-03
80. Hu S, Chang S, Li X, Yu G. Outcome comparison of Lisfranc injuries treated through dorsal plate fixation versus screw fixation. *Acta Ortopédica Bras.* 2014. doi:10.1590/1413-78522014220600576
81. Koperen PJ Van, Jong VM De, Luitse JSK, Schepers T. Functional Outcomes After Temporary Bridging With Locking Plates in Lisfranc Injuries. *J Foot Ankle Surg.* 2016;55(5):922-926. doi:10.1053/j.jfas.2016.04.005
82. Cochran G, Renninger C, Tompane T, Bellamy J, Kuhn K. Primary Arthrodesis versus Open Reduction and Internal Fixation for Low-Energy Lisfranc Injuries in a Young Athletic Population. *Foot Ankle Int.* 2017;38(9):957-963. doi:10.1177/1071100717711483

83. Kitaoka HB, Alexander IJ, Adelaar RS, Nunley JA, Myerson MS, Sanders M. Clinical Rating Systems for the Ankle-Hindfoot, Midfoot, Hallux, and Lesser Toes. *Foot Ankle Int.* 1994;15(7):349-353. doi:10.1177/107110079401500701
84. Garratt AM, Stavem K. Measurement properties and normative data for the Norwegian SF-36: Results from a general population survey. *Health Qual Life Outcomes.* 2017;15(1):1-10. doi:10.1186/s12955-017-0625-9
85. Ware JE, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care.* 1992;30(6):473-483. <http://www.ncbi.nlm.nih.gov/pubmed/1593914>. Accessed October 29, 2019.
86. Myerson M. The diagnosis and treatment of injuries to the Lisfranc joint complex. *Orthop Clin North Am.* 1989;20:655-664.
87. Hunt KJ, Hurwit D. Use of patient-reported outcome measures in foot and ankle research. *J Bone Jt Surg - Ser A.* 2013;95(16):e118 (1-9). doi:10.2106/JBJS.L.01476
88. Ponkilainen VT, Uimonen M, Repo JP, Mattila VM, Haapasalo HH. Validity and internal consistency of the American Orthopaedic Foot and Ankle Society Midfoot Scale in patients with Lisfranc injury. *Foot Ankle Surg.* 2020;26(5):523-529. doi:10.1016/j.fas.2019.06.005
89. Kitaoka HB, Meeker JE, Phisitkul P, Adams SB, Kaplan JR, Wagner E. AOFAS Position Statement Regarding Patient-Reported Outcome Measures. *Foot Ankle Int.* 2018;39(12):1389-1393. doi:10.1177/1071100718809066
90. Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain: Visual Analog Scale for Pain (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire

(MPQ), Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF-36 BPS), and Measure of Intermittent and Constant Osteoarthritis Pain (ICOAP). *Arthritis Care Res.* 2011;63(SUPPL. 11). doi:10.1002/acr.20543

91. Biswas D, Bible JE, Bohan M, Simpson AK, Whang PG, Grauer JN. Radiation exposure from musculoskeletal computerized tomographic scans. *J Bone Jt Surg - Ser A.* 2009. doi:10.2106/JBJS.H.01199
92. Kuo RS, Tejawani NC, Digiovanni CW, et al. Outcome After Open Reduction and Internal Fixation of Lisfranc Joint Injuries. *J Bone Jt Surg.* 2000;82-A(11):1609-1618.
93. Sripanich Y, Steadman J, Krähenbühl N, et al. Asymmetric lambda sign of the second tarsometatarsal joint on axial weight-bearing cone-beam CT scans of the foot: preliminary investigation for diagnosis of subtle ligamentous Lisfranc injuries in a cadaveric model. *Skeletal Radiol.* 2020;49(10):1615-1621. doi:10.1007/s00256-020-03445-5

Contents lists available at ScienceDirect

Foot and Ankle Surgery

journal homepage: www.elsevier.com/locate/fas

Lisfranc injuries: Incidence, mechanisms of injury and predictors of instability

Are H. Stølle^{a,c,*}, Kjetil H. Hvaal^a, Martine Enger^{a,c}, Helga Brøgger^b, Jan Erik Madsen^{a,c}, Elisabeth Ellingsen Husebye^a^a Division of Orthopaedic Surgery, Oslo University Hospital, Norway^b Department of Radiology and Nuclear Medicine, Oslo University Hospital, Norway^c Institute of Clinical Medicine, University of Oslo, Norway

ARTICLE INFO

Article history:

Received 21 November 2018

Received in revised form 15 May 2019

Accepted 13 June 2019

Keywords:

Lisfranc injury

Tarsometatarsal joint injury

Midfoot injury

ABSTRACT

Background: In Lisfranc injuries the stability of the tarsometatarsal joints guides the treatment of the injury. Determining the stability, especially in the subtle Lisfranc injuries, can be challenging. The purpose of this study was to identify incidence, mechanisms of injury and predictors for instability in Lisfranc injuries.

Methods: Eighty-four Lisfranc injuries presenting at Oslo University Hospital between September 2014 and August 2015 were included. The diagnosis was based on radiologically verified injuries to the tarsometatarsal joints. Associations between radiographic findings and stability were examined.

Results: The incidence of Lisfranc injuries was 14/100,000 person-years, and only 31% were high-energy injuries. The incidence of unstable injuries was 6/100,000 person-years, and these were more common in women than men ($P = 0.016$). Intraarticular fractures in the two lateral tarsometatarsal joints increased the risk of instability ($P = 0.007$). The height of the second tarsometatarsal joint was less in the unstable injuries than in the stable injuries ($P = 0.036$).

Conclusion: The incidence of Lisfranc injuries in the present study is higher than previously published. The most common mechanism of injury is low-energy trauma. Intraarticular fractures in the two lateral tarsometatarsal joints, female gender and shorter second tarsometatarsal joint height increase the risk of an unstable injury.

Level of Evidence: Level III, cross-sectional study.

© 2019 European Foot and Ankle Society. Published by Elsevier Ltd. All rights reserved.

1. Introduction

Lisfranc (tarsometatarsal joint) injuries are complex, and can lead to high morbidity and substantial disability if not adequately treated [1–7]. The incidence has been reported to be 1/60,000 person-years [5]; however, these injuries may be underreported as up to 24% are missed on primary radiographs [8]. An increased awareness of these injuries combined with increased use of MRI, CT scans, weightbearing radiographs and stress fluoroscopy to detect them, seem to have increased the incidence compared to that reported in previous publications [2,9–13].

High-energy trauma (motor vehicle accidents, falls from height and crush injuries) has been reported to account for the majority of the cases [4,5,10,14,15]. Low-energy injuries are most often caused

by axial and/or rotational forces on a foot fixed in equinus, and these injuries tend to be more subtle [3,16,17].

The stability of the midfoot is provided by osseous and ligamentous structures. Injuries to these structures may cause instability and progress to displacement of the tarsometatarsal joints [18]. In the subtle Lisfranc injuries with no displacement on radiographs or CT scans, it can be challenging to determine the stability of the injury. Weightbearing radiographs and stress fluoroscopy have been advocated to reveal an occult instability, preferably with images of the non-injured foot for comparison [3,19–24]. Lisfranc injuries without detectable displacement on weightbearing radiographs or on stress fluoroscopy should generally be treated non-operatively, whereas for the unstable injuries anatomic reduction and stable fixation is recommended [3–5,14,20,21,25,26]. In the acute phase both stress fluoroscopy and weightbearing radiographs can be painful. Furthermore, the interpretation of these examinations are also often subjective and examiner dependent. CT scans can be useful in the evaluation of Lisfranc injuries and many patients will be subjected to a CT scan of

* Corresponding author at: Ortopedisk avdeling, Oslo Universitetssykehus, Kirkeveien 166, 0450 Oslo, Norway.
E-mail address: arhauk@ous-hf.no (A.H. Stølle).

their foot. Therefore, determining radiological predictors of instability on CT images can be valuable in improving diagnosis of these injuries.

The aim of this study was to survey the incidence, the most common mechanisms of injury and to evaluate radiological predictors of instability in Lisfranc injuries. The hypothesis was that Lisfranc injuries and especially low-energy injuries are more frequent than previously reported, and that CT scans can help predict instability.

2. Materials and methods

The study was approved by the Regional Ethics Committee (2014/853/REK) and the patients signed an informed consent form. Between the 1st of September 2014 and the 31st of August 2015 all patients with Lisfranc joint injuries treated at Oslo University Hospital (a level one trauma center) and Oslo Accident and Emergency Department were registered. A Lisfranc injury was defined as injury to tarsometatarsal joint with avulsion fractures, intra-articular fractures and/or displacement of tarsometatarsal joint. Injuries to the tarsometatarsal joint were identified using radiographs, CT scans, MRI, stress fluoroscopy and/or weightbearing radiographs. Patients with isolated fracture of the fifth metatarsal and patients with Charcot arthropathy were excluded from the study.

Demographic data were recorded at presentation, as well as mechanism of injury and clinical findings. To determine the incidence of Lisfranc injuries all patients referred for treatment from other hospitals were excluded and only the patients with a permanent address in the Oslo University Hospital catchment area were included. On January 1st 2015 the hospital had a local catchment population of 399,665.

High-energy injuries were defined as injuries caused by motor vehicle accidents (MVA), fall from height (>3 m) and crush injuries. Low-energy injuries were fall from own height, twisting injury of the foot, falling down stairs, bike accidents, kicking into an object. Sports related injuries were categorized separately.

The diagnostic algorithm is presented in Fig. 1. Ten patients did not have a primary radiograph when admitted, as they had already had a CT or MRI scan. In patients without joint displacement on the CT scan, a stress fluoroscopy of both injured and non-injured foot was performed 7–14 days after the injury. Stress fluoroscopy could be performed without anesthesia in the majority of the patients. If stress fluoroscopy was not possible due to pain, general anesthesia was applied. Weightbearing radiographs of both feet (AP, lateral and 30° oblique views) were also used for evaluation when the stress fluoroscopy was inconclusive regarding stability.

Radiographs and CT scans were analyzed using Syngo Studio VB36E (Siemens Healthcare GmbH, Erlangen, Germany). The images were evaluated by two foot and ankle consultants and one radiology consultant experienced in musculoskeletal imaging. Fractures were categorized as intraarticular, extraarticular or avulsion fractures. Joint displacement of 2 mm or more were registered. The Lisfranc injuries were defined as unstable if there was a displacement of ≥ 2 mm in a tarsometatarsal, intercuneiform or naviculocuneiform joint on any of the initial non-weightbearing radiographs, CT scans or weightbearing radiographs, or if the patient had a positive stress fluoroscopy with joint incongruity.

The second metatarsal base is recessed between the medial and lateral cuneiforms in a “mortise”. The medial and lateral depth of the Lisfranc mortise as well as the height of the second tarsometatarsal joint, were measured on the CT scans by the radiology consultant (Fig. 2).

The findings on radiographs and CT scans were correlated to the fluoroscopically evaluated stability to reveal any radiographic predictors of instability. All patients with unstable Lisfranc injuries

Fig. 1. Diagnostic algorithm for suspected Lisfranc injuries. Displacement is defined as ≥ 2 mm displacement of a tarsometatarsal, intercuneiform or naviculocuneiform joint on radiographs or CT scans, or obvious displacement on stress fluoroscopy.

were recommended operative treatment, while the patients with stable injuries were treated with a below knee cast for 6 weeks and then examined with weightbearing radiographs of both feet.

2.1. Statistics

Descriptive statistical analyses were used to determine frequencies of categorical variables and the group mean and standard deviation of continuous variables. The independent samples t-test was used to compare group means for continuous variables and for categorical variables the odds ratio and Pearson Chi-square test was used. The correlation between fracture pattern and stability was assessed using logistic regression. The interrater reliability when evaluating fractures and dislocations was calculated using the intraclass correlation coefficient. The statistical analyses were performed using SPSS version 25 (IBM, Armonk, New York). A threshold of $p < 0.05$ was set for statistical significance.

3. Results

3.1. Patient demographics

Eighty-nine Lisfranc injuries were registered prospectively during the one-year period. Eighty-four patients consented to participate in the study. One patient had bilateral Lisfranc injuries. There was an equal distribution between genders (Table 1). The mean age was 41.0 (range, 14–83) years and the men were on average 10 years younger than the women (36.0 vs 45.8, $P = 0.05$).

Fifty-four of the 89 patients with Lisfranc injury lived in the Oslo University Hospital catchment area, and resulted in an incidence of all Lisfranc injuries of 14/100,000 person-years. Twenty-two of these patients had injuries with instability, resulting in an incidence for unstable Lisfranc injuries of 6/100,000 person-years.

3.2. Mechanism of injury

The mechanisms of injury are presented in Table 2. High-energy mechanisms (motor vehicle accidents (MVA), falls from more than

Fig. 2. (a) CT scan sagittal plane, left foot. Height of the second tarsometatarsal joint (arrow). Red line indicating coronal plane centered in second tarsometatarsal joint (2b). (b) CT scan left foot, coronal plane centered in the second tarsometatarsal joint as shown in picture 2a. Distance A represent the medial Lisfranc mortise depth, and distance B represent the lateral Lisfranc mortise depth (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article).

three meters height and crush injuries) accounted for 31% of the injuries. The single most common mechanism of injury was fall from own height / twisting injury of the foot, occurring in 31% of the cases. In 21% percent, the injuries were sports related.

3.3. Radiological assessment and stability

Seventy-four feet (87%) had a primary nonweightbearing radiograph and 21 (28%) of these radiographs were described as normal. All patients except one had a CT scan of their injured foot (84 ft), all with findings consistent with Lisfranc injury. The

intratester reliability of evaluating the fractures and displacements on radiographs and CT scans, was 0.83 (95% CI, 0.81–0.84), determined with the intraclass correlation coefficient.

Thirty-eight (45%) Lisfranc injuries were defined as unstable and 47 (55%) were stable (Table 2). Joint displacement as a sign of instability, was mainly detected on CT scans (17 ft) or a positive stress fluoroscopy (14 ft) (Table 3). In one patient an increased diastasis between the medial and middle cuneiform was detected on weightbearing radiographs, this was not detected on the stress fluoroscopy. Two other patients had an instability that was overlooked on initial stress fluoroscopy, but detected on weight-bearing radiographs at the 6 weeks follow-up.

The distribution of avulsion fractures, intraarticular fractures and extraarticular fractures is shown in Table 4. The only fracture pattern on CT scans that was correlated to instability in Lisfranc injuries was an intraarticular fracture of the fourth and/or fifth tarsometatarsal joint (OR = 6.0, 95% CI = 1.6–21.5).

When evaluating the Lisfranc mortise measurements, an increased height of the second tarsometatarsal (TMT) joint in the feet with a stable injury compared to those with an unstable injury was observed (21.2 vs 20.1 mm, $p=0.04$). The medial Lisfranc mortise depth was larger in the group with stable Lisfranc injuries compared to the unstable group (mean 7.3 vs 6.6 mm), but this finding was not statistically significant ($p=0.07$). Women had more shallow mortise depths and lower TMT-2 heights compared to men (Table 5).

4. Discussion

The most important findings of the present study are that we observed a higher incidence of Lisfranc injuries than previously reported in the literature, and that the majority of the injuries are low-energy or sports-related. Furthermore, intraarticular fractures of the lateral tarsometatarsal joints, female gender and a lower second tarsometatarsal joint height increase the risk of having an unstable Lisfranc injury.

In the present study, all Lisfranc injuries during a one year period were prospectively registered. The overall incidence of both stable and unstable Lisfranc injuries was 14/100,000 person-years, whereas the incidence of unstable injuries was 6/100,000 person-years. The incidence of Lisfranc injuries has been reported to be 1/60,000 person-years or 0.2 percent of fractures based on older studies [5,27,28]. Recently Ponkilainen et al. published a CT based study where they retrospective examined all CT scans of midfoot fractures during a 5-year period. They found the CT based incidence of Lisfranc injuries to be 9.2/100,000 person-years. The findings of the present study and those of Ponkilainen et al. suggest that the incidence of Lisfranc injuries is probably higher than previously reported. The high incidence found in the present study may be caused by the prospective design, a higher awareness of these injuries and the use of more advanced diagnostic tools such as CT scans, MRIs, stress fluoroscopy and weightbearing radiographs, thereby also detecting the more subtle injuries.

We found high-energy trauma to be the cause of injury in only 31% of patients and low-energy trauma to be the most common injury mechanism. Numerous authors have reported Lisfranc injuries primarily being caused by high-energy trauma [4,5,10,14]. More recently, however, Renninger et al. found that 60% of the surgically treated Lisfranc injuries at their institution resulted from low energy trauma [17]. Ponkilainen et al. also reported the majority of Lisfranc injuries to be caused by low-energy trauma and only 36.5% of the injuries being caused by high-energy trauma mechanisms [29]. This emphasizes that one should have a high suspicion of Lisfranc injuries even in patients with midfoot pain after a low-energy trauma as these injuries may lead to severe disability if they are missed or treated inadequately [30,31].

Table 1
Patient demographics.

	Stable injury	Unstable injury	Total	P-value	Odds ratio
Mean age (SD)	38.4 (16.5)	44.2 (15.6)	41.0 (16.3)	0.10	
Gender (male/female)	29/18	13/25 ^a	42/43 ^a	0.016	OR = 3.1 (1.3–7.6)
Side (right/left)	21/26	17/21	38/47	1.0	OR = 1.0 (0.4–2.4)
Days to diagnosis	3.2 (7.8)	3.5 (9.7)	3.4 (8.7)	0.92	
Ipsilateral FA fx ^b	9	5	14	0.46	OR = 0.6 (0.2–2.1)

^a One female patient with bilateral unstable Lisfranc injuries.^b Other ipsilateral foot and ankle fractures.**Table 2**
Mechanism of injury.

Mechanism of injury	Lisfranc Injuries		
	Stable	Unstable	Total
- Fall from own height/ twisting of foot	14	12	26
- Bike accident	2	2	4
- Fall down stairs	1	3	4
- Kicked into an object	2	1	3
- Sports related injuries	7	11	18
Soccer	4	3	7
Gymnastics	2	2	4
Martial arts	0	3	3
Windsurfing/kiting	1	1	2
Snowboard	0	1	1
Skateboard	1	0	1
- Motor vehicle accident	7	3	10
- Fall > 3 m	3	3 ^a	6
- Crush injury	10	1	11
- Unknown ^b	1	2	3
Total	47	38	85

^a One patient with bilateral injuries.^b Unknown due to alcohol intoxication.**Table 3**
Detection of joint displacement.

	No. of feet	Negative	Joint displacement detected
Primary radiographs (non-WB)	74	21 ^a	4
CT scan	84	0 ^a	17
Stress-test under fluoroscopy	67	45 ^b	14
Primary weightbearing radiographs	19	17 ^b	1
Follow-up weightbearing radiographs	30	28 ^b	2
Sum			38

Weight-bearing radiographs were compared to the non-injured side. CT scans and radiographs were registered as positive if there were any fractures (including minor avulsion fractures) or joint displacements.

^a Negative in terms of no joint displacement or fracture (including small avulsion fractures).^b No joint displacement detected.

Evaluating the stability of Lisfranc injuries is essential in treating these injuries as nonoperative treatment is recommended in stable injuries and operative treatment in unstable injuries [19,26,32]. Occult instability in a Lisfranc injury can be detected by either weightbearing radiographs or stress fluoroscopy [4,19,20]. The stress fluoroscopy has been criticized for being subjective and examiner dependent [24]. On the other hand, Kaar et al. demonstrated in a cadaver study that stress fluoroscopy had better sensitivity in detecting instability compared to weightbearing radiographs [22]. Both stress fluoroscopy and weightbearing radiographs present a challenge in the acute setting, as they can

Table 4
CT scan findings related to stability.

	No. feet	Stable injuries	Unstable injuries	Odds ratio (95% CI)	P-value
CT scan	84	46	38		
"Fleck" sign	39	18	21	2.00 (0.64– 6.21)	0.23
Medial column					
No fracture	16	11	5	Ref	
Avulsion fractures	32	19	13	1.88 (0.39– 9.01)	0.43
Intraarticular fractures	35	15	20	2.51 (0.53– 11.94)	0.25
Extraarticular fractures	1	1	0	NA	
Middle column					
No fracture	8	5	3	Ref	
Avulsion fractures	9	8	1	0.14 (0.01– 2.15)	0.16
Intraarticular fractures	56	25	31	0.81 (0.13– 4.89)	0.82
Extraarticular fractures	11	8	3	0.45 (0.05– 3.94)	0.47
Lateral column					
No fracture	36	26	10	Ref	
Avulsion fractures	2	0	2	NA	
Intraarticular fractures	31	9	22	5.95 (1.64– 21.54)	0.007
Extraarticular fractures	15	11	4	1.13 (0.25– 5.22)	0.87

Fracture patterns detected on CT scans related to stability of the Lisfranc injury. Statistical significant finding highlighted. Ref = reference group. NA = not applicable.

be painful examinations. Since we delayed the stress fluoroscopy until 7–14 days after injury, we were able to perform the testing without anesthesia in most patients. However, two of the 45 Lisfranc injuries initially evaluated as stable after stress fluoroscopy had a positive weightbearing radiograph indicating midfoot instability on the 6 weeks follow-up. This emphasizes the importance of follow-up with weightbearing radiographs in patients with injuries that initially are evaluated as stable, as also recommended by Myerson and Cerrato [16].

As both stress fluoroscopy and weightbearing radiographs are challenging to perform in the acute setting, identifying predictors of instability on CT scans could be very helpful in diagnosing these injuries. By comparing the CT findings to the stability of the injuries, we found that intraarticular fractures in the two lateral tarsometatarsal joints increased the risk of having an unstable Lisfranc injury. An avulsion fracture of the Lisfranc ligament (fleck sign) has in previous articles been interpreted as a sign of instability [17,19]. We were not able to correlate any other fracture pattern (including fleck sign) to the stability of the Lisfranc injuries.

Several authors have reported Lisfranc injuries to be more common in men compared to women [1,4,5,14,21,33]. In the current study the distribution between genders was equal, as also reported by both Crates et al. and Komenda et al. [15,25] We found, however, a higher proportion of unstable injuries in women. Also, women had a

Table 5
Lisfranc mortise measures related to instability.

	Lisfranc injuries		P-value	Gender		P-value
	Stable injuries	Unstable injuries		Male	Female	
Medial mortise depth (mm)						
Mean (SD)	7.3 (1.8)	6.6 (1.7)	0.072	7.3 (1.9)	6.6 (1.6)	0.057
Lateral mortise depth (mm)						
Mean (SD)	3.6 (1.5)	3.7 (1.2)	0.785	4.0 (1.5)	3.3 (1.1)	0.024
TMT-2 height (mm)						
Mean (SD)	21.2 (2.3)	20.1 (2.4)	0.036	21.6 (2.3)	19.8 (2.1)	0.001

TMT-2 height = tarsometatarsal joint 2 height.

Measurements are described in Fig. 2.

Statistical significant findings are marked with bold values.

decreased Lisfranc mortise depth and second tarsometatarsal joint height compared to the men. Peicha et al. have previously reported that a shallow medial mortise depth is a risk factor for Lisfranc injuries, and this is also supported by Yu-Kai et al., who observed that women had a more shallow medial mortise depth and a shorter height of the second metatarsal base than men [34,35]. As the lateral aspect of the medial cuneiform is the origin of the interosseous and plantar part of the Lisfranc ligament and the medial and plantar aspect of the second metatarsal base is the attachment area, one might speculate that the feet with a deeper medial mortise and a higher second tarsometatarsal joint might have a broader and stronger Lisfranc ligament, and thereby a decreased risk of obtaining an unstable Lisfranc injury [18].

The present study has some inherent weaknesses. First of all, a larger patient number would have increased study power. There is some degree of uncertainty regarding the epidemiological data, as some patients with Lisfranc injuries from the Oslo University Hospital population might have been treated elsewhere. This would lead to an underestimated injury incidence. Furthermore, we were not able to compare stress fluoroscopy with weightbearing radiographs, as most patients did not have weightbearing radiographs.

The strengths of the study include Oslo University Hospital being the primary trauma center in the region and Oslo A&E Department is the only public primary health care walk-in facility in Oslo. In addition, all patients were included in the study in a prospective manner when presenting with the injury, evaluated by an orthopaedic surgeon specialized in Foot and Ankle Surgery and a diagnostic algorithm was used. All patients, except one, had a CT scan of the injured foot. Over 90 percent of patients with stable injuries were followed up with weightbearing radiographs at 6 weeks, thereby any occult instability could be detected.

5. Conclusion

In the present study we observed a higher incidence of Lisfranc injuries than previously reported, and low-energy trauma was the most common mechanism of injury. Women had a shallower Lisfranc mortise than men and a higher proportion of unstable injuries. We also found that shorter second tarsometatarsal joint height and intraarticular fractures in the two lateral tarsometatarsal joints increased the risk of having an unstable Lisfranc injury.

Conflict of interest

The authors declare no potential conflicts of interests.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- [1] Goossens M, De Stoop N. Lisfranc's fracture-dislocations: etiology, radiology, and results of treatment. *Clin Orthop Relat Res* 1983.
- [2] Haapamäki V, Kiuru M, Koskinen S. Lisfranc fracture-dislocation in patients with multiple trauma: diagnosis with multidetector computed tomography. *Foot Ankle Int* 2004;25:614–9, doi: <http://dx.doi.org/10.1177/107110070402500903>.
- [3] Curtis MJ, Myerson M, Szura B. Tarsometatarsal joint injuries in the athlete. *Am J Sports Med* 1993;21:497–502, doi: <http://dx.doi.org/10.1177/036354659302100403>.
- [4] Myerson MS, Fisher RT, Burgess AR, Kenzora JE. Fracture dislocations of the tarsometatarsal joints: end results correlated with pathology and treatment. *Foot Ankle* 1986;6:225–42, doi: <http://dx.doi.org/10.1177/107110078600600504>.
- [5] Hardcastle P, Reschauer R, Kutscha-Lissberg E, Schoffmann W. Injuries to the tarsometatarsal joint. Incidence, classification and treatment. *J Bone Jt Surg* 1982;64-B:349–56.
- [6] Faciszewski T, Burks RT, Manaster BJ. Subtle injuries of the Lisfranc joint. *J Bone Jt Surg* 1990;72:1519–22, doi: <http://dx.doi.org/10.1016/j.jpcad.2015.11.006>.
- [7] Wilppula E. Tarsometatarsal fracture-dislocation. Late results in 26 patients. *Acta Orthop Scand* 1973;44:335–45.
- [8] Haapamäki VV, Kiuru MJ, Koskinen SK. Ankle and foot injuries: analysis of MDCT findings. *AJR Am J Roentgenol* 2004;183:615–22, doi: <http://dx.doi.org/10.2214/ajr.183.3.1830615>.
- [9] Raikin SM, Elias I, Dheer S, Besser MP, Morrison WB, Zoga AC. Prediction of midfoot instability in the subtle Lisfranc injury. Comparison of magnetic resonance imaging with intraoperative findings. *J Bone Joint Surg Am* 2009;91:892–9, doi: <http://dx.doi.org/10.2106/JBJS.H.01075>.
- [10] Vuori JP, Aro HT. Lisfranc joint injuries: trauma mechanisms and associated injuries. *J Trauma* 1993;35:40–5, doi: <http://dx.doi.org/10.1097/00005373-199307000-00007>.
- [11] Watson TS, Shurnas PS, Denker J. Treatment of Lisfranc joint injury: current concepts. *J Am Acad Orthop Surg* 2010;18:718–28.
- [12] Myerson MS. The diagnosis and treatment of injury to the tarsometatarsal joint complex. *J Bone Joint Surg Br* 1999;81:756–63, doi: <http://dx.doi.org/10.1302/0301-620X.81B5.10369>.
- [13] Stavlas P, Roberts CS, Xypnitos FN, Giannoudis PV. The role of reduction and internal fixation of Lisfranc fracture-dislocations: a systematic review of the literature. *Int Orthop* 2010;34:1083–91, doi: <http://dx.doi.org/10.1007/s00264-010-1101-x>.
- [14] Kuo RS, Tejwani NC, Digiovanni CW, Holt SK, Benirschke SK, Hansen ST, et al. Outcome after open reduction and internal fixation of Lisfranc joint injuries. *J Bone Joint Surg Am* 2000;82-A:1609–18.
- [15] Komenda GA, Myerson MS, Biddinger KR. Results of arthrodesis of the tarsometatarsal joints after traumatic injury. *J Bone Jt Surg* 1996;78:1665–76.
- [16] Myerson MS, Cerrato RA. Current management of tarsometatarsal injuries in the athlete. *J Bone Jt Surg* 2008;90-A:2522–33.
- [17] Renninger CH, Cochran G, Tompane T, Bellamy J, Kuhn K. Injury characteristics of low-energy Lisfranc injuries compared with high-energy injuries. *Foot Ankle Int* 2017;38:964–9, doi: <http://dx.doi.org/10.1177/1071100717709575>.
- [18] Solan MC, Moorman CT, Miyamoto RG, Jasper LE, Belkoff SM. Ligamentous restraints of the second tarsometatarsal joint: a biomechanical evaluation. *Foot Ankle Int* 2001;22:637–41.
- [19] Seybold JD, Coetzee JC. Lisfranc injuries: when to observe, fix, or fuse. *Clin Sports Med* 2015;34:705–23, doi: <http://dx.doi.org/10.1016/j.csm.2015.06.006>.
- [20] Nunley JA, Vertullo CJ. Classification, investigation, and management of midfoot sprains Lisfranc injuries in the athlete. *Am J Sports Med* 2002;30:871–8.
- [21] Arntz CT, Veith RG, Hansen ST, Holt SK, Benirschke SK, Hansen ST, et al. Fractures and fracture-dislocations of the tarsometatarsal joint. *J Bone Joint Surg Am* 1988;70:173–81, doi: <http://dx.doi.org/10.1097/00005373-197403000-00001>.
- [22] Kaar S, Femino J, Morag Y. Lisfranc joint displacement following sequential ligament sectioning. *J Bone Joint Surg Am* 2007;89:2225–32, doi: <http://dx.doi.org/10.2106/JBJS.F.00958>.
- [23] Coss HS, Manos RE, Buoncristiani A, Mills WJ. Abduction stress and AP weightbearing radiography of purely ligamentous injury in the tarsometatarsal joint. *Foot Ankle Int* 1998;19:537–41.
- [24] Shakked RJ. Lisfranc injury in the athlete. *JBJS Rev* 2017;5:e4, doi: <http://dx.doi.org/10.2106/JBJS.RVW.17.00025>.
- [25] Crates JM, Barber FA, Sanders EJ. Subtle Lisfranc subluxation: results of operative and nonoperative treatment. *J Foot Ankle Surg* 2015;54:350–5, doi: <http://dx.doi.org/10.1053/j.jfas.2014.07.015>.
- [26] Weatherford BM, Bohay DR, Anderson JG. Open reduction and internal fixation versus primary arthrodesis for Lisfranc injuries. *Foot Ankle Clin* 2017;22:1–14, doi: <http://dx.doi.org/10.1016/j.fcl.2016.09.002>.
- [27] English T. Dislocations of the metatarsal bone and adjacent toe. *J Bone Jt Surg* 1964;46B:700–4.
- [28] Aitken A, Poulson D. Dislocations of the tarsometatarsal joint. *J Bone Jt Surg* 1963;45-A:246–60.
- [29] Ponkilainen VT, Laine H-J, Mäenpää HM, Mattila VM, Haapasalo HH. Incidence and characteristics of midfoot injuries. *Foot Ankle Int* 2019;40(1):105–12, doi: <http://dx.doi.org/10.1177/1071100718799741>.
- [30] Philbin T, Rosenberg G, Sferia JJ. Complications of missed or untreated Lisfranc injuries. *Foot Ankle Clin* 2003;8:61–71, doi: [http://dx.doi.org/10.1016/S1083-7515\(03\)00003-2](http://dx.doi.org/10.1016/S1083-7515(03)00003-2).
- [31] Rammelt S, Schneiders W, Schikore H, Holch M, Heineck J, Zwipp H. Primary open reduction and fixation compared with delayed corrective arthrodesis in

- the treatment of tarsometatarsal (Lisfranc) fracture dislocations. *J Bone Joint Surg Br* 2008;2008(November): 1499.
- [32] Nunley JA, Vertullo CJ, Carolina N. Classification, investigation, and management of midfoot sprains Lisfranc injuries in the athlete. *Am J Sports Med* 2002;30:871–8.
- [33] Coetzee JC, Ly TV. Treatment of primarily ligamentous Lisfranc joint injuries: primary arthrodesis compared with open reduction and internal fixation. Surgical technique. *J Bone Joint Surg Am* 2007;89(Suppl 2):122–7, doi:<http://dx.doi.org/10.2106/JBJS.F.01004>.
- [34] Peicha G, Labovitz J, Seibert FJ, Grechenig W, Weiglein A, Preidler KW, et al. The anatomy of the joint as a risk factor for Lisfranc dislocation and fracture-dislocation. An anatomical and radiological case control study. *J Bone Joint Surg Br* 2002;84:981–5.
- [35] Yu-kai Y, Shiu-bii L. The journal of foot & ankle surgery anatomic parameters of the Lisfranc joint complex in a radiographic and cadaveric comparison. *J Foot Ankle Surg* 2015;54:883–7, doi:<http://dx.doi.org/10.1053/j.jfas.2015.02.021>.

Outcome after nonoperative treatment of stable Lisfranc injuries.

A prospective cohort study.

Authors:

Are H. Stødle, MD^{1,3}

Kjetil H. Hvaal, MD, PhD¹

Jan Erik Madsen, MD, PhD^{1,3}

Helga M. Brøgger, MD²

Elisabeth Ellingsen Husebye, MD, PhD¹

¹Division of Orthopaedic Surgery, Oslo University Hospital, Norway

²Department of Radiology and Nuclear medicine, Oslo University Hospital, Norway

³Institute of Clinical Medicine, University of Oslo, Norway

Corresponding author:

Are H. Stødle

Division of Orthopaedic Surgery

Oslo University Hospital

Pb 4950 Nydalen

0424 Oslo

ABSTRACT

Background

Unstable Lisfranc injuries are best treated by anatomic reduction and stable fixation, while the stable injuries can be treated nonoperatively. The aim of this study was to evaluate the outcome after nondisplaced and stable Lisfranc injuries.

Methods

26 patients with injuries to the Lisfranc joint complex detected on CT scans, but without displacement were tested to be stable using a fluoroscopic stress test and prospectively included in the study. The patients were immobilized in a non-weightbearing short leg cast for 6 weeks. Weightbearing radiographs were obtained at 6, 12 and 52 weeks. The final follow-up was 55 (IQR 53 - 60) months after injury. The main outcome was the American Foot and Ankle Society midfoot score (AOFAS score). Secondary outcomes were visual analog scale for pain (VAS pain), 36-Item Short Form Health Survey (SF-36) and radiologically detected osteoarthritis.

Results

All the Lisfranc injuries were confirmed to be stable on follow-up weightbearing radiographs at a minimum of 3 months after injury. Median AOFAS score at 1-year follow-up was 89 (IQR 84 - 97) and at final follow-up 100 (IQR 90 - 100); The AOFAS score continued to improve after 1-year ($P=.005$). The median VAS pain score both at rest and during walking were 0 (IQR 0 - 0) at the final follow-up. The SF-36 scores at 1-year follow up were comparable to general population with a median physical component summary (PCS) score of 51.8 (IQR 46.0 - 54.6) and a mental component summary (MCS) score of 52.9 (IQR 40.9 - 58.4). One patient had radiological signs of osteoarthritis at 1-year follow-up.

Conclusion

Stable Lisfranc injuries treated nonoperatively had an excellent outcome in this study with a median follow-up of 55 months. The AOFAS score continued to improve after 1 year.

INTRODUCTION

The Lisfranc joint complex comprise of the tarsometatarsal, intercuneiform and naviculocuneiform joints with their associated ligaments.^{22,36} Lisfranc injuries can vary from severe fracture-dislocations to stable nondisplaced injuries.^{7,20,24,32,39,40} The incidence is previously reported to be 1/60 000 person/years, but recent studies report a higher incidence ranging from 9.2-14/100 000 person/years.^{12,25,39} Stødle et al. reported the incidence of stable Lisfranc injuries to be 8/100 000 person/years.³⁹ Up to 24% of Lisfranc injuries have been reported to be missed on initial radiographs.^{11,26,39} The increase in reported incidence seems to be related to an increased awareness of this entity, together with more advanced diagnostic tools such as CT scan, MRI, weightbearing radiographs and fluoroscopic stress test.^{25,39,40}

Several studies have shown operative treatment with anatomic reduction and stable fixation to be the most important factors in achieving a favorable outcome in the unstable Lisfranc injuries.^{1,16,17,22} In the nondisplaced and stable Lisfranc injuries nonoperative treatment is recommended, and good outcome scores have been reported.^{3,6,24} A stable Lisfranc injury has been defined as an injury to the Lisfranc joint complex with less displacement than 2 mm on weightbearing radiographs and/or no obvious displacement on stress fluoroscopy, when compared to the non-injured foot.^{2,4,7,28,39,40}

The aim of the present study was to evaluate the outcome after nonoperative treatment of stable Lisfranc injuries.

Material and Method

A prospective cohort study including stable Lisfranc injuries was conducted at Oslo University Hospital (a level one trauma center) and Oslo Accident and Emergency Department. The patients in the present study were also included in a previously published study reporting on incidence, mechanism of injury and predictors of instability.³⁹ The study was approved by The Regional Committee for Medical and Health Research (2014/849/REK) and approved by the data protection officer at the university hospital. An informed consent form was signed by the patients prior to enrollment. Patients presenting with an isolated stable Lisfranc injury between September 1, 2014 and August 31, 2015 were invited to participate.

An acute, stable Lisfranc injury was defined as:

- acute trauma (Lisfranc injury diagnosed within 4 weeks)
- midfoot pain on weightbearing and manipulation of the midfoot
- injury to the tarsometatarsal joint line with intraarticular or avulsion fractures, but absence of displacement of 2mm or more on radiographs or CT scans, or obvious displacement on stress test under fluoroscopy.

Exclusion criteria were:

- Unstable Lisfranc injuries
- Concomitant lower extremity injuries
- Charcot arthropathy
- Isolated fifth metatarsal fracture
- Non-compliant patient
- Patient not available for follow-up

Patient characteristics and mechanism of injury were registered at enrollment in the study. The patients had their injured foot tested for occult instability using a fluoroscopic stress test 7-14 days after injury. The opposite non-injured foot was tested for comparison. The stress test was positive if joint displacement was detected.

The patients were immobilized in a nonweightbearing short leg cast for 6 weeks. If no pain on weightbearing at 6 weeks, they started full weightbearing in normal shoes, whereas if pain was present on weightbearing they were offered a Walker boot with full weightbearing for the next 4 weeks. Weightbearing AP, oblique and lateral radiographs of both the injured and non-injured foot were obtained at 6 weeks, 3 months and 12 months follow-up. 4-5 years after injury the patients were interviewed by phone and invited to a final follow-up.

Radiographs and CT scans were evaluated using Syngo Studio VB36E (Siemens Healthcare GmbH, Erlangen, Germany). Two orthopaedic surgeons specialized in foot and ankle surgery and one radiology consultant experienced in musculoskeletal imaging evaluated the images.

Injuries to the tarsometatarsal joints were categorized as avulsion fractures or intraarticular fractures and located to the medial, middle and lateral column according to the columnar theory as described by Chiodo and Myerson.⁴

The primary outcome measure was the American Orthopaedic Foot and Ankle Society midfoot score (AOFAS score), consisting of three main components (pain, function, and alignment). The score ranges from 0 to 100 with the best score being 100.¹⁵ The secondary outcome measures were the 36-Item Short Form Health Survey (SF-36); a patient-reported survey of patient health, and the visual analog scale for pain (VAS pain) at rest and during walking.

Patients not able to attend final follow-up were interviewed by phone and the AOFAS score, Numeric rating scale (NRS) for pain at rest and during walking, as well as return to activities were addressed. With regards to gait abnormality and alignment part of the AOFAS score, the patients interviewed by phone were asked about any gait abnormality or if the alignment of their foot had changed (compared to the non-injured foot). The NRS pain has been shown to highly correlate to the VAS pain.¹³

Secondary displacement and signs of posttraumatic osteoarthritis (OA) were evaluated on the radiographs obtained at the follow-ups. Radiographs were not routinely obtained at the final follow-up.

Statistics

Statistical analyses were conducted using SPSS version 26 (SPSS Inc, IBM, Chicago, IL). Data were tested for normality and normally distributed data are presented with a mean value and standard deviation (SD). The one sample t-test was used for statistical analysis. Non-parametric data are presented with median values and the interquartile range (IQR) and the Wilcoxon signed rank test was used for statistical analysis. The significance level was set to $P < .05$.

When examining the change in AOFAS score over time, a one-sample T-test was conducted to determine whether the average slope (change in AOFAS score per one-month time unit) was different from zero (no change). To be able to evaluate change in AOFAS score over the whole study period, only patients who had AOFAS score measured at 3 and/or 12 months and at the final follow-up were included. The slope was computed between either “final follow-up - 3 months”, or “final follow-up - 1 year”, depending on what data was available for each subject and then divided by the number of months occurring between these two measurements.

Results

Twenty-six patients with isolated stable Lisfranc injuries were included in the study. The patient characteristics at enrollment and mechanisms of injury are presented in Table 1 a and b. A plantar ecchymosis was registered in 20/26 patients.

Twenty-three of the patients had initial nonweightbearing radiographs and nine of these radiographs were evaluated as normal. CT scans were obtained of all injured feet prior to inclusion, all of which showed avulsion fractures or intraarticular fractures in the TMT joint line,

indicating a Lisfranc injury. No displacement of 2mm or more were detected in any joints of the Lisfranc joint complex. Radiological findings are presented in Table 2.

A stress test under fluoroscopy was performed in 25 of the 26 patients, at median 10 (IQR 8-14) days after injury, to evaluate stability. All stress tests were negative. One patient did not have a stress test (an 80-years old female). On the follow-up weightbearing radiographs, none of the patients had any subsequent displacement in the Lisfranc joint complex. When comparing the injured foot to the non-injured foot, none of the 26 patients had a difference of 2 mm or more in the C1-M2 interval or the M1-M2 interval on the follow-up weightbearing radiographs at 6, 12 and 52 weeks. Only one patient had radiological signs of degenerative joint disease at the 1-year follow-up, the joint degeneration was located to the TMT-1 joint where the patient had an intraarticular fracture on presentation. The patient presented no symptoms from the injured foot at the final follow-up.

The AOFAS score, VAS pain scores and SF-36 are listed in Tables 3 and 4.

14/26 patients attended the final follow-up, eight patients were interviewed by phone and four patients were lost to follow-up. Median follow-up time was 55 (IQR 53 – 60) months. Of the eight patients interviewed by phone, seven patients reported no pain or discomfort from their injured foot. One patient reported mild pain at rest (NRS 1/10) and during walking (NRS 1/10). None of the patients interviewed by phone on the final follow-up reported limitations during activities, walking distance, footwear or walking surface. No patients reported any gait

abnormality or were able to detect any change in appearance or alignment of their injured foot compared to the non-injured foot.

Twenty patients had at least one earlier AOFAS score in addition to the AOFAS score at the final follow-up. The scores from these patients were used to calculate the slopes of the AOFAS score over time. The slopes were normally distributed as assessed by Shapiro-Wilk's test of normality ($p=0.228$). Mean slope (0.24, SD = 0.17) was statistically different from zero, $t(19)= 6.57$, $p<.001$, indicating a positive change in AOFAS score over time. With regard to the time period from 1 year after injury until the final follow-up (median 55 months), there was also an improvement of the AOFAS score (median 89 vs 100, $P=.005$) indicating a positive change in the AOFAS score between 1 year and final follow-up.

None of the patients reported work-related limitations due to their Lisfranc injury at final follow-up, but 2/22 patients reported some limitations in recreational sports activities (skateboarding and skiing). None of the patients had undergone any surgeries to their foot at final follow-up.

Discussion

The outcome of nonoperatively treated stable Lisfranc injuries in 26 patients are presented in this study. All injuries had CT verified avulsion fractures or intraarticular fracture in the tarsometatarsal joint line. The stability was verified using fluoroscopic stress test and follow-up weight bearing radiographs at a minimum of 3 months after injury. The outcome after a median follow-up time of 55 (IQR 53 – 60) months was excellent with a median AOFAS midfoot score of

100 (IQR 90-100), improving from 89 (IQR 84-97) at 1 year ($P = .005$). Osteoarthritis does not seem to be frequent after a stable Lisfranc injury as degenerative joint disease was only detected radiologically at 1-year in one patient. Only 2/22 patients reported limitations in recreational sports activity caused by their Lisfranc injury at the final follow-up. Stable Lisfranc injuries are easily overlooked as 9/23 injuries were not detected on initial nonweightbearing radiographs.

Most previous papers on Lisfranc injuries focus on the unstable fracture-dislocations and only few report on the stable and nondisplaced Lisfranc injuries.^{3,6,7,24} The stable injuries have been reported to have good outcomes when being treated nonoperatively, while the unstable injuries should be treated surgically.^{4,9,21,24,31,40} One of the most challenging issues concerning Lisfranc injuries is to adequately evaluate stability of the nondisplaced Lisfranc injuries and thereby properly select the right patients for nonoperative treatment. To detect occult instability in the nondisplaced Lisfranc injuries, the feet can be examined using either weightbearing radiographs or a fluoroscopic stress test.^{2,5,14,21,24} MRI has also been shown to have a high accuracy in detecting instability in ligamentous Lisfranc injuries.^{28,30} When evaluating both weightbearing radiographs and fluoroscopic stress tests the injured foot should be compared to the non-injured foot, as the M1-M2 and C1-M2 distances show variance between individuals and also between the nonweightbearing and weightbearing state.^{9,21,24,28,36,40} The fluoroscopic stress test has been criticized for lack of reliability in detecting unstable Lisfranc injuries, even though it is reported to be more sensitive compared to weightbearing radiographs in cadaver studies.^{5,14,23,33}

In a study by Chen et al. on minimally displaced Lisfranc injuries evaluated by weightbearing radiographs at presentation, 14/26 patients showed subsequent displacement at follow-up, where 9/14 were detected within 14 days.³ Preidler et al. showed that obtaining weightbearing radiographs on the day of injury did not add any information to nonweightbearing radiographs.²⁹ This emphasizes that the assessment of stability in these nondisplaced injuries is difficult, especially during the first days after injury. Hence, testing the stability of the midfoot should be postponed until 10-14 days, when pain is reduced.^{3,4,21,39} The patients should also be reassessed with weightbearing radiographs at follow-ups to detect any subsequent displacement.^{3,21} In the present study, the Lisfranc injuries were tested for instability using a fluoroscopic stress test at a median of 10 (IQR 8, 14) days. Weightbearing radiographs were obtained at 6 and 12 weeks and 1 year. None of the injured feet showed any subsequent displacement.

Although the stable Lisfranc injuries are uniformly recommended to be treated conservatively, only few authors have reported on the outcome of nondisplaced Lisfranc injuries after evaluating the stability using weightbearing radiographs or fluoroscopic stress tests.^{3,6,7,24}

Nunley et al. reported excellent outcome after Lisfranc injuries that were tested to be stable using weightbearing radiographs.²⁴ In their study on minimally displaced Lisfranc injuries, Chen et al. reported on 12/26 patients with a stable Lisfranc injury treated nonoperatively.³ The patient reported AOFAS score at mean follow-up of 54 months was 78.0 (95% CI, 68.6 – 87.4) and Manchester-Oxford Foot Questionnaire (MOXFQ) 24.8 (95% CI, 11.1 – 38.5). The AOFAS

score was poorer than the excellent results we present at a median of 55 months, median 100 (IQR 90 – 100). This may be explained by a difference in nature of the Lisfranc injuries included in the two studies. In Chen's study, the inclusion criteria were less than 2mm diastasis between the bases of the first and second metatarsal (M1-M2) and no displacement in the third, fourth and fifth tarsometatarsal joint. Thus, an increased diastasis between the medial cuneiform and the base of the second metatarsal (C1-M2) in addition to displacement of the first and second tarsometatarsal joint could be present. In our study only injuries with less than 2mm difference in the C1-M2 interval and the M1-M2 interval, and no displacement in any tarsometatarsal joints were included. Crates et al. reported the outcomes of 36 patients with subtle Lisfranc injuries, including patients with 2 mm or less diastasis in the M1-M2 interval on weightbearing radiographs. 16/36 patients were treated successfully nonoperatively, while 20 patients failed nonoperative treatment. Failed nonoperative treatment were based on judgement by the senior author, and neither timing of initial weightbearing radiographs , nor the time from injury to failed nonoperative treatment or the use of weightbearing radiographs on follow-up were specified.⁶ These factors and the poor AOFAS score in the group who failed nonoperative treatment (mean AOFAS score of 63.5) compared to the group with successful nonoperative treatment (mean AOFAS score = 90.2, calculated from the subgroups reported) suggest that the injuries failing nonoperative treatment were in fact unstable injuries. The AOFAS score of the patients treated successfully with nonoperative treatment on the other hand, corresponds to the AOFAS score of the stable Lisfranc injuries that we present.

We believe that subtle displacement on weightbearing radiographs indicating an unstable Lisfranc injury is best detected by an increase in the C1-M2 interval (where the Lisfranc

ligament originates and inserts). In contrast to measuring the M1-M2 interval, where the first metatarsal can displace laterally together with the second metatarsal, a disruption in the Lisfranc ligament might not be detected.

The AOFAS score in the present study continued to improve even after 1 year (median 89 versus 100, $P = .005$), implying that the rehabilitation period can be prolonged over years.

The VAS pain score at rest and during walking at the final follow-up in the current study were both 0 (IQR: 0-0), corresponding well with an excellent outcome. Also, the SF-36 summary scores (PCS 51.8 and MCS 52.9) at the 1-year follow-up were equivalent to the scores of the general population (40-49 years old, PCS mean 50.8 (SD 9.1) and MCS mean 52.6 (SD 9.1)) indicating return to the same quality of life as the Norwegian general population.¹⁰ Only 2/22 patients reported limitations in recreational sports activities (skiing and skateboarding) at the final follow-up. We have not encountered any other study reporting VAS pain score or SF-36 after stable Lisfranc injuries.

A high incidence of radiographic post traumatic OA is reported in patients treated for unstable Lisfranc injuries.^{8,16,19,37,38} The frequency of post traumatic OA after stable Lisfranc injuries is largely unknown. Chen et al. reported on twelve stable Lisfranc injuries treated conservatively with a follow-up of 54 months and follow-up radiographs at minimum 24 weeks; None of the stable injuries showed signs of degenerative joint disease.³ In the current study only one patient had radiological signs of posttraumatic OA at 1 year and no patients had clinical signs of OA at 5 years.

Several previous studies have shown that Lisfranc injuries are easily missed (up to 28%) on initial nonweightbearing radiographs.^{11,26,34,39} In the present study 9/23 of the initial nonweightbearing radiographs were evaluated as normal. This demonstrates, not surprisingly, that non-displaced Lisfranc injuries are more difficult to detect on initial radiographs than displaced Lisfranc injuries, as also pointed out by Ponkilainen et al.²⁶

The present study has several strengths, firstly the prospective design with a median follow-up of 55 months. All, but one patient, were tested for stability and all patients had weightbearing radiographs obtained at a minimum of 3 months. The injuries were all assessed using CT scans to verify injury to the Lisfranc joint complex (fractures, including avulsion fractures) and rule out any subtle displacements often not detected on conventional radiographs.¹⁸ All patients followed the same nonoperative treatment protocol. Although the study included only 26 patients, it is to our knowledge the largest study on stable Lisfranc injuries treated nonoperatively throughout the entire study period.

The study also has some inherent weaknesses. Only 14 patients were able to come to the outpatient clinic for the final follow-up, further 8 patients were interviewed by phone and 4 patients were lost to follow-up. As 8 patients only were interviewed by phone at the final follow-up, any gait abnormality or malalignment not detected by the patients themselves might have been missed, even though this was not detected in these patients on previous follow-ups. The patients did not have radiographs obtained at final follow-up and any radiological OA without clinical manifestation at that point was thereby not detected. The AOFAS midfoot scale

has been shown to have a ceiling effect and therefore a limited ability to differentiate between patients with high outcome scores.²⁷ We used fluoroscopic stress test for the initial evaluation of stability of non-displaced Lisfranc injuries and weightbearing radiographs at follow-ups. In the future, weightbearing CT scans might be an even more precise method to detect any occult instability.³⁵

Conclusion:

Stable Lisfranc injuries treated nonoperatively had an excellent outcome 55 months after injury in this prospective cohort study. The AOFAS midfoot score continued to improve even after 1 year. Assessment of stability in Lisfranc injuries should preferably be done 7-14 days after the injury. None of the patients had clinical symptoms of posttraumatic osteoarthritis at the final follow-up, one patient had radiologically detected degenerative joint disease at the 1-year follow-up.

Table 1a

Patient characteristics	
Number of patients	26
Gender (Male/Female)	15/11
Side (Right/Left)	14/12
Age (mean, SD)	40.0 (15.7)
Time injury-diagnosis in days (mean, SD)	2.2 (5.7)
Plantar ecchymosis	20

Caption: Patient characteristics at time of inclusion.

Table 1b

Mechanism of injury	
Fall from own height /twisting injury of foot	7
Fall down stairs	1
Bike accident	2
Kicked into an object	1
Sports related injury	4
Motor vehicle accident	2
Fall > 3 meters	2
Crush injury	7

Table 2

Radiological findings	
Initial non-WB radiographs (pos /neg)*	14/9
Positive CT scans**	26
Medial column	
Negative	8
Avulsion fx	10
Intraarticular fx	8
Middle column	
Negative	5
Avulsion fx	5
Intraarticular fx	13
Lateral column	
Negative	15
Avulsion fx	1
Intraarticular fx	7

Caption:

* Positive = Fractures detected on initial nonweightbearing radiographs

** Intraarticular fractures or avulsion fractures TMT joints

WB= weightbearing. Fx= fracture

Table 3

	3 months (19 patients)	1 year (20 patients)	Final FU* (22 patients**)
AOFAS midfoot score	85 (77-87)	89 (84-97)	100 (90-100)
VAS pain score			
At rest	0 (0-2)	0 (0-1.1)	0 (0-0)
During walking	3 (2-5)	0.5 (0-2.4)	0 (0-0)

Caption:

American Foot and Ankle Society (AOFAS) midfoot score and the visual analog scale for pain.

(VAS pain). The median and interquartile range (IQR) are reported.

*Final follow-up (FU) = Median 55 (IQR 53-60) months

**8 of the 22 patients were interview by phone

Table 4

SF-36	1 year (20 patients)
Physical function	95 (86.3-100)
Role physical	100 (75-100)
Bodily pain	78.8 (60-90)
General health	70 (57.5-80)
Vitality	65 (47.5-70)
Social function	100 (78.1-100)
Role emotional	100 (41.7-100)
Mental health	80 (66-88)
Physical component summary (PCS)	51.8 (46-54.6)
Mental component summary (MCS)	52.9 (40.9-58.4)

Caption:

36-Item Short Form Health Survey (SF-36) score at 1 year. The median and interquartile range

(IQR) are reported.

Figure 1 a, b, c,

Caption:

36 years old male who fell during jogging, was unable to weightbear on his forefoot and had midfoot pain and plantar ecchymosis. A CT scan revealed a nondisplaced intraarticular fracture at the base of the 2nd metatarsal (1a and b). A dorsolateral avulsion fracture of the third cuneiform and extraarticular nondisplaced fractures of the proximal third and fourth metatarsal was present (not shown). The foot was stable when stress tested under fluoroscopy. No displacements were detected on weightbearing radiographs of both feet at the 3-month follow-up (1c).

References

1. Arntz CT, Veith RG, Hansen ST, et al. Fractures and fracture-dislocations of the tarsometatarsal joint. *J Bone Joint Surg Am.* 1988;70(2):173–81.
2. Aronow MS. Treatment of the missed Lisfranc injury. *Foot Ankle Clin.* 2006;11(1):127–142, ix.
3. Chen P, Ng N, Snowden G, Mackenzie SP, Nicholson JA, Amin AK. Rates of Displacement and Patient-Reported Outcomes Following Conservative Treatment of Minimally Displaced Lisfranc Injury. *Foot Ankle Int.* 2020;41(4):387–391.
4. Chiodo CP, Myerson MS. Developments and advances in the diagnosis and treatment of

- injuries to the tarsometatarsal joint. *Orthop Clin North Am*. 2001;32(1):11–20.
5. Coss HS, Manos RE, Buoncristiani a, Mills WJ. Abduction stress and AP weightbearing radiography of purely ligamentous injury in the tarsometatarsal joint. *Foot Ankle Int*. 1998;19(8):537–41.
 6. Crates JM, Barber FA, Sanders EJ. Subtle lisfranc subluxation: results of operative and nonoperative treatment. *J Foot Ankle Surg*. 2015;54(3):350–5.
 7. Curtis MJ, Myerson M, Szura B. Tarsometatarsal joint injuries in the athlete. *Am J Sports Med*. 1993;21(4):497–502.
 8. Dubois-Ferrière V, Lübbecke A, Chowdhary A, Stern R, Dominguez D, Assal M. Clinical Outcomes and Development of Symptomatic Osteoarthritis 2 to 24 Years After Surgical Treatment of Tarsometatarsal Joint Complex Injuries. *J Bone Joint Surg Am*. 2016;98(9):713–20.
 9. Eleftheriou KI, Rosenfeld PF, Calder JDF. Lisfranc injuries: an update. *Knee Surg Sports Traumatol Arthrosc*. 2013;21(6):1434–46.
 10. Garratt AM, Stavem K. Measurement properties and normative data for the Norwegian SF-36: Results from a general population survey. *Health Qual Life Outcomes*. 2017;15(1):1–10.
 11. Haapamaki V, Kiuru M, Koskinen S. Lisfranc Fracture-dislocation in Patients with Multiple Trauma: Diagnosis with Multidetector Computed Tomography. *Foot Ankle Int*. 2004;25(9):614–619.
 12. Hardcastle P, Reschauer R, Kutscha-Lissberg E, Schoffmann W. Injuries to the tarsometatarsal joint. Incidence, classification and treatment. *J Bone Jt Surg*. 1982;64-

B(3):349–356.

13. Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain: Visual Analog Scale for Pain (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF-36 BPS), and Measure of Intermittent and Constant Osteoarthritis Pain (ICOAP). *Arthritis Care Res.* 2011;63(SUPPL. 11).
14. Kaar S, Femino J, Morag Y. Lisfranc joint displacement following sequential ligament sectioning. *J Bone Joint Surg Am.* 2007;89(10):2225–32.
15. Kitaoka HB, Alexander IJ, Adelaar RS, Nunley JA, Myerson MS, Sanders M. Clinical Rating Systems for the Ankle-Hindfoot, Midfoot, Hallux, and Lesser Toes. *Foot Ankle Int.* 1994;15(7):349–353.
16. Kuo RS, Tejwani NC, Digiovanni CW, et al. Outcome After Open Reduction and Internal Fixation of Lisfranc Joint Injuries. *J Bone Jt Surg.* 2000;82-A(11):1609–1618.
17. Lau S, Howells N, Millar M, De Villiers D, Joseph S, Oppy A. Plates, Screws, or Combination? Radiologic Outcomes After Lisfranc Fracture Dislocation. *J Foot Ankle Surg.* 2016;55(4):799–802.
18. Lu J, Ebraheim NA, Skie M, Porshinsky B, Yeasting RA. Radiographic and computed tomographic evaluation of Lisfranc dislocation: a cadaver study. *Foot ankle Int / Am Orthop Foot Ankle Soc [and] Swiss Foot Ankle Soc.* 1997;18:351–355.
19. Mulier T, Reynders P, Dereymaeker G, Broos P. Severe Lisfrancs injuries: primary arthrodesis or ORIF? *Foot ankle Int / Am Orthop Foot Ankle Soc [and] Swiss Foot Ankle Soc.* 2002;23(10):902–905.

20. Myerson MS. The diagnosis and treatment of injury to the tarsometatarsal joint complex. *J Bone Jt Surg - Ser B*. 1999;81(5):756–763.
21. Myerson MS, Cerrato R. Current management of tarsometatarsal injuries in the athlete. *Instr Course Lect*. 2009;58:583–594.
22. Myerson MS, Fisher RT, Burgess AR, Kenzora JE. Fracture dislocations of the tarsometatarsal joints: end results correlated with pathology and treatment. *Foot Ankle*. 1986;6:225–242.
23. Naguib S, Meyr AJ. Reliability, Surgeon Preferences, and Eye-Tracking Assessment of the Stress Examination of the Tarsometatarsal (Lisfranc) Joint Complex. *J Foot Ankle Surg*. 2019;58(1):93–96.
24. Nunley JA, Vertullo CJ. Classification , Investigation , and Management of Midfoot Sprains Lisfranc Injuries in the Athlete. *Am J Sports Med*. 2002;30(6):871–878.
25. Ponkilainen VT, Laine HJ, Mäenpää HM, Mattila VM, Haapasalo HH. Incidence and Characteristics of Midfoot Injuries. *Foot Ankle Int*. 2019;40(1):105–112.
26. Ponkilainen VT, Partio N, Salonen EE, et al. Inter- and intraobserver reliability of non-weight-bearing foot radiographs compared with CT in Lisfranc injuries. *Arch Orthop Trauma Surg*. 2020.
27. Ponkilainen VT, Uimonen M, Repo JP, Mattila VM, Haapasalo HH. Validity and internal consistency of the American Orthopaedic Foot and Ankle Society Midfoot Scale in patients with Lisfranc injury. *Foot Ankle Surg*. 2020;26(5):523–529.
28. Potter HG, Deland JT, Gusmer PB, Carson E, Warren RF. Magnetic resonance imaging of the Lisfranc ligament of the foot. *Foot Ankle Int*. 1998;19(7):438–446.

29. Preidler KW, Peicha G, Lajtai G, et al. Conventional radiography, CT, and MR imaging in patients with hyperflexion injuries of the foot: Diagnostic accuracy in the detection of bony and ligamentous changes. *Am J Roentgenol*. 1999;173(6):1673–1677.
30. Raikin SM, Elias I, Dheer S, Besser MP, Morrison WB, Zoga AC. Prediction of midfoot instability in the subtle Lisfranc injury: Comparison of magnetic resonance imaging with intraoperative findings. *J Bone Jt Surg - Ser A*. 2009;91(4):892–899.
31. Robertson GAJ, Ang KK, Maffulli N, Keenan G, Wood AMD. Return to sport following Lisfranc injuries: A systematic review and meta-analysis. *Foot Ankle Surg*. 2019;25(5):654–664.
32. Scolaro J, Ahn J, Mehta S. Lisfranc fracture dislocations. *Clin Orthop Relat Res*. 2011;469(7):2078–80.
33. Shakked RJ. Lisfranc injury in the athlete. *JBJS Rev*. 2017;5(9):e4.
34. Sherief TI, Mucci B, Greiss M. Lisfranc injury: how frequently does it get missed? And how can we improve? *Injury*. 2007;38(7):856–60.
35. Sripanich Y, Steadman J, Krähenbühl N, et al. Asymmetric lambda sign of the second tarsometatarsal joint on axial weight-bearing cone-beam CT scans of the foot: preliminary investigation for diagnosis of subtle ligamentous Lisfranc injuries in a cadaveric model. *Skeletal Radiol*. 2020;49(10):1615–1621.
36. Sripanich Y, Weinberg MW, Krähenbühl N, et al. Imaging in Lisfranc injury: a systematic literature review. *Skeletal Radiol*. 2001;49(1):31–53.
37. Stavlas P, Roberts CS, Xypnitos FN, Giannoudis P V. The role of reduction and internal fixation of Lisfranc fracture-dislocations: A systematic review of the literature. *Int Orthop*.

2010;34(8):1083–1091.

38. Stødle AH, Hvaal KH, Brøgger HM, Madsen JE, Husebye EE. Temporary Bridge Plating vs Primary Arthrodesis of the First Tarsometatarsal Joint in Lisfranc Injuries: Randomized Controlled Trial. *Foot Ankle Int.* 2020;41(8):901–910.
39. Stødle AH, Hvaal KH, Enger M, Brøgger H, Madsen JE, Ellingsen Husebye E. Lisfranc injuries: Incidence, mechanisms of injury and predictors of instability. *Foot Ankle Surg.* 2020;26(5):535–540.
40. Watson TS, Shurnas PS, Denker J. Treatment of Lisfranc joint injury: current concepts. *J Am Acad Orthop Surg.* 2010;18(12):718–728.

Temporary Bridge Plating vs Primary Arthrodesis of the First Tarsometatarsal Joint in Lisfranc Injuries: Randomized Controlled Trial

Foot & Ankle International®
1–10

© The Author(s) 2020

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/1071100720925815

journals.sagepub.com/home/fai

Are H. Stødle, MD^{1,3} , Kjetil H. Hvaal, MD, PhD¹, Helga M. Brøgger, MD², Jan Erik Madsen, MD, PhD^{1,3} , and Elisabeth Ellingsen Husebye, MD, PhD¹

Abstract

Background: Unstable Lisfranc injuries are best treated with anatomic reduction and stable fixation. There are controversies regarding which type of stabilization is best. In the present study, we compared primary arthrodesis of the first tarsometatarsal (TMT) joint to temporary bridge plating in unstable Lisfranc injuries.

Methods: Forty-eight patients with Lisfranc injuries were included and followed for 2 years. Twenty-four patients were randomized to primary arthrodesis (PA) of the medial 3 TMT joints, whereas 24 patients were randomized to temporary bridge plate (BP) over the first TMT joint and primary arthrodesis of the second and third TMT joints. The main outcome parameter was the American Orthopaedic Foot & Ankle Society (AOFAS) midfoot scale and the secondary outcome parameters were the 36-Item Short Form Health Survey (SF-36) and visual analog scale for pain (VAS pain). Computed tomography (CT) scans pre- and postoperatively were obtained. Radiographs were obtained at follow-ups. Pedobarographic examination was performed at the 2-year follow-up. Twenty-two of 24 patients in the PA and 23/24 in the BP group completed the 2-year follow-up.

Results: The mean AOFAS midfoot score 2 years postoperatively was 89 (SD 9) in the PA group and 85 (SD 15) in the BP group ($P = .32$). There were no significant differences between the groups with regard to SF-36 or VAS pain scores. The alignment of the first metatarsal was better in the BP group than in the PA group measured by the anteroposterior Meary angle ($P = .04$). The PA group had a reduced peak pressure under the fifth metatarsal ($P = .047$). In the BP group, 11/24 patients had radiologic signs of osteoarthritis in the first TMT joint.

Conclusion: Both treatment groups had good outcome scores. The first metatarsal was better aligned in the BP group; however, there was a high incidence of radiographic osteoarthritis in this group.

Level of Evidence: Therapeutic level I, prospective randomized controlled study.

Keywords: Lisfranc injury, tarsometatarsal joint injury, primary arthrodesis, bridge plate

Introduction

Lisfranc injuries are defined as injuries to the tarsometatarsal joint complex, which includes the tarsometatarsal (TMT) joints, intercuneiform and naviculocuneiform joints.^{16,21} These injuries consist of a wide spectrum of injuries from nondisplaced, stable injuries to severe fracture dislocations.^{6,21,24,31,32,35} Several studies have shown anatomic reduction and stable fixation to be the most important factors in achieving a good functional outcome in acute Lisfranc fracture dislocations.^{2,5,14,17,23,28,36} The best technique used to achieve an anatomic and stable fixation of the joints is still debated. Open reduction and transarticular screw fixation has been the standard approach for many

years.^{2,3,14,21,32,36} In recent years, good results have been reported on dorsal bridge plating of the TMT joints after Lisfranc injuries.^{1,9,11,17,33} In contrast to transarticular screw fixation, the dorsal bridge plate technique avoids additional

¹Division of Orthopaedic Surgery, Oslo University Hospital, Oslo, Norway

²Department of Radiology and Nuclear medicine, Oslo University Hospital, Oslo, Norway

³Institute of Clinical Medicine, University of Oslo, Oslo, Norway

Corresponding Author:

Are H. Stødle, MD, Division of Orthopaedic Surgery, Oslo University Hospital, Pb 4950 Nydalen, Oslo, 0424, Norway.
Email: arhauk@ous-hf.no

damage to the cartilage and might reduce the high incidence of post-traumatic osteoarthritis in these patients.^{1,9,11,14,20,25,32} Primary arthrodesis of the 3 medial TMT joints has also been advocated in treating unstable Lisfranc injuries. Two randomized controlled trials comparing transarticular screw fixation to primary arthrodesis in high-energy Lisfranc injuries have shown favorable results in the primary arthrodesis group.^{8,18} In addition, one retrospective study comparing primary arthrodesis to transarticular screw fixation in young patients with low-energy Lisfranc injuries also showed favorable results in the arthrodesis group.⁴

In the present study, we have compared the first TMT joint-preserving technique using a temporary dorsal bridge plate, to primary arthrodesis in acute Lisfranc injuries. A primary arthrodesis of the second and third TMT joint was performed in all patients.

Our hypothesis was that the joint-sparing procedure using a temporary dorsal bridge plate would lead to a better functional outcome compared to primary arthrodesis.

Methods

A randomized controlled trial comparing primary arthrodesis (PA) to temporary bridge plating (BP) of the first TMT joint in acute Lisfranc injuries was conducted (Figure 1). The study was approved by the National Committee for Medical and Health Research, registered in Clinical trials.org (ID: NCT01448941), and approved by the data protection officer at the university hospital. All patients signed an informed consent form prior to inclusion. The patients were enrolled from October 2011 to August 2015 (Figure 1). The trial was performed at a level 1 trauma center.

An acute Lisfranc injury was defined as injury to the TMT joints with avulsion fractures, intra-articular fractures, and/or displacement of 1 or more TMT joints. Injuries to the TMT joint were identified using primary radiographs, CT scans, stress fluoroscopy, and/or weightbearing radiographs.

The joints were considered unstable if radiographs or CT scans showed a displacement of 2 mm or more in any direction or a stress test under fluoroscopy showed displacement of the joints.

Inclusion criteria were Lisfranc injuries with instability of the medial 3 TMT joints and no fractures in relation to the first TMT joint, in patients between 18 and 65 years old. Minor capsular avulsions in relation to the first TMT joint was accepted as a primarily ligamentous injury.

The exclusion criteria were concomitant other major lower extremity injuries / polytrauma, open injuries, previous foot pathology, diabetes mellitus, neuropathy, and peripheral vascular disease.

Randomization

After informed consent was obtained, the patients were randomly assigned to one of the 2 groups using the random

allocation rule. Allocations were kept in sealed, opaque envelopes containing "Primary Arthrodesis" or "Bridge Plate" and were manually shuffled. Neither the orthopedic surgeon performing the follow-up examinations nor the patients were blinded.

Demographics. Twenty-four patients were randomized to the BP group, and 24 patients were randomized to the PA group. The patient characteristics at time of inclusion, mechanisms of injury, and injury characteristics are presented in Table 1. No between-group differences were observed at baseline, except from a higher rate of ligamentous injuries in the lateral column in the BP group ($P = .049$). The time from injury to surgery was 16.5 days in the PA group and 15 days in the BP group ($P = .68$). The bridge plate over the first TMT joint in the BP group was removed at a mean of 138 (SD 22) days after primary surgery. The "homerun-screw" and the hardware used for the TMT joint arthrodesis were not routinely removed.

Operative Technique and Rehabilitation

The surgeries were performed by orthopedic surgeons experienced in foot and ankle surgery. A 2-incision technique was used, with one longitudinal dorsomedial incision over the first TMT joint and a second incision over the third TMT joint. A skin bridge of at least 4 cm between the incisions was preserved to reduce the risk of wound complications. The 3 medial TMT joints were exposed. As all of the patients were treated with a primary arthrodesis of the second and third TMT joint, the cartilage was removed from these joints and the subchondral bone was multiperforated using a 2-mm drill bit to enhance fusion. No bone graft was used. If the patient was randomized to the PA group, the same procedure was performed in the first TMT joint. In the patients randomized to dorsal bridge plating, the cartilage of the first TMT joint was left intact. The TMT joints were then reduced anatomically, starting with the first TMT joint and temporary fixed using 1.6-mm Kirschner wires. The reduction was confirmed both visually and by fluoroscopy. In the BP group, the first TMT joint was bridged with a 2.7-mm locking plate (LCP Compact Foot 2.4/2.7; DePuy Synthes, Oberdorf, Switzerland), while in the arthrodesis group the arthrodesis of the first TMT joint was fixed with two 2.7- or 3.5-mm fully threaded screws with interfragmentary compression. The primary arthrodesis of the second and third TMT joints was either fixed using 2.7- or 3.5-mm fully threaded screws or, in case of a severely comminuted joint, a locking plate was used. A "homerun-screw" was then placed from the medial cuneiform to the base of the second metatarsal securing the Lisfranc mortise. After reduction and fixation of the 3 medial TMT joints, the reduction of the fourth and fifth TMT joints was assessed. If displaced, the 2 lateral TMT joints were reduced and stabilized using 1.6-mm Kirschner wires (Figure 2).

Figure 1. CONSORT flowchart of the trial enrollment and analysis. TMT, tarsometatarsal.

Postoperative Care

The patients remained non-weight bearing for the first 6-8 weeks. Postoperatively the patients had a short leg splint that was removed at 2-3 weeks together with the sutures, and a short leg cast was applied. Cast immobilization was discontinued at 6-8 weeks postoperatively, and any K-wires transfixing the fourth and fifth TMT joints were removed. The patients then started weight-bearing as tolerated in a walker boot, which was used during weight-bearing until 12

weeks after surgery. In the BP group, the dorsal bridge plate over the first TMT joint was removed 4-5 months after surgery.

Outcome Measures

The patients returned for follow-up at 2-3 weeks, 6-8 weeks, 12 weeks, 6 months, 1 and 2 years postoperatively. The main outcome measure was the American Orthopaedic Foot & Ankle Society (AOFAS) midfoot scale, which consists of

Table 1. Patient Characteristics at Time of Enrollment, Mechanism of Injury, and Injury Characteristics.

Characteristics	PA (n= 24)	BP (n=24)
Gender, male/female	11/13	11/13
Age ^a	30 (23-40)	34 (28-40)
Side, right/left	13/11	7/17
Smoking, yes/no	3/21	5/19
BMI ^a	24 (21-28)	25 (22-27)
Mechanism of injury		
Fall <1 m or twisting of foot	12	7
Fall 1-3 m	2	1
Fall >3 m	0	1
Crush injury	0	1
Bike accident	0	2
Sports-related injury	10	12
Injury characteristics		
Ligamentous injuries		
Medial column	24	24
Central column	2	6
Lateral column ^b	2	9
Intra-articular fractures		
Medial column	0	0
Central column	22	18
Lateral column ^b	15	9

Abbreviations: BP, bridge plate; PA, primary arthrodesis.

^aPresented as median and interquartile range in parentheses.

^bThe only statistically significant difference between the PA and BP group at enrollment was a higher rate of ligamentous injuries in the lateral column in the BP group, $P = .049$.

3 main parts (pain, function, and alignment) resulting in a score ranging from 0 to 100, the best score being 100.¹² The secondary outcome parameters were the visual analog scale (VAS) for pain at rest and during walking, and the 36-Item Short Form Health Survey (SF-36) health survey, a patient-reported survey of patient health, at the 1- and 2-year follow-ups.³⁵ The same orthopedic surgeon conducted all of the 1- and 2-year follow-up examinations.

Radiographic Measurements

Postoperative CT scans and radiographs were obtained within the first 3 days after surgery. Weight-bearing anteroposterior (AP), lateral, and 30-degree oblique radiographs were obtained at the 6-week, 12-week, 12-month, and 24-month follow-ups. Reduction was evaluated on the postoperative CT scans. An anatomic reduction was defined as <2-mm displacement in any direction of the TMT joints, and a distance of no more than 2 mm between the medial cuneiform and the base of the second metatarsal.

At the 2-year follow-up, the alignment of the foot was assessed with weight-bearing radiographs. A displacement of 2 mm or more compared with the postoperative CT scans

was considered as secondary displacement. The Meary angle (talus-first metatarsal angle) was measured on the AP and lateral weightbearing radiographs to assess any difference in the alignment of the first metatarsal between the 2 groups.¹⁵ Osteoarthritis (reduced joint space, sclerosis, cysts, or osteophytes) of the first TMT joint in the BP group, as well as any nonunion of the fused TMT joints, were recorded.

Pedobarography

At 2-year follow-up, the plantar pressure of the patients was evaluated with the Tekscan HR mat (Tekscan Inc, South Boston, MA) and the Tekscan research software. Contact length, peak pressure and contact area were recorded for the following areas: medial heel (MH), lateral heel (LH), mid-foot (MF), first metatarsal (M1), second metatarsal (M2), third metatarsal (M3), fourth metatarsal (M4), fifth metatarsal (M5), and first toe (T1).

Statistical Analysis

A superiority study was conducted and power analysis prior to the study enrollment showed that 44 patients (22 in each group) had to be included to show a difference of 10 points or more on the AOFAS scale, with an estimated standard deviation of 10. This would give a power of 0.9 and a significance level of 5%. Taking into account any loss to follow-up we planned for 24 patients included in each group. The statistical analyses were performed using the Statistical Package for the Social Science (SPSS) software, version 25 (SPSS Inc, Chicago, IL). Data were tested for normality. Normal distributed data are presented as means with standard deviations and the 2-sided t test for independent samples was used for statistical analysis. The Mann-Whitney U test was used for nonparametric data, which are presented as median and interquartile range (IQR). The primary outcome (AOFAS midfoot scale), which was used for power analysis, showed a nonparametric distribution. These data are presented as means and standard deviation (SD), and the t test has been applied. In addition, the data were analyzed using nonparametric tests (Mann-Whitney U test), gaining similar results. The Fischer exact test was used to analyze categorical data. Spearman rank correlation was used to calculate correlation between variables. The significance level was $P \leq .05$.

Results

Clinical Outcomes

The mean AOFAS midfoot score and median VAS pain scores were not significantly different between the 2 groups at the 1- and 2-year follow-up (Table 2).

Table 2. Outcome Measures.

Outcome Measure	PA	BP	P Value
AOFAS midfoot scale ^a			
1 y (PA: n=23, BP: n=23)	85 (10)	79 (16)	.12
2 y (PA: n=22, BP: n=23)	89 (9)	85 (15.0)	.32
VAS pain at rest ^b			
1 y (PA: n=23, BP: n=23)	0 (0-0)	0 (0-2.4)	.14
2 y (PA: n=22, BP: n=23)	0 (0-0.4)	0 (0-0.8)	.72
VAS pain during walking ^b			
1 y (PA: n=23, BP: n=23)	1.4 (0-4.2)	2.5 (0-4.4)	.45
2 y (PA: n=22, BP: n=23)	0 (0-2.0)	0.9 (0-3.0)	.42
SF-36 1-y ^b (PA: n=23, BP: n=23)			
Physical function	90 (85-100)	90 (69-96)	.44
Role physical	100 (75-100)	88 (25-100)	.05
Bodily pain	90 (70-90)	70 (45-90)	.18
General health	80 (65-95)	80 (65-90)	.91
Vitality	60 (45-75)	60 (54-66)	.62
Social function	100 (63-100)	100 (72-100)	.93
Role emotional	100 (67-100)	100 (0-100)	.29
Mental health	80 (48-80)	80 (50-80)	.62
Physical component	54 (39-54)	51 (33-51)	.27
Mental component	54 (30-54)	54 (31-54)	.46
SF-36 2-y ^b (PA: n=22, BP: n=22)			
Physical function	95 (89-100)	95 (89-100)	.98
Role physical	100 (44-100)	100 (88-100)	.61
Bodily pain	90 (62-100)	90 (82-100)	>.99
General health	83 (70-95)	75 (64-90)	.25
Vitality	60 (44-75)	63 (45-71)	.88
Social function	94 (72-100)	100 (84-100)	.37
Role emotional	100 (25-100)	100 (58-100)	.54
Mental health	78 (60-93)	80 (72-84)	.69
Physical component	55 (49-58)	54 (46-58)	.92
Mental component	50 (36-58)	52 (47-55)	.75

Abbreviations: AOFAS, American Orthopaedic Foot & Ankle Society midfoot scale; BP, bridge plate; IQR, interquartile range; PA, primary arthrodesis; SF-36, 36-Item Short Form Health Survey; VAS pain, visual analog scale for pain.

^aData are presented as mean, with the SD in parentheses.

^bData are presented as median, with the IQR in parentheses.

The SF-36 at one- and 2-year follow-up did not show any significant difference between the groups in any of the 8 subgroups nor in the physical and mental component summary (Table 2).

Radiographic Outcomes

The radiographic findings are presented in Table 3. In both the PA and the BP group, 21 of the 24 patients had an anatomical reduction evaluated on the postoperative CT scan. At the 2-year follow-up, no patients had secondary displacement on weightbearing radiographs. In the PA group, the AP Meary angle was significantly increased compared with the BP group (12 degrees; IQR, 5-15 vs 5 degrees; IQR, 2-13, $P = .04$).

After 2 years, 11/24 patients in the BP group had radiographic osteoarthritis in the first TMT joint. Only 1 of these patients required conversion to arthrodesis of the first TMT joint at 24 months because of painful osteoarthritis.

Pedobarography

The pedobarographic results are presented in Figure 3. The only statistically significant difference between the 2 groups was the reduced peak pressure under the fifth metatarsal in the PA group. The peak pressure under the fifth metatarsal was also reduced compared with the non-injured side in the PA group (16 N/cm², IQR, 13-24, vs 26 N/cm², IQR, 15-31; $P = .047$); this reduction was not seen in the BP group. The fifth metatarsal peak pressure was

Table 3. Radiographic Findings.

Radiographic Findings	PA Group	BP Group	P Value
K-wire fixation TMT 4+5	6 (24)	2 (24)	.245
Anatomic reduction postop CT scan ^a	21 (24)	21 (24)	>.99
Anatomic reduction radiographs at 2 y	21 (22)	23 (23) ^b	.49
AP Meary angle, degrees, median (IQR)	12 (5-15)	5 (2-13)	.042^d
Lateral Meary angle, degrees, median (IQR)	1 (0-3)	3 (0-6)	.23
Nonunion (all third TMT joint)	1 (22)	2 (23)	>.99
Osteoarthritis TMT-I		11 (24) ^c	
Spontaneous fusion TMT-I		4 (23)	

Abbreviations: AP, anteroposterior; BP, bridge plate; CT, computed tomography; IQR, interquartile range; PA, primary arthrodesis; TMT, tarsometatarsal.

^aThere was 1 nonanatomic reduction in the first TMT joint in each group.

^bTwo of the TMT joint displacements seen on the postoperative CT scans were not detected on the 2-year follow-up radiographs.

^cThe one patient who did not attend the 2-year follow-up had osteoarthritis of the first TMT joint on the 1-year follow-up.

^dBold indicates statistically significant findings.

Figure 2. (A) Primary arthrodesis of the 3 medial TMT joints. (B) Temporary bridge plating of the first TMT joint using locking plate and primary arthrodesis of the second and third TMT joint. (C) Same patient as in panel B, but after removal of the temporary bridging plate. TMT, tarsometatarsal.

correlated to the lateral Meary angle in the PA group ($r_s = -0.65$, $P = .002$).

Complications and Additional Surgeries

One patient had a superficial wound infection in the BP group treated successfully with oral antibiotics. Two

patients in the BP group developed complex regional pain syndrome. In the PA group, 1 patient had persistent pain because of irritation of the medial branch of the superficial peroneal nerve. One patient in the BP group experienced loosening of a screw from the plate, which led to a rupture of the extensor hallucis longus tendon. The patient was treated with a transfer of the second toe extensor digitorum

Figure 3. Pedobarographic data of injured side presented as boxplots. The only statistically significant finding (Mann-Whitney U test) was the reduced peak pressure under the fifth metatarsal in the PA group ($P = .01$). BP, bridge plate; M1, first metatarsal; M2, second metatarsal; M3, third metatarsal; M4, fourth metatarsal; M5, fifth metatarsal; MF, midfoot; MH, medial heel; LH, lateral heel; PA, primary arthrodesis; TF, total foot.

longus to extensor hallucis longus with an excellent outcome. No statistical difference in complication rate was observed between the 2 groups ($P = .348$).

Nonunions developed in the third TMT joint in 3 patients (1 in the PA and 2 in the BP group). In addition to the planned removal of the bridge plate in the BP group, 7/23 patients in the PA and 5/24 patients in the BP group had hardware removed ($P = .450$) at a median of 302 days (IQR, 149-725).

Discussion

In this randomized controlled trial comparing temporary bridge plating (BP) to primary arthrodesis (PA) of the first TMT joint in acute Lisfranc injuries, we were not able to detect superiority of the BP group compared to the PA group using AOFAS midfoot score, VAS pain score, or SF-36 at 2-year follow-up. A high incidence of radiologic osteoarthritis was present in the first TMT joint in the BP group; however, only 1 patient required a fusion due to painful osteoarthritis. The first metatarsal was better aligned in the AP radiographs in the BP group evaluated by the AP Meary angle. Pedobarographic examination revealed a reduced peak pressure under the fifth metatarsal in the PA group. We had a trend toward more complications in the BP group ($P = .348$).

The medial 3 TMT joints are considered nonessential joints, and fusion of these joints is well tolerated by patients.^{8,13,18,19,22} The first tarsometatarsal joint is more mobile than the central TMT joints, and preserving this joint might be beneficial.²¹ As there is almost no motion in the second and third TMT joints and these joints have a tendency to develop painful osteoarthritis, these joints were fused in the present study.¹³ The lateral column is the most mobile and plays an important role in the foot's ability to adapt to the surface and should not be fused.^{13,20,22} In the current study, we have compared preserving the first TMT joint to fusing it.

We did not find superiority of the BP group compared to the PA group with regards to the AOFAS midfoot score or VAS pain score (mean AOFAS 89 vs 85, $P = .32$ and median VAS pain during walking 0 vs 1.0, $P = .42$). Both groups had good outcome scores, comparable to the study by Ly et al, in which the primary arthrodesis group had a mean AOFAS midfoot score of 88 and VAS pain score of 1.2 at 2-year follow-up.¹⁸ No studies have, to our knowledge, investigated the outcome after combining temporary bridge plating of the first TMT joint with primary arthrodesis of the second and third TMT joints. However, the AOFAS midfoot scores were similar to the results of temporary bridge plating reported by Hu et al in their prospective comparative study with an average follow-up of 31 months (mean AOFAS midfoot score in plate group of 83.1), and in the retrospective study by Kirzner et al (mean AOFAS midfoot score in plate group of 82.5).^{9,11} Stern and Assal

reported a mean AOFAS midfoot score of 85 at 1-year follow-up in their retrospective case series of 15 patients treated with dorsal bridge plating.³³ In all these studies the first, second, and/or third TMT joints were stabilized using bridge plates whereas the 2 lateral TMT joints were stabilized using K-wires, if unstable.

With regard to quality of life, Schepers et al reported normalization of the SF-36 score at a median follow-up of 76 months after isolated Lisfranc injuries.³⁰ The SF-36 score in both groups of our study was comparable to reported data from the Norwegian general population aged 30-39 years, both with regard to the physical component summary (mean 53, SD 8) and mental component summary (mean 52, SD 9) score, suggesting a return to quality of life at the same level as the general population.⁷

Osteoarthritis after Lisfranc injuries is reported in up to 25%-94%.^{14,20,25,32} The bridge plate technique is a joint-sparing technique that might reduce the risk of osteoarthritis.^{1,11,33} Lau et al was, however, not able to relate osteoarthritis to the method of fixation when comparing bridge plating to transarticular screw fixation.¹⁷ Previous studies using dorsal bridge plating have reported a high rate of mild post-traumatic osteoarthritis on follow-up radiographs.^{17,33} In the present study 11/24 patients had degenerative changes on the follow-up radiographs at 2 years, but only 1 patient required an arthrodesis of the first TMT joint due to pain. The long-term incidence and severity of osteoarthritis after temporary bridge plating, and how many of these patients will require an arthrodesis, are still not known.

Hu et al reported a spontaneous fusion rate of 5/32 (15.6%) after treating Lisfranc injuries with joint-preserving dorsal bridge plates, but 4 of the patients did not have any discomfort.⁹ In the present study 4/24 patients treated with a bridge plate of the first TMT joint developed a spontaneous fusion of the joint within 2 years. The patients with a spontaneous fusion had a favorable outcome with a median AOFAS midfoot score at 2 years of 92 (range, 80-100).

In the 2 previously published randomized controlled trials comparing primary arthrodesis to transarticular screw fixation in Lisfranc injuries, low rates of nonunions were reported (1/21 and 1/19).^{8,18} Altogether, there were 3 nonunions after arthrodesis in our present study, all of these occurred in the third TMT joint (one in the PA group and 2 in the BP group). These patients had only minor complaints and did not want further surgery with revision arthrodesis.

Anatomic reduction of the TMT joints and a stable fixation are considered the most important factors in achieving a good outcome when treating unstable Lisfranc injuries.^{11,14,23,29,32} Kirzner et al found a trend toward better reduction of Lisfranc injuries using bridge plating compared with transarticular screw fixation ($P = .06$).¹¹ On the other hand, Lau et al were not able to confirm any

differences in the quality of reduction when comparing transarticular screw fixation to bridge plating.¹⁷ In the present study, we could not find a difference between the 2 groups in the anatomic reduction of the TMT joints or in the distance between the medial cuneiform and the base of the second metatarsal on postoperative CT scan or on the radiographs at the 2-year follow-up. However, examining the alignment of the first metatarsal we did observe an increased AP Meary angle in the PA group compared to the BP group (12 vs 5 degrees, $P = .042$), indicating that the first metatarsal was fixed in an abducted position. Lamm et al reported the AP Meary angle in normal subjects to have a median value of 6.5 degrees (range, 0-18) which is comparable to the findings in our BP group.¹⁵ No statistical difference was found between the 2 groups when comparing the lateral Meary angle, although the PA group had a slightly reduced angle compared with the BP group (1 vs 3 degrees, $P = .23$). Pedobarographic examination showed a reduced fifth metatarsal peak pressure in the PA group compared with the BP group ($P = .047$). The fifth metatarsal peak pressure correlated to the lateral Meary angle in the PA group ($r_s = -0.65$, $P = .002$). In addition, the medial column is shortened when performing an arthrodesis, which also may elevate the head of the first metatarsal. Both the shortening of the medial column and the slightly reduced lateral Meary angle can lead to increased pronation of the foot. We believe that the reduced fifth metatarsal pressure was caused by a change in the length and alignment of the first metatarsal in the PA group.

Loss of reduction has been reported as a complication to bridge plating of Lisfranc injuries. Kirzner and colleagues reported a loss of reduction in 11% of their patients treated with bridge plates vs 24% in the group treated with transarticular screws.¹¹ We found no patients with a secondary displacement at the 2-year follow-up. This might be because only the first TMT joint was bridge plated and the second and third TMT joints were fused, in contrast to the bridge plate group in the study by Kirzner, where all the unstable central and medial joints were bridge plated.¹¹

All the patients in the BP group had the locking plate bridging the first TMT joint removed at a mean of 138 (22) days after the primary surgery. This additional surgery can be one of the causes for the higher number of complications, although not statistically significant, seen in the BP group.

The strength of the present study is first and foremost its randomized controlled design. The study is, to our knowledge, the only study comparing temporary bridge plating to primary arthrodesis in Lisfranc injuries. All the surgeries were performed by orthopedic surgeons experienced in foot and ankle surgery and the surgeons were familiar with both operative techniques prior to study inclusion. Anatomic reduction was evaluated with postoperative CT scans, and there was a low rate of loss to follow-up in the study.

The study does have some inherent limitations. A majority of the patients sustained low-energy trauma as polytrauma patients were excluded. Low-energy injuries are known to have better results than the high-energy injuries in terms of outcome scores, postoperative reduction, and post-traumatic osteoarthritis.^{17,27,34} Only the patients with clinically and/or radiologically suspected nonunion were evaluated with CT scans. We thereby might have missed nonunions with only minor symptoms. The patients and examiner were not blinded for the treatment, with a potential for introducing bias. In addition, the AOFAS midfoot scale has been shown to have an acceptable validity and internal consistency when used to evaluate Lisfranc injuries, but has a ceiling effect and a lack of coverage and targeting.²⁶ Even so, a review of patient-reported outcome measures in foot and ankle surgery showed the AOFAS midfoot scale to be the most commonly used outcome measure in studies of Lisfranc injuries together with the VAS for pain and SF-36.¹⁰ This makes these 3 outcome scores suitable for comparing the results with previously published studies.

Conclusion

Both the temporary bridge plate group and the primary arthrodesis group yielded good outcome scores when treating Lisfranc injuries. We did not find superiority of the BP group compared to the PA group according to the AOFAS midfoot score. The first metatarsal was better aligned in the BP group. Despite avoiding transarticular screw damage by bridge plating the first TMT joints, there was a high rate of radiologically detected osteoarthritis in the first TMT joint. The long-term effects of post-traumatic osteoarthritis is still unknown and longer follow-up is required.

Declaration of Conflicting Interests

The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: Are H. Stødle reports grants from Sofies Mindes Ortopedi AS, Oslo, Norway, during the conduct of the study. ICMJE forms for all authors are available online.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Sophies Minde Ortopedi.

ORCID iDs

Are H. Stødle, MD, <https://orcid.org/0000-0002-4173-2748>
Jan Erik Madsen, MD, PhD, <https://orcid.org/0000-0002-9002-1213>
Elisabeth Ellingsen Husebye, MD, PhD, <https://orcid.org/0000-0002-3297-3224>

References

1. Alberta FG, Aronow MS, Barrero M, Diaz-Doran V, Sullivan RJ, Adams DJ. Ligamentous Lisfranc joint injuries: a biomechanical comparison of dorsal plate and transarticular screw fixation. *Foot Ankle Int.* 2005;26(6):462-473.
2. Arntz CT, Veith RG, Hansen ST, et al. Fractures and fracture-dislocations of the tarsometatarsal joint. *J Bone Joint Surg Am.* 1988;70(2):173-181.
3. Benirschke SK, Kramer PA. High energy acute lisfranc fractures and dislocations. *Tech Foot Ankle Surg.* 2010;9(3):82-91.
4. Cochran G, Renninger C, Tompane T, Bellamy J, Kuhn K. Primary arthrodesis versus open reduction and internal fixation for low-energy lisfranc injuries in a young athletic population. *Foot Ankle Int.* 2017;38(9):957-963.
5. Coetzee JC. Making sense of lisfranc injuries. *Foot Ankle Clin.* 2008;13(4):695-704.
6. Curtis MJ, Myerson M, Szura B. Tarsometatarsal joint injuries in the athlete. *Am J Sports Med.* 1993;21(4):497-502.
7. Garratt AM, Stavem K. Measurement properties and normative data for the Norwegian SF-36: results from a general population survey. *Health Qual Life Outcomes.* 2017;15(1):1-10.
8. Henning JA, Jones CB, Sietsema DL, Bohay DR, Anderson JG. Open reduction internal fixation versus primary arthrodesis for lisfranc injuries: a prospective randomized study. *Foot Ankle Int.* 2009;30(10):913-922.
9. Hu S, Chang S, Li X, Yu G. Outcome comparison of Lisfranc injuries treated through dorsal plate fixation versus screw fixation. *Acta Orthop Bras.* 2014;22(6):315-320.
10. Hunt KJ, Hurwit D. Use of patient-reported outcome measures in foot and ankle research. *J Bone Joint Surg Am.* 2013;95(16):e118(1-9).
11. Kirzner N, Zotov P, Goldbloom D, Curry H, Bedi H. Dorsal bridge plating or transarticular screws for Lisfranc fracture dislocations. *Bone Joint J.* 2018;100-B(4):468-474.
12. Kitaoka HB, Alexander IJ, Adelaar RS, Nunley JA, Myerson MS, Sanders M. Clinical rating systems for the ankle-hindfoot, midfoot, hallux, and lesser toes. *Foot Ankle Int.* 1994;15(7):349-353.
13. Komenda GA, Myerson MS, Biddinger KR. Results of arthrodesis of the tarsometatarsal joints after traumatic injury. *J Bone Joint Surg Am.* 1996;78:1665-1676.
14. Kuo RS, Tejwani NC, Digiovanni CW, et al. Outcome after open reduction and internal fixation of Lisfranc joint injuries. *J Bone Joint Surg Am.* 2000;82-A:1609-1618.
15. Lamm BM, Stasko PA, Gesheff MG, Bhav A. Normal foot and ankle radiographic angles, measurements, and reference points. *J Foot Ankle Surg.* 2016;55(5):991-998.
16. Lau S, Bozin M, Thillainadesan T. Lisfranc fracture dislocation: a review of a commonly missed injury of the midfoot. *Emerg Med J.* 2017;34(1):52-56.
17. Lau S, Howells N, Millar M, De Villiers D, Joseph S, Oppy A. Plates, screws, or combination? Radiologic outcomes after lisfranc fracture dislocation. *J Foot Ankle Surg.* 2016;55(4):799-802.
18. Ly TV, Coetzee JC. Treatment of primarily ligamentous lisfranc joint injuries: primary arthrodesis compared with open reduction and internal fixation: A prospective, randomized study. *J Bone Joint Surg.* 2006;88(3):514.
19. Mulier T, Haan J, Vriesendorp P, Reynders P. The treatment of lisfranc injuries: review of current literature. *Eur J Trauma Emerg Surg.* 2010;36(3):206-216.
20. Mulier T, Reynders P, Dereymaeker G, Broos P. Severe Lisfrancs injuries: primary arthrodesis or ORIF? *Foot ankle Int.* 2002;23(10):902-905.
21. Myerson MS. The diagnosis and treatment of injury to the tarsometatarsal joint complex. *J Bone Joint Surg Br.* 1999;81(5):756-763.
22. Myerson MS, Cerrato RA. Current management of tarsometatarsal injuries in the athlete. *J Bone Joint Surg Am.* 2008;90(11):2522-2533.
23. Myerson MS, Fisher RT, Burgess AR, Kenzora JE. Fracture dislocations of the tarsometatarsal joints: end results correlated with pathology and treatment. *Foot Ankle.* 1986;6(5):225-242.
24. Nunley JA, Vertullo CJ. Classification, investigation, and management of midfoot sprains. *Am J Sports Med.* 2002;30(6):871-878.
25. Pereira CJ, Espinosa EG, Miranda I, Pereira MB, Canto RSDT. Evaluation of the surgical treatment of lisfranc joint fracture-dislocation. *Acta Orthop Bras.* 2008;16:93-97.
26. Ponkilainen VT, Uimonen M, Repo JP, Mattila VM, Haapasalo HH. Validity and internal consistency of the American Orthopaedic Foot & Ankle Society Midfoot Scale in patients with Lisfranc injury [published online ahead of print June 21, 2019]. *Foot Ankle Surg.* 2019.
27. Renninger CH, Cochran G, Tompane T, Bellamy J, Kuhn K. Injury characteristics of low-energy lisfranc injuries compared with high-energy injuries. *Foot Ankle Int.* 2017;38(9):964-969.
28. Richter M, Thermann H, Huefner T, Schmidt U, Krettek C. Aetiology, treatment and outcome in Lisfranc joint dislocations and fracture dislocations. *Foot Ankle Surg.* 2002;8(1):21-32.
29. Richter M, Wippermann B, Krettek C, Schratz HE, Hufner T, Thermann H. Fractures and fracture dislocations of the midfoot: Occurrence, causes and long-term results. *Foot Ankle Int.* 2001;22(5):392-398.
30. Schepers T, Kieboom BCT, Kieboom B, van Diggele P, Patka P, Van Lieshout EMM. Pedobarographic analysis and quality of life after Lisfranc fracture dislocation. *Foot Ankle Int.* 2010;31(10):857-864.
31. Scolaro J, Ahn J, Mehta S. Lisfranc fracture dislocations. *Clin Orthop Relat Res.* 2011;469(7):2078-2080.
32. Stavlas P, Roberts CS, Xypnitos FN, Giannoudis PV. The role of reduction and internal fixation of Lisfranc fracture-dislocations: a systematic review of the literature. *Int Orthop.* 2010;34(8):1083-1091.
33. Stern RE, Assal M. Dorsal multiple plating without routine transarticular screws for fixation of lisfranc injury. *Orthopedics.* 2014;37(12):815-819.
34. van Koperen PJ, de Jong VM, Luitse JSK, Schepers T. Functional outcomes after temporary bridging with locking plates in lisfranc injuries. *J Foot Ankle Surg.* 2016;55(5):922-926.
35. Ware JE, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care.* 1992;30(6):473-483.
36. Watson TS, Shurnas PS, Denker J. Treatment of Lisfranc joint injury: current concepts. *J Am Acad Orthop Surg.* 2010;18(12):718-728.

AOFAS Midfoot Scale (100 Points Total)

Pain (40 points)

- None.....40
- Mild, occasional.....30
- Moderate, daily.....20
- Severe, almost always present.....0

Function (45 points)

Activity limitations, support

- No limitations, no support.....10
- No limitation of daily activities, limitation of recreational activities, no support...7
- Limited daily and recreational activities, cane.....4
- Severe limitation of daily and recreational activities, walker, crutches, wheelchair..... 0

Footwear requirements

- Fashionable, conventional shoes, no insert required....5
- Comfort footwear, shoe insert.....3
- Modified shoes or brace.....0

Maximum walking distance, blocks

- Greater than 6.....10
- 4-6.....7
- 1-3.....4
- Less than 1.....0

Walking surfaces

- No difficulty on any surface.....10
- Some difficulty on uneven terrain, stairs, inclines, ladders.....5
- Severe difficulty on uneven terrain, stairs, inclines, ladders.....0

Gait abnormality

- None, slight.....10
- Obvious.....5
- Marked.....0

Alignment (15 points)

- Good, plantigrade foot, midfoot well aligned.....15
- Fair, plantigrade foot, some degree of midfoot malalignment observed, no symptoms.....8
- Poor, nonplantigrade foot, severe malalignment, symptoms....0

Visuell Analog Skala (VAS)

(Pasienten markerer selv)

0: Ingen smerter

10: Verst tenkelige smerter

Smerter i hvile:

Smerter ved gange:

SF-36 SPØRRESKJEMA OM HELSE

INSTRUKSJON: Dette spørreskjemaet handler om hvordan du ser på din egen helse. Disse opplysningene vil hjelpe oss til å få vite hvordan du har det og hvordan du er i stand til å gjennomføre dine daglige gjøremål.

Hvert spørsmål skal besvares ved å sette en ring rundt det tallet som passer best for deg. Hvis du er usikker på hva du skal svare, vennligst svar så godt du kan.

1. Stort sett, vil du si din helse er:

(sett ring rundt ett tall)

Utmerket	1
Meget god	2
God	3
Nokså god	4
Dårlig	5

2. Sammenliknet med for ett år siden, hvordan vil du si at din helse stort sett er nå?

(sett ring rundt ett tall)

Mye bedre nå enn for ett år siden	1
Litt bedre nå enn for ett år siden	2
Omtrent den samme som for ett år siden	3
Litt dårligere nå enn for ett år siden	4
Mye dårligere nå enn for ett år siden	5

3. De neste spørsmålene handler om aktiviteter som du kanskje utfører i løpet av en vanlig dag. Er din helse slik at den begrenser deg i utførelsen av disse aktivitetene nå? Hvis ja, hvor mye?

(sett ring rundt ett tall på hver linje)

AKTIVITETER	Ja, begrenser meg mye	Ja, begrenser meg litt	Nei, begrenser meg ikke i det hele tatt
a. Anstrengende aktiviteter som å løpe, løfte tunge gjenstander, delta i anstrengende idrett	1	2	3
b. Moderate aktiviteter som å flytte et bord, støvsuge, gå en tur eller drive med hagearbeid	1	2	3
c. Løfte eller bære en handlekurv	1	2	3
d. Gå opp trappen flere etasjer	1	2	3
e. Gå opp trappen en etasje	1	2	3
f. Bøye deg eller sitte på huk	1	2	3
g. Gå mer enn to kilometer	1	2	3
h. Gå noen hundre meter	1	2	3
i. Gå hundre meter	1	2	3
j. Vaske deg eller kle på deg	1	2	3

4. I løpet av de siste 4 ukene, har du hatt noen av følgende problemer i ditt arbeid eller i andre dine daglige gjøremål på grunn av din fysiske helse?

(sett ring rundt ett tall på hver linje)

	JA	NEI
a. Du har måttet redusere tiden du har brukt på arbeid eller på andre gjøremål	1	2
b. Du har utrettet mindre enn du hadde ønsket	1	2
c. Du har vært hindret i å utføre visse typer arbeid eller gjøremål	1	2
d. Du har hatt problemer med å gjennomføre arbeidet eller andre gjøremål (for eksempel fordi det krevde ekstra anstrengelser)	1	2

5. I løpet av de siste 4 ukene, har du hatt noen av følgende problemer i ditt arbeid eller i andre dine daglige gjøremål på grunn av følelsesmessige problemer (som for eksempel å være deprimentert eller engstelig)?

(sett ring rundt ett tall på hver linje)

	JA	NEI
a. Du har måttet redusere tiden du har brukt på arbeid eller på andre gjøremål	1	2
b. Du har utrettet mindre enn du hadde ønsket	1	2
c. Du har utført arbeidet eller andre gjøremål mindre grundig enn vanlig	1	2

6. I løpet av de siste 4 ukene, i hvilken grad har din fysiske helse eller følelsesmessig problemer hatt innvirkning på din vanlige sosiale omgang med familie, venner, naboer eller foreninger?

(sett ring rundt ett tall)

Ikke i det hele tatt	1
Litt	2
En del	3
Mye	4
Svært mye	5

7. Hvor sterke kroppslige smerter har du hatt i løpet av de siste 4 ukene?

(sett ring rundt ett tall)

Ingen	1
Meget svake	2
Svake	3
Moderate	4
Sterke	5
Meget sterke	6

8. I løpet av de siste 4 ukene, hvor mye har smerter påvirket ditt vanlige arbeid (gjelder både arbeid utenfor hjemmet og husarbeid)?

(sett ring rundt ett tall)

Ikke i det hele tatt	1
Litt	2
En del	3
Mye	4
Svært mye	5

9. De neste spørsmålene handler om hvordan du har følt deg og hvordan du har hatt det de siste 4 ukene. For hvert spørsmål, vennligst velg det svaralternativet som best beskriver hvordan du har hatt det. Hvor ofte i løpet av de siste 4 ukene har du:

(sett ring rundt ett tall på hver linje)

	Hele tiden	Nesten hele tiden	Mye av tiden	En del av tiden	Litt av tiden	Ikke i det hele tatt
a. Følt deg full av tiltakslyst	1	2	3	4	5	6
b. Følt deg veldig nervøs?	1	2	3	4	5	6
c. Vært så langt nede at ingenting har kunnet muntre deg opp?	1	2	3	4	5	6
d. Følt deg rolig og harmonisk?	1	2	3	4	5	6
e. Hatt mye overskudd?	1	2	3	4	5	6
f. Følt deg nedfor og trist?	1	2	3	4	5	6
g. Følt deg sliten?	1	2	3	4	5	6
h. Følt deg glad?	1	2	3	4	5	6
i. Følt deg trett?	1	2	3	4	5	6

10. I løpet av de siste 4 ukene, hvor mye av tiden har din fysiske helse eller følelsesmessige problemer påvirket din sosiale omgang (som det å besøke venner, slektninger osv.)?

(sett ring rundt ett tall)

Hele tiden	1
Nesten hele tiden	2
En del av tiden	3
Litt av tiden	4
Ikke i det hele tatt	5

11. Hvor RIKTIG eller GAL er hver av de følgende påstander for deg?

(sett ring rundt ett tall på hver linje)

	Helt riktig	Delvis riktig	Vet ikke	Delvis gal	Helt gal
a. Det viker som jeg blir syk litt lettere enn andre	1	2	3	4	5
b. Jeg er like frisk som de fleste jeg kjenner	1	2	3	4	5
c. Jeg tror at helsen min vil forverres	1	2	3	4	5
d. Jeg har utmerket helse	1	2	3	4	5