Snow avalanche controls, monitoring strategies, and hazard management in Svalbard

Holt John Hancock

Thesis submitted for the degree of Philosophiae Doctor (PhD)

Department of Geosciences
Faculty of Mathematics and Natural Sciences
University of Oslo
Oslo, Norway
2021

© **Holt John Hancock, 2021**

*Series of dissertations submitted to the
Faculty of Mathematics and Natural Sciences, University of Oslo
No. 2440*

ISSN 1501-7710

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

A super blue blood moon over Hiortfjellet during the January 31st, 2018 lunar eclipse.

Contents

Abstract.....	vi
Acknowledgements	viii
List of Publications	x
Introduction.....	2
Thesis scope and research objectives.....	5
Thesis structure	6
Background	8
Physiographic setting	8
Climate and meteorology	10
Snow climate and avalanche regime.....	13
Knowledge gaps.....	17
Methods.....	20
Terrestrial laser scanning	20
regObs	25
Weather and snow depth data	26
Research summary.....	28
Paper I.....	28
Paper II.....	29
Paper III	30
Synthesis and discussion.....	32
The avalanche forecasting framework	32
Meteorological controls on avalanche activity in central Spitsbergen	35
<i>The influence of snow and wind on avalanche activity in central Spitsbergen</i>	<i>35</i>
<i>Loading events and the high-Arctic maritime snow climate.....</i>	<i>37</i>
Methods for snow and avalanche observation in Arctic settings.....	38
<i>Terrestrial laser scanning.....</i>	<i>38</i>
<i>Regional snow and avalanche observations</i>	<i>40</i>
Hazard management implications.....	41
Conclusions.....	44
Future perspectives	46
References.....	48
Papers.....	56
Paper I.....	58
Paper II.....	74
Paper III	98

Abstract

Snow avalanches are a common natural hazard on Spitsbergen, the largest island in the Svalbard archipelago. The climatological setting here as the warmest and wettest region in the high-Arctic and the distinctive regional topography result in a unique snow environment with marked avalanche hazards. Recent destructive avalanche events in Longyearbyen, Svalbard's administrative center located in central Spitsbergen, highlighted the need for better avalanche process understanding and improved avalanche hazard management strategies in this avalanche setting. To better understand the demonstrated avalanche hazard in this high-Arctic environment, this work examines snow and avalanche processes across a range of spatiotemporal scales in central Spitsbergen and contextualizes the results from a hazard management perspective.

The research contained in the following thesis addresses knowledge gaps from previous avalanche-related studies in Svalbard by 1) employing and further developing terrestrial laser scanning (TLS) techniques to illustrate and quantify how specific meteorological conditions control avalanche activity at the slope and catchment spatial scales near Longyearbyen, and 2) investigating the synoptic-scale meteorological controls on regional avalanche activity in central Spitsbergen. Specific methodological contributions from this work include developing a workflow for investigating snow cornice and cornice-fall avalanche processes with TLS and establishing a foundation to consider the broader-scale controls on – and future changes to – avalanching in this location.

The findings in aggregate help demonstrate how snow and wind associated with winter storms serve as key controls on avalanche activity at all spatial scales in high-Arctic central Spitsbergen. Low-pressure activity near Svalbard promotes a snowy, windy synoptic environment and broadly controls dry avalanche activity in the study area. Regional wind direction controlled by the low pressure's position relative to Spitsbergen then determines the aspects on which cornices accrete and cornice-fall avalanches occur during these events at the catchment scale. At the slope scale, local topography modifies the regional wind direction to determine snow accumulation and distribution patterns which influence destructive avalanche release.

Hazard managers can apply the methods and results from this work to better forecast avalanches in Svalbard via synoptic-scale weather forecasts, develop more specific cornice-fall avalanche forecasting and management strategies, employ TLS data to aid in slope-scale avalanche forecasting, and design more effective permanent mitigation strategies. Finally, this work provides important information regarding avalanche activity in a rapidly changing environmental setting which can help inform future avalanche research and hazard management efforts in this and other areas where climatic changes are expected to impact avalanche regimes.

Acknowledgements

I am very grateful for Markus (Max) Eckerstorfer, Jordy Hendrikx, and Thomas Schuler's support and supervision throughout the duration of this work. Max's knowledge of snow and avalanches in Svalbard and elsewhere in the world, availability for assistance, and motivation (in life, at work, on skis, on bikes) as a friend and mentor have made this journey possible. Jordy introduced me to Svalbard as a master student and has helped guide my work since 2014. This work would not have been possible without Jordy's perspective and (especially) the fact he introduced me to coffee on our first trip here. Thomas has kindly assisted with the development of this research and the thesis writing, while also serving as an important link to the mainland university system. Alexander Prokop's help with terrestrial laser scanning was crucial for establishing this project in its current form. Finally, I am thankful for support from Chris Borstad during a particularly difficult stretch at UNIS.

I am deeply grateful for the environment in the Arctic Geology Department provided by Hanne H. Christiansen. Hanne's dedication to her co-workers is, I believe, unmatched and I am so thankful for her constant support as department head and as a friend throughout both the ups and downs of the last four years. Martin Indreiten's work with the Arctic Safety Centre at UNIS provided an opportunity to get out of the office and attempt to keep my operational skills somewhat honed, for which I am grateful.

I especially want to thank Sara Mollie Cohen for her friendship – help with work, dinners, trips, an open-door policy at home and work, being there during meltdowns, texting each other from our neighboring offices etc. – over the last half-decade. Suffice it to say we are all exceedingly lucky to have Sara Mollie's presence at UNIS and in Longyearbyen! And: Wes, Hrefna, Lilli, Martine, Kamilla, Karl, Micke, Kathinka, Oskar, Kieran, Tim, Erik, Natasza, Bryndis, Lise, Audun, Even, Graham, TK, Konrad – thank you all. In Montana, I'd like to thank Ty and Erin for continued and undeserved hospitality when I'm home, and Tristan and Ryan (and their partners) for constantly bearing the brunt of my “digital only” socialization techniques.

Siiri Wickström supplied 110+ support, owls, and unbelievable tolerance during this PhD. Looking forward for more ski trips in the future! I am always grateful for my family: Eliza Caldwell who continues to be an inspirational sibling with her kindness, generosity, and life management skills, and my parents, Scott and Julie Hancock, who probably took me skiing too much as a child and have no one but themselves to blame for my fascination with snow. Can't wait to see you guys again!

Holt Hancock

Longyearbyen, May 2021

List of Publications

Paper I

Hancock, H., Prokop, A., Eckerstorfer, M., and Hendrikx, J., 2018. Combining high spatial resolution snow mapping and meteorological analyses to improve forecasting of destructive avalanches in Longyearbyen, Svalbard, Cold Regions Science and Technology, 154, 120-132, doi:<https://doi.org/10.1016/j.coldregions.2018.05.011>.

Paper II

Hancock, H., Eckerstorfer, M., Prokop, A., and Hendrikx, J., 2020. Quantifying seasonal cornice dynamics using a terrestrial laser scanner in Svalbard, Norway, Nat. Hazards Earth Syst. Sci., 20, 603-623, doi:[10.5194/nhess-20-603-2020](https://doi.org/10.5194/nhess-20-603-2020).

Paper III

Hancock, H., Hendrikx, J., Eckerstorfer, M., and Wickström, S., 2021: Synoptic control on snow avalanche activity in central Spitsbergen, The Cryosphere, 15, 3813-3837, doi:[10.5194/tc-15-3813-2021](https://doi.org/10.5194/tc-15-3813-2021).

Introduction

Snow avalanches are broadly defined as snow masses rapidly descending steep slopes (e.g. Schweizer et al., 2003a) and are included in the suite of gravitational, mountain slope processes alongside landslides, rockfalls, and debris flows (McClung and Schaerer, 2006). Occurring in snow-covered areas with high topographic relief, snow avalanches – often referred to simply as avalanches – are a natural hazard impacting human life and infrastructure in mountainous locations throughout the world. Worldwide, avalanches presently account for approximately 250 human fatalities annually (Schweizer, 2008), and infrastructure damage estimates from single avalanche disasters in the last 30 years have on occasion exceeded 500 million euros (Fuchs and McAlpin, 2005).

Although avalanches are noteworthy agents of geomorphic change (e.g. André, 1990; Decaulne and Saemundsson, 2006; Luckman, 1977), ecological disturbance (e.g. Bebi et al., 2009; Peitzsch et al., 2021), and glacial accumulation (e.g. Scherler et al., 2011), avalanche process investigations have historically been motivated by mitigating avalanche-related hazards. The contemporary understanding of avalanches is thus largely based on operational and scientific experience in mid-latitude regions with concentrated human activity – and the resources to address avalanche hazard – in avalanche-prone locations. Improved avalanche process understanding has helped inform avalanche forecasts, updated avalanche hazard zonation and building regulations, and installations of permanent mitigation measures (Bründl and Margreth, 2014; Sauermoser, 2006; Stethem et al., 2003). In locations such as the European Alps and western North America, improvements to avalanche hazard management strategies over the last century have reduced fatalities and infrastructure damage in settlements and along transportation corridors. Recreational accidents now accordingly account for most avalanche-related fatalities in developed countries, with roughly 100 and 30 deaths annually in Europe and the United States, respectively, over the last decade (EAWS, 2021; CAIC, 2021). Catastrophic avalanche events involving settlements still do occur, however, in developing countries without the resources to adequately manage avalanche hazards (e.g. Chabot and Kaba, 2016), in remote locations without formalized avalanche hazard management strategies (e.g. Germain, 2016), or given particularly severe weather and snow conditions (e.g. Höller, 2009).

Figure 1. Svalbard's physical location at the interface between the Atlantic and Arctic Oceans (top panel). The shaded area in the inset map indicates the Nordenskiöld Land region in central Spitsbergen of primary interest for this research. The photo taken by Markus Eckerstorfer in the bottom left panel shows the aftermath of a fatal recreational avalanche accident near Longyearbyen in 2009. The photo taken by Erik Næss in the bottom right panel shows the fatal December 2015 avalanche in Longyearbyen investigated in further detail in this work.

Such a destructive avalanche event struck Longyearbyen, a small town in the remote, high-Arctic Svalbard archipelago, just prior to this doctoral research's initiation (Figure 1). Longyearbyen at the time lacked formalized avalanche hazard management routines and residents were not evacuated from their homes during a particularly potent winter storm in late 2015. Tragically, two residents died in their homes when a large avalanche naturally released and struck residential infrastructure the morning following the storm. Avalanches had, unfortunately, previously resulted in human fatalities and infrastructure damages in this area. Of particular note, three people were killed in Longyearbyen when a slush avalanche destroyed several buildings including the hospital in 1953 (Hestnes and Kristensen, 2010), and nine people have lost their lives in recreational avalanche accidents in Svalbard since 2000 (Arctic Safety Centre, 2020; Engeset et al., 2020; Svalbardposten, 2019).

The documented history of avalanche accidents and damaged infrastructure on Spitsbergen, the largest island in the archipelago encompassing the region immediately surrounding Longyearbyen (Figure 1), even prior to the 2015 avalanche event demonstrated the clear need for understanding avalanches as a natural hazard in Svalbard. Formal investigations of avalanches as a natural hazard in this location have therefore taken place since the early 1980s (e.g. Hestnes, 2000). More contemporary scientific research built upon this early work to help establish a framework for understanding avalanches in Svalbard's unique environmental setting (Eckerstorfer, 2013). Here, a novel suite of snowpack conditions characterized by the "high-Arctic maritime" snow climate classification (Eckerstorfer and Christiansen, 2011a), a primarily direct-action slab avalanche regime (Eckerstorfer and Christiansen, 2011b), and prevalent cornice-related avalanche activity (Eckerstorfer and Christiansen, 2011c; Vogel et al., 2012) form the conceptual context for this doctoral research.

Although this previous work in central Spitsbergen established a foundation for investigating avalanches as a natural hazard (e.g. Hestnes, 2000) and formally described the region's snow and avalanche climate (e.g. Eckerstorfer, 2013), relatively little avalanche research exists for this region compared to other mountainous locations in more accessible, lower-latitude locations. Ample opportunity to address knowledge gaps in the body of scientific avalanche literature both specific to Svalbard and of more global relevance therefore existed upon the initiation of this doctoral work. Terrestrial laser scanning (TLS) methodology developed in lower latitudes and supporting quantitative analyses of snow accumulation and avalanche activity (e.g. Deems et al., 2015; Prokop, 2008) had, for example, not specifically been applied to Arctic avalanche processes. Similarly, investigations of the synoptic-scale meteorological controls on avalanches had not been conducted for the Svalbard region. Such studies often form the foundation for understanding avalanche regimes and broader climate-avalanche interactions in locations such as, for instance, western North America (Birkeland et al., 2001) and mainland Norway (Fitzharris and Bakkehøi, 1986). Well-studied cornice systems near Longyearbyen (Vogel et al., 2012), finally, presented an opportunity to develop TLS methods to better quantify cornice processes and contribute to the relatively limited body of cornice-specific scientific literature (e.g. Montagne et al., 1968; Munroe, 2018).

Paralleling the research contained in this thesis, avalanche hazard management strategies in Longyearbyen evolved in response to the December 2015 accident. Avalanche forecasting in the area, implemented historically on an ad hoc basis, has since developed into a system where both local, infrastructure-based and regional, recreational advisories are issued daily (Engeset et al., 2020). Construction of permanent structural protections and delineation of new avalanche hazard zones in Longyearbyen have further complemented forecasting efforts in recent years. However, challenges associated with developing hazard management strategies suitable for a remote location with unique avalanche problems highlighted – and continue to emphasize – the operational utility of avalanche research in this high-Arctic setting.

Thesis scope and research objectives

The research included in this doctoral thesis explicitly builds upon previous scientific work from central Spitsbergen – primarily the studies included in Eckerstorfer (2013) – by examining snow and avalanche processes across a range of spatiotemporal scales and presenting the results in the context of improved avalanche hazard management strategies. The focus in this thesis on avalanche hazard stems from two key factors. First, implementation of avalanche forecasting routines and construction of permanent avalanche defense structures in Longyearbyen have paralleled this doctoral work and offer accessible real-world applications for these results. Second, formalized avalanche forecasting provides an established framework for considering the various information and data pertaining to avalanche processes across a range of spatial and temporal scales (LaChapelle, 1980; McClung, 2002b). Consisting of a structured approach to synthesizing weather, snowpack, and avalanche data across the spatial scales relevant for avalanche processes, avalanche forecasting's framework has strongly influenced this work's overall design and implementation. This structure is therefore also useful to help integrate the results of this research via this thesis.

While previous research provided a sound conceptual foundation from which to consider avalanche process and associated hazards in central Spitsbergen, opportunities clearly still existed for conceptual refinement and process quantification – particularly with respect to slope-scale avalanche activity and cornice processes. Furthermore, the existing conceptual framework lacked a broader, synoptic-scale perspective valuable for assessing the impacts of dramatic climatic changes (e.g. Hanssen-Bauer et al., 2019) on the region's snow climate and avalanche regime. Primary research objectives (RO) for this work therefore include:

- (RO1) Demonstrating and quantifying how specific meteorological conditions control avalanche activity in central Spitsbergen across a range of spatial scales;
- (RO2) Upscaling the understanding of surface level meteorological controls on avalanche activity in this location to the synoptic scale; and
- (RO3) Developing and employing monitoring techniques to investigate, document, and quantify snow and avalanche processes, specifically cornices and cornice-fall avalanches, of particular importance in central Spitsbergen, but with global relevance.

Finally, this thesis also seeks to address challenges associated with managing avalanche hazard in remote and Arctic locations such as Svalbard by highlighting and discussing the operational hazard management implications of the results from this research.

Thesis structure

Research presented in three scientific manuscripts comprises this thesis' foundation. Snow avalanche processes are investigated with progressively increasing spatial and temporal coverage in these manuscripts (Figure 2). Paper I concentrates on two specific destructive avalanches in Longyearbyen and covers the finest spatial and shortest temporal scale of the manuscripts. Paper II's research involves monitoring cornice dynamics and cornice-fall avalanches with a terrestrial laser scanner (TLS) at a broader, catchment scale over entire winter seasons. Finally, Paper III encompasses the broadest spatial scale addressed in this work by exploring the relationships between regional avalanche activity and synoptic-scale atmospheric circulation patterns encompassing the broadest spatial scale addressed in this work.

Figure 2. Schematic showing the relative spatial and temporal coverage of the research included in the three papers comprising this thesis.

Background

The present work investigates snow avalanches in the Nordenskiöld Land region surrounding Longyearbyen in central Spitsbergen, Svalbard (Figures 1 and 3). This chapter first presents Svalbard's physiographic and climatological setting before introducing the conceptual snow and avalanche framework on which this research is based. The final section of this chapter further details the knowledge gaps addressed by this doctoral work.

Figure 3. Topographic map of Nordenskiöld Land showing the region's physiographic setting and the locations of settlements and automated weather stations (AWSs) important for this work.

Physiographic setting

Svalbard is an archipelago under Norwegian administration situated 600 km east of Greenland roughly halfway between the Norwegian mainland and the North Pole (Figure 1). The southern Arctic Ocean borders Svalbard to the north, the Barents Sea to the south and east, and the Fram Strait in the Greenland Sea to the west (Figure 1). The archipelago is generally mountainous and glaciated, with approximately 34,000 km² of Svalbard's 62,000 km² land area presently covered by glaciers (Nuth et al., 2013). Spitsbergen ("pointed mountains" in Dutch), the archipelago's largest and only permanently inhabited island, was named by the Dutch explorer Willem Barentsz after sighting the jagged mountain peaks on island's western coast. Such high-relief, alpine topography characterizes the island's northwestern, western coastal, and southern regions, whereas northeastern Spitsbergen is typified by broad glaciated highlands. Nordenskiöld Land, the focus of this thesis, comprises much of Spitsbergen's least glaciated central region, approximately centered on 78°N and 16°E (Figure 3). Svalbard's two largest settlements, Longyearbyen and Barentsburg, are located in north-central and western

Nordenskiöld Land, respectively, and most human activity in the archipelago is consequently also concentrated here.

Broad plateaus punctuated by higher summits and incised by glacio-fluvial valleys largely characterize Nordenskiöld Land's topography (Figure 4). Resistant strata in the horizontally-bedded sedimentary bedrock of Tertiary and late-Cretaceous age control plateau summit elevations near 450 m (Dallmann, 2019), while the region's highest peaks exceed elevations of 1000 m. Glaciation is most extensive in the region's eastern and southern higher elevations and is more confined to isolated cirque glaciers in the northwest. The broadest deglaciated fjord-valleys run generally east-west and are fed by smaller, north-south oriented valleys such as Longyeardalen ("the Longyear valley") – the valley which contains Longyearbyen and is the study area for much of this work. Deglaciation has resulted in steep, concave slopes descending from the plateaus to the valley floors where protruding, resistant bedrock buttresses have been incised fluvial and gravitational slope processes (Larsson, 1982). Continuous permafrost exceeding thicknesses of 500 m in the alpine areas underlies the entire region (Humlum et al., 2003), and no high, woody vegetation grows in Nordenskiöld Land – or elsewhere in Svalbard – due to the harsh climate.

Figure 4. Overview of the topographic setting in Nordenskiöld Land. The top left panel shows the broad plateaus and higher summits east of Longyearbyen, the top right panel shows the slopes descending to a broad fjord-valley south of Longyearbyen, and the bottom panel shows Longyeardalen and the adjacent plateaus under 2020's first sunrise.

Climate and meteorology

Spitsbergen's climate reflects the island's location bordering the North Atlantic and Arctic regions at the interface between ocean currents and air masses with differing thermal characteristics (e.g. Hanssen-Bauer et al., 1990). Here, Spitsbergen's proximity to oceanic and atmospheric poleward heat transport pathways results in the warmest and wettest climate in the high-Arctic (Serreze and Barry, 2014). The West Spitsbergen Current – the Gulf Stream's northernmost extension – conveys warm Atlantic northwards in the eastern Fram Strait and promotes perennial ice-free conditions off Spitsbergen's western coast (Walczowski and Piechura, 2011). Warm Atlantic water flows further from Spitsbergen in the Barents Sea to the east, and sea ice forms annually along the island's northern and eastern coasts. Southern and western Spitsbergen's climate is thus moderated by the open ocean in the Fram Strait while sea ice limits oceanic heat availability and promotes colder conditions in the north and east. Air temperatures across the archipelago thus generally decrease along a southwesterly-to-northeasterly gradient (Figure 5; Vikhamar-Schuler et al., 2019). Similarly, winter and spring precipitation generally decreases with distance from the open ocean to the west, with the highest amounts in the western mountains (Figure 5).

Figure 5. Long-term (1971 – 2000) average seasonal air temperature (°C – note the different color scales) and average cumulative seasonal precipitation (mm) for the winter and spring seasons of importance for this thesis. Figures are from Vikhamar-Schuler et al. (2019) based on Sval_Imp (Schuler and Østby, 2020) data.

Interactions between cyclonic activity along the North Atlantic storm track and semi-permanent high-pressure centers over the Arctic Ocean, Greenland, and Siberia largely control weather patterns over Svalbard. Atmospheric heat and moisture reaching Svalbard during the winter primarily originate in the mid-latitudes and is conveyed northeastwards along the North Atlantic storm track via extra-tropical cyclones (e.g. Hanssen-Bauer et al., 2019; Serreze et al., 2007). Weather patterns, especially in winter, are thus characterized by stable, cold, clear periods during which high pressure dominates punctuated by warm, wet low-pressure systems conveying heat and moisture from lower latitudes. Dominant low pressure in the Norwegian Sea results in easterly prevailing winds over Spitsbergen (Isaksen et al., 2016; Przybylak and Maszewski, 2012). Although observational precipitation data for Spitsbergen is generally sparse and can be unreliable, recent modeling efforts have indicated annual precipitation is highest in Spitsbergen's western mountains and lowest in the fjord-valleys of central Spitsbergen (Figure 5; Humlum, 2002; Vikhamar-Schuler et al., 2019).

Figure 6. Monthly mean air temperature and accumulated precipitation at the Svalbard Airport AWS located 4 km west of Longyearbyen (see Figure 3) for the duration of this doctoral work (MET Norway data).

Observations from the Svalbard Airport AWS taken at the mouth of one such fjord-valley near Longyearbyen represent one of the longest meteorological time-series in the Arctic (Nordli et al., 2020), and are used here to represent conditions in Nordenskiöld Land (Figure 6). For the 1971-2000 climate normal period, annual air temperature at the Svalbard Airport averaged -5.9°C , with the 3-month winter (DJF) and spring (MAM) temperatures averaging -13.9°C and -9.6°C , respectively (Vikhamar-Schuler et al., 2019). Annual precipitation for the same period averages 196 mm and is the lowest value measured at stations in Svalbard. Roughly half this precipitation comes in the winter months of primary interest for this work. The combined effects of systematic undercatch of solid precipitation in Arctic environments and the existence of vertical precipitation gradients in Nordenskiöld Land (e.g. Goodison et al., 1998; Humlum, 2002) indicate the measured precipitation value at the low-elevation Svalbard Airport AWS can be viewed as a minimum value for the mountainous locations investigated in this work. Svalbard's regional easterly prevailing winds also dominate at the Svalbard Airport AWS and are strongest in the winter season (Christiansen et al., 2013; Hanssen-Bauer et al., 2019)

Svalbard's climate is, however, undergoing some of the most pronounced changes in the world. Situated at a warming hotspot even in the context of Arctic Amplification where warming in Arctic regions is enhanced relative to lower-latitudes (Cohen et al., 2014), temperatures across the Svalbard have increased between $3\text{--}5^{\circ}\text{C}$ from 1971 and 2017 (Hanssen-Bauer et al., 2019). Both the warmest year (2016) and the highest ever air temperature recorded at the Svalbard Airport (21.7°C on 25 July 2020) occurred during the duration of this doctoral research (MET Norway data). Precipitation trends are less clear for the most recent decades, but data from the Svalbard Airport shows a statistically significant increase in autumn precipitation during the 1971-2017 period (Vikhamar-Schuler et al., 2019). Rain-on-snow event frequency appears to be increasing (Peeters et al., 2019; Vikhamar-Schuler et al., 2016; Wickström et al., 2020) and extreme precipitation event frequency is also expected to increase in the future (Isaksen et al., 2017). Sea-ice concentration and extent is decreasing (e.g. Onarheim et al., 2014), and many of western Spitsbergen's fjords are now perennially ice-free (Muckenhuber et al., 2016). These changes are projected to continue, with Svalbard becoming progressively warmer and wetter throughout the 21st century (Hanssen-Bauer et al., 2019).

Snow climate and avalanche regime

Avalanches result from complex interactions between an area's terrain, weather, and snowpack (e.g. Schweizer et al., 2003a). Nordenskiöld Land's avalanche regime thus stems from the region's distinctive combination of topographic, meteorological, and snowpack factors. Terrain functions as the fundamental control on avalanche activity by determining the physical locations where avalanches can potentially occur (e.g. slopes steeper than 30 degrees in the case of slab avalanches) but is typically considered static over the timescales of primary relevance for avalanche processes. Weather patterns, then, interact with the underlying topography to control the physical processes which accumulate the seasonal snowpack through snowfall and wind redistribution (e.g. Lehning et al., 2008). Surface meteorological conditions continue to influence snowpack development following snow accumulation by controlling the temperatures differences and vapor movement in the snowpack which lead to snow metamorphism (Armstrong and Brun, 2008). Snow metamorphism can either strengthen a snow layer by promoting bonding between snow crystals or weaken a snow layer by fostering crystal growth such that inter-crystal bonds are much more difficult to form. These weak layers can fail – potentially resulting in a slab avalanche – when overloaded with new or wind-drifted snow (e.g. Davis et al., 1999; Hendrikx et al., 2005), if further destabilized by liquid water from rain or melting (e.g. Baggi and Schweizer, 2009), or due to changing characteristics over the overlying snow slab with increased temperatures (e.g. Reuter and Schweizer, 2012).

Central Spitsbergen's unique climate leads to a novel assemblage of snowpack characteristics previously described as the high-Arctic maritime snow climate (Eckerstorfer and Christiansen, 2011a). Here, relatively dry conditions and generally low air temperatures result in a cold, thin snowpack (Figure 7). Depth hoar develops early in most winter seasons and persists as a structural weakness in the snowpack throughout the 8-10 months snow covers the ground (Eckerstorfer and Christiansen, 2011a). Wind readily transports snow across the barren landscape (Jaedicke, 2002), and wind slabs are accordingly a dominant stratigraphic feature in the snowpack (Figure 7, Figure 8). These characteristics are suggestive of more commonly recognized snow climates such as the continental snow climate (Mock and Birkeland, 2000) or tundra snow cover type (Sturm et al., 1995). However, warm winter storms accompanied by rain and the resulting prevalent ice layering in the Svalbard snowpack represent a strong maritime influence typically absent from continental or tundra snow climates (Figure 7, Figure 8). The unique combination of maritime and continental snowpack characteristics differentiates high-Arctic maritime conditions from other, more commonly recognized snow climates. A representative high-Arctic maritime snowpack is therefore composed of basal depth hoar overlain by dense wind slabs and ice layers separated by faceted grains developing during cold periods of stable weather (Figure 7).

Figure 7. An idealized snow profile showing a typical high-Arctic maritime snowpack based on the findings of Eckerstorfer and Christiansen (2011a) and observed snow stratigraphy during this work with symbology after Fierz et al. (2009). The hand hardness index is a field measurement of snow hardness representative of the object which can be pushed into the snow with moderate force (e.g. a knife hardness layer is harder than a fist layer because a knife can be pushed into the layer with moderate force, but a fist cannot).

Nordenskiöld Land's avalanche regime stems from the region's distinctive combination of topographic, meteorological, and snowpack factors. Snow transported by the wind across the broad plateau summits accumulates as cornices at the plateau margins, with the largest cornices forming on the westerly aspects situated leeward from the prevailing southeasterly winds (Figure 8). Cornice-fall avalanches – avalanches occurring due to cornice failure – comprised the most commonly observed avalanche type in previous avalanche monitoring campaigns (Figure 9; Eckerstorfer and Christiansen, 2011c). Previous studies observed cornice-fall avalanches to cluster near the end of the winter season as air temperatures warmed (Eckerstorfer and Christiansen, 2011c; Vogel et al., 2012). Dry slab avalanche activity (Figure 8, Figure 9) is understood to be largely direct-action, coinciding with winter storms during which precipitation as snow and strong winds result in snowdrift and rapid wind slab development in leeward areas (Eckerstorfer and Christiansen, 2011b). Finally, warm winter storms during which precipitation falls as rain can result in wet avalanches and slush avalanches at any point during the snow season (Figure 9; Eckerstorfer and Christiansen, 2012).

Figure 8. Photos illustrating the high-Arctic maritime snow climate and avalanche regime. Counter-clockwise from top left: snow blowing over the edge of the Gruvefjellet plateau, an ice layer in the snowpack, a wind slab overlying faceted snow, a slab avalanche releasing under a cornice, and the cornices on Gruvefjellet.

This conceptual model for the high-Arctic maritime snow climate and avalanche regime described above is based primarily on research conducted nearly a decade ago. Given Svalbard's changing climatic situation, changes to the snow climate can also be expected – and have likely occurred since conceptual model's development. Interactions between increased winter air temperatures (Hanssen-Bauer et al., 2019), more frequent rain-on-snow events (Peeters et al., 2019; Vikhamar-Schuler et al., 2016; Wickström et al., 2020), shorter snow seasons (e.g. López-Moreno et al., 2016), and changing winter storm characteristics and frequency (e.g. Wickström et al., 2019) are shaping and – will continue to shape – the snow climate and avalanche regime in the future. While recent investigations have documented changes to snowpack physical properties related to climatic shifts (Peeters et al., 2019), changes to the avalanche regime have not yet been specifically studied.

Figure 9. Photos showing representative snow avalanche activity in central Spitsbergen including a cornice-fall avalanche (top panel), a slab avalanche (bottom left), and a slush avalanche (bottom right, Markus Eckerstorfer photo).

Knowledge gaps

Motivation for this work stems from a lack of modern scientific avalanche research in Svalbard and other polar locations, the importance of cornices – a globally understudied snow phenomena – to central Spitsbergen’s avalanche regime, and the value of understanding the impacts of Svalbard’s rapidly changing climate on avalanche activity here. Specifically, this work helps address scientific knowledge gaps related to 1) limited work linking avalanche activity to well-quantified snow accumulation and distribution patterns in Arctic settings, 2) lacking scientific studies focused on snow cornice process quantification, and 3) synoptic-scale atmospheric controls on avalanche activity in central Svalbard. Finally, the December 2015 avalanche underscored the clear need for hazard management improvements near Longyearbyen to which this research could contribute.

The dearth of modern, quantitative cornice research offered a compelling knowledge gap in which TLS techniques employed in Svalbard could be used to quantitatively refine existing conceptual cornice process models (Montagne et al., 1968; Vogel et al., 2012). TLS had also, prior to the initiation of the present work, been used in other locations to precisely measure snow surface changes resulting from avalanches to determine dynamic avalanche modeling parameters (Prokop et al., 2015) and to better understand and model how the wind redistributes snow (Schön et al., 2015). In Svalbard, snow drift processes in complex terrain had been studied and modeled (Jaedicke, 2001; Sauter et al., 2013) but were not specifically linked to avalanche activity. TLS techniques again offered possibilities to close this knowledge gap by better quantifying snow drift accumulations in avalanche terrain and linking these accumulations to avalanche release.

The impact of ongoing, dramatic climatic changes on Svalbard’s snow climate and avalanche regime also presented a knowledge gap upon initiating this research. Even in areas with better snow monitoring systems and longer historical avalanche activity records, resolving research questions related to climate change and avalanche activity is challenging due to complex relationships between air temperatures, precipitation, snowpack characteristics, and snow stability (e.g. Bellaire et al., 2016). Svalbard previously lacked scientific research relating broad, synoptic-scale atmospheric circulation patterns to increased avalanche activity. Such studies in other regions (e.g. Birkeland et al., 2001; Fitzharris and Bakkehøi, 1986; García et al., 2009) form a basis for understanding avalanche activity at the broader spatial scales where links between climate models and snow and avalanche processes are best resolved. Addressing the knowledge gap regarding synoptic controls on avalanches in Svalbard also provided a way to link avalanche processes with work in Svalbard studying relationships between synoptic atmospheric circulation and recent warming (e.g. Isaksen et al., 2016) and precipitation patterns (Serreze et al., 2015). Thus, providing a framework for understanding the synoptic controls on

snow avalanches in Spitsbergen can serve as a first-order attempt at more specifically understanding climate change's effects on avalanche activity here.

Prior to the December 2015 avalanche event and, subsequently, the initiation of this doctoral project, avalanche hazard management in Longyearbyen primarily involved contractual hazard assessments performed by the Norwegian Geotechnical Institute (NGI) on an ad hoc basis (Hestnes, 2000; Hestnes et al., 2016). More permanent hazard management strategies included construction of an earthen deflecting dam shortly after the 1953 accident to protect infrastructure from the slush avalanches which run down a side valley into the eastern portion of Longyearbyen. Additionally, a bulldozer removed – and continues to remove – snow in the drainage channel of this side valley each season prior to the onset of the spring melt. However, despite numerous reports authored by NGI staff evaluating avalanche hazard, defining hazard zones, and recommending suitable structural defenses and avalanche forecasting routines (e.g. Hestnes, 2000; Hestnes et al., 2016), limited implementation of recommended strategies occurred. As a result, avalanche forecasting routines were not in place when the December 2015 occurred, and evacuations were not ordered (Hestnes et al., 2016).

Local authorities and the Norwegian Water Resources and Energy Directorate (NVE) again contracted NGI to establish a daily forecasting service for threatened infrastructure in Longyearbyen in the aftermath of the December 2015 avalanche (Engeset et al., 2020; Jaedicke et al., 2016). This arrangement continued through January 2016, when NVE's national avalanche forecasting service expanded to include Nordenskiöld Land in their regional avalanche bulletins. NVE also took responsibility for issuing local avalanche forecasts for the infrastructure in Longyearbyen from January 2016 to January 2019, when the consulting firm Skred AS took over responsibility for the local forecasts (Engeset et al., 2020). At present, NVE issues daily regional forecasts daily for Nordenskiöld Land throughout the winter season, while Skred AS issues daily local, infrastructure-specific forecasts for Longyearbyen.

The research presented in this thesis thus coincided with the implementation of formalized avalanche forecasting routines in central Spitsbergen. Furthermore, a second destructive avalanche struck Longyearbyen in February 2017, reinforcing the challenges associated with avalanche hazard management in this location and prompting the development of additional permanent structural mitigation measures. In addition to addressing the scientific objectives guiding this research, this work has therefore also sought to support improved avalanche hazard management strategies in Longyearbyen by refining the understanding of Arctic avalanche processes, developing new monitoring routines, and improving operational data availability.

Methods

Terrestrial laser scanning

Terrestrial laser scanning (TLS) methods comprised the primary fieldwork and resulted in the datasets unique to Papers I and II. TLS, or ground-based lidar, is a remote sensing technique for measuring distances with light pulses and creating accurate 3D representations of the target surface. As an active remote sensing technology, the TLS emits a pulse of light and accurately measures the time it takes the emitted pulse to travel to the surface to be measured, reflect off the surface, and travel back to the TLS (Figure 10). This two-way travel time in combination with information about the direction in which the pulse was emitted can be used to determine the precise location of a point on the surface. The scanner rotates through a “scan window” on a vertical axis, scanning a swath of the surface and recording the position of millions of points. The final output from the scanner is a dense point cloud representing the measured surface.

Figure 10. Terrestrial laser scanning methods in Longyearbyen. The top panel shows the conceptual basis for TLS methods in a modified image of the scanner at the Sukkertoppen scan position. The bottom left and right panels show the TLS scanning Gruvefjellet and Platåberget, respectively.

Research included in this thesis involved repeatedly scanning the three mountain slopes – Sukkertoppen, Gruvefjellet, and Platåberget – surrounding Longyearbyen during the 2016/2017 and 2017/2018 winters, with one scan of Sukkertoppen from January 2016 also included (Figure 11). We used a Riegl® VZ-6000 scanner for all scans included in this work (Figure 10). The VZ-6000 is an ultra-long range TLS, particularly suitable for acquiring data from snow and ice covered surfaces due to its near-infrared operating laser wavelength and high scanning speed. In total, five scanned surfaces of Sukkertoppen (one bare-earth and four snow-covered scans), 25 scans of Gruvefjellet (one bare-earth and 24 snow-covered scans), and 22 scans of Platåberget (one bare-earth scan and 21 snow-covered scans) comprise the basis for Papers I (Sukkertoppen) and II (Gruvefjellet and Platåberget).

Figure 11. Overview of the scan locations and scanned areas in Longyeardalen included in this thesis. Snow depth stations used in Paper II are also indicated.

TLS has been increasingly used to gather information about snow-covered surfaces (e.g. Deems et al., 2013; Prokop, 2008). Repeatedly scanning the same area allows for the quantification of changes to the snow cover by differencing co-registered surfaces, represented either as a processed data product like a digital elevation model (Figure 12), or on the raw point cloud itself (Figure 13). Snow depth maps, a common TLS-based snow data product, are traditionally derived by first interpolating digital elevation models (DEMs) from the georeferenced raw point clouds acquired at times when the surface lacked snow (bare-earth) and those acquired at different times with snow cover. The resulting DEMs represent the surface elevations without snow (bare-earth) and the surface elevations of snow-covered surfaces. Snow depths are then calculated for each collocated grid point in the DEMs by subtracting bare-earth elevations from the those of the snow-covered surface (Figure 12). Differential snow depth data products can similarly be derived by calculating elevation differences between snow-covered surfaces taken at different times, and are useful for investigating the impacts of, for example, specific meteorological conditions or avalanche activity on snow distribution patterns (e.g. the avalanche shown in Figure 12).

Figure 12. The basic workflow concepts used to derive spatially-distributed snow depth (HS) and change in snow depth (dHS) data from surfaces scanned with a TLS. The included maps taken from Paper I illustrate the snow depth on Sukkertoppen following a destructive avalanche in 2017 (left panel) and the change in snow depth in the week leading up to and including the avalanche (right panel).

Vertical snow surface changes calculated with the techniques described above are suitable for most snow monitoring applications concerned primarily with snow depth. Changes to the cornices investigated in Paper II, however, are very often non-vertical. The most interesting and relevant surface changes include horizontal growth which the workflow shown in Figure 12 would not adequately represent. Quantifying changes to the cornices therefore required development of separate, novel workflow which nevertheless follows the same principle of comparing bare-earth and various snow-covered surfaces (Figure 13). Georeferenced point clouds representing the cornice surfaces with considerable horizontal extension (Figure 13a) are first imported into CloudCompare software (Figure 13b; CloudCompare, 2019). Cross-sectional profiles can be selected from this point cloud to visualize cornice morphology or calculate horizontal changes to the cornices at specific locations in the cornice system (Figure 13c, Figure 13d). The real benefit of TLS data acquisition, however, comes from acquiring this type of data across an entire slope (Figure 13e and Figure 13f). Employing the Multiscale Model-to-Model Cloud Comparison (M3C2) algorithm (Lague et al., 2013) plugin in CloudCompare allows for the calculation of three-dimensional (including both horizontal and vertical changes) change across the entire scanned area (Figure 13g). Further detailed information related to these methods is found in Paper II.

Figure 13. Conceptual overview of the workflow used to quantify changes to the cornices and underlying slopes in Longyeardalen for Paper II. The photograph in panel (a) shows the front view of a cornice with considerable horizontal extension, and Panel (b) shows the same cornice represented with TLS data (here the point cloud is interpolated to a surface for visualization purposes). Panel (c) shows the raw point cloud (colored for visualization) with the profile displayed in Panel (d) highlighted. Panel (d) displays the profile indicated in Panels (a), (b), and (c) with the green line compared to the bare-earth surface in white. Panels (e) and (f) expand panels (a) and (b), the spatial extent of which are indicated with red rectangles. Panel (g) displays the results of an M3C2 calculation from a before and after scan of a cornice-fall avalanche on Gruvefjellet.

regObs

All three papers included in this thesis incorporate data from the regObs database. RegObs, which stands for *registration* of *observations*, is the Norwegian Water Resources and Energy Directorate's (NVE) system for sourcing natural hazards observations from both professional and amateur observers. RegObs consists of a website (<https://www.regobs.no>), a mobile app, and a database accessed via an application programming interface (Engeset et al., 2018). Of interest for this work were the snow and avalanche observations available on the regObs database. These observations are employed both by the Norwegian Avalanche Warning Service to develop regional avalanche forecasts and by local hazard managers to manage avalanche hazard in Longyearbyen. Observations submitted to regObs can include information related to snow cover, snow stratigraphy in snow profiles, stability tests, danger signs, avalanche activity, avalanche hazard assessments, and weather. Avalanche observations from regObs form the basis of the avalanche activity record developed for Paper III, snow profiles registered on regObs from the December 2015 and February 2017 avalanche crowns facilitated idealized snow profile creation in Paper I, and avalanche observations helped better temporally constrain the cornice-fall avalanches investigated in Paper II

Weather and snow depth data

Data from the Svalbard Airport and Gruvefjellet automatic weather stations (AWS) near Longyearbyen support all three papers (Figure 3). The Svalbard Airport AWS situated 4 km west of Longyearbyen at 28 m elevation has been recording standardized meteorological parameters since 1975 and is the only station in the region with long-term precipitation time series measurements. Data from this station is available through the Norwegian Meteorological Institutes data clearinghouse at www.eklima.no. Data from the Gruvefjellet AWS located on the Gruvefjellet plateau at 464 m better represents wind and air temperature conditions at higher elevations, but do not include precipitation measurements. The Gruvefjellet AWS data are available for download at <https://www.unis.no/resources/weather-stations>. Paper II also includes data from a network of automated snow monitoring stations in Longyeardalen which remained in place only for the 2017/2018 winter season (Figure 11; Prokop et al., 2018).

ERA5 reanalysis data served as the primary meteorological input for Paper III. ERA5 is a gridded global reanalysis product from the European Centre for Medium-Range Weather Forecasts (ECMWF) within the Copernicus Climate Change Service. ERA5 reanalysis products are available at a 31 km spatial resolution with hourly temporal resolution (Hersbach et al., 2020). Paper III incorporates ERA5 data regarding mean sea level pressure (MSLP), air temperature at 2 m above the surface, accumulated precipitation, and wind speed at 10 m above the surface to in combination with the Niedźwiedź Classification to characterize synoptic conditions around Svalbard. The Niedźwiedź Classification (Niedźwiedź, 2013) is a daily calendar of common synoptic situations over Svalbard defined by Tadeusz Niedźwiedź from the University of Silesia in Katowice, Poland. These situations are defined based on the geostrophic wind direction assessed via the MSLP pressure field and whether the airflow is primarily driven by cyclonic or anti-cyclonic circulation.

Research summary

Paper I

Combining high spatial resolution snow mapping and meteorological analyses to improve forecasting of destructive avalanches in Longyearbyen, Svalbard

Hancock, H., Prokop, A., Eckerstorfer, M., and Hendrikx, J., 2018. Combining high spatial resolution snow mapping and meteorological analyses to improve forecasting of destructive avalanches in Longyearbyen, Svalbard, *Cold Regions Science and Technology*, 154, 120-132, doi:<https://doi.org/10.1016/j.coldregions.2018.05.011>.

This work, published in *Cold Regions Science and Technology* in 2018, explores the naturally triggered avalanches which released from Sukkertoppen in December 2015 and February 2017. Both avalanches struck inhabited infrastructure in Longyearbyen and were preconditioned by similar meteorological conditions. The December 2015 event tragically resulted in the deaths of two residents in their homes, while major injuries were fortunately avoided in February 2017. We investigated these events as case studies to specifically address the meteorological controls on slope-scale snow distribution in this location (RO1), while demonstrating TLS as an effective technique for event documentation and analysis in an Arctic setting (RO3). Using snow surface measurements obtained via TLS, we developed snow depth and/or differential snow depth maps for Sukkertoppen's western aspect for each avalanche event. These data, in combination with more traditional weather and snowpack analyses, allowed us to assess, describe, and illustrate the slope-scale meteorological controls on destructive avalanche activity on this avalanche path.

We identified strong easterly winds combined with moderate to heavy snowfall and considerable snow available for transport as the meteorological precursors to destructive avalanche activity here. Relatively minor differences in wind direction, snowfall rate, and air temperature prior to the events helped to explain differing snow accumulation patterns and resulting avalanche activity between the two events. These results demonstrate the critical control wind exerts on snow accumulation and avalanche activity near Longyearbyen, with implications for understanding these processes in other Arctic and alpine locations. Finally, we used these avalanche events to illustrate how limited scale-appropriate data and poor observation density combined with uncertainty due to climatic changes and inadequate historical avalanche records have impacted avalanche forecasting in Longyearbyen.

Paper II

Quantifying seasonal cornice dynamics using a terrestrial laser scanner in Svalbard, Norway

Hancock, H., Eckerstorfer, M., Prokop, A., and Hendrikx, J., 2020. Quantifying seasonal cornice dynamics using a terrestrial laser scanner in Svalbard, Norway, *Nat. Hazards Earth Syst. Sci.*, 20, 603-623, doi:10.5194/nhess-20-603-2020.

Published in *Natural Hazards and Earth System Sciences* in 2020, this research focused on developing effective techniques to monitor and quantify seasonal cornice dynamics (RO3). We employed TLS over two winters (2016/2017 and 2017/2018) to repeatedly scan the cornice systems which form each winter season on the plateau margins comprising Longyeardalen's eastern and western borders. TLS techniques produced snow surface data at fine enough spatial and temporal resolutions to allow us to observe and quantify changes to the cornice systems in detail not previously achieved. Spatially distributed snow surface data with sub-decimeter accuracy provided a robust basis for investigating the meteorological controls on cornice dynamics and associated avalanche activity at the catchment spatial scale (RO1). We found horizontal cornice accretion rates can exceed 15 mmh^{-1} under ideal conditions in which wind speeds are sufficient to transport snow from directions placing the plateau edge in the lee. Cornice accretion preceded five of the six cornice-fall avalanche events observed in this study, and none of the events included avalanche activity on both the west-facing and east-facing cornice systems.

These results demonstrated a noteworthy relationship between meteorological conditions related to winter storms, cornice accretion, and cornice-fall avalanches. Here, winter storms provide the necessary snow available for transport and wind to promote cornice accretion, which in turn can result in cornice failure and cornice-fall avalanches. Wind direction exerts a critical control on the specific aspects on which cornice accretion occurs. This relationship can help hazard managers seeking to forecast cornice-fall avalanches near Longyearbyen better anticipate the conditions and locations cornice accretion and associated cornice failure. Methodological contributions from this work included demonstrating TLS's utility for monitoring cornice processes –especially in locations where the long polar night precludes direct visual observation – and presenting new methods for quantifying cornice surface and volume changes which can serve as a foundation for future cornice-related studies.

Paper III

Synoptic control on snow avalanche activity in central Spitsbergen

Hancock, H., Hendriks, J., Eckerstorfer, M., and Wickström, S., 2021: Synoptic control on snow avalanche activity in central Spitsbergen, *The Cryosphere*, 15, 3813-3837, doi:10.5194/tc-15-3813-2021.

The third manuscript included in this thesis, published in *The Cryosphere*, investigated the synoptic-scale meteorological controls on avalanche activity in Nordenskiöld Land (RO2). This work involved two main research tasks. First, we constructed a daily avalanche activity record for Nordenskiöld Land over four winter seasons (2016/2017 – 2019/2020) from crowd-sourced observations available on the regObs portal, the Norwegian Water Resources and Energy Directorate's (NVE) online natural hazards observation registration system. We then analyzed the synoptic atmospheric conditions across a broad portion of the North Atlantic on days with differing avalanche activity situations using ERA5 reanalysis data in combination with a well-established subjective atmospheric circulation classification system developed for Svalbard.

The results from this work largely support the understanding of meteorological controls on avalanche activity for central Spitsbergen developed at finer spatial scales (RO1). We found synoptic atmospheric conditions conducive to windier, wetter, and warmer weather near Svalbard are associated with increased avalanche activity in Nordenskiöld Land (RO2). Different synoptic circulation patterns and associated upwind conditions help explain differences between days characterized by dry and wet avalanche activity. We identified anomalously low sea-level pressures just west of Spitsbergen – and the resulting increased wind speeds and precipitation – as the aggregate synoptic signal for dry high avalanche activity days. Wet high-activity days were related to very warm air diverted northwards along a high-pressure ridge over Fennoscandia in combination with isolated positive precipitation anomalies near Spitsbergen's west coast. These results provide a synoptic framework helpful both for avalanche forecasting based on synoptic-scale weather forecasts and for placing avalanches within the broader context of Arctic environmental change.

Synthesis and discussion

The three papers comprising this thesis investigate the controls on avalanche activity in central Spitsbergen via different methods and at different spatial and temporal scales. This chapter synthesizes and discusses the results from the papers to provide a cohesive overview of this doctoral work. The first section places this research within the established framework used in avalanche forecasting to consider avalanche-related information. The second and third sections detail how this work addressed the desired research objectives and discuss how the findings relate to existing snow avalanche research. Finally, the fourth section discusses the realized and potential contributions from this work towards improving avalanche hazard management strategies in Svalbard.

The avalanche forecasting framework

Formalized avalanche forecasting involves a structured approach to evaluating current and predicting future avalanche conditions. This process entails synthesizing weather, snowpack, and avalanche occurrence data across the spatial and temporal scales relevant for avalanche processes (LaChapelle, 1980; McClung, 2002b). Although developed primarily for operational applications, the avalanche forecasting framework strongly influenced this research's overall design and implementation. Thus, this framework is also valuable for contextualizing and synthesizing the scientific contribution from this research across the spatiotemporal scales investigated in this work.

Figure 14. Modified figures from McClung (2002b) showing the conceptual framework, organization, and relative influence at different spatial scales of the data types used in avalanche forecasting. The left panel shows the classes into which information about avalanches can be separated. The right panel shows how information from each class is incorporated to develop an avalanche forecast for different spatial scales.

Avalanche forecasters seek to recognize instabilities in the snowpack which lead to avalanche release by assessing direct snow stability observations (Class I information), snowpack structure (Class II information), and meteorological data (Class III information; LaChapelle, 1980; McClung, 2002b). These data sources have traditionally been separated into three classes, with descending class number indicating increasingly direct evidence of instability (Figure 14). Forecasting avalanche conditions necessitates integrating data from all three classes across a range of spatial and temporal scales. Typical spatial scales relevant to avalanche forecasting are the synoptic, meso, and micro scales. These scales are roughly analogous to entire mountain ranges or regions, mountain massifs or valley catchments, and individual slopes, respectively (McClung, 2000; McClung, 2002b). Avalanche processes and their associated forecasts similarly occur along a temporal continuum including the seconds in which avalanche release occurs, the hours and days which snow loading can occur during winter storms, and the weeks and months over which snowpack conditions conducive to avalanche release develop during the winter season (e.g. Hägeli and McClung, 2004; Statham et al., 2018). The avalanche forecasting problem's spatiotemporal scale thus largely dictates the relative influence of the three data classes – based largely on scale-appropriate data availability – to understanding the physical processes leading to avalanche release at, and developing a forecast for, the given spatial scale. As a result, readily available synoptic-scale Class III weather data will be of greater relative importance to a regional avalanche forecast than to a slope-scale forecast where Class II and Class I instability data must be incorporated to achieve adequate forecasting precision (Figure 14).

Placed in a similar framework, the three papers comprising this work cover a broad portion of the spatiotemporal continuum (Figure 15). Paper I investigates two destructive avalanches on Sukkertoppen's western aspect corresponding to a micro, slope-scale study. The avalanches studied in Paper I happened in a matter of seconds, the snow loading during winter storms took place the hours and days prior to the avalanche release, and weak layer development occurred over the preceding weeks and months. The Longyeardalen cornice systems encompass a catchment, meso-scale area explored in Paper II. Cornice development monitored for this publication spans entire winter seasons, with winter storms occurring over hours to days exerting a strong control on the seasonal cornice dynamics. Although catastrophic cornice failure and associated avalanche release occurs over seconds, our data's temporal resolution for these events is not as fine as for the avalanches in Paper I and the minimum time scale for this work is thus defined as hourly. Paper III focuses on a much broader spatial scale by analyzing how synoptic-scale atmospheric circulation controls regional avalanche activity across Nordenskiöld Land. This manuscript primarily relies on daily atmospheric circulation patterns and avalanche occurrence. Thus, while the analyses are conducted across multiple complete winter seasons, the key temporal scale investigated in Paper III is daily.

Figure 15. The spatiotemporal setting for the three manuscripts included in this thesis. Primary data from each data class employed in each paper are indicated, with the relative influence of the different data types to the conclusions in each work represented as in Figure 15.

Meteorological controls on avalanche activity in central Spitsbergen

Two research objectives addressed by this work (RO1 and RO2) involve investigating the meteorological controls on avalanche activity in central Spitsbergen. Understanding how weather controls avalanche activity is critical to evaluating future avalanche conditions because numerical weather forecasting products are typically the best resolved information related to future instability available to avalanche forecasters. Analyzing how previous weather conditions (Class III information) have influenced historical avalanche activity (Class I information) and past snowpack development (Class II information) can therefore help clarify how forecasted weather might influence future avalanche activity in a particular location. Papers I and II employ TLS-derived Class I and II data to build upon previous work investigating how weather conditions influence avalanche activity in central Spitsbergen at the slope and catchment spatial scales (RO1), while Paper III uses data with broader spatial coverage to upscale this understanding to the synoptic, regional scale (RO2; Figure 15). The following first synthesizes the primary results related to RO1 and RO2 and then places these findings more broadly in the context of the high-Arctic maritime snow climate.

The influence of snow and wind on avalanche activity in central Spitsbergen

The findings from this research show interactions between precipitation and wind act as a fundamental control on avalanching in central Spitsbergen at all spatial scales by influencing where and when snow loading events occur. Central Spitsbergen's dry, windy climate promotes a snow-limited environment. Here, wind consistently in excess of the $5\text{--}7\text{ ms}^{-1}$ threshold wind speed for snow transport (e.g. Pomeroy, 1989) readily redistributes and hardens the existing surface snow, limiting future transport. Fresh snowfalls, typically also coinciding with wind, largely control when snow is available for transport and when hazardous avalanche conditions can develop from loading events. Previous work established the strong relationship between blowing and drifting snow and avalanche activity in central Spitsbergen (Eckerstorfer and Christiansen, 2011b; Eckerstorfer and Christiansen, 2011c) but the present work refines this understanding by specifically demonstrating this relationship at the synoptic scale (RO2) and quantitatively illustrating this relationship with TLS data at the catchment and slope spatial scales (RO1). This discussion focuses primarily on dry avalanches, as only Paper III includes considerable research into the meteorological controls on wet avalanches.

At the broadest spatial scale investigated in this work, Paper III's results show how cyclonic activity near Svalbard promotes the windy, snowy synoptic environment associated with elevated regional avalanche activity in Nordenskiöld Land. Low pressure systems following the North Atlantic storm track transport moisture from lower latitudes to the Svalbard region which, combined with the wind characteristic of these systems, create the winter storm conditions commonly related to avalanche release in Nordenskiöld Land. Avalanche research has identified atmospheric circulation patterns leading to winter storms (when wind and

snowfall are plentiful) as a synoptic control on avalanche activity in other locations such as western North America (Birkeland et al., 2001) and the Eastern Pyrenees (García et al., 2009). These results are unsurprising as new snowfall and wind are well-established contributory factors to avalanche release in most snow climates (Davis et al., 1999; Hendrikx et al., 2005; Schweizer et al., 2003a). In Paper III, specifically linking avalanche activity in central Spitsbergen to snow and wind at the synoptic scale helps demonstrate how winter storms also generally determine avalanche timing here via a direct-action avalanche regime. Paper III's results have implications for understanding avalanche activity at finer spatial scales (discussed below), but also help tie a high-Arctic avalanche setting into the body of work investigating synoptic-scale meteorological controls on avalanche activity (e.g. Birkeland et al., 2001; Germain et al., 2009; Hansen and Underwood, 2012; Keylock, 2003; Schauer et al., 2021).

Snow and wind arriving to Svalbard via low pressure systems further control avalanche activity at finer spatial scales. The low pressure's position in relation to Spitsbergen will generally control the regional wind direction characterizing conditions on Nordenskiöld Land's broad plateaus where topographic channeling is minimal (e.g. Christiansen et al., 2013). Paper II's results show cornices accrete – and cornice-fall avalanches occur – on the slope leeward to the dominant regional wind direction during winter storms, while cornice scouring occurs on the windward slope. This finding links cornice-fall avalanches and cornice accretion to the regional winds – and snow available for transport – controlled largely by synoptic-scale cyclonic activity. Combined with previous work indicating cornice-fall avalanches cluster near the end of the snow season due to rising air temperatures (Vogel et al., 2012), these results indicate cornice failures in catchments near Longyearbyen can be differentiated between temperature-induced failures and accretion-induced failures (e.g. Munroe, 2018). Thus, different meteorological conditions control 1) the prolonged elevated air temperatures responsible for temperature-induced failures near the snow season's end and, 2) accretion-induced failures related to drifting snow at any point in the season once cornices have begun to form.

Finally, Paper I's results show how relatively minor differences in otherwise similar meteorological conditions can have demonstrable effects on snow accumulation and resulting avalanche activity at the slope scale. The regional winds – or regional winds modified via valley channeling (Christiansen et al., 2013) – and snow available for transport interact with local topography to control snow accumulation at this spatial scale. The resulting snow accumulation patterns have important implications for avalanche release, with specific slab release locations observed in Paper I governed, in part, by locations with the greatest snow accumulations. Although previous work found a clear temporal trend with avalanche activity clustered near the snow season's end (Eckerstorfer and Christiansen, 2011c), this work shows destructive avalanches can, superimposed upon the overall seasonal trend, release at any time given strong enough winds, sufficient snow available for transport, and the presence of a suitable weak layer in the underlying snowpack.

Loading events and the high-Arctic maritime snow climate

Central Spitsbergen's avalanche regime broadly reflects the dichotomy in the region's winter weather: dry slab avalanches typically occur when wind-drifted snow accumulates during maritime winter storms and overburdens weak layers which have formed in the snowpack during colder, stable weather patterns. As shown in Paper I, the December 2015 event illustrated an archetypal, albeit dramatic, high-Arctic maritime avalanche scenario in which rapid, intense loading on a thin, structurally weak snowpack resulted in the failure of a deep persistent weak layer. More gradual, less intense loading during the February 2017 event, by contrast, triggered a layer of newly formed facets under the storm snow. For this event, a thick melt-freeze crust which developed during a mid-winter rain event several weeks prior to the avalanche prevented failures on deeper instabilities. Rapid faceting likely occurred a subsequent cold snowfall buried this crust (e.g. Jamieson, 2006) and contributed to widespread slab propagation.

Both destructive avalanche events described in Paper I highlight how interactions between the existing snowpack – which reflects antecedent weather – and the specific loading patterns during a winter storm interact to control avalanche activity at the slope scale in this snow climate. Although incorporating antecedent weather and snowpack data into the analyses in Paper III fell outside the scope of this research, interactions between the snowpack and the specific loading events demonstrated in Paper I may be broadly reflected at the regional scale. A sequence of synoptic atmospheric circulation patterns promoting weak layer development – for example, via creation of a facet over crust stratigraphy as in the February 2017 scenario – prior to the loading event could be expected to produce widespread avalanche activity. A loading event with the same synoptic signal occurring on a snowpack preconditioned for stability, on the other hand, will likely result in fewer avalanches regionally. Future studies in Svalbard could reflect work by Schauer et al. (2021) on the synoptics of deep slab avalanches in the western United States to explore how synoptic atmospheric circulation patterns precondition the snowpack prior to the loading events which ultimately trigger the avalanches.

The underlying snowpack also likely helps control cornice-fall avalanche size, although lacking Class II snow stratigraphy data in Paper II makes resolving this relationship difficult in the context of this research. Cornice-fall avalanches coinciding with loading events documented in Paper II did not exceed destructive size 2, however, as none of the observed cornice failures triggered avalanches on deeper instabilities or resulted in widely propagating slabs. Similar cornice failures impacting a snowpack with widespread, sensitive weak layers or reactive deep instabilities can thus be expected to trigger larger slabs and entrain more snow, resulting in considerably more destructive cornice-fall avalanches. Future cornice-fall avalanche work should thus seek to integrate either observed or modeled snowpack data and quantitative cornice data to better resolve the relationship between cornice failure, underlying snowpack stratigraphy, and the resulting cornice-fall avalanche activity.

Methods for snow and avalanche observation in Arctic settings

Terrestrial laser scanning

This work employed TLS methods to investigate, document, and quantify snow and avalanche processes at the catchment and slope scales to address RO3. Snow depth data products with high spatial resolution and slope-to-catchment scale spatial coverage have been a primary focus for many previous snow-related TLS studies (e.g. Deems et al., 2013; Prokop, 2008). Demonstrated applications for TLS-derived snow depth data products include assessing slope and catchment scale snow distribution (Fey et al., 2019; Prokop et al., 2008), quantifying snow drift processes (e.g. Schneiderbauer and Prokop, 2011; Schön et al., 2015), and deriving dynamic avalanche modeling parameters from accurate avalanche measurements (Prokop et al., 2015). Building upon these and other previous works, the methods used in Papers I and II demonstrate TLS as a valuable method for acquiring snow surface information in central Spitsbergen. TLS data acquisition techniques specifically provided unique snow surface measurements of destructive avalanche activity (Paper I) and cornice dynamics (Paper II). A novel workflow developed to process the latter measurements serves as a major methodological contribution from this work.

TLS offers notable advantages over other remote sensing methods – including visual observation – in central Spitsbergen and other high-latitude environments where much of the snow season is shrouded in the dark polar night. The lack of visible light during the early snow season precludes effective snow and avalanche observation through alternative measurement techniques employed in lower latitudes at the slope and catchment scale such as unmanned aerial system photogrammetry (Adams et al., 2018) or time lapse photography from stationary cameras (Parajka et al., 2012). TLS thus can be used to gather spatially-distributed snow surface data across the snow season's entirety in Svalbard, providing an avenue to complement or refine – as in Paper II – studies using time lapse photography to monitor seasonal cornice development at lower latitudes (Munroe, 2018; van Herwijnen and Fierz, 2014) or in Svalbard (Vogel et al., 2012).

Future work should attempt to develop automated workflows to build upon the TLS methods employed in this research. Automating TLS data acquisition with a permanently installed TLS programmed to scan at regular intervals would help reduce the time interval between scans which introduced uncertainty and limited result interpretation in both Papers I and II. Such systems have been employed for high temporal resolution (sub-hourly) landslide change detection in France (Kromer et al., 2017), to monitor the Helheim Glacier terminus in Greenland (Gadomski et al., 2017), and to quantify snow redistribution on the Hintereisferner glacier in the Austrian Alps (Voordendag et al., 2021). The latter two examples are particularly compelling models on which to base future work in Longyearbyen, as they employ the same Riegl VZ-6000 TLS model used in this research installed in a permanent housing and set to

scan at regular intervals or via remote access. Automated TLS systems such as these examples present an opportunity to refine the results and conclusions from Paper II regarding snow cornices by considerably increasing the temporal resolution of data acquisition. This in turn would allow for better temporal constraints on cornice accretion rate calculations and cornice-fall avalanche events (i.e. pre- and post-event scans are closer temporally to the actual event) Furthermore, automating the TLS data processing and visualization workflows would permit data product availability in an operationally-relevant timeframe at regular intervals throughout the winter season.

Regional snow and avalanche observations

Addressing the meteorological controls on avalanche activity at the synoptic scale (RO2) required avalanche observations at a broader spatial scale than available via TLS methods. The workflow using regObs data to characterize regional avalanche activity in Paper III presented a first-order attempt at monitoring regional avalanches (e.g. RO3). These data rely on manual observer input, and lack the objectivity and spatial coverage of more quantitative regional avalanche monitoring methods such as satellite observations (Bühler et al., 2019; Eckerstorfer et al., 2019). However, regObs data presented an alternative approach to characterizing regional avalanche conditions made necessary because Svalbard currently lacks satellite coverage with adequate temporal resolution and sensors – such as the the Sentinel-1 synthetic aperture radar (Eckerstorfer et al., 2019) suitable for avalanche monitoring in polar environments. Given adequate satellite coverage, regional avalanche monitoring via satellite data would present a clear possibility for further refinement of the work in Paper III.

Since the research included in Paper III specifically focused on the direct, synoptic-scale meteorological controls on regional avalanche activity, the analyses lack viable Class II data (Figure 15). This is partially because spatial variability in snowpack (Class II) information can be difficult to resolve at broader spatial scales (Clark et al., 2011; Schweizer et al., 2008). Recently published work in Canada used forecasted avalanche problems as a proxy for different types of instabilities in the snowpack (Haegeli et al., 2021). Such an approach may also make sense for future work with regObs data, given the data's primary use is to develop an avalanche forecast via avalanche problems. Chained atmosphere-snowpack-snow stability modeling efforts at lower latitudes have similarly shown promise in addressing broad-scale controls and climate change's influence on avalanching (e.g. Castebrunet et al., 2014). Although this style of research was outside the scope of this work, combining similar modeling efforts with an objective atmospheric circulation classification approach (e.g. Schauer et al., 2021) is a logical next step to refining synoptic-scale avalanche process understandings in Svalbard and applying the conceptual framework from Paper III to other locations without a subjective atmospheric classification scheme.

Hazard management implications

The implementation of both daily regional public avalanche bulletins and local, slope-scale avalanche forecasts for endangered infrastructure in Longyearbyen over the last five years has coincided with this doctoral research (Engeset et al., 2020). Work included in this thesis defining and clarifying the meteorological controls on avalanche activity in central Spitsbergen reinforces the understanding of the Class III data on which avalanche forecasts are initially based. Regional avalanche forecasts in Nordenskiöld Land relying heavily on Class III data from synoptic-scale weather forecasts can benefit from the early identification of atmospheric circulation patterns associated with avalanche activity in Paper III. Early recognition of these synoptic weather patterns in long-range forecasts can help both regional and local forecasters better anticipate periods of increased hazard and target collection of more relevant snowpack (Class II) and instability (Class I) information accordingly.

Managing the distinct cornice-fall avalanche hazard in central Spitsbergen provides an example of a potential forecasting workflow benefiting from the methods and findings presented here. Paper II's results show larger cornice-fall avalanches can be anticipated when synoptic weather forecasts indicate a winter storm will produce the blowing snow conditions which lead to cornice accretion and wind slab development in the avalanche paths under the cornices. The regional wind direction, largely controlled by the synoptic atmospheric conditions, will in turn control the aspect on which cornice failures are more likely. Finer-scale Class II and Class I data collection can then be targeted to the aspect of concern, which is particularly important for local slope-scale forecasts requiring increased forecast precision (McClung, 2002b). At the slope scale, TLS methods developed in this work can then be employed to, for example, determine specific cornice structures more prone to failure and locate avalanche paths with potentially hazardous snow accumulations available for entrainment in the event of a cornice collapse. Targeted snow profile site selection informed by TLS snow cover data can help reduce the uncertainty associated with slope-scale weak layer spatial variability (e.g. Kronholm and Schweizer, 2003; Schweizer et al., 2003b). This in turn helps better forecast the effect of cornice failure on the snowpack— e.g. to determine the likelihood of cornice collapse triggering a larger slab avalanche on the slope below. To fully integrate TLS data products into an operational setting in this way, however, future work must strive to provide useable data products in an operationally relevant timeframe (e.g. Deems et al., 2015).

Uncertainty involved in the forecasting process means it is presently impossible to predict the precise location and timing of avalanche release (Schweizer, 2004). Permanent avalanche mitigation measures such as avalanche hazard zoning and structural defense installation must therefore be implemented in situations where forecasting-related uncertainty is deemed unacceptable (Rudolf-Miklau et al., 2014). In response to the December 2015 and February 2017 destructive avalanche events, authorities elected to redraw avalanche hazard zones in

Longyearbyen (Gundersen et al., 2018). Dynamic avalanche modeling efforts employed to determine the redefined hazard zones employed the TLS data acquired for Paper I, demonstrating a realized hazard management application of this research. Ultimately, authorities chose permanent structural defenses as a strategy to manage avalanche hazard on Sukkertoppen and forego dealing with the uncertainty inherent in the forecasting process (Figure 16). With continued permanent structural defense construction planned, TLS methods can potentially be further integrated into these strategies using promising workflows from other locations (Prokop and Procter, 2016)

Figure 16. Avalanche barriers installed on the slope upon which the December 2015 avalanche on Sukkertoppen released.

Conclusions

Primary contributions addressing the research objectives of this work include:

- Using TLS methods to document and quantify destructive avalanches in Longyearbyen to better understand the meteorological controls on, and thus forecast for, slope-scale avalanche activity in Arctic locations (RO1);
- Developing TLS methods to monitor and quantify cornice and cornice-fall avalanche processes (RO3), and;
- Establishing a foundation for understanding the meteorological controls on avalanche activity in Nordenskiöld Land at the synoptic scale (RO2).

These contributions help refine the conceptual model for avalanche activity in central Spitsbergen by:

- More specifically resolving the relationship between snow accumulation and distribution and destructive avalanche activity at the slope-scale (RO1);
- Demonstrating and quantifying the link between cornice accretion and mid-winter cornice-fall avalanches (RO1), and;
- More directly connecting winter storms at the synoptic scale to regional avalanche activity in Nordenskiöld Land (RO2).

An improved knowledge of central Spitsbergen's high-Arctic maritime snow climate and avalanche regime has implications for both local and global snow avalanche process understanding and hazard management routines. Locally, this work helps clarify how winter storms influence avalanche activity in Nordenskiöld Land and presents methods appropriate for observing snow and avalanche processes in this unique environmental setting. The methods developed in this work, particularly the TLS methodology for cornice monitoring, are applicable worldwide and provide an avenue for better understanding cornice dynamics in other snow climates as well. Hazard managers can furthermore use the findings from this work to better forecast avalanche conditions or plan other mitigation strategies in Svalbard or other locations with similar (e.g. cornice-fall) avalanche hazards.

Globally, perhaps this work's most important contribution is providing a first-order attempt at contextualizing snow avalanches within Svalbard's rapidly changing environmental setting. The synoptic controls on avalanche activity in Svalbard identified in this work provide important insight into the broader relationships between atmospheric circulation and snow avalanches in a location experiencing some of the world's most dramatic environmental changes. Future work investigating how central Spitsbergen's avalanche regime responds to climatic and cryospheric change will thus be useful for anticipating future avalanche regime changes at lower latitudes where environmental change has thus far been less significant. Strategies to better manage and forecast for avalanche hazard in Svalbard's changing climate can help inform similar efforts in other locations in preparation for anticipated climate and avalanche regime changes.

Future perspectives

As of this writing in the spring of 2021, Nordenskiöld Land is experiencing a regional, delayed-action avalanche cycle with large avalanches releasing on deep persistent weak layers. Hazard managers have at the same time elected to indefinitely evacuate the portion of Longyearbyen most threatened by cornice-fall avalanches due to the uncertainty associated with accurately forecasting cornice failure. These circumstances highlight two avenues to build upon the findings from this work with continued research to further improve our understanding of snow avalanche processes here. Automating TLS data acquisition and processing methods would increase the temporal resolution and operational availability of the TLS-derived data products which would help better resolve, and potentially forecast for, the processes leading up to cornice failure. At broader spatial scales, incorporating snowpack structure – either observed or modeled – into the synoptic-scale analyses would help detail atmospheric circulation's control on weak layer development. This would allow more robust analyses including a seasonal progression of atmospheric circulation patterns and avalanche conditions, rather than the simple day-of analyses as presented here in Paper III. A coherent seasonal progression of atmospheric circulation (Class III data), snowpack development (Class II data), and avalanche occurrences (Class I data) would clarify the relationships between snowpack development, loading during winter storms and direct-action avalanches, and less common delayed-action avalanche release in central Spitsbergen. In Svalbard and other avalanche-prone regions, better resolving the relationships between atmospheric circulation, snowpack, and avalanches along seasonal time scales will furthermore improve predictions regarding avalanche activity along the longer temporal scales required for climate change studies.

References

- Adams, M.S., Bühler, Y. and Fromm, R., 2018. Multitemporal Accuracy and Precision Assessment of Unmanned Aerial System Photogrammetry for Slope-Scale Snow Depth Maps in Alpine Terrain. *Pure and Applied Geophysics*, 175(9): 3303-3324.
- André, M.F., 1990. Geomorphic impact of spring avalanches in Northwest Spitsbergen (79 N). *Permafrost and Periglacial Processes*, 1(2): 97-110.
- Arctic Safety Centre, 2020. Accident Report, Avalanche Accident, Fridtjovbreen - Marcussenfjellet, Spitsbergen, Svalbard, 20 February 2020, The Arctic Safety Centre.
- Armstrong, R.L. and Brun, E., 2008. Snow and climate, Physical Processes, Surface Energy Exchange and Modeling. Cambridge University Press, Cambridge, pp. 222.
- Baggi, S. and Schweizer, J., 2009. Characteristics of wet-snow avalanche activity: 20 years of observations from a high alpine valley (Dischma, Switzerland). *Natural Hazards*, 50(1): 97-108.
- Bebi, P., Kulakowski, D. and Rixen, C., 2009. Snow avalanche disturbances in forest ecosystems—State of research and implications for management. *Forest Ecology and Management*, 257(9): 1883-1892.
- Bellaire, S., Jamieson, B., Thumlert, S., Goodrich, J. and Statham, G., 2016. Analysis of long-term weather, snow and avalanche data at Glacier National Park, B.C., Canada. *Cold Regions Science and Technology*, 121: 118-125.
- Birkeland, K.W., Mock, C. and Shinker, J., 2001. Avalanche extremes and atmospheric circulation patterns. *Annals of Glaciology*, 32(1): 135-140.
- Bründl, M. and Margreth, S., 2014. Integrative risk management: the example of snow avalanches. In: W. Haeberli and C. Whiteman (Editors), *Snow and Ice-Related Hazards, Risks, and Disasters*. Elsevier, pp. 263-301.
- Bühler, Y., Hafner, E.D., Zweifel, B., Zesiger, M. and Heisig, H., 2019. Where are the avalanches? Rapid SPOT6 satellite data acquisition to map an extreme avalanche period over the Swiss Alps. *The Cryosphere*, 13(12): 3225-3238.
- Castebrunet, H., Eckert, N., Giraud, G., Durand, Y. and Morin, S., 2014. Projected changes of snow conditions and avalanche activity in a warming climate: the French Alps over the 2020–2050 and 2070–2100 periods. *The Cryosphere*, 8(5): 1673-1697.
- Chabot, D. and Kaba, A., 2016. Avalanche Forecasting in the Central Asian countries of Afghanistan, Pakistan and Tajikistan, International Snow Science Workshop 2016, Breckenridge, CO, USA.
- Christiansen, H.H., Humlum, O. and Eckerstorfer, M., 2013. Central Svalbard 2000-2011 meteorological dynamics and periglacial landscape response. *Arctic, Antarctic, and Alpine Research*, 45(1): 6-18.
- Clark, M.P., Hendrikx, J., Slater, A.G., Kavetski, D., Anderson, B., Cullen, N.J., Kerr, T., Örn Hreinsson, E. and Woods, R.A., 2011. Representing spatial variability of snow water equivalent in hydrologic and land-surface models: A review. *Water Resources Research*, 47(7).
- CloudCompare, 2019. CloudCompare 2.10.2, Retrieved from <http://www.cloudcompare.org>, last access 17 August 2019.
- Cohen, J., Screen, J.A., Furtado, J.C., Barlow, M., Whittleston, D., Coumou, D., Francis, J., Dethloff, K., Entekhabi, D., Overland, J. and Jones, J., 2014. Recent Arctic amplification and extreme mid-latitude weather. *Nature Geoscience*, 7(9): 627-637.
- Dallmann, W.K., 2019. Geoscience Atlas of Svalbard. Norsk Polarinstitutt, Oslo, 292 pp.
- Davis, R.E., Elder, K., Howlett, D. and Bouzaglou, E., 1999. Relating storm and weather factors to dry slab avalanche activity at Alta, Utah, and Mammoth Mountain,

- California, using classification and regression trees. *Cold Regions Science and Technology*, 30(1–3): 79-89.
- Decaulne, A. and Saemundsson, T., 2006. Geomorphic evidence for present-day snow-avalanche and debris-flow impact in the Icelandic Westfjords. *Geomorphology*, 80(1): 80-93.
- Deems, J.S., Gadowski, P.J., Vellone, D., Evanczyk, R., LeWinter, A.L., Birkeland, K.W. and Finnegan, D.C., 2015. Mapping starting zone snow depth with a ground-based lidar to assist avalanche control and forecasting. *Cold Regions Science and Technology*, 120: 197-204.
- Deems, J.S., Painter, T.H. and Finnegan, D.C., 2013. Lidar measurement of snow depth: a review. *Journal of Glaciology*, 59(215): 467-479.
- Eckerstorfer, M., 2013. Snow avalanches in central Svalbard: a field study of meteorological and topographical triggering factors and geomorphological significance. Doctoral thesis Thesis, University of Oslo, 56 pp.
- Eckerstorfer, M. and Christiansen, H.H., 2011a. The "High Arctic Maritime Snow Climate" in Central Svalbard. *Arctic, Antarctic, and Alpine Research*, 43(1): 11-21.
- Eckerstorfer, M. and Christiansen, H.H., 2011b. Relating meteorological variables to the natural slab avalanche regime in High Arctic Svalbard. *Cold Regions Science and Technology*, 69(2–3): 184-193.
- Eckerstorfer, M. and Christiansen, H.H., 2011c. Topographical and meteorological control on snow avalanching in the Longyearbyen area, central Svalbard 2006–2009. *Geomorphology*, 134(3): 186-196.
- Eckerstorfer, M. and Christiansen, H.H., 2012. Meteorology, Topography and Snowpack Conditions causing Two Extreme Mid-Winter Slush and Wet Slab Avalanche Periods in High Arctic Maritime Svalbard. *Permafrost and Periglacial Processes*, 23(1): 15-25.
- Eckerstorfer, M., Vickers, H., Malnes, E. and Grahm, J., 2019. Near-Real Time Automatic Snow Avalanche Activity Monitoring System Using Sentinel-1 SAR Data in Norway. *Remote Sensing*, 11(23): 2863.
- Engeset, R.V., Ekker, R., Humstad, T. and Landrø, M., 2018. Varsom:regobs - A common real-time picture of the hazard situation shared by mobile information technology, International Snow Science Workshop, Innsbruck, Austria, pp. 1573-1577.
- Engeset, R.V., Landrø, M., Indreiten, M., Müller, K., Mikkelsen, O.-A. and Hoseth, K.I.A., 2020. Avalanche warning in Svalbard, Norwegian Water Resources and Energy Directorate, Oslo, Norway.
- Fey, C., Schattan, P., Helfricht, K. and Schöber, J., 2019. A compilation of multitemporal TLS snow depth distribution maps at the Weisssee snow research site (Kaunertal, Austria). *Water Resources Research*, 55(6): 5154-5164.
- Fierz, C., Armstrong, R.L., Durand, Y., Etchevers, P., Greene, E., McClung, D.M., Nishimura, K., Satyawali, P.K. and Sokratov, S.A., 2009. The international classification for seasonal snow on the ground, 25. UNESCO/IHP Paris.
- Fitzharris, B. and Bakkehøi, S., 1986. A synoptic climatology of major avalanche winters in Norway. *Journal of climatology*, 6(4): 431-446.
- Fuchs, S. and McAlpin, M.C., 2005. The net benefit of public expenditures on avalanche defence structures in the municipality of Davos, Switzerland. *Nat. Hazards Earth Syst. Sci.*, 5(3): 319-330.
- Gadowski, P.J., Finnegan, D.C., LeWinter, A., Stearns, L.A., Kershner, C.M. and Hanlon, G., 2017. Three summers of high-resolution, high-accuracy velocity data of Helheim Glacier, as measured by an automated terrestrial LiDAR scanner: methods, challenges, and applications, pp. C41B-1215.

- García, C., Martí, G., Oller, P., Moner, I., Gavalda, J., Martínez, P. and Peña, J.C., 2009. Major avalanches occurrence at regional scale and related atmospheric circulation patterns in the Eastern Pyrenees. *Cold Regions Science and Technology*, 59(2–3): 106-118.
- Germain, D., 2016. Snow avalanche hazard assessment and risk management in northern Quebec, eastern Canada. *Natural Hazards*, 80(2): 1303-1321.
- Germain, D., Fillion, L. and Héту, B., 2009. Snow avalanche regime and climatic conditions in the Chic-Choc Range, eastern Canada. *Climatic Change*, 92(1): 141-167.
- Goodison, B., Louie, P. and Yang, D., 1998. The WMO solid precipitation measurement intercomparison. World Meteorological Organization-Publications: WMO/TD-No. 872: 65-70.
- Gundersen, J., Mikkelsen, O.-A., Johnsen, I. and Nikolaisen, R.U., 2018. Skredrapport Sukkertoppen, Dimensjonerende skred fra Sukkertoppen og faresoner, Norwegian Water Resources and Energy Directorate, Oslo, Norway.
- Haegeli, P., Shandro, B. and Mair, P., 2021. Using avalanche problems to examine the effect of large-scale atmosphere–ocean oscillations on avalanche hazard in western Canada. *The Cryosphere*, 15(3): 1567-1586.
- Hägel, P. and McClung, D.M., 2004. Hierarchy theory as a conceptual framework for scale issues in avalanche forecast modeling. *Annals of Glaciology*, 38: 209-214.
- Hansen, C. and Underwood, S.J., 2012. Synoptic Scale Weather Patterns and Size-5 Avalanches on Mt. Shasta, California. *Northwest Science*, 86(4): 329-341.
- Hanssen-Bauer, I., Førland, E., Hisdal, H., Mayer, S., Sandø, A. and Sorteberg, A., 2019. Climate in Svalbard 2100 - a knowledge base for climate adaptation. 1/2019, Norwegian Centre for Climate Services.
- Hanssen-Bauer, I., Solås, M.K. and Steffensen, E., 1990. The Climate of Spitsbergen, 39. Det Norske Meteorologiske Institutt, 40 pp.
- Hendrikx, J., Owens, I., Carran, W. and Carran, A., 2005. Avalanche activity in an extreme maritime climate: The application of classification trees for forecasting. *Cold Regions Science and Technology*, 43(1–2): 104-116.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R.J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S. and Thépaut, J.-N., 2020. The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, n/a(n/a).
- Hestnes, E., 2000. Impact of rapid mass movement and drifting snow on the infrastructure and development of Longyearbyen, Svalbard, Proceedings of the International Workshop on Permafrost Engineering, pp. 259-282.
- Hestnes, E., Bakkehøi, S. and Jaedicke, C., 2016. Longyearbyen, Svalbard - Vulnerability and risk management of an Arctic settlement under changing climate - a challenge to authorities and experts, International Snow Science Workshop 2016, Breckenridge, CO, USA, pp. 363-370.
- Hestnes, E. and Kristensen, K., 2010. The diversities of large slushflows illustrated by selected cases, International Snow Science Workshop, Lake Tahoe, CA, USA, pp. 17-22.
- Höller, P., 2009. Avalanche cycles in Austria: an analysis of the major events in the last 50 years. *Natural Hazards*, 48(3): 399-424.
- Humlum, O., 2002. Modelling late 20th-century precipitation in Nordenskiöld Land, Svalbard, by geomorphic means. *Norsk geografisk tidsskrift*, 56(2): 96-103.

- Humlum, O., Instanes, A. and Sollid, J.L., 2003. Permafrost in Svalbard: a review of research history, climatic background and engineering challenges. *Polar Research*, 22(2): 191-215.
- Isaksen, K., Førland, E., Dobler, A., Benestad, R., Haugen, J.E. and Mezghani, A., 2017. Klimascenarier for Longyearbyen-området, Svalbard. No. 15/2017, Norwegian Meteorological Institute.
- Isaksen, K., Nordli, Ø., Førland, E.J., Łupikasza, E., Eastwood, S. and Niedźwiedź, T., 2016. Recent warming on Spitsbergen—Influence of atmospheric circulation and sea ice cover. *Journal of Geophysical Research: Atmospheres*, 121(20): 11,913-11,931.
- Jaedicke, C., 2001. Drifting snow and snow accumulation in complex Arctic terrain: field experiments and numerical modelling. Doctoral Thesis, Geophysical Institute, University of Bergen.
- Jaedicke, C., 2002. Snow drift losses from an Arctic catchment on Spitsbergen: an additional process in the water balance. *Cold Regions Science and Technology*, 34(1): 1-10.
- Jaedicke, C., Hestnes, E., Bakkehøi, S., Mørk, T. and Brattlien, K., 2016. Forecasting the fatal Longyearbyen avalanche. Possibilities and challenges., *International Snow Science Workshop*, Breckenridge, CO, USA, pp. 379-382.
- Jamieson, B., 2006. Formation of refrozen snowpack layers and their role in slab avalanche release. *Reviews of Geophysics*, 44(2).
- Keylock, C.J., 2003. The North Atlantic Oscillation and snow avalanching in Iceland. *Geophysical Research Letters*, 30(5): n/a-n/a.
- Kromer, R.A., Abellán, A., Hutchinson, D.J., Lato, M., Chanut, M.A., Dubois, L. and Jaboyedoff, M., 2017. Automated terrestrial laser scanning with near-real-time change detection – monitoring of the Séchilienne landslide. *Earth Surf. Dynam.*, 5(2): 293-310.
- Kronholm, K. and Schweizer, J., 2003. Snow stability variation on small slopes. *Cold Regions Science and Technology*, 37(3): 453-465.
- LaChapelle, E.R., 1980. The fundamental processes in conventional avalanche forecasting. *Journal of Glaciology*, 26: 75-84.
- Lague, D., Brodu, N. and Leroux, J., 2013. Accurate 3D comparison of complex topography with terrestrial laser scanner: Application to the Rangitikei canyon (N-Z). *ISPRS Journal of Photogrammetry and Remote Sensing*, 82: 10-26.
- Larsson, S., 1982. Geomorphological effects on the slopes of Longyear valley, Spitsbergen, after a heavy rainstorm in July 1972. *Geografiska Annaler. Series A. Physical Geography*: 105-125.
- Lehning, M., Löwe, H., Ryser, M. and Raderschall, N., 2008. Inhomogeneous precipitation distribution and snow transport in steep terrain. *Water Resources Research*, 44(7): W07404.
- López-Moreno, J.I., Boike, J., Sanchez-Lorenzo, A. and Pomeroy, J.W., 2016. Impact of climate warming on snow processes in Ny-Ålesund, a polar maritime site at Svalbard. *Global and Planetary Change*, 146: 10-21.
- Luckman, B.H., 1977. The Geomorphic Activity of Snow Avalanches. *Geografiska Annaler. Series A, Physical Geography*, 59(1/2): 31-48.
- McClung, D., 2000. Predictions in avalanche forecasting. *Annals of Glaciology*, 31(1): 377-381.
- McClung, D., 2002b. The Elements of Applied Avalanche Forecasting, Part II: The Physical Issues and the Rules of Applied Avalanche Forecasting. *Natural Hazards*, 26(2): 131-146.
- McClung, D. and Schaerer, P.A., 2006. *The Avalanche Handbook*. The Mountaineers Books, Seattle, 288 pp.

- Mock, C.J. and Birkeland, K.W., 2000. Snow avalanche climatology of the western United States mountain ranges. *Bulletin of the American Meteorological Society*, 81(10): 2367.
- Montagne, J., McPartland, J., Super, A. and Townes, H., 1968. The Nature and control of snow cornices on the Bridger Range, Southwestern Montana, Alta Avalanche Study Center, Miscellaneous Report.
- Muckenhuber, S., Nilsen, F., Korosov, A. and Sandven, S., 2016. Sea ice cover in Isfjorden and Hornsund, Svalbard (2000–2014) from remote sensing data. *The Cryosphere*, 10(1): 149-158.
- Munroe, J.S., 2018. Monitoring snowbank processes and cornice fall avalanches with time-lapse photography. *Cold Regions Science and Technology*, 154: 32-41.
- Niedźwiedź, T., 2013. The atmospheric circulation. In: A.A. Marsz and A. Styszyńska (Editors), *Climate and Climate Change at Hornsund, Svalbard*. Gdynia Maritime University, Gdynia, Poland, pp. 57-74.
- Nordli, Ø., Wyszyński, P., Gjeltén, H.M., Isaksen, K., Łupikasza, E., Niedźwiedź, T. and Przybylak, R., 2020. Revisiting the extended Svalbard Airport monthly temperature series, and the compiled corresponding daily series 1898–2018. *Polar Research*, 39.
- Nuth, C., Kohler, J., König, M., von Deschanden, A., Hagen, J.O., Kääb, A., Moholdt, G. and Pettersson, R., 2013. Decadal changes from a multi-temporal glacier inventory of Svalbard. *The Cryosphere*, 7(5): 1603-1621.
- Onarheim, I.H., Smedsrud, L.H., Ingvaldsen, R.B. and Nilsen, F., 2014. Loss of sea ice during winter north of Svalbard. *Tellus A: Dynamic Meteorology and Oceanography*, 66(1): 23933.
- Parajka, J., Haas, P., Kirnbauer, R., Jansa, J. and Blöschl, G., 2012. Potential of time-lapse photography of snow for hydrological purposes at the small catchment scale. *Hydrological Processes*, 26(22): 3327-3337.
- Peeters, B., Ønvik Pedersen, Å., Loe, L.E., Isaksen, K., Veiberg, V., Stien, A., Kohler, J., Gallet, J.-C., Aanes, R. and Hansen, B., 2019. Spatiotemporal patterns of rain-on-snow and basal ice in high Arctic Svalbard: detection of a climate-cryosphere regime shift.
- Peitzsch, E., Hendriks, J., Stahle, D., Pederson, G., Birkeland, K. and Fagre, D., 2021. A regional spatiotemporal analysis of large magnitude snow avalanches using tree rings. *Nat. Hazards Earth Syst. Sci.*, 21(2): 533-557.
- Pomeroy, J.W., 1989. A process-based model of snow drifting. *Ann. Glaciol*, 13: 237-240.
- Prokop, A., 2008. Assessing the applicability of terrestrial laser scanning for spatial snow depth measurements. *Cold Regions Science and Technology*, 54(3): 155-163.
- Prokop, A., Hancock, H., Praz, M. and Jahn, E., 2018. Slope scale avalanche forecasting in the arctic (Svalbard), *International Snow Science Workshop*, Innsbruck, Austria, pp. 1035-1039.
- Prokop, A. and Procter, E.S., 2016. A new methodology for planning snow drift fences in alpine terrain. *Cold Regions Science and Technology*, 132: 33-43.
- Prokop, A., Schirmer, M., Rub, M., Lehning, M. and Stocker, M., 2008. A comparison of measurement methods: terrestrial laser scanning, tachymetry and snow probing for the determination of the spatial snow-depth distribution on slopes. *Annals of glaciology*, 49(1): 210-216.
- Prokop, A., Schön, P., Singer, F., Pulfer, G., Naaim, M., Thibert, E. and Soruco, A., 2015. Merging terrestrial laser scanning technology with photogrammetric and total station data for the determination of avalanche modeling parameters. *Cold Regions Science and Technology*, 110: 223-230.

- Przybylak, R. and Maszewski, R., 2012. Atmospheric circulation and dynamic conditions. Topoclimatic Diversity in Forlandsundet Region (NW Spitsbergen) in *Global Warming Conditions*: 27-51.
- Reuter, B. and Schweizer, J., 2012. The effect of surface warming on slab stiffness and the fracture behavior of snow. *Cold Regions Science and Technology*, 83-84: 30-36.
- Rudolf-Miklau, F., Skolaut, C. and Sauermoser, S., 2014. Avalanche hazard assessment and planning of protection measures, *The Technical Avalanche Protection Handbook*, pp. 91-126.
- Sauermoser, S., 2006. Avalanche hazard mapping—30 years experience in Austria, *Proceedings of the 2006 International Snow Science Workshop*, Telluride, Colorado, ISSW USA, Colorado, USA, pp. 314-321.
- Sauter, T., Möller, M., Finkelnburg, R., Grabiec, M., Scherer, D. and Schneider, C., 2013. Snowdrift modelling for the Vestfonna ice cap, north-eastern Svalbard. *The Cryosphere*, 7(4): 1287-1301.
- Schauer, A.R., Hendrikx, J., Birkeland, K.W. and Mock, C.J., 2021. Synoptic atmospheric circulation patterns associated with deep persistent slab avalanches in the western United States. *Nat. Hazards Earth Syst. Sci.*, 21(2): 757-774.
- Scherler, D., Bookhagen, B. and Strecker, M.R., 2011. Hillslope-glacier coupling: The interplay of topography and glacial dynamics in High Asia. *Journal of Geophysical Research: Earth Surface*, 116(F2).
- Schneiderbauer, S. and Prokop, A., 2011. The atmospheric snow-transport model: SnowDrift3D. *Journal of Glaciology*, 57(203): 526-542.
- Schön, P., Prokop, A., Vionnet, V., Guyomarc'h, G., Naaim-Bouvet, F. and Heiser, M., 2015. Improving a terrain-based parameter for the assessment of snow depths with TLS data in the Col du Lac Blanc area. *Cold Regions Science and Technology*, 114: 15-26.
- Schuler, T.V. and Østby, T.I., 2020. Sval_Imp: a gridded forcing dataset for climate change impact research on Svalbard. *Earth Syst. Sci. Data*, 12(2): 875-885.
- Schweizer, J., 2004. Snow Avalanches. *Water Resources IMPACT*, 6(1): 12-18.
- Schweizer, J., 2008. Snow avalanche formation and dynamics. *Cold Regions Science and Technology*, 54(3): 153-154.
- Schweizer, J., Bruce Jamieson, J. and Schneebeil, M., 2003a. Snow avalanche formation. *Reviews of Geophysics*, 41(4).
- Schweizer, J., Kronholm, K., Jamieson, J.B. and Birkeland, K.W., 2008. Review of spatial variability of snowpack properties and its importance for avalanche formation. *Cold Regions Science and Technology*, 51(2): 253-272.
- Schweizer, J., Kronholm, K. and Wiesinger, T., 2003b. Verification of regional snowpack stability and avalanche danger. *Cold Regions Science and Technology*, 37(3): 277-288.
- Serreze, M.C., Barrett, A.P., Slater, A.G., Steele, M., Zhang, J. and Trenberth, K.E., 2007. The large-scale energy budget of the Arctic. *Journal of Geophysical Research: Atmospheres*, 112(D11).
- Serreze, M.C. and Barry, R.G., 2014. *The Arctic Climate System*. Cambridge Atmospheric and Space Science Series. Cambridge University Press, Cambridge.
- Serreze, M.C., Crawford, A.D. and Barrett, A.P., 2015. Extreme daily precipitation events at Spitsbergen, an Arctic Island. *International Journal of Climatology*, 35(15): 4574-4588.
- Statham, G., Haegeli, P., Greene, E., Birkeland, K., Israelson, C., Tremper, B., Stethem, C., McMahon, B., White, B. and Kelly, J., 2018. A conceptual model of avalanche hazard. *Natural Hazards*, 90(2): 663-691.
- Stethem, C., Jamieson, B., Schaerer, P., Liverman, D., Germain, D. and Walker, S., 2003. Snow avalanche hazard in Canada—a review. *Natural Hazards*, 28(2): 487-515.

- Sturm, M., Holmgren, J. and Liston, G.E., 1995. A seasonal snow cover classification system for local to global applications. *Journal of Climate*, 8(5): 1261-1283.
- Svalbardposten, 2019. To omkommet i snøskred i Hornsund, Svalbardposten, Longyearbyen, NO.
- van Herwijnen, A. and Fierz, C., 2014. Monitoring snow cornice development using time-lapse photography, *Proceedings of the International Snow Science Workshop*, 2014, Banff, Canada, pp. 865-869.
- Vikhamar-Schuler, D., Førland, E.J., Lutz, J. and Gjelten, H.M., 2019. Evaluation of downscaled reanalysis and observations for Svalbard -- Background report for Climate in Svalbard 2100, (www.klimaservicesenter.no).
- Vikhamar-Schuler, D., Isaksen, K., Haugen, J.E., Tømmervik, H., Luks, B., Schuler, T.V. and Bjerke, J.W., 2016. Changes in winter warming events in the Nordic Arctic region. *Journal of Climate*, 29(17): 6223-6244.
- Vogel, S., Eckerstorfer, M. and Christiansen, H., 2012. Cornice dynamics and meteorological control at Gruvefjellet, Central Svalbard. *The Cryosphere*, 6(1): 157-171.
- Voordendag, A.B., Goger, B., Klug, C., Prinz, R., Rutzinger, M. and Kaser, G., 2021. AUTOMATED AND PERMANENT LONG-RANGE TERRESTRIAL LASER SCANNING IN A HIGH MOUNTAIN ENVIRONMENT: SETUP AND FIRST RESULTS. *ISPRS Ann. Photogramm. Remote Sens. Spatial Inf. Sci.*, V-2-2021: 153-160.
- Walczowski, W. and Piechura, J., 2011. Influence of the West Spitsbergen Current on the local climate. *International Journal of Climatology*, 31(7): 1088-1093.
- Wickström, S., Jonassen, M.O., Cassano, J.J. and Vihma, T., 2020. Present Temperature, Precipitation and Rain-on-Snow Climate in Svalbard. *Journal of Geophysical Research: Atmospheres*, n/a(n/a): e2019JD032155.
- Wickström, S., Jonassen, M.O., Vihma, T. and Uotila, P., 2019. Trends in cyclones in the high-latitude North Atlantic during 1979–2016. *Quarterly Journal of the Royal Meteorological Society*, n/a(n/a).

Papers

Paper I

Combining high spatial resolution snow mapping and meteorological analyses to improve forecasting of destructive avalanches in Longyearbyen, Svalbard

Combining high spatial resolution snow mapping and meteorological analyses to improve forecasting of destructive avalanches in Longyearbyen, Svalbard

Holt Hancock^{a,b,*}, Alexander Prokop^a, Markus Eckerstorfer^{c,a}, Jordy Hendrikx^d

^a Department of Arctic Geology, University Centre in Svalbard, N-9171 Longyearbyen, Norway

^b Department of Geosciences, University of Oslo, N-0371 Oslo, Norway

^c Earth Observation Division, Northern Research Institute (Norut), N-9294 Tromsø, Norway

^d Snow and Avalanche Lab, Department of Earth Science, Montana State University, 59717-3480 Bozeman, Montana, USA

ARTICLE INFO

Keywords:

Geohazards

High-Arctic maritime snow climate

Terrestrial laser scanning

ABSTRACT

Two naturally triggered snow avalanches occurred on 19 December 2015 and 21 February 2017 in the town of Longyearbyen, Svalbard in the Norwegian high-Arctic. These events resulted in two fatalities, numerous injuries, and rendered fourteen residential buildings uninhabitable. Both avalanches occurred on the west-facing slope of the Sukkertoppen Mountain and were preconditioned by similar meteorological conditions. We investigate these two events by combining traditional weather and snowpack analyses with snow distribution data acquired via terrestrial laser scanning (TLS). As limited snow data exists on Svalbard, the TLS-derived snow depth and differential snow depth maps are the primary viable method for the description and analysis of destructive avalanche activity in this location. These TLS-derived surfaces permit detailed assessment of slope-scale snow distribution patterns both prior to and following avalanche activity. We identify strong easterly winds and moderate to heavy snowfall as precursors to destructive avalanche activity on this slope. The results of our investigation help clarify the relationship between winter storm characteristics and avalanche activity in high-Arctic environments and demonstrate the importance of scale-appropriate snow data for avalanche forecasting with increased precision at finer spatial scales. These results have implications for avalanche forecasting in this setting and other data sparse, high-relief Arctic settings where snow distribution patterns are controlled by wind.

1. Introduction

Fatal avalanches occurring in settlements have become increasingly rare in Europe and North America due to refined operational forecasting services, updated avalanche danger zonation and regulations, and increased permanent mitigation measures (e.g. Bründl and Margreth, 2014; Laternser and Schneebeli, 2002; Sauermoser, 2006; Stethem et al., 2003). In these regions, avalanche forecasts serve as a critical public safety measure and hazard management strategy in settlements exposed to snow avalanches.

Avalanche forecasting involves analyzing data sources from multiple spatial and temporal scales to identify instabilities within the snowpack, evaluate current avalanche conditions, and predict future avalanche activity (LaChapelle, 1980). Conventional avalanche forecasting techniques rely on recognizing instabilities within the snowpack based on direct snow stability observations (e.g. observations of avalanches and stability test results), snowpack structure (e.g. snowpack

stratigraphy and snow temperatures), and meteorological data (LaChapelle, 1980; McClung, 2002). The availability of scale-appropriate data determines, to a large degree, the ability to produce a robust avalanche forecast for a given area (McClung, 2000b; McClung, 2002).

Modern remote sensing techniques complement traditional field methods for avalanche forecasting and allow the efficient collection of information concerning snow and avalanches from remote locations or at broader spatial scales (Bühler et al., 2016; Deems et al., 2013; Eckerstorfer et al., 2016; Wesselink et al., 2017). Terrestrial laser scanning (TLS) is an emerging remote sensing tool for monitoring snow distribution and avalanche occurrence at the slope-scale, allowing users to gather data from safe locations well removed from avalanche hazards (Deems et al., 2015; Prokop, 2008; Prokop, 2009; Prokop et al., 2008). TLS is particularly advantageous over other snow observation methods in Arctic environments where the polar night can preclude direct visual observation of snow and avalanche conditions. Recent studies have demonstrated the potential of TLS in operational avalanche forecasting

* Corresponding author at: Department of Arctic Geology, University Centre in Svalbard, N-9171 Longyearbyen, Norway.

E-mail address: Holt.Hancock@unis.no (H. Hancock).

<https://doi.org/10.1016/j.coldregions.2018.05.011>

Received 14 December 2017; Received in revised form 2 May 2018; Accepted 6 May 2018

Available online 10 July 2018

0165-232X/ © 2018 Elsevier B.V. All rights reserved.

settings (Deems et al., 2015), to validate the performance of snow drift modeling (Schneiderbauer and Prokop, 2011; Schön et al., 2015; Vionnet et al., 2014), to retrieve parameters for dynamic avalanche modeling (Prokop et al., 2015), and as a tool in planning structural avalanche defenses (Prokop and Procter, 2016).

The challenges associated with accurate, high precision avalanche forecasting in Arctic settings where relevant data sources are sparse or non-existent are highlighted by two recent slab avalanche events on the Sukkertoppen Mountain (hereafter Sukkertoppen) in Longyearbyen, Svalbard, Norway. The first avalanche occurred on 19 December 2015, trapping 25 people in 11 residential buildings. This avalanche tragically resulted in two fatalities. On 21 February 2017, a second avalanche released on the same slope and destroyed two more residential buildings.

While operational avalanche forecasting services were not regularly implemented for Longyearbyen prior to the 19 December 2015 avalanche, lack of scale-specific data contributed to inaccurate hazard assessments (and failure to evacuate endangered buildings) during the 21 February 2017 event (Landrø et al., 2017). This lack of data substantially reduces the precision of any hazard assessment which, given the lack of adherence to mapped avalanche paths and known hazard zones in Longyearbyen, culminated in infrastructure destruction during these two events.

We investigate these two destructive avalanche events as case studies to better understand – and thus in the future forecast for – avalanche activity in remote, data sparse, and wind-dominated Arctic regions. The specific objectives are to:

- (1) combine a unique spatial snow distribution dataset obtained via TLS with meteorological and snowpack data to characterize conditions leading to these events;
- (2) demonstrate how meteorological conditions and associated slope-scale snow distribution patterns influence avalanche activity on this slope, and;
- (3) illustrate how data availability and associated uncertainty affect the ability to forecast for slope-scale destructive avalanches in Longyearbyen.

The implications from this investigation will be of significance for other data sparse high-Arctic settlements and high-relief areas where wind is an important agent for the redistribution of snow.

1.1. Study area

Svalbard is a high-Arctic archipelago located 700 km north of the Norwegian mainland on the northwestern margin of the Barents Sea. Longyearbyen, Svalbard's administrative center, is located in the Nordenskiöld Land region of central Spitsbergen, Svalbard's largest island (Fig. 1). Longyearbyen is nestled between steep mountain slopes in the Longyear valley (Longyeardalen). Longyeardalen is a north-south oriented, glacially-formed U-shaped valley extending from the termini of two mountain glaciers to the Advent fjord.

The geomorphological setting around Longyearbyen strongly influences local avalanche activity (Eckerstorfer and Christiansen, 2011c). The plateau summits serve as large source areas for blowing snow. Snow entrained across these areas is deposited as cornices on the leeward plateau margins and forms wind slabs on the slopes below (Vogel et al., 2012). Cornice fall and slab avalanche activity dominate the avalanche regime in the study area, with the majority of the avalanche activity confined to avalanche paths between the protruding bedrock buttresses (Eckerstorfer and Christiansen, 2011c). On Sukkertoppen's west-northwestern aspect, however, the open, planar slope morphology is conducive to connected slab development and failure, with minor slab avalanches observed almost every year.

1.2. Climate and meteorology

Svalbard's setting near the confluence of various air masses and ocean currents with differing thermal characteristics is recognized as one of the most climatically sensitive regions in the world (Hanssen-Bauer et al., 1990; Rogers et al., 2005). Synoptic airflow is primarily controlled by interactions between the Icelandic Low and anticyclones over the Arctic Ocean, Greenland, and Siberia resulting in a pattern of alternating high and low pressure systems over Svalbard (Christiansen et al., 2013). Historically, this pattern has been most pronounced during the winter season when weather in Svalbard is characterized by long periods of cold, stable high pressure interspersed with warm, wet, low pressure systems transported meridionally from the south-south-west along the North Atlantic cyclone track (Dickson et al., 2000; Rogers et al., 2005). The warm open ocean along Spitsbergen's western coast exerts considerable maritime influence on the region's climate during all months, with inter-annual fluctuations in Arctic and local sea ice extent substantially affecting local climatic conditions (Cottier et al., 2007; Førland et al., 2011).

The mean winter season (DJF) air temperature measured at the Svalbard Airport during the period from 1981 to 2010 was -11.7°C , with annual measured winter season precipitation during this same period of 55 mm (Førland et al., 2011). Wind measurements on Gruefjellet (Fig. 1) from the past 15 years show prevailing southeasterly winds on the mountain plateaus, while winds are redirected along the major valleys axes (Christiansen et al., 2013). Since 1990, the mean annual precipitation has increased by as much as 20–30% (Vikhamar-Schuler et al., 2016) and air temperatures, particularly during the winter, have increased during the same period (Førland et al., 2011; Nordli et al., 2014).

Central Svalbard's snow and avalanche climate has been classified as “high-Arctic maritime” as a result of this unique climatological setting (Eckerstorfer and Christiansen, 2011a). The high-Arctic maritime snow climate is typified by a thin, cold snowpack that persists for 8–10 months of the year in which structural weaknesses – typically depth hoar near the ground – develop early in the winter and persist throughout the season. Warm winter storms are increasingly frequent and associated with rain-on snow-events and ice layering within the snowpack. Strong, persistent winds form dense wind slabs in leeward areas. Slab avalanches typically occur as direct action avalanches during or immediately after winter storms (Eckerstorfer and Christiansen, 2011b; Eckerstorfer and Christiansen, 2011c).

1.3. Avalanche forecasting

Longyearbyen has until recently lacked – or failed to employ – suitable hazard zoning, monitoring programs, and data acquisition infrastructure for conventional avalanche forecasting at the slope scale. Prior to the 19 December 2015 avalanche event, snow avalanche hazard assessments in the region were primarily performed by the Norwegian Geotechnical Institute (NGI) on a contractual basis. Despite numerous reports authored by NGI staff evaluating avalanche hazard and providing suggestions for structural defenses and avalanche forecasting routines, limited progress was made to developing long-term avalanche protection solutions (e.g. Hestnes, 2000; Hestnes et al., 2016; Jaedicke et al., 2016). Local authorities again contracted NGI to establish a daily forecasting service for Longyearbyen following the 19 December 2015 avalanche, which was continued until the end of January 2016 when the Norwegian national avalanche forecasting service expanded to include Nordenskiöld Land (Jaedicke et al., 2016). Currently, public advisories are issued daily by the Norwegian Water Resources and Energy Directorate (NVE) for the Nordenskiöld Land region. NVE is also contracted by local authorities to provide hazard assessments for the slopes immediately above Longyearbyen to be used in making evacuation decisions.

Fig. 1. Overview of the Longyearbyen area indicating meteorological stations and key locations mentioned in text. Contour lines in panel (a) are spaced at 100 m. The location and direction of the photograph in panel (b) is indicated (a). The location and extent of the 19 December 2015 and 21 February 2017 Sukkertoppen avalanches are indicated in both panels.

2. Data sources and methods

2.1. Meteorology

We obtained data from the Gruvefjellet automated weather station (AWS), located on the Gruvefjellet plateau at 464 m a.s.l. approximately 2 km south of Sukkertoppen to characterize meteorological conditions in the region's higher elevation terrain. This AWS records hourly meteorological data. We used data from the Adventdalen AWS for high temporal resolution meteorological measurements representative of lower elevation locations during storm events. The Adventdalen AWS is situated at 15 m a.s.l. approximately 4 km east of Sukkertoppen. Finally, we accessed precipitation data from the Svalbard Airport AWS, located 4 km west of Longyearbyen at the Svalbard Airport at 28 m a.s.l. The Svalbard Airport AWS is the only weather station in the region with long-term precipitation measurements. However, its location is poorly suited to characterize spatially variable precipitation patterns in the higher elevation, mountainous areas of particular concern for snow avalanche processes due to the region's considerable vertical precipitation gradients (e.g. Humlum, 2002).

2.2. Snow and avalanche data

2.2.1. Terrestrial laser scanning (TLS)

We scanned with a TLS regularly throughout the 2016/2017 winter season, with particular emphasis placed on obtaining scans of pre- and post-event snow surfaces for major storm and avalanche events. For our repeated slope-scale snow measurements we used the Riegl® Laser

Measurement Systems VZ-6000 laser scanner that is particularly suitable for measuring snow surfaces due to an operating laser wavelength of 1064 nm and a high scanning speed (www.riegl.com; Prokop, 2008). We georeferenced and positioned repeated scans using a differential global positioning system (DGPS) and an iterative closest point algorithm in RiSCAN PRO, Riegl's proprietary data processing software (Prokop and Panholzer, 2009). We filtered non-ground or snow surface points using the wedge filtering approach (Panholzer and Prokop, 2013). We resampled the original georeferenced point clouds to a 0.25 m grid for analysis and display purposes, as this resolution has been shown to retain snow surface roughness features while reducing the influence of artifacts relative to higher resolution grids (Deems et al., 2015). By computing the difference in vertical distance of two digital elevation models, we derived final elevation differences between two snow and terrain surfaces resulting in spatially distributed maps of snow depth or changes in snow depth.

We use data from five scans to quantify and analyze changes in snow distribution for this study: (S1) a bare-earth scan prior to the onset of seasonal snow cover in September 2016, (S2) a post-event scan of the 19 December 2015 avalanche taken 29 January 2016, (S3) a pre-event scan from 14 February 2017, with the 21 February 2017 post-event snow depths obtained by merging scans taken on (S4) 22 February 2017 and (S5) 24 February 2017.

Laser scanner availability precluded pre- and timely post-event scanning for the 19 December 2015 event, but we scanned the avalanche area on 29 January 2016 (S2) as soon as the laser scanner was available. Differential snow depths could not be calculated for the December 2015 avalanche due to the lack of a pre-event scanned snow

surface, but we calculated 29 January 2016 snow depths by subtracting the September 2016 bare-earth data (S1) from the 29 January surface (S2).

For the February 2017 event, we performed S4 as soon as visibility allowed for accurate scanning to take place following the avalanche. We took an additional scan (S5) to gather data from the avalanche deposition area that was obscured by buildings in S4. We merged these two scans together in RiSCAN Pro to create a single post-event snow surface. This snow surface was used in the snow depth and differential snow depth calculations presented in the results below.

2.2.2. Snowpack development and snow stratigraphy

We obtained snow stratigraphy data from NVE's observation platform regObs, (www.regobs.no). In order to analyze the snowpack development prior to both avalanches, we created idealized snow profiles for Sukkertoppen's western aspect using available regObs snow observations and weather data from different automated weather stations in the region. Snow profiles from the starting zone on Sukkertoppen were only available on regObs as crown profiles obtained after both avalanches released.

3. Results

3.1. 19 December 2015

3.1.1. Meteorology and snowpack development

Snowpack data near Longyearbyen were sparse in the early 2015/2016 winter season, with only six snow profiles in the region available prior to 19 December 2015. Warm (above 0°C), wet weather in the first half of November (Fig. 2) formed a pair of melt-freeze crusts sandwiching a layer of weak faceted snow, with well-developed basal facets under this crust-facet-crust sequence (Fig. 3a). A series of weak winter storms in early December buried the deeper structural instabilities

under a denser layer of wind packed snow (Fig. 3b).

At ca. 1100 UTC on 17 December 2015, winds near Longyearbyen began to increase from an easterly direction as a large low pressure system west of Svalbard intensified (Fig. 4a). Heavy snowfall began on 18 December, accompanied by rising air temperatures and continued strong easterly winds. Easterly winds persisted at Gruvefjellet and Adventdalen for the 48 hours prior to the avalanche, with maximum wind speeds approaching 30 m s⁻¹ at both weather stations (Fig. 5, Table 1). By the time the storm abated on the morning of 19 December, 18.1 mm (W.E.) of precipitation – which fell as snow – had been officially recorded at the Svalbard Airport, the majority of which fell during the period with the strongest easterly winds (avg. 19.3 m s⁻¹ at Gruvefjellet) in the 24 hours prior to the avalanche release (Table 1).

The strong easterly winds and heavy snowfall resulted in a period of intense wind loading on the lower, northerly portion of Sukkertoppen's west aspect. Over two meters of snow accumulated on the weak existing snowpack, with knife (K) hardness snow at the bottom of the new wind slab transitioning to four finger (4F) hardness at the snow surface (Fig. 3c). The avalanche released at the interface between the lower melt-freeze crust and the overlying facet layer approximately 25 cm above the ground (Fig. 3c; regObs #71681).

3.1.2. Terrestrial Laser Scanning

As no pre-event or timely post-event TLS data are available for the 19 December 2015 avalanche, much of the data in Table 2 comes from field observations immediately following the event. TLS data products derived from a scanning campaign on 29 January 2016 are affected by the considerable time elapsed post-avalanche, during which a major rain-on-snow event and several small snow storms deteriorated and refilled the original avalanche surface.

The 29 January 2016 snow depth map (Fig. 6a) reflects the topographical and meteorological controls on Sukkertoppen's seasonal snow cover. The deepest snow accumulations occur on aspects immediately

Fig. 2. Meteorological summary of the early snow season preceding and immediately following the 19 December 2015 avalanche event. Maximum wind speed and air temperature are hourly data. Snow depth (HS) and precipitation data are measured at the Svalbard Airport AWS, with snow depth measured once daily at 0600 UTC and precipitation twice daily at 0600 and 1800 UTC. The light grey shaded vertical bars in each graph panel highlight the 48 hours preceding the avalanche event.

Fig. 3. Snow profile observation summaries showing the evolution of the snowpack through November and December 2015 on Sukkertoppen's western aspect. Panels (a) and (b) represent idealized snow profiles created from snow observations and weather data for the specified time periods. Panel (c) is an idealized crown profile of the avalanche based on snow pits and crown profile investigations following avalanche release, with the faceted weak layer on which the slab failed outlined in red. Clarification for the hand hardness index (American Avalanche Association, 2016) and snow grain form symbology (Fierz et al., 2009) included in the figure is provided in the legend.

Fig. 4. 850 mb geopotential heights illustrating the synoptic situation and low pressure positioning for (a) 19 December 2015 at 0000 UTC and (b) 21 February 2017 at 0000 UTC. Longyearbyen's location is indicated by the red dot in each panel, with northern mainland Norway located in the lower right. Data courtesy of the NCEP/NCAR Reanalysis Project (Kalnay et al., 1996)

leeward of the prevailing east-southeasterly wind direction along Sukkertoppen's northerly ridge and in the debris flow channels incised in the mountain's western face. Wind-scoured areas of bare or nearly bare ground exist on Sukkertoppen's north aspect, where easterly valley winds from Adventdalen regularly strip any snow available for transport, depositing much of this snow in the deeper accumulation areas.

The avalanche release area boundaries digitized referencing the 29 January snow surface hillshade (Fig. 6b) and event photographs (Fig. 6c) are less obvious in the snow depth map (Fig. 6a). Despite considerable refill and rain-induced deterioration of the original crown line, the release area's right flank is still clearly distinguishable and the crown's location can be inferred by abrupt snow depth decreases near the release area's perimeter. The stauchwall visible in post-event photographs has deteriorated in the scanned surface, and much of the debris was removed by local authorities in the time between the avalanche event and the TLS scan.

3.2. 21 February 2017

3.2.1. Meteorology and snowpack development

By early January 2017, a combination of low density facets and

higher density wind slabs had been deposited on depth hoar and melt forms resulting from a slow snow cover onset and multiple rain-on-snow events earlier in the season (Fig. 7a).

Positive air temperatures at most elevations and precipitation falling as rain near Longyearbyen from 4 February to 10 February followed by cold air intrusion on 11 and 12 February formed a pencil (P) to knife (K) hardness melt-freeze crust at most elevations, capping deeper basal instabilities and the January stratigraphy (Figs. 8, 7b). As temperatures continued to fall on 14 and 15 February, disturbances within the northerly airflow deposited modest snow totals at all elevations. Continued northerly airflow ushered in a period of calm, clear, cold weather on 16 February that persisted through 18 February, providing ideal conditions to promote faceting in the new snow at the surface.

On 19 February, a low pressure system over the Norwegian Sea began to impact central Spitsbergen, with moderate snowfall accompanying generally light northwesterly winds (Figs. 8 and 9). The snow deposited on 14 and 15 February was readily transported together with the new storm snow, with notable soft slab avalanche activity observed on the eastern aspects of Longyeardalen (regObs #115494 and #115556).

Fig. 5. Wind summary of the 48 hours prior to the 19 December 2015 avalanche event. The red vertical line in panels (a) and (b) indicates the timing of the avalanche release. The shaded grey background in panels (a) and (b) designates the 48 hours prior to the avalanche event and corresponds to the same time period as the vertical grey bars in Fig. 2. Panels (c) and (d) display wind rose diagrams for the 48 hours prior to avalanche release at Gruvefjellet and Adventdalen, respectively

Light snowfall continued through the night of 19 February and re-intensified under calmer conditions on 20 February, with the Svalbard Airport recording 7.5 mm (W.E.) of precipitation from 0600 UTC on 19 February to 0600 UTC on 21 February (Table 1). By the evening of 20 February, more than 20 cm of new snow was observed in Long-yearbyen, with an observer reporting “abnormal amounts of dry loose snow” available for transport throughout the region (regObs #115671). With elongated low pressure situated south of Spitsbergen at 0000 UTC on 21 February (Fig. 4b), surface winds increased once again from a prevailing easterly direction (Fig. 9). East-northeasterly winds blew strongly throughout the night, with an average wind direction of 86.4° and averaged wind speeds of 10.2 m s⁻¹ at Gruvefjellet in the 24 hour period prior to the avalanche (Table 1). Strong winds and light snowfall continued into the morning of 21 February and throughout the afternoon following the avalanche release. The regional avalanche hazard rating was increased to 4 (high) at 0735 UTC 21 February due to the strong winds and the amount of snow available for transport.

The idealized snowpack stratigraphy displayed in Fig. 7c from the crown profile shows the facets formed during the clear, cold weather from 16 February – 18 February overlain by a wind slab formed on 20

and 21 February. The faceted weak layer failed naturally after widespread wind loading, with the underlying melt forms formed earlier in February acting as the bed surface.

3.2.2. Terrestrial laser scanning

Slope-scale snow distribution patterns on 14 February 2017 (Fig. 10a) show similar general features to those on 29 January 2016, with major accumulation areas situated on the leeward aspect of the northern ridge and in the location of the 19 December 2015 avalanche release area.

The post-event snow depth and differential snow depth maps (Fig. 10b and c) show the changes to the snow cover resulting from the 19 – 21 February storm and the 21 February avalanche event. Areas of accumulation over this time period are primarily associated with avalanche deposition (annotations 1, 2, 3, and 4 on Fig. 10c), while snow depth decreases result from avalanche release (annotations 5, 6, and 7 on Fig. 10c) and wind scouring under the strong east-northeasterly winds during the storm (annotation 8 on Fig. 10c).

A rather complex avalanche morphology is displayed in the differential snow depth data (Fig. 10c), with multiple avalanche release areas

Table 1
Summary of the meteorological conditions preceding the 19 December 2015 and 21 February 2017 avalanche events. Averages represent the 24 or 48 hour averaging periods, while maximum wind speeds represent 48 hour maximums.

		19 December 2015		21 February 2017	
		24 hrs prev.	48 hrs prev.	24 hrs prev.	48 hrs prev.
Event time (UTC)		0923		1035	
Airport accumulated precipitation (mm)		18.1	18.1	4.6	7.5
Average Gruvefjellet air temp. (°C)		−5.9	−10.0	−10.0	−12.5
Average Adventdalen air temp. (°C)		−3.0	−6.8	−7.4	−9.6
Gruvefjellet average wind speed (m s ^{−1})	Vector	19.3	15.2	8.8	4.1
	Scalar	19.3	15.3	10.2	6.6
Gruvefjellet maximum wind speed (m s ^{−1})		28.8		22.7	
Gruvefjellet average wind direction (°)		95.3	96	86.4	77.5
Adventdalen average wind speed (m s ^{−1})	Vector	18.0	15.1	9.1	4.4
	Scalar	18.0	15.2	9.2	6.2
Adventdalen maximum wind speed (m s ^{−1})		28.3		26.9	
Adventdalen average wind direction (°)		90.1	90.0	82.8	6.2

Table 2
Event characteristics from the two avalanches. Displayed avalanche characteristics and abbreviations presented parenthetically are after the avalanche observation guidelines outlined in American Avalanche Association (2016). Characteristics from the February 21st avalanche event are only for the primary avalanche that released from near Sukkertoppen's summit.

		19 December 2015	21 February 2017
Path characteristics			
Aspect		Northwest	Northwest
Starting zone slope angle (°)		35	35–38
Crown face elevation (m a.s.l.)		120	255 – 355
Event characteristics			
Avalanche type		Hard slab (HS)	Soft slab (SS)
Trigger		Natural (N)	Natural (N)
Size		(R4 D2.5)	(R3 D2.5)
Snow properties			
Bed surface		Old snow (O)	Old/new snow interface (I)
Weak layer snow grain type		Facets (FC)	Facets (FC)
Slab snow grain type		Rounded grains (RG)	Rounded grains (RG)
Dimensions			
Maximum crown height (m)		2.5	1.25
Width (m)		200	120 – 150
Vertical (m)		90	315
Alpha-angle (°)		16	25
Beta-angle (°)		19	28

and three primary deposition zones. The greatest snow depth decreases occur along the northwestern ridge, where the large drift in the 14 February scan released in the largest 21 February avalanche (annotation 5 on Fig. 10c). A portion of this drift that did not release (annotation 9 on Fig. 10c) shows localized accumulations exceeding one meter between 14 February and 21 February, representing the sizable snow load deposited during the storm and giving an indication of the avalanche's maximum crown height (Table 2). The release area's snow depth up to 0.5 m below the 14 February surface indicates this avalanche also ran on the 10 February crust, entraining snow down to this surface as it traveled downslope. While a secondary deposition area exists at a mid-path bench feature where the slope decreases from 35° to below 25° for approximately 50 m (annotation 3 on Fig. 10c), the primary deposition area for this avalanche had snow depths in excess of 2

m (annotation 1 on Fig. 10c).

At the same time, to the north of this main event, a separate avalanche released in the same location as the 19 December 2015 event, also running on the 10 February rain crust. While snow accumulations in this avalanche's deposition zone generally exceed 2 m (annotation 2 on Fig. 10c), insufficient snow existed in the release area (annotation 6 on Fig. 10c) compared to the December 2015 event to reach the location of infrastructure destroyed by that prior avalanche. Upslope of this avalanche, a third and smaller avalanche ran approximately 70 m downslope (annotations 4 and 7 on Fig. 10c).

4. Discussion

4.1. Snow climate considerations

Investigations focusing on snow avalanches as natural hazards in central Svalbard over the last 30 years (e.g. Eckerstorfer, 2013; Hestnes, 2000; Hestnes et al., 2016) have established a conceptual framework for understanding avalanche activity in this location. The novel suite of meteorological and snowpack conditions typifying the high-Arctic maritime snow climate result in a persistently weak snowpack that is nevertheless conditionally stable through much of winter due to the “bridging effects” (Thumlert and Jamieson, 2014) of crusts and hard wind slabs in addition to a lack of snow available for slab formation (Eckerstorfer and Christiansen, 2011a). Natural slab avalanche release is primarily direct action and results from winter storm conditions providing the snow available for transport and strong winds necessary for storm slab formation and overburden of existing structural weaknesses (Eckerstorfer and Christiansen, 2011b).

Both the 19 December 2015 and 21 February 2017 avalanche events in Longyearbyen generally support this conceptual framework. Snowpack conditions prior to each event were characteristically “high-Arctic maritime” and can be considered relatively normal – for the last decade – for both events considering their temporal situation within the snow season. The snowpack prior to the 17 – 19 December 2015 storm typified early season conditions in central Svalbard, with generally shallow snow, basal faceting, and persistent instabilities in the upper snowpack sandwiched between melt-freeze crusts. Similarly, the meteorological situation in early 2017 resulted in a fairly typical – based on modern climatic conditions – mid-season Svalbard snowpack prior to the 21 February avalanche, with warm weather and rain followed by cold, stable conditions effectively capping and bridging pre-existing instabilities with a thick melt-freeze crust (e.g. Tremper, 2008).

Despite antecedent snowpack conditions within the range currently expected in this snow climate, atypical winter storm characteristics resulted in the historically unprecedented nature of these avalanche events. Although smaller avalanches regularly run from Sukkertoppen's lower release area following light to moderate snowfall and southeasterly winds (Hestnes et al., 2016), avalanches from Sukkertoppen's upper release area had not been documented prior to the 21 February 2017 avalanche (Landrø et al., 2017). Furthermore, no previous avalanches from Sukkertoppen had resulted in major infrastructure damage. While other work has shown major winter precipitation events in central Spitsbergen usually coincide with southwesterly winds (Jaedicke et al., 2016), both destructive avalanche events investigated here occurred following winter storms during which plentiful snow available for transport and snowfall coincided with prevailing easterly winds. This suggests that the relatively rare snow and weather situation in central Spitsbergen prior to 19 December 2015 and 21 February 2017 set the stage for destructive avalanche activity on Sukkertoppen's westerly aspect.

Snow climate characteristics and mountain range specific terrain attributes have also been shown to affect avalanche runout distances (McClung, 2000a; Mears, 1989). Previous work in Svalbard has documented avalanche runouts with alpha-angles as low as 15.5° (Hestnes et al., 1999) and indicates up to ten percent of naturally released

Fig. 6. Overview of the slope-scale snow distribution patterns for the 19 December 2015 avalanche event. Panel (a) shows the snow depth in meters on Sukkertoppen's western aspect derived from a TLS scan on 29 January 2016. Panel (b) shows the interpreted avalanche release area (in red) and runout (in green) from the avalanche displayed on a hillshade map of the 29 January 2016 snow surface. The black rectangle in panel (a) indicates the extent of panel (b). Panel (c) is a photo of the avalanche release area taken on 22 December 2015 (photo courtesy Erik Myrum Næss).

Fig. 7. Snow profile observation summaries showing the evolution of the snowpack through January and February 2017 on Sukkertoppen's western aspect. Panels (a) and (b) represent idealized snow profiles created from snow observations and weather data for the specified time periods. Panel (c) is an idealized crown profile of the avalanche based on snow pits and crown profile investigations following avalanche release, with the faceted weak layer on which the slab failed outlined in red. Clarification for the hand hardness index (American Avalanche Association, 2016) and snow grain form symbology (Fierz et al., 2009) included in the figure is provided in the legend.

avalanches in Svalbard may have alpha-angles of less than 20° (Hestnes et al., 2016). Although the 16° alpha-angle from the December 2015 avalanche is very low (e.g. Issler et al., 2016) compared to the expected value based on the alpha-beta model calibrated on the Norwegian mainland (Lied and Bakkehoi, 1980), similarly long runout distances are not unprecedented in Svalbard. The 25° alpha-angle from the February 2017 event is very similar to the 25.5° mean alpha angle derived from 76 avalanches previously recorded in the region (Hestnes et al., 1999) and cannot be considered extreme in this snow climate. In this context, while the December 2015 event is likely a lower frequency scenario than the February 2017 avalanche based on runout length, avalanche forecasting and hazard zonation in Svalbard should take unexpectedly long avalanche runouts into consideration.

4.2. Event comparison

Despite the general similarities in the local meteorological situations between the two events, notable differences in observed wind and precipitation affected the slope-scale snow distribution patterns and resulting avalanche activity. Winds during both storms blew from a prevailing easterly direction during peak storm intensity, but subtle differences in air temperature, precipitation rate, and the direction and duration of the strongest winds likely exerted considerable control on the slope-scale snow distribution patterns influencing avalanche release. Strong winds blew from a remarkably consistent easterly direction at both the Adventdalen and Gruvefjellet stations in the 48 hours prior to the 19 December 2015 avalanche. During the same period, heavy snowfall supplied a substantial amount of snow available for transport across the region. The consistently strong easterly winds placed Sukkertoppen's lower release area directly leeward of

Fig. 8. Meteorological summary of the early snow season preceding and immediately following the 21 February 2017 avalanche event. Wind speed and air temperature are hourly data. Snow depth (HS) and precipitation data are measured at the Svalbard Airport AWS, with snow depth measured once daily at 0600 UTC and precipitation twice daily at 0600 and 1800 UTC. The light grey shaded vertical bars in each graph panel highlight the 48 hours preceding the avalanche event.

Adventdalen's expansive snow-covered fetch, providing a virtually limitless supply of snow to a relatively specific accumulation area. This localized deposition of a massive storm snow load with minimal time for the snowpack to adjust ultimately resulted in the catastrophic release of a single slab.

By contrast, winds prior to the 21 February 2017 avalanche were considerably more variable, with light to moderate northwesterly winds 48 hours prior to avalanche release transitioning to strong east-northeasterly winds as the storm intensified. More variable wind directions combined with plentiful loose snow throughout the terrain prior to the onset of winter storm conditions resulted in a more complex loading pattern for the February event, with accumulation areas more widely distributed across Sukkertoppen's westerly aspect. In particular, the slight northeasterly component to the strongest winds in addition to prior light loading under easterly and northwesterly winds appear to have shifted the deepest snow storm snow accumulations to Sukkertoppen's upper release area. This loading occurred on a facet-over-crust stratigraphy associated with the variation between warm-wet and cold-dry conditions characteristic of Svalbard's modern climatic setting. These weak snow conditions were unable to adjust to the wind loading during the night and early morning hours of 21 February, with widespread but discontinuous patterns of simultaneous slab release demonstrating the sensitivity of this storm slab instability. Furthermore, the 10 February crust provided an effective bed surface on which the avalanche from Sukkertoppen's upper release area could entrain remaining loose snow and run a relatively long distance into the buildings below (Landrø et al., 2017).

Previous work has shown precipitation and snowdrift indices to be the best meteorological predictors of natural slab avalanche activity near Longyearbyen (Eckerstorfer and Christiansen, 2011b). In Svalbard's snow climate where threshold wind speeds for snow transport are regularly exceeded (Christiansen et al., 2013; Eckerstorfer and Christiansen, 2011a), available snow for transport is a critical limiting factor in the development of hazardous avalanche conditions. The

relatively unique combination of strong winds from the eastern quadrant and considerable snow available for transport appear to be prerequisites for destructive avalanche activity on this slope (Jaedicke et al., 2016), but differing slope-scale snow distribution patterns resulting from subtle variations in wind direction govern specific slab release locations. Based on these case studies, hazardous slab development has occurred in Sukkertoppen's lower release area following snowfall and sustained strong due easterly winds, while Sukkertoppen's upper release area presented a hazard to infrastructure following intense wind loading under east-northeasterly winds. Since these two potentially discrete release areas may threaten infrastructure under different loading patterns, effectively forecasting for destructive avalanche activity in this location without excessive evacuations requires supplementing an understanding of the general meteorological and snowpack situation with more slope-specific snow accumulation data available in an operationally relevant timeframe.

4.3. Local forecasting

As it is presently impossible to predict the precise location and timing of an avalanche release, avalanche forecasting at all scales entails dealing with uncertainty (Schweizer, 2004). In general, the avalanche forecasting problem becomes more challenging and specific as the spatial extent of a forecasting region decreases, requiring more direct, less uncertain data obtained at a finer spatial scale (McClung and Schaerer, 2006). Forecasting precision is especially important in Longyearbyen where, unlike the majority of settlements in North America and Europe, many structures are located in avalanche paths without the added safety provided by, for example, structural avalanche defenses employed extensively in the Alps (Höller, 2007). Until strategies including structural defenses, rezoning, or relocation of endangered infrastructure are employed in Longyearbyen, resident safety in buildings threatened by avalanches is primarily dependent on accurate and precise avalanche hazard assessments to determine if and

Fig. 9. Wind summary of the 48 hours prior to the 21 February 2017 avalanche event. The red vertical line in panels (a) and (b) indicates the timing of the avalanche release. The shaded grey background in panels (a) and (b) designates the 48 hours prior to the avalanche event and corresponds to the same time period as the vertical grey bars in Fig. 6. Panels (c) and (d) are wind rose diagrams for the 48 hours prior to avalanche release at Gruvefjellet and Adventdalen, respectively.

when to evacuate threatened infrastructure.

The failure to evacuate threatened infrastructure prior to the 21 February 2017 event despite a high regional avalanche hazard rating illustrates the challenges associated with integrating multiple data sources across a range of spatial scales while compensating for limited data availability. At the regional scale, a verified synoptic weather forecast for moderate to heavy snowfall accompanied by strong winds and considerable amounts of loose, new snow throughout the region allowed for an accurate regional avalanche hazard assessment. On Sukkertoppen's western aspect, however, a lack of operationally available slope-specific meteorological and snow data precluded an accurate assessment of localized avalanche conditions. Here, hazard assessments were based on field observations in the lower release area taken the evening before the avalanche released, prior to the increase in east-northeasterly winds during the night before the avalanche (Landrø et al., 2017), and no operationally relevant snow depth data existed for the upper release area. Based on the available data, forecasters believed snow accumulation on Sukkertoppen was insufficient to present a danger to infrastructure below despite the increased regional hazard rating, and no evacuations were ordered by the local authorities.

The lack of scale-appropriate total snow depth and new snowfall data is of particular concern in this scenario, as snow depth and snow depth changes in avalanche release areas have long been recognized as crucial indicators of developing avalanche hazard (Atwater, 1954; Davis et al., 1999; Eckerstorfer and Christiansen, 2011b; Hendrikx et al., 2005; Schweizer et al., 2009). Unfortunately, the only continuous, automated precipitation or snow measurements available for the region come from the Svalbard Airport AWS. While these data can serve as a basis for rough post-event precipitation estimates, measuring solid precipitation in windy Arctic environments is fraught with documented uncertainty (Goodison et al., 1998), and this AWS' location near sea level is unrepresentative of the mountainous areas. As such, these data are not suitable for slope-scale hazard management purposes requiring high spatial and temporal resolution snow depth measurements.

To better manage avalanche hazard in Longyearbyen, attempts must be made to reduce forecast uncertainty through improved scale-appropriate data acquisition infrastructure. Here, we have employed TLS data to map slope-scale snow distribution patterns in this environment, allowing for a better understanding of the processes influencing destructive avalanche activity in Longyearbyen. While TLS is clearly a

Fig. 10. Overview of the slope-scale snow distribution patterns for 21 February 2017 avalanche event. Panels (a) and (b) display snow depth (HS) information derived from TLS data acquired 14 February 2017 (pre-event) and 22 and 24 February 2017 (post-event), respectively. The outline of the avalanche is also displayed in panel (b). Panel (c) shows the change in snow depth (differential snow depth – dHS) for the period 14 – 21 February 2017, with specific snow surface features annotated as they are referred to in the text. The photo in panel (d) taken at 1700 UTC on 21 February 2017 shows the avalanche from a similar angle as shown in panels (a) to (c), with the avalanche perimeter indicated where visible in the photo.

powerful tool for post-event documentation and analysis, scan timing and visibility constraints limit its operational applicability and derived data product availability. Furthermore, the equipment costs and labor involved in data processing are currently still prohibitive for most operational programs. TLS data products are limited for the 19 December 2015 event by the time interval between the avalanche and the scanning campaign, although such a time interval is unlikely to occur in an operational setting. In the 21 February 2017 case study, poor visibility during the storm period prior to and immediately after the avalanche event prohibited scanning until almost 24 hours following the avalanche. The differential snow depth and snow depth maps derived from the post-event scans are valuable for understanding and analyzing snow distribution patterns following the event, but could not have been available for forecasting purposes prior to avalanche release. Nevertheless, TLS viability studies in other environments have shown promising operational possibilities (Deems et al., 2015) and continued work should strive to improve strategies integrating TLS derived data products into operational settings by reducing data processing time through workflow automation. Additionally, equipment and labor costs for TLS equipped forecasting programs should be considered with respect to other hazard reduction strategies (i.e. structural defenses and/or relocation of some buildings) with cost analyses similar to those employed in other regions (e.g. Fuchs et al., 2007).

When lacking more direct evidence, such as snow depth, to assess avalanche hazard, forecasters must rely on other, less direct data types. For example, information about past situations – referred to as distributional data (McClung, 2000b) – can be used to predict avalanche

activity under similar conditions in the present. However, reliance on distributional data requires a robust record of past avalanche activity with which to compare current conditions. This is particularly problematic in Svalbard, where a relatively short and non-continuous avalanche observation record does not provide a sufficient framework from which to base future forecasts. Furthermore, documented climatic changes over the past decade and modeled future climate change scenarios for the 21st century indicate Svalbard's climate is undergoing dramatic change (Isaksen et al., 2016; Nordli et al., 2014; Vikhamar-Schuler et al., 2016). Svalbard's snow and avalanche climate will thus also likely undergo considerable change, with the potential for more novel and previously undocumented avalanche situations arising from, for example, increased winter precipitation and likelihood of mid-winter rain on snow events (Bintanja and Andry, 2017; Vikhamar-Schuler et al., 2016). Such changes will introduce further uncertainty into the forecasting process, as past snow and avalanche situations will have limited relevance for forecasting under previously undocumented climatic scenarios.

Fatal avalanche events in settlements during the 1990s in northern Iceland and northern Quebec, Canada further highlight the difficulties associated with forecasting in high-latitude settings where snow depth measurements are sparse, records of avalanche activity are missing or incomplete, and existing precipitation measurements are often unrepresentative of snow accumulation in avalanche release areas (Bernhardsdóttir, 2001; Egilsson, 1996; Germain, 2016). Similar to the destructive avalanche events investigated in this work, these events have been attributed to anomalous snow accumulation patterns due

intense wind-loading prior to avalanche release, with slab avalanches releasing as a result of wind-induced snow accumulations of previously undocumented magnitude or in unexpected locations (Egilsson, 1996; Germain, 2016). Continued difficulty anticipating snow accumulation and forecasting avalanche activity in these situations demonstrates the necessity of continued research in Arctic and other high-latitude snow and avalanche climates, as most avalanche knowledge and applied forecasting strategies have been developed in lower-latitude settings. With amplified Arctic warming (e.g. Cohen et al., 2014) likely inducing dramatic shifts in Arctic snow and avalanche climates preceding similar changes at lower latitudes, developing improved observational and forecasting strategies in Arctic environments can provide a framework to better forecast avalanche activity in destabilized climatic conditions where climate changes introduce further uncertainty into the forecasting process.

5. Conclusions

Destructive avalanche activity on Sukkertoppen in consecutive winter seasons resulted from a combination of prevailing easterly winds and moderate to heavy snowfall with considerable snow available for transport. Small changes in air temperature, snowfall rate, and direction within the easterly winds had an impact on slope-scale snow accumulation patterns and associated avalanche activity, highlighting the importance of documenting the wind's influence on snow distribution patterns when forecasting for avalanche activity at the slope scale in Arctic environments and other mountainous areas where wind controls snow accumulation.

Unfortunately, severe limitations on scale-appropriate data availability have precluded effective hazard assessment and avalanche forecasting in Longyearbyen. Accurate avalanche forecasting is further hindered by uncertainty related to short periods of record, unreliable precipitation data, a changing climate, and poor observation density. This lack of data and high uncertainty within traditionally available data sources highlight the need for a more robust snow and avalanche data acquisition infrastructure, but expansion of structural defenses, rezoning, and relocating the most endangered infrastructure is still likely necessary even with improved data. Here, we have demonstrated the applicability of TLS techniques for event documentation and analysis, but limitations on TLS data acquisition for operational forecasting purposes still exist. While access to a TLS in this work provided an advantageous avenue – especially for the 21 February 2017 avalanche – to acquire post-event data that is often lacking from destructive avalanches, continued and improved observation and data collection in this and other Arctic snow and avalanche settings is necessary to better forecast for avalanches under the unique climatic conditions characterizing these locations.

Declarations of interest

None.

Acknowledgements

This work was greatly improved by comments on an early version of this manuscript by Graham Lewis Gilbert and numerous discussions with Wesley Randall Farnsworth. We also thank the editor, Dr. Peter Gauer, and two anonymous reviewers for helpful suggestions and comments during the review process.

References

- American Avalanche Association, 2016. Snow, Weather and Avalanches: Observation Guidelines for Avalanche Programs in the United States, 3rd ed. (Victor, ID, 104 pp.).
- Atwater, M.M., 1954. Snow avalanches. *Sci. Am.* 190, 26–31.
- Bernhardsdóttir, Á.E., 2001. Learning From Past Experiences: The 1995 Avalanches in Iceland. CRISMART, Försvarshögskolan, Stockholm, pp. 69.
- Bintanja, R., Andry, O., 2017. Towards a rain-dominated Arctic. *Nat. Clim. Chang.* 7 (4), 263–267.
- Bründl, M., Margreth, S., 2014. Integrative risk management: the example of snow avalanches. In: Haeblerli, W., Whiteman, C. (Eds.), *Snow and Ice-related Hazards, Risks, and Disasters*. Elsevier, pp. 263–301.
- Bühler, Y., Adams, M.S., Bosch, R., Stoffel, A., 2016. Mapping snow depth in alpine terrain with unmanned aerial systems (UAS): potential and limitations. *Cryosphere* 10 (3), 1075–1088.
- Christiansen, H.H., Humlum, O., Eckerstorfer, M., 2013. Central Svalbard 2000–2011 meteorological dynamics and periglacial landscape response. *Arct. Antarct. Alp. Res.* 45 (1), 6–18.
- Cohen, J., Screen, J.A., Furtado, J.C., Barlow, M., Whittleston, D., Coumou, D., Francis, J., Dethloff, K., Entekhabi, D., Overland, J., Jones, J., 2014. Recent Arctic amplification and extreme mid-latitude weather. *Nat. Geosci.* 7 (9), 627–637.
- Cottier, F.R., Nilsen, F., Inall, M.E., Gerland, S., Tverberg, V., Svendsen, H., 2007. Wintertime warming of an Arctic shelf in response to large-scale atmospheric circulation. *Geophys. Res. Lett.* 34 (10), L10607.
- Davis, R.E., Elder, K., Howlett, D., Bouzaglou, E., 1999. Relating storm and weather factors to dry slab avalanche activity at Alta, Utah, and Mammoth Mountain, California, using classification and regression trees. *Cold Reg. Sci. Technol.* 30 (1–3), 79–89.
- Deems, J.S., Painter, T.H., Finnegan, D.C., 2013. Lidar measurement of snow depth: a review. *J. Glaciol.* 59 (215), 467–479.
- Deems, J.S., Gadowski, P.J., Vellone, D., Evanczyk, R., LeWinter, A.L., Birkeland, K.W., Finnegan, D.C., 2015. Mapping starting zone snow depth with a ground-based lidar to assist avalanche control and forecasting. *Cold Reg. Sci. Technol.* 120, 197–204.
- Dickson, R.R., Osborn, T.J., Hurrell, J.W., Meincke, J., Blindheim, J., Adlandsvik, B., Vinje, T., Alekseev, G., Maslowski, W., 2000. The Arctic ocean response to the north Atlantic oscillation. *J. Clim.* 13 (15), 2671–2696.
- Eckerstorfer, M., 2013. Snow Avalanches in Central Svalbard: A Field Study of Meteorological and Topographical Triggering Factors and Geomorphological Significance. Doctoral thesis Thesis. University of Oslo (56 pp.).
- Eckerstorfer, M., Christiansen, H.H., 2011a. The “High Arctic Maritime Snow Climate” in central svalbard. *Arct. Antarct. Alp. Res.* 43 (1), 11–21.
- Eckerstorfer, M., Christiansen, H.H., 2011b. Relating meteorological variables to the natural slab avalanche regime in high arctic Svalbard. *Cold Reg. Sci. Technol.* 69 (2–3), 184–193.
- Eckerstorfer, M., Christiansen, H.H., 2011c. Topographical and meteorological control on snow avalanching in the Longyearbyen area, central Svalbard 2006–2009. *Geomorphology* 134 (3), 186–196.
- Eckerstorfer, M., Bühler, Y., Frauenfelder, R., Malnes, E., 2016. Remote sensing of snow avalanches: recent advances, potential, and limitations. *Cold Reg. Sci. Technol.* 121, 126–140.
- Egilsson, J.G., 1996. Two Destructive Avalanches in Iceland, International Snow Science Workshop (ISSW), Banff, Canada.
- Fierz, C., Armstrong, R.L., Durand, Y., Etchevers, P., Greene, E., McClung, D., Nishimura, K., Satyawali, P.K., Sokratov, S.A., 2009. The international classification for seasonal snow on the ground. In: IHP-VII Technical Documents in Hydrology N°83, IACS Contribution N°1, 25. UNESCO-IHP Paris, Paris.
- Førland, E., Benestad, R., Hanssen-Bauer, I., Haugen, J.E., Skaugen, T.E., 2011. Temperature and precipitation development at Svalbard 1900–2100. *Adv. Meteorol.* 2011.
- Fuchs, S., Thöni, M., McAlpin, M.C., Gruber, U., Bründl, M., 2007. Avalanche hazard mitigation strategies assessed by cost effectiveness analyses and cost benefit analyses—evidence from Davos, Switzerland. *Nat. Hazards* 41 (1), 113–129.
- Germain, D., 2016. Snow avalanche hazard assessment and risk management in northern Quebec, eastern Canada. *Nat. Hazards* 80 (2), 1303–1321.
- Goodison, B., Louie, P., Yang, D., 1998. The WMO Solid Precipitation Measurement Intercomparison. World Meteorological Organization-Publications: WMO/TD-No. 872, pp. 65–70.
- Hanssen-Bauer, I., Solås, M.K., Steffensen, E., 1990. The Climate of Spitsbergen, 39. Det Norske Meteorologiske Institutt (40 pp.).
- Hendrikx, J., Owens, I., Carran, W., Carran, A., 2005. Avalanche activity in an extreme maritime climate: the application of classification trees for forecasting. *Cold Reg. Sci. Technol.* 43 (1–2), 104–116.
- Hestnes, E., 2000. Impact of rapid mass movement and drifting snow on the infrastructure and development of Longyearbyen, Svalbard. In: *Proceedings of the International Workshop on Permafrost Engineering*, pp. 259–282.
- Hestnes, E., Bakkehoi, S., Brattlien, K., Onarheim, K., 1999. Snø og snøskred på Svalbard. Registrering av snøforhold, snøskred og utløpslengder. NGL-Report 589100-4. Norwegian Geotechnical Institute, Oslo.
- Hestnes, E., Bakkehoi, S., Jaedicke, C., 2016. Longyearbyen, Svalbard - Vulnerability and Risk Management of an Arctic Settlement Under Changing Climate - A Challenge to Authorities and Experts, International Snow Science Workshop 2016, Breckenridge, CO, USA, pp. 363–370.
- Höller, P., 2007. Avalanche hazards and mitigation in Austria: a review. *Nat. Hazards* 43 (1), 81–101.
- Humlum, O., 2002. Modelling Late 20th-Century Precipitation in Nordenskiöld Land, Svalbard, by Geomorphic means. *Norsk Geografisk tidsskrift*. vol. 56(2). pp. 96–103.
- Isaksen, K., Nordli, Ø., Førland, E.J., Łupikasza, E., Eastwood, S., Niedźwiedz, T., 2016. Recent warming on Spitsbergen—Influence of atmospheric circulation and sea ice cover. *J. Geophys. Res. Atmos.* 121 (20), 11,913–11,931.
- Issler, D., Jónsson, Á., Gauer, P., Domaas, U., 2016. Vulnerability of Houses and Persons Under Avalanche Impact—The Avalanche at Longyearbyen on 2015-12-19. International Snow Science Workshop 2016, Breckenridge, CO, USA, pp. 371–378.

- Jaedicke, C., Hestnes, E., Bakkehøi, S., Mørk, T., Brattlien, K., 2016. Forecasting the Fatal Longyearbyen Avalanche. Possibilities and Challenges. International Snow Science Workshop, Breckenridge, CO, USA, pp. 379–382.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K.C., Ropelewski, C., Wang, J., Jenne, R., Joseph, D., 1996. The NCEP/NCAR 40-year reanalysis project. *Bull. Am. Meteorol. Soc.* 77 (3), 437–471.
- LaChapelle, E.R., 1980. The fundamental processes in conventional avalanche forecasting. *J. Glaciol.* 26, 75–84.
- Landrø, M., Mikkelsen, O.-A., Jaedicke, C., 2017. Gjennomgang og Evaluering av skredhendelsen i Longyearbyen 21.02.2017. Oslo, Norway.
- Latenser, M., Schneebeli, M., 2002. Temporal trend and spatial distribution of avalanche activity during the last 50 years in Switzerland. *Nat. Hazards* 27 (3), 201–230.
- Lied, K., Bakkehøi, S., 1980. Empirical calculations of snow-avalanche run-out distance based on topographic parameters. *J. Glaciol.* 26 (94), 165–177.
- McClung, D., 2000a. Extreme avalanche runout in space and time. *Can. Geotech. J.* 37 (1), 161–170.
- McClung, D., 2000b. Predictions in avalanche forecasting. *Ann. Glaciol.* 31 (1), 377–381.
- McClung, D., 2002. The elements of applied avalanche forecasting, part ii: the physical issues and the rules of applied avalanche forecasting. *Nat. Hazards* 26 (2), 131–146.
- McClung, D., Schaerer, P.A., 2006. The Avalanche Handbook. The Mountaineers Books, Seattle, pp. 288.
- Mears, A.I., 1989. Regional comparisons of avalanche-profile and runout data. *Arct. Alp. Res.* 21 (3), 283–287.
- Nordli, Ø., Przybylak, R., Ogilvie, A.E.J., Isaksen, K., 2014. Long-term Temperature Trends and Variability on Spitsbergen: The Extended Svalbard Airport Temperature Series, 1898–2012. pp. 2014.
- Panholzer, H., Prokop, A., 2013. Wedge-filtering of geomorphologic terrestrial laser scan data. *Sensors* 13 (2), 2579.
- Prokop, A., 2008. Assessing the applicability of terrestrial laser scanning for spatial snow depth measurements. *Cold Reg. Sci. Technol.* 54 (3), 155–163.
- Prokop, A., 2009. Terrestrial laser scanning for snow depth observations: an update on technical developments and applications. In: Schweizer, J., Van Herwijnen, A. (Eds.), *Proceedings of the International Snow Science Workshop, 27 September - 2 October 2009, Davos, Switzerland*, pp. 192–196.
- Prokop, A., Panholzer, H., 2009. Assessing the capability of terrestrial laser scanning for monitoring slow moving landslides. *Nat. Hazards Earth Syst. Sci.* 9 (6), 1921–1928.
- Prokop, A., Procter, E.S., 2016. A new methodology for planning snow drift fences in alpine terrain. *Cold Reg. Sci. Technol.* 132, 33–43.
- Prokop, A., Schirmer, M., Rub, M., Lehning, M., Stocker, M., 2008. A comparison of measurement methods: terrestrial laser scanning, tachymetry and snow probing for the determination of the spatial snow-depth distribution on slopes. *Ann. Glaciol.* 49 (1), 210–216.
- Prokop, A., Schön, P., Singer, F., Pulfer, G., Naaim, M., Thibert, E., Soruco, A., 2015. Merging terrestrial laser scanning technology with photogrammetric and total station data for the determination of avalanche modeling parameters. *Cold Reg. Sci. Technol.* 110, 223–230.
- Rogers, J.C., Yang, L., Li, L., 2005. The role of Fram Strait winter cyclones on sea ice flux and on Spitsbergen air temperatures. *Geophys. Res. Lett.* (6), 32.
- Sauermoser, S., 2006. Avalanche hazard mapping—30 years experience in Austria. In: *Proceedings of the 2006 International Snow Science Workshop, Telluride, Colorado, ISSW USA, Colorado, USA*, pp. 314–321.
- Schneiderbauer, S., Prokop, A., 2011. The atmospheric snow-transport model: snow-drift3d. *J. Glaciol.* 57 (203), 526–542.
- Schön, P., Prokop, A., Vionnet, V., Guyomarc'h, G., Naaim-Bouvet, F., Heiser, M., 2015. Improving a terrain-based parameter for the assessment of snow depths with TLS data in the Col du Lac Blanc area. *Cold Reg. Sci. Technol.* 114, 15–26.
- Schweizer, J., 2004. Snow avalanches. *Water Resour. IMPACT* 6 (1), 12–18.
- Schweizer, J., Mitterer, C., Stoffel, L., 2009. On forecasting large and infrequent snow avalanches. *Cold Reg. Sci. Technol.* 59 (2–3), 234–241.
- Stethem, C., Jamieson, B., Schaerer, P., Liverman, D., Germain, D., Walker, S., 2003. Snow avalanche hazard in Canada—a review. *Nat. Hazards* 28 (2), 487–515.
- Thumlert, S., Jamieson, B., 2014. Stress measurements in the snow cover below localized dynamic loads. *Cold Reg. Sci. Technol.* 106, 28–35.
- Tremper, B., 2008. *Staying Alive in Avalanche Terrain*. The Mountaineers Books, Seattle, WA, USA, pp. 318.
- Vikharnar-Schuler, D., Isaksen, K., Haugen, J.E., Tømmervik, H., Luks, B., Schuler, T.V., Bjerke, J.W., 2016. Changes in winter warming events in the Nordic Arctic region. *J. Clim.* 29 (17), 6223–6244.
- Vionnet, V., Martin, E., Masson, V., Guyomarc'h, G., Naaim Bouvet, F., Prokop, A., Durand, Y., Lac, C., 2014. Simulation of wind-induced snow transport and sublimation in alpine terrain using a fully coupled snowpack/atmosphere model. *Cryosphere* 8, 395–415.
- Vogel, S., Eckerstorfer, M., Christiansen, H., 2012. Cornice dynamics and meteorological control at Gruvefjellet, Central Svalbard. *Cryosphere* 6 (1), 157–171.
- Wesselink, D.S., Malnes, E., Eckerstorfer, M., Lindenberg, R.C., 2017. Automatic detection of snow avalanche debris in central Svalbard using C-band SAR data. *Polar Res.* 36 (1), 1333236.

Web references

- regObs registration #71681, <http://www.regobs.no/Registration/71681>, observer: Kjetil Brattlien, observation date and time: 20/12/2015 12:00:00
- regObs registration #115494, <http://www.regobs.no/Registration/115494>, observer: Jens Abild, observation date and time: 19/02/2017 14:43
- regObs registration #115556, <http://www.regobs.no/Registration/115556>, observer: Kjetil@obskorps, observation date and time: 19/02/2017 22:44
- regObs registration #115671, <http://www.regobs.no/Registration/115671>, observer: Martini@unis, observation date and time: 20/02/2017 17:30
- Nordenkiöld Land avalanche Forecast, <http://www.varsom.no/snoskredvarsling/varsel/Nordenski%C3%B6ld%20Land/>, provided by The Norwegian Water Resour. Energy Directorate (NVE)
- http://www.riegl.com/uploads/tx_pxpriegldownloads/RIEGL_VZ-6000_Datasheet_2017-09-01.pdf.

Paper II

*Quantifying seasonal cornice dynamics using a
terrestrial laser scanner in Svalbard, Norway*

Quantifying seasonal cornice dynamics using a terrestrial laser scanner in Svalbard, Norway

Holt Hancock^{1,2}, Markus Eckerstorfer³, Alexander Prokop^{4,5,1}, and Jordy Hendrikx⁶

¹Department of Arctic Geology, University Centre in Svalbard, 9171 Longyearbyen, Norway

²Department of Geosciences, University of Oslo, 0371 Oslo, Norway

³Earth Observation Group, NORCE, 9294 Tromsø, Norway

⁴Department of Geodynamics and Sedimentology, University of Vienna, 1090 Vienna, Austria

⁵Snow Scan GmbH, Research, Engineering, Education, Stadlauerstrasse 31, 1220 Vienna, Austria

⁶Snow and Avalanche Lab, Department of Earth Sciences, Montana State University, Bozeman, Montana 59717, USA

Correspondence: Holt Hancock (holt.hancock@unis.no)

Received: 8 October 2019 – Discussion started: 18 November 2019

Revised: 23 January 2020 – Accepted: 3 February 2020 – Published: 26 February 2020

Abstract. Snow cornices develop along mountain ridges, edges of plateaus, and marked inflections in topography throughout regions with seasonal and permanent snow cover. Despite the recognized hazard posed by cornices in mountainous locations, limited modern research on cornice dynamics exists and accurately forecasting cornice failure continues to be problematic. Cornice failures and associated cornice fall avalanches comprise a majority of observed avalanche activity and endanger human life and infrastructure annually near Longyearbyen in central Svalbard, Norway. In this work, we monitored the seasonal development of the cornices along the plateaus near Longyearbyen with a terrestrial laser scanner (TLS) during the 2016–2017 and 2017–2018 winter seasons. The spatial resolution at which we acquired snow surface data with TLS enabled us to observe and quantify changes to the cornice systems in detail not previously achieved. We focused primarily on the evolution and failure of the lower cornice surfaces where accessibility has precluded previous research. We measured cornice accretion rates in excess of 10 mm h^{-1} during several accretion events coinciding with winter storms. We observed five cornice fall avalanche events following periods of cornice accretion and one event following a warm period with mid-winter rain. The results of our investigation provide quantitative reinforcement to existing conceptual models of cornice dynamics and illustrate cornice response to specific meteorological events. Our results demonstrate the utility of TLS for monitoring cornice processes and as a viable method for

quantitative cornice studies in this and other locations where cornices are of scientific or operational interest.

1 Introduction

Snow cornices are overhanging projections of snow that form due to the deposition of wind-transported snow in the lee of ridgelines or sharp slope inflections (Montagne et al., 1968; Seligman, 1936). Cornices have attracted interest for their hydrologic implications (e.g., Anderton et al., 2004) and as agents of geomorphic change in periglacial environments (Eckerstorfer et al., 2013; Humlum et al., 2007), but they are perhaps best recognized as a snow and avalanche hazard in mountainous terrain (Montagne et al., 1968; Vogel et al., 2012). Cornices pose an avalanche hazard when they fail either as a full cornice failure with the entire cornice detaching from the ground or as a partial failure with a smaller cornice mass separating from the rest of the cornice. The detached cornice blocks travel downslope under the influence of gravity and become a cornice fall avalanche by entraining loose surface snow or triggering a snow slab on the slope below (e.g., Vogel et al., 2012). In ski areas or where cornices and cornice fall avalanches endanger infrastructure, both explosives (Farizy, 2013; McCarty et al., 1986) and structural defenses (e.g., Montagne et al., 1968) are employed operationally to mitigate cornice hazards. Most cornice-related fatalities, however, occur in recreational backcountry settings

and result from the victim's weight triggering cornice failure.

Despite the well-recognized hazards and operational challenges associated with cornices and cornice fall avalanches, specific cornice research is relatively scarce. Early cornice studies summarized by Vogel et al. (2012) focused on qualitative descriptions of cornice formation processes and resulting cornice structures (e.g., Montagne et al., 1968; Seligman, 1936). Later studies investigated mechanisms by which individual snow crystals adhere during cornice accretion (Latham and Montagne, 1970), the physical snow characteristics at various structural locations on individual cornices (Naruse et al., 1985), and the specific interactions between wind-drifted snow and cornice morphology during cornice formation (Kobayashi et al., 1988).

Recent work has refined the conceptual model of seasonal cornice dynamics established by these earlier studies primarily by employing time-lapse photography to examine cornice responses to the meteorological factors controlling the development and failure of cornices (Munroe, 2018; van Herwijnen and Fierz, 2014; Vogel et al., 2012). Vogel et al. (2012) observed cornice processes over two winter seasons on a single mountain slope in central Svalbard and proposed a conceptual model of seasonal cornice dynamics including cornice accretion, deformation, and failure. Their results indicated cornice accretion occurs during or immediately following winter storms with wind speeds in excess of 10 m s^{-1} from a direction perpendicular to the ridgeline, while cornice scouring resulted from strong winds oriented towards the cornice's leading edge. Smaller cornice failures observed by Vogel et al. (2012) were clustered in June near the end of the snow season and coincided with increasing air temperatures and decreased snow strength. Less frequent failures in the earlier part of the winter often involved the entire cornice mass and resulted in some of the largest cornice fall avalanches observed in the study.

Later work in an alpine setting also linked cornice accretion to strong winds during or soon after a snowfall and found the SNOWPACK wind drift index correlated well with cornice width estimates (van Herwijnen and Fierz, 2014). No cornice failures or cornice fall avalanches were observed in this study, however. Munroe (2018) used time-lapse photography to observe the growth and repeated failure of a cornice in Utah, USA. He also found cornice accretion to primarily coincide with periods of snowdrift. He divided the 19 cornice fall avalanches observed in his study into two distinct groups: snow-caused cornice fall avalanches where failure primarily resulted from snow loading on the cornice and temperature-caused avalanches where failure was related to rapid temperature increases presumably leading to destabilization of the cornice through the loss of snow strength.

We build upon the observational understanding and conceptual model of seasonal cornice dynamics established in these previous works by monitoring cornice systems in Longyeardalen – including one site previously examined by

Vogel et al. (2012) – with a terrestrial laser scanner (TLS). TLS – or ground-based lidar (light detection and ranging) – is an active remote sensing technology with documented applications for observing and monitoring various slope processes and hazards including landslides (Jaboyedoff et al., 2012; Prokop and Panholzer, 2009), coastal cliff erosion (e.g., Caputo et al., 2018), and rock slope instability (Abellán et al., 2014). TLS is being increasingly employed in snow and avalanche research to map snow depth and snow depth change (e.g., Deems et al., 2013; Fey et al., 2019; Prokop, 2008; Schirmer et al., 2011). Other specific snow-related applications include quantifying snow drift processes to verify physical models (Mott et al., 2011; Schön et al., 2015; Vionnet et al., 2014), observing avalanche activity to calibrate dynamic avalanche models (Prokop et al., 2015), assisting avalanche control operations (Deems et al., 2013), and planning and designing snow fences that limit hazardous snow accumulation in avalanche release areas (Prokop and Procter, 2016).

We monitored cornice accretion, deformation, failure, and associated cornice fall avalanche activity near Longyearbyen, Svalbard, with TLS technology over two winter seasons (2016–2017 and 2017–2018). To our knowledge TLS has not been employed to specifically monitor cornice dynamics, so our primary objectives are to use the high-spatial-resolution snow surface data acquired via TLS to

1. demonstrate the utility of TLS to observe cornice processes;
2. observe and quantify cornice accretion, deformation, failure, and associated cornice fall avalanches and link these processes to their controlling meteorological factors;
3. use our findings to provide suggestions for forecasting cornice fall avalanches in this and other locations threatened by cornices.

2 Study area

The present study focuses on the cornices forming above Longyeardalen (“the Longyear valley”) in central Svalbard (Fig. 1). Longyeardalen is a glacially sculpted, U-shaped valley with a northeast–southwest-oriented axis running approximately 3 km from the termini of two small mountain glaciers to a fjord. The Gruvefjellet and Platåberget plateaus border Longyeardalen to the west and east, respectively, with Svalbard's administrative center, Longyearbyen, situated in the valley bottom. The Gruvefjellet and Platåberget slopes lie within the horizontally bedded, lower-Tertiary Van Mijenfjord Group of sandstones and shales (Major et al., 2001). Resistant strata within this group form the area's extensive plateau topography. The entire region is underlain by continuous permafrost ranging in thickness from 100 m near the

Figure 1. Overview of Longyearfjorden and key locations, including automated weather stations (AWSs), mentioned in the text. Contour lines in (a) are spaced at 100 m. The location and direction from which the photo in (b) was taken is indicated by POV (point of view) in (a). The location and extent of the Gruvefjellet and Platåberget study sites are indicated in (a) and (b) with green and red shading, respectively. Locations of scan positions SP1, SP2, SP3, and SP4 as well as the orientation of the scanner at each position are also indicated.

coasts to over 500 m in the higher mountains (Humlum et al., 2003).

We investigated seasonal cornice dynamics and cornice fall avalanches along and under the Gruvefjellet and Platåberget plateau margins, respectively (Fig. 2). The steep valley walls descending from the broad plateau summits (approximately 450 m elevation) are characterized in their upper portions by protruding resistant bedrock buttresses and transport couloirs incised by fluvial and gravitational slope processes. The Gruvefjellet slope described in detail by Eckerstorfer et al. (2013) consists of a 50–70 m near-vertical bedrock cliff situated under the plateau margin and over a 40–50° slope that serves as a slab avalanche release area. This broad slope transitions into the transport couloirs which in turn feed extensive avalanche fan deposits downslope. Similar morphology exists on the Platåberget slope, but the plateau margin transitions directly into discrete 45–55° release areas leading into the couloirs and lacks the near-vertical bedrock face present on Gruvefjellet.

Central Svalbard's climate is cold and arid, with a mean annual air temperature of -4.6°C and mean annual pre-

cipitation of 191 mm at the Svalbard Airport automated weather station (AWS) for the 1981–2010 period of record (Førland et al., 2011). Combined mean winter (DJF) and mean spring (MAM) precipitation for 1981–2010 is 86 mm (Førland et al., 2011). Mean winter air temperature for the same period is -11.7°C and mean spring air temperature is -8.3°C (Førland et al., 2011). Rapidly increasing air temperatures in the winter and spring in response to decreased sea ice extent (Isaksen et al., 2016) create difficulties establishing representative baseline temperature conditions, with recent reports indicating warming on the order of 3–5 °C for Svalbard as a whole from 1971 to 2017 (Hanssen-Bauer et al., 2019). Less clear changes are apparent in the precipitation trends (e.g., Hanssen-Bauer et al., 2019), but midwinter rain-on-snow events are dramatically increasing in frequency (e.g., Vikhamar-Schuler et al., 2016).

The climate of Svalbard prohibits the growth of woody vegetation, and snow distribution across the landscape is thus strongly controlled by the wind (e.g., Jaedicke and Sandvik, 2002). Southeasterly winds generally prevail across the region's plateau mountains but often switch to westerly or southwesterly during winter storms and are frequently redirected along the major valley axes at lower elevations (Christiansen et al., 2013). Winter weather in central Svalbard fluctuates between extended periods of cold, stable high pressure punctuated by warm, wet low-pressure systems conveyed northwards along the North Atlantic cyclone track (Hanssen-Bauer et al., 1990; Rogers et al., 2005). This is reflected in the region's snow and avalanche climate, where the snow-pack typically consists of persistent weak layers formed during high pressure interspersed with wind slabs or ice layers formed during snowstorms or rain-on-snow events (Eckerstorfer and Christiansen, 2011a). Avalanche activity here displays a strong topographical and meteorological control, with direct action slab avalanches clustered around winter storms and the region's plateaus serving as source areas for the extensive cornice systems that contribute to frequent cornice fall avalanches (Eckerstorfer and Christiansen, 2011b).

3 Methods

3.1 Automated snow and weather data

We obtained wind and air temperature data from the Gruvefjellet automated weather station (AWS), precipitation data from the Svalbard Airport AWS, and a limited time series of snow depth data from a pair of ultrasonic snow depth sensors placed in avalanche release areas on Gruvefjellet and Platåberget during the 2017–2018 winter season (Figs. 1 and 2). We defined the winter season as 1 December to 30 June for the purposes of this study. The Gruvefjellet AWS is located less than 500 m east of the Gruvefjellet cornice system at an elevation of 464 m and records hourly meteorological data. The Svalbard Airport AWS is situated approx-

Figure 2. Overview of the cornice systems and locations of the primary spatial data employed in this work. Panel (a) shows Gruvefjellet from SP1 taken on 21 March 2017, with the white rectangle approximating the 600 m horizontal extent of (b). Panel (b) indicates the location of the 2-D cross-sectional profiles GF1, GF2, and GF3. Panel (c) shows Platåberget from SP4 taken on 24 May 2017, with the white rectangle approximating the 600 m horizontal extent of (d). Panel (d) indicates the location of the 2-D cross-sectional profiles PB1, PB2, and PB3.

imately 5 km northwest of the study area at 28 m and is the only weather station in the region with long-term precipitation measurements.

As part of the installation of a network of automated snow monitoring stations in Longyeardalen (Prokop et al., 2018), we mounted two ultrasonic snow depth sensors in avalanche release areas under the cornice systems in autumn 2017. These sensors were located at 350 and 450 m elevation on Gruvefjellet and Platåberget, respectively (Fig. 2). We employed the Campbell Scientific SR50A ultrasonic distance sensor to measure snow depth at each location. The snow sensors began recording reliable snow depth data on 15 November 2017 and continued until the end of the 2017–2018 season at 10 min intervals.

3.2 Terrestrial laser scanning (TLS) and post-processing

We used a Riegl® Laser Measurement Systems VZ-6000 ultra-long-range terrestrial laser scanner to repeatedly scan the Gruvefjellet and Platåberget cornice systems throughout the 2016–2017 and 2017–2018 winter seasons. The VZ-6000's 1064 nm operating wavelength is particularly well-suited for measuring snow surfaces, while the high scanning speed and measurement range up to 6 km with a 30 kHz pulse repetition rate ensured adequate data acquisition capabilities across the study area in a variety of atmospheric conditions (Riegl® Laser Measurement Systems, 2019; Prokop, 2008).

We use data from 25 scans of Gruvefjellet and 22 scans of Platåberget during the duration of the study (Appendix A). Of these, one scan from Gruvefjellet and Platåberget each is

a snow-free surface taken on 16 September 2016. For Gruvefjellet, we acquired usable snow surface data from 18 scans during the 2016–2017 season and seven scans during 2017–2018. We acquired 14 snow surface scans of Platåberget during 2016–2017 and seven scans during 2017–2018. The TLS was unfortunately damaged in late April 2018, and we were unable to acquire any scans after our final scan on 13 April 2018.

We preprocessed the raw point clouds in RiSCAN Pro, Riegl's proprietary data processing software. We established a suite of ground control points on both Platåberget and Gruvefjellet using a differential global positioning system (DGPS) which we used to georeference individual point clouds. We then aligned repeated snow-covered scans to the snow-free scans established in September 2016 using these ground control points and the “Multi-Station Adjustment” plugin in RiSCAN Pro following the approach outlined by Prokop and Panholzer (2009). We then manually filtered non-ground points or points above the snow surface. Finally, we applied an octree filter with a 0.10 m increment and exported to an XYZ text file.

We imported individual point clouds into CloudCompare (CloudCompare, 2019) for further analyses (Fig. 3). To create 2-D cornice profile cross sections, we extracted point cloud profile sections along manually defined axes using the polyline extraction tool native to CloudCompare (Fig. 3c and d). This tool requires user-defined inputs for profile type, section thickness, and maximum edge length which we set to “both”, 0.6 and 0.2 m. We then manually edited and digitized the resulting shapefiles in the ArcScene 3-D Editing environ-

Figure 3. Visualization of the point cloud processing methods in CloudCompare. Panel (a) shows a photo of the cornice represented by profile GF1 on 21 March 2017. Panel (b) shows the same surface as represented by the 0.10 m point cloud. The manually defined axis of GF1 is indicated by the white line. Panels (c) and (d) show the 21 March scanned surface and extracted profile from two vantage points. Panel (e) displays both the 21 March (colored points) and bare-earth (white points) surfaces oriented parallel to the projection direction, with the 21 March profile (green) and bare-earth profile (white) also indicated. Panels (f) and (h) display similar data but with the surfaces oriented roughly perpendicular to the projection direction (shown with red arrows), and (h) shows a cross section of the surface shown in (e) and (f). The 1 m grid showing the horizontal differences between the 21 March and bare-earth scans is displayed in (g). All scale bars are in meters.

ment (ArcGIS 10.4.1) to create the vertical cornice profile schematics as 3-D shapefiles.

We calculated representative volumes for selected areas from both the Platåberget and Gruvefjellet cornice systems using the “compute 2.5-D volume” tool in CloudCompare. This tool computes the volume between two 2.5-D point clouds by rasterizing the point clouds to a specified grid size and then computing volumes based on the differences in a specified projection direction between the rasterized values (Fig. 3e–h). In our case, we rasterized our point clouds to a 1 m grid and calculated horizontal distance differences along the “X” projection direction, which in our georeferenced point clouds corresponds to east–west (i.e., the slope

fall lines). For each cornice system, we computed the volume of snow in a 40 m × 8 m area of the plateau margin for each usable snow surface scan by subtracting the bare-earth surface from the scanned snow surface (Fig. 3e–g). We chose this areal extent to maximize coverage of an individual cornice throughout its development during the season (i.e., to completely capture the vertical extension of the leading edge) while minimizing volume changes related to other snow on the slope.

We used the Multiscale Model-to-Model Cloud Comparison (M3C2) algorithm developed by Lague et al. (2013) and implemented as a plugin in CloudCompare to quantify changes in the cornices and snow surfaces on the slopes be-

low in 3-D. The M3C2 algorithm allows for direct comparison of point clouds in 3-D and is specifically developed to handle 3-D differences and detect changes to complex surfaces where both vertical and horizontal changes exist (Lague et al., 2013). This functionality requires the user to input the following parameters: the normal scale, the projection scale, and the maximum depth (e.g., Lague et al., 2013; Watson et al., 2017). We selected a normal scale of 2 m oriented positively to the scan position (i.e., the normals “face” the scan position), a projection scale of 1 m, and a maximum depth of 10 m for all M3C2 calculations.

TLS-based snow surface measurement accuracy generally decreases with increasing distance from the scanner to the measured snow surface and is affected by the manner in which the laser beam interacts with the snow surface, the local terrain characteristics, the stability of the scanner while scanning, and the quality of the scan data registration process (Fey et al., 2019; Hartzell et al., 2017; Prokop et al., 2008). The relative accuracy – the deviation between measurements of an unchanged surface taken under different measurement conditions – can be assessed to quantify uncertainties related to both registration errors and positional errors from the interaction of the laser beam with the surface (Fey et al., 2019; Prokop and Panholzer, 2009). We assessed relative accuracy for our data by measuring M3C2 distances between each snow-covered scan and the snow-free scan on a 10 m × 10 m area of stable, snow-free rock faces near the cornices on both Gruvefjellet and Platåberget (Fig. 2). We were unfortunately limited to this single area on which to conduct accuracy assessments because all other surfaces near the cornices became snow covered at some point during the study. We report relative accuracy for each snow-covered scan as the mean of all M3C2 distances on the 10 m × 10 m area (Appendix A). This location for the relative accuracy assessment was selected based on its ability to remain functionally snow-free throughout the study and because it was not otherwise used in the registration process. As both registration and positional errors can be spatially variable across the scanned area (Fey et al., 2019; Hartzell et al., 2017; Prokop, 2008), we used this location in close proximity to the cornices of interest to best represent the relative accuracy near the cornices. Mean M3C2 distance values are smaller than 80 mm for all scans, with standard deviations ranging from < 10 to 72 mm (Appendix A). Uncertainty associated with the relative volume metric, calculated by multiplying the relative accuracy of each scan by the surface area considered in the volume calculations (369 m²) thus ranged from less than 1 to 28 m³ (Appendix A).

3.3 Supplemental observational data

We relied on snow and avalanche observations from Platåberget and Gruvefjellet from the Norwegian Water Resources and Energy Directorate’s (NVE) online observation platform regObs (<https://www.regobs.no/>, last access: 6 October 2019)

to supplement our TLS observations. Local observers conduct snow and avalanche assessments on the Gruvefjellet and Platåberget slopes on a sub-weekly basis, so we were able to much better constrain avalanche cycle timing than with the temporal resolution available from the TLS data.

4 Results

4.1 Seasonal summaries of meteorological conditions and cornice dynamics

4.1.1 2016–2017

We compare seasonal meteorological conditions (Fig. 4) with cross-sectional cornice profiles derived from eight scanned snow surfaces on Gruvefjellet and seven surfaces on Platåberget. We selected these profiles from a pool of 18 usable scans from Gruvefjellet and 14 from Platåberget (Appendix A) to represent key points in the development of the cornice systems.

Small cornices had accumulated on Gruvefjellet by 2 December 2016. Maximum horizontal cornice growth prior to this scan occurred in the vicinity of profile GF2, where both vertical and horizontal cornice growth exceeded 1 m from the edge of the plateau (Fig. 5). The representative cornice volume in the vicinity of profile GF1 already approached 200 m³. Temperatures remained below freezing over the next month, and daily averaged wind speeds exceeded 10 m s^{−1} only on 29 December 2016. By 12 January 2017, the representative cornice volume on Gruvefjellet had more than tripled relative to early December to over 600 m³. Horizontal cornice extension along the Gruvefjellet cornice system exceeded 4 m in most locations, with maximum horizontal extension near profile GF1 exceeding 5 m (Fig. 5). The representative cornice volume of just over 300 m³ from the Platåberget cornices on the same date shows considerably less cornice growth (Fig. 4).

Heavy snowfall followed by strong westerly winds preceded several cornice fall avalanches on 21 January on Platåberget (Fig. 4, Table 1). Representative cornice volume on Platåberget nearly doubled from roughly 300 to over 600 m³ between the 12 and 21 January scans. Horizontal accretion on profile PB2 exceeded 3.5 m, resulting in an accretion rate of 17 mm h^{−1} (Table 2). The cornice represented by profile PB1 failed, triggering a cornice fall avalanche (Size D2, R3 after American Avalanche Association, 2016) which reached the road at the foot of the slope. The failure plane represented by the 21 January profile on PB1 does not extend back to the 12 January surface, suggesting newly accreted snow comprised the bulk of the failure (Fig. 5). Cornices on Gruvefjellet experienced comparably minor changes during this event, with the representative volumes decreasing by just 30 m³ and minimal changes evident in the profiles (Fig. 5).

Figure 4. Summary of the representative cornice volume progression and meteorological conditions for the 2016–2017 winter season. Wind speed and air temperature are daily averaged values from the Gruvefjellet AWS, and precipitation data are daily values from the Svalbard Airport AWS measured at 06:00 UTC. Shaded blue vertical bars indicate well-constrained cornice accretion periods for which we were able to calculate horizontal cornice accretion rates (Table 2). Shaded grey vertical bars indicate 48 h periods with observed noteworthy cornice fall avalanche activity (Table 1).

Table 1. Summary of avalanche cycles.

Event date	Area	Trigger	Pre-event scan date	Post-event scan date	No. of observed cornice fall avalanches
21 January 2017	Platåberget	Accretion	12 January 2017	22 January 2017	> 3
9 April 2017	Gruvefjellet	Accretion	21 March 2017	25 April 2017	1
21 April 2017	Gruvefjellet	Accretion	21 March 2017	25 April 2017	1
29 April 2017	Platåberget	Accretion	25 April 2017	1 May 2017	> 3
14 January 2018	Gruvefjellet	Temperature (rain)	15 December 2017	24 January 2018	1
18 March 2018	Gruvefjellet	Accretion	2 March 2018	23 March 2018	2

A major accretion event in mid-February 2017 followed several weeks of unseasonably high temperatures at cornice elevation during early February (Fig. 4). Locally heavy snowfall and strong easterly winds accompanying a vigorous winter storm impacted the region between 19 and 21 February. Profile GF1’s horizontal extension increased by nearly 3 m between the 17 and 24 February scans, resulting in horizontal accretion rates exceeding 15 mm h^{-1} (Table 2). The representative volume increased by approximately 100 m^3 during the same timeframe. The strong easterly winds stripped the Platåberget cornice system on the windward side of the valley reflected by the abrupt decrease of over 100 m^3 to the representative volume there.

Representative volumes for both cornice systems gradually increased in the following month, and profiles from 21 March 2017 show considerable rounding and downslope creep of the cornices’ leading edges in profiles GF1 and GF3 (Fig. 5). Cornices continued to grow on Platåberget, with horizontal growth exceeding 2 m on portions of the PB1 and PB2 profiles and PB3’s vertical extent increasing by over 2 m. The Platåberget cornices did not deform downslope to the same degree as the Gruvefjellet cornices during this time period.

A representative volume decrease of over 500 m^3 (roughly 50 % of the volume) on Gruvefjellet in April is related to a major cornice failure near profile GF1, while Platåberget’s representative volume increased by 150 m^3 in an accretion event near the end of the month (see Sect. 4.2.1). Considerable cornice accretion is evident in all cornice profiles between 21 March and 1 May except for profiles GF1 and PB2 where we documented cornice failures. Representative volumes continued to increase in early May as light precipitation coincided with continued subzero temperatures. Representative cornice volume on Gruvefjellet gradually increased through 31 May and then dramatically decreased with the onset of sustained positive temperatures at the Gruvefjellet AWS. Cornices on Platåberget continued to accrete through the 18 May scan before beginning to melt away between 18 May and 9 June.

4.1.2 2017–2018

We gathered seven scanned snow surfaces from both Gruvefjellet and Platåberget for the 2017–2018 season with which to compare to meteorological conditions. Cornice develop-

Table 2. Summary of well-constrained accretion events.

Area	Pre-event scan date and time (UTC)	Post-event scan date and time (UTC)	Between interval scan (h)	Profile with max horizontal accretion	Max horizontal accretion (m)	Accretion rate (mm h ^{−1})
Platåberget	12 January 2017 19:30	21 January 2017 21:00	217.5	PB2	3.6	17
Gruvefjellet	17 February 2017 09:00	24 February 2017 11:00	170	GF1	2.9	17
Platåberget	25 April 2017 13:15	1 May 2017 09:45	140.5	PB1	2.0	14

Figure 5. The 2-D cornice profiles showing cornice progression for selected scan dates throughout the 2016–2017 winter season. Each profile is labeled as it is referred to in the text and corresponds to the location and POV depicted in Fig. 2.

ment during the 2017–2018 winter season differed considerably from the 2016–2017 winter season despite relatively similar seasonal meteorological conditions (Table 3). Gruvefjellet profiles from 15 December 2017 show over 5 m of horizontal cornice growth in all profiles, and representative volume approached 1000 m³ (Figs. 6 and 7). Contrastingly, the Platåberget plateau margin remained functionally free of snow. Cornices continued to grow over the following 5 weeks

on Gruvefjellet up to the 24 January scan, with profiles GF1 and GF3 reaching their maximum horizontal extensions for the season of nearly 7 and over 8 m, respectively (Fig. 7). Cornice fall avalanches observed on 13 January 2018 are evident in the decreased cornice extension in GF2 in the 24 January scan relative to the 15 December 2017 surface and were associated with positive air temperatures and rain at cornice elevation (Table 1). Profiles on Platåberget on 24 January 2018 do not show cornice development, with snow accumulating relatively parallel to the underlying topography.

Representative volume doubled on Platåberget between the 31 January and 22 February scans from 400 to 900 m³. This coincided with a 0.34 m increase in snow depth at the snow sensor during a snowstorm on 5 and 6 February where 14 mm of precipitation was measured at the Svalbard Airport AWS (Fig. 6). Cornice system changes were more minimal on Gruvefjellet, with a subtle increase of 100 m³ in representative volume. Measured snow depth on Gruvefjellet increased from 1.45 m on 31 January to a maximum of 1.77 m on 13 February, before slowly decreasing back to 1.48 m by 22 February (Fig. 6). A minor decrease in horizontal extension (< 1 m) and slight downslope deformation exhibited in profile GF1 are the main observed changes to the cornice profiles between 31 January and 22 February (Fig. 7).

Snow depths increased by 0.20 and 0.28 m on Gruvefjellet and Platåfjellet, respectively, on 26 and 27 February 2018 as over 7 mm precipitation was recorded at the airport (Fig. 7). A marked increase in representative volume of 230 m³ on Platåberget between the 22 February and 2 March scans coincides with an increase in snow depth of 0.28 cm over 26 and 27 February. Although a small cornice is evident in profile PB3 on 2 March, increased volume during this time illustrates slope-normal snow depth increase rather than cornice accretion in the representative volume area in the vicinity of PB2 (Fig. 7). On Gruvefjellet, downslope creep of the cornice masses continued, with maximum vertical deformation close to 0.80 m for the leading edge of profile GF1. A winter storm on 18 March 2018 resulted in cornice failures seen in both GF1 and GF2 and decreased representative volume on

Figure 6. Summary of the representative cornice volumes and meteorological conditions for the 2017–2018 winter season. Wind speed and air temperature are daily averaged values from the Gruvefjellet AWS, precipitation data are daily values from the Svalbard Airport AWS measured at 06:00 UTC, and snow depths are daily averages from the snow sensors on Gruvefjellet and Platåfjellet. Shaded grey vertical bars indicate 48 h periods with observed noteworthy cornice fall avalanche activity (Table 2).

Table 3. Seasonal summaries. All parameters are measured at the Gruvefjellet AWS except for precipitation, which is measured at the Svalbard Airport AWS.

	2016–2017	2017–2018
Mean seasonal air temperature (°C)	−9.3	−7.5
Accumulated precipitation (mm)	125.6	124.5
Percentage of hours in season with accretion winds on Platåberget	5.7	5.1
Percentage of hours in season with accretion winds on Platåberget and daily precipitation > 0.2 mm	3.0	1.7
Percentage of hours in season with accretion winds on Gruvefjellet	13.5	21.7
Percentage of hours in season with accretion winds on Gruvefjellet and daily precipitation > 0.2 mm	4.0	4.1
Accretion winds on Platåberget: wind speed > 5 m s ^{−1} ; 225° < wind direction < 315°		
Accretion winds on Gruvefjellet: wind speed > 5 m s ^{−1} ; 45° < wind direction < 135		

Gruvefjellet, while scouring reduced volume during this time on Platåberget (see Sect. 4.2.2). Minimal further changes are evident in season’s final scan (due to scanner failure) taken on 13 April 2018.

4.2 Case studies

4.2.1 Cornice accretion and failure in April 2017

We documented three periods of cornice fall avalanche activity with TLS data in April 2017. In the first, a small portion of the cornice between profiles GF1 and GF2 failed on 9 April following a period of precipitation falling as snow and easterly winds in excess of 10 m s^{−1} (Figs. 8a and 9a, annota-

tion 1). The cornice represented by profile GF1 then failed completely on 21 April 2017 coincident with trace precipitation falling as snow and 2 d of moderate to strong easterly winds (Figs. 8b and 10). Negative M3C2 distances displaying changes to the Gruvefjellet cornice system between the 21 March and 25 April 2017 scans show the largest portion of the failed cornice along the axis of profile GF1 (Fig. 9a, annotation 2). This failure extended northwards almost 40 m along the cornice. Negative M3C2 distances on the vertical rock face immediately downslope of both the 9 and 21 April cornice failures show how the falling cornice blocks remove snow from the rock face before impacting avalanche release areas below (Fig. 9a, annotation 3). Here, cornice impact

Figure 7. The 2-D cornice profiles showing cornice progression for 2017–2018 winter season scan dates. Each profile is labeled as it is referred to in the text and corresponds to the location and POV depicted in Fig. 2.

craters and small slab avalanche releases are apparent in negative M3C2 distances (Fig. 9a, annotation 4). Lower on the slope, the cornice fall avalanche deposition – complete with intact cornice chunks in the avalanche debris – is apparent in strongly positive M3C2 distances (Fig. 9a, annotation 5). Other positive M3C2 distances along the cornices (Fig. 9a, annotation 6) and horizontal and vertical extent increases on profiles GF2 and GF3 (Fig. 10) show cornice accretion occurred elsewhere along Gruvefjellet during this time span. The easterly winds stripped the cornices on Platåberget, evidenced by representative volume decreases of nearly 200 m³ and vertical extension decreases of up to 1.5 m at profile PB3 (Fig. 10).

A warm winter storm accompanied by 4.5 mm of precipitation, southwesterly winds, and air temperatures approaching 0 °C at cornice level resulted in a period of major cornice accretion and associated cornice fall avalanche activity on the Platåberget cornice system between the 25 April and 1 May 2017 scans (Fig. 8c). Widespread cornice failures are shown by negative M3C2 distances along the Platåberget plateau margin (Fig. 9b, annotation 1). These failures coincide with positive M3C2 distances in excess of 1.5 m indicative of cornice accretion elsewhere along the plateau margin (Fig. 9b, annotation 2). Profile PB3, for example, ex-

perienced over a meter of increased vertical cornice extension (Fig. 10) just south of a cornice failure shown in the M3C2 distances (Fig. 9b, annotation 3). In profile PB1, 2 m maximum increases in horizontal extension resulted in accretion rates of 17 mm h⁻¹ (Table 2). The semi-vertical profile surface shown in profile PB2 (Fig. 10) combined with the M3C2 distance decreases in the profile's immediate surroundings (Fig. 9b, annotation 4) indicate cornice failure here occurred after some vertical cornice accretion, as the failure plane extends above the cornice roof from the 25 April snow surface. Cornice blocks released from this cornice and the one immediately to the north poured over cliffs further downslope and gouged impact craters (Fig. 9b, annotation 5) before releasing slab avalanches lower on the slope (Fig. 9b, annotation 6). Minimal changes to the Gruvefjellet cornices occurred during this event.

4.2.2 Cornice accretion and failure in March 2018

A storm in mid-March 2018 punctuated a month of otherwise stable weather and resulted in cornice fall avalanches on Gruvefjellet (Fig. 11a). From 15 to 19 March, 5.6 mm of precipitation accumulated at the airport AWS, snow depths at the Gruvefjellet sensor increased by a maximum of 18 cm while those at the Platåberget sensor decreased by approximately 0.25 m, and strong winds blew from the ENE for 24 h on 17–18 March. Two large cornice failures on Gruvefjellet visible as strongly negative M3C2 distances near profile GF1 and slightly to the north (Fig. 12a, annotation 1) triggered avalanches on the slope below (Fig. 12a, annotation 2). Similar to the morphology observed in the April 2017 cornice fall avalanches, the failed cornice blocks stripped snow off the vertical rock face and created impact craters while entraining snow as they moved downslope. The cornice chunks from these cornice failures also remained intact throughout the event and ran further than the rest of the avalanche debris (Fig. 12a, annotation 3). A cornice block approximately 5 m in horizontal extension detached from the cornice represented by profile GF1, while a smaller (< 1 m horizontal extension) piece detached near GF2 (Fig. 13). The GF3 profile did not fail, but over 1 m of snow accreted vertically on the cornice's leading edge. By contrast, Platåberget's plateau margin lost snow, with snow depth decreases in excess of 0.20 m measured at the snow station and strongly negative M3C2 distances across the upper portion of the Platåberget release areas (Fig. 12b, annotation 1).

5 Discussion

5.1 Seasonal cornice dynamics

TLS-derived cornice data from the 2016–2017 and 2017–2018 winter seasons provide quantitative reinforcement to the conceptual models of cornice dynamics developed in previous studies (e.g., Montagne et al., 1968; Vogel et al., 2012).

Figure 8. Meteorological summary of the April 2017 case study. Wind speed, wind direction, and air temperature are hourly values from the Gruvefjellet AWS, and precipitation data are daily values from the Svalbard Airport AWS measured at 06:00 UTC. Colored vertical lines in the time series indicate the scan timing corresponding to the profiles in Fig. 9. Vertical grey bars marked (a)–(c) correspond to 48 h time periods with noteworthy avalanche activity discussed in the text.

Figure 9. M3C2 distances displaying changes to the snow cover on Gruvefjellet between the 21 March and 25 April 2017 scans (a) and on Platåberget between the 25 April and 1 May 2017 scans (b). Red rectangles in both panels indicate the locations of the cornice profiles. Specific snow surface features are annotated as they are referred to in the text.

Figure 10. Cornice profiles illustrating cornice dynamics during the April 2017 case study, with each profile labeled as it is referred to in the text. Dashed lines indicate interpolated data where overhanging cornice structure shadowed the snow surface from the TLS.

Figure 11. Meteorological summary of the March 2018 case study. Wind speed, wind direction, and temperature are hourly values from the Gruvefjellet AWS, and precipitation data are daily values from the Svalbard Airport AWS measured at 06:00 UTC. Colored vertical lines in the time series indicate the scan timing corresponding to the profiles in Fig. 12, and the grey vertical bar annotated with (a) corresponds to the 48 h time period with noteworthy avalanche activity mentioned in the text.

Figure 12. M3C2 distances displaying changes to the snow cover on Gruvefjellet (a) and Platåberget (b) between the 2 and 23 March 2018 scans. Red rectangles in both panels indicate the locations of the cornice profiles. Specific snow surface features are annotated as they are referred to in the text, and snow depth sensors are marked and labeled.

Figure 13. Cornice profiles illustrating cornice dynamics during the March 2018 case study, with each profile labeled as it is referred to in the text. Dashed lines indicate interpolated data where overhanging cornice structure shadowed the snow surface from the TLS.

In these models, cornices accrete through relatively discrete events and begin to deform under their own weight before either failing or melting away towards the end of the snow season.

Our data show cornices can rapidly accrete at any point in the snow season given abundant snow available for wind transport, wind speeds sufficient to mobilize surface snow, and wind directions oriented relatively perpendicular to the ridgeline. We documented accretion rates in excess of 15 mm h^{-1} at various times throughout the 2016–2017 season on each side of the valley, which can be considered a minimum given the relatively poor temporal constraints on the accretion events provided by the TLS data. Each of these periods of accretion coincided with measured precipitation at the airport, wind speeds in excess of 5 m s^{-1} , and wind directions roughly placing the plateau margin in the lee. The relatively small proportion of the winter seasons characterized by meteorological conditions conducive for accretion suggests just a few accretion events play a key role in cornice development each season (Table 3). Asynchronous cornice responses on Gruvefjellet and Platåberget to specific weather events further illustrate the importance of wind direction in controlling cornice dynamics. During the February 2017 event, for example, cornices on Gruvefjellet rapidly accreted and gained volume with plentiful snow available for transport and strong easterly winds. Cornices on Platåberget lost volume, however, as they were eroded by the same easterly winds. Similar out-of-phase behavior was exhibited in late April 2017, when precipitation and westerly winds resulted in considerable cornice growth on Platåberget accompanied by slight decreases to horizontal and vertical extension in profiles GF2 and GF3 and minimal representative volume changes near profile GF1.

Following initial accretion, the cornices' leading edges begin to deform downslope. Deformation becomes more pronounced later in the season, presumably as increased air temperatures and solar radiation begin to warm the snow, decreasing the stiffness of the cornices and increasing creep (e.g., Schweizer et al., 2003). Further accretion events can then be superimposed on this deformation as the season progresses, with short accretion events interspersed by longer periods of downslope creep. This can be seen in the minor increases in horizontal extension and continued downslope deformation in GF1 and GF3 through the latter portion of the 2017–2018 season (Fig. 7). Cornice accretion and downslope deformation can also occur almost simultaneously with air temperatures approaching or even exceeding freezing at cornice level, as evidenced by the rapid accretion and downslope creep shown in profile PB1 for the 25 April–1 May scan interval (Fig. 10).

While meteorological conditions control the specific timing of cornice accretion and downslope deformation, the underlying topography appears to act as a fundamental control on cornice structure and seasonal cornice dynamics. The presence of the steep bedrock face directly beneath the

Gruvefjellet plateau margin limits the support provided by the underlying topography compared to the gentler sloping Platåberget margin. The result is a more overhung cornice structure on Gruvefjellet, while Platåberget's topography allows for more slope-normal snow accumulation. Profiles PB1 and PB2 failed to develop cornices at all during the 2017–2018 winter season (Fig. 7). The presence of cornices with horizontal extension approaching 5 m in these locations during the 2016–2017 winter season (Fig. 5), however, shows the topography can support cornice development given the right meteorological conditions. Differences in meteorological conditions between the 2016–2017 and 2017–2018 winter seasons may provide a partial explanation for differing seasonal snow cover responses on Platåberget (Table 3). Winds in excess of 5 m s^{-1} – a conservative estimate for threshold wind speeds required to mobilize loose snow (Li and Pomeroy, 1997) – from the western quadrant conducive to cornice accretion on Platåberget were slightly less prevalent during the 2017–2018 season, and these winds also coincided with precipitation roughly half as frequently as during the 2016–2017 season. Easterly winds exceeding 5 m s^{-1} were considerably more prevalent during the 2017–2018 season, which may have increased cornice scouring or limited snow available for transport – and thus accretion – on Platåberget. Nevertheless, the meteorological differences between the two winter seasons are subtle enough when compared to the noteworthy differences in cornice dynamics to suggest specific interactions between meteorology and topography not necessarily captured by our analyses meaningfully impact cornice development.

Topography also seems to control the relative size of cornice failures. Vogel et al. (2012) describe a “geomorphologically determined sedimentary step approximately 3 m below the plateau that most likely acts as the cornice pivot point” on Gruvefjellet. This pivot point is most evident in profile GF1, where in both winter seasons the downslope creep of the overhung cornice beyond this pivot point ultimately became overburdened during an accretion event and caused the cornice to fail completely. The cornice represented by GF1 has the least topographic support and developed the most overhanging cornice structure of the specific cornices we investigated, and also failed completely both seasons. By contrast, the topographic support provided by Platåberget does not promote overhanging cornices to the same degree, instead promoting a thicker slope-normal snowpack which in itself supports the cornice structure. Here, observed cornice failures such as that shown in PB2 during the 25 April–1 May 2017 scan interval (Fig. 10) are limited to the recently accreted snow and did not involve the entire cornice mass. Similarly, profile GF2 failed in March 2018 within hours of profile GF1's full failure but involved a much smaller portion of the cornice predating the 2 March scan – potentially related to increased topographic support to this cornice relative to GF1 (Fig. 13).

5.2 Cornice fall avalanches

Previous work has differentiated cornice fall avalanche types by the inferred mechanism of cornice failure – via either increased snow load from accretion or decreased snow strength in the cornice related to increased snow and air temperatures. Five of the six cornice fall avalanche events observed in this study coincided with winter storms leading to accretion just prior to cornice failure (Table 1). This is in contrast to previous findings from this location, where no cornice failures were observed in direct response to snow loading caused by a snowstorm (Vogel et al., 2012). The lone cornice fall avalanche event we cannot link to cornice accretion occurred in January 2018. This event coincided with heavy precipitation, but positive temperatures at the Gruvefjellet AWS and decreasing snow depths at the Gruvefjellet snow sensor indicate this precipitation fell as rain (Fig. 6). Our truncated TLS observation record in late spring 2018 unfortunately omits the May–June period found by Vogel et al. (2012) to be critical for air-temperature-induced cornice failures in this location, but observational records throughout this time do not indicate further cornice fall avalanches. Accretion's role in determining cornice failure is also reflected in the asynchronous timing of cornice failures on Gruvefjellet and Platåberget during our study. None of the observed avalanche events included activity on both Gruvefjellet and Platåberget simultaneously as would be expected with air-temperature-induced failures, with avalanches instead occurring only on the leeward aspect.

Observed cornice fall avalanche size appears to be controlled largely by the snow conditions in the underlying release area. Cornice fall avalanches on Gruvefjellet follow a pattern exemplified by the April 2017 case study in which the cornice fails and removes snow from the steep bedrock face below as it descends before impacting the release areas at the base of the cliff (e.g., Fig. 9a). The cornice block can then, depending on the snow conditions in the release area, entrain snow from its impact crater and the avalanche path below or trigger a larger slab avalanche. Cornice failures near profile GF3 in both April 2017 and March 2018 triggered small slab avalanches, but the majority of the avalanche debris resulted from entrainment as the cornice blocks bounced downslope.

Platåberget's topography promotes slightly different avalanche dynamics. The gentler slope at the plateau edge allows snow to accumulate directly under the cornices such that failed cornice masses land directly on the snow to be released as an avalanche. Release areas on Platåberget collect snow during accretion events much more efficiently than those on Gruvefjellet, where blowing snow mass losses due to suspension are promoted by the separation created between the cornices and the release areas by the bedrock cliff. Accumulation in the upper release areas on Platåberget coinciding with accretion events primes these locations for slab avalanche release with even small cornice fail-

ures. Relatively small cornice failures triggering larger slab avalanches on Platåberget in April 2017 resulted in magnitude avalanches (D2, R2–R3) comparable to those releasing from much larger cornice failures but less entrainable snow on Gruvefjellet in March 2018 (Figs. 9b and 13a).

5.3 Hazard management implications

Cornice fall avalanches are the most common avalanche type observed in the portion of central Svalbard surrounding our study area where the broad plateau summits and steep valley walls of Longyeardalen's topography are recurrent across the region (Eckerstorfer and Christiansen, 2011b). Cornice fall avalanches observed in this study thus represent processes occurring elsewhere throughout central Svalbard – and to a lesser extent other locations throughout the world – and provide an opportunity to reinforce existing forecasting frameworks with detailed cornice data. The conceptual model of avalanche hazard in North America treats cornice failure both as an individual avalanche problem to be considered by forecasters and as a potential trigger when assessing the likelihood of other avalanche types releasing in a given forecasting area and time period (Statham et al., 2018). Cornice fall avalanche hazard assessments should thus consider both the likelihood of cornice failure and the nature of the snow conditions in the release area to best judge cornice fall avalanche hazard. Our limited dataset, especially in the absence of multiple air-temperature-induced failures, is insufficient to make broad generalizations linking cornice failure type and resulting cornice fall avalanche activity. As a specific example, however, fairly widespread wind slab avalanche activity throughout the region accompanied each of the accretion-induced avalanche events observed in this work. The conditions leading to cornice accretion – strong winds and available snow for wind transport – also promote the development of wind slab problems. Thus, when conditions are favorable for cornice accretion and accretion-induced cornice failures, conditions are also favorable for the development of more widespread – and potentially more sensitive – slab avalanche problems. In this scenario, the chance of a cornice failure triggering a secondary slab avalanche would rise, subsequently amplifying the cornice fall avalanche hazard by also increasing the expected size of the resulting cornice fall avalanche. Furthermore, in all cornice fall avalanches observed on Gruvefjellet the main cornice blocks traveled further downslope than the rest of the avalanche debris. This pattern is apparent on larger failures on Platåberget as well, but is in some cases less obvious, likely due to the smaller cornice blocks being functionally indistinguishable from the avalanche debris. While the dataset presented here is insufficient to draw more quantitative conclusions regarding the runout distance of these cornice blocks, hazard management strategies should consider the destructive potential and extended runout of these blocks relative to the other entrained snow.

5.4 Uncertainties and limitations

The TLS data acquisition and processing techniques employed in this work allowed us to illustrate and quantify changes to the observed cornice systems in detail not previously achieved, but our results and subsequent interpretations are nonetheless limited by several factors. Measurement uncertainties specifically related to measuring snow surfaces with TLS are well-discussed in previous research (Deems et al., 2015; Prokop, 2008), but we introduced additional uncertainty to our results and interpretations due to the scan timing. Our TLS data acquisition scheme involved time-intensive manual input, so we were unable to achieve the temporal resolution required to better constrain individual accretion and cornice failure events. Decreasing time between scans would allow for more continuous and robust accretion rate calculations and could better constrain failure and avalanche snow surfaces, especially pre-event. Sufficiently decreasing the between-scan interval to a sub-daily resolution for such applications would likely require some degree of automation, and future work should consider employing a permanently installed TLS acquiring data automatically similar to systems employed for mining applications or slope stability assessments.

Uncertainties in cornice volume calculations are also affected by occasionally lengthy inter-scan intervals. Volume changes corresponding to specific meteorological conditions are in these cases aggregated across the entire scan interval, making disentangling the specific contributions to volume changes difficult. These conceptual uncertainties are magnified by the technical uncertainties related to TLS data acquisition. The TLS accuracy is of increased importance for volume quantification as measurement uncertainties are propagated throughout the volume calculation process. However, calculated volume uncertainties (Appendix A) are sufficiently low to instill a degree of confidence in the volume calculation process presented here. Finally, volume calculations are perhaps least robust in this study for times when the lack of obvious cornice structure makes calculating volumes particularly challenging (e.g., Platåberget during the 2017–2018 season).

Our experimental design focused on investigating the evolution and failure of the lower cornice surfaces from scan positions underneath the cornices where accessibility has precluded previous research. These scan positions did not, however, allow for systematic monitoring of the cornice roof. The orientation of the cornices' leading edges frequently shielded the cornice roof from the scanner, and our profiles often do not include the complete cornice roof. This also has implications for representative volume calculations, as uncertainty in the location of the cornice roof can result in inaccurate horizontal difference calculations in these specific locations. By failing to capture the cornice roof in our data, we also limit comparisons with earlier work on Gruevfjellet relating downslope cornice deformation and cornice failure to the

appearance of tension cracks between the cornice roof and the plateau anchoring point (Vogel et al., 2012). Future work should pair TLS data with some form of tension crack observation, and approaches combining TLS and unmanned aerial vehicle (UAV) photogrammetry present intriguing possibilities for future work in this and other locations.

TLS was shown to be a particularly suitable remote sensing tool for cornice monitoring in Svalbard where we were able to obtain useful data during the early winter seasons when the polar night precludes direct visual observation and cornice photography. Svalbard's unique environmental characteristics – such as the polar night – limit to a degree the applicability of our results to lower latitudes where more diurnal variations in radiation and temperature may influence cornice dynamics in ways not represented in Svalbard (e.g., Munroe, 2018). It is also unclear how representative the two winter seasons for which we present data are for the cornice systems in Longyear dalen, as previous research has also noted considerable differences in cornice dynamics between seasons (Vogel et al., 2012). Continued cornice monitoring in this and other lower-latitude settings would help clarify such uncertainties.

6 Conclusions

We monitored seasonal cornice dynamics and associated cornice fall avalanche activity with a TLS over two winter seasons in high-Arctic Svalbard. The spatial and temporal resolution at which we acquired snow surface data with the TLS allowed us to quantify changes to the cornices with sub-decimeter accuracy. These data provide quantitative reinforcement to existing conceptual models of cornice dynamics and further strengthen the validity of these models. Notable quantitative contributions from this work include documentation of conservatively calculated horizontal accretion rates well in excess of 10 mm h^{-1} and a methodology for calculating cornice volumes from TLS data.

This study demonstrated the viability of TLS methods for monitoring cornice dynamics. TLS methods for obtaining snow surface data are appropriate in Svalbard where the long polar night precludes data acquisition via other methods (e.g., photogrammetry), but techniques presented in this work are also suitable for cornices in other lower-latitude environments. Future work should investigate automated TLS data acquisition as an avenue to improve the temporal resolution of the measurements and better constrain cornice dynamics to specific meteorological conditions.

Our findings show complex interactions between topography, wind speed and direction, snow available for transport, existing snowpack, and cornice structure govern the growth, failure, and associated avalanche activity of the cornices in Longyear dalen. In particular, we show cornices rapidly accrete given winds strong enough to mobilize surface snow from a direction roughly perpendicular to the plateau edge,

placing the cornices in the lee. Our findings also reinforce previous work indicating an increased likelihood of cornice failure and associated avalanche activity during these periods of cornice accretion. This is encouraging for hazard managers seeking to forecast cornice fall avalanches, as anticipating the relatively infrequent conditions leading to cornice accretion can help predict periods of elevated cornice fall avalanche hazard. We observed the largest failures in our dataset in areas with minimal topographic support, demonstrating knowledge of the topography underlying the cornices can be beneficial when considering the specific location of cornice failure. Nevertheless, our limited dataset of cornice failures hinders conclusions drawn from this work, and continued work in a variety of environments is needed to better understand the specific mechanisms and dynamics of cornice fall avalanches.

Appendix A

Table A1. TLS data summary for the 2016–2017 winter season. n/a denotes “not applicable”.

Date and time (UTC)	Area	Scan position	Mean relative error (m)	Standard deviation (m)	Representative volume (m ³)	Volume uncertainty
16 September 2016 12:00	Gruvefjellet	S1	n/a	n/a	n/a	n/a
16 September 2016 13:00	Platåberget	S3	n/a	n/a	n/a	n/a
2 December 2016 10:15	Gruvefjellet	S1	0.036	0.001	179	13.173
5 December 2016 12:50	Gruvefjellet	S1	0.019	0.016	159	6.827
12 January 2017 17:30	Gruvefjellet	S4	0.047	0.028	649	17.343
12 January 2017 19:30	Platåberget	S3	0.067	0.039	328	24.612
21 January 2017 21:00	Platåberget	S3	0.032	0.044	619	11.956
22 January 2017 14:00	Gruvefjellet	S2	0.030	0.025	633	11.144
27 January 2017 08:50	Gruvefjellet	S1	0.024	0.016	689	8.930
3 February 2017 08:00	Platåberget	S4	0.000	0.044	688	0.148
14 February 2017 14:00	Platåberget	S4	0.066	0.039	1145	24.170
14 February 2017 15:00	Gruvefjellet	S1	0.022	0.020	789	7.970
17 February 2017 09:00	Gruvefjellet	S1	0.032	0.030	823	11.771
17 February 2017 10:00	Platåberget	S4	0.055	0.039	1190	20.111
22 February 2017 10:45	Platåberget	S3	0.042	0.037	1072	15.350
24 February 2017 11:00	Gruvefjellet	S1	0.027	0.031	970	9.926
12 March 2017 16:00	Platåberget	S4	0.003	0.046	1224	1.033
12 March 2017 17:30	Gruvefjellet	S2	0.034	0.027	1043	12.435
21 March 2017 13:10	Gruvefjellet	S1	0.031	0.027	1117	11.402
21 March 2017 14:10	Platåberget	S4	0.064	0.042	1255	23.616
25 April 2017 10:00	Gruvefjellet	S1	0.068	0.027	581	25.018
25 April 2017 13:15	Platåberget	S4	0.007	0.042	1291	2.731
1 May 2017 09:45	Platåberget	S4	0.045	0.071	1440	16.753
1 May 2017 10:25	Gruvefjellet	S1	0.034	0.020	591	12.472
8 May 2017 10:15	Platåberget	S4	0.015	0.045	1563	5.535
9 May 2017 08:25	Gruvefjellet	S1	0.041	0.023	654	15.166
18 May 2017 12:05	Gruvefjellet	S1	0.076	0.016	677	28.044
18 May 2017 13:00	Platåberget	S4	0.011	0.039	1620	3.948
31 May 2017 10:40	Gruvefjellet	S1	0.061	0.020	693	22.546
1 June 2017 13:15	Platåberget	S4	0.033	0.062	1599	11.993
9 June 2017 12:20	Gruvefjellet	S1	0.018	0.015	485	6.642
9 June 2017 13:30	Platåberget	S4	0.014	0.039	1379	5.203
14 June 2017 14:35	Gruvefjellet	S1	0.005	0.010	401	1.845

n/a = not applicable.

Table A2. TLS data summary for the 2017–2018 winter season.

Date and time (UTC)	Area	Scan position	Mean relative error ($\pm$ m)	Standard deviation (m)	Representative volume (m ³)	Volume uncertainty
15 December 2017 10:20	Gruvefjellet	S1	0.018	0.024	929	6.458
15 December 2017 11:00	Platåberget	S4	0.016	0.039	5	5.978
24 January 2018 11:20	Gruvefjellet	S1	0.049	0.031	938	18.081
24 January 2018 12:25	Platåberget	S4	0.069	0.062	419	25.277
31 January 2018 15:45	Gruvefjellet	S1	0.009	0.024	947	3.321
31 January 2018 17:00	Platåberget	S4	0.070	0.052	415	25.978
22 February 2018 10:30	Gruvefjellet	S1	0.020	0.025	1051	7.528
22 February 2018 11:30	Platåberget	S4	0.032	0.056	902	11.734
2 March 2018 11:55	Gruvefjellet	S1	0.003	0.037	1031	1.144
2 March 2018 12:30	Platåberget	S4	0.024	0.072	1130	8.819
23 March 2018 13:15	Gruvefjellet	S1	0.041	0.023	539	15.092
23 March 2018 14:00	Platåberget	S4	0.032	0.049	889	11.845
13 April 2018 11:00	Gruvefjellet	S1	0.024	0.021	593	8.745
13 April 2018 13:20	Platåberget	S4	0.000	0.044	1053	0.037

Data availability. Data from the Svalbard Airport AWS are freely available through the Norwegian Meteorological Institute's web portal, eKlima (eKlima, 2019). Data from the Gruvefjellet AWS are freely available through the University Centre in Svalbard (UNIS) via <https://www.unis.no/resources/weather-stations/> (UNIS, 2019). All TLS, snow depth, and other auxiliary data can be obtained by contacting the corresponding author by email (holt.hancock@unis.no).

Author contributions. HH was responsible for the majority of the data acquisition, analyses, and interpretation of the results. ME helped develop the conceptual framework for the study and contextualized and interpreted the results within a broader snow and avalanche perspective. AP provided technical guidance with regards to TLS data acquisition and analysis techniques and assisted in the development of the study's technical framework in addition to assisting in data acquisition. JH provided advice and supervision relating to study design, data analysis, and interpretation of the results. HH and ME were responsible for manuscript preparation with input from AP and JH.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We thank Christine Fey and Jeffrey Munroe for their thorough and constructive reviews which greatly improved this work. Andreas Günther is thanked for serving as editor for this work.

Review statement. This paper was edited by Andreas Günther and reviewed by Christine Fey and Jeffrey Munroe.

References

- Abellán, A., Oppikofer, T., Jaboyedoff, M., Rosser, N. J., Lim, M., and Lato, M. J.: Terrestrial laser scanning of rock slope instabilities, *Earth Surf. Proc. Land.*, 39, 80–97, <https://doi.org/10.1002/esp.3493>, 2014.
- American Avalanche Association: Snow, Weather and Avalanches: Observation Guidelines for Avalanche Programs in the United States, 3rd Edn., Victor, ID, 104 pp., 2016.
- Anderton, S. P., White, S. M., and Alvera, B.: Evaluation of spatial variability in snow water equivalent for a high mountain catchment, *Hydrol. Process.*, 18, 435–453, <https://doi.org/10.1002/hyp.1319>, 2004.
- Caputo, T., Marino, E., Matano, F., Somma, R., Troise, C., and De Natale, G.: Terrestrial Laser Scanning (TLS) data for the analysis of coastal tuff cliff retreat: application to Coroglio cliff, Naples, Italy, *Ann. Geophys.*, 61, SE110, <https://doi.org/10.4401/ag-7494>, 2018.
- Christiansen, H. H., Humlum, O., and Eckerstorfer, M.: Central Svalbard 2000–2011 meteorological dynamics and periglacial landscape response, *Arct. Antarct. Alp. Res.*, 45, 6–18, <https://doi.org/10.1657/1938-4246-45.16>, 2013.
- CloudCompare: version 2.10.2 [GPL Software], available at: <http://www.cloudcompare.org>, last access: 17 August 2019.
- Deems, J. S., Painter, T. H., and Finnegan, D. C.: Lidar measurement of snow depth: a review, *J. Glaciol.*, 59, 467–479, <https://doi.org/10.3189/2013JoG12J154>, 2013.
- Deems, J. S., Gadowski, P. J., Vellone, D., Evanczyk, R., LeWinter, A. L., Birkeland, K. W., and Finnegan, D. C.: Mapping starting zone snow depth with a ground-based lidar to assist avalanche control and forecasting, *Cold Reg. Sci. Technol.*, 120, 197–204, <https://doi.org/10.1016/j.coldregions.2015.09.002>, 2015.
- Eckerstorfer, M. and Christiansen, H. H.: The “High Arctic Maritime Snow Climate” in Central Svalbard, *Arct. Antarct. Alp. Res.*, 43, 11–21, <https://doi.org/10.1657/1938-4246-43.1.11>, 2011a.
- Eckerstorfer, M. and Christiansen, H. H.: Topographical and meteorological control on snow avalanching in the Longyearbyen area, central Svalbard 2006–2009, *Geomorphology*, 134, 186–196, <https://doi.org/10.1016/j.geomorph.2011.07.001>, 2011b.
- Eckerstorfer, M., Christiansen, H. H., Rubensdotter, L., and Vogel, S.: The geomorphological effect of cornice fall avalanches in the Longyearfjorden valley, Svalbard, *The Cryosphere*, 7, 1361–1374, <https://doi.org/10.5194/tc-7-1361-2013>, 2013.
- eKlima: Free access to weather- and climate data from the Norwegian Meteorological Institute from historical data to real time observations, The Norwegian Meteorological Institute, available at: http://sharki.oslo.dnmi.no/portal/page?_pageid=73,39035,73_39049&_dad=portal&_schema=PORTAL, last access: 7 August 2018.
- Farizy, B.: Cornice control by helicopter, in: International Snow Science Workshop, 7–11 October 2013, Grenoble – Chamonix Mont-Blanc, 845–847, 2013.
- Fey, C., Schattan, P., Helfricht, K., and Schöber, J.: A compilation of multitemporal TLS snow depth distribution maps at the Weisssee snow research site (Kaunertal, Austria), *Water Resour. Res.*, 55, 5154–5164, <https://doi.org/10.1029/2019wr024788>, 2019.
- Førland, E., Benestad, R., Hanssen-Bauer, I., Haugen, J. E., and Skaugen, T. E.: Temperature and precipitation development at Svalbard 1900–2100, *Adv. Meteorol.*, 2011, 1–14, <https://doi.org/10.1155/2011/893790>, 2011.
- Hanssen-Bauer, I., Solås, M. K., and Steffensen, E.: The Climate of Spitsbergen, Report 39/90, Det Norske Meteorologiske Institutt, Oslo, Norway, 40 pp., 1990.
- Hanssen-Bauer, I., Førland, E., Hisdal, H., Mayer, S., Sandø, A., and Sorteberg, A.: Climate in Svalbard 2100 – a knowledge base for climate adaptation, 1/2019, Norwegian Centre for Climate Services, 207 pp., available at: <https://www.miljodirektoratet.no/globalassets/publikasjoner/M1242/M1242.pdf> (last access: 25 February 2020), 2019.
- Hartzell, P. J., Gadowski, P. J., Glennie, C. L., Finnegan, D. C., and Deems, J. S.: Rigorous error propagation for terrestrial laser scanning with application to snow volume uncertainty, *J. Glaciol.*, 61, 1147–1158, <https://doi.org/10.3189/2015JoG15J031>, 2017.
- Humlum, O., Instanes, A., and Sollid, J. L.: Permafrost in Svalbard: a review of research history, climatic background and engineering challenges, *Polar Res.*, 22, 191–215, 2003.

- Humlum, O., Christiansen, H. H., and Juliussen, H.: Avalanche-derived rock glaciers in Svalbard, *Permafrost Periglac. Process.*, 18, 75–88, 2007.
- Isaksen, K., Nordli, Ø., Førland, E. J., Łupikasza, E., Eastwood, S., and Niedźwiedz, T.: Recent warming on Spitsbergen – Influence of atmospheric circulation and sea ice cover, *J. Geophys. Res.-Atmos.*, 121, 11913–11931, <https://doi.org/10.1002/2016JD025606>, 2016.
- Jaboyedoff, M., Oppikofer, T., Abellán, A., Derron, M.-H., Loye, A., Metzger, R., and Pedrazzini, A.: Use of LIDAR in landslide investigations: a review, *Nat. Hazards*, 61, 5–28, <https://doi.org/10.1007/s11069-010-9634-2>, 2012.
- Jaedicke, C. and Sandvik, A. D.: High resolution snow distribution data from complex Arctic terrain: a tool for model validation, *Nat. Hazards Earth Syst. Sci.*, 2, 147–155, <https://doi.org/10.5194/nhess-2-147-2002>, 2002.
- Kobayashi, D., Ishikawa, N., and Nishio, F.: Formation process and direction distribution of snow cornices, *Cold Reg. Sci. Technol.*, 15, 131–136, [https://doi.org/10.1016/0165-232X\(88\)90059-6](https://doi.org/10.1016/0165-232X(88)90059-6), 1988.
- Lague, D., Brodu, N., and Leroux, J.: Accurate 3D comparison of complex topography with terrestrial laser scanner: Application to the Rangitikei canyon (N-Z), *ISPRS J. Photogram. Remote Sens.*, 82, 10–26, <https://doi.org/10.1016/j.isprsjprs.2013.04.009>, 2013.
- Latham, J. and Montagne, J.: The Possible Importance of Electrical Forces in the Development of Snow Cornices, *J. Glaciol.*, 9, 375–384, <https://doi.org/10.3189/S0022143000022899>, 1970.
- Li, L. and Pomeroy, J. W.: Estimates of Threshold Wind Speeds for Snow Transport Using Meteorological Data, *J. Appl. Meteorol.*, 36, 205–213, [https://doi.org/10.1175/1520-0450\(1997\)036<0205:EOTWSF>2.0.CO;2](https://doi.org/10.1175/1520-0450(1997)036<0205:EOTWSF>2.0.CO;2), 1997.
- Major, H., Haremo, P., Dallmann, W. K., Andresen, A., and Salvigsen, O.: Geological map of Svalbard 1 : 100,000, sheet C9G Adventdalen, Norsk Polarinstitut, Tromsø, Norway, 2001.
- McCarty, D., Brown, R., and Montagne, J.: Cornices: their growth, properties, and control, in: *International Snow Science Workshop, Lake Tahoe, CA, USA*, 41–45, 1986.
- Montagne, J., McPartland, J., Super, A., and Townes, H.: The Nature and control of snow cornices on the Bridger Range, Southwestern Montana, *Miscellaneous Report, Alta Avalanche Study Center, Bozeman, MT*, 1968.
- Mott, R., Schirmer, M., and Lehning, M.: Scaling properties of wind and snow depth distribution in an Alpine catchment, *J. Geophys. Res.-Atmos.*, 116, D06106, <https://doi.org/10.1029/2010JD014886>, 2011.
- Munroe, J. S.: Monitoring snowbank processes and cornice fall avalanches with time-lapse photography, *Cold Reg. Sci. Technol.*, 154, 32–41, <https://doi.org/10.1016/j.coldregions.2018.06.006>, 2018.
- Naruse, R., Nishimura, H., and Maeno, N.: Structural Characteristics of Snow Drifts and Cornices, *Ann. Glaciol.*, 6, 287–288, 1985.
- Prokop, A.: Assessing the applicability of terrestrial laser scanning for spatial snow depth measurements, *Cold Reg. Sci. Technol.*, 54, 155–163, <https://doi.org/10.1016/j.coldregions.2008.07.002>, 2008.
- Prokop, A. and Panholzer, H.: Assessing the capability of terrestrial laser scanning for monitoring slow moving landslides, *Nat. Hazards Earth Syst. Sci.*, 9, 1921–1928, <https://doi.org/10.5194/nhess-9-1921-2009>, 2009.
- Prokop, A. and Procter, E. S.: A new methodology for planning snow drift fences in alpine terrain, *Cold Reg. Sci. Technol.*, 132, 33–43, <https://doi.org/10.1016/j.coldregions.2016.09.010>, 2016.
- Prokop, A., Schirmer, M., Rub, M., Lehning, M., and Stocker, M.: A comparison of measurement methods: terrestrial laser scanning, tachymetry and snow probing for the determination of the spatial snow-depth distribution on slopes, *Ann. Glaciol.*, 49, 210–216, <https://doi.org/10.3189/172756408787814726>, 2008.
- Prokop, A., Schön, P., Singer, F., Pulfer, G., Naaim, M., Thibert, E., and Soruco, A.: Merging terrestrial laser scanning technology with photogrammetric and total station data for the determination of avalanche modeling parameters, *Cold Reg. Sci. Technol.*, 110, 223–230, 2015.
- Prokop, A., Hancock, H., Praz, M., and Jahn, E.: Slope scale avalanche forecasting in the arctic (Svalbard), in: *International Snow Science Workshop, Innsbruck, Austria*, 1035–1039, 2018.
- Riegl® Laser Measurement Systems: VZ-6000 Datasheet, available at: http://www.riegl.com/uploads/tx_pxpriegl/downloads/RIEGL_VZ-6000_Datasheet_2019-05-28_01.pdf, last access: 6 October 2019.
- Rogers, J. C., Yang, L., and Li, L.: The role of Fram Strait winter cyclones on sea ice flux and on Spitsbergen air temperatures, *Geophys. Res. Lett.*, 32, L06709, <https://doi.org/10.1029/2004GL022262>, 2005.
- Schirmer, M., Wirz, V., Clifton, A., and Lehning, M.: Persistence in intra-annual snow depth distribution: 1. Measurements and topographic control, *Water Resour. Res.*, 47, W09516, <https://doi.org/10.1029/2010WR009426>, 2011.
- Schön, P., Prokop, A., Vionnet, V., Guyomarc'h, G., Naaim-Bouvet, F., and Heiser, M.: Improving a terrain-based parameter for the assessment of snow depths with TLS data in the Col du Lac Blanc area, *Cold Reg. Sci. Technol.*, 114, 15–26, <https://doi.org/10.1016/j.coldregions.2015.02.005>, 2015.
- Schweizer, J., Bruce Jamieson, J., and Schneebeil, M.: Snow avalanche formation, *Rev. Geophys.*, 41, 1016, <https://doi.org/10.1029/2002RG000123>, 2003.
- Seligman, G.: *Snow structure and ski fields: being an account of snow and ice forms met with in nature and a study on avalanches & snowcraft*, Macmillan, London, 1936.
- Statham, G., Haegeli, P., Greene, E., Birkeland, K., Israelson, C., Tremper, B., Stethem, C., McMahon, B., White, B., and Kelly, J.: A conceptual model of avalanche hazard, *Nat. Hazards*, 90, 663–691, <https://doi.org/10.1007/s11069-017-3070-5>, 2018.
- UNIS: Weather Station Resources – Gruvefjellet, The University Centre in Svalbard, available at: <https://www.unis.no/resources/weather-stations/>, last access: 7 August 2018.
- van Herwijnen, A. and Fierz, C.: Monitoring snow cornice development using time-lapse photography, in: *Proceedings of the International Snow Science Workshop 2014, Banff, Canada*, 865–869, 2014.
- Vikhamar-Schuler, D., Isaksen, K., Haugen, J. E., Tømmervik, H., Luks, B., Schuler, T. V., and Bjerke, J. W.: Changes in winter warming events in the Nordic Arctic region, *J. Climate*, 29, 6223–6244, <https://doi.org/10.1175/jcli-d-15-0763.1>, 2016.

- Vionnet, V., Martin, E., Masson, V., Guyomarc'H, G., Naaim Bouvet, F., Prokop, A., Durand, Y., and Lac, C.: Simulation of wind-induced snow transport and sublimation in alpine terrain using a fully coupled snowpack/atmosphere model, *The Cryosphere*, 8, 395–415, <https://doi.org/10.5194/tc-8-395-2014>, 2014.
- Vogel, S., Eckerstorfer, M., and Christiansen, H.: Cornice dynamics and meteorological control at Gruvefjellet, Central Svalbard, *The Cryosphere*, 6, 157–171, <https://doi.org/10.5194/tc-6-157-2012>, 2012.
- Watson, C. S., Quincey, D. J., Smith, M. W., Carrivick, J. L., Rowan, A. V., and James, M. R.: Quantifying ice cliff evolution with multi-temporal point clouds on the debris-covered Khumbu Glacier, Nepal, *J. Glaciol.*, 63, 823–837, <https://doi.org/10.1017/jog.2017.47>, 2017.

Paper III

*Synoptic control on snow avalanche activity in central
Spitsbergen*

Synoptic control on snow avalanche activity in central Spitsbergen

Holt Hancock^{1,2}, Jordy Hendriks³, Markus Eckerstorfer^{4,5}, and Siiri Wickström⁶

¹Department of Arctic Geology, University Centre in Svalbard, 9171 Longyearbyen, Norway

²Department of Geosciences, University of Oslo, 0371 Oslo, Norway

³Snow and Avalanche Lab, Department of Earth Sciences, Montana State University, P.O. Box 173480, Bozeman, MT 59717, USA

⁴Regional Climate Lab., Climate Department, NORCE Norwegian Research Centre, 5838 Bergen, Norway

⁵Bjerknes Centre for Climate Research, 5007 Bergen, Norway

⁶Department of Arctic Geophysics, University Centre in Svalbard, 9171 Longyearbyen, Norway

Correspondence: Holt Hancock (holt.hancock@unis.no)

Received: 10 February 2021 – Discussion started: 17 March 2021

Revised: 19 July 2021 – Accepted: 21 July 2021 – Published: 18 August 2021

Abstract. Atmospheric circulation exerts an important control on a region's snow avalanche activity by broadly determining the mountain weather patterns that influence snowpack development and avalanche release. In central Spitsbergen, the largest island in the High Arctic Svalbard archipelago, avalanches are a common natural hazard throughout the winter months. Previous work has identified a unique snow climate reflecting the region's climatically dynamic environmental setting but has not specifically addressed the synoptic-scale control of atmospheric circulation on avalanche activity here. In this work, we investigate atmospheric circulation's control on snow avalanching in the Nordenskiöld Land region of central Spitsbergen by first constructing a four-season (2016/2017–2019/2020) regional avalanche activity record using observations available on a database used by the Norwegian Water Resources and Energy Directorate (NVE). We then analyze the synoptic atmospheric conditions on days with differing avalanche activity situations. Our results show atmospheric circulation conducive to elevated precipitation, wind speeds, and air temperatures near Svalbard are associated with increased avalanche activity in Nordenskiöld Land, but different synoptic signals exist for days characterized by dry, mixed, and wet avalanche activity. Differing upwind conditions help further explain differences in the frequency and nature of avalanche activity resulting from these various atmospheric circulation patterns. We further employ a daily atmospheric circulation calendar to help contextualize our results in the growing body of literature related to climate change in this location. This

work helps expand our understanding of snow avalanches in Svalbard to a broader spatial scale and provides a basis for future work investigating the impacts of climate change on avalanche activity in Svalbard and other locations where avalanche regimes are impacted by changing climatic and synoptic conditions.

1 Introduction

Snow avalanches are natural hazards occurring in mountainous environments where complex interactions between the terrain, weather, and snowpack allow for masses of snow to rapidly descend steep slopes (e.g., Schweizer et al., 2003a). Avalanche forecasts thus seek to reduce risks associated with snow avalanche hazards by predicting avalanche conditions by integrating information related to an area's terrain, past and current meteorological conditions, and snowpack (LaChapelle, 1980; McClung, 2002b). Of the three factors – terrain, weather, and snowpack – contributing to avalanche release and considered in avalanche forecasts, terrain is generally considered static (e.g., Schweizer et al., 2003a), while the snowpack exhibits spatial heterogeneity that is difficult to resolve at broader spatial scales (Schweizer et al., 2008; Clark et al., 2011). Previous work investigating the controls on avalanching at broader, regional to sub-continental spatial scales has therefore focused on understanding the linkages between synoptic-scale atmospheric circulation patterns and periods of increased avalanche activity (e.g., Birkeland et al.,

2001; Fitzharris, 1987; Fitzharris and Bakkehøi, 1986; Keylock, 2003; Martin and Germain, 2017; Schauer et al., 2020).

The synoptic scale in meteorology refers to weather phenomena – such as extratropical cyclones – with horizontal extents on the order of 1000 km (e.g., Yarnal, 1993). Atmospheric processes at this scale are roughly analogous to a mountain range when considering avalanche processes and related operational hazard forecasts (McClung, 2002b). As atmospheric circulation is the primary expression of meteorological conditions at this spatial scale, studying the interactions between atmospheric circulation and the surface environment is known as synoptic climatology (Yarnal, 1993). A synoptic avalanche climatology therefore relates a region's avalanche activity to synoptic-scale atmospheric circulation patterns and can serve as a basis for understanding – and thus forecasting for – avalanche activity at the regional scale. The foundation established by a synoptic avalanche climatology represents the mean atmospheric conditions associated with avalanche activity in a particular location and provides a basis for considering the discrete meteorological conditions resulting from a specific synoptic event which lead to avalanche release.

Numerous studies have addressed avalanche activity at the synoptic scale through analyses of atmospheric circulation patterns. Synoptic avalanche climatologies have been developed for Iceland (Björnsson, 1980), the Norwegian mainland (Fitzharris and Bakkehøi, 1986), western Canada (Fitzharris, 1987), the western United States (Birkeland et al., 2001; Schauer et al., 2020), the Spanish Pyrenees (García et al., 2009), Mt. Shasta in northern California (Hansen and Underwood, 2012), and portions of the northeastern United States (Martin and Germain, 2017). Other works investigating synoptic-scale meteorological controls on regional avalanche activity include analyses of the North Atlantic Oscillation's influence on the avalanche regimes in Iceland (Keylock, 2003) and the Eastern Pyrenees (García-Sellés et al., 2010), as well as the effects of the El Niño–Southern Oscillation on avalanche patterns in western Canada and Chile (McClung, 2013).

In the High Arctic archipelago of Svalbard, atmospheric circulation classifications have been established through subjective (Niedźwiedź, 2013) and objective (Käsmacher and Schneider, 2011) classification methods. Previous work from this region has employed the subjective classification from Niedźwiedź (2013) to relate atmospheric circulation patterns to recent climatic warming across the archipelago (Isaksen et al., 2016), to analyze meteorological conditions and snow distribution on selected glacial systems (Laska et al., 2017; Małeck, 2015), and to characterize wintertime rain-on-snow events (Wickström et al., 2020). Additional research on synoptic-scale meteorological processes in Svalbard includes works examining the effects of atmospheric circulation on air temperatures (e.g., Bednorz and Fortuniak, 2011; Bednorz and Kolendowicz, 2013), precipitation patterns (e.g., Serreze et al., 2015), cloudiness (Bednorz et al.,

2016), and extratropical cyclone activity (Rinke et al., 2017; Rogers et al., 2005; Wickström et al., 2019).

This study investigates the relation between atmospheric circulation patterns and avalanche activity in the Nordenskiöld Land region of central Spitsbergen, the largest island in the Svalbard archipelago (Fig. 1). Spitsbergen is situated at the climatically dynamic interface between the Arctic and North Atlantic regions, where its proximity to poleward oceanic and atmospheric heat transport pathways results in a warmer, wetter climate than expected at these latitudes. The West Spitsbergen Current off the island's western coast – the northernmost extension of the warm Atlantic Gulf Stream – has a moderating effect on regional temperatures and results in perennial ice-free conditions in the eastern Fram Strait (Walczowski and Piechura, 2011). Atmospheric heat and moisture reach Svalbard during winter via extratropical cyclones traveling northeastwards along the North Atlantic storm track, and temperatures can rise far above freezing, even in winter (e.g., Serreze et al., 2015; Vikhamar-Schuler et al., 2016). Spatiotemporal fluctuations in sea ice concentration and extent (e.g., Muckenhuber et al., 2016; Onarheim et al., 2014), atmospheric and oceanic circulation (e.g., Cottier et al., 2007), and storm track location (e.g., Rogers et al., 2005; Wickström et al., 2019) dramatically affect climatic and meteorological conditions in Spitsbergen, making the region one of the most climatically sensitive in the world.

Nordenskiöld Land's High Arctic maritime snow climate reflects central Svalbard's unique environmental setting (Eckerstorfer and Christiansen, 2011a). Winter weather is characterized by prolonged periods of relatively cold, stable high pressure interrupted by warmer, wetter low-pressure systems traveling northwards along the North Atlantic storm track (Hanssen-Bauer et al., 1990; Rogers et al., 2005). Wind slabs and ice layers form in the snowpack during these warm, wet winter storms and are often separated by persistent weak layers of faceted grains developing during colder, stable periods (Eckerstorfer and Christiansen, 2011a). Snow is readily transported by the wind across the region's broad plateau summits and expansive glaciofluvial valleys in the absence of woody vegetation, and large cornices develop annually along the edges of the plateaus (Vogel et al., 2012; Hancock et al., 2020). Avalanche activity in this environment clusters temporally around winter storms, where precipitation and strong winds result in modest snow fall amounts rapidly accumulating in leeward areas (Eckerstorfer and Christiansen, 2011b). Although Nordenskiöld Land's winter climate and thin snowpack promote persistent weak layer development throughout the season (Eckerstorfer and Christiansen, 2011a), the avalanche regime here is nevertheless primarily direct action, with avalanches – even those releasing on deeper instabilities – typically occurring in direct response to snow loading during winter storms (Eckerstorfer and Christiansen, 2011b, c; Hancock et al., 2018).

Dramatic recent changes have been superimposed onto the region's baseline climatic variability (e.g., Hanssen-Bauer

Figure 1. Panel (a) displays Svalbard's location (extent of panel (b) is shown by the red box) in the North Atlantic. Panel (b) presents an overview of Svalbard with the Nordenskiöld Land region, detailed in panel (c), highlighted in green. © OpenStreetMap contributors 2020. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.

et al., 2019), however, and likely portend similarly shifting norms in the snow avalanche climate. Air temperatures over Svalbard have increased on the order of 3–5 °C between 1971 and 2017, with warming particularly pronounced during the winter months (Hanssen-Bauer et al., 2019). Trends in precipitation are generally increasing but do so less clearly (e.g., Førland et al., 2020; Hanssen-Bauer et al., 2019), and increased extreme precipitation event frequency has been observed in recent decades (Hanssen-Bauer et al., 2019; Serreze et al., 2015). These changes have implications for Nordenskiöld Land's snow climate, with observed and projected increases in winter temperatures (Hanssen-Bauer et al., 2019) and rain-on-snow (ROS) event frequency (Peeters et al., 2019; Vikhamar-Schuler et al., 2016; Wickström et al., 2020), combined with shorter snow seasons (e.g., López-Moreno et al., 2016), affecting winter storm characteristics, snowpack duration and composition, and – inferentially, at least – avalanche activity.

The present study builds on this growing body of climatological research to investigate the influence exerted by

synoptic-scale atmospheric circulation on avalanche activity in Nordenskiöld Land. Our specific aims are as follows:

- construct a record of recent avalanche activity using crowd-sourced avalanche observation data from the Norwegian Water Resources and Energy Directorate's (NVE) online snow and avalanche observation registration portal (<https://www.regobs.no>, last access: 7 February 2021);
- investigate the synoptic controls on avalanche activity in Nordenskiöld Land with reference to a well-established subjective atmospheric circulation classification for the Svalbard region;
- place the observed synoptic controls on regional avalanche activity within the larger context of environmental change in the High Arctic.

2 Data and methods

2.1 Avalanche observations and avalanche activity index calculations

A record of avalanche activity spanning four winter seasons (2016/2017–2019/2020) for the Nordenskiöld Land region of central Spitsbergen comprises the primary dataset unique to this work. We used snow and avalanche observations stored in the regObs (an abbreviation for registration of observations) registry and made publicly – and freely – available by the Norwegian Water Resources and Energy Directorate (NVE) at <https://www.regobs.no/> (last access: 7 February 2021). RegObs is NVE's system for acquiring both crowd-sourced and professional observations for natural hazard forecasting in Norway and consists of a mobile app, a website, and a database accessed via an application programming interface (API) (Engeset et al., 2018). Users upload an observation via an interface on either the mobile app or the website that is then immediately displayed and stored for others to view and/or access later via the API. We focus on the snow and avalanche observations registered in regObs for the Nordenskiöld Land region of central Svalbard. These observations form the basis for public avalanche advisories issued by the Norwegian Avalanche Warning Service (NAWS) for Nordenskiöld Land, the only portion of Svalbard for which public avalanche bulletins are regularly issued due to the concentration of human activity near Longyearbyen (Fig. 1). Observation quantity and quality became reliable in Nordenskiöld Land for the 2016/2017 winter season coincident with the initiation of public avalanche forecasts issued by the NAWS and the hiring of trained observers to provide regular snow and avalanche observations via regObs. We accordingly chose this winter season to begin our avalanche activity analyses, with the winter season in this work defined as 1 December–31 May and corresponding to the period for which daily avalanche advisories are issued.

We considered regObs observations classified as avalanche activity, avalanche incidents, avalanche accidents, and danger signs and located within the Nordenskiöld Land forecasting area when constructing our avalanche records (Fig. 2). Avalanche activity observations can report multiple avalanches of similar avalanche types and size for a specified timeframe, single avalanches, or no observed avalanches (“no avalanche activity observed”). These observations typically include a specification of avalanche type and destructive size, a categorical estimate of the number of observed avalanches (0, 1, 2–5, 6–10, > 10), and (ideally) a photo. Avalanche incident observations report single avalanche occurrences and are often used for detailed descriptions of single avalanche events. Avalanche accidents are reported similarly to avalanche incidents but typically involve human activity either through triggering or impact of the avalanche (i.e., danger to infrastructure).

Recent avalanches can also be recorded as a danger sign observation.

We used the API implemented in Python made available by NVE (Egger, 2019) to retrieve all avalanche activity, avalanche incident, and avalanche accident observations in addition to records where recent avalanches were listed as a danger sign from the regObs database for the Nordenskiöld Land forecasting region to create our avalanche activity record. From the API's output including the registration ID, we manually checked all 678 observations on the <https://regobs.no> (last access: 7 February 2021) website to identify individual avalanches. Of these 678 observations, 385 reported avalanches, while 293 were avalanche activity observations with “no avalanches observed.” Manual checks allowed us to discard avalanches registered on multiple occasions or resulting from unreliable observations, correct egregious errors in observed avalanche sizes via visual checks of avalanche photos, and precisely identify the number of individual avalanches included in avalanche activity observations. For the latter, we identified individual avalanches either through visual inspection of photos included with the observation or via the text description of the avalanche activity where individual avalanches may have been mentioned. In situations where neither photos nor the text sufficed to identify individual avalanches, we conservatively estimated the number of avalanches based on the lower limits of the number category such that an observation recording “several” (i.e., 6–10) size 1 avalanches would be recorded as six individual size 1 avalanches. We discarded 110 of the 385 (31 %) records of all observation types reporting avalanches in which the observed avalanches were duplicated in other observations or where we could not determine an avalanche date, size, type, or number either via the avalanche parameters listed by the user or via our own inspection of the observation online. Our final database included 632 individual avalanches recorded in 275 separate avalanche observations (Fig. 2). Avalanches are primarily observed on the date they occur, but in some cases the observer has changed the date based on their knowledge of the avalanche situation or we have changed the date based on our records or the information included in the observation.

We calculated a daily avalanche activity index (AAI) following Schweizer et al. (2003b) using this daily avalanche activity record. The daily AAI represents the total number of all observed avalanches, with each individual avalanche's contribution to the daily sum weighted based on the avalanche's destructive size and trigger type. We assigned typical weights (e.g., Schweizer et al., 2003b) of 0.01, 0.1, and 1 for avalanches of destructive sizes 1, 2, and 3, respectively (we had no destructive 4 or 5 avalanches in our record). We further assigned naturally triggered avalanches a weight of 1 and human-triggered avalanches a weight of 0.5, and we assumed avalanches with an unknown or unspecified trigger to be natural and thus assigned a weight of 1. An example day on which two naturally triggered size 1 avalanches (2

Figure 2. Overview of the regObs observations employed in this work. Panel (a) shows a schematic displaying the regObs observation types retrieved in this work, the parameters we analyzed from each observation type, and the values for each parameter available for user selection. Note that avalanche accidents are not displayed in this figure as they have the same parameters as avalanche incidents and, as of June 2020, are only available for input on the website and not the mobile application. Panel (b) shows the distribution of observations reporting avalanche activity used in this work, while panel (c) highlights observation density near Longyearbyen in the region indicated in panel (b).

avalanches $\times$ 0.01 size weight $\times$ 1 trigger weight = 0.02) and one human-triggered size 3 avalanche (1 avalanches $\times$ 1 size weight $\times$ 0.5 trigger weight = 0.5) occurred would result in a daily AAI of 0.52 (0.02 + 0.5). We further sub-classified wet and dry AAI components such that dry avalanches contributed to the daily dry AAI and wet avalanches contributed to the daily wet AAI, which were then summed to produce the daily AAI. We considered days with only a dry AAI as a “dry” avalanche type day, days with only a wet AAI as “wet”, and days with both dry and wet AAI contributions as “mixed.”

We divided the 166 d in our database with observed avalanche activity into low-activity and high-activity avalanche days based on an AAI threshold value of 0.4 (Fig. 3). We classified the 34 d with an AAI greater than or equal to 0.4 ($0.4 \leq \text{AAI}$) as high-activity days, and we classified the 132 d with an AAI greater than 0 but less than 0.4 ($0 < \text{AAI} < 0.4$) as low-activity days. Knowledge of Svalbard’s avalanche regime (where avalanche activity is generally more limited relative to other locations) and an analysis of the daily AAI distribution (Fig. 3) formed the basis for our use of an AAI value of 0.4 as the threshold to differentiate between low- and high-activity days. Previous

work in Svalbard used a threshold of two size 2 avalanches (equivalent to an AAI of 0.2) to differentiate between a “non-avalanche day” and an “avalanche day” (Eckerstorfer and Christiansen, 2011b). Using a value double the 0.2 threshold used in this previous work, we explored AAI values of both 0.4 and 0.5 as potential thresholds for the differentiation between low-activity days and high-activity days. The 0.5 threshold represents roughly the 86th percentile on our daily AAI distribution, and results in 26 d classified as high-activity days. The 0.4 threshold corresponds to the 83rd percentile of days with avalanche activity, resulting in 34 d classified as high-activity days. We ultimately selected 0.4 as the threshold after detailed analyses indicated the differing threshold values had relatively little impact on the final results, but lowering the threshold to 0.4 increased the number of high-activity days in our analyses, which aided in, for example, more robust composite analyses and significance calculations as described in Sect. 2.2.1. High-activity days above the 0.4 threshold ($0.4 \leq \text{AAI}$) represent 20 % of all days with observed avalanche activity and 5 % of all 729 winter days included in these analyses. The 132 low-activity days ($0 < \text{AAI} < 0.4$) below the 0.4 threshold represent 80 %

Figure 3. Distribution of the calculated daily AAI for days where $AAI > 0$. The dashed black vertical line illustrates the 0.4 AAI threshold above which days were classified as high-activity days ($n = 34$) and below which days were classified as low-activity days ($n = 132$).

of all 166 d with observed avalanche activity and 18 % of all 729 winter days included in these analyses.

2.2 Atmospheric circulation

We employ a calendar of synoptic types known as the Niedźwiedź classification (Niedźwiedź, 2013) to characterize atmospheric circulations patterns in the Svalbard region. The Niedźwiedź classification applies a methodology similar to Lamb's (1972) classification for the British Isles to manually define common synoptic situations over Svalbard. The Niedźwiedź classification assesses the mean sea level pressure (MSLP) field to define common synoptic situations based on the geostrophic wind direction over Svalbard (Niedźwiedź, 2013; Isaksen et al., 2016). We use a version of the Niedźwiedź classification synoptic types denoted by the approximate geostrophic wind direction (N+NE, E+SE, S+SW, W+NW) and whether the situation is dominantly cyclonic (c) or anti-cyclonic (a). A situation with northeasterly geostrophic airflow driven by low-pressure, cyclonic activity would be thus be denoted as N+NEc. Additional cyclonic situations denoted as Cc+Bc indicate a closed low-pressure center (Cc) near Svalbard or a low-pressure trough in the vicinity (Bc), while anti-cyclonic situations with a high-pressure center (Ca) or a ridge (Ka) near Svalbard are denoted as Ca+Ka. Finally, situations where the synoptic situation cannot be explicitly determined are assigned an undefined type (x) such that every day since 1950 is assigned to one of these 11 synoptic types.

We use interpolated MSLP, air temperature (2 m), precipitation, and wind speed data from the ERA5 Reanalysis dataset available at 31 km horizontal resolution (Hersbach

et al., 2020) to analyze meteorological conditions associated with these atmospheric circulation patterns. In combination with the Niedźwiedź classification types, the 1 h ERA5 re-analysis data aggregated to daily mean values (MSLP, temperature, wind speed) and accumulated daily values (precipitation) form the basis for our atmospheric circulation pattern analyses. We apply composite analysis on the MSLP, air temperature, precipitation, and wind speed fields to study the atmospheric conditions associated with each synoptic type based on the ERA5 data (Appendix A). Finally, we use data from two automated weather stations (AWSs), the Svalbard Airport AWS (28 m elevation, 4 km west of Longyearbyen) and the Gruvefjellet AWS (464 m, 2 km southeast of Longyearbyen) to characterize the same parameters in terms of locally observed meteorological conditions (Fig. 1).

2.2.1 Statistical significance

We employed Monte Carlo simulations to evaluate the statistical significance of the anomalies between the mean winter conditions and various composites presented in our results. We considered mean winter conditions to be the overall mean for each parameter taken across all 729 winter (December–May) days between 1 December 2016 and 31 May 2020 (Appendix B). The Monte Carlo simulation process involved selecting a random sample of days the same length as the composite of interest from all 729 winter days and repeating this process 1000 times to generate a synthetic probability distribution following Wickström et al. (2019). In an approach similar to Hendrikx et al. (2009), we determined anomalies to be statistically significant if they fell either below the 5th percentile for negative anomalies or above the 95th percentile for positive anomalies.

We tested significance in the differing distributions of the locally measured meteorological parameters using pairwise a Wilcoxon rank sum test with a Holm p -value adjustment method. We considered distributions to be significantly different for p values less than 0.05.

3 Results

3.1 Avalanche activity

We identified 632 individual avalanches from the regObs observations. Observed avalanches are primarily destructive size 1 and 2 and naturally released (Fig. 4), with no size 4 or 5 avalanches included in our database. Avalanches were observed on 166 of the 729 d included in our four-season record (2016/2017–2019/2020), with 34 of these 166 d further classified as high-activity days (with an $AAI > 0.4$) (Table 1). Just under half (48 %) of all observed avalanches and 63 % of observed size D2 and D3 avalanches occurred on high-activity days. Dry avalanches comprise a majority of observed avalanche activity (75 % of all avalanches), with wet activity dominating only the loose and slush – wet by

definition – avalanche types. Dry slab avalanches are the most observed avalanche type, while only one slushflow (size 3) was observed. High-activity days are primarily dry (27), with mixed (4) and wet (3) high-activity days comprising just 20 % of all high-activity days.

Days without observed avalanches in Nordenskiöld Land are typified at the synoptic scale by low pressure limited to the study domain's southern portions (over northern Fennoscandia), with MSLP conditions near Svalbard differing insignificantly from mean winter conditions (Fig. 5). This indicates the region is situated in the colder, Arctic air mass with the storm track to the south. Weakly negative temperature anomalies over Svalbard extend southwards to Fennoscandia, with negative precipitation and wind speed anomalies focused more directly over Svalbard and the surrounding ocean. While Svalbard is still under mean winter MSLP conditions on low-activity days, high-pressure anomalies dominate the southern study domain. Low-activity days are further characterized by a narrow swath of positive precipitation anomalies extending southwards from Svalbard to the mainland, with localized positive precipitation and wind anomalies concentrated near Spitsbergen's western coast (Fig. 5). The MSLP field during high-activity days is dominated by anomalous low pressure extending from Svalbard to the northeast across the Fram Strait. This circulation pattern is associated with stronger and more widespread positive air temperature, precipitation, and wind speed anomalies over Spitsbergen relative to low-activity days (Fig. 5).

Differences in meteorological parameters between different type of days are broadly reflected in conditions measured locally near Longyearbyen (Fig. 6). Daily air temperatures at the Gruvefjellet AWS are higher on low-activity days and high-activity days than on days without observed avalanches, but no statistical difference exists between air temperatures on low-activity days and high-activity days. Precipitation at the Svalbard Airport AWS is greater on high-activity days than on low-activity days and days without observed activity but does not statistically differ between avalanche days and days without observed activity. Finally, daily averaged wind speeds at the Gruvefjellet AWS are significantly greater on avalanche days than on no observed activity days and greater still on high-activity days.

The MSLP fields differ markedly between low-activity days and high-activity days characterized by dry, mixed, and wet avalanche activity (Fig. 7). Dry low-activity days do not significantly differ from mean MSLP conditions near Svalbard, while significant low-pressure anomalies are centered over Svalbard during dry high-activity days. These circulation patterns are reflected in other meteorological parameters, with dry low-activity days only significantly differing from mean conditions with a limited area of slightly positive wind speed anomalies southeast of Svalbard (Fig. C3). Dry high-activity days show widespread significantly positive precipitation and wind speed anomalies in conjunction with the low-pressure anomaly over the region (Figs. C2

and C3). On wet low-activity days, by contrast, Svalbard is encompassed by strongly positive MSLP anomalies stretching from Fennoscandia across the Barents Sea. This pattern is more pronounced during wet high-activity days, although the region over which the positive anomalies are significant decreases likely due to the limited sample size (Fig. 7). Both wet low-activity and high-activity days exhibit strongly positive air temperature anomalies over Svalbard (Fig. C1), wet high-activity days show a thin strip of significant positive precipitation anomalies extending southwest from central Spitsbergen (Fig. C2), and wind speeds do not significantly differ from mean winter conditions on either wet low- or high-activity days (Fig. C3). MSLP patterns on low- and high-activity days with mixed avalanche activity closely resemble those associated with wet avalanche activity, but the high-pressure ridge over Fennoscandia is somewhat flattened just south of Svalbard and does not extend as dramatically northwards into the Barents Sea (Fig. 7). Mixed low- and high-activity days both show significant positive temperature anomalies (Fig. C1), but other significant anomalies are restricted to a very limited positive precipitation anomaly region in northwestern Spitsbergen and a region of positive wind speed anomalies southeast of the island during mixed high-activity days.

At the AWSs near Longyearbyen, higher air temperatures are observed on mixed and wet low- and high-activity days than on dry low- and high-activity days (Fig. 8). Wind speeds and daily accumulated precipitation, however, do not statistically differ between dry, mixed, and wet low- and high-activity days.

3.2 Connection with established synoptic classifications

We characterize 729 winter days by synoptic type and avalanche activity (Table 1). The most common synoptic types are Types 6 (N+NEc), 7 (E+SEc), and 10 (Cc+Bc). The least frequent synoptic types are Types 4 (W+NWa), 9 (W+NWc), and 3 (S+SWa). Types 7 (E+SEc), 9 (W+NWc), and 6 (N+NEc) result in the most avalanches (Table 1, Fig. 9a), and Types 9 (W+NWc), 4 (W+NWa), and 8 (S+SWc) result in the most avalanches per day (Table 1). Types 9 (W+NWc), 4 (W+NWa), and 10 (Cc+Bc) result in the highest normalized AAI values. Synoptic types dominated by cyclonic airflow – Types 6 (N+NEc), 7 (E+SEc), 8 (S+SWc), 9 (W+NWc), and 10 (Cc+Bc) – occur frequently, result in most observed avalanches, and comprise most high-activity days (Fig. 9, Table 1). Synoptic types dominated by anti-cyclonic airflow – Types 1 (N+NEa), 2 (E+SEa), 3 (S+SWa), 4 (W+NWa), and 5 (Ca+Ka) – occur less frequently, result in fewer observed avalanches, and in some cases do not contain a single high-activity day (Types 2 (E+NEa) and 5 (Ca+Ka)). Dry avalanches dominate observed avalanche activity for all synoptic types except Types 3 (S+SWa) and 9 (W+NWc), during which wet avalanches are more frequently observed (Fig. 9b). Similarly,

Figure 4. Summary of individual avalanches in the regObs data. Panel (a) shows the proportion of observed avalanches triggered naturally and artificially by destructive size, panel (b) shows the proportion of observed avalanches falling into low- and high-activity days by destructive size class, and panel (c) shows the proportion of observed avalanche wetness by avalanche type.

Figure 5. Composite anomaly maps for the meteorological parameters investigated in this study by the daily avalanche activity type classification. Anomalies represent the departure from mean winter conditions (December–May, 2016–2020; see Appendix 1) for the composite mean from daily avalanche activity type and parameter combination. Anomalies that are not statistically significant from mean winter conditions are indicated with dense hatching.

high-activity days are dominated by dry activity, with the exceptions of Types 3 (S+SWa), 9 (W+NWc), 10 (Cc+Bc), and 11 (x – unclassified), which all include mixed or wet high-activity days. Destructive size 3 avalanches – the largest avalanches observed in this study – most frequently occur with Type 10 (Cc+Bc) conditions (seven avalanches),

with Type 9 (W+NWc) and Type 7 (E+SEc) each resulting in four size 3 avalanches. Types 9 (W+NWc) and Types 7 (E+SEc) both result in the highest total number of observed avalanches, but the differing frequencies with which these synoptic types occur contribute to markedly different avalanches per day and normalized AAI values. The follow-

Figure 6. Boxplots showing the distributions for locally measured meteorological parameters by avalanche day type. Air temperature and wind speed are observed at the Gruvefjellet AWS, while precipitation and MSLP are observed at the Svalbard Airport AWS.

ing subsections investigate these synoptic types in more detail.

3.2.1 Synoptic types with a high number of avalanches per day and high normalized AAI values

Low-pressure anomalies situated northeast of Svalbard and high-pressure anomalies over Fennoscandia characterize Type 9 (W+NWc) circulation (Fig. 10). During winter, the resulting west-northwesterly airflow reaches Spitsbergen from across the ice-free Fram Strait. Relative to mean winter conditions, a thin band of positive precipitation anomalies concentrated along Spitsbergen's western coast, positive air temperature anomalies across most of the domain, and positive wind speed anomalies south of Spitsbergen typify Type 9 (W+NWc) days. Type 9 (W+NWc) conditions result in the highest total number of avalanches, equalled only by the number of avalanches occurring on Type 7 (E+SEc) days. Type 9 (W+NWc) days occur much less frequently than Type 7 (E+SEc) days, however, with only Type 4 (W+NWa)

days occurring less frequently. Type 9 (W+NWc) conditions therefore result in the highest number of avalanches per day and the highest normalized AAI of all the synoptic types. Avalanches occurring on Type 9 (W+NWc) days are primarily size 1 (Fig. 9a) and roughly evenly split between dry and wet avalanche types (Fig. 9b). Most of these avalanches occur on high-activity days (Fig. 9c) which can be classified as either dry, mixed, or wet (Fig. 9d). Synoptic conditions on Type 9 (W+NWc) high-activity days differ insignificantly from mean Type 9 (W+NWc) winter conditions (Fig. 10), suggesting mean Type 9 (W+NWc) conditions are sufficient to regularly produce avalanches. Westerly airflow in general appears to be particularly conducive to avalanche activity, as Type 4 (W+NWa) and Type 8 (S+SWc) conditions produce the second and third highest number of avalanches per day, respectively.

Type 10 (Cc+Bc) airflow is characterized by a low-pressure center or trough over Spitsbergen and is not associated with a specific wind direction here (e.g., Appendix A). Calmer conditions are expected with the low centered over Spitsbergen, with northerly flow west and southerly flow east of the island. The low pressure's specific location and depth ultimately determine the resulting wind field. While explicitly westerly airflow types (Types 9, 4, and 8) result in the highest number of avalanches per day, Type 10 (Cc+Bc) conditions result in a greater number of avalanches than either Types 4 (W+NWa) or 8 (S+SWc) and a higher normalized AAI than Type 8 (S+SWc). Type 10 (Cc+Bc) days also result in the highest number of size 3 avalanches of all the synoptic types (Fig. 9a), which contributes to the relatively high normalized AAI value.

3.2.2 Synoptic types with a high total number of avalanches and low normalized AAI values

Type 7 (E+SEc) days result in the same number of avalanches as Type 9 (W+NWc) days. However, the east-southeasterly airflow over Spitsbergen resulting from low pressure situated southwest of Svalbard characteristic of Type 7 (E+SEc) conditions occur much more commonly than Type 9 (W+NWc) conditions. As a result, avalanches per day and the normalized AAI are much lower for Type 7 (E+SEc) conditions (Table 1). Avalanche activity on Type 7 (E+SEc) days is primarily dry (Fig. 9b), and more avalanches greater than size 1 occur on Type 7 (E+SEc) days than on Type 9 (W+NWc) days. Synoptic conditions differ significantly from mean Type 7 (E+SEc) days on high-activity Type 7 (E+SEc) days, with the low-pressure center deepening and shifting to the northeast (Fig. 10). This results in significantly wetter and windier conditions over Spitsbergen than the mean Type 7 (E+SEc) state (Fig. 10). Synoptic conditions on high-activity Type 7 (E+SEc) days thus represent infrequent but significant departures from the mean Type 7 (E+SEc) state. Avalanche activity on Type 7 (E+SEc) high-activity days is dry. Fewer total avalanches (but more avalanches greater than size 1)

Figure 7. Maps of composite MSLP anomalies versus mean winter MSLP conditions (December–May, 2016–2020) by dry, mixed, and wet avalanche activity classifications. Anomalies which are not statistically significantly different from mean winter conditions are indicated with dense hatching. Additional parameter plots are found in Appendix C.

Table 1. Distribution of total winter days and avalanche activity by synoptic type after Niedźwiedź (2013). Maximum values for each column are displayed in bold. Appendix A shows the composite parameter field anomalies for each synoptic type over our period of record.

Synoptic type (numeric identifier)	Synoptic type (airflow descriptor)	Total number of winter days	Proportion of winter days classified as synoptic type	No. of high- activity days	Proportion of days classified as high- activity days	Total number of avalanches	Normalized number of avalanches (avalanches per day)	Normalized AAI (cumulative AAI/ number of days of synoptic type)
1	N+NEa	73	0.10	2	0.03	36	0.49	0.03
2	E+SEa	48	0.07	0	0	14	0.29	0.02
3	S+SWa	26	0.04	2	0.08	24	0.92	0.08
4	W+NWa	11	0.02	1	0.09	23	2.1	0.14
5	Ca+Ka	69	0.09	0	0	16	0.23	0.01
6	N+NEc	174	0.24	6	0.03	106	0.61	0.04
7	E+SEc	120	0.16	5	0.04	107	0.89	0.08
8	S+SWc	59	0.08	3	0.05	80	1.36	0.08
9	W+NWc	17	0.02	4	0.24	107	6.3	0.44
10	Cc+Bc	89	0.12	8	0.09	89	1.0	0.12
11	x	43	0.06	3	0.07	30	0.70	0.07
Total		729	NA	34	0.05	632	0.87	0.07

occur on Type 7 (E+SEc) high-activity days than on Type 9 (W+NWc) high-activity days. These avalanche activity patterns are reflected by the other synoptic type characterized by cyclonically driven, generally easterly airflow – Type 6 (N+NEc). By far the most commonly occurring synoptic type in this study, Type 6 (N+NEc) circulation results in just one fewer avalanche than Types 9 (W+NWc) and 7 (E+SEc). Avalanches on Type 6 (N+NEc) days are almost exclusively dry, and the largest total number of avalanches greater than

size 2 occur under this synoptic type. However, the number of avalanches per day and the normalized AAI for Type 6 (N+NEc) is the lowest of all the cyclonically driven circulation types. Only anti-cyclonic easterly airflow types (Types 1 (N+NEa) and 2 (E+Sea)) and Type 5 (Ca+Ka) (high pressure over Spitsbergen) have lower normalized AAI values than Type 6 (N+NEc).

Figure 8. Boxplots showing the distributions for locally measured meteorological parameters by dry, mixed, and wet avalanche activity classifications. Air temperature and wind speed are observed at the Gruvefjellet AWS, while precipitation is observed at the Svalbard Airport AWS.

4 Discussion

4.1 Synoptic controls on avalanche activity

Our results indicate atmospheric circulation patterns leading to wet, windy, and warm conditions over Nordenskiöld Land are associated with increased observed regional avalanche activity. The conditions are commonly related to cyclonic airflow related to low-pressure systems transported northwards along the North Atlantic storm track. In aggregate, high-activity days display a clear low-pressure signal, with cyclonic activity in the Fram Strait drawing in the warm, wet air represented by positive air temperature and precipitation anomalies (Fig. 5). These controls are further demonstrated by the inverse situation where days without observed avalanche activity show decreased wind speeds, precipitation, and air temperatures over Svalbard relative to mean winter conditions. Cyclonic control is also broadly evident when considering avalanche activity by synoptic type, as synoptic types driven by cyclonic airflow dominate both in terms of total observed avalanches (Fig. 9a) and in the number of high-activity days (Table 1, Fig. 9d). Furthermore, the specific synoptic circulation type characterized by a low-pressure center or trough directly over Svalbard – Type 10

(Cc+Bc) – results in the most high-activity days and the most size 3 avalanches of all synoptic types. The Type 5 (Ca+Ka) synoptic type – high pressure over Svalbard characterizing a drier, calmer synoptic environment – by contrast results in the lowest avalanches per day and normalized AAI values and did not result in a single high-activity day. These results largely confirm previous characterizations of Nordenskiöld Land's direct-action avalanche regime where snowfall, increased wind speeds, and increased air temperatures related to winter storms are precursors to widespread avalanche activity (Eckerstorfer and Christiansen, 2011b, c; Hancock et al., 2018).

While atmospheric circulation conducive to increased precipitation, wind speeds, and air temperatures in the vicinity of Svalbard generally controls avalanche timing here, air-flow direction considerably influences the nature of observed avalanche activity. We found observed wet avalanche activity to be primarily associated with airflow from the southerly through northwesterly airflow synoptic types (e.g., Types 3 (S+SWa), 4 (W+NWa), 8 (S+SWc), and 9 (W+NWc)), while dry activity dominates northerly and easterly airflows (e.g., Types 1 (N+NEa), 6 (N+NEc), and 7 (E+SEc)). These findings align well with previous work employing the Niedźwiedź (2013) classification, showing that synoptic types resulting in southerly and westerly airflow over Spitsbergen result in higher temperatures. They are most frequently associated with precipitation at the Svalbard Airport AWS (Isaksen et al., 2016; Wickström et al., 2020). Synoptic types with an easterly or northerly airflow component were found to be colder and drier than their southerly and westerly counterparts (Førland et al., 2020; Wickström et al., 2020).

The Type 7 (E+SEc) and Type 9 (W+NWc) synoptic types further reflect these synoptic controls. The significant low-pressure anomaly due south of Spitsbergen during Type 7 (E+SEc) high-activity days (relative to Type 7 (E+SEc) mean conditions) promotes strong and wet enough easterly airflow to convey precipitation across eastern Spitsbergen's orographic barrier to Nordenskiöld Land. Sea ice limits upwind sea to air heat exchange in this scenario, and the low pressure's positioning places Spitsbergen in the colder, Arctic air mass with the jet stream to the south. Precipitation thus falls as snow during Type 7 (E+SEc) high-activity days despite a tendency towards positive temperature anomalies relative to mean Type 7 (E+SEc) conditions, and avalanche activity is dry.

By contrast, low pressure to the northeast of Svalbard and high pressure to the south typifies Type 9 (W+NWc) conditions and results in westerly airflow across Nordenskiöld Land. Upwind conditions in this synoptic environment differ considerably from Type 7 (E+SEc) conditions, with warm Atlantic water off Spitsbergen's west coast serving as a heat and moisture source for the westerly, onshore airflow. Nordenskiöld Land is situated on Spitsbergen's windward portion in this scenario (and other synoptic types with westerly airflow), and precipitation is orographically enhanced.

Figure 9. Avalanche activity by synoptic type denoted by the numeric identifier. Panel (a) displays the total number of observed avalanches of each destructive size for each synoptic type. Panel (b) shows the total number of wet and dry avalanches observed during days of each synoptic type. Panel (c) displays the total number of observed avalanches of each destructive size on high-activity days of each synoptic type, and panel (d) displays the number of high-activity days classified by avalanche activity wetness for each synoptic type.

Figure 10. Composite anomalies for all Type 9 and Type 7 days versus mean winter conditions are shown in the first and third columns. Anomalies for Type 9 and Type 7 high-activity days versus the Type 9 and Type 7 composites are shown in the second and fourth columns. High-activity day anomalies that are not statistically significantly different from the respective synoptic type’s composite are indicated with dense hatching.

Type 9 (W+NWc) mean state conditions are therefore more readily conducive to supplying the ingredients for avalanches than mean Type 7 (E+SEc) conditions, as Type 9 (W+NWc) high-activity synoptics do not appear to differ significantly from the Type 9 (W+NWc) mean state (e.g., Fig. 10). The prevalent wet component observed in Type 9 (W+NWc) avalanche activity – suggesting positive air temperatures and liquid precipitation falling in Nordenskiöld Land – reflects the heat available to the air mass from the upwind open ocean. One Type 9 (W+NWc) high-activity day contained only dry avalanches, however, indicating this synoptic situation is not necessarily diagnostic of wet avalanche activity. Westerly airflow is particularly conducive to avalanche activity, as further evidenced by the high normalized AAI and normalized number of avalanches per day from Type 4 (W+NWa) circulation, despite the overall anti-cyclonic dominance of this synoptic type (Table 1).

The composite MSLP fields for dry, mixed, and wet low- and high-activity days (Fig. 7) further reinforce the synoptic controls on avalanching in Nordenskiöld Land. Statistically significant low-pressure anomalies in Svalbard's vicinity characterize the MSLP fields for dry high-activity days, while high pressure over Fennoscandia and the Barents Sea dominates during mixed and wet high-activity days. Low pressure near Svalbard during dry high-activity days again emphasizes the clear cyclonic control on dry avalanche activity in Nordenskiöld Land and represents the most frequently observed synoptic situation leading to avalanches here. This situation provides the snow available for transport and wind speeds necessary to develop dry avalanche conditions. Less frequently, strongly positive MSLP anomalies in the study domain's southern portion lead to mixed and wet avalanche activity days. This finding suggests the Scandinavian blocking pattern, where high pressure over Fennoscandia directs warm and wet air masses to Svalbard, leads to wet avalanches in Nordenskiöld Land. These results encouragingly align with other work that has linked similar synoptic patterns characterized by south-southwesterly airflow over Svalbard with extreme precipitation events in western Spitsbergen (Serreze et al., 2015) and rain-on-snow events throughout the archipelago (Wickström et al., 2020). Such a synoptic environment allows heat and moisture from lower-latitude Atlantic source regions to be efficiently transported to Svalbard, providing the warm air and liquid precipitation required to develop conditions conducive to wet avalanche release. While an association between extreme precipitation and rain-on-snow events with wet avalanche activity is intuitive, it is helpful to specifically place these linked processes within the same synoptic context.

4.2 Implications

4.2.1 Environmental change

Documented increases in air temperature (e.g., Isaksen et al., 2016; Nordli et al., 2020), increasing precipitation and shifting precipitation regimes (e.g., Førland et al., 2020), changing storm tracks (e.g., Wickström et al., 2019), and declining sea ice (e.g., Muckenhuber et al., 2016; Onarheim et al., 2014) have broad implications for Svalbard's physical environment. Previous research on Spitsbergen has investigated the influence of climatic changes on snowpack characteristics in the recent past (e.g., Peeters et al., 2019; van Pelt et al., 2016), but a lack of robust avalanche activity records impedes avalanche-specific analyses over longer temporal scales. Projecting future changes to avalanche regimes in this and other locations is further hindered by the complex interactions between the meteorological and snowpack parameters influencing avalanche release and the difficulty resolving finer-scale snow and avalanche processes with the resolution currently available in climate models (e.g., Beniston et al., 2018). Predictions related to future avalanche activity have therefore primarily focused on broader spatial scales and changes to the overall avalanche regime, with expectations of increasing wet avalanche activity dominating projections in other snow avalanche climates (e.g., Ballesteros-Cánovas et al., 2018; Castebrunet et al., 2014; Dyrørdal et al., 2020). Although we are unable to specifically investigate changes to central Spitsbergen's avalanche regime given our avalanche activity record's considerable limitations, we argue that connecting observed avalanche activity to synoptic atmospheric circulation patterns helps frame avalanche processes within the larger context of environmental and climatic changes. As an example, we link wet avalanches in Nordenskiöld Land to specific synoptic conditions allowing warm air advection to central Svalbard from the south and west. Isaksen et al. (2016), however, noted a correlation between decreasing wintertime sea ice north and east of Svalbard with warmer airflow from these locations, and Wickström et al. (2020) discuss how the ice-free ocean here promotes enhanced precipitation during the early winter season with northerly airflow. With continued background warming superimposed on warmer, moister upwind conditions related to decreased sea ice in the Barents Sea, one might expect to see more frequent and prevalent wet avalanche activity with easterly airflow – airflow that currently results in primarily dry avalanche activity – should air temperatures rise sufficiently. Future research can therefore build upon this work to address questions related to the changing avalanche regime in central Svalbard by, for example, disentangling the relative influence of specific atmospheric circulation patterns, changes to their associated upwind conditions, and general background warming on future wet avalanche activity in this location.

4.2.2 Hazard management

Avalanches present documented hazards to human life and infrastructure in Svalbard, with numerous fatalities occurring in both recreational accidents and infrastructure-related events over the past decade (Engeset et al., 2020; Hancock et al., 2018). Currently, formalized avalanche hazard assessments throughout the winter months in Svalbard include daily regional public avalanche hazard bulletins for Nordenskiöld Land issued by the NAWS and a daily local avalanche hazard bulletin for local officials issued for the infrastructure in Longyearbyen by Skred AS, a consulting firm (Engeset et al., 2020). Our results can help inform avalanche forecasters and hazard managers in this setting by helping resolve avalanche processes at the synoptic-scale that, in combination with more scale-appropriate data for a specific forecasting problem (e.g., slope-scale data for a slope-specific avalanche forecast), form the basis for accurate and precise hazard forecasts (LaChapelle, 1980; McClung, 2002b). For example, we identify westerly, onshore airflow as a synoptic situation particularly conducive to avalanche activity in Nordenskiöld Land. Long-range synoptic-scale weather forecasts can then be used to identify these situations well in advance and refine the hazard assessment with finer spatial- and temporal-scale weather and snowpack data as it becomes available. Our results showing dry high-activity days are related to cyclonic activity (Fig. 5) can be linked to other findings demonstrating more frequent low pressure near Svalbard in recent years. (Wickström et al., 2019). More frequent low pressure may thus result in more frequent dry avalanche activity, which can help hazard managers better prepare for changing frequencies of avalanche events and plan hazard management strategies accordingly.

This work benefited from the availability of the Niedźwiedź classification. The Niedźwiedź classification provided an avenue to easily classify synoptic conditions related to avalanche activity using an existing and well-established classification method for the Svalbard region. Benefits of using the Niedźwiedź classification for this work include (1) the ability to directly link our results to previous work using this classification to investigate regional warming (Isaksen et al., 2016) and rain-on-snow events (Wickström et al., 2020) and (2) the intuitive methodology—where synoptic types are based primarily on the regional wind direction and the cyclonic or anti-cyclonic dominance of the situation—that helps to communicate the results to a broader audience including hazard managers. However, such subjective (or manual) classifications are specific to a single area and are difficult to reproduce (Yarnal, 1993). In locations where a subjective atmospheric circulation pattern classification such as the Niedźwiedź classification employed in this work is unavailable, objective classification techniques offer researchers and hazard managers an avenue to improve the synoptic-scale understanding of avalanche processes. Self-organizing maps, for example, have recently been used to

classify atmospheric circulation patterns leading to deep slab avalanches in the western United States (Schauer et al., 2020). The synoptic types identified using such an objective approach can then be used, as in this study, to identify the synoptic controls on avalanche activity and better inform hazard management decisions.

4.3 Study limitations

This study represents a first-order attempt at understanding the synoptic controls on snow avalanching in central Svalbard, but conclusions drawn from this work are limited by the study's restricted temporal coverage and uncertainties in our avalanche record. Using the regObs database from 2016 onwards gave us access to the best available avalanche data for Nordenskiöld Land, but the resulting four-season record is insufficient for researchers to consider it a true climatological study. This is especially true in Svalbard, where current conditions are not representative of conditions even a decade or two prior (Hanssen-Bauer et al., 2019). Our results are thus best interpreted as a snapshot of the contemporary avalanche setting likely differing both from even recent past and near-future avalanche conditions. We also chose to use mean winter conditions derived from these four winter seasons rather than from across the entire 1979–2020 timeframe for which ERA5 data are available when calculating anomalies. Using composite anomalies calculated from selected days (e.g., no observed activity days, low-activity days, or high-activity days) compared to mean winter conditions from the 1979–2020 period resulted in statistically significant positive air temperature anomalies across the entire domain regardless of the selected day type. We thus elected to consider mean winter conditions as the four-season mean to better differentiate between the primary day types of interest to this work, as our research objectives focused primarily on differentiating the synoptic controls on avalanche activity rather than specifically investigating the impacts of climatic changes across a longer time series.

Considerable uncertainties in the avalanche activity record also limit interpretation of our results. Our avalanche activity record relies on manual, visual avalanche observations from users in the field. Avalanche activity observations thus cluster near Longyearbyen, where most of the human activity occurs (e.g., Fig. 2). The spatial distribution of observations therefore implies our results are most representative for the Longyearbyen area and may be less robust for other, more remote areas in Nordenskiöld Land. Furthermore, the polar night dramatically hinders direct visual avalanche observation during the early snow season, and inclement weather can preclude travel and avalanche observations outside of Longyearbyen at any point during the winter. Following winter storms that prevent travel out of Longyearbyen, avalanches are often reported by observers on the first day when the weather has cleared enough to allow snowmobile travel. In these situations, observers usually either in-

dicating the avalanche occurred prior to the observation date (e.g., the avalanche date is selected to be prior to the observation date) or comment that they believe the avalanche occurred previously. We elected to generally accept the observer's assessment of the avalanche date but have in some cases been forced to discard observations for which we cannot reliably determine an avalanche date. Furthermore, there are very likely days with relatively high avalanche activity that are not captured by this dataset – particularly during the dark season (December–early February) when travel throughout the region and visual observations are both very limited. As a result, avalanche activity recorded on regObs likely represents only a portion of the actual avalanche activity in Nordenskiöld Land during a given period, and days lacking observed activity (“no observed activity days”) do not necessarily indicate that avalanches did not occur in Nordenskiöld Land. Since our analyses include relatively few days with increased avalanche activity, missing even a small number of high-activity days can considerably impact our results. Studies in other locations employing satellite data to detect and quantify avalanche activity have shown manual field observations identify far fewer avalanches than more comprehensive satellite-based approaches (e.g., Bühler et al., 2019; Hafner et al., 2021). More objectively detecting avalanche activity across Nordenskiöld Land through, for example, satellite observations would help improve the robustness of our results and resulting interpretations (e.g., Bühler et al., 2019; Eckerstorfer et al., 2019). Such a record would also help objectively quantify avalanche parameters used in our analyses (e.g., avalanche size, avalanche type, date of avalanche release) currently derived from subjective, judgment-based observations from potentially untrained users. Until satellite products with sufficient temporal resolution and spatial coverage to reliably detect avalanches are available for Svalbard, however, we believe the regObs data are the best alternative for investigating regional avalanche activity in this location.

We differentiated between low and high avalanche activity days in this study based on an AAI threshold of 0.4. This value corresponds to a day with, for example, 40 size 1 avalanches, four size 2 avalanches, or less than one size 3 avalanche (AAI 0.1). Compared to studies in other locations employing AAI values to characterize avalanche activity (e.g., Hägeli and McClung, 2003; Schweizer et al., 2003b), the values in our AAI distribution (Fig. 3) and our selected threshold value are very low. The low AAI values in this work likely stem from two factors. First, manual observations recorded in regObs and used in the AAI calculations likely encompass only a portion of the actual avalanche activity. However, lacking a comprehensive verification dataset (e.g., satellite data), we cannot reliably scale the manual observations to generate a more representative AAI value. Second, our results and past work indicate relatively infrequent avalanche activity, and small avalanches generally characterize central Spitsbergen's avalanche regime (e.g., Eckerstorfer

and Christiansen, 2011b). Thus, we selected an admittedly low threshold to differentiate low- and high-activity days – although we argue this low value is representative of overall low avalanche activity in central Spitsbergen relative to other locations. Finally, although AAI could be used as a continuous variable thus avoiding a threshold selection, we see value in distinguishing between low and high avalanche activity periods, despite limited avalanche activity in our record. Employing a threshold to generate categorical avalanche activity data is furthermore also consistent with the categorical approach avalanche forecasters use to assign, for example, regional avalanche hazard ratings (i.e., a five-tier scale from 1–5).

We have attempted to reduce uncertainties related to observation quality by manually checking each observation as described in Sect. 3.1 and discarding observations for which the quality is too suspect, but uncertainties likely remain. Uncertainty related to avalanche timing, however, has arguably the greatest potential to affect our analyses when attempting to link avalanche release to specific meteorological conditions. While previous work in Svalbard and the results in this work suggest a strong direct-action control on avalanche activity in this location, future work could more specifically target the seasonal progression of avalanche conditions (e.g., the synoptic controls on weak layer development) and should also examine conditions in the 48 and 72 h prior to observed avalanche activity to improve the current analyses. Despite these limitations, this work demonstrates the important role atmospheric circulation plays in the avalanche regime in this unique setting and provides a framework for future analysis on the impacts of climate change on avalanche occurrence.

5 Conclusions

We explored the relationship between atmospheric circulation patterns across a broad portion of the North Atlantic and Arctic regions and observed avalanches in Nordenskiöld Land to investigate the synoptic-scale meteorological controls on avalanche activity in central Spitsbergen. We employed crowd-sourced avalanche occurrence data available on the regObs natural hazard database from four winter seasons (2016/2017–2019/2020) to identify 632 individual avalanches occurring on 166 d, 34 of which we classified as high-activity days. Analyzing the synoptic environment associated with various avalanche activity situations allowed us to build upon previous snow avalanche work in this location by specifically addressing controls on avalanching at a broader spatial scale.

Our findings generally support the existing understanding of snow avalanching in Nordenskiöld Land by showing synoptic conditions leading to increased precipitation, stronger winds, and elevated air temperatures are associated with increased avalanche activity. We identified low pressure near Svalbard resulting in positive precipitation

anomalies and positive wind speed anomalies as the aggregate synoptic signal for dry high-activity days. Warm air diverted northwards by a high-pressure ridge over Fennoscandia and isolated positive precipitation anomalies near Spitsbergen's west coast characterize mixed and wet high-activity days. Further linking our avalanche activity record and analyses to a well-established subjective classification of synoptic situations over Svalbard provided an avenue to consider avalanche activity in Svalbard within the growing body of cryospheric, meteorological, and climatological research for the region. In this classification scheme, circulation patterns characterized by westerly airflow (e.g., Types 9 (W+NWc), 4 (W+NWa)) occur infrequently during the winter, but tend to produce avalanches when they do occur. By contrast, anomalous conditions within common easterly airflow (e.g., Type 7 (E+SEc)) appear to be important to generate high avalanche activity days during these circulation patterns. We therefore argue upwind conditions play a key role in controlling the presence or nature of avalanche activity resulting from a particular circulation pattern. This finding affords future opportunities to link this work to studies of past and future atmospheric and cryospheric change in this location. For example, continued sea ice loss north and east of Svalbard may result in warmer, wetter upwind conditions during periods with generally easterly airflow over Svalbard (e.g., Type 6 (N+NEc) or Type 7 (E+SEc)). Warm, wet conditions and wet avalanche activity currently associated with westerly airflow (e.g., Type 9 (W+NWc) or Type 3 (S+SWa)) may thus begin to occur in the future under the easterly airflow currently associated with dry avalanches. Finally, our results can help hazard managers anticipate future periods of increased avalanche activity based on synoptic-scale weather forecasts. Although limited by a short analysis period and uncertainties in the avalanche record, this work provides a foundation on which to base future studies more specifically investigating the role of Svalbard's rapidly changing climate on avalanche activity here and helps place avalanches within the broader context of environmental change in the Arctic.

Appendix A: Composite parameter field anomalies by synoptic type

Figure A1. Composite daily mean air temperature anomaly by synoptic type, calculated versus mean daily air temperature conditions (Fig. B1). Composite MSLP isobars by synoptic type are included for reference.

Figure A2. Composite daily accumulated precipitation anomaly by synoptic type, calculated versus mean daily accumulated precipitation conditions (Fig. B2). Composite MSLP isobars by synoptic type are included for reference.

Figure A3. Composite daily mean wind speed anomaly by synoptic type, calculated versus mean daily wind speed conditions (Fig. B3). Composite MSLP isobars by synoptic type are included for reference.

Appendix B: Composite mean winter condition fields

Figure B1. Mean daily air temperature composite field for all winter days ($n = 729$) included in these analyses. This represents the mean daily air temperature conditions from which anomalies are calculated. Composite MSLP isobars for all winter days are included for reference.

Figure B2. Mean daily accumulated precipitation composite field for all winter days ($n = 729$) included in these analyses. This represents the mean daily accumulated precipitation conditions from which anomalies are calculated. Composite MSLP isobars for all winter days are included for reference.

Figure B3. Mean daily wind speed field for all winter days ($n = 729$) included in these analyses. This represents the mean daily wind speed conditions from which anomalies are calculated. Composite MSLP isobars for all winter days are included for reference.

Appendix C: Additional composite fields for dry, mixed, and wet avalanche activity

Figure C1. Daily mean air temperature anomalies, relative to mean winter conditions (Fig. B1), for dry, mixed, and wet avalanche low- and high-activity days. Anomalies that are not significantly statistically different from mean winter conditions are indicated with dense hatching.

Figure C2. Daily accumulated precipitation anomalies, relative to mean winter conditions (Fig. B2), for dry, mixed, and wet low- and high-activity days. Anomalies that are not significantly statistically different from mean winter conditions are indicated with dense hatching.

Figure C3. Daily mean wind speed anomalies, relative to mean winter conditions (Fig. B3), for dry, mixed, and wet avalanche low- and high-activity days. Anomalies that are not significantly statistically different from mean winter conditions are indicated with dense hatching.

Data availability. The dataset containing all individual avalanches retrieved from regObs and the calculated daily AAI used in these analyses is available at <https://doi.org/10.17632/dv4m9bbn9y.1>. RegObs data are freely available from NVE on the <http://regobs.no> (last access: 7 February 2021) website. We used the methods contained in <https://github.com/ragnarekker/varsomdata> (last access: 3 June 2021) to access the regObs data. ERA5 data are available through the Copernicus Climate Change Service (CS3) Climate Data Store at <https://cds.climate.copernicus.eu/> (last access: 22 June 2020). Data for the Svalbard Airport AWS are available through Norwegian Meteorological Institute's online data accessibility platform (<https://seklima.met.no/>, last access: 9 June 2020). Data from the Gruvefjellet AWS are freely available through the University Centre in Svalbard via <https://www.unis.no/resourcer/weather-stations/> (last access: 9 June 2020). Tadeusz Niedźwiedź kindly provided us his atmospheric circulation classification calendar upon request.

Author contributions. HH, JH, and ME developed the idea for the study. HH processed and analyzed the data with assistance from SW. HH prepared the manuscript with contributions from all co-authors.

Competing interests. The authors declare they have no conflict of interest.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. We are grateful for Tadeusz Niedźwiedź's contribution of his atmospheric circulation calendar, without which this work would not have been possible in its current form. Constructive comments from Karsten Müller and an anonymous referee and editorial assistance from Masashi Niwano considerably improved the manuscript. We are also thankful for NVE's work to allow user access to the considerable wealth of information contained in the regObs database.

Review statement. This paper was edited by Masashi Niwano and reviewed by Karsten Müller and one anonymous referee.

References

- Ballesteros-Cánovas, J. A., Trappmann, D., Madrigal-González, J., Eckert, N., and Stoffel, M.: Climate warming enhances snow avalanche risk in the Western Himalayas, *P. Natl. Acad. Sci. USA*, 115, 3410–3415, <https://doi.org/10.1073/pnas.1716913115>, 2018.
- Bednorz, E. and Fortuniak, K.: The occurrence of coreless winters in central Spitsbergen and their synoptic conditions, *Polar Res.*, 30, 12218, <https://doi.org/10.3402/polar.v30i0.12218>, 2011.
- Bednorz, E. and Kolendowicz, L.: Summer mean daily air temperature extremes in Central Spitsbergen, *Theor. Appl. Climatol.*, 113, 471–479, 2013.
- Bednorz, E., Kaczmarek, D., and Dudlik, P.: Atmospheric conditions governing anomalies of the summer and winter cloudiness in Spitsbergen, *Theor. Appl. Climatol.*, 123, 1–10, <https://doi.org/10.1007/s00704-014-1326-5>, 2016.
- Beniston, M., Farinotti, D., Stoffel, M., Andreassen, L. M., Coppola, E., Eckert, N., Fantini, A., Giacona, F., Hauck, C., Huss, M., Huwald, H., Lehning, M., López-Moreno, J.-I., Magnusson, J., Marty, C., Morán-Tejeda, E., Morin, S., Naaim, M., Provenzale, A., Rabatel, A., Six, D., Stötter, J., Strasser, U., Terzago, S., and Vincent, C.: The European mountain cryosphere: a review of its current state, trends, and future challenges, *The Cryosphere*, 12, 759–794, <https://doi.org/10.5194/tc-12-759-2018>, 2018.
- Birkeland, K. W., Mock, C., and Shinker, J.: Avalanche extremes and atmospheric circulation patterns, *Ann. Glaciol.*, 32, 135–140, 2001.
- Björnsson, H.: Avalanche activity in Iceland, climatic conditions, and terrain features, *J. Glaciol.*, 26, 13–23, 1980.
- Bühler, Y., Hafner, E. D., Zweifel, B., Zesiger, M., and Heisig, H.: Where are the avalanches? Rapid SPOT6 satellite data acquisition to map an extreme avalanche period over the Swiss Alps, *The Cryosphere*, 13, 3225–3238, <https://doi.org/10.5194/tc-13-3225-2019>, 2019.
- Castebrunet, H., Eckert, N., Giraud, G., Durand, Y., and Morin, S.: Projected changes of snow conditions and avalanche activity in a warming climate: the French Alps over the 2020–2050 and 2070–2100 periods, *The Cryosphere*, 8, 1673–1697, <https://doi.org/10.5194/tc-8-1673-2014>, 2014.
- Clark, M. P., Hendrikx, J., Slater, A. G., Kavetski, D., Anderson, B., Cullen, N. J., Kerr, T., Örn Hreinsson, E., and Woods, R. A.: Representing spatial variability of snow water equivalent in hydrologic and land-surface models: A review, *Water Resour. Res.*, 47, W07539, <https://doi.org/10.1029/2011WR010745>, 2011.
- Cottier, F. R., Nilsen, F., Inall, M. E., Gerland, S., Tverberg, V., and Svendsen, H.: Wintertime warming of an Arctic shelf in response to large-scale atmospheric circulation, *Geophys. Res. Lett.*, 34, L10607, <https://doi.org/10.1029/2007GL029948>, 2007.
- Dyrddal, A. V., Isaksen, K., Jacobsen, J. K. S., and Nilsen, I. B.: Present and future changes in winter climate indices relevant for access disruptions in Troms, northern Norway, *Nat. Hazards Earth Syst. Sci.*, 20, 1847–1865, <https://doi.org/10.5194/nhess-20-1847-2020>, 2020.
- Eckerstorfer, M. and Christiansen, H. H.: The “High Arctic Maritime Snow Climate” in Central Svalbard, *Arct. Antarct. Alp. Res.*, 43, 11–21, <https://doi.org/10.1657/1938-4246-43.1.11>, 2011a.
- Eckerstorfer, M. and Christiansen, H. H.: Relating meteorological variables to the natural slab avalanche regime in High Arctic Svalbard, *Cold Reg. Sci. Technol.*, 69, 184–193, <https://doi.org/10.1016/j.coldregions.2011.08.008>, 2011b.
- Eckerstorfer, M. and Christiansen, H. H.: Topographical and meteorological control on snow avalanching in the Longyearbyen area, central Svalbard 2006–2009, *Geomorphology*, 134, 186–196, <https://doi.org/10.1016/j.geomorph.2011.07.001>, 2011c.
- Eckerstorfer, M., Vickers, H., Malnes, E., and Grahm, J.: Near-Real Time Automatic Snow Avalanche Activity Monitoring System

- Using Sentinel-1 SAR Data in Norway, *Remote Sens.*, 11, 2863, 2019.
- Ekker, R.: varsomdata: GitHub, <https://github.com/ragnarekker/varsomdata>, last access: 6 June 2020, 2019.
- Engeset, R. V., Ekker, R., Humstad, T., and Landrø, M.: Varsom:regobs – A common real-time picture of the hazard situation shared by mobile information technology, International Snow Science Workshop, Innsbruck, Austria, 7–12 October 2018, 1573–1577, 2018.
- Engeset, R. V., Landrø, M., Indreiten, M., Müller, K., Mikkelsen, O.-A., and Hoseth, K. I. A.: Avalanche warning in Svalbard, Norwegian Water Resources and Energy Directorate, Oslo, Norway, 25 pp., 2020.
- Fitzharris, B.: A climatology of major avalanche winters in western Canada, *Atmos. Ocean*, 25, 115–136, 1987.
- Fitzharris, B. and Bakkehøi, S.: A synoptic climatology of major avalanche winters in Norway, *J. Climatol.*, 6, 431–446, 1986.
- Førland, E. J., Isaksen, K., Lutz, J., Hanssen-Bauer, I., Schuler, T. V., Dobler, A., Gjeltén, H. M., and Vikhamar-Schuler, D.: Measured and Modeled Historical Precipitation Trends for Svalbard, *J. Hydrometeorol.*, 21, 1279–1296, <https://doi.org/10.1175/jhm-d-19-0252.1>, 2020.
- García, C., Martí, G., Oller, P., Moner, I., Gavalda, J., Martínez, P., and Peña, J. C.: Major avalanches occurrence at regional scale and related atmospheric circulation patterns in the Eastern Pyrenees, *Cold Reg. Sci. Technol.*, 59, 106–118, <https://doi.org/10.1016/j.coldregions.2009.07.009>, 2009.
- García-Sellés, C., Peña, J. C., Martí, G., Oller, P., and Martínez, P.: WeMOI and NAOi influence on major avalanche activity in the Eastern Pyrenees, *Cold Reg. Sci. Technol.*, 64, 137–145, <https://doi.org/10.1016/j.coldregions.2010.08.003>, 2010.
- Hafner, E. D., Techel, F., Leinss, S., and Bühler, Y.: Mapping avalanches with satellites – evaluation of performance and completeness, *The Cryosphere*, 15, 983–1004, <https://doi.org/10.5194/tc-15-983-2021>, 2021.
- Hägeli, P. and McClung, D. M.: Avalanche characteristics of a transitional snow climate – Columbia Mountains, British Columbia, Canada, *Cold Reg. Sci. Technol.*, 37, 255–276, [https://doi.org/10.1016/S0165-232X\(03\)00069-7](https://doi.org/10.1016/S0165-232X(03)00069-7), 2003.
- Hancock, H., Prokop, A., Eckerstorfer, M., and Hendriks, J.: Combining high spatial resolution snow mapping and meteorological analyses to improve forecasting of destructive avalanches in Longyearbyen, Svalbard, *Cold Reg. Sci. Technol.*, 154, 120–132, <https://doi.org/10.1016/j.coldregions.2018.05.011>, 2018.
- Hancock, H., Eckerstorfer, M., Prokop, A., and Hendriks, J.: Quantifying seasonal cornice dynamics using a terrestrial laser scanner in Svalbard, Norway, *Nat. Hazards Earth Syst. Sci.*, 20, 603–623, <https://doi.org/10.5194/nhess-20-603-2020>, 2020.
- Hansen, C. and Underwood, S. J.: Synoptic Scale Weather Patterns and Size-5 Avalanches on Mt. Shasta, California, *Northwest Sci.*, 86, 329–341, 2012.
- Hanssen-Bauer, I., Solås, M. K., and Steffensen, E.: The Climate of Spitsbergen, Det Norske Meteorologiske Institutt, 40 pp., 1990.
- Hanssen-Bauer, I., Førland, E., Hisdal, H., Mayer, S., Sandø, A., and Sorteberg, A.: Climate in Svalbard 2100 – a knowledge base for climate adaptation, Norwegian Centre for Climate Services 1/2019, 207, 2019.
- Hendriks, J., Birkeland, K., and Clark, M.: Assessing changes in the spatial variability of the snowpack fracture propagation propensity over time, *Cold Reg. Sci. Technol.*, 56, 152–160, <https://doi.org/10.1016/j.coldregions.2008.12.001>, 2009.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. Roy. Meteor. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Isaksen, K., Nordli, Ø., Førland, E. J., Łupikasza, E., Eastwood, S., and Niedźwiedz, T.: Recent warming on Spitsbergen – Influence of atmospheric circulation and sea ice cover, *J. Geophys. Res.-Atmos.*, 121, 11913–11931, <https://doi.org/10.1002/2016JD025606>, 2016.
- Käsmacher, O. and Schneider, C.: An objective circulation pattern classification for the region of Svalbard, *Geogr. Ann. A*, 93, 259–271, <https://doi.org/10.1111/j.1468-0459.2011.00431.x>, 2011.
- Keylock, C. J.: The North Atlantic Oscillation and snow avalanching in Iceland, *Geophys. Res. Lett.*, 30, 1254, <https://doi.org/10.1029/2002GL016272>, 2003.
- LaChapelle, E. R.: The fundamental processes in conventional avalanche forecasting, *J. Glaciol.*, 26, 75–84, 1980.
- Lamb, H. H.: British Isles weather types and a register of the daily sequence of circulation patterns, 1861–1971, *Geophysical Memoirs: HMSO, London*, 116, p. 85, 1972.
- Laska, M., Grabiec, M., Ignatiuk, D., and Budzik, T.: Snow deposition patterns on southern Spitsbergen glaciers, Svalbard, in relation to recent meteorological conditions and local topography, *Geogr. Ann. A*, 99, 262–287, <https://doi.org/10.1080/04353676.2017.1327321>, 2017.
- López-Moreno, J. I., Boike, J., Sanchez-Lorenzo, A., and Pomeroy, J. W.: Impact of climate warming on snow processes in Ny-Ålesund, a polar maritime site at Svalbard, *Global Planet. Change*, 146, 10–21, <https://doi.org/10.1016/j.gloplacha.2016.09.006>, 2016.
- Małeck, J.: Glacio- meteorology of Ebbabreen, Dickson Land, central Svalbard, during 2008–2010 melt seasons, *Pol. Polar Res.*, 36, 145–161, <https://doi.org/10.1515/popore-2015-0010>, 2015.
- Martin, J.-P. and Germain, D.: Large-scale teleconnection patterns and synoptic climatology of major snow-avalanche winters in the Presidential Range (New Hampshire, USA), *Int. J. Climatol.*, 37, 109–123, <https://doi.org/10.1002/joc.4985>, 2017.
- McClung, D.: The Elements of Applied Avalanche Forecasting, Part II: The Physical Issues and the Rules of Applied Avalanche Forecasting, *Nat. Hazards*, 26, 131–146, <https://doi.org/10.1023/a:1015604600361>, 2002.
- McClung, D. M.: The effects of El Niño and La Niña on snow and avalanche patterns in British Columbia, Canada, and central Chile, *J. Glaciol.*, 59, 783–792, <https://doi.org/10.3189/2013JoG12J192>, 2013.
- Muckenhuber, S., Nilsen, F., Korosov, A., and Sandven, S.: Sea ice cover in Isfjorden and Hornsund, Svalbard (2000–2014) from remote sensing data, *The Cryosphere*, 10, 149–158, <https://doi.org/10.5194/tc-10-149-2016>, 2016.

- Niedźwiedź, T.: The atmospheric circulation, in: *Climate and Climate Change at Hornsund, Svalbard*, Marsz, A. A. and Styszyńska, A. (Eds.), Gdynia Maritime University, Gdynia, Poland, 57–74, 2013.
- Nordli, Ø., Wyszyński, P., Gjeltén, H. M., Isaksen, K., Łupikasza, E., Niedźwiedź, T., and Przybylak, R.: Revisiting the extended Svalbard Airport monthly temperature series, and the compiled corresponding daily series 1898–2018, *Polar Res.*, 39, <https://doi.org/10.33265/polar.v39.3614>, 2020.
- Onarheim, I. H., Smedsrud, L. H., Ingvaldsen, R. B., and Nilsen, F.: Loss of sea ice during winter north of Svalbard, *Tellus A*, 66, 23933, <https://doi.org/10.3402/tellusa.v66.23933>, 2014.
- Peeters, B., Pedersen, Å. Ø., Loe, L. E., Isaksen, K., Veiberg, V., Stien, A., Kohler, J., Gallet, J.-C., Aanes, R., and Hansen, B. B.: Spatiotemporal patterns of rain-on-snow and basal ice in high Arctic Svalbard: detection of a climate-cryosphere regime shift, *Environ. Res. Lett.*, 14, 015002, <https://doi.org/10.1088/1748-9326/aaefb3>, 2019.
- Rinke, A., Maturilli, M., Graham, R. M., Matthes, H., Handorf, D., Cohen, L., Hudson, S. R., and Moore, J. C.: Extreme cyclone events in the Arctic: Wintertime variability and trends, *Environ. Res. Lett.*, 12, 094006, <https://doi.org/10.1088/1748-9326/aa7def>, 2017.
- Rogers, J. C., Yang, L., and Li, L.: The role of Fram Strait winter cyclones on sea ice flux and on Spitsbergen air temperatures, *Geophys. Res. Lett.*, 32, <https://doi.org/10.1029/2004GL022262>, 2005.
- Schauer, A. R., Hendrikx, J., Birkeland, K. W., and Mock, C. J.: Synoptic atmospheric circulation patterns associated with deep persistent slab avalanches in the western United States, *Nat. Hazards Earth Syst. Sci.*, 21, 757–774, <https://doi.org/10.5194/nhess-21-757-2021>, 2021.
- Schweizer, J., Bruce Jamieson, J., and Schneebeli, M.: Snow avalanche formation, *Rev. Geophys.*, 41, L06709, <https://doi.org/10.1029/2002RG000123>, 2003a.
- Schweizer, J., Kronholm, K., and Wiesinger, T.: Verification of regional snowpack stability and avalanche danger, *Cold Reg. Sci. Technol.*, 37, 277–288, [https://doi.org/10.1016/S0165-232X\(03\)00070-3](https://doi.org/10.1016/S0165-232X(03)00070-3), 2003b.
- Schweizer, J., Kronholm, K., Jamieson, J. B., and Birkeland, K. W.: Review of spatial variability of snowpack properties and its importance for avalanche formation, *Cold Reg. Sci. Technol.*, 51, 253–272, <https://doi.org/10.1016/j.coldregions.2007.04.009>, 2008.
- Serreze, M. C., Crawford, A. D., and Barrett, A. P.: Extreme daily precipitation events at Spitsbergen, an Arctic Island, *Int. J. Climatol.*, 35, 4574–4588, <https://doi.org/10.1002/joc.4308>, 2015.
- van Pelt, W. J. J., Kohler, J., Liston, G. E., Hagen, J. O., Luks, B., Reijmer, C. H., and Pohjola, V. A.: Multidecadal climate and seasonal snow conditions in Svalbard, *J. Geophys. Res.-Earth*, 121, 2100–2117, <https://doi.org/10.1002/2016jf003999>, 2016.
- Vikhamar-Schuler, D., Isaksen, K., Haugen, J. E., Tømmervik, H., Luks, B., Schuler, T. V., and Bjerke, J. W.: Changes in winter warming events in the Nordic Arctic region, *J. Climate*, 29, 6223–6244, <https://doi.org/10.1175/jcli-d-15-0763.1>, 2016.
- Vogel, S., Eckerstorfer, M., and Christiansen, H. H.: Cornice dynamics and meteorological control at Gruvefjellet, Central Svalbard, *The Cryosphere*, 6, 157–171, <https://doi.org/10.5194/tc-6-157-2012>, 2012.
- Walczowski, W. and Piechura, J.: Influence of the West Spitsbergen Current on the local climate, *Int. J. Climatol.*, 31, 1088–1093, <https://doi.org/10.1002/joc.2338>, 2011.
- Wickström, S., Jonassen, M. O., Vihma, T., and Uotila, P.: Trends in cyclones in the high-latitude North Atlantic during 1979–2016, *Q. J. Roy. Meteor. Soc.*, 146, 762–779, <https://doi.org/10.1002/qj.3707>, 2019.
- Wickström, S., Jonassen, M. O., Cassano, J. J., and Vihma, T.: Present Temperature, Precipitation and Rain-on-Snow Climate in Svalbard, *J. Geophys. Res.-Atmos.*, 125, e2019JD032155, <https://doi.org/10.1029/2019jd032155>, 2020.
- Yarnal, B.: *Synoptic Climatology in Environmental Analysis: a Primer*, Studies in Climatology Series, edited by: Gregory, S., Bellhaven Press, London, UK, 1993.