Symptomatic lumbar disc herniation treated in hospitals

Prognosis of unfavourable surgical events and persistent leg-pain

Doctoral thesis by
Olaf R. Fjeld

2020

Institute of Clinical Medicine, Faculty of Medicine, University of Oslo

Research and Communication Unit for Musculoskeletal Health (FORMI), Department of
Research and Innovation, Division of Clinical Neuroscience, Oslo University Hospital

© Olaf R. Fjeld, 2021

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-884-7

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

CONTENTS

ACKNOWLEDGEMENTS.....	3
LIST OF PAPERS.....	4
ABBREVIATIONS	5
SUMMARY IN ENGLISH	6
SAMMENDRAG PÅ NORSK.....	8
PREFACE	10
1. INTRODUCTION	11
1.1 EPIDEMIOLOGY.....	11
1.2 PATHOMECHANISMS AND CLINICAL FEATURES.....	11
1.3 TREATMENT	13
1.4 PROGNOSIS AND PROGNOSTIC FACTOR RESEARCH.....	15
1.4.1 Unfavourable event as an outcome in lumbar disc surgery.....	15
1.4.2 Self-reported outcome measures in sciatica	16
1.4.3 Prognostic factors for poor outcome in sciatica	17
1.4.4 Conditioned pain modulation in sciatica: a potential prognostic factor.....	17
2. AIMS OF THE STUDY	21
3. MATERIALS AND METHODS	22
3.1 STUDY DESIGNS	22
3.2 STUDY POPULATIONS	24
3.3 MEASUREMENTS	24
3.3.1 Outcome measures	24
3.3.2 The potential prognostic factors considered.....	26
3.4 STATISTICAL ANALYSIS	28
3.4.1 A general overview of statistical methods and hardware.....	28
3.4.2 A description and rationale for the statistical methods	29
3.4.3 Sample Size and missing values.....	29
3.5 ETHICS.....	30
4. SUMMARY OF THE RESULTS.....	31
5. DISCUSSION	33
5.1 METHODOLOGICAL CONSIDERATIONS	33
5.1.1 Strengths and disadvantages of study designs.....	33
5.1.2 Internal validity	35
5.1.2.1 Random associations.....	35
5.1.2.2 Selection bias	35
5.1.2.3 Information bias (misclassification)	36
5.1.2.4 Confounding	37
5.1.2.5 Errors specific to regression analysis.....	38
5.1.3 External validity	40
5.2 MAIN RESULTS COMPARED WITH OTHER PUBLISHED STUDIES	42
5.2.1 Unfavourable events in lumbar disc surgery	42
5.2.2 Persistent leg-pain	44
5.3 CLINICAL IMPLICATIONS	45
5.4 FUTURE PERSPECTIVES	46
6. CONCLUSIONS	49
REFERENCES	50
PAPERS I-III.....	60

ACKNOWLEDGEMENTS

The work presented in this thesis was financially supported by the South-Eastern Norway Regional Health Authority grant number 2013030. The work was performed at the Research and Communication Unit for Musculoskeletal Health (FORMI), Department of Research and Innovation, and the Department of Neurology, Division of Clinical Neuroscience, Oslo University Hospital, and Institute of Clinical Medicine, Faculty of Medicine, University of Oslo. Many a thanks goes to these institutions.

First of all, I want to thank my main supervisor Margreth Grotle for including me in her projects and introducing me to her world of research and fellow researchers. She has generously shared her comprehensive knowledge and great interest in the field of back pain research. Her attention to detail and commitment has been invaluable. Equally, a special thanks goes to my second supervisor John-Anker Zwart for his enthusiasm and immense optimism, always offering his support and insight in team meetings and paper discussions, but also in the hallway and in the office kitchen. I would also especially like to thank Kristian Bernhard Nilsen for all the hours spent teaching me about CPM testing and pain research.

I want to give a big thanks to Kjersti Storheim, Mathias Toft and Hanne Harbo for making it possible for me to work as a researcher and at the same time making room for me to work as a clinician.

I am sincerely grateful to my co-authors who have contributed in various ways: Audun Stubhaug, Ingrid Heuch, Jon Helgeland, Lars Grøvle, Linda Pedersen, Marie Udnesseter Lie, Milada Cvancarova Småstuen, Tore Kristian Solberg and Vibeke Siewers.

To the patients and all you guys at FORMI and at the Neurological department, let's keep up the good work! I absolutely love coming to work because of you.

Finally, thanks to my mother and father that nurtured my interest in research and writing throughout my life. And, above all, I would like to thank Ingrid for sharing her life with me.

LIST OF PAPERS

The following original articles are referred to in the text by their Roman numerals:

Paper I

Complications, reoperations, readmissions, and length of hospital stay in 34 639 surgical cases of lumbar disc herniation.

Fjeld O.R., Grøvle L., Helgeland J., Småstuen M.C., Solberg T.K., Zwart J.A., Grotle M.
Bone Joint J. 2019;101(4):470-477.

Paper II

Prognostic Factors for Persistent Leg-Pain in Patients Hospitalized With Acute Sciatica.

Fjeld O., Grotle M., Siewers V., Pedersen L.M., Nilsen K.B., Zwart J.A.
Spine (Phila Pa 1976) 2017;42(5):272-279.

Paper III

Predicting the Outcome in Persistent Sciatica Using Conditioned Pain Modulation: 1-year results from a prospective cohort study

Fjeld O., Grotle M., Matre D., Pedersen L.M., Lie M.U., Småstuen M.C., Storheim K., Heuch I., Stubhaug A., Zwart J.A., Nilsen K.B.
Scand. J. Pain. 2019;20(1):69-75.

ABBREVIATIONS

BMI	Body Mass Index
CCI	Charlson Comorbidity Index
CI	Confidence Interval
CPM	Conditioned Pain Modulation
ICD-10	International Classification Diseases number 10
MRI	Magnetic Resonance Imaging
NRS	Numeric Rating Scale
PROMS	Patient-reported Outcome Measures
ODI	Oswestry Disability Index
OR	Odds Ratio
REC	Regional Ethics Committee
SD	Standard Deviation
VAS	Visual Analogue Scale
ÖMSPQ	Örebro Musculoskeletal Pain Questionnaire

SUMMARY IN ENGLISH

Background

Low-back pain is the single leading cause for disability worldwide causing activity limitation and work absence with subsequently enormous economic burden on individuals, families, communities, industry, health services and governments. In about 5-10% low-back pain is accompanied by nerve root impingement causing radiating pain into one or both legs, described as radiculopathy or sciatica. Patients with severe symptoms or refractory radiating pain are typically offered surgery, making lumbar disc surgery a common procedure worldwide. Thus, it is important that patients are informed of the different types and rates of unfavorable events that may occur during and after surgery.

Although the prognosis is generally favorable, some patients unfortunately go on to develop persistent sciatic pain regardless of treatment. It is believed that a combination of biological and psychosocial properties among the patients is responsible for the transition from acute to chronic pain (the biopsychosocial model). However, at present, it is difficult to predict which patients will go on to develop chronic symptoms and which will not. Prognostic factor research enables us to better understand who these patients are and provide insight into what mechanisms are responsible for unsuccessful healing.

Aims/Objectives

The overall aim of the presented work was to assess the frequency of unfavorable surgical events after lumbar disc herniation surgery, and in addition to study potential prognostic factors for the persistence of leg-pain in surgical and non-surgical patients hospitalized with sciatica.

This was sought through an epidemiological analysis of two large national databases complemented by two in-depth prospective cohort studies.

The aim of Paper I presented in this thesis was to elicit the rates for unfavorable events such as complications, reoperations and readmissions in lumbar disc surgery.

Paper II was aimed at identifying sociodemographic, lifestyle, psychosocial and clinical prognostic factors for persistent leg-pain at 12 months in both surgically and non-surgically treated patients hospitalized with acute severe sciatica.

The aim of Paper III was to test whether Conditioned Pain Modulation (CPM) effect measured 6 weeks after hospital discharge could be associated with self-reported leg-pain at 12 months in a cohort consisting of both surgically and non-surgically treated sciatica patients.

Patients and methods

Paper I was a longitudinal observation study. Data were retrieved using a combination of procedure and diagnosis codes from two large medical administrative registries covering all lumbar disc surgeries performed in the public sector in Norway in the years 1999-2013. The impact of age, gender, geographical affiliation, education, civil status, income, and comorbidity on unfavorable surgical events were analyzed by logistic regression.

Paper II was based on data from a cohort study with a total of 210 patients acutely admitted to hospital for either surgical or nonsurgical treatment of sciatica. The patients were consecutively recruited upon hospital admission and received follow-up assessments at 6 weeks, 6 months, and 12 months post hospital discharge. Potential prognostic factors were measured at hospital admission and at 6 weeks. The impact of these factors on leg-pain measured at 12 months was analyzed by multiple linear regression modeling.

Paper III was a prospective cohort study in which CPM effect was measured in 111 patients who continued to suffer from sciatica 6 weeks after being discharged from hospital. The patients received a follow up assessment of leg-pain 12 months after hospital discharge. The impact of CPM effect on self-reported leg-pain at 12 months was analyzed using logistic regression.

Results

In 34 639 lumbar disc surgeries, 2.7% had a *surgical complication*, 2.1% had *repeat surgery* within 90 days, 2.4% had a *non-surgical readmission* within 90 days and 6.7% experienced at least one of these unfavorable events (in italics). There were no registered mortalities associated with surgery. The most common surgical complications were dural tears/punctures, infections and hemorrhages. Advanced age and comorbidity were associated with a higher rate of complications.

In patients hospitalised with sciatica as their main complaint, sciatic pain regressed rapidly during the first 6 weeks after discharge, but their pain remained comparatively static after this time point. The following 4 factors were associated with their self-reported leg pain measured 1 year after hospital discharge: 1) High risk according to a psychosocial screening tool, 2) not receiving surgical treatment, 3) not actively employed prior to hospital admission and 4) self-reported leg-pain measured 6 weeks after hospital discharge.

The prognostic factors: age, sex, smoking, BMI, education, social status, analgesic use, previous lumbar surgery, clinical neurological examination of the lower extremities and CPM effect were all found to have poor predictive capacity.

Conclusions

Less than 7% of patients that undergo simple lumbar disc surgery experience an unfavorable event such as a reoperation, complication, or readmission. Whether the rate of unfavorable events is acceptable must be considered in relation to the health gains achieved by choosing surgical over conservative care, but patients can be informed that simple lumbar disc surgery is safe with low risk of mortality.

Sociodemographic characteristics, clinical examination, lifestyle factors and CPM effect were found to be poor predictors of long-term leg-pain in patients hospitalised with sciatica. Simply measuring leg-pain 6 weeks after hospital discharge could be an efficient way of establishing prognosis for patients hospitalized with sciatica as their main complaint. A patient's psychosocial profile was found to have some predictive value in the early stages of disease, but there is still a prognostic void that needs to be filled in order for us to reach the goal of early prognostication in patients hospitalized with sciatica.

SAMMENDRAG PÅ NORSK

Bakgrunn

Korsryggsmerter er den fremste årsaken til funksjonsbegrensning i verden. Smertene forårsaker redusert livskvalitet og sykefravær hos den enkelte, og kan være til stor økonomisk byrde for enkeltindividet og familien, og utgjør en stor kostnad for samfunnet som helhet. Rundt 5-10% av pasienter med ryggsmarter har i tillegg nerverots kompresjon som forårsaker utstrålende smerter i et eller begge ben, noe som gjerne blir betegnet som radikulære smerter eller isjias. I de tilfeller der symptomene er svært uttalte eller smerten vedvarer over lengre tid uten å gå i regress blir pasienten ofte tilbudt kirurgisk behandling. Prolapskirurgi er derfor en utbredt prosedyre og gjøres over store deler av verden. I forkant av prolapskirurgi er det viktig at pasientene informeres inngående om risikoen knyttet til selve operasjonen. Dessverre er vil en andel av pasientene ikke oppleve bedring, uavhengig av behandlingsmetode. Årsaken til at noen pasienter utvikler kroniske smerter er antatt å være en kombinasjon av biologiske og psykososiale faktorer. Derimot er det vanskelig å predikere hvilke pasienter som utvikler kroniske smerter og hvilke pasienter som blir friske. Forskning på prognostiske faktorer hjelper oss å bedre forstå hvem dette gjelder og hvilke mekanismer som ligger til grunn for mangel på tilfriskning.

Mål

Det overordnede målet med denne avhandlingen var å kartlegge graden av uheldige hendelser i forbindelse med prolapskirurgi, samt identifisere potensielle prognostiske faktorer assosiert med persisterende bensmerte blant kirurgiske og konservativt behandlede pasienter innlagt på sykehus med isjias. Dette arbeidet ble gjort gjennom epidemiologiske analyser av to store administrative sykehusregistre og to kohortstudier.

Målet med Artikkel I var å kartlegge graden av komplikasjoner, reoperasjoner og reinnleggelser på sykehuset etter prolapskirurgi.

Målet med Artikkel II var å identifisere potensielle prognostiske faktorer assosiert med selvrapportert bensmerte tolv måneder etter sykehusutskrivelse, hos kirurgiske og konservativt behandlede pasienter innlagt med isjias.

Målet med Artikkel III var å teste hvorvidt betinget smertemodulering (Conditioned Pain Modulation), målt 6 uker etter sykehusutskrivelse, var assosiert med selvrapportert bensmerte 12 måneder senere hos kirurgiske og konservativt behandlede pasienter innlagt med isjias.

Metode

Artikkel I var en longitudinell observasjonsstudie. Relevante medisinsk og kirurgisk data ble identifisert gjennom å kombinere prosedyre- og diagnosekoder fra administrative sykehusregistre, som inneholdt alle kirurgiske prosedyrer utført i det offentlige helsevesenet i perioden 1999-2013. Logistisk regresjon ble brukt til å analysere forholdet mellom pasientenes alder, kjønn, geografisk tilhørighet, utdanning, sivilstand, inntekt og komorbiditet opp mot oddsen for negative hendelser ved prolapskirurgi.

Artikkel II var en kohortstudie som fulgte 210 pasienter fra de ble innlagt akutt på sykehus for kirurgisk eller konservativ behandling av deres isjias. Pasientene ble rekruttert ved innleggelse og vurdert etter 6 uker, 6 måneder og et år etter utskrivelse. Potensielle assosiasjoner mellom de potensielle prognostiske faktorene målt tidlig i forløpet (innleggelse og 6 uker) opp mot selvrapportert bensmerte et år etter sykehusutskrivelse ble analysert ved lineær regresjon.

Artikkel III var basert på en kohortestudie hvor betinget smertemodulering var målt hos 111 isjiaspasienter som fremdeles hadde vedvarende bensmerte 6 uker etter utskrivelse fra sykehus. Pasientenes bensmerte ble også målt 12 måneder senere. En potensiell sammenheng mellom pasientenes betingede smertemoduleringsevne og selvrapporterte bensmerte ved et år ble analysert med logistisk regresjon.

Resultater

Blant de 34 639 lumbale prolapskirurgiske prosedyrene fant vi at 2,7 % hadde en kirurgisk *komplikasjon*, 2,1 % hadde en *reoperasjon* innen 90 dager, 2,4 % hadde en *reinnleggelse* innen 90 dager uten at det ble utført kirurgi og 6,7% hadde minst en av de overfor nevnte uheldige hendelsene (i kursiv). Det var ingen dødelighet assosiert med kirurgi, men høy alder og komorbiditet var assosiert med høyere grad av kirurgiske komplikasjoner.

Bensmerten hos pasienter innlagt med isjias ble raskt bedre i løpet av de første 6 ukene etter sykehusutskrivelse, men det var lite endring i smertebilde etter dette tidspunktet. Følgende fire prognostiske faktorer var assosiert med selvrapportert bensmerte målt ett år etter utskrivelse fra sykehus: 1) Høy risiko for kronisitet målt med et psykososialt screeningverktøy, 2) Ikke-kirurgisk behandling, 3) Ikke i arbeid forut for sykehusinnleggelsen og 4) Selvrapportert bensmerte målt 6 uker etter utskrivelse fra sykehus.

Alder, kjønn, røyking, BMI, registret partner, bruk av smertestillende, tidligere ryggkirurgi, nevrologisk undersøkelse av bena og betinget smertemodulering var alle funnet å ikke ha prognostiske egenskaper.

Konklusjon

Denne avhandlingen viser at færre enn 7 % av de 34 639 lumbale prolapsoperasjonene medførte en komplikasjon, reoperasjon og/eller reinnleggelse. Hvorvidt dette er en akseptabel rate må måles opp mot potensiell helsegevinst ved kirurgisk behandling i forhold til konservativ behandling, men pasienter kan informeres at enkel lumbal prolapskirurgi er trygt med liten risiko for mortalitet.

Sosiodemografiske karakteristika, klinisk undersøkelse, livsstilsfaktorer og betinget smertemodulering ble funnet å være dårlige prediktorer for langvarig bensmerte hos pasienter innlagt på sykehus med isjias. Ved å måle bensmerte seks uker etter utskrivelse gir derimot et grunnlag for å kunne prognostisere pasienter med isjias. Pasientens psykososiale profil ved sykehusinnleggelse ble funnet å ha noe prognostisk verdi tidlig i forløpet, men det er fremdeles et prognostisk tomrom som må fylles for at vi skal nå et endelig mål om tidlig prognose av pasienter innlagt sykehus med isjias.

PREFACE

In the years prior to conducting spinal research, I spent most of my working days as a doctor with the patients on the neurological ward at Ullevål University Hospital. A large proportion of these patients suffered from severe sciatica. Most of them were in great pain and desired some form of confirmation that their current situation would not become permanent. In the cases where lumbar surgery was offered, additional questions concerning adverse events and failed surgery rates were common. I was therefore always in search of literature that could provide answers to these questions. However, the literature was sparse, outdated, and often conflicting, especially in regards to surgical complication and reoperation rates. I often avoided questions concerning these rates by suggesting that the patient wait until the surgeon arrived, only to later find that the surgeons also struggled to provide concrete evidence-based answers on the subject. In my experience, patients and clinicians decide upon surgery based on the risk of unfavourable events and the risk of failed surgery against the patients overall prognosis. Therefore, it was frustrating for both the patients and myself that clear answers were not accessible. These unanswered questions were the motivation for this PhD project.

1. INTRODUCTION

1.1 Epidemiology

Low-back pain (LBP) is the single leading cause for disability worldwide causing activity limitation and work absence with subsequently enormous economic burden on individuals, families, communities, industry, health services and governments [1, 2]. Patients suffering from sciatica are estimated to account for 5-10% of patients that present to general practice with low back pain [3, 4]. A wide range of different definitions and interchangeable terms are used to describe pain in the distribution of the sciatic nerve, also known as sciatica. For instance, lumbosacral radicular syndrome, sciatic neuralgia and lumbar radiculopathy are other terms used interchangeably with sciatica. Sciatica is a clinical description based on symptoms of radiating pain or radicular pain in the leg [5, 6], most often (90%), but not exclusively, caused by a herniated intervertebral disc [6]. Despite this, the term “sciatica” is commonly used both in scientific literature and daily clinical practice to refer to the condition of radiating leg pain caused by a lumbar disc herniation [5, 6]. This thesis will therefore use the term “sciatica” in this manner, to indicate a lumbar disc herniation with radicular leg pain.

The annual prevalence of sciatica varies considerably from 1.2% to 43% [7]. The variance cannot only be explained by the differences in the populations studied, but also by the different sciatica definitions and methods of data collection. Stricter self-report definitions, the application of clinical assessments, and the use of point estimates as opposed to period estimates gives a more conservative estimate (2% - 5%), than when a wider definition such as “radiating leg-pain associated with low back pain” is accepted as sciatica [7]. Patients that suffer from sciatica report a higher degree of pain, have a poorer prognosis and show lower rates of return to work when compared to patients with LBP only [8, 9].

Most cases of symptomatic disc herniation are successfully managed in primary care, but an unclear percentage of patients are referred to hospital and other secondary care units. Referral is most often prompted by progressive muscular weakness, bladder dysfunction, a need for surgical treatment or intractable sciatic pain that is not manageable in an outpatient setting [5, 6]. The treatment of lumbar spinal disorders constitutes a large portion of hospital costs, but the major economic impact on society resides in health care utilization and work absenteeism before and after hospital discharge [10-12], thus early identification and intervention of this subgroup of patients is warranted.

1.2 Pathomechanisms and clinical features

In clinical practice, the human spine is divided into 4 segments, the cervical, thoracic, lumbar and sacral. Although this thesis is only concerned with the lumbar segment, the top three segments share some important anatomic principles. Each segment consists of several stacked vertebral bones separated by intervertebral discs. The intervertebral discs alleviate shear pressure and deterioration of the vertebrae. They also facilitate a better range of movement by allowing vertebrae to tilt forward without being obstructed by the vertebra directly below. The intervertebral discs consist of an inner gel-like nucleus consisting of proteoglycan and water held together by an irregular network of collagen type II and elastin, the nucleus pulposus. The nucleus is encircled by a series of collagen type I fibers, the annulus fibrosus. The annulus

fibrosus acts as a surrounding band, containing the viscous nucleus pulposus within the boundaries of the disc. The cartilaginous vertebral endplates act as the roof and floor of the disc, Fig.1.

Figure 1: The nomenclature and anatomy of the intervertebral disc

A disc herniation is defined as a displacement of the disc material (nucleus pulposus, annular tissue, cartilage, or any combination thereof) beyond the limits of the intervertebral disc space [13]. Disc herniation can occur in normal discs [14], but most often it proceeded by degenerative changes to the disc such as a dehydrated nucleus pulposus, a fissured, bulging, stiffened and weaker annulus fibrosus, and/or a sclerotic endplate with inflammatory changes [13, 15-17]. Aging is found to be highly associated with degenerative changes to the disc as it seems to increase the level of oxidative stress and reduce vascular channelling [18, 19]. Depending on the shape of the herniated material, herniated discs are commonly classified as either protrusions or extrusions [13]. Disc herniation in a posterior or posterolateral direction in the lumbar segment may impinge on the nearby (escaping) nerve roots, and, in cases where the nerve's motor fibres are sufficiently impinged, a reduction or loss of motor function in the innervated muscles ensues.

However, it is not the mechanical pressure on the nerve root that is thought to be the main cause of associated pain, but rather nerve compression in combination with a surrounding inflammatory response [20-22] dominated by cytokines such as IL-1 and $\text{TNF}\alpha$, both of which are expressed by the cells in the nucleus pulposus [23]. The cytokine activation is thought to effectively cause a catabolic response inducing an increase of vascularisation, in addition to nerve innervation and growth to the annulus fibrosus [24]. It also causes an immune cell response, recruiting macrophages and mast cells to the disc, causing phagocytosis, secretion of proteolytic enzymes that deteriorate and lead to resorption of the herniated material and further increase the cytokine levels [25, 26].

Though no definite consensus on the underlying molecular mechanism specific to the development of pain, the process of nerve compression coupled with inflammation is widely regarded as an essential part of the sharp burning, stabbing and radiating pain in the dermatomal distribution of the specific nerve affected, also known as sciatica. However, cellular or molecular mechanisms that explain why some patients continue to suffer from sciatic pain and others successfully heal have not yet been convincingly identified. In addition, disc herniation which

compresses nerve roots does not always result in clinical symptoms, and may remain undetected [27]. The mechanisms behind symptomatic vs. asymptomatic disc herniation are poorly understood. Clinically silent disc herniations have been reported to occur in 20%-30% of previously asymptomatic individuals [28, 29]. Since 10-15 % of all cases of sciatic pain are not due to a herniated lumbar disc, the large proportion of asymptomatic disc herniation may present diagnostic challenges. Clinicians may mistakenly attribute a herniated disc seen on a magnetic resonance imaging (MRI) scan as the cause of a patient's leg pain [30].

1.3 Treatment

It is common to divide treatment for sciatica into invasive and non-invasive treatments. In accordance with the NICE guidelines, the non-invasive treatments are further divided into pharmacological and non-pharmacological treatment. As for the use of invasive treatments, only lumbar disc surgery and epidural injections are recommended (not spinal injections and radiofrequency denervation) [31].

Non-invasive interventions are often referred to as conservative treatment in the hospital care setting. These treatments usually aim to reduce sciatic pain through the use of analgesics while the natural process of resorption of extruded disc material ensues. These treatments are often supplemented by non-pharmaceutical measures such as physical therapy modalities, which most often include exercise therapy and patient education. In general, patients are encouraged to continue with normal activities and try to avoid prolonged bedrest [31, 32]. The NICE guidelines recommends that patients be offered a supervised exercise program (biomechanical, aerobic, mind-body or a combination of approaches) where their specific needs, preferences and capabilities are taken into account [31]. These recommendations have been further supported in a Danish paper from 2018 aimed at developing clinical guidelines for non-surgical treatment in patients with lumbar radiculopathy [33]. The paper recognized 6 RCT's [34-39] where supervised exercise therapy in addition to usual care was compared to advice [35, 37, 38], sham exercises [36], rest [39] or usual GP care [34]. Exercise therapy was defined as exercises or physical activities, which had a therapeutic focus, tailored and adjusted to the individual patient, and delivered by a trained healthcare professional. The type of exercises presented in the studies varied considerably from general exercises [34], to more specific neuromuscular control exercises [36]. The Danish Guidelines recommended the use of exercise in sciatica. This was based on an observation that there was a trend found in all the six included studies, which favored supervised exercise. Despite this uniform trend was neither statistically significant nor clinically, the positive effect of supervised exercise could not be conclusively dismissed. In addition, the guideline emphasized that exercise had a potential positive effect on the patients' general health, and it could not out rule that exercise may also prevent recurrent episodes of sciatica and that serious adverse events were rare.

Spinal manipulation provided by manual therapists is widely used, but ultimately there is no definite documentation of a clinically relevant difference regarding the use of spinal manual therapy over supervised exercise therapy. Thus, manual therapy (spinal manipulation, mobilisation or soft tissue techniques such as massage) should only be considered as part of a treatment package (add-on to usual treatment) including exercise, with or without psychological therapy [31, 33]. Traction, acupuncture or electrotherapies (percutaneous electrical nerve stimulation and transcutaneous electrical nerve stimulation) are not recommended in acute or chronic sciatica [31, 33]. In addition to exercise, a psychological programme, incorporating a

cognitive behavioural approach should be considered in cases with significant psychosocial obstacles such as avoiding normal activities based on inappropriate beliefs about their condition preventing their recovery [31]. There is great practice variation with regards the type and extent of non surgical treatment, and a more systematic delivery has been warranted. However, it has been shown difficult to develop a successful stratification algorithm for allocating the different patients to specific non-surgical treatment regimes based on their psychosocial and pain profile. A recent study showed that a stratified care model for patients with sciatica consulting in primary care was not better than usual care for both clinical and health economic outcomes[40].

The evidence concerning the efficacy of commonly used analgesics for sciatica such as non-steroidal anti-inflammatory drugs, paracetamol, opioids, neuroleptics and spinal injections is not convincing when compared with a placebo [41]. Unfortunately, high quality randomised controlled trials are limited, and the significant overuse of analgesics continues despite the risk of adverse events [41-43]. The NICE guidelines recommends that sciatica patients should not be offered gabapentinoids, or other antiepileptics, oral corticosteroids or benzodiazepines as there is no overall evidence of benefit, and there is evidence of harm. Opioids should not be offered in the management of chronic sciatica, and if a person is already taking opioids, gabapentinoids or benzodiazepines for sciatica, the risks of continuing these medicines should be explained. In addition, patients should be made aware of the risk of harm and limited evidence of benefit in relation to the use of non-steroidal anti-inflammatory drugs (NSAIDs) in sciatica. If prescribed, it should be in the lowest effective dose for the shortest possible period of time, and the physician must take into account potential differences in gastrointestinal, liver and cardio-renal toxicity, as well as the person's risk factors, including age [31]

Surgical treatment is most often aimed at removing the area of the herniated disc that affects the nerve in order to alleviate pain and, if lost, to restore sensory or motor function. Standard open discectomy or micro discectomy are the two surgical treatments of choice, and have been shown to be equally effective [44, 45]. In cases of Cauda Equina Syndrome, immediate surgery is warranted, and there is also a consensus that in cases of severe or progressive paresis prompt surgery is the preferred treatment [30]. The likelihood of a patient undergoing lumbar disc surgery as opposed to conservative treatment varies between countries and the geographical regions within each country [46, 47]. This is thought to reflect the accessibility of surgery and differences in opinions among health care providers as to the benefit of lumbar disc surgery when compared to conservative care [46, 47].

A systematic review from 2011 identified only 5 randomised controlled trials where surgery was compared to conservative management in sciatica due to a lumbar herniated disc [48]. One “low-risk-of-bias” study showed that faster pain relief was achieved when surgery was offered to patients suffering from radicular pain for 6-12 weeks when compared to prolonged conservative treatment [49]. However, this same study [49], in addition to another high quality study [50], concluded that there were no significant differences in regards to the long-term outcome beyond 2 years. This finding was further supported by two other studies of lesser quality [51, 52]. Two of the studies also found no difference in outcome at one 1 year follow-up [49, 51]. The systematic review identified Weber’s pioneer trial from 1983 as the only trial that showed a statistically significant better result among the surgically treated patients at the 1-year follow-up, but this difference was no longer significant at 4 and 10 years [53]. Unfortunately, it has proven difficult to perform randomised controlled trials because of the high rate of non-adherence to group allocation [44, 48-50, 53]. There is a general consensus that lumbar disc surgery should in most cases only be offered to patients with muscle paresis or still suffering from severe sciatic pain beyond 8-12 weeks of conservative treatment [5, 30, 33]. A recent large single-centre trial

involving patients with severe sciatic pain lasting between 4-12 months caused by lumbar disk herniation found that lumbar disc surgery was superior to nonsurgical care with respect to pain intensity at 6 months of follow-up [54]. However, definite surgical selection criteria and timing of surgery is still a debatable issue warranting continued high quality research on the subject.

Standard open or micro discectomy have been the surgical treatment of choice for lumbar disc herniation for over 20 years, and are the most frequent lumbar spine operations in the world [55]. These surgeries are conducted under general anaesthetic, and are performed by either an orthopaedic surgeon or a neurosurgeon.

1.4 Prognosis and prognostic factor research

Prognosis means foreseeing, predicting, or estimating the probability or risk of future conditions [56]. In spinal research, prognosis is related to the probability or risk that a patient would develop a particular outcome based on a set of measured patient characteristics thought to influence this outcome. Outcomes can be specific events such as death or surgical complications, but they may also be quantities, such as changes in self-reported back or leg-pain, health-related quality-of-life measures, clinician-based grading of a paresis and other neurological symptoms and signs, as well as records of sickness absence and return to work. In prognosis research it is common to use patient-reported outcome measures (PROMs) of pain, disability, and health-related quality-of-life in addition to other measurements of unfavorable events.

Prognostic factors are any measure (such as age, blood type or education) with a given start point that is associated with a subsequent outcome (such as death, diagnosis of chronic pain or surgical complication). Prognostic factors can be used alone or in combination with other prognostic factors as part of a multivariate prognostic model. A single prognostic factor, however, rarely gives an adequate estimate of prognosis, as outcomes are most likely influenced by a multitude of different factors [56]. Regression models are commonly used in prognostic research, and use numeric values for each predictor in order to estimate an absolute risk or probability that a patient will experience a specific outcome. The most common reasons for developing a prognostic model are to better inform patients about the future course of their illness and to guide decisions on further treatment [56]. Regarding sciatica, there is a demand for more personalized prognostic models that more accurately predict outcome and the unfavourable surgical events associated with surgery. Although a prognostic model may provide insight into causality or pathophysiology of the studied outcome, a model is not synonymous with explaining the underlying cause [56].

1.4.1 Unfavourable event as an outcome in lumbar disc surgery

In this thesis, surgical complications, reoperations and readmissions are referred to as unfavourable surgical events. The term “unfavourable surgical event” lacks a clear universal definition, thus it should be further explained and defined when used outside the context of this thesis. Further, the term does not include other entities such as “a lack of improvement after surgery” or other measures used to explain unfortunate outcomes after surgery. Reports with a large sample size that describe the rates of unfavourable events in lumbar disc surgery in detail are limited and do not cover the long-term impact on health economy and patient pain and disability.

Though a variety of different definitions and classifications exist [57, 58], a *surgical complication* can be defined as “any deviation from the normal postoperative course” [59].

Lumbar disc surgery may be well executed without any complications, but still fail to relieve the patient of sciatic pain. This is often referred to in the literature as “failure to cure” [59, 60]. Complications may vary from minor events, requiring limited resources and exerting minimal long-term disability, to serious events requiring and exerting the opposite [59, 61]. In general, complications are said to contribute to an increase in length of hospital stay, mortality and level of patient care required at hospital discharge [61]. Unfortunately, underreporting of complications and lack of follow-up are suspected to be common among surgeons [62, 63], even though these complications are predictors for poor psychosocial long-term outcome [64]. It is estimated that 50% of surgical complications occur as a result of a provider error (avoidable), ranging from a technical blunder to missed opportunities to deliver optimal care such as “a patient with an increased risk of infection not receiving adequate prophylactic antibiotic therapy before surgery, for then to develop a postoperative infection” [65, 66]. However, complications can also occur because of a patient’s underlying comorbidity, despite receiving optimal surgical care. An example would be if a patient with an underlying comorbidity developed an unforeseen postoperative infection despite having been given adequate antibiotic therapy [65]. Distinguishing between complications due to patient disease and those due to provider error is important when discussing preventive measures.

Complication rates are often closely related to *readmission and reoperation* rates as many complications are discovered after hospital discharge and ultimately require subsequent hospital treatment [67, 68]. Readmissions can also be caused by patients having intractable postoperative pain, a suspected complication, the need for repeat diagnostics or a lack of expected improvement. Readmissions such as these may ultimately lead to revision surgery, which can be the result of diagnostic errors, relapse of disc herniation or a failure to remove enough of the herniated disc during the initial operation. It is, however, important to acknowledge that despite being successfully treated for sciatica, either through surgery or conservative care, there is a future risk of re-herniating the same disc or herniating another.

1.4.2 Self-reported outcome measures in sciatica

There are few objective means available to measure severity of pain and disability in sciatica. Therefore, what is considered a poor or expectable outcome may vary between patients and between clinicians. This is evident in the many different clinical study definitions of what is considered a poor outcome in sciatica. For instance, leg-pain, return to work, and disability are all potential outcomes which can all be scored and measured in different ways. Thus, finding a common definition for what should be considered a poor outcome is very difficult. In addition, full recovery with no persistent pain or disability is evident in a considerable proportion of sciatica patients; however, studies show that patients may be satisfied with their condition despite a degree of persistent pain and disability [69]. Although there are validated specific outcome measurement tools for sciatica, such as the Sciatica Bothersomeness Index, Sciatica Frequency Index [70] and Maine-Seattle Bothersomeness Index [71], that offer a clear definition of poor outcome, they are not used regularly in clinical studies addressing the outcome of sciatica. Interestingly, the less sciatica-specific outcomes such as; self-reported back pain, leg pain and the Oswestry Disability Index [72] are more frequently used in clinical studies, often together with outcome specific questions local to the individual study [73]. Thus, reaching a consensus on the definition of a poor outcome in sciatica remains a challenge [73].

Determining the rate at which patients who suffer from sciatica can expect a poor outcome is further complicated by the wide range of sciatica definitions and patient samples. A large proportion of patients (30-70%) recruited from a hospital or a secondary care unit using a

stringent clinical and radiological definition of sciatica are shown to have persistent symptoms of sciatica one year after hospital discharge [69, 74, 75]. The prognosis for full recovery in a patient visiting his/her general practitioner for the first time within a broad definition of sciatica, is generally considered more favourable, yet studies suggest that only just over half of primary care patients improve substantially over a period of 12 months [76].

1.4.3 Prognostic factors for poor outcome in sciatica

There is an extensive body of literature regarding prognostic factors for the development of chronic LBP and other chronic pain conditions, but there are far fewer studies concerning sciatica specifically. The biopsychosocial model stresses the importance of psychological and social factors in addition to biological factors as key players in the transition from acute to chronic pain [77-79]. Multiple studies and resources have been devoted to further show that psychosocial factors such as fear-avoidance beliefs, catastrophizing, somatization and kinesiophobia are associated with poor outcomes in LBP [80-86]. However, it is unclear whether psychosocial factors have the same level of prognostic impact for sciatica specifically. Studies suggest that psychosocial factors have a prognostic value, but are less dominant predictors than what is observed in non-specific LBP [73, 87, 88]. This has warranted further studies aimed at examining the role of psychosocial factors in sciatica, since such psychological characteristics are potentially modifiable. Patients hospitalised with sciatica are especially compelling in this context as it is suggested that this group have particularly poor outcomes compared to those with LBP [89, 90].

This prognostic and explanatory void among sciatica patients has also catalysed researchers to explore the role of biological factors such as genetic, transcriptomic, proteomic, metabolomic, neurophysiologic, radiologic, and physiological factors in chronic sciatica, rather than settling entirely upon psychological and cultural factors as the main determinants for poor outcomes. Unfortunately, there are currently no biological markers that can adequately predict any poor outcome in sciatica either alone or in combination with other factors. A recent systematic review concluded that multiple inflammatory markers such as the interleukins, tumor necrosis factor- α and phospholipases were found to be present in sciatica, but there was insufficient evidence to draw any firm conclusions regarding the level of inflammatory activity in relation to clinical symptoms [91]. Similarly, multiple genetic variants such as the OPRM1 rs1799971 G allele, COMT rs4680 G allele, MMP1 rs1799750 2G allele, IL1 α rs1800587 T allele, and IL1RN rs2234677 A allele have been found to be associated with persistent sciatica, but their prognostic value in a clinical setting is undetermined [92]. Multiple blood biomarkers such as keratin sulfate, hyaluronan and cartilage oligomeric matrix protein have not been shown to have any diagnostic value [93]. In addition, a small study found that depressed levels of fibrinolytic activity were associated with 12-month outcome in surgically treated sciatica patients [94].

Further, clinical and lifestyle parameters such as long duration of symptoms, specific findings on clinical neurological examination [90, 95, 96], specific MRI findings [97-99], being female [100] and smoking [88] were found to be associated with poor outcome in sciatica in some individual studies. Nonetheless, due to the limited strength of impact and generalisability of these factors, they have yet to equip clinicians with useful prognostic tools that can direct treatment at an early stage of the disease [73, 91].

1.4.4 Conditioned pain modulation in sciatica: a potential prognostic factor

The subjective experience and reporting of pain intensity vary greatly, and do not necessarily match with clinical findings [101]. The mismatch in objective findings and pain perception can

be partially explained by endogenous pain modulation, that is, the central nervous system's modulation of nociceptive pain signals [102]. Nociceptive pain signals from injured/ affected peripheral tissue ascend through the spinal cord and brainstem to reach the cerebral cortex where the nociceptive signals are ultimately perceived as painful. At several points along this pathway, the incoming nociceptive signals are inhibited or facilitated along their way to the brain. The final perception of pain is indirectly affected by these mechanisms through a series of ascending (bottom up) and descending (top down) pain-controlling pathways along the spinal cord, brain stem and cortex. The descending pain-controlling pathways (from brain to brainstem), can for example be modulated by psychological interventions. The ascending pain-controlling pathways (from peripheral nerves and spinal cord to brainstem) also have the potential for modulation [102].

One of these proposed methods for pain modulation can be demonstrated through a psychophysical experimental measure termed 'conditioned pain modulation' abbreviated as CPM [102]. The concept of CPM can be demonstrated by a painful stimulus being perceived as less painful in the presence of or shortly thereafter, a second painful stimulus. The concept of CPM or "pain inhibiting pain" has been recognized since ancient times. It was previously used in relation to painful surgical procedures on domesticated animals, such as castrations of cattle and horses by applying a painful clamp to the animal's nose in order to diminish pain associated with the castration itself [103]. This early example of Conditioned Pain Modulation reveals the inhibitory effect on one nociceptive stimulus (the castration) when another concomitant nociceptive stimulus is inflicted (the nose clamp) [102]. The difference in perceived pain of the first painful stimulus with and without the second painful stimulus is termed the CPM effect. It can be studied experimentally in laboratories by measuring the pain intensity of a noxious test stimulus (TS) given one time as a stand-alone, and at another time during or immediately after a second noxious conditioning stimulus (CS) [102].

All types of nociceptive stimuli are said to trigger an inhibitory effect (i.e. thermal, mechanical, chemical and electrical stimulus) and the degree of inhibition is also proportional to the intensity and duration of the applied nociceptive stimulus [104]. In addition, greater pain inhibition occurs when additional distraction tasks are used compared to CPM testing alone, which indicates that CPM acts independently from distraction, but with possible overlaps and psychological properties [105]. However, whether CPM can change during or due to the presence of chronic pain or if CPM changes in the acute stages disease is still largely unknown. Nevertheless, impaired CPM levels have been found both in acute low back pain patients as well as in chronic low back pain patients, indicating alterations in CPM might already be present in acute stages of disease [106]. Studies have also shown that impaired CPM in osteoarthritis was restored once the painful joint was replaced suggesting that CPM changes in the presence or absence of clinical pain [102].

Increasing evidence shows that populations with chronic pain disorders have less efficient CPM effect compared to healthy individuals [107]. This suggests an association between chronic pain disorders and alterations in pain modulation [102]. Having impaired or reduced inert CPM effect has been particularly associated with chronic pain disorders such as irritable bowel syndrome, migraine, tension-type headache, temporomandibular joint (TMJ) disorders, osteoarthritis and muscle pain, whiplash-associated disorders and fibromyalgia [102]. Less efficient CPM is most commonly observed in idiopathic pain syndromes such as fibromyalgia, irritable bowel syndrome, temporomandibular disorder and headaches [107]. In addition, reduced or absent CPM effect has been associated with non-response in different pharmacological and non-pharmacological pain treatments [102].

Exercise therapy has been known to induce pain relief in many pain conditions. During strenuous exercise, an increased concentration of lactic acids and potassium (K⁺) activates nociceptors and to a certain extent, causes exercise to be perceived as painful [108]. Individuals with more efficient CPM, have been previously shown to experience greater reduction in pain after exercise compared to those with less efficient CPM, suggesting greater exercise-induced hypoalgesia can be predicted by CPM. With this in mind, CPM has been forwarded as a possible explanation for exercise-induced hypoalgesia [109]. However, these findings are inconsistent with a Finnish study from 2014, which found no correlation between CPM and exercise-induced hypoalgesia, thus suggesting CPM and exercise-induced hypoalgesia are based on different endogenous modulation mechanisms [110]. Similarly, CPM has been proposed as a useful tool to determine patient response and explain the effect of transcutaneous electrical nerve stimulation (TENS). TENS is mediated by endogenous inhibition, and CPM is used to measure the individual's ability to produce endogenous inhibition [111, 112].

Several CPM studies performed on LBP patients have investigated the association between low CPM effect and chronic LBP [106, 113, 114], but no studies have investigated its association with persistent sciatic pain. A potential association might suggest that measuring CPM effect could be a valuable prognostic tool in the management of sciatica.

Currently, the greatest challenge regarding CPM is the significant variation in the experimental protocols [115]. CPM can be tested by a parallel- or a sequential paradigm. In the parallel paradigm, the test stimulus (TS) is first given alone and then in parallel with conditioning stimulus (CS), while in the sequential method, TS is first given alone and then immediately after CS. The parallel method has shown to give greater pain reduction compared to a sequential paradigm. However, in the parallel method, it can be difficult to distinguish between a clean physiological inhibitory effect due to nociceptive stimulation and the psychological influences due to distraction [102, 115].

It is broadly accepted that the testing stimulus and conditioning stimulus are of different modalities [115]. The most commonly used conditioning stimulus is an upper extremity immersed in cold water [116]. Cold water has been shown to be more reliable as a conditioning stimulus compared to ischemic pain and pressure pain in relation to both inter- and intra-individual variance [117]. Heat stimulation is also frequently used as a conditioning stimulus by employing an electrical contact heat thermode or water bath [116]. The conditioning stimulus is usually given as a continuous stimulation over a period of 30 seconds to 2 minutes. It is widely accepted that CS needs to be sufficiently painful in order to provide an inhibitory effect on the testing stimulus [102]. Furthermore, an increase in the conditioning stimulus intensity increases the magnitude of the CPM effect [118].

With regards to the testing stimulus, thermal and mechanical stimuli are the recommended modalities [115], but ischemic and electrical stimuli applied to either upper or lower extremities is also used frequently [116]. The duration of stimuli among different CPM studies varies from milliseconds to persistent stimulation over several minutes [102]. In addition, the stimulation can be either tonic (a persistent continuous stimulation from 30-120 seconds) or phasic (repetitive stimulation of shorter duration) [119]. The intensity of the test stimulus is commonly set to an individual pain level, where the intensity is painful enough to detect a CPM effect, but not intolerable or harmful in any way [102]. Most studies use a visual analogue scale or a numeric rating scale to measure changes in pain. However previously, neurophysiological measures such as the nociceptive withdrawal reflex recorded by electromyography (EMG) have also been used [116]. In addition, there is a consensus that pain inhibition should always be reported by a negative value, and results denoting pain facilitation reported by a positive value.

Lastly, the change in pain should be presented as absolute values, percent change or, preferably both [115].

2. AIMS OF THE STUDY

The overall aim of this thesis was to assess the frequency of unfavorable events such as complications, reoperations and hospital readmissions in lumbar disc surgery as well as to study potential prognostic factors for the persistence of leg-pain one year after hospital discharge in surgically and non-surgically treated patients with sciatica.

Our hypothesis was that the rates of unfavorable events are lower than previously thought, and that complete prognostication of poor outcome in sciatica is dependent on a multitude of different prognostic factors rather than a single biological or psychosocial factor.

Aim of Paper I:

To determine the rate of prolonged hospital admission and unfavourable events such as complications, reoperations, and readmissions for patients operated for lumbar disc herniation over a 15-year period from 1999 to 2014. The secondary objective was to investigate the impact of patient comorbidity and sociodemographic factors on the four outcomes.

Aim of Paper II:

To identify neurological signs and psychosocial characteristics as potential prognostic factors for persistent leg-pain one year after hospital discharge in surgically and non-surgically treated patients with acute sciatica.

Aim of Paper III:

To determine whether CPM effect measured 6 weeks after hospital discharge was associated with leg-pain measured at 12 months in surgically and non-surgically treated patients with acute sciatica.

3. MATERIALS AND METHODS

3.1 Study Designs

Table 1 shows an overview of the study design of the three papers in the present thesis.

Table 1: Overview of the study design in Paper I-III

	Paper I	Paper II	Paper III
Type of study	Retrospective descriptive study using large administrative medical database	Prospective cohort study	
Patients	Patients were identified using surgical codes and diagnostic codes from all public hospitals in Norway. Symptomatic lumbar disc herniation	Patients were recruited upon acute hospital admission with symptomatic lumbar disc herniation. Oslo University Hospital only. Both surgical and non-surgical patients	
Main statistical method used	Logistic regression	Linear regression	Logistic regression
Main factor(s) of interest	Sociodemographic factors	Sociodemographic, psychosocial, lifestyle, Neurological signs	Conditioned Pain Modulation (CPM) effect
Main outcomes	Surgical complication Reoperation within 90 days Readmission within 90 days Unfavourable surgical event	Self-reported leg-pain 12 months after hospital discharge	

Paper I was a longitudinal observation study. Data from herniated lumbar disc operations were retrieved using a large medical database and a national population registers from the period 1999 to 2013.

The study design in Paper II was a prospective cohort study modeled after the Prognosis research strategy (PROGRESS) framework for prognostic factor research [120-123], Fig.2. Patients were evaluated at 4 time points (baseline, 6 weeks, 6 months and 12 months). The 6-month follow-up was conducted via a telephone interview and the 12-month follow-up through a postal questionnaire. The possible association between the prognostic factors tested and the 12-month outcome was analysed by linear regression modelling.

Figure 2: Flow diagram of the study population in Paper II

The study design in Paper III was a prospective cohort study with assessments at two time points: 1) CPM testing 6 weeks after discharge from hospital with sciatica as the main complaint 2) a follow-up postal questionnaire at 12 months post hospital discharge, Fig. 3. The possible association between CPM effect and 12-month outcome was analyzed by logistic regression modelling.

Figure 3: Flow diagram of the study population in Paper III

3.2 Study Populations

Norway, with a population of 4.9 million, has a large, government-funded national healthcare system divided and run by four regional health authorities: the South-Eastern, Western, Central, and Northern Health Regions. Prior to 2003 these public regional health authorities were responsible for almost all spinal surgery in Norway, but from 2007 the public sector was only responsible for an estimated 80% of lumbar disc surgery due to an increase in private surgery [124]. Papers I-III were based on Norwegian citizens ≥ 18 years of age admitted to public hospitals in Norway with disc herniation.

In Paper I, patient selection was performed retrospectively using two national medical databases for the years 1999-2009 and 2010-2013 respectively. Patient selection was performed through a combination of procedure and diagnosis codes. First, the relevant surgical procedure codes were retrieved. To ensure that the procedure codes were specific for lumbar disc herniation, only patients with a diagnosis code indicating degenerative disc herniation were included. The dataset is thought to represent all surgeries for lumbar disc herniation carried out through the public health system in Norway during the period 1999-2013.

Paper II studied patients acutely admitted to Oslo University Hospital for either surgical or nonsurgical treatment of sciatica. The study population was representative of and captured almost all patients living in Oslo that required (for any reason) an acute hospital admission for (clinically and radiologically confirmed) sciatica between 2012 and 2014. The hospital provides care to all Oslo residents who require either acute surgical or non-surgical treatment of sciatica. Sciatica was defined in the paper as “radiating leg-pain with dermatomal distribution and a radiological confirmation of disc herniation on magnetic resonance imaging”. The physicians responsible for patient admission to the hospital were working independently of the study program. Upon hospital admission, most patients were treated with a combination of paracetamol, non-steroidal anti-inflammatory drugs and mild or strong opioids. All clinical exams and physical measurements were performed by the same three physiotherapists. Lumbar surgery was typically offered to patients with severe motor deficits or unbearable sciatic pain refractory to any analgesic.

The patient population in Paper III shared many of the same patients with Paper II. The main difference was that only patients that continued to suffer from leg pain 6 weeks after hospital discharge were eligible for inclusion and subsequent CPM testing.

3.3 Measurements

3.3.1 Outcome measures

Surgical complication (yes/no): One of four main outcomes presented in Paper I. The outcome contributes important information regarding the quality and costs of surgical and hospital care [125]. Further, knowledge of the complication rate is valuable to both the patient and clinician when discussing the pros and cons of surgical treatment for sciatica. Healthcare workers registered the complications using the International Classification of Diseases number 10 (ICD-10) complication codes either on initial hospital admission or subsequent readmissions within 90 days of initial surgery.

Reoperation (yes/no): This outcome is important in determining quality of lumbar disc surgery, but is also vulnerable to misinterpretation. There is a risk of overestimating the reoperation rate by interpreting subsequent operations occurring at different lumbar levels than the original surgery as reoperations rather than separate entities. The term “reoperation” is often used interchangeably in the literature for both immediate correction of a failed lumbar disc surgery because of a complication and revision surgery due to re-herniation, a process that often occurs long after the original surgery [126-128]. In this context, we defined reoperation as “all disc herniation surgeries within 90 days of initial surgery (immediate reoperation included) ensuring that the timeframe between the two surgeries was short, thus indicating that the primary surgery had failed and that the second surgery had failed, creating the need for a second surgery.

Non-surgical readmission (yes/no): Defined in Paper I as “a patient readmitted within 90 days of lumbar surgery with a lumbar complaint, but not re-operated”. The outcome is of relevance as it reflects the multifactorial and diverse issues that ensue post-surgery and has economic implications in terms of planning available hospital beds, nurses and physicians. This outcome is strongly dependent upon local hospital practice with regards to how they deal with possible reasons for readmission such as unexpected strong postoperative pain, need for immediate repeat radiology and issues concerning poor postoperative function.

Unfavourable surgical event (yes/no): Though there is no universal definition of this term. It is defined in Paper I as “surgery with a complication, reoperation, and/or non-surgical readmission”. The outcome measure was made specifically for Paper I and is meant to provide a collective term for the overall rate of experiencing a complication, reoperation and/or a readmission. The outcome “unfavourable events” divides surgeries into two groups: those seemingly without any major obstacles, and those that require additional resources. The term “adverse event”, defined as an “unintended injury or complication resulting in a prolonged hospital stay, disability at the time of discharge or death caused by healthcare management and not by the patients” is closely related to an “unfavourable event”, and this more established term could arguably have been used instead [129]. However, with no measures of postoperative disability or patient related outcomes available, and the wish to include a readmission as part of a collective term for poor surgical outcome, the less established outcome “unfavourable events” was used as the main outcome.

Leg-pain (below the knee) experienced in the last week, measured on a self-reported Numeric Rating scale from 0-10 (0=no pain, 10=worst pain ever) was used as the final outcome in Paper II. Leg pain has been found to be one of the most responsive outcomes for capturing disabling symptoms for patients suffering from sciatica [130, 131].

Even though the final outcome measurements in Paper III were collected in the same manner as in Paper II, the condition of normality could not be fulfilled and the 0-10 self-reported Numeric Rating Scale had to be dichotomized into the following; a leg pain score of ≥ 1 the past week. There is no universal way of dichotomizing such a NRS scale for sciatic leg-pain, and therefore a cut-off value that unquestionably separated pain free individuals from those with continued leg-pain (regardless of magnitude) was preferred (they either have/do not have pain). The cut-off value for the outcome would be easy to understand from a patient’s, clinician’s or a researcher’s perspective.

3.3.2 The potential prognostic factors considered

Potential prognostic factors tested in Papers I, II and III are presented in Table 2. The selection of potential prognostic factors in each paper was based on previous research evidence in sciatica.

Table 2: Potential prognostic factors tested in Papers I-III

Potential prognostic factors	Paper nr.		
	I	II	III
Age	x	x	x
Sex	x	x	x
Civil status	x	x	
Education	x		
Current employment status		x	
Income	x		
Body Mass Index (BMI).		x	
Comorbidity	x		
Current smoking status		x	
Hypertension		x	
Use of analgesics		x	
Duration of leg-pain prior to inclusion		x	
Leg-pain intensity upon inclusion		x	x
Örebro Musculoskeletal Pain Questionnaire (ÖMSPQ)		x	
Oswestry Disability Index		x	
Clinical examination		x	
Previous lower back surgery		x	x
Surgical treatment within study		x	x
Year of hospital discharge	x		
Conditioned Pain Modulation (CPM)			x

Though poorly amendable to intervention, age and sex as potential prognostic factors were tested in all 3 papers. The general prevalence of chronic pain has previously been shown to be higher among women and the elderly thus a requisite to be included in pain research [132, 133]. The 3 age groups (the first used as reference category) in Paper I were meant to reflect the young (18-39), middle-aged (40-59) and elderly (≥ 60) populations of sciatica patients based on the age distribution from other published sciatica cohorts [127, 134].

Depression, anxiety and catastrophizing behaviour have been shown to be associated with chronic pain conditions [135, 136]. The Örebro Musculoskeletal Pain Questionnaire (ÖMSPQ), used as a prognostic factor in Paper II, has previously been used to identify patients with LBP in general practice who were at risk of developing persistent back pain by: measuring days off work, anxiety, tension, depression, pain, activities of daily life, ability to cope, job satisfaction,

fear avoidance beliefs and the patients' recovery expectations [137, 138]. A potentially strong correlation between the questionnaire scores and persistent sciatica among the hospitalised sciatica patients would indicate that the questionnaire could also have value; (1) not only for non-specific low back pain, but sciatica patients specifically, (2) in the secondary care setting and (3) among patients treated with lumbar disc surgery.

Comorbidity was tested in Paper I using the Charlson Comorbidity Index (CCI) [139]. In the Charlson Comorbidity Index (CCI) each comorbid condition (heart disease, diabetes, chronic lung disease, cancer etc., a total of 22 conditions) is weighted differently on a scale from 1 to 6 depending on severity, eventually resulting in a final "comorbid score" based on the patients predicted one-year mortality. The index is very useful for condensing the different comorbidities rather than trying to analyse and interpret the effect of multiple individual comorbidities.

Prognosis for revision surgery is dependent on outcome of primary surgery, but there is no convincing evidence that prior lumbar surgery in itself is correlated with a poor outcome of repeat surgery [140]. Therefore, previous lumbar surgery was elicited as a potential prognostic factor in Paper II.

BMI, hypertension and smoking are lifestyle factors that have been shown to be directly or indirectly associated with chronic pain conditions [141]. The literature also suggests that acute pain management has preventive properties in relation to later development of chronic pain conditions. The duration of pain prior to inclusion, the grade of pain at baseline and the use of analgesics were therefore important as potential prognostic factors in Paper II [96, 142, 143].

Though only a few extensive studies are available, there is limited evidence that specific neurological findings are accurate predictors of pain chronicity in sciatica [73, 144]. However, several studies have found a slight prognostic value in neurological findings [90, 100, 145, 146]. A battery of different clinical neurological tests (power, sensation, reflexes and straight leg test) on the lower limbs was performed in Paper II. All four tests were awarded equal value when added together, resulting in a potential minimum score of 0 and maximum score of 4. In order to preserve test reliability, the same three physiotherapists were responsible for all clinical testing.

Figure 4 gives an overview of how CPM testing was conducted. At the time of inclusion, there was insufficient data to identify a superior CPM protocol [115, 147]. However, the CPM testing in Paper III adhered to most of the recommendations later published in a consensus paper on the matter [115]. The body's limbs were recommended as superior sites for testing of the CPM paradigm. Further, it was recommended that temperature stimulus be individually calibrated rather than fixed at a specific temperature. This was done to avoid floor and ceiling effects in patients that perceived the fixed temperatures as either extremely painful or not painful at all. Results denoting pain inhibition were to be reported with negative values, and effect presented as absolute values and/or percentage change. The recommendation paper concluded that the conditioning stimulus should be applied immediately after the test stimulus (sequential order), rather than applied simultaneously as done in Paper III (parallel order).

Figure 4: Conditioned Pain Modulation (CPM) testing

3.4 Statistical analysis

3.4.1 A general overview of statistical methods and hardware

All papers were co-authored by a medical statistics expert. Regression analysis was used as the main statistical method in all three papers. Chi-square (χ^2) tests and independent t-tests were used to explore potential differences between patient stratified subgroups (male vs. female). An overview of all the main statistical methods used in the three papers is presented in Table 1. All data analyses were performed using SPSS 22.00 (SPSS Inc., Chicago IL.) (Paper II and III) and SPSS version 25.0 (IBM, Armonk, NY, USA) (Paper I). Significance level was set to $p \leq 0.05$ (two-sided) and all model assumptions were checked and fulfilled unless otherwise stated in the article.

3.4.2 A description and rationale for the statistical methods

Paper I.

A prognostic model using *logistic* regression analysis can only have a dichotomous (binary) outcome [148]. The outcome of interest in Paper I was dichotomous in nature (surgical event yes/no), and thus dictated the choice of statistical method. Multivariate logistic regression analysis was used to investigate the association between the independent variables (age, sex, comorbidity, education, civil status, income, and regional health authority affiliation) and the dependent variable unfavorable event. All independent variables were entered simultaneously, and all significant associations were described with Odds Ratios (OR) and 95% Confidence Intervals (CI).

Paper II.

The main outcome variable in Paper II was continuous, therefore linear and not logistic regression was used as the main statistical method. The impact of the potential prognostic factors (independent variables) on leg pain (dependent variable) was analysed using linear regression modelling. Results were expressed with unstandardized beta coefficient (the regression slope), level of significance (p -value) and the Nagelkerke R^2 [149]. The prognostic factors in the final model were selected using a “stepwise” approach [150]. Each potential prognostic factor was individually tested by simple linear regression. If the individual factor was found to be associated with the outcome by a p -value <0.1 (lax criterion) it would be entered into a *multivariate* regression model. Prognostic factors that lost their association with the outcome once put into the multivariate model (p -values >0.05) were removed, with the highest p -value being removed first. However, a strict p -value driven approach for selecting variables into a multivariate model is not optimal because factors can have an impact in a multivariate model even if not statistically significant on the univariate level [151, 152]. Non-significant factors often change the effect of other factors once included in a multivariate model, or have prognostic properties when included with other factors, especially if they are not truly independent of each other [153]. With this in mind, age, sex, and “surgical treatment yes/no” were added to Paper II’s multivariate model regardless of p -value when tested individually. A variance inflation factor (VIF) was used when testing for multicollinearity and Durbin Watson when testing for serial correlation [149]. Repeated Measures Analysis of Variance (rmANOVA) was used to show and analyse changes in leg-pain measured throughout the study [149].

Paper III.

Again, the choice of outcome variable influenced the choice of statistical method. In Paper III, the outcome was initially measured on a continuous scale (leg pain on a Numeric scale (0-10)), most suitable for linear regression analysis. Unfortunately, the outcome data was not normally distributed, but rather skewed towards the left, residing in the extreme ends of the scale, and was not sufficiently impacted by any attempts of scale transformation. Therefore, the linear outcome measurements had to be binary, and, accordingly, logistic regression, not linear regression, was the main statistical method used to investigate the association between the independent variables (CPM effect, age, sex) and the dependent variable of leg-pain. All independent variables were entered simultaneously, and all significant associations were described with OR and 95% CI.

3.4.3 Sample Size and missing values

Literature indicates that data should contain at least 10 events for each variable entered into a regression model [56, 154]. Incorporating too many explanatory variables compared to the size of

the study has been estimated to occur in roughly 1 of 5 medical research publications [155]. Sample size calculations for the regression models in this thesis were based on the principle that at least 10 participants were required for each prognostic factor that was studied. With roughly 2000 events per prognostic factor tested, sample size was not an issue in Paper I. In Paper II, 210 patients provided sufficient statistical power to test approximately 10-15 prognostic factors. In Paper III, a sample of 110 patients was estimated to provide sufficient statistical power to show any clinically relevant association between CPM effect and the final outcome, when adjusting for 3 other variables.

A multiple regression model is mathematically dependant on each patient having a complete set of values for the different factors tested in the model [149, 150]. A patient with a missing value (for example age) would have to be subsequently dropped from analysis, unless the missing value was to be imputed. Missing values were imputed (with mean values) in Paper II when single questions in the ÖMSPQ and ODI questionnaire were not answered. Missing values were almost non-existent in Paper I and Paper III, thus no imputation was needed. Missing data were assumed to be missing at random.

3.5 Ethics

All papers were funded by the South-Eastern Norway Regional Health Authority and approved by the Norwegian Regional Ethics Committee (REC) (2013/1662^(paper I), 2012/1108^(paper II and III) REC south-east D). The storage of data in Paper I was approved by the Norwegian Data Inspectorate (2014/14413) and the Ministry of Health. Data was received through an encrypted PIN, generated by Statistics Norway. This did not enable us to have access to patient names, addresses or personal identification numbers, thus ensuring patient privacy. For Papers II and III, written informed consent was obtained prior to participation and the patients were thoroughly informed about the study. Human testing in Paper III was performed in accordance with the Declaration of the World Medical Association guidelines for experimental investigation [156]. We ensured that no patient was at risk of any permanent harm, or felt unwanted pressure to conduct and follow through with the pain experiments. The test method and instrumentation was previously widely tested without any indication of inflicting an instant or delayed pain condition, or any other form of permanent harm. No authors declared any conflict of interest with regards to the 3 papers.

4. SUMMARY OF THE RESULTS

Paper I. *Complications, reoperations, readmissions, and length of hospital stay in 34 639 surgical cases of lumbar disc herniation*

From 1999 to 2013, 34 639 simple herniated lumbar disc surgeries were performed in Norwegian public hospitals, 90% discectomies and 10% decompressions. Microdiscectomy was specified in roughly 70%, the remaining 30% were specified neither as microdiscectomy or standard/open procedures. Of the 34 639 operations, 2.7% (95% CI 2.6-2.9) had a *surgical complication*, 2.1% (95% CI 2.0-2.3) had *repeat surgery* within 90 days, 2.4% (95% CI 2.2-2.5) had a *non-surgical readmission* within 90 days and 6.7% (95% CI 6.4-6.9) experienced at least one of these unfavorable events (in cursive). The most common surgical complications were dural tears/punctures, infections, and hemorrhages, with 30% of the complications being registered on a subsequent readmission. At least 14% of the reoperations were due to surgical complications. Through logistic regression analysis, unfavorable surgical events were found to be associated with advanced age and comorbidity, but no association was detected for the following sociodemographic factors: gender, geographical affiliation, education, civil status and income.

Paper II. *Prognostic Factors for Persistent Leg-Pain in Patients Hospitalized With Acute Sciatica*

Of the 210 patients included in the study, 151 (72%) completed the study. The dropouts differed from the patients completing the study in that they had a higher smoking percentage (37% vs. 37%) and less education (33% vs. 48% with <12 years). Mean self-reported leg-pain intensity was measured on a 0 to 10 Numeric Rating Scale (0= no pain, 10=worst leg-pain imaginable), and was measured at 7.3 (95% CI 6.9-7.8) on hospital admission (baseline). It was measured at only 2.7 (95% CI 2.1-3.1) 6 weeks after hospital discharge. No significant change in mean leg-pain was observed between 6 weeks 2.03 (95% CI 1.6-2.4), 6 months 2.03 (95% CI 1.6-2.4), and 12 months 2.5 (95% CI 1.9-2.8), Fig.5. A total of 93 patients (62%) received surgical treatment during the study. Upon hospital admission, the patients that received surgical treatment had a significantly higher level of leg-pain when compared to patients that did not receive surgery, 7.9 (95% CI 7.5-8.3) versus 6.3 (95%CI 5.6-7.1) ($p<0.01$). The overall reduction in mean leg-pain from baseline to 6 weeks was also significantly higher for patients receiving surgery. There were two predictive multivariate linear regression models. The first one was used during hospital admission; the second was used 6 weeks after hospital discharge. The first model found the following 3 factors to be significantly associated with leg-pain at 12 months post hospital discharge: 1) High psychosocial risk according to the Örebro Musculoskeletal Pain Questionnaire, 2) not receiving surgical treatment and 3) not actively employed prior to hospital admission, with the unstandardized beta coefficients of 1.55 (95% CI 0.72–2.38, $p<0.001$), 1.11 (95%CI 0.29–1.93, $P=0.01$), and 1.47 (95%CI 0.63–2.31, $p<0.01$). The second model found that self-reported leg-pain measured 6 weeks after hospital discharge was associated with leg-pain at 12 months, 0.49 (95% CI 0.34–0.63, $p<0.001$). The Örebro Musculoskeletal Pain Questionnaire was shown to have most of its prognostic value among the non-surgically treated patients. Age, sex, smoking, education, social status, analgesic use, previous lumbar surgery and clinical

neurological examination of the lower extremities were all found to have poor predictive capacity on the outcome.

Figure 5: Leg-pain as a function of time

Paper III. Predicting the Outcome in Persistent Sciatica Using Conditioned Pain Modulation: 1-year results from a prospective cohort study

Of the 149 patients included in the study, 111 of them completed the study. A total of 51 patients (54%) received lumbar disc surgery 6 weeks prior to CPM testing. Mean CPM value was measured to -2.4 cm (95% CI -2.8 , -2.1) on a 10 cm visual analogue pain scale. No statistically significant difference was detected when comparing surgically and non-surgically treated patients, -2.3 cm vs. -2.5 cm, $t(109)=0.56$, $p=0.59$. Mean NRS leg pain was measured to 4.0 (SD 2.1) for the participants at baseline, and was subsequently measured to 2.5 (SD 2.5) at 12 months. Logistic regression analysis did not reveal any statistically significant association between the CPM effect and the persistence of any patient reported leg pain 12 months later, OR 0.87, 95% CI 0.7–1.1, $p=0.23$. Further, no significant association was found when adjusting individually or concurrently for the following: age, sex, baseline leg-pain and surgical treatment.

5. DISCUSSION

5.1 Methodological considerations

5.1.1 Strengths and disadvantages of study designs

Though prospective cohort studies are common and most often the preferred design in prognostic factor research, large medical databases offer an opportunity to research rare events over a long time period within a large sample [157]. The outcomes in Paper I are examples of such rare events on a large scale, but this type of database would require significant resources to explore them in a cohort study. The retrospective use of large medical databases in prognostic factor research also has its challenges, for example, determining whether exposure or outcome came first. Therefore, there is a limit to the variety of different potential prognostic variables that can be tested in such a database. Large administrative databases are also historically limited to non-modifiable categorical variables that are unethical to use when directing treatment and may not be accurate for individual cases. It is also typical that the design used for Paper I will not have patients at the same stage of their disease, as an inception cohort study would, thus the results are less applicable for prognosis and potential intervention in a clinical setting. This is especially true for potential prognostic factor research meant to influence a strategy for future treatment, for example, in studies examining what type of patients benefit from disc herniation surgery. In such cases, the study would ideally require a well-defined cohort at the same stage of the disease (first incidence of sciatica) receiving randomized treatment [158]. With this in mind, the study design in Paper I warrants some caution regarding the use of its potential prognostic factors in clinical decision-making. However, with limited resources, the design is ideal for stating overall rates with small confidence intervals, from multiple study centres and different surgeons.

In contrast, Paper II covered large number of different patient variables believed to be necessary to compensate for the heterogeneity that exists in such a diverse group of patients in different stages of their disease and with different underlying reasons for hospital admission. The paper had the advantage of testing potentially modifiable prognostic factors. Psychological characteristics and mental health in contrast to sociodemographic factors can be modified through specific interventions [159]. BMI, hypertension and smoking are also potentially modifiable factors that have previously been shown to be directly or indirectly associated with chronic pain conditions [141]. However, there is limited evidence that interventional smoking secession, weight loss and blood pressure control improve chronic pain outcomes. It is possible that these lifestyle factors may exhibit their effects through secondary prevention of vascular disease and general comorbidity [141].

Comorbidity as a prognostic factor was tested in Paper I, but would also have value if tested in Paper II since chronic pain conditions are shown to be associated with other chronic medical conditions [160]. There are several ways of testing and defining comorbidity either individually or through a series of measures. The Elixhauser Score and the Charlson Comorbidity Index (used in Paper I) are examples of the latter [139, 161]. The Charlson Comorbidity Index index is very useful for condensing the different comorbidities rather than trying to analyse and interpret the effect of multiple individual comorbidities. Whether to use such scoring systems as the CCI or test the different comorbidities separately is however debatable [162]. It is the lack of accuracy behind the weighting/scoring-values for each of the different comorbid conditions that has been criticized. This is especially valid in Paper I where the final outcome was not mortality,

and the study population was very different from that which the original CCI was developed. The grade of severity within a single comorbid condition runs the risk of being lost, but more importantly, there may be comorbid conditions that are correctly scored in regards to one-year mortality, but are incorrectly weighted in regards to surgical outcomes such as surgical complications and re-operations.

The magnitude of follow-up evaluations and the bulk of questionnaires that must be filled out by the patients and subsequently sent back to the researchers made the study design in Paper II vulnerable and ultimately the end results were potentially affected by “loss of follow-up”. In addition, the observational design with a non-randomized surgical treatment selection precluded us from comparing the outcomes of non-surgically and surgically treated patients. A randomized selection of surgery would have certainly given the study additional clinical value, possibly drawing causal inferences concerning treatment effects and/or directing surgical treatment based on our prognostic models. This would have been particularly compelling as non-surgically treated high-risk ÖMSPQ patients were found to have an especially high level of self-reported leg-pain at 12 months.

Furthermore, the final outcome in paper II (pain on a NRS scale) does not measure functional disability directly, nor does it measure other important life quality measures such as depression, anxiety, work absence or mobility. In light of this, we originally planned to include Oswestry Disability Index scores at 12 months as a secondary outcome measure, but chose not to publish these results in order to meet the demands of the editor’s and journal’s word count, especially since the results were much the same for the two outcomes.

Paper III can arguably be criticized for sharing many of the same patients as Paper II. More importantly, the wide inclusion criteria ran the risk of favoring no association between CPM and sciatic chronicity, as the study did not attempt to exclude patients with traits known to negatively impact CPM results such as long duration of pain (weeks), analgesic use, hypertension, previous lumbar surgery, previous episode of sciatica or having psychosocial traits known to be associated with pain chronicity [163]. One could further argue that with the limited previous research on CPM effect on chronic sciatica, a study population with the optimal conditions for observing an association would have been preferable in this novel stage of CPM testing in sciatica. With this argument in mind, the optimal sciatica patient would be one who had sciatic pain for only a short period of time (preferably tested even before developing the condition), non-surgically treated, not treated with analgesics or antidepressants prior to CPM testing, no previous history of pain condition and a prior minimal of psychosocial traits known to be associated with pain chronicity [115, 163]. However, the use of such a narrow inclusion criteria would considerably limit both study testing and future clinical use to a minimum of hospitalised patients. In addition, CPM testing was not performed in the acute stages of hospital admission but rather 6 weeks later. The patients may arguably have undergone both psychological and physical changes during the time from hospital admission to CPM testing that could have affected their CPM profile negatively and subsequently diminished any possible predictive properties by the time they were tested. Testing the patients upon admission may have identified CPM as a potent prognostic factor in sciatica. However, at the time of planning, it was considered more practical, accurate and economically feasible to test patients in an outpatient laboratory setting, rather than a crowded hospital ward where they were believed to be both stressed, medicated, immobile and facing a limited number of days before discharge. Thus the design was constructed in such a way that CPM testing could be implemented for all patients in an outpatient setting if the study were to reveal a positive result despite the wide inclusion criteria and late testing.

5.1.2 Internal validity

5.1.2.1 Random associations

Internal validity refers to the question of whether results from a study hold true for the study population, i.e. alternative explanations for its findings can be ruled out [164]. In prognostic factor research there is always a risk that observed associations have occurred at random and do not represent true effects. This is especially true when a large number of potential prognostic factors are tested within a small sample size. In Paper II, nearly 20 prognostic factors were tested with simple linear regression with the significance level set to 5%, indicating that it is likely that one statistical association among the prognostic factors and outcome could occur by chance. The problem of association at random in prognostic factor research stresses the importance of a predetermined rationale behind choosing to test a potential prognostic factor. This calls for further study replication and experimental validation before any definite conclusions can be made [164].

5.1.2.2 Selection bias

In contrast to random error, bias is a systematic error in the recruitment of participants, the measurements of their risk factors and outcomes, or reporting of their results. Observational studies are prone to three broad types of biases: *selection bias*, *information bias* and *confounding*. Selection bias is said to have occurred when the study sample is not representative of the intended population of the study as a result of the procedures used to select subjects and the factors that influence study participation [164]. Paper I runs the risk of selection bias in that only operations performed in the public sector were included in the study, whereas the generally healthier and less complicated patients found in the private sector were not included. This may induce falsely high complication rates and thus affect the prognostic models. One could further argue that there might also be differences between public and private hospitals with regards to the surgeons, operating theatres, and patient care. However, a previous Norwegian study has shown that, on a national basis, these differences appear to be minimal [124].

Attrition bias, a type of selection bias, occurs when the patients lost to follow-up systematically differ in some way from patients that complete the cohort study [165]. Most cohort studies assume that patients drop out of the study at random. However, studies have shown that this is often not the case [166, 167]. Paper II can be criticized for its high dropout rate of 28% and subsequent vulnerability to attrition bias [165]. The magnitude of follow-up evaluations and the bulk of questionnaires that had to be filled out by the patients and subsequently sent back to the researchers by mail made the study design vulnerable in this regard. There were more smokers and a generally lower level of education among the dropouts compared with the patients who completed the study: two factors known to influence pain perception negatively [100, 141]. There is a clear possibility that other such influencing factors were either overrepresented or underrepresented among the “dropouts”, but were undisclosed at baseline. Also the outcome itself could influence the patients’ willingness to complete the study. Patients suffering from sciatica at 12 months could be less likely to complete the 12-month questionnaire because of their reduced mobility, psychological factors related to chronic pain or distrust in the medical profession. Further, patients with complete regression of symptoms may also be associated with loss to follow-up, as these patients may have resumed normal daily activities and dealing with hospital and study related issues is no longer a priority as their sciatic pain no longer

has an active role in their daily lives. Even if one knew all the differences between the groups, statistical correction of attrition bias would nevertheless be difficult, especially given the studies' small sample size. It is therefore important to acknowledge that in studies such as Papers II and III, attrition bias should be prevented during the study rather than dealt with post hoc. However, with a larger sample size, sampling weights and tailored replenishment samples can help to compensate for the effects of attrition bias [168].

Non-response bias, a type of selection bias that occurs when patients decline to participate systematically, differs from patients who do want to participate in the study [169]. It is imaginable that non-response bias is a common threat to all studies involving pain testing, thus that 16% of the patients declined to participate in Paper III was not surprising. However, 14% of the patients in Paper II also declined to participate for various reasons. Again, it was statistically assumed that these patients were no different than the patients included in the study in relation to clinical features, sociodemographic characteristics and pain perception. However, it is more likely these patients have traits in common that deviate from those who decided to participate. One could argue that the results in Paper III only apply to sciatica patients that “do not mind being CPM (pain) tested” until one either has shown that typical non-participants do not systematically differ from patients willing to participate in experimental pain studies or one has mapped out any differences and adjusted the results accordingly. Non-response bias was fortunately not believed to be an issue with regards to Paper I, as registering surgical information was a prerequisite for reimbursement.

5.1.2.3 Information bias (misclassification)

Information bias, or misclassification, is said to occur when the estimated effect is inaccurate because of measurement error or due to misclassification of subjects for the exposure outcome variables [169]. Information bias serves as an umbrella term for the many ways in which measurements can be systematically distorted. It is important that it is distinguished from random error which is an inherent part of all measurements [164].

Medical doctors intentionally underreporting complications is a well-known example of information bias, and might be relevant in light of the low complication rates found in Paper I [62, 63]. However, there was no obvious incentive for the physicians not to register the complications. There may also be a difference in opinion among surgeons as to what surgical events constitute a complication, and whether or not to factor in the severity of the condition and the patient's general health, age and comorbidity. The potential variation in the coding practices, experience, and thoroughness of physicians could further affect the accuracy and completeness of the complication coding.

Surgical databases in which data is volunteered for the purpose of measuring and improving the quality of surgical care most often include specific outcomes such as pain and function scores, implant information and neurological findings. However, these surgical registries are limited in that they often include only surgical centres or surgeons that specifically make an effort to report into the registry [170]. In contrast, databases designed for hospital-administrative or reimbursement purposes (like the one in Paper I), often have a larger variability with regards to surgical centres and surgeons, but they do not give insight into functional outcomes, explain the severity of the complication or the underlying reason for a readmission or a reoperation. Most importantly, there is often poor reliability in the diagnostic coding in such registries [157]. It is very unlikely that Paper I is an exception. In addition, data associated with governmental

reimbursement is also vulnerable to being skewed in favor of maximal economic refunding as certain complications or diagnoses require greater compensation than others [171]. This is problematic since systematic and random errors related to coding are difficult to uncover and account for statistically, despite large sample sizes [171].

Underreporting of smoking habits and analgesic use are well known examples of information bias that might be relevant to Paper II. Further, it is not inconceivable that prior knowledge of the CPM paradigm among the participants could influence CPM test results in favour of experiencing a CPM effect. Systematic variation in patient expectations and prior knowledge of CPM effect is therefore an important source of potential information bias related to Paper III.

Patient information bias can either be dependent (differential) or independent (non-differential) of the exposure and the outcome variables [164]. Most cohort studies are vulnerable to the latter (non-differential), since information regarding the predictor of interest is collected prospectively in relation to the outcome. This will, in turn, lead to the level of misclassification to be the same among those experiencing the outcome and those not experiencing it. Non-differential information bias will, for the most part, lead to an underestimation of the association with outcome [164]. However, in Papers II and III, the outcome of interest (leg-pain) was measured at both the beginning and the end of the study, and it is conceivable that the initial pain scored at baseline had an effect on how the patients scored their pain at 12 months. In this way, the predictor and outcome measurements were not irrespective of each other and any information bias within pain reporting would be subject to *differential* misclassification. Differential misclassification may result in over or underestimation of the association between predictor and outcome, and it often affects the results more severely than non-differential misclassification [164]. Though few and insufficient, there are methods available for correcting information bias [172], but it is best to try to prevent it during the planning phase of data collection.

5.1.2.4 Confounding

In aetiological research, a confounder is a third factor that is associated with both the exposure and the outcome, but is not the causal pathway between them [164]. In order to find the causal relationship between exposure and outcome, the effects of the confounder must be controlled for. The classic example is the comparison between subjects with and without yellow nicotine staining on the hands as a predictor of lung cancer. A comparison would most likely reveal an association between yellowness on the fingers and lung cancer, only to be confounded by smoking. Though prognostic factors that are causal to the outcome will naturally have the strongest prognostic capacity, a prognostic factor's main function is to predict an outcome and can do so to some extent independent of causation. Because prognostic factors do not have to be causal, confounders can act as and are referred to as prognostic factors in prognostic factor research. In the example above, the nicotine staining has high value as a prognostic factor for lung cancer, but would be misleading in regards to causation of lung cancer. To look for nicotine staining on the patients hands as part of routine clinical examination is an example of a prognostic factor that has clinical importance, despite its lack of causality to the outcome. The use and willingness to accept such "black box" associations as causal, is unfortunately a growing problem and has been heavily criticized as a serious threat to the life of aetiological research epidemiology [173]. In this context, researchers have argued that over- and unnecessary adjustment of available factors can give misleading results [174]. Further, even if researchers

adjust for all known confounding factors, different distributions of unknown factors may still bias the results, a problem known as residual confounding [173]. In Papers I and III the final prognostic models were adjusted for age and sex by default, regardless of their impact on the model. One may argue that this was redundant, and that such an adjustment misleads the reader to believe that results imply causality, rather than a result of model building tradition as was the case in the present papers. However, by showing no responsibility with regards to the underlying causality, it is particularly important that the study population has an adequate sample size and that information and selection bias is severely limited.

5.1.2.5 Errors specific to regression analysis

The validity of the regression model is often dependent on the number and suitability of the factors tested [148]. Ideally all relevant factors should be included. In the case of Paper I, the type of surgical instruments used, the number of surgeons per operation, the surgeons' experience and the preoperative status of the patient would arguably add prognostic value to the model; without knowledge of these factors the model's clinical use is severely limited. Further, most regression models require that the prognostic factors examined in the model are independent of each other, and do not convey closely related information [148]. In such a case of "collinearity" between the prognostic factors, the logistic regression model will tend to outweigh the significance of these factors [148]. This problem relates especially to Paper I where both age and comorbidity were found to be associated with the outcome, and the two were also closely dependent on each other. Sociodemographic variables such as age, sex, income and education are also closely connected and warrant caution within this context. The issue of "interaction", which is the effect of one prognostic factor being influenced by the value of another predictor can further influence the validity and limit the use of a regression model [148].

Odds ratios (OR) were used to show the strength of association between the prognostic factors and the outcome in the logistic regression models [175]. The OR in logistic regression can be interpreted as whether someone with a certain trait is more (or less) likely than someone without that trait to experience the outcome of interest. Unfortunately, the OR provides no information about the absolute frequency or the probability of the outcome occurring [175]. Caution must therefore be exercised when using the odds ratios from studies where the overall frequency of the studied outcome is rare [175, 176]. The outcome in Paper I can be considered as such, where the high odds ratios poorly reflect the marginal differences in frequency of surgical events within the tested patient groups. The difference in complication rate in comorbid vs. non-comorbid patients was 6% (6% vs. 12%), yet the odds ratio of 1.6 signified that the odds of the negative outcome occurring were 60% higher amongst the comorbid patients than the non-comorbid patients. Intuitively, one can look at the odds ratio in isolation and wrongly conclude that "comorbid patients should not undergo lumbar disc surgery". However, if the general complication rate was higher (and the proportion of comorbid patients and the odds ratio remained the same), the above conclusion would have value.

The level of statistical significance was set to $p < 0.05$ for all models. When interpreting the regression models, it is important to acknowledge that the highly statistically significant p -values ($p < 0.001$) found in regression models do not automatically reflect the strength of the association between the predictor and outcome. It is therefore possible to have a statistically significant ($p < 0.001$) OR of only 1.1, indicating poor prognostic value despite its high level of significance. OR cannot be converted directly to relative risk, but the two are often mistakenly used interchangeably by unwary researchers or the news media, resulting in an overestimation of

the predictive impact of the prognostic factor of interest. This is especially true when the prevalence of the outcome is common and exceeds 10%. If the outcome is rare, Risk ratio and OR are nearly identical, which could arguably be the case in Paper I [175, 176].

In paper II, one could argue that more of the potential prognostic factors from the univariate analysis should have been added to the multiple regression models despite their lack of significance in the univariate testing. There is a strong argument that a full backward elimination approach (i.e. no initial univariate testing) beginning with all the candidate factors included in a univariate model before subsequently removing them, would be a better way to avoid overfitting the model and selection bias. This would arguably provide more correct standard errors and *p*-values [153]. Variable selection is further complicated when prognostic factors are statistically significant in the univariate level but show no statistical significance once put in a multivariate model. This is often the case when they are so strongly dependent upon the other factors in the model that they make no additional contribution towards explaining the outcome. However, which of these correlated factors end up being dismissed and which of them remain in the final multivariate model is partly dependant on the method of selection [150]. Therefore caution is warranted when adopting a strict *p*-value driven selection in the final stages of model building, as in Paper II. Further, there is an argument for retaining all the candidate factors in the final model as this could improve the overall predictive power of the model and reduce the danger of rejecting a true hypothesis as false [153]. On the other hand, a model with many intercorrelated non-significant factors runs the risk of multicollinearity.

With this in mind, the use of the ÖMSPQ as a prognostic factor is interesting, as it captures a wide range of pain related entities condensed into a single prognostic factor. Possible strong associations between the outcome variable (sciatic pain) and the specific entities (for example, fear avoidance) in the questionnaire may be masked by the remaining entities in the questionnaire because of the totality of questionnaire scoring. One could argue that the questions in the screening tool should have been tested individually rather than grouped together in a single questionnaire. This would also prevent potential statistical interactions and potential masking of results as some of the potential prognostic factors were tested both as independent variables and indirectly through the ÖMSPQ screening tool (pain distribution, intensity and longevity). The problem with this approach would be that testing the ÖMSPQ items individually would overflow the model with independent variables, and thus induce large statistical challenges with regards to sample size of the study.

In Paper III, the linear outcome had to be converted to binary outcome, and accordingly, logistic regression and not linear regression was used as the main statistical method. The action of forced conversion from a linear to a categorical outcome is vulnerable to type II error, especially if split into possibly unjustified, oversimplified, or clinically irrelevant categories [177]. Though multiple sensitivity analyses using different cut-off values were performed (with no different end result), the low cut-off value in Paper III ran the risk of overestimating the number of patients with a negative outcome, since suffering from leg-pain measured as 2 on a 0-10 NRS can arguably be considered a favourable outcome by many clinicians and researchers.

The difficult issue of selecting the appropriate cut-off values for regression models also applies for the prognostic factors collected as scores or in a continuum. In Paper I, age was not used on a continuous scale as in Papers II and III, but had to be dichotomised into 3 age groups to meet the logistic regression model assumption of a “constant magnitude of association across the range of values for that variable” [148]. For example, in examining the relationship between age and unfavourable surgical events, the association needed to be the same when comparing 30- and 40-year-olds as it were when comparing 70- and 80-year-olds if the prognostic factor was not to

be dichotomised into ranges that met the assumption requirements. With this in mind one could argue that the age groups in Paper I were too large and that there was no previous evidence to suggest that the age cut-offs represent either important milestones in general human aging or the aging/degradation of the lumbar discs specifically. Further, “education” was split into university vs. non-university education, with the purpose of showing an association between outcome and the possible health related protective properties of a university education. One could, however, argue that its prognostic value would most likely reside in the patients who were not able to complete secondary school, as their possibly poor socioeconomic capital would have a greater potential to be associated with a poor final outcome. An objection to the use of this potential cut-off would be that such an unevenly distributed categorization would have very little practical implication in a Norwegian patient population where most inhabitants complete secondary school.

There is an argument that cohorts with a proper sample size are fairly robust to violations of the normality assumption, and that outlying values, high leverage, heteroscedasticity, correlated errors, non-linearity and interactions pose a larger threat to the study’s validity [178]. Further, logistic regression commonly requires more data points per predictor than linear regression in order to achieve stable and meaningful results. Thus, the transition from a linear to a categorical outcome also affected statistical power in Paper III.

One could argue that sample sizes for both Papers II and III were in the lower range of what is preferred, especially since the patients were later stratified into surgical and non-surgical cohorts as part of each study’s sensitivity analysis. Ideally, prognostic studies require at least several hundred participants completing a study. In smaller studies, a higher number of prognostic factors tested per patient will increase the risk of overestimating the predictive performance of the model due to “overfitting” [153, 154]. Therefore, the low sample size in Paper III allowed only a few variables to be included in the prognostic model without losing statistical power. For this reason, many of the factors known to confound CPM results were not included in the prognostic model. Cohort studies aimed at aetiological research often require a high number of confounder variables and subsequently high number of participants in order to prove any causal effects. The relatively small sample size in Paper III was evaluated as sufficient to detect any crude prognostic value of CPM effect, but the sample size limited the study’s ability to adjust for multiple confounders. The small sample size also had the disadvantage that the CPM effect required a strong association to the outcome in order to be statistically significant, whereas a larger sample size might have detected a weak association between CPM and chronic pain. However, detecting such an association would have little practical prognostic value unless the increase in sample size and addition of confounders revealed an unforeseen dramatic change in the association between CPM and chronic sciatica.

5.1.3 External validity

Generalizability is the extent to which study results can be generalized to persons outside the original study population [164]. A prognostic model needs to be externally validated before practitioners can confidently apply a model to their patients. A prognostic model can only be claimed valuable if previously externally validated using independent data from a different location than the original development data [179]. This process requires unambiguous definitions of the prognostic factors and outcomes. How the model is derived is arguably of less importance if the model consequently performs well over different populations and settings [123]. However, a prerequisite for achieving external validity is that the results hold true for the original study

population (internal validity). In addition, impact studies should be conducted to evaluate effectiveness compared with usual care without the model [180].

During the data collection period, the incidence of lumbar disc surgery in Norway was 58/100,000 inhabitants compared to the 29 per 100,000 in Sweden and 46 per 100,000 in Denmark [181]. A study comparing back surgery rates in 11 mostly high-income countries found that the overall surgery rate was at least 40% higher in the USA than in any other country, and it was more than five times than in Great Britain [182]. Differences in health economic systems, population size and governance can arguably affect the incentive, threshold and, possibly, indication for surgery, and this may explain the large variations in operation rates even between high-income countries. These differences in surgical rates may, in turn, create international differences in surgical results as the populations differ by virtue of selection rather than the surgical procedure itself being performed differently. Caution is therefore warranted whenever transcribing surgical results from a single national study of any kind to an international level, especially if surgery rates vastly differ but the disease incidence is identical.

With this in mind, and despite the on-going attempt to streamline hospital procedures and surgical indication throughout the world [47, 183], it is more natural to discuss external validity in the context of the other Nordic countries, as their populations are relatively small and they have strong public health care systems that provide lumbar disc surgery on indication, free of charge. The surgeries presented in Paper I most likely varied to some degree with regards to surgical indication, pre- and post-operative work-up practice, surgical technique, equipment, training, experience, antibiotic use and anaesthesia [124]. This national variability arguably makes the study somewhat robust in relation to the same preferential differences that likely exist between and within the Nordic countries. In addition, the surgeries are considered to be performed in a relatively similar comportment by the fact that the surgical circles governing surgical etiquette are closely connected through language, culture and research. However, a previous study comparing lumbar disc surgery in Norway, Sweden and Denmark showed that preoperative disability, duration of symptoms and lifestyle factors such as smoking varied significantly between the Scandinavian countries [181]. Despite these differences in baseline characteristics, no significant differences between the Nordic countries have been found with regards to the patients' self-reported 1-year outcomes (disability/pain/and quality of life) [181]. In addition, the indication for surgery did not differ despite there being a difference in surgery rates among the countries. These similar findings indicate that the complication, readmission and reoperation rates are also somewhat similar.

It is also important to question how long the study results are valid in the context of the constant improvement in surgical techniques, hospital services and public health. With regard to these temporal changes, it is most important to evaluate whether or not significant changes to surgical practice have occurred since the study was conducted. If no such changes have taken place, there is most likely no great change in complication rates, and the study complication rates will have value despite being decades old. However, over time, multiple subtle changes in surgical indication, patient health and surgical performance can have such a large impact that previously used complication rates would be falsely high. This may explain why Paper I's complication rates were lower than earlier complication rates, despite there being no great change in surgical techniques or practice prior to the study which could offer an explanation. It is also important to acknowledge that prognostic models should be revisited over time with regards to the potential for model improvement as new biomarkers and other prognostic factors are developed or discovered [180].

It is common that prognostic models developed in secondary care are also intended for use in primary care. Unfortunately, this usually decreases the model's strength [180]. The sciatica patients recruited in Papers II and III were included upon hospital admission, thus generalizing any results from these studies to general practice or even a secondary care outpatient clinic warrants caution. In general, national and international differences in hospital availability, local hospital practices and health economic policies dictate to a large extent what treatment sciatica patients receive, and within what speciality they are treated (orthopaedic, neurosurgeon, rheumatologist, neurologist, general surgeon or medicinal specialist). Further, what leads to a hospital admission in Oslo may vary considerably from what is required for a hospital admission in smaller more rural hospitals where admission may require lengthy transport, surgical treatment options may be limited and bed capacity may be over or under what is required by the patient population. It is also conceivable that there are differences among physicians regarding the level of pain and neurologic symptoms required for hospital acceptance. Such perceptive differences among physicians coupled with resource availability warrants caution when generalizing hospitalized sciatica both nationally and internationally.

In Papers II and III the patients were recruited upon hospital admission, but the study itself had no influence over whether patients were admitted to hospital or not. The inclusion criteria only ensured that the patients were in fact suffering from painful sciatica. However, the criteria gave no instruction as to reasoning behind the hospital admission, or the underlying premise for why the patients needed hospital admission rather than primary or secondary outpatient care treatment. The wide selection criteria in the papers would therefore be easy to implement upon a reproduction study based in another hospital. However, unless the bed capacity, surgical treatment availability, and the hospital physicians were to have the same opinion on what constitutes a valid hospital admission for sciatica, the prognostic capacity of the models has the potential to deviate considerably upon study reproduction in another clinical setting. A narrowing of the inclusion criteria such as "only include patients strictly admitted for pain management", "only include patients admitted for possible surgery/surgical evaluation" or "only include patients admitted for examination of recent acquired paresis" may have strengthened the study population's generalizability at the expense of its inclusiveness.

5.2 Main results compared with other published studies

5.2.1 Unfavourable events in lumbar disc surgery

Comparable large data registers concerning surgical complications specifically related to lumbar disc surgery are unfortunately limited. In Paper I, we found that the complication rate (2.7%) was the same [184] or 2-4% lower than what preceding literature has stated. [50, 134, 185-188]. Rather than this reduction in unfavourable surgical events being a result of a single advance in one area of surgical care, the lower rates are likely the result of multiple small contributions in improved surgical technique, instrumentation, preoperative/postoperative care, hospital quality and improved general health (for example smoking cessation). This explanation is further strengthened by acknowledging the falling rates of complications seen in most surgical fields today. However, it is important to recognize that many of the rates referenced above have been drawn from cohort studies with smaller sample sizes and larger variability in their complication rates than the small confidence intervals presented in Paper I. Variability aside, registry datasets (Paper I) are vulnerable to showing falsely low surgical complication rates, as they most often do

not track complications treated outside the domain of the hospital administration. The patients in an intervention study, however, will often be followed up closely for complications as part of research protocol. The Spine Patient Outcomes Research Trial (SPORT) [50] with a slightly higher registered rate of complications (5%) is an example of such. However, one can argue that surgical centres that conduct intervention studies are more experienced and their patients are to a greater extent selected for successful surgery than would be the case for surgery performed outside a study trial setting. Thus, the complication rates reported from intervention studies are comparatively low, as may be the case in a randomized controlled trial from 2007 where only 1.6% of the patients experienced a complication [49]. Further, patients generally report more complications than surgeons do, which may explain why the rates for wound infection are reported as 1-2% higher in the Norwegian Registry for Spine Surgery (NORspine), a national Norwegian registry that includes PROMS and self-reported wound infection [189].

Reoperation rates as opposed to complication rates are more robust concerning such reporting biases. However, comparing studies in relation to reoperation rates is not without its difficulties. Reoperation rates can vary because of professional uncertainty about criteria or indications for repeat surgery, surgical complications, patient expectations and local practice philosophy and patterns [68]. The cut-off value for what is considered a reoperation varies in the literature from 90 days to 5 years. In accordance with previous findings, Paper I found that the 90 day reoperation rate (2.1%) was the same [68] or 2-4% lower than what earlier literature has stated [190]. Reoperations were also found to be associated with the rate of complications [68]. It is our understanding that a reoperation occurring shortly after the time of the original operation most often signifies a complication or faulty original operation. In contrast, revision surgery performed 6 months after the original operation is likely the result of a recurrence, a progression or a lack of expected improvement of the lumbar disc herniation – conditions that are estimated to occur in 5-15% of the patients treated with disc herniation surgery [98, 191]. Long-term reoperation rates may therefore also reflect the willingness of the surgeons to perform a reoperation and not necessarily the quality of the initial procedure [68]. It is therefore the author's opinion that one should have a greater appreciation for the distinction between a reoperation that occurs in close proximity to the original operation (<90 days) and a revision surgery 1-5 years later, since they are interchangeably referred to as reoperations both in the literature and clinical practice. Hence, due to this lack of distinction, the different underlying reasons behind a second surgery are somewhat distorted. Although both short-term and long-term repeat surgery rates may serve as a potential marker of the quality of surgical care [68], they may also elicit different targets for practice improvements and it is the author's opinion that the distinction between the two should be conveyed to the patient as part of preoperative consultation.

Though age and comorbidity have been shown to be associated with repeat surgery and complications in spine surgery [185, 192], Paper I reaffirms that the strength of this association was much the same when applied only to simple micro discectomy specifically. The literature suggests that comorbid patients are less likely to receive surgical treatment [193]. It is reasonable to assume that older and comorbid patients are carefully selected for surgery as lumbar discectomy is seldom performed on vital indication and there is often a choice between surgical and non-surgical care. It is therefore the author's belief that a randomization of treatment (conservative or surgical) could yield an even stronger association between age/comorbidity and unfavorable surgical events, since it is more than likely that only the "healthiest" of elderly and comorbid patients are advised/granted surgery. Therefore, caution is warranted when applying rates from non-randomized retrospective studies, such as that of Paper I, as the basis for a change in clinical practice, for example, implementing a lower threshold for operating on comorbid and

aged patients. However, if the results had shown a considerably higher number of unfavorable events among this specific patient group, there would be a better rationale for a practice change.

Though comparable updated literature on the impact of sociodemographic factors such as income, education and marital status on the rate of unfavorable surgical events is scarce, Paper I concluded that no such convincing associations exist. As these prognostic factors are non-modifiable and there is no obvious theoretical rationale as to why they should have a large impact on surgical performance other than indirectly affecting patient general health, it is the author's opinion that future replicative testing of these factors should be considered redundant. However, their role in prognostication of functional outcomes (pain and disability) and long-term-reoperation rates is unclear, and sociodemographic factors may play a larger role in outcomes with greater patient-subjective properties (pain and comorbidity). Previous studies have shown that women are subject to poorer surgical outcomes than men [194]. Despite this finding not being replicated in the present study, patient's sex as a prognostic factor in the context of failed surgery should not be entirely dismissed at the present time.

5.5.2 Persistent leg-pain

Paper II found that mean leg-pain measured 1 year after hospital discharge was generally low (2,5 on a 10 point NRS scale) for both patients that received surgical and non-surgical care. The rate of mean leg-pain as a function of time from hospital admission to 12 months post discharge had much the same graph configuration as leg-pain recordings found in other sciatica cohorts, both in specialised spine centres and in general practice [49, 50, 76, 131, 195], Fig.5 (page 32). The mean sciatic pain resided in the top 30% of the measurement scale, but rapidly dropped to the lower 30% of the pain scale within 6-12 weeks. The curves then remained stable without any large changes from this period onward. On average, the surgically treated patients had a higher level of pain at baseline compared to non-surgically treated patients and a subsequently larger and a more rapid drop in mean leg-pain from baseline to the second follow-up (6-12 weeks). Based on this universally recognized graph, it should be no surprise that Paper II found that leg-pain recorded 6 weeks post hospital discharge was found to be a predictor of leg-pain at 12 months. The fact that these graphs are similar across different sciatica cohorts also indicates that leg-pain measured at 6-8 weeks has potential clinical value.

Unfortunately, the disadvantage of measuring leg-pain 6 weeks after hospital discharge is that it has no predictive value in the immediate and early stages of the disease when prediction would have great potential for directing treatment or in some way changing current practice [56]. Therefore, the paper also presented a more complex model, which could be used upon hospital admission. The model found that using the ÖMSPQ tool in combination with registering leg-pain, patient employment status and whether or not they received surgical treatment gave the most accurate prognostication of outcome using the data available. However, in line with other similar models, applying clinical, sociodemographic and psychosocial patient characteristics produced only moderate prognostic value for long-term sciatica [73, 144]. Nevertheless, that a high risk score on the ÖMSPQ was found to be associated with persistent leg-pain must not be overlooked. There is evidence to support the association between psychosocial factors and LBP chronicity in general practice [138]. Our results indicated that this also applies to patients in hospital suffering from severe sciatica. The association between a high psychosocial risk profile and leg-pain is especially prominent in those patients who do not receive surgery, which again poses the question of whether some of these patients would have benefited from surgical intervention. The observational design of the study precluded us from answering this question, but the possibility of surgeons unfairly refraining from operating on patients with pronounced

psychosocial disadvantages must be considered. The true prognostic value of assessing psychosocial traits in a hospital setting has yet to be fully mapped. Not only is it likely that simply using the short 10-item version of the ÖMSPQ could yield the same prognostic value as was obtained in Paper II using the long 25-item version of the questionnaire, it is possible that selecting only some of the items in the questionnaire and using them as individual prognostic factors could have greater prognostic value than that obtained when using the entire questionnaire. It is conceivable that one (or more) of the items in the questionnaire could have a sizable prognostic impact on its own, but that its value was contaminated by the other 24 items dictating the questionnaire score. However, despite the possibility of diagnostic improvement through selecting more specific and sensitive psychosocial prognostic factors, Paper II confirmed previous suppositions that early and accurate prognostication in sciatica is difficult and the key to success does not lie solely in clinical examination, sociodemographic affiliation, lifestyle factors or a patient's psychosocial profile, nor in the combination these factors [73, 144].

Neurophysiological measures, such as electromyography, have previously been tested in sciatica without yielding any evident prognostic value [90]. Similarly, Paper III found no prognostic value in measuring the CPM effect in sciatica patients. This result was neither consistent nor inconsistent with the body of studies aimed at showing CPM as a biomarker of chronic pain, as roughly 70 % of the studies in a recent systematic review reported a negative result [196]. Although there is a strong argument that this inconsistency in results was due to large methodological differences between the studies, publishing bias may further underestimate the percentage of studies with nonsignificant correlations between CPM effect and pain. In addition, it is also possible that the mechanisms responsible for the transition from acute to chronic pain are specific to different pain conditions and that there exists no single universal mechanism that can explain all types of pain chronicity across the large spectre of pain related conditions. This may explain why studies have found CPM to be associated with conditions such as chronic widespread pain and irritable bowel syndrome [197, 198], but Paper III failed to show such an association in sciatica.

5.3 Clinical Implications

Paper I concluded that simple lumbar disc surgery was safe, based on having found a low rate of unfavourable events and no surgery related deaths. Because of the retrospective nature of the study and lack of functional outcomes, the study cannot be used to argue in favour of conducting more surgery as opposed to conservative care. However, its clinical value lies in its ability to convey the risks involved in surgery, especially in cases where indication for surgery is clear. Patients that would benefit from surgery, but are uncertain or unwilling to commit to surgery because of an unfounded fear of unfavourable events or as a result of being introduced to outdated high rates, may be reassured by the study results and be better able to give their informed consent. Patients are also apprised of the different types of complications and of their current rates. Individual rates presented for elderly and comorbid patients will give both the patient and clinician a better understanding of the increased risks involved when operating on these specific subgroups of patients. In addition, the present study indicates that age and comorbidity are essential building blocks for future prognostic models aimed at predicting unfavourable surgical outcomes. From a healthcare perspective, Paper I creates an opportunity for surgical centres and individual surgeons not only to compare rates, but to gain insight into the

secondary costs involved in lumbar disc surgery as a result of reoperations and readmissions. This information is important within the economic context of surgical vs. prolonged conservative care in the treatment of sciatica.

Paper II concluded that sciatic pain experienced 6 weeks after hospital discharge was associated with sciatic pain at 1 year. Although there is a risk that patients suffering non-pain-related issues will be forsaken or that the prognostic impact is too small and inaccurate, a follow-up telephone consultation rating patients' leg-pain 6 weeks after discharge could be a practical way of predicting long-term sciatic pain, as it would not be time-consuming, expensive or impound upon surgical resources. Patients that report persistent leg-pain at 6 weeks could be offered changes in their care, and as a consequence, possibly alter their long-term pain trajectory. Reconsideration for surgery, adjustments in analgesic use, scheduling of future follow-up consultations, or the offer of participating in clinical studies targeting patients with a prognosticated poor outcome are all measures that could be implemented if sciatic pain were to persist beyond 6 weeks. However, it is important to acknowledge that the study did not show that any of the measures suggested above would have an effect on the final outcome, and that the only real therapeutic value of the paper lies in that it has established a time point and measure with predictive capacity. Clinicians and spinal centres can, however, choose to use these findings to provide information about general prognosis and implement different therapeutic strategies and follow-up regimes for patients at risk of symptomatic persistence. Researchers will also have a prognostic tool that can be used to select target patients with a projected poor outcome for interventional studies. This is of particular importance if the studied intervention is expensive or involves any risk to the participants, since many sciatica patients have good outcomes with the current treatments, in many cases, additional treatment would be redundant.

Paper III concluded that CPM effect measured at 6 weeks did not predict sciatic pain at 1 year. Unfortunately, the negative result does not have the same clinical implications as if the results were positive, especially since at present there is no widespread clinical use of CPM testing on spinal patients. However, had we found CPM to be of prognostic value, caution would still have to be exercised if one were to implement such a prognostic factor into clinical practice without the results being replicated. Validating a prognostic model in other populations is almost always a prerequisite before any clinical use. This also holds true for the prognostic factors found in Papers I and II.

5.4 Future Perspectives

In Paper I, we presented a detailed rate of unfavourable events in simple lumbar disc surgery, but no information was available concerning how these events affected the patients in terms of pain and disability. Paper I also gave no account of the seriousness of the complications, costs or any possible explanation from the surgeon as to why the complication occurred. The same is true for the reoperation and readmission rates, thus we can only speculate on the underlying causes. These questions need answers if we are to fully appreciate the rates of unfavourable surgical events presented in future research. Large international multicentre cohort studies and databases with homogeneous design and methods would be ideal to answer these questions.

There are several ongoing Scandinavian spine surgery registers that would be ideal in relation to future mapping of the different; 1) causes, 2) the type of patients and surgeons involved, and 3) the patient implications, of surgery with unfavourable events. The Norwegian

Registry for Spine Surgery (NorSpine), DaneSpine (Denmark) and SweSpine (Sweden) [181] comprises lumbar disc surgeries with a relatively high completeness rate both in the public and private sector. The three registries have methodologically comparable parameters, recording patient demographics, lifestyle factors and patient-reported outcome measures (pain and ODI) prior to surgery as well as 12 months post-surgery for thousands of spine surgeries annually. In addition, the surgeons record preoperative diagnosis, clinical presentation, patient comorbidity and different surgical parameters such as operation time, antibiotic use, type of procedure (micro or standard open) and anatomical level. It is worth noting that the original plan when designing paper I was to merge the original administrative data with clinical data from the NorSpine register. However, due to privacy considerations this was not made possible.

In the future, these registries or similar ones would be ideal for a more in-depth description of the type of patients involved, the surgery itself and the ultimate clinical consequence (long-term pain and disability) in the case of an unfavourable event. Further, it is possible to elicit the prognostic impact of future key factors such as the patient's clinical presentation, surgical indication, experience of the surgeon, surgeon workload and hours, surgical techniques and instrumentation, detailed stratification of MRI findings alongside patient age and comorbidity. These potential prognostic factors elicited through this type of registered data could potentially be tested later in clinical studies concerning their use in patient selection for surgery. In addition, national surgical databases that have relatively similar structures could be combined in order to create even larger international databases ideal for conducting pattern recognition and machine learning using artificial intelligence. This could identify patients with a specific clinical, sociodemographic, radiological and lifestyle profile at risk for unfavourable events or the opposite, a high likelihood of surgical success, aiding clinical decision-making. However, it is the author's opinion that these surgical registries would benefit greatly in having a comparable registry available for patients who were conservatively treated both in primary and secondary care. In this way, specific patient profiles that either are at risk of poor outcome or success upon surgery could be compared to the same profile receiving non-surgical care. Further, this would be compelling information in relation to prolonged non-surgical treatment for patients that have persistent sciatic pain beyond 4-12 months. This group has an especially high rate of poor outcome [54], and may arguably be in greater risk of other types of complications such as gastric, liver, kidney and dependency conditions, because of prolonged use of different types of analgesics.

The Danish "SpineData" [199] is one of few large scale clinical registries that encompasses conservative care sciatica patients. It could possibly be available for a merger with surgical registries for future comparison and patient pattern recognition. However, a more holistic future register that encompasses both surgical and nonsurgical patients from their initial stages of care, their clinical presentation, treatment and their long-term outcome, as well as surgical information detailing reasons for surgery, complications and reoperations would be preferable. Such a registry, reflecting clinical presentation and outcome as well as rationale for treatment (conservative/surgery) in addition to any complication of treatment would allow us to better compare the treatment modalities, in addition to identifying specific subgroups of patients that ultimately have a high or low risk of unfavourable events or poor clinical outcome (pain and disability) depending or regardless of treatment modality.

Since lumbar disc surgery is usually not performed on an absolute indication, it is important to consider the rates of unfavourable events relative to health gains achieved through conservative care. Future randomised control trials comparing surgical and non-surgical care will indirectly put the complication and reoperation rates in perspective, in that high complication and

reoperation rates will become less acceptable the closer the treatment modalities are. Future collaboration between Scandinavian and possibly other European countries could be essential in order to achieve a robust sample size for subgroup analysis and definite conclusions.

It is argued that the spinal literature has been exhausted in the repeated attempts to identify lifestyle, psychosocial and work tendencies as reasons for why people suffer from chronic back pain conditions, and that future emphasis should shift towards genetic and basic research if there is to be a breakthrough finding. Not only will any biological or genetic causative mechanism give additional prognostic value, but this could ultimately form the basis for new and better personalized treatments. With this in mind, future prognostic studies exploring possible biomarkers would bring new life to previous prognostic studies, as new and old prognostic factors combined could establish even stronger models for prognostication.

Paper III was an attempt to explore an underlying pathophysiological basal mechanism in a clinical context. The study required professional experience outside of what a normal spinal clinician could offer. The process was time consuming, expensive and yielded a negative result, all common attributes of basal and investigative research. However, spinal researchers should not be overly anxious or hesitant in using discoveries and advances within basic pain research and implementing them in clinical research. Though a future study replication might be redundant, the results from Paper III should not discourage future studies of CPM in sciatica, since paper III left us wondering whether CPM should have been tested in the infant stages of sciatica rather than several weeks into the condition. A study exploring the predictive properties of CPM among first time presenters with sciatica in general practice or directly upon hospital admission may find CPM to have prognostic value. Further, CPM may also aid future decisions related to tailoring medical treatment as well as giving us a better understanding of the mechanisms underlying pain chronicity in general.

6. CONCLUSIONS

Sciatica is a common complaint causing temporary and persistent pain worldwide. At present, the mechanisms that cause sciatic pain to persist rather than follow the favourable trajectory of pain regression after prolonged conservative care or lumbar disc surgery are not sufficiently understood. The studies covered in this thesis aim to aid clinicians in predicting these unfortunate outcomes, and at the same time provide patients with information concerning their long-term prognosis, and, if electing to be treated with lumbar disc surgery, make clear the rate at which they may expect an unfavourable surgical event.

The first paper in this thesis revealed that fewer than 7% of 34 639 lumbar disc surgeries had at least one unfavourable event, such as a complication (2.7%), a reoperation (2.1%), and/or a hospital readmission (2.4%). There were no mortalities associated with surgery. Whether this rate of unfavourable events is acceptable must be considered in relation to the health gains achieved by choosing surgical over conservative care. This paper suggests that surgical complications are less frequent than previously suggested and that patients can be informed that lumbar disc surgery is safe. However, caution must be exercised when surgery is performed on patients of advanced age or with comorbidity, as these factors were shown to be associated with a higher complication rate. In the future, mapping of unfavourable events should be performed using registries that not only capture a patient's preoperative sociodemographic profile and comorbidity, but also detail clinical, radiologic, and intra operative entities, in addition to a subsequent patient related outcome. Though the administrative dataset used in Paper I lacked many of these clinical data, the dataset's value in comparison to other more specific spinal surgery registers lies in that the dataset encompassed all public patients that received lumbar disc surgery and their respective surgeons. The surgeons or patients had no option to not register the surgery or respective readmissions.

In the second and third papers, we found that leg-pain in patients hospitalized with sciatica as their main complaint regressed rapidly during the first 6 weeks after discharge, but that their pain remained comparatively static after this time point regardless of treatment (surgical/non-surgical). Results from our prognostic model indicate that simply measuring leg-pain 6 weeks after hospital discharge is an efficient way of establishing prognosis, regardless of whether the patient received surgical or non-surgical treatment. A patient's psychosocial profile was also found to have some predictive value, and should therefore always be included in future prognostic modelling within the field of sciatica. However, there is still a large void of unknown prognostic factors that need to be addressed in order for us to reach the goal of early prognostication in patients hospitalised with sciatica. This warrants a continued search for a better understanding of the biological mechanisms involved in refractory sciatica as sociodemographic properties, clinical examination and lifestyle factors were found have limited prognostic value with regards to the persistence of sciatic pain.

In an attempt to explore the more biological mechanisms involved in pain chronicity Paper III was based on the hypothesis that patients with an inert low CPM-effect would ultimately have a subsequent poor long-term recovery from their sciatica. The study failed to show such an effect. With this in mind, future study designs must consider the heterogeneity that exists in sciatica cohorts with regards to symptom debut, clinical presentation and treatment received, as the prognostic value of biologic factors such as the CPM-effect may be negatively impacted by such entities. Thus, future CPM studies on sciatica patients should be conducted at an early stage of disease, preferentially before any treatment has been implemented.

REFERENCES

1. Wu, A., et al., *Global low back pain prevalence and years lived with disability from 1990 to 2017: estimates from the Global Burden of Disease Study 2017*. Ann Transl Med, 2020. **8**(6): p. 299.
2. Maher, C., M. Underwood, and R. Buchbinder, *Non-specific low back pain*. Lancet, 2017. **389**(10070): p. 736-747.
3. Koes, B.W., M.W. van Tulder, and W.C. Peul, *Diagnosis and treatment of sciatica*. Bmj, 2007. **334**(7607): p. 1313-7.
4. Hartvigsen, L., et al., *Leg pain location and neurological signs relate to outcomes in primary care patients with low back pain*. BMC Musculoskelet Disord, 2017. **18**(1): p. 133.
5. Jensen, R.K., et al., *Diagnosis and treatment of sciatica*. Bmj, 2019. **367**: p. 16273.
6. Valat, J.P., et al., *Sciatica*. Best Pract Res Clin Rheumatol, 2010. **24**(2): p. 241-52.
7. Konstantinou, K. and K.M. Dunn, *Sciatica: review of epidemiological studies and prevalence estimates*. Spine (Phila Pa 1976), 2008. **33**(22): p. 2464-72.
8. Konstantinou, K., et al., *The impact of low back-related leg pain on outcomes as compared with low back pain alone: a systematic review of the literature*. Clin J Pain, 2013. **29**(7): p. 644-54.
9. Jensen, O.K., C.V. Nielsen, and K. Stengaard-Pedersen, *One-year prognosis in sick-listed low back pain patients with and without radiculopathy. Prognostic factors influencing pain and disability*. Spine J, 2010. **10**(8): p. 659-75.
10. Lambeek, L.C., et al., *The trend in total cost of back pain in The Netherlands in the period 2002 to 2007*. Spine (Phila Pa 1976), 2011. **36**(13): p. 1050-8.
11. van Tulder, M.W., B.W. Koes, and L.M. Bouter, *A cost-of-illness study of back pain in The Netherlands*. Pain, 1995. **62**(2): p. 233-40.
12. Dagenais, S., J. Caro, and S. Haldeman, *A systematic review of low back pain cost of illness studies in the United States and internationally*. Spine J, 2008. **8**(1): p. 8-20.
13. Fardon, D.F., et al., *Lumbar disc nomenclature: version 2.0: Recommendations of the combined task forces of the North American Spine Society, the American Society of Spine Radiology and the American Society of Neuroradiology*. Spine J, 2014. **14**(11): p. 2525-45.
14. ADAMS, M.A. and W.C. HUTTON, *Prolapsed Intervertebral Disc: A Hyperflexion Injury*. 1982. **7**(3): p. 184-191.
15. Vergroesen, P.P., et al., *Mechanics and biology in intervertebral disc degeneration: a vicious circle*. Osteoarthritis Cartilage, 2015. **23**(7): p. 1057-70.
16. Kepler, C.K., et al., *The molecular basis of intervertebral disc degeneration*. Spine J, 2013. **13**(3): p. 318-30.
17. Adams, M.A. and P.J. Roughley, *What is intervertebral disc degeneration, and what causes it?* Spine (Phila Pa 1976), 2006. **31**(18): p. 2151-61.
18. Boos, N., et al., *Classification of age-related changes in lumbar intervertebral discs: 2002 Volvo Award in basic science*. Spine (Phila Pa 1976), 2002. **27**(23): p. 2631-44.
19. Nerlich, A.G., et al., *Immunomorphological analysis of RAGE receptor expression and NF-kappaB activation in tissue samples from normal and degenerated intervertebral discs of various ages*. Ann N Y Acad Sci, 2007. **1096**: p. 239-48.
20. Mulleman, D., et al., *Pathophysiology of disk-related sciatica. I.--Evidence supporting a chemical component*. Joint Bone Spine, 2006. **73**(2): p. 151-8.
21. Omarker, K. and R.R. Myers, *Pathogenesis of sciatic pain: role of herniated nucleus pulposus and deformation of spinal nerve root and dorsal root ganglion*. Pain, 1998. **78**(2): p. 99-105.

22. Takahashi, N., et al., *Pathomechanisms of nerve root injury caused by disc herniation: an experimental study of mechanical compression and chemical irritation*. Spine (Phila Pa 1976), 2003. **28**(5): p. 435-41.
23. Binch, A.L., et al., *Expression and regulation of neurotrophic and angiogenic factors during human intervertebral disc degeneration*. Arthritis Res Ther, 2014. **16**(5): p. 416.
24. Lee, J.M., et al., *Interleukin-1 β induces angiogenesis and innervation in human intervertebral disc degeneration*. J Orthop Res, 2011. **29**(2): p. 265-9.
25. Kim, J.H., et al., *Activated macrophage-like THP-1 cells modulate anulus fibrosus cell production of inflammatory mediators in response to cytokines*. Spine (Phila Pa 1976), 2008. **33**(21): p. 2253-9.
26. Autio, R.A., et al., *Determinants of spontaneous resorption of intervertebral disc herniations*. Spine (Phila Pa 1976), 2006. **31**(11): p. 1247-52.
27. Brinjikji, W., et al., *Systematic literature review of imaging features of spinal degeneration in asymptomatic populations*. AJNR Am J Neuroradiol, 2015. **36**(4): p. 811-6.
28. Jensen, M.C., et al., *Magnetic resonance imaging of the lumbar spine in people without back pain*. N Engl J Med, 1994. **331**(2): p. 69-73.
29. Boden, S.D., et al., *Abnormal magnetic-resonance scans of the lumbar spine in asymptomatic subjects. A prospective investigation*. J Bone Joint Surg Am, 1990. **72**(3): p. 403-8.
30. Deyo, R.A. and S.K. Mirza, *CLINICAL PRACTICE. Herniated Lumbar Intervertebral Disk*. N Engl J Med, 2016. **374**(18): p. 1763-72.
31. *Low back pain and sciatica in over 16s: assessment and management. NICE guideline [NG59]*
Published date: 30 November 2016 Last updated: 11 December 2020
<https://www.nice.org.uk/guidance/NG59/chapter/Recommendations#non-invasive-treatments-for-low-back-pain-and-sciatica>.
32. Vroomen, P.C., et al., *Lack of effectiveness of bed rest for sciatica*. N Engl J Med, 1999. **340**(6): p. 418-23.
33. Stochkendahl, M.J., et al., *National Clinical Guidelines for non-surgical treatment of patients with recent onset low back pain or lumbar radiculopathy*. Eur Spine J, 2018. **27**(1): p. 60-75.
34. Luijsterburg, P.A., et al., *Physical therapy plus general practitioners' care versus general practitioners' care alone for sciatica: a randomised clinical trial with a 12-month follow-up*. Eur Spine J, 2008. **17**(4): p. 509-17.
35. Paatelma, M., et al., *Orthopaedic manual therapy, McKenzie method or advice only for low back pain in working adults: a randomized controlled trial with one year follow-up*. J Rehabil Med, 2008. **40**(10): p. 858-63.
36. Albert, H.B. and C. Manniche, *The efficacy of systematic active conservative treatment for patients with severe sciatica: a single-blind, randomized, clinical, controlled trial*. Spine (Phila Pa 1976), 2012. **37**(7): p. 531-42.
37. Ye, C., et al., *Comparison of lumbar spine stabilization exercise versus general exercise in young male patients with lumbar disc herniation after 1 year of follow-up*. Int J Clin Exp Med, 2015. **8**(6): p. 9869-75.
38. Bakhtiary, A.H., Z. Safavi-Farokhi, and A. Rezasoltani, *Lumbar stabilizing exercises improve activities of daily living in patients with lumbar disc herniation*. Journal of Back and Musculoskeletal Rehabilitation, 2005. **18**: p. 55-60.
39. Huber, J., et al., *The effect of early isometric exercises on clinical and neurophysiological parameters in patients with sciatica: An interventional randomized single-blinded study*. Isokinetics and Exercise Science, 2011. **19**: p. 207-214.

40. Konstantinou, K., et al., *Stratified care versus usual care for management of patients presenting with sciatica in primary care (SCOPIC): a randomised controlled trial*. *Lancet Rheumatol*, 2020. **2**(7): p. e401-e411.
41. Pinto, R.Z., et al., *Drugs for relief of pain in patients with sciatica: systematic review and meta-analysis*. *Bmj*, 2012. **344**: p. e497.
42. Pinto, R.Z., A.J.H. Verwoerd, and B.W. Koes, *Which pain medications are effective for sciatica (radicular leg pain)?* *Bmj*, 2017. **359**: p. j4248.
43. Rasmussen-Barr, E., et al., *Nonsteroidal Anti-inflammatory Drugs for Sciatica: An Updated Cochrane Review*. *Spine (Phila Pa 1976)*, 2017. **42**(8): p. 586-594.
44. Gibson, J.N. and G. Waddell, *Surgical interventions for lumbar disc prolapse: updated Cochrane Review*. *Spine (Phila Pa 1976)*, 2007. **32**(16): p. 1735-47.
45. Rasouli, M.R., et al., *Minimally invasive discectomy versus microdiscectomy/open discectomy for symptomatic lumbar disc herniation*. *Cochrane Database Syst Rev*, 2014(9): p. Cd010328.
46. Du Bois, M., M. Szpalski, and P. Donceel, *A decade's experience in lumbar spine surgery in Belgium: sickness fund beneficiaries, 2000-2009*. *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society*, 2012. **21**(12): p. 2693-2703.
47. Gadjradj, P.S., et al., *Management of Symptomatic Lumbar Disk Herniation: An International Perspective*. *Spine (Phila Pa 1976)*, 2017. **42**(23): p. 1826-1834.
48. Jacobs, W.C., et al., *Surgery versus conservative management of sciatica due to a lumbar herniated disc: a systematic review*. *Eur Spine J*, 2011. **20**(4): p. 513-22.
49. Peul, W.C., et al., *Surgery versus prolonged conservative treatment for sciatica*. *N Engl J Med*, 2007. **356**(22): p. 2245-56.
50. Weinstein, J.N., et al., *Surgical vs nonoperative treatment for lumbar disk herniation: the Spine Patient Outcomes Research Trial (SPORT): a randomized trial*. *Jama*, 2006. **296**(20): p. 2441-50.
51. Buttermann, G.R., *Treatment of lumbar disc herniation: epidural steroid injection compared with discectomy. A prospective, randomized study*. *J Bone Joint Surg Am*, 2004. **86**(4): p. 670-9.
52. Osterman, H., et al., *Effectiveness of microdiscectomy for lumbar disc herniation: a randomized controlled trial with 2 years of follow-up*. *Spine (Phila Pa 1976)*, 2006. **31**(21): p. 2409-14.
53. Weber, H., *Lumbar disc herniation. A controlled, prospective study with ten years of observation*. *Spine (Phila Pa 1976)*, 1983. **8**(2): p. 131-40.
54. Bailey, C.S., et al., *Surgery versus Conservative Care for Persistent Sciatica Lasting 4 to 12 Months*. *N Engl J Med*, 2020. **382**(12): p. 1093-1102.
55. Deyo, R.A. and J.N. Weinstein, *Low back pain*. *N Engl J Med*, 2001. **344**(5): p. 363-70.
56. Moons, K.G., et al., *Prognosis and prognostic research: what, why, and how?* *Bmj*, 2009. **338**: p. b375.
57. Lebude, B., et al., *Defining "complications" in spine surgery: neurosurgery and orthopedic spine surgeons' survey*. *J Spinal Disord Tech*, 2010. **23**(8): p. 493-500.
58. Sanford, D.E., et al., *Variations in definition and method of retrieval of complications influence outcomes statistics after pancreatoduodenectomy: comparison of NSQIP with non-NSQIP methods*. *J Am Coll Surg*, 2014. **219**(3): p. 407-15.
59. Dindo, D., N. Demartines, and P.A. Clavien, *Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey*. *Ann Surg*, 2004. **240**(2): p. 205-13.
60. Clavien, P.A., et al., *The Clavien-Dindo classification of surgical complications: five-year experience*. *Ann Surg*, 2009. **250**(2): p. 187-96.
61. Tevis, S.E. and G.D. Kennedy, *Postoperative complications and implications on patient-centered outcomes*. *J Surg Res*, 2013. **181**(1): p. 106-13.

62. Mannion, A.F., et al., *The assessment of complications after spine surgery: time for a paradigm shift?* Spine J, 2013. **13**(6): p. 615-24.
63. Ohrn, A., et al., *Reporting of sentinel events in Swedish hospitals: a comparison of severe adverse events reported by patients and providers.* Jt Comm J Qual Patient Saf, 2011. **37**(11): p. 495-501.
64. Pinto, A., et al., *Surgical complications and their impact on patients' psychosocial well-being: a systematic review and meta-analysis.* BMJ Open, 2016. **6**(2): p. e007224.
65. Healey, M.A., et al., *Complications in surgical patients.* Arch Surg, 2002. **137**(5): p. 611-7; discussion 617-8.
66. Weiser, T.G., et al., *An estimation of the global volume of surgery: a modelling strategy based on available data.* Lancet, 2008. **372**(9633): p. 139-44.
67. Woodfield, J., et al., *Patient reporting of complications after surgery: what impact does documenting postoperative problems from the perspective of the patient using telephone interview and postal questionnaires have on the identification of complications after surgery?* BMJ Open, 2019. **9**(7): p. e028561.
68. Martin, B.I., et al., *Repeat surgery after lumbar decompression for herniated disc: the quality implications of hospital and surgeon variation.* Spine J, 2012. **12**(2): p. 89-97.
69. Nykvist, F., et al., *Severe sciatica: a 13-year follow-up of 342 patients.* Eur Spine J, 1995. **4**(6): p. 335-8.
70. Patrick, D.L., et al., *Assessing health-related quality of life in patients with sciatica.* Spine (Phila Pa 1976), 1995. **20**(17): p. 1899-908; discussion 1909.
71. Atlas, S.J., et al., *The Maine-Seattle back questionnaire: a 12-item disability questionnaire for evaluating patients with lumbar sciatica or stenosis: results of a derivation and validation cohort analysis.* Spine (Phila Pa 1976), 2003. **28**(16): p. 1869-76.
72. Fairbank, J.C. and P.B. Pynsent, *The Oswestry Disability Index.* Spine (Phila Pa 1976), 2000. **25**(22): p. 2940-52; discussion 2952.
73. Ashworth, J., K. Konstantinou, and K.M. Dunn, *Prognostic factors in non-surgically treated sciatica: a systematic review.* BMC Musculoskelet Disord, 2011. **12**: p. 208.
74. Weber, H., I. Holme, and E. Amlie, *The natural course of acute sciatica with nerve root symptoms in a double-blind placebo-controlled trial evaluating the effect of piroxicam.* Spine (Phila Pa 1976), 1993. **18**(11): p. 1433-8.
75. Vroomen, P.C., et al., *Conservative treatment of sciatica: a systematic review.* J Spinal Disord, 2000. **13**(6): p. 463-9.
76. Konstantinou, K., et al., *Prognosis of sciatica and back-related leg pain in primary care: the ATLAS cohort.* Spine J, 2018. **18**(6): p. 1030-1040.
77. Waddell, G., *The Back pain revolution, 2nd ed.* 2004, Churchill Livingstone: New York.
78. Engel, G.L., *The need for a new medical model: a challenge for biomedicine.* Science, 1977. **196**(4286): p. 129-36.
79. Linton, S.J. and W.S. Shaw, *Impact of psychological factors in the experience of pain.* Phys Ther, 2011. **91**(5): p. 700-11.
80. Grotle, M., et al., *Clinical course and prognostic factors in acute low back pain: patients consulting primary care for the first time.* Spine (Phila Pa 1976), 2005. **30**(8): p. 976-82.
81. Linton, S.J., *Do psychological factors increase the risk for back pain in the general population in both a cross-sectional and prospective analysis?* Eur J Pain, 2005. **9**(4): p. 355-61.
82. Pincus, T., et al., *Fear avoidance and prognosis in back pain: a systematic review and synthesis of current evidence.* Arthritis Rheum, 2006. **54**(12): p. 3999-4010.
83. Vlaeyen, J.W. and S.J. Linton, *Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art.* Pain, 2000. **85**(3): p. 317-32.

84. Waddell, G., et al., *A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability*. Pain, 1993. **52**(2): p. 157-68.
85. Burton, A.K., et al., *Psychosocial predictors of outcome in acute and subchronic low back trouble*. Spine (Phila Pa 1976), 1995. **20**(6): p. 722-8.
86. Crombez, G., et al., *Pain-related fear is more disabling than pain itself: evidence on the role of pain-related fear in chronic back pain disability*. Pain, 1999. **80**(1-2): p. 329-39.
87. den Boer, J.J., et al., *A systematic review of bio-psychosocial risk factors for an unfavourable outcome after lumbar disc surgery*. Eur Spine J, 2006. **15**(5): p. 527-36.
88. Haugen, A.J., et al., *Prognostic factors for non-success in patients with sciatica and disc herniation*. BMC Musculoskelet Disord, 2012. **13**: p. 183.
89. Selim, A.J., et al., *The importance of radiating leg pain in assessing health outcomes among patients with low back pain. Results from the Veterans Health Study*. Spine (Phila Pa 1976), 1998. **23**(4): p. 470-4.
90. Balagué, F., et al., *Recovery of severe sciatica*. Spine (Phila Pa 1976), 1999. **24**(23): p. 2516-24.
91. Jungen, M.J., et al., *Inflammatory biomarkers in patients with sciatica: a systematic review*. BMC Musculoskelet Disord, 2019. **20**(1): p. 156.
92. Bjorland, S., et al., *Genes associated with persistent lumbar radicular pain; a systematic review*. BMC Musculoskelet Disord, 2016. **17**(1): p. 500.
93. Balagué, F., et al., *The potential value of blood biomarkers of intervertebral disk metabolism in the follow-up of patients with sciatica*. Eur Spine J, 2006. **15**(5): p. 627-33.
94. Haaland, A.K., et al., *Fibrinolytic activity as a predictor of the outcome of prolapsed intervertebral lumbar disc surgery with reference to background variables: results of a prospective cohort study*. Spine (Phila Pa 1976), 1992. **17**(9): p. 1022-7.
95. Vroomen, P.C., M.C. de Krom, and J.A. Knottnerus, *Predicting the outcome of sciatica at short-term follow-up*. Br J Gen Pract, 2002. **52**(475): p. 119-23.
96. Rihn, J.A., et al., *Duration of symptoms resulting from lumbar disc herniation: effect on treatment outcomes: analysis of the Spine Patient Outcomes Research Trial (SPORT)*. J Bone Joint Surg Am, 2011. **93**(20): p. 1906-14.
97. Carragee, E.J. and D.H. Kim, *A prospective analysis of magnetic resonance imaging findings in patients with sciatica and lumbar disc herniation. Correlation of outcomes with disc fragment and canal morphology*. Spine (Phila Pa 1976), 1997. **22**(14): p. 1650-60.
98. Carragee, E.J., et al., *Clinical outcomes after lumbar discectomy for sciatica: the effects of fragment type and anular competence*. J Bone Joint Surg Am, 2003. **85**(1): p. 102-8.
99. El Barzouhi, A., et al., *Prognostic value of magnetic resonance imaging findings in patients with sciatica*. J Neurosurg Spine, 2016. **24**(6): p. 978-85.
100. Peul, W.C., et al., *Influence of gender and other prognostic factors on outcome of sciatica*. Pain, 2008. **138**(1): p. 180-91.
101. Waddell, G., *1987 Volvo award in clinical sciences. A new clinical model for the treatment of low-back pain*. Spine (Phila Pa 1976), 1987. **12**(7): p. 632-44.
102. Yarnitsky, D., *Role of endogenous pain modulation in chronic pain mechanisms and treatment*. Pain, 2015. **156** Suppl 1: p. S24-31.
103. Le Bars, D., et al., *Diffuse noxious inhibitory controls (DNIC) in animals and in man*. Patol Fiziol Eksp Ter, 1992(4): p. 55-65.
104. Le Bars, D., A.H. Dickenson, and J.M. Besson, *Diffuse noxious inhibitory controls (DNIC). I. Effects on dorsal horn convergent neurones in the rat*. Pain, 1979. **6**(3): p. 283-304.
105. Moont, R., et al., *'Pain inhibits pain' mechanisms: Is pain modulation simply due to distraction?* Pain, 2010. **150**(1): p. 113-120.

106. Mlekusch, S., et al., *Conditioned Pain Modulation in Patients With Acute and Chronic Low Back Pain*. Clin J Pain, 2016. **32**(2): p. 116-21.
107. Lewis, G.N., D.A. Rice, and P.J. McNair, *Conditioned pain modulation in populations with chronic pain: a systematic review and meta-analysis*. J Pain, 2012. **13**(10): p. 936-44.
108. Koltyn, K.F., et al., *Mechanisms of exercise-induced hypoalgesia*. J Pain, 2014. **15**(12): p. 1294-1304.
109. Lemley, K.J., S.K. Hunter, and M.K. Bement, *Conditioned pain modulation predicts exercise-induced hypoalgesia in healthy adults*. Med Sci Sports Exerc, 2015. **47**(1): p. 176-84.
110. Vaegter, H.B., G. Handberg, and T. Graven-Nielsen, *Similarities between exercise-induced hypoalgesia and conditioned pain modulation in humans*. 2014. **155**(1): p. 158-167.
111. Liebano, R.E., et al., *Transcutaneous electrical nerve stimulation and conditioned pain modulation influence the perception of pain in humans*. Eur J Pain, 2013. **17**(10): p. 1539-46.
112. Dailey, D.L., et al., *Transcutaneous electrical nerve stimulation reduces pain, fatigue and hyperalgesia while restoring central inhibition in primary fibromyalgia*. Pain, 2013. **154**(11): p. 2554-2562.
113. Mlekusch, S., et al., *Do central hypersensitivity and altered pain modulation predict the course of chronic low back and neck pain?* Clin J Pain, 2013. **29**(8): p. 673-80.
114. LeResche, L., et al., *Psychophysical tests as predictors of back pain chronicity in primary care*. J Pain, 2013. **14**(12): p. 1663-70.
115. Yarnitsky, D., et al., *Recommendations on practice of conditioned pain modulation (CPM) testing*. Eur J Pain, 2015. **19**(6): p. 805-6.
116. Pud, D., Y. Granovsky, and D. Yarnitsky, *The methodology of experimentally induced diffuse noxious inhibitory control (DNIC)-like effect in humans*. Pain, 2009. **144**(1-2): p. 16-9.
117. Oono, Y., et al., *The inter- and intra-individual variance in descending pain modulation evoked by different conditioning stimuli in healthy men*. Scand J Pain, 2011. **2**(4): p. 162-169.
118. van Wijk, G. and D.S. Veldhuijzen, *Perspective on diffuse noxious inhibitory controls as a model of endogenous pain modulation in clinical pain syndromes*. J Pain, 2010. **11**(5): p. 408-19.
119. Granot, M., et al., *Contact heat-evoked temporal summation: tonic versus repetitive-phasic stimulation*. Pain, 2006. **122**(3): p. 295-305.
120. Hemingway, H., et al., *Prognosis research strategy (PROGRESS) 1: a framework for researching clinical outcomes*. Bmj, 2013. **346**: p. e5595.
121. Hingorani, A.D., et al., *Prognosis research strategy (PROGRESS) 4: stratified medicine research*. Bmj, 2013. **346**: p. e5793.
122. Riley, R.D., et al., *Prognosis Research Strategy (PROGRESS) 2: prognostic factor research*. PLoS Med, 2013. **10**(2): p. e1001380.
123. Steyerberg, E.W., et al., *Prognosis Research Strategy (PROGRESS) 3: prognostic model research*. PLoS Med, 2013. **10**(2): p. e1001381.
124. Grotle, M., et al., *Public and private health service in Norway: a comparison of patient characteristics and surgery criteria for patients with nerve root affections due to discus herniation*. Eur Spine J, 2014. **23**(9): p. 1984-91.
125. Sherman, J., et al., *Economic impact of improving outcomes of lumbar discectomy*. Spine J, 2010. **10**(2): p. 108-16.
126. Osterman, H., et al., *Risk of multiple reoperations after lumbar discectomy: a population-based study*. Spine (Phila Pa 1976), 2003. **28**(6): p. 621-7.
127. Jansson, K.A., et al., *Surgery for herniation of a lumbar disc in Sweden between 1987 and 1999. An analysis of 27,576 operations*. J Bone Joint Surg Br, 2004. **86**(6): p. 841-7.

128. Kim, C.H., et al., *The Long-term Reoperation Rate Following Surgery for Lumbar Herniated Intervertebral Disc Disease: A Nationwide Sample Cohort Study With a 10-year Follow-up*. Spine (Phila Pa 1976), 2019. **44**(19): p. 1382-1389.
129. Brennan, T.A., et al., *Incidence of adverse events and negligence in hospitalized patients. Results of the Harvard Medical Practice Study I*. N Engl J Med, 1991. **324**(6): p. 370-6.
130. Grøvle, L., et al., *The bothersomeness of sciatica: patients' self-report of paresthesia, weakness and leg pain*. Eur Spine J, 2010. **19**(2): p. 263-9.
131. Haugen, A.J., et al., *Estimates of success in patients with sciatica due to lumbar disc herniation depend upon outcome measure*. Eur Spine J, 2011. **20**(10): p. 1669-75.
132. Elliott, A.M., et al., *The epidemiology of chronic pain in the community*. Lancet, 1999. **354**(9186): p. 1248-52.
133. Greenspan, J.D., et al., *Studying sex and gender differences in pain and analgesia: a consensus report*. Pain, 2007. **132 Suppl 1**(Suppl 1): p. S26-45.
134. Deyo, R.A., et al., *Morbidity and mortality in association with operations on the lumbar spine. The influence of age, diagnosis, and procedure*. J Bone Joint Surg Am, 1992. **74**(4): p. 536-43.
135. Von Korff, M., L. Le Resche, and S.F. Dworkin, *First onset of common pain symptoms: a prospective study of depression as a risk factor*. Pain, 1993. **55**(2): p. 251-8.
136. Boersma, K. and S.J. Linton, *Expectancy, fear and pain in the prediction of chronic pain and disability: a prospective analysis*. Eur J Pain, 2006. **10**(6): p. 551-7.
137. Linton, S.J. and K. Halldén, *Can we screen for problematic back pain? A screening questionnaire for predicting outcome in acute and subacute back pain*. Clin J Pain, 1998. **14**(3): p. 209-15.
138. Hockings, R.L., J.H. McAuley, and C.G. Maher, *A systematic review of the predictive ability of the Orebro Musculoskeletal Pain Questionnaire*. Spine (Phila Pa 1976), 2008. **33**(15): p. E494-500.
139. Charlson, M.E., et al., *A new method of classifying prognostic comorbidity in longitudinal studies: development and validation*. J Chronic Dis, 1987. **40**(5): p. 373-83.
140. Lehmann, T.R. and H.S. LaRocca, *Repeat lumbar surgery. A review of patients with failure from previous lumbar surgery treated by spinal canal exploration and lumbar spinal fusion*. Spine (Phila Pa 1976), 1981. **6**(6): p. 615-9.
141. van Hecke, O., N. Torrance, and B.H. Smith, *Chronic pain epidemiology - where do lifestyle factors fit in?* Br J Pain, 2013. **7**(4): p. 209-17.
142. Hanley, M.A., et al., *Preamputation pain and acute pain predict chronic pain after lower extremity amputation*. J Pain, 2007. **8**(2): p. 102-9.
143. Elliott, A.M., et al., *The course of chronic pain in the community: results of a 4-year follow-up study*. Pain, 2002. **99**(1-2): p. 299-307.
144. Verwoerd, A.J., et al., *Systematic review of prognostic factors predicting outcome in non-surgically treated patients with sciatica*. Eur J Pain, 2013. **17**(8): p. 1126-37.
145. Woertgen, C., et al., *Variability of outcome after lumbar disc surgery*. Spine (Phila Pa 1976), 1999. **24**(8): p. 807-11.
146. Asch, H.L., et al., *Prospective multiple outcomes study of outpatient lumbar microdiscectomy: should 75 to 80% success rates be the norm?* J Neurosurg, 2002. **96**(1 Suppl): p. 34-44.
147. Kennedy, D.L., et al., *Reliability of conditioned pain modulation: a systematic review*. Pain, 2016. **157**(11): p. 2410-2419.
148. Tolles, J. and W.J. Meurer, *Logistic Regression: Relating Patient Characteristics to Outcomes*. Jama, 2016. **316**(5): p. 533-4.
149. Pallant, J., *SPSS Survival Manual. 5th edition*. 2013: Open University press.
150. Schneider, A., G. Hommel, and M. Blettner, *Linear regression analysis: part 14 of a series on evaluation of scientific publications*. Dtsch Arztebl Int, 2010. **107**(44): p. 776-82.

151. Sun, G.W., T.L. Shook, and G.L. Kay, *Inappropriate use of bivariable analysis to screen risk factors for use in multivariable analysis*. J Clin Epidemiol, 1996. **49**(8): p. 907-16.
152. Harrell, F.E., Jr., K.L. Lee, and D.B. Mark, *Multivariable prognostic models: issues in developing models, evaluating assumptions and adequacy, and measuring and reducing errors*. Stat Med, 1996. **15**(4): p. 361-87.
153. Royston, P., et al., *Prognosis and prognostic research: Developing a prognostic model*. Bmj, 2009. **338**: p. b604.
154. Riley, R.D., et al., *Calculating the sample size required for developing a clinical prediction model*. Bmj, 2020. **368**: p. m441.
155. Nojima, M., M. Tokunaga, and F. Nagamura, *Quantitative investigation of inappropriate regression model construction and the importance of medical statistics experts in observational medical research: a cross-sectional study*. BMJ Open, 2018. **8**(5): p. e021129.
156. Puri, K.S., et al., *Declaration of Helsinki, 2008: implications for stakeholders in research*. J Postgrad Med, 2009. **55**(2): p. 131-4.
157. Deyo, R.A., et al., *Analysis of automated administrative and survey databases to study patterns and outcomes of care*. Spine (Phila Pa 1976), 1994. **19**(18 Suppl): p. 2083s-2091s.
158. Altman, D.G., *Systematic reviews of evaluations of prognostic variables*. Bmj, 2001. **323**(7306): p. 224-8.
159. Kroenke, K., et al., *Optimized antidepressant therapy and pain self-management in primary care patients with depression and musculoskeletal pain: a randomized controlled trial*. Jama, 2009. **301**(20): p. 2099-110.
160. Barnett, K., et al., *Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study*. Lancet, 2012. **380**(9836): p. 37-43.
161. van Walraven, C., et al., *A modification of the Elixhauser comorbidity measures into a point system for hospital death using administrative data*. Med Care, 2009. **47**(6): p. 626-33.
162. Austin, S.R., et al., *Why Summary Comorbidity Measures Such As the Charlson Comorbidity Index and Elixhauser Score Work*. Med Care, 2015. **53**(9): p. e65-72.
163. Hermans, L., et al., *Inventory of Personal Factors Influencing Conditioned Pain Modulation in Healthy People: A Systematic Literature Review*. Pain Pract, 2016. **16**(6): p. 758-69.
164. Rothman, K.J., S. Greenland, and T.L. Lash, *Modern Epidemiology*. 2015: Wolters Kluwer Health/Lippincott Williams & Wilkins.
165. Kristman, V., M. Manno, and P. Côté, *Loss to follow-up in cohort studies: how much is too much?* Eur J Epidemiol, 2004. **19**(8): p. 751-60.
166. Gades, N.M., et al., *Dropout in a longitudinal, cohort study of urologic disease in community men*. BMC Med Res Methodol, 2006. **6**: p. 58.
167. Howe, C.J., et al., *Selection Bias Due to Loss to Follow Up in Cohort Studies*. Epidemiology, 2016. **27**(1): p. 91-7.
168. Nunan, D., J. Aronson, and C. Bankhead, *Catalogue of bias: attrition bias*. BMJ Evid Based Med, 2018. **23**(1): p. 21-22.
169. Sedgwick, P., *Bias in observational study designs: prospective cohort studies*. Bmj, 2014. **349**: p. g7731.
170. Lønne, G., et al., *Lumbar spinal stenosis: comparison of surgical practice variation and clinical outcome in three national spine registries*. Spine J, 2019. **19**(1): p. 41-49.
171. Sebastian, A.S., *National Database Research in Spine Surgery: Limitations in the Current Literature*. Clin Spine Surg, 2017. **30**(1): p. 27-29.
172. Lash, T.L., et al., *Good practices for quantitative bias analysis*. Int J Epidemiol, 2014. **43**(6): p. 1969-85.

173. Smith, G.D., *Reflections on the limitations to epidemiology*. J Clin Epidemiol, 2001. **54**(4): p. 325-31.
174. Schisterman, E.F., S.R. Cole, and R.W. Platt, *Overadjustment bias and unnecessary adjustment in epidemiologic studies*. Epidemiology, 2009. **20**(4): p. 488-95.
175. Norton, E.C., B.E. Dowd, and M.L. Maciejewski, *Odds Ratios-Current Best Practice and Use*. Jama, 2018. **320**(1): p. 84-85.
176. Cummings, P., *The relative merits of risk ratios and odds ratios*. Arch Pediatr Adolesc Med, 2009. **163**(5): p. 438-45.
177. Streiner, D.L., *Breaking up is hard to do: the heartbreak of dichotomizing continuous data*. Can J Psychiatry, 2002. **47**(3): p. 262-6.
178. Schmidt, A.F. and C. Finan, *Linear regression and the normality assumption*. J Clin Epidemiol, 2018. **98**: p. 146-151.
179. Altman, D.G., et al., *Prognosis and prognostic research: validating a prognostic model*. Bmj, 2009. **338**: p. b605.
180. Moons, K.G., et al., *Prognosis and prognostic research: application and impact of prognostic models in clinical practice*. Bmj, 2009. **338**: p. b606.
181. Lagerbäck, T., et al., *Effectiveness of surgery for sciatica with disc herniation is not substantially affected by differences in surgical incidences among three countries: results from the Danish, Swedish and Norwegian spine registries*. Eur Spine J, 2019. **28**(11): p. 2562-2571.
182. Cherkin, D.C., et al., *An international comparison of back surgery rates*. Spine (Phila Pa 1976), 1994. **19**(11): p. 1201-6.
183. Bath, M., T. Bashford, and J.E. Fitzgerald, *What is 'global surgery'? Defining the multidisciplinary interface between surgery, anaesthesia and public health*. BMJ Glob Health, 2019. **4**(5): p. e001808.
184. Peul, W.C., et al., *Prolonged conservative care versus early surgery in patients with sciatica caused by lumbar disc herniation: two year results of a randomised controlled trial*. Bmj, 2008. **336**(7657): p. 1355-8.
185. Strömqvist, F., et al., *Surgical treatment of lumbar disc herniation in different ages-evaluation of 11,237 patients*. Spine J, 2017. **17**(11): p. 1577-1585.
186. Stadler, J.A., 3rd, et al., *Complications associated with posterior approaches in minimally invasive spine decompression*. Neurosurg Clin N Am, 2014. **25**(2): p. 233-45.
187. Nandyala, S.V., et al., *Complications after lumbar spine surgery between teaching and nonteaching hospitals*. Spine (Phila Pa 1976), 2014. **39**(5): p. 417-23.
188. Støttrup, C.C., et al., *Increasing reoperation rates and inferior outcome with prolonged symptom duration in lumbar disc herniation surgery - a prospective cohort study*. Spine J, 2019. **19**(9): p. 1463-1469.
189. *Norwegian Registry for Spine Surgery (NORspine) September 2019*
<https://www.kvalitetsregistre.no/registers/530/resultater>.
190. Kim, C.H., et al., *Reoperation rate after surgery for lumbar herniated intervertebral disc disease: nationwide cohort study*. Spine (Phila Pa 1976), 2013. **38**(7): p. 581-90.
191. Tanavalee, C., et al., *A comparison between repeat discectomy versus fusion for the treatment of recurrent lumbar disc herniation: Systematic review and meta-analysis*. J Clin Neurosci, 2019. **66**: p. 202-208.
192. Campbell, P.G., et al., *Patient comorbidity score predicting the incidence of perioperative complications: assessing the impact of comorbidities on complications in spine surgery*. J Neurosurg Spine, 2012. **16**(1): p. 37-43.
193. Lurie, J.D., et al., *Surgical versus nonoperative treatment for lumbar disc herniation: eight-year results for the spine patient outcomes research trial*. Spine (Phila Pa 1976), 2014. **39**(1): p. 3-16.

194. Graver, V., et al., *Seven-year clinical follow-up after lumbar disc surgery: results and predictors of outcome*. Br J Neurosurg, 1999. **13**(2): p. 178-84.
195. Weinstein, J.N., et al., *Surgical vs nonoperative treatment for lumbar disk herniation: the Spine Patient Outcomes Research Trial (SPORT) observational cohort*. Jama, 2006. **296**(20): p. 2451-9.
196. Fernandes, C., et al., *Conditioned pain modulation as a biomarker of chronic pain: a systematic review of its concurrent validity*. Pain, 2019. **160**(12): p. 2679-2690.
197. Marcuzzi, A., et al., *Conditioned Pain Modulation (CPM) is Reduced in Irritable Bowel Syndrome: A Systematic Review and Meta-Analysis of CPM and the Role of Psychological Factors*. J Clin Gastroenterol, 2019. **53**(6): p. 399-408.
198. Gerhardts, A., et al., *Conditioned pain modulation in patients with nonspecific chronic back pain with chronic local pain, chronic widespread pain, and fibromyalgia*. Pain, 2017. **158**(3): p. 430-439.
199. Kent, P., et al., *SpineData - a Danish clinical registry of people with chronic back pain*. Clin Epidemiol, 2015. **7**: p. 369-80.

PAPERS I-III

■ SPINE

Complications, reoperations, readmissions, and length of hospital stay in 34 639 surgical cases of lumbar disc herniation

O. R. Fjeld,
L. Grøvre,
J. Helgeland,
M. C. Småstuen,
T. K. Solberg,
J-A. Zwart,
M. Grotle

From Oslo University
Hospital, Oslo,
Norway

■ O. R. Fjeld, MD, PhD Student,
Doctor in Specialization
(Neurology), Research and
Communication Unit for
Musculoskeletal Health, Oslo
University Hospital, Oslo, Norway;
Department of Neurology,
Oslo University Hospital, Oslo,
Norway; Institute of Clinical
Medicine, Faculty of Medicine,
University of Oslo, Norway.

■ L. Grøvre, MD, PhD,
Rheumatologist, Department of
Rheumatology, Østfold Hospital
Trust, Grålum, Norway.

■ J. Helgeland, MSc, Head of
Research, Division for Health
Services, Norwegian Institute of
Public Health, Oslo, Norway.

■ M. Grotle, PT, PhD, Professor
■ M. C. Småstuen, PhD,
Associate Professor
Research and Communication
Unit for Musculoskeletal Health,
Oslo University Hospital, Oslo,
Norway; Faculty of Health
Science, OsloMet – Oslo
Metropolitan University, Oslo,
Norway.

■ T. K. Solberg, MD,
PhD, Associate Professor,
Neurosurgeon, Department of
Neurosurgery, University Hospital
of Northern Norway, Tromsø,
Norway; Institute of Clinical
Medicine, The Arctic University
(UiT) of Norway, Tromsø,
Norway; Norwegian Registry
for Spine Surgery (NORspine),
University Hospital of Northern
Norway, Tromsø, Norway.

■ J-A. Zwart, MD, PhD,
Professor, Neurologist, Institute
of Clinical Medicine, Faculty of
Medicine, University of Oslo,
Oslo, Norway; Division of Clinical
Neuroscience, Oslo University
Hospital, Oslo, Norway.

Correspondence should be sent
to O. R. Fjeld; email: fjeldo@tcd.ie

©2019 Author(s) et al
doi:10.1302/0301-620X.101B4.
BJJ-2018-1184.R1 \$2.00

Bone Joint J
2019;101-B:470–477.

Aims

The aims of this study were to determine the rates of surgical complications, reoperations, and readmissions following herniated lumbar disc surgery, and to investigate the impact of sociodemographic factors and comorbidity on the rate of such unfavourable events.

Patients and Methods

This was a longitudinal observation study. Data from herniated lumbar disc operations were retrieved from a large medical database using a combination of procedure and diagnosis codes from all public hospitals in Norway from 1999 to 2013. The impact of age, gender, geographical affiliation, education, civil status, income, and comorbidity on unfavourable events were analyzed by logistic regression.

Results

Of 34 639 operations, 2.7% (95% confidence interval (CI) 2.6 to 2.9) had a surgical complication, 2.1% (95% CI 2.0 to 2.3) had repeat surgery within 90 days, 2.4% (95% CI 2.2 to 2.5) had a non-surgical readmission within 90 days, and 6.7% (95% CI 6.4 to 6.9) experienced at least one of these unfavourable events. Unfavourable events were found to be associated with advanced age and comorbidity.

Conclusion

The results suggest that surgical complications are less frequent than previously suggested. There are limited associations between sociodemographic patient characteristics and unfavourable events.

Cite this article: *Bone Joint J* 2019;101-B:470–477.

Surgery for lumbar disc herniation is a well-recognized and effective treatment in properly selected patients.¹ Measuring rates of complications, reoperations, readmissions, and length of hospital stay contributes important information regarding the quality and costs of surgical and hospital care.^{2,3} This information is valuable, not only in relation to general economic issues, cost-effectiveness, and the framing of clinical guidelines on a population level, but also for conversations between patients and clinicians when evaluating treatment options. The complication and reoperation rates may influence both physicians' recommendations and patients' final decisions, thus these rates need to be kept updated, accessible, and patient-personalized.^{4,5}

Although the length of hospital stay has been decreasing steadily despite the trend of an increasingly old surgical population,^{6,7} there are few reports of time trends relating to complication and reoperation rates. Large national medical

databases offer an opportunity to research rare events, such as complications and reoperations, over a long-time period within a large sample of operative procedures.⁸

The primary objective of this study was to determine the rate of unfavourable events such as surgical complications, reoperations, and readmissions for patients undergoing surgery for lumbar disc herniation, over a 15-year period from 1999 to 2013. The secondary objective was to investigate the impact of age, gender, comorbidity, education, civil status, income, and regional health authority affiliation on the rate of such events.

Patients and Methods

Design. This was a longitudinal observation study using data from a national medical database and a national population register from 1999 to 2013.

Data source. Hospital administrative data were retrieved from a national database at the Norwegian Knowledge Centre for Health Services for

the years between 1999 and 2009, and from the National Patient Registry (NPR) for the years between 2010 to 2013. A detailed description of the methods employed in data collection is published elsewhere.⁹ The data set contained medical and surgical procedures performed, final medical diagnosis at discharge, and patient comorbidity according to the Charlson Comorbidity Index.¹⁰ The complications at discharge were coded by the physicians according to the International Classification of Diseases (ICD-10 Norwegian version)¹¹ for each hospital admission. It also included data on the date and time of hospital admission and discharge, as well as the name(s) of the hospital(s) involved. A unique national registration number connects Norwegian residents to a hospital stay and to the sociodemographic data found in the National Population Register and other databases, such as Statistics Norway. These data sources cover the whole Norwegian population and provide information on age, gender, civil status, income, education level, and residence for each individual patient. The merging of data from the hospitals and the National Population Register was performed using a unique 11-digit personal identifier assigned to each Norwegian citizen.

Norway, with a population of 4.9 million, has a large, government-funded national healthcare system divided and run by four regional health authorities: the South-Eastern, Western, Central, and Northern Health Regions. Prior to 2007, these public regional health authorities were responsible for almost all spinal surgery in Norway, but from 2007 the public sector was only responsible for an estimated 80% of lumbar disc surgery, due to the increase of private surgery.^{12,13}

Study sample. The inclusion criteria for this study was patients older than or equal to 18 years of age discharged from Norwegian public hospitals following surgery for a herniated lumbar disc between 1 January 1999 and 31 December 2013. A detailed description of the patient selection and coding clarification is presented in Figure 1. Data were retrieved through identification of the relevant surgical procedure codes in the NPR, which employs the Norwegian version of the Nordic Medico-Statistical Committee (NOMESCO) Classification of Surgical Procedures (NCSP),¹⁴ a surgical coding system used in most Nordic countries. To ensure that the selected NCSP codes were specific for lumbar disc herniation surgery and not for other lumbar disorders (e.g. spinal stenosis) only operations with an ICD-10 diagnosis code (Norwegian version) indicating disc herniation were included. Operations for cancer, trauma, spinal infection, ankylosing spondylitis, and vertebral fractures were excluded from the study.

Once the relevant herniated lumbar disc surgery was identified, each patient's subsequent hospital admissions were retrieved, including surgical complications and follow-up surgical procedures. To make sure that all the patients could be followed for potential readmissions, reoperations, and complications for at least 90 days, initial operations performed in the time period between 1 September 2013 and 31 December 2013 were omitted from the analysis, as were operations performed in the first 90 days of the study period. All relevant surgical procedures with an added NSCP code indicating a more complex procedure (such as fusions and disc prosthesis) were omitted from analysis (Fig. 1). The raw data set contained seven

endoscopic procedures (NCSP code: ABC 07), which were not included in the study.

Definition of outcomes. Surgical complication: each patient was given one of the following ICD-10 complication diagnoses either upon initial surgical admission or upon readmission within 90 days: T81.0-T81.9, K91, N99, I97.8-9.

Reoperation: patients reoperated on within 90 days of initial surgery (immediate reoperation included).¹⁵ The reoperations were found using the same NSCP codes as the study inclusion; however, additional NSCP codes indicating a more complex procedure (such as fusions and disc prosthesis) were included.

Non-surgical readmission: patients readmitted within 90 days of lumbar surgery with an ICD-10 diagnosis of disc herniation, without reoperation. Such readmissions are usually due to intractable postoperative neurogenic pain, need of a repeat MRI, lack of expected clinical improvement, or other complications.

Unfavourable event: surgery with a surgical complication, reoperation, or a non-surgical readmission. The rate of unfavourable events represents the overall rate of the patient experiencing a surgical complication, reoperation, or readmission for surgery. Each patient may experience more than one of the above, thus the sum of the three individual rates is higher than the rate of a single unfavourable event.

Prolonged hospitalization: a patient's total hospital stay exceeding six days from day of admission to hospital discharge. This cut-off was based on the authors' clinical experience and has been used previously discussed.¹⁶ A hospital stay exceeding six days is judged highly indicative of issues such as intractable postoperative pain, other complications, need of a repeat MRI, poor general condition, or lack of expected improvement.

Statistical analysis and missing data. All data analyses were performed using SPSS version 24.00 (IBM Corp., Armonk, New York). The significance level was set to 5%. Continuous data were described with median and interquartile range (IQR) or mean and range. The 95% confidence intervals (CI) are shown for all estimates. Associations between pairs of categorical data were analyzed using the chi-squared test. Multivariate logistic regression analysis was used to investigate the association between the independent variables (age, gender, comorbidity, education, civil status, income, and regional health authority affiliation) and the dependent variable (unfavourable event: surgical complication, reoperation, or a non-surgical readmission). All independent variables were entered simultaneously, and all significant associations were described with odds ratios (OR) and 95% CI. The data set contains almost no missing data ($n < 20$), except for the sociodemographic data concerning income (19% missing) and education (8% missing). Therefore, no missing values were imputed.

Ethics. The study was approved by the Norwegian Data Inspectorate (2014/14413) and the Norwegian Regional Ethics Committee (2013/1662, REC south-east D). Individual patient consent was not required.

Results

Between 1999 and 2013, there were 34 639 herniated lumbar disc operations and 735 reoperations. Of the initial operations, 31 155 (90%) were coded as surgical discectomies and 3484 (10%) as surgical decompressions, for a total of

Fig. 1

Flow diagram of patient selection (NCSP, Nordic Medico-Statistical Committee (NOMESCO) Classification of Surgical Procedures; ICD, International Classification of Diseases).

Table I. Rates for hospital duration, complications, repeat surgery, and readmissions, according to the patient's age, gender, civil status, income, and year and region of operation

Variable	Procedures, n	Median hospital duration, days (IQR)	Prolonged hospitalization ≥ 7 days, % (95% CI)	Surgical complication, % (95% CI)	Repeat surgery ≤ 90 days, % (95% CI)	Readmission, no surgery ≤ 90 days, % (95% CI)*	Unfavourable event, % (95% CI) [†]
Total	34 639	4.1 (2.3 to 6.3)	31.7 (31.2 to 32.2)	2.7 (2.6 to 2.9)	2.1 (2.0 to 2.3)	2.4 (2.2 to 2.5)	6.7 (6.4 to 6.9)
Age, yrs							
18 to 29	3336	3.7 (2.2 to 5.8)	25.4 (24.0 to 26.9)	1.7 (1.2 to 2.1)	1.4 (1.0 to 1.8)	2.1 (1.6 to 2.6)	4.9 (4.2 to 5.7)
30 to 39	9362	4.1 (2.3 to 6.1)	28.8 (27.9 to 29.8)	2.4 (2.1 to 2.7)	2.0 (1.7 to 2.3)	2.3 (2.0 to 2.6)	6.1 (5.6 to 6.7)
40 to 49	10 456	4.1 (2.3 to 6.2)	31.2 (30.3 to 32.1)	2.7 (2.4 to 3.0)	2.6 (2.3 to 2.9)	2.5 (2.2 to 2.8)	7.4 (6.9 to 7.9)
50 to 59	6702	4.1 (2.4 to 6.7)	32.8 (31.7 to 33.9)	2.8 (2.4 to 3.2)	2.0 (1.7 to 2.4)	2.3 (2.0 to 2.7)	6.5 (5.9 to 7.1)
60 to 69	3311	4.2 (2.3 to 7.1)	35.0 (33.4 to 36.7)	3.8 (3.2 to 4.5)	1.9 (1.4 to 2.3)	2.5 (2.1 to 3.0)	7.6 (6.7 to 8.5)
70 to 79	1212	6.1 (3.2 to 10.1)	52.1 (49.3 to 55.0)	4.8 (3.6 to 6.0)	2.2 (1.3 to 3.0)	2.2 (1.4 to 3.1)	8.3 (6.8 to 9.9)
≥ 80	260	7.2 (4.2 to 13.1)	63.1 (57.2 to 69.0)	6.2 (3.2 to 9.1)	1.2 (0.0 to 3.0)	2.3 (0.5 to 4.1)	8.1 (4.7 to 11.4)
Gender							
Male	19 623	4.0 (2.2 to 6.1)	28.9 (28.3 to 29.6)	2.6 (2.4 to 2.9)	2.0 (1.8 to 2.2)	2.4 (2.2 to 2.6)	6.5 (6.2 to 6.9)
Female	15 016	4.2 (2.8 to 7.1)	35.2 (34.5 to 36.0)	2.8 (2.6 to 3.1)	2.3 (2.0 to 2.5)	2.3 (2.1 to 2.6)	6.9 (6.5 to 7.3)
CCI							
0	33 328	4.1 (2.3 to 6.2)	31.2 (30.8 to 31.7)	2.6 (2.5 to 2.8)	2.1 (1.9 to 2.2)	2.3 (2.1 to 2.4)	6.5 (6.2 to 6.7)
1 to 2	1191	5.0 (3.1 to 8.2)	40.9 (38.1 to 43.7)	5.1 (3.9 to 6.4)	3.3 (2.3 to 4.3)	4.5 (3.3 to 5.6)	11.3 (9.5 to 13.4)
≥ 3	120	6.1 (3.2 to 10.3)	54.2 (45.1 to 63.2)	1.7 (0.0 to 4.0)	3.3 (0.1 to 6.6)	6.7 (2.1 to 11.2)	11.7 (5.8 to 17.5)
Education[‡]							
Basic	24 781	4.1 (2.3 to 6.3)	31.8 (31.2 to 32.4)	2.8 (2.6 to 3.0)	2.1 (1.9 to 2.3)	2.5 (2.3 to 2.7)	6.8 (6.5 to 7.1)
Higher	7019	4.0 (2.2 to 6.2)	29.8 (28.7 to 31.0)	2.6 (2.3 to 3.0)	2.3 (2.0 to 2.6)	2.0 (1.7 to 2.3)	6.5 (5.9 to 7.1)
Missing data	2839	4.2 (2.9 to 6.8)	35.3 (33.5 to 37.0)	2.3 (1.7 to 2.8)	1.5 (1.1 to 2.0)	2.3 (1.8 to 2.9)	5.9 (5.0 to 6.7)
Civil status[§]							
Partner	16 573	4.1 (2.4 to 6.5)	32.4 (31.7 to 33.1)	2.6 (2.4 to 2.8)	2.0 (1.8 to 2.2)	2.2 (2.0 to 2.4)	6.3 (5.9 to 6.6)
No partner	18 066	4.1 (2.3 to 6.2)	31.0 (30.3 to 31.6)	2.9 (2.6 to 3.1)	2.3 (2.0 to 2.5)	2.5 (2.3 to 2.7)	7.0 (6.7 to 7.4)
Income, NOK[¶]							
< 125k	3915	4.2 (3.0 to 7.1)	35.5 (34.0 to 37.0)	2.5 (2.0 to 3.0)	1.6 (1.2 to 2.0)	2.9 (2.4 to 3.5)	6.1 (5.7 to 7.3)
125k to 399k	17 024	4.1 (2.4 to 6.2)	30.5 (29.8 to 31.2)	2.7 (2.4 to 2.9)	2.1 (1.9 to 2.3)	2.3 (2.1 to 2.5)	6.5 (6.2 to 6.9)
400k to 700k	5960	3.2 (2.1 to 5.2)	23.3 (22.2 to 24.4)	2.4 (2.0 to 2.8)	2.5 (2.1 to 2.9)	1.8 (1.5 to 2.2)	6.2 (5.6 to 6.9)
> 700k	1283	3.1 (2.0 to 5.1)	22.1 (19.8 to 24.3)	3.3 (2.3 to 4.3)	3.1 (2.2 to 4.1)	1.9 (1.1 to 2.6)	7.7 (6.3 to 9.2)
Missing data	6457	5.1 (3.1 to 8.0)	42.0 (40.8 to 43.2)	3.2 (2.7 to 3.6)	1.9 (1.6 to 2.3)	2.9 (2.5 to 3.3)	6.5 (6.2 to 6.9)
Year							
1999 to 2001	7262	5.3 (3.9 to 8.0)	46.6 (45.5 to 47.8)	1.2 (1.0 to 1.5)	1.3 (1.1 to 1.6)	2.5 (2.1 to 2.9)	4.8 (4.3 to 5.2)
2002 to 2004	7470	4.4 (3.1 to 7.1)	37.0 (35.9 to 38.1)	2.0 (1.7 to 2.3)	1.6 (1.3 to 1.9)	2.3 (2.0 to 2.7)	5.5 (5.0 to 6.0)
2005 to 2007	6295	4.1 (2.7 to 6.2)	30.1 (28.9 to 31.2)	3.5 (3.1 to 4.0)	1.9 (1.5 to 2.2)	2.3 (1.9 to 2.6)	7.0 (6.4 to 7.7)
2008 to 2010	6406	3.2 (2.1 to 5.2)	23.6 (22.6 to 24.6)	3.4 (3.0 to 3.9)	2.5 (2.1 to 2.8)	2.5 (2.1 to 2.9)	7.7 (7.1 to 8.4)
2011 to 2013	7206	2.9 (2.0 to 5.0)	19.6 (18.7 to 20.5)	3.7 (3.3 to 4.2)	3.4 (3.0 to 3.8)	2.2 (1.9 to 2.6)	8.6 (7.9 to 9.2)
Demographics**							
Region 1	19 643	4.1 (2.3 to 6.2)	30.5 (29.9 to 31.2)	2.9 (2.7 to 3.1)	1.9 (1.7 to 2.1)	2.3 (2.1 to 2.5)	6.5 (6.2 to 6.9)
Region 2	9190	4.7 (3.1 to 7.1)	36.6 (35.6 to 37.6)	2.9 (2.5 to 3.2)	2.1 (1.8 to 2.4)	2.2 (1.9 to 2.5)	6.6 (6.1 to 7.1)
Region 3	4468	2.8 (1.3 to 4.9)	21.2 (20.0 to 22.4)	1.7 (1.4 to 2.1)	3.0 (2.5 to 3.5)	2.7 (2.3 to 3.2)	7.1 (6.4 to 7.9)
Region 4	1338	5.9 (3.8 to 8.1)	49.0 (46.4 to 51.7)	3.0 (2.1 to 3.9)	1.8 (1.1 to 2.5)	3.3 (2.3 to 4.3)	7.6 (6.2 to 9.1)

*Patient was readmitted within 90 days of lumbar surgery with a lumbar diagnosis, but was not reoperated

[†]Surgical complication, repeat surgery ≤ 90 days from discharge, or a readmission within 90 days of lumbar surgery (lumbar diagnosis, no reoperation)[‡]Basic education = high school/vocational education; higher education = college/university for three or more years[§]Partner = living registered partner or spouse[¶]NOK = Norwegian currency: 1NOK $\approx$ US\$0.12 $\approx$ £0.10; < 125k = low 10% percentile, 400k = median, > 700k = top 10% percentile**Region 1 = South-Eastern Health Region; region 2 = Western Health Region; region 3 = Central Health Region; region 4 = Northern Health Region
IQR, interquartile range; CI, confidence interval; CCI, Charlson Comorbidity Index

30 961 individual patients. Microdiscectomy was specified in 23 929 patients (69%); the remaining 10 710 (31%) were specified neither as microdiscectomy or standard/open procedures.

The annual public hospital surgery rate was stable throughout the course of the study, with an average of 50 lumbar disc operations per 100 000 Norwegian inhabitants.

Table I shows rates of unfavourable events according to patient characteristics and year of operation. The patients' mean age was 45 years (18 to 92) and the median duration of hospitalization was 4.1 days (IQR 2.3 to 6.3). There was a decrease in median hospitalization from 5.3 days (IQR 3.9 to 8.0) to 2.9 days (IQR 2.0 to 5.0) during the course of this study. The mean age increased from 42.9 years (18 to 87) to 46.6 years (18 to 92)

Table II. Surgical complications within 30 days after 34 639 herniated lumbar disc surgeries*

ICD-10 codes	Description of complication	Operations, n	Percentage of all operations (n = 34 639)
T81.2	Unintentional dural puncture/laceration	376	1.1
T81.4	Infection following a procedure	179	0.5
T81.0	Bleeding or haematoma	172	0.5
T81.8-9	Unspecified and other complications of procedure	134	0.4
T88.4-5	Complications during anaesthesia	47	0.1
T81.3	Disruption of wound/wound dehiscence	20	< 0.1
T81.1	Post-procedural shock	5	< 0.1
T81.5	Foreign body accidentally left in body following procedure	4	< 0.1

*In no patients were the following complications identified: T81.6, acute reaction to foreign body accidentally left during a procedure; T81.7, vascular complications following a procedure; K91, post-procedural disorders of the digestive system; N99, post-procedural disorders of the genitourinary system; G97.8-9, post-procedural disorder of nervous system; I97.8-9, post-procedural disorder of the circulatory system

ICD, International Classification of Diseases

and the proportion of patients over 60 years of age gradually increased from 7.3% (95% CI 6.7 to 7.9) in the first three years to 17% (95% CI 16.5 to 18.2) in the last three years. There were 7% more operations on male than female patients during the study period. The rate of experiencing an unfavourable event (complication, reoperation, or a readmission) was 6.7% (95% CI 6.4 to 6.9). The rate was 6.4% (95% CI 5.9 to 6.9) in operations coded specifically as microdiscectomy and 6.8% (95% CI 6.5 to 7.1) among the remaining, not specified as microdiscectomy or standard/open. No patients died during an operation or within 30 days after hospital discharge.

About one-third of the patients (35% female and 29% male) experienced a prolonged period of hospitalization which increased with the patient's age. Overall, however, there was a decrease in the period of hospitalization over time. Only 3% of the operations were performed on patients with one or more comorbidity according to the Charlson Comorbidity Index. These operations had a significantly higher proportion of prolonged hospitalization, compared with operations on patients with no prior comorbidity.

The surgical complication rate was 2.7% (95% CI 2.6 to 2.9), with 30% being identified on readmission. The most common surgical complications were dural tears/punctures, infections, and haemorrhages (Table II). There were no obvious gender differences, but the complication rate increased with the patient's age and comorbidity. The complication rate was significantly lower in the study's first six years, 1.6% (95% CI 1.4 to 1.8), compared with the following nine years, 3.5% (95% CI 3.2 to 3.8).

The rate of repeat surgery was 2.1% (95% CI 2.0 to 2.3), 14% of the reoperations being performed during the initial admission and the remaining 86% within 90 days of hospital discharge. A total of 14% of the reoperations were coded as reoperations due to a surgical complication, but no cause was specified in the remaining 86%. Age did not significantly impact the rate of repeat surgery, but the rate was higher in patients with prior comorbidities. From 1999 to 2013, there was an increase in the reoperation rate from 1.3% (95% CI 1.1 to 1.6) to 3.4% (95% CI 3.0 to 3.8). In total, 1856 of all operations (5.4%) were followed by a second operation within a year of initial discharge.

The rate of non-surgical readmissions within 90 days was 2.4% (95% CI 2.2 to 2.5). The rate was not significantly affected by stratifying by age, gender, type of operation, or year

of surgery (Table I), but was higher amongst patients with prior comorbidities.

Table III shows the impact of patient comorbidity, sociodemographic factors, and the year of operation on unfavourable events, analyzed with logistic regression modelling. The model showed significant p-values, but, at the same time, low OR (95% CI) for the following factors: 40 to 59 years of age (age 18 to 39 as reference group), greater than or equal to 60 years of age, and comorbidity according to the Charlson Comorbidity Index. There was a statistically significant association and an increase in OR in patients when operated on between the years 2005 to 2007, 2008 to 2010, and 2011 to 2013 (with the years between 1999 to 2001 used as the reference group).

A total of 151 patients were admitted to hospital within 30 days after surgery with medical conditions that may be complications of lumbar surgery, but their hospital stays were not registered as such (Table IV). Their readmissions and medical conditions were, therefore, not included in the study's complication rate analysis. However, if these readmissions are considered as a complication of the patient's lumbar disc surgery, the overall complication rate was 3.2% (95% CI 3.0 to 3.4).

Discussion

In this study of 34 639 herniated lumbar disc operations performed over a 15-year period, 2.7% had a surgical complication, 2.1% had a repeat surgery, 2.4% had a non-surgical readmission, and 6.6% had at least one of these unfavourable events (surgical complication, repeat surgery, or a non-surgical readmission). There was a slight increase in complication rates over the course of the study. Unfavourable events were found to be significantly associated with advanced age and comorbidity but there was limited association with other sociodemographic characteristics such as gender, geographical affiliation, education, civil status, and income.

The present study has several strengths, one being that the data set represents almost all operations for lumbar disc herniation carried out through the public health system in Norway during the defined period. There was, therefore, no selection bias in terms of only using procedures performed by specific surgeons or hospitals.¹⁷ Thus, the study was independent of surgeons' and hospitals' prior experience, competence, and general willingness to take part in the study. Private sector procedures

Table III. Multivariate logistic regression model showing the associations of sociodemographic factors, comorbidity, and year of operation on the occurrence of unfavourable events in lumbar disc surgery

Patient characteristics, predictors	OR for an unfavourable event (95% CI) (n = 27 872)	p-value
Age group, yrs		
18 to 39	1	Referent
40 to 59	1.23 (1.11 to 1.37)	< 0.001
≥ 60	1.22 (1.02 to 1.48)	0.03
Female gender	1.06 (0.96 to 1.17)	0.23
Comorbidity (a score of ≥ 1 according to the Charlson Comorbidity Index)	1.62 (1.27 to 2.06)	< 0.001
Education (basic education only, high school/vocational education)	1.06 (0.95 to 1.19)	0.30
Civil status (no living registered partner or spouse)	1.18 (0.98 to 1.41)	0.09
Low income*	1.05 (0.91 to 1.21)	0.53
Demographics (health regions in Norway)		
South-Eastern	1	Referent
Western	0.97 (0.87 to 1.09)	0.61
Central	1.10 (0.95 to 1.27)	0.19
Northern	1.23 (0.96 to 1.56)	0.10
Year of operation		
1999 to 2001	1	Referent
2002 to 2004	1.17 (0.98 to 1.40)	0.08
2005 to 2007	1.50 (1.28 to 1.75)	< 0.001
2008 to 2010	1.61 (1.38 to 1.89)	< 0.001
2011 to 2013	1.84 (1.59 to 2.15)	< 0.001

* < 125 000 NOK yearly income in the year prior to surgery. This number represents the lowest 10% percentile of the Norwegian population median; NOK = Norwegian currency, 1 NOK ≈ US\$0.12 ≈ £0.10

OR, odds ratio; CI, confidence interval

Table IV. Description, number, and frequency of medical conditions not registered as complications of surgery, but leading to hospital readmission within 30 days of initial discharge*

Description of medical condition	Admissions, n (n = 151)	Percentage of all operations (n = 34 639)
Unspecified abdominal/chest pain	26	0.08
Cystitis	19	0.05
Pneumonia	10	0.03
Pulmonary embolism	9	< 0.03
Arrhythmia	8	< 0.03
Gastroenteritis	7	< 0.03
Sepsis	6	< 0.03
Heart attack	6	< 0.03
Gastrointestinal ulcer	5	< 0.02
Cholelithiasis	5	< 0.02
Headache	5	< 0.02
Urolithiasis	5	< 0.02
Skin infection	5	< 0.02
Brain infarct	4	< 0.02
Intraspinal abscess	4	< 0.02
Appendicitis	4	< 0.02
Deep venous thrombosis	3	< 0.01
Subarachnoid haemorrhage, epilepsy, delirium, depression/ anxiety, transient ischemic attack, dehydration, diabetes, lung abscess, pneumothorax, pancreatitis	≤ 2	< 0.01
Total	151	0.4

* Since these conditions have not been coded directly as complications to surgery, it is not possible to know with certainty whether these medical conditions were, or were not, related to the surgical procedure

accounted for 20% of the lumbar disc operations in Norway during the second half of the study period (2007 to 2013) but hardly any private surgery was performed during the first half (1999 to 2006). Private surgical patients are on average both younger

and healthier than patients operated through the public sector.¹² This introduces a selection bias in the years 2007 to 2013, in that the public sector (the subject of this study) is left with a greater proportion of patients with a higher risk of unfavourable events.

Except for data concerning income and education, almost no data were missing, and there is little reason to doubt the accuracy of time of admission and discharge, as their registration falls under the legal responsibilities of the hospital system as a prerequisite for reimbursement. No information regarding the identity of the surgeons was recorded, thus there is no obvious incentive for a physician to not record complications. Nevertheless, the accuracy and completeness of the diagnostic and complication coding is difficult to assess, due to potential variation in the coding practices, experience, and thoroughness of physicians and surgeons. It is, therefore, important to acknowledge that the diagnostic codes may not accurately reflect what was documented in the patients' medical records.¹⁸ Moreover, the data set provided no information regarding prior lumbar surgery outside the study period, a factor known to influence reoperation rates.¹⁹

Although the number of comparable studies is limited, their estimates and trends are similar to those revealed in the present study. The median length of hospital stay for lumbar surgery continues to decrease in Scandinavia.⁶ The 90-day reoperation rate is 1% to 3% and can be associated with the rate of complications.¹⁵ The elderly are more prone to complications,²⁰ of which most common complications are dural tears/punctures, infections, and haemorrhage.¹⁶

Bearing in mind that complications tend to be under-reported by health care workers,²¹ the present study's mean surgical complication rate was generally (but not exclusively (1.6%))²² lower than the rate reported in other studies (5.7%, 3.9%, 3% to 5%, and 5%).^{16,20,23,24} Despite improvements in surgical techniques and hospital care that evolved during the study period, the especially low complication rate of 1.6% for the first six years of the study compared with the more commonly reported rate of 3.5% for the remaining nine years may be explained by the increase of private surgery along with the general trend of operating on older patients. Moreover, the present study only accounted for surgical complications that occurred during primary hospitalization or complications that led to a subsequent hospital readmission. It is quite possible that minor complications were treated in General Practice, without being recorded in the NPR, which might also explain the low surgical complication rate. Furthermore, we did not include discectomies and decompressions undertaken for more comprehensive procedures such as a spinal fusion or disc prosthesis surgery. Previous studies have included a range of different surgical procedures for lumbar disc herniation, which could explain their slightly higher rates of surgical complications and mortality (0.5%, 0.7%, 0.8%).^{5,6,23}

The low ORs in the logistic regression model suggest that advanced age and comorbidity contribute only a small additive increment to the risk of an unfavourable event. However, in our study, patients were all carefully selected for surgery with the general understanding that age and comorbidity will influence their surgical outcome. Thus, the study's comorbid and elderly patients may have a higher level of physical health than the elderly and comorbid in the general population. Therefore, the ORs may be misleadingly low, and age and comorbidity to be greater influencers of non-favourable events than our data suggests.

The present study showed that surgery for lumbar disc herniation has very low mortality (< 0.001%) and fewer than 7% of

patients experience an unfavourable event such as a reoperation, complication, or readmission. Whether this rate is acceptable must be considered in relation to the health gains achieved, but patients can in general be informed that lumbar disc surgery is safe, although advanced age and comorbidity may slightly affect the complication rate. While the readmission and reoperation rates presented in the article are very accurate, the surgical complication rates may unfortunately be falsely low due to under reporting by the hospital physicians and surgeons.

Take home message

- To our knowledge, this is the largest study that specifically quantifies these unfavourable events over such a large sample size.
- We believe that these updated rates are important from a health-economic perspective and in relation to the conversations between patient and surgeon that take place in daily clinical practice.

References

1. Lurie JD, Tosteson TD, Tosteson AN, et al. Surgical versus nonoperative treatment for lumbar disc herniation: eight-year results for the spine patient outcomes research trial. *Spine* 2014;39:3–16.
2. Weinstein MC, Stason WB. Foundations of cost-effectiveness analysis for health and medical practices. *N Engl J Med* 1977;296:716–721.
3. Tosteson AN, Skinner JS, Tosteson TD, et al. The cost effectiveness of surgical versus nonoperative treatment for lumbar disc herniation over two years: evidence from the Spine Patient Outcomes Research Trial (SPORT). *Spine* 2008;33:2108–2115.
4. Ramaswami R, Ghogawala Z, Weinstein JN. Management of Sciatica. *N Engl J Med* 2017;376:1175–1177.
5. Pearson A, Lurie J, Tosteson T, et al. Who should have surgery for an intervertebral disc herniation? Comparative effectiveness evidence from the spine patient outcomes research trial. *Spine* 2012;37:140–149.
6. Jansson KA, Németh G, Granath F, Blomqvist P. Surgery for herniation of a lumbar disc in Sweden between 1987 and 1999. An analysis of 27,576 operations. *J Bone Joint Surg [Br]* 2004;86-B:841–847.
7. Sivasubramaniam V, Patel HC, Ozdemir BA, Papadopoulos MC. Trends in hospital admissions and surgical procedures for degenerative lumbar spine disease in England: a 15-year time-series study. *BMJ Open* 2015;5:e009011.
8. Deyo RA, Taylor VM, Diehr P, et al. Analysis of automated administrative and survey databases to study patterns and outcomes of care. *Spine* 1994;19(Suppl):2083s–2091s.
9. Hassani S, Lindman AS, Kristoffersen DT, Tomic O, Helgeland J. 30-Day Survival Probabilities as a Quality Indicator for Norwegian Hospitals: Data Management and Analysis. *PLoS One* 2015;10:e0136547.
10. Sundararajan V, Henderson T, Perry C, et al. New ICD-10 version of the Charlson comorbidity index predicted in-hospital mortality. *J Clin Epidemiol* 2004;57:1288–1294.
11. No authors listed. International Statistical Classification of Diseases and Related Health Problems, 10th Revision (ICD-10 Norwegian version). <https://finnkode.ehelse.no> (date last accessed 11 February 2019).
12. Grotle M, Solberg T, Storheim K, Laerum E, Zwart JA. Public and private health service in Norway: a comparison of patient characteristics and surgery criteria for patients with nerve root affections due to disc herniation. *Eur Spine J* 2014;23:1984–1991.
13. No authors listed. The Norwegian Quality Register for Back Surgery. Last updated 9 January 2019. <https://unn.no/fag-og-forskning/medisinske-kvalitetsregistre/nasjonalt-kvalitetsregister-for-ryggkirurgi> (date last accessed 14 January 2019).
14. No authors listed. NOMESCO Classification of Surgical Procedures (NCSP). Nordic Centre for Classifications in Health Care. http://www.nordclass.se/ncsp_e.htm (date last accessed 14 January 2019).
15. Martin BI, Mirza SK, Flum DR, et al. Repeat surgery after lumbar decompression for herniated disc: the quality implications of hospital and surgeon variation. *Spine J* 2012;22:89–97.
16. Deyo RA, Cherkin DC, Loeser JD, Bigos SJ, Ciol MA. Morbidity and mortality in association with operations on the lumbar spine. The influence of age, diagnosis, and procedure. *J Bone Joint Surg [Am]* 1992;74-A:536–543.
17. Gavrielov-Yusim N, Friger M. Use of administrative medical databases in population-based research. *J Epidemiol Community Health* 2014;68:283–287.

18. **Cherkin DC, Deyo RA, Volinn E, Loeser JD.** Use of the International Classification of Diseases (ICD-9-CM) to identify hospitalizations for mechanical low back problems in administrative databases. *Spine* 1992;17:817–825.
19. **Osterman H, Sund R, Seitsalo S, Keskimäki I.** Risk of multiple reoperations after lumbar discectomy: a population-based study. *Spine* 2003;28:621–627.
20. **Strömqvist F, Strömqvist B, Jönsson B, Karlsson MK.** Surgical treatment of lumbar disc herniation in different ages-evaluation of 11,237 patients. *Spine J* 2017;17:1577–1585.
21. **Ohn A, Elfström J, Liedgren C, Rutberg H.** Reporting of sentinel events in Swedish hospitals: a comparison of severe adverse events reported by patients and providers. *Jt Comm J Qual Patient Saf* 2011;37:495–501.
22. **Peul WC, van den Hout WB, Brand R, et al.** Prolonged conservative care versus early surgery in patients with sciatica caused by lumbar disc herniation: two year results of a randomised controlled trial. *BMJ* 2008;336:1355–1358.
23. **Nandyala SV, Marquez-Lara A, Fineberg SJ, Hassanzadeh H, Singh K.** Complications after lumbar spine surgery between teaching and nonteaching hospitals. *Spine* 2014;39:417–423.
24. **Weinstein JN, Tosteson TD, Lurie JD, et al.** Surgical vs nonoperative treatment for lumbar disk herniation: the Spine Patient Outcomes Research Trial (SPORT): a randomized trial. *JAMA* 2006;296:2441–2450.

Author contributions:

O. R. Fjeld: Designed the study methodology, Curated and analyzed the data, Drafted and wrote the manuscript.

L. Grøve: Curated the data, Supervised the study, Wrote the manuscript.

J. Helgeland: Curated the data, Wrote the manuscript.

M. C. Småstuen: Curated and analyzed the data, Wrote the manuscript.

T. K. Solberg: Supervised the study, Wrote the manuscript.

J-A. Zwart: Supervised the study, Acted as project administrator, Wrote the manuscript.

M. Grotle: Designed the study methodology, Supervised the study, Acted as project administrator, Wrote the manuscript.

Funding statement:

Governmental funds from the South-Eastern Norway Regional Health Authority (project number 2013030) were received in support of this work. No relevant financial activities outside the submitted work.

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article.

Acknowledgements:

The authors would like to thank the South-Eastern Norway Regional Health Authority for the funds received in support of this work.

Ethical review statement:

The study was approved by the Norwegian Data Inspectorate (2014/14413) and the Norwegian Regional Ethics Committee (2013/1662, REC south-east D). No individual patient consent was required.

Open access statement:

This is an open-access article distributed under the terms of the Creative Commons Attributions license (CC-BY-NC), which permits unrestricted use, distribution, and reproduction in any medium, but not for commercial gain, provided the original author and source are credited.

This article was primary edited by S. P. F. Hughes.

Clinical pain research

Olaf R. Fjeld*, Margreth Grotle, Dagfinn Matre, Linda M. Pedersen, Marie U. Lie, Milada C. Småstuen, Kjersti Storheim, Ingrid Heuch, Audun Stubhaug, John-Anker Zwart and Kristian Bernhard Nilsen

Predicting the outcome of persistent sciatica using conditioned pain modulation: 1-year results from a prospective cohort study

<https://doi.org/10.1515/sjpain-2019-0112>

Received August 8, 2019; revised August 26, 2019; accepted August 28, 2019; previously published online September 17, 2019

Abstract

Background and aims: Recovery in patients hospitalised with severe sciatica is unpredictable. Prognostic tools to aid clinicians in the early identification of patients at risk of developing chronic sciatic pain are warranted.

***Corresponding author: Olaf R. Fjeld**, Research and Communication Unit for Musculoskeletal Health (FORMI), Oslo University Hospital, Ullevål, Oslo, Norway; Department of Neurology, Oslo University Hospital, Oslo, Norway; and Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway, E-mail: fjeld@tcd.ie

Margreth Grotle: Research and Communication Unit for Musculoskeletal Health (FORMI), Oslo University Hospital, Ullevål, Oslo, Norway; and Faculty of Health Science, OsloMet - Oslo Metropolitan University, Oslo, Norway

Dagfinn Matre: Department of Work Psychology and Physiology, National Institute of Occupational Health, Oslo, Norway

Linda M. Pedersen, Marie U. Lie, Kjersti Storheim and Ingrid Heuch: Research and Communication Unit for Musculoskeletal Health (FORMI), Oslo University Hospital, Ullevål, Oslo, Norway

Milada C. Småstuen: Research and Communication Unit for Musculoskeletal Health (FORMI), Oslo University Hospital, Ullevål, Oslo, Norway; and Institute of Physiotherapy, Faculty of Health Science, Oslo and Akershus University College, Oslo, Norway

Audun Stubhaug: Department of Pain Management and Research, Faculty of Medicine, University of Oslo, Oslo, Norway

John-Anker Zwart: Research and Communication Unit for Musculoskeletal Health (FORMI), Oslo University Hospital, Ullevål, Oslo, Norway; Department of Neurology, Oslo University Hospital, Oslo, Norway; and Norway Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway

Kristian Bernhard Nilsen: Research and Communication Unit for Musculoskeletal Health (FORMI), Oslo University Hospital, Ullevål, Oslo, Norway; Department of Neurology, Oslo University Hospital, Oslo, Norway; and Department of Neuromedicine and Movement Science, Norwegian University of Science and Technology, Trondheim, Norway

Conditioned pain modulation (CPM) is a psychophysical measure of the endogenous pain modulatory pathways. Several studies have suggested CPM as a potentially important predictive biomarker for the development of chronic pain. The aim of the study was to determine whether CPM effect in patients still suffering from leg pain 6 weeks after hospital discharge for severe sciatica is associated with persistent leg pain at 12 months. A potential association would suggest that measuring CPM effect could be a valuable prognostic tool in the hospital management of sciatica.

Methods: A prospective cohort study in which CPM effect was measured 6 weeks after hospital discharge following an acute admission with sciatica as the main complaint. The impact of CPM effect on the outcome was analysed using logistic regression. The outcome measured was self-reported leg pain score of ≥ 1 in the past week on a 0–10 numeric rating scale (NRS) at 12 months post discharge.

Results: A total of 111 patients completed the entire study, 51 of whom received non-randomised surgical treatment. Crude and confounder adjusted analyses showed no significant association between CPM effect and leg-pain measured at 12 months, crude Odds Ratio 0.87, 95% CI 0.7–1.1, $p = 0.23$.

Conclusions: Our results suggest that CPM assessment has limited prognostic value for the long-term outcome in severe sciatica when measured 6 weeks after hospital discharge.

Implications: The present study adds important knowledge concerning the limited clinical use of late CPM testing in sciatica patients. The heterogeneity in patients, the wide range of treatments received and a generally favourable outcome are factors that may affect CPM's clinical value as a prognostic factor for severe sciatica.

Keywords: conditioned pain modulation; sciatica; disc herniation; pain; prognostic factor; neuropathic pain.

1 Introduction

The treatment of lumbar spinal disorders constitutes a large portion of hospital costs, but the economic impact of work-absenteeism continues long after patients have been discharged [1]. Long-term recovery in patients hospitalised with severe sciatica is unpredictable and warrants better prognostic tools to aid clinicians in the identification of patients at risk of developing chronic sciatic pain and disability [2, 3].

Conditioned pain modulation (CPM) effect is a psychophysical measure of the endogenous pain modulatory pathways. This is demonstrated by a painful test stimulus (TS) being perceived as less painful in the presence of, or shortly after, a second painful stimulus [conditioning stimulus (CS)]. The difference in perceived pain of the TS with and without the CS is termed the CPM effect [4]. Impaired inhibitory CPM effect has been associated with a variety of chronic pain disorders [5], and has been suggested as a potentially important predictive biomarker for the development of chronic pain and a predictor of analgesic response to neuropathic pain [6, 7]. To our knowledge, the predictive value of CPM in sciatica has not been studied. The primary aim of this study was to determine whether CPM effect in patients still suffering from leg pain 6 weeks after hospital discharge for severe sciatica is associated with persistent leg pain at 12 months post hospital discharge. A potential association would suggest that measuring CPM effect could be a valuable prognostic tool in the hospital management of sciatica.

2 Methods

2.1 Study design

This was a prospective cohort study with assessments at three different time points: (1) questionnaire, clinical examination and lumbar MRI upon hospital admission, (2) CPM testing and questionnaire 6 weeks after hospital discharge, (3) a postal questionnaire at 12 months (Fig. 1).

2.2 Participants

Between December 2012 and March 2018, eligible patients were consecutively recruited from a larger cohort study aimed at patients suffering from sciatica that required acute hospitalisation at the Neurological Department of Oslo University Hospital [8]. This hospital provides care to all Oslo residents who require acute surgical or non-surgical treatment of sciatica. All physicians responsible for hospital admission were working independently of the study program and patients were not hospitalized for the purpose of inclusion in the study. Upon hospital admission, most patients were treated with a combination of paracetamol, non-steroidal anti-inflammatory drugs and mild or strong opioids. Lumbar surgery was typically offered to patients with severe sciatic pain refractory to analgesics or with a severe motor deficit. No other forms

Fig. 1: Flow diagram of the study population.

of treatment were offered. Patients (surgical and non-surgical) that still suffered from sciatic pain [≥ 2 on a numeric rating scale (NRS)] 6 weeks post hospital discharge (baseline of the study/CPM testing) were further for the present study following a series of inclusion and exclusion criteria.

Inclusion criteria: Age 18–65; suffering from sciatica, defined as radiating leg-pain with dermatomal distribution graded ≥ 2 on a self-reported NRS ranging from 0 to 10 (0=no pain, 10=worst pain imaginable) at baseline; hospitalised with acute sciatica as their major complaint with self-reported leg pain of ≥ 4 on a NRS 6 weeks prior to baseline testing; and radiological confirmation of disc herniation on magnetic resonance imaging (MRI).

Exclusion criteria: Cauda-equina syndrome; pregnant or breastfeeding; drug or alcohol addiction; primary psychiatric disease; use of antipsychotics or antidepressants; limited Norwegian proficiency; chronic non-lumbar pain rated ≥ 4 on NRS for ≥ 3 months in the last 2 years; lumbar surgery between baseline CPM testing and 12 months post hospital discharge.

2.3 The predictor of interest (independent variable): mean CPM effect

All participants received written information concerning the test program, but were blinded to the study hypothesis and that CPM would be tested. Each test was conducted by experimenters wearing the same clothing in a laboratory

with a consistent placement of instruments. During the experiment, the tester read slowly from a standardized instruction protocol, and all sessions followed the same procedure. The study procedure was adapted from a previously published protocol [9]. The CPM effect was measured following a 4-step procedure using a computerized temperature stimulation device [10] with a 30×30 mm Peltier thermode as TS and a container [11] of circulating water at 7°C as CS (Fig. 2).

Step 1 (identifying TS temperature): After identifying heat pain threshold and heat pain tolerance level by the methods of limit (baseline 32°C , rate of change 1°C/s), the experimenter identified the individual patient's TS temperature ($^\circ\text{C}$) scored as 6 cm on a 10 cm computerized visual analogue scale (VAS) with endpoints “no pain” and “worst imaginable pain”.

Step 2 (TS): Patients received 120 s of continuous TS (baseline temperature: 32°C , increase rate: 2°C/s) to their right forearm. The perceived TS pain intensity was concurrently scored on a 10 cm computerized horizontal VAS (endpoints: “no pain” and “worst pain imaginable”), with the patients continuously scrolling the wheel on a computer mouse according to perceived pain. The TS pain scoring was sampled at 1 Hz.

Step 3 (TS and CS): After a 300 s break, patients received another 120 s of TS on the same arm while the other arm was immersed in a container of 7°C circulating water (CS). As in step 2, TS pain was scored simultaneously.

Fig. 2: The method of testing conditioned pain modulation (CPM). ^{a)}TS: painful heat applied by a peltier thermode on the participants right forearm. ^{b)}Pain rating: participants scrolled the wheel on a computer mouse according to perceived TS heat pain (10 cm horizontal VAS on a computer). ^{c)}CS: left hand immersed in a container of painful cold (7°C) circulating water. ^{d)}Timeline. TS=test stimulus; VAS=visual analogue scale; CS=conditioning stimulus.

Step 4 (CPM calculation): The average pain score from the 120 s of TS in both step 2 and step 3 was applied when calculating the CPM effect, defined as the difference between the two means ($VAS_{(TS \text{ and } CS)} - VAS_{(TS)}$). Accordingly, a negative value represents an inhibitory CPM effect.

2.4 Main outcome (dependent variable): leg pain at 12 months post hospital discharge

At 12 months, a leg pain score of ≥ 1 in the past week on a self-reported 0–10 NRS was set as the cut-off value for persistent leg pain and the present study's final outcome. Patients with a leg pain NRS score of 0 were considered to have recovered from leg-pain. Leg pain does not give a direct measure of patient function or quality of life, but it is found to be one of the most responsive outcomes for capturing disabling symptoms in patients suffering from sciatica [12].

2.5 Statistical analysis

All data analyses were performed using SPSS 24.00 (IBM, Armonk, NY, USA). Significance level was set to 5%.

Continuous data were described with mean and standard deviation (SD). Associations between pairs of categorical data were analysed using the χ^2 test and associations between pairs of continuous data were analysed using independent samples *t*-tests. Logistic regression analysis was used to investigate the association between the independent variables (CPM, age, sex, baseline leg pain) and the dependent variable (leg pain ≥ 1 scored on a 0–10 NRS at 12 months). All independent variables were entered simultaneously, and all significant associations were described with Odds Ratios (OR) and 95% confidence intervals (95% CI). No missing values were imputed. Finally, three sensitivity regression analyses were carried out in which more severe cut-off values for the main outcome were investigated; (1) leg pain ≥ 2 on a 0–10 NRS measured at 12 months, (2) leg pain ≥ 6 on a 0–10 NRS measured at 12 months, (3) a reduction in leg pain of ≥ 2 on a 0–10 NRS at 12 months.

3 Results

Of the 149 patients included in the study, 111 patients responded to the follow-up questionnaire at 12 months (Fig. 1). Table 1 shows the baseline patient characteristics

Table 1: Characteristics and measured CPM effect for the different patient subgroups.

	N (% of all patients)	Mean CPM effect in cm (CI)
All patients	111	−2.4 (−2.8, −2.1)
Males	72 (65)	−2.4 (−2.8, −2.0)
Females	39 (35)	−2.4 (−3.1, −1.7)
Education >12 years	79 (72)	−2.2 (−2.6, −1.8)
Education <12 years	32 (28)	−2.9 (−3.6, −2.1)
Living with spouse/partner	85 (77)	−2.3 (−2.7, −1.9)
No spouse/partner	26 (23)	−2.7 (−3.7, −2.0)
Actively employed	83 (75)	−2.5 (−2.9, −2.1)
Not actively employed	28 (25)	−2.3 (−3.0, −1.5)
European Caucasian	103 (94)	−2.7 (−2.7, −2.0)
Not European Caucasian	6 (6)	−3.5 (−5.3, −1.7)
Disc herniation surgery within 6 weeks prior to CPM testing	51 (54)	−2.5 (−3.1, −1.9)
No disc herniation surgery within 6 weeks prior to CPM testing	60 (46)	−2.3 (−2.8, −1.9)
Any previous lumbar surgery	45 (41)	−2.3 (−2.8, −1.8)
No previous lumbar surgery	66 (59)	−2.5 (−3.0, −2.0)
Sporadic use of analgesics for leg-pain	91 (82)	−2.3 (−2.7, −1.9)
No use of analgesics for leg pain	20 (20)	−2.8 (−3.5, −2.1)
Daily use of analgesics	47 (42)	−2.4 (−2.9, −1.9)
Non daily use of analgesics	64 (58)	−2.4 (−2.9, −2.0)
Current smoker	13 (12)	−3.2 (−4.4, −2.0)
Non smoker	97 (88)	−2.3 (−2.7, −1.9)
Poor quality of life last week, yes	24 (22)	−2.5 (−3.3, −1.8)
Poor quality of life last week, no	85 (78)	−2.4 (−2.8, −2.0)

CI = confidence interval; CPM = conditioned pain modulation.

Fig. 3: Dispersion of CPM scores for each of the 111 patients.

and the different CPM values. The patients were hospitalized for a mean (SD) of 5.2 (4.4) days. A total of 51 patients (54%) received surgical treatment in the 6-week period between hospital admission and CPM testing (baseline). Mean CPM value was measured to -2.4 cm (95% CI $-2.8, -2.1$).

Figure 3 shows the general dispersion of CPM scores among the patients. No statistically significant difference in mean CPM effect was evident when comparing surgically and non-surgically treated patients, -2.3 cm vs. -2.5 cm, $t(109)=0.56$, $p=0.59$. Mean NRS leg pain was 4.0 (SD 2.1) measured at baseline and 2.5 (SD 2.5) at 12 months (Fig. 4). The mean (SD) change in leg pain measured between baseline and 12 months for each patient ($\text{NRS score}_{(\text{baseline})} - \text{NRS score}_{(12 \text{ months})}$) was 1.5 (2.8) (Fig. 5).

3.1 Association between CPM effect and main outcome

Crude analysis did not reveal any statistically significant association between CPM effect and leg pain ≥ 1

Fig. 4: Mean NRS leg pain measured at baseline and 12 months.

Fig. 5: The change in leg pain measured between baseline and 12 months ($\text{NRS score}_{(\text{baseline})} - \text{NRS score}_{(12 \text{ months})}$) for each patient. A positive value represents a decrease in pain from baseline to 12 months.

on a 0–10 NRS measured at 12 months, OR 0.87, 95% CI 0.7–1.1, $p=0.23$. Further, no significant association was found when adjusting individually or concurrently for the following: age, sex, baseline leg-pain and surgical treatment (Table 2). Sensitivity analysis of the main outcome revealed no association between CPM effect and; (1) leg pain ≥ 2 on a 0–10 NRS measured at 12 months, (2) leg pain ≥ 6 on a 0–10 NRS measured at 12 months, (3) a reduction in 0–10 NRS leg pain of ≥ 2 at 12 months (logistic regression analysis, results not shown).

4 Discussion

This prospective cohort study on patients admitted to hospital due to acute sciatica revealed no statistically significant

Table 2: The association between CPM effect and persistent leg-pain at 12 months, adjusted for age, sex, baseline leg pain and surgical treatment.

Outcome (dependent variable)=leg pain score of ≥ 1 the past week on a 0–10 NRS at 12 months, $n=111$, $R^2=0.05$				
Factors tested (independent variables entered simultaneously)	B	Wald statistic	p-Value	OR (95% CI)
CPM effect (cm)	−0.14	1.31	0.25	0.87 (0.69, 1.10)
Age (years)	0.01	0.26	0.61	1.01 (0.97, 1.05)
Sex (male)	0.36	0.65	0.42	1.43 (0.60, 3.42)
Baseline 0–10 NRS leg pain score	0.06	0.28	0.60	1.06 (0.86, 1.31)
Surgery prior to CPM testing	0.43	0.95	0.33	1.54 (0.65, 3.68)

R^2 = Nagelkerke R^2 ; NRS = 0–10 numeric rating scale; B = unstandardized β coefficient; OR = odds ratio; CI = confidence interval; CPM = conditioned pain modulation.

association between CPM effect measured 6 weeks post hospital discharge and persistent leg-pain at 12 months. However, it is important to bear in mind that the purpose of the study was not to explore the causal inference of CPM effect in sciatic pain chronicity or test the validity of the CPM paradigm, but to test its crude prognostic value in a clinical setting of heterogeneous sciatic patients whose pain persists despite pharmaceutical or surgical treatment. It is possible that further stratified analysis of different subgroups of sciatica patients or implementation of stricter inclusion criteria in terms of symptom debut, treatment and pain characteristics may demonstrate an association between the CPM effect and the outcomes that the present study failed to show. However, stricter inclusion criteria and stratification of this heterogeneous group of patients would limit the use of CPM to only a few selected patients, decreasing its value as a clinically useful prognostic tool, especially since few patients have persistent sciatica 6 weeks after hospital discharge.

One can argue that a limitation in the present study is that patients were CPM tested 6 weeks post admission rather than during the acute phase of the disease or prior to lumbar surgery. The present study's patients may have undergone both psychological and physical changes that could affect their CPM effect and its possible predictive properties. However, it is more practical, accurate and economically feasible to test patients in an outpatient laboratory setting, rather than a crowded hospital ward where patients are stressed, medicated, immobile and facing a limited number of days before discharge.

Currently, there is insufficient data to identify a superior CPM protocol [13–15]. Until such data is available, study validity and comparability of different CPM protocols will continue to be questioned. This may also apply to the present study. One can argue that the present study's CPM protocol may introduce attention bias in the form of distraction, since the second TS is applied in parallel to the CS and not in sequence as in other CPM protocols [16]. Further, the

0–10 NRS pain scale as the final outcome could not be preserved as a continuous variable in the regression analysis without violating the assumption of multivariate normality. Though other cutoffs values were tested and showed no significant results, the present study runs the risk of losing valuable information and making type II errors when data collected as continua were split into categories [17].

There is a limited and conflicting body of studies exploring CPM effect as a potential prognostic factor for persistent pain and disability in patients with sciatica and other spinal disorders. Supporting the present article's main findings is a cohort-study of 156 primary care patients with non-specific low-back pain who were CPM tested and subsequently assessed for chronic pain 4 months later [18]. The study failed to demonstrate CPM effect as a predictor for pain chronicity. Though the patient population was different in terms of disease mechanism and symptom severity, this previous study bears a methodological resemblance to the present study. The present findings are further supported by two additional CPM studies on low-back pain patients, which indicated that altered CPM is not a major determinant of hyperalgesia and pain in their patient population [19–21]. In contrast, CPM effect was shown to have predictive value for persistent postoperative pain in patients who had undergone surgical thoracotomy and abdominal surgery [22]. The obvious methodological difference in this case is that the CPM testing was performed prior to pain debut, and not after or during ongoing pain as in the present study. This suggests that measures of CPM effect have predictive value if measured prior to pain debut, but not after.

5 Conclusions

Our results suggest that CPM effect measured 6 weeks post hospital discharge has limited prognostic value for the

long-term outcome in persistent sciatic pain. Despite this negative finding, CPM effect as a biomarker in the management of sciatica should not be entirely dismissed. The clinical value of CPM testing may lie in its ability to direct treatment or predict outcomes in more carefully selected sciatica patients than in the present study.

Authors' statements

Research funding: Governmental funding. South-Eastern Norway Regional Health Authority (project number 2012/1108) funds were received in support of this work.

Conflict of interest: Authors state no conflict of interest.

Informed consent: Written informed consent was obtained prior to participation.

Ethical approval: Human testing was performed in accordance with the Declaration of Helsinki, and the Norwegian Regional Ethics Committee (REK) approved the study (project number 2012/1108).

References

- [1] van Tulder MW, Koes BW, Bouter LM. A cost-of-illness study of back pain in The Netherlands. *Pain* 1995;62:233–40.
- [2] Balague F, Nordin M, Sheikhzadeh A, Echegoyen AC, Brisby H, Hoogewoud HM, Fredman P, Skovron ML. Recovery of severe sciatica. *Spine* 1999;24:2516–24.
- [3] Ashworth J, Konstantinou K, Dunn KM. Prognostic factors in non-surgically treated sciatica: a systematic review. *BMC Musculoskeletal Disord* 2011;12:208.
- [4] Yarnitsky D, Arendt-Nielsen L, Bouhassira D, Edwards RR, Fillingim RB, Granot M, Hansson P, Lautenbacher S, Marchand S, Wilder-Smith O. Recommendations on terminology and practice of psychophysical DNIC testing. *Eur J Pain* 2010;14:339.
- [5] Lewis GN, Rice DA, McNair PJ. Conditioned pain modulation in populations with chronic pain: a systematic review and meta-analysis. *J Pain* 2012;13:936–44.
- [6] Yarnitsky D, Granot M, Nahman-Averbuch H, Khamaisi M, Granovsky Y. Conditioned pain modulation predicts duloxetine efficacy in painful diabetic neuropathy. *Pain* 2012;153:1193–8.
- [7] Niesters M, Proto PL, Aarts L, Sarton EY, Drewes AM, Dahan A. Tapentadol potentiates descending pain inhibition in chronic pain patients with diabetic polyneuropathy. *Br J Anaesth* 2014;113:148–56.
- [8] Fjeld O, Grotle M, Siewers V, Pedersen LM, Nilsen KB, Zwart JA. Prognostic factors for persistent leg-pain in patients hospitalized with acute sciatica. *Spine* 2017;42:E272–9.
- [9] Matre D, Andersen MR, Knardahl S, Nilsen KB. Conditioned pain modulation is not decreased after partial sleep restriction. *Eur J Pain* 2016;20:408–16.
- [10] PATHWAY – Pain & Sensory Evaluation System, Model ATS. Medoc Ltd., Ramat Yishai, Israel.
- [11] LAUDA RM6-RMS low temperature thermostat. LAUDA-Brinkmann LP., United States.
- [12] Haugen AJ, Grøvre L, Brox JJ, Natvig B, Keller A, Soldal D, Grotle M. Estimates of success in patients with sciatica due to lumbar disc herniation depend upon outcome measure. *Eur Spine J* 2011;20:1669–75.
- [13] Pud D, Granovsky Y, Yarnitsky D. The methodology of experimentally induced diffuse noxious inhibitory control (DNIC)-like effect in humans. *Pain* 2009;144:16–9.
- [14] Yarnitsky D, Bouhassira D, Drewes AM, Fillingim RB, Granot M, Hansson P, Landau R, Marchand S, Matre D, Nilsen KB, Stubhaug A, Treede RD, Wilder-Smith OH. Recommendations on practice of conditioned pain modulation (CPM) testing. *Eur J Pain* 2015;19:805–6.
- [15] Kennedy DL, Kemp HI, Ridout D, Yarnitsky D, Rice AS. Reliability of conditioned pain modulation: a systematic review. *Pain* 2016;157:2410–9.
- [16] Tousignant-Laflamme Y, Page S, Goffaux P, Marchand S. An experimental model to measure excitatory and inhibitory pain mechanisms in humans. *Brain Res* 2008;1230:73–9.
- [17] Streiner DL. Breaking up is hard to do: the heartbreak of dichotomizing continuous data. *Can J Psychiatry* 2002;47:262–6.
- [18] LeResche L, Turner JA, Saunders K, Shortreed SM, Von Korff M. Psychophysical tests as predictors of back pain chronicity in primary care. *J Pain* 2013;14:1663–70.
- [19] Mlekusch S, Schliessbach J, Camara RJ, Arendt-Nielsen L, Juni P, Curatolo M. Do central hypersensitivity and altered pain modulation predict the course of chronic low back and neck pain? *Clin J Pain* 2013;29:673–80.
- [20] Mlekusch S, Neziri AY, Limacher A, Juni P, Arendt-Nielsen L, Curatolo M. Conditioned pain modulation in patients with acute and chronic low back pain. *Clin J Pain* 2016;32:116–21.
- [21] Goubert D, Danneels L, Graven-Nielsen T, Descheemaeker F, Meeus M. Differences in pain processing between patients with chronic low back pain, recurrent low back pain, and fibromyalgia. *Pain Physician* 2017;20:307–18.
- [22] Yarnitsky D, Crispel Y, Eisenberg E, Granovsky Y, Ben-Nun A, Sprecher E, Best LA, Granot M. Prediction of chronic post-operative pain: pre-operative DNIC testing identifies patients at risk. *Pain* 2008;138:22–8.