Thesis for the degree of *Doctor philosophiae* (dr. philos.) submitted to the
Faculty of Medicine, University of Oslo

Re-appraisal of prognostic factors in primary gastrointestinal stromal tumour. Refining definitions.

TOTO HØLMEBAKK

Department of Abdominal and Paediatric Surgery
The Norwegian Radium Hospital | Oslo University Hospital

© Toto Hølmebakk, 2021

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-889-2

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

In memory of my parents

LIST OF PAPERS

PAPER I

Hølmekbakk T, Bjerkehagen B, Boye B, Bruland ØS, Stoldt S, Sundby Hall K. Definition and clinical significance of tumour rupture in gastrointestinal stromal tumours of the small intestine. *Br J Surg* 2016; **103**: 684-691.

PAPER II

Hølmekbakk T, Hompland I, Bjerkehagen B, Stoldt S, Bruland SØ, Sundby Hall K, Boye K. Recurrence-free survival after resection of gastric gastrointestinal stromal tumors classified according to a strict definition of tumor rupture: a population-based study. *Ann Surg Oncol* 2018; **25**: 1133-1139.

PAPER III

Hølmekbakk T, Bjerkehagen B, Hompland I, Stoldt S, Boye K. Relationship between R1 resection, tumour rupture and recurrence in resected gastrointestinal stromal tumour. *Br J Surg* 2019; **106**: 419-426.

PAPER IV

Hølmekbakk T, Bjerkehagen B, Lobmaier IVK, Hompland I, Stoldt S, Boye K. Is peritoneal tumor penetration of prognostic importance in gastrointestinal stromal tumors? *Ann Surg Oncol* 2019; **26**: 4730-4736.

PAPER V

Boye K, Berner JM, Hompland I, Bruland ØS, Stoldt S, Sundby Hall K, Bjerkehagen B, Hølmekbakk T. Genotype and risk of tumour rupture in gastrointestinal stromal tumour. *Br J Surg* 2018; **105**: e169-e175. doi:10.1002/bjs10743.

PAPER VI

Hølmekbakk T, Wiedswang AM, Meza-Zepeda LA, Hompland I, Lobmaier IVK, Berner JM, Stoldt S, Boye K. Integrating anatomical, molecular and clinical risk factors in gastrointestinal stromal tumor of the stomach. *Ann Surg Oncol*. In press.

ABBREVIATIONS

ACOSOG	American College of Surgeons Oncology Group
AFIP	Armed Forces Institute of Pathology (US)
AIO	Arbeitsgemeinschaft Internistische Onkologie (Ge)
AJCC	American Joint Committee on Cancer
c.i.	confidence interval
CT	computed tomography
del557/558	<i>KIT</i> exon 11 deletions and insertion-deletions involving codons 557 and 558
DFS	disease-free survival
DOG1	discovered on GIST-1 (immunohistochemical marker)
EORTC	European Organisation for Research and Treatment of Cancer
ESMO	European Society for Medical Oncology
EUS	endoscopic ultrasonography
FDG-PET	fluorodeoxyglucose positron emission tomography
FNCLCC	Fédération nationale des centres de lutte contre le cancer (Fr)
GIST	gastrointestinal stromal tumour
HPF	high power field (of the microscope)
HR	hazard ratio
NCCN	National Comprehensive Cancer Network (US)
NCI	National Cancer Institute (US)
KIT	derived from <i>kitten</i> ; detected in a virus causing sarcoma in infected cats (<i>gene/protein</i>)
MI	mitotic index
mNIH	modified National Institutes of Health (classification system)
MRI	magnetic resonance imaging
NF1	neurofibromatosis type 1 (<i>gene/protein/disease</i>)
NGS	next-generation sequencing
NIH	National Institutes of Health (US)
OS	overall survival
<i>P</i>	probability (of retaining the null hypothesis)
PDGFRA	platelet-derived growth factor receptor- α (<i>gene/protein</i>)
PFS	progression-free survival

PP	peritoneal penetration
PR	partial response
R0	no residual tumour (after resection)
R1	possible residual tumour (microscopically involved surgical margins)
R2	macroscopic residual tumour (after resection)
RECIST	response evaluation criteria in solid tumours
RFS	recurrence-free survival, relapse-free survival
RR	relative risk
SD	stable disease
SDH	succinate dehydrogenase (<i>gene/protein</i>)
SSG	Scandinavian Sarcoma Group
TKI	tyrosine kinase inhibitor
TNM	tumour node metastasis (classification system)
UICC	Unio internationalis contra cancrum/Union for International Cancer Control
US	ultrasonography, ultrasound; United States of America

ACKNOWLEDGEMENT

Two persons have contributed to this study more than anybody else, in widely different ways: Kjetil Boye, Senior Consultant at the Department of Oncology and researcher at the Department of Tumour Biology, and Stephan Stoldt, my colleague at the Section for Surgical Oncology. Kjetil Boye has compensated for all my shortcomings in molecular biology, scientific analysis and methodology, while acknowledging, in a most generous fashion, my experience in surgery. We have been in almost daily contact for years and our cooperation has been an unalloyed pleasure. Stephan Stoldt is a surgeon with an exceptionally analytical attitude towards all aspects of surgical oncology. Probably without being aware of it himself, he is the origin of ideas pursued in these investigations.

My co-authors, besides their invaluable participation in these studies, are equally appreciated for their daily efforts in the Oslo Sarcoma Group, headed by Kirsten Sundby Hall, along with other members of various specialities and professions. Øyvind S. Bruland, Professor and Senior Consultant oncologist, has been my guide in the academic maze.

Stein G. Larsen, Head of the Section for Surgical Oncology and a genuinely kind person, accorded sabbatical leave to complete this work. Being part of his team of accomplished surgical oncologists inspires both humility and pride.

On this occasion I would also like to extend my gratitude to former colleagues at Aker Hospital: Thorstein B. Harbitz, who introduced me to surgical oncology, and Arild Nesbakken, the first truly academic surgeon I encountered in my career.

My wife Rita, a physician and researcher herself, has been a persistent believer in this project. Without her encouragement and counsel it may well have ended where it once started, on a scrap of paper with a pencilled suggestion for a definition of tumour rupture in GIST, long lost by now.

During the period I was engaged in these studies, I lost both my parents. None of them were scientifically minded; my mother, besides so much else, was a translator, my father a literary

publisher. They dealt with words, their meaning and significance. This thesis is dedicated to their memory.

Oslo, December 2020

TH

CONTENTS

List of papers 4

Abbreviations 5

Acknowledgements 7

Introduction 11

Definitions in clinical medicine 11

The concept of radicality in surgical oncology 12

PART ONE: BACKGROUND

Gastrointestinal stromal tumour: an outline 15

Epidemiology and clinical presentation 15

Pathogenesis and molecular biology 16

Diagnosis and staging 18

Risk stratification 18

Surgery for primary disease 22

Tyrosine kinase inhibitors 23

Adjuvant treatment 25

Neoadjuvant treatment 27

Medical treatment for metastatic disease 30

Surgery for metastatic disease 31

Established and controversial prognostic factors in primary GIST 33

Age and sex 33

Tumour size 33

Mitotic index 34

Tumour site – anatomical location 34

Tumour rupture 35

Genotype 38

R1 resection 38

Percutaneous biopsy 39

PART TWO: THE STUDY

Aims 42

Patients and methods 43

Surgical treatment	43
Adjuvant and neoadjuvant treatment	45
Follow-up and recurrence	45
Ethical approval	46
Statistical analysis	46
Issues pertaining to the individual studies	46
Summary of results and implications	50
Paper I	50
Paper II	51
Paper III	52
Paper IV	53
Paper V	55
Paper VI	56
Methodological discussion	58
Retrospective cohort studies	60
Time dependent variation	60
Endpoints in survival analyses	61
The confounding role of adjuvant treatment	62
Mutation analysis	62
Statistical analysis	63
Follow-up	64
General discussion	65
Tumour rupture in GIST	65
Minor defects of tumour integrity	69
Tumour genotype	72
Anatomical gastric subsite and tumour growth pattern	73
Adjuvant treatment in gastric GIST	75
The enigma of peritoneal recurrence in GIST	76
Conclusions	77
References	78
Papers I to VI	
Errata	

INTRODUCTION

DEFINITIONS IN CLINICAL MEDICINE

In 1791, the French National Convention introduced the precursor of the metric system, *le mètre*. They defined its length as one 10^{7th} of the distance from the North Pole to the equator, measured along a line passing through Paris, and a prototype was made in platinum. In 1889, a second prototype was consigned to the headquarter of the *Conférence général des poids et mesures* in Sèvres. Since, the definition has changed, and one meter is now defined as the distance travelled by light in empty space in one 299 792 458th of a second. However, the length has not changed, only the definition has been made more accurate.

A current definition of pain is ‘an unpleasant sensory or emotional experience associated with, or resembling that associated with, actual or potential tissue damage’ ¹.

The definition of the meter is an *operational* definition referring to a *method* and based on *agreement*; it has to be explained or instructed, otherwise it cannot be perceived; and refining it makes sense. The definition of pain is a *semantic* definition referring to *sense*, based on *experience* or some sort of *inference*. Without ever having heard the definition proposed by the International Association for the Study of Pain quoted above, almost anyone would be able to define pain, however awkwardly. Refining such a definition often does not make much sense.

Definitions in clinical medicine are in general neither operational nor semantic but share elements of either. They are *pragmatic*, made for a purpose - tools. The criteria of the *Diagnostic and Statistical Manual of Mental Disorders* ², preposterous as some of them may seem to the surgeon, are crucial for communication among professionals, science, prognostication, therapy, reimbursement etc., in the same manner as are criteria for the diagnosis of myocardial infarction, despite strikingly different epistemological bases ^{3,4}.

Not only do definitions change over time, also their significance changes. The Beaufort Wind Force Scale was developed during the 19th century and defines wind speed by its related effects at sea and on land. At a time when all international trade was carried out by sail ships, the difference between a high wind and a gale was a significant one. Today, nobody cares much. Surgery is still a prerequisite for cure in most solid malignancies. Hopefully, less

invasive therapies will evolve. Then, the definition of tumour rupture will probably be as irrelevant as the twigs breaking off, defining a fresh gale.

THE CONCEPT OF RADICALITY IN SURGICAL ONCOLOGY

Surgical oncologists manage macroscopic disease with mechanical measures. What they see with their eyes and feel with their fingers, or what is made visible by diagnostic imaging, they try to remove completely. With this treatment alone, two out of three patients with localised solid tumours - by very approximate estimation - will be cured. When malignant disease recurs after surgery it is in the form either of a local recurrence or distant metastasis. A local recurrence is generally considered a surgical failure whereas distant metastasis, even when occurring years after the extirpation of the primary tumour, reflects hidden disease at the time of diagnosis. Manipulation of the tumour during surgery has been considered a possible cause of systemic dissemination but remains a theoretical issue ^{5,6}; however, in gastrointestinal and retroperitoneal malignancies, it is accepted that spontaneous or iatrogenic spillage of tumour cells to the abdominal cavity induces recurrent peritoneal disease ⁷⁻¹⁰. Radical surgery therefore rests on the concept that the primary tumour is localised and that resection should be performed completely without iatrogenic spillage of malignant cells.

That solid tumours represent a confined process was distrusted already by Halsted, one of the pioneers of modern surgery. Guided by a notion about centrifugal spread he introduced the radical mastectomy for breast cancer in 1894, an extremely wide – and disfiguring – resection whereby the breast, axillary and pectoral lymph nodes, and the pectoral muscles were removed ¹¹. This operation remained the standard of care until the 1950s. Epithelial malignancies spread by way of the lymphatics, and the extent of lymph node dissection is a continuous debate in surgical oncology. For decades, upper gastrointestinal surgeons have argued about local lymphadenectomy (D1), regional lymphadenectomy, (D2) and extended regional lymphadenectomy (D3) for gastric cancer ^{12,13}, and in colonic cancer, having settled for D2 lymphadenectomy, the surgical community now moves towards D3 dissection ¹⁴. At the core of this discussion is the question whether surgical removal of affected regional lymph nodes prevents recurrence and improves prognosis. Breast cancer surgery has moved in the opposite direction of gastrointestinal cancer surgery. From the radical mastectomy advocated by Halsted, via the modified mastectomy, breast conserving treatment with adjuvant radiotherapy and sentinel node biopsy has become standard. Within this frame, lymph node

removal is a *staging* procedure, guiding further adjuvant therapy. This is indeed also the case in most gastrointestinal cancer, in gynaecological cancer, and in lung cancer. Lymph node metastasis is a proof of locally advanced disease and a predictor of distant metastasis which is best treated with oncological measures - radiotherapy and/or chemotherapy for local control, medical treatment for the prevention of distant relapse. Thus, adjuvant or neoadjuvant regimens are integrated in guidelines for the treatment of most cancer types. However, the modest effect on overall survival should be emphasised, only in the order of 10 per cent.

Sarcomas rarely metastasise to lymph nodes and lymphadenectomy is not standard. Therefore, in surgery on abdominal organs, vessels can be spared and complete excisions achieved with local or restricted segmental resections. Contrary to carcinomas, which grow by infiltration, mesenchymal tumours tend to have an expansive growth pattern, and close but careful dissection is characteristic of abdominal sarcoma surgery. Traditionally, only directly involved neighbouring organs are removed ¹⁵. In retroperitoneal sarcoma there are surgeons who advocate ‘compartmental’ resection of contiguous organs irrespective of signs of infiltration ^{16,17}. However, only dispensable organs can be removed even with a compartmental strategy, and for large abdominal or retroperitoneal sarcomas, the end result is usually a marginal excision.

For the individual surgical oncologist, the ability to achieve a complete resection is a personal matter, and most harbour a ‘margin complex’. The description of surgical margins was systematised in the late 1970s, by Enneking for mesenchymal tumours of the extremities and trunk wall ¹⁸ and by the Union for International Cancer Control (UICC)/American Joint Committee on Cancer (AJCC) for solid tumours in general ¹⁹. These systems, which are not interchangeable, make distinctions between a safe resection, a failed resection, and an ambiguous resection. The latter, termed ‘R1’ in the residual tumour classification ²⁰, indicates a macroscopic complete resection with microscopically involved margins and possible residual disease. For a number of solid tumours, carcinomas and sarcomas alike, an R1 resection is a predictor of poor outcome compared to an R0 resection ²¹⁻²⁶. However, there are tumours in which an R1 resection does not unequivocally confer an inferior prognosis ^{27,28}, and even when associated with an increased risk of recurrence, R1 resection may be far from an omen of certain relapse. Clearly, there is more to this than mechanics.

Myriads of cancer cells from the primary tumour enter the circulation daily ²⁹, by active invasion or passive shedding. In circulation, they are short-lived and difficult to detect, but

circulating tumour cells have been demonstrated in more than 90 per cent of women with non-metastatic breast cancer ³⁰. How and why some of these cells extravasate is poorly understood, but they may do so even at a very early stage ³¹. However, these disseminated tumour cells do not necessarily progress to clinically detectable metastases. In glioblastoma multiforme, a disease that hardly ever gives rise to extracranial metastasis, circulating tumour cells have been found in a considerable fraction of patients ³², and in immunocompromised organ recipients with glioblastoma, distant metastasis occurs ³³, indicating dormant cells that normally do not proliferate. An enigmatic trait of distant metastasis is its organotropism – the predilection for certain organs of metastasis from a given type of cancer. In distant tumour colonisation, there is interaction between the invasive cell and local factors at a specific site. This secondary site – often referred to as the premetastatic niche - does not receive the intrusive cell passively; it is primed by signals from the primary tumour itself ³⁴.

It is hard to believe, at least for the author of this thesis, that such an extremely complex picture of dissemination, extravasation, colonisation, and proliferation seen in haematogenous metastasis is not in some respect mirrored in iatrogenic, surgical dissemination of tumour cells. Gastrointestinal stromal tumours (GISTs) are special among solid malignancies, even among soft tissue sarcomas of which they constitute the majority. Most of them are slowly-growing, but some are highly aggressive; they are prone to rupture; and they metastasise almost uniquely to the peritoneum and liver. The rest of this thesis is devoted to them.

PART ONE: BACKGROUND

GASTROINTESTINAL STROMAL TUMOUR: AN OUTLINE

GIST accounts for less than 1 per cent of gastrointestinal tumours, is less frequent than adenocarcinomas and lymphomas, but is the most frequent non-epithelial tumour of the digestive tract ³⁵. GISTs arise from the interstitial cells of Cajal or their progenitors, which are situated along the autonomous nerves in the wall of the entire alimentary tract, where they have a pacemaker function ³⁶. Some 60 per cent arise in the stomach, approximately 30 per cent are situated in the small intestine and about 5 per cent are rectal ³⁷. GISTs of the oesophagus or colon are very rare. Extra-gastrointestinal GISTs, anatomically not connected with the gastrointestinal tract, form a minute subclass. Whether they have a genuine extra-gastrointestinal origin or at some point in their development have been secluded from the alimentary tract, is unsettled. GIST constitutes the largest group of sarcomas, approximately 20 per cent ³⁸, with a diversity in clinical behaviour ranging from small indolent nodules to rapidly growing, frankly malignant neoplasms. Consequently, until the discovery in the late 1990s that GISTs share a common and unique immunophenotype expressing the KIT antigen ³⁹, they were variously classified as benign, indeterminate or malignant mesenchymal – mostly myogenic - tumours, based on morphologic criteria (light microscopy).

EPIDEMIOLOGY AND CLINICAL PRESENTATION

The annual incidence is 7 to 19 cases per 1 million inhabitants as reported in Scandinavian studies ⁴⁰⁻⁴³. These are clinical cases; in autopsy series as many as 35 per cent among the elderly are found to harbour small, less than 1 cm, GISTs ⁴⁴. The true incidence is therefore largely unknown, and geographical discrepancies in incidence probably reflect differences in clinical and diagnostic practice more than epidemiological differences.

GISTs have an even sex distribution. They are commonly detected in middle and old age with a median age at diagnosis of 63 to 66 years ³⁷. Paediatric GIST accounts for less than 1 per cent of all cases and represent GIST syndromes - separate entities with different biomolecular aetiologies ⁴⁵. In sporadic GIST, no predisposing factors are known. There are publications indicating an association between GIST and other malignancies (genitourological and

gastrointestinal carcinomas)¹⁵⁻¹⁷. However, as a considerable proportion of GISTs are incidental findings during staging, surgery or follow-up of other cancers⁴⁶⁻⁴⁸, a detection bias is more than likely.

Gastrointestinal bleeding is a common presentation. Life-threatening haemorrhage and other surgical emergency states – bowel perforation or obstruction - do occur. Large tumours may cause discomfort and be detected as a palpable mass. Fatigue, wasting and general symptoms indicate disseminated disease, present in 10 to 20 per cent at diagnosis⁴⁹. Approximately 25 per cent have their GIST detected as an incidental discovery at imaging or endoscopy initiated for unrelated reasons⁴⁵. As stated above, GIST may be a finding made during evaluation or treatment of other malignancies, which is a fact that should be kept in mind when appraising studies on GIST where unspecific death is an endpoint in survival analyses.

PATHOGENESIS AND MOLECULAR BIOLOGY

In 85 to 90 per cent of patients, the initial oncogenic event is the gain-of-function mutation of the proto-oncogene *KIT* (75 per cent) or its homologous counterpart *platelet-derived growth factor receptor-α* (*PDGFRA*) (10 to 15 per cent)^{35,50,51}. *KIT* and *PDGFRA* mutations are mutually exclusive. These aberrations lead to constitutive, ligand-independent receptor activation and ultimately to apoptosis inhibition and cell proliferation. In primary GIST, *KIT* mutations most frequently involve exons 11 and 9; *PDGFRA* mutations involve exon 18, rarely exon 12 or 14^{52,53} (*Table 1*). Among the *KIT* exon 11 mutations, deletions affecting codons 557 and 558 (del557/558) deserve special mention as they are frequent and confer an aggressive phenotype; moreover, these tumours more often become imatinib resistant⁵⁴⁻⁵⁶. Among the *PDGFRA* mutated tumours, exon 18 D842V substitutions are most common, and contrary to other *PDGFRA* mutations, tumours harbouring this mutation do not respond to established tyrosine kinase inhibitors (TKIs)⁵⁷ (see section on TKI below).

Certain genotypes have their intrinsic phenotypical characteristics: the aggressiveness associated with the del557/558 genotype has already been mentioned; *KIT* exon 9 mutated GISTs are mostly non-gastric and are hyposensitive to imatinib⁵⁸; by contrast, *PDGFRA*-mutated tumours almost exclusively affect the stomach⁵². *KIT* exons 13 and 17 mutations are essentially secondary mutations in patients treated with TKIs^{59,60}.

In 10 to 15 per cent of GIST patients, no mutation in *KIT* or *PDGFRA* can be detected ⁴⁵. These tumours were commonly – and by many still are – referred to as ‘wild type’. Mutations or epigenetic changes in genes encoding one of the four subunits of the succinate dehydrogenase (SDH) complex can be found in approximately half of these patients ^{45,61}. Rare mutations in GIST involve *BRAF*, *HRAS*, *NRAS* or *PIK3CA* ^{45,62}.

The GIST syndromes Carney-Stratakis syndrome, Carney triad and neurofibromatosis type 1 (NF1) constitute separate entities. The Carney-Stratakis syndrome (gastric GIST and paraganglioma) affects young individuals, mostly females, and is caused by a germline mutation in one of the four *SDHs* encoding the SDH complex ⁶³. The Carney triad (multiple gastric GISTs, paraganglioma and pulmonary chondroma) likewise affects young females ⁶⁴. It is a non-heritable condition depending on epigenetic changes in *SDH* encoding the SDH(C) subunit. Contrary to *KIT* and *PDGFRA* mutated GISTs, SDH-deficient GISTs frequently metastasise to lymph nodes, and they are usually quiescent, even with lymphatic dissemination ⁶¹. Twenty-five per cent of individuals with NF1 have GIST at autopsy ⁶⁵. In these patients, tumours are typically small, multiple, and located in the duodenum or small intestine. The GIST syndromes are rare, comprising less than 5 per cent of GIST patients, and even rarer are germline mutations in *KIT* and *PDGFRA*.

Table 1. Approximate frequencies of genotypes in primary GIST

Mutation/deficiency	Frequency (per cent)	Comment
<i>KIT</i> mutation	75	
Exon 11	65	
del557-558	25	
Exon 9	10	Predominantly small intestinal
Exon 13	1+	
Exon 17	1-	
<i>PDGFRA</i> mutation	10-15	Predominantly gastric
Exon 18	10-15	
D842V	5-10	20 per cent of gastric GIST
Exon 12	1+	
Exon 14	1-	
SDH deficient	5+	90 per cent of paediatric GIST
<i>NF1</i> mutation	1+	
<i>BRAF</i> mutation	1-	
<i>KIT/PDGFRA</i> germline mutation	1-	Familial GIST, gastric

PDGFRA, platelet-derived growth factor receptor- α ; *SDH*, succinate dehydrogenase; *NF1*, neurofibromatosis type 1

DIAGNOSIS AND STAGING

Histologically, the diagnosis relies on immunohistochemical staining for KIT (positive in 95 per cent) and anoctamin-1 (DOG1, positive in 98 per cent) in tumours morphologically compatible with GIST ⁶⁶. Most GISTs have a characteristic appearance on imaging (contrast-enhanced CT), and in patients who are not surgically at risk, resection can be performed without biopsy when restricted procedures are sufficient, e.g. local gastric excision or small bowel resection. With diagnostic uncertainty, in frail patients or when surgery entails major resections, biopsy is mandatory. In the oesophagus, stomach and rectum, this can be done by the endoscopic route, ultrasound (US) guided if necessary; small intestinal tumours can be reached and safely punctured by US guided core needle. Mutation analysis is required only when neoadjuvant medical treatment is considered.

Liver and peritoneal metastases are excluded by abdominal and pelvic CT (contrast-enhanced, portal venous phase). Isolated lung metastases are exceedingly rare in GIST ⁶⁷, and chest X-ray is sufficient in staging, although in Norwegian practice, X-ray is likely to be superseded by CT. Magnetic resonance imaging (MRI) provides additional information in the preoperative assessment of rectal GIST. Fluorodeoxyglucose positron emission tomography (FDG-PET) has a limited role, mainly in the evaluation of the early response to TKI treatment ⁶⁸.

RISK STRATIFICATION

Risk stratification in GIST differs in important aspects from that of gastrointestinal malignancies in general. In most primary carcinomas of the alimentary tract, prognosis depends on three parameters: (i) clinical stage, as expressed by the TNM classification system ⁶⁹ assessing the primary tumour (T), nodal (N) and distant (M) metastasis; (ii) histological grade; and (iii) the completeness of surgical excision, reported according to the residual tumour (R) classification ²⁰. In GIST, risk of recurrence is stipulated in classification systems or nomograms that combine both clinical and histological variables. Contrary to other soft tissue sarcomas, GIST is not histologically graded. Mitotic index (MI), recorded per 50 high power fields (HPF) of the microscope, was chosen by Fletcher and colleagues in 2002 as the sole morphological variable and combined with tumour size in the National Institutes of Health (NIH) consensus criteria ⁷⁰, a four-tiered system (*Table 2*). An alternate model, published side by side in the same journal, includes tumour location (gastric or non-gastric) in

a three-tiered system ⁷¹. In the latter model, mitotic activity is divided into two categories only, below or above 5 per 50 HPF, whereas the former model has two cut-off values, at 5 and 10.

Table 2. NIH consensus criteria (Fletcher 2002 ⁷⁰)

Risk category	Tumour size (cm)	Mitotic index (per 50 HPF)
Very low	<2	<5
Low	2-5	<5
Intermediate	<5	6-10
	5-10	<5
High	>10	Any
	Any	>10

NIH, National Institutes of Health; HPF, high power field of the microscope. Note that for mitotic index, the value 5 is not included in any risk category.

Based on 1 939 patients followed long-term, Miettinen and Lasota of the US Armed Forces Institute of Pathology (AFIP), in 2006 introduced an elaborate version of the alternate NIH criteria ⁷². In addition to MI (with one cut-off value at 5) and tumour size, they considered anatomical location more specifically: stomach, duodenum, jejunum or ileum, and rectum. These factors were assembled in a complex table yielding eight categories and five tiers, the AFIP classification (*Table 3*).

Table 3. The AFIP classification (Miettinen-Lasota 2006 ⁷²)

Group	Tumour parameters		Risk category			
	Size (cm)	Mitotic index (per 50 HPF)	Gastric	J/I	Duodenal	Rectal
1	≤2	≤5	None	None	None	None
2	>2-5	≤5	Very low	Low	Low	Low
3a	>5-10	≤5	Low	-	-	-
3b	>10	≤5	Moderate	High	High	High
4	≤2	>5	-	-	-	High
5	>2-5	>5	Moderate	High	High	High
6a	>5-10	>5	High	High	-	-
6b	>10	>5	High	High	High	High

AFIP, Armed Forces Institute of Pathology; HPF, high power field of the microscope; J/I, jejunal or ileal. Blank spaces indicate anatomical locations with small number of patients.

In 2008, Joensuu proposed a modification of the NIH consensus criteria (mNIH) that included anatomical location (gastric or non-gastric), specified the cut-off value for mitotic activity, and added a fourth variable – tumour rupture ⁷³. In the original publication, non-gastric

tumours between 2 and 5 cm with more than 5 mitoses per 50 HPF could be classified as either intermediate or high-risk. This ambiguity was amended in a later version where non-gastric tumours were excluded altogether from the intermediate category ⁷⁴. *Table 4* shows the original partition.

Table 4. Modified NIH consensus criteria (Joensuu 2008 ⁷³)

Risk category	Tumour size (cm)	Mitotic index (per 50 HPF)	Tumour site	Rupture
Very low risk	≤2	≤5	Any	No
Low risk	>2-5	≤5	Any	No
Intermediate*	>2-5	>5-10	Gastric	No
	<5	6-10	Any	No
	>5-10	≤5	Gastric	No
High	>2-5	>5	Non-gastric	No
	>5-10	≤5	Non-gastric	No
	>5	>5	Any	No
	>10	Any	Any	No
	Any	>10	Any	No
	Any	Any	Any	Yes

NIH, National Institutes of Health; HPF, high power field of the microscope. *Non-gastric tumours between 2 and 5 cm with mitotic index above 5 could be classified as either intermediate or high-risk, amended in a later version.

The UICC/AJCC TNM classification also comprises a section on GIST ⁶⁹. Here, the T category simply reflects size (three cut-off values), and the N category is included – *pro forma*, as lymph node metastasis hardly ever occurs in sporadic GIST. However, MI is added to the system with one cut-off at 5 mitoses per 5 mm². Thus, the variables are those of the alternate NIH consensus criteria ⁷¹, but they are combined in the TNM fashion into stages: six stages for gastric tumours, five stages for non-gastric tumours.

These classification systems have been validated in a number of studies, and several modifications have been proposed ⁷⁵. Neither the NIH consensus criteria nor the AFIP classification do discriminate risk of recurrence between their two low-risk categories, and the high-risk categories of either system are heterogeneous, encompassing approximately 50 per cent of the patients ⁷⁶. The mNIH criteria identify patients at high risk and are therefore suited for selecting patients for adjuvant imatinib treatment. However, they do not differentiate well between the intermediate and low-risk groups; here, the AFIP classification performs better. Sensitivity and accuracy (for predicting recurrence) were assessed in a study by Yanagimoto and colleagues ⁷⁷. The mNIH criteria showed the highest sensitivity, 93 per cent, whereas

accuracy was highest by the UICC/AJCC TNM classification, 83 per cent. Yet for some reason, the TNM classification seems not to appeal to the GIST community, which sticks to the AFIP classification and the mNIH criteria.

Stratification implies categorising patients into broad groups – strata –, and small differences on either side of a cut-off value may allocate patients with almost similar prognoses to very different risk categories. Stratification systems are therefore adapted to scientific purposes rather than individual prognostication. Prognostic nomograms are two-dimensional diagrams using statistical models that incorporate variables with additive prognostic information to predict outcomes in individual patients. Such a nomogram, including the variables tumour size, MI, and anatomical location (stomach, colon/rectum or small intestine), was created by Gold and colleagues and published in 2009 ⁷⁸. Based on a small number of patients from the Memorial Sloan-Kettering Cancer Center, it was validated in two external cohorts and compared favourably to both the NIH classifications, concordance probabilities being similar to those of the AFIP classification. However, the nomogram uses MI not as a continuous but as a categorical parameter and underestimates RFS in several studies ^{79,80}. More recently, Joensuu and colleagues ³⁷ presented another diagram, a contour map constructed with data from 2 560 patients, using Bayesian statistics and non-linear modelling of tumour size and mitoses, taking account also of anatomical location and tumour rupture (*Figure 1*). Finally, sex, tumour size, MI and anatomical location (gastric or non-gastric) are variables included in a nomogram devised by Bischof and colleagues ⁸¹.

Based on data from patients treated with surgery alone, none of these models applies with equal accuracy to patients who have received adjuvant therapy. Such a model has been developed with data from the Scandinavian Sarcoma Group (SSG) XVIII trial and estimates the probability of recurrence-free survival (RFS) in high-risk patients who have received adjuvant imatinib for 1 or 3 years ⁸².

Figure 1. Prognostic nomogram for estimating the risk of recurrence after complete excision of primary GIST without adjuvant imatinib. [Printed with permission from Elsevier³⁷]

SURGERY FOR PRIMARY DISEASE

All GISTs larger than 1 cm are considered to have a proliferative and metastatic potential, and complete removal is the only curative measure. Tumours less than 1 cm – so called micro-GISTs – harbour the same oncogenic mutations^{48,83}, but when located in the stomach they do not progress, and excision or surveillance is stated to be unnecessary⁸⁴. In gastric and oesophageal tumours less than 2 cm, the metastatic potential is negligible, and active surveillance (CT or endoscopy/EUS) is justified. However, in young individuals with gastric lesions, the costs and inconvenience of long-term surveillance must be weighed against the low risk associated with surgery.

The aim of surgery is the complete excision of the tumour without undue loss of gastrointestinal function. For all practical purposes, GIST in adults does not metastasise to

lymph nodes, and organ-preserving procedures will suffice in most instances. In certain anatomical locations, e.g. the gastroesophageal junction, the second part of the duodenum, and the rectum, formal resections are sometimes required, and with tumour involvement of neighbouring structures, multivisceral resections, reported in 15 to 20 per cent ^{85,86}. Neoadjuvant treatment with imatinib, discussed in the following section, probably reduces the extent of surgery in some of these patients.

Minimally invasive techniques in oncological surgery, hotly debated at the turn of the century, have been established for colorectal cancer, upper gastrointestinal cancer and many hepatobiliary and pancreatic procedures. In GIST, it is still a matter of some controversy. A meta-analysis comparing open and laparoscopic surgery, including 2 140 patients, found that RFS was longer for patients treated with laparoscopy, who also benefited from the usual short-term advantages - less intraoperative blood loss, reduced time to oral diet, shorter hospital stay etc. ⁸⁷ However, none of the studies were randomised controlled trials, there was significant heterogeneity between the studies, and tumours were smaller in the laparoscopic group. Surgical techniques are continuously evolving, and so is the proficiency of the surgeons who practise them. Mini-invasive techniques are essentially new ways of gaining surgical access, and compared to the primary objective in oncological surgery - the safe removal of the malignant lesion – choices of access and instruments are, in the view of this author, of secondary importance.

Endoscopic excision, another minimally invasive procedure, is an established treatment for small and medium-sized gastric GISTs, mostly practised in East Asian countries ⁸⁸. The procedure can be performed by submucosal dissection, muscular dissection, or full-thickness wall resection, laparoscopically assisted or not. Recurrence rates of 0 to 6 per cent have been reported ^{88,89}. For GISTs of the lower rectum, transanal endoscopic microsurgery has emerged as an alternative to perineal or abdominal procedures.

TYROSINE KINASE INHIBITORS

Tyrosine kinases (TKs) phosphorylate tyrosine in proteins by transferring a phosphate group from ATP to tyrosine, thereby influencing intracellular signalling and eventually cellular proliferation, differentiation and apoptosis. The majority are receptor TKs situated at the level of the cellular membrane, where KIT and PDGFRA bind the ligands stem cell factor or

platelet-derived growth factor. When aberrant, subsequent to genetic or epigenetic changes, TKs are possible targets for specific therapy. Such an aberration is the BCR-ABL expressed in chronic myelogenous leukaemia, a disease which responds to the TKI imatinib. In 2001, imatinib was tried for the first time in a patient with metastatic GIST, and her immediate response revolutionised therapy ⁹⁰. In GIST, imatinib is a competitive inhibitor of the receptor TK KIT (or PDGFRA) (see section on pathogenesis and molecular biology above). Since 2001, a number of TKIs have been introduced for the treatment of metastatic GIST - sunitinib, regorafenib, nilotinib, pazopanib, sorafenib among others, but imatinib is by far the most effective and best tolerated and the only TKI approved for adjuvant and neoadjuvant treatment. However, there are genotypes that are associated with reduced sensitivity or resistance to imatinib, notably mutations in exon 18 D842V (*Table 5*). For these patients, avapritinib has recently been approved in the US, and crenolanib and ripretinib are promising agents which are currently being tested in randomised trials.

Table 5. Genotypes and their associated sensitivity to imatinib

Genotype	Sensitivity
<i>KIT</i> mutation	
Exon 11	Sensitive
Exon 9	Hyposensitive, double dose recommended in metastatic disease
Exon 13	Sensitive as primary mutation, resistant as secondary mutation
Exon 17	Uncertain as primary mutation, resistant as secondary mutation
<i>PDGFRA</i> mutation	
Exon 18 D842V	Resistant
Exon 18, others	Sensitive
Exon 12	Sensitive
Exon 14	Uncertain
SDH-deficient	Mostly resistant
<i>NF1</i> mutation	Mostly resistant
<i>BRAF</i> mutation	Resistant

PDGFRA, platelet-derived growth factor receptor- α ; SDH, succinate dehydrogenase; *NF1*, neurofibromatosis type 1.

Imatinib will be further treated in the following sections on adjuvant and neoadjuvant therapy; other TKIs in the section on metastatic disease.

ADJUVANT TREATMENT

The efficacy and tolerability of adjuvant imatinib in GIST have been explored in eight trials ⁹¹ of which these four are the most important:

1. The American College of Surgeons Oncology Group (ACOSOG) Z9001 trial ⁹² enrolled 713 patients with primary GIST at least 3 cm in size completely excised. Participants were randomised to 1 year of imatinib 400 mg once daily or to placebo. The primary endpoint was RFS; overall survival (OS) a secondary endpoint. RFS was longer for participants in the treatment arm, 98 per cent *versus* 83 per cent at 1 year; hazard ratio (HR) 0.35 (95 per cent confidence interval [c.i.] 0.22 to 0.53; $P < 0.001$). OS was similar. These results persisted at follow-up at median 74 months, HR for RFS 0.6 (95 per cent c.i. 0.43 to 0.75; $P < 0.001$). OS remained similar in both arms.
2. The Scandinavian Sarcoma Group XVIII/Arbeitsgemeinschaft Internistische Onkologie (SSGXVIII/AIO) trial ⁹³ enrolled 397 patients with primary high-risk GIST (mNIH). Participants were randomised to 1 or 3 years of imatinib 400 mg once daily. The primary endpoint was RFS; OS a secondary endpoint. RFS was improved for participants treated for 3 years, 71 *versus* 52 per cent at 5 years; HR 0.60 (95 per cent c.i. 0.44 to 0.81; $P < 0.001$); OS was marginally better after 5 years, 92 *versus* 85 per cent; HR 0.60 (95 per cent c.i. 0.37 to 0.97; $P = 0.036$) ⁹⁴. Ten-year OS was 79 *versus* 65 per cent; HR 0.55 (95 per cent c.i. 0.37 to 0.84; $P = 0.004$) ⁹⁵.
3. The European Organisation for Research and Treatment of Cancer/Soft Tissue and Bone Sarcoma Group Intergroup (EORTC/Intergroup) trial ⁹⁶ enrolled 835 patients with primary GIST completely excised, classified as high or intermediate risk according to the NIH consensus criteria. Participants were randomised to 2 years of imatinib 400 mg once daily or to observation. The primary endpoint was OS, amended to imatinib failure-free survival (IFFS), a surrogate for OS; RFS was a secondary endpoint. There was no difference in IFFS at 5 years, 87 *versus* 84 per cent; HR 0.79 (98.5 c.i. 0.50 to 1.25; $P = 0.21$). RFS at 5 years was improved for participants in the treatment arm, 69 *versus* 63 per cent; $P < 0.001$.
4. The PERSIST-5, a phase II, single arm trial ⁹⁷, enrolled 91 participants with GIST at intermediate or high risk according to the AFIP classification, completely excised. Participants received imatinib 400 mg once daily for 5 years. Forty-five (49 per cent) patients discontinued treatment. One patient relapsed during treatment (*PDGFRA*

D842V mutation); 6 patients relapsed after discontinuation. Estimated 5-year RFS was 90 per cent (95 per cent c.i. 80 to 95 per cent).

The European Society for Medical Oncology (ESMO), the National Comprehensive Cancer Network (NCCN), and Asian guidelines all recommend adjuvant imatinib for 3 years⁹⁸⁻¹⁰⁰. Imatinib is reasonably well tolerated with adverse events grade 3 or 4 reported in 20 to 50 per cent^{93,97}. Nevertheless, a substantial number of patients discontinue treatment at a drop-out rate increasing from 13 to 27 per cent after 1 year to 49 per cent after 5 years^{92,93,97}. Several important issues are still unsettled:

- (i) *At what estimated risk should adjuvant therapy be recommended?* The SSGXVIII/AIO trial recruited patients at high risk according to the mNIH criteria. This group comprises approximately 50 per cent of the patients in reference series^{37,77} and they have a risk of recurrence at 10 years of 64 per cent; in the intermediate group, the risk is only 13 per cent¹⁰¹. The intermediate group of the AFIP classification includes a larger proportion, approximately 35 per cent, who are at a higher risk of recurrence, 20 to 50 per cent at 10 years; in the high-risk group, the risk is 75 to 90 per cent¹⁰¹. The guidelines state that adjuvant therapy is recommended for patients at high risk and that patients at intermediate risk should be counselled individually. The multidisciplinary conference of the Oslo Sarcoma Group rarely recommends adjuvant treatment for patients with less than a 50 per cent risk of recurrence.
- (ii) *Should mutation status influence decisions on adjuvant treatment?* The trials mentioned above, being initiated shortly after the advent of imatinib, also included patients with imatinib insensitive tumours. Since, there is agreement that adjuvant therapy should not be offered to patients with tumours harbouring a *PDGFRA* D842V mutation or are *KIT/PDGFR*A wild type (*Table 5*). In metastatic disease, a higher dose of imatinib is recommended for GISTs that are *KIT* exon 9 mutated¹⁰², but whether this should apply in the adjuvant setting, is unclear. Neither in the ACOSOG trial nor in the SSGXVIII/AIO trial (both imatinib 400 mg daily) was RFS improved in the experimental arms in patients with *KIT* exon 9 mutated tumours. In Oslo, these patients receive 800 mg daily if tolerated. Subgroup analyses of the SSGXVIII/AIO trial indicate that only patients with *KIT* exon 11 mutant GISTs benefit from 3 years of treatment⁵³. Whether patients with non-exon 11 mutated tumours benefit from adjuvant therapy at all, is a legitimate

question. As a matter of fact, only patients with *KIT* exon 11 deletions improved their RFS in the ACOSOG Z9001 trial ¹⁰³.

- (iii) *Does adjuvant imatinib cure or just prolong time to recurrence?* The SSGXVIII/AIO trial is the only trial that has demonstrated improved overall survival with adjuvant treatment – by a small margin after 5 years' follow-up ⁹⁴, more convincingly after 10 years' follow-up ⁹⁵. Probably there are patients at high risk of recurrence who are cured by prolonged adjuvant imatinib therapy. These patients have to be identified, and the exploratory analysis of the SSGXVIII/AIO trial by Joensuu and colleagues, indicates *KIT* exon 11 del557/558 mutants with high mitotic activity ⁵³. There seems to be no difference in prognosis between an imatinib naïve patient who relapses and one who relapses after adjuvant treatment provided that the recurrence is detected at an early stage ⁹⁵, and in the opinion of this author, close follow-up without treatment would be an equally safe strategy for many high-risk patients.
- (iv) *Should adjuvant therapy for patients with tumour rupture be extended in time?* Tumour rupture carries a particularly sinister prognosis. In the SSGXVIII/AIO trial, patients with rupture had a poor outcome uninfluenced by the length of treatment ⁹⁴. This issue will be considered in the general discussion.

NEOADJUVANT TREATMENT

The spectacular response to imatinib in metastatic disease immediately prompted its use in the neoadjuvant setting. Several reports were published – small studies or case reports that included unresectable tumours, locally recurrent disease, metastasis, ordinary intermediate and high-risk tumours, and true locally advanced primary tumours – demonstrating that significant size reduction followed by complete resection could be achieved after imatinib treatment in a number of patients, with a reasonable prognosis in primary disease ¹⁰⁴. The ESMO and NCCN guidelines now recommend neoadjuvant imatinib in primary GIST where R0 resection is deemed unlikely at diagnosis, where response to treatment could facilitate less extensive surgery or reduce the risk of tumour rupture and complications ^{98,99}. There are no randomised trials to support these recommendations, and most likely there will never be. These four studies, comprising 285 patients, constitute the best evidence there is:

1. The North American RTOG (Radiation Therapy Oncology Group) 0132 phase II trial¹⁰⁵ recruited patients with locally advanced primary, recurrent or metastatic GIST. 'Locally advanced' was defined as tumours exceeding 5 cm. Patients were assigned to receive imatinib 600 mg daily for 8 to 12 weeks preoperatively and 2 years postoperatively. Thirty patients with primary disease were included. Median tumour size was 9 (range 5 to 25) cm. Mutational analysis was not performed. Only 2 patients (7 per cent) had partial response (PR) as assessed with RECIST; 83 per cent had stable disease (SD), and in 10 per cent, response was unknown. R0, R1, and R2 resections were performed in 77, 15, and 8 per cent, respectively. Tumour rupture was not reported. Four patients did not undergo surgery. The estimated 2-year progression-free survival (PFS) was 83 per cent, and 3 out of 9 recurrences occurred while on adjuvant imatinib therapy.
2. The APOLLON phase II trial¹⁰⁶, published in abstract only, recruited 41 patients with locally advanced non-metastatic GIST. Participants were assigned to receive imatinib 400 mg daily for 6 to 12 months before surgery; postoperative treatment was not part of the protocol. Tumour genotype was not considered. Median tumour size was 11 cm. Two patients progressed; 1 patient died from treatment related toxicity. Eighty-two per cent had PR; median tumour size was reduced to 5 cm; and 36 patients (88 per cent) proceeded to surgery. R0 resection was achieved in 30 patients (83 per cent). In 72 per cent, surgery was less extensive than envisaged before neoadjuvant treatment. Tumour rupture was not reported. PFS at 3 years was 85 per cent.
3. The retrospective EORTC STBSG (Soft Tissue and Bone Sarcoma Group) pooled analysis¹⁰⁴ included 161 patients with locally advanced GIST from 10 European centres. Patients received imatinib 400 mg daily for median 40 (range 6 to 190) weeks before surgery. Fifty-six per cent continued imatinib postoperatively for at least 1 year. Mutational analysis was not performed on biopsies. Thirty per cent were duodenal or rectal GISTs. PR rate was 80 per cent; only 2 patients progressed. R0 resection was achieved in 83 per cent. Tumour rupture was not reported. Estimated 5-year DFS was 65 per cent (95 per cent c.i. 59 to 71 per cent).
4. The Japanese-Korean UMIN00000003114 phase II trial¹⁰⁷ included 53 patients with gastric GIST at least 10 cm. Participants received imatinib 400 mg daily for 6 to 9 months preoperatively and for 1 year postoperatively. Mutational analysis was performed on biopsies, and 89 per cent were *KIT* exon 11 mutated GISTs. Median tumour size was 12 (range 10 to 23) cm. No patient progressed; PR rates were 62 and

98 per cent according to RECIST and the Choi criteria, respectively; median tumour regression was 35 (range 0 to 87) per cent. Three patients refused surgery. R0 resection was achieved in all but 2 patients. One patient had tumour rupture. Only 3 patients underwent a total gastrectomy, and at least half of the stomach was preserved in 84 per cent. Multivisceral resection was performed in 24 per cent. The 2-year PFS rate was 89 per cent (95 per cent c.i. 81 to 98 per cent).

Upon the administration of imatinib, the metabolic response is immediate, as can be recognised *in vitro*, on functional MRI or FDG-PET, or appreciated clinically ^{90,108}. However, volumetric response takes longer, as reflected in the low response rate reported in the RTOG 0132 trial (8 to 12 weeks of treatment) ¹⁰⁵. In tumours sensitive to imatinib, progression is exceedingly rare ¹⁰⁴, and the risk of secondary resistance is considered to be negligible within the first year. Guidelines recommend that surgery be performed when maximum reduction has been achieved, after 6 to 12 months. Contrary to other TKIs, imatinib does not interfere with surgery; it can be taken until the day before operation and resumed immediately afterwards, and complications do not seem to be increased ^{98,104}. Life-threatening tumour bleeding induced by imatinib has been reported in less than 5 per cent ¹⁰⁴.

Despite the lack of formal evidence for the benefit of its use, neoadjuvant therapy is intuitively sound in locally advanced disease. The recommendations are vague, and the decision whether to operate up front or after treatment lies in the end with the surgeon. The studies cited above indicate that neoadjuvant treatment facilitates the preservation of organs, which is important, both for gastrointestinal function, and in case of the stomach, for the uptake of imatinib in adjuvant treatment, or later in the event of recurrent disease. However, the extent of surgery also depends upon the individual surgeon – the surgeon's attitude, training, skills etc., often more than upon any response to preoperative treatment. Moreover, a substantial reduction in tumour size does not necessarily entail reduced involvement of the organ of origin. Finally, imatinib being more of a cytostatic than a cytotoxic drug, it cannot be taken for granted that an R0 resection after neoadjuvant therapy warrants complete removal of microscopically dormant disease, even though reported local recurrence rates are low ¹⁰⁴.

Imatinib decreases tumoural vascularisation, and it is stated that a pre-treated GIST is more resistant to iatrogenic rupture ¹⁰⁴. A pre-treated tumour assumes a characteristic gelatinous consistency, unmistakable in the surgical field, and, in the experience of this author, they are

still as prone to rupture. Any reduced risk of rupture would primarily be related to the reduction in size.

MEDICAL TREATMENT FOR METASTATIC DISEASE

At diagnosis, 10 to 20 per cent of GIST patients will have disseminated disease, with metastases primarily affecting the liver and/or the peritoneal cavity ^{49,109}. Twenty-six to 40 per cent relapse after radical surgery for primary disease ^{37,85}; moreover, a small number of patients will have primary tumours not amenable to surgery. For these patients, treatment with a TKI is the first option. GIST is virtually resistant to chemotoxic agents ¹¹⁰, and before the introduction of imatinib in 2001, median survival with metastasis was a grim 9 months ¹¹¹.

Imatinib is administered orally at a standard dose of 400 mg once daily, side effects are usually few and tolerable, and treatment should be continued until progression, which occurs after median 2 to 3 years ^{111,112}. However, there are patients who are still in complete remission after more than 15 years ¹¹³. Less than 15 per cent of patients with metastatic GIST have tumours with primary resistance to imatinib ^{111,112}. For patients with *PDGFRA* exon 18 D842V mutated GIST, avapritinib has been approved by the US Food and Drug Administration and other agents will probably soon be available. Patients who progress on imatinib will usually have developed secondary resistance, typically involving *KIT* exon 13 or 17 ^{59,60,114}. Some may benefit from a higher dose of imatinib, others will go on to second-line treatment, which is sunitinib, a multitarget inhibitor ¹¹⁵. Unfortunately, response rates are considerably lower for sunitinib than imatinib, and median PFS is approximately 6 months; toxicity is also more pronounced ¹¹⁵. For third-line treatment, only regorafenib has been approved, with median PFS approximately 5 months and even more toxic side-effects ¹¹⁶. In trials, several other TKIs have been tested as second or third-line agents, e.g. pazopanib, sorafenib, masitinib and dovitinib, with PFS rates in the range of 3 to 6 months ¹¹⁷⁻¹²⁰.

As seen, once treatment with imatinib fails, hope for long-time survival is lost. However, by dose escalation, new treatment lines, and even by re-challenging with imatinib, median survival for patients with metastatic GIST has now passed 6 years ^{95,113}.

SURGERY FOR METASTATIC DISEASE

The tremendous effect of imatinib has relegated surgery to an unsettled secondary role in metastatic GIST. A randomised phase III trial sponsored by EORTC, assigning patients who responded to imatinib to continued medical treatment alone or cytoreductive surgery plus medical treatment, was prematurely closed due to poor accrual (Clin. Gov: NCT00956072). Another randomised trial, enrolling only 41 patients with peritoneal metastasis, showed improved PFS and OS for patients undergoing surgery ¹²¹. However, the trial was designed for 210 patients, and underpowered, it also had other important methodological limitations. Prospective studies are lacking. Theoretically, cytoreductive surgery should delay secondary mutations and drug resistance, and in several retrospective series, factors associated with improved prognosis after surgery for metastasis have been identified ¹²²⁻¹²⁷. Fairweather and colleagues reviewed 323 patients undergoing surgery for disseminated disease while on treatment with TKI at two large US sarcoma centres ¹²⁷. Patients with radiographic responsive disease (RD) before surgery had better outcomes than patients with SD, unifocal progressive disease (UPD) or multifocal progressive disease (MPD): median PFS from date of surgery was 36, 30, 11, and 6 months, respectively. Correspondingly, median OS was not reached for patients with RD; with SD it was 110 months, with UPD 59 months, and with MPD 24 months. In another large US series, unifocal disease and short duration of TKI (less than 1 year) were predictive of improved outcomes ¹²⁶.

Patients subjected to cytoreductive surgery in retrospective series are highly selected, and whether their prognosis is influenced by surgery is impossible to say. The fact that patients on second-line TKI and longer duration of TKI preoperatively do worse, indicate that resistance to TKI is the limiting factor. Interestingly, An and colleagues retrospectively compared 35 patients who had undergone cytoreduction of at least 75 per cent to 214 patients with no cytoreduction: the cytoreduction group was younger, had a superior performance status, a smaller tumour burden, and presented with recurrent rather than primary metastatic disease; nevertheless, PFS was not superior ¹²⁸.

Surgery for recurrent and metastatic GIST is performed at all major sarcoma centres, doubtlessly to the benefit of some patients. These patients seem to be those with (i) responsive disease on first-line TKI (imatinib), (ii) short duration of TKI (6 to 24 months), (iii) unifocal or oligometastatic disease, and (iv) lesions amenable to R0/R1 resection ¹²⁹. Patients who have unifocal progression on imatinib may have second-line treatment delayed by surgery.

However, this respite is limited to less than 6 months ^{115,127}. Ablative treatment, i.e. radio frequency ablation or stereotactic radiation, is an alternative to surgery for some patients.

TKI should be continued after cytoreduction, irrespective of R status. Hompland and colleagues have demonstrated that patients with oligometastatic disease, i.e. less than four lesions, could be considered a particular subgroup with a favourable prognosis among metastatic patients ¹¹³. Patients with oligometastatic disease, treated with imatinib for at least 5 years, who have at some point had all lesions removed, are now enrolled in a prospective multicentre single-arm phase II trial where imatinib is discontinued, the SSGXXV trial (Clin. Gov: NCT02924714). The primary endpoint is 3-year PFS.

ESTABLISHED AND CONTROVERSIAL PROGNOSTIC FACTORS IN PRIMARY GIST

In the following chapter, the main clinical, pathological, and molecular factors of known or presumed influence on prognosis in localised GIST are presented. For an analysis of their importance, readers are also referred to the general discussion.

AGE AND SEX

Age does not seem to influence prognosis in sporadic GIST ^{37,74,77,86,130}. The only important study known to this author showing a slight (negative) impact of increasing age upon DFS in multivariable analysis, not considering Paper I, is the one published by Bischof and colleagues ⁸⁵ (HR 1.03 [95 per cent c.i. 1.01 to 1.06]; $P = 0.005$). By contrast, male sex has been associated with decreased DFS or RFS in large series ^{37,67,77,81,85,86}. Paper VI indicates that this could be explained by differences in RFS between males and females with gastric GIST. None of the prevailing prognostic systems or models includes sex as a variable.

TUMOUR SIZE

Tumour size, two-dimensionally reported in cm or mm, and erroneously referred to as ‘diameter’, should be conceived as a surrogate measure for tumour volume. It is a longstanding parameter in the staging of soft tissue sarcomas and carcinomas alike, although for gastrointestinal adenocarcinomas, depth of infiltration rather than size or volume indicates the stage of the primary tumour ⁶⁹. Size was chosen by Fletcher and colleagues ⁷⁰ as one of two variables in their initial classification of GIST and has remained one of the most robust prognostic indicators (*Figure 2A*). Its relationship with recurrence is non-linear, i.e. an increase in size exerts a larger prognostic impact at low levels ³⁷. In most series, size is a statistically independent factor ^{37,77,85,86,131}, and also in the adjuvant ACOSOG Z9001 trial ¹⁰³, size was independently related to prognosis; in the SSGXVIII/AIO trial, this analysis was not performed. However, increased size is often associated with an elevated MI. Still, large GISTs with a low MI are encountered, and in Paper VI, 34 per cent of gastric GISTs larger than 10 cm were recorded with a MI of 5 or less. Interestingly, in a study only including GISTs

exceeding 10 cm, Wada and colleagues found that tumour size was no longer independently related to RFS, MI was the only independent factor ¹³².

MITOTIC INDEX

Mitoses, readily identified by the trained pathologist upon light microscopy, indicate cell proliferation and are characteristic of malignant tumours. In the grading of soft tissue sarcomas, the presence of mitoses is incorporated in both the US NCI (National Cancer Institute) and the French FNCLCC (Fédération nationale des centres de lutte contre le cancer) systems ^{133,134}, expressed as a ratio of mitoses per 10 HPF – MI or mitotic count. MI was adopted by Fletcher and colleagues for the NIH consensus criteria along with tumour size, however, with an extended denominator, 50 HPF ⁷⁰. Selected *a priori*, it has repeatedly reclaimed its position as the most important prognostic indicator in GIST (*Figure 2B*). Compared with a MI less than 2, the risk of recurrence with an index above 10 increases more than 20 times; with an index between 6 and 10, the increase is 10-fold; and with an index between 2 and 5, 4-fold ³⁷. At 20 mitoses, the risk seems to reach a plateau ⁶⁷.

Microscopes differ. With a traditional field size, like with the ones originally used by Fletcher and Miettinen, 50 HPF equal an area of 5 mm² at a magnification of 40 times. With modern wide-field oculars, the same number of fields cover approximately 10 mm². It is therefore recommended that mitoses be reported per 5 mm², or, if the denominator 50 HPF is retained, that the number of fields be adjusted to the field size. Nevertheless, to count mitoses per 50 HPF still seems to be common practice, irrespective of field size ¹³⁵.

Areas with high mitotic activity should be selected for counting ⁷¹. Random needle biopsies or cell blocks are therefore not suited for assessing MI, and consequently, risk stratification cannot be done prior to surgery. In specimens from patients subjected to neoadjuvant imatinib, mitoses are reduced, often dramatically, with the result that proper staging is precluded.

TUMOUR SITE – ANATOMICAL LOCATION

That malignant tumours at different subsites within the same organ may behave differently has long been recognised: gastric carcinomas of the distal stomach and the cardia can be

considered separate entities ¹³⁶, and more recently ‘sidedness’ in colonic cancer has become a subject of research with important biological and clinical disparities between the right and left hemicolon becoming evident ¹³⁷. If the entire alimentary tract be regarded as the organ of origin of GIST, a similar approach could be applied.

The risk of recurrence for gastric GIST is roughly half the one for small intestinal GIST ^{37,77,85,138} (*Figure 2C*). Tumours of the small intestine tend to be larger and have a higher MI than tumours of the stomach ⁵², but tumours of similar size and mitotic activity still have a worse prognosis when located in the small bowel ¹³⁹. Rupture is more frequent in tumours of the small intestine ⁷⁴, but the difference in outcome remains when ruptures are excluded ³⁷. There is no explanation that sufficiently accounts for this discrepancy. There are important genetic variations, which will be discussed in a following section, but they do not provide a complete answer. Nor does the recurrence pattern, which is similar for gastric and small intestinal GISTs (Oslo sarcoma database, unpublished data), indicate possible explanations.

Duodenal GIST only accounts for approximately 5 per cent of all GISTs ^{48,72} and they are rarely analysed separately. Grouped with other small intestinal or non-gastric tumours their prognosis has been considered similar ^{72,101}. However, as Wozniak and colleagues point out, DFS is probably closer to gastric than small intestinal GISTs ⁵². Moreover, among duodenal tumours, the number of NF1 related GISTs is not negligible. The outcome for rectal GIST have been likened to those of the small bowel ³⁷, but with improved surgery after neoadjuvant imatinib good long-term results have been achieved, which suggests that tumours at this site may not be inherently as aggressive as previously regarded ^{76,140}.

TUMOUR RUPTURE

Originating outside the mucosa and growing expansively rather than by infiltration, GISTs may reach considerable dimensions. Their size, attenuated serosal lining, and fragile – often necrotic – consistency lend them prone to rupture into the peritoneal cavity, spontaneously or at operation, in a manner not encountered in carcinomas of the gastrointestinal tract. The term ‘rupture’ with respect to sarcomas was probably coined a long time ago, and Rutkowski and colleagues, in their important paper of 2007, introduced the concept as a variable in their retrospective assessment of 335 GIST patients, demonstrating its negative impact on DFS in multivariable analysis ⁶⁷. The parameter was then incorporated in the mNIH criteria ⁷³, before

Joensuu and colleagues, in their large study of pooled population-based cohorts published in 2012, re-stated its significance as a prognostic factor ³⁷ (Figure 2D).

Figure 2. Recurrence-free survival (years) related to established risk factors. **A** tumour size, **B** mitotic index, **C** anatomical site, **D** tumour integrity. [Printed with permission from Elsevier³⁷]

Widely accepted as a prognosticator, it has since been surprisingly hard to prove its independent impact (Table 6). The reason for this is probably three-fold: (i) retrospective data collection, (ii) under-reporting, and (iii) lack of an established definition. The definition of tumour rupture in GIST is one of the main subjects of the present thesis, and these issues will be addressed in the general discussion.

Table 6. Studies reporting tumour rupture and related prognosis in GIST

STUDY	CHARACTERISTICS	N	RUPTURE (per cent)	DEFINITION	PROGNOSTIC SIGNIFICANCE
Rutkowski 2007 ⁶⁷	2002–2006; Polish clinical GIST registry; retrospective on rupture	335	75 (22)	Unspecified, R1 resection included	In multivariable analysis
Takahashi 2007 ¹³⁰	1987-2003; Japan, Osaka region; retrospective on rupture	303	12 (4)	Unspecified	In univariable analysis
Rutkowski 2011 ⁷⁴	2001-2010; Polish Clinical Registry; retrospective on rupture	640	46 (7)	Spontaneous and intraoperative, otherwise unspecified	In univariable analysis only
Joensuu 2012 ³⁷	1972-2010; Merged data from European/Japanese studies; retrospective	1198	71 (6)	Unspecified	In multivariable analysis
Yanagimoto 2014 ⁷⁷	1980-2010; Japan, Osaka region; retrospective	711	34 (5)	Unspecified, adjacent infiltration included	In multivariable analysis
Wozniak 2014 ⁵²	1985-2012; ConticaGIST Registry; prospective	854	54 (6)	Unspecified	In univariable analysis only
Bischof 2015 ⁸⁵	1998-2012; Academic centres, North America; retrospective	502	7 (1)	Unspecified	None
Casali 2015 ⁹⁶	2004-2008; EORTC/Australasian Trial; high/intermediate risk GIST	908	97 (11)	Unspecified, R1 resection included	None
Kim 2015 ⁸⁶	2000-2007; Korean multi-centre; gastric GIST; retrospective	1057	17 (2)	Intraoperative, otherwise unspecified	In univariable analysis only
Paper I	2000-2012; Small intestinal GIST; retrospective	71	19 (27)	See Table 9	In multivariable analysis
Paper II	2000-2015; Gastric GIST; retrospective	242	22 (9)	See Table 9	In multivariable analysis
Nishida 2018 ¹³¹	2003-2007; Kinki GIST Study Group, Japan; retrospective	665	21 (3)	Fracture and bleeding, otherwise individually defined	In multivariable analysis
Paper III	2000-2017; retrospective	410	52 (13)	See Table 9	In multivariable analysis

Paper VI	2000-2018; Gastric GIST; retrospective	295	24 (8)	See Table 9	In multivariable analysis
----------	--	-----	--------	-------------	---------------------------

GIST, gastrointestinal stromal tumour; EORTC, European Organisation for Research and Treatment of Cancer. [Adapted from Nishida *et al.* ¹³¹]

GENOTYPE

As described in a previous section, the initial event in the oncogenesis of GIST is the gain-of-function mutation of the *KIT* or *PDGFRA* oncogene. Thus, *KIT* exon 11, *KIT* exon 9, or *PDGFRA* exon 18 mutations are detected in 85 to 90 per cent of GISTs (*Table 1*). Tumour genotype is of importance both as a prognostic and predictive factor. Its *predictive* significance – briefly, non-sensitivity to imatinib of *PDGFRA* D842V mutated and *KIT*/*PDGFRA* wild-type tumours and hyposensitivity of *KIT* exon 9 mutated tumours – are summarised in *Table 5*. The *prognostic* significance has been investigated in several important studies ^{52,54,103,135,142}. Among these, the results made by Corless and colleagues, in their exploratory analysis of the ACOSOG Z9001 trial, diverge as they were unable to detect any independent impact of tumour genotype on RFS in patients receiving placebo ¹⁰³. From other studies, the following conclusions can be drawn: (i) patients with *PDGFRA* mutated tumours have a favourable prognosis ^{52,135}, as have (ii) patients with tumours wild-type for *KIT*/*PDGFRA* ¹³⁵ and (iii) tumours with *KIT* exon 11 duplications ¹³⁵. By contrast, (iv) *KIT* exon 11 del557/558 indicate an unfavourable prognosis ^{52,135}, as do (v) *KIT* exon 9 mutation ^{52,142}. However, for some mutations there is a close link to anatomical location, and this relationship between genotype and anatomy will be addressed in the general discussion. Let it just be pointed out that tumours with identical genotype may have strikingly different phenotypical characteristics, illustrated by the common occurrence of the *KIT* del557/558 genotype in clinically irrelevant micro-GISTs ^{48,83}.

R1 RESECTION

Complete tumour removal with negative microscopic margins is considered to be the optimal result and aim of surgery for localised malignant disease - anything less is either due to necessity or a surgical failure. This equation of margin status with surgical quality makes the ‘R1 issue’ almost an identity question among surgeons ¹⁴³. In breast cancer, ‘ink on tumour’ has become a byword for inadequate surgery ¹⁴⁴, and in soft tissue sarcomas of the extremities

as well as in carcinomas of the gastrointestinal tract, there is ample evidence that both RFS and OS are reduced following R1 resection ^{21,22,25}. This attitude has been transferred to the surgery of GIST, and in guidelines it is stated that an R0 resection is the goal of surgical treatment ^{98,145}. Positive resection margins are reported in 5 to 20 per cent of GIST specimens in large series ^{93,131,143,146}, and there are indeed studies indicating worse outcomes after R1 surgery ^{147,148}. However, neither McCarter and colleagues, in their exploratory analysis of the adjuvant ACOSOG Z9001 trial, nor Gronchi and colleagues, in a post hoc analysis of the EORTC/Intergroup trial, were able to detect differences in RFS related to margin status ^{143,146}.

In surgery for tumours embedded in healthy tissue, the resection margin simply equals the dissection surface ^{18,20}. In segmental resection for abdominal carcinomas of the alimentary tract, the resection margin equals the cut surface of the organ - and in extraperitoneal locations like the rectum or oesophagus, also the circumferential margin ²⁰. The same definition applies to surgical specimens for GIST, but the size of GISTs, their extended relationship to the serosal surface and their tendency to rupture have given rise to confusion about this definition, and a peritoneal breach, spontaneous or iatrogenic, is sometimes recorded as an R1 or even an R2 resection ^{76,147,149}. This is probably the most important confounder in the assessment of margin status in GIST, and the relationship between R1 resection and tumour rupture is the subject of Paper III, presented and discussed in Part Two.

PERCUTANEOUS BIOPSY

The reasons to perform a diagnostic biopsy in suspected malignancy are many: some tumours should be treated with oncological, not surgical measures; in some tumours, neoadjuvant treatment is recommended; sometimes the extent of surgery depends on an exact diagnosis; mutilating or risky procedures are hard to justify without a verified diagnosis; or if malignancy is not confirmed, tumours might be left in place. Nevertheless, practising biopsy also seems to be a matter of tradition: in breast cancer, a biopsy is mandatory ¹⁵⁰, as in soft tissue sarcomas of the extremities ⁹⁹. In retroperitoneal sarcoma, biopsy is recommended ⁹⁹; tumours of the respiratory or alimentary tract often lend themselves to endoscopic biopsy; but pancreaticoduodenectomy, hepatectomy, nephrectomy and major gynaecological resections are regularly performed without a biopsy. There is a risk of needle tract seeding and implantation - and in transperitoneal procedures - of peritoneal tumour dissemination. However, in most cancers, this risk seems to be very low ¹⁵¹. In a pooled analysis of 547

patients with primary retroperitoneal sarcoma subjected to percutaneous core needle biopsy, there were only two instances of presumed needle tract recurrence ¹⁵². GIST patients enrolled in the SSGXVIII/AIO adjuvant trial were post hoc analysed for a possible increased risk of recurrence after percutaneous needle biopsy, and there was no difference in RFS or OS between 47 patients who underwent biopsy and 346 patients who did not ¹⁵³. Van Houdt and colleagues reported one needle tract recurrence in 70 GIST patients who had had percutaneous biopsy ¹⁵⁴.

PART TWO: THE STUDY

The present study is an assemblage of five papers published between 2016 and 2019 and one paper in press. They are all retrospective cohort studies of patients identified in the sarcoma database of Oslo University Hospital, and they all relate to prognostic risk factors in resected primary GIST without metastases. To identify prognostic factors and assess their significance have always been a major discipline in oncological research and they are essential in clinical practice: in deciding the extent of surgical treatment, in selecting patients for neoadjuvant or adjuvant treatment, in follow-up and counselling. In GIST, with an unusual range in metastatic potential, these considerations are particularly pertinent. Adjuvant treatment with imatinib for 3 years is currently recommended for patients at presumed increased risk of recurrence, and even if generally well tolerated, such extended treatment should be well founded when prescribed to persons without evidence of disease. The present studies all investigated established and potential risk factors with the aim of improving prognostication and making it more specific.

AIMS

Paper I: To assess whether tumour rupture, when precisely defined as a major defect of tumour integrity as opposed to specific minor defects, is an independent risk factor for overall and peritoneal recurrence in GIST of the small intestine.

Paper II: (i) To investigate the prognostic value of the definitions applied in Paper I in a larger cohort of patients with gastric GIST. (ii) If confirmed, to assess the contribution of tumour rupture, compared to other established risk factors, to the risk of recurrence in gastric GIST.

Paper III: Having proposed a strict definition of tumour rupture, to assess the prognostic significance of positive resection margins (R1 resection) in GIST patients with and without tumour rupture.

Paper IV: To assess the prognostic significance of peritoneal tumour penetration in GIST, corresponding to stage T4a in gastrointestinal carcinoma, in patient without tumour rupture defined according to the criteria proposed in Paper I.

Paper V: To investigate the relationship between tumour genotype and rupture in GIST.

Paper VI: (i) To provide a detailed description of anatomical (growth pattern and subsite), molecular, and clinical features of gastric GIST in a population-based cohort. (ii) To investigate their associations and possible impact on prognosis.

PATIENTS AND METHODS

Patients were identified in the sarcoma database of Oslo University Hospital. Oslo University Hospital is the sarcoma centre for the South-East Region of Norway, counting 2·9 million inhabitants. According to national guidelines and longstanding tradition, patients have been referred for diagnosis, treatment and follow-up. Patients undergoing emergency surgery or surgery for unsuspected sarcoma at local hospitals were generally referred at the time of diagnosis. In these cases, the diagnosis was confirmed by a sarcoma pathologist at Oslo University Hospital ⁶⁶. Mutation analysis was performed routinely on all intermediate and high-risk tumours from 2003 and selectively on tumours at low or very low risk. Genomic DNA was extracted from fresh frozen or paraffin-embedded tumour tissue and exons 9, 11, 13 and 17 of *KIT* and exons 12, 14 and 18 of *PDGFRA* were analysed by Sanger sequencing. Additional mutation analyses were performed in Paper VI (see below).

On two occasions the authors of the present studies contacted the Cancer Registry of Norway, making inquiries about the number of patients undergoing surgery for localised GIST within the South-East Region reported to the Cancer Registry. The corresponding figures for the same periods of patients recorded in the sarcoma database of Oslo University Hospital represented 93 per cent (Paper III) and 98 per cent (Paper VI) of the regional cohort. However, only crude numbers were compared, no corroboration was made at an individual level.

From the sarcoma database, demographical data and basic clinical information were extracted. These data were confirmed and retrospectively supplemented from medical records, and for one study, Paper VI, additional analyses were performed. For the complete study, Papers I to VI, the total number of 433 patients were included. Their demographical and treatment characteristics are presented in *Table 7*.

SURGICAL TREATMENT

Two-thirds of the patients underwent surgery at the sarcoma centre (*Table 7*). During the study period, from January 2000 to August 2018, four consultant surgeons were engaged in the treatment of GIST patients at Oslo University Hospital, one of them for the complete

interval. They had a uniform surgical strategy and technique and a uniform way of reporting. Great care was taken to avoid tumour rupture. Organ-preserving resections were preferred whenever possible. Multivisceral resections were performed when malignant infiltration was evident or suspected or when necessary for function. A complete resection of the tumour with microscopically negative margins was the aim of surgery, but no standard as to the minimum distance was set. Patients undergoing incomplete (R2) resections were excluded from the investigations, as were patients with synchronous metastases, even if completely removed.

Table 7. Demographical and clinical characteristics of patients included in Papers I to VI

Characteristic	Number	Per cent
Total	433	100
Sex ratio (males : females)	218 : 215	50 : 50
Age at surgery, median (range), years	65 (14-93)	
Follow-up period, median (range), months	46 (0-185)	
Surgery at sarcoma centre		
Yes	289	67
No	144	33
Elective surgery	387	89
Emergency surgery	31	7
Incidental finding at surgery	15	3
Surgical mortality	4	1
R1 resection	47	11
Multivisceral resection	65	15
Tumour site		
Gastric	298	69
Non-gastric	135	31
Tumour rupture	54	13
Spontaneous	24	
Iatrogenic	25	
Spontaneous and iatrogenic	5	
Risk category (mNIH)		
High	163	38
Intermediate	74	17
Low or very low	191	45
Unspecified	6	
Adjuvant treatment	82	19
Duration, median (range), months	25 (1-60)	
Neoadjuvant treatment	20	5
Duration, median (range), months	8 (1-16)	

Lymphadenectomy was not routinely performed. During the study period, laparoscopic resection was introduced for gastric GISTs of small to intermediate size. The use of cutting

stapling devices greatly increased. For patients undergoing surgery at local hospitals or in other departments of Oslo University Hospital, surgical reports were available in all cases.

ADJUVANT AND NEOADJUVANT TREATMENT

From May 2004 to August 2008, patients with high-risk GIST according to the mNIH criteria⁷³ were considered for inclusion in the SSGXVIII/AIO trial (1 *versus* 3 years of adjuvant imatinib, 400 mg daily)⁹³. From September 2008 to May 2011, high-risk patients received adjuvant imatinib for 1 year; from June 2011 for 3 years. Since then, patients with *KIT* exon 9 mutated tumours received 800 mg daily if tolerated. Since May 2015, patients with gastric GIST with more than 10 mitoses per 5 mm², non-gastric GIST with more than 5 mitoses per 5 mm², or patients with tumour rupture were considered for the SSGXXII trial (Clin.gov: NCT02413736) (3 *versus* 5 years of adjuvant imatinib).

Neoadjuvant treatment with imatinib was given at the discretion of the multidisciplinary team, usually for 6 to 12 months (*Table 7*), to patients with marginally resectable tumours, tumours at high risk for rupture, or in cases where anticipated tumour shrinkage would reduce the extent of surgery or secure margins. The use of neoadjuvant treatment expanded during the study interval. Mitoses were not counted in specimens from patients subjected to neoadjuvant treatment.

FOLLOW-UP AND RECURRENCE

Standard follow-up included clinical examination and abdominal and pelvic CT and chest X-ray or CT every 3 to 6 months for 5 to 10 years. Patients without recurrence, including patients who died from an unrelated cause, were censored at the date of the latest CT. Patients who died from surgical complications were censored at the date of surgery. Recurrence was recorded if verified on biopsy or indisputable on CT. RFS was measured from the date of surgery. There were 82 patients who experienced a recurrence at a median time of 24 months since surgery (range 3 to 164). Estimated 5 and 7½-year RFS was 78 and 73 per cent.

ETHICAL APPROVAL

All studies were approved by the Data Protection Officer at Oslo University Hospital (Paper I, No. 2015/15603; Paper II, No. 2016/16853; Paper III, No. 2018/05487; Paper IV, Nos. 2016/16853 and 2018/05487; Paper V, Nos. 2015/15603, 2016/16853 and 2018/05487; Paper VI, Nos. 2016/16853 and 2018/05487). In Paper VI, mutation analysis using next-generation sequencing (NGS) required approval by the Regional Committee for Medical and Health Research Ethics of South-East Norway, which was obtained (No. 2010/1244). Written informed consent was given by the patients for information to be registered and used in the sarcoma database.

STATISTICAL ANALYSIS

In all studies where survival analysis was performed, RFS was estimated by the Kaplan-Meier method and compared using the log-rank test. Multivariable survival analysis was performed using the Cox proportional hazards regression model. Associations between variables were investigated with two-tailed Fisher's exact test or Pearson's χ^2 test for categorical variables and independent Mann-Whitney *U* test or Kruskal-Wallis test for continuous variables. Relative risks (RRs) and HRs were reported with 95 per cent c.i. Interobserver agreement was evaluated with Cohen's κ test (Paper VI). $P < 0.05$ was considered statistically significant. IBM SPSS Statistics for Windows (Armonk, NY, US) was used.

ISSUES PERTAINING TO THE INDIVIDUAL STUDIES

Paper I: Patients undergoing complete resection for localised GIST of the small intestine from January 2000 to December 2012 were included. Definitions of major and minor defects of tumour integrity were formulated *a priori* and retrospectively assessed from surgical and histopathological reports. *Minor defects of tumour integrity* were (i) core needle biopsy, (ii) peritoneal tumour penetration (corresponding to T4a in the TNM classification of gastrointestinal carcinoma ⁶⁹), (iii) iatrogenic peritoneal laceration, or (iv) microscopically involved intestinal resection margins (R1) (*Figure 3A*). Histological slides from all tumours except those that were removed intralesionally were re-examined for possible penetration of the peritoneum. *Major defects of tumour integrity* were (i) tumour fracture or spillage, (ii) piecemeal resection or intralesional dissection, (iii) bowel perforation at the tumour site, (iv)

blood-tinged ascites at laparotomy, (v) surgical (incisional) biopsy, or (vi) microscopic adjacent organ infiltration (corresponding to T4b in the TNM classification of gastrointestinal carcinoma ⁶⁹) (Figure 3B).

Figure 3A. Minor defects of tumour integrity. From left to right: superficial peritoneal laceration; peritoneal tumour penetration; core needle biopsy; microscopically involved resection margin.

Figure 3B. Criteria for tumour rupture proposed in Paper I and applied in Papers I-VI. From left to right: tumour fracture; tumour spillage; blood-tinged ascites; gastrointestinal perforation at the tumour site; microscopic adjacent organ infiltration; piecemeal resection; incisional biopsy. [Reproduced from Nishida *et al.* ¹⁴¹]

Paper II: Patients undergoing complete resection for localised gastric GIST from January 2000 to December 2015 were included. The definition of ‘tumour rupture’ corresponded to the definition of major defects of tumour integrity in Paper I; the definition of minor defects of tumour integrity was retained with the specification that mucosal defects or possible tumour spillage contained within the gastric lumen were *per se* neither classified as tumour rupture nor minor defects of tumour integrity. Criteria for the assessment of resection margins (R status) applied equally to specimens removed surgically and endoscopically.

Paper III: Patients undergoing complete resection for localised GIST from January 2000 to September 2017 were included. The definition of ‘tumour rupture’ corresponded to the definition of major defects of tumour integrity in Paper I. Residual tumour status was defined according to the R classification ²⁰ with the following specifications: complete resections with microscopically positive or ‘uncertain’ margins were coded R1; margins less than 1 mm but explicitly negative were coded R0. For GIST in the stomach, small intestine and colon, the organ transection surface equalled the resection margin; tumour relation to the peritoneal (serosal) surface was without implication for R status. For extraperitoneal GISTs (oesophagus, duodenum, rectum), both the transection and dissection surfaces were considered, and for extra-gastrointestinal GISTs, the dissection surface equalled the resection margin. Tumour rupture did not influence R status.

Paper IV: Patients undergoing complete resection for localised GIST from January 2000 to September 2017 were included except patients with extraperitoneal (oesophagus, rectum) GIST or tumour rupture according to the definition cited above. Only peritoneal defects or tumour extension to the peritoneal surface were considered peritoneal penetration; incomplete peritoneal penetration was not. Peritoneal penetration with microscopic adjacent organ infiltration was a criterion of tumour rupture, and these patients were excluded. During the study interval, the Department of Pathology, Oslo University Hospital, introduced a reporting template that included a required field for recording serosal involvement. Before the introduction of this template, the tumour-peritoneum relationship was described in an individual fashion. Slides were re-examined if the original report was inconclusive.

Paper V: Patients were included who had undergone complete resection of localised GIST of the stomach or small intestine from January 2000 to April 2017 and in whom mutation analysis was performed as part of clinical routine (see above). Mutations were categorised as deletions, insertions-deletions, substitutions or duplications/insertions according the amino acid changes predicted from the nucleotide sequence. An insertion-deletion refers to a deletion of amino acids and a concomitant insertion of at least one amino acid, whereas a deletion is an exclusive deletion of at least one amino acid. Tumour rupture was defined according to the criteria cited above.

Paper VI: Patients undergoing complete resection for localised gastric GIST from January 2000 to August 2018 were included. Gastric subsite was divided into the upper, middle and lower thirds. Macroscopic growth pattern was classified as *luminal*, not affecting the peritoneal contour of the stomach; *exophytic*, not affecting the mucosal contour; or *transmural*, affecting both the peritoneal and mucosal contours (*Figure 4*). The assessment was done on CT imaging independently by a radiologist and a surgeon (interobserver agreement 90 per cent, $\kappa=0.85$) and resolved by consensus when not in agreement. If the surgical report indicated a growth pattern different from the images, the surgical assessment was chosen. Patients with a history of mucosal ulceration (bleeding) or ulceration detected on microscopy/endoscopy were by definition classified with a luminal component; similarly, microscopic serosal penetration was equivalent to an exophytic component.

Figure 4. **A** The partition of the stomach into an upper, middle and lower third. **B** Macroscopic growth pattern: luminal, exophytic and transmural tumours.

Tumours not analysed routinely were analysed using the AmpliSeq for Illumina Cancer Hotpot Panel version 2, examining hotspot regions of 50 cancer genes, including *KIT* exons 2, 9, 10, 11, 13, 15, 17 and 18, as well as *PDGFRA* exons 12, 14, 15 and 18. NGS libraries were generated from 100 ng of genomic DNA following Illumina's protocol. Libraries were normalised using the AmpliSeq Library Equalizer kit from Illumina and sequenced paired end 2 x 150 base pairs on an Illumina MiSeq instrument. Sequence reads were mapped to the Human UCSC hg19 reference genome and variants called using the DNA Amplicon workflow version 2.1.0.19 using the Illumina's Local Run Manager software. Mutation calls were manually inspected on the Integrative Genomics Viewer.

SUMMARY OF RESULTS AND IMPLICATIONS

PAPER I: DEFINITION AND CLINICAL SIGNIFICANCE OF TUMOUR RUPTURE IN GASTRO-INTESTINAL STROMAL TUMOURS OF THE SMALL INTESTINE

The study included 71 patients with localised GIST of the small intestine completely resected (R0/R1). The median follow-up period was 58 (7 to 122) months. A major defect of tumour integrity was recorded in 20 patients and a minor defect in 21. Estimated 5-year recurrence rates were 64 per cent, 29 per cent and 31 per cent in patients with major, minor and no defect, respectively ($P = 0.001$). HR for major defect compared to no defect was 3.55 (95 per cent c.i. 1.51 to 8.35) (Figure 5A). Peritoneal recurrence rates for major, minor and no defect were 52 per cent, 25 per cent and 19 per cent ($P = 0.002$), and the HR for major defect compared to no defect 4.98 (1.69 to 14.68) (Figure 5B). On multivariable analysis, MI, major defect of tumour integrity, tumour size, and age were independently associated with the risk of recurrence.

Implications: Defined according to these criteria, a major defect of tumour integrity was independently associated with recurrence of GISTs of the small intestine. Reproduced in gastric GIST, these criteria would serve as a general definition of tumour rupture in GIST.

Figure 5 (Paper I). Estimated recurrence (Kaplan-Meier) after surgery for non-metastatic gastrointestinal stromal tumours of the small intestine. **A** overall recurrence, **B** peritoneal recurrence.

PAPER II: RECURRENCE-FREE SURVIVAL AFTER RESECTION OF GASTRIC GASTROINTESTINAL STROMAL TUMORS CLASSIFIED ACCORDING TO A STRICT DEFINITION OF TUMOR RUPTURE: A POPULATION-BASED STUDY

Among 242 patients with localised gastric GIST included in the study, tumour rupture occurred in 22 (9 per cent) and minor defects of tumour integrity in 81 (33 per cent). Estimated 5-year RFS for patients with tumour rupture, minor defects of tumour integrity, and no defect was 37, 91, and 96 per cent, respectively ($P < 0.001$) (*Figure 6A*). In the high-risk group (mNIH), 5-year RFS for patients with rupture was 37 per cent *versus* 77 per cent without rupture, HR 3.56 (95 per cent c.i. 1.57 to 8.08; $P = 0.001$) (*Figure 6D*). On multivariable analysis, tumour rupture and MI were independently associated with recurrence. Of 13 patients who received adjuvant imatinib after tumour rupture, 11 relapsed.

Implications: The definition of tumour rupture identified patients at particularly high risk of recurrence also in gastric GIST and would serve as a general definition of rupture in GIST. In patients with tumour rupture, the disease tended to recur regardless of adjuvant therapy, and extended treatment – possibly for life – should be considered. Patients with gastric GIST without rupture had a good prognosis, even in the high-risk category, and in this group, further investigations should try to identify patients for whom adjuvant treatment could be omitted.

Figure 6 (Paper II). Estimated recurrence-free survival (Kaplan-Meier) for patients with localised gastric GIST. **a** by defects of tumour integrity, **b** by tumour rupture in tumours > 10 cm, **c** by tumour rupture in tumours with mitotic index > 5 per 50 HPF, **d** by tumour rupture in high-risk patients (mNIH).

PAPER III: RELATIONSHIP BETWEEN R1 RESECTION, TUMOUR RUPTURE AND RECURRENCE IN GASTROINTESTINAL STROMAL TUMOUR

Among 410 patients with localised GISTs, there were 47 (12 per cent) with R1 resection and 52 (13 per cent) with tumour rupture. The RR of R1 resection with rupture was 3.55 (95 per cent c.i. 2.09 to 6.03; $P < 0.001$). In patients without rupture, there was no difference in estimated 5-year RFS after R0 *versus* R1 resection (88 *versus* 93 per cent; HR 0.71, 95 per cent c.i. 0.17 to 2.98; $P = 0.638$) (Figure 7C); nor was there any difference in patients with rupture (37 *versus* 31 per cent; HR 1.31, 95 per cent c.i. 0.68 to 2.54; $P = 0.420$) (Figure 7D). In multivariable analysis, tumour rupture but not R1 resection was independently associated with recurrence. Twenty-four patients at very low, low or intermediate risk did not receive adjuvant imatinib after R1 resection and remained recurrence-free.

Implications: R1 resection did not confer an inferior prognosis. In the absence of high-risk features, these patients will not benefit from additional treatment – adjuvant therapy or re-excision.

Figure 7 (Paper III). Estimated RFS (Kaplan-Meier) in patients with localised GISTs by **A** R status ($P = 0.001$), **B** tumour integrity ($P < 0.001$), **C** R status in patients without rupture ($P = 0.638$), **D** R status in patients with rupture ($P = 0.420$).

PAPER IV: IS PERITONEAL TUMOR PENETRATION OF PROGNOSTIC IMPORTANCE IN GASTRO-INTESTINAL STROMAL TUMOR?

The study included 341 patients with abdominally located, localised GISTs without rupture. The median follow-up period was 51 (range 0 to 175) months. Peritoneal penetration (PP) was recorded in 82 (24 per cent) patients. There were 32 recurrences, 9 in patients with PP, 23 in patients without PP. Despite statistically significant associations between PP and established

risk factors (size, mitotic index, non-gastric location), there was no difference in 5-year RFS in patients with and without PP, 86 and 90 per cent, respectively (HR 1.25, 95 per cent c.i. 0.58 to 2.70; $P = 0.577$) (Figure 8). Of 97 patients in the high-risk category (mNIH), 53 received adjuvant imatinib. There was no statistically significant difference in recurrence rates between PP-positive and PP-negative patients in either group.

Implications: The fact that PP did not have a negative impact on prognosis separates GISTs from gastrointestinal carcinomas and indicates biological differences between tumours of mesenchymal and epithelial origin that ought to be investigated in experimental studies. PP should not be confounded with tumour rupture and should not entail adjuvant treatment.

Figure 8 (Paper IV). Estimated RFS (Kaplan-Meier) in patients with abdominally located, localised GIST without rupture. **A** the complete cohort ($P = 0.577$), **B** gastric tumours ($P = 0.389$), **C** non-gastric tumours ($P = 0.425$).

PAPER V: GENOTYPE AND RISK OF TUMOUR RUPTURE IN GASTROINTESTINAL STROMAL TUMOUR

Two-hundred and nine patients with localised GIST and available mutation data were identified. Tumour rupture occurred in 37 (18 per cent) patients. Among the 155 patients with *KIT* exon 11 mutations, an increased risk of rupture was observed with a deletion or insertion/deletion (29 per cent) compared to substitutions (10 per cent) or duplications/insertions (11 per cent; $P = 0.014$) (Table 8). Notably, 37 per cent with *KIT* exon 11 del557/558 had rupture *versus* 14 per cent with other exon 11 mutations ($P = 0.002$). This association was confined to gastric tumours, of whom 35 per cent with del557/558 had rupture *versus* 8 per cent with other exon 11 mutations ($P = 0.001$). In multivariable logistic regression analysis, del557/558 and tumour size were associated with an increased likelihood of tumour rupture while MI was not.

Implications: In gastric GIST, tumours with a del557/558 genotype were at increased risk of rupture. This high-risk feature can be identified in the diagnostic evaluation and should be taken into account when neoadjuvant treatment is considered.

Table 8 (Paper V). Genotypes and selected clinical and histopathological characteristics

	All patients	Stomach	Small intestine	Size (cm)*	MI (/50 HPF)*	Rupture
All patients	209	148 (71)	61 (29)	6.2 (0.5-30.0)	3 (0-178)	37 (18)
<i>PDGFRA</i> exon 18	27	27 (100)	0 (0)	6.0 (2.5-21.0)	1 (0-19)	2 (7)
D842V	21	21 (100)	0 (0)	6.0 (3.0-21.0)	1 (0-4)	1 (5)
Other	6	6 (100)	0 (0)	n.d.	n.d.	1 (17)
<i>PDGFRA</i> exon 12	2	2 (100)	0 (0)	n.d.	n.d.	0 (0)
<i>KIT</i> exon 9	7	0 (0)	7 (100)	n.d.	n.d.	0 (0)
<i>KIT</i> exon 11	155	111 (72)	44 (28)	6.5 (0.5-30.0)	5 (0-178)	32 (21)
del557/558	46	34 (74)	12 (26)	9.9 (2.8-30.0)	12 (0-178)	17 (37)
del not 557/558	40	22 (55)	18 (45)	4.2 (0.5-20.0)	4 (0-106)	8 (20)
Substitutions	50	37 (74)	13 (26)	6.0 (1.3-15.0)	3 (0-24)	5 (10)
Dupl/insert	19	18 (95)	1 (5)	6.0 (2.0-18.0)	4 (0-16)	2 (11)
<i>KIT</i> exon 13	2	0 (0)	2 (100)	n.d.	n.d.	1 (50)
<i>KIT</i> exon 17	3	3 (100)	0 (0)	n.d.	n.d.	0 (0)
No mutation detected	13	5 (38)	8 (62)	3.8 (1.7-19.0)	1 (0-8)	2 (15)

Values in parenthesis are percentages unless indicated otherwise. *Values are median (range). MI, mitotic index; HPF, high-power field; n.d., not determined in groups with fewer than 10 patients; del, insertions-deletions; dupl/insert, duplications/insertions.

The cohort comprised 295 patients with localised gastric GIST, representing > 95 per cent of the patients reported to the Cancer Registry of Norway from the South-East Region. Of 292 tumours suitable for classification, 122 were located in the upper (42 per cent), 120 in the middle (41 per cent), and 50 in the lower third (17 per cent). The number of luminal, exophytic and transmural tumours was even. *KIT* mutations were detected in 183 of 256 tumours (72 per cent), *PDGFRA* mutations in 62 (24 per cent) and *KIT/PDGFRA* wild-type tumours in 11 patients (4 per cent). *KIT* mutated tumours had a predilection for the upper two-thirds; *PDGFRA* mutated tumours for the lower two-thirds (*Figure 9*). In the upper third, 92 per cent of the tumours harboured mutations conferring sensitivity to imatinib. Transmural tumours were larger, more often located in the upper third, had higher mitotic activity, and were more frequently *KIT* del557/558 mutated. Estimated 5-year RFS was 71 per cent for patients with transmural tumours *versus* 96 per cent with luminal/exophytic tumours, HR 8.45 (95 per cent c.i. 3.69 to 19.36; $P < 0.001$) (*Figure 10B*); in high-risk patients, estimated 5-year RFS was 46 per cent for patients with transmural tumours *versus* 83 per cent for patients with luminal/exophytic tumours, HR 4.47 (95 per cent c.i. 1.71 to 11.66; $P = 0.001$). In the 66 patients with exophytic or luminal tumours > 5 cm without rupture, 5-year RFS was 98 per cent. Fifty-five of these 66 patients (83.3 per cent) did not receive adjuvant treatment.

Implications: In the stomach, tumour genotypes had a characteristic anatomical distribution. In the upper third, over 90 per cent of tumours were sensitive to imatinib. Without facilities for molecular testing, neoadjuvant treatment could be started for tumours in this location, and even adjuvant treatment could be contemplated. A transmural growth pattern was a predictor of poor outcome, related to an elevated MI. In lack of information on mitotic activity after neoadjuvant treatment, this variable could act as a surrogate when considering continued postoperative treatment. Exophytic and luminal tumours, irrespective of size, were associated with an excellent prognosis without rupture, and these patients may not benefit from adjuvant treatment.

Figure 9 (Paper VI). Relationships between tumour subsite, growth pattern, tumour size, mitotic index and tumour genotype in gastric GIST. GIST, gastrointestinal stromal tumour; MI, mitotic index; HPF, high-power fields of the microscope; PDGFRA, platelet-derived growth factor receptor- α .

Figure 10 (Paper VI). Estimated RFS (Kaplan-Meier) in patients with localised gastric GIST by **A** gastric subsite ($P = 0.020$); **B** macroscopic growth pattern ($P < 0.001$); **C** tumour genotype ($P < 0.001$). GIST, gastrointestinal stromal tumour; PDGFRA, platelet-derived growth factor receptor- α .

METHODOLOGICAL DISCUSSION

These investigations, despite their unity, did not have a preconceived structure. The question stated in Paper I originated from clinical curiosity and an obvious need for a definition of tumour rupture in GIST; it was pursued in Papers II to V, and in Paper VI, other but related problems were introduced. The thesis centres on tumour rupture and on prognostic factors in gastric GIST. The definition examined in Paper I was proposed *a priori* by the investigators. So far, it constitutes the only strict and systematically applied definition, and it has already gained a certain acceptance ^{141,143}, but a slightly different definition, suggested in the first place, may have yielded similar results in the same patients. Most importantly, the criteria remain to be validated in other cohorts.

In the present work, associations were explored – associations between risk factors and associations between risk factors and outcomes. It should be emphasised at this point, elementary though it is, that *associations* are not *causal* relationships, and that conclusions on pathogenesis cannot be drawn from associations, e.g. the fact that tumour rupture is associated with recurrence does not signify that rupture leads to recurrence; or that del557/558 mutations are associated with a transmural growth pattern does not mean that tumours are transmural because they harbour a del557/558 mutation.

RETROSPECTIVE COHORT STUDIES

All papers were retrospective cohort studies with inherent limitations. These limitations are – if not compensated – to a certain degree counterbalanced by the fact that they are population-based. Patients with soft tissue sarcomas in the South-East Region of Norway are referred to the Sarcoma Group at Oslo University Hospital for diagnostic investigations, treatment and follow-up; and patients undergoing surgery for undiagnosed sarcoma are usually referred at the time of diagnosis. Since the early 1990s, information from clinicians and pathologists at Oslo University Hospital has been prospectively recorded in a sarcoma database, including second opinions on pathology from local laboratories. For the present studies, all patients were identified in this database. By legal regulation, cancer patients are reported to the Cancer Registry of Norway from clinical and pathology departments. As stated in the Methods section, the Cancer Registry was contacted by the investigators on two occasions (Papers III

and VI). The number of patients recorded in the sarcoma database corresponded to 93 and 98 per cent, respectively, of the number reported to the Cancer Registry during the same time periods. However, no corroboration at an individual level was done, and in the Cancer Registry, some patients with GIST were probably not detected because they were erroneously recorded with benign disease. Moreover, in the Cancer Registry, patients are identified by domicile, and the sarcoma database contains a small number of patients living outside the South-East Region. Therefore, a certain selection bias must be assumed; e.g. missing patients with small GISTs incidentally discovered during surgery or electively removed in local hospitals and not referred to the Sarcoma Group.

Only basic information was retrieved from the sarcoma database, and most variables were retrospectively assessed from medical records, i.e. surgical and pathology reports, diagnostic imaging, case notes etc. As described in the Methods section, only four consultant surgeons with a uniform manner of reporting treated the patients at the sarcoma centre during the study period. The presence of ascites or disseminated disease, tumour location and growth pattern, adjacent infiltration, macroscopic peritoneal penetration and resection margins were generally reported, but in the absence of such information, positive findings cannot be excluded. Moreover, one third of the operations were performed in local hospitals or other departments of Oslo University Hospital with reports sometimes of inferior quality. With a prospective design, information would have been more reliable. In this respect, the present study shares the deficiencies of the vast majority of studies on tumour rupture in GIST (*Table 6*).

All pathology examinations were performed in clinical routine by a limited number of pathologists specialised in sarcoma diagnostics. The counting of mitoses was done according to established criteria ⁶⁶, and the assessment of resection margins was straightforward, even if by nature uncertain in individual cases. Explicitly uncertain margins were coded as R1 whereas explicitly negative margins, whatever the width, were coded as R0 (Paper III). ‘Tumour rupture’ was a clinical category, but the tumour surface was assessed during the pathology examination. A template was introduced late in the study interval with a required field for reporting the tumour-peritoneal relationship; otherwise, the serosal surface was described in an individual fashion, and for Papers I and II, slides were re-examined. However, artificial, iatrogenic and true tumoural penetrations of the peritoneum were sometimes undistinguishable under the microscope. In Paper I, an attempted distinction was made between iatrogenic and spontaneous peritoneal penetration, but as this distinction seemed arbitrary, these categories were merged in the subsequent studies. They were both classified

as minor defects of tumour integrity; thus, no consequence for the analysis ensued. Again, a prospective protocol with specifications for the handling, processing and examination of the specimens would have been preferable.

Retrospective assessment of diagnostic images was not hampered in the same manner as were evaluations of surgery and histopathology. The classification of growth pattern relied on CT imaging (Paper VI). (For the few patients who lacked preoperative images, the evaluation relied on the surgical report.) After adjusting their evaluations in pilot cases, scans were examined independently by a radiologist and a surgeon. Interobserver agreement was satisfactory (90 per cent, $\kappa = 0.85$). Still, these assessments are inherently subjective. For small gastric tumours amenable to endoscopic resection, a classification of growth pattern has been proposed based on the relationship of the tumour to the muscular layer, as visualised on endoscopic ultrasonography⁸⁸. A similar classification based on histopathology is conceivable. However, in a cohort with 25 per cent of the tumours exceeding 7.5 cm, these assessments would also be highly observer dependent.

TIME DEPENDENT VARIATION

The study period spans 18 years. In 2000, GIST was barely recognised as a nosological entity, and experience and knowledge acquired since then have brought about changes in diagnostics, surgery and medical treatment. For instance, among the patients included in the present study, those undergoing total gastrectomy were all treated before 2004, and transperineal extirpation of rectal GIST was not performed after 2010. Adjuvant imatinib was introduced for trial patients in 2004; from 2008 to 2011, this treatment was given for 1 year; since 2011, for 3 years; and since 2015, patients at particularly high risk have been enrolled in the SSGXXII trial (5 *versus* 3 years). Neoadjuvant treatment was given for the first time to a patient in 2005, to a second patient in 2009, and in 2015, to four patients. In the early years, one has to assume that some patients were misdiagnosed and missed. Molecular diagnostics were introduced in 2003. Moreover, the proportion of incidental gastric GISTs increased during the study period from 23 per cent in the first half to 42 per cent in the second half ($P < 0.001$), probably due to an expanded use of diagnostic CT for unrelated symptoms. Correspondingly, median tumour size decreased from 5.4 to 4.2 cm ($P = 0.006$). Exactly how these shifts have influenced the present analyses is difficult to determine, but an impact cannot be neglected.

ENDPOINTS IN SURVIVAL ANALYSES

In clinical cancer studies and cancer trials, overall survival (OS) is the most objectively defined endpoint, generally understood as the time to the patients' death from any cause. For other endpoints during this period - from intervention to death - a number of terms are used: in the metastatic setting, progression-free survival and time to progression; in the non-metastatic setting, cancer-specific survival, disease-free survival, recurrence-free survival, relapse-free survival, recurrence-free interval, cancer-free interval, to mention only a few. In the non-metastatic setting, these terms denote time to cancer recurrence – sometimes the occurrence of any cancer, and some of the definitions include death – from any cause, from unknown cause, from any cancer, from index cancer or from cancer treatment. The definitions of these endpoints vary from protocol to protocol, from study to study, and the same term may signify different events, or different terms may signify the same events. This makes cross-study comparison difficult - or impossible - and obscures the interpretation of results. In GIST, with extraordinarily effective treatment for advanced disease, OS reflects the response to this treatment, its quality, and the general health of the study population rather than characteristics of the primary tumour or the treatment of localised disease. Methodologically it is therefore a forbidding endpoint, illustrated by its substitution in the EORTC/Intergroup trial by the makeshift surrogate 'imatinib failure-free survival' (see Introduction, section on adjuvant treatment) ⁹⁶.

Hompland and colleagues have shown that for patients with metastatic GIST treated by the Oslo Sarcoma Group, median OS is 7 years ¹¹³. In none of the studies in the present thesis was OS an endpoint. In survival analysis, only RFS was analysed, uniformly defined as time from surgery to any GIST recurrence – local, regional or distant. Death was not counted as an event under any circumstance. Patients who died from surgical complications were censored at the date of surgery; one patient who died from complications related to neoadjuvant imatinib treatment never reached surgery and was not included. Defined accordingly, the DATECAN initiative (Definition of the Assessment of Time-to-Event Endpoints in Cancer Trials) recommends that this endpoint be termed 'recurrence-free interval' ¹⁵⁵. However, the more commonly used 'RFS' was chosen and uniformly defined for each study. Patients were censored at the latest CT even when subsequent clinical data regarding unrelated conditions existed. The sarcoma database is linked to the National Population Register, and all deaths were recorded. In patients under follow-up who died between controls, death from an undiagnosed recurrence can be excluded. In patients who died after completing the follow-up

programme, death from GIST recurrence is very unlikely. In patients with a short follow-up – chiefly low-risk or very old patients, death from undiagnosed recurrence cannot be ruled out but is unlikely (see section on follow-up below). In this closely followed cohort of patients with a long-term RFS of approximately 75 per cent, including death from unknown cause as an event would not improve the quality of the data; and including death from any cause – in a cohort where other malignancies are overrepresented, would make their quality inferior.

THE CONFOUNDING ROLE OF ADJUVANT TREATMENT

The characterisation of GIST and the discovery of the effect of imatinib happened within a few years ^{36,39,90}. The first adjuvant trial was launched in 2002 ⁹², and since the very beginning, the confounding role of adjuvant TKI has haunted population-based investigations. From the pooled, population-based series analysed and published by Joensuu and colleagues in 2012, an unknown number of patients treated with adjuvant imatinib were excluded, thereby introducing a bias ³⁷. Most analyses of prognostic factors in TKI-naïve patients have been performed either in early series or in the placebo arms of randomised trials in selected patients ^{37,52,67,74,77,78,98,103,109,130,135,138,143,146,156,157}.

In the present study, 82 patients received adjuvant imatinib for a median time of 25 months (*Table 7*), exactly half of the patients in the high-risk category (mNIH). For patients without recurrence, follow-up was similar, median 50 months (range 6 to 134) for treated patients and 54 months (range 0 to 143) for high-risk patients not treated. Estimated 5-year RFS was 60 per cent for treated patients compared to 46 per cent for patients not treated ($P = 0.05$). The difference dropped at 7 years to 52 per cent *versus* 44 per cent. With a longer follow-up period, more recurrences would be expected in the adjuvant group. Moreover, the adjuvant group possibly contained patients at higher risk. A masking effect of adjuvant imatinib must therefore be suspected. Subgroup analyses were only indicative as their small size precluded valid statistics.

MUTATION ANALYSIS

Mutation analysis was introduced in clinical routine in 2003 and performed by Sanger sequencing on high and intermediate risk tumours (mNIH), discriminately on lower risk tumours. All recurrent tumours were genotyped, either on tissue from the primary tumour or

on biopsy from a metastasis. Sanger sequencing is an established and robust methodology. For the present investigations, all tumours classified as wild-type for *KIT/PDGFRA* were re-analysed in order to exclude a technical issue, and a few cases were reclassified as ‘material unfit for analysis’, a category which finally included 11 of 266 samples (4 per cent). In Paper VI, supplementary analyses were performed by NGS (99 cases, 34 per cent). As a matter of course, more tumours in the very low and low-risk categories and fewer tumours in the intermediate and high-risk categories were analysed by this method. NGS was not performed on tissue from tumours resected outside the sarcoma centre (23 cases, 8 per cent of patients included in Paper VI). There was no difference in the distribution across risk categories between tumour analysed by NGS and tumours not subjected to analysis ($P = 0.209$). Material was unfit for analysis in 10 cases (10 per cent) subjected to NGS. This method has an inherent risk of missing long deleted or duplicated sequences; therefore, sequences were manually inspected. Still, we cannot exclude that certain mutations could have been missed.

KIT exon 11 deletions and insertion-deletions involving codons 557 and/or 558 have been associated with poor outcomes in a number of studies. However, there is no consensus regarding neither definition nor nomenclature. For the present studies (Papers V and VI), the authors chose to include *KIT* exon 11 *deletions and insertion-deletions* involving codons 557 and 558, designated ‘del557/558’.

STATISTICAL ANALYSIS

In none of the investigations were sophisticated statistical methods applied. Associations between categorical and continuous variables were analysed with standard methods, survival was analysed with the Kaplan-Meier method and differences compared with the log-rank test. In multivariable analysis, the basic epidemiological co-variable ‘sex’ was always included; age was included in all but one analysis. No more than six co-variables were entered in any model due to the limited number of patients or events. In multivariable analyses, tumour size and mitoses were treated as categorical co-variables with specified cut-off values. However, analyses were also performed with these as continuous co-variables yielding similar results (unpublished data). All major conclusions were supported by statistically highly significant results. Nevertheless, confidence intervals tended to be wide due to low number of patients or events.

FOLLOW-UP

Patients were followed at the sarcoma centre by a consultant surgeon or oncologist, 20 patients according to the SSGXVIII/AIO protocol ⁹³, others according to internal guidelines (see Methods section). However, doctors had the latitude to individualise, and patients at very low risk (mNIH) had shorter follow-up than patients at low to high risk, median 21 months (range 0 to 129) compared to 48 months (range 0 to 186) ($P = 0.001$). There was no difference in follow-up between the low, intermediate and high-risk groups for patients without recurrence ($P = 0.148$). Undetected recurrences in the very low risk group, which included 37 patients, are unlikely.

Two out of three patients underwent surgery at the sarcoma centre; all medical treatment was given there, and patients were discussed at multidisciplinary meetings with pathologists and radiologists in the sarcoma group. Even if this is not, strictly speaking, a single centre study, it strongly reflects the attitudes of the Oslo Sarcoma Group, which may differ from practice in other sarcoma centres.

GENERAL DISCUSSION

TUMOUR RUPTURE IN GIST

In 2007, tumor rupture was first identified as a risk factor in retrospective Polish and Japanese studies ^{67,130}. In the former, the concept included R1 resection, but the term was not otherwise defined. The following year, Joensuu proposed the mNIH consensus criteria including spontaneous and iatrogenic tumour rupture as a high-risk feature ⁷³. Joensuu and colleagues subsequently confirmed the negative prognostic impact of tumour rupture in a pivotal study merging data from four population-based cohorts ³⁷. However, there was no pre-defined definition of tumour rupture, tumour rupture was not always considered in the original publications, the concept variously included R1 resection and mucosal perforation, and data on rupture were retrospectively recorded. With no established definition of tumour rupture, the reported incidence varies greatly, from 1 to 22 per cent ^{67,85}, and its independent impact on prognosis has been hard to demonstrate in multivariate analysis, even in large series (*Table 6*).

Anatomical considerations. The great majority of GISTs are located in the abdominal portion of the gastrointestinal tract - the stomach or the small intestine. At these sites, tumour rupture causes contamination of the entire abdominal cavity, and the peritoneum – in surgical terminology, ‘the serosal surface’ – is the barrier breached spontaneously or at operation. The so-called pseudocapsule of compressed normal tissue, often referred to in GIST, is virtually absent in small intestinal GIST and also in a number of gastric GISTs with an exophytic tumour component. Gastric GISTs, however, may have an entirely intraluminal tumour with minimal infiltration of the muscular layer ⁸⁸. For this and other less understood reasons, tumour rupture is more common in small intestinal than in gastric GISTs, with a risk more than twice as high in non-gastric compared to gastric tumours ^{52,74,131}. Anatomical issues regarding gastric GIST were explored in Paper VI (see below).

Extraperitoneal GIST, i.e. oesophageal and rectal GISTs and GIST in the retroperitoneal portion of the duodenum, do not rupture spontaneously to the abdominal cavity. Here, the risk of rupture is related to the surgical procedure, and rupture corresponds to an intralesional dissection according to the Enneking system ¹⁸. However, following a transabdominal procedure, the whole cavity is at risk.

Biological considerations. Intuitively, tumour rupture is a mechanical phenomenon related to mass, movement, surgical handling etc. However, there is also a biological aspect. GISTs that rupture have a higher MI, as shown in Paper III, and their predilection for the small intestine indicates a different biology. Other factors than size and location certainly are of importance: fragility, necrosis, vascularisation – which may all be related to an increased growth rate. In Papers V and VI, a clear association between rupture and KIT del557/558 mutations was documented in gastric GIST.

The association between tumour rupture and recurrence is well documented and by most authors the latter is taken to be a consequence of the former, which is possible but a scientifically flawed conclusion. Apart from the paper published by Rutkowski and colleagues in 2011 ⁷⁴, few studies have analysed tumours with rupture, and the present data are among the most detailed. There was no difference in time to first recurrence between tumours with and without rupture, median 27 months (range 2 to 164) *versus* 32 months (range 4 to 126), $P = 0.504$. While tumours of equal size or with equal MI generally have a better prognosis in the stomach, ruptured tumours had an equally sinister prognosis in gastric and non-gastric positions, as shown in Papers I and II. Tumour size (> 10 cm *versus* ≤ 10 cm) and tumour genotype (del557/558 *versus* other mutations) did not influence prognosis in ruptured tumours, $P = 0.280$ and 0.872 , and MI (> 10 *versus* ≤ 10) had a statistically significant but not dramatic impact, $P = 0.035$ (unpublished data). Moreover, there was no difference in RFS between patients with spontaneous and iatrogenic rupture ($P = 0.990$), which is surprising as one would expect a biological difference (*Table 7* and *Table 9*). Tumours with spontaneous and iatrogenic rupture were remarkably similar – in size (median size 11.3 cm [range 2.5 to 28.0] *versus* 12.0 cm [range 2.0 to 24.0]), in mitotic activity (median MI 11 [range 1 to 178] *versus* 9 [range 0 to 53]), anatomically (38 per cent gastric *versus* 44 per cent), and genetically (35 per cent del557/558 *versus* 40 per cent) (unpublished data). The generally accepted assumption that tumour rupture leads to recurrence, is not gainsaid by any of these observations, but the inverted logic, that some recurring tumours may have aggressive biological features that also lend them prone to rupture, is an equally valid hypothesis.

Neoadjuvant imatinib treatment for locally advanced GIST is increasing both with the objective to reduce the extent of surgery and the risk of rupture – by a reduction in size and possibly by inducing biological alterations (see Introduction, section on neoadjuvant treatment). GISTs treated with imatinib become metabolically silent, and if recurrence after tumour rupture is indeed caused by implantation and proliferation of seeded cells, the risk

could be attenuated in pre-treated tumours that rupture. This hypothesis could probably be tested in laboratory studies. Clinical data from the present study are scarce and somewhat discouraging. Of 20 patients undergoing surgery after neoadjuvant treatment, four had tumour rupture. Three of these experienced a recurrence, one peritoneal, one hepatic, and one concomitantly at both sites.

Defining ‘tumour rupture’ in GIST. So far, there has been no generally accepted definition of rupture in GIST. Conceptually, any breach in the healthy tissue containing the tumour could be termed rupture. In Paper I, a distinction was made between major defects of tumour integrity – called rupture – and minor defects, not considered rupture, listed in *Table 9*. *Tumour fracture or spillage* would unreservedly be considered rupture by most surgeons and when iatrogenic, it corresponds to an intralesional dissection ¹⁸. Tumour fracture has been described following trauma ¹⁵⁸. In the SSGXVIII/AIO trial, spillage at surgery was defined as intraoperative rupture ⁹³. Most patients reported to have unspecified tumour rupture probably fall into this category, e.g. in the original Polish and Japanese reports cited above. *Piecemeal resection* is a variation of the preceding criterion and is invariably associated with spillage. *Incisional biopsy* is usually a premeditated diagnostic procedure performed at laparotomy or laparoscopy. Fortunately, it is a rare event in sarcoma centres, and the evidence regarding this criterion is lacking. However, followed by extirpation it is technically a piecemeal resection, and it is considered rupture in the latest version of the NCCN guidelines ⁹⁹. *Blood-tinged ascites*, when present, is easily evident at laparotomy or laparoscopy. It is sometimes associated with spontaneous tumour fracture, but macroscopic signs of rupture may be missing. It strongly influences prognosis ^{131,158}, and in the SSGXVIII/AIO trial, it was a criterion of preoperative rupture ⁹³. *Gastrointestinal perforation at the tumour site*, caused by an increase in luminal pressure, tumour fragility, or transmural tumour necrosis, leads to spillage of gastrointestinal content and tumour tissue to the abdominal cavity. Of note, a contained perforation, i.e. gastrointestinal content or gas penetrating the tumour but without communication to the abdominal cavity on imaging, at surgery or during pathology review – was not considered rupture. *Microscopic adjacent infiltration* is, strictly speaking, not rupture, but represents longstanding peritoneal exposure of a biologically aggressive tumour, and even if resected *en bloc* with free margins the prognosis is poor ¹³⁰. Fibrous or inflammatory adhesions without microscopic infiltration were not considered rupture according to this criterion. In gastrointestinal carcinoma, peritoneal tumour penetration (T4a) is separated from adjacent infiltration (T4b) in the classification system, but they both confer

an increased risk of recurrence ^{159,160}. Peritoneal penetration in GIST was investigated in Paper IV (see below) and found to be without independent prognostic influence. By contrast, the present studies included eight patients with adjacent microscopic infiltration and they all relapsed. This fact should be accounted for. Neither the TNM classification nor other stratification systems make this distinction for GIST, and in the present studies, adjacent infiltration has been termed ‘rupture’, albeit semantically it is not precisely so.

Table 9. Criteria defining defects of tumour integrity in Papers I and II

Rupture	
CRITERION	CLINICAL SETTING
Fracture, spillage or intralesional dissection	Iatrogenic or spontaneous
Piecemeal resection	Iatrogenic
Incisional biopsy	Iatrogenic
Blood-tinged ascites	Spontaneous
Gastrointestinal perforation at tumour site	Spontaneous
Microscopic adjacent infiltration	Spontaneous
Minor defects of tumour integrity, not rupture	
CRITERION	CLINICAL SETTING
Percutaneous needle biopsy	Iatrogenic
Mucosal defect, contained luminal spillage or bleeding	Spontaneous
Peritoneal penetration or iatrogenic peritoneal laceration	Spontaneous or iatrogenic
R1 resection	Iatrogenic

Defined according to these criteria, tumour rupture was a strong and independent predictor of recurrence even in comparatively small series (Papers I, II, III and VI). Some of the criteria are clinically and semantically uncontroversial, like tumour spillage and intralesional dissection; some are semantically not as evident but clearly confer an inferior prognosis, like blood-tinged ascites and microscopic adjacent infiltration; and some situations would have to be termed ‘rupture’ by any reasonable definition, like incisional biopsy and spontaneous perforation through tumour, in spite of lacking evidence for a prognostic significance in the present cohort. Asare and Feig at M.D. Anderson Cancer Center, US, have deplored the weak evidence supporting the definition and objected that some criteria are contrary to common sense ^{161,162}. Nevertheless, they constitute the only clear-cut definition of ‘tumour rupture’ in GIST and have been proposed for general application ¹⁴¹. Adherence to this definition would

facilitate cross-study comparison, homogenise inclusion in prospective studies, guide clinical decisions, and perhaps of equal importance – inspire surgeons to improve their observations and reporting. Hopefully, the definition will be validated in other population-based cohorts and revised.

MINOR DEFECTS OF TUMOUR INTEGRITY

Introduced as a separate category distinct from tumour rupture in Paper I, minor defects of tumour integrity were shown to be without prognostic significance (Papers I and II).

R1 resection is associated with increased recurrence-rates in soft tissue sarcomas of most anatomical locations ²⁵, but as mentioned in the Introduction, McCarter and colleagues were unable to confirm this association in their exploratory investigation of the ACOSOG Z9001 trial ¹⁴⁶. Their analysis indicated that progressive disease could be ascribed to tumour rupture rather than involved margins. Certainly, the seeding and possible implantation associated with tumour rupture outweighs any contribution of a positive margin, and in patients with concomitant rupture and R1 resection, recurrence is more likely to be a consequence of the former. The ACOSOG Z9001 trial enrolled patients at intermediate and high risk. In Paper III, the relationship between rupture and R1 resection was explored in a population-based series. A strong association was documented, but independently, R1 resection was not an adverse factor neither among patients with rupture nor in patients without. The analysis made a sharp distinction between rupture and R1 resection. This distinction is specified in the TNM residual tumour (R) classification, which states that peritoneal breaches do not influence R status ²⁰. However, on this point, confusion prevails, even among sarcoma surgeons, and inferior outcomes after R1 resection, reported in some studies, probably derives from this ambiguity ^{147,148}. Recently, the significance of positive resection margins in GIST was explored in the EORTC/Intergroup trial, which randomised GIST patients at intermediate or high risk to adjuvant imatinib for 2 years or surgery alone ¹⁴³. Among 697 patients available for central review, there was a clear association between tumour rupture and R1 resection, but in patients without rupture, there was no difference in RFS or OS between those undergoing R0 and R1 surgery, neither in the treatment nor in the control arm. In this study, a similar but not identical definition of tumour rupture was applied. These results confirm those of Paper III and reduce the uncertainty related to a possible masking effect of adjuvant imatinib in high-risk patients undergoing R1 resection.

In sarcomas and carcinomas negatively affected by R1 surgery, local recurrence-rates are increased, reflecting residual microscopic disease at the resection margin ^{21,25}. In GIST, true local recurrence – in the gastric or bowel wall – is extremely rare, with a possible exception regarding extraperitoneal rectal GIST ¹⁶³. Recurrence in GIST is peritoneal or distant metastatic, indicating other mechanisms than local disease left behind by the surgeon.

Peritoneal penetration (PP). In gastrointestinal carcinoma, the prognostic significance of peritoneal tumour penetration, accorded a specific stage in the TNM classification – T4a - is a settled issue ^{159,160}. It is strongly associated with recurrent peritoneal disease, which increases from approximately 10 to approximately 50 per cent both in gastric and colonic cancer ^{9,160}. In GIST, a peritoneal recurrence pattern is more pronounced than in carcinomas, accounting for two thirds of recurrences, evenly distributed between isolated peritoneal disease and peritoneal disease with concurrent organ metastases ⁶⁷. One would therefore expect a negative impact on prognosis of PP in GIST like in carcinoma. The frequency and significance of PP have been investigated in a few small institutional series ^{47,164-166}. The largest, comprising 171 patients, reported PP in 32 per cent ⁴⁷. In this study, with a majority of high-risk tumours (58 per cent according to the mNIH), PP was associated with an increased risk of recurrence on multivariable analysis together with an elevated mitotic activity, tumour size, and tumour necrosis. In Paper IV, PP was present in 24 per cent. There was a significant association between PP and established risk factors (MI, tumour size, tumour site), but in this study, which excluded patients with tumour rupture, no impact on prognosis was demonstrated, neither in gastric nor in non-gastric tumours. The study included 243 patients not at high risk (71 per cent), and also in this group, PP was a common occurrence, even if significantly less frequent than in the high-risk group – 18 *versus* 39 per cent. The high frequency of peritoneal dissemination in GIST apparently unrelated to PP of the primary tumour indicates unexplored differences between mesenchymal and epithelial malignancies. Exfoliated cells from a primary tumour are thought to be the origin of peritoneal surface disease. However, a complex sequence of adhesion, penetration, proliferation and angiogenesis must be induced for carcino or sarcomatosis to become evident ¹⁶⁷. Carcinomas and sarcomas may behave differently in this respect. Moreover, there are experimental data suggesting a haematogenous route in peritoneal metastasis ¹⁶⁸, and such a route – if existent - may be of greater importance in GIST than in carcinomas, a hypothesis possibly supported by the frequent occurrence of concurrent peritoneal and liver metastases in GIST. PP may also be a dynamic phenomenon, occurring at some point in all exophytic GISTs, but present at surgery only occasionally.

However, strikingly different outcomes in gastric GISTs with exophytic and transmural growth patterns, shown in Paper VI, make this explanation less probable.

Mucosal defects or contained luminal spillage or bleeding. Among the series constituting the evidence for the mNIH classification ³⁷, one study defined mucosal perforation as tumour rupture ⁴¹. Mucosal defects with or without clinically significant bleeding are common even in low-risk gastric GIST. To consider this tumour rupture is contrary to biological and surgical sense. It is, however, *de facto* a defect of tumour integrity. In large tumours, necrotic material may spill into the gastric or intestinal lumen. If the spillage is contained during the operation without contaminating the abdominal cavity, it should not be considered tumour rupture. Endoscopic resection for small and medium sized gastric GISTs is increasing (see Introduction). The R classification applies equally to these resections, and intralesional dissections or tumour spillage without peritoneal perforation should be classified accordingly but not considered rupture. Interestingly, Song and colleagues published a series of 195 patients with endoscopically removed gastric GISTs, of whom 61 had inadvertent gastric perforation or full thickness resections ⁸⁹. Among these, there were 14 tumour ruptures, but no peritoneal recurrences were detected, neither in the rupture group nor in the group without rupture. However, these results are only indicative as the follow-up time was short.

Percutaneous core needle biopsy was included as a criterion because it technically and biologically signifies a minor defect of tumour integrity. However, performed by an experienced operator for an adequate reason it is an uncontroversial procedure not associated with increased risk of recurrence in GIST or retroperitoneal sarcomas, as stated in the Introduction ¹⁵²⁻¹⁵⁴. In Papers I and II, 34 patients (11 per cent) underwent core needle biopsy.

In Papers I and II, tumour rupture, defined according to the proposed definition, was an independent risk factor for recurrence, confirming the hypothesis for these studies. Just as important, and perhaps of greater clinical interest, was the evidence that minor defects of tumour integrity did not negatively influence outcomes. In Papers I and II, adjuvant imatinib was administered to 26 of 106 (25 per cent) patients with minor defects of tumour integrity, somewhat confounding the analysis (see Methodological discussion). In all the studies, a total number of 319 patients without rupture did not receive adjuvant treatment. Of these, 100 patients (31 per cent) had minor defects of tumour integrity. Estimated 5-year RFS was 92 per cent with minor defects *versus* 89 per cent with no defect, HR 0.70 (95 per cent c.i. 0.28 to 1.77; $P = 0.450$).

In Paper V, the relationship between tumour genotype and tumour rupture was examined in 148 patients with gastric and 61 patients with small intestinal GIST. There was an increased risk of rupture for tumours with a *KIT* exon 11 del557/558 mutations compared to other *KIT* exon 11 mutations. Rutkowski and colleagues also reported more frequent ruptures in del557/558 mutated tumours ⁷⁴. By contrast, Wozniak and colleagues were unable to confirm this association in a study with missing information on rupture in 20 per cent of the patients ⁵². In the present study, a statistically significant association was confined to gastric tumours. The frequency of del557/558 mutated tumours was similar in gastric and small intestinal tumours (23 and 20 per cent), but for small intestinal tumours harbouring other mutations, rates of rupture were higher than for gastric tumours. The small number of non-gastric del557/558 mutated tumours could explain this lack of relationship with rupture in this location. However, different outcomes for the same genotype depending on anatomical site have been demonstrated for other genotypes. *KIT* exon 11 other than del557/558 mutants have a significantly better prognosis in gastric than in non-gastric tumours ⁵². *KIT* exon 9 mutations hardly ever arise in the stomach ^{52,135}, and *PDGFRA* exon 18 mutations are extremely rare in the small bowel ^{52,135,157}. Still, their associated inferior - respectively superior prognoses - seem related to anatomical site more than genotype *per se*. Del557/558 mutations are exceptional in that they are associated with an *equally* poor prognosis in either location – gastric or non-gastric ⁵².

Tumour genotype was further examined in Paper VI, a large study of gastric GISTs with both a descriptive and a prognostic aim. Several novel findings were made: (i) *PDGFRA* mutated tumours were located in the lower two-thirds of the stomach, (ii) conversely, *KIT* mutated tumours were located in the upper two-thirds, and among these, (iii) del557/558 mutated tumours had a predilection for the upper third and was practically absent in the lower third, (iv) the proportion of tumours wild type for *KIT/PDGFRA* in this population-based cohort was much smaller than previously reported ^{37,52}, only 4 per cent, and (v) in the upper third of the stomach, over 90 per cent of tumours harboured mutations conferring sensitivity to imatinib. This latter point has an immediate clinical relevance. Mutation analysis is considered a prerequisite for adjuvant or neoadjuvant therapy. However, in parts of the world, facilities for biomolecular analysis are lacking, or services may be too expensive for routine practice ¹⁶⁹. For tumours in the upper third, starting neoadjuvant treatment without this

information seems safe, and for patients with high-risk tumours in this location, even adjuvant treatment could be contemplated.

Del557/558 mutated tumours were larger and had a higher mitotic activity than tumours harbouring other genotypes, which has also been described by others ⁵², and their increased risk of rupture, demonstrated in Paper V, was confirmed in this extended study. RFS was significantly shorter for del557/558 mutants, while RFS for *KIT* other than del557/558 mutants and *PDGFRA* mutants was similar – and excellent. From early studies ⁸³, the belief that *KIT* mutations are associated with reduced RFS compared to *PDGFRA* mutations still persists. *KIT* mutated tumours occur throughout the alimentary tract whereas *PDGFRA* mutations are confined to gastric tumours, and in the stomach they carry the same prognosis when del557/558 mutations are excluded. However, the increased risk of recurrence for del557/558 mutants could just as well be ascribed to their increased mitotic activity, and on multivariable analysis, only MI and tumour rupture were independently associated with a poor prognosis. Among 14 patients with del557/558 mutated tumours not classified at high risk, only one relapsed, once again underscoring the fact that genotype by itself does not sufficiently explain prognosis in GIST ¹³⁵. Therefore, recommending adjuvant imatinib for patients with del557/558 mutated tumours outside the high-risk category, proposed by some authors ⁵², is not supported by these data.

ANATOMICAL GASTRIC SUBSITE AND TUMOUR GROWTH PATTERN

In his seminal paper from 2005, Miettinen reports on ‘the distinct differences of gastric GISTs at different sites, including the predominance of malignant tumors and more common progressive disease for tumors in the fundus and cardia-gastroesophageal junction, in 36 per cent and 53 per cent of patients [...]. In contrast, patients with tumors located in the antrum developed progressive disease in only 8 per cent of cases...’ ¹⁵⁶. Since then, studies on tumour location within the stomach have been few, but in Paper VI, Miettinen’s investigations were pursued and his findings confirmed. For patients with tumours in the upper third, estimated 5-year RFS was 81 per cent compared to 93 per cent for patients with tumours in the lower third ($P = 0.025$). In the upper third, mitotic activity was significantly increased, tumour rupture was more frequent, and 31 per cent of the tumours were classified as high-risk compared to 12 per cent in the lower third ($P = 0.008$). The distribution of tumour genotypes was strikingly different as pointed out in the previous paragraph. Thus, the upper and lower thirds may

almost be considered two separate organs as far as GIST is concerned. The anatomical correlate for this difference is not apparent. Admittedly, the upper part of the stomach acts as a reservoir whereas the pyloric antrum churns the gastric content and has a thickened circular muscular layer and an attenuated or absent inner oblique layer ¹⁷⁰. (The gastric mucosa need not be considered in this respect.) However, no difference regarding the myenteric nerve plexuses is known to this author.

Related to these anatomical, phenotypical and genetic differences were differences in macroscopic growth patterns, examined in Paper VI. The fraction of transmural tumours was higher in the upper than in the lower third, 41 per cent *versus* 18 per cent ($P = 0.004$), whereas luminal and exophytic tumours were evenly distributed throughout the stomach. GISTs originate from the interstitial cells of Cajal, situated at different levels of the gastric wall ³⁶. *A priori*, their direction of growth would be decided by the depth of the proliferating cells and the tenacity of the surrounding tissue, and unrelated to inherent biological factors. The low number of transmural tumours in the lower third could be explained by the thickness of the muscular tunic. Nevertheless, there were associations between growth patterns and risk factors that suggest a more complex explanation: transmural tumours had significantly higher mitotic activity than luminal and exophytic tumours; 45 per cent of transmural tumours belonged to the high-risk category compared to 16 per cent of luminal/exophytic tumours ($P < 0.001$); and 40 per cent of transmural tumours harboured a del557/558 mutation compared to 8 per cent of luminal/exophytic tumours ($P < 0.001$). Most importantly, patients with transmural tumours were at increased risk of recurrence with an estimated 5-year RFS of 71 per cent compared to 96 per cent for patients with luminal or exophytic tumours ($P < 0.001$). This difference was retained when corrections were made for tumour subsite, tumour size, tumour rupture, and tumour genotype, and persisted within the high-risk category. However, a transmural growth pattern was associated with a high MI, and in multivariable analysis, only MI and tumour rupture were independently related to RFS. Thus, there seem to be factors commonly shared by tumours with a transmural growth pattern and those with an elevated mitotic activity. These factors – unknown so far - more frequently exert their influence in tumours of the upper part of the stomach.

Clinicians generally find adjuvant treatment in GIST justified when the risk of recurrence approaches 50 per cent, and based on results from the SSGXVIII/AIO trial, treatment is recommended for 3 years^{98,99,145}. In this trial, 5-year RFS increased from 48 per cent to 66 per cent in high-risk patients treated for 3 years compared to 1 year. In other malignancies, toxic therapy is accepted in the adjuvant setting even for small benefits in overall survival¹⁷¹⁻¹⁷³. However, in GIST, the overall survival benefit is uncertain. No such benefit was documented in the ACOSOG Z9001 or EORTC/Intergroup trials, but in the SSGXVIII/AIO trial, a recent report showed improved survival in patients treated for 3 years compared to 1 year⁹⁵. Treatment is long and expensive, and even if generally well tolerated, it is discontinued in approximately 25 per cent of the patients⁹³.

The classification systems are sensitive – they identify patients at risk of recurrence – but specificity is low. There are few studies reporting separate data for gastric GIST, but in a Korean-Japanese study including more than 1000 gastric GIST patients of whom only 9 per cent received adjuvant therapy, estimated 5-year RFS was 75 per cent in high-risk patients⁸⁶. In Paper II, one important finding was that patients with tumour rupture relapsed despite adjuvant treatment, a finding recently confirmed in a Polish study¹⁷⁴. With rupture, treatment should therefore be extended or possibly given for life. Without rupture, high-risk patients with $MI \leq 10$ included in Paper VI, had an estimated 5-year RFS of 95 per cent; with a luminal or exophytic growth pattern, 96 per cent; and with other than del557/558 mutated tumours, 85 per cent. Less than half of these patients received adjuvant imatinib. Therefore, prescribing adjuvant therapy indiscriminately to patients with high-risk gastric GIST cannot be justified. Based on data in Paper VI, high-risk patients without rupture and with $MI \leq 10$, regardless of size and genotype, would have approximately 10 per cent risk of recurrence without adjuvant treatment. This risk is acceptable, and the recent survival data from the SSGXVIII/AIO⁹⁵ trial do not, in the opinion of this author, alter this conclusion: with adjuvant treatment, nine patients would have to pay a considerable price for postponing recurrence in one patient. Neoadjuvant treatment for locally advanced tumours presumably at high risk is increasing, and usually precludes the assessment of MI. In lack of information on mitoses, Paper VI indicates that tumour growth pattern can serve as a surrogate marker and that adjuvant treatment could be reserved for patients with transmural tumours.

Randomised controlled trials to investigate these strategies are tempting to design but probably impossible to conduct. Prospective protocols in a multi-centre setting are the second best alternative, but most high-risk patients will have to have their treatment carefully considered and decided based on information on mitotic activity, growth pattern, size, gastric subsite and genotype.

THE ENIGMA OF PERITONEAL RECURRENCE IN GIST

Arising from this study is a question not answered: how to explain peritoneal dissemination in GIST? Two-thirds of recurrences are peritoneal, and it has been taken for granted, with reference to gastrointestinal carcinoma, that peritoneal disease is caused by transcoelomic exfoliation of tumour cells. The poor prognosis associated with tumour rupture could substantiate this: Nishida and colleagues found peritoneal metastases in 86 per cent of recurrences after tumour rupture compared to 30 per cent when rupture did not occur ¹³¹. However, the site of first recurrence does not necessarily reflect the true pattern of dissemination. Patients with tumour rupture are also at increased risk of organ metastasis, and among patients included in the present studies, the recurrence pattern was not significantly different in patients with and without rupture (unpublished data). Biologically aggressive tumours are also the ones most prone to rupture. In Papers I to IV, the prognostic significance of minor defects of tumour integrity was investigated. These are factors usually associated with increased risk of local and peritoneal recurrence in gastrointestinal carcinoma, but no relationship was evident in GIST. Furthermore, as shown in Paper VI, there are important clinical and genetic differences between exophytic and transmural tumours, but they do not differ in their ability to shed malignant cells to the peritoneal cavity. However, recurrence rates were significantly different. Therefore, exfoliation does not seem to be a sufficient factor for peritoneal sarcomatosis in GIST – and perhaps not even a necessary factor. Other unknown factors are needed. Whether these are the same as those involved in distant organ dissemination is not at all certain.

CONCLUSIONS

- (i) Tumour rupture, defined as tumour spillage or fracture, piecemeal resection, incisional biopsy, gastrointestinal perforation at the tumour site, blood-tinged ascites at laparotomy, or microscopic adjacent organ infiltration, is an independent risk factor for recurrence in GIST.
- (ii) Minor defects of tumour integrity, i.e. transabdominal core needle biopsy, peritoneal tumour penetration or iatrogenic peritoneal laceration, R1 resection, and contained luminal spillage or bleeding, are not associated with an increased risk of recurrence in GIST.
- (iii) In gastric GIST, patients without rupture have a good prognosis, even when classified at high-risk according to established stratification systems.
- (iv) Gastric GISTs harbouring a *KIT* exon 11 del557/558 mutation are at increased risk of rupture.
- (v) GISTs in the upper part of the stomach more frequently recur than do GISTs in the lower part.
- (vi) In gastric GIST, tumour genotypes have a characteristic anatomical distribution, i.e. *PDGFRA* mutated tumours are located in the lower two-thirds, *KIT* mutated tumours in the upper two-thirds, and *KIT* exon 11 del557/558 in the upper third. In the upper third, over 90 per cent of GISTs harbour mutations conferring sensitivity to imatinib. In lack of facilities for mutation analysis, neoadjuvant treatment can be started without such information for tumours in this location, and also adjuvant treatment can be considered.
- (vii) Gastric GISTs with a transmural growth pattern, as opposed to luminal or exophytic tumours, have unfavourable phenotypical and genetic characteristics and are associated with increased risk of recurrence. Luminal and exophytic tumours without rupture have an excellent prognosis irrespective of size and genotype and may not benefit from adjuvant therapy.

REFERENCES

1. International Association for the Study of Pain (IASP). <https://www.iasp-pain.org/PublicationsNews/NewsDetail.aspx?ItemNumber=10475> [accessed 7 December 2020].
2. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. American Psychiatric Publishing: Arlington VA, 2013.
3. Hyman SE. The diagnosis of mental disorders: the problem of reification. *Annu Rev Clin Psychol* 2010; **6**: 155-79.
4. Thygesen K, Alpert JS, Jaffe AS *et al*. Fourth universal definition of myocardial infarction (2018). *Eur Heart J* 2019; **40**: 237-69.
5. Turnbull RB, Kyle K, Watson FR, Spratt J. Cancer of the colon: the influence of the no-touch isolation technic on survival rates. *Ann Surg* 1967; **166**: 420-7.
6. Takii Y, Mizusawa J, Kanemitsu Y *et al*. A randomized controlled trial of the conventional technique versus the no-touch isolation technique for primary tumor resection in patients with colon cancer: Primary analysis of Japan Clinical Oncology Group study JCOG1006. *J Clin Oncol* 2019; **37** (Suppl 15): 3515.
7. Gronchi A, Strauss DC, Miceli R *et al*. Variability in Patterns of Recurrence After Resection of Primary Retroperitoneal Sarcoma (RPS): A Report on 1007 Patients From the Multi-institutional Collaborative RPS Working Group. *Ann Surg* 2016; **263**: 1002-9.
8. Eriksen MT, Wibe A, Syse A, Haffner J, Wiig JN. Inadvertent perforation during rectal cancer resection in Norway. *Br J Surg* 2004; **91**: 210-6.
9. Hompes D, Tiek J, Wolthuis A *et al*. HIPEC in T4a colon cancer: a defensible treatment to improve oncologic outcome? *Ann Oncol* 2012; **23**: 3123-9.
10. Roviello F, Marrelli D, de Manzoni G *et al*. Prospective study of peritoneal recurrence after curative surgery for gastric cancer. *Br J Surg* 2003; **90**: 1113-9.
11. Halsted WS. The Results of Operations for the Cure of Cancer of the Breast Performed at the Johns Hopkins Hospital from June, 1889, to January, 1894. *Ann Surg* 1894; **20**: 497-555.
12. Songun I, Putter H, Kranenbarg EM-K, Sasako M, van de Velde CJH. Surgical treatment of gastric cancer: 15-year follow-up results of the randomised nationwide Dutch D1D2 trial. *Lancet Oncol* 2010; **11**: 439-49.
13. Sasako M, Sano T, Yamamoto S *et al*. D2 lymphadenectomy alone or with para-aortic nodal dissection for gastric cancer. *N Engl J Med* 2008; **359**: 453-62.
14. Emmanuel A, Haji A. Complete mesocolic excision and extended (D3) lymphadenectomy for colonic cancer: is it worth that extra effort? A review of the literature. *Int J Colorect Dis* 2016; **31**: 797-804.
15. Russo P, Kim Y, Ravindran S, Huang W, Brennan MF. Nephrectomy during operative management of retroperitoneal sarcoma. *Ann Surg Oncol* 1997; **4**: 421-4.
16. Gronchi A, Lo Vullo S, Fiore M *et al*. Aggressive surgical policies in a retrospectively reviewed single-institution case series of retroperitoneal soft tissue sarcoma patients. *J Clin Oncol* 2009; **27**: 24-30.
17. Bonvalot S, Rivoire M, Castaing M *et al*. Primary retroperitoneal sarcomas: a multivariate analysis of surgical factors associated with local control. *J Clin Oncol* 2009; **27**: 31-7.

18. Enneking WF, Spanier SS, Goodman MA. A system for the surgical staging of musculoskeletal sarcoma. *Clin Orthop Relat Res* 1980; **153**: 106-20.
19. Beahrs O CD, Rubin P. *Manual for staging of cancer, American Joint Committee for Cancer Staging and End Results Reporting*. 1st Edition. Whiting Press: Chicaco IL, 1977.
20. Wittekind C, Compton CC, Greene FL, Sobin LH. TNM residual tumor classification revisited. *Cancer* 2002; **94**: 2511-6.
21. PelvEx Collaborative. Surgical and Survival Outcomes Following Pelvic Exenteration for Locally Advanced Primary Rectal Cancer: Results From an International Collaboration. *Ann Surg* 2019; **269**: 315-21.
22. Raziee HR, Cardoso R, Seevaratnam R *et al*. Systematic review of the predictors of positive margins in gastric cancer surgery and the effect on survival. *Gastric Cancer* 2012; **15** (Suppl 1): S116-24.
23. Viswanathan AN, Lee H, Hanson E, Berkowitz RS, Crum CP. Influence of margin status and radiation on recurrence after radical hysterectomy in Stage IB cervical cancer. *Int J Radiat Oncol Biol Phys* 2006; **65**: 1501-7.
24. Gundle KR, Kafchinski L, Gupta S *et al*. Analysis of Margin Classification Systems for Assessing the Risk of Local Recurrence After Soft Tissue Sarcoma Resection. *J Clin Oncol* 2018; **36**: 704-9.
25. Stojadinovic A, Leung DH, Hoos A, Jaques DP, Lewis JJ, Brennan MF. Analysis of the prognostic significance of microscopic margins in 2,084 localized primary adult soft tissue sarcomas. *Ann Surg* 2002; **235**: 424-34.
26. Karakiewicz PI, Eastham JA, Graefen M *et al*. Prognostic impact of positive surgical margins in surgically treated prostate cancer: multi-institutional assessment of 5831 patients. *Urology* 2005; **66**: 1245-50.
27. Singer S, Antonescu CR, Riedel E, Brennan MF. Histologic subtype and margin of resection predict pattern of recurrence and survival for retroperitoneal liposarcoma. *Ann Surg* 2003; **238**: 358-70; discussion 70-1.
28. Verbeke CS. Resection margins in pancreatic cancer. *Surg Clin North Am* 2013; **93**: 647-62.
29. Allard WJ, Matera J, Miller MC *et al*. Tumor cells circulate in the peripheral blood of all major carcinomas but not in healthy subjects or patients with nonmalignant diseases. *Clin Cancer Res* 2004; **10**: 6897-904.
30. Fischer JC, Niederacher D, Topp SA *et al*. Diagnostic leukapheresis enables reliable detection of circulating tumor cells of nonmetastatic cancer patients. *Proc Natl Acad Sci U S A* 2013; **110**: 16580-5.
31. Sanger N, Effenberger KE, Riethdorf S *et al*. Disseminated tumor cells in the bone marrow of patients with ductal carcinoma in situ. *Int J Cancer* 2011; **129**: 2522-6.
32. Muller C, Holtschmidt J, Auer M *et al*. Hematogenous dissemination of glioblastoma multiforme. *Science translational medicine* 2014; **6**: 247ra101.
33. Chen H, Shah AS, Girgis RE, Grossman SA. Transmission of glioblastoma multiforme after bilateral lung transplantation. *J Clin Oncol* 2008; **26**: 3284-5.
34. Peinado H, Zhang H, Matei IR *et al*. Pre-metastatic niches: organ-specific homes for metastases. *Nat Rev Cancer* 2017; **1**: 302-17.
35. Miettinen M, Lasota J. Gastrointestinal stromal tumors - definition, clinical, histological, immunohistochemical, and molecular genetic features and differential diagnosis. *Virchows Arch* 2001; **438**: 1-12.
36. Kindblom LG, Remotti HE, Aldenborg F, Meis-Kindblom JM. Gastrointestinal pacemaker cell tumor (GIPACT): gastrointestinal stromal tumors show phenotypic characteristics of the interstitial cells of Cajal. *Am J Pathol* 1998; **152**: 1259-69.

37. Joensuu H, Vehtari A, Riihimäki J *et al.* Risk of recurrence of gastrointestinal stromal tumour after surgery: an analysis of pooled population-based cohorts. *Lancet Oncol* 2012; **13**: 265-74.
38. Ducimetière F, Lurkin A, Ranchère-Vince D *et al.* Incidence of sarcoma histotypes and molecular subtypes in a prospective epidemiological study with central pathology review and molecular testing. *PLoS One* 2011; **6**: e20294.
39. Hirota S, Isozaki K, Moriyama Y *et al.* Gain-of-function mutations of c-kit in human gastrointestinal stromal tumors. *Science* 1998; **279**: 577-80.
40. Nilsson B, Bümming P, Meis-Kindblom JM *et al.* Gastrointestinal stromal tumors: the incidence, prevalence, clinical course, and prognostication in the preimatinib mesylate era - a population-based study in western Sweden. *Cancer* 2005; **103**: 821-9.
41. Tryggvason G, Gislason HG, Magnusson MK, Jonasson JG. Gastrointestinal stromal tumors in Iceland, 1990-2003: the icelandic GIST study, a population-based incidence and pathologic risk stratification study. *Int J Cancer* 2005; **117**: 289-93.
42. Sandvik OM, Søreide K, Kvaløy JT, Gudlaugsson E, Søreide JA. Epidemiology of gastrointestinal stromal tumours: single-institution experience and clinical presentation over three decades. *Cancer Epidemiol* 2011; **35**: 515-20.
43. Steigen SE, Eide TJ, Wasag B, Lasota J, Miettinen M. Mutations in gastrointestinal stromal tumors--a population-based study from Northern Norway. *APMIS : acta pathologica, microbiologica, et immunologica Scandinavica* 2007; **115**: 289-98.
44. Kawanowa K, Sakuma Y, Sakurai S *et al.* High incidence of microscopic gastrointestinal stromal tumors in the stomach. *Hum Pathol* 2006; **37**: 1527-35.
45. Corless CL, Barnett CM, Heinrich MC. Gastrointestinal stromal tumours: origin and molecular oncology. *Nat Rev Cancer* 2011; **11**: 865-78.
46. Hassan I, You YN, Shyyan R *et al.* Surgically managed gastrointestinal stromal tumors: a comparative and prognostic analysis. *Ann Surg Oncol* 2008; **15**: 52-9.
47. Goh BK, Chow PK, Yap WM *et al.* Which is the optimal risk stratification system for surgically treated localized primary GIST? Comparison of three contemporary prognostic criteria in 171 tumors and a proposal for a modified Armed Forces Institute of Pathology risk criteria. *Ann Surg Oncol* 2008; **15**: 2153-63.
48. Rossi S, Gasparotto C, Toffolatti L *et al.* Molecular and clinicopathologic characterization of GISTs of small size. *Am J Surg Pathol* 2010; **34**: 1480-91.
49. Emile JF, Brahimi S, Coindre JM, *et al.* Frequencies of KIT and PDGFRA mutations in the MolecGIST prospective population-based study differ from those of advanced GISTs. *Medical Oncology* 2012; **29**: 1765-72.
50. Corless CL, Fletcher JA, Heinrich MC. Biology of gastrointestinal stromal tumors. *J Clin Oncol* 2004; **22**: 3813-25.
51. Lasota J, Miettinen M. KIT and PDGFRA mutations in gastrointestinal stromal tumors (GISTs). *Semin Diagn Pathol* 2006; **23**: 91-102.
52. Wozniak A, Rutkowski P, Schoffski P *et al.* Tumor genotype is an independent prognostic factor in primary gastrointestinal stromal tumors of gastric origin: a european multicenter analysis based on ConticaGIST. *Clin Cancer Res* 2014; **20**: 6105-16.
53. Joensuu H, Wardelmann E, Sihto H *et al.* Effect of KIT and PDGFRA Mutations on Survival in Patients With Gastrointestinal Stromal Tumors Treated With Adjuvant Imatinib: An Exploratory Analysis of a Randomized Clinical Trial. *JAMA Oncol* 2017; **3**: 602-9.
54. Martin J, Poveda A, Llombart-Bosch A *et al.* Deletions affecting codons 557-558 of the c-KIT gene indicate a poor prognosis in patients with completely

- resected gastrointestinal stromal tumors: a study by the Spanish Group for Sarcoma Research (GEIS). *J Clin Oncol* 2005; **23**: 6190-8.
55. Wardelmann E, Losen I, Hans V *et al.* Deletion of Trp-557 and Lys-558 in the juxtamembrane domain of the c-kit protooncogene is associated with metastatic behavior of gastrointestinal stromal tumors. *Int J Cancer* 2003; **106**: 887-95.
 56. Wozniak A, Rutkowski P, Piskorz A *et al.* Prognostic value of KIT/PDGFRA mutations in gastrointestinal stromal tumours (GIST): Polish Clinical GIST Registry experience. *Ann Oncol* 2012; **23**: 353-60.
 57. Cassier PA, Fumagalli E, Rutkowski P *et al.* Outcome of patients with platelet-derived growth factor receptor alpha-mutated gastrointestinal stromal tumors in the tyrosine kinase inhibitor era. *Clin Cancer Res* 2012; **18**: 4458-64.
 58. Heinrich MC, Owzar K, Corless CL *et al.* Correlation of kinase genotype and clinical outcome in the North American Intergroup Phase III Trial of imatinib mesylate for treatment of advanced gastrointestinal stromal tumor: CALGB 150105 Study by Cancer and Leukemia Group B and Southwest Oncology Group. *J Clin Oncol* 2008; **26**: 5360-7.
 59. Antonescu CR, Besmer P, Guo T *et al.* Acquired resistance to imatinib in gastrointestinal stromal tumor occurs through secondary gene mutation. *Clin Cancer Res* 2005; **11**: 4182-90.
 60. Nishida T, Kanda T, Nishitani A *et al.* Secondary mutations in the kinase domain of the KIT gene are predominant in imatinib-resistant gastrointestinal stromal tumor. *Cancer Sci* 2008; **99**: 799-804.
 61. Miettinen M, Wang ZF, Sarlomo-Rikala M, Osuch C, Rutkowski P, Lasota J. Succinate dehydrogenase-deficient GISTs: a clinicopathologic, immunohistochemical, and molecular genetic study of 66 gastric GISTs with predilection to young age. *Am J Surg Pathol* 2011; **35**: 1712-21.
 62. Lasota J, Felisiak-Golabek A, Wasag B *et al.* Frequency and clinicopathologic profile of PIK3CA mutant GISTs: molecular genetic study of 529 cases. *Modern Pathology* 2016; **29**: 275-82.
 63. Stratakis CA, Carney JA. The triad of paragangliomas, gastric stromal tumours and pulmonary chondromas (Carney triad), and the dyad of paragangliomas and gastric stromal sarcomas (Carney-Stratakis syndrome): molecular genetics and clinical implications. *J Intern Med* 2009; **266**: 43-52.
 64. Carney JA, Sheps SG, Go VL, Gordon H. The triad of gastric leiomyosarcoma, functioning extra-adrenal paraganglioma and pulmonary chondroma. *N Engl J Med* 1977; **296**: 1517-8.
 65. Ricci R. Syndromic gastrointestinal stromal tumors. *Hered Cancer Clin Pract* 2016; **14**: 15.
 66. Fletcher CDM, Bridge JA, Hoogendorn PCW, Mertens F (eds). *WHO Classification of Tumours of Soft Tissue and Bone*. 4th Edition. WHO/IARC Classification of Tumours. IARC: Lyon, 2013.
 67. Rutkowski P, Nowecki ZI, Michej W *et al.* Risk criteria and prognostic factors for predicting recurrences after resection of primary gastrointestinal stromal tumor. *Ann Surg Oncol* 2007; **14**: 2018-27.
 68. Revheim ME, Hole KH, Bruland ØS *et al.* Multimodal functional imaging for early response assessment in GIST patients treated with imatinib. *Acta Oncol* 2014; **53**: 143-8.
 69. Brierley JD, Gospodarowicz MK, Wittekind C (eds). *TNM classification of malignant tumours (UICC)*. 8th Edition. Wiley: Oxford, 2017.
 70. Fletcher CD, Berman JJ, Corless C *et al.* Diagnosis of gastrointestinal stromal tumors: A consensus approach. *Hum Pathol* 2002; **33**: 459-65.
 71. Miettinen M, El-Rifai W, L HLS, Lasota J. Evaluation of malignancy and prognosis of gastrointestinal stromal tumors: a review. *Hum Pathol* 2002; **33**: 478-83.

72. Miettinen M, Lasota J. Gastrointestinal stromal tumors: pathology and prognosis at different sites. *Semin Diagn Pathol* 2006; **23**: 70-83.
73. Joensuu H. Risk stratification of patients diagnosed with gastrointestinal stromal tumor. *Hum Pathol* 2008; **39**: 1411-9.
74. Rutkowski P, Bylina E, Wozniak A *et al*. Validation of the Joensuu risk criteria for primary resectable gastrointestinal stromal tumour - the impact of tumour rupture on patient outcomes. *Eur J Surg Oncol* 2011; **37**: 890-6.
75. Khoo CY, Chai X, Quek R, Teo MCC, Goh BKP. Systematic review of current prognostication systems for primary gastrointestinal stromal tumors. *Eur J Surg Oncol* 2018; **44**: 388-94.
76. Cavnar MJ, Seier K, Curtin C *et al*. Outcome of 1000 Patients With Gastrointestinal Stromal Tumor (GIST) Treated by Surgery in the Pre and Post-imatinib Eras. *Ann Surg* 2019. DOI: 10.1097/SLA.0000000000003277.
77. Yanagimoto Y, Takahashi T, Muguruma K *et al*. Re-appraisal of risk classifications for primary gastrointestinal stromal tumors (GISTs) after complete resection: indications for adjuvant therapy. *Gastric Cancer* 2015; **18**: 426-33.
78. Gold JS, Gönen M, Gutiérrez A *et al*. Development and validation of a prognostic nomogram for recurrence-free survival after complete surgical resection of localised primary gastrointestinal stromal tumour: a retrospective analysis. *Lancet Oncol* 2009; **10**: 1045-52.
79. Chok AY, Goh BK, Koh YX *et al*. Validation of the MSKCC Gastrointestinal Stromal Tumor Nomogram and Comparison with Other Prognostication Systems: Single-Institution Experience with 289 Patients. *Ann Surg Oncol* 2015; **22**: 3597-605.
80. Racz JM, Brar SS, Cleghorn MC *et al*. The accuracy of three predictive models in the evaluation of recurrence rates for gastrointestinal stromal tumors. *J Surg Oncol* 2015; **111**: 371-6.
81. Bischof DA, Kim Y, Behman R *et al*. A nomogram to predict disease-free survival after surgical resection of GIST. *J Gastrointest Surg* 2014; **18**: 2123-9.
82. Joensuu H, Eriksson M, Sundby Hall K *et al*. Risk factors for gastrointestinal stromal tumor recurrence in patients treated with adjuvant imatinib. *Cancer* 2014; **120**: 2325-33.
83. Corless CL, McGreevey L, Haley A, Town A, Heinrich MC. KIT mutations are common in incidental gastrointestinal stromal tumors one centimeter or less in size. *Am J Pathol* 2002; **160**: 1567-72.
84. von Mehren M, Joensuu H. Gastrointestinal Stromal Tumors. *J Clin Oncol* 2018; **36**: 136-43.
85. Bischof DA, Kim Y, Dodson R *et al*. Conditional disease-free survival after surgical resection of gastrointestinal stromal tumors: a multi-institutional analysis of 502 patients. *JAMA Surg* 2015; **150**: 299-306.
86. Kim MC, Yook JH, Yang HK *et al*. Long-Term Surgical Outcome of 1057 Gastric GISTs According to 7th UICC/AJCC TNM System: Multicenter Observational Study From Korea and Japan. *Medicine (Baltimore)* 2015; **94**(41): e1526.
87. Xiong H, Wang J, Jia Y *et al*. Laparoscopic surgery versus open resection in patients with gastrointestinal stromal tumors: An updated systematic review and meta-analysis. *Am J Surg* 2017; **214**: 538-46.
88. Kim HH. Endoscopic treatment for gastrointestinal stromal tumor: Advantages and hurdles. *World J Gastrointest Endosc* 2015; **7**: 192-205.
89. Song S, Ren W, Wang Y *et al*. Tumor rupture of gastric gastrointestinal stromal tumors during endoscopic resection: a risk factor for peritoneal metastasis? *Endosc Int Open* 2018; **6**: E950-E6.
90. Joensuu H, Roberts PJ, Sarlomo-Rikala M *et al*. Effect of the tyrosine kinase inhibitor STI571 in a patient with a

- metastatic gastrointestinal stromal tumor. *N Engl J Med* 2001; **344**: 1052-6.
91. Eisenberg BL, Trent JC. Adjuvant and neoadjuvant imatinib therapy: current role in the management of gastrointestinal stromal tumors. *Int J Cancer* 2011; **129**: 2533-42.
 92. DeMatteo RP, Ballman KV, Antonescu CR *et al.* Adjuvant imatinib mesylate after resection of localised, primary gastrointestinal stromal tumour: a randomised, double-blind, placebo-controlled trial. *Lancet* 2009; **373**: 1097-104.
 93. Joensuu H, Eriksson M, Sundby Hall K *et al.* One vs three years of adjuvant imatinib for operable gastrointestinal stromal tumor: a randomized trial. *Jama* 2012; **307**: 1265-72.
 94. Joensuu H, Eriksson M, Sundby Hall K *et al.* Adjuvant Imatinib for High-Risk GI Stromal Tumor: Analysis of a Randomized Trial. *J Clin Oncol* 2016; **34**: 244-50.
 95. Joensuu H, Eriksson M, Sundby Hall K *et al.* Survival Outcomes Associated With 3 Years vs 1 Year of Adjuvant Imatinib for Patients With High-Risk Gastrointestinal Stromal Tumors: An Analysis of a Randomized Clinical Trial After 10-Year Follow-up. *JAMA Oncol* 2020.
 96. Casali PG, Le Cesne A, Poveda Velasco A *et al.* Time to Definitive Failure to the First Tyrosine Kinase Inhibitor in Localized GI Stromal Tumors Treated With Imatinib As an Adjuvant: A European Organisation for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group Intergroup Randomized Trial in Collaboration With the Australasian Gastro-Intestinal Trials Group, UNICANCER, French Sarcoma Group, Italian Sarcoma Group, and Spanish Group for Research on Sarcomas. *J Clin Oncol* 2015; **33**: 4276-83.
 97. Raut CP, Espat NJ, Maki RG *et al.* Efficacy and Tolerability of 5-Year Adjuvant Imatinib Treatment for Patients With Resected Intermediate- or High-Risk Primary Gastrointestinal Stromal Tumor: The PERSIST-5 Clinical Trial. *JAMA Oncol* 2018; **4**: e184060.
 98. Casali PG, Abecassis N, Aro HT *et al.* Gastrointestinal stromal tumours: ESMO-EURACAN Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2018; **29** (Suppl 4): iv68-iv78.
 99. National Comprehensive Cancer Network Guidelines. *NCCN Clinical Practice Guidelines in Oncology: Soft Tissue Sarcoma*; 2020. http://www.nccn.org/professionals/physician_gls/f_guidelines.asp [accessed 10 July 2020].
 100. Nishida T, Blay JY, Hirota S, Kitagawa Y, Kang YK. The standard diagnosis, treatment, and follow-up of gastrointestinal stromal tumors based on guidelines. *Gastric Cancer* 2016; **19**: 3-14.
 101. Joensuu H, Hohenberger P, Corless CL. Gastrointestinal stromal tumour. *Lancet* 2013; **382**: 973-83.
 102. Gastrointestinal Stromal Tumor Meta-Analysis Group (MetaGIST). Comparison of two doses of imatinib for the treatment of unresectable or metastatic gastrointestinal stromal tumors: a meta-analysis of 1,640 patients. *J Clin Oncol* 2010; **28**: 1247-53.
 103. Corless CL, Ballman KV, Antonescu CR *et al.* Pathologic and molecular features correlate with long-term outcome after adjuvant therapy of resected primary GI stromal tumor: the ACOSOG Z9001 trial. *J Clin Oncol* 2014; **32**: 1563-70.
 104. Rutkowski P, Gronchi A, Hohenberger P *et al.* Neoadjuvant imatinib in locally advanced gastrointestinal stromal tumors (GIST): the EORTC STBSG experience. *Ann Surg Oncol* 2013; **20**: 2937-43.
 105. Eisenberg BL, Harris J, Blanke CD *et al.* Phase II trial of neoadjuvant/adjuvant imatinib mesylate (IM) for advanced primary and metastatic/recurrent operable gastrointestinal stromal tumor (GIST): early results of RTOG

- 0132/ACRIN 6665. *J Surg Oncol* 2009; **99**: 42-7.
106. Hohenberger P, Langer C, Wendtner CM *et al.* Neoadjuvant treatment of locally advanced GIST: results of APOLLON, a prospective, open label phase II study in KIT- or PDGFRA-positive tumors. *J Clin Oncol* 2012; **30** (Suppl 15): 100031.
 107. Kurokawa Y, Yang HK, Cho H *et al.* Phase II study of neoadjuvant imatinib in large gastrointestinal stromal tumours of the stomach. *Br J Cancer* 2017; **117**: 25-32.
 108. Heinrich MC, Griffith DJ, Druker BJ, Wait CL, Ott KA, Zigler AJ. Inhibition of c-kit receptor tyrosine kinase activity by STI 571, a selective tyrosine kinase inhibitor. *Blood* 2000; **96**: 925-32.
 109. DeMatteo RP, Lewis JJ, Leung D, Mudan SS, Woodruff JM, Brennan MF. Two hundred gastrointestinal stromal tumors: recurrence patterns and prognostic factors for survival. *Ann Surg* 2000; **231**: 51-8.
 110. Plaat BE, Hollema H, Molenaar WM *et al.* Soft tissue leiomyosarcomas and malignant gastrointestinal stromal tumors: differences in clinical outcome and expression of multidrug resistance proteins. *J Clin Oncol* 2000; **18**: 3211-20.
 111. Blanke CD, Demetri GD, von Mehren M *et al.* Long-term results from a randomized phase II trial of standard- versus higher-dose imatinib mesylate for patients with unresectable or metastatic gastrointestinal stromal tumors expressing KIT. *J Clin Oncol* 2008; **26**: 620-5.
 112. Verweij J, Casali PG, Zalcberg J *et al.* Progression-free survival in gastrointestinal stromal tumours with high-dose imatinib: randomised trial. *Lancet* 2004; **364**: 1127-34.
 113. Hompland I, Bruland ØS, Hølmekak T *et al.* Prediction of long-term survival in patients with metastatic gastrointestinal stromal tumor: analysis of a large, single-institution cohort. *Acta Oncol* 2017; **56**: 1317-23.
 114. Li GZ, Raut CP. Targeted therapy and personalized medicine in gastrointestinal stromal tumors: drug resistance, mechanisms, and treatment strategies. *Onco Targets Ther* 2019; **12**: 5123-33.
 115. Demetri GD, van Oosterom AT, Garrett CR *et al.* Efficacy and safety of sunitinib in patients with advanced gastrointestinal stromal tumour after failure of imatinib: a randomised controlled trial. *Lancet* 2006; **368**: 1329-38.
 116. Demetri GD, Reichardt P, Kang YK *et al.* Efficacy and safety of regorafenib for advanced gastrointestinal stromal tumours after failure of imatinib and sunitinib (GRID): an international, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet* 2013; **381**: 295-302.
 117. Mir O, Cropet C, Toulmonde M *et al.* Pazopanib plus best supportive care versus best supportive care alone in advanced gastrointestinal stromal tumours resistant to imatinib and sunitinib (PAZOGIST): a randomised, multicentre, open-label phase 2 trial. *Lancet Oncol* 2016; **17**: 632-41.
 118. Montemurro M, Gelderblom H, Bitz U *et al.* Sorafenib as third- or fourth-line treatment of advanced gastrointestinal stromal tumour and pretreatment including both imatinib and sunitinib, and nilotinib: A retrospective analysis. *Eur J Cancer* 2013; **49**: 1027-31.
 119. Adenis A, Blay JY, Bui-Nguyen B *et al.* Masitinib in advanced gastrointestinal stromal tumor (GIST) after failure of imatinib: a randomized controlled open-label trial. *Ann Oncol* 2014; **25**: 1762-9.
 120. Joensuu H, Blay JY, Comandone A *et al.* Dovitinib in patients with gastrointestinal stromal tumour refractory and/or intolerant to imatinib. *Br J Cancer* 2017; **117**: 1278-85.
 121. Du CY, Zhou Y, Song C *et al.* Is there a role of surgery in patients with recurrent or metastatic gastrointestinal stromal tumours responding to imatinib: a prospective randomised trial in China. *Eur J Cancer* 2014; **50**: 1772-8.

122. Bauer S, Rutkowski P, Hohenberger P *et al.* Long-term follow-up of patients with GIST undergoing metastasectomy in the era of imatinib -- analysis of prognostic factors (EORTC-STBSG collaborative study). *Eur J Surg Oncol* 2014; **40**: 412-9.
123. Bauer S, Hartmann JT, de Wit M *et al.* Resection of residual disease in patients with metastatic gastrointestinal stromal tumors responding to treatment with imatinib. *Int J Cancer* 2005; **117**: 316-25.
124. Rutkowski P, Nyckowski P, Grzesiakowska U *et al.* The clinical characteristics and the role of surgery and imatinib treatment in patients with liver metastases from c-Kit positive gastrointestinal stromal tumors (GIST). *Neoplasma* 2003; **50**: 438-42.
125. Mussi C, Ronellenfitsch U, Jakob J *et al.* Post-imatinib surgery in advanced/metastatic GIST: is it worthwhile in all patients? *Ann Oncol* 2010; **21**: 403-8.
126. Roland CL, Bednarski BK, Watson K *et al.* Identification of preoperative factors associated with outcomes following surgical management of intra-abdominal recurrent or metastatic GIST following neoadjuvant tyrosine kinase inhibitor therapy. *J Surg Oncol* 2018; **117**: 879-85.
127. Fairweather M, Balachandran VP, Li GZ *et al.* Cytoreductive Surgery for Metastatic Gastrointestinal Stromal Tumors Treated With Tyrosine Kinase Inhibitors: A 2-institutional Analysis. *Ann Surg* 2018; **268**: 296-302.
128. An HJ, Ryu MH, Ryoo BY *et al.* The effects of surgical cytoreduction prior to imatinib therapy on the prognosis of patients with advanced GIST. *Ann Surg Oncol* 2013; **20**: 4212-8.
129. Schrage Y, Hartgrink H, Smith M *et al.* Surgical management of metastatic gastrointestinal stromal tumour. *Eur J Surg Oncol* 2018; **44**: 1295-300.
130. Takahashi T, Nakajima K, Nishitani A *et al.* An enhanced risk-group stratification system for more practical prognostication of clinically malignant gastrointestinal stromal tumors. *Int J Clin Oncol* 2007; **12**: 369-74.
131. Nishida T, Cho H, Hirota S *et al.* Clinicopathological Features and Prognosis of Primary GISTs with Tumor Rupture in the Real World. *Ann Surg Oncol* 2018; **25**: 1961-9.
132. Wada N, Kurokawa Y, Nishida T *et al.* Subgroups of patients with very large gastrointestinal stromal tumors with distinct prognoses: a multicenter study. *J Surg Oncol* 2014; **109**: 67-70.
133. Costa J, Wesley RA, Glatstein E, Rosenberg SA. The grading of soft tissue sarcomas. Results of a clinicohistopathologic correlation in a series of 163 cases. *Cancer* 1984; **53**: 530-41.
134. Guillou L, Coindre JM, Bonichon F *et al.* Comparative study of the National Cancer Institute and French Federation of Cancer Centers Sarcoma Group grading systems in a population of 410 adult patients with soft tissue sarcoma. *J Clin Oncol* 1997; **15**: 350-62.
135. Joensuu H, Rutkowski P, Nishida T *et al.* KIT and PDGFRA mutations and the risk of GI stromal tumor recurrence. *J Clin Oncol* 2015; **33**: 634-42.
136. Marrelli D, Polom K, de Manzoni G, Morgagni P, Baiocchi GL, Roviello F. Multimodal treatment of gastric cancer in the west: Where are we going? *World J Gastroenterol* 2015; **21**: 7954-69.
137. Ghidini M, Petrelli F, Tomasello G. Right Versus Left Colon Cancer: Resectable and Metastatic Disease. *Curr Treat Options in Oncol* 2018; **19**: 31.
138. Dematteo RP, Gold JS, Saran L *et al.* Tumor mitotic rate, size, and location independently predict recurrence after resection of primary gastrointestinal stromal tumor (GIST). *Cancer* 2008; **112**: 608-15.
139. Emory TS, Sobin LH, Lukes L, Lee DH, O'Leary TJ. Prognosis of gastrointestinal smooth-muscle (stromal) tumors: dependence on anatomic site. *Am J Surg Pathol* 1999; **23**: 82-7.

140. Bai X, Zhou W, Li Y, Lin G. Transanal endoscopic microsurgery with alternative neoadjuvant imatinib for localized rectal gastrointestinal stromal tumor: a single center experience with long-term surveillance. *Surg Endosc* 2020. DOI: <https://doi.org/10.1007/s00464-020-07837-5>
141. Nishida T, Hølmekbakk T, Raut CP, Rutkowski P. Defining Tumor Rupture in Gastrointestinal Stromal Tumor. *Ann Surg Oncol* 2019; **26**: 1669-75.
142. Antonescu CR, Sommer G, Sarran L *et al.* Association of KIT exon 9 mutations with nongastric primary site and aggressive behavior: KIT mutation analysis and clinical correlates of 120 gastrointestinal stromal tumors. *Clin Cancer Res* 2003; **9**: 3329-37.
143. Gronchi A, Bonvalot S, Poveda Velasco A *et al.* Quality of Surgery and Outcome in Localized Gastrointestinal Stromal Tumors Treated Within an International Intergroup Randomized Clinical Trial of Adjuvant Imatinib. *JAMA Surg* 2020: e200397.
144. Moran MS, Schnitt SJ, Giuliano AE *et al.* Society of Surgical Oncology-American Society for Radiation Oncology consensus guideline on margins for breast-conserving surgery with whole-breast irradiation in stages I and II invasive breast cancer. *Ann Surg Oncol* 2014; **21**: 704-16.
145. Nishida T. Asian consensus guidelines for gastrointestinal stromal tumor: what is the same and what is different from global guidelines. *Transl Gastroenterol Hepatol* 2018; **3**: 11.
146. McCarter MD, Antonescu CR, Ballman KV *et al.* Microscopically positive margins for primary gastrointestinal stromal tumors: analysis of risk factors and tumor recurrence. *J Am Coll Surg* 2012; **215**: 53-9; discussion 59-60.
147. Åhlén J, Karlsson F, Wejde J, Nilsson IL, Larsson C, Bränström R. Wide Surgical Margin Improves the Outcome for Patients with Gastrointestinal Stromal Tumors (GISTs). *World J Surg* 2018; **42**: 2512-21.
148. Zhi X, Jiang B, Yu J *et al.* Prognostic role of microscopically positive margins for primary gastrointestinal stromal tumors: a systematic review and meta-analysis. *Sci Rep* 2016; **6**: 21541.
149. Rutkowski P, Skoczylas J, Wisniewski P. Is the Surgical Margin in Gastrointestinal Stromal Tumors Different? *Visc Med* 2018; **34**: 347-52.
150. Joe BN, Essman LJ. Breast biopsy. UpToDate. <https://www.uptodate.com/contents/breast-biopsy> [accessed 17 June 2020].
151. Robertson EG, Baxter G. Tumour seeding following percutaneous needle biopsy: the real story! *Clin Radiol* 2011; **66**: 1007-14.
152. Berger-Richardson D, Swallow CJ. Needle tract seeding after percutaneous biopsy of sarcoma: Risk/benefit considerations. *Cancer* 2017; **123**: 560-7.
153. Eriksson M, Reichardt P, Sundby Hall K *et al.* Needle biopsy through the abdominal wall for the diagnosis of gastrointestinal stromal tumour - Does it increase the risk for tumour cell seeding and recurrence? *Eur J Cancer* 2016; **59**: 128-33.
154. van Houdt WJ, NS IJ, Marjolein Schrijver A, *et al.* Oncological Outcome After Diagnostic Biopsies in Gastrointestinal Stromal Tumors: A Retrospective Cohort Study. *Ann Surg* 2019. DOI: 10.1097/sla.0000000000003744.
155. Gourgou-Bourgade S, Cameron D, Poortmans P *et al.* Guidelines for time-to-event end point definitions in breast cancer trials: results of the DATECAN initiative (Definition for the Assessment of Time-to-event Endpoints in CANcer trials). *Ann Oncol* 2015; **26**: 873-9.
156. Miettinen M, Sobin LH, Lasota J. Gastrointestinal stromal tumors of the stomach: a clinicopathologic, immunohistochemical, and molecular genetic study of 1765 cases with long-term follow-up. *Am J Surg Pathol* 2005; **29**: 52-68.

157. Miettinen M, Makhoul H, Sobin LH, Lasota J. Gastrointestinal stromal tumors of the jejunum and ileum: a clinicopathologic, immunohistochemical, and molecular genetic study of 906 cases before imatinib with long-term follow-up. *Am J Surg Pathol* 2006; **30**: 477-89.
158. Hohenberger P, Ronellenfitsch U, Oladeji O *et al.* Pattern of recurrence in patients with ruptured primary gastrointestinal stromal tumour. *Br J Surg* 2010; **97**: 1854-9.
159. Gunderson LL, Jessup JM, Sargent DJ, Greene FL, Stewart AK. Revised TN categorization for colon cancer based on national survival outcomes data. *J Clin Oncol* 2010; **28**: 264-71.
160. Sano T, Coit DG, Kim HH *et al.* Proposal of a new stage grouping of gastric cancer for TNM classification: International Gastric Cancer Association staging project. *Gastric Cancer* 2017; **20**: 217-25.
161. Asare EA, Feig BW. Raining Frogs, Flying Horses, and Defining Tumor Rupture in GIST. *Ann Surg Oncol* 2019; **26**: 1601-3.
162. Hølmekjær T, Nishida T, Rutkowski P, Raut CP. Defining Rupture in Gastrointestinal Stromal Tumor: Semantics and Prognostic Value. *Ann Surg Oncol* 2019; **26**: 2304-5.
163. Cavnar MJ, Wang L, Balachandran VP *et al.* Rectal Gastrointestinal Stromal Tumor (GIST) in the Era of Imatinib: Organ Preservation and Improved Oncologic Outcome. *Ann Surg Oncol* 2017; **24**: 3972-80.
164. Vallböhmer D, Marcus HE, Baldus SE *et al.* Serosal penetration is an important prognostic factor for gastrointestinal stromal tumors. *Oncology Reports* 2008; **20**: 779-83.
165. Agaimy A, Vassos N, Wunsch PH, Hohenberger W, Hartmann A, Croner RS. Impact of serosal involvement/extramural growth on the risk of synchronous and metachronous peritoneal spread in gastrointestinal stromal tumors: proposal for a macroscopic classification of GIST. *Int J Clin Exp Pathol* 2012; **5**: 12-22.
166. Cananzi FC, Lorenzi B, Belgaumkar A, Benson C, Judson I, Mudan S. Prognostic factors for primary gastrointestinal stromal tumours: are they the same in the multidisciplinary treatment era? *Langenbecks Arch Surg* 2014; **399**: 323-32.
167. Jayne D. Molecular biology of peritoneal carcinomatosis. In: Ceelen WP (ed.). *Peritoneal carcinomatosis. A multidisciplinary approach*. Springer: New York, 2007.
168. Pradeep S, Kim SW, Wu SY *et al.* Hematogenous metastasis of ovarian cancer: rethinking mode of spread. *Cancer Cell* 2014; **26**: 77-91.
169. Florindez J, Trent J. Low Frequency of Mutation Testing in the United States: An Analysis of 3866 GIST Patients. *Am J Clin Oncol* 2020; **43**: 270-8.
170. Standring S (ed.) *Gray's Anatomy*. 41st Edition. Elsevier: 2016.
171. André T, Boni C, Mounedji-Boudiaf L *et al.* Oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for colon cancer. *N Engl J Med* 2004; **350**: 2343-51.
172. Sarcoma Meta-analysis Collaboration. Adjuvant chemotherapy for localised resectable soft tissue sarcoma in adults: meta-analysis of individual data. *Lancet* 1997; **350**: 1647-54.
173. Neoptolemos JP, Stocken DD, Friess H *et al.* A randomized trial of chemoradiotherapy and chemotherapy after resection of pancreatic cancer. *N Engl J Med* 2004; **350**: 1200-10.
174. Rutkowski P, Ziętek M, Cybulska-Stopa B *et al.* The analysis of 3-year adjuvant therapy with imatinib in patients with high-risk molecular profiled gastrointestinal stromal tumors (GIST) treated in routine practice. *Eur J Surg Oncol* 2020. <https://doi.org/10.1016/j.ejso.2020.08.004>.

Annals of Surgical Oncology. Article in press.

Title: Integrating anatomical, molecular and clinical risk factors in gastrointestinal stromal tumor of the stomach

Running title: Anatomical risk factors in gastric GIST

Toto Hølmekbakk, MD¹, Anne Marit Wiedswang, MD², Leonardo A. Meza-Zepeda, PhD^{3,4},
Ivar Hompland, MD, PhD⁵, Ingvild V. K. Lobmaier, MD⁶, Jeanne-Marie Berner, PhD⁶,
Stephan Stoldt, MD¹, Kjetil Boye, MD, PhD^{5,3}

¹Department of Abdominal and Pediatric Surgery, Oslo University Hospital, The Norwegian Radium Hospital, Oslo, Norway

²Department of Radiology, Oslo University Hospital, The Norwegian Radium Hospital, Oslo, Norway

³Department of Tumor Biology, Institute for Cancer Research, Oslo University Hospital, The Norwegian Radium Hospital, Oslo, Norway

⁴Genomics Core Facility, Department of Core Facilities, Institute for Cancer Research, Oslo University Hospital, The Norwegian Radium Hospital, Oslo, Norway

⁵Department of Oncology, Oslo University Hospital, The Norwegian Radium Hospital, Oslo, Norway

⁶Department of Pathology, Oslo University Hospital, The Norwegian Radium Hospital, Oslo, Norway

Corresponding Author: Dr Toto Hølmekbakk, Department of Abdominal and Pediatric Surgery, Oslo University Hospital, The Norwegian Radium Hospital, Box 4953 Nydalen, 0424 Oslo, Norway. Telephone: +47 92 82 44 29. E-mail: toto.holmekbakk@ous-hf.no

Disclosure: The authors declare no conflicts of interest.

Synopsis: Genotypes have a characteristic anatomical distribution in gastric GIST. Patients with exophytic or luminal tumors without rupture, irrespective of size, have an excellent prognosis and may not benefit from adjuvant therapy.

Abstract

Background: Adjuvant imatinib for three years is recommended to patients with high-risk gastrointestinal stromal tumor (GIST). Risk stratification is inaccurate, and risk assessments are further complicated by the increased use of neoadjuvant treatment. Anatomical criteria for prognostication have not been investigated.

Methods: Clinical, molecular and anatomical variables were retrospectively studied in a population-based cohort of 295 patients with gastric GIST resected between 2000 and 2018. Gastric subsite was divided into the upper, middle and lower thirds. Growth pattern was classified as luminal, exophytic or transmural based on imaging and surgical reports.

Results: Of 113 tumors in the upper third of the stomach, 103 (91.2%) were *KIT* mutated, 7 (6.2%) were *PDGFRA* mutated, and 104 (92.0%) harbored genotypes sensitive to imatinib. Transmural tumors were strongly associated with a high mitotic index (MI). Five-year recurrence-free survival (RFS) was 71% for patients with transmural tumors vs. 96% with luminal or exophytic tumors, hazard ratio 8.45 (95% confidence interval [CI] 3.69-19.36), $P<0.001$; in high-risk patients 46% with transmural tumors vs. 83% with luminal or exophytic tumors, hazard ratio 4.47 (95% CI 1.71-11.66), $P=0.001$. Among 134 patients with tumors >5 cm, there were 29 recurrences. Only 5 patients with exophytic or luminal tumors had recurrent disease, of whom 4 had tumor rupture. Five-year RFS for patients with exophytic/luminal tumors >5 cm without rupture was 98%.

Conclusions: In the upper third, over 90% of tumors were sensitive to imatinib. Patients with exophytic or luminal tumors without rupture, irrespective of size, had an excellent prognosis and may not benefit from adjuvant therapy.

Introduction

Adjuvant imatinib treatment for three years is recommended to patients with localized gastrointestinal stromal tumor (GIST) at increased risk of disease recurrence¹⁻³. These patients are identified by the established stratification systems with high sensitivity⁴⁻⁷. However, specificity is low, and most patients with high-risk gastric GIST will not relapse without adjuvant therapy. Neoadjuvant treatment for six to 12 months is increasingly used for locally advanced tumors and tumors in surgically difficult positions, e.g. the gastric cardia, and effectively reduces tumor volume⁸. Upon neoadjuvant treatment, mitoses disappear in the surgical specimen, precluding accurate risk stratification, and mitotic index (MI) cannot be properly assessed in preoperative biopsies. Nevertheless, continued imatinib treatment postoperatively is recommended and generally practiced in patients submitted to neoadjuvant therapy^{3,8}. Mutation analysis is considered a prerequisite for adjuvant or neoadjuvant therapy, as approximately 25% of GISTs harbor a genotype not sensitive to imatinib^{9,10}. However, in parts of the world, facilities for biomolecular analysis are lacking, or services may be too expensive for routine practice¹¹.

Gastric GISTs carry a better prognosis than tumors of the small intestine¹⁰. Smaller size and lower mitotic activity partly explain this, but gastric GISTs still have a favorable prognosis compared to non-gastric GISTs of similar size and with comparable MI¹². Molecular characteristics have predictive as well as prognostic implications in GIST, and they differ in gastric and non-gastric tumors. Tumors wild type for *KIT* or *platelet-derived growth factor receptor-α* (*KIT/PDGFRα*) and *PDGFRα* exon 18 D842V mutated tumors are not sensitive to imatinib, and the latter is almost exclusively found in the stomach⁹. These genotypes have a good prognosis, whereas tumors with *KIT* exon 11 deletions involving codons 557 and 558 (del557/558) are associated with an aggressive phenotype in the stomach⁹.

Mutation status is not incorporated in the staging systems, neither are anatomical features. Organ subsite is a recognized prognostic variable in gastrointestinal carcinomas^{13,14}, but has hardly been investigated in GIST. Characteristic of gastric GIST is their macroscopic growth pattern: luminal, exophytic, or transmural. Whether these patterns represent similar neoplasms arising from different layers of the gastric wall or are associated with different clinical or biological properties, is so far not known.

The present study was undertaken to give a comprehensive picture of gastric GIST in a population-based series with the aim of improving prognostic accuracy, thereby making adjunctive medical treatment more specific.

Methods

Patients, treatment and follow-up

Patients who underwent complete excision (R0/R1) of primary gastric GIST between January 1st 2000 and August 31st 2018 were identified in the sarcoma database of the Oslo University Hospital. Patients with synchronous metastases and multifocal primary GIST were excluded. Oslo University Hospital is a sarcoma center for the South-East Health Region of Norway with a population of 2.9 million. From all hospitals and laboratories of pathology, reporting cancer cases to the Norwegian Cancer Registry is mandatory. To verify the completeness of the sarcoma database at Oslo University Hospital this registry was contacted. The number of patients with resected, localized gastric GIST in the sarcoma database corresponded to 98% of the patients reported to the Norwegian Cancer Registry from the South-East Region.

The diagnosis of GIST was made by sarcoma pathologists according to established criteria¹⁵.

The administration of adjuvant imatinib and follow-up have been described previously¹⁶.

Recurrence was recorded if verified on biopsy or indisputable on computed tomography (CT).

Recurrence-free survival (RFS) was measured from the date of surgery to recurrence. Patients were censored at the date of the latest CT, and for patients who died from surgical complications, at the date of surgery. The study was approved by the institutional Data Protection Officer and mutation analysis using next-generation sequencing (NGS) by the Regional Committee for Medical and Health Research Ethics of South-East Norway (No. 2010/1244). Written informed consent was obtained.

Clinical data and definitions

Prospectively collected data from the database were supplemented from medical records and additional mutation analyses were performed. Mitoses were only counted in imatinib-naïve patients and not in biopsies. Risk of recurrence was stratified according to the modified National Institutes of Health (mNIH) criteria⁵. For patients who received neoadjuvant imatinib treatment, only those with tumors larger than 10 cm on initial imaging or tumor rupture were stratified (high risk) (Tables 1 and 2). Tumor rupture was defined according to the Oslo definition¹⁶. Gastric subsite was divided into the upper, middle and lower thirds (Fig. 1a). Macroscopic growth pattern was classified as *luminal*, not affecting the peritoneal contour of the stomach; *exophytic*, not affecting the mucosal contour; or *transmural*, affecting both the peritoneal and mucosal contours (Fig. 1b). The assessment was done on CT imaging independently by a radiologist (AMW) and a surgeon (TH) (interobserver agreement 90·0%, $\kappa=0\cdot85$) and resolved by consensus when not in agreement. If the surgical report indicated a growth pattern different from the images, the surgical assessment was chosen. Patients with a history of mucosal ulceration (bleeding) or ulceration detected on microscopy/endoscopy were by definition classified with a luminal component; similarly, microscopic serosal penetration was equivalent to an exophytic component.

Mutation analysis

Mutation analysis was performed routinely on all intermediate and high-risk tumors and selectively on tumors at low or very low risk. Genomic DNA was extracted from fresh frozen or paraffin-embedded tumor tissue and exons 9, 11, 13 and 17 of *KIT* and exons 12, 14 and 18 of *PDGFRA* were analyzed by Sanger sequencing and categorized as described previously¹⁷. Tumors not analyzed routinely were analyzed using the AmpliSeq for Illumina Cancer Hotpot Panel version 2, examining hotspot regions of 50 cancer genes, including *KIT* exons 2, 9, 10, 11, 13, 15, 17 and 18, as well as *PDGFRA* exons 12, 14, 15 and 18. NGS libraries were generated from 100 ng of genomic DNA following Illumina's protocol. Libraries were normalized using the AmpliSeq Library Equalizer kit from Illumina and sequenced paired end 2 x 150 base pairs on an Illumina MiSeq instrument. Sequence reads were mapped to the Human UCSC hg19 reference genome and variants called using the DNA Amplicon workflow version 2.1.0.19 using the Illumina's Local Run Manager software. Mutation calls were manually inspected on the Integrative Genomics Viewer.

Statistical analysis

RFS was estimated by the Kaplan-Meier method and compared using the log-rank test. Multivariable survival analysis was performed using the Cox proportional hazards regression model. Associations between variables were investigated with two-tailed Fisher's exact test or Pearson's χ^2 test for categorical variables and independent Mann-Whitney *U* test or Kruskal-Wallis test for continuous variables. Interobserver agreement was evaluated with Cohen's κ test. $P < 0.05$ was considered statistically significant. IBM SPSS Statistics for Windows version 25.0 (Armonk, NY, USA) was used.

Results

During the study interval, 297 patients underwent complete resection of primary gastric GIST. Two patients had multifocal gastric disease and were not included in the study cohort, which therefore comprised 295 patients. According to the mNIH criteria, 77 tumors (26.2%) were classified as high-risk (Table 1). Forty-seven patients (15.9%), all at high risk, received adjuvant imatinib. In 14 (29.8%) patients, treatment was prematurely discontinued. Eleven patients (3.7%) received neoadjuvant treatment, and only one of these had a tumor ≤ 10 cm without rupture and was assigned to the unspecific risk category (Tables 1 and 2). Treatment was continued postoperatively in 10 patients. Three patients were still treated by the end of the study period. Further characteristics are shown in Table 1.

Tumor location

One hundred twenty-two tumors were located in the upper third of the stomach (41.8%), 120 in the middle (41.1%), and 50 in the lower third (17.1%) (Table 1 and Fig. 2). There was no difference in size between tumors at different sites (Fig. 2 and Supplementary Fig. S1a).

Mitotic activity was higher in tumors of the upper third, with a median MI of 3 (range 0-130) vs. 2 (range 0-166) for those of the middle, and 2 (range 0-12) for the lower third ($P = 0.020$) (Supplementary Fig. S1d). Non-high-risk tumors were more common in the lower third compared to the upper two thirds: 88.0 vs. 71.8% ($P = 0.019$) (Table 2).

Tumor growth pattern

The distribution of luminal, exophytic and transmural tumors was even (Table 1). The proportion of transmural tumors was higher in the upper third (41.0% vs. 28.4% in the middle/lower third, $P = 0.032$) (Table 2 and Fig. 2). Transmural tumors had a higher MI, median 3 (range 0-166), than exophytic and luminal tumors, median 2 (range 0-32) and 2 (range 0-14), $P < 0.001$ (Supplementary Fig. S1e). Transmural tumors were also larger

(median size 6.0 cm [range 2.0-30.0]) than exophytic (median size 5.3 cm [range 0.5-34.0]) and luminal tumors (median size 4.0 cm [range 0.5-10.2], $P < 0.001$) (Supplementary Fig. S1b). Of ruptured tumors, 15 were transmural, 7 exophytic and 1 luminal ($P = 0.001$). Forty-five per cent of transmural tumors were high-risk compared to 16.5% of exophytic/luminal tumors ($P < 0.001$).

Tumor genotype

Mutation analysis was successfully performed in 256 patients (86.8%) (Table 1). Only 11 (4.3%) patients had tumors wild type for *PDGFRA/KIT*. The majority of *PDGFRA* mutated tumors were located in the lower two thirds of the stomach, whereas *KIT* mutated tumors had a predilection for the upper end (Table 2 and Fig. 2). This pattern was even more pronounced for del557/558 mutated tumors: 33 (70.2%) were located in the upper third and only one (2.1%) in the lower third. In the upper third, only 9 of 113 patients (8.0%) had tumors with genotypes insensitive to imatinib (*PDGFRA* exon 18 D842V or *PDGFRA/KIT* wild-type). *PDGFRA* and other than del557/558 *KIT* mutated tumors typically presented a luminal or exophytic pattern, whereas 72.3% of tumors with a del557/558 mutation were transmural ($P < 0.001$) (Table 2). Del557/558 mutated tumors had a higher mitotic activity than tumors with other *KIT* and *PDGFRA* mutations, median 12 (range 0-166) vs. median 2 (range 0-53), ($P < 0.001$) (Table 2 and Supplementary Fig. S1f). Rupture was recorded in 11 tumors (23.4%) with del557/558 and in 12 tumors (5.7%) of other genotypes ($P = 0.001$). Among 134 males with known mutation status, 31 had del557/558 mutated tumors (23.1%), whereas this genotype was detected in only 17 of 122 females (13.9%), ($P = 0.037$).

Recurrence

All patients were included in survival analysis. After a median follow-up period of 46 months (range 0-185), 35 recurrences were documented. Estimated 5-year RFS was 88%. RFS was shorter for patients with GIST in the upper third than in the middle or lower third, 81% vs.

92% at 5 years, hazard ratio (HR) 2.25 (95% confidence interval [CI] 1.12-4.52), $P = 0.020$ (Fig. 3a). Only seven recurrences were seen in patients with luminal or exophytic tumors, corresponding to a 5-year RFS of 96%. With transmural tumors, 5-year RFS was 71%, HR 8.45 (95% CI 3.69-19.36), $P < 0.001$ (Fig. 3b). In the high-risk category, 5-year RFS was 46% with transmural tumors vs. 83% with luminal or exophytic tumors, HR 4.47 (95% CI 1.71-11.66), $P = 0.001$. Patients with del557/558 mutated tumors had a 5-year RFS of 58%, which was inferior to patients with other than del557/558 *KIT* mutations, 94%, HR 8.43 (95% CI 3.71-19.17), $P < 0.001$; and to patients with *PDGFRA* mutations, 95%, HR 9.65 (95% CI 2.86-32.59), $P < 0.001$ (Fig. 3c). With tumor rupture, 5-year RFS was 36% vs. 93% without rupture, HR 15.29 (95% CI 7.78-30.01), $P < 0.001$. In multivariable survival analysis with sex, tumor size, MI, tumor rupture, growth pattern and genotype as covariates, only MI and tumor rupture were independently related to recurrence, HR 22.23 (95% CI 5.06-97.60), $P < 0.001$, and HR 7.53 (95% CI 1.50-37.76), $P = 0.014$ (Supplementary Table S1).

Tumor growth pattern as a supplement to established risk stratification criteria

In lack of information on mitotic activity, e.g. after neoadjuvant therapy, tumor size and rupture are the only established criteria for risk stratification. However, transmural growth was strongly associated with high MI, and the results reported above indicate that patients with luminal or exophytic tumors without rupture have a good prognosis despite an unfavorable tumor size. We therefore investigated patients with tumors larger than 5 cm. This group included 60 patients with transmural tumors (44.8%), 73 patients with exophytic or luminal tumors (54.5%), and one patient with unspecified tumor growth pattern. Among the 73 patients with exophytic or luminal tumors, 7 patients had tumor rupture: only 5 recurrences were observed, and 4 of these had tumor rupture. In the 66 patients with exophytic or luminal tumors > 5 cm without rupture, 5-year RFS was 98%. Fifty-five of these 66 patients (83.3%) did not receive adjuvant treatment. By contrast, there were 45 patients with transmural tumors

> 5 cm without rupture, and among these, 10 recurrences were detected, corresponding to an estimated 5-year RFS of 77% ($P < 0.001$). In this group, 26 patients (57.8%) did not receive adjuvant treatment.

Discussion

The present investigation has documented that tumor genotypes have a characteristic anatomical distribution in gastric GIST. This finding is clinically relevant, as over 90 per cent of tumors in the upper third harbored mutations sensitive to imatinib. Furthermore, a transmural growth pattern was strongly associated with high mitotic activity and recurrent disease. By contrast, patients with exophytic or luminal tumors rarely relapsed even when assigned to the high-risk category

A rare disease, and still a young entity, much of our understanding of GIST has been provided by academic centers with biased referral patterns or trials with selected participants. In these reports, the proportion of tumors wild type for *KIT/PDGFR*A is stated to be 10 to 20% - higher than in the present series, 4%^{9,10}. This discrepancy may also reflect technical shortcomings in old studies. *PDGFR*A mutated tumors had a strong predilection for the middle and lower third of the stomach. In the upper third, 92% of the tumors harbored genotypes sensitive for imatinib, and neoadjuvant treatment might be started in lack of mutation analysis. In countries and centers without facilities for molecular testing, adjuvant treatment could also be considered for high-risk tumors in this location.

For patients with GIST in the upper end of the stomach, RFS was shorter than with tumors in the lower end. This observation was made by Miettinen and colleagues in 2005, but has been given little attention¹⁸. In the current study, among patients with tumors in the lower third, there were only three recurrences, representing an estimated 5-year RFS of 93% vs. 81% for

patients with tumors in the upper third. This difference seems related to mitotic activity and the distribution of tumor genotypes, more specifically to tumors with a del557/558 mutation. The association between del557/558 mutated tumors and an elevated MI has been described by others⁹, but their concurrence in the upper part of the stomach – as well as the overrepresentation of del557/558 mutated tumors in gastric GISTs of men - are to our knowledge novel findings.

GISTs originate from the interstitial cells of Cajal, situated at different levels of the gastric wall. *A priori*, their direction of growth would be decided by the depth of the proliferating cells and the tenacity of the surrounding tissue, and unrelated to inherent biological factors. Nevertheless, there were associations between growth patterns and risk factors that suggest a more complex explanation: transmural tumors both had a higher MI and a higher frequency of del557/558 mutations than luminal or exophytic tumors. Most importantly, patients with transmural tumors had significantly reduced RFS. This difference remained when tumors of the low-risk lower third were excluded ($P < 0.001$), when ruptured tumors were excluded ($P < 0.001$), when del557/558 mutated tumors were excluded ($P < 0.001$), and when tumors > 10 cm were excluded ($P < 0.001$). However, there was a strong association between a transmural growth pattern and high mitotic activity. Neoadjuvant treatment with imatinib is increasing and precludes the assessment of MI. In these patients, risk stratification is uncertain, often impossible. In lack of information on mitoses, our data suggest that growth pattern can be used as a surrogate variable after neoadjuvant therapy. Such treatment is rarely indicated for tumors smaller than 5 cm, but in larger tumors without rupture, withholding continued adjuvant treatment for those with exophytic or luminal tumors seems safe. In the present cohort, this group would have comprised 66 patients, including 22 truly at high risk, and there was only one recurrence. A minority of these patients, 17%, received adjuvant treatment.

Del557/558 mutations were closely associated with established risk factors of greater intrinsic impact, and recommending adjuvant imatinib for patients with del557/558 mutated tumors outside the high-risk category, proposed by some authors⁹, is not supported by the present data.

This study has limitations, most importantly its retrospective design and the inconsistent use of adjuvant imatinib, which confounds the analysis of recurrence. Of 35 patients who relapsed, 20 received adjuvant therapy. However, among the 15 who did not, two patients had low-risk tumors and would not have been treated by any standard; three patients at high risk had imatinib insensitive tumors; and of the remaining 10 patients, six had high-risk transmural tumors with imatinib sensitive genotypes and would have received treatment according to our suggested recommendations.

Conclusions

In this population-based study of patients with gastric GIST, tumor genotypes had a characteristic anatomical distribution. In the upper third, over 90% of tumors were sensitive to imatinib. A transmural growth pattern was a predictor of poor outcome, related to an elevated MI. Exophytic and luminal tumors, irrespective of size, were associated with an excellent prognosis without rupture, and these patients may not benefit from adjuvant treatment.

Acknowledgement: The authors thank Ms Karen-Marie Heintz for excellent technical assistance and Dr Ellen Tenstad, Science Shaped, for designing Figures 1 and 2. Sequencing services were provided by the Genomics Core Facility (oslo.genomics.no) at Oslo University Hospital. In the present study, information from the Cancer Registry of Norway has been used. The interpretation and reporting of these data are the sole responsibility of the authors, and no endorsement by the Cancer Registry of Norway should be inferred.

Funding: The study was supported by The Norwegian Cancer Society (grant number 5790283 to K.B.), the South-East Norway Regional Health Authority (grant number 2019064 to K.B.), and the Legacy of the Norwegian Radium Hospital (grant number 182007 to T.H. and K.B.).

References

1. Casali PG, Abecassis N, Aro HT, et al. Gastrointestinal stromal tumours: ESMO-EURACAN Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2018;29(Suppl 4):iv68-iv78.
2. Nishida T. Asian consensus guidelines for gastrointestinal stromal tumor: what is the same and what is different from global guidelines. *Transl Gastroenterol Hepatol.* 2018;3:11.
3. National Comprehensive Cancer Network Guidelines. NCCN Clinical Practice Guidelines in Oncology: Soft Tissue Sarcoma; 2020. Available at: http://www.nccn.org/professionals/physician_gls/f_guidelines.asp. Accessed 10 July 2020.
4. Miettinen M, Lasota J. Gastrointestinal stromal tumors: pathology and prognosis at different sites. *Semin Diagn Pathol.* 2006;23(2):70-83.
5. Joensuu H. Risk stratification of patients diagnosed with gastrointestinal stromal tumor. *Hum Pathol.* 2008;39(10):1411-9.
6. Rutkowski P, Bylina E, Wozniak A, et al. Validation of the Joensuu risk criteria for primary resectable gastrointestinal stromal tumour - the impact of tumour rupture on patient outcomes. *Eur J Surg Oncol.* 2011;37(10):890-6.
7. Khoo CY, Chai X, Quek R, Teo MCC, Goh BKP. Systematic review of current prognostication systems for primary gastrointestinal stromal tumours. *Eur J Surg Oncol.* 2018;44(4):388-94.
8. Rutkowski P, Gronchi A, Hohenberger P, et al. Neoadjuvant imatinib in locally advanced gastrointestinal stromal tumors (GIST): the EORTC STBSG experience. *Ann Surg Oncol.* 2013;20(9):2937-43.

9. Wozniak A, Rutkowski P, Schoffski P, et al. Tumor genotype is an independent prognostic factor in primary gastrointestinal stromal tumors of gastric origin: a European multicenter analysis based on ConticaGIST. *Clin Cancer Res.* 2014;20(23):6105-16.
10. Joensuu H, Vehtari A, Riihimäki J, et al. Risk of recurrence of gastrointestinal stromal tumour after surgery: an analysis of pooled population-based cohorts. *Lancet Oncol.* 2012;13(3):265-74.
11. Florindez J, Trent J. Low Frequency of Mutation Testing in the United States: An Analysis of 3866 GIST Patients. *Am J Clin Oncol.* 2020;43(4):270-8.
12. Emory TS, Sobin LH, Lukes L, Lee DH, O'Leary TJ. Prognosis of gastrointestinal smooth-muscle (stromal) tumors: dependence on anatomic site. *Am J Surg Pathol.* 1999;23(1):82-7.
13. Marrelli D, Polom K, de Manzoni G, Morgagni P, Baiocchi GL, Roviello F. Multimodal treatment of gastric cancer in the west: Where are we going? *World J Gastroenterol.* 2015;21(26):7954-69.
14. Ghidini M, Petrelli F, Tomasello G. Right Versus Left Colon Cancer: Resectable and Metastatic Disease. *Curr Treat Options Oncol.* 2018;19(6):31.
15. Fletcher CDM, Bridge JA, Hagendoorn PCW, Mertens F (eds). WHO Classification of Tumours of Soft Tissue and Bone. WHO/IARC: Classifications of tumors, 4th Edition. IARC: Lyon, 2013.
16. Hølmekjær T, Hompland I, Bjerkehaugen B, et al. Recurrence-Free Survival After Resection of Gastric Gastrointestinal Stromal Tumors Classified According to a Strict Definition of Tumor Rupture: A Population-Based Study. *Ann Surg Oncol.* 2018;25(5):1133-9.

17. Boye K, Berner J-M, Hompland I, et al. Genotype and risk of tumour rupture in gastrointestinal stromal tumor. *Br J Surg*. 2018;105(2):e169-e175.
18. Miettinen M, Sobin LH, Lasota J. Gastrointestinal stromal tumors of the stomach: a clinicopathologic, immunohistochemical, and molecular genetic study of 1765 cases with long-term follow-up. *Am J Surg Pathol*. 2005;29(1):52-68.

Figure legends

Fig. 1. **a** The partition of the stomach into an upper, middle and lower third; **b** macroscopic growth pattern: luminal, exophytic and transmural tumors.

Fig. 2. Relationships between tumor subsite, growth pattern, tumor size, mitotic index and tumor genotype in gastric GIST. GIST, gastrointestinal stromal tumor; MI, mitotic index; HPF, high-power fields of the microscope; *PDGFRA*, platelet-derived growth factor receptor- α .

Fig. 3. Estimated recurrence-free survival (Kaplan-Meier) after resection of gastric GIST by **a** gastric subsite ($P = 0.020$); **b** macroscopic growth pattern ($P < 0.001$); **c** tumor genotype ($P < 0.001$). GIST, gastrointestinal stromal tumor; *PDGFRA*, platelet-derived growth factor receptor- α .

Table 1. Clinical, pathological and biological characteristics	
Total number of patients	295
Sex (males:females)	149:146 (50.5:49.5)
Age at surgery, years ^a	66 (14-93)
Syndromic GIST	1 (0.3)
Presentation	
Symptomatic	193 (65.4)
Bleeding/anemia	99 (33.6)
Pain/discomfort	57 (19.3)
Other	37 (12.5)
Incidental finding	102 (34.6)
At imaging/endoscopy	90 (30.5)
At surgery	12 (4.1)
R1 resection	28 (9.5)
Multivisceral resection	41 (13.9)
Tumor rupture	24 (8.1)
Gastric subsite	
Upper third	122 (41.8)
Middle third	120 (41.1)
Lower third	50 (17.1)
Borderline or unspecified	3
Macroscopic growth pattern	
Luminal	100 (34.1)
Exophytic	94 (32.1)
Transmural	99 (33.8)
Unspecified	2
Tumor size, cm ^a	4.8 (0.5-34.0)
≤5	161 (54.6)
5.1-10.0	83 (28.1)
>10	51 (17.3)
Mitotic index, per 50 HPF ^a	2 (0-166)
0-5	221 (77.9)
6-10	25 (8.8)
>10	38 (13.4)
Unspecified	11
Modified NIH consensus criteria	
Very low risk	23 (7.8)
Low risk	120 (40.8)
Intermediate risk	74 (25.2)
High risk	77 (26.2)
Unspecified	1
Mutation analysis ^b	256 (86.8)
<i>KIT</i> exon 11	177 (69.1)
Substitution	63 (24.6)
Duplication/insertion	28 (10.9)
Deletion/insertion-deletion	86 (33.6)
Del557/558	47 (18.4)
Not del557/558	39 (15.2)
<i>KIT</i> exon 17	6 (2.3)
<i>PDGFRA</i> exon 12	9 (3.5)

Table 1 continued	
<i>PDGFRA</i> exon 14	3 (1·2)
<i>PDGFRA</i> exon 18	50 (19·5)
D842V	40 (15·6)
Other	10 (3·9)
No <i>PDGFRA</i> or <i>KIT</i> mutation detected	11 (4·3)
SDHB expression positive	6
SDHB expression negative	2
Analysis not performed	3
Adjuvant imatinib	47 (15·9)
Duration, months ^a	24 (1-60)
Neoadjuvant imatinib	11 (3·7)
Duration, months ^a	8 (1-14)

Values are numbers (%) unless indicated otherwise. ^aValues are median (range). ^b23 patients in whom mutation analysis was not performed and 16 patients in whom tumour tissue was unfit for analysis were excluded. GIST, gastrointestinal stromal tumor; HPF, high-power field of the microscope; NIH, National Institutes of Health; *PDGFRA*, platelet-derived growth factor receptor- α ; SDHB, succinate dehydrogenase complex subunit B

Table 2. Associations between clinical, anatomical and molecular characteristics

	Gastric subsite ^a				Growth pattern ^b				Genotype ^c			
	Upper	Middle	Lower	P value	Luminal	Exophytic	Transmural	P value	Del557/558	KIT not del557/558	PDGFRA	P value
Sex				0.089 ^f				0.698				0.010
Male	70 (47.3)	52 (35.1)	26 (17.6)		49 (33.3)	45 (30.6)	53 (36.1)		31 (24.0)	60 (46.5)	38 (29.5)	
Female	52 (36.1)	68 (47.2)	24 (16.7)		51 (35.0)	49 (33.6)	46 (31.5)		16 (13.8)	76 (65.5)	24 (20.7)	
Tumor size				0.214				<0.001				<0.001
≤10 cm	103 (42.4)	95 (39.1)	45 (18.5)		99 (40.7)	71 (29.2)	73 (30.0)		28 (14.0)	123 (61.5)	49 (24.5)	
>10 cm	19 (38.8)	25 (51.0)	5 (10.2)		1 (2.0)	23 (46.0)	26 (52.0)		19 (42.2)	13 (28.9)	13 (28.9)	
Mitotic index				0.006				<0.001				<0.001
0-5	82 (37.4)	90 (41.1)	47 (21.5)		91 (41.6)	75 (34.2)	53 (24.2)		13 (7.4)	105 (60.0)	57 (32.6)	
6-10	11 (44.0)	12 (48.0)	2 (8.0)		7 (28.0)	9 (36.0)	9 (36.0)		2 (9.1)	19 (86.4)	1 (4.5)	
>10	24 (64.9)	12 (32.4)	1 (2.7)		2 (5.3)	6 (15.8)	30 (78.9)		24 (64.9)	10 (27.0)	3 (8.1)	
Tumor rupture				0.092 ^g				0.001				0.001
No	111 (41.1)	109 (40.3)	50 (18.5)		99 (36.7)	87 (32.2)	84 (31.1)		36 (16.1)	129 (57.8)	58 (26.0)	
Yes	11 (50.0)	11 (50.0)	0		1 (4.3)	7 (30.4)	15 (65.2)		11 (50.0)	7 (31.8)	4 (18.2)	
Risk category ^{de}				0.029				<0.001				<0.001
Non-high-risk	83 (38.2)	90 (41.5)	44 (20.3)		94 (43.5)	68 (31.5)	54 (25.0)		14 (8.1)	112 (64.7)	47 (27.2)	
High-risk	38 (51.3)	30 (40.5)	6 (8.1)		6 (7.9)	26 (34.2)	44 (57.9)		32 (45.1)	24 (33.8)	15 (21.1)	
Gastric subsite ^a								0.018				<0.001
Upper	-	-	-		40 (32.8)	32 (26.2)	50 (41.0)		33 (30.0)	70 (63.6)	7 (6.4)	
Middle	-	-	-		35 (29.4)	45 (37.8)	39 (32.8)		13 (13.5)	50 (52.1)	33 (34.4)	
Lower	-	-	-		24	17 (34.0)	9 (18.0)		1 (2.6)	15 (39.4)	22 (57.9)

Fig. 1. a The partition of the stomach into an upper, middle and lower third; **b** macroscopic growth pattern: luminal, exophytic and transmural tumors.

Fig. 2. Relationships between tumor subsite, growth pattern, tumor size, mitotic index and tumor genotype in gastric GLST. GLST, gastrointestinal stromal tumor; MI, mitotic index; HPF, high-power fields of the microscope; PDGFRA, platelet-derived growth factor receptor- α .

Fig. 3. Estimated recurrence-free survival (Kaplan-Meier) after resection of gastric GIST by a gastric subsite ($P = 0.020$); **b** macroscopic growth pattern ($P < 0.001$); **c** tumor genotype ($P < 0.001$). GIST, gastrointestinal stromal tumor; PDGFRA, platelet-derived growth factor receptor- α .

Supplementary Table S1. Multivariable Cox regression analysis of recurrence-free survival

	HR	95% CI	<i>P</i> value
Sex			0.922
Female	1.00 (reference)		
Male	1.05	0.40-2.75	
Tumor size			0.278
≤ 10 cm	1.00 (reference)		
> 10 cm	0.40	0.08-2.10	
Mitotic index			< 0.001
≤ 10 per 50 HPF	1.00 (reference)		
> 10 per 50 HPF	22.23	5.06-97.60	
Tumor rupture			0.014
No	1.00 (reference)		
Yes	7.53	1.50-37.76	
Tumor growth pattern			0.120
Luminal or exophytic	1.00 (reference)		
Transmural	2.90	0.76-11.06	
Tumor genotype			0.349
<i>PDGFRA</i>	1.00 (reference)		
<i>KIT</i> not del557/558	0.41	0.09-1.82	
<i>KIT</i> del557/558	0.30	0.06-1.54	

HR, hazard ratio; CI, confidence interval; HPF, high-power field of the microscope; *PDGFRA*, platelet-derived growth factor receptor- α .

Supplementary Fig. S1. Boxplots showing tumor diameter (**A-C**) and mitotic index (**D-F**) based on tumor location (**A, D**), tumor growth pattern (**B, E**) and tumor genotype (**C, F**). Boxes indicate the median, the 25th and 75th percentile, and whiskers represent maximum and minimum values. Outliers are censored.

ERRATA

Changes made after submission and approved by the committee.

PAGE	LINE	TEXT	SUBMITTED VERSION	CORRECTED VERSION
15	19	Body	...cases per one million...	...cases per 1 million...
15	21	Body	...less than one cm, ...	...less than 1 cm, ...
17	16	Body	...less than five per cent...	...less than 5 per cent...
33	24	Body	...was size independently...	..., size was independently...
34	2	Body	...RFS, only MI was.	...RFS, MI was the only independent factor.
38	last	Body	...soft tissue sarcoma of the extremities...	...soft tissue sarcomas of the extremities...
39	26	Body	...tissue sarcoma of the extremities...	...tissue sarcomas of the extremities...
52	Figure 6, line 3	Legend	...> 5 per 50 HPFs,	...> 5 per 50 HPF,
53	Figure 7, line 2	Legend	...(P < 0.001),...	...(P < 0.001),...
62	11	Body	...pooled, allegedly population-based...	...pooled, population-based...