DETRIMENTAL SYNERGY; DECIPHERING THE ROLES OF AQUAPORINS IN PARKINSON'S DISEASE

By Agnete Prydz MD

Thesis for the degree of Philosophiae Doctor (PhD)

Division of Anatomy

Department of Molecular Medicine

Institute of Basic Medical Sciences

University of Oslo

2021

Principal supervisor:

Prof. Mahmood Amiry-Moghaddam MD PhD

Co-supervisor:

Dr. Katja Stahl PhD

© Agnete Prydz, 2021

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-927-1

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

Jeg holder ditt hode

*Jeg holder ditt hode
i mine hender, som du holder
mitt hjerte i din ømhet
slik allting holder og blir
holdt av noe annet enn seg selv
Slik havet løfter en sten
til sine strender, slik treet
holder høstens modne frukter, slik
kloden løftes gjennom kloders rom
Slik holdes vi begge av noe og løftes
dit gåte holder gåte i sin hånd*

Stein Mehren (1935-2017)
Norwegian poet,
suffered from Parkinson's disease

ACKNOWLEDGEMENTS

This work was carried out at the Laboratory of Molecular Neuroscience, Division of Anatomy, Department of Molecular Medicine, Institute of Basic Medical Sciences at the University of Oslo. This research would not have been possible without the financial support from the Olav Thon Foundation.

As a curious, and perhaps overly ambitious, first-year medical student, I came across a soon-to-be Professor who, with his enthusiastic and witty personality, encouraged me to explore the world of neuroscience. He – Mahmood Amiry-Moghaddam – came to be my main supervisor throughout the Medical Student Research Program – a decision I am sure he has regretted several times, when finding out that his new student was not about to miss out on anything medical school had to offer, and ended up working somewhat unconventionally at evenings/nights/weekends/holidays in Oslo, Bergen, South Africa, and Finnmark. Even though our common enemy of ‘available time’ has been working against us for over eight years, we have managed to produce some great papers, of which I am truly proud. This would not have been possible without his ability to constantly challenge and motivate me, his vast scientific knowledge, and inspiring persona, qualities for which I am sincerely grateful.

When first introduced to the research group, I met Katja Stahl – an exceptionally bright, hardworking, and cool person whom I am now delighted to call my friend. Katja became my outstanding co-supervisor. I am deeply grateful for her availability regardless of the hour of the day, thoroughness, and deep investment in my research. I would also like to extend a special thanks to the hard work and fruitful collaboration with Soulmaz Zahl.

I owe many thanks to the great technical staff in our group; Bjørg Riber, Karen Marie Gujord, Jorunn Knutsen, Johannes Helm, and Bashir Hakim for technical assistance and kindness. I thank Trygve Leergaard for assistance with the stereology program and Finn-Mogens Haug for assistance with the Analysis program. I particularly thank Ole Petter Ottersen for his invaluable help throughout my work. Special thanks to Maja Puchades and Vidar Gundersen for their enjoyable collaboration. Also, many thanks to Carina Knutsen and Gunnar Lothe for help with the graphical work.

Special thanks go to my colleagues Nadia Skauli, Laura Camassa, Eystein Hoddevik, Gry Helen Enger Syverstad, Shreyas Rao, and Shirin Katoozi for creating a fun and stimulating working environment and rewarding scientific discussions. I thank all my good friends and especially Lara Pasovic, Rishab Chawla, Gry Fluge Vindedal, and Nina Obad for shared frustrations, endless support, and numerous laughs along the way. I am also very grateful to Paul who fought his way through this thesis in order to give valuable feedback on the English language.

I was fortunate to meet Mahmood and Magnus at about the same time. As Mahmood has lifted me off the ground with inspiring and ambitious ideas, my husband Magnus has grounded me, constantly reminding me that there are more important things in life than medical research. His endless love and support, encouragement, and inexhaustible reserves of patience have kept me going on this rocky road, for which I am forever grateful.

Dedicated to my father, whom I never truly got to know due to the cruel ways of brain disease.

Agnete Prydz

Oslo, December 2020

TABLE OF CONTENTS

ABSTRACT	7
SAMMENDRAG	9
ABBREVIATIONS	11
PAPERS INCLUDED IN THE THESIS	13
THE CANDIDATE'S SCIENTIFIC CONTRIBUTION	14
INTRODUCTION	15
PARKINSON'S DISEASE	15
Epidemiology	15
Clinical features and diagnosis	15
Basal ganglia circuits	16
Etiology	18
Pathology	19
Treatment	23
BRAIN MICROENVIRONMENT IN PARKINSON'S DISEASE	24
Roles of glial cells in normal and PD brain	25
Roles of brain aquaporins in normal and PD brain	31
HYPOTHESES AND AIMS OF THE THESIS	37
SUMMARY OF RESULTS	38
Paper I: Subcellular expression of aquaporin-4 in substantia nigra of normal and MPTP-treated mice.	38
Paper II: Pro-inflammatory role of AQP4 in mice subjected to intrastriatal injections of the parkinsonogenic toxin MPP+	39
Paper III: Targeted deletion of the aquaglyceroporin AQP9 is protective in a mouse model of Parkinson's disease.	40
METHODOLOGICAL CONSIDERATIONS	41
TRANSGENIC ANIMAL MODELS	41
MODELING PARKINSON'S DISEASE	43
Animal models of PD	43
IMMUNOCYTOCHEMISTRY	48

Immunocytochemistry for light microscopy	48
Immunocytochemistry for electron microscopy	48
Antibody specificity and sensitivity	50
QUANTITATIVE REAL TIME PCR	51
STEREOLOGICAL CELL QUANTITATION	51
DISCUSSION	54
PHYSIOLOGICAL AND PATHOPHYSIOLOGICAL IMPLICATIONS	54
AQP4 – MICROENVIRONMENTAL LIABILITY	55
AQP4 – TOXIN EXACERBATOR	57
AQP9 – TOXIN FACILITATOR	62
IMPLICATIONS FOR PARKINSON’S DISEASE	65
CONCLUSION AND FUTURE PERSPECTIVES	67
REFERENCES	68
PAPER I-III	86

ABSTRACT

Parkinson's disease (PD) is a neurodegenerative disorder characterized by progressive selective loss of dopaminergic neurons in substantia nigra pars compacta (SNpc) in the midbrain [1]. The etiology of PD is to date unknown, though several studies have identified environmental toxins as potential key factors in the pathogenesis [2, 3]. However, the mechanism by which these toxins function, and other underlying factors contributing to the selective vulnerability of these dopaminergic neurons, remains elusive. Hence, the microenvironment surrounding the dopaminergic neurons in SNpc has been greatly overlooked, as the available research has largely focused on the neurons. Yet, recent studies point out neuroinflammation as a pathological hallmark of PD [4].

Three aquaporins (AQPs) have been proven expressed in the mammalian brain [5], where two of these protein channels – aquaporin 4 (AQP4) and aquaporin 9 (AQP9) – are expressed in a variety of brain regions, including the mesencephalon [6]. AQP4 is the predominant AQP in the central nervous system (CNS), expressed in astrocytes throughout all brain regions, mainly localized in a polarized manner in the perivascular astrocytic endfeet [7]. AQP9 is less generally expressed with a dominance in the mesencephalon [8], both in astrocytes and dopaminergic neurons [9]. The architectural configuration of AQP4 makes the channel selective for water transportation, unlike AQP9, which has proven permeable to a diversity of solutes including the parkinsonogenic compound arsenite [10]. AQP4 plays a critical role in water homeostasis/dyshomeostasis [5] and has recently been coupled to the pathogenesis of several neurodegenerative diseases such as Alzheimer's disease (AD) [11], amyotrophic lateral sclerosis (ALS) [12], and Huntington's disease (HD) [13], as well as neuroinflammation [14-16]. Furthermore, MRI studies have shown increased accumulation of water in SNpc of patients with PD [17].

To investigate microenvironmental factors contributing to the selective death of dopaminergic neurons in PD, this thesis evaluates the potential pathophysiological role of AQP4 and AQP9 in PD, specifically focusing on glial aspects of neuroinflammation and toxic contribution.

In order to illuminate the potential role of AQP4 in PD, we investigated the expression pattern of AQP4 in SNpc compared to the cerebral cortex under physiological conditions, uncovering significantly higher expression of AQP4 both in the perivascular astrocytic membrane, as well

as in the astrocytic processes found in the neuropil. This high expression of AQP4 under normal conditions was further exacerbated in a sub-acute mouse model of PD, where the parkinsonogenic toxin MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) was subcutaneously (sc.) injected. This raised the question of whether SN is more prone to toxic exposure and neuroinflammation due to the considerable physiological astrocytic expression of AQP4. In order to further understand the pathophysiological – and potentially neuroinflammatory – role of AQP4 in PD, we used an acute mouse model through intrastriatal injections of MPP⁺ (1-methyl-4-phenylpyridinium), the active toxic compound of MPTP, to investigate the consequence of targeted deletion of *Aqp4* on dopaminergic cell death and microglial activation. We show that intrastriatal injections of MPP⁺ resulted in increased expression of AQP4 in SNpc as well as astrocytic endfoot swelling. We further demonstrate that treatment with MPP⁺ led to considerable upregulation of genes known to activate microglia in SNpc in wild-type (WT) mice, however not in *Aqp4*^{-/-} mice. WT mice also showed a significantly higher degree of dopaminergic cell death. Thus, our findings are consistent with the hypothesis that AQP4 plays a pro-inflammatory role in PD, plausibly due to astrocytic dyshomeostasis.

To investigate the theory that AQP9 functions as a gateway channel for parkinsonogenic toxins, we established AQP9 permeability to MPP⁺ by using *Xenopus* oocytes expressing AQP9. Furthermore, targeted deletion of *Aqp9* proved protective of dopaminergic cell death in both an *in vitro* organotypic cell culture model and in an *in vivo* acute MPP⁺ mouse model. These results show that AQP9 contributes to the selective vulnerability of dopaminergic neurons in PD by enabling transportation of endogenous toxin into the cells.

In summary, this thesis has pinpointed AQP4 and AQP9 as detrimental factors in the pathogenesis of PD, where AQP9 functions as a toxin facilitator enabling access to dopaminergic neurons, whereas AQP4 plays the role of toxin exacerbator, amplifying the toxic effect in SNpc through neuroinflammation. This novel insight into the pathogenesis of PD may help to pave the way for creating new therapeutic targets in PD.

SAMMENDRAG

Parkinson's sykdom (PD) er en nevrodegenerativ sykdom som kjennetegnes av et progredierende tap av dopaminerge nevroner i substantia nigra pars compacta (SNpc) i mesencephalon i hjernen [1]. Etiologien til PD er hittil ukjent, men flere studier har siste tiden identifisert giftstoffer fra miljøet som en potensiell nøkkelfaktor i patogenesen [2, 3]. Virkningsmekanismene til disse toksinene, samt andre underliggende årsaker som potensielt kan bidra til den selektive sårbarheten til disse dopaminerge nevronene er foreløpig uavklart. I den sammenheng har mikromiljøet som omgir de dopaminerge nevronene i SNpc fått påfallende lite oppmerksomhet der tilgjengelig litteratur i stor grad har fokusert på selve nevronene. Dog er det nå generell konsensus om at nevroinflammasjon er et patologisk kjennetegn i PD [4], hvilket gjør mikromiljøet i SNpc ytterligere interessant.

Tre typer aquaporiner (AQP) uttrykkes i pattedyrhjernen [5], hvorav to av disse proteinkanalene – aquaporin 4 (AQP4) og aquaporin 9 (AQP9) – uttrykkes i et bredt spekter av regioner i hjernen, inkludert i mesencephalon [6]. AQP4 er den vanligste AQPen i sentralnervesystemet (CNS) og uttrykkes på en polarisert måte hovedsakelig i astrocyttenes perivaskulære endeføtter gjennom hele CNS [7]. Sammenliknet med AQP4, er ekspresjonen av AQP9 mindre generalisert, der kanalen i størst grad finnes i mesencephalon [8], både i astrocytter og i dopaminerge nevroner [9]. Den arkitektoniske utformingen til AQP4 gjør at kanalen kun er permeabel for vann, dette i motsetning til AQP9 som har vist seg å være permeabel for et bredt spekter av stoffer, inkludert arsenitt som har blitt assosiert med PD [10]. AQP4 spiller en sentral rolle i vannhomestase/dyshomeostase [5] og har nylig blitt koplet til flere nevrodegenerative lidelser som for eksempel Alzheimers sykdom (AD) [11], amyotrofisk lateral sklerose (ALS) [12] og Huntingtons sykdom (HD) [13], i tillegg til nevroinflammasjon [14-16]. Videre har MR-studier vist økt akkumulering av fritt vann i SNpc hos pasienter med PD [17].

Målet med denne doktorgradsavhandlingen er å undersøke faktorer i mikromiljøet i SNpc som kan bidra til den selektive celledøden av dopaminerge nevroner i PD. Avhandlingen evaluerer de potensielle patofysiologiske rollene til AQP4 og AQP9 i PD med et spesielt fokus på gliacelleaspekter ved nevroinflammasjon og toksisitet i sykdomforløpet.

For å belyse de potensielle rollene til AQP4 i PD, har vi undersøkt ekspresjonsmønsteret til AQP4 i SNpc sammenliknet med cerebral cortex under fysiologiske forhold. Vi avdekket

signifikant høyere ekspresjon av AQP4 i både perivaskulære astrocyttmembraner og i astrocyttprosesser i nevrofil. Denne høye ekspresjonen av AQP4 økte ytterligere i en subakutt musemodell av PD, hvor det parkinsonassosierte toksinet MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) ble injisert subkuttant i musene. På grunnlag av dette stilte vi spørsmål ved hvorvidt SN er mer utsatt for toksisk eksponering og nevroinflammasjon gitt den fysiologisk betydelige høye ekspresjonen av AQP4 i astrocytter i denne regionen. For å bedre kunne forstå den patofysiologiske – og potensielt nevroinflammatoriske – rollen til AQP4 i PD, benyttet vi en akutt musemodell av PD der musene gjennomgikk injeksjoner direkte i striatum av MPP⁺ (1-methyl-4-phenylpyridinium), hvilket er den aktive, toksiske komponenten av MPTP. Forsøkene ble gjennomført på WT og *Aqp4*^{-/-} mus for å belyse hvilke konsekvenser knock-out av *Aqp4* har på dopaminerg celledød og mikrogliaaktivering. Vi viser at MPP⁺ injeksjoner i striatum resulterer i økt ekspresjon av AQP4 i SNpc, samt økt hevelse i astrocyttendeføtter. Videre viser vi at behandling med MPP⁺ fører til betydelig oppregulering av gener som er kjent for å aktivere mikroglia i SNpc i WT-mus, hvilket vi ikke i samme grad fant i *Aqp4*^{-/-} mus. I tillegg fant vi at WT-musene gjennomgikk signifikant høyere grad av dopaminerg celledød. Følgelig er disse resultatene samsvarende med hypotesen om at AQP4 spiller en pro-inflammatorisk rolle i PD, sannsynligvis grunnet dyshomeostase i astrocyttene.

For å undersøke hypotesen om at AQP9 kan fungere som en inngangskanal for parkinsonassosierte toksiner, brukte vi *Xenopus* oocytter som uttrykket AQP9 for å vise at AQP9 er permeabel for MPP⁺. Deretter viste vi at knock-out av *Aqp9* virket beskyttende mot dopaminerg celledød i både en *in vitro* organotypisk cellekulturmodell og i en *in vivo* MPP⁺ akutt musemodell. Disse resultatene viser at AQP9 bidrar til den selektive sårbarheten til dopaminerge nevroner i PD ved å fungere som en inngangskanal for endogene toksiner inn i cellene.

Vi konkluderer med at denne doktorgradsavhandlingen har pekt på AQP4 og AQP9 som skadelige faktorer i patogenesen i PD, hvor AQP9 fungerer som en tilrettelegger for toksiner ved å etablere tilgang inn i de dopaminerge nevronene, mens AQP4 fungerer som en toksinforsterker ved å intensivere de toksiske effektene i SNpc via nevroinflammasjon. Denne avhandlingen gir følgelig ny innsikt i patogenesen i PD og vil således kunne bane vei for nye terapeutiske muligheter for denne nevrodegenerative sykdommen.

ABBREVIATIONS

AD	Alzheimer's disease
ALS	Amyotrophic lateral sclerosis
AQP	Aquaporins
AQP4	Aquaporin-4
AQP9	Aquaporin-9
BBB	Blood brain barrier
CNS	Central nervous system
CSF	Cerebrospinal fluid
Cx43	Connexin 43
DA	Dopamine
DAB	3,3'-Diaminobenzidin
DAPC	Dystrophin-associated protein complex
DAT	Dopamine transporter
DBS	Deep brain stimulation
DJ-1	Protein deglycase
DM	Diabetes mellitus
ECM	Extracellular matrix
En1	Engrailed 1
ETC	Electron transport chain
GABA	Gamma-aminobutyric acid
GDNF	Glial-derived neurotrophic factor
GFAP	Glial fibrillary acidic protein
GLAST	Glutamate aspartate transporter
GNR	Glia/neuron –ratio
GSH	Glutathione
GPe	Globus pallidus external
GPe	Globus pallidus external
GPe	Globus pallidus internal
HD	Huntington's disease
HEK	Human embryonic kidney
HVA	Homovanillic acid
ICAM-1	Intracellular adhesion molecule 1
IL-1 β	Interleukin-1 beta
IL-6	Interleukin-6
IL-12	Interleukin-12
INF- γ	Interferon- γ
IP ₃	Inositol triphosphate
K-ATP	Adeno triphosphate sensitive potassium channel
Kir4.1	Inwardly rectifying potassium channel 4.1
KI	Knock in
KO	Knock out
LB	Lewy bodies
L-DOPA	Levodopa
LPS	Lipopolysaccharide
LPC	Lysophosphatidylcholine
LRRK2	Leucine-rich repeat kinase 2
MAO-B	Monamine oxidase B
MPTP	1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine
MPP+	1-methyl-4-phenylpyridinium

MRI	Magnetic resonance imaging
MS	Multiple sclerosis
MTLE	Mesial temporal lobe epilepsy
NCAM	Neural cell adhesion molecule
NET	Norepinephrine transporter
NF- κ B	Nuclear factor kappa light chain enhancer of activated B cells
NMDA	N-methyl-D-aspartate
NMO	Neuromyelitis optica
NPH	Normal pressure hydrocephalus
NO	Nitric oxide
PD	Parkinson's disease
ROS	Reactive oxygen species
ROI	Region of interest
RT-qPCR	Real time quantitative reverse transcription
RRF	Retrorubral field
SMA	Prefrontal supplementary motor area
sc	Subcutaneous
siAQP4	Small interference RNA against AQP4
SN	Substantia nigra
SNpc	Substantia nigra pars compacta
SNpr	Substantia nigra pars reticulata
SPECT	Single photon emission computerized tomography
SE	Spongiform encephalopathy
SMA	Supplementary motor area
SOCS1	Suppressor of cytokine signaling 1
SRS	Systematically random sampled sections
STN	Subthalamic nucleus
TBI	Traumatic brain injury
TH	Thyrosine hydroxylase
TLR	Toll like receptor
TNF- α	Tumor necrosis factor alpha
UPDRS	Unified Parkinson's Disease Rating Scale
UKPDSBB	UK Parkinson's Disease Society Brain Bank
VTA	Ventral tegmental area
WT	Wild type
6-OHDA	6-hydroxydopamine

PAPERS INCLUDED IN THE THESIS

Paper I

Subcellular expression of aquaporin-4 in substantia nigra of normal and MPTP-treated mice

Prydz A, Stahl K, Puchades M, Davarpaneh N, Nadeem M, Ottersen OP, Gundersen V, Amiry-Moghaddam M. *Neuroscience*. 2017; Sep 17;359:258-266. doi: 10.1016/j.neuroscience.2017.07.029.

Paper II

Pro-inflammatory role of AQP4 in mice subjected to intrastriatal injections of the parkinsonogenic toxin MPP+

Prydz A*, Stahl K*, Zahl S, Skauli N, Skare Ø, Ottersen OP, Amiry-Moghaddam M., *Cells*. 2020 Nov 5;9(11):E2418. doi: 10.3390/cells9112418.

* Both authors have contributed equally.

Paper III

Targeted deletion of the aquaglyceroporin AQP9 is protective in a mouse model of Parkinson's disease

Stahl K, Rahmani S, **Prydz A**, Skauli N, MacAulay N, Mylonakou MN, Torp R, Skare Ø, Berg T, Leergaard TB, Paulsen RE, Ottersen OP, Amiry-Moghaddam M. *PLoS One* 2018. Mar 22;12(3):e0194896. doi: 10.1371/journal.pone.0194896

THE CANDIDATE'S SCIENTIFIC CONTRIBUTION

Paper I: Subcellular expression of aquaporin-4 in substantia nigra of normal and MPTP-treated mice

- Planning and performing immunocytochemistry experiments of MPTP-treated animals
- Planning and performing immunogold labelling of MPTP-treated animals
- Confocal microscopy; image acquisition and analyses of MPTP-treated animals
- Electron microscopy; image acquisition, analyses, and quantitation of MPTP-treated animals
- Statistical analyses of immunogold quantitation of MPTP-treated animals
- Preparation of figures 3, 4 and 5
- Writing the original draft, editing and submitting the manuscript

Paper II: Pro-inflammatory role of AQP4 in mice subjected to intrastriatal injections of the parkinsonogenic toxin MPP+

- Animal care pre- and postoperatively
- Post-operative behavioral/clinical assessment of animals
- Dissection and tissue preparation for immunocytochemistry
- Planning and performing immunocytochemistry experiments
- Planning and performing immunogold labelling
- Electron microscopy; image acquisition, analyses, and quantitation
- Stereological quantification
- Statistical analysis of immunogold and stereological quantification
- Preparation of figures 1-5
- Writing the original draft and editing the manuscript

Paper III: Targeted deletion of the aquaglyceroporin AQP9 is protective in a mouse model of Parkinson's disease.

- All post-operative animal care
- All post-operative behavioral/clinical assessment of animals
- Performing and analyzing motoric testing of animals (apomorphine and cylinder -test)
- Crucial contribution to stereological quantification
- Writing parts of the manuscript, editing the manuscript

INTRODUCTION

PARKINSON'S DISEASE

Epidemiology

In 1817 James Parkinson discovered a disease which he named “The Shaking Palsy”, today widely known as Parkinson’s disease (PD) [18]. PD is the second most common neurodegenerative disorder, affecting 1.4% of the population with the prevalence increasing with age [20] to an estimation of a global prevalence of 3% in the population exceeding 80 years of age [21]. The Norwegian ParkWest study concluded that the incidence rates of PD in Norway are comparable to incidence rates in Western European and Northern American countries [22]. In these progressively ageing populations, the number of people affected is expected to double from 2005 till 2030 [23], collated with increased years of disability, PD is expected to cause a significant increase in the personal and socioeconomic burdens associated with the disease [23]. Only approximately 20% of patients with PD have a familial history of PD while the exact number of known genetic causes is unknown [24] and up to 95% of PD-cases are considered idiopathic [25], making the question of environmental risk factors highly relevant in the pursuit of unravelling the pathogenesis of PD.

Clinical features and diagnosis

PD is a progressive, neurodegenerative disorder and the leading cause of the clinical syndrome parkinsonism. Parkinsonism is clinically characterized by motor dysfunctions such as bradykinesia, resting tremor, rigidity, flexed posture, freezing and postural instability [26]. The onset of motor symptoms is typically asymmetrical in PD, persisting throughout the disease. Motor disability is most commonly monitored using the Unified Parkinson’s Disease Rating Scale (UPDRS) [27]. In PD, non-motor symptoms are also common, such as depression, lack of motivation, impaired control of impulses, passivity, fatigue, cognitive impairment, orthostatic hypotension, obstipation, nausea, dysphagia, and urinary dysfunction [28][29]. The overall clinical picture of the PD patient regarding motor and non-motor symptoms makes it challenging to diagnose the patients correctly, especially early in the disease. Assessed against the gold standard post-mortem pathological examination, the use of standard clinical indicators such as the UK Parkinson’s Disease Society Brain Bank (UKPDSBB) criteria can improve the accuracy of clinical diagnosis [30]. In the past decade, molecular neuroimaging is becoming increasingly relevant in the diagnostics of PD. For

instance, dopamine-transporter scan/single-photon emission computerized tomography (DAT scan/SPECT) is an important tool in distinguishing atypical parkinsonism from actual PD [31]. It has been shown that the onset of clinical symptoms emerges when the loss of dopaminergic neurons reaches 60-80% [32, 33], making the prodromal phase ideal for pre-symptomatic diagnosis, and thus possibly for pre-symptomatic treatment in the future. Currently, there is no available verified diagnostic test for PD based on cerebrospinal fluid (CSF) or blood biomarkers, even though several studies have addressed the issue, especially regarding levels of α -synuclein species [34]. The need for diagnostic tests and biomarkers to advance early diagnostic testing is thus clearly evident.

Basal ganglia circuits

The extrapyramidal or basal ganglia system (figure 1) contains the substantia nigra pars compacta (SNpc) and pars reticulata (SNpr), dorsal striatum including caudate and putamen, globus pallidus, the subthalamic nucleus (STN), and thalamus. The excitatory cortical input to the basal ganglia is mediated by the neurotransmitter glutamate, and is derived from the prefrontal supplementary motor area (SMA), as well as the amygdala and hippocampus [35].

The dopaminergic neurons of the ventral mesencephalon comprise over 90% of the dopaminergic cell bodies in the brain, where the catecholamine neurotransmitter dopamine (DA) is produced in SN, ventral tegmental area (VTA), and retrorubral field (RRF) [36]. The neurons in SNpc provide major dopaminergic excitatory and inhibitory modulatory input to the striatum and its output system, which is mediated by the inhibitory neurotransmitter gamma-aminobutyric acid (GABA) [35]. Five different DA receptors have been characterized (D1-D5), where the D1 and D2 receptors being the most relevant in the PD pathophysiology. These receptors are highly concentrated in the dorsal striatum (motor striatum) and are activated through the dopaminergic pathway from the SNpc [37]. There are two nigrostriatal output pathways, subdivided into the direct and the indirect pathway (figure 1). The direct pathway excites the putamen through dopaminergic excitatory mediation on D1 DA receptors, which further projects directly to the globus pallidus interna (GPi). The indirect pathway on the other hand, is mediated through dopaminergic inhibition on D2 DA receptors. This pathway projects from the putamen to the globus pallidus external (GPe), which in turn projects to the STN. These projections are GABAergic. The last projection of the indirect pathway from the STN to the GPi provides excitatory glutamatergic input. Finally, the

neurons of the GPi are GABAergic synapsing in the thalamus in turn provide excitatory thalamic input to the cortex. To summarize, the release of DA thus leads to inhibition of the GPi through both the direct and indirect pathways, resulting in reduced inhibition of the thalamus and thus increased stimulation of the motor cortex [35-37].. In PD, the selective loss of dopaminergic neurons in SNpc leads to DA depletion in the SN and thus in the nigrostriatal pathway. The consequence is a relative overactivity of the indirect pathway and a decreased inhibition of the direct pathway leading to increased output from GPi, ultimately resulting in increased inhibition of the thalamus and reduced excitatory input to the motor cortex, and leading to the cardinal symptoms of PD brady/hypokinesia [38].

Figure 1: Basal ganglia dysfunction in PD.

D1, dopamine receptor 1; D2, dopamine receptor 2; GPe, globus pallidus external; GPi, globus pallidus internal; SNc, substantia nigra pars compacta; SNr, substantia nigra pars reticulata; STN, nucleus subthalamicus; Thal, thalamus.

Adapted from Watts RL *et al.*, 2011 [39].

Etiology

The etiology of PD is multifactorial with both genetic and environmental factors playing central roles. The biggest risk factor is age [21]. Additionally, the prevalence of PD is unevenly distributed among genders and geographical areas, with a higher prevalence in men [40] as well as in Western countries, and especially rural areas [41].

Genetics

PD is generally considered an idiopathic disorder, where approximately 15-20% of cases report a family history, and about 5% are deemed to be caused by Mendelian inheritance [42]. The genes associated with PD or parkinsonism are called PARK and to date, 23 PARK genes have been linked to PD [42] where mutations in these genes are inherited through autosomal dominant, recessive inheritance, and possibly x-linked [43]. The numerically most important mutations are those found in the genes coding for parkin, leucine-rich repeat kinase 2 (LRRK2), α -synuclein, and protein deglycase (DJ-1) [42]. These mutations are associated with protein aggregation, mitochondrial dysfunction, oxidative stress, and reduced protein clearance and give rise to parkinsonism indistinguishable from idiopathic PD [44].

Environmental risk factors

The suspicion of environmental toxins as risk factors for developing PD was fully manifested in 1983 when people developed irreversible parkinsonism identical to the idiopathic PD after injecting a drug contaminated with 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP), a complex-I inhibitor of the electron transport chain (ECT) [45, 46] (figure 2). The drug-induced selective death of dopaminergic neurons in SN and the symptoms and pathological findings were reproduced in primate [47] and rodent models of PD [48, 49]. Studies on the herbicide paraquat [50] and the pesticide rotenone [51] also show selective complex-I inhibition in dopaminergic neurons in animal models. Since the discovery of MPTP, several studies have supported a link between exposure to environmental neurotoxins as a substantial risk factor for developing PD, including a case-control study demonstrating a correlation between pesticide exposure and late-onset PD [52], as well as consumption of well water containing arsenite [53]. However, this hypothesis has been considered controversial and is disputed as some studies failed to find a correlation [54-56]. Of note, these studies have, among others, been criticized for relying solely on self-reports, rather than focusing on more objective semi-quantitative methods when assessing exposure. Recent large meta-analyses of

epidemiological studies have demonstrated a clear link between exposure to environmental neurotoxins and PD [2, 3, 57].

Interestingly, a large meta-analysis has shown that both cigarette smoking and coffee drinking are inversely correlated with PD [58], though the causative role remains as yet unclear.

Figure 2: MPTP metabolism and effect.

Illustration showing that MPTP crosses the BBB and is further converted to MPP⁺ by monamine-oxidase-B (MAO-B) in non-dopaminergic cells such as astroglia. MPP⁺ enters the dopaminergic neurons through the dopamine transporter (DAT) where it inhibits complex I in the ETC in the mitochondria.

Pathology

The two characteristic pathological hallmarks of PD are selective loss of pigmented dopaminergic neurons in SNpc (specifically the A9 dopaminergic neuron group) intracellular accumulation of the protein α -synuclein called Lewy bodies (LBs) [59], and more recently neuroinflammation has now also commonly been accepted as a hallmark of the disease [4].

Lewy bodies

The normal function of α -synuclein is not understood, but it has been suggested to play a role in synaptic vesicle dynamics, as well as mitochondrial function, and possibly as a potential chaperone in intracellular trafficking [60, 61]. The underlying mechanism for formation of

neurotoxic protein accumulation and aggregation is believed to be multifactorial, e.g. protein misfolding and/or overproduction. On the other hand, LBs are not restricted to PD. LBs are observed in LB dementia [62] and normal ageing [63], among others. It is further evident that the absence of LB formation is observed in both sporadic and familial PD, and is thus not pathognomonic for the disease [64].

Selective death of dopaminergic neurons in SNpc

In PD, SNpc undergoes selective and dramatic loss of the A9 dopaminergic neuronal group (figure 2), where > 60% of these neurons are lost at the onset of clinical symptoms [32, 33]. This begs the question as to why these neurons show such exceptional and consistent vulnerability, both in familial and idiopathic forms of PD, especially when compared to other groups of dopaminergic neurons in adjacent regions, such as A10 in the VTA and A8 in the RRF [65] (figure 3). It is recognized that these A9 dopaminergic neurons, in order to massively innervate striatum, provide a vast and largely unmyelinated axonal arbor [66]. Suggested consequences are exceptionally high energy demands to maintain normal cell functions such as cytoskeleton maintenance, axonal transport, protein synthesis, and synaptic transmission, and transmission of action potentials [37]. This massive metabolic load has been shown to increase the risk of oxidative stress in these neurons compared to other dopaminergic neuronal groups [67, 68]. Further, the dopaminergic neurons in SNpc differ greatly from other neurons in the brain, as they are autonomously active through a pacemaker activity. The tonic activity is driven by L-type Cav 1.3 Ca^{2+} channels that elevate the cytoplasmic Ca^{2+} level [69, 70], resulting in high metabolic activity and thus increased production of reactive oxygen species (ROS). The A9 dopaminergic neurons lack the protective Ca^{2+} buffering protein calbindin, thus making the cells more vulnerable to calcium overload [71], although the link between calbindin and dopaminergic neuroprotection has not yet been clarified [72, 73].

In summary, these A9 dopaminergic neurons have been suggested to be especially prone to metabolic and oxidative stress, with convincing evidence of increased oxidative stress in post mortem brain tissue of PD patients [74]. One compelling theory is increased oxidative stress as a consequence of mitochondrial dysfunction.

Figure 3: Topographical organization of the midbrain.

Schematic illustration of the midbrain showing a sagittal section that displays retrorubal field with the A8 dopaminergic neurons, in close proximity to substantia nigra with the A9 dopaminergic neurons and the ventral tegmental area with the A10 dopaminergic neurons.

Modified from Matsuda, 2012 [75]

Mitochondrial dysfunction has been connected to the pathogenesis of PD in a multitude of studies [76-78]. Reduced activity of the complex I of the electron transport chain (ETC) in the mitochondria is found post mortem in PD brain tissue [76]. Adenosine triphosphate (ATP) is generated in the mitochondria through the process of oxidative phosphorylation, where electrons are transferred through different complexes forming the ETC. When electrons leak from these complexes throughout the process of oxidative phosphorylation, ROS can be formed, including superoxide and hydrogen peroxide [78]. It has been suggested that accumulation of α -synuclein leads to deficits of the mitochondrial complex I, increased electron leakage, and thus oxidative stress [79]. Further, animals injected with toxins that impair mitochondrial function, in large mimic features of PD [77]. In MitoPark mice – mice in which mitochondrial transcription factor A is depleted in the dopaminergic neurons – the electron transport chain of the dopaminergic neurons in SNpc becomes defective, leading to dopaminergic neuronal degeneration [80]. Further, mice lacking an allele of En1 (encoding for engrailed 1 – a transcription factor vital for development and survival of dopaminergic neurons) show decreased activity of the mitochondrial complex I and hence progressive degeneration of dopaminergic neurons in the midbrain when approaching adulthood [81]. These mice display several important characteristics of PD e.g. autophagy, neuroinflammation, and dopaminergic neuronal death in SNpc following retrograde axonal

degeneration [81]. The retrograde axonal degeneration has been demonstrated through imaging of the human brain showing changes in the striatum in patients with PD several years before diagnosis [82, 83]. One theory explaining the axonal degeneration is energy deficiency, whereas another explanation is whether α -synuclein aggregation could become an obstacle to axonal transport [84]. Thus, as mitochondrial dysfunction and oxidative stress stands out as plausibly involved in the pathogenesis of PD, the underlying molecular pathogenesis of the selective death of dopaminergic neurons in SNpc is probably multifactorial, involving multiple pathways and mechanisms such as neuroinflammation, which will both be thoroughly discussed in the preceding sections.

In addition to the hallmark of loss of dopaminergic neurons in SNpc, loss of other neurons have also been observed, e.g. dopaminergic neurons in VTA [85], the olfactory bulb [86], in the vagus nerve, and the sympathetic ganglia [86], noradrenergic neurons in locus coeruleus [87], cholinergic neurons in the Meynert basalis nucleus [88], and in the enteric nervous system [89]. This cell loss could potentially account for some of the non-motor symptoms observed in PD.

Neuroinflammation

The CNS has long been considered an organ, except for vast immune responses due to the lack of dendritic cells, as well as the protection of brain parenchyma due to the blood brain barrier (BBB) [90]. However, it is now commonly accepted that the CNS does initiate immune responses when exposed to injury, toxins, pathogens, and cytokines, through activation of the CNS macrophage – microglia – and astrocytes [91] a process called neuroinflammation. This immune response can give rise to chronic CNS inflammation if the source of activation persists or if parts involved in maintaining the inflammatory response is inadequately terminated [92]. In PD, potential disruption of the BBB can enable the crossing of pro-inflammatory molecules from the periphery, however this thesis will focus on the inner CNS immune response.

Neuroinflammation is rather a double-edged sword, encompassing both positive and negative connotations [93]. In recent years, chronic neurodegeneration has become a manifest common denominator in neurodegenerative diseases, including PD, where neurodegeneration is now considered a pathological hallmark [4] central in the disease progression [94-96]. Studies have shown that systemic injections of the endotoxins lipopolysaccharides (LPS) causes the degeneration of dopaminergic neurons in SNpc [97]. Moreover, dopaminergic neurons

cultured with reactive astrocytes induced 25% dopaminergic cell death [98]. As post-mortem analyses of brain from PD patients, as well as AD, HD, ALS and multiple sclerosis (MS) commonly show large amounts of reactive astrocytes and microglia, one must ask whether neuroinflammation could be decisive for the neurodegeneration observed in these diseases. The pathophysiological role of the inner CNS immune response in PD is however not nearly close to being fully understood, and in dire need of further investigation. Even so, the theory of a toxic-loop of initiated and persistent neuroinflammation driven by a cross-talk between the glial cells astrocytes and microglia has gained considerable ground, and will be further elaborated upon and discussed in the following pages.

Treatment

The gold standard for treatment of PD is the substitution of DA in the striatum through systemic administration of levodopa (L-DOPA), thus addressing the core mechanism underlying the cardinal motoric dysfunction of PD [99]. However, as the disease progresses, L-DOPA administration gives rise to motoric complications such as motor response fluctuations (referred to as ‘on’ and ‘off’ state) and iatrogenic dyskinesias [100]. Due to the short half-life of L-DOPA, the problem has been addressed through alternative, and continuous drug-administration, e.g. through percutaneous gastro-jejunostomy tubes [101].

When the STN was identified as part of the basal ganglia circuitry, this gave rise to Deep Brain Stimulation (DBS) as a novel treatment concept for PD [102]. DBS mimics a destructive brain lesion without physically destroying brain tissue using high-frequency electrical stimulation, typically 100-200 Hz, via a surgically implanted electrode [103]. DBS has proven most effective in patients with good response to L-DOPA treatment, but who struggle with motoric complications and fluctuations due to long term medical treatment [103].

Nevertheless, the options for treatment remain purely symptomatic. Recently, cell therapy through transplantation of dopaminergic neuroblasts into the striatum has been suggested as a possible non-pharmacological treatment option. Yet, these experiments raised an important question regarding the microenvironment surrounding these dopaminergic neurons. Where the scientific focus had been mainly directed at the dopaminergic neurons, it was now shifting towards the adjacent environment, which potentially could be directly harmful to the

transplanted dopaminergic neuroblasts. This hypothesis became evident when Li *et al.* showed that grafted embryonic dopaminergic neurons in PD patients developed LBs [104, 105]. They carried out experiments with transplantation of fetal mesencephalic dopaminergic neurons, where development of α -synuclein-positive LBs were observed in long-surviving grafted dopaminergic neurons, morphologically indistinguishable from those seen in SNpc in PD [104], evidently illustrating that the SNpc microenvironment harmfully affects the grafts. This suggests that these grafted dopaminergic neurons were exposed to an unfavorable microenvironment in SNpc. In an attempt to overcome the direct pathological effect of the harmful microenvironment, astrocytes were co-grafted together with neural progenitor cells [106]. This study by Song *et al.* proved that the co-grafted astrocytes improved differentiation of the neuronal progenitor cells into dopaminergic neurons, dopaminergic synaptic maturation, function, and resistance against toxins clinically resulting in virtually complete motoric restoration in the PD rat model, as well as widespread dopaminergic neuronal engraftments. It thus became evident that the microenvironment surrounding the dopaminergic neurons likely plays an important role in the selective vulnerability of these neurons in PD and gaining knowledge and a deeper understanding of the area is crucial.

BRAIN MICROENVIRONMENT IN PARKINSON'S DISEASE

To this point, most of the research on PD has been focused on the dopaminergic neurons and how vulnerability of these neurons potentially could play a role in the PD pathogenesis. It has recently been suggested, however, that as neuronal death is a rapid process, it can hardly explain the gradual neurodegeneration of PD [107]. A plausible theory is that glial dysfunction induces and/or enhances slow and gradual deterioration of the brain microenvironment, ultimately leading to the endpoint of neuronal death, as illustrated by the above-mentioned work of Li *et al.* [104], proving the detrimental effect of the SN microenvironment on the transplanted fetal mesencephalic dopaminergic neurons. This is consistent with the hypothesis that glial cells – key ingredients in the brain microenvironment – play crucial roles in initiation and maintenance of neuroinflammation in PD, proven to be crucial for disease progression [94].

Roles of glial cells in normal and PD brain

Astrocytes in normal and PD brain

Astrocytes are a highly heterogeneous group of neuronal cells, which name translating ‘star-like cell’ was introduced by Michael von Lenhossék in 1895 [108]. However, the majority of astrocytes do not present with such morphology – a somewhat misguided concept born from immunostaining artefacts [109]. Astrocytes have been proposed to increase in number, complexity, and size from lower-order to higher-order species, which have given rise to the question of whether highly developed astrocytes are a necessity for the functioning of higher-order neuronal networks [110, 111]. Furthermore, the glia/neuron –ratio (GNR) differs from species to species, and even though the topic has been wildly debated, the consensus to date is that GNR in the human brain is 1:1, but the ratio differs within different parts of the brain [112]. Animals such as horse, rhesus monkey, Göttingen minipig and even in some parts of the snail *Planorbis corneus*’s brain all display a GNR quite similar to humans [113, 114], whereas elephants and whales show a markedly higher GNR compared to humans [115, 116]. Thus, there is no hard evidence underlining the long-standing misconception of the so called ‘Glial Doctrine’; stating evolutionary glial preservation and glial outnumbering of neurons as potentially fundamental for human intelligence [117, 118].

The astrocytes can functionally and morphologically be divided into protoplasmic astrocytes in the grey matter, encompassing neurons, fibrous astrocytes in white matter, where they are involved in the process of myelination, and radial glia involved in development found in the retina and in the cerebellum [111]. The protoplasmic astrocytes each extend 5-10 processes in a radial manner [119-121]. Each astrocyte radiates 1-5 processes encompassing capillaries, arterioles and venules, forming an overlapping, perivascular glial sheath [122, 123] while other processes separate into finer leaflets enwrapping synapses [124]. Characteristically, astrocytes form a functional cellular syncytium due to their extensive coupling to each other via gap junctions, which is an essential difference in neuroglial and neuronal networking [125]. The astrocytic processes are named according to the structures which they abut; perisynaptic leaflets and perivascular or sub-pial endfeet [126-128] (figure 4). The different membrane domains of the astrocyte cell contain different molecules such as the water channel

AQP4 and the inward-rectifier potassium channel Kir4.1; this is referred to as astrocyte polarization [129-132].

The astrocytes play a major role in the maintenance of homeostasis in the brain microenvironment, e.g. by maintaining the BBB, regulating extracellular water content through the water channel AQP4 [128, 133, 134], through uptake of glutamate through the excitatory amino acid transporter [135], of potassium through the inward-rectifier potassium channel Kir4.1 [136, 137] and modulation of oxidative stress by the production of glutathione (GSH) [124]. Recently, it has been suggested that astrocytes play a major role in elimination of waste products in the brain through the glymphatic system [138, 139]. Further, astrocytes provide important growth factors for neurons stimulating cell survival and synaptic formation [140]. Finally, astrocytes also regulate phagocytosis through secretion of cytokines, inflammatory molecules, and ROS [141, 142]. To enable the execution of this vast spectrum of functions, astrocytes form a functional network of communicating cells connected by gap junctions, which allows for the transfer of ions and signaling molecules by forming a functional syncytium [130].

When the brain is hit by injury or disease, astrocytes respond by changing into their reactive state of astrogliosis [143]. The cells undergo functional and morphological changes and become hypertrophic and sometimes proliferate, as well as changing their molecular expression by increasing the expression of Kir4.1, glutamate aspartate transporter -1 (GLAST), and glial fibrillary acidic protein (GFAP) [144], the latter of which is a cardinal feature of astrogliosis [145, 146]. However, increased levels of GFAP is also recognized in physiological conditions such as physical activity [147] as well as fluctuating levels due to circadian changes [148, 149]. Thus, elevated numbers of GFAP-positive immunoreactivity can represent physiological stimuli and not necessarily represent a response to pathological incitements [149]. These reactive features facilitate the removal of increased extracellular levels of K^+ and glutamate, which are released from damaged cells [150]. Astrocytes respond to ROS and oxidative stress by releasing GSH [151] and adjusting inflammatory response through regulation of microglial activation [152, 153] and are thus reported to be crucial for neuronal protection and survival in an injured or diseased brain. This is underlined by several studies e.g. where intracranial N-methyl-D-aspartate (NMDA) [150] and ATP-injections [154] lead to acute neuronal death, not seen in the penumbra region rich in activated astrocytes. Multiple studies support this by unambiguously reporting neuronal death in the

absence of activated astrocytes and survival in the presence of activated astrocytes [155, 156]. Further, in mice that do not exhibit astrogliosis, such as GFAP- and vimentin knock-out (KO) -mice, both intermediate filaments forming the astroglial cytoskeleton, induced spinal cord injury lead to more severe destruction compared to wild-type (WT) mice [157]. Astrocytes have also been shown to suppress excessive and harmful inflammatory response through the release of soluble factors inhibiting interferon- γ (IFN- γ) -induced microglial expression of inflammatory mediators [153]. Recent studies suggest that astrocytes express growth factors and extracellular matrix proteins supporting axonal growth and thus participate in the repair of an injured brain [158, 159]. Interestingly, it has been suggested that reactive astrocytes in the injured brain could de-differentiate into stem-like cells and re-differentiate into neurons [160]. The overall consensus to date regarding reactive astrocytes is well described by *Escartin et al. 2021*, stating that astrocytes can be classified as ‘diseased astrocytes’ when the astrocytes explicitly initiate disease and disease progression, for example in astrocytopathies such as Alexander disease [161] caused by mutation of GFAP, as well as implicit impact on disease development through astrocytic genetic polymorphisms, exemplified in AD [149, 162].

Increased age is considered the most important risk factor for PD. It has been reported that the glial functions are both altered and decreased with age [163]. Interestingly, the density of astrocytes has been shown to be uncharacteristically low in SN compared to other brain areas such as VTA and neocortex [164]. As the dopaminergic neurons in SN are (as previously described) highly metabolically active, one hypothesis regarding their selective vulnerability in PD is that these cells are less protected by astrocytes by their ability to scavenge ROS by GSH, inhibition of excessive microglial expression, and expressing growth factors such as glial-derived neurotrophic factor (GDNF), shown to be especially important for the survival of dopaminergic neurons [165]. On the other hand, genes associated with PD, such as Parkin, PINK1, DJ-1 and LRRK2, are strongly expressed in astrocytes [166-168]. These genes have been proposed to intervene in the initiation and regulation of astrocyte activation in response to inflammatory mediators and thus disrupting inflammatory signaling pathways. Furthermore, intracellular adhesion molecule 1 (ICAM-1) mediates sustained inflammation and has been found in large quantities in reactive astrocytes in SNpc in post mortem PD brains [169], thus potentially promoting disease progression through prolonged sustained inflammation. The literature is to a large extent conflicted regarding the detrimental or beneficial role of reactive astrogliosis, however the work of *Zamanian et al.* demonstrates the

dualism of these cells. They show that reactive astrocytes can be divided into two groups based on the genes expressed; namely A1 and A2 [170]. The astrocytes expressing genes known to induce inflammatory cascade [171] were termed A1 and were observed in *in vivo* mouse models where animals were treated with systemic LPS -injections. The A2 astrocytes expressed high levels of neurotrophic factors, and these cells were observed in a mouse model of ischemic stroke through transient occlusion of the cerebral middle artery (MCAO), thus believed to be neuroprotective. This is consistent with another study uncovering increased neuronal cell death when culturing neurons with the so called A1 astrocytes [98]. However, *Escartin et. al. 2021* strongly argues against this binary nomenclature, reasoning that the A1/A2, neuroprotectant/neurotoxic compartmentalization of astrocytes simplifies the complexity of reactive astrocytes, as the A1/A2 hallmark genes seem to be upregulated in a mixed manner, where the function of these genes to a large degree is not understood [149]. The authors further proclaim that the functional changes that astrocytes undergo when transitioning in to the state of reactivity, such as loss and/or gain of protective and detrimental functions, could be a simultaneous process where the endpoint of implications for the various pathological states, depends on the balance of the nature of acquired or lost functions, be it detrimental or protective.

Though, it must be mentioned that there is currently no hard evidence for a decisive role of reactive astrogliosis in the pathology of PD, as the literature is conflicting, reporting of high levels of reactive gliosis in both animals studies and in post-mortem human PD patients on one side [172-175], whereas other studies report the opposite [176-178]. Some studies also suggest astrogliosis as a secondary event to neuronal death, indicating that it is not the primary cause of neuronal degeneration.

To conclude, the role of astrocytes in the pathogenesis of PD appears dual in the sense that reactive astrogliosis can be both detrimental and neuroprotective, potentially protecting against oxidative stress and degeneration on one side, while possibly promoting progression of the disease through sustained neuroinflammation on the other. Single-cell RNA sequencing data indicates that astrocytes are a more heterogeneous population than originally anticipated and more subgroups of astrocytes may be identified [179], thus underlining the argument of abandoning the misleading, binary nomenclature [149]. Hence, the mechanisms are not nearly fully understood and the characteristics of the astrocyte population in the SNpc is in need of

closer study in order to further understand how these cells potentially protect or contribute to the neurodegeneration of the dopaminergic neurons in SNpc in PD.

Microglia in normal and PD brain

Microglial cells are immune-cells functioning as CNS tissue macrophages migrating to the brain during embryonic development [180]. They are dispersed throughout the CNS in a resting state and will undergo a transformation when a pathological event occurs, acquiring an amoeboid morphology and the capacity to migrate, proliferate and phagocytize [181]. The resting microglia have highly motile processes which they extend and retract in order to survey the neuronal environment [182]. By continuously pruning the synapses, microglia play a key role in maintaining, and fine-tuning neural circuits [183], which if disrupted is suggested to be of great importance in the development of neurodegenerative disorders [184, 185]. Stimuli such as neuronal death, mechanical injury, and toxins transform the microglia to their activated state with morphologically thicker processes [181], in response to which their extended processes isolate the lesions [186], suggested to prevent progression of further injury as a consequence of disordered microenvironmental homeostasis [187].

The reactive state of microglia, termed reactive microgliosis, is originally divided into the M1 (pro-inflammatory) and M2 (anti-inflammatory) activation state [95], where the M1 state is considered the classical activation state linked to the production of pro-inflammatory cytokines such as tumor necrosis factor (TNF- α), IL-1 β , interleukin-6 (IL-6), interleukin-12 (IL-12), several chemokines as well as superoxide, ROS and nitric oxide (NO) [188, 189]. The alternative activation state M2 is in contrast to the pro-killing state of M1, associated with reconstruction of extracellular matrix (ECM), tissue repair, and induction of neuronal survival by the release of neurotrophic factors, suppressor of cytokine release 1 (SOCS1) and polyamines and interleukin-10 (IL-10) which block pro-inflammatory cytokine production [190-192]. However, similar to the nomenclature describing reactive astrogliosis, the M1/M2 characterization was developed in order to simplify data interpretation, and such schematic polarization of reactive microglia has not been sufficiently supported by research [193]. The overall consensus thus appears in line with the previously mentioned paper by *Escarlin et al. 2021*, moving beyond this binary characterization [149].

The density of microglia is especially high in SN compared to other brain regions [194]. Based on findings in post-mortem PD brains of elaborate microgliosis, pro-inflammatory

cytokines, and clear signs of tissue damage caused by oxidative stress, a significant role of microglia in the pathogenesis of PD is proposed [195-197]. PD is a chronic, not acute disease of the brain, and it is thus suggested that an initial inflammatory response may be beneficial, later becoming detrimental as the disease progresses. It is commonly believed that the dopaminergic neurons in SN in PD are particularly vulnerable to the secretion of factors and solutes associated with reactive microgliosis [198, 199]. This is demonstrated in studies showing extensive loss of dopaminergic neurons in SNpc following injection of the bacterial endotoxin LPS, not seen in nearby regions harbouring large populations of dopaminergic neurons e.g. VTA [200, 201]. In the MPTP mouse model of PD, blocking the microglial inflammation through minocycline treatment resulted in significant dopaminergic neuroprotection [202]. Further, as age is considered one of the most important risk factors in PD, it is interesting that immune cells such as microglia have been shown to be more reactive with ageing, and thus more susceptible to inflammatory stimuli and resistant to regulation [203, 204]. This could potentially amplify neuroinflammatory response. Unfortunately, little is known about hypothetically neuroprotective and anti-inflammatory effects of reactive microgliosis in PD. However, data from the neurodegenerative disease MS as well as in spinal cord injury shows reactive microgliosis displaying mixed features of both neuroprotective and neurodetrimental traits within the site of injury [205], and further experiments where subgroups of reactive microglia characterized by neuroprotective functions are downregulated, show increased lesion size and neuronal death [205]. A better understanding of the reactive microglial response in PD is needed to unravel possible inflammatory therapeutic targets in PD.

The role of glial interaction in neuroinflammation in PD

Activation of microglia is presumed to be influenced by astrocytes and neurons, through a cell-to-cell cross-talk mediated by inflammatory molecules. Recent findings have demonstrated that activation of reactive A1 astrocytes through LPS-injections were dependent of microglia, as cultured astrocytes without the presence of microglia did not fully reach the reactive A1 state [98]. Microglia, on the other hand, interact with neurons through immunoregulatory molecules such as Cx3CL1, CD22, and neuronal cell adhesion molecule (NCAM) released or expressed by neurons, for which microglia cells in large extent solely express receptors [206]. Neurons undergoing the process of neurodegeneration will secrete molecules such as ATP, α -synuclein, and matrix metalloproteinase-3 (MMP-3) further activating microglia [207]. A cross-talk between astrocytes and microglia is commonly

accepted and occurs by molecular signaling through among others ATP and cytokines [208], creating bidirectional feedback suggested by several researchers to play a key role in amplifying a cascade of neuroinflammatory response [209-211]. Such a bidirectional toxic-loop will potentially sustain reactive astro- and microgliosis, inducing chronic neuroinflammation as seen in neurodegenerative disorders such as PD.

Roles of brain aquaporins in normal and PD brain

AQPs are membrane intrinsic proteins that allow passive movement of water and solutes across the plasma membrane, first discovered as a membrane protein in red blood cells facilitating bidirectional water transportation [212]. 13 subtypes of AQPs have been identified in humans, namely AQP0-12 [133] where AQP0,1,2,4,5,6 and 8 are characterized as orthodox AQPs being selectively permeable for water, whereas AQP3,7,9 and 10 are called the aquaglyceroporins, and show permeability for small solutes as well such as glycerol, urea and arsenite [213]. AQP11 and 12 are labeled supra-aquaporins, about which we have scarce information, although suggested to be involved in intracellular water transport [214, 215]. To date, the expression of AQP1,2,3,4,5,8,9 and 11 have been described in rodent brain [216-222]. AQP2,3,5 and 8 are not well described at the protein level, possibly due to deficiency of functioning antibodies. The only AQPs proven localized in the human brain at both mRNA and protein level are AQP1, 4, and 9 [9, 133]. AQP1 is located in ependymal cells of the choroid plexus participating in the formation of CSF [217]. AQP4 and 9 will in the following be described in detail.

Aquaporin-4 in CNS physiology and pathology

AQP4 is the predominant water channel in the brain, especially expressed in the brain fluid interfaces in the astrocytic endfeet encompassing microvessels in the brain [223]. In the CNS, AQP4 water channels are expressed in the astrocyte cell membrane as heterotetramers and exist as M1 (AQPa) and M23 (AQPc) isoforms, where M23 is the predominant isoform [224]. AQP4 is made out of four monomers forming a tetramer with a central pore, where each monomer has a narrow water pore [225]. AQP4 is distributed throughout the CNS, however, there is a significant heterogeneity regarding its expression level and the perivascular density in different brain regions [226]. The polarized expression pattern of AQP4, where the protein mostly is expressed in astrocyte endfeet membrane domains abutting microvessels and the subpial interface [5] (figure 4), can likely be ascribed to the intracellular anchoring

mechanism of AQP4. There is evidence that AQP4 in astroglial paravascular endfeet co-localizes with the dystrophin-associated protein complex (DAPC) with the protein α -syntrophin suggested to directly anchor AQP4 to the DAPC [227-229]. Studies have shown a reverse polarization pattern of AQP4 in α -syntrophin KO-mice where these animals presented significantly reduced AQP4 expression in perivascular astrocyte endfeet and higher expression levels in astrocyte membrane domains facing neuropil [229].

Figure 4: Illustration of astrocyte membrane domains and the distribution pattern of AQP4 in the brain.

The highest density of AQP4 is seen in perivascular and subpial endfeet membrane domains, however astrocytic membranes (leaflets) abutting synapses also show AQP4 expression. Inset displays cross section of astrocytic perivascular endfeet encompassing the capillary. A: Astrocyte; E: Endothelial cell; L: Capillary lumen. Black dots represent AQP4.

AQP4 is widely distributed in the CNS. Multiple papers suggest a key role of AQP4 in the maintenance of a favorable microenvironment through regulation of extracellular volume, circulation of CSF, and buffering of potassium, cell migration, osmosensation, waste clearance and calcium signaling [137, 230] (figure 5). In mice deficient of AQP4, increase of the extracellular volume by 25% has been demonstrated [231]. In addition to clearance of

water from the brain, AQP4 has recently been associated with waste clearance through bulk flow, via a lymphatic system equivalent of the CNS. This particular structure, called the glymphatic system, proposes clearance of water and waste from the brain through a paravascular route mediated by the astrocytic paravascular AQP4 [232]. Depletion of α -syntrophin, the immediate anchoring protein of AQP4 in the endfoot membrane of astrocytes, as well as deletion of AQP4 itself, results in a delayed clearance of potassium following high-frequency synaptic activation [137], suggesting an important role of AQP4 in potassium homeostasis. Since AQP4 is not permeable to potassium or other ions, it is believed that AQP4's contribution to potassium homeostasis is mediated by regulation of the extracellular space volume [134]. AQP4 is further suggested to participate in cell migration, where rapid changes in the shape of the cells occur due to retractions of filopodia and lamellipodia, accompanied by a change in cell volume where AQP4 presents a key role in facilitating rapid water flow into and out of the cell [233].

Figure 5 AQP4 in brain homeostasis

Schematic illustration of the most central roles of AQP4 in brain homeostasis. Middle image adapted from *Nagelhus et al.* [5] illustrating AQP4 structure and tetrameric organization with lipids in ball-and-stick illustration and the water molecules displayed as red spheres.

Most of the data regarding the roles of AQP4 in pathological conditions is based on studies in mice lacking the perivascular AQP4 [234, 235], or knockouts of AQP4 [236]. AQP4 has been

shown to play an important role in CNS conditions involving volume regulations and edema; a dreaded complication of multiple brain diseases such as stroke, inflammation, head trauma, and brain tumors characterized by water accumulation [235, 237]. There are two types of brain edema: vasogenic and cytotoxic. In cytotoxic brain edema, intracellular osmolytes leak into the extracellular space e.g., due to ischemic stroke [236]. Mice lacking AQP4 show reduced water accumulation and mortality in animal models of cytotoxic edema [235], suggesting that lack of AQP4 plays a protective role due to decreased water permeability of the BBB. Leakage of the BBB will, on the other hand, cause vasogenic edema where AQP4 facilitates a transcellular water clearance of the excess water and thus plays a major role in neuronal protection [238]. Further, studies on traumatic brain injury (TBI) where AQP4 was targeted by small interference RNA against AQP4 (siAQP4) show a protective effect in the acute phase three days post injury, with improved motoric function as well as spatial memory, decreased edema formation, increased microglia activation, reduced astrogliosis, and neuronal death in the animals treated with siAQP4 compared to the control animals [239]. Recent studies on normal pressure hydrocephalus (NPH) revealed decreased clearance of CSF-tracers in NPH patients, as well as significantly reduced expression of AQP4 at the paravascular astrocyte membrane domains [240, 241]. Thus AQP4 proves critical in conditions involving regulations of volume and formations of brain edema, where AQP4 functions as an entry route for water flux to the brain when the BBB is intact, and a major exit route for water in the resolution phase of the edema and in conditions with BBB disruption.

AQP4 has also been shown to play an important role in conditions in the CNS where neuroinflammation is the common denominator. Research on neuroinflammatory diseases has shown altered astrocyte expression of AQP4, e.g. increased AQP4 expression in MS and reduced expression in neuromyelitis optica (NMO) [242, 243]. A mouse model of pneumococcal meningitis showed a significant upregulation of AQP4 expression leading to enhanced brain edema, concluding that inhibition of AQP4 function or upregulation could possibly improve clinical outcomes in pneumococcal meningitis [244]. In contrast, a mouse model of cerebral malaria demonstrated increased mortality among the AQP4 KO animals, stating a protective role of AQP4 in cerebral malaria [245]. Although the role of AQP4 in neuroinflammatory processes is not sufficiently clarified, new data present findings indicative of increased expression of AQP4 in acute inflammatory mouse models induced by injections of lysophosphatidylcholine (LPC), LPS, and pro-inflammatory cytokines [246, 247]. Simultaneous results show reduced neuroinflammation and astrocytic release of pro-

inflammatory cytokines (TNF- α and interleukin-6 (IL-6)) in AQP4-KO animals compared to WT [247, 248]. Furthermore, studies have revealed that the known microglia activating ATP is released from astrocytes through an AQP4-coupled process [249, 250].

Neuroinflammation is additionally a common denominator in neurodegenerative diseases [251]. Involvement of AQP4 in neurodegenerative diseases such as ALS has recently been demonstrated [252]. Further, the role of AQP4 in the most common neurodegenerative disorder AD has been shown in a mouse model of AD [253]. The findings demonstrated a redistribution of AQP4 from endfoot membrane domains of the astrocyte to non-endfoot membrane domains encompassing the amyloid- β plaques, indicating a role of AQP4 in the formation of amyloid plaques in AD. Mice lacking AQP4 have shown delayed clearance of amyloid- β in the glymphatic system [254]. These findings indicate that AQP4 may play a role in the pathophysiology of neurodegenerative disorders, although our knowledge so far is limited. The potential role of AQP4 in the second most common neurodegenerative disorder PD thus remains largely unexplored. However, some studies suggest a connection; significant upregulation of AQP4 mRNA was found in a 6-hydroxydopamine (6-OHDA) mouse model of PD [255] and in an MPTP mouse model of PD, increased dopaminergic cell loss, as well as microglial activation in the SN, was detected in the AQP4-KO animals. The results indicate that AQP4 may play a role in the dopaminergic neuronal vulnerability in SNpc [256]. Further studies regarding the potential role of AQP4 in PD pathology are sorely needed.

Aquaporin-9 in CNS physiology and pathology

The aquaglyceroporin AQP9 is broadly expressed in the liver, leukocytes, lung, spleen, bone marrow, adipose tissue and the brain, among others [9, 221, 257-260]. Using Western blotting on mitochondrial membrane fractions from rat brain, in 1995 Amiry-Moghaddam *et al.* demonstrated AQP9-positive bands at the inner mitochondrial membrane and by immunogold cytochemistry showed AQP9 immunopositive mitochondria in a subpopulation of astrocytes and neurons (including dopaminergic neurons) in SN, VTA, and arcuate nucleus [261]. As these experiments were carried out using antibody-dependent techniques such as immunohistochemical labeling, it has been debated whether this resulted from unspecific AQP9 labeling which had been shown in AQP9-KO animals [262]. However, conclusive evidence of AQP9 expression in brain and especially in nigral neurons was obtained by using both immunocytochemistry and *in situ* hybridization techniques demonstrating correspondence with AQP9 mRNA and AQP9 immunoreactive cells, as well as

demonstrating correspondence of tetrameric AQP9 to the functional form of AQP9 through blue native gel analysis [9]. AQP9 expression has further been shown in primate brain, both in astrocytes and neurons [263].

The AQP9 channel permits passage of a broad spectrum of solutes including water, carbamides, polyols, purines, pyrimidines, monocarboxylates such as lactate, ammonia [257, 264, 265] and, interestingly arsenite, a toxic chemical compound as a result of arsenic oxidation [266] that has been linked to PD [53, 267]. There is severely limited knowledge available regarding the physiological role of AQP9 in the brain. In the liver, hepatic AQP9 has shown significant upregulation both on mRNA and protein level in mouse models of Diabetes Mellitus (DM) and insulin resistance, indicative of a role in the pathogenesis of this widespread disease [10, 268]. It has been suggested that AQP9 in the brain, as in the liver [10, 260], could play a role in brain metabolism functioning as a glucose channel and providing passage for alternative energy sources e.g. glycerol, lactate and ketone bodies [269, 270].

As AQP9 facilitates water transportation, it is natural to ask whether this aquaporin, like AQP4, is involved in water homeostasis and the formation of brain edema. AQP9 expressed in astrocytes has proven upregulated in areas surrounding the ischemic lesion following brain ischemia in a mouse model of ischemic stroke [258], proposing a role of AQP9 not only in the regulation of edema following an ischemic stroke, but in the process of clearing the affected area of accumulated lactate [258].

Due to the broad spectrum of solutes permeability and the localization of AQP9 in dopaminergic neurons, our group has suggested a potential role for this aquaglyceroporin in the selective vulnerability of dopaminergic neurons in the pathogenesis of PD [261]. Notably, recent studies have proposed a link between DM and PD [271], again suggesting AQP9 as a factor in the pathogenesis of this neurodegenerative disease. However, this hypothesis is to date merely speculative and is thus in need of further research and experimental substantiation.

HYPOTHESES AND AIMS OF THE THESIS

This thesis endeavors to evaluate the pathophysiological role of AQP4 and AQP9 in the brain by focusing on the brain microenvironment and neuroinflammation in PD and hereby illuminating neuronal and glial factors potentially contributing to the selective vulnerability of dopaminergic neurons in SNpc.

Hypotheses:

- 1) Astrocytic AQP4 contributes to the selective vulnerability of dopaminergic neurons in SNpc by making SN more prone to inflammation.
- 2) AQP9 plays a key role in the dopaminergic neuronal cell death in SNpc by functioning as a gateway channel for neurotoxins into dopaminergic neurons in the SNpc.

Aims:

- 1) To unravel the expression and distribution pattern of AQP4 in SNpc under normal physiological conditions in mice and two different mouse models of PD.
- 2) To investigate whether AQP4 plays a pro-inflammatory role in a mouse model of PD.
- 3) To reveal the potential AQP9 permeability to the neurotoxin MPP⁺ and to investigate susceptibility to MPP⁺ and other parkinsonian toxins such as arsenite in models of PD both *in vitro* and *in vivo* in WT and *Aqp9*^{-/-} animals.

SUMMARY OF RESULTS

Paper I: Subcellular expression of aquaporin-4 in substantia nigra of normal and MPTP-treated mice.

This study was carried out in order to investigate the distribution and expression of AQP4 in SNpc under physiological conditions in untreated animals as well as in a sub-acute mouse model of PD.

Untreated animals showed significantly higher expression of AQP4 in SN compared to the cerebral neocortex, demonstrating especially strong labelling in the perivascular as well as in the non-endfeet membrane domains in the neuropil – a distribution pattern not observed in the neocortex. The distribution pattern was illustrated through confocal and electron microscopic images and proven statistically significant by quantitation.

MPTP-injections in the sub-acute mouse model of PD prompted significant loss of tyrosine hydroxylase (TH)-positive dopaminergic neurons in SNpc compared to saline controls. This was simultaneously accompanied by a statistically significant increase in AQP4 protein expression both in perivascular endfeet membranes of astrocytes, as well as in non-endfeet membrane domains abutting the neuropil. A significant increase of AQP4 expression was also observed in the neocortex. Further, confocal imaging showed clearly increased levels of GFAP in the mice treated with MPTP, compared to untreated animals and saline controls.

We concluded that AQP4 is highly expressed in SNpc astrocytes, both in perivascular membranes and in the neuropil under physiological conditions, and its expression level is significantly increased following the same expression pattern in a sub-acute mouse model of PD. This is supplemented by increased levels of GFAP indicative of astrogliosis.

Paper II: Pro-inflammatory role of AQP4 in mice subjected to intrastriatal injections of the parkinsonogenic toxin MPP⁺

The study was conducted in order to investigate whether the astrocytes in SN, proven in Paper I to greatly express AQP4 compared to cortical astrocytes and further increase after MPTP injections, play a detrimental or protective role in the pathogenesis of PD. This was carried out using an acute PD mouse model through intrastriatal injections of MPP⁺. We further asked whether MPP⁺ treatment led to increased microglial activation and if so, if this was amplified or impaired after the targeted deletion of AQP4.

Our experiments revealed that targeted deletion of AQP4 restrained dopaminergic cell loss in SNpc after intrastriatal injections of the parkinsonogenic toxin MPP⁺ compared to WT littermates. Treatment with MPP⁺ further induced increased expression of genes known to activate microglia, however, compared to WT animals, *Aqp4*^{-/-} animals showed significantly lower expression of these microglial activating genes. Our experiments further revealed an increased width of astrocytic endfeet in WT animals compared to *Aqp4*^{-/-} animals, both under physiological conditions and after MPP⁺ treatment.

We conclude that targeted deletion of AQP4 protects against MPP⁺ toxicity and that AQP4 plays a pro-inflammatory role in our acute mouse model of PD and contributes to the selective vulnerability of DAergic neurons in PD.

Paper III: Targeted deletion of the aquaglyceroporin AQP9 is protective in a mouse model of Parkinson's disease.

This study was undertaken in order to test the hypothesis of AQP9-permeability to the neurotoxin MPP⁺ as well as to conclude whether WT animals are more susceptible to dopaminergic neuronal death in SNpc after MPP⁺ injections than AQP9-deficient mice.

Xenopus oocytes expressing AQP9 showed significantly greater permeability for radiolabeled MPP⁺ and radiolabeled urea compared to oocytes expressing AQP4 and un-injected oocytes. Injections of MPP⁺ demonstrated a significantly higher degree of dopaminergic neuronal death in SNpc in WT animals compared to *Aqp9*^{-/-} animals both *in vitro* using organotypic cultures and *in vivo* using stereotactic intrastriatal injections, with respectively approx. 65% and 46% dopaminergic neuronal cell loss. The *Aqp9*^{-/-} animals also accomplished better scores on the motoric behaviour tests and displayed improved post-operative clinical appearance.

Moreover, *Aqp9*^{-/-} animals proved higher levels of DA and DA metabolites in the injected striatum compared to WT littermates. Furthermore, human embryonic kidney (HEK) cells expressing AQP9 proved more sensitive to MPP⁺ and arsenite toxicity than HEK cells expressing DAT. There were no significant up- or downregulation of DAT, DA receptor 2, or antioxidants in the *Aqp9*^{-/-} animals.

In conclusion, AQP9 is permeable to the neurotoxin MPP⁺ and deletion of AQP9 protects against MPP⁺ neurotoxicity both *in vitro* and *in vivo* in a mouse model of PD. We thus conclude that AQP9 plays a role in the selective vulnerability of dopaminergic neurons in SNpc by functioning as a gateway channel for neurotoxins into dopaminergic neurons in this area.

METHODOLOGICAL CONSIDERATIONS

TRANSGENIC ANIMAL MODELS

The use of mice when modeling human disease is based on the predicted physiological similarities between these species [272, 273]. Combined with easy maintenance and the possibility of many inbred strains, the use of mice in modeling human disease is widely applied [274]. The use of mouse models increased drastically as the method for creating transgenic mouse models with KO and knock-in (KI) mice developed. However, there are numerous examples of different responses in mice to experimental interventions compared to humans. One example is Endostatin, an anticancer drug that cures cancer in mice but has a limited effect in humans [275]. Another example is the toxin used in this thesis (MPTP/MPP+), which shows far greater susceptibility in humans compared to mice [276]. Another important factor to consider when using mouse models is variations between different strains of mice. Common strains of laboratory mice show different traits compared to wild mice and between different inbred strains, possibly due to selection based on early sexual maturity, reproduction frequency, and the lack of genetic variation in the different inbred strains [277, 278]. In Paper II, in addition to C57BL/6J WT mice bred in our animal facility, we also investigated additional animals with the C57BL/6J -background in the shape of externally purchased WT mice from *The Jackson Laboratory*® (n = 12). Surprisingly, the most pronounced difference in TH-ir cell loss was found between the animals purchased externally and the *Aqp4*^{-/-} animals and their WT littermates (WT-INT), with approximately 75%, 38%, and 45% cell loss, respectively. Even though these animals allegedly were genetically identical, their response to MPP+ toxicities did differ from the inbred WT animals and thus had to be used as controls. It is well known that a vast limitation in the use of animal models is the fact that the toxin susceptibility varies in different animals due to differences in strain, age, gender, and general aspects attached to breeding- and animal care conditions [279-282]. Despite this common knowledge, the use of additional WT mice externally purchased due to a shortage of WT littermates is widespread practice.

KO and KI models enable the isolated study of removing or adding a protein, as well as allowing for antibody validation through negative controls. However, the physiological and functional roles of isolated genes studied through KO/KI models comes with potential problems of genetic redundancy, as well as possible effects of genetic background on the phenotypes [283-285]. The transgenic mice in this thesis were created by replacement of

neomycin phosphotransferase expression cassette exon 2 for *Aqp9*^{-/-} [262] and downstream insertion of a flippase recognition target (FRT)-neomycin-FRT-LoxPD cassette of exon 3 and an upstream insertion of a LoxP site of exon 1 for *Aqp4*^{-/-} [249]. In this thesis we validate the KO-models through qRT-PCR, confirming the absence of AQP9 and AQP4 mRNA in the KO-animals. Further, anti-AQP4 incubation in *Aqp4*^{-/-} mice show no immunoreactivity. Antibodies against AQP9 have, on the other hand, shown some unspecific labeling. As the sequence in one of the initial exons is replaced, however, the possibility of translation of any functional form of the protein is unlikely. A potential secondary effect with upregulation or downregulation of other proteins with similar traits is possible due to the plastic nature of the brain, e.g. upregulation of astrocytic connexins in *Aqp4*^{-/-} mice as demonstrated by Katoozi *et al.* [286]. Further, one of the available *Aqp4*^{-/-} strains used by other laboratories, have shown alteration in BBB integrity [287], though evidence against this effect in the *Aqp4*^{-/-} strain used in this thesis has been proven [288].

It should be noted that neither *Aqp4*^{-/-} or *Aqp9*^{-/-} mice show any specific phenotypic characteristics under normal conditions, including development, survival, and gross brain anatomy and morphology [262, 289]. However, *Aqp4*^{-/-} mice have shown less apparent abnormalities such as hearing deficits, enlarged extracellular space, and retinal abnormalities [5, 134, 290]. This raises the question of the functional and physiological importance of these proteins. Noteworthy, they have been well conserved through evolution [291, 292] and have been proven crucial in a stressed or pathological environment such as TBI and ischemic stroke [236, 238], suggesting that aquaporins in the brain may primarily play a role in certain pathological environments.

MODELING PARKINSON'S DISEASE

Animal models of PD

A variety of animal models have been used to create a better understanding of the molecular mechanisms, etiology, and pathophysiology of PD. The animal models of PD can generally be divided into genetic models expressing genetic mutations related to PD, or animal models using environmental or synthetic neurotoxins such as 6-OHDA, MPTP/MPP⁺, rotenone, and paraquat [293]. The animal models using MPTP and MPP⁺ will be the focus in the following, as these are the models that were used in this thesis.

MPTP is highly lipophilic and can thus cross the BBB, after which it is immediately converted to the active toxic metabolite MPP⁺ by monamine oxidase B (MAO-B) in astrocytes and serotonergic neurons and thereafter taken up into the dopaminergic neurons through the DAT [294]. DAT has proven an entry route for parkinsonogenic toxins including MPP⁺, and the selective MPP⁺ sensitivity of the dopaminergic neurons in SNpc has been ascribed to this transporter [295]. On the other side, it is worth highlighting that DAT is not exclusively expressed in dopaminergic neurons. MPP⁺ does show an affinity for DAT on other cell groups and other transporters such as the norepinephrine transporter (NET), however to a much lower extent [295, 296]. This can potentially account for non-dopaminergic cell loss as seen in idiopathic PD e.g. in nucleus coeruleus [297-299], as well as in our models. However, this was not addressed in our experiments. The toxicity of MPP⁺ is induced through inhibition of complex I and III of the mitochondrial electron transport chain, thus resulting in oxidative stress and neurodegeneration of the A9 dopaminergic neurons and depletion of the nigrostriatal tract [300], although the reasons for the selective targeting and high affinity of these A9 dopaminergic neurons are not understood.

The sub-acute model: Systemic MPTP injections

Mimicking PD in rodents, which do not naturally develop the disorder, raises many questions of validity and transferability of the animal data to the human disease. The MPTP/MPP⁺ mouse models of PD mimic a variety of aspects of human PD, such as induction of selective dopaminergic neuronal death in SNpc, leaving VTA largely unaffected [145, 279]. The models have thus been, and still are, important tools for understanding the underlying pathophysiology of PD. As MPTP crosses the BBB after subcutaneous injection (figure 6),

strict safety procedures must be carefully followed when working with this substance. Another limitation with MPTP-injections as a mouse model for PD is variable susceptibility to the toxin in mice due to age, gender, and strains which induces highly variable results [279-281]. Moreover, it has been demonstrated that primates and humans show far greater susceptibility to MPP⁺ than mice [301] and that the dopaminergic neuronal loss in SNpc is transient in contrast to humans and primates exposed to the toxin [276], indicating unknown differences in the neurodegenerative processes in mice compared to humans. Another drawback with this MPTP sub-acute model is the fact that subcutaneous MPTP-injections are associated with cardiac failure attributed to the systemic injections [85]. The mortality rate in our project (Paper I) was approximately 50%. Further, MPP⁺ can show affinity to DAT localized in other cells than dopaminergic neurons, although with a much lower affinity [101], which can explain why we detected a significant upregulation of AQP4 in the adluminal membranes of astrocytes in neocortex as well (Paper I).

Figure 6: Illustration of subcutaneous injection of MPTP in mice.

The acute model: Stereotaxic intrastriatal MPP⁺ injections

MPP⁺ does not cross the BBB and has to be injected directly into the brain in our *in vivo* mouse model. The MPP⁺ model is thus safer (for the investigator) to work with. Using the MPP⁺ model, we avoided possible systemic effects of MPTP on the peripheral organs and the immune system. In contrast to the MPTP model, mice subjected to intrastriatal MPP⁺ (figure 7) in this thesis revealed a considerably lower mortality rate (15%) and a stable dopaminergic cell loss around 65%, corresponding to the clinical onset in human PD, regardless of variations in gender and to some degree age [302]. In addition, one has the opportunity to use

the contralateral hemisphere as an internal control, thereby corresponding to our organotypic model (Paper III). This model further selectively targets the dopaminergic neuron population affected in PD as it mimics the retrograde neurodegeneration by intrastriatal injections in the dorso-lateral striatum (putamen), which is the striatal area receiving dopaminergic projections from the ventral tier of SNpc, which is mostly affected in PD [35, 303].

The most pronounced limitation of the acute MPP⁺ mouse model of PD is the fact that the single injection of the toxin causes massive acute brain damage in the striatum that does not resemble the progressive degenerative course of PD seen in humans. This model leads to more inflammation around the injection site, which can spread to areas beyond the striatum. This must be taken into account when assessing our behavioral studies (Paper III), where the results likely are reflected not only by dopaminergic cell loss in SNpc, but also by striatal tissue destruction. The changes observed in the SN must be attributed to the retrograde transport of MPP⁺ through the nigrostriatal pathway, and so should be regarded as specific. However, injury to striatal axons may in itself cause retrograde unspecific cell death in SNpc. To account for the possibility of hemispheric crossover of nigrostriatal projections, saline controls were included in all the studies.

Figure 7: Illustration of intrastriatal injection of MPP+.

Upper panel: MPP+ was injected into the striatum 0.6 mm anteriorly, 2.2 mm laterally and 3.2 mm ventrally to Bregma (the anatomical point located on the mouse cranium at which the coronal suture intersects with the sagittal and lamda sutures).

Lower panel: Schematic anatomical coronal illustration of injection site.

CPu: Caudate putamen; LV: Lateral ventricle.

Both the MPTP and MPP⁺ models have been shown to produce transient neurotoxic effects to the nigrostriatal dopaminergic neurons in rodents as shown through a transient loss of TH, with no long-term permanent loss of dopaminergic neurons [102]. This is evident in sub-acute models, whereas chronic MPTP models have shown the persistent loss of dopaminergic neurons and striatal dopamine uptake six months after toxin administration [276], thus demonstrating that the neurotoxic effect depends on dosage, species, and type of model [276, 304]. As our experiments lack staining with co-localization of TH and Nissl to ensure actual cell death as well as euthanizing of animals at day 7, i.e., without the possibility of controlling for possible transient TH-ir cell decline, we unfortunately cannot rule out this possibility in our sub-acute and acute model. Also, accumulation of the protein α -synuclein, which is considered one of the pathological hallmarks of PD but not at diagnostic criteria, is only detected in the chronic MPTP model, which was not used in this thesis [102]. Thus, even though in general the MPTP/MPP⁺ models are the golden standard of mouse models in PD, it should be taken into account that a chronic MPTP mouse model using osmotic pumps may be better suited when studying the pathophysiology of a progressive neurodegenerative disease as is the case with PD, especially with regards to a possible loss of polarity in astrocytes. Studies demonstrating loss of astrocyte polarity in mesial temporal lobe epilepsy (MTLE) [305] and AD [253] were mainly performed in chronic mouse models. However, loss of polarized expression of AQP4 was observed in the latent phase of kainite-induced epilepsy [305]. All things considered, despite its limitations, the MPTP/MPP⁺ mouse model is, to this date, considered the best and most widely used PD model. Even though the sub-acute and acute MPTP/MPP⁺ models used in this thesis have obvious limitations when investigating the chronic disorder PD, the models do show two out of three pathological hallmarks of PD – namely selective loss of dopaminergic neurons in SNpc and neuroinflammation. As this thesis endeavors to investigate the dopaminergic neuronal vulnerability to toxins and the role of AQP4 in neuroinflammation, these models thus served their purpose.

IMMUNOCYTOCHEMISTRY

Immunocytochemistry for light microscopy

Immunohistochemistry provides the ability to detect specific proteins and their distribution at the cellular or tissue level by using specific antibodies to specific epitopes in the target protein. With this technique, several proteins within the same section can be labeled, thus enabling visualisation of the proteins in relationship to each other. In this thesis, indirect immunofluorescence was used, meaning the use of two antibodies; a primary antibody directed to the target protein and a secondary antibody attached to a fluorescent dye (fluorophore) directed to the primary antibody. When the fluorescent dye is exposed to light, it will emit light with a specific wavelength which can be visualised by a fluorescent or confocal microscope. The main limitations of the immunofluorescence technique include non-specific binding and background staining (which are common for all immunocytochemical techniques). All the experiments involving antibodies in this thesis have included negative controls incubating the tissue section only in the secondary antibody, ensuring that the labeling visualized can be ascribed to the primary antibody. Moreover, we have previously used AQP4 KO mice to show the specificity of the antibody [306]. Another limitation with immunofluorescence technique, compared with the immunogold technique, is that it is a rather qualitative method and is not suitable for detailed quantitative analyses. However, immunoperoxidase cytochemistry with 3,3'-Diaminobenzidine (DAB) and confocal fluorescence microscopy were particularly useful when evaluating and describing the pattern for dopaminergic neuronal cell loss though visualisation of protein expression on a cellular level.

Immunocytochemistry for electron microscopy

Immunogold cytochemistry for electron micron microscopy provides the opportunity to visualize and quantify proteins at the sub-cellular level [307, 308]. A primary antibody is directed to the target protein, and then a secondary antibody, subsequently coupled to a gold particle, is directed to the primary antibody. The gold particle is electron-dense and can thus be visualised and photographed in the electron microscope. A general concern regarding the immunogold cytochemistry technique is the discrepancy between the localization of the gold particle and the actual protein of interest, as the distance from the center of the gold particle to the target antigen can be 23.5 nm due to the primary-secondary antibody complex when using

15 nm gold particles, as done in our experiments [308] (figure 8). In this project, visualisation of the target protein was performed using post-embedding immunogold cytochemistry. The fixed tissue is embedded in a plastic resin and subjected to the immunogold procedure. The technique is well established for visualisation and quantitation of AQP4, and the antibody used in this thesis is highly specific and extensively tested in earlier projects [306, 309].

Anatomical orientation is another issue that deserves attention when using electron microscopy to analyse relatively small regions such as SNpc. It should be noted that in Paper I the entire SN was included. This problem was addressed by optimizing the protocol in Paper II where sub-regions of interest in the ventral midbrain were dissected out prior to embedding, based on anatomical landmarks. This may have led to the inclusion of some areas where the dopaminergic cell loss is less than the one observed in SN. In addition, even in SNpc there are some differences in degeneration of dopaminergic neurons of the dorsal and ventral tier and the lateral and medial part of SN through the pathophysiological neurodegenerative process of PD. These differences could lead to some variations in our analyses.

Figure 8: Principles underlying a two-step, post-embedding immunogold procedure.

Triangles on the surface of ultrathin sections (hatched) represent the antigen against which the primary antibody (A) was raised. The secondary antibody (B) is attached to a colloidal gold particle (Au); in this case, a gold particle with a radius (r) of 7.5 nm. With an effective IgG diameter of 8 nm, the maximum projected distance between epitope and gold particle will be in the range of 23 nm (two IgG diameters plus gold particle radius). Adapted from Amiry-Moghaddam and Ottersen, 2013 [308].

Antibody specificity and sensitivity

Since Coons *et al.* [310] first described immunohistochemical staining in 1958, the method has been standardised in most laboratories working with cellular localisation of proteins. However, there is no guarantee that an antibody manufactured to localise precise molecular targets will be both specific and sensitive. Thus, there are many pitfalls regarding the use of antibodies. One such pitfall is the possibility of cross-reactions where the antibody might recognize other similar epitopes and therefore cause non-specific labelling. Further, the antibody can stick to other structures than the epitope against which it was raised, such as structures in the embedding material or encompassing tissue [311]. To minimise the probability of non-specific labelling, the antibodies must undergo thorough specificity testing before use in experiments.

The antibodies used in this thesis are both polyclonal and monoclonal. Polyclonal antibodies refer to many clones that will bind the target epitopes on the antigen of interest, and thus are considered to produce stronger labelling intensity compared to monoclonal antibodies that only recognise one epitope. The specificity of anti-AQP4 was tested by comparing the labelling in WT mice to *Aqp4*^{-/-} mice, which is the state-of-the-art when it comes to antibody specificity testing. When KO tissue is unavailable, specificity testing can be carried out by Western blots taken together with labelling patterns of the antigen in question. The specificity for the antibody against TH, Iba1 and GFAP is well documented [312-314]. On the other side, AQP9 lacks specific antibodies which raises difficulties when assessing with certainty which cell types express AQP9. A cross-reaction with AQP9 and GFAP antibodies has been observed [9]. Further, there is to date no evidence for exclusive expression of AQP9 in the A9 dopaminergic neurons, as the A8 neurons in the RRF and the A10 neurons in VTA are not fully studied. Even though our qRT-PCR results from Paper III do show that the expression of AQP9 is significantly higher in the midbrain compared to other brain regions such as the striatum, cortex, and hippocampus, we cannot rule out the idea that other cell types in the midbrain express AQP9, something which must be addressed in future studies. To conclude, the overview regarding AQP9 expression is to date not satisfactory, both in regard to the lack of specific antibodies, and due to the fact that studies on AQP9 expression in the human brain are approximately non-existent. This demands further study.

QUANTITATIVE REAL TIME PCR

Quantitative polymerase chain reaction (qPCR) is a reliable and objective way of detecting and quantifying specific gene expression present at a particular time point at the cellular level. Using reverse transcription qPCR, the RNA sample is reversely transcribed to cDNA using reverse transcriptase. In real-time PCR, amplified DNA is detected throughout the progression of the reaction and not only at the endpoint as seen in conventional PCR [315]. Reverse transcription qPCR makes it possible to rapidly and precisely assess changes in the gene expression as a result of e.g., pathological conditions [316]. The use of TaqMan probes increases the specificity of the detection due to concomitant fluorometric quantification of the PCR product during PCR. When comparing the amount of mRNA from a reference gene to the amount of mRNA generated from the target gene, quantification of the relative mRNA level is possible. Hence, it is of importance that the target gene is stable. To ensure the use of a stable reference gene, several housekeeping genes (*Gapdh*, *Tbp*, *Hprt1*, β -actin) were used as internal, endogenous controls as well as standardisation of the procedure, using NormFinder, *Gapdh* and *Hprt1* turned out to be the most stable. One pitfall using this method is the obligatory use of calibration curves when comparing DNA standards to the target DNA. Here, it is crucial that the amplification efficiencies are the same, and even if the same amplification efficiencies are assumed, amplification of a longer product will naturally generate more signal than a shorter product, hence an overestimation of the product is possible [315]. However, it must be stressed that the mRNA level of a gene does not automatically reflect the actual functional level of protein. The steady-state of protein concentration is determined by transcription, mRNA decay, translation, and protein degradation, which are factors that largely contribute to variations in protein quantities [317]. Further, the cell can keep mRNA in a pool for later use, for instance in a stress condition. All things considered; mRNA levels ideally should be interpreted together with quantitative protein experiments.

STEREOLOGICAL CELL QUANTITATION

Stereology is a method for systematic and unbiased quantitative analysis and three-dimensional interpretation of two-dimensional planar consecutive tissue sections [318]. In this thesis, the optical fractionator method was used [319] as this method allows for the estimation of the total number of cells and cell density in any three-dimensional volume, regardless of

the shape of the region of interest (ROI). The method is not sensitive to factors influencing the size, shape, spatial orientation, or spatial distribution of the cells, such as tissue shrinkage after staining and dehydration of the tissue [319]. The optical fractionator uses the fractionator principle and selects series of systematically random sampled sections with a thickness of 15-30 μm and then, by using the fractionator principle, samples each section in the X, Y, and Z-axis (figure 9). By using this method to sample and then extrapolating a subset of the total number of cells within the three-dimensional volume of the ROI, an unbiased estimate of the total number of cells can be generated.

One concern is the chosen method for defining the ROI, which must be reproducible and validated. We approached this by using anatomical landmarks when identifying each ROI (SNpc, SNpr, VTA) to separate the TH-positive cells in A8 (RRF), A9 (SNpc), and A10 (VTA), using the tractus opticus in the anterior sections and the medial lemniscus in the posterior sections. We further copied the ROI templates in the control hemisphere to the ipsilateral injected hemisphere to ensure the same volume size and create an internal control.

Our sections were cut at 40 μm , allowing them to shrink to approximately 25 μm post-staining and dehydration, and we chose every third section sequentially through SN. Due to previous experience from Stahl *et al.* (unpublished), where cell density along the anterior-posterior axis of SN was found to differ significantly, we included 10 sections to ensure coverage of the entire SN and thus reinforcing the statistical analysis. Our inclusion criteria were well defined, including only cells in which the cell nucleus was clearly visible, as well as an intact cell membrane visibly separated from the nucleus and extracellular environment. When surveying the literature, the reported estimates of TH-positive neurons in the bilateral SNpc ranges from 5200 [320] to 23670 [321], with an average of 8000-12000 neurons [322-324]. Using the sampling procedure, our estimates of the total number of TH-positive neurons in the control hemisphere of WT mice were currently 3000, corresponding to the previously reported range. Our analyses further showed a Gundersen's coefficient of errors at < 10%, indicative of low variance [325].

Figure 9: Stereological quantitation of tissue sample.

Illustration of microscopic view of tissue sample. Optical dissection probes were randomly positioned within the ROI with a thickness of 15-30 μm . TH-positive neurons with visible and intact nuclei were counted if their soma were within or touching the inclusion lines.

DISCUSSION

PHYSIOLOGICAL AND PATHOPHYSIOLOGICAL IMPLICATIONS

This thesis provides novel insight into the roles of the brain AQPs AQP4 and AQP9 in the neurodegenerative disorder PD. Moreover, these findings may also shed light on the roles of AQP4 and 9 in brain physiology and as potential common denominators in a vast spectrum of brain diseases.

As extensively elaborated in the introduction, both AQP4 and AQP9 facilitates transportation of water and the latter is deeply involved in passage of energy sources such as glycerol, lactate and ketone bodies [269, 270]. Paper I provides evidence that AQP4 is highly expressed in SNpc even under physiological conditions, which is an area of the brain where AQP9 also is detected [261]. The physiologically high metabolic rate in this brain region [68] logically explains the expression pattern of these AQPs, as a high demand in energy will result in substantial amounts of metabolic water and also plausibly require a considerable need for alternative energy substrates and lactate buffering. The expression pattern of AQP4 in the brain is quite well mapped, showing increased levels particularly in areas with high metabolic rate such as the visual cortex and the cerebellum [364]. It should be stressed that the studies on AQP9 expression in brain only have been carried out in rodents, and mapping of the human brain is crucial.

Both AQP4 and AQP9 are suggested to be involved in a broad range of brain pathologies [6, 131, 237, 258, 365] where this thesis has focused on novel roles of these channels in the neurodegenerative disease PD and in neuroinflammation. Reactive astro- and microgliosis are common features of several neurodegenerative diseases including PD, AD, HD, ALS and MS [251]. The main risk factor for development of neurodegenerative disorders is age.

Interestingly, older mice show higher expression levels of AQP4 [354] as well as increased sensitivity to the toxin MPTP [353]. Thus, one could ask whether a high expression level of AQP4 and AQP9, for example in areas of the brain with high metabolic demands, could predispose to brain conditions characterized by neuroinflammation; a common feature of numerous brain diseases, including neurodegenerative disorders, and secondary brain injury following TBI.

Thus, AQP4 and AQP9 could potentially predispose to and play key roles in development and progression of neuroinflammation; a common denominator of a vast spectrum of brain pathology. Consequently, therapeutic targeting these protein channels, could potentially redefine neurological disease and treatment.

AQP4 – MICROENVIRONMENTAL LIABILITY

Research on PD has been overwhelmingly focused on the dopaminergic neurons, whereas the abutting microenvironment and its potential role in the PD pathogenesis and underlying mechanisms of the selective vulnerability of the dopaminergic neurons in SNpc has gained little attention, and thus the available information is limited. The theory of microenvironment and hereunder glial dysfunction truly gained significant ground when the work of J. Y. Li *et al.* showed that long-surviving transplanted dopaminergic graphs developed α -synuclein-positive LBs, illustrating a plausible detrimental effect on the graphs by the surrounding microenvironment [104]. Further, water homeostasis has shown increasing relevancy in neurodegenerative diseases including PD. Using diffusion MRI, increased levels of free water in SNpc has been observed in patients with PD where the level of free water positively correlated with clinical measures of cognitive and motoric symptoms [17, 326]. Thus, glial microenvironment and water homeo-/dyshomeostasis becomes increasingly relevant in PD. In that regard, it is of great interest that newly acquired information points directly at the astrocytic water channel AQP4 as a potentially important component in the pathophysiology of neurodegenerative diseases such as AD [11, 253, 327, 328], ALS [329], SE [330] and HD [13], as well as in neuroinflammation [246-248]. Taken together, the scarce information on the mesencephalic microenvironment, and especially astrocytes and AQP4, demands further investigation.

Nigral astroglia are highly enriched in AQP4

In Paper I, we carried out the first detailed study analysing AQP4 localisation and distribution in SN. We found that AQP4 is significantly more abundantly expressed in the SN than in neocortex, both in astrocytic endfoot membranes surrounding microvessels and in the non-endfoot membranes in the neuropil under physiological conditions, with 70% and 135% higher expression in SN, respectively. The physiologically high level of AQP4 in SNpc suggests an important role for this water channel in maintaining homeostasis in this area. As

extensively elaborated in the introduction, dopaminergic neurons in SNpc demand a considerable supply of energy due to their pacemaker activity, resulting in substantial amounts of metabolic water. Furthermore, AQP4 is a homeotetramer with a central pore, where each monomer contains a water channel only able to transport water molecules due to the architecture of the central pore [133]. Interestingly, it has been proposed that the tetrameric central pore of aquaporins can transport other molecules such as CO₂ [331, 332] and possibly also ROS [333] which could further explain the high levels of AQP4 in a brain region with possibly high levels of reactive oxygen species generated, among others, by DA.

AQP4 is expressed in astrocytes in the CNS, and although astrocytes are known to be essential in maintenance of neuronal function, little is known about the morphology and function of these cells in the midbrain and they have gained minor attention regarding the pathophysiology in PD. It has however been demonstrated that the density of astrocytes in the SNpc is particularly low [334]. As astrocytes release a variety of growth factors under normal conditions that are likely to affect neuronal survival [335], one could ask if the low density of astrocytes in SNpc would increase the dopaminergic neuronal vulnerability in this region compared to for example VTA. Further, one of the primary functions of astrocytes is to optimize the interstitial space for synaptic transmission by strict control of water and ionic homeostasis [128] through, among others, AQP4 and the inward rectifying potassium channel Kir4.1 [133]. It has been suggested that the consecutive astrocytic uptake of potassium after neuronal firing, which causes the extracellular potassium level to rise, is coupled to a simultaneous water flux through AQP4 [137], where the two channels work in concert [336]. This could explain the extensive perivascular pool in astrocytic endfeet of AQP4 in SNpc and begs the question of whether a sub-optimal water-potassium homeostasis could be a reason for the selective vulnerability of dopaminergic neurons in SNpc. Further, it is commonly accepted that astrocytes play an important role in neuroinflammation; a pathophysiological hallmark of PD [208] affecting this specific area of the brain. Thus, an interesting take, is whether the high levels of AQP4 in SNpc under physiological conditions shown in Paper I, potentially could lower the threshold for immune response and consequently play a pro-inflammatory role in this particular area of the brain. The work of Li *et al.* [247] demonstrated a pro-inflammatory mechanism mediated by AQP4 linked to the release of cytokines, involving AQP4-dependent astrocyte water permeability and consequently cell swelling followed by release of cytokines *in vitro* and in a mouse model of autoimmune encephalopathy (EAE). In the study, both models, *in vitro* and *in vivo*, showed lower degree

of inflammation in *Aqp4*^{-/-} animals compared to WT mice. Their experiments further revealed release of the cytokines TNF- α and IL-6 from astrocytes after intracerebral injections of LPS and that this cytokine release was AQP4-dependant, possibly due to astrocytic swelling. In the process of neuroinflammation, these cytokines activate microglia [188, 189]. The hypothesis of AQP4-mediated inflammation fits well with the pathogenesis of PD where neuroinflammation has gained ground as a pathological hallmark as the disease is intimately linked to inflammation and associated with elevated levels of cytokines, both in post mortem PD brains as well as in animal models [195-197, 337, 338].

In conclusion, the high levels of AQP4 expression in nigral astrocytes under physiological conditions, may make this area of the brain more prone to neuroinflammation and thus the selective vulnerability of dopaminergic neurons in SNpc seen in PD.

AQP4 – TOXIN EXACERBATOR

After establishing the high levels of AQP4 in SNpc under physiological conditions, we aimed to investigate the expression pattern of AQP4 in two mouse models of PD, keeping in mind that several neurodegenerative disorders have been found to exhibit loss of the polarized expression of AQP4, i.e., in AD [11, 253], HD [13], MTLE [305], ALS [329] as well as in normal ageing [339]. In Paper I and II, we show that this high level of AQP4 in SNpc was further enhanced after systemic injections of the parkinsonogenic neurotoxin MPTP and intrastriatal injections of the active metabolic compound MPP⁺, suggesting that the perivascular pool of AQP4 is additionally increased when exposed to MPTP/MPP⁺. The results in Paper I and II indicate comparable trends in the two mouse models, however some differences were observed. In Paper I, immunogold labeling visualized by electron micrographs corresponding with confocal imaging showed significantly increased expression of AQP4 paravascularly, perineuronally, and in the non-endfoot membranes of astrocytes in the neuropil following systemically MPTP injections. In Paper II, however, the increased AQP4 expression was more pronounced in the neuropil with only small differences when quantitatively compared to saline controls. However, retrospectively, the number of saline controls was low and can probably account for some degree of variation. The results from qRT-PCR were on the other hand highly significant in favor of increased level of AQP4 mRNA in the MPP⁺ treated hemisphere compared both to saline controls and the contralateral hemisphere. The reasons for these discrepancies are likely a result of multiple factors. The

sub-acute MPTP model mimics the chronic neurodegenerative process of PD in a higher degree than the acute MPP⁺ model which creates a severe toxic lesion in the striatum, as discussed under methodological considerations. Further, upregulation of AQP4 is an unrapid process depending on intracellular signaling cascades and phosphorylation [320], a process which plausibly could be interfered with due to the toxic environment characterised by oxidative stress following intracranial MPP⁺ injections. Additionally, perhaps the most important factor separating the two models is the fact that the animals used in the sub-acute MPTP model were significantly younger than the animals used in the acute MPP⁺ model, 6-8 weeks and 5-8 months, respectively. The young animals had a 60% higher baseline level of AQP4 in the saline-treated animals than older mice used in the acute model. This apparent age-related decline in AQP4 expression is in line with previous findings [339].

Having ascertained that the physiologically high level of AQP4 in SNpc is additionally enhanced after MPTP/MPP⁺ exposure, we investigated the potentially harmful or neuroprotective role of this channel in this toxic microenvironment by quantifying dopaminergic cell loss in *Aqp4*^{-/-} animals compared to their WT littermates. We found that intrastriatal unilateral injections of MPP⁺ resulted in significantly reduced dopaminergic cell loss in *Aqp4*^{-/-} animals compared to their WT littermates. However, our results differ from four studies conducted by the same group stating that *Aqp4*^{-/-} animals were more prone to dopaminergic cell death following MPTP-treatment, both in a chronic and sub-acute model [256, 340-342]. However, both their chronic and sub-acute model produced a dopaminergic cell loss at 36.8-42% in the WT-animals, well below cell loss observed in post mortem PD human brains at > 61% [85, 145]. Further, the *Aqp4*^{-/-} animals used in these studies hold fragmented BBB both in MPTP and saline-treated animals and further show baseline motor disturbances [287], neither present in the *Aqp4*^{-/-} animals used in our experiments [289]. A disrupted BBB indicates exposure of the brain microenvironment and neurons to elements such as proteins, molecules and immune cells normally only present in the blood, thus confounding the results. Importantly, there is evidence that the toxin MPTP in itself does not disrupt the BBB [343].

The detrimental role of AQP4 after exposure to the parkinsonogenic toxin MPP⁺ could be coupled to the potassium channel Kir4.1, the expression and spatial potassium buffering of which are closely linked to AQP4 [137, 336]. It has been demonstrated that phasic activation of dopaminergic neurons in SNpc is activated by ATP-sensitive potassium (K-ATP) channels

that enable NMDA-mediated spontaneous burst firing [344]. It has been proposed that K-ATP channel gated burst firing in already metabolically challenged dopaminergic neurons in SN, due to elevated energetic requirements [68, 345], could induce excitotoxicity [325].

Interestingly, K-ATP channels have been shown to be selectively activated in response to complex I inhibition in SN dopaminergic neurons, but not in VTA dopaminergic neurons [346, 347]. Furthermore, genetic inactivation of the K-ATP channel Kir6.2 demonstrated a selective rescue of SN dopaminergic neurons [347]. Thus, genetic deletion of *Aqp4* and consequently preventing the synergetic AQP4/Kir4.1 water/potassium flux, could play a protective role in the toxic microenvironment in SNpc.

Encouraged by the previously mentioned work of Li *et al.* suggesting a pro-inflammatory role of AQP4 mediated by a linked release of inflammatory cytokines as a response to astrocytic cell swelling [247], we hypothesised that the detrimental role of AQP4 in the acute MPP+ PD mouse model could possibly be coupled to inflammation through astrogliosis, microgliosis or even a crosstalk between astrocytes and microglia. We investigated this hypothesis by exploring whether intrastriatal injections of the parkinsonogenic toxin MPP+ resulted in increased expression of genes known to play central roles in reactive microgliosis. We found a highly significant increased expression of genes known to cause reactive microgliosis in WT animals after intrastriatal injections of MPP+. We further detected that targeted deletion of *Aqp4* greatly confines this MPP+ induced gene upregulation, especially genes relevant to microglial migration and phagocytosis. Microglial cells are transformed into their reactive state when exposed to pathological stimuli such as inflammatory cytokines typically released from immune cells, astrocytes, and damaged/dying neurons [188, 189, 191]. As it is well established that brain AQP4 is only expressed in astrocytes, we deduce that the effect on gene expression following MPP+ injection in *Aqp4*^{-/-} animals, is arbitrated by astrocytes (figure 10). The literature is compelling regarding astrocytic activation of microglia, where several lines of evidence state that reactive astrocytes play a crucial role in microglial activation [152, 153] – a key factor in the neuroinflammatory process in PD, which now is considered a pathological hallmark of PD [348]. This has further been demonstrated after astrocytic exposure to α -synuclein [349, 350] and was additionally underlined by the newly published work of Hammond *et al.* [351]. They proved that deleting the astrocyte-specific inhibitor of nuclear factor kappa-B kinase subunit β (IKK2), an activator of the pro-inflammatory nuclear kappa B (NF- κ B) [333], resulted in reduced microglial activation and neuroinflammation after MPTP injections.

As previously described, Li *et al.* showed that astrocytic release of the pro-inflammatory cytokines TNF- α and IL-6 was coupled to AQP4-dependant astrocyte swelling [247]. Interestingly, in line with these findings, it has previously been shown that ATP, known to cause microglia reactivation [352], is dependent on AQP4 when released from astrocytes [249]. In Paper II, we found that mice subjected to targeted deletion of *Aqp4* showed significantly less swelling of astrocyte endfeet after intrastriatal MPP+ injections, compared to WT mice. We further demonstrated that WT animals show a greater baseline endfeet width under physiological conditions compared to *Aqp4*^{-/-} animals. Thus, we draw the conclusion that the high baseline expression of AQP4 in the mesencephalic astrocytes makes a pro-inflammatory liability in this area of the brain, resulting in a possibly stronger inflammatory response when exposed to toxins, plausibly representing a key factor in the selective vulnerability of dopaminergic neurons in SN. This conclusion is further supported by the fact that the sensitivity to the toxin MPTP is provenly higher in older mice [353], which demonstrably show higher expression levels of AQP4 [354], probably inducing their proneness to neuroinflammation.

On the other hand, it should be discussed that astrocytic endfoot swelling indicates cytotoxic edema. Our results correspond to previous findings where *Aqp4*^{-/-} animals showed reduced cytotoxic edema and lower mortality rates in a model of bacterial meningitis [14]. We cannot rule out that parts of the cell loss observed in our experiments, might be due to necrotic cell death induced by cytotoxic edema [355], perhaps as a result of ischemia [356], which in part could explain the significant difference in dopaminergic cell loss, disfavoring WT animals. One note, formation of cytotoxic edema peaks at 48h, thus in the hyperacute phase [357]. Tourdias *et al.* demonstrated a two-phase course in regard to edema build-up using a mice model of neuroinflammation by injecting lysolecithin, where phase one of active inflammation lasted from day 1-7 with peak edema formation at day 3 correlating with increased AQP4 expression, whereas phase two, which lasted from day 7-20 post-injection, showed edema resolution and glial scar formation. This second phase was accompanied by a much stronger AQP4 upregulation [246]. Interestingly, studies on TBI in *Aqp4*^{-/-} mice, showed that AQP4 deficiency plays a protective role in the acute phase, however these animals experienced more severe delayed lesions accompanied by increased microglial inflammatory response [358]. Further, it is suggested that astrocyte recruitment and astrocyte-

mediated neuroprotective pathways are crucial for pre-symptomatic compensation and preservation of the remaining neurons in the early stages of PD [33, 359]. However, our results clearly demonstrate a detrimental, pro-inflammatory role of the astroglial water channel AQP4 in our PD mouse model 7 days post toxin injection. Though, it must be stressed that an acute mouse model assessing neuroinflammation at day 7 presumptively is a sub-optimal way of assessing a chronic disease. However, even at this early stage, we show significantly higher degree of dopaminergic cell loss in the WT animals, which is coherent with the work carried out by Noelker *et al.* [342], demonstrating that toll-like receptor 4 (TLR4) deficient mice showed a provably lower degree of dopaminergic cell loss 7 days after injections of MPTP [360]. TLR4 is known to participate in microglial reactivity in PD [361]. Overall, neuroinflammation, possibly exacerbated by astroglial AQP4, seems to have an overwhelmingly detrimental role in the early stages of the disease.

Figure 10: Pro-inflammatory role of AQP4 after MPP⁺ injection.

In Paper II, we show that intrastriatal injection of the parkinsonigenic toxin MPP⁺ results in significantly higher expression levels of genes known to cause microglial reactivity in SNpc in WT animals. This was however not observed in *Aqp4*^{-/-} animals. Further, WT-mice subjected to MPP⁺ injections, demonstrate higher levels of AQP4 expressed in astrocytes, as well as swelling of astrocytic endfeet. We conclude that AQP4 plays a pro-inflammatory role through AQP4-dependant astrocytic swelling inducing cytokine release, thus activating microglia. Prydz *et al.*, 2020.

To summarize, the results in Paper I and II clearly point to a pro-inflammatory role of AQP4, exacerbating the toxic effect of MPP⁺ following intrastriatal injections in an area of the brain seemingly vulnerable to such toxic exposure, given high baseline expression of this water channel even at physiological conditions. As it appears in Paper II, however, other factors than AQP4 are likely more important in the degenerative process and dopaminergic cell death in SNpc, as proven by substantial dopaminergic cell death also in *Aqp4*^{-/-} animals. Such factors can include high expression of DAT and AQP9, facilitating uptake of MPP⁺ into dopaminergic neurons. This will be more thoroughly discussed in the following section.

AQP9 – TOXIN FACILITATOR

It is an established hypothesis that external factors most likely have an impact on the neurodegenerative process in PD. Such external factors include exposure to environmental neurotoxins; a theory that made considerable ground after the discovery of MPTP [48]. The mechanism of action of the environmental toxins and the cause of the selective vulnerability of these dopaminergic neurons remains unelucidated. Though the DA transporter DAT long ago was established as an entry route for neurotoxins [295], this transporter is for critical physiological reasons excluded as a potential therapeutic target. In Paper III we demonstrate that the aquaglyceroporin AQP9 is permeable to the parkinsonogenic neurotoxin MPP⁺, a protein channel that previously, by our research group, has been shown to be expressed in dopaminergic neurons [9, 261]. We further show that deletion of AQP9 protects against dopaminergic neuronal death in SNpc induced by the neurotoxin MPP⁺, both *in vitro* and *in vivo*. Further, when comparing HEK-cells expressing AQP9 and DAT, we found that cells expressing AQP9 proved more sensitive to the neurotoxic effect of MPP⁺ and arsenite – another known parkinsonogenic toxin. As AQP9 has proven non-critical for brain function [262], it makes for an alluring therapeutic target for pharmacological intervention in PD.

As our experiments established AQP9-permeability to MPP⁺, we aimed to investigate the significance of toxin permeability compared to other known toxic transporters, most relevantly DAT. We discovered that HEK-cells – non-dopaminergic cells that normally do not express AQP9 or DAT – were more sensitive to the toxic effect of MPP⁺ when expressing AQP9 or DAT. The toxicity of arsenite, however, was only exacerbated by AQP9 at very high concentrations of the toxin. These findings underline the relevance of AQP9 as a potential factor in the underlying causes of the selective vulnerability of dopaminergic

neurons in SNpc. This suggests that the localisation of AQP9 in TH-positive neurons in the A9 dopaminergic neuronal group in SNpc, more accurately at the plasma membrane and inner mitochondrial membrane [261], is indisputable. Several studies have identified AQP9 expression in catecholaminergic midbrain neurons [8, 9, 263] as well as in astroglial cells in the brain [258, 362]. However, the immunohistochemical studies must be interpreted with caution, as the lack of specific AQP9 antibodies is evident. Studies have shown that AQP9 antibodies cross-react with GFAP, demonstrated using *Aqp9*^{-/-} animals [262]. On the other side, AQP9 has been demonstrated localised in the SN using *in situ* hybridization techniques and qPCR [262]. Taken together with immunocytochemical studies using WT and *Aqp9*^{-/-} animals establishing AQP9 expression in dopaminergic neuronal cell membranes in SN, as well as in the inner mitochondrial membrane of these neurons [9, 261], the data is reliable (figure 11). In Paper III, we find high levels of AQP9 mRNA in the midbrain, exceeding the mRNA levels in the striatum and cortex. This discovery is in line with previous data stating a neuronal and astroglial pool of AQP9 [9], potentially in dopaminergic neuronal nerve terminals, thus enabling retrograde degeneration of dopaminergic neurons through toxic transportation similar to DAT [363]. Still, it must be stated that the studies on AQP9 expression and localisation in the brain have only been executed in rodents, and that data does not automatically translate to the human brain. Yet, aquaporins have traditionally been well preserved across species throughout evolution [245, 291, 292] and it is not without reason to assume that the highly metabolically active A9 dopaminergic neurons in the human brain hold mechanisms supporting effective exploitation of alternative energy substrates such as lactate, glycerol and ketone bodies – all which AQP9 allows passage of [269, 270]. Therefore, it is plausible that AQP9 plays a beneficial role in the brain metabolism in SNpc, while on the other side is also a detrimental component in the selective vulnerability of these neurons by allowing the passage of neurotoxic compounds.

Figure 11: MPP⁺ transportation and effect

Illustration showing that MPTP crosses the BBB and is further converted to MPP⁺ by monamine-oxidase-B (MAO-B) in non-dopaminergic cells such as astroglia as described in the introduction. MPP⁺ enters the dopaminergic neurons through the dopamine transporter (DAT) and AQP9. Paper III shows that the latter is permeable to MPP⁺ and is located at the cell membrane of dopaminergic neurons, as well as at the inner mitochondrial membrane.

Upon establishing MPP⁺ permeability, we studied the toxic effect on dopaminergic neurons in SNpc comparing WT and *Aqp9*^{-/-} animals. Our results prove that AQP9 deletion causes an evidently protective effect against the neurotoxicity of MPP⁺, using both organotypic cultures *in vitro* as well as *in vivo* using intrastriatal stereotactic injections. Our stereological quantitative analyses showed a loss of TH-positive neurons of 67% in WT mice compared to 48% in *Aqp9*^{-/-} mice, the latter being below cell loss observed in post mortem parkinsonian human brains, shown to be greater than 61% [85]. Interestingly, our organotypic experiments using the known AQP9 blocker phloretin in order to mimic the AQP9-KO effect, showed similar results as seen in the KO-animals, indicating that the results reliably can be ascribed to the AQP9 deletion, rather than a possible off-target effect. In line with these results, we found more severe motor dysfunction as well as worse clinical appearance in the WT animals. These quantitative results were accompanied by 40% higher levels of the DA metabolite homovanillic acid (HVA) in the ipsilateral striatum in *Aqp9*^{-/-} animals, further indicating a

protective effect when genetically deleting or, hypothetically, therapeutically targeting the AQP9 channel.

It must be noted that the *in vivo* studies showed significant TH-positive cell loss not only in the WT animals, but also in *Aqp9*^{-/-} animals. The likely assumption here is that the cell loss observed in the *Aqp9*^{-/-} animals, can be attributed to MPP⁺ transportation through DAT, making AQP9 likely not solely responsible for neurotoxic transportation, but a significant element in a multifactorial process (figure 11). Interestingly, we observed a more pronounced protective effect of AQP9 deletion in the *in vitro* model. This can likely be ascribed to the fact that the model induces severing of the axons, potentially making AQP9 the predominate transportation method for MPP⁺ in this model, as this channel is highly expressed in the somatic plasma membrane.

Having established that AQP9 contributes to the selective vulnerability of dopaminergic neurons in the SNpc and thus likely plays a damaging role in the pathogenesis of PD, it must be emphasized that this channel not only is expressed in neurons, but also in astrocytes in SNpc [362]. Thus, the detrimental effect mediated by AQP9 can just as likely be a result of secondary responses due to astrocytic uptake of neurotoxins.

IMPLICATIONS FOR PARKINSON'S DISEASE

To this date, therapeutic options in PD are solely symptomatic. This shows the importance of detecting and identifying molecules and factors contributing to the selective vulnerability of the dopaminergic neurons in SNpc, thus possibly enabling development of therapeutic targets in PD. In this thesis, two protein channels were identified as molecules potentially contributing to this vulnerability and disease initiation and/or progression, namely AQP4 and AQP9. Systemic blockage of these channels may prove problematic as these AQPs are expressed in organs throughout the body. Development of antagonists specific for the brain could be a solution, although this is merely hypothetical. KO-animals of AQP9 have yet to reveal phenotypic traits and several AQP9-antagonists have already been identified, though preliminary results indicate that they may affect other AQPs as well [257, 366].

Another potential therapeutic approach is targeting neuroinflammation. As we have identified harmful pro-inflammatory compounds that show that AQP4 potentially aggravates the

inflammatory process, one therapeutic approach could be immunosuppression and -modulation through suppression of pro-inflammatory factors, simultaneously potentially engaging neuroprotective immune cells. Several studies have investigated the use of non-steroidal anti-inflammatory drugs (NSAIDs), however, the results have been non-conclusive and splayed [367-370] and more experimental contribution is needed before reaching a conclusion. Interestingly, the immunomodulatory compound Pomalidomide has recently been shown to reduce dopaminergic cell loss in a *Drosophila* LRRK2 model of PD [371]. Nonetheless, the important identification of microenvironmental vulnerability factors for the selective dopaminergic cell death in SNpc in PD may lay the groundwork for future therapeutic discoveries in PD.

CONCLUSION AND FUTURE PERSPECTIVES

This thesis aimed to evaluate the mesencephalic microenvironment and neuroinflammation in PD, especially focusing on the glial and neuronal contributions - including the protein channels AQP4 and AQP9 - to the selective vulnerability of dopaminergic neurons in PD. Using two different mouse models of PD, we uncovered that these channels play detrimental roles in the pathogenesis of PD.

We show that nigral astrocytes express higher levels of AQP4 compared to cortical astrocytes under physiological conditions and that this strong expression is additionally enhanced after subcutaneous injections of MPTP and intrastriatal injections of MPP+. We provide evidence that AQP4 plays a pro-inflammatory role in experimental PD, plausibly linked to AQP4-dependent astrocytic swelling and thus swelling-induced astrocytic release of components causing inflammation and microglia reactivity.

We further demonstrate that AQP9 operates as a gateway channel for neurotoxins into dopaminergic neurons in SNpc, proving a dramatic protective effect on dopaminergic cell loss after targeted deletion of *Aqp9*. This makes AQP9 an intriguing potential therapeutic target.

All in all, we have elucidated the roles of AQP4 and AQP9 in experimental PD, suggesting that these AQPs in a synergetic manner promotes disease initiation and progression where AQP9 facilitates toxic transportation into dopaminergic neurons whereas AQP4 exacerbates toxic effects through inflammatory progression. Future work will illuminate alterations in AQP4- and AQP9-expression in regard to toxin exposure, age and genetic predisposition – both in mice and humans. The experiments carried out in this thesis, should also be conducted using a chronic model of PD as the sub-acute and acute models used in our experiments to a lesser degree mimics the true nature of PD. Further, our data are based on animal studies. We still lack a thorough analysis of human brain and especially SN with regard to the expression of AQPs. Such an analysis on brain samples from PD patients and controls has the highest priority.

REFERENCES

1. Fahn, S., *Description of Parkinson's disease as a clinical syndrome*. Ann N Y Acad Sci, 2003. **991**: p. 1-14.
2. Priyadarshi, A., et al., *A meta-analysis of Parkinson's disease and exposure to pesticides*. Neurotoxicology, 2000. **21**(4): p. 435-40.
3. van der Mark, M., et al., *Is pesticide use related to Parkinson disease? Some clues to heterogeneity in study results*. Environ Health Perspect, 2012. **120**(3): p. 340-7.
4. Troncoso-Escudero, P., et al., *Outside in: Unraveling the Role of Neuroinflammation in the Progression of Parkinson's Disease*. Frontiers in Neurology, 2018. **9**(860).
5. Nagelhus, E.A. and O.P. Ottersen, *Physiological roles of aquaporin-4 in brain*. Physiol Rev, 2013. **93**(4): p. 1543-62.
6. Badaut, J., et al., *Aquaporins in brain: distribution, physiology, and pathophysiology*. J Cereb Blood Flow Metab, 2002. **22**(4): p. 367-78.
7. Nielsen, S., et al., *Specialized membrane domains for water transport in glial cells: high-resolution immunogold cytochemistry of aquaporin-4 in rat brain*. J Neurosci, 1997. **17**(1): p. 171-80.
8. Badaut, J., et al., *Distribution of Aquaporin 9 in the adult rat brain: preferential expression in catecholaminergic neurons and in glial cells*. Neuroscience, 2004. **128**(1): p. 27-38.
9. Mylonakou, M.N., et al., *Analysis of mice with targeted deletion of AQP9 gene provides conclusive evidence for expression of AQP9 in neurons*. J Neurosci Res, 2009. **87**(6): p. 1310-22.
10. Carbrey, J.M., et al., *Aquaglyceroporin AQP9: solute permeation and metabolic control of expression in liver*. Proc Natl Acad Sci U S A, 2003. **100**(5): p. 2945-50.
11. Hoshi, A., et al., *Characteristics of aquaporin expression surrounding senile plaques and cerebral amyloid angiopathy in Alzheimer disease*. J Neuropathol Exp Neurol, 2012. **71**(8): p. 750-9.
12. Kaiser, M., et al., *Progressive loss of a glial potassium channel (KCNJ10) in the spinal cord of the SOD1 (G93A) transgenic mouse model of amyotrophic lateral sclerosis*. J Neurochem, 2006. **99**(3): p. 900-12.
13. Wu, T.T., et al., *Mesenchymal stem cells alleviate AQP-4-dependent glymphatic dysfunction and improve brain distribution of antisense oligonucleotides in BACHD mice*. Stem Cells, 2019.
14. Papadopoulos, M.C. and A.S. Verkman, *Aquaporin-4 gene disruption in mice reduces brain swelling and mortality in pneumococcal meningitis*. J Biol Chem, 2005. **280**(14): p. 13906-12.
15. Promeneur, D., et al., *Protective role of brain water channel AQP4 in murine cerebral malaria*. Proc Natl Acad Sci U S A, 2013. **110**(3): p. 1035-40.
16. Lennon, V.A., et al., *IgG marker of optic-spinal multiple sclerosis binds to the aquaporin-4 water channel*. J Exp Med, 2005. **202**(4): p. 473-7.
17. Ofori, E., et al., *Increased free water in the substantia nigra of Parkinson's disease: a single-site and multi-site study*. Neurobiol Aging, 2015. **36**(2): p. 1097-104.
18. Parkinson, J., *An essay on the shaking palsy. 1817*. J Neuropsychiatry Clin Neurosci, 2002. **14**(2): p. 223-36; discussion 222.
19. Parkinson, J., *AN ESSAY ON THE SHAKING PALSY*. Archives of Neurology, 1969. **20**(4): p. 441-445.
20. De Rijk, M.C., et al., *Prevalence of Parkinson's disease in the elderly: The Rotterdam Study*. Neurology, 1995. **45**(12): p. 2143-2146.

21. Pringsheim, T., et al., *The prevalence of Parkinson's disease: A systematic review and meta-analysis*. 2014. p. 1583-1590.
22. Alves, G., et al., *Incidence of Parkinson's disease in Norway: the Norwegian ParkWest study*. Journal of Neurology, Neurosurgery & Psychiatry, 2009. **80**(8): p. 851.
23. Dorsey, R.E., et al., *Projected number of people with Parkinson disease in the most populous nations, 2005 through 2030*. Neurology, 2007. **68**(5): p. 384-386.
24. Nalls, M.A., et al., *Diagnosis of Parkinson's disease on the basis of clinical and genetic classification: a population-based modelling study*. Lancet Neurol, 2015. **14**(10): p. 1002-9.
25. Fabio, C., *Genetics and Epigenetics of Parkinson's Disease*. The Scientific World Journal, 2012. **2012**.
26. Fahn, S., *Description of Parkinson's disease as a clinical syndrome*. Ann.NY Acad.Sci., 2003. **991**: p. 1-14.
27. Goetz, C.G., et al., *Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): Scale presentation and clinimetric testing results*. Movement Disorders, 2008. **23**(15): p. 2129-2170.
28. Bayulkem, K. and G. Lopez, *Nonmotor fluctuations in Parkinson's disease: clinical spectrum and classification*. J Neurol Sci, 2010. **289**(1-2): p. 89-92.
29. Georgescu, D., et al., *Nonmotor gastrointestinal disorders in older patients with Parkinson's disease: is there hope?* Clin Interv Aging, 2016. **11**: p. 1601-1608.
30. Rizzo, G., et al., *Accuracy of clinical diagnosis of Parkinson disease: A systematic review and meta-analysis*. Neurology, 2016. **86**(6): p. 566-576.
31. Stoessl, A.J., S. Lehericy, and A.P. Strafella, *Imaging insights into basal ganglia function, Parkinson's disease, and dystonia*. Lancet, 2014. **384**(9942): p. 532-44.
32. Gonera, E.G., et al., *Symptoms and duration of the prodromal phase in Parkinson's disease*. Mov Disord, 1997. **12**(6): p. 871-6.
33. Blesa, J., et al., *Compensatory mechanisms in Parkinson's disease: Circuits adaptations and role in disease modification*. Exp Neurol, 2017. **298**(Pt B): p. 148-161.
34. Chen-Plotkin, A.S., *Unbiased approaches to biomarker discovery in neurodegenerative diseases*. Neuron, 2014. **84**(3): p. 594-607.
35. Gerfen, C.R., *Molecular effects of dopamine on striatal-projection pathways*. Trends Neurosci, 2000. **23**(10 Suppl): p. S64-70.
36. Haber, S.N., *The place of dopamine in the cortico-basal ganglia circuit*. Neuroscience, 2014. **282**: p. 248-57.
37. Bolam, J.P., et al., *Synaptic organisation of the basal ganglia*. Journal of anatomy, 2000. **196** (Pt 4): p. 527.
38. Albin, R.L., A.B. Young, and J.B. Penney, *The functional anatomy of basal ganglia disorders*. Trends Neurosci, 1989. **12**(10): p. 366-75.
39. Watts RL, S.D., Obeso JA, *Movement disorders* 3ed. 2011, McGraw-Hill Education. 1052.
40. Grundl, W., et al., *Parkinson's syndrome: cranial computed-tomography findings. Their dependence on sex and age*. Acta Med Hung, 1991. **48**(3-4): p. 127-36.
41. Kalia, L.V. and A.E. Lang, *Parkinson's disease*. Lancet, 2015. **386**(9996): p. 896-912.
42. Schulte, C. and T. Gasser, *Genetic basis of Parkinson's disease: inheritance, penetrance, and expression*. Appl Clin Genet, 2011. **4**: p. 67-80.
43. Coppede, F., *Genetics and epigenetics of Parkinson's disease*. ScientificWorldJournal, 2012. **2012**: p. 489830.

44. Trinh, J. and M. Farrer, *Advances in the genetics of Parkinson disease*. Nat Rev Neurol, 2013. **9**(8): p. 445-54.
45. Xu, K., et al., *Estrogen prevents neuroprotection by caffeine in the mouse 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine model of Parkinson's disease*. J Neurosci, 2006. **26**(2): p. 535-41.
46. Langston, J.W., et al., *Chronic Parkinsonism in humans due to a product of meperidine-analog synthesis*. Science, 1983. **219**(4587): p. 979-80.
47. Chiueh, C.C., et al., *Primate model of parkinsonism: selective lesion of nigrostriatal neurons by 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine produces an extrapyramidal syndrome in rhesus monkeys*. Life Sci, 1985. **36**(3): p. 213-8.
48. Forno, L.S., et al., *Similarities and differences between MPTP-induced parkinsonism and Parkinson's disease. Neuropathologic considerations*. Adv Neurol, 1993. **60**: p. 600-8.
49. Heikkila, R.E., et al., *Some features of the nigrostriatal dopaminergic neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) in the mouse*. Mol Chem Neuropathol, 1989. **10**(3): p. 171-83.
50. Di Monte, D., et al., *Comparative studies on the mechanisms of paraquat and 1-methyl-4-phenylpyridine (MPP+) cytotoxicity*. Biochem Biophys Res Commun, 1986. **137**(1): p. 303-9.
51. Betarbet, R., et al., *Chronic systemic pesticide exposure reproduces features of Parkinson's disease*. Nat Neurosci, 2000. **3**(12): p. 1301-6.
52. Elbaz, A., et al., *Professional exposure to pesticides and Parkinson disease*. Ann Neurol, 2009. **66**(4): p. 494-504.
53. Fan, S.F., P.L. Chao, and A.M. Lin, *Arsenite induces oxidative injury in rat brain: synergistic effect of iron*. Ann N Y Acad Sci, 2010. **1199**: p. 27-35.
54. Li, A.A., et al., *Evaluation of epidemiologic and animal data associating pesticides with Parkinson's disease*. J Occup Environ Med, 2005. **47**(10): p. 1059-87.
55. Nuti, A., et al., *Environmental factors and Parkinson's disease: a case-control study in the Tuscany region of Italy*. Parkinsonism Relat Disord, 2004. **10**(8): p. 481-5.
56. Behari, M., et al., *Risk factors of Parkinson's disease in Indian patients*. J Neurol Sci, 2001. **190**(1-2): p. 49-55.
57. Tanner, C.M., et al., *Rotenone, paraquat, and Parkinson's disease*. Environ Health Perspect, 2011. **119**(6): p. 866-72.
58. Hernan, M.A., et al., *A meta-analysis of coffee drinking, cigarette smoking, and the risk of Parkinson's disease*. Ann Neurol, 2002. **52**(3): p. 276-84.
59. Dickson, D.W., et al., *Neuropathological assessment of Parkinson's disease: refining the diagnostic criteria*. Lancet Neurol, 2009. **8**(12): p. 1150-7.
60. Vekrellis, K., et al., *Pathological roles of alpha-synuclein in neurological disorders*. Lancet Neurol, 2011. **10**(11): p. 1015-25.
61. Wales, P., et al., *Limelight on alpha-synuclein: pathological and mechanistic implications in neurodegeneration*. J Parkinsons Dis, 2013. **3**(4): p. 415-59.
62. McKeith, I.G., et al., *Diagnosis and management of dementia with Lewy bodies: Fourth consensus report of the DLB Consortium*. Neurology, 2017. **89**(1): p. 88-100.
63. Buchman, A.S., et al., *Nigral pathology and parkinsonian signs in elders without Parkinson disease*. Ann Neurol, 2012. **71**(2): p. 258-66.
64. Wakabayashi, K., et al., *Restricted occurrence of Lewy bodies in the dorsal vagal nucleus in a patient with late-onset parkinsonism*. J Neurol Sci, 1999. **165**(2): p. 188-91.
65. Lammel, S., et al., *Unique properties of mesoprefrontal neurons within a dual mesocorticolimbic dopamine system*. Neuron, 2008. **57**(5): p. 760-73.

66. Matsuda, W., et al., *Single nigrostriatal dopaminergic neurons form widely spread and highly dense axonal arborizations in the neostriatum*. J Neurosci, 2009. **29**(2): p. 444-53.
67. Pacelli, C., et al., *Elevated Mitochondrial Bioenergetics and Axonal Arborization Size Are Key Contributors to the Vulnerability of Dopamine Neurons*. Curr Biol, 2015. **25**(18): p. 2349-60.
68. Pissadaki, E.K. and J.P. Bolam, *The energy cost of action potential propagation in dopamine neurons: clues to susceptibility in Parkinson's disease*. Front Comput Neurosci, 2013. **7**: p. 13.
69. Grace, A.A. and B.S. Bunney, *Intracellular and extracellular electrophysiology of nigral dopaminergic neurons--2. Action potential generating mechanisms and morphological correlates*. Neuroscience, 1983. **10**(2): p. 317-31.
70. Surmeier, D.J., J.N. Guzman, and J. Sanchez-Padilla, *Calcium, cellular aging, and selective neuronal vulnerability in Parkinson's disease*. Cell Calcium, 2010. **47**(2): p. 175-82.
71. Catoni, C., T. Cali, and M. Brini, *Calcium, Dopamine and Neuronal Calcium Sensor 1: Their Contribution to Parkinson's Disease*. Front Mol Neurosci, 2019. **12**: p. 55.
72. Airaksinen, M.S., H. Thoenen, and M. Meyer, *Vulnerability of midbrain dopaminergic neurons in calbindin-D28k-deficient mice: lack of evidence for a neuroprotective role of endogenous calbindin in MPTP-treated and weaver mice*. Eur J Neurosci, 1997. **9**(1): p. 120-7.
73. Yuan, H.H., et al., *The neuroprotective effect of overexpression of calbindin-D(28k) in an animal model of Parkinson's disease*. Mol Neurobiol, 2013. **47**(1): p. 117-22.
74. Dias, V., E. Junn, and M.M. Mouradian, *The role of oxidative stress in Parkinson's disease*. J Parkinsons Dis, 2013. **3**(4): p. 461-91.
75. Matsuda, W., *Imaging of Dopaminergic Neurons and the Implications for Parkinson's Disease*, in *Systems Biology of Parkinson's Disease*, P. Wellstead and M. Cloutier, Editors. 2012, Springer New York: New York, NY. p. 1-17.
76. Schapira, A.H., *Mitochondrial dysfunction in Parkinson's disease*. Cell Death Differ, 2007. **14**(7): p. 1261-6.
77. Bose, A. and M.F. Beal, *Mitochondrial dysfunction in Parkinson's disease*. J Neurochem, 2016. **139 Suppl 1**: p. 216-231.
78. Schapira, A.H., *Mitochondria in the aetiology and pathogenesis of Parkinson's disease*. Lancet Neurol, 2008. **7**(1): p. 97-109.
79. Devi, L., et al., *Mitochondrial import and accumulation of alpha-synuclein impair complex I in human dopaminergic neuronal cultures and Parkinson disease brain*. J Biol Chem, 2008. **283**(14): p. 9089-100.
80. Ekstrand, M.I., et al., *Progressive parkinsonism in mice with respiratory-chain-deficient dopamine neurons*. Proc Natl Acad Sci U S A, 2007. **104**(4): p. 1325-30.
81. Nordstroma, U., et al., *Progressive nigrostriatal terminal dysfunction and degeneration in the engrailed1 heterozygous mouse model of Parkinson's disease*. Neurobiol Dis, 2015. **73**: p. 70-82.
82. Sossi, V., et al., *Dopamine turnover increases in asymptomatic LRRK2 mutations carriers*. Mov Disord, 2010. **25**(16): p. 2717-23.
83. Sossi, V., et al., *Changes of dopamine turnover in the progression of Parkinson's disease as measured by positron emission tomography: their relation to disease-compensatory mechanisms*. Journal of Cerebral Blood Flow & Metabolism, 2004. **24**(8): p. 869-876.

84. Lamberts, J.T., E.N. Hildebrandt, and P. Brundin, *Spreading of alpha-synuclein in the face of axonal transport deficits in Parkinson's disease: a speculative synthesis*. Neurobiol Dis, 2015. **77**: p. 276-83.
85. German, D.C., et al., *Midbrain dopaminergic cell loss in Parkinson's disease: computer visualization*. Ann Neurol, 1989. **26**(4): p. 507-14.
86. Braak, H., et al., *Idiopathic Parkinson's disease: possible routes by which vulnerable neuronal types may be subject to neuroinvasion by an unknown pathogen*. J Neural Transm (Vienna), 2003. **110**(5): p. 517-36.
87. Joutsa, J., et al., *Localizing parkinsonism based on focal brain lesions*. Brain, 2018. **141**(8): p. 2445-2456.
88. Candy, J.M., et al., *Pathological changes in the nucleus of Meynert in Alzheimer's and Parkinson's diseases*. J Neurol Sci, 1983. **59**(2): p. 277-89.
89. Derkinderen, P., et al., *Parkinson disease: the enteric nervous system spills its guts*. Neurology, 2011. **77**(19): p. 1761-7.
90. Louveau, A., T.H. Harris, and J. Kipnis, *Revisiting the Mechanisms of CNS Immune Privilege*. Trends Immunol, 2015. **36**(10): p. 569-577.
91. Li, Q. and B.A. Barres, *Microglia and macrophages in brain homeostasis and disease*. Nat Rev Immunol, 2018. **18**(4): p. 225-242.
92. Sochocka, M., B.S. Diniz, and J. Leszek, *Inflammatory Response in the CNS: Friend or Foe?* Mol Neurobiol, 2017. **54**(10): p. 8071-8089.
93. DiSabato, D.J., N. Quan, and J.P. Godbout, *Neuroinflammation: the devil is in the details*. J Neurochem, 2016. **139 Suppl 2**(Suppl 2): p. 136-153.
94. Chao, Y., S.C. Wong, and E.K. Tan, *Evidence of inflammatory system involvement in Parkinson's disease*. Biomed Res Int, 2014. **2014**: p. 308654.
95. Moehle, M.S. and A.B. West, *M1 and M2 immune activation in Parkinson's Disease: Foe and ally?* Neuroscience, 2015. **302**: p. 59-73.
96. Cebrián, C., J.D. Loike, and D. Sulzer, *Neuroinflammation in Parkinson's disease animal models: a cell stress response or a step in neurodegeneration?* Curr Top Behav Neurosci, 2015. **22**: p. 237-70.
97. Qin, L., et al., *Systemic LPS causes chronic neuroinflammation and progressive neurodegeneration*. Glia, 2007. **55**(5): p. 453-62.
98. Liddelow, S.A., et al., *Neurotoxic reactive astrocytes are induced by activated microglia*. Nature, 2017. **541**(7638): p. 481-487.
99. LeWitt, P.A. and S. Fahn, *Levodopa therapy for Parkinson disease: A look backward and forward*. Neurology, 2016. **86**(14 Suppl 1): p. S3-12.
100. Olanow, C.W., J.A. Obeso, and F. Stocchi, *Continuous dopamine-receptor treatment of Parkinson's disease: scientific rationale and clinical implications*. Lancet Neurol, 2006. **5**(8): p. 677-87.
101. Poewe, W. and A. Antonini, *Novel formulations and modes of delivery of levodopa*. Mov Disord, 2015. **30**(1): p. 114-20.
102. Limousin, P., et al., *Effect of parkinsonian signs and symptoms of bilateral subthalamic nucleus stimulation*. Lancet, 1995. **345**(8942): p. 91-5.
103. Bronstein, J.M., et al., *Deep brain stimulation for Parkinson disease: an expert consensus and review of key issues*. Arch Neurol, 2011. **68**(2): p. 165.
104. Li, J.Y., et al., *Lewy bodies in grafted neurons in subjects with Parkinson's disease suggest host-to-graft disease propagation*. Nat Med, 2008. **14**(5): p. 501-3.
105. Kordower, J.H., et al., *Lewy body-like pathology in long-term embryonic nigral transplants in Parkinson's disease*. Nat Med, 2008. **14**(5): p. 504-6.
106. Song, J.J., et al., *Cografting astrocytes improves cell therapeutic outcomes in a Parkinson's disease model*. J Clin Invest, 2018. **128**(1): p. 463-482.

107. Joe, E.-H., et al., *Astrocytes, Microglia, and Parkinson's Disease*. Experimental neurobiology, 2018. **27**(2): p. 77-87.
108. Lenhossék, M., *Der feinere Bau des Nervensystems im Lichte neuester Forschungen: eine allgemeine Betrachtung der Strukturprinzipien des Nervensystems, nebst einer Darstellung des feineren Baues des Rückenmarkes*. 1895: Fischer.
109. Parpura, V. and A. Verkhratsky, *Astrocytes revisited: concise historic outlook on glutamate homeostasis and signaling*. Croat Med J, 2012. **53**(6): p. 518-28.
110. Oberheim, N.A., et al., *Uniquely hominid features of adult human astrocytes*. The Journal of neuroscience : the official journal of the Society for Neuroscience, 2009. **29**(10): p. 3276.
111. Oberheim, N.A., S.A. Goldman, and M. Nedergaard, *Heterogeneity of astrocytic form and function*. Methods Mol Biol, 2012. **814**: p. 23-45.
112. Herculano-Houzel, S., *The glia/neuron ratio: How it varies uniformly across brain structures and species and what that means for brain physiology and evolution*. 2014. p. 1377-1391.
113. Verkhratsky, A., Butt, Arthur M., *The History of the Decline and Fall of the Glial Numbers Legend*. Neuroglia, 2018. **1**(1): p. 188-192.
114. Pentreath, V.W., et al., *The glial cells and glia-neuron relations in the buccal ganglia of Planorbis corneus (L.): cytological, qualitative and quantitative changes during growth and ageing*. Philos Trans R Soc Lond B Biol Sci, 1985. **307**(1133): p. 399-455.
115. Hawkins, A. and J. Olszewski, *Glia/nerve cell index for cortex of the whale*. Science, 1957. **126**(3263): p. 76-7.
116. Tower, D.B., *Structural and functional organization of mammalian cerebral cortex; the correlation of neurone density with brain size; cortical neurone density in the fin whale (Balaenoptera physalus L.) with a note on the cortical neurone density in the Indian elephant*. J Comp Neurol, 1954. **101**(1): p. 19-51.
117. Herculano-Houzel, S., Dos Santos, Sandra E., *You Do Not Mess with the Glia*. Neuroglia, 2018. **1**(1): p. 193-219.
118. Nagelhus, E.A., et al., *The glia doctrine: Addressing the role of glial cells in healthy brain ageing*. Mechanisms of Ageing and Development, 2013. **134**(10): p. 449-459.
119. Oberheim, N.A., S.A. Goldman, and M. Nedergaard, *Heterogeneity of astrocytic form and function*. Methods in molecular biology (Clifton, N.J.), 2012. **814**: p. 23.
120. Calì, C., et al., *3D cellular reconstruction of cortical glia and parenchymal morphometric analysis from Serial Block-Face Electron Microscopy of juvenile rat*. Progress in neurobiology, 2019. **183**: p. 101696.
121. Moye, S.L., et al., *Visualizing astrocyte morphology using lucifer yellow iontophoresis*. JoVE (Journal of Visualized Experiments), 2019(151): p. e60225.
122. Mathiisen, T.M., et al., *The perivascular astroglial sheath provides a complete covering of the brain microvessels: an electron microscopic 3D reconstruction*. Glia, 2010. **58**(9): p. 1094-103.
123. Kacem, K., et al., *Structural organization of the perivascular astrocyte endfeet and their relationship with the endothelial glucose transporter: a confocal microscopy study*. Glia, 1998. **23**(1): p. 1-10.
124. Ventura, R. and K.M. Harris, *Three-dimensional relationships between hippocampal synapses and astrocytes*. Journal of Neuroscience, 1999. **19**(16): p. 6897-6906.
125. Verkhratsky, A. and A. Butt, *Astroglia*. 2013, Chichester, UK: Chichester, UK: John Wiley & Sons, Ltd. 105-244.
126. Simard, M. and M. Nedergaard, *The neurobiology of glia in the context of water and ion homeostasis*. Neuroscience, 2004. **129**(4): p. 877-896.

127. Barres, B.A., *The mystery and magic of glia: a perspective on their roles in health and disease*. Neuron, 2008. **60**(3): p. 430-40.
128. Simard, M. and M. Nedergaard, *The neurobiology of glia in the context of water and ion homeostasis*. Neuroscience, 2004. **129**(4): p. 877-96.
129. Derouiche, A., et al., *Beyond Polarity: Functional Membrane Domains in Astrocytes and Müller Cells*. Neurochemical Research, 2012. **37**(11): p. 2513-2523.
130. Giaume, C. and X. Liu, *From a glial syncytium to a more restricted and specific glial networking*. J Physiol Paris, 2012. **106**(1-2): p. 34-9.
131. Amiry-Moghaddam, M., D.S. Frydenlund, and O.P. Ottersen, *Anchoring of aquaporin-4 in brain: molecular mechanisms and implications for the physiology and pathophysiology of water transport*. Neuroscience, 2004. **129**(4): p. 999-1010.
132. Nedergaard, M., B. Ransom, and S.A. Goldman, *New roles for astrocytes: redefining the functional architecture of the brain*. Trends Neurosci, 2003. **26**(10): p. 523-30.
133. Amiry-Moghaddam, M. and O.P. Ottersen, *The molecular basis of water transport in the brain*. Nat Rev Neurosci, 2003. **4**(12): p. 991-1001.
134. Haj-Yasein, N.N., et al., *Glial-conditional deletion of aquaporin-4 (Aqp4) reduces blood-brain water uptake and confers barrier function on perivascular astrocyte endfeet*. Proc Natl Acad Sci U S A, 2011. **108**(43): p. 17815-20.
135. Rothstein, J.D., et al., *Knockout of glutamate transporters reveals a major role for astroglial transport in excitotoxicity and clearance of glutamate*. Neuron, 1996. **16**(3): p. 675-86.
136. Olsen, M.L., et al., *Functional expression of Kir4.1 channels in spinal cord astrocytes*. Glia, 2006. **53**(5): p. 516-28.
137. Amiry-Moghaddam, M., et al., *Delayed K⁺ clearance associated with aquaporin-4 mislocalization: phenotypic defects in brains of alpha-syntrophin-null mice*. Proc Natl Acad Sci U S A, 2003. **100**(23): p. 13615-20.
138. Iliff, J.J. and M. Nedergaard, *Is there a cerebral lymphatic system?* Stroke, 2013. **44**(6 Suppl 1): p. S93-5.
139. Nedergaard, M., *Neuroscience. Garbage truck of the brain*. Science, 2013. **340**(6140): p. 1529-30.
140. Muller, H.W., U. Junghans, and J. Kappler, *Astroglial neurotrophic and neurite-promoting factors*. Pharmacol Ther, 1995. **65**(1): p. 1-18.
141. Chung, W.S., et al., *Astrocytes mediate synapse elimination through MEGF10 and MERTK pathways*. Nature, 2013. **504**(7480): p. 394-400.
142. Martin, D.L., *Synthesis and release of neuroactive substances by glial cells*. Glia, 1992. **5**(2): p. 81-94.
143. Burda, J.E. and M.V. Sofroniew, *Reactive gliosis and the multicellular response to CNS damage and disease*. Neuron, 2014. **81**(2): p. 229-48.
144. Eng, L.F. and R.S. Ghirnikar, *GFAP and astrogliosis*. Brain Pathol, 1994. **4**(3): p. 229-37.
145. Agid, Y. and J. Blin, *Nerve cell death in degenerative diseases of the central nervous system: clinical aspects*. Ciba Found Symp, 1987. **126**: p. 3-29.
146. Yang, Z. and K.K. Wang, *Glial fibrillary acidic protein: from intermediate filament assembly and gliosis to neurobiomarker*. Trends Neurosci, 2015. **38**(6): p. 364-74.
147. Rodríguez, J.J., et al., *Enriched environment and physical activity reverse astroglial degeneration in the hippocampus of AD transgenic mice*. Cell Death Dis, 2013. **4**(6): p. e678.
148. Gerics, B., F. Szalay, and F. Hajós, *Glial fibrillary acidic protein immunoreactivity in the rat suprachiasmatic nucleus: circadian changes and their seasonal dependence*. J Anat, 2006. **209**(2): p. 231-7.

149. Escartin, C., et al., *Reactive astrocyte nomenclature, definitions, and future directions*. Nat Neurosci, 2021. **24**(3): p. 312-325.
150. Jeong, H.K., et al., *Astrogliosis is a possible player in preventing delayed neuronal death*. Mol Cells, 2014. **37**(4): p. 345-55.
151. Gegg, M.E., et al., *Differential effect of nitric oxide on glutathione metabolism and mitochondrial function in astrocytes and neurones: implications for neuroprotection/neurodegeneration?* J Neurochem, 2003. **86**(1): p. 228-37.
152. Yang, M.S., K.J. Min, and E. Joe, *Multiple mechanisms that prevent excessive brain inflammation*. J Neurosci Res, 2007. **85**(11): p. 2298-305.
153. Kim, J.H., et al., *Astrocytes in injury states rapidly produce anti-inflammatory factors and attenuate microglial inflammatory responses*. J Neurochem, 2010. **115**(5): p. 1161-71.
154. Jeong, H.K., I. Jou, and E.H. Joe, *Absence of Delayed Neuronal Death in ATP-Injected Brain: Possible Roles of Astrogliosis*. Exp Neurobiol, 2013. **22**(4): p. 308-14.
155. Min, K.J., et al., *Spatial and temporal correlation in progressive degeneration of neurons and astrocytes in contusion-induced spinal cord injury*. J Neuroinflammation, 2012. **9**: p. 100.
156. Jeong, H.K., et al., *Brain inflammation and microglia: facts and misconceptions*. Exp Neurobiol, 2013. **22**(2): p. 59-67.
157. Faulkner, J.R., et al., *Reactive astrocytes protect tissue and preserve function after spinal cord injury*. J Neurosci, 2004. **24**(9): p. 2143-55.
158. Tom, V.J., et al., *Astrocyte-associated fibronectin is critical for axonal regeneration in adult white matter*. J Neurosci, 2004. **24**(42): p. 9282-90.
159. White, R.E., F.Q. Yin, and L.B. Jakeman, *TGF-alpha increases astrocyte invasion and promotes axonal growth into the lesion following spinal cord injury in mice*. Exp Neurol, 2008. **214**(1): p. 10-24.
160. Duan, C.L., et al., *Striatal astrocytes transdifferentiate into functional mature neurons following ischemic brain injury*. Glia, 2015. **63**(9): p. 1660-70.
161. Messing, A., et al., *Alexander disease*. J Neurosci, 2012. **32**(15): p. 5017-23.
162. Lin, Y.T., et al., *APOE4 Causes Widespread Molecular and Cellular Alterations Associated with Alzheimer's Disease Phenotypes in Human iPSC-Derived Brain Cell Types*. Neuron, 2018. **98**(6): p. 1141-1154.e7.
163. Streit, W.J., *Microglial senescence: does the brain's immune system have an expiration date?* Trends Neurosci, 2006. **29**(9): p. 506-10.
164. Damier, P., et al., *Glutathione peroxidase, glial cells and Parkinson's disease*. Neuroscience, 1993. **52**(1): p. 1-6.
165. Xing, B., et al., *Glial cell line-derived neurotrophic factor protects midbrain dopaminergic neurons against lipopolysaccharide neurotoxicity*. J Neuroimmunol, 2010. **225**(1-2): p. 43-51.
166. Dodson, M.W. and M. Guo, *Pink1, Parkin, DJ-1 and mitochondrial dysfunction in Parkinson's disease*. Curr Opin Neurobiol, 2007. **17**(3): p. 331-7.
167. Giasson, B.I., *Mitochondrial injury: a hot spot for parkinsonism and Parkinson's disease?* Sci Aging Knowledge Environ, 2004. **2004**(48): p. pe42.
168. Saha, S., et al., *LRRK2 modulates vulnerability to mitochondrial dysfunction in Caenorhabditis elegans*. J Neurosci, 2009. **29**(29): p. 9210-8.
169. Miklossy, J., et al., *Role of ICAM-1 in persisting inflammation in Parkinson disease and MPTP monkeys*. Exp Neurol, 2006. **197**(2): p. 275-83.
170. Zamanian, J.L., et al., *Genomic analysis of reactive astrogliosis*. J Neurosci, 2012. **32**(18): p. 6391-410.

171. Stephan, A.H., B.A. Barres, and B. Stevens, *The complement system: an unexpected role in synaptic pruning during development and disease*. Annu Rev Neurosci, 2012. **35**: p. 369-89.
172. Lastres-Becker, I., et al., *alpha-Synuclein expression and Nrf2 deficiency cooperate to aggravate protein aggregation, neuronal death and inflammation in early-stage Parkinson's disease*. Hum Mol Genet, 2012. **21**(14): p. 3173-92.
173. Thannickal, T.C., Y.Y. Lai, and J.M. Siegel, *Hypocretin (orexin) cell loss in Parkinson's disease*. Brain, 2007. **130**(Pt 6): p. 1586-95.
174. Renkawek, K., G.J. Stege, and G.J. Bosman, *Dementia, gliosis and expression of the small heat shock proteins hsp27 and alpha B-crystallin in Parkinson's disease*. Neuroreport, 1999. **10**(11): p. 2273-6.
175. Knott, C., G.P. Wilkin, and G. Stern, *Astrocytes and microglia in the substantia nigra and caudate-putamen in Parkinson's disease*. Parkinsonism Relat Disord, 1999. **5**(3): p. 115-22.
176. Kurowska, Z., et al., *Signs of degeneration in 12-22-year old grafts of mesencephalic dopamine neurons in patients with Parkinson's disease*. J Parkinsons Dis, 2011. **1**(1): p. 83-92.
177. Mirza, B., et al., *The absence of reactive astrocytosis is indicative of a unique inflammatory process in Parkinson's disease*. Neuroscience, 2000. **95**(2): p. 425-32.
178. Song, Y.J., et al., *Degeneration in different parkinsonian syndromes relates to astrocyte type and astrocyte protein expression*. J Neuropathol Exp Neurol, 2009. **68**(10): p. 1073-83.
179. Miller, S.J., *Astrocyte Heterogeneity in the Adult Central Nervous System*. Front Cell Neurosci, 2018. **12**: p. 401.
180. Kettenmann, H., *Neuroscience: the brain's garbage men*. Nature, 2007. **446**(7139): p. 987.
181. Verkhratsky, A. and A. Butt, *Microglia*. 2013, Chichester, UK: Chichester, UK: John Wiley & Sons, Ltd. 343-380.
182. Tremblay, M.-È., et al., *The role of microglia in the healthy brain*. The Journal of neuroscience : the official journal of the Society for Neuroscience, 2011. **31**(45): p. 16064.
183. Paolicelli, R.C., et al., *Synaptic pruning by microglia is necessary for normal brain development*. Science, 2011. **333**(6048): p. 1456-8.
184. Zhan, Y., et al., *Deficient neuron-microglia signaling results in impaired functional brain connectivity and social behavior*. Nat Neurosci, 2014. **17**(3): p. 400-6.
185. Hong, S., et al., *Complement and microglia mediate early synapse loss in Alzheimer mouse models*. Science, 2016. **352**(6286): p. 712-716.
186. Davalos, D., et al., *ATP mediates rapid microglial response to local brain injury in vivo*. Nat Neurosci, 2005. **8**(6): p. 752-8.
187. Hines, D.J., et al., *Microglia processes block the spread of damage in the brain and require functional chloride channels*. Glia, 2009. **57**(15): p. 1610-8.
188. Nau, G.J., et al., *Human macrophage activation programs induced by bacterial pathogens*. Proc Natl Acad Sci U S A, 2002. **99**(3): p. 1503-8.
189. Martinez, F.O. and S. Gordon, *The M1 and M2 paradigm of macrophage activation: time for reassessment*. F1000Prime Rep, 2014. **6**: p. 13.
190. Martinez, F.O., et al., *Genetic programs expressed in resting and IL-4 alternatively activated mouse and human macrophages: similarities and differences*. Blood, 2013. **121**(9): p. e57-69.
191. Edwards, J.P., et al., *Biochemical and functional characterization of three activated macrophage populations*. J Leukoc Biol, 2006. **80**(6): p. 1298-307.

192. Lu, J., et al., *Discrete functions of M2a and M2c macrophage subsets determine their relative efficacy in treating chronic kidney disease*. *Kidney Int*, 2013. **84**(4): p. 745-55.
193. Ransohoff, R.M., *A polarizing question: do M1 and M2 microglia exist?* *Nat Neurosci*, 2016. **19**(8): p. 987-91.
194. Lawson, L.J., et al., *Heterogeneity in the distribution and morphology of microglia in the normal adult mouse brain*. *Neuroscience*, 1990. **39**(1): p. 151-70.
195. Boka, G., et al., *Immunocytochemical analysis of tumor necrosis factor and its receptors in Parkinson's disease*. *Neurosci Lett*, 1994. **172**(1-2): p. 151-4.
196. McGeer, P.L., et al., *Reactive microglia are positive for HLA-DR in the substantia nigra of Parkinson's and Alzheimer's disease brains*. *Neurology*, 1988. **38**(8): p. 1285-91.
197. Mogi, M., et al., *Interleukin-1 beta, interleukin-6, epidermal growth factor and transforming growth factor-alpha are elevated in the brain from parkinsonian patients*. *Neurosci Lett*, 1994. **180**(2): p. 147-50.
198. Hirsch, E.C., et al., *The role of glial reaction and inflammation in Parkinson's disease*. *Ann N Y Acad Sci*, 2003. **991**: p. 214-28.
199. Gonzalez-Hernandez, T., et al., *Vulnerability of mesostriatal dopaminergic neurons in Parkinson's disease*. *Front Neuroanat*, 2010. **4**: p. 140.
200. Kim, W.G., et al., *Regional difference in susceptibility to lipopolysaccharide-induced neurotoxicity in the rat brain: role of microglia*. *J Neurosci*, 2000. **20**(16): p. 6309-16.
201. Castano, A., et al., *The degenerative effect of a single intranigral injection of LPS on the dopaminergic system is prevented by dexamethasone, and not mimicked by rh-TNF-alpha, IL-1beta and IFN-gamma*. *J Neurochem*, 2002. **81**(1): p. 150-7.
202. Wu, D.C., et al., *Blockade of microglial activation is neuroprotective in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson disease*. *J Neurosci*, 2002. **22**(5): p. 1763-71.
203. Eggen, B.J., et al., *Microglial phenotype and adaptation*. *J Neuroimmune Pharmacol*, 2013. **8**(4): p. 807-23.
204. Norden, D.M. and J.P. Godbout, *Review: microglia of the aged brain: primed to be activated and resistant to regulation*. *Neuropathol Appl Neurobiol*, 2013. **39**(1): p. 19-34.
205. Shechter, R., et al., *Infiltrating blood-derived macrophages are vital cells playing an anti-inflammatory role in recovery from spinal cord injury in mice*. *PLoS Med*, 2009. **6**(7): p. e1000113.
206. Le, W., J. Wu, and Y. Tang, *Protective Microglia and Their Regulation in Parkinson's Disease*. *Front Mol Neurosci*, 2016. **9**: p. 89.
207. Block, M.L., L. Zecca, and J.S. Hong, *Microglia-mediated neurotoxicity: uncovering the molecular mechanisms*. *Nat Rev Neurosci*, 2007. **8**(1): p. 57-69.
208. Jha, M.K., et al., *Microglia-Astrocyte Crosstalk: An Intimate Molecular Conversation*. *Neuroscientist*, 2019. **25**(3): p. 227-240.
209. Kirkley, K.S., et al., *Microglia amplify inflammatory activation of astrocytes in manganese neurotoxicity*. *J Neuroinflammation*, 2017. **14**(1): p. 99.
210. Litwin, M., et al., *The role of FGF2 in migration and tubulogenesis of endothelial progenitor cells in relation to pro-angiogenic growth factor production*. *Mol Cell Biochem*, 2015. **410**(1-2): p. 131-42.
211. Pascual, O., et al., *Microglia activation triggers astrocyte-mediated modulation of excitatory neurotransmission*. *Proc Natl Acad Sci U S A*, 2012. **109**(4): p. E197-205.
212. Preston, G.M., et al., *Appearance of water channels in Xenopus oocytes expressing red cell CHIP28 protein*. *Science*, 1992. **256**(5055): p. 385-7.

213. Gonen, T. and T. Walz, *The structure of aquaporins*. Q Rev Biophys, 2006. **39**(4): p. 361-96.
214. Ishibashi, K., *Aquaporin superfamily with unusual npa boxes: S-aquaporins (superfamily, sip-like and subcellular-aquaporins)*. Cell Mol Biol (Noisy-le-grand), 2006. **52**(7): p. 20-7.
215. Ishibashi, K., et al., *The role of a group III AQP, AQP11 in intracellular organelle homeostasis*. J Med Invest, 2009. **56 Suppl**: p. 312-7.
216. Jung, J.S., et al., *Molecular characterization of an aquaporin cDNA from brain: candidate osmoreceptor and regulator of water balance*. Proc Natl Acad Sci U S A, 1994. **91**(26): p. 13052-6.
217. Agre, P., et al., *Aquaporin CHIP: the archetypal molecular water channel*. Am J Physiol, 1993. **265**(4 Pt 2): p. F463-76.
218. Mobasher, A., S. Wray, and D. Marples, *Distribution of AQP2 and AQP3 water channels in human tissue microarrays*. J Mol Histol, 2005. **36**(1-2): p. 1-14.
219. Kuwahara, S., et al., *Expression of aquaporin water channels in the rat pituitary gland*. J Vet Med Sci, 2007. **69**(11): p. 1175-8.
220. Oshio, K., et al., *Expression of aquaporin water channels in mouse spinal cord*. Neuroscience, 2004. **127**(3): p. 685-93.
221. Elkjaer, M., et al., *Immunolocalization of AQP9 in liver, epididymis, testis, spleen, and brain*. Biochem Biophys Res Commun, 2000. **276**(3): p. 1118-28.
222. Gorelick, D.A., et al., *Aquaporin-11: a channel protein lacking apparent transport function expressed in brain*. BMC Biochem, 2006. **7**: p. 14.
223. Nielsen, S., et al., *Specialized membrane domains for water transport in glial cells: high-resolution immunogold cytochemistry of aquaporin-4 in rat brain*. The Journal of neuroscience : the official journal of the Society for Neuroscience, 1997. **17**(1): p. 171.
224. Smith, A.J., et al., *Aggregation state determines the localization and function of M1- and M23-aquaporin-4 in astrocytes*. The Journal of cell biology, 2014. **204**(4): p. 559.
225. Neely, J.D., et al., *Heterotetrameric composition of aquaporin-4 water channels*. Biochemistry, 1999. **38**(34): p. 11156.
226. Hoddevik, E.H., et al., *Factors determining the density of AQP4 water channel molecules at the brain-blood interface*. Brain Struct Funct, 2016.
227. Blake, D.J., et al., *Function and genetics of dystrophin and dystrophin-related proteins in muscle*. Physiol Rev, 2002. **82**(2): p. 291-329.
228. Amiry-Moghaddam, M., et al., *An alpha-syntrophin-dependent pool of AQP4 in astroglial end-feet confers bidirectional water flow between blood and brain*. Proc Natl Acad Sci U S A, 2003. **100**(4): p. 2106-11.
229. Neely, J.D., et al., *Syntrophin-dependent expression and localization of Aquaporin-4 water channel protein*. Proc Natl Acad Sci U S A, 2001. **98**(24): p. 14108-13.
230. Nagelhus, E.A. and O.P. Ottersen, *Physiological roles of aquaporin-4 in brain*. Physiological reviews, 2013. **93**(4): p. 1543.
231. Yao, X., et al., *Aquaporin-4-deficient mice have increased extracellular space without tortuosity change*. Journal of Neuroscience, 2008. **28**(21): p. 5460-5464.
232. Nedergaard, M., *Neuroscience. Garbage truck of the brain*. Science (New York, N.Y.), 2013. **340**(6140): p. 1529.
233. Papadopoulos, M.C., S. Saadoun, and A.S. Verkman, *Aquaporins and cell migration*. Pflugers Arch, 2008. **456**(4): p. 693-700.
234. Vajda, Z., et al., *Delayed onset of brain edema and mislocalization of aquaporin-4 in dystrophin-null transgenic mice*. Proc Natl Acad Sci U S A, 2002. **99**(20): p. 13131-6.

235. Amiry-Moghaddam, M., D.S. Frydenlund, and O.P. Ottersen, *Anchoring of aquaporin-4 in brain: Molecular mechanisms and implications for the physiology and pathophysiology of water transport*. Neuroscience, 2004. **129**(4): p. 997-1008.
236. Manley, G., et al., *Aquaporin-4 deletion in mice reduces brain edema after acute water intoxication and ischemic stroke*. Nature Medicine, 2000. **6**(2): p. 159-63.
237. Thrane, A.S., V. Rangroo Thrane, and M. Nedergaard, *Drowning stars: reassessing the role of astrocytes in brain edema*. Trends in Neurosciences, 2014. **37**(11): p. 620-628.
238. Papadopoulos, M.C., et al., *Aquaporin-4 facilitates reabsorption of excess fluid in vasogenic brain edema*. Faseb J., 2004. **18**(9).
239. Andrew, M.F., et al., *Posttraumatic reduction of edema with aquaporin-4 RNA interference improves acute and chronic functional recovery*. Journal of Cerebral Blood Flow & Metabolism, 2013. **33**(10): p. 1621.
240. Hasan-Olive, M.M., et al., *Loss of perivascular aquaporin-4 in idiopathic normal pressure hydrocephalus*. Glia, 2019. **67**(1): p. 91-100.
241. Lindstrom, E.K., et al., *Cerebrospinal fluid volumetric net flow rate and direction in idiopathic normal pressure hydrocephalus*. Neuroimage Clin, 2018. **20**: p. 731-741.
242. Sinclair, C., et al., *Absence of aquaporin-4 expression in lesions of neuromyelitis optica but increased expression in multiple sclerosis lesions and normal-appearing white matter*. Acta Neuropathologica, 2007. **113**(2): p. 187-194.
243. Kawachi, I. and H. Lassmann, *Neurodegeneration in multiple sclerosis and neuromyelitis optica*. J Neurol Neurosurg Psychiatry, 2017. **88**(2): p. 137-145.
244. Papadopoulos, M.C. and M.C. Verkman, *Aquaporin-4 gene disruption in mice reduces brain swelling and mortality in pneumococcal meningitis*. Journal of Biological Chemistry, 2005. **280**(14): p. 13906-13912.
245. Promeneur, D., et al., *Protective role of brain water channel AQP4 in murine cerebral malaria*. Proceedings of the National Academy of Sciences of the United States of America, 2013. **110**(3): p. 1035.
246. Tourdias, T., et al., *Differential aquaporin 4 expression during edema build-up and resolution phases of brain inflammation*. J Neuroinflammation, 2011. **8**: p. 143.
247. Li, L., et al., *Proinflammatory role of aquaporin-4 in autoimmune neuroinflammation*. Faseb j, 2011. **25**(5): p. 1556-66.
248. Sisto, M., D. Ribatti, and S. Lisi, *Aquaporin water channels: New perspectives on the potential role in inflammation*. Adv Protein Chem Struct Biol, 2019. **116**: p. 311-345.
249. Thrane, A.S., et al., *Critical role of aquaporin-4 (AQP4) in astrocytic Ca²⁺ signaling events elicited by cerebral edema*. Proc Natl Acad Sci U S A, 2011. **108**(2): p. 846-51.
250. Koizumi, S., et al., *UDP acting at P2Y₆ receptors is a mediator of microglial phagocytosis*. Nature, 2007. **446**(7139): p. 1091-5.
251. Guzman-Martinez, L., et al., *Neuroinflammation as a Common Feature of Neurodegenerative Disorders*. Front Pharmacol, 2019. **10**: p. 1008.
252. Nicaise, C., et al., *Aquaporin-4 as a potential marker of BBB disruption in ALS models*. Amyotrophic Lateral Sclerosis, 2010, Vol.11(1-2), p.253-254, 2010. **11**(1-2): p. 253-254.
253. Yang, J., et al., *Loss of astrocyte polarization in the tg-ArcSwe mouse model of Alzheimer's disease*. Journal of Alzheimer's disease : JAD, 2011. **27**(4): p. 711.
254. Iliff, J.J., et al., *A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid β* . Science translational medicine, 2012. **4**(147): p. 147ra111.

255. Wachter, B., et al., *6-Hydroxydopamine leads to T2 hyperintensity, decreased claudin-3 immunoreactivity and altered aquaporin 4 expression in the striatum*. Behavioural Brain Research, 2012. **232**(1): p. 148-158.
256. Chi, Y., et al., *Novel role of aquaporin-4 in CD4 + CD25 + T regulatory cell development and severity of Parkinson's disease*. Aging Cell, 2011. **10**(3): p. 368-382.
257. Tsukaguchi, H., et al., *Functional and molecular characterization of the human neutral solute channel aquaporin-9*. Am J Physiol, 1999. **277**(5): p. F685-96.
258. Badaut, J., et al., *Astrocyte-specific expression of aquaporin-9 in mouse brain is increased after transient focal cerebral ischemia*. J Cereb Blood Flow Metab, 2001. **21**(5): p. 477-82.
259. Nicchia, G.P., et al., *Tissue distribution and membrane localization of aquaporin-9 water channel: evidence for sex-linked differences in liver*. J Histochem Cytochem, 2001. **49**(12): p. 1547-56.
260. Kishida, K., et al., *Aquaporin adipose, a putative glycerol channel in adipocytes*. J Biol Chem, 2000. **275**(27): p. 20896-902.
261. Amiry-Moghaddam, M., et al., *Brain mitochondria contain aquaporin water channels: evidence for the expression of a short AQP9 isoform in the inner mitochondrial membrane*. Faseb j, 2005. **19**(11): p. 1459-67.
262. Rojek, A.M., et al., *Defective glycerol metabolism in aquaporin 9 (AQP9) knockout mice*. Proc Natl Acad Sci U S A, 2007. **104**(9): p. 3609-14.
263. Arcienega, II, et al., *Cell locations for AQP1, AQP4 and 9 in the non-human primate brain*. Neuroscience, 2010. **167**(4): p. 1103-14.
264. Ishibashi, K., et al., *Cloning and functional expression of a new aquaporin (AQP9) abundantly expressed in the peripheral leukocytes permeable to water and urea, but not to glycerol*. Biochem Biophys Res Commun, 1998. **244**(1): p. 268-74.
265. Holm, L.M., et al., *NH₃ and NH₄⁺ permeability in aquaporin-expressing Xenopus oocytes*. Pflugers Arch, 2005. **450**(6): p. 415-28.
266. Liu, Z., et al., *Arsenite transport by mammalian aquaglyceroporins AQP7 and AQP9*. Proc Natl Acad Sci U S A, 2002. **99**(9): p. 6053-8.
267. Koller, W., et al., *Environmental risk factors in Parkinson's disease*. Neurology, 1990. **40**(8): p. 1218-21.
268. Kuriyama, H., et al., *Coordinated regulation of fat-specific and liver-specific glycerol channels, aquaporin adipose and aquaporin 9*. Diabetes, 2002. **51**(10): p. 2915-21.
269. Davis, J.D., et al., *Sustained intracerebroventricular infusion of brain fuels reduces body weight and food intake in rats*. Science, 1981. **212**(4490): p. 81-3.
270. McKenna, M.C., et al., *Competition of glycerol with other oxidizable substrates in rat brain*. Biochem J, 1986. **237**(1): p. 47-51.
271. Cereda, E., et al., *Diabetes and risk of Parkinson's disease*. Mov Disord, 2013. **28**(2): p. 257.
272. Waterston, R.H., et al., *Initial sequencing and comparative analysis of the mouse genome*. Nature, 2002. **420**(6915): p. 520-62.
273. Perlman, R.L., *Mouse models of human disease: An evolutionary perspective*. Evol Med Public Health, 2016. **2016**(1): p. 170-6.
274. Morse, H., *The Mouse in Biomedical Research* (eds Foster, HL, Small, JD & Fox, JG) 1-16. 1981, Academic, New York.
275. Kerbel, R.S., *What is the optimal rodent model for anti-tumor drug testing?* Cancer Metastasis Rev, 1998. **17**(3): p. 301-4.
276. Petroske, E., et al., *Mouse model of Parkinsonism: a comparison between subacute MPTP and chronic MPTP/probenecid treatment*. Neuroscience, 2001. **106**(3): p. 589-601.

277. Austad, S.N., *A mouse's tale*. Natural history, 2002. **111**(3): p. 64-71.
278. Paigen, K., *One hundred years of mouse genetics: an intellectual history. I. The classical period (1902-1980)*. Genetics, 2003. **163**(1): p. 1-7.
279. Muthane, U., et al., *Differences in nigral neuron number and sensitivity to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine in C57/bl and CD-1 mice*. Exp Neurol, 1994. **126**(2): p. 195-204.
280. Giovanni, A., et al., *Correlation between the neostriatal content of the 1-methyl-4-phenylpyridinium species and dopaminergic neurotoxicity following 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine administration to several strains of mice*. J Pharmacol Exp Ther, 1991. **257**(2): p. 691-7.
281. Giovanni, A., et al., *Studies on species sensitivity to the dopaminergic neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. Part 1: Systemic administration*. J Pharmacol Exp Ther, 1994. **270**(3): p. 1000-7.
282. Jarvis, M.F. and G.C. Wagner, *Age-dependent effects of 1-methyl-4-phenyl-1,2,5,6-tetrahydropyridine (MPTP)*. Neuropharmacology, 1985. **24**(6): p. 581-3.
283. Crusio, W.E., *Flanking gene and genetic background problems in genetically manipulated mice*. Biol Psychiatry, 2004. **56**(6): p. 381-5.
284. Nowak, M.A., et al., *Evolution of genetic redundancy*. Nature, 1997. **388**(6638): p. 167-71.
285. Utriainen, A., et al., *Structurally altered basement membranes and hydrocephalus in a type XVIII collagen deficient mouse line*. Hum Mol Genet, 2004. **13**(18): p. 2089-99.
286. Katoozi, S., et al., *Targeted deletion of Aqp4 promotes the formation of astrocytic gap junctions*. Brain Struct Funct, 2017. **222**(9): p. 3959-3972.
287. Zhou, J., et al., *Altered blood-brain barrier integrity in adult aquaporin-4 knockout mice*. Neuroreport, 2008. **19**(1): p. 1-5.
288. Eilert-Olsen, M., et al., *Deletion of aquaporin-4 changes the perivascular glial protein scaffold without disrupting the brain endothelial barrier*. Glia, 2012. **60**(3): p. 432-40.
289. Saadoun, S., et al., *AQP4 gene deletion in mice does not alter blood-brain barrier integrity or brain morphology*. Neuroscience, 2009. **161**(3): p. 764-72.
290. Yao, X., et al., *Aquaporin-4-deficient mice have increased extracellular space without tortuosity change*. J Neurosci, 2008. **28**(21): p. 5460-4.
291. Johansson, I., et al., *The role of aquaporins in cellular and whole plant water balance*. Biochim Biophys Acta, 2000. **1465**(1-2): p. 324-42.
292. Santoni, V., et al., *The high diversity of aquaporins reveals novel facets of plant membrane functions*. Curr Opin Plant Biol, 2000. **3**(6): p. 476-81.
293. Blesa, J., et al., *Classic and New Animal Models of Parkinson's Disease*. Journal of Biomedicine and Biotechnology, 2012. **2012**.
294. Langston, J.W., et al., *Chronic Parkinsonism in Humans due to a Product of Meperidine-Analog Synthesis*. Science, 1983. **219**(4587): p. 979-980.
295. Javitch, J.A., et al., *Parkinsonism-inducing neurotoxin, N-methyl-4-phenyl-1,2,3,6-tetrahydropyridine: uptake of the metabolite N-methyl-4-phenylpyridine by dopamine neurons explains selective toxicity*. Proc Natl Acad Sci U S A, 1985. **82**(7): p. 2173-7.
296. Sonsalla, P.K., G.D. Zeevalk, and D.C. German, *Chronic intraventricular administration of 1-methyl-4-phenylpyridinium as a progressive model of Parkinson's disease*. Parkinsonism Relat Disord, 2008. **14 Suppl 2**(Suppl 2): p. S116-8.
297. Braak, H. and K. Del Tredici, *Neuropathological Staging of Brain Pathology in Sporadic Parkinson's disease: Separating the Wheat from the Chaff*. J Parkinsons Dis, 2017. **7**(s1): p. S71-s85.
298. Kim, S., et al., *Transneuronal Propagation of Pathologic α -Synuclein from the Gut to the Brain Models Parkinson's Disease*. Neuron, 2019. **103**(4): p. 627-641.e7.

299. Tildon, J.T. and L.M. Roeder, *Glycerol oxidation in rat brain: subcellular localization and kinetic characteristics*. J Neurosci Res, 1980. **5**(1): p. 7-17.
300. Nicklas, W.J., et al., *MPTP, MPP+ and mitochondrial function*. Life sciences, 1987. **40**(8): p. 721.
301. Jackson-Lewis, V. and S. Przedborski, *Protocol for the MPTP mouse model of Parkinson's disease*. Nat Protoc, 2007. **2**(1): p. 141-51.
302. Halliday, G., A. Lees, and M. Stern, *Milestones in Parkinson's disease--clinical and pathologic features*. Mov Disord, 2011. **26**(6): p. 1015-21.
303. Gibb, W.R. and A.J. Lees, *Anatomy, pigmentation, ventral and dorsal subpopulations of the substantia nigra, and differential cell death in Parkinson's disease*. J Neurol Neurosurg Psychiatry, 1991. **54**(5): p. 388-96.
304. Gibrat, C., et al., *Differences between subacute and chronic MPTP mice models: investigation of dopaminergic neuronal degeneration and alpha-synuclein inclusions*. J Neurochem, 2009. **109**(5): p. 1469-82.
305. Alvestad, S., et al., *Mislocalization of AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy*. Epilepsy Res, 2013. **105**(1-2): p. 30-41.
306. Hoddevik, E.H., et al., *Factors determining the density of AQP4 water channel molecules at the brain-blood interface*. Brain Struct Funct, 2017. **222**(4): p. 1753-1766.
307. Ottersen, O.P., *Quantitative electron microscopic immunocytochemistry of neuroactive amino acids*. Anat Embryol (Berl), 1989. **180**(1): p. 1-15.
308. Amiry-Moghaddam, M. and O.P. Ottersen, *Immunogold cytochemistry in neuroscience*. Nat Neurosci, 2013. **16**(7): p. 798-804.
309. Gundersen, G.A., et al., *Evidence that pericytes regulate aquaporin-4 polarization in mouse cortical astrocytes*. Brain Struct Funct, 2014. **219**(6): p. 2181-6.
310. Coons, A.H., *Fluorescent antibody methods*. Gen Cytochem Methods, 1958. **1**: p. 399-422.
311. Holmseth, S., et al., *Specificity of antibodies: unexpected cross-reactivity of antibodies directed against the excitatory amino acid transporter 3 (EAAT3)*. Neuroscience, 2005. **136**(3): p. 649-60.
312. Bianchini, D., et al., *GFAP expression of human Schwann cells in tissue culture*. Brain Res, 1992. **570**(1-2): p. 209-17.
313. Pandya, H., et al., *Differentiation of human and murine induced pluripotent stem cells to microglia-like cells*. Nat Neurosci, 2017. **20**(5): p. 753-759.
314. Xu, J., et al., *Dopamine-dependent neurotoxicity of alpha-synuclein: a mechanism for selective neurodegeneration in Parkinson disease*. Nat Med, 2002. **8**(6): p. 600-6.
315. Bustin, S.A. and T. Nolan, *Pitfalls of quantitative real-time reverse-transcription polymerase chain reaction*. J Biomol Tech, 2004. **15**(3): p. 155-66.
316. Sanders, R., et al., *Considerations for accurate gene expression measurement by reverse transcription quantitative PCR when analysing clinical samples*. Anal Bioanal Chem, 2014. **406**(26): p. 6471-83.
317. Vogel, C., et al., *Sequence signatures and mRNA concentration can explain two-thirds of protein abundance variation in a human cell line*. Mol Syst Biol, 2010. **6**: p. 400.
318. Mouton, P.R., et al., *Unbiased estimation of cell number using the automatic optical fractionator*. J Chem Neuroanat, 2017. **80**: p. A1-a8.
319. West, M.J., L. Slomianka, and H.J. Gundersen, *Unbiased stereological estimation of the total number of neurons in the subdivisions of the rat hippocampus using the optical fractionator*. Anat Rec, 1991. **231**(4): p. 482-97.

320. Coban, A. and N.M. Filipov, *Dopaminergic toxicity associated with oral exposure to the herbicide atrazine in juvenile male C57BL/6 mice*. J Neurochem, 2007. **100**(5): p. 1177-87.
321. Jakowec, M.W., et al., *Tyrosine hydroxylase and dopamine transporter expression following 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced neurodegeneration of the mouse nigrostriatal pathway*. J Neurosci Res, 2004. **76**(4): p. 539-50.
322. McCormack, A.L., et al., *Decreased susceptibility to oxidative stress underlies the resistance of specific dopaminergic cell populations to paraquat-induced degeneration*. Neuroscience, 2006. **141**(2): p. 929-37.
323. Walters, T.L., et al., *Diethyldithiocarbamate causes nigral cell loss and dopamine depletion with nontoxic doses of MPTP*. Exp Neurol, 1999. **156**(1): p. 62-70.
324. Liberatore, G.T., et al., *Inducible nitric oxide synthase stimulates dopaminergic neurodegeneration in the MPTP model of Parkinson disease*. Nat Med, 1999. **5**(12): p. 1403-9.
325. Glaser, E.M. and P.D. Wilson, *The coefficient of error of optical fractionator population size estimates: a computer simulation comparing three estimators*. J Microsc, 1998. **192**(Pt 2): p. 163-71.
326. Ofori, E., et al., *Free water improves detection of changes in the substantia nigra in parkinsonism: A multisite study*. Mov Disord, 2017. **32**(10): p. 1457-1464.
327. Pérez, E., et al., *Aquaporin expression in the cerebral cortex is increased at early stages of Alzheimer disease*. Brain Res, 2007. **1128**(1): p. 164-74.
328. Moftakhar, P., et al., *Aquaporin expression in the brains of patients with or without cerebral amyloid angiopathy*. J Neuropathol Exp Neurol, 2010. **69**(12): p. 1201-9.
329. Nicaise, C., et al., *Aquaporin-4 overexpression in rat ALS model*. Anat Rec (Hoboken), 2009. **292**(2): p. 207-13.
330. Costa, C., et al., *Aquaporin 1 and aquaporin 4 overexpression in bovine spongiform encephalopathy in a transgenic murine model and in cattle field cases*. Brain Res, 2007. **1175**: p. 96-106.
331. Herrera, M. and J.L. Garvin, *Aquaporins as gas channels*. Pflugers Arch, 2011. **462**(4): p. 623-30.
332. Musa-Aziz, R., et al., *Relative CO₂/NH₃ selectivities of AQP1, AQP4, AQP5, AmtB, and RhAG*. Proc Natl Acad Sci U S A, 2009. **106**(13): p. 5406-11.
333. Miller, E.W., B.C. Dickinson, and C.J. Chang, *Aquaporin-3 mediates hydrogen peroxide uptake to regulate downstream intracellular signaling*. Proc Natl Acad Sci U S A, 2010. **107**(36): p. 15681-6.
334. Ji, K.A., et al., *Differential neutrophil infiltration contributes to regional differences in brain inflammation in the substantia nigra pars compacta and cortex*. Glia, 2008. **56**(10): p. 1039-47.
335. Chen, Y. and R.A. Swanson, *Astrocytes and brain injury*. J Cereb Blood Flow Metab, 2003. **23**(2): p. 137-49.
336. Nagelhus, E.A., T.M. Mathiisen, and O.P. Ottersen, *Aquaporin-4 in the central nervous system: cellular and subcellular distribution and coexpression with KIR4.1*. Neuroscience, 2004. **129**(4): p. 905-13.
337. McGeer, P.L., K. Yasojima, and E.G. McGeer, *Association of interleukin-1 beta polymorphisms with idiopathic Parkinson's disease*. Neurosci Lett, 2002. **326**(1): p. 67-9.
338. Kohutnicka, M., et al., *Microglial and astrocytic involvement in a murine model of Parkinson's disease induced by 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)*. Immunopharmacology, 1998. **39**(3): p. 167-80.

339. Kress, B.T., et al., *Impairment of paravascular clearance pathways in the aging brain*. Ann Neurol, 2014. **76**(6): p. 845-61.
340. Zhang, J., et al., *Aquaporin-4 deficiency diminishes the differential degeneration of midbrain dopaminergic neurons in experimental Parkinson's disease*. Neurosci Lett, 2016. **614**: p. 7-15.
341. Fan, Y., et al., *Hypersensitivity of aquaporin 4-deficient mice to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine and astrocytic modulation*. Neurobiol Aging, 2008. **29**(8): p. 1226-36.
342. Sun, H., et al., *Aquaporin-4 mediates communication between astrocyte and microglia: Implications of neuroinflammation in experimental Parkinson's disease*. Neuroscience, 2016. **317**: p. 65-75.
343. García-Domínguez, I., et al., *Peripheral Inflammation Enhances Microglia Response and Nigral Dopaminergic Cell Death in an in vivo MPTP Model of Parkinson's Disease*. Front Cell Neurosci, 2018. **12**: p. 398.
344. Schiemann, J., et al., *K-ATP channels in dopamine substantia nigra neurons control bursting and novelty-induced exploration*. Nat Neurosci, 2012. **15**(9): p. 1272-80.
345. Bolam, J.P. and E.K. Pissadaki, *Living on the edge with too many mouths to feed: why dopamine neurons die*. Mov Disord, 2012. **27**(12): p. 1478-83.
346. Dragicevic, E., J. Schiemann, and B. Liss, *Dopamine midbrain neurons in health and Parkinson's disease: emerging roles of voltage-gated calcium channels and ATP-sensitive potassium channels*. Neuroscience, 2015. **284**: p. 798-814.
347. Liss, B., et al., *K-ATP channels promote the differential degeneration of dopaminergic midbrain neurons*. Nat Neurosci, 2005. **8**(12): p. 1742-51.
348. Vivekanantham, S., et al., *Neuroinflammation in Parkinson's disease: role in neurodegeneration and tissue repair*. Int J Neurosci, 2015. **125**(10): p. 717-25.
349. Farina, C., F. Aloisi, and E. Meinl, *Astrocytes are active players in cerebral innate immunity*. Trends Immunol, 2007. **28**(3): p. 138-45.
350. Lee, H.J., et al., *Direct transfer of alpha-synuclein from neuron to astroglia causes inflammatory responses in synucleinopathies*. J Biol Chem, 2010. **285**(12): p. 9262-72.
351. Hammond, S.L., et al., *NF- κ B signaling in astrocytes modulates brain inflammation and neuronal injury following sequential exposure to manganese and MPTP during development and aging*. Toxicol Sci, 2020.
352. Domercq, M., N. Vázquez-Villoldo, and C. Matute, *Neurotransmitter signaling in the pathophysiology of microglia*. Front Cell Neurosci, 2013. **7**: p. 49.
353. Gupta, M., et al., *Aged mice are more sensitive to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine treatment than young adults*. Neurosci Lett, 1986. **70**(3): p. 326-31.
354. Gupta, R.K. and M. Kanungo, *Glial molecular alterations with mouse brain development and aging: up-regulation of the Kir4.1 and aquaporin-4*. Age (Dordr), 2013. **35**(1): p. 59-67.
355. Liang, D., et al., *Cytotoxic edema: mechanisms of pathological cell swelling*. Neurosurg Focus, 2007. **22**(5): p. E2.
356. Simard, J.M., et al., *Brain oedema in focal ischaemia: molecular pathophysiology and theoretical implications*. Lancet Neurol, 2007. **6**(3): p. 258-68.
357. Forbes, K.P., J.G. Pipe, and J.E. Heiserman, *Evidence for cytotoxic edema in the pathogenesis of cerebral venous infarction*. AJNR Am J Neuroradiol, 2001. **22**(3): p. 450-5.

358. Shi, W.-Z., et al., *Aquaporin-4 deficiency attenuates acute lesions but aggravates delayed lesions and microgliosis after cryoinjury to mouse brain*. Neuroscience Bulletin, 2012. **28**(1): p. 61-68.
359. Werner, C.J., et al., *Proteome analysis of human substantia nigra in Parkinson's disease*. Proteome Sci, 2008. **6**: p. 8.
360. Noelker, C., et al., *Toll like receptor 4 mediates cell death in a mouse MPTP model of Parkinson disease*. Sci Rep, 2013. **3**: p. 1393.
361. Fiebich, B.L., et al., *Role of Microglia TLRs in Neurodegeneration*. Frontiers in Cellular Neuroscience, 2018. **12**(329).
362. Hirt, L., et al., *Increase of aquaporin 9 expression in astrocytes participates in astrogliosis*. J Neurosci Res, 2018. **96**(2): p. 194-206.
363. Dauer, W. and S. Przedborski, *Parkinson's disease: mechanisms and models*. Neuron, 2003. **39**(6): p. 889-909.
364. Hubbard, J.A., et al., *Expression of the Astrocyte Water Channel Aquaporin-4 in the Mouse Brain*. ASN Neuro, 2015. **7**(5).
365. Han, X., et al., *Changes in the Expression of AQP4 and AQP9 in the Hippocampus Following Eclampsia-Like Seizure*. Int J Mol Sci, 2018. **19**(1).
366. Tsukaguchi, H., et al., *Molecular characterization of a broad selectivity neutral solute channel*. J Biol Chem, 1998. **273**(38): p. 24737-43.
367. Ton, T.G., et al., *Nonsteroidal anti-inflammatory drugs and risk of Parkinson's disease*. Mov Disord, 2006. **21**(7): p. 964-9.
368. Wahner, A.D., et al., *Nonsteroidal anti-inflammatory drugs may protect against Parkinson disease*. Neurology, 2007. **69**(19): p. 1836-42.
369. Becker, C., S.S. Jick, and C.R. Meier, *NSAID use and risk of Parkinson disease: a population-based case-control study*. Eur J Neurol, 2011. **18**(11): p. 1336-42.
370. Ren, L., et al., *Nonsteroidal anti-inflammatory drugs use and risk of Parkinson disease: A dose-response meta-analysis*. Medicine (Baltimore), 2018. **97**(37): p. e12172.
371. Casu, M.A., et al., *Neuroprotection by the Immunomodulatory Drug Pomalidomide in the Drosophila LRRK2(WD40) Genetic Model of Parkinson's Disease*. Front Aging Neurosci, 2020. **12**: p. 31.

PAPER I-III

PAPER I

Subcellular expression of aquaporin-4 in substantia nigra of normal and MPTP-treated mice.

Prydz A, Stahl K, Puchades M, Davarpaneh N, Nadeem M, Ottersen OP, Gundersen V, Amiry-Moghaddam M. *Neuroscience*. 2017; Sep 17;359:258-266. doi: 10.1016/j.neuroscience.2017.07.029.

SUBCELLULAR EXPRESSION OF AQUAPORIN-4 IN SUBSTANTIA NIGRA OF NORMAL AND MPTP-TREATED MICE

AGNETE PRYDZ,^a KATJA STAHL,^a MAJA PUCHADES,^a
NINA DAVARPANEH,^a MARIA NADEEM,^a
OLE PETTER OTTERSEN,^a VIDAR GUNDERSEN^{a,b} AND
MAHMOOD AMIRY-MOGHADDAM^{a*}

^a Division of Anatomy and Healthy Brain Ageing Center Regional Research Network, Department of Molecular Medicine, Institute of Basic Medical Sciences, University of Oslo, PO Box 1105, Blindern, 0317 Oslo, Norway

^b Department of Neurology, Oslo University Hospital, Rikshospitalet, PO Box 4950 Nydalen, 0424 Oslo, Norway

Abstract—Aquaporin-4 (AQP4) is the predominant water channel in mammalian CNS where it is localized at the perivascular astrocytic foot processes abutting brain microvessels. Several lines of evidence suggest that AQP4 is involved in important homeostatic functions and that mislocalization of the perivascular pool of AQP4 is implicated in several different brain disorders. A recent study suggests that the differential susceptibility of midbrain dopaminergic neurons to the parkinsonogenic toxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) depends on the expression of AQP4. Further, MRI studies of patients with Parkinson's disease (PD) point to an excessive water accumulation in the substantia nigra (SN). This prompted us to investigate the cellular and subcellular distribution of AQP4 in mouse SN using immunofluorescence and quantitative immunogold cytochemistry. Compared with neocortex, SN exhibits a higher concentration of AQP4. Specifically, judged by electron microscopic immunogold analysis, the perivascular density of AQP4 in SN exceeds by 70% the perivascular density of AQP4 in the neocortex. An even larger difference in AQP4 labeling was found for astrocytic processes in the neuropil. Treatment with MPTP further increased (by >30%) the perivascular AQP4 density in SN, but also increased AQP4 labeling in the neocortex. Our data indicate that the perivascular AQP4 pool in SN is high in normal animals and even higher after treatment with MPTP. This would leave the SN more prone to water accumulation and supports the idea that AQP4 could be involved in the pathogenesis of PD. © 2017 IBRO. Published by Elsevier Ltd. All rights reserved.

Key words: aquaporin-4, astrocyte, MPTP, Parkinson's disease, immunogold histochemistry.

*Corresponding author.

E-mail address: mahmo@medisin.uio.no (M. Amiry-Moghaddam).

Abbreviations: AQP4, aquaporin-4; DA, dopaminergic; GFAP, glial fibrillary acidic protein; MPTP, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; PD, Parkinson's disease; SN, substantia nigra; SNpc, substantia nigra pars compacta; TH, tyrosine hydroxylase.

<http://dx.doi.org/10.1016/j.neuroscience.2017.07.029>

0306-4522/© 2017 IBRO. Published by Elsevier Ltd. All rights reserved.

INTRODUCTION

Parkinson's disease (PD) is the second most common neurodegenerative disorder in the Western hemisphere, characterized by progressive neuronal cell death predominantly of the dopaminergic neurons (DA) in substantia nigra pars compacta (SNpc) (Hirsch et al., 2013). More than 90% of all cases with PD have unknown etiology and the mechanisms underlying the selective loss of nigral DA neurons are not understood. Evidence is accumulating to suggest that astrocytes might play a role in the pathophysiology of PD (Mena and García De Yébenes, 2008) and recent studies in animal models of PD have indicated an involvement of the astrocytic water channel aquaporin-4 (AQP4) (Fan et al., 2008; Yang et al., 2011; Sun et al., 2016; Zhang et al., 2016). Moreover, MRI studies of patients with PD have revealed an abnormal water accumulation in the SN (Ofori et al., 2015a). However, the cellular and subcellular expression of AQP4 in the SN is unknown, as is the expression level of AQP4 in the parkinsonian brain.

Studies of other brain regions have revealed that AQP4 is concentrated in astrocytic endfeet that encompass the blood–brain and brain–liquid interfaces (Nielsen et al., 1997; Rash et al., 1998; Amiry-Moghaddam and Ottersen, 2003). The polarized expression of AQP4 in astrocytes is crucial for homeostatic processes maintaining normal function, including water and potassium homeostasis (Nagelhus et al., 1999). Further, polarized expression of AQP4 has been shown to be important for cell migration (Saadoun et al., 2005) and possibly astrocytic Ca^{2+} signaling (Thrane et al., 2011). Loss of the polarized expression of AQP4 has been shown in several pathological conditions, such as stroke (De Castro Ribeiro et al., 2006; Frydenlund et al., 2006), temporal lobe epilepsy (Alvestad et al., 2013) and Alzheimer's disease (Yang et al., 2011). Here we used immune-microscopy in healthy mice and mice treated with the parkinsonogenic toxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) to ask whether the SN differs from other brain regions with respect to the expression of AQP4 and whether the expression of AQP4 in SN is sensitive to MPTP.

Our data indicate that astrocytes in the SN differ from those in neocortex by showing a higher level of AQP4, particularly in those endfoot membrane domains that mediate water exchange between brain and blood. Nine

days after treatment with MPTP the AQP4 pool in these membrane domains has increased further, and more so than the AQP4 pools in other domains of the astrocyte membrane. AQP4 over expression in specific astrocyte membrane domains may contribute to the water dyshomeostasis observed in patients with PD.

EXPERIMENTAL PROCEDURES

Animals

Male wild-type C57BL/6 mice 8–12 weeks were used in this study. Experimental protocols were approved by the Institutional Animal Care and Use Committee and conform to National Institutes of Health guidelines for the care and use of animals.

Tissue preparation

Untreated animals. Animals were anesthetized deeply with hypnorm dormicum (0.105 mg/kg) followed by intracardial perfusion. For light microscopy of untreated SN and neocortex, the animals ($n = 5$) were perfusion-fixed with 4% formaldehyde (FA) in 0.1 M phosphate buffer (PB) (Sigma–Aldrich, St. Louis, MO, USA) using pH-shift (Eilert-Olsen et al., 2012) starting with pH 6.0 for 5 min, then shifting to pH 10.5 for 10 min. The brains were dissected out and post-fixed overnight in 4% FA with pH 10.5. Animals used for electron microscopy ($n = 3$) were perfusion-fixed with 4% FA and 0.1% glutaraldehyde (GA) in 0.1 PB (Sigma–Aldrich), and then post-fixed in the same solution overnight. All brains were stored in a 1:10 dilution of their respective fixative at 4°C until further preparations.

MPTP- and saline-treated animals. Mice were treated with either MPTP or saline, following an acute protocol as previously described (Jackson-Lewis et al., 1995; Przedborski and Jackson-Lewis, 2007; Puchades et al., 2013). One series of MPTP-treated mice and controls was used for EM studies and another series was used for light microscopic immunohistochemistry. For EM experiments, male C57 BL/6 mice ($n = 5$; 8 weeks of age; Taconic, Denmark) received MPTP–HCl free base (Sigma–Aldrich) in an acute paradigm (four injections subcutaneously at 2-h intervals, 18 mg/kg for each injection) (Przedborski and Jackson-Lewis, 2007). The control mice ($n = 5$) received injections of saline (4 mL/kg). For light microscopic immunohistochemistry, male C57 BL/6 mice ($n = 4$; 12 weeks of age; Taconic, Denmark) received MPTP–HCl free base (Sigma–Aldrich) in an acute paradigm (three subcutaneous injections of 15 mg/kg at 2-h intervals). The control mice ($n = 3$) received injections of saline (4 mL/kg). The MPTP dosage was reduced in the second series in order to reduce the acute mortality associated with the higher MPTP concentration. Based on Jackson-Lewis et al. (Jackson-Lewis et al., 1995) who showed that the degeneration of dopamine neurons started within the first day after MPTP treatment, reached a plateau at day 4, and was thereafter stable until day 28, we chose a survival time of 9 days after the last

MPTP injection. The choice of this time point is also in line with the recommendations of Przedborski and Jackson-Lewis (2007). From day 7 until day 28 TH-immunoreactivity can be used to determine accurately the number of living neurons in SNpc (Jackson-Lewis et al., 1995). Nine days after the last injection, both groups of animals were deeply anesthetized with pentobarbital (0.1 mL/50 g body weight) followed by transcardial perfusion of either 4% FA and 0.1% GA or 4% FA for 10 min and post-fixed *in situ* overnight (Puchades et al., 2013). Tissue samples from the 4% FA-fixed brains were used for immunofluorescence analysis and samples from both fixatives were used for the immunogold analysis. The difference in the fixatives did not have any impact on the density of the immunogold labeling (data not shown). The brains were dissected out and either processed for electron microscopy or cut into 30 μ m coronal sections using a Vibratome (Microm, Jacksonville, FL, USA) for immunohistochemistry and stored at 4°C in 1/10 fixative until further preparations.

Immunofluorescence and confocal microscopy. The immunofluorescence labeling was performed as described previously (Yang et al., 2011). In brief, the sections were treated with 10% normal goat serum (NGS) and 1% bovine serum albumin (BSA) upon incubation with primary antibodies overnight in 3% NGS, 1% BSA, 0.5% Triton-X-100 and 0.05% sodium azide in 0.01 M phosphate-buffered saline (all chemicals from Sigma–Aldrich). Following washing, the sections were incubated with secondary antibodies, washed in PBS, and mounted with Prolong Gold Antifade Reagent with DAPI (Invitrogen, Norway). A complete list of the primary and secondary antibodies used in this study is presented in Table 1. Images were collected with a LSM 510 META Confocal Microscope (Zeiss, Jena, Germany) using a

Table 1. Antibodies

Method	Primary antibody	Secondary antibody
Immunogold	AQP4, host: rabbit, Sigma-Aldrich, diluted 1:400	Goat anti-rabbit 15 nm gold, Abcam, diluted 1:20
Immunofluorescence	AQP4, host: rabbit, Sigma-Aldrich, diluted 1:400	Cy3 donkey anti-rabbit, Jackson, diluted 1:500
	TUJ1, host: mouse, Covance, diluted 1:1000	Alexa 488 donkey anti-mouse, Invitrogen, diluted 1:500
	TH, host mouse, Chemicon, diluted 1:1000	Alexa 488 donkey anti-mouse, Invitrogen, diluted 1:500
	GFAP, host chicken, Nordic BioSite, diluted 1:500	Cy2 donkey anti-chicken, Jackson, diluted 1:500

Fig. 1. Distribution of AQP4 in substantia nigra (SN; A–D) and cerebral cortex (CX; E–H). Strong AQP4 immunofluorescent staining (red) is present in the astrocyte processes in neuropil (asterisk) and around blood vessels (white arrows) in SN (A). In the cortex, AQP4 expression is restricted to astrocytic endfeet facing blood vessels and the pial surface (white arrow head) (E). Tyrosine hydroxylase (TH; green) is used as a marker for dopaminergic neurons in SNpc (B), and tubulin class III (TuJ1; green) is used as a general neuronal marker in CX (F). Astrocytic processes with endfoot membranes abutting microvessels and the pial surface (white arrow head, G) are visualized by GFAP (orange). Merged images show co-localization of AQP4 with GFAP in both SN (D) and CX (H). SNpc; Substantia nigra pars compacta, and SNpr; Substantia nigra pars reticulata. $n = 5$, scale bar: 50 μm . (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

40 $\times$ oil objective. DAPI fluorescence was captured at 405 nm, Alexa 488 at 510 nm, Cy3 at 568 nm and Cy5 at 670 nm.

Electron microscopic immunogold cytochemistry. Tissue blocks of 1 $\times$ 1 mm from SNpc and parietal cortex

were dissected out using a dissection microscope (Olympus SZX12, Tokyo, Japan). The tissue blocks were subjected to freeze substitution procedure as previously described (van Lookeren Campagne et al., 1991) for the untreated animals and for the MPTP- and saline-treated animals. In brief, the tissue blocks were cryo-protected in glycerol and rapidly frozen in liquid propane at -170°C . The frozen tissue was immersed into anhydrous methanol containing 1.5% uranyl acetate at 90°C in an automatic freeze substitution unit (EMAFS, Leica, Vienna, Austria), infiltrated with Lowicryl HM20 Resin (Lowy, Waldkraiburg, Germany) at -30°C and polymerized by ultraviolet light. Ultrathin sectioning and immunogold labeling were performed as described previously (Hoddevik et al., 2016). Briefly, the blocks were cut into ultrathin sections of 90–100 nm using an ultramicrotome (Leica EM UC6, Vienna, Austria), transferred to mesh grids and incubated in drops of Tris-buffered saline (HCl 0.3%, NaCl 0.01%) with 0.1% triton (TBST) and 50 mM glycine for 10 min, washed in double-distilled water and then incubated for 10 min in TBST containing 2% w/v human serum albumin (HSA). The grids were then incubated in a solution of 2% w/v HSA in TBST and the primary antibody (Table 1) over night at room temperature. The labeling was visualized with a secondary antibody coupled to 15 nm colloidal gold particles (Table 1) diluted in 1:20 TBST with 2% w/v HSA and polyethylene glycol. The sections were contrasted with 1% uranyl acetate and 0.3% lead citrate for 1.5 min and rinsed in double-distilled water in between. The sections were finally examined with an electron microscope (Tecnai 12, FEI Company, Eindhoven, Netherlands). Random microvessels ($n = 25$) were selected from each section at low magnification and one electron microscopic image of the perivascular endfoot was acquired per capillary. For the analysis of the non-endfoot AQP4 immunogold labeling, 20 electron micrographs of the neuropil were randomly obtained throughout each section at 20,500 $\times$ magnification. The analyzer was blind to treatment and area.

Quantification and statistical analysis

Postembedding immunogold microscopy and quantification of AQP4 labeling. For the quantification of the AQP4 distribution in untreated and MPTP-/saline-treated SNpc and parietal cortex, images from capillaries were selected in a blinded manner from each section (one section for each animal) and quantified with an image analysis program, analySIS Pro version 3.2 (Soft Imaging System GmbH, Munster, Germany) for the untreated sections and analySIS IMGAP version Beta cell[®]p (Soft Imaging Systems GmbH, Munster, Germany) for the MPTP-/saline-treated sections. Larger vessels and distorted or mechanically damaged basal lamina were excluded, as well as non-neuropil components such as neurons and vessels for the neuropil electron micrographs. Membrane segments of interest and membrane curves were drawn in the overlay interactively; the membrane segment of the astrocyte endfoot facing the capillary lumen was called

Fig. 2. Immunogold labeling and quantitative analysis of AQP4-labeling in substantia nigra (SN) and neocortex (CX) in untreated animals. The AQP4 immunogold labeling at the perivascular astrocyte membrane domains (arrow) is stronger in the SN (A) than in the CX (B). Quantitative analysis of the perivascular immunogold labeling shows a significantly higher mean linear density of gold particles in the SN, compared to the CX (*** $p < 0.001$) (C). Non-endfoot membranes show significantly higher density of AQP4 immunogold particles in the SN compared to CX (*** $p < 0.001$) (D). $n = 3$, scale bar: 1 μm.

adluminal and the membrane segment facing neuropil was called abluminal. Both membrane segments were analyzed in the MPTP-/saline-treated sections and only the adluminal segment was analyzed in the untreated sections. Gold particles near each membrane curve were detected semi-automatically, and the distance between each particle and its membrane curve was calculated by the program. Particles localized within 23.5 nm from their membrane curve (Amiry-Moghaddam and Ottersen, 2013) were included in the calculation of the number of particles per μm of membrane (linear particle density). In the neuropil area, gold particles per μm² were calculated by the program after drawing an area of interest in the overlay for each section. The non-endfoot astrocyte processes in the neuropil were called non-end. Groups were analyzed statistically by Student's t -test (two-tailed) using SPSS version 18.0 for the untreated sections and version 22.0 for the MPTP-/saline-treated sections. The significance level was set to 0.05.

RESULTS

AQP4 in normal, untreated animals

Immunofluorescence analysis demonstrated highly polarized expression of AQP4 around blood vessels in SNpc (Fig. 1A–D) and neocortex (Fig. 1E–H). In neocortex, AQP4 is scarce in the neuropil and around profiles positive for the neuronal marker TuJ1 (Fig. 1E–H).

In the SNpc, AQP4 is abundantly expressed with a particularly pronounced labeling in perivascular astrocyte processes. AQP4 immunolabeling is also found in non-endfoot membranes in the neuropil. A similar pattern occurs in the pars reticulata, where the density of tyrosine hydroxylase (TH)-positive cells is lower and the density of glial fibrillary acidic protein (GFAP)-positive cells is higher than in the pars compacta (Fig. 1A–D).

In line with the immunofluorescence labeling, electron micrograph images demonstrated a particularly

Fig. 3. Expression of AQP4, TH-positive cells and GFAP in SN in the MPTP mouse model of PD (A–D) compared to the saline control (E–H). Subcutaneous injections of MPTP resulted in strong upregulation of AQP4 (red) in the SN, both perivascularly (white arrow) and in non-endfoot (asterisk) membranes (A). Further, the toxin induced dopaminergic cell death, as judged by loss of TH-positive cells (green, B). Upregulation of GFAP (orange) was also observed, indicating astrogliosis (C, D). This reaction was not seen in control animals injected with saline (G, H). MPTP, $n = 4$; saline, $n = 3$. Scale bar: 50 μm . (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

dense immunolabeling of AQP4 in the SN (Fig. 2A) compared to neocortex (Fig. 2B). AQP4 immunogold particles were concentrated in perivascular endfoot membranes, and were also numerous in fine astrocytic processes in the neuropil. Compared to the neocortex, the SN showed stronger AQP4 immunolabeling both perivascularly (Fig. 2C) and in neuropil (Fig. 2D). In the SN, perivascular labeling was 70% higher than in the neocortex, with a mean linear density of 28.4 ± 1.0 gold

particles/ μm compared to 16.8 ± 0.7 gold particles/ μm in the neocortex ($p < 0.001$). Non-endfoot labeling was measured as areal density of gold particles in randomly selected images from neuropil. In the SN, density of AQP4 immunogold particles was 12.7 ± 1.6 gold particles/ μm^2 , compared to 5.5 ± 0.5 gold particles/ μm^2 in the neocortex ($p < 0.001$) (values are mean $\pm$ SEM). These results complement and extend the immunofluorescence data.

AQP4 in MPTP-treated animals

MPTP induces dopaminergic cell death, as seen by loss of TH-positive cells. The method of quantification and the level of degeneration of TH-positive neurons in the SNpc in the MPTP-treated mice were previously described (Puchades et al., 2013). MPTP treatment led to a 42% reduction in the density of TH-positive cell bodies (Puchades et al., 2013). Immunofluorescence analysis of SN revealed that MPTP injections led to strong upregulation of AQP4, both perivascularly and in non-endfoot membranes (Fig. 3A,D). Upregulation of GFAP was also observed, indicating reactive astrogliosis (Fig. 3C–D). This is consistent with previous findings (Puchades et al., 2013). No change was found in control animals injected with saline (Fig. 3E–H).

In accordance with the immunofluorescence labeling, immunoelectron microscopy of SNpc of MPTP-treated animals showed an increased expression of AQP4 both in the perivascular endfoot membranes and in the non-endfoot membranes in the neuropil (Fig. 4).

The quantitative analysis of MPTP-treated animals revealed in SNpc a significantly higher expression of AQP4 in the *adluminal* as well as the *abluminal* membranes of perivascular endfeet. The *adluminal* labeling was 30% higher in MPTP-treated animals than in saline controls, with a mean linear density of 30.3 ± 1.21 gold particles/ μm and 23.5 ± 1.05 ($p < 0.05$), respectively (Fig. 5A). The *abluminal* labeling was 34% higher in the MPTP-treated animals than in the saline controls with a mean linear density of 15.3 ± 1.02 gold particles/ μm compared to 11.4 ± 0.71 in the saline controls ($p < 0.01$) (Fig. 5A). In line with the immunofluorescence data, the quantitative immunogold analysis revealed an increased expression of AQP4 in the non-endfoot astrocytic membranes following MPTP injections. The neuropil labeling was 21% higher in the animals treated with MPTP, with a mean linear density of 6.3 ± 0.36 gold particles/ μm^2 compared to 5.2 ± 0.28 ($p = 0.019$) in saline controls (Fig. 5B).

Administration of MPTP also caused a modest increase in perivascular AQP4 immunolabeling in the neocortex. The changes were restricted to the *adluminal* membrane domains (Fig. 5C).

DISCUSSION

Information is scant on the microscopic and fine structural localization of AQP4 in the SN. This void must be filled, as aquaporins and water homeostasis have become increasingly relevant in the context of PD. Notably, experimental data suggest that the toxin susceptibility of

Fig. 4. Immunogold labeling of AQP4 in SN in animals exposed to MPTP (A, B) and saline injections (C, D). Electron micrograph imaging showed increased expression of AQP4 labeling in astrocytic perivascular endfoot membranes (A), both in the abluminal (black arrow) and in the abluminal (black arrow head) membrane, as well as in the astrocytic non-endfoot membranes in the neuropil (B, black arrow) compared to saline controls (C, D). L = lumen, E = endothelial cell. $n = 5$ for each experimental group, scale bar: 1 μm .

dopaminergic neurons depends on the level of AQP4 expression (Zhang et al., 2016) and a recent report based on a diffusion MRI technique has shown increased levels of free water in SNpc of patients with PD (Ofori et al., 2015b). There was a positive correlation between Montreal Cognitive Assessment scores and free water values in SNpc (Ofori et al., 2015a, 2015b).

Our study is the first detailed study of AQP4 in the SN and shows that the perivascular AQP4 pool – which is

relatively large even in normal animals – is further increased after treatment with MPTP. This suggests an involvement of AQP4 in PD and may explain why PD patients exhibit water accumulation in SN. These data contrast with findings in Alzheimer's disease, stroke and epilepsy, in which the perivascular AQP4 pool is reduced (Frydenlund et al., 2006; Yang et al., 2011; Alvestad et al., 2013). After experimental stroke, the endfoot pool of AQP4 is unaltered or increased in the initial phase, but then decreases sharply to very low levels at ~24 h of the ischemic episode (Frydenlund et al., 2006; De Castro Ribeiro et al., 2006).

Our results show that AQP4 is more plentiful in the SN than in the neocortex, both in endfoot membranes encompassing microvessels and in non-endfoot astrocytic membranes in the neuropil. This high density of AQP4 was further increased following injections of MPTP. Our conclusions were based on a postembedding immunogold technique that has been modified so as to enable a precise localization and quantification of membrane-bound AQP4 (Takumi et al., 1998; Amiry-Moghaddam and Ottersen, 2013).

Attempts to identify mechanisms underlying the selective vulnerability of dopaminergic cells in the SNpc have focused primarily on the neurons. Although astrocytes are known to be essential in maintenance of homeostatic processes and neuronal function, little is known about their roles in the pathophysiology of PD. The density of astrocytes in the SNpc is low compared to that in other regions (Ji et al., 2008). As astrocytes release a variety of growth factors under normal conditions (Swanson et al., 2002), one could ask if this low density of astrocytes in SNpc would contribute

to the susceptibility of local dopaminergic neurons. Another question is whether nigral astrocytes differ from other astrocytes when it comes to their roles in maintaining key homeostatic functions. The present analysis of AQP4 was prompted by this question.

To shed light on the potential role of astrocytes and AQP4 in the pathophysiology of PD, we studied AQP4 expression pattern in an acute MPTP mouse model of PD. MPTP induces a parkinsonian syndrome by

Fig. 5. Quantitative immunogold analysis of AQP4 in the SN and CX of animals injected with MPTP ($n = 5$) compared to animals injected with saline ($n = 5$; A–C). Quantitative analysis of the immunogold labeling at the perivascular astrocytic endfoot membrane domains in SN revealed a significantly higher linear density of AQP4 labeling in both the adluminal ($p < 0.05$) and abluminal ($p < 0.01$) membrane domains of the MPTP-treated animals compared to the saline controls (A). The AQP4 immunogold labeling of the non-endfoot astrocyte processes in the SN is significantly higher in the MPTP-treated mice compared to the saline controls (B, $p < 0.05$). Immunogold labeling of AQP4 in the CX shows a small but significant increase in the perivascular astrocytic adluminal membrane domains ($p = 0.041$). No significant alteration in the AQP4 expression was detected in the astrocytic abluminal membrane domains (C).

inactivating complex I of the electron transfer chain (Nicklas et al., 1987). It has also been reported that MPTP treatment leads to astrogliosis in SNpc, as well as in striatum (Puchades et al., 2013). In this respect, it should be noted that the death of dopamine neurons increases between days 1 and 4 after MPTP treatment, and the number of degenerated neurons is thereafter stable for at least one month (Jackson-Lewis et al., 1995). Astrogliosis follows a similar time course; starting at day 1 and increasing until day 7 after MPTP, when it reaches a plateau where the number of reactive astrocytes is more or less stable for one month (Hou et al., 2017).

Our study revealed a general upregulation of AQP4 expression in SNpc in animals treated with MPTP. The high resolution immunogold analysis allowed us to distinguish between different subcellular AQP4 pools and identified a pronounced increase in the density of AQP4 in the astrocytic endfoot membranes that enwrap capillaries and microvessels. This perivascular pool of AQP4 is the one that is directly engaged in water exchange between blood and brain and that controls the influx of water in pathological conditions such as stroke and hyponatremic brain edema (Amiry-Moghaddam et al., 2003, 2004). An increased perivascular pool of AQP4 would make the SN more prone to water accumulation – thus providing a possible mechanistic explanation for the abnormal water contents seen in SN of patients with PD (Ofori et al., 2015a).

The present data call for further studies to disentangle the possible coupling between astrocytic dysfunction and the propensity for developing PD – be it in its idiopathic form or in response to known parkinsonogenic drugs. A number of neurological conditions have been associated with perturbations of water and ion homeostasis, and mislocalization of astrocytic AQP4 occurs in human temporal lobe epilepsy as well as in animal models of stroke and Alzheimer's disease (Eid et al., 2005; Yang et al., 2011). In the kainate model of epilepsy, mislocalization of AQP4 was found to precede the development of overt seizures, suggesting a role in pathogenesis (Alvestad et al., 2013). In terms of AQP4 mislocalization, PD differs from the above conditions in showing an increase, rather than a decrease, in AQP4 in perivascular membranes.

It should be emphasized that the increase in the expression of AQP4 which is observed in our study cannot be explained solely by gliosis. Indeed, gliosis is observed in the sclerotic hippocampus of patients with mesial temporal lobe epilepsy and in mouse models of Alzheimer's disease. In both conditions gliosis is associated with a loss of rather than an increase in endfoot AQP4 (Eid et al., 2005; Yang et al., 2011). Also in a model of cerebral malaria there is an overall decrease in AQP4 expression, despite extensive gliosis (Promeneur et al., 2013). Further studies are needed to unravel the mechanistic link between AQP4 expression

and proneness to PD and – in a longer time perspective – to ascertain whether astrocytes could serve as targets of new therapies.

Acknowledgments—The authors would like to thank Bjørge Riber, Karen Marie Gujord, Jorunn Knutsen and Bashir Hakim for technical assistance and Gunnar Lothe for graphics. This study was supported by M.D. Ph.D Curriculum of the medical faculty at the University of Oslo – Norway.

REFERENCES

- Alvestad S, Hammer J, Hoddevik EH, Skare Ø, Sonnewald U, Amiry-Moghaddam M, Ottersen OP (2013) Mislocalization of AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy. *Epilepsy Res* 105(1–2):30–41. <http://dx.doi.org/10.1016/j.eplepsyres.2013.01.006>.
- Amiry-Moghaddam M, Ottersen OP (2003) The molecular basis of water transport in the brain. *Nat Rev Neurosci* 4(12):991. <http://dx.doi.org/10.1038/nrn1252>.
- Amiry-Moghaddam M, Ottersen OP (2013) Immunogold cytochemistry in neuroscience. *Nat Neurosci* 16(7):798–804. <http://dx.doi.org/10.1038/nn.3418>.
- Amiry-Moghaddam M, Otsuka T, Hurn PD, Traaystman RJ, Haug F-M, Froehner SC, Adams ME, Neely JD, Agre P, Ottersen OP, Bhardwaj A (2003) An alpha-syntrophin-dependent pool of AQP4 in astroglial end-feet confers bidirectional water flow between blood and brain. *Proc Natl Acad Sci USA* 100(4):2106.
- Amiry-Moghaddam M, Xue R, Haug F-M, Neely JD, Bhardwaj A, Agre P, Adams ME, Froehner SC, Mori S, Ottersen OP (2004) Alpha-syntrophin deletion removes the perivascular but not endothelial pool of aquaporin-4 at the blood-brain barrier and delays the development of brain edema in an experimental model of acute hyponatremia. *FASEB J* 18(3):542.
- De Castro Ribeiro M, Hirt L, Bogousslavsky J, Regli L, Badaut J (2006) Time course of aquaporin expression after transient focal cerebral ischemia in mice. *J Neurosci Res* 83(7):1231–1240. <http://dx.doi.org/10.1002/jnr.20819>.
- Eid T, Lee TS, Thomas MJ, Amiry-Moghaddam M, Bjornsen LP, Spencer DD, Agre P, Ottersen OP, de Lanerolle NC (2005) Loss of perivascular aquaporin 4 may underlie deficient water and K⁺ homeostasis in the human epileptogenic hippocampus. *Proc Natl Acad Sci USA* 102(4):1193. <http://dx.doi.org/10.1073/pnas.0409308102>.
- Eilert-Olsen M, Haj-Yasein NN, Vindedal GF, Enger R, Gundersen GA, Hoddevik EH, Petersen PH, Haug F-M, Skare Ø, Adams ME, Froehner SC, Burkhardt JM, Thoren AE, Nagelhus EA (2012) Deletion of aquaporin-4 changes the perivascular glial protein scaffold without disrupting the brain endothelial barrier. *Glia* 60(3):432. <http://dx.doi.org/10.1002/glia.22277>.
- Fan Y, Kong H, Shi X, Sun X, Ding J, Wu J, Hu G (2008) Hypersensitivity of aquaporin 4-deficient mice to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine and astrocytic modulation. *Neurobiol Aging* 29(8):1226–1236. <http://dx.doi.org/10.1016/j.neurobiolaging.2007.02.015>.
- Frydenlund DS, Bhardwaj A, Otsuka T, Mylonakou MN, Yasumura T, Davidson KGV, Zeynalov E, Skare Ø, Laake P, Haug F-M, Rash JE, Agre P, Ottersen OP, Amiry-Moghaddam M (2006) Temporary loss of perivascular aquaporin-4 in neocortex after transient middle cerebral artery occlusion in mice. *Proc Natl Acad Sci USA* 103(36):13532–13536. doi:www.pnas.org/cig/doi/10.1073/pnas.0605796103.
- Hirsch EC, Jenner P, Przedborski S (2013) Pathogenesis of Parkinson's Disease. *Mov Disord* 28. <http://dx.doi.org/10.1002/mds.25032>.
- Hoddevik EH, Khan FH, Rahmani S, Ottersen OP, Boldt HB, Amiry-Moghaddam M (2016) Factors determining the density of AQP4 water channel molecules at the brain–blood interface. *Brain Struct Funct* 1–14. <http://dx.doi.org/10.1007/s00429-016-1305-y>.
- Hou L, Zhou X, Zhang C, Wang K, Liu X, Che Y, Sun F, Li H, Wang Q, Zhang D, Hong J-S (2017) NADPH oxidase-derived H₂O₂ mediates the regulatory effects of microglia on astrogliosis in experimental models of Parkinson's disease. *Redox Biol*. <http://dx.doi.org/10.1016/j.redox.2017.02.016>.
- Jackson-Lewis V, Jakowec M, Burke RE, Przedborski S (1995) Time course and morphology of dopaminergic neuronal death caused by the neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *Neurodegeneration* 4(3):257–269. [http://dx.doi.org/10.1016/1055-8330\(95\)90015-2](http://dx.doi.org/10.1016/1055-8330(95)90015-2).
- Ji K, Eu M, Kang S, Gwag B, Jou I, Joe E (2008) Differential neutrophil infiltration contributes to regional differences in brain inflammation in the substantia nigra pars compacta and cortex. *Glia* 56(10):1039–1047. <http://dx.doi.org/10.1002/glia.20677>.
- Mena MA, Garcia De Yébenes J (2008) Glial cells as players in parkinsonism: the “good,” the “bad,” and the “mysterious” glia. *Neuroscientist* 14(6):544–560. <http://dx.doi.org/10.1177/1073858408322839>.
- Nagelhus EA, Horio Y, Inanobe A, Fujita A, Haug FM, Nielsen S, Kurachi Y, Ottersen OP (1999) Immunogold evidence suggests that coupling of K⁺ siphoning and water transport in rat retinal Müller cells is mediated by a coenrichment of Kir4.1 and AQP4 in specific membrane domains. *Glia* 26(1):47–54. [http://dx.doi.org/10.1002/\(sici\)1098-1136\(199903\)26:1<47::aid-glia5>3.0.co;2-5](http://dx.doi.org/10.1002/(sici)1098-1136(199903)26:1<47::aid-glia5>3.0.co;2-5).
- Nicklas WJ, Youngster SK, Kindt MV, Heikkila RE (1987) MPTP, MPP⁺ and mitochondrial function. *Life Sci* 40(8):721.
- Nielsen S, Nagelhus EA, Amiry-Moghaddam M, Bourque C, Agre P, Ottersen OP (1997) Specialized membrane domains for water transport in glial cells: high-resolution immunogold cytochemistry of aquaporin-4 in rat brain. *J Neurosci* 17(1):171.
- Ofori E, Pasternak O, Planetta PJ, Burciu R, Snyder A, Febo M, Golde TE, Okun MS, Vaillancourt DE (2015) Increased free water in the substantia nigra of Parkinson's disease: a single-site and multi-site study. *Neurobiol Aging* 36(2):1097–1104. <http://dx.doi.org/10.1016/j.neurobiolaging.2014.10.029>.
- Ofori E, Pasternak O, Planetta PJ, Li H, Burciu RG, Snyder AF, Lai S, Okun MS, Vaillancourt DE (2015) Longitudinal changes in free-water within the substantia nigra of Parkinson's disease. *Brain* 138(8):2322–2331. <http://dx.doi.org/10.1093/brain/awv136>.
- Promeneur D, Lunde LK, Amiry-Moghaddam M, Agre P (2013) Protective role of brain water channel AQP4 in murine cerebral malaria. *Proc Natl Acad Sci USA* 110(3):1035–1040. <http://dx.doi.org/10.1073/pnas.1220566110>.
- Puchades M, Sogn CJ, Maehlen J, Bergersen LH, Gundersen V (2013) Unaltered lactate and glucose transporter levels in the MPTP mouse model of Parkinson's disease. *J Parkinson's Dis* 3(3):371. <http://dx.doi.org/10.3233/jpd-130190>.
- Rash JE, Yasumura T, Hudson CS, Agre P, Nielsen S (1998) Direct immunogold labeling of aquaporin-4 in square arrays of astrocyte and ependymocyte plasma membranes in rat brain and spinal cord. *Proc Natl Acad Sci USA* 95(20):11981–11986.
- Saadoun S, Papadopoulos MC, Hara-Chikuma M, Verkman AS (2005) Impairment of angiogenesis and cell migration by targeted aquaporin-1 gene disruption. *Nature* 434(7034):786–792. <http://dx.doi.org/10.1038/nature03460>.
- Sun H, Liang R, Yang B, Zhou Y, Liu M, Fang F, Ding J, Fan Y, Hu G (2016) Aquaporin-4 mediates communication between astrocyte and microglia: implications of neuroinflammation in experimental Parkinson's disease. *Neuroscience* 317:65–75. <http://dx.doi.org/10.1016/j.neuroscience.2016.01.003>.
- Swanson RA, Ying W, Alano CC, Chen Y (2002) PARP-1 activation causes mitochondrial dysfunction. *J Neurochem* 81:108–111. <http://dx.doi.org/10.1046/j.1471-4159.81.s1.42.5.x>.
- Takumi Y, Nagelhus E, Eidet J, Matsubara A, Usami S, Shinkawa H, Nielsen S, Ottersen OP (1998) Select types of supporting cell in the inner ear express aquaporin-4 water channel protein. *Eur J Neurosci* 10(12):3584–3595.
- Thrane AS, Rappold PM, Fujita T, Torres A, Bekar LK, Takano T, Peng W, Wang F, Rangroo Thrane V, Enger R, Haj-Yasein NN, Skare Ø, Holen T, Klungland A, Ottersen OP, Nedergaard M,

- Nagelhus EA (2011) Critical role of aquaporin-4 (AQP4) in astrocytic Ca²⁺ + signaling events elicited by cerebral edema. *Proc Natl Acad Sci USA* 108(2):846. <http://dx.doi.org/10.1073/pnas.1015217108>.
- van Lookeren Campagne M, Oestreicher AB, van Der Krift TP, Gispen WH, Verkleij AJ (1991) Freeze-substitution and Lowicryl HM20 embedding of fixed rat brain: suitability for immunogold ultrastructural localization of neural antigens. *J Histochem Cytochem* 39(9):1267.
- Przedborski S, Jackson-Lewis V (2007) Protocol for the MPTP mouse model of Parkinson's disease. *Nat Protoc* 2(1):141. <http://dx.doi.org/10.1038/nprot.2006.342>.
- Yang J, Lunde L, Nuntagij P, Oguchi T, Camassa L, Nilsson LNG, Lannfelt L, Xu Y, Amiry-Moghaddam M, Ottersen OP, Torp R (2011) Loss of Astrocyte Polarization in the Tg-ArcSwe Mouse Model of Alzheimer's Disease. *J Alzheimers Dis* 27(4):711–722. <http://dx.doi.org/10.3233/jad-2011-110725>.
- Zhang J, Yang B, Sun H, Zhou Y, Liu M, Ding J, Fang F, Fan Y, Hu G (2016) Aquaporin-4 deficiency diminishes the differential degeneration of midbrain dopaminergic neurons in experimental Parkinson's disease. *Neurosci Lett* 614:7–15. <http://dx.doi.org/10.1016/j.neulet.2015.12.057>.

(Received 30 March 2017, Accepted 12 July 2017)
(Available online 19 July 2017)

PAPER II

Pro-inflammatory role of AQP4 in mice subjected to intrastriatal injections of the parkinsonogenic toxin MPP+.

Prydz A*, Stahl K*, Zahl S, Skauli N, Skare Ø, Ottersen OP, Amiry-Moghaddam M., *Cells*. 2020 Nov 5;9(11):E2418. doi: 10.3390/cells9112418.

* Both authors have contributed equally.

Article

Pro-Inflammatory Role of AQP4 in Mice Subjected to Intrastriatal Injections of the Parkinsonogenic Toxin MPP+

Agnete Prydz ^{1,†}, Katja Stahl ^{1,†}, Soulmaz Zahl ¹, Nadia Skauli ¹, Øivind Skare ² , Ole Petter Ottersen ³ and Mahmood Amiry-Moghaddam ^{1,*}

¹ Laboratory of Molecular Neuroscience, Division of Anatomy, Department of Molecular Medicine, Institute of Basic Medical Sciences, University of Oslo, P.O. Box 1105, Blindern, 0317 Oslo, Norway; agnpry@gmail.com (A.P.); katja.stahl@gmail.com (K.S.); soulmaz.r@gmail.com (S.Z.); nadia.skauli@medisin.uio.no (N.S.)

² Department of Occupational Medicine and Epidemiology, National Institute of Occupational Health, Gydas vei 8, 0363 Oslo, Norway; Oivind.Skare@stami.no

³ President's Office, Karolinska Institutet, Nobels väg 6, 171 77 Stockholm, Sweden; ole.petter.ottersen@ki.se

* Correspondence: mahmo@medisin.uio.no; Tel.: +47-91742177

† These authors have contributed equally.

Received: 14 October 2020; Accepted: 2 November 2020; Published: 5 November 2020

Abstract: Aquaporin-4 (AQP4) is critically involved in brain water and volume homeostasis and has been implicated in a wide range of pathological conditions. Notably, evidence has been accrued to suggest that AQP4 plays a proinflammatory role by promoting release of astrocytic cytokines that activate microglia and other astrocytes. Neuroinflammation is a hallmark of Parkinson's disease (PD), and we have previously shown that astrocytes in substantia nigra (SN) are enriched in AQP4 relative to cortical astrocytes, and that their complement of AQP4 is further increased following treatment with the parkinsonogenic toxin MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine). Here, we investigated the effect of *Aqp4* deletion on microglial activation in mice subjected to unilateral intrastriatal injection of 1-methyl-4-phenylpyridinium (MPP+, the toxic metabolite of MPTP). Our results show that MPP+ injections lead to a pronounced increase in the expression level of microglial activating genes in the ventral mesencephalon of wild type (WT) mice, but not *Aqp4*^{−/−} mice. We also show, in WT mice, that MPP+ injections cause an upregulation of nigral AQP4 and swelling of astrocytic endfeet. These findings are consistent with the idea that AQP4 plays a pro-inflammatory role in Parkinson's disease, secondary to the dysregulation of astrocytic volume homeostasis.

Keywords: AQP4; Parkinson's disease; MPP+; astrocyte; microglia; neuroinflammation

1. Introduction

Of the 13 mammalian aquaporins (AQP) that have been discovered to date, three are expressed and functional in brain tissue [1]. One of these (AQP1) shows a very restricted expression, limited to the choroid plexus, while the other two (AQP4 and AQP9) are found in many brain regions, including the SN [2,3]. Other AQPs, including AQP2, 3, 5, 6 and 8, have been reported to be expressed in the brain, but so far only at the mRNA level [4].

AQP4 is by far the most prevalent aquaporin in brain—being expressed across all brain regions—and is particularly abundant at the brain-blood interface as a major component of the molecular assembly in perivascular astrocytic endfeet [5–7]. AQP9 is more weakly expressed, but with

a preponderance in the mesencephalon [3]. Thus, both AQP4 and AQP9 are relevant in the context of Parkinson's disease.

AQP9 has a broad substrate profile and has been shown to mediate transport of parkinsonogenic compounds like arsenite and 1-methyl-4-phenylpyridinium (MPP+) [8,9]. In our previous study, we showed that the targeted deletion of AQP9 protects against MPP+ induced degeneration of nigral neurons [8]. Based on this finding, we hypothesized that AQP9 could be parkinsonogenic by acting as an influx route of compounds that are toxic to nigral dopaminergic neurons.

Unlike AQP9, AQP4 is strikingly selective for water, due to the dimensional constraint and configuration of the channel pore, and plays an important role in brain water homeostasis and dyshomeostasis, including brain edema [10–13]. However, evidence has accrued to show that AQP4 is engaged in a host of brain functions and dysfunctions that are only indirectly coupled to or even independent of water transport [7,14].

The list of neurological conditions in which AQP4 might play a pathophysiological role includes epilepsy [15,16] and neurodegenerative diseases, such as Alzheimer's disease (AD) [17–20], amyotrophic lateral sclerosis (ALS) [21,22], Huntington's disease (HD) [23] and spongiform encephalopathy (SE) [24]. Studies have also highlighted possible links between AQP4 and infectious and neuroinflammatory diseases, most notably meningitis [25], malaria [26] and neuromyelitis optica (NMO) [27].

Several neurological conditions—including epilepsy and a number of neurodegenerative diseases—also show hallmarks of neuroinflammation [28,29]. This is also true of Parkinson's disease (PD) [30]. Here, we use an animal model to explore our hypothesis that AQP4 could have a proinflammatory impact in PD. If so, this would further expand the number of roles attributed to AQP4 in neurological conditions.

The present work was inspired by study of Li et al. [31], which provided evidence of a proinflammatory role of AQP4 both in a mouse model of experimental autoimmune encephalopathy (EAE) and in an in vitro model. In both experimental approaches, mice or murine cell cultures with targeted deletion of AQP4 showed lower inflammation scores than wild type (WT) mice used as controls. The in vitro model showed that lipopolysaccharide (LPS) induced release of the cytokines TNF- α and IL-6 from astrocytes in an AQP4-dependent manner, possibly as a consequence of astrocytic swelling. These cytokines are known to activate cerebral microglia during neuroinflammation [32,33].

The goal of the present study was twofold. First, we set out to investigate whether intrastriatal injections of 1-methyl-4-phenylpyridinium (MPP+)—a mouse model of PD developed in our laboratory—upregulate expression of genes involved in microglial activation. Second, if such an effect could be demonstrated, we asked whether it is amplified by targeted deletion of AQP4. An answer in the affirmative would encourage further studies to resolve whether AQP4 serves a proinflammatory role in PD.

The timelines of this study is highlighted by recent investigations on AQP4 in brain. Notably, we have shown that nigral astrocytes are enriched in AQP4 relative to cortical astrocytes, and that their complement of AQP4 is further increased following treatment with MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) [34]. MPP+—the parkinsonogenic drug used in the present study—is the active metabolite of MPTP [35]. Further, imaging studies have revealed abnormal accumulation of water in mesencephala of patients with advanced PD [36].

Here, we demonstrate that MPP+ injections lead to an increase in microglial activation in the ventral mesencephalon, and that this increase is curbed by deletion of *Aqp4*. We also show, in WT mice, that MPP+ injections cause an upregulation of nigral AQP4 and swelling of astrocytic endfeet. These findings are consistent with the idea that upregulation of AQP4 plays a pro-inflammatory role in Parkinson's disease—possibly secondary to dysregulation of astrocytic volume homeostasis.

2. Materials and Methods

2.1. Animals

Mixed gender and adult WT C57BL/6J mice (Jackson Laboratories, Boulder, CO, USA) bred in the local animal facility ($n = 20$) and constitutive *Aqp4*^{−/−} mice backcrossed for more than 10 generations on C57BL/6J background ($n = 22$) were used in this study. *Aqp4*^{−/−} mice, generated as described by Thrane et al. [37], carry a mutated *Aqp4* gene where exon 1–3 is replaced by a flippase recognition target (FRT)-neomycin-FRT-LoxP-validated cassette inserted downstream of exon 3, and a LoxP site inserted upstream of exon 1, thus, avoiding any expression of putative splice variants. Both experimental groups (*Aqp4*^{−/−} and WT controls) were bred and kept at the local animal facility under the same conditions. For control purposes, we also purchased WT mice with the C57BL/6J genetic background from The Jackson Laboratory ($n = 12$). These animals had to be discarded as controls, as they differed from our internal controls (WT mice bred in the local facility) in regard to cell counts and their response to MPP+ (Supplementary Figure S1). The difference between these two cohorts of presumptive genetically identical animals is potentially interesting and will be the subject of further studies. Experimental protocols were approved by The Norwegian Animal Research Authority (NARA) with license number FOTS 4012 and carried out in accordance with the European Directive 2010/63/EU.

2.2. Stereotaxic Surgery and Injection of MPP+

The procedure was carried out as previously described [8]. In short, the animals were anesthetized with zoletil mixture (Zoletil Forte (250 mg/mL), Rompun (20 mg/mL) and Fentanyl (50 µg/mL); 0.1 mL/10 g; intraperitoneally) prior to fixing the head to a stereotaxic frame (TSE Systems, Bad Homburg, Germany). A total of 7.5 µg MPP+ (Sigma-Aldrich, St. Louis, MO, USA) dissolved in 1 µL of 0.9% saline, or 1 µL of 0.9% saline alone, was injected into the striatum +0.6 mm anterior to bregma, +2.2 mm laterally and −3.2 mm ventrally [38] at 12 µL/hr, using a syringe pump (Kd Scientific, Holliston, MA, USA) Body temperature was maintained at 36.5 ± 0.5 °C by a heating pad (PanLab, Barcelona, Spain), and the eyes were covered with Vaseline during the procedure.

2.3. Post-Operative Animal Care

At 24 h post-surgery, all animals were put on a heating pad overnight and were treated with Rimadyl (0.1 mL/10 g; s.c. every 12 h) to prevent pain, and a mixture of 0.9% saline and 0.9% sucrose (s.c. every 24 h) to restore water and energy balance. The animals were fed moisture pellets to improve food and water uptake. This treatment was continued for as long as needed, until the animals were sacrificed 7 days after surgery. The animals were scored daily for their weight and clinical appearance to evaluate the post-operative condition.

2.4. Tissue Preparation for Light and Electron Microscopic Cytochemistry

At 7 days post-surgery, the animals (WT; $n = 20$, *Aqp4*^{−/−}; $n = 22$) were deeply anesthetized with zoletil mixture (0.2 mL/10 g; i.p.), then euthanized by intra-cardial perfusion fixation with 2% ice cold dextran, followed by 15 min of 4% formaldehyde (FA), freshly depolymerized from paraformaldehyde (PFA), dissolved in 0.1 M phosphate buffer (PB) for light microscopic cytochemistry, or 4% FA and 0.1% glutaraldehyde in 0.1 M PB for electron microscopic cytochemistry (all from Sigma-Aldrich). Brains were dissected out and post-fixed for 24 h in the fixation solution.

2.5. Immunocytochemistry

The tissue used for light microscopic cytochemistry was cryo-protected by sequential immersions in 10%, 20% and 30% sucrose solutions. The midbrain was then isolated and sectioned into coronal sections of 40 µm on a HM 450 freeze microtome (Microm, Walldorf, Germany). From each animal, 10 sections were systematically selected for further immunohistochemical analyses (immunofluorescence;

WT ($n = 3$), $Aqp4^{-/-}$ ($n = 3$) and stereological analysis; WT ($n = 7$), $Aqp4^{-/-}$ ($n = 9$), with a slice interval of 120 μm , covering the entire substantia nigra pars compacta (SNpc), from Bregma -2.70 to -3.80 [38]. Immunocytochemistry for stereological quantifications has been previously described [8]. In brief, free-floating sections were blocked and permeabilized before incubation in primary antibody (Table 1) overnight, followed by incubation with secondary and tertiary antibody. Finally, sections were incubated with 3,3'-diaminobenzidine (DAB; Sigma-Aldrich) dissolved in 0.1 M PBS, then treated with a DAB-solution containing 0.03% H_2O_2 (Applichem, Darmstadt, Germany), before washing and mounting with glycerin-gelatin.

For immunofluorescence and confocal imaging, free floating sections were blocked, permeabilized and incubated with primary antibody (Table 1), prior to incubation with species-specific secondary antibodies prior to mounting. Images were collected with a LSM 510 META Confocal Microscope (Zeiss, Jena, Germany) using a 40 $\times$ oil objective. 4',6-diamidino-2-phenylindol (DAPI) fluorescence was captured at 405 nm, Alexa 488 at 510 nm, Cy3 at 568 nm and Cy5 at 670 nm.

Table 1. Primary secondary and tertiary antibodies for immunocytochemistry and -fluorescence.

Methods	Primary Antibody	Secondary Antibody	Tertiary Antibody
Immunocytochemistry	Rabbit anti-onized calcium-binding adapter molecule 1, Iba1, 1:200, Wako	Biotinylated anti-rabbit IgG (H + L) produced in goat, 1:100, Vector, Burlingame	Streptavidin-Biotinylated horse radish peroxidase complex, 1:100, GE Healthcare
	Mouse anti-glial fibrillary acidic protein, GFAP, 1:1000, Nordic BioSite AB	Biotinylated anti-mouse IgG (H + L) produced in goat, 1:100, Vector, Burlingame	Streptavidin-Biotinylated horse radish peroxidase complex, 1:100, GE Healthcare
	Mouse anti-tyrosine hydroxylase, 1:1000, Chemicon	Biotinylated anti-mouse IgG (H + L) produced in goat, 1:100, Vector, Burlingame	Streptavidin-Biotinylated horse radish peroxidase complex, 1:100, GE Healthcare
Immunofluorescence	Mouse anti-tyrosine hydroxylase, 1:1000, Chemicon	Alexa 488, 1:500, Thermo Fisher Scientific	
	Rabbit anti-AQP4, 1:400, Sigma-Aldrich	Cy3 anti-rabbit, 1:500, Jackson ImmunoResearch Laboratories	
	Chicken anti-glial fibrillary acidic protein, GFAP, 1:1000, Nordic BioSite AB	Cy5 anti-chicken, 1:500, Jackson ImmunoResearch Laboratories	

2.6. Stereological Quantifications of Dopaminergic Density in the Midbrain

Stereological quantifications of dopaminergic cells in midbrain subregions was performed bilaterally in WT and $Aqp4^{-/-}$ mice injected with MPP+ or saline. The investigator was blinded to genotype and treatment. The procedure has been described in detail [8].

2.7. Electron Microscopy

Tissue blocks $<1 \text{ mm}^3$ of SNpc from both the ipsilateral and the contralateral hemisphere of WT and $Aqp4^{-/-}$ mice were dissected out, cryoprotected in glycerol and quickly frozen in liquid propane (-170°C). The tissue was then subjected to freeze substitution [39], embedded in Lowicryl HM20 Resin (Lowy, Waldkraiburg, Germany) at -30°C and polymerized by UV light. Ultrathin sections of 90–100 nm were cut from the blocks using an ultramicrotome (Leica EM UC6, Vienna, Austria) and transferred to mesh grids [40]. Each section was contrasted with uranyl acetate and lead citrate for 1.5 min and rinsed in ddH₂O in-between. The sections were examined using an electron microscope (Tecnai 12, FEI Company, Eindhoven, The Netherlands).

2.8. Electron Microscopic Analysis

For the quantification of endfoot width, as an indicator for swelling, 25 electron micrographs of randomly selected capillaries from SN of WT ($n = 4$) and *Aqp4*^{-/-} mice ($n = 4$) were obtained from each section at 26,500× magnification. Endfoot width was assessed by calculating endfoot area per unit length of adluminal membrane using the software program ImageJ (U. S. National Institutes of Health, Bethesda, MD, USA). The analyzer was blind to genotype and area.

2.9. RNA Isolation and Quantitative Real Time qPCR Analysis

At 7 days post-surgery, the animals injected with MPP+ (WT ($n = 6$), *Aqp4*^{-/-} ($n = 5$) or saline WT ($n = 3$), *Aqp4*^{-/-} ($n = 3$)) were decapitated under isofluorane (Baxter, Deerfield, IL, USA) anesthesia and the brains were dissected out. The brains were cut along the midsagittal line to separate the injected and control hemispheres. The midbrain and striatum were dissected out from both hemispheres, snap frozen and stored at -80 °C until analysis. The total RNA was isolated from the regional brain samples using the RNeasy Plus Mini Kit (QIAGEN, Hilden, Germany). The RNA concentration and integrity were determined using the NanoDrop 2000c spectrophotometer (Thermo Scientific, Oslo, Norway) and agarose gel electrophoresis. The cDNA was synthesized using GoScript Reverse Transcription System (Promega, Madison, WI, USA) using Oligo (dT)15. The gene expression was evaluated by using the qPCR absolute quantification. The qPCR was performed in a total volume of 20 µL, containing the Power SYBR Green PCR Master Mix (Applied Biosystems, Oslo, Norway), forward and reverse primers (10 µM), and 10 ng of total cDNA per reaction. Thermal cycling was performed on the StepOnePlus system (Applied Biosystems) with the following conditions: 95 °C for 10 min, followed by 40 cycles at 95 °C for 15 s and 60 °C for 1 min and at the end the melting curve step. No reverse transcriptase (NRT) and no template control (NTC) negative controls were included in the study. Using the NormFinder software (Aarhus, Denmark), the expression variation of *Actb*, *H2afz*, *Pgk1* and *Ubc* as endogenous reference genes were evaluated. *Ubc* was selected as the most stable endogenous control for normalization across groups and regions with a stability value of 0.184. One-way ANOVA with Bonferroni post hoc test was used for statistical analysis of the data. The data are presented as mean ± 2SE, and $p < 0.05$ was considered as significant. A list of the primer pairs used in the experiment is shown in Table 2.

Table 2. Primer pairs.

Gene Name	Forward Primer	Reverse Primer	Product Size	PCR Efficiency
<i>Aqp4</i>	AGCAATGGATTTTCCGTTC	TGAGCTCCACATCAGGACAG	203 bp	96%
<i>Aif1</i>	CTGCCAGCCTAAGACAACCA	GGAATGTGCTGTGATCCCCT	128 bp	99%
<i>Cx3cr1</i>	TGCTCAGGACCTCACCATGTC	CTCAAGGCCAGGTCAGGAG	246 bp	98%
<i>Itgam</i>	TGCGCGAAGGAGATATCCAG	GCCTGCGTGTGTGTCTTT	108 bp	94%
<i>Cd14</i>	CAGAGAACACCACCGCTGTA	CACGCTCCATGGTCGGTAGA	97 bp	96%
<i>Cxcl10</i>	ATGACGGGCCAGTGAGAATG	TCGTGGCAATGATCTCAACAC	80 bp	97%
<i>Tgfb1</i>	AATTCCTGGCGTTACCTTGG	AGTGAGCGCTGAATCGAAAG	139 bp	100%
<i>Trem2</i>	CTGGAGGACCCTCTAGATGAC	CCACAGGATGAAACCTGCCT	116 bp	98%
<i>Gfap</i>	GCACTCAATACGAGGCAGTG	GCTCTAGGGACTCGTTCGTG	207 bp	97%
<i>Ubc</i>	CGTCGAGCCCACTGTTACCAACAAGAAG	CCCCATCACACCAAGAACAAGCACAAG	112 bp	92%

3. Results

3.1. Survival Rate and Clinical Appearance Post-Surgery

All animals were scored daily for their clinical appearance, including the parameters weight loss, inactivity, appearance and reaction patterns, giving a total score ranging from 0–12. Animals reaching a score of 10 or above were considered severely affected by the toxin and were euthanized, due to ethical concerns. Among the total number of animals treated with MPP+ ($n = 32$), five animals died or were euthanized. Among those animals, three animals were WT, while two animals were *Aqp4*^{-/-}. There was no difference in behavioral assessments between WT and *Aqp4*^{-/-} mice.

3.2. Unilateral Intrastratial Injections of MPP+ Lead to Astrogliosis and Microgliosis in the Ipsilateral SN

Astrogliosis and microgliosis have been demonstrated in the ventral midbrain of many animal models of PD, including systemic treatment with MPTP [34,41]. We investigated if unilateral intrastratial MPP+ injections induce astrogliosis and microgliosis in the ventral midbrain. Light microscopic immunohistochemistry of midbrain sections of animals subjected to unilateral intrastratial MPP+ injections showed stronger glial fibrillary acidic protein (GFAP) and ionized calcium-binding adapter molecule 1 (Iba1) staining in the injected hemisphere, compared to the control hemisphere (Figure 1A–H). WT and *Aqp4*^{−/−} animals did not differ in this regard. In both groups of animals, GFAP-positive astrocytes were most abundant in a distinct area corresponding to the SN of the injected hemisphere (Figure 1B,D). In both genotypes, the SN of the injected hemispheres showed strong immunostaining for Iba1, with WT mice displaying a slightly higher staining intensity than *Aqp4*^{−/−} animals (Figure 1E–H). Microglia stained for Iba1 exhibited enlarged cell bodies and chubby processes (Figure 1F,H) indicative of a microglial reactive state [42].

Figure 1. Increased expression of glial fibrillary acidic protein (GFAP) and (ionized calcium-binding adapter molecule 1) Iba1 in substantia nigra (SN) following treatment with MPP+. Representative micrographs showing midbrain sections immunostained with antibodies to the astrocyte marker GFAP upper panel (A–D) or the microglial marker Iba1 lower panel (E–H). Unilateral injection of 1-methyl-4-phenylpyridinium (MPP+) in the striatum resulted in increased expression of GFAP (arrows B,D) in the ipsilateral substantia nigra pars compacta (SNpc) (injected, B,D) compared to the contralateral hemisphere (Ctrl, A,C) in wild type (WT) (A,B) as well as in *Aqp4*^{−/−} animals (C,D). An increased expression of Iba1 (arrow H) is also visible in the ipsilateral SNpc (F), compared to the contralateral hemisphere (E) in WT, as well as in *Aqp4*^{−/−} animals, respectively, (H,G). Iba1 showed a slightly higher staining intensity in WT compared to *Aqp4*^{−/−} animals (lower panel, compared the area around SNpc in the injected hemispheres). Scale bar: 1000 μ m; inset: 20 μ m.

3.3. Unilateral Intrastratial MPP+ Injections Lead to a Strong Increase in the Transcript Levels of Microglial Activating Genes in Ipsilateral Midbrain of Wild Type but Not *Aqp4*^{-/-} Mice

We performed quantitative real time PCR analysis on samples from midbrain of injected and control hemispheres of WT and *Aqp4*^{-/-} mice (Figure 2). Confirming the immunohistochemical data, our analysis showed increased expression level for *Gfap* in the midbrain of the MPP+ injected hemisphere compared to midbrain of the control hemisphere in both WT (control: 1035 ± 134 injected: 3487 ± 542 $p < 0.001$) and *Aqp4*^{-/-} mice (control: 753 ± 448 , injected: 4182 ± 1438 $p < 0.001$). There was no significant difference between the two genotypes. The expression level of the general microglial marker *Iba1* (*Aif1*) was significantly increased in the injected hemisphere of both genotypes. Notably, the transcript levels of *Aif1* in the ipsilateral midbrain were significantly higher in WT than in *Aqp4*^{-/-} mice (WT: 2217.4 ± 242.2 vs *Aqp4*^{-/-}: 1295.3 ± 220.8 $p < 0.001$).

To resolve whether AQP4 plays a pro-inflammatory role, we analyzed the expression levels of key genes coding for microglial membrane receptors known to be upregulated during inflammation. We specifically investigated the expression levels of *Cd14*, a key organizer of microglial inflammatory response [43]; *Trem2*, a receptor promoting microglial survival, proliferation and phagocytic activity [44]; *Cx3cr1*, a chemokine receptor involved in microglial inflammatory response [45,46]; and *Itgam*, the gene coding for the integrin receptor CD11b involved in microglial adhesion and migration [47]. In WT animals, our qPCR analysis revealed a several fold increase in the expression levels of each of these genes in midbrain samples ipsilateral to MPP+ injections. In *Aqp4*^{-/-} mice, in contrast, none of these genes differed significantly between injected and control hemispheres in regard to levels of expression (Figure 2).

We also analyzed the expression levels of the two cytokines, *Tgfb1* and *Cxcl10*, previously shown to be upregulated in MPTP models of PD [48,49]. Under normal conditions in the midbrain, TGF- β 1 is mainly expressed by neurons, while under pathological conditions, it is, in addition, highly expressed in activated microglia [50], and its secretion leads to the upregulation of microglial CX3CR1 [51,52]. CXCL10 is expressed in and secreted by astrocytes and to some extent by microglia [53,54]. Earlier studies have shown that release of CXCL10 from astrocytes is involved in early microglial activation, initiating migration and inducing a pro-inflammatory phenotype [55]. Expression levels of both chemokines were strongly increased in the midbrain of the injected hemisphere of WT mice, compared to the control side (*Tgfb1*, control: 60.4 ± 6 , injected: 202.9 ± 41.4 , $p < 0.001$; *Cxcl10*, control: 42.8 ± 18.2 , injected: 1492.7 ± 340 , $p < 0.001$). However, in the *Aqp4*^{-/-} midbrain samples, only the increase in the expression level of *Cxcl10* reached significant levels (control: 83.9 ± 37.8 , injected: 1128.6 ± 342 , $p < 0.001$), while *Tgfb1* failed to increase significantly, following unilateral intrastratial MPP+ injection (injected: 104.8 ± 23.7 , control: 66.4 ± 17.6 , $p > 0.05$), indicating reduced microglial activation.

Quantitative real-time PCR analysis showed a significant increase in the expression levels of *Gfap* and *Aif1* in the midbrain of MPP+ injected hemispheres in both the WT and *Aqp4*^{-/-} mice. The increase observed in the injected hemispheres of the WT mice was significantly higher for *Aif1* than the increase in the injected hemisphere of *Aqp4*^{-/-} mice ($p < 0.01$). In the case of *Gfap*, there was a significant increase in the transcript levels in the injected hemisphere compared to the control hemisphere of both genotypes $p < 0.001$, with no significant difference between the genotypes.

Expression of *Cd14* was increased from 70.2 ± 9.2 mRNA copies per ng RNA in the control midbrain to 174.5 ± 26.4 copies in the injected midbrain of the WT mice ($p < 0.001$), compared to 49.5 ± 10.4 in the control and 83.8 ± 26 in the injected hemispheres of the *Aqp4*^{-/-} mice ($p = 1.0$). The trend was similar for *Trem2* (WT control: 76.4 ± 13.6 , WT injected: 568.3 ± 116 , $p < 0.001$ vs *Aqp4*^{-/-} control: 111.1 ± 39.4 , *Aqp4*^{-/-} injected: 222.0 ± 64 , $p = 1.0$) *Cx3cr1* (WT control: 243.3 ± 40.6 , WT injected: 743.9 ± 171.6 , $p < 0.001$ vs *Aqp4*^{-/-} control: 296.2 ± 176.8 , *Aqp4*^{-/-} injected: 315.7 ± 169.2 , $p = 1.0$), *Itgam* (WT control: 20.3 ± 3 , WT injected: 41.9 ± 10 , $p = 0.001$ vs *Aqp4*^{-/-} control: 6.6 ± 4.6 , *Aqp4*^{-/-} injected: 17.6 ± 11.2 , $p > 0.05$). The mRNA expression of *Tgfb1* and *Cxcl10*, were significantly increased in the ipsilateral hemisphere of the WT animals compared to control hemisphere, and the increase was significantly higher than that observed in the *Aqp4*^{-/-} animals, at $p < 0.05$ and $p < 0.01$, respectively.

In the *Aqp4*^{−/−} mice, the increase in the expression levels of *Cxcl10* and not *Tgfb1*, reached significant levels in the midbrain ipsilateral to the MPP+ injection, compared to the control midbrain.

Figure 2. Increased mRNA levels of microglial activating genes in ipsilateral midbrain of WT but not *Aqp4*^{−/−} mice, following unilateral intrastriatal MPP+ injections. Bars are mean ± 2SEM. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

3.4. Expression of AQP4 in SN after MPP+ Injections

We have previously shown that systemic MPTP treatment causes an upregulation of AQP4 expression in SN [34]. Here, we investigated whether unilateral intrastratial injections of MPP+ have a similar effect. Our immunofluorescence analysis revealed an increased expression of AQP4 ipsilateral to, but not contralateral to MPP+ injections, particularly around larger vessels and in the neuropil of the SN (Figure 3A,B). Ipsilateral and contralateral SN showed no difference in AQP4 immunofluorescence labeling intensity when MPP+ injections were replaced by unilateral intrastratial injections of saline (Figure 3C,D).

Quantitative real time PCR analysis showed that *Aqp4* expression in the midbrain was significantly increased in the hemisphere treated with MPP+, as compared to the contralateral control hemisphere and saline controls (Figure 3E). The transcript levels of *Aqp4* mRNA in hemispheres treated with MPP+ was 4976 ± 1110 mRNA copies per ng RNA, compared with 1397 ± 124.4 in the control hemisphere ($p < 0.001$).

Figure 3. Increased expression of Aquaporin-4 (AQP4) protein and mRNA in the substantia nigra following treatment with MPP+ (A–D). Unilateral injection of MPP+ in the striatum results in loss of TH-positive cells (green) in the ipsilateral SNpc (A, injected) compared to the contralateral SNpc (B, ctrl) and animals receiving unilateral injections of saline (C,D). Strong upregulation of AQP4 (red) is further observed in the ipsilateral SN, particularly around larger vessels (double arrows) (A) compared to larger vessels in the control hemisphere (B) and saline controls (C,D). Increased AQP4 labeling is also evident in non-endfeet membranes of the neuropil (asterisk) following treatment with MPP+ (A). The toxin also induced upregulation of GFAP (orange) in the ipsilateral hemisphere (single arrow), (A). The injected hemisphere showed significantly fewer TH-positive cells (green, A) compared to the control hemisphere (B). Increased levels of *Aqp4* mRNA following treatment with MPP+ (E). Quantitative real-time PCR analysis showed more than two-fold higher expression of *Aqp4* mRNA in midbrain samples ipsilateral to the MPP+ injection compared to midbrain samples of the control hemisphere (ctrl). Bars are mean $\pm$ 2SEM. *** $p < 0.001$. Scale bar: 50 μ m.

3.5. Electron Microscopic Analysis of Endfoot Width

EM analysis showed an increased width of perivascular astrocyte endfeet in WT and *Aqp4*^{−/−} mice injected with MPP+ compared with saline controls (Figure 4). Endfoot width was larger in MPP+ injected WT mice ($1.28 \mu\text{m} \pm 0.28$) than in MPP+ injected *Aqp4*^{−/−} animals ($0.8 \mu\text{m} \pm 0.2$; $p < 0.01$). Additionally, after saline injections did the two genotypes differ in regard to endfoot width (Figure 4).

Figure 4. Perivascular endfoot width following treatment of MPP+ and saline. Electron micrographs of perivascular astrocytic endfeet in animals exposed to MPP+ treatment (A,B) and saline controls (C,D). Arrows mark the membrane domains of the astrocytic endfoot facing endothelial basal lamina, arrowheads mark membrane facing neuropil, thus, identifying the astrocyte endfoot. Asterisks marks basal lamina between the endothelial cell and astrocyte endfoot. The statistical analysis of the perivascular astrocytic endfoot width in SNpc of MPP+ injected hemispheres (E) showed a larger endfoot width in both genotypes, compared to the saline controls ($p < 0.01$). The endfoot width in the WT animals was larger compared to the *Aqp4*^{−/−} animals in both the MPP+ and saline injected hemispheres ($p < 0.01$). E, endothelial cell; L, capillary lumen. Bars are mean $\pm$ 2SEM. ** $p < 0.01$. Scale bar: 500 nm.

3.6. Stereological Quantification of Dopaminergic Cell Density

Dopaminergic cells (TH-ir cells) in the ipsi- (Figure 5A,C) and contralateral (Figure 5B,D) hemispheres of SNpc were quantified in *Aqp4*^{−/−} and WT animals subjected to unilateral intrastratial injections of MPP+ or saline. The number of TH-ir cells in the MPP+ injected hemisphere was significantly higher in *Aqp4*^{−/−} mice than in WT, with average cell counts of respectively 1789 ± 262 and 1401 ± 82 ($p < 0.05$, Figure 5E).

In animals treated with saline, no difference in TH-ir cell counts was observed between genotypes or between the ipsi- and contralateral side within genotypes (Figure 5F).

TH-ir cell counts in ventral tegmental area (VTA) revealed no significant differences between ipsi- and contralateral midbrain following unilateral MPP+ injections. Further, cell counts in VTA did not differ between WT and *Aqp4*^{−/−} mice after MPP+ injections, nor after injections of saline (Figure 5G).

Figure 5. Dopaminergic neuronal cell count after unilateral treatment with MPP⁺ or saline in the SNpc and ventral tegmental area (VTA) of *Aqp4*^{-/-} and WT animals. Following unilateral treatment with MPP⁺, loss of TH-immunoreactive (TH-ir) cells is evident in the ipsilateral hemisphere (A,C) compared to the contralateral side (B,D) in both *Aqp4*^{-/-} and WT animals. In the SNpc, the average TH-ir cell count reduced from 2742 ± 490 in the control hemisphere to 1401 ± 82 in the injected hemisphere of the WT animals ($p < 0.001$) (E). In *Aqp4*^{-/-} animals, the average TH-ir cell count for the control and injected hemispheres were 2940 ± 856 and 1789 ± 262, respectively ($p = 0.052$). The number of TH-ir cells in the MPP⁺ injected hemisphere was significantly higher in *Aqp4*^{-/-} mice than in WT (E). Unilateral treatment with saline did not result in reduction in the dopaminergic cell count in the ipsilateral hemisphere of *Aqp4*^{-/-} ($p = 0.948$) or WT animals ($p = 0.916$), (F). There were no differences between the injected hemispheres or the control hemispheres of the two genotypes, respectively $p = 0.732$ and $p = 0.693$ (F). In the VTA, no significant differences were observed in the cell count of TH-positive cells in the two hemispheres of the two genotypes, ($p = 0.607$ for *Aqp4*^{-/-} and $p = 0.0215$ for WT, G). There were no differences between the genotypes in the injected side ($p = 0.860$) or the control sides ($p = 0.864$) (G). Bars are mean ± 2SEM. Scale bars: 1000 μm; inset: 20 μm. * $p < 0.05$.

4. Discussion

The question explored in the present study is whether AQP4—the most prevalent aquaporin in brain—is involved in the pathophysiology of PD, and if so—is it protective or does it exacerbate disease progression?

Recent studies in our laboratory and others have shown that AQP4 is more strongly expressed in nigral astrocytes than in astrocytes in cortex or hippocampus, and that its expression is further increased after MPTP injections in mice [34], and after 6-hydroxydopamin (6-OHDA) injections in

rats [56]. It is also interesting to note that imaging studies have revealed abnormal accumulation of water in mesencephala of patients with advanced Parkinson's disease [36]. Further, several studies have pointed to a putative coupling between AQP4 and inflammation [31,57]. These findings prompted us to investigate whether AQP4 could be involved in neuroinflammatory processes relevant for the development of PD. We chose a model that was successfully used to identify a putative link between AQP9 and loss of dopaminergic neurons. Our results show that AQP4 deletion restrains MPP⁺ induced upregulation of genes involved in microglia activation. Specifically, compared with WT mice, mice with targeted deletion of *Aqp4* revealed significantly lower expression of microglial genes known to be involved in microglial migration and phagocytosis.

Under pathological conditions, microglial activation is triggered by signal substances and inflammatory mediators released by peripheral immune cells, damaged neurons and astrocytes [32,33,58]. Of these cell types, only astrocytes are known to express AQP4. Thus, it is likely that the effects of *Aqp4* deletion presently reported are mediated by astrocytes.

Neuroinflammation is a hallmark of PD and of animal models of PD [59]. Microglia have been shown to be the main actors in the neuroinflammatory process contributing to dopaminergic cell loss, and several lines of evidence indicate that reactive astrocytes play an essential role in microglia activation. Upon exposure to neuronal α -synuclein, astrocytes express high levels of proinflammatory cytokines involved in migration and activation of microglia [53,60]. Moreover, the key role of astrocytes in MPTP-induced neuroinflammation and microglial activation was demonstrated in a recent study where astrocyte-specific deletion of inhibitor of nuclear factor kappa-B kinase subunit β (IKK2), a central upstream activator of NF- κ B, led to reduced neuroinflammation and microglial activation following MPTP treatment [61]. These findings are in line with the idea that astrocytes may sustain neuroinflammation in PD models.

While a proinflammatory role of AQP4 is consistent with the early study of Li et al. [23], the present findings are not easily reconciled with recent studies suggesting that AQP4 may play an immunosuppressive role in models of MPTP and LPS induced neuroinflammation [62,63]. The latter studies reported that *Aqp4*^{−/−} mice treated with MPTP or LPS revealed an increased microglial inflammatory response and a more pronounced loss of TH-neurons in the SN, compared with WT mice. The *Aqp4*^{−/−} mouse line used in these studies was found to have an impaired blood brain barrier, even in the absence of any experimental intervention [64]. A leaky blood brain barrier implies that the microenvironment of neurons and glia is perturbed and that neural cells are exposed to immune cells and molecules that are normally retained in blood. This makes results difficult to interpret. The *Aqp4*^{−/−} mouse line used in the present study has an intact blood brain barrier [65]. It is worth noting that MPTP alone does not lead to a disruption of blood brain barrier function [66].

As to the mechanisms underlying the present findings, Li et al. found that release of inflammatory mediators such as interferon gamma (IFN γ) and IL-1 β by astrocytes was coupled to AQP4-dependent astrocyte swelling [31]. We have previously shown that astrocytes release ATP, another activator of microglia, in an AQP4-dependent manner [37,67]. The exact mechanism of this AQP4-dependent ATP release is not known, but it could occur as part of the regulatory volume decrease where AQP4 and other AQPs have been shown to be involved [68,69].

In the present study, we demonstrate that *Aqp4*^{−/−} mice subjected to MPP⁺ injections show less swelling of astrocytic endfeet than MPP⁺ injected WT mice. We conclude that the most salient explanation of our findings is that *Aqp4* deletion curbs swelling induced release of compounds that normally cause microglial activation and inflammation.

Importantly, we show, in the current study, that astrocytic endfeet in SN are more swollen than endfeet in *Aqp4*^{−/−} mice both under basic conditions and after MPP⁺ exposure. The enrichment of AQP4 in nigral astrocytes and its further upregulation after toxin exposure may contribute to stronger inflammatory response and represent an additional factor contributing to the selective vulnerability of SN to toxins. Upregulation of AQP4 similar to that found here has been reported in several studies of chronic neurodegenerative diseases, including PD [34], AD [17,18,20,70], ALS [21,22] and SE [24],

indicating that increased expression of AQP4 is a common feature of neurodegeneration. Interestingly the sensitivity to MPTP (and inflammatory response) is increased in aged mice [71], which express higher levels of AQP4 than younger mice [72].

The present study was designed to assess the effect of *Aqp4* deletion on microglial activation in ventral mesencephalon, a process thought to occur upstream of cell degeneration. However, even at this early time point, the loss of dopaminergic neurons after MPP+ injections was less pronounced in *Aqp4*^{−/−} mice, compared with WT controls (38% vs 45%). Our data are consistent with the findings by other groups showing that dopaminergic cell loss is lower in TLR4-deficient mice vs WT one week after treatment with MPTP (34% vs 45%) [73]. TLR4 is member of Toll-like receptor family that mediates microglial activation in neurodegenerative diseases, including PD [74].

5. Conclusions

It is well established that PD is associated with neuroinflammation through elevation of cytokines [75], some of which are released by reactive astrocytes [53,60] and lead to activation of microglia [76].

This is the first study to show that *Aqp4* deletion reduces microglial activation indicating that AQP4 plays a proinflammatory role in a model of PD. The mechanisms underlying this effect remain to be identified, but the most likely explanation is that AQP4 promotes swelling of astrocytes and hence swelling-induced release of ATP and other compounds that, in turn, activate nigral microglia. Taken together with previous studies, our data suggest that two of the three aquaporins that are functional in brain—i.e., AQP4 and AQP9—are implicated in PD, albeit in fundamentally different ways. While AQP9 may serve to facilitate toxin access [8], AQP4 is likely to exacerbate the effect of toxin exposure. We hypothesize that differences in AQP4 or AQP9 expression—related to age, environmental exposure or genetic factors—may affect vulnerability to PD. This hypothesis will be explored in future studies.

Supplementary Materials: The following are available online at <http://www.mdpi.com/2073-4409/9/11/2418/s1>, Figure S1: Stereological Quantification of Dopaminergic Cell Density comparing the wild type (WT) mice bred in the local facility (WT-INT) and the WT mice purchased from Jacksons (WT-EXT).

Author Contributions: Conceptualization, M.A.-M., A.P. and K.S.; methodology, M.A.-M., K.S., A.P.; validation, M.A.-M., K.S., A.P., S.Z. and Ø.S.; formal analysis, A.P., K.S., S.Z. and Ø.S.; investigation, A.P., K.S., S.Z. and N.S.; data curation, A.P. and K.S.; writing—original draft preparation, M.A.-M., O.P.O., A.P. and K.S.; writing—review and editing, M.A.-M., O.P.O., K.S., A.P., S.Z. and N.S.; visualization, A.P., K.S. and N.S.; supervision, M.A.-M.; project administration, M.A.-M.; funding acquisition, M.A.-M. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by The Olav Thon Foundation, grant number 16-1606.

Acknowledgments: We are grateful to Bjørge Riber, Karen-Marie Gujord and Bashir Hakim for technical assistance. Finn-Mogens Haug for assistance with the analysis program, and Trygve Leergaard for assistance with the stereology program.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Nagelhus, E.A.; Ottersen, O.P. Physiological roles of Aquaporin-4 in brain. *Physiol. Rev.* **2013**, *93*, 1543–1562. [CrossRef] [PubMed]
2. Mylonakou, M.N.; Petersen, P.H.; Rinvik, E.; Rojek, A.; Valdimarsdottir, E.; Zelenin, S.; Zeuthen, T.; Nielsen, S.; Ottersen, O.P.; Amiry-Moghaddam, M. Analysis of mice with targeted deletion of AQP9 gene provides conclusive evidence for expression of AQP9 in neurons. *J. Neurosci. Res.* **2009**, *87*, 1310–1322. [CrossRef] [PubMed]
3. Badaut, J.; Petit, J.-M.; Brunet, J.-F.; Magistretti, P.J.; Charriaud-Marlangue, C.; Regli, L. Distribution of Aquaporin 9 in the adult rat brain: Preferential expression in catecholaminergic neurons and in glial cells. *Neuroscience* **2004**, *128*, 27–38. [CrossRef] [PubMed]

4. Badaut, J.; Lasbennes, F.; Magistretti, P.J.; Regli, L. Aquaporins in brain: Distribution, physiology, and pathophysiology. *Br. J. Pharmacol.* **2002**, *22*, 367–378. [[CrossRef](#)] [[PubMed](#)]
5. Nielsen, S.; Nagelhus, E.A.; Amiry-Moghaddam, M.; Bourque, C.; Agre, P.; Ottersen, O.P. Specialized membrane domains for water transport in glial cells: High-resolution immunogold cytochemistry of aquaporin-4 in rat brain. *J. Neurosci.* **1997**, *17*, 171–180. [[CrossRef](#)] [[PubMed](#)]
6. Amiry-Moghaddam, M.; Frydenlund, D.; Ottersen, O. Anchoring of aquaporin-4 in brain: Molecular mechanisms and implications for the physiology and pathophysiology of water transport. *Neuroscience* **2004**, *129*, 997–1008. [[CrossRef](#)] [[PubMed](#)]
7. Amiry-Moghaddam, M.; Ottersen, O.P. The molecular basis of water transport in the brain. *Nat. Rev. Neurosci.* **2003**, *4*, 991–1001. [[CrossRef](#)]
8. Stahl, K.; Rahmani, S.; Prydz, A.; Skauli, N.; Macaulay, N.; Mylonakou, M.N.; Torp, R.; Skare, O.; Berg, T.; Leergaard, T.B.; et al. Targeted deletion of the aquaglyceroporin AQP9 is protective in a mouse model of Parkinson's disease. *PLoS ONE* **2018**, *13*, e0194896. [[CrossRef](#)]
9. Carbrey, J.M.; Gorelick-Feldman, D.A.; Kozono, D.; Praetorius, J.; Nielsen, S.; Agre, P. Aquaglyceroporin AQP9: Solute permeation and metabolic control of expression in liver. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 2945–2950. [[CrossRef](#)]
10. Manley, G.T.; Fujimura, M.; Ma, T.; Noshita, N.; Filiz, F.; Bollen, A.W.; Chan, P.; Verkman, A. Aquaporin-4 deletion in mice reduces brain edema after acute water intoxication and ischemic stroke. *Nat. Med.* **2000**, *6*, 159–163. [[CrossRef](#)] [[PubMed](#)]
11. Vajda, Z.; Pedersen, M.; Füchtbauer, E.-M.; Wertz, K.; Stødkilde-Jørgensen, H.; Sulyok, E.; Dóczy, T.; Neely, J.D.; Agre, P.; Frokiaer, J.; et al. Delayed onset of brain edema and mislocalization of aquaporin-4 in dystrophin-null transgenic mice. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 13131–13136. [[CrossRef](#)] [[PubMed](#)]
12. Frydenlund, D.S.; Bhardwaj, A.; Otsuka, T.; Mylonakou, M.N.; Yasumura, T.; Davidson, K.G.V.; Zeynalov, E.; Skare, O.; Laake, P.; Haug, F.-M.; et al. Temporary loss of perivascular aquaporin-4 in neocortex after transient middle cerebral artery occlusion in mice. *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 13532–13536. [[CrossRef](#)]
13. Hirt, L.; Ternon, A.; Price, M.; Mastour, N. Protective role of early aquaporin 4 induction against postischemic edema formation. *Br. J. Pharmacol.* **2008**, *29*, 423–433. [[CrossRef](#)]
14. Papadopoulos, M.C.; Verkman, A.S. Aquaporin water channels in the nervous system. *Nat. Rev. Neurosci.* **2013**, *14*, 265–277. [[CrossRef](#)] [[PubMed](#)]
15. Alvestad, S.; Hammer, J.; Hoddevik, E.H.; Skare, O.; Sonnewald, U.; Amiry-Moghaddam, M.; Ottersen, O.P. Mislocalization of AQP4 precedes chronic seizures in the kainate model of temporal lobe epilepsy. *Epilepsy Res.* **2013**, *105*, 30–41. [[CrossRef](#)] [[PubMed](#)]
16. Eid, T.; Lee, T.-S.W.; Thomas, M.J.; Amiry-Moghaddam, M.; Bjørnsen, L.P.; Spencer, D.D.; Agre, P.; Ottersen, O.P.; De Lanerolle, N.C. Loss of perivascular aquaporin 4 may underlie deficient water and K⁺ homeostasis in the human epileptogenic hippocampus. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 1193–1198. [[CrossRef](#)]
17. Yang, J.; Lunde, L.L.; Paworn, N.; Tomohiro, O.; Camassa, L.M.A.; Nilsson, N.G.; Lannfelt, L.; Xu, Y.; Amiry-Moghaddam, M.; Ottersen, O.P.; et al. Loss of astrocyte polarization in the tg-ArcSwe mouse model of Alzheimer's disease. *J. Alzheimers Dis.* **2011**, *27*, 711–722. [[CrossRef](#)]
18. Pérez, E.; Barrachina, M.; Rodríguez, A.; Torrejón-Escribano, B.; Boada, M.; Hernández, I.; Sánchez, M.; Ferrer, I. Aquaporin expression in the cerebral cortex is increased at early stages of Alzheimer disease. *Brain Res.* **2007**, *1128*, 164–174. [[CrossRef](#)]
19. Moftakhar, P.; Lynch, M.D.; Pomakian, J.L.; Vinters, H.V. Aquaporin expression in the brains of patients with or without cerebral amyloid angiopathy. *J. Neuropathol. Exp. Neurol.* **2010**, *69*, 1201–1209. [[CrossRef](#)]
20. Hoshi, A.; Yamamoto, T.; Shimizu, K.; Ugawa, Y.; Nishizawa, M.; Takahashi, H.; Kakita, A. Characteristics of aquaporin expression surrounding senile plaques and cerebral amyloid angiopathy in Alzheimer disease. *J. Neuropathol. Exp. Neurol.* **2012**, *71*, 750–759. [[CrossRef](#)]
21. Kaiser, M.; Maletzki, I.; Hülsmann, S.; Holtmann, B.; Schulz-Schaeffer, W.; Kirchhoff, F.; Bähr, M.; Neusch, C. Progressive loss of a glial potassium channel (KCNJ10) in the spinal cord of the SOD1 (G93A) transgenic mouse model of amyotrophic lateral sclerosis. *J. Neurochem.* **2006**, *99*, 900–912. [[CrossRef](#)]
22. Nicaise, C.; Soyfoo, M.S.; Authalet, M.; De Decker, R.; Bataveljić, D.; Delporte, C.; Roland, P. Aquaporin-4 overexpression in rat ALS model. *Anat. Rec. Adv. Integr. Anat. Evol. Biol.* **2009**, *292*, 207–213. [[CrossRef](#)]

23. Wu, T.-T.; Su, F.-J.; Feng, Y.-Q.; Liu, B.; Li, M.-Y.; Liang, F.-Y.; Li, G.; Li, X.-J.; Zhang, Y.; Cai, Z.-Q.; et al. Mesenchymal stem cells alleviate AQP-4-dependent glymphatic dysfunction and improve brain distribution of antisense oligonucleotides in BACHD mice. *STEM CELLS* **2019**. [\[CrossRef\]](#)
24. Costa, C.; Tortosa, R.; Rodriguez, A.; Ferrer, I.; Torres, J.M.; Bassols, A.; Pumarola, M.B. Aquaporin 1 and aquaporin 4 overexpression in bovine spongiform encephalopathy in a transgenic murine model and in cattle field cases. *Brain Res.* **2007**, *1175*, 96–106. [\[CrossRef\]](#)
25. Papadopoulos, M.C.; Verkman, A.S. Aquaporin-4 gene disruption in mice reduces brain swelling and mortality in pneumococcal meningitis. *J. Biol. Chem.* **2005**, *280*, 13906–13912. [\[CrossRef\]](#) [\[PubMed\]](#)
26. Promeneur, D.; Lunde, L.K.; Amiry-Moghaddam, M.; Agre, P. Protective role of brain water channel AQP4 in murine cerebral malaria. *Proc. Natl. Acad. Sci. USA* **2012**, *110*, 1035–1040. [\[CrossRef\]](#)
27. Lennon, V.A.; Kryzer, T.J.; Pittock, S.J.; Verkman, A.S.; Hinson, S.R. IgG marker of optic-spinal multiple sclerosis binds to the aquaporin-4 water channel. *J. Exp. Med.* **2005**, *202*, 473–477. [\[CrossRef\]](#) [\[PubMed\]](#)
28. Gershen, L.D.; Zanotti-Fregonara, P.; Dustin, I.H.; Liow, J.-S.; Hirvonen, J.; Kreisl, W.C.; Jenko, K.J.; Inati, S.K.; Fujita, M.; Morse, C.L.; et al. Neuroinflammation in temporal lobe epilepsy measured using positron emission tomographic imaging of translocator protein. *JAMA Neurol.* **2015**, *72*, 882–888. [\[CrossRef\]](#) [\[PubMed\]](#)
29. Amor, S.; Puentes, F.; Baker, D.; Van Der Valk, P. Inflammation in neurodegenerative diseases. *Immunology* **2010**, *129*, 154–169. [\[CrossRef\]](#)
30. Troncoso-Escudero, P.; Parra, A.; Nassif, M.; Vidal, R.L. Outside in: Unraveling the role of neuroinflammation in the progression of Parkinson's disease. *Front. Neurol.* **2018**, *9*, 860. [\[CrossRef\]](#)
31. Li, L.; Zhang, H.; Varrin-Doyer, M.; Zamvil, S.S.; Verkman, A.S. Proinflammatory role of aquaporin-4 in autoimmune neuroinflammation. *FASEB J.* **2011**, *25*, 1556–1566. [\[CrossRef\]](#)
32. Martinez, F.O.; Gordon, S. The M1 and M2 paradigm of macrophage activation: Time for reassessment. *F1000Prime Rep.* **2014**, *6*, 13. [\[CrossRef\]](#)
33. Nau, G.J.; Richmond, J.F.L.; Schlesinger, A.; Jennings, E.G.; Lander, E.S.; Young, R.A. Human macrophage activation programs induced by bacterial pathogens. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 1503–1508. [\[CrossRef\]](#)
34. Prydz, A.; Stahl, K.; Puchades, M.; Davarpaneh, N.; Nadeem, M.; Ottersen, O.P.; Gundersen, V.; Amiry-Moghaddam, M. Subcellular expression of aquaporin-4 in substantia nigra of normal and MPTP-treated mice. *Neuroscience* **2017**, *359*, 258–266. [\[CrossRef\]](#)
35. Dauer, W.; Przedborski, S. Parkinson's disease: Mechanisms and models. *Neuron* **2003**, *39*, 889–909. [\[CrossRef\]](#)
36. Ofori, E.; Pasternak, O.; Planetta, P.J.; Burciu, R.; Snyder, A.; Febo, M.; Golde, T.E.; Okun, M.S.; Vaillancourt, D.E. Increased free water in the substantia nigra of Parkinson's disease: A single-site and multi-site study. *Neurobiol. Aging* **2015**, *36*, 1097–1104. [\[CrossRef\]](#)
37. Thrane, A.S.; Rappold, P.M.; Fujita, T.; Torres, A.; Bekar, L.K.; Takano, T.; Peng, W.; Wang, F.; Thrane, V.R.; Enger, R.; et al. Critical role of aquaporin-4 (AQP4) in astrocytic Ca²⁺ signaling events elicited by cerebral edema. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 846–851. [\[CrossRef\]](#)
38. Paxinos, G.A.F.K. *The Mouse Brain in Stereotaxic Coordinates*, 3rd ed.; Elsevier Inc.: San Diego, CA, USA, 2007.
39. Van Lookeren Campagne, M.; Oestreicher, A.B.; Buma, P.; Verkleij, A.J.; Gispen, W.H. Ultrastructural localization of adrenocorticotrophic hormone and the phosphoprotein B-50/growth-associated protein 43 in freeze-substituted, Lowicryl HM20-embedded mesencephalic central gray substance of the rat. *Neuroscience* **1991**, *42*, 517–529. [\[CrossRef\]](#)
40. Amiry-Moghaddam, M.; Lindland, H.; Zelenin, S.; Røberg, B.Å.; Gundersen, B.B.; Petersen, P.; Rinvik, E.; Torgner, I.A.; Ottersen, O.P. Brain mitochondria contain aquaporin water channels: Evidence for the expression of a short AQP9 isoform in the inner mitochondrial membrane. *FASEB J.* **2005**, *19*, 1459–1467. [\[CrossRef\]](#)
41. Wu, D.C.; Jackson-Lewis, V.; Vila, M.; Tieu, K.; Teismann, P.; Vadseth, C.; Choi, D.-K.; Ischiropoulos, H.; Przedborski, S. Blockade of microglial activation is neuroprotective in the 1-Methyl-4-Phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson disease. *J. Neurosci.* **2002**, *22*, 1763–1771. [\[CrossRef\]](#)
42. Davis, E.; Foster, T.; Thomas, W. Cellular forms and functions of brain microglia. *Brain Res. Bull.* **1994**, *34*, 73–78. [\[CrossRef\]](#)
43. Janova, H.; Böttcher, C.; Holtman, I.R.; Regen, T.; Van Rossum, D.; Götz, A.; Ernst, A.-S.; Fritsche, C.; Gertig, U.; Saiepour, N.; et al. CD14 is a key organizer of microglial responses to CNS infection and injury. *Glia* **2015**, *64*, 635–649. [\[CrossRef\]](#)

44. Cignarella, F.; Filipello, F.; Bollman, B.; Cantoni, C.; Locca, A.; Mikesell, R.; Manis, M.; Ibrahim, A.; Deng, L.; Benitez, B.A.; et al. TREM2 activation on microglia promotes myelin debris clearance and remyelination in a model of multiple sclerosis. *Acta Neuropathol.* **2020**, *140*, 513–534. [\[CrossRef\]](#)
45. Chen, G.; Zhou, Z.; Sha, W.; Wang, L.; Yan, F.; Yang, X.; Qin, X.; Wu, M.; Li, D.; Tian, S.; et al. A novel CX3CR1 inhibitor AZD8797 facilitates early recovery of rat acute spinal cord injury by inhibiting inflammation and apoptosis. *Int. J. Mol. Med.* **2020**, *45*, 1373–1384. [\[CrossRef\]](#)
46. Ewolf, Y.; Eyona, S.; Ekim, K.-W.; Jung, S. Microglia, seen from the CX3CR1 angle. *Front. Cell. Neurosci.* **2013**, *7*, 26. [\[CrossRef\]](#)
47. Jeetle, J.K.; Hagger, G.N.; Topps, S.S.; Male, D.; Rezaie, P. Microglial colonization of the developing mouse brain: The effect of CD11b deletion. *Neuropathol. Appl. Neurobiol.* **2002**, *28*, 164. [\[CrossRef\]](#)
48. Kalkonde, Y.V.; Morgan, W.W.; Sigala, J.; Maffi, S.K.; Condello, C.; Kuziel, W.; Ahuja, S.S.; Ahuja, S.K. Chemokines in the MPTP model of Parkinson's disease: Absence of CCL2 and its receptor CCR2 does not protect against striatal neurodegeneration. *Brain Res.* **2007**, *1128*, 1–11. [\[CrossRef\]](#)
49. Xue, X.; Zhang, W.; Zhu, J.; Chen, X.; Zhou, S.; Xu, Z.; Hu, G.; Su, C. Aquaporin-4 deficiency reduces TGF- β 1 in mouse midbrains and exacerbates pathology in experimental Parkinson's disease. *J. Cell. Mol. Med.* **2019**, *23*, 2568–2582. [\[CrossRef\]](#) [\[PubMed\]](#)
50. Lehrmann, E.; Kiefer, R.; Finsen, B.; Diemer, N.H.; Zimmer, J.; Hartung, H.P. Cytokines in cerebral ischemia: Expression of transforming growth factor beta-1 (TGF- β 1) mRNA in the postischemic adult rat hippocampus. *Exp. Neurol.* **1995**, *131*, 114–123. [\[CrossRef\]](#)
51. Spittau, B. Transforming growth factor β 1-mediated anti-inflammation slows progression of midbrain dopaminergic neurodegeneration in Parkinson's disease? *Neural Regen. Res.* **2015**, *10*, 1578–1580. [\[CrossRef\]](#)
52. Chen, S.; Luo, D.; Streit, W.J.; Harrison, J.K. TGF- β 1 upregulates CX3CR1 expression and inhibits fractalkine-stimulated signaling in rat microglia. *J. Neuroimmunol.* **2002**, *133*, 46–55. [\[CrossRef\]](#)
53. Farina, C.; Aloisi, F.; Meinl, E. Astrocytes are active players in cerebral innate immunity. *Trends Immunol.* **2007**, *28*, 138–145. [\[CrossRef\]](#)
54. Guedes, J.R.; Lao, T.; Cardoso, A.L.; El Khoury, J. Roles of microglial and monocyte chemokines and their receptors in regulating Alzheimer's disease-associated amyloid- β and Tau pathologies. *Front. Neurol.* **2018**, *9*, 549. [\[CrossRef\]](#)
55. Clarner, T.G.; Janssen, K.; Nellessen, L.; Stangel, M.; Skripuletz, T.; Krauspe, B.; Hess, F.-M.; Denecke, B.; Beutner, C.; Linnartz-Gerlach, B.; et al. CXCL10 triggers early microglial activation in the cuprizone model. *J. Immunol.* **2015**, *194*, 3400–3413. [\[CrossRef\]](#)
56. Dong, Y.; Yuan, Y.; Fang, Y.; Zheng, T.; Du, D.; Gao, D.; Du, J.; Liu, L.; He, Q. Effect of aquaporin 4 protein overexpression in nigrostriatal system on development of Parkinson's disease. *Int. J. Neurosci.* **2020**, 1–8. [\[CrossRef\]](#)
57. Tourdias, T.; Mori, N.; Dragonu, I.; Cassagno, N.; Boiziau, C.; Aussudre, J.; Brochet, B.; Moonen, C.T.; Klaus, P.; Dousset, V. Differential aquaporin 4 expression during edema build-up and resolution phases of brain inflammation. *J. Neuroinflamm.* **2011**, *8*, 143. [\[CrossRef\]](#)
58. Edwards, J.P.; Zhang, X.; Frauwirth, K.A.; Mosser, D.M. Biochemical and functional characterization of three activated macrophage populations. *J. Leukoc. Biol.* **2006**, *80*, 1298–1307. [\[CrossRef\]](#)
59. Vivekanantham, S.; Shah, S.; Dewji, R.; Dewji, A.; Khatri, C.; Ologunde, R. Neuroinflammation in Parkinson's disease: Role in neurodegeneration and tissue repair. *Int. J. Neurosci.* **2015**, *125*, 717–725. [\[CrossRef\]](#)
60. Lee, H.-J.; Suk, J.-E.; Patrick, C.; Bae, E.-J.; Cho, J.-H.; Rho, S.; Hwang, D.; Masliah, E.; Lee, S.-J. Direct transfer of α -synuclein from neuron to astroglia causes inflammatory responses in synucleinopathies. *J. Biol. Chem.* **2010**, *285*, 9262–9272. [\[CrossRef\]](#)
61. Hammond, S.L.; Bantle, C.M.; Popichak, K.A.; Wright, K.A.; Thompson, D.; Forero, C.; Kirkley, K.S.; Damale, P.U.; Chong, E.K.P.; Tjalkens, R.B. NF- κ B signaling in astrocytes modulates brain inflammation and neuronal injury following sequential exposure to manganese and MPTP during development and Aging. *Toxicol. Sci.* **2020**, *177*, 506–520. [\[CrossRef\]](#)
62. Zhang, J.; Yang, B.; Sun, H.; Zhou, Y.; Liu, M.; Ding, J.; Fang, F.; Fan, Y.; Hu, G. Aquaporin-4 deficiency diminishes the differential degeneration of midbrain dopaminergic neurons in experimental Parkinson's disease. *Neurosci. Lett.* **2016**, *614*, 7–15. [\[CrossRef\]](#)
63. Liang, R.; Yong, S.; Huang, X.; Kong, H.; Hu, G.; Fan, Y. Aquaporin-4 mediates the suppressive effect of Lipopolysaccharide on hippocampal neurogenesis. *Neuroimmunomodulation* **2016**, *23*, 309–317. [\[CrossRef\]](#)

64. Zhou, J.; Kong, H.; Hua, X.; Xiao, M.; Ding, J.; Hu, G. Altered blood–brain barrier integrity in adult aquaporin-4 knockout mice. *NeuroReport* **2008**, *19*, 1–5. [\[CrossRef\]](#)
65. Eilert-Olsen, M.; Haj-Yasein, N.N.; Vindedal, G.F.; Enger, R.; Gundersen, G.A.; Hoddevik, E.H.; Petersen, P.H.; Haug, F.-M.S.; Skare, O.; Adams, M.E.; et al. Deletion of aquaporin-4 changes the perivascular glial protein scaffold without disrupting the brain endothelial barrier. *Glia* **2011**, *60*, 432–440. [\[CrossRef\]](#)
66. García-Domínguez, I.; Veselá, K.; García-Revilla, J.; Carrillo-Jiménez, A.; Roca-Ceballos, M.A.; Santiago, M.; De Pablos, R.M.; Venero, J.L. Peripheral inflammation enhances microglia response and nigral dopaminergic cell death in an in vivo MPTP model of Parkinson’s disease. *Front. Cell. Neurosci.* **2018**, *12*, 398. [\[CrossRef\]](#)
67. Koizumi, S.; Shigemoto-Mogami, Y.; Nasu-Tada, K.; Shinozaki, Y.; Ohsawa, K.; Tsuda, M.; Joshi, B.V.; Jacobson, K.A.; Kohsaka, S.; Inoue, K. UDP acting at P2Y6 receptors is a mediator of microglial phagocytosis. *Nat. Cell Biol.* **2007**, *446*, 1091–1095. [\[CrossRef\]](#)
68. Pizzoni, A.; Bazzi, Z.; Di Giusto, G.; Alvarez, C.L.; Rivarola, V.; Capurro, C.; Schwarzbaum, P.J.; Ford, P. Release of ATP by TRPV4 activation is dependent upon the expression of AQP2 in renal cells. *J. Cell. Physiol.* **2020**. [\[CrossRef\]](#) [\[PubMed\]](#)
69. Benfenati, V.; Caprini, M.; Dovizio, M.; Mylonakou, M.N.; Ferroni, S.; Ottersen, O.P.; Amiry-Moghaddam, M. An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex is essential for cell-volume control in astrocytes. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 2563–2568. [\[CrossRef\]](#)
70. Meshorer, E.; Biton, I.E.; Ben-Shaul, Y.; Ben-Ari, S.; Assaf, Y.; Soreq, H.; Cohen, Y. Chronic cholinergic imbalances promote brain diffusion and transport abnormalities. *FASEB J.* **2005**, *19*, 910–922. [\[CrossRef\]](#)
71. Gupta, M.; Gupta, B.; Thomas, R.; Bruemmer, V.; Sladek, J.; Felten, D. Aged mice are more sensitive to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine treatment than young adults. *Neurosci. Lett.* **1986**, *70*, 326–331. [\[CrossRef\]](#)
72. Gupta, R.; Kanungo, M. Glial molecular alterations with mouse brain development and aging: Up-regulation of the Kir4.1 and aquaporin-4. *AGE* **2011**, *35*, 59–67. [\[CrossRef\]](#) [\[PubMed\]](#)
73. Noelker, C.; Morel, L.; Lescot, T.; Osterloh, A.; Alvarez-Fischer, D.; Breloer, M.; Henze, C.; Depboylu, C.; Skrzydelski, D.; Michel, P.P.; et al. Toll like receptor 4 mediates cell death in a mouse MPTP model of Parkinson disease. *Sci. Rep.* **2013**, *3*, 1393. [\[CrossRef\]](#)
74. Fiebich, B.L.; Batista, C.R.A.; Saliba, S.W.; Yousif, N.M.; De Oliveira, A.C.P. Role of microglia TLRs in neurodegeneration. *Front. Cell. Neurosci.* **2018**, *12*, 329. [\[CrossRef\]](#)
75. Mogi, M.; Harada, M.; Kondo, T.; Riederer, P.; Inagaki, H.; Minami, M.; Nagatsu, T. Interleukin-1 beta, interleukin-6, epidermal growth factor and transforming growth factor-alpha are elevated in the brain from parkinsonian patients. *Neurosci. Lett.* **1994**, *180*, 147–150. [\[CrossRef\]](#)
76. McGeer, P.L.; Kawamata, T.; Walker, D.G.; Akiyama, H.; Tooyama, I.; McGeer, E.G. Microglia in degenerative neurological disease. *Glia* **1993**, *7*, 84–92. [\[CrossRef\]](#)

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

PAPER III

Targeted deletion of the aquaglyceroporin AQP9 is protective in a mouse model of Parkinson's disease.

Stahl K, Rahmani S, Prydz A, Skauli N, MacAulay N, Mylonakou MN, Torp R, Skare Ø, Berg T, Leergaard TB, Paulsen RE, Ottersen OP, Amiry-Moghaddam M. *PLoS One* 2018. Mar 22;12(3):e0194896. doi: 10. 1371/journal.pone.0194896

RESEARCH ARTICLE

Targeted deletion of the aquaglyceroporin AQP9 is protective in a mouse model of Parkinson's disease

Katja Stahl¹, Soulmaz Rahmani¹, Agnete Prydz¹, Nadia Skauli¹, Nanna MacAulay², Maria N. Mylonakou³, Reidun Torp¹, Øivind Skare⁴, Torill Berg⁵, Trygve B. Leergaard¹, Ragnhild E. Paulsen⁶, Ole P. Ottersen^{1,7}, Mahmood Amiry-Moghaddam^{1*}

1 Division of Anatomy, Department of Molecular Medicine, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway, **2** Department of Neuroscience and Pharmacology, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark, **3** Centre for Molecular Medicine Norway, Nordic EMBL Partnership, Norway Biotechnology Centre, University of Oslo, Oslo, Norway, **4** Department of Occupational Medicine and Epidemiology, National Institute of Occupational Health, Oslo, Norway, **5** Division of Physiology, Department of Molecular Medicine, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway, **6** Department of Pharmaceutical Sciences, School of Pharmacy, University of Oslo, Oslo, Norway, **7** Karolinska Institutet, Stockholm, Sweden

* mahmo@medisin.uio.no

OPEN ACCESS

Citation: Stahl K, Rahmani S, Prydz A, Skauli N, MacAulay N, Mylonakou MN, et al. (2018) Targeted deletion of the aquaglyceroporin AQP9 is protective in a mouse model of Parkinson's disease. PLoS ONE 13(3): e0194896. <https://doi.org/10.1371/journal.pone.0194896>

Editor: Zhicheng Carl Lin, Harvard Medical School, UNITED STATES

Received: January 14, 2018

Accepted: March 12, 2018

Published: March 22, 2018

Copyright: © 2018 Stahl et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: The present work is funded by University of Oslo, Institute of Basic Medical Sciences. The funder approved the initial project description but had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abstract

More than 90% of the cases of Parkinson's disease have unknown etiology. Gradual loss of dopaminergic neurons of substantia nigra is the main cause of morbidity in this disease. External factors such as environmental toxins are believed to play a role in the cell loss, although the cause of the selective vulnerability of dopaminergic neurons remains unknown. We have previously shown that aquaglyceroporin AQP9 is expressed in dopaminergic neurons and astrocytes of rodent brain. AQP9 is permeable to a broad spectrum of substrates including purines, pyrimidines, and lactate, in addition to water and glycerol. Here we test our hypothesis that AQP9 serves as an influx route for exogenous toxins and, hence, may contribute to the selective vulnerability of nigral dopaminergic (tyrosine hydroxylase-positive) neurons. Using *Xenopus* oocytes injected with *Aqp9* cRNA, we show that AQP9 is permeable to the parkinsonogenic toxin 1-methyl-4-phenylpyridinium (MPP⁺). Stable expression of AQP9 in HEK cells increases their vulnerability to MPP⁺ and to arsenite—another parkinsonogenic toxin. Conversely, targeted deletion of *Aqp9* in mice protects nigral dopaminergic neurons against MPP⁺ toxicity. A protective effect of *Aqp9* deletion was demonstrated in organotypic slice cultures of mouse midbrain exposed to MPP⁺ *in vitro* and in mice subjected to intrastriatal injections of MPP⁺ *in vivo*. Seven days after intrastriatal MPP⁺ injections, the population of tyrosine hydroxylase-positive cells in substantia nigra is reduced by 48% in *Aqp9* knockout mice compared with 67% in WT littermates. Our results show that AQP9—selectively expressed in catecholaminergic neurons—is permeable to MPP⁺ and suggest that this aquaglyceroporin contributes to the selective vulnerability of nigral dopaminergic neurons by providing an entry route for parkinsonogenic toxins. To our knowledge this is the first evidence implicating a toxin permeable membrane channel in the pathophysiology of Parkinson's disease.

Abbreviations: AQP4, aquaporin-4; AQP9, aquaglyceroporin-9; DAB, 3,3'-diaminobenzidine; DAT, dopamine transporter; DMSO, dimethyl sulfoxide; DOPAC, 3,4-dihydroxyphenylacetic acid; EGFP, enhanced green fluorescent protein; HEK, human embryonic kidney cells 293; HPLC, High performance liquid chromatography; HVA, homovanillic acid; MPP⁺, 1-methyl-4-phenylpyridinium; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; PBS, phosphate buffered saline; SN, substantia nigra; SNpc, substantia nigra pars compacta; SNpr, substantia nigra pars reticulata; TH, tyrosine hydroxylase; VTA, ventral tegmental area; WT, wild type; YFP, yellow fluorescent protein.

Introduction

Parkinson's disease is one of the most common neurodegenerative disorders, affecting 1–2% of the population over 50 years [1–4]. This disease is characterized by loss of dopaminergic neurons in the SNpc with decreased dopamine levels in the basal ganglia [5], and is clinically manifested by motor dysfunction, including bradykinesia, resting tremor, rigidity and postural instability, and non-motor symptoms including, but not limited to, cognitive impairment, mood disorders, sleep disorders, constipation, bladder dysfunction and loss of sense of smell [6]. Less than 10% of the cases have a strict familial etiology [7]. For sporadic Parkinson's disease, the cause of the selective dopaminergic cell death remains unknown, thus restricting the development of drugs which may halt, stop or reverse the degenerative process. However, research has identified a possible link to environmental factors targeting mitochondria, and Parkinson's disease has been associated with exposure to herbicides, arsenite, and other environmental toxins [8–21]. Mechanisms for the selective vulnerability of dopaminergic neurons to such toxins remain to be elucidated, although there is ample evidence that toxin uptake may occur through the dopamine transporter [22].

Studies have shown that AQP9 like the dopamine transporter DAT, is expressed in TH-positive neurons of the SN in rat, mouse and primates [23–26]. At the subcellular level, AQP9 is localized in the plasma membrane as well as in the inner mitochondrial membrane [23,26]. AQP9 belongs to the aquaglyceroporin subfamily of water channels and is permeable to a broad range of substrates including glycerol and urea [27–29], monocarboxylates (lactate and β -hydroxybutyrate) [28], purines [28], ammonia [30] and arsenite [31].

Here, we show that the parkinsonogenic toxin MPP⁺ permeates *Xenopus* oocytes injected with *Aqp9* cRNA. Second, we demonstrate that stable expression of AQP9 in HEK cells increases their vulnerability to this toxin and to arsenite—another toxin associated with PD. Third, we provide data suggesting that the expression of this water channel contributes to the selective vulnerability of nigral dopaminergic neurons in mice. Specifically, after intrastriatal injections of MPP⁺, *Aqp9*^{-/-} mice retain a higher number of nigral cells positive for TH than do WT littermate controls. This is the first evidence suggesting that dopaminergic neurons are equipped with a plasma membrane channel that contributes to their vulnerability. Our findings may open new avenues for prevention and therapy.

Materials and methods

Heterologous expression in *Xenopus laevis* oocytes and radio-isotope permeability

Oocytes were surgically removed from *Xenopus laevis* frogs (Nasco, USA or National Center for Scientific Research, France) anesthetized with 2 g/l Tricain, 3-aminobenzoic acid ethyl ester, (Sigma-Aldrich A-5040) as previously described [32]. The protocol complies with the European Community guidelines for the use of experimental animals and the experiments were approved by The Danish National Committee for Animal Studies (2015-15-0201-00560).

The cDNA encoding rAqp4 and rAqp9 (obtained from Søren Nielsen, Aalborg University) were subcloned into the expression vector pXOOM [33]. The constructs were linearized down-stream from the poly-A segments and *in vitro* transcribed to cRNA using mMessage mMachin T7 transcription kit (Ambion, Roskilde, Denmark) followed by transcript purification using MEGAclear (Ambion). Oocytes were micro-injected with 50 ng cRNA with a

Nanoject micro injector (Drummond Scientific, Broomall, PA, USA) and kept in Kulori medium at 18°C for 3–6 days prior to experiments.

Five to ten uninjected, AQP4-, or AQP9-expressing oocytes were placed in 0.5 ml uptake solution (in mM: 100 NaCl, 2 KCl, 1 CaCl₂, 1 MgCl₂, 10 HEPES (pH 7.4)), and either ¹⁴[C] urea (1 μCi/ml, 9.25 MBq/ml, Perkin Elmer, MA, USA) or ³[H]MPP⁺ (3 μCi/ml, 1.85 MBq/ml (GE Healthcare, Little Chalfont, England), and incubated with slight agitation at room temperature for 3 minutes. The oocytes were washed in wash solution (in mM: 100 choline chloride, 2 KCl, 1 CaCl₂, 1 MgCl₂, 10 HEPES, pH 7.4) to remove residual isotope, and subsequently lysed individually in scintillation vials with 200 μl SDS. Scintillation fluid (Opti-Fluor, Perkin Elmer, Skovlunde, Denmark) was added and the radioactive samples counted (Tri-Carb 2900TR, Perkin Elmer). Data were obtained as counts per minute (CPM)/oocyte and averaged for each construct. As basal uptake displays batch variation, the average CPM/oocyte for each batch of oocytes was normalized to that obtained for AQP4-expressing oocytes and averaged across all seven experiments.

Cell cultures and MTT cell viability assay

Native HEK cells (ATCC, Manassas, VA) and HEK cells stably transfected with YFP-*hDAT* [34] were cultured in Dulbecco's Modified Eagle Medium high glucose (Gibco Life Technologies, Paisley, UK) with 10% fetal bovine serum (Biowest, Nuaillé, France) and 1% Penicillin-streptomycin (Gibco). For HEK cells stably transfected with tetracycline inducible EGFP-*hAQP9* (pTRE-EGFP-*hAQP9*) [35,36], expression of EGFP-*hAQP9* was induced using 1 μg/ml doxycycline (Clontech Labs, Takara, CA, USA) overnight, and 0.1 mg/ml Hygromycin B (Invitrogen, Carlsbad, CA, USA) and 0.3 mg/ml G418 Geneticin (Gibco) were added for maintenance of the expression.

To confirm protein expression of AQP9 and DAT, WT HEK293 and stably transfected EGFP-*hAQP9* or YFP-*hDAT* HEK293 cells were plated on coverslips, the cell layer washed with 0.01 M PBS after 24 hours, then fixed in 4% formaldehyde and permeabilized with 0.1% Triton X-100 (Sigma Aldrich, St. Louis, MO, USA) in PBS. Following blocking with 10% normal donkey serum in PBS, cells were incubated with the primary antibodies rabbit anti-AQP9 (1:500, AQP91-A, Alpha Diagnostics, San Antonio, TX, USA) or rat anti-DAT (1:800, Merck Millipore, Billerica, MA, USA), and then with donkey anti-rabbit Cy3 or anti-rat Cy2 secondary antibodies (Jackson ImmunoResearch Laboratories, West Grove, USA) diluted 1:1000. Coverslips were mounted with EverBrite™ Hardset Mounting Media with DAPI (4',6-diamidino-2-phenylindole) (Biotium Inc., Fremont, CA, USA). A Zeiss LSM5 PASCAL confocal microscope was used to acquire the images.

HEK293 and YFP-*hDAT* expressing cells were seeded in 96-well plates at a density of 25 000–35 000 per well. HEK293 cells stably transfected with EGFP-*hAQP9* were seeded in media containing 1 μg/mL doxycycline (Clontech Labs, Mountain View, CA, USA) for induction of AQP9 expression at a density of 50 000 cells per well due to impaired growth. The cultures were grown for 24 hours before MPP⁺ iodide (D048, Sigma Aldrich) or arsenite (S7400, Sigma) was added in 100 μL medium at various concentrations.

Cell metabolic activity was assessed by the MTT assay. After 24 hours incubation, 100 μL 0.5 mg/mL MTT reagent was added to each well in the respective media and the plate incubated at 37°C 5% CO₂ for 3 hours. MTT-containing medium was replaced with 100 μL DMSO (Sigma) and incubated for 30 minutes at room temperature.

Absorbance was measured at 570 nm with a CLARIOstar plate reader (BMG Labtech, Ortenberg, Germany). Blank well values without MTT were subtracted as background.

Animals

All mice were bred and maintained at the animal facility of the Institute of basic medical sciences, University of Oslo, Norway. *Aqp9*^{-/-} mice were a kind gift from Søren Nielsen's laboratory, University of Aarhus, Denmark [37]. *Aqp9*^{-/-} mice backcrossed onto a C57BL/6J background (Jacksons Laboratories) and littermate WT mice of mixed gender were used for organotypic cultures (5 days old) and stereotaxic surgery (2–10 months old). All animal experiments were prospectively approved by The Norwegian Animal Research Authority (NARA), project license no FOTS 3730 and 4012, and conducted in accordance with the European Directive 2010/63/EU. Animals were maintained under standard animal facility conditions on a 12 hour light/dark cycle, with food and water available *ad libitum*.

Organotypic slice cultures of the ventral mesencephalon

WT (n = 8) and *Aqp9*^{-/-} (n = 7) mice were used to prepare organotypic cultures of ventral mesencephalon as previously described [38]. In brief, animals were anesthetized with isoflurane upon decapitation, and the ventral midbrain was isolated and sectioned into slices of 400 μ m. The SNpc was unilaterally exposed to either 30 μ M MPP⁺ (Sigma Aldrich) diluted in 0.01 M PBS, 60 μ M MPP⁺ in PBS followed by 100 μ M phloretin (Sigma) diluted in DMSO (Sigma) or only PBS (control) [38,39]. Following fixation in 4% formaldehyde (Sigma Aldrich) overnight, slices were incubated with rabbit anti-TH (1:1000, Chemicon, Billerica, MA, USA) overnight, and then with Cy3 conjugated donkey anti-rabbit (1:1000, Jackson). Images from the SNpc were collected with a LSM 5 Pascal Confocal Microscope (Zeiss, Oberkochen, Germany) using a 20 $\times$ objective, as previously described [39]. Cells immunostained for TH were counted using ImageJ.

Intrastriatal injections of MPP⁺

C57BL/6J WT (n = 34) and *Aqp9*^{-/-} mice (n = 29) were deeply anesthetized with zoletil mixture (Zoletil Forte (250 mg/ml), Rompun (20 mg/ml) and Fentanyl (50 μ g/ml); 0.1 ml/10 g; intraperitoneally) and then subjected to stereotaxic (TSE systems, Bad Homburg, Germany) intrastriatal injections of MPP⁺ (7.5 μ g dissolved in saline) or saline. 1 μ l MPP⁺ solution or saline was injected into the striatum 0.6 mm anterior to Bregma, 2.2 mm laterally and 3.2 mm ventrally [40] at 12 μ l/hr using a syringe pump (Kd Scientific, Holliston, MA, USA). MPP⁺ was protected against light during the procedure. The injector was left for 5 minutes to allow diffusion before suture. A heating pad (PanLab, Barcelona, Spain) was used to maintain the body temperature at 36.5 $\pm$ 0.5°C during the surgery, and the eyes were covered with Vaseline to avoid drying of the cornea.

Post-operative animal care, behavioral assessments and humane endpoints

The first 24 hrs post-surgery, all animals (total n = 84, MPP⁺: n = 63, saline: n = 21) were put on a heating pad overnight and were treated with Rimadyl (0.1 ml/10 g; subcutaneously every 12 hours) to prevent pain, and a mixture of 0.9% saline and 0.9% sucrose (subcutaneously, every 24 hours) to restore water and energy balance. Animals were fed moisture pellets to improve food- and water uptake. This treatment was continued for as long as needed until the animals were sacrificed 7 days after surgery.

All animals were scored daily to evaluate their post-operative condition. Briefly, animals were assessed daily on a scale from 0–3 for their clinical appearance, including a) weight loss (0%, <5%, 5–10%, 10–15%) b) inactivity (normal, minor changes, less mobile and isolated, inactive) c) external appearance (normal, reduced self-care, nose/eye redness and discolored

fur, curved posture and bristling fur) and d) reaction patterns to external stimuli (normal, slightly reduced, moderately reduced and less alert, weak or precomatose) giving a total score ranging from 0–12. Animals reaching a score of ≥ 10 were euthanized immediately, whereas the remaining animals were sacrificed at post-operative day seven (see following sections). The animal welfare protocol was established in collaboration with the leading veterinary at the animal department of the host institution.

Apomorphine rotation test. Six days after the lesion, the animals ($n = 70$) were treated with apomorphine [41]. The animal was placed in a circular bowl (diameter 20 cm) for 5 minutes to adapt to the environment, then subcutaneously injected with an ice cold and light protected solution of 0.1 mg/kg apomorphine and 0.2 mg/ml ascorbic acid in saline (all from Sigma Aldrich). The animal was video recorded for 15 minutes immediately after the injection. Animals were scored for 360° rotational behavior ipsilaterally and contralaterally to the injection. Data are expressed as net ipsilateral turns (ipsilateral turns—contralateral turns).

Histological and stereological evaluation of MPP⁺ lesions and TH-positive neurons

Seven days post-surgery, animals ($n = 27$) were deeply anaesthetized (zoletil mixture; 0.1 ml/10 g; intraperitoneally) and perfusion-fixed with 4% formaldehyde. Brains were dissected, post-fixed, cryo-protected and coronally sectioned into 40 μm . Sections through striatum and midbrain were stained with hematoxylin/eosin for histological analysis (S1 Fig). Successive sections through the SN were immunostained with mouse anti-TH (1:1000, Chemicon) followed by incubation with Streptavidin-Biotinylated horseradish peroxidase complex (1:100; GE Healthcare). Unbiased stereological counting of TH-positive neurons was performed bilaterally in the SNpc, SNpr, and VTA of *Aqp9*^{-/-} and WT mice injected with MPP⁺ (*Aqp9*^{-/-}, $n = 8$; WT, $n = 7$) or saline (*Aqp9*^{-/-}, $n = 3$; WT, $n = 3$) as described elsewhere [42,43]. A detailed description of the stereological counting procedure is provided in S1 File.

High-performance liquid chromatography

HPLC was used to measure the amount of dopamine and its metabolites HVA and DOPAC. WT and *Aqp9*^{-/-} animals ($n = 6$ for each genotype) were anesthetized with isoflurane and sacrificed by decapitation seven days post-surgery, and the striatum of each hemisphere (MPP⁺ injected and control) were dissected out, flash frozen in liquid N₂ and stored at -80°C. The tissue samples were homogenized in 250 μl 0.2 M perchloric acid (4°C), mixed with an equal amount of 0.454 μM 2,3-dihydroxybenzoic acid containing 0.135 mM ascorbic acid and centrifuged (20,800 g, 20 minutes, 4°C). The pellet was dissolved in 0.6 ml 0.1M NaOH and protein concentration measured by the BCA Assay (Thermo Scientific, Waltham, MA, USA). The supernatant was mixed with an equal volume of hexane, and allowed 15 minutes to separate (4°C). The bottom layer was collected and filtered through a 0.2 μm nylon filter and stored at -80°C until analyzed for dopamine on a Shimadzu HPLC monoamines analyzer system. The samples were run in duplicates at an isocratic flow rate of 0.8 ml/min.

RNA isolation and real time qPCR

Seven days post-surgery, animals (MPP⁺, $n = 7$ for each genotype; saline, $n = 3$ for each genotype) were decapitated under isoflurane anesthesia. The brains were dissected out and cut along the midsagittal line into the injected and control hemisphere. Midbrain, striatum and neocortex were dissected out from both hemispheres and snap frozen. Total RNA was isolated from the regional brain samples using the RNeasy Mini Kit (QIAGEN, Hilden, Germany) and

the relative change in gene expression ($\Delta\Delta\text{Ct}$) in the MPP⁺-treated versus control hemispheres of WT and *Aqp9*^{-/-} mice, was assessed as described elsewhere [44]. List of the TaqMan probes used in the study are shown in [S1 Table](#) and a detailed description of the procedure is available under [S1 File](#).

Statistics

For stereological, HPLC and qPCR analyses, Student's *t*-test and Mann–Whitney U tests were performed using SPSS Statistics (Release 24.0.0.2, IBM Corporation, USA). For oocyte experiments, differences between groups were determined with one-way ANOVA and Dunnett's *post hoc* test in SPSS.

For MTT assays, statistical analysis was performed in SPSS and GraphPad Prism (Ver. 7.03, GraphPad Software, Inc.). OD570 nm was expressed as percentage of the untreated controls. Statistical testing was performed by one-way ANOVA with LSD *post hoc* tests. Additionally, independent samples *t*-tests were performed comparing each group. Inhibitory concentration 50% values for arsenite were calculated by nonlinear regression, log(inhibitor) vs response (three parameters) and a curve fitted. For log transformed data, the concentration 0 was set to 1 nM.

The behavioral tests were analyzed in R (version 3.2.2). For apomorphine rotation test, ordinary linear regression was used where interaction terms between genotype and treatment were added. To obtain model robust estimates of standard errors, *p*-values and confidence intervals, we used the *sandwich* function (*sandwich* package, version 2.3–3) in combination with the *coefest* function (*lmtest* package, version 0.9–34). For clinical appearance, a linear mixed model was used, where interaction terms were used to see how the treatment (MPP⁺ vs saline) effect differs between genotypes (*Aqp9*^{-/-} vs WT). The mixed model included a random intercept for animals. The variance of random effect was allowed to vary with treatment. The mixed model allowed the variance of the residual term to vary between treatments, and between days.

For the organotypic data, a linear mixed model was applied. The observations come in pairs, one observation for each hemisphere of each slice, where one of the hemispheres is treated (with MPP⁺ or PBS) and the other is untreated. The effect of the treatment is seen as the difference in outcome value between these two sides. An interaction term between treatment, side and genotype was used to estimate how the treatment effect of MPP⁺ and PBS differed between the two genotypes. The outcome values were ln-transformed prior to analysis to obtain a better fit to the normal distribution. Random effects were added for section and slide, with slide nested in section. The residual variance was allowed to vary with treatment and side.

Data were analyzed using the function *lme* (*nlme* package, version 3.1–122). A *p*-value <0.05 was considered statistically significant.

Experimental design

Data was blinded for analyses susceptible to human bias. For cell quantitation analysis of the organotypic cultures, stereology and behavior tests the investigator was blinded to the groups by third party concealment of treatments.

A minimum of six animals per group was considered as a minimum to estimate statistical power due to within-group variations. The actual number of included animals was calculated based on the high mortality rates following exposure to MPP⁺.

Results

AQP9 but not AQP4 is permeable to MPP⁺

To determine if MPP⁺ is able to permeate AQP9, we expressed AQP9 in *Xenopus* oocytes and exposed these oocytes to radiolabeled MPP⁺ (³[H]MPP⁺). As controls, we included uninjected and AQP4-expressing oocytes and determined in parallel uptake of ¹⁴[C]-urea which is known to permeate AQP9 but not AQP4 [28,45]. Following ³[H]MPP⁺ exposure, AQP9-expressing oocytes showed a twofold higher level of radioactivity than AQP4-expressing oocytes (Fig 1). An even larger difference between the two groups of oocytes was found after incubation with ¹⁴[C]-urea (ratio AQP9 vs AQP4 containing oocytes: 5.51 ± 0.79). Uninjected and AQP4-expressing oocytes did not differ in regard to the level of accumulated ³[H]-MPP⁺ or ¹⁴[C]-urea (Fig 1). In parallel experiments comparing the swelling rates of AQP4-expressing to native oocytes, we confirmed the functionality of the expressed AQP4.

Stable expression of AQP9 or DAT in HEK cells increases their vulnerability to MPP⁺

We compared the toxin sensitivity of HEK293 cells stably expressing YFP-*hDAT* with that of cells stably expressing EGFP-*hAqp9* (Fig 2A and 2B). Neither DAT nor AQP9 is normally expressed in HEK293 cells. Expression of EGFP-*hAQP9* made these cells sensitive to MPP⁺, as did a stable expression of YFP-*hDAT* (Fig 2C–2G). For *hAQP9* expressing cells, a statistically significant toxic effect was observed already at 0.1 μM MPP⁺ (Fig 2G). By comparison, an MPP⁺ concentration of 1 μM was required to exert a significant toxic effect on *hDAT* expressing cells (Fig 2F). HEK293 cells expressing *hAQP9* were clearly more sensitive to the parkinsonogenic compound arsenite than were HEK293 cells expressing *hDAT* (Fig 3).

Aqp9 gene deletion protects nigral dopaminergic cells against MPP⁺ neurotoxicity *ex vivo*

Organotypic slice cultures containing SN were used to assess the effect of MPP⁺ on neurons expressing TH (Fig 4A). To resolve whether knockout of the *Aqp9* gene conferred protection, MPP⁺ was applied unilaterally to midbrain slice cultures from WT and *Aqp9*^{−/−} mice (Fig 4). In slices of WT animals, application of MPP⁺ led to loss of TH-positive neurons on the ipsilateral side (Fig 4B and 4F). Only a few immunopositive elements remained, and most of these were delicate processes that were scattered in the neuropil of SNpc (Fig 4B). In contrast, MPP⁺ did not cause any obvious damage or loss of TH-positive neurons in slices obtained from *Aqp9*^{−/−} mice (Fig 4D). In the latter mice, there was no significant difference in the number of TH-positive neurons between the ipsi- and contralateral sides (Fig 4G). The density of TH-positive neurons contralateral to MPP⁺ exposure was lower in slices of WT mice than in slices of *Aqp9*^{−/−} mice. This might be attributed to spillover of the toxin to the contralateral side.

Application of phloretin—a blocker of AQP9—mimicked the effect of *Aqp9* gene deletion. Thus, in the presence of 100 μM phloretin, MPP⁺ did not cause any significant loss of TH-positive neurons (Fig 4H). Application of PBS in lieu of MPP⁺ did not cause any cell loss (Fig 4I). In sum, these experiments indicated that deletion of *Aqp9* protects against MPP⁺ toxicity *in vitro*.

Aqp9 gene deletion protects nigral dopaminergic cells against MPP⁺ neurotoxicity *in vivo*

Stereological cell counts in the ventral midbrain. *Aqp9*^{−/−} and littermate WT mice were subjected to unilateral intrastriatal injections of MPP⁺ or saline (control). The density of

Fig 1. *Xenopus* oocytes expressing AQP9 reveal higher uptake of MPP $^{+}$. *Xenopus* oocytes expressing AQP4 or AQP9 and uninjected oocytes were exposed to ^{14}C urea or ^3H MPP $^{+}$. Data were obtained as counts per minute (CPM)/oocyte and averaged for each construct. The uptake was normalized to that of the AQP4-expressing oocytes prior to averaging across the $n = 7$ experiments with five to ten oocytes included for each experimental condition. Compared with uninjected oocytes and AQP4 expressing oocytes, oocytes expressing AQP9 accumulate significantly higher amounts of urea as well as MPP $^{+}$. Bars are mean $\pm$ SEM; * $p < 0.05$, *** $p < 0.001$.

<https://doi.org/10.1371/journal.pone.0194896.g001>

dopaminergic neurons was quantified in the SNpc, SNpr and VTA of *Aqp9* $^{-/-}$ ($n = 8$ for MPP $^{+}$, $n = 3$ saline) and WT animals ($n = 7$ for MPP $^{+}$, $n = 3$ saline) by unbiased stereological counting of TH-positive cells in each region [46]. Cell counts for SNpc and VTA are shown in the S2 Table. Gundersen coefficients of error $m = 1$; [43] ranged from 5–9% for both SNpc and VTA.

Fig 2. HEK293 cells expressing *hAQP9* or *hDAT* reveal higher sensitivity to MPP⁺. A-B) Immunofluorescence images of HEK293 cells expressing EGFP-*hAQP9* (A) or YFP-*hDAT* (B) grown on coverslips confirm plasma membrane localization of the respective constructs (identified by antibodies to AQP9 or DAT). The cells were counterstained with Hoechst to visualize nuclei. C-G) Native HEK293 cells and HEK293 cells expressing EGFP-*hAQP9* or YFP-*hDAT* were grown in 96-well plates and exposed to different concentrations of MPP⁺ (four wells for each concentration). Cell viability was assessed after 24 hours using the MTT assay. Data were collected from independent plates ($n = 3$ for each construct) and normalized to respective untreated cells. Native HEK293 cells show sensitivity to MPP⁺ only at very high concentrations ($\sim 100 \mu\text{M}$). Cells expressing *hDAT* become sensitive at $1 \mu\text{M}$ MPP⁺, compared with $0.1 \mu\text{M}$ for cells expressing *hAQP9*. Overlay of the dose/response curve for the three groups (D) and the individual curves for native HEK293 cells (E), YFP-*hDAT* expressing (F) and EGFP-*hAQP9* expressing HEK293 cells (G) are shown. Bars are mean $\pm$ SEM. Asterisks: significantly different from untreated controls; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; crosses: significantly different from previous data point: ++ $p < 0.01$.

<https://doi.org/10.1371/journal.pone.0194896.g002>

Fig 3. HEK293 cells expressing EGFP-*hAQP9* are more sensitive to arsenite than HEK293 cells expressing YFP-*hDAT*. A-C) Native HEK293 cells and HEK293 cells expressing EGFP-*hAQP9* or YFP-*hDAT* were grown in 96-well plates and exposed to different concentrations of arsenite (eight wells for each concentration). Cell viability was assessed after 24 hours using the MTT assay. Data were collected from independent plates ($n = 3$ for each construct) and normalized to respective untreated cells. Both EGFP-*hAQP9* and YFP-*hDAT* expressing cells showed higher sensitivity to arsenite, than native HEK293 cells, with EGFP-*hAQP9* cells being the most sensitive. At the arsenite concentration of $10 \mu\text{M}$, stably transfected EGFP-*hAQP9* were the only cells showing toxin sensitivity (A). The curve showing IC₅₀ values for arsenite calculated by nonlinear regression, log(inhibitor) vs response (three parameters) is shown (B). For log transformed data, the concentration 0 was set to 1 nM . Comparison of the IC₅₀ values shows a significantly lower IC₅₀ value for the HEK293 cells expressing EGFP-*hAQP9* compared to the native HEK293 cells or HEK293 cells expressing YFP-*hDAT* (C). Bars are mean $\pm$ SEM. Asterisks: significantly different from untreated controls; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; crosses: significantly different from previous data point: ++ $p < 0.01$.

<https://doi.org/10.1371/journal.pone.0194896.g003>

Fig 4. Deletion of *Aqp9* protects against MPP⁺ toxicity *in vitro*. A) Photomicrograph (left) and schematic representation (right) of a representative midbrain slice. Site of MPP⁺ application is indicated in red. B-E) Immunofluorescence staining of representative midbrain slices from WT (B, C) and *Aqp9*^{-/-} mice (D, E) showing TH-positive neurons in the MPP⁺ treated side (ipsilateral; B, D), and in the control side (contralateral, C, E). Note the extensive loss of ipsilateral TH-positive neurons in the WT slice (B). F-H) Quantitative analyses of the TH-positive cell count in slices treated with 30 μ M MPP⁺ (F, G) show significant loss of TH-positive neurons in the ipsilateral hemisphere of the WT slice ($n = 8$) (F). No significant difference was observed between the ipsi- and contralateral hemisphere of *Aqp9*^{-/-} slices treated with 30 μ M MPP⁺ ($n = 7$) (G), WT slices treated with a combination of 60 μ M MPP⁺ and 100 μ M phloretine ($n = 11$) (H), or WT slices treated with saline ($n = 8$) (G). In WT mice, the TH-positive cell count contralateral to MPP⁺ application (F) was lower than the TH-positive cell count in saline treated slices (H), suggesting spillover of MPP⁺ from the ipsilateral side. RRF: retrorubral field; Aq: aqueduct; V: ventricle; A9: population of dopaminergic neurons. Bars are mean $\pm$ SEM; *** $p < 0.001$. Scale bar: 100 μ m; scale bar inset: 50 μ m.

<https://doi.org/10.1371/journal.pone.0194896.g004>

Both genotypes showed a significant reduction in the number of TH-positive neurons in the ipsilateral SNpc following MPP⁺ injections (S2 Table, Fig 5A–5C). In the ipsilateral SNpc, the reduction in TH cell number (using the cell count in the contralateral hemisphere as reference) was significantly lower in *Aqp9*^{-/-} animals than in WT littermates (47.59% and 67.02%, respectively; $p < 0.001$) (Fig 5B). In the ipsilateral VTA we found a small but significant reduction in TH-positive neurons in WT animals, but no reduction in *Aqp9*^{-/-} mice (Fig 5E). As expected, SNpr contained few TH-positive cells in either mouse line. No significant change was observed in animals injected with saline, regardless of genotype ($p > 0.05$) (Fig 5D).

Clinical appearance and behavioral assessment. All animals (MPP⁺: $n = 63$; saline: $n = 21$) were evaluated daily and given a score based on post-operative clinical condition, where increasing scores reflected progressive symptoms of disease. Following unilateral striatal treatment with MPP⁺, both *Aqp9*^{-/-} and WT mice showed a higher score than control animals injected with saline (Fig 6A). In total, 10 MPP⁺ treated animals (6 WT and 4 *Aqp9*^{-/-}) were euthanized ($n = 4$) or died during the three days due to the toxin ($n = 6$) and were not included in the analysis.

Animals were tested for ipsilateral rotation behavior day six post-surgery after systemic treatment with the dopamine agonist apomorphine. The frequency of rotations is expected to correlate with the extent of dopaminergic cell loss [41]. Significant differences were observed in net rotations (ipsilateral minus contralateral turns) between animals treated with MPP⁺ ($n = 53$) and saline controls ($n = 17$). The significance levels were $p < 0.001$ for WT and $p = 0.007$ for *Aqp9*^{-/-}. In the MPP⁺ treated group, net rotational behavior was significantly lower for *Aqp9*^{-/-} animals than for WT littermates (2.69 ± 1.91 and 6.33 ± 2.15 , respectively ($p = 0.014$)) (Fig 6B).

HPLC of dopamine and dopamine metabolites. The levels of dopamine and its metabolites HVA and DOPAC were quantified by HPLC in tissue samples from the ipsi- and contralateral striatum in *Aqp9*^{-/-} ($n = 6$) and WT littermates ($n = 6$), seven days after intrastriatal injections of MPP⁺ (Fig 7). MPP⁺ significantly reduced the mean concentration of dopamine (ng/mg protein) on the ipsilateral hemisphere, compared to the contralateral side, by ~83% in WT ($p = 0.004$) and ~74% in *Aqp9*^{-/-} mice ($p = 0.005$) (Fig 7A). In WT mice, the reductions in HVA and DOPAC were ~68% ($p = 0.003$) and ~79% ($p = 0.009$), respectively. In *Aqp9*^{-/-} mice, metabolite levels were reduced to ~57% ($p = 0.002$) for HVA and ~57% ($p = 0.009$) for DOPAC (Fig 7B and 7C). The ipsilateral reduction in HVA was significantly more pronounced in WT mice than in *Aqp9*^{-/-} mice ($p = 0.038$) (Fig 7B). The same tendency was observed for dopamine and DOPAC although the difference was not significant ($p = 0.063$ and 0.059 , respectively) (Fig 7A and 7C).

Gene expression levels in *Aqp9*^{-/-} animals

In the knockout animals, the *Aqp9* mRNA level was close to the detection limit, as expected. In WT animals, semi-quantitative real time PCR revealed significantly higher *Aqp9* mRNA levels in the midbrain and striatum (79% of the level in midbrain, $p = 0.053$) than in the neocortex (59% of the level in midbrain, $p = 0.002$) (Fig 8A and 8B).

To resolve whether the reduced dopaminergic cell count in the *Aqp9*^{-/-} mice could be attributed to compensatory up- or downregulation of other genes, we analyzed the expression of 12 genes (*Aqp4*, *Gfap*, *Kir4.1*, *mTOR*, *Prph*, *Cat*, *Ppard*, *Slc6a3* (DAT), *Drd2*, *Bcl2*, *Bax* and *Sod2*; Fig 6C). Only one of these genes (peripherin, *Prph*) showed an increase in expression level (~30%) after *Aqp9* deletion. For five genes (*Slc6a3* (DAT), *Drd2*, *Bcl2*, *Bax* and *Sod2*), mRNA levels were assessed in the ipsi- and contralateral midbrain and striatum of animals injected with MPP⁺. The relative levels of these transcripts did not differ between *Aqp9*^{-/-} and WT

Fig 5. Deletion of *Aqp9* protects against MPP⁺ toxicity *in vivo*. A) Representative midbrain sections of WT and *Aqp9*^{-/-} mice, seven days after treatment with unilateral striatal injection of 7.5 μ g MPP⁺. Cell bodies and processes of the dopaminergic neurons are identified by TH immunostaining. The visible reduction of the TH immunostained cell bodies in the ipsilateral SNpc is more pronounced in WT than in *Aqp9*^{-/-} mice. B) Quantitation of the TH-positive neurons in SNpc is described in the text. TH-positive cell loss, calculated as [(n contra – n ipsi): (n contra)] is significantly lower in *Aqp9*^{-/-} mice, where 47.59% of the cells are lost, compared to 67.02% in WT littermates ($p < 0.001$). C) Compared with WT mice, *Aqp9*^{-/-} mice show a significantly higher count of TH-positive neurons on the injected side ($p < 0.001$). D) Animals treated with saline showed no significant loss of TH-positive cells in SNpc, regardless of genotype. E) In WT mice, VTA showed a slight but statistically significant decrease in number of TH-positive neurons on the ipsilateral side ($p = 0.033$). No change was observed in VTA of *Aqp9*^{-/-} mice. The MPP⁺ treated group consisted of 15 animals (*Aqp9*^{-/-}, $n = 8$; WT, $n = 7$), and the saline group consisted of six animals (*Aqp9*^{-/-}, $n = 3$; WT, $n = 3$). ML, medial lemniscus. Bars are mean $\pm$ 2 SEM; $n = 6$; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Scale bar, 1000 μ m; scale bar inset, 20 μ m.

<https://doi.org/10.1371/journal.pone.0194896.g005>

Fig 6. Clinical appearance and behavioral tests of WT and *Aqp9*^{-/-} with unilateral intrastratial MPP⁺ injection. A) Clinical appearance of animals following intrastratial injections of MPP⁺ or saline. All animals (MPP⁺: n = 63; saline: n = 21) were assessed daily on a scale from 1–3 for their clinical appearance, including weight loss, inactivity, and reaction patterns, giving a total score ranging from 0–12 (see Results). Following unilateral striatal treatment with MPP⁺, both *Aqp9*^{-/-} and WT mice showed a significantly higher score than control animals injected with saline. *Aqp9*^{-/-} mice treated with MPP⁺ showed a lower score than the WT littermates throughout the observation period. Statistical analysis of data from individual observations revealed significant difference on day 2. B) Following a systemic injection apomorphine, *Aqp9*^{-/-} and WT littermates treated with a unilateral striatal injection of 7.5 µg MPP⁺ tend to spin towards the injected side. *Aqp9*^{-/-} (n = 26) show significantly less net rotations (ipsilateral—contralateral turns) compared to WT mice (n = 27), with a net rotation number of 2.69 ± 1.91 compared to 6.33 ± 2.15 (mean ± 2 SEM; p = 0.014, n = 53). The rotational behavior was not present in animals treated with saline (n = 17); Boxes are mean ± 95% CI, bars are maximum and minimum values, *p < 0.05.

<https://doi.org/10.1371/journal.pone.0194896.g006>

Fig 7. HPLC analysis of dopamine and its metabolites. A–C) After unilateral injections of MPP⁺ in the striatum, the ipsilateral striatum shows a significant reduction in the concentration of DA (A), HVA (B) and DOPAC (C) compared to the contralateral hemisphere (*Aqp9*^{-/-}, n = 6; WT, n = 6). The ipsilateral reduction in HVA is significantly more pronounced in WT mice than in *Aqp9*^{-/-} mice (p = 0.038). Corresponding p-values for DA and DOPAC are 0.063 and 0.059, respectively. Bars are mean ± 2 SEM; *p < 0.05, ** p < 0.01, ***p < 0.001.

<https://doi.org/10.1371/journal.pone.0194896.g007>

Fig 8. Semiquantitative PCR analyses of gene expression in *Aqp9*^{-/-} and WT mice brain. A) Semi-quantitative Real-Time PCR revealed significantly higher *Aqp9* mRNA levels in the midbrain and striatum than in the neocortex. The level of *Aqp9* mRNA in neocortex is 59% of that in midbrain ($p = 0.002$). The level of *Aqp9* mRNA in the *Aqp9*^{-/-} mice was close to the detection limit ($n = 7$ for each genotype). B) Representative DNA agarose gel electrophoresis of midbrain samples from WT and *Aqp9*^{-/-} mice (upper panel), and of three different regions in WT mice (lower panel). These data support the PCR analysis in A. C) In order to rule out that the reduced dopaminergic cell loss in the *Aqp9*^{-/-} mice could be attributed to compensatory up- or downregulation of other genes, an analysis was done of the expression levels of *Aqp4*, *Gfap*, *Kir4.1*, *mTOR*, *Prph*, *Cat*, *Ppard*, *Slc6a3* (DAT), *Drd2*, *Bcl2*, *Bax* and *Sod2*. The relative levels of these transcripts did not differ between *Aqp9*^{-/-} ($n = 7$) and WT animals ($n = 7$). D-H) For selected genes the expression levels were analyzed in the treated and untreated hemispheres. In both groups of animal, the transcript levels of *Drd2* were lower in the striatum on the injected side than in the striatum on the contralateral side (E). In contrast, in both groups of animals, the level of *Bax* was higher in the ipsilateral striatum than in the contralateral one (H). The values indicated in the graphs for DAT, *Bcl2* and *Sod2* show the values for midbrain, and the values for *Drd2* are for striatum. All values are relative to the values for the corresponding samples from the control hemisphere of the saline treated mice. Bars are mean $\pm$ 2 SEM; ** $p < 0.005$.

<https://doi.org/10.1371/journal.pone.0194896.g008>

animals. In both animals groups, transcript levels of DAT and *Drd2* were lower on the injected side than on the contralateral side (Fig 8D–8E). In contrast, in both groups of animals, the level of *Bax* was higher in the ipsilateral striatum than in the striatum contralateral to the injection (Fig 8H).

Discussion

There is an urgent need to identify the mechanisms underlying the selective vulnerability of dopaminergic cells, the root cause of Parkinson's disease. In the majority of cases, Parkinson's disease has no clear genetic etiology, thus emphasizing the importance of unravelling the contribution of environmental factors. Here we show that the parkinsonogenic toxin MPP⁺ permeates AQP9, an aquaglyceroporin water channel that is selectively expressed in dopaminergic neurons, and that a targeted deletion of this water channel affords protection *in vitro* and *in vivo*. We also show that stable expression of AQP9 in HEK cells exacerbates their vulnerability to MPP⁺ and arsenite—another toxin associated with the development of PD. Our data open for the possibility that toxins and other parkinsonogenic substances access dopaminergic neurons through AQP9 and imply that this aquaglyceroporin should be explored as a new target for pharmacological intervention.

The dopamine transporter DAT has long been known to provide an entry route for environmental and experimental toxins [22]. For obvious reasons, this transporter is not relevant as a therapeutic target. AQP9, on the other hand, is not critical to brain function [37] and can be targeted by extant drugs. AQP9 is the first membrane channel that has been associated with the entry of parkinsonogenic toxins.

Our experiments in frog oocytes established that AQP9 is permeable to MPP⁺. To compare the relative significance of AQP9 and DAT we had to resort to cells that normally do not express these molecules, and that by inference are non-dopaminergic. We chose to use HEK cells that have been applied extensively for this purpose in previous studies. We found that AQP9 exacerbates MPP⁺ toxicity, as does DAT. By contrast, the toxicity of arsenite—another parkinsonogenic compound—was potentiated by AQP9 only. The expression of DAT failed to potentiate arsenite toxicity, except at arsenite concentrations exceeding those normally used.

The most salient observation in the present study is that deletion of *Aqp9* affords protection against MPP⁺. We demonstrate a protective effect in organotypic midbrain cultures, as well as after MPP⁺ injections *in vivo*. Stereological quantification of the density of TH-positive neurons in the SNpc after MPP⁺ injection *in vivo* showed a loss of TH-positive neurons of 67% in WT animals, compared with 48% in *Aqp9*^{-/-} animals. The reduced cell loss was accompanied by less pronounced motor dysfunction. The level of the dopamine metabolite HVA in the ipsilateral striatum was about 40% higher in *Aqp9*^{-/-} animals than in WT, supporting the idea that deletion of *Aqp9* helps protect the nigrostriatal pathway against the deleterious effect of MPP⁺.

In organotypic cultures we showed that the protective effect of *Aqp9* gene deletion was mimicked by the application of phloretin—an effective blocker of AQP9. This finding indicates that the reduced vulnerability to MPP⁺ in *Aqp9*^{-/-} vs. WT mice is due to the deletion of the *Aqp9* gene and not a result of any off-target effect. In agreement, targeted deletion of *Aqp9* did not alter the expression level of select genes associated with MPP⁺-induced cell death.

Taken together with our data showing increased permeability to MPP⁺ in AQP9-expressing oocytes, these results suggest that AQP9 is involved in mediating the toxicity of MPP⁺ by enabling this toxin to enter dopaminergic neurons. Importantly, as AQP9 is found in the inner mitochondrial membrane as well as in the plasma membrane [23], this channel may allow MPP⁺ and other toxins to permeate the mitochondrial matrix. This is relevant, as the electron transport chain has been identified as one of several targets of parkinsonogenic toxins [47,48].

Methodological considerations

Here we used an *in vivo* model of Parkinson's disease where MPP⁺ was injected unilaterally into the striatum of *Aqp9* deficient and WT mice. The injections covered a large part of the projection field of nigral dopaminergic neurons and thus allowed retrograde transport of toxin to these neurons. In designing these experiments we aimed for a cell loss of more than 60% in the SNpc of WT mice, comparable to the cell loss seen in postmortem analyses of patients with Parkinson's disease [49]. The dose of MPP⁺ produced a rather extensive and unspecific damage around the injection site. In midbrain, however, the deleterious effect of MPP⁺ was largely restricted to the SN. Specifically, neither genotype showed a substantial loss of dopaminergic cells in the VTA. The most likely explanation is that the MPP⁺ injection was discrete enough to encompass the projection area of nigral neurons without encroaching upon the projections of those dopaminergic neurons that reside in the VTA.

Compared with WT, *Aqp9*^{-/-} mice developed less motor dysfunction when exposed to MPP⁺. In the apomorphine rotation test, *Aqp9*^{-/-} mice showed significantly less ipsilateral spinning than WT littermates. Apomorphine is thought to induce contralateral spinning at a high rate due to supersensitivity of dopaminergic D₂ receptors in the striatal nerve terminals [41].

However, several studies have shown that severe damage to the striatum—similar to that seen here—induces ipsilateral turning behavior following administration of apomorphine [50–54]. In agreement, we observed a pronounced decrease in the expression level of dopamine D₂ receptors in the striatum of both genotypes, reflecting the extensive damage and explaining the ipsilateral rotation.

In line with the outcome of the apomorphine rotation test, the clinical appearance of MPP⁺ injected animals, as well as the pronounced drop in dopamine and dopamine metabolites in the ipsilateral striatum, point to a rather severe local effect of MPP⁺ injections. Histological analyses concurred by showing areas of cell loss and immune cell infiltration involving large parts of the striatum and overlying cortex. The lesion did not extend into the ventral striatum. The nonspecific nature of the local lesion was distinct from the lesion in midbrain, which likely is mediated by retrograde transport of the injected toxin. Thus, the midbrain lesion specifically involved TH-positive neurons in the SNpc, leaving nearby VTA neurons (that normally project to the ventral striatum) virtually unharmed.

AQP9 is expressed in dopaminergic midbrain neurons [23,24,26,55] but is also found in astroglia throughout the brain [23,25,55,56]. This explains why we found *Aqp9* mRNA in other brain regions, including the neocortex. However, our regional analysis showed higher levels of *Aqp9* mRNA in midbrain and striatum than in neocortex. This is consistent with previous immunocytochemical data and with the finding that these structures contain a neuronal as well as an astroglial pool of AQP9 [26].

Acute injections of MPP⁺ remain a rather crude model for Parkinson's disease. This calls for due care when it comes to the interpretation of the present results. In a clinical setting Parkinson's disease develops over years, even decades, implying that the present study should be followed up in more chronic models. However, even with long time exposure to minute amounts of a putative neurotoxin the vulnerability may be determined by the same factors as in the acute model *i.e.*, the transport capacity for the toxin or toxins in question. Thus, the expression level of AQP9 in plasma and mitochondrial membranes might not only explain the selective vulnerability of dopaminergic neurons within an individual, but possibly also the differences in vulnerability between individuals. This issue will be pursued in future studies.

AQP9 vs. DAT

In earlier works the selective sensitivity of SN neurons to parkinsonogenic toxins has been attributed to the activity of the dopamine transporter DAT [22]. It is well documented that DAT transports MPP⁺ [57]. This begs the question whether DAT is the predominant mediator of MPP⁺ toxicity, leaving only a minor role for AQP9. We addressed this issue by use of the MTT assay, which is widely employed to assess the effects of toxins, including MPP⁺.

We performed the MTT assay in HEK293 cells as these cells do not normally express AQP9 or DAT. HEK293 cells stably transfected with *hAQP9* were more sensitive to MPP⁺ than were cells stably transfected with *hDAT*. Notably, while 1 μM MPP⁺ was required to produce significant effects in *hDAT* expressing cells, 0.1 μM MPP⁺ was sufficient to induce toxicity in cells expressing *hAQP9*. HEK293 cells expressing *hAQP9* were also clearly more sensitive to the parkinsonogenic compound arsenite than were HEK293 cells expressing *hDAT*. At concentrations up to 10 μM, arsenite had no toxic effects in *hDAT* expressing HEK293 cells. This is in line with previous studies that did not detect any difference in arsenite sensitivity between *hDAT* expressing HEK293 cells and native HEK293 cells [58]. Only at very high arsenite concentrations do *hDAT* expressing cells differ from native HEK293 cells in terms of their viability.

In our *in vivo* model, MPP⁺ caused a significant dopaminergic cell loss also in the *Aqp9*^{-/-} animals. This can probably be attributed to the expression of DAT in the nigrostriatal terminals, facilitating uptake of MPP⁺ also into dopaminergic neurons that are depleted of AQP9 [22]. In midbrain organotypical cultures, the nigrostriatal axons are severed, implying that MPP⁺ uptake through AQP9, expressed at high levels in dopaminergic cell bodies [23,26], may predominate over MPP⁺ uptake by DAT. This may explain why the protective effect of *Aqp9* deletion was more pronounced in the slice cultures than in the *in vivo* model.

We conclude that both DAT (a transporter) and AQP9 (a channel) are likely to be involved in mediating MPP⁺ toxicity. However, while DAT transports MPP⁺, it has a rather narrow substrate specificity that does not encompass the wide range of toxins that have been associated with the development of Parkinson's disease. Unlike DAT, which is essential for normal brain function, AQP9 is not critically involved in any known physiological process in brain. Thus, *Aqp9*^{-/-} mice have a rather mild phenotype largely limited to the peripheral organs such as the liver [37]. Little is known about the roles of AQP9 in human brain as most studies of AQP9 in humans have focused on the reproductive system and placenta [59–61]. If applicable to humans, the present data suggest that pharmacologic inhibition of AQP9 could be considered a viable approach to curb the progression of Parkinson's disease in cases where exposure to exogenous toxins is documented or suspected as a cause of morbidity.

Importantly, targeted deletion of *Aqp9* did not affect the expression level of DAT. Our finding that phloretin treatment mimicked the effect of *Aqp9* deletion lends further support to the idea that the protection observed is a bona fide effect of the targeted deletion and not due to compensatory up- or downregulation of other genes.

Conclusions and prospects for new therapy

Earlier studies have shown that AQP9 is expressed in midbrain neurons that contain TH—a marker of dopaminergic neurons [23,24,26,55]. Here, using both *in vitro* and *in vivo* models, we provide evidence that AQP9 contributes to the selective vulnerability of dopaminergic neurons to exogenous toxins. This is the first study to couple this vulnerability to a specific membrane channel allowing toxin influx. The broad substrate specificity of AQP9 implies that this channel might permit entry of a number of compounds that compromise the viability of dopaminergic neurons. It is of interest in this regard that the *Aqp9* gene contains a negative insulin response element, and that insulin downregulates its expression level [62–64]. Studies on experimental diabetes in rodents as well as tissue samples from patients with diabetes have shown an increase in the level of AQP9 mRNA and protein [65,66]. The possibility should be considered that an upregulation of *Aqp9* contributes to the increased prevalence of Parkinson's disease among patients with diabetes mellitus type 2 [67].

As pointed out in our discussion, the rather short survival times that are used in such models do not match the protracted course of disease development in the human condition. This is a general shortcoming of most current animal models of PD and calls for due care when it comes to the interpretation of the results. Inspired by the comments of the reviewers we deepen our discussion of this point in our revised text. As we state in our discussion a natural follow up of our study is to subject our *Aqp9*^{-/-} animals to other and more chronic experimental approaches that more faithfully mimic the progressive and protracted nature of human disease.

Supporting information

S1 Fig. Tissue damage and immune cell infiltration following ipsilateral intrastriatal injection of MPP⁺. A-D) Semi-thin sections immunostained for TH (A) or hematoxylin/eosin (B-D) showing local effect of the MPP⁺ injections. The site of MPP⁺ injection (Ipsi) in the

anterior striatum is indicated by arrow (A). The toxin induces tissue damage in the entire ipsilateral striatum (CPu). Cell loss and immune cell infiltration extended into parts of the overlying neocortex (B-D). Reflecting the tissue damage, the TH immunosignal was blurred, with loss of the sharp contrast between the corpus callosum (cc), overlying cortex (Cx) and striatum (CPu) seen on the contralateral (Cont) side. Stippled line outlines the damaged area. Boxed areas in B are enlarged in C and D. cc, Corpus Callosum; CPu, Caudate-Putamen; Cx, frontal cortex; LV, Lateral Ventricle. Scale bar: 1000 μ m.
(TIF)

S1 File. Methods.

(DOCX)

S2 File. Humane endpoints checklist.

(DOCX)

S1 Table. List of the TaqMan probes used for quantitative RT-PCR.

(DOCX)

S2 Table. Overview of the estimated total cell count of TH-positive neurons.

(DOCX)

Acknowledgments

We thank Ngoc-Trang Thi Huynh for HPLC analysis, Hong Qu for technical support on light microscopic imaging, Carina Knudsen for graphic support and Bjørge Riber, Karen Marie Gujord, Grazyna Babinska, Mona Gaarder and Vesa Loitto for technical assistance. *Aqp9*^{-/-} mice were a kind gift from Søren Nilsen and Aleksandra Rojek, University of Aarhus. HEK-DAT cells were a kind gift from Dr. Habibeh Khoshbouei, Department of Neuroscience, University of Florida, USA, and HEK-*Aqp9* cells were a kind gift from Professor Karl-Eric Magnusson, Linköping University, Sweden.

Author Contributions

Conceptualization: Ole P. Ottersen, Mahmood Amiry-Moghaddam.

Data curation: Katja Stahl, Soulmaz Rahmani, Agnete Prydz, Nadia Skauli, Nanna MacAulay, Maria N. Mylonakou, Øivind Skare, Torill Berg, Ragnhild E. Paulsen.

Formal analysis: Katja Stahl, Soulmaz Rahmani, Nadia Skauli, Nanna MacAulay, Maria N. Mylonakou, Øivind Skare, Torill Berg, Trygve B. Leergaard, Ragnhild E. Paulsen, Mahmood Amiry-Moghaddam.

Funding acquisition: Mahmood Amiry-Moghaddam.

Investigation: Katja Stahl, Soulmaz Rahmani, Agnete Prydz, Nadia Skauli, Nanna MacAulay, Maria N. Mylonakou, Torill Berg, Ragnhild E. Paulsen, Mahmood Amiry-Moghaddam.

Methodology: Katja Stahl, Soulmaz Rahmani, Nadia Skauli, Nanna MacAulay, Øivind Skare, Torill Berg, Trygve B. Leergaard, Ragnhild E. Paulsen, Mahmood Amiry-Moghaddam.

Project administration: Katja Stahl, Mahmood Amiry-Moghaddam.

Supervision: Reidun Torp, Trygve B. Leergaard, Ole P. Ottersen, Mahmood Amiry-Moghaddam.

Validation: Ole P. Ottersen, Mahmood Amiry-Moghaddam.

Visualization: Katja Stahl, Soulmaz Rahmani, Nanna MacAulay, Maria N. Mylonakou, Øivind Skare, Torill Berg, Trygve B. Leergaard, Ragnhild E. Paulsen.

Writing – original draft: Katja Stahl, Ole P. Ottersen, Mahmood Amiry-Moghaddam.

Writing – review & editing: Katja Stahl, Soulmaz Rahmani, Agnete Prydz, Nadia Skauli, Nanna MacAulay, Maria N. Mylonakou, Reidun Torp, Øivind Skare, Torill Berg, Trygve B. Leergaard, Ragnhild E. Paulsen, Ole P. Ottersen, Mahmood Amiry-Moghaddam.

References

1. de Rijk MC, Breteler MM, Graveland GA, Ott A, Grobbee DE, van der Meche FG, et al. (1995) Prevalence of Parkinson's disease in the elderly: the Rotterdam Study. *Neurology* 45: 2143–2146. PMID: [8848182](#)
2. de Rijk MC, Launer LJ, Berger K, Breteler MM, Dartigues JF, Baldereschi M, et al. (2000) Prevalence of Parkinson's disease in Europe: A collaborative study of population-based cohorts. *Neurologic Diseases in the Elderly Research Group. Neurology* 54: S21–23.
3. Hirsch L, Jette N, Frolkis A, Steeves T, Pringsheim T (2016) The Incidence of Parkinson's Disease: A Systematic Review and Meta-Analysis. *Neuroepidemiology* 46: 292–300. <https://doi.org/10.1159/000445751> PMID: [27105081](#)
4. Pringsheim T, Jette N, Frolkis A, Steeves TD (2014) The prevalence of Parkinson's disease: a systematic review and meta-analysis. *Mov Disord* 29: 1583–1590. <https://doi.org/10.1002/mds.25945> PMID: [24976103](#)
5. Gibb WR, Lees AJ (1991) Anatomy, pigmentation, ventral and dorsal subpopulations of the substantia nigra, and differential cell death in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 54: 388–396. PMID: [1865199](#)
6. Fahn S (2003) Description of Parkinson's disease as a clinical syndrome. *Ann N Y Acad Sci* 991: 1–14.
7. Payami H, Zarepari S (1998) Genetic epidemiology of Parkinson's disease. *J Geriatr Psychiatry Neurol* 11: 98–106. <https://doi.org/10.1177/089198879801100207> PMID: [9877530](#)
8. Davis GC, Williams AC, Markey SP, Ebert MH, Caine ED, Reichert CM, et al. (1979) Chronic Parkinsonism secondary to intravenous injection of meperidine analogues. *Psychiatry Res* 1: 249–254. PMID: [298352](#)
9. Langston JW, Ballard P, Tetrud JW, Irwin I (1983) Chronic Parkinsonism in humans due to a product of meperidine-analog synthesis. *Science* 219: 979–980. PMID: [6823561](#)
10. Burns RS, Chiueh CC, Markey SP, Ebert MH, Jacobowitz DM, Kopin IJ (1983) A primate model of parkinsonism: selective destruction of dopaminergic neurons in the pars compacta of the substantia nigra by N-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *Proc Natl Acad Sci U S A* 80: 4546–4550. PMID: [6192438](#)
11. Forno LS, DeLanney LE, Irwin I, Langston JW (1993) Similarities and differences between MPTP-induced parkinsonism and Parkinson's disease. *Neuropathologic considerations. Adv Neurol* 60: 600–608. PMID: [8380528](#)
12. Heikkila RE, Sieber BA, Manzino L, Sonsalla PK (1989) Some features of the nigrostriatal dopaminergic neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) in the mouse. *Mol Chem Neuropathol* 10: 171–183. PMID: [2669769](#)
13. Gorell JM, Johnson CC, Rybicki BA, Peterson EL, Kortsha GX, Brown GG, et al. (1999) Occupational exposure to manganese, copper, lead, iron, mercury and zinc and the risk of Parkinson's disease. *Neurotoxicology* 20: 239–247. PMID: [10385887](#)
14. Koller W, Vetere-Overfield B, Gray C, Alexander C, Chin T, Dolezal J, et al. (1990) Environmental risk factors in Parkinson's disease. *Neurology* 40: 1218–1221. PMID: [2381528](#)
15. Ascherio A, Chen H, Weisskopf MG, O'Reilly E, McCullough ML, Calle EE, et al. (2006) Pesticide exposure and risk for Parkinson's disease. *Ann Neurol* 60: 197–203. <https://doi.org/10.1002/ana.20904> PMID: [16802290](#)
16. Frigerio R, Sanft KR, Grossardt BR, Peterson BJ, Elbaz A, Bower JH, et al. (2006) Chemical exposures and Parkinson's disease: a population-based case-control study. *Mov Disord* 21: 1688–1692. <https://doi.org/10.1002/mds.21009> PMID: [16773614](#)
17. Fan SF, Chao PL, Lin AM (2010) Arsenite induces oxidative injury in rat brain: synergistic effect of iron. *Ann N Y Acad Sci* 1199: 27–35. <https://doi.org/10.1111/j.1749-6632.2009.05170.x> PMID: [20633106](#)
18. Priyadarshi A, Khuder SA, Schaub EA, Shrivastava S (2000) A meta-analysis of Parkinson's disease and exposure to pesticides. *Neurotoxicology* 21: 435–440. PMID: [11022853](#)

19. van der Mark M, Brouwer M, Kromhout H, Nijssen P, Huss A, Vermeulen R (2012) Is pesticide use related to Parkinson disease? Some clues to heterogeneity in study results. *Environ Health Perspect* 120: 340–347. <https://doi.org/10.1289/ehp.1103881> PMID: [22389202](#)
20. Pezzoli G, Cereda E (2013) Exposure to pesticides or solvents and risk of Parkinson disease. *Neurology* 80: 2035–2041. <https://doi.org/10.1212/WNL.0b013e318294b3c8> PMID: [23713084](#)
21. Heikkilä RE, Nicklas WJ, Vyas I, Duvoisin RC (1985) Dopaminergic toxicity of rotenone and the 1-methyl-4-phenylpyridinium ion after their stereotaxic administration to rats: implication for the mechanism of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine toxicity. *Neurosci Lett* 62: 389–394. PMID: [3912685](#)
22. Javitch JA, D'Amato RJ, Strittmatter SM, Snyder SH (1985) Parkinsonism-inducing neurotoxin, N-methyl-4-phenyl-1,2,3,6-tetrahydropyridine: uptake of the metabolite N-methyl-4-phenylpyridine by dopamine neurons explains selective toxicity. *Proc Natl Acad Sci U S A* 82: 2173–2177. PMID: [3872460](#)
23. Amiry-Moghaddam M, Lindland H, Zelenin S, Røberg BA, Gundersen BB, Petersen P, et al. (2005) Brain mitochondria contain aquaporin water channels: evidence for the expression of a short AQP9 isoform in the inner mitochondrial membrane. *FASEB J* 19: 1459–1467. <https://doi.org/10.1096/fj.04-3515com> PMID: [16126913](#)
24. Arcienega II, Brunet JF, Bloch J, Badaut J (2010) Cell locations for AQP1, AQP4 and 9 in the non-human primate brain. *Neuroscience* 167: 1103–1114. <https://doi.org/10.1016/j.neuroscience.2010.02.059> PMID: [20226845](#)
25. Badaut J, Hirt L, Granziera C, Bogousslavsky J, Magistretti PJ, Regli L (2001) Astrocyte-specific expression of aquaporin-9 in mouse brain is increased after transient focal cerebral ischemia. *J Cereb Blood Flow Metab* 21: 477–482. <https://doi.org/10.1097/00004647-200105000-00001> PMID: [11333357](#)
26. Mylonakou MN, Petersen PH, Rinvik E, Rojek A, Valdimarsdóttir E, Zelenin S, et al. (2009) Analysis of mice with targeted deletion of AQP9 gene provides conclusive evidence for expression of AQP9 in neurons. *J Neurosci Res* 87: 1310–1322. <https://doi.org/10.1002/jnr.21952> PMID: [19115411](#)
27. Ishibashi K, Kuwahara M, Gu Y, Tanaka Y, Marumo F, Sasaki S (1998) Cloning and functional expression of a new aquaporin (AQP9) abundantly expressed in the peripheral leukocytes permeable to water and urea, but not to glycerol. *Biochem Biophys Res Commun* 244: 268–274. <https://doi.org/10.1006/bbrc.1998.8252> PMID: [9514918](#)
28. Tsukaguchi H, Shayakul C, Berger UV, Mackenzie B, Devidas S, Guggino WB, et al. (1998) Molecular characterization of a broad selectivity neutral solute channel. *J Biol Chem* 273: 24737–24743. PMID: [9733774](#)
29. Tsukaguchi H, Weremowicz S, Morton CC, Hediger MA (1999) Functional and molecular characterization of the human neutral solute channel aquaporin-9. *Am J Physiol* 277: F685–696. PMID: [10564231](#)
30. Holm LM, Jahn TP, Møller AL, Schjøerring JK, Ferri D, Klaerke DA, et al. (2005) NH₃ and NH₄⁺ permeability in aquaporin-expressing *Xenopus* oocytes. *Pflügers Arch* 450: 415–428. <https://doi.org/10.1007/s00424-005-1399-1> PMID: [15988592](#)
31. Liu Z, Shen J, Carbrey JM, Mukhopadhyay R, Agre P, Rosen BP (2002) Arsenite transport by mammalian aquaglyceroporins AQP7 and AQP9. *Proc Natl Acad Sci U S A* 99: 6053–6058. <https://doi.org/10.1073/pnas.092131899> PMID: [11972053](#)
32. Moeller HB, Fenton RA, Zeuthen T, Macaulay N (2009) Vasopressin-dependent short-term regulation of aquaporin 4 expressed in *Xenopus* oocytes. *Neuroscience* 164: 1674–1684. <https://doi.org/10.1016/j.neuroscience.2009.09.072> PMID: [19800950](#)
33. Jespersen T, Grønnet M, Angelo K, Klaerke DA, Olesen SP (2002) Dual-function vector for protein expression in both mammalian cells and *Xenopus laevis* oocytes. *Biotechniques* 32: 536–538, 540. PMID: [11911656](#)
34. Richardson BD, Saha K, Krout D, Cabrera E, Felts B, Henry LK, et al. (2016) Membrane potential shapes regulation of dopamine transporter trafficking at the plasma membrane. *Nat Commun* 7: 10423. <https://doi.org/10.1038/ncomms10423> PMID: [26804245](#)
35. Karlsson T, Bolshakova A, Magalhaes MA, Loitto VM, Magnusson KE (2013) Fluxes of water through aquaporin 9 weaken membrane-cytoskeleton anchorage and promote formation of membrane protrusions. *PLoS One* 8: e59901. <https://doi.org/10.1371/journal.pone.0059901> PMID: [23573219](#)
36. Loitto VM, Huang C, Sigal YJ, Jacobson K (2007) Filopodia are induced by aquaporin-9 expression. *Exp Cell Res* 313: 1295–1306. <https://doi.org/10.1016/j.yexcr.2007.01.023> PMID: [17346701](#)
37. Rojek AM, Skowronski MT, Fuchtbauer EM, Fuchtbauer AC, Fenton RA, Agre P, et al. (2007) Defective glycerol metabolism in aquaporin 9 (AQP9) knockout mice. *Proc Natl Acad Sci U S A* 104: 3609–3614. <https://doi.org/10.1073/pnas.0610894104> PMID: [17360690](#)

38. Stahl K, Skare O, Torp R (2009) Organotypic cultures as a model of Parkinson's disease. A twist to an old model. *ScientificWorldJournal* 9: 811–821. <https://doi.org/10.1100/tsw.2009.68> PMID: 19705040
39. Stahl K, Mylonakou MN, Skare O, Amiry-Moghaddam M, Torp R (2011) Cytoprotective effects of growth factors: BDNF more potent than GDNF in an organotypic culture model of Parkinson's disease. *Brain Res* 1378: 105–118. <https://doi.org/10.1016/j.brainres.2010.12.090> PMID: 21236244
40. Paxinos GaF K (2007) *The Mouse Brain in Stereotaxic Coordinates*: Elsevier Inc.
41. Ungerstedt U, Arbuthnott GW (1970) Quantitative recording of rotational behavior in rats after 6-hydroxy-dopamine lesions of the nigrostriatal dopamine system. *Brain Res* 24: 485–493. PMID: 5494536
42. Gundersen HJ, Bagger P, Bendtsen TF, Evans SM, Korbo L, Marcussen N, et al. (1988) The new stereological tools: disector, fractionator, nucleator and point sampled intercepts and their use in pathological research and diagnosis. *APMIS* 96: 857–881. PMID: 3056461
43. Gundersen HJ, Jensen EB, Kieu K, Nielsen J (1999) The efficiency of systematic sampling in stereology—reconsidered. *J Microsc* 193: 199–211. PMID: 10348656
44. Katoozi S, Skauli N, Rahmani S, Camassa LMA, Boldt HB, Ottersen OP, et al. (2017) Targeted deletion of Aqp4 promotes the formation of astrocytic gap junctions. *Brain Struct Funct* 222: 3959–3972. <https://doi.org/10.1007/s00429-017-1448-5> PMID: 28551776
45. Fenton RA, Moeller HB, Zelenina M, Snaebjornsson MT, Holen T, MacAulay N (2010) Differential water permeability and regulation of three aquaporin 4 isoforms. *Cell Mol Life Sci* 67: 829–840. <https://doi.org/10.1007/s00018-009-0218-9> PMID: 20013023
46. Smeyne RJ, Breckenridge CB, Beck M, Jiao Y, Butt MT, Wolf JC, et al. (2016) Assessment of the Effects of MPTP and Paraquat on Dopaminergic Neurons and Microglia in the Substantia Nigra Pars Compacta of C57BL/6 Mice. *PLoS One* 11: e0164094. <https://doi.org/10.1371/journal.pone.0164094> PMID: 27788145
47. Betarbet R, Sherer TB, MacKenzie G, Garcia-Osuna M, Panov AV, Greenamyre JT (2000) Chronic systemic pesticide exposure reproduces features of Parkinson's disease. *Nat Neurosci* 3: 1301–1306. <https://doi.org/10.1038/81834> PMID: 11100151
48. Nicklas WJ, Youngster SK, Kindt MV, Heikkila RE (1987) MPTP, MPP+ and mitochondrial function. *Life Sci* 40: 721–729. PMID: 3100899
49. German DC, Manaye K, Smith WK, Woodward DJ, Saper CB (1989) Midbrain dopaminergic cell loss in Parkinson's disease: computer visualization. *Ann Neurol* 26: 507–514. <https://doi.org/10.1002/ana.410260403> PMID: 2817827
50. Ghorayeb I, Fernagut PO, Hervier L, Labattu B, Bioulac B, Tison F (2002) A 'single toxin-double lesion' rat model of striatonigral degeneration by intrastratial 1-methyl-4-phenylpyridinium ion injection: a motor behavioural analysis. *Neuroscience* 115: 533–546. PMID: 12421620
51. Dunnett SB, Isacson O, Sirinathsinghji DJ, Clarke DJ, Bjorklund A (1988) Striatal grafts in rats with unilateral neostriatal lesions—III. Recovery from dopamine-dependent motor asymmetry and deficits in skilled paw reaching. *Neuroscience* 24: 813–820. PMID: 3380301
52. Kafetzopoulos E, Vlaha V, Konitsiotis S (1988) Different patterns of rotational behavior in rats after dorsal or ventral striatal lesions with ibotenic acid. *Pharmacol Biochem Behav* 29: 403–406. PMID: 3362934
53. Nakao N, Brundin P (1997) Effects of alpha-phenyl-tert-butyl nitron on neuronal survival and motor function following intrastratial injections of quinolinic acid or 3-nitropropionic acid. *Neuroscience* 76: 749–761. PMID: 9135048
54. Ghorayeb I, Puschban Z, Fernagut PO, Scherfler C, Rouland R, Wenning GK, et al. (2001) Simultaneous intrastratial 6-hydroxydopamine and quinolinic acid injection: a model of early-stage striatonigral degeneration. *Exp Neurol* 167: 133–147. <https://doi.org/10.1006/exnr.2000.7535> PMID: 11161601
55. Badaut J, Petit JM, Brunet JF, Magistretti PJ, Charriaut-Marlangue C, Regli L (2004) Distribution of Aquaporin 9 in the adult rat brain: preferential expression in catecholaminergic neurons and in glial cells. *Neuroscience* 128: 27–38. <https://doi.org/10.1016/j.neuroscience.2004.05.042> PMID: 15450351
56. Hirt L, Price M, Mastour N, Brunet JF, Barriere G, Friscourt F, et al. (2018) Increase of aquaporin 9 expression in astrocytes participates in astrogliosis. *J Neurosci Res* 96: 194–206. <https://doi.org/10.1002/jnr.24061> PMID: 28419510
57. Kitayama S, Mitsuhata C, Davis S, Wang JB, Sato T, Morita K, et al. (1998) MPP+ toxicity and plasma membrane dopamine transporter: study using cell lines expressing the wild-type and mutant rat dopamine transporters. *Biochim Biophys Acta* 1404: 305–313. PMID: 9739158
58. Shavali S, Sens DA (2008) Synergistic neurotoxic effects of arsenic and dopamine in human dopaminergic neuroblastoma SH-SY5Y cells. *Toxicol Sci* 102: 254–261. <https://doi.org/10.1093/toxsci/kfm302> PMID: 18079140

59. Vilarino-Garcia T, Perez-Perez A, Dietrich V, Fernandez-Sanchez M, Guadix P, Duenas JL, et al. (2016) Increased Expression of Aquaporin 9 in Trophoblast From Gestational Diabetic Patients. *Horm Metab Res* 48: 535–539. <https://doi.org/10.1055/s-0042-105152> PMID: 27082037
60. Winterbottom EF, Koestler DC, Fei DL, Wika E, Capobianco AJ, Marsit CJ, et al. (2017) The aquaglyceroporin AQP9 contributes to the sex-specific effects of in utero arsenic exposure on placental gene expression. *Environ Health* 16: 59. <https://doi.org/10.1186/s12940-017-0267-8> PMID: 28615018
61. Vilarino-Garcia T, Perez-Perez A, Dietrich V, Guadix P, Duenas JL, Varone CL, et al. (2018) Leptin upregulates aquaporin 9 expression in human placenta in vitro. *Gynecol Endocrinol* 34: 175–177. <https://doi.org/10.1080/09513590.2017.1380184> PMID: 28942694
62. Badaut J, Regli L (2004) Distribution and possible roles of aquaporin 9 in the brain. *Neuroscience* 129: 971–981. <https://doi.org/10.1016/j.neuroscience.2004.06.035> PMID: 15561412
63. Carbre JM, Gorelick-Feldman DA, Kozono D, Praetorius J, Nielsen S, Agre P (2003) Aquaglyceroporin AQP9: solute permeation and metabolic control of expression in liver. *Proc Natl Acad Sci U S A* 100: 2945–2950. <https://doi.org/10.1073/pnas.0437994100> PMID: 12594337
64. Kuriyama H, Shimomura I, Kishida K, Kondo H, Furuyama N, Nishizawa H, et al. (2002) Coordinated regulation of fat-specific and liver-specific glycerol channels, aquaporin adipose and aquaporin 9. *Diabetes* 51: 2915–2921. PMID: 12351427
65. Castro Parodi M, Farina M, Dietrich V, Aban C, Szpilbarg N, Zotta E, et al. (2011) Evidence for insulin-mediated control of AQP9 expression in human placenta. *Placenta* 32: 1050–1056. <https://doi.org/10.1016/j.placenta.2011.09.022> PMID: 22018417
66. Fan Z, Yuan Y, Wang F, Qi Y, Han H, Wu J, et al. (2017) Diabetes mitigates the recovery following intracranial hemorrhage in rats. *Behav Brain Res* 320: 412–419. <https://doi.org/10.1016/j.bbr.2016.10.047> PMID: 27818237
67. Driver JA, Smith A, Buring JE, Gaziano JM, Kurth T, Logroscino G (2008) Prospective cohort study of type 2 diabetes and the risk of Parkinson's disease. *Diabetes Care* 31: 2003–2005. <https://doi.org/10.2337/dc08-0688> PMID: 18599528