A Novel Approach to Inner Cancer Treatment through the Activation of Photosensitizers by Protons

Maria Mastrangelopoulou

Department of Radiation Biology
Institute for Cancer Research
The Norwegian Radium Hospital
Oslo University Hospital

Dissertation submitted for the degree of Ph.D.

Faculty of Medicine
Institute for Clinical Medicine
University of Oslo

© Maria Mastrangelopoulou, 2021

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-936-3

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

Acknowledgements

The work included in this thesis was carried out at the Department of Radiation Biology, Institute for Cancer Research at the Norwegian Radium Hospital in the laboratory of Professor Kristian Berg between the year 2017 and 2021. The financial support received from the South-Eastern Norway Regional Health Authority (Helse Sør-Øst) is gratefully acknowledged.

First and foremost I would like to thank my supervisor and Project leader in the Protonic group, Dr Theodossis Theodossiou, for giving me the opportunity to work with such an exciting and cutting-edge scientific project! I am so grateful for all your guidance and assistance throughout all these years. Your enthusiasm and creativity has been an enormous inspiration for the progress of this project, which would have not been possible without your support and encouragement. Thank you so much for caring, helping, and listening to my ideas. Thank you for your “brain-storming” sessions, which have taught me so much about the field of Oncology and Photobiology. I could have not have asked for a better supervisor!

I would also like to acknowledge the important contributions of Dr Mantas Grigalavicius to this work. Thank you Mantas for being an excellent collaborator. Your insightful input have been substantial contributor to the success of these projects. I am very grateful for everything you have taught me about Photobiology and for sharing your knowledge with me. I would also like to thank Dr Beata Grallert for filling me with creative ideas, sharing her outstanding lab knowledge, and always being so helpful. Our scientific talks have always been a great pleasure and a source of inspiration for me throughout the years.

I would also like to thank Professor Kristian Berg for being an immense source of knowledge in the field of Photomedicine and Photobiology and for always keeping your door open for discussions. Your insightful input has been substantial contributors to the success of these projects. I further wish to express my gratitude to all my collaborators for their significant effort, exceptional work, and fundamental guidance, including Professor Eirik Malinen, Associate Professor Nina Edin, senior engineer Efim Brondz, and Dr Ellen Skarpen.

I would like to thank everyone in the Protonic, PCI group, and the Department of Radiation Biology, for being exceptional colleagues and providing such a great working environment. A special thanks to Judith Wong, Tine Raabe, and Tuva Martin for all the scientific (and non-scientific) discussions and who have been amazing companions on this academic journey. It has been a pleasure having you in my everyday work life!

Finally, I would like to thank my parents and my partner Christoffer, for all their love, support, and patience. The faith that you have in me always encouraged and inspired me to do the best I can. This thesis is dedicated to you!

-My PhD work

Table of Contents

Acknowledgements	III
Table of Contents.....	1
Norwegian Summary of Thesis	3
List of Abbreviations	5
List of Publications	9
Aims of Study.....	11
Introduction	13
Chapter 1: Cancer and Glioblastoma	15
Chapter 2: Photodynamic Therapy as a treatment of Cancer	17
2.1 Photodynamic Therapy.....	17
2.2 Photophysics and Photochemistry.....	18
2.3 Intracellular targets for PDT	20
2.4 Photosensitizers for PDT	21
2.5 Photosensitizers used in the current work	22
Cercosporin	22
5-Aminolevulinic acid	23
2.6 Other relevant Photosensitizers for Photodynamic Therapy	26
Hypericin.....	26
Hypocrellin A.....	27
2.7 Photochemical Internalization (PCI).....	28
Basic mechanism of PCI	28
2.8 Adverse reactions and current limitations of PDT and PCI technology	30
Chapter 3: Ionizing Radiation.....	33
3.1 Types of radiation.....	33
Proton Therapy	34
3.2 Mechanisms of biological damage.....	35
Principles of “Photodynamic Therapy”	37
Chapter 4: Experimental Conditions	39
4.1 Cell lines	39
4.2 PDT	40
4.3 Cytotoxicity assays.....	41
4.4 Flow cytometry	41
4.5 Microscopy	42
4.6 Metabolic studies.....	43
4.7 Signal normalization and statistics.....	44
Summary of Publications	45
Paper I.....	47
Paper II.....	49
Paper III.....	51
Paper IV	53

Discussion.....	55
Chapter 5: Protodynamic Therapy – A combination of Proton Therapy and conventional PDT.....	57
5.1 Protodynamic Therapy	57
5.2 PDT in combination with gamma-radiation and X-rays	58
5.3 Current advances in Proton Therapy	58
5.4 5-ALA as a prodrug and PpIX cytotoxic effects in cancer	60
5.5 GSH and other ROS scavengers.....	62
5.6 Enzymes in heme synthesis and degradation.....	63
5.7 PpIX effects on cell metabolism.....	64
5.8 Cercosporin: A PS with promising applications outside conventional PDT.....	64
5.9 Cercosporin accumulation and intracellular localization.....	66
5.10 Cercosporin cellular metabolic effects.....	67
5.11 Excitation of PSs by protons	68
5.12 Effects of Protodynamic Therapy on cells	69
5.13 Protodynamic Therapy limitations	70
Chapter 6: Conclusions	71
6.1 Synopsis	71
6.2 Future Perspectives	72
Bibliography.....	73

Norwegian Summary of Thesis

Kreft i hjernen som glioblastoma multiforme (GBM), er praktisk talt uhelbredelig på grunn av dets beliggenhet og infiltrerende natur. Circa 28,000 nye tilfeller av ondartede gliomer diagnostiseres hvert år i EU og USA (Kilde: US National Cancer Registry). Dagens standarbehandling består av kirurgi, etterfulgt av strålebehandling og cellegift. Imidlertid gir denne behandlingen en begrenset overlevelsesfordel og kun 3% av pasientene lever lenger enn fem år etter første diagnose, noe som indikerer et høyt, uoppfylt medisinsk behov.

Fotodynamisk terapi (PDT) induserer behandlingseffekt gjennom synergien av tre viktige, men likevel individuelt ikke-kjemotoksiske komponenter: (i) det fotosensibiliserende legemiddelet (PS), (ii) lys med passende bølgelengde for å aktivere PS, og (iii) oksygen som omdannes til reaktive oksygenforbindelser (ROS) etter aktivering av PS med lys (Dougherty et al., 1998). PS-ene akkumulerer preferensielt i kreftvev og i særlig stor grad protoporfyrin IX (PpIX) som syntetiseres intracellulært ved administrasjon av 5-aminolevulinsyre (5-ALA). Fluorescensen fra GBM-spesifikke PS-ere brukes for tiden intraoperativt for å veilede reseksjon av GBM-svulster, mens behandling av GBM med PDT står overfor to problemer: (i) dårlig lysinntrengning i vev og (ii) manglende evne til å adressere spredt sykdom i hjernen.

Protonterapi benytter en ekstern stråle av akselererte protoner. Når protonene kommer inn i vevet, mister de energien ved elektromagnetiske interaksjoner i nesten konstant hastighet. Så snart de når en lavere terskelenergi, deponerer de denne gjenværende energien over en liten region, beskrevet som Bragg-toppen. På grunn av denne funksjonen, kan protonbehandling være lesjonsspesifikk, da den kan justeres for å avsette store mengder energi i kreftvevet og unngå eller redusere skade på normalvev. Protonbehandling kan imidlertid etterlate noen infiltrative grenseområder og nærliggende øyer med spredt sykdom ubehandlet. Dessuten, selv om protonbehandling er mindre skadelig for normalt vev enn gamma- eller røntgenstrålebehandling, er det fortsatt en ekstern strålebehandling med iboende risiko og bivirkninger tilsvarende andre strålebehandlinger.

For å kombinere styrkene til fotodynamisk terapi og protonterapi og samtidig overvinne deres individuelle begrensninger, har vi under utvikling proton-dynamisk terapi, en hybrid behandlingsmodalitet, hvor akselererte protoner brukes til å stimulere PS ved Coulomb-interaksjoner og fremkalle en 'PDT-lignende' effekt på toppen av proton-indusert ionisering, både til primær og spredt sykdom. Denne kombinasjonen kan forbedre effekten av kreftbehandlingen, gjøre den mer spesifikk og potensielt redusere bivirkninger ettersom PS akkumulerer spesifikt i kreftvevet. I arbeidet beskrevet i denne avhandlingen, forsøkte vi å dokumentere prinsippet om eksitering av elektronrike PS-ere med akselererte protoner, og åpnet for ny innsikt i og forbedring av behandling med protonterapi, spesielt siden PS-ene akkumulerer preferensielt i kreftvev. I denne doktorgraden ble 5-ALA-indusert protoporfyrin IX (PpIX) evaluert på grunn av den høye spesifisiteten for kreftvev og pågående klinisk utnyttelse. I denne studien, undersøkte vi mekanismer som potensielt kunne transformere 5-ALA-PDT fra en universell behandling til en mer krefttilpasset presisjonsterapi, for å forbedre effekten både i nåværende klinisk PDT-behandling, men også i nye applikasjoner som

‘Protodynamisk’ terapi. Disse mekanismene inkluderte cellespesifikk PpIX-produksjon, PpIX-utskillelse, cellemetabolisme, iboende antioksidant celleforsvar, enzymet som er ansvarlig for chelering av jern ved PpIX (ferrochelatase, FECH) og enzymet som er ansvarlig for initiell degradering av heme (heme oksygenase, HO-1). Vi har følgelig foreslått og validert viktige biomarkører, spesifikke for 5-ALA-PDT, som potensielt kan forutsi behandlingsresultatene og fungere som terapeutiske forsterkere av 5-ALA-PDT eller protodynamisk terapi etter hemning, spesielt for resistente cellerlinjer.

I denne studien, ønsket vi i tillegg å evaluere alternative tilgjengelige PS-ere som effektivt kunne fungere med protodynamisk terapi. Vi valgte blant annet cercosporin, en fotosensibiliserende forbindelse som dannes i enkelte arter i soppsekten *Cercospora*. Cercosporin ble funnet å være en kraftig singlet-oksygenprodusent ved lysaktivering sammenlignet med andre kjente PS-ere, og viste forbedret fotocytotoksitet ved lyseksitering i to GBM cellerlinjer (T98G, U87) og en brystadenokarsinom (MCF7) cellerlinje. Dette var et forventet resultat, siden vi registrerte lokaliseringen både i mitokondriene og endoplasmatisk retikulum, som er blant de mest fremtredende PDT-målene. Videre forårsaket cercosporin-PDT en metabolsk kollaps i alle undersøkte cellerlinjer, og reduserte både respiratoriske og glykolytiske aktiviteter. Vi undersøkte i tillegg effekten av cercosporin-PDT i GBM 3D-cellekulturer (sfæroider) som et modellsystem intermediært mellom 2D-cellekulturer og solide svulster. Igjen fant vi at cercosporin var en effektiv PDT PS, ikke bare i små, men også i større (1mm diameter) sfæroider som demonstrerte det store potensialet som PS. Cercosporin har i liten grad blitt evaluert for bruk i PDT pga. manglende absorpsjon av lys i det optiske vinduet hvor lys penetrerer tilfredstillende i vev (630-800 nm). Dette er imidlertid ingen begrensning ved protodynamisk behandling.

I arbeidet beskrevet i denne avhandlingen, demonstrerte vi for første gang at akselererte protoner kan aktivere PS-ere for å generere fluorescens og singlet oksygen i løsninger og geler. Vi dokumenterte også prinsippet om ‘protodynamiske’ effekter på to GBM-cellerlinjer, og la grunnlaget for en ny hybridterapi basert på protonterapi. Denne nye hybridmodaliteten benyttet i tillegg ROS generert av samspillet mellom akselererte protoner og PS-elektroner, for å eliminere kreftceller i synergisme med ioniserende bestråling. Vi er for tiden på et tidlig stadium i utviklingen av denne teknologien. Ytterligere validering i 3D-kulturer og *in vivo*-studier vil kaste mer lys over effekten av denne behandlingsmodaliteten, men ser for oss at denne nye terapien kan integreres i eksisterende protonterapieregimer.

List of Abbreviations

5-ALA	5-aminolevulinic acid
ABCG2	ATP-binding cassette super-family G member 2
AK	Actinic keratosis
ALAD	ALA dehydratase
ALAS	ALA synthetase
ATP	Adenosine triphosphate
BBB	Blood-brain barrier
BSO	Buthionine sulfoximine
BCC	Basal cell carcinoma
CNS	Central nervous system
CPOX	Coproporphyrinogen III oxidase
CRET	Complete resection of enhancing tumour
CT	Computed tomography
DABCO	1,4-diazabicyclo[2.2.2]octane
DAMPs	Damage-associated molecular patterns
DMSO	Dimethyl Sulfoxide
DNA	Deoxyribonucleic acid
DNA-PK	DNA-dependent protein kinase
DSBs	Double strand breaks
ECAR	Extracellular acidification rate
EPR	Enhanced Permeation and Retention
ER	Endoplasmic reticulum
ER	Estrogen receptor
ETC	Electron transport chain
FCCP	Carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazine
FDA	U.S. Food and Drug administration
FECH	Ferrochelatase
GPX4	Glutathione peroxidase 4

GSH	Glutathione
GSTP1	Glutathione S-transferase P1
GTR	Gross total resection
HER2	Human epidermal growth factor receptor 2
HMB	Hydroxymethylbilane
HP	Hematoporphyrin
HPD	Hematoporphyrin derivative
HPde	Hematoporphyrin dimethyl ether
IR	Ionizing radiation
LDH	Lactate dehydrogenase
LDL	Low-density lipoprotein
LED	Light-emitting diode
MAL	Methyl-5-aminolevulinate
MTT	3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
NAD(P)H	Nicotinamide adenine dinucleotide (phosphate)
OCR	Oxygen consumption rate
PII	Photofrin II
PBG	Porphobilinogen
PBGD	Porphobilinogen deaminase
PCI	Photochemical internalization
PDD	Photodynamic diagnosis
PDT	Photodynamic therapy
PpIX	Protoporphyrin IX
PPOX	Protoporphyrinogen III oxidase
PPpIX	Photoporphyrin IX
PR	Progesterone receptor
PrDT	Protodynamic therapy
ProCI	Proton-chemical internalization
PS	Photosensitizer

RNS	Reactive nitrogen species
ROS	Reactive oxygen species
RS	Radiosensitizer
RT	Radiation therapy
SCC	Squamous cell carcinoma
SDS-PAGE	Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis
SOD1	Superoxide dismutase 1
SOD2	Superoxide dismutase 2
SSBs	Single strand breaks
TNBC	Triple-negative breast cancer
UROD	Uroporphyrinogen III decarboxylase
UROS	Uroporphyrinogen III synthetase
UV	Ultraviolet
WHO	World health organization

List of Publications

- (I) **Cytotoxic and Photocytotoxic Effects of Cercosporin on Human Tumour Cell Lines.**
 Mastrangelopoulou M., Grigalavicius M., Berg K., Ménard M., Theodossiou T.A. Photochem Photobiol. 2019 Jan; 95(1):387-396

- (II) **Proton-dynamic therapy following photosensitiser activation by accelerated protons demonstrated through fluorescence and singlet oxygen production.**
 Grigalavicius M., Mastrangelopoulou M., Berg K., Arous D., Ménard M., Raabe-Henriksen T., Brondz E., Siem S., Görgen A., Edin N. F. J., Malinen E. & Theodossiou T. A. Nature Communications volume 10, Article number: 3986 (2019)

- (III) **Predictive biomarkers for 5-ALA-PDT can lead to personalized treatments and overcome tumour-specific resistances.**
 Mastrangelopoulou M., Grigalavicius M., Raabe-Henriksen T., Skarpen E., Juzenas P., Peng Q., Berg K., Theodossiou T.A. Cancer Reports (2020)

- (IV) **Photodynamic efficacy of cercosporin in 3D tumour cell cultures.**
 Grigalavicius M., Mastrangelopoulou M., Arous D., Juzeniene A., Ménard M., Skarpen E., Berg K., Theodossiou T.A. Photochem Photobiol. 2020 Mar 3.

Aims of Study

Photodynamic therapy (PDT) is a promising anticancer treatment, which combines visible light and a chemically non-toxic drug (photosensitizer, PS), which in the presence of oxygen can lead to the generation of reactive oxygen species (ROS), and subsequently cell death and tissue destruction at the tumour site where the light is directed. Although PDT has been known practically since ancient times, it is only in the recent decade that is becoming widely used. Its application varies according to the ailment and is also relevant to diseases other than cancer such as inflammatory and microbial diseases as well as photoaging/photorejuvenation (Wan and Lin, 2014). While PDT is an efficient treatment modality, it is facing limitations mostly related to the low penetration depth of light into tissue: (i) invasive intervention may be required for deep seated lesions, not accessible by light, (ii) partial eradication of tumour, and (iii) non applicability to non-diagnosed, disseminated disease since it is not irradiated by light.

Proton therapy is a form of tumour radiotherapy, which is using a targeted external beam of accelerated protons (ionizing irradiation), and which is recently becoming a mature technology. The protons mainly release this energy through their interaction with the electrons of matter, through Coulomb interactions. These interactions mainly lead to electron ionization, but also to electron excitation, depending on the energy transfer from the protons. In this sense, proton therapy can be cancer specific as accelerated protons can be adjusted to deposit high energy at the cancer site via the so-called Bragg peak.

Even though electron ionization from a proton beam traversing through matter and leading to radiotherapy is a well-studied phenomenon, electron excitation by accelerated protons is somewhat of an understudied process and not well described in bibliography.

The aim of the work leading to the present thesis was to access the possibility of the excitation of electron rich PSs by accelerated protons, opening a new perspective in proton therapy, especially since PSs have been shown to exhibit preferential accumulation in tumours. This preferential accumulation varies with tumour type and PS. In the work presented in this thesis, we have chosen the brain cancer glioblastoma multiforme (GBM) as our model, due to its difficult-to-access location and infiltrative nature, which makes it practically incurable. We have also identified two PSs for our study: (i) the naturally occurring dye cercosporin, which is very potent, yet not suitable for conventional PDT due to its short wavelength of activation and hence very shallow light penetration into tissue, and (ii) protoporphyrin IX (PpIX) derived through heme biosynthesis following the administration of the prodrug 5-ALA.

In this sense, the two main objectives in the studies leading to this thesis were: (i) to study the conventional PDT response of these PSs in GBM cell models, and (ii) provide proof of principle of the feasibility of PS activation during proton radiotherapy towards an additional "protondynamic" therapy effect.

The work demonstrating the achievements of these objectives can be found in the following publications:

Paper I

Cercosporin is a perylenequinone with photosensitizing characteristics, naturally occurring in fungi of the genus *Cercospora*, and it has only been studied for the purposes of phytopathology. The aim of this paper was to evaluate the photocytotoxic effects of cercosporin in human cancer cell lines (objective i).

Paper II

Proton therapy is an established tumour therapy that has been in clinical use for decades. Despite the fact that proton therapy is less harmful to normal tissue than gamma- or X-ray radiotherapy, it is still based on ionizing radiation and could potentially have radiation-related risks and side effects. The combination of proton radiotherapy and proton-induced PS activation could lead to a more effective tumour extermination without additional radiotoxicity or radiation-related carcinogenesis, as the additional killing would be through ROS production and not ionizing irradiation. The aim of this seminal publication was to provide proof of principle of the excitation and activation of PSs by accelerated protons to generate fluorescence, ROS, and ultimately tumour cell death (objective ii).

Paper III

5-aminolevulinic acid (5-ALA) is a natural precursor of the endogenously synthesized PS PpIX, in the heme biosynthetic pathway. 5-ALA-PDT is based on the external administration of 5-ALA to enhance PpIX production in cell mitochondria. Due to its higher accumulation in tumour cells compared to the normal cells, 5-ALA has been used as a prodrug to enable PDT and fluorescence guided surgery. The aim of this project was to highlight mechanisms that could potentially make ALA-based PDT more efficient not only in the clinical setting, but also in novel applications like "protondynamic" therapy (objective i).

Paper IV

In paper I, we explored the efficacy of cercosporin PDT in 2D cell cultures. In paper IV, we wanted to explore the efficacy of cercosporin PDT in a model system of intermediate complexity between solid tumours and 2D cell cultures: Cell spheroids (objective i).

Introduction

Chapter 1

Cancer and Glioblastoma

Cancer is one of the most life-threatening diseases of the 21st century with increasing incidence worldwide. It is the second leading cause of death globally with an estimate of 9.6 million deaths in 2018. According to the World Health Organization (WHO), the most prevalent types of cancer are lung cancer (1.76 million deaths), colorectal cancer (862 000 deaths), stomach (783 000 deaths), liver (782 000 deaths), and breast (627 000 deaths) (www.who.int).

Tumorigenesis in humans is a multistep process that drives the progressive transformation of normal human cells into malignant cells. It has been proposed that cancer cells have distortions in regulatory pathways that govern normal cell proliferation and homeostasis (Hanahan and Weinberg, 2000). This tumour progression advances through a process analogous to Darwinian evolution, in which a series of genetic alterations may confer growth advantages, ultimately leading to the progressive conversion of normal human cells into malignant cells (Foulds, 1954; Hanahan & Weinberg, 2000; Nowell, 1976). It has further been proposed that the transformation of normal cells to cancer cells is associated with six essential hallmark transformations in cell physiology (Fig. 1): (i) evading growth and proliferation suppressors, (ii) self-sustainability in growth signals, (iii) escaping apoptotic cell death, (iv) infinite replication capability, (v) angiogenic potential, and (vi) invasive and metastatic nature (Hanahan and Weinberg, 2000). In addition, two enabling characteristics for tumour progression are: (i) genome instability, which facilitates the generation of genetic diversity with selective advantages, and (ii) inflammation, which paradoxically enhances tumorigenesis and progression by assisting incipient lesions to obtain hallmark capabilities (Hanahan and Weinberg, 2011). With new developments in the past decade, there were two new emerging hallmark characteristics that were added in the concept: (i) deregulated control of energy metabolism that fuels cell growth, and (ii) division and ability to escape destruction by the host's immune system (Hanahan and Weinberg, 2011).

Gliomas are the most common tumours of the central nervous system (CNS) with glial cell origin, and they account for almost 80% of all malignant primary tumours in the brain (Agnihotri et al., 2013; Holland, 2000; Maher et al., 2001). According to the international standard by WHO, gliomas are classified into grade I to IV based on the level of malignancy, which are dependent on histopathological criteria. Grade I gliomas refer to lesions that have low proliferative potential and the possibility of cure solely following surgical resection. Grade II to IV gliomas are progressively more malignant and invasive with grade IV being most aggressive, invasive, and undifferentiated as designated by WHO (Jovčevska et al., 2013; Louis et al., 2007).

GBM is a grade IV glioma, virtually incurable due to its location and infiltrative nature, which makes it the deadliest malignant brain tumour with poor prognosis (Schucht et al., 2012; Tetard et al., 2014). Despite the advances in clinical oncology, GBM treatment is still the most challenging task (Mrugala, 2013). The current standard of care for GBM involves gross total surgical resection followed by radiation therapy and

chemotherapy with the median survival for primary GBM being 15 months (Stupp et al., 2005; Tetard et al., 2014). The poor prognosis of GBM is mainly linked to: (i) the extensive infiltrative nature of the tumour cells, impeding the efficacy of surgical resection, (ii) the blood-brain barrier (BBB) hindering the optimal delivery of chemotherapeutic agents, and (iii) various mechanisms of drug resistance (Haar et al., 2012; Tetard et al., 2014). It is evident from the above that developing an effective and safe treatment for GBM represents an urgent unmet medical need.

GBM is the selected cancer model for the present study, as we were interested to apply our new hybrid technology to difficult-to-cure and inaccessible cancers. "Protodynamic" therapy is expected to combine proton radiotherapy with singlet oxygen (ROS) mediated cell death, benefiting from the deep penetration of protons into tissue.

Figure 1.1: The hallmarks of cancer

Schematic illustration of the acquired functional capabilities associated with the transformation of normal cells into cancerous cells. Adapted from (Hanahan and Weinberg, 2000) and (Hanahan and Weinberg, 2011).

Chapter 2

Photodynamic Therapy as a treatment of Cancer

2.1 Photodynamic Therapy

Photodynamic therapy of cancer (PDT) is a clinically approved, minimally invasive therapeutic procedure that utilizes three key components- the photosensitizer (PS), light of the appropriate wavelength to excite the PS, and oxygen. Upon PS activation by light, these components initiate a photochemical reaction, resulting in cell photodamage, which may lead to cell death via apoptosis or necrosis (Fig. 2.1) (Agostinis et al., 2011; Dolmans et al., 2003; Dougherty et al., 1998). Apart from its main cytotoxic mode of action, PDT can eradicate cancerous lesions by two additional intertwined mechanisms. The first mechanism is the vascular shutdown of the capillary tumour vessels either by permanently destroying the vessel endothelium and obstructing the transport of oxygen and nutrients (Fingar, 1996), or by temporarily shutting the vessels down (e.g., by temporarily inactivating the vasodilation) (Korbelik et al., 2000), consequently leading to additional damage upon vessel reopening by reperfusion injury (Korbelik et al., 2003). The second mechanism is the induction of treatment-induced systemic immune response (Dougherty et al., 1998; Garg et al., 2012a; Garg et al., 2012b; Korbelik, 2006; Kousis et al., 2007; Zitvogel et al., 2011), through the release of damage-associated molecular pattern (DAMPs) molecules, induction of inflammation, and recruitment of immune cells. This may lead to abscopal immunity against the specific cancer.

Modern PDT application dates back to the early 1900s when workers used dyes in conjunction with light for the treatment of skin cancer (Castano et al., 2004). After the findings of Oscar Raab that acridine in combination with light could sensitize parameria, von Tappeiner and colleagues carried out extensive studies, which led to the introduction of the term 'photodynamic action' by von Tappeiner in 1904. Hematoporphyrin (HP) was used to that end shortly after the original studies by Raab, with the first study being conducted by Hausmann in 1908 (Lipson and Baldes, 1960). Thereafter and until 1960s, there have been sporadic studies both examining the selective localization of porphyrins in tumours and relapse after exposure to visible light (Castano et al., 2004; Figge et al., 1948). It was not before the discovery of hematoporphyrin derivative (HPD) in 1960 by Lipson and Baldes (Lipson and Baldes, 1960) that marked the modern era of PDT, followed by an eruption of interest fuelled by the pioneering studies of Dougherty (Dougherty, 1974; Dougherty et al., 1978; Dougherty et al., 1979). Clinically used PDT for cancer treatment dates back to the late 1970s, when Photofrin (HPD) was used in a clinical study involving patients with bladder cancer (Kelly and Snell, 1976). In the first large successful series of PDT using HPD, there was a complete or partial response observed in 111 of 113 malignant lesions, with none of the examined tumours being unresponsive (Dougherty et al., 1978).

The first PS to gain regulatory approval for the treatment of diverse cancers in many countries worldwide, including USA, was a semi-purified preparation of HPD referred to as Photofrin® (Castano et al., 2004). Despite being a promising substance,

Photofrin, being a "complicated mixture of ambiguous structure" (Kessel and Thompson, 1987), was presented with a number of shortcomings: (i) prolonged skin sensitivity to sunlight (Baas et al., 1995), (ii) inferior tumour selectivity (Orenstein et al., 1996), and (iii) poor light penetration into tissue due to its short activation wavelength (~ 630 nm) (Spikes, 1990). Recently, there have been many studies on developing new PSs, with great potential for clinical application. Some PSs are naturally produced (Daub et al., 2005), while some others can be produced by partial synthesis from ample natural basic components, such as protoheme, chlorophyll, and bacteriochlorophyll, which is both economically and environmentally advantageous, compared to total chemically synthesized drugs (Nyman and Hynninen, 2004).

Figure 2.1: Mechanisms of Photodynamic action

Following PS administration, the PS is systemically distributed in the tissue and selectively accumulates in the cancer cells. Light illumination of a specific wavelength in conjunction with the presence of O₂ activates the PS, triggering a series of reactions leading to the production of ROS and ¹O₂, via Type I and Type II reactions, respectively.

2.2 Photophysics and Photochemistry

Under dark ambient conditions, PSs exist in their singlet ground state (paired electrons of antiparallel spins in the low energy molecular orbital, Fig. 2.2). Once the photons of the appropriate energy interact with the PS, they excite its electrons to the first singlet excited state where the electrons are still paired (antiparallel spins). This short-lived species (nanoseconds) can decay back to the ground state giving the energy back as light (fluorescence) or by internal conversion (non-radiative decay). The excited singlet state PS may also undergo intersystem crossing to its relatively long-lived (microseconds) triplet excited state, where the electrons are not paired (parallel spins)

(Turro, 1991). The long lifetime of the PS triplet-state is due to the fact that the transition back to the singlet ground state, by the emission of light (phosphorescence), is 'spin-forbidden' (Turro, 1991).

In the triplet state, the PS can undergo two kinds of reactions. It can interact with molecular oxygen within the tissue either via charge transfer (type I reactions) or energy transfer (type II reactions). In type I reactions, one electron is usually donated to oxygen thus reducing it to superoxide anion and consequently creating a cascade of ROS. As the superoxide is dismutated by superoxide dismutase (SOD) to H_2O_2 and subsequently transition metals (e.g., Cu^+ or Fe^{2+}), it can generate hydroxyl radicals through Fenton reactions. In type II reactions, the PS in its triplet state interacts with oxygen, which is the only element found in a triplet ground state, in a collisional manner transferring energy and causing a spin flip to both the PS, which returns to its singlet ground state, and the triplet oxygen, which transitions to its excited singlet state. Singlet oxygen is a type of ROS with high toxicity and is the most prominent therapeutic species in PDT (Turro, 1991). Due to the high reactivity and short half-life of the singlet oxygen and hydroxyl radicals produced in biological environments, only areas adjacent to the PS are directly affected by PDT. In particular, singlet oxygen has a half-life time of <40 ns in biological systems, thus its radius of action is short-ranged, of the order of 20 nm (Nowell, 1976).

Figure 2.2: Jablonski Diagram

Light with sufficient energy and wavelength matching the absorption spectrum of the PS can excite the PS from its ground state (S_0) by moving electrons from a low energy-state to a high energy-state. The activated PS (S_1) can either undergo intersystem crossing entering its triplet state (T_1), or lose its energy by emission of fluorescence or non-radiative decay. Subsequently, the PS may transition from T_1 to S_0 , with the simultaneous release of phosphorescence, or react with other molecules (e.g., O_2) via Type I and Type II reactions.

2.3 Intracellular targets for PDT

The most critical factor governing the direct cytotoxic outcome of PDT is the interaction of the PS with crucial intracellular targets within the tumour tissue, which depends on the intracellular localization of the PS. ROS including singlet oxygen have a relatively short half-life and short range from the site of their generation, due to the density of targets along the path of ROS propagation. It is therefore important for the ROS to be generated close to crucial intracellular targets, damage to which may prove fatal to the cells.

Mitochondria, endoplasmic reticulum (ER), lysosomes, plasma membrane, and nuclei of the tumour cells have been assessed as potential PDT targets (Dougherty et al., 1998). The localization of the PSs inside the cells is mainly dictated by the physicochemical properties of PSs; PSs, which are hydrophobic, may diffuse across the plasma membrane, and then relocate to other intracellular membranes, while PSs that are positively charged and hydrophobic can localize in the mitochondria (Castano et al., 2004). Mitochondria have been considered a key subcellular target for many

PSs used in PDT (Morgan and Oseroff, 2001). This is due to the tendency of many PSs to induce apoptosis via mitochondrial damage (Castano et al., 2004), however high PDT doses, tend to move the balance towards a necrotic cell death (Kessel and Reiners, 2007). The concentration and physicochemical properties of the PS, the concentration of oxygen, the wavelength and intensity of light, and the cell type specific properties may all influence the mode and extent of cell death (Castano et al., 2005).

2.4 Photosensitizers for PDT

A great number of natural and synthetic photoactive compounds have photosensitizing potential (Ebermann et al., 1996). The majority of the PSs used in cancer treatment are based on a tetrapyrrole structure, comparable to that of protoporphyrin, precursor of heme. An ideal PS compound should possess the following properties: (i) it should be a pure compound with good stability in storage; (ii) it should have a strong absorption in the red to deep red region of the spectrum (600-800 nm), considering that the penetration of light into tissue increases with its wavelength (Fig. 2.3); (iii) it should have no dark toxicity and relatively rapid excretion from normal tissues, consequently minimizing phototoxic side effects. Other desirable characteristics of an ideal photosensitizer include: (iv) low chemical (dark) toxicity; (v) high ratio of cancer to normal tissue accumulation; (vi) high quantum yield of singlet oxygen and high production of singlet oxygen in vivo; (viii) cost effectiveness and commercial availability; and (viii) solubility compatible with easy systemic delivery, e.g., through injections (Agostinis et al., 2011; Allison et al., 2004; Konopka and Goslinski, 2007).

Figure 2.3: Light penetration into tissue

Blue light penetrates less efficiently through tissue, while red and infrared light penetrate deeper into tissue. Each point in the graph represents the tissue depth at which the incident light is attenuated to $1/e$ of its original intensity. Adapted from (Macrobert, 2005).

2.5 Photosensitizers used in the current work

Cercosporin

One family of photosensitizers are the perylenequinones. Perylenequinone PSs share a core phenolic quinone chromophore and predominantly differ in their side chains. Upon light activation, they produce various types of ROS such as singlet oxygen ($^1\text{O}_2$), hydroxyl radical ($\text{OH}\cdot$), superoxide ($\text{O}_2^{\cdot-}$), and hydrogen peroxide (H_2O_2) (Daub and Ehrenshaft, 2000; Daub et al., 2005). Perylenequinones have gained attention because of their therapeutic potential and photodynamic activity (Olivo and Chin, 2006). Some of the perylenequinones studied extensively with regard to their photodynamic properties are hypericin (Agostinis et al., 2002; Theodossiou et al., 2009) and hypocrellins A and B (Ali and Olivo, 2003; Zhenjun and Lown, 1990). Numerous phytopathogenic fungi produce nonspecific, light activated perylenequinone toxins involved in pathogenesis of their hosts. These toxins act as PSs, absorbing light energy and generating ROS that damage the membranes of the host cells. Cercosporin is a non-host-selective photosensitizing compound produced by members of the fungal genus *Cercospora* and it is crucial to the ability of these fungi to parasitize plants (Daub, 1982). *Cercospora* species give rise to disease on a variety of plants worldwide including corn, sugar beet, tobacco, coffee, soybean, and diverse ornamental and weed species causing a major economic burden (Daub et al., 2005). Cercosporin was first isolated in the 1950s from *Cercospora kikuchii*, a soybean pathogen (Kuyama and Tamura, 1957), and since then it has been isolated from numerous *Cercospora* species and infected plants (Daub and Ehrenshaft, 2000). Cercosporin is a deep red pigment firstly reported for its photosensitizing abilities by Yamazaki in 1975, who demonstrated that cercosporin had toxic effects on both mice and bacteria, and that the toxicity was dependent on light and oxygen availability (Yamazaki et al., 1975).

Cercosporin has been shown to produce both singlet oxygen and superoxide when irradiated by light, with singlet oxygen being the dominant species, generated both in vitro and in vivo (Daub, 1992; Daub and Hangarter, 1983). Singlet oxygen and superoxide are highly toxic to cells leading to oxidation of fatty acids, sugars, guanine, and several amino acids consequently resulting in DNA damage, inactivation of enzymes, and cell membrane destruction (CS, 1976; Daub, 1982). As a result of the production of ROS, cercosporin is in essence universally phototoxic. Besides its well documented toxicity to plants, cercosporin has also been shown to be toxic to mice, gram-positive and gram-negative bacteria, fungi, and Oomycetes (Daub, 1987b; Yamazaki et al., 1975). In addition, cercosporin has been demonstrated to have antiviral activity and to be an effective inhibitor of protein kinase C (Hudson et al., 1997; Tamaoki and Nakano, 1990). The photodynamic activity of cercosporin has been studied in various types of natural and artificial membranes and its toxicity has been attributed to its ability to cause cellular lipid peroxidation (Cavallini et al., 1979). This is due to the fact that cells that are being damaged by cercosporin demonstrate an accumulation of lipid peroxidation products, a pronounced increase in saturated fatty acids as compared to unsaturated, and a decrease in the plant membrane fluidity (Daub and Briggs, 1983). Cercosporin exhibits a short excitation wavelength (max ~470 nm), and therefore it could only be clinically relevant to superficial, shallow

lesions. Nevertheless, for applications where the PS excitation does not depend on the limitations of light penetration into tissue, cercosporin can prove an invaluable asset.

5-Aminolevulinic acid

5-aminolevulinic acid (5-ALA) is a metabolite in the heme biosynthetic pathway in humans (Fig. 2.4). This pathway is catalyzed by eight enzymes, and it takes place partially in mitochondria and partially in the cytosol of nearly all cells containing a nucleus (Herceg et al., 2017; Ponka, 1999; Yang et al., 2015). The first step of the pathway takes place in the mitochondrion, and it involves the generation of 5-ALA from glycine and succinyl coenzyme A by ALA synthetase (ALAS). ALA then migrates to the cytosol, where two molecules of ALA are combined to make porphobilinogen (PBG), a reaction catalyzed by ALA dehydratase (ALAD). Subsequently, four molecules of PBG are merged to form a linear tetrapyrrole, hydroxymethylbilane (HMB), by porphobilinogen deaminase (PBGD). Thereafter, HMB is circularized to form uroporphyrinogen III by uroporphyrinogen III synthetase (UROS). Uroporphyrinogen III is then decarboxylated by uroporphyrinogen III decarboxylase (UROD) forming coproporphyrinogen III, which is imported back into the mitochondria. Coproporphyrinogen III in turn undergoes oxidative decarboxylation by coproporphyrinogen III oxidase (CPOX), to produce protoporphyrinogen III, which is then oxidized by protoporphyrinogen III oxidase (PPOX) to generate PpIX. PpIX, which is the 5-ALA-derived PS, with the assistance of ferrochelatase (FECH), subsequently incorporates ferrous iron to form heme, which becomes "muted", both with respect to fluorescence and singlet oxygen generation, and is therefore not a PS (Ponka, 1999; Yang et al., 2015).

Figure 2.4: Schematic representation of the heme biosynthetic pathway

5-ALA is a natural precursor of heme in the heme biosynthetic pathway that takes place partially in the mitochondria and partially in the cytosol. Porphobilinogen (PBG), uroporphobilinogen III (URO III), coproporphyrinogen III (COPRO III), protoporphyrinogen III (PROTO III), protoporphyrin IX (PpIX).

5-Aminolevulinic acid in clinics

5-ALA-PDT has been among the most active agents/prodrugs of clinical PDT with promising outcomes, including outstanding cosmetic results (Peng et al., 1997b). One of the main advantages of 5-ALA-PDT is its great specificity for cancer cells, since PpIX production is much higher in tumour cells (Peng et al., 1997b). Its additional selectivity may be attributed by its administration as a precursor. 5-ALA, in contrast to other exogenously administered PSs such as Photofrin, is a photodynamically inactive, non-selective, and non-toxic compound that is intracellularly metabolized to the photodynamically active PpIX (Wachowska et al., 2011). Although nearly all human cell types express the enzymes involved in the heme biosynthetic pathway, a differential activity of the enzymes in the tumour cells, as compared to normal cells,

leads to a higher PpIX accumulation within the neoplastic cells (Collaud et al., 2004; Wachowska et al., 2011). This has primarily been attributed to: (i) the significantly lower FECH situated in the inner mitochondrial membrane (Teng et al., 2011), leading to reduced efficacy of PpIX-to-heme conversion and subsequently PpIX accumulation within cancer cells, and (ii) the notably enhanced PBGD activity, which is leading to augmented PpIX production (el-Sharabasy et al., 1992; Kondo et al., 1993; Rubino and Rasetti, 1966; Schoenfeld et al., 1988; Van Hillegersberg et al., 1992). Nevertheless, other factors, such as limited availability of free iron, may also contribute to increased PpIX production in cancer cells (Rittenhouse-Diakun et al., 1995). PpIX is converted to heme within 24 hours after 5-ALA administration, eliminating prolonged photosensitivity of patients, which is one of the side-effects associated with PDT (Wachowska et al., 2011). In addition, 5-ALA is a relatively small (<200Da) and hydrophilic molecule (Donnelly et al., 2007), making its systemic administration easier than other typically larger, lipophilic PSs routinely used in clinical PDT.

Due to the effectiveness of PpIX as a PS, 5-ALA was initially introduced into the clinic in 1990 by Kennedy (Kennedy et al., 1990), who first reported the treatment of 80 cases of basal cell carcinoma (BCC) using topical 5-ALA-PDT with success. Since then, 5-ALA-PDT has been used for the treatment of various neoplastic and non-neoplastic diseases, and has become the treatment option primarily for dermatological malignancies. 5-ALA-PDT has been successful in the treatment of BCC, actinic keratosis, Bowen's disease, vulval intraepithelial neoplasia, vulval Paget's disease, and cervical intraepithelial neoplasia (Donnelly et al., 2007). Due to the high accumulation of PpIX in neoplastic cells, 5-ALA-elicited photodynamic diagnosis (PDD) has been clinically applied to many cancer types (Koizumi et al., 2016). These include mouth (Leunig et al., 2001), endometrium (Wyss-Desserich et al., 1996), cervix (Keefe et al., 2001), bladder (Inoue et al., 2012), prostate (Fukuhara et al., 2011), and brain malignancies (Stummer et al., 2000; Stummer et al., 1998). In addition, it has been clinically used for the fluorescence-guided resection of brain and bladder tumours with promising results (Jichlinski and Leisinger, 2001; Tetard et al., 2014; Yang et al., 2015). Systematic GBM intraoperative 5-ALA-PDT has the potential to facilitate excision of more cancer cells, and hence significantly prolong patients' survival, since according to clinical studies, the recurrence of gliomas is common due to the residual malignant cells omitted after treatment (Teng et al., 2011). 5-ALA was approved by the U.S. Food and Drug administration (FDA) in 2017, as an adjunct for the visualization of tissue malignancies, in grade III or IV gliomas (NDA 208630/SN0014).

5-ALA can be applied topically or systemically for PDT of skin and other tumours such as BCC and gastrointestinal adenocarcinoma (Cairnduff et al., 1994; Calzavara-Pinton, 1995; Fromm et al., 1996; Kennedy et al., 1990; Regula et al., 1995; Svanberg et al., 1994; Wolf et al., 1993). Topical 5-ALA-PDT offers several advantages over therapies with conventional PSs and other mainstream treatments: (i) it produces excellent cosmetic results in comparison to surgery, (ii) it is non-invasive, (iii) it is well tolerated by patients, (iv) it can be applied to treat multiple superficial lesions in short treatment sessions, (v) it can be offered to patients who decline to undergo surgery or have pacemakers and bleeding tendency, (vi) it can be applied to treat lesions in

specific sites, unlike most chemotherapeutic regimens, (vii) it can be practiced as a palliative treatment to relieve symptoms, and (viii) it can be repeated as many times as required due to its minimally toxic effects, low risk of treatment resistance, and low carcinogenic potential (Peng et al., 1997b). For its numerous advantages 5-ALA-PDT has received worldwide approval and is currently routinely used in dermatology (Zhao and He, 2010). Its effectiveness in treating both other types of cancer and non-cancer related diseases are being actively investigated as well (Nokes et al., 2013; Szeimies et al., 2002).

Despite the benefits of 5-ALA-PDT, there are some limitations that need to be overcome to enhance its therapeutic potential. 5-ALA is a polar compound and exists primarily as a charged zwitterion in physiological pH, which attributes its low lipid solubility and reduced bioavailability. As a hydrophilic molecule, 5-ALA does not penetrate efficiently through the skin or cell membranes. Consequently, PpIX production after topical ALA administration is restricted to a 2-3 mm skin surface penetration, which may be inefficient to evoke satisfactory photosensitization (Peng et al., 1997a). This could be overcome by ALA ester derivatives, such as methyl-5-aminolevulinate (MAL), which would improve the bioavailability of both systemically and topically administered 5-ALA. MAL is currently used in the clinics similarly to 5-ALA (Gold, 2009). Due to its methyl group, MAL is more lipophilic and has better selectivity for tissue, comprising a good alternative particularly in the case of deep lesions such as in squamous cell carcinoma (SCC). To increase drug penetration through the skin, ethyl-ALA, propyl-ALA, and butyl-ALA have also been produced with improved skin penetration after topical application in vivo (Lopez et al., 2004). In addition, two other prodrugs hexyl-ALA and octyl-ALA have been developed and tested in vitro (De Rosa et al., 2003; Gaullier et al., 1997; Gederaas et al., 2001; Rud et al., 2000) and in vivo (Neittaanmäki-Perttu et al., 2016) showing promising results.

Although 5-ALA-PDT has high specificity to cancers, differences in treatment outcomes impose the need for predictive biomarkers to better stratify patients, and to personalize the treatment depending on each individual's genotypic and phenotypic characteristics in their lesions. Therefore, in our paper III, we sought to identify and highlight key biomarkers that may predict treatment outcome and concomitantly be exploited to overcome cancer-specific resistances to 5-ALA-PDT.

2.6 Other relevant Photosensitizers for Photodynamic Therapy

Hypericin

Hypericin is a phenanthroperylenequinone, naturally occurring in plants of the *Hypericum* genus, in particular *Hypericum perforatum* (St. John's wort) (Jendželovská et al., 2016; Theodossiou et al., 2009). It has similar structure to the perylenequinones but with a fused octacyclic core (Mulrooney et al., 2012). Hypericin has a broad pharmacological activity, among others, antidepressant and anti-viral capabilities as well as being a powerful PS, that is suitable as a PDT agent of various oncological diseases. Hypericin is not only a potent PS for PDT but also has been proven useful in the context of PDD, as an effective fluorescent marker for tumour visualization.

Hypericin has several advantages that makes it a powerful fluorescent PS suitable for applications in both PDT and PDD: (i) it shows minimal or no toxicity under dark conditions (Feruszová et al., 2016; Jendželovská et al., 2014), (ii) it accumulates preferentially in tumorigenic tissues (Kamuhabwa et al., 2002; Noell et al., 2011), and (iii) in the presence of light and oxygen, it produces ROS (Diwu and Lown, 1993). Imaging studies have revealed that hypericin accumulates in the membranes of the ER, Golgi apparatus, lysosomes, and mitochondria (Agostinis et al., 2002; Ali and Olivo, 2002; Galanou et al., 2008; Mikeš et al., 2011), although cellular uptake and intracellular localization may be dependent on its lipophilicity, incubation concentrations, and/or interaction with serum lipoproteins (Jendželovská et al., 2016).

The cytotoxic effects of hypericin upon photoactivation has been demonstrated in vitro in various cancer cell lines (Kleemann et al., 2014; Koval et al., 2010; Mikešová et al., 2013; Theodossiou et al., 2009; Theodossiou et al., 2006; Theodossiou et al., 2017). Furthermore, recent preclinical and clinical studies have demonstrated that PDT with hypericin might offer effective treatment of some skin tumours, carcinomas, and sarcomas (Jendželovská et al., 2016). Due to its fluorescent properties and specificity for neoplastic tissue, hypericin-PDD has been assessed for diverse clinical practices such as optical tumour imaging as well as targeting, monitoring, and detecting tumour stages and grades. Currently, hypericin-PDD has been clinically evaluated in bladder, head and neck cancers, and gliomas with promising results in all but head and neck carcinomas (Jendželovská et al., 2016). Thus, hypericin might be a very potent agent in cancer treatment and diagnosis, and further studies are needed to unveil the full potential of this PS.

Hypocrellin A

One class of perylenequinones that has been extensively studied is the *Hypocrellin* family, isolated from the species *Hypocrellin Bambusae* (Mulrooney et al., 2012). The naturally occurring hypocrellin compounds are of great interest due to their potential anticancer properties, along with their possible benefits in treating skin conditions, such as psoriasis and acne (Zhenjun and Lown, 1990). Hypocrellins possess most of the characteristics of an ideal PS including: (i) easy preparation and purification, (ii) high quantum yield of singlet oxygen, (iii) high phototoxicity, (iv) low dark toxicity, and (v) rapid elimination from normal tissues. This confirms that hypocrellins are a promising phototherapeutic regiment.

Natural hypocrellins primarily include two components, hypocrellin A and hypocrellin B, with the former comprising 95% of hypocrellins (Jiang and He, 2001). Following hypocrellin A photosensitization, the production of multiple ROS has been observed such as singlet oxygen ($^1\text{O}_2$), superoxide anion ($\text{O}_2^{\cdot-}$), hydrogen peroxide (H_2O_2), and hydroxyl radicals ($\text{OH}^\cdot$) (Zhenjun and Lown, 1990). Thus, both type II and type I reactions are thought to be involved in photosensitization by hypocrellins. Hypocrellins are lipophilic organic compounds, which enhance cellular uptake (Engelhardt et al., 2011) but hinder drug-delivery and bioavailability (Ishikawa and Hashimoto, 2011). In comparison to other PSs, hypocrellin fluorescence is very prone to microenvironment changes because of the intramolecular H-atom transfer process signifying that

alterations in the intramolecular hydrogen bond will affect fluorescence properties (Xu et al., 2003). Taken that hypocrellins exhibit differential fluorescence responses when exposed to various biomolecules and pH values, pinpoints the potential of hypocrellins as probes to monitor microenvironmental changes due to the unique properties tumour cells and tissues possess in relation to healthy tissue (Deng, 2015).

2.7 Photochemical Internalization (PCI)

PCI, initially proposed in 1999, is a novel concept for the specific cytosolic delivery of biological macromolecules bypassing the intracellular barrier of endosomes and lysosomes (Berg et al., 1999). PCI is based on the use of PSs specifically located in the membrane of endocytic vesicles. Upon light activation, the PSs cause endosomal membrane rupture and subsequent release of the endocytosis-loaded contents of endocytic vesicles into the cytoplasmic matrix (Berg et al., 2011). This has made possible the enhanced delivery of a number of macromolecules, including ribosome-inactivating toxins, gene encoding plasmids, adenoviruses and oligonucleotides, nanoparticles and certain chemotherapeutics (Jerjes et al., 2020). PCI has been evaluated for use not only in the treatment of cancer but also other diseases, such as intracellular bacterial infections (Zhang et al., 2018) and rheumatoid arthritis (Dietze et al., 2005).

Basic mechanism of PCI

PSs may enter the cells either directly through the plasma membrane or indirectly by several endocytic pathways (Berg et al., 1999; Jerjes et al., 2020). Endocytosis is a biological process that takes place in all nucleated cells (Grant and Donaldson, 2009) and involves the formation of vesicles intracellularly, which evolve into early and late endosomes. Subsequently, late endosomes may fuse with lysosomes leading to the breakdown of the endocytosed material by means of hydrolytic enzymes. Some PSs are not able to permeate the plasma membrane, and therefore they enter the cell through endocytosis. Endocytosis may occur via phagocytosis, receptor-mediated (clathrin-mediated) and non-clathrin/caveolae-mediated endocytosis, pinocytosis or adsorptive endocytosis. The most hydrophilic PSs, such as tetracarboxylic and tetrasulfonated PSs, cannot efficiently infringe into the plasma membrane and are typically taken up by cells through pinocytosis, which is a low efficacy uptake mechanism (Jerjes et al., 2020). Amphiphilic PSs burrow into the plasma membrane with their lipophilic parts, while their hydrophilic parts “pull” them towards the aqueous phase, out of the cell. These PSs integrate into the outer leaflet of the plasma membrane and subsequently end up in the inner leaflet of the endocytic vesicles through adsorptive endocytosis (Norum et al., 2009). Cellular uptake of PSs can also occur through receptor-mediated endocytosis when linked to targeting moieties with affinity for a plasma membrane receptor.

The mechanism of PCI is illustrated in Fig. 2.5. As mentioned earlier, the amphiphilic PSs, intruding into the plasma membrane but unable to penetrate it, will enter the cells via adsorptive endocytosis and will consequently accumulate in the inner leaflet of the

endocytic vesicles. Upon light irradiation, the PS induces the formation of ROS and predominantly $^1\text{O}_2$ within the endocytic membranes leading to lipid peroxidation and rupture. Consequently, this perforation leads to the cytosolic release of the cargo loaded inside the vesicles by endocytosis. In the absence of light exposure, the endocytosed agents remain entrapped within the endocytic vesicles. Upon the maturation and/or fusion of the latter with lysosomes, their contents will eventually become degraded by hydrolytic enzymes (Jerjes et al., 2020). Of note, hydrophilic PSs localized in the matrix of the endocytic vesicles are unable to induce any 'PCI-effect', underlining the importance of the intravesicular localization of the PS (Norum et al., 2009). Initially, it was assumed that for the endocytosed cargo to be released, it had to be localized in the same endocytic compartments with the PS, simultaneously with light irradiation. However, studies have shown that photochemical treatment up to 8 h prior to the cargo delivery does not result in reduction of the PCI effect (Prasmickaite et al., 2002), suggesting that newly formed vesicles can fuse with photochemically damaged vesicles.

Figure 2.5: Schematic illustration of the PCI technology

Administered drugs of choice are taken up by the cell through endocytosis. At the same time, amphiphilic PSs “lodge” in the cell plasma membrane, which subsequently becomes the endosomal membrane, and if left untreated both endocytic cargo and PS end up in the lysosomes with which endocytic vehicles fuse. Inside the lysosomes, the drugs are exposed to enzymatic hydrolases and hence are degraded and deactivated. Alternatively, generation of ROS through the activation of the membrane-anchored PS by light will disrupt the endocytic vesicle membranes before degradation and release the active drug in the cytosol. The released drugs can thereby reach their intracellular targets and exert their therapeutic effects.

2.8 Adverse reactions and current limitations of PDT and PCI technology

The major limitation of PDT and PCI is the low depth of penetration of visible light in tissue (<2cm) (Benov, 2015), thereby rendering the treatment of deep-seated tumours challenging and potentially leading to incomplete eradication of tumour cells. Since tissue oxygenation is essential for the photodynamic reaction to take place, the PDT effect in hypoxic tumours, tumours surrounded by necrotic tissue or dense tumour masses is severely reduced (Calixto et al., 2016). In the case of PCI, even though a significant fraction of sequestered molecules are released to the cytosol, some of the

drug molecules proximal to the membranes of the endocytic vesicles may be photoxidized by the singlet oxygen produced, rendering them inactive. This can also be the case when the carrier is lipophilic, e.g., in the case of gene therapy (Hellum et al., 2003; Prasmickaite et al., 2000), and may be overcome with the use of more hydrophilic gene vectors. Another possible limitation of PDT and PCI, as in the case of surgery and radiation therapy, is its inability to eradicate occult, undiagnosed metastasis, since the light is selectively applied to the primary lesion. Interestingly, however, PDT may confer enhanced anti-tumour immunity as shown in pre-clinical models (Agostinis et al., 2011), suppressing occult, disseminated cancers.

Compared to surgery and radiation therapy, PDT is a less invasive treatment modality and the adverse reactions are rather mild and not enduring. Depending on the PS and the therapeutic protocol, adverse events associated with PDT, which are predictable and treatable (Brown et al., 2004), include erythema, fever, pleural effusion, constipation, anaemia, and respiratory insufficiency (Ochsner, 1997). The prominent side effects in the immediate post-PDT phase include treatment-site pain and swelling (Attili et al., 2011; Ibbotson, 2011; Ochsner, 1997), which arise around 24-48 h, and can last up to 14 days. Another common side effect when using a systemic PS is residual photosensitivity, which can last for months (Benov, 2015). Of note, most side effects can be alleviated by the proper selection of PS and PS dosage, parameters of illumination, and other factors of the PDT treatment protocol. The ideal PS and light doses as well as drug-light treatment time interval may differ between patients or treatment indications, which prevents the application of standardized protocols and the attainment of optimal response rates (Benov, 2015).

Chapter 3

Ionizing Radiation

3.1 Types of radiation

Radiation is present in our everyday lives, arising both from natural and artificial sources. Radiation greatly affects both healthy and diseased tissues with a wide range of responses. These responses are determined by a number of parameters including: (i) the radiation type/source, (ii) radiation dose, (iii) duration of exposure, and (iv) the individual genetic and epigenetic composition (Reisz et al., 2014).

Radiation is classified into two major categories, namely ionizing (IR) and non-ionizing (non-IR) (Fig. 3.1). The largest part of the environmental radiation is non-ionizing, such as ultraviolet (UV) radiation and visible light from the sun, and electromagnetic radiation resulting from microwaves and radio waves. It has become evident that sources of non-IR, such as UV rays from the sun or tanning sunbeds, may cause damage to biomolecules (Karagas et al., 2002; Zhang et al., 2012). However, the interaction of IR with biological molecules is significantly more aggressive than non-IR due to its ability to induce atom ionization.

IR is defined as the radiation, which can ionize atoms, i.e., strip away one or more electrons from the atomic or molecular orbitals into the continuum (Han and Yu, 2009). Due to this ability of IR, its interaction with biological systems is considerably more vigorous than non-IR, and therefore resulting in profoundly more damage. The major classes of IR are alpha particles, beta particles, X-rays, and gamma-rays. Even though neutrons are not directly ionizing, they are classified as ionizing radiation due to the cascade of secondary events in their interaction with matter leading to ionization. Alpha radiation possesses the lowest penetration capability, and even though it can be stopped by physical barriers, such as paper or skin (El Ghissassi et al., 2009; Reisz et al., 2014), it still has the greatest ionizing ability as compared to beta or gamma-radiation. Beta radiation consists of high energy electrons, it is highly ionizing, and although it has a somewhat higher penetration than alpha radiation, it can still be stopped by relatively thin physical barriers. On the other hand, X- and gamma-rays are more penetrating, and therefore environmental exposure (cosmic sources, atmospheric exposure, food and drinks, and ground and building materials) to gamma-rays, which have higher energy than X-rays, leads to more extensive biological damage than exposure to alpha or beta particles (Reisz et al., 2014). All the four types of radiation mentioned are valuable tools in current medicine either for diagnostic or therapeutic purposes (Reisz et al., 2014). X-ray radiation is commonly used as a diagnostic tool, e.g., in plain X-ray images or computed tomography (CT) scanning, while both gamma- and X-rays are used in radiotherapy as external-beam radiation therapies for cancer (Leibel et al., 2003). Targeted alpha particle therapy (Wadas et al., 2014) and beta emitters (Jødal, 2009) are also of increasing use in cancer treatment.

Currently, external-beam radiation cancer therapy aims to destroy neoplastic formations, sparing the adjacent normal tissue. However, radiation-driven damage

and treatment-induced secondary carcinogenesis are major concerns. Another type of radiotherapy is proton beam radiotherapy. Anticancer treatment by accelerated protons allows for specific targeting of therapeutic doses [of about 60-80 Gy, depending on the indication] to the lesions and no exit dose. These attributes make proton therapy an attractive choice, particularly for the treatment of deep-seated tumours (Loeffler et al., 1997), tumours located adjacent to critical structures, such as spinal cord, eyes, and critical brain structures as well as for paediatric malignancies (Levin et al., 2005).

THE ELECTROMAGNETIC SPECTRUM

Figure 3.1: The electromagnetic spectrum

The electromagnetic radiation spectrum may be divided into multiple categories according to the frequency and/or amplitude of the electromagnetic fields. The spectrum may also be divided into two main ranges, namely IR and non-IR depending on the amount of energy the radiation possesses and the potency of changes in physical matter.

Proton Therapy

The history of proton therapy dates back in 1946 when Robert Wilson hypothesized that an accelerator-produced beam of protons could be utilized to treat deep-seated tumours in humans while minimizing radiation exposure of adjacent normal tissues. The first studies on proton irradiation to test this hypothesis were initiated at the Lawrence Berkeley Laboratory (Olsen et al., 2007), with the first human being treated with protons in 1954 (Lawrence et al., 1958).

Protons, as is the case for all charged particles, experience energy loss upon their entrance into tissue (Levin et al., 2005). Once the interaction with matter begins, the proton beams produce energy depositions with a relatively low dose in the initial part of their path. When the protons reach close to the end of their range, they deposit all their remaining energy as a sharply localized energy peak, namely the Bragg peak and promptly after that, the energy drops to zero (Smith, 2006). The Bragg peak is the therapeutic-cytotoxic window, which needs to overlap with the cancerous lesion. The depth of the Bragg peak occurrence is directly related to the proton's initial energy. Therefore, the Bragg peak can be precisely positioned over the target tissue (Levin et al., 2005). A high dose of IR can thus be delivered to a deep-seated tumour with minimal damage to the normal tissue around it. It is estimated that the dose deposition before the Bragg peak is approximately 30% of the maximum peak dose, and beyond the Bragg peak, the dose drops to zero, thereby helping to spare the normal tissue. The lower integral doses received by the surrounding normal tissues together with the steeper dose gradient, where beyond the Bragg peak the dose diminishes, are among the important advantages of proton therapy (Leroy et al., 2016). In contrast, in conventional radiotherapy, gamma- or X-rays deposit their peak dose in the tissue below the skin surface and lose energy exponentially along their path through the tissue, thus unavoidably also affecting normal tissue. A comparative treatment planning study from Lomax and colleagues (Lomax et al., 1999) documented that the use of scanned protons could result in a substantial reduction (50%-67%) of the integral dose and in a reduction of the mean and mid-to-low doses to critical structures in relation to standard photon techniques and intensity-modulated photon plans. In addition, protons have been shown to be more efficient in causing biological damage -approximately 10% more damage per unit dose than their photon counterparts (Terasawa et al., 2009). These advantages make proton therapy ideal for deep-seated, hard-to-reach tumours, cancers that are near critical structures, and oncological treatment for children.

However, even though proton therapy is less damaging to normal tissue than gamma- or X-ray based radiotherapies, potential risks and side effects similar to other radiotherapy regimes should still be taken under consideration. Clinical application of proton therapy has still some limitations and disadvantages, which require further investigation and need to be overcome (Leroy et al., 2016) including: (i) the magnitude of the lateral penumbra (Flanz and Bortfeld, 2013), (ii) the relative biological effectiveness (Miller et al., 2013), (iii) range inaccuracies (Flanz and Bortfeld, 2013), and (iv) uncertainties with relation to tumour location and organs at risk (Flanz and Bortfeld, 2013). Further data on the clinical efficacy of proton therapy are of utmost importance to allow for better understanding of the advantages of proton therapy over conventional radiotherapy.

3.2 Mechanisms of biological damage

The deleterious effects of IR implicate a highly complex sequence of events that are augmented by endogenous signalling, and which altogether lead to oxidative damage

of a wide number of biological molecules including DNA, lipids, proteins, and metabolites (Reisz et al., 2014).

After irradiation with gamma-rays, there is a rapid onset of intracellular ROS production, $\cdot\text{OH}$ and ionized water (H_2O^+), and formation of $\text{O}_2^{\cdot-}$ and H_2O_2 (Singh and Singh, 1982). Subsequently, numerous changes in the intracellular stoichiometry of these reactive species take place together with the generation of additional cell-impairing molecules, including the formation of peroxynitrite (ONOO^-) from the rapid reaction of nitric oxide ($\cdot\text{NO}$) with $\text{O}_2^{\cdot-}$ (Darley-Usmar et al., 1992), peroxynitrous acid (ONOOH), nitrogen dioxide ($\text{NO}_2^{\cdot}$), and nitrogen trioxide (N_2O_3), collectively referred to as reactive nitrogen species (RNS). The increased formation and propagation of ROS and RNS are damaging to the cells as they are highly reactive with biomolecules (Reisz et al., 2014). The time window of cellular damage caused by IR ranges from chemical reactions as rapid as 0.01 ps following irradiation, to more widespread and significant cellular events, such as the delay or arrest of mitosis, damage to protein signalling networks, and membrane rupture that can occur within minutes or even hours after irradiation (Singh and Singh, 1982).

Cellular macromolecules are altered directly through interactions with ionizing particles or indirectly via free radical production from water radiolysis (Ward, 1994). Nucleic acid damage in response to IR is a classic example of the harmful biological effects of IR. These include the detrimental alterations of bases and sugars, DNA cross-link formation, single- and double-strand breaks (SSBs/DSBs), and DNA clustering (El-Missiry et al., 2007; Reisz et al., 2014; Thompson, 2012). Hydroxyl radicals formed during water radiolysis are also particularly deleterious to nucleic acids. Depending on the extent of cellular damage after IR exposure and the capabilities of the insulted cells, the biological consequences may include cell-cycle arrest, initiation of the DNA repair response, or ultimately the induction of apoptosis or necrosis (Han and Yu, 2009).

Proton therapy undoubtedly has reduced the radiation doses to normal tissues and organs in relation to conventional radiotherapy. The ultimate goal in radiation oncology is to improve survival and quality of life, with a balance between tumour eradication and minimization of the (late) effects of radiation, such as the development of secondary malignancies. Even relatively low doses of radiation, once considered to be 'safe', may lead to increased probability of secondary cancer development (Leroy et al., 2016). Nevertheless, several cancers may be radio-resistant, requiring high radiation doses, which may put the patient at risk for enhanced side and late effects. In addition, targeted proton treatment of tumours may have the risk of leaving some infiltrative margins partially treated and neighbouring diseased islets untreated. Despite proton therapy becoming an increasingly popular modality, no one has yet explored the interaction of protons with PSs for induction of ROS to synergistically eradicate tumours. PDT and/or PCI could be powerful treatment modalities, but they are limited by restricted light penetration into tissue. Replacing light with protons to activate PSs is the key novelty that we have pursued in this project. This fusion of PDT and PCI with proton therapy could make protons induce a PDT or a PCI effect both to primary and disseminated disease. Furthermore, this hybrid therapy could be used

postoperatively, as a non-invasive treatment approach to eradicate leftovers of the disease in the resection cavity, which could otherwise lead to recurrence.

Principles of “Protodynamic Therapy”

The background of Protodynamic Therapy (PrDT) lies in the radical hybridization of the PDT principles with those of proton radiotherapy. As mentioned earlier, PDT and PCI are very efficient treatments in eradicating cancer by ROS production but their limiting factor is light accessibility. On the other hand, proton therapy can be targeted through the Bragg peak, but could be enhanced by a secondary mode of action and also by the potential to target occult disseminated disease.

To overcome the limitations of current anticancer, photomedical modalities like PDT and PCI, we sought to combine the therapeutic action of photomedical therapies with the deep penetration of proton therapy. This has led to the conception of a novel bimodal therapy, which we have named Protodynamic therapy (PrDT), and which could also be extended to Proton-Chemical Internalization (ProCI) without the need for an external light source, but driven by accelerated protons. Our hypothesis was based on the assumption that electron rich PSs could be excited by protons, through Coulomb interactions, concurrently to proton therapy giving rise to cytotoxic ROS and eradicating cells with preferentially accumulated PS as in the case of malignant tissues. This tendency of many PSs' to preferentially localize in tumours has been mostly attributed to: (i) the increased vascular permeability to macromolecules of the tumour neovasculature (Yuan et al., 1993), which is important in mediating the uptake of intravenously administered drugs in solid tumours, a phenomenon known as the enhanced permeation and retention (EPR) effect (Wong et al., 2015); (ii) PS binding to LDL in malignant cells overexpressing LDL receptor (Jori and Reddi, 1993); (iii) poorly developed tumorigenic lymphatic drainage in tumours (Roberts and Hasan, 1992), which may increase retention of bound PSs; (iv) infiltrating tumour-associated macrophages (Hamblin and Newman, 1994), which have been shown to accumulate up to 13 times the amount of some PSs compared to cancer cells (Korbelik and Krosli, 1995) and has been attributed either to the phagocytosis of PS aggregates (Korbelik et al., 1991) or the preferential uptake of lipoproteins altered by porphyrin-binding by macrophages (Hamblin and Newman, 1994); (v) low-pH conditions, typically found in tumours, may cause trapping of some of the anionic PSs, which are otherwise ionized at physiological pH (Pottier and Kennedy, 1990); and (vi) the articular tissue's tendency to accumulate lipid droplets (Freitas, 1990).

Brain tumours are an optimal model for PrDT as they are deep seated, often of diffusive and infiltrative nature like for example GBM, which we have chosen as a model for the current project. In this context, using protons can be an advantageous treatment, since they can reach deep into tissue and deposit a large amount of energy before they stop. The added modality of protodynamic therapy, i.e., protons transferring enough energy to PSs to activate them into producing enough singlet oxygen, in addition to their radiotherapeutic action, would contribute to a major breakthrough towards eliminating practically incurable tumours such as GBM.

Up to the present project, proton excitation of photoactive substances had not been used for therapeutic purposes. It had however been shown that low-energy protons (0.5-2 MeV) could excite rare earth complexes (Hollerman et al., 2011; Hollerman et al., 1992; Toh et al., 2009), as well as organic compounds, e.g., anthracene (Birks, 1950, 1951; Birks and Black, 1951). Anthracene can be considered as an interesting model since it is an organic, aromatic compound with a conjugated electronic configuration like PSs. Rare earth compounds also possess atomic electron-rich configurations and they are, therefore, also excited by visible light. Excitation of both the rare earth complexes and anthracene by protons was accompanied by fluorescence emission back to their ground states. Even though all the above mentioned cases were studied in solid form and with the sole purpose of finding an ideal radiodosimetric system, they were construed as a reasonable evidence in support of our hypothesis: Accelerated protons can activate PSs in solutions, gels or biological systems to generate sufficient cytotoxic ROS through interactions with ambient molecular oxygen to enhance the elimination of cancerous cells. The present PhD thesis is a part of an exploratory project granted by the Norwegian Health Southeast in 2016, "Protonic".

Chapter 4

Experimental Conditions

4.1 Cell lines

Cell lines constitute an essential research tool that has revolutionized scientific research and has become valuable in providing a minimal human model to help address fundamental biological questions. Immortalized (continuous) cell lines have the ability to divide indefinitely, unlike primary cells, and they offer a great number of advantages. They are easy to use and manipulate, they are cost effective, and they provide an unlimited supply of biological material. They constitute good model systems, with characterized genetic backgrounds, that offer reproducible results without the requirement for ethical approvals, which are now essential for work with animal and/or human material (Kaur and Dufour, 2012).

Even though cell work has numerous advantages, there are also some limitations. Accumulation of mutations through multiple rounds of in vitro culture can lead to genotypic changes of the cell lines with marked differences from their ancestors. In addition, using 2D-monolayer cell cultures is an oversimplified representation of the biological processes under investigation. In the human body, cells are part of more complex systems, taking part in 3D cell-cell interactions and being surrounded by a microenvironment of different cell types, nutrients, and substrates.

The human GBM cell lines, T98G and U87, and the human adenocarcinoma cell line MCF7 were employed to investigate the photocytotoxic effects of cercosporin on human cancer cell lines (**paper I**). As mentioned earlier, the current standard therapy of GBM consists of surgery, followed by radiotherapy and chemotherapy. However, these therapies offer limited overall patient survival: The median survival for GBM, following surgical resection alone, is approximately 3 months (Walker et al., 1978). The combination of surgery with post-operative radiotherapy increases median survival to approximately 8 months (Walker et al., 1978). Additional chemotherapy with temozolomide extends survival for a further period of 6 months, offering a significant survival benefit (Stupp et al., 2005). Population-based studies report that overall survival of patients, newly diagnosed for GBM, is 17-30% at 1 year and only 3-5% at 2 years, despite access to state-of-the-art modalities of therapy (Adamson et al., 2009), showing a high unmet medical need.

T98G cells are commonly used as an experimental model for this type of cancer and are a polyploid derivative of the T98 cell line, which was acquired from GBM tissue of a 61-year-old adult male (Kiseleva et al., 2016). U87MG is a commonly studied grade IV glioma cell line (Clark et al., 2010) and was obtained from a 44-year-old female patient. Both cell lines are characteristic GBM models and have been extensively used in GBM research, therefore, they came as a natural choice. The third cell line in the paper, MCF7, was used to represent breast adenocarcinoma and was initially isolated from a 69-year-old Caucasian woman. MCF7 cells are estrogen- and progesterone receptor (ER, PR) positive and they have been extensively used as a model of hormone therapy responsive breast cancers, in breast cancer research.

To study whether PSs could be activated by protons and cause cytotoxic death in human GBM (**paper II**), the T98G, M059K, and U87 GBM cell lines were used. Both T98G and U87MG cells have been shown to exhibit high intrinsic resistance against radiation-induced apoptosis (Rübel et al., 2006). M059K cells were firstly isolated from a 33-year-old male patient with untreated GBM (Allalunis-Turner et al., 1993). M059K cells are also known to be irradiation resistant due to their functional DNA-dependent protein kinase (DNA-PK), which plays an important part in the repair of DNA double-strand breaks (Daido et al., 2005). Therefore, all three cell lines constitute great models of radio-resistant GBM and were thus chosen for our protondynamic therapy experiments. In addition, M059K have a tumorigenic potential making them excellent candidates for establishing GBM mouse models for the future in vivo validation of protondynamic therapy.

To explore tumour-specific variations that affect the 5-ALA-PDT outcome and could serve either as predictive biomarkers or even cell-type-specific treatment options, we used a panel of five different cell lines (**paper III**): T98G and U87 cells as models for GBM, and MCF7, MDA-MB-231, and T47D as models of breast adenocarcinoma. T47D, like MCF7, are estrogen receptor positive cells that were once isolated from a 54-year-old female patient. MDA-MB-231 are triple-negative breast cancer (TNBC) cells, meaning that they are lacking expression of ER, PR, and human epidermal growth factor receptor 2 (HER2) (Palasuberniam et al., 2015). TNBC patients do not respond to chemotherapy, while they experience substantial normal tissue toxicity, which makes the development of an effective treatment against TNBC a medical need. MDA-MB-231 cells were first isolated from a 51-year-old Caucasian female patient.

For the cercosporin PDT studies in 3D-cultures, to simulate an environment closer to that of tumours (**paper IV**), we used again two GBM cell lines (T98G, U87) and one adenocarcinoma cell line (T47D). These three cell lines were found to form good, reproducible 3D spheroid models in culture.

4.2 PDT

The PS cercosporin exhibits an absorption band in the range 420-540 nm with a tail extending up to 600 nm and a broad fluorescence with a peak at 620 nm when dissolved in acetone. Although cercosporin absorbs light in the blue spectrum where light cannot propagate deep into tissue (1-2 mm) (Agostinis et al., 2011; Macrobert, 2005), it is an excellent singlet oxygen producer, and therefore could be an ideal PS for applications not requiring external irradiation or for PDT treatment of superficial malignancies. In our work, the cells were seeded in 96-well plates and incubated overnight in complete media at 37°C, in a 5% CO₂ humidified atmosphere. Cells were incubated with various concentrations of the PSs at various time points. Prior to irradiation, the cells were washed twice with PBS with calcium and magnesium, and complete media were added to the cells. Illumination of cells was carried out through the underside of the plates, by means of a Lumisource® lamp (PCI Biotech AS, Oslo, Norway) equipped with four OSRAM L 18/67 fluorescent tubes, which emitted broad blue light band with a peak at approximately 450 nm and an irradiance of 4.7 mW cm⁻². The lamp was turned on 15 min prior to treatment to assure stable and uniform

irradiance. For the 3D spheroid experiments, a yellow lamp was used in addition, since yellow light penetrates deeper into tissue (in our case into spheroids) than blue light (Agostinis et al., 2011; Macrobert, 2005). Yellow light irradiations were performed by a custom-made lamp equipped with an array of 1 W LEDs and an irradiance of 0.9 mW/cm². The spectrum of the yellow lamp peaked at 590 nm. Varying irradiation times were used (0-120 s), resulting in different light doses. In all experiments, a light positive control without the PS and a dark control, with and without the PS, were included.

PpIX displays five different absorption peaks with the largest one at approximately 405 nm (405 nm - Soret band), and four smaller peaks at about 510, 543, 583, and 633 nm (red Q-band). In addition, it exhibits two fluorescent peaks, with the biggest at approximately 635 nm and a smaller one at 705 nm. In these experiments, the cells were irradiated by means of an in-house built lamp equipped with LEDs at 405 nm, while the irradiance at the cell level was 5.6 mW/cm². All cell groups treated with PSs were kept at dark ambient conditions until either illuminated or assayed for viability.

4.3 Cytotoxicity assays

Cellular toxicity was assessed using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. The MTT assay is a colorimetric assay that measures the redox activity of the cells as an indicator of cell viability, proliferation, and/or cytotoxicity. It is based on the conversion of a yellow tetrazolium salt (MTT) to purple formazan crystals by metabolically active cells (Maehara et al., 1987; Vistica et al., 1991) and that enzymatic reduction is executed by NAD(P)H-dependent oxidoreductase enzymes, which reduce MTT to formazan (Berridge and Tan, 1993). Subsequently, the formazan crystals are dissolved using a solubilizing solution, such as DMSO in our case, and thereafter, their endpoint absorbance at 561 nm is measured using the Tecan spark M10 plate reader (Tecan Group Ltd., Männedorf, Switzerland). In all cases, blank subtraction was performed using wells without cells. The association of the MTT with either cell death or proliferation is dependent on the experimental conditions.

4.4 Flow cytometry

Flow cytometry is a broadly used method for quantitatively analyzing cell characteristics, such as the expression of surface and intracellular molecules, differentiation between different cell types in heterogeneous populations as well as analysis of cell size and volume. It can be used for the simultaneous analysis of a variety of parameters on both live and fixed cells, with the possibility to utilize several fluorescent probes in the same run.

Flow cytometry is primarily used to measure fluorescence intensity produced by fluorescent-labeled antibodies bound to their target or by fluorescent substances including PSs (**paper I-IV**), which give a characteristic emission when excited by a laser of the appropriate wavelength. In publications **paper I-IV**, we used live cells, which were incubated with the PSs of choice, and thereafter were analyzed as a

single-cell suspension in the flow-cytometer at various time points versus cells incubated with media only.

Efficient intracellular accumulation is an important index for a PS, since it can be translated to lower amounts of the PS required for PDT activation and subsequently lower chemical (dark) toxicity. It can also partly explain why different cells are either more sensitive or resistant to PDT depending on their ability to internalize PSs. Nevertheless, flow cytometry cannot in principle provide information on the intracellular localization of the PSs. Therefore, in our studies, we complemented the flow cytometry, quantitative studies with qualitative live-cell imaging, to also visualize the subcellular localization of the PSs investigated.

4.5 Microscopy

In the work presented in this thesis, we have used different types of microscopy including confocal and epifluorescence microscopy in live and fixed cells. Sample fixation terminates all enzymatic and metabolic activities, in principle stopping post-fixation changes, while at the same time allows the opportunity to permeabilize cellular membranes for labelled antibodies to engage their intracellular targets. In that context, fixation makes it possible to examine multiple specimens at the same time. Nevertheless, erroneous fixation (under-fixation or over-fixation) could lead to: (i) loss of morphology, (ii) loss of signal, (iii) artifacts, (iv) increased unspecific background signals ('noise'), and (v) impairment of antigenicity (Fischer et al., 2008). During our work (**paper IV**), spheroids were fixed with 2% formaldehyde, "sandwiched" between a microscope slide and a cover slip and visualized by confocal microscopy.

By contrast, live-cell imaging is a broadly used method in biology. The samples are not exposed to fixatives and detergents, keeping the cells intact/functional, and therefore offering the ability of imaging dynamic processes. In our work (**paper I, III, IV**), cells were examined with a Zeiss LSM 880 Airyscan microscope (Carl Zeiss MicroImaging GmbH, Jena, Germany) equipped with a temperature, CO₂, and humidity controller incubator for live-cell imaging. One major limitation that we should take into consideration when executing live-cell imaging is the phototoxicity, which frequently occurs, especially when working with PSs, as extensive illumination can damage living cells and cause cell death. Therefore, limiting phototoxicity is a necessity for obtaining reproducible quantitative data on biological procedures; this can be achieved, e.g., by reducing the illumination at the focal plane and have appropriate controls for the phototoxicity effects (Icha et al., 2017). Another limitation that has to be taken into account, is the photobleaching of the fluorescent dyes/PSs used for microscopy. Photobleaching is the phenomenon of a gradual, and in some cases rapid, loss of fluorescence of the dye/PS upon illumination due to the chemical modification of the molecule, making fluorescence imaging difficult and in some cases even impossible.

For qualitative evaluation of subcellular localization of the PSs (**paper I, III, IV**), we employed confocal microscopy. With confocal microscopy, we can get high resolution images in the x-, y-, and z- axes based on the axiom of signal exclusion from outside

the focal plane. Confocal microscopy provides a powerful tool for studying subcellular localization, as long as measures have been taken to avoid artifacts such as cross talk between the various channels and signal saturation.

4.6 Metabolic studies

Metabolic studies were conducted using a Seahorse XFe96 metabolic analyzer (Agilent, Santa Clara, CA, USA), a one-of-a-kind instrument that allows the performance of metabolic studies, such as measurement of the oxygen consumption rate (OCR, related to cellular respiration) and the extracellular acidification rate (ECAR, related to glycolysis) in intact cells. In the present work (**paper I, III, IV**), the Seahorse XF Cell Mito Stress assay was employed to assess the mitochondrial respiratory and glycolytic functions of GBM and breast adenocarcinoma cells. In the so-called “mito-stress” assay, which we implemented, four chemical compounds are added sequentially in real time: Oligomycin to induce the inhibition of F1F0 ATPase and consequently of ATP production from respiration, carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP) to uncouple the electron transport chain from oxidative phosphorylation, and finally, antimycin A plus rotenone (or alternatively myxothiazol), to completely inhibit the mitochondrial electron transport.

Following the initial basal measurements, oligomycin is injected into the samples, causing a reduction of the OCR. This decrease is associated with the oxygen consumption for ATP production. In the second injection following oligomycin, FCCP uncouples, as delineated above, electron transport from oxidative phosphorylation. Consequently, the cell increases its effort to produce some ATP from respiration, by enhancing the electron transport, and hence the OCR at complex IV reaches its maximum capacity, which can be used to calculate the spare respiratory capacity (difference between maximal respiration and basal respiration). Next, myxothiazol, or a cocktail of antimycin A plus rotenone is added to completely inhibit the mitochondrial electron transport between complex III and IV. As a consequence, any residual respiration ceases being related to oxygen consumption and is driven by processes non-related to the electron transport chain (Divakaruni et al., 2014).

With respect to the measurement of ECAR concurrently to OCR, after the injection of oligomycin, there is usually a concomitant increase in ECAR, as the cells have to compensate for the loss of respiratory ATP production, unveiling the cellular maximum glycolytic capacity.

Even though Seahorse has notable advantages, such as easy-to-use, high resolution, and high-throughput capability, it also has its limitations. The main Seahorse limitation is associated with the selection of the appropriate normalization method to account for cell number, cell size etc. In our studies we have chosen to use a seeding-to-experiment time considerably shorter than the cell doubling time, and report our results per cell line, comparing cell lines irrespective of cell size.

4.7 Signal normalization and statistics

The data from viability and protein expression assays were normalized to a non-treated control, which was nominated as 100%. All other treatments were normalized relative to this internal control. For flow cytometry, the data were normalized to the PS-treated sample with the highest fluorescence.

The statistical analysis amongst parallels for MTT was performed using Microsoft Office Excel 2007. To further assess significance between data groups, a Student's t-test was used, and variations were considered significant when $p \leq 0.05$. Synergism was found when the result of the combination treatment was significantly different ($p \leq 0.05$) from the additive effect of the individual treatments, i.e., $\text{Survival (PS+ treatment)} < \text{Survival (PS)} \times \text{Survival (treatment)}$.

Summary of Publications

Paper I

Cytotoxic and Photocytotoxic Effects of Cercosporin on Human Tumour Cell Lines.

Authors

Mastrangelopoulou M., Grigalavicius M., Berg K., Ménard M., Theodossiou T.A.

Main findings

Cercosporin is a naturally occurring perylenequinone produced by fungi of the genus *Cercospora*. Although other perylenequinones, such as hypericin and hypocrellins, have been extensively studied as photosensitizers in cancer PDT, most studies on cercosporin are focused on its role in phytopathology. In this publication, we set out to determine the photocytotoxic effects of cercosporin in two GBM (T98G and U87) and one breast adenocarcinoma (MCF7) human cell lines.

Cercosporin, dissolved in acetone, exhibited an absorption band in the range 420-540 nm and a broad fluorescence with peak at 620nm. Upon excitation at 532nm, cercosporin was found to be a potent singlet oxygen producer compared to well-known photosensitizer standards with high singlet oxygen quantum yield. The intracellular accumulation of cercosporin was three times higher in T98G cells than in MCF7 or U87 cells, with a loading plateau onset after 2 h of incubation for the 2 and 4 μ M concentrations. Confocal analysis showed that cercosporin accumulated both in mitochondria and endoplasmic reticulum within the cells, which constitute prominent PDT targets. Cercosporin showed diverse survival trends following PDT treatment, which was dependent on the irradiation time, cercosporin concentration, and cell line studied. There was no evident dark toxicity 24 h following 4h incubation for all concentrations tested (0.8-8 μ M) and across all the cell lines investigated. We further looked into the metabolic profiles of intact cells following cercosporin PDT, with the use of a "Seahorse" XFe96 metabolic analyzer. T98G and MCF7 seemed to exhibit the highest respiration with similar basal OCR. By contrast, U87 cells exhibited 2-3 times lower OCR under media-only conditions as compared to T98G and MCF7 cells. One hour after irradiation, there was a complete collapse of the respiratory activity in all cells treated with cercosporin (PDT groups). Regarding the cells' glycolytic activities, T98G exhibited a much higher glycolytic activity than the other two cell lines. Light irradiation caused again a glycolytic collapse in all the cells treated with cercosporin (PDT groups), which could be mostly due to the photo-inactivation of intracellular LDH. Copper is a transition metal essential for the function of key enzymes and protection against oxidative stress, while it is also involved in primary metabolism such as mitochondrial oxidative phosphorylation (Horn and Barrientos, 2008). In excess quantities copper can, however, lead to the generation of hydroxyl radicals by Fenton reactions ultimately leading to cell damage (Pham et al., 2013). Our data showed a synergistic dark toxicity by co-incubation of cercosporin with copper in MCF7 and T98G cells as opposed to a merely additive effect in U87 cells.

According to the results of our study, the PS cercosporin exhibited enhanced photocytotoxicity against the GBM and breast cancer cell lines studied. In spite of its

short wavelength of activation, incompatible with clinical PDT, cercosporin could prove to be suitable for superficial PDT treatments or applications, which do not require external light sources, e.g., intracellular chemiluminescence PDT.

Published

Photochem Photobiol 95 (2019) 387-396 (Mastrangelopoulou et al., 2019)

Paper II

Proton-dynamic therapy following photosensitiser activation by accelerated protons demonstrated through fluorescence and singlet oxygen production

Authors

Grigalavicius M., **Mastrangelopoulou M.**, Berg K., Arous D., Ménard M., Raabe-Henriksen T., Brondz E., Siem S., Görgen A., Edin N. F. J., Malinen E. & Theodossiou T. A.

Main findings

PDT is a clinically approved, minimally invasive therapeutic modality that offers targeted cancer treatment through the synergy of three components: (i) the PS, (ii) light of the appropriate wavelength to excite the PS, and (iii) oxygen, leading to the generation of ROS (Agostinis et al., 2011). On the other hand, proton therapy (Levin et al., 2005) is a targeted, external-beam tumour radiotherapy using accelerated protons that can penetrate deep into tissue and be tuned to deliver a large dose of energy specifically to the tumour site. This can be achieved, as tuning the initial kinetic energy of protons can determine the depth in which they deposit this high amount of radiant energy within a narrow space, the Bragg peak. As the protons lose their energy by electromagnetic interactions while traversing the tissue, once they reach low energies, they increase their cross section of interaction with the surrounding electrons, and consequently they release their remaining energy in a big burst within the Bragg peak. This can overcome the main limitation of PDT, i.e., the fact that the penetration depth of light is limited only to a few millimeters in tissue (Macrobert, 2005). The combination of the photodynamic and proton therapies by applying proton induced PS activation eliminates the need for an external light source, while it can extend the treatment activity deep within the tissue, overcoming the limitations of PDT and potentially leading to a bimodal killing of tumour cells through: (i) ionizing irradiation and (ii) ROS and primarily singlet oxygen production.

In this study, we sought to deliver a proof-of-concept for the anti-cancer efficacy of the activation of PSs by protons in synergy with the ionizing radiation of proton therapy. Proton-induced fluorescence spectra were studied in PS solutions, with their emission spectra matching their light-induced fluorescence counterparts. Porphyrin- and chlorin-type PSs exhibited the strongest fluorescence as opposed to quinone-type compounds that exhibited weaker fluorescence. PSs excited in dry gelatin led to enhanced phosphorescence emission as compared to solutions, suggesting an efficient PSs triplet state population. We performed singlet oxygen measurements optically around 1270 nm by registering the phosphorescence of singlet oxygen in acetone d_6 . In addition, we measured singlet oxygen production indirectly by studying the PpIX singlet-oxygen-associated photoproduct (photo-PpIX) formation at ~670nm. Both methods showed singlet oxygen generation by proton excitation of PSs. The singlet oxygen-specific scavenger 1,4-diazabicyclo[2.2.2]octane (DABCO) offered a nearly complete abolishment of the fluorescence bleaching after photon irradiation.

On the other hand, there was only a partial recovery of the fluorescence degradation after proton irradiation, suggesting structural damage to the PSs most probably through ionization. Finally, we examined the efficacy of the proton-dynamic technology on 2D GBM cell cultures. M059K and T98G cells showed a statistically significant enhancement of cytotoxicity after addition of the PS cercosporin on top of proton irradiation, while U87 cells exhibited no additional effect from PS activation at any proton irradiation dose examined.

In the current work, we demonstrated the excitation of PSs by accelerated protons in solutions and gels, and the generation of fluorescence and singlet oxygen. We also exhibited for the first time the efficacy of the proton-dynamic effects on GBM cell lines. Our results pave the way for a new hybrid therapy, namely 'proton-dynamic therapy', as ROS generated by this interaction could demolish malignant cells in synergy with ionizing irradiation, leading to efficient tumour eradication.

Published

Nature Communications 10 (2019) (Grigalavicius et al., 2019)

Paper III

Predictive biomarkers for 5-ALA-PDT can lead to personalized treatments and overcome tumour-specific resistances.

Authors

Mastrangelopoulou M., Grigalavicius M., Raabe-Henriksen T., Skarpen E., Juzenas P., Peng Q., Berg K., Theodossiou T.A.

Main findings

5-ALA (see Chapter 2.5) is a natural precursor of heme in the heme biosynthetic pathway and it is metabolically converted to PpIX, a photosensitizing porphyrin with a relatively high fluorescence yield [$\Phi_f = 0.085$ when dissolved in DMSO (Gotardo et al., 2017)] and potent photosensitizing activity (Kennedy and Pottier, 1992; Kennedy et al., 1990; Wachowska et al., 2011). Due to its great tumour specificity, reduced nonspecific and cutaneous photosensitivity, and inherent metabolism into non-fluorescent heme, 5-ALA has shown great promise as a prodrug in the clinic (Herceg et al., 2017; Yang et al., 2015). Due to the efficient fluorescence quantum yield of PpIX, 5-ALA has received a lot of attention not only as a PDT agent but also as a fluorescent agent that can be used in PDD (Krammer and Plaetzer, 2008). 5-ALA-PDD is already in use in the clinic for the diagnosis of various tumours, such as prostate, bladder, and brain malignancies (Fukuhara et al., 2011; Inoue et al., 2012; Stummer et al., 2000; Stummer et al., 1998), and it is already being used for the guided resection of brain and bladder tumours with promising clinical outcomes (Jichlinski and Leisinger, 2001; Tetard et al., 2014; Yang et al., 2015).

In the current work, we present an in vitro study of 5-ALA-PDT on two human GBM cell lines (T98G and U87) and three breast adenocarcinoma cell lines (MCF7, MDA-MB-231, and T47D) for comparison. After cell survival studies at different 5-ALA concentrations and irradiation times, we showed that the GBM cells were more resistant to 5-ALA-PDT than breast cancer cells, and we established the LD₃₀ for the various cell lines. Next, we set to examine 5-ALA uptake, also in conjunction with Ko143, an inhibitor of the PpIX transporter ABCG2. The cells showed large heterogeneity in PpIX fluorescence, and treatment with Ko143 significantly increased PpIX fluorescence in all cell lines, except T47D.

We examined the role of different anti-oxidant proteins on 5-ALA-PDT using western blotting and showed that U87 exhibited significantly higher expression of SOD2, the mitochondrial SOD, and the antioxidant enzyme GPX4. These two findings demonstrate the importance of cell mitochondria as a protective mechanism for U87, despite displaying the highest glycolysis/respiration ratio, implying a predominantly Warburg-type cellular metabolism. With respect to antioxidant defenses, we showed that T98G exhibited high levels of SOD1 and GSTP1 and the highest cellular glutathione (GSH) expression as compared to the rest of the cell lines. To further evaluate the mechanisms of 5-ALA-PDT action on the various cell lines, we evaluated the impact on GSH as a defense mechanism against oxidative stress induced by PDT by pretreating the cells with buthionine sulfoximine (BSO-an inhibitor of GSH

synthesis) prior to 5-ALA PDT. We found that BSO-induced depletion of GSH significantly enhanced the 5-ALA-PDT killing effect in T98G and MDA-MB-231 cells. T98G exhibited an adaptive expression of heme oxygenase (HO-1), the enzyme responsible for heme degradation, which increased with 5-ALA administration and decreased following 5-ALA-PDT. Furthermore, inhibition of HO-1 by OB24 caused effective 5-ALA-PDT killing only in T98G cells. The rest of the cell lines that demonstrated moderate (U87) or minimal HO-1 levels (MCF7, MDA-MB-231, and T47D), did not exhibit augmented PDT cytotoxicity following OB24 pretreatment. Inhibition of FECH, the enzyme converting PpIX to heme, sensitized T98G, MCF7, and T47D cells to 5-ALA-PDT, and the degree of photosensitization was analogous to their corresponding FECH levels. However, the cells that demonstrated lower FECH levels (U87 and MDA-MB-231) were found to be less responsive to 5-ALA-PDT following FECH inhibition. With respect to the ABCG2 transporter, we found it to be expressed in all cell lines with highest expression in T98G cells. To get a deeper understanding of the transporter's influence on 5-ALA-PDT, we inhibited ABCG2 with Ko143 and performed cytotoxic assays and microscopy analysis. T98G, U87, and to a lesser extent MDA-MB-231 cells were found to be quite susceptible to 5-ALA-PDT following ABCG2-inhibition, despite the fact that the two latter cell lines exhibited low expression levels of ABCG2. On the contrary, MCF7 cells displayed a moderate PDT response, despite having higher levels of ABCG2 than U87 and MDA-MB-231. There was no notable effect in the 5-ALA-PDT response following ABCG2 inhibition on T47D cells, despite expressing ABCG2 at a comparable level to MDA-MB-231. At the microscopy level, treatment with Ko143 significantly enhanced the mitochondrial accumulation of PpIX in all cells with the exception of T47D where there was no marked difference. Finally, we studied the metabolic profiles of the cells before and after 5-ALA-PDT treatment, and we showed that 5-ALA-PDT caused both a respiratory and glycolytic collapse in all the cell lines examined.

Our study demonstrates the response heterogeneity to 5-ALA-PDT across the cell lines and highlights novel biomarkers that may ultimately provide patient-specific prognosis for treatment outcomes. Furthermore, it may pave the way for personalized treatments based on 5-ALA-PDT resulting in profoundly higher treatment efficacy, and potentially overcoming resistances to this treatment modality.

Published

Cancer Reports (2020) (Mastrangelopoulou et al., 2020)

Paper IV

Photodynamic efficacy of cercosporin in 3D tumour cell cultures

Authors

Grigalavicius M., Mastrangelopoulou M., Arous D., Juzeniene A., Ménard M., Skarpen E., Berg K., Theodossiou T.A.

Main findings

Cercosporin (see Chapter 2.5) is a naturally-derived perylenequinone PS that is produced by the *Cercospora* family of plant pathogens and that has been extensively studied for its photodynamic properties within the realms of phytopathology, as it is a potent phototoxin for plants and crops (Daub et al., 2013). Previously, we have shown that cercosporin is a potent PDT agent against GBM and breast cancer cell lines in 2D cultures causing enhanced phototoxicity and a significant metabolic collapse in both the respiratory and glycolytic activity [see above, paper I, (Mastrangelopoulou et al., 2019)]. Cercosporin also exhibited enhanced cytotoxicity against two GBM cancer cell lines upon proton activation [see above, paper II, (Grigalavicius et al., 2019)], rendering it a valuable candidate for proton dynamic therapy.

In the current work, we were interested in studying the photodynamic action of cercosporin on human cancer multicellular spheroids of three mammalian cancer cell lines, namely T47D (human breast adenocarcinoma), and U87 and T98G (GBM). The 3D culture model has been established as a system of intermediate complexity between solid tumours in vivo and cell monolayers in 2D cultures (Sutherland, 1981). Spheroids provide a good model system for studying the 3D cell-cell interactions, structural barriers to nutrients, substrates, and oxygen. Flow cytometry studies revealed that U87 spheroids displayed double the cercosporin uptake as compared to their T47D and T98G counterparts. Confocal fluorescence microscopy studies showed that cercosporin was uniformly and efficiently taken up by the spheroids in all cell lines studied. Cytotoxicity studies demonstrated that U87 spheroids exhibited the least vulnerability to cercosporin-PDT, despite their higher cercosporin uptake. Furthermore, the PDT efficacy of cercosporin was also dependent on the spheroid size. Despite the fact that 300 μm diameter spheroids were shrinking and becoming necrotic after PDT, bigger spheroids (>500 μm) started to re-grow after PDT, at least the ones irradiated with blue light and exhibited higher viability. We also examined the PDT efficacy on larger spheroids reaching up to 1 mm in diameter and showed that there was a severe growth inhibition after yellow-light irradiation (~590 nm). Finally, we examined the metabolic profiles of intact spheroids from the cell lines under investigation and found that they exhibited comparable OCR. Nevertheless, after cercosporin-PDT, there was a respiratory collapse in all three spheroid types. The glycolytic activity of T98G and T47D spheroids was also depleted following cercosporin-PDT, but U87 spheroids retained approximately half of their glycolytic rate post-PDT.

In paper IV, we demonstrated that cercosporin is a very potent photosensitizer not only in 2D cultures but also in a 3D model, which is more similar to solid tumours. We

showed the limitations of shorter wavelength (blue) light in penetrating tissue as smaller spheroids could activate the PS relatively in the entire volume of the spheroid, but not in larger spheroids. Yellow light, however, despite being much less absorbed by cercosporin, may cause far more detrimental effects in cercosporin treated-spheroids, even at a lower effective dose than blue light.

Published

Photochem Photobiol 96 (2020) 699-707 (Grigalavicius et al., 2020)

Discussion

Chapter 5

Protodynamic Therapy – A combination of Proton Therapy and conventional PDT

5.1 Protodynamic Therapy

Brain cancers like GBM are essentially incurable due to their deep-lying, highly infiltrative and metastatic nature, stealthily drifting between different parts of the brain by forming little islets of disease around the primary lesion. Approximately 28,000 new cases of malignant glioma are diagnosed every year in the EU and USA (Source: US National Cancer Registry) with the relative survival rate of adults diagnosed with GBM being less than 30% within one year of diagnosis (Source: Central Brain Tumour Registry of the United States), and only 3% of patients exceeding five years survival after initial diagnosis, indicating a highly unmet medical need. Accelerated protons can reach deep into tissue where they deposit substantial amounts of energy at the Bragg peak, but there is a risk of not treating islets of diffuse disease with the appropriate dose. Furthermore, brain cancers are frequently radio-resistant, requiring high radiation doses causing enhanced side effects to the patient. Both PDT and/or PCI could be very beneficial alongside proton therapy, but their main constraint is the depth of light penetration into tissue. To overcome the limitations of proton therapy and PDT/PCI treatment modality, we propose Protodynamic therapy, a hybrid treatment modality, where we combine PDT and PCI with proton therapy, using protons to induce a PDT or a PCI effect both to primary and disseminated disease.

Our study had a focus on obtaining proof of principle of the excitation and activation of PSs by accelerated protons to generate fluorescence, ROS, and ultimately tumour cell death (paper II). To achieve that, we chose PSs that are either well established in the PDT bibliography or already in clinical use, such as PpIX (already used both in clinical PDT and as a PDD agent to guide tumour resection on malignant tissues during GBM surgery), mTHPC (temoporfin), and TPCS2a (Amphinex®). Given that PpIX is a potent PS and has multiple uses in the treatment of cancer, we sought to investigate mechanisms that could potentially transform 5-ALA-PDT into a more efficient treatment modality both in current clinical settings and in novel applications such as PrDT (paper III). We also wanted to explore new options for PSs that are so far not used in the clinics, yet they are well-known for being efficient singlet oxygen producers such as cercosporin. In that context, in paper I, we studied the photocytotoxic effects of cercosporin in vitro using GBM and breast adenocarcinoma cell lines in 2D cell cultures. In a follow-up study, we investigated the efficacy of cercosporin as a PDT agent in 3D spheroid cultures (paper IV), which represent a model of higher complexity and hence more relevant to tumours than 2D cultures. The purpose of this project was to set up the basis for a new hybrid treatment based on proton therapy, and at the same time, using PSs to eradicate malignant lesions through the synergistic effects of ROS (singlet oxygen) generation and proton ionization.

5.2 PDT in combination with gamma-radiation and X-rays

Radiation therapy (RT) constitutes a crucial therapeutic alternative in clinical oncology with over 50% of all cancer patients receiving RT as a treatment modality during their curative process (Bernier et al., 2004). However, not all cancer cells are adequately responsive to RT, while tumours that were originally responsive may develop secondary resistance during the therapy (Begg et al., 2011). Meanwhile, PDT and RT have different targets of action, and it has been shown that some radiation-resistant tumour cells are sensitive to PDT and vice versa (Xu et al., 2016). In that sense, there have been several attempts to investigate the synergism of PDT and ionizing radiation (X-rays or gamma-rays). Although several in vitro studies have been published on the combination of PDT and RT, there is no clear consensus on the overall benefit of such a combinatorial treatment approach. Some investigators concluded in favour of a synergistic interaction (Boegheim et al., 1987; Kostron et al., 1986), while other authors showed a merely additive interaction (Bellnier and Dougherty, 1986; Ben-Hur et al., 1988; Schnitzhofer and Krammer, 1996). Some other studies concluded that there is a possibility of either a synergistic or an antagonistic effect depending on the experimental conditions (Berg et al., 1995; Ma et al., 1993). This controversy on the efficacy of the combination of radiotherapy and PDT has been attributed to various factors: (i) cell line dependent variation to PDT sensitivity (Prinsze et al., 1992) and (ii) dose and timing variations between the two treatments (Berg et al., 1995; Kavarnos et al., 1994; Ma et al., 1993).

PSs that act as radiosensitizers (RSs) have been studied by a number of groups to gain a greater understanding on the combinatorial effects of PDT and RT, and ultimately induce greater cytotoxicity when treating deep-seated tumours. Luksiene et al. demonstrated that Hematoporphyrin dimethyl ether (HPde) acts as a RS in mice with transplanted tumours. The combination of PDT and RT led to an additional effect, compared to radiation only (Luksiene et al., 1999), not attributed to the excitation of the PS. In a following study, the RS properties of three different porphyrin-type PSs (HPde, Photofrin II (PII), and HPD) were evaluated (Luksiene et al., 2006). It was demonstrated that HPde was the most effective RS, followed by PII and HPD, with the RS effect of these PSs being cell line dependent. In addition, the data indicated that the efficacy of radiosensitization is strongly dependent on chemical homogeneity and accumulation potential of the compound. The low efficacy of PSs to act as RSs under direct excitation has led to the loss of interest for the development of dual PS and RS agents and has triggered the development of alternative approaches to excite the PSs in deep tissue.

5.3 Current advances in Proton Therapy

In theory, proton therapy offers a considerable clinical advantage over conventional photon therapy. This is due to the exceptional depth-dose properties of protons, which can be exploited to offer substantial reduction on the doses deposited on normal tissues proximal and distal to the target volume (Mohan and Grosshans, 2017). Upon entry into tissue, the proton beam energy loss is almost constant with depth and when the protons reach low energies, they deposit all their remaining energy at the Bragg

peak. The therapeutic Bragg peak is what gives proton beams an advantage over X- and gamma-rays, where the highest energy loss is upon entrance of the X- or gamma-photons in the matter, and the energy is lost exponentially as they transverse the tissue. This advantage of accelerated protons may allow increase of the dose at the tumour site, sparing of normal tissues adjacent to the tumour, hence possibly enhancing local control and survival, while simultaneously diminishing toxic side effects and improving life quality (Mohan and Grosshans, 2017). Unfortunately, due to the high cost of establishing and running proton therapy centers, proton therapy is not at present readily available to patients in contrast to photon therapy. Nevertheless, due to the increased construction of proton centers internationally, there are several clinical indications where proton therapy is becoming the therapy of choice. Furthermore, as the cost of the proton therapy systems decreases and clinical interest evolves, there will be a higher amount of patients receiving proton therapy in the future.

Children are particularly susceptible to late adverse effects of radiation such as secondary cancers, cardiac disease, endocrinopathies, and cognitive dysfunction. Therefore, proton therapy has become widely accepted for use in pediatric cancers due to its tissue sparing capacity (Mohan and Grosshans, 2017). In addition to childhood malignancies, proton therapy is increasingly used for intracranial and sino-nasal malignancies. That is due to the close proximity of critical tissues, such as optic structures and brainstem, to the sites of irradiation where high doses are frequently required. Outstanding results have been achieved for patients with skull-based lesions (Ares et al., 2009) and improved survival outcomes for patients with sino-nasal tumours (Patel et al., 2014).

Proton therapy shows great promise also in the treatment of brain tumours. The precise area of irradiation and minimal lateral scattering in tissue, ensuring minimal exposure of the normal tissue surrounding the tumour, makes proton therapy ideal for treating CNS malignancies in order to minimize neurocognitive dysfunction due to radiological damage in healthy tissue (Alan Mitteer et al., 2015) as well as potential dose escalation for radiation resistant tumours such as GBM (Mohan and Grosshans, 2017). Numerous studies have taken place studying the therapeutic outcome of proton therapy with promising results (Noël et al., 2005; Weber et al., 2012; Wenkel et al., 2000). However, the sharp distal fall-off with proton therapy could lead to some infiltrative margins being left partially treated and neighboring islets of disseminated disease untreated, and therefore cause higher rates of disease recurrence.

Our new proposed technology, PrDT, can potentially make proton therapy more targeted and effective, also against infiltrative tumour margins and/or small foci of disseminated disease. The proton beam will electronically excite the PSs and activate them to produce cytotoxic ROS. Consequently, the generated ROS will eliminate not only the surrounding tumour tissue but also the undiagnosed disseminated GBM lesion islets around the primary GBM tumour, where the PS has also accumulated, causing the premature formation of Bragg peaks, specifically on these islets. For this purpose, we need electron rich PSs that exhibit preferential accumulation in tumours, potent singlet oxygen production, and low levels of dark toxicity. The choice of PS is expected to be critical for successful treatment, therefore we selected two PSs for further studies in PrDT: (i) PpIX derived through heme biosynthesis following the

administration of the prodrug 5-ALA, a potent and widely used PS both in PDT and PDD, and (ii) the natural occurring pigment cercosporin, which is a potent singlet oxygen producer, yet due to its short wavelength of activation has not been explored for conventional, light-based PDT clinical use.

5.4 5-ALA as a prodrug and PpIX cytotoxic effects in cancer

5-ALA (see Chapter 2.5) is a natural precursor of heme in the heme biosynthetic pathway that takes place in virtually all cells containing a nucleus (Herceg et al., 2017). 5-ALA-PDT exhibits many advantages over conventional PSs used in PDT and other conventional cancer treatments such as surgery, chemotherapy, and/or radiotherapy (Wachowska et al., 2011). Exogenous administration of 5-ALA causes preferential accumulation of the natural PS PpIX in tumour cells due to differences in the heme biosynthetic pathway between malignant and non-malignant cells (Wachowska et al., 2011). Since 5-ALA is administered as a prodrug and it does not possess PS properties, its administration does not elicit major photosensitivity issues caused by PS accumulation in non-targeted tissues. This makes repeated 5-ALA administrations possible with minimal side effects (Herceg et al., 2017). Furthermore, PpIX produced, following 5-ALA administration, may be used not only for PDT purposes but also for PDD, making 5-ALA a very attractive prodrug for various clinical uses. In paper III, we explored different biomarkers that could be potentially used in the clinic to predict 5-ALA-PDT treatment outcomes but also as therapeutic modulators of the treatment efficacy, thus transforming 5-ALA-PDT into a more personalized and efficient treatment modality both in conventional PDT practice as well as in our newly proposed PrDT technology.

We demonstrated that there was a vast divergence of 5-ALA-PDT-related cytotoxicity, with the GBM cell lines (T98G and U87) being more resistant compared to the breast adenocarcinoma cell lines (MCF7, MDA-MB-231, and T47D). There were also differences in PDT sensitivity within the cell lines in each group with T98G being more resistant than U87, and MCF7 being more resistant than MDA-MB-231 and T47D cells. The higher resistance of MCF7 compared to MDA-MB-231 cells is in agreement with previous hypericin-PDT studies (Theodossiou et al., 2019; Theodossiou et al., 2017).

5-ALA-induced PpIX synthesis, intracellular accumulation, and its subcellular localization are determined by several factors that differ among cell types. The first factor that needs to be taken into account is the PS mechanism and consequently rate of uptake. Another factor is the intracellular fate of the PS and metabolism depending on the differential expressions of various enzymes in different cell lines. One mechanism responsible for divergent responses to PDT could be differential drug uptake and efflux rates or intracellular trafficking (Casas et al., 2011). A major function of ABC transporters is to protect the cells from many toxic insults such as endogenous and exogenous molecules entering the cells. This protective mechanism of ABC transporter-mediated efflux of such substances, which can be metabolic waste products, naturally occurring substances or drugs, is what offers the cancer cells resistance to the toxic effects of numerous chemotherapeutic agents (Doyle et al.,

1998; Robey et al., 2011). ABCG2, a member of this family of transporters, has an exceptionally wide range of substrates and has been associated with drug resistance in breast cancer and leukemia (Doyle et al., 1998; Robey et al., 2011).

The ATP-binding cassette transporter ABCG2 plays a crucial role in regulating the intracellular level of porphyrins in cancer cells. In this sense the cell-to-cell differences in ABCG2 expression and function may affect the efficacy of both PDT and PDD. Nevertheless, blocking the ABCG2-mediated porphyrin efflux from cells can potentially enhance PDT and PDD outcomes making the transporter a great target for future inhibitor development (Ishikawa et al., 2015). Our data show that T98G exhibited the highest PpIX production contradicting their relatively low PDT cytotoxicity results. However, our protein studies show that T98G exhibited the highest ABCG2 expression compared to the rest of the cell lines, which could account for the controversy above. Accordingly, inhibition of the ABCG2 transporter by Ko143 significantly decreased the efflux of PpIX from T98G, resulting in a profound increase in cell death following PDT as compared to cells without Ko143 treatment. U87 and MDA-MB-231 were also quite susceptible to PDT after co-incubation with Ko143, while MCF7 cells exhibited a moderate response. Interestingly, T47D cells showed no difference in PpIX production or viability after PDT following co-incubation with Ko143. This suggests the onset of other defense mechanisms against PDT in these cells.

Subcellular localization of PpIX is critical because ROS that are formed during PDT have a very short half-life; therefore, the mechanisms of action and efficacy of PDT will rely on the intracellular localization of PpIX (Herceg et al., 2017). PpIX fluorescence was observed throughout the cytoplasm of all the cell lines, while in some (T98G, MCF7, and MDA-MB-231) PpIX fluorescence was also present in the plasma membrane. The observation that PpIX fluorescence displayed a diffuse and cell membrane bound pattern before Ko143 treatment suggests possible PpIX efflux transport. Inhibition of the ABCG2 transporter with Ko143 significantly enhanced intracellular PpIX fluorescence in all but T47D cells. Of note, the fluorescence pattern after addition of Ko143 was primarily in the perinuclear area, while it was not noticeable in the plasma membrane. As a consequence of PpIX intracellular accumulation after Ko143 treatment, both the GBM (T98G and U87) and the TNBC cells (MDA-MB-231) became sensitized to 5-ALA-PDT. These results show that enhanced ABCG2 transporter activity renders GBM cell resistance to 5-ALA-PDT by reducing intracellular PpIX levels, which could be reversed by inhibiting ABCG2 activity. This is in agreement with another study on breast cancer cells where they showed that TNBC cells were sensitized to 5-ALA-PDT after ABCG2 transporter inhibition (Palasuberniam et al., 2015).

Overall, our data indicate that the various cell lines had differential ABCG2 expression, and its inhibition by Ko143 caused significant changes in PDT sensitivity and PS intracellular localization in some cell lines (T98G, U87, MCF7, and MDA-MB-231) but had no effect in others (T47D). Therefore, it became evident that in some cell lines PpIX accumulation and ABCG2 inhibition are not enough to account for their PDT resistance.

5.5 GSH and other ROS scavengers

Glutathione (GSH) plays a key role in a wide variety of cellular processes, including cell differentiation, proliferation, and apoptosis. Disruptions in GSH homeostasis may lead to the progression of many human diseases including cancer (Traverso et al., 2013). A balance in GSH levels is of major importance since GSH deficiency leads to an increased vulnerability to oxidative stress, while increased GSH levels enhance the antioxidant capacity and subsequently the resistance to oxidative stress as is the case for many cancer cells (Traverso et al., 2013).

We saw treatment with 5-ALA caused a number of changes in various cellular ROS scavengers, suggesting that they may have a protective function against PDT-induced ROS. Localization of the ROS scavengers is of utmost importance, as the primary ROS produced by the PS is short-lived, and therefore has a limited migration capacity (Moan and Berg, 1991). A ROS scavenger, namely GPX4, has been reported to localize in the cytosol, mitochondria, nucleus, and ER (Brigelius-Flohé and Maiorino, 2013; Liang et al., 2009). We suggested that T47D defensive mechanisms to PDT could be associated with the high membrane-resident GPX4 levels, which despite the low levels of intracellular GSH, might help to catalyze the detoxification of lipid hydroperoxides produced by PpIX induced peroxidation. Nevertheless, when we reduced the intracellular GSH levels by BSO, the most prominent GSH-depleting agent studied in both preclinical and clinical trials (Traverso et al., 2013), it did not affect significantly their PDT response, suggesting yet other potential defence mechanisms.

Another family of enzymes, which are involved in the flux of various compounds, potentially offering resistance to their hosts, are the glutathione S-transferases (GSTs), which play a crucial role in detoxification of xenobiotics and endogenous metabolites by catalysing the conjugations of various compounds with reduced glutathione (Laborde, 2010). GSTP1 has enhanced expression in tumour cells, which are resistant to a wide variety of anticancer drugs (Laborde, 2010). It is critical in expelling xenobiotics, in the management of lipid oxidation as well as S-glutathiolated proteins produced by oxidative stress (Awasthi et al., 2004; Listowsky, 2005). Our results showed that T98G cells displayed the highest GSTP1 levels and together with MDA-MB-231 cells were found to be the most responsive to GSH depletion after 5-ALA-PDT. This is in good agreement with previous results from our group (Theodossiou et al., 2017) where it was shown that MDA-MB-231 cells utilized GSTP1 and GSH as defence mechanisms to detoxify hypericin and alleviate its PDT effects.

One additional type of defence mechanism the cells have to prevent or limit intracellular damage and alleviate the harmful effects of ROS, apart from glutathione peroxidases (GPXs), include antioxidant enzymes such as superoxide dismutases (SODs) (Deavall et al., 2012). The antioxidant enzymes SOD1 and SOD2 are localized in the cytoplasm and mitochondria respectively, and therefore they would be expected to offer protection against 5-ALA-PDT. U87 cells exhibited the highest expression of SOD2 and also GPX4. Indeed, we can see from our work that U87 were quite resistant to 5-ALA-PDT, indicating the major importance of cell mitochondria protection for U87 cells, even though they displayed the lowest OCR rates and highest ECAR/OCR

(glycolysis/respiration) ratio, pointing to a predominantly Warburg-type cellular metabolism. Altogether, our data demonstrate that ROS scavengers may provide partial resistance to 5-ALA-induced cell death but there are possibly other mechanisms that also have to be taken into consideration.

5.6 Enzymes in heme synthesis and degradation

PpIX preferentially accumulates in tumour cells as compared to normal cells, which can be partly explained by differences in the heme biosynthetic pathway between normal and tumour cells. Decreased activity of FECH (Hefti et al., 2011; Ohgari et al., 2005) and limited availability of iron (Stout and Becker, 1990) in tumour cells contribute to enhanced PpIX accumulation. FECH forms the final step in the pathway by catalyzing the conversion of PpIX to heme through insertion of ferrous iron to PpIX (Franken et al., 2011). On the heme degradation pathway, heme oxygenases are metabolic enzymes that catalyze the conversion of heme to biliverdin, carbon monoxide, and iron (Wegiel et al., 2014). Previously, it has been shown that HO-1 can play a protective role against 5-ALA-PDT mediated cytotoxicity so that inhibition of HO-1 activity might be used to increase the efficacy of 5-ALA-PDT (Frank et al., 2007).

In paper III, we show that even though Ko143 inhibition of the ABCG2 transporter had no effect on T47D cells, inhibition of FECH by Noc-18 elicited a profound sensitivity to 5-ALA-PDT. As mentioned earlier, FECH has a key role in the production of heme by catalyzing iron incorporation into PpIX, hence converting it into heme, and at the same time reducing the amount of PpIX available for PDT. T98G, MCF7, and T47D cells exhibited enhanced PDT-induced cytotoxicity following FECH inhibition, which was in good agreement with their reciprocal FECH protein levels.

The lack of a multifaceted defense in T47D cells suggests that their main defense to PDT might be their increased FECH expression, and may provide an explanation as to why T47D cells are the most responsive to 5-ALA-PDT. Furthermore, T47D were found to lack HO-1, a member of the heat shock protein family that relieves oxidative stress in tumour cells (Alaoui-Jamali et al., 2009; Frank et al., 2007). T47D response to 5-ALA-PDT was hence unaffected following inhibition of HO-1 by OB24. On the other hand, T98G demonstrated a higher HO-1 expression, which increased after 5-ALA administration and decreased after PDT. In addition, T98G were the only cell line to show an enhanced 5-ALA-PDT response following inhibition of HO-1.

Overall, we suggest that ABCG2, FECH, and HO-1 could be potentially used as predictive biomarkers for the treatment outcome of 5-ALA-PDT, and their inhibitors can be adopted as clinical modulators to elicit far greater efficacy in 5-ALA-PDT treatments. SOD2 could also be used as a complementary marker, since 5-ALA-PDT is relevant to cell mitochondria. Finally, GPX4, GSTP1, and intracellular levels of GSH could make additional markers for 5-ALA-PDT, as they have previously been shown to be crucial modulators of hypericin PDT (Theodossiou et al., 2017).

5.7 PpIX effects on cell metabolism

Mitochondria are the cell's powerhouse organelles and they play a key role in cell respiration, survival, and metabolism (Rinaldi et al., 2018). Mitochondria contribute to the development and progression of a wide variety of human cancers. Actively replicating cancer cells undergo metabolic changes in order to get sufficient energy for their increased metabolic rates, providing the cells with building blocks for the assembly of cellular organelles, cytoskeleton, and membranes. Blocking this metabolic reprogramming would have a significant impact on the growth and propagation of various human cancers (Altieri, 2017; Loureiro et al., 2017).

Our cell metabolism studies demonstrated that 5-ALA-PDT had a detrimental effect on the OCR in all of the cell lines studied (T98G, MCF7, MDA-MB-231, and T47D) with the exception of U87 cells, which did not undergo profound changes on their basal respiration but exhibited a substantial, 5-ALA-PDT-induced decrease in their maximal respiratory capacity. The milder response of U87 cells is in line with our observation on their reinforced mitochondrial defences. With regard to the reduction in the maximal respiratory capacity, we hypothesized that since PpIX is produced in the mitochondria, 5-ALA-PDT may be causing a damage to some key mitochondrial component, e.g., the electron transport chain (ETC) complex III, as previously shown for hypericin PDT (Theodossiou et al., 2019; Theodossiou et al., 2008).

With respect to the glycolytic pathway of the cells, there was a detrimental PDT effect in all but U87 and MDA-MB-231 cells, which endured the least damage to their glycolytic machinery among the cell lines. Interestingly, both of these two cell lines exhibit a more “Warburg” than “Pasteur” type of metabolism. Following our results, it is possible that clinical inhibition of glycolysis prior- and/or post-PDT could lead to a profound enhancement of the PDT therapeutic outcome. Indeed, reversal of the Warburg effect has been shown to decrease cell proliferation and arrest the cell cycle in S phase, making it an attractive therapeutic option for GBM (Poteet et al., 2013).

5.8 Cercosporin: A PS with promising applications outside conventional PDT

Cercosporin (see Chapter 2.5) is a phototoxin produced by the members of the fungal genus *Cercospora*, playing a critical role in the capability of these fungi to parasitize plants (Daub, 1987a). It is structurally similar to several photosensitizing compounds such as hypericin as well as hypocrellin A and B (Jendželovská et al., 2016). Cercosporin toxicity to plants has been attributed to its ability to generate ROS and predominantly singlet oxygen upon activation by light (Daub et al., 2005), and as a result cercosporin has been found to be almost universally toxic. Interestingly, the only organisms that have shown resistance to cercosporin toxicity are *Cercospora* species themselves and other perylenequinone producing fungi such as *Alternaria* and *Cladosporium* (Daub, 1987b).

Although the effects of cercosporin on plant physiology and disease have been extensively studied, there is no documented knowledge on the effects of cercosporin on human cancer cell lines. It is therefore of great significance that paper I showed cercosporin to be highly toxic in three different cancer cell lines (T98G, U87, and

MCF7). As shown in the same study, cercosporin has its main absorption peak around 470 nm. This places cercosporin outside the PDT 'therapeutic window' between 600-800 nm, where the energy is sufficient for the generation of singlet oxygen and where light can penetrate deeper into the tissue (Agostinis et al., 2011). However, blue-shifted PSs, like cercosporin, have been shown to be of great use: (i) in antimicrobial photoinactivation (PDI) (Abrahamse and Hamblin, 2016), (ii) in various types of superficial and skin lesions, such as actinic keratosis (AK) (Gholam et al., 2018), basal cell carcinoma (BCC) (Maytin et al., 2018), melanoma (Akasov et al., 2019), and (iii) occasions where the probability of perforations are of major concern (e.g., esophageal cancer, cholangiocarcinoma, colon, stomach, bladder, and lung cancer), where shallower light penetration would be beneficial. Makdoui and colleagues showed that blue light activation of the naturally occurring PS riboflavin eradicated methicillin-resistant *Staphylococcus aureus* (MRSA), which is one of the most predominant bacterial pathogens associated with morbidity and mortality. He also suggested that blue light in combination with PSs could be further exploited for the treatment of skin diseases associated with keratinocyte hyperproliferation, such as psoriasis and atopic dermatitis, since it could effectively inactivate human keratinocytes (Makdoui et al., 2019). Comparative studies between blue and red light PDT for BCC (Maytin et al., 2018) and AK (Gholam et al., 2018) have shown that both types of PDT appear to be equally safe and virtually equally effective in treating BCC and thin AK, while blue light irradiation was found to be significantly less painful as rated by VAS. One possible explanation could be that blue light, with a relatively lower fluence rate (energy delivery over a longer time period), may cause less stimulation of PpIX-loaded nerve fibers as compared to red light (Babes et al., 2016). Indeed, daylight PDT with topical MAL application was shown to be completely pain-free during illumination (2.5 h of sunlight) and effective (>90% of 30 BCC tumours were eradicated at 3 months), supporting the idea that low-intensity PDT is less painful while remaining effective (Wiegell et al., 2014). Another explanation of this finding could be the increased penetration depth of up to 6mm by red light, which would probably reflect nerve stimulation and/or tissue damage by ROS (Morton et al., 2002).

Cercosporin exhibited a high singlet oxygen quantum yield in relation to other commonly used PSs as demonstrated in paper I. In this context, it could be a very valuable PS in future and emerging PDT applications, where external light administration is not required for the activation of the PS, and in particular in the treatment of hard-to-reach cancers. One of these cases is presented in paper II where accelerated protons were able to activate cercosporin internalized by two GBM cell lines (T98G, MO59K), yielding significantly higher cytotoxicity than in cells irradiated without the addition of cercosporin. These results were of great importance not only because we demonstrated that cercosporin may have a potential role in the clinic but also as we showed for the first time that PSs can be activated by accelerated protons. Another application where cercosporin could be very useful, without the need of external irradiation, is chemi- or bio-luminescence PDT (Theodossiou et al., 2003), where the PS is activated directly by chemi/bio-luminescent molecules when they come into close proximity within the cell.

Although the maximal absorption of cercosporin was found to be at 470 nm, it also exhibited a secondary shoulder towards the end of the primary absorption peak extending to approximately 580nm. This secondary peak can be used to achieve a deeper treatment as yellow light can penetrate the skin much deeper than blue light. In paper IV, we used yellow light to irradiate cercosporin-treated 3D cell cultures, which are more structurally similar to solid tumours, and we demonstrated that yellow-light treatment had a more profound effect on larger spheroids as compared to that of blue light.

5.9 Cercosporin accumulation and intracellular localization

PS uptake by cancer cells is critical for effective PDT treatment. Cercosporin uptake studies showed that there was a substantial difference in the three cell lines studied, with T98G exhibiting the highest intracellular accumulation of cercosporin, at least three times higher than U87 or MCF7 cells. The more efficient PS accumulation of T98G cells may also be one of the reasons behind their enhanced phototoxicity. On the other hand, in our 3D study (paper IV), incubation with cercosporin under the same conditions demonstrated that U87 cells showed the most efficient loading at the single cell level, which was two times more than T98G and T47D cells. In paper I, U87 cells exhibited the lowest cercosporin loading, compared to the other cell lines investigated. This could partly be explained by the size (volume) of the spheroids, which was differential among the cell lines and displayed a ratio of T98G:T47D:U87~1: 2.9: 3.6. U87 spheroids were 3 times larger in size, and therefore might have had the ability to accumulate more dye in the interstitium between cells, thereby forming a reservoir for cellular uptake. It was also shown by microscopy that U87 spheroids are more loosely arranged, in contrast with T47D and T98G, and possibly facilitating the retention and diffusion of cercosporin.

Subcellular localization of the PS is of utmost importance since ROS has a short half-life, and therefore acts in the PS's immediate vicinity (Dougherty et al., 1998). In that context, the type of photodamage that occurs in cells loaded with a PS depends, to an extent, on the localization of the PS within the cell. Mitochondria and ER are undoubtedly among the most prominent targets in PDT, and cercosporin was found to localize in both organelles in all cell lines studied (T98G, U87, and MCF7).

There are multiple reports on the central role of mitochondria in apoptotic cell death, where overlapping signaling pathways through intrinsic and extrinsic cell death stimuli converge to induce permeabilization of the mitochondrial membranes, leading to mitochondrial depolarization and cytosolic release of several apoptogenic proteins (Buytaert et al., 2007; Kroemer et al., 2007). The ER has a pivotal role in the maintenance of intracellular calcium homeostasis, protein synthesis, and posttranslational modifications as well as the proper folding, sorting, and trafficking of proteins (Moserova and Kralova, 2012). PDT-induced oxidative damage to the ER can lead to cell death due to unmanageable calcium release in the cytosol or through activation of the unfolded protein response (UPR), which may initiate the downstream cell death machinery (Moserova and Kralova, 2012). Previously, it has been shown that PDT with hypericin is damaging both the mitochondria, by critically depleting

mitochondrial aconitase (Theodossiou et al., 2009; Theodossiou et al., 2006), and the ER, by causing irreversible photodamage to the SERCA-2 calcium ATPases, resulting in a lethal overload of the cytosol with calcium (Buytaert et al., 2006; Theodossiou et al., 2009).

5.10 Cercosporin cellular metabolic effects

We also looked at the metabolic profiles, both for respiration and glycolysis, of the three cell lines investigated, before and after cercosporin PDT, to uncover the effects of cercosporin on cellular metabolism (paper I). T98G and MCF7 were the most respiratory cells, while U87 on the other hand displayed 2-3 times lower basal respiration as measured through their OCR. Interestingly, when cells were incubated with cercosporin under dark conditions, there was a slight reduction in the basal respiration, and in particular for MCF7 cells, which is a different response from what we had previously seen with hypericin (Theodossiou et al., 2019), where the basal respiration increased after addition of the PS. In previous studies, it was shown that hypericin is a substrate of mitochondrial ETC complex III, facilitating the transport of electrons at $\frac{1}{2}$ the V_{max} of the complex III natural substrate, namely ubiquinone (Theodossiou et al., 2008). We hypothesize that cercosporin also localizes in the mitochondrial complex III, which is supported by the findings that the cytotoxic synergy of cercosporin with copper is abrogated when pre-incubated with the complex III inhibitor antimycin A. The significant decrease in the maximal respiratory capacity of all the cell lines down to the level of the basal respiration after incubation with cercosporin under dark conditions could be attributed to the possibility of cercosporin being an inefficient complex III substrate.

Following PDT treatment, all the cells incubated with cercosporin showed a complete respiratory collapse and remained completely irresponsive to the stimuli of oligomycin, FCCP, and antimycin A/rotenone, which indicates either complete inhibition or photodamage of the electron transport chain. Alternatively, the TCA cycle inside the mitochondria could also be damaged. In paper IV, we demonstrated that the metabolic profiles of 3D cultures varied considerably in comparison to their corresponding 2D cultures in paper I. Even though U87 cells displayed the lowest respiratory capacity among the cells examined, in the 3D cell cultures all spheroids (T98G, U87, and T47D) exhibited similar OCR. Following PDT, there was a respiratory collapse in all spheroids examined, which is in good agreement with our 2D data.

With respect to the glycolytic activity, in paper I, we showed that T98G cells had the highest glycolytic rate. MCF7 and U87 cells had similar glycolytic rates about $\frac{1}{2}$ that of T98G. After cercosporin addition under dark conditions, the glycolytic rate of T98G cells was not affected but MCF7 and U87 rates dropped by 35%. After PDT, T98G cells had a dramatic decrease in their glycolytic rate but it was not completely abolished, maintaining $\sim\frac{1}{3}$ of their glycolytic rate. On the contrary, MCF7 and U87 cells experienced a complete collapse of their glycolytic levels. These results are in agreement with one of our recent publications where we demonstrated that hypericin-PDT caused a dramatic decrease of both the respiratory and glycolytic rates in MCF7 cells (Theodossiou et al., 2019). On the other hand, in the 3D cultures (paper IV), the

glycolytic rates of T98G and T47D spheroids collapsed, but not those of U87 spheroids, which remained at almost half capacity following PDT. This could be explained by the looser adhesion of U87 spheroids, which could possibly grant the spheroid inner layers easier access to glycolysis substrates from the media. This could also be linked to the differential spheroid viability, as in paper I, U87 cells presented the lowest OCR compared to the other cells lines (T98G, MCF7); however, their glycolytic activity was comparable to that of MCF7, indicating that U87 cells rely more on the glycolytic pathway for ATP production and hence survival (Liberti and Locasale, 2016).

5.11 Excitation of PSs by protons

In paper II, we sought to demonstrate that PSs can be excited by accelerated protons. The goal of this study was to establish a new hybrid treatment based on proton therapy, using generation of singlet oxygen from proton-activated PSs to eliminate cancerous lesions in synergy with ionizing radiation. We found the first indication that this would be feasible within the works by Birks et al. who studied, among other things, the scintillation of organic crystals, and particularly anthracene, an aromatic organic compound reminiscent of PSs when excited by protons (Birks, 1951).

In our study, apart from the seminal proof of principle of PS activation, we demonstrated that the use of a PS could accelerate the onset of the Bragg peak in places of high PS concentration, through the proton-induced fluorescence profile. We additionally showed that a panel of PSs could produce fluorescence and singlet oxygen, which is the major PDT cytotoxic mediator, upon proton excitation. The addition of the PS to the microenvironment can significantly increase the density of π -electrons, and hence greatly force an earlier onset of the Bragg peak. This would be of great value in cancer therapy, given the specific accumulation of PSs in tumours, in particular in the cases of brain cancers such as GBM, where the blood-brain barrier (BBB) is impaired, making the tumour-specific accumulation even higher. This preferential accumulation has been previously shown in the case of 5-ALA by an increased accumulation of PpIX in tumour and inflamed tissues, as compared to normal tissues (Olivo and Wilson, 2004). In addition, 5-ALA derived PpIX fluorescence has been demonstrated to accomplish higher sensitivity and diagnostic accuracy for prediction of the presence of diffuse gliomas than standard gadolinium-enhanced magnetic resonance imaging (Samkoe et al., 2011). Finally, PpIX has been successfully used intraoperatively to guide the GBM resection, resulting in improved complete resection of enhancing tumour (CRET) and gross total resection (GTR), without increasing the incidence of neurological morbidities (Schucht et al., 2012). Even though PpIX is a prominent candidate since it is produced endogenously as part of the heme biosynthetic pathway, other PSs have similarly shown preferential accumulation in the tumours such as temoporfin (mTHPC) (Zimmermann et al., 2001). The accumulation of the PS in GBM primary and secondary cancer lesions can force the onset of the Bragg peak at the target tissues. In any case, proton therapy has already been shown to more efficiently eliminate glioma stem cells, which significantly contribute to GBM progression and disease relapse through a ROS-dependent

mechanism (Alan Mitteer et al., 2015). That together with the ROS produced by PSs could significantly improve the clinical outcome for GBM patients.

To register the PS triplet state population, we measured the phosphorescence in dry gels. There was a prominent phosphorescence emission in dry gels and extended triplet state lifetime, due to the lack of water, oxygenation, and collisional quenching. Following the pronounced phosphorescence emissions that we obtained, we hypothesized that the triplet state was also populated in solutions, but possibly quenched through collisions with ambient oxygen. These collisions of the triplet PS with triplet, ground-state oxygen led to an energy transfer and a mutual spin flip, resulting in the formation of singlet oxygen and ground state (singlet) PS. The singlet oxygen generation was measured in PS solutions of acetone d_6 in which the singlet oxygen lifetime is maximal (singlet oxygen lifetime in acetone d_6 is $\sim 1046 \mu s$ as compared to $\sim 3.45 \mu s$ in water, $\sim 30.4 \mu s$ in benzene h_6 , $\sim 45.8 \mu s$ in acetone h_6 at $25^\circ C$ (Bregnhøj et al., 2016), and $\sim 19 \mu s$ in DMSO (Dvoranová et al., 2014)). These direct measurements were performed around $\sim 1270 nm$ with the use of a near infrared PMT (Hamamatsu R5509-73, Hamamatsu, Japan). This phosphorescence corresponds to the spin-forbidden radiative transition of singlet oxygen down to its triplet ground state. Apart from the direct singlet oxygen measurements, the singlet oxygen production was also verified in the PS PpIX, which is well known for its singlet oxygen-related photoproduct (photoporphyrin IX, pPpIX) (Cox et al., 1982a; Theodossiou and MacRobert, 2002). In this sense, with the use of the singlet oxygen quencher DABCO (Wilkinson et al., 1995), we demonstrated the efficient generation of pPpIX in the absence of DABCO, upon proton irradiation. Given that the photoproduct generation is directly related to the quantity of singlet oxygen produced (Cox et al., 1982b), it can possibly be used as a measure of dosimetry in PpIX PDT, but potentially also in PpIX PrDT.

5.12 Effects of Protodynamic Therapy on cells

Having established that protons can excite PSs in solutions and gels, we also sought to provide a proof-of-principle in 2D cell cultures. There was a differential response to protodynamic therapy in the three GBM cell lines examined (T98G, M059K, and U87). We used cercosporin, an efficient singlet oxygen producer (paper I and IV), and since protons transverse through matter, the short wavelength of cercosporin activation is not a matter of concern. From the cell lines studied, we observed that M059K exhibited the best results in terms of cytotoxicity for lower doses (2-10 Gy), while in T98G the highest efficacy was observed in higher doses (≥ 10 Gy). We speculate that this is likely due to M059K cells being markedly more sensitive to proton irradiation than T98G without PS added, and thus at higher radiation doses, leading to excessive cell death, it can be difficult to differentiate the cytotoxic contribution of protodynamic therapy from that of ionizing radiation. However, when we examined U87 cells, we did not observe any protodynamic mediated cell death. This could be due to the fact that U87 cells displayed a markedly lower cercosporin uptake than T98G cells (paper I) and M059K cells. Indeed, this was shown to be the case during the excitation of porphyrins by gamma-irradiation where radiosensitization efficiency

was dependent on PS intracellular accumulation (Luksiene et al., 2006). From our cell culture results, we could see that we got approximately 20% protodynamic additive effect as the mean effect, depending on the cell type and proton irradiation dose. This is a substantial in vitro observation, which if further translated could prove beneficial to cancer patients, resulting in more efficient eradication of the cancer cells, particularly in deep-lying lesions such as GBM. Our results lay the foundation for a novel treatment, combining the principles of proton therapy and PDT for more systematic tumour elimination.

5.13 Protodynamic Therapy limitations

In this project, we only investigated the PrDT technology in 2D cell cultures, and therefore we may have possibly missed the full potential of the treatment. The determination of the Bragg peak position is critical for our set up and had to be done independently for each experiment. The cell monolayers used in these experiments have very small dimensions along the proton beam, and therefore precise positioning is of utmost importance, while small deviations may mean substantially large divergence from optimal conditions. We are planning to overcome this limitation by examining the effects of protodynamic therapy in 3D cell cultures, since we expect to obtain the optimal efficacy depth without the necessity for calculating the exact position of the Bragg peak. This is because in longitudinal 3D cultures (e.g., in hydrogels), the maximal protodynamic cytotoxicity position will be determined by comparing the viability amongst different slices along the axis of beam propagation. In addition, during our work, we experienced variations in the beam flux and/or energy as a consequence of using a research-grade cyclotron. Experimental parameters had to be readjusted between experiments leading to possible discrepancies in experimental outcomes, which may consequently affect the statistics on outcomes from different runs. For more accuracy in the experimental conditions in the future, we intend to advance our research to automated clinical facilities in big proton centers, like the Oslo Proton Centre, which is currently being built within the Radium Hospital campus and expected to be operational in 2024.

One of the main limitations in cancer therapy, causing up to 90% of cancer related deaths, is metastatic disease (Weigelt et al., 2005). Overcoming the limitation of proton therapy and PDT to treat metastatic cancer, by either ROS-induced cytotoxicity or by in situ induction of an 'ectopic' proton Bragg peak, would be revolutionary for the field.

In this sense, we need to determine whether the PS accumulation in secondary disease lesion islets will provide a high enough density of delocalized electrons to trigger the formation of Bragg peaks on these lesions, resulting in efficient eradication of disseminated disease. Another challenge we need to address in vivo is the determination of the radiation dose required to activate the PSs of choice, and the optimal time interval between administration of PS and the irradiation. Furthermore, even though some data are available on the complications of PDT and proton therapy as monotherapies, it will be of outmost importance to identify the toxicity of the hybrid treatment we have introduced, i.e., protodynamic therapy.

Chapter 6

Conclusions

6.1 Synopsis

In the work included in this thesis, we demonstrated, for the first time, that accelerated protons can activate PSs to produce fluorescence and singlet oxygen in solutions and gels. We identified new PSs, which are very efficacious in PrDT and studied them in conventional PDT in vitro regimes, to gain knowledge of their capacity. Furthermore, we provided a proof of principle of the protondynamic therapy effects on two GBM cell lines. These findings may lay the foundations for a new hybrid therapy, based on proton therapy, for more efficient tumour eradication, utilizing, in synergy with ionizing irradiation, the ROS generated by the proton-PS interaction. Proton therapy shows particular promise in the treatment of brain tumours. The potential advantages of proton therapy in the treatment of brain tumours rely on the reduction of adverse effects such as cognitive dysfunction (by sparing critical regions around the tumour), in conjunction with potential dose escalation for radiation resistant tumours such as GBM. The advantages of proton therapy together with the strengths of PDT could help eliminate the requirement of an external light source, while simultaneously treating infiltrative tumour margins and small foci of disseminated disease. We are currently at a very early stage of this journey but we predict that, if successful, this therapy could proceed relatively quickly. With the increasing popularity of proton therapy and new proton facilities being built worldwide, protondynamic treatment is expected to be economical for both the health system and the patients, and it is anticipated to cost less than recurrent chemotherapeutic schemes, since the only additional cost to the proton therapy is the administration of the PS.

We also identified a new potential PS for PrDT, namely cercosporin. In our study, we demonstrated that cercosporin displays enhanced photocytotoxicity against two GBM and one breast adenocarcinoma cell lines, in part due to its strategic localization in the mitochondria and the ER, which are both vulnerable PDT targets. That makes cercosporin an ideal PS for off-shoot PDT applications, where light penetration depth is not relevant, such as chemi- or bio-luminescence PDT or PrDT/ProCI.

Cercosporin was shown to be effective as a PS not only in 2D but also in 3D cell cultures, a model that resembles more closely solid tumours where the light penetration in tissue is limited. Indeed, irradiation with yellow light was more efficient in larger spheroids than blue light.

The therapeutic outcome of PDT might be influenced by several resistance mechanisms. Increasing our knowledge of these resistance mechanisms, unveiling new prognostic markers for therapeutic outcome, together with adjuvant treatments and combination treatments to improve the curative effects of PDT are of utmost importance. In our study, some cell lines were found to be intrinsically less sensitive to 5-ALA-PDT than others, and we identified several mechanisms that may confer 5-

ALA-PDT-resistance. These included differential expression of cellular transporters, such as ABCG2 and GSTP1, and altered expression of enzymes, important in the rate limiting step in the heme biosynthetic pathway (FECH) as well as the heme degradation pathway (HO-1). Furthermore, we showed that the various cell lines exhibit diverse ROS scavenging capacities and that in some cases increased scavenging capacity did not protect against 5-ALA-PDT cytotoxicity, suggesting a multifactorial interplay.

6.2 Future Perspectives

In this thesis, we have demonstrated that a proton beam can electronically excite PSs and activate them to produce cytotoxic ROS. In a clinical setting, the therapeutic Bragg peak could offer a synergistic curative effect in conjunction with the ROS resulting from proton excitation of the PS. The next steps would require the treatment to be validated initially in 3D cultures and subsequently with tests on animals. It is widely accepted that multicellular spheroids provide a system of intermediate complexity between solid tumours in vivo and cell monolayers in culture (Sutherland et al., 1981); therefore, it would be an adequate starting point before initialization of animal experiments.

In future studies, we will also seek to provide proof of principle of the efficacy of pro-CI effects initially on GBM cell cultures and 3D spheroids, before we subsequently conduct in vivo experiments. This could potentially improve the specificity and efficacy of the therapeutic benefit as compared to PrDT, and it will be performed based on existing optimal protocols for PCI.

The results from our 5-ALA-PDT study propose the establishment of predictive biomarkers/therapy enhancers, converting it from a universal treatment to a tumour-specific oncological therapy. Depending on the status of the biomarkers, in our future studies, we will investigate the PpIX efficacy in PrDT, in combination with the above mentioned therapy enhancers, in order to restrain tumour resistance and enhance the treatment. We will monitor the PrDT responses in 2D and 3D cultures and subsequently in GBM mice models.

As mentioned earlier, cercosporin is a very potent singlet oxygen producer and exhibits enhanced photocytotoxicity not only in 2D and 3D cell cultures, but also in PrDT. Nevertheless, there is a need for the identification, synthesis, and validation of optimal, customised PSs for PrDT and ProCI.

We are currently at the very moment of emergence of a new technology for which our results may provide the foundation. This treatment would require proton beam irradiation with the supplementary administration of a PS, combining the strengths of proton therapy and PDT, without the necessity for an external light source. Our future efforts will be to translate this technology from the bench to the (proton therapy) bedside.

Bibliography

- Abrahamse, H., and Hamblin, M.R. (2016). New photosensitizers for photodynamic therapy. *Biochem J* 473, 347-364.
- Adamson, C., Kanu, O.O., Mehta, A.I., Di, C., Lin, N., Mattox, A.K., and Bigner, D.D. (2009). Glioblastoma multiforme: a review of where we have been and where we are going. *Expert opinion on investigational drugs* 18, 1061-1083.
- Agnihotri, S., Burrell, K.E., Wolf, A., Jalali, S., Hawkins, C., Rutka, J.T., and Zadeh, G. (2013). Glioblastoma, a brief review of history, molecular genetics, animal models and novel therapeutic strategies. *Arch Immunol Ther Exp (Warsz)* 61, 25-41.
- Agostinis, P., Berg, K., Cengel, K.A., Foster, T.H., Girotti, A.W., Gollnick, S.O., Hahn, S.M., Hamblin, M.R., Juzeniene, A., Kessel, D., *et al.* (2011). Photodynamic therapy of cancer: an update. *CA Cancer J Clin* 61, 250-281.
- Agostinis, P., Vantieghem, A., Merlevede, W., and de Witte, P.A. (2002). Hypericin in cancer treatment: more light on the way. *Int J Biochem Cell Biol* 34, 221-241.
- Akasov, R.A., Sholina, N.V., Khochenkov, D.A., Alova, A.V., Gorelkin, P.V., Erofeev, A.S., Generalova, A.N., and Khaydukov, E.V. (2019). Photodynamic therapy of melanoma by blue-light photoactivation of flavin mononucleotide. *Sci Rep* 9, 9679.
- Alan Mitteer, R., Wang, Y., Shah, J., Gordon, S., Fager, M., Butter, P.P., Jun Kim, H., Guardiola-Salmeron, C., Carabe-Fernandez, A., and Fan, Y. (2015). Proton beam radiation induces DNA damage and cell apoptosis in glioma stem cells through reactive oxygen species. *Scientific reports* 5, 13961.
- Alaoui-Jamali, M.A., Bismar, T.A., Gupta, A., Szarek, W.A., Su, J., Song, W., Xu, Y., Xu, B., Liu, G., Vlahakis, J.Z., *et al.* (2009). A novel experimental heme oxygenase-1-targeted therapy for hormone-refractory prostate cancer. *Cancer research* 69, 8017-8024.
- Ali, S.M., and Olivo, M. (2002). Bio-distribution and subcellular localization of Hypericin and its role in PDT induced apoptosis in cancer cells. *International journal of oncology* 21, 531-540.
- Ali, S.M., and Olivo, M. (2003). Mechanisms of action of phenanthroperylenequinones in photodynamic therapy (review). *International journal of oncology* 22, 1181-1191.
- Allalunis-Turner, M.J., Barron, G.M., Day, R.S., 3rd, Dobler, K.D., and Mirzayans, R. (1993). Isolation of two cell lines from a human malignant glioma specimen differing in sensitivity to radiation and chemotherapeutic drugs. *Radiation research* 134, 349-354.
- Allison, R.R., Downie, G.H., Cuenca, R., Hu, X.H., Childs, C.J., and Sibata, C.H. (2004). Photosensitizers in clinical PDT. *Photodiagnosis and photodynamic therapy* 1, 27-42.
- Altieri, D.C. (2017). Mitochondria on the move: emerging paradigms of organelle trafficking in tumour plasticity and metastasis. *British journal of cancer* 117, 301-305.
- Ares, C., Hug, E.B., Lomax, A.J., Bolsi, A., Timmermann, B., Rutz, H.P., Schuller, J.C., Pedroni, E., and Goitein, G. (2009). Effectiveness and safety of spot scanning proton radiation therapy for chordomas and chondrosarcomas of the skull base: first long-term report. *International journal of radiation oncology, biology, physics* 75, 1111-1118.
- Attili, S.K., Dawe, R., and Ibbotson, S. (2011). A review of pain experienced during topical photodynamic therapy--our experience in Dundee. *Photodiagnosis and photodynamic therapy* 8, 53-57.
- Awasthi, Y.C., Yang, Y., Tiwari, N.K., Patrick, B., Sharma, A., Li, J., and Awasthi, S. (2004). Regulation of 4-hydroxynonenal-mediated signaling by glutathione S-transferases. *Free radical biology & medicine* 37, 607-619.
- Baas, P., van Mansom, I., van Tinteren, H., Stewart, F.A., and van Zandwijk, N. (1995). Effect of N-acetylcysteine on Photofrin-induced skin photosensitivity in patients. *Lasers Surg Med* 16, 359-367.
- Babes, A., Sauer, S.K., Moparthy, L., Kichko, T.I., Neacsu, C., Namer, B., Filipovic, M., Zygmunt, P.M., Reeh, P.W., and Fischer, M.J. (2016). Photosensitization in Porphyrrias and Photodynamic Therapy Involves TRPA1 and TRPV1. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 36, 5264-5278.

- Begg, A.C., Stewart, F.A., and Vens, C. (2011). Strategies to improve radiotherapy with targeted drugs. *Nature reviews Cancer* 11, 239-253.
- Bellnier, D.A., and Dougherty, T.J. (1986). Haematoporphyrin derivative photosensitization and gamma-radiation damage interaction in Chinese hamster ovary fibroblasts. *International journal of radiation biology and related studies in physics, chemistry, and medicine* 50, 659-664.
- Ben-Hur, E., Kol, R., Marko, R., Riklis, E., and Rosenthal, I. (1988). Combined action of phthalocyanine photosensitization and gamma-radiation on mammalian cells. *International journal of radiation biology* 54, 21-30.
- Benov, L. (2015). Photodynamic therapy: current status and future directions. *Medical principles and practice : international journal of the Kuwait University, Health Science Centre* 24 Suppl 1, 14-28.
- Berg, K., Luksiene, Z., Moan, J., and Ma, L. (1995). Combined treatment of ionizing radiation and photosensitization by 5-aminolevulinic acid-induced protoporphyrin IX. *Radiation research* 142, 340-346.
- Berg, K., Nordstrand, S., Selbo, P.K., Tran, D.T., Angell-Petersen, E., and Høgset, A. (2011). Disulfonated tetraphenyl chlorin (TPCS2a), a novel photosensitizer developed for clinical utilization of photochemical internalization. *Photochemical & photobiological sciences : Official journal of the European Photochemistry Association and the European Society for Photobiology* 10, 1637-1651.
- Berg, K., Selbo, P.K., Prasmickaite, L., Tjelle, T.E., Sandvig, K., Moan, J., Gaudernack, G., Fodstad, O., Kjølsrud, S., Anholt, H., *et al.* (1999). Photochemical internalization: a novel technology for delivery of macromolecules into cytosol. *Cancer research* 59, 1180-1183.
- Bernier, J., Hall, E.J., and Giaccia, A. (2004). Radiation oncology: a century of achievements. *Nature reviews Cancer* 4, 737-747.
- Berridge, M.V., and Tan, A.S. (1993). Characterization of the cellular reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT): subcellular localization, substrate dependence, and involvement of mitochondrial electron transport in MTT reduction. *Archives of biochemistry and biophysics* 303, 474-482.
- Birks, J.B. (1950). The Properties of Ferromagnetic Compounds at Centimetre Wavelengths. *Proceedings of the Physical Society of London Section B* 63, 65-74.
- Birks, J.B. (1951). Scintillations from Organic Crystals - Specific Fluorescence and Relative Response to Different Radiations. *P Phys Soc Lond A* 64, 874-877.
- Birks, J.B., and Black, F.A. (1951). Deterioration of Anthracene under α -Particle Irradiation. *Proceedings of the Physical Society Section A* 64, 511-512.
- Boegheim, J.P., Dubbelman, T.M., Mullenders, L.H., and Van Steveninck, J. (1987). Photodynamic effects of haematoporphyrin derivative on DNA repair in murine L929 fibroblasts. *The Biochemical journal* 244, 711-715.
- Bregnhøj, M., Westberg, M., Jensen, F., and Ogilby, P.R. (2016). Solvent-dependent singlet oxygen lifetimes: temperature effects implicate tunneling and charge-transfer interactions. *Physical Chemistry Chemical Physics* 18, 22946-22961.
- Brigelius-Flohé, R., and Maiorino, M. (2013). Glutathione peroxidases. *Biochimica et Biophysica Acta (BBA) - General Subjects* 1830, 3289-3303.
- Brown, S.B., Brown, E.A., and Walker, I. (2004). The present and future role of photodynamic therapy in cancer treatment. *The Lancet Oncology* 5, 497-508.
- Buytaert, E., Callewaert, G., Hendrickx, N., Scorrano, L., Hartmann, D., Missiaen, L., Vandenheede, J.R., Heirman, I., Grooten, J., and Agostinis, P. (2006). Role of endoplasmic reticulum depletion and multidomain proapoptotic BAX and BAK proteins in shaping cell death after hypericin-mediated photodynamic therapy. *Faseb J* 20, 756-758.
- Buytaert, E., Dewaele, M., and Agostinis, P. (2007). Molecular effectors of multiple cell death pathways initiated by photodynamic therapy. *Biochim Biophys Acta* 1776, 86-107.
- Cairnduff, F., Stringer, M.R., Hudson, E.J., Ash, D.V., and Brown, S.B. (1994). Superficial photodynamic therapy with topical 5-aminolaevulinic acid for superficial primary and secondary skin cancer. *British journal of cancer* 69, 605-608.

- Calixto, G.M., Bernegossi, J., de Freitas, L.M., Fontana, C.R., and Chorilli, M. (2016). Nanotechnology-Based Drug Delivery Systems for Photodynamic Therapy of Cancer: A Review. *Molecules (Basel, Switzerland)* **21**, 342.
- Calzavara-Pinton, P.G. (1995). Repetitive photodynamic therapy with topical delta-aminolaevulinic acid as an appropriate approach to the routine treatment of superficial non-melanoma skin tumours. *Journal of photochemistry and photobiology B, Biology* **29**, 53-57.
- Casas, A., Di Venosa, G., Hasan, T., and Al, B. (2011). Mechanisms of resistance to photodynamic therapy. *Current medicinal chemistry* **18**, 2486-2515.
- Castano, A.P., Demidova, T.N., and Hamblin, M.R. (2004). Mechanisms in photodynamic therapy: part one-photosensitizers, photochemistry and cellular localization. *Photodiagnosis and photodynamic therapy* **1**, 279-293.
- Castano, A.P., Demidova, T.N., and Hamblin, M.R. (2005). Mechanisms in photodynamic therapy: part two-cellular signaling, cell metabolism and modes of cell death. *Photodiagnosis and photodynamic therapy* **2**, 1-23.
- Cavallini, L., Bindoli, A., Macri, F., and Vianello, A. (1979). Lipid peroxidation induced by cercosporin as a possible determinant of its toxicity. *Chem Biol Interact* **28**, 139-146.
- Clark, M.J., Homer, N., O'Connor, B.D., Chen, Z., Eskin, A., Lee, H., Merriman, B., and Nelson, S.F. (2010). U87MG decoded: the genomic sequence of a cytogenetically aberrant human cancer cell line. *PLoS Genet* **6**, e1000832.
- Collaud, S., Juzeniene, A., Moan, J., and Lange, N. (2004). On the selectivity of 5-aminolevulinic acid-induced protoporphyrin IX formation. *Current medicinal chemistry Anti-cancer agents* **4**, 301-316.
- Cox, G.S., Bobillier, C., and Whitten, D.G. (1982a). PHOTOOXIDATION AND SINGLET OXYGEN SENSITIZATION BY PROTOPORPHYRIN IX AND ITS PHOTOOXIDATION PRODUCTS. *Photochemistry and photobiology* **36**, 401-407.
- Cox, G.S., Krieg, M., and Whitten, D.G. (1982b). Photochemical reactivity in organized assemblies. 30. Self-sensitized photooxidation of protoporphyrin IX derivatives in aqueous surfactant solutions; product and mechanistic studies. *Journal of the American Chemical Society* **104**, 6930-6937.
- CS, F. (1976). Photosensitized oxidation and singlet oxygen: consequences in biological systems. *Free radicals in biology* **2**, 85-133.
- Daido, S., Yamamoto, A., Fujiwara, K., Sawaya, R., Kondo, S., and Kondo, Y. (2005). Inhibition of the DNA-dependent protein kinase catalytic subunit radiosensitizes malignant glioma cells by inducing autophagy. *Cancer research* **65**, 4368-4375.
- Darley-Usmar, V.M., Hogg, N., O'Leary, V.J., Wilson, M.T., and Moncada, S. (1992). The simultaneous generation of superoxide and nitric oxide can initiate lipid peroxidation in human low density lipoprotein. *Free Radic Res Commun* **17**, 9-20.
- Daub, M.E. (1982). Cercosporin, a photosensitizing toxin from *Cercospora* species. *Phytopathology* **72**, 370-374.
- Daub, M.E. (1987a). The Fungal Photosensitizer Cercosporin and Its Role in Plant Disease. In *Light-Activated Pesticides (American Chemical Society)*, pp. 271-280.
- Daub, M.E. (1987b). Resistance of fungi to the photosensitizing toxin, cercosporin. *Phytopathology* **77**, 1515-1520.
- Daub, M.E. (1992). Singlet oxygen yields, optical properties, and phototoxicity of reduced derivatives of the photosensitizer cercosporin. *Photochemistry and photobiology* **55**, 373-379.
- Daub, M.E., and Briggs, S.P. (1983). Changes in Tobacco Cell Membrane Composition and Structure Caused by Cercosporin. *Plant Physiology* **71**, 763.
- Daub, M.E., and Ehrenshaft, M. (2000). The Photoactivated *Cercospora* Toxin Cercosporin: Contributions to Plant Disease and Fundamental Biology. *Annu Rev Phytopathol* **38**, 461-490.
- Daub, M.E., and Hangarter, R.P. (1983). Light-induced production of singlet oxygen and superoxide by the fungal toxin, cercosporin. *Plant Physiol* **73**, 855-857.
- Daub, M.E., Herrero, S., and Chung, K.R. (2005). Photoactivated perylenequinone toxins in fungal pathogenesis of plants. *FEMS Microbiol Lett* **252**, 197-206.

- Daub, M.E., Herrero, S., and Chung, K.R. (2013). Reactive oxygen species in plant pathogenesis: the role of perylenequinone photosensitizers. *Antioxidants & redox signaling* **19**, 970-989.
- De Rosa, F.S., Tedesco, A.C., Lopez, R.F., Pierre, M.B., Lange, N., Marchetti, J.M., Rotta, J.C., and Bentley, M.V. (2003). In vitro skin permeation and retention of 5-aminolevulinic acid ester derivatives for photodynamic therapy. *Journal of controlled release : official journal of the Controlled Release Society* **89**, 261-269.
- Deavall, D.G., Martin, E.A., Horner, J.M., and Roberts, R. (2012). Drug-induced oxidative stress and toxicity. *Journal of toxicology* **2012**, 645460.
- Deng, H. (2015). Drug-delivery and multifunction possibilities of hypocrellin photosensitizers. *Journal of Innovative Optical Health Sciences* **8**, 13.
- Dietze, A., Engesaeter, B., and Berg, K. (2005). Transgene delivery and gelonin cytotoxicity enhanced by photochemical internalization in fibroblast-like synoviocytes (FLS) from rheumatoid arthritis patients. *Photochemical & photobiological sciences : Official journal of the European Photochemistry Association and the European Society for Photobiology* **4**, 341-347.
- Divakaruni, A.S., Paradyse, A., Ferrick, D.A., Murphy, A.N., and Jastroch, M. (2014). Analysis and interpretation of microplate-based oxygen consumption and pH data. *Methods Enzymol* **547**, 309-354.
- Diwu, Z., and Lown, J.W. (1993). Photosensitization with anticancer agents. 17. EPR studies of photodynamic action of hypericin: formation of semiquinone radical and activated oxygen species on illumination. *Free Radic Biol Med* **14**, 209-215.
- Dolmans, D.E., Fukumura, D., and Jain, R.K. (2003). Photodynamic therapy for cancer. *Nature reviews Cancer* **3**, 380-387.
- Donnelly, R.F., McCarron, P.A., and Woolfson, A.D. (2007). Derivatives of 5-aminolevulinic Acid for photodynamic therapy. *Perspectives in medicinal chemistry* **1**, 49-63.
- Dougherty, T.J. (1974). Activated dyes as antitumor agents. *J Natl Cancer Inst* **52**, 1333-1336.
- Dougherty, T.J., Gomer, C.J., Henderson, B.W., Jori, G., Kessel, D., Korbelik, M., Moan, J., and Peng, Q. (1998). Photodynamic therapy. *J Natl Cancer Inst* **90**, 889-905.
- Dougherty, T.J., Kaufman, J.E., Goldfarb, A., Weishaupt, K.R., Boyle, D., and Mittleman, A. (1978). Photoradiation therapy for the treatment of malignant tumors. *Cancer research* **38**, 2628-2635.
- Dougherty, T.J., Lawrence, G., Kaufman, J.H., Boyle, D., Weishaupt, K.R., and Goldfarb, A. (1979). Photoradiation in the treatment of recurrent breast carcinoma. *J Natl Cancer Inst* **62**, 231-237.
- Doyle, L.A., Yang, W., Abruzzo, L.V., Krogmann, T., Gao, Y., Rishi, A.K., and Ross, D.D. (1998). A multidrug resistance transporter from human MCF-7 breast cancer cells. *Proceedings of the National Academy of Sciences of the United States of America* **95**, 15665-15670.
- Dvoranová, D., Barbieriková, Z., and Brezová, V. (2014). Radical intermediates in photoinduced reactions on TiO₂ (an EPR spin trapping study). *Molecules (Basel, Switzerland)* **19**, 17279-17304.
- Ebermann, R., Alth, G., Kreitner, M., and Kubin, A. (1996). Natural products derived from plants as potential drugs for the photodynamic destruction of tumor cells. *Journal of photochemistry and photobiology B, Biology* **36**, 95-97.
- El-Missiry, M.A., Fayed, T.A., El-Sawy, M.R., and El-Sayed, A.A. (2007). Ameliorative effect of melatonin against gamma-irradiation-induced oxidative stress and tissue injury. *Ecotoxicol Environ Saf* **66**, 278-286.
- el-Sharabasy, M.M., el-Waseef, A.M., Hafez, M.M., and Salim, S.A. (1992). Porphyrin metabolism in some malignant diseases. *British journal of cancer* **65**, 409-412.
- El Ghissassi, F., Baan, R., Straif, K., Grosse, Y., Secretan, B., Bouvard, V., Benbrahim-Tallaa, L., Guha, N., Freeman, C., Galichet, L., *et al.* (2009). A review of human carcinogens--part D: radiation. *The Lancet Oncology* **10**, 751-752.
- Engelhardt, V., Kiesslich, T., Berlanda, J., Hofbauer, S., Krammer, B., and Plaetzer, K. (2011). Lipophilic rather than hydrophilic photosensitizers show strong adherence to standard cell culture microplates under cell-free conditions. *Journal of photochemistry and photobiology B, Biology* **103**, 222-229.

- Feruszová, J., Imreová, P., Bodnárová, K., Ševčovičová, A., Kyzek, S., Chalupa, I., Gálová, E., and Miadoková, E. (2016). Photoactivated hypericin is not genotoxic. *Gen Physiol Biophys* 35, 223-230.
- Figge, F.H., Weiland, G.S., and Manganiello, L.O. (1948). Cancer detection and therapy; affinity of neoplastic, embryonic, and traumatized tissues for porphyrins and metalloporphyrins. *Proc Soc Exp Biol Med* 68, 640.
- Fingar, V.H. (1996). Vascular effects of photodynamic therapy. *J Clin Laser Med Surg* 14, 323-328.
- Fischer, A.H., Jacobson, K.A., Rose, J., and Zeller, R. (2008). Fixation and permeabilization of cells and tissues. *CSH protocols* 2008, pdb.top36.
- Flanz, J., and Bortfeld, T. (2013). Evolution of technology to optimize the delivery of proton therapy: the third generation. *Semin Radiat Oncol* 23, 142-148.
- Frank, J., Lornejad-Schafer, M.R., Schoffl, H., Flaccus, A., Lambert, C., and Biesalski, H.K. (2007). Inhibition of heme oxygenase-1 increases responsiveness of melanoma cells to ALA-based photodynamic therapy. *International journal of oncology* 31, 1539-1545.
- Franken, A.C.W., Lokman, B.C., Ram, A.F.J., Punt, P.J., van den Hondel, C.A.M.J.J., and de Weert, S. (2011). Heme biosynthesis and its regulation: towards understanding and improvement of heme biosynthesis in filamentous fungi. *Appl Microbiol Biotechnol* 91, 447-460.
- Freitas, I. (1990). Lipid accumulation: the common feature to photosensitizer-retaining normal and malignant tissues. *Journal of photochemistry and photobiology B, Biology* 7, 359-361.
- Fromm, D., Kessel, D., and Webber, J. (1996). Feasibility of photodynamic therapy using endogenous photosensitization for colon cancer. *Archives of surgery (Chicago, Ill : 1960)* 131, 667-669.
- Fukuhara, H., Inoue, K., Satake, H., Tamura, K., Karashima, T., Yamasaki, I., Tatsuo, I., Kurabayashi, A., Furihata, M., and Shuin, T. (2011). Photodynamic diagnosis of positive margin during radical prostatectomy: preliminary experience with 5-aminolevulinic acid. *Int J Urol* 18, 585-591.
- Galanou, M.C., Theodossiou, T.A., Tsiourvas, D., Sideratou, Z., and Paleos, C.M. (2008). Interactive transport, subcellular relocation and enhanced phototoxicity of hypericin encapsulated in guanidinylated liposomes via molecular recognition. *Photochemistry and photobiology* 84, 1073-1083.
- Garg, A.D., Krysko, D.V., Vandenabeele, P., and Agostinis, P. (2012a). Hypericin-based photodynamic therapy induces surface exposure of damage-associated molecular patterns like HSP70 and calreticulin. *Cancer Immunol Immunother* 61, 215-221.
- Garg, A.D., Krysko, D.V., Verfaillie, T., Kaczmarek, A., Ferreira, G.B., Marysael, T., Rubio, N., Firczuk, M., Mathieu, C., Roebroek, A.J., *et al.* (2012b). A novel pathway combining calreticulin exposure and ATP secretion in immunogenic cancer cell death. *EMBO J* 31, 1062-1079.
- Gaullier, J.M., Berg, K., Peng, Q., Anholt, H., Selbo, P.K., Ma, L.W., and Moan, J. (1997). Use of 5-aminolevulinic acid esters to improve photodynamic therapy on cells in culture. *Cancer research* 57, 1481-1486.
- Gederaas, O.A., Holroyd, A., Brown, S.B., Vernon, D., Moan, J., and Berg, K. (2001). 5-Aminolaevulinic acid methyl ester transport on amino acid carriers in a human colon adenocarcinoma cell line. *Photochemistry and photobiology* 73, 164-169.
- Gholam, P., Bosselmann, I., Enk, A., and Fink, C. (2018). Impact of red versus blue light on tolerability and efficacy of PDT: a randomized controlled trial. *J Dtsch Dermatol Ges* 16, 711-717.
- Gold, M.H. (2009). Therapeutic and Aesthetic Uses of Photodynamic Therapy Part five of a five-part series: ALA-PDT and MAL-PDT What Makes Them Different. *J Clin Aesthet Dermatol* 2, 44-47.
- Gotardo, F., Cocca, L.H.Z., Acunha, T.V., Longoni, A., Toldo, J., Gonçalves, P.F.B., Iglesias, B.A., and De Boni, L. (2017). Investigating the intersystem crossing rate and triplet quantum yield of Protoporphyrin IX by means of pulse train fluorescence technique. *Chemical Physics Letters* 674, 48-57.
- Grant, B.D., and Donaldson, J.G. (2009). Pathways and mechanisms of endocytic recycling. *Nat Rev Mol Cell Biol* 10, 597-608.

- Grigalavicius, M., Mastrangelopoulou, M., Arous, D., Juzeniene, A., Ménard, M., Skarpen, E., Berg, K., and Theodossiou, T.A. (2020). Photodynamic Efficacy of Cercosporin in 3D Tumor Cell Cultures. *Photochemistry and photobiology* 96, 699-707.
- Grigalavicius, M., Mastrangelopoulou, M., Berg, K., Arous, D., Ménard, M., Raabe-Henriksen, T., Brondz, E., Siem, S., Görgen, A., Edin, N.F.J., *et al.* (2019). Proton-dynamic therapy following photosensitizer activation by accelerated protons demonstrated through fluorescence and singlet oxygen production. *Nat Commun* 10, 3986.
- Haar, C.P., Hebbbar, P., Wallace, G.C.t., Das, A., Vandergrift, W.A., 3rd, Smith, J.A., Giglio, P., Patel, S.J., Ray, S.K., and Banik, N.L. (2012). Drug resistance in glioblastoma: a mini review. *Neurochem Res* 37, 1192-1200.
- Hamblin, M.R., and Newman, E.L. (1994). On the mechanism of the tumour-localising effect in photodynamic therapy. *Journal of photochemistry and photobiology B, Biology* 23, 3-8.
- Han, W., and Yu, K.N. (2009). Response of cells to ionizing radiation. In *Advances in Biomedical Sciences and Engineering* (Bentham Science Publishers Ltd.), pp. 204-262.
- Hanahan, D., and Weinberg, R.A. (2000). The hallmarks of cancer. *Cell* 100, 57-70.
- Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of cancer: the next generation. *Cell* 144, 646-674.
- Hefti, M., Holenstein, F., Albert, I., Looser, H., and Luginbuehl, V. (2011). Susceptibility to 5-aminolevulinic acid based photodynamic therapy in WHO I meningioma cells corresponds to ferrochelatase activity. *Photochem Photobiol* 87, 235-241.
- Hellum, M., Høgset, A., Engesaeter, B.O., Prasmickaite, L., Stokke, T., Wheeler, C., and Berg, K. (2003). Photochemically enhanced gene delivery with cationic lipid formulations. *Photochemical & photobiological sciences : Official journal of the European Photochemistry Association and the European Society for Photobiology* 2, 407-411.
- Herceg, V., Lange, N., Allemann, E., and Babic, A. (2017). Activity of phosphatase-sensitive 5-aminolevulinic acid prodrugs in cancer cell lines. *Journal of photochemistry and photobiology B, Biology* 171, 34-42.
- Holland, E.C. (2000). Glioblastoma multiforme: the terminator. *Proc Natl Acad Sci U S A* 97, 6242-6244.
- Hollerman, W.A., Fisher, J.H., Ila, D., Jenkins, G.M., and Holland, L.R. (2011). Proton-induced fluorescence properties of terbium gallium garnet. *Journal of Materials Research* 10, 1861-1863.
- Hollerman, W.A., Fisher, J.H., Shelby, G.A., Holland, L.R., and Jenkins, G.M. (1992). Spectroscopic analysis of proton-induced fluorescence from yttrium and gadolinium oxysulfide phosphors. *IEEE Transactions on Nuclear Science* 39, 2295-2297.
- Horn, D., and Barrientos, A. (2008). Mitochondrial Copper Metabolism and Delivery to Cytochrome c Oxidase. *IUBMB life* 60, 421-429.
- Hudson, J.B., Imperial, V., Haugland, R.P., and Diwu, Z. (1997). Antiviral activities of photoactive perylenequinones. *Photochemistry and photobiology* 65, 352-354.
- Ibbotson, S.H. (2011). Adverse effects of topical photodynamic therapy. *Photodermatology, photoimmunology & photomedicine* 27, 116-130.
- Icha, J., Weber, M., Waters, J.C., and Norden, C. (2017). Phototoxicity in live fluorescence microscopy, and how to avoid it. *Bioessays* 39.
- Inoue, K., Fukuhara, H., Shimamoto, T., Kamada, M., Iiyama, T., Miyamura, M., Kurabayashi, A., Furihata, M., Tanimura, M., Watanabe, H., *et al.* (2012). Comparison between intravesical and oral administration of 5-aminolevulinic acid in the clinical benefit of photodynamic diagnosis for nonmuscle invasive bladder cancer. *Cancer* 118, 1062-1074.
- Ishikawa, M., and Hashimoto, Y. (2011). Improvement in aqueous solubility in small molecule drug discovery programs by disruption of molecular planarity and symmetry. *Journal of medicinal chemistry* 54, 1539-1554.
- Ishikawa, T., Kajimoto, Y., Inoue, Y., Ikegami, Y., and Kuroiwa, T. (2015). Critical role of ABCG2 in ALA-photodynamic diagnosis and therapy of human brain tumor. *Adv Cancer Res* 125, 197-216.

- Jendželovská, Z., Jendželovský, R., Hiřovská, L., Kovař, J., Mikeš, J., and Fedoročko, P. (2014). Single pre-treatment with hypericin, a St. John's wort secondary metabolite, attenuates cisplatin- and mitoxantrone-induced cell death in A2780, A2780cis and HL-60 cells. *Toxicol In Vitro* 28, 1259-1273.
- Jendželovská, Z., Jendželovský, R., Kuchárová, B., and Fedoročko, P. (2016). Hypericin in the Light and in the Dark: Two Sides of the Same Coin. *Front Plant Sci* 7, 560.
- Jerjes, W., Theodossiou, T.A., Hirschberg, H., Høgset, A., Weyergang, A., Selbo, P.K., Hamdoon, Z., Hopper, C., and Berg, K. (2020). Photochemical Internalization for Intracellular Drug Delivery. From Basic Mechanisms to Clinical Research. *J Clin Med* 9.
- Jiang, L., and He, Y. (2001). Photophysics, photochemistry and photobiology of hypocrellin photosensitizers. *Chinese Science Bulletin* 46, 6-16.
- Jichlinski, P., and Leisinger, H.J. (2001). Photodynamic therapy in superficial bladder cancer: past, present and future. *Urol Res* 29, 396-405.
- Jødal, L. (2009). Beta emitters and radiation protection. *Acta oncologica (Stockholm, Sweden)* 48, 308-313.
- Jori, G., and Reddi, E. (1993). The role of lipoproteins in the delivery of tumour-targeting photosensitizers. *The International journal of biochemistry* 25, 1369-1375.
- Jovčevska, I., Kočvar, N., and Komel, R. (2013). Glioma and glioblastoma - how much do we (not) know? *Mol Clin Oncol* 1, 935-941.
- Kamuhabwa, A.A., Cosserat-Gerardin, I., Didelon, J., Notter, D., Guillemin, F., Roskams, T., D'Hallewin, M.A., Baert, L., and de Witte, P.A. (2002). Biodistribution of hypericin in orthotopic transitional cell carcinoma bladder tumors: implication for whole bladder wall photodynamic therapy. *Int J Cancer* 97, 253-260.
- Karagas, M.R., Stannard, V.A., Mott, L.A., Slattery, M.J., Spencer, S.K., and Weinstock, M.A. (2002). Use of tanning devices and risk of basal cell and squamous cell skin cancers. *J Natl Cancer Inst* 94, 224-226.
- Kaur, G., and Dufour, J.M. (2012). Cell lines: Valuable tools or useless artifacts. *Spermatogenesis* 2, 1-5.
- Kavarnos, G., Nath, R., and Bongiorno, P. (1994). Visible-light and X irradiations of Chinese hamster lung cells treated with hematoporphyrin derivative. *Radiation research* 137, 196-201.
- Keefe, K.A., Chahine, E.B., DiSaia, P.J., Krasieva, T.B., Lin, F., Berns, M.W., and Tadir, Y. (2001). Fluorescence detection of cervical intraepithelial neoplasia for photodynamic therapy with the topical agents 5-aminolevulinic acid and benzoporphyrin-derivative monoacid ring. *American journal of obstetrics and gynecology* 184, 1164-1169.
- Kelly, J.F., and Snell, M.E. (1976). Hematoporphyrin derivative: a possible aid in the diagnosis and therapy of carcinoma of the bladder. *J Urol* 115, 150-151.
- Kennedy, J.C., and Pottier, R.H. (1992). Endogenous protoporphyrin IX, a clinically useful photosensitizer for photodynamic therapy. *Journal of photochemistry and photobiology B, Biology* 14, 275-292.
- Kennedy, J.C., Pottier, R.H., and Pross, D.C. (1990). Photodynamic therapy with endogenous protoporphyrin IX: basic principles and present clinical experience. *Journal of photochemistry and photobiology B, Biology* 6, 143-148.
- Kessel, D., and Reiners, J.J., Jr. (2007). Apoptosis and autophagy after mitochondrial or endoplasmic reticulum photodamage. *Photochemistry and photobiology* 83, 1024-1028.
- Kessel, D., and Thompson, P. (1987). Purification and analysis of hematoporphyrin and hematoporphyrin derivative by gel exclusion and reverse-phase chromatography. *Photochemistry and photobiology* 46, 1023-1025.
- Kiseleva, L.N., Kartashev, A.V., Vartanyan, N.L., Pinevich, A.A., and Samoilovich, M.P. (2016). CHARACTERISTICS OF A172 AND T98G CELL LINES. *Tsitologiya* 58, 349-355.
- Kleemann, B., Loos, B., Scriba, T.J., Lang, D., and Davids, L.M. (2014). St John's Wort (*Hypericum perforatum* L.) photomedicine: hypericin-photodynamic therapy induces metastatic melanoma cell death. *PLoS one* 9, e103762.

- Koizumi, N., Harada, Y., Minamikawa, T., Tanaka, H., Otsuji, E., and Takamatsu, T. (2016). Recent advances in photodynamic diagnosis of gastric cancer using 5-aminolevulinic acid. *World journal of gastroenterology* 22, 1289-1296.
- Kondo, M., Hirota, N., Takaoka, T., and Kajiwar, M. (1993). Heme-biosynthetic enzyme activities and porphyrin accumulation in normal liver and hepatoma cell lines of rat. *Cell Biol Toxicol* 9, 95-105.
- Konopka, K., and Goslinski, T. (2007). Photodynamic therapy in dentistry. *J Dent Res* 86, 694-707.
- Korbelik, M. (2006). PDT-associated host response and its role in the therapy outcome. *Lasers Surg Med* 38, 500-508.
- Korbelik, M., and Kros, G. (1995). Photofrin accumulation in malignant and host cell populations of a murine fibrosarcoma. *Photochemistry and photobiology* 62, 162-168.
- Korbelik, M., Kros, G., and Chaplin, D.J. (1991). Photofrin Uptake by Murine Macrophages. *Cancer research* 51, 2251.
- Korbelik, M., Parkins, C.S., Shibuya, H., Cecic, I., Stratford, M.R., and Chaplin, D.J. (2000). Nitric oxide production by tumour tissue: impact on the response to photodynamic therapy. *Br J Cancer* 82, 1835-1843.
- Korbelik, M., Sun, J., and Zeng, H. (2003). Ischaemia-reperfusion injury in photodynamic therapy-treated mouse tumours. *Br J Cancer* 88, 760-766.
- Kostron, H., Swartz, M.R., Miller, D.C., and Martuza, R.L. (1986). The interaction of hematoporphyrin derivative, light, and ionizing radiation in a rat glioma model. *Cancer* 57, 964-970.
- Kousis, P.C., Henderson, B.W., Maier, P.G., and Gollnick, S.O. (2007). Photodynamic therapy enhancement of antitumor immunity is regulated by neutrophils. *Cancer Res* 67, 10501-10510.
- Koval, J., Mikes, J., Jendzelovský, R., Kello, M., Solár, P., and Fedorocko, P. (2010). Degradation of HER2 receptor through hypericin-mediated photodynamic therapy. *Photochemistry and photobiology* 86, 200-205.
- Krammer, B., and Plaetzer, K. (2008). ALA and its clinical impact, from bench to bedside. *Photochemical & photobiological sciences : Official journal of the European Photochemistry Association and the European Society for Photobiology* 7, 283-289.
- Kroemer, G., Galluzzi, L., and Brenner, C. (2007). Mitochondrial membrane permeabilization in cell death. *Physiol Rev* 87, 99-163.
- Kuyama, S., and Tamura, T. (1957). Cercosporin. A Pigment of *Cercosporina Kikuchii* Matsumoto et Tomoyasu. I. Cultivation of Fungus, Isolation and Purification of Pigment. *Journal of the American Chemical Society* 79, 5725-5726.
- Laborde, E. (2010). Glutathione transferases as mediators of signaling pathways involved in cell proliferation and cell death. *Cell Death Differ* 17, 1373-1380.
- Lawrence, J.H., Tobias, C.A., Born, J.L., Mc, C.R., Roberts, J.E., Anger, H.O., Low-Beer, B.V., and Huggins, C.B. (1958). Pituitary irradiation with high-energy proton beams: a preliminary report. *Cancer research* 18, 121-134.
- Leibel, S.A., Fuks, Z., Zelefsky, M.J., Hunt, M., Burman, C.M., Mageras, G.S., Chui, C.S., Jackson, A., Amols, H.I., and Ling, C.C. (2003). Technological advances in external-beam radiation therapy for the treatment of localized prostate cancer. *Semin Oncol* 30, 596-615.
- Leroy, R., Benahmed, N., Hulstaert, F., Van Damme, N., and De Ruyscher, D. (2016). Proton Therapy in Children: A Systematic Review of Clinical Effectiveness in 15 Pediatric Cancers. *International journal of radiation oncology, biology, physics* 95, 267-278.
- Leunig, A., Mehlmann, M., Betz, C., Stepp, H., Arbogast, S., Grevers, G., and Baumgartner, R. (2001). Fluorescence staining of oral cancer using a topical application of 5-aminolevulinic acid: fluorescence microscopic studies. *Journal of photochemistry and photobiology B, Biology* 60, 44-49.
- Levin, W.P., Kooy, H., Loeffler, J.S., and DeLaney, T.F. (2005). Proton beam therapy. *British journal of cancer* 93, 849-854.
- Liang, H., Ran, Q., Jang, Y.C., Holstein, D., Lechleiter, J., McDonald-Marsh, T., Musatov, A., Song, W., Van Remmen, H., and Richardson, A. (2009). Glutathione peroxidase 4 differentially regulates the release of apoptogenic proteins from mitochondria. *Free Radical Biology and Medicine* 47, 312-320.

- Liberti, M.V., and Locasale, J.W. (2016). The Warburg Effect: How Does it Benefit Cancer Cells? *Trends in Biochemical Sciences* 41, 211-218.
- Lipson, R.L., and Baldes, E.J. (1960). The photodynamic properties of a particular hematoporphyrin derivative. *Arch Dermatol* 82, 508-516.
- Listowsky, I. (2005). Proposed intracellular regulatory functions of glutathione transferases by recognition and binding to S-glutathiolated proteins. *The journal of peptide research : official journal of the American Peptide Society* 65, 42-46.
- Loeffler, J.S., Smith, A.R., and Suit, H.D. (1997). The potential role of proton beams in radiation oncology. *Semin Oncol* 24, 686-695.
- Lomax, A.J., Bortfeld, T., Goitein, G., Debus, J., Dykstra, C., Tercier, P.A., Coucke, P.A., and Mirimanoff, R.O. (1999). A treatment planning inter-comparison of proton and intensity modulated photon radiotherapy. *Radiother Oncol* 51, 257-271.
- Lopez, R.F., Lange, N., Guy, R., and Bentley, M.V. (2004). Photodynamic therapy of skin cancer: controlled drug delivery of 5-ALA and its esters. *Advanced drug delivery reviews* 56, 77-94.
- Louis, D.N., Ohgaki, H., Wiestler, O.D., Cavenee, W.K., Burger, P.C., Jouvet, A., Scheithauer, B.W., and Kleihues, P. (2007). The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol* 114, 97-109.
- Loureiro, R., Mesquita, K.A., Magalhaes-Novais, S., Oliveira, P.J., and Vega-Naredo, I. (2017). Mitochondrial biology in cancer stem cells. *Semin Cancer Biol* 47, 18-28.
- Luksiene, Z., Juzenas, P., and Moan, J. (2006). Radiosensitization of tumours by porphyrins. *Cancer letters* 235, 40-47.
- Luksiene, Z., Kalvelyte, A., and Supino, R. (1999). On the combination of photodynamic therapy with ionizing radiation. *Journal of photochemistry and photobiology B, Biology* 52, 35-42.
- Ma, L., Iani, V., and Moan, J. (1993). Combination therapy: photochemotherapy; electric current; and ionizing radiation. Different combinations studied in a WiDr human colon adenocarcinoma cell line. *Journal of photochemistry and photobiology B, Biology* 21, 149-154.
- Macrobert, A.J.T., T. (2005). Photodynamic therapy of cancer. *Encyclopedia of Modern Optics* 1, 53–62
- Maehara, Y., Anai, H., Tamada, R., and Sugimachi, K. (1987). The ATP assay is more sensitive than the succinate dehydrogenase inhibition test for predicting cell viability. *European Journal of Cancer and Clinical Oncology* 23, 273-276.
- Maher, E.A., Furnari, F.B., Bachoo, R.M., Rowitch, D.H., Louis, D.N., Cavenee, W.K., and DePinho, R.A. (2001). Malignant glioma: genetics and biology of a grave matter. *Genes Dev* 15, 1311-1333.
- Makdoui, K., Hedin, M., and Bäckman, A. (2019). Different photodynamic effects of blue light with and without riboflavin on methicillin-resistant *Staphylococcus aureus* (MRSA) and human keratinocytes in vitro. *Lasers Med Sci* 34, 1799-1805.
- Mastrangelopoulou, M., Grigalavicius, M., Berg, K., Menard, M., and Theodossiou, T.A. (2019). Cytotoxic and Photocytotoxic Effects of Cercosporin on Human Tumor Cell Lines. *Photochemistry and photobiology* 95, 387-396.
- Mastrangelopoulou, M., Grigalavicius, M., Raabe, T.H., Skarpen, E., Juzenas, P., Peng, Q., Berg, K., and Theodossiou, T.A. (2020). Predictive biomarkers for 5-ALA-PDT can lead to personalized treatments and overcome tumor-specific resistances. *Cancer Reports n/a*, e1278.
- Maytin, E.V., Kaw, U., Ilyas, M., Mack, J.A., and Hu, B. (2018). Blue light versus red light for photodynamic therapy of basal cell carcinoma in patients with Gorlin syndrome: A bilaterally controlled comparison study. *Photodiagnosis and Photodynamic Therapy* 22, 7-13.
- Mikeš, J., Hýžďalová, M., Kočí, L., Jendželovský, R., Koval, J., Vaculová, A., Hofmanová, J., Kozubík, A., and Fedoročko, P. (2011). Lower sensitivity of FHC fetal colon epithelial cells to photodynamic therapy compared to HT-29 colon adenocarcinoma cells despite higher intracellular accumulation of hypericin. *Photochemical & photobiological sciences : Official journal of the European Photochemistry Association and the European Society for Photobiology* 10, 626-632.

- Mikešová, L., Mikeš, J., Kovaľ, J., Gyurászová, K., Culka, L., Vargová, J., Valeková, B., and Fedoročko, P. (2013). Conjunction of glutathione level, NAD(P)H/FAD redox status and hypericin content as a potential factor affecting colon cancer cell resistance to photodynamic therapy with hypericin. *Photodiagnosis and photodynamic therapy* 10, 470-483.
- Miller, R.C., Lodge, M., Murad, M.H., and Jones, B. (2013). Controversies in clinical trials in proton radiotherapy: the present and the future. *Semin Radiat Oncol* 23, 127-133.
- Moan, J., and Berg, K. (1991). The photodegradation of porphyrins in cells can be used to estimate the lifetime of singlet oxygen. *Photochemistry and photobiology* 53, 549-553.
- Mohan, R., and Grosshans, D. (2017). Proton therapy - Present and future. *Advanced drug delivery reviews* 109, 26-44.
- Morgan, J., and Oseroff, A.R. (2001). Mitochondria-based photodynamic anti-cancer therapy. *Advanced drug delivery reviews* 49, 71-86.
- Morton, C.A., Brown, S.B., Collins, S., Ibbotson, S., Jenkinson, H., Kurwa, H., Langmack, K., McKenna, K., Moseley, H., Pearse, A.D., *et al.* (2002). Guidelines for topical photodynamic therapy: report of a workshop of the British Photodermatology Group. *Br J Dermatol* 146, 552-567.
- Moseroova, I., and Kralova, J. (2012). Role of ER stress response in photodynamic therapy: ROS generated in different subcellular compartments trigger diverse cell death pathways. *PloS one* 7, e32972.
- Mrugala, M.M. (2013). Advances and challenges in the treatment of glioblastoma: a clinician's perspective. *Discov Med* 15, 221-230.
- Mulrooney, C.A., O'Brien, E.M., Morgan, B.J., and Kozlowski, M.C. (2012). Perylenequinones: Isolation, Synthesis, and Biological Activity. *European J Org Chem* 2012, 3887-3904.
- Neittaanmäki-Perttu, N., Grönroos, M., Karppinen, T.T., Tani, T.T., and Snellman, E. (2016). Hexyl-5-aminolaevulinate 0.2% vs. methyl-5-aminolaevulinate 16% daylight photodynamic therapy for treatment of actinic keratoses: results of a randomized double-blinded pilot trial. *The British journal of dermatology* 174, 427-429.
- Noël, G., Feuvret, L., Calugaru, V., Dhermain, F., Mammar, H., Haie-Méder, C., Ponvert, D., Hasboun, D., Ferrand, R., Nauraye, C., *et al.* (2005). Chordomas of the base of the skull and upper cervical spine. One hundred patients irradiated by a 3D conformal technique combining photon and proton beams. *Acta oncologica (Stockholm, Sweden)* 44, 700-708.
- Noell, S., Mayer, D., Strauss, W.S., Tatagiba, M.S., and Ritz, R. (2011). Selective enrichment of hypericin in malignant glioma: pioneering in vivo results. *International journal of oncology* 38, 1343-1348.
- Nokes, B., Apel, M., Jones, C., Brown, G., and Lang, J.E. (2013). Aminolevulinic acid (ALA): photodynamic detection and potential therapeutic applications. *J Surg Res* 181, 262-271.
- Norum, O.J., Selbo, P.K., Weyergang, A., Giercksky, K.E., and Berg, K. (2009). Photochemical internalization (PCI) in cancer therapy: from bench towards bedside medicine. *Journal of photochemistry and photobiology B, Biology* 96, 83-92.
- Nowell, P.C. (1976). The clonal evolution of tumor cell populations. *Science* 194, 23-28.
- Nyman, E.S., and Hynninen, P.H. (2004). Research advances in the use of tetrapyrrolic photosensitizers for photodynamic therapy. *Journal of photochemistry and photobiology B, Biology* 73, 1-28.
- Ochsner, M. (1997). Photophysical and photobiological processes in the photodynamic therapy of tumours. *Journal of photochemistry and photobiology B, Biology* 39, 1-18.
- Ohgari, Y., Nakayasu, Y., Kitajima, S., Sawamoto, M., Mori, H., Shimokawa, O., Matsui, H., and Taketani, S. (2005). Mechanisms involved in delta-aminolevulinic acid (ALA)-induced photosensitivity of tumor cells: relation of ferrochelatase and uptake of ALA to the accumulation of protoporphyrin. *Biochem Pharmacol* 71, 42-49.
- Olivo, M., and Chin, W. (2006). Perylenequinones in photodynamic therapy: cellular versus vascular response. *J Environ Pathol Toxicol Oncol* 25, 223-237.

- Olivo, M., and Wilson, B.C. (2004). Mapping ALA-induced PPIX fluorescence in normal brain and brain tumour using confocal fluorescence microscopy. *International journal of oncology* 25, 37-45.
- Olsen, D.R., Bruland, O.S., Frykholm, G., and Norderhaug, I.N. (2007). Proton therapy - a systematic review of clinical effectiveness. *Radiother Oncol* 83, 123-132.
- Orenstein, A., Kostenich, G., Roitman, L., Shechtman, Y., Kopolovic, Y., Ehrenberg, B., and Malik, Z. (1996). A comparative study of tissue distribution and photodynamic therapy selectivity of chlorin e6, Photofrin II and ALA-induced protoporphyrin IX in a colon carcinoma model. *British journal of cancer* 73, 937-944.
- Palasuberniam, P., Yang, X., Kraus, D., Jones, P., Myers, K.A., and Chen, B. (2015). ABCG2 transporter inhibitor restores the sensitivity of triple negative breast cancer cells to aminolevulinic acid-mediated photodynamic therapy. *Scientific reports* 5, 13298.
- Patel, S.H., Wang, Z., Wong, W.W., Murad, M.H., Buckley, C.R., Mohammed, K., Alahdab, F., Altayar, O., Nabhan, M., Schild, S.E., *et al.* (2014). Charged particle therapy versus photon therapy for paranasal sinus and nasal cavity malignant diseases: a systematic review and meta-analysis. *The Lancet Oncology* 15, 1027-1038.
- Peng, Q., Berg, K., Moan, J., Kongshaug, M., and Nesland, J.M. (1997a). 5-Aminolevulinic acid-based photodynamic therapy: principles and experimental research. *Photochemistry and photobiology* 65, 235-251.
- Peng, Q., Warloe, T., Berg, K., Moan, J., Kongshaug, M., Giercksky, K.E., and Nesland, J.M. (1997b). 5-Aminolevulinic acid-based photodynamic therapy. Clinical research and future challenges. *Cancer* 79, 2282-2308.
- Pham, A.N., Xing, G., Miller, C.J., and Waite, T.D. (2013). Fenton-like copper redox chemistry revisited: Hydrogen peroxide and superoxide mediation of copper-catalyzed oxidant production. *Journal of Catalysis* 301, 54-64.
- Ponka, P. (1999). Cell biology of heme. *Am J Med Sci* 318, 241-256.
- Poteet, E., Choudhury, G.R., Winters, A., Li, W., Ryou, M.-G., Liu, R., Tang, L., Ghorpade, A., Wen, Y., Yuan, F., *et al.* (2013). Reversing the Warburg effect as a treatment for glioblastoma. *J Biol Chem* 288, 9153-9164.
- Pottier, R., and Kennedy, J.C. (1990). The possible role of ionic species in selective biodistribution of photochemotherapeutic agents toward neoplastic tissue. *Journal of photochemistry and photobiology B, Biology* 8, 1-16.
- Prasmickaite, L., Høgset, A., Selbo, P.K., Engesaeter, B., Hellum, M., and Berg, K. (2002). Photochemical disruption of endocytic vesicles before delivery of drugs: a new strategy for cancer therapy. *British journal of cancer* 86, 652-657.
- Prasmickaite, L., Høgset, A., Tjelle, T.E., Olsen, V.M., and Berg, K. (2000). Role of endosomes in gene transfection mediated by photochemical internalisation (PCI). *The Journal of Gene Medicine* 2, 477-488.
- Prinsze, C., Penning, L.C., Dubbelman, T.M., and VanSteveninck, J. (1992). Interaction of photodynamic treatment and either hyperthermia or ionizing radiation and of ionizing radiation and hyperthermia with respect to cell killing of L929 fibroblasts, Chinese hamster ovary cells, and T24 human bladder carcinoma cells. *Cancer research* 52, 117-120.
- Regula, J., MacRobert, A.J., Gorchein, A., Buonaccorsi, G.A., Thorpe, S.M., Spencer, G.M., Hatfield, A.R., and Bown, S.G. (1995). Photosensitisation and photodynamic therapy of oesophageal, duodenal, and colorectal tumours using 5 aminolaevulinic acid induced protoporphyrin IX--a pilot study. *Gut* 36, 67-75.
- Reisz, J.A., Bansal, N., Qian, J., Zhao, W., and Furdui, C.M. (2014). Effects of ionizing radiation on biological molecules--mechanisms of damage and emerging methods of detection. *Antioxidants & redox signaling* 21, 260-292.
- Rinaldi, L., Delle Donne, R., Borzacchiello, D., Insabato, L., and Feliciello, A. (2018). The role of compartmentalized signaling pathways in the control of mitochondrial activities in cancer cells. *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer* 1869, 293-302.

- Rittenhouse-Diakun, K., Van Leengoed, H., Morgan, J., Hryhorenko, E., Paszkiewicz, G., Whitaker, J.E., and Oseroff, A.R. (1995). The role of transferrin receptor (CD71) in photodynamic therapy of activated and malignant lymphocytes using the heme precursor delta-aminolevulinic acid (ALA). *Photochemistry and photobiology* 61, 523-528.
- Roberts, W.G., and Hasan, T. (1992). Role of neovasculature and vascular permeability on the tumor retention of photodynamic agents. *Cancer research* 52, 924-930.
- Robey, R.W., Ierano, C., Zhan, Z., and Bates, S.E. (2011). The challenge of exploiting ABCG2 in the clinic. *Curr Pharm Biotechnol* 12, 595-608.
- Rübel, A., Handrick, R., Lindner, L.H., Steiger, M., Eibl, H., Budach, W., Belka, C., and Jendrossek, V. (2006). The membrane targeted apoptosis modulators erucylphosphocholine and erucylphosphohomocholine increase the radiation response of human glioblastoma cell lines in vitro. *Radiat Oncol* 1, 6.
- Rubino, G.F., and Rasetti, L. (1966). Porphyrin metabolism in human neoplastic tissues. *Panminerva Med* 8, 290-292.
- Rud, E., Gederaas, O., Høgset, A., and Berg, K. (2000). 5-aminolevulinic acid, but not 5-aminolevulinic acid esters, is transported into adenocarcinoma cells by system BETA transporters. *Photochemistry and photobiology* 71, 640-647.
- Samkoe, K.S., Gibbs-Strauss, S.L., Yang, H.H., Khan Hekmatyar, S., Jack Hoopes, P., O'Hara, J.A., Kauppinen, R.A., and Pogue, B.W. (2011). Protoporphyrin IX fluorescence contrast in invasive glioblastomas is linearly correlated with Gd enhanced magnetic resonance image contrast but has higher diagnostic accuracy. *J Biomed Opt* 16, 096008.
- Schnitzhofer, G.M., and Krammer, B. (1996). Photodynamic treatment and radiotherapy: combined effect on the colony-forming ability of V79 Chinese hamster fibroblasts. *Cancer letters* 108, 93-99.
- Schoenfeld, N., Epstein, O., Lahav, M., Mamet, R., Shaklai, M., and Atsmon, A. (1988). The heme biosynthetic pathway in lymphocytes of patients with malignant lymphoproliferative disorders. *Cancer letters* 43, 43-48.
- Schucht, P., Beck, J., Abu-Isa, J., Anderegg, L., Murek, M., Seidel, K., Stieglitz, L., and Raabe, A. (2012). Gross total resection rates in contemporary glioblastoma surgery: results of an institutional protocol combining 5-aminolevulinic acid intraoperative fluorescence imaging and brain mapping. *Neurosurgery* 71, 927-935; discussion 935-926.
- Singh, A., and Singh, H. (1982). Time-scale and nature of radiation-biological damage: approaches to radiation protection and post-irradiation therapy. *Prog Biophys Mol Biol* 39, 69-107.
- Smith, A.R. (2006). Proton therapy. *Phys Med Biol* 51, R491-504.
- Spikes, J.D. (1990). Chlorins as photosensitizers in biology and medicine. *Journal of photochemistry and photobiology B, Biology* 6, 259-274.
- Stout, D.L., and Becker, F.F. (1990). Heme synthesis in normal mouse liver and mouse liver tumors. *Cancer Res* 50, 2337-2340.
- Stummer, W., Novotny, A., Stepp, H., Goetz, C., Bise, K., and Reulen, H.J. (2000). Fluorescence-guided resection of glioblastoma multiforme by using 5-aminolevulinic acid-induced porphyrins: a prospective study in 52 consecutive patients. *Journal of neurosurgery* 93, 1003-1013.
- Stummer, W., Stocker, S., Wagner, S., Stepp, H., Fritsch, C., Goetz, C., Goetz, A.E., Kieffmann, R., and Reulen, H.J. (1998). Intraoperative detection of malignant gliomas by 5-aminolevulinic acid-induced porphyrin fluorescence. *Neurosurgery* 42, 518-525; discussion 525-516.
- Stupp, R., Mason, W.P., van den Bent, M.J., Weller, M., Fisher, B., Taphoorn, M.J., Belanger, K., Brandes, A.A., Marosi, C., Bogdahn, U., *et al.* (2005). Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med* 352, 987-996.
- Sutherland, R., Carlsson, J., Durand, R., and Yuhas, J. (1981). Spheroids in Cancer Research. *Cancer Research* 41, 2980.
- Sutherland, R., J (1981). Spheroids in cancer research. *Cancer Res* 41, 2980-2984.
- Svanberg, K., Andersson, T., Killander, D., Wang, I., Stenram, U., Andersson-Engels, S., Berg, R., Johansson, J., and Svanberg, S. (1994). Photodynamic therapy of non-melanoma malignant tumours

- of the skin using topical delta-amino levulinic acid sensitization and laser irradiation. *The British journal of dermatology* 130, 743-751.
- Szeimies, R.M., Landthaler, M., and Karrer, S. (2002). Non-oncologic indications for ALA-PDT. *The Journal of dermatological treatment* 13 Suppl 1, S13-18.
- Tamaoki, T., and Nakano, H. (1990). Potent and Specific Inhibitors of Protein Kinase C of Microbial Origin. *Bio/Technology* 8, 732-735.
- Teng, L., Nakada, M., Zhao, S.G., Endo, Y., Furuyama, N., Nambu, E., Pyko, I.V., Hayashi, Y., and Hamada, J.I. (2011). Silencing of ferrochelatase enhances 5-aminolevulinic acid-based fluorescence and photodynamic therapy efficacy. *British journal of cancer* 104, 798-807.
- Terasawa, T., Dvorak, T., Ip, S., Raman, G., Lau, J., and Trikalinos, T.A. (2009). Systematic review: charged-particle radiation therapy for cancer. *Ann Intern Med* 151, 556-565.
- Tetard, M.C., Vermandel, M., Mordon, S., Lejeune, J.P., and Reyns, N. (2014). Experimental use of photodynamic therapy in high grade gliomas: a review focused on 5-aminolevulinic acid. *Photodiagnosis and photodynamic therapy* 11, 319-330.
- Theodossiou, T., Hothersall, J.S., Woods, E.A., Okkenhaug, K., Jacobson, J., and MacRobert, A.J. (2003). Firefly luciferin-activated rose bengal: in vitro photodynamic therapy by intracellular chemiluminescence in transgenic NIH 3T3 cells. *Cancer Res* 63, 1818-1821.
- Theodossiou, T., and MacRobert, A.J. (2002). Comparison of the photodynamic effect of exogenous photoporphyrin and protoporphyrin IX on PAM 212 murine keratinocytes. *Photochemistry and photobiology* 76, 530-537.
- Theodossiou, T.A., Ali, M., Grigalavicius, M., Grallert, B., Dillard, P., Schink, K.O., Olsen, C.E., Walchli, S., Inderberg, E.M., Kubin, A., *et al.* (2019). Simultaneous defeat of MCF7 and MDA-MB-231 resistances by a hypericin PDT-tamoxifen hybrid therapy. *NPJ Breast Cancer* 5, 13.
- Theodossiou, T.A., Hothersall, J.S., De Witte, P.A., Pantos, A., and Agostinis, P. (2009). The multifaceted photocytotoxic profile of hypericin. *Mol Pharm* 6, 1775-1789.
- Theodossiou, T.A., Noronha-Dutra, A., and Hothersall, J.S. (2006). Mitochondria are a primary target of hypericin phototoxicity: synergy of intracellular calcium mobilisation in cell killing. *Int J Biochem Cell Biol* 38, 1946-1956.
- Theodossiou, T.A., Olsen, C.E., Jonsson, M., Kubin, A., Hothersall, J.S., and Berg, K. (2017). The diverse roles of glutathione-associated cell resistance against hypericin photodynamic therapy. *Redox Biol* 12, 191-197.
- Theodossiou, T.A., Papakyriakou, A., and Hothersall, J.S. (2008). Molecular modeling and experimental evidence for hypericin as a substrate for mitochondrial complex III; mitochondrial photodamage as demonstrated using specific inhibitors. *Free Radic Biol Med* 45, 1581-1590.
- Thompson, L.H. (2012). Recognition, signaling, and repair of DNA double-strand breaks produced by ionizing radiation in mammalian cells: the molecular choreography. *Mutat Res* 751, 158-246.
- Toh, K., Sakasai, K., Nakamura, T., Soyama, K., Nagata, S., Tsuchiya, B., and Shikama, T. (2009). Proton-induced fluorescence and long lasting emission of Sr₄Al₁₄O₂₅:Eu²⁺, Dy³⁺. Paper presented at: 2009 IEEE Nuclear Science Symposium Conference Record (NSS/MIC).
- Traverso, N., Ricciarelli, R., Nitti, M., Marengo, B., Furfaro, A.L., Pronzato, M.A., Marinari, U.M., and Domenicotti, C. (2013). Role of glutathione in cancer progression and chemoresistance. *Oxidative medicine and cellular longevity* 2013, 972913.
- Turro, N.J. (1991). *Modern Molecular Photochemistry*. Univ Science Books.
- Van Hillegersberg, R., Van den Berg, J.W., Kort, W.J., Terpstra, O.T., and Wilson, J.H. (1992). Selective accumulation of endogenously produced porphyrins in a liver metastasis model in rats. *Gastroenterology* 103, 647-651.
- Vistica, D.T., Skehan, P., Scudiero, D., Monks, A., Pittman, A., and Boyd, M.R. (1991). Tetrazolium-based assays for cellular viability: a critical examination of selected parameters affecting formazan production. *Cancer research* 51, 2515-2520.

- Wachowska, M., Muchowicz, A., Firczuk, M., Gabrysiak, M., Winiarska, M., Wańczyk, M., Bojarczuk, K., and Golab, J. (2011). Aminolevulinic Acid (ALA) as a Prodrug in Photodynamic Therapy of Cancer. *Molecules* (Basel, Switzerland) *16*, 4140-4164.
- Wadas, T.J., Pandya, D.N., Solingapuram Sai, K.K., and Mintz, A. (2014). Molecular targeted α -particle therapy for oncologic applications. *AJR Am J Roentgenol* *203*, 253-260.
- Walker, M.D., Alexander, E., Jr., Hunt, W.E., MacCarty, C.S., Mahaley, M.S., Jr., Mealey, J., Jr., Norrell, H.A., Owens, G., Ransohoff, J., Wilson, C.B., *et al.* (1978). Evaluation of BCNU and/or radiotherapy in the treatment of anaplastic gliomas. A cooperative clinical trial. *Journal of neurosurgery* *49*, 333-343.
- Wan, M.T., and Lin, J.Y. (2014). Current evidence and applications of photodynamic therapy in dermatology. *Clin Cosmet Investig Dermatol* *7*, 145-163.
- Ward, J.F. (1994). DNA damage as the cause of ionizing radiation-induced gene activation. *Radiation research* *138*, S85-88.
- Weber, D.C., Schneider, R., Goitein, G., Koch, T., Ares, C., Geismar, J.H., Schertler, A., Bolsi, A., and Hug, E.B. (2012). Spot scanning-based proton therapy for intracranial meningioma: long-term results from the Paul Scherrer Institute. *International journal of radiation oncology, biology, physics* *83*, 865-871.
- Wegiel, B., Nemeth, Z., Correa-Costa, M., Bulmer, A.C., and Otterbein, L.E. (2014). Heme oxygenase-1: a metabolic nuke. *Antioxid Redox Signal* *20*, 1709-1722.
- Weigelt, B., Peterse, J.L., and van 't Veer, L.J. (2005). Breast cancer metastasis: markers and models. *Nature reviews Cancer* *5*, 591-602.
- Wenkel, E., Thornton, A.F., Finkelstein, D., Adams, J., Lyons, S., De La Monte, S., Ojeman, R.G., and Munzenrider, J.E. (2000). Benign meningioma: partially resected, biopsied, and recurrent intracranial tumors treated with combined proton and photon radiotherapy. *International journal of radiation oncology, biology, physics* *48*, 1363-1370.
- Wiegell, S.R., Skødt, V., and Wulf, H.C. (2014). Daylight-mediated photodynamic therapy of basal cell carcinomas - an explorative study. *Journal of the European Academy of Dermatology and Venereology : JEADV* *28*, 169-175.
- Wilkinson, F., Helman, W.P., and Ross, A.B. (1995). Rate Constants for the Decay and Reactions of the Lowest Electronically Excited Singlet State of Molecular Oxygen in Solution. An Expanded and Revised Compilation. *Journal of Physical and Chemical Reference Data* *24*, 663-677.
- Wolf, P., Rieger, E., and Kerl, H. (1993). Topical photodynamic therapy with endogenous porphyrins after application of 5-aminolevulinic acid. An alternative treatment modality for solar keratoses, superficial squamous cell carcinomas, and basal cell carcinomas? *Journal of the American Academy of Dermatology* *28*, 17-21.
- Wong, A.D., Ye, M., Ulmschneider, M.B., and Searson, P.C. (2015). Quantitative Analysis of the Enhanced Permeation and Retention (EPR) Effect. *PloS one* *10*, e0123461.
- Wyss-Desserich, M.T., Sun, C.H., Wyss, P., Kurlawalla, C.S., Haller, U., Berns, M.W., and Tadir, Y. (1996). Accumulation of 5-aminolevulinic acid-induced protoporphyrin IX in normal and neoplastic human endometrial epithelial cells. *Biochemical and biophysical research communications* *224*, 819-824.
- Xu, J., Gao, J., and Wei, Q. (2016). Combination of Photodynamic Therapy with Radiotherapy for Cancer Treatment. *Journal of Nanomaterials* *2016*, 8507924.
- Xu, S., Chen, S., Zhang, M., and Shen, T. (2003). First synthesis of methylated hypocrellin and its fluorescent excited state: a cautionary tale. *The Journal of organic chemistry* *68*, 2048-2050.
- Yamazaki, S., Okubo, A., Akiyama, Y., and Fuwa, K. (1975). Cercosporin, a Novel Photodynamic Pigment Isolated from *Cercospora kikuchii*. *Agricultural and Biological Chemistry* *39*, 287-288.
- Yang, X., Palasuberniam, P., Kraus, D., and Chen, B. (2015). Aminolevulinic Acid-Based Tumor Detection and Therapy: Molecular Mechanisms and Strategies for Enhancement. *Int J Mol Sci* *16*, 25865-25880.

- Yuan, F., Leunig, M., Berk, D.A., and Jain, R.K. (1993). Microvascular permeability of albumin, vascular surface area, and vascular volume measured in human adenocarcinoma LS174T using dorsal chamber in SCID mice. *Microvascular research* 45, 269-289.
- Zhang, M., Qureshi, A.A., Geller, A.C., Frazier, L., Hunter, D.J., and Han, J. (2012). Use of tanning beds and incidence of skin cancer. *J Clin Oncol* 30, 1588-1593.
- Zhang, X., de Boer, L., Heiligers, L., Man-Bovenkerk, S., Selbo, P.K., Drijfhout, J.W., Høgset, A., and Zaat, S.A.J. (2018). Photochemical internalization enhances cytosolic release of antibiotic and increases its efficacy against staphylococcal infection. *Journal of controlled release : official journal of the Controlled Release Society* 283, 214-222.
- Zhao, B., and He, Y.Y. (2010). Recent advances in the prevention and treatment of skin cancer using photodynamic therapy. *Expert Rev Anticancer Ther* 10, 1797-1809.
- Zhenjun, D., and Lown, J.W. (1990). Hypocrellins and their use in photosensitization. *Photochemistry and photobiology* 52, 609-616.
- Zimmermann, A., Ritsch-Marte, M., and Kostron, H. (2001). mTHPC-mediated photodynamic diagnosis of malignant brain tumors. *Photochemistry and photobiology* 74, 611-616.
- Zitvogel, L., Kepp, O., and Kroemer, G. (2011). Immune parameters affecting the efficacy of chemotherapeutic regimens. *Nat Rev Clin Oncol* 8, 151-160.

Research Papers

ARTICLE

<https://doi.org/10.1038/s41467-019-12042-7>

OPEN

Proton-dynamic therapy following photosensitiser activation by accelerated protons demonstrated through fluorescence and singlet oxygen production

M. Grigalavicius¹ , M. Mastrangelopoulou¹, K. Berg¹, D. Arous¹, M. Ménard¹ , T. Raabe-Henriksen¹, E. Brondz², S. Siem², A. Görgen² , N.F.J. Edin² , E. Malinen^{2,3} & T.A. Theodossiou¹

We demonstrate excitation of photosensitisers (PSs) by accelerated protons to produce fluorescence and singlet oxygen. Their fluorescence follows a pattern similar to the proton energy loss in matter, while proton-derived fluorescence spectra match the photon-induced spectra. PSs excited in dry gelatin exhibit enhanced phosphorescence, suggesting an efficient PSs triplet state population. Singlet oxygen measurements, both optically at ~1270 nm and through the photoproduct of protoporphyrin IX (PpIX), demonstrate cytotoxic singlet oxygen generation by proton excitation. The singlet oxygen-specific scavenger 1,4-diazabicyclo [2.2.2]octane (DABCO) abrogates the photoproduct formation under proton excitation, but cannot countermand the overall loss of PpIX fluorescence. Furthermore, in two cell lines, M059K and T98G, we observe differential cell death upon the addition of the PS cercosporin, while in U87 cells we see no effect at any proton irradiation dose. Our results pave the way for a novel treatment combining proton therapy and “proton-dynamic therapy” for more efficient tumour eradication.

¹Department of Radiation Biology, Institute for Cancer Research, Radium Hospital, Oslo University Hospital, Montebello 0379 Oslo, Norway. ²Department of Physics, University of Oslo, Blindern 0315 Oslo, Norway. ³Department of Medical Physics, Oslo University Hospital, Montebello 0379 Oslo, Norway. Correspondence and requests for materials should be addressed to T.A.T. (email: ththeo@rr-research.no)

Photodynamic therapy (PDT)^{1,2} provides targeted cancer treatment through the synergy of three individually non-chemotoxic components: the photosensitiser (PS), i.e. a light-activated drug; light of the appropriate wavelength to excite the PS; and oxygen as the terminal generator of toxic species upon interaction with the excited PS^{3,4}. The photodynamic action is effected through the generation of reactive oxygen species (ROS) either by: (i) charge transfer between PS and molecules, which could involve oxygen superoxide anion and hydrogen peroxide ultimately leading to the formation of hydroxyl radicals⁵ (type I mechanism) or (ii) energy transfer, resulting in the generation of deleterious singlet oxygen [$O_2(^1\Delta_g)$ or 1O_2] (type II mechanism). The deexcitation of singlet oxygen to its triplet ground state, which is a spin forbidden transition, can be detected optically as low intensity phosphorescence at ~ 1270 nm.

The main limitation of PDT is the penetration depth of light, which can at most reach a few millimetres in tissue¹.

However, proton therapy⁶, the treatment of cancer with accelerated protons, can be tumour-specific, as protons with appropriate initial kinetic energy can deposit a high amount of radiant energy at any selected depth. When the proton beam enters and traverses tissue, the protons mainly lose energy by electromagnetic interactions with atomic electrons, initially at an almost constant rate. When the protons reach low energies, they rapidly deposit their remaining energy over a small region called the Bragg peak. This peak is the therapeutic/cytotoxic window that needs to be spatially-matched to the targeted cancerous lesion. The Bragg peak can be tuned to occur over a defined range within the tissue by selecting an appropriate span of initial proton energies. Thus, the depth of penetration of protons can be increased (or decreased) by amplifying (or reducing) the proton energy to include the target lesion in the Bragg peak. The main advantage of protons over high energy therapeutic X-rays is the lack of exit dose, as all their energy is delivered before, and especially at, the Bragg peak, while the main benefit of protons over visible therapeutic light is that protons can excite atoms over a broad energy range because of the continuous electromagnetic interactions with atomic electrons.

In the present study, we show the excitation of organic PSs in solutions or gels by accelerated protons to generate fluorescence and ROS. Most importantly, we present proof of principle of our hypothesis on two glioblastoma multiforme (GBM) cell lines. These findings are the first step towards a novel hybrid cancer treatment combining proton therapy with singlet-oxygen and other ROS-mediated cytotoxicity.

Results

Proton-induced fluorescence imaging. The experimental set-up used for the present work is shown in Fig. 1. The PSs fluorescence induced by protons, was imaged by a digital camera through a red filter. Analysis of the resulting images with FLUKA Monte Carlo software revealed that the spatial fluorescence pattern (Fig. 2) matched the simulated proton dose deposition pattern (Supplementary Fig. 1). The estimated proton energy at the cuvette entrance was 12.5 MeV, giving a simulated Bragg peak at around 1.8 mm in the PS solution. Correspondingly, the maximal fluorescence intensities were found at the depths of 1.6 mm (*meso*-tetrahydroxyphenyl chlorin—mTHPC, 55 μ M), 1.6 (Erythrosin B—ErB, 110 μ M), and 1.7 mm (protoporphyrin IX—PpIX, 70 μ M) from the cuvette entrance interface. In all cases (Fig. 2) there was a deviation between the fluorescence pattern and the proton dose deposition profile (elevated fluorescence prior to the Bragg peak onset), which will be further discussed.

Fig. 1 Experimental setup. Cyclotron and the main components in the proton beam path to the sample, including quadrupole magnets (Qs) to shape and focus the beam, switching magnets (SWM) to choose the exit port and an analysing magnet to select the energies by bending the beam 90° and the sample (S). The detection system comprised a grating spectrograph coupled to a synapse charge-coupled device (CCD) camera and a near-infrared photomultiplier tube (PMT)

Proton-induced fluorescence spectra. Proton-induced fluorescence spectra were also registered in PS solutions and compared to the corresponding spectra obtained by photonic excitation (Fig. 3). The proton-induced emissions in all cases matched spectrally the light-induced fluorescence. From a panel of nine PSs presented in Fig. 3, porphyrin- and chlorin-type PSs gave the strongest fluorescence while quinone-type compounds hypericin (Hyp) and cercosporin (Cerceo) produced weaker fluorescence at the same concentration and proton fluence rate. Of the PSs tested, hypocrellin A generated the weakest signal, observable only at concentrations higher than 20 μ M and with 2–4 times increased proton fluence rate (data not shown). We also studied two UV dyes, coumarin 102 and 6,8-difluoro-7-hydroxy-4-methylcoumarin (DiFMU) (Supplementary Fig. 2a, b). As expected, coumarin 102 yielded a ~ 4 -fold higher fluorescence intensity, even though its concentration was 25 times smaller (200 μ M vs. 5 mM), since coumarin 102 is a laser dye with a high fluorescence quantum yield in polar solvents⁷.

With the same detection system, we studied the fluorescence intensity dependence on PS concentration (Fig. 4a) and proton fluence (Fig. 4b) for selected PSs. Hypericin (Hyp, dimethyl sulfoxide [DMSO]) exhibited a linear response to higher concentrations and was employed to probe the relation between proton-induced fluorescence and proton fluence, which was found to be linear in the range of fluences studied. To determine the most efficient PS for fluorescence generation, the proton-induced fluorescence area was integrated for each PS (10μ M, 8×10^8 protons $\times$ cm⁻² $\times$ sec⁻¹) and divided by their corresponding integral of the extinction coefficients over the wavelengths from 350 to 750 nm. These values were plotted versus the integrated fluorescence for each PS (Fig. 4c). By normalising the integrated proton-induced fluorescence by the integrated extinction coefficient for each PS, we delineate the dependence on conjugated and resonant π electron systems, as these are responsible for visible light absorbance. In this sense Hyp, Cerceo, ErB and Rose Bengal (RoseB) are more efficient in utilising their electrons for interactions with protons, while mTHPC and aluminium 1,4-di

Fig. 2 Imaging of proton-induced fluorescence. **a–c** Fluorescence images captured during proton irradiation by a Samsung NX1000 camera (F-stop f/5.6, ISO speed 200) after a long pass filter with a cut-off wavelength of 610 nm. The resulting images were processed with MATLAB® to yield heat map representations of the fluorescence emission. **a** mTHPC 55 μM in DMSO. **b** PpIX 70 μM in dimethyl sulfoxide (DMSO) **c** Erythrosin B 110 μM in DMSO. **d–f** Top-view 2D contour heat map representations of **a–c** in an expanded scale. The black lines cross at the estimated position of fluorescence maximum. **g–i** Corresponding 3D representations of (**d–f**). Again, the black lines denote the estimated position of the maximum

(sulfonyloxy)phthalocyanine (AlPcS2a) possess more delocalised electrons and hence can produce more proton-induced fluorescence.

Proton-induced phosphorescence in dry gels. Proton-induced luminescence was also studied in anhydrous PS gels made from gelatin, which served as samples depleted of oxygen and with minimal intermolecular collisions (Fig. 5). The absence of collisional quenching (lack of oxygen) in dry gels led to enhanced phosphorescence emissions compared to solutions. The spectral profile and position of the proton-induced fluorescence and phosphorescence emission bands were verified by photonic

excitation. The peak intensity of ErB phosphorescence (Fig. 5b) was found to be at least 40 times higher than for RoseB (Fig. 5a), despite the fact that both sensitizers gave relatively similar intensity phosphorescence under photonic excitation (right axes).

Proton-induced singlet oxygen generation. We also registered the production of singlet oxygen from PS solutions by use of the near infrared PMT around ~ 1270 nm, despite the low intensity of the signal and low signal-to-noise ratio. Singlet oxygen luminescence registration was achieved for ErB and RoseB (Fig. 6a, b, respectively) in acetone d₆ (singlet oxygen lifetime in acetone d₆ is ~ 1046 μs vs 3.45 μs in water at 25 $^{\circ}\text{C}$)⁸. The generation of singlet

Fig. 3 Spectral profiles of proton-induced fluorescence. Emission spectra for nine photosensitisers (PSs) obtained under proton excitation are shown in separate panels in black. All PSs were studied at 10 μ M concentration in dimethyl sulfoxide (DMSO). The corresponding spectra of the PSs under light excitation were recorded from 10 μ M DMSO solutions and are shown in red

oxygen by PpIX was also shown through the registration of its singlet-oxygen-derived photoproduct. We followed the PpIX photoproduct evolution by using fluorescence spectroscopy after proton irradiation, in the absence and presence of 1,4-diazabicyclo [2.2.2]octane (DABCO), a specific singlet oxygen scavenger⁹. In contrast to the absence of DABCO, in the presence of DABCO no photoproduct could be detected within our irradiation regime (Fig. 6c). In addition, we studied the photo- vs. proton-degradation of PpIX fluorescence (Fig. 6d). In the case of light irradiation (Xenon lamp 350–750 nm, 16 mW $\times$ cm⁻¹), the fluorescence was only moderately reduced through photobleaching (max ~20%, Fig. 6d), and this was almost completely abrogated by the application of DABCO. However, in proton irradiation, the fluorescence loss was profound (max ~80%, Fig. 6d), and DABCO only partially protected (max ~25%, Fig. 6d, Supplementary Fig. 3) against this fluorescence depletion.

Proton-dynamic therapy in 2D cell cultures. Having established the proof of principle of the proton-dynamic technology in PS solutions and gels, we tested its efficacy on 2D cell cultures. Three human GBM cell lines were employed for these pilot experiments (M059K, T98G and U87), and their differential death following proton irradiation was assayed in the absence and presence of the potent PS Cerco¹⁰. The positive results for M059K and T98G cells are shown in Fig. 7; no differential proton-dynamic cell death was observed in the two U87 experiments performed.

In the case of M059K cells, as seen in the representative results (Fig. 7a), the differential death from the addition of the PS to the cells before irradiation was statistically significant for 2 ($P = 0.045$) and 5 Gy ($P = 0.02$), while for 10 Gy we had marginal statistical significance ($P = 0.065$). Correspondingly, in the case of T98G we had statistically significant enhancement of cytotoxicity due to proton-dynamic processes for 10 Gy ($P = 0.04$) and 20 Gy ($P = 0.006$). In the case of 30 Gy for T98G and 15 Gy for M059K, we could not perform any statistical analyses as these doses were tested once only due to their high toxicity. In Fig. 7c we have represented the differential gain in cytotoxicity due to proton-dynamic effects, i.e. due to the addition of PS prior to irradiation. From these data, we can see that the overall mean of gain from all five M059K experiments and for all doses is ~19%, while the median is ~16%. The corresponding values for T98G (4 experiments) were ~17% (mean) and ~10% (median). These results suggest that the M059K cells were more sensitive to the proton-dynamic effect at lower proton doses (2–10 Gy), while T98G cells were more responsive at higher radiation doses (10–30 Gy). Finally, as stated earlier, U87 cells were not responsive to proton-dynamic death at any of the doses of irradiation administered (data not shown).

Discussion

To the best of our knowledge, the excitation of PSs used in PDT by accelerated protons is a novel concept. This study was inspired

Fig. 4 Proton-induced fluorescence dependence. Proton-induced fluorescence dependence on photosensitiser (PS) concentration, proton flux and proton-induced fluorescence efficiency by PS. **a** Concentration dependence of the peak fluorescence intensity of Hyp, PpIX and mTHPC. **b** Fluorescence intensity dependence on proton fluence for Hyp (10 μM , dimethyl sulfoxide). **c** Comparison of the calculated proton-induced fluorescence efficiency between selected PSs scaled to methylene blue. The black squares show the integrated proton-induced fluorescence for each PS; the red circles show the same values normalised by the integrated extinction coefficient (350–750 nm) for each PS. Source data for c are provided as a Source Data file

Fig. 5 Proton-induced emission in dry photosensitiser (PS) gels. **a** Rose Bengal (RoseB) gel and **b** Erythrosin B (ErB) gel. The black lines correspond to the proton-induced emission of the gels (1 g gelatin (per 4 mL water, 500 μM PS), the blue lines denote the fluorescence emission from photonic excitation and the red lines show photon-induced phosphorescence ($\lambda_{\text{ex}} = 530 \text{ nm}$ for RoseB and 470 nm for ErB)

by the work of Birks¹¹, who explored the scintillation of organic crystals, and in particular anthracene as a solid, under the excitation of various types of radiation, including protons. Similar scintillation studies for the detection, discrimination and dosimetry of proton beams have used proton excitation but were

mainly on rare-earth complexes in crystal form^{12–14} and plastic-based scintillators¹⁵.

We looked for the first time at the excitation and activation of a panel of PSs in solutions and gels by accelerated protons through the registration of their fluorescence and singlet oxygen

Fig. 6 Proton-beam induced singlet oxygen and fluorescence degradation. Singlet oxygen phosphorescence, registered during proton beam irradiation at a fluence rate of $3.2 \times 10^9 \text{ protons cm}^{-2} \text{ sec}^{-1}$ in acetone d_6 solutions of **a** Erythrosin B (ErB) (60 μM) and **b** Rose Bengal (RoseB) (50 μM). **c** Photoproduct (photoproduct) production in the presence and absence of the singlet oxygen-specific scavenger 1,4-diazabicyclo[2.2.2]octane (DABCO) (200 mM) following proton irradiation of 50 μM PpIX in dimethyl sulfoxide (DMSO). The spectra were taken with a spectrofluorometer at 405 nm excitation. **d** Time evolution of PpIX (50 μM , DMSO) fluorescence intensity at 635 nm against irradiation dose either by a proton beam (crossed symbols) or by light (Xenon lamp, 350–750 nm, open symbols), in the presence (red squares) or absence (black circles) of DABCO (200 mM). In both cases, data were normalised to the unirradiated 635 nm maximum. The fluorescence data of panels **c** and **d** were acquired at 405 nm excitation

generation. The purpose of the study was to establish the basis for a new treatment concomitant with proton therapy, using PSs to destroy cancerous lesions by the generation of singlet oxygen in synergy to ionisations caused by the protons.

There have been a few attempts to use radiosensitisation by other forms of radiation such as γ -radiation and X-rays in combination with PSs. In those studies, PSs were employed as adjuvants with and without light excitation, while there was no indication of direct radiodynamic sensitisation of the PSs, e.g. through the registration of concomitant singlet oxygen generation^{16,17}.

Protons have a clear advantage over other radiation types such as γ -radiation and X-rays (photons) or β -radiation (electrons). As they traverse matter, they lose energy through interactions with the electrons they encounter on their course. Initially, the energy loss is almost constant with depth, up to the point where the cross section of interaction with electrons becomes very high and a high amount of energy is released in a spatially confined Bragg peak. This is in contrast with β - and γ - radiation and X-rays,

where the highest energy loss is upon entrance and the energy loss subsequently decreases with distance. This significant discrepancy allows the high-energy dissipation of protons to be tuned to a specific spatial position, such as the cancer lesion site in proton therapy. In addition, there is no energy left in the proton beam after the Bragg peak that could cause damage to normal tissue.

Our main hypotheses were that the use of a PS would: (i) accelerate the occurrence of the Bragg peak in places of high PS concentration (i.e. cancer lesions) and (ii) generate fluorescence and singlet oxygen upon proton excitation, which is the main PDT cytotoxic mediator. With respect to the first hypothesis, the analysis of the images in Fig. 2 clearly shows that the profile of proton-induced fluorescence is similar to the proton dose deposition profile (Supplementary Fig. 1). However, the spatial fluorescence pattern is distorted (Supplementary Fig. 1, solid lines), showing the induction of high-intensity fluorescence earlier than the Bragg peak-like maximum (Supplementary Fig. 1, black dots—simulated). This suggests that the PS may indeed be

Fig. 7 Proof of principle of proton-dynamic cell death in glioblastoma cell lines using cercosporin as the photosensitiser (PS). Differential cell death of M059K (**a**) and T98G (**b**) human glioblastoma cell lines, for varying radiation doses (2–5 Gy) with/without PS (cercosporin 4–10 μ M, 4 h incubation), vs. their respective controls. The grey bars depict cell groups without PS while the magenta bars depict cell groups with PS. These results showcase the best experiments so far (3 parallels). Errors are represented as standard errors of the mean. **c** Box plot illustrating the increase in proton-dynamic differential cell death (cells without PS vs with PS) from all experiments conducted on M059K and T98G cells. The magenta line shows the mean of all experiments while the black line represents the median. The error bars represent s.e.m. Source data for **a**, **b** are provided as Source Data files

accelerating the high-energy depositions prior to the Bragg peak by highly increasing the interaction cross sections (ErB, 110 μ M, Supplementary Fig. 1, red trace). The concentration of 110 μ M in solutions may, at first sight, seem excessive. If we, however, consider that the average volume of normal or neoplastic mononuclear cells is $\sim 200 \mu\text{m}^3$ or about 0.2 femtolitres (2×10^{-16} L)¹⁸, it would take the internalisation of a mere 13000 ErB molecules to reach 110 μ M intracellularly. Furthermore, if the PS had an affinity for subcellular organelles, e.g. the mitochondria, then the same concentration might be reached by 10–100 molecules per mitochondrion.

Regarding the second hypothesis, we have shown that accelerated protons were able to excite a panel of PSs to produce fluorescence and generate singlet oxygen. As protons traverse the medium with the PS, they undergo Coulomb interactions with delocalised (π) electrons, either ionising these electrons or exciting them to their higher, metastable electronic states from which

they return to the ground state emitting fluorescence. While in the singlet excited state, however, they can undergo intersystem crossing to the triplet state where they can interact with molecular oxygen (originally in triplet state). Through this collisional interaction both the PS and the oxygen undergo spin flip to their respective singlet states, and thus cytotoxic singlet oxygen is formed (Supplementary Fig. 4).

Two main factors determine the probability for ionisation or excitation of π electrons upon interaction with protons: (i) the kinetic energy of protons and (ii) the distance between the proton and the delocalised electron. With respect to kinetic energy, the faster the protons (higher initial energy), the shorter the time of interaction with the electrons and consequently the lower the probabilities, predominantly for electron ionisation, but also for excitation. The distance between the passing proton and the electron also determines the intensity of the Coulomb interaction, as this is (in the classical frame) inversely proportional to the

square of the distance. The closer the proton passes by the electron the stronger the probability for ionisation, whereas as the distance increases the probability of excitation vs. ionisation increases. As the Bragg peak is reached, the kinetic energy of the protons decreases to a point where the interaction time with bystander π electrons increases to an optimal value, giving a high cross section for interactions, leading both to electronic ionisations and excitations. This is probably the reason for the broadened profile of the fluorescence (Supplementary Fig. 1 red trace) with respect to the simulated Bragg peak in the absence of PS (Supplementary Fig. 1 black dotted trace). The addition of the PS increases the density of π electrons quite profoundly compared to the plain medium (solvent), and thus the probability for excitation is much higher even before the onset of the Bragg peak. This opens up the possibility of lesion-specific cancer therapy, due to the specific accumulation of PSs in tumours, especially in cases of brain cancers (e.g. GBM) in which the blood-brain barrier has been breached, making cancer-specific accumulation even higher. This has been previously shown for the PS PpIX, which is endogenously generated following the administration of 5-aminolevulinic acid (5-ALA)¹⁹. In fact, 5-ALA derived PpIX fluorescence has been found to have higher diagnostic accuracy for GBM than gadolinium-enhanced magnetic resonance imaging contrast²⁰, and PpIX has been used intraoperatively to define the of GBM lesions resection margins²¹. Even though PpIX has the advantage of being endogenously synthesised in the haem biosynthetic cycle, a remarkable selectivity for GBM has also been shown in the clinical context for other external PSs such as temoporfin (mTHPC)²². In light of this information, the specific accumulation of PSs in the cancerous lesion can expedite the onset of a Bragg peak on the lesion site and use the ROS derived from the proton-induced PS activation to kill the cancer cells in synergy with proton therapy.

In our study, we registered efficient fluorescence production following the excitation of a panel of PSs by accelerated protons. The spectral profiles of proton-induced fluorescence in all cases matched closely the corresponding profiles obtained by photonic excitation at 532 nm using the same acquisition parameters and registration system (CCD-spectrograph). The spikes seen in every luminescence registration in the present work are artefacts from the ionising radiation, as it is well known that CCDs are very sensitive even to cosmic rays. These spikes represent visible light scintillations in response to ionising radiation, which can be present both in charge-coupled silicon devices and metal-oxide semiconductors.

Looking at the luminescence from dry gels (Fig. 5), it is obvious that, in the absence of water and hence oxygenation but also collisional quenching, the proton-induced phosphorescence emission (verified spectrally by photonic excitation) is quite prominent, since the triplet state lifetime is significantly extended by the ErB and RoseB examples. In turn, this indicates that the triplet state is populated also in solutions, yet is very quickly quenched by collisions with ambient oxygen, which may lead to singlet oxygen formation via energy transfer and mutual spin flip. This is evidence that the PSs occupy their triplet state when excited by accelerated protons and is also an initial indication that from there they are available to interact with molecular oxygen to generate singlet oxygen. The fact that the proton-induced phosphorescence of ErB was 40-fold that of RoseB, albeit their light-induced phosphorescences and proton-induced fluorescences were comparable, might suggest that the spin-forbidden direct pumping to the triplet state in the case of ErB is much more favourable.

The registration of singlet oxygen from proton excitation of PSs is a very encouraging sign for the feasibility of a hybrid synergistic treatment using proton therapy and ROS. The level of

singlet oxygen production is quite low, but we have in the past registered cytotoxic results from chemiluminescence photosensitisation²³, in which singlet oxygen production was significantly lower. Additionally, in our experimental setup the proton energies were quite small (~ 15 – 16 MeV), and could only achieve a small penetration into matter (~ 2 mm in solvent). However, in therapeutic facilities, where energies in excess of 200 MeV are used, tissue penetration is quite substantial, reaching much deeper than light (Supplementary Fig. 5).

The singlet oxygen-associated PpIX photoproduct is another way of registering singlet oxygen generation²⁴. Using the data initially presented in Fig. 6d, we further analysed the formation of the PpIX photoproduct by light and proton irradiation. Naturally, the photoproduct undergoes bleaching much like the parent molecule (PpIX), since it is also a quite potent PS²⁵. By making the assumption that the photoproduct degradation is similar to that of PpIX, we have deconvoluted the photoproduct's evolution from its destruction in both proton and photon irradiation (Supplementary Fig. 6). In protonic and photonic excitation and for the fluences studied, the photoproduct formation is comparable between photon and proton irradiation and higher in the case of protons. Since the photoproduct formation is directly related to the amount of singlet oxygen generated²⁶, it can be used as dosimetry in PpIX PDT.

The fluorescence depletion versus irradiation time was much more efficient in the case of protons, as $\sim 10^{11}$ protons conferred the same depletion in fluorescence as $\sim 10^{20}$ photons. Moreover, in photonic irradiation, DABCO conferred an almost complete abrogation of the fluorescence bleaching, whilst in proton irradiation only a partial recovery of the fluorescence degradation was observed, despite the fact that DABCO completely suppressed the photoproduct formation (Fig. 6c). This implies that, although singlet oxygen is also partly involved in the fluorescence degradation by proton bombardment, mechanisms other than PS bleaching (mainly electron ionisation) are the predominant mechanisms of fluorescence loss and chemical modification.

In the clinic, prescription doses are based on the photon experience, adjusted for the radiobiological effectiveness of the protons (relative biological effectiveness = 1.1). In this sense, if we consider a clinically relevant daily dose of 50–60 Gy in 25–30 fractions, in our settings this would be equivalent to a continuous irradiation time of ~ 90 s (~ 0.66 Gy sec⁻¹ within the Bragg peak). This corresponds to a mere $\sim 10\%$ fluorescence loss (Fig. 6d). Moreover, in the cells our therapeutic doses varied from 2–30 Gy, which is much lower than in the clinical setting. Therefore, we do not expect the proton destruction of the PSs to be a limiting factor in the clinic.

In addition to solutions and gels, in the present study, we also present proof-of-principle in 2D cell cultures. From the results presented herein, we can see that the proton-dynamic effect is different in the three GBM cell lines investigated. The best overall effect was observed in the M059K cells for lower radiation doses (2–10 Gy), while in T98G the best effect was observed at higher doses (≥ 10 Gy). This is most probably due to the fact that the M059K cells are considerably more sensitive to proton irradiation (without PS) than T98G, and hence at high radiation-induced cytotoxicities in M059K it is difficult to differentiate the proton-dynamic effect from radiation cytotoxicity. U87, on the other hand, did not produce any detectable proton-dynamic cytotoxicity. This, however, can be related to our previous results¹⁰ showing that the cercosporin uptake of U87 cells was profoundly lower (~ 3 times) than that of T98G. We have reproduced these results to also include M059K cells (Supplementary Fig. 7). These data confirm the ~ 3 -times higher uptake of cerco by T98G in relation to U87, while M059K exhibit a ~ 2 -fold higher uptake than U87 cells.

The results presented in Fig. 7 show a ~20% additive proton-dynamic effect in the best case, depending on the tumour cell type. Although this enhancement is already very significant and could benefit patients with more effective elimination of the cancer cells in deep-lying lesions such as GBM, the full potential could be substantially higher than the results presented in 2D cultures. Cell monolayers have very small dimensions in the plane of proton beam incidence and hence require very precise positioning (in the range of μm) for optimisation of the proton-dynamic cytotoxicity. In our future research, we plan to evaluate the proton-dynamic efficacy in 3D cell cultures, as we expect that the depth effect will be seen more clearly and hence the optimal efficacy depth will be determined without any need for positioning in relation to the Bragg peak profile.

The determination of the Bragg peak position for cell experiments had to be done independently for each experiment in our setting. In addition, as expected for research-grade cyclotrons, we encountered variant beam conditions between experiments, which necessitated readjustment of the experimental parameters. These variations can account for the discrepancies in the outcomes of separate experiments, which in turn affect the overall statistical profiles of the average outcomes. For this reason, we plan to translate our research to clinical setups in big proton centres, where the experimental conditions are accurately controlled.

In this seminal study, we present the generation of fluorescence and associated singlet oxygen from the excitation of PSs in solutions and gels by accelerated protons. We also provide proof of principle of the efficacy of the proton-dynamic effects on GBM cell lines. Our results may lay the foundations for a new hybrid therapy, adjuvant to proton therapy, as the ROS generated by this interaction could destroy cancer cells in synergy with ionising radiation. Such a treatment would require proton beam irradiation at existing proton therapy facilities, with the additional administration of a PS, and would benefit from the therapeutic effects of proton therapy and PDT, without the requirement of an external light source. We are currently at the very beginning of this journey; however, we envisage the therapy could be integrated into current proton therapy practices fairly quickly (10–15 years). The next steps require the treatment to be validated in 3D cultures and tests on animals. At the same time an optimal, customised PS has to be identified, synthesised, validated and approved for clinical use.

Our findings could have significant applications not only in medicine but also in other fields, such as radiation dosimetry.

Methods

Chemicals and reagents. Dimethyl sulfoxide (DMSO), acetone d₆, 1,4-diazabicyclo[2.2.2]octane (DABCO), thiazolyl blue tetrazolium bromide (MTT), coumarin 102, cercosporin (Cerc), hematoporphyrin (HPD), protoporphyrin IX (PpIX), temoporfin (mTHPC), hypericin (Hyp), erythrosin B (ErB), Rose Bengal (RoseB) and methylene blue (MetBlue) were purchased from Sigma-Aldrich Norway AS (Oslo, Norway). Amphinex® (TPCS2a) was kindly provided by PCI Biotech AS (Oslo, Norway). *Meso*-tetra(p-hydroxyphenyl)porphyrin (mTHPP), aluminium 1,4-di(sulfonyloxy)phthalocyanine (AlPcS2a) and *N*-aspartyl chlorin e6 (NPe6) were purchased from Frontier Scientific (Logan, UT, USA). 6,8-difluoro-7-hydroxy-4-methylcoumarin (DiFMU) was obtained from ThermoFisher Scientific (Oslo, Norway). The gelatin powder was regular food grade, of animal origin (swine), (Dr. Oetker Norge AS, Kolbotn, Norway).

Proton irradiation. Protons were accelerated by an MC-35 Scanditronix cyclotron (Scanditronix, Uppsala, Sweden). The proton energies were selected at 16 MeV by the analysing magnet. An ionization chamber (Advanced Markus type, PTW, Freiburg, Germany) was used for estimating the absorbed proton dose D in units of Gy as in a previously established beamline for radiobiological experiments²⁷. The fluence Φ was calculated from the equation $D = \Phi \times (S/p)$, where (S/p) is the mass stopping power. Proton energy and stopping power were estimated as a function of depth using the residual range method. The fluence rate entering the sample was 8×10^8 protons $\times \text{cm}^{-2} \times \text{sec}^{-1}$ (5.3 Gy/s) in the case of spectral

measurements of fluorescence and phosphorescence as calculated. When capturing fluorescence images with a conventional Samsung NX1000 (Samsung Electronics Euro, Yateley, U.K.) camera, fluence rate was increased by a factor 40 to 32×10^9 protons $\times \text{cm}^{-2} \times \text{sec}^{-1}$. The sample was placed in all cases at a distance of 30 cm from the beam exit window and the lateral beam profile at the sample was each time verified by the use of Gafchromic film EBT3 (Radiation Products Design Inc., Albertville, MN, USA). The experimental setup simplified to a linear layout comprising the cyclotron, the light-sealed sample compartment and light detection system is shown in Fig. 1.

Proton-induced luminescence in solutions. The sample was placed at the focal point of a Horiba iHR 320 grating spectrograph cavity (HORIBA Europe GmbH, Gothenburg, Sweden). The solutions measurements were performed in horizontally-oriented quartz or a vertically-placed plastic cuvette at a 90-degree angle. Protons entered the cuvette through a polystyrene film of 8 μm thickness. For fluorescence collection, a 1200 groove per mm diffraction grating blazed at 630 nm (visible—VIS) was used to disperse the light, which was consequently collected by a Synapse 1024 $\times$ 256 VOE CCD camera (Horiba Instruments Inc., Edison, NJ, USA). For the registration of singlet oxygen luminescence, a 600 groove/mm diffraction grating blazed at 1500 nm (near infrared—NIR) was used to disperse the light, which was collected by a Hamamatsu R5509-73 liquid nitrogen-cooled photomultiplier tube (PMT, Hamamatsu, Japan). With regard to the singlet oxygen luminescence, proton irradiation generated singlet oxygen in the acetone d₆ solution without PS at a detectable level, which was however lower than the amount generated by the PS both in RB and ErB. In order to extract the pure singlet oxygen signal from the PS we had to (i) subtract the NIR PMT dark current and (ii) subtract the solvent contribution from that of the PS. The fluorescence images during proton irradiation were captured by a Samsung NX1000 camera (F-stop $f/5.6$, ISO speed 200) using a coloured long pass filter, with a cut-off wavelength of 610 nm placed in front of the objective. For the requirements of fluorescence imaging, the following PS concentrations were used: mTHPC 55 μM , PpIX 70 μM and ErB 110 μM . In the fluorescence spectral response measurements, all PSs were studied at 10 μM , while the UV dyes, coumarin 102 and 6,8-difluoro-7-hydroxy-4-methylcoumarin were studied at 200 μM and 5 mM, respectively. Singlet oxygen luminescence measurements were performed on 50 μM RB and 60 μM ErB, while PpIX photoproduct studies were conducted on a 50 μM solution.

Dry gels. 7.5 g of gelatin powder was dissolved into 30 mL of 80 °C Milli-Q® water by stirring. Aqueous stock solutions of either ErB or RoseB were mixed with the liquidised gelatin to reach 500 μM PS concentration before the water evaporated. The air-dried gels of 3 mm thickness were irradiated face-on with $\sim 10^9$ protons $\times \text{cm}^{-2} \times \text{sec}^{-1}$, and the overall photonic emission was registered by the iHR 320/Synapse CCD. Following light excitation (λ_{ex} = 530 nm for RoseB and 470 nm for ErB), the fluorescence and phosphorescence emission spectra were registered with the Varian Carry Eclipse spectrofluorometer (Agilent, Santa Clara, CA, USA), using a 5 ms gating time and 0.2 ms delay in the phosphorescence mode.

Light irradiation. Monochromatic excitation of PSs was facilitated by a 532 nm diode laser with variable output power (0–300 mW). The intensity of a collimated beam at the position of the sample was measured using an optic power meter, Gentec Solo2 (Gentec Electro-Optics Inc., Quebec, QC, Canada), equipped with a silicon photodetector, PH100 SiUV (Laser Components Nordic AS, Gothenburg, Sweden).

PpIX photoproduct formation under photonic excitation was achieved using a solar simulator, model 16S-300-002 (Solar Light Co., Inc., Glenside, PA, USA), equipped with a 150 W xenon arc lamp. KG-3 and appropriate dichroic filters were employed to ascertain a broadband spectrum of irradiation (350–750 nm). Spectral irradiance was measured using a portable Avantes AvaSpec 2048 $\times$ 14 FiberOptic spectrometer (Azpect Photonics AS, Södertälje, Sweden). By integrating the spectral irradiance, a total irradiance of 16 mW $\times \text{cm}^{-2}$ was obtained at the sample position. In both laser and Xenon lamp irradiation, the number of photons was calculated by multiplying the intensity of light at the particular wavelength by the exposure time in seconds and dividing by the energy of one single photon. In the case of the Xenon lamp, the number of photons at individual wavelengths was summed.

Cell experiments. 10⁶ cells (M059K CRL-2365™, T98G ATCC CRL-1690™, and U87 ATCC HTB-14™ human GBM cell lines, all purchased from and authenticated by ATCC, Manassas, Virginia, USA), were seeded in 60 mm petri dishes 24 h prior to the experiments and incubated overnight at 37 °C in a 5% CO₂ humidified atmosphere. Before experiments, the cells were tested negative for mycoplasma using Mycoalert®, Mycoplasma detection kit (Lonza, Rockland, ME, USA). The cells were divided into groups according to the irradiation dose (0–30 Gy) and the presence or absence of PS (cercosporin 4–10 μM for 4 h). Following PS, incubation media were removed from the cells, and they were irradiated by protons through the petri dish underside, within the Bragg peak as determined by the combination of a transmission chamber (Monitor Chamber 7862, PTW, Freiburg, Germany) before the petri dish and an ionisation chamber (Advanced Markus® Electron Chamber, PTW, Freiburg, Germany) after the petri dish. The cells never remained

without media for longer than 10 min, while all controls were treated in the same manner, even when not irradiated. All cell groups were carefully shielded from ambient light, while the proton irradiations were performed in dark ambient surroundings. The dosimetry was performed with respect to the values obtained with the calibrated Markus[®] ionisation chamber¹¹. During the irradiation, the cells were maintained at 37 °C using customised heated housing. Subsequently, fresh media were added to the cells and they were incubated for 20 h prior to the cytotoxicity assessment. The cytotoxicity was evaluated using a standard MTT assay: The cell media were replaced with 1 ml complete media containing 0.5 mg ml⁻¹ MTT. The cells were incubated for 1.5–2 h with the MTT media and then the media were removed and replaced with a 1 ml DMSO/dish. The dissolved formazan crystals were transferred to new plates that were read for absorbance, at 561 nm, by a Tecan spark M10 plate reader (Tecan Group Ltd., Männedorf, Switzerland). Blank values (wells with no cells) were in all cases subtracted. Cell viability was determined by comparison to the respective controls (irradiated cells without PS were compared to media controls, while PS-incubated and irradiated cells were compared to PS dark controls). In all experiments, each cell group consisted of at least 3 parallels.

Flow cytometry. U87, M059K and T98G cells were seeded in 60 mm-petri dishes (10⁶ cells per dish) and incubated overnight in their normal media at 37 °C, in a 5% CO₂ humidified atmosphere. After 4-h incubation with cercosporin, cells were trypsinized, washed in PBS, resuspended in 500 µL PBS containing Ca²⁺ and Mg²⁺ and measured with flow excitation at 488 nm. The fluorescence was registered in the PerCP channel (685/35-nm bandpass filter) of an LSRII (BD) flow cytometer. Flow cytometry data were analysed in FlowJo v.7.6.1 (Treestar Inc., Ashland, OR)

Software and data processing. To depict the fluorescence intensity of selected PSs versus sample depth in 2D and 3D modes, two RGB (true colour) images were acquired (camera model Samsung NX1000, F-stop f/5.6, ISO speed 200, focal length 50 mm, resolution 5472 × 3648 and bit depth 24) of the PSs in the custom-made cuvette with and without proton irradiation. The two corresponding images were spatially superimposed on each other using MATLAB[®] (MathWorks, Natick, MA, USA). The illuminated true colour image was converted to grayscale by eliminating the hue and saturation information while retaining intensity information for each pixel and thereby showing intensity vs depth in the solution.

With reference to Supplementary Figs. 1 and 5, the geometrical scheme of the proton beam system was defined by using Flair v2.3-0 (FLUKA Advanced Interface). The absorbed dose profiles of 16 MeV (in DMSO) and 200 MeV protons (in a head phantom) were simulated in FLUKA v2011.2×-4. In both, the Gaussian energy spread was set to ±0.1% full width at half maximum (FWHM) with a Gaussian FWHM lateral spread of 10 mm. The profile of 800 nm light penetration into the skin was reproduced by using the AccuRT radiative transfer model (Geminor Inc., Maplewood, NJ, USA), in which total irradiance has both diffuse and direct light components.

Computer code availability. All computer codes used for data analysis are available from the corresponding author on reasonable request.

Reporting summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Data availability

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Received: 28 March 2019 Accepted: 12 August 2019

Published online: 04 September 2019

References

- Macrobert, A. J. & Theodossiou, T. in *Encyclopedia of Modern Optics* Vol. 1 (ed R. D. Guenther) p. 53–62 (Elsevier, Amsterdam, 2005).
- Agostinis, P. et al. Photodynamic therapy of cancer: an update. *CA: a cancer J. Clin. Oncol.* **61**, 250–281 (2011).
- Dougherty, T. J. et al. Photodynamic therapy. *J. Natl. Cancer Inst.* **90**, 889–905 (1998).
- Dolmans, D. E., Fukumura, D. & Jain, R. K. Photodynamic therapy for cancer. *Nat. Rev.* **3**, 380–387 (2003).
- Fuchs, J. & Thiele, J. The role of oxygen in cutaneous photodynamic therapy. *Free Radic. Biol. Med.* **24**, 835–847 (1998).
- Levin, W. P., Kooy, H., Loeffler, J. S. & DeLaney, T. F. Proton beam therapy. *Br. J. Cancer* **93**, 849–854 (2005).
- Rurack, K. & Spieles, M. Fluorescence quantum yields of a series of red and near-infrared dyes emitting at 600–1000 nm. *Anal. Chem.* **83**, 1232–1242 (2011).
- Bregnhøj, M., Westberg, M., Jensen, F. & Ogilby, P. R. Solvent-dependent singlet oxygen lifetimes: temperature effects implicate tunneling and charge-transfer interactions. *Phys. Chem. Chem. Phys.: PCCP* **18**, 22946–22961 (2016).
- Wilkinson, F., Helman, W. P. & Ross, A. B. Rate constants for the decay and reactions of the lowest electronically excited singlet state of molecular oxygen in solution. An expanded and revised compilation. *J. Phys. Chem. Ref. Data* **24**, 663–677 (1995).
- Mastrangelopoulou, M., Grigalavicius, M., Berg, K., Menard, M. & Theodossiou, T. A. Cytotoxic and photocytotoxic effects of cercosporin on human tumor cell lines. *Photochem. Photobiol.* **95**, 387–396 (2019).
- Birks, J. B. Scintillations from organic crystals: specific fluorescence and relative response to different radiations. *Proc. Phys. Soc. A* **64**, 874–877 (1951).
- Hollerman, W. A. et al. Spectroscopic analysis of proton-induced fluorescence from yttrium and gadolinium oxysulfide phosphors. *IEEE Trans. Nucl. Sci.* **39**, 2295–2297 (1992).
- Hollerman, W. A., Fisher, J. H., Ila, D., Jenkins, G. M. & Holland, L. R. Proton-induced fluorescence properties of terbium gallium garnet. *J. Mater. Res.* **10**, 1861–1863 (1995).
- Toh, K. et al. Proton-induced fluorescence and long lasting emission of Sr₄Al₁₄O₂₅:Eu²⁺, Dy³⁺. *IEEE Nuc. Sci. Sym. Con. Rec.* **N25–84**, 1427–1430 (2009).
- Preedom, B. M., Saltmarsh, M. J., Ludemann, C. A. & Alster, J. The response of plastic scintillator to stopping protons in the energy range 70 to 100 MeV. *Nucl. Instrum. Methods* **133**, 311–314 (1976).
- Berg, K., Luksiene, Z., Moan, J. & Ma, L. Combined treatment of ionizing radiation and photosensitization by 5-aminolevulinic acid-induced protoporphyrin IX. *Radiat. Res.* **142**, 340–346 (1995).
- Luksiene, Z., Juzenas, P. & Moan, J. Radiosensitization of tumours by porphyrins. *Cancer Lett.* **235**, 40–47 (2006).
- Chapman, E. H., Kurec, A. S. & Davey, F. R. Cell volumes of normal and malignant mononuclear cells. *J. Clin. Pathol.* **34**, 1083–1090 (1981).
- Olivo, M. & Wilson, B. C. Mapping ALA-induced PPIX fluorescence in normal brain and brain tumour using confocal fluorescence microscopy. *Int. J. Oncol.* **25**, 37–45 (2004).
- Samkoe, K. S. et al. Protoporphyrin IX fluorescence contrast in invasive glioblastomas is linearly correlated with Gd enhanced magnetic resonance image contrast but has higher diagnostic accuracy. *J. Biomed. Opt.* **16**, 096008 (2011).
- Schucht, P. et al. Gross total resection rates in contemporary glioblastoma surgery: results of an institutional protocol combining 5-aminolevulinic acid intraoperative fluorescence imaging and brain mapping. *Neurosurgery* **71**, 927–935 (2012).
- Zimmermann, A., Ritsch-Marte, M. & Kostron, H. mTHPC-mediated photodynamic diagnosis of malignant brain tumors. *Photochem. Photobiol.* **74**, 611–616 (2001).
- Theodossiou, T. et al. Firefly luciferin-activated rose bengal: in vitro photodynamic therapy by intracellular chemiluminescence in transgenic NIH 3T3 cells. *Cancer Res.* **63**, 1818–1821 (2003).
- Cox, G. S., Bobillier, C. & Whitten, D. G. Photooxidation and singlet oxygen sensitization by Protoporphyrin IX and its photooxidation products. *Photochem. Photobiol.* **36**, 401–407 (1982).
- Theodossiou, T. & MacRobert, A. J. Comparison of the photodynamic effect of exogenous photoporphyrin and protoporphyrin IX on PAM 212 murine keratinocytes. *Photochem. Photobiol.* **76**, 530–537 (2002).
- Cox, G. S., Krieg, M. & Whitten, D. G. Photochemical reactivity in organized assemblies. 30. Self-sensitized photooxidation of protoporphyrin IX derivatives in aqueous surfactant solutions; product and mechanistic studies. *J. Am. Chem. Soc.* **104**, 6930–6937 (1982).
- Dahle, T. J. et al. Monte Carlo simulations of a low energy proton beamline for radiobiological experiments. *Acta Oncol.* **56**, 779–786 (2017).

Acknowledgements

The present work was financially supported by the South-Eastern Norway Regional Health Authority (Helse Sør Øst, project “protonic” no. 2017116). We would also like to thank Pawel A. Sobas, Victor Modamio and Jan C. Müller for the technical support with the cyclotron.

Author contributions

M.G., Experimental design/experimental planning/ experimental work/ bespoke technical constructions for proton-dynamic experiments, data analysis, manuscript writing and figures production; M.ArM., Experimental work/data analysis for proton-dynamic experiments, manuscript writing; K.B., experimental design/ data analysis for proton-dynamic experiments, manuscript writing; A.D., experimental work/image analysis for proton-dynamic experiments, Monte Carlo simulations, manuscript writing; M.atM., experimental work, manuscript editing; T.H.R., experimental work; E.B, bespoke technical constructions for proton-dynamic experiments; A.G., Experimental work (proton irradiation), manuscript editing; S.S., Experimental design input (cyclotron-irradiations),

manuscript editing, N.F.J.E., Experimental design input/experimental work (proton irradiations/ dosimetry), manuscript editing; E.M., Experimental design input/data analysis (proton irradiations/ dosimetry), manuscript editing; T.A.T., proton-dynamic concept, funding acquisition, project management, experimental design/experimental planning/ experimental work for proton-dynamic experiments, data analysis, manuscript workflow and writing, correspondence.

Additional information

Supplementary Information accompanies this paper at <https://doi.org/10.1038/s41467-019-12042-7>.

Competing interests: The authors declare no competing interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

Peer review information: *Nature Communications* thanks Jean-Olivier Durand and other anonymous reviewer(s) for their contribution to the peer review of this work. Peer reviewer reports are available

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019

**“Proton-dynamic therapy following photosensitiser activation by
accelerated protons demonstrated through fluorescence and singlet
oxygen production”**

Grigalavicius, M., et al.

Supplementary information

Supplementary Figure 1. Monte Carlo simulated dose deposition of 16 MeV protons in dimethyl sulfoxide (dotted line) vs proton-triggered fluorescence in a solution of mTHPC (55 μM , blue line), protoporphyrin IX (70 μM , green line) and erythrosin B (110 μM , red line).

Supplementary Figure 2. Spectral profiles of proton-induced fluorescence in UV dyes (coumarins). Emission spectra for **a** coumarin 102 and **b** 6,8-difluoro-7-hydroxy-4-methylcoumarin obtained under proton excitation are shown in separate panels in black. The two dyes were studied at **a** 200 μM and **b** 5mM concentration in dimethyl sulfoxide. The corresponding spectra of the photosensitisers under light excitation (407 nm) are shown in red.

Supplementary Figure 3. Relative protection (%) from fluorescence degradation by singlet oxygen scavenger DABCO. The data were derived from Fig. 6d.

Supplementary Figure 4. A simplified Jablonski scheme describing the main processes in the formation of singlet oxygen ($^1\Delta_g$) from solutions of photosensitisers under proton excitation.

Supplementary Figure 5. Light (800 nm) penetration into skin vs dose deposition of protons (200 MeV) in a head phantom simulating the skin, cranium, and brain.

Supplementary Figure 6. Evolution of fluorescence of photo-protoporphyrin IX at 670 nm following proton or photon irradiation, after deconvolution of protoporphyrin IX photobleaching by each time normalising spectra to 635 nm.

Supplementary Figure 7. Representative dot plots and adjunct histograms of cercosporin uptake by T98G (green), M059K (red), and U87 cells (cyan) in cercosporin (cerco) untreated and treated (4 μ M, 4h) cells. Uptake was measured by flow cytometry after incubation with cerco.

Predictive biomarkers for 5-ALA-PDT can lead to personalized treatments and overcome tumor-specific resistances

Maria Mastrangelopoulou¹ | Mantas Grigalavicius¹ | Tine H. Raabe¹ |
Ellen Skarpen² | Petras Juzenas³ | Qian Peng³ | Kristian Berg¹ |
Theodossis A. Theodossiou¹

¹Department of Radiation Biology, Institute for Cancer Research, Oslo University Hospital, Oslo, Norway

²Department of Molecular Cell Biology, Institute for Cancer Research, Oslo University Hospital, Oslo, Norway

³Department of Pathology, The Norwegian Radium Hospital, Oslo University Hospital, Oslo, Norway

Correspondence

Theodossis A. Theodossiou, Department of Radiation Biology, Institute for Cancer Research, Oslo University Hospital, Montebello, 0379 Oslo, Norway.
Email: ththeo@rr-research.no

Funding information

H2020 Future and Emerging Technologies, Grant/Award Number: 712921; Helse Sør-Øst RHF, Grant/Award Number: 2017116

Abstract

Background: Photodynamic therapy (PDT) is a minimally invasive, clinically approved therapy with numerous advantages over other mainstream cancer therapies. 5-aminolevulinic acid (5-ALA)-PDT is of particular interest, as it uses the photosensitizer PpIX, naturally produced in the heme pathway, following 5-ALA administration. Even though 5-ALA-PDT shows high specificity to cancers, differences in treatment outcomes call for predictive biomarkers to better stratify patients and to also diversify 5-ALA-PDT based on each cancer's phenotypic and genotypic individualities.

Aims: The present study seeks to highlight key biomarkers that may predict treatment outcome and simultaneously be exploited to overcome cancer-specific resistances to 5-ALA-PDT.

Methods and Results: We submitted two glioblastoma (T98G and U87) and three breast cancer (MCF7, MDA-MB-231, and T47D) cell lines to 5-ALA-PDT. Glioblastoma cells were the most resilient to 5-ALA-PDT, while intracellular production of 5-ALA-derived protoporphyrin IX (PpIX) could not account for the recorded PDT responses. We identified the levels of expression of ABCG2 transporters, ferrochelatase (FECH), and heme oxygenase (HO-1) as predictive biomarkers for 5-ALA-PDT. GPX4 and GSTP1 expression vs intracellular glutathione (GSH) levels also showed potential as PDT biomarkers.

For T98G cells, inhibition of ABCG2, FECH, HO-1, and/or intracellular GSH depletion led to profound PDT enhancement. Inhibition of ABCG2 in U87 cells was the only synergistic adjuvant to 5-ALA-PDT, rendering the otherwise resistant cell line fully responsive to 5-ALA-PDT.

ABCG2 or FECH inhibition significantly enhanced 5-ALA-PDT-induced MCF7 cytotoxicity, while for MDA-MB-231, ABCG2 inhibition and intracellular GSH depletion

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Cancer Reports* published by Wiley Periodicals LLC.

conferred profound synergies. FECH inhibition was the only synergism to ALA-PDT for the most susceptible among the cell lines, T47D cells.

Conclusion: This study demonstrates the heterogeneity in the cellular response to 5-ALA-PDT and identifies biomarkers that may be used to predict treatment outcome. The study also provides preliminary findings on the potential of inhibiting specific molecular targets to overcome inherent resistances to 5-ALA-PDT.

KEYWORDS

5-aminolevulinic acid, ABCG2 transporters, antioxidant defenses, ferrochelatase, heme oxygenase, metabolism, personalized therapy, photodynamic therapy

1 | INTRODUCTION

Current anticancer treatment modalities, such as ionizing radiation, chemotherapeutics, hormone-based therapies, and immunological treatments, risk failing as they encounter primary and secondary resistance, which varies not only from tumor to tumor but also within the same tumor, due to intrinsic heterogeneity.

Photodynamic therapy (PDT) is a minimally invasive, clinically approved therapy that uses three key components: a photoreactive drug (photosensitizer, PS), light of the appropriate wavelength, and oxygen. Upon light-activation, the PS interacts with ambient molecular oxygen, generating highly reactive oxygen species (ROS) and prominently singlet oxygen (1O_2), which facilitate cancer cell death.^{1–4}

PDT has numerous advantages over other mainstream cancer therapies, including cosmetic outcome, curative potential, and palliation to ensure improved quality of life, while it can be repeated as many times as required due to its minimal toxicity, low risk of treatment resistance, high treatment specificity, and low carcinogenic potential.⁵ Differences in treatment outcomes seen both preclinically and in patients call for predictive biomarkers to better stratify patients for PDT or alternative treatment modalities.

5-aminolevulinic acid (5-ALA)-PDT is a particularly attractive form of PDT, since 5-ALA is a natural intermediate in the heme synthesis that induces accumulation of the PS protoporphyrin IX (PpIX) upon administration as a prodrug to neoplastic cells. Iron (Fe^{2+}) is incorporated into PpIX to form heme by means of the enzyme ferrochelatase (FECH). Heme, unlike PpIX, lacks PS properties, since the chelation of iron mutes both fluorescence and singlet oxygen generation. The main asset of 5-ALA-PDT lies in its specificity for cancer cells, as PpIX formation is much higher in tumor cells.⁵ This has been mainly attributed to the significantly lower FECH, located in the inner mitochondrial membrane,⁶ and enhanced porphobilinogen deaminase (PBGD) activity in cancer cells, leading to reduced conversion of PpIX into heme.^{7–11} However, other factors, such as limited access to free iron, may also contribute to the high specificity for PpIX accumulation in cancer cells.¹² PpIX is converted to heme within 24 hours after 5-ALA administration, causing no long-term skin photosensitivity to the patients, one of the side-effects associated with PDT. Moreover, 5-ALA, being a hydrophilic and relatively small molecule (<200 Da), is

in principle much easier to administer systemically than other larger and lipophilic PSs frequently used for clinical PDT.

Due to these attractive features, 5-ALA-PDT was first introduced to the clinic in 1990 (Kennedy et al¹³) and has since become an established, yet underrated, clinical cancer treatment option primarily for dermatological malignancies. In addition, 5-ALA-derived PpIX photodynamic diagnosis has already been clinically applied for the diagnosis of bladder,¹⁴ prostate,¹⁵ and brain malignancies.^{16,17} 5-ALA was approved by the U.S. Food and Drug Administration (FDA) in 2017, as an adjunct for the visualization of malignant tissue, in grade III or IV glioma (NDA 208630/SN0014).

Glioblastoma multiforme (GBM), is an orphan, incurable brain cancer with poor patient prognosis and consequent high mortality. The main reasons behind the unfavorable GBM progression are: (a) extensive infiltration of the tumor cells into the surrounding brain tissue limiting the efficiency of surgical resection and (b) innate or rapidly acquired tumor resistance to chemotherapy and radiotherapy.^{18,19} At present, the standard of care for GBM comprises surgery, radiotherapy, and chemotherapy, giving a limited survival benefit to the patient however (approximately 15 months median survival vs 3 months when untreated).²⁰ Early clinical studies indicate that 5-ALA-PDT is safe and potentially beneficial in treating GBM.^{21,22}

Breast cancer, by contrast, is the most common cancer in women, accounting for 30% of all female cancers, and an eminent cause of female morbidity and mortality.²³ Standard treatment regimens for breast cancer include surgery, chemotherapy, immunotherapy, and radiation therapy, which have offered significant advances in primary disease treatment; however, refractory disease and recurrence remain prevalent.²⁴ Depending on their expression of therapy-related biomarkers, breast cancers can be classified as estrogen- and/or progesterone-receptor positive, human epidermal growth factor-receptor 2-positive, and triple-negative breast cancer (TNBC), which lacks the aforementioned receptors/markers.²⁵ TNBC is the most difficult to cure, displaying a higher risk for drug resistance and cancer recurrence.²⁴

We present an in vitro investigation of the outcome of 5-ALA-PDT on two human GBM cell lines, T98G and U87, and three breast adenocarcinoma cell lines, MCF7, MDA-MB-231, and T47D, and correlate the outcomes with the inherent characteristics of these cell lines. More specifically, we look into how 5-ALA-PDT-induced cell death is affected by

critical factors modulating the production of PpIX and its conversion into heme, cell metabolism, inherent antioxidant cell defenses, and mechanisms facilitating or inhibiting the expulsion of PpIX from the cells. The aim of the study is to establish crucial biomarkers, specific for 5-ALA-PDT, which will ultimately provide patient-specific prognosis for treatment outcomes and which may lead to personalized treatments based on 5-ALA-PDT with profoundly higher efficacy, even in cells initially non-responsive to 5-ALA-PDT alone.

2 | METHODS

2.1 | Chemicals and reagents

5-ALA, dimethyl sulfoxide (DMSO), oligomycin (oligo), carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP), myxothiazol (MYXO), sodium pyruvate, β -Nicotinamide adenine dinucleotide reduced (NADH), buthionine sulfoximine (BSO), Triton X-100, polyethylene glycol sorbitan monolaurate (TWEEN 20), thiazolyl blue tetrazolium bromide (MTT), metaphosphoric acid (MPA), triethanolamine (TEAM), anti- γ -tubulin, Ko143 hydrate, protease inhibitor cocktail, phosphatase inhibitor cocktail I and II, RPMI 1640 without phenol red, L-glutamine, penicillin-streptomycin, trypsin and Dulbecco's phosphate-buffered saline (PBS) with or without calcium chloride and magnesium chloride were purchased from Sigma-Aldrich Norway AS (Oslo, Norway). Noc-18, anti-FECH, and anti-superoxide dismutase 2 (SOD2) were purchased from Santa Cruz Biotechnology, Inc., (Dallas, Texas), and OB24 hydrochloride from Bio-Techne Ltd (Abingdon, UK). Fetal bovine serum (FBS), Pierce lane marker reducing sample buffer, TrypLE express enzyme, and MitoTracker red CMXRos were purchased from Thermo Fisher Scientific (Oslo, Norway). Precision Plus Protein dual color standard and 12% Mini-PROTEAN TGX precast proteins gels were purchased from Bio-Rad (Hercules, California). Immobilon Western Chemiluminescent HRP Substrate was purchased from Merck (Oslo, Norway). Anti-HO-1, anti-GPX4, and anti-BCRP/ABCG2 antibodies were purchased from Abcam (Cambridge, UK). Anti-superoxide dismutase 1 (SOD1) and anti-glutathione-S-transferase 1 (GSTP1) were purchased from Cell Signaling Technology, Inc. (Danvers, Massachusetts). The total glutathione (GSH) assay kit was purchased from Cayman Chemical Company (Ann Arbor, Michigan). The consumables used for the Seahorse XFe96 metabolic analysis were purchased from Agilent Technologies (Santa Clara, California).

2.2 | Cell culture

The cells used in this study were the human GBM cell lines T98G (ATCC CRL-1690) and U87 (ATCC HTB-14) and the human breast cancer cell lines MCF7 (ATCC HTB-22), MDA-MB-231 (ATCC HTB-26), and T47D (ATCC HTB-133). Since all cell lines were purchased from ATCC (LGC Standards GmbH, Wesel, Germany), no additional authentication was carried out; all cells were tested negative for mycoplasma (MycAlert, Lonza Group Ltd, Basel, Switzerland). The

cells were grown in RPMI 1640 without phenol red, supplemented with 10% FBS, 100 U/mL penicillin/100 μ g/mL streptomycin and 2 mM L-Glutamine at 37°C in a 5% CO₂ humidified atmosphere.

2.3 | PDT experiments and light irradiation

The cells were seeded in 96-well plates (Nunc, 1.5×10^4 cells/100 μ L media/well or Agilent Seahorse, 3×10^4 cells/100 μ L media/well) and incubated overnight in complete media. The cells were incubated with various concentrations of 5-ALA (0.1–2 mM) in Opti-MEM (Thermo Fisher Scientific) for 4 hours. Prior to irradiation, the cells were washed twice with PBS containing calcium and magnesium, and complete media were added to the cells. The cells were irradiated from the underside of the plate with blue light by means of an in-house-built lamp that was equipped with LEDs at 405 nm. The irradiance at the cell level was 5.6 mW/cm², as measured by an irradiance-calibrated AvaSpec-2048x14-SPU2 FiberOptic Spectrometer (Avantes, Apeldoorn, The Netherlands). Varying irradiation times were used (0–120 seconds), resulting in different light doses. In all the experiments, a light positive control without 5-ALA and dark controls with and without 5-ALA were included.

2.4 | Cell treatment with various inhibitors prior to 5-ALA-PDT

BSO treatment: BSO (100 μ M) was added to the cells overnight prior to incubation with 5-ALA and maintained in the respective treatment groups until the assessment of cytotoxicity (20 hours after irradiation). The cells were incubated with 5-ALA (0.2 mM T47D and MDA-MB-231, 0.5 mM MCF7, 1 mM T98G, and 2 mM U87) for 4 hours, washed twice in PBS, and irradiated at different time points per cell line to achieve LD₃₀ for each cell line (120 seconds T47D, 90 seconds MDA-MB-231, 60 seconds MCF7, 60 seconds U87, and 120 seconds T98G).

Treatment with other inhibitors: Ko143 (1 or 5 μ M) was added to the cells 1 hour prior to incubation with 5-ALA and maintained in the respective groups until the assessment of cytotoxicity (20 hours after irradiation). Noc-18 (0.5 mM) or OB24 (20 μ M) were added to the cells either overnight or 1 hour prior to 5-ALA addition and removed immediately prior to irradiation. The cell lines were incubated with different 5-ALA concentrations and irradiated at different times to achieve LD₃₀ conditions (see BSO treatment). A synergistic cytotoxicity of 5-ALA-PDT with the inhibitors was determined when $Viability (5-ALA + inhibitor) < Viability (5-ALA) \times Viability (inhibitor) / Viability (CTRL)$. In the cases where $Viability (5-ALA + inhibitor) \sim Viability (5-ALA) \times Viability (inhibitor) / Viability (CTRL)$, the effects were merely additive.

2.5 | Flow cytometry on 5-ALA-derived PpIX

The cells were seeded in 6-well plates (Nunc, 7×10^5 cells per well) and incubated overnight. Ko143 (1 μ M) was added to the cells 1 hour

prior to 5-ALA addition and maintained in the respective groups until the end of the treatment (20 hours after irradiation). The cells were incubated with 5-ALA (0.2 and 0.5 mM T47D and MDA-MB-231, 0.5 mM MCF7, 1 and 0.5 mM T98G, and 2 and 0.5 mM U87) for 4 hours, washed in PBS with calcium and magnesium, and trypsinated with 1 mL TrypLE Express. Next, the cells were resuspended in ice-cold PBS containing calcium and magnesium and measured using an LSRII (BD) flow cytometer with excitation at 405 nm. The fluorescence was registered at the BV650 channel (635 nm cut-on longpass filter and 660/20 nm bandpass filter). All flow cytometry data were analyzed in FlowJo v.10.6.2 (Treestar Inc., Ashland, Oregon).

2.6 | Cell viability assessment

Twenty hours following irradiation, the redox function of all the cell groups was assessed by MTT assay. Briefly, 100 μ L of complete media containing 0.5 mg/mL MTT was added to the cells and incubated for 3 hours at 37°C in a 5% CO₂ incubator. Subsequently, the MTT media were removed, and 100 μ L DMSO was added per well. The plates were shaken for 10 minutes at 350 rpm on a HeidolphTitramax 101 orbital shaker (Heidolph Instruments GmbH & Co. KG, Schwabach, Germany), and an endpoint absorbance measurement was performed at 561 nm using the Tecan spark M10 plate reader (Tecan Group Ltd., Männedorf, Switzerland). Blank values measured in wells with DMSO and no cells were subtracted in all cases.

2.7 | Metabolic assay

T98G, U87, MCF7, MDA-MB-231, and T47D cells were seeded in Seahorse XFe96 96-well plates (3×10^4 cells per well) and incubated overnight in a 37°C, 5% CO₂ humidified atmosphere. The next day, cells were incubated with various concentrations of 5-ALA (0.5 mM T47D and MDA-MB-231, 1 mM MCF7, and 2 mM T98G and U87) for 4 hours and irradiated for 60 seconds for each cell line. One hour prior to metabolic measurement, the cells were placed in Seahorse XF RPMI media, pH 7.4 supplemented with 10 mM glucose, 2 mM sodium pyruvate, and 2 mM glutamine and incubated at 37°C in the absence of CO₂. Cell respiration was monitored through the oxygen consumption rate (OCR) with a Seahorse XFe96 analyzer (Agilent, Santa Clara, California), before (basal rate) and after the sequential addition of respiration modulators (1 μ M oligo, 1 μ M FCCP, and 2 μ M MYXO). The cells were also studied for their glycolytic activity by measuring the rate of extracellular acidification before (basal rate) and after the addition of 1 μ M oligo.

2.8 | Glutathione assay

The total GSH was measured using a glutathione assay kit (Cayman Chemical), following the Tietze recycling assay method²⁶ and according to the manufacturer's instructions. Briefly, the cells were

seeded in 96-well plates (Nunc, 40×10^3 cells/100 μ L media/well) and incubated overnight in their normal media in a 37°C, 5% CO₂ humidified atmosphere. BSO (100 μ M) was added to cells overnight where appropriate. The cells were incubated with 5-ALA (0.2 mM T47D and MDA-MB-231, 0.5 mM MCF7, 1 mM T98G, and 2 mM U87) for 4 hours in the presence and absence of BSO, washed twice in PBS with calcium and magnesium, and finally left in complete medium. One hour following the addition of the medium, the cells were either treated with 0.5% MPA (for the glutathione assay) or 0.1% (v/v) Triton-X 100 aqueous solution for protein quantification. The cells were then kept at –20°C until the GSH assay was performed. Prior to the glutathione assay, 2 μ L of 4 M TEAM was added to each well treated with MPA to reinstate the physiological pH (7.4). Part of the supernatant (50 μ L) from each well was moved to a new 96-well plate and assayed for GSH content. The kinetic absorbance measurement was performed at 405 nm using the Tecan spark M10 plate reader. Blank values measured in wells with no cells were subtracted in all cases.

2.9 | Live cell imaging

The cells were seeded on glass bottom 35-mm dishes (MatTek Corp., Ashland, Massachusetts) containing 2×10^5 cells per dish and incubated overnight. The next day, Ko143 (1 μ M) was added to cells 1 hour prior to 5-ALA addition (1.5 mM ALA, 1 hour or 4 hours) and maintained in the respective groups until imaging. The mitochondrial probe MitoTracker Green FM (100 nm) was added to the cells 15 minutes prior to live cell imaging. The cells were examined with a Zeiss LSM 880 Airyscan microscope (Carl Zeiss MicroImaging GmbH, Jena, Germany), equipped with an Ar Laser Multiline (458/488/514 nm), a DPSS 561 10 (561 nm), a Laser diode 405-30 CW (405 nm), and a HeNe Laser (633 nm). The objective used was a Zeiss plan-Apochromat 63xNA/1.4 oil DICII. Image acquisition, processing and analysis were performed with ZEN 2.3 SP1 basic software (Carl Zeiss), ZEN 2.3 Blue (Carl Zeiss), or Imaris 7.7.2 (Bitplane AG, Zürich, Switzerland). The Pearson's colocalization measurements were performed in ZEN 2.3 Blue with Costes' automatic thresholding.

2.10 | Western blotting

The cells were seeded on 60 mm dishes (Corning, 5×10^5 cells per dish) and incubated overnight in complete media in a 37°C, 5% CO₂ humidified atmosphere. The following day, the cells were incubated with 5-ALA (0.2 mM T47D and MDA-MB-231, 0.5 mM MCF7, 1 mM T98G, and 2 mM U87) for 4 hours, washed twice with PBS with calcium and magnesium, and complete medium was reintroduced. The cells were irradiated (through the petri dish undersides) for different times to achieve LD₅₀ for each cell line (120 seconds T47D, 90 seconds MDA, 60 seconds MCF7, 60 seconds U87, and 120 seconds T98G). Two hours following irradiation, the cells were lysed with RIPA buffer (also containing 10 μ L/mL protease inhibitor

cocktail, 10 $\mu\text{L/mL}$ phosphatase inhibitor cocktail I and II, 10 $\mu\text{L/mL}$ 2 M β -glycerol phosphate, and 5 $\mu\text{L/mL}$ NaF). Finally, lane marker sample buffer was added. The lysates were heated (95°C for 5 minutes), then applied on 18-well 12% TGX gels together with the precision plus protein dual color standard, and subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (120 V, 90 minutes). Subsequently, the proteins were blotted using the Trans-Blot Turbo system onto PVDF transfer membranes (Bio-Rad, Hercules, California). The membranes were washed with tris-buffered saline containing 0.1% Tween 20 (TBST) and blocked in 5% milk for 1 hour at room temperature. The membranes were subsequently incubated with the primary antibodies in 5% milk overnight at 4°C . The following day, the membranes were washed in TBST and incubated with secondary antibodies in TBST for 1.5 hours at room temperature. The membranes were then washed with TBST and incubated with Immobilon Western Chemiluminescent HRP Substrate. Luminescence was measured with a ChemiDoc MP Imaging system (Bio-Rad), and γ -tubulin was used as the housekeeping gene. Data analysis and processing were performed using Image Lab 4.1 and Fiji software.

2.11 | Data analysis and statistics

The data are shown as averages of at least three independent experiments with error bars representing 1 standard deviation (SD). Statistical analysis was assessed using the single tailed *t* test.

3 | RESULTS

The breast cancer cells were more-responsive to 5-ALA PDT than the GBM cells. We initially studied 5-ALA-PDT efficacy on the five selected cell lines through cell viability following treatment. The cell lines demonstrated diverse survival trends, which were found to be 5-ALA concentration- and light dose-dependent (Figure 1A-C).

As shown in Figure 1A-C, T98G (LD_{50} at 2 minutes irradiation and 1 mM 5-ALA) and U87 (LD_{50} at 1 minute irradiation and 2 mM 5-ALA) were the least responsive to 5-ALA-PDT. The MCF7 cells were the least responsive of the breast cancer cell lines (LD_{50} at 1 minute irradiation and 0.5 mM 5-ALA), followed by MDA-MB-231 cells (LD_{50} at 1.5 minutes and 0.2 mM 5-ALA). The T47D cells were the most susceptible to 5-ALA-PDT, as all the cells were killed at 60 seconds of irradiation, following incubation with 1 mM 5-ALA. The collective results from the PDT experiments (Figure 1A-C), confirm that 5-ALA-PDT was profoundly more photocytotoxic in the breast adenocarcinoma cell lines than in the GBM cell lines, with an order of cytotoxicity $\text{T47D} > \text{MDA-MB-231} > \text{MCF7} > \text{U87} > \text{T98G}$. The dark chemical toxicity of 5-ALA and/or the photosensitizer (PpIX) on the cells (without irradiation) is presented in Figure 1D. These data demonstrate that all the 5-ALA concentrations studied were not toxic to the cell lines following 4 hours incubation in the dark. The data in Figure S1 portray the overlap between the emission spectrum of the lamp used for the cell irradiation and the spectral profile of the molar

FIGURE 1 Viability of T98G, U87, MCF7, MDA-MB-231 (MDA), and T47D cells following 5-ALA-PDT vs 5-ALA concentration and time of irradiation. Cells were incubated with 0–2 mM 5-ALA for 4 hours, washed twice in PBS and irradiated with blue light at different time points: A, 60 seconds; B, 90 seconds; C, 120 seconds; and D, 0 s. Cell survival was measured by the MTT assay 20 hours after irradiation. The data represent the average of three individual experiments, while the error bars correspond to 1 standard deviation (SD). E, Representative histograms of 5-ALA-derived PpIX accumulation in T98G, U87, MCF7, MDA-MB-231, and T47D cells. F, Averaged quantitative data for 5-ALA-derived, intracellular PpIX accumulation from three independent experiments. The cells were incubated with 0.5 mM 5-ALA for 4 hours, while the ABCG2 inhibitor Ko143 (1 μM) was introduced to the corresponding treatment groups 1 hour prior to the addition of 5-ALA. Fluorescence was registered by flow cytometry at the BV650 channel (635 nm cut-on longpass filter and 660/20 nm bandpass filter) following excitation at 405 nm. Statistical significance was assessed using the single tailed *t* test: (ns) $P > .05$; (*) $P < .05$; (**) $P < .01$; (***) $P < .001$; (****) $P < .0001$

extinction coefficient of PpIX. The lamp's spectrum had an emission peak at around 405 nm, closely matching the PpIX absorbance Soret band (405 nm).

From these initial PDT experiments, the most suitable light exposure time across all cell lines was determined as 60 seconds, since the viability trends for the various 5-ALA concentrations were well discriminated (Figure 1A).

3.1 | Levels of 5-ALA-derived PpIX and the influence of Ko143

Subsequently, we investigated by flow cytometry the 5-ALA-derived PpIX production following 4 hours incubation with 0.5 mM 5-ALA. The effect of pretreatment with Ko143, a specific inhibitor of the ABCG2 transporter on the accumulation of PpIX, was also studied. The results are presented in Figure 1E, F. Figure 1E shows the relative changes in fluorescence as histograms for each cell line and for the various treatments. The quantitative results are shown in Figure 1F and reveal considerable differences in PpIX fluorescence across the cell lines studied. The fluorescence intensity was highest in T98G (about 4 × that of U87 and MCF7, 2 × that of MDA-MB-31, and 1.6 × that of T47D). The addition of Ko143 increased the PpIX content in all the cell lines apart from T47D. Specifically, following incubation with 0.5 mM 5-ALA (4 hours) the amount of PpIX was 1.8 × ($P < .0001$) higher in the presence of Ko143 for T98G, 1.7 × ($P < .01$) for U87, 1.5 × ($P < .01$) for MCF7, 1.4 × ($P < .05$) for MDA-MB-231, and 1 × ($P > .05$) for T47D (unaffected).

3.2 | Effect of ABCG2 inhibitor Ko143 on 5-ALA-PDT

Following our findings from the PpIX production experiments, we investigated the ABCG2 transporter expression across our chosen cell line panel, incubated with 5-ALA and before or after PDT.

ABCG2 is an ATP-binding cassette (ABC) transporter G2 that has been connected with regulation of the intracellular accumulation of porphyrin derivatives in cancer cells and consequently a modulator of the efficacy of PDT.²⁷ Using Western blotting, we found ABCG2 expression across all the cell lines examined (Figure 2A). T98G exhibited the highest expression levels at about double that of MCF7. U87 cells expressed 4 times less ABCG2 than T98G, whereas MDA-MB-231 and T47D cells expressed about 8 times less ABCG2 than T98G. The ABCG2 expression levels did not change significantly following incubation with 5-ALA in the dark or LD₃₀ 5-ALA-PDT treatment in any of the cell lines. Following these findings, we studied the effect of the specific ABCG2 inhibitor Ko143 (1 and 5 μM, 1 hour before 5-ALA) on 5-ALA-PDT across the cell lines (Figure 2B). We observed a synergistic killing effect between 5-ALA-PDT and Ko143 pre-conditioning in T98G ($p[1 \mu\text{M}] < 0.001$, $p[5 \mu\text{M}] < 0.001$), U87 ($p[1 \mu\text{M}] < 0.0001$, $p[5 \mu\text{M}] < 0.0001$), MCF7 ($p[1 \mu\text{M}] < 0.001$, $p[5 \mu\text{M}] = 0.001$), and MDA-MB-231 ($p[1 \mu\text{M}] = 0.0001$, $p[5 \mu\text{M}] < 0.0001$) cells. On the contrary, the application of 5-ALA-PDT under

FIGURE 2 A, Western blotting analysis of ABCG2 protein levels in T98G, U87, MCF7, MDA-MB-231, and T47D cells after 4 hours treatment with 5-ALA at LD₃₀ concentrations for each cell line and γ-tubulin as the loading control. There are three blotting bands per cell line: first control, second 5-ALA-PDT, and third 5-ALA dark. B, Cell viability assay after 4 hours 5-ALA treatment with or without addition of the ABCG2 inhibitor Ko143 (1 and 5 μM) 1 hour prior to 5-ALA addition and either following PDT or in the dark. The data shown represent the average of three independent experiments, while the error bars correspond to 1 SD. Statistical significance was assessed using the single tailed t test: (ns) $P > .05$; (*) $P < .05$; (**) $P < .01$; (***) $P < .001$; (****) $P < .0001$

Ko143 inhibition yielded no differential cytotoxicity in T47D cells. Incubation with Ko143 alone (1 or 5 μM) was not toxic to any of the cell lines examined.

3.3 | Effect of ABCG2 inhibitor Ko143 on PpIX intracellular accumulation and localization

To further complement these promising results, we investigated the subcellular localization of PpIX in the absence or presence of Ko143. The cells were incubated with 5-ALA in Opti-MEM for 4 hours, followed by the addition of MitoTracker Green FM 15 minutes before confocal imaging. As seen from the representative micrographs in Figure 3, PpIX fluorescence was observed in the cytoplasm of all the cell lines, while in some (T98G, MCF7, and MDA-MB-231), the fluorescence was also notable in the plasma membrane. The addition of Ko143 significantly increased the intracellular PpIX accumulation in the T98G ($P < 0.0001$), U87 ($P < .0001$), MCF7 ($P < .0001$), and MDA-MB-231 ($P < .0001$) cells but conferred no notable changes to PpIX fluorescence intensity in the T47D ($P > 0.05$) cells. Upon addition of Ko143, the fluorescence pattern became more punctate, primarily localizing in the perinuclear area, while the plasma membrane fluorescence component was no longer noticeable. The MitoTracker (green) fluorescence partially overlapped with the PpIX fluorescence as determined by Pearson's correlation, suggesting a slight PpIX colocalization with mitochondria in the T98G (Pearson's coefficient = 0.08, Figure 3A), U87 (Pearson's coefficient = 0.04, Figure 3B), MCF7 (Pearson's coefficient = 0.09, Figure 3C), MDA-MB-231 (Pearson's coefficient = 0.14, Figure 3D), and T47D (Pearson's coefficient = 0.03, Figure 3E) cells. Notably, the corresponding Pearson's coefficients were all negative in the cell groups only incubated with 5-ALA for 4 hours, indicating no mitochondrial localization, except for the T47D cells [-0.001 for T98G, -0.007 for U87, -0.02 for MCF7, -0.02 for MDA-MB-231, and 0.011 for T47D].

3.4 | Role of HO-1 in 5-ALA-PDT

HO-1, which is induced by oxidative cellular stress,²⁸ is known to catalyze heme degradation, hence mediating PDT resistance.²⁹ HO-1 expression was hardly noticeable in the non-treated (control) T98G cells (C, 100 relative units [RU], Figure 4A), but there was a dramatic increase as a consequence of 5-ALA administration both after light irradiation (L, $\times 10$, $P = .005$) and under dark conditions (D, $\times 23$, $P = .001$). HO-1 expression in the U87 cells was nearly equal in all the groups examined (C 417[L 452][D 467 RU). HO-1 expression was lower in the MCF7 (C 45[L 115][D 101 RU), MDA-MB-231 (C 28[L 51][D 43 RU), and T47D (C 35[L 90][D 77 RU) cells for the conditions under investigation.

To determine the effect of HO-1 on 5-ALA-PDT, the cells were incubated with OB24, a HO-1 inhibitor, either overnight or 1 hour prior to co-incubation with 5-ALA (Figure 4B). Interestingly, a synergistic effect with 5-ALA-PDT was only registered in the T98G cells

FIGURE 3 Intracellular localization of the 5-ALA-derived photosensitizer PpIX for the various cell lines: A, T98G; B, U87; C, MCF7; D, MDA-MB-231; and E, T47D. The cells were incubated with 1.5 mM 5-ALA for 4 hours with (odd rows) or without (even rows) pretreatment with Ko143 (1 μM). The cells were incubated with 100 nm MitoTracker Green 15 minutes prior to imaging. PpIX fluorescence is represented in red (left column) and MitoTracker fluorescence in green (second column). An overlay of PpIX fluorescence with MitoTracker fluorescence is shown in the overlap panel (third column) where colocalization can be seen in yellow. Pure colocalization between PpIX and MitoTracker is shown in the fourth panel in white

that were preincubated overnight with 20 μM OB24 ($P = .007$) prior to 5-ALA treatment. There was no differential cytotoxicity either in

FIGURE 4 A, Western blotting analysis of HO-1 protein levels in T98G, U87, MCF7, MDA-MB-231, and T47D cells after 4 hours treatment with 5-ALA at LD₃₀ concentrations for each cell line and γ-tubulin as the loading control. There are three blotting bands per cell line: first control, second 5-ALA-PDT, and third 5-ALA dark. b) Cell viability assay after 4 hours 5-ALA treatment with or without preincubation with the HO-1 inhibitor OB24 overnight or 1 hour prior to 5-ALA addition, either following PDT or in the dark. The data shown represent the average of three independent experiments, while the error bars correspond to 1 SD. Statistical significance was assessed using the single tailed t test: (ns) $P > .05$; (*) $P < .05$; (**) $P < .01$; (***) $P < .001$; (****) $P < .0001$

T98G for 1 hour preincubation with OB24 or in any other of the cell lines examined for both overnight and 1 hour OB24 preincubation ($P > .05$), which is in agreement with our protein expression studies and suggests that HO-1 induction by 5-ALA protects T98G cells against 5-ALA-PDT's cytotoxic effect.

3.5 | FECH expression and the effect of its inhibition on 5-ALA-PDT

Since FECH can potentially decrease the PpIX content of the cells and consequently their responsiveness to 5-ALA-PDT, FECH expression was studied across the cell lines, and profound differences were found in the expression levels (Figure 5A). T98G showed the highest FECH expression (C 100|D 106|L 93 RU), MCF7 came next with C 70|L 59|D 56 RU, comparable to T47D (C 58|L 54|D 54 RU), followed by U87 (C 28|L 23|D 20 RU), and MDA-MB-231 (C 9|L 7|D 7 RU). There were no notable differences in the FECH expression between the control cells and the 5-ALA treated cells either in the dark or following PDT, for any of the cell lines ($P > .05$), except between the T98G control and 5-ALA-PDT groups ($P < .05$), which was, however, due to the very low SDs.

To investigate the effect of FECH on 5-ALA-derived-PpIX-PDT, the cells were incubated either overnight or 1 hour prior to PDT

treatment with the FECH inhibitor Noc-18. The results are shown in Figure 5B. There was a synergistic effect between 5-ALA-PDT and co-incubation with Noc-18 in the T98G (overnight $P < .01$, 1 hour $P = .01$), MCF7 (overnight $P < .0001$, 1 hour $P < .001$) and T47D (overnight $P < .0001$, 1 hour $P < .0001$) cells, while the effects on the U87 cells and MDA-MB-231 were merely additive. Incubation with Noc-18 alone was subtoxic when administered 1 hour prior to 5-ALA treatment (0-12%), but moderate cytotoxicity (11-35%) was observed when cells were incubated overnight with Noc-18.

The effects of enzymatic inhibitors Ko143, Noc 18, OB24 and BSO on the 5-ALA-PDT outcomes has been summarized for all cell lines in Table 1.

3.6 | Role of intracellular anti-oxidants on 5-ALA-PDT

The enzymes from the superoxide dismutase (SOD) family are essential to detoxify the ROS generated by the cells³⁰ by catalyzing the dismutation of superoxide radicals into H₂O₂. We were interested to see the influence of the level of expression of superoxide dismutases (Zn, Cu SOD [SOD1] and Mn SOD [SOD2]) on 5-ALA-PDT and vice versa in the cell lines examined. Figure 6A shows that the T98G cells exhibited the highest expression of cytosolic SOD1 (C 100|L 109|D

FIGURE 5 A, Western blotting analysis of FECH levels in T98G, U87, MCF7, MDA-MB-231, and T47D cells after 4 hours treatment with 5-ALA at LD₃₀ concentrations for each cell line and γ -tubulin as the loading control. There are three blotting bands per cell line: first control, second 5-ALA-PDT, and third 5-ALA dark. B, Cell viability after 4 hours 5-ALA treatment with or without addition of the FECH inhibitor Noc-18 overnight or 1 hour prior to 5-ALA and either following PDT or in the dark. The data shown represent the average of three independent experiments, while the error bars correspond to 1 SD. Statistical significance was assessed using the single tailed *t* test: (ns) *P* > .05; (*) *P* < .05; (**) *P* < .01; (***) *P* < .001; (****) *P* < .0001

TABLE 1 Enhancement of 5-ALA-PDT cytotoxicity across the cell lines by the inhibitors used in the study

Cell lines	Protein/Inhibitor Enhancement of 5-ALA-PDT cytotoxicity (%)			
	ABCG2/Ko143	HO-1/OB24	FECH/Noc-18	γ GCS/BSO
T98G	58 ± 3	33 ± 18	44 ± 5	52 ± 14
U87	78 ± 9	7 ± 10	25 ± 6	15 ± 7
MCF7	45 ± 14	4 ± 18	50 ± 10	9 ± 8
MDA	63 ± 8	4 ± 10	36 ± 9	65 ± 11
T47D	1 ± 10	-4 ± 9	64 ± 9	16 ± 14

141 RU), comparable to T47D (C 104|L 107|D 106 RU) and then MCF7 (C 90|L 67|D 87 RU), while the U87 (C 43|L 33|D 43) and MDA-MB-231 (C 27|L 28|D 30 RU) cells had the lowest expression.

With respect to SOD2 (mitochondrial MnSOD, Figure 6B), the U87 cells had the highest expression of SOD2 (C 467|L 513|D 443 RU) compared to the other cell lines, namely T98G (C 100|L 126|D 115 RU), MCF7 (C 70|L 59|D 53 RU), MDA-MB-231 (C 70|L 84|D 79 RU), and T47D (C 58|L 56|D 47 RU) (Figure 6B). These data demonstrate that there were no statistically significant differences in SOD expression between the control, dark PpIX control, and PDT treatment groups in all cell lines.

We assessed the impact of glutathione peroxidase as an antioxidant defense mechanism against 5-ALA-PDT and hence a modulator

of the PDT outcomes. The U87 cells had the highest expression of GPX4 (C 319|L 215|D 228 RU, Figure 6C). The T47D (C 230|L 166|D 178 RU) and MCF7 (C 157|L 107|D 135 RU) cells followed in the level of expression, while the T98G (C 100|L 29|D 121 RU) and MDA-MB-231 (C 88|L 75|D 114 RU) cells exhibited the lowest GPX4 expression. These results further suggest that the decrease in GPX4 expression between the 5-ALA-dark and 5-ALA-PDT groups was only significant in the T98G cells (*P* < .05). All the other differences in expression across the treatment groups were statistically non-significant across all the cell lines.

In relation to the level of expression of GSTP1 across the cell lines (Figure 6D), there were no major differences in GSTP1 expression between the various treatment groups. The T98G cells exhibited the

FIGURE 6 Western blotting analysis of protein levels in T98G, U87, MCF7, MDA-MB-231, and T47D cells after 4 hours treatment with 5-ALA at LD30 concentrations for each cell line. γ -tubulin was used as the loading control. The proteins examined were A, SOD1; B, SOD2; C, GSTP1; and D, GPX4. There are three blotting bands per cell line: first control, second 5-ALA-PDT, and third 5-ALA dark. E, Total GSH measurement in T98G, U87, MCF7, MDA-MB-231, and T47D cells treated overnight with 100 μ M BSO. F, Cell viability assay following 4 hours 5-ALA treatment with or without overnight preincubation with 100 μ M BSO, either following 5-ALA-PDT or G, in the dark. The data shown represent the average of three independent experiments, while the error bars correspond to 1 SD. Statistical significance was assessed using the single tailed *t* test: (ns) $P > .05$; (*) $P < .05$; (**) $P < .01$; (***) $P < .001$; (****) $P < .0001$

highest levels of GSTP1 expression (C 100|L 99|D 104 RU), followed by the U87 (C 22|L 16|D 24 RU) and MDA-MB-231 (C 7|L 8|D 8 RU) cells. The GSTP1 expression levels for the MCF7 (C 1|L 0.9|D 1 RU) and T47D (C 0.3|L 0.3|D 0.3 RU) cells were barely detectable.

3.7 | Effect of intracellular GSH levels on 5-ALA-PDT

The total glutathione levels were measured in dark conditions with and without overnight BSO (an inhibitor of the *de-novo* intracellular GSH synthesis³¹) pretreatment and with or without 5-ALA incubation (4 hours)

on all the cell lines examined. The overnight treatment with 100 μ M BSO profoundly depleted intracellular GSH in all the cell lines (~100% reduction, Figure 6E), without significantly affecting cell survival in the absence of oxidative stress (Figure 6G), with the exception of MCF7, where the reduction in GSH levels was approximately 90%. Following 5-ALA-PDT, as shown in Figure 6F, the treatment of T98G and MDA-MB-231 cells with BSO had a profound synergistic effect on 5-ALA-PDT cytotoxicity (T98G: 70% survival $\rightarrow$ 18% survival, $P < .001$, MDA-MB-231: 72% survival $\rightarrow$ 7% survival, $P < .01$). Correspondingly, there was a 10% difference in cell death in the T47D ($P < .01$) and MCF7 ($P < .01$) cells, while there was no statistically significant effect of GSH depletion by BSO on the outcome of 5-ALA-PDT in the U87 cells ($P > .05$).

FIGURE 7 Metabolic studies 1 hour after 5-ALA-PDT, measured by the Seahorse XFe96 analyzer. 30 000 cells were seeded in 96 well plates. T98G, U87, MCF7, MDA-MB-231, and T47D cells were treated with media only or various concentrations of 5-ALA (see methods section) for 4 hours and subsequently irradiated for 60 seconds. A, Basal oxygen consumption rate (OCR) for all cell lines and treatment groups. B, Maximal respiratory capacity as represented by the OCR following the addition of the protonophore uncoupler FCCP. C, Basal ECAR. D, Glycolytic capacity as represented by the ECAR following the addition of the F-ATPase inhibitor oligomycin. The data shown represent the average of three independent experiments, while the error bars correspond to 1 SD. Statistical significance was assessed using the single tailed *t* test: (ns) $P > .05$; (*) $P < .05$; (**) $P < .01$; (***) $P < .001$; (****) $P < .0001$

3.8 | Effect of 5-ALA-PDT on cell metabolism

To better understand the effect of 5-ALA-PDT on cell bioenergetics, we performed metabolic analysis using a Seahorse metabolic analyzer (XFe96). Under dark conditions (Figure 7A), the basal (media only) OCR of MCF7 (107 RU) was almost two times higher than that of the T98G (52 RU) and T47D (47 RU) cells, 5–6 times higher than U87 (16 RU), and 4 times higher than the MDA-MB-231 (23 RU) cells. The addition of 5-ALA to cells without irradiation only led to a substantial decrease in respiration in the T47D cells (47 → 26 RU, $P = .009$). The maximal respiratory capacity, upon addition of the protonophore FCCP, followed a similar trend, only this time apart from T47D (37 → 18 RU, $P = .007$) there was a marked difference between the dark controls and 5-ALA dark also in T98G (120 → 78 RU, $P = .0002$) (Figure 7B). Following light irradiation in the absence of 5-ALA, the OCR levels were in agreement with their dark counterparts. However, this was not the case for the 5-ALA-PDT treated cells, where OCR measurements 1 hour following irradiation demonstrated a profound drop in all the cell lines. Specifically, the basal respiratory rates (as reflected through the OCRs) exhibited reductions of 90% ($P < .0001$) for T98G, approximately 45% ($P < .01$) for U87, approximately 90% ($P < .0001$) for MCF7, approximately 75% ($P < .001$) for

MDA-MB-231, and approximately 94% ($P < .0001$) for the T47D cells. In all cases, the cells were not responsive to the stimuli of oligo, FCCP, or MYXO 1 hour following 5-ALA-PDT (Supplementary Figure 3).

With regard to the glycolytic profiles of the five cell lines, the results are shown in Figure 7C. The control (media only) T98G cells were found to be the most glycolytic cell line with an extracellular acidification rate (ECAR) (52 RU) almost two times higher than the U87 (33 RU, $P \leq .001$), MCF7 (32 RU, $P \leq .01$), MDA-MB-231 (23 RU, $P < .0001$), and T47D (30 RU, $P \leq .001$) cells. No significant drop in the extracellular acidification rate was noted after 4 hours incubation with 5-ALA in the T98G ($P > .05$), U87 ($P > .05$), MDA-MB-231 ($P > .05$), and MCF7 ($P > .05$) cells. However, there was a measurable drop in the glycolytic activity of the T47D cells (30 → 21 RU, $P < .01$). After irradiation, we observed a dramatic decrease in the glycolytic rates (Figure 7D) 1 hour following 5-ALA-PDT: the decline was approximately 98% ($P < .0001$) for T98G, approximately 62% ($P < .0001$) for U87, approximately 88% ($P < .0001$) for MCF7, approximately 62% ($P < .0001$) for MDA-MB-231 and approximately 94% ($P < .0001$) for the T47D cells.

An overview of the biochemical interactions and our proposed interventions to enhance the outcome of 5-ALA-PDT, as presented herein, is schematically summarized in Figure 8.

FIGURE 8 Overview of the basic biochemical processes and regulatory interventions enhancing the outcome of 5-ALA-PDT. The production of PpIX, the main photosensitizer in 5-ALA-PDT, is enhanced by the administration of exogenous 5-ALA. The scheme above illustrates the various biochemical interventions applied in the present work aiming to enhance the 5-ALA-PDT outcome. BSO is shown to inhibit GSH for reasons of simplicity, whereas in reality BSO inhibits glutathione synthetase, resulting in the suppression of the *de-novo* production of GSH. GSH is a substrate for both the antioxidant activity of glutathione peroxidase (GPX) and the expulsion of xenobiotics via the GS-X pump. Ko143 is a specific inhibitor of the ABCG2 transporters which are responsible for the expulsion of PpIX from cells and cell mitochondria. NOC-18 is an inhibitor of the enzyme ferrochelatase (FECH) which is responsible for the introduction of a Fe^{2+} into the PpIX structure transforming it into the PDT-inert heme. OB-24 is an inhibitor of heme oxygenase (HO-1). HO-1 cleaves the heme pushing the PpIX-heme equilibrium towards the reduction of PpIX

4 | DISCUSSION

In the present study, two GBM cell lines (T98G and U87) and three breast adenocarcinoma cell lines (MCF7, MDA-MB-231, and T47D) exhibited different 5-ALA-PDT-related cytotoxicities, with the GBM cell lines being in general more resistant (practically unresponsive). There was also a differential PDT response between the GBM cells, with T98G being more resistant than the U87 cells, which also exhibited a very limited response to 5-ALA-PDT. With regard to the breast cancer cell lines, MCF7 was more resistant than MDA-MB-231, while T47D presented as the most responsive cell line to 5-ALA-PDT. This confirms previous observations regarding the higher vulnerability of MDA-MB-231 in comparison to MCF7.^{31,32} In general, the sensitivity of cell lines to PDT was found to be variant, due to phenotypic and genotypic differences as also reported elsewhere.³³ Furthermore, there was no direct correlation between the intracellular amount of PpIX and 5-ALA-PDT cytotoxicity, across the cell lines (Figure S4). We further investigated these differential cytotoxicities delineated above for three reasons: (a) to understand the underlying mechanisms for these dissimilar responses; (b) to identify biomarkers for the prediction of PDT treatment outcome; and (c) to implement methods for enhancing

the tumoricidal effect of 5-ALA-PDT based on specific tumor characteristics.

Our experimental observations warranted the investigation of possible ways to sensitize the GBM cells to 5-ALA-PDT, especially since it is well established that 5-ALA-derived PpIX fluorescence is used to precisely guide the surgical resection of GBM and bladder lesions due to its high specificity.^{16,17,19,34-37} Efficient GBM intraoperative 5-ALA-PDT could significantly prolong patients' disease-free survival, as clinical studies have revealed that the recurrence of gliomas is frequently due to the residual cancer cells left behind after treatment.⁶ Apart from the apparent benefits for GBM cancers, the present work could also contribute by increasing the susceptibility of already responsive tumors, as illustrated through the breast cancer cell study results, auguring a more efficient treatment of a wide range of cancers in the clinical setting.

Even though the T98G cells exhibited much higher PpIX production than all the other cell lines, following administration of the same amount of 5-ALA, they presented the lowest cytotoxic response to 5-ALA-PDT (Figure 1A-D). These contradictory results were later reconciled by our finding that T98G exhibited the highest expression of ABCG2 transporters. These are known to play an essential role in the

regulation of the intracellular accumulation of porphyrins in cancer cells by pumping the porphyrins out of the cells and therefore reducing the efficacy of PDT.¹⁸ Indeed, preincubation of T98Gs with the ABCG2 inhibitor Ko143 led to a approximately 2-fold increase in the intracellular levels of PpIX. Consequently, the T98G cells were profoundly sensitized to 5-ALA-PDT, following preincubation with Ko143 (Figure 2B). Apart from T98Gs, the U87, and to a lesser extent MDA-MB-231, cells were found to be significantly responsive to 5-ALA-PDT following ABCG2 inhibition by Ko143, despite the fact that these cell lines (U87 and MDA-MB-231) exhibited disproportionately low expression of ABCG2s. Conversely, the MCF7 cells demonstrated a moderate PDT response to ABCG2 inhibition by Ko143, despite their levels of ABCG2 expression being higher than these of U87 and MDA-MB-231 cells. No changes in the 5-ALA-PDT response was recorded for the T47D cells following ABCG2-inhibition, even though T47Ds also expressed ABCG2 at a level comparable to MDA-MB-231. This can to an extent be explained by the micrographs shown in Figure 3 and Figure S2. It is evident from these representative images that different cell lines responded differently to Ko143 pretreatment. The T98G, MDA-MB-231, U87, and MCF7 cells exhibited enhanced mitochondrial retention of PpIX 4 hours (or even 1 hour) after Ko143 administration, while T47D showed no marked difference in the mitochondrial retention of PpIX with and without Ko143. Furthermore, the 5-ALA-PDT insensitivity to Ko143 also indicates that T47D recruited defense mechanisms against PDT other than ABCG2. These include FECH, as our data (Figure 5B) showed that inhibition of the latter by Noc-18 induced a profound T47D cytotoxicity in response to 5-ALA-PDT. FECH is important for the production of heme since it catalyzes iron chelation by PpIX, as delineated above, but at the same time it reduces the level of PpIX available for PDT. Other mechanisms of T47D resistance to 5-ALA-PDT, different from those of T98G could include the high membrane-resident GPX4 levels, which despite the relatively low levels of T47D intracellular GSH, may help catalyze the detoxification of lipid hydroperoxides, products of 5-ALA-PDT induced lipid peroxidation. However, even significantly reducing the GSH levels in T47Ds by overnight pretreatment with BSO does not substantially affect the PDT response, suggesting that the main defense of T47D to PDT may indeed be FECH. This lack of a multifaceted defense may also be the reason why T47D cells are most responsive to 5-ALA-PDT among the cell lines investigated. Other factors that are relatively enhanced in T47D cells are the antioxidant enzymes SOD1 and SOD2. By contrast, T47D seem to be relatively deficient in GSTP1 that conjugates xenobiotics with GSH exocytosis by the GS-X pump.³⁸ GSTP1 is overexpressed in a number of tumor cell lines resistant to several anticancer drugs that confer a cytoprotective role.³⁹ In addition, T47Ds seem quite deficient in heme oxygenase 1 (HO-1), a member of the heat shock protein family that has been shown to alleviate oxidative stress in tumor cells.^{28,40} HO-1 can cleave heme, thus releasing iron, which is then available to bind even more PpIX at the expense of 5-ALA-PDT induced photocytotoxicity.

T98Gs, however, demonstrated an adaptive HO-1 expression, which increased with 5-ALA administration and decreased following

5-ALA-PDT. Indeed, upon HO-1 inhibition by OB24, T98G was the only cell line that exhibited enhancement of 5-ALA-PDT induced photocytotoxicity. In contrast, U87s, which also exhibited a moderate HO-1 expression (not dependent upon 5-ALA administration), did not exhibit augmented PDT cytotoxicity following OB24 pretreatment. The other cell lines only expressed minimal HO-1 levels and no notable responses to 5-ALA-PDT upon OB24 pre-conditioning. T98G, much like T47D, seemed to be responsive to FECH inhibition, as did the MCF7 cells, and their corresponding increases in PDT-induced cell death were analogous to their reciprocal FECH levels. The U87 and MDA-MB-231 cells demonstrated lower FECH levels and were hence found less responsive to FECH inhibition with regard to 5-ALA-PDT.

With respect to antioxidant defenses, T98Gs exhibited high GSTP1 and SOD-1 levels and the highest GSH levels on cell lines. In addition, together with MDA-MB-231, they were the most responsive to GSH depletion by BSO with regard to the 5-ALA-PDT outcome. MDA-MB-231 cells were found in a previous study to mostly utilize GSTP1 with GSH to detoxify hypericin³¹ and mitigate its PDT action, rather than GPX4, of which they express low levels. U87 had the highest level of SOD2, MnSOD, which is the mitochondrial SOD, and also of GPX4. These two findings indicate the high importance of cell mitochondria protection for U87, despite the fact that they exhibited the lowest OCR rates and the highest ECAR/OCR (glycolysis/respiration) ratio (approximately 2), implying a predominantly Warburg-type cellular metabolism.

In summary, for T98G cells, adjuvant inhibition of ABCG2 alone led to a profound PDT effect, and this could also be achieved by inhibition of FECH and HO-1 as well as intracellular GSH depletion. In the case of U87 cells, only inhibition of the ABCG2 was found to provide a synergistic adjuvant effect to 5-ALA-PDT (see also Table 1).

Even though the breast cancer cell lines proved to be more susceptible to 5-ALA-PDT, in the case of MCF7 cells, inhibition of ABCG2 or FECH greatly enhanced 5-ALA-PDT cytotoxicity, while for MDA-MB-231s, ABCG2 inhibition, and intracellular GSH depletion conferred the most profound synergies to 5-ALA-PDT. As also delineated earlier, the only substantial synergism to ALA-PDT in the already responsive T47D cells was afforded by FECH inhibition. Turning these observations (also summarized in Table 1) around, the levels of expression of ABCG2, FECH primarily, but also of HO-1, can be used as strategic biomarkers for the prediction of the treatment outcome of 5-ALA. SOD2 can also be used as an auxiliary predictive marker, since 5-ALA-PDT is relevant to cell mitochondria. Finally, the triad GPX4 and GSTP1 expression vs intracellular GSH levels that have been shown to be relevant markers for PDT³¹ can be also used for 5-ALA-PDT. However, ABCG2, FECH, and HO-1 are the main indicators of 5-ALA-PDT treatment outcome, and their inhibitors can be used in the clinic to elicit far more effective 5-ALA-PDT treatments.

There are numerous other inhibitors of ABCG2 besides Ko143.⁴¹ Many have been used in the clinic in an effort to increase the bioavailability of chemotherapeutics through barriers like the blood-brain barrier, but they also maintain a high intracellular level of these chemotherapeutics.⁴¹ Apart from 5-ALA-PDT, ABCG2 inhibition has

been found to enhance PDT for various other PSs,^{42–44} while the sensitization of a panel of TNBC cells to 5-ALA-PDT following ABCG2 inhibition has been shown elsewhere.²⁵ Correspondingly, the inhibition of FECH has been previously shown to enhance 5-ALA-PDT in PC3 prostate cancer,⁴⁵ while genetically silencing FECH has been proved to synergize with 5-ALA-PDT on the glioma cells SNB19.⁶ Finally, the inhibition of HO-1 in a melanoma cell model increased their responsiveness to 5-ALA-PDT.²⁸ Apart from these sporadic and isolated reports, however, and up to the present work, a systematic study on the various factors that could specifically synergize with 5-ALA-PDT to make it a more potent cancer treatment and applicable to a wider range of cancer tissues, overcoming their resistance, has been lacking.

Our study on the effect of 5-ALA-PDT on cell metabolism shows that there was a detrimental effect on the OCR for all cell lines with the exception of U87 cells, which did not suffer a notable loss on their basal respiration but underwent a sizable decrease in their maximal respiratory capacity. This is well in line with the observation of the fortified mitochondrial defenses of U87 (see above). It is quite possible, since PpIX is produced in mitochondria, that 5-ALA-PDT caused damage to one of the electron transport chain components, and severely impaired the ETC complex III much like hypericin-PDT.^{32,46} In terms of glycolysis, the U87 and MDA-MB231 cells suffered the least damage to their glycolytic machinery among the cell lines, and since they are both metabolically more “Warburg” than “Pasteur” type of cells in terms of metabolism, perhaps clinical suppression of glycolysis prior to and even post 5-ALA-PDT, either pharmacologically or by regulated fasting, could lead to profound enhancement of the PDT treatment outcome.

The results of our study could provide a paradigm shift for 5-ALA-PDT as an anticancer treatment for two main reasons: (a) we have proposed and validated specific biomarkers that can potentially predict the outcome of 5-ALA-PDT treatments in the clinic based on the analysis of tumor biopsies; and (b) at the same time our results can be used to set up personalized 5-ALA-PDT treatments with improved efficacy based on tumor profiling with respect to our proposed biomarkers.

To translate our results into clinical practice, a two-phase plan should be adopted. Initially, biopsies derived from the tumors of patients to be treated with 5-ALA-PDT should be derived and analyzed with regard to their expression of ABCG2, FECH, and/or HO-1 against well-known negative and positive cell lines for these markers. During that phase, the results of 5-ALA-PDT should be reported and correlated with the lack or abundance of the suggested biomarkers. In the second phase, an example of a proposed protocol would be to exploit the genomic analysis of the biopsy and apply clinically tested and approved inhibitors (eg, febuxostat or elacridar for ABCG2) either alone or in combination, depending on how many of the predictive biomarkers come out positive prior to PDT in order to curb the main tumor resistance with respect to the treatment.

ACKNOWLEDGEMENTS

The present work was financially supported by the European Union's Horizon 2020 framework program for research and innovation under

grant agreement No 712921 and by the South-Eastern Norway Regional Health Authority (Helse Sør Øst, project no. 2017116).

CONFLICT OF INTEREST

The authors declare no potential conflicts of interest.

AUTHOR CONTRIBUTIONS

Maria Mastrangelopoulou Data curation-Lead, Formal analysis-Equal, Investigation-Lead, Validation-Equal, Writing-original draft-Lead.

Mantas Grigalavicius Data curation-Equal, Formal analysis-Equal, Investigation-Equal, Methodology-Equal, Validation-Equal, Writing-review and editing-Equal.

Tine Henriksen-Raabe Data curation-Equal, Investigation-Equal, Validation-Equal.

Ellen Skarpen Formal analysis-Equal, Investigation-Equal, Writing-review and editing-Equal.

Petras Juzenas Investigation-Supporting, Writing-review and editing-Supporting.

Qian Peng Investigation-Supporting, Writing-review and editing-Supporting.

Kristian Berg Formal analysis-Equal, Funding acquisition-Equal, Project administration-Supporting, Resources-Equal, Writing-review and editing-Equal.

Theodossis A. Theodossiou Conceptualization-Lead, Formal analysis-Lead, Funding acquisition-Lead, Methodology-Lead, Project administration-Lead, Resources-Lead, Supervision-Lead, Investigation-Supporting, Validation-Equal, Visualization-Lead, Writing-original draft-Lead, Writing-review and editing-Equal.

ETHICS STATEMENT

Only studies on commercially available cell lines were performed in the present work and hence no ethical considerations apply.

DATA AVAILABILITY STATEMENT

The data supporting the findings of the current study will be stored in an OUS repository and be made available on reasonable request.

ORCID

Theodossis A. Theodossiou <https://orcid.org/0000-0002-1982-8372>

REFERENCES

1. Agostinis P, Berg K, Cengel KA, et al. Photodynamic therapy of cancer: an update. *CA Cancer J Clin*. 2011;61:250–281. <https://doi.org/10.3322/caac.20114>.
2. Dolmans DE, Fukumura D, Jain RK. Photodynamic therapy for cancer. *Nat Rev Cancer*. 2003;3:380–387. <https://doi.org/10.1038/nrc1071>.
3. Dougherty TJ, Gomer CJ, Henderson BW, et al. Photodynamic therapy. *J Natl Cancer Inst*. 1998;90:889–905. <https://doi.org/10.1093/jnci/90.12.889>.
4. Mastrangelopoulou M, Grigalavicius M, Berg K, Menard M, Theodossiou TA. Cytotoxic and photocytotoxic effects of cercosporin on human tumor cell lines. *Photochem Photobiol*. 2019;95:387–396. <https://doi.org/10.1111/php.12997>.
5. Peng Q, Warloe T, Berg K, et al. 5-Aminolevulinic acid-based photodynamic therapy. Clinical research and future challenges. *Cancer*.

- 1997;79:2282-2308. [https://doi.org/10.1002/\(sici\)1097-0142\(19970615\)79:12<2282::aid-cncr2>3.0.co;2-o](https://doi.org/10.1002/(sici)1097-0142(19970615)79:12<2282::aid-cncr2>3.0.co;2-o).
6. Teng L, Nakada M, Zhao SG, et al. Silencing of ferrochelatase enhances 5-aminolevulinic acid-based fluorescence and photodynamic therapy efficacy. *Br J Cancer*. 2011;104:798-807. <https://doi.org/10.1038/bjc.2011.12>.
7. el-Sharabasy MM, el-Waseef AM, Hafez MM, Salim SA. Porphyrin metabolism in some malignant diseases. *Br J Cancer*. 1992;65:409-412. <https://doi.org/10.1038/bjc.1992.83>.
8. Kondo M, Hirota N, Takaoka T, Kajiwaru M. Heme-biosynthetic enzyme activities and porphyrin accumulation in normal liver and hepatoma cell lines of rat. *Cell Biol Toxicol*. 1993;9:95-105. <https://doi.org/10.1007/bf00755143>.
9. Rubino GF, Rasetti L. Porphyrin metabolism in human neoplastic tissues. *Panminerva Med*. 1966;8:290-292.
10. Schoenfeld N, Epstein O, Lahav M, Mamet R, Shaklai M, Atsmon A. The heme biosynthetic pathway in lymphocytes of patients with malignant lymphoproliferative disorders. *Cancer Lett*. 1988;43:43-48. [https://doi.org/10.1016/0304-3835\(88\)90211-x](https://doi.org/10.1016/0304-3835(88)90211-x).
11. Van Hillegersberg R, Van den Berg JW, Kort WJ, Terpstra OT, Wilson JH. Selective accumulation of endogenously produced porphyrins in a liver metastasis model in rats. *Gastroenterology*. 1992;103:647-651. [https://doi.org/10.1016/0016-5085\(92\)90860-2](https://doi.org/10.1016/0016-5085(92)90860-2).
12. Rittenhouse-Diakun K, Leengoed HVAN, Morgan J, et al. The role of transferrin receptor (CD71) in photodynamic therapy of activated and malignant lymphocytes using the heme precursor delta-aminolevulinic acid (ALA). *Photochem Photobiol*. 1995;61:523-528. <https://doi.org/10.1111/j.1751-1097.1995.tb02356.x>.
13. Kennedy JC, Pottier RH, Pross DC. Photodynamic therapy with endogenous protoporphyrin IX: basic principles and present clinical experience. *J Photochem Photobiol B Biol*. 1990;6:143-148. [https://doi.org/10.1016/1011-1344\(90\)85083-9](https://doi.org/10.1016/1011-1344(90)85083-9).
14. Inoue K, Fukuhara H, Shimamoto T, et al. Comparison between intravesical and oral administration of 5-aminolevulinic acid in the clinical benefit of photodynamic diagnosis for nonmuscle invasive bladder cancer. *Cancer*. 2012;118:1062-1074. <https://doi.org/10.1002/cncr.26378>.
15. Fukuhara H, Inoue K, Satake H, et al. Photodynamic diagnosis of positive margin during radical prostatectomy: preliminary experience with 5-aminolevulinic acid. *Int J Urol*. 2011;18:585-591. <https://doi.org/10.1111/j.1442-2042.2011.02789.x>.
16. Stummer W, Novotny A, Stepp H, Goetz C, Bise K, Reulen HJ. Fluorescence-guided resection of glioblastoma multiforme by using 5-aminolevulinic acid-induced porphyrins: a prospective study in 52 consecutive patients. *J Neurosurg*. 2000;93:1003-1013. <https://doi.org/10.3171/jns.2000.93.6.1003>.
17. Stummer W, Stocker S, Wagner S, et al. Intraoperative detection of malignant gliomas by 5-aminolevulinic acid-induced porphyrin fluorescence. *Neurosurgery*. 1998;42:518-525; discussion 525-516. <https://doi.org/10.1097/00006123-199803000-00017>.
18. Ishikawa T, Kajimoto Y, Inoue Y, Ikegami Y, Kuroiwa T. Critical role of ABCG2 in ALA-photodynamic diagnosis and therapy of human brain tumor. *Adv Cancer Res*. 2015;125:197-216. <https://doi.org/10.1016/bs.acr.2014.11.008>.
19. Tetard MC, Vermandel M, Mordon S, Lejeune JP, Reyns N. Experimental use of photodynamic therapy in high grade gliomas: a review focused on 5-aminolevulinic acid. *Photodiagnosis Photodyn Ther*. 2014;11:319-330. <https://doi.org/10.1016/j.pdpdt.2014.04.004>.
20. Faraz S, Pannullo S, Rosenblum M, Smith A, Wernicke AG. Long-term survival in a patient with glioblastoma on antipsychotic therapy for schizophrenia: a case report and literature review. *Ther Adv Med Oncol*. 2016;8:421-428. <https://doi.org/10.1177/1758834016659791>.
21. Hefti M, von Campe G, Moschopoulos M, Siegner A, Looser H, Landolt H. 5-aminolevulinic acid induced protoporphyrin IX fluorescence in high-grade glioma surgery: a one-year experience at a single institution. *Swiss Med Weekly*. 2008;138:180-185.
22. Stummer W, Reulen H-J, Meinel T, et al. Extent of resection and survival in glioblastoma multiforme: identification of and adjustment for bias. *Neurosurgery*. 2008;62:564-576; discussion 564-576. <https://doi.org/10.1227/01.neu.0000317304.31579.17>.
23. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin*. 2020;70:7-30. <https://doi.org/10.3322/caac.21590>.
24. Aniogo EC, Plackal Adimuriyl George B, Abrahamse H. The role of photodynamic therapy on multidrug resistant breast cancer. *Cancer Cell Int*. 2019;19:91. <https://doi.org/10.1186/s12935-019-0815-0>.
25. Palasuberniam P, Yang X, Kraus D, Jones P, Myers KA, Chen B. ABCG2 transporter inhibitor restores the sensitivity of triple negative breast cancer cells to aminolevulinic acid-mediated photodynamic therapy. *Sci Rep*. 2015;5:13298. <https://doi.org/10.1038/srep13298>.
26. Tietze F. Enzymic method for quantitative determination of nanogram amounts of total and oxidized glutathione: applications to mammalian blood and other tissues. *Anal Biochem*. 1969;27:502-522. [https://doi.org/10.1016/0003-2697\(69\)90064-5](https://doi.org/10.1016/0003-2697(69)90064-5).
27. Kobuchi H, Moriya K, Ogino T, et al. Mitochondrial localization of ABC transporter ABCG2 and its function in 5-aminolevulinic acid-mediated protoporphyrin IX accumulation. *PLoS One*. 2012;7:e50082. <https://doi.org/10.1371/journal.pone.0050082>.
28. Frank J, Lornejad-Schäfer MR, Schöffel H, Flaccus A, Lambert C, Biesalski HK. Inhibition of heme oxygenase-1 increases responsiveness of melanoma cells to ALA-based photodynamic therapy. *Int J Oncol*. 2007;31:1539-1545.
29. Nowis D, Legat M, Grzela T, et al. Heme oxygenase-1 protects tumor cells against photodynamic therapy-mediated cytotoxicity. *Oncogene*. 2006;25:3365-3374. <https://doi.org/10.1038/sj.onc.1209378>.
30. Fridovich I. Superoxide dismutases: anti-versus pro-oxidants? *Anti-cancer Agents Med Chem*. 2011;11:175-177. <https://doi.org/10.2174/187152011795255966>.
31. Theodossiou TA, Olsen CE, Jonsson M, Kubin A, Hotherhall JS, Berg K. The diverse roles of glutathione-associated cell resistance against hypericin photodynamic therapy. *Redox Biol*. 2017;12:191-197. <https://doi.org/10.1016/j.redox.2017.02.018>.
32. Theodossiou TA, Ali M, Grigalavicius M, et al. Simultaneous defeat of MCF7 and MDA-MB-231 resistances by a hypericin PDT-tamoxifen hybrid therapy. *NPJ Breast Cancer*. 2019;5:13. <https://doi.org/10.1038/s41523-019-0108-8>.
33. Olsen CE, Sellevold S, Theodossiou T, Patzke S, Berg K. Impact of genotypic and phenotypic differences in sarcoma models on the outcome of photochemical internalization (PCI) of bleomycin. *Photodiagnosis Photodyn Ther*. 2017;20:35-47. <https://doi.org/10.1016/j.pdpdt.2017.08.010>.
34. Coburger J, Engelke J, Scheuerle A, et al. Tumor detection with 5-aminolevulinic acid fluorescence and Gd-DTPA-enhanced intraoperative MRI at the border of contrast-enhancing lesions: a prospective study based on histopathological assessment. *Neurosurg Focus*. 2014;36:E3. <https://doi.org/10.3171/2013.11.focus13463>.
35. Jichlinski P, Leisinger HJ. Photodynamic therapy in superficial bladder cancer: past, present and future. *Urol Res*. 2001;29:396-405. <https://doi.org/10.1007/s002400100215>.
36. Samkoe KS, Gibbs-Strauss SL, Yang HH, et al. Protoporphyrin IX fluorescence contrast in invasive glioblastomas is linearly correlated with Gd enhanced magnetic resonance image contrast but has higher diagnostic accuracy. *J Biomed Opt*. 2011;16:096008. <https://doi.org/10.1117/1.3622754>.
37. Stummer W, Stepp H, Wiestler OD, Pichlmeier U. Randomized, prospective double-blinded study comparing 3 different doses of 5-Aminolevulinic acid for fluorescence-guided resections of malignant gliomas. *Neurosurgery*. 2017;81:230-239. <https://doi.org/10.1093/neuros/nyx074>.

38. Ishikawa T. The ATP-dependent glutathione S-conjugate export pump. *Trends Biochem Sci.* 1992;17:463-468. [https://doi.org/10.1016/0968-0004\(92\)90489-v](https://doi.org/10.1016/0968-0004(92)90489-v).
39. Laborde E. Glutathione transferases as mediators of signaling pathways involved in cell proliferation and cell death. *Cell Death Differ.* 2010;17:1373-1380. <https://doi.org/10.1038/cdd.2010.80>.
40. Alaoui-Jamali MA, Bismar TA, Gupta A, et al. A novel experimental heme oxygenase-1-targeted therapy for hormone-refractory prostate cancer. *Cancer Res.* 2009;69:8017-8024. <https://doi.org/10.1158/0008-5472.can-09-0419>.
41. Robey RW, Ierano C, Zhan Z, Bates SE. The challenge of exploiting ABCG2 in the clinic. *Curr Pharm Biotechnol.* 2011;12:595-608. <https://doi.org/10.2174/138920111795163913>.
42. Abdel Gaber SA, Müller P, Zimmermann W, et al. ABCG2-mediated suppression of chlorin e6 accumulation and photodynamic therapy efficiency in glioblastoma cell lines can be reversed by KO143. *J Photochem Photobiol B, Biol.* 2018;178:182-191. <https://doi.org/10.1016/j.jphotobiol.2017.10.035>.
43. Kim JH, Park JM, Roh YJ, Kim IW, Hasan T, Choi MG. Enhanced efficacy of photodynamic therapy by inhibiting ABCG2 in colon cancers. *BMC Cancer.* 2015;15:504. <https://doi.org/10.1186/s12885-015-1514-4>.
44. Robey RW, Steadman K, Polgar O, Bates SE. ABCG2-mediated transport of photosensitizers: potential impact on photodynamic therapy. *Cancer Biol Ther.* 2005;4:187-194.
45. Fukuhara H, Inoue K, Kurabayashi A, et al. The inhibition of ferrochelatase enhances 5-aminolevulinic acid-based photodynamic action for prostate cancer. *Photodiagnosis Photodyn Ther.* 2013;10:399-409. <https://doi.org/10.1016/j.pdpdt.2013.03.003>.
46. Theodossiou TA, Papakyriakou A, Hothersall JS. Molecular modeling and experimental evidence for hypericin as a substrate for mitochondrial complex III; mitochondrial photodamage as demonstrated using specific inhibitors. *Free Radic Biol Med.* 2008;45:1581-1590. <https://doi.org/10.1016/j.freeradbiomed.2008.09.015>.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Mastrangelopoulou M, Grigalavicius M, Raabe TH, et al. Predictive biomarkers for 5-ALA-PDT can lead to personalized treatments and overcome tumor-specific resistances. *Cancer Reports.* 2020;e1278. <https://doi.org/10.1002/cnr2.1278>

“Predictive biomarkers for 5-ALA-PDT can lead to personalized treatments and overcome tumor-specific resistances”

Maria Mastrangelopoulou¹, Mantas Grigalavicius¹, Tine Henriksen Raabe¹, Ellen Skarpen², Petras Juzenas³, Qian Peng³, Kristian Berg¹, Theodossis A. Theodossiou^{1*}

¹Department of Radiation Biology, Institute for Cancer Research, Oslo University Hospital, Oslo, Norway.

²Department of Molecular Cell Biology, Institute for Cancer Research, Oslo University Hospital, Oslo, Norway.

³Department of Pathology, The Norwegian Radium Hospital, Oslo University Hospital, 0379 Oslo, Norway

Supplementary information

Supplementary Fig. 1. Spectral profile of PpIX and emission profile of irradiation source.

Supplementary Fig. 2. Intracellular localization of the 5-ALA-derived photosensitizer PpIX for the various cell lines.

Supplementary Fig. 3. Metabolic studies 1 h after 5-ALA-PDT, measured by the Seahorse XFe96 analyzer.

Supplementary Figure 1

Supplementary Fig.1. Spectral profile of the molar extinction coefficient of PpIX in DMSO (blue line) vs. the emission spectrum of our lamp built in-house (yellow line). The PpIX absorbance profile also exhibited four Q bands at ~510, 543, 583, and 633 nm.

Supplementary Fig. 2. Intracellular localization of the 5-ALA-derived photosensitizer PpIX for the various cell lines: a) T98G, b) U87, c) MCF7, d) MDA-MB-231, and e) T47D. Cells were incubated with 1.5 mM 5-ALA for 1 h with (even rows) or without (odd rows) pretreatment with Ko143 (1 μ M). The cells were incubated with 100 nm MitoTracker Green 15 min prior to imaging. MitoTracker Green fluorescence is represented in green (left column) and PpIX fluorescence in red (second column). An overlay of PpIX fluorescence with MitoTracker fluorescence is shown in the overlap panel (third column) where colocalization can be seen in yellow. Pure colocalization between PpIX and MitoTracker is shown in the colocalization panel in white (fourth panel).

Supplementary Fig. 3. Metabolic studies 1 h following 5-ALA-PDT, measured by the Seahorse XFe96 analyzer. 30,000 cells (T98G, U87, MCF7, MDA-MB-231, and T47D) per 96-well plate were treated with media only or various concentrations of 5-ALA corresponding to LD₃₀ for 4 h

and subsequently irradiated for 60 sec. Basal oxygen consumption rates (OCRs) for all cell lines without and with light irradiation are presented in panels a and b, respectively. The basal extracellular acidification rate (ECAR) for all cell lines without and with light irradiation are presented in panels c and d, respectively. The data shown represent the average of three independent experiments, while the error bars correspond to 1 SD.

Supplementary Fig. 4. Intracellular production of PpIX following 4h incubation with 0.5mM 5-ALA (taken from the flow data, Fig.1F) vs. cell viability following 60s irradiation again after incubation with 0.5mM 5-ALA (data taken from Fig. 1A). Data shown represent the average of three independent experiments, while the error bars correspond to 1 SD.

IV

Special Issue Research Article

Photodynamic Efficacy of Cercosporin in 3D Tumor Cell Cultures[†]Mantas Grigalavicius¹, Maria Mastrangelopoulou¹, Delmon Arous^{1,2}, Asta Juzeniene¹, Mathilde Ménard³, Ellen Skarpen⁴, Kristian Berg¹ and Theodossis A. Theodossiou^{1*} ¹Department of Radiation Biology, Institute for Cancer Research, The Norwegian Radium Hospital, Oslo University Hospital, Oslo, Norway²Department of Physics, University of Oslo, Oslo, Norway³Institut Charles Gerhardt Montpellier, UMR-5253 CNRS-UM-ENSCM cc 1701, Montpellier cedex 05, France⁴Department of Core Facilities, Institute for Cancer Research, The Norwegian Radium Hospital, Oslo University Hospital, Oslo, Norway

Received 28 October 2019, accepted 2 February 2020, DOI: 10.1111/php.13257

ABSTRACT

In the present work, we study the photodynamic action of cercosporin (cerco), a naturally occurring photosensitizer, on human cancer multicellular spheroids. U87 spheroids exhibit double the uptake of cerco than T47D and T98G spheroids as shown by flow cytometry on the single cell level. Moreover, cerco is efficiently internalized by cells throughout the spheroid as shown by confocal microscopy, for all three cell lines. Despite their higher cerco uptake, U87 spheroids show the least vulnerability to cerco-PDT, in contrast to the other two cell lines (T47D and T98G). While 300 μm diameter spheroids consistently shrink and become necrotic after cerco PDT, bigger spheroids (>500 μm) start to regrow following blue-light PDT and exhibit high viability. Cerco-PDT was found to be effective on bigger spheroids reaching 1 mm in diameter especially under longer exposure to yellow light (~590 nm). In terms of metabolism, T47D and T98G undergo a complete bioenergetic collapse (respiration and glycolysis) as a result of cerco-PDT. U87 spheroids also experienced a respiratory collapse following cerco-PDT, but retained half their glycolytic activity.

INTRODUCTION

Photodynamic therapy of cancer (PDT) (1) is cancer treatment modality which utilizes a photoactivatable drug (photosensitizer, PS), light of the appropriate wavelength ensuring PS activation, and interstitial molecular oxygen. The light-activated sensitizer reacts with the surrounding oxygen, generating toxic reactive oxygen species (ROS) and prominently singlet oxygen. *In vivo*, PDT can destroy cancer through a combination of (1) cell phototoxicity, (2) vascular shut down leading to tumor starvation and (3) immunogenicity triggered by the primary treatment-induced inflammation. In *in vitro* PDT studies, however, only the photocytotoxic action

is evaluated, except if a specialized model has been designed. The main restraint of PDT is the limited depth of light penetration into tissue. Light in the red end of the visible spectrum can penetrate a few millimeters (~2 mm) into tissue while in the blue end the corresponding penetration depth is a couple of hundred micrometers (~300 μm at ~450 nm, (1)). In that context, an ideal clinical PDT photosensitizer would need to exhibit strong absorbance at or longer than 650 nm (deep red).

Cercosporin (cerco) is a naturally derived perylenequinone PS, like hypericin and the hypocrellins. It is produced by the *Cercospora* family of plant pathogens, however, its total synthesis in the lab has also been reported (2). Cerco has not been extensively studied as an anticancer photosensitising agent due to its short wavelength of activation (peak at ~470 nm). Nevertheless, it has been studied for its photodynamic activity within the remit of phytopathology (3) as it is a very powerful phototoxin for plants and crops. In that context, cerco has been reported to have a high singlet oxygen quantum yield, ranging from 0.84 in D_2O to 0.97 in dioxane (4).

In our recent work on 2D cell cultures (5), we evaluated the capacity of cerco as a photosensitizer on three mammalian cancer cell lines, namely MCF7 (human breast adenocarcinoma), T98G and U87 (both human glioblastoma multiforme). We found that cerco is a very potent PS, efficiently killing all the cell lines investigated at different light doses applied. The cell killing essentially followed the PS loading patterns: U87 that presented half the cerco uptake of the other two cell lines, also demonstrated the lowest sensitivity to cerco-PDT, as compared to the other two cell lines in the study. In the same work, we also investigated the metabolic profiles of the three cell lines before and after treatment with cerco-PDT. We found that PDT treatment collapsed the metabolism in MCF7 and U87 cells while T98G cells, despite of their respiratory activity also dropping to background levels, retained about 1/3 of their glycolytic activity following cerco-PDT.

In the present study, we evaluated the effect of cerco-PDT in 3D (spheroid) cultures of three mammalian cancer cell lines, namely T47D (human breast adenocarcinoma), U87 and T98G (both human glioblastoma multiforme). It is well known that multicellular spheroids provide a system of intermediate complexity between solid tumors *in vivo* and cell monolayers in

*Corresponding author email: ththeo@rr-research.no (Theodossis A. Theodossiou)

[†]This article is part of a Special Issue dedicated to Dr. Thomas Dougherty.© 2020 The Authors. *Photochemistry and Photobiology* published by Wiley Periodicals, Inc. on behalf of the American Society for Photobiology.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

culture (6), as they model the 3D cell–cell interactions and also the 3D structural barriers to cell nutrients, substrates and oxygen. In that context, the rationale and motivation for conducting the present study was to compare our recent 2D cell culture results with the corresponding cerco-PDT effects in 3D cultures.

MATERIALS AND METHODS

Chemicals and reagents. Agarose, cercosporin from *Cercospora hayii* (cerco), antimycin A, oligomycin (Oligo), carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP), myxothiazol (Myxo), RPMI 1640 without phenol red, L-glutamine, penicillin/streptomycin, trypsin, poly-D-lysine hydrobromide, fluorescein diacetate (FDA) and Dulbecco's phosphate-buffered saline (PBS) with or without calcium chloride and magnesium chloride were purchased from Sigma-Aldrich Norway AS (Oslo, Norway). Corning™ Cell-Tak cell and tissue adhesive, Corning™ Matrigel® Matrix, and Corning™ 96-well clear ultra low attachment microplates were purchased from Fisher Scientific (Göteborg, Sweden). TrypLE™ Express Enzyme without phenol red, propidium iodide (PI), LIVE/DEAD™ cell stain kit and fetal bovine serum (FBS) were purchased from Thermo Fisher Scientific (Oslo, Norway). All consumables used for Seahorse XFe96 metabolic analysis were purchased from Agilent Technologies (Santa Clara, CA).

Generation of spheroids. One human breast cancer (T47D) and two human brain cancer (T98G and U87) cell lines were employed to initiate the spheroids. Both cells in 2D culture and spheroids were maintained in RPMI 1640 without phenol red, supplemented with 10% FBS, 100 U mL⁻¹ penicillin/100 µg mL⁻¹ streptomycin and 2 mM L-glutamine at 37°C in a 5% CO₂ humidified atmosphere. The spheroids were generated in agarose-coated flat bottom or ultra low attachment U-shaped 96-well microplates. The agarose was boiled in PBS for 10 min under continuous stirring prior dispensing into 96-well microplates and allowed to cool down for at least 1 h before setting the cells. The flat bottom microplates were coated with 1.5% (wt/vol) agarose and following the transfer of 100 µL cell suspension into each well they were centrifuged at 800 g for 20 min. The initial cell number required to create spheroids of similar size at day 4 after seeding varied from 500 to 5000.

PDT experiments. Cercosporin was dissolved in 100% DMSO to yield a 4 mM stock solution and then further diluted in RPMI complete medium as required for each experiment. Spheroids were incubated with cercosporin (6 µM) in complete medium for 4 h and washed prior to irradiation with phosphate buffer saline (PBS) containing calcium and magnesium. The blue light irradiations were performed in PBS from the top of 96-well plates using a Lumisource® lamp (PCI Biotech AS, Oslo, Norway) with four OSRAM L 18/67 fluorescent tubes while yellow light irradiations were carried out with a custom-made lamp equipped with an array of 1W LEDs. The Lumisource® lamp had an irradiance of 5.3 mW cm⁻² while the irradiance of the yellow lamp was 0.9 mW cm⁻², both measured by a photodetector PH100-SiUV coupled to a calibrated optic power meter Gentec-EO SOLO2 (Gentec-EO, Quebec, Canada). Exposure of spheroids to 4 min of blue light corresponded to a total dose of 1.3 J cm⁻² while exposure of 30 min to yellow light gave a total dose of 1.6 J cm⁻². The spectrum of blue and yellow lamp peaked at approximately 450 and 590 nm, respectively (Fig. 1a), as measured by an irradiance-calibrated AvaSpec-2048x14-SPU2 FiberOptic Spectrometer (Avantes, Apeldoorn, The Netherlands).

Metabolic studies. T98G, U87 and T47D spheroids of 300–550 µm size were transferred from Corning™ U-shaped microplates to Agilent Seahorse XFe96 Spheroid microplates coated with poly-D-lysine or Cell-Tak™ adhesives according to the manufacturers' recommendations. 175 µL of XF RPMI Medium pH 7.4 (Agilent Technologies) supplemented with 10 mM glucose, 2 mM L-glutamine and 2 mM sodium pyruvate was added to each well following pH adjustment to 7.4. The spheroids were incubated for an extra hour at 37°C, without CO₂. The changes in spheroid oxygen consumption and media acidification rates (OCR and ECAR, respectively) with and without cercosporin incubation (6 µM) following light irradiation were registered using Seahorse XFe96 analyzer (Agilent Technologies). The mitostress assay was used according to a modified version of the manufacturer's protocol, employing serial injections of Oligo, FCCP and Myxo at final concentrations of 1, 1.3 and 2 µM, respectively. The obtained results were also normalized for relative spheroid volumes between the cell groups.

Confocal and epifluorescence microscopy. The spheroids were examined with a Zeiss LSM 880 Airyscan microscope (Carl Zeiss MicroImaging GmbH, Jena, Germany), equipped with an Ar-Laser Multiline (458/488/514 nm), a DPSS-561 10 (561 nm), a Laser diode 405-30 CW (405 nm) and a HeNe laser (633 nm). The objectives used were the Zeiss plan-Apochromat 20× NA/0.8 air, the 25× NA/0.8 oil DICII or the 10× NA/1.4 air DICII for a larger field of view. Image acquisition, processing and visualization were performed with the ZEN 2.3 SP1 basic software (Carl Zeiss). The spheroids were either fixed with 2% formaldehyde and sandwiched between a thin glass plate and cover slip or transferred to a cloning ring mounted on a MatTek glass bottom dish without fixation. In both cases, spheroids were immobilized using a Matrigel® Matrix:media mixture (50:50, vol/vol).

In case of epifluorescence microscopy, green- and red-filtered images (live/dead or FDA/PI assay) were obtained by using an inverted microscope Axiovert 200M (Zeiss, Germany) in reflection mode while the images without optical filtering were acquired in transmission mode (achroplan objective with magnification/numeric aperture 4×/0.1, resolution 2560 × 1920, spatial resolution 1.208 pixels µm⁻¹, bit depth 24, xenon lamp voltage 2.5–10.0 V). Camera exposure times were the same for PDT and control (CTRL) groups for the same spheroid (green FDA channel 50–150 ms, red PI channel 50–900 ms, nonfluorescent imaging 750–1000 µs). The spheroids were incubated with FDA (green, 8 µg mL⁻¹) and PI (red, 20 µg mL⁻¹) for 5 min and then washed with PBS containing Ca and Mg just before the imaging.

Spheroid image analysis. The spheroid areas were automatically delineated and quantified with in-house made software developed in Matlab (MathWorks, Natick, MA). The L channel in the La*b* space of each RGB image were enhanced by performing contrast-limited adaptive histogram equalization (CLAHE) before being converted into grayscale images by eliminating the hue and saturation information while retaining the luminance. Utilizing the imbinarize Matlab function, grayscale images were then binarized to appropriately mask the whole spheroids by iteratively sweeping through different threshold sensitivities. Locally adaptive thresholding, based on mean intensity in the neighborhood of each pixel (Bradley's method (7)), was necessary as nonuniform background illumination varied across the images due to spheroid spatial displacement over time. Of the spheroids that the Matlab program was inadequately able to automatically delineate, polygon structures were manually outlined and measured on the raw spheroid images using the Zeiss microscopy software AxioVision release 4.8.2.

Spectroscopic and flow cytometric evaluation of loaded cercosporin. In order to simulate the absorbance-fluorescence profile of cercosporin in spheroids, 40 000–60 000 cells were inoculated into 96-well plates 24 h prior to 4 µM cerco incubation for 4 h. Cells were subsequently washed twice with PBS and scanned for their 2D excitation–emission (20 nm bandwidth) in the Tecan spark M10 plate reader.

In order to determine the loading of spheroids with cerco, twenty to thirty spheroids were incubated with 6 µM cercosporin in complete RPMI media for 4 h before transferring them from agarose-coated microplates. Dissociation of these 3D cell structures into single cell suspensions was performed in 1 mL of TrypLE™ Express Enzyme after a single wash in PBS. The resulting cell suspensions were washed again with PBS and strained through a 35 µm mesh screen before the flow cytometry experiments. The flow cytometric analyses were performed using a LSRII (BD Biosciences, Franklin Lakes, NJ) and at least of 10 000 events for each sample. Cercosporin fluorescence was excited at 488 nm and registered after a bandpass filter at 670/14 nm. Flow cytometry data were analyzed using the FlowJo v.7.6.1 software (Treestar Inc., Ashland, OR).

Data analysis. A statistical analysis of the data obtained was performed using SigmaPlot version 12.5 (Systat Software, Inc., Chicago, IL) and RStudio version 1.1.463 (8) with a preinstalled R package version 3.6.1 (9).

RESULTS

Spectral response and relative quantity of loaded cercosporin

Initially, we incubated the cells with 4 µM cerco for 4 h. At the end of the incubation, the cells were scanned from the plate underside to produce a 2D excitation vs emission spectra. A representative pseudocolor contour map is shown in Fig. 1b for

Figure 1. Spectroscopic and flow cytometric analyses. (a) Emission spectra of the lamps used for PDT (blue lamp—blue line, yellow lamp—orange line) (b) 2D excitation–emission profile of cercosporin in cell culture after incubation for 4 h with $4 \mu\text{M}$, (c) Representative histograms of cercosporin uptake by T98G, T47D and U87 spheroids after incubating them with $6 \mu\text{M}$ cercosporin for 4 h.

T98G cells. From that graph, it can be seen that the maximal absorbance of cell-internalized cerco is at $\sim 480 \text{ nm}$ whereas maximal emission is at $\sim 630 \text{ nm}$. There is also a secondary shoulder at the end of the primary excitation peak extending to $\sim 580 \text{ nm}$. With regard to spheroid irradiation with blue light, the emission spectrum of the lamp covered the most pronounced excitation peak of cerco while the yellow lamp excited cerco over its extended secondary shoulder (Fig. 1a). The effective irradiance of the blue lamp, $E(t)$, was calculated as the product of cerco excitation, $\phi(\lambda)$, and lamp emission, $I(\lambda, t)$, and found to be 3.9 mW cm^{-2} , when the integration was performed over the wavelength (λ) range of the lamp emission spectrum: $\int I(\lambda, t) \cdot \phi(\lambda) \cdot d\lambda$. A 4-min exposure time, t , yielded an effective dose of 0.9 J cm^{-2} . The effective dose and irradiance of the yellow lamp were 0.44 J cm^{-2} and 0.24 mW cm^{-2} , respectively.

In order to evaluate the cerco uptake of the spheroids, we incubated them for 4 h with $6 \mu\text{M}$ cerco and then disrupted the spheroids into single cell suspensions using TrypLE[™] as elaborated in the materials and methods section. The cells then were analyzed by flow cytometry for their cerco content at the single cell level, and the results are shown in Fig. 1c. It can be seen that T98G and T47D spheroids took up half the cerco than U87.

Study of spheroid uptake of cercosporin by confocal microscopy

All spheroids were incubated with $4 \mu\text{M}$ cerco for 4 h. Then, the spheroids were imaged in two ways as explained in materials and methods. Representative micrographs are shown in Fig. 2. In the top row, the spheroids were squeezed between the slide and the coverslip to reduce the spheroid z-dimensions so that the uniformity of cerco fluorescence could be evaluated without losing light due to spheroid size. On top and right side of each micrograph, the representative orthogonal projections are shown. The confocal slice selected in the each main micrograph is represented as a blue line in the orthogonal projections. From these micrographs, it can be seen that in all spheroids (U87, T47D and T98G) the fluorescence is uniform throughout the x- or y-dimensions, even though the U87 spheroids were bigger.

The bottom row micrographs were obtained without any z squeezing of the spheroids and thus depict the actual z

dimensions of the spheroids. In this case, as can be seen in the micrographs, the light could not penetrate through the whole spheroid height so the spheroid dimensions can be deduced by assuming symmetry in the orthogonal projections.

Once more as can be seen in the representative confocal slices, the fluorescence is uniform throughout the confocal slice area (even perhaps slightly more intense in the spheroid core), indicating that cerco was taken up well even $\sim 150 \mu\text{m}$ deep in the spheroids.

Spheroid responses to PDT treatment

PDT-treated spheroids were incubated with $6 \mu\text{M}$ cerco for 4 h and then irradiated either with 1.3 J cm^{-2} of blue light or 1.6 J cm^{-2} yellow light.

In a first set of PDT experiments, T47D spheroids were grown from different cell inoculation numbers (500, 1000 and 2000) to different sizes (see below) and subjected to cerco-PDT. On day 17 from irradiation, fluorescence images were obtained with use of two dyes, PI (red for dead) and FDA (green for live). From those endpoint images, it can be seen that there is a profound difference in spheroid area between the two groups. Furthermore, the control spheroids consisted in their vast majority of live cells (green) while the PDT-treated spheroids almost exclusively consisted of dead cells (red), showing no sign of regrowth. The growth curves (Fig. 3a) confirm that the PDT-treated T47D spheroids exhibited sustained cell inhibition for all seeding densities (0.049 ± 0.003 to $0.043 \pm 0.002 \text{ mm}^2$ for 500 cells, 0.074 ± 0.005 to $0.067 \pm 0.005 \text{ mm}^2$ for 1000 cells and 0.104 ± 0.004 to $0.096 \pm 0.009 \text{ mm}^2$ for 2000 cells), while the control spheroids exhibited continuous growth (0.050 ± 0.003 to $0.456 \pm 0.024 \text{ mm}^2$ for 500 cells, 0.072 ± 0.003 to $0.534 \pm 0.033 \text{ mm}^2$ for 1000 cells and 0.100 ± 0.005 to $0.642 \pm 0.017 \text{ mm}^2$ for 2000 cells).

In a second set of PDT experiments, cell spheroids were irradiated at three different sizes in case of T47D (0.27 ± 0.02 , 0.57 ± 0.02 and $0.81 \pm 0.04 \text{ mm}^2$) and one size in case of U87 ($0.47 \pm 0.03 \text{ mm}^2$). The growth of both control (irradiated in the absence of cerco) and cerco-PDT treated spheroids was followed in time through measurement of their cross-sectional area. Representative images at selected time points are shown for T47D and U87 spheroids in Fig. 4a and b correspondingly while

Figure 2. Confocal images of spheroids incubated with 4 μM cercosporin. The images in the top panels (a–c) combining both fluorescence and transmitted light channels represent spheroids between two glass slides while the bottom panels (d–f) show confocal images of nonsqueezed spheroids in a single fluorescence channel. All the panels contain orthogonal projections generated from Z-stacks. Blue lines within the orthogonal projections denote the Z-position of presented cross section.

the full sets of cross-sectional area curves are shown in Fig. 4c. From Fig. 4c, it can be seen that in the case of T47D there is a growth inhibition following PDT, where spheroid area dropped from 0.27 ± 0.02 to $0.21 \pm 0.01 \text{ mm}^2$ in eight days. Conversely, the nontreated T47D spheroids expanded in size from 0.22 ± 0.02 to $0.43 \pm 0.02 \text{ mm}^2$. The results were quite different for U87 cells: While cerco PDT conferred a temporary growth inhibition until day 3 after the PDT treatment from 0.47 ± 0.03 to $0.32 \pm 0.02 \text{ mm}^2$, the treated cells eventually caught up with the non treated cells which exhibited a slower (nonlinear) growth unlike the T47D (0.42 ± 0.05 to $0.57 \pm 0.05 \text{ mm}^2$).

Having observed that PDT-treated, larger U87 spheroids caught up in area with their control counterparts, we also followed bigger T47D spheroids of 0.81 ± 0.04 and $0.57 \pm 0.02 \text{ mm}^2$ area after irradiating them with blue (Fig. 4d) and yellow (Fig. 4e) light correspondingly to see if they also exhibit a similar delayed trend. The growth curves from bigger spheroids in Fig. 4f show that the application of blue light on cerco-incubated spheroids resulted only in a weak PDT response when compared to their control group (0.57 ± 0.02 to $0.73 \pm 0.06 \text{ mm}^2$ vs 0.50 ± 0.02 to $0.82 \pm 0.03 \text{ mm}^2$). The irradiation with yellow light, however, caused a significant growth inhibition from 0.81 ± 0.04 to $0.73 \pm 0.04 \text{ mm}^2$ in 9 days while the control spheroids exhibited continuous growth (0.82 ± 0.03 to $1.25 \pm 0.04 \text{ mm}^2$).

We subsequently assessed the live-dead fractions of the T47D and U87 spheroids as the follow-up endpoint. The results can be seen in Fig. 5: In the first panel (a) the T47D spheroids with/without PDT treatment at day 8 are shown. It can be seen that apart from their profound difference in area, the control spheroids are in the vast majority consisted of live cells (green) while the PDT-treated spheroids are almost exclusively consisted of dead cells (red). The corresponding micrographs for U87 cells corroborate the observations in Fig. 4b. The end sizes of control and PDT-treated spheroids are comparable. Additionally, while the treated spheroids contain a significantly larger fraction of dead cells (red) they also contain a large fraction of viable cells. What is interesting in this case (PDT-treated spheroids) is that following PDT treatment, viable U87 cell groups detach from the main spheroid body and eventually start forming new spheroids (Fig. 4b, day 8, and Fig. 5b, white circles). Figure 5c represents yellow light-irradiated spheroids after 9 days of cerco-PDT treatment (initial size in diameter before treatment was $1010 \pm 26 \mu\text{m}$). Here, the treated spheroids experienced an observable shrinkage and only a small fraction of viable cells was observed in comparison with either the yellow-light control group or blue light-irradiated spheroids (Fig. 5d). Blue light irradiation had a marginal effect and induced only a delay in spheroid regrowth over the first week (Fig. 4f) that was followed by an increase in viable cells in the periphery of the spheroids in the next two weeks (Fig. 5d, irradiated at initial size of $802 \pm 12 \mu\text{m}$).

Figure 3. The effect of blue-light cerco PDT on T47D spheroid growth, 7 days following the inoculation with different cell densities. (a) 17-day follow-up growth curves. Dotted lines represent untreated controls, solid lines correspond to growth following cerco PDT. (b) Spheroid endpoint images: The spheroids were incubated with FDA (green, 8 $\mu\text{g mL}^{-1}$) and PI (red, 20 $\mu\text{g mL}^{-1}$) for 5 min and then washed with PBS containing Ca and Mg just before the imaging. Scale bars = 100 μm .

Metabolic studies in spheroids

All spheroids were treated with 6 μM cerco and incubated for 4 h, before being subjected to PDT. One hour following the PDT treatment, control and treated spheroids were evaluated by the seahorse XFe96 analyzer for their oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) representing their respiration and glycolytic profiles correspondingly. The OCR trends in non-PDT-treated spheroids are presented in Fig. 6a. It can be seen that T47D spheroids exhibited higher basal OCR ($\sim 40 \text{ pmol min}^{-1}$) than the other cell lines with U87 second ($\sim 30 \text{ pmol min}^{-1}$) and T98G presenting much lower respiratory activity ($\sim 10 \text{ pmol min}^{-1}$). Nevertheless, when the respective spheroid volumes were taken into account (T47D: U87:T98G $\sim 3.6:2.9:1$), the three cell lines exhibited comparable OCR (the normalized OCR profiles are shown in Fig. 6b).

It should be noted that the data in Fig. 6b reveal that T47D displayed the smallest reduction in oxygen consumption upon application of Oligo, while T98G spheroids exhibited the highest respiratory capacity among the three cell lines, when FCCP was added. In Fig. 6a, it can be seen that the respiration of all three spheroid types was profoundly subdued as early as one hour following PDT and was not responsive to any of the modulator stimuli (Oligo, FCCP, Myxo).

The glycolytic profiles of the three spheroid types can be seen in Fig. 6c and again in Fig. 6d following volume correction. It is evident from non-PDT-treated spheroids (Fig. 6d) that U87 exhibited the highest glycolytic activity ($\sim 23 \text{ mpH min}^{-1}$) with T98G a close second ($\sim 20 \text{ mpH min}^{-1}$) and T47D exhibiting the lowest basal glycolysis at $\sim 13 \text{ mpH min}^{-1}$. All glycolytic profiles were responsive to the addition of Oligo, blocking ATP production from respiration. The application of PDT also abolished the glycolytic function in both T47D and T98G, while in U87, even though there was a profound reduction in the glycolytic rate, it was found to be at about 40% the glycolytic rate of control cells. The metabolic phenotypes of the spheroids (OCR vs ECAR) are shown in Fig. 6e and also in Fig. 6f following corrections for spheroid volumes, providing an overview of the findings in Fig. 6a–d.

DISCUSSION

The present study on the effects of cerco PDT in 3D spheroid cultures is an extension/follow-up of our previous study (5) where we studied the efficacy of cercosporin as a photosensitizer in 2D cultures.

Even though cercosporin is a very potent photosensitizer as concluded from both the studies, it is not used in conventional

Figure 4. The effect of cerco PDT on spheroid growth. Spheroid sizes by means of cross-sectional areas followed for 8–9 days (a and b) and 17 (e) days. Microscope images (a, b and d, e) taken at different time points (days 0–1, 3 and 8–9 after exposure to blue (a, b and e) or yellow (f) light) correspond to panels a and b and depict the actual differences between cerco (+C) and noncerco (–C) groups. The cerco-incubated spheroids were imaged through a red filter to avoid further excitation of cerco. The T47D long follow-up curves (e) exhibited first order exponential growth. Error bars: standard deviation (SD).

PDT, due to its short wavelength of activation (~470 nm, Fig. 1b) which very much limits the depth of light penetration into tissue in contrast to clinically used photosensitizers absorbing in the red end of the visible spectrum (mTHPC, PpIX, chlorine e6, etc.). Nevertheless, cercosporin possesses a high singlet oxygen quantum yield (from 0.84 in D₂O to 0.97 in dioxane, (4)) and could be very valuable in cases where external light administration is not required for the activation of the photosensitizer. One such case is our recently published work (10), where accelerated protons were shown to activate photosensitizers. More specifically cerco was activated by proton irradiation in M059K and T98G glioblastoma multiforma cells achieving significant differential cytotoxicity in comparison to cells irradiated in the absence of cerco. Another example of an application where cerco could also be very useful, is chemi- or bioluminescence PDT (11) where the photosensitizer is excited directly by a chemi/bioluminescent molecule in close proximity within the cell.

Despite the fact that both our 2D cell study (5) and our present work in 3D spheroid cultures are in agreement that cerco is a quite potent photosensitizer, there are several discrepancies in the effect of cerco-PDT on cell monolayers and 3D cultures. In 2D cell cultures, we found that U87 cells were the least efficient in uploading cercosporin as compared to MCF7 breast adenocarcinoma and T98G GBM cell lines following incubation with 4 µM cerco for 4 h. In the case of the 3D spheroids, however, incubation under the same conditions led to U87 cells showing the most efficient loading in the single cell level, double than both the T98G and T47D cells. One possible explanation for that is that the size (volume) of the spheroids was not the same for all cell lines but as delineated above it followed a ratio of T98G:T47D:U87 ~ 1:2.9:3.6. The substantially (three-fold) larger U87 spheroids may have the ability to hold more dye in the interstitium between cells, providing the actual cells with a bigger reservoir for uptake. It can also be seen from Fig. 2 that U87

spheroids are not that tightly packed with cells as T47D or T98G spheroids, probably making the diffusion of cerco easier.

In contrast to the findings of the 2D study, blue light irradiation was not very effective in the bigger spheroid models at the regime used (6 µM cerco for 4 h, with 1.3 J cm⁻² blue light at 450 nm), while the smaller T47D were found to be vulnerable (growth inhibited and necrotic) to PDT in the same conditions. The small T47D spheroids remain in a compact spheroid form and continuously shrink with time into a necrotic core, while the bigger T47D spheroids are more or less resistant to the blue light treatment, even more than similar size U87 spheroids. This is in agreement with our previous 2D cerco-PDT study (5), the extent of vulnerability tends to correlate with the uptake of cerco as U87 spheroids experienced a stronger initial insult of cerco-PDT than T47D spheroids of similar size (Fig. 4c and f). On the other hand, the bigger size T47D spheroids (~1000 µm) were quite effectively inhibited with yellow light irradiation (1.6 J cm⁻² at ~590 nm) at a lower effective dose than the blue light (blue:yellow ~ 0.9 J cm⁻²:0.44 J cm⁻²). This clearly demonstrates the limitations of shorter wavelength (blue) light in penetrating tissue; in smaller spheroids, blue light can more or less activate the PS in the entire volume of the spheroid, but not in larger spheroids. This is also evident from the spheroid microscopy experiments: When the spheroids are imaged in Matrigel (no suppression of their z dimension), the bottom end of the spheroids in the z stack image is dark, especially for U87 which are much larger. Yellow light conversely, although much less absorbed by cerco, has a far more detrimental effect on cerco-treated spheroids, even at a lower effective dose than blue light.

Another notable trend of the U87 spheroids is that following cerco-PDT with blue light which is more effective in the surface, they tend to shed cell clusters which then form new viable spheroids (Fig. 4b, white circles). This is a sign of motility upon insult (of PDT in this case) and could be associated with the documented ability of GBM cells to readily metastasize mainly

Figure 5. Spheroid endpoint images of short (a–c) and long (d) follow-ups. Camera exposure times were the same for PDT and control (CTRL) groups for the same spheroid (green FDA channel 50–150 ms, red PI channel 50–900 ms). The spheroids were incubated with FDA (green, $8 \mu\text{g mL}^{-1}$) and PI (red, $20 \mu\text{g mL}^{-1}$) for 5 min and then washed with PBS containing Ca and Mg just before the imaging. Scale bars = 100 μm .

Figure 6. The changes in metabolic activity of spheroids after cercosporin treatment. The left-sided panels (a, c and e) represent the raw (uncorrected) data while the panels on the right side depict the normalized results by spheroid volume. Oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) per one spheroid in 96-well Seahorse microplates were measured by the Seahorse XFe96 analyzer. T47D, T98G and U87 spheroids were treated with media only (–C) or 6 μ M cercosporin (+C) for 4 h and subsequently irradiated for 3 min after washing and spheroid transfer steps. The vertical lines show the injection of oligomycin (Oligo), carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP) and myxothiazol (Myxo). The OCR vs ECAR values in panels a and b were taken from the fifth measurement representing equilibrated baseline respiration levels. Circle symbols represent T47D, diamonds—T98G, triangles up—U87. Figure legends are located within the panels e (cercosporin-treated) and f (non-cercosporin-treated). The vertical arrows in e represent the drop in OCR following PDT, while the horizontal arrows represent the corresponding drop in ECAR. The arrows in f represent change in metabolism for the non-cercosporin-treated spheroids following normalization for spheroid volume. The normalization factors appear next to each arrow. Error bars: standard error of the mean (SE).

within the brain forming new islets (12). U87 cells form spheroids easier than both T98G and T47D, due to their natural tendency even in 2D cultures to form clusters. The U87 spheroids however are quite loose than the more “tightly packed” T47D spheroids.

In all spheroid cases, the outer layer of the spheroids remains more viable as they grow and that appears as a ring in the microscopy micrographs. As the spheroid size grows, the inner cell layers suffer from the deprivation of nutrients and poor oxygen diffusion and gradually become necrotic. This is closer to the reality in tumors (6), that is why 3D cultures are much better tumor models than 2D cell monolayers. Nevertheless the real

tumors are much more heterogeneous in their constitution (various malignant cells, auxiliary normal cells and extracellular matrix) than the spheroids which purely consist of only the tumor cell line.

The metabolic profile of the spheroids is also very different to that in 2D cell cultures (5). In cell monolayers, U87 cells exhibited much lower respiratory activity than T98G cells. When assayed in 3D formation, however, after volume corrections all three spheroids exhibited similar OCR. It must be noted that in the 2D studies, the cells were not corrected for size. Indeed, when adhered on the plates and stretched out T98G look much bigger than U87, however when their sizes were measured by a

Beckman-Coulter counter (Beckman-Coulter, Miami), these differences were not so profound (median volume: U87 $\sim 3.3 \times 10^3 \mu\text{m}^3$, T47D $\sim 3.3 \times 10^3 \mu\text{m}^3$ and T98G $\sim 5.2 \times 10^3 \mu\text{m}^3$) and normalization would only reduce the U87/T98G OCR ratio from 1/3 to 1/2. This indicates that the spheroid respiration is regulated mainly by spheroid volume and not cell type. Similarly there were also discrepancies in the glycolytic rates between the 2D and 3D cultures. In cell monolayers, again T98G had a much higher glycolytic rate than U87 ($\sim$ three-fold) while in the 3D case the rates are comparable with U87 spheroids being marginally more glycolytic than T98G. Cerco-PDT collapsed the metabolism of both T98G and T47D, and also the respiration of U87, but not the glycolytic rate of the latter which continued at almost half capacity. This again is in disagreement with the 2D scenario where the cell lines were crashed metabolically (including U87) except for T98G which maintained their glycolytic rate at $\sim$ 1/3. One more factor that contributes to the above observations is the much looser adhesion of U87 spheroids (*vide infra*), which may apparently allow both glycolysis substrates and oxygen easier access to the spheroid inner layers. As it has been shown in our previous work (5), cerco is subcellularly located in the endoplasmic reticuli/cell mitochondria. As a result, the PDT activation of cerco may confer considerable damage to the cellular respiration by damaging the electron transport chain (ETC) constituent complexes. In our previous work (13) using a similar naturally occurring perylenequinone PS, hypericin (HYP), we found that HYP-PDT damages irreversibly complex III of the ETC in DU145, human prostate carcinoma cells. In a more recent publication (14), we further saw that HYP-PDT caused a profound decrease of both the respiration and the glycolytic rate of MCF7 human breast adenocarcinoma cells, 1 h following irradiation. This is in good agreement with our 2D cerco findings (5), where cerco-PDT collapsed the respiration and profoundly diminished the glycolytic action in U87, T98G human glioblastoma multiforme and MCF7 human breast carcinoma cells.

From the above, it is obvious that the metabolic collapse of T47D cells 1 h following cerco-PDT is in good agreement with the concomitant growth inhibition/shrinkage of the T47D spheroids which gradually become necrotic. In contrast the U87 spheroids, although their respiration breaks down following cerco-PDT, apparently their quasi-functional glycolytic ability, could be one key factor that helped the spheroids retain their viability and their capacity to grow up to the size of the nontreated controls by day 8, despite their initial decrease in size up to day 3. This is also supported from the results of our previous work on cerco PDT (5) where U87 present with the lowest respiration of the 3 cell lines investigated (T98G, MCF7 and U87) yet their glycolytic activity is comparable to that of MCF7 indicating that they rely more on glycolysis for ATP production (Warburg effect) (15) and hence survival since ATP deficiency leads to bioenergetic collapse and cell death (16,17).

Once again, in the present work, cercosporin was found to be a very potent photosensitizer. In the PDT field in general, the vast majority of the *in vitro* studies have been conducted in cell monolayers where cell killing, and cell-cell interactions where 3D structural considerations do not apply. It is therefore very important, especially for PDT applications where the light penetration into tissue is limited, to use 3D cultures which are

structurally more similar to solid tumors. Even though the short activation wavelengths of cerco are limiting for classical PDT application in the clinic, it may prove very useful for emerging applications as for example proton-dynamic therapy (10).

Acknowledgements—The present work was financially supported by the European Union's Horizon 2020 framework program for research and innovation under grant agreement No 712921 as well as by the SouthEastern Norway Regional Health Authority (Helse Sør Øst, project no. 2017116).

REFERENCES

- MacRobert, A. J. and T. Theodosiou (2005) Photodynamic therapy of cancer. In *Encyclopedia of Modern Optics* (Edited by R. D. Guenther), pp. 53–62. Elsevier, Amsterdam.
- Morgan, B. J., C. A. Mulrooney and M. C. Kozlowski (2010) Perylenequinone natural products: evolution of the total synthesis of cercosporin. *J. Org. Chem.* **75**, 44–56.
- Daub, M. E., S. Herrero and K. R. Chung (2013) Reactive oxygen species in plant pathogenesis: the role of perylenequinone photosensitizers. *Antioxid. Redox Signal.* **19**, 970–989.
- Daub, M. E., M. Li, P. Bilski and C. F. Chignell (2000) Dihydrocercosporin singlet oxygen production and subcellular localization: a possible defense against cercosporin phototoxicity in *Cercospora*. *Photochem. Photobiol.* **71**, 135–140.
- Mastrangelopoulou, M., M. Grigalavicius, K. Berg, M. Menard and T. A. Theodosiou (2019) Cytotoxic and photocytotoxic effects of cercosporin on human tumor cell lines. *Photochem. Photobiol.* **95**, 387–396.
- Sutherland, R., J. Carlsson, R. Durand and J. Yuhas (1981) Spheroids in cancer research. *Cancer Res.* **41**, 2980–2984.
- Bradley, D. and G. Roth (2007) Adaptive thresholding using the integral image. *J. Graphics Tools* **12**, 13–21.
- RStudio Team (2016) *RStudio: Integrated Development of R*. RStudio Inc., Boston, MA.
- R Core Team (2019) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Grigalavicius, M., M. Mastrangelopoulou, K. Berg, D. Arous, M. Menard, T. Raabe-Henriksen, E. Brondz, S. Siem, A. Gorgen, N. F. J. Edin, E. Malinen and T. A. Theodosiou (2019) Proton-dynamic therapy following photosensitizer activation by accelerated protons demonstrated through fluorescence and singlet oxygen production. *Nat. Commun.* **10**, 3986.
- Theodosiou, T., J. S. Hotherhall, E. A. Woods, K. Okkenhaug, J. Jacobson and A. J. MacRobert (2003) Firefly luciferin-activated rose bengal: in vitro photodynamic therapy by intracellular chemiluminescence in transgenic NIH 3T3 cells. *Cancer Res.* **63**, 1818–1821.
- Parsa, A. T., S. Wachhorst, K. R. Lamborn, M. D. Prados, M. W. McDermott, M. S. Berger and S. M. Chang (2005) Prognostic significance of intracranial dissemination of glioblastoma multiforme in adults. *J. Neurosurg.* **102**, 622–628.
- Theodosiou, T. A., A. Papakiriakou and J. S. Hotherhall (2008) Molecular modeling and experimental evidence for hypericin as a substrate for mitochondrial complex III; mitochondrial photodamage as demonstrated using specific inhibitors. *Free Radic. Biol. Med.* **45**, 1581–1590.
- Theodosiou, T. A., M. Ali, P. Dillard, K. O. Schink, K. E. Olsen, S. Wälchli, E. M. Inderberg, A. Kubin, Q. Peng and K. Berg (2018) Simultaneous defeat of mutually exclusive MCF7 and MDA-MB-231 treatment resistances by a hypericin PDT – tamoxifen hybrid therapy: HYPERTAM. *npj Breast Cancer* **5**, 13.
- Liberti, M. V. and J. W. Locasale (2016) The Warburg effect: How does it benefit cancer cells? *Trends Biochem. Sci.* **41**, 211–218.
- Lieberthal, W., S. A. Menza and J. S. Levine (1998) Graded ATP depletion can cause necrosis or apoptosis of cultured mouse proximal tubular cells. *Am. J. Physiol.* **274**, F315–F327.
- Kushnareva, Y. and D. D. Newmeyer (2010) Bioenergetics and cell death. *Ann N Y Acad. Sci.* **1201**, 50–57.