Research paper

RFU derived LR s for activity level assignments using Bayesian Networks Peter Gill ^{a,b,*}, Øyvind Bleka ^a, Ane Elida Fonnelløp ^a^a Forensic Genetics Research Group, Oslo University Hospital, Oslo, Norway^b Department of Clinical Medicine, University of Oslo, Oslo, Norway

ARTICLE INFO

Keywords:

Direct transfer
Secondary transfer
Mixtures
Likelihood ratio (LR)
Evidence evaluation
Bayesian Network
ALTRaPht
EuroForMix

ABSTRACT

A comparative study has been carried out, comparing two different methods to estimate activity level likelihood ratios (LR_a) using Bayesian Networks. The first method uses the sub-source likelihood ratio ($\log_{10} LR_\phi$) as a 'quality indicator'. However, this has been criticised as introducing potential bias from population differences in allelic proportions. An alternative method has been introduced that is based upon the total RFU of a DNA profile that is adjusted using the mixture proportion (M_x) which is calculated from quantitative probabilistic genotyping software (EuroForMix). Bayesian logistic regressions of direct transfer data showed that the two methods were comparable. Differences were attributed to sampling error, and small sample sizes of secondary transfer data. The Bayesian approach facilitates comparative studies by taking account of sampling error; it can easily be extended to compare different methods.

1. Introduction

In Gill et al. [1], we used sub-source likelihood ratios ($\log_{10} LR_\phi$) as a 'quality-indicator' to help assign activity level likelihood ratios (LR_a). Calculations of $\log_{10} LR_\phi$ s are partly dependent upon population allele frequency databases, with potential bias introduced by rare alleles; secondly, mixture composition may itself have an effect on the $\log_{10} LR_\phi$ without affecting the DNA contribution measured by RFU. It was proposed that a measurement based upon the quantity of DNA from a person of interest (POI) may be preferable. The purpose of this study is to compare such a method with the published $\log_{10} LR_\phi$ measurement. Probabilities of direct (T) and secondary transfer (S) are modelled by Bayesian logistic regression. An advantage is that sampling uncertainties are taken into account, which is important because sample sizes are small. We show how to compare different methods of measurement.

2. Materials and methods

Samples were processed and analysed as described in section 2. of [1]; a brief description follows:

1. A total of 33 participants were included in the study in different pair combinations
2. Direct transfer: Participant B contacted the arm of participant A, a minimum of one hour after participant B had washed hands. After the contact, participant A went about normal activities and samples were collected between 0–24 h after the initial

contact (only one sample collected per experiment). A total of 88 samples were collected.

3. Secondary transfer: A minimum of one hour after washing hands, participants A and B shook hands for 30 s and immediately stroked their own upper arm. Samples were taken between 0–24 h after contact
4. Powerplex® Fusion 6C multiplex was used to analyse the samples.

2.1. Notation

Details provided in [1] are briefly summarised here:

1. A sample is collected at time h_0 . All preceding transfer events are measured relative to the time of sampling.
2. A direct transfer event (T) conditioned under H_p , but disputed by H_d , occurs at time h_1 in hours.
3. Secondary transfer event(s) (S) occur at time h_2 in hours. This may be a range where the number of transfer events (n) are specified.
4. A direct transfer event (T') occurs at time h_3 . This is a social interaction such as a handshake and is accepted under H_p and H_d .
5. Activity level LR_a is used generically.
6. Activity level LR_{a_x} is conditioned on $\log_{10} LR_\phi > x|h_1$ where x is the logistic function decision threshold value.

* Corresponding author at: Forensic Genetics Research Group, Oslo University Hospital, Oslo, Norway.

E-mail address: peterd.gill@gmail.com (P. Gill).

7. Activity level LR_{a_y} is conditioned on sub-source $RFU_{POI} > y|h_1$, where y is the logistic function decision threshold value.

2.2. Quantitation of the DNA profile from peak height information

From [1], there are two sets of data simulating direct skin to skin contact that may occur in an assault. The first relates to direct transfer experiments, whereas the second data-set relates to the secondary transfer experiments. The data-sets have been expanded (Supplements 5,6) to include details of DNA quantity estimated from the sum of RFU values. To carry out this measurement, probabilistic genotyping software *EuroForMix* v.3.3.1 [2] was used to carry out mixture analysis. Mixture proportions (M_x) for the POI are automatically calculated by the software. For a given evidence profile, a script was used to summate all alleles and peaks (RFU_{tot}) above the analytical threshold of 100 RFU , across all (m) STR loci. This was converted into the mean RFU_{tot}/m per locus. Then the contribution to the mean RFU_{POI} was calculated by multiplying RFU_{tot}/m with the POI specific M_x . If the sample is diluted, then multiplication by the dilution factor will compensate the decreased RFU_{POI} , e.g. if one third of the sample has been aliquoted and loaded into the PCR thermocycler, then the dilution factor $d_l = 3$: note that none of the samples in this study were diluted ($d_l = 1$).

$$RFU_{POI} = M_x \times \frac{RFU_{tot}}{m} \times d_l \quad (1)$$

2.3. Bayesian logistic regression

All programming was carried out in R v. 4.0.3 [3]. To analyse the direct transfer results, Bayesian logistic regression, using the R *stan.glm* function [4] from package (*rstanarm*), utilising MCMC to generate 4000 pairs of coefficients per logistic regression to find $Pr(RFU_{POI} > y|h_1)$ where y is the threshold value tested and h_1 is the time-interval in hours between the transfer and collection of samples.

2.4. Exponential distribution fits to secondary transfer data

The combined data between time zero and one hours after deposition were modelled using an exponential distribution to determine $Pr(S|h_2 \in 0,1)$. The exponential distribution was determined to be the best fit after testing different options with the R package (*fitdistrplus*) [5] (supplement S3; Fig. S2).

2.5. Likelihood ratio calculations

Activity level LR_{a_y} was calculated using a modified Activity Level, Transfer, Results and Persistence program [1] (ALTRaPht); source files available at <https://sites.google.com/view/altrap/home>, along with the data-sets and user manual. The program can be launched directly from the 'shinyapps.io' server site <https://activitylevel.shinyapps.io/shinyaltrapht/> without any need to load R.

3. Results

3.1. Linear regression

Linear regression of the direct transfer data (where $\log_{10}LR_\phi > 0$ vs RFU_{POI}) was carried out (Fig. 1); Intercept = 21.7; Slope = 24.6. $R^2 = 0.8$. Higher values are typically associated with short time intervals between DNA transfer and sampling.

Table 1

$RFU_{POI} > y$ equivalents to $\log_{10}LR_\phi > x$ logistic regression thresholds, with mean square error (MSE) of the residuals.

$\log_{10}LR_\phi > x$	$RFU > y$ equivalent	MSE
1	75	3.0E-06
2	106	1.0E-05
3	137	2.0E-04
4	146	7.0E-04
6	172	3.0E-04
8	182	3.0E-04
10	237	5.0E-04
12	266	1.0E-04

3.2. Logistic regression

Linear regression does not take into account the time since transfer (h), and was insufficient to fully characterise the relationship of RFU and $\log_{10}LR_\phi$ relative to h . In the next step, logistic regressions of $RFU > y|h_1$ were fitted to regressions of $\log_{10}LR_\phi > x|h_1$. This was carried out by optimisation as follows:

Step 1: A table of logistic regression probabilities for $x = 1, \dots, 12$ was prepared (Supplement 7).

Step 2: To find the equivalent of $RFU_{POI} > y|h_1$ for a given x threshold (Table 1), optimisation was carried out in order to determine the values of y that gave the best fit to $\log_{10}LR_\phi > x|h_1$ logistic regressions. This was carried out by iterating y for a specific value of x in order to minimise the mean square error (MSE) i.e. the mean of the sum of squared residuals across all time points in hours (0,1, ...,24). The resultant logistic regressions were all good fits (Fig. 2; supplement S2; Table S1). The "Optimizer.R" program used to carry out the optimisation is available at <https://github.com/peterdggill/altrapHt.git>.

Step 3: The Activity Level, Transfer Recovery and Persistence (ALTRaP) program [1] was modified to accept RFU_{POI} values instead of $\log_{10}LR_\phi$ values.

Some further data exploration using optimisation is described in Supplement S1.

4. Discussion

4.1. Reworked example

In Gill et al. [1], the ALTRaP program was used to calculate activity level LR_{a_x} based upon sub-source $\log_{10}LR_\phi$ values. To illustrate the method a number of examples were described. Details of the first example are briefly summarised as follows: Sampling was at time $h = 0$, following an assault at time $h_1 = 3$ and social contact at $h_2 = 6, \dots, 10$ (one contact by secondary transfer each hour, i.e. $n = 1$); direct innocent contact e.g. handshake at time $h_3 = 10$. Activity level LR_{a_x} vs. LR_{a_y} , were calculated and compared (Table 2).

Activity level LR_{a_x} s are comparable for logistic thresholds $x = 1, 2, 3$ and y equivalents, but for higher values of x , the LR_{a_y} results plateau. Logistic regression coefficients for direct transfer are similar when $x > 8$ [1], because there are few observations, especially when $h_1 > 9$ h.

4.2. Investigation into effect of small sample size

An investigation was carried out to determine the reasons for the divergences between LR_{a_x} and LR_{a_y} values. LR_a is calculated from Bayesian logistic regression analysis of direct and secondary transfer data. Because of the very low success rate of secondary transfer, especially for time intervals > 1 h, only data immediately following the secondary transfer opportunity was analysed ($h_2 = 0, 1$) as described in section 8 of Gill et al. [1], except that data were fitted to an exponential distribution rather than Pareto. Preferred distributions, and associated

Fig. 1. Linear Regression analysis of direct transfer data $\log_{10} LR_{\phi}$ vs. M_x adjusted mean peak height of the POI (RFU_{POI}).

Fig. 2. Logistic Regression analysis of direct transfer data, Time vs. Transfer probability, where each panel compares $\log_{10} LR_{\phi} > x|h_1$ with $RFU_{POI} > y|h_1$ equivalents listed in Table 1.

parameters, were determined by diagnostic tests using the *fitdistrplus* R-package (Supplement S3; Fig. S2).

From Bayesian logistic regressions, quantiles were tabulated in rows (section 14.5 of [1]), corresponding to quantiles derived from 1000x bootstraps of the secondary transfer probabilities $Pr(S)$ (see section

14.6 of [1]). Briefly: data collected from zero — one hour after the secondary transfer experiment, were subject to bootstrapping [6] in order to generate 1000 new randomised sets of samples. For each data-set, new exponential distribution parameters were estimated, and $Pr(S|y)$ was determined as described in [1] (section 8 and fig 6).

Table 2

Comparison of activity level LR_{a_x} and LR_{a_y} using logistic regression thresholds $\log_{10} LR_{\phi} > x$ and the equivalent $RFU_{POI} > y$. Direct transfer (T) $h_1 = 3$; secondary transfer (S) $h_2 = 6, \dots, 10$, with one contact per hour ($n = 1$); direct (innocent) transfer (T ϵ) $h_3 = 10$. For full details see Gill et al [1] Table 3.

$Pr(E Hp)$ proposition	Type of transfer		T and S		(T, S and T ϵ)	
	$\log_{10} LR_{\phi} > x$	$RFU > y$	LR_{a_x}	LR_{a_y}	LR_{a_x}	LR_{a_y}
POI only	1	75	6	8	3	4
	2	106	10	12	4	5
	3	137	18	16	5	5
	4	146	33	19	6	5
	6	172	43	22	5	5
	8	182	72	24	5	4
POI and unknown(s)	1	75	1	2	1	1
	2	106	3	3	1	1
	3	137	5	5	1	1
	4	146	10	6	2	2
	6	172	16	8	2	2
	8	182	29	9	2	2

Table 3

Secondary transfer data were bootstrapped to generate 1000 new randomised sets of samples. Exponential distributions were estimated for each to determine $Pr(S|x)$ and the range was expressed by quantiles in the rows of the table. Respective $Pr(S|x, y)$ values were plugged into Bayesian logistic regressions that generated the quantiles denoted by the columns. Figures highlighted in grey are the median LR_{a_y} results. The figures highlighted in orange indicate the LR_{a_x} equivalent values.

Logistic regression threshold	Secondary transfer quantiles	Logistic regression quantiles		
		0.25	0.5	0.75
$\log_{10} LR_{\phi} > 4$; 146 RFU	0.25	22	31	43
	0.50	14	19	27
	0.75	10	14	19
$\log_{10} LR_{\phi} > 6$; 172 RFU	0.25	29	41	59
	0.50	16	23	32
	0.75	11	16	22
$\log_{10} LR_{\phi} > 8$; 182 RFU	0.10	59	82	119
	0.25	30	42	60
	0.50	17	23	33
	0.75	11	16	22

Quantiles that describe the range of $Pr(S|y)$ were generated from the 1000x bootstrapped samples. The second step was to substitute a value of $Pr(S|y)$ in the Bayesian logistic regression calculation, where 4000 pairs of data-dependent simulated coefficients were generated in order to calculate likelihood ratios. These data are also described by quantiles. Following these two steps, a plausible range of likelihood ratios can be prepared. In particular, we are interested in the range of LR_{a_y} defined by 0.25–0.75 quantiles (Table 3) which can be described as encompassing the plausible range of activity level likelihood ratios.

Table 3 shows an analysis for the example described in Table 2: the grey cells are median values of LR_{a_y} ; the orange cells show the nearest values associated with the LR_{a_x} for a given logistic regression threshold (x) that lies within the 0.25–0.75 quantiles. Greater divergence is observed when $x = 8$; $y = 182$, but there are few values observed in the data sets. In conclusion, the differences in LR_{a_x} vs. LR_{a_y} can be explained by sampling error since their respective values fall within the 0.25–0.75 quantiles of direct transfer logistic regressions and secondary transfer bootstrapped samples.

4.3. Reporting

Here we have described two alternative methods to assign LR_a from the same data. Differences may be attributed to sampling.

There are a number of ways that we could proceed: the simplest option would be to adopt a single method, i.e. the most informative method chosen after further comparative studies. Using both methods concurrently is possible, but this raises the question of how to report?

As an analogy with probabilistic genotyping, it is useful to refer to the debate arising from [7,8] and subsequent discussions by Taylor et al. [9], where the latter suggest reporting an average LR where two or more (comparable) methods are employed. When sensitivity analysis is carried out, reporting the median value has a good consensus — see discussion in [1], section 14.9.

As explained by Taroni et al. [10], the issue is not uncertainty of the likelihood ratio, rather it is a question of precision and communication. The use of 0.25 – 0.75 quantiles can be used as a sensitivity analysis to provide a plausible range of LR_a values. The authors suggest that one way to simplify communication, would be to avoid the use of numbers completely. However, this idea runs counter to the ISFG DNA commission [11] which recommends that the numeric likelihood ratio as compulsory, and the verbal analogue as optional. The precision of the results from sensitivity analysis suggests that the reported value of LR_a should be rounded down to multiples of ten, unless it is less than ten.

In Supplement S4 further examples are explored, taken directly from [1]. The trend for lower LR_{a_y} compared to LR_{a_x} values is repeated. If this is mainly attributable to sampling error of secondary transfer data-sets, then larger sample sizes should improve the comparability of the results and reduce the observed variance, thereby increasing precision.

4.4. Conclusion

This is part of a study to lay the ground-work for a much larger project which will begin in 2022, entitled: “Establishment of a Trace DNA Transfer Rate Repository and Bayes Nets to Calculate LRs (ReAct)”. Funded by the European Network of Forensic Science Institutes (ENFSI) monopoly project, there are more than 30 participating laboratories. This will provide an opportunity to generate new sets of data that will help to optimise the methodology and experimental design, in addition to facilitating much needed tests of reproducibility. It is anticipated that the “Optimizer.R” program may be useful to compare logistic regressions from different laboratories to determine equivalence factors as described in Section 3.2. Taking the mean RFU measurement across loci facilitates comparisons across different multiplexes. To measure M_x , a laboratory will need to use qualitative (continuous) probabilistic genotyping software. Comparative studies are needed to address complex issues related to standardisation: where the aim is to show whether different laboratories following similar methods, report similar LR_a s for given conditions.

Appendix A. Supplementary data

Supplementary material related to this article can be found online at <https://doi.org/10.1016/j.fsigen.2021.102608>.

References

- [1] P. Gill, Ø. Bleka, A. Roseth, A.E. Fonneløp, An LR framework incorporating sensitivity analysis to model multiple direct and secondary transfer events on skin surface, *Forensic Sci. Int. Genet.* 53 (2021) 102509.
- [2] Ø. Bleka, G. Storvik, P. Gill, *EuroForMix*: an open source software based on a continuous model to evaluate STR dna profiles from a mixture of contributors with artefacts, *Forensic Sci. Int. Genet.* 21 (2016) 35–44.
- [3] R Core Team, R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, 2013, URL <http://www.R-project.org/>.
- [4] C. Muth, Z. Oravecz, J. Gabry, User-friendly Bayesian regression modeling: A tutorial with rstanarm and shiny, *Quant. Methods Psychol.* 14 (2) (2018) 99–119.
- [5] M.L. Delignette-Muller, C. Dutang, et al., Fitdistrplus: An R package for fitting distributions, *J. Stat. Softw.* 64 (4) (2015) 1–34.
- [6] B. Efron, R.J. Tibshirani, *An Introduction to the Bootstrap*, CRC Press, 1994.
- [7] E. Alladio, M. Omedei, S. Cisana, G. D’Amico, D. Caneparo, M. Vincenti, P. Garofano, DNA mixtures interpretation—a proof-of-concept multi-software comparison highlighting different probabilistic methods’ performances on challenging samples, *Forensic Sci. Int. Genet.* 37 (2018) 143–150.

- [8] P. Garofano, D. Caneparo, G. D'Amico, M. Vincenti, E. Alladio, An alternative application of the consensus method to DNA typing interpretation for low template-DNA mixtures, *Forensic Sci. Int. Genet. Suppl. Ser. 5* (2015) e422–e424.
- [9] D.A. Taylor, J.S. Buckleton, J.-A. Bright, Comment on “DNA mixtures interpretation—A proof-of-concept multi-software comparison highlighting different probabilistic methods’ performances on challenging samples” by Alladio et al., *Forensic Sci. Int. Genet.* 40 (2019) e248–e251.
- [10] F. Taroni, S. Bozza, A. Biedermann, C. Aitken, Dismissal of the illusion of uncertainty in the assessment of a likelihood ratio, *Law, Probab. Risk* 15 (1) (2016) 1–16.
- [11] P. Gill, T. Hicks, J.M. Butler, E. Connolly, L. Gusmão, B. Kokshoorn, N. Morling, R.A. van Oorschot, W. Parson, M. Prinz, et al., DNA commission of the international society for forensic genetics: Assessing the value of forensic biological evidence-guidelines highlighting the importance of propositions: Part I: evaluation of DNA profiling comparisons given (sub-) source propositions, *Forensic Sci. Int. Genet.* 36 (2018) 189–202.