PHD thesis

**Secondary hyperparathyroidism after obesity surgery
- exploring the impact of vitamin D and calcium,
and aspects on bone health**

Based on 4 observational studies

by Stephen Hewitt

Institute of Clinical Medicine

Faculty of Medicine

University of Oslo

Department of Endocrinology, Morbid Obesity and Preventive Medicine

Oslo University Hospital

© **Stephen Hewitt, 2022**

Secondary hyperparathyroidism after obesity surgery
- exploring the impact of vitamin D and calcium, and aspects on bone health

With basis on 4 observational studies

*Series of dissertations submitted to the
Faculty of Medicine, University of Oslo*

ISBN 978-82-8377-973-8

All rights reserved. No part of this publication may be
reproduced or transmitted, in any form or by any means, without permission.

Cover: Hanne Baadsgaard Utigard.
Print production: Reprosentralen, University of Oslo.

Table of contents

Table of contents	i
Foreword.....	ii
Acknowledgements.....	iv
Abbreviations:	vi
List of publications	vii
Sammendrag (Norwegian abstract)	viii
 Part I	 1
1. Perspective.....	3
2. Background	5
3. Secondary hyperparathyroidism after obesity surgery.....	15
 Part II	 23
4. Aims of thesis	24
5. Materials and methods	27
6. Summary of papers	33
 Part III	 43
7. Methodological considerations.....	45
8. Discussion.....	55
9. Conclusions	69
10. References.....	71
 Supplementary materials.....	 87
 Papers included – Study 1-4.....	 91

Foreword

I may admit that some of my thoughts around the theme of this thesis, arose during my early years in training in internal medicine. In most of these years, parathyroid hormone (PTH) was tested rather occasionally. The blood test results usually arrived after the patients had recovered and were discharged from the hospital, and sometimes they may have drowned among all the other work to be done. Often, an elevated PTH received no further attention with normal calcium and vitamin D levels, and it remained somewhat peculiar to me.

Following patients after obesity surgery, certainly reactivated previous wondering, obliged as I was to handle the results that flowed into my office. Frequent among these, were the results of PTH, vitamin D and calcium levels, which arrived on separate printings from the Hormone Laboratory.

The elevations of PTH were frequent in these patients. Reasoning back to previous teaching and reading, I had learned that blood calcium was the major determinant of PTH secretion. Moreover, vitamin D was subject for growing attention at our endocrinologic department at the time. Both calcium and vitamin D levels were suspected to influence the hyperparathyroidism I was to investigate.

I found myself pondering over the blood test results. My real scientific interest had started in late years of medical school. This interest increased during my early years with Leiv Ose and colleagues at the Lipid Clinic in Oslo, and my collaboration with Sidsel Graff-Iversen at the National Institute of Public Health. Clinical practice had also taken much of my concentration and effort the previous years.

Sitting in my office, I started to group the PTH results by vitamin D status. The discussion around secondary hyperparathyroidism (elevated PTH) after obesity surgery these years was about vitamin D and bone health, and calcium was rather seldomly coupled with PTH. Although the main hypothesis in the first study turned negative, it did pave the way for exciting research.

There has been development within the field during these years. First of all, several guidelines for obesity surgery have adopted or mentioned higher levels for vitamin D sufficiency, in line with our main research objective. Yet, our subsequent studies are large in the field and still

among few indicating difference by the specified vitamin D levels. Secondly, several reviews were published in the years 2012-2017, adding valuable thoughts around bone issues after obesity surgery, but all were limited to the first 2-3 postoperative years. Quite few have concentrated on vitamin D and PTH. In addition, the consecutive studies of later bone effects, were highly relevant to discuss in this thesis.

Writing a PhD thesis is a journey. The world looks different compared to the time we started the studies, and the literature and scientific world have expanded further to me in the writing process. Some new facts and nuances in my interpretations were natural to include in the introduction part. Most that was not yet acknowledged in our first years, has been reserved for later chapters. I may have wanted to present the situation how it looked like, and let the readers follow the reasoning I had to follow.

Part II presents our hypotheses, objectives, methods, materials and study results. The studies 1-3 analysed the potential role of higher vitamin D levels on secondary hyperparathyroidism, and they included considerations of calcium levels. Study 2-4 investigated the potential impact on bone health. It was natural to delimit study 2-4 for patients with gastric bypass only, being the dominant procedure, and due to the differences between procedures.

The discussion evaluates the results in our current perspective and intends to give a relevant update on the topic. Calcium and vitamin D levels are considered important factors in secondary hyperparathyroidism, which in turn may impact bone health.

It is hoped that our studies provide a valuable scientific approach to this topic. The thesis also aims to present and discuss the relevant debate around secondary hyperparathyroidism in the context of obesity surgery, and gastric bypass in particular. Some of the findings could possibly apply to other populations.

Acknowledgements

The thesis is based on my clinical work at the Centre of Morbid Obesity and Bariatric Surgery, a section of The Department of Endocrinology, Morbid Obesity and Preventive Medicine at Oslo University Hospital Aker. I was privileged to be positioned here for such a project. Researchers around were enthusiastic, and questions around many. The barriers to start registration was low, as the patient logistics and allowances were all ready. I am also grateful to the Faculty of Medicine of approving this as a PhD project.

My greatest gratitudes go to all my supervisors, who formed the sphere I needed to do this project. They embraced my work from early on, and showed full trust during the period. This was particularly important after study 1, which disclosed no difference by vitamin D status.

Main supervisor Tom Mala, now professor, has showed extraordinary skills and dedication to science, besides being a full time clinical surgeon. He managed responsibly all our duties towards the Faculty. His qualities of supportiveness, gentleness and follow-up are of few.

Centre leader and co-supervisor Jon Kristinsson has revealed a fundamental commitment to research. He readily welcomed the project and facilitated for all my needs. Decisive was his faithfulness and protection over the project. He made smart remarks and showed patience.

Co-supervisor professor Erik Fink Eriksen deserves special appreciation for bringing the research issue of vitamin D into my mind. His exciting lectures and knowledge created a seed that was so essential in my new conundrum. I am grateful for his engagement and expertise.

Professor Jørgen Jahnsen at Department of Gastroenterology (later at Ahus) receives my best gratitude for providing experienced feedback from his background in PTH and bone. Professor Thomas Bøhmer participated with great interest and shared willingly from his field of nutrition and endocrinology, and provided extensive feedback. Co-operating with Grethe Støa-Birketvedt was also of great value the first years.

Torgeir Søvik and Erlend Aasheim ended their PhDs during my first years at the centre. Much of the drive can be attributed to their spirits and achievements. It put our centre on the world map. Torgeir set a tone of enthusiasm and dedication, while the devotion of Erlend displayed internist considerations. As co-authors, they shared of their records.

During the years, I was inspired by Hilde Risstad, being open-minded, energetic, clever and a quick learner. Ingvild Blom-Høgestøl received a devoted role in the 10-year bone study. Our tone and collaboration are still highly appreciated, and made this study a part of my thesis. Huge credit goes also to Monica Chahal-Kummen who organized the patient logistics in this 10-year study, being a sub-study of a more comprehensive project. All doing a PhD.

The Department of Endocrinology has been a particular important arena for clinical learning and scientific discussion. Much of my current understanding has been formed after meetings with individuals here. Working at the Osteoporosis Clinic has been fundamental for my clinical perspective on bone issues. My special gratitude goes to Head of Department Lene Seland, to previous Head Kåre Birkeland, and to Gøril Vinje and Maida Huskovic who carried out the DXA scans.

The Hormone Laboratory performed the main blood determinations in these studies, and our close connection was essential for the work. Per Thorsby and Sandra Dahl contributed in some of the laboratory issues. Solve Tjora also helped from the Central Laboratory.

The statisticians at Oslo Centre for Biostatistics and Epidemiology (OCBE) receive my best acknowledgements. Lien My Diep supported with analyses in study 2. Ole Klungsøyr helped me with mixed model analyses in study 3, and professor Magne Thoresen acknowledged the final model. Cathrine Brunborg helped analyse in study 4.

Our clinical dieticians are to be acclaimed for their role in the nutritional handling of our patients. Susanna Hanvold and Eline Birkeland have worked with me throughout my period, and both completed their PhDs. Eline receives particular tribute for her willingly sharing of formality and technical issues in the final months.

Special nurses have an important role in our centre milieu. The laughter and unifying jokes of Ann Steen survives years after retirement. Inger Eribe displays mother's instinct towards us all, and has contributed with essential support in the research. Anita Sunde has received a trusted role in our patient care. The psychologists, Eva Hjalmsdotter and Tina Avantis, cared for selected patients. Among the secretaries, Marianne Sæter receives honour for her role as the logistic chief, handling several practical issues in close co-operation with Anneli Skulbørstad, Iren Johannessen and Allette Dahler.

Abbreviations:

BMI: Body mass index

PTH: Parathyroid hormone

SHPT: secondary hyperparathyroidism

Ca²⁺: free ionized calcium

S-25(OH)D: serum concentrations of 25-hydroxyvitamin D (calcifediol)

1,25(OH)₂D: 1,25-dihydroxyvitamin D, active vitamin D

ALP: Alkaline phosphatase (total)

B-ALP: Bone specific alkaline phosphatase

PINP: procollagen type 1 propeptides

CTX: carboxy-terminal collagen crosslinks

LC-MS/MS: liquid chromatography-mass spectrometry

CRF: Case report form

DXA: dual-energy X-ray absorptiometry

BMD: bone mineral density

Low BMD: DXA based t-/z-score < -1.0 according to menopausal status or male age

QCT: quantitative computer tomography

HR-qCT: High-resolution quantitative computer tomography

OR: Odds ratio

RR: Relative ratio

GEE: Generalized estimated equations

ASMBS: American Society for Metabolic and Bariatric Surgery

BOMSS: British Obesity and Metabolic Surgery Society

SOREG: Scandinavian Obesity surgery Registry

List of publications

Paper 1 (Study 1)

Secondary Hyperparathyroidism, Vitamin D Sufficiency, and serum Calcium 5 years after Gastric Bypass and Duodenal Switch.

Hewitt S, Søvik TT, Aasheim ET, Kristinsson J, Jahnsen J, Birketvedt GS, Bøhmer T, Eriksen EF, Mala T.

Obes Surg. 2013 Mar; 23(3): 384-90. doi: 10.1007/s11695-012-0772-3.

Paper 2 (Study 2)

Relationships of serum 25-hydroxyvitamin D, ionized calcium and parathyroid hormone after obesity surgery.

Hewitt S, Aasheim ET, Søvik TT, Jahnsen J, Kristinsson J, Eriksen EF, Mala T.

Clin Endocrinol (Oxf). 2018 Mar; 88(3): 372-379. doi: 10.1111/cen.13531.

Paper 3 (Study 3)

Relationships between Vitamin D Status and PTH over 5 years after Roux-en-Y Gastric Bypass: A Longitudinal Cohort Study.

Hewitt S, Kristinsson J, Aasheim ET, Blom-Høgestøl IK, Aaseth E, Jahnsen J, Eriksen EF, Mala T.

Obes Surg. 2020 Sep; 30(9) 3426-34. doi: 10.1007/s11695-020-04582-5

Paper 4 (Study 4)

Bone metabolism, bone mineral density and low-energy fractures 10 years after Roux-en-Y gastric bypass.

Blom-Høgestøl IK, Hewitt S, Chahal-Kummen M, Brunborg C, Gulseth HL, Kristinsson JA, Eriksen EF, Mala T.

Bone. 2019 Oct; 127: 436-445. doi: 10.1016/j.bone.2019.07.014.

Sammendrag (Norwegian abstract)

I 2004 ble fedmekirurgi introdusert i Norge, med gastrisk bypass som en dominerende prosedyre. Ved gastrisk bypass blir en liten magesekk på ca 25 ml konstruert og koplet på tynntarmen, og maten passerer over i tarmen lenger ned enn normalt. Den forminskede magesekken begrenser matinntaket, og absorpsjon forhindres i den forbikopledede tarmdelen. Duodenal omkopling er en prosedyre med større forbikopling av tynntarmen og mer nedsatt absorpsjon, og ble utført hos noen utvalgte med størst grad av fedme.

Vi fant ofte økte blodverdier av parathyroideahormon (PTH) etter slike inngrep. Ved normal regulering av kalsiummetabolismen vil lave nivåer av kalsium og/eller lave nivåer av vitamin D øke PTH og forårsake sekundær hyperparathyroidisme (SHPT). SHPT opprettholder stabile kalsiumnivåer via 3 mekanismer: 1) redusert utskillelse av kalsium i nyrer, 2) økt absorpsjon av kalsium i tarmen, og 3) mobilisering av kalsium fra skjelettet.

Litteraturen var uklar på hvordan vi skulle håndtere pasienter i forhold til vitamin D, kalsium og SHPT. Ofte var kalsium- og D-vitaminsnivåene normale og tilfredsstillende. Vi kartla derfor forekomsten av SHPT i vår populasjon, og undersøkte om det var noen sammenheng med disse nivåene. Hovedhypotesen var at forekomsten av SHPT ville være lavere med nivåer av 25-hydroxyvitamin D (S-25(OH)D) ≥ 75 nmol/l, sammenlignet med nivåer under dette. Vi undersøkte videre om det var en sammenheng mellom kalsium- og PTH-nivåer, og hvorvidt økte PTH-verdier påvirket benmarkører og skjelettstatus.

I den første studien (studie 1) fant vi SHPT hos 40% av våre 114 pasienter 5 år etter gastrisk bypass, og blant samtlige 11 etter duodenal omkopling. Vi fant ingen forskjell i PTH etter nivåene av D-vitamin, men SHPT forekom hyppigere ved kalsiumnivåer i det nedre området.

Studie 2 viste en sammenheng mellom SHPT og nivåene av både vitamin D og kalsium blant 502 pasienter 2 år etter gastrisk bypass. SHPT forekom hos 34% av pasientene. Pasienter med S-25(OH)D nivåer < 50 nmol/l hadde høyest forekomst av SHPT (45%). Nivåer med S-25(OH)D ≥ 50 og ≥ 75 nmol/l var begge assosiert med lavere og sammenlignbar forekomst av SHPT (henholdsvis 29% og 28%). Lavest forekomst hadde pasienter med S-25(OH)D nivåer ≥ 100 nmol/l (16%), og disse hadde de mest normale kalsiumverdiene. Kalsiumnivåer i det nedre normalområdet og lavere var assosiert med økt forekomst av SHPT også i denne studien.

S-25(OH)D < 50 nmol/l og SHPT var begge assosierte med høyere alkalisk fosfatase (ALP), forenlig med benpåvirkning.

I studie 3 studerte vi PTH og SHPT over 5 år hos 554 gastrisk bypass pasienter. Forekomsten av SHPT var 38% etter 5 år. Både kalsium- og D-vitaminnivåene viste en tydelig sammenheng med endringer i PTH. PTH var gjennomgående høyest og økte mest hos pasienter med S-25(OH)D < 50 nmol/l fra 6 måneder til 5 år postoperativt. PTH og SHPT økte i begge gruppene med S-25(OH)D 50-74 nmol/l og S-25(OH)D 75-99 nmol/l, der forskjellene imellom disse var små. Pasienter med S-25(OH)D nivåer ≥ 100 nmol/l hadde også i denne studien den laveste forekomsten av SHPT, og PTH forble lav over tid. SHPT var hyppigere ved kalsiumverdier i nedre tredjedel av normalområdet eller lavere. Det var en klar sammenheng mellom SHPT og høyere total-ALP og benspesifikk ALP, og vedvarende økt benspesifikk ALP over 5 år.

Studie 4 viste lavere bentetthet enn normalt (forventet) blant 122 pasienter 10 år etter gastrisk bypass. Vi registrerte SHPT hos 31%. Bentettheten var lavere ved SHPT, økende alder, postmenopause, høyere benmarkører og preoperativ hypotyreose. Osteoporose forelå hos 27% av postmenopausale kvinner og menn ≥ 50 år, og 6% av de yngre hadde en tilsvarende lav bentetthet. 15% rapporterte lavenergibrudd i løpet av de 10 årene postoperativt, og 8% fikk påvist virvelbrudd på sidebildet. Pasientene med osteoporose hadde gjennomgått flere lavenergibrudd.

Samlet viste funnene en sammenheng mellom D-vitamin/kalsiumnivåene og forekomsten av SHPT etter gastrisk bypass. S-25(OH)D nivåer ≥ 50 nmol/l viste lavere forekomst av SHPT og bedre kalsiumverdier, mens forekomsten av SHPT var lavest ved nivåer ≥ 100 nmol/l. Økt PTH var assosiert med høyere benomsetning og lavere bentetthet.

Å sikre S-25(OH)D nivåer ≥ 50 nmol/l, er nødvendig for å begrense forekomsten av SHPT, og har positive effekter på kalsiumbalansen og benhelsen. Våre studier ga lite støtte til at nivåer ≥ 75 nmol/l var mer effektive enn nivåer ≥ 50 nmol/l med hensyn på PTH, kalsium og ALP. Dataene taler imidlertid for at S-25(OH)D nivåer ≥ 100 nmol/l kan senke forekomsten av SHPT ytterligere etter gastrisk bypass, som igjen kan ha positive effekter på fremtidig benhelse.

Part I

Introduction

1. Perspective

The prevalence of overweight and obesity have increased in Norway and worldwide over the last decades, reaching epidemic proportions (1-3). Obesity is a major health challenge, as it increases the risk of metabolic disorders and obesity related diseases (2, 4). Weight loss surgery (obesity surgery) was introduced as a treatment option for patients with morbid obesity in Norway in 2004. Surgical complications were reduced, and studies showed significant and sustained weight loss (4, 5).

The classical aim of obesity surgery is to limit the food intake and the absorption of caloric nutrients from the intestinal tract. Reduced intake and reduced absorption both contribute to weight loss. The surgery related consequences include risk of nutritional deficiencies due to the altered gastro-intestinal anatomy (6).

A frequent observation in our follow-up after obesity surgery, was the elevated blood concentrations of parathyroid hormone (PTH). Calcium levels were typically normal, as were vitamin D levels and other factors influencing PTH secretion. The clinical workup of high PTH was challenging, and it had a potential to cause bone loss (7, 8).

Vitamin D status influences PTH secretion, and is important for bone health. The ongoing vitamin D debate accelerated from around 2011, with new vitamin D recommendations. In the US, the Endocrine Society defined a higher threshold level for vitamin D sufficiency, with 25-hydroxyvitamin D levels ≥ 75 nmol/l, while others defined sufficiency by levels ≥ 50 nmol/l (9-12).

Different guidelines adopted both these target levels. The evidence and clinical relevance underlying these cut-off values were however uncertain after obesity surgery, and some characterized the vitamin D recommendations as expert opinions. We postulated that the higher vitamin D levels of at least 75 nmol/l would have beneficial effects on PTH.

2. Background

Parathyroid hormone

Parathyroid hormone (PTH) is a peptide hormone produced in 4 small pea-sized parathyroid glands, located at the throat behind the thyroid gland (Figure 1). PTH is stored in the chief cells of the glands. *Intact PTH* is the active natural form of PTH. It consists of 84 amino acids (PTH 1-84) and is rapidly released into the blood when ionized (free) calcium levels decrease below a certain setpoint (7, 8).

Figure 1. Parathyroid hormone (PTH) is produced and stored in the parathyroid glands and secreted into the blood. The four small parathyroid glands are located at the throat on the backside of the thyroid gland. From Servier Medical Art, with permission.

The most important effect of PTH in human physiology is to increase calcium levels (8). The kidneys and bone cells are the prime target organs, exhibiting abundant receptors which can bind PTH. These receptors traverse the cell membrane, and are coupled with an intracellular G-protein complex. The extracellular binding of PTH thereby induces signalling within the cells via second messengers, such as calcium, cyclic AMP and inositol triphosphate (13). PTH can via such mechanisms regulate the calcium levels (8). Within minutes, higher PTH levels can increase blood calcium, by reducing loss of calcium in urine and mobilizing calcium from bone (14, 15). On the longer term, PTH stimulates calcium absorption in the intestine due to its ability to activate vitamin D.

Calcium

Calcium is an essential trace element in our body, and one of the 118 chemical elements in the periodic system. It is typically ingested in milk and dairy products. Most is bound in bone (99%). Less than 1 % is solubilized in blood, with about a half as free ionized calcium (Ca^{2+}), and the other half is bound to proteins (mostly albumin, 40%) or other compounds (15, 16). Calcium is crucial for a wide array of cellular functions, such as membrane function,

intracellular signal transduction, muscle contraction, cardiac function, nerve signals and hormone secretion (7). The concentrations in blood and extracellular fluid are therefore kept stable within a narrow range of about 0.02 mmol/l. Calcium is analysed by determinations of the blood concentrations of total calcium, albumin-corrected calcium or ionized calcium.

Calcium is absorbed throughout the intestine by active and passive transport (17, 18). The active transport occurs predominantly in the small intestine influenced by hormones and nutrients. High amounts of calcium within the intestine increase the passive absorption by paracellular diffusion, while active transport may dominate particularly when calcium intake is low (17).

It is well established that calcium levels in the extracellular fluid influence PTH secretion from the parathyroid glands. This occurs as ionized calcium binds to receptors on the surface of the parathyroid cells (chief cells) (7, 14, 19). Calcium sensing receptors are also prevalent in the kidney tubular system, and they are present in bone cells and in the gastrointestinal mucosa (15, 20). Small changes in the serum calcium levels can thereby induce compensatory responses via receptors in all these organs to keep calcium levels stable. As PTH can increase calcium levels, it plays a major role in the fine-tuned feedback mechanisms that stabilize calcium concentrations and support normal physiology (14).

Vitamin D

Vitamin D is primarily produced from cholesterol in the human body. Sunlight (UVB radiation) is a critical rate limiting catalyzer in this process, and cholesterol is transformed in the skin to vitamin D (cholecalciferol, vitamin D₃), which passes into the blood (17, 21, 22). Around 10-20% of vitamin D is ingested in food products such as fat fish, fish products, egg yolk, enriched dairy products and nutritional supplements, as vitamin D₂ or vitamin D₃. Vitamin D is thereafter converted to 25-hydroxycholecalciferol (25-hydroxyvitamin D) in the liver by a hydroxylation (22, 23). This form of vitamin D is the most stable, and is transported in the blood bound to proteins. The functional form of vitamin D is formed in the kidneys by a second hydroxylation to *active vitamin D*, 1,25-dihydroxyvitamin D (1,25(OH)₂D, called calcitriol) (22). PTH impacts this conversion. Active vitamin D binds to vitamin D receptors at the cell membranes. The binding induces rapid responses via second messengers such as cyclic AMP

in the cells, and the vitamin D receptor complex becomes internalized into the cells, and can interact with nuclear receptors and influence genomic pathways (18, 21, 22). Determination of the serum concentration of *25-hydroxyvitamin D (S-25(OH)D)* is the standard when assessing vitamin D status, due to its long half-life in blood (11, 24).

The classical effects of vitamin D are to increase calcium levels and promote mineralization of bone (7, 22). Active vitamin D increases gastro-intestinal absorption of calcium and reduces calcium excretion in the urine, which both increase calcium levels (21, 22, 25). Low vitamin D therefore predisposes for low calcium levels, while an increase in vitamin D levels can raise the calcium levels, as a consequence of changes in intestinal absorption and excretion in kidneys. It follows that vitamin D can influence PTH secretion indirectly via the calcium concentrations. There is also a direct effect of active vitamin D binding to receptors on the parathyroid cells, exerting a direct negative feedback on PTH secretion (8, 22). Consequently, low vitamin D levels stimulate production and secretion of PTH, while high levels have an inhibitory effect on PTH.

There are also vitamin D receptors located on several cells throughout the body, which can convert 25-hydroxyvitamin D to active vitamin D intracellularly. This may explain the many additional vitamin D effects, that have emerged in recent years.

Figure 2. Main effects of vitamin D influencing PTH. Active vitamin D increases absorption of calcium in the intestine, it decreases loss of calcium in the kidneys, and it facilitates the mineralization of calcium into bone. Secretion of PTH increases in response to low vitamin D or low calcium levels, and can mobilize calcium from bone to blood. Changes in vitamin D levels can influence PTH and bone by these mechanisms. From Servier Medical Art, with permission.

PTH disorders and secondary hyperparathyroidism

Several disorders influence the balance of PTH. PTH disorders can be categorized as *hyperparathyroidism* in cases with high PTH levels, or hypoparathyroidism when PTH is low. Hypoparathyroidism is further classified as primary or secondary, and hyperparathyroidism as primary, secondary or tertiary, depending on the nature of the abnormality (7).

In hypoparathyroidism, PTH secretion and serum PTH levels are low. Defects in the parathyroid gland function may lower or stop PTH secretion (primary hypoparathyroidism). In these cases, calcium is low. When the gland function is normal, high calcium levels can suppress PTH production and secretion via the negative feedback mechanism (secondary hypoparathyroidism)

Of the hyperparathyroid disorders, primary hyperparathyroidism has received most attention, due to the clinical effects with renal stones and bone loss, and much of our knowledge of the PTH effects comes from research of this disorder. Its feature consists of a combination of high concentrations of both PTH and calcium (7). Enlarged parathyroid glands (adenomas) can secrete excessive amounts of PTH, which increase calcium levels. In *secondary hyperparathyroidism (SHPT)*, the parathyroid gland function is normal, and the secretion of PTH is increased in response to external factors, such as low calcium, and/or low vitamin D (7, 19). Tertiary hyperparathyroidism may be considered a combination of primary and secondary hyperparathyroidism, in which enlarged hyperplastic glands or adenomas lose some of the negative feedback mechanism after longstanding stimulation with low calcium (7, 15). This is primarily seen in renal failure patients.

SHPT is the focus of this thesis. SHPT is commonly found in clinical practice, typically with vitamin D deficiency, malnutrition and disorders with reduced gastro-intestinal absorption and renal failure (7). It occurs more often with higher age. In Tromsø (Northern Norway), a study described a prevalence of 10% in a control-group of postmenopausal women aged 54-94 years without fractures (26).

Other factors

Substances such as phosphate and magnesium can also affect PTH secretion (14). Studies of renal failure support that high phosphate levels increase PTH. Hyperparathyroidism may on the contrary increase the renal excretion of phosphate. Active vitamin D increases the intestinal absorption of phosphate, together with calcium. Magnesium may influence PTH via several mechanisms, which seems related with its ability to modulate both the calcium receptor and the PTH receptor responses, but the response pattern on PTH remains somewhat inconsistent.

Bone physiology and bone health

The skeleton supports the body's full stature and protects softer and more vulnerable organs, and it secures mobility through anchoring of muscles. The bones and the bone cells play important roles in mineral homeostasis, including calcium metabolism (15). Around 99% of mineralized bone consists of calcium-hydroxyapatite crystals. The bone marrow also houses environments for blood cell formation, development and storage. Bone cells, blood cells and adipocytes share common stem cells that mature differently from fetal life.

Bone and remodelling

Bone undergoes dynamic changes throughout life through the process of bone remodelling. Bone cells replace damaged bone after strain and microfractures to preserve skeletal integrity (27). It is held that the entire skeleton is renewed by the remodelling process over a period of about 10 years. The osteoblasts and osteoclasts are bone cells playing a major role in this (Figure 3) (27, 28). Osteoclasts initiate the bone remodelling as they adhere to a wounded bone surface and secrete degrading acids and enzymes, which resorb bone and release calcium. Osteoblasts get recruited to synthesize and deposit new bone material (mainly type-1-collagen), and to build an osteoid framework, which is mineralized by calcium.

The bone remodelling is an integral part of the calcium-phosphate homeostasis, with blood calcium in equilibrium with calcium-hydroxyapatite (15, 18, 29). Sufficient calcium levels are necessary for its crystallization. The functions of osteoblasts and osteoclasts are tightly coupled. In a balanced bone homeostasis, which is the case in most individuals aged below 50, bone formation equals bone resorption. These processes are regulated by a complex interplay between bone, bone cells and signalling pathways (27).

Figur 3. The osteoblasts and osteoclasts are bone cells which continuously replace microdamaged bone with new bone material. This remodelling of bone renews the skeleton. An imbalance in this process degrade or build new bone. From Servier Medical Art, with permission.

The best described effects of vitamin D and PTH on bone, go through the osteoblastic cells. Insufficient vitamin D leads to decreased osteoblastic function and osteomalacia (22, 30). The feedback mechanisms increase PTH, which leads to tunneled bone resorption, woven bone and fibrosis (first phase osteomalacia). The second phase of osteomalacia involves defective mineralization. Over time, all bone surfaces can become replaced with unmineralized osteoid (third phase).

Hyperparathyroidism and an increase in $1,25(\text{OH})\text{D}_2$ recruit macrophages and osteoclasts, as a result of increased expression of the receptor activator NF- κB ligand (RANKL) at osteoblastic cells, and downregulation of the decoy receptor osteoprotegerin (27, 29). Osteoclasts exhibit specific receptors for RANKL, and can become numerous and induce bone loss. Macrophage colony-stimulating factor is another essential co-factor in this process. Calcium sensing receptors are present on bone cells, particularly on osteoblasts, and can modulate the PTH effect, and adjust the tight calcium balance and local bone remodelling (20).

PTH has additional effects on bone as it can activate the wnt-signalling pathway and inhibit the sost-gene/sclerostin pathway, which are important regulators of stem cell maturation into osteoblasts (27). The latter pathway mediates loading effects on bone, together with the osteocytes. Insulin growth factors (IGF) may have a role in these effects (31).

Bone health and assessments

Dual-energy X-ray absorptiometry (DXA) is the gold standard to evaluate bone health in clinical practice (32). It is usually assessed at the lumbar spine and hip region. Vertebral fracture assessment can help identify undiagnosed vertebral fractures (33). The outcome measure bone mineral density (BMD) is predominantly based on the amount of mineralized tissue per area. BMD is an important predictor of fracture risk (32).

For decades, alkaline phosphatase (ALP) has been used to assess the bone remodelling activity, and to evaluate metabolic bone disease and osteomalacia (34). More specific bone turnover markers have become available, such as bone specific ALP (B-ALP), procollagen type 1 propeptides (PINP) and carboxy-terminal collagen crosslinks (CTX), as were used in our studies. Recent advances can determine osteoblastic bone formation (PINP) and osteoclastic bone resorption (CTX) more specifically. The bone turnover markers are related with BMD, and they predict fracture risk independently of BMD (35).

The bones consist of an outer cortex covering the inner trabecular bone structure. The cortex is highly mineralized and provides most of the strength. Hyperparathyroidism has been linked primarily to degradation of cortical bone, while effects on trabecular bone have been less apparent (26). Whereas DXA can not separate cortical bone from trabecular bone, the recent three dimensional quantitative computer tomography (QCT) imaging has enabled more advanced bone characterization. Some QCT studies indicate that both cortical and trabecular bone are affected in primary hyperparathyroidism, which may match with an increased fracture risk observed in both compartments (36). Hyperparathyroidism seems also to have negative effects on bone structure and bone strength, according to QCT and trabecular bone score (TBS) (37, 38).

Obesity

The obesity prevalence has nearly tripled in Norway and worldwide since the 1980s (39-41). It has become a growing health issue, and is related with several co-morbidities, such as gastro-oesophageal reflux disease, dyslipidemia, type 2 diabetes, hypertension, cardiovascular diseases, osteoarthritis and certain cancers (41). WHO acknowledged obesity as a disease in 1998, and it is rated among the top preventable non-communicable health diseases (40, 42).

Obesity is characterized by an abnormal accumulation of body fat. This accumulation is considered a result of an energy imbalance. The traditional obesity definition is based on a body mass index (BMI) $\geq 30 \text{ kg/m}^2$. Later studies indicate an obesity prevalence of 18-25% in Norwegian adults depending on age and gender (3, 39). Our specialized national health service offers treatment for morbid obesity, defined as BMI $\geq 40 \text{ kg/m}^2$, or BMI $\geq 35 \text{ kg/m}^2$ combined with co-morbidities (5, 43-45). The numbers with morbid obesity have increased more rapidly than obesity over the last decades, which probably apply also to Norway (1, 46). HUNT 3 data found a prevalence around 2-3 % of the adult population in 2006-2008 (1).

The calory excess is explained by a shift towards increased consumption of energy dense food products, usually easy accessible at low cost, together with lower physical activity and energy expenditure. The energy balance are further influenced by hormones and neuronal pathways. Typical hormones are leptin and adiponectin, being secreted from adipose tissue, and gastro-intestinal hormones like ghrelin, peptide YY (PPY) and glucagon-like peptide-1 (GLP1). They may all impact appetite and sensations of hunger and satiety (47-49). The energy homeostasis is a result of physiological adaptations over years, and compensatory responses are activated to counteract changes in body weight (49). Achieving sustained weight loss is therefore a great challenge for most people.

Excessive food consumption does unfortunately not assure sufficient intake of essential nutrients, as many high-caloric food products are nutrient poor (41). Nutrient disorders are therefore common in obesity, and vitamin D and calcium intake are frequently reported lower than recommended (50).

Vitamin D and calcium

Vitamin D levels are frequently low in obesity, and the levels typically decrease with higher BMI (6, 51, 52). Reports of *vitamin D deficiency* vary, with prevalence rates up to over 90% in populations before obesity surgery, using a threshold of S-25(OH)D $< 50 \text{ nmol/l}$ (53). In addition to a lower intake of vitamin D, the skin biosynthesis is often lower in obese persons, who exhibit less outdoor activities and avoid sun exposure. One study found that S-25(OH)D levels increased in a normal population during summertime, but not in morbid obese subjects (54). Furthermore, vitamin D is fat soluble and may sequester into fat, lowering the circulating

levels. The bioavailability in blood may be further impaired due to altered activity in hydroxylase enzymes which metabolize S-25(OH)D (51, 55). Fatty liver disease is one mechanistic factor which can affect vitamin D metabolism adversely (53).

There are limited reports that calcium levels are altered in the obese compared with others. A left-shifted relationship between calcium and PTH was demonstrated in a Swedish study (56), which might suggest subnormal calcium levels. Low vitamin D levels in obesity can decrease calcium levels, but a compensatory increase in PTH and active vitamin D will minimize the changes. Although calcium intake is low in obesity, active calcium absorption can be high by other mechanisms, and calcium levels may also become higher than in normal weight (51, 57-59). Nutrients such as fat, proteins and fiber seem to modulate calcium absorption and influence calcium levels, together with hormones such as PTH/1,25(OH)₂D, estrogen, growth hormone (GH) and glucocorticoids (17, 57).

PTH

PTH increases with higher BMI, and it is often elevated in obesity (51, 60). The prevalence of SHPT seems often to be in the range 20-35% in morbid obesity, but has been reported as high as 75% (4, 61, 62). Hjeltnes and colleagues reported a prevalence of 24% in a Norwegian population of morbid obese, with 51% exhibiting vitamin D deficiency (63). Lower vitamin D levels and disturbances in calcium homeostasis may help explain SHPT. It is otherwise poorly understood how obesity impacts PTH. There may exist a hormonal cross-talk between fat cells and parathyroid cells, and a correlation between the adiposity hormone leptin and PTH has been observed (58, 64).

Bone turnover

Bone turnover is often low in obesity (65-68). Resorption markers seem relatively lower than formation markers, and the balance favours bone formation (66). The mechanisms underlying these phenomena are uncertain. Bone turnover and particularly bone resorption is suppressed postprandially, and higher food intake in obesity might be an explanation (69). A postprandial decrease in PTH could be a mediator. The elevated PTH levels in obesity,

suggest other mechanisms. Higher leptin and estrogen levels are hormones which can suppress bone resorption (65, 70, 71). Inflammation and additional hormones such as adiponectin, insulin and sclerostin may likely also impact bone turnover (65, 72, 73).

Bone health

Obesity consistently increases bone mass (61, 74). The validity of DXA determinations were for some time questioned, as increased amounts of soft adipose tissue can induce artefacts (75). It appears that QCT assessments confirm the DXA findings (66, 76).

Several mechanisms stimulate bone formation and may increase bone mass (65, 66). While immobility decreases bone mass, mobilization and increased muscular activity exert anabolic effects on bone. Higher body weight induces similar loading effects on the skeleton. Deficiencies in vitamin D and calcium may on the contrary increase PTH and bone resorption, and it is suggested that higher PTH pose a particular risk for cortical bone in the morbid obese (77). Among relevant hormones, is estrogen a major bone hormone, being produced in high amounts in adipose tissue due to the aromatase activity (78, 79). Adiposity hormones, gastrointestinal hormones, insulin and androgen hormones can also exert anabolic bone effects (78). Hormones may have multiple actions, and some effects on bone could be mediated via PTH (58, 80-82). Obesity, and particularly abdominal obesity, may also increase bone marrow fat and impact the biomechanical properties of bone (71, 83-85).

With higher bone mass, obesity was considered protective for bone health, and studies found osteoporosis and fractures frequent in the underweight (84, 86). More recent studies found less fractures at traditional osteoporotic fracture sites in the obese, including hip and wrist, but higher rates at “atypical” sites, such as ankle, leg and upper arm (84, 86). A high bone mass may therefore not fully compensate for the increased strain induced by excessive weight (66, 83). Increases in fractures may apply particularly to cases with abdominal obesity, and some studies indicate a protective effect of lean mass (71, 83, 87, 88). Although soft fat tissue can be protective (“padding effect”), fall risk and fracture rate can rise in obese with relatively less muscle strength, imbalance and greater fall impact (66, 71, 84, 89).

3. Secondary hyperparathyroidism after obesity surgery

Obesity surgery – history and procedures

Until the 1990s, surgery of obesity remained in obscurity (90). The first procedure in Scandinavia was probably performed in Göteborg in 1952, with a large intestinal resection (91). The jejunio-ileal bypass was introduced in 1954 in Iowa, US, with a bypass over large parts of the small intestine. There was an option to reverse this operation, as the bypassed intestinal segment remained in the abdomen. Jejunio-ileal bypass became the most common procedure for severe obesity worldwide in the 1970s (90). It was also performed in Norway, but its use stopped in the early 1980s due to serious complications (92). Mason developed the *gastric bypass* in the US during the years 1965-67 as an alternative procedure. It was adjusted over the years, and mimicked the Bilroth II operation, which was used in cases with gastric ulcer and cancer. Gastric bypass was introduced in Norway in the 1970s, and vertical banded gastroplasty (VBG) and gastric banding in the 1980s. These procedures were ended in Norway around 1990 due to complications, but they were continued in other countries. Over the years, the modified Roux-en-Y gastric bypass (gastric bypass) received a position as a reference procedure in obesity surgery (also called bariatric surgery) (93).

Early in the 2000s, the National health service offered obesity surgery to some few morbid obese patients abroad, and some few in Førde. In 2004, obesity surgery was reintroduced as a treatment option in all 5 Norwegian health regions as a part of the public health care (5, 45). Gastric bypass was the dominant procedure. Large studies reported at this time substantial and sustained weight loss after obesity surgery, together with benefits in obesity related co-morbidities, improved health-related quality of life and improved life-expectancy (4, 94, 95). Most procedures in Norway were now performed laparoscopically, with less surgical complications. A few selected patients with severe obesity ($\text{BMI} \geq 50 \text{ kg/m}^2$) got the *duodenal switch* operation for greater weight loss. This was a more malabsorptive procedure, performed in some centres. Obesity surgery increased in number, passing 1000 patients annually in year 2009, 2000 in year 2010, and stabilized around 3000 per year from around 2012 (96). The studies in this thesis focus on gastric bypass, which was the leading procedure these years. *Sleeve gastrectomy* was developed around year 2000 by Gagner as a part of the

duodenal switch operation, and was introduced as a “stand-alone” operation some years later (90). This predominant restrictive operation became increasingly popular and is currently the most common procedure, performed in nearly 50% of the operations in Norway in 2019 (97).

Obesity surgery – mechanisms and consequences

Obesity surgery classically targets 2 mechanisms to induce weight loss (6). The first involves a reduced stomach volume, restricting the amount of food intake and the calories that can be ingested. Gastric banding, variants of gastroplasty and sleeve gastectomy are considered to be predominantly restrictive procedures. The second mechanism, malabsorption, is induced by a bypass and shortening of the functional intestine, which reduce the area for absorption of food contents. Digestion and absorption is further impaired in malabsorptive procedures, as the food mixes with bile and pancreatic secretions lower down in the intestine.

Gastric bypass and duodenal switch are both combined restrictive and malabsorptive procedures. In gastric bypass, a very small stomach is created of around 25 ml (gastric pouch), and the food bypasses the first segment of the intestine (duodenum and proximal jejunum, about 50 cm) (Figure 4). The duodenal switch (also called biliopancreatic diversion with duodenal switch) is another combined procedure, which is created with a sleeve gastrectomy and a longer bypass of the small intestine than gastric bypass, inducing more malabsorption. In addition to the mechanisms of restriction and malabsorption, increasing attention has been directed to additional postoperative neuro-endocrine and gastro-intestinal responses (48). Alterations in GLP1 and PYY,

Figur 4. Illustration of gastric bypass. A small stomach is created (gastric pouch), and coupled directly to the small intestine. The remnant stomach is kept intact, and the outlet is coupled lower to the intestine. (From Hewitt & Høgestøl, NGF-nytt nr 4, 2015, reprint with permission)

and possibly ghrelin, can influence the weight result (47). They may alter taste, food preferences, appetite and energy homeostasis, although the relative impact on postoperative weight loss has yet to be established. The potential role of GLP1 is exemplified by the new GLP1 analogs, which have been introduced as medications for weight loss.

The changes in the gastro-intestinal anatomy and physiology disrupt the nutritional balance (21, 23). In addition to a decreased intake of calories, the smaller stomach limits the intake of important nutrients. Absorption of nutrients is similarly affected. Some data indicated that procedures with a large degree of malabsorption might involve a higher risk of nutritional deficiencies (4, 98).

Studies had reported nutritional disturbances after malabsorptive procedures such as jejuno-ileal bypass and biliopancreatic diversion, with low calcium, low vitamin D and increased PTH levels (4, 99-101). Disturbances were also recognized after gastrectomy, with findings of osteomalacia and bone loss (102). Studies after gastric bypass should show similar results, but most studies were of small size and probably attracted a small audience.

In 2005, De Prisco reasoned that little was published about the consequences of obesity surgery on calcium homeostasis, SHPT and bone health (102). The concern of postoperative nutritional deficiencies was increasing, and nutritional supplements were implemented gradually based on experience.

Calcium

Crowley et al had reported in 1984 that calcium intake was low and calcium absorption compromised in a follow-up of 41 patients 6-9 years after gastric bypass (103). Bone turnover with ALP was increased in this study, suggesting metabolic bone disease. In line with this, Ott et al found in 1992 that calcium and vitamin D levels were low, and PTH and osteocalcin were high, in a report of 29 patients 10 years after gastric bypass (104).

Prior to the studies of this thesis, the systematic longitudinal studies usually reported unchanged or normal calcium levels after gastric bypass (105-108). Coates et al observed for instance stable concentrations of calcium, vitamin D and PTH the first year after gastric bypass with daily supplementation of calcium 1200 mg and vitamin D 400-800 IE (109). This compared

with studies after gastropasty and medical weight loss (110). Riedt and co-authors found that calcium absorption was reduced 6 months after gastric bypass, but as the preoperative absorption was high, the postoperative absorption and blood levels were normal (59). After 2010, we reasoned that the calcium levels were most often normal, although the data was limited (102).

Calcium levels were frequently lower after predominant malabsorptive procedures. The Norwegian study by Våge and co-authors found low calcium levels in 8 (45%) of the 18 females 25 years after the malabsorptive jejunal-ileal bypass procedure (101). This corresponded with a serie by Marceau et al, finding decreased calcium levels 10 years after duodenal switch (111).

Vitamin D

If vitamin D levels were low in obesity, the levels might be further deranged postoperatively (108, 112, 113). Some studies concluded that the absorption of vitamin D was reduced after gastric bypass (114, 115). Given the high preoperative prevalence of vitamin D deficiency, the postoperative population was at high risk (21, 53, 105).

Until the early years of 2000, studies had found postoperative decreases in 25(OH)D levels, and some indicated a progressive decline over years (116-119). Later studies reported more variable responses. It might be ascribed to the supplementation of vitamin D that was gradually introduced (112, 116).

Most of the studies at this time were after gastric bypass or procedures with a degree of malabsorption. Few studies had compared procedures specifically, but procedures with more malabsorption seemed to have higher risk of further vitamin D derangements (4, 113, 120). Søvik and colleagues from our centre might be the first showing lower vitamin D levels after duodenal switch in a randomized controlled trial, in comparison with standard gastric bypass (120).

There are several surgery specific factors which can reduce vitamin D absorption and cause vitamin D derangements (112, 114). In addition to the reduced amount of food per meal, new food preferences limit intake of fat and vitamin D rich nutrients (103). The digestion process is further disrupted with a smaller stomach, and undigested food passes directly into the

intestine, and may pass through faster. Vitamin D is a lipophilic compound, which absorption is impaired with the delayed mixing of bile and pancreatic enzymes after malabsorptive procedures.

Some physiological mechanisms may tend to increase vitamin D levels during weight loss. Reduced amounts of adipose tissue may release some vitamin D into the blood, as well as other causes, opposite to those described for obesity (112).

Supplementation

Whereas one tablet of multivitamin a day was insufficient (115, 121-123), studies indicated that vitamin D doses of 400-800 IU might prevent further derangements of calcium and vitamin D the first years after gastric bypass, and in some cases improve the levels (45, 62, 109, 113, 116, 117, 124, 125). These doses were recognized in the first US official guidelines of supplementation in 2008, being more than the double of those recommended for the general population (150-200 IU) (4, 121).

The general focus on vitamin D status was increasing these years. With the Endocrine Society raising the threshold levels for vitamin D deficiency to $S-25(OH)D < 50 \text{ nmol/l}$, and defining vitamin D sufficiency as $S-25(OH)D \geq 75 \text{ nmol/l}$, this topic became subject for discussion (9, 10). This applied particularly for the population being operated with obesity surgery, with emerging concerns around vitamin D deficiency and bone health. It might not surprise that Heber and co-authors from the Endocrine Society were the first to address these thresholds for the bariatric population (126).

The evidence of the higher vitamin D target levels was very limited in the bariatric population (98, 127). Vitamin D sufficiency was also defined as $S-25(OH)D \geq 50 \text{ nmol/l}$ (11, 12, 128). We found that much of the focus concentrated around the high prevalence of vitamin D deficiency, and a call for higher vitamin D doses (98, 127, 129). Some characterized the vitamin D recommendations as expert opinions due to the lack of evidence (112, 130). The risk of derangements after procedures with more malabsorption, supported that patients undergoing these needed a more aggressive approach (4, 120).

PTH and SHPT

Managing SHPT was challenging, and we found our strategy to oppose SHPT insufficient. The prevalence of SHPT appeared high, even with sufficient vitamin D levels, and the supplementation regimens showed little impact (116, 117, 131, 132). PTH elevations over time were of concern, being able to increase bone turnover and induce bone loss (6, 98).

When we started our studies, the prevalence of SHPT was usually reported in the range 30-50% after gastric bypass, and higher after other more malabsorptive procedures (100, 105, 117, 133-135). SHPT appeared more frequent after gastric bypass, compared with non-operated morbid obese and after predominately restrictive procedures (118, 134, 136). According to a limited number of studies, there were indications of an increase in PTH over time (106, 116, 119, 132, 137). Some few studies reported a transient decrease in PTH the first 6 months, being followed by a subsequent increase (133).

Calcium and vitamin D levels were both thought to impact the PTH changes (98, 129, 138, 139). Deficiencies in these were early suspected to increase PTH concentrations and predispose for postoperative bone disease (103, 104). There were no obvious deficiencies in many of our cases. Supplementation was one new factor. The role of fat mass was unclear.

We found limited evidence of a relationship between calcium and PTH after gastric bypass in the beginning of this millennium (136, 140). Rather seldomly, the PTH concentrations were reported to be related with calcium (59, 109, 118, 136). Most of the PTH concern concentrated around the highly prevalent vitamin D deficiency (129, 141). De Prisco et al concluded that an inverse relationship existed between these (102). This conclusion might probably have been drawn from the general field of vitamin D and PTH regulation, as studies after obesity surgery would have statistical limitations. In the following years, some studies reported a correlation between vitamin D levels and PTH (117, 119, 142). We identified only one study that supported favourable effects of S-25(OH)D levels ≥ 75 nmol with regards to SHPT (119).

Bone health

Observational studies found small differences in bone mass between operated patients and controls; in some cases, the operated had higher BMD at the spine (93, 116). Yet, bone turnover markers were usually high.

Coates et al was probably the first in 2004 to report longitudinal bone loss the first year after gastric bypass, which corresponded with another study 1 year after vertical banded gastroplasty (109, 143). Bone loss was also observed in some later longitudinal studies, but not in all. In a review of 2012, BMD was found reduced by 10% at the hip and 5-6% at the spine 1 year postsurgery (93). These studies also confirmed an increase in bone turnover (107, 109).

A few studies at the time indicated that bone turnover remained high and that BMD decreased further to 2-3 years after gastric bypass, but data beyond was unclear (74, 106, 136, 137). The new bone turnover markers, showed a shift towards higher bone resorption, exceeding bone formation already at one month postoperatively (59, 107, 109, 144, 145).

Controversy existed whether the increase in bone turnover and decrease in BMD were pathological, or simply normal physiological adaptations to the lower body weight (45, 70, 74, 78). As markers of bone turnover peaked after 6-12 months, together with the weight loss, the weight loss could itself induce the bone loss (117, 140, 145). Parallel changes had been shown after conservative weight loss, immobilization with bed rest and after space tours (70, 74, 146). And the postoperative studies usually described normal BMD (131, 137).

The high prevalence of SHPT and high bone turnover state suggested that pathological mechanisms could be involved, and potentially affect bone health and fracture risk (4, 93). Some studies from around 1980 had indicated a relationship for vitamin D and calcium on bone (4, 99, 104, 147). In some recent studies, PTH correlated statistically with bone turnover markers and BMD (107, 119, 136, 140, 141, 148). The findings were however very inconsistent, limiting the evidence of PTH (70). Other factors such as higher age and postmenopausal status were highlighted in the studies, and an impact of additional hormonal changes was suggested.

Additional uncertainties existed around the DXA determinations. Obesity related artefacts and the weight changes might confound the results (75, 76). Older DXA techniques were less precise and inappropriate when comparing with the new DXA machines (109).

Often, the weight loss was related with degree of BMD change, but the data comparing different procedures was limited (107, 109, 131, 137, 149). Bone turnover markers seemed higher after the malabsorptive procedures and less affected after restrictive procedures (74, 117, 150-152).

The first report of fractures in 2009, found a fracture incidence of 5% in a cross-sectional sample 1-5 years after gastric bypass. A population based study from UK found that the fracture risk was comparable between surgical patients and matched controls (153). This study indicated a trend towards higher fracture risk with large weight loss and follow-up after 3-5 years. Most of these patients were operated with adjustable gastric banding (60%), and 29% with gastric bypass. In 2014, Nakamura et al reported high fracture rates in a US cohort study after a median of 7.7 years postoperatively (most gastric bypass) (154).

Part II

Review of studies

4. Aims of thesis

SHPT was a frequent observation after obesity surgery. Calcium and vitamin D levels were both assumed to influence PTH, but these levels were often normal. There were no other obvious explanations for the presence of SHPT, and the elevated PTH levels were suspected to impact bone health. We found the existing literature insufficient in the clinical management of SHPT.

Most experts agreed that S-25(OH)D levels ≥ 50 nmol/l were necessary (155). Some studies suggested that higher vitamin D levels might lower PTH further. The new guidelines from the Endocrine Society defined vitamin D sufficiency as S-25(OH)D ≥ 75 nmol/l (10, 126). We postulated that these levels would reduce SHPT more effectively after obesity surgery. Other factors might possibly also raise PTH levels, factors such as calcium, phosphate, magnesium and creatinine concentrations, and obesity. The effects of SHPT on bone health were poorly elucidated.

Hypotheses

1. The prevalence of SHPT would be lower with vitamin D sufficiency defined by S-25(OH)D levels ≥ 75 nmol compared with lower levels ≥ 50 nmol after obesity surgery.
2. SHPT would induce higher bone turnover and lower BMD after obesity surgery.

Primary objectives:

1. To investigate whether the prevalence of SHPT was lower with S-25(OH)D ≥ 75 nmol/l compared with lower vitamin D thresholds after obesity surgery.
2. To assess the prevalence of osteoporosis and low BMD 10 years after gastric bypass.

Secondary objectives:

1. To explore the prevalence of SHPT in our population after obesity surgery.
2. To investigate relationships between S-25(OH)D, ionized calcium concentrations and PTH/SHPT after obesity surgery.
3. To assess relationships between PTH/SHPT and bone turnover after gastric bypass.
4. To investigate if SHPT was associated with lower BMD 10 years after gastric bypass.
5. To explore the occurrence of low-energy fractures over 10 years after gastric bypass.
6. To explore relationships of other factors thought to influence PTH regulation, factors such as magnesium, phosphate, creatinine, BMI, age and gender.
7. To explore potential effects of our supplementation on calcium homeostasis.

5. Materials and methods

Patients and study design

The first 3 studies included patients who attended standard follow-up visits after obesity surgery at our department at 2 years and 5 years postoperatively. The fourth was an extended follow-up study 10 years postoperatively, with a predefined focus on bone health. Study 1, 2 and 4 were cross-sectional studies, while study 3 was a longitudinal cohort study up to 5 years.

Obesity surgery

Centre for Morbid Obesity and Bariatric Surgery, at the Department of Endocrinology, Morbid Obesity and Preventive Medicine at Oslo University Hospital, Aker offered an outpatient follow-up programme after obesity surgery. The centre is among the largest public bariatric centres in Norway, operating around 300 per year. A multidisciplinary team was engaged, including surgeons, physicians, specialist nurses, clinical dietists and psychologists. A research group at the centre focused on research related to the treatment.

Morbid obesity was defined as a BMI ≥ 40 kg/m², or a BMI ≥ 35 kg/m² in presence of obesity-related co-morbidities (44). The co-morbidities included type 2 diabetes mellitus, cardiovascular disease, obstructive sleep apnoea, osteoarthritis, liver steatohepatitis, and risk factors such as hypertension and dyslipidemia. Surgery for weight loss was offered after failed attempts of sustained weight loss by dietary and lifestyle measures.

Laparoscopic standard Roux-en-Y gastric bypass (gastric bypass) was the dominant surgical procedure at the department in the period being studied. A small stomach (gastric pouch) of 25-30 ml was constructed. This was coupled to the intestine about 50 cm downward from near the pylorus (ie. from ligament of Treitz; gastro-jejunal anastomosis). The bypassed segment (duodenum and proximal jejunum) was anastomosed 150 cm more distally along the intestine. Thus, the food first passes a 150 cm “alimentary limb” before it mixes with the bile and pancreatic enzymes secreted via the “biliopancreatic limb”. Selected patients with BMI ≥ 50 kg/m² were offered duodenal switch in the early period, performed with a sleeve

gastrectomy along a nasogastric tube of 30–32 F, and a subsequent construction of an alimentary limb of 200 cm and a common channel of 100 cm. No other obesity surgery procedures were included in these studies.

Examinations

All patients were evaluated preoperatively and postoperatively with weight, height and blood samples. Follow-up visits were offered after 6 weeks, 6 months, 1 year, 2 years, 3-4 years and 5 years. At 5 years, all were invited by letter or contacted by telephone at least twice. Body weight was measured with an electronic platform weight (Seca 635 0-300 kg, Class III), and height with a wall-fixed adjustable steel measure. Study 4 included a DXA assessment to determine BMD at 10 year follow-up, in addition to an extended blood sample panel.

Laboratory analyses

Blood samples were drawn after an overnight fast. Standard serum analyses during follow-up included determinations of intact PTH, S-25(OH)D and ionized calcium, which were also analysed non-systematically preoperatively. B-ALP was also analyzed. These concentrations were all determined at the Hormone Laboratory, Oslo University Hospital. PTH (1-84) was analysed by a chemiluminoimmunometric assay (Immulite 2000/2500, Siemens Health Diagnostics, Los Angeles, CA, USA) (reference range: 1.5-7.0, coefficient of variation (CV) 7%). S-25(OH)D was analysed by radioimmunoassay (Dia-Sorin, Stillwater, MN, USA) (reference range: 37-131, CV 14%) up to August 2012, and from September 2012 with liquid chromatography-mass spectrometry (LC-MS/MS) (identical reference range, CV 9%). The determinations of vitamin D levels with these methods were comparable.

The determinations of ionized calcium differed over the years: (a) first CIBA Corning, instrument 634 Ca²⁺/pH Analyzer (Bayer); from December 2005 to December 2007 with (b) Rapidlab 348 pH/Blood Gas Analyser (Instru-Med Inc., Atlanta, GA, USA) (both reference range 1.15-1.35 mmol/l, CV 1% and 2%, respectively); from January 2008 to January 2012 with (c) Rapidlab 348 pH/Blood Gas Analyser (Instru-Med Inc., Atlanta, GA, USA) (reference range 1.18-1.35 mmol/l); from February 2012 with (d) Cobas b221 (Roche Diagnostics GmbH,

Mannheim, Germany) (reference range: 1.15-1.33 mmol/l, CV 2%). Reference for iCa in the first 3 studies was based on (c) Rapidlab 348 determinations with reference range 1.18-1.35 mmol/l. Values from (a) and (b) were transformed to values of (c) Rapidlab adding 0.015 mmol/l (differences between reference ranges). Values from (d) Cobas b221 were transformed to (c) Rapidlab 348 adding 0.025 mmol/l in study 3.

The radioimmunoassay (Diasorin) used by the Hormone Laboratory at Oslo University Hospital, compared very well with the protocols of the Vitamin D Standardization Programme and the LC-MS/MS technology, being considered as a gold standard of vitamin D quantification (156, 157). The laboratory participated in the Vitamin D External Quality Assessment Scheme for the LC-MS/MS determinations to validate the determinations of total 25(OH)D concentrations (158). The Hormone Laboratory is approved by Norwegian Accreditation and complies with the requirements of the NS-EN ISO/IEC 17025 standards.

The concentrations of phosphate, magnesium, creatinine and total ALP were analysed in plasma at the Central Laboratory, Oslo University Hospital Aker, with a Modular (Roche) analyser with reagents: Phosphate and magnesium were determined photometrically, creatinine enzymatically and ALP by an enzymatic calorimetric measurement. Serum was periodically also analysed for ionized calcium at the Central Laboratory over the years with unchanged reference range (1.15-1.33 mmol/l).

Bone turnover was assessed in the last 3 studies. Total ALP was included in the second and third study. It derives from bone and liver, each contributing about 50% of the blood levels in normal physiology, and was for decades used to diagnose metabolic bone disease or osteomalacia (34, 159, 160). B-ALP was included in the study 3 and 4. Most B-ALP analyses in study 3 were performed in the years 2006-2009, with immune-extraction and enzymatically determinations (Metra Biosystems Inc, Mountain View, Ca, USA) (reference range: women 12-31 U/l, men 15-41 U/l (CV 12%).

Some extended blood test analyses were performed at the Hormone Laboratory in study 4. These included LIAISON chemiluminescence immunoassay to determine B-ALP, and Roche electrochemiluminescence immunoassay for procollagen type 1 aminoterminal propeptide (PINP) and carboxyl terminal telopeptide of type 1 collagen (CTX-1), all being modern bone

turnover markers, and in addition thyroid stimulating hormone (TSH) and follicle stimulating hormone (FSH).

Bone data

Bone status was assessed in study 4 with DXA and complementary medical history. The DXA scans assessed BMD for the total body and for the lumbar spine (L1-L4), total hip, and femoral neck. GE Lunar Prodigy technology was used until 26th August 2016, which was subsequently replaced by GE Lunar iDXA. The two scanners were cross-calibrated with 16 volunteers with very high correlation (intercorrelation coefficient 0.99).

Fracture history was recorded at the 10-year follow-up in a case record form (CRF), with information of fracture site, type of trauma and time of incidence. We reported low-energy fractures, ie. fractures after falls from standing height. Vertebral fractures were inspected visually by semiquantitative technique from DXA based lateral spine X-rays, and reported as present with at least one vertebral height reduction $> 25\%$ (ie. $VQ \geq 2$) (161).

Study inclusion

Patients were recruited during scheduled follow-up routines, and non-attending patients were called by extra invitations and eventual phone calls. All studies included gastric bypass patients, and the first study included a few patients after duodenal switch. Signed consent was a criteria for study enrollment. The study populations were defined by time of follow-up in the first 2 studies, ie. at 2 and 5 years postoperatively, and recruited consecutively at follow-up over a defined time period. These studies excluded patients with missing values of PTH, S-25(OH)D or ionized calcium, or patients having primary hyperparathyroidism, elevated creatinine or other procedures than primary gastric bypass or duodenal switch. In study 3 and 4, the study populations were defined by time of operation, and only those who attended follow-up were eligible for study inclusion. In study 3, patients with missing values of PTH or S-25(OH)D at 5 years follow-up were excluded, or elevated creatinine or suspicious primary hyperparathyroidism during the period. Study 4 included patients with a valid DXA.

Definitions and subgroups

In all 4 studies, SHPT was defined as PTH above reference range with ionized calcium up to the upper limit of the normal reference range. All studies used cut-offs of S-25(OH)D < 50, 50-74 and ≥ 75 nmol/l. We defined vitamin D deficiency as S-25(OH)D levels < 50 nmol/l according to the Endocrine Society, and vitamin D sufficiency by S-25(OH)D ≥ 75 nmol/l, thresholds which were subject to investigation in this work (10, 126). The second and third study also investigated S-25(OH)D levels < 25 and ≥ 100 nmol/l. Ionized calcium was evaluated by 3 groups in study 1-3; the first study with cut-off values defined by tertiles of the population, and study 2 and 3 applied cut-offs by tertiles of the normal reference range (1.18-1.35 mmol/l).

The WHO definitions of osteoporosis are not fully acknowledged for premenopausal women and males younger than 50 years, and our BMD analyses were separated by menopausal status and age in study 4. We defined osteoporosis using a t-score ≤ -2.5 in postmenopausal females and males ≥ 50 years according to WHO (32). For premenopausal women and men aged < 50 years, we defined a corresponding BMD using z-score ≤ 2.0 , in accordance with the criteria of The International Society of Clinical Densitometry (ISCD) (33). For simplicity reasons, we refer to these as *osteoporosis or z-score ≤ 2.0* . The same guidelines of WHO and ISCD were applied to define *low BMD* in postmenopausal females and males ≥ 50 years with a t-score < -1.0 (ie. osteopenia), and in the premenopausal females and males aged < 50 years with a z-score < -1.0.

Supplementation after gastric bypass

Standard recommended supplements during the period included one daily tablet of multivitamin containing cholecalciferol (vitamin D) 200 IU and one tablet twice a day containing calcium carbonate 500 mg and cholecalciferol 400 IU (Calcigran forte®, Takeda Nycomed, Norway). This was in addition to iron 65-100 mg daily and vitamin B12 injections 1 mg (intramuscular) every 3 months. Compliance was defined by supplement use at least 5 days a week, and non-compliance as less or no use. The vitamin D content in multivitamins was included in our dose reports of vitamin D. During the follow-up programme, supplements were adjusted to keep blood values within the normal reference range, from 2012 to maintain

S-25(OH)D $\geq$ 50 nmol/l, or S-25(OH)D $\geq$ 75 nmol/l in cases with SHPT. Ursodeoxycholic acid was prescribed the first 6 months to patients with a gall bladder intact.

Statistical analyses

All statistical analyses were performed with SPSS for Windows, versions 15-25 (IBM). Continuous data were compared with t-test and Pearson correlation test, and categorical values with Chi-square test as appropriate in all studies. The first study applied logistic regression analysis to explore differences in SHPT by stratified groups of S-25(OH)D, iCa, phosphate, magnesium and creatinine. The second study included linear regression analyses to investigate relationships between continuous data, with a stepwise exclusion of non-significant variables. The longitudinal analyses in the third study were performed with linear mixed model for individual repeated data, diagonal covariance matrices, random intercept, and stepwise backward exclusion. Generalized estimating equations (GEE) with unstructured covariance matrices were used for categorical SHPT analyses. Gender, age and BMI were included as confounders in multivariate analyses of PTH in the second and third study, and relevant interactions between S-25(OH)D, ionized calcium, PTH, B-ALP and time were tested. Continuous data were presented with means and standard deviations ($\pm$ SD), categorical variables with percentages, and odds ratios (OR) and relative ratios (RR) with 95% confidence intervals (95%-CI).

Ethics

Included patients were registered in an institutional database, licensed by The Norwegian Data Inspectorate. They had provided written informed consent, allowing their data to be registered. The studies were approved by the Regional Ethical Committee (REK) and the Data Protection Officer at Oslo University Hospital (Personvernombudet), and were performed in accordance with the Helsinki Declaration.

6. Summary of papers

Paper 1 - Secondary Hyperparathyroidism, Vitamin D Sufficiency and Serum Calcium 5 Years After Gastric Bypass and Duodenal Switch.

The first study was an exploratory cross-sectional study of patients attending a 5-year follow-up programme after gastric bypass (N=114) and duodenal switch (N=11). The procedures were performed from September 2004 to June 2006, and the patients were recruited consecutively at 5-year follow-ups during 2010 to 2012.

We tested the hypothesis that the prevalence of SHPT would be lower with S-25(OH)D $\geq$ 75 nmol, compared with lower levels. The prevalence of SHPT was analysed by 3 categories of vitamin D sufficiency, defined by S-25(OH)D < 50 nmol/l, 50-74 nmol/l and $\geq$ 75 nmol/l. We also explored strata of other factors supposed to influence PTH, such as ionized calcium, magnesium, phosphate, creatinine and supplements of calcium and vitamin D. We analysed the total population and separately for each procedure.

Strikingly, SHPT occurred in all patients after duodenal switch, and in 40% after gastric bypass. The main result was no difference in the prevalence of SHPT by the 3 vitamin D groups. The most important observation was the relationship between SHPT and ionized calcium, with a lower prevalence of SHPT in the upper two calcium tertiles compared with the lower tertile (OR 0.35, 95-CI 0.15-0.79). We demonstrated a left shifted calcium distribution in the population, with 79% exhibiting ionized calcium values below the middle of the normal reference range (Figure). The inverse relationship between ionized calcium and PTH was statistically significant for gastric bypass, but not for the 11 duodenal switch patients (P=0.083). Perhaps, the second most important observation might be the exploratory finding of a numerically but non-significantly lower prevalence of SHPT (21%) with S-25(OH)D $\geq$ 95 nmol/l after gastric bypass. Very few had levels above the more natural cut-off 100 nmol/l.

The study supported that the calcium concentration was a strong determinant of PTH levels and SHPT after obesity surgery, and not the notion of vitamin D. We reasoned that these findings needed to be retested, together with a new hypothesis that even higher S-25(OH)D levels might be needed to suppress SHPT.

Figure 5. Distribution of serum ionized calcium in 125 patients after obesity surgery with gastric bypass (N=114) or duodenal switch (N=11). The distribution is shifted towards the left, with the majority (79%) having calcium levels below the middle (dotted line) of the normal population reference range (ie. ≤ 1.26 mmol/l).

Paper 2 - Relationships of serum 25 hydroxyvitamin D, ionized calcium and parathyroid hormone after obesity surgery.

The second study was a similar cross-sectional study on a larger group of patients. We included 502 gastric bypass patients, at 2-year follow-up from June 2010 to May 2013. The study was planned during the first study.

The main objective was to test whether the prevalence of SHPT was lower with S-25(OH)D levels ≥ 75 nmol/l, compared with lower thresholds. We aimed to include a larger sample, with a higher number of patients exhibiting S-25(OH)D levels ≥ 100 nmol/l to allow further exploration of this category. We also postulated an effect of ionized calcium on PTH levels and the prevalence of SHPT, and we questioned a potential impact of PTH on bone.

Relationships of SHPT were analysed over a wider range of S-25(OH)D with the 5 subgroups: < 25 nmol/l, 25-49 nmol/l, 50-74 nmol/l, 75-99 nmol/l and ≥ 100 nmol/l, and the upper 3 groups were compared with vitamin D deficiency (S-25(OH)D < 50 nmol/l). We explored relationships for PTH with ionized calcium, and other factors such as phosphate, magnesium, creatinine, BMI and supplements. Bone turnover was assessed with ALP.

SHPT was prevalent in 34% of the population 2 years after gastric bypass. In this study, a strong inverse relationship was demonstrated between vitamin D and PTH levels across the spectrum of S-25(OH)D (Figure 6). The main finding was that the prevalence of SHPT was lower with the threshold levels S-25(OH)D ≥ 50 nmol/l (29%, RR 0.64, 95%-CI 0.50-0.81) and S-25(OH)D ≥ 75 nmol/l (28%, RR 0.61, 95%-CI 0.44-0.84), both compared with S-25(OH)D below 50 nmol/l (45%); whereas the prevalence of SHPT was lowest with high S-25(OH)D ≥ 100 nmol/l (16%, RR 0.35, 95%-CI 0.14-0.88). The study confirmed a left-shifted calcium distribution, with a large proportion (77%) having ionized calcium values below the middle of the reference range. There was a strong relationship between ionized calcium and PTH. This relationship was also found in the patients exhibiting ionized calcium within the reference range.

Vitamin D levels correlated positively with the calcium levels, as patients with the highest S-25(OH)D levels exhibited the highest and most normal ionized calcium levels. A synergistic association existed between these two on the prevalence of SHPT, and there was no SHPT in subjects with S-25(OH)D ≥ 100 nmol/l and ionized calcium ≥ 1.30 mmol/l. In subanalyses, the association of vitamin D with SHPT was only significant with calcium in the lower tertile. The study demonstrated a positive relationship between BMI and PTH. We also found higher ALP with SHPT and with S-25(OH)D < 50 nmol/l.

This study supported that vitamin D levels together with calcium levels were important determinants of PTH and SHPT after gastric bypass, and that vitamin D levels influenced the calcium levels. The threshold level of S-25(OH)D ≥ 75 nmol/l was not found superior to S-25(OH)D ≥ 50 nmol/l with regards to SHPT. The study did, however, support the findings from the first study, that S-25(OH)D levels ≥ 100 nmol/l might suppress PTH and SHPT more effectively. SHPT correlated positively with ALP, indicating that elevated PTH increased bone turnover.

Figure 6. SHPT and ionized calcium concentrations varied by vitamin D levels. The prevalence of SHPT was highest with vitamin D deficiency and lowest with high vitamin D levels. Ionized calcium was highest and most normal with high vitamin D levels.

Paper 3 - Relationships between Vitamin D Status and PTH over 5 years after Roux-en-Y Gastric Bypass: A Longitudinal Cohort Study

The third paper reported results obtained in another large longitudinal cohort study which included 554 gastric bypass patients after attending a 5-year follow-up. It was planned at the end of the first study to assess the relationship between vitamin D and PTH over time. Based on the previous studies, we aimed to include at least 500 patients, and all patients operated at our hospital from June 2004 to December 2009 (N = 823) were candidates. Inclusion was ended by August 2016.

We applied subgroups of S-25(OH)D as in study 2 when analysing the prevalence of SHPT. The prevalence was postulated to be lower with the higher S-25(OH)D levels up to 5 years postoperatively. We also explored how ionized calcium was related with PTH and SHPT, and if S-25(OH)D influenced ionized calcium postoperatively. The potential role of PTH on bone turnover was investigated.

Our emphasis was put on the period from 6 months postoperatively, as S-25(OH)D and PTH were routinely analysed. Specifically, we investigated the 4 subgroups of S-25(OH)D: < 50 nmol/l, 50-74 nmol/l, 75-99 nmol/l and ≥ 100 nmol/l, comparing the upper groups with vitamin D deficiency. Additional relationships were analysed, such as for S-25(OH)D with iCa, iCa with PTH, and PTH with B-ALP, and for supplements.

PTH increased from 6 months to 5 years, and the prevalence of SHPT increased from 18% to 38%. In parallel, S-25(OH)D and ionized calcium decreased, while BMI increased. PTH and the prevalence of SHPT were lower with S-25(OH)D levels ≥ 50 nmol/l and ≥ 75 nmol/l, and they were lowest with S-25(OH)D ≥ 100 nmol/l over the 5 years. At 5 years, SHPT occurred in 50% with S-25(OH)D < 50 nmol/l. The prevalence was lower with S-25(OH)D levels ≥ 50 nmol/l (31%, RR 0.63) and ≥ 75 nmol/l (31%, RR 0.63), and lowest with S-25(OH)D ≥ 100 nmol/l (19%, RR 0.37), all compared with vitamin D deficiency. PTH increased in all vitamin D groups, except in patients with S-25(OH)D ≥ 100 nmol/l, and the increase was larger with lower S-25(OH)D levels (Table 1). A similar relationship was documented between ionized calcium and PTH levels, and the prevalence of SHPT was most pronounced with ionized calcium in the lower third of normal range or below over the 5 years. The synergistic

relationship between S-25(OH)D and iCa was also confirmed in this study, with the highest prevalence of SHPT in patients with concomitant low S-25(OH)D and ionized calcium levels. Furthermore, no SHPT was seen with S-25(OH)D ≥ 100 nmol/l and ionized calcium in the upper 2/3 of normal range at 5 years. Ionized calcium decreased in all vitamin D groups. Supplements with calcium and vitamin D showed a small but significant relationship with higher S-25(OH)D and lower PTH over time, but not at 5 years. B-ALP was higher and remained high in the patients with SHPT over the 5 years.

The study displayed a longitudinal relationship for both S-25(OH)D and ionized calcium with PTH over time. The prevalence of SHPT was consistently lower with S-25(OH)D ≥ 50 nmol/l, and most suppressed with S-25(OH)D ≥ 100 nmol/l up to 5 years. PTH were positively related with bone turnover over the 5 years after gastric bypass.

Table 1. Relationships for S-25(OH)D on PTH between individuals and by change from 6 months to 5 years after gastric bypass (N=554).

25(OH)D (nmol/l)	Between			Change			I
	B	SE	P	B	SE	P	
D < 50	-0.078	0.02	<0.001	-0.068	0.01	<0.001	10
D 50-99	-0.024	0.01	0.002	-0.030	0.01	<0.001	7.6
D ≥ 75	-0.020	0.01	0.031	-0.022	0.01	<0.001	7.1
D ≥ 100	-0.0001	0.01	0.99	-0.003	0.01	0.74	4.8
All	-0.034	0.01	<0.001	-0.031	0.003	<0.001	8.4

Linear mixed model analysis. PTH, parathyroid hormone (pmol/l); B, regression coefficient; SE, standard error; 95%-CI, 95% confidence interval; P, significance level; I, intercept; S-25(OH)D, stratified serum 25-hydroxyvitamin D levels.

Paper 4 - Bone metabolism, bone mineral density and low-energy fractures 10 years after Roux-en-Y gastric bypass

Study 4 was a cross-sectional study of patients who attended a 10-year follow-up after gastric bypass. The study was planned after study 1 with a predefined focus on postoperative bone status determined with DXA. Included were 122 of the 203 which had been operated from June 2004 to December 2006. The same cohort had previously attended a 5-year follow-up (162).

We specifically aimed to investigate the prevalence of osteoporosis, and to explore a potential relationship between SHPT and BMD. We also tested whether other factors were related with bone health, and we recorded low-energy fractures.

Osteoporosis was identified in 16 (27%) of the 59 postmenopausal women or men ≥ 50 years, and in 4 (6%) of the 63 younger had a corresponding z-score ≤ 2.0 at 10 years. Low BMD occurred in 70 (57%) of the participants, 24 in the younger group, and 46 among the elder. SHPT was prevalent in 37 (31%). Vitamin D deficiency was found in 40 (33%), and 91 (75%) had S-25(OH)D levels < 75 nmol/l. The calcium distribution was left-shifted as in the previous studies.

SHPT was associated with low BMD after adjustments (OR 2.8). Higher PTH levels showed comparable linear relationships with lower BMD (z-score) at all skeletal sites, but were only statistically significant at the femoral neck. The bone turnover markers PINP and CTX were found higher than in preoperative cohorts of similar age, and they related with low BMD. PTH was not significantly related with bone turnover markers in this study. The other factors related with low BMD were higher age, postmenopausal status, BMI < 35 kg/m² and preoperative hypothyreosis.

There were 18 subjects (15%) who reported low-energy fractures during the 10 years, occurring at mean 8.4 years postoperatively. Most fractures were in the lower limbs or ribs. Lateral spine x-ray revealed additional 10 (8%) significant morphometric vertebral fractures. The low-energy fractures occurred most frequently among subjects with osteoporosis, compared with the non-osteoporotic (Table 2).

Consequently, a large proportion exhibited low BMD 10 years after gastric bypass, and SHPT was an apparent modifiable factor associated with low BMD. This may increase the risk for fractures after obesity surgery, particularly in those developing osteoporosis.

Table 2. Overview of low-energy fractures by BMD status 10 years after gastric bypass (N=122).							
	All	Low-energy fractures		No low-energy fractures		RR	P
	N	N	%	N	%		
<i>Premenopausal & men < 50 years</i>	63						
z-score $\leq$ -2.0	4	1/4	25%	3/4	75%	2.5	0.34
z-score $>$ -2.0	59	6/59	10%	53/59	91.5%		
<i>Postmenopausal & men $\geq$ 50 years</i>	59						
t-score $\leq$ -2.5	16	6/16	38%	10/16	62%	3.2	0.027
t-score $>$ -2.5	43	5/43	12%	38/43	88%		
<i>All (N=122)</i>							
Osteoporosis or z-score $\leq$ -2.0	20	7	35%	13	65%	3.2	0.005
No osteoporosis or z-score $>$ -2.0	102	11	11%	91	89%		
RR, relative risk for low-energy fracture (compared with higher BMD group).							

Part III

Evaluations

7. Methodological considerations

A study's ability to analyse a research question relies on multiple factors. Measurements must be reliable and valid to determine what is to be determined. Populations, study designs and statistical methods differ, and may impact interpretations of the results.

Study design

This thesis combines observational data from 3 cross-sectional studies and 1 longitudinal study. The designs have inherent flaws. Cross-sectional studies are useful to determine prevalence of a characteristic, disease or exposure at a given time, and they may suit to analyse associations between an exposure variable and an outcome variable (163). These studies can provide information of relative relationships between the variables, and they may suggest a potential effect and effect size, as fundamentals for further investigation. A cross-sectional design does however not permit causal conclusions, as the variables of interest are measured at the same time, and no temporal direction can be tested (163, 164). In contrast to cross-sectional studies, longitudinal studies can provide information of changes over time and time-dependent dynamic relationships between variables of interest, and they may improve insight into coherent changes (164). Still, disentangling causation from association is a major challenge in any observational study. A causal relationship requires that the cause precedes the outcome, that there is a relationship between the two, and that there are no other plausible explanations for the effect. Even a well-planned prospective longitudinal design does not exclude the possibility that other external factors influence inter-relationships, and that a relationship is of non-causal nature. Multivariate analyses and stratification may help take other factors into account. Randomized controlled trials remain the gold standard study design to address questions of causation (163).

Study population

Our studies included only patients with morbid obesity who were operated during 2004-2011. Oslo University Hospital, Aker is a part of the Norwegian public health service, which performed about 2/3 of all procedures of obesity surgery in Norway during these years. The population was overall Caucasians, and very few were of other ethnicities. The standard surgical procedure at the centre was gastric bypass, which is quite standardized in Norway and other Scandinavian countries. The duodenal switch procedure was offered to selected patients with BMI ≥ 50 kg/m², and was only subject for analysis in study 1. All operated patients were invited for regular follow-up visits, and all patients operated with gastric bypass in the respective periods were candidates for inclusion. Patients with a suspicious primary hyperparathyroidism or elevated creatinine were excluded.

The results presented in this thesis may apply to our centre in general, and the internal validity is probably high. The results cannot automatically be generalized to other procedures of obesity surgery or other bariatric populations outside our centre, as they might differ in demographic characteristics, surgical procedural aspects, or follow-up regimens. Differences may exist between ethnic groups, and changes over the years may induce new challenges in the interpretations.

Participation

Achieving high participation is considered important in studies to reduce selection bias and improve validity (163). The attendance rates are typically highest the first 1-2 years postoperatively, and decreases over time. Performing long-term follow-up studies has been challenging after obesity surgery, with some reporting very low follow-up rates 5-10 years postoperatively (165). There are several factors that influence participation. Satisfaction and a feeling of well-being may explain less interest or a feeling of less need of follow-up in some patients, while health problems, life dissatisfaction, poor weight loss or geographical distance may reduce attendance in others. Non-attendance is generally assumed to be associated with lower compliance and poorer weight loss, and may relate with nutritional status (166).

The attendance rate was relatively high in all 4 studies, representing 63-78% of the potential candidates. Our centre put particularly effort into achieving high attendance in the gastric

bypass patients operated 2004-2006 at both 5 year and 10 year follow-up (162, 167, 168). Study 2 recruited patients at regular 2-year follow-up visits, with no further attempts to increase participation, thus representing a more normal population attending follow-up. Effort was also made for high participation at 5 years in those operated in 2009 (169). The positive attitude is an important characteristic of the attenders, with approximately 99% providing written signed informed consent.

The hospital region had relatively low geographical mobility and limited travel distances in the period, and most patients could still be reached within our region. The follow-up programme was an integrated part of the public health care and in large part financed by the National Health Insurance, which may favour attendance. Less organized health systems, larger individual payments, longer travel distances and decentralized follow-up may all influence the follow-up rate.

Measurements and potential biases

Fasting blood specimens were collected routinely at a specialized station for blood sampling, and the determinations of S-25(OH)D, ionized calcium and PTH concentrations were performed at the Hormone Laboratory with established processing.

Vitamin D levels were determined as total S-25(OH)D, which was reported in all of our studies. From 2012, the vitamin D analyses included separate determinations vitamin D₂ and D₃ with the new LC-MS/MS technique, with vitamin D₃ making up the predominant part. Determinations of S-25(OH)D by the Hormone Laboratory were validated according to the Vitamin D Standardization Programme (156, 158, 170). Even with standardization, variability and measurement error of vitamin D may be substantial, and one single measurement may be insufficient at the individual level. There is also a great variability for PTH, which has no standardization (171).

We were not able to demonstrate significant differences between the methods of serum determinations of S-25(OH)D or PTH at the Hormone Laboratory during the period. Alterations in the calcium methodology in study 3 were more likely to induce measurement bias, but the development of calcium was comparable in some parallel ionized calcium analyses available from the Central Laboratory, with no methodologic change (172).

The studies were initiated and planned to study the postoperative relationships between S-25(OH)D and PTH/SHPT, as preoperative assessments were often missing. During the data registration, some preoperative data were observed and recorded. We were thus able to include some analyses of preoperative S-25(OH)D (N=158) and PTH (N=160) at baseline in study 3 only. They deviated from the linear assumption, and were not included in the longitudinal 5 year analyses.

The relationships between S-25(OH)D and PTH levels remained consistent in the large studies 2 and 3, also when including confounders such as BMI, age and gender in multivariate analyses. We found support for an impact of sunlight on S-25(OH)D levels in study 2, but we found no seasonal differences for PTH (173).

ALP was used as a bone turnover marker in the studies 2 and 3. With the tight coupling between osteoblasts and osteoclasts in bone remodelling, this single marker provides useful information of the bone turnover state, and possibly osteomalacia (160, 174). The B-ALP analyses in study 3 were *bone specific*, but these analyses were performed non-systematically, increasing risk of bias. Future studies may clarify whether SHPT maintains B-ALP over time.

This work was delimited with regards to phosphate. Phosphate levels were primarily analysed aiming to identify determinants of PTH concentrations and SHPT. The consistent inverse correlation, indicated that it was the opposite relationship between these that dominated, with PTH increasing the urinary phosphate excretion (reverse causation).

DXA is the gold standard to assess bone mass and determine bone health, and defining osteoporosis and fracture risk in clinical practice (75, 175). Change of DXA machines in study 4 could induce measurement bias. The inter-correlation coefficient of 0.99 was however very high between the machines (167). Preoperative DXA assessments were not available. QCT techniques are not readily available, but may reduce the bias of adipose tissue, as volumetric bone density measurements can overcome the two compartment assumption in DXA (66, 76).

Most of the results presented, relies on objective measurements and determinations of blood concentrations and BMD. Some report bias might exist regarding use of supplements and fracture history. Self-reported supplement use related with higher S-25(OH)D levels in study 2 and 3, but its impact on PTH was limited. Supplement users generally used a combination of both vitamin D and calcium, making it impossible to differentiate the effects of each. Intake

of energy and macro- and micronutrients may be under-reported, and subjects may have presented themselves more desirable with regards to adherence of the supplementation (176, 177). Clinical fractures were questioned specifically at the 10-year follow-up, and noted in the CRF with further specifications.

Categorization issues

Results from blood tests and data of BMD are continuous variables. Defining what is normal and optimal values can be subject of debate. Commonly, the reference range has served to define what is normal, but clinical outcomes may not depend of these cut-offs.

An important example for our purpose, was the different vitamin D definitions of vitamin D sufficiency, $S-25(OH)D \geq 50$ nmol/l and ≥ 75 nmol/l, which were both within the reference range 35 - 130 nmol/l (9-11). With the limited data available, the common thread in our work was to investigate if the prevalence of SHPT was lower with the higher S-25(OH)D levels.

The optimal PTH levels are to be defined (178, 179). The high variability of PTH and differences between assays can help explain conflicting results between studies. It has also been suggested that slightly elevated PTH levels may be an appropriate adaptive response to underlying imbalances (178). In line with some others, our study 4 supported a continuous relationship between PTH and BMD (z-score) 10 years after gastric bypass. We also found support for using the upper reference range of PTH as a cut-off, as the slightly elevated PTH concentrations were related with low BMD. Most of the PTH elevations were mild and just above reference range (ie 7.1-10 pmol/l).

Calcium levels within the normal reference range have most often been referred to as normal. Striking in our studies 1-3, was the clear positive relationship between SHPT and serum calcium concentrations, even across the normal reference range. Our exploratory calcium tertiles applied in study 1, proved well in the consecutive studies 2 and 3. Hence, the findings for the lower calcium tertile, may have revealed the most pronounced calcium disturbances. Future studies may hopefully help to differentiate adverse calcium levels better. Cases with suspected primary hyperparathyroidism were excluded, due to the causative role of PTH on calcium.

The WHO categorization of BMD is established to predict fractures in postmenopausal women and men aged ≥ 50 years. Comparing our younger subjects, we found very small differences categorizing BMD by t-scores or z-scores, defined by WHO and ISCD. The DXA also helped identify vertebral fractures.

Statistical considerations

Sample size is a critical factor for statistical analyses, as statistical power increases with increasing number of observations. Small samples are often insufficient to reject a null hypothesis (type 2 error) (180). In contrast, extensive exploratory studies have a risk of finding differences just by random (type 1 error). Study 1 and 4 were primarily exploratory studies with a limited number of participants, and a risk of type 1 and type 2 errors. The non-significant relationship between ionized calcium and PTH levels after duodenal switch in study 1 was likely a type 2 error. We also questioned whether the non-significant SHPT finding with $S-25(OH)D \geq 95$ nmol/l in study 1 could be a type 2 error, and the relationship of high $S-25(OH)D$ levels turned significant in the larger studies 2 and 3. We were surprised to find lower BMD in subjects with preoperative hypothyreosis in study 4, and a confounder or a type 1 error should not be ruled out.

The distribution of PTH was not fully parametric. PTH was therefore logarithmically transformed when analysing this continuous variable in study 1. Deviation from normality was of less concern when analyzing the dichotomized SHPT variable by categories with chi-square test and logistic regression analyses.

Initially, PTH was transformed logarithmically in the T-tests and regression analyses in study 2, to account for a non-parametric distribution. The central limit theorem states that the mean distribution approximates the normal distribution with increasing sample size, unless the distribution is too skewed (181). Thus, we used PTH itself as an easier variable, which compared fairly equal. The same reasoning was utilized in the consecutive study 3.

In study 2, we reported RR as the effect estimate, in contrast with OR in study 1. The two ratios have several similarities. Both are typically used to compare prevalences of an outcome by an exposure, and they perform equally in determinations of statistical differences. When the outcome prevalence is low, they are close to each other. However, the ratios diverge as the prevalence increases, and typically when it exceeds 10-20%. Figure 7 illustrates how they perform at different prevalences, given a fixed ratio between the prevalences of the two groups studied. RR remains constant irrespective of the prevalence, while OR increases excessively with higher prevalence, and with a larger ratio. In such analyses, RR can be considered more robust irrespective of prevalence, and it provides an easy and direct understanding of the relationship being studied. OR may on the other hand be a more flexible statistical tool (182).

Figure 7. OR and RR compare the prevalence of a categorized outcome variable by two categories of an exposure variable. They perform equally in significance testing. Given a constant ratio between the prevalences, RR remains stable, and OR will divert with increasing prevalence of the outcome, and with larger differences between the populations.

Linear mixed model was applied in the longitudinal analyses of PTH and B-ALP in study 3. A major advantage with this model is the ability to account for individual covariances and thereby increase statistical power to reveal true differences (183). Introducing a random individual intercept, showed that individual characteristics made up around half of the observed variation of PTH. Linear mixed model also enables analyses with missing values in the data set, which would be excluded with other methods (183). In study 3, we found that the continuous S-25(OH)D levels, but only a cut-off defined by S-25(OH)D $\geq$ 50 nmol/l, preceded and predicted the subsequent PTH levels independently, with a declining

correlation over the following 1-2 years. The maximal response of vitamin D was expected to be quicker than the time interval between our follow-ups (with blood tests) (171). We therefore defined a data matrix with S-25(OH)D and PTH values at each time point to describe the development over time. Table 1 illustrates the findings of PTH differences between individuals and the individual changes by this model. The analyses of mean changes performed equally compared with paired sample T-tests, linear regression and interaction analyses. Corresponding analyses of SHPT were made with GEE analyses. The GEE analyses also enable multivariate analyses, take into account individual covariance and include subjects with missing data.

Generalizability

Study 2 and 3 were reported according to the Strobe guidelines of standardized reporting, making comparisons more easy across populations (184). With the standardization of S-25(OH)D determinations, these may be compared worldwide. The Hormone Laboratory is further approved by the Norwegian Accreditation and complies with requirements of the NS-EN ISO/IEC 17025 standards (158). There is no worldwide standardization for the PTH determinations, as for vitamin D (171). Generalizations for PTH and calcium are therefore somewhat more difficult. No clear definitions exist of the threshold levels to be used, and the methodological determinations and population reference ranges show variations. Relating ionized calcium and PTH with the population reference range was considered to be most robust and transferable. The DXA determinations were calibrated regularly with a phantom, being acknowledged and recognized across institutions and countries.

Surgical procedures seem to have their distinct effects on absorption and nutritional status. Although surgical technique and patient follow-up were standardized at our institution, the external validity may be impacted by several issues. The procedure is quite standardized in Norway and Scandinavia, but the bypass of the proximal 50 cm intestine at our department is shorter than the bypass in some of the other reports. A longer bypass can induce more malabsorption of calcium and vitamin D. Preoperative and postoperative routines may vary between centres, not at least the supplementation regimens, and may influence the relationships studied. Routines and characteristics also change over time. Cautiousness is

therefore necessary when evaluating results across bariatric populations. The guidelines which have been developed over the years, may help improve inequalities between centres and countries.

The results may help improve our insights into some aspects behind SHPT across populations. However, they should not be generalized to obese subjects or others, with normal intestinal anatomy and function. Subjects who seek and undergo obesity surgery may not be similar to obese in general. Such aspects may also vary between centres.

8. Discussion

The prevalence of SHPT was high after gastric bypass and duodenal switch. SHPT was most consistently related with calcium levels in the lowest range. There was also a clear inverse relationship for S-25(OH)D with PTH and SHPT in the larger studies 2 and 3 after gastric bypass, with the highest prevalence of SHPT in vitamin D deficient patients. Applying the thresholds S-25(OH)D ≥ 50 nmol/l and ≥ 75 nmol/l showed comparable PTH levels and prevalence of SHPT, whereas both these endpoints were lowest with S-25(OH)D ≥ 100 nmol/l. The longitudinal study 3 found an initial decline in PTH and the prevalence of SHPT the first 6 months, which was followed by an increase up to 5 years. SHPT related with higher bone turnover markers in study 2 and 3. In study 4, bone mass was lower with SHPT at 10 years, and low-energy fractures were most frequent in the osteoporotic patients.

Table 3. Overview of the 4 studies of secondary hyperparathyroidism and factors related after gastric bypass, and patient characteristics.

	Study 1	Study 2	Study 3	Study 4
SHPT, %	40	34	38	30
Patients included (N)	114	502	554	122
Time postop (years)	5y	2y	5y	10y
Participation rate, %	78	64	70	63
Age (years)	45	46	47	50
BMI (kg/m ²)	34	32	35	36
Ionized calcium (mmol/l)	*	*	*	
25(OH)D (nmol/l)		*	*	*
Phosphate (mmol/l)		*	*	
BMI (kg/m ²)		*	*	
ALP (U/L)		*	*	
B-ALP (U/L)			*	
Low BMD				*
BMD femoral neck, z-score				^a
Supplementation (Ca + vit D)			^a	
SHPT, secondary hyperparathyroidism; 25(OH)D, 25-hydroxyvitamin D; BMI, body mass index; ALP, alkaline phosphatase; B-ALP, bone specific alkaline phosphatase; Low BMD, defined by z- or t-score < 0.1 according to age and menopausal status; * Relationship with secondary hyperparathyroidism; ^a Relationship with PTH.				

SHPT after obesity surgery

The prevalence of SHPT in the studies included, compares well with other reports after obesity surgery, but the short-term prevalence of 18 % at 6 months in study 3 is of the lowest reported. Study 2 and 3 should supply relevant data on SHPT, being large studies in the field, and the inverse relationships for S-25(OH)D and calcium levels with SHPT are in line with the concept of PTH regulation (7, 14).

The increases in PTH and SHPT up to 5 years correspond with other long-term reports after gastric bypass (185-190). The increases were most pronounced with vitamin D deficiency, as has been found in some few studies (185, 188). Studies indicate that SHPT is less frequent by elevating S-25(OH)D levels ≥ 50 nmol/l, while the data is not conclusive regarding levels ≥ 75 nmol/l (191-195). Our cross-sectional findings may suggest that a plateau effect exists for PTH within the S-25(OH)D range 50-99 nmol/l, whereas the longitudinal effects in study 3, lend support to PTH variation also within this range of S-25(OH)D (Table 1). Study 2 and 3 both suggested that S-25(OH)D levels ≥ 100 nmol/l are most effective in maintaining calcium balance. These levels revealed the greatest PTH suppression over time in study 3. PTH did not increase with S-25(OH)D ≥ 100 nmol/l, but increased in groups with lower S-25(OH)D.

Few studies have published on SHPT by S-25(OH)D of 100 nmol/l. Two studies from Norway and Sweden reported recently that PTH and the prevalence of SHPT were lowest applying a threshold of S-25(OH)D ≥ 100 nmol/l in the general population, and neither the merged randomized controlled trials by Jorde and Grimnes disclosed an upper vitamin D limit (171, 196).

The relationship between PTH and calcium was present within the normal range of calcium, as illustrated in study 2 (Figure 8). This was also shown in a Swedish population study some few years ago (197). Consequently, we may suppose a continuous relationship between these within the reference range, and the lower levels may exhibit the greatest risk of SHPT.

The left-shifted calcium distribution seems relevant. Small calcium changes of 0.02-0.03 mmol/l can explain this shift and a PTH increase. Few studies have reported on the prevalence of hypocalcemia after gastric bypass the last 10-20 years. A Brazilian study reported numbers

in the range 5-10% at several time-points up to 10 years after gastric bypass (185). Ionized calcium was below reference range in 6% of our patients at 2 years postoperatively in study 2. Another report from US found low calcium in 1.9% up to 1 year after gastric bypass, and 9-10% after sleeve gastrectomy and duodenal switch (198). One may annotate that 2.5% is the expected number at normal.

Figure 8. In study 2 there was a relationship for PTH with ionized calcium levels. As illustrated, this was also seen in the individuals with ionized calcium within the normal reference range.

It is held that the major vitamin D dependent active absorption of calcium takes place in the proximal intestine, of which most is functionally removed in gastric bypass. A recent study indicated that vitamin D and calcium, together with PTH, explained 57% of the variation in calcium absorption 1 year after gastric bypass and sleeve gastrectomy, the latter with the full intestine being intact (199). These findings suggest that vitamin D can increase active calcium uptake after both procedures in the functional intestine. Some of the vitamin D effect on active absorption may thus persist postoperatively, in addition to the role in facilitating the passive calcium transport across the intestinal border, as previously described. The active absorption is particularly important when calcium intake is low, which is common after obesity surgery (17). The passive absorption also depends on the amount of calcium within the gastrointestinal lumen.

First in 2016, a meta-analysis concluded that the calcium concentrations were decreased and PTH concentrations were increased after gastric bypass (139). The study mentioned above, suggests that calcium absorption can be compromised after sleeve gastrectomy as well, which has no intestinal bypass or malabsorptive component (199). There are several mechanisms that influence calcium absorption and blood levels. In addition to lower calcium intake, the postoperative calcium absorption is affected by factors such as the digestion process, the intestinal transit time, hormonal changes including estrogen loss, and the dietary composition.

It is plausible that insufficient nutritional supplementation can lead to a PTH increase over time (200). Reduced functional length of the small intestine in more malabsorptive procedures, can increase the disturbances. Two studies from our centre by Risstad and colleagues, found a larger increase in PTH after both duodenal switch and gastric bypass with an enlarged intestinal bypass, both being compared with standard gastric bypass (201, 202). The S-25(OH)D and ionized calcium levels were also found lower after duodenal switch. Such disturbances seem less pronounced after the predominantly restrictive procedures, but the prevalence of SHPT may increase after these procedures as well (200, 203).

PTH and further considerations

Supplements of calcium and vitamin D are advised to maintain calcium balance after obesity surgery. In addition, vitamin D can be liberated from adipose tissue into the blood during fat loss, and calcium is released from bone during bone loss. In study 3, we studied a subpopulation with valid values of S-25(OH)D and PTH from baseline to 6 months. PTH decreased during this period in all vitamin D subgroups, except in the non-compliant subjects with 25(OH)D < 50 nmol/l (figure 9, not published). Some minimum of vitamin D in blood may therefore maintain calcium homeostasis and suppress PTH in this early postoperative period. A recent study by Schafer et al found that calcium absorption was still reduced and PTH increased with keeping S-25(OH)D levels ≥ 75 nmol/l over 6 months after gastric bypass (204). We found a small effect of

gastric bypass. PTH decreased in all groups except in the patients with S-25(OH)D < 50 nmol/l who not used supplements.

supplementation on S-25(OH)D levels at 2 years in study 2, and on S-25(OH)D and PTH levels over the 5 years in study 3.

The higher prevalence of SHPT in patients with calcium levels in the lower range, may apply particularly to populations with a left-shifted calcium distribution. Ionized calcium is a more sensitive measure of calcium status than total calcium and albumin-adjusted calcium (205). Use of other calcium analyses, may explain that some found no calcium-PTH relationship. The small calcium changes within the normal range that affect PTH, might also help explain the clinical challenge in interpreting SHPT. We noted that SHPT varied most by S-25(OH)D levels in the lowest calcium tertile in study 2 and 3. It may suggest that PTH secretion is more sensitive to the vitamin D effect in the lower calcium range.

The regulation of PTH is insufficiently understood, and the great variability in S-25(OH)D and PTH makes such determinations challenging. With all the factors in study 2, the model explained 16% of the variation. Study 3 revealed that around 50% of the variability in PTH was due to individual characteristics, such as genetics and environmental factors. The impact of other factors is to be explored. Fat mass may influence PTH by leptin or by other unknown adiposity related mechanisms. The inverse relationship we found between phosphate and PTH, indicates that the dominant pathway between these, is that of PTH increasing phosphodiuresis, partly via its effect increasing fibroblast growth factor (FGF-23). Although phosphate exerts a small effect on PTH, its potential impact on bone must not be neglected. We were not able to demonstrate any relationship between magnesium in blood and PTH. Changes in magnesium levels can blunt the PTH response, and we should not ignore its potential impact (206). Additional factors such as age, gender and renal function seems to have an effect, while other factors are less established.

Bone health

In study 3 and study 4, bone turnover was increased up to 5 years and at 10 years after gastric bypass. This compares with a few other recent longitudinal studies up to 5-7 years, which also indicate a balance towards higher bone resorption (190, 207-209). A publication by Yu et al from 2016, support that bone resorption may precede bone formation already at 10 days

postoperatively (144). A recent case-control study concluded that bone turnover was increased 10 years after gastric bypass, but not after gastric banding (210).

Longitudinal studies the last years have observed that the bone loss continues after weight stabilization. A Swedish study by Raoof and colleagues reported in 2016 a persistent decline in BMD from 2 to 5 years after gastric bypass at both hip and spine (0-5y: 25% and 19%, respectively) (186). These patients received no standard vitamin D and calcium supplementation. Lindeman et al reported 2 years later more moderate declines with both DXA (15% at hip and 8% at spine) and QCT (12% at spine) after 5 years (208). Volumetric QCT determinations seem less influenced by obesity related artefacts; the real spine changes may therefore be closer to the hip changes, than have been observed with DXA (211, 212). Corresponding reductions have been observed at the tibia and the radius with HR-qCT, at both cortical and trabecular compartments up to 7 years after gastric bypass (208, 209, 211-214). The 10-year follow-up by Raoof et al with DXA reported a slower decline beyond 5 years (215). The QCT studies have generally reported adverse effects on additional bone characteristics, such as increased cortical porosity, trabecular deterioration, and reduced estimated bone strength (208, 209, 214). Two studies assessed trabecular bone score over 1-3 years after gastric bypass and sleeve gastrectomy, and found contradictory results on trabecular spine properties (192, 216).

The bone loss has been reported in the range 1-3 % per year beyond weight stabilization, which is more than normal (186, 208, 209, 214, 215, 217). While studies up to 2-3 years postoperatively usually observed normal bone mass, the longer term studies have reported mean or median t-scores or z-scores in the negative range (0 to -1.0 SD) 5-10 years after gastric bypass (167, 186, 209, 210, 215). Osteoporosis results consist primarily of sporadic cases. In study 4, we found an osteoporosis prevalence of 27% in postmenopausal women and men aged ≥ 50 years, and a corresponding z-score ≤ -2.0 in 6% of the younger group 10 years after gastric bypass. Most bone studies have been performed after gastric bypass procedures. Results so far indicate that bone loss is less pronounced at the hip after gastric banding and sleeve gastrectomy, whereas the spine data is inconsistent (61, 218-220).

Factors related with postoperative bone health

Our understanding of bone loss after obesity surgery is incomplete. The reviews are mostly based on studies up to 2 years postoperatively, a period skeletal unloading contributes significantly. Several studies have reported correlations between postoperative weight loss and lower BMD (93, 214). Correspondingly, 2 of the 3 studies evaluating bone mass with peripheral HR-qCT, found that the weight bearing tibia was more affected than the radius 1 year after gastric bypass (151, 211).

The BMD reductions are generally greater after obesity surgery than after non-surgical weight loss (139). The continued bone loss after 2 years, also suggests that surgery specific mechanisms are involved. Two studies by Hofsø and colleagues indicate a specific effect of gastric bypass on bone turnover and BMD (220, 221). As the weight bearing and the non-weight bearing skeleton both appear to be affected, there are likely systemic factors involved (218). One likely such factor is SHPT, which was early linked with osteomalacia and osteoporosis. But its impact on bone health needs to be better clarified.

Whereas some previous studies found a relationship between PTH and BMD, the studies 2-4 are among the few showing significant relationships of the categorical endpoint SHPT with higher bone turnover and lower BMD (202). Two additional longitudinal studies have reported that bone turnover was increased up to 5 years after gastric bypass (190, 208). In study 3, the B-ALP elevations persisted up to 5 years in the subjects with SHPT, whereas the trend was declining with no SHPT (Figure 10). Elevations in PTH and ALP may both reflect osteomalacia. Persistent SHPT can

explain the findings by others that bone resorption dominates over formation, and bone degradation. If calcium levels decrease, it may impair the crystallization of calcium and phosphate. In study 4, both SHPT and bone turnover were associated with low BMD. In the

Figur 10. B-ALP was increased at 6 months after gastric bypass and remained high over time with secondary hyperparathyroidism, while there was a decline in patients without

few other long-term bone studies, PTH tends to correlate more frequently with BMD 4-10 years after gastric bypass (186, 208, 210, 214).

Some studies found correlations between vitamin D levels and bone mass after obesity surgery, but reports of a potential threshold effect of S-25(OH)D are very limited. The vitamin D threshold for bone health parameters is generally most robust for 25(OH)D $\geq$ 50 nmol/l (155). We found support for this threshold with regard to bone turnover in study 2 and 3. A recent study by Luger et al also found higher BMD and lower bone loss in the patients with S-25(OH)D $\geq$ 50 nmol/l 1 year after one-anastomosis gastric bypass (222).

Accumulating observational evidence support that fracture risk increases after obesity surgery, with most data available after gastric bypass (223) (Supplementary Table 1). The fracture risk appears to increase by time from 2-4 years postoperatively, and is highest after predominantly malabsorptive procedures (224). At present, it is uncertain whether the procedures gastric banding and gastric sleeve have an increased risk for fractures, both being predominantly restrictive procedures (223, 225). In study 4, low-energy fractures were more frequent in the subjects presenting with osteoporosis at 10 years. Fractures have so far not been significantly related with SHPT after obesity surgery. Only the Olmsted study provided information that hypocalcemia and vitamin D deficiency were both prevalent in the population, suggesting a high prevalence of SHPT (154). The clinical relevance of PTH was underlined in two meta-analyses this year, which concluded that mild primary hyperparathyroidism were related with an increased fracture risk (226, 227).

In study 4, postmenopausal status and preoperative hypothyreosis were related with low BMD. Estrogen is important for skeletal integrity, and the estrogen loss in women transitioning into menopause reduce bone mass and calcium levels (137). Estrogen levels are also reduced in obese undergoing weight loss, due to lower aromatase activity, and may affect bone similarly. Thyroid hormones maintain bone mass, but an association with low BMD need further exploration.

Several adiposity related and gastro-intestinal hormones are postulated to have effects on the postoperative bone loss (61, 74, 218). Reduced leptin levels were related with bone loss in a few studies, and changes in hormones such as adiponectin, ghrelin and PYY may impact bone. Lower insulin may decrease BMD, while increases in GLP1 and glucose-dependent

insulinotropic polypeptide (GIP) can act positively. Growth hormone affects calcium and bone, and several hormones interact with the GH-IGF pathway. Little is known about androgen effects on bone in men and in females with polycystic ovary syndrome, which is prevalent in morbid obesity.

Some previous studies indicated a protective effect of lean body mass on BMD and fractures. The stimulatory effect of lean mass on bone was also supported in 2 recent studies, which showed that physical activity mitigated postoperative bone loss (228, 229). Muscular strain likely mediates a “loading effect” on the skeleton, whereas muscle mass and muscular activity can stimulate bone formation by mutual interactions and myokines like irisin (230). Recent meta-analyses indicate that physical training can preserve and even increase bone mass during conservative weight reduction (231, 232). Some intervention studies do also support that protein substitution can have a positive impact on lean mass and BMD during weight loss (192, 233-235).

Vitamin D supplementation

Most data from the general population support a benefit of S-25(OH)D levels ≥ 50 nmol/l with regard to bone health, and some studies on PTH may suggest an additional effect with S-25(OH)D levels ≥ 75 -100 nmol/l. The doses will necessarily need to be increased in many to reach the higher vitamin D levels. As the American ASMBS adopted the target of S-25(OH)D ≥ 75 nmol/l, they proposed start doses of 3000 IU per day after gastric bypass and sleeve gastrectomy (6). This target threshold was referred to as “optimal” in the Scandinavian guidelines of 2018, while values below were considered more difficult to interpret (236). The very recent British BOMSS guidelines state both S-25(OH)D ≥ 50 nmol/l and S-25(OH)D ≥ 75 nmol/l in their recent recommendations, and point out the lack of evidence (130).

Standard vitamin D supplementation doses of 400-800 IU have generally showed small effects on endpoints. Vitamin D has been used in higher peroral doses in four randomized intervention studies after gastric bypass (Supplementary Table 2). Goldner compared daily doses of 2000 IU and 5000 IU with 800 IU over 1 year (193). Carlin added a weekly dose of 50,000 IU to standard care (191). Two other studies included patients after both gastric bypass and sleeve gastrectomy; Muschitz et al supplied preoperative weekly loading doses of 28,000

IU over 8 weeks followed by a weekly maintenance dose of 16,000 IU vitamin D up to 2 years postoperatively, and Smelt et al provided peroral liquified cholecalciferol 50,000 IU every month over 6 months (192, 194). The intervention of Muschitz added calcium 1000 mg per day, together with an individualized protein substitution and exercise programme. All these studies had a high proportion of vitamin D deficiency preoperatively, and a mean/median S-25(OH)D < 50 nmol/l. The S-25(OH)D levels increased to 60-95 nmol/l. Goldner found that more than one half reached S-25(OH)D $\geq$ 75 nmol/l with the intervention, and observed a transient correlation between vitamin D and PTH at 6 months, but not later. PTH was reduced in 2 of the other studies, by 17% and 24% (192, 194). In one study, bone turnover was reduced, and two of the studies reported positive effects on BMD (191, 192). Higher vitamin D doses are recommended after duodenal switch, but there are few studies available.

The benefits of elevating the S-25(OH)D threshold higher than $\geq$ 50 nmol/l are debated (155). Traditionally, S-25(OH)D levels up to 220-250 nmol/l have been considered safe with regards to calcium toxicity. While some studies suggest improved health outcomes with S-25(OH)D $\geq$ 100 nmol/l, other studies have raised safety concerns finding increased cardiovascular diseases, cancers and mortality with such high levels and the high vitamin D doses (9, 11, 237-240). The concerns may at least partly be confounded by the cross-sectional study designs. Prospective observational studies and supplementation trials may possibly not support excess mortality with S-25(OH)D $\geq$ 100 nmol/l, and the mortality increase was reversed after standardization of S-25(OH)D to the gold standard assay in NHANES III (241-243).

Randomized controlled vitamin D trials have neither found advantages of vitamin D alone on BMD and fractures in the general population, but vitamin D doses of 400-800 IU in combination with calcium 500-1200 per day seem protective in high risk individuals (244-246). Higher doses have not been superior to 800 IU in meta-analyses of fractures. While doses up to 2000 IU per day are generally considered safe, higher doses such as 4000-10,000 IU were related with lower BMD in one recent randomized trial (247). As with low vitamin D levels, some intervention studies reported increased falls and fractures with vitamin D doses of 4000 IU per day, 48,000-60,000 IU per month and annual doses of 300,000-500,000 IU (248-251). These trials were primarily performed in the elderly, but call for cautiousness regarding intermittent boluses and high-dose supplements.

Vitamin D deficient patients appear to be more responsive to vitamin D substitution, compared to patients with S-25(OH)D levels ≥ 50 nmol/l. Calculations from study 3 indicate that S-25(OH)D levels need to be increased by 35-50 nmol/l in the non-deficient, to reduce PTH by 1 pmol/l. Accordingly, S-25(OH)D levels ≥ 100 nmol/l may often be needed to counteract PTH concentrations of around 8 pmol/l or higher. Estimations indicate that *additional* doses of around 2000 IU per day are needed to raise the S-25(OH)D levels by 40-50 nmol/l in the normal population (244). It may involve even larger doses after gastric bypass.

Calcium supplementation

To assure adequate amounts of calcium for gastrointestinal absorption, guidelines recommend 1000-1500 mg calcium per day after gastric bypass and sleeve gastrectomy (6, 236). We recommend a standard calcium dose of 1000 mg per day, and an increase to 1500 mg in cases with SHPT, and all duodenal switch patients are prescribed 2000 mg. Single doses up to 600 mg are advised for absorptional reasons (6). Higher calcium doses could potentially increase the diffusion gradient and passive absorption, but are often associated with discomfort.

There have been variable results in studies of calcium supplementation after gastric bypass. In two randomized intervention studies, PTH levels were reduced (252, 253). The citrate formulations may be more effective than carbonate, especially in less acidic environments (252, 254). One study found higher bioavailability of calcium with citrate (252). A Danish study found no differences on calcium and PTH levels, but the bone turnover markers were lower with citrate (255). Another study observed equal reductions in PTH with both calcium citrate and calcium carbonate in doses of 600 mg twice a day (253). Better studies can hopefully clarify potential differences between these formulations (256). In Norway, calcium carbonate is most available, and calcium citrate may be less tolerable (255, 256). Peroral liquified solutions are possibly more effective than tablets, but are less in use (257).

Clinical implications

All 4 studies found a high prevalence of SHPT, which may have adverse bone effects. The studies together support that calcium and vitamin D levels are important determinants for SHPT after gastric bypass, and they may impact bone health. Assuring sufficient vitamin D levels, can support calcium homeostasis, reduce SHPT and improve bone health after obesity surgery. It follows that lifelong supplementation, monitoring and dose adjustments are essential. Our findings indicate that S-25(OH)D levels < 100 nmol/l are insufficient to optimize calcium homeostasis in a substantial portion of the patients, and many of these may need S-25(OH)D levels of at least 95-100 nmol/l to have an additional effect on SHPT.

Two of the high-dose vitamin D intervention studies found that PTH was reduced in the vitamin D deficient, and that it impacted bone health parameters positively. In both studies, 25(OH)D levels were below 50 nmol/l and reached levels ≥ 75 nmol/l, which might advocate both these target levels. None of these studies did compare the groups reaching 25(OH)D levels under or above 75 nmol/l, and no intervention study has provided data of vitamin D levels ≥ 100 nmol/l.

Our studies support that recommending vitamin D doses of 600-1000 IU per day can prevent vitamin D deficiency in the majority of the patients after gastric bypass, but some need higher doses. Suppressing SHPT with 25(OH)D levels ≥ 75 -100 nmol/l will demand higher doses for most subjects. It needs to be discussed to whom higher doses should be recommended, and which doses to be used. Obvious candidates would be those on supplementation with persistent SHPT and 25(OH)D levels in the range 50-99 nmol/l. Advicing S-25(OH)D levels ≥ 75 nmol/l for all, can be a stepwise approach to reduce the overall prevalence of SHPT, but such an intensified approach might benefit a small part of the population. Our data indicate that S-25(OH)D levels ≥ 100 nmol/l would be necessary to achieve a substantial further reduction in SHPT up to 5 years, but this goal must be balanced with the benefits on clinical outcomes and safety risk.

Individual risk estimations may be evaluated. Monitoring PTH, calcium, vitamin D and bone turnover is relevant for evaluation of SHPT, and may help predict bone risk. SHPT, low range calcium levels and/or high bone turnover might all indicate a need of higher 25(OH)D levels.

Factors such as age, gender, menopausal status, family history and previous fractures is also relevant to account for the bone risk.

A bone assessment after weight stabilization 1-2 years postoperatively is preferable to identify the most affected, and to predict future bone risk. The observed bone loss over time may indicate another DXA at 5-10 years, and with some intervals later. Some high-risk patients will probably benefit from an upfront assessment. Algorithms of bone treatment need also to be reviewed for the population having undergone obesity surgery.

The bone health data imply an evaluation of the postoperative follow-up program. By now, specialized centres have provided standard follow-up for up to 5 years after gastric bypass. We recommend blood tests every half year up to 5 years, and thereafter annually if stable. A DXA is recommended after the 5-year follow-up. When general practitioners take over the long-term management, co-operation with the specialists seems necessary.

Future research

This work has identified multiple research questions around calcium homeostasis, vitamin D and bone health, and the questions involving obesity surgery likely exceed the answers. Despite the mechanistic understanding of PTH regulation and its effects, the knowledge may be insufficiently implemented in clinical practice. PTH may further be explained by additional factors to be explored. Future measures and research aiming to counteract SHPT, may help improve bone health beyond the population undergoing obesity surgery.

Large studies would be needed to better quantify the relationships between the vitamin D cut-offs and SHPT. Is there a plateau effect for PTH, or is there a continuous PTH effect in the S-25(OH)D range 50-100 nmol/l? Future studies need to retest whether S-25(OH)D $\geq$ 100 nmol/l is most optimal for calcium homeostasis. The risk of harm needs to be established. And what is the role of adipose tissue on SHPT?

There appear to be insufficient knowledge of calcium homeostasis after obesity surgery. PTH differences within the calcium reference range need more investigation. Are there differences in changes of vitamin D and calcium in the lower range with regard to PTH, compared with the upper range? Or is the increased SHPT prevalence in the low range a result of more subjects

exhibiting a small decrease into this range? Causes of subnormal calcium may differ between populations, and may be influenced by variations in diet, supplementation, vitamin D status and other hormones. Mechanisms of intestinal uptake may need further characterization, including those of the classical active vitamin D dependent uptake of calcium in the proximal intestine.

Future studies should investigate how higher vitamin D levels and SHPT impact bone turnover and bone health. DXA can help to predict long-term bone health in the postoperative population. Risk factors for SHPT and bone health need more clarification, to develop treatment modalities for bone health. The processes of bone resorption, formation and mineralization should be investigated. Bone histomorphometry and bone quality assessments can improve our understanding of these. Adipose tissue, body mass and lean mass have effects to be explored. Adiposity related hormones, appetite regulating hormones, gut hormones and other signalling pathways need also more research.

We need improved evidence based guidelines to handle SHPT and bone health after obesity surgery. Substitution regimens must be investigated and optimized for calcium homeostasis and bone parameters. Phosphate need to be explored closer, being related with vitamin D, PTH and bone health. Future studies may help identify a relevance of magnesium, which we could not demonstrate. Protein substitution and physical activity studies may reveal positive effects on bone health. Treatment regimens with vitamin D analogs, estrogens and anti-osteoporotic drugs should also be explored.

As follows, the vitamin D and calcium levels are relevant for SHPT, but insufficiently explored. Our understanding of SHPT and bone health after obesity surgery is also incomplete. Resolving these issues has relevance for the future success of obesity surgery, and its application.

9. Conclusions

We can conclude that the prevalence of SHPT was relatively high 2-10 years after gastric bypass (study 1-4). A transient decrease in SHPT was observed the first 6 months, but this was no longer seen after 2 years (study 3). The prevalence of SHPT was very high after duodenal switch (study 1).

SHPT was consistently related with ionized calcium concentrations in the lower normal range or below. Mean ionized calcium was lower than normal after gastric bypass (and duodenal switch) (study 1-3), and the concentrations decreased over time after gastric bypass (study 3).

The prevalence of SHPT varied across the range of S-25(OH)D levels, with the highest prevalence in the vitamin D deficient patients (study 2 and 3). We observed small differences in SHPT comparing subjects exhibiting S-25(OH)D levels ≥ 75 nmol/l with levels ≥ 50 nmol/l after gastric bypass (study 1-3). The prevalence of SHPT was lowest with high S-25(OH)D levels ≥ 100 nmol/l in study 2 and 3, and the longitudinal data support that these high vitamin D levels can suppress PTH up to 5 years after gastric bypass (study 3).

Vitamin D levels were positively related with ionized calcium levels (study 2 and 3), but the ionized calcium levels decreased in all vitamin D subgroups up to 5 years (study 3).

SHPT was related with higher bone turnover markers, ALP and B-ALP (study 2 and 3). BMD was lower than expected 10 years after gastric bypass, and SHPT and bone turnover markers correlated with low BMD (study 4). Low-energy fractures occurred most often in the patients presenting with osteoporosis at 10 years.

Together these studies support that the vitamin D and calcium levels impact SHPT after obesity surgery, and that they can influence bone health. Keeping S-25(OH)D levels ≥ 50 nmol/l are essential for PTH and bone issues. S-25(OH)D levels ≥ 100 nmol/l may be necessary in some patients to suppress SHPT further, but how they impact bone parameters remain unanswered.

10. References

1. Midthjell K, Lee CM, Langhammer A, Krokstad S, Holmen TL, Hveem K, et al. Trends in overweight and obesity over 22 years in a large adult population: the HUNT Study, Norway. *Clin Obes*. 2013;3(1-2):12-20.
2. WHO. Obesity and overweight. Factsheet. <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight2020> [
3. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* (London, England). 2014;384(9945):766-81.
4. Mechanick JI, Kushner RF, Sugerman HJ, Gonzalez-Campoy JM, Collazo-Clavell ML, Spitz AF, et al. American Association of Clinical Endocrinologists, The Obesity Society, and American Society for Metabolic & Bariatric Surgery medical guidelines for clinical practice for the perioperative nutritional, metabolic, and nonsurgical support of the bariatric surgery patient. *Obesity*. 2009;17 Suppl 1:S1-70, v.
5. Helsedirektoratet. Utredning og behandling av sykelig overvekt i spesialisthelsetjenesten. Voksne. In: Hdir, editor. 2007.
6. Mechanick JI, Youdim A, Jones DB, Timothy Garvey W, Hurley DL, Molly McMahon M, et al. Clinical practice guidelines for the perioperative nutritional, metabolic, and nonsurgical support of the bariatric surgery patient--2013 update: cosponsored by American Association of Clinical Endocrinologists, the Obesity Society, and American Society for Metabolic & Bariatric Surgery. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*. 2013;9(2):159-91.
7. Løvås K, Husebye ES. *Endokrinologi*. 2013.
8. Gardella TJ, Nissenson RA, Jüppner H. Parathyroid hormone. In: Bilezikian JP, editor. *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism*, Ninth Edition: John Wiley & Sons, Inc; 2019. p. 205-11.
9. Ross AC, Manson JE, Abrams SA, Aloia JF, Brannon PM, Clinton SK, et al. The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *The Journal of clinical endocrinology and metabolism*. 2011;96(1):53-8.
10. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *The Journal of clinical endocrinology and metabolism*. 2011;96(7):1911-30.
11. Lamberg-Allardt C, Brustad M, Meyer HE, Steingrimsdottir L. Vitamin D - a systematic literature review for the 5th edition of the Nordic Nutrition Recommendations. *Food & nutrition research*. 2013;57.
12. Francis R, Aspray T, Fraser W, Gittoes N, Javaid K, Macdonald H, et al. *Vitamin D and Bone Health: A Practical Clinical Guideline for Patient Management*. 1 ed: National Osteoporosis Society; 2013.
13. Meena A, Tovey SC, Taylor CW. Sustained signalling by PTH modulates IP3 accumulation and IP3 receptors through cyclic AMP junctions. *J Cell Sci*. 2015;128(2):408-20.
14. Gardella TJ, Jüppner H, Brown EM, Kronenberg HM, Jr JTP. Parathyroid hormone and parathyroid hormone-related peptide in the regulation of calcium homeostasis and bone development. In: Jr JTP, editor. *The parathyroid gland and bone metabolism*. Philadelphia, US: Elsevier; 2013. p. 1-34.

15. Vautour L, Goltzman D. Regulation of Calcium Homeostasis. In: Bilezikian JP, editor. *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism*. Ninth Edition: Wiley & Sons, Inc; 2019. p. 165-72.
16. Biokjemi NsfM. Nasjonal brukerhåndbok i medisinsk biokjemi. In: Lindberg M, Hov GG, Hardang I, Monsen A-LB, editors. 2.0 ed. <https://brukerhandboken.no/2019>.
17. Schafer AL. Vitamin D and intestinal calcium transport after bariatric surgery. *The Journal of steroid biochemistry and molecular biology*. 2017;173:202-10.
18. Lips P. Vitamin D physiology. *Prog Biophys Mol Biol*. 2006;92(1):4-8.
19. Saleh F, Jorde R, Sundsfjord J, Haug E, Figenschau Y. Causes of secondary hyperparathyroidism in a healthy population: the Tromsø study. *Journal of bone and mineral metabolism*. 2006;24(1):58-64.
20. Hendy GN. Calcium-Sensing Receptor. In: John PB, editor. *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism* Ninth Edition: Wiley & Sons, Inc; 2019. p. 221-9.
21. Peterson LA, Zeng X, Caufield-Noll CP, Schweitzer MA, Magnuson TH, Steele KE. Vitamin D status and supplementation before and after bariatric surgery: a comprehensive literature review. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*. 2016;12(3):693-702.
22. Bikle DD, Adams JS, Christiakos S. Vitamin D: Production, Metabolism, Action, and Clinical Requirements. In: Bilezikian JP, editor. *Primer of Metabolic Bone Diseases and Disorders of Mineral Metabolism*, Ninth Edition: John Wiley & Sons, Inc; 2019. p. 230-40.
23. Frame-Peterson LA, McGill RD, Carobrese S, Schweitzer M. Nutrient Deficiencies Are Common Prior to Bariatric Surgery. *Nutrition in clinical practice : official publication of the American Society for Parenteral and Enteral Nutrition*. 2017;32(4):463-9.
24. Rejnmark L, Vestergaard P, Heickendorff L, Mosekilde L. Determinants of plasma PTH and their implication for defining a reference interval. *Clinical endocrinology*. 2011;74(1):37-43.
25. Khundmiri SJ, Murray RD, Lederer E. PTH and Vitamin D. *Compr Physiol*. 2016;6(2):561-601.
26. Osima M, Borgen TT, Lukic M, Grimnes G, Joakimsen RM, Eriksen EF, et al. Serum parathyroid hormone is associated with increased cortical porosity of the inner transitional zone at the proximal femur in postmenopausal women: the Tromsø Study. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2018;29(2):421-31.
27. Eriksen EF. Cellular mechanisms of bone remodeling. *Reviews in endocrine & metabolic disorders*. 2010;11(4):219-27.
28. Dirckx N, Maes C. Local and Circulating Osteoprogenitor Cells and Lineages. In: Bilezikian JP, editor. *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism*. Ninth: John Wiley & Son Inc; 2019. p. 20-30.
29. Holick MF. Vitamin D deficiency. *The New England journal of medicine*. 2007;357(3):266-81.
30. Parfitt AM, Oliver I, Villanueva AR. Bone histology in metabolic bone disease: the diagnostic value of bone biopsy. *Orthop Clin North Am*. 1979;10(2):329-45.
31. Ma YL, Zeng Q, Donley DW, Ste-Marie LG, Gallagher JC, Dalsky GP, et al. Teriparatide increases bone formation in modeling and remodeling osteons and enhances IGF-II immunoreactivity in postmenopausal women with osteoporosis. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2006;21(6):855-64.
32. WHO. Assessment of Fracture Risk and its application to screening for Postmenopausal Osteoporosis. WHO Technical Report Series 843. 1994.
33. (ISCD) TISfCD. 2019 ISCD Official Positions - Adult. ISCD; 2019.
34. Parfitt AM, Pødenphant J, Villanueva AR, Frame B. Metabolic bone disease with and without osteomalacia after intestinal bypass surgery: a bone histomorphometric study. *Bone*. 1985;6(4):211-20.
35. Rosen H. Use of biochemical markers of bone turnover in osteoporosis 2019 5th Nov 2020.
36. Stein EM, Silva BC, Boutroy S, Zhou B, Wang J, Udesky J, et al. Primary hyperparathyroidism is associated with abnormal cortical and trabecular microstructure and reduced bone stiffness in

postmenopausal women. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2013;28(5):1029-40.

37. Muñoz-Torres M, Manzanares Córdova R, García-Martín A, Avilés-Pérez MD, Nieto Serrano R, Andújar-Vera F, et al. Usefulness of Trabecular Bone Score (TBS) to Identify Bone Fragility in Patients with Primary Hyperparathyroidism. *Journal of clinical densitometry : the official journal of the International Society for Clinical Densitometry*. 2019;22(2):162-70.

38. Hansen S, Hauge EM, Rasmussen L, Jensen JE, Brixen K. Parathyroidectomy improves bone geometry and microarchitecture in female patients with primary hyperparathyroidism: a one-year prospective controlled study using high-resolution peripheral quantitative computed tomography. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2012;27(5):1150-8.

39. FHI. Overvekt og fedme blant voksne (indikator 14)

<https://www.fhi.no/nettpub/ncd/overvekt/voksne/>; Folkehelseinstituttet; 2017 [

40. Shekar M, (Ed.) BP. Obesity. Health and Economic Consequences of an Impending Global Challenge. Human Development Perspectives World Bank Group. 2020.

41. Branca F, Nikogosian H, (Ed.) TL. The challenge of obesity in the WHO Region and the strategies for response. In: WHO, editor. 2007.

42. WHO. Obesity: preventing and managing the global epidemic. Report of a WHO Consultation. WHO Technical Report Series 894. 2000.

43. Helsedirektoratet. Forebygging, utredning og behandling av overvekt og fedme hos voksne. Nasjonale retningslinjer for primærhelsetjenesten. 2010;IS-1735.

44. NIH. Gastrointestinal surgery for severe obesity. Proceedings of a National Institutes of Health Consensus Development Conference. March 25-27, 1991, Bethesda, MD. *The American journal of clinical nutrition*. 1992;55(2 Suppl):615s-9s.

45. Hofsø D, Aasheim ET, Søvik TT, Jakobsen GS, Johnson LK, Sandbu R, et al. [Follow-up after bariatric surgery]. *Tidsskrift for den Norske lægeforening : tidsskrift for praktisk medicin, ny række*. 2011;131(19):1887-92.

46. Sturm R, Hattori A. Morbid obesity rates continue to rise rapidly in the United States. *International journal of obesity (2005)*. 2013;37(6):889-91.

47. Pournaras DJ, Le Roux CW. The effect of bariatric surgery on gut hormones that alter appetite. *Diabetes Metab*. 2009;35(6 Pt 2):508-12.

48. Miras AD, le Roux CW. Mechanisms underlying weight loss after bariatric surgery. *Nat Rev Gastroenterol Hepatol*. 2013;10(10):575-84.

49. Greenway FL. Physiological adaptations to weight loss and factors favouring weight regain. *International journal of obesity (2005)*. 2015;39(8):1188-96.

50. Sánchez A, Rojas P, Basfi-Fer K, Carrasco F, Inostroza J, Codoceo J, et al. Micronutrient Deficiencies in Morbidly Obese Women Prior to Bariatric Surgery. *Obesity surgery*. 2016;26(2):361-8.

51. Himbert C, Ose J, Delphan M, Ulrich CM. A systematic review of the interrelation between diet- and surgery-induced weight loss and vitamin D status. *Nutrition research (New York, NY)*. 2017;38:13-26.

52. Aasheim ET, Hofsø D, Hjelmessaeth J, Birkeland KI, Bøhmer T. Vitamin status in morbidly obese patients: a cross-sectional study. *The American journal of clinical nutrition*. 2008;87(2):362-9.

53. Grace C, Vincent R, Aylwin SJ. High prevalence of vitamin D insufficiency in a United Kingdom urban morbidly obese population: implications for testing and treatment. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*. 2014;10(2):355-60.

54. Grethen E, McClintock R, Gupta CE, Jones R, Cacucci BM, Diaz D, et al. Vitamin D and hyperparathyroidism in obesity. *The Journal of clinical endocrinology and metabolism*. 2011;96(5):1320-6.

55. Pereira-Santos M, Costa PR, Assis AM, Santos CA, Santos DB. Obesity and vitamin D deficiency: a systematic review and meta-analysis. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2015;16(4):341-9.

56. Hultin H, Edfeldt K, Sundbom M, Hellman P. Left-shifted relation between calcium and parathyroid hormone in obesity. *The Journal of clinical endocrinology and metabolism*. 2010;95(8):3973-81.
57. Shapses SA, Sukumar D, Schneider SH, Schluskel Y, Brolin RE, Taich L. Hormonal and dietary influences on true fractional calcium absorption in women: role of obesity. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2012;23(11):2607-14.
58. Grethen E, Hill KM, Jones R, Cacucci BM, Gupta CE, Acton A, et al. Serum leptin, parathyroid hormone, 1,25-dihydroxyvitamin D, fibroblast growth factor 23, bone alkaline phosphatase, and sclerostin relationships in obesity. *The Journal of clinical endocrinology and metabolism*. 2012;97(5):1655-62.
59. Riedt CS, Brolin RE, Sherrell RM, Field MP, Shapses SA. True fractional calcium absorption is decreased after Roux-en-Y gastric bypass surgery. *Obesity*. 2006;14(11):1940-8.
60. Sneve M, Emaus N, Joakimsen RM, Jorde R. The association between serum parathyroid hormone and bone mineral density, and the impact of smoking: the Tromsø Study. *European journal of endocrinology*. 2008;158(3):401-9.
61. Gagnon C, Schafer AL. Bone Health After Bariatric Surgery. *JBMR plus*. 2018;2(3):121-33.
62. Avgerinos DV, Leitman IM, Martínez RE, Liao EP. Evaluation of markers for calcium homeostasis in a population of obese adults undergoing gastric bypass operations. *J Am Coll Surg*. 2007;205(2):294-7.
63. Hjelmsaeth J, Hofsvø D, Aasheim ET, Jenssen T, Moan J, Hager H, et al. Parathyroid hormone, but not vitamin D, is associated with the metabolic syndrome in morbidly obese women and men: a cross-sectional study. *Cardiovasc Diabetol*. 2009;8:7.
64. Guglielmi V, Bellia A, Gentileschi P, Lombardo M, D'Adamo M, Lauro D, et al. Parathyroid hormone in surgery-induced weight loss: no glucometabolic effects but potential adaptive response to skeletal loading. *Endocrine*. 2018;59(2):288-95.
65. Walsh JS, Bowles S, Evans AL. Vitamin D in obesity. *Current opinion in endocrinology, diabetes, and obesity*. 2017;24(6):389-94.
66. Evans AL, Paggiosi MA, Eastell R, Walsh JS. Bone density, microstructure and strength in obese and normal weight men and women in younger and older adulthood. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2015;30(5):920-8.
67. Ivaska KK, Huovinen V, Soinio M, Hannukainen JC, Saunavaara V, Salminen P, et al. Changes in bone metabolism after bariatric surgery by gastric bypass or sleeve gastrectomy. *Bone*. 2017;95:47-54.
68. Viljakainen H, Ivaska KK, Paldanius P, Lipsanen-Nyman M, Saukkonen T, Pietiläinen KH, et al. Suppressed bone turnover in obesity: a link to energy metabolism? A case-control study. *The Journal of clinical endocrinology and metabolism*. 2014;99(6):2155-63.
69. Fuglsang-Nielsen R, Rakvaag E, Vestergaard P, Hartmann B, Holst JJ, Hermansen K, et al. Consumption of nutrients and insulin resistance suppress markers of bone turnover in subjects with abdominal obesity. *Bone*. 2020;133:115230.
70. Stein EM, Silverberg SJ. Bone loss after bariatric surgery: causes, consequences, and management. *Lancet Diabetes Endocrinol*. 2014;2(2):165-74.
71. Savvidis C, Tournis S, Dede AD. Obesity and bone metabolism. *Hormones (Athens, Greece)*. 2018;17(2):205-17.
72. Chen XX, Yang T. Roles of leptin in bone metabolism and bone diseases. *Journal of bone and mineral metabolism*. 2015;33(5):474-85.
73. Turcotte AF, Grenier-Larouche T, Ung RV, Simonyan D, Carreau AM, Carpentier AC, et al. Effects of Biliopancreatic Diversion on Bone Turnover Markers and Association with Hormonal Factors in Patients with Severe Obesity. *Obesity surgery*. 2019;29(3):990-8.
74. Yu EW. Bone metabolism after bariatric surgery. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2014;29(7):1507-18.

75. Knapp KM, Welsman JR, Hopkins SJ, Fogelman I, Blake GM. Obesity increases precision errors in dual-energy X-ray absorptiometry measurements. *Journal of clinical densitometry : the official journal of the International Society for Clinical Densitometry*. 2012;15(3):315-9.
76. Yu EW, Thomas BJ, Brown JK, Finkelstein JS. Simulated increases in body fat and errors in bone mineral density measurements by DXA and QCT. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2012;27(1):119-24.
77. Sukumar D, Schluskel Y, Riedt CS, Gordon C, Stahl T, Shapses SA. Obesity alters cortical and trabecular bone density and geometry in women. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2011;22(2):635-45.
78. Folli F, Sabowitz BN, Schwesinger W, Fanti P, Guardado-Mendoza R, Muscogiuri G. Bariatric surgery and bone disease: from clinical perspective to molecular insights. *International journal of obesity (2005)*. 2012;36(11):1373-9.
79. Khosla S, Oursler MJ, Monroe DG. Estrogen and the skeleton. *Trends in endocrinology and metabolism: TEM*. 2012;23(11):576-81.
80. Joseph F, Ahmad AM, Ul-Haq M, Durham BH, Whittingham P, Fraser WD, et al. Effects of growth hormone administration on bone mineral metabolism, PTH sensitivity and PTH secretory rhythm in postmenopausal women with established osteoporosis. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2008;23(5):721-9.
81. Reid IR. Relationships between fat and bone. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2008;19(5):595-606.
82. Cosman F, Shen V, Xie F, Seibel M, Ratcliffe A, Lindsay R. Estrogen protection against bone resorbing effects of parathyroid hormone infusion. Assessment by use of biochemical markers. *Annals of internal medicine*. 1993;118(5):337-43.
83. Shapses SA, Pop LC, Wang Y. Obesity is a concern for bone health with aging. *Nutrition research (New York, NY)*. 2017;39:1-13.
84. De Laet C, Kanis JA, Oden A, Johanson H, Johnell O, Delmas P, et al. Body mass index as a predictor of fracture risk: a meta-analysis. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2005;16(11):1330-8.
85. Blom-Høgestøl IK, Mala T, Kristinsson JA, Hauge EM, Brunborg C, Gulseth HL, et al. Changes in Bone Marrow Adipose Tissue One Year After Roux-en-Y Gastric Bypass: A Prospective Cohort Study. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2019;34(10):1815-23.
86. Compston J. Obesity and fractures in postmenopausal women. *Current opinion in rheumatology*. 2015;27(4):414-9.
87. Sjøgaard AJ, Holvik K, Omsland TK, Tell GS, Dahl C, Schei B, et al. Abdominal obesity increases the risk of hip fracture. A population-based study of 43,000 women and men aged 60-79 years followed for 8 years. *Cohort of Norway*. *Journal of internal medicine*. 2015;277(3):306-17.
88. Paik JM, Rosen HN, Katz JN, Rosner BA, Rimm EB, Gordon CM, et al. BMI, Waist Circumference, and Risk of Incident Vertebral Fracture in Women. *Obesity*. 2019;27(9):1513-9.
89. Wilsaard T, Jacobsen BK, Ahmed LA, Joakimsen RM, Stormer J, Jorgensen L. BMI change is associated with fracture incidence, but only in non-smokers. The Tromsø Study. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2011;22(4):1237-45.
90. Faria GR. A brief history of bariatric surgery. *Porto biomedical journal*. 2017;2(3):90-2.
91. Løvig T. Overvektskirurgi med spesielt fokus på utviklingen i Norge. *Kirurgen*. 2008(Nr 4):8-12.
92. Våge V, Solhaug JH, Berstad A, Svanes K, Viste A. Jejunoileal bypass in the treatment of morbid obesity: a 25-year follow-up study of 36 patients. *Obesity surgery*. 2002;12(3):312-8.

93. Scibora LM, Ikramuddin S, Buchwald H, Petit MA. Examining the link between bariatric surgery, bone loss, and osteoporosis: a review of bone density studies. *Obesity surgery*. 2012;22(4):654-67.
94. Sjostrom L, Lindroos AK, Peltonen M, Torgerson J, Bouchard C, Carlsson B, et al. Lifestyle, diabetes, and cardiovascular risk factors 10 years after bariatric surgery. *The New England journal of medicine*. 2004;351(26):2683-93.
95. Sjostrom L, Narbro K, Sjostrom CD, Karason K, Larsson B, Wedel H, et al. Effects of bariatric surgery on mortality in Swedish obese subjects. *The New England journal of medicine*. 2007;357(8):741-52.
96. Kunnskapssenteret. Langtidseffekter etter fedmekirurgi. 2014;2014.
97. Våge V, Lyyjynen H. Årsrapport for 2019 med plan for forbedringstiltak. <https://helse-bergen.no/avdelinger/laboratorieklinikken/medisinsk-biokjemi-og-farmakologi/hormonlaboratoriet/norsk-kvalitetsregister-for-fedmekirurgi-soreg-norge#arsrapport>: Norsk kvalitetsregister for fedmekirurgi (Soreg-N); 2020.
98. Isom KA, Andromalos L, Ariagno M, Hartman K, Mogensen KM, Stephanides K, et al. Nutrition and metabolic support recommendations for the bariatric patient. *Nutrition in clinical practice : official publication of the American Society for Parenteral and Enteral Nutrition*. 2014;29(6):718-39.
99. Rickers H, Christiansen C, Balslev I, Rødbro P. Impairment of vitamin D metabolism and bone mineral content after intestinal bypass for obesity. A longitudinal study. *Scand J Gastroenterol*. 1984;19(2):184-9.
100. Newbury L, Dolan K, Hatzifotis M, Low N, Fielding G. Calcium and vitamin D depletion and elevated parathyroid hormone following biliopancreatic diversion. *Obesity surgery*. 2003;13(6):893-5.
101. Våge V, Gjesdal CG, Eide GE, Halse JI, Viste A. Bone mineral density in females after jejunoileal bypass: a 25-year follow-up study. *Obesity surgery*. 2004;14(3):305-12.
102. De Prisco C, Levine SN. Metabolic bone disease after gastric bypass surgery for obesity. *The American journal of the medical sciences*. 2005;329(2):57-61.
103. Crowley LV, Seay J, Mullin G. Late effects of gastric bypass for obesity. *The American journal of gastroenterology*. 1984;79(11):850-60.
104. Ott MT, Fanti P, Malluche HH, Ryo UY, Whaley FS, Strodel WE, et al. Biochemical Evidence of Metabolic Bone Disease in Women Following Roux-Y Gastric Bypass for Morbid Obesity. *Obesity surgery*. 1992;2(4):341-8.
105. Jin J, Robinson AV, Hallowell PT, Jasper JJ, Stellato TA, Wilhem SM. Increases in parathyroid hormone (PTH) after gastric bypass surgery appear to be of a secondary nature. *Surgery*. 2007;142(6):914-20; discussion -20.
106. Johnson JM, Maher JW, Samuel I, Heitshusen D, Doherty C, Downs RW. Effects of gastric bypass procedures on bone mineral density, calcium, parathyroid hormone, and vitamin D. *J Gastrointest Surg*. 2005;9(8):1106-10; discussion 10-1.
107. Fleischer J, Stein EM, Bessler M, Della Badia M, Restuccia N, Olivero-Rivera L, et al. The decline in hip bone density after gastric bypass surgery is associated with extent of weight loss. *The Journal of clinical endocrinology and metabolism*. 2008;93(10):3735-40.
108. Hamoui N, Kim K, Anthone G, Crookes PF. The significance of elevated levels of parathyroid hormone in patients with morbid obesity before and after bariatric surgery. *Archives of surgery (Chicago, Ill : 1960)*. 2003;138(8):891-7.
109. Coates PS, Fernstrom JD, Fernstrom MH, Schauer PR, Greenspan SL. Gastric bypass surgery for morbid obesity leads to an increase in bone turnover and a decrease in bone mass. *The Journal of clinical endocrinology and metabolism*. 2004;89(3):1061-5.
110. Guney E, Kisakol G, Ozgen G, Yilmaz C, Yilmaz R, Kabalak T. Effect of weight loss on bone metabolism: comparison of vertical banded gastroplasty and medical intervention. *Obesity surgery*. 2003;13(3):383-8.
111. Marceau P, Biron S, Lebel S, Marceau S, Hould FS, Simard S, et al. Does bone change after biliopancreatic diversion? *J Gastrointest Surg*. 2002;6(5):690-8.

112. Chakhtoura M, Rahme M, El-Hajj Fuleihan G. Vitamin D Metabolism in Bariatric Surgery. *Endocrinology and metabolism clinics of North America*. 2017;46(4):947-82.
113. Aasheim ET, Björkman S, Søvik TT, Engström M, Hanvold SE, Mala T, et al. Vitamin status after bariatric surgery: a randomized study of gastric bypass and duodenal switch. *The American journal of clinical nutrition*. 2009;90(1):15-22.
114. Aarts E, van Groningen L, Horst R, Telting D, van Sorge A, Janssen I, et al. Vitamin D absorption: consequences of gastric bypass surgery. *European journal of endocrinology*. 2011;164(5):827-32.
115. Gasteyger C, Suter M, Gaillard RC, Giusti V. Nutritional deficiencies after Roux-en-Y gastric bypass for morbid obesity often cannot be prevented by standard multivitamin supplementation. *The American journal of clinical nutrition*. 2008;87(5):1128-33.
116. Goode LR, Brolin RE, Chowdhury HA, Shapses SA. Bone and gastric bypass surgery: effects of dietary calcium and vitamin D. *Obesity research*. 2004;12(1):40-7.
117. DiGiorgi M, Daud A, Inabnet WB, Schroepe B, Urban-Skuro M, Restuccia N, et al. Markers of bone and calcium metabolism following gastric bypass and laparoscopic adjustable gastric banding. *Obesity surgery*. 2008;18(9):1144-8.
118. El-Kadre LJ, Rocha PR, de Almeida Tinoco AC, Tinoco RC. Calcium metabolism in pre- and postmenopausal morbidly obese women at baseline and after laparoscopic Roux-en-Y gastric bypass. *Obesity surgery*. 2004;14(8):1062-6.
119. Johnson JM, Maher JW, DeMaria EJ, Downs RW, Wolfe LG, Kellum JM. The long-term effects of gastric bypass on vitamin D metabolism. *Annals of surgery*. 2006;243(5):701-4; discussion 4-5.
120. Søvik TT, Aasheim ET, Taha O, Engström M, Fagerland MW, Björkman S, et al. Weight loss, cardiovascular risk factors, and quality of life after gastric bypass and duodenal switch: a randomized trial. *Annals of internal medicine*. 2011;155(5):281-91.
121. Aills L, Blankenship J, Buffington C, Furtado M, Parrott J. ASMBS Allied Health Nutritional Guidelines for the Surgical Weight Loss Patient. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*. 2008;4(5 Suppl):S73-108.
122. Diniz Mde F, Diniz MT, Sanches SR, Salgado PP, Valadão MM, Araújo FC, et al. Elevated serum parathormone after Roux-en-Y gastric bypass. *Obesity surgery*. 2004;14(9):1222-6.
123. da Rosa CL, Dames Olivieri Saubermann AP, Jacqueline J, Pereira SE, Saboya C, Ramalho A. Routine supplementation does not warrant the nutritional status of vitamin d adequate after gastric bypass Roux-en-Y. *Nutr Hosp*. 2013;28(1):169-72.
124. Fish E, Beverstein G, Olson D, Reinhardt S, Garren M, Gould J. Vitamin D status of morbidly obese bariatric surgery patients. *The Journal of surgical research*. 2010;164(2):198-202.
125. Aasheim ET, Johnson LK, Hofsø D, Bøhmer T, Hjelmæsæth J. Vitamin status after gastric bypass and lifestyle intervention: a comparative prospective study. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*. 2012;8(2):169-75.
126. Heber D, Greenway FL, Kaplan LM, Livingston E, Salvador J, Still C. Endocrine and nutritional management of the post-bariatric surgery patient: an Endocrine Society Clinical Practice Guideline. *The Journal of clinical endocrinology and metabolism*. 2010;95(11):4823-43.
127. Lespessailles E, Toumi H. Vitamin D alteration associated with obesity and bariatric surgery. *Exp Biol Med (Maywood)*. 2017;242(10):1086-94.
128. O'Kane M, Pinkney J, Aasheim ET, Barth J, Batterham RL, Welbourn R. BOMSS Guidelines on perioperative and postoperative biochemical monitoring and micronutrient replacement for patients undergoing bariatric surgery. <https://www.bomss.org.uk/bomss-nutritional-guidance/> (accessed): BOMSS; 2014.
129. Chakhtoura MT, Nakhoul NN, Shawwa K, Mantzoros C, El Hajj Fuleihan GA. Hypovitaminosis D in bariatric surgery: A systematic review of observational studies. *Metabolism: clinical and experimental*. 2016;65(4):574-85.
130. O'Kane M, Parretti HM, Pinkney J, Welbourn R, Hughes CA, Mok J, et al. British Obesity and Metabolic Surgery Society Guidelines on perioperative and postoperative biochemical monitoring

- and micronutrient replacement for patients undergoing bariatric surgery-2020 update. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2020.
131. Vilarrasa N, de Gordejuela AG, Gómez-Vaquero C, Pujol J, Elio I, San José P, et al. Effect of bariatric surgery on bone mineral density: comparison of gastric bypass and sleeve gastrectomy. *Obesity surgery*. 2013;23(12):2086-91.
 132. Switzer NJ, Marcil G, Prasad S, Debru E, Church N, Mitchell P, et al. Long-term hypovitaminosis D and secondary hyperparathyroidism outcomes of the Roux-en-Y gastric bypass: a systematic review. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2017;18(5):560-6.
 133. Sinha N, Shieh A, Stein EM, Strain G, Schulman A, Pomp A, et al. Increased PTH and 1.25(OH)(2)D levels associated with increased markers of bone turnover following bariatric surgery. *Obesity*. 2011;19(12):2388-93.
 134. Toelle P, Peterli R, Zobel I, Noppen C, Christoffel-Courtin C, Peters T. Risk factors for secondary hyperparathyroidism after bariatric surgery: a comparison of 4 different operations and of vitamin D-receptor-polymorphism. *Exp Clin Endocrinol Diabetes*. 2012;120(10):629-34.
 135. Slater GH, Ren CJ, Siegel N, Williams T, Barr D, Wolfe B, et al. Serum fat-soluble vitamin deficiency and abnormal calcium metabolism after malabsorptive bariatric surgery. *J Gastrointest Surg*. 2004;8(1):48-55; discussion 4-5.
 136. Valderas JP, Velasco S, Solari S, Liberona Y, Viviani P, Maiz A, et al. Increase of bone resorption and the parathyroid hormone in postmenopausal women in the long-term after Roux-en-Y gastric bypass. *Obesity surgery*. 2009;19(8):1132-8.
 137. Vilarrasa N, San José P, García I, Gómez-Vaquero C, Miras PM, de Gordejuela AG, et al. Evaluation of bone mineral density loss in morbidly obese women after gastric bypass: 3-year follow-up. *Obesity surgery*. 2011;21(4):465-72.
 138. Stein J, Stier C, Raab H, Weiner R. Review article: The nutritional and pharmacological consequences of obesity surgery. *Alimentary pharmacology & therapeutics*. 2014;40(6):582-609.
 139. Liu C, Wu D, Zhang JF, Xu D, Xu WF, Chen Y, et al. Changes in Bone Metabolism in Morbidly Obese Patients After Bariatric Surgery: A Meta-Analysis. *Obesity surgery*. 2016;26(1):91-7.
 140. Balsa JA, Botella-Carretero JI, Peromingo R, Zamarrón I, Arrieta F, Muñoz-Malo T, et al. Role of calcium malabsorption in the development of secondary hyperparathyroidism after biliopancreatic diversion. *J Endocrinol Invest*. 2008;31(10):845-50.
 141. Youssef Y, Richards WO, Sekhar N, Kaiser J, Spagnoli A, Abumrad N, et al. Risk of secondary hyperparathyroidism after laparoscopic gastric bypass surgery in obese women. *Surg Endosc*. 2007;21(8):1393-6.
 142. Clements RH, Yellumahanthi K, Wesley M, Ballem N, Bland KI. Hyperparathyroidism and vitamin D deficiency after laparoscopic gastric bypass. *The American surgeon*. 2008;74(6):469-74; discussion 74-5.
 143. Cundy T, Evans MC, Kay RG, Dowman M, Wattie D, Reid IR. Effects of vertical-banded gastroplasty on bone and mineral metabolism in obese patients. *The British journal of surgery*. 1996;83(10):1468-72.
 144. Yu EW, Wewalka M, Ding SA, Simonson DC, Foster K, Holst JJ, et al. Effects of Gastric Bypass and Gastric Banding on Bone Remodeling in Obese Patients With Type 2 Diabetes. *The Journal of clinical endocrinology and metabolism*. 2016;101(2):714-22.
 145. Bruno C, Fulford AD, Potts JR, McClintock R, Jones R, Cacucci BM, et al. Serum markers of bone turnover are increased at six and 18 months after Roux-en-Y bariatric surgery: correlation with the reduction in leptin. *The Journal of clinical endocrinology and metabolism*. 2010;95(1):159-66.
 146. Harper C, Pattinson AL, Fernando HA, Zibellini J, Seimon RV, Sainsbury A. Effects of obesity treatments on bone mineral density, bone turnover and fracture risk in adults with overweight or obesity. *Hormone molecular biology and clinical investigation*. 2016;28(3):133-49.
 147. Compston JE, Horton LW, Laker MF, Ayers AB, Woodhead JS, Bull HJ, et al. Bone disease after jejuno-ileal bypass for obesity. *Lancet (London, England)*. 1978;2(8079):1-4.

148. Balsa JA, Botella-Carretero JJ, Peromingo R, Caballero C, Muñoz-Malo T, Villafruela JJ, et al. Chronic increase of bone turnover markers after biliopancreatic diversion is related to secondary hyperparathyroidism and weight loss. Relation with bone mineral density. *Obesity surgery*. 2010;20(4):468-73.
149. Vilarrasa N, Gómez JM, Elio I, Gómez-Vaquero C, Masdevall C, Pujol J, et al. Evaluation of bone disease in morbidly obese women after gastric bypass and risk factors implicated in bone loss. *Obesity surgery*. 2009;19(7):860-6.
150. Brzozowska MM, Sainsbury A, Eisman JA, Baldock PA, Center JR. Bariatric surgery, bone loss, obesity and possible mechanisms. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2013;14(1):52-67.
151. Stein EM, Carrelli A, Young P, Bucovsky M, Zhang C, Schrope B, et al. Bariatric surgery results in cortical bone loss. *The Journal of clinical endocrinology and metabolism*. 2013;98(2):541-9.
152. Coupaye M, Breuil MC, Rivière P, Castel B, Bogard C, Dupré T, et al. Serum vitamin D increases with weight loss in obese subjects 6 months after Roux-en-Y gastric bypass. *Obesity surgery*. 2013;23(4):486-93.
153. Lalmohamed A, de Vries F, Bazelier MT, Cooper A, van Staa TP, Cooper C, et al. Risk of fracture after bariatric surgery in the United Kingdom: population based, retrospective cohort study. *BMJ (Clinical research ed)*. 2012;345:e5085.
154. Nakamura KM, Haglund EG, Clowes JA, Achenbach SJ, Atkinson EJ, Melton LJ, 3rd, et al. Fracture risk following bariatric surgery: a population-based study. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2014;25(1):151-8.
155. Henry HL, Bouillon R, Norman AW, Gallagher JC, Lips P, Heaney RP, et al. 14th Vitamin D Workshop consensus on vitamin D nutritional guidelines. *The Journal of steroid biochemistry and molecular biology*. 2010;121(1-2):4-6.
156. Cashman KD, Dowling KG, Škrabáková Z, Kiely M, Lamberg-Allardt C, Durazo-Arvizu RA, et al. Standardizing serum 25-hydroxyvitamin D data from four Nordic population samples using the Vitamin D Standardization Program protocols: Shedding new light on vitamin D status in Nordic individuals. *Scand J Clin Lab Invest*. 2015;75(7):549-61.
157. Bresson J-L, Burlingame B, Dean T, Fairweather-Tait S, Heinonen M, Hirsch-Ernst K-I, et al. Dietary reference values for vitamin D. EFSA NDA Panel, 2016. *EFSA Journal* 2016. p. 4547.
158. Frigstad SO, Høivik M, Jahnsen J, Dahl SR, Cvancarova M, Grimstad T, et al. Vitamin D deficiency in inflammatory bowel disease: prevalence and predictors in a Norwegian outpatient population. *Scand J Gastroenterol*. 2017;52(1):100-6.
159. Delmas PD, Eastell R, Garnero P, Seibel MJ, Stepan J. The use of biochemical markers of bone turnover in osteoporosis. Committee of Scientific Advisors of the International Osteoporosis Foundation. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2000;11 Suppl 6:S2-17.
160. Chinoy MA, Javed MI, Khan A, Sadruddin N. Alkaline phosphatase as a screening test for osteomalacia. *J Ayub Med Coll Abbottabad*. 2011;23(1):23-5.
161. Genant HK, Cooper C, Poor G, Reid I, Ehrlich G, Kanis J, et al. Interim report and recommendations of the World Health Organization Task-Force for Osteoporosis. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 1999;10(4):259-64.
162. Aftab H, Risstad H, Sjøvik TT, Bernklev T, Hewitt S, Kristinsson JA, et al. Five-year outcome after gastric bypass for morbid obesity in a Norwegian cohort. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*. 2014;10(1):71-8.
163. Veierød MB, Lydersen S, Laake P. Design and analysis. In: Veierød MB, Lydersen S, Laake P, editors. *Medical statistics in clinical and epidemiological research*. 1 ed. Oslo: Gyldendal Norsk Forlag; 2012.

164. Fitzmaurice GM, Laird NM, Ware JH. Introduction to Longitudinal and Clustered Data. In: Fitzmaurice GM, Laird NM, Ware JH, editors. *Applied Longitudinal Analysis*. 2nd ed. Hoboken, New Jersey: John Wiley & Sons, Inc; 2011. p. 1-18.
165. Higa K, Ho T, Tercero F, Yunus T, Boone KB. Laparoscopic Roux-en-Y gastric bypass: 10-year follow-up. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*. 2011;7(4):516-25.
166. Harper J, Madan AK, Ternovits CA, Tichansky DS. What happens to patients who do not follow-up after bariatric surgery? *The American surgeon*. 2007;73(2):181-4.
167. Blom-Hogestol IK, Hewitt S, Chahal-Kummen M, Brunborg C, Gulseth HL, Kristinsson JA, et al. Bone metabolism, bone mineral density and low-energy fractures 10years after Roux-en-Y gastric bypass. *Bone*. 2019;127:436-45.
168. Chahal-Kummen M, Salte OBK, Hewitt S, Blom-Høgestøl IK, Risstad H, Kristinsson J, et al. Health benefits and risks during 10 years after Roux-en-Y gastric bypass. *Surg Endosc*. 2020.
169. Høgestøl IK, Chahal-Kummen M, Eribe I, Brunborg C, Stubhaug A, Hewitt S, et al. Chronic Abdominal Pain and Symptoms 5 Years After Gastric Bypass for Morbid Obesity. *Obesity surgery*. 2017;27(6):1438-45.
170. Meyer HE, Falch JA, Sogaard AJ, Haug E. Vitamin D deficiency and secondary hyperparathyroidism and the association with bone mineral density in persons with Pakistani and Norwegian background living in Oslo, Norway, The Oslo Health Study. *Bone*. 2004;35(2):412-7.
171. Jorde R, Grimnes G. Serum PTH is not a good marker for defining a threshold for vitamin D deficiency. *Endocr Connect*. 2020;9(5):396-404.
172. Hewitt S, Kristinsson J, Aasheim ET, Blom-Høgestøl IK, Aaseth E, Jahnsen J, et al. Relationships Between Vitamin D Status and PTH over 5 Years After Roux-en-Y Gastric Bypass: a Longitudinal Cohort Study. *Obesity surgery*. 2020;30(9):3426-34.
173. Hewitt S, Aasheim ET, Sovik TT, Jahnsen J, Kristinsson J, Eriksen EF, et al. Relationships of serum 25-hydroxyvitamin D, ionized calcium and parathyroid hormone after obesity surgery. *Clinical endocrinology*. 2018;88(3):372-9.
174. Watts NB. Clinical utility of biochemical markers of bone remodeling. *Clin Chem*. 1999;45(8 Pt 2):1359-68.
175. Lewiecki M. Osteoporotic fracture risk assessment 2020 30th Oct 2020.
176. Hanvold SE. Health benefits two years after Roux-en-Y gastric bypass surgery and the effect of lifestyle intervention two to four years after surgery on weight regain and metabolic disturbances. [PhD Thesis]: University of Oslo; 2018.
177. Bergh I. The role of psychosocial factors in predicting weight loss and adherence to health behaviours in patients undergoing obesity surgery - A prospective cohort study. [PhD Thesis]. Oslo: University of Oslo; 2017.
178. Ketteler M, Block GA, Evenepoel P, Fukagawa M, Herzog CA, McCann L, et al. Executive summary of the 2017 KDIGO Chronic Kidney Disease-Mineral and Bone Disorder (CKD-MBD) Guideline Update: what's changed and why it matters. *Kidney Int*. 2017;92(1):26-36.
179. Hawley CM, Holt SG. Parathyroid hormone targets in chronic kidney disease and managing severe hyperparathyroidism. *Nephrology (Carlton)*. 2017;22 Suppl 2:47-50.
180. Harris M, Taylor G. *Medical statistics made easy*. 2nd ed. Oxfordshire, UK: Scion Publishing Ltd; 2008.
181. le Cessie S, Goeman JJ, Dekkers OM. Who is afraid of non-normal data? Choosing between parametric and non-parametric tests. *European journal of endocrinology*. 2020;182(2):E1-e3.
182. *Introduction to Statistics*. Course MF9130: University of Oslo; 2015.
183. Fitzmaurice GM, Laird NM, Ware JH. Linear Mixed Effects Models. In: Fitzmaurice GM, Laird NM, Ware JH, editors. *Applied Longitudinal Analysis*. 2nd ed. Hoboken, New Jersey: John Wiley & Sons, Inc; 2011.
184. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet (London, England)*. 2007;370(9596):1453-7.

185. Mônico-Ferreira DV, Leandro-Merhi VA, Aranha NC, Brandalise A, Brandalise NA. VITAMIN D DEFICIENCY AND PARATHYROID HORMONE INCREASE IN LATE POSTOPERATIVE GASTRIC BYPASS IN ROUX-EN-Y. *Arquivos brasileiros de cirurgia digestiva : ABCD = Brazilian archives of digestive surgery*. 2018;31(4):e1407.
186. Raoof M, Naslund I, Rask E, Szabo E. Effect of Gastric Bypass on Bone Mineral Density, Parathyroid Hormone and Vitamin D: 5 Years Follow-up. *Obesity surgery*. 2016;26(5):1141-5.
187. Karefylakis C, Naslund I, Edholm D, Sundbom M, Karlsson FA, Rask E. Vitamin D status 10 years after primary gastric bypass: gravely high prevalence of hypovitaminosis D and raised PTH levels. *Obesity surgery*. 2014;24(3):343-8.
188. Fox A, Slater C, Ahmed B, Ammori BJ, Senapati S, Akhtar K, et al. Vitamin D Status After Gastric Bypass or Sleeve Gastrectomy over 4 Years of Follow-up. *Obesity surgery*. 2020;30(4):1473-81.
189. Obeid NR, Malick W, Concors SJ, Fielding GA, Kurian MS, Ren-Fielding CJ. Long-term outcomes after Roux-en-Y gastric bypass: 10- to 13-year data. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*. 2016;12(1):11-20.
190. Crawford MR, Pham N, Khan L, Bena JF, Schauer PR, Kashyap SR. INCREASED BONE TURNOVER IN TYPE 2 DIABETES PATIENTS RANDOMIZED TO BARIATRIC SURGERY VERSUS MEDICAL THERAPY AT 5 YEARS. *Endocr Pract*. 2018;24(3):256-64.
191. Carlin AM, Rao DS, Yager KM, Parikh NJ, Kapke A. Treatment of vitamin D depletion after Roux-en-Y gastric bypass: a randomized prospective clinical trial. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*. 2009;5(4):444-9.
192. Muschitz C, Kocijan R, Haschka J, Zendeli A, Pirker T, Geiger C, et al. The Impact of Vitamin D, Calcium, Protein Supplementation, and Physical Exercise on Bone Metabolism After Bariatric Surgery: The BABS Study. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2016;31(3):672-82.
193. Goldner WS, Stoner JA, Lyden E, Thompson J, Taylor K, Larson L, et al. Finding the optimal dose of vitamin D following Roux-en-Y gastric bypass: a prospective, randomized pilot clinical trial. *Obesity surgery*. 2009;19(2):173-9.
194. Smelt HJM, Pouwels S, Smulders JF. The Influence of Different Cholecalciferol Supplementation Regimes on 25(OH) Cholecalciferol, Calcium and Parathyroid Hormone after Bariatric Surgery. *Medicina (Kaunas)*. 2019;55(6).
195. Salazar DA, Ferreira MJS, Neves JS, Pedro JMP, Guerreiro VA, S ESV, et al. Variable Thresholds of Vitamin D Plasma Levels to Suppress PTH: the Effect of Weight and Bariatric Surgery. *Obesity surgery*. 2020;30(4):1551-9.
196. Kontogeorgos G, Welin L, Fu M, Hansson PO, Landin-Wilhelmsen K, Laine CM. Hyperparathyroidism in men - morbidity and mortality during 21 years' follow-up. *Scand J Clin Lab Invest*. 2020;80(1):6-13.
197. Kontogeorgos G, Trimpou P, Laine CM, Olerod G, Lindahl A, Landin-Wilhelmsen K. Normocalcaemic, vitamin D-sufficient hyperparathyroidism - high prevalence and low morbidity in the general population: A long-term follow-up study, the WHO MONICA project, Gothenburg, Sweden. *Clinical endocrinology*. 2015;83(2):277-84.
198. Shah M, Sharma A, Wermers RA, Kennel KA, Kellogg TA, Mundi MS. Hypocalcemia After Bariatric Surgery: Prevalence and Associated Risk Factors. *Obesity surgery*. 2017;27(11):2905-11.
199. Carrasco F, Basfi-Fer K, Rojas P, Csendes A, Papapietro K, Codoceo J, et al. Calcium absorption may be affected after either sleeve gastrectomy or Roux-en-Y gastric bypass in premenopausal women: a 2-y prospective study. *The American journal of clinical nutrition*. 2018;108(1):24-32.
200. Wei JH, Lee WJ, Chong K, Lee YC, Chen SC, Huang PH, et al. High Incidence of Secondary Hyperparathyroidism in Bariatric Patients: Comparing Different Procedures. *Obesity surgery*. 2018;28(3):798-804.
201. Risstad H, Sovik TT, Engstrom M, Aasheim ET, Fagerland MW, Olsen MF, et al. Five-year outcomes after laparoscopic gastric bypass and laparoscopic duodenal switch in patients with body mass index of 50 to 60: a randomized clinical trial. *JAMA surgery*. 2015;150(4):352-61.

202. Svanevik M, Risstad H, Hofso D, Blom-Hogestol IK, Kristinsson JA, Sandbu R, et al. Bone Turnover Markers After Standard and Distal Roux-en-Y Gastric Bypass: Results from a Randomized Controlled Trial. *Obesity surgery*. 2019.
203. Gillon S, Jeanes YM, Andersen JR, Våge V. Micronutrient Status in Morbidly Obese Patients Prior to Laparoscopic Sleeve Gastrectomy and Micronutrient Changes 5 years Post-surgery. *Obesity surgery*. 2017;27(3):606-12.
204. Schafer AL, Weaver CM, Black DM, Wheeler AL, Chang H, Szefc GV, et al. Intestinal Calcium Absorption Decreases Dramatically After Gastric Bypass Surgery Despite Optimization of Vitamin D Status. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2015;30(8):1377-85.
205. Ong GS, Walsh JP, Stuckey BG, Brown SJ, Rossi E, Ng JL, et al. The importance of measuring ionized calcium in characterizing calcium status and diagnosing primary hyperparathyroidism. *The Journal of clinical endocrinology and metabolism*. 2012;97(9):3138-45.
206. Gunnarsson O, Indridason OS, Franzson L, Sigurdsson G. Factors associated with elevated or blunted PTH response in vitamin D insufficient adults. *Journal of internal medicine*. 2009;265(4):488-95.
207. Melo TL, Froeder L, Baia LDC, Heilberg IP. Bone turnover after bariatric surgery. *Arch Endocrinol Metab*. 2017;61(4):332-6.
208. Lindeman KG, Greenblatt LB, Rourke C, Boussein ML, Finkelstein JS, Yu EW. Longitudinal 5-Year Evaluation of Bone Density and Microarchitecture After Roux-en-Y Gastric Bypass Surgery. *The Journal of clinical endocrinology and metabolism*. 2018;103(11):4104-12.
209. Hansen S, Jørgensen NR, Hermann AP, Støving RK. Continuous decline in bone mineral density and deterioration of bone microarchitecture 7 years after Roux-en-Y gastric bypass surgery. *European journal of endocrinology*. 2020;182(3):303-11.
210. Lindeman KG, Rushin CC, Cheney MC, Boussein ML, Hutter MM, Yu EW. Bone Density and Trabecular Morphology at Least 10 Years After Gastric Bypass and Gastric Banding. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2020.
211. Yu EW, Boussein ML, Putman MS, Monis EL, Roy AE, Pratt JS, et al. Two-year changes in bone density after Roux-en-Y gastric bypass surgery. *The Journal of clinical endocrinology and metabolism*. 2015;100(4):1452-9.
212. Schafer AL, Kazakia GJ, Vittinghoff E, Stewart L, Rogers SJ, Kim TY, et al. Effects of Gastric Bypass Surgery on Bone Mass and Microarchitecture Occur Early and Particularly Impact Postmenopausal Women. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2018;33(6):975-86.
213. Shanbhogue VV, Støving RK, Frederiksen KH, Hanson S, Brixen K, Gram J, et al. Bone structural changes after gastric bypass surgery evaluated by HR-pQCT: a two-year longitudinal study. *European journal of endocrinology*. 2017;176(6):685-93.
214. Krez A, Agarwal S, Bucovsky M, McMahon DJ, Hu Y, Bessler M, et al. Long-term Bone Loss and Deterioration of Microarchitecture After Gastric Bypass in African American and Latina Women. *The Journal of clinical endocrinology and metabolism*. 2020.
215. Raoof M, Näslund I, Rask E, Szabo E. Bone Mineral Density, Parathyroid Hormone, and Vitamin D After Gastric Bypass Surgery: a 10-Year Longitudinal Follow-Up. *Obesity surgery*. 2020.
216. Marengo AP, Guerrero Pérez F, San Martín L, Monseny R, Casajoana A, Valera R, et al. Is Trabecular Bone Score Valuable in Bone Microstructure Assessment after Gastric Bypass in Women with Morbid Obesity? *Nutrients*. 2017;9(12).
217. Emaus N, Wilsgaard T, Ahmed LA. Impacts of body mass index, physical activity, and smoking on femoral bone loss: the Tromsø study. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2014;29(9):2080-9.
218. Saad R, Habli D, El Sabbagh R, Chakhtoura M. Bone Health Following Bariatric Surgery: An Update. *Journal of clinical densitometry : the official journal of the International Society for Clinical Densitometry*. 2020;23(2):165-81.

219. Paccou J, Caiazzo R, Lespessailles E, Cortet B. Bariatric Surgery and Osteoporosis. *Calcified tissue international*. 2021.
220. Hofsø D, Hillestad TOW, Halvorsen E, Fatima F, Johnson LK, Lindberg M, et al. Bone Mineral Density and Turnover After Sleeve Gastrectomy and Gastric Bypass: A Randomized Controlled Trial (Oseberg). *The Journal of clinical endocrinology and metabolism*. 2021;106(2):501-11.
221. Hofsø D, Bollerslev J, Sandbu R, Jørgensen A, Godang K, Hjelvesæth J, et al. Bone resorption following weight loss surgery is associated with treatment procedure and changes in secreted Wnt antagonists. *Endocrine*. 2016;53(1):313-21.
222. Luger M, Kruschitz R, Winzer E, Schindler K, Grabovac I, Kainberger F, et al. Changes in Bone Mineral Density Following Weight Loss Induced by One-Anastomosis Gastric Bypass in Patients with Vitamin D Supplementation. *Obesity surgery*. 2018;28(11):3454-65.
223. Beavers KM, Greene KA, Yu EW. MANAGEMENT OF ENDOCRINE DISEASE: Bone complications of bariatric surgery: updates on sleeve gastrectomy, fractures, and interventions. *European journal of endocrinology*. 2020;183(5):R119-r32.
224. Rousseau C, Jean S, Gamache P, Lebel S, Mac-Way F, Biertho L, et al. Change in fracture risk and fracture pattern after bariatric surgery: nested case-control study. *BMJ (Clinical research ed)*. 2016;354:i3794.
225. Paccou J, Martignène N, Lespessailles E, Babykina E, Pattou F, Cortet B, et al. Gastric Bypass But Not Sleeve Gastrectomy Increases Risk of Major Osteoporotic Fracture: French Population-Based Cohort Study. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2020;35(8):1415-23.
226. Narayanan N, Palui R, Merugu C, Kar SS, Kamalanathan S, Sahoo J, et al. The Risk of Fractures in Primary Hyperparathyroidism: A Meta-Analysis. *JBMR plus*. n/a(n/a):e10482.
227. Ejlsmark-Svensson H, Rolighed L, Harsløf T, Rejnmark L. Risk of fractures in primary hyperparathyroidism: a systematic review and meta-analysis. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2021.
228. Murai IH, Roschel H, Dantas WS, Gil S, Merege-Filho C, de Cleve R, et al. Exercise Mitigates Bone Loss in Women With Severe Obesity After Roux-en-Y Gastric Bypass: A Randomized Controlled Trial. *The Journal of clinical endocrinology and metabolism*. 2019;104(10):4639-50.
229. Diniz-Sousa F, Veras L, Boppre G, Sa-Couto P, Devezas V, Santos-Sousa H, et al. The Effect of an Exercise Intervention Program on Bone Health After Bariatric Surgery: A Randomized Controlled Trial. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2020.
230. Lara-Castillo N, Johnson ML. Bone-Muscle Mutual Interactions. *Current osteoporosis reports*. 2020;18(4):408-21.
231. Yarizadeh H, Asadi S, Baharloo H, Setayesh L, Kakavandi NR, Hambly C, et al. Beneficial impact of exercise on bone mass in individuals under calorie restriction: a systematic review and Meta-analysis of randomized clinical trials. *Crit Rev Food Sci Nutr*. 2021;61(4):553-65.
232. Soltani S, Hunter GR, Kazemi A, Shab-Bidar S. The effects of weight loss approaches on bone mineral density in adults: a systematic review and meta-analysis of randomized controlled trials. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2016;27(9):2655-71.
233. Romeijn MM, Holthuijsen DDB, Kolen AM, Janssen L, Schep G, van Dielen FMH, et al. The effect of additional protein on lean body mass preservation in post-bariatric surgery patients: a systematic review. *Nutr J*. 2021;20(1):27.
234. Sukumar D, Ambia-Sobhan H, Zurfluh R, Schluskel Y, Stahl TJ, Gordon CL, et al. Areal and volumetric bone mineral density and geometry at two levels of protein intake during caloric restriction: a randomized, controlled trial. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2011;26(6):1339-48.

235. Schollenberger AE, Karschin J, Meile T, Küper MA, Königsrainer A, Bischoff SC. Impact of protein supplementation after bariatric surgery: A randomized controlled double-blind pilot study. *Nutrition*. 2016;32(2):186-92.
236. Sandvik J, Laurenus A, Naslund I, Videhult P, Wiren M, Aasheim ET. [Nordic guidelines for follow-up after obesity surgery]. *Tidsskrift for den Norske lægeforening : tidsskrift for praktisk medicin, ny raekke*. 2018;138(4).
237. Rosen CJ, Taylor CL. Common misconceptions about vitamin D--implications for clinicians. *Nat Rev Endocrinol*. 2013;9(7):434-8.
238. Durup D, Jørgensen HL, Christensen J, Tjønneland A, Olsen A, Halkjær J, et al. A Reverse J-Shaped Association Between Serum 25-Hydroxyvitamin D and Cardiovascular Disease Mortality: The CopD Study. *The Journal of clinical endocrinology and metabolism*. 2015;100(6):2339-46.
239. Michaëlsson K, Baron JA, Snellman G, Gedeberg R, Byberg L, Sundström J, et al. Plasma vitamin D and mortality in older men: a community-based prospective cohort study. *The American journal of clinical nutrition*. 2010;92(4):841-8.
240. Charoenngam N, Shirvani A, Holick MF. Vitamin D for skeletal and non-skeletal health: What we should know. *J Clin Orthop Trauma*. 2019;10(6):1082-93.
241. Durazo-Arvizu RA, Dawson-Hughes B, Kramer H, Cao G, Merkel J, Coates PM, et al. The Reverse J-Shaped Association Between Serum Total 25-Hydroxyvitamin D Concentration and All-Cause Mortality: The Impact of Assay Standardization. *Am J Epidemiol*. 2017;185(8):720-6.
242. Fan X, Wang J, Song M, Giovannucci EL, Ma H, Jin G, et al. Vitamin D Status and Risk of All-Cause and Cause-Specific Mortality in a Large Cohort: Results From the UK Biobank. *The Journal of clinical endocrinology and metabolism*. 2020;105(10).
243. Zhang Y, Fang F, Tang J, Jia L, Feng Y, Xu P, et al. Association between vitamin D supplementation and mortality: systematic review and meta-analysis. *BMJ (Clinical research ed)*. 2019;366:l4673.
244. Yao P, Bennett D, Mafham M, Lin X, Chen Z, Armitage J, et al. Vitamin D and Calcium for the Prevention of Fracture: A Systematic Review and Meta-analysis. *JAMA Netw Open*. 2019;2(12):e1917789.
245. Bolland MJ, Grey A, Avenell A. Effects of vitamin D supplementation on musculoskeletal health: a systematic review, meta-analysis, and trial sequential analysis. *Lancet Diabetes Endocrinol*. 2018;6(11):847-58.
246. Reid IR, Bolland MJ, Grey A. Effects of vitamin D supplements on bone mineral density: a systematic review and meta-analysis. *Lancet (London, England)*. 2014;383(9912):146-55.
247. Burt LA, Billington EO, Rose MS, Raymond DA, Hanley DA, Boyd SK. Effect of High-Dose Vitamin D Supplementation on Volumetric Bone Density and Bone Strength: A Randomized Clinical Trial. *Jama*. 2019;322(8):736-45.
248. Sanders KM, Stuart AL, Williamson EJ, Simpson JA, Kotowicz MA, Young D, et al. Annual high-dose oral vitamin D and falls and fractures in older women: a randomized controlled trial. *Jama*. 2010;303(18):1815-22.
249. Smith H, Anderson F, Raphael H, Maslin P, Crozier S, Cooper C. Effect of annual intramuscular vitamin D on fracture risk in elderly men and women--a population-based, randomized, double-blind, placebo-controlled trial. *Rheumatology (Oxford)*. 2007;46(12):1852-7.
250. Bischoff-Ferrari HA, Dawson-Hughes B, Orav EJ, Staehelin HB, Meyer OW, Theiler R, et al. Monthly High-Dose Vitamin D Treatment for the Prevention of Functional Decline: A Randomized Clinical Trial. *JAMA Intern Med*. 2016;176(2):175-83.
251. Smith LM, Gallagher JC, Suiter C. Medium doses of daily vitamin D decrease falls and higher doses of daily vitamin D3 increase falls: A randomized clinical trial. *The Journal of steroid biochemistry and molecular biology*. 2017;173:317-22.
252. Tondapu P, Provost D, Adams-Huet B, Sims T, Chang C, Sakhaee K. Comparison of the absorption of calcium carbonate and calcium citrate after Roux-en-Y gastric bypass. *Obesity surgery*. 2009;19(9):1256-61.

253. Baretta GA, Cambi MP, Rodrigues AL, Mendes SA. SECONDARY HYPERPARATHYROIDISM AFTER BARIATRIC SURGERY: TREATMENT IS WITH CALCIUM CARBONATE OR CALCIUM CITRATE? *Arquivos brasileiros de cirurgia digestiva : ABCD = Brazilian archives of digestive surgery*. 2015;28 Suppl 1(Suppl 1):43-5.
254. Sakhaee K, Bhuket T, Adams-Huet B, Rao DS. Meta-analysis of calcium bioavailability: a comparison of calcium citrate with calcium carbonate. *Am J Ther*. 1999;6(6):313-21.
255. Ring Madsen L, Espersen R, Rejnmark L, Richelsen B. Effect of calcium citrate vs calcium carbonate on elevated parathyroid hormone after Roux-en-Y gastric bypass. A double-blinded, randomized trial. *Clinical endocrinology*. 2018;89(6):734-41.
256. Smelt HJ, Pouwels S, Smulders JF. The Clinical Dilemma of Calcium Supplementation After Bariatric Surgery: Calcium Citrate or Calcium Carbonate That Is the Question? *Obesity surgery*. 2016;26(11):2781-2.
257. Sakhaee K, Pak C. Superior calcium bioavailability of effervescent potassium calcium citrate over tablet formulation of calcium citrate after Roux-en-Y gastric bypass. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*. 2013;9(5):743-8.
258. Berarducci A, Haines K, Murr MM. Incidence of bone loss, falls, and fractures after Roux-en-Y gastric bypass for morbid obesity. *Appl Nurs Res*. 2009;22(1):35-41.
259. Maghrabi AH, Wolski K, Abood B, Licata A, Pothier C, Bhatt DL, et al. Two-year outcomes on bone density and fracture incidence in patients with T2DM randomized to bariatric surgery versus intensive medical therapy. *Obesity*. 2015;23(12):2344-8.
260. Lu CW, Chang YK, Chang HH, Kuo CS, Huang CT, Hsu CC, et al. Fracture Risk After Bariatric Surgery: A 12-Year Nationwide Cohort Study. *Medicine*. 2015;94(48):e2087.
261. Douglas IJ, Bhaskaran K, Batterham RL, Smeeth L. Bariatric Surgery in the United Kingdom: A Cohort Study of Weight Loss and Clinical Outcomes in Routine Clinical Care. *PLoS Med*. 2015;12(12):e1001925.
262. Yu EW, Lee MP, Landon JE, Lindeman KG, Kim SC. Fracture Risk After Bariatric Surgery: Roux-en-Y Gastric Bypass Versus Adjustable Gastric Banding. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2017;32(6):1229-36.
263. Javanainen M, Pekkarinen T, Mustonen H, Scheinin T, Leivonen M. Two-Year Nutrition Data in Terms of Vitamin D, Vitamin B12, and Albumin After Bariatric Surgery and Long-term Fracture Data Compared with Conservatively Treated Obese Patients: a Retrospective Cohort Study. *Obesity surgery*. 2018;28(9):2968-75.
264. Axelsson KF, Werling M, Eliasson B, Szabo E, Näslund I, Wedel H, et al. Fracture Risk After Gastric Bypass Surgery: A Retrospective Cohort Study. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2018;33(12):2122-31.
265. Fashandi AZ, Mehaffey JH, Hawkins RB, Schirmer B, Hallowell PT. Bariatric surgery increases risk of bone fracture. *Surg Endosc*. 2018;32(6):2650-5.
266. Yu EW, Kim SC, Sturgeon DJ, Lindeman KG, Weissman JS. Fracture Risk After Roux-en-Y Gastric Bypass vs Adjustable Gastric Banding Among Medicare Beneficiaries. *JAMA surgery*. 2019;154(8):746-53.
267. Blom-Høgestøl IK, Hewitt S, Chahal-Kummen M, Brunborg C, Gulseth HL, Kristinsson JA, et al. Bone metabolism, bone mineral density and low-energy fractures 10 years after Roux-en-Y gastric bypass. *Bone*. 2019;127:436-45.
268. Ahlin S, Peltonen M, Sjöholm K, Anveden Å, Jacobson P, Andersson-Assarsson JC, et al. Fracture risk after three bariatric surgery procedures in Swedish obese subjects: up to 26 years follow-up of a controlled intervention study. *Journal of internal medicine*. 2020;287(5):546-57.

Supplementary materials

Supplementary Table 1. Overview of major fracture studies after obesity surgery.

Author, year	Study	Procedure	Follow-up (y, mean)	N	Mean time to fracture	Adj RR (all)
Berarducci 2009 (258)	Cohort	RYGB	2.4 y	167		
Lalmohamed UK 2012 (153)	Cohort Matched ctrl	All Matched ctrl AGB 60%/RYGB 29%	2.2 y	2079 10,442		
Nakamura USA 2013 (154)	Cohort 1985-2004	Bariatric vs community RYGB	8.9 y	258 243 (94%)	13 y median	2.3
Maghrabi USA 2015 (259)	RCT DM pts	Bariatric lifestyle RYGB/sleeve	2 y	37 17 18/19		
Lu Taiwan, 2015 (260)	Cohort matched obese	All bariatric Ctrl - matched obese Malabsorptiv/restrictiv	4.8 y	2064 5027 289/1775		1.21 Ref 1.47 / 1.17
Douglas UK 2015 (261)	cohort	Bariatric surg vs obese ctrl ABG/sleeve (63%) RYGB (37%)	3.0 y median	3447 3535 2442 1421		1.25 (unadj.) (ns)
Rosseau Canada 2016 (224)	Cohort matched ctrl	All bariatric Obese (ctrl) Non-obese (ctrl) RYGB / BPD	4.4 y	12,676 38,028 126,760	3.9 y	1.44 1.04 (ns) Ref 1.13 (ns)/1.60
Yu, USA 2017 (262)	Cohort	RYGB vs AGB matched	2.3 y	7516 7516		1.43
Javanainen, Finland 2018 (263)	Cohort	RYGB Sleeve Control, obese	6.8 y 6.9 y 13.3 y	253 142 199	4.5 y 3.9 y 6.2 y	5.49
Axelsson Sweden 2018 (264)	Cohort vs matched ctrl	RYGB (DM+/-)	3.2 y 3.1 y	7758/31,213 7758/31,213		1.26/1.32
Fashandi USA 2018 (265)	RYGB Obes match	RYGB All surg Obese matched	7.6 y	2729 3439 3880	7.6y vs GS All 8.2y vs ctrl	OR 2.2 (adj) vs SG HR 2.4 (unadj.)
Yu, USA 2019 (266)	Cohort	RYGB vs AGB	3.3 y 3.9 y	29,624 12,721		1.73
Blom-Høgestøl, Norway 2019 (267)	Cohort	RYGB	10 y	122	8.4 y	
Ahlin Sweden 2020 (268)	Matched ctrl	RYGB Banding VBG control	15.1-17.9y (median)	266 376 1365 2040		2.6 1.3 (ns) 1.2
Paccou France 2020 (225)	Matched ctrl	Banding sleeve VBG RYGB All surg Ctrl, obese	6.8 y 5.8 y 7.6 y 6.2 y 6.2 y 5.3 y	5178 18635 2647 14532 40992 40992		0.95 (ns) 0.95 (ns) 0.95 (ns) 1.70 1.22

RYGB, Roux-en-Y gastric bypass; AGB, adjustable gastric banding; BPD, biliopancreatic diversion (not specified duodenal switch); VBG, vertical banded gastroplasty (restrictive); SG, sleeve gastrectomy; RR, relative ratio; OR, odds ratio; HR, hazard ratio.

Supplementary Table 2. Overview of randomized vitamin D intervention studies after Roux-en-Y gastric bypass (RYGB)

Study	Carlin, 2009	Goldner, 2009*	Muschitz, 2016	Smelt, 2019
Procedure	RYGB	RYGB	RYGB/sleeve	RYGB/sleeve
Number (Interv, Ctrl)	30 + 30	45	110 + 110	50 + 50
<i>Baseline</i>				
S-25(OH)D, nmol/l	49	38 or 57	44	47
PTH, pmol/l	7.1	10.6 or 7.1	7.2	6.8
SHPT (%)	n.a.	46%	n.a	
<i>Intervention</i>				
Duration	1y	1y*	2y	6m
Loading dose			28,000 IU/w x8	
Maintenance dose	50,000 IU/w		16,000 IU/w	50,000 IU/m
Maintenance dose/d	7143 IU	2000 IU or 5000 IU	2,286 IU	1667 IU/d
Daily dose	800 IU (all)	800 IU (ctrl only)	200 IU (all)	800 IU (all)
<i>Postop</i>				
25(OH)D, nmol/l	95 ^a	60 or 66 ^a	80 ^a	94 ^b
Vit D def (<50 nmol/l)	14%	n.a.	n.a.	20%
Vit D suff (≥75 nmol/l)	71%	78% and 70%	n.a	n.a.
PTH, pmol/l	7,1	7.4 or 4.6 ^a	6.0 ^a	5.2 ^b
SHPT (%)	40%.	n.a	n.a.	n.a
Compliance		63% and 83%		

RYGB, gastric bypass; sleeve, sleeve gastrectomy; S-25(OH)D, serum 25-hydroxyvitamin D; PTH, parathyroid hormone; SHPT, secondary hyperparathyroidism; * planned for 2 years, analysed at 1 year; ^a difference between groups; ^b difference with baseline.

Papers included – Study 1-4

Secondary Hyperparathyroidism, Vitamin D Sufficiency, and Serum Calcium 5 Years After Gastric Bypass and Duodenal Switch

Stephen Hewitt · Torgeir T. Søvik · Erlend T. Aasheim ·
Jon Kristinsson · Jørgen Jahnsen · Grethe S. Birketvedt ·
Thomas Bøhmer · Erik F. Eriksen · Tom Mala

Published online: 28 September 2012
© Springer Science+Business Media, LLC 2012

Abstract

Background The prevalence of secondary hyperparathyroidism (SHPT) is high after bariatric surgery. Vitamin D is supplied to counteract SHPT and bone disease, and we studied vitamin D associations with SHPT.

Methods We measured serum levels of 25-OH vitamin D and parathyroid hormone (PTH) 5 years after gastric bypass and duodenal switch. One hundred twenty-five patients were included, of whom 114 (91 %) had undergone gastric bypass and 11 (9 %) had undergone duodenal switch. SHPT was defined as PTH > 7.0 pmol/l in the absence of hypercalcemia. 25-OH vitamin D levels were divided into three categories: <50, 50–74, and ≥75 nmol/l. Serum ionized

calcium, magnesium, phosphate, and creatinine were divided into tertiles.

Results Mean age ± SD was 44 ± 9 years at 5 years follow-up. Ninety out of 125 (72 %) patients were women. SHPT was present in 45 out of 114 (40 %) gastric bypass patients and in 11 out of 11 (100 %) duodenal switch patients. The prevalence was high in all vitamin D categories studied. An inverse association between ionized calcium and PTH was found. For the gastric bypass patients, the odds ratio for SHPT in the upper two tertiles of ionized calcium was 0.35; 95 % CI, 0.15–0.79; $p=0.011$, compared with the lowest tertile. Supplements of vitamin D and calcium were not associated with a lower prevalence of SHPT at 5 years follow-up.

Conclusions The prevalence of SHPT was high 5 years after gastric bypass and duodenal switch. SHPT was inversely associated with serum ionized calcium, but not with vitamin D. The supplementation used was insufficient to compensate for the impaired calcium absorption after surgery.

Keywords Secondary hyperparathyroidism · Parathyroid hormone · Vitamin D · Calcium · Hypocalcemia · Bariatric surgery · Gastric bypass · Duodenal switch

S. Hewitt (✉) · J. Kristinsson · G. S. Birketvedt · T. Mala
Department of Morbid Obesity and Bariatric Surgery,
Department of Medicine,
Oslo University Hospital Aker,
P.O. Box 4959, Nydalen,
0424 Oslo, Norway
e-mail: stehewi@online.no

T. T. Søvik · J. Kristinsson · T. Mala
Department of Gastrointestinal Surgery, Department of Surgery,
Oslo University Hospital Aker,
Oslo, Norway

T. T. Søvik · E. F. Eriksen
Faculty of Medicine, University of Oslo,
Oslo, Norway

E. T. Aasheim · T. Bøhmer · E. F. Eriksen
Department of Endocrinology, Department of Medicine,
Oslo University Hospital Aker,
Oslo, Norway

J. Jahnsen
Department of Gastroenterology, Department of Medicine,
Oslo University Hospital Aker,
Oslo, Norway

Background

Bariatric surgery provides effective and sustainable weight loss in morbidly obese patients [1]. Basically, the concept of bariatric surgery consists of a restrictive component and a malabsorptive component, reducing the intake and absorption of food. Due to this and the fact that obesity itself is often accompanied by nutritional deficiencies, bariatric patients are followed up especially with regards to deficiencies in iron, folic acid, calcium, and vitamins after surgery [2–8].

Several studies have shown a high prevalence of secondary hyperparathyroidism (SHPT) after gastric bypass and duodenal switch [8–20]. Most of these have been performed within 2–3 years after surgery. A few studies evaluating parathyroid hormone (PTH) and SHPT have included patients with longer follow-up and suggest that PTH elevations and SHPT continue and may actually worsen over time [7, 15–18].

Vitamin D sufficiency is a highly discussed topic after bariatric surgery. An inverse relationship has been described between vitamin D and PTH or hyperparathyroidism [9, 12, 17, 21]. In most studies, serum levels of 25-OH vitamin D at 50 nmol/l have been considered adequate, mainly with regards to bone health [22]. Other reports have found 25-OH vitamin D levels at 75 nmol/l or above more beneficial [16, 21, 23]. There has also been a general impression that increased vitamin D supplementation is needed to achieve sufficient vitamin D levels after bariatric surgery [24–27]. However, only one intervention study with vitamin D has reported the reversal of SHPT [26].

To our knowledge, this is the first standardized 5-year follow-up study after gastric bypass and duodenal switch studying SHPT. In this study, our main objective was to test whether levels of 25-OH vitamin D at 75 nmol/l or above were associated with a lower prevalence of SHPT, than 25-OH vitamin D levels below, 5 years after gastric bypass and duodenal switch. We also aimed to study the associations of other factors thought to be involved in SHPT, including the effect of supplementation on SHPT.

Materials and Methods

This study was a cross-sectional 5-year follow-up study of patients who had undergone gastric bypass or biliopancreatic diversion with duodenal switch at the Department of Morbid Obesity and Bariatric Surgery, Oslo University Hospital Aker from September 2004 and throughout June 2006. Blood samples, anthropometric data, and clinical information were retrieved prospectively, and the study was formulated in the early phase of the 5-year follow-up. A total of 133 out of 171 patients (78 %) participated in the scheduled 5-year follow-up. Of these, eight patients were excluded: one patient with distal gastric bypass; six (four gastric bypass and two duodenal switch) due to at least one missing value of ionized calcium, 25-OH vitamin D, or PTH in serum; and one gastric bypass patient due to ionized calcium above the upper normal reference range (1.18–1.35 mmol/l) to avoid primary hyperparathyroidism, leaving 125 patients available for analyses, of whom 114 (91 %) had undergone standard Roux-en-Y gastric bypass and 11 (9 %) had undergone duodenal switch.

Eligibility criteria for surgery were a body mass index (BMI) ≥ 40 kg/m² or a BMI ≥ 35 kg/m² in patients with obesity-related comorbidities [28]. Gastric bypass was performed by constructing a gastric pouch of about 25 ml, an alimentary limb length of 150 cm, and a biliopancreatic limb of 50 cm. In selected super obese patients (BMI ≥ 50 kg/m²), duodenal switch was applied: a sleeve gastrectomy was performed along a nasogastric tube of 30–32 F and an alimentary limb of 200 cm and a common channel of 100 cm were constructed. All procedures were performed laparoscopically and have been described previously [29].

At the preoperative consultation and at the 5-year follow-up, body weight was measured with a platform weight, Seca 635 Class III (0–300 kg) and height was measured with a wall-fixed adjustable steel measure. Blood samples were drawn after an overnight fast. Serum was analyzed for 25-OH vitamin D by radioimmunoassay (DiaSorin, Stillwater, MN, USA), intact PTH by chemiluminoimmunoassay (Immulite 2500, Siemens Healthcare Diagnostics, Los Angeles, CA, USA), and ionized calcium with a Rapidlab 348 analyzer (Instrumed Inc., Atlanta, GA, USA) at the Hormone Laboratory of Oslo University Hospital Aker. Analysis of magnesium, phosphate, and creatinine were performed at the Central Laboratory of Oslo University Hospital Aker with a Modular (Roche®) analyzer and with Roche® reagents. Levels of magnesium and phosphate were measured photometrically. Creatinine levels were determined by an enzymatic method. Levels of 25-OH vitamin D were divided into three categories: <50 , 50–74, and ≥ 75 nmol/l. SHPT was defined as serum PTH above the upper limit of the normal reference range (>7.0 pmol/l) in the absence of elevated ionized calcium (i.e., >1.35 mmol/l). Tertiles of serum ionized calcium, phosphate, magnesium, and creatinine were identified. The cutoff for the lowest ionized calcium tertile was found to be <1.22 mmol/l.

The patients had follow-up visits with the multidisciplinary team 6 weeks, 6 months, 1 year, and 2 years after surgery. They were also encouraged to have regular consultations by their general physician. All 5-year follow-up visits were performed by a physician. Patients who did not show up for the scheduled 5-year follow-up were called at least twice more. Blood samples, which included serum PTH, 25-OH vitamin D, and ionized calcium, were drawn at every visit, and supplementation of vitamins and minerals was adjusted according to laboratory results.

Standard supplementation of vitamin D included one multivitamin tablet (containing 200 IU cholecalciferol) daily on a permanent basis together with 1,000 mg calcium carbonate combined with 800 IU cholecalciferol daily. The latter was recommended permanently after

duodenal switch and for at least 1 year after gastric bypass, and later individualized. At the 5-year follow-up, the patients were asked standardized questions about the intake of supplements of vitamin D and calcium and the number of days per week the supplements were taken. Patients taking supplements at least 5 days a week were registered as compliant. The doses reported for vitamin D included the 200 IU in the multivitamin tablet.

The patient data were registered in a database approved and licensed by The Norwegian Data Inspectorate. All patients gave informed written consent to participate.

Statistical analyses were performed with SPSS version 15. Student's *t* test was applied for comparisons of continuous variables. If normal distribution was not fulfilled, logarithmic transformation was performed. Pearson's correlation test was used for continuous variables. Proportions were compared with chi-square test. Logistic regression analyses were used to estimate the odds ratio (OR) between categorical values. Gastric bypass and duodenal switch are considered separately due to malabsorptive differences between the two procedures.

Results

Demographic characteristics, laboratory findings, and the use of supplements at 5 years follow-up are shown in Tables 1 and 2. SHPT was prevalent in 45 (40 %) of the gastric bypass patients and in all 11 (100 %) duodenal switch patients.

In SHPT patients, 80 % of the gastric bypass patients and 64 % of the duodenal switch patients had 25-OH vitamin D ≥ 50 nmol/l and 36 % of the gastric bypass patients and 27 % of the duodenal switch patients had 25-OH vitamin D ≥ 75 nmol/l (Table 3).

There was no association between levels of 25-OH vitamin D and prevalence of SHPT in gastric bypass patients (Table 4).

Table 1 Demographic characteristics at 5-year follow-up

	Gastric bypass (<i>n</i> =114)	Duodenal switch (<i>n</i> =11)	<i>p</i> value
Age (years)	45 (9)	44 (9)	0.84
Female, <i>n</i> (%)	83 (73)	7 (64)	0.53
Weight (kg)	100 (24)	103 (19)	0.69
Height (cm)	171.3 (10.0)	171.6 (7.8)	0.50
BMI (kg/m ²)	33.8 (6.1)	34.9 (5.8)	0.58
Weight loss (kg)	37.3 (15.5)	67.2 (13.4)	<0.001
Weight loss (%)	27.3 (10.6)	39.7 (7.5)	<0.001

Continuous variables are shown as mean (SD)

BMI body mass index

Table 2 Laboratory findings and use of supplements at 5-year follow-up

	Gastric bypass (<i>n</i> =114)	Duodenal switch (<i>n</i> =11)	<i>p</i> value
PTH>7.0 pmol/l, <i>n</i> (%)	45 (40)	11 (100)	<0.001
PTH (pmol/l)	7.1 (4.1)	15.5 (4.9)	<0.001
25(OH)D (nmol/l)	65 (22)	62 (33)	0.67
Ionized Ca (mmol/l)	1.24 (0.04)	1.19 (0.05)	<0.001
Phosphate (mmol/l)	1.06 (0.17)	1.09 (0.11)	0.55
Magnesium (mmol/l)	0.82 (0.07)	0.82 (0.06)	0.95
Creatinine (μmol/l)	64 (13)	60 (19)	0.38
Vit D supplementation, <i>n</i> (%) ^a	46 (40)	11 (100)	<0.001
Ca supplementation, <i>n</i> (%) ^b	47 (41)	9 (82)	0.010

Continuous variables are shown as mean (SD)

PTH parathyroid hormone, PTH>7.0 pmol/l secondary hyperparathyroidism, 25(OH)D 25-OH vitamin D

^a Daily use of vitamin D supplementation of at least 400 IU

^b Daily use of calcium supplementation (any dose)

However, the OR for SHPT was lower in the upper two tertiles of serum ionized calcium compared with the lowest tertile in gastric bypass patients (OR, 0.35; 95 % CI, 0.15–0.79; *p*=0.011). Further subanalyses showed the same tendency in all the three vitamin D categories studied. Ionized calcium was within the normal reference range in 106 out of 114 (93 %) of the gastric bypass patients. However, there was a left-shifted calcium distribution with 90 out of 114 (79 %) of the population having calcium levels below the middle of the reference range (1.27 mmol/l). A suspected association of SHPT by phosphate tertiles (OR, 0.48; 95 % CI, 0.22–1.04; *p*=0.064) was attenuated when calcium was added in a multivariate model (data not shown). Assessing a possible role of sunlight exposure, we also included month of follow-up in a multivariate model, but this did not affect the associations of SHPT with vitamin D or calcium. The left shift in calcium distribution was even more pronounced in the duodenal switch patients as 10 out of 11 (91 %) had levels below the middle of the reference range.

Furthermore, there was an inverse correlation between ionized calcium and PTH levels in gastric bypass patients (*r*=−0.28; *p*=0.003) (Fig. 1a); a corresponding, albeit non-significant, correlation was also found for the duodenal switch patients (*r*=−0.55; *p*=0.083) (Fig. 1b). Significant correlations between 25-OH vitamin D and PTH were not demonstrable for neither gastric bypass patients nor duodenal switch patients.

Supplementation of vitamin D and calcium did not show any association with the prevalence of SHPT after gastric bypass (data not shown). Similarly, serum PTH did not differ by vitamin D doses in gastric bypass patients, while

Table 3 Vitamin D status according to the presence of SHPT at 5-year follow-up

	Total (n)	25-OH vitamin D					
		<50 nmol/l		50–74 nmol/l		≥75 nmol/l	
		Number	Percent	Number	Percent	Number	Percent
Gastric bypass (n=114)							
PTH>7 pmol/l	45	9	20	20	44	16	36
PTH≤7 pmol/l	69	18	26	27	39	24	35
Duodenal switch (n=11)							
PTH>7 pmol/l	11	4	36	4	36	3	27
PTH≤7 pmol/l	0						

PTH parathyroid hormone,
PTH>7.0 pmol/l secondary hyperparathyroidism

PTH was somewhat lower in those of the duodenal switch patients who used the largest doses (Fig. 2).

Discussion

This study revealed a high prevalence of SHPT 5 years after gastric bypass, and this was more pronounced after duodenal switch. Prevalence of SHPT was not associated with different levels of serum 25-OH vitamin D. In the gastric bypass patients, there was a tendency of a left-shifted calcium distribution in this population, and we found SHPT to be significantly lower in the upper two tertiles of ionized calcium. Serum ionized calcium was even lower in the duodenal switch patients, who all had SHPT. Therefore, a left shift in the distribution of serum ionized calcium may at least

partly explain the high prevalence of SHPT after these two bariatric procedures.

Findings in Relation to Other Studies

The prevalence of SHPT is generally increased after bariatric surgery. Left uncorrected, it will promote bone loss and increase the risk of osteopenia and osteoporosis [4]. Most of the studies of PTH and SHPT have short-term follow-up, although

Table 4 OR for SHPT 5 years after gastric bypass (n=114)

	Number	OR	95 % CI	p value
25(OH)D≥50 nmol/l ^a	114	1.41	0.57–3.50	0.46
25(OH)D≥75 nmol/l ^a	114	1.24	0.55–2.77	0.61
Ionized Ca ^b	114	0.35	0.15–0.79	0.011
25(OH)D<50 nmol/l ^c	27	0.51	0.10–2.57	0.41
25(OH)D 50–74 nmol/l ^c	47	0.10	0.02–0.45	0.003
25(OH)D≥75 nmol/l ^c	40	0.79	0.18–3.54	0.79
Phosphate ^b	114	0.48	0.22–1.04	0.064
Magnesium ^b	114	1.43	0.60–3.43	0.42
Creatinine ^b	114	0.57	0.26–1.26	0.16

CI confidence interval, 25(OH)D 25-OH vitamin D

^a OR for SHPT were calculated for 25-OH vitamin D comparing levels of ≥50 and ≥75 nmol/l, respectively, with levels below

^b For other variables, OR were calculated comparing the upper two tertiles with the lowest tertile

^c OR for SHPT in the upper two calcium tertiles compared with the lowest tertile in different categories of 25-OH vitamin D

Fig. 1 Correlation of PTH and ionized calcium 5 years after **a** gastric bypass (n=114) and **b** duodenal switch (n=11)

Fig. 2 PTH by daily doses of vitamin D (in international units per day) after gastric bypass and duodenal switch

a few have included patients up to 20 years after surgery [8–21]. After gastric bypass, the prevalence of SHPT is typically 30–50 %, and after biliopancreatic diversion, 50–80 %. In this study, the prevalence of SHPT after gastric bypass was consistent with other studies and high after duodenal switch.

Recent literature has focused on vitamin D in the evaluation of SHPT after bariatric surgery. Some of the studies have identified an inverse association of PTH and vitamin D [9, 12, 17, 21], while others have failed to show this. In our study, the majority of the patients had adequate levels of vitamin D. Levels of 25-OH vitamin D below 50 nmol/l were not associated with higher prevalence of SHPT. Neither did we find a lower prevalence of SHPT with 25-OH vitamin D ≥ 75 nmol/l, which is increasingly referred to as vitamin D sufficiency [23].

There are no conclusive data regarding the association of altered calcium and vitamin D homeostasis after gastric bypass, but calcium deficiency and bone disease can occur [4, 30]. As in this study, calcium is usually found normal in serum [4]. To our knowledge, this is the first study to show a left-shifted calcium distribution in the gastric bypass population and the first study to show that serum ionized calcium levels in the lower normal range were associated with elevated PTH in the gastric bypass patients. This observation also seemed consistent across different levels of vitamin D. There is agreement of an increased risk for calcium deficiency with SHPT after malabsorptive procedures, such as biliopancreatic diversion with duodenal switch [4, 30].

Accordingly, SHPT primarily seems associated with low ionized calcium and less so to vitamin D 5 years after gastric bypass. This may also apply to the duodenal switch patients, but they were few in this study.

Causes of SHPT

Anatomic changes after gastric bypass and duodenal switch can cause calcium and vitamin D deficiencies, with duodenal switch being the most malabsorptive procedure [4]. As calcium is the main stimulator of PTH secretion and in light

of our findings, we suspect that too little attention has been directed to a low normal serum calcium.

To our knowledge, other factors such as magnesium, phosphate, and creatinine have not been reported to be associated with SHPT in other bariatric studies. Magnesium deficiency may also occur after bariatric surgery and may affect SHPT by reducing calcium absorption and synthesis of PTH [7, 31]. Phosphate and creatinine may also play a role in SHPT, especially in patients with serious renal insufficiency, but they are very few in bariatric populations. Little attention has been paid to a possible mechanism of tertiary hyperparathyroidism with a changed calcium-sensing set point, which also could be involved after a longstanding relative hypocalcemia after bariatric surgery [32].

Intestinal cells may adapt to environmental changes and increase their uptake of deficient minerals, implying that the need for supplements may be reduced by time [33]. However, it should be discussed whether and when supplements are to be reduced postoperatively, as the prevalence of SHPT continues to be high.

Future Challenges

There is a need for an optimization of the calcium homeostasis after bariatric surgery. In this regard, more intervention studies are required to improve the effectiveness of the supplementation regimens, aiming to prevent SHPT and bone loss in the long term.

In accordance with this study, standard vitamin D doses of 400–800 IU postoperatively seem to be insufficient to achieve vitamin D sufficiency and to suppress PTH in the bariatric population [19]. The intervention studies of Goldner et al., with daily vitamin D doses of 2,000 or 5,000 IU, and Carlin et al., with weekly doses of 50,000 IU, indicate that larger doses may be necessary in most gastric bypass patients [24, 25]. However, we are only aware of one intervention study after gastric bypass which have described a reverse in SHPT with an aggressive treatment regimen of

vitamin D and calcium citrate [26]. For duodenal switch patients, daily doses of vitamin D up to 50,000 IU two or three times a day are recommended, with limited documentation on SHPT [4, 30].

In the present study, vitamin D levels ≥ 75 nmol/l was not associated with less SHPT. However, in a subanalysis of the gastric bypass patients with vitamin D ≥ 95 nmol/l, SHPT appeared in only 3 of 14 (21 %), suggesting that 25-OH vitamin D levels of around 100 nmol/l or higher could be more beneficial in these patients.

The calcium doses implemented in guidelines are now typically 1,200–1,500 mg after gastric bypass surgery and up to 2,000 mg after duodenal switch [4, 30]. We are not aware of any studies showing less SHPT by calcium supplementation. In this study, calcium supplementation was not associated with differences in SHPT. Calcium citrate is the preferred formulation, exhibiting better intestinal absorption than calcium carbonate [34]. However, the citrate formulation is often less available and more expensive for the patients and implies a larger number of capsules a day.

Strengths and Limitations

This standardized 5-year follow-up provides important long-term data of SHPT after surgery. The study had a relatively high follow-up rate of 78 % with blood sample results for nearly all of the patients. Patients lost to follow-up after bariatric surgery have shown less favorable weight loss compared to those who participated [35]. Less weight loss is associated with noncompliance with dietary and lifestyle instructions [30]. Therefore, the prevalence of SHPT reported in this population might be somewhat underestimated.

The main limitation of this study is the cross-sectional design, making us unable to conclude about cause–effect relationships. Furthermore, values of 1,25-(OH)₂ vitamin D and urinary calcium were not available for analyses; and there were no preoperative values of PTH, vitamin D, or ionized calcium available. The use of supplements was self-reported at the consultation, and compliance could be lower than reported. The results of the duodenal switch patients should be evaluated cautiously, as there were only 11 patients available for analyses, and the comparisons between the two procedures in Tables 1 and 2 may be affected by some uncertainties and type 2 statistical errors.

Summary

Vitamin D sufficiency, with 25-OH vitamin D ≥ 75 nmol/l, was associated with the same high prevalence of SHPT 5 years after gastric bypass and duodenal switch, compared with 25-OH vitamin D levels below. More attention to calcium levels in the lower end of the normal reference range can be valuable in the understanding of SHPT. The findings suggest that the

supplementation used was unable to compensate for the impaired intestinal absorptive capacity following surgery. There is a need for further exploration of the mechanisms behind SHPT in this population, and the supplementation regimens seem to need improvements to counteract SHPT and the risk of bone disease in the long term after surgery.

Acknowledgments We are especially thankful to the whole multidisciplinary team at the Department of Morbid Obesity and Bariatric Surgery, Oslo University Hospital Aker for their participation and follow-up before and after surgery. We are also thankful to Peter Abusdal Torjesen and Solve Tjora at the Hormone Laboratory and the Central Laboratory, respectively, for reviewing the section with the methodological aspects of the serum analyses.

Conflict of Interest Torgeir T Søvik received travel support from Johnson & Johnson. Jon Kristinsson received travel support from Covidien and Johnson & Johnson. Thomas Bøhmer received royalties from Nycomed Pharma. Erik Fink Eriksen has received travel support and speaker fees from Lilly, Novartis, Amgen, and IDS. Tom Mala has received speaker fees from Johnson & Johnson and travel support from Covidien. Stephen Hewitt, Erlend T Aasheim, Jørgen Jahnsen, and Grethe S Birketvedt did not report any form of conflicts of interest.

References

1. Sjöström L, Narbro K, Sjöström CD, et al. Effects of bariatric surgery on mortality in Swedish obese subjects. *N Engl J Med*. 2007;8:741–52.
2. Tucker ON, Szomstein S, Rosenthal RJ. Nutritional consequences of weight-loss surgery. *Med Clin N Am*. 2007;91:499–514.
3. Ziegler O, Sirveaux MA, Brunaud L, et al. Medical follow-up after bariatric surgery: nutritional and drug issues. General recommendations for the prevention and treatment of nutritional deficiencies. *Diabetes Metab*. 2009;35:544–57.
4. Mechanick JI, Kushner RF, Sugerman HJ, et al. AACE/TOS/ASMBS guidelines. *Obesity*. 2009;17:S1–S70.
5. Aasheim ET, Hofso D, Hjelmæsæth J, et al. Vitamin status in morbidly obese patients: a cross-sectional study. *Am J Clin Nutr*. 2008;87:862–9.
6. Aasheim ET, Björkman S, Søvik TT, et al. Vitamin status after bariatric surgery: a randomised study of gastric bypass and duodenal switch. *Am J Clin Nutr*. 2009;90:15–22.
7. Gasteyger C, Suter M, Gaillard RC, et al. Nutritional deficiencies after Roux-en-Y gastric bypass for morbid obesity often cannot be prevented by standard multivitamin supplementation. *Am J Clin Nutr*. 2008;87:1128–33.
8. Søvik TT, Aasheim ET, Taha O, et al. Weight loss, cardiovascular risk factors, and quality of life after gastric bypass and duodenal switch. *Ann Intern Med*. 2011;155:281–91.
9. DiGiorgi M, Daud A, Inabnet WB, et al. Markers of bone and calcium metabolism following gastric bypass and laparoscopic adjustable gastric banding. *Obes Surg*. 2008;18:1144–8.
10. Signori C, Zalesin KC, Franklin B, et al. Effect of gastric bypass on vitamin D and secondary hyperparathyroidism. *Obes Surg*. 2010;20:949–52.
11. Youssef Y, Richards WO, Sekhar N, et al. Risk of secondary hyperparathyroidism after laparoscopic gastric bypass surgery in obese women. *Surg Endosc*. 2007;21:1393–6.
12. Clements RH, Yellumhanhi K, Wesley M, et al. Hyperparathyroidism and vitamin D deficiency after laparoscopic gastric bypass. *Am Surg*. 2008;74(6):469–74.

13. Ybarra J, Sanches-Hernandez J, Gich I, et al. Unchanged hypovitaminosis D and secondary hyperparathyroidism in morbid obesity after bariatric surgery. *Obes Surg*. 2005;3:330–5.
14. Vilarrasa N, San Jose P, Garcia I, et al. Evaluation of bone mineral density loss in morbidly obese women after gastric bypass: 3-year follow-up. *Obes Surg*. 2011;21:465–72.
15. Valderas JP, Soledad V, Solari S, et al. Increase of bone resorption and the parathyroid hormone in postmenopausal women in the long-term after Roux-en-Y gastric bypass. *Obes Surg*. 2009;19:1132–8.
16. Balsa JA, Botella-Carratero JI, Peromingo R, et al. Role of calcium malabsorption in the development of secondary hyperparathyroidism after biliopancreatic diversion. *J Endocrinol Invest*. 2008;31:845–50.
17. Topart P, Becouarn G, Salle A. Five-year follow-up after biliopancreatic diversion with duodenal switch. *Surg Obes Relat Dis*. 2011;7:199–205.
18. Johnson JM, Maher JW, DeMaria EJ, et al. The long-term effects of gastric bypass on vitamin D metabolism. *Ann Surg*. 2006;243:701–5.
19. Compher CW, Badellino K, Boullata JI. Vitamin D and the bariatric surgical patient: a review. *Obes Surg*. 2008;18:220–4.
20. Hamoui N, Kim K, Anthone G, et al. The significance of elevated levels of parathyroid hormone in patients with morbid obesity before and after bariatric surgery. *Arch Surg*. 2003;138:891–7.
21. Grethen E, McClintock R, Gupta CE, et al. Vitamin D and hyperparathyroidism in obesity. *J Clin Endocrinol Metab*. 2011;96:1320–6.
22. Ross AC, Manson JE, Abrams SA, et al. The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *J Clin Endocrinol Metab*. 2011;96:53–8.
23. Bischoff-Ferrari HA. Optimal serum 25-OH-hydroxyvitamin D levels for multiple health outcomes. *Adv Exp Med Biol*. 2008;624:55–71.
24. Goldner WS, Stoner JA, Lyden E, et al. Finding the optimal dose of vitamin D following Roux-en-Y gastric bypass: a prospective, randomized pilot clinical trial. *Obes Surg*. 2009;19:173–9.
25. Carlin AM, Rao DS, Yager KM, et al. Treatment of vitamin D depletion after Roux-en-Y gastric bypass: a randomized prospective clinical trial. *Surg Obes Relat Dis*. 2009;5:444–9.
26. Srikanth MS, Quddusi S, Fox SR, et al. Correction of secondary hyperparathyroidism after distal gastric bypass using oral high dose vitamin D, calcitriol and calcium replacement. *Surg Obes Relat Dis*. 2005;1:271.
27. Nelson ML, Bolduc LM, Toder ME, et al. Correction of preoperative vitamin D deficiency after Roux-en-Y gastric bypass surgery. *Surg Obes Relat Dis*. 2007;3:434–7.
28. Anonymous. Gastrointestinal surgery for severe obesity: National Institutes of Health Consensus Development Conference Statement. *Am J Clin Nutr* 1992; 55:615S–619S.
29. Søvik TT, Taha O, Aasheim ET, et al. Randomized clinical trial of laparoscopic gastric bypass versus laparoscopic duodenal switch for superobesity. *Br J Surg*. 2010;97(2):160–6.
30. Heber D, Greenway FL, Kaplan LM, et al. Endocrine and nutritional management of the post-bariatric surgery patient: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2010;95:4823–43.
31. Bøhmer T. Magnesium deficiency. *Tidsskr Nor Lægeforen*. 1986;106:548–51.
32. Parfitt AM. The hyperparathyroidism of chronic renal failure: a disorder of growth. *Kidney Int*. 1997;52:3–9.
33. McDuffie LA, Bucher BT, Erwein CR, et al. Intestinal adaptation after small bowel resection in human infants. *J Pediatr Surg*. 2011;46:1045–51.
34. Tandapu P, Provost D, Adams-Huet B, et al. Comparison of the absorption of calcium carbonate and calcium citrate after Roux-en-Y gastric bypass. *Obes Surg*. 2009;19:1256–61.
35. Harper J, Madan AK, Ternovits CA, et al. What happens to patients who do not follow-up after bariatric surgery? *Am Surg*. 2007;73:181–4.

ORIGINAL ARTICLE

Relationships of serum 25-hydroxyvitamin D, ionized calcium and parathyroid hormone after obesity surgery

Stephen Hewitt^{1,2} | Erlend Tuseth Aasheim^{1,3} | Torgeir Thorson Søvik⁴ | Jørgen Jahnsen^{2,5} | Jon Kristinsson^{1,4} | Erik Fink Eriksen^{1,2} | Tom Mala^{1,4}

¹Department of Endocrinology, Morbid Obesity and Preventive Medicine, Oslo University Hospital, Oslo, Norway

²Institute of Clinical Medicine, University of Oslo, Oslo, Norway

³Department of Health Management and Health Economics, University of Oslo, Oslo, Norway

⁴Department of Gastrointestinal Surgery, Oslo University Hospital, Oslo, Norway

⁵Department of Gastroenterology, Akershus University Hospital, Lørenskog, Norway

Correspondence

Stephen Hewitt, Department of Endocrinology, Morbid Obesity and Preventive Medicine, Oslo University Hospital, Oslo, Norway.
Email: stehewi@online.no

Summary

Objective: The high prevalence of secondary hyperparathyroidism (SHPT) after obesity surgery is a concern for long-term bone health. Limited knowledge exists about optimal vitamin D and suppression of parathyroid hormone (PTH) after these procedures. The aim of this study was to investigate the prevalence of SHPT and its relation to vitamin D status.

Design: A cross-sectional study at Oslo University Hospital, Norway.

Patients: A total of 502 consecutive patients, age 22–64 years, attending 2-year follow-up after Roux-en-Y gastric bypass.

Measurements: A serum intact PTH >7.0 pmol/L in the absence of elevated serum ionized calcium (iCa) was considered as SHPT. Vitamin D status was defined by serum concentrations of 25-hydroxyvitamin D (S-25(OH)D).

Results: Altogether, 171 patients (34%) had SHPT. The prevalence of SHPT varied across the range of S-25(OH)D ($P < 0.001$), being highest (71%) with S-25(OH)D < 25 nmol/L. Compared with S-25(OH)D < 50 nmol/L, the prevalence of SHPT was lower with S-25(OH)D ≥ 50 nmol/L (29.0%; RR = 0.64 (95%-CI:0.50–0.81)) and S-25(OH)D ≥ 75 nmol/L (27.7%; RR = 0.61 (95%-CI:0.44–0.84)). S-25(OH)D ≥ 100 nmol/L was associated with the lowest PTH and the lowest prevalence of SHPT (16.0%; RR = 0.35 (95%-CI:0.14–0.88) compared with S-25(OH)D < 50 nmol/L) and the most normal calcium distribution. These associations were most pronounced with iCa in the lower range. A synergistic association was found for S-25(OH)D and iCa on SHPT.

Conclusions: Vitamin D deficient patients had the highest prevalence of SHPT 2 years after gastric bypass. PTH and the prevalence of SHPT were notably lower with S-25(OH)D ≥ 100 nmol/L, compared with lower target levels.

KEYWORDS

bariatric surgery, calcium, morbid obesity, parathyroid hormone, vitamin D

1 | INTRODUCTION

The prevalence of obesity and the number of patients operated for morbid obesity have increased worldwide over the last decades.^{1–3}

Although surgical weight loss improves several obesity-related outcomes, there are concerns about unintended consequences, such as the high prevalence of secondary hyperparathyroidism (SHPT) and long-term bone health.^{3–11}

Food restriction and reduced intestinal absorption post-operatively can explain nutritional deficiencies of calcium and vitamin D, and the increase in serum parathyroid hormone (PTH).^{7,10-12} Often, PTH remains elevated in spite of supplementation with calcium and vitamin D, implying a negative impact on bone metabolism.^{5-8,13,14}

Vitamin D is crucial for sufficient calcium absorption, PTH regulation and bone mineralization^{8,9,15-19}; but yet, there is no consensus on optimal vitamin D after obesity surgery.^{7,14-16}

Some guidelines have recommended to substitute vitamin D aiming at serum concentrations of 25-hydroxyvitamin D (S-25(OH)D) $\geq$ 75 nmol/L.^{3-5,20} Such a target level has not been adopted widely, due to the limited evidence.²¹⁻²³ In a previous study, we found no benefit with these vitamin D levels, as only ionized calcium (iCa) was associated with SHPT.²⁴

In this study, we investigated whether the prevalence of SHPT was lower with S-25(OH)D $\geq$ 75 nmol/L in a large population 2 years after gastric bypass. We also explored other factors related to PTH.

2 | MATERIALS AND METHODS

2.1 | Patients and study design

The study objective of this cross-sectional study was conceived as we worked with our above-mentioned study.

Obesity surgery was offered to selected patients seeking surgical weight loss at the Department of Morbid Obesity and Bariatric Surgery, Oslo University Hospital. Criteria were a body mass index (BMI) ≥ 40 kg/m², or BMI ≥ 35 kg/m² in patients with obesity-related comorbidities, after failed attempts of weight loss by dietary and lifestyle measures.²⁵ Roux-en-Y gastric bypass has been the main surgical procedure at the department. It was performed laparoscopically by constructing a gastric pouch of 25-30 mL, a 150 cm antegastric, antecolic alimentary limb and a biliopancreatic limb of 50 cm.²⁶

All patients were offered standard outpatient follow-up visits and were called for after 6 weeks, 6 months, 1 year and 2 years. From June 2010 to May 2013, we recruited 530 consecutive subjects as they attended a 2-year follow-up consultation after a gastric bypass. In total, 822 morbid obese patients were operated with gastric bypass 2 years earlier. Body weight was measured using an electronic platform weight scale (Seca 635 0-300 kg (Class III)), and height with a wall-fixed adjustable steel measure. Blood samples were drawn for serum analyses after an overnight fast.

Only patients operated at our hospital with valid analyses of PTH, S-25(OH)D and iCa were eligible. Patients with elevated creatinine and patients with a suspicion of primary hyperparathyroidism were excluded (Figure 1).

The study was approved by the Regional Ethical Committee. All patients provided written informed consent. Patient data were registered in a database, licensed by The Norwegian Data Inspectorate.

2.2 | Laboratory analyses

Determinations of intact PTH (1-84), S-25(OH)D and iCa in serum were performed at the Hormone Laboratory, Oslo University Hospital. PTH was analysed by a chemiluminoimmunometric assay (Immulite 2000

XPI, Siemens Healthcare Diagnostics, Los Angeles, CA, USA) (reference range: 1.5-7.0 pmol/L, coefficient of variation (CV) 7%). S-25(OH)D was analysed by radioimmunoassay (Dia-Sorin, Stillwater, MN, USA) (reference range: 37-131 nmol/L, CV 14%), from September 2012 with a liquid chromatography-mass spectrometry (LC-MS/MS) method (identical reference range, CV 9%). The measurements of the two analytic methods were comparable, with mean $\pm$ SD concentrations of S-25(OH)D 63 ± 22 nmol/L vs 60 ± 26 nmol/L ($P = .36$), respectively (Figure S1, found online in the Supporting Information). The reference population for the assessment of S-25(OH)D is described previously.²⁷ Serum iCa was analysed with a Rapidlab 348 analyser (Instru-Med Inc., Atlanta, GA, USA) (reference range: 1.18-1.35 mmol/L, CV 2%), from February 2012 with a Cobas b221 (Roche Diagnostics GmbH, Mannheim, Germany) (reference range: 1.15-1.33 mmol/L, CV 2%). The reference population for iCa was 746 Norwegian women and men aged 45-75 years with PTH and S-25(OH)D within normal reference range (J.P. Berg, unpublished data). Values of iCa from Cobas b221 measurements were transformed to corresponding values of Rapidlab 348 by adding 0.025 mmol/L, according to the difference between the reference ranges. In sensitivity analyses, relationships between S-25(OH)D, iCa, PTH and SHPT were consistent irrespective of analytic methodology of S-25(OH)D and iCa.

Determinations of phosphate, magnesium, creatinine, gamma-glutamyltransferase (γ -GT) and total alkaline phosphatase (ALP) in serum were performed at the Central Laboratory, Oslo University Hospital with a Modular (Roche) analyser and Roche reagents: Phosphate and magnesium photometrically by an end-point measurement, creatinine by an enzymatic method, γ -GT and ALP by an enzymatic calorimetric measurement.

SHPT was defined as a serum concentration of PTH above the reference range (ie >7.0 pmol/L) in the absence of an elevated iCa. S-25(OH)D levels were divided into five categories: <25 , 25-49, 50-74, 75-99 and ≥ 100 nmol/L. We used the common cut point of S-25(OH)D < 50 nmol/L to define insufficient vitamin D status.^{17,18} The term high vitamin D in this text denotes S-25(OH)D ≥ 100 nmol/L with most cases within the range 100-125 nmol/L. Serum iCa was correspondingly divided into five categories: the lowest with values below the normal reference range, and the highest with values above, yielding the following iCa groups: <1.18 , 1.18-1.23, 1.24-1.29, 1.30-1.35 and >1.35 mmol/L. ALP (reference range: 35-105 U/L, CV 8%) is closely linked with PTH and bone loss and was applied to assess bone turnover.^{5,14,20} With normal liver function, 50% of ALP in serum derives from bone in adults.²⁸ We excluded additional 14 patients with elevated γ -GT when analysing ALP, as a disturbance in liver function could not be ruled out, and one was excluded with missing ALP.

Standard recommended dietary supplementation included one tablet daily of multivitamin (200 IU cholecalciferol) and one tablet containing both 500 mg calcium carbonate and 400 IU cholecalciferol (Calcigran Forte®, Takeda Nycomed, Norway) twice a day.

2.3 | Statistical analyses

Statistical analyses were performed with SPSS version 23. P -values $< .05$ were considered significant. Student's t test was applied to

compare independent mean variables. Independent proportions were compared with chi-square test or Fischer's exact test. Relative ratio (RR) for SHPT was calculated between strata of S-25(OH)D.

We examined associations between PTH as the dependent variable and factors of interest in univariate and multivariate linear regression analyses. Explanatory variables with $P < .10$ in univariate analyses were entered in the multiple linear regression, and a backward stepwise procedure was used to eliminate nonsignificant factors. Magnesium was included due to its postulated effects, together with age, gender and BMI as possible confounders.

3 | RESULTS

The flow diagram illustrates recruitment and selection of patients for analyses (Figure 1).

Blood sample analyses and patient characteristics are shown in Table 1. Mean time of follow-up was 26 ± 3 months post-operatively. Women (65%) had lower serum creatinine and higher phosphate, but

PTH did not differ by gender. Women had also lower BMI and a larger percentage weight loss.

In total, 171 patients (34%) had SHPT. PTH associated inversely with S-25(OH)D, and there was a large variation in the prevalence of SHPT over the range of vitamin D levels (both $P < .001$ for linearity) (Table 2). SHPT was most prevalent (71%) in patients with S-25(OH)D < 25 nmol/L and occurred in 45% with S-25(OH)D < 50 nmol/L. The prevalence of SHPT was significantly lower in all groups of S-25(OH)D ≥ 50 nmol/L compared with levels below this cut point: 29.0% and RR 0.64 (95%-CI: 0.50-0.81) with S-25(OH)D ≥ 50 nmol/L; 27.7% and RR 0.61 (95%-CI: 0.44-0.84) with S-25(OH)D ≥ 75 nmol/L; and 16.0% and RR 0.35 (95%-CI: 0.14-0.88) with S-25(OH)D ≥ 100 nmol/L. As illustrated, no differences were found for PTH and SHPT within the S-25(OH)D range 50-99 nmol/L. Compared with S-25(OH)D 50-99 nmol/L, both PTH and the prevalence of SHPT were lower with S-25(OH)D ≥ 100 nmol/L ($P = .007$ for PTH), although nonsignificantly for SHPT (RR 0.53 (95%-CI: 0.21-1.33)). Serum iCa was lowest in subjects with low vitamin D levels and increased with increasing S-25(OH)D, being highest and most normal with S-25(OH)D ≥ 100 nmol/L ($P = .004$ compared with S-25(OH)D < 50 nmol/L, and $P = .056$ compared with S-25(OH)D 50-99 nmol/L).

Serum iCa was inversely associated with PTH and with the prevalence of SHPT (data not shown). Altogether, 473 patients (94%) were normocalcemic, with iCa within the reference range. Serum iCa was below the middle of the normal reference range (ie iCa ≤ 1.26 mmol/L) in 386 subjects (77%), and 263 (52%) had values in the lower third of the reference range or lower (iCa ≤ 1.23 mmol/L). Hypocalcaemia was present in 28 subjects (6%). In SHPT, iCa was below the middle of the normal range in 148 of 171 (87%). The distribution of iCa was most normal with S-25(OH)D ≥ 100 nmol/L.

TABLE 1 Blood chemistry and patient characteristics 2 years after gastric bypass (N = 502)

	Mean	SD
PTH, pmol/L	6.4	3.3
S-25(OH)D, nmol/L	62	23
iCa, mmol/L	1.24	0.04
Phosphate, mmol/L	1.10	0.15
Magnesium, mmol/L	0.82	0.06
Creatinine, μ mol/L	62	11
ALP, U/L	76	28
γ -GT, U/L	28	69
Age, years	46	9
Weight, kg	94	19
Height, cm	172	10
BMI, kg/m ²	32	5
Weight loss, kg	42	16
Total weight loss, %	31	9

PTH, parathyroid hormone; S-25(OH)D, 25-OH vitamin D; iCa, ionized calcium; ALP, alkaline phosphatase; U, Units; γ -GT, γ -glutamyltransferase; BMI, body mass index.

FIGURE 1 Flow diagram of patient inclusion. PTH, parathyroid hormone; S-25(OH)D, serum 25-OH vitamin D; iCa, ionized calcium

TABLE 2 Secondary hyperparathyroidism (%), parathyroid hormone and ionized calcium by serum concentrations of 25-OH vitamin D 2 years after gastric bypass (N = 502)

	N	%	SHPT		PTH (pmol/L)		iCa (mmol/L)	
			N	%	Mean	SD	Mean	SD
S-25(OH)D ^a								
<25	17	3	12	71	9.8	4.3	1.22	0.04
25-49	140	28	59	42	7.0	3.6	1.23	0.03
50-74	204	41	61	30**	6.1**	3.1	1.24 ^{ns}	0.04
75-99	116	23	35	30*	6.0**	2.8	1.24 ^{ns}	0.05
≥100	25	5	4	16**	4.7***	2.8	1.25**	0.04
All	502	100	171	34	6.4	3.3	1.24	0.04

SHPT, secondary hyperparathyroidism (PTH > 7.0 pmol/L); PTH, parathyroid hormone; iCa, ionized calcium.

^aS-25(OH)D, 25-OH vitamin D (nmol/L); ^{ns}P ≥ .05, *P < .05, **P < .01, ***P < .001, each group compared with 25(OH)D < 50 nmol/L.

There was a strong synergistic association of S-25(OH)D and iCa on the prevalence of SHPT (Figure 2). SHPT was present in all subjects (100%) within the lowest categories of both S-25(OH)D and iCa, and no SHPT (0%) was found within the highest categories of both. In subgroup analyses of iCa, the associations of SHPT by vitamin D status were only significant within the lowest two categories of iCa (ie iCa ≤ 1.23 mmol/L) with RR 0.64 (95%-CI: 0.48-0.85) for S-25(OH)D ≥ 50 nmol/L, RR 0.58 (95%-CI: 0.38-0.89) for S-25(OH)D ≥ 75 nmol/L, and RR 0.20 (95%-CI: 0.03-1.33) for S-25(OH)D ≥ 100 nmol/L, all compared with S-25(OH)D < 50 nmol/L.

In linear regression analyses, S-25(OH)D, iCa, phosphate, creatinine, gender and BMI were all independently related to PTH (Table 3). Magnesium, age and weight loss (%) were not found significant. As illustrated in Table 3, the standardized regression coefficients for both S-25(OH)D and iCa were higher than for BMI. Analysing PTH as the dependent, the model was R^2 0.091 with the combination of S-25(OH)D and iCa, and R^2 0.032 with BMI. S-25(OH)D was higher and PTH lower during the months May to September, but the seasonal variation

of S-25(OH)D was not significantly associated with PTH in multivariate analyses.

Figure 3A-C illustrates the relationships of PTH, S-25(OH)D and iCa separately with ALP. ALP was higher with SHPT (81 ± 39 U/L, vs 73 ± 19 U/L in non-SHPT, $P = .007$) and with S-25(OH)D < 50 nmol/L (79 ± 22 U/L, vs 74 ± 29 U/L with S-25(OH)D ≥ 50 nmol/L, $P = .022$).

4 | DISCUSSION

The prevalence of SHPT was particularly high in vitamin D deficient patients 2 years after gastric bypass. Serum PTH was inversely associated with S-25(OH)D, and the prevalence of SHPT showed a large variation across the range of vitamin D levels. Patients with S-25(OH)D ≥ 100 nmol/L had a markedly lower prevalence of SHPT, compared with lower vitamin D levels, and they exhibited the most normal calcium distribution. In addition to S-25(OH)D, PTH was independently related to iCa, phosphate and creatinine, and with gender and BMI.

The overall prevalence of SHPT was high and is comparable to other studies, which may raise concerns about current substitution regimens and future bone health in this population.^{8,14,16}

A relation between vitamin D and PTH is well recognized.^{17,18} Consequently, vitamin D insufficiency is among the most cited mechanisms underlying SHPT and bone loss after obesity surgery.^{6,7,14} In this study, SHPT was reduced with S-25(OH)D ≥ 50 nmol/L. Evidence of vitamin D optimization above this target level is very limited. The present study might suggest a plateau effect for PTH with S-25(OH)D levels in the range 50-99 nmol/L, with a marked decline in PTH and SHPT with vitamin D levels above this range. We are not aware of other reports of S-25(OH)D ≥ 100 nmol/L in these patients.

Of importance for SHPT after gastric bypass surgery, is probably the anatomical changes influencing gastrointestinal uptake of nutrients. Schafer et al recently showed a decrease in calcium absorption after these procedures.²⁹ Serum calcium also seems to be lower after gastric bypass.¹² In this study, there was a left-shifted calcium distribution, corresponding to a state of relative calcium insufficiency. As serum calcium is tightly regulated, low or subnormal levels are

FIGURE 2 Prevalence of secondary hyperparathyroidism (%) by serum concentrations of 25-OH vitamin D and ionized calcium 2 years after gastric bypass (N = 502)

	B	SE	95%-CI	β	P
25-OH vitamin D (nmol/L)	-0.031	0.006	-0.044, -0.018	-0.21	<.001
Ionized calcium (mmol/L)	-15.7	3.5	-22.6, -8.7	-0.19	<.001
Phosphate (mmol/L)	-3.3	1.0	-5.2, -1.4	-0.15	<.001
Magnesium (mmol/L)	3.6	2.2	-0.7, 8.0	0.07	.10
Creatinine (μ mol/L)	0.048	0.015	0.019, 0.077	0.16	.001
BMI (kg/m^2)	0.092	0.027	0.039, 0.15	0.14	.001
Age (years)	0.020	0.016	-0.011, 0.052	0.06	.20
Gender ^b	-0.91	0.34	-1.57, -0.25	-0.13	.007

PTH, parathyroid hormone; B, regression coefficient; SE, standard error; 95%-CI, 95% confidence interval; β , standardized regression coefficient.

^aMultivariate analysis, with nonsignificant factors excluded, R^2 (adjusted) 0.16.

^bGender; women = 1, men = 2. Intercept = 22.7.

TABLE 3 Multivariate linear regression analysis of PTH and potential modulating factors in 502 patients 2 years after gastric bypass^a

counteracted by an increase in PTH, at the expense of mobilizing calcium from bone.⁹ This probably explains that a large proportion of the SHPT cases were normocalcemic.

The primary effect of vitamin D is to stimulate calcium absorption in the intestine and increase serum calcium, with a subsequent negative feedback on PTH secretion.^{15,17-19} This was illustrated as iCa was lowest in subjects with low vitamin D levels, increased with higher vitamin D, and was highest and most normal with high vitamin D. Vitamin D also exerts a direct inhibitory effect on the PTH synthesis in the parathyroids.¹⁵

A combined evaluation of both S-25(OH)D and iCa may prove useful in SHPT, illustrated by their synergistic association. Patients with low or subnormal calcium levels showed the most pronounced benefit on SHPT with higher vitamin D levels. Optimal vitamin D may thus depend on individual calcium status and differ between patients. Differences between individuals can also be explained by a high variability between serum vitamin D and its responses.¹⁶⁻¹⁸

Nutritional deficiencies and inadequate supplementation of calcium and vitamin D are probable mechanisms of SHPT after obesity surgery.⁵⁻⁹ As several others, we did not find differences in PTH and SHPT by supplementation of vitamin D and/or calcium. Higher doses of vitamin D are therefore suggested.

Little is known regarding electrolyte deficiencies and their relationships with PTH after gastric bypass.⁵ In contrast to renal failure patients,³⁰ we found phosphate inversely related to PTH. It may reflect nutritional disturbances and reduced absorption, and elevated PTH levels may reduce serum phosphate by its phosphaturic effects.^{9,30} Phosphate did not differ by S-25(OH)D status (data not shown). Magnesium was included in the model as studies support an effect on PTH secretion, particularly in patients with moderately low serum calcium.³¹ Such a relationship was not found significant in this study. The association found between PTH and creatinine is in accordance with studies in other populations.

The relationship between PTH and obesity needs further investigation.³² Possibly, persistent obesity may induce some increase in PTH in our population, and PTH was observed higher in the patients with BMI $\geq 35 \text{ kg}/\text{m}^2$ 2 years after gastric bypass (data not shown). In our analyses, however, we considered BMI as a confounder, influencing

both S-25(OH)D and PTH.³² Evaluating the relative influence of the factors associated with PTH, we found that the combination of S-25(OH)D and iCa explained more of the variation with PTH, than BMI.

Controversies exist in regard to bone health in the obese population and after obesity surgery. Traditionally, a higher bone mass in the obese has been considered protective for skeletal health. This view is modified recently towards an understanding of skeletal site-specific differences between the obese and nonobese, as total number of fractures seems comparable.³³

It is debated whether the bone loss observed after obesity surgery is solely an adaptation to weight reduction, or whether additional procedure-related mechanisms may contribute, such as intestinal malabsorption and hormonal changes. Based on limited evidence, bone turnover seems higher and bone loss continued although weight stabilizes the second year after gastric bypass, suggesting that skeletal homeostasis has not yet been reached.^{6,34} In our study, ALP was 11% higher in SHPT patients 2 years after gastric bypass, indicating a higher rate of bone turnover and bone loss in the individuals with elevated PTH.^{5,8,14} ALP was also higher (7%) with S-25(OH)D < 50 nmol/L.

There is limited data available about fracture rates after obesity surgery. Some recently published long-term observational studies indicate an increased incidence of atypical fractures following obesity surgery, which may be most pronounced after malabsorptive procedures.³⁵ The role of SHPT needs to be investigated with regard to skeletal end-points.

Strengths of the study include the large single centre population, which made us able to analyse PTH and SHPT within a wide range of S-25(OH)D and to make evaluations of several potential modulating factors. All patients had undergone the same procedure, and data were collected at the same time point post-operatively. Analyses of iCa in serum have higher sensitivity than other analyses, like total and albumin-adjusted calcium.³⁶ Sensitivity analyses showed similar relationships of S-25(OH)D, iCa, PTH and SHPT, irrespective of analytic method in use for these.

Among limitations, nonparticipation may associate with reduced adherence to supplementation, which could influence the parameters studied. An attendance rate of 64% is, however, relatively high after obesity surgery. The cross-sectional design is

FIGURE 3 A-C, Serum alkaline phosphatase (ALP) by serum concentrations of (A) parathyroid hormone, (B) 25-OH vitamin D and (C) ionized calcium in 487 patients 2 years after gastric bypass

inappropriate for causal conclusions between the parameters. A directional effect of vitamin D and calcium on PTH is, however, recognized. The reduction in SHPT by S-25(OH)D ≥ 100 nmol/L was nonsignificant compared with S-25(OH)D in the range 50-99 nmol/L. As PTH was significantly lower, it may be explained with type II error. The notable difference in SHPT observed for the high vitamin D levels was also consistent with exploratory observations in our previous study.²⁴ Due to limited evidence from other populations, the effects we found with S-25(OH)D ≥ 100 nmol/L may primarily apply for specific populations, such as gastric bypass patients exhibiting a relative calcium insufficiency. The linear model explained 16% of the variation with PTH. Variability of the parameters may contribute to this uncertainty. Change in the laboratory methods to

determine the S-25(OH)D and iCa concentrations during the study period may introduce some bias between the methods, but the relationships between S-25(OH)D, iCa, PTH and SHPT were consistent irrespective of methodology. S-25(OH)D, iCa and PTH were available post-operatively only, so analysing the effect of the procedure on these was not possible. Additional bone turnover markers like bone-specific ALP and collagen crosslink peptides could have provided improved insights.

In this study, we found small differences in PTH and SHPT by applying the two thresholds of S-25(OH)D ≥ 50 nmol/L and S-25(OH)D ≥ 75 nmol/L. Some populations may need higher S-25(OH)D concentrations to prevent SHPT.³⁷ In our gastric bypass population, SHPT was most effectively suppressed with S-25(OH)D ≥ 100 nmol/L. More prospective data pertaining vitamin D optimization are needed.

It is hoped that the study will add knowledge relevant to the understanding of SHPT in patients who have undergone obesity surgery, in particular gastric bypass. The effect observed with S-25(OH)D concentrations ≥ 100 nmol/L to improve calcium homeostasis and suppress PTH needs further investigation.

Of particular interest, a large variation in the prevalence of SHPT was observed across the spectrum of S-25(OH)D 2 years after gastric bypass, including a marked reduction with S-25(OH)D concentrations ≥ 100 nmol/L. The data suggest a potential to reduce the prevalence of SHPT by 40%-50% with such high vitamin D concentrations, which may apply particularly to patients with low or subnormal serum calcium. Data on skeletal end-points are, however, needed.

According to two recent studies, high doses of vitamin D supplementation may help to reduce the prevalence of SHPT and resist bone loss after gastric bypass.^{38,39} A large number of vitamin D supplementation studies in different populations have, however, not shown benefits on end-points such as fractures, and some well-designed high-dose vitamin D studies have resulted in increased number of falls and fractures.^{40,41} For almost all of our subjects with high vitamin D, the concentrations of S-25(OH)D were in the range 100-125 nmol/L. According to large observational studies, there are concerns about potential adverse effects of elevating S-25(OH)D above 125 nmol/L.¹⁷

ACKNOWLEDGEMENTS

Solve Tjora, Department of Laboratory Medicine, and Sandra Dahl and the Hormone Laboratory, Oslo University Hospital made valuable help with laboratory analyses. We received statistical support by Lien My Diep, Oslo Centre for Biostatistics and Epidemiology (OCBE), Oslo University Hospital. Thomas Böhmer, Department of Endocrinology, Morbid Obesity and Preventive Medicine, Oslo University Hospital contributed with review and comments to the manuscript.

ORCID

Stephen Hewitt <http://orcid.org/0000-0001-8900-8436>

REFERENCES

- Ng M, Fleming T, Robinson M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014;384:766-781.
- Angrisani L, Santonicola A, Iovino P, Formisano G, Buchwald H, Scopinaro N. Bariatric Surgery Worldwide 2013. *Obes Surg*. 2015; 25:1822-1832.
- Mechanick JI, Kushner RF, Sugerman HJ, et al. AACE/TOS/ASMBS guidelines. American Association of Clinical Endocrinologists, The Obesity Society, and American Society for Metabolic & Bariatric Surgery Medical Guidelines for Clinical Practice for the Perioperative Nutritional, Metabolic, and Nonsurgical Support of the Bariatric Surgery Patient. *Obesity*. 2009;17:S1-S70.
- Mechanick JI, Youdim A, Jones DB, et al. Clinical Practice Guidelines for the Perioperative Nutritional, Metabolic, and Nonsurgical Support of the Bariatric Surgery Patient-2013 Update: Cosponsored by American Association of Clinical Endocrinologists, The Obesity Society, and American Society for Metabolic & Bariatric Surgery. *Obesity*. 2013;21:S1-S27.
- Isom KA, Andromalos L, Ariagno M, et al. Nutrition and metabolic support recommendations for the bariatric patient. *Nutr Clin Pract*. 2014;29:718-739.
- Yu EW. Bone metabolism after bariatric surgery. *J Bone Miner Res*. 2014;29:1507-1518.
- Liu C, Wu D, Zhang J-F, et al. Changes in bone metabolism in morbidly patients after bariatric surgery: a meta-analysis. *Obes Surg*. 2016;26:91-97.
- Bal BS, Finelli FC, Shope TR, Koch TR. Nutritional deficiencies after bariatric surgery. *Nat Rev Endocrinol*. 2012;8:544-556.
- Pramyothin P, Holick MF. Vitamin D supplementation: guidelines and evidence for subclinical deficiency. *Curr Opin Gastroenterol*. 2012;28:139-150.
- Jahnsen J, Falch JA, Mowinckel P, Aadland E. Vitamin D status, parathyroid hormone and bone mineral density in patients with inflammatory bowel disease. *Scand J Gastroenterol*. 2002;37:192-199.
- Karefylakis C, Näslund I, Edholm D, Sundbom M, Karlsson FA, Rask E. Vitamin D status after primary gastric bypass: gravely high prevalence of hypovitaminosis D and raised PTH levels. *Obes Surg*. 2014;24:343-348.
- Risstad H, Søvik TT, Engström M, et al. Five-year outcomes after laparoscopic gastric bypass and laparoscopic duodenal switch in patients with body mass index of 50 to 60: a randomized clinical trial. *JAMA Surg*. 2015;26:353-361.
- Aaseth E, Fagerland MW, Hewitt S, et al. Vitamin concentrations 5 years after gastric bypass. *Eur J Clin Nutr*. 2015;69:1249-1255.
- Peterson LA, Zeng X, Caufield-Noll CP, Schweitzer MA, Magnuson TH, Steele KE. Vitamin D status and supplementation before and after bariatric surgery: a comprehensive literature review. *Surg Obes Relat Dis*. 2016;12:693-704.
- Muscogiuri G, Mitri J, Mathieu C, et al. Vitamin D as a potential contributor in endocrine health and disease. *Eur J Endocrinol*. 2014;171:R101-R110.
- Chakhtoura MT, Nakhoul NN, Shawwa K, Mantzoros C, El Hajj Fuleihan GA. Hypovitaminosis D in bariatric surgery: a systematic review of observational studies. *Metabolism*. 2016;65:574-585.
- The Institute of Medicine. Food and Nutrition Board. *Dietary Reference Intakes for Calcium and Vitamin D*. Washington, D.C.: National Academy Press; 2011.
- Francis R, Aspray T, Fraser W, et al. Vitamin D and bone health: A practical clinical guideline for patient management. National Osteoporosis Society. 2013. <https://nos.org.uk/media/2073/vitamin-d-and-bone-health-adults.pdf>. Latest accessed July 22, 2017.
- Lips P, Bouillon R, van Schoor NM, et al. Reducing fracture risk with calcium and vitamin D. *Clin Endocrinol*. 2010;73:277-285.
- Heber D, Greenway FL, Kaplan LM, Livingston E, Salvador J, Still C. Endocrine and nutritional management of the post-bariatric surgery patient: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab*. 2010;95:4823-4843.
- O'Kane M, Pinkney J, Aasheim ET, Barth JH, Batterham RL. BOMSS Guidelines on peri-operative and postoperative biochemical monitoring and micronutrient replacement for patients undergoing bariatric surgery. <http://www.bomss.org.uk/bomss-nutritional-guidance/>. Latest accessed July 22, 2017.
- Fried M, Yumuk V, Oppert JM, et al. Interdisciplinary European guidelines on metabolic and bariatric surgery. *Obes Surg*. 2014;24:42-55.
- Johnson JM, Maher JW, DeMaria EJ, Downs RW, Wolfe LG, Kellum JM. The long-term effects of gastric bypass on vitamin D metabolism. *Ann Surg*. 2006;243:701-705.
- Hewitt S, Søvik TT, Aasheim ET, et al. Secondary hyperparathyroidism, vitamin D sufficiency, and serum calcium 5 years after gastric bypass and duodenal switch. *Obes Surg*. 2013;23:384-390.
- Anonymous. Gastrointestinal surgery for severe obesity: national Institutes of Health Consensus Development Conference Statement. *Am J Clin Nutr* 1992; 55: 615S-619S.
- Søvik TT, Taha O, Aasheim ET, et al. T. Randomized clinical trial of laparoscopic gastric bypass versus laparoscopic duodenal switch for superobesity. *Br J Surg*. 2010;97:160-166.
- Meyer HE, Falch JA, Sogaard AJ, Haug E. Vitamin D deficiency and secondary hyperparathyroidism and the association with bone mineral density in persons with Pakistani and Norwegian background living in Oslo, Norway, The oslo health study. *Bone*. 2004;35:412-417.
- Delmas PD, Eastell R, Garnero P, Seibel MJ, Stephan J, for the Committee of Scientific Advisors of the International Osteoporosis Foundation. The use of biochemical markers of bone turnover in osteoporosis. *Osteoporos Int*. 2000;11(suppl 6):S2-S17.
- Schafer AL, Weaver CM, Black DM, et al. Intestinal calcium absorption decreases dramatically after gastric bypass surgery despite optimization of vitamin D status. *J Bone Miner Res*. 2015;8:1377-1385.
- Silver J, Naveh-Many T. Phosphate and the parathyroid. *Kidney Int*. 2009;75:898-905.
- Rodríguez-Ortiz ME, Canalejo A, Herencia C, et al. Magnesium modulates parathyroid hormone secretion and upregulates parathyroid receptor expression at moderately low calcium concentration. *Nephrol Dial Transplant*. 2014;29:282-289.
- Grethen E, McClintock R, Gupta CE, et al. Vitamin D and hyperparathyroidism in obesity. *J Clin Endocrinol Metab*. 2011;96:1320-1326.
- Compston J. Obesity and fractures in postmenopausal women. *Curr Opin Rheumatol*. 2015;27:414-419.
- Raoof M, Näslund I, Rask E, Szabo E. Effect of gastric bypass on bone mineral density, parathyroid hormone and vitamin D: 5 years follow-up. *Obes Surg*. 2016;26:1141-1145.
- Lu C-W, Chang Y-K, Chang H-H, et al. Fracture risk after bariatric surgery: a 12-year nationwide cohort study. *Medicine*. 2015;94:e2087.
- Ong GSY, Walsh JP, Stuckey BGA, et al. The importance of measuring ionized calcium in characterizing calcium status and diagnosing primary hyperparathyroidism. *J Clin Endocrinol Metab*. 2012;97:3138-3145.
- Heaney RP. What is vitamin D insufficiency? And does it matter? *Calcif Tissue Int*. 2013;92:177-183.
- Muschitz C, Kocijan R, Haschka J, et al. The impact of vitamin D, calcium, protein supplementation, and physical exercise on bone metabolism after bariatric surgery: the BABS study. *J Bone Miner Res*. 2016;31:672-682.
- Luger M, Kruschitz R, Kienbacher C, et al. Vitamin D₃ loading is superior to conventional supplementation after weight loss surgery in vitamin D-deficient morbidly obese patients: a double-blind randomized placebo-controlled trial. *Obes Surg*. 2017;27:1196-1207.

40. Sanders KM, Stuart AL, Williamson EJ, et al. Annual high-dose oral vitamin D and falls and fractures in older women. A randomized controlled trial. *JAMA*. 2010;303:1815-1822.
41. Bishoff-Ferrari HA, Dawson-Hughes B, Orav EJ, et al. Monthly High-Dose Vitamin D Treatment for the Prevention of Functional Decline. A Randomized Clinical Trial. *JAMA Intern Med*. 2016;176:175-183.

How to cite this article: Hewitt S, Aasheim ET, Søvik TT, et al. Relationships of serum 25-hydroxyvitamin D, ionized calcium and parathyroid hormone after obesity surgery. *Clin Endocrinol (Oxf)*. 2018;88:372–379. <https://doi.org/10.1111/cen.13531>

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

Relationships Between Vitamin D Status and PTH over 5 Years After Roux-en-Y Gastric Bypass: a Longitudinal Cohort Study

Stephen Hewitt^{1,2} · Jon Kristinsson^{1,3} · Erlend Tuset Aasheim^{1,4} · Ingvild Kristine Blom-Høgestøl^{1,2} · Eirik Aaseth⁵ · Jørgen Jahnsen^{2,6} · Erik Fink Eriksen^{1,2} · Tom Mala^{1,3}

Published online: 18 April 2020

© The Author(s) 2020

Abstract

Purpose Secondary hyperparathyroidism (SHPT) after obesity surgery may affect bone health. Optimal vitamin D levels have not been established to prevent SHPT postoperatively. We investigated whether SHPT differed across threshold levels of serum 25-hydroxyvitamin D (S-25(OH)D) from 6 months up to 5 years after Roux-en-Y gastric bypass (RYGB).

Materials and Methods We included 554 patients at follow-up 5 years postoperatively. Blood samples were analysed for S-25(OH)D, ionized calcium (iCa) and parathyroid hormone (PTH) during follow-up.

Results PTH and prevalence of SHPT increased from 6 months to 5 years postoperatively, while S-25(OH)D and iCa decreased (all $P < 0.001$). PTH and SHPT development are related with S-25(OH)D, and PTH differed between all subgroups of S-25(OH)D. SHPT occurred less frequently across all subgroups of S-25(OH)D ≥ 50 nmol/l during follow-up: odds ratio (OR) 0.44 (95% CI 0.36–0.54) in patients with S-25(OH)D ≥ 50 nmol/l, OR 0.38 (0.30–0.49) with S-25(OH)D ≥ 75 nmol/l and OR 0.19 (0.12–0.31) with S-25(OH)D ≥ 100 nmol/l, all compared with S-25(OH)D < 50 nmol/l. At 5 years, 208/554 patients (38%) had SHPT; SHPT was found in 94/188 patients (50%) with S-25(OH)D < 50 nmol/l, in 69/222 (31%) with S-25(OH)D 50–74 nmol/l, in 40/117 (34%) with S-25(OH)D 75–99 nmol/l and in 5/27 (19%) with S-25(OH)D ≥ 100 nmol/l. An interaction existed between S-25(OH)D and iCa. Bone alkaline phosphatase remained increased with SHPT.

Conclusions A significant relationship existed between S-25(OH)D and development of PTH and SHPT. The prevalence of SHPT was lower with threshold levels S-25(OH)D ≥ 50 nmol/l and ≥ 75 nmol/l over the 5 years, and lowest with S-25(OH)D ≥ 100 nmol/l.

Keywords Bariatric surgery · Morbid obesity · Vitamin D · Calcium · Parathyroid hormone · Bone turnover

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11695-020-04582-5>) contains supplementary material, which is available to authorized users.

✉ Stephen Hewitt
stehewi@online.no

Jon Kristinsson
joakri@ous-hf.no

Erlend Tuset Aasheim
erlend.aasheim@gmail.com

Ingvild Kristine Blom-Høgestøl
ingvild.hogestol@gmail.com

Eirik Aaseth
eirik.aaseth@sykehuset-innlandet.no

Jørgen Jahnsen
jorgen.jahnsen@medisin.uio.no

Erik Fink Eriksen
efinkeriksen@gmail.com

Tom Mala
tommenmala@gmail.com

¹ Department of Endocrinology, Morbid Obesity and Preventive Medicine, Oslo University Hospital HF, Aker, P.O. Box 4950 Nydalen, 0424 Oslo, Norway

² Institute of Clinical Medicine, University of Oslo, 0450 Oslo, Norway

³ Department of Gastrointestinal Surgery, Oslo University Hospital, P.O. Box 4950, Nydalen, 0407 Oslo, Norway

⁴ Department of Global Health and Documentation, Norwegian Directorate of Health, P.O. Box 220, Skøyen, 0213 Oslo, Norway

⁵ Department of Medicine, Innlandet Hospital Trust, Elverum, Norway

⁶ Department of Gastroenterology, Akershus University Hospital, 1474 Lørenskog, Norway

Introduction

Obesity surgery provides effective weight loss in morbid obesity [1, 2]. Changes in gastrointestinal anatomy and physiology may also influence intestinal uptake and nutritional status, including for calcium and vitamin D [1–6]. Calcium and vitamin D are central in parathyroid hormone (PTH) regulation [7, 8]. Frequently, PTH is elevated after obesity surgery, and long-term studies indicate that PTH increases over time [4, 9–15].

Relationships of calcium and vitamin D with PTH should be explored further, as PTH may impact bone turnover and bone mineral density (BMD) postoperatively [14–24]. More focus on these are needed during follow-up. Supplementation of calcium and vitamin D is recommended after obesity surgery, but secondary hyperparathyroidism (SHPT) remains prevalent [22, 25]. Few studies have shown differences in prevalence of SHPT with 25-hydroxyvitamin D (S-25(OH)D) threshold levels higher than ≥ 50 nmol/l after obesity surgery [9, 20]. It could be that optimal vitamin D should be higher after obesity surgery than in other nonsurgical populations [7, 8, 24, 26].

We aimed to study the development of PTH and SHPT over 5 years after Roux-en-Y gastric bypass (RYGB), and relationships between these and different threshold levels of S-25(OH)D ≥ 50 nmol/l. We assessed whether the prevalence of SHPT would be lower among patients with higher S-25(OH)D up to 5 years postoperatively.

Materials and Methods

Patients and Study Design

This longitudinal observational cohort study was analysed prospectively, and the report was written to comply with the STROBE checklist [27].

Morbid obesity was defined as BMI ≥ 40 kg/m², or BMI ≥ 35 kg/m² with obesity-related comorbidities [28]. Obesity surgery was offered at Oslo University Hospital, Aker, after failed weight loss by other means. Laparoscopic RYGB was the preferred procedure in the period, with construction of a gastric pouch 25–30 ml, a 150-cm antegastric, antecolic alimentary limb and a 50-cm biliopancreatic limb [29, 30].

We aimed for minimum 500 patients in a population with high follow-up. Candidates were RYGB patients operated 2004–2009. They were evaluated preoperatively and postoperatively with weight, height and blood samples. Follow-up visits were after 6 weeks, 6 months, 1 year, 2 years, 3–4 years and 5 years. At 5 years, all were contacted by letter and eventually by telephone. Body weight was measured electronically (platform weight, Seca 635 0–300 kg, class III), height with

wall-fixed steel measure and blood samples were drawn after overnight fast.

Patients with signed consent and valid PTH and S-25(OH)D at 5 years were candidates, while patients with primary hyperparathyroidism and elevated creatinine were excluded.

Laboratory Analyses

Serum intact PTH (1–84), S-25(OH)D, ionized calcium (iCa) and bone specific alkaline phosphatase (B-ALP) were determined at the Hormone Laboratory, Oslo University Hospital, Aker. PTH was analysed by a chemiluminoimmunoassay (Immulite 2000/2500, Siemens Health Care Diagnostics) (reference range 1.5–7.0 pmol/l, coefficient of variation (CV) 7%). S-25(OH)D was determined by radioimmunoassay (Dia-Sorin) (reference range 37–131 nmol/l, CV 14%), from September 2012 with liquid chromatography-mass spectrometry (LC-MS/MS) (identical reference range, CV 9%); measurements were comparable. Different methods were used for iCa: first, (a) CIBA Corning, instrument 634 Ca²⁺/pH Analyser (Bayer); from December 2005 to December 2007, (b) Rapidlab 348 pH/Blood Gas Analyser (Instru-Med) (both reference range 1.15–1.35 mmol/l, CV 1% and 2%, respectively); from January 2008 to January 2012, (c) Rapidlab 348 pH/Blood Gas Analyser (Instru-Med) (reference range 1.18–1.35 mmol/l); from February 2012, (d) Cobas b221 (Roche Diagnostics) (reference range 1.15–1.33 mmol/l, CV 2%). Reference for iCa was (c) with reference range 1.18–1.35 mmol/l. Values from (a) and (b) were transformed to (c) adding 0.015 mmol/l (differences between reference ranges). Similarly were values from (d) transformed to (c) adding 0.025 mmol/l. B-ALP was determined enzymatically after immune-extraction (Metra Biosystems) (reference range: women 12–31 U/l, men 15–41 U/l (CV 12%). Standard analyses included S-25(OH)D, iCa and PTH during follow-up, which were analysed sporadically preoperatively. Phosphate, magnesium, creatinine and total alkaline phosphatase (ALP) in plasma were analysed at the Central Laboratory, Oslo University Hospital, Aker, with a Modular (Roche) analyzer: Phosphate and magnesium were determined photometrically, creatinine enzymatically and ALP by an enzymatic calorimetric measurement.

Definitions and Subgroups

SHPT was defined as PTH > 7.0 pmol/l with no elevation of iCa (i.e. ≤ 1.35 mmol/l). S-25(OH)D was grouped into 4 categories, < 50 , 50–74, 75–99 and ≥ 100 nmol/l. S-25(OH)D < 50 nmol/l is commonly acknowledged as vitamin D deficiency (or insufficiency) [7, 8]. Calcium levels were grouped into 3 by normal reference range (iCa ≤ 1.23 , 1.24–1.29, ≥ 1.30 mmol/l); all iCa values ≤ 1.23 mmol/l are termed in the

“lower range”. Values below normal reference range (< 1.18 mmol/l) are termed “Low iCa”.

Supplementation

Recommended daily supplements included one multivitamin (cholecalciferol 200 IU) and two combination tablets, each containing calcium carbonate 500 mg and cholecalciferol 400 IU. Compliance was defined by use of calcium ≥ 500 mg and vitamin D ≥ 600 IU minimum 5 days a week, noncompliance as less or no use. Supplements were adjusted to keep blood values within normal reference range, from 2012 to maintain S-25(OH)D ≥ 50 nmol/l, or S-25(OH)D ≥ 75 nmol/l in cases with SHPT [7, 8]. Our supplementation regimen also included oral iron (100 mg daily) and intramuscular vitamin B12 injections (1 mg per 3 months).

Statistical Analyses

Statistical analyses were performed with IBM SPSS for Windows, version 25. Continuous and categorical variables were tested with *t* test and chi-square test as appropriate. Regression analyses were performed with linear mixed model, diagonal covariance matrices for PTH and B-ALP using individual repeated measurements, time-dependent covariates and random intercept. Variables were included in multivariate analyses of PTH with stepwise backward elimination of non-significant variables. Gender, age and BMI were included as covariates. We tested multiplicative interactions for S-25(OH)D, iCa and time on PTH, and PTH and time on B-ALP. B-ALP was adjusted for gender. Generalized estimating equations (GEE), unstructured covariance matrices were used for SHPT. Missing data were not imputed. We analysed two periods: from baseline to 6 months postoperatively and from 6 months to 5 years. The second period was the main focus, as S-25(OH)D and PTH were routinely assessed. Continuous variables are presented with means and standard deviations ($\pm$ SD), categorical variables in percentages, and odds ratios (OR) and relative ratios (RR) with 95% confidence intervals (95% CI).

Results

Of 823 operated, 584 (71%) attended 5-year follow-up. Included were 554 patients (67%), after exclusion of 4 with no signed consent, 10 with a suspicious primary hyperparathyroidism, 3 with elevated creatinine, and 13 with missing data of S-25(OH)D and PTH at 5 years. Follow-up period was 5.3 ± 0.4 years. Three patients had moved from our region and could not be contacted, and 7 had died during the 5 years. Table 1 summarizes preoperative characteristics.

Table 1 Preoperative characteristics ($N = 554$)

	<i>N</i>	Mean	SD
Women	384		
Age (y)	554	41.7	9.1
Weight (kg)	554	136	22
Height (cm)	554	172	10
BMI (kg/m ²)	554	46.2	5.3
Blood samples			
PTH (pmol/l)	160	6.4	2.8
S-25(OH)D (nmol/l)	158	53	22
iCa (mmol/l)	245	1.25	0.04
Phosphate (mmol/l)	204	1.02	0.19
Magnesium (mmol/l)	247	0.81	0.07
Creatinine (μ mol/l)	549	68	13
B-ALP (U/l)	44	26.3	6.6
SHPT	56	(35%)	

BMI, body mass index; *PTH*, parathyroid hormone; *S-25(OH)D*, serum 25-hydroxyvitamin D; *iCa*, ionized calcium; *B-ALP*, bone alkaline phosphatase; *SHPT*, secondary hyperparathyroidism

The First 6 Months

BMI decreased 27% in the first 6 months from 46.2 ± 5.3 to 33.8 ± 5.1 kg/m², PTH decreased from 6.4 ± 2.8 to 5.2 ± 2.5 pmol/l and prevalence of SHPT from 35 to 18% (all $P < 0.001$). S-25(OH)D and iCa were independently related with PTH development ($P < 0.001$).

PTH decreased in all subgroups of S-25(OH)D. Patients with S-25(OH)D < 50 nmol/l had highest PTH preoperatively and 6 months postoperatively, and they had the largest decrease in PTH ($P < 0.001$). PTH related positively with B-ALP ($P < 0.001$).

Six Months to 5 Years Postoperatively

Figure 1 illustrates development of PTH and SHPT from 6 months to 5 years postoperatively. PTH increased to 6.8 ± 3.5 pmol/l, and the prevalence of SHPT increased to 38%, while S-25(OH)D decreased to 59 ± 24 nmol/l, and iCa to 1.22 ± 0.04 mmol/l (all $P < 0.001$). BMI increased to 34.5 ± 6.2 kg/m² ($P = 0.005$).

Vitamin D, PTH and SHPT

S-25(OH)D was inversely related with development of PTH and occurrence of SHPT from 6 months to 5 years ($P < 0.001$), and PTH differed between all subgroups of S-25(OH)D ($P < 0.001$) (Fig. 1). Mean PTH response by change in vitamin D levels was.

$$\Delta\text{PTH}(\text{pmol/l}) = -0.031 * \Delta\text{25(OH)D}(\text{nmol/l})$$

Fig. 1 Observed development of parathyroid hormone (PTH) and secondary hyperparathyroidism (%) from 6 months to 5 years after Roux-en-Y gastric bypass ($N = 554$) to the left, and for subgroups of 25-hydroxyvitamin D (S-25(OH)D) to the right

expressing that each 1 nmol/l increase in S-25(OH)D related with a PTH decrease of 0.031 pmol/l during follow-up, and vice versa. From 6 months, PTH and prevalence of SHPT increased most with S-25(OH)D < 50 nmol/l ($P < 0.001$), but not significantly with S-25(OH)D ≥ 100 nmol/l.

Over the 5 years, SHPT differed between subgroups of S-25(OH)D (Fig. 2). Higher S-25(OH)D levels were associated with lower occurrence of SHPT: OR 0.44 (0.36–0.54) with S-25(OH)D ≥ 50 nmol/l, OR 0.38 (0.30–0.49) with S-25(OH)D ≥ 75 nmol/l, OR 0.19 (0.12–0.31) with S-25(OH)D ≥ 100 nmol/l, all compared with S-25(OH)D < 50 nmol/l.

50 nmol/l. Compared with S-25(OH)D 50–99 nmol/l, OR was 0.40 (0.26–0.62) with S-25(OH)D ≥ 100 nmol/l.

At 5 years, 208 of 554 patients (38%) had SHPT. The prevalence was 50% with S-25(OH)D < 50 nmol/l (Fig. 3/ Table 2); 61% with S-25(OH)D < 25 nmol/l. SHPT was less frequent with S-25(OH)D ≥ 50 nmol/l (31%, RR 0.63 (0.51–0.77)), S-25(OH)D ≥ 75 nmol/l (31%, RR 0.63 (0.47–0.83)), and lowest with S-25(OH)D ≥ 100 nmol/l (19%, RR 0.37 (0.17–0.83)), all compared with S-25(OH)D < 50 nmol/l. Compared with S-25(OH)D 50–99 nmol/l, PTH was lower with S-25(OH)D ≥ 100 nmol/l

Fig. 2 Odds ratio (OR) for secondary hyperparathyroidism (SHPT) from 6 months to 5 years postoperatively in 554 Roux-en-Y gastric bypass patients by categories of serum 25-hydroxyvitamin D (S-25(OH)D, nmol/l) and ionized calcium (iCa, mmol/l) compared with reference (ref.). S-25(OH)D was adjusted for time (y): OR 1.11 (1.02–1.22), with additional adjustments for baseline S-25(OH)D and iCa as given

Fig. 3 Relative ratio (RR) for secondary hyperparathyroidism (SHPT) 5 years after Roux-en-Y gastric bypass ($N = 554$) by categories of serum 25-hydroxyvitamin D (S-25(OH)D or D, nmol/l) and ionized calcium (iCa, mmol/l) compared with reference (ref.)

(5.0 ± 2.1 compared with 6.4 ± 3.3 pmol/l, $P = 0.032$), but SHPT not significantly (RR 0.57 (0.25–1.28)) at 5 years. These relationships were not significant with iCa above lower range (iCa ≥ 1.24 mmol/l).

Ionized Calcium and BMI

Serum iCa decreased in all subgroups of vitamin D from 6 months to 5 years postoperatively, and percentage with iCa in the lower range (iCa ≤ 1.23) increased from 26% to 61% ($P < 0.001$) (Fig. 4). PTH increased in all 3 subgroups of iCa over 5 years, and was higher in the lower range. iCa was inversely related with PTH development ($P < 0.001$). Correspondingly, SHPT was consistently lower with iCa above lower range during follow-up.

An interaction existed between S-25(OH)D and iCa. At 5 years, SHPT was most prevalent with combined vitamin D deficiency and iCa in the lower range, and there was no SHPT

with S-25(OH)D ≥ 100 nmol/l and iCa above the lower range (Fig. 3).

Multivariate Analyses

In multivariate analyses of PTH from 6 months to 5 years (Table 3), the relationship with S-25(OH)D and iCa remained robust, also with BMI. Interactions for S-25(OH)D and iCa with time over the 5 years were not significant in multivariate analyses. SHPT occurred more often in men, with lower age and with higher BMI over the period (Supplementary Table). The trend of lower OR with higher S-25(OH)D remained across strata of gender, age and BMI.

Supplementation, PTH/SHPT and Bone Turnover

360 of 475 (77%) used supplements of calcium and vitamin D at 6 months, and 278 (50%) at 5 years. S-25(OH)D was higher

Table 2 Observed numbers with secondary hyperparathyroidism by vitamin D status at follow-up from 6 months to 5 years after Roux-en-Y gastric bypass ($N = 554$)

	6 months		1 year		2 years		3–4 years		5 years	
	SHPT	All	SHPT	All	SHPT	All	SHPT	All	SHPT	All
S-25(OH)D										
< 50	25	77	38	100	57	121	39	86	94	188
50–74	20	125	49	178	57	171	50	147	69	222
75–99	17	126	35	134	28	117	20	72	40	117
≥ 100	6	51	2	39	3	30	2	14	5	27
All	68	379	124	451	145	439	111	319	208	554
Missing		175		103		115		235		0

S-25(OH)D, serum 25-hydroxyvitamin D; SHPT, secondary hyperparathyroidism; All, attenders

Fig. 4 Observed development of ionized calcium by subgroups of 25-hydroxyvitamin D (S-25(OH)D), and proportions of low calcium and calcium within the lower range from 6 months to 5 years after Roux-en-Y gastric bypass ($N = 552$) (upper), and development of parathyroid hormone (PTH) and proportion with secondary hyperparathyroidism by subgroups of ionized calcium (lower)

Table 3 Regression analysis^a of parathyroid hormone and modulating factors from 6 months to 5 years after Roux-en-Y gastric bypass ($N = 554$)

	<i>B</i>	SE	<i>P</i>
Univariate			
S-25(OH)D	− 0.032	0.0025	< 0.001
iCa	− 23.9	1.46	< 0.001
BMI	0.12	0.014	< 0.001
Phosphate	− 3.57	0.38	< 0.001
Magnesium	− 0.12	0.031	0.69
Creatinine	0.023	0.0075	0.003
Time	0.32	0.028	< 0.001
Preop Vit D	− 0.034	0.0091	< 0.001
Gender ^b	0.35	0.23	0.13
Age ^c	− 0.0069	0.012	0.56
Multivariate^d			
I	42.2	4.86	< 0.001
S-25(OH)D	− 0.248	0.064	< 0.001
iCa	− 30.65	3.81	< 0.001
BMI	0.065	0.014	< 0.001
S-25(OH)D × iCa	0.182	0.051	< 0.001
Creatinine	0.020	0.0076	0.009
Time	0.073	0.030	0.015

^a Linear mixed model; ^b men 1, women 0; ^c Age at 5 years; ^d Adjusted for age and gender; ×, interaction

RYGB, Roux-en-Y gastric bypass; *I*, intercept; *B*, regression coefficient; *SE*, standard error; *S-25(OH)D*, serum 25-hydroxyvitamin D (nmol/l); *iCa*, ionized calcium (mmol/l); *BMI*, body mass index (kg/m²);

Creatinine (μmol/l); Time, time postoperatively (years); Preop Vit D, S-25(OH)D preoperatively

and PTH lower in users over the period ($P < 0.001$), but not significantly at 5 years.

PTH related positively with B-ALP and ALP over the 5 years ($P < 0.001$), with interaction for PTH and time. B-ALP declined by 1.3 U/l per year, but remained increased with SHPT ($P < 0.001$ for group difference). Total ALP declined slower with SHPT ($P = 0.004$).

Discussion

PTH and the prevalence of SHPT decreased the first 6 months after RYGB, and thereafter increased up to 5 years. S-25(OH)D was related with PTH development. Patients with vitamin D deficiency had the largest increase in PTH from 6 months, while patients with high S-25(OH)D ≥ 100 nmol/l had lowest PTH and prevalence of SHPT.

Vitamin D and SHPT

This study is among few longitudinal long-term reports of SHPT by vitamin D and calcium status after RYGB. Vitamin D deficiency is usually defined by S-25(OH)D < 30 – 50 nmol/l, while target levels of S-25(OH)D ≥ 50 – 75 nmol/l are adopted in most recommendations after obesity surgery [1, 2, 4–8]. However, evidence supporting these target levels is limited [22, 24].

The observed increase in PTH from 6 months is in accordance with other long-term evaluations [9–13]. S-25(OH)D was strongly related with PTH over time, however, with limited differences in SHPT by traditional target thresholds,

defined by $S-25(OH)D \geq 50$ nmol/l and ≥ 75 nmol/l. The findings with $S-25(OH)D \geq 100$ nmol/l corresponded with our cross-sectional study 2 years postoperatively [20].

Calcium and SHPT

Calcium absorption seems reduced after RYGB [31, 32]. However, few have reported a relationship between calcium and PTH [18, 20, 23]. Extracellular calcium is a determinant of PTH secretion, and even calcium within the lower normal range may increase PTH [20, 33, 34]. With the feedback mechanisms involved, PTH may increase above reference range, while serum calcium still remains within normal reference range.

Our observations suggest a role of iCa on PTH. The proportion of patients with iCa in the lower range increased over time, and iCa was related with PTH development. This relationship was independent of $S-25(OH)D$. An interaction existed between $S-25(OH)D$ and iCa on PTH. Still, iCa declined in all subgroups of $S-25(OH)D$ but more slowly with higher levels.

Supplementation, SHPT and Bone Effects

Several studies have failed to document benefits of vitamin D and calcium supplementation, and the regimens have been questioned [22, 24]. This study supports a modest effect with supplements of calcium ≥ 500 mg and vitamin D ≥ 600 IU on PTH. These doses are however lower than recommended by many [2, 4, 6].

We also found higher B-ALP in patients with SHPT up to 5 years postoperatively, suggesting higher bone turnover. SHPT might therefore help explain increased bone turnover, which is observed up to 5 years after RYGB [18, 19, 22]. SHPT may also lead to reduced BMD, which is observed after weight stabilizes 1–2 years postoperatively [15, 19]. We recently found SHPT related with lower BMD 10 years after RYGB [21].

Implications

In clinical practice, SHPT may be considered as a marker of vitamin D and calcium insufficiency, and it is of concern after RYGB. Further research should address whether increasing $S-25(OH)D$ levels can suppress SHPT and improve bone health. Higher doses seem necessary to achieve sufficient vitamin D levels and suppress SHPT after RYGB [2, 4, 24, 35].

Attention to calcium status seems relevant to identify risk for SHPT. SHPT was more frequent with iCa in the lower range. Higher $S-25(OH)D$ can increase calcium levels and lower PTH, and some individuals may need higher vitamin D levels than others [20, 26, 36]. The interaction between $S-25(OH)D$ and iCa may be relevant in clinical practice.

SHPT was not prevalent with $S-25(OH)D \geq 100$ nmol/l and iCa in the upper two tertiles of reference range at 5 years, which we previously reported 2 years postoperatively [20].

Optimal $S-25(OH)D$ levels are not established after RYGB and obesity surgery in general. Achieving $S-25(OH)D \geq 100$ nmol/l may be needed to suppress SHPT more effectively in some individuals, with an aim to improve long-term bone health.

Strengths and Limitations

The main strength of this study was a large sample size with high 5-year follow-up rate and repeated measurements, providing statistical strength. Bias of primary hyperparathyroidism was minimized. The single centre study design with standard surgical and follow-up procedures strengthen internal validity, but the findings need testing in other populations. As relationships up to 5 years were the primary focus, only patients with 5-year data on $S-25(OH)D$ and PTH were included. Prevalence of SHPT in nonattenders may be higher than observed, assuming less compliance [37]. Data on pre-operative $S-25(OH)D$ and PTH, and $S-25(OH)D \geq 100$ nmol/l and B-ALP, during follow-up were limited. Supplemental use was self-reported. iCa determinations at the Hormone Laboratory were adjusted during the period; however, the drop in iCa was also found in some parallel analyses performed at the Central Laboratory, with unchanged methodology.

Conclusions

$S-25(OH)D$ levels related inversely with PTH development and occurrence of SHPT up to 5 years after RYGB. The prevalence of SHPT was lower with $S-25(OH)D$ thresholds ≥ 50 nmol/l and ≥ 75 nmol/l. Some patients may need $S-25(OH)D \geq 100$ nmol/l to suppress SHPT more effectively.

Acknowledgements Open access funding provided by Oslo University & Oslo University Hospital. Per Thorsby at the Hormon Laboratory and Solve Tjora at the Central Laboratory assisted with laboratory data, Inger Eribe and Marianne Sæter at Center for Morbid Obesity and Bariatric Surgery assisted with patient data, all located at Oslo University Hospital, Aker.

Compliance with Ethical Standards

Disclosures, Ethics and Consent The Regional Ethical Committee approved the study. All procedures performed were in accordance with the ethical standards of the institutional and national research committee and with the 1964 Helsinki declaration and its later amendments. Informed consent was obtained from all individual participants included in the study.

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Mechanick JJ, Kushner RF, Sugerman HJ, et al. American Association of Clinical Endocrinologists, The Obesity Society, and American Society for Metabolic & Bariatric Surgery medical guidelines for clinical practice for the perioperative nutritional, metabolic, and nonsurgical support of the bariatric surgery patient. *Obesity*. 2009;17(Suppl 1):S1–S70.
- Busetto L, Dicker D, Azran C, et al. Obesity Management Task Force of the European Association for the study of obesity released “practical recommendations for the post-bariatric surgery medical management”. *Obes Surg*. 2018;28(7):2117–21.
- Via MA, Mechanick JJ. Nutritional and micronutrient care of bariatric surgery patients: current evidence update. *Curr Obes Rep*. 2017;6(3):286–96.
- Mechanick JJ, Youdim A, Jones DB, et al. Clinical practice guidelines for the perioperative nutritional, metabolic, and nonsurgical support of the bariatric surgery patient–2013 update: cosponsored by American Association of Clinical Endocrinologists, the Obesity Society, and American Society for Metabolic & Bariatric Surgery. *Surg Obes Relat Dis*. 2013;9(2):159–91.
- Parrott J, Frank L, Rabena R, et al. American Society for Metabolic and Bariatric Surgery Integrated Health Nutritional Guidelines for the surgical weight loss patient 2016 update: micronutrients. *Surg Obes Relat Dis*. 2017;13(5):727–41.
- Sandvik J, Laurenius A, Naslund I, Videhult P, Wiren M, Aasheim ET. [Nordic guidelines for follow-up after obesity surgery]. *Tidsskr Nor Lægeforen* 2018; 138(4).
- Ross AC, Manson JE, Abrams SA, et al. The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *J Clin Endocrinol Metab*. 2011;96(1):53–8.
- Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2011;96(7):1911–30.
- Johnson JM, Maher JW, DeMaria EJ, et al. The long-term effects of gastric bypass on vitamin D metabolism. *Ann Surg*. 2006;243(5):701–4. discussion 4–5.
- Switzer NJ, Marcil G, Prasad S, et al. Long-term hypovitaminosis D and secondary hyperparathyroidism outcomes of the Roux-en-Y gastric bypass: a systematic review. *Obes Rev*. 2017;18(5):560–6.
- Monaco-Ferreira DV, Leandro-Merhi VA, Aranha NC, et al. Vitamin D deficiency and parathormone increase in late post-operative gastric bypass in Roux-En-Y. *Arq Bras Cir Dig*. 2018;31(4):e1407.
- Aaseth E, Fagerland MW, Aas AM, et al. Vitamin concentrations 5 years after gastric bypass. *Eur J Clin Nutr*. 2015;69(11):1249–55.
- Risstad H, Sovik TT, Engstrom M, et al. Five-year outcomes after laparoscopic gastric bypass and laparoscopic duodenal switch in patients with body mass index of 50 to 60: a randomized clinical trial. *JAMA Surgery*. 2015;150(4):352–61.
- Karefylakis C, Naslund I, Edholm D, et al. Vitamin D status 10 years after primary gastric bypass: gravely high prevalence of hypovitaminosis D and raised PTH levels. *Obes Surg*. 2014;24(3):343–8.
- Raoof M, Naslund I, Rask E, et al. Effect of gastric bypass on bone mineral density, parathyroid hormone and vitamin D: 5 years follow-up. *Obes Surg*. 2016;26(5):1141–5.
- Liu C, Wu D, Zhang JF, et al. Changes in bone metabolism in morbidly obese patients after bariatric surgery: a meta-analysis. *Obes Surg*. 2016;26(1):91–7.
- Svanevik M, Risstad H, Hofso D, et al. Bone turnover markers after standard and distal Roux-en-Y gastric bypass: results from a randomized controlled trial. *Obes Surg*. 2019;29:2886.
- Yu EW. Bone metabolism after bariatric surgery. *J Bone Miner Res*. 2014;29(7):1507–18.
- Lindeman KG, Greenblatt LB, Rourke C, et al. Longitudinal 5-year evaluation of bone density and microarchitecture after Roux-en-Y gastric bypass surgery. *J Clin Endocrinol Metab*. 2018;103(11):4104–12.
- Hewitt S, Aasheim ET, Sovik TT, et al. Relationships of serum 25-hydroxyvitamin D, ionized calcium and parathyroid hormone after obesity surgery. *Clin Endocrinol*. 2018;88(3):372–9.
- Blom-Hogestol IK, Hewitt S, Chahal-Kummen M, et al. Bone metabolism, bone mineral density and low-energy fractures 10 years after Roux-en-Y gastric bypass. *Bone*. 2019;127:436–45.
- Gagnon C, Schafer AL. Bone health after bariatric surgery. *JBM plus*. 2018;2(3):121–33.
- Hewitt S, Sovik TT, Aasheim ET, et al. Secondary hyperparathyroidism, vitamin D sufficiency, and serum calcium 5 years after gastric bypass and duodenal switch. *Obes Surg*. 2013;23(3):384–90.
- Chakhtoura MT, Nakhoul NN, Shawwa K, et al. Hypovitaminosis D in bariatric surgery: a systematic review of observational studies. *Metabolism*. 2016;65(4):574–85.
- Peterson LA, Zeng X, Caufield-Noll CP, et al. Vitamin D status and supplementation before and after bariatric surgery: a comprehensive literature review. *Surg Obes Relat Dis*. 2016;12(3):693–702.
- Rosen CJ, Abrams SA, Aloia JF, et al. IOM committee members respond to Endocrine Society vitamin D guideline. *J Clin Endocrinol Metab*. 2012;97(4):1146–52.
- von Elm E, Altman DG, Egger M, et al. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. 2007;370(9596):1453–7.
- Gastrointestinal surgery for severe obesity: National Institutes of Health Consensus Development Conference Statement. *Am J Clin Nutr*. 1992;55(2 Suppl):615s–9s.
- Sovik TT, Taha O, Aasheim ET, et al. Randomized clinical trial of laparoscopic gastric bypass versus laparoscopic duodenal switch for superobesity. *Br J Surg*. 2010;97(2):160–6.
- Salte OB, Sovik TT, Risstad H, Skattum J, Blom-Hogestol IK, Eribe IEL, et al. Bariatric surgery at Oslo University Hospital 2004–14. *Tidsskr Nor Lægeforen* 2019; 139(10).
- Schafer AL, Weaver CM, Black DM, et al. Intestinal calcium absorption decreases dramatically after gastric bypass surgery despite optimization of vitamin D status. *J Bone Miner Res*. 2015;30(8):1377–85.
- Riedt CS, Brolin RE, Sherrell RM, et al. True fractional calcium absorption is decreased after Roux-en-Y gastric bypass surgery. *Obesity*. 2006;14(11):1940–8.
- Kontogeorgos G, Trimpou P, Laine CM, et al. Normocalcaemic, vitamin D-sufficient hyperparathyroidism - high prevalence and low morbidity in the general population: a long-term follow-up

- study, the WHO MONICA project, Gothenburg, Sweden. *Clin Endocrinol*. 2015;83(2):277–84.
34. Bollerslev J, Schalin-Jantti C, Rejnmark L, et al. Management of endocrine disease: unmet therapeutic, educational and scientific needs in parathyroid disorders. *Eur J Endocrinol*. 2019;181(3):P1–p19.
35. Muschitz C, Kocijan R, Haschka J, et al. The impact of vitamin D, calcium, protein supplementation, and physical exercise on bone metabolism after bariatric surgery: the BABS study. *J Bone Miner Res*. 2016;31(3):672–82.
36. Heaney RP. What is vitamin D insufficiency? And does it matter? *Calcif Tissue Int*. 2013;92(2):177–83.
37. Harper J, Madan AK, Ternovits CA, et al. What happens to patients who do not follow-up after bariatric surgery? *Am Surg*. 2007;73(2):181–4.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Full Length Article

Bone metabolism, bone mineral density and low-energy fractures 10 years after Roux-en-Y gastric bypass

Ingvild Kristine Blom-Høgestøl^{a,b,*}, Stephen Hewitt^{a,b}, Monica Chahal-Kummen^{a,b}, Cathrine Brunborg^c, Hanne Løvdal Gulseth^{a,d}, Jon A. Kristinsson^{a,e}, Erik Fink Eriksen^{a,b}, Tom Mala^{a,e}

^a Department of Endocrinology, Morbid Obesity and Preventive Medicine, Oslo University Hospital, Oslo, Norway

^b Institute of Clinical Medicine, University of Oslo, Oslo, Norway

^c Oslo Centre for Biostatistics and Epidemiology, Research Support Services, Oslo University Hospital, Oslo, Norway

^d Department of Chronic Diseases and Ageing, Norwegian Institute of Public Health, Norway

^e Department of Gastrointestinal and Pediatric Surgery, Oslo University, Norway

ARTICLE INFO

Keywords:

Obesity
Bariatric surgery
Bone health, fracture
Secondary hyperparathyroidism
Weight loss

ABSTRACT

Background: Roux-en-Y gastric bypass (RYGB) is a common surgical procedure for treatment of morbid obesity. RYGB induces considerable and sustained weight loss, and remission of obesity related-comorbidities. While studies have suggested negative effects of RYGB on bone health, long-term data are lacking. We aimed to evaluate the prevalence of aBMD below the expected range for age, osteopenia, osteoporosis and low-energy fractures in a defined patient cohort 10 years after RYGB. Secondly, we wanted to identify factors associated with increased risk of aBMD z-score or t-score of -1.1 or lower 10 years after RYGB.

Methods: Patients undergoing RYGB surgery from June 2004 to December 2006 at the Department of Morbid Obesity and Bariatric Surgery, Oslo University Hospital, a tertiary referral centre for treatment of morbid obesity, were invited to a 10 year follow-up. Follow-up visits included morning fasting blood samples, clinical examination, anthropometric measures and dual energy X-ray absorptiometry (DXA).

Results: Out of 194 patients eligible for the study, 124 attended the 10 year follow-up and 122 (63%) were examined with DXA. Mean (SD) age was 50.3 (9.0) years, 118 (97%) were of Caucasian ethnicity, 94 were females (77%), of whom 41 (44%) were postmenopausal.

Secondary hyperparathyroidism (SHPT) was noted in 37 participants (31%) and vitamin D deficiency (value below 50 nmol/L) and insufficiency (value below 75 nmol/L) in 40 (33%) and 91 (75%), respectively. Among the 63 participants who were premenopausal females or males 49 years or younger the prevalence of areal bone mineral density (aBMD) in the lower range of normal (z-score -1.1 to -1.9) was 30% ($n = 19$) and aBMD below the expected range for age (z-score ≤ -2.0) was noted in 8% ($n = 5$). Among the 59 participants who were postmenopausal females or males 50 years or older, the prevalence of osteopenia (t-score -1.1 to -2.4) was 51% ($n = 30$) and osteoporosis (t-score ≤ -2.5) was 27% ($n = 16$). The bone resorption markers CTX-1 and PINP were higher in participants with aBMD z-score or t-score of -1.1 or lower compared to participants with aBMD z-score or t-score of -1.0 or higher. Preoperative hypothyroidism, or higher age, postmenopausal status, BMI < 35 kg/m², SHPT or higher PINP levels at 10 year follow-up were independently associated with aBMD z-score or t-score of -1.1 or lower 10 years after RYGB. Eighteen participants (15%) reported a clinical low-energy fracture after RYGB. In addition, vertebral fracture assessment by DXA revealed that 10 participants (8%) had experienced at least one moderate to severe morphometric vertebral fracture.

Conclusion: Ten years after RYGB 27% of postmenopausal females and males 50 years or older were osteoporotic, and 8% of premenopausal females and males 49 years or younger exhibited aBMD below the expected range for age. The prevalence of fragility fractures was high. SHPT, higher age, postmenopausal status or higher

Abbreviations: aBMD, Areal bone mineral density; BALP, Bone specific alkaline phosphatase; CKD, Chronic kidney disease; CTX-1, Carboxyl terminal telopeptide of type 1 collagen; eGFR, estimated glomerular filtration rate; FSH, Follicular stimulating hormone; DXA, Dual energy X-ray absorptiometry; PINP, Procollagen type 1 N-terminal propeptide; RYGB, Roux-en-Y gastric bypass; SHPT, Secondary hyperparathyroidism; T2D, Type 2 diabetes; TSH, Thyroid stimulating hormone; 25-hydroxyvitamin D, 25(OH) vitamin D

* Corresponding author at: Lilleåsen 15C, 1396 Billingstad, Norway.

E-mail address: ingvild.hogestol@gmail.com (I.K. Blom-Høgestøl).

<https://doi.org/10.1016/j.bone.2019.07.014>

Received 14 March 2019; Received in revised form 10 July 2019; Accepted 12 July 2019

Available online 16 July 2019

8756-3282/ © 2019 Elsevier Inc. All rights reserved.

PINP levels at 10 years and preoperative hypothyroidism were all independent risk factors for aBMD z-score or t-score of -1.1 or lower 10 years after RYGB.

1. Introduction

Laparoscopic Roux-en-Y gastric bypass (RYGB) surgery is a widely applied surgical technique for treatment of morbid obesity following failed attempts of sustained weight loss [1]. RYGB induces considerable and sustained weight loss, remission of obesity related comorbidities and reduces mortality [2–4]. However, changes to gastrointestinal anatomy and physiology may pose negative effects on bone health.

Shortly after RYGB, increased levels of bone turnover markers and reduced bone mineral density, assessed by dual energy X-ray absorptiometry (DXA) or quantitative computed tomography, have been noted [5–8]. Persistent elevation of serum bone turnover markers and continuous reductions in areal bone mineral density (aBMD) following weight stabilization indicate effects of RYGB on bone metabolism beyond that of adaptation to a reduced weight [9]. In line with the high turnover state increased fracture rates have been reported in several studies [10,11]. The high turnover state after RYGB is probably multifactorial. The high prevalence of secondary hyperparathyroidism (SHPT) [7,12], likely contributes to this state of high bone turnover.

These findings emphasize the need for long-term evaluations of bone health after RYGB. Unfortunately studies describing bone health five to 10 years after RYGB are few, characterised by small samples of females only, and with low or undefined rates of follow-up [13,14].

We aimed to evaluate the prevalence of aBMD below the expected range for age, osteopenia, osteoporosis and low-energy fractures in a defined patient cohort 10 years after RYGB. Secondly, we wanted to identify factors associated with an increased risk of aBMD z-score or t-score of -1.1 or lower 10 years after RYGB. We hypothesized that SHPT would be associated with an increased risk of aBMD z-score or t-score of -1.1 or lower 10 years after RYGB.

2. Methods

2.1. Study population

Patients undergoing RYGB surgery between June 2004 and December 2006 at the Department of Morbid Obesity and Bariatric Surgery, Oslo University Hospital, a tertiary referral centre for treatment of morbid obesity, were invited to a 10 year follow-up. Eligibility criteria for RYGB were body mass index (BMI) ≥ 40 kg/m² or BMI ≥ 35 kg/m² with obesity-related comorbidity, age between 20 and 60 years, and failed attempts of sustained weight loss. Results from the five year follow-up have been published previously [15]. Ten year follow-up visits were conducted from June 2016 to March 2018.

2.2. Surgery, follow-up and study visits

Prior to surgery all patients had one or more individual consultations with a nurse, a dietician and a surgeon. If indicated they also had consultations with an internist or a psychologist. Preoperative data were registered on predefined forms. Until December 2005 preoperative data were registered retrospectively, with subsequent prospective registration. Patients underwent a laparoscopic RYGB with a gastric pouch of about 25 mL, a 150 cm antecolic alimentary limb and a 50 cm biliopancreatic limb [16]. Postoperative routine clinical follow-up visits were conducted six-eight weeks, six months, one, two and five years after surgery. After surgery all patients were prescribed oral supplementation therapy consisting of daily doses of calcium carbonate (500 mg $\times$ 2 daily), vitamin D3 (400 IU $\times$ 2 daily), Nycoplus multi-vitamin – Nycomed® (one tablet), iron (100–200 mg) as well as vitamin

B12 injections (1 mg) every three months. At routine clinical visits, vitamin levels were monitored and additional supplements were recommended if appropriate.

Ten year follow-up visit included morning fasting blood samples, clinical examination, anthropometric measures and DXA. All, but four, 10 year follow-up visits were performed by one clinician (SH). Data concerning comorbidity, medications, nutritional supplements, menstrual status and fracture history were recorded in a predefined case report form.

DXA scans were performed for assessment of lumbar spine (L₁–L₄), hip, proximal femur and total body aBMD and body composition (including percent body fat). Vertebral fracture assessment by DXA was evaluated for the presence of vertebral fractures. All DXA scans were performed by the same nurse. GE Lunar Prodigy was used until August 26, 2016 when it was replaced by GE Lunar iDXA. The two DXA scanners were cross-calibrated by scanning 16 volunteers with both machines and revealed lumbar spine (L1–L4) intra-class correlation coefficient (ICC) (95% CI) of 0.989 (0.968 to 0.996), and for femoral neck and total hip, ICC (95% CI), was 0.994 (0.982 to 0.998) and 0.996 (0.988 to 0.999), respectively. Furthermore, studies have shown excellent *in vivo* precision of the GE Lunar iDXA for the measurement of lumbar spine, hip and total body aBMD in adults with normal and obese BMI [17,18]. Our DXA machine is calibrated daily against the standard calibration phantom, supplied by the manufacturer, with estimated short-term precision errors of $< 1.0\%$ for aBMD at the lumbar spine and at the femoral neck.

Blood samples were drawn before 10 am after an overnight fast, centrifuged and analyzed shortly after retrieval. The Hormone Laboratory, Oslo University Hospital analyzed: thyroid stimulating hormone (TSH) using a non-competitive immunofluorometric method from DELFIA (reference range *TSH mIU/L*: 0.5–3.6). Carboxyl terminal telopeptide of type 1 collagen (CTX-1) and procollagen type 1 N-terminal propeptide (PINP) using Roche® electrochemiluminescence immunoassay (ECLIA) (reference ranges based on information from the manufacturer; *CTX-1 µg/L*: females 25–49 years: ≤ 0.57 , ≥ 50 years: ≤ 1.01 , males 30–50 years: ≤ 0.58 , 51–70 years: ≤ 0.7 and *PINP µg/L*: females > 25 years: 11–94, males > 25 years: 20–91). Bone specific alkaline phosphatase (BALP) and osteocalcin using LIAISON® chemiluminescence immunoassay (CLIA) (reference range based on information from the manufacturer; *BALP µg/L*: 5.5–24.6, *osteocalcin nmol/L*: females ≥ 21 years: 1.5–5.4, males ≥ 21 years 1.6–4.3 [19]). Serum 25-hydroxyvitamin D (25(OH) vitamin D) levels were analyzed by liquid chromatography-mass spectrometry (LC-MS/MS) method, Calcitriol (1.25(OH)₂D3) vitamin D with enzymatic immunoassay from IDS Nordic (reference range 1.25(OH)₂D3 *pmol/L*: 39–193). Serum parathyroid hormone (PTH) by Immulite 2000 XPI, Siemens Healthineers a chemiluminometric assay (reference range *PTH pmol/L*: 1.5–7.0). Serum ionized calcium using Roche® Cobas b221 (reference range *ionized calcium mmol/L*: 1.15–1.33). FSH was analyzed using Immulite 2000 XPI, Siemens Healthineers, a non-competitive immunoluminometric assay. The coefficient of variance for CTX-1, PINP, osteocalcin, 25(OH) vitamin D and PTH is 5%, 5%, 6%, 11% and 7%, respectively.

The Central Laboratory of Oslo University Hospital analyzed HbA_{1c} using Tosoh G8 high-performance liquid chromatography and calculated estimated glomerular filtration rate (eGFR) using CKD-EPI formula (*mL/min/1.73m²*); were categorized into the following stages of chronic kidney disease (CKD): CKD stage I: > 90 , CKD stage II: 60–90, CKD stage III: 30–59, stage IV: 5–29, stage V: < 15 .

2.3. Outcomes

2.3.1. Osteopenia, osteoporosis and aBMD below expected range for age

Areal BMD t-scores represent the number of standard deviations an actual aBMD deviates from the peak bone mass of young women. The aBMD z-score represents the number of standard deviations an actual BMD deviates from the expected aBMD of age, gender and ethnicity. Both scores are based on the reference population from the NHANES and Lunar studies given by the manufacturer. T-scores and Z-scores of the lumbar spine L1-L4 were calculated after exclusion of vertebrae with osteoarthritic changes (spondylosis) or compression fractures. The percent estimates of aBMD z-score were referred to as percent of expected aBMD.

In order to explore aBMD outcomes the population was split according to gender, age and menopausal status:

i. Premenopausal females and males 49 years or younger

aBMD in the higher range of normal: 1.0 standard deviations lower than the expected aBMD of age, gender and ethnicity or higher (z-score > -1.0).

aBMD in the lower range of normal: 1.1 to 1.9 standard deviations

lower than the expected aBMD of age, gender and ethnicity (z-score < -1.0 to > -2.0).

aBMD below expected range for age: 2.0 standard deviations lower than the expected aBMD of age, gender and ethnicity (z-score ≤ -2.0) [20].

ii. Postmenopausal females and males 50 years or older [21]:

Normal aBMD: 1.0 standard deviations lower than the peak bone mass of young women or higher (t-score > -1.0).

Osteopenia: 1.1 to 2.4 standard deviations lower than the peak bone mass of young women (t-score < -1.0 to > -2.5).

Osteoporosis: aBMD 2.5 standard deviations or more lower than the peak bone mass of young women (t-score ≤ -2.5).

2.3.2. Fractures

Vertebral fractures were assessed on lateral X-rays recorded on the DXA scanner and by visual semiquantitative technique (VFA). Moderate (reduction of vertebral height of > 25 –40%) and severe (reduction of vertebral height of > 40 %) vertebral fractures are reported [22]. Vertebral fractures revealed by DXA are referred to as morphometric vertebral fractures.

Table 1

Participant characteristics 10 years after Roux-en-Y gastric bypass (RYGB).

	All n = 122	Premenopausal females and males younger than 50 years of age n = 63	Postmenopausal females and males 50 years or older n = 59	p-Value
Caucasian, n	118 (97%)	62 (98%)	56 (95%)	0.47
Age, years	50.3 $\pm$ 9.0	44.2 $\pm$ 6.7	56.8 $\pm$ 6.1	< 0.001
Body mass index, kg/m ²	35.6 $\pm$ 7.2	37.4 $\pm$ 7.0	33.6 $\pm$ 6.6	0.003
% fat mass	45.4 $\pm$ 7.8	46.4 $\pm$ 7.2	44.2 $\pm$ 7.1	0.043
% total weight loss after RYGB	24.6 $\pm$ 13.6	22.6 $\pm$ 11.5	26.8 $\pm$ 15.4	0.093
Loss of height, cm	1.5 $\pm$ 1.9	0.91 $\pm$ 1.6	2.1 $\pm$ 2.0	0.001
First degree relative with osteoporosis	22 (18%)	8 (13%)	14 (24%)	0.11
First degree relative with hip fracture	17 (14%)	6 (10%)	11 (19%)	0.15
TSH, mIU/L	1.7 (< 0.001 –7.4)	1.5 (< 0.001 –7.4)	1.9 (0.2–4.8)	0.026
eGFR, mL/min/1.73m ²	98.4 $\pm$ 16.0	104.8 $\pm$ 12.8	91.5 $\pm$ 16.4	< 0.001
Ionized calcium, mmol/L	1.2 $\pm$ 0.039	1.2 $\pm$ 0.032	1.2 $\pm$ 0.045	0.70
25(OH) vitamin D, nmol/L	60.0 $\pm$ 22.9	56.1 $\pm$ 21.8	64.2 $\pm$ 23.5	0.052
1.25(OH) ₂ D ₃ , pmol/L	135.1 $\pm$ 41.9	139.0 $\pm$ 38.9	133.2 $\pm$ 41.6	0.44
PTH, pmol/L	5.9 $\pm$ 2.8	5.9 $\pm$ 2.4	5.9 $\pm$ 3.2	0.96
CTX-1, μ g/L	0.48 $\pm$ 0.24	0.41 $\pm$ 0.18	0.54 $\pm$ 0.27	0.002
PINP, μ g/L	55.4 $\pm$ 26.3	58.6 $\pm$ 28.5	52.1 $\pm$ 23.5	0.19
BALP, μ g/L	14.3 $\pm$ 5.4	14.0 $\pm$ 5.5	14.6 $\pm$ 5.3	0.51
Osteocalcin, nmol/L	4.1 $\pm$ 1.2	3.9 $\pm$ 1.1	4.2 $\pm$ 1.4	0.22
Areal bone mineral density				
Lumbar spine (L1-L4), g/cm ²	1.17 $\pm$ 0.16	1.21 $\pm$ 0.14	1.12 $\pm$ 0.17	0.001
Femoral neck (left), g/cm ²	0.92 $\pm$ 0.13	0.97 $\pm$ 0.11	0.86 $\pm$ 0.13	< 0.001
Total hip, g/cm ²	0.97 $\pm$ 0.16	1.02 $\pm$ 0.13	0.93 $\pm$ 0.18	0.002
Total body, g/cm ²	1.17 $\pm$ 0.15	1.22 $\pm$ 0.12	1.11 $\pm$ 0.16	< 0.001
T-score				
Lumbar (L1-L4) ^a	-0.9 $\pm$ 1.3	-0.55 $\pm$ 1.17	-1.3 $\pm$ 1.34	0.001
Femoral neck (left)	-0.8 $\pm$ 0.98	-0.36 $\pm$ 0.85	-1.19 $\pm$ 0.96	< 0.001
Total hip	-0.4 $\pm$ 1.1	0.07 $\pm$ 1.01	-0.79 $\pm$ 1.09	< 0.001
Total body	0.64 $\pm$ 1.4	1.19 $\pm$ 1.2	0.016 $\pm$ 1.3	< 0.001

The results are given as number (proportion in per cent) for categorical variables, mean (SD) for continuous variables with normal distribution and median (range) for other variables $\geq$.

Loss of height is the difference between the measured preoperative height and the measured height at 10 year follow-up.

Abbreviations; Thyroid stimulation hormone (TSH), estimated glomerular filtration rate (eGFR), 25-hydroxyvitamin D (25(OH) vitamin D), Calcitriol (1.25(OH)₂D₃), parathyroid hormone (PTH), carboxyl terminal telopeptide of type 1 collagen (CTX-1), procollagen type 1 N-terminal propeptide (PINP), bone specific alkaline phosphatase (BALP).

Missing (n): FSH (4), % total weight loss (3), TSH (2), eGFR (1), ionized calcium (2), 1.25(OH)₂ vitamin D (5), PTH (1), CTX-1 (4), PINP (5), BALP (7), Osteocalcin (10), aBMD lumbar spine (1), aBMD left femoral neck (6), aBMD total hip (5), aBMD total body (3).

Reference ranges: TSH mIU/L: 0.5–3.6; eGFR (mL/min/1.73 m²): Stages of chronic kidney disease (CKD): CKD stage I: > 90 , CKD stage II: 60–90, CKD stage III: 30–59, stage IV: 5–29, stage V: < 15 ; ionized calcium mmol/L: 1.15–1.33; 25(OH) vitamin D nmol/L: 37–132; 1.25(OH)₂D₃ pmol/L: 39–193; PTH pmol/L: 1.5–7.0; CTX-1 μ g/L: Females 25–49 years: ≤ 0.57 , ≥ 50 years: ≤ 1.01 , males 30–50 years: ≤ 0.58 , 51–70 years: ≤ 0.7 ; PINP μ g/L: Females > 25 years: 11–94, males > 25 years: 20–91; BALP μ g/L: 5.5–24.6; osteocalcin nmol/L: Females ≥ 21 years: 1.5–5.4, males ≥ 21 years: 1.6–4.3.

^a Vertebrae with osteoarthritic changes (spondylosis) or compression fractures were excluded from calculation.

Clinical fractures were self-reported and registered in predefined case report forms. A low energy fracture was defined as a fracture resulting from minimal trauma quantified as forces equivalent to a fall from standing height or less [21]. All low energy fractures, except digital fractures, acquired during the 10 years after RYGB are reported.

2.3.3. Calcitropic hormones and supplements

Vitamin D deficiency was defined as serum 25(OH) vitamin D levels below 50 nmol/L and insufficiency as levels below 75 nmol/L [23,24]. Seasons were defined as; Summer: June to August, fall: September to November, winter: December to February and spring: March to May. SHPT was defined as a serum concentration of PTH above 7.0 pmol/L in the absence of serum ionized calcium above 1.33 mmol/L. Intake of vitamin supplements was self-reported. Patients were classified as taking calcium and vitamin D supplements if they reported intake of 1000 mg and 800 IU, respectively, or more, at least five days a week.

2.3.4. Menopausal status

Hormonal intrauterine devices made clinical evaluation of menstrual cycle challenging. For this reason a postmenopausal status was defined as a serum follicle stimulating hormone (FSH) of > 25 IU/L or higher [25].

2.3.5. Comorbidities

Preoperative type 2 diabetes (T2D) was defined as $HbA_{1c} \geq 6.5\%$ and/or the use of one or more oral glucose lowering drugs with or without the use of insulin. Preoperative hypothyroidism was based on information given by the participant preoperatively, stated in referrals or based on use of levothyroxine substitution treatment.

2.4. Statistical analysis

Normally distributed continuous variables are presented as mean $\pm$ SD, others are presented as median (range). Categorical data are reported as proportions (percentage). Subgroups were compared with an independent sample *t*-test, the Chi Square test or Fisher exact test as appropriate. Intra-class correlation coefficient (ICC) with 95% CI was used to assess concordance between the two DXA scanners (GE Lunar Prodigy and GE Lunar iDXA). ICC values of 0.75 or higher were considered excellent [26]. One way-ANOVA was used to explore possible seasonal differences in 25(OH) vitamin D values. To explore the association between PTH and aBMD z-scores, linear regression analyses were applied. The results from the linear regression analyses are presented as regression coefficients (β) with 95% CI.

Logistic regression analyses were performed to identify factors associated with aBMD z-score or t-score of -1.1 or lower 10 years after RYGB. All variables with a $p < 0.25$ in the univariable analysis were entered into the multivariable logistic regression model using a manual backward stepwise elimination procedure. Multivariable analyses were preceded by estimation of correlation between risk factors. Predictors that correlated > 0.7 , were not included in the model in order to avoid multicollinearity. The associations between factors and aBMD z-score or t-score of -1.1 or lower 10 years after RYGB at 10 year follow-up visit were quantified by calculating odds ratios (OR) with 95% CI. In linear regression analysis all continuous variables were checked for deviation from normality, non-linear effects, multicollinearity, and homoscedasticity. For logistic regression analysis they were checked for deviation of linearity of the logit and multicollinearity. Evaluation of the models predictive accuracy was assessed by calibration (Hosmer and Lemeshow goodness-of-fit test) and by the discriminatory capability (area under the ROC curve). Two tailed *p*-values < 0.05 were considered statistically significant. All statistical analyses were performed using the IBM SPSS statistics version 25.0 (IBM SPSS Inc., Armonk, NY: IBM Corp).

2.5. Ethics

The study was conducted in accordance with the Declaration of Helsinki and approved by the Regional Committee for Medical and Health Research Ethics South-East Norway; 2015/142. Written informed consent was obtained from all participants. Preoperative data were registered in a database licensed by the Norwegian Data Inspectorate.

3. Results

Nine out of 203 patients died prior to the 10 year follow-up; thus 194 were eligible for the study. Of these, 124 attended the 10 year follow-up, and 122 (63%) were examined with DXA scans and were included in this study.

3.1. Participant characteristics

Participant characteristics are presented in Table 1. Mean age was 50.3 (SD 9.0) years, 118 (97%) were of Caucasian ethnicity. There were 94 females (77%), of whom 41 (44%) were postmenopausal. Of the 122 participants 63 (52%) were premenopausal females or males 49 years or younger, and 59 (48%) were postmenopausal females or males 50 years or older.

At 10 year follow-up, five participants received bone specific treatment for osteoporosis (oral bisphosphonates (2), intravenous bisphosphonates (2) or denosumab (1): one male older than 50 years of age and four postmenopausal females, they were all classified as osteoporotic). These participants had lower CTX-1 levels, $0.23 \pm 0.11 \mu\text{g/L}$, compared to $0.49 \pm 0.23 \mu\text{g/L}$ for the remaining cohort, $p = 0.034$, but their PINP levels of $51.0 \pm 14.7 \mu\text{g/L}$ were comparable to $55.6 \pm 26.6 \mu\text{g/L}$ for the remaining cohort, $p = \text{n.s.}$ Three participants reported treatment for rheumatoid arthritis, one had been treated for prostate cancer and one had undergone surgery for primary hyperparathyroidism. No participants had been diagnosed with breast cancer or Cushing's disease. Twenty-five participants (21%) had CKD stage II and three had CKD stage III (3%), no participant had CKD stage IV or higher. At the 10 year follow-up seven participants presented with low TSH levels ($< 0.5 \text{ mIU/L}$), five had known preoperative hypothyroidism, while two had no known thyroid condition.

3.2. Calcium and calcitropic hormones

A total of 29 participants (24%) reported regular intake of calcium supplements and 38 (31%) reported regular intake of vitamin D supplements. Individual measurements of serum ionized calcium, PTH and 25(OH) vitamin D are shown in Fig. 1. Forty participants (33%) were vitamin D deficient and 91 (75%) insufficient, respectively. Among the participants reporting regular intake of vitamin D supplements, five (12.5%) had vitamin D deficiency, a significantly lower fraction when compared to participants not taking supplements where 33 participants (41%) were deficient, $p = 0.002$. 25(OH) vitamin D levels drawn during summer, fall, winter and spring season were 61.9 ± 20.5 , 65.4 ± 22.4 , 52.8 ± 24.4 and 56.8 ± 22.9 , respectively. No statistical difference was noted between the 25(OH) vitamin D levels of the different seasons.

No participant had serum ionized calcium or 25(OH) vitamin D levels above the reference range. SHPT was noted in 37 participants (31%). Participants with SHPT exhibited numerically lower levels of 25(OH) vitamin D compared to those without SHPT, 54.0 ± 3.9 and 62.7 ± 2.4 , respectively, $p = 0.055$.

3.3. Markers of bone turnover

Mean values for the bone turnover markers; CTX-1, PINP, BALP and osteocalcin, are given in Table 1 and individual measurements are

Fig. 1. Individual measurements of serum ionized calcium, parathyroid hormone and 25(OH) vitamin D in 122 participants 10 years after Roux-en-Y gastric bypass. Reference range is marked with lines. Reference ranges: Serum ionized calcium mmol/L: 1.15–1.33; 25(OH) vitamin D nmol/L: 37–132; PTH pmol/L: 1.5–7.0.

presented in Fig. 2. Participants with an aBMD z-score or t-score of -1.1 or lower 10 years after RYGB revealed higher levels of CTX-1 and PINP compared to participants with an aBMD z-score or t-score of -1.1 or lower 10 years after RYGB. However, only the association with PINP remained significant in the multivariable analysis (Table 2).

3.4. Bone mineral density

The percent estimates of aBMD z-scores, relative to expected aBMD of age, gender, and ethnicity, were below 90% in 35%, 43% and 27% of participants, in the total hip, femoral neck and lumbar spine, respectively (Fig. 3). An inverse association was noted between PTH and aBMD z-score of femoral neck; $\beta -0.051$ 95% CI (-0.099 to -0.002), $p = 0.040$, while the association with lumbar spine and total hip aBMD z-score were non-significant (Fig. 4).

3.4.1. Osteopenia, osteoporosis and aBMD below expected range for age
3.4.1.1. Premenopausal females and males 49 years or younger. Of the 63 participants who were premenopausal females or males 49 years or younger, 5 (8%) revealed aBMD below the expected range for age (z-score ≤ -2.0) and 19 (30%) had osteopenia (z-score -1.1 to -1.9) of the spine, femoral neck or total hip 10 years after RYGB.

3.4.1.2. Postmenopausal females and males 50 years or older. Of the 59 participants who were postmenopausal females or males 50 years or older 16 (27%) had osteoporosis (t-score ≤ -2.5) and 30 (51%) had aBMD in the osteopenic range (t-score -1.1 to -2.4) of the spine, femoral neck or total hip 10 years after RYGB.

The prevalence of aBMD below the expected range for age and/or osteoporosis in the different skeletal locations (spine, femoral neck and total hip) are presented in Table 3. Of the total cohort 70 (57%) participants exhibited an aBMD z-score or t-score of -1.1 or lower. Pre-operative hypothyroidism, or higher age, postmenopausal status, BMI $< 35 \text{ kg/m}^2$, higher PINP value or SHPT at 10 year follow-up were all independent risk factors for aBMD z-score or t-score of -1.1 or lower 10 years after RYGB (Table 2). The Hosmer-Lemeshow's goodness of fit test for the multivariable model indicated a satisfactory fit of the model ($p = 0.35$). The discriminatory capability (area under the ROC) between participants with and without a z-score or t-score of -1.1 10 years after RYGB was 0.81 (95% CI: 0.73–0.89). When exploring risk factors for aBMD below the expected range for age, osteoporosis and/or fragility fractures (low energy fracture and morphometric vertebral fractures) only age was detected as a risk factor (Supplementary Table 1).

3.5. Fractures

Clinical low-energy fractures after RYGB were reported by 18 participants (15%) during the 10 years of follow-up, seven (11%) premenopausal females or males 49 years or younger and 11 (19%) postmenopausal females or males 50 years or older. Lower limb fractures were the most prevalent, followed by rib fractures (Table 4). The mean duration from RYGB to first low-energy fracture was $8.4 \text{ years} \pm 1.8$. Participants with osteoporotic aBMD had a significantly higher prevalence of low energy fractures when compared to participants with normal or osteopenic aBMD ($p = 0.023$). Vertebral fracture assessments of DXA revealed that three male (11%) and seven female (7%) participants had experienced one or more moderate to severe morphometric vertebral fractures. Of these 10 participants, one had presented with symptoms of a vertebral fracture.

4. Discussion

4.1. Areal bone mineral density below the expected range for age and osteoporosis

Among premenopausal females and males 49 years or younger the prevalence of aBMD below expected range for age was 8% and among the postmenopausal females and males 50 years or older the prevalence of osteoporosis was 27%. Duran et al. observed osteoporosis (t-score ≤ -2.5) in 13% in a study of 30 females, with a median age of 46 years, eight years after RYGB [13]. However, without knowing the prevalence of postmenopausal status it is difficult to directly compare this study to ours. The Tromsø study, a Norwegian population based study, reported a prevalence of osteoporosis of the femoral neck of 4.8% in males and 6.1% in females for subjects aged 50–69 years in contrast to our findings of 13% and 33%, respectively [27]. Furthermore, we noted that more than one in three had aBMD measures below 90% of expected for age, gender and ethnicity. Collectively, our findings suggest that RYGB may be a risk factor for aBMD below expected range for age and osteoporosis.

Fig. 2. Individual measurements of bone turnover markers in 122 participants 10 years after Roux-en-Y gastric bypass. Reference range is marked with lines. Reference ranges: Carboxyl terminal telopeptide of type 1 collagen (CTX-1), $\mu\text{g/L}$: females 25–49 years: ≤ 0.57 , ≥ 50 years: ≤ 1.01 , males 30–50 years: ≤ 0.58 , 51–70 years: ≤ 0.7 ; procollagen type 1 N-terminal propeptide (PINP), $\mu\text{g/L}$: females > 25 years: 11–94, males > 25 years: 20–91; bone specific alkaline phosphatase (BALP), $\mu\text{g/L}$: 5.5–24.6; osteocalcin, nmol/L : females ≥ 21 years: 1.5–5.4, males ≥ 21 years: 1.6–4.3. Missing (n): CTX-1 (4), PINP (5), BALP (7), Osteocalcin (10).

4.2. Bone turnover

We observed higher levels of bone turnover markers (CTX-1 and PINP) than those reported in preoperative cohorts of similar age [6,28,29] indicating elevated bone turnover 10 years after RYGB. Participants with an aBMD z-score or t-score of -1.1 or lower 10 years after RYGB had higher levels of both CTX-1 and PINP, suggesting that longstanding accelerated bone turnover is a central mechanism underlying the development of decreasing aBMD after RYGB. Several studies comparing changes in aBMD and bone turnover markers, have shown that high turnover states increase bone loss and fracture rates [30].

4.3. Mechanisms

The pathophysiology underlying the high rates of an aBMD z-score or t-score of -1.1 or lower 10 years after RYGB is likely multifactorial. We noted that participants with SHPT had approximately three times higher odds for having an aBMD z-score or t-score of -1.1 or lower 10 years after RYGB, and we also observed an inverse association between PTH and aBMD z-scores of the femoral neck. Together these findings support our hypothesis that SHPT cause negative effects on skeletal health after RYGB. Studies have revealed significantly reduced intestinal calcium absorption after RYGB [7,31]. To what extent regular intake of calcium and vitamin D supplements could have prevented SHPT in our population is beyond the scope of the study. Notably, the standard recommendation of calcium supplementation was in the form of calcium carbonate, in contradiction to recent European and American recommendations of calcium citrate. Furthermore, the standard doses of 800 IU vitamin D3 daily is in line with Norwegian and European guidelines for post-bariatric management [32,33], but are considerably lower than the 3000 IU daily recommended by the American guidelines [24]. Additionally, we found that only one of three participants reported regular intake of the recommended dose of calcium and vitamin D supplements, that vitamin D

non-compliance was associated with vitamin D deficiency, and that participants with SHPT had numerically lower 25(OH) vitamin D levels compared to the remaining cohort. Other cohorts have also reported limited compliance in regard to use of recommended supplements after bariatric surgery [34,35].

Participants with preoperative hypothyroidism revealed four times higher odds for having an aBMD z-score or t-score of -1.1 or lower 10 years after RYGB. In adulthood, the thyroid hormone is important for bone mass maintenance and strength [36]. Untreated hypothyroidism is associated with reduced bone turnover and a positive bone balance [37]; which generally protects against bone loss and fracture. Due to simple, reliable biochemical diagnostic tools, patients are quickly diagnosed and put on thyroid substitution therapy. Studies of subjects receiving substitution treatment for hypothyroidism have revealed discordant results regarding whether or not it is associated with a lower aBMD [38,39], but supraphysiological doses of substitution therapy have been shown to exert negative effects on bone in hypothyroid patients [40]. In addition a hyperthyroid state preceding hypothyroidism could have negatively affected bone health in this group. We therefore favor the notion that excessive doses of thyroid hormone or possible hyperthyroidism preceding hypothyroidism, after thyroiditis or surgical/radioiodine treatment of hyperthyroidism, constitute plausible reasons for our findings.

Mechanical loading of bone plays a key role in determining bone mass, strength and size [41,42]. In line with this we noted that participants with $\text{BMI} < 35 \text{ kg/m}^2$ 10 years after RYGB had a doubled risk of having an aBMD z-score or t-score of -1.1 or lower 10 years after RYGB, however this association did not remain significant in the multivariable analysis, indicative of limited importance in the long-term after RYGB. As for the general population, postmenopausal status and older age were associated with an increased risk of having an aBMD z-score or t-score of -1.1 or lower 10 years after RYGB [21]. Our findings are supported by Schafer et al. who described that postmenopausal females had a larger decrease in bone mineral density after RYGB when compared to premenopausal females and males [29]. Older age was the

Table 2Factors associated with aBMD z-score or t-score -1.1 or lower 10 years after Roux-en-Y gastric bypass (RYGB).

	Univariable				Multivariable	
	aBMD z-score or t-score of -1.0 or higher 10 years after RYGB n = 52	aBMD z-score or t-score of -1.1 or lower 10 years after RYGB n = 70	Odds ratio (95% CI)	p-value	Odds ratio (95% CI)	p-Value
Gender, female	39 (75%)	55 (79%)	1.2 (0.52–2.9)	0.64		
Heredity osteoporosis	5 (10%)	17 (24%)	3.0 (1.0–8.8)	0.044		
Preoperative						
Hypothyroidism	4 (8%)	18 (27%)	4.4 (1.4–14.0)	0.012	4.1 (1.1–16.3)	0.041
Type 2 diabetes	13 (26%)	21 (34%)	1.5 (0.66–3.4)	0.34		
BMI > 50	19 (37%)	21 (31%)	0.79 (0.37–1.7)	0.55		
10 years after RYGB						
Age, years	47.4 ± 8.0	52.5 ± 9.1	1.1 (1.0–1.1)	0.002	1.1 (1.0–1.1)	0.016
Postmenopausal	7 (14%)	35 (52%)	6.7 (2.6–16.9)	< 0.001	4.7 (1.6–14.1)	0.005
Δ BMI	-10.0 ± 6.6	-11.6 ± 6.7	0.96 (0.91–1.0)	0.18*		
BMI < 35 kg/m ²	20 (39%)	40 (57%)	2.1 (1.0–4.4)	0.043		
Loss of height, cm	-1.4 ± 1.9	-1.6 ± 1.9	0.95 (0.78–1.1)	0.57		
Smoking	15 (29%)	26 (37%)	1.5 (0.67–3.2)	0.34		
Muscular or skeletal pain	32 (62%)	47 (67%)	1.3 (0.63–2.8)	0.52		
Secondary hyperparathyroidism	13 (25%)	24 (35%)	1.6 (0.72–3.6)	0.25	2.8 (1.0–7.9)	0.045
25(OH) vitamin D insufficiency	41 (79%)	50 (71%)	0.67 (0.29–1.6)	0.35		
Low-energy fracture after RYGB	7 (13%)	11 (16%)	1.2 (0.43–3.3)	0.73		
CTX-1, µg/L	0.39 ± 0.17	0.54 ± 0.26	34.1 (3.9–295)	< 0.001		
PINP, µg/L	48.1 ± 17.2	60.7 ± 30.3	1.0 (1.0–1.0)	0.016	1.0 (1.0–1.1)	0.006

aBMD z-score or t-score of -1.1 or lower 10 years after includes:

i. Premenopausal females and males 49 years or younger

aBMD in the lower range of normal: 1.1 to 1.9 standard deviations lower than the expected aBMD of age, gender and ethnicity (z-score < -1.0 to > -2.0).aBMD below expected range for age: 2.0 standard deviations lower than the expected aBMD of age, gender and ethnicity (z-score ≤ -2.0) [20].

ii. Postmenopausal females and males 50 years or older [21]

Osteopenia: 1.1 to 2.4 standard deviations lower than the peak bone mass of young women (t-score < -1.0 to > -2.5).Osteoporosis: aBMD 2.5 standard deviations or more lower than the peak bone mass of young women (t-score ≤ -2.5).

Heredity of osteoporosis was defined as having a first degree relative with osteoporosis diagnosis.

The results are given as number (proportion in per cent) for categorical variables, mean (SD) for continuous variables with normal distribution. Patient characteristics associated with aBMD < -1.0 10 years after RYGB were studied with univariate regression. Any variable associated with $p \leq 0.25$ from the univariable analysis were entered into a multivariable logistic regression model using a manual backward stepwise elimination procedure. In total, 8 potential predictors were examined (p-values written in bold). * Δ BMI not included in multivariable analysis due to high degree of association with BMI < 35 kg/m² at 10 year.

Abbreviations; Body mass index (BMI), carboxyl terminal telopeptide of type 1 collagen (CTX-1), procollagen type 1 N-terminal propeptide (PINP), bone specific alkaline phosphatase (BALP).

Missing: CTX-1 (4), PINP (5), BALP (7), Osteocalcin (10).

Definitions: Postmenopausal; follicular stimulating hormone > 25 IU/L, 25(OH) vitamin D insufficiency; serum 25(OH) vitamin D < 75.0 nmol/L.

Postmenopausal females are compared to premenopausal females and men.

only independent risk factor for having aBMD below the expected range for age, osteoporosis or a fragility fracture, stressing the importance of screening older subjects in a bariatric population.

4.4. Fractures

Emerging data indicate that obesity is associated with a site specific

increased fracture rate [43,44] despite a normal aBMD [45], implicating that obesity is associated with reduced bone quality. A study comparing subjects pursuing bariatric surgery to obese controls revealed an increased fracture rate in the bariatric population, prior to the surgical intervention [46]. To what extent the RYGB surgery may have contributed to the high fracture rate noted cannot be fully answered by our study. However, two years or more after RYGB patients have a 43%

Fig. 3. Percent of expected areal bone mineral density in 122 patients 10 years after Roux-en-Y gastric bypass when matched with a reference population (NHANES and Lunar given by manufacturer GE Lunar) in regard to age, gender and ethnicity.

Fig. 4. Inverse association between parathyroid hormone and areal bone mineral density (aBMD) z-scores of lumbar spine ($\beta -0.38$ 95% CI $(-0.12$ to $0.044)$, $p = 0.36$), femoral neck ($\beta -0.051$ 95% CI $(-0.099$ to $-0.002)$, $p = 0.040$) and total hip ($\beta -0.054$ 95%CI $(-0.11$ to $0.005)$, $p = 0.072$) in 122 participants 10 years after Roux-en-Y gastric bypass.

higher risk of skeletal fractures compared to subjects after adjustable gastric banding surgery, supporting the notion that RYGB itself may pose a negative effect on bone health [10].

During follow-up 15% of our participants experienced clinical low-energy fractures, similar rates have been reported for subjects exposed

Table 3

Prevalence of areal bone mineral density below the expected range for age or osteoporosis assessed by dual energy X-ray absorptiometry in 122 patients 10 years after Roux-en-Y gastric bypass for subgroups based on menopausal status or age, subdivided for gender.

	Areal bone mineral density below the expected range for age in premenopausal females and males younger than 50 years of age n = 63						Osteoporosis in postmenopausal females and males 50 years or older n = 59					
	Total		Male		Female		Total		Male		Female	
	n	%	n	%	n	%	n	%	n	%	n	%
Any skeletal location	5/ 63	8%	1/ 12	8%	4/ 51	8%	16/ 59	27%	2/ 16	13%	14/ 43	33%
Lumbar spine ^a	3/ 63	5%	1/ 12	8%	2/ 51	4%	11/ 59	19%	1/ 16	6%	10/ 43	23%
Femoral neck	0/ 61	0%	0/ 11	0%	0/ 50	0%	6/ 57	9%	1/ 15	7%	5/ 42	12%
Total hip	2/ 61	3%	0/ 11	0%	2/ 50	4%	5/ 57	9%	0/ 15	0%	5/ 42	12%

Premenopausal females and males younger than 50 years of age: Areal bone mineral density below expected range for age; z-score -2.0 or lower.

Postmenopausal females and males 50 years or older: Osteoporosis; t-score -2.5 or lower.

Missing: Femoral neck and total hip; total n = 4, male younger than 50 years of age (1), male 50 years or older (1), premenopausal female (1), postmenopausal female (1).

^a Vertebrae with osteoarthritic changes (spondylosis) or compression fractures were excluded from calculation.

to long-term glucocorticoid treatment [47]. In addition 11% of male participants and 7.4% of females had a moderate or severe morphometric vertebral fracture detected by vertebral fracture assessments of DXA. These rates are higher compared to prevalence of 7.5% in males and 3% in females reported in the general Norwegian population (38–59 years) [48]. The presence of morphometric vertebral fractures is a known risk factor for future fracture [49], yielding clinical relevance to our finding. Median duration to low energy fracture post RYGB was more than eight years. Together, our findings indicate an increased risk of fracture 10 years after RYGB and support the need for long term follow-up focusing on bone health.

4.5. Strengths and limitations

Strengths of our study include the large and well defined cohort, the comprehensive clinical examination during individual consultations, clinical interviews according to predefined case report forms, DXA scans and retrieval of bone turnover markers of all participants. The single center design may be a strength of the study as all patients received the same type of surgery, supplementation, counseling, and follow up, but this design may also limit the external validity of our findings.

The study is limited by the lack of preoperative bone specific medical history, bone turnover markers and DXA measurements. Possible effects of changes in neuro-hormonal axis on bone health were not evaluated. A total of 37% of potential participants were lost to follow-up. In the letter inviting patients to participate in the study we informed that the 10 year follow-up would include evaluation of bone health. This might have influenced subjects suspecting bone disease to participate in the study. The introduction of a new DXA machine during the study could have affected the aBMD results. Of the 122 participants, 51 had their examination with the GE Lunar Prodigy and 71 with the GE Lunar iDXA, but proper cross calibration was performed during transition from one machine to another. Additionally, the definition of aBMD z-score of -1.1 to -1.9 as aBMD in the lower range of normal

Table 4

Prevalence of low-energy fractures acquired during 10 years of follow up after Roux-en-Y gastric bypass (RYGB) in 122 participants.

Fracture site	Premenopausal females and males younger than 50 years of age n = 63			Postmenopausal females and males 50 years or older n = 59		
	aBMD higher range of normal n = 39	aBMD in the lower range of normal n = 20	aBMD below expected range for age n = 4	Normal aBMD n = 13	Osteopenia n = 30	Osteoporosis n = 16
Spine	1	1	0	–0	–0	2
Femoral neck/hip	–0	–0	–0	–0	1	1
Rib	–0	–0	–0	–0	1	1
Upper limb	1	–0	–0	–0	–0	2
Lower limb	2	–0	1	1	1	–0
More than one location	1	–0	–0	1	–0	–0
Proportion of participants with clinical low energy fracture after RYGB	5 (13%)	1 (5%)	1 (25%)	2 (15%)	3 (10%)	6 (38%)
Morphometric fractures of spine	2 (%)	3 (15%)			3 (10%)	2 (13%)

Definitions: aBMD: areal bone mineral density.

Premenopausal females and males younger than 50 years of age.

aBMD in the higher range of normal: 1.0 standard deviations lower than the expected aBMD of age, gender and ethnicity or higher (z-score > –1.0).*aBMD in the lower range of normal*: 1.1 to 1.9 standard deviations lower than the expected aBMD of age, gender and ethnicity (z-score < –1.0 to > –2.0).*aBMD below expected range for age*: 2.0 standard deviations lower than the expected aBMD of age, gender and ethnicity (z-score ≤ –2.0).

Postmenopausal females and males 50 years or older:

Normal aBMD: 1.0 standard deviations lower than the peak bone mass of young women or higher (t-score > –1.0).*Osteopenia*: 1.1 to 2.4 standard deviations lower than the peak bone mass of young women (t-score < –1.0 to > –2.5).*Osteoporosis*: aBMD 2.5 standard deviations or more lower than the peak bone mass of young women (t-score ≤ –2.5).

Upper limb fractures: Distal radius (n = 1), humerus (n = 2).

Lower limb fractures: Proximal tibia (n = 1), ankle (n = 1), tarsal (n = 1), metatarsal (n = 2).

More than one location: Ankle and distal radius (n = 1), carpal bone and costa (n = 1).

Morphometric fractures of spine revealed by vertebral fracture assessments by DXA.

was defined by the authors and is not a part of ICDS guidelines. This new definition may have inflated the number of patients considered to have abnormal aBMD. Lastly, information concerning etiology of hypothyroidism and repeated TSH measurements could have further elaborated the association between an aBMD z-score or t-score of –1.1 or lower 10 years after RYGB and preoperative hypothyroidism.

5. Conclusion

Ten years after RYGB 27% of postmenopausal females and males 50 years or older were osteoporotic, and 8% of premenopausal females and males 49 years or younger exhibited aBMD below the expected range for age. The prevalence of low-energy fractures was high. Hypothyroidism prior to, or SHPT, higher PINP levels, older age or postmenopausal status at 10 year follow-up were all risk factors for aBMD z-score or t-score of –1.1 or lower 10 years after RYGB.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bone.2019.07.014>.

Authors' roles

Conceptualization; IKBH, TM, SH, MCK, and JAK. Data curation; SH, IKBH and MCK. Formal analysis; IKBH and CB. Funding acquisition; EFE. Investigation; IKBH. Methodology; IKBH. Project administration; MCK, TM, SH, JAK, IKBH and HLG. Supervision; TM, JAK, EFE, HLG, Visualization; IKBH, Writing - original draft; Writing; IKBH, review & editing; IKBH, SH, MCK, HLG, EFE, TM, JAK, CB.

Acknowledgments

We deeply appreciate the support from the staff at Department of Endocrinology, Preventive medicine and Morbid Obesity, Oslo University Hospital. Especially we are thankful for the support of Gøril Vinje, Åse Halsne, and Kristine Bech Holte. This work was supported by the South-Eastern Norway Regional Health Authority (grant number 2014/1403) and Oslo University Hospital.

References

- [1] L. Angrisani, A. Santonicola, P. Iovino, A. Vitiello, N. Zundel, H. Buchwald, N. Scopinaro, Bariatric surgery and endoluminal procedures: IFSO worldwide survey 2014, *Obes. Surg.* 27 (9) (2017) 2279–2289.
- [2] H. Buchwald, Y. Avidor, E. Braunwald, M.D. Jensen, W. Pories, K. Fahrback, K. Schoelles, Bariatric surgery: a systematic review and meta-analysis, *Jama* 292 (14) (2004) 1724–1737.
- [3] T.D. Adams, L.E. Davidson, S.E. Litwin, J. Kim, R.L. Kolotkin, M.N. Nanjee, J.M. Gutierrez, S.J. Frogley, A.R. Ibele, E.A. Brinton, P.N. Hopkins, R. McKinlay, S.C. Simper, S.C. Hunt, Weight and metabolic outcomes 12 years after gastric bypass, *N. Engl. J. Med.* 377 (12) (2017) 1143–1155.
- [4] T.D. Adams, R.E. Gress, S.C. Smith, R.C. Halverson, S.C. Simper, W.D. Rosamond, M.J. Lamonte, A.M. Stroup, S.C. Hunt, Long-term mortality after gastric bypass surgery, *N. Engl. J. Med.* 357 (8) (2007) 753–761.
- [5] C. Liu, D. Wu, J.F. Zhang, D. Xu, W.F. Xu, Y. Chen, B.Y. Liu, P. Li, L. Li, Changes in bone metabolism in morbidly obese patients after bariatric surgery: a meta-analysis, *Obes. Surg.* 26 (1) (2016) 91–97.
- [6] E.W. Yu, M.L. Boussein, M.S. Putman, E.L. Monis, A.E. Roy, J.S. Pratt, W.S. Butsch, J.S. Finkelstein, Two-year changes in bone density after roux-en-Y gastric bypass surgery, *J. Clin. Endocrinol. Metab.* 100 (4) (2015) 1452–1459.
- [7] A.L. Schafer, C.M. Weaver, D.M. Black, A.L. Wheeler, H. Chang, G.V. Szefc, L. Stewart, S.J. Rogers, J.T. Carter, A.M. Posselt, D.M. Shoback, D.E. Sellmeyer, Intestinal calcium absorption decreases dramatically after gastric bypass surgery despite optimization of vitamin D status, *J. Bone Min. Res.* 30 (8) (2015) 1377–1385.
- [8] M.T. Blom-Høgestøl, Kristinsson IK, Brunborg JA, H.L. Gulseth, E.F. Eriksen, Changes in Bone Quality after Roux-En-Y Gastric Bypass: A Prospective Cohort Study in Subjects with and without Type 2 Diabetes, (2019) (Submitted for publication in Bone).
- [9] K.G. Lindeman, L.B. Greenblatt, C. Rourke, M.L. Boussein, J.S. Finkelstein, E.W. Yu, Longitudinal 5-year evaluation of bone density and microarchitecture after roux-en-Y gastric bypass surgery, *J. Clin. Endocrinol. Metab.* 103 (11) (2018) 4104–4112.
- [10] E.W. Yu, M.P. Lee, J.E. Landon, K.G. Lindeman, S.C. Kim, Fracture risk after bariatric surgery: roux-en-Y gastric bypass versus adjustable gastric banding, *J. Bone Min. Res.* 32 (6) (2017) 1229–1236.
- [11] K.F. Axelsson, M. Werling, B. Eliasson, E. Szabo, I. Naslund, H. Wedel, D. Lundh, M. Lorentzon, Fracture risk after gastric bypass surgery: a retrospective cohort study, *J. Bone Min. Res.* 33 (12) (2018) 2122–2131.
- [12] S. Hewitt, E.T. Aasheim, T.T. Sovik, J. Jahnsen, J. Kristinsson, E.F. Eriksen, T. Mala, Relationships of serum 25-hydroxyvitamin D, ionized calcium and parathyroid hormone after obesity surgery, *Clin. Endocrinol.* 88 (3) (2018) 372–379.
- [13] C. Duran de Campos, L. Dalcaneale, D. Pajecchi, A.B. Garrido Jr., A. Halpern, Calcium intake and metabolic bone disease after eight years of Roux-en-Y gastric bypass, *Obes. Surg.* 18 (4) (2008) 386–390.
- [14] M.T. Ott, P. Fanti, H.H. Malluche, U.Y. Ryo, F.S. Whaley, W.E. Strodel, T.A. Colacchi, Biochemical evidence of metabolic bone disease in women following

- roux-Y gastric bypass for morbid obesity, *Obes. Surg.* 2 (4) (1992) 341–348.
- [15] H. Aftab, H. Rissstad, T.T. Sovik, T. Bernklev, S. Hewitt, J.A. Kristinsson, T. Mala, Five-year outcome after gastric bypass for morbid obesity in a Norwegian cohort, *Surg. Obes. Relat. Dis.* 10 (1) (2014) 71–78.
- [16] P.R. Schauer, S. Ikramuddin, G. Hamad, G.M. Eid, S. Mattar, D. Cottam, R. Ramanathan, W. Gourash, Laparoscopic gastric bypass surgery: current technique, *J. Laparoendosc. Adv. Surg. Tech. A* 13 (4) (2003) 229–239.
- [17] K. Hind, B. Oldroyd, J.G. Truscott, In vivo precision of the GE Lunar iDXA densitometer for the measurement of total-body, lumbar spine, and femoral bone mineral density in adults, *J. Clin. Densitom.* 13 (4) (2010) 413–417.
- [18] T.E. Carver, N.V. Christou, O. Court, H. Lemke, R.E. Andersen, In vivo precision of the GE lunar iDXA for the assessment of lumbar spine, total hip, femoral neck, and total body bone mineral density in severely obese patients, *J. Clin. Densitom.* 17 (1) (2014) 109–115.
- [19] M.T. Vietri, M. Sessa, P. Pilla, M. Misso, D. Di Troia, A. Sorriento, N. Parente, A.M. Molinari, M. Cioffi, Serum osteocalcin and parathyroid hormone in healthy children assessed with two new automated assays, *J. Pediatr. Endocrinol. Metab.* 19 (12) (2006) 1413–1419.
- [20] ISCD, Official Positions of the International Society for Clinical Densitometry, (2015).
- [21] Assessment of Fracture Risk and its Application to Screening for Postmenopausal Osteoporosis. Report of a WHO Study Group, World Health Organization Technical Report Series, vol. 843, (1994), pp. 1–129.
- [22] H.K. Genant, C. Cooper, G. Poor, I. Reid, G. Ehrlich, J. Kanis, B.E. Nordin, E. Barrett-Connor, D. Black, J.P. Bonjour, B. Dawson-Hughes, P.D. Delmas, J. Dequeker, S. Ragi Eis, C. Gennari, O. Johnell, C.C. Johnston Jr., E.M. Lau, U.A. Liberman, R. Lindsay, T.J. Martin, B. Masri, C.A. Mautalen, P.J. Meunier, N. Khaltaev, et al., Interim report and recommendations of the World Health Organization task-force for osteoporosis, *Osteoporos. Int.* 10 (4) (1999) 259–264.
- [23] M.F. Holick, N.C. Binkley, H.A. Bischoff-Ferrari, C.M. Gordon, D.A. Hanley, R.P. Heaney, M.H. Murad, C.M. Weaver, Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline, *J. Clin. Endocrinol. Metab.* 96 (7) (2011) 1911–1930.
- [24] J.I. Mechanick, A. Youdim, D.B. Jones, W.T. Garvey, D.L. Hurley, M.M. McMahon, L.J. Heinberg, R. Kushner, T.D. Adams, S. Shikora, J.B. Dixon, S. Brethauer, Clinical practice guidelines for the perioperative nutritional, metabolic, and nonsurgical support of the bariatric surgery patient–2013 update: cosponsored by American Association of Clinical Endocrinologists, the Obesity Society, and American Society for Metabolic & Bariatric Surgery, *Endocr. Pract.* 19 (2) (2013) 337–372.
- [25] S.D. Harlow, M. Gass, J.E. Hall, R. Lobo, P. Maki, R.W. Rebar, S. Sherman, P.M. Sluss, T.J. de Villiers, Executive summary of the stages of reproductive aging workshop + 10: addressing the unfinished agenda of staging reproductive aging, *J. Clin. Endocrinol. Metab.* 97 (4) (2012) 1159–1168.
- [26] P.E. Shrout, J.L. Fleiss, Intraclass correlations: uses in assessing rater reliability, *Psychol. Bull.* 86 (2) (1979) 420–428.
- [27] N. Emaus, T.K. Omsland, L.A. Ahmed, G. Grimnes, M. Sneve, G.K. Berntsen, Bone mineral density at the hip in Norwegian women and men—prevalence of osteoporosis depends on chosen references: the Tromsø study, *Eur. J. Epidemiol.* 24 (6) (2009) 321–328.
- [28] M.A. Bredella, L.B. Greenblatt, A. Ejazi, M. Torriani, E.W. Yu, Effects of Roux-en-Y gastric bypass and sleeve gastrectomy on bone mineral density and marrow adipose tissue, *Bone* 95 (2017) 85–90.
- [29] A.L. Schafer, G.J. Kazakia, E. Vittinghoff, L. Stewart, S.J. Rogers, T.Y. Kim, J.T. Carter, A.M. Posselt, C. Pasco, D.M. Shoback, D.M. Black, Effects of gastric bypass surgery on bone mass and microarchitecture occur early and particularly impact postmenopausal women, *J. Bone Miner. Res. Off. J. Am. Soc. Bone Miner. Res.* 33 (6) (2018) 975–986.
- [30] R. Civitelli, R. Armamento-Villareal, N. Napoli, Bone turnover markers: understanding their value in clinical trials and clinical practice, *Osteoporos. Int.* 20 (6) (2009) 843–851.
- [31] C.S. Riedt, R.E. Brodin, R.M. Sherrell, M.P. Field, S.A. Shapses, True fractional calcium absorption is decreased after Roux-en-Y gastric bypass surgery, *Obesity* (Silver Spring, Md.) 14 (11) (2006) 1940–1948.
- [32] D. Hofso, E.T. Aasheim, T.T. Sovik, G.S. Jakobsen, L.K. Johnson, R. Sandbu, A.T. Aas, J. Kristinsson, J. Hjelmestaeth, Follow-up after bariatric surgery, *Tidsskr. Nor. Laegeforen.* 131 (19) (2011) 1887–1892.
- [33] L. Busetto, D. Dicker, C. Azran, R.L. Batterham, N. Farpour-Lambert, M. Fried, J. Hjelmestaeth, J. Kinzl, D.R. Leitner, J.M. Makaronidis, K. Schindler, H. Toplak, V. Yumuk, Practical recommendations of the obesity management task force of the European Association for the Study of obesity for the post-bariatric surgery medical management, *Obes. Facts* 10 (6) (2017) 597–632.
- [34] K.K. Mahawar, K. Clare, M. O’Kane, Y. Graham, L. Callejas-Diaz, W.R.J. Carr, Patient perspectives on adherence with micronutrient supplementation after bariatric surgery, *Obes. Surg.* 29 (5) (2019) 1551–1556 May.
- [35] A.C. Modi, M.H. Zeller, S.A. Xanthakos, T.M. Jenkins, T.H. Inge, Adherence to vitamin supplementation following adolescent bariatric surgery, *Obesity* (Silver Spring, Md.) 21 (3) (2013) E190–E195.
- [36] J.H. Bassett, G.R. Williams, Role of thyroid hormones in skeletal development and bone maintenance, *Endocr. Rev.* 37 (2) (2016) 135–187.
- [37] E.F. Eriksen, L. Mosekilde, F. Melsen, Kinetics of trabecular bone resorption and formation in hypothyroidism: evidence for a positive balance per remodeling cycle, *Bone* 7 (2) (1986) 101–108.
- [38] E. Moser, T. Sikjaer, L. Mosekilde, L. Rejnmark, Bone indices in thyroidectomized patients on long-term substitution therapy with levothyroxine assessed by DXA and HR-pQCT, *J. Thyroid. Res.* 2015 (2015) 796871.
- [39] A.W. Kung, K.K. Pun, Bone mineral density in premenopausal women receiving long-term physiological doses of levothyroxine, *Jama* 265 (20) (1991) 2688–2691.
- [40] T.L. Paul, J. Kerrigan, A.M. Kelly, L.E. Braverman, D.T. Baran, Long-term L-thyroxine therapy is associated with decreased hip bone density in premenopausal women, *Jama* 259 (21) (1988) 3137–3141.
- [41] D.T. Felson, Y. Zhang, M.T. Hannan, J.J. Anderson, Effects of weight and body mass index on bone mineral density in men and women: the Framingham study, *J. Bone Miner. Res. Off. J. Am. Soc. Bone Miner. Res.* 8 (5) (1993) 567–573.
- [42] J.M. Hughes, M.A. Petit, Biological underpinnings of Frost’s mechanostat thresholds: the important role of osteocytes, *J. Musculoskelet. Neuronal Interact.* 10 (2) (2010) 128–135.
- [43] H. Johansson, J.A. Kanis, A. Oden, E. McCloskey, R.D. Chapurlat, C. Christiansen, S.R. Cummings, A. Diez-Perez, J.A. Eisman, S. Fujiwara, C.C. Gluer, D. Goltzman, D. Hans, K.T. Khaw, M.A. Krieg, H. Kroger, A.Z. LaCroix, E. Lau, W.D. Leslie, D. Mellstrom, L.J. Melton 3rd, T.W. O’Neill, J.A. Pasco, J.C. Prior, D.M. Reid, F. Rivadeneira, T. van Staa, N. Yoshimura, M.C. Zillikens, A meta-analysis of the association of fracture risk and body mass index in women, *J. Bone Miner. Res.* 29 (1) (2014) 223–233.
- [44] D. Bergkvist, K. Hekmat, T. Svensson, L. Dahlberg, Obesity in orthopedic patients, *Surg. Obes. Relat. Dis.* 5 (6) (2009) 670–672.
- [45] W.D. Leslie, S.N. Morin, S.R. Majumdar, L.M. Lix, Effects of obesity and diabetes on rate of bone density loss, osteoporosis international: a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA 29 (1) (2018) 61–67.
- [46] C. Rousseau, S. Jean, P. Gamache, S. Lebel, F. Mac-Way, L. Biertho, L. Michou, C. Gagnon, Change in fracture risk and fracture pattern after bariatric surgery: nested case-control study, *BMJ* 354 (2016) i3794.
- [47] J.R. Curtis, A.O. Westfall, J. Allison, J.W. Bijlsma, A. Freeman, V. George, S.H. Kovac, C.M. Spettell, K.G. Saag, Population-based assessment of adverse events associated with long-term glucocorticoid use, *Arthritis Rheum.* 55 (3) (2006) 420–426.
- [48] S. Waterloo, L.A. Ahmed, J.R. Center, J.A. Eisman, B. Morseth, N.D. Nguyen, T. Nguyen, A.J. Sogaard, N. Emaus, Prevalence of vertebral fractures in women and men in the population-based Tromsø study, *BMC Musculoskelet. Disord.* 13 (2012) 3.
- [49] P. Chen, J.H. Krege, J.D. Adachi, J.C. Prior, A. Tenenhouse, J.P. Brown, E. Papadimitropoulos, N. Kreiger, W.P. Olszynski, R.G. Josse, D. Goltzman, Vertebral fracture status and the World Health Organization risk factors for predicting osteoporotic fracture risk, *J. Bone Miner. Res.* 24 (3) (2009) 495–502.