1 *Pancreatology (Print)* 2021; 21(3), pp. 530-543 (DOI: [10.1016/j.pan.2021.01.025](https://doi.org/10.1016/j.pan.2021.01.025))

2

3 **Gene expression profiling of morphological subtypes of pancreatic**
4 **ductal adenocarcinoma using surgical and EUS-FNB specimens**

5 Lukas Gammelgaard Rasmussen^{1,2,3}, Caroline Sophie Verbeke^{4,5}, Mia Dahl Sørensen^{1,3}, Per
6 Pfeiffer^{2,3,6}, Qihua Tan⁷, Michael Bau Mortensen^{2,3,8}, Claus Frstrup^{2,8}, Sönke Detlefsen^{1,2,3}

7 1: Department of Pathology, Odense University Hospital, Odense, Denmark

8 2: Odense Pancreas Center (OPAC), Odense University Hospital, Odense, Denmark

9 3: Department of Clinical Research, Faculty of Health Sciences, University of Southern Denmark,
10 Odense, Denmark

11 4: Department of Pathology, Oslo University Hospital, Oslo, Norway

12 5: Institute of Clinical Medicine, University of Oslo, Oslo, Norway

13 6: Department of Oncology, Odense University Hospital, Odense, Denmark

14 7: Epidemiology and Biostatistics, Institute of Public Health and Unit of Human Genetics, Clinical
15 Institute, Faculty of Health Science, University of Southern Denmark, Odense, Denmark

16 8: Department of Surgery, Upper GI and HPB Section, Odense University Hospital, Odense,
17 Denmark

18

19 **Corresponding author:**

20 Sönke Detlefsen

21 Department of Pathology, Odense University Hospital

22 J.B. Winsløvs Vej 15, 5000 Odense C, Denmark

23 E-mail: Sonke.Detlefsen@rsyd.dk

24 Phone: +45 6541 4806, Fax: +45 6591 2943

25

26 **Keywords:** pancreatic cancer; subtyping; gene expression analysis; immunohistochemistry;
27 differentiation grading

28 **Running head:** Gene expression analysis of pancreatic cancer

29 **Conflicts of interest:** The authors declare no conflict of interest.

30 **Funding:** This study was supported by funding from Odense Pancreas Center (OPAC) and
31 Guldsmed A. L. Rasmussen og Hustrus Mindefond (grant number A4001).

32 **Word count:** 4.997

1 **Abstract**

2 **Background/Objectives:** Various classifications of pancreatic ductal adenocarcinoma (PDAC)
3 based on RNA-profiling resulted in two main subtypes. Kalimuthu and coworkers proposed a
4 morphology-based classification that concurred with these subtypes. Immune therapy approaches
5 were so far disappointing in PDAC. Morphological PDAC subtypes may differ regarding key
6 immune-oncology pathways. We aimed to examine the reproducibility and prognostic value of
7 Kalimuthu's subtypes, and to evaluate differences regarding gene expression related to tumor
8 biology and immune-oncology.

9 **Methods:** 108 consecutive chemotherapy-naïve surgical specimens and 88 endoscopic ultrasound-
10 guided fine needle biopsies (EUS-FNBs) from 196 PDAC patients were evaluated by two
11 pancreatic pathologists as per Kalimuthu, resulting in Group A and Group B tumors. Digital mRNA
12 expression profiling was performed, on the surgical cohort using the NanoString IO360 panel of
13 770 key tumor biology related and 30 custom-genes, and on the EUS-FNB cohort using a targeted
14 panel of 123 genes.

15 **Results:** Morphological subtyping reached substantial agreement. In the surgical and EUS-FNB
16 cohorts, 44.4% and 38.6% were Group A, which was associated with improved survival.
17 Hierarchical clustering based on genes significantly different expressed in Group A and Group B
18 revealed clusters with prognostic value. One of these showed accumulation of Group A tumors and
19 upregulation of genes related to immune system and cytokine/chemokine/interleukin signaling.

20 **Conclusions:** Morphological subtyping according to Kalimuthu is reproducible and holds
21 prognostic value, in surgical as well as EUS-FNB specimens. As upregulation of immune
22 regulatory genes was found in Group A, future studies should evaluate the potential of immune
23 therapy in this particular subtype of PDAC.

1 **Introduction**

2 Pancreatic ductal adenocarcinoma (PDAC) is a highly lethal cancer with an overall 5-year survival
3 rate of 5% and median survival of 5.9 months [1]. PDAC has been estimated to be the third- and
4 second-leading cause of cancer-related death in the European Union and US by 2025 and 2030 [1].
5 In the last decade, subtyping of PDAC has been investigated in various studies, aiming to stratify
6 patients regarding treatment and outcome. One important obstacle to the management of PDAC is
7 the lack of predictive biomarkers for effective personalized treatment, including immunotherapy.

8 So far, classification of PDAC has been mainly based on transcriptional profiling, resulting in
9 several prognostically relevant classification systems [2-5]. Most studies have used RNA
10 sequencing (RNAseq) and RNA array techniques, which are time consuming and relatively difficult
11 to implement in clinical practice. Recently, Kalimuthu et al. proposed a histomorphology-based
12 classification of PDAC, the two subtypes of which showed considerable overlap with the two main
13 molecular subtypes as per Bailey, Collison, and Moffitt [6]. However, our knowledge regarding the
14 potential therapeutic implications of this morphological classification is very limited.

15 In contrast to other types of cancer, immunotherapeutic strategies for the treatment of PDAC have
16 so far not been successful [7-9]. It is tempting to speculate whether the morphological subtyping of
17 PDAC as per Kalimuthu may be able to help identify subsets of patients where targeted treatment,
18 such as immunotherapy, may be useful, but this has so far not been elucidated. A hurdle for the
19 implementation of a transcription-based classification in clinical practice is the fact that RNA
20 sequencing requires access to frozen tissue samples, which are more difficult to obtain than
21 formalin-fixed paraffin embedded (FFPE) tissues. Digital mRNA gene expression profiling may
22 overcome this hurdle, as it is specifically designed to detect RNA molecules in FFPE tissue
23 samples. Compared to RNA sequencing, it also has the advantage of measuring directly without the
24 need for prior amplification or cloning, thereby avoiding introduction of no gene-specific or 3'
25 biases. In addition, the reaction is driven to completion, allowing for a higher level of sensitivity
26 than in micro-arrays across many target genes with lower amounts of starting material. Moreover,
27 the system provides a digital readout of the amount of transcript in a sample. Finally, the time,
28 effort, and sample requirements of dedicated digital mRNA gene expression are more scalable than
29 real-time PCR or microarrays [10-12].

30 The aims of this study were to examine the reproducibility and prognostic value of the
31 morphological subtyping of PDAC as per Kalimuthu, using surgical specimens as well as
32 histological endoscopic ultrasound-guided fine-needle biopsies (EUS-FNBs). We further aimed to

- 1 evaluate whether the morphological subtypes differ regarding canonical pathways related to tumor
- 2 biology and key immune-oncology pathways.

3

1 **Materials and methods**

2 **Study cohort**

3 A series of 108 consecutive patients with chemotherapy-naïve PDAC who underwent surgical
4 resection at Odense University Hospital (OUH), Odense, Denmark, was identified by searching the
5 Danish Pathology Registry for the period 01.01.2015-30.04.2018. We also included a cohort of 88
6 patients who had been diagnosed with PDAC using EUS-FNB in the period from 01.01.2018–
7 31.12.2019. The inclusion criteria were a tissue area $\geq 8 \text{ mm}^2$ and a tumor/normal tissue ratio ≥ 55
8 %.

9 This study was approved by the Ethics Committee of the Region of Southern Denmark (project-ID:
10 S-20190175) and the Danish Data Protection Agency (project-ID 19/45478). Patients had not
11 advocated against the use of their tissue in the Danish registry for the use of tissue in research
12 ('Vævsanvendelsesregisteret').

13 **Digitalization of H&E-stained slides**

14 H&E-stained slides were scanned using a 20x objective on a NanoZoomer 2.0HT whole-slide
15 scanner (Hamamatsu Photonics, Hamamatsu, Japan). One of the pathologists (SDE) measured the
16 tumor-bearing areas on each slide, by marking them using the NDP.view 2.7.25 software
17 (Hamamatsu Photonics). The digitalized slides were uploaded to a pseudonymized web library,
18 with personalized access.

19 **Morphological subtyping according to Kalimuthu et al.**

20 All H&E-stained slides with tumor tissue from the 108 surgical PDAC and 88 EUS-FNB specimens
21 were re-evaluated by two pancreatic pathologists, referred to as observer A and observer B, and
22 subtyped as described by Kalimuthu et al. [6]. This classification recognizes four different
23 morphological patterns (conventional, tubulopapillary, composite, and squamous), giving rise to
24 two different morphological subtypes (Group A and Group B).

25 The conventional morphological pattern shows well-differentiated glands with a tubular, stellate
26 configuration. The tubulopapillary morphological pattern shows rounded and dilated glands that are
27 generally larger and often lined by a combination of foveolar-gastric type and pancreaticobiliary-
28 type epithelium. The composite morphological pattern shows a range of morphological features
29 traditionally associated with poor differentiation, such as angulated glands, cribriform complexes,

1 sheets, nests/islands, ribbons, cords, angulated glands, or scattered buds and single cells. The
2 squamous morphological pattern consists of well-to-poorly differentiated squamous structures. The
3 conventional and tubulopapillary patterns are merged into Group A and, correspondingly, given the
4 lack of well-formed glands, the composite and squamous patterns are merged to represent Group B
5 [6].

6 Each resection specimen with tumor was semi-quantitatively assessed for the presence of the four
7 patterns as follows: The area of tumor-containing tissue on each slide was divided into four
8 quadrants. For each quadrant, the dominant pattern was identified. Subsequently, the percentage of
9 quadrants dominated by each of the four morphological patterns was calculated [6]. Group A is
10 defined as dominance of conventional and/or tubulopapillary pattern in at least 60% of the
11 quadrants. Group B is defined as dominance of the composite and/or squamous pattern in at least
12 40% of the quadrants. With other words, tumors showing a dominant conventional/tubulopapillary
13 pattern in 0% to 59% or 60% to 100% of the quadrants were categorized as Group A or Group B
14 [6].

15 **Tumor differentiation grading**

16 The grade of histological differentiation was assessed for all resection specimens by both
17 pathologists based on the grading system recommended by the WHO [13, 14]. The tumor grade was
18 determined by the lowest grade. For the purpose of this study, it was decided that this lowest grade
19 had be present in at least 20% of a given tumor.

20 **RNA extraction**

21 For the surgical cohort, the FFPE block most representative of the morphological subtype was
22 selected for RNA isolation and macrodissected, for removal of non-tumor tissue. Sections of 10 μm
23 thickness were cut, their number depending on the size of the tumor area on the slide. At least 150
24 mm^2 total tumor area was included from all tumors/EUS-FNBs. The sections were prepared and
25 processed according to the Prosigna® Breast Cancer Prognostic Gene Signature Assay package
26 insert (2016-09 LBL-C0223-06). Total RNA was extracted using High Pure FFPE RNA Isolation
27 kit (Roche Diagnostics GmbH, Mannheim, Germany, 06650775001), according to the Prosigna®
28 protocol. RNA concentration was measured by Thermo Scientific™ NanoDrop™ One^C
29 Spectrophotometer (Thermo Scientific) to meet the quality control of RNA concentration (≥ 12.5
30 $\text{ng}/\mu\text{L}$) and purity (A260/A280 ratio between 1.7 and 2.3). If the RNA concentration exceeded 200

1 ng/ μ L, they were diluted with molecular grade RNase- and DNase-free water (Sigma-Aldrich,
2 7732-18-5) prior to downstream hybridization assay according to NanoString protocol. The RNA
3 samples were stored at -80°C.

4 **Digital mRNA expression profiling and data processing**

5 For the surgical cohort, mRNA gene expression levels were assessed using 100 ng RNA and the
6 PanCancer immune-profiling panel, Immuno-Oncology 360 (IO-360, NanoString Technologies,
7 Seattle, WA), designed to give a unique 360 degree view of gene expression in the tumors. The
8 panel consists of 770 genes (hereof, 20 housekeeping genes), falling into eight functional
9 categories: tumor immunogenicity, tumor sensitivity to immune attack, inhibitory immune
10 mechanisms, stromal factors, inhibitory metabolism, anti-tumor immune activity, inhibitory
11 immune signaling, and immune cell population abundance. Thirty custom genes were added, related
12 to fibroblasts, endothelial cells, and extracellular matrix (ECM), resulting in a total of 800 genes:
13 *ACTA2*, *ANO1*, *CALD1*, *CD34*, *CEACAM5*, *COL3A1*, *COL4A1*, *CYGB*, *FNI*, *GPC1*, *HAS2*, *INS*,
14 *KCNH2*, *KRT7*, *KRT8*, *LGALS1*, *MME*, *MUC1*, *NES*, *PDPN*, *POSTN*, *PRSSI*, *S100A4*, *SLC16A3*,
15 *SMAD4*, *SPARC*, *SYP*, *TNC*, *VCL* and *VIM* [15]. *CALD1*, *INS*, *KRT8*, *PRSSI*, and *SYP* were not
16 included in the hierarchical clustering analysis.

17 Normalization of RNA was performed using the geometric mean of internal negative controls,
18 positive controls, and 20 housekeeping genes. RNA samples were aligned with a synthetic panel
19 standard and hybridized in a 12-strip tube well with gene-specific probes (reporter and capture)
20 following the manufacturer's protocol. Subsequently, the target-probe complexes were read,
21 counted, and processed within the Counter Digital Analyzer (NanoString Technologies, Seattle,
22 WA) [11]. The raw digital counts of expression were exported to the nSolver v4.0 software
23 (NanoString) for downstream analysis following the manufacturer's protocol. The raw data from
24 the NanoString nCounter Digital Analyser were imported to the nSolver software, where the data
25 was investigated to check quality of reads, according to manufacturer's instructions. The data was
26 normalized and grouped according to the morphological subtype. Differential gene expression was
27 analyzed, where the normalized data was used to calculate the false discovery rate (FDR) for each
28 gene. The genes showing significantly different expression between Group A and Group B were
29 used to create a heat map with hierarchical clustering data, and the expression of these genes and of
30 Moffitt's top 50 basal-like and classical genes was examined in the EUS-FNB cohort [4].

1 **Pathway analysis**

2 The significant differently expressed genes (FDR<0.05) identified in the surgical cohort were
3 evaluated for over-representation of gene-sets or pathways in the Molecular Signatures Database
4 (MSigDB). The over-representation analysis compares a reference set of genes to a test gene-set
5 using the hypergeometric test. The probability of finding the number of significant genes belonging
6 to a particular gene-set or pathway can be calculated using hypergeometric distribution. The
7 analysis was performed using the analytical tool provided by Gene Set Enrichment Analysis
8 (GSEA) (<http://www.broadinstitute.org/gsea/index.jsp>) [16, 17].

9 **Tissue microarrays and immunohistochemistry (IHC)**

10 Multi-punch tissue microarrays (TMA) were used to validate the level of four representative
11 proteins encoded by differently expressed genes. The TMAs had been produced in a previous study,
12 as described elsewhere [18]. Included in the IHC analysis were 64/108 randomly chosen PDACs
13 (32 each of Group A and B). TMA sections were immunostained for CD5, CD10, FOXP3, and
14 TNC using the BenchMark Ultra immunostainer (Ventana Medical Systems, Tucson, AZ).
15 Antibody specifications, dilutions, incubation times, and epitope retrieval procedures are specified
16 in [Supplementary Table 1](#). Stained slides were scanned using a 40x objective on a Hamamatsu
17 NanoZoomer 2.0-HT whole slide scanner (Hamamatsu Photonics, Hamamatsu, Japan). Automated
18 digital image analysis was performed using the Visiopharm Image Analysis Software, version
19 2020.01 (Hoersholm, Denmark) as previously described [15].

20 **Statistical analysis**

21 The interobserver agreement of pattern assignment, subtype classification, and WHO grading was
22 investigated by utilizing Krippendorff's alpha, according to Landis J.R. and Koch G.G [19], which
23 calculates coefficients 0 (or <0 in extreme cases) to 1. A coefficient 0 indicates no agreement, while
24 coefficient 1 represents perfect agreement. The coefficient values are interpreted as follows: <0:
25 poor/systematic disagreement, 0 - 0.2: slight agreement, 0.21 - 0.40: fair agreement, 0.41 - 0.60:
26 moderate agreement, 0.61 - 0.80: substantial agreement, and 0.81 - 1.0: almost perfect agreement
27 [19]. Survival was analyzed using Kaplan-Meier and Cox regression using gender, age, tumor stage,
28 and area of tumor as covariates. Survival was calculated from date of surgery or date of diagnosis.
29 Patients alive on February 1, 2020 (surgical cohort) or December 1, 2020 (EUS-FNB cohort) were

1 censored at this date. Median survival and categorical data are presented with 95% confidence
2 intervals. A *P*-value < 0.05 was considered significant. Analysis was made using Stata v. 16,
3 StataCorp, USA.

4 The normalized gene expression data were first transformed by taking the logarithm with base 2.
5 The log transformation has the advantage of making the data approach a normal distribution.
6 Association between gene expression and subtypes was investigated using the binomial logistic
7 regression to the log odds of subtype as a function of gene expression. Significance of the gene
8 expression was determined by the likelihood ratio test comparing the likelihood of the current
9 model with that of the reduced model without gene expression variable, with the test statistic
10 distributed as a chi-squared random variable with 1 degree of freedom. The logistic regression
11 analysis was performed by the glm function in R (<https://www.r-project.org>) with a binomial link
12 function. To adjust for the statistical significance due to multiple testing, a false discovery rate
13 (FDR) was calculated for each gene tested using the Benjamini and Hochberg method [20].
14 Statistical significance was assigned to genes with FDR<0.05.

15 The two-tailed Mann-Whitney test was used to compare the immunohistochemical expression of
16 the selected markers in the gland-forming vs. non-gland-forming subtypes, and scatter plots were
17 generated in GraphPad Prism, version 5.01.

18 **Results**

19 **Clinical data**

20 Characteristics of the patients included in the surgical cohort of 108 PDACs and general
21 pathological findings are presented in [Table 1](#).

22 **Morphological subtyping as per Kalimuthu et al.**

23 In average, 7.50 ± 3.40 (range 2-23) slides containing tumor tissue were included in the
24 morphological scoring. The average tumor-containing area per slide was $94.26 \pm 87.88 \text{ mm}^2$ (range
25 $0.24\text{--}744.3 \text{ mm}^2$). There was substantial agreement between both observers regarding
26 morphological subtypes (Krippendorff's alpha coefficient 0.68 ± 0.07 ; 95%CI 0.54-0.82). For
27 assignment of the dominant morphological patterns, the agreement was moderate (Krippendorff's
28 alpha coefficient 0.58 ± 0.07 ; 95%CI 0.44-0.71). Considering that agreement was substantial for the

subtype classification, further analyses were conducted based on observer A's scoring, as decided during the design of this study.

Observer A classified 48/108 of the tumors as Group A and 60/108 as Group B. Obviation of slides that contained very small amounts of tumour (less than 20 mm² and 30 mm², respectively) resulted in minimal changes in the ratio between Group A and Group B tumors (Supplementary Table 2). A cut-off of 20 mm² and 30 mm² resulted in the change of 2 tumors from Group A to Group B and of 3 tumors from Group B to Group A and the Group A/Group B ratio was 49/59 for both cut-off values. The dominant morphological pattern was conventional in 46/108 (42.59%), tubulopapillary in 10/108 (9.26%), composite in 46/108 (42.59%), and squamous in 6/108 (5.56%). Morphological heterogeneity was observed, as two or more patterns were present in 95/108 (87.96%) (Fig. 1A). Thirteen of 108 tumors (12.04%) showed exclusively one morphological pattern. These patterns were conventional (n=2), tubulopapillary (n=1), and composite (n=10). In some instances, small cancer cell clusters or abortive glands were associated with larger glandular formations but were less conspicuous than the latter (Fig. 1B-E).

WHO grading

Regarding the grade of differentiation, interobserver agreement was substantial, with a Krippendorff's alpha coefficient of 0.78±0.05 (95% CI 0.68-0.89). Given this result, further (survival) analyses were based on observer A's grading, as decided during the design of this study. Twenty of 108 (18.52%) tumors were well-differentiated (G1), 38/108 (35.18%) moderately differentiated (G2), and 50/108 (46.30%) poorly differentiated (G3).

Survival analysis

The median survival for the entire surgical cohort was 18.83±1.56 months (95% CI 15.31-25.82) (Fig. 1F). At the time of survival analysis, 87/108 patients (80.56%) had died. Group A (n=48) and Group B (n=60) showed a median survival of 25.82±8.65 months (95% CI 15.34-38.28) and 17.15±2.84 months (95% CI 12.75-19.78). There was a significantly longer survival of patients in Group A compared to Group B ($P = 0.026$, Fig 1G). Median survival associated with the four morphological patterns is given in Figure 1H. The squamous pattern was associated with a significantly worse survival ($P = 0.020$), and there was a trend towards poorer outcome for the composite pattern ($P = 0.052$) (Fig. 1H). Median survival based on the WHO differentiation

grading is shown in Fig. 1I: Tumors showing G3 differentiation had a significantly shorter survival than G1 ($P = 0.044$), whereas G2 tumors did not differ significantly ($P = 0.71$) (Fig. 1I).

Gene expression profiling

In the surgical cohort, 53 genes were significantly different expressed in Group A versus Group B ($FDR < 0.05$) (Supplementary Table 3). Thirty-nine genes were upregulated in Group A: *CCR2*, *CD2*, *CD27*, *CD40LG*, *CD5*, *CD6*, *CD74*, *CX3CR1*, *FOXP3*, *FUT4*, *GLI1*, *ICOSLG*, *IL10RA*, *IL22RA1*, *JAK2*, *KLRB1*, *LY9*, *MMP1*, *MMP7*, *NLRC5*, *PDZK1IP1*, *PIK3CG*, *PRKACB*, *PRLR*, *PSMB10*, *P2RY13*, *RASAL1*, *RIPK3*, *RORC*, *SHC2*, *SLAMF7*, *SYK*, *TICAM1*, *TLR3*, *TNFRSF11A*, *TNFRSF17*, *TRAF1*, *WNT2B*, and *ZAP70*. Fourteen genes were upregulated in Group B: *ANGPTL4*, *ANLN*, *BIRC5*, *CCNB1*, *CDK2*, *ENO1*, *FLNB*, *IL11*, *MME*, *NT5E*, *PLOD2*, *PTGS2*, *TNC*, and *UBE2C*.

Expression of these 53 differentially expressed genes served as the basis for hierarchical clustering, resulting in a heat map, where four clusters could be identified (Fig. 2A). Cluster 1 contained 64 cases, 43 (67.19%) of which were Group A. Cluster 2 contained 27 tumors, 22 (81.48%) of which were Group B. Clusters 3 and 4 contained 10 and 7 tumors, all of which (100%) were in Group B. Six tumors (case 6, 60, 64, 73, 93, and 107) were in Group B with a dominant squamous pattern, three of which were located in cluster 4 (42.86%, 3/7) and one case each in clusters 1 (1.56%), 2 (3.70%), and 3 (10.00%). Cluster 1 and 3 had the best survival, whereas cluster 2 and cluster 4 had significantly poorer survival, with hazard ratios of 2.3 and 12 (Fig. 2B-C).

GSEA analysis identified the top-20 canonical pathways for each morphological subtype (Table 2). Group A showed higher expression of immune-related genes and cytokine/chemokine/interleukin signaling, whereas Group B showed higher expression of genes related to cancer cell proliferation and cell cycle regulation.

Microsatellite deficiency

The tumors included in the present study were also evaluated in a previous study, revealing that one tumor was microsatellite instable (case 33) and another showed loss mismatch repair protein MSH6 (case 96) [18]. All other tumors were microsatellite stable. Case 33 was assigned to cluster 2 and case 96 to cluster 1.

1 **Immunohistochemical protein expression**

2 CD5 protein was expressed significantly higher in the Group A than Group B, in accordance with
3 the corresponding gene expression levels ($P = 0.037$, Fig. 3A-B). There was a similar trend for
4 FOXP3 ($P = 0.12$, Fig. 3C-D). Likewise, the protein expression of both CD10 ($P = 0.042$) (Fig. 3E-
5 F) and TNC ($P < 0.0001$) (Fig. 3G-H) was significantly higher in the Group B than in Group A.

6 **Morphological subtyping, survival analysis and gene expression profiling in 88 EUS-FNBs** 7 **from patients with pancreatic ductal adenocarcinoma**

8 Eighty-eight EUS-FNBs were included, where 46/88 (52.3%) were from males, mean age 73 years
9 (range 53-88). There was substantial agreement between both observers regarding morphological
10 subtypes (Krippendorff's alpha coefficient 0.79 ± 0.06 ; 95%CI 0.66-0.92). Observer A classified
11 34/88 (38.6%) and 54/88 (62.4%) of the tumors as Group A and B (Fig. 4A). The survival analysis
12 showed an improved survival for Group A vs. Group B (Fig. 4B). Twenty of the top 53 genes
13 identified in the surgical cohort were significantly different expressed in Group A vs. Group B.
14 Hierarchical clustering of these genes identified gene clusters similar to those in the surgical cohort
15 (Fig. 4C-D), albeit with different relative frequencies, probably due to the clinically very different
16 populations studied. Figure 4E illustrates the survival analysis regarding the clusters in the EUS-
17 FNB cohort.

18 When we examined Moffitt's top 50 genes for the classical and basal-like subtypes of PDAC, 26 of
19 these genes were significantly different expressed between Group A and Group B. Hierarchical
20 clustering showed that Group A tumors were more often classical (28/34, 82%) and that Group B
21 tumors were more often basal-like (35/54, 65%) (Fig. 5A). Of the 41 basal-like tumors, 35 (85%)
22 were Group B. Of the 47 classical tumors, 28 (60%) were Group A. A survival analysis showed a
23 tendency towards poorer survival for the basal-like tumors ($P=0.026$) (Fig. 5B).

1 **Discussion**

2 In the present study, we were able to reproduce the morphological subtyping of PDAC, as per
3 Kalimuthu and coworkers, with substantial agreement between two pancreatic pathologists, in
4 surgical and EUS-FNB specimens. Our data confirm that the morphological subtyping holds
5 prognostic value. Using digital RNA expression analysis, we found that tumors categorized in
6 Group A had a higher expression of genes related to the immune system and
7 cytokine/chemokine/interleukin signaling, whereas Group B had a higher expression of genes
8 related to cancer cell proliferation and cell cycle regulation. Hence, it may be hypothesized that
9 immune therapeutic approaches may be more promising in some patients with the Group A subtype
10 of PDAC.

11 This appears to be the first study testing the reproducibility of the histology-based classification of
12 PDAC as per Kalimuthu et al in surgical specimens, and the first study using this approach in EUS-
13 FNBs. In the surgical cohort, we identified a slightly higher relative number of tumors located to
14 Group B (n=60, 55.6%), compared to the original report (n=37, 43%) [6]. A possible explanation
15 could be the selection of included tumors. IPMN-associated PDACs, for example, were excluded
16 from the current study, in contrast to Kalimuthu et al [6]. There may also be differences between the
17 populations from which the cases were selected. The interobserver agreement was substantial,
18 indicating the robustness and applicability of the morphological criteria that define the patterns and
19 subtypes. Regarding the grade of histological tumor differentiation, we found a higher rate of G3
20 tumors (46.3%), in contrast to Kalimuthu et al. (18.6%), but in agreement with other studies [6, 21,
21 22]. In our cohort, more than 15% were G1, in agreement with Kalimuthu et al. and Moffitt et al. [4,
22 6]. Given the histomorphological criteria that define both subtypes of the classification, low-grade
23 PDACs are likely to be assigned to the gland-forming subtype, and high-grade PDACs to the non-
24 gland-forming subtype. The co-variation between morphological and molecular subtypes, as
25 revealed by the Kalimuthu study, seems therefore to bring us a step closer to the molecular basis for
26 the morphological concept of grade of tumor differentiation, which hitherto has remained
27 unexplored [6].

28 Group A showed a better median overall survival than Group B. The squamous morphological
29 pattern conferred the worst prognosis (statistically significant) and was associated with a poorer
30 overall survival compared to the conventional pattern, while there was a trend towards poorer
31 prognosis for the composite pattern. Also Kalimuthu and coworkers reported a trend for difference

1 in outcome between both subtypes, a significant worse outcome for the composite pattern, and a
 2 trend towards worse outcome for the squamous pattern [6]. We found that poorly differentiated
 3 tumors (G3) were associated with a worse outcome. In the study conducted by Kalimuthu et al., the
 4 association of G3 with poor outcome did not reach full statistical significance[6].
 5 We used the morphological subtyping as per Kalimuthu as framework for hierarchical clustering-
 6 based gene expression analysis. Fifty-three significantly different expressed genes were identified
 7 in the surgical cohort. Ten of these 53 genes were also included in several gene sets that have been
 8 reported in earlier studies, which have investigated RNA-based PDAC classification [3, 4, 6, 23,
 9 24]: *ICOSLG*, *IL22RA1*, *MME*, *NT5E*, *PDZKIIP1*, *RASAL1*, *TNC*, *TNFRSF11A*, *ANGPTL4*, and
 10 *MMP7*. Thirty-nine genes were expressed at significantly higher levels in Group A, related to
 11 immune system and to cytokine/chemokine/interleukin signaling. Kalimuthu et al. demonstrated
 12 that Group A shared most commonalities with the classical subtype as described by Moffitt and
 13 Collison, respectively, and the pancreatic progenitor subtype proposed by Bailey et al. [3-5]. Our
 14 own data support this, as 82% of Group A tumors were classical. Interestingly, also Bailey et al.
 15 (immunogenic subtype) and Puleo et al. (immune classical subtype) identified a subgroup of PDAC
 16 that was characterized by high expression of genes related to immunosuppressive checkpoint
 17 pathways and gene signatures of multiple executors of immunity [2, 5]. Some these factors can
 18 potentially be targeted with immune modulators [2]. Despite the wide range of upregulated genes,
 19 only anti-PD-1 and anti-CTLA4 treatments have been tested in PDAC so far, with poor results [7-
 20 9]. In accord with this, we also found a trend for higher expression of *CTLA-4* in the group of
 21 gland-forming tumors. However, the relative proportion of the immunogenic and immune classical
 22 tumors in Bailey et al.'s and Puleo et al.'s studies were much lower than Group A in the present
 23 study [2, 5]. Le et al. observed that MSI PDAC was responsive to PD-1/PDL-1 antagonist treatment
 24 [7]. MSI has, however, a low frequency in PDAC with about 1-3%, which also was found in our
 25 cohort (2/108 (1.85%)) [18]. It may be hypothesized that immune therapeutic approaches may have
 26 potential in some patients, particularly in those with the Group A subtype. For example, we found
 27 that *FOXP3*, *ICOSLG*, *IL10RA*, *RORC*, *TICAM1*, and *TLR3* are expressed at higher levels in this
 28 subtype, and these genes were considered treatable targets [25-30]. *RORC* encodes the proteins
 29 ROR γ and ROR γ t, hormone receptors that upon stimulation function as transcription factors. Low
 30 expression of ROR γ has been associated with poorly differentiated tumors and negative outcome of
 31 aggressive basal-like breast cancer and bladder cancer, in agreement with our findings on PDAC
 32 [31-33]. Several studies indicate that targeting of ROR γ and ROR γ t may be promising as a new

1 immunotherapeutic approach [25, 31-33]. Cancer cells are able to activate TLR3 and TICAM-1 to
2 increase release of cytokines and chemokines that stimulate immune suppressive cells leading to
3 tumor progression [29, 30, 34]. A phase 1 study investigated a TLR3 agonist and dendritic cell-
4 based vaccine, and found an increased median survival of 2.8 months [35]. Future studies should
5 evaluate whether the individual RNA expression profile is able to predict response to certain
6 therapeutic approaches, for instance immune therapy. This could be based on digital mRNA
7 oncogenic profiling, the approach used in the present study, including an EUS-FNB cohort in
8 addition to the surgical cohort. Only one previous study used a similar approach, examining 13
9 EUS-FNBs with PDAC and identifying possible targets for combined immunomodulatory
10 therapeutics [12].

11 Fourteen genes were expressed significantly higher in Group B, related to cancer cell proliferation
12 and cell cycle regulation, in agreement with Kalimuthu et al., who reported that Group B was
13 similar to the subtypes known as basal-like, squamous or QM-PDA [3-6]. Also the present study
14 found that 85% of the basal-like tumors in the EUS-FNB cohort were Group B subtype. Moffitt et
15 al. reported that basal-like tumors had faster growth rates than the classical subtype [4]. Collisson et
16 al. reported a higher proliferation in QM-PDA tumors [3]. QM-PDA/squamous/basal-like PDACs
17 had a worse outcome compared to classical/pancreatic progenitor, immunogenic, ADEX/classical
18 subtypes [3-5]. Early results from the COMPASS trial suggest that first-line chemotherapy is
19 associated with significantly better outcome in the classical subtype, whereas basal-like tumors tend
20 to be more resistant [24]. We found *ANGPTL4*, *ENO1*, *PLOD*, *PTGS2*, *TNC*, and *UBE2C* among
21 the genes that were upregulated in Group B, some of which are under scrutiny for targeted therapy
22 [36-41].

23 While the high degree of interobserver agreement testifies to the applicability of the morphological
24 classification system proposed by Kalimuthu et al., it has several weaknesses. While the quadrant-
25 based approach considerably facilitates scoring, the fact that scoring is done irrespective of the
26 tumor area may be a limitation. The patterns in small tumor areas have a disproportionately strong
27 impact on the overall scoring of the tumor. In addition, the “size” of a tumor area in a section may
28 be entirely fortuitous, determined by the way the specimen slices were divided to fit into the tissue
29 cassettes. A small rim of tumor along the edge of tissue block has as much impact on the overall
30 score as a block consisting mainly of tumor. While identifying the dominant tumor pattern seems
31 fairly straightforward, several practical issues make this at times challenging. First, it is not clearly
32 defined whether “dominance” is determined by the area occupied by tumor of a certain pattern or

1 by the number of tumor cells. As cancer cell density often varies significantly from one spot to the
 2 next, it is difficult to compare the “burden” of tumor of various morphologies within a given
 3 quadrant. Furthermore, the presence of small cancer cell clusters or abortive glands less
 4 conspicuous than larger glandular formations are easily underestimated at low power. In order to
 5 identify all of them, one needs higher magnification, but then the quantification, *i.e.*, assessment of
 6 the proportion with respect to the remainder of the tumor, becomes difficult. It is tempting to
 7 speculate whether artificial intelligence could be used for this quantification in the future.

8 The Kalimuthu classification takes into account morphological heterogeneity by semiquantitative
 9 assessment of the different morphologies in a given tumor. By translating this into a binary system,
 10 the subtypes themselves do not reflect the wide morphological variety that exists in PDAC but serve
 11 as a framework that may ease the translation into clinical practice. The RNA-based classifications
 12 with discrete categories, including morphological subtyping, are probably also somewhat rigid, and
 13 one should probably rather think of a "molecular grading" system, with each PDAC having more or
 14 less of either subtypes (*i.e.* a sliding scale between the extremes of pure classical and pure basal-
 15 like) [42]. This approach is supported by Kalimuthu et al.’s and our own data, showing a high level
 16 of morphological heterogeneity within most of the PDACs. These findings are in line with studies
 17 based on single-cell sequencing, showing that basal-like and classical tumor cells frequently co-
 18 exist [23, 43].

19 Regarding the morphological subtyping as per Kalimuthu, it should be mentioned that the term
 20 “non-glandular” may appear a bit confusing, as tumors categorized as “composite” do often show
 21 lumina. A further challenge lies in the change in growth pattern that not uncommonly occurs when
 22 tumor cells invade particular microanatomic compartments. For example, a cancer of composite
 23 pattern in the bulk of the tumor mass, often assumes a conventional pattern when invading the
 24 duodenal muscle layer, a well-known phenomenon coined as "intestinal mimicry" [44]. Foci of so-
 25 called duct cancerization, that is, the growth of invasive PDAC along the wall of preexisting
 26 pancreatic ducts often present a cystic-papillary pattern, irrespective of the pattern in the remainder
 27 of the tumor mass. While these observations likely represent cancer “plasticity”, *i.e.* the reversible
 28 transition from one transcriptional subtype to the other, it is not clear whether this has the same
 29 oncological implications as the tumor phenotype outside these microcompartments.

30 In conclusion, our data indicate that the morphological classification as per Kalimuthu et al. is
 31 reproducible and holds prognostic value, in surgical as well as EUS-FNB specimens. Using digital
 32 mRNA expression profiling, we identified key canonical pathways that are differentially regulated

1 when comparing the two morphological subtypes of PDAC. These findings may have therapeutic
2 implications, as immunotherapeutic strategies may be more promising in at least some patients with
3 Group A subtype. Future studies should evaluate the potential of immune therapy with special
4 emphasis on this subtype of PDAC, and these could also include the use of EUS-FNB specimens.

References

- [1] Rahib, L., B.D. Smith, R. Aizenberg, A.B. Rosenzweig, J.M. Fleshman, and L.M. Matrisian, Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States. *Cancer Res*, 2014. 74(11): p. 2913-21.
- [2] Puleo, F., R. Nicolle, Y. Blum, J. Cros, L. Marisa, P. Demetter, et al., Stratification of Pancreatic Ductal Adenocarcinomas Based on Tumor and Microenvironment Features. *Gastroenterology*, 2018. 155(6): p. 1999-2013.e3.
- [3] Collisson, E.A., A. Sadanandam, P. Olson, W.J. Gibb, M. Truitt, S. Gu, et al., Subtypes of pancreatic ductal adenocarcinoma and their differing responses to therapy. *Nat Med*, 2011. 17(4): p. 500-3.
- [4] Moffitt, R.A., R. Marayati, E.L. Flate, K.E. Volmar, S.G. Loeza, K.A. Hoadley, et al., Virtual microdissection identifies distinct tumor- and stroma-specific subtypes of pancreatic ductal adenocarcinoma. *Nat Genet*, 2015. 47(10): p. 1168-78.
- [5] Bailey, P., D.K. Chang, K. Nones, A.L. Johns, A.M. Patch, M.C. Gingras, et al., Genomic analyses identify molecular subtypes of pancreatic cancer. *Nature*, 2016. 531(7592): p. 47-52.
- [6] Kalimuthu, S.N., G.W. Wilson, R.C. Grant, M. Seto, G. O'Kane, R. Vajpeyi, et al., Morphological classification of pancreatic ductal adenocarcinoma that predicts molecular subtypes and correlates with clinical outcome. *Gut*, 2020. 69(2): p. 317-28.
- [7] Le, D.T., J.N. Durham, K.N. Smith, H. Wang, B.R. Bartlett, L.K. Aulakh, et al., Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science*, 2017. 357(6349): p. 409-13.
- [8] Brahmer, J.R., S.S. Tykodi, L.Q. Chow, W.J. Hwu, S.L. Topalian, P. Hwu, et al., Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. *N Engl J Med*, 2012. 366(26): p. 2455-65.
- [9] Royal, R.E., C. Levy, K. Turner, A. Mathur, M. Hughes, U.S. Kammula, et al., Phase 2 trial of single agent Ipilimumab (anti-CTLA-4) for locally advanced or metastatic pancreatic adenocarcinoma. *J Immunother*, 2010. 33(8): p. 828-33.
- [10] Robinson, J.M., S. Turkington, S.A. Abey, N. Kenea, and W.A. Henderson, Differential gene expression and gene-set enrichment analysis in Caco-2 monolayers during a 30-day timeline with Dexamethasone exposure. *Tissue Barriers*, 2019. 7(3): p. e1651597.
- [11] Geiss, G.K., R.E. Bumgarner, B. Birditt, T. Dahl, N. Dowidar, D.L. Dunaway, et al., Direct multiplexed measurement of gene expression with color-coded probe pairs. *Nat Biotechnol*, 2008. 26(3): p. 317-25.
- [12] Gleeson, F.C., M.J. Levy, R.A. Jackson, S.J. Murphy, K.C. Halling, B.R. Kipp, et al., Endoscopic ultrasound may be used to deliver gene expression signatures using digital mRNA detection methods to immunophenotype pancreatic ductal adenocarcinoma to facilitate personalized immunotherapy. *Pancreatology*, 2020. 20(2): p. 229-38.
- [13] Lüttges, J., S. Schemm, I. Vogel, J. Hedderich, B. Kremer, and G. Klöppel, The grade of pancreatic ductal carcinoma is an independent prognostic factor and is superior to the immunohistochemical assessment of proliferation. *J.Pathol.*, 2000. 191(2): p. 154-61.
- [14] Hruban, R.H., V.N. Adsay, I. Esposito, N. Fukushima, T. Furukawa, G. Klöppel, et al., The 2019 WHO classification of tumours of the digestive system, in *Digestive System Tumours*, W.C.o.T.E. Board, Editor. 2019, IARC. p. 322-33.

- 1 [15] Nielsen, M.F.B., M.B. Mortensen, M.D. Sørensen, M. Wrenfeldt, B.W. Kristensen, H.D.
2 Schröder, et al., Spatial and phenotypic characterization of pancreatic cancer-associated
3 fibroblasts after neoadjuvant treatment. *Histology and histopathology*, 2020: p. 18201-01.
- 4 [16] Subramanian, A., P. Tamayo, V.K. Mootha, S. Mukherjee, B.L. Ebert, M.A. Gillette, et al.,
5 Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide
6 expression profiles. *Proceedings of the National Academy of Sciences*, 2005. 102(43): p.
7 15545.
- 8 [17] Mootha, V.K., C.M. Lindgren, K.-F. Eriksson, A. Subramanian, S. Sihag, J. Lehar, et al.,
9 PGC-1 α -responsive genes involved in oxidative phosphorylation are coordinately
10 downregulated in human diabetes. *Nature Genetics*, 2003. 34(3): p. 267-73.
- 11 [18] Ahmad-Nielsen, S.A., M.F. Bruun Nielsen, M.B. Mortensen, and S. Detlefsen, Frequency of
12 mismatch repair deficiency in pancreatic ductal adenocarcinoma. *Pathol Res Pract*, 2020.
13 216(6): p. 152985.
- 14 [19] Landis, J.R. and G.G. Koch, The measurement of observer agreement for categorical data.
15 *Biometrics*, 1977. 33(1): p. 159-74.
- 16 [20] Benjamini, Y. and Y. Hochberg, Controlling the False Discovery Rate: A Practical and
17 Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B*
18 (Methodological), 1995. 57(1): p. 289-300.
- 19 [21] Muckenhuber, A., A.K. Berger, A.M. Schlitter, K. Steiger, B. Konukiewicz, A. Trumpp, et
20 al., Pancreatic Ductal Adenocarcinoma Subtyping Using the Biomarkers Hepatocyte
21 Nuclear Factor-1A and Cytokeratin-81 Correlates with Outcome and Treatment Response.
22 *Clin Cancer Res*, 2018. 24(2): p. 351-59.
- 23 [22] Schlitter, A.M., A. Segler, K. Steiger, C.W. Michalski, C. Jager, B. Konukiewicz, et al.,
24 Molecular, morphological and survival analysis of 177 resected pancreatic ductal
25 adenocarcinomas (PDACs): Identification of prognostic subtypes. *Sci Rep*, 2017. 7: p.
26 41064.
- 27 [23] Chan-Seng-Yue, M., J.C. Kim, G.W. Wilson, K. Ng, E.F. Figueroa, G.M. O'Kane, et al.,
28 Transcription phenotypes of pancreatic cancer are driven by genomic events during tumor
29 evolution. *Nat Genet*, 2020. 52(2): p. 231-40.
- 30 [24] Aung, K.L., S.E. Fischer, R.E. Denroche, G.H. Jang, A. Dodd, S. Creighton, et al.,
31 Genomics-Driven Precision Medicine for Advanced Pancreatic Cancer: Early Results from
32 the COMPASS Trial. *Clin Cancer Res*, 2018. 24(6): p. 1344-54.
- 33 [25] Lytle, N.K., L.P. Ferguson, N. Rajbhandari, K. Gilroy, R.G. Fox, A. Deshpande, et al., A
34 Multiscale Map of the Stem Cell State in Pancreatic Adenocarcinoma. *Cell*, 2019. 177(3): p.
35 572-86.e22.
- 36 [26] Solinas, C., C. Gu-Trantien, and K. Willard-Gallo, The rationale behind targeting the ICOS-
37 ICOS ligand costimulatory pathway in cancer immunotherapy. *ESMO open*, 2020. 5(1): p.
38 e000544.
- 39 [27] Ruffell, B., D. Chang-Strachan, V. Chan, A. Rosenbusch, C.M.T. Ho, N. Pryer, et al.,
40 Macrophage IL-10 blocks CD8⁺ T cell-dependent responses to chemotherapy by
41 suppressing IL-12 expression in intratumoral dendritic cells. *Cancer cell*, 2014. 26(5): p.
42 623-37.
- 43 [28] Wang, X., X. Li, X. Wei, H. Jiang, C. Lan, S. Yang, et al., PD-L1 is a direct target of
44 cancer-FOXP3 in pancreatic ductal adenocarcinoma (PDAC), and combined
45 immunotherapy with antibodies against PD-L1 and CCL5 is effective in the treatment of
46 PDAC. *Signal Transduction and Targeted Therapy*, 2020. 5(1): p. 38.
- 47 [29] Cen, X., S. Liu, and K. Cheng, The Role of Toll-Like Receptor in Inflammation and Tumor
48 Immunity. *Frontiers in Pharmacology*, 2018. 9(878).

- 1 [30] Galluzzi, L., E. Vacchelli, A. Eggermont, W.H. Fridman, J. Galon, C. Sautès-Fridman, et
2 al., Trial Watch: Experimental Toll-like receptor agonists for cancer therapy.
3 Oncoimmunology, 2012. 1(5): p. 699-716.
- 4 [31] Hu, X., K. Majchrzak, X. Liu, M.M. Wyatt, C.J. Spooner, J. Moisan, et al., In Vitro Priming
5 of Adoptively Transferred T Cells with a ROR γ Agonist Confers Durable Memory and
6 Stemness In Vivo. Cancer Res, 2018. 78(14): p. 3888-98.
- 7 [32] Cao, D., Z. Qi, Y. Pang, H. Li, H. Xie, J. Wu, et al., Retinoic Acid-Related Orphan Receptor
8 C Regulates Proliferation, Glycolysis, and Chemoresistance via the PD-L1/ITGB6/STAT3
9 Signaling Axis in Bladder Cancer. Cancer Res, 2019. 79(10): p. 2604-18.
- 10 [33] Mahalingam, D., J.S. Wang, E.P. Hamilton, J. Sarantopoulos, J. Nemunaitis, G. Weems, et
11 al., Phase 1 Open-Label, Multicenter Study of First-in-Class ROR γ Agonist LYC-55716
12 (Cintirorgon): Safety, Tolerability, and Preliminary Evidence of Antitumor Activity. Clin
13 Cancer Res, 2019. 25(12): p. 3508-16.
- 14 [34] Zambirinis, C.P. and G. Miller, Signaling via MYD88 in the pancreatic tumor
15 microenvironment: A double-edged sword. Oncoimmunology, 2013. 2(1): p. e22567-e67.
- 16 [35] Mehrotra, S., C.D. Britten, S. Chin, E. Garrett-Mayer, C.A. Cloud, M. Li, et al., Vaccination
17 with poly(IC:LC) and peptide-pulsed autologous dendritic cells in patients with pancreatic
18 cancer. Journal of hematology & oncology, 2017. 10(1): p. 82-82.
- 19 [36] Kirby, M.K., R.C. Ramaker, J. Gertz, N.S. Davis, B.E. Johnston, P.G. Oliver, et al., RNA
20 sequencing of pancreatic adenocarcinoma tumors yields novel expression patterns
21 associated with long-term survival and reveals a role for ANGPTL4. Molecular oncology,
22 2016. 10(8): p. 1169-82.
- 23 [37] Nawrocki, S.T., J.S. Carew, M.S. Pino, R.A. Highshaw, K. Dunner, Jr., P. Huang, et al.,
24 Bortezomib sensitizes pancreatic cancer cells to endoplasmic reticulum stress-mediated
25 apoptosis. Cancer Res, 2005. 65(24): p. 11658-66.
- 26 [38] Hill, R., Y. Li, L.M. Tran, S. Dry, J.H. Calvopina, A. Garcia, et al., Cell intrinsic role of
27 COX-2 in pancreatic cancer development. Molecular cancer therapeutics, 2012. 11(10): p.
28 2127-37.
- 29 [39] Raavé, R., T.H. van Kuppevelt, and W.F. Daamen, Chemotherapeutic drug delivery by
30 tumoral extracellular matrix targeting. Journal of Controlled Release, 2018. 274: p. 1-8.
- 31 [40] Sada, M., K. Ohuchida, K. Horioka, T. Okumura, T. Moriyama, Y. Miyasaka, et al.,
32 Hypoxic stellate cells of pancreatic cancer stroma regulate extracellular matrix fiber
33 organization and cancer cell motility. Cancer Lett, 2016. 372(2): p. 210-8.
- 34 [41] Principe, M., P. Ceruti, N.-Y. Shih, M.S. Chattaragada, S. Rolla, L. Conti, et al., Targeting
35 of surface alpha-enolase inhibits the invasiveness of pancreatic cancer cells. Oncotarget,
36 2015. 6(13): p. 11098-113.
- 37 [42] Nicolle, R., Y. Blum, P. Duconseil, C. Vanbrugghe, N. Brandone, F. Poizat, et al.,
38 Establishment of a pancreatic adenocarcinoma molecular gradient (PAMG) that predicts the
39 clinical outcome of pancreatic cancer. EBioMedicine, 2020. 57: p. 102858.
- 40 [43] Juiz, N., A. Elkaoutari, M. Bigonnet, O. Gayet, J. Roques, R. Nicolle, et al., Basal-like and
41 Classical cells coexistence in pancreatic cancer revealed by single cell analysis. bioRxiv,
42 2020: p. 2020.01.07.897454.
- 43 [44] Hruban, R.H., M.B. Pitman, and D. Klimstra, Ductal adenocarcinoma, in Tumors of the
44 pancreas, A.A.o.T.P.t.S.F. 6, Editor. 2007, Amer Registry of Pathology: ARP Press. p. 422.
- 45 [45] Sobin, L.H., M.K. Gospodarowicz, and C. Wittekind, TNM classification of malignant
46 tumours. 7 ed. 2009, Oxford (UK): Wiley Blackwell.
- 47 [46] Verbeke, C.S., D. Leitch, K.V. Menon, M.J. McMahon, P.J. Guillou, and A. Anthoney,
48 Redefining the R1 resection in pancreatic cancer. Br J Surg, 2006. 93(10): p. 1232-7.

1
2

3 Acknowledgments

4 Parts of this study were part of Lukas Gammelgaard Rasmussen's Master's thesis, where parts of
5 Fig. 1-3, Tab. 1-2 and Supplementary Table 1 and 3 also appear.

6 Legends

7 **Fig. 1.** Morphological subtyping of 108 consecutive, chemotherapy-naïve, surgically resected
8 pancreatic ductal adenocarcinomas (PDACs) as per Kalimuthu. **A.** Bar chart visualizing the
9 morphological pattern composition of each of the 108 tumors (in % of all tumor areas of a given
10 tumor). Each bar represents one tumor. **B–E.** Risk of underestimation of Group B. **B.** At low
11 magnification, the presence of larger glandular tumor formations characteristic of the Group A
12 subtype is conspicuous (H&E). **C.** Higher magnification is required to identify small tumor cell
13 clusters with a solid or abortive glandular pattern consistent with Group B, and to distinguish them
14 from disrupted remnants of non-neoplastic parenchyma. **D–E.** Immunohistochemical staining for
15 broad-spectrum cytokeratin CKAE1/AE3 helps with identifying the less conspicuous non-gland-
16 forming tumor component, but semiquantification, which requires an overview at low power,
17 remains difficult. **F–I.** Survival data for the cohort of surgically resected PDACs, based on Kaplan-
18 Meier and Cox regression analysis. **F.** Survival of the entire cohort. **G.** Survival related to
19 morphological subtypes Group A vs. Group B. **H.** Survival related to the four morphological
20 patterns as per Kalimuthu. **I.** Survival related to WHO differentiation grades.

21 **Fig. 2.** **A.** Hierarchical clustering based on gene expression analysis of 108 surgically resected
22 pancreatic ductal adenocarcinomas (PDACs). The analysis is based on 53 genes, 39 and 14 of
23 which were upregulated in Group A and Group B of PDAC. Bar A) indicates the four identified
24 clusters: Cluster 1 (n=64), Cluster 2 (n=27), Cluster 3 (n=10) and Cluster 4 (n=7). Bar B) indicates
25 the WHO differentiation grades: G1 (grey), G2 (green), and G3 (brown). Bar C) indicates the
26 dominant morphological pattern: Conventional (orange), Tubulopapillary (green), Composite
27 (blue), and Squamous (brown). Bar D) indicates the morphological subtype as per Kalimuthu:
28 Group A (red) and Group B (blue). Case numbers are stated at the bottom of the heat map. **B–C.**

Survival data for the surgical PDAC cohort, based on Kaplan-Meier and Cox regression analysis. **B.** Survival related to the gene clusters 1-4. **C.** Survival related to the clusters 1+3 vs. 2+4.

Fig. 3. Immunohistochemical expression of four representative proteins encoded by genes differentially expressed when comparing Group A with Group B subtypes of PDAC. (A) CD5 is strongly expressed in immune cells and neoplastic glands in a case of the Group A subtype, but almost lacking in Group B (CD5 immunostaining). Scale bar = 250 μ m. (B) CD5 levels are significantly increased in Group A compared to Group B. (C) Higher frequency of FOXP3-positive cells in a Group A case, compared to a Group B case (FOXP3 immunostaining). Scale bar = 250 μ m. (D) FOXP3 tends to be higher expressed in Group A compared to Group B tumors. (E) Strong expression of CD10 in cancer-associated fibroblasts in a Group B, while a Group A cases are almost negative (CD10 immunostaining). Scale bar: 100 μ m. (F) CD10 protein expression levels are significantly higher in Group B compared to Group A. (G) Strong expression of tenascin c (TNC) in cancer-associated fibroblasts and extracellular matrix in a Group B compared to a Group A tumor (TNC immunostaining). Scale bar: 250 μ m (upper panel) and 100 μ m (lower panel). (H) TNC protein expression levels are significantly higher in the Group B compared to the Group A. *Horizontal lines indicate the median. * signifies $P < 0.05$.*

Fig. 4. Morphological subtyping, survival analysis, and gene expression profiling of 88 patients who underwent endoscopic ultrasound-guided fine needle biopsy (EUS-FNB) with a diagnosis of pancreatic ductal adenocarcinoma (PDAC). **A.** EUS-FNBs showing Group A features (upper panel) and Group B features (lower panel) (H&E). **B.** Survival data for the EUS-FNB PDAC cohort, based on Kaplan-Meier and Cox regression analysis (Group A vs. Group B). **C.** Hierarchical clustering of genes in the EUS-FNB PDAC cohort, based on 21 genes, 15 and 6 of which are upregulated in Group A and Group B subtypes of PDAC. **D.** Relative expression of the 21 genes, shown as mean values for each cluster, in the EUS-FNB cohort (n=88) and the surgical PDAC cohort (n=108). For each gene and each cohort, the cluster with the highest and lowest expression is highlighted in green and red. Clusters in-between are highlighted in white/light green/light red. **E.** Survival data for the EUS-FNB PDAC cohort (Cluster 1 vs. Cluster 2/4 vs. Cluster 3). Twelve patients who underwent surgery after pancreatic biopsy were excluded from the survival analyses.

Fig. 5. Expression profiling of genes characteristic of the basal-like and classical subtypes of

1 pancreatic ductal adenocarcinoma (PDAC) in 88 patients who underwent endoscopic ultrasound
2 guided fine needle biopsy (EUS-FNB). **A.** Hierarchical clustering shows that most Group A tumors
3 correspond to the classical subtype (28/34, 82%). Most Group B tumors, on the other hand,
4 correspond to the basal-like subtype (35/54, 65%). **B.** Survival data for the EUS-FNB PDAC
5 cohort, based on Kaplan-Meier and Cox regression analysis (basal-like vs. classical). Twelve
6 patients who underwent surgery after pancreatic biopsy were excluded from the survival analyses.

Tables

Table 1 - Clinicopathological characteristics of the study cohort of 108 preoperative chemotherapy-naïve pancreatic ductal adenocarcinomas (PDACs), specified for the entire cohort and per subtype.

		Entire cohort		Group A		Group B	
		n	%	n	%	n	%
Subjects (n)		108	100.00%	48	44.44%	60	55.56%
Gender	Male	52	48.15%	22	45.83%	30	50.00%
	Female	56	51.85%	26	54.17%	30	50.00%
Age in years – average, (range):		68.8±8.7 (47-86)		69.75±8.62 (49-86)		67.05±8.47 (47-81)	
Tumor site	Pancreatic head	81	75.00%	37	77.08%	44	73.33%
	Pancreatic body	2	1.85%	1	2.08%	1	1.67%

	Pancreatic tail	25	23.15%	10	20.84%	15	25.00%
Resection type	Whipple	71	65.74%	32	66.67%	39	65.00%
	Distal pancreatectomy	27	25.00%	12	25.00%	15	25.00%
	Total pancreatectomy	10 ^a	9.26%	4	8.33%	6	10.00%
Tumor stage [45]	T1	1	0.93%	1	2.08%	0	0.00%
	T2	3	2.78%	0	0.00%	3	5.00%
	T3	103	95.37%	46	95.84%	57	95.00%
	T4	1	0.93%	1	2.08%	0	0.00%
Nodal stage [45]	N0	26	24.07%	13	27.08%	13	21.67%
	N1	82	75.93%	35	72.92%	47	79.33%
Grade of differentiation [13, 14]	Well differentiated (G1)	20	18.52%	18	37.50%	2	3.33%
	Moderately differentiated (G2)	38	35.18%	24	50.00%	14	23.33%

	Poorly differentiated (G3)	50	46.30%	6	12.50%	44	73.33%
Perineural invasion	Yes	64	59.26%	27	56.25%	37	61.67%
	No	10	9.26%	5	10.42%	5	8.33%
	Not available	34	31.48%	16	33.33%	18	30.00%
Vascular invasion	Yes	59	54.63%	25	52.08%	34	56.67%
	No	19	17.59%	9	18.75%	10	16.67%
	Not available	30	27.78%	14	29.67%	16	26.67%
Pancreatic transection margin	Positive	2 ^b	2.78%	1	2.08%	1 ^b	1.67%
	Negative	106 ^c	97.22%	47 ^c	97.92%	59	78.33%
Shortest distance to one of the four resection margins^d	0	36	33.33%	14	29.17%	22	36.67%
	0.5	20	18.52%	12	25.00%	8	13.33%
	1	24	22.22%	12	25.00%	12	20.00%

	1.5	3	2.78%	3	6.25%	0	0.00%
	2	12	11.11%	3	6.25%	9	15.00%
	2.5	1	0.93%	0	0.00%	1	1.67%
	3	11	10.19%	4	8.33%	7	11.67%
	Not available	1	0.93%	0	0.00%	1	1.67%
Mismatch repair status	Mismatch repair deficient	2	1.85%	0	0.00%	2	3.33%
	Mismatch repair stable	106	98.15%	48	100.00%	58	96.67%
Postoperative adjuvant chemotherapy	Gemcitabine ^{e,f}	50	46.30%	26	54.17%	24	40.00%
	Gemcitabine/S1 ^e	18	16.67%	6	12.50%	12	20.00%
	Gemcitabine/capecitabine ^g	3	2.78%	1	2.08%	2	3.33%

	Gemcitabine <4 months ^{h,i}	11	10.18%	4	8.33%	7	11.67%
	None ^j	26	24.07%	11	22.92%	15	25.00%
Overall survival - median (months)		18.83±1.56 (95% CI; 15.31-25.82)		25.8±8.65 (95% CI; 15.34-38.28)		17.15±2.85 (95% CI; 12.75-19.78)	

a: Six total pancreatectomies were Whipple resections followed by left-sided resection, due to a positive lateral transection margin at frozen section. b: One patient had high grade dysplasia in the lateral transection margin, but not PDAC. c: One patient had a NET G1 at the lateral margin, but not PDAC. d: The resection margins are the anterior, posterior, superior mesenteric vein (SMV) and superior mesenteric artery (SMA) margins, according to Verbeke et al. [46]. e: Median duration was 6 months. f: One patient received gemcitabine plus nab-paclitaxel and one patient received 6 months of preoperative chemotherapy. g: Median duration was 4 months. h: Median duration was 2 months. i: Six patients received gemcitabine, three patients received gemcitabine/S1, and 2 patients received gemcitabine/capecitabine. j: Patients did not receive therapy because of poor performance or very old age. Two patients decided not to receive adjuvant therapy.

Table 2 - The top-20 canonical pathways identified in the Group A and Group B subtypes of PDAC using expression analysis of 800 genes.

Description	Number of Genes in Overlap (k), gene name(s)	k/K	p-value	FDR q-value
Group A subtype of PDAC				
Cytokine Signaling in Immune system	15 <i>CCR2, CD27, CD40LG, IL10RA, IL22RA1, JAK2, MMP1, PRLR, PSMB10, RASAL1, RORC, SHC2, SYK, TNFRSF11A, TNFSF17</i>	0.1215	2.33E-15	5.21E-12
Cytokine-cytokine receptor interaction	9 <i>CCR2, CD27, CD40LG, CX3CR1, IL10RA, IL22RA1, PRLR, TNFRSF11A, TNFSF17</i>	0.2361	5.47E-12	6.11E-9

Signaling by Interleukins	8 <i>CCR2, IL10RA, IL22RA1, JAK2, MMP1, PSMB10, RORC, SYK</i>	0.1201	1.83E-8	1.08E-5
TLR3-mediated TICAM1-dependent programmed cell death	3 <i>RIPK3, TICAM1, TLR3</i>	3.4722	1.93E-8	1.08E-5
Chemokine signaling pathway	6 <i>CCR2, CX3CR1, JAK2, PIK3CG, PRKACB, SHC2</i>	0.2201	3.74E-8	1.67E-5
TNFR2 non-canonical NF-kB pathway	5 <i>CD27, CD40LG, TNFRSF11A, TNFSF17, PSMB10</i>	0.3438	6.1E-8	2.27E-5
Adaptive Immune System	9 <i>CD40LG, CD74, ICOSLG, PRKACB, PSMB10, KLRB1, SLAMF7, SYK, ZAP70</i>	0.0764	1.03E-7	3.29E-5

Jak-STAT signaling pathway	5 <i>IL10RA, IL22RA1, JAK2, PIK3CG, PRLR</i>	0,2243	5.17E-7	1.44E-4
RIP-mediated NFkB activation via ZBP1	3 <i>RIPK3, TICAM1, TLR3</i>	1.2257	6.52E-7	1.62E-4
TICAM1, RIP1-mediated IKK complex recruitment	3 <i>RIPK3, TICAM1, TLR3</i>	1.0965	9.28E-7	2.07E-4
ZBP1(DAI) mediated induction of type I IFNs	3 <i>RIPK3, TICAM1, TLR3</i>	0.9924	1.27E-6	2.58E-4
Hemostasis	7 <i>CD2, CD74, JAK2, MMP1, PIK3CG, PRKACB, SYK</i>	0.0715	4.88E-6	9.07E-4
GMCSF-mediated signaling events	3 <i>JAK2, PRKACB, SYK</i>	0.5785	6.75E-6	1.16E-3
Cell adhesion molecules (CAMs)	4 <i>CD2, CD6, CD40LG, ICOSLG</i>	0.2090	1.03E-5	1.64E-3

Natural killer cell mediated cytotoxicity	4 <i>PIK3CG, SHC2, SYK, ZAP70</i>	0.2028	1.16E-5	1.7E-3
MAPK family signaling cascades	5 <i>JAK2, SHC2, PIK3CG, PSMB10, RASAL1</i>	0.1174	1.22E-5	1.7E-3
Innate Immune System	8 <i>CCR2, NLRC5, PRKACB, PSMB10, RIPK3, SYK, TICAM1, TLR3,</i>	0.0072	1.34E-5	1.76E-3
Intestinal immune network for IgA production	3 <i>CD40LG, ICOSLG, TNFSF17</i>	0.4340	1.62E-5	1.91E-3
Class I PI3K signaling events	3 <i>PIK3CG, SYK, ZAP70</i>	0.4340	1.62E-5	1.91E-3
Pathways in cancer	5 <i>GLI1, MMP1, PIK3CG, TRAF1, WNT2B</i>	0.1069	1.91E-5	2.14E-3

Genes not related to top 20 canonical pathways	7 <i>CD5, FOXP3, FUT4, LY9, MMP7, PDZK1IP1, P2RY13</i>	-	-	-
--	---	---	---	---

Description	Number of Genes in Overlap (k), gene name	k/K	p-value	FDR value	q-value
Group B subtype of PDAC					
FOXM1 transcription factor network	3 <i>BIRC5, CCNB1, CDK2</i>	0.5208	3.78E-7	8.44E-4	
Validated targets of C-MYC transcriptional activation	3 <i>BIRC5, CCNB1, ENO1</i>	0.2639	3,00E-06	2.16E-3	
Cell Cycle Checkpoints	4 <i>BIRC5, CCNB1, CDK2, UBE2C</i>	0.0951	3.04E-6	2.16E-3	

APC/C-mediated degradation of cell cycle proteins	3 <i>CCNB1, CDK2, UBE2C</i>	0.2424	3.88E-6	2.16E-3
Estrogen-responsive protein Efp controls cell cycle and breast tumors growth	2 <i>CCNB1, CDK2</i>	0.8681	1.48E-5	6.59E-3
Phosphorylation of the APC/C	2 <i>CCNB1, UBE2C</i>	0.7313	2.1E-5	7.82E-3
Cyclins and Cell Cycle Regulation	2 <i>CCNB1, CDK2</i>	0.6042	3.11E-5	8.67E-3
APC/C:Cdc20 mediated degradation of Cyclin B	2 <i>CCNB1, UBE2C</i>	0.6042	3.11E-5	8.67E-3

Cyclin A/B1/B2 associated events during G2/M transition	2 <i>CCNB1, CDK2</i>	0.5556	3.68E-5	8.9E-3
Cell Cycle, Mitotic	4 <i>BIRC5, CCNB1, CDK2, UBE2C</i>	0.0071	3.99E-5	8.9E-3
Nicotinate metabolism	2 <i>NT5E, PTGS2</i>	0.4479	5.7E-5	1.16E-2
Mitotic Metaphase and Anaphase	3 <i>BIRC5, CCNB1, UBE2C</i>	0.0889	7.83E-5	1.4E-2
Cell Cycle	4 <i>BIRC5, CCNB1, CDK2, UBE2C</i>	0.0059	8.18E-5	1.4E-2
FoxO family signaling	2 <i>CCNB1, CDK2</i>	0.2833	1.44E-4	2.14E-2

TP53 Regulates Transcription of Cell Cycle Genes	2 <i>CCNB1, CDK2</i>	0.2833	1.44E-4	2.14E-2
Pathways in cancer	3 <i>BIRC5, CDK2, PTGS2</i>	0.0092	2.04E-4	2.72E-2
Cytokine Signaling in Immune system	4 <i>BIRC5, FLNB, IL11, PTGS2</i>	0.0047	2.07E-4	2.72E-2
HIF-1-alpha transcription factor network	2 <i>ENO1, NT5E</i>	0.2104	2.61E-4	3.2E-2
p53 signaling pathway	2 <i>CCNB1, ENO1</i>	0.2042	2.77E-4	3.2E-2
Transcriptional Regulation by TP53	3 <i>BIRC5, CCNB1, CDK2</i>	0.0082	2.87E-4	3.2E-2
Genes not related to top 20 canonical pathways	5 <i>ANGPTL4, ANLN, MME, TNC, PLOD2</i>	-	-	-