

Mathematical Proceedings of the Cambridge Philosophical Society

<http://journals.cambridge.org/PSP>


Additional services for *Mathematical Proceedings of the Cambridge Philosophical Society*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Characterizing connected K -theory by homotopy groups

John Rognes

Mathematical Proceedings of the Cambridge Philosophical Society / Volume 114 / Issue 01 / July 1993, pp 99 - 102

DOI: 10.1017/S0305004100071437, Published online: 24 October 2008

Link to this article: http://journals.cambridge.org/abstract_S0305004100071437

How to cite this article:

John Rognes (1993). Characterizing connected K -theory by homotopy groups. *Mathematical Proceedings of the Cambridge Philosophical Society*, 114, pp 99-102 doi:10.1017/S0305004100071437

Request Permissions : [Click here](#)

Characterizing connected K -theory by homotopy groups

BY JOHN ROGNES

Matematisk Institutt, Universitetet i Oslo, Norway

(Received 31 July 1992; revised 18 November 1992)

In this note we provide proofs for two facts related to connected topological K -theory, expected by Ib Madsen.

Let bu denote the connected K -theory spectrum, and let $\text{Im } J$ denote the connected image of J spectrum. For an odd prime p we prove that bu_p^\wedge is characterized by its homotopy groups and v_1 -periodic structure. Thereafter we prove that spectrum maps from bu_p^\wedge to $B \text{Im } J_p^\wedge$ are detected on homotopy groups.

The purpose of proving these Lemmas characterizing properties of bu by homotopy group data is to complete the calculation of M. Bökstedt, W. C. Hsiang and I. Madsen's cyclotomic trace invariant $TC(\mathbb{Z}_p^\wedge, p)$, getting a map from $K(\mathbb{Z}_p^\wedge)$ for odd p . See Corollary 3, whose proof is due to Bökstedt and Madsen. Their proof of its hypothesis will appear in [2].

LEMMA 1. *Suppose X is a p -complete spectrum, with $\pi_* X \cong \pi_* bu_p^\wedge$ abstractly, and*

$$\pi_*(X; \mathbb{F}_p) \cong \pi_*(bu; \mathbb{F}_p)$$

as $\mathbb{F}_p[v_1]$ -modules. Then $X \simeq bu_p^\wedge$.

Let S denote the sphere spectrum, and $H\mathbb{F}_p$ the mod p Eilenberg–MacLane spectrum. For any spectrum Y , let Y/p denote the cofibre of the ‘multiplication by p ’ map. In all proofs, let all spectra be implicitly completed at p .

Proof. We first prove that $X/p \simeq bu/p$. The map $\Sigma^{2p-2} S/p \xrightarrow{v_1} S/p$ smashed with X induces the v_1 -module structure on mod p homotopy, and appears in the bottom line of:

$$\begin{array}{ccccc}
 \Sigma^{2p-2} bu/p & \xrightarrow{v_1} & bu/p & \longrightarrow & \bigvee_{i=1}^{p-1} \Sigma^{2i} H\mathbb{F}_p \\
 & & \vdots & & \downarrow = \\
 & & f & & \\
 & & \vdots & & \\
 \Sigma^{2p-2} X/p & \xrightarrow{v_1} & X/p & \longrightarrow & C(v_1) \longrightarrow \Sigma^{2p-1} X/p
 \end{array}$$

where the horizontal lines are (co-)fibration sequences (cf. [1], corollary 4·8). By the hypothesis about the $\mathbb{F}_p[v_1]$ -module structure on $\pi_*(X/p)$, the mapping cofibre $C(v_1)$ has the same homotopy groups as $\bigvee_{i=1}^{p-1} \Sigma^{2i} H\mathbb{F}_p$. The k -invariants of its Postnikov tower thus sit in degrees 3 to $2p-3$ in the mod p Steenrod algebra, where all the groups vanish. Hence $C(v_1)$ is a wedge of Eilenberg–MacLane spectra, and we can choose an equivalence as in the diagram above.

Consider the composite $bu/p \rightarrow \vee_i \Sigma^{2i} H\mathbb{F}_p \rightarrow C(v_1) \rightarrow \Sigma^{2p-1} X/p$. By the usual Atiyah–Hirzebruch spectral sequence argument (all groups sit in odd degrees), this map must induce a null-homotopic map of underlying spaces.

For the corresponding spectrum level conclusion, recall that the spectrum bu is the connected cover of the mapping telescope for the canonical degree two map (multiply by the Bott class) on the suspension spectrum of its zeroth space. Also note that there is an equivalence between maps from a periodic spectrum or its connected cover to a connected periodic spectrum (cf. [4], proposition II:3:8).

Similarly, the spectrum bu/p can be recovered as the connected cover of the mapping telescope for the v_1 -map on the suspension spectrum of its zeroth space. Hence by a Milnor exact sequence argument there are no essential spectrum maps from bu/p to $\Sigma^{2p-1} X/p$. (The \lim^1 -term vanishes by completeness.)

Thus there exists a lifting f making the diagram above commute. On homotopy, f induces a homomorphism of isomorphic $\mathbb{F}_p[v_1]$ -modules which is an isomorphism modulo the ideal generated by v_1 . Hence $\pi_*(f)$ is an isomorphism and f is an equivalence.

Next consider the diagram :

$$\begin{array}{ccccccc}
 bu & \xrightarrow{p^*} & bu & \xrightarrow{q} & bu/p & \longrightarrow & su \\
 & & \vdots & & \downarrow f & & \\
 & & X & \longrightarrow & X/p & \longrightarrow & \Sigma X \\
 X & \xrightarrow{p^*} & X & & & &
 \end{array}$$

By an Atiyah–Hirzebruch spectral sequence argument as above, there are no essential maps $bu \rightarrow \Sigma X$. Hence the composite $f \circ q$ lifts to a map $h : bu \rightarrow X$. This is a map between p -complete spectra with abstractly isomorphic homotopy groups, which induces an equivalence with mod p -coefficients, and is hence an equivalence. █

I thank Marcel Bökstedt for noting the need for the spectrum level extensions in the above argument.

Let q be a topological generator for the p -adic units $\mathbb{Z}_p^\times \subset \mathbb{Z}_p^\wedge$. Extend the Adams operation ψ^q to a spectrum map $bu_p^\wedge \rightarrow bu_p^\wedge$ mapping as $q^{-n} \psi^q$ on the $2n$ th space, and let $\text{Im } J_p^\wedge$ be the fibre of $\psi^q - 1 : bu_p^\wedge \rightarrow bu_p^\wedge$.

LEMMA 2. *The spectrum maps from bu_p^\wedge to $B \text{Im } J_p^\wedge$ are precisely the p -adic multiples of the cofibre map corresponding to $\psi^q - 1$. They are all detected on homotopy groups.*

Proof. Consider the diagram :

$$\begin{array}{ccccccc}
 & & bu & & & & \\
 & & \vdots & \searrow & & & \\
 & & \phi & & & & \\
 & & \vdots & & & & \\
 bu & \xrightarrow{\psi^q - 1} & bu & \longrightarrow & B \text{Im } J & \longrightarrow & su
 \end{array}$$

Again, there are no essential maps from bu to su , so any spectrum map $bu \rightarrow B \text{Im } J$

lifts to a spectrum map $\phi: bu \rightarrow bu$. This is a reduced K -theory operation, and by [3] (formula 2.3 and remark 2.10), ϕ is expressible as a series:

$$\phi = \sum_{r \geq 0} \sum_{i \geq 0} a_{r,i} (\psi^{p^{r+i}} - \psi^i)$$

with $a_{r,i} \in \mathbb{Z}_p^\wedge$, where $a_{r,i} = 0$ if $i \geq p^{r+1} - p^r$ or $p \mid i$, and the sum may be infinite in r . Note that $\psi^{p^{r+i}} - \psi^i$ may be expressed as $(\psi^{p^{r+i}} - 1) - (\psi^i - 1)$ with both p^{r+i} and i prime to p in the cases where $a_{r,i} \neq 0$, except when $r = i = 0$.

We claim that any operation $\psi^n - 1$ with n a p -adic unit factors through $\psi^q - 1$. This follows for $n = q^i$ with i a natural number from the factorization $\psi^{q^i} - 1 = (\psi^q - 1)(\psi^{q^{i-1}} + \dots + \psi^q + 1)$, and for negative i using the operation ψ^{-q} . As q is a topological generator, the general case follows by completion.

On the other hand, any operation ϕ factoring through $\psi^q - 1$ has $a_{0,0} = 0$. For if it were nonzero, let $j = v_p(a_{0,0}) < \infty$ be the p -adic valuation, let $i = (p-1)p^j$, and consider homotopy groups in degree $2i$. For any p -adic unit n , $v_p(n^i - 1) \geq j + 1$, so

$$v_p(\pi_{2i}(\phi)) = v_p(a_{0,0}) = j$$

which precludes ϕ from mapping into the image of $\psi^q - 1$, as

$$v_p(\pi_{2i}(\psi^q - 1)) = v_p(q^i - 1) \geq j + 1$$

by the above (q is a p -adic unit).

Hence the subspace of operations ϕ which factor through $\psi^q - 1$ is contained in the closed ideal where $a_{0,0} = 0$ in the compact Hausdorff ring of all stable operations, and contains the dense subset of such operations which are finite sums of Adams operations. As the former subspace is a continuous image of the whole compact ring, it is itself compact, thus closed, and is therefore the whole ideal.

In conclusion, the coefficient $a_{0,0}$ precisely classifies the spectrum maps $bu \rightarrow B \operatorname{Im} J$, as we wanted to prove. \blacksquare

Let j denote the connective image of J spectrum, i.e. the connective cover of the fibre of $\psi^q - 1: K \rightarrow K$, or equivalently the connective cover of the K -localization of S , where K is the periodic K -theory spectrum.

COROLLARY 3. (*Bökstedt–Madsen*). *Suppose there is a map of p -complete spectra $j_p^\wedge \vee Bj_p^\wedge \rightarrow TC(\mathbb{Z}_p^\wedge, p)$ with cofibre X , injective on homotopy groups, such that $\pi_* X \cong \pi_* su_p^\wedge$ abstractly, and $\pi_*(X; \mathbb{F}_p) \cong \pi_*(su; \mathbb{F}_p)$ as $\mathbb{F}_p[v_1]$ -modules. Then*

$$TC(\mathbb{Z}_p^\wedge, p) \simeq j_p^\wedge \vee Bj_p^\wedge \vee su_p^\wedge.$$

Proof. By suspending Lemma 1 once, we find $X \simeq su$. Thus there is a fibration sequence of spectra

$$j \vee Bj \longrightarrow TC(\mathbb{Z}_p^\wedge, p) \longrightarrow su \xrightarrow{\Sigma e} Bj \vee BBj$$

Desuspending the map Σe once, we are led to consider $e: bu \rightarrow j \vee Bj$, which is assumed zero on homotopy groups. There are no essential maps $bu \rightarrow j$ which are zero on homotopy groups, owing to the corresponding fact for maps $bu \rightarrow bu$. Any map $bu \rightarrow Bj$ which is zero on homotopy groups factors through the 1-connected cover $B \operatorname{Im} J$, so by Lemma 2 the map e (and thus Σe) is null-homotopic, from which the splitting follows (lift the identity map of su). \blacksquare

REFERENCES

- [1] J. F. ADAMS. *Lectures on Generalized Cohomology*. Category Theory, Homology Theory and their Applications III. Lecture Notes in Math., vol. 99 (Springer, 1969), pp. 1–138.
- [2] M. BÖKSTEDT and I. MADSEN. Topological Cyclic Homology of the Integers, to appear in *Astérisque* (1993).
- [3] I. MADSEN, V. P. SNAITH and J. TORNEHAVE. Infinite loop maps in geometric topology. *Math. Proc. Camb. Phil. Soc.* **81** (1977), 399–429.
- [4] J. P. MAY *et al.* E_∞ ring spaces and E_∞ ring spectra. Lecture Notes in Math., vol. 577 (Springer, 1977).