



Antihypertensive therapy prevents new-onset atrial fibrillation in patients with and without isolated systolic hypertension: the LIFE Study

Journal:	<i>Blood Pressure</i>
Manuscript ID	BP-2019-OR-0051.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Larstorp, Anne C; University of Oslo, Departments of Medical Biochemistry, Oslo University Hospital Stokke, Ildri Maria; University of Oslo, Institute for Clinical Medicine Kjeldsen, S. E.; Oslo universitetssykehus Ullevål, Hecht Olsen, Michael; Department of Internal Medicine, Holbaek Hospital, and Centre for Individualized Medicine in Arterial Diseases (CIMA), Odense University Hospital Okin, Peter; Weill Cornell Medical College, Medicine Devereux, Richard B; Weill Cornell Medicine, Greenberg Division of Cardiology Wachtell, Kristian; University of Oslo, Departments of Cardiology, Oslo University Hospital
Keywords:	atrial fibrillation, blood pressure, left ventricular hypertrophy, isolated systolic function

SCHOLARONE™
Manuscripts

1
2
3
4 **Antihypertensive therapy prevents new-onset atrial fibrillation in patients**
5
6
7 **with and without isolated systolic hypertension: the LIFE Study**
8
9

10
11
12 A.C.K. Larstorp ^a, I.M. Stokke ^a, S.E. Kjeldsen ^a, M.H. Olsen ^b, P.M. Okin ^c, R.B. Devereux ^c
13
14 and K. Wachtell ^a
15

16
17
18
19 ^a University of Oslo, Departments of Medical Biochemistry and Cardiology, Oslo University
20
21 Hospital; Oslo, Norway; ^b Department of Internal Medicine, Holbaek Hospital, and Centre
22
23 for Individualized Medicine in Arterial Diseases (CIMA), Odense University Hospital,
24
25 Denmark; and ^c Weill Cornell Medicine, Greenberg Division of Cardiology, New York, NY,
26
27 U.S.A.
28
29

30
31
32 Contact_Sverre E. Kjeldsen

33 s.e.kjeldsen@medisin.uio.no

34
35 Department of Cardiology, Oslo University Hospital, Ullevaal,
36
37 Oslo, Norway. Telephone No. +47 22119100, Telefax No. +47 22119181
38
39

40
41 **Clinical Trials Registration Information:**

42
43 URL: <http://www.clinicaltrials.gov/ct2/show/NCT00338260> (Identifier NCT00338260)
44
45

46
47
48 **Word Count: Abstract 260 and Ms. including references 46203743**
49

50
51 **KEYWORDS: Atrial fibrillation; blood pressure; left ventricular hypertrophy, isolated**
52
53 **systolic hypertension**

54
55 **Abbreviations:** AF, atrial fibrillation; BP, blood pressure; ECG, electrocardiography; ECG-
56
57 LVH, electrocardiographic left ventricular hypertrophy; ISH, isolated systolic hypertension;
58
59 LIFE, Losartan Intervention For Endpoint reduction in hypertension; LVH, left ventricular
60
hypertrophy; SBP, systolic blood pressure

ABSTRACT

Aims: Atrial fibrillation (AF) is associated with increased cardiovascular risk and the incidence increases with age, hypertension and left ventricular hypertrophy (LVH). Reducing in-treatment systolic blood pressure (SBP) prevents new-onset AF but has previously not been studied in patients with isolated systolic hypertension (ISH). We aimed to investigate the effect on preventing new-onset AF by decreased in-treatment SBP in patients with ISH compared to patients with non-ISH.

Methods and results: Double-blind, randomized, parallel-group study of 1,320 patients with ISH and electrocardiographic (ECG) LVH, included among the 9,193 patients in the Losartan Intervention For Endpoint reduction in hypertension (LIFE) study. Annual ECGs were Minnesota coded centrally, and new-onset AF were evaluated in 1,248 ISH patients and compared with 7,583 non-ISH patients during mean 4.8 ± 0.9 years follow-up. Cox regression analyses were used to assess the effect of reduced in-treatment SBP. New-onset AF occurred in 61 (4.9%) ISH patients and 292 (3.9%) non-ISH patients. In multivariate analysis lower in-treatment SBP was associated with 17% risk reduction ($p=0.008$) for new-onset AF in ISH patients and 9% risk reduction ($p=0.006$) in non-ISH patients per 10 mmHg decrease in in-treatment SBP, independent of treatment modality, baseline risk factors, baseline SBP and in-treatment heart rate and ECG-LVH. There was a significant interaction ($p=0.041$) in favor of SBP reduction and AF prevention in ISH vs. non-ISH patients.

Conclusion: Our data suggest that the effect of in-treatment SBP reduction in preventing new-onset AF is stronger in ISH compared to non-ISH patients with hypertension and ECG-LVH. However, the principal findings were the same in ISH and non-ISH patients.

Introduction

Isolated systolic hypertension (ISH), usually the most common form of hypertension in the elderly, is associated with increased risk of cardiovascular morbidity and mortality compared to systolic-diastolic and isolated diastolic (non-ISH) hypertension [1,2], and antihypertensive treatment is efficacious in preventing stroke and myocardial infarction in ISH patients [3].

The Losartan Intervention For Endpoint reduction in hypertension (LIFE) study comprised of 9,193 patients aged 55–80 yrs., with hypertension and electrocardiographic left ventricular hypertrophy (ECG-LVH), followed for 4.8 ± 0.9 yrs. who were randomized to losartan vs. atenolol targeting systolic blood pressure (SBP) <140 mmHg [4]. Losartan reduced new-onset atrial fibrillation (AF), a secondary endpoint and subsequent stroke compared to atenolol [5] and lower Cornell product and lower heart rate over time were associated with less new-onset AF [6,7].

~~Isolated systolic hypertension (ISH), usually the most common form of hypertension in the elderly, is associated with increased risk of cardiovascular morbidity and mortality compared to systolic-diastolic and isolated diastolic (non-ISH) hypertension [5,6], and antihypertensive treatment is efficacious in preventing stroke and myocardial infarction in ISH patients [7].~~

Patients with ISH comprised a pre-specified subgroup of special interest in the LIFE Study [8]. We have shown that the LIFE patients with ISH benefitted from regression of LVH in response to the antihypertensive treatment [9,10] and that pulse pressure at baseline and during treatment were predictors of incident AF [11]. Whether lower in-treatment SBP is associated with less AF in patients with ISH and ECG-LVH has, however, not been specifically investigated.

Based on the benefit of LVH regression in preventing incident AF [3], the benefit in the ISH

1
2
3 patients of regression of LVH on endpoint protection [9,10] and the association between
4
5 lower pulse pressure during treatment and incident AF [11], we hypothesized and
6
7 investigated whether patients with ISH would benefit from the antihypertensive treatment in
8
9 preventing AF compared to non-ISH patients who participated in the LIFE study. Competing
10
11 risk was additionally analyzed with incident AF and mortality as a combined endpoint.
12
13
14
15
16
17

18 **Material and methods**

19 **Participants and study design**

20
21 As described in detail elsewhere [1, 12-14], the LIFE study enrolled patients with
22
23 hypertension having ECG-LVH determined by Cornell voltage-duration product [15-17]
24
25 and/or Sokolow-Lyon voltage criteria [18] on a screening ECG in a prospective, double-blind,
26
27 randomized, parallel group study large enough (n=9,193) to have sufficient power (80%) to
28
29 detect a difference of at least 15% in the incidence of combined cardiovascular morbidity and
30
31 mortality with use of losartan as opposed to atenolol. The primary end point was a composite
32
33 of cardiovascular death, non-fatal myocardial infarction, and non-fatal stroke, and investigator
34
35 reported end points were verified by an expert end point committee. Patients included were
36
37 men and women aged 55 to 80 years with previously untreated or treated essential
38
39 hypertension with mean seated BP in the range of 160 to 205 mmHg systolic, or 95 to 115
40
41 mmHg diastolic, or both, after 1 to 2 weeks of receiving placebo and who had not experienced
42
43 a myocardial infarction or stroke within 6 months and did not require treatment with a β -
44
45 blocker, angiotensin-converting enzyme inhibitor, or angiotensin-receptor blocker. The trial
46
47 protocol was approved by all ethics committees concerned, was overseen by an independent
48
49 data and safety monitoring board and all patients gave written informed consent. Targeting a
50
51 BP of 140/90 mmHg or lower, double-blind treatment was initiated with losartan, 50 mg, or
52
53 atenolol, 50 mg, daily and matching placebo of the other agent. Study therapy was up-titrated
54
55
56
57
58
59
60

1
2
3 by addition of hydrochlorothiazide, 12.5 mg, followed by increase in blinded losartan or
4
5 atenolol to 100 mg/day. If this regimen was not sufficient to reach the target BP, additional
6
7 open-label upward titration of hydrochlorothiazide and institution of therapy with in most
8
9 cases calcium channel blocker or additional other medication (excluding β -blockers,
10
11 angiotensin-converting enzyme inhibitors, and angiotensin-receptor blockers) was added to
12
13 the treatment regimen [12].
14
15

16
17 A total of 8,831 patients with neither a history of AF or AF on their baseline ECG
18
19 were included in the present study. Of these, 1,248 patients (14.1%) had ISH, a priori defined
20
21 as systolic BP of 160 to 205 mmHg and diastolic BP <90 mmHg [8] and 7,583 (85.9%) had
22
23 non-ISH. Characteristics of ISH patients are presented in Table 1. Characteristics of the non-
24
25 ISH patients have been previously published [9,10].
26
27

28 **Electrocardiography and endpoint determination**

29
30 ECGs were obtained at study baseline, at 6 months, and at yearly follow-up intervals until study
31
32 termination or patient death as previously reported in detail [2-4]. New-onset AF was a pre-
33
34 specified secondary endpoint and was identified by Minnesota coding of annual in-study ECGs
35
36 at the core laboratory at Sahlgrenska University Hospital/Östra, Göteborg, Sweden [2,12] by
37
38 experienced readers blinded to clinical information.
39
40

41 **Statistical analyses**

42
43 Data management and statistical analyses were performed by the investigators with SPSS
44
45 version 22.0 (IBM, Inc, Armonk, NY). Data are presented as mean (SD) for continuous
46
47 variables and as proportions for categorical variables. Differences in prevalences were
48
49 compared using χ^2 analyses, and differences in mean values were compared using
50
51 independent-samples *t*-tests. Associations between SBP during antihypertensive therapy and
52
53 the occurrence of new-onset AF were analyzed using Cox proportional hazard models [19,20]
54
55 and based on the intention-to-treat principle. Univariate Cox regression analyses were
56
57
58
59
60

1
2
3 performed initially to identify important and significant background predictors of AF in the
4 ISH population. The final multivariate models included significant univariate predictors that
5 remained significant in multivariate analyses by using stepwise forward regression. The final
6 model could not include too many predictors due to the number of new-onset atrial fibrillation
7 events. Therefore, Framingham risk score was used to adjust for several cardiovascular risk
8 factors in a one propensity score. Baseline and subsequent determinations of SBP were entered
9 as a time-varying covariate in the Cox models. History of coronary heart disease was entered
10 as a standard covariate as it has previously been shown to predict atrial fibrillation [52]. The
11 same multivariate models were used on both ISH and non-ISH patients. for the purpose of
12 comparison. In Model 1, in-treatment SBP was adjusted for a treatment group indicator
13 entered as a standard covariate to account for possible effects of treatment with losartan vs.
14 atenolol, and Framingham risk score and history of coronary heart disease included as
15 standard covariates, and in-treatment heart rate and in-treatment ECG-LVH determined by
16 Cornell voltage-duration product entered as time-varying covariates. In Model 2, in-treatment
17 SBP was adjusted for baseline SBP, in addition to the above-mentioned predictors in Model 1.
18 The hazard ratios (HRs) for incident AF were computed for 10 mmHg decrements of in-
19 treatment SBP treated as a continuous variable. The 95% confidence intervals (CI) of HRs
20 were calculated from the estimated coefficients and their standard errors [21], and Wald χ^2
21 statistics and probability values were calculated. In parallel analyses competing risk was
22 analyzed with incident AF and mortality as a combined endpoint and adjusted for the same
23 variables as in Model 2 in Table 3. In order to test if the effect of reduced SBP on reduced
24 incident AF was significantly different in ISH patients compared to non-ISH patients,
25 interaction analyses were performed using Cox proportional hazard models with all patients
26 and entering ISH status as a categorical covariate and in-treatment SBP as a time-varying
27 covariate and the cross-product of ISH status and time-varying SBP. The *P* value of this
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 interaction cross-product was used to determine if there was a quantitative interaction, i.e.
4
5 whether ISH increased or decreased the outcome effect of the reduction in SBP. A two tailed
6
7 *P* value of <0.05 was required for statistical significance. All study data reside in a database
8
9 with the authors.
10

11 Results

12
13
14 New-onset AF occurred in 61 patients (4.9 %) of 1,248 patients with ISH during the 4.8 ± 0.9
15
16 yrs. of mean follow-up (Fig. 1). The patients with incident AF were slightly older than those
17
18 who remained in sinus rhythm, and fewer were treated with losartan, they were taller (though
19
20 no difference in body mass index), they had higher SBP, pulse pressure and Sokolow-Lyon
21
22 voltage, and their serum potassium was slightly lower (Table 1). New-onset AF occurred in
23
24 292 patients (3.9 %) of 7,583 non-ISH patients during the same mean follow-up time (Fig. 1).
25
26
27
28

29
30 Reductions in SBP were approximately the same in ISH and non-ISH patients whether they
31
32 had new-onset AF or not (Fig. 2a). ISH patients without new-onset AF had SBP values (mean
33
34 (SD)) at baseline and year 1 to year 5 of follow-up of 174.1 (10.8), 151.9 (17.0), 150.1 (16.4),
35
36 148.3 (16.8), 147.7 (16.9) and 148.6 (16.5) mmHg compared to 179.0 (11.5), 153.7 (16.7),
37
38 151.1 (20.3), 149.9 (17.8), 147.0 (18.9) and 139.1 (14.8) mmHg in ISH patients with new-
39
40 onset AF. Non-ISH patients without new-onset AF had SBP values (mean (SD)) at baseline
41
42 and year 1 to year 5 of follow-up of 174.2 (14.8), 150.0 (16.7), 148.0 (16.2), 146.7 (16.4),
43
44 145.4 (15.8) and 144.5 (15.3) mmHg compared to 177.0 (14.2), 152.8 (17.9), 151.0 (17.8),
45
46 149.3 (19.9), 145.1 (18.2) and 144.9 (15.4) mmHg in non-ISH patients with new-onset AF.
47
48
49

50
51 Overall reductions in diastolic BPs were also approximately the same in ISH and non-ISH
52
53 patients (Fig. 2b), with a possible larger decrease in mean diastolic BP from baseline to year 1
54
55 in non-ISH patients. ISH patients without new-onset AF had diastolic DBP values (mean
56
57 (SD)) at baseline and year 1 to year 5 of follow-up of 82.7 (5.8), 76.9 (8.9), 76.1 (8.8), 75.2
58
59 (9.3), 74.7 (9.0) and 74.7 (8.7) mmHg compared to 83.4 (4.8), 78.4 (8.5), 78.7 (9.9), 75.7
60

1
2
3 [\(10.4\), 75.1 \(9.6\) and 71.1 \(9.2\) mmHg in ISH patients with new-onset AF. Non-ISH patients](#)
4 [without new-onset AF had DBP values \(mean \(SD\)\) at baseline and year 1 to year 5 of](#)
5 [follow-up of 100.4 \(6.4\), 86.5 \(8.5\), 85.2 \(8.2\), 84.1 \(8.5\), 82.8 \(8.5\) and 82.2 \(8.1\) compared](#)
6 [to 99.7 \(6.1\), 86.1 \(8.8\), 84.9 \(8.3\), 83.7 \(8.8\), 82.1 \(9.6\) and 83.3 \(8.9\) mmHg in non-ISH](#)
7 [patients with new-onset AF.](#) Cornell voltage product as a measure of ECG-LVH was
8 however higher throughout the study in both ISH and non-ISH patients who developed new-
9 onset AF compared to the patients who remained without AF (Fig. 3).

10
11
12
13
14
15
16
17
18
19
20 We analyzed univariate predictors of new-onset AF in the ISH patients (Table 2a). There was
21 a 4 % increased risk of new-onset AF per 1 mmHg rise in SBP at baseline. Furthermore,
22 height, baseline Sokolow-Lyon voltage, potassium, treatment allocation, time-varying heart
23 rate and age were significant univariate predictors of incident AF, but smoking, time-varying
24 Cornell voltage product and gender were not. [Univariate predictors of new-onset AF in the](#)
25 [non-ISH patients are shown in Table 2b. We have not elaborated on these in detail as the sample](#)
26 [size and thus statistical power were much higher and stronger than in the ISH subset of](#)
27 [patients.](#)

28
29
30
31
32
33
34
35
36
37
38
39 The effect of reduction of in-treatment SBP is further explored in multivariate analysis (Table
40 3). While the reduction in new-onset AF in ISH patients was borderline significant in
41 univariate analysis ($p=0.070$), and when adjusted for treatment effect only ($p=0.066$), it was
42 highly significant when adjusted for baseline SBP and in the multivariate model 2 ($p=0.008$
43 for both). Lower in-treatment SBP was associated with a 17% risk reduction for new-onset
44 AF per 10 mmHg decrease in SBP in ISH patients ($p=0.008$), and 9% risk reduction
45 ($p=0.006$) in non-ISH patients, independent of treatment modality, baseline risk factors,
46 baseline SBP and in-treatment heart rate and ECG-LVH.

1
2
3 There was a significant interaction ($p=0.041$) in favor of stronger effect of SBP reduction on
4 preventing new onset AF in ISH vs. non-ISH patients.
5
6

7
8 New-onset AF or death occurred in 190 (15.2%) of the ISH patients and in 858 (11.3%) of the
9 non-ISH patients (Fig. 1). Lower in-treatment SBP was associated with lower risk of new-
10 onset AF or death in the non-ISH patients ($p<0.001$, Model 2), however the effect of lower in-
11 treatment SBP on the combined endpoint new-onset AF and death was only borderline
12 significant in the ISH patients ($p=0.087$) in Model 2 which included adjustment for baseline
13 SBP (Table 4).
14
15
16
17
18
19
20
21
22

23 Discussion

24
25 New-onset AF occurred in 61 (4.9%) of 1,248 patients with ISH and ECG-LVH. Lower in-
26 treatment SBP was associated with a 17% risk reduction for new-onset AF per 10 mmHg
27 decrease in SBP, independent of treatment modality, baseline risk factors, baseline SBP and
28 in-treatment heart rate and ECG-LVH in patients with ISH. There was a significant
29 interaction in favor of stronger impact of SBP reduction in preventing incident AF in ISH vs.
30 non-ISH patients. When taking death and competing risk into account the overall findings
31 confirmed the treatment benefits on preventing new-onset AF both in patients with ISH and in
32 patients with non-ISH.
33
34
35
36
37
38
39
40
41
42
43

44
45 Our findings suggest that reduced SBP per se may reduce the incidence of AF in patients
46 above 55 years of age with ISH and ECG-LVH. The impact of SBP reduction in preventing
47 incident AF is even slightly stronger in patients with ISH compared to non-ISH patients
48 recruited from the same population. These findings are in line with our previous findings
49 showing benefits of SBP lowering therapy and regression of ECG-LVH on preventing
50 cardiovascular mortality, myocardial infarction, cerebral stroke and heart failure in LIFE
51 study participants with ISH [8-10]. The patients who develop new-onset AF and other
52
53
54
55
56
57
58
59
60

1
2
3 complications remained with higher Cornell voltage product ECG-LVH throughout the study
4
5 as depicted in Fig. 3.
6
7

8 Atrial fibrillation is associated with increased risk of cardiovascular morbidity and mortality.
9
10 It is important to identify modifiable risk factors as both men and women have an
11
12 approximate 25% overall lifetime risk of AF (22). To our knowledge this is the first study to
13
14 report an effect of lower in-treatment SBP and reduced new-onset AF in patients with ISH
15
16 and ECG-LVH.
17
18

19
20 AF is the most prevalent sustained cardiac arrhythmia and the prevalence is increasing [23].
21
22 In the Rotterdam study, the prevalence of AF varied from 0.7% in the age group 55-59 years
23
24 to 17.8% in those aged 85 years and above [24]. AF incidence increases with age but also
25
26 with other risk factors [25], including diabetes, obesity, hypertension, LVH, coronary heart
27
28 disease, congestive heart failure, valvular heart disease and increased left atrial size by
29
30 echocardiography [26-28]. AF is associated with a 4 to 5 fold increased risk of ischemic
31
32 stroke [29-30] and with a near doubled cardiovascular mortality risk [31]. Prevention of AF is
33
34 thus of major importance and hypertension is currently the most prevalent, potentially
35
36 modifiable risk factor, accounting for approximately 14 to 22% of AF cases [26,32,33].
37
38

39
40 ISH is closely related to increased pulse pressure, a marker of advanced vascular ageing (34) and
41
42 arterial stiffness [35,36], which may contribute in the structural and electrical remodeling of the
43
44 myocardium leading to the development of AF, possibly through increased pulsatile load on the
45
46 heart and increased left atrial size [37]. Studies have shown that reduced distensibility of large
47
48 arteries parallel cardiac hypertrophy and remodeling in hypertensive patients [38,39]. Large
49
50 artery stiffness may increase the work load on the heart similar to volume overload and may thus
51
52 represent one of the mechanisms by which hypertension leads to eccentric hypertrophy and left
53
54 atrial enlargement [39]. In a LIFE sub-study, there was a significant correlation between baseline
55
56
57
58
59
60

1
2
3 brachial pulse pressure and left atrial size, independent of age, gender and body surface area
4
5 [40]. Furthermore, there is evidence for linking brachial pulse pressure to microvascular damage
6
7 in the heart and other target organs, which again may lead to increased peripheral resistance and
8
9 blood pressure, further increasing arterial stiffness and central pulse pressure. Increased central
10
11 pulse pressure may then further damage small arteries and lead to LVH [41]. Studies have found
12
13 brachial pulse pressure to be a powerful predictor of cardiovascular morbidity and mortality [42-
14
15 49], and the predictive effect increases with age [46-48]. The present study evaluated ISH in
16
17 parallel with brachial pulse pressure and not central pulse pressure. Non-invasive central pulse
18
19 pressure has been shown to better predict cardiovascular outcomes than brachial pulse pressure,
20
21 and to be closer associated with extent of atherosclerosis (carotid plaque burden and intimal-
22
23 medial thickness, and vascular mass) [50].
24
25
26
27

28
29 Some strengths and limitations of the present study should be mentioned. This is a
30
31 post hoc analysis of data collected prospectively in a randomized clinical trial. Atrial
32
33 fibrillation was a pre-specified secondary endpoint and the ISH patients defined as a subgroup
34
35 of patients of special interest. In the present analysis a strict and conservative approach was
36
37 used in as much as new-onset AF was ascertained on yearly study ECGs. This ensured
38
39 objective documentation of all incident cases of AF, however, the true incidence may have
40
41 been underestimated by missing possible cases of paroxysmal AF. The patients recruited in
42
43 the LIFE study all had ECG-LVH and results cannot automatically be extrapolated to
44
45 hypertensive patients at lower risk. Also we excluded patients with AF at baseline known to
46
47 have especially high cardiovascular risk and strong benefit of SBP reduction. Such exclusion
48
49 of study participants confounded the relationship between the reduction of SBP and all-cause
50
51 mortality in the ISH group due to the higher prevalence of AF at baseline in this group.
52
53
54
55

56
57 Further, we investigated a fairly homogenous study population of mostly Caucasian
58
59 patients with LVH. Systolic blood pressure at baseline when patients were in the untreated
60

1
2
3 state was rather strongly predicting new-onset AF despite quite aggressive treatment and
4
5 lowering of SBP by about 30 mmHg throughout. Blood pressures were taken standardized
6
7 and regularly which allowed calculation of in-treatment SBP, also strongly predicting less
8
9 new-onset AF. Certainly, statistical interference happened between SBP at baseline, changes
10
11 in SBP during treatment and achieved in-treatment SBP during the study. However, despite
12
13 these rather complicated relations between various SBPs, adjusted for in multivariate models,
14
15 quite strong relationships appeared between SBP and risk for new-onset AF and benefits of
16
17 treatment.
18
19
20
21

22 **Conclusions**

23
24
25 The impact of in-treatment SBP in preventing incidence AF was stronger in patients with ISH
26
27 compared to non-ISH patients in the same population. This finding is in line with our previous
28
29 findings showing benefits of BP lowering therapy and regression of ECG-LVH on preventing
30
31 cardiovascular mortality, myocardial infarction, cerebral stroke and heart failure in LIFE
32
33 study participants with ISH. Overall our findings are principally the same in ECG-LVH
34
35 hypertensive patients with and without ISH.
36
37
38
39

40 **Disclosure statement**

41
42 The LIFE study (Losartan Intervention For Endpoint reduction in hypertension) was
43
44 originally sponsored by Merck and Co. Inc., Whitehouse Station, NJ, USA. Outside the
45
46 present work SEK has received ad hoc honoraria for lecturing from [Bayer](#), Merck KGaA,
47
48 Merck & Co., Sanofi, and Takeda, and honoraria from Takeda for study committee work
49
50 within the past 3 years. Richard B. Devereux has received honoraria and grant from Merck.
51
52 Kristian Wachtell has received honoraria from Merck. The other authors report no conflicts.
53
54
55

56 **Funding**

57
58 The study was supported by grant C=Z-368 from Merck and Co., Inc, West Point, PA, USA.
59
60

References

1. Kannel WB, Gordon T, Schwartz MJ. Systolic versus diastolic blood pressure and risk of coronary heart disease. The Framingham study. Am J Cardiol 1971; 27: 335-346.
2. Kannel WB, Wolf PA, McGee DL et al. Systolic blood pressure, arterial rigidity, and risk of stroke. The Framingham study. JAMA. 1981; 245: 1225-1229.
3. SHEP Cooperative Research Group. Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension. JAMA. 1991; 265: 3255-3264.
4. Dahlöf B, Devereux RB, Kjeldsen SE et al for the LIFE Study Group. Cardiovascular morbidity and mortality in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomised trial against atenolol. Lancet. 2002; 359: 995-1003.
5. Wachtell K, Lehto M, Gerds E et al. Angiotensin II receptor blockade reduces new-onset atrial fibrillation and subsequent stroke compared to atenolol: The LIFE Study. JACC. 2005; 45: 712-719.
6. Okin PM, Wachtell K, Devereux RB et al for the LIFE Study Investigators. Regression of Electrocardiographic Left Ventricular Hypertrophy and Decreased Incidence of New-Onset Atrial Fibrillation in Hypertensive Patients: The LIFE Study. JAMA. 2006; 296: 1242-1248.
7. Okin PM, Wachtell K, Kjeldsen SE et al. Incidence of atrial fibrillation in relation to changing heart rate over time in hypertensive patients: The LIFE Study. Circ Arrhythmia Electrophysiol. 2008; 1: 337-343.
8. ~~Kannel WB, Gordon T, Schwartz MJ. Systolic versus diastolic blood pressure and risk of coronary heart disease. The Framingham study. Am J Cardiol 1971; 27: 335-346.~~
9. ~~Kannel WB, Wolf PA, McGee DL et al. Systolic blood pressure, arterial rigidity, and risk of stroke. The Framingham study. JAMA. 1981; 245: 1225-1229.~~
10. ~~SHEP Cooperative Research Group. Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension. JAMA. 1991; 265: 3255-3264.~~

- 1
2
3 11. Kjeldsen SE, Dahlöf B, Devereux RB et al for the LIFE Study Group. Effects of losartan
4 on cardiovascular morbidity and mortality in patients with isolated systolic hypertension
5 and left ventricular hypertrophy. A losartan intervention for endpoint reduction (LIFE)
6 and left ventricular hypertrophy. A losartan intervention for endpoint reduction (LIFE)
7 substudy. JAMA. 2002; 288: 1491-1498.
8
9
- 10 12. Larstorp ACK, Okin PM, Devereux RB et al. Regression of Electrocardiographic Left
11 Ventricular Hypertrophy during Antihypertensive Therapy and the Prediction of Major
12 Cardiovascular Events in Patients with Isolated Systolic Hypertension. J Hum
13 Hypertens. 2011; 25: 178-185.
14
- 15 13. Larstorp ACK, Okin PM, Devereux RB et al. Reduced ECG-LVH during
16 Antihypertensive Therapy is Associated with less New-Onset Heart Failure and
17 Mortality in Patients with Isolated Systolic Hypertension. The LIFE Study. Am J
18 Hypertens. 2012; 25: 1101-1109.
19
- 20 14. Larstorp ACK, Ariansen I, Gjesdal K et al. Association of pulse pressure with new-onset
21 atrial fibrillation in hypertensive patients with left ventricular hypertrophy: the LIFE
22 Study. Hypertension. 2012; 60: 347-353.
23
- 24 15. Dahlöf B, Devereux R, de Faire U et al. The Losartan Intervention For Endpoint
25 reduction (LIFE) in Hypertension study: rationale, design, and methods. The LIFE Study
26 Group. Am J Hypertens. 1997; 10: 705-713.
27
- 28 16. Dahlöf B, Devereux RB, Julius S et al. Characteristics of 9194 patients with left
29 ventricular hypertrophy: the LIFE study. Losartan Intervention For Endpoint Reduction
30 in Hypertension. Hypertension. 1998; 32: 989-997.
31
- 32 17. Kjeldsen SE, Dahlof B, Devereux RB et al. Lowering of blood pressure and predictors of
33 response in patients with left ventricular hypertrophy: the LIFE study. Losartan
34 Intervention For Endpoint. Am J Hypertens. 2000; 13: 899-906.
35
- 36 18. Molloy TJ, Okin PM, Devereux RB, Kligfield P. Electrocardiographic detection of left
37 ventricular hypertrophy by the simple QRS voltage-duration product. J Am Coll Cardiol.
38 1992; 20: 1180-1186.
39
- 40 19. Okin PM, Roman MJ, Devereux RB, Kligfield P. Electrocardiographic identification of
41 increased left ventricular mass by simple voltage-duration products. J Am Coll Cardiol.
42 1995; 25: 417-423.
43
- 44 20. Okin PM, Roman MJ, Devereux RB et al. Time-Voltage QRS Area of the 12-Lead
45 Electrocardiogram: Detection of Left Ventricular Hypertrophy. Hypertension. 1998; 31:
46 937-942.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
- 2
- 3 21. Sokolow M, Lyon TP. The ventricular complex in left ventricular hypertrophy as
- 4 obtained by unipolar precordial and limb leads. *Am Heart J.* 1949; 37: 161-186.
- 5
- 6 22. Cox DR. Regression Models and Life-Tables. *J R Stat Soc, B.* 1972;34:187-220.
- 7
- 8 23. Kalbfleisch JD, Prentice RL. The Statistical Analysis of Failure Time Data. New York,
- 9 NY: John Wiley & Sons, 1980.
- 10
- 11 24. Kaplan EL, Meier P. Nonparametric Estimation from Incomplete Observations. *J Am*
- 12 *Stat Assoc.* 1958;53:457-481.
- 13
- 14 25. Lloyd-Jones DM, Wang TJ, Leip EP et al. Lifetime Risk for Development of Atrial
- 15 Fibrillation: The Framingham Heart Study. *Circulation.* 2004;110:1042-1046.
- 16
- 17 26. Roger VL, Go AS, Lloyd-Jones DM et al. Heart Disease and Stroke Statistics-2012
- 18 Update: A Report From the American Heart Association. *Circulation.* 2012;125:e2-e220.
- 19
- 20 27. Heeringa J, van der Kuip DA, Hofman A et al. Prevalence, incidence and lifetime risk of
- 21 atrial fibrillation: the Rotterdam study. *Eur Heart J.* 2006;27:949-953.
- 22
- 23 28. Khairallah F, Ezzedine R, Ganz LI et al. Epidemiology and determinants of outcome of
- 24 admissions for atrial fibrillation in the United States from 1996 to 2001. *Am J Cardiol.*
- 25 2004;94:500-504.
- 26
- 27 29. Benjamin EJ, Levy D, Vaziri SM et al. Independent risk factors for atrial fibrillation in a
- 28 population-based cohort. The Framingham Heart Study. *JAMA.* 1994;271:840-844.
- 29
- 30 30. Psaty BM, Manolio TA, Kuller LH et al. Incidence of and Risk Factors for Atrial
- 31 Fibrillation in Older Adults. *Circulation.* 1997;96:2455-2461.
- 32
- 33 31. Wang TJ, Parise H, Levy D et al. Obesity and the risk of new-onset atrial fibrillation.
- 34 *JAMA.* 2004;292:2471-2477.
- 35
- 36 32. Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for
- 37 stroke: the Framingham Study. *Stroke.* 1991;22:983-988.
- 38
- 39 33. Go AS. The epidemiology of atrial fibrillation in elderly persons: the tip of the iceberg.
- 40 *Am J Ger Cardiol.* 2005;14:56-61.
- 41
- 42 34. Benjamin EJ, Wolf PA, D'Agostino RB et al. Impact of atrial fibrillation on the risk of
- 43 death: the Framingham Heart Study. *Circulation.* 1998;98:946-952.
- 44
- 45 35. Aksnes TA, Flaa A, Strand A, Kjeldsen SE. Prevention of new-onset atrial fibrillation
- 46 and its predictors with angiotensin II-receptor blockers in the treatment of hypertension
- 47 and heart failure. *J Hypertens.* 2007;25:15-23.
- 48
- 49 36. Huxley RR, Lopez FL, Folsom AR et al. Absolute and Attributable Risks of Atrial
- 50 Fibrillation in Relation to Optimal and Borderline Risk Factors. *Circulation.*
- 51 2011;123:1501-1508.
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
2
3 37. Aviv A. Hypothesis: pulse pressure and human longevity. *Hypertension*. 2001;37:1060-
4 1066.
- 5
6 38. Darne B, Girerd X, Safar M et al. Pulsatile versus steady component of blood pressure: a
7 cross-sectional analysis and a prospective analysis on cardiovascular mortality.
8 *Hypertension*. 1989;13:392-400.
- 9
10 39. Franklin SS, Sutton-Tyrrell K, Belle SH et al. The importance of pulsatile components
11 of hypertension in predicting carotid stenosis in older adults. *J Hypertens*. 1997;15:1143-
12 1150.
- 13
14 40. Vaziri SM, Larson MG, Lauer MS et al. Influence of blood pressure on left atrial size.
15 The Framingham Heart Study. *Hypertension*. 1995;25:1155-1160.
- 16
17 41. Bouthier JD, De Luca N, Safar ME, Simon AC. Cardiac hypertrophy and arterial
18 distensibility in essential hypertension. *Am Heart J*. 1985;109:1345-1352.
- 19
20 42. Boutouyrie P, Laurent S, Girerd X et al. Common Carotid Artery Stiffness and Patterns
21 of Left Ventricular Hypertrophy in Hypertensive Patients. *Hypertension*. 1995;25:651-
22 659.
- 23
24 43. Gerds E, Papademetriou V, Palmieri V et al. Correlates of pulse pressure reduction
25 during antihypertensive treatment (losartan or atenolol) in hypertensive patients with
26 electrocardiographic left ventricular hypertrophy (the LIFE study). *Am J Cardiol*.
27 2002;89:399-402.
- 28
29 44. Laurent S, Briet M, Boutouyrie P. Large and small artery cross-talk and recent
30 morbidity-mortality trials in hypertension. *Hypertension*. 2009;54:388-392.
- 31
32 45. Benetos A, Rudnichi A, Safar M, Guize L. Pulse pressure and cardiovascular mortality
33 in normotensive and hypertensive subjects. *Hypertension*. 1998;32:560-564.
- 34
35 46. Domanski MJ, Davis BR, Pfeffer MA et al. Isolated systolic hypertension : prognostic
36 information provided by pulse pressure. *Hypertension*. 1999;34:375-380.
- 37
38 47. Millar JA, Lever AF. Implications of pulse pressure as a predictor of cardiac risk in
39 patients with hypertension. *Hypertension*. 2000;36:907-911.
- 40
41 48. Franklin SS, Khan SA, Wong ND et al. Is pulse pressure useful in predicting risk for
42 coronary heart disease? The Framingham Heart Study. *Circulation*. 1999;100:354-360.
- 43
44 49. Vaccarino V, Berger AK, Abramson J et al. Pulse pressure and risk of cardiovascular
45 events in the systolic hypertension in the elderly program. *Am J Cardiol*. 2001;88:980-
46 986.
- 47
48 50. Glynn RJ, Chae CU, Guralnik JM et al. Pulse pressure and mortality in older people.
49 *Arch Intern Med*. 2000;160:2765-2772.
- 50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 51. Khattar RS, Swales JD, Dore C et al. Effect of aging on the prognostic significance of
4 ambulatory systolic, diastolic, and pulse pressure in essential hypertension. *Circulation*.
5 2001;104:783-789.
6
7
8 52. Fyhrquist F, Dahlof B, Devereux RB et al for the LIFE Study Group. Pulse pressure and
9 effects of losartan or atenolol in patients with hypertension and left ventricular
10 hypertrophy. *Hypertension*. 2005;45:580-585.
11
12 53. Roman MJ, Devereux RB, Kizer JR et al. Central pressure more strongly relates to
13 vascular disease and outcome than does brachial pressure: the Strong Heart Study.
14 *Hypertension*. 2007;50:197-203.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Figure Legends

Figure 1 Flow [diagram-chart](#) of the patients who participated in the present study

Figure 2 Development of systolic blood pressure (Fig. 2a, upper panel) and diastolic blood pressure (Fig. 2b, lower panel) in the ISH patients and the non-ISH patients who had incident AF and the patients who did not have incident AF. [The y-axes have been truncated.](#)

Figure 3 Cornell product as indicator of ECG-LVH at baseline and yearly during the study in the ISH patients and the non-ISH patients with new-onset AF (upper lines) and in patients without new-onset AF during the study (lower lines). [The y-axis has been truncated.](#)

Fig. 1




Fig. 2a




Fig. 2b





Fig. 3



Peer Review Only

Table 1. Baseline demographic and clinical characteristics in relation to new-onset atrial fibrillation in 1,248 patients with isolated systolic hypertension

Variables	Sinus Rhythm (n=1,187)	New-Onset Atrial Fibrillation (n=61)	<i>P</i> Value
Age, yr.	70.1 (6.4)	71.9 (5.4)	0.03
Sex, % female	60.6	52.5	0.21
Race, % black	6.5	1.6	0.17
Treatment with losartan, %	50.9	36.1	0.024
Diabetes, %	17.3	14.8	0.61
History of ischaemic heart disease, %	19.6	23.0	0.53
History of myocardial infarction, %	8.5	9.8	0.72
History of heart failure, %	2.1	4.9	0.15
History of cerebrovascular disease, %	11.4	11.5	0.98
History of TIA, %	5.8	4.9	1.00
History of peripheral vascular disease, %	8.1	6.6	0.81
Current smokers, %	15.4	6.7	0.064
Weight, kg	75.1 (14.7)	77.1 (13.7)	0.31
Height, cm	165.2 (9.3)	167.9 (8.6)	0.031
Body mass index, kg/m ²	27.5 (4.9)	27.4 (4.9)	0.89
Systolic blood pressure, mmHg	174.1 (10.8)	179.0 (11.5)	0.001
Diastolic blood pressure, mmHg	82.7 (5.8)	83.4 (4.8)	0.34
Pulse pressure, mmHg	91.4 (11.7)	95.6 (11.1)	0.006
Heart rate, beats/min	71.7 (10.8)	70.8 (11.3)	0.53
Sokolow-Lyon voltage, mm	30.8 (10.2)	33.7 (12.0)	0.03
Cornell voltage-duration product, mm x msec	2772 (1053)	2920 (1063)	0.29
Hemoglobin, mmol/L	138.9 (12.4)	138.8 (13.0)	0.97
Sodium, mmol/L	140.3 (2.7)	140.4 (2.3)	0.91
Potassium, mmol/L	4.2 (0.4)	4.1 (0.4)	0.018
Creatinine, mmol/L	88.0 (21.1)	84.0 (16.0)	0.15
Total cholesterol, mmol/L	5.98 (1.15)	5.89 (1.08)	0.56
HDL cholesterol, mmol/L	1.49 (0.44)	1.52 (0.44)	0.64
Uric acid, μmol/L	324 (77)	329 (77)	0.58
Glucose, mmol/L	6.22 (2.46)	6.17 (2.14)	0.90
Urine albumin/creatinine ratio, mg/μmol	6.2 (22.2)	5.2 (12.5)	0.74
Framingham risk score, %	23.1 (9.9)	24.0 (9.8)	0.47

Data are presented as mean (SD) unless otherwise indicated. Abbreviations: TIA, transitory ischaemic attack; HDL, high density lipoprotein.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Table 2a. Predictors of new-onset atrial fibrillation in 1,248 patients with isolated systolic hypertension

Univariate Predictor Variable	Hazard Ratio (95 % CI)	P Value
Baseline systolic BP (per 1 mm-Hg increase)	1.04 (1.01-1.06)	0.001
Height (per 1 cm increase)	1.04 (1.01-1.06)	0.014
Baseline Sokolow-Lyon (per 10.5 -mm increase)	1.36 (1.06-1.75)	0.015
Potassium (per 1 mmol/L increase)	0.39 (0.18-0.86)	0.019
Treatment (atenolol vs. losartan)	1.84 (1.09-3.11)	0.022
Time-varying heart rate (per 1 beat/min increase)	1.025 (1.003-1.047)	0.023
Age (per 1 yr. increase)	1.05 (1.01-1.10)	0.029
Smoking (yes vs. no)	0.41 (0.15-1.14)	0.087
Time-varying Cornell product (per 1,050 mm x msec increase)	1.17 (0.96-1.43)	0.11
Gender (male vs. female)	1.49 (0.90-2.48)	0.12

Table 2b. Predictors of new-onset atrial fibrillation in 7,583 patients with non-isolated systolic hypertension

Univariate Predictor Variable	Hazard Ratio (95 % CI)	P Value
Age (per 1 yr. increase)	1.10 (1.08-1.12)	<0.001
Baseline pulse pressure (per 1 mmHg increase)	1.02 (1.01-1.03)	<0.001
Framingham risk score (per 1 percentage point increase)	1.02 (1.01-1.03)	<0.001
Time-varying heart rate (per 1 beat/min increase)	1.03 (1.02-1.04)	<0.001
Log urine albumin-creatinine ratio (mg/mmol)	1.39 (1.17-1.65)	<0.001
Time-varying Cornell product (per 1,050 mm x msec increase)	1.17 (1.06-1.29)	0.001
Baseline systolic BP (per 1 mmHg increase)	1.01 (1.00-1.02)	0.002
Treatment (atenolol vs. losartan)	1.43 (1.13-1.80)	0.003
Baseline Cornell product (per 1,050 mm x msec increase)	1.16 (1.05-1.28)	0.005
Race (black)	0.31 (0.13-0.74)	0.009
Baseline cholesterol (per 1 mmol/L increase)	0.88 (0.79-0.98)	0.015
Baseline diastolic BP (per 1 mmHg increase)	0.981 (0.963-0.999)	0.036

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

<u>Gender (male vs. female)</u>	<u>0.81 (0.64-1.02)</u>	<u>0.066</u>
<u>History of coronary artery disease (yes vs. no)</u>	<u>1.27 (0.94-1.72)</u>	<u>0.120</u>

For Peer Review Only

Table 3. Time dependent (time to event) multivariate Cox regression analyses to assess the risk of incident atrial fibrillation related to in-treatment variation of 10 mm Hg in systolic BP in patients with isolated systolic hypertension and non- isolated systolic hypertension

In-treatment SBP per 10 mm Hg decrease	ISH (n=1,248)			Non-ISH (n=7,583)		
	Hazard Ratio	95 % CI	P Value	Hazard Ratio	95 % CI	P Value
Univariate	0.87	0.75-1.01	0.070	0.96	0.89-1.02	0.19
Adjusted for treatment	0.87	0.76-1.01	0.066	0.95	0.89-1.02	0.16
Adjusted for baseline SBP	0.82	0.71-0.95	0.008	0.92	0.86-0.99	0.023
Multivariate Model 1	0.87	0.75-1.00	0.054	0.93	0.87-1.00	0.043
Multivariate Model 2	0.83	0.73-0.95	0.008	0.91	0.85-0.97	0.006

Model 1: Adjusted for treatment effect, Framingham risk score, history of coronary heart disease, and in-treatment heart rate and in-treatment ECG-LVH determined by Cornell voltage-duration product

Model 2: Adjusted for baseline SBP, treatment effect, Framingham risk score, history of coronary heart disease, and in-treatment heart rate and in-treatment ECG-LVH determined by Cornell voltage-duration product

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Table 4. Time dependent (time to event) multivariate Cox regression analyses to assess the risk of the combined endpoint of incident atrial fibrillation or all-cause mortality related to in-treatment variation of 10 mm Hg in systolic BP in patients with isolated systolic hypertension and non- isolated systolic hypertension

In-treatment SBP per 10 mm Hg decrease	ISH (n=1,248)			Non-ISH (n=7,583)		
	Hazard Ratio	95 % CI	P Value	Hazard Ratio	95 % CI	P Value
Univariate	0.91	0.84-0.98	0.015	0.89	0.86-0.93	<0.001
Adjusted for treatment	0.92	0.85-0.99	0.021	0.90	0.86-0.93	<0.001
Adjusted for baseline SBP	0.94	0.87-1.02	0.133	0.91	0.88-0.95	<0.001
Multivariate Model 1	0.91	0.84-0.98	0.014	0.92	0.89-0.96	<0.001
Multivariate Model 2	0.93	0.87-1.01	0.087	0.93	0.89-0.96	<0.001

Model 1: Adjusted for treatment effect, Framingham risk score, history of coronary heart disease, and in-treatment heart rate and in-treatment ECG-LVH determined by Cornell voltage-duration product

Model 2: Adjusted for baseline SBP, treatment effect, Framingham risk score, history of coronary heart disease, and in-treatment heart rate and in-treatment ECG-LVH determined by Cornell voltage-duration product